



(51) International Patent Classification:

A61K 9/00 (2006.01) *A61K 47/12* (2006.01)
A61K 47/10 (2006.01)

(21) International Application Number:

PCT/US2014/072319

(22) International Filing Date:

24 December 2014 (24.12.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/921,361 27 December 2013 (27.12.2013) US

(71) Applicant: **CELGENE CORPORATION** [US/US]; 86
Morris Avenue, Summit, NJ 07901 (US).

(72) Inventors: **HUANG, Lianfeng**; 1 Wellington Drive, Bask-
ing Ridge, NJ 07920 (US). **HUI, Ho-Wah**; 10 Heritage
Court, Basking Ridge, NJ 07920 (US). **NARINGREKAR,**
Vijay; 12 Courtside Lane, Princeton, NJ 08540 (US).
YANG, Gang; 56 Huntley Way, Bridgewater, NJ 08807
(US).

(74) Agents: **BRITVA, Irina** et al.; Jones Day, 222 East 41st
Street, New York, NY 10017-6702 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: ROMIDEPSIN FORMULATIONS AND USES THEREOF

(57) Abstract: Provided herein are liquid concentrate formulations of romidepsin. Also provided are methods for producing these formulations and uses thereof. In one embodiment, the formulation comprises romidepsin, polyethylene glycol, ethanol and a citrate buffer. In another embodiment, the formulation comprises romidepsin, polyethylene glycol, ethanol and an acetate buffer.



WO 2015/100399 A1

ROMIDEPSIN FORMULATIONS AND USES THEREOF**CROSS REFERENCE TO RELATED APPLICATIONS****[0001]**

This application claims the benefit of priority to U.S. Provisional Patent Application Serial No. 61/921,361 filed December 27, 2013, the disclosure of which is incorporated by reference herein in its entirety.

FIELD

[0002] Provided herein are liquid formulations of romidepsin. Also provided are methods for producing these formulations and uses thereof.

BACKGROUND

[0003] Cancer is a major public health problem in the United States and in the world. Currently, one in 4 deaths in the United States is due to cancer. Each year, the American Cancer Society estimates the numbers of new cancer cases and deaths expected in the United States in the current year and compiles the most recent data on cancer incidence, mortality, and survival based on incidence data from the National Cancer Institute, the Centers for Disease Control and Prevention, and the North American Association of Central Cancer Registries and mortality data from the National Center for Health Statistics. A total of 1,596,670 new cancer cases and 571,950 deaths from cancer were projected to occur in the United States in 2011. Overall cancer incidence rates were stable since late 1990s. The reduction in the overall cancer death rates since 1990 in men and 1991 in women translated to the avoidance of about 898,000 deaths from cancer. Despite an obvious progress, approximately 560,000 people died of cancer in 2006 in the United States alone. Aging of the general population and development of new forms of cancer contribute to the problem.

[0004] Romidepsin has been shown to have anticancer activities. The drug is approved in the U.S. for treatment of cutaneous T-cell lymphoma (CTCL) and peripheral T-cell lymphoma (PTCL), and is currently being tested, for example, for use in treating patients with other hematological malignancies (*e.g.*, multiple myeloma, *etc.*) and solid tumors (*e.g.*, prostate cancer, pancreatic cancer, *etc.*). It is thought to act by selectively inhibiting deacetylases (*e.g.*, histone deacetylase, tubulin deacetylase), promising new targets for development of a new class of anti-cancer therapies (Bertino & Otterson, *Expert Opin Investig Drugs* 20(8):11151-1158, 2011). One mode of action involves the inhibition of one or more classes of histone deacetylases (HDAC).

[0005] As cancer remains a major worldwide public health problem, there is a continued need for effective therapies to treat cancer.

SUMMARY

[0006] In one embodiment, provided herein is a romidepsin formulation. In one embodiment, the formulation is a liquid concentrate formulation. In one embodiment, the formulation is a liquid concentrate formulation for dilution. In one embodiment, a liquid concentrate formulation for dilution is formulated in a solvent system. In one embodiment, the solvent system comprises a citrate buffer. In another embodiment, the solvent system comprises an acetate buffer. In one embodiment, a romidepsin formulation is an injectable formulation.

[0007] In one embodiment, provided herein are methods to treat proliferative diseases using a romidepsin formulation provided herein. In some embodiments, provided herein are methods to treat cancer. In some embodiments, cancers include, but are not limited to, carcinomas, sarcomas, leukemias, lymphomas and the like. In certain embodiments, the cancer is a hematological malignancy. In certain embodiments, the cancer is a solid tumor.

[0008] In one embodiment, provided are methods of producing a romidepsin formulation.

DETAILED DESCRIPTION

Definitions

[0009] As used in the specification and the accompanying claims, the indefinite articles “a” and “an” and the definite article “the” include plural as well as singular referents, unless the context clearly dictates otherwise.

[0010] As used herein, and unless otherwise specified, the term “about” or “approximately” means an acceptable error for a particular value as determined by one of ordinary skill in the art, which depends in part on how the value is measured or determined. In certain embodiments, the term “about” or “approximately” means within 1, 2, 3, or 4 standard deviations. In certain embodiments, the term “about” or “approximately” means within 30%, 25%, 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.1%, or 0.05% of a given value or range.

[0011] As used herein, and unless otherwise specified, the terms “treat,” “treating” and “treatment” refer to the eradication or amelioration of a disease or disorder, or of one or more symptoms associated with the disease or disorder. In certain embodiments, the terms refer to minimizing the spread or worsening of the disease or disorder resulting from the administration of one or more prophylactic or therapeutic agents to a subject with such a disease or disorder. In some embodiments, the terms refer to the administration of a compound or dosage form provided herein, with or without one or more additional active agent(s), after the diagnosis or the onset of symptoms of the particular disease.

[0012] As used herein, and unless otherwise specified, the terms “prevent,” “preventing” and “prevention” refer to the prevention of the onset, recurrence or spread of a disease or disorder, or of one or more symptoms thereof. In certain embodiments, the terms refer to the treatment with or administration of a compound or dosage form provided herein, with or without one or more other additional active agent(s), prior to the onset of symptoms, particularly to subjects at risk of disease or disorders provided herein. The terms encompass the inhibition or reduction of a symptom of the particular disease. In certain embodiments, subjects with familial history of a disease are potential candidates for preventive regimens. In certain embodiments, subjects who have a history of recurring symptoms are also potential candidates for prevention. In this regard, the term “prevention” may be interchangeably used with the term “prophylactic treatment.”

[0013] As used herein, and unless otherwise specified, the terms “manage,” “managing” and “management” refer to preventing or slowing the progression, spread or worsening of a disease or disorder, or of one or more symptoms thereof. Often, the beneficial effects that a subject derives from a prophylactic and/or therapeutic agent do not result in a cure of the disease or disorder. In this regard, the term “managing” encompasses treating a subject who had suffered from the particular disease in an attempt to prevent or minimize the recurrence of the disease.

[0014] As used herein, and unless otherwise specified, “amelioration” of the symptoms of a particular disorder by administration of a particular pharmaceutical composition refers to any lessening, whether permanent or temporary, lasting or transient, that can be attributed to or associated with the administration of the composition.

[0015] As used herein, and unless otherwise specified, the term “therapeutically effective amount” or “effective amount” of a compound means an amount sufficient to

provide a therapeutic benefit in the treatment or management of a disease or disorder, or to delay or minimize one or more symptoms associated with the disease or disorder. A “therapeutically effective amount” or “effective amount” of a compound means an amount of therapeutic agent, alone or in combination with one or more other agent(s), which provides a therapeutic benefit in the treatment or management of the disease or disorder. The terms “therapeutically effective amount” and “effective amount” can encompass an amount that improves overall therapy, reduces, delays, or avoids symptoms or causes of disease or disorder, or enhances the therapeutic efficacy of another therapeutic agent.

[0016] As used herein, and unless otherwise specified, a “prophylactically effective amount” of a compound is an amount sufficient to prevent a disease or disorder, or prevent its recurrence. A prophylactically effective amount of a compound means an amount of therapeutic agent, alone or in combination with one or more other agent(s), which provides a prophylactic benefit in the prevention of the disease. The term “prophylactically effective amount” can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent.

[0017] As used herein, and unless otherwise specified, the term “subject” is defined herein to include animals such as mammals, including, but not limited to, primates (*e.g.*, humans), cows, sheep, goats, horses, dogs, cats, rabbits, rats, mice, and the like. In specific embodiments, the subject is a human. The terms “subject” and “patient” are used interchangeably herein in reference, for example, to a mammalian subject, such as a human. In particular embodiments, a subject having cancer is a subject who has been previously diagnosed as having cancer.

[0018] As used herein, and unless otherwise specified, “neoplasm” is an abnormal mass of tissue as a result of neoplasia. The growth of neoplastic cells exceeds and is not coordinated with that of the normal tissues around it. The growth persists in the same excessive manner even after cessation of the stimuli. It usually causes a lump or tumor. Neoplasms may be benign, pre-malignant (carcinoma in situ) or malignant (cancer).

[0019] As used herein, and unless otherwise specified, “tumor” refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues. As used herein, and unless otherwise specified, “neoplastic” refers to any form of deregulated or unregulated cell growth, whether malignant or benign,

resulting in abnormal tissue growth. Thus, “neoplastic cells” include malignant and benign cells having deregulated or unregulated cell growth.

[0020] As used herein, and unless otherwise specified, the terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include, but are not limited to, lymphoma, leukemia, and solid tumors, such as, for example, lung cancer. In one embodiment, the term “cancer” as used herein includes, but is not limited to, solid tumors and blood-borne tumors. The term “cancer” refers to disease of skin tissues, organs, blood, and vessels, including, but not limited to, cancers of the bladder, bone or blood, brain, breast, cervix, chest, colon, endometrium, esophagus, eye, head, kidney, liver, lymph nodes, lung, mouth, neck, ovaries, pancreas, prostate, rectum, stomach, testis, throat, and uterus. Specific cancers include, but are not limited to, advanced malignancy, amyloidosis, neuroblastoma, meningioma, atypical meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor prognosis malignant brain tumor, malignant glioma, recurrent malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi’s sarcoma, karyotype acute myeloblastic leukemia, Hodgkin’s lymphoma, non-Hodgkin’s lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, metastatic melanoma (localized melanoma, including, but not limited to, ocular melanoma), malignant mesothelioma, malignant pleural effusion mesothelioma syndrome, peritoneal carcinoma, papillary serous carcinoma, gynecologic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma, fibrodysplasia ossificans progressiva, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unresectable hepatocellular carcinoma, Waldenstrom’s macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer, papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, and leiomyoma. In a specific embodiment, the cancer is metastatic. In another embodiment, the cancer is refractory or resistant to chemotherapy or radiation.

[0021] As used herein, and unless otherwise specified, the term “proliferative” disorder or disease refers to unwanted cell proliferation of one or more subset of cells in a multicellular organism resulting in harm (*i.e.*, discomfort or decreased life expectancy) to the multicellular organism. For example, as used herein, proliferative disorder or disease includes neoplastic disorders and other proliferative disorders.

[0022] As used herein, and unless otherwise specified, the term “relapsed” refers to a situation where a subject, that has had a remission of cancer after a therapy, has a return of cancer cells.

[0023] As used herein, and unless otherwise specified, the term “refractory” or “resistant” refers to a circumstance where a subject, even after intensive treatment, has residual cancer cells in the body.

[0024] As used herein, and unless otherwise specified, the term “drug resistance” refers to the condition when a disease does not respond to the treatment of a drug or drugs. Drug resistance can be either intrinsic, which means the disease has never been responsive to the drug or drugs, or it can be acquired, which means the disease ceases responding to a drug or drugs that the disease had previously responded to. In certain embodiments, drug resistance is intrinsic. In certain embodiments, the drug resistance is acquired.

[0025] As used herein, and unless otherwise specified, the term “anticancer agent” or “cancer therapeutic agent” is meant to include histone deacetylase (HDAC) inhibitors, including, but not limited to, romidepsin, anti-proliferative agents and chemotherapeutic agents, including, but not limited to, antimetabolites (*e.g.*, 5-fluoro uracil, methotrexate, fludarabine, cytarabine (also known as cytosine arabinoside or Ara-C), and high dose cytarabine), antimicrotubule agents (*e.g.*, vinca alkaloids, such as vincristine and vinblastine; and taxanes, such as paclitaxel and docetaxel), alkylating agents (*e.g.*, mechlorethamine, chlorambucil, cyclophosphamide, melphalan, ifosfamide, carmustine, azacitidine, decitabine, busulfan, cyclophosphamide, dacarbazine, ifosfamide, and nitrosoureas, such as carmustine, lomustine, bischloroethylnitrosurea, and hydroxyurea), platinum agents (*e.g.*, cisplatin, carboplatin, oxaliplatin, satraplatin (JM-216), and CI-973), anthracyclines (*e.g.*, doxorubicin and daunorubicin), antitumor antibiotics (*e.g.*, mitomycin, bleomycin, idarubicin, adriamycin, daunomycin (also known as daunorubicin, rubidomycin, or cerubidine), and mitoxantrone), topoisomerase inhibitors (*e.g.*, etoposide and camptothecins), purine antagonists or pyrimidine antagonists (*e.g.*, 6-mercaptopurine, 5-fluorouracil, cytarabine,

clofarabine, and gemcitabine), cell maturing agents (*e.g.*, arsenic trioxide and tretinoin), DNA repair enzyme inhibitors (*e.g.*, podophyllotoxines, etoposide, irinotecan, topotecan, and teniposide), enzymes that prevent cell survival (*e.g.*, asparaginase and pegaspargase), histone deacetylase inhibitors (*e.g.*, vorinostat), any other cytotoxic agents (*e.g.*, estramustine phosphate, dexamethasone, prednimustine, and procarbazine), hormones (*e.g.*, dexamethasone, prednisone, methylprednisolone, tamoxifen, leuprolide, flutamide, and megestrol), monoclonal antibodies (*e.g.*, gemtuzumab ozogamicin, alemtuzumab, rituximab, and yttrium-90-ibritumomab tiuxetan), immuno-modulators (*e.g.*, thalidomide and lenalidomide), Bcr-Abl kinase inhibitors (*e.g.*, AP23464, AZD0530, CGP76030, PD180970, SKI-606, imatinib, BMS354825 (dasatinib), AMN107 (nilotinib), and VX-680), hormone agonists or antagonists, partial agonists or partial antagonists, kinase inhibitors, surgery, radiotherapy (*e.g.*, gamma-radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and systemic radioactive isotopes), endocrine therapy, biological response modifiers (*e.g.*, interferons, interleukins, and tumor necrosis factor), hyperthermia and cryotherapy, and agents to attenuate any adverse effects (*e.g.*, antiemetics).

[0026] As used herein, and unless otherwise specified, the terms “co-administration” and “in combination with” include the administration of two or more therapeutic agents simultaneously, concurrently or sequentially within no specific time limits unless otherwise indicated. In one embodiment, the agents are present in the cell or in the subject’s body at the same time or exert their biological or therapeutic effect at the same time. In one embodiment, the therapeutic agents are in the same composition or unit dosage form. In other embodiments, the therapeutic agents are in separate compositions or unit dosage forms. In certain embodiments, a first agent can be administered prior to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), essentially concomitantly with, or subsequent to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapeutic agent.

[0027] As used herein, and unless otherwise specified, the terms “composition,” “formulation,” and “dosage form” are intended to encompass products comprising the specified ingredient(s) (in the specified amounts, if indicated), as well as any product(s)

which result, directly or indirectly, from combination of the specified ingredient(s) in the specified amount(s).

[0028] As used herein, and unless otherwise specified, the term “excipient” refers to a pharmacologically inactive substance used as a carrier for the active ingredient of a medication or as a bulking agent to allow for convenient and accurate dosage of an active ingredient.

[0029] As used herein, and unless otherwise specified, the term “pharmaceutically acceptable carrier,” “pharmaceutically acceptable excipient,” “physiologically acceptable carrier,” or “physiologically acceptable excipient” refers to a pharmaceutically-acceptable material, composition, or vehicle, such as a liquid or solid filler, diluent, excipient, solvent, or encapsulating material. In one embodiment, each component is “pharmaceutically acceptable” in the sense of being compatible with the other ingredients of a pharmaceutical formulation, and suitable for use in contact with the tissue or organ of humans and animals without excessive toxicity, irritation, allergic response, immunogenicity, or other problems or complications, commensurate with a reasonable benefit/risk ratio. In one embodiment, by “pharmaceutical” or “pharmaceutically acceptable” it is meant that any diluent(s), excipient(s) or carrier(s) in the composition, formulation, or dosage form are compatible with the other ingredient(s) and not deleterious to the recipient thereof. *See, e.g.,* Remington, *The Science and Practice of Pharmacy*, 21st Edition; Lippincott Williams & Wilkins: Philadelphia, PA, 2005; *Handbook of Pharmaceutical Excipients*, 5th Edition; Rowe *et al.*, ed., The Pharmaceutical Press and the American Pharmaceutical Association: 2005; and *Handbook of Pharmaceutical Additives*, 3rd Edition; Ash and Ash ed., Gower Publishing Company: 2007; *Pharmaceutical Preformulation and Formulation*, Gibson ed., CRC Press LLC: Boca Raton, FL, 2004.

[0030] As used herein, and unless otherwise specified, the term “pharmaceutically acceptable salts” is meant to include salts of active compounds which are prepared with relatively nontoxic acids. Acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the

salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (see, for example, Berge, *et al.* (1977) *J. Pharm. Sci.* 66:1-19).

[0031] A pharmaceutically acceptable salt form of a compound can be prepared *in situ* during the final isolation and purification of the compound, or separately by reacting the free base functionality with a suitable organic or inorganic acid. Examples of typical pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, and perchloric acid, or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid, or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts can include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, harnisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts can include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, lower alkyl sulfonate, and aryl sulfonate.

[0032] The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

[0033] As used herein, and unless otherwise specified, the terms, “polymorphs” and “polymorphic forms” and related terms refer to one of a variety of different crystal structures that can be adopted by a particular compound. In some embodiments, polymorphs occur when a particular chemical compound can crystallize in more than one structural arrangement. Different polymorphs may have different physical properties such as, for example, melting temperatures, heats of fusion, solubilities, dissolution rates and/or vibrational spectra as a result of the arrangement or conformation of the molecules in the crystal lattice. The differences in physical properties exhibited by polymorphs affect pharmaceutical parameters such as storage stability, compressibility and density (important in formulation and product manufacturing), and dissolution rates (an important factor in determining bioavailability). Differences in stability can result from changes in chemical reactivity (*e.g.*, differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical changes (*e.g.*, tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both (*e.g.*, tablets of one polymorph are more susceptible to breakdown at high humidity). As a result of solubility/dissolution differences, in the extreme case, some polymorphic transitions may result in lack of potency or, at the other extreme, toxicity. In addition, the physical properties of the crystal may be important in processing, for example, one polymorph might be more likely to form solvates or might be difficult to filter and wash free of impurities (*i.e.*, particle shape and size distribution might be different between one polymorph relative to the other).

[0034] As used herein, and unless otherwise specified, the term, “solvate” refers to a crystal form of a substance which contains solvent.

[0035] As used herein, and unless otherwise specified, the term “hydrate” refers to a crystal form adopted by a particular compound in which either a stoichiometric or non-stoichiometric amount of water is incorporated into the crystal lattice.

[0036] As used herein, and unless otherwise specified, the term “prodrug” refers to structurally modified forms of the compound that readily undergo chemical changes under physiological conditions to provide the compound. Additionally, prodrugs can be converted to the compound by chemical or biochemical methods in an *ex vivo* environment. Prodrugs are often useful because, in some situations, they may be easier to administer than the compound, or parent drug. They may, for instance, be bioavailable by oral administration

whereas the parent drug is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. A wide variety of prodrug derivatives are known in the art, such as those that rely on hydrolytic cleavage or oxidative activation of the prodrug. An example, without limitation, of a prodrug would be a compound which is administered as an ester (the “prodrug”), but then is metabolically hydrolyzed to the carboxylic acid, the active entity.

[0037] As used herein, and unless otherwise specified, the term “anhydrous” refers to a form of a compound that is substantially free of water. One of skill in the art will appreciate that an anhydrous solid can contain various amounts of residual water wherein that water is not incorporated in the crystalline lattice. Such incorporation of residual water can depend upon a compound’s hygroscopicity and storage conditions.

[0038] As used herein, and unless otherwise specified, the term “isostructural” or “isostructure” refers to two or more solid forms of a compound containing essentially the same three-dimensional arrangement of geometrically similar structural units. In some embodiments, “isostructural” forms show with similar or identical unit cell dimensions, the same space group, and similar or identical atomic coordinates for common atoms. In some embodiments, “isostructural” forms have the same structure, but not the same cell dimensions nor the same chemical composition, and have comparable variability in their atomic coordinates to that of the cell dimensions and chemical composition.

[0039] As used herein, and unless otherwise specified, the term “lyophilize” refers to the process of isolating a solid substance from solution and/or removal of solvent. In some embodiments, this may be achieved by various techniques known to one of skill in the art, including, for example, evaporation (*e.g.*, under vacuum, for example by rotary evaporation), freeze drying, and/or freezing the solution and vaporizing frozen solvent away under vacuum conditions, *etc.*

[0040] As used herein, and unless otherwise specified, the term “parenteral” includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques.

[0041] As used herein, and unless otherwise specified, the term “substantially free of” means containing no more than an insignificant amount. In some embodiments, a composition or preparation is “substantially free of” a recited element if it contains less than 5%, 4%, 3%, 2%, or 1%, by weight of the element. In some embodiments, the composition

or preparation contains less than 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1% or less of the recited element. In some embodiments, the composition or preparation contains an undetectable amount of the recited element.


[0042] As used herein, and unless otherwise specified, the expression “unit dose” refers to a physically discrete unit of a formulation appropriate for a subject to be treated (*e.g.*, for a single dose); each unit containing a predetermined quantity of an active agent selected to produce a desired therapeutic effect (it being understood that multiple doses may be required to achieve a desired or optimum effect), optionally together with a pharmaceutically acceptable carrier, which may be provided in a predetermined amount. The unit dose may be, for example, a volume of liquid (*e.g.*, an acceptable carrier) containing a predetermined quantity of one or more therapeutic agents, a predetermined amount of one or more therapeutic agents in solid form, a sustained release formulation or drug delivery device containing a predetermined amount of one or more therapeutic agents, etc. It will be appreciated that a unit dose may contain a variety of components in addition to the therapeutic agent(s). For example, acceptable carriers (*e.g.*, pharmaceutically acceptable carriers), diluents, stabilizers, buffers, preservatives, etc., may be included as described *infra*. It will be understood, however, that the total daily usage of a formulation of the present disclosure will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular subject or organism may depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of specific active compound employed; specific composition employed; age, body weight, general health, sex and diet of the subject; time of administration, and rate of excretion of the specific active compound employed; duration of the treatment; drugs and/or additional therapies used in combination or coincidental with specific compound(s) employed, and like factors well known in the medical arts.

[0043] As used herein, and unless otherwise specified, a compound described herein is intended to encompass all possible stereoisomers, unless a particular stereochemistry is specified. Where structural isomers of a compound are interconvertible *via* a low energy barrier, the compound may exist as a single tautomer or a mixture of tautomers. This can take the form of proton tautomerism; or so-called valence tautomerism in the compound.

Romidepsin

[0044] Romidepsin is a natural product which was isolated from *Chromobacterium violaceum* by Fujisawa Pharmaceuticals (Published Japanese Patent Application Hei 7 (1995)-64872; and U.S. Patent 4,977,138, issued December 11, 1990, each of which is incorporated herein by reference). Various preparations and purifications of romidepsin are described in PCT Publication WO 02/20817, which is incorporated herein by reference.

[0045] Romidepsin is a bicyclic peptide consisting of four amino acid residues (D-valine, D-cysteine, dehydrobutyrine, and L-valine) and a novel acid (3-hydroxy-7-mercapto-4-heptenoic acid), which contains both amide and ester bonds. Romidepsin can be obtained from *C. violaceum* using fermentation. It can also be prepared by synthetic or semi-synthetic means. The total synthesis of romidepsin reported by Kahn *et al.* (*J. Am. Chem. Soc.* 118:7237-7238, 1996) involves 14 steps and yields romidepsin in 18% overall yield. The structure of romidepsin is shown below (formula I):




[0046] Romidepsin has been shown to have anti-microbial, immunosuppressive, and anti-tumor activities. In the US, it is approved for the treatment of patients with cutaneous T-cell lymphoma (CTCL) and peripheral T-cell lymphoma (PTCL). It is currently being tested for multiple myeloma and solid tumors (*e.g.*, prostate cancer, pancreatic cancer, *etc.*) and is thought to act by selectively inhibiting deacetylases (*e.g.*, histone deacetylase, tubulin deacetylase) (Nakajima *et al.*, *Exp Cell Res* 241:126-133, 1998). One mode of action of romidepsin involves the inhibition of one or more classes of histone deacetylases (HDAC). Preparations and purification of romidepsin is described, for example, in U.S. Patent 4,977,138 and International PCT Application Publication WO 02/20817, each of which is incorporated herein by reference.

[0047] Exemplary forms of romidepsin include, but are not limited to, salts, esters, pro-drugs, isomers, stereoisomers (*e.g.*, enantiomers, diastereomers), tautomers, protected forms, reduced forms, oxidized forms, derivatives, and combinations thereof, with the desired

activity (*e.g.*, deacetylase inhibitory activity, aggressive inhibition, cytotoxicity). In certain embodiments, romidepsin is a pharmaceutical grade material and meets the standards of the U.S. Pharmacopoeia, Japanese Pharmacopoeia, or European Pharmacopoeia. In certain embodiments, the romidepsin is at least 95%, at least 98%, at least 99%, at least 99.9%, or at least 99.95% pure. In certain embodiments, the romidepsin is at least 95%, at least 98%, at least 99%, at least 99.9%, or at least 99.95% monomeric. In certain embodiments, no impurities are detectable in the romidepsin materials (*e.g.*, oxidized material, reduced material, dimerized or oligomerized material, side products, *etc.*). Romidepsin typically includes less than 1.0%, less than 0.5%, less than 0.2%, or less than 0.1% of total other unknowns. The purity of romidepsin may be assessed by appearance, HPLC, specific rotation, NMR spectroscopy, IR spectroscopy, UV/Visible spectroscopy, powder x-ray diffraction (XRPD) analysis, elemental analysis, LC-mass spectroscopy, or mass spectroscopy.

[0048] In one embodiment, the formulation contains a derivative of romidepsin.

[0049] In one embodiment, the derivative of romidepsin is of the formula (II):



(II)

wherein

n is 1, 2, 3 or 4;

n is 0, 1, 2 or 3;

p and q are independently 1 or 2;


X is 0, NH, or NR₈;

R₁, R₂, and R₃ are independently hydrogen, unsubstituted or substituted, branched or unbranched, cyclic or acyclic aliphatic; unsubstituted or substituted, branched or unbranched, cyclic or acyclic heteroaliphatic; unsubstituted or substituted aryl; or unsubstituted or substituted heteroaryl; and R₄, R₅, R₆, R₇ and R₈ are independently hydrogen, or substituted

or unsubstituted, branched or unbranched, cyclic or acyclic aliphatic; and pharmaceutically acceptable forms thereof.

[0050] In one embodiment, m is 1, n is 1, p is 1, q is 1, X is 0, R₁, R₂, and R₃ are unsubstituted or substituted, branched or unbranched acyclic aliphatic. In one embodiment, R₄, R₅, R₆ and R₇ are all hydrogen.

In one embodiment, the derivative of romidepsin is of the formula (III):



wherein:

m is 1, 2, 3 or 4;

n is 0, 1, 2 or 3;

q is 2 or 3;

X is 0, NH, or NR₈;


Y is ORB, or SR_8 ;

R₂ and R₃ are independently hydrogen, unsubstituted or substituted, branched or unbranched, cyclic or acyclic aliphatic, unsubstituted or substituted, branched or unbranched, cyclic or acyclic heteroaliphatic, unsubstituted or substituted aryl or unsubstituted or substituted heteroaryl;

[0051] R₄, R₅, R₆, R₇ and R₈ are independently selected from hydrogen or substituted or unsubstituted, branched or unbranched, cyclic or acyclic aliphatic, and pharmaceutically acceptable forms thereof.

[0052] In one embodiment, m is 1, n is 1, q is 2, X is NH and R₂ and R₃ are unsubstituted or substituted, branched or unbranched, acyclic aliphatic. In one embodiment, R₄, R₅, R₆ and R₇ are all hydrogen.

[0053] In one embodiment, the derivative of romidepsin is of the formula (IV):



wherein:


A is a moiety that is cleaved under physiological conditions to yield a thiol group and includes, for example, an aliphatic or aromatic acyl moiety (to form a thioester bond), an aliphatic or aromatic thioxy (to form a disulfide bond), or the like, and pharmaceutically acceptable forms thereof. Such aliphatic or aromatic groups can include a substituted or unsubstituted, branched or unbranched, cyclic or acyclic aliphatic group, a substituted or unsubstituted aromatic group, a substituted or unsubstituted heteroaromatic group, or a substituted or unsubstituted heterocyclic group. A can be, for example, $-\text{COR}_1$, $-\text{SC}(=\text{O})-\text{O}-\text{R}_1$, or $-\text{SR}_2$;

R_1 is independently hydrogen, substituted or unsubstituted amino, substituted or unsubstituted, branched or unbranched, cyclic or acyclic aliphatic, substituted or unsubstituted aromatic group, substituted or unsubstituted heteroaromatic group, or a substituted or unsubstituted heterocyclic group. In one embodiment, R_1 is hydrogen, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, benzyl, or bromobenzyl;


R_2 is a substituted or unsubstituted, branched or unbranched, cyclic or acyclic aliphatic group, a substituted or unsubstituted aromatic group, a substituted or unsubstituted heteroaromatic group, or a substituted or unsubstituted heterocyclic group.

[0054] In one embodiment, R_2 is methyl, ethyl, 2-hydroxyethyl, isobutyl, a fatty acid, a substituted or unsubstituted benzyl, a substituted or unsubstituted aryl, cysteine, homocysteine, or glutathione.

[0055] In one embodiment, the derivatives of romidepsin are of formulae (V) or (V'):



(V),



(V'),

wherein:

each of R₁, R₂, R₃ and R₄ is the same or different and represent an amino acid side chain moiety;

each R₆ is the same or different and represents hydrogen or (C₁-C₄)alkyl; and

Pr¹ and Pr² are the same or different and represent hydrogen or thiol-protecting group.

[0056] In one embodiment, the amino acid side chain moieties are those derived from natural amino acids. In one embodiment, the amino acid side chain moieties are those derived from unnatural amino acids.

[0057] In one embodiment, each amino acid side chain is a moiety selected from hydrogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, -L-O-C(0)-R', -L-C(0)-O-R'', -L-A, -L-NR''R'', -L-Het-C(0)-Het-R'', and -L-Het-R'', wherein L is a (C₁-C₆)alkylene group, A is phenyl or a 5- or 6-membered heteroaryl group, each R' is the same or different and represents (C₁-C₄)alkyl, each R'' is the same or different and represent H or (C₁-C₆)alkyl, each -Het- is the same or different and is a heteroatom spacer selected from -O-, -N(R''')-, and -S-, and each R''' is the same of different and represents hydrogen or (C₁-C₄)alkyl.

[0058] In one embodiment, R₆ is hydrogen.

[0059] In one embodiment, Pr¹ and Pr² are the same or different and are selected from hydrogen and a protecting group selected from a benzyl group which is optionally substituted by (C₁-C₆)alkoxy, (C₁-C₆)acyloxy, hydroxy, nitro, picolyl, picolyl-N-oxide, anthrylmethyl, diphenylmethyl, phenyl, t-butyl, adamantyl, (C₁-C₆)acyloxymethyl, (C₁-C₆)alkoxymethyl, tetrahydropyranyl, benzylthiomethyl, phenylthiomethyl, thiazolidine, acetamidomethyl, benzamidomethyl, tertiary butoxycarbonyl (BOC), acetyl and its derivatives, benzoyl and its derivatives, carbamoyl, phenylcarbamoyl, and (C₁-C₆)alkylcarbamoyl.

[0060] Various romidepsin derivatives of formula (V) and (V') are disclosed in PCT application publication WO 2006/129105, published December 7, 2006, which is

incorporated herein by reference.

[0061] In one embodiment, a formulation of romidepsin is a liquid concentrate formulation for dilution. In some embodiments, provided are liquid formulations comprising one or more additional components. In some such embodiments, additional components are selected from the group consisting of, for example, solubilizers, stabilizers, buffers, tonicity modifiers, bulking agents, viscosity enhancers/reducers, surfactants, chelating agents, adjuvants, diluents, solvents, or other pharmaceutical additives.

[0062] In one embodiment, the additional component is a buffer. In one embodiment, the buffer is a citrate buffer. In another embodiment, the buffer is an acetate buffer.

PHARMACEUTICAL FORMULATIONS

[0063] In one embodiment, provided herein are pharmaceutical formulations of romidepsin. In one embodiment, the formulation is a liquid concentrate formulation. In one embodiment, the formulation is a liquid concentrate formulation for dilution. In one embodiment, a liquid concentrate formulation for dilution is formulated in a solvent or a solvent system. In one embodiment, the solvent system comprises a citrate buffer. In another embodiment, the solvent system comprises an acetate buffer. In one embodiment, a romidepsin formulation is an injectable formulation.

[0064] In one embodiment, the romidepsin liquid concentrate for dilution is formulated in a solvent system. It was found that the solubility issues for developing a liquid IV formulation of romidepsin presented a problem. Romidepsin is practically insoluble in water. Therefore, solvent based formulations were explored. Solubility of romidepsin was studied in the following solvents and solvent mixtures: glycerin:EtOH (1:1); PEG 300; PEG 400; Tween 80:EtOH (1:1); dimethylacetamide (DMA); SolutolHS15:EtOH (1:1); NMP; Water; EtOH:PEG 400 (1:1); EtOH;water (4:6); EtOH; EtOH:PG (80:20); EtOH:PG (60:40); EtOH:PG (40:60); EtOH:PG (20:80); PG; EtOH:PG (40:60) + 10% water; EtOH:PG (40:60) + butylated hydroxyanizole (BHA)/butylated hydroxytoluebe (BHT); and EtOH:PG (40:60) + ascorbic acid.

[0065] Solvents suitable for use in the formulations provided herein include, but are not limited to, propylene glycol (PG), ethanol (EtOH) and a buffer. In one embodiment, the buffer is a citrate buffer. In another embodiment, the buffer is an acetate buffer.

[0066] In one embodiment, the solvent system is a combination of 70% PG, 20% EtOH and 10% citrate buffer. In some embodiments, the solvent system comprises an organic acid, in particular a fatty acid. In a particular embodiment, the fatty acid is oleic acid. In one embodiment, the solvent system comprises 0.05 % oleic acid. In another embodiment, the solvent system comprises 0.10 % oleic acid. In yet another embodiment, the solvent system comprises 0.025 % oleic acid.

[0067] In one embodiment, the solvent system is a combination of 30% PG, 30% EtOH and 40% acetate buffer. In another embodiment, the solvent system is a combination of 70% PG, 20% EtOH and 10% citrate buffer.

[0068] In one embodiment, a liquid concentrate formulation is diluted into normal saline.

[0069] In one embodiment, the pharmaceutical formulations provided herein may be formulated in various dosage forms for parenteral administration. In one embodiment, the pharmaceutical formulation provided herein may be provided in a unit-dosage form or multiple-dosage form. A unit-dosage form, as used herein, refers to a physically discrete unit suitable for administration to human and animal subjects, and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of the active ingredient(s) sufficient to produce the desired therapeutic effect, in association with the required pharmaceutical carriers or excipients. Examples of a unit-dosage form include an ampoule, a vial, a prefilled syringe, a cartridge, or an IV bag. A unit-dosage form may be administered in fractions or multiples thereof. A multiple-dosage form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dosage form. Examples of a multiple-dosage form include a vial.

[0070] In one embodiment, the pharmaceutical formulations provided herein may be administered at once or multiple times at intervals of time. It is understood that the precise dosage and duration of treatment may vary with the age, weight, and condition of the patient being treated, and may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test or diagnostic data. It is further understood that for any particular individual, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the formulations.

Parenteral Administration

[0071] In one embodiment, the pharmaceutical formulations provided herein may be administered parenterally by injection or infusion, for local or systemic administration. Parenteral administration, as used herein, include intravenous, intra-arterial, intraperitoneal, intrathecal, intraventricular, intraurethral, intrasternal, intracranial, intramuscular, intrasynovial, and subcutaneous administration.

[0072] In one embodiment, the pharmaceutical formulations provided herein may be formulated in any dosage forms that are suitable for parenteral administration, including solutions, suspensions, emulsions, micelles, liposomes, microspheres, nanosystems, and solid forms suitable for solutions or suspensions in liquid prior to injection. Such dosage forms can be prepared according to conventional methods known to those skilled in the art of pharmaceutical science (*see, e.g., Remington, The Science and Practice of Pharmacy, supra*).

[0073] In one embodiment, the pharmaceutical formulations intended for parenteral administration may include one or more pharmaceutically acceptable carriers and excipients, including, but are not limited to, aqueous vehicles, water-miscible vehicles, non-aqueous vehicles, antimicrobial agents or preservatives against the growth of microorganisms, stabilizers, solubility enhancers, isotonic agents, buffering agents, antioxidants, local anesthetics, suspending and dispersing agents, wetting or emulsifying agents, complexing agents, sequestering or chelating agents, cryoprotectants, lyoprotectants, thickening agents, pH adjusting agents, and inert gases.

[0074] In one embodiment, suitable aqueous vehicles include, but are not limited to, water, saline, physiological saline or phosphate buffered saline (PBS), citrate buffer, acetate buffer, sodium chloride injection, Ringers injection, isotonic dextrose injection, sterile water injection, dextrose and lactated Ringers injection.

[0075] Non-aqueous vehicles include, but are not limited to, fixed oils of vegetable origin, castor oil, corn oil, cottonseed oil, olive oil, peanut oil, peppermint oil, safflower oil, sesame oil, soybean oil, hydrogenated vegetable oils, hydrogenated soybean oil, and medium-chain triglycerides of coconut oil, and palm seed oil.

[0076] Water-miscible vehicles include, but are not limited to, ethanol, 1,3-butanediol, liquid polyethylene glycol (*e.g.,* polyethylene glycol 300 and polyethylene glycol 400), propylene glycol, glycerin, *N*-methyl-2-pyrrolidone, *N,N*-dimethylacetamide, and dimethyl sulfoxide.

[0077] In one embodiment, suitable antimicrobial agents or preservatives include, but are not limited to, phenols, cresols, mercurials, benzyl alcohol, chlorobutanol, methyl and propyl p-hydroxybenzoates, thimerosal, benzalkonium chloride (*e.g.*, benzethonium chloride), methyl- and propyl-parabens, and sorbic acid.

[0078] Suitable isotonic agents include, but are not limited to, sodium chloride, glycerin, and dextrose.

[0079] Suitable buffering agents include, but are not limited to, phosphate, acetate and citrate.

[0080] Suitable antioxidants include, but are not limited to, bisulfite and sodium metabisulfite.

[0081] Suitable local anesthetics include, but are not limited to, procaine hydrochloride.

[0082] Suitable suspending and dispersing agents include, but are not limited to, sodium carboxymethylcellulose, hydroxypropyl methylcellulose, and polyvinylpyrrolidone.

[0083] Suitable emulsifying agents include, but are not limited to, polyoxyethylene sorbitan monolaurate, polyoxyethylene sorbitan monooleate 80, and triethanolamine oleate.

[0084] Suitable sequestering or chelating agents include, but are not limited to, EDTA.

[0085] Suitable pH adjusting agents include, but are not limited to, sodium hydroxide, hydrochloric acid, acetic acid, citric acid, lactic acid, and sodium citrate.

[0086] Suitable complexing agents include, but are not limited to, cyclodextrins, including α -cyclodextrin, β -cyclodextrin, hydroxypropyl- β -cyclodextrin, sulfobutylether- β -cyclodextrin, and sulfobutylether 7- β -cyclodextrin (CAPTISOL[®], CyDex, Lenexa, KS).

[0087] In one embodiment, a vehicle suitable for a liquid concentrate romidepsin formulation is propylene glycol (70) : ethanol (20) : citrate buffer (10) (pH- 3.0; 25 mM). In one embodiment, a vehicle suitable for a liquid concentrate romidepsin formulation is propylene glycol (70) : ethanol (20) : acetate buffer (10) (pH- 3.0; 25 mM).

[0088] In one embodiment, the pharmaceutical formulations provided herein are formulated for single or multiple dosage administration. In one embodiment, the single dosage formulations are packaged in an ampoule, a vial, or a syringe. In one embodiment, the multiple dosage formulations are packaged in a vial. In one embodiment, the multiple dosage parenteral formulations contain an antimicrobial agent at bacteriostatic or fungistatic concentrations. All parenteral formulations are sterile, as known and practiced in the art.

[0089] In one embodiment, the pharmaceutical formulations are provided as ready-to-use sterile solutions. In another embodiment, the pharmaceutical formulations are provided as sterile dry soluble products, including lyophilized powders to be reconstituted with a vehicle prior to use.

Combination Therapy

[0090] In some embodiments, romidepsin is administered in combination with one or more other pharmaceutical agents. In some embodiments, romidepsin is administered in combination with one or more other chemotherapeutic agents and/or in combination with one or more other pharmaceutical agents (*e.g.*, pain relievers, anti-inflammatories, antibiotics, steroidal agents, anti-folates, kinase inhibitors, methyl transferase inhibitors, antibodies, *etc.*).

[0091] In certain embodiments, romidepsin is administered in combination with one or more cytotoxic agents. Exemplary cytotoxic agents include, but are not limited to, gemcitabine, decitabine, and flavopiridol. In certain embodiments, romidepsin is administered in combination with one or more taxanes and/or one or more proteasome inhibitors. Exemplary proteasome inhibitors include, but are not limited to, bortezomib (VELCADE[®]), peptide boronates, salinosporamide A (NPI-0052), lactacystin, epoxomicin (Ac(Me)-Ile-Ile-Thr-Leu-EX), MG-132 (Z-Leu-Leu-Leu-al), PR-171, PS-519, eponemycin, aclacinomycin A, CEP-1612, CVT-63417, PS-341 (pyrazylcarbonyl-Phe-Leu-boronate), PSI (Z-Ile-Glu(OtBu)-Ala-Leu-al), MG-262 (Z-Leu-Leu-Leu-bor), PS-273 (MNLB), omuralide (*clasto*-lactacystin- β -lactone), NLVS (Nip-Leu-Leu-Leu-vinyl sulfone), YLVS (Tyr-Leu-Leu-Leu-vs), dihydroeponemycin, DFLB (dansyl-Phe-Leu-boronate), ALLN (Ac-Leu-Leu-Nle-al), 3,4-dichloroisocoumarin, 4-(2-aminoethyl)-benzenesulfonyl fluoride, TMC-95A, gliotoxin, EGCG ((-)-epigallocatechin-3-gallate), YU101 (Ac-hFLFL-ex), and combinations thereof.

[0092] In certain embodiments, romidepsin is administered in combination with one or more anti-folates. In some such embodiments, romidepsin is administered in combination with one or more of: folinic acid (leucovorin), methotrexate, pralatrexate, premextred, triazinate, or combinations thereof.

[0093] In certain embodiments, romidepsin is administered in combination with one or more kinase inhibitors (*e.g.*, tyrosine kinase inhibitors). In some embodiments, romidepsin is administered in combination with one or more antibodies that act as a kinase inhibitor. In

some embodiments, romidepsin is administered in combination with one or more of ABT-869, AC220, AZD7762, BIBW 2992, BMS-690154, CDKIAT7519, CYC116, ISIS3521, GSK690693, GSK-461364, MK-0457, MLN8054, MLN8237, MP470, ON 01910.Na, OSI-930, PHA-739358, R935788, SNS-314, TLN-232, XL147, XL228, XL281, XL418, or XL765.

[0094] In certain embodiments, romidepsin is administered in combination with one or more methyl transferase inhibitors.

[0095] In certain embodiments, romidepsin is administered in combination with one or more therapeutic antibodies. In some embodiments, the therapeutic antibodies include, but are not limited to, bevacizumab, cetuximab, dasatinib, erlotinib, gefitinib, imatinib, lapatinib, nilotinib, panitumumab, pegaptanib, ranibizumab, sorafenib, sunitinib, trastuzumab, or any antibody that binds to an antigen bound by one of these moieties.

[0096] In some embodiments, romidepsin is administered in combination with an anti-inflammatory agent, pain reliever, anti-nausea medication, or anti-pyretic. Anti-inflammatory agents useful in the methods provided herein include, but are not limited to, aspirin, ibuprofen, and acetaminophen.

[0097] In certain embodiments, romidepsin is administered in combination with a steroidal agent. In certain embodiments, romidepsin is administered in combination with a steroidal agent selected from the group consisting of alclometasone dipropionate, amcinonide, beclomethasone dipropionate, betamethasone, betamethasone benzoate, betamethasone dipropionate, betamethasone sodium phosphate, betamethasone sodium phosphate and acetate, betamethasone valerate, clobetasol propionate, clocortolone pivalate, cortisol (hydrocortisone), cortisol (hydrocortisone) acetate, cortisol (hydrocortisone) butyrate, cortisol (hydrocortisone) cypionate, cortisol (hydrocortisone) sodium phosphate, cortisol (hydrocortisone) sodium succinate, cortisol (hydrocortisone) valerate, cortisone acetate, desonide, desoximetasone, dexamethasone, dexamethasone acetate, dexamethasone sodium phosphate, diflorasone diacetate, fludrocortisone acetate, flunisolide, fluocinolone acetonide, fluocinonide, fluorometholone, flurandrenolide, halcinonide, medrysone, methylprednisolone, methylprednisolone acetate, methylprednisolone sodium succinate, mometasone furoate, paramethasone acetate, prednisolone, prednisolone acetate, prednisolone sodium phosphate, prednisolone tebutate, prednisone, triamcinolone, triamcinolone acetonide, triamcinolone diacetate, triamcinolone hexacetonide, or

combinations thereof. In one embodiment, romidepsin is administered in combination with dexamethasone.

[0098] In certain embodiments, romidepsin is administered in combination with an agent to treat gastrointestinal disturbances such as nausea, vomiting, and diarrhea. Such agents include, but are not limited to, anti-emetics, anti-diarrheals, fluid replacements, electrolyte replacements, *etc.*

[0099] In certain embodiments, romidepsin is administered in combination with electrolyte replacement or supplementation such as potassium, magnesium, or calcium. In certain embodiments, romidepsin is administered in combination with electrolyte replacement or supplementation such as potassium, and/or magnesium.

[00100] In certain embodiments, romidepsin is administered in combination with an anti-arrhythmic agent.

[00101] In certain embodiments, romidepsin is administered in combination with an agent that increases the production of platelets.

[00102] In certain embodiments, romidepsin is administered in combination with an agent to boost the production of blood cells. In certain embodiments, the agent is erythropoietin.

[00103] In some embodiments, romidepsin is administered in combination with an agent to prevent hyperglycemia.

[00104] In certain embodiments, romidepsin is administered with another HDAC or DAC inhibitor.

METHODS OF USE

[00105] In one embodiment, provided is a method for treating, preventing, or managing cancer in a patient comprising administering to said patient an effective amount of a liquid concentrate formulation provided herein.

[00106] In some embodiments, cancers treatable by the methods provided herein include, but are not limited to, carcinomas, sarcomas, haematological malignancies and the like. In certain embodiments, cancer is a hematological malignancy. In certain embodiments, cancer is a solid tumor.

[00107] In one embodiment, hematological malignancies that can be treated by the methods provided herein include, but are not limited to, lymphomas, leukemias, multiple myeloma, plasma cell-derived cancers, relapsed hematological malignancies, and refractory

hematological malignancies. In one embodiment, lymphomas that can be treated by the methods provided herein include, but are not limited to, mature B-cell lymphomas, mature T-cell and natural killer cell lymphomas, Hodgkin's lymphomas and immunodeficiency-associated lymphoproliferative disorders. In another embodiment, lymphomas that can be treated by the methods provided herein include, but are not limited to, small lymphocytic lymphoma, follicular lymphoma, Mantle cell lymphoma, diffuse large B-cell lymphoma, Burkitt lymphoma, B-cell lymphoblastic lymphoma, small cleaved B-cell lymphoma, non-cleaved B-cell lymphoma, cutaneous T-cell lymphoma (CTCL), and peripheral T-cell lymphoma (PTCL). In another embodiment, leukemias that can be treated by the methods provided herein include, but are not limited to, acute lymphoid leukemia (ALL), chronic lymphocytic leukemia (CLL), acute myeloid leukemia (AML), chronic myeloid leukemia (CML), MLL-rearranged ALL, including leukemias that are relapsed, refractory or resistant to conventional therapy, multiple myeloma, and plasma cell-derived cancer.

[00108] In one embodiment, solid cancers that can be treated by the methods provided herein include, but are not limited to, cancer of the skin; lymph node; breast; cervix; uterus; gastrointestinal tract; pancreas, lung; ovary; prostate; colon; rectal; mouth; brain; head and neck; throat; testes; kidney; pancreas; bone; spleen; liver; bladder; larynx; or nasal passages, and relapsed or refractory cancer.

[00109] In one embodiment, an effective amount of romidepsin to be used is a therapeutically effective amount. In one embodiment, the amounts of romidepsin to be used in the methods provided herein include an amount sufficient to cause improvement in at least a subset of patients with respect to symptoms, overall course of disease, or other parameters known in the art. Precise amounts for therapeutically effective amounts of romidepsin in the pharmaceutical compositions will vary depending on the age, weight, disease, and condition of the patient.

[00110] In one embodiment, romidepsin is administered intravenously. In one embodiment, romidepsin is administered intravenously over a 1-6 hour period. In one embodiment, romidepsin is administered intravenously over a 3-4 hour period. In one embodiment, romidepsin is administered intravenously over a 5-6 hour period. In one embodiment, romidepsin is administered intravenously over a 4 hour period.

[00111] In one embodiment, romidepsin is administered in a dose ranging from 0.5 mg/m² to 28 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 0.5

mg/m² to 5 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 1 mg/m² to 25 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 1 mg/m² to 20 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 1 mg/m² to 15 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 2 mg/m² to 15 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 2 mg/m² to 12 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 4 mg/m² to 12 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 6 mg/m² to 12 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 8 mg/m² to 12 mg/m². In one embodiment, romidepsin is administered in a dose ranging from 8 mg/m² to 10 mg/m². In one embodiment, romidepsin is administered in a dose of about 8 mg/m². In one embodiment, romidepsin is administered in a dose of about 9 mg/m². In one embodiment, romidepsin is administered in a dose of about 10 mg/m². In one embodiment, romidepsin is administered in a dose of about 11 mg/m². In one embodiment, romidepsin is administered in a dose of about 12 mg/m². In one embodiment, romidepsin is administered in a dose of about 13 mg/m². In one embodiment, romidepsin is administered in a dose of about 14 mg/m². In one embodiment, romidepsin is administered in a dose of about 15 mg/m².

[00112] In one embodiment, romidepsin is administered in a dose of 14 mg/m² over a 4 hour iv infusion on days 1, 8 and 15 of the 28 day cycle. In one embodiment, the cycle is repeated every 28 days.

[00113] In one embodiment, increasing doses of romidepsin are administered over the course of a cycle. In one embodiment, the dose of about 8 mg/m² followed by a dose of about 10 mg/m², followed by a dose of about 12 mg/m² is administered over a cycle.

[00114] In some embodiments, unit doses of romidepsin are within the range of about 0.5 mg/m² to about 28 mg/m². In certain embodiments, unit doses are in the range of about 1 mg/m² to about 25 mg/m². In certain embodiments, unit doses are in the range of about 0.5 mg/m² to about 15 mg/m². In certain embodiments, unit doses are in the range of about 1 mg/m² to about 15 mg/m². In certain embodiments, unit doses are in the range of about 1 mg/m² to about 8 mg/m². In certain embodiments, unit doses are in the range of about 0.5 mg/m² to about 5 mg/m². In certain embodiments, the unit doses are in the range of about 2 mg/m² to about 10 mg/m². In some embodiments, unit doses are in the range of about 10 mg/m² to about 20 mg/m². In certain embodiments, unit doses are in the range of about 5 mg/m² to

about 10 mg/m². In some embodiments, unit doses are in the range of about 10 mg/m² to about 15 mg/m². In some embodiments, unit doses are in the range of about 6 to about 19 mg/m². In some embodiments, unit doses are approximately 8 mg/m². In still other embodiments, the unit doses are approximately 9 mg/m². In still other embodiments, unit doses are approximately 10 mg/m². In still other embodiments, unit doses are approximately 11 mg/m². In still other embodiments, unit doses are approximately 12 mg/m². In still other embodiments, unit doses are approximately 13 mg/m². In still other embodiments, unit doses are approximately 14 mg/m². In still other embodiments, unit doses are approximately 15 mg/m². In still other embodiments, unit doses are approximately 30 mg/m².

[00115] In certain embodiments, different individual unit doses within the romidepsin therapy regimen are different. In some embodiments, increasing doses of romidepsin are administered over the course of a cycle. In certain embodiments, a dose of approximately 8 mg/m² is administered, followed by a dose of approximately 10 mg/m², followed by a dose of approximately 12 mg/m² may be administered over a cycle.

[00116] An amount of romidepsin administered in individual unit doses varies depending on the form of romidepsin being administered. In certain embodiments, individual unit doses of romidepsin are administered on one day followed by several days on which romidepsin is not administered. In certain embodiments, romidepsin is administered twice a week. In certain embodiments, romidepsin is administered once a week. In other embodiments, romidepsin is administered every other week.

[00117] In some embodiments, romidepsin is administered daily (for example for 2 weeks), twice weekly (for example for 4 weeks), thrice weekly (for example for 4 weeks), or on any of a variety of other intermittent schedules (*e.g.*, on days 1, 3, and 5; on days 4 and 10; on days 1 and 15; on days 5 and 12; or on days 5, 12, and 19 of 21 or 28 day cycles).

[00118] In certain embodiments, romidepsin is administered on days 1, 8, and 15 of a 28 day cycle. In certain particular embodiments, an 8 mg/m² dose of romidepsin is administered on day 1, a 10 mg/m² dose of romidepsin is administered on day 8, and a 12 mg/m² dose of romidepsin is administered on day 15. In certain embodiments, romidepsin is administered on days 1 and 15 of a 28 day cycle with day 8 being skipped. A 28 day dosing cycle may be repeated. In certain embodiments, a 28 day cycle is repeated 2-10, 2-7, 2-5, or 3-10 times. In certain embodiments, the treatment includes 5 cycles. In certain embodiments, the treatment includes 6 cycles. In certain embodiments, the treatment includes 7 cycles. In certain

embodiments, the treatment includes 8 cycles. In certain embodiments, 10 cycles are administered. In certain embodiments, greater than 10 cycles are administered.

Dosing

[00119] In some embodiments, romidepsin and/or compositions comprising romidepsin are administered according to a standard dosing regimen. In some embodiments, romidepsin and/or compositions comprising romidepsin are administered according to an accelerated dosing regimen.

Standard Dosing for Romidepsin

[00120] In some embodiments, unit doses of romidepsin are within the range of about 0.5 mg/m² to about 28 mg/m² body surface area. In some embodiments, the range of about 6 to about 18 mg/m² is used. In some embodiments, the range is about 10 mg/m² to about 17 mg/m². In some embodiments, particular unit doses are 10 mg/m², 12 mg/m², 13 mg/m², 14 mg/m², and 15 mg/m².

[00121] In some embodiments, intravenous dosing regimens include daily dosing for 2 weeks, twice weekly dosing for 4 weeks, thrice weekly dosing for 4 weeks, and various other intermittent schedules (*e.g.*, on days 1, 3, and 5; on days 4 and 10; on days 1, 8 and 15; on days 1 and 15; on days 5 and 12; or on days 5, 12, and 19 of 21 or 28 day cycles).

[00122] In some embodiments, romidepsin is administered in individual unit doses over 4 hours on days 1, 8, and 15, with courses repeating every 28 days. Often, several courses (*e.g.*, at least 4, at least 6, or more) are administered. Indeed, instances have been reported of as many as 72 courses being administered. In some embodiments, individual unit doses are administered by 4 hour infusion.

Accelerated Dosing for Romidepsin

[00123] Accelerated dosing regimens for romidepsin may be utilized, in which one or more individual unit doses is administered intravenously over a period of time that is less than or equal to about one hour. In some embodiments, one or more individual doses are administered intravenously over a period of time that is less than about 50 minutes, 40 minutes, 30 minutes, 20 minutes, or less. Any regimen that includes at least one unit dose administered over a period of time that is less than about one hour (60 minutes) may be considered an accelerated dosing regimen in accordance with the present disclosure.

[00124] In some embodiments, all unit doses within a regimen are administered intravenously over a time period that is less than or equal to about one hour. In some embodiments, only some of the unit doses within a regimen are administered over a time period that is less than or equal to about one hour. In some embodiments, one or more unit doses within a regimen are administered by a route other than intravenous administration (*e.g.*, oral, subcutaneous, nasal, topical, *etc.*).

[00125] Accelerated dosing regimens of romidepsin can be administered without a significant increase in toxicity or adverse events, particularly in serious adverse events, as compared with a comparable regimen (*e.g.*, an otherwise identical regimen) in which individual unit doses are administered intravenously over a 4-hour period. In one embodiment, accelerated dosing regimens can be administered without a significant increase in toxicity or adverse events, particularly in serious adverse events, as compared with a standard regimen of romidepsin administered by 4-hour intravenous infusion of a dose of about 6-14 mg/m² on days 1, 8, and 15 of a 28 day cycle.

[00126] In some embodiments, romidepsin is administered in an accelerated dosing regimen that is identical to a standard dosing regimen except that one or more unit doses is administered over a time period that is less than about 1 hour (*e.g.*, rather than over a time period of about 4 hours).

[00127] As will be appreciated by one of skill in the art, the dosage, timing and/or routes of administration of particular unit doses of romidepsin may vary depending on the patient and condition being treated. In certain embodiments, the cycles are continued as long as the patient is responding. Therapy may be terminated once there is disease progression, a cure or remission is achieved, or side effects become intolerable. Adverse side effects may also call for lowering the dosage of romidepsin administered, or for adjusting the schedule by which doses are administered.

Kits

[00128] In one embodiment, a kit comprises a dosage form of romidepsin liquid concentrate formulation for dilution. Kits can further comprise a pharmacologically active derivative of romidepsin.

[00129] In other embodiments, kits can further comprise devices that are used to administer the active ingredients. Examples of such devices include, but are not limited to, syringes, and drip bags.

[00130] In one embodiment, kits can further comprise a pharmaceutically acceptable vehicle that can be used to administer one or more active ingredients. For example, if an active ingredient is provided as sterile dry soluble products, including lyophilized powders to be reconstituted with a vehicle prior to use, the kit can comprise a sealed container of a suitable vehicle in which the active ingredient can be reconstituted to form a particulate-free sterile solution that is suitable for parenteral administration. Examples of pharmaceutically acceptable vehicles include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

[00131] Unless otherwise defined, all technical and scientific terms used herein are accorded the meaning commonly known to one of skill in the art. All publications, patents, published patent applications, and other references mentioned herein are hereby incorporated by reference in their entirety. The embodiments of the disclosure should not be deemed to be mutually exclusive and can be combined.

EXAMPLES

[00132] The following examples are provided by way of illustration, not limitation.

Example 1. Romidepsin Liquid Concentrate Injectable Formulations

[00133] Romidepsin liquid concentrate injectable formulation (citrate buffer) and romidepsin liquid concentrate injectable formulation (acetate buffer) were manufactured at the Celgene, Melrose Park, IL Pilot Laboratory using romidepsin raw material (Manufacturer: Sandoz). The product information for the two formulations is summarized in Table 1 (Citrate Buffer) and Table 2 (Acetate Buffer).

Table 1

Product name	Romidepsin Injection (Citrate Buffer)
Strength	5 mg Romidepsin/mL
Banch Size	1L
Vehicle	Propylene Glycol:Ethanol:Citrate buffer (pH 4.5, 25 mM) = 70:20:10
Fill Volume Claim	2 mL
Container and Closure	Vials, Schott, P.O. # BL-026545, Item # 5010-2095, Lot # 6102291228; Stoppers, Westar RS, 13 mm Teflon 4416/50; Grey; Caps 13mm, light blue
RM (active)	Romidepsin
Excipient	Propylene Glycol, Ethanol, Citric Acid Anhydrous, Sodium Citrate Dihydrate

Table 2

Product name	Romidepsin Injection (Acetate Buffer)
Strength	5 mg Romidepsin/mL
Banch Size	1L
Vehicle	Propylene Glycol:Ethanol:Acetate buffer (pH 4.0, 25 mM) = 30:30:40
Fill Volume Claim	2 mL
Container and Closure	Vials, Schott, P.O. # BL-026545, Item # 5010-2095, Lot # 6102291228; Stoppers, Westar RS, 13 mm Teflon 4416/50; Grey; Caps 13mm, light blue
RM (active)	Romidepsin
Excipient	Propylene Glycol, Ethanol, Acetic Acid Glacial

Example 2. Stability of Romidepsin Liquid Concentrate Injectable Formulations

Storage Conditions

[00134] Finished romidepsin liquid concentrate injectable formulation (citrate buffer) and romidepsin liquid concentrate injectable formulation (acetate buffer) were injected in vials and stored inverted and upright per stability testing protocols in storage chambers maintained at -85°C to -70°C and -25°C to -10°C, 5°C ± 3°C, 25°C ± 2°C/60%RH ± 5% RH, and 40°C ± 2°C/75%RH ± 5% RH. A pre-determined number of vials was removed from storage after specified time periods for testing.

Test Methods

[00135] The testing parameters included: visual appearance/color of solution, pH, romidepsin assay, individual impurities, and total impurities by HPLC.

Results

[00136] Based on formulation composition design of experiment study of the potentially promising compositions, five formulations were selected. The romidepsin liquid concentrate formulations were prepared at 100 mL batch size. Two mL aliquots of each formulation were filled into 2 mL clear ampoule vials and sealed. The compositions of romidepsin liquid concentrate injectable formulations are listed in Table 3.

Table 3

Formulation 1	Romidepsin 5 mg/mL PG:EtOH:citrate buffer (pH 4.5; 25 mM)::70:20:10 with 0.05% oleic acid
Formulation 2	Romidepsin 5 mg/mL PG:EtOH:citrate buffer (pH 4.5; 25 mM)::70:20:10
Formulation 3	Romidepsin 5 mg/mL PG:EtOH:citrate buffer (pH 4.5; 50 mM)::70:20:10 with 0.10% oleic acid
Formulation 4	Romidepsin 5 mg/mL PG:EtOH:acetate buffer (pH 4.0 ; 20 mM)::30:30:40
Formulation 5	Romidepsin 5 mg/mL PG:EtOH:citrate buffer (pH 4.5; 25 mM)::70:20:10 with 0.025% oleic acid

[00137] Table 4 shows the stability data of five romidepsin liquid concentrate injectable formulations stored at 5°C.

Table 4

Samples stored at 5°C			Relative peak area @ RRT						
		LC (%)	0.34	0.45	1.00	1.36	1.89	2.04	2.85
F1-1	initial	102.2			100.00				
	1M	102.1			100.00				
	2M	102.2			100.00				
	3M	103.1			100.00				
	6M	104.0			100.00				
F1-2	initial	103.8			100.00				
	1M	102.4			100.00				
	2M	103.1			100.00				
	3M	103.1			100.00				
	6M	105.3			100.00				
F2-1	initial	104.7			100.00				
	1M	104.1			100.00				
	2M	102.8			100.00				
	3M	104.4			100.00				
	6M	105.0			100.00				
F2-2	initial	104.7			100.00				

	1M	103.9			100.00				
	2M	103.6			100.00				
	3M	104.7			100.00				
	6M	106.1			100.00				
F3-1	initial	104.3			100.00				
	1M	102.7			100.00				
	2M	103.1			100.00				
	3M	103.4			100.00				
	6M	104.1			100.00				
F3-2	initial	103.2			100.00				
	1M	103.2			100.00				
	2M	102.6			100.00				
	3M	103.6			100.00				
	6M	104.4			100.00				
F4-1	initial	108.3			100.00				
	1M	107.1			100.00				
	2M	106.5			100.00				
	3M	108.3			100.00				
	6M	108.2			100.00				
F4-2	initial	108.4			100.00				
	1M	107.0			100.00				
	2M	106.5			100.00				
	3M	107.7			100.00				
	6M	108.4			100.00				
F5-1	initial	101.7			100.00				
	1M	102.2			100.00				
	2M	102.4			100.00				
	3M	103.2			100.00				
	6M	103.6			100.00				
F5-2	initial	102.7			100.00				
	1M	102.4			100.00				
	2M	102.1			100.00				
	3M	103.6			100.00				
	6M	103.7			100.00				

[00138] Table 5 shows the stability data of five romidepsin liquid concentrate injectable formulations stored at controlled ambient temperature.

Table 5

Samples stored at RT			Relative peak area @ RRT						
		LC (%)	0.34	0.45	1.00	1.36	1.89	2.04	2.85
F1-1	initial	102.2			100.00				
	1M	103.8			100.00				
	2M	103.9			100.00				
	3M	104.2			100.00				
	6M	105.7			100.00				
F1-2	initial	103.8			100.00				
	1M	102.8			100.00				
	2M	104.3			100.00				
	3M	102.7			100.00				
	6M	106.3			100.00				
F2-1	initial	104.7			100.00				
	1M	103.5			100.00				
	2M	104.3			100.00				
	3M	104.1			100.00				
	6M	107.4			100.00				
F2-2	initial	104.7			100.00				
	1M	103.6			100.00				
	2M	104.0			100.00				

	3M	103.8			100.00				
	6M	106.0			100.00				
F3-1	initial	104.3			100.00				
	1M	103.6			100.00				
	2M	103.4			100.00				
	3M	103.9			100.00				
	6M	104.6			100.00				
F3-2	initial	103.2			100.00				
	1M	104.1			100.00				
	2M	103.6			100.00				
	3M	103.8			100.00				
	6M	105.6			100.00				
F4-1	initial	108.3			100.00				
	1M	108.2			100.00				
	2M	107.9			100.00				
	3M	108.0			100.00				
	6M	110.0			100.00				
F4-2	initial	108.4			100.00				
	1M	107.4			100.00				
	2M	108.1			100.00				
	3M	107.9			100.00				
	6M	109.1			100.00				
F5-1	initial	101.7			100.00				
	1M	103.2			100.00				
	2M	103.0			100.00				
	3M	103.4			100.00				
	6M	104.9			100.00				
F5-2	initial	102.7			100.00				
	1M	103.3			100.00				
	2M	103.0			100.00				
	3M	103.4			100.00				
	6M	104.3			100.00				

[00139] Table 6 shows the stability data of five romidepsin liquid concentrate injectable formulations stored at 40°C.

Table 6

Samples stored at 40°C			Relative peak area @ RRT										
		LC	0.34	0.37	0.45	0.53	0.60	0.67	1.00	1.36	1.89	2.04	2.85
F1-1	initial	102.2							100.00				
	1M	102.8							100.00				
	2M	103.0							99.84		0.07		0.09
	3M	106.6	0.06		0.04				99.76		0.04		0.09
	6M	102.2	0.13	0.03	0.09		0.04		99.40		0.09		0.22
F1-2	initial	103.8							100.00				
	1M	103.0							100.00				
	2M	110.4							99.91		0.04		0.06
	3M	103.3	0.03		0.03				99.78		0.06		0.10
	6M	113.2	0.28	0.05	0.19	0.07	0.09		99.03		0.06		0.20
F2-1	initial	104.7							100.00				
	1M	103.5							100.00				
	2M	103.5							99.91		0.04		0.05
	3M	103.6	0.04		0.03				99.79		0.05		0.10
	6M	103.0	0.15	0.05	0.09	0.03	0.05		99.33		0.09		0.21
F2-2	initial	104.7							100.00				
	1M	103.1							100.00				
	2M	105.9							99.89		0.04		0.07
	3M	103.4	0.04		0.03				99.78		0.06		0.10
	6M	103.9	0.13	0.03	0.08		0.03		99.43		0.09		0.20

F3-1	initial	104.3							100.00				
	1M	102.3							100.00				
	2M	102.3							99.95				0.05
	3M	103.2	0.04		0.03				99.82		0.04		0.07
	6M	102.5	0.15	0.05	0.09	0.03	0.05		99.39		0.07		0.18
F3-2	initial	103.2							100.00				
	1M	102.9							100.00				
	2M	103.4							99.92		0.04		0.04
	3M	103.0	0.03		0.04				99.81		0.04		0.07
	6M	102.6	0.15	0.05	0.10		0.05		99.37		0.08		0.18
F4-1	initial	108.3							100.00				
	1M	106.9							100.00				
	2M	109.7							99.90		0.04		0.05
	3M	107.7							99.84		0.08		0.08
	6M	108.3						0.04	99.64		0.16		0.16
F4-2	initial	108.4							100.00				
	1M	107.2							100.00				
	2M	108.6							99.90		0.05		0.05
	3M	107.4							99.88		0.07		0.05
	6M	107.6						0.03	99.68		0.14		0.15
F5-1	initial	101.7							100.00				
	1M	102.9							100.00				
	2M	103.9							99.95				0.05
	3M	102.1	0.03		0.03				99.81		0.05		0.08
	6M	101.0	0.18		0.11	0.04	0.07		99.34		0.08		0.19
F5-2	initial	102.7							100.00				
	1M	103.0							100.00				
	2M	104.3							99.89		0.04		0.07
	3M	102.4	0.03						99.83		0.06		0.08
	6M	101.6	0.16	0.05	0.11	0.03	0.07		99.32		0.08		0.19

[00140] Table 7 shows the stability data of romidepsin liquid concentrate injectable formulations after exposure to light.

Table 7

Stressed at RT under light			Relative peak area @ RRT						
		LC (%)	0.34	0.45	1.00	1.36	1.89	2.04	2.85
F1	initial	102.2			100.00				
	1M	102.4			100.0				
	2M	101.7			99.85		0.05		0.10
	3M	102.4			99.70	0.03	0.08	0.03	0.14
	6M	101.7			99.73		0.09		0.18
F2	initial	104.7			100.0				
	1M	102.8			100.0				
	2M	102.1			99.82		0.07		0.12
	3M	103.4			99.66	0.02	0.11	0.02	0.18
	6M	102.0			99.50	0.03	0.17		0.29
F3	initial	104.3			100.0				
	1M	103.2			100.0				

	2M	101.7			99.83		0.06		0.12
	3M	102.9			99.76	0.02	0.07	0.02	0.13
	6M	102.1			99.78		0.07		0.14
F4	initial	108.3			100.0				
	1M	106.7			99.95		0.05		
	2M	105.4			99.64		0.16		0.20
	3M	107.0			99.57	0.04	0.17	0.02	0.20
	6M	104.8			99.35	0.06	0.27		0.33
F5	initial	101.7			100.0				
	1M	102.3			100.0				
	2M	101.7			99.83		0.06		0.11
	3M	102.4			99.63	0.04	0.12	0.02	0.20
	6M	100.8			99.55		0.15		0.30

[00141] No loss of potency was observed for all tested formulations over the stability test period as demonstrated by the consistency of the % label claim results. Individual related substances were found to be less than 0.05% or not detected at all time points up to 6 months at 5°C and controlled ambient conditions. A few related substances were observed with the stability samples under accelerated conditions at 40°C. The highest level of related substance observed was no more than 0.30% at 6 months at 40°C. Samples exposed to light showed higher levels of related substances when compared with samples stored in dark at controlled ambient temperature.

[00142] Dilution study was performed to evaluate the compatibility of the tested formulations with normal saline. A dose of 22.5 mg (4.5 mL of formulation) was diluted into 250 mL of normal saline. After gentle mixing, the diluted formulation was observed for appearance. Formulations with oleic acid as a component (formulations 1, 3 and 5) showed haziness in the diluted samples while formulations without oleic acid (formulations 2 and 4) appeared clear. These results indicate that the romidepsin liquid concentrate injectable formulations 2 and 4 are compatible with normal saline upon dilution and were used for further studies.

[00143] Summary of stability test results for romidepsin liquid concentrate injection in citrate buffer (formulation 2) and romidepsin liquid concentrate injection in acetate buffer (formulation 4) are shown in Tables 8-15.

[00144] Tables 8 and 9 demonstrate test results for romidepsin liquid concentrate injection in citrate buffer and romidepsin liquid concentrate injection in acetate buffer stored at -85°C to -70°C and at -25°C to -10°C up to 1 month, accordingly.

Table 8

<i>n</i> =2	Citrate						Acetate					
	Upright			Inverted			Upright			Inverted		
	0	1W	1M	0	1W	1M	0	1W	1M	0	1W	1M
Label Claim (%)	103.5	104.1	104.8	103.5	104.1	105.0	105.2	107.4	107.7	105.2	107.4	108.0
Impurities (%)	0.09											
Total Impurities (%)	0.09											
Appearance	Sample solution is transparent, no visible particulate matter											
pH	4.5	4.6	4.7	4.5	4.7	4.7	4.0	4.1	4.1	4.0	4.1	4.1

Table 9

<i>n</i> =2	Citrate						Acetate					
	Upright			Inverted			Upright			Inverted		
	0	1W	1M	0	1W	1M	0	1W	1M	0	1W	1M
Label Claim (%)	103.5	104.8	104.8	103.5	101.7	104.1	105.2	107.5	108.2	105.2	107.9	107.7
Impurities (%)	0.09											
Total Impurities (%)	0.09											
Appearance	Sample solution is transparent, no visible particulate matter											
pH	4.5	4.7	4.7	4.5	4.7	4.7	4.0	4.1	4.2	4.0	4.1	4.2

[00145] Tables 10 and 11 show the test results for romidepsin liquid concentrate injection in citrate buffer (Table 10) and romidepsin liquid concentrate injection in acetate buffer (Table 11) stored at 5°C ± 3°C up to 12 months.

Table 10

n=2	Citrate											
	Upright						Inverted					
	0	1M	3M	6M	9M	12M	0	1M	3M	6M	9M	12M
Label Claim (%)	103.5	105.0	NA	102.6	NA	104.7	103.5	105.7	105.7	103.5	103.3	105.0
Impurities (%) ¹	0.09	0.09	NA	0.09	NA	0.09	0.09	0.09	0.09	0.10	0.09	0.10
Total Impurities (%)	0.09	0.09	NA	0.09	NA	0.09	0.09	0.09	0.09	0.10	0.09	0.10
Appearance	Sample solution is transparent, no visible particulate matter			Sample solution is transparent, no visible particulate matter	NA	Clear solution, no visible particulate matter	Sample solution is transparent, no visible particulate matter				Clear solution, no visible particulate matter	
pH	4.5	4.7	NA	4.7	NA	4.5	4.5	4.7	4.6	4.7	4.6	4.5

Table 11

n=2	Acetate											
	Upright						Inverted					
	0	1M	3M	6M	9M	12M	0	1M	3M	6M	9M	12M
Label Claim (%)	105.2	107.8	NA	107.9	NA	108.6	105.2	108.2	110.1	108.1	107.1	108.2
Impurities (%)	0.09	0.09	NA	0.10	NA	0.09	0.09	0.09	0.09	0.10	0.09	0.09
Total Impurities (%)	0.09	0.09	NA	0.10	NA	0.09	0.09	0.09	0.09	0.10	0.09	0.09
Appearance	Sample solution is transparent, no visible particulate matter			Sample solution is transparent, no visible particulate matter	NA	Clear solution, no visible particulate matter	Sample solution is transparent, no visible particulate matter				Clear solution, no visible particulate matter	
pH	4.0	4.2	NA	4.1	NA	4.0	4.0	4.3	4.1	4.1	4.1	4.0

[00146] As can be seen, romidepsin liquid concentrate injection in citrate buffer and romidepsin liquid concentrate injection in acetate buffer maintained physical and chemical stabilities at -85°C to -70°C for up to 1 month (Table 8), at -25°C to -10°C for up to 1 month (Table 9), and at 5°C ± 3°C up to 12 months (Tables 10 and 11).

[00147] Table 12 demonstrates the test results for romidepsin liquid concentrate injection in citrate buffer stored at $25^{\circ}\text{C} \pm 2^{\circ}\text{C}/60\% \text{RH} \pm 5\% \text{RH}$.

Table 12

n=2	Citrate											
	Upright						Inverted					
	0	1M	3M	6M	9M	12M	0	1M	3M	6M	9M	12M
Label Claim (%)	103.5	104.0	NA	103.9	NA	104.6	103.5	105.7	106.7	104.2	102.7	104.4
Impurities (%) ¹	0.09	0.09	NA	0.09 0.06	NA	0.09 0.09	0.09	0.09	0.09	0.09 0.06	0.09 0.08	0.09 0.09
Total Impurities (%)	0.09	0.09	NA	0.15	NA	0.18	0.09	0.09	0.09	0.15	0.17	0.18
Appearance	Sample solution is transparent, no visible particulate matter			Sample solution is transparent, no visible particulate matter	NA	Clear solution, no visible particulate matter	Sample solution is transparent, no visible particulate matter				Clear solution, no visible particulate matter	
pH	4.5	4.7	NA	4.8	NA	4.6	4.5	4.7	4.7	4.8	4.8	4.6

[00148] The results demonstrate that the romidepsin formulation in citrate buffer at $25^{\circ}\text{C} \pm 2^{\circ}\text{C}/60\% \text{RH} \pm 5\% \text{RH}$ maintained its physical stability for up to 6 months. No visible particulate matter was observed up to 12 months. The label claim remained essentially unchanged while the total impurities increased from 0.09% to 0.16% and 0.15% at 6 months and further to 0.18% at 12 months for upright and inverted storage condition, respectively (Table 12).

[00149] Table 13 demonstrates the test results for romidepsin liquid concentrate injection in acetate buffer stored at $25^{\circ}\text{C} \pm 2^{\circ}\text{C}/60\% \text{RH} \pm 5\% \text{RH}$ up to 12 months.

Table 13

n=2	Acetate											
	Upright						Inverted					
	0	1M	3M	6M	9M	12M	0	1M	3M	6M	9M	12M
Label Claim (%)	105.2	107.8	NA	107.5	NA	106.9	105.2	107.8	109.6	107.6	106.6	107.8
Impurities (%)	0.09	0.09	NA	0.09	NA	0.09 0.07	0.09	0.09	0.09	0.09	0.09 0.05	0.09 0.06
Total Impurities (%)	0.09	0.09	NA	0.09	NA	0.16	0.09	0.09	0.09	0.09	0.14	0.15
Appearance	Sample solution is		NA	Sample solution is	¹	Clear solution,	Sample solution is				Clear solution,	Clear solution,

	transparent, no visible particulate matter			transparent, no visible particulate matter	²	no visible particul ate matter	transparent, no visible particulate matter				no visible particul ate matter	no visible particula te matter
pH	4.0	4.2	NA	4.2	NA	4.1	4.0	4.2	4.1	4.2	4.1	4.2

¹ – Sample solution is transparent. No visible particulate matter.

² – Some visible particulate matter are observed.

[00150] The results demonstrate that the romidepsin formulation in acetate buffer at 25°C ± 2°C/60% RH ± 5% RH maintained its physical stability for up to 6 months. Visible particulate matter was observed at 9 months, however, no visible particulate matter was observed at 12 months. The label claim remained essentially unchanged for up to 6 months. The total impurities remained at the level of 0.09% after 6 months of storage at 25°C ± 2°C/60% RH ± 5% RH, and increased to 0.16% and 0.15% at 12 months for upright and inverted storage conditions, respectively (Table 13).

[00151] Table 14 demonstrates the test results for romidepsin liquid concentrate injection in citrate buffer stored at 40°C ± 2°C/75% RH ± 5% RH up to 6 months.

Table 14

n=2	Citrate									
	Upright					Inverted				
	0	2W	1M	3M	6M	0	2W	1M	3M	6M
Label Claim (%)	103.5	103.3	104.7	104.9	96.9	103.5	103.3	105.2	104.8	97.7
Total Impurities (%)	0.09	0.09	0.16	0.50	2.43	0.09	0.09	0.17	0.52	2.21
Appearance	Sample solution is transparent, no visible particulate matter					Sample solution is transparent, no visible particulate matter				
pH	4.5	4.7	4.9	4.9	4.8	4.5	4.7	4.9	4.9	4.7

[00152] The results indicate that no visible particulate matter was observed for the romidepsin formulation in citrate buffer maintained at 40°C ± 2°C/75% RH ± 5% RH for up to 6 month. However, the pH value increased from 4.5 to 4.7 at 2 weeks and further to 4.9 at 1 month. Label claim of the romidepsin formulation in citrate buffer remained essentially unchanged up to 3 months, but decreased from 103.5% at 0 time to 96.9% and 97.7% at 6 months for upright and inverted storage conditions, respectively. The total impurities remained 0.09% at 2 weeks, but increased to 0.16% and 0.17% at 1 month and further to 2.43% and 2.21% at 6 months for upright and inverted storage conditions, respectively (Table 14).

[00153] Table 15 demonstrates the test results for romidepsin liquid concentrate injection in acetate buffer stored at $40^{\circ}\text{C} \pm 2^{\circ}\text{C}/75\% \text{ RH} \pm 5\% \text{ RH}$ up to 6 months.

Table 15

n=2	Acetate									
	Upright					Inverted				
	0	2W	1M	3M	6M	0	2W	1M	3M	6M
Label Claim (%)	105.2	106.7	107.3	108.7	104.3	105.2	107.0	106.3	108.3	104.8
Total Impurities (%)	0.09	0.09	0.09	0.36	1.43	0.09	0.09	0.09	0.35	1.21
Appearance	Sample solution is transparent, no visible particulate matter			Some visible particulate matter		Sample solution is transparent, no visible particulate matter			Some visible particulate matter	
pH	4.0	4.2	4.2	4.2	4.1	4.0	4.2	4.2	4.2	4.1

[00154] The results have shown that visible particulate matter was observed for the romidepsin formulation in acetate buffer at 3 months and 6 months stored at $40^{\circ}\text{C} \pm 2^{\circ}\text{C}/75\% \text{ RH} \pm 5\% \text{ RH}$. The pH value remained essentially unchanged. Label claim of the romidepsin formulation in acetate buffer remained essentially unchanged at $40^{\circ}\text{C} \pm 2^{\circ}\text{C}/75\% \text{ RH} \pm 5\% \text{ RH}$ for up to 6 months. The total impurities remained at 0.09% level after 1 month of storage at $40^{\circ}\text{C} \pm 2^{\circ}\text{C}/75\% \text{ RH} \pm 5\% \text{ RH}$, but increased to 0.36% and 0.35% at 3 month and further to 1.43% and 1.21% at 6 months for upright and inverted storage conditions, respectively (Table 15).

[00155] Based on 12 months stability data, the romidepsin liquid concentrate formulation in acetate buffer is more stable than the romidepsin liquid concentrate formulation in citrate buffer due to lower total impurity content at accelerated storage condition.

[00156] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

[00157] The present disclosure has been described above with reference to exemplary embodiments. However, those skilled in the art, having read this disclosure, will recognize that changes and modifications may be made to the exemplary embodiments without departing from the scope of the present disclosure. The changes or modifications are intended to be included within the scope of the present disclosure, as expressed in the following claims.

CLAIMS

1. A formulation comprising romidepsin in a concentration of 1 mg/mL to 10 mg/mL and a citrate buffer.
2. The formulation of claim 1, further comprising propylene glycol (PG) and ethanol (EtOH).
3. The formulation of claim 2, wherein the ratio of PG, EtOH and the citrate buffer is 70% of PG, 20% of EtOH, and 10% of the citrate buffer.
4. The formulation of claim 3, wherein a pH of the solution is in a range of 3.0 and 6.0.
5. The formulation of claim 4, wherein the pH is 4.5.
6. The formulation of claim 1, wherein the concentration of romidepsin is 5 mg/mL.
7. A formulation comprising romidepsin in a concentration of 1 mg/mL to 10 mg/mL and an acetate buffer.
8. The formulation of claim 7, further comprising propylene glycol (PG) and ethanol (EtOH).
9. The formulation of claim 8, wherein the ratio of PG, EtOH and the acetate buffer is 30% of PG, 30% of EtOH, and 40% of the acetate buffer.
10. The formulation of claim 9, wherein a pH of the solution is in a range of 2.0 and 6.0.
11. The formulation of claim 10, wherein the pH is 4.0.

12. The formulation of any one of claims 1 to 11, wherein the formulation is a unit dosage form.

13. The formulation of claim 12, wherein the amount of romidepsin is about 2 to about 20 mg per vial.

14. A method of treating cancer comprising administering to a subject the formulation of any of claims 1 to 13.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2014/072319

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K9/00 A61K47/10 A61K47/12
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/106696 A1 (CELGENE CORP [US]) 18 July 2013 (2013-07-18) paragraph [0005] - paragraph [0007] paragraph [0067] paragraph [0111] - paragraph [0115] paragraph [00165] - paragraph [00168]; example 2 claims	1-14
X	WO 2012/037008 A2 (GLOUCESTER PHARMACEUTICALS INC [US]; STAM RONALD [NL]; STUMPEL DOMINIQ) 22 March 2012 (2012-03-22) paragraph [00157] - paragraph [00158] paragraph [00167] - paragraph [00168] ----- -/--	1-14



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

27 February 2015

Date of mailing of the international search report

09/03/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Muller, Sophie

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2014/072319

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2008/083290 A1 (GLOUCESTER PHARMACEUTICALS [US]; VERDINE GREGORY L [US]; VROLIJK NICHOL) 10 July 2008 (2008-07-10) page 28, line 1 - line 8 -----	1-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2014/072319

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013106696 A1	18-07-2013	AU 2013201937 A1 CA 2863291 A1 CN 104168908 A EP 2802340 A1 JP 2015503623 A US 2013184225 A1 US 2015051157 A1 WO 2013106696 A1	01-08-2013 18-07-2013 26-11-2014 19-11-2014 02-02-2015 18-07-2013 19-02-2015 18-07-2013
WO 2012037008 A2	22-03-2012	EP 2616054 A2 JP 2013540734 A US 2012094927 A1 WO 2012037008 A2	24-07-2013 07-11-2013 19-04-2012 22-03-2012
WO 2008083290 A1	10-07-2008	AU 2007342030 A1 BR PI0720734 A2 CA 2674313 A1 CN 101687010 A CN 103497238 A EP 2124990 A1 EP 2450049 A1 EP 2556838 A1 EP 2815761 A1 JP 2010514802 A JP 2013240332 A US 2010093610 A1 US 2013236928 A1 WO 2008083290 A1	10-07-2008 07-01-2014 10-07-2008 31-03-2010 08-01-2014 02-12-2009 09-05-2012 13-02-2013 24-12-2014 06-05-2010 05-12-2013 15-04-2010 12-09-2013 10-07-2008