



(86) Date de dépôt PCT/PCT Filing Date: 1994/07/15
(87) Date publication PCT/PCT Publication Date: 1995/02/09
(45) Date de délivrance/Issue Date: 2007/04/24
(85) Entrée phase nationale/National Entry: 1996/01/08
(86) N° demande PCT/PCT Application No.: US 1994/007989
(87) N° publication PCT/PCT Publication No.: 1995/004057
(30) Priorité/Priority: 1993/07/29 (US08/099019)



(51) Cl.Int./Int.Cl. *C07D 487/04* (2006.01),
A61K 31/55 (2006.01), *C07D 519/00* (2006.01)

(72) Inventeurs/Inventors:
BLACKBURN, BRENT K., US;
ROBARGE, KIRK, US;
SOMERS, TODD C., US

(73) Propriétaire/Owner:
GENENTECH, INC., US

(74) Agent: DENNISON ASSOCIATES

(54) Titre : INHIBITEURS TRICYCLIQUES DU RECEPTEUR GPII_B III_A
(54) Title: TRICYCLIC INHIBITORS OF THE GPII_B III_A RECEPTOR



(57) **Abrégé/Abstract:**

A tricyclic beazodiazepine derivative that acts as a nonpeptidyl platelet aggregation inhibitor is provided. This inhibitor potently inhibits fibrinogen binding to the GPII_b III_a receptor and is provided in therapeutic compositions for the treatment of diseases for which blocking platelet aggregation is indicated. These nonpeptidyl inhibitors are provided in combination with thrombolytics and anticoagulants.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 487/04, A61K 31/55, C07D 519/00 // (C07D 487/04, 243:00, 235:00) (C07D 487/04, 249:00, 243:00) (C07D 487/04, 243:00, 239:00) (C07D 487/04, 257:00, 243:00)	A1	(11) International Publication Number: WO 95/04057 (43) International Publication Date: 9 February 1995 (09.02.95)
(21) International Application Number: PCT/US94/07989 (22) International Filing Date: 15 July 1994 (15.07.94) (30) Priority Data: 08/099,019 29 July 1993 (29.07.93) US (60) Parent Application or Grant (63) Related by Continuation US Filed on 08/099,019 (CIP) 29 July 1993 (29.07.93) (71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 460 Point San Bruno Boulevard, South San Francisco, CA 94080-4990 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BLACKBURN, Brent, K. [US/US]; 154 Yerba Buena, San Francisco, CA 94127 (US). ROBARGE, Kirk [US/US]; Apartment 4, 1260 Haight Street, San Francisco, CA 94114 (US). SOMERS, Todd, C.		[US/US]; 706 Bounty Drive #610, Foster City, CA 94404 (US). (74) Agents: WINTER, Daryl, B. et al.; Genentech, Inc., 460 Point San Bruno Boulevard, South San Francisco, CA 94080 (US). (81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report.

(54) Title: TRICYCLIC INHIBITORS OF THE GPII_bIII_a RECEPTOR

(57) Abstract

A tricyclic benzodiazepine derivative that acts as a nonpeptidyl platelet aggregation inhibitor is provided. This inhibitor potently inhibits fibrinogen binding to the GPII_bIII_a receptor and is provided in therapeutic compositions for the treatment of diseases for which blocking platelet aggregation is indicated. These nonpeptidyl inhibitors are provided in combination with thrombolytics and anticoagulants.



TRICYCLIC INHIBITORS OF THE GPII_bIII_a RECEPTOR

FIELD OF THE INVENTION

5

The present invention relates to tricyclic integrin receptor inhibitors, especially inhibitors of the platelet GPII_bIII_a receptor. Specifically, the invention is directed to 1,4-benzodiazepines having one or more 5- or 6-member heterocycle fused to the diazepine moiety that act as antagonists of the final common pathway of platelet aggregation. These antagonists act as potent antithrombotics. The invention further relates to therapeutic applications of these tricyclic inhibitors in diseases for which blocking platelet aggregation is indicated.

10

BACKGROUND OF THE INVENTION

15 A. Platelets

Platelets are particles found in whole blood that initiate and provide the structural basis for the hemostatic plug necessary to stop bleeding. Platelets depend on adhesive interactions with extracellular proteins and other cells for proper function (see Hawiger, J. *Atherosclerosis Reviews* 21:165-186 (1990) and Roth J. R. *Immunology Today* 13(2):100-105 (1992)).

The external platelet plasma membrane surface is covered with a variety of membrane bound glycoproteins, many of which have recognition and adhesive functions. Perhaps the most abundant platelet membrane adhesive proteins belong to the integrin superfamily which include the glycoproteins; GPII_bIII_a, GPI_aII_a, GPI_cII_a, GPI_bIX, and the fibronectin and vitronectin receptors (Hynes, R. O., *Cell*, 48: 549 (1987). Each integrin receptor is an $\alpha\beta$ heterodimer displaying characteristic affinity and specificity toward various protein ligands found in serum and/or the extracellular matrix including; von Willebrand factor (vWF), collagen, entactin, tenascin, fibronectin (Fn), vitronectin (Vn), and laminin, as well as fibrinogen (Fg) and thrombospondin (see Kieffer *et al.*, *Ann. Rev. Cell Biol.* 6:329-357(1990) and Ruoslahti, *J. Clin. Invest.*, 87:1-5 (1991)).

The most abundant integrin found on the surface of normal platelets is GPII_bIII_a comprising about 50,000 molecules per platelet and representing about 2% of the total platelet protein. GPII_bIII_a is a non-covalent, calcium ion dependent heterodimeric complex (Jennings, *et al.*, *J. Biol. Chem.* 257: 10458 (1982)) that is restricted in distribution to platelets and other cells of the megakaryocytic lineage (Kieffer *et al.*, *supra*). On activated platelets, GPII_bIII_a promiscuously binds a number of protein ligands with varying affinities, including; fibrinogen, fibronectin, von Willebrand factor, vitronectin and thrombospondin (Plow *et al.*, *Biochemistry of Platelets*, Phillips and Shuman eds., p. 225-256, Orlando: Academic Press [1986]). Each of these protein ligands contain at least one tripeptide sequence Arg-Gly-Asp which is commonly referred to as the "recognition sequence". It is believed the most

20

25

30

35

important interactions mediating platelet aggregation involve GPII_bIII_a binding with the trimeric fibrinogen and, to a lesser extent, with the filamentous von Willebrand factor (Kieffer *et al.*, *supra* and Albeda *et al.*, *The FASEB Journal*, 4:2868-2880 [1990]).

5 GPII_bIII_a binding to its natural ligands can be inhibited to varying degrees by peptides and proteins containing the amino acid recognition sequences; Arg-Gly-Asp (RGD) (Ruoslahti, *supra* and EPO 0368486, assigned to Merck & Co.), Lys-Gly-Asp (KGD), and the fibrinogen γ -chain carboxy-terminal dodecapeptide HHLGGAKQAGDV and analogues thereof (Timmons *et al.*, *Biochemistry*, 28:2919-2922 [1989]).

10 B. The Hyperthrombotic State

Many common human disorders are characteristically associated with a hyperthrombotic state leading to intravascular thrombi and emboli. These are a major cause of medical morbidity, leading to infarction, stroke and phlebitis, and of mortality from stroke and pulmonary and cardiac emboli. Patients with atherosclerosis are predisposed to arterial thromboembolic phenomena for a variety of reasons. Atherosclerotic plaques form niduses for platelet plugs and thrombi that lead to vascular narrowing and occlusion, resulting in myocardial and cerebral ischemic disease. This may happen spontaneously or following procedures such as angioplasty or endarterectomy. Thrombi that break off and are released into the circulation cause infarction of different organs, especially the brain, extremities, heart and kidneys.

20 In addition to being involved in arterial thrombosis, platelets may also play a role in venous thrombosis. A large percentage of such patients have no antecedent risk factors and develop venous thrombophlebitis and subsequent pulmonary emboli without a known cause. Other patients who form venous thrombi have underlying diseases known to predispose them to these syndromes. Some of these patients may have genetic or acquired deficiencies of factors that normally prevent hypercoagulability, such as antithrombin-3. Others have mechanical obstructions to venous flow, such as tumor masses, that lead to low flow states and thrombosis. Patients with malignancy have a high incidence of thrombotic phenomena for unclear reasons. Antithrombotic therapy in this situation with currently available agents is dangerous and often ineffective.

30 Patients whose blood flows over artificial surfaces, such as prosthetic synthetic cardiac valves or through extracorporeal perfusion devices, are also at risk for the development of platelet plugs, thrombi and emboli. It is standard practice that patients with artificial cardiac valves be treated chronically with anti-coagulants. However, in all instances, platelet activation and emboli formation may still occur despite adequate anticoagulation treatment.

35 Thus, a large category of patients, including those with atherosclerosis, coronary artery disease, artificial heart valves, cancer, and a history of stroke, phlebitis, or pulmonary emboli, are candidates for limited or chronic antithrombotic therapy. The number of available


therapeutic agents is limited and these, for the most part, act by inhibiting or reducing levels of circulating clotting factors. These agents are frequently not effective against the patient's underlying hematologic problem, which often concerns an increased propensity for platelet aggregation and adhesion. They also cause the patient to be susceptible to abnormal bleeding.

- 5 Available antiplatelet agents, such as aspirin, inhibit only part of the platelet activation process and are therefore often inadequate for therapy and also cause the patient to be susceptible to abnormal bleeding. An ideal anti-thrombotic drug would have many properties currently not available (see e.g. Sixma, *et al. Thrombosis Research* 67: 305-311 [1992]).

10 C. Therapeutic Agents

An agent which effectively inhibits the final common pathway of platelet activation, namely fibrinogen binding to the GP II_bIII_a receptor, should accordingly be useful in a large group of disorders characterized by a hyperthrombotic state described above.


- Such agents include anti-thrombotic peptides and pseudopeptides capable of inhibiting platelet aggregation. Ruoslahti *et al.* (U.S. patent no. 4,578,079) suggest that tetrapeptides containing the RGD sequence may be used to effect platelet aggregation. Zimmerman *et al.* (U.S. patent no. 4,683,291) disclose that positively charged amino acid residues (eg Arg and Lys) and homologues located before or toward the amino terminus of the RGD sequence are superior for inhibiting fibrinogen-platelet binding. Adams *et al.* (U.S. patent no. 4,857,508) describe superior results for *in-vitro* inhibition of human platelet aggregation in platelet-rich plasma for linear tetrapeptides containing O-methyl-Tyr-amide immediately following the RGD (or homo-RGD) sequence. Barker *et al.*, WO 90/01331, disclose substantially rigid RGD cyclic peptides possessing high affinity for the GP II_bIII_a receptor. A particularly efficacious rigid RGD cyclic peptide described by Barker *et al.* is represented by the following structure:




- Tjoeng *et al.* (U.S. patent no. 4,879,313) describe peptide mimetic platelet aggregation inhibitors in which the first two residues of the RGD sequence are replaced by the pseudodipeptidyl 8-guanidino-octanoyl moiety. Other peptidomimetics in which the Arg of the RGD sequence has been altered include; WO89/07609 (homo-Arg[Har]), EP 341 915 (Har

and alkyl-Arg), WO90/15620 (H₂N and amidino derivatives eg. imidazolyl, imidazolyl, and substituted imidazolyl), EP 422 937 (aryl-, arylalkyl-, and cycloalkyl-amines), WO91/07976 (alkylamidino and alkylamino derivatives), WO91/04247 (alkylamino and alkylguanidino proline derivatives), and WO93/07867 (*p*-(amidino)benzoyl amino substituted β amino acid derivatives). See also EP 384 362 (glycine derivatives) and EP 381 033.


Complete replacement of all residues in the RGD sequence has been described in EP 372 486, assigned to Hoffmann La Roche, where platelet aggregation inhibitors that are derivatives of benzoic and phenylacetic acid are presented. A benzoic acid derivative inhibitor having a particular low IC₅₀ in an ELISA measurement of fibrinogen GPII_bIII_a binding has the following structure:




EP 542 708 A1 assigned to Monsanto/Searle discloses related compounds that may be considered *p*-amidinoaniline derivatives having structures represented by:




EP 537 980 A1 assigned to Glaxo Ltd. discloses cyclohexanecarboxylic acids linked through a piperidyl or piperazyl group to a *p*-amidinophenyl moiety, e.g.



Fibrinogen receptor antagonists possessing alternative structures can be found in EP 478,362, EP 478,363, and EP 478,328, all assigned to Merck. A representative Merck compound has the following structure:




Also of interest are biphenyl derivatives described in EP 483,667 and EP 496,378, the latter publication providing a representative compound having the structure;



5

Quinazoline-3-alkanoic acid derivatives are also reported to have an inhibitory effect on platelet aggregation (although possibly by a different mechanism) in EP 456,835 A1. A generic formula representing these compounds is given by:




10

where n is 1 to 3 and R^2 and R^3 may be *interalia* hydrogen, lower alkyl, lower alkoxy, aralkyl groups that may be substituted with *interalia* $-NR^4R^5$, where R^4 or R^5 may be hydrogen, lower alkyl, or connected with each other to make five- or six-membered heterocycles which may contain another heteroatom, and $A-R^1$ may be lower alkyl.

15


Substituted benzodiazepines have been reported to be platelet aggregation inhibitors. WO 93/00095 (Smith-Kline Beecham) discloses bicyclic benzodiazepine fibrinogen antagonists in which the benzo moiety is preferably substituted with *p*-(amidino)benzoyl amino and N-4 may be substituted with carboxyethyl as represented below

20




Benzodiazepinedione platelet aggregation inhibitors are described in WO 93/08174 (Genentech). A representative disclosed compound having the *p*-(amidino)benzoyl amino benzo substituent is

25



D. Tricyclic 1,4-benzodiazepines

1,4-benzodiazepines with fused 6-membered heterocyclic rings used as CCK antagonists for treatment of panic or anxiety disorders are disclosed in EP 0 519 678 and U.S. patent no. 5,210,082(Merck).



where X is *inter alia* NH₂, R² is *inter alia* loweralkyl or carboxyl substituted phenyl, R³ is *inter alia* NHC(=O)CH(NH₂)benzyl, R¹⁰ is H, OH, or methyl, W is CH, CH(alkyl), or C=O, Y is N(alkyl), C(alkyl), or C=O, and Z is N, N(alkyl), C(alkyl), or C=O. None of these tricyclic 1,4-benzodiazepines is disclosed as having platelet aggregation inhibition activity.

E. Objects

It is an object of this invention to produce tricyclic 1,4-benzodiazepines having potent antithrombotic activity. It is another object of the invention to produce such compounds that are essentially free of peptide bonds, substantially rigid, and stable to degradation. It is a further object to produce potent tricyclic 1,4-benzodiazepine antithrombotics that specifically inhibit the GPIIb/IIIa-Fg interaction but do not strongly inhibit other RGD sensitive integrin interactions including the Vn-VnR, Fn-FnR, and GPIIb/IIIa-vWF interactions. It is still a further object to produce potent tricyclic 1,4-benzodiazepine platelet aggregation inhibitors that are selective and do not significantly increase cutaneous bleeding time or diminish other hemodynamic factors. It is also an object of this invention to produce tricyclic 1,4-benzodiazepine compounds having a long half-life and a large therapeutic range. Finally it is an object of this invention to produce platelet aggregation inhibitors that have good oral bioavailability.



It is still a further object to produce potent tricyclic 1,4-benzodiazepine inhibitors that are capable of selectively inhibiting other integrin interactions such as the Vn-VnR interaction.

These and other objects of this invention will be apparent from consideration of the invention as a whole.

5

SUMMARY OF THE INVENTION

The objects of this invention are accomplished by providing a tricyclic compound represented by structural Formula I or II:



where R¹ and R² are each one or two substituents typically selected from; hydrogen, halo(F, Cl, Br, I), cyano, carboxamido, carboxy, carbamoyloxy, aminocarbonyl, formyloxy, formyl, azido, nitro, imidazolyl, ureido, thioureido, thiocyanato, hydroxy, mercapto, sulfonamido, and an

optionally substituted radical selected from; C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂alkynyl, C₃-C₁₂cycloalkyl, C₆-C₁₄aryl, C₆-C₁₀aryl-C₁-C₈alkyl, C₁-C₁₂alkyloxy, C₁-C₁₂alkyloxyC₁-C₁₂alkyl, C₁-C₁₂alkyloxycarbonyl, C₆-C₁₄ aryloxy, C₁-C₁₂acylamino, N,N-di(C₁-C₁₂)acylamino, N-(C₁-C₁₂)alkyl-N-(C₁-C₁₂)-acylamino, C₁-C₁₂alkylsulfonamido, N-(C₁-C₁₂)alkyl-N-(C₁-C₁₂)alkylsulfonamido, C₁-C₁₂alkylthiocarbonyl, C₁-C₁₂alkylthio, C₁-C₁₂alkylthioC₁-C₁₂alkyl, C₁-C₁₂alkylsulfinyl, C₁-C₆alkylsulfinylC₁-C₆alkyl, C₁-C₁₂alkylsulfonyl, C₁-C₆alkylsulfonylC₁-C₆alkyl, C₁-C₁₂alkylsulfonato, N-(C₁-C₁₂)alkylsulfonamido, N,N-di-(C₁-C₁₂)sulfonamido, N-(C₁-C₁₂) alkyl-N-thioformylamino, C₁-C₁₂thioacylamino, N-(C₁-C₁₂)alkyl-N-(C₁-C₁₂) thioacylamino, C₁-C₁₂alkylsulfinamido, N-(C₁-C₁₂)alkyl-N-(C₁-C₁₂)alkylsulfinylamino, C₁-C₁₂carbalkoxy, C₁-C₁₂alkylcarbonyl, C₁-C₁₂alkyloxycarbonyl, C₁-C₁₂alkanoyloxy, N-(C₁-C₁₂)alkylcarboxamido, N,N-di-(C₁-C₁₂)carboxamido, N,N-di-(C₁-C₁₂)alkylaminocarbonyl, N-(C₁-C₁₂)alkylcarbamoxyloxy, N,N-di-(C₁-C₁₂)carbamoxyloxy, and heterocycloalkyl or heteroaryl having from 1 to 3 rings, each ring having from 5 to 7 atoms with from 0-3 heteroatoms selected from N, O, and S, provided that at least one ring contains a heteroatom, where the substituents are typically selected from halo (F, Cl, Br, I), amino, guanidino, imidazolyl, indolyl, cyano, azido, nitro, hydroxy, mercapto, morpholino, morpholinyl, piperazinyl, piperidinyl, pyrrolinyl, sulfonamido, ureido, thioureido, carboxy, aminocarbonyl, C₁-C₄aminocarbonyl, carboxamido, carbamoxyloxy, formyloxy, formyl, C₁-C₄alkyloxycarbonyl, C₁-C₄alkyl, C₁-C₄alkylamino, C₁-C₄alkoxy, phenyl, hydroxyphenyl, and phenoxy. Optionally, R¹ and R² when bonded to adjacent carbon atoms may join to form an unsubstituted or substituted aryl ring, where the substituents are typically selected from halo (F, Cl, Br, I), cyano, azido, nitro, hydroxy, mercapto, sulfonamido, ureido, thioureido, carboxamido, carbamoxyloxy, formyloxy, formyl, C₁-C₄alkyl, C₁-C₄alkoxy, phenyl, and phenoxy.

Q is a substituted or unsubstituted moiety selected from:

(A) an amino group including;

(1) -NH₂,

(2) -NR³H,

(3) -NR³R⁴, and

(4) -NR³R⁴R⁵,

where R³, R⁴, and R⁵ are typically selected from; (i) an optionally substituted radical selected from (a) -NR⁶R⁷, (b) -C(=NR⁸)-NR⁶R⁷, (c) -N=CR⁹-NR⁶R⁷, (d) -NR¹⁰-CR⁹=NR⁸, and (e) -NR¹⁰-C(=NR⁸)-NR⁶R⁷ where each R⁶, R⁷, R⁸, R⁹, and R¹⁰ is independently selected from hydrogen, hydroxy, C₁-C₄alkoxy, C₁-C₄alkyl, cyano, and halo(F, Cl, Br, I)C₁-C₄alkyl, (ii) optionally substituted C₁-C₁₂alkyl, (iii) optionally substituted C₃-C₇alkenyl, (iv) optionally substituted C₃-C₇alkynyl, (v) optionally substituted C₃-C₁₂cycloalkyl, (vi) optionally substituted C₅-C₁₂cycloalkenyl, (vii) optionally substituted C₆-C₁₄aryl, (viii) optionally

- substituted C₁-C₆alkylC₆-C₁₄aryl, (ix) optionally substituted C₃-C₆alkenyl-C₆-C₁₀aryl, (x) optionally substituted heterocyclyl, (xi) optionally substituted C₁-C₆alkyl-heterocyclyl, (xii) optionally substituted C₁-C₈alkoxy, (xiii) optionally substituted C₁-C₈alkoxycarbonyl, (xiv) optionally substituted C₁-C₈thioalkoxy, (xv) optionally substituted C₃-C₁₀alkenoxy, and (xvi) optionally substituted C₆-C₁₄aryloxy, (xvii) optionally substituted C₆-C₁₄aryloxycarbonyl, (xviii) optionally substituted C₆-C₁₄arylC₁-C₆alkyloxycarbonyl, where the substituents are usually one to three R¹¹, each R¹¹ typically selected from (a) optionally substituted C₆-C₁₂aryloxy, (b) optionally substituted C₆-C₁₂arylamino, (c) optionally substituted C₆-C₁₂aroyl, (d) optionally substituted C₆-C₁₂arylthio, where the substituents are usually one to three R¹², each R¹² typically selected from nitro, amino, C₁-C₈alkylamino, di-(C₁-C₈)alkylamino, amidino, aminomethyleneimino, imino, imino-C₁-C₄alkyl, iminomethyleneamino, guanidino, C₆-C₁₀arylamino, C₁-C₈acylamino, C₁-C₄alkylsulfonamino, azido, cyano, hydroxy, hydroxyC₁-C₈alkyl, C₁-C₈alkoxy, phenyloxy, C₁-C₈alkanoyloxy, C₁-C₈alkanoyl, C₆-C₁₂aroyl, benzamido, phenyl, halo(F, Cl, Br, I), haloC₁-C₈alkyl, and C₁-C₈alkyl, (e) C₁-C₈alkoxy (f) C₁-C₈alkthio (g) halo(F, Cl, Br, I), (h) hydroxy, (i) mercapto, (j) C₁-C₈alkylcarbonyl, (k) carbamoyl, (l) formyl, (m) formyloxy, (n) carboxy, (o) carb-C₁-C₈alkyloxy, (p) C₁-C₈alkanoyloxy, (q) N-(C₁-C₈)alkylcarboxamido, (r) N-(C₁-C₈),N-(C₁-C₈)dialkylcarboxamido, (s) carbamoyloxy, (t) N-(C₁-C₈)alkylcarbamoyloxy, (u) N-(C₁-C₈),N-(C₁-C₈)dialkylcarbamoyloxy, (v) C₁-C₈alkylsulfinyl, (w) C₁-C₈alkylsulfonyl, (x) C₁-C₈alkylsulfonato, (y) sulfo, (z) sulfonamido, (aa) N-(C₁-C₈)alkylsulfonamido, (ab) N-(C₁-C₈),N-(C₁-C₈)dialkylsulfonamido, (ac) amino, (ad) C₁-C₈alkylamino, (ae) C₁-C₈dialkylamino, (af) C₁-C₈acylamino, (ag) N-(C₁-C₈),N-(C₁-C₈)diacylamino, (ah) N-(C₁-C₈)-alkyl-N-(C₁-C₈)-acylamino, (ai) formylamino, (aj) ureido, (ak) isothioureido, (al) amino-C₂-C₈alkylthio, (am) amino-C₂-C₈alkoxy, (an) amidino, (ao) guanidino, (ap) aminomethyleneimino, (aq) imino, (ar) imino-C₁-C₄alkyl, (as) iminomethyleneamino, (at) glycyllamino, (au) glycyll, (av) phthalimido, (aw) succinimido, (ax) morpholino, (ay) C₁-C₈alkylsulfonamido, (az) N-(C₁-C₈)-alkyl-N-(C₁-C₈)alkyl sulfonoylamino, (ba) C₁-C₈alkylsulfinamino, (bb) N-(C₁-C₈)alkyl-N-(C₁-C₈)alkylsulfinamino, (bc) C₁-C₈alkoxyamino, (bd) C₁-C₈alkoxyamino, (be) N-(C₁-C₈)alkyl-N-(C₁-C₈)alkoxyamino, (bf) C₃-C₇cycloalkyl, (bg) oxo, and (bh) heterocyclyl, optionally any one or two pairs of R³-R¹⁰ may independently be joined to form one or two optionally substituted heterocyclic rings, each ring optionally fused with one or two optionally substituted homocyclic or heterocyclic rings of from four to seven atoms where any heterocyclic ring contains from one to four heteroatoms selected from N, O, and S and where any ring may be substituted with from one to three R¹²,
- (B) an amidino (aminoiminomethyl) group including;
- (1) -C(=NH)-NH₂,
 - (2) -C(=NH)-NHR³,

(3) $-C(=NR^4)-NHR^3$,

(4) $-C(=NH)-NR^3R^4$, and

(5) $-C(=NR^5)-NR^3R^4$,

where R^3 , R^4 , and R^5 are defined above,

5

(C) an aminoalkyleneamino group including;

(1) $-N=CH-NH_2$,

(2) $-N=CH-NHR^3$,

(3) $-N=CH-NR^3R^4$, and

(4) $-N=CR^5-NR^3R^4$,

10

where R^3 , R^4 , and R^5 are defined above,

(D) an iminoalkyleneamino group, including;

(1) $-NH-CH=NH$,

(2) $-NH-CH=NR^3$,

(3) $-NH-CR^4=NR^3$, and

15

(4) $-NR^5-CR^4=NR^3$,

where R^3 , R^4 , and R^5 are defined above,

(E) a guanidino (aminoiminomethyleneamino) group including;

(1) $-NH-C(=NH)-NH_2$,

(2) $-NH-C(=NH)-NR^3H$,

20

(3) $-NH-C(=NH)-NR^3R^4$,

(4) $-NH-C(=NR^5)-NR^3R^4$,

(5) $-NR^3-C(=NR^3)-NR^3R^4$,

(6) $-NR^3-C(=NH)-NR^3R^4$,

(7) $-NR^3-C(=NR^3)-NH_2$,

25

(8) $-NR^3-C(=NH)-NH_2$,


(9) $-NR^3-C(=NR^3)-NHR^4$, and


(10) $-NR^3-C(=NH)-NHR^4$,

where R^3 , R^4 , and R^5 are defined above,

(F) an optionally substituted saturated heterocyclic group including;


30






where (1) n is 0, 1, 2, or 3, (2) R^{13} is selected from; R^6 , $-\text{CR}^9=\text{NR}^8$, $-\text{CR}^9(=\text{NR}^8)-\text{NR}^6\text{R}^7$,
 5 $-\text{C}(=\text{NR}^8)-\text{NR}^6\text{R}^7$, $-\text{N}=\text{CR}^9-\text{NR}^6\text{R}^7$, $-\text{NR}^{10}-\text{CR}^9=\text{NR}^8$, and $-\text{NR}^{10}-(\text{C}=\text{NR}^8)-\text{NR}^6\text{R}^7$ where R^6 -
 R^{10} are defined above, (3) Z^2 is O, S, or NR^{13} , and (4) the substituents are independently one to
 three R^{12} ,


(G) an optionally substituted unsaturated (nonaromatic) heterocyclyl including;



where (1) m is 1, 2, or 3, (2) Z^2 and R^{13} are defined above, and (3)
 the substituents are typically one to three R^{12} ,

(H) an optionally substituted unsaturated (aromatic) heterocyclyl including;


















5

where (1) Z^3 , Z^4 , and Z^5 are typically selected from O, S, N, and NH, provided at least one Z^3 , Z^4 , or Z^5 is N or NH, (2) R^{13} is defined above, and the substituents are independently one to three R^{12} . Exemplary heterocyclic Q groups include; isoindoliny, quinuclidiny, morpholiny, piperidine, piperazine, pyridine, pyrimidine, and 1,3-diazacyclohex-4-ene. Optionally, any of the nitrogen containing heterocycles described above may be substituted at any accessible position with amino, imino, amidino, aminomethyleneimino, iminomethyleneamino, guanidino, N^G -aminoguanidino, alkylamino, dialkylamino, trialkylamino, or alkylideneamino groups.

L is a bivalent radical containing from 3 to 9 methylene groups where any methylene group or groups may be replaced with one or more alkene, alkyne, aryl, or functional groups containing the heteroatoms selected from the group N, O, and S. Typically, L is phenyl or a 6-member heteroaryl bonded through two optionally substituted atoms selected from O, N, and C (e.g. $-C(=O)NH-$) to the benzodiazepine nucleus. More specifically, typical L's are benzoylamino and pyridylcarbonylamino. Preferably, L is an optionally substituted bivalent radical including; (A) C3-C7-alkylene, (B) C3-C7-cycloalkylene, (C) C3-C7-alkenylene, (D) C3-C7-alkadienylene, (E) C3-C7-alkynylene, (F) C4-C7-alkadiynylene, (G) C4-C7-alkenynylene, (H) C6-C14-arylene, (I) C6-C14-aryl-C2-C4-alkynylene, (J) C1-C3-alkyl-C6-C14-aryl-C2-C4-alkynylene, (K) C6-C14-aryl-C2-C4-alkenylene, (L) C1-C3-alkyl-C6-C14-arylene, (M) C1-C3-

- alkyl-C₆-C₁₄-aryl-C₂-C₄-alkenylene, (N) C₆-C₁₄-aryl-C₁-C₃-alkylene, (O) C₆-C₁₄-aryl-C₁-C₃-alkyloxyene, (P) C₁-C₂-alkyl-C₆-C₁₄-aryl-C₁-C₂-alkylene, (Q) C₁-C₃-alkyloxy-C₆-C₁₄-arylene, (R) C₂-C₆-alkyloxyene, (S) C₁-C₅-alkyloxy-C₁-C₅-alkylene, (T) C₆-C₁₀-aryloxyene, (U) C₆-C₁₀-aryloxy-C₁-C₅-alkylene, (V) C₂-C₆-alkylthioene, (W) C₁-C₅-alkylthio-C₁-C₅-alkylene, (X) C₆-C₁₀-arylthioene, (Y) C₆-C₁₀-arylthio-C₁-C₅-alkylene, (Z) C₁-C₅-alkylsulfoxide-C₁-C₅-alkylene, (AA) C₁-C₅-alkylsulfone-C₁-C₅-alkylene,





5 where

R^{14} is selected from; (i) a chemical bond, (ii) C₁-C₈alkyl, (iii) C₁-C₈alkyloxy, (iv) C₃-C₇-cycloalkyl, (v) C₂-C₅-alkenyl, (vi) C₃-C₅-alkynyl, (vii) C₆-C₁₀-aryl, (viii) C₁-C₂-alkyl-C₆-C₁₀-aryl-C₁-C₂-alkyl, (ix) C₁-C₃-alkyl-C₆-C₁₂-aryl, (x) C₆-C₁₀-aryl-C₁-C₂-alkyl, (xi) C₆-C₁₀-aryloxy-C₁-C₂-alkyl, (xii) HET, and (xiii) piperiziny;

10 R^{15} is selected from; (i) a chemical bond, (ii) C₁-C₄-alkyl, (iii) C₂-C₄-alkenyl, (iv) C₂-C₄-alkynyl, (v) C₁-C₃-alkyl-C₆-C₁₂-aryl, and (vi) C₆-C₁₀-aryl;

R^{16} is selected from; (i) a chemical bond, (ii) C₁-C₅-alkyl, (iii) C₃-C₇-cycloalkyl, (iv) C₃-C₅-alkenyl, (v) C₃-C₅-alkynyl, (vi) C₆-C₁₀-aryl, (vii) C₁-C₃-alkyl-C₆-C₁₂-aryl, (viii) C₆-C₁₀-aryl-C₁-C₂-alkyl, (ix) HET, and (x) piperiziny;

15 R^{17} is selected from; (i) C₃-C₄-alkenyl, (ii) C₃-C₄-alkynyl, (iii) C₆-C₁₀-aryl, and (iv) benzyl; HET is a saturated or unsaturated heterocycle having from 5-14 atoms in the cycle(s) and from 1-3 heteroatoms selected from N, O, and S, where the substituents are selected from one to three R^{12} ;

20 Where any aryl group comprising substituent L is unsubstituted or substituted with one to four of the groups selected from (i) nitro, (ii) halo(F, Cl, Br, I), (iii) C₁-C₄alkoxy, (iv) halo(F, Cl, Br, I)C₁-C₄alkyl, and (v) amino;

R^{18} , R^{19} , R^{20} and R^{21} are independently selected from hydrogen, C₁-C₅alkyl, halo(F, Cl, Br, I)C₁-C₅alkyl, C₁-C₅alkoxy, halo(F, Cl, Br, I), cyano, carboxy, hydroxy, C₁-C₅alkoxycarbonyl, and C₁-C₅alkylsulfonylC₀-C₃alkyl;

25 R^{22} is selected from (a) hydroxy, (b) C₁-C₈-alkoxy, (c) C₃-C₁₂-alkenoxy, (d) C₆-C₁₂-aryloxy, (e) C₁-C₆-alkyl-C₆-C₁₂-aryloxy, (f) di-C₁-C₈-alkylamino-C₁-C₈-alkoxy, (g) acylamino-C₁-C₈-alkoxy selected from the group (i) acetylaminoethoxy, (ii) nicotinoylaminoethoxy, and (iii) succinamidoethoxy, (h) C₁-C₈-alkoxyloxy-C₁-C₈-alkoxy, (i) C₆-C₁₂-aryl-C₁-C₈-alkoxy, where any aryl groups are unsubstituted or substituted with one to three of the groups (i) nitro, (ii) halo (F, Cl, Br, I), (iii) C₁-C₄-alkoxy, and (iv) amino, (j) hydroxy-C₂-C₈-alkoxy, (k) dihydroxy-C₃-C₈-alkoxy, and (l) $\text{NR}^{23}\text{R}^{24}$;

30 R^{23} and R^{24} are independently selected from the group (a) hydrogen, (b) C₁-C₈-alkyl, (c) C₃-C₈-alkenyl, (d) C₆-C₁₂-aryl, (e) C₆-C₁₂-aryl-C₁-C₈-alkyl where any aryl groups are

unsubstituted or substituted with one to three of the groups (i) nitro, (ii) halo (F, Cl, Br, I), (iii) C₁-C₄-alkoxy, and (iv) amino;

A¹ is selected from CR¹ N, and NR²⁵;

A² is selected from CR², N, O=S=O, S=O, S, O, C=O, C-OR²⁶, and C=N-R²⁵;

5 B¹ is selected from CR¹, N, NR²⁵, or C=O;

B² is selected from CR², NR²⁵, O=S=O, S=O, S, O, or C=O;


B³ is selected from, CR¹, CHR², or C=O;

R²⁵ is selected from R⁶ and (CH₂)_mR¹, where m is 1, 2, or 3; and


R²⁶ is selected from hydrogen, C₁-C₈-alkyl, C₆-C₁₂-aryl, and C₆-C₁₂-aryl-C₁-C₈-alkyl

10 where any aryl or alkyl groups are unsubstituted or substituted with one to three of the groups nitro, and halo (F, Cl, Br, I).


Preferred tricyclic compounds of this invention are represented by structural Formulae IIIa-VIIIb.




IIIa




IIIb




IVa




IVb




Va




Vb




VIa



VII



VIIIa




VIIIb

5

where R^1 , R^2 , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , Q , and L are as defined above.

More preferred compounds of this invention are represented by structural formula Ia



Ia

10

where each R^1 and R^2 is one or two substituents selected from hydrogen, fluoro, trifluoromethyl and chloro; R^{18} , R^{19} , R^{20} and R^{21} are selected from R^1 and methyl; A^1 is CH or N; A^2 is CH, C-CH₃, C-CH₂-CH₃ or N; -X-Y- is selected from -NH-C(=O)-, -C(=O)-NH-, -CC-, -CH₂-O-, -O-CH₂- and -CH₂-CH₂-; and Z^1 is selected from N and CR¹.

15

Most preferred compounds of this invention may be selected from:

- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 5 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-methyl-6-oxo, acetate,
- 10 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-phenyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-phenyl-6-oxo, acetate,
- 15 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-phenyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-phenyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methyl-6-oxo, acetate,
- 20 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-methyl-6-oxo, acetate,
- 25 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-methyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-hydroxy-6-oxo, acetate,

- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-hydroxy-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-hydroxy-6-oxo, acetate,
- 5 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-hydroxy-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methoxy-6-oxo, acetate,
- 10 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methoxy-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-methoxy-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-methoxy-6-oxo, acetate,
- 15 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-phenyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-phenyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-phenyl-6-oxo, acetate,
- 20 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-phenyl-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-6-oxo, acetate,
- 25 4H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-6-oxo, acetate,

- 4H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)picolinoyl]amino]-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 5 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 10 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 15 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 20 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 25 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,

- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 5 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-]pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 10 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-]pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 15 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 20 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-3-methyl, propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 25 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino-6-oxo, acetate,

4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino]-6-oxo, acetate,

4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate,

5 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino]-6-oxo, acetate,

4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate,

10 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino]-6-oxo, acetate,

4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate,

4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-3-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino]-6-oxo, acetate,

15 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-3-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate,

20 The invention further provides a pharmaceutical composition comprising a pharmaceutically acceptable excipient and/or pharmaceutically acceptable hydrates, solvates, and salts of Formulae I and II.

25 The invention provides a method for inhibiting platelet aggregation or reducing platelet aggregation in a mammal comprising administering a platelet aggregation inhibiting amount of the pharmaceutical composition of Formulae I and II. Optionally, the method includes administering the pharmaceutical composition in combination with a peptidyl antithrombotic and/or a thrombolytic agent and/or an anticoagulant.

DETAILED DESCRIPTION OF THE INVENTION

A. Definitions

The term "alkyl" means a branched or unbranched, saturated aliphatic hydrocarbon radical, having the number of carbon atoms specified, or if no number is specified, having up to 12 carbon atoms. Examples of alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, 2-methylbutyl, 2,2-dimethylpropyl, n-hexyl, 2-methylpentyl, 2,2-dimethylbutyl, n-heptyl, 2-methylhexyl, and the like. The terms "lower alkyl" "C₁-C₆ alkyl" and "alkyl of 1 to 6 carbon atoms" are synonymous and used interchangeably. A preferred "C₁-C₆ alkyl" group is methyl.

The term "substituted C_n-C_m alkyl" where m and n are integers identifying the range of carbon atoms contained in the alkyl group, denotes the above alkyl groups that are substituted by one, two or three; halogen, hydroxy, protected hydroxy, amino, protected amino, C₁-C₇ acyloxy, nitro, carboxy, protected carboxy, carbamoyl, carbamoyloxy, cyano, methylsulfonylamino, morpholino, or C₁-C₄ alkoxy groups. The substituted alkyl groups may be substituted once, twice or three times with the same or with different substituents.

Examples of the above substituted alkyl groups include but are not limited to; cyanomethyl, nitromethyl, hydroxymethyl, hydroxyethyl, trityloxymethyl, propionyloxymethyl, aminomethyl, carboxymethyl, alkyloxycarbonylmethyl, allyloxycarbonylaminomethyl, carbamoyloxymethyl, methoxymethyl, ethoxymethyl, t-butoxymethyl, acetoxymethyl, chloromethyl, bromomethyl, iodomethyl, trifluoromethyl, 2-morpholinoethyl, 6-hydroxyhexyl, 2,4-dichloro(n-butyl), 2-amino(iso-propyl), 2-carbamoyloxyethyl and the like. A preferred group of examples within the above "C₁-C₁₂ substituted alkyl" group includes the substituted methyl group, e.g. a methyl group substituted by the same substituents as the "substituted C_n-C_m alkyl" group. Examples of the substituted methyl group include groups such as hydroxymethyl, protected hydroxymethyl (e.g. tetrahydropyranyloxymethyl), acetoxymethyl, carbamoyloxymethyl, trifluoromethyl, chloromethyl, bromomethyl and iodomethyl.

The terms "C₁-C₁₂ alkyloxy" or "C₁-C₁₂ alkoxy" are used interchangeably herein and denote groups such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, t-butoxy and like groups. Substituted alkoxy groups include those substituents described above for "alkyl".

The terms "C₁-C₁₂ acyloxy" or "C₁-C₁₂ alkanoyloxy" are used interchangeably and denote herein groups such as formyloxy, acetoxy, propionyloxy, butyryloxy, pentanoyloxy, hexanoyloxy, heptanoyloxy, and the like.

The terms "C₁-C₁₂ alkylcarbonyl", "C₁-C₁₂ alkanoyl" and "C₁-C₁₂ acyl" are used interchangeably herein encompass groups such as formyl, acetyl, propionyl, butyryl, pentanoyl, hexanoyl, heptanoyl, benzoyl and the like.

The term "cycloalkyl" as used herein refers to a mono-, bi-, or tricyclic aliphatic ring having 3 to 14 carbon atoms and preferably 3 to 7 carbon atoms.

5 The term "alkenyl" means a branched or unbranched hydrocarbon radical having the number of carbon atoms designated containing one or more carbon-carbon double bonds, each double bond being independently cis, trans, or a nongeometric isomer.

The term "alkynyl" means a branched or unbranched hydrocarbon radical having the number of carbon atoms designated containing one or more carbon-carbon triple bonds.

10 The terms "C₁-C₁₂ alkylthio" and "C₁-C₁₂ substituted alkylthio" denote C₁-C₁₂ alkyl and C₁-C₁₂ substituted alkyl groups, respectively, attached to a sulfur which is in turn the point of attachment for the alkylthio or substituted alkylthio group to the group or substituent designated.

15 The term "aryl" when used alone means a homocyclic aromatic radical, whether or not fused, having the number of carbon atoms designated. Preferred aryl groups include phenyl, naphthyl, biphenyl, phenanthrenyl, naphthacenyl, and the like (see e.g. *Lang's Handbook of Chemistry* (Dean, J. A., ed) 13th ed. Table 7-2 [1985]).

20 The term "substituted phenyl" or "substituted aryl" denotes a phenyl group or aryl group substituted with one, two or three substituents chosen from halogen(F, Cl, Br, I), hydroxy, protected hydroxy, cyano, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, carboxy, protected carboxy, carboxymethyl, protected carboxymethyl, hydroxymethyl, protected hydroxymethyl, aminomethyl, protected aminomethyl, trifluoromethyl N-(methylsulfonylamino) or other groups specified.

25 Examples of the term "substituted phenyl" includes but is not limited to a mono- or di(halo)phenyl group such as 4-chlorophenyl, 2,6-dichlorophenyl, 2,5-dichlorophenyl, 3,4-dichlorophenyl, 3-chlorophenyl, 3-bromophenyl, 4-bromophenyl, 3,4-dibromophenyl, 3-chloro-4-fluorophenyl, 2-fluorophenyl and the like; a mono- or di(hydroxy)phenyl group such as 4-hydroxyphenyl, 3-hydroxyphenyl, 2,4-dihydroxyphenyl, the protected-hydroxy derivatives thereof and the like; a nitrophenyl group such as 3- or 4-nitrophenyl; a cyanophenyl group, for example, 4-cyanophenyl; a mono- or di(lower alkyl)phenyl group such as 4-methylphenyl, 2,4-dimethylphenyl, 2-methylphenyl, 4-(iso-propyl)phenyl, 4-ethylphenyl, 3-(n-propyl)phenyl and
30 the like; a mono- or di(alkoxy)phenyl group, for example, 2,6-dimethoxyphenyl, 4-methoxyphenyl, 3-ethoxyphenyl, 4-(isopropoxy)phenyl, 4-(t-butoxy)phenyl, 3-ethoxy-4-methoxyphenyl and the like; 3- or 4-trifluoromethylphenyl; a mono- or dicarboxyphenyl or (protected carboxy)phenyl group such as 4-carboxyphenyl; a mono- or di(hydroxymethyl)phenyl or (protected hydroxymethyl)phenyl such as 3-(protected hydroxymethyl)phenyl or 3,4-di(hydroxymethyl)phenyl; a mono- or di(aminomethyl)phenyl or (protected aminomethyl)phenyl such as 2-(aminomethyl)phenyl or 2,4-(protected aminomethyl)phenyl; or
35 a mono- or di(N-(methylsulfonylamino))phenyl such as 3-(N-methylsulfonylamino)-phenyl. Also, the term "substituted phenyl" represents disubstituted phenyl groups wherein the

substituents are different, for example, 3-methyl-4-hydroxyphenyl, 3-chloro-4-hydroxyphenyl, 2-methoxy-4-bromophenyl, 4-ethyl-2-hydroxyphenyl, 3-hydroxy-4-nitrophenyl, 2-hydroxy-4-chlorophenyl and the like. Preferred substituted phenyl groups include the 2- and 3-trifluoromethylphenyl, the 4-hydroxyphenyl, the 2-aminomethylphenyl and the 3-(N-(methylsulfonylamino))phenyl groups.

The term "arylalkyl" or "aralkyl" means one, two, or three aryl groups having the number of carbon atoms designated, appended to an alkyl radical having the number of carbon atoms designated including but not limited to; benzyl, naphthylmethyl, phenethyl, benzhydryl (diphenylmethyl), trityl, and the like. A preferred arylalkyl group is the benzyl group.

The term "substituted C₆-C₁₀aryl-C₁-C₈alkyl" denotes a C₁-C₈alkyl group substituted at any carbon with a C₆-C₁₀aryl group bonded to the alkyl group through any aryl ring position and substituted on the C₁-C₈alkyl portion with one, two or three groups chosen from halogen (F, Cl, Br, I), hydroxy, protected hydroxy, amino, protected amino, C₁-C₇acyloxy, nitro, carboxy, protected carboxy, carbamoyl, carbamoyloxy, cyano, C₁-C₆alkylthio, N-(methylsulfonylamino) or C₁-C₄alkoxy. Optionally the aryl group may be substituted with one, two, or three groups chosen from halogen, hydroxy, protected hydroxy, nitro, C₁-C₆alkyl, C₁-C₄alkoxy, carboxy, protected carboxy, carboxymethyl, protected carboxymethyl, hydroxymethyl, protected hydroxymethyl, aminomethyl, protected aminomethyl, or an N-(methylsulfonylamino) group. As before, when either the C₁-C₈alkyl portion or the aryl portion or both are disubstituted, the substituents can be the same or different.

Examples of the term "substituted C₆-C₁₀aryl-C₁-C₈alkyl" include groups such as 2-phenyl-1-chloroethyl, 2-(4-methoxyphenyl)ethyl, 2,6-dihydroxy-4-phenyl(n-hexyl), 5-cyano-3-methoxy-2-phenyl(n-pentyl), 3-(2,6-dimethylphenyl)n-propyl, 4-chloro-3-aminobenzyl, 6-(4-methoxyphenyl)-3-carboxy(n-hexyl), 5-(4-aminomethyl phenyl)-3-(aminomethyl)(n-pentyl), and the like.

The term "carboxy-protecting group" as used herein refers to one of the ester derivatives of the carboxylic acid group commonly employed to block or protect the carboxylic acid group while reactions are carried out on other functional groups on the compound. Examples of such carboxylic acid protecting groups include 4-nitrobenzyl, 4-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, pentamethylbenzyl, 3,4-methylenedioxybenzyl, benzhydryl, 4,4'-dimethoxybenzhydryl, 2,2',4,4'-tetramethoxybenzhydryl, t-butyl, t-amyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, 4,4',4''-trimethoxytrityl, 2-phenylprop-2-yl, trimethylsilyl, t-butyl dimethylsilyl, phenacyl, 2,2,2-trichloroethyl, β -(trimethylsilyl)ethyl, β -(di(n-butyl)methylsilyl)ethyl, p-toluenesulfonylethyl, 4-nitrobenzylsulfonylethyl, allyl, cinnamyl, 1-(trimethylsilylmethyl)prop-1-en-3-yl, and like moieties. The species of carboxy-protecting group employed is not critical so long as the derivatized carboxylic acid is stable to the condition of subsequent reaction(s) on other positions of the benzodiazepinedione molecule and can be removed at the appropriate point

without disrupting the remainder of the molecule. In particular, it is important not to subject the carboxy-protected benzodiazepinedione molecule to strong nucleophilic bases or reductive conditions employing highly activated metal catalysts such as Raney nickel. (Such harsh removal conditions are also to be avoided when removing amino-protecting groups and hydroxy-protecting groups, discussed below.) Preferred carboxylic acid protecting groups are the allyl and p-nitrobenzyl groups. Similar carboxy-protecting groups used in the cephalosporin, penicillin and peptide arts can also be used to protect a carboxy group substituents of the benzodiazepinedione. Further examples of these groups are found in E. Haslam, "Protective Groups in Organic Chemistry", J. G. W. McOmie, Ed., Plenum Press, New York, N.Y., 1973, Chapter 5, and T.W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, NY, 1981, Chapter 5. The term "protected carboxy" refers to a carboxy group substituted with one of the above carboxy-protecting groups.

As used herein the term "amide-protecting group" refers to any group typically used in the peptide art for protecting the peptide nitrogens from undesirable side reactions. Such groups include p-methoxyphenyl, 3,4-dimethoxybenzyl, benzyl, O-nitrobenzyl, di-(p-methoxyphenyl)methyl, triphenylmethyl, (p-methoxyphenyl)diphenylmethyl, diphenyl-4-pyridylmethyl, m-2-(picolyl)-N'-oxide, 5-dibenzosuberyl, trimethylsilyl, t-butyl dimethylsilyl, and the like. Further descriptions of these protecting groups can be found in "Protective Groups in Organic Synthesis", by Theodora W. Greene, 1981, John Wiley and Sons, New York.

Unless otherwise specified, the terms "heterocyclic group" or "heterocyclic" or "HET" or "heterocyclyl" are used interchangeably herein and refer to any mono-, bi-, or tricyclic saturated, unsaturated, or aromatic ring having the number of atoms designated where at least one ring is a 5-, 6- or 7-membered ring containing from one to four heteroatoms selected from nitrogen, oxygen, and sulfur. Preferably at least one heteroatom is nitrogen (*Lang's Handbook of Chemistry, supra*). Typically, the 5-membered ring has 0 to 2 double bonds and the 6- or 7-membered ring has 0 to 3 double bonds and the nitrogen or sulfur heteroatoms may optionally be oxidized, and any nitrogen heteroatom may optionally be quaternized. Included in the definition are any bicyclic groups where any of the above heterocyclic rings are fused to a benzene ring. Heterocyclics in which nitrogen is the heteroatom are preferred.

The following ring systems are examples of the heterocyclic (whether substituted or unsubstituted) radicals denoted by the term "heterocyclic": thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, thiatriazolyl, oxatriazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, thiazinyl, oxazinyl, triazinyl, thiadiazinyl, oxadiazinyl, dithiazinyl, dioxazinyl, oxathiazinyl, tetrazinyl, thiatriazinyl, oxatriazinyl, dithiadiazinyl, imidazolinyl, dihydropyrimidyl, tetrahydropyrimidyl, tetrazolo[1,5-b]pyridazinyl and purinyl, as well as benzo-fused derivatives, for example benzoxazolyl, benzthiazolyl, benzimidazolyl and indolyl.

Heterocyclic 5-membered ring systems containing a sulfur or oxygen atom and one to three nitrogen atoms are also suitable for use in the instant invention. Examples of such preferred groups include thiazolyl, in particular thiazol-2-yl and thiazol-2-yl N-oxide, thiadiazolyl, in particular 1,3,4-thiadiazol-5-yl and 1,2,4-thiadiazol-5-yl, oxazolyl, preferably oxazol-2-yl, and oxadiazolyl, such as 1,3,4-oxadiazol-5-yl, and 1,2,4-oxadiazol-5-yl. A group of further preferred examples of 5-membered ring systems with 2 to 4 nitrogen atoms include imidazolyl, preferably imidazol-2-yl; triazolyl, preferably 1,3,4-triazol-5-yl; 1,2,3-triazol-5-yl, 1,2,4-triazol-5-yl, and tetrazolyl, preferably 1H-tetrazol-5-yl.

Further suitable specific examples of heterocyclic ring systems are 6-membered ring systems containing one to three nitrogen atoms. Such examples include pyridyl, such as pyrid-2-yl, pyrid-3-yl, and pyrid-4-yl; pyrimidyl, preferably pyrimid-2-yl and pyrimid-4-yl; triazinyl, preferably 1,3,4-triazin-2-yl and 1,3,5-triazin-4-yl; pyridazinyl, in particular pyridazin-3-yl, and pyrazinyl. The pyridine N-oxides and pyridazine N-oxides and the pyridyl, pyrimid-2-yl, pyrimid-4-yl, pyridazinyl and the 1,3,4-triazin-2-yl radicals, are a preferred group.

An alternative suitable group of "heterocyclics" or "HET" include; 1,3-thiazol-2-yl, 4-(carboxymethyl)-5-methyl-1,3-thiazol-2-yl, 4-(carboxymethyl)-5-methyl-1,3-thiazol-2-yl sodium salt, 1,2,4-thiadiazol-5-yl, 3-methyl-1,2,4-thiadiazol-5-yl, 1,3,4-triazol-5-yl, 2-methyl-1,3,4-triazol-5-yl, 2-hydroxy-1,3,4-triazol-5-yl, 2-carboxy-4-methyl-1,3,4-triazol-5-yl sodium salt, 2-carboxy-4-methyl-1,3,4-triazol-5-yl, 1,3-oxazol-2-yl, 1,3,4-oxadiazol-5-yl, 2-methyl-1,3,4-oxadiazol-5-yl, 2-(hydroxymethyl)-1,3,4-oxadiazol-5-yl, 1,2,4-oxadiazol-5-yl, 1,3,4-thiadiazol-5-yl, 2-thiol-1,3,4-thiadiazol-5-yl, 2-(methylthio)-1,3,4-thiadiazol-5-yl, 2-amino-1,3,4-thiadiazol-5-yl, 1H-tetrazol-5-yl, 1-methyl-1H-tetrazol-5-yl, 1-(1-(dimethylamino)eth-2-yl)-1H-tetrazol-5-yl, 1-(carboxymethyl)-1H-tetrazol-5-yl, 1-(carboxymethyl)-1H-tetrazol-5-yl sodium salt, 1-(methylsulfonic acid)-1H-tetrazol-5-yl, 1-(methylsulfonic acid)-1H-tetrazol-5-yl sodium salt, 2-methyl-1H-tetrazol-5-yl, 1,2,3-triazol-5-yl, 1-methyl-1,2,3-triazol-5-yl, 2-methyl-1,2,3-triazol-5-yl, 4-methyl-1,2,3-triazol-5-yl, pyrid-2-yl N-oxide, 6-methoxy-2-(n-oxide)-pyridaz-3-yl, 6-hydroxypyridaz-3-yl, 1-methylpyrid-2-yl, 1-methylpyrid-4-yl, 2-hydroxypyrimid-4-yl, 1,4,5,6-tetrahydro-5,6-dioxo-4-methyl-as-triazin-3-yl, 1,4,5,6-tetrahydro-4-(formylmethyl)-5,6-dioxo-as-triazin-3-yl, 2,5-dihydro-5-oxo-6-hydroxy-as-triazin-3-yl, 2,5-dihydro-5-oxo-6-hydroxy-as-triazin-3-yl sodium salt, 2,5-dihydro-5-oxo-6-hydroxy-2-methyl-as-triazin-3-yl sodium salt, 2,5-dihydro-5-oxo-6-hydroxy-2-methyl-as-triazin-3-yl, 2,5-dihydro-5-oxo-6-methoxy-2-methyl-as-triazin-3-yl, 2,5-dihydro-5-oxo-as-triazin-3-yl, 2,5-dihydro-5-oxo-2-methyl-as-triazin-3-yl, 2,5-dihydro-5-oxo-2,6-dimethyl-as-triazin-3-yl, tetrazolo[1,5-b]pyridazin-6-yl and 8-aminotetrazolo[1,5-b]-pyridazin-6-yl.

An alternative group of "heterocyclics" includes; piperidyl, piperazinyl, morpholinyl, morpholino, 4-(carboxymethyl)-5-methyl-1,3-thiazol-2-yl, 4-(carboxymethyl)-5-methyl-1,3-thiazol-2-yl sodium salt, 1,3,4-triazol-5-yl, 2-methyl-1,3,4-triazol-5-yl, 1H-tetrazol-5-yl, 1-methyl-1H-tetrazol-5-yl, 1-(1-(dimethylamino)eth-2-yl)-1H-tetrazol-5-yl, 1-(carboxymethyl)-1H-

tetrazol-5-yl, 1-(carboxymethyl)-1H-tetrazol-5-yl sodium salt, 1-(methylsulfonic acid)-1H-tetrazol-5-yl, 1-(methylsulfonic acid)-1H-tetrazol-5-yl sodium salt, 1,2,3-triazol-5-yl, 1,4,5,6-tetrahydro-5,6-dioxo-4-methyl-as-triazin-3-yl, 1,4,5,6-tetrahydro-4-(2-formylmethyl)-5,6-dioxo-as-triazin-3-yl, 2,5-dihydro-5-oxo-6-hydroxy-2-methyl-as-triazin-3-yl sodium salt, 2,5-dihydro-5-oxo-6-hydroxy-2-methyl-as-triazin-3-yl, tetrazolo[1,5-b]pyridazin-6-yl, and 8-aminotetrazolo[1,5-b]pyridazin-6-yl.

Bivalent radicals L whether branched or unbranched, derived from alkanes, alkenes, alkadienes, alkynes, alkadiynes, and arenes optionally containing O, N and/or S atoms, or homo- and heterocycles either aromatic or aliphatic, are designated by adding the suffix "ene" to the corresponding monovalent radical. Atoms bearing the free valences may include any C, O, N or S.

Each substituent or term used in any formula or expression herein, e.g. R^n , A^n , B^n , Z, X, C_1 - C_4 alkyl, etc., when it appears more than once, is independent of its definition elsewhere in that or any other formula or structure.

"Pharmaceutically acceptable salts" include both acid and base addition salts.

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases and which are not biologically or otherwise undesirable; formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, carbonic acid, phosphoric acid and the like, and organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic, and sulfonic classes of organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, gluconic acid, lactic acid, pyruvic acid, oxalic acid, malic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, aspartic acid, ascorbic acid, glutamic acid, anthranilic acid, benzoic acid, cinnamic acid, mandelic acid, embonic acid, phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

"Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Salts derived from pharmaceutically acceptable organic nontoxic bases includes salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-diethylaminoethanol, trimethamine, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic non-toxic bases are isopropylamine, diethylamine, ethanolamine, trimethamine, dicyclohexylamine, choline, and caffeine.

The term "prodrug" as used herein means a derivative of a parent drug molecule that enhances pharmaceutically desirable characteristics or properties (e.g. transport, bioavailability, pharmacodynamics, etc.) and that requires biotransformation, either spontaneous or enzymatic, within the organism to release the active parent drug.


5 In the description that follows standard abbreviations as recommended by the *Journal of Organic Chemistry* (see "Guidelines for Authors" in any volume) are employed unless otherwise specified. For convenience, the following standard abbreviations used herein have the meanings provided below.

ABBREVIATIONS

10	ACN	acetonitrile
	Boc or BOC	<i>tert</i> -butoxycarbonyl
	BOP	benzotriazolyloxy-trisdimethylamino-phosphonium hexafluorophosphate
15	DCC	dicyclohexylcarbodiimide
	DCM	dichloromethane
	DEAD	diethylazodicarboxylate
	DIPC	diisopropylcarbodiimide
	DIPEA	diisopropylethylamine
20	DMA	dimethylacetamide
	DMAP	dimethylaminopyridine
	DMF	dimethylformamide
	DMSO	dimethylsulfoxide
	DTT	dithiothreitol
25	EDC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
	EtN(iPr) ₂	diisopropylethylamine
	Fmoc or FMOC	fluorenylmethoxycarbonyl
	HOBt	N-hydroxybenzotriazole
	HPLC	high pressure liquid chromatography
30	LDA	lithium diisopropylamine
	MBHA	methylbenzhydramine
	NMM	N-methylmorpholine
	Ph ₃ P	triphenylphosphine
	PyBrOP	bromo-tri-pyrrolidinophosphoniumhexafluorophosphate
35	SPPS	solid phase peptide synthesis
	<i>t</i> -BuO ₂ C	<i>t</i> -butoxycarbonyl
	TFA	trifluoroacetic acid

B. Utility

The present invention is the result of the unexpected discovery that substituted fused 6-7-6 or 6-7-5 membered benzodiazepine compounds defined by Formula I and II, referred to hereafter as "substituted tricyclic benzodiazepines", are capable of acting as fibrinogen receptor antagonists inhibiting both platelet aggregation and binding of fibrinogen to the platelet receptor, GPII_bIII_a, at concentrations comparable to many of the more potent anti-aggregatory peptides of the prior art (see e.g. Ali *et al.* EP 341 915). Thus, the instant fibrinogen receptor antagonists are particularly useful when they inhibit the binding of fibrinogen to the GPII_bIII_a receptor at an IC₅₀ in a Fg/GPII_bIII_a ELISA as described herein of at least about 35nM, and an IC₅₀ in a platelet aggregation (PA) assay, also described herein, of at least about 3μM. Since there is some variability in the values obtained for ELISA and PA assays in different laboratories, preferred fibrinogen receptor antagonists represented by Formula I and II will have IC₅₀'s for Fg/GPII_bIII_a ELISA's and PA assays equal to or greater than that of the prior art compound N-[3-(4-amidinobenzamido)benzoyl]-3-aminopropionic acid,




This compound is used as an internal standard and has been found to have an Fg/GPII_bIII_a ELISA IC₅₀ of about 25 nM and a PA inhibition assay IC₅₀ of about 3μM when measured according to the assays described herein. Most preferred fibrinogen receptor antagonists represented by Formula I and II have IC₅₀'s in Fg/GPII_bIII_a ELISA of at least about 15 nM and in platelet aggregation inhibition assays of at least about 1μM or three times more potent in a platelet aggregation inhibition assay than the reference compound N-[3-(4-amidinobenzamido)benzoyl]-3-aminopropionic acid. This reference compound is readily prepared by standard methods from commercially available precursors (e.g. β-alanine, *m*-aminobenzoic acid, and *p*-amidinobenzoic acid) or may be synthesized according to the procedures set forth in EP 372 486 A2.


The compounds described in the present invention may therefore inhibit the binding of fibrinogen to its receptor on platelets, GPII_bIII_a, and thus prevent the aggregation of platelets and the formation of platelet plugs, emboli and thrombi in the circulatory system in mammals. Thromboembolic disorders have been shown to be directly related to the susceptibility of blood platelets to aggregate. Mammals exposed to medical procedures such as angioplasty and thrombolytic therapy are particularly susceptible to thrombus formation. The compounds of the present invention can be used to inhibit thrombus formation following angioplasty. They may also be used in combination with antithrombolytic agents such as tissue plasminogen activator and its derivatives (US patents 4,752,603; 4,766,075; 4,777,043; EP 199 574; EP 238 304;

subsequent to the antithrombolytic agent. Mammals exposed to renal dialysis, blood oxygenation, cardiac catheterization and similar medical procedures as well as mammals fitted with certain prosthetic devices are also susceptible to thromboembolic disorders. Physiologic conditions, with or without known cause may also lead to thromboembolic disorders. Thus, the compounds described herein may be useful in treating thromboembolic disorders in mammals. The compounds described herein may also be used as adjuncts to anticoagulant therapy, for example in combination with aspirin, heparin or warfarin and other anticoagulant agents. The application of the compounds described herein for these and related disorders will be apparent to those skilled in the art.

Compounds of this invention are also useful as intermediates generally, or as precursors of inhibitors to GPIIb/IIIa or other integrin receptors and thus in addition to treating cardiovascular disease, these compounds may be usefully employed in metastatic disease, or for any disease where antagonism of RGD binding integrin receptors is indicated.

Compounds represented by formula I or II are capable of antagonizing the interaction between vitronectin (Vn) and the vitronectin receptor (VnR) and thus are useful in situations where such antagonism is indicated. Thus it is contemplated the instant compounds are useful in treating bone disorders such as osteoporosis. Vn-VnR antagonism is particularly pronounced when substituent Q-L- is bonded to position 8 or 9 based on the benzodiazapinedione numbering system. Exemplary Vn-VnR antagonists are represented by structural formula Ib and IIb below, where the substituents are as previously defined.










C. Preferred Embodiments


Nonpeptidyl GPIIb/IIIa Inhibitors

- 5 One embodiment of the invention comprises a compound represented by Formula I and II, capable of inhibiting binding of the platelet GPIIb/IIIa receptor to its native *in vivo* ligands. Preferred nonpeptidyl inhibitors include compounds represented by structural Formulae III-XXI and prodrug forms thereof:







VII




VIII




IX




X




XI




XII




XIII




XIV




XV




XVI




XVII




XVIII



XIX



XX



XXI

10 where R¹, R², R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²⁵, R²⁶, Q, and L are defined above.

Referring to Formulae I-XXI the following structural features of the instant nonpeptidyl antithrombotic inhibitors can be identified:

- a. The positively charged Q moiety;
- b. The linking moiety L;
- 5 c. The flat aromatic "benzo" ring ;
- d. The 7-member "diazepine" ring;
- e. The fused 5 or 6 member "amidino containing " ring; and
- f. The negatively charged acidic moiety COR²².

10 a. Positively charged Q


Suitable groups Q contain one or more nitrogen atoms and have a pK_b sufficiently high so that they are at least 10% positively charged at physiological pH. Q may be one or more primary, secondary, tertiary, or quaternary amines or imines either isolated or conjugated with other nitrogen atoms to form groups including but not limited to; aminomethyleneimino, 15 amidino, and guanidino groups and multiples thereof. Alternatively, Q may be a saturated or unsaturated (including aromatic) heterocyclic group provided the group bears a positive charge at physiological pH. In one embodiment of the invention, Q is preferably selected from; amino (H₂N-), imino (=NH), amidino (H₂NC(=NH)-), aminomethyleneamino (H₂N-CH=N-), iminomethylamino (HN=CH-NH-), guanidino (H₂N-C(=NH)-NH-), N^G-aminoguanidino 20 (H₂N-HN-C(=NH)-NH-), alkylamino (R¹NH-), dialkylamino (R¹₂N-), trialkylamino (R¹₃N-), alkylideneamino (R¹₂C=N-), pyranyl, pyrrolyl, imadazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3H-indolyl, indolyl, 1H-indazolyl, purinyl, 4H-quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnoliny, pteridinyl, 4aH-carbazolyl, carbazolyl, b-carboliny, phenanthridinyl, 25 acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, phenarsazinyl, phenothiazinyl, pyrrolinyl, imidazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, indolinyl, isoindolinyl, quinuclidinyl, morpholinyl, morpholino, 1,3-diazacyclohex-4-ene, and combinations thereof, where R¹ is selected from; hydrogen, C₁-C₆-alkyl, C₆-C₁₀-aryl, and C₃-C₁₀-cycloalkyl. Optionally, any of the nitrogen containing heterocycles described above may 30 be substituted with amino, imino, amidino, aminomethyleneamino, iminomethylamino, guanidino, N^G-amino-guanidino, alkylamino, dialkylamino, trialkylamino, or alkylideneamino groups.

Exemplary preferred Q groups include the following:



(A) Amino and ammonio groups;

35







(B) Amidino groups;




5




10

(C) Aminoalkyleneimino groups;



15



(D) Iminoalkyleneamino groups;


20



25



5



(E) Guanidino groups;

10






15



20







25







(F) Saturated heterocyclic groups;





5







10





15




20







, and





5

(G) Unsaturated (nonaromatic) heterocyclic groups;





10





N(CH₃)


15



20





, and



25

(H) Unsaturated aromatic heterocyclic groups:





















5

10





15

20





25






5



10





15








(I) Bicycloheterocyclic groups;

20



25





5 Most preferred Q groups are amino, amidino, and piperazinyl groups.

To increase oral bioavailability the positively charged Q moiety can be made neutral. For example R^3 , R^4 , or R^5 may be a substituent that transforms compounds represented by structural formulae I-XXI into a prodrug. Exemplary prodrugs include hydroxyamidines and carbamates. Preferred R^3 , R^4 , or R^5 groups of this embodiment include;



10

HO-,






C₁-C₁₀alkyl-O-(C=O)-,

C₁-C₁₀alkyl-S-(C=O)-,


Cl₃CCH₂O-(C=O)-,



15




20



5 **b. The linking group L**


10 The length of the bivalent radical L appears to be important to biological activity. By length is meant the distance between the "benzo" ring and the first charge bearing N of substituent Q. For example, suitable lengths for bivalent radical L range from about 3 to about 9 and preferably 5 or 6 methylene equivalents, where any methylene group or groups may be replaced with one or more or a combination of; alkene, alkyne, aryl, heterocycle or a functional group or groups containing the heteroatoms N, O, and S, so long as the overall length is equivalent to from 3 to 9 methylene groups. These functional groups include one or more of the following:




15 and may be isolated within the linker (e.g. forming ethers, thioethers, ketones, sulfoxides and the like) or combined in any combination, provided only that the compounds so produced are stable in aqueous solution and do not exceed the above stated length requirements. For
20 example, combining these functional groups produces esters, amides, ureidos, carbamates, carbonates, sulfonamides, sulfoxides, sulfones, and the like. The linker may optionally be substituted with one or more substituents selected from loweralkyl, haloloweralkyl, loweralkoxy, cyano, hydroxy, halo, haloloweralkoxy and phenyl optionally substituted with halo or haloloweralkyl, provided only that the linker is stable in aqueous solution and does not
25 exceed the length requirements.

Preferred L's are selected from substituted or unsubstituted; C₃-C₇-alkylene, C₃-C₇-cycloalkylene, C₃-C₇-alkenylene, C₄-C₇-cycloalkenylene, C₅-C₈-cycloalkadienylene, C₃-C₇-alkadienylene, C₃-C₇-alkynylene, C₄-C₇-alkenynylene, C₆-C₁₄-arylene, C₆-C₁₄-aryl-C₂-C₄-alkynylene, C₁-C₃-alkyl-C₆-C₁₄-aryl-C₂-C₄-alkynylene, C₆-C₁₄-aryl-C₂-C₄-alkenylene, C₁-C₃-alkyl-C₆-C₁₄-arylene, C₁-C₃-alkyl-C₆-C₁₄-aryl-C₂-C₄-alkenylene, C₆-C₁₄-aryl-C₁-C₃-alkylene, C₆-C₁₄-aryl-C₁-C₃-alkyloxyene, C₁-C₃-alkyl-C₆-C₁₄-aryl-C₁-C₂-alkylene, C₁-C₃-alkyloxy-C₆-C₁₄-arylene, C₂-C₈-alkyloxyene, C₁-C₅-alkyloxy-C₁-C₅-alkylene, C₆-C₁₀-aryloxyene, C₆-C₁₀-aryloxy-C₁-C₅-alkylene, C₆-C₁₀-arylthio-C₁-C₅-alkylene,


10




15





20



25



30







5





where R^{14} is selected from; a chemical bond, C₁-C₈-alkyl, C₁-C₈-alkyloxy, C₃-C₇-cycloalkyl C₂-C₅-alkenyl, C₃-C₅-alkynyl, C₆-C₁₀-aryl, C₁-C₃-alkyl-C₆-C₁₂-aryl, C₁-C₂-alkyl-C₆-C₁₀-aryl-
 10 C₁-C₂-alkyl, C₆-C₁₀-aryl-C₁-C₂-alkyl, and C₆-C₁₀-aryloxy-C₁-C₂-alkyl. R^{15} is selected from; a chemical bond, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₆-C₁₀-aryl, and C₁-C₃-alkyl-C₆-C₁₂-aryl. R^{16} is selected from; a chemical bond, C₁-C₅-alkyl, C₃-C₇-cycloalkyl C₃-C₅-alkenyl, C₃-C₅-alkynyl, C₆-C₁₀-aryl, C₁-C₃-alkyl-C₆-C₁₂-aryl, and C₆-C₁₀-aryl-C₁-C₂-alkyl. The substituents are preferably selected from one to three R^{12} defined above.

15



More preferred bivalent radicals L are selected from the following:






20




25






5




Most preferred Q-L- combinations are selected from:












10



15



20



5

10









15

2166811

2166811

WO 95/04057

PCT/US94/07989







5




10

15


5




10



15






, and





where any amino, amidino, or guanidino nitrogen is optionally derivitized with C₁-
20 C₆alkoxycarbonyl.

Very most preferred Q-L- combinations are selected from;




5




10




15



20






5

c. The "benzo" ring

Preferred substituents of the "benzo" ring are R^1 and R^2 , where each R^1 and R^2 , is independently selected from hydrogen, halogen(F, Cl, Br, I), cyano, nitro, carboxyl, mercaptocarbonyl, mercapto, phthalimido and a substituent having from 1 to 12 carbon atoms selected from: alkyl, hydroxyalkyl, formyl, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, aralkyl, aralkylhaloalkyl, aryl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkyl, carboxylalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, alkylthiocarbonyl, aralkylthiocarbonylthio, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, and cycloheteroalkylcarbonylalkyl where each heteroaryl- and cyclohetero-containing groups has one or more ring hetero atoms selected from oxygen, sulfur and nitrogen. Optionally, each R^1 and R^2 also may be independently selected from sulfonyl, sulfonylamido, amino and amido radicals of the formula:




30

1780015

2166811

WO 95/04057

PCT/US94/07989




where X is oxygen or sulfur, each "a" is a number independently selected from zero to six, inclusive, and where each of R³ through R⁵ is independently selected from the groups described above. Optionally, R³ and R⁴ taken together, R⁴ and R⁵ taken together and R³ and R⁵ taken together may each form a heterocyclic having from five to seven ring members including the hetero atom of the sulfonyl, amino or amido radical and which heterocyclic may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen, and sulfur and which heterocyclic group may be saturated or partially saturated. R²⁸ is selected from the groups defined below.

d. The 7-member "diazepine" ring;

Substituents of the 7-member "diazepine" ring are preferably hydrogen or halogen. Optionally any oxo group (=O) may be substituted with an =S group.

e. The fused 5- or 6-member "amidino containing" ring;

Preferred 5-member "amidino containing" rings of Formula I are selected from;




20

where R¹ is preferably hydrogen, lower alkyl, lower alkyl-morpholino, hydroxy lower alkyl, carboxy lower alkyl, amino lower alkyl, and aminocarbonyl lower alkyl; R² is preferably hydrogen, lower alkyl-morpholino, the side chain of a naturally occurring α -amino acid, or the D stereoisomer thereof; R²⁵ is preferably hydrogen, or lower alkyl; and R²⁶ is preferably lower alkyl.

25

Preferred 6-member "amidino containing" rings of Formula II are selected from;

5




10 where R^1 , R^2 , R^{25} , and R^{26} are defined above.

15 f. The negatively charged acidic moiety COR^{22} ;
 R^{22} is selected from the group consisting of (a) hydroxy, (b) C_1 - C_8 -alkoxy, (c) C_3 - C_{12} -alkenoxo, (d) C_6 - C_{12} -aryloxy, (e) C_1 - C_6 -alkyl- C_6 - C_{12} -aryloxy, (f) di- C_1 - C_8 -alkylamino- C_1 - C_8 -alkoxy, (g) acylamino- C_1 - C_8 -alkoxy selected from the group (i) acetylaminooethoxy, (ii) nicotinoylaminoethoxy, and (iii) succinamidoethoxy, (h) C_1 - C_8 -alkoxyloxy- C_1 - C_8 -alkoxy, (i) C_6 - C_{12} -aryl- C_1 - C_8 -alkoxy where the aryl group is unsubstituted or substituted with one to three of the groups (i) nitro, (ii) halo (F, Cl, Br, I), (iii) C_1 - C_4 -alkoxy, and (iv) amino, (j) hydroxy- C_2 - C_8 -alkoxy, (k) dihydroxy- C_3 - C_8 -alkoxy, and (l) $NR^{23}R^{24}$.



R^{23} and R^{24} are independently selected from the group (a) hydrogen, (b) C_1 - C_8 -alkyl, (c) C_3 - C_8 -alkenyl, (d) C_6 - C_{12} -aryl where the aryl group is unsubstituted or substituted with one to three of the groups (i) nitro, (ii) halo (F, Cl, Br, I), (iii) C_1 - C_4 -alkoxy, and (iv) amino, and (e) C_6 - C_{12} -aryl- C_1 - C_8 -alkyl where the aryl group is unsubstituted or substituted with one to three of the groups (i) nitro, (ii) halo (F, Cl, Br, I), and (iii) C_1 - C_4 -alkoxy.

R^{22} is preferably a substituent that transforms compounds represented by structural formulae I-XXI into prodrugs. For example, preferred prodrug forms include simple esters, α -acyloxyalkyl esters, and amides. Preferred esters include compounds where R^{22} is selected from;


10










15









20






5

10















, and















n= 2- 4

Preferred compounds of this invention are represented by structural Formula I and II and are selected from Tables I and II below, where R^1 is hydrogen or chlorine and R^{18} , R^{19} , R^{20} and R^{21} are hydrogen or methyl, and the other substituents are as set forth in the tables.

TABLE I

1180015

2166811

WO 95/04057

PCT/US94/07989

11888193

2166811

WO 95/04057

PCT/US94/07989

186312

2166811

WO 95/04057

PCT/US94/07989

1182215

2166811

WO 95/04057












PCT/US94/07989













2166811













2166811

WO 95/04057

PCT/US94/07989

780013

2166811

WO 95/04057









PCT/US94/07989

180015

2166811

WO 95/04057

PCT/US94/07989












	
	
	
	
	













20012












2166811













WO 95/04057

PCT/US94/07989

$Q-L-$	
	
	
	
	
	
















	
	
	
	
	
	














180015

2166811

WO 95/04057

PCT/US94/07989

$Q-L-$	
	
	
	
	
	
	
	








$Q-L-$	
	
	
	
	
	
	

180015

2166811

WO 95/04057

PCT/US94/07989










	
	
	
	
	












186015












2166811














WO 95/04057

PCT/US94/07989












	
	
	
	
	
	
	














2166811










2166811













WO 95/04057













PCT/US94/07989















	
	
	
	
	
	

1188815

2166811

WO 95/04057

PCT/US94/07989

188815

2166811

WO 95/04057

PCT/US94/07989

1182015

2166811

WO 95/04057















PCT/US94/07989

188815

2166811

WO 95/04057

PCT/US94/07989










$Q-L-$	
	
	
	
	
	
	
	










180015

2166811

WO 95/04057

PCT/US94/07989












$Q-L-$	
	
	
	
	

















180015












2166811

WO 95/04057

PCT/US94/07989

1180015

2166811

WO 95/04057











PCT/US94/07989

1180015

2166811

WO 95/04057

PCT/US94/07989











	
	
	
	
	

118001S

2166811

WO 95/04057

PCT/US94/07989

1188819

2166811

WO 95/04057

PCT/US94/07989









1880155

2166811

WO 95/04057

PCT/US94/07989

TABLE II

118001S

2166811

WO 95/04057

PCT/US94/07989

180815

2166811

WO 95/04057










PCT/US94/07989














1180012











2166811















WO 95/04057










PCT/US94/07989








	
	
	
	
	










	
	
	
	
	
	
	

$Q-L-$	
	
	
	
	











	
	
	
	
	

















118001S

2166811

WO 95/04057

PCT/US94/07989










	
	
	
	
	
	
	
	













111800015














2166811











WO 95/04057














PCT/US94/07989










	
	
	
	
	














	
	
	
	
	
	










	
	
	
	
	
	
	










	
	
	
	
	












	
	
	
	
	
	
	















	
	
	
	
	

$Q-L-$	
	
	
	
	
	
	

$Q-L-$	
	
	
	
	

$Q-L-$	
	
	
	
	
	


D. Methods of Making

Compounds of the present invention can be prepared by methods employing standard chemical methodologies described and referenced in standard textbooks (e.g. March, J. "Advanced Organic Chemistry" McGraw-Hill, New York, 1977; Collman, J.P., Hegedus, L.S., Norton, J.R., Finke, R.G. "Principles and Applications of Organotransition Metal Chemistry" University Science, Mill Valley, 1987; Larock, R.C. "Comprehensive Organic Transformations" Verlag, New York, 1989). For a detailed description of the preparation of the intermediate, substituted 3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione **4** and of adding the substituent Q-L, see WO 93/08174.

1. 5-member heterocycles fused to 1,4-benzodiazepine

The key intermediate for synthesis of 5-member heterocycles fused to 1,4-benzodiazepines is the substituted 3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione **4**. The benzodiazepinedione **4** may be prepared from isatoic anhydride **1**, a β -alanine ester or a β -amino acid and an α -haloacetyl halide, according to Scheme 1. Methods for preparation of the isatoic anhydrides **1** are known in the art and are available from commercial sources such as Aldrich Chemical Co.

Scheme I



Briefly, isatoic anhydride is converted to N-(2-aminobenzoyl)- β -alanine ester **2** by allowing **1** to react with a β -alanine ester or its salt, in the presence an organic base. The isatoic anhydride **1** may be substituted or unsubstituted, but the 5-iodo isatoic anhydride is preferred ($1, X^*=I$). The β -alanine ester may also be substituted or unsubstituted as the free


amine or, more conveniently, as its salt (e.g. HCl). Generally, the reaction is conducted in a dry polar aprotic solvent, such as dimethylformamide or the like, in the presence of an equimolar amount, or up to 30% excess, of a tertiary amine as the organic base, e.g. triethylamine or 2,6-lutidine. Alternatively, the reaction may be catalysed by
5 dimethylaminopyridine (Venuti, M.C. *Synthesis* 266 (1982)). The reaction requires between about 0.5 to 2 days at temperatures between about room temperature and 200°C. Preferably, the reaction is conducted at 80°C for about 16 hours in dimethylformamide as the solvent. The product is isolated and purified by conventional methods, typically by solvent extraction into a compatible solvent. The product may be further purified by column chromatography or
10 other appropriate methods.

Acylation of 2 with an α -haloacyl halide is carried out in methylene chloride under a dry inert atmosphere, such as nitrogen. The reaction is allowed to run at temperatures between 0°C and room temperature for about 2 hours, and the product 3 is isolated or purified by conventional methods.

15 Cyclization of the acylated N-[2-(α -bromoacetyl)benzoyl]- β -alanine ester 3 may be achieved by reaction with a base such as an alkali metal alkoxide, hydride, or carbonate in a polar solvent at temperatures between about 0°C and 100°C for about 0.5 to 2 hours. For example, a solution of the ester 3 is commonly added over a period of approximately 15 to 60 minutes to a slurry of an alkali metal carbonate in an appropriate
20 solvent, cooled to 0°C. It is preferable that the solvent will be a polar aprotic solvent such as dimethylformamide and the alkali metal carbonate be sodium carbonate, calcium carbonate, or cesium carbonate. Once the addition is complete, the reaction mixture is generally allowed to warm to room temperature and run for an additional 60 to 180 minutes after which the reaction is neutralized by the addition of a solution of an acid, such as 10% citric acid or the
25 like, and the solvent evaporated. The product is then isolated by solvent extraction and further purified by column chromatography.

Preparation of 5-member rings fused to the substituted 3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione 4 is carried out according to Scheme II-III. First, the benzodiazepine-dione 4 is treated with Lawesson's reagent to produce the thiolactam 5.
30 Alkylation of the thiolactam 5 is carried out in a bi-phasic mixture of CH₂Cl₂ and basic H₂O by treatment with methyl iodine and a catalytic amount of tetrabutylammonium hydrogen sulfate to produce the intermediate thioimide ester 6.

Scheme II





5

The thioimidate ester intermediate 6 may then be reacted with a variety of reagents to produce the fused 5-member rings of this invention according to Scheme III.

Scheme III

10







Briefly, 6 may be heated to reflux in a toluene/pyridine•HCl solution with a substituted propargylamine to produce the substituted imidazo-benzodiazepine 7. Alternatively, 6 may be similarly refluxed with an appropriately substituted acylhydrazide to give the triazolo-benzodiazepine 8. Optionally, the thioimide ester intermediate 6 may be refluxed in toluene/pyridine•HCl with a substituted carbazate ester or α -amino acid ester to produce respectively the oxotriazolo-benzodiazepine 9 or oxoimidazolo-benzodiazepine 10. Alternatively, the thiolactam 5 (Scheme II) may be reacted with an appropriately protected α -amino acid and cyclized with dicyclohexylcarbodiimide to yield 10 (see Ager *et al.*, *J. Med. Chem.* 20(8):1035-1040 (1977)). Finally, 6 may be heated in DMF with sodium azide to form the tetrazolo-benzodiazepine 11. The tetrazolo-benzodiazepine 11 can also be prepared directly from 4 by conversion of the secondary amide directly into a tetrazole using triphenylphosphine and trimethylsilylazide (see Duncia *et al.*, *J. Org. Chem.* 56(7):2395-2400 (1991)).

2. 6-member heterocycles fused to 1,4-benzodiazepine

6-member heterocycles fused to 1,4-benzodiazepines can be prepared by methods known in the art (see e.g. EP 0 519 678 A2). More specifically, beginning with the substituted 3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione 4 (Scheme I) and reacting it with Lawesson's reagent as described in Scheme II produces the thiolactam 5. This intermediate can be further reacted to produce substituted oxodiazines according to Scheme IV.

Scheme IV






Briefly, the thiolactam 5 may be treated with ammonia in the presence of a mercury (II) salt to form the intermediate amidine 12. Reacting 12 with diketene followed by cyclization with heat treatment yields the substituted pyridinone-benzodiazepine 14. Alternatively, the thiolactam 5 may be treated with methyl iodide and refluxed in toluene/pyridine•HCl with an optionally substituted anthranilate ester or a substituted β -alanine ester to produce respectively the substituted quanzolino-benzodiazepine 15 or substituted pyrimidone-benzodiazepine 16.

Other substituted pyrimidone-benzodiazepines are prepared from the intermediate amidine 12 according to reaction Scheme V.

Scheme V




5
10 Briefly, the intermediate amidine 12 is heated with ethoxymalonic acid diester to yield the carboxyethyl substituted pyrimidineone-benzodiazepine 17. Saponification of 17 yields the free acid 18 which may be decarboxylated by heating to give compound 19. Alternatively, the carboxyethyl substituted pyrimidineone-benzodiazepine 17 may be reduced with sodium cyanoborohydride to give the corresponding alcohol 20, which can be

alkylated with 2,2-dimethyl-1,3-dioxolan-4-ylmethyl-*p*-toluenesulfonate to yield the glycerol-isopropylidene adduct 21.

The carboxyethyl substituted pyrimidone-benzodiazepine 17 also provides access to a wide variety of primary, secondary, and tertiary amine adducts as shown in Scheme VI.

5

Scheme VI




10

For example, the hydroxyethyl substituted pyrimidineone-benzodiazepine 20 can be treated with a mild oxidizing agent such as manganese dioxide to yield the corresponding aldehyde 23 which in turn can be reductively aminated to produce the primary and tertiary amines 25, 26, and 27. Optionally a Wolff-Kishner reaction converts aldehyde 23 to the methyl compound 24.

The amidine 12 of Scheme V may also be converted into triazinodiones according to Scheme VII.


Scheme VII



Briefly, amidine 12 is treated with ethoxycarbonyl isocyanate to yield compound 28. Upon heating, 28 is cyclized to the triazinodione 29. Optionally, 29 may be alkylated with an alkyl halide to give the alkyl substituted triazinodione 30.

Triazinobenzodiazepines can be prepared from benzodiazepines according to Scheme VIII.

Scheme VIII




5 Briefly, 1,4-benzodiazepines 4 are alkylated with ethylbromoacetate and sodium hydride to give the N-1 alkylated carboxyethyl ester 31. Treatment of 31 with Lawesson's reagent produces the N-1 alkylated thiolactam 32 which, upon treatment with an alkyl substituted hydrazine, cyclizes to give the triazinobenzodiazepines 33. Alternatively, reaction of thiolactam 32 with hydrazine followed by alkylation with an alkyl halide as shown in Scheme VII yields 33.


10

Other triazinobenzodiazepines can be prepared from the intermediate thiolactam 5 (Scheme II) according to Scheme IX.

Scheme IX





15



Briefly, thiolactam 5 is treated with hydrazine to give the intermediate
 5 amidrazone 34. Amidrazone 34 may be reacted with haloaldehydes and ketones to produce
 hydrazones 35 which, upon heating, gives triazinobenzodiazepine 36. Alternatively,
 treatment of 34 with α -keto esters yields hydrazones 37 which are cyclized by heating in
 acetic acid to give oxo-triazinobenzodiazepines 38. Additionally, treatment of 34 with
 ketoacetals gives hydrazones 39 which upon treatment with concentrated sulfuric acid gives
 10 hydroxy-triazinobenzodiazepines 40. Finally, treatment of amidrazone 34 with
 chlorocarbonyl ethylformate produces the oxalic ester hydrazid 41 which is cyclized in
 refluxing pyridine to 1,2-dicarbonyl-triazinobenzodiazepines 42 according to Scheme X.

Scheme X






Other oxo-triazinobenzodiazepines and 5-member sulfonamides can be prepared from the intermediate thioimide ester 6 according to Scheme XI.

5

Scheme XI



10 Treatment of 6 with hydrazinoesters yields amidrazonoesters 45 which upon heating in acetic acid gives substituted oxo-triazinobenzodiazepines 46. Compound 6 may also be reacted with α -amino sulfonic acid esters or sulfonamidrazone esters in pyridine/HCl to yield sulfonamides 43 and 44 respectively.

15

3. The Q-L Group

The nitrogen containing substituents Q or a precursor thereof may be added to the linker L and the combination Q-L-, usually in protected form, bonded to the "benzo" ring.


Alternatively, Q or Q plus a portion of L, referred to below as L', may be added to L or a portion thereof after the "benzo" ring and the rest of the molecule have been formed. Q itself may be prepared and bonded to the "benzo" ring by standard methods published in both the scientific and patent literature (see e.g. US patents; 4,992,542, 4,997,936, 4,194,047, 5,003,076, 5,063,207, 5,063,208, and 5,079,357, and references cited therein).

In the description following, addition of Q-L- (or Q-L') to the benzene moiety of the imidazo-benzodiazepine 7 where R¹ and R² are both hydrogen is specified. It will be understood that these same procedures may be applied to other ring systems described in schemes I-IX above and encompassed by formula I and II. For a more comprehensive description of methods for adding Q-L- (or Q-L') to the benzene moiety of the imidazo-benzodiazepine see WO 93/08174.

Generally, conversion of the compounds having the structural formula 7 to compounds having the structural formula 48 may be accomplished by allowing 7 to react with an alkyne 47 in the presence of a palladium (II) salt, a copper (I) salt, and an organic base. Preferably, the alkyne 47 is substituted in a way such that the Y group can be synthetically converted to the positively charged Q group of structural Formula I and II. More preferably, the Y group is a protected form of the positively charged Q group. For example, alkyne 47 may be a N-Boc-amino alkyne, a benzo or alkynitrile alkyne, a nitrobenzo alkyne, or the like.

Referring to Scheme XII, the reaction is allowed to proceed in a dry organic polar aprotic solvent, such as ethyl acetate or the like, with the exclusion of oxygen, at from room temperature to 180°C for times between about 2 and 48 hours.

Scheme XII



Preferably, the reaction will run with a catalytic amount of palladium(II) salt at 10 molar percent and copper (I) salt at 5 molar percent. A 2 fold excess of the alkyne 47 and a 5 fold excess of a tertiary amine as the organic base, such as triethylamine or the like, is preferred.

- 5 The product 48 can be isolated by solvent extraction in to a suitable organic solvent, such as ethyl acetate, and washed with a solution of 10% ethylenediaminetetraacetic acid and the solvent evaporated. 48 may be further purified by column chromatography.

- Preparation of compounds 49 and 50 may be achieved by selective reduction of the alkyne moiety in compounds of the general formula 48 to yield alkene 49 and alkane 50 by
10 allowing the alkyne to stir under an atmosphere of hydrogen in the presence of a small amount of palladium on carbon. Typically, the reaction is run in an inert solvent, such as ethyl acetate, with a 5-10 molar percent by weight of 10% palladium on carbon at temperatures between about room temperature and 50°C for times between about 15 and 240 minutes. Preferably, the reaction is carried out at room temperature for 1 hour. The products
15 are isolated by filtration of the mixture through a filter agent, such as Celite®, and evaporation of solvent.

- Conversion of the compounds of the general formula 48, 49, or 50 in which Y is a N-(*tert*-butoxycarbonyl) amino protected moiety (N-BOC) to the amino ester can be accomplished by allowing the material to react with a strong acid. Generally, the reaction is
20 carried out by mixing the N-BOC amino ester with a large excess of concentrated solution of a strong acid, such as hydrogen chloride, dissolved in an appropriate inert solvent, such as ethyl acetate. The reaction can be conducted at a temperature between about minus 30°C and room temperature for about 0.5 to 24 hours. The reaction can be run in the presence of a molar amount of trialkylsilane, such as triethylsilane or the like. Preferably, the reaction will be
25 carried out at room temperature for 2 hours with a molar equivalent of triethylsilane in ethyl acetate. The solvent is evaporated and the products can be further purified by high-pressure liquid chromatography using a reverse phase column.

- Conversion of compounds of the general formula 48, 49, or 50 in which the Y functional group is an aryl nitrile to the benzylamine ester can be achieved by allowing the aryl nitriles
30 to react with dicobalt octacarbonyl in the presence of trimethylsilane. The reduction of a benzonitrile to a benzylamine is a known reaction (Murai, T.; Sakane, T.; Kato, S. *Tetrahedron Lett.* 26: 5145-5148[1985]) and is generally carried out in an inert solvent, such as toluene or the like, at 60°C for 20 hours with an 8 molar percent of dicobalt octacarbonyl and a 10 fold excess of trimethylsilane. The solvent is evaporated and the resulting material
35 diluted with methanol and allowed to react with a 5 fold excess of potassium fluoride. The products are then isolated by solvent extraction and further purified high-pressure liquid chromatography using a reverse phase column. Conversion of the same aryl nitrile compounds to their corresponding amidino esters can be accomplished by sequential reaction with

hydrogen sulphide, methyl iodide, and ammonium acetate. The resulting amidino esters may be purified by either column chromatography, crystallization, or preparative high-pressure liquid chromatography using a reverse phase column.

From commercially available precursors, alkynyl benzonitriles containing substituents on the nitrile containing benzo ring can be prepared by known methods. These substituents specifically include a chlorine (positioned either *ortho* or *meta* to the aryl nitrile), an amino (suitably protected, positioned *ortho* to the aryl nitrile), fluorine (tetra, *i.e.*, fluorines at both *ortho* and *meta* positions relative to the aryl nitrile), and a methyl group (positioned *ortho* and *meta* to the aryl nitrile). Palladium mediated coupling of these alkynyl benzonitriles containing the substituents described to 7, followed by treatment with hydrogen sulfide, methyl iodide, ammonium acetate and purification via silica, reverse phase chromatography, or recrystallization would afford amidino esters of interest. It is also possible to prepare from a commercially available material an amidino benzodiazepinedione ester where L' is a pyridine ring rather than a benzene ring. In the synthesis of the alkynyl pyridyl nitrile which would be reacted with 7, it may be preferable to use a nickel catalyst (rather than palladium) to effect the coupling of 2-chloro-5-cyano-pyridine with the acetylene synthon (most likely trimethylsilyl acetylene).

Conversion of the compounds of the general formula 48, 49, or 50 in which the Y functional group is a nitroarene, to the anilino ester can be executed by a selective reduction (Bellamy, F.D.; Ou, K. *Tetrahedron Lett.* 25: 839-842[1984]). The reaction is generally run with a five molar excess of stannous chloride dihydrate in either ethyl acetate or ethanol as the solvent at temperatures between about 50 and 100°C for times between about 15-120 minutes under an inert atmosphere, such as nitrogen. Preferably, the reaction is carried out at 70°C in ethanol for approximately 30 minutes. The products are then isolated by solvent extraction and further purified by high-pressure liquid chromatography using a reverse phase column.

Conversion of the amino esters to their corresponding amino acids of the structural Formula I and II involves saponification using well known conditions and reagents. For example, an aqueous solution of a strong alkali metal base, such as sodium hydroxide, lithium hydroxide or the like, is added to an alcohol-water (1:3) solution of the ester. Alcohols which may be used as the solvent for this reaction may include, for example, methanol, ethanol, and isopropanol, but, methanol is preferred. The preferred base is sodium hydroxide at a concentration between about 1 to 6N, though 2N is preferred. The reaction may be conducted at a temperature between about 0 to 50°C for times between about 10 to 60 minutes. Preferably, the reaction is carried out at room temperature for 30 minutes after which the reaction is neutralized with a concentrated solution of a strong acid, such as hydrochloric acid or the like, and the solvent evaporated. The products are isolated by high-pressure liquid chromatography using a reverse phase column.

The conversion of the amino acids to their corresponding guanidino acids of the structural Formula I and II is a known reaction (Kim, K. Lin, Y.-L.; Mosher, H.S. *Tetrahedron Lett.* 3183-3186 [1988]). The reaction can be accomplished by allowing the amino acid to react with aminoiminomethanesulfonic acid. Generally, the reaction can be conducted with a

5 equimolar to a 10 fold molar excess of aminoiminomethanesulfonic acid at temperatures between 0 to 50°C for times between about 15 to 120 minutes in a polar protic solvent, such as methanol, water or the like. The solution, prior to the addition of aminoiminomethanesulfonic acid, may be made neutral or basic by the addition of weak base, such as an alkali metal carbonate. Preferably, for alkyl amines the amino acid will be


10 allowed to react with a 5 fold excess aminoiminomethanesulfonic acid at room temperature for 30 minutes with 5% potassium bicarbonate in water as the reaction medium, whereas for aryl amines the amino acid will be allowed to react with an equimolar amount of aminoiminomethanesulfonic acid at room temperature for 1 hour in methanol. Generally, the reaction mixture will be made acidic by the addition of a dilute solution of an acid, such as


15 acetic acid and the solvent evaporated. The products are isolated by high-pressure liquid chromatography using a reverse phase column.

Preparation of other tricyclic compounds containing preferred L-Q groups bonded to the benzo moiety may also be prepared from key intermediate 7 according to Schemes XIII and XIV below.

20

Scheme XIII






Conversion of the iodoarene 7 to a synthetic precursors of 59 may be accomplished by carbonylation in an alcoholic solvent (R'OH) under a carbon monoxide atmosphere utilizing a palladium(0) catalyst. The alcoholic solvent (R'OH) must be chosen so as to allow for the selective removal of the R' group of the diester 51. For example, when R' = CH_3 and $\text{R}^{22} = \text{tert-butyl}$, the alcoholic solvent is methanol, and this methyl ester functionality in 51 can be removed by mild basic hydrolysis to afford acid 52.

For preparation of the biaryl adducts 53, 7 may be allowed to react with substituted iodoaryls or arylboronates, where Y and L' are defined as above, in the presence of palladium(0). Alternatively, synthetic precursors of 58 and 60 may be prepared by converting iodoarene 7 into the aniline 56 or phenol 57 by standard procedures. Phenol 57 may be converted to ether 58 by combining with the appropriately protected alcohol and dehydrating with diethyl azodicarboxylate-triphenylphosphene (DEAD).

Precursor carboxylates 52 and anilines 56 may be converted into amides 59 and 60 in a dehydration reaction utilizing dicyclohexylcarbodiimide (DCC) with the appropriate amine or acid according to Scheme XIV.

Scheme XIV



- 5 The latter reaction preferably is carried out with an acyl chloride in the presence of triethylamine and 4-dimethylaminopyridine (DMAP).

4. Isomeric Products

- 10 In products of Formula I and II carbon atoms bonded to four nonidentical substituents are asymmetric. Accordingly, the compounds may exist as diastereoisomers, enantiomers or mixtures thereof. The syntheses described above may employ racemates, enantiomers or diastereomers as starting materials or intermediates. Diastereomeric products resulting from such syntheses may be separated by chromatographic or crystallization methods. Likewise, enantiomeric product mixtures may be separated using the same techniques or by other
- 15 methods known in the art. Each of the asymmetric carbon atoms, when present in compounds of Formula I and II, may be in one of two configurations (R or S) and both are within the scope of the present invention.

E. Pharmaceutical Compositions

- 20 The compounds described in this invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of this invention. Examples of such salts include ammonium, metal salts like sodium, potassium, calcium and magnesium; salts with organic bases like dicyclohexylamine, N-methyl-D-glucamine and the like; and salts with amino acids like arginine or lysine. Salts
- 25 with inorganic and organic acids may be likewise prepared, for example, using hydrochloric, hydrobromic, sulfuric, phosphoric, trifluoroacetic, methanesulfonic, malic, maleic, fumaric and the like. Non-toxic and physiologically compatible salts are particularly useful although other less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. For example, reaction of the free acid or free base form of a compound of Formula I and II with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble; or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free acid or base form of the product may be passed over an ion exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

In the management of thromboembolic disorders the compounds of this invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration; suppositories for rectal administration; sterile solutions or suspensions for injectable administration, and the like. Animals in need of treatment using compounds of this invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from animal to animal and be dependent upon such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize.

Dosage formulations of the nonpeptidyl inhibitors of the present invention are prepared for storage or administration by mixing the inhibitor having the desired degree of purity with physiologically acceptable carriers, excipients, or stabilizers. Such materials are non-toxic to the recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, acetate and other organic acid salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) peptides such as polyarginine, proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium and/or nonionic surfactants such as Tween, Pluronic or polyethyleneglycol.

Dosage formulations of the nonpeptidyl inhibitors of the present invention to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes such as 0.2 micron membranes. Nonpeptidyl inhibitor formulations ordinarily will be stored in lyophilized form or as an aqueous solution. The pH of the cyclic inhibitor preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of cyclic polypeptide salts. While the preferred route of administration is by hypodermic injection needle, other methods of administration are also anticipated such as suppositories, aerosols,

oral dosage formulations and topical formulations such as ointments, drops and dermal patches.

Therapeutic nonpeptidyl inhibitor formulations generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by hypodermic injection needle.

Therapeutically effective dosages may be determined by either *in vitro* or *in vivo* methods. For each particular nonpeptidyl inhibitor of the present invention, individual determinations may be made to determine the optimal dosage required. The range of therapeutically effective dosages will naturally be influenced by the route of administration. For injection by hypodermic needle it may be assumed the dosage is delivered into the body's fluids. For other routes of administration, the absorption efficiency must be individually determined for each inhibitor by methods well known in pharmacology.

The range of therapeutic dosages is from about 0.001 nM to 1.0 mM, more preferably from 0.1 nM to 100 mM, and most preferably from 1.0 nM to 50 mM.

Typical formulation of compounds of Formula I and II as pharmaceutical compositions are discussed below.

About 0.5 to 500 mg of a compound or mixture of compounds of Formula I and II, as the free acid or base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., as called for by accepted pharmaceutical practice. The amount of active ingredient in these compositions is such that a suitable dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules and the like are a binder such as acacia, corn starch or gelatin; an excipient such as microcrystalline cellulose; a disintegrating agent like corn starch or alginic acid; a lubricant such as magnesium stearate; a sweetening agent such as sucrose or lactose; a flavoring agent such as peppermint, wintergreen or cherry. When the dosage form is a capsule, in addition to the above materials it may also contain a liquid carrier such as a fatty oil. Other materials of various types may be used as coatings or as modifiers of the physical form of the dosage unit. A syrup or elixir may contain the active compound, a sweetener such as sucrose, preservatives like propyl paraben, a coloring agent and a flavoring agent such as cherry. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example, dissolution or suspension of the active compound in a vehicle such as water or naturally occurring vegetable oil like sesame, peanut, or cottonseed oil or a synthetic fatty vehicle like ethyl oleate or the like may be desired. Buffers, preservatives, antioxidants and the like can be incorporated according to accepted pharmaceutical practice.

F. Platelet Inhibition Assays

Evaluation of the Formula I and II inhibitors of the fibrinogen-platelet interaction is guided by *in vitro* receptor binding assays and *in vitro* platelet aggregation inhibition assays.

In-vitro biological activity of the compounds of Formula I and II are monitored using
 5 a modified fibrinogen-GPIIb/IIIa ELISA based on the method of Nachman and Leung (*J. Clin. Invest.* 69:263-269 (1982)) which measures the inhibition of fibrinogen binding to purified human platelet GPIIb/IIIa receptor. Human fibrinogen is prepared by the method of Lipinska, *et al.* (*J. Lab. Clin. Med.* 84:509-516 (1974)). Platelet GPIIb/IIIa is prepared by the method of Fitzgerald, *et al.*, *Anal. Biochem.*, 151:169-177 (1985).


10 Briefly, microtiter plates are coated with fibrinogen (10 mg/ml) and then blocked with TACTS buffer containing 0.5% bovine serum albumin (BSA). (TACTS buffer contains 20mM Tris.HCl, pH 7.5, 0.02% sodium azide, 2 mM calcium chloride, 0.05% Tween 20, 150 mM sodium chloride.) The plate is washed with phosphate buffered saline (PBS) containing
 15 GP IIb/IIIa receptor (40 mg/ml) in TACTS, 0.5% BSA. After incubation, the plate is washed and 1 mg/ml of murine anti-platelet monoclonal antibody AP3 (Newman *et al.*, *Blood* 65:227-232 (1985)) is added. After another wash a goat anti-mouse IgG conjugated to horseradish peroxidase is added. A final wash is performed and developing reagent buffer (10 mg o-phenylenediamine dihydrochloride, 0.0212% hydrogen peroxide, 0.22 mM citrate, 50 mM
 20 phosphate, pH 5.0) is added and then incubated until color develops. The reaction is stopped with 1N sulfuric acid and the absorbance at 492 nm is recorded.

In addition to the GPIIb/IIIa ELISA assay, platelet aggregation assays may be performed in human platelet rich plasma (PRP). Fifty milliliters of whole human blood (9 parts) is drawn on 3.6% sodium citrate (1 part) from a donor who has not taken aspirin or
 25 related medications for at least two weeks. The blood is centrifuged at 160 x g for 10 min at 22° C and then allowed to stand for 5 min after which the PRP is decanted. Platelet poor plasma (PPP) is isolated from the remaining blood after centrifugation at 2000 x g for 25 min. The platelet count of the PRP is adjusted to ca. 300,000 per microliter with PPP.

A 225 mL aliquot of PRP plus 25 mL of either a dilution of the test sample or a control
 30 (PBS) is incubated for 5 min in a Chrono-log Whole Blood Aggregometer at 25° C. An aggregating agent (collagen, 1 mg/ml; U46619, 100 ng/ml; or ADP, 8 mM) is added and the platelet aggregation recorded.

Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and illustrative examples, make and utilize the present
 35 example to the fullest extent. The following working examples therefore, specifically point out preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

Example 1



5 **4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate**

a) A slurry of 1-(diphenylmethyl)-4-(2-carboxyethyl)-7-iodo-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester (15grams, 0.0264mol), (Blackburn, B.K., Barker, P.;
 10 Gadek, T; McDowell, R.S.; McGee, L.; Somers, T.; Webb, R.; Robarge, K. International Patent Application WO 93/08174), 15mL anisole, and 3mL ethyl methyl sulfide in a teflon tube was cooled to -196°C (N₂) and condensed 70mL hydrogen fluoride. The reaction was stirred for 2hrs and concentrated *in vacuo*. The resulting residue was purified on column chromatography (SiO₂, using 30% ethyl acetate in hexane to 80% ethyl acetate in hexane). The material can
 15 be further purified by crystalization using a minimum volume of ethyl acetate (heating was required to dissolve the material) followed by the addition of an equal volume of hexanes to yield 8.5 grams (80%) of m.p. 129-130°C 4-(2-carboxyethyl)-7-iodo-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester. ¹H NMR (CDCl₃, δTMS) 9.60 (1H, s, NH), 8.21 (1H, d, ⁴J_{HH}= 2Hz, C6 Ar-H), 7.72 (1H, dd, ⁴J_{HH}=2Hz, ³J_{HH}=9Hz, C8 Ar-H), 6.81 (1H, d, ³J_{HH}=9Hz, C9 ArH), 4.12 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.97 (2H, s, NCH₂CO), 3.91 (2H, t, ³J_{HH}=7Hz, NCH₂CH₂CO₂), 2.72 (2H, t, ³J_{HH}=7Hz, CH₂CO₂), 1.23 (3H, t, ³J_{HH}=7Hz, CH₃). ¹³C NMR (CDCl₃) 171.3, 170.7, 165.7, 141.1, 140.1, 135.6, 127.9, 122.4, 88.5, 60.8, 51.5, 45.7, 32.8, 14.1.

Using the above procedure, but substituting the appropriate 1-(diphenylmethyl)-4-(2-carboxyethyl)-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester for 1-(diphenylmethyl)-4-(2-carboxyethyl)-7-iodo-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester there may be prepared, for example, the following compounds:
 4-(2-carboxyethyl)-7-[3-(4-piperidine)-propoxy]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester,
 25 4-(2-carboxyethyl)-7-[4-(1-piperazine)-phenyl]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester,
 30

- 4-(2-carboxyethyl)-7-[3-(1-piperazine)-propoxy]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester,
- 4-(2-carboxyethyl)-7-[4-(4-piperidine)-2-oxobutyl]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester,
- 5 4-(2-carboxyethyl)-7-[2-(4-cyanophenyl)-2-oxoethyl]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester,
- 4-(2-carboxyethyl)-7-[(4-cyanoanilino)-carbonyl]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester,
- 4-(2-carboxyethyl)-7-[[4-(cyano)-phenylsulphonyl]-amino]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester,
- 10 7-[[4-(cyano)benzoyl]amino]-1H-[1,4]benzodiazepine-2,5-dione-4-propanoic acid ethyl ester.

- b) To a solution of 4-(2-carboxyethyl)-7-iodo-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester (1.0 grams, 2.49mmol), in 10mL THF at room temperature under an atmosphere of nitrogen was added 1.0 equiv. of Lawesson's reagent (1.0 grams) and the reaction heated to 50°C for 2hrs. The reaction mixture was allowed to cool to room temperature and concentrated *in vacuo*. The resulting residue was purified on column chromatography (SiO₂, using 40% ethyl acetate in hexanes to 60% ethyl acetate in hexanes, TLC (1:1, ethyl acetate/hexane): R_f=0.44, μ v positive) to yield 0.95 grams (91%) of 4-(2-carboxyethyl)-7-iodo-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester (m.p. 187-189°C). ¹H NMR (CDCl₃, δ TMS) 8.21 (1H, d, ⁴J_{HH}= 2Hz, C6 Ar-H), 7.78 (1H, dd, ⁴J_{HH}=2Hz, ³J_{HH}=9Hz, C8 Ar-H), 6.88 (1H, d, ³J_{HH}=9Hz, C9 Ar-H), 4.28 (2H, s, NCH₂CS), 4.15 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.94 (2H, t, ³J_{HH}=7Hz, NCH₂CH₂CO₂), 2.72 (2H, t, ³J_{HH}=7Hz, CH₂CO₂), 1.27 (3H, t, ³J_{HH}=7Hz, CH₃). ¹³C NMR (CDCl₃) 198.8, 171.4, 165.3, 141.2, 140.4, 135.9, 128.9, 121.9, 90.4, 60.8, 58.0, 45.4, 32.8, 14.1.
- 25

- Using the above procedure, but substituting the appropriate 4-(2-carboxyethyl)-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester for 4-(2-carboxyethyl)-7-iodo-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester there may be prepared, for example, the following compounds:
- 30

- 4-(2-carboxyethyl)-7-[3-(1-(N-Boc)-4-piperidine)-propoxy]-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester,
- 4-(2-carboxyethyl)-7-[4-(4-(N-Boc)-1-piperazine)-phenyl]-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester,
- 35 4-(2-carboxyethyl)-7-[3-(4-(N-Boc)-1-piperazine)-propoxy]-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester,
- 4-(2-carboxyethyl)-7-[4-(1-(N-Boc)-4-piperidine)-2-oxobutyl]-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester,

- 4-(2-carboxyethyl)-7-[2-(4-cyanophenyl)-2-oxoethyl]-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester,
 4-(2-carboxyethyl)-7-[(4-cyanoanilino)-carbonyl]-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester,
 5 4-(2-carboxyethyl)-7-[[4-(cyano)-phenylsulphonyl]-amino]-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester,
 7-[[4-(cyano)benzoyl]amino]-1H-[1,4]benzodiazepine-2-thione-5-one-4-propanoic acid ethyl ester.

- 10 c) To a biphasic solution of the 4-(2-carboxyethyl)-7-iodo-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester (0.95 grams, 2.27mmol), CH₂Cl₂ (10mL), water (10mL), methyl iodide (0.2 grams) and a catalytic amount of tetrabutylammonium hydrogen sulfate at room temperature with vigorous stirring was added 1.2mL 2N sodium hydroxide. The reaction was monitored by TLC. After 2hrs, the layers were separated and the aqueous
 15 layer was washed 2X25mL CH₂Cl₂. The combined organics were dried over sodium sulfate, decanted and concentrated in vacuo. The resulting residue was dissolved in 10mL toluene and allowed to react with propargyl amine (4-fold excess, 0.64mL) and pyridine hydrogen chloride (1 molar equiv., 0.23 grams). The reaction was heated to reflux and the reaction monitored by TLC. After 6 hrs, the reaction mixture was allowed to cool to room temperature,
 20 concentrated in vacuo, and purified by column chromatography (SiO₂, 100% ethyl acetate), to yield 0.653 grams (66% yield) of 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-iodo-1-methyl-6-oxo, ethyl ester. ¹H NMR (CDCl₃, δTMS) 8.33 (1H, d, ⁴J_{HH}=2Hz, C7 Ar-H), 7.90 (1H, dd, ⁴J_{HH}=2Hz, ³J_{HH}=9Hz, C9 Ar-H), 7.00 (1H, d, ³J_{HH}=9Hz, C10 ArH), 6.89 (1H, bq, ⁴J_{HH}=1Hz), 4.49 (1H, d, ²J_{HH}=15Hz, NCHHCN), 4.32 (1H, d, ²J_{HH}=15Hz, NCHHCN) 4.17 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.94 (2H, m, NCH₂CH₂CO₂), 2.69 (2H, m, CH₂CO₂), 2.33 (3H, d, ⁴J_{HH}=1Hz, CH₃), 1.27 (3H, t, ³J_{HH}=7Hz, CH₃). ¹³C NMR (CDCl₃) 171.3, 145.9, 140.7, 140.5, 140.4, 132.8, 128.1, 124.6, 91.9, 60.8, 45.7, 45.1, 32.8, 14.1, 11.0.

- Using the above procedure, but substituting the appropriate 4-(2-carboxyethyl)-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester for 4-(2-carboxyethyl)-7-iodo-3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one ethyl ester there may be prepared, for example, the following compounds:

- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[1-(N-Boc)-4-piperidine]-propoxy]-1-methyl-6-oxo, ethyl ester,
 35 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[4-(N-Boc)-1-piperazine]-phenyl]-1-methyl-6-oxo, ethyl ester,
 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[4-(N-Boc)-1-piperazine]-propoxy]-1-methyl-6-oxo, ethyl ester,

- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[1-(N-Boc)-4-piperidine]-2-oxobutyl]-1-methyl-6-oxo, ethyl ester,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[2-(4-cyanophenyl)-2-oxoethyl]-1-methyl-6-oxo, ethyl ester,
- 5 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[(4-cyanoanilino)-carbonyl]-1-methyl-6-oxo, ethyl ester,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(cyano)phenylsulphonyl]-amino]-1-methyl-6-oxo, ethyl ester,
- 10 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(cyano)benzoyl]amino]-1-methyl-6-oxo, ethyl ester.

d) To a slurry of the 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-iodo-1-methyl-6-oxo, ethyl ester (0.15 grams, 0.34mmol), 4mL ethyl acetate (degassed), 4-cyanophenylacetylene (2 molar equiv. 0.135 grams),

15 bis(triphenylphosphine)palladium dichloride (10mgs), copper iodide (10mgs), at room temperature under an atmosphere of nitrogen was added triethylamine (0.24 grams), and the reaction allowed to stir overnight. The reaction was concentrated in vacuo and the residue purified by column chromatography (SiO₂, using 100% ethyl acetate) to yield 0.157 grams (78%) of 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-

20 cyanophenyl]ethynyl]-1-methyl-6-oxo, ethyl ester. ¹H NMR (CDCl₃, δTMS) 8.01 (1H, d, ⁴J_{HH}=2Hz, C7 Ar-H), 7.92 (2H, d, ³J_{HH}=8Hz, *o*-CN ArH), 7.84 (1H, dd, ⁴J_{HH}=2Hz, ³J_{HH}=9Hz, C9 Ar-H), 7.78 (2H, d, ³J_{HH}=8Hz, *m*-CN ArH), 7.61 (1H, d, ³J_{HH}=9Hz, C10 ArH), 6.82 (1H, bq, ⁴J_{HH}=1Hz), 4.49 (1H, d, ²J_{HH}=15Hz, NCHHCN), 4.35 (1H, d, ²J_{HH}=15Hz, NCHHCN) 4.00 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.77 (2H, bt, NCH₂CH₂CO₂), 2.58

25 (2H, m, CH₂CO₂), 2.30 (3H, d, ⁴J_{HH}=1Hz, CH₃), 1.13 (3H, t, ³J_{HH}=7Hz, CH₃). ¹³C NMR (CDCl₃) 171.3, 166.1, 135.5, 134.2, 133.1, 132.1, 128.1, 127.3, 121.6, 118.3, 111.9, 91.4 89.5, 60.7, 45.7, 45.1, 32.7, 14.1, 11.1.

e) To a solution of 0.15 grams of 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanophenyl]ethynyl]-1-methyl-6-oxo, ethyl ester and pyridine (2mL)

30 and diethylamine (2mL) was bubbled via a pipette for 5 mins. hydrogen sulfide (gas). The reaction was stirred for 2hrs at room temperature, concentrated *in vacuo*, and purified by column chromatography (SiO₂, 100% ethyl acetate). The yellow compound was further allowed to react with methyl iodide (2mL) in methylene chloride (4mL), heating to 50°C for

35 30 mins. The mixture was concentrated *in vacuo* and the resulting residue allowed to react with ammonium acetate (0.5 grams) in ethanol (5mL). The mixture was again heated to 50°C for 30 mins. After cooling to room temperature the reaction was purified by HPLC (O-10 mins. (30% acetonitrile(0.5% acetic acid)/70% water(0.5% acetic acid) 10-50 mins. ramp to 70%

acetonitrile (0.5% acetic acid)/30% water (0.5% acetic acid), 15mL/min., Detector: 254nm, R_f =27.6 mins.) to yield 0.025 grams of 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, ethyl ester acetate (16%). MS (FAB) cald. for $C_{26}H_{25}N_5O_3$ 455.1, found MH^+ 456.1.

5

Using the above procedure, but substituting the appropriate 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 1-methyl-6-oxo, ethyl ester for 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, ethyl ester there may be prepared, for example, the following compounds:


- 10 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[2-[4-(aminoiminomethyl)]-2-oxoethyl]-1-methyl-6-oxo, ethyl ester,
 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)-anilino]-carbonyl]-1-methyl-6-oxo, ethyl ester,
 15 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenylsulphonyl]-amino]-1-methyl-6-oxo, ethyl ester,
 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)benzoyl]-amino]-1-methyl-6-oxo, ethyl ester.

- f) 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, ethyl ester acetate taken from part (e) above was saponified in 2mL methanol with 2mL 2N sodium hydroxide for 20 mins. The reaction mixture was quenched with acetic acid and purified by HPLC (O-10 mins. (30% acetonitrile(0.5% acetic acid)/70% water(0.5% acetic acid) 10-50 mins. ramp to 70% acetonitrile (0.5% acetic acid)/30% water (0.5% acetic acid), Detector: 254nm, R_f =23.5 mins.)
 20 to yield 20mgs (85%) of 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate. Exact mass (FAB, MH^+) cald for $C_{24}H_{25}N_5O_3$: 428.1723, found 428.1705.

- Using the above procedure, but substituting the appropriate 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 1-methyl-6-oxo, ethyl ester for 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, ethyl ester there may be prepared, for example, the following compounds:
 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[2-[4-(aminoiminomethyl)]-2-oxoethyl]-1-methyl-6-oxo, acetate,
 35 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)anilino]-carbonyl]-1-methyl-6-oxo, acetate,
 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenylsulphonyl]-amino]-1-methyl-6-oxo, acetate

4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)benzoyl]amino]-1-methyl-6-oxo, acetate.

Example 2



4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[6-aminohex-1-ynyl]-1-methyl-6-oxo, acetate

10

a) To a slurry 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-iodo-1-methyl-6-oxo, ethyl ester, prepared as described in part (c) of example 1, (0.15 grams, 0.34 mmol), 4mL ethyl acetate (degassed), N-Boc-6-aminohexyne (0.135 grams, 0.68 mmol), bistrisphenylphosphine palladium dichloride (10 mgs), copper iodide (10 mgs), at room temperature under an atmosphere of nitrogen was added triethylamine (0.16mL, 5X excess) and the reaction allowed to stir overnight. The reaction was concentrated *in vacuo* and the residue chromatographed (60mL SiO₂, ethyl acetate) to yield the desired product, 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[N-Boc]6-aminohex-1-ynyl]-1-methyl-6-oxo, ethyl ester (0.113 grams, 65%). ¹H NMR (CDCl₃, δTMS) 7.84 (1H, d, ⁴J_{HH}=2Hz, C7 Ar-H), 7.40 (1H, dd, ⁴J_{HH}=2Hz, ³J_{HH}=9Hz, C8 Ar-H), 7.04 (1H, d, ³J_{HH}=9Hz, C9 Ar-H), 6.64 (1H, s, C2-H), 4.45 (1H, bq, NH), 4.32 (1H, d, ²J_{HH}=16Hz, C4H), 4.13 (1H, d, ²J_{HH}=16Hz, C4H), 3.97 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.76 (2H, m, NCH₂CH₂), 3.03 (2H, bq, NHCH₂), 2.51 (2H, m, CH₂CO₂), 2.3 (2H, t, ³J_{HH}=7Hz, C≡CCH₂), 2.17 (3H, s, Me), 1.49 (4H, m, CH₂CH₂), 1.28 (9H, s, t-Bu), 1.08 (3H, t, ³J_{HH}=7Hz, CH₂CH₃); ¹³C NMR (CDCl₃) 171.3, 166.4, 145.9, 135.1, 134.2, 131.9, 130.3, 127.9, 123.5, 123.0, 92.2, 79.2, 60.7, 45.6, 44.9, 32.8, 29.3, 28.4, 25.7, 19.0, 14.1, 11.1; MS (FAB) cald. for C₂₈H₃₆N₄O₅ 508.3, found MH⁺ 509.3.

25

b) A solution of 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[N-Boc]6-aminohex-1-ynyl]-1-methyl-6-oxo, ethyl ester (0.113 grams, 0.22 mmol) in 2mL ethyl acetate and 0.5mL triethylsilane was added 2mL of a saturated solution of hydrogen

30


chloride in ethyl acetate and the reaction monitored by TLC. After 30 mins., the reaction was concentrated *in vacuo*, diluted with 2mL ethanol and allowed to react with 0.5mL 2N sodium hydroxide. After 15 mins. the reaction was quenched with 0.5mL glacial acetic acid and purified by HPLC (C18, 0-10 mins. (30% acetonitrile(0.5% acetic acid)/70% water(0.5% acetic acid) 10-50 mins. ramp to 70% acetonitrile (0.5% acetic acid)/30% water (0.5% acetic acid), Detector: 254nm) to yield 44.2mgs (52%) of 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[6-aminohept-1-ynyl]-1-methyl-6-oxo, acetate. ¹H NMR (D₂O, δHOD) 7.63 (1H, d, ⁴J_{HH}=2Hz, C7 Ar-H), 7.42 (1H, dd, ⁴J_{HH}=2Hz, ³J_{HH}=9Hz, C8 Ar-H), 7.29 (1H, d, ³J_{HH}=9Hz, C9 Ar-H), 6.98 (1H, bs, C2H), 4.50 (1H, bd, ²J_{HH}=16Hz, C4H), 4.24 (1H, bd, ²J_{HH}=16Hz, C4H), 3.64 (2H, bs, NCH₂CH₂), 2.92 (2H, t, ³J_{HH}=7Hz, NH₃CH₂), 2.41 (2H, m, CH₂CO₂), 2.2 (5H, m, C≡CCH₂, Me), 1.63 (2H, m, NH₃CH₂CH₂CH₂), 1.42 (2H, m, NH₃CH₂CH₂CH₂). Exact mass (FAB, MH⁺) calcd for C₂₁H₂₄N₄O₃: 381.1927, found 381.1943.

15 Using the above procedure, but substituting the appropriate 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 1-methyl-6-oxo, ethyl ester for 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[N-Boc]6-aminohept-1-ynyl]-1-methyl-6-oxo, ethyl ester there may be prepared, for example, the following compounds:

- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-(4-piperidine)-propoxy]-1-methyl-6-oxo, acetate,
- 20 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-(1-piperazine)-phenyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-(1-piperazine)-propoxy]-1-methyl-6-oxo, acetate,
- 25 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-(4-piperidine)-2-oxobutyl]-1-methyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[4-piperidine]-propoxy]-1-methyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[1-piperazine]-phenyl]-1-methyl-6-oxo, acetate,
- 30 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[1-piperazine]-propoxy]-1-methyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[4-piperidine]-2-oxobutyl]-1-methyl-6-oxo, acetate,
- 35 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[4-piperidine]-propoxy]-1-hydroxy-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[1-piperazine]-phenyl]-1-hydroxy-6-oxo, acetate,

- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[1-piperazine]-propoxy]-1-hydroxy-6-oxo, acetate,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[4-piperidine]-2-oxobutyl]-1-hydroxy-6-oxo, acetate,
 5 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[4-piperidine]-propoxy]-6-oxo, acetate,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[1-piperazine]-phenyl]-16-oxo, acetate,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[1-piperazine]-propoxy]-6-oxo, acetate,
 10 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[4-piperidine]-2-oxobutyl]-6-oxo, acetate,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[3-[4-piperidine]-propoxy]-7,13-dihydro-5,13-dioxo, acetate,
 15 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[4-[piperazine]-phenyl]-7,13-dihydro-5,13-dioxo, acetate,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[3-[1-piperazine]-propoxy]-7,13-dihydro-5,13-dioxo, acetate,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[4-[4-piperidine]-2-oxobutyl]-7,13-dihydro-5,13-dioxo, acetate.
 20

Example 3



25
 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methyl-6-oxo, acetate

- 30 a) A solution of 1.44 g (7.15 mmol, 1.1 eq.) bromoacetyl bromide in 10 mL of methylene chloride was added over 10 mins. to a stirred biphasic solution of 2.38 g (6.5 mmol,

1.0 eq.) of N-(5-(4-cyanobenzyloxy)-2-aminobenzoyl)- β -alanine ethyl ester in 100 ml of methylene chloride and 30 mL of water cooled to 0°C. The reaction was complete in 1/2 hr. (TLC, 96 CH₂Cl₂:4 MeOH, product R_f=0.71) and was partitioned between dilute aqueous sodium bicarbonate and methylene chloride. The organic phase was dried over sodium sulfate, filtered and concentrated to give 3.11 g (98%) of off white crystals of N-(5-(4-cyanobenzyloxy)-2-(bromoacetyl)aminobenzoyl)- β -alanine ethyl ester which was used unpurified but may be recrystallized from ethyl acetate/hexane. ¹H NMR (CDCl₃, δ TMS) 8.41(1H,d,ArH), 7.60 (4H,abq,ArH), 7.02(2H,d,ArH), 6.98(1H,t,NH), 5.12 (2H,s,ArCH₂O), 4.16 (2H,q,COOCH₂), 3.97 (2H,s,CH₂Br), 3.68 (2H,q,NCH₂), 2.61 (2H,t,CH₂COO), 1.25 (3H,t,CH₃). IR (KBr): 3389(NH), 3355(NH), 2226(CN), 1735(ester), 1675,1642(amides),1609,1523,1423,1244,1184,819. MS (FAB,M+H⁺) calcd for C₂₂H₂₂N₃O₅Br 488.3, found (M+H⁺) 489.

b) A mixture of 3.11g (6.37 mmol, 1.0 eq.) of N-(5-(4-cyanobenzyloxy)-2-(bromoacetyl)aminobenzoyl)- β -alanine ethyl ester, 1.32 g (9.55 mmol, 1.5 eq.) powdered potassium carbonate and 350 mL of dry DMF were combined and stirred at 50°C for 2 hrs. TLC (70 EtOAc:30 hexane, product R_f=0.31) indicated complete reaction. The reaction mixture was partitioned between ethyl acetate and water, the organic phase washed twice with water, once with brine, dried over sodium sulfate and evaporated to yield the crystalline product. The crystals were slurried in ether/ethyl acetate and filtered to give 1.5g of 4-(2-carboxyethyl)-7-(4-cyanobenzyloxy)-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester. The residual ether/ethyl acetate solution was concentrated and chromatographed on silica to recover an additional 0.5 g of product. Total yield=2.0 g (77%). ¹H NMR (300 MHz, acetone-d₆): 9.32 (1H,s,NH), 7.78 (4H,abq,ArH), 7.41 (1H,d,Ar), 7.20 (1H,d,Ar), 7.15 (1H,d,Ar), 5.24 (2H,s,ArCH₂O), 4.10 (2H,q,COOCH₂), 3.98 (2H,s,COCH₂N), 3.88 (2H,t,NCH₂), 3.77 (HOD), 2.66 (2H,t,CH₂COO), 2.03(acetone), 1.20(3H,t,CH₃). IR(KBr): 3521(NH), 2226(CN), 1735(ester), 1682, 1642, 1609, 1503, 1277, 1204, 819. Exact mass: (FAB,M+H⁺) calcd for C₂₂H₂₂N₃O₅: 408.1559, found: 408.1538. A 19% yield of dimer was observed (TLC, 70 EtOAc: 30 hexane, R_f=0.23).

30

c) To a solution of 7-[[4-cyanophenyl]methoxy]-4-propanoic acid, 3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione, ethyl ester (1.0 grams, 2.45 mmol), in THF (20mL) at room temperature under an atmosphere of nitrogen was added 1.0 molar equivalent of Lawesson's reagent (1.0 grams) and the mixture heated to 50°C for 2 hours. The mixture was allowed to cool to room temperature, concentrated *in vacuo*, and purified by column chromatography (SiO₂, 4:6 ethyl acetate:hexanes to 6:4 ethyl acetate:hexanes) to yield 1.0 grams (96%) of 7-[[4-cyanophenyl]methoxy]-4-propanoic acid, 3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one, ethyl ester. ¹H NMR (CDCl₃, δ TMS) 9.65 (1H, s, NH), 7.71 (2H, d, ³J_{HH}=8Hz,

35


3,5H:4-NCC₆H₄), 7.55 (2H, ³J_{HH}=8Hz, 2,6H:4-NCC₆H₄), 7.51 (1H, d, ⁴J_{HH}=3Hz, C6 ArH), 7.13 (1H, dd, ⁴J_{HH}=3Hz, ³J_{HH}=9Hz, C8 ArH), 7.01 (1H, d, ³J_{HH}=9Hz, C9 ArH), 5.17 (2H, s, OCH₂Ar), 4.29 (2H, s, C(S)CH₂), 4.16 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.96 (2H, t, ³J_{HH}=7Hz, NCH₂), 2.79 (2H, t, ³J_{HH}=7Hz, CH₂CO₂), 1.27 (3H, t, ³J_{HH}=7Hz, CH₂CH₃); MS (FAB) cald. for C₂₂H₂₁N₃O₄S 423.1, found MH⁺ 424.1.

d) To a biphasic solution of 7-[[4-cyanophenyl]methoxy]-4-propanoic acid, 3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one, ethyl ester (1 gram, 2.36 mmol), methylene chloride (10mL), water (10mL), methyl iodide (0.2mL) and a catalytic amount of tetrabutyl ammonium hydrogen sulphate at room temperature with vigorous stirring was added 1.2mL of a 2N solution of sodium hydroxide. After 16 hours (overnight) the layers were separated, the aqueous layer was washed 2X60mL methylene chloride, and the combined organics were dried over sodium sulfate, decanted, and concentrated *in vacuo*. The resulting residue, 7-[[4-cyanophenyl]methoxy]-4-propanoic acid, 3,4-dihydro-1,4-benzodiazepine-2-methylmercaptyl-5-one, ethyl ester (¹H NMR (CDCl₃, δTMS) 7.70 (2H, d, ³J_{HH}=8Hz, 3,5H:4-NCC₆H₄), 7.56 (2H, ³J_{HH}=8Hz, 2,6H:4-NCC₆H₄), 7.49 (1H, d, ⁴J_{HH}=3Hz, C6 ArH), 7.18 (1H, d, ³J_{HH}=9Hz, C9 ArH), 7.15 (1H, dd, ⁴J_{HH}=3Hz, ³J_{HH}=9Hz, C8 ArH), 5.18 (2H, s, OCH₂Ar), 4.16 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.94 (2H, t, ³J_{HH}=7Hz, NCH₂), 3.91 (2H, s, C(S)CH₂), 2.78 (2H, t, ³J_{HH}=7Hz, CH₂CO₂), 2.52 (3H, s, SMe), 1.24 (3H, t, ³J_{HH}=7Hz, CH₂CH₃); MS (FAB) cald. for C₂₃H₂₃N₃O₄S 437.1, found MH⁺ 438.1), was dissolved in dry toluene (15mL) and allowed to react with 0.23 grams pyridinium hydrogen chloride and 0.64 mL propargyl amine. The reaction was heated to reflux. After 5 hours, the reaction was allowed to cool to room temperature and the reaction concentrated *in vacuo*. The crude material was purified by column chromatography (SiO₂, using ethyl acetate as the eluting solvent, R_f=0.1) to yield 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-cyanophenyl]methoxy]-1-methyl-6-oxo, ethyl ester (0.23 grams, 22%). ¹H NMR (CDCl₃, δTMS) 7.71 (2H, d, ³J_{HH}=8Hz, 3,5H:4-NCC₆H₄), 7.56 (2H, ³J_{HH}=8Hz, 2,6H:4-NCC₆H₄), 7.53 (1H, d, ⁴J_{HH}=3Hz, C6 ArH), 7.24 (1H, d, ³J_{HH}=9Hz, C9 ArH), 7.18 (1H, dd, ⁴J_{HH}=3Hz, ³J_{HH}=9Hz, C8 ArH), 5.21 (2H, dd, OCH₂Ar), 4.52 (1H, d, ²J_{HH}=16Hz, C5H), 4.30 (1H, d, ²J_{HH}=16Hz, C5H), 4.16 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.89 (2H, m, NCH₂), 2.66 (2H, m, CH₂CO₂), 2.29 (3H, s, C1-Me), 1.22 (3H, t, ³J_{HH}=7Hz, CH₂CH₃).

e) To a solution of 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-cyanophenyl]methoxy]-1-methyl-6-oxo, ethyl ester (0.1 grams, 0.22 mmol), in 2mL pyridine and 2mL triethylamine was added hydrogen sulfide gas *via* a pipette to saturation and the reaction stirred for 20 mins. A stream of nitrogen was passed over the reaction mixture for 20 mins. The mixture was concentrated *in vacuo*, diluted with methylene chloride (2mL) and allowed to react with methyl iodide (2mL). The mixture was heated

briefly to 50°C, stirred for 30 mins., concentrated *in vacuo*, diluted with methanol and the reaction allowed to react with ammonium acetate (0.25 grams). Again the reaction was heated briefly to 50°C and stirred for an additional 30 mins. The mixture was concentrated *in vacuo* and purified by HPLC (solvent gradient 1:4 acetonitrile:water(0.5% acetic acid) (time 0 to 10 mins.) to 3:2 acetonitrile:water(0.5% acetic acid) (time 50). The major component isolated by HPLC was diluted in methanol (2mL) and allowed to react 2N sodium hydroxide (2mL). After 15 mins. the mixture was quenched with glacial acetic acid (0.5mL) and purified by HPLC (same eluting conditions as those described above) to yield 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methyl-6-oxo, acetate (0.005 grams, 5%). ¹H NMR (D₂O, δHOD) 7.63 (2H, d, ³J_{HH}=8Hz, 3,5H:4-NCC₆H₄), 7.53 (2H, ³J_{HH}=8Hz, 2,6H:4-NCC₆H₄), 7.44 (1H, d, ³J_{HH}=9Hz, C6 ArH), 7.36 (1H, d, ⁴J_{HH}=3Hz, C9 ArH), 7.26 (1H, dd, ⁴J_{HH}=3Hz, ³J_{HH}=9Hz, C8 ArH), 7.17 (1H, bq, C2H), 5.21 (2H, dd, OCH₂Ar), 3.78 (1H, d, ²J_{HH}=16Hz, C4H), 3.79 (1H, dt, NCHH), 3.63 (1H, dt, NCHH), 2.54 (2H, m, CH₂CO₂), 2.22 (3H, s, C1-Me). Exact mass (FAB, MH⁺) calcd for C₂₃H₂₄N₅O₄: 434.1828, found 434.1794.

Example 4



4H-[1,2,4]Triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methyl-6-oxo, acetate

a) To a biphasic solution of 7-[[4-cyanophenyl]methoxy]-4-propanoic acid, 3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one, ethyl ester (0.75 grams, 1.77 mmol), prepared as described in part (c) of Example 3, methylene chloride (20mL), water (20mL), 0.2mL methyl iodide, and a catalytic amount of tetrabutyl ammonium hydrogen sulfate was added 1.2mL 2N sodium hydroxide. After 16 hrs, overnight, of vigorous stirring the layers were separated. The aqueous layer was washed 3X30mL methylene chloride and the combined organics were dried over sodium sulfate, decanted, and concentrated *in vacuo*. the resulting

residue was dissolved in toluene (15mL) and allowed to react with 0.23 grams (2.0 mmol) pyridinium hydrogen chloride and 0.64 grams (8.6 mmol) acetic hydrazide and the reaction was heated to reflux. After 5 hours, the reaction was allowed to cool to room temperature and concentrated *in vacuo*. The resulting residue was purified by column chromatography (SiO₂, using 5:95 methanol:methylene chloride as the eluting solvent) to yield 0.5 grams (63%) of 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanophenyl]methoxy]-1-methyl-6-oxo, ethyl ester. ¹H NMR (CDCl₃, δTMS) 7.59 (2H, d, ³J_{HH}=8Hz, 3,5H:4-NCC₆H₄), 7.46 (2H, ³J_{HH}=8Hz, 2,6H:4-NCC₆H₄), 7.45 (1H, d, ⁴J_{HH}=3Hz, C6 ArH), 7.17 (1H, d, ³J_{HH}=9Hz, C9 ArH), 7.12 (1H, dd, ⁴J_{HH}=3Hz, ³J_{HH}=9Hz, C8 ArH), 5.12 (2H, dd, OCH₂Ar), 4.61 (1H, d, ²J_{HH}=16Hz, C5H), 4.25 (1H, d, ²J_{HH}=16Hz, C5H), 4.01 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.77 (2H, m, NCH₂), 2.57 (2H, m, CH₂CO₂), 2.46 (3H, s, C1-Me), 1.13 (3H, t, ³J_{HH}=7Hz, CH₂CH₃); ¹³C NMR (CDCl₃) 171.1, 165.9, 157.6, 152.4, 150.5, 141.0, 132.4, 131.2, 127.5, 124.1, 119.4, 118.3, 116.8, 111.9, 69.3, 60.6, 45.2, 43.3, 32.4, 14.0, 12.1.

15

Using the above procedure, but substituting the appropriate 4-propanoic acid, 3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one, ethyl ester for 7-[[4-cyanophenyl]methoxy]-4-propanoic acid, 3,4-dihydro-1H-1,4-benzodiazepine-2-thione-5-one, ethyl ester and hydrazide or azide for acetic hydrazide there may be prepared, for example, the following compounds:

20

4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[1-(N-Boc)-4-piperidine]-propoxy]-1-methyl-6-oxo, ethyl ester,

4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[4-(N-Boc)-1-piperazine]-phenyl]-1-methyl-6-oxo, ethyl ester,

25

4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[4-(N-Boc)-1-piperazine]-propoxy]-1-methyl-6-oxo, ethyl ester,

4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[1-(N-Boc)-4-piperidine]-2-oxobutyl]-1-methyl-6-oxo, ethyl ester,

30

4H-[1,2,4]triazolo[4,3-a][1,4]benzo-diazepine-5(6H)-propanoic acid, 8-[2-(4-cyanophenyl)-2-oxoethyl]-1-methyl-6-oxo, ethyl ester,

4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[(4-cyanoanalino)-carbonyl]-1-methyl-6-oxo, ethyl ester,

4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(cyano)-phenylsulphonyl]-amino]-1-methyl-6-oxo, ethyl ester,

35

4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[1-(N-Boc)-4-piperidine]-propoxy]-1-hydroxy-6-oxo, ethyl ester,

4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[4-(N-Boc)-1-piperazine]-phenyl]-1-hydroxy-6-oxo, ethyl ester,

- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[4-(N-Boc)-1-piperazine]-propoxy]-1-hydroxy-6-oxo, ethyl ester,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[1-(N-Boc)-4-piperidine]-2-oxobutyl]-1-hydroxy-6-oxo, ethyl ester,
 5 4H-[1,2,4]triazolo[4,3-a][1,4]benzo-diazepine-5(6H)-propanoic acid, 8-[2-(4-cyanophenyl)-2-oxoethyl]-1-hydroxy-6-oxo, ethyl ester,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[(4-cyanoanalino)-carbonyl]-1-hydroxy-6-oxo, ethyl ester,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(cyano)-
 10 phenylsulphonyl]-amino]-1-hydroxy-6-oxo, ethyl ester,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(cyano)benzoyl]amino]-1-hydroxy-6-oxo, ethyl ester,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[1-(N-Boc)-4-piperidine]-propoxy]-6-oxo, ethyl ester,
 15 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[4-(N-Boc)-1-piperazine]-phenyl]-6-oxo, ethyl ester,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[3-[4-(N-Boc)-1-piperazine]-propoxy]-1-methyl-6-oxo, ethyl ester,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[4-[1-(N-Boc)-4-piperidine]-2-oxobutyl]-1-methyl-6-oxo, ethyl ester,
 20 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzo-diazepine-5(6H)-propanoic acid, 8-[2-(4-cyanophenyl)-2-oxoethyl]-6-oxo, ethyl ester,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[(4-cyanoanalino)-carbonyl]-6-oxo, ethyl ester,
 25 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(cyano)-phenylsulphonyl]-amino]-6-oxo, ethyl ester,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(cyano)benzoyl]-amino]-6-oxo, ethyl ester,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[3-[1-(N-Boc)-4-piperidine]-propoxy]-7,13-dihydro-5,13-dioxo, ethyl ester,
 30 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[4-[4-(N-Boc)-1-piperazine]-phenyl]-7,13-dihydro-5,13-dioxo, ethyl ester,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[3-[4-(N-Boc)-1-piperazine]-propoxy]-7,13-dihydro-5,13-dioxo, ethyl ester,
 35 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[4-[1-(N-Boc)-4-piperidine]-2-oxobutyl]-7,13-dihydro-5,13-dioxo, ethyl ester,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[2-(4-cyanophenyl)-2-oxoethyl]-7,13-dihydro-5,13-dioxo, ethyl ester,

quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[(4-cyanoanilino)-carbonyl]-7,13-dihydro-5,13-dioxo, ethyl ester,

quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(cyano)phenylsulphonyl]amino]-7,13-dihydro-5,13-dioxo, ethyl ester,

- 5 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(cyano)benzoyl]amino]-7,13-dihydro-5,13-dioxo, ethyl ester.

- b) A solution of 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanophenyl]methoxy]-1-methyl-6-oxo, ethyl ester (grams, mmol) in 3mL
 10 pyridine and 3mL diethylamine was saturated with hydrogen sulfide gas, bubbled through the solution *via* a pipette. The reaction was monitored by TLC (SiO₂, 100% ethyl acetate) and was complete after 3 hours. The mixture was concentrated *in vacuo* and dissolved in 3mL methylene chloride. The mixture was allowed to react with an excess of methyl iodide (2mL), heating to 50°C for 30 mins. The reaction was concentrated *in vacuo*, dissolved in
 15 ethanol (5mL) and heated to 50°C with ammonium acetate (0.5 grams) for 30 mins. The reaction was cooled to room temperature and purified by HPLC (solvent gradient 1:4 acetonitrile:water(0.5% acetic acid) (time 0 to 10 mins.) to 3:2 acetonitrile:water(0.5% acetic acid) (time 50 mins.), 20mL/min., C18 2" column, R_f=36.1 mins.) to yield 4H-[1,2,4]triazolo[4,3-
 20 a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenyl]methoxy]-1-methyl-6-oxo, ethyl ester. ¹H NMR (D₂O, δHOD) 7.60 (2H, d, ³J_{HH}=8Hz, 3,5H:4-NCC₆H₄), 7.46 (2H, ³J_{HH}=8Hz, 2,6H:4-NCC₆H₄), 7.33 (1H, d, ³J_{HH}=9Hz, C9 ArH), 7.23 (1H, d, ⁴J_{HH}=3Hz, C6 ArH), 7.18 (1H, dd, ⁴J_{HH}=3Hz, ³J_{HH}=9Hz, C8 ArH), 5.05 (2H, s, OCH₂Ar), 4.48 (1H, d, ²J_{HH}=16Hz, C5H), 4.27 (1H, d, ²J_{HH}=16Hz, C5H), 3.93 (1H, m, NCHH), 3.74 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.48 (1H, dt, ²J_{HH}=14Hz, ³J_{HH}=7Hz, NCHH), 2.55 (1H, m, CHHCO₂), 2.40 (1H, dt, ²J_{HH}=15Hz, ³J_{HH}=6Hz, CHHCO₂), 2.33 (3H, s, C1-Me), 0.85 (3H, t, ³J_{HH}=7Hz, CH₂CH₃).

- Using the above procedure, but substituting the appropriate 4H-[1,2,4]triazolo[4,3-
 30 a][1,4]benzodiazepine-5(6H)-propanoic acid, 1-methyl-6-oxo, ethyl ester for 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanophenyl]methoxy]-1-methyl-6-oxo, ethyl ester there may be prepared, for example, the following compounds:
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[2-[4-(iminoaminomethyl)phenyl]-2-oxoethyl]-1-methyl-6-oxo, ethyl ester,
 35 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)anilino]carbonyl]-1-methyl-6-oxo, ethyl ester,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenylsulphonyl]-amino]-1-methyl-6-oxo, ethyl ester,


- 4H-[1,2,4]triazolo[4,3-a][1,4]benzo-diazepine-5(6H)-propanoic acid, 8-[2-[4-(iminoaminomethyl)phenyl]-2-oxoethyl]-1-hydroxy-6-oxo, ethyl ester,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)analino]-carbonyl]-1-hydroxy-6-oxo, ethyl ester,
 5 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenylsulphonyl]-amino]-1-hydroxy-6-oxo, ethyl ester,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzo-diazepine-5(6H)-propanoic acid, 8-[2-[4-(iminoaminomethyl)phenyl]-2-oxoethyl]-6-oxo, ethyl ester,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)analino]-carbonyl]-6-oxo, ethyl ester,
 10 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenylsulphonyl]-amino]-6-oxo, ethyl ester,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)benzoyl]-amino]-6-oxo, ethyl ester,
 15 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[2-[4-(iminoaminomethyl)phenyl]-2-oxoethyl]-7,13-dihydro-5,13-dioxo, ethyl ester,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(iminoaminomethyl)analino]-carbonyl]-7,13-dihydro-5,13-dioxo, ethyl ester,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenylsulphonyl]-amino]-7,13-dihydro-5,13-dioxo, ethyl ester,
 20 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(iminoaminomethyl)benzoyl]-amino]-7,13-dihydro-5,13-dioxo, ethyl ester.

- c) To a solution of 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic
 25 acid, 8-[[4-(iminoaminomethyl)phenyl]methoxy]-1-methyl-6-oxo, ethyl ester in 3mL ethanol at room temperature was added 0.75 mL 2N sodium hydroxide and the reaction stirred for 45 mins. The mixture was quenched 0.5mL glacial acetic acid and purified by HPLC (solvent gradient 1:4 acetonitrile:water(0.5% acetic acid) (time 0 to 10 mins.) to 3:2 acetonitrile:water(0.5% acetic acid) (time 50 mins.), 20mL/min, C18 2" column, $R_f=33.2$
 30 mins.) to yield 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenyl]methoxy]-1-methyl-6-oxo, acetate. Exact mass (FAB, MH^+) calcd for $C_{22}H_{23}N_6O_4$: 435.1781, found 435.1744.

- Using the above procedure, but substituting the appropriate 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 1-methyl-6-oxo, ethyl ester for 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenyl]methoxy]-1-methyl-6-oxo, ethyl ester there may be prepared, for example, the following compounds:

- 4H-[1,2,4]triazolo[4,3-a][1,4]benzo-diazepine-5(6H)-propanoic acid, 8-[2-[4-(iminoaminomethyl)phenyl]-2-oxoethyl]-1-methyl-6-oxo, acetate,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)anilino]carbonyl]-1-methyl-6-oxo, acetate,
 5 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenylsulphonyl]-amino]-1-methyl-6-oxo, acetate,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzo-diazepine-5(6H)-propanoic acid, 8-[2-[4-(iminoaminomethyl)phenyl]-2-oxoethyl]-1-hydroxy-6-oxo, acetate,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)anilino]-carbonyl]-1-hydroxy-6-oxo, acetate,
 10 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenylsulphonyl]-amino]-1-hydroxy-6-oxo, acetate,
 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)benzoyl]amino]-1-hydroxy-6-oxo, acetate,
 15 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzo-diazepine-5(6H)-propanoic acid, 8-[2-[4-(iminoaminomethyl)phenyl]-2-oxoethyl]-6-oxo, acetate,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)anilino]-carbonyl]-6-oxo, acetate,
 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenylsulphonyl]amino]-6-oxo, acetate,
 20 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)benzoyl]amino]-6-oxo, acetate,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[2-[4-(iminoaminomethyl)phenyl]-2-oxoethyl]-7,13-dihydro-5,13-dioxo, acetate,
 25 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(iminoaminomethyl)anilino]-carbonyl]-7,13-dihydro-5,13-dioxo, acetate,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenylsulphonyl]amino]-7,13-dihydro-5,13-dioxo, acetate,
 quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(iminoaminomethyl)benzoyl]amino]-7,13-dihydro-5,13-dioxo, acetate,
 30 (iminoaminomethyl)benzoyl]amino]-7,13-dihydro-5,13-dioxo, acetate.

Example 5




5 *Quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]methoxy]-7,13-dihydro-5,13-dioxo, acetate*

a) Quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-cyanophenyl]methoxy]-7,13-dihydro-5,13-dioxo, ethyl ester was prepared using the method
 10 described in part (a) of example 4, substituting methyl anthranilate (1.42mL, 11.0mmol) for acetic hydrazide; yield 0.3 grams (33%). ¹H NMR (CDCl₃, δTMS) 8.33 (1H, d, ³J_{HH}=8Hz, ArH), 7.78 (2H, m, ArH), 7.71 (2H, d, ³J_{HH}=8Hz, 3,5H:4-NCC₆H₄), 7.56 (4H, m, 2,6H:4-NCC₆H₄, ArH), 7.43 (1H, d, ⁴J_{HH}=3Hz, C6 ArH), 7.17 (1H, d, ³J_{HH}=9Hz, C9 ArH), 5.20 (2H, s, OCH₂Ar), 4.52 (1H, d, ²J_{HH}=16Hz, C5H), 4.34 (1H, d, ²J_{HH}=16Hz, C5H), 4.04 (2H, q, ³J_{HH}=7Hz, OCH₂), 3.93 (2H, bt, NCH₂), 2.78 (1H, dt, ²J_{HH}=15Hz, ³J_{HH}=6Hz, CHHCO₂),
 15 2.62 (1H, dt, ²J_{HH}=15Hz, ³J_{HH}=6Hz, CHHCO₂), 1.18 (3H, t, ³J_{HH}=7Hz, CH₂CH₃).

b) Quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenyl]methoxy]-7,13-dihydro-5,13-dioxo, ethyl ester was prepared
 20 using the method described in part (b) of example 4. Thus, from 0.3 grams (0.6mmol) quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 3-[[4-cyanophenyl]methoxy]-7,13-dihydro-5,13-dioxo, ethyl ester was prepared quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid, 3-[[4-(aminoiminomethyl)phenyl]methoxy]-7,13-dihydro-5,13-dioxo, ethyl ester (0.22 grams, 71%, mp=235-238°C). The product precipitates from the crude
 25 reaction mixture and was isolated by suction filtration. The white powder was washed with a 1:1 mixture of ethanol:ether, and then ether.

c) Quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid,3-[[4-(aminoiminomethyl)phenyl]methoxy]-7,13-dihydro-5,13-dioxo, acetate was prepared by dissolving quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid,3-[[4-(iminoaminomethyl)phenyl]methoxy]-7,13-dihydro-5,13-dioxo, ethyl ester (50mg, 0.085 mmol), in 4mL THF and 2mL water at room temperature and sequentially adding 0.77mL 30% solution of hydrogen peroxide (9X excess) and 0.006 grams lithium hydroxide (0.256mmol, 3X excess). The mixture was stirred for 2 hours, quenched with a saturated solution of sodium bisulfite and 0.25mL glacial acetic acid. The mixture was concentrated *in vacuo* and purified by HPLC (solvent gradient 1:4 acetonitrile:water(0.5% acetic acid) (time 0 to 10 mins.) to 3:2 acetonitrile:water(0.5% acetic acid) (time 50 mins.), 20mL/min, C18 2" column) to yield quinazolino[3,2-a][1,4]benzodiazepine-6(5H)-propanoic acid,3-[[4-(aminoiminomethyl)phenyl]methoxy]-7,13-dihydro-5,13-dioxo, acetate (0.005 grams, 10%).

Example 6



4H-[1,2,3,4]Tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-6-oxo, acetate

20

a) A solution of 4-(2-carboxyethyl)-7-[(4-cyanophenyl)methoxy]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester (prepared in part (b) of Example 3, 1 g, 2.5mmol) diethyl diazodicarboxylate (DEAD, 0.39 mL, 2.5 mmol), triphenylphosphine (0.65 g, 2.5 mmol), and trimethylsilylazide (0.33 mL, 2.5 mmol) in 20 mL dry tetrahydrofuran was magnetically stirred for for 24 hrs. An additional equivalent of DEAD, triphenylphosphine, and trimethylsilylazide were added and the mixture stirred for an additional 48hrs. The mixture was concentrated *in vacuo* and purified by column chromatography (SiO₂, elution gradient of 30% ethyl acetate / 70% hexane to 100% ethyl acetate) followed by a second column (SiO₂, elution gradient 5% acetone / methylene chloride to 10% acetone / methylene chloride) to yield 0.787 g (72%) of 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-


30

propanoic acid, 8-[(4-cyanophenyl)methoxy]-6-oxo ethyl ester: mp = 64-67°C; TLC (SiO₂, 1:1 ethyl acetate: hexane) R_f = 0.11; ¹H NMR (CDCl₃, δTMS) 7.87 (1H, d, ³J_{HH} = 9Hz, ArH C10-H), 7.72 (2H, d, ³J_{HH} = 8Hz, ArH), 7.66 (1H, d, ⁴J_{HH} = 3Hz, ArH C7-H), 7.59 (2H, d, ³J_{HH} = 8Hz, ArH), 7.34 (1H, dd, ³J_{HH} = 9Hz, ⁴J_{HH} = 3Hz, ArH C9-H), 5.27 (2H, s, OCH₂Ar) 4.85 (2H, s, C4H), 4.13 (2H, q, ³J_{HH} = 7Hz, OCH₂), 3.93 (2H, t, ³J_{HH} = 6Hz, NCH₂CH₂CO₂), 2.70 (2H, t, ³J_{HH} = 6Hz, CH₂CO₂), 1.24 (3H, s, CH₃); ¹³C NMR (CDCl₃) 171.4, 165.2, 158.7, 151.7, 141.0, 132.5, 128.5, 127.6, 121.0, 123.8, 120.7, 118.4, 117.0, 112.1, 69.4, 61.0, 46.2, 41.3, 33.0, 30.9, 14.1. HRMS (FAB, M+H⁺) calcd. for C₂₂H₂₁N₆O₄ 433.1624, found 433.1596.

10 b) The amidino ester 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[(4-(iminoaminomethyl)phenyl) methoxy]-6-oxo ethyl ester, acetate was prepared using the method described in part (e) of Example 1. Thus 0.5 grams of 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[(4-cyanophenyl)methoxy]-6-oxo ethyl ester (1.16mmol) yielded 0.343 grams (66%) of the
15 amidino ester acetate salt. [Analytical HPLC Microsorb Short-One 80-200-C3, 9 mm x 85 mm ; elution gradient 0-100% acetonitrile (0.1% TFA)/water (0.1% TFA) over 11 min] R_t = 5.51 min: mp = 213-216°C; ¹H NMR (CD₃OD, δTMS) 7.88 (1H, d, ³J_{HH} = 9Hz, ArH C10-H), 7.84 (2H, d, ³J_{HH} = 8Hz, ArH) 7.73 (2H, d, ³J_{HH} = 8Hz, ArH), 7.62 (1H, d, ⁴J_{HH} = 3Hz, ArH C7-H), 7.47 (1H, dd, ³J_{HH} = 9Hz, ⁴J_{HH} = 3Hz, ArH C9-H), 5.36 (2H, s, OCH₂Ar) 4.90 (2H, s, C4H), 4.01 (2H, q, ³J_{HH} = 7Hz, OCH₂), 3.93 (2H, t, ³J_{HH} = 6Hz, NCH₂CH₂CO₂), 2.64 (2H, t, ³J_{HH} = 6Hz, CH₂CO₂), 1.14 (3H, t, ³J_{HH} = 7Hz, CH₃); ¹³C NMR (CDCl₃) 172.9, 168.2, 167.3, 160.4, 154.0, 144.4, 129.9, 129.2, 125.5, 125.1, 121.6, 118.3, 70.6, 61.9, 47.1, 41.7, 33.9, 22.6, 14.5; HRMS (FAB, M+H⁺) calc'd. for C₂₂H₂₁H₇O₄ 450.1890, found 450.1869.

25 c) 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[(4-(iminoaminomethyl)phenyl) methoxy]-6-oxo ethyl ester, acetate was saponified using the conditions described in part (f) of Example 1. Thus 50 mgs of the amidino ester (0.1mmol) yielded 30mgs of 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[(4-(iminoaminomethyl)phenyl) methoxy]-6-oxo, acetate. [Analytical HPLC Microsorb Short-
30 One 80-200-C3, 9 mm x 85 mm ; elution gradient 0-100% acetonitrile (0.1% TFA)/water (0.1% TFA) over 11 min] R_t = 4.60 min. ; mp = 238-240°C (dec.); ¹H NMR (DMSO-d₆, δTMS) 9.32(1H,bs, NH), 9.06 (1H, bs, NH), 7.87 (1H, d, ³J_{HH} = 9.0 Hz, ArH C10-H), 7.84 (2H, d, ³J_{HH} = 8.0 Hz, ArH), 7.71(2H,d,³J_{HH} = 8 Hz, ArH), 7.58 (1H, d, ⁴J_{HH} = 3Hz, ArH C7-H), 7.50 (1H, dd, ³J_{HH} = 9 Hz, ⁴J_{HH} = 3 Hz, ArH, C9-H), 5.40 (2H, s, C4H), 4.90 (2H,s, OCH₂Ar),
35 3.78 (2H, t, ³J_{HH} = 7 Hz, NCH₂CH₂CO₂H), 2.49 (2H, m, NCH₂CH₂CO₂H, coincident with reference peak for DMSO-d₆); HRMS (FAB, M+H⁺) calc'd for C₂₀H₂₀N₇O₄ 422.1577, found 422.1547.


Example 7



5 **4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-[N-(ethylcarbamoyl)iminoaminomethyl]phenyl]methoxy]-6-oxo ethyl ester**

- 10 a) To a solution of 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenyl]methoxy]-6-oxo ethyl ester, acetate (0.15 grams, 0.334 mmol), sodium bicarbonate (70 mgs, 2.5 molar equiv.) tetrahydrofuran(4 mL), and water (2 mL) was added ethyl chloroformate (48μL, 1.5 molar equiv.). The reaction was stirred for 2hrs, concentrated *in vacuo*, and purified by column chromatography (SiO₂, elution gradient 10% acetone / methylene chloride to 20% acetone / methylene chloride to yield 69.4 mg (40%)
- 15 of 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-[N-(ethylcarbamoyl)iminoaminomethyl]phenyl]methoxy]-6-oxo ethyl ester: TLC (SiO₂, 100% ethyl acetate) R_f = 0.42 uv positive; mp = 78-82°C; ¹H NMR (CDCl₃, δTMS) 9.57 (1H, bs, NH), 7.92 (2H, d, ³J_{HH} = 8Hz, ArH), 7.82 (1H, d, ³J_{HH} = 9Hz, ArH C10-H), 7.61 (1H, d, ⁴J_{HH} = 3Hz, ArH C7-H), 7.48 (2H, d, ³J_{HH} = 8Hz, ArH), 7.30 (1H, dd, ³J_{HH} = 9Hz, ⁴J_{HH} = 3Hz, ArH C9-H), 5.22 (2H, s, OCH₂Ar) 4.82 (2H, s, C4H), 4.19 (2H, q, ³J_{HH} = 7 Hz, H₃CCH₂OCONH), 4.12 (2H, q, ³J_{HH} = 7Hz, OCH₂), 3.91 (2H, t, ³J_{HH} = 6Hz, NCH₂CH₂CO₂), 2.69 (2H, t, ³J_{HH} = 6Hz, CH₂CO₂), 1.32 (3H, t, ³J_{HH} = 7Hz, CH₃), 1.22 (3H, t, ³J_{HH} = 7Hz CH₃); ¹³C NMR (CDCl₃) 171.4, 165.3, 158.9, 151.7, 139.9, 134.5, 128.3, 127.7, 127.2, 123.7, 120.7, 117.1, 69.8, 61.3, 61.0, 46.2, 41.3, 33.0, 14.4, 14.0; HRMS (FAB, M+H⁺)
- 25 calc'd for C₂₅H₂₈N₇O₆ 522.2101, found 522.2088.

Example 8



5 *4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid,
8-[[4-(iminoaminomethyl)phenyl]acetylene]-6-oxo, acetate*

a) *4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid,*


8-[[4-cyanophenyl]acetylene]-6-oxo was prepared from 4-(2-carboxyethyl)-7-[(4-cyanophenyl)acetylene]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester (see part
10 (b) of Example 1 for prep of the ethyl ester) using the method described in part (a) of Example 6 for conversion of the secondary amide into the tetrazole ring. Thus 1 gram (2.49mmol) of the 7-[(4-cyanophenyl)acetylene] benzodiazepinedione yielded 0.5 gram (47%) of the tetrazole: TLC (SiO₂, 1:1 ethyl acetate / hexane) R_f = 0.21, uv positive; mp =167-169°C; ¹H NMR (CDCl₃, δTMS) 8.30 (1H, d, ⁴J_{HH} = 2Hz, ArH C7-H), 7.97 (1H, d, ³J_{HH} = 9Hz, ArH C10-H), 7.86 (1H, dd, ⁴J_{HH} = 2Hz, ³J_{HH} = 9Hz, ArH C9-H), 7.68 (2H, d, ³J_{HH} = 8Hz, ArH), 7.64 (2H, d, ³J_{HH} = 8Hz, ArH), 4.88 (2H, s, C4H), 4.15 (2H, q, ³J_{HH} = 7Hz, OCH₂), 3.95 (2H, t, ³J_{HH} = 6Hz, NCH₂CH₂CO₂), 2.73 (2H, t, ³J_{HH} = 6Hz, CH₂CO₂), 1.25 (3H, s, CH₃); ¹³C NMR (CDCl₃) 171.5, 164.8, 152.1, 136.3, 135.8, 132.2, 129.7, 127.3, 126.9, 124.5, 122.3, 118.2,
15 112.4, 90.7, 64.1, 46.5, 41.3, 33.0, 14.2. HRMS (FAB, M+H⁺) calc'd. for C₂₃H₁₈N₆O₃ 427.1519, found 427.1489.

b) *4H-[1,2,3,4] tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8 - [[4-(iminoaminomethyl)phenyl]acetylene] -6-oxo, ethyl ester , acetate* was prepared as in part
25 (e) of Example 1. Thus, *4H-[1,2,3,4] tetrazolo [4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8- [[4-cyanophenyl]acetylene]- 6-oxo ethyl ester, acetate* (0.5 grams, 1.17mmol) yielded the desired amidino ester (0.175 grams, 34%) : ¹H NMR (CD₃OD, δTMS) 8.23 (1H, d, ⁴J_{HH} = 2Hz, ArH C7-H), 8.03 (1H, d, ³J_{HH} = 9Hz, ArH C10-H), 7.97 (1H, dd, ⁴J_{HH} = 2Hz, ³J_{HH} = 9Hz, ArH C9-H), 7.84 (2H, d, ³J_{HH} = 9Hz, ArH), 7.74 (2H, d, ³J_{HH} = 9Hz, ArH), 4.96 (2H, s,

C4H), 4.04 (2H, q, $^3J_{HH} = 7\text{Hz}$, OCH₂), 3.97 (2H, t, $^3J_{HH} = 6\text{Hz}$, NCH₂CH₂CO₂), 2.67 (2H, t, $^3J_{HH} = 6\text{Hz}$, CH₂CO₂), 1.15 (3H, s, CH₃); ¹³C, HRMS

- 5 c) Saponification of the amidino ester 4H - [1,2,3,4] tetrazolo [4,3a][1,4] benzodiazepine-5(6H)-propanoic acid, 8-[[4-(iminoaminomethyl)phenyl]acetylene]- 6-oxo ethyl ester, acetate was accomplished using the method described in part (f) of Example 1 to yield 4H - [1,2,3,4] tetrazolo [4,3a][1,4] benzodiazepine- 5(6H)- propanoic acid, 8- [[4-(iminoaminomethyl)phenyl]acetylene]- 6-oxo, acetate in 95% yield; ¹H NMR (DMSO-d₆, δ TMS) 8.64 (1H, bs, ArH C7-H), 8.44 (2H, bs, ArH C9,10-H), 8.26 (4H, bs, ArH), 5.33 (2H, s, C4H), 3.03 (2H, t, $^3J_{HH} = 6\text{Hz}$, NCH₂CH₂CO₂), 2.45 (2H, t, $^3J_{HH} = 6\text{Hz}$, CH₂CO₂); HRMS (M+H⁺) calc'd. for C₂₁H₁₈N₇O₃ 416.1471, found 416.1448.
- 10

Example 9



15


*4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid,
8-[[4-[N-(ethylcarbamoyl)iminoaminomethyl]phenyl]acetylene]-6-oxo, ethyl ester*

- 20 a) The carbamate ester 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-[N-(ethylcarbamoyl)iminoaminomethyl]phenyl]acetylene]-6-oxo, ethyl ester was prepared as described in part (a) of Example 7 to afford the carbamate ester in a 30% yield; TLC (SiO₂, 100% ethyl acetate) R_f = 0.73, uv positive; mp = 91-95°C; ¹H NMR (CDCl₃, δ TMS) 9.6 (1H, bs, NH), 7.92 (2H, d, $^3J_{HH} = 8\text{Hz}$, ArH), 8.26 (1H, d, $^4J_{HH} = 2\text{Hz}$, ArH C7-H), 7.93 (2H, d, $^3J_{HH} = 8\text{Hz}$, ArH), 7.86 (2H, d, $^3J_{HH} = 8\text{Hz}$, ArH), 7.82 (1H, dd, $^3J_{HH} = 9\text{Hz}$, $^4J_{HH} = 3\text{Hz}$, ArH C9-H), 7.59 (1H, d, $^3J_{HH} = 9\text{Hz}$, ArH C10-H), 4.87 (2H, s, C4H), 4.23 (2H, q, $^3J_{HH} = 7\text{Hz}$, H₃CCH₂OCONH), 4.15 (2H, q, $^3J_{HH} = 7\text{Hz}$, OCH₂), 3.94 (2H, t, $^3J_{HH} = 6\text{Hz}$, NCH₂CH₂CO₂), 2.70 (2H, t, $^3J_{HH} = 6\text{Hz}$, CH₂CO₂), 1.38 (3H, t, $^3J_{HH} = 7\text{Hz}$, CH₃), 1.22 (3H, t, $^3J_{HH} = 7\text{Hz}$ CH₃); ¹³C NMR (CDCl₃) 171.5, 164.9, 152.1, 136.1, 135.8,
- 25

134.8, 131.9, 129.5, 127.4, 126.1, 125.1, 122.3, 91.8, 89.3, 61.5, 61.1, 46.4, 41.4, 33.0, 14.4, 14.1;
HRMS (M+H⁺) calc'd. for C₂₆H₂₆N₇O₅ 516.1995, found 516.1976.

Example 10

5



4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)benzoyl] amino]-1-methyl-6-oxo, trifluoroacetate

10

a) A slurry of 2-amino-5-[4-cyanobenzoyl]amino-1-benzoyl-(ethyl-3-aminopropanoate) (8 grams, 2.12 mmol) in 1,2-dichloroethane (100 ml) was charged with 2,6-lutidine (2.5 ml, 21.2 mmol) and heated to 60°C. Over a 3 hour period diphenylmethylbromide (10 grams, 40 mmol, Aldrich) dissolved in 1,2-dichloroethane (50 ml) was added in 4 equal portions. The reaction mixture was allowed to cool to room temperature and concentrated *in vacuo*. The residue was dissolved in ethyl acetate (50 ml), transferred to a separatory funnel and washed with water (50 ml), 5% sodium bicarbonate solution (50 ml) and brine (50 ml). The organic phase was dried over magnesium sulfate, filtered, concentrated *in vacuo*, and purified by column chromatography (SiO₂, 1:1 ethyl acetate:hexane) to yield 10 grams (86% yield) of N-(2-diphenylmethylamino)-5-[4-cyanobenzoyl] amino -1-benzoyl-(ethyl-3-aminopropanoate). ¹H NMR (CDCl₃, δTMS) 8.25 (1H, s, Ar-CONH-Ar), 7.88 (2H, d, ³J_{HH} = 7.8 Hz, *o*-CN ArH), 7.74 (1H, br s, C6 aniline ArH), 7.65 (2H, d, ³J_{HH} = 7.8 Hz, *m*-CNArH), 7.15-7.35 (12H, m, ArCHAr, ArNH(CH)Ar₂, C4 aniline ArH), 6.96 (1H, br t, ³J_{HH} = 6.0 Hz, Ar CONHCH₂), 6.48 (1H, d, ³J_{HH} = 9 Hz, C3 aniline ArH), 5.54 (1H, s, ArCHAr), 4.11 (2H, q, ³J_{HH} = 6.8 Hz, CO₂CH₂CH₃), 3.59 (2H, q, ³J_{HH} = 5.9 Hz, CONHCH₂CH₂CO₂Et), 2.56 (2H, t, CONHCH₂CH₂CO₂Et), 1.23 (3H, t, ³J_{HH} = 6.8 Hz, CO₂CH₂CH₃). ¹³C NMR (CDCl₃) 172.6, 169.2, 163.9, 146.2, 142.4, 138.7, 132.3, 128.8, 128.4, 127.8, 127.7, 127.3, 127.2, 126.3, 125.6, 120.8, 118.0, 115.2, 114.9, 113.5, 62.1, 60.8, 35.2, 34.0, 14.1. Exact mass (FAB M+H⁺) C₃₃H₃₀N₄O₄ calc'd. 547.2346, found 547.2309

30

b) To a biphasic $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ (50 ml/50 ml) solution of N-(2-diphenyl methylamino)-5-[4-cyanobenzoyl]amino-1-benzoyl-(ethyl-3-aminopropanoate) (9 grams, 16 mmol), bromoacetyl bromide (5.90 grams, 29.2 mmol) and potassium carbonate (4.14 grams, 30 mmol) were added in 3 equal portions over 25 minutes. The pH was maintained at 7 - 8. After stirring at room temperature for 1 hour, the reaction was transferred to a separatory funnel and the layers separated. The organic phase was dried over magnesium sulfate, filtered, concentrated *in vacuo*, and purified by column chromatography (SiO_2 , 2:1 ethyl acetate:hexane) to yield 6.68 grams (61% yield) of N-[2-diphenylmethylamino]-N-bromoacetyl-5-[4-cyanobenzoyl]amino-1-benzoyl-(ethyl-3-aminopropanoate). ^1H NMR (CDCl_3 , δTMS) 9.75 (1H, s, Ar-CONH-Ar), 8.05 (2H, d, $^3J_{\text{HH}} = 7.2$ Hz, o -CN ArH), 7.98 (1H, br s, C6 aniline ArH), 7.71, (2H, d, $^3J_{\text{HH}} = 7.2$ Hz, m -CNArH), 7.58 (1H, br d, $^3J_{\text{HH}} = 8.7$ Hz, C4 aniline ArH), 7.30-7.00 (10H, m, ArCHAr), 6.82 (1H, s, ArCHAr), 6.74 (1H, d, $^3J_{\text{HH}} = 8.7$ Hz, C3 aniline ArH), 5.73 (1H, br t, $^3J_{\text{HH}} = 5.4$ Hz, Ar CONHCH₂), 4.18 (2H, q, $^3J_{\text{HH}} = 7.2$ Hz, $\text{CO}_2\text{CH}_2\text{CH}_3$), 3.83 (2H, br s, NCOCH_2HBr), 3.47 (1H, m, $^3J_{\text{HH}} = 6.9$ Hz, CONHCHHCH₂CO₂Et), 3.13 (1H, m, $^3J_{\text{HH}} = 6.9$ Hz, CONHCHHCH₂CO₂Et), 2.46 (2H, br t, CONHCH₂CH₂CO₂Et), 1.28 (3H, t, $^3J_{\text{HH}} = 7.2$ Hz, $\text{CO}_2\text{CH}_2\text{CH}_3$). ^{13}C NMR (CDCl_3) 172.0, 169.3, 166.5, 164.5, 139.6, 139.3, 138.1, 136.6, 135.4, 132.7, 132.3, 131.3, 128.3, 128.2, 128.0, 127.4, 127.2, 122.6, 119.2, 118.0, 115.0, 64.4, 60.8, 35.3, 33.5, 29.9, 14.1. Mass spec. (FAB $\text{M}+\text{H}^+$) calc'd 667.15, found 667.2 Exact mass (FAB $\text{M}+\text{H}^+$) $\text{C}_{35}\text{H}_{32}\text{BrN}_4\text{O}_5$ calc'd 667.1556, found 667.1538.

c) To a stirring DMF (400 ml) solution of N-[2-diphenylmethylamino]-N-bromoacetyl-5-[4-cyanobenzoyl]amino-1-benzoyl-(ethyl-3-aminopropanoate) (6.56 grams, 9.83 mmol) at room temperature, solid cesium carbonate (6.41 grams, 19.6 mmol) was added in one portion. The heterogeneous solution was stirred at rt for 2 hours, then poured into a separatory funnel containing ethyl acetate (500 ml) and water (500 ml). After thorough mixing, the aqueous phase was removed and the organic phase was washed with water (200 ml), brine (300 ml), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give 5.36 grams (93 % yield) of 1-(diphenylmethyl)-4-(2-carboxyethyl)-7-[4-[cyano)benzoyl]-amino] -3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester. ^1H NMR (CDCl_3 , δTMS) 9.47 (1H, s, ArCONHAr), 7.83 (2H, d, $^3J_{\text{HH}} = 7.8$ Hz, o -CNArH), 7.77 (1H, s, C6 ArH), 7.46 (2H, d, $^3J_{\text{HH}} = 8.4$ Hz, m -CNArH), 7.25-7.0 (11H, m, ArCHAr, upon expansion of this region C8 ArH appears @ 7.09 as a dd, $^3J_{\text{HH}} = 7.8$ Hz, $^4J_{\text{HH}} = 2.0$ Hz), 6.97 (1H, d, $^3J_{\text{HH}} = 8.4$ Hz, C9 ArH), 6.62 (1H, s, ArCHAr), 4.09 (1H, d, $^2J_{\text{HH}} = 14.4$ Hz, $\text{ArN(R')COCHHNR(R)CO}$), 3.96 (2H, q, $^3J_{\text{HH}} = 7.2$ Hz, $\text{CO}_2\text{CH}_2\text{CH}_3$), 3.81 (1H, m, $^3J_{\text{HH}} = 7.2$ Hz, CONHCHHCH₂CO₂Et), 3.70 (1H, d, $^2J_{\text{HH}} = 14.4$ Hz, $\text{ArN(R')COCHHNR(R)CO}$), 3.55 (1H, m, $^3J_{\text{HH}} = 7.2$ Hz, (CONHCHHCH₂CO₂Et), 2.39 (2H, br t, CONHCH₂CH₂CO₂Et), 1.10, (3H, t, $^3J_{\text{HH}} = 7.2$ Hz, $\text{CO}_2\text{CH}_2\text{CH}_3$). ^{13}C NMR (CDCl_3) 170.8, 168.2, 166.6, 164.2, 138.1, 138.0, 137.6, 136.4, 135.5,

132.0, 130.3, 129.1, 128.2, 128.1, 128.0, 127.9, 127.6, 127.4, 124.4, 123.1, 121.9, 117.8, 114.9, 67.0, 60.6, 44.6, 32.0, 14.0. Exact mass (FAB M+H⁺) C₃₅H₃₀N₄O₅ calc'd 587.2295, found 587.2324

- d) A slurry of 1-(diphenylmethyl)-4-(2-carboxyethyl)-7-[4-(cyano)benzoyl]-amino]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester (5.36 grams, 9.14 mmol), anisole (5 ml) and ethyl methyl sulfide (5 ml) in a teflon tube was cooled to -196°C (N₂) and hydrogen fluoride (30 ml) was condensed into the teflon tube. The reaction was stirred for 30 min at -196°C, allowed to warm to 0° C, and concentrated *in vacuo* over 2 hrs. The resulting residue was triturated with ethyl ether, collected in a filter and dried *in vacuo*, to give 3.22 grams (56% yield) of 4-(2-carboxyethyl)-7-[4-(cyano)benzoyl]-amino]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester. TLC R_f = 0.46 (SiO₂, 1:1 hexane:ethyl acetate, uv positive), mp = 253-257°C (dec.) ¹H NMR (DMSO-d₆, δTMS) 10.63 (1H, s, ArCONHAr), 10.45 (1H, s, ArNHCOCH₂N(R)CO), 8.16 (1H, d, ⁴J_{HH} = 3.3 Hz, C6 ArH), 8.12 (2H, d, ³J_{HH} = 8 Hz, *o*-CN ArH), 8.03 (2H, d, ³J_{HH} = 8 Hz, *m*-CN ArH), 7.95 (1H, dd, ³J_{HH} = 8.7 Hz, ⁴J_{HH} = 3.3 Hz, C8 ArH), 7.10 (1H, d, ³J_{HH} = 8.7 Hz, C9 ArH), 4.06 (2H, q, ³J_{HH} = 7.2 Hz, CO₂CH₂CH₃), 3.91 (2H, s, ArNHCOCH₂N(R)CO), 3.81 (2H, t, ³J_{HH} = 7.2 Hz, NCH₂CH₂CO₂Et), 2.62 (2H, t, ³J_{HH} = 7.2 Hz, NCH₂CH₂CO₂Et), 1.18 (3H, t, ³J_{HH} = 7.2 Hz, CO₂CH₂CH₃). ¹³C NMR (DMSO- d₆) 170.9, 170.0, 166.2, 164.0, 138.6, 134.8, 133.2, 132.5, 128.5, 126.2, 124.4, 122.3, 121.2, 118.3, 114.0, 60.1, 50.9, 44.5, 32.4, 14.0. Mass spec. (FAB M+H⁺) calc'd 421.4, found 421.4. Exact mass (FAB M+ H⁺) C₂₂ H₂₁ N₄ O₅ calc'd 421.1512, found 421.1501.

- e) 4-(2-Carboxyethyl)-7-[4-(cyano)benzoyl]-amino]-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester, (1.5 grams, 3.6 mmol), Lawesson's reagent (2,4-bis(4-methoxyphenyl)1,3,2,4-dithiadiphosphetane 2,4-disulfide, 1.43 grams, 3.6 mmol) and anhydrous THF (20 ml) were combined and heated to 50°C. The slurry was stirred for 2 hours, cooled to room temperature and the resulting solid was collected and washed with THF to give 1.1 grams of 4-(2-carboxyethyl)-7-[4-(cyano)benzoyl]-amino)-3,4-dihydro-1H-1,4-benzodiazepine-2-thio-5-oxo ethyl ester (70% yield). TLC R_f = 0.47 (SiO₂, 1:2 hexane:ethyl acetate, uv positive) ¹H NMR (DMSO-d₆, δTMS) 10.72 (1H, s, ArCONHAr), 8.24 (1H, br s, C6 ArH), 8.14 (2H, d, ³J_{HH} = 7.8 Hz, *o*-CN ArH), 8.04 (2H, d, ³J_{HH} = 7.8 Hz, *m*-CN ArH), 8.01 (1H, br d, ³J_{HH} = 8.7 Hz, C8 ArH), 7.26 (1H, d, ³J_{HH} = 8.7 Hz, C9 ArH), 4.25 (2H, s, ArNHCSCH₂N(R)CO), 4.06 (2H, q, ³J_{HH} = 6.9 Hz, CO₂CH₂CH₃), 3.80 (2H, t, ³J_{HH} = 6.9 Hz, NCH₂CH₂CO₂Et), 2.71 (2H, t, ³J_{HH} = 6.9 Hz, NCH₂CH₂CO₂Et), 1.18 (3H, t, ³J_{HH} = 6.9 Hz, CO₂CH₂CH₃). Mass spec. (FAB M+H⁺) calc'd 437.1 found 437.1. Exact mass (FAB M+ H⁺) C₂₂ H₂₁ N₄ SO₄ calc'd 437.1284, found 437.1269.

f) A slurry of 4-(2-carboxyethyl)-7-[4-(cyano)benzoyl]-amino-3,4-dihydro-1H-1,4-benzodiazepine-2-thio-5-oxo ethyl ester (1 gram, 2.3 mmol) in anhydrous acetonitrile (60 mL) was treated with methyl iodide (0.65 grams, 4.6 mmol) and heated to reflux. The reaction flask was charged with additional methyl iodide (0.65 grams, 4.6 mmol) at six, seven and
 5 eight hours of reflux. Soon after the last charge with methyl iodide (t = 8h) the reaction turned clear. The solution was dried *in vacuo*, taking care to avoid exposure to moisture. The crude reaction mixture was treated with a solution of propargyl amine (0.73 ml, 11.5 mmol) and pyridinium chloride (0.27 grams, 2.3 mmol) in anhydrous toluene (250 ml). The resulting
 10 slurry was heated to reflux for 6 hours and the warm toluene solution decanted from the solids to give 350 mg of the crude imidazole. The crude product was purified (RP - Dynamax 60A C18, methanol/water/ 0.1% TFA, gradient elution 5 % methanol for 15 minutes, 5 to 90% methanol in 70 minutes, product collected from 60-70 minutes, UV @ 254 nm) to give 4H
 15 -imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanobenzoyl] amino -1-methyl-6-oxo, ethyl ester, (240 mg, 23 % yield) after lyophilization. ¹H NMR (DMSO-d₆, δTMS) 10.89 (1H, s, ArCONHAr), 8.36 (1H, d, ⁴J_{HH} = 2.4 Hz, C7 ArH), 8.22 (1H, dd, ³J_{HH} = 9 Hz, ⁴J_{HH} = 2.4 Hz, C9 ArH), 8.16 (2H, d, ³J_{HH} = 8.2 Hz, o -CN ArH), 8.08 (2H, d, ³J_{HH} = 8.2 HZ, m -CN ArH), 7.74 (1H, d, ³J_{HH} = 9 Hz, C10 ArH), 7.30 (1H, s, C2 imidazole CH), 4.68 (1H, d, ²J_{HH} = 16 Hz, ArNCNCHHN(R)CO), 4.51 (1H, d, ²J_{HH} = 16 Hz, ArNCNCHHN(R)CO), 4.03 (2H, q, ³J_{HH} = 6.9 Hz, CO₂CH₂CH₃), 3.79 (2H, m, NCH₂CH₂CO₂Et), 2.62 (2H, t, ³J_{HH} = 7.2 Hz, NCHCH₂CO₂Et), 2.39 (3H, s, imidazole CH₃), 1.14 (3H, t, ³J_{HH} = 6.9 Hz, CO₂CH₂CH₃). Mass spec. (FAB M+H⁺) calc'd 458.2, found 458.1 Exact mass (FAB M+ H⁺) C₂₅ H₂₄ N₅ O₄ calc'd 458.1828, found 458.1819.

g) 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanobenzoyl] amino -1-methyl-6-oxo, ethyl ester (80 mg, 0.17 mmol) was dissolved in anhydrous pyridine/triethylamine (2ml/1 ml), transferred to a pressure tube, and the pyridine/Et₃N solution saturated with hydrogen sulfide. The pressure tube was sealed, heated to 65°C and stirred for 30 min. After cooling to room temperature, the reaction solution was poured into a
 25 separatory funnel containing ethyl acetate (30 ml) and water (30 ml). After thorough mixing, the aqueous phase was removed and the organic phase was washed with water (20 ml), brine (30 ml), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give 75 mg (90 % yield) 4H -imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-thioamidobenzoyl] amino -1-methyl-6-oxo, ethyl ester. ¹H NMR (CDCl₃, δTMS) 9.47 (1H, s, ArCONHAr), 8.46 (1H, dd, ³J_{HH} = 8.7 Hz, ⁴J_{HH} = 2.4 Hz, C9 ArH), 8.23 (1H, brs, NH), 8.11
 30 (1H, d, ⁴J_{HH} = 2.7 Hz, C7 ArH), 8.03 (1H, brs, NH), 7.71 (2H, d, ³J_{HH} = 8.0 Hz, o -H₂NCS ArH), 7.67 (1H, m, C10 ArH), 7.30 (2H, m, m -H₂NCS ArH), 6.88 (1H, s, C2 imidazole CH), 4.45 (1H, d, ²J_{HH} = 15.6 Hz, ArNCNCHHN(R)CO), 4.33 (1H, d, ²J_{HH} = 15.6 Hz, ArNCNCHHN(R)CO), 4.13 (2H, q, ³J_{HH} = 7.0 Hz, CO₂CH₂CH₃), 3.68 (2H, m,


NCH₂CH₂CO₂Et), 2.57 (2H, t, ³J_{HH} = 6.9 Hz, NCH₂CH₂CO₂Et), 2.33 (3H, s, imidazole CH₃), 1.25 (3H, t, ³J_{HH} = 7.0 Hz, CO₂CH₂CH₃). Mass spec. (FAB M+H⁺) calc'd 492, found 492. Exact mass (FAB M+ H⁺) C₂₅ H₂₆ N₅SO₄ calc'd 492.1706, found 492.1715.

- 5 h) 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-thioamidobenzoyl]amino-1-methyl-6-oxo, ethyl ester (75 mg, 0.16 mmol) was suspended in anhydrous acetonitrile (3 ml) in a pressure tube. The slurry was treated with methyl iodide (15 ml, 0.25 mmol), the pressure tube sealed, and the stirred heterogeneous solution heated to 85°C (oil bath) for 1 hour. After heating for one hour the reaction mixture was homogeneous.
- 10 The solution was dried *in vacuo*, taking care to avoid exposure to moisture. The crude product was dissolved in ethanol (3 ml), treated with ammonium acetate (52 mg, 0.68 mmol), stoppered, and stirred overnight at room temperature. The reaction mixture was concentrated *in vacuo* and the crude product purified by prep RP HPLC (RP-Dynamax 60A C18, methanol/water/0.1% TFA, gradient elution 5 % methanol for 15 minutes, 5 to 90% methanol
- 15 in 70 minutes, retention time 33-40 minutes, UV @ 254 nm) to give 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl) benzoyl] amino] -1-methyl-6-oxo, ethyl ester, trifluoroacetate (24.5 mg, 24% yield) after lyophilization. ¹H NMR (CDCl₃, δTMS) 10.8 (1H, s, ArCONHAr), 9.45 (1H, brs, NH), 9.12 (2H, brs, NH), 8.33 (1H, d, ⁴J_{HH} = 2.7 Hz, C7 ArH), 8.21 (1H, dd, ³J_{HH} = 11.7 Hz, ⁴J_{HH} = 2.7 Hz, C9 ArH), 8.20
- 20 (2H, d, ³J_{HH} = 8.4 Hz, *o*-H₂NCNH ArH), 7.97 (2H, d, ³J_{HH} = 8.4 Hz, *m*-H₂NCNH ArH), 7.66 (1H, d, ³J_{HH} = 11.7 Hz, C10 ArH), 7.06 (1H, s, C2 imidazole CH), 4.58 (2H, d, ²J_{HH} = 15.9 Hz, ArNCNCHHN(R)CO), 4.41 (2H, d, ²J_{HH} = 15.9 Hz, ArNCNCHHN(R)CO), 4.03 (2H, q, ³J_{HH} = 7.2 Hz, CO₂CH₂CH₃), 3.77 (2H, t, ³J_{HH} = 7.5 Hz, NCH₂CH₂CO₂Et), 2.59 (2H, t, ³J_{HH} = 7.5 Hz, NCH₂CH₂CO₂Et), 2.35 (3H, s, imidazole CH₃), 1.15 (3H, t, ³J_{HH} = 7.2 Hz, CO₂CH₂CH₃). Mass spec. (FAB M+H⁺) C₂₅H₂₇N₆O₄ calc'd 475 found 475.2 Exact mass (FAB M+ H⁺) C₂₅ H₂₇ N₆O₄ calc'd 475.2090, found 475.2096.

- i) 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl) benzoyl] amino]-1-methyl-6-oxo, ethyl ester, trifluoroacetate (23 mg, 0.40 mmol) was dissolved in 3:2:1 tetrahydrofuran:methanol:water (0.5 ml), treated with 1 N sodium hydroxide solution (1.25 ml, 3.0 equiv) and stirred at room temperature overnight. The reaction mixture was concentrated *in vacuo*. and the crude product purified (RP - Dynamax 60A C18, methanol/water/0.1% TFA, gradient elution 0 % methanol for 20 minutes, 0 to 24% methanol in 40 minutes, product collected from 50 - 52.5 minutes, UV @ 254 nm) to give 4H
- 35 -imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl) benzoyl]amino]1-methyl-6-oxo, trifluoroacetate (24.5 mg, 24% yield). ¹H NMR (DMSO- d₆, δTMS) 10.8 (1H, s, ArCONHAr), 9.45 (1H, brs, NH), 9.11 (2H, brs, NH₂), 8.33 (1H, d, ⁴J_{HH} = 2.4 Hz, C7 ArH), 8.20 (2H, d, ³J_{HH} = 8.4 Hz, *o*-H₂NCNH ArH plus hidden under the

resonance @ 8.20 , 1H, dd, J values could not be determ'd C9 ArH), 7.97 (2H, d, $^3J_{HH}$ = 8.4 Hz, *m*-H₂NCNH ArH), 7.66 (1H, d, $^3J_{HH}$ = 11.7 Hz, C10 ArH), 7.03 (1H, s, C2 imidazole CH), 4.58 (2H, d, $^2J_{HH}$ = 16.2 Hz, ArNCNCHHN(R)CO), 4.41 (2H, d, $^2J_{HH}$ = 16.2 Hz, ArNCNCHHN(R)CO), ~3.7 hidden partially by water peak, (2H, t, NCH₂CH₂CO₂H), ~2.5 hidden under DMSO, (2H, t, $^3J_{HH}$ = 7.5 Hz, NCH₂CH₂CO₂H), 2.35 (3H, s, imidazole CH₃). Exact mass (FAB M+H⁺) C₂₃H₂₃N₆O₄ calc'd 447.1781, found 447.1764. Fg IIbIIIa ELISA IC₅₀ = 9 nM. PA (citrate)IC₅₀ = 85 nM

Example 11



4H-[1,2,3,4] tetrazolo [4,3a][1,4] benzodiazepine-5 (6H)-propanoic acid, 8-[[4-(hydroxyiminoaminomethyl)- benzoyl] amino] -6-oxo,

15


a) An ethanolic solution (0.6 ml, anhydrous absolute EtOH, over 3A molecular sieves for 48 h prior to use) of 4H-imidazo[1,2-a][1,4] benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanobenzoyl]amino-1-methyl-6-oxo, ethyl ester (see Example 11, part (f), 113 mg, 0.25 mmol) was treated with hydroxylamine hydrochloride (28 mg, 0.41 mmol) followed by sodium ethoxide (24.5 mg, 0.360 mmol). After stirring at room temperature for five minutes the reaction flask was stoppered and placed in a 50°C oil bath for 16 hours. TLC (SiO₂, 4% MeOH, 15% acetone in EtOAc) indicated complete consumption of the nitrile. The reaction mixture was diluted to a volume of approximately 2.5 ml by additon of EtOH(approx 0.4 ml), H₂O(approx.1.0 ml), and acetic acid(approx. 0.5 ml) and the crude purified directly by prep RP HPLC (Dynamax 60A, C18 83-221C, acetonitrile/water/0.5% HOAc, gradient elution 5% CH₃CN to 31% CH₃CN in 55 minutes, retention time 31-42 minutes, UV @ 254) to afford 70 mg (58%) of the desired amidooxime/ ethyl ester, 4H -imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(hydroxyaminoiminomethyl)benzoyl]amino]-1-methyl-6-oxo, ethyl ester. TLC R_f = 0.04 (SiO₂, 4% MeOH, 15% acetone in EtOAc); ¹H NMR (CD₃OD, δTMS) 8.26 (1H, d, $^4J_{HH}$ = 2.0 Hz , C7 ArH), 8.19 (1H, dd, $^3J_{HH}$ = 9.0 Hz, $^4J_{HH}$ =

30

2.0 Hz, C9 ArH), 8.0 (2H, d, $^3J_{HH} = 8.4$ Hz, *o*-HONHCNH ArH), 7.82 (2H, d, $^3J_{HH} = 8.4$ Hz, *m*-HONHCNH ArH), 7.55 (1H, d, $^3J_{HH} = 9.0$ Hz, C10 ArH), 6.87 (1H, brs, C2, imidazole CH), 4.46 (2H, brs, ArNCNCHHN(R)CO), 4.10 (2H, q, $^3J_{HH} = 7.2$ Hz, CO₂CH₂CH₃), 3.93 (1H, m, NCHHCH₂CO₂Et), 3.86 (1H, m, NCHHCH₂CO₂Et), 2.67 (2H, m, NCH₂CH₂CO₂Et), 2.38 (3H, s, imidazole CH₃), 1.21 (3H, t, $^3J_{HH} = 7.2$ Hz, CO₂CH₂CH₃). ¹³C NMR (CD₃OD) 172.87, 168.63, 168.11, 154.33, 147.50, 139.62, 137.62, 136.64, 131.67, 130.11, 130.01, 129.42, 128.83, 128.32, 127.64, 127.34, 125.46, 125.32, 124.10, 61.79, 49.83, 45.88, 33.5, 14.40, 10.93. Mass spec. (FAB M+H⁺) calc'd 491.2, found 491.2. Exact mass (FAB M+H⁺)C₂₅H₂₇N₆O₅ calc'd 491.2042, found 491.2043.

10

Example 12



15

4H-[1,2,3,4] tetrazolo [4,3a][1,4] benzodiazepine-5 (6H)-propanoic acid, 8-[[4-(iminoaminomethyl)- benzoyl] amino]- 6-oxo, trifluoroacetate

a) 4-(2-Carboxyethyl)-7-[4-(cyano)benzoyl]amino-3,4-dihydro-1H-1,4-benzodiazepine-2,5-dione ethyl ester (Example 10, part (d), 0.312 grams, 0.74 mmol), triphenylphosphine (0.126 grams, 1.48 mmol), trimethylsilylazide (0.200 ml, 1.48 mmol) diethyl azodicarboxylate and anhydrous glyme (2 ml) were stirred at room temperature for 16 hours. The reaction mixture was concentrated *in vacuo* and the crude product purified by sequential column chromatography (SiO₂, 1:1 hexane:ethyl acetate, then RP C18 Dynamax 60A 83-221 acetonitrile/water/0.1% TFA gradient elution 15 to 50% acetonitrile in 75 minutes, retention time 50-56 minutes, UV @ 254 nm) to give 4-H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanobenzoyl]amino]-6-oxo, ethyl ester (170 mg, 52% yield). TLC R_f = 0.3 (SiO₂, 1:2 hexane:ethyl acetate, uv positive); ¹H NMR (DMSO-d₆, δTMS) 10.94 (1H, s, ArCONHAr), 8.46 (1H, d, $^4J_{HH} = 2.4$ Hz, C7 ArH), 8.30 (1H, dd, $^4J_{HH} = 2.4$ Hz, $^3J_{HH} = 8.7$ Hz, C9 ArH), 8.17 (2H, d, $^3J_{HH} = 8.1$ Hz, *o*-CN ArH), 8.06 (2H, d, $^3J_{HH} = 8.1$ Hz, *m*-CN ArH), 7.97 (1H, d, $^3J_{HH} = 8.7$ Hz, C10 ArH), 4.97 (2H, s, ArNCNCH₂N(R)CO), 3.95 (2H, q, $^3J_{HH} = 6.9$ Hz, CO₂CH₂CH₃), 3.86 (2H, t, $^3J_{HH} = 6.9$ Hz,

20

25

30

NCH₂CH₂CO₂Et), 2.59 (2H, t, ³J_{HH} = 6.9 Hz, NCH₂CH₂CO₂Et), 1.09 (3H, t, ³J_{HH} = 6.9 Hz, CO₂CH₂CH₃). ¹³C NMR (CDCl₃) 171.4, 165.7, 164.5, 151.8, 139.8, 137.8, 132.3, 128.3, 127.2, 125.9, 125.0, 123.8, 122.9, 117.8, 115.5, 61.1, 46.3, 41.3, 32.9, 14.0. Mass spec. (FAB M+H⁺) calc'd 446.1, found 446.1. Exact mass (FAB M+H⁺)C₂₂H₂₀N₇O₄ calc'd 446.1577, found 446.1561.

5

b) 4-H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanobenzoyl]amino]-6-oxo, ethyl ester (93 mg, 0.17 mmol) was dissolved in anhydrous pyridine (2 ml)/ triethylamine (1 ml), transferred to a pressure tube, and the pyridine/triethylamine solution saturated with hydrogen sulfide. The vessel was sealed, heated to 50°C and stirred for 90 min. After cooling to room temperature, the reaction mixture was concentrated *in vacuo* and transferred to a separatory funnel containing ethyl acetate (30 ml) and water (30 ml). After thorough mixing, the aqueous phase was removed and the organic phase was washed with water (20 ml), brine (30 ml), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give 60 mg (76 % yield) 4-H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-thioamidobenzoyl]amino]-6-oxo, ethyl ester. TLC R_f = 0.55 (SiO₂, ethyl acetate, uv positive). Analytical HPLC Dynamax 300 A, C8 83-303-C5, elution gradient 30-100% acetonitrile (0.1% TFA) over 10 minutes, retention time 4.27 min (UV @ 214 nm). ¹H NMR (CD₃OD, δ TMS) 8.46 (1H, d, ⁴J_{HH} = 2.4 Hz, C7 ArH), 8.26 (1H, dd, ⁴J_{HH} = 2.4 Hz, ³J_{HH} = 9.0 Hz, C9 ArH), 8.00 (2H, s, *o*-H₂NCS ArH), 7.99 (2H, s, *m*-H₂NCS ArH), 7.96 (1H, d, ³J_{HH} = 9.0 Hz, C10 ArH), 4.94 (2H, s, NCNCH₂N(R)CO), 4.05 (2H, q, ³J_{HH} = 7.2 Hz, CO₂CH₂CH₃), 3.97 (2H, t, ³J_{HH} = 6.3 Hz, NCH₂CH₂CO₂Et), 2.69 (2H, t, ³J_{HH} = 6.3 Hz, NCH₂CH₂CO₂Et), 1.09 (2H, t, ³J_{HH} = 7.2 Hz, CO₂CH₂CH₃). Mass spec. FAB (M+H⁺) calc'd 480, found 480. Exact mass (FAB M+H⁺)C₂₂H₂₂N₇SO₄ calc'd 480.1454, found 480.1450.

25

c) Treated 4-H[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-thioamidobenzoyl]amino]-6-oxo, ethyl ester (0.21 gm, 0.47 mmol) with methyl iodide (20 ml) and addition of N-methyl pyrrolidinone (2 ml) was necessary to achieve a homogeneous solution (the thioamide is sparingly soluble in most typical organic solvents). The reaction was stirred overnight at room temperature. The crude reaction was concentrated *in vacuo* to a volume of 2 ml, taking care to avoid exposure to moisture. Chloroform (5 ml) was added and removed *in vacuo* to chase off traces of methyl iodide and the resulting crude residue placed under high vacuum for 15 minutes. The crude methyl iodide adduct was dissolved in ethanol (3 ml), saturated with anhydrous ammonium acetate (decanted from the 3A sieves and undissolved ammonium acetate), stoppered and stirred overnight at room temperature (anhydrous ammonium acetate was prepared by stirring ammonium acetate (1.2 gm, 15.0 mmol) in anhydrous ethanol (9 ml) with freshly activated 3A sieves for 16h at room temperature). The reaction mixture was concentrated *in vacuo* and the crude product purified


35

by prep RP HPLC (RP-Dynamax 60A C18 83-221-C, acetonitrile/water/0.1% TFA, gradient elution 0 % acetonitrile for 20 minutes, 0 to 50 % acetonitrile in 60 minutes, retention time 46-52 minutes, UV @ 254 nm) to give 4-*H*-[1,2,3,4]tetrazolo[4,3-*a*][1,4]benzodiazepine-5(6*H*)-propanoic acid, 8-[[4-(iminoaminomethyl)benzoyl]amino]-6-oxo, ethyl ester, trifluoroacetate (24.5 mg, 24% yield). Analytical HPLC Dynamax 300 A, C8 83-303-C5, isocratic elution @ 35% acetonitrile (0.1% TFA) over 6 minutes, retention time 2.78 min (UV @ 214 nm). ¹H NMR (DMSO-*d*₆, δTMS) 11.0 (1H, bs, ArCONHAr), 8.46 (1H, d, ⁴J_{HH} = 2.4 Hz, C7 ArH), 8.31 (1H, dd, ³J_{HH} = 8.7 Hz, ⁴J_{HH} = 2.4 Hz, C9 ArH), 8.17 (2H, d, ³J_{HH} = 8.4 Hz, *o*-H₂NCNH ArH), 8.06 (2H, d, ³J_{HH} = 8.4 Hz, *m*-H₂NCNH ArH), 7.96 (1H, d, ³J_{HH} = 8.7 Hz, C10 ArH), 4.97 (2H, br s, NCNCHHN(R)CO), 3.95 (2H, q, ³J_{HH} = 7.0 Hz, CO₂CH₂CH₃), 3.86 (2H, t, ³J_{HH} = 7.0 Hz, NCH₂CH₂CO₂Et), 2.59 (2H, t, ³J_{HH} = 7.5 Hz, NCH₂CH₂CO), 2.59 (2H, t, ³J_{HH} = 7.0 Hz, NCH₂CH₂CO₂Et), 1.09 (3H, t, ³J_{HH} = 7.0 Hz, CO₂CH₂CH₃) Mass spec. FAB (M+H⁺) calc'd 463.1 found 463.1. Exact mass FAB (M+H⁺) C₂₂H₂₃N₈O₄ calc'd 463.1842, found 463.1835.

15

d) 4-*H*-[1,2,3,4]tetrazolo[4,3-*a*][1,4]benzodiazepine-5(6*H*)-propanoic acid, 8-[[4-(iminoaminomethyl)benzoyl]amino]-6-oxo, ethyl ester, trifluoroacetate (7.0 mg, 0.015 mmol) was dissolved in THF:MeOH:H₂O (3:2:1, 0.25ml:17ml:0.08 ml, total volume = 0.5 ml) and treated with 1.0 N NaOH (15 μl, 0.015mmol). The reaction mixture was stirred at room temperature for 2 hr and additional 1.0 N NaOH was added (15 μl, 0.015mmol). After stirring additionally for 16h at room temperature another one equivalent of 1.0N NaOH (15 μl, 0.015mmol) was added and the reaction mixture heated to 60° C for 1 hr. After cooling to room temperature, the reaction mixture was acidified (neat TFA), and purified by prep RP HPLC (RP-Dynamax 60A C18 83-221-C, acetonitrile/water/0.1% TFA, gradient elution 0 % acetonitrile for 20 minutes, 0 to 30 % acetonitrile in 60 minutes, retention time 63-64 minutes, UV @ 254 nm) to give 4-*H*-[1,2,3,4]tetrazolo[4,3-*a*][1,4]benzodiazepine-5(6*H*)-propanoic acid, 8-[[4-(iminoaminomethyl)benzoyl]amino]-6-oxo, trifluoroacetate (1.7 mg, 17% yield). Analytical HPLC Dynamax 300 A, C8 83-303-C5, isocratic elution @ 15% acetonitrile (0.1% TFA) over 10 minutes, retention time 4.07 min (UV @ 214 nm). ¹H NMR (D₂O, δTMS) 8.23 (1H, apparent t due to two sets of overlapping doublets in a 1:1 ratio, ⁴J_{HH} = 2.1 Hz, C7 ArH), 8.09 (1H, d of apparent t, ³J_{HH} = 9.0 Hz, ⁴J_{HH} = 2.1 Hz, C9 ArH), 8.06, 8.05 (2H, d, ³J_{HH} = 8.7, 8.4 Hz, *o*-H₂NCNH ArH), 7.98, 7.97 (2H, d, ³J_{HH} = 9.3, 8.7 Hz, C10 ArH), 7.86, 7.85 (2H, d, ³J_{HH} = 8.10 Hz, *m*-H₂NCNH), 4.98 (2H, br s, NCNCHHN(R)CO), 3.97 (2H, t, ³J_{HH} = 6.0 Hz, NCH₂CH₂CO₂H), 2.71 (2H, t, ³J_{HH} = 6.0 Hz, NCH₂CH₂CO₂H). Mass spec. FAB (M+H⁺) calc'd 435.1 found 435.1. Exact mass FAB (M+H⁺) C₂₀H₁₉N₈O₄ calc'd 435.1529, found 435.1509. F_g IIbIIIa ELISA IC₅₀ = 1.2 nM, PA (citrate) IC₅₀ = 23 nM.


Example 13



5 *4H-[1,2,3,4] tetrazolo [4,3a][1,4] benzodiazepine- 5 (6H)-propanoic acid, 8-[[4-(hydroxyiminoaminomethyl)-benzoyl] amino]- 6-oxo, ethyl ester*

a) *4H-[1,2,3,4]tetrazolo[4,3a][1,4] benzodiazepine-5(6H)-propanoic acid, 8-[[4-(hydroxyiminoaminomethyl)-benzoyl]amino]-6-oxo, ethyl ester* was prepared by the method described in Example 11, part (a). *4-H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-cyanobenzoyl] amino]-6-oxo, ethyl ester* (see Example 12, part (a), for preparation, 64.5 mg, 0.115 mmol) yielded the desired hydroxyamidine/tetrazole (29 mg, 53% yield) after treatment with hydroxylamine hydrochloride (17.1 mg, .246 mmol)/NaOEt (14.8 mg, .217 mmol) in EtOH (60°C) and purification by prep RP HPLC (RP-Dynamax 60A C18 83-221-C, acetonitrile/water/0.1% TFA, gradient elution 5 % acetonitrile to 31 % acetonitrile in 60 minutes, retention time 30 - 44 minutes, UV @ 254 nm). Analytical HPLC Dynamax 300 A, C8 83-303-C5, isocratic elution @ 35% acetonitrile (0.1% TFA) over 10 minutes, retention time 2.90 min (UV @ 214 nm). ¹H NMR (DMSO-d₆) 10.78 (1H, brs, ArCONHAr), 9.85 (1H, brs, HONHCNH), 8.48 (1H, brs, C7 ArH), 8.23 (1H, br dd, ⁴J_{HH} = 1.5 Hz, ³J_{HH} = 8.7 Hz, C9 ArH), 8.02 (2H, d, ³J_{HH} = 8.10 Hz, *o*-HONHCNH ArH), 7.95 (2H, d, ³J_{HH} = 8.70 Hz, C10 ArH), 7.86 (2H, d, ³J_{HH} = 9.0 Hz, *m*-HONHCNH ArH), 5.94 (br s, 1H, HONHCNH), 4.97 (br s, 2H, ArNCNCHH N(R)CO), 3.93 (2H, q, ³J_{HH} = 6.9 Hz, CO₂CH₂CH₃), 3.86 (2H, t, ³J_{HH} = 6.9 Hz, NCH₂CH₂CO₂Et), 2.59 (2H, t, ³J_{HH} = 6.9 Hz, NCH₂CH₂CO₂Et), 1.09 (3H, t, ³J_{HH} = 6.9 Hz, CO₂CH₂CH₃). Mass spec. FAB (M+H⁺) C₂₂H₂₃N₈O₅ calc'd 479.0 found 479.0. Exact mass FAB (M+H⁺) calc'd 479.1791, found 479.1796.

Example 14




5 *4H-[1,2,3,4] tetrazolo [4,3a][1,4] benzodiazepine- 5 (6H)-propanoic acid, 8-[[4- [N-(ethylcarbamoyl)iminoaminomethyl]-benzoyl] amino] - 6-oxo, ethyl ester*

- a) A THF/H₂O (2:1, 0.2ml: 0.1 ml) solution of 4-H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4- (iminoaminomethyl)benzoyl]amino]-6-oxo, ethyl ester, trifluoroacetate (mono) (Example 12, part (c), 36.2 mg, 0.062 mmol) was treated with sodium bicarbonate (19.6 mg, 0.234 mmol) and ethyl chloroformate (9.32 μ l, 0.098 mmol). After stirring at room temperature for 45 minutes, analytical HPLC analysis (Dynamax 300 A, C8 83-303-C5, isocratic elution @ 35% acetonitrile (0.1% TFA) over 10 minutes) indicated complete consumption of the amidino ester. The THF/H₂O solution was transferred to a separatory funnel and shaken with ethyl acetate and water. The layers were separated and the organic layer washed with water (2x), brine, dried (MgSO₄), filtered, and concentrated *in vacuo* to afford 4H-[1,2,3,4]tetrazolo[4,3a][1,4] benzodiazepine-5(6H)-propanoic acid, 8-[[4-[N-(ethylcarbamoyl) iminoaminomethyl]-benzoyl] amino] - 6-oxo, ethyl ester (27 mg , 79% based on the amidinoester being a mono TFA salt). Analytical HPLC (Dynamax 300 A, C8 83-303-C5, isocratic elution @ 35% acetonitrile (0.1% TFA) over 6 minutes, retention time 2.90 min (UV @ 214 nm). ¹H NMR (CDCl₃/CD₃OD, δ TMS) 8.39 (1H, dd, ³J_{HH}= 9.0 Hz, ⁴J_{HH} = 2.4 Hz, C9 ArH), 8.26 (1H, d, ⁴J_{HH} = 2.4 Hz, C7 ArH), 8.01(2H, d, ³J_{HH}= 8.7 Hz, *o* -RO₂CHNHCNHAr), 7.97 (2H, d, ³J_{HH}= 8.7 Hz, *m* -RO₂CHNHCNHAr), 7.96 (1H, d, ³J_{HH} = 9.0 Hz, C10 ArH), 4.89 (2H, br s, NCNCHHN(R)CO), 4.24 (2H, q, ³J_{HH} = 7.0 Hz, H₃CCH₂OCONH), 4.15 (2H, q, ³J_{HH} = 7.0 Hz, CO₂CH₂CH₃), 3.96 (2H, t, ³J_{HH} = 6.3 Hz, NCH₂CH₂CO₂Et), 2.73 (2H, t, ³J_{HH} = 6.3 Hz, NCH₂CH₂CO₂Et), 1.37 (3H, t, ³J_{HH} = 7.5 Hz, H₃CCH₂OCONH), 1.25 (3H, t, ³J_{HH} = 7.0 Hz, CO₂CH₂CH₃). ¹³C NMR (CDCl₃/CD₃OD) 171.49, 166.32, 165.69, 151.86, 140.18, 137.59, 137.22, 127.69, 127.61, 127.19, 125.62, 125.18, 123.43, 122.75, 61.28, 60.99, 41.04, 32.76, 14.13, 13.81. Mass spec. FAB

(M+H⁺) C₂₅H₂₇N₈O₆ calc'd 535.2 found 535.2. Exact mass FAB (M+H⁺) calc'd 535.2053, found 535.2056.

5


Example 15

ELISA and platelet aggregation assay results

10

O-L	IC ₅₀ (μM)	IC ₅₀ (μM)	IC ₅₀ (μM)
	Eg/IIbIIIa	PA (citrate)	PA (heparin)
H ₂ N(CH ₂) ₄ CC	0.057	8.29	ND
<i>p</i> -H ₂ N(NH)CC ₆ H ₄ CC	0.004	0.093	0.24
<i>p</i> -H ₂ N(NH)CC ₆ H ₄ CH ₂ O	0.009	0.133	0.22
<i>p</i> -H ₂ N(NH)CC ₆ H ₄ CONH	0.009	0.086	ND


15



20


<u>Q-L</u>	IC ₅₀ (μ M)	IC ₅₀ (μ M)
	PA (citrate)	PA (heparin)
<i>p</i> -H ₂ N(NH)CC ₆ H ₄ CH ₂ O	0.54	1.7

5



10

<u>Q-L</u>	IC ₅₀ (μ M)	IC ₅₀ (μ M)
	PA (citrate)	PA (heparin)
<i>p</i> -H ₂ N(NH)CC ₆ H ₄ CH ₂ O	0.005	0.086





15

<u>Q-L</u>	IC ₅₀ (μ M)	IC ₅₀ (μ M)
	PA (citrate)	PA (heparin)
<i>p</i> -H ₂ N(NH)CC ₆ H ₄ CH ₂ O	0.116	ND
<i>p</i> -H ₂ N(NH)CC ₆ H ₄ CC	0.449	ND
<i>p</i> -H ₂ N(NH)CC ₆ H ₄ CONH	0.025	ND

20

What is claimed is:

- 1) A compound represented by structural Formula I or II:



where

- 10 R¹ and R² are each one or two groups independently selected from
hydrogen,
halo(F, Cl, Br, I),
cyano,
carboxy,
15 aminocarbonyl,
carboxamido,

- carbamoyloxy,
 formyloxy,
 formyl,
 azido,
 5 nitro,
 imidazolyl,
 ureido,
 thioureido,
 thiocyanato,
 10 hydroxy,
 mercapto,
 sulfonamido, and
 an optionally substituted radical selected from
- 15 C₁-C₁₂alkyl,
 C₂-C₁₂alkenyl,
 C₃-C₁₂alkynyl,
 C₃-C₁₂cycloalkyl,
 C₆-C₁₄aryl,
 C₆-C₁₀arylC₁-C₈alkyl,
 20 C₁-C₁₂alkyloxy,
 C₁-C₁₂alkyloxyC₁-C₁₂alkyl,
 C₆-C₁₄aryloxy,
 C₁-C₁₂acylamino,
 N,N-di(C₁-C₁₂)acylamino,
 25 N-(C₁-C₁₂)alkyl-N-(C₁-C₁₂)alkylsulfonylamino,
 C₁-C₁₂alkylthiocarbonyl,
 C₁-C₁₂alkylthio,
 C₁-C₁₂alkylthioC₁-C₁₂alkyl,
 C₁-C₁₂alkylsulfinyl,
 30 C₁-C₁₂alkylsulfinylC₁-C₁₂alkyl,
 C₁-C₁₂alkylsulfonyl,
 C₁-C₁₂alkylsulfonylC₁-C₁₂alkyl,
 C₁-C₁₂alkylsulfonato,
 N-(C₁-C₁₂)alkylsulfonamido,
 35 N,N-di-(C₁-C₁₂)sulfonamido,
 N-(C₁-C₁₂)alkyl-N-thioformylamino,
 C₁-C₁₂thioacylamino,
 N-(C₁-C₁₂)alkyl-N-(C₁-C₁₂)thioacylamino,

C₁-C₁₂alkylsulfinamido,
 N-(C₁-C₁₂)alkyl-N-(C₁-C₁₂)alkylsulfinylamino,
 C₁-C₁₂carbalkoxy,
 C₁-C₁₂alkylcarbonyl,
 5 C₁-C₁₂alkyloxycarbonyl,
 C₁-C₁₂alkanoyloxy,
 N-(C₁-C₁₂)alkylcarboxamido,
 N-(C₁-C₁₂)alkylaminocarbonyl,
 10 N,N-di-(C₁-C₁₂)carboxamido,
 N,N-di-(C₁-C₁₂)alkylaminocarbonyl,
 N-(C₁-C₁₂)alkylcarbamoyloxy,
 N,N-di-(C₁-C₁₂)carbamoyloxy, and
 heterocycloalkyl or heteroaryl having from 1 to 3 rings, each ring having
 15 from 5 to 7 atoms with from 0-3 heteroatoms selected from N, O, and
 S, provided that at least one ring contains a heteroatom,
 where the substituents are one to three groups selected from
 halo (F, Cl, Br, I),
 amino,
 20 amidino,
 guanidino,
 imidazolyl,
 indolyl,
 cyano,
 azido,
 25 nitro,
 hydroxy,
 mercapto,
 morpholino,
 morpholinyl,
 30 piperazinyl,
 piperidinyl,
 pyrrolinyl,
 sulfonamido,
 ureido,
 35 thioureido,
 carboxy,
 aminocarbonyl,
 C₁-C₄aminocarbonyl,

5 carboxamido,
 carbamoyloxy,
 formyloxy,
 formyl,
 C₁-C₄alkyloxycarbonyl,
 C₁-C₄alkyl,
 C₁-C₄alkylamino,
 di-(C₁-C₄)alkylamino,
 C₁-C₄alkoxy,
 10 hydroxyphenyl,
 phenyl, and
 phenoxy,

optionally, R¹ and R² when bonded to adjacent carbon atoms may join to
 form a fused unsubstituted or substituted C₆-C₁₀aryl ring, where the
 15 substituents are selected from
 halo (F, Cl, Br, I),
 cyano,
 azido,
 nitro,
 20 hydroxy,
 mercapto,
 sulfonamido,
 ureido,
 thioureido,
 25 carboxamido,
 carbamoyloxy,
 formyloxy,
 formyl,
 C₁-C₄alkyl,
 30 haloC₁-C₄alkyl,
 C₁-C₄alkoxy,
 phenyl, and
 phenoxy;

Q is selected from the group consisting of
 35 an amino group selected from
 NH₂,
 NR³H,
 NR³R⁴, and

NR³R⁴R⁵,

R³, R⁴, and R⁵ are independently selected from

hydrogen,

cyano,

5

NR⁶R⁷,

C(=NR⁸)-NR⁶R⁷,

N=CR⁹-NR⁶R⁷,

NR¹⁰-CR⁹=NR⁸,

NR¹⁰-C(=NR⁸)-NR⁶R⁷,

10

halo(F, Cl, Br, I)-C₁-C₄alkyl, and

an optionally substituted group selected from

C₁-C₁₂alkyl,

C₂-C₇alkenyl,

C₃-C₁₂cycloalkyl,

15

C₅-C₁₂cycloalkenyl,

C₆-C₁₄aryl,

C₁-C₆ alkylC₆-C₁₄aryl,

C₂-C₆alkenylC₆-C₁₀aryl,

heterocyclyl,

20

C₁-C₆alkyl-heterocyclyl,

C₁-C₈alkoxy,

C₁-C₈alkoxycarbonyl,

C₆-C₁₄aryloxy,

C₆-C₁₄aryloxycarbonyl, and

25

C₆-C₁₄arylC₁-C₆alkyloxycarbonyl,

where each R⁶, R⁷, R⁸, R⁹, and R¹⁰ is independently selected from

hydrogen,

cyano,

C₁-C₆alkyl, and

30

R¹¹,

and where the substituents are one to three R¹¹,

R¹¹ is selected from optionally substituted

C₆-C₁₂aryloxy,

C₆-C₁₂arylamino,

35

C₆-C₁₂aroyl,

C₁-C₈alkoxy,

C₁-C₈alkoxyalkyl,

- C₁-C₈alkoxyalkyloxy,
 C₁-C₈alkoxycarbonyl,
 C₁-C₈alkylcarbonyl,
 C₁-C₈aralkylcarbonyl,
 5 C₁-C₈alkylthiocarbonyl,
 C₆-C₁₂aralkylthiocarbonyl,
 C₂-C₈alkoxythiocarbonyl,
 C₆-C₁₂aryl,
 C₁-C₄alkanoylamino,
 10 C₁-C₆alkoxycarbonyl-C₀-C₆alkylamino,
 C₁-C₈alkylsulfonylamino,
 C₆-C₁₀aralkylsulfonylamino,
 C₆-C₁₀aralkyl,
 C₆-C₁₀alkaryl,
 15 C₁-C₈alkylthio,
 C₆-C₁₀aralkylthio,
 C₁-C₈alkylsulfinyl,
 C₆-C₁₀aralkylsulfinyl,
 C₁-C₈alkylsulfonyl,
 20 C₆-C₁₀aralkylsulfonyl,
 C₁-C₈alkylaminosulfonyl,
 C₆-C₁₀aralkylsulfonylamino,
 C₂-C₈alkenyl, and
 5- or 6-member heterocycle containing 1-4 hetero atoms
 25 selected from N, O, and S,
 where the substituents are one to three R¹²,
 R¹² is selected from
 nitro,
 amino,
 30 C₁-C₈alkylamino,
 di-(C₁-C₈)alkylamino,
 amidino,
 aminomethyleneimino,
 imino,
 35 iminoC₁-C₄alkyl,
 iminomethyleneamino,
 guanidino,
 C₆-C₁₀arylamino,










- 5 C₁-C₈acylamino,
C₁-C₄alkylsulfonamino,
azido,
cyano,
hydroxy,
hydroxyC₁-C₈alkyl,
C₁-C₈alkoxy,
phenyloxy,
C₁-C₈alkanoyloxy,
10 C₁-C₈alkanoyl,
C₆-C₁₂aroyl,
benzamido,
phenyl,
halo(F, Cl, Br, I),
15 haloC₁-C₈alkyl, and
C₁-C₈alkyl,
aminosulfonyl,
oxo,
thio,
20 thiocarbonyl,
hydroxy,
mercapto,
formyl,
formyloxy,
25 carboxy,
ureido,
glycyl,
phthalimido,
succinimido,
30 morpholino,
C₃-C₇cycloalkyl;
optionally R³ and R⁴ taken together may form optionally substituted
tetramethylene,
pentamethylene,
35 3-oxopentamethylene, and
3-azapentamethylene,

where the substituents are selected from one to three R¹²;
an amidino group selected from


$C(=NH)-NH_2$,
 $C(=NH)-NHR^3$,
 $C(=NR^4)-NHR^3$,
 $C(=NH)-NR^3R^4$, and
 $C(=NR^5)-NR^3R^4$,

where R^3 , R^4 , and R^5 are defined above or are selected from the group

$HO-$,
 $C_1-C_{10}alkyl-O-(C=O)-$,
 $C_1-C_{10}alkyl-S-(C=O)-$,
 $Cl_3CCH_2O-(C=O)-$,



, and



an aminoalkyleneimino group selected from

- 5
- $\text{N}=\text{CH}-\text{NH}_2$,
 - $\text{N}=\text{CH}-\text{NHR}^3$,
 - $\text{N}=\text{CH}-\text{NR}^3\text{R}^4$, and
 - $\text{N}=\text{CR}^5-\text{NR}^3\text{R}^4$,

where R^3 , R^4 , and R^5 are defined above;

an iminoalkyleneamino group selected from

- 10
- $\text{NH}-\text{CH}=\text{NH}$,
 - $\text{NH}-\text{CH}=\text{NR}^3$,
 - $\text{NH}-\text{CR}^4=\text{NR}^3$, and
 - $\text{NR}^5-\text{CR}^4=\text{NR}^3$,


where R^3 , R^4 , and R^5 are defined above;

a guanidino group selected from


- 15
- $\text{NH}-\text{C}(=\text{NH})-\text{NH}_2$,
 - $\text{NH}-\text{C}(=\text{NH})-\text{NR}^3\text{H}$,
 - $\text{NH}-\text{C}(=\text{NH})-\text{NR}^3\text{R}^4$,
 - $\text{NH}-\text{C}(=\text{NR}^5)-\text{NR}^3\text{R}^4$,
 - $\text{NR}^3-\text{C}(=\text{NR}^3)-\text{NR}^3\text{R}^4$,
 - 20 $\text{NR}^3-\text{C}(=\text{NH})-\text{NR}^3\text{R}^4$,
 - $\text{NR}^3-\text{C}(=\text{NR}^3)-\text{NH}_2$,
 - $\text{NR}^3-\text{C}(=\text{NH})-\text{NH}_2$,
 - $\text{NR}^3-\text{C}(=\text{NR}^3)-\text{NHR}^4$, and
 - $\text{NR}^3-\text{C}(=\text{NH})-\text{NHR}^4$,

25 where R^3 , R^4 , and R^5 are defined above;

an optionally substituted saturated heterocyclic group selected from



30



where n is 0, 1, 2, or 3,

5 R^{13} is selected from R^6 ,

$CR^9=NR^8$,

$CR^9(=NR^8)-NR^6R^7$,

$C(=NR^8)-NR^6R^7$,

$N=CR^9-NR^6R^7$,


10 $NR^{10}-CR^9=NR^8$, and

$NR^{10}-(C=NR^8)-NR^6R^7$,

where R^6 - R^{10} are defined above, Z^2 is O, S, or NR^{13} , and the substituents are independently one to three R^{12} ;

an optionally substituted unsaturated (nonaromatic) heterocyclyl selected from

15



20

where







m is 1, 2, or 3,

Z^2 and R^{13} are defined above, and

25

the substituents are independently one to three R^{12} ;

an optionally substituted unsaturated (aromatic) heterocyclyl selected from



where Z^3 , Z^4 , and Z^5 are selected from O, S, N, and NR^6 , provided that at least one Z^3 , Z^4 , or Z^5 is N or NR^6 ,

R^{13} is defined above, and the substituents are independently one to three R^{12} ,

L is an optionally substituted bivalent radical selected from the group

C3-C7-alkylene,

C3-C7-cycloalkylene,




C3-C7-alkenylene,










C3-C7-alkadienylene,



C3-C7-alkynylene,

- C₄-C₇-alkadiynylene,
 C₄-C₇-alkenynylene,
 C₆-C₁₄-arylene,
 C₆-C₁₄-aryl-C₂-C₄-alkynylene,
 5 C₁-C₃-alkyl-C₆-C₁₄-aryl-C₂-C₄-alkynylene,
 C₆-C₁₄-aryl-C₂-C₄-alkenylene,
 C₁-C₃-alkyl-C₆-C₁₄-arylene,
 C₁-C₃-alkyl-C₆-C₁₄-aryl-C₂-C₄-alkenylene,
 C₆-C₁₄-aryl-C₁-C₃-alkylene,
 10 C₆-C₁₄-aryl-C₁-C₃-alkyloxyene,
 C₁-C₂-alkyl-C₆-C₁₄-aryl-C₁-C₂-alkylene,
 C₁-C₃-alkyloxy-C₆-C₁₄-arylene,
 C₂-C₆-alkyloxyene,
 C₁-C₅-alkyloxy-C₁-C₅-alkylene,
 15 C₆-C₁₀-aryloxyene,
 C₆-C₁₀-aryloxy-C₁-C₅-alkylene,
 C₂-C₆-alkylthioene,
 C₁-C₅-alkylthio-C₁-C₅-alkylene,
 C₆-C₁₀-arylthioene,
 20 C₆-C₁₀-arylthio-C₁-C₅-alkylene,
 C₁-C₅-alkylsulfoxide-C₁-C₅-alkylene,
 C₁-C₅-alkylsulfone-C₁-C₅-alkylene,


25








5



10



where

15 R^{14} is selected from

a chemical bond,

C₁-C₈-alkyl,C₁-C₈-alkyloxy,C₃-C₇-cycloalkyl,

20

C₂-C₅-alkenyl,C₃-C₅-alkynyl,C₆-C₁₀-aryl,C₁-C₃-alkyl-C₆-C₁₂-aryl,C₁-C₂-alkyl-C₆-C₁₀-aryl-C₁-C₂-alkyl,

25

C₆-C₁₀-aryl-C₁-C₂-alkyl,C₆-C₁₀-aryloxy-C₁-C₂-alkyl,

HET, and

piperizinyl;

 R^{15} is selected from

30

a chemical bond,

C₁-C₄-alkyl,

C₂-C₄-alkenyl,
 C₂-C₄-alkynyl,
 C₆-C₁₀-aryl, and
 C₁-C₃-alkyl-C₆-C₁₂-aryl;

5 R¹⁶ is selected from

a chemical bond,
 C₁-C₅-alkyl,
 C₃-C₇-cycloalkyl,
 C₃-C₅-alkenyl,
 10 C₃-C₅-alkynyl,
 C₆-C₁₀-aryl,
 C₁-C₃-alkyl-C₆-C₁₂-aryl,
 C₆-C₁₀-aryl-C₁-C₂-alkyl,
 HET, and
 15 pyridyl;

R¹⁷ is selected from

C₃-C₄-alkenyl,
 C₃-C₄-alkynyl,
 C₆-C₁₀-aryl,
 20 benzyl, and
 HET,

where HET is a heterocycle, saturated or unsaturated, having from 5-14 atoms in from
 1-3 cycle(s), each cycle having from 1-3 heteroatoms selected from N, O, and
 S,

25 and where the substituents are selected from one to three R¹²;

R¹⁸, R¹⁹, R²⁰ and R²¹ are independently selected from

hydrogen,
 C₁-C₄alkyl,
 halo(F, Cl, Br, I)C₁-C₄alkyl,
 30 C₁-C₄alkoxy,
 halo(F, Cl, Br, I),
 cyano,
 carboxy,
 hydroxy,
 35 C₁-C₄alkoxycarbonyl, and
 C₁-C₄alkylsulfonylC₀-C₃alkyl;

R²² is selected from the group consisting of
 hydroxy,

- C₁-C₈-alkoxy,
 C₃-C₁₂-alkenoxy,
 C₆-C₁₂-aryloxy,
 C₁-C₆-alkyl-C₆-C₁₂-aryloxy,
 5 di-C₁-C₈-alkylamino-C₁-C₈-alkoxy,
 acylamino-C₁-C₈-alkoxy selected from the group
 acetylaminooethoxy,
 nicotinoylaminoethoxy, and
 succinamidoethoxy,
 10 C₁-C₈-alkoxyloxy-C₁-C₈-alkoxy, and
 C₆-C₁₂-aryl-C₁-C₈-alkoxy,
 where any aryl groups are unsubstituted or substituted with
 one to three of the groups
 nitro,
 15 halo (F, Cl, Br, I),
 C₁-C₄-alkoxy, and
 amino,
 hydroxy-C₂-C₈-alkoxy,
 dihydroxy-C₃-C₈-alkoxy,
 20 morpholino-C₂-C₈-alkoxy,
 morpholinyl-C₁-C₈-alkoxy, and
 NR²³R²⁴,
 R²³ and R²⁴ are independently selected from the group
 hydrogen,
 25 C₁-C₈-alkyl,
 C₃-C₈-alkenyl,
 C₆-C₁₂-aryl,
 C₆-C₁₂-aryl-C₁-C₈-alkyl
 where any aryl groups are unsubstituted or substituted with
 30 one to three of the groups
 nitro,
 halo (F, Cl, Br, I),
 C₁-C₄-alkoxy, and
 amino;
 35 A¹ is selected from
 CR¹,
 CHR²,
 N, and


- NR²⁵;
- A² is selected from
- CHR¹,
- CR²,
- 5 N,
- O=S=O,
- S=O,
- S,
- O,
- 10 C=O,
- C-OR²⁶, and
- C=N-R²⁵;
- B¹ is selected from
- CR¹,
- 15 CHR²,
- N,
- NR²⁵, and
- C=O;
- B² is selected from
- 20 CR²,
- CHR¹,
- NR²⁵,
- O=S=O,
- S=O,
- 25 S,
- O, and
- C=O;
- B³ is selected from
- CR¹,
- 30 CHR², and
- C=O;
- R²⁵ is selected from
- R⁶, and
- (CH₂)_mR¹, where m is 1, 2, or 3;
- 35 R²⁶ is selected from
- hydrogen,
- C₁-C₈-alkyl,

C₆-C₁₂-aryl, and
C₆-C₁₂-aryl-C₁-C₈-alkyl


where any aryl or alkyl groups are unsubstituted or substituted with one to three of the groups
nitro, and

5 halo (F, Cl, Br, I); and
pharmaceutically acceptable salts thereof.


2) The compound of Claim 1 represented by Formula III-VIII:




III




IV




V



VI



VII



VIII

where

R¹ and R² are independently selected from
hydrogen,

- halo(F, Cl, Br, I),
 cyano,
 carboxy,
 aminocarbonyl,
 5 carboxamido,
 carbamoyloxy,
 formyloxy,
 formyl,
 azido,
 10 nitro,
 imidazolyl,
 ureido,
 thioureido,
 thiocyanato,
 15 hydroxy,
 mercapto,
 sulfonamido, and
 an optionally substituted radical selected from
 C₁-C₁₂alkyl,
 20 C₁-C₁₂alkyloxycarbonyl,
 C₁-C₁₂alkyloxyC₁-C₁₂alkyl,
 C₂-C₁₂alkenyl,
 C₂-C₁₂alkynyl,
 C₃-C₁₂cycloalkyl,
 25 C₆-C₁₄aryl,
 C₆-C₁₀aryl-C₁-C₈alkyl,
 C₁-C₁₂alkyloxy,
 C₆-C₁₄aryloxy, and
 C₁-C₁₂acylamino,
 30 where the substituents are selected from
 halo (F, Cl, Br, I),
 amino,
 amidino,
 guanidino,
 35 imidazolyl,
 indolyl,
 cyano,
 azido,

WO 95/04057

PCT/US94/07989

- 5 nitro,
 hydroxy,
 mercapto,
 morpholino,
 morpholinyl,
 piperazinyl,
 piperidinyl,
 pyrrolinyl,
 sulfonamido,
10 ureido,
 thioureido,
 carboxy,
 aminocarbonyl,
 C₁-C₄aminocarbonyl,
15 carboxamido,
 carbamoyloxy,
 formyloxy,
 formyl,
 C₁-C₄alkyloxycarbonyl,
20 C₁-C₄alkyl,
 C₁-C₄alkylamino,
 di-(C₁-C₄)alkylamino,
 C₁-C₄alkoxy,
 C₁-C₄alkthio,
25 hydroxyphenyl,
 phenyl, and
 phenoxy;

Q is selected from the group consisting of
an amino group selected from



- 30 NH₂,
 NR³H,
 NR³R⁴, and
 NR³R⁴R⁵,

R³, R⁴, and R⁵ are independently an optionally substituted radical selected from

- 35 hydrogen,
 C₁-C₄ alkoxy,
 C₁-C₄ alkyl,
 halo(F, Cl, Br, I)C₁-C₄-alkyl,

- optionally substituted C₁-C₁₂ alkyl,
 optionally substituted C₃-C₁₂ cycloalkyl,
 optionally substituted C₆-C₁₄ aryl,
 optionally substituted C₁-C₆alkylC₆-C₁₄aryl,
 optionally substituted C₁-C₈ alkoxy,
 optionally substituted C₁-C₈ alkoxycarbonyl,
 optionally substituted C₆-C₁₄ aryloxy,
 optionally substituted C₆-C₁₄ aryloxycarbonyl,
 NR⁶R⁷,
 C(=NR⁸)-NR⁶R⁷,
 N=CR⁹-NR⁶R⁷,
 NR¹⁰-CR⁹=NR⁸, and
 NR¹⁰-C(=NR⁸)-NR⁶R⁷,
 where each R⁶, R⁷, R⁸, R⁹, and R¹⁰ is independently selected from
 hydrogen,
 C₁-C₄ alkyl, and
 R¹¹,
 and where the substituents are one to three R¹¹,
 R¹¹ is independently selected from
 optionally substituted C₆-C₁₂ aryloxy,
 optionally substituted C₆-C₁₂ arylamino,
 optionally substituted C₆-C₁₂ aroyl, and
 optionally substituted C₆-C₁₂ aryl,
 where the substituents are one to three R¹²,
 R¹² is selected from
 nitro,
 amino,
 C₁-C₈alkylamino,
 di-(C₁-C₈)alkylamino,
 amidino,
 aminomethyleneimino,
 imino,
 imino-C₁-C₄alkyl,
 iminomethyleneamino,
 guanidino,
 C₆-C₁₀arylamino,
 C₁-C₈acylamino,
 C₁-C₄alkylsulfonamino,

- azido,
 cyano,
 hydroxy,
 hydroxy-C₁-C₈alkyl,
 5 C₁-C₈alkoxy,
 phenyloxy,
 C₁-C₈alkanoyloxy,
 C₁-C₈alkanoyl,
 C₆-C₁₂aroyl,
 10 benzamido,
 phenyl,
 halo(F, Cl, Br, I),
 halo-C₁-C₈alkyl,
 C₁-C₈alkyl,
 15 aminosulfonyl,
 oxo,
 thio,
 thiocarbonyl,
 hydroxy,
 20 mercapto,
 formyl,
 formyloxy,
 carboxy,
 ureido,
 25 glycylyl,
 phthalimido,
 succinimido,
 morpholino, and
 C₃-C₇cycloalkyl;
 30 optionally R³ and R⁴ taken together may form optionally substituted
 tetramethylene,
 pentamethylene,
 3-oxopentamethylene, and
 3-azapentamethylene,
 35 where the substituents are selected from one to three R¹²,
 an amidino group selected from
 C(=NH)-NH₂,
 C(=NH)-NHR³,









where R^3 , R^4 , and R^5 are defined above or are selected from the group




5




10



15




, and




;

an aminoalkyleneamino group selected from




where R^3 , R^4 , and R^5 are defined above,

an iminoalkyleneamino group selected from



where R^3 , R^4 , and R^5 are defined above,

a guanidino group selected from



where R^3 , R^4 , and R^5 are defined above;

L is an optionally substituted bivalent radical selected from the group

C_3 - C_7 -alkylene,

C_3 - C_7 -cycloalkylene,

C_3 - C_7 -alkenylene,

C_4 - C_7 -cycloalkenylene,

C_5 - C_8 -cycloalkadienylene,

C_3 - C_7 -alkadienylene,

C_3 - C_7 -alkynylene,

C_4 - C_7 -alkenynylene,

5





10




15

20

25

C₆-C₁₄-arylene,
 C₆-C₁₄-aryl-C₂-C₄-alkynylene,
 C₁-C₃-alkyl-C₆-C₁₄-aryl-C₂-C₄-alkynylene,
 C₆-C₁₄-aryl-C₂-C₄-alkenylene,
 C₁-C₃-alkyl-C₆-C₁₄-arylene,
 C₁-C₃-alkyl-C₆-C₁₄-aryl-C₂-C₄-alkenylene,
 C₆-C₁₄-aryl-C₁-C₃-alkylene,
 C₆-C₁₄-aryl-C₁-C₃-alkyloxyene,
 C₁-C₂-alkyl-C₆-C₁₄-aryl-C₁-C₂-alkylene,
 C₁-C₃-alkyloxy-C₆-C₁₄-arylene,
 C₂-C₈-alkyloxyene,
 C₁-C₅-alkyloxy-C₁-C₅-alkylene,
 C₆-C₁₀-aryloxyene,
 C₆-C₁₀-aryloxy-C₁-C₅-alkylene,
 C₆-C₁₀-arylthio-C₁-C₅-alkylene,





5

where

10 R^{14} is selected from a chemical bond,C₁-C₈-alkyl,C₁-C₈-alkyloxy,C₃-C₇-cycloalkyl,C₂-C₅-alkenyl,

15

C₃-C₅-alkynyl,C₆-C₁₀-aryl,C₁-C₃-alkyl-C₆-C₁₂-aryl,C₁-C₂-alkyl-C₆-C₁₀-aryl-C₁-C₂-alkyl,C₆-C₁₀-aryl-C₁-C₂-alkyl,

20

HET, and

pyridyl;

 R^{15} is selected from a chemical bond,C₁-C₄-alkyl,C₂-C₄-alkenyl,

25

C₂-C₄-alkynyl, andC₆-C₁₀-aryl; R^{16} is selected from a chemical bond,C₁-C₅-alkyl,C₃-C₇-cycloalkyl,

30

C₃-C₅-alkenyl,C₃-C₅-alkynyl,C₆-C₁₀-aryl,C₁-C₃-alkyl-C₆-C₁₂-aryl,

- C₁-C₃-alkyl-C₆-C₁₂-aryl,
 C₆-C₁₀-aryl-C₁-C₂-alkyl,
 HET, and
 pyridyl;
- 5 R¹⁸, R¹⁹, R²⁰ and R²¹ are independently selected from
 hydrogen,
 C₁-C₄alkyl, and
 halo(F, Cl, Br, I)C₁-C₄alkyl,
- R²² is selected from the group consisting of
- 10 hydroxy,
 C₁-C₈-alkoxy,
 C₃-C₁₂-alkenoxy,
 C₆-C₁₂-aryloxy,
 C₁-C₆-alkyl-C₆-C₁₂-aryloxy,
- 15 di-C₁-C₈-alkylamino-C₁-C₈-alkoxy,
 acylamino-C₁-C₈-alkoxy selected from the group
 acetylaminooethoxy,
 nicotinoylaminoethoxy, and
 succinamidoethoxy,
- 20 C₁-C₈-alkoxyloxy-C₁-C₈-alkoxy,
 C₆-C₁₂-aryl-C₁-C₈-alkoxy where the aryl group is unsubstituted or
 substituted with one to three of the groups
 nitro,
- 25 halo (F, Cl, Br, I),
 C₁-C₄-alkoxy, and
 amino,
 hydroxy-C₂-C₈-alkoxy,
 dihydroxy-C₃-C₈-alkoxy,
 morpholino-C₂-C₈-alkoxy, and
- 30 NR²³R²⁴,
- R²³ and R²⁴ are independently selected from
 hydrogen,
 C₁-C₈-alkyl,
 C₃-C₈-alkenyl,
- 35 C₆-C₁₂-aryl
- where the aryl group is unsubstituted or substituted with one to three of the groups
 nitro,
 halo (F, Cl, Br, I),

C₁-C₄-alkoxy, and
 amino, and
 C₆-C₁₂-aryl-C₁-C₈-alkyl

where the aryl group is unsubstituted or substituted with one to three of the groups

5 nitro,
 halo (F, Cl, Br, I), and
 C₁-C₄-alkoxy;

R²⁵ is selected from

 R⁶, and
 10 (CH₂)_mR¹, where m is 1, 2, or 3;



R²⁶ is selected from

 hydrogen,
 C₁-C₈-alkyl,
 C₆-C₁₂-aryl, and
 15 C₆-C₁₂-aryl-C₁-C₈-alkyl


where any aryl or alkyl groups are unsubstituted or substituted with one to three of the groups

 nitro, and
 halo (F, Cl, Br, I); and
 20 pharmaceutically acceptable salts thereof.


3) The compound of Claim 1 represented by Formula IX-XX:




25




XI




XII




XIII




XIV




XV





XVI



XVII



XVIII



where

R¹ and R² are independently selected from

- 5 hydrogen,
- halo(F, Cl, Br, I),
- cyano,
- carboxy,
- aminocarbonyl,
- 10 carboxamido,
- carbamoyloxy,
- formyloxy,
- formyl,
- azido,
- 15 nitro,
- imidazolyl,
- ureido,
- thioureido,
- thiocyanato,
- 20 hydroxy,
- mercapto,
- sulfonamido, and
- an optionally substituted radical selected from
- C₁-C₁₂alkyl,
- 25 C₁-C₁₂alkyloxycarbonyl,
- C₁-C₁₂alkyloxyC₁-C₁₂alkyl,
- C₂-C₁₂alkenyl,
- C₂-C₁₂alkynyl,
- C₃-C₁₂cycloalkyl,
- 30 C₆-C₁₄aryl,
- C₆-C₁₀aryl-C₁-C₈alkyl,

C₁-C₁₂alkyloxy,
C₆-C₁₄aryloxy, and
C₁-C₁₂acylamino,

where the substituents are selected from

5 halo (F, Cl, Br, I),
 amino,
 amidino,
 guanidino,
 imidazolyl,
10 indolyl,
 cyano,
 azido,
 nitro,
 hydroxy,
15 mercapto,
 morpholino,
 morpholinyl,
 piperazinyl,
 piperidinyl,
20 pyrrolinyl,
 sulfonamido,
 ureido,
 thioureido,
 carboxy,
25 aminocarbonyl,
 C₁-C₄aminocarbonyl,
 carboxamido,
 carbamoyloxy,
 formyloxy,
30 formyl,
 C₁-C₄alkyloxycarbonyl,
 C₁-C₄alkyl,
 C₁-C₄alkylamino,
 di-(C₁-C₄)alkylamino,
35 C₁-C₄alkoxy,
 C₁-C₄alkthio,
 hydroxyphenyl,
 phenyl, and

phenoxy;

Q is selected from the group consisting of
an amino group selected from

- 5 NH₂,
 NR³H,
 NR³R⁴, and
 NR³R⁴R⁵,

R³, R⁴, and R⁵ are independently an optionally substituted radical selected from

- 10 hydrogen,
 C₁-C₄alkoxy,
 C₁-C₄alkyl,
 halo(F, Cl, Br, I)C₁-C₄alkyl,
 optionally substituted C₁-C₁₂alkyl,
15 optionally substituted C₃-C₁₂cycloalkyl,
 optionally substituted C₆-C₁₄aryl,
 optionally substituted C₁-C₆alkylC₆-C₁₄aryl,
 optionally substituted C₁-C₈alkoxy,
 optionally substituted C₁-C₈alkoxycarbonyl,
20 optionally substituted C₆-C₁₄aryloxy,
 optionally substituted C₆-C₁₄aryloxycarbonyl,
 NR⁶R⁷,
 C(=NR⁸)-NR⁶R⁷,
 N=CR⁹-NR⁶R⁷,
25 NR¹⁰-CR⁹=NR⁸, and
 NR¹⁰-C(=NR⁸)-NR⁶R⁷,

where each R⁶, R⁷, R⁸, R⁹, and R¹⁰ is independently selected from

- hydrogen,
 C₁-C₄ alkyl, and
30 R¹¹,

and where the substituents are one to three R¹¹,

R¹¹ independently selected from

- optionally substituted C₆-C₁₂aryloxy,
 optionally substituted C₆-C₁₂arylamino,
35 optionally substituted C₆-C₁₂aroyl,
 optionally substituted C₆-C₁₂aryl,

where the substituents are one to three R¹²,

R¹² is selected from



5 nitro,
 amino,
 C₁-C₈alkylamino,
 di-(C₁-C₈)alkylamino,
10 amidino,
 aminomethyleneimino,
 imino,
 imino-C₁-C₄alkyl,
 iminomethyleneamino,
15 guanidino,
 C₆-C₁₀arylamino,
 C₁-C₈acylamino,
 C₁-C₄alkylsulfonamino,
20 azido,
 cyano,
 hydroxy,
 hydroxy-C₁-C₈alkyl,
 C₁-C₈alkoxy,
 phenyloxy,
25 C₁-C₈alkanoyloxy,
 C₁-C₈alkanoyl,
 C₆-C₁₂aroyl,
 benzamido,
 phenyl,
30 halo(F, Cl, Br, I),
 halo-C₁-C₈alkyl,
 C₁-C₈alkyl,
 aminosulfonyl,
 oxo,
35 thio,
 thiocarbonyl,
 hydroxy,
 mercapto,
 formyl,
 formyloxy,
 carboxy,
 ureido,
 glycyl,



- phthalimido,
succinimido,
morpholino,
C₃-C₇cycloalkyl;
- 5 optionally R³ and R⁴ taken together may form optionally substituted
tetramethylene,
pentamethylene,
3-oxopentamethylene, and
3-azapentamethylene,
- 10 where the substituents are selected from one to three R¹²,
an amidino group selected from
C(=NH)-NH₂,
C(=NH)-NHR³,
C(=NR⁴)-NHR³,
15 C(=NH)-NR³R⁴, and
C(=NR⁵)-NR³R⁴,
where R³, R⁴, and R⁵ are defined above,
an aminoalkyleneamino group selected from
N=CH-NH₂,
20 N=CH-NHR³,
N=CH-NR³R⁴, and
N=CR⁵-NR³R⁴,
where R³, R⁴, and R⁵ are defined above,
an iminoalkyleneamino group selected from
25 NH-CH=NH,
NH-CH=NR³,
NH-CR⁴=NR³, and
NR⁵-CR⁴=NR³,
where R³, R⁴, and R⁵ are defined above,
- 30 a guanidino group selected from
NH-C(=NH)-NH₂,
NH-C(=NH)-NR³H,
NH-C(=NH)-NR³R⁴,
NH-C(=NR⁵)-NR³R⁴,
35 NR³-C(=NR³)-NR³R⁴,
NR³-C(=NH)-NR³R⁴,
NR³-C(=NR³)-NH₂,
NR³-C(=NH)-NH₂,

$\text{NR}^3\text{-C(=NR}^3\text{)-NHR}^4$, and
 $\text{NR}^3\text{-C(=NH)-NHR}^4$,
 where R^3 , R^4 , and R^5 are defined above;


- 5 L is an optionally substituted bivalent radical selected from the group
- C3-C7-alkylene,
 - C3-C7-cycloalkylene,
 - C3-C7-alkenylene,
 - C4-C7-cycloalkenylene,
 - 10 C5-C8-cycloalkadienylene,
 - C3-C7-alkadienylene,
 - C3-C7-alkynylene,
 - C4-C7-alkenynylene,
 - C6-C14-arylene,
 - 15 C6-C14-aryl-C2-C4-alkynylene,
 - C1-C3-alkyl-C6-C14-aryl-C2-C4-alkynylene,
 - C6-C14-aryl-C2-C4-alkenylene,
 - C1-C3-alkyl-C6-C14-arylene,
 - C1-C3-alkyl-C6-C14-aryl-C2-C4-alkenylene,
 - 20 C6-C14-aryl-C1-C3-alkylene,
 - C6-C14-aryl-C1-C3-alkyloxyene,
 - C1-C2-alkyl-C6-C14-aryl-C1-C2-alkylene,
 - C1-C3-alkyloxy-C6-C14-arylene,
 - C2-C8-alkyloxyene,
 - 25 C1-C5-alkyloxy-C1-C5-alkylene,
 - C6-C10-aryloxyene,
 - C6-C10-aryloxy-C1-C5-alkylene,
 - C6-C10-arylthio-C1-C5-alkylene,

30






5



10



15



where

R¹⁴ is selected from a chemical bond,

20

C₁-C₈-alkyl,C₁-C₈-alkyloxy,C₃-C₇-cycloalkyl,C₂-C₅-alkenyl,C₃-C₅-alkynyl,C₆-C₁₀-aryl,

25

C₁-C₃-alkyl-C₆-C₁₂-aryl,C₁-C₂-alkyl-C₆-C₁₀-aryl-C₁-C₂-alkyl,C₆-C₁₀-aryl-C₁-C₂-alkyl,

HET, and

pyridyl;

R¹⁵ is selected from a chemical bond,

5 C₁-C₄-alkyl,
C₂-C₄-alkenyl,
C₂-C₄-alkynyl, and
C₆-C₁₀-aryl;

R¹⁶ is selected from a chemical bond,

10 C₁-C₅-alkyl,
C₃-C₇-cycloalkyl,
C₃-C₅-alkenyl,
C₃-C₅-alkynyl,
C₆-C₁₀-aryl,
C₁-C₃-alkyl-C₆-C₁₂-aryl,
C₁-C₃-alkyl-C₆-C₁₂-aryl,
15 C₆-C₁₀-aryl-C₁-C₂-alkyl,
HET, and
pyridyl;

R²² is selected from the group consisting of

20 hydroxy,
C₁-C₈-alkoxy,
C₃-C₁₂-alkenoxy,
C₆-C₁₂-aryloxy,
C₁-C₆-alkyl-C₆-C₁₂-aryloxy,
di-C₁-C₈-alkylamino-C₁-C₈-alkoxy,
25 acylamino-C₁-C₈-alkoxy selected from the group
acetylaminooethoxy,
nicotinoylaminoethoxy, and
succinamidoethoxy,

30 C₁-C₈-alkoxyloxy-C₁-C₈-alkoxy,
C₆-C₁₂-aryl-C₁-C₈-alkoxy where the aryl group is unsubstituted or
substituted with one to three of the groups

nitro,
halo (F, Cl, Br, I),
C₁-C₄-alkoxy, and
35 amino,

hydroxy-C₂-C₈-alkoxy,
dihydroxy-C₃-C₈-alkoxy,
morpholino-C₂-C₈-alkoxy, and

NR²³R²⁴,

R²³ and R²⁴ are independently selected from

C₁-C₈-alkyl,

C₃-C₈-alkenyl,

5

hydrogen,

C₆-C₁₂-aryl

where the aryl group is unsubstituted or substituted with one to three of the groups

nitro,

halo (F, Cl, Br, I),

10

C₁-C₄-alkoxy, and

amino, and

C₆-C₁₂-aryl-C₁-C₈-alkyl

where the aryl group is unsubstituted or substituted with one to three of the groups

nitro,

15

halo (F, Cl, Br, I), and

C₁-C₄-alkoxy;

R²⁵ is selected from

R⁶, and

(CH₂)_mR¹, where m is 1, 2, or 3;

20

R²⁶ is selected from

hydrogen,

C₁-C₈-alkyl,

C₆-C₁₂-aryl, and

C₆-C₁₂-aryl-C₁-C₈-alkyl

25

where any aryl or alkyl groups are unsubstituted or substituted with one to three of the groups


nitro, and

halo (F, Cl, Br, I); and


pharmaceutically acceptable salts thereof.

30


- 4) The compound of Claim 2 selected from the group:




IIIa




IIIb




Va




Vb




VIa



VIb



VIIa



VIII

- 10 where
R¹ is selected from
hydrogen,

- halo(F, Cl, Br, I),
 cyano,
 carboxy,
 aminocarbonyl,
 5 carboxamido,
 carbamoyloxy,
 formyloxy,
 formyl,
 azido,
 10 nitro,
 imidazolyl,
 ureido,
 thioureido,
 thiocyanato,
 15 hydroxy,
 mercapto,
 sulfonamido, and
 an optionally substituted radical selected from
 C₁-C₁₂alkyl,
 20 C₁-C₁₂alkyloxy, carbonyl,
 C₁-C₁₂alkyloxyC₁-C₁₂alkyl,
 C₂-C₁₂alkenyl,
 C₂-C₁₂alkynyl,
 C₃-C₁₂cycloalkyl,
 25 C₆-C₁₄aryl,
 C₆-C₁₀aryl-C₁-C₈alkyl,
 C₁-C₁₂alkyloxy,
 C₆-C₁₄aryloxy, and
 C₁-C₁₂acylamino,
 30 where the substituents are selected from
 halo (F, Cl, Br, I),
 amino,
 amidino,
 guanidino,
 35 imidazolyl,
 indolyl,
 cyano,
 azido,

5 nitro,
 hydroxy,
 mercapto,
 morpholino,
 morpholinyl,
 piperazinyl,
 piperidinyl,
 pyrrolinyl,
 10 sulfonamido,
 ureido,
 thioureido,
 carboxy,
 aminocarbonyl,
 15 C₁-C₄aminocarbonyl,
 carboxamido,
 carbamoyloxy,
 formyloxy,
 formyl,
 20 C₁-C₄alkyloxycarbonyl,
 C₁-C₄alkyl,
 C₁-C₄alkylamino,
 di-(C₁-C₄)alkylamino,
 C₁-C₄alkoxy,
 25 C₁-C₄alkthio,
 hydroxyphenyl,
 phenyl, and
 phenoxy;

30 Q is selected from the group consisting of
 an amino group selected from

 NH₂,
 NR³H,
 NR³R⁴, and
 NR³R⁴R⁵,

35 R³, R⁴, and R⁵ are independently an optionally substituted radical selected from
 hydrogen,
 C₁-C₄ alkoxy,
 C₁-C₄ alkyl,

- halo(F, Cl, Br, I)C₁-C₄-alkyl,
 optionally substituted C₁-C₁₂ alkyl,
 optionally substituted C₃-C₁₂ cycloalkyl,
 optionally substituted C₆-C₁₄ aryl,
 5 optionally substituted C₁-C₆alkylC₆-C₁₄aryl,
 optionally substituted C₁-C₈ alkoxy,
 optionally substituted C₁-C₈ alkoxycarbonyl,
 optionally substituted C₆-C₁₄ aryloxy,
 optionally substituted C₆-C₁₄ aryloxycarbonyl,
 10 NR⁶R⁷,
 C(=NR⁸)-NR⁶R⁷,
 N=CR⁹-NR⁶R⁷,
 NR¹⁰-CR⁹=NR⁸, and
 NR¹⁰-C(=NR⁸)-NR⁶R⁷
 15 R⁶, R⁷, R⁸, R⁹, and R¹⁰ are independently selected from
 hydrogen,
 C₁-C₄ alkyl, and
 R¹¹,

- and where the substituents are one to three R¹¹,
 20 R¹¹ is independently selected from
 optionally substituted C₆-C₁₂ aryloxy,
 optionally substituted C₆-C₁₂ arylamino,
 optionally substituted C₆-C₁₂ aroyl, and
 optionally substituted C₆-C₁₂ aryl,
 25 where the substituents are one to three R¹²,
 R¹² is selected from

- nitro,
 amino,
 C₁-C₈alkylamino,
 30 di-(C₁-C₈)alkylamino,
 amidino,
 aminomethyleneimino,
 imino,
 imino-C₁-C₄alkyl,
 35 iminomethyleneamino,
 guanidino,
 C₆-C₁₀arylamino,
 C₁-C₈acylamino,

- 5 C₁-C₄alkylsulfonamino,
 azido,
 cyano,
 hydroxy,
 hydroxy-C₁-C₈alkyl,
 C₁-C₈alkoxy,
 phenyloxy,
 C₁-C₈alkanoyloxy,
 C₁-C₈alkanoyl,
 10 C₆-C₁₂aroyl,
 benzamido,
 phenyl,
 halo(F, Cl, Br, I),
 halo-C₁-C₈alkyl,
 15 C₁-C₈alkyl,
 aminosulfonyl,
 oxo,
 thio,
 thiocarbonyl,
 20 hydroxy,
 mercapto,
 formyl,
 formyloxy,
 carboxy,
 25 ureido,
 glycyI,
 phthalimido,
 succinimido,
 morpholino,
 30 C₃-C₇cycloalkyl, and
 optionally R³ and R⁴ taken together may form optionally substituted
 tetramethylene,
 pentamethylene,
 3-oxopentamethylene, and
 35 3-azapentamethylene,

where the substituents are selected from one to three R¹²,
 an amidino group selected from
 C(=NH)-NH₂,






$C(=NH)-NHR^3$,
 $C(=NR^4)-NHR^3$,
 $C(=NH)-NR^3R^4$, and
 $C(=NR^5)-NR^3R^4$,

5





where R^3 , R^4 , and R^5 are defined above or are selected from the group

$HO-$,
 $C_1-C_{10}alkyl-O-(C=O)-$,
 $C_1-C_{10}alkyl-S-(C=O)-$,
 $Cl_3CCH_2O-(C=O)-$,


10



15





, and




;

an aminoalkyleneamino group selected from




where R^3 , R^4 , and R^5 are defined above,

an iminoalkyleneamino group selected from



where R^3 , R^4 , and R^5 are defined above,

a guanidino group selected from



where R^3 , R^4 , and R^5 are defined above;

L is an optionally substituted bivalent radical selected from the group

C₃-C₇-alkylene,

C₃-C₇-cycloalkylene,

C₃-C₇-alkenylene,



C₄-C₇-cycloalkenylene,

C₅-C₈-cycloalkadienylene,


C₃-C₇-alkadienylene,

C₃-C₇-alkynylene,

- 5 C₄-C₇-alkenynylene,
 C₆-C₁₄-arylene,
 C₆-C₁₄-aryl-C₂-C₄-alkynylene,
 C₁-C₃-alkyl-C₆-C₁₄-aryl-C₂-C₄-alkynylene,
 C₆-C₁₄-aryl-C₂-C₄-alkenylene,
 C₁-C₃-alkyl-C₆-C₁₄-arylene,
 C₁-C₃-alkyl-C₆-C₁₄-aryl-C₂-C₄-alkenylene,
 C₆-C₁₄-aryl-C₁-C₃-alkylene,
 C₆-C₁₄-aryl-C₁-C₃-alkyloxyene,
 10 C₁-C₂-alkyl-C₆-C₁₄-aryl-C₁-C₂-alkylene,
 C₁-C₃-alkyloxy-C₆-C₁₄-arylene,
 C₂-C₈-alkyloxyene,
 C₁-C₅-alkyloxy-C₁-C₅-alkylene,
 C₆-C₁₀-aryloxyene,
 15 C₆-C₁₀-aryloxy-C₁-C₅-alkylene,
 C₆-C₁₀-arylthio-C₁-C₅-alkylene,






20



25





10 where

R^{14} is selected from a chemical bond,

15 C₁-C₈-alkyl,
C₁-C₈-alkyloxy,
C₃-C₇-cycloalkyl,
C₂-C₅-alkenyl,
C₃-C₅-alkynyl,
C₆-C₁₀-aryl,
C₁-C₃-alkyl-C₆-C₁₂-aryl,
C₁-C₂-alkyl-C₆-C₁₀-aryl-C₁-C₂-alkyl,
20 C₆-C₁₀-aryl-C₁-C₂-alkyl,
HET, and
pyridyl;

R^{15} is selected from a chemical bond,

25 C₁-C₄-alkyl,
C₂-C₄-alkenyl,
C₂-C₄-alkynyl, and
C₆-C₁₀-aryl;

R^{16} is selected from a chemical bond,

30 C₁-C₅-alkyl,
C₃-C₇-cycloalkyl,
C₃-C₅-alkenyl,
C₃-C₅-alkynyl,
C₆-C₁₀-aryl,


- C₁-C₃-alkyl-C₆-C₁₂-aryl,
 C₁-C₃-alkyl-C₆-C₁₂-aryl,
 C₆-C₁₀-aryl-C₁-C₂-alkyl,
 HET, and
 5 pyridyl;
 R¹⁸, R¹⁹, R²⁰ and R²¹ are independently selected from
 hydrogen,
 C₁-C₄alkyl, and
 halo(F, Cl, Br, I)C₁-C₄alkyl;
 10 R²² is selected from the group consisting of
 hydroxy,
 C₁-C₈-alkoxy,
 C₃-C₁₂-alkenoxy,
 C₆-C₁₂-aryloxy,
 15 C₁-C₆-alkyl-C₆-C₁₂-aryloxy,
 di-C₁-C₈-alkylamino-C₁-C₈-alkoxy,
 acylamino-C₁-C₈-alkoxy selected from the group
 acetylaminoethoxy,
 nicotinoylaminoethoxy, and
 20 succinamidoethoxy,
 C₁-C₈-alkoxyloxy-C₁-C₈-alkoxy,
 C₆-C₁₂-aryl-C₁-C₈-alkoxy where the aryl group is unsubstituted or
 substituted with one to three of the groups
 nitro,
 25 halo (F, Cl, Br, I),
 C₁-C₄-alkoxy, and
 amino,
 hydroxy-C₂-C₈-alkoxy,
 dihydroxy-C₃-C₈-alkoxy,
 30 morpholino-C₂-C₈-alkoxy, and
 NR²³R²⁴,
 R²³ and R²⁴ are independently selected from
 hydrogen,
 C₁-C₈-alkyl,
 35 C₃-C₈-alkenyl,
 C₆-C₁₂-aryl
 where the aryl group is unsubstituted or substituted with one to three of the groups
 nitro,

halo (F, Cl, Br, I),
 C₁-C₄-alkoxy, and
 amino, and
 C₆-C₁₂-aryl-C₁-C₈-alkyl



5 where the aryl group is unsubstituted or substituted with
 nitro,
 halo (F, Cl, Br, I), and
 C₁-C₄-alkoxy; and
 pharmaceutically acceptable salts thereof.

10




5) The compound of Claim 2 where Q-L- is selected from






15





20






25




5










10







15










5




10



15




, and



where any amino, amidino, or guanidino nitrogen is optionally derivitized with C₁-
5 C₆alkoxycarbonyl.

6) The compound of Claim 2 represented by structural formula Ia



Ia

where

R¹ and R² are each one or two substituents independently selected from hydrogen, fluoro, trifluoromethyl and chloro;

R¹⁸, R¹⁹, R²⁰ and R²¹ are independently selected from R¹ and methyl;

A¹ is CH or N;

A² is CH, C-CH₃, C-CH₂-CH₃, or N;

-X-Y- is selected from -NH-C(=O)-, -C(=O)-NH-, -CC-, -CH₂-O-, -O-CH₂- and -CH₂-CH₂-; and

Z¹ is selected from N and CR¹.

7) The compound of Claim 2 selected from the group consisting of

4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,

- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methyl-6-oxo, acetate,
- 5 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-methyl-6-oxo, acetate,
- 10 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-phenyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-phenyl-6-oxo, acetate,
- 15 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-phenyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-phenyl-6-oxo, acetate,
- 20 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 25 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-methyl-6-oxo, acetate,
- 30 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-methyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-hydroxy-6-oxo, acetate,
- 35 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-hydroxy-6-oxo, acetate,

- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-hydroxy-6-oxo, acetate,
- 5 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-hydroxy-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-methoxy-6-oxo, acetate,
- 10 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methoxy-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-methoxy-6-oxo, acetate,
- 15 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-methoxy-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-1-phenyl-6-oxo, acetate,
- 20 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-phenyl-6-oxo, acetate,
- 25 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)benzoyl]amino]-1-phenyl-6-oxo, acetate,
- 4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)picolinoyl]amino]-1-phenyl-6-oxo, acetate,
- 30 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy]-6-oxo, acetate,
- 35 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]ethynyl]-6-oxo, acetate,

- 4H-[1,2,3,4]tetrazolo[4,3-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)benzoyl]amino]-6-oxo, acetate,
- 5 4H-[1,2,3,4]tetrazolo[1,5-a][1,4]benzodiazepine-5(6H)-propanoic acid, 8-[[4-(aminoiminomethyl)picolinoyl]amino]-6-oxo, acetate,
- 10 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 15 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 20 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 25 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 30 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 35 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,

- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(R)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 5 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 10 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-2-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-]pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 15 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 20 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-3-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-2-chloro-phenyl]ethynyl]-1-methyl-6-oxo, acetate,
- 25 4H-imidazo[1,2-a][1,4]benzodiazepine-5(6H)-(S)-3-methyl propanoic acid, 8-[[4-(aminoiminomethyl)-]pyridyl]ethynyl]-1-methyl-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 30 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 35 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,

- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 5 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-3-methyl, propanoic acid,8-[[4-(aminoiminomethyl)phenyl]methoxy-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino-6-oxo, acetate,
- 10 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl,(S)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino]-6-oxo, acetate,
- 15 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl,(R)-3-methyl-propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino-6-oxo, acetate,
- 20 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(R)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino-6-oxo, acetate,
- 25 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-2-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate,
- 30 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-3-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-7-chloro-benzoyl]amino-6-oxo, acetate,
- 4H-[1,2,3,4]tetrazolo[4,3a][1,4]benzodiazepine-5(6H)-(S)-3-methyl, propanoic acid,8-[[4-(aminoiminomethyl)-benzoyl]amino]-6-oxo, acetate.
- 35

8) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and the compound of Claim 1.

9) The use of a platelet aggregation inhibiting amount of the composition of Claim 8A for inhibiting platelet aggregation.

10) The use of a pharmaceutically effective amount of the composition of Claim 8
5 for reducing platelet aggregation in a mammal.

11) The use of a pharmaceutically effective amount of the composition of Claim 8 for treating a mammal having an increased propensity for thrombus formation.

10 12) The use of Claim 11 wherein the composition further includes a thrombolytic agent.


13) The use of Claim 11 wherein the composition further includes an
anticoagulant.

15


20

25

30



I



II