

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 October 2008 (16.10.2008)

PCT

(10) International Publication Number
WO 2008/124157 A1

(51) International Patent Classification:
A61K 31/70 (2006.01) **A61K 31/519** (2006.01)

(74) Agent: **COLEMAN, Henry, D.**; Coleman Sudol Sapone,
P.C., 714 Colorado Avenue, Bridgeport, CT 06605-1601
(US).

(21) International Application Number:
PCT/US2008/004568

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date: 9 April 2008 (09.04.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/922,701 10 April 2007 (10.04.2007) US

(71) Applicant (*for all designated States except US*): **UNIVERSITY OF GEORGIA RESEARCH FOUNDATION, INC.** [US/US]; Boyd Graduate Studies Research Center, Room No. 633, Athens, GA 30602-7411 (US).

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **CHU, David** [US/US]; 662 Garden Cir., Statham, GA 30666 (US).
RAO, Jagadeeshwar [IN/US]; 2091 S. Milledge Ave., Apt. B-45, Athens, GA 30605 (US).

Published:

— with international search report

(54) Title: CARBOCYCLIC COMPOUNDS AND METHODS FOR TREATING EMERGING DISEASES, INCLUDING INFLUENZA AND VENEZUELA EQUINE ENCEPHALITIS VIRUS

FIGURE 1

Table 1 COMPARISON OF the Activity of CARBODINE ENANTIOMERS

VIRUS			VISUAL			NEUTRAL			VYR- EC90
VIRUS	DILUTION	DRUG	EC50	IC50	SI	EC50	IC50	SI	
DUCK	1/1000	Neg. Enantiomer	0.57	>100	175	0.87	>100	>115	0.22
DUCK	1/1000	Pos. Enantiomer	>100	>100	>1	>100	>100	>1	>100
DUCK	1/1000	Racemic							
DUCK	1/1000	Carbodine	2.7	>100	>37	5.3	>100	>19	11.4
DUCK	1/1000	ribivirin	5.7	>320	>56	5.8	>320	>55	6.5
GULL	1/100	Neg. Enantiomer	0.27	>100	370	0.31	>100	323	0.22
GULL	1/100	Pos. Enantiomer	>100	>100	>1	>100	>100	>1	>100
GULL	1/100	Racemic							
GULL	1/100	Carbodine	1.8	>100	>56	1.9	>100	>53	1.3
GULL	1/100	ribivirin	5.7	>320	>56	5.9	>320	>54	3.7
HK/2003	1/100	Neg. Enantiomer	0.18	>100	556	0.3	>	333	0.35
HK/2003	1/100	Pos. Enantiomer	>100	>100	>1	>100	>100	>1	>100
HK/2003	1/100	Racemic							
HK/2003	1/100	Carbodine	0.7	>100	>143	1.7	>100	>59	4.6
HK/2003	1/100	ribivirin	0.57	>320	>561	1.4	>320	>229	3.5
VN	1/500	Neg. Enantiomer	0.18	>100	>556	0.19	>100	526	0.11
VN	1/500	Pos. Enantiomer	>100	>100	>1	>100	>100	>1	>100
VN	1/500	Racemic							
VN	1/500	Carbodine	0.57	>100	>175	0.92	>100	>109	1.9
VN	1/500	ribivirin	0.39	>320	>821	0.59	>320	>542	0.44

(57) Abstract: The present invention relates to the use of carbodine and 5-F carbodine and analogs thereof for use in the treatment or prophylaxis of influenza, in particular the H5N1 strain of Avian Influenza A virus or "bird flu" strain of influenza as well as the treatment or prophylaxis of Venezuela equine encephalitis virus or VEE.

WO 2008/124157 A1

Carbocyclic Compounds and Methods for Treating Emerging Diseases, Including Influenza and Venezuela Equine Encephalitis Virus

Field of the Invention

The present invention relates to the use of carbodine and 5-F carbodine and analogs thereof for use in the treatment or prophylaxis of influenza, in particular the H5N1 strain of Avian Influenza A virus or "bird flu" strain of influenza as well as the treatment or prophylaxis of Venezuela equine encephalitis virus or VEE.

Claim of Priority and Government Rights

This application claims the benefit of priority of US provisional application serial number US60/922,701, filed April 10, 2007, which is incorporated by reference in its entirety herein.

The work which gave rise to this patent application was supported by a government grant, NIH (1 UO19 AI056540). Consequently, the government retains certain rights in the invention.


Background of the Invention

In 1997, highly pathogenic avian influenza H5N1 was transmitted from poultry to humans in Hong Kong, resulting in eighteen infected people and six deaths, and reemerged in 2003 causing two similar cases with one fatality. In 2003-2005, extensive outbreaks of H5N1 influenza occurred in nine Asian countries resulting in 19 human cases in Thailand, 91 in Vietnam, seven in Indonesia, and four in Cambodia, with a total of 62 reported deaths. Furthermore, H5N1 infections in family clusters have raised the possibility of human-to-human transmission. As human exposure to and infection with H5N1 viruses continues to increase, so, too, does the likelihood of the generation of an avian-human reassortment virus that may be transmitted efficiently within the global human population, which currently lacks H5N1 specific immunity. Such reassortment events between avian-human and swine-human influenza A viruses have been associated with the 1957 and 1968 influenza pandemics; the 1918 pandemic events remain unclear.

Concern over the potential for the generation of a pandemic H5 strain and its concomitant morbidity and mortality are spurring the search for effective agents against same.

Brief Description of the Invention

The present invention relates to carbocyclic nucleoside compounds according to the structure:



Wherein R is H or F;

R¹ and R^{1a} are each independently H, an acyl group, a C₁—C₂₀ alkyl or ether group, an amino acid residue (D or L), a phosphate, diphosphate, triphosphate or phosphodiester group or together R¹ and R^{1a} form a carbodiester or phosphodiester group with the oxygen atoms to which they are bonded;

R² is H, an acyl group, a C₁—C₂₀ alkyl or ether group or an amino acid residue;

And pharmaceutically acceptable salts, enantiomers, hydrates and solvates thereof.

Preferably R^{1a} is H. Also preferably, R¹ and R² are both H.

The present invention also relates to pharmaceutical compositions comprising an effective amount of a compound as described above, optionally in combination with a pharmaceutically acceptable carrier, additive or excipient.

Methods of treatment represent further embodiments according to the present invention. In this aspect a method of treating or reducing the likelihood of a an Avian

influenza A H5N1 strain viral infection or a Venezuelan equine encephalitis viral infection comprising administering to a patient in need of therapy or at risk for infection thereof an effective amount of a compound as otherwise described above.

Detailed Description of the Invention

The following definitions are used to describe the invention. If a term is not specifically defined herein, the meaning given to the term is that which one of ordinary skill would apply to the term within the context of the term's use.

The term "compound", as used herein, unless otherwise indicated, refers to any specific chemical compound disclosed herein, generally refers to β -D nucleoside analogs, but may include, within context, tautomers, regioisomers, geometric isomers, and where applicable, optical isomers (enantiomers) thereof, as well as pharmaceutically acceptable salts thereof, solvates and/or polymorphs. Within its use in context, the term compound generally refers to a single compound, but also may include other compounds such as stereoisomers, regioisomers and/or optical isomers (including racemic mixtures) as well as specific enantiomers or enantiomerically enriched mixtures of disclosed compounds.

The term "patient" or "subject" is used throughout the specification to describe an animal, preferably a human, to whom treatment, including prophylactic treatment, with the compositions according to the present invention is provided. For treatment of those infections, conditions or disease states which are specific for a specific animal such as a human patient, the term patient refers to that specific animal. In general, in the present invention, the term patient refers to a human patient unless otherwise stated. In the present invention, in addition to humans, domesticated animals (e.g., horses, cows, dogs, cats, etc.) also may be commonly treated.

The term "H5N1 influenza" is used to describe a highly pathogenic subtype or strain of influenza A ("HPAI") originating in birds which is particularly lethal to humans. Humans can be infected with influenza types A, B, and C viruses. Subtypes of influenza A that are currently circulating among people worldwide include H1N1, H1N2, and H3N2 viruses. Wild birds are the natural host for all known subtypes of influenza A viruses. Typically, wild

birds do not become sick when they are infected with avian influenza A viruses. However, domestic poultry, such as turkeys and chickens, can become very sick and die from avian influenza, and some avian influenza A viruses also can cause serious disease and death in wild birds.

Avian influenza A virus strains are further classified as low pathogenic (LPAI) or highly pathogenic (HPAI) on the basis of specific molecular genetic and pathogenesis criteria that require specific testing. Most avian influenza A viruses are LPAI viruses that are usually associated with mild disease in poultry. In contrast, HPAI viruses can cause severe illness and high mortality in poultry. More recently, some HPAI viruses (e.g., H5N1) have been found to cause no illness in some poultry, such as ducks. LPAI viruses have the potential to evolve into HPAI viruses and this has been documented in some poultry outbreaks. Avian influenza A viruses of the subtypes H5 and H7, including H5N1, H7N7, and H7N3 viruses, have been associated with HPAI, and human infection with these viruses have ranged from mild (H7N3, H7N7) to severe and fatal disease (H7N7, H5N1).

In general, direct human infection with avian influenza viruses occurs very infrequently, and has been associated with direct contact (e.g., touching) infected sick or dead infected birds (domestic poultry).

The term “Venezuelan equine encephalitis virus” or “encephalomyelitis (VEE) is a mosquito-borne viral pathogen that causes Venezuelan equine encephalitis or encephalomyelitis (VEE). VEE can affect all equine species, such as horses, asses, and zebras. After infection, equines may suddenly die or show progressive central nervous system disorders. Humans also can contract this disease. Healthy adults who become infected by the virus may experience flu-like symptoms, such as high fevers and headaches. People with weakened immune systems and the young and the elderly can become severely ill or die from this disease.

The virus that causes VEE is transmitted primarily by mosquitoes that bite an infected animal and then bite and feed on another animal or human. The speed with which the disease spreads depends on the subtype of the VEE virus and the density of mosquito populations. Enzootic subtypes of VEE are diseases endemic to certain areas. Generally these serotypes do not spread to other localities. Enzootic subtypes are associated with the rodent-mosquito

transmission cycle. These forms of the virus can cause human illness but generally do not affect equine health.

Epizootic subtypes, on the other hand, can spread rapidly through large populations. These forms of the virus are highly pathogenic to equines and can also affect human health. Equines, rather than rodents, are the primary animal species that carry and spread the disease. Infected equines develop an enormous quantity of virus in their circulatory system. When a blood-feeding insect feeds on such animals, it picks up this virus and transmits it to other animals or humans. Although other animals, such as cattle, swine, and dogs, can become infected, they generally do not show signs of the disease or contribute to its spread.

Serology testing has been performed on this virus and has shown the presence of six different subtypes (classified I to VI). There are seven different variants in subtype I, and three of these variants, A, B, and C are the epizootic strains. The present application relates to the treatment or prophylaxis (reducing the likelihood of prevention) of the six different subtypes and the variants within the subtypes.

The term "compound", as used herein, unless otherwise indicated, refers to any specific chemical compound disclosed herein and includes within the context of a compound's use, tautomers, regioisomers, geometric isomers, and where applicable, optical isomers thereof, as well as pharmaceutically acceptable salts, solvates and polymorphs thereof. Within its use in context, the term compound generally refers to a single compound, but also may include other compounds such as stereoisomers, regioisomers and/or optical isomers (including in some instances, racemic mixtures) as well as specific enantiomers or enantiomerically enriched mixtures of disclosed compounds. In aspects of the present invention, the β -D nucleoside analog is used, especially in a highly purified form.

The term "pharmaceutically acceptable salt" is used throughout the specification to describe, where applicable, a salt form of one or more of the compounds described herein which are presented to increase the solubility of the compound in the gastric juices of the patient's gastrointestinal tract in order to promote dissolution and the bioavailability of the compounds. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids, where applicable. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as

calcium, magnesium and ammonium salts, among numerous other acids well known in the pharmaceutical art. Sodium and potassium salts are particularly preferred as neutralization salts of the phosphates according to the present invention.

The term "pharmaceutically acceptable derivative" is used throughout the specification to describe any pharmaceutically acceptable prodrug form (such as an ester, ether or amide or other prodrug group) which, upon administration to a patient, provides directly or indirectly the present compound or an active metabolite of the present compound.

The term "alkyl" shall mean within its context a C₁-C₂₀, preferably a C₁-C₁₀ linear, branch-chained or cyclic fully saturated hydrocarbon radical, which may be optionally substituted. The term "ether" shall mean an optionally substituted C₁ to C₂₀ ether group, formed from an oxygen and an alkyl group, or alternatively, may also contain at least one oxygen within the alkyl or alkylene chain.


The term "aromatic" or "aryl" shall mean within its context a substituted or unsubstituted monovalent carbocyclic aromatic radical having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl, anthracene, phenanthrene). Other examples include optionally substituted heterocyclic aromatic ring groups ("heteroaromatic" or "heteroaryl") having one or more nitrogen, oxygen, or sulfur atoms in the ring. The preferred aryl group in compounds according to the present invention is a phenyl or a substituted phenyl group.

The term "heterocycle" shall mean an optionally substituted moiety which is cyclic and contains at least one atom other than a carbon atom, such as a nitrogen, sulfur, oxygen or other atom, which ring may be saturated and/or unsaturated.

The term "unsubstituted" shall mean substituted only with hydrogen atoms. The term "substituted" shall mean, within the chemical context of the compound defined, a substituent (each of which substituent may itself be substituted) selected from a hydrocarbyl (which may be substituted itself, preferably with an optionally substituted alkyl or fluoro group, among others), preferably an alkyl (generally, no greater than about 3 carbon units in length), including CF₃, an optionally substituted aryl, halogen (F, Cl, Br, I), thiol, hydroxyl, carboxyl, C₁-C₃ alkoxy, alkoxycarbonyl, CN, nitro or an optionally substituted amine (e.g., an

alkyleneamine or a C₁-C₃ monoalkyl or dialkyl amine). Various optionally substituted moieties may be substituted with 3 or more substituents, preferably no more than 3 substituents and preferably with 1 or 2 substituents.

The term "acyl" is used throughout the specification to describe a group at the 5' position of the nucleoside analog (i.e., at the free hydroxyl position in the carbocyclic moiety) which contains a C₁ to C₂₀ linear, branched or cyclic alkyl chain. The acyl group at the 5' position, in combination with the 5' hydroxyl group results in an ester, which, after administration, may be cleaved to produce the free nucleoside form of the present invention. Acyl groups according to the present invention are represented by the structure:





where R⁴ is a C₁ to C₂₀ linear, branched or cyclic alkyl group, alkoxyalkyl (including an ethylene oxide chain which may end in a free hydroxyl group or a C₁-C₁₀ alkyl group and ranges in molecular weight from about 50 to about 40,000 or about 200 to about 5,000), such as phoxymethyl, aryl, alkoxy, alkoxy carbonyloxy groups (e.g., [(isopropoxycarbonyl)oxy]-methoxy), aryloxyalkyl, among others, all of which groups may be optionally substituted. Preferred acyl groups are those where R⁴ is a C₁ to C₁₀ alkyl group. Acyl groups according to the present invention also include, for example, those acyl groups derived from benzoic acid and related acids, 3-chlorobenzoic acid, succinic, capric and caproic, lauric, myristic, palmitic, stearic and oleic groups, among numerous others and may include such related groups as sulfone groups such as mesylate groups. All groups may be appropriately substituted within context as otherwise described herein. One of ordinary skill in the art will recognize the acyl groups which will have utility in the present invention, either to synthesize the target pharmaceutical compounds or as prodrug of the nucleosides according to the present invention.

The term "amino acid" or "amino acid residue" shall mean, within context, a radical of a D- or L-amino acid which is covalently bound to a nucleoside analog at the 4' exocyclic amine position of the cytosine base or the 5'- or 3'-OH position of the sugar synthon (R², R¹ or R^{1a}) through a carboxylic acid moiety of the amino acid, thus forming respectively, an amide or ester group linking the nucleoside to the amino acid. Representative amino acids

include both natural and unnatural amino acids, preferably including, for example, alanine, β -alanine, arginine, asparagine, aspartic acid, cysteine, cystine, glutamic acid, glutamine, glycine, phenylalanine, histidine, isoleucine, lysine, leucine, methionine, proline, serine, threonine, valine, tryptophan or tyrosine, among others.

The term "phosphate ester" or "phosphodiester" is used throughout the specification to describe mono-phosphate groups at the 5' position of the carboxylic sugar synthon which are mono- or diesterified such that the phosphate group is negatively charged or is rendered neutral, i.e., has a neutral charge. Phosphate esters for use in the present invention include those represented by the structures:



where each R^5 , R^6 and $R^{\text{''}}$ is independently selected from H, a C_1 to C_{20} linear, branched or cyclic alkyl group, alkoxyalkyl, aryloxyalkyl, such as phenoxymethyl, aryl and alkoxy, among others, including alkoxycarbonyloxy groups (e.g., (isopropoxycarbonyl)oxy]-methoxy) with the proviso that at least one R^5 group is other than H, or the two R^5 groups together form a five- or six-membered heterocyclic group, B is a direct bond (N directly bonded to C of the ester/carboxylic acid group) or a C_1 - C_3 alkylene group optionally substituted with a C_1 - C_3 alkyl group, preferably a methyl group and R^7 is a C_1 to C_{20} linear, branched or cyclic alkyl or acyl group, alkoxyalkyl, aryloxyalkyl, such as phenoxymethyl, aryl and alkoxy, among others, each of which groups previously mentioned may be optionally substituted. Preferred monophosphate esters for use in prodrug forms according to

the present invention are those where R⁵ is a C₁ to C₂₀ linear or branched chain alkyl group, more preferably a C₁ to C₃ alkyl group, all of which groups may be optionally substituted.

The term "effective amount" shall mean an amount or concentration of a compound according to the present invention which is effective within the context of its administration, which may be inhibitory, prophylactic and/or therapeutic. Within context, all active compounds which are used in the present invention are used in effective amounts. The present compound also relates to combinations of compounds which contain effective amounts of each of the compounds used, whether that combination is additive or synergistic in effect, provided that the overall effect of the combination of compounds is to inhibit the growth, reduce the likelihood of or treat viral infections in patients.

The term "D-configuration" as used in the context of the present invention refers to the configuration of the nucleoside compounds according to the present invention which mimics the natural configuration of sugar moieties as opposed to the unnatural occurring nucleosides or "L" configuration. The term "β" or "β anomer" is used to describe nucleoside analogs according to the present invention in which the nucleoside base (in this case cytosine or 5-fluorocytosine) is configured (disposed) above the plane of the carbocyclic moiety in the carbodine or 5-fluoro carbodine analog.


The term "enantiomerically enriched" is used throughout the specification to describe a nucleoside which includes at least about 95%, preferably at least about 96%, more preferably at least about 97%, even more preferably, at least about 98%, and even more preferably at least about 100% or more of a single enantiomer of that nucleoside. Carbodine and 5-F carbodine compounds according to the present invention are generally β-D-nucleoside compounds. When the present carbodine compounds according to the present invention are referred to in this specification, it is presumed that the nucleosides have the D-nucleoside configuration and are enantiomerically enriched (preferably, about 100% of the D-nucleoside), unless otherwise stated.

The terms "coadminister" and "coadministration" are used synonymously to describe the administration of at least one of the carbodine or 5-F carbodine nucleoside compounds according to the present invention in combination with at least one other agent, preferably at least one additional anti-viral agent (such as oseltamavir phosphate or tamiflu), including

other nucleoside anti-viral agents which are specifically disclosed herein in amounts or at concentrations which would be considered to be effective amounts at or about the same time. While it is preferred that coadministered agents be administered at the same time, agents may be administered at times such that effective concentrations of both (or more) agents appear in the patient at the same time for at least a brief period of time. Alternatively, in certain aspects of the present invention, it may be possible to have each coadministered agent exhibit its inhibitory effect at different times in the patient, with the ultimate result being the inhibition of influenza A, especially H5N1 influenza or VEE and the treatment of the aforementioned infections. Of course, when more than one viral or other infection or other condition is present, the present compounds may be combined with agents to treat that other infection or condition as required.

The term “independently” is used herein to indicate that the variable, which is independently applied, varies independently from application to application.

The present invention relates to carbocyclic nucleoside compounds according to the structure:



Wherein R is H or F;

R¹ and R^{1a} are each independently H, an acyl group, a C₁—C₂₀ alkyl or ether group, a phosphate, diphosphate, triphosphate or phosphodiester group or together R¹ and R^{1a} form a carbodiester or phosphodiester group with the oxygen atoms to which they are bonded;

R² is H, an acyl group, a C₁—C₂₀ alkyl or ether group;

And pharmaceutically acceptable salts, enantiomers, hydrates and solvates thereof.

In certain aspects of the invention, R^{1a} is H. Also, in certain aspects of the invention, R^1 and R^2 are both H. R is preferably H.

The present invention also relates to pharmaceutical compositions comprising an effective amount of a compound as described above, optionally in combination with a pharmaceutically acceptable carrier, additive or excipient.

Methods of treatment represent further embodiments according to the present invention. In this aspect a method of treating or reducing the likelihood of an Avian influenza A infection, in particular, a H5N1 strain viral infection or a Venezuelan equine encephalitis viral infection in a patient comprising administering to a patient in need of therapy or at risk for infection thereof an effective amount of a compound as otherwise described above.

Chemical Synthesis


Compositions according to the present invention are synthesized readily from D-ribose according to scheme I, which is presented below. In this scheme, D-ribose is first converted through a series of chemical steps to a protected carbocyclic five-membered ring as depicted below (compounds 9 and 10). Compound 10 (mesyl group on 1' hydroxyl position) is then converted to an azide group (compound 11), which is subsequently converted to an amine group (compound 12). Compound 12 is subsequently converted to a carbocyclic uridine analog 14, which may be converted, through an additional series of steps to carbodine (compound 16, R=H) or to 5-Fluorocarbodine (compound 20, R=F).

Each of carbodine and 5-Fluorocarbodine may be readily converted to prodrug or alternative forms of the present invention (e.g., acylated, phosphate or phosphodiester derivatives, etc. as otherwise described herein) utilizing standard synthetic chemistry for introducing various groups onto the hydroxyl positions at 2', 3' and/or 5' positions of the carbocyclic moiety or alternatively, at the exocyclic amine position at the 4-position of the cytosine base. Acylation proceeds through well known synthetic methods (acyl anhydride, acyl halide, etc.) and phosphorylation may be performed using standard chemical techniques which are well-known in the art. One of ordinary skill may readily synthesize compounds according to the present invention by utilizing specific chemical steps which are presented

herein or by way of analogy, through the use of chemical steps which are in the literature, by way of analogy.

(-)-Carbodine and 5-Fluorocarbodine

Scheme 1: Synthesis of (1*R*,2*S*,3*R*,4*R*)-(-)-1-[2,3-Dihydroxy-4-(hydroxymethyl)-cyclopentan-1-yl]cytosine



Reagents and conditions: (a) 2,2-DMP, *p*-toluenesulfonic acid, acetone, 0 °C to rt, 2 h; (b) TBDMSCl, imidazole, CH₂Cl₂, rt, 4 h; (c) vinylmagnesium bromide, THF, -78 °C to rt, 2 h; (d) TBAF, THF, rt, 2 h; (e) NaIO₄, H₂O, rt, 2 h; (f) KO^tBu, methyltriphenylphosphonium bromide, THF, 0 °C to rt, 6 h; (g) 1st gen. Grubb's catalyst, CH₂Cl₂, rt, 7 h; (h) PDC, 4 Å molecular sieves, AcOH, CH₂Cl₂, rt, 12 h; (i) (*t*-BuOCH₂)₂CuLi, *t*-BuOMe/THF, -30 °C, 30 min; (j) CeCl₃·7H₂O, NaBH₄, MeOH, -78 °C to rt, 1 h; (k) MsCl, Et₃N, CH₂Cl₂, 0 °C, 1 h; (l) NaN₃, DMF, 140 °C, 4 h; (m) 10% Pd/C, EtOH, rt, 30 psi, 5 h; (n) β-methoxyacryloyl isocyanate, DMF, -20 °C to rt, 10 h; (o) 30 % NH₄OH, EtOH/dioxane (1:1), 100 °C, 20 h; (p) 2,4,6-triisopropylbenzenesulfonyl chloride, DMAP, Et₃N, 30 % NH₄OH, rt, 12 h; (q) TFA/H₂O (2:1), 60 °C, 4 h; (r) Ac₂O, Et₃N, DMAP, CH₃CN, rt, 30 min; (s) i. selectfluor, AcOH, CH₃CN, 95 °C, 3h; ii. Et₃N, 95 °C, 3h.

Pharmaceutical compositions based upon the nucleoside compounds according to the present invention comprise one or more of the above-described compounds in a therapeutically effective amount for treating a viral, especially an Avian Influenza A strain viral infection, in particular, a H5N1 viral infection or a Venezuelan equine encephalitis (VEE) infection in a patient in need of therapy thereof, optionally in combination with a pharmaceutically acceptable additive, carrier or excipient. One of ordinary skill in the art will recognize that a therapeutically effective amount will vary with the infection or condition to be treated, its severity, the treatment regimen to be employed, the pharmacokinetics of the agent used, as well as the patient or subject (animal or human) to be treated.

In the pharmaceutical aspect according to the present invention, the compound according to the present invention is formulated preferably in admixture with a pharmaceutically acceptable carrier. In general, it is preferable to administer the pharmaceutical composition in orally-administrable form, but certain formulations may be administered via a parenteral, intravenous, intramuscular, transdermal, buccal, subcutaneous, suppository or other route. Intravenous and intramuscular formulations are preferably administered in sterile saline. In certain instances, transdermal administration may be preferred. Of course, one of ordinary skill in the art may modify the formulations within the teachings of the specification to provide numerous formulations for a particular route of administration without rendering the compositions of the present invention unstable or compromising their therapeutic activity. In particular, the modification of the present compounds to render them more soluble in water or other vehicle, for example, may be easily

accomplished by minor modifications (salt formulation, esterification, etc.) which are well within the ordinary skill in the art. It is also well within the routineer's skill to modify the route of administration and dosage regimen of a particular compound in order to manage the pharmacokinetics of the present compounds for maximum beneficial effect in patients.

In certain pharmaceutical dosage forms, the pro-drug form of the compounds, especially including acylated (acetylated or other) and ether (alkyl and related) derivatives, phosphate esters and various salt forms of the present compounds, are preferred. One of ordinary skill in the art will recognize how to readily modify the present compounds to pro-drug forms to facilitate delivery of active compounds to a targeted site within the host organism or patient. The routineer also will take advantage of favorable pharmacokinetic parameters of the pro-drug forms, where applicable, in delivering the present compounds to a targeted site within the host organism or patient to maximize the intended effect of the compound.

The amount of compound included within therapeutically active formulations according to the present invention is an effective amount for treating the infection or condition, especially an Avian influenza A infection, in particular, a H5N1 strain viral infection or a Venezuelan equine encephalitis viral infection. In general, a therapeutically effective amount of the present compound in pharmaceutical dosage form usually ranges from about 0.05 mg/kg to about 100 mg/kg per day or more, more preferably, slightly less than about 1 mg/kg to about 25 mg/kg per day of the patient or considerably more, depending upon the compound used, the condition or infection treated and the route of administration. The active nucleoside compound according to the present invention is preferably administered in amounts ranging from about 0.5 mg/kg to about 25 mg/kg per day of the patient, depending upon the pharmacokinetics of the agent in the patient. This dosage range generally produces effective blood level concentrations of active compound which may range from about 0.05 to about 100 micrograms/cc of blood in the patient. For purposes of the present invention, a prophylactically or preventive effective amount of the compositions according to the present invention falls within the same concentration range as set forth above for therapeutically effective amount and is usually the same as a therapeutically effective amount.

Administration of the active compound may range from continuous (intravenous drip) to several oral administrations per day (for example, Q.I.D.) or transdermal administration and may include oral, topical, parenteral, intramuscular, intravenous, sub-cutaneous,

transdermal (which may include a penetration enhancement agent), buccal and suppository administration, among other routes of administration. Enteric coated oral tablets may also be used to enhance bioavailability of the compounds from an oral route of administration. The most effective dosage form will depend upon the bioavailability/pharmacokinetics of the particular agent chosen as well as the severity of disease in the patient. Oral dosage forms are particularly preferred, because of ease of administration and prospective favorable patient compliance.

To prepare the pharmaceutical compositions according to the present invention, a therapeutically effective amount of one or more of the compounds according to the present invention is preferably intimately admixed with a pharmaceutically acceptable carrier according to conventional pharmaceutical compounding techniques to produce a dose. A carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral. In preparing pharmaceutical compositions in oral dosage form, any of the usual pharmaceutical media may be used. Thus, for liquid oral preparations such as suspensions, elixirs and solutions, suitable carriers and additives including water, glycols, oils, alcohols, flavouring agents, preservatives, colouring agents and the like may be used. For solid oral preparations such as powders, tablets, capsules, and for solid preparations such as suppositories, suitable carriers and additives including starches, sugar carriers, such as dextrose, mannitol, lactose and related carriers, diluents, granulating agents, lubricants, binders, disintegrating agents and the like may be used. If desired, the tablets or capsules may be enteric-coated or sustained release by standard techniques. The use of these dosage forms may significantly enhance the bioavailability of the compounds in the patient.

For parenteral formulations, the carrier will usually comprise sterile water or aqueous sodium chloride solution, though other ingredients, including those which aid dispersion, also may be included. Of course, where sterile water is to be used and maintained as sterile, the compositions and carriers must also be sterilized. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed.

Liposomal suspensions (including liposomes targeted to viral antigens) may also be prepared by conventional methods to produce pharmaceutically acceptable carriers. This

may be appropriate for the delivery of free nucleosides, acyl/alkyl nucleosides or phosphate ester pro-drug forms of the nucleoside compounds according to the present invention.

In particularly preferred embodiments according to the present invention, the compounds and compositions are used to treat, prevent or delay the onset of an Avian influenza A infection, in particular, a H5N1 strain viral infection or a Venezuelan equine encephalitis viral infection. Preferably, to treat, prevent or delay the onset of these infections, the compositions will be administered in oral dosage form in amounts ranging from about 250 micrograms up to about 500 mg or more at least once a day, preferably, up to four times a day. The present compounds are preferably administered orally, but may be administered parenterally, topically or in suppository form.

In the case of the co-administration of the present compounds in combination with another compound used to treat an Avian influenza A infection, in particular, a H5N1 strain viral infection and/or a Venezuelan equine encephalitis viral infection, such as Tamiflu, the amount of the carboline or 5-fluorcarboline nucleoside compound to be administered ranges from about 1 mg/kg. of the patient to about 500 mg/kg. or more of the patient or considerably more, depending upon the second agent to be co-administered and its potency against influenza (e.g. H5N1) or the VEE virus to be inhibited, the condition or infection treated and the route of administration. In the case of influenza A (e.g. H5N1 infections) the other agent may be preferably administered in amounts ranging from about 100 ug/kg (micrograms per kilogram) to about 500 mg/kg. In certain preferred embodiments, these compounds may be preferably administered in an amount ranging from about 1 mg/kg to about 50 mg/kg or more (usually up to about 100 mg/kg), generally depending upon the pharmacokinetics of the two agents in the patient. These dosage ranges generally produce effective blood level concentrations of active compound in the patient.

The compounds according to the present invention, may advantageously be employed prophylactically to prevent or reduce the likelihood of a viral infection or to prevent or reduce the likelihood of the occurrence of clinical symptoms associated with the viral infection or to prevent or reduce the likelihood of the spread of a viral infection to another person. Thus, the present invention also encompasses methods for the prophylactic treatment of an avian influenza A strain (e.g. H5N1) viral infection or a VEE viral infection. In this aspect according to the present invention, the present compositions are used to prevent, reduce the

likelihood of or delay the onset of a viral infection or a virus related disease or condition or the spread of infection to other people. This prophylactic method comprises administering to a patient in need of such treatment or who is at risk for the development of an avian influenza A strain (e.g. H5N1) viral infection or a VEE viral infection or an infected patient who wishes to prevent or reduce the likelihood of a viral infection from spreading to another person, an amount of a compound according to the present invention alone or in combination with another anti-viral effective for alleviating, preventing or delaying the onset of the viral infection. In the prophylactic treatment according to the present invention, it is preferred that the antiviral compound utilized should be as low in toxicity and preferably non-toxic to the patient. It is particularly preferred in this aspect of the present invention that the compound which is used should be maximally effective against the virus and should exhibit a minimum of toxicity to the patient. In the case of compounds of the present invention for the prophylactic treatment of viral infections, these compounds may be administered within the same dosage range for therapeutic treatment (i.e., about 250 micrograms up to about 500 mg. or more from one to four times per day for an oral dosage form) as a prophylactic agent to prevent the proliferation of the viral infection or alternatively, to prolong the onset of or reduce the likelihood of a patient contracting a virus infection which manifests itself in clinical symptoms.

In addition, compounds according to the present invention may be administered alone or in combination with other agents, including other compounds of the present invention. Certain compounds according to the present invention may be effective for enhancing the biological activity of certain agents according to the present invention by reducing the metabolism, catabolism or inactivation of other compounds and as such, are co-administered for this intended effect.

The present invention is now described, purely by way of illustration, in the following examples. It will be understood by one of ordinary skill in the art that these examples are in no way limiting and that variations of detail can be made without departing from the spirit and scope of the present invention.

Examples

Experimental (Chemical Synthesis)

2,3-*O*-Isopropylidene-D-ribofuranose (1). A suspension of D-ribose (300 g, 1.99 mol) in acetone (3000 mL) was cooled to 0 °C and treated with *p*-toluenesulfonic acid (11.4 g, 0.059 mol) and 2,2-dimethoxypropane (268 mL, 2.19 mol). After stirred at room temperature for 2 h, the clear resulting mixture was neutralized with NaHCO₃ (6.7 g), filtered and the filtrate was concentrated in vacuo. The residue was purified by silica gel column chromatography (EtOAc : hexane = 1:2) to afford **1** (342 g, 90%) as a colorless oil. ¹H NMR (CDCl₃) δ 5.62 (s, 1H), 4.83 (d, *J* = 5.9 Hz, 1H), 4.73 (d, *J* = 5.9 Hz, 1H), 4.41 (s, 1H), 3.72 (m, 2H), 1.48 (s, 3H), 1.32 (s, 3H); ¹³C NMR (CDCl₃) δ 112.05, 102.61, 87.53, 86.60, 81.53, 63.37, 26.23, 24.58.

5-(*tert*-Butyldimethylsilyloxy)-2,3-isopropylidenedioxy-D-ribofuranose (2). A solution of **1** (320 g, 1.68 mol) in CH₂Cl₂ (3000 mL), *tert*-Butyldimethylsilyl chloride (304 g, 2.01 mol) and imidazole (172 g, 2.52 mol) were added at 0 °C. After stirred for 4 h at room temperature, H₂O (1000 mL) was added to the reaction mixture. The water layer was separated and the organic layer was dried over Na₂SO₄ and filtered. The filtrate was concentrated in vacuo and the residue was purified by column chromatography on a silica gel (EtOAc : hexane = 1:20) to give **2** (415 g, 81%) as a colorless oil. ¹H NMR (CDCl₃) δ 5.27 (d, *J* = 11.7 Hz, 1H), 4.77 (d, *J* = 11.8 Hz, 1H), 4.69 (d, *J* = 5.9 Hz, 1H), 4.49 (d, *J* = 5.9 Hz, 1H), 4.35 (s, 1H), 1.54 (s, 0.3H), 1.47 (s, 2.7H), 1.38 (s, 0.3H), 1.31 (s, 2.7H), 0.92 (s, 8H), 0.87 (s, 1H), 0.16 (d, *J* = 2.8 Hz, 0.6H), 0.13 (d, *J* = 2.8 Hz, 5.4H); ¹³C NMR (CDCl₃) δ 112.00, 103.59, 103.43, 97.94, 88.70, 87.98, 87.58, 86.94, 81.87, 81.73, 81.18, 79.39, 65.47, 64.78, 63.83, 26.44, 25.74, 25.51, 24.89, -5.72.

(4*R*,5*S*)-1-{4-[2-(*tert*-Butyldimethylsilyloxy)-1-hydroxyethyl]-2,2-dimethyl-1,3-dioxolan-5-yl}-(*S*)-prop-2-en-1-ol (3). A solution of **2** (250 g 0.82 mol) in anhydrous tetrahydrofuran (1700 mL) was cooled to -78 °C and vinylmagnesium bromide (1M solution in tetrahydrofuran, 2.62 L, 2.62 mol) was added dropwise under -78 °C. After addition was completed, the reaction mixture was allowed to stir at room temperature for 2 h. The reaction mixture was poured in to a mix of saturated NH₄Cl (aq) and ether (1:2) (3 L). The organic and layer was separated and aqueous layer was extracted with ether (2 x 600 mL). The combined organic layers was washed with brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by silica gel column chromatography (EtOAc : hexane = 1:20) to give **3** (251 g, 92%) as colorless oil. $[\alpha]_D^{23} +6.86^\circ$ (C 0.59, CHCl₃); ¹H NMR (400

MHz, CDCl₃) δ 6.02 (m, 2H), 5.43 (d, J = 17.2 Hz, 1H), 5.26 (d, J = 10.4 Hz, 1H), 4.35 (bs, OH, D₂O exchangeable, 1H), 4.31 (bs, 1H), 4.07 (m, 2H), 3.86 (m, 2H), 3.65 (dd, J = 6.7 and 9.9 Hz, 1H), 3.36 (bs, OH, D₂O exchangeable, 1H), 1.39 (s, 3H), 1.32 (s, 3H), 0.91 (s, 9H), 0.10 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 137.31, 115.93, 108.65, 80.62, 76.53, 69.61, 69.22, 64.14, 27.86, 25.73, 25.29, -5.53; Anal. Calcd for C₁₆H₃₂O₅Si: C, 57.79; H, 9.70. Found: C, 58.13; H, 9.80.

(4R,5S)-1-[4-(1,2-dihydroxyethyl)-2,2-dimethyl-[1,3]dioxolan-5-yl]-(S)-prop-2-en-1-ol (4).

Tetrabutylammonium fluoride (1M solution in tetrahydrofuran, 1128 mL, 1.12 mol) was added to a solution of **3** (250 g, 0.75 mol) in tetrahydrofuran (1000 mL) and stirred at room temperature for 2 hr. The resulting brown mixture was concentrated in vacuo and the residue was purified by column chromatography on a silica gel (EtOAc : Hexane = 2:1) to give **4** (161 g, 98%) as a white solid. mp 73-74 °C; [α]_D²³ -31.33° (C 1.00, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 6.03 (m, 1H), 5.40 (dd, J = 0.8 and 17.2 Hz, 1H), 5.31 (dd, J = 0.8 and 10.5 Hz, 1H), 4.34 (t, J = 8.1 Hz, 1H), 4.16 (dd, J = 5.4 and 9.4 Hz, 1H), 4.06 (dd, J = 5.4 and 9.2 Hz, 1H), 3.95-3.87 (m, 1H D₂O exchangeable, 3H), 2.15 (bs, OH, D₂O exchangeable, 1H), 1.40 (s, 3H), 1.34 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 137.64, 117.13, 79.98, 77.83, 69.31, 64.56, 27.96, 25.43; Anal. Calcd for C₁₀H₁₈O₅·0.03 hexane: C, 55.37; H, 8.41. Found: C, 55.60; H, 8.31.

(1S,2S,3S)-2,2-Dimethyl-6-vinyltetrahydrofuro[3,4-*d*]-1,3-dioxol-4-ol (5). A solution of triol **4** (161 g, 737.6 mmol) in H₂O (1600 mL) was cooled to 0 °C and NaIO₄ (237 g, 1.106 mol) was added portion wise. After stirring at room temperature for 2 h, the reaction mixture was extracted with EtOAc (500 mL x 3) and the extracts were dried over MgSO₄, filtered, and concentrated to dryness under reduced pressure. The residue was purified by silica gel column chromatography (EtOAc : hexane = 1:10) to give **5** (129 g, 94%) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 6.01 (m, 0.8H), 5.79 (m, 0.2H), 5.50 (d, J = 2.8 Hz, 0.8H), 5.43-5.16 (m, 2.2H), 4.70-4.56 (m, 3H), 3.93 (d, J = 10.4 Hz, 0.2H), 2.66 (d, J = 2.8 Hz, 0.8H), 1.59 (s, 0.6H), 1.51 (s, 2.4H), 1.39 (s, 0.6H), 1.33 (s, 2.4H); ¹³C NMR (100 MHz, CDCl₃) δ 137.94, 134.43, 117.31, 117.01, 114.32, 112.45, 102.95, 102.91, 96.15, 88.54, 86.62, 84.73, 80.58, 79.01, 26.42, 26.19, 24.95; Anal. Calcd for C₉H₁₄O₄: C, 58.05 H, 7.58. Found: C, 58.38; H, 7.74.

(1S,2S,3R)-1-(2,2-Dimethyl-5-vinyl[1,3]dioxolan-4-yl)-(S)-prop-2-en-1-ol (6). A suspension of methyltriphenylphosphonium bromide (562 g, 1.55 mol) in tetrahydrofuran (1000 mL) was cooled to 0 °C and KO^tBu (210 g, 1.86 mol) was added portion wise under

nitrogen atmosphere. The reaction mixture was stirred at room temperature for 2 h and then recooled to 0 °C. A solution of lactol **5** (116 g, 0.622 mol) in tetrahydrofuran (300 mL) was added to the resulting reaction mixture at 0 °C. The mixture was stirred for 6 h at rt, diethyl ether (1500 mL) was added to the reaction mixture, and washed with H₂O (500 mL) and brine (500 mL). The organic layer was dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by silica gel column chromatography (EtOAc : hexane = 1:10) to give **6** (101 g, 88%) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 6.05 (m, 2H), 5.36 (m, 4H), 4.70 (t, *J* = 6.8 Hz, 1H), 4.19 (m, 1H), 4.04 (dd, *J* = 6.8 and 7.6 Hz, 1H), 1.79 (bs, 1H), 1.50 (s, 3H), 1.38 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 137.89, 134.24, 118.47, 116.63, 109.06, 80.72, 78.88, 71.20, 27.80, 25.46; Anal. Calcd for C₁₀H₁₆O₃: C, 65.19; H, 8.75. Found: C, 65.34; H, 8.43.

(4*R*,5*R*)-4,5-*O*-Isopropylidene-cyclopent-2-enone (7). To a solution of first generation Grubb's catalyst (4.198 g, 5.10 mmol, 1 mol%) in anhydrous CH₂Cl₂ (300 mL), diene **6** (94 g, 510.2 mmol) in anhydrous CH₂Cl₂ (1500 mL) was added and stirred for 7 h at rt under nitrogen atmosphere. To the mixture 4 Å molecular sieves (94 g), pyridinium dichromate (288 g, 765.3 mmol) and acetic acid (1.53 mL, 25.5 mmol) were added. The resulting brown mixture was stirred at rt for 12 h and filtered through a celite pad with CH₂Cl₂. The filtrate was concentrated in vacuo and the residue was purified by column chromatography on a silica gel (EtOAc:hexane = 1:10) to give **7** (72 g, 91%) as a white crystalline solid. mp 68.5-70.3 °C; [α]_D²³ -69.3 (c 0.60, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.61 (dd, *J* = 2.4, 6.0 Hz, 1H), 6.22 (d, *J* = 6.0 Hz, 1H), 5.28 (dd, *J* = 2.4, 5.6 Hz, 1H), 4.47 (d, *J* = 5.6 Hz, 1H), 1.42 (s, 3H), 1.41 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 159.56, 134.64, 115.36, 78.58, 76.49, 27.40, 26.14; Anal. Calcd for C₁₀ H₁₀O₃: C, 62.33; H, 6.54. Found: C, 62.15; H, 6.52.

(1*S*,2*S*,3*R*,4*R*)-4-(*tert*-Butoxymethyl)-2,3-(isopropylidenedioxy)-1-cyclopentanone (8).

A suspension of potassium *tert*-butoxide (41.6 g, 0.35 mol) in anhydrous *tert*-butylmethyl ether (1200 mL) was cooled to -78 °C under nitrogen. The well stirring mixture was treated with *sec*-butyllithium (1.6 M solution in hexane, 250 mL, 0.35 mol) over 20 min. After the mixture was stirred for 2 h at -78 °C, a solution of LiBr (61 g, 0.70 mol) in THF (450 mL) was added drop wise over 20 min at -78 °C and the resulting solution was stirred at -15 °C for 30 min. Upon recooling to -78 °C, a solution of CuBr·SMe₂ (37.3 g, 0.14 mol) in diisopropyl sulfide (150 mL) was added drop wise over 20 min. The resulting viscous dark solution was stirred for 1 h at -78 °C and treated with a solution of enone **7** (18.1 g, 0.117 mol) in THF (150 mL) over 15 min. The reaction mixture was allowed to warm to -30 °C over 15 min and

stirred for 30 min at -30 °C. The mixture was recooled to -78 °C and 1:1 acetic acid-methanol (415 mL) was added and the mixture poured in to 3000 mL of NH₄Cl/NH₄OH (1:1). After removal of the aqueous layer, the organic layer was washed with a saturated aqueous NH₄Cl solution (300 mL x 2) and brine (400 mL). The organic phase was dried (Na₂SO₄), filtered, concentrated and purified through silica gel column chromatography with 5 % EtOAc in hexanes to give **8** (25 g, 88%) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ 1.10 (s, 9H), 1.34 (s, 3H), 1.43 (s, 3H), 2.06 (d, *J* = 22 Hz, 1H), 2.55 (d, *J* = 12.5 Hz, 1H), 2.72 (dd, *J* = 11, 22 Hz, 1H), 3.34 (dd, *J* = 3.5, 10.5 Hz, 1H), 3.52 (dd, *J* = 3.5, 10.5 Hz, 1H), 4.22 (d, *J* = 7 Hz, 1H), 4.61 (d, *J* = 7 Hz, 1H); Anal. Calcd for C₁₃H₂₂O₄: C, 64.44; H, 9.15. Found: C, 64.19; H, 9.14.

(1S,2S,3R,4R)-4-(tert-Butoxymethyl)-2,3-(isopropylidenedioxy)-cyclopentan-1-ol (9). A solution of compound **8** (25 g, 0.103 mol) and CeCl₃·7H₂O (49.96 g, 0.134 mol) in MeOH (200 mL) was stirred 15 min at rt. The mixture was cooled to -78 °C, NaBH₄ (5.07 g, 0.134 mol) was slowly added and allowed the mix warm to 0 °C. After stirring 1 h at 0 °C, cold water (20 mL) was added and the methanol was removed under reduced pressure. The residue was extracted with EtOAc (2 X 600 mL), the combined organic phase was washed with brine (500 mL) and then dried over anhydrous Na₂SO₄, and filtered. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (15% EtOAc:hexane) to give **9** (24 g, 95%) as a syrup. [α]_D²⁶ -16.95 (c 1.59, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 1.13 (s, 9H), 1.34 (s, 3H), 1.48 (s, 3H), 1.83 (m, 2H), 2.19 (m, 1H), 2.44 (d, *J* = 9.0 Hz, 1H, D₂O exchangeable), 3.20 (dd, *J* = 4.5, 8.8 Hz, 1H), 3.31 (dd, *J* = 4.5, 8.8 Hz, 1H), 4.23 (m, 1H), 4.44 (m, 2H); Anal. Calcd for C₁₃H₂₄O₄: C, 63.91; H, 9.90. Found: C, 64.09; H, 9.87.

(1S,2S,3R,4R)-4-(tert-Butoxymethyl)-2,3-(isopropylidenedioxy)-1-(methylsulfonyloxy)-cyclopentane (10). A mixture of compound **9** (22 g, 90.04 mmol) and triethylamine (37.4 mL, 270.14 mmol) in CH₂Cl₂ (400 mL), methane sulfonyl chloride (10.41 mL, 135.07 mmol) was added drop wise at 0 °C. After stirring for 1 h at 0 °C, the reaction mixture was quenched with cold water (300 mL) and extracted with CH₂Cl₂ (2 x 500 mL). The combined organic layers were washed with brine (400 mL) and dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure and purified using column chromatography with EtOAc:hexane (1:1) to give **10** (quantitative yield) as a colorless oil.

(1R,2S,3R,4R)-1-Azido-4-(tert-butoxymethyl)-2,3-(isopropylidenedioxy)-cyclopentane (11). A solution of **10** (30 g, 93.04 mmol) in dry DMF (400 mL) in the presence of sodium

azide (60.49 g, 930.4 mmol) was heated at 140 °C for 5 h with stirring. After cooling to room temperature, the reaction mixture was filtered, and the filtrate was concentrated to dryness. The residue was diluted with EtOAc (500 mL), washed with water (2 x 300 mL) and brine (200 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated to dryness. The resulting oil was purified by silica gel column chromatography (4% EtOAc in hexane) to give **11** (22.2 g, 89%) as a colorless oil. $[\alpha]_D^{26} -46.94^0$ (c 1.11, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 1.18 (s, 9H), 1.30 (s, 3H), 1.46 (s, 3H), 1.71 (m, 1H), 2.29 (m, 2H), 3.29 (dd, *J* = 6.7, 8.8 Hz, 1H), 3.37 (dd, *J* = 7.0, 8.8 Hz, 1H), 3.96 (m, 1H), 4.40 (dd, *J* = 2.3, 6.1 Hz, 1H), 4.48 (dd, *J* = 2.0, 6.1 Hz, 1H); Anal. Calcd for C₁₃H₂₃N₃O₃·0.13EtOAc: C, 57.95; H, 8.65; N, 14.99. Found: C, 58.25; H, 8.71; N, 14.76.

(1*R*,2*S*,3*R*,4*R*)-4-(*tert*-Butoxymethyl)-2,3-(isopropylidenedioxy)-1-cyclopentanamine

(12). A suspension of compound **11** (20 g) and 10% Pd/C (5.0 g) in absolute EtOH (250 mL) was shaken under 35 psi of H₂ for 5 h. The reaction mixture was filtered and the filtrate was evaporated to give crude **12** (quantitative yield) which was used for next step without further purification. ¹H NMR (400 MHz, CDCl₃) δ 1.18 (s, 9H), 1.28 (s, 3H), 1.36 (m, 1H), 1.45 (s, 3H), 1.89 (br s, 2H, NH₂), 2.24-2.36 (m, 2H), 3.34-3.43 (m, 3H), 4.21 (dd, *J* = 2.6, 6.2 Hz, 1H), 4.48 (dd, *J* = 2.8, 6.2 Hz, 1H); Anal. Calcd for C₁₃H₂₅NO₃·0.16H₂O: C, 63.41; H, 10.37; N, 5.69. Found: C, 63.09; H, 10.16; N, 5.59.

N-[(1*R*,2*S*,3*R*,4*R*)-4-(*tert*-Butoxymethyl)-2,3-(isopropylidenedioxy)-cyclopentyl]-

aminocarbonyl}-3-methoxy-2-propenamide (13). A solution of β-methoxy acrylic acid in a mixture of CH₂Cl₂ : benzene (2:1) (600 mL) was cooled to 0 °C, then anhydrous DMF (1.2 mL) was added followed by oxalyl chloride (90 mL) was added at 0 °C under nitrogen atmosphere. The mix was stirred at 0 °C for 1 h and the solution was concentrated under reduced pressure to give the crude β-methoxy acryloyl chloride. Silver cyanate (77 g) was added to a solution of above obtained crude β-methoxy acryloyl chloride in anhydrous benzene (300 mL). The resulting mixture was heated under reflux for 1 h and allowed to cool to room temperature. After the solid phase had settled, the supernatant solution was contained with β-methoxy acryloyl isocyanate was added during 15 min to a solution of amine **12** (18 g) in anhydrous DMF (50 mL) at -15 to -20 °C under nitrogen. The reaction mixture was stirred for 2 h at -15 °C and then for 10 h at room temperature under nitrogen. The solvent was removed under reduced pressure and purified using silica gel column chromatography to afford **13** (19.8 g, 72%) as a solid. $[\alpha]_D^{27} -26.62$ (c 0.57, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 1.17 (s, 9H), 1.28 (s, 3H), 1.47 (s, 3H), 1.58 (m, 1H), 2.28 (m, 1H), 2.36-2.43 (m,

1H), 3.33-3.42 (m, 2H), 3.73 (s, 1H), 4.20 (m, 1H), 4.45 (m, 2H), 5.35 (d, $J = 12.3$ Hz, 1H), 7.67 (d, $J = 12.3$ Hz, 1H), 8.72 (br s, 1H, NH), 9.35 (br s, 1H, NH); Anal. Calcd for $C_{18}H_{30}N_2O_6$: C, 58.36; H, 8.16; N, 7.56. Found: C, 58.28; H, 8.13; N, 7.60.

(1R,2S,3R,4R)-1-[4-(*tert*-Butoxymethyl)-2,3-(isopropylidenedioxy)-cyclopentan-1-yl]-uracil (14). Compound 13 (18 g, 11.34 mmol) was dissolved in ethanol (80 mL), dioxane (60 mL) and ammonium hydroxide (30%, 80 mL). The reaction mixture was heated at 100 °C in a steel bomb for 20 h. After cooling, the solution was evaporated to dryness. The residue was purified by silica gel column chromatography (50% EtOAc in hexane) to give 14 (14 g, 85%) as a white foam. $[\alpha]_D^{27}$ -41.53 (c 0.88, $CHCl_3$); UV (MeOH) λ_{max} 266.0 nm; 1H NMR (400 MHz, $CDCl_3$) δ 1.19 (s, 9H), 1.30 (s, 3H), 1.54 (s, 3H), 1.97 (m, 1H), 2.32-2.41 (m, 2H), 3.43-3.50 (m, 2H), 4.48 (dd, $J = 4.1, 6.5$ Hz, 1H), 4.65-4.75 (m, 2H), 5.72 (d, $J = 8.0$ Hz, 1H), 7.35 (d, $J = 8.0$ Hz, 1H), 8.63 (br s, 1H, NH); Anal. Calcd for $C_{17}H_{26}N_2O_5$: C, 60.34; H, 7.74; N, 8.28. Found: C, 60.06; H, 7.70; N, 8.14.

(1R,2S,3R,4R)-1-[4-(*tert*-Butoxymethyl)-2,3-(isopropylidenedioxy)-cyclopentan-1-yl]-cytosine (15). A mixture of 14 (8 g, 23.6 mmol) and 4-dimethylamino-pyridine (5.78 g, 47.3 mmol) in anhydrous acetonitrile (250 mL), triethylamine (6.59 mL, 47.3 mmol) followed by 2,4,6-triisopropylbenzenesulfonyl chloride (14.33 g, 47.3 mmol) was added at rt under nitrogen. After being stirred for 12 h, NH_4OH (30%, 200 mL) was added and stirred for 5 hours. The mixture was diluted with chloroform and washed with saturated aqueous NH_4Cl (3 x 600 mL) solution and followed by washed with water (3 x 600 mL). The organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified using silica gel column chromatography (CH_2Cl_2 :MeOH = 96:4) to give 15 (6.22 g, 78%) as a solid. UV (MeOH) λ_{max} 275.0 nm; 1H NMR (500 MHz, $DMSO-d_6$) δ 1.14 (s, 9H), 1.22 (s, 3H), 1.44 (s, 3H), 1.80 (m, 1H), 2.10 (m, 2H), 3.30 (dd, $J = 6, 9$ Hz, 1H), 3.40 (dd, $J = 5.5, 9.5$ Hz, 1H), 4.38 (dd, $J = 5, 7$ Hz, 1H), 4.60 (m, 1H), 4.75 (dd, $J = 5, 6.5$ Hz, 1H), 5.71 (d, $J = 7.5$ Hz, 1H), 7.13 (brs, 1H), 7.22 (brs, 1H), 7.70 (d, $J = 7.5$ Hz, 1H); Anal. Calcd for $C_{17}H_{27}N_3O_4$: C, 60.51; H, 8.07; N, 12.45. Found: C 60.42; H 8.03; N 12.45.

(1R,2S,3R,4R)-1-[2,3-Dihydroxy-4-(hydroxymethyl)-cyclopentan-1-yl]cytosine (16).

Compound 15 (5.5 g) was dissolved in CF_3CO_2H/H_2O (2:1) (150 mL) and heated to 70 °C for 4 h. The solution was then concentrated under reduced pressure and coevaporated three times with methanol (3 x 200 mL). The residue was dissolved in water (300 mL) and washed with CH_2Cl_2 (4 x 100 mL). The water layer was concentrated under vacuum and the residue was dissolved in methanol and neutralized with IRA-400 (OH) basic resin. The resin was filtered,

washed with methanol and the filtrate was concentrated under reduced pressure to give **16** (2.75g, 82%) as a white solid. mp 214 °C; $[\alpha]_D^{27}$ -74.5 (c 0.52, H₂O); UV (H₂O) $\lambda_{\max}$ 275.0 nm (pH 7), 285 (pH 2), 275 (pH 11); ¹H NMR (500 MHz, DMSO-d₆) δ 1.23 (m, 1H), 1.94 (m, 1H), 2.01 (m, 1H), 3.42 (m, 2H), 3.80 (dd, J = 4.5, 9.0 Hz, 1H), 4.01 (m, 1H), 4.54 (d, J = 4.5 Hz, 1H), 4.59 (q, 1H), 4.65 (t, J = 5.0 Hz, 1H), 4.72 (d, J = 7 Hz, 1H), 5.69 (d, J = 7.5 Hz, 1H), 6.96 (brs, 1H), 7.01 (brs, 1H) 7.60 (d, J = 7.5 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 165.7, 156.6, 143.8, 93.8, 73.8, 72.1, 63.4, 61.7, 45.4, 29.0; Anal. Calcd for C₁₀H₁₅N₃O₄: C, 49.77; H, 6.27; N, 17.42. Found: C 49.67; H 6.31; N 17.43.

(1R,2S,3R,4R)-1-[2,3-Diacetoxy-4-(acetoxymethyl)-cyclopentan-1-yl]uracil (18).

Compound **14** (5.7 g) was dissolved in CF₃CO₂H/H₂O (2:1) (150 mL) and heated to 70 °C for 4 h. The solution was then concentrated under reduced pressure and coevaporated three times with methanol (3 x 200 mL). The residue containing **17** was dried under vacuum for overnight and it is used for next step directly. To a stirring solution of compound **17** (4.08, 16.85 mmol) in acetonitrile (100 mL), triethylamine (9.34 mL, 67.43 mmol), DMAP (41 mg, 0.33 mmol) followed by Ac₂O (6.38 mL) were added at rt and stirred for 30 min. To mixture methanol (5 mL) was added and stirred for 10 min. The solvent was evaporated by vacuum to obtain white crude product. This was purified using silica gel column chromatography (EtOAc:Hexane = 1:1) to give **18** (5.58 g, 90 %) as a foam. ¹H NMR (500 MHz, CDCl₃) δ 1.73 (m, 1H), 2.04 (s, 3H), 2.10 (s, 6H), 2.39 (m, 1H), 2.52 (m, 1H), 2.51 (m, 2H), 4.79 (q, 1H), 5.24 (d, J = 5 Hz, 1H), 5.43 (t, J = 8 Hz, 1H), 5.77 (d, J = 6 Hz, 1H), 7.20 (d, J = 7.5 Hz, 1H), 9.35 (s, 1H); ¹³C NMR (500 MHz, CDCl₃) δ 170.8, 170.1, 169.8, 163.0, 150.7, 141.7, 103.0, 73.1, 72.0, 64.2, 60.8, 40.0, 27.7, 20.8, 20.7, 20.6; HRMS-ESI (m/z): (M+H)⁺ calcd for C₁₆H₂₀N₂O₈, 369.1299; found, 369.1298.

(1R,2S,3R,4R)-1-[2,3-Diacetoxy-4-(acetoxymethyl)-cyclopentan-1-yl]-5-fluorouracil (19).

To a mixture of **18** (1.92 g, 5.21 mmol), selectfluor (2.03 g, 5.73 mmol) in acetonitrile (50 mL), AcOH (4 mL) was added at rt and stirred at 95 °C for 3 h. The mixture was diluted with ethyl acetate (200 mL), washed with H₂O (100 mL) and saturated NaHCO₃ solution (2 X 100 mL). The organic layer was dried over Na₂SO₄ and concentrated under reduced pressure and dried over night with high vacuum. The crude residue was treated with Et₃N (100 mL) and stirred at 95 °C for 3 h. The solvent was evaporated and purified using silica gel column chromatography (EtOAc:hexane = 4:6) to give **19** (1.44 g, 72 %) as a white foam. ¹H NMR (500 MHz, CDCl₃) δ 1.71 (m, 1H), 2.05 (s, 3H), 2.10 (s, 3H), 2.12 (s, 3H), 2.39 (m, 1H), 2.52 (m, 1H), 2.51 (m, 2H), 4.17 (dd, J = 5.5, 12 Hz, 1H), 4.21 (dd, J = 5.5, 12 Hz, 1H), 4.82 (q,

1H), 5.21 (t, $J = 6$ Hz, 1H), 5.38 (m, 1H), 7.33 (d, $J = 6$ Hz, 1H), 9.21 (s, 1H); ^{19}F NMR (CDCl_3 , 500 MHz) δ -164; HRMS-ESI (m/z): ($\text{M}+\text{H}$) $^+$ calcd for $\text{C}_{16}\text{H}_{19}\text{FN}_2\text{O}_8$, 387.1204; found, 387.1208.

(1R,2S,3R,4R)-1-[2,3-Dihydroxy-4-(hydroxymethyl)-cyclopentan-1-yl]-5-fluorocytosine (20). A mixture of **19** (3.5 g, 9.06 mmol) and 4-dimethylamino-pyridine (2.21 g, 18.13 mmol) in anhydrous acetonitrile (100 mL), triethylamine (2.5 mL, 18.13 mmol) followed by 2,4,6-triisopropylbenzenesulfonyl chloride (5.49 g, 18.13 mmol) was added at rt under nitrogen. After being stirred for 12 h, NH_4OH (30%, 150 mL) was added and stirred for 5 hours. The mix was concentrated under vacuum and the residue was purified using amine functionalized silica gel column chromatography ($\text{CH}_2\text{Cl}_2:\text{MeOH} = 5:1$) followed by C_{18} silica gel chromatography (water:methanol = 8:2) to give **20** (1.77 g, 76%) as a foam. UV (H_2O) λ_{max} 284 nm (pH 7), 293 nm (pH 2), 284 nm (pH 11); ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 1.21 (m, 1H), 1.92 (m, 1H), 2.01 (m, 1H), 3.42 (m, 2H), 3.72 (q, 1H), 3.99 (m, 1H), 4.54 (d, $J = 4.5$ Hz, 1H), 4.59 (q, 1H), 4.65 (t, $J = 5.5$ Hz, 1H), 4.76 (d, $J = 6.5$ Hz, 1H), 7.36 (brs, 1H), 7.60 (brs, 1H) 7.96 (d, $J = 7.5$ Hz, 1H); HRMS-ESI (m/z): ($\text{M}+\text{H}$) $^+$ calcd for $\text{C}_{17}\text{H}_{14}\text{FN}_3\text{O}_4$, 260.1047; found, 260.1044.

Biological Data

H5N1 Data

(-)-Carbodine (the D- nucleoside compound) and other enantiomers (the L- nucleoside compound as well as a racemic mixture) were tested in animals infected with Avian influenza A strain H5N1 virus at dilutions ranging from 1/100 to 1/1000 to determine inhibitory effect on the virus. The virus was that found in ducks, seagulls, a Hong Kong strain (HK/2003) ducks and a Vietnamese strain of H5N1 bird flu. In each of the four strains of H5N1 bird flu, (-)-Carbodine, (+)-Carbodine, a racemic mixture of the two enantiomers of Carbodine and ribivirin (an antiviral agent) were used. The results from the experiment appear in Table 1, attached in Figure 1. The experiment shows that (-) carbodine exhibited good activity as an inhibitor of Avian Influenza A strain H5N1.

In a separate experiment, the activity of 5-F carbodine was tested against avian influenza A H5N1 viruses or hybrid viruses. The results, which appear in attached Figure 2, Table 2, evidence that 5-F carbodine is quite active *in vitro* against avian Influenza A H5N1 strain.

VEE Data

In Vitro

(-) Carbodine was also tested *in vitro* against Venezuelan Equine Encephalitis (VEE) as well as yellow fever virus and H5N1. The results of that *in vitro* experiment appear in Figure 3, Table 3. (-) carbodine evidenced good activity against VEE as well as H5N1 in the *in vitro* experiment.

In Vivo

Effect of Later Post-virus Administration of i.p. (-) Carbodine on Survival of Mice Infected with Venezuelan Equine Encephalitis Virus.

[(-)-carbodine] is an enantiomerically pure D-nucleoside that has been shown to be active against the vaccine strain (TC-83) of Venezuelan equine encephalitis virus (VEEV) in a C3H/HeN mouse model. An important consideration in the treatment of viral encephalitis is whether an effective compound will be useful in treatment after the establishment of virus infection in the brain has occurred. Virus is detectable in the brain 2 days post-virus installation (dpi), peaks at 4 dpi, and continues at high titers until death at around 9 dpi in this model of VEEV disease. A similar course of virus infection of the brain is seen in mouse models utilizing a molecular clone of the pathogenic Trinidad Donkey (TrD) VEEV strain (1, 2). The main objective of this experiment is to determine if treatment initiated at the time of viral entry into the brain or when virus titers peak in the brain is effective in reducing mortality and other disease parameters in mice challenged intranasally (i.n.) with VEEV.

In previous experiments, ampligen has been used as a positive control compound. Positive results were generally obtained, but occasionally ampligen failed to improve disease parameters. This was likely due to incomplete mixing and solubilization of stock before aliquoting resulting in an unequal distribution of RNA between tubes. Another objective of this study was to reduce the experiment-to-experiment variability by preparing homogenous stock aliquots in sterile tubes.

Materials and Methods

Animals: Female C3H/HeN mice obtained from Charles River Laboratories (Wilmington, MA) between 16 and 18 g were used. Animals were randomly assigned to cages and individually marked with eartags. Mice were fed standard mouse chow and tap water *ad libitum*.

Test article: The compound (-) Carbodine was dissolved initially in DMSO, after which the solution was diluted in sterile saline to a final DMSO concentration of 10% and stored at 4°C until use. Ampligen was obtained as a viscous solution from Hemispherx Biopharma (Philadelphia, PA) and was used undiluted at 12 mg/kg/d. Ampligen was heated to 50°C prior to use to ensure a homogenous mixture of compound.

Venezuelan equine encephalitis virus: The TC-83 vaccine strain of VEE was obtained from ATCC and used after passage in vero cells. A 10^{-2} dilution ($10^{7.4}$ 50% cell culture infectious doses/ml) of the virus was prepared, and animals were inoculated i.n. with 0.05 ml of the diluted virus.

Experimental design: Mice were treated i.p. with (-) carbodine at a dose of 200 mg/kg/day starting 2 or 4 days post-intranasal virus installation (dpi). Placebo control mice were treated with 10% DMSO in saline on the same schedule as GYS-VI-16-23 treatment initiated -4 h prior to virus challenge. Mortality was checked daily for 21 days. Mice were weighed on the day of virus challenge and on 7 and 9 dpi. Ampligen was used as a positive control at 12 mg/treatment given -4 h and 2 dpi. A group of sham-infected mice treated bid, 2 to 9 dpi with 200 mg/kg/d of GYS-VI-16-23 was included as a toxicity control and weights were taken at 0, 7, and 9 dpi.

To reduce the variability of results obtained with ampligen treatment, ampligen stock solution was heated to 50°C for 15 minutes prior to aliquoting to ensure a homogenous mixture of RNA. The ampligen solution was gently mixed after heating by inverting the stock tube several times, aliquoted to sterile tubes, and frozen at -20°C.

Statistical analysis: Survival data were analyzed using the Wilcoxon log-rank survival analysis (JMP™ software, The Statistical Discovery Software, SAS Institute, Inc). All other statistical analysis was done using one-way Students T-test.

Results and Discussion

Treatment i.p. with 200 mg/kg/d of GYS-VI-16-23, beginning 2 dpi and continuing with twice-daily treatments through 9 dpi, was efficacious in significantly improving survival, mean day to death, and weight gain (Table 4, Fig 4). Later bid treatment with 200 mg/kg/d for 8 days beginning 4 dpi was also effective in significantly improving all disease signs measured. Virus peaks in the brain 4 dpi, so it is very remarkable that treatment initiated on the day of peak brain virus titer was effective in significantly improving disease. Virus titers remain constant in the brain after the peak at 4 dpi, and it would be interesting to see if treatment beginning later than 4 dpi is effective. Another study will be required to determine how long after virus challenge this compound will be efficacious.

It would also be interesting to see if treatment on or after 4 dpi would affect the brain titers in mice at later times during the course of infection. In a previous experiment (NIA-673, this report), titration of brain virus 4 dpi showed a 1-log reduction in virus titer. It would also be interesting to see if the titer continued to drop, if the reduction was sustained, or if virus rebounded. Carbodine is a known inhibitor of CTP synthase that converts UTP to CTP (3, 4) and results in a decrease of CTP pools, which affects replication of different viruses (5). If CTP pools are reduced, a steady decline in virus replication might be expected over the course of infection. Further studies will try to delineate the effect of treatment on the virus replication kinetics in the brain.

No toxicity was observed in this experiment with 200 mg/kg/d treatment beginning 2 or 4 days after virus infection as determined by weight change or mortality. In a previous experiment (NIA-673) weight loss was observed in sham infected mice treated with 200 mg/kg/d GYS-VI-16-23 given bid beginning 4 h prior to virus challenge. This could possibly be due to the timing of administration of the compound, with earlier treatment interacting in an unfavorable manner with virus infection parameters or host defense. This could also indicate a possible alternative mode of action of the compound, which may involve host

parameters, making the timing of treatment administration an important consideration. It could also be simply an artifactual result due to the low number of mice used in toxicity control groups.

Ampligen treatment was effective as determined by a significant improvement in survival, mean day to death, and weight change. Further experiments using the aliquots prepared with the same stock used in this experiment determine if the preparation measures reduce the variability seen in past experiments.

Conclusions


- (-) Carboline 200 mg/kg/d, given i.p., bid for 8 days was effective in significantly improving survival, weight gain, and mean day to death when treatment was initiated 2 or 4 dpi.
- No toxicity was apparent with (-) Carboline treatment in sham-infected toxicity controls as determined by weight change and mortality.
- Ampligen was also effective in significantly improving survival, mean day to death and weight change when administered i.p. -4 h and 2 dpi at a dose of 12 mg/treatment.

References

1. Schoneboom, B.A., K.M. Catlin, A.M. Marty, and F.B. Grieder. 2000. Inflammation is a component of neurodegeneration in response to Venezuelan equine encephalitis virus infection in mice. *J Neuroimmunol.* 109: 132-46.
2. Charles, P.C., J. Trgovcich, N.L. Davis, and R.E. Johnston. 2001. Immunopathogenesis and immune modulation of Venezuelan equine encephalitis virus-induced disease in the mouse. *Virology.* 284: 190-202.
3. De Clercq, E., J. Murase, and V.E. Marquez. 1991. Broad-spectrum antiviral and cytotoxic activity of cyclopentenylcytosine, a carbocyclic nucleoside targeted at CTP synthetase. *Biochem Pharmacol.* 41: 1821-9.
4. Neyts, J., A. Meerbach, P. McKenna, and E. De Clercq. 1996. Use of the yellow fever virus vaccine strain 17D for the study of strategies for the treatment of yellow fever virus infections. *Antiviral Res.* 30: 125-32.
5. De Clercq, E., R. Bernaerts, Y.F. Shealy, and J.A. Montgomery. 1990. Broad-spectrum antiviral activity of carbodine, the carbocyclic analogue of cytidine. *Biochem Pharmacol.* 39: 319-25.

Claims:

1. A nucleoside compound according to the structure:




Wherein R is H or F;

R¹ and R^{1a} are each independently H, an acyl group, a C₁—C₂₀ alkyl or ether group, an amino acid residue, a phosphate, diphosphate, triphosphate or phosphodiester group or together R¹ and R^{1a} form a carbodiester or phosphodiester group with the oxygen atoms to which they are bonded;

R² is H, an acyl group, a C₁—C₂₀ alkyl or ether group or an amino acid residue; or a pharmaceutically acceptable salt, enantiomer, hydrate or solvate thereof.


2. The compound according to claim 1 wherein R^{1a} is H and R¹ and R² are both H.
3. The compound according to claim 1 or 2 wherein R is F.
4. The compound according to claim 1 or 2 wherein R is H.
5. The compound according to claim 1, 3 or 4 wherein R^{1a} and R² are H.
6. The compound according to claim 1 or 3-5 wherein R¹ is an acyl group, a phosphate or phosphodiester group.

7. The compound according any of claims 1 or 3-6 wherein R^1 together with the nucleoside to which it is attached forms a group according to the structure:



where each R^5 , R^6 and $R^{\text{''}}$ is independently selected from H, a C_1 to C_{20} linear, branched or cyclic alkyl group, alkoxyalkyl, aryloxyalkyl, such as phenoxymethyl, aryl and alkoxy, with the proviso that at least one R^5 group is other than H, or the two R^5 groups together form a five- or six-membered heterocyclic group, B is a direct bond or a C_1 - C_3 alkylene group optionally substituted with a C_1 - C_3 alkyl group and R^7 is a C_1 to C_{20} linear, branched or cyclic alkyl or acyl group, alkoxyalkyl, aryloxyalkyl, such as phenoxymethyl, aryl and alkoxy, among others, each of which groups previously mentioned may be optionally substituted.

8. The compound according any of claims 1 or 3-6 wherein R^1 and R^2 together with the nucleoside to which those groups are attached form a group according to the structure:



where R^6 is selected from H, a C_1 to C_{20} linear, branched or cyclic alkyl group, alkoxyalkyl, aryloxyalkyl, aryl and alkoxy.

9. A pharmaceutical composition comprising an effective amount of a compound according to any of claims 1-8, optionally in combination with a pharmaceutically acceptable carrier, additive or excipient.

10. A method of treating an influenza viral infection or a Venezuelan equine encephalitis viral infection in a patient in need of therapy or at risk for infection thereof comprising administering to said patient an effective amount of a compound according to any of claims 1-8 to said patient.

11. The method according to claim 10 wherein said infection is an Avian influenza A H5N1 strain viral infection.

12. The method according to claim 10 wherein said infection is a Venezuelan equine encephalitis viral infection.

13. A method of reducing the likelihood of an influenza viral infection or a Venezuelan equine encephalitis viral infection in a patient in need of therapy or at risk for infection thereof comprising administering to said patient an effective amount of a compound according to any of claims 1-8 to said patient.

14. The method according to claim 13 wherein said infection is an Avian influenza A H5N1 strain viral infection.

15. The method according to claim 13 wherein said infection is a Venezuelan equine encephalitis viral infection.

16. Use of a compound according to any of claims 1-8 for the manufacture of a medicament for the treatment of an influenza A viral infection or a Venezuelan equine encephalitis viral infection.

17. Use according to claim 16 wherein said viral infection is an influenza A strain H5N1 viral infection.

18. Use according to claim 16 wherein said viral infection is a Venezuelan equine encephalitis viral infection.

19. Use of a compound according to any of claims 1-8 for the manufacture of a medicament for reducing the likelihood of an influenza A viral infection or a Venezuelan equine encephalitis viral infection in a patient at risk.
20. Use according to claim 19 wherein said viral infection is an influenza A strain H5N1 viral infection.
21. Use according to claim 16 wherein said viral infection is a Venezuelan equine encephalitis viral infection.
22. The method according to any of claims 11-15 wherein said patient is a human.
23. The method according to any of claims 11-15 wherein said patient is a domesticated animal.
24. The use according to any of claims 16-21 wherein said patient is a human.
25. The composition according to claim 9 further including an additional anti-viral agent.
26. The composition according to claim 25 wherein said additional anti-viral agent is oseltamavir phosphate.
27. The method according to claim 14 wherein said compound according to any of claims 1-8 is coadministered with oseltamavir phosphate.
28. Use according to claim 17 wherein said compound is combined with oseltamavir phosphate.
29. Use according to claim 20 wherein said compound is combined with oseltamavir phosphate.

FIGURE 1

Table 1 COMPARISON OF the Activity of CARBODINE ENANTIOMERS

VIRUS	DILUTION	DRUG	VISUAL			NEUTRAL			VYR- EC90
			EC50	IC50	SI	N. RED	RED	N. RED	
DUCK	1/1000	Neg, Enantiomer	0.57	>100	175	0.87	IC50	SI	0.22
DUCK	1/1000	Pos. Enantiomer	>100	>100	>1	>100	>100	>1	>100
DUCK	1/1000	Racemic							
DUCK	1/1000	Carbodine	2.7	>100	>37	5.3	>100	>19	11.4
DUCK	1/1000	ribivirin	5.7	>320	>56	5.8	>320	>55	6.5
GULL	1/100	Neg, Enantiomer	0.27	>100	370	0.31	>100	323	0.22
GULL	1/100	Pos. Enantiomer	>100	>100	>1	>100	>100	>1	>100
GULL	1/100	Racemic							
GULL	1/100	Carbodine	1.8	>100	>56	1.9	>100	>53	1.3
GULL	1/100	ribivirin	5.7	>320	>56	5.9	>320	>54	3.7
HK/2003	1/100	Neg, Enantiomer	0.18	>100	556	0.3	>	333	0.35
HK/2003	1/100	Pos. Enantiomer	>100	>100	>1	>100	>100	>1	>100
HK/2003	1/100	Racemic							
HK/2003	1/100	Carbodine	0.7	>100	>143	1.7	>100	>59	4.6
HK/2003	1/100	ribivirin	0.57	>320	>561	1.4	>320	>229	3.5
VN	1/500	Neg, Enantiomer	0.18	>100	>556	0.19	>100	526	0.11
VN	1/500	Pos. Enantiomer	>100	>100	>1	>100	>100	>1	>100
VN	1/500	Racemic							
VN	1/500	Carbodine	0.57	>100	>175	0.92	>100	>109	1.9
VN	1/500	ribivirin	0.39	>320	>821	0.59	>320	>542	0.44


FIGURE 2

Table 2 Effects of 5-F-carbodine on avian influenza A H5N1 viruses or hybrid viruses.

H5N1 Influenza A Strain	Visual Assay			Neutral Red Uptake Assay			Virus Yield Reduction Assay
	EC50 (µg/ml)	IC50 (µg/ml)	SI	EC50 (µg/ml)	IC50 (µg/ml)	SI	
Duck MN/1525/81 Trial 1	2.4	>100	>42	2	>100	>50	ND
Duck MN/1525/81 Trial 2	1.1	>100	>91	2.4	>100	>42	24
Vietnam/1203/04 X H1N1 Ann Arbor/6/60	0.81	>100	>120	0.86	>100	>120	ND

Figure 3

Table 3. *In vitro* antiviral activity of (-)-Carbodine²⁰

Structure	Viruses	EC ₅₀ (μM)	IC ₅₀ (μM)
	Venezuelan Equine Encephalitis	0.2	>100
	Yellow Fever	1.8	>100
	Flu A (DUCK)-H5N1	0.87	>100

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 08/04568

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/70, 31/519 (2008.04)

USPC - 514/49; 514/264.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC - 514/49; 514/264.1 (see search terms below)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC - 514/43; 514/256 (see search terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

USPTO-WEST - PGPB,USPT,USOC,EPAB,JPAB keywords: carbodine, influenza, avian, antiviral activity, type A, treatment, humans, birds, ducks, cyclopentan-1-yl, cytosine, infect, 5-fluorocarbodine, 5-fluorocytosine, nucleosides.

INTERNET search - Google - same as for USPTO-WEST.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2003/0087873 A1 (STUYVER et al.) 08 May 2003 (08.05.2003), para [0106]-[0107]; [0109]; [0113]-[0123]; [0128]-[0131].	1-4

☐

Further documents are listed in the continuation of Box C.

☐

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

06 July 2008 (06.07.2008)

Date of mailing of the international search report

14 JUL 2008

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 08/04568

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 5-29
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.