



US00PP29154P2

(12) **United States Plant Patent**  
**Koot**

(10) **Patent No.:** **US PP29,154 P2**

(45) **Date of Patent:** **Mar. 27, 2018**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED ‘DUETAMARCHERBLOS’**

(50) Latin Name: *Impatiens hawkeri*  
Varietal Denomination: **Duetamarcherblos**

(71) Applicant: **DUMMEN GROUP B.V.**, De Lier (NL)

(72) Inventor: **Arjan Koot**, Oeffelt (NL)

(73) Assignee: **Dümmen Group B.V.**, De Lier (NL)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/530,621**

(22) Filed: **Feb. 9, 2017**

(51) **Int. Cl.**  
**A01H 5/02** (2006.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./318.1**

(58) **Field of Classification Search**  
USPC ..... Plt./318.1, 318.4, 318.5, 318.6  
CPC ..... A01H 5/0261  
See application file for complete search history.

*Primary Examiner* — Kent L Bell

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Impatiens* plant named ‘Duetamarcherblos’ characterized by its upright, outwardly spreading and mounding plant habit; moderately vigorous growth habit; medium in size; freely branching habit; dark green-colored leaves; freely and early flowering habit; large red purple and light red purple bi-colored flowers; and good garden performance.

**1 Drawing Sheet**

**1**

Botanical designation: *Impatiens hawkeri*.  
Cultivar denomination: ‘DUETAMARCHERBLOS’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of New Guinea *Impatiens* plant, botanically known as *Impatiens hawkeri* and hereinafter referred to by the name ‘Duetamarcherblos’.

The new *Impatiens* plant is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. The objective of the breeding program is to create new medium-sized New Guinea *Impatiens* plants with large attractive flowers.

The new *Impatiens* plant originated from a cross-pollination made by the Inventor in July, 2011 in Rheinberg, Germany of a proprietary selection of *Impatiens hawkeri* identified as code number NN09-002148-001, not patented, as the female, or seed, parent with a proprietary selection of *Impatiens hawkeri* identified as code number NN-1041, not patented, as the male, or pollen, parent. The new *Impatiens* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rheinberg, Germany in May, 2016.

Asexual reproduction of the new *Impatiens* plant by terminal cuttings in a controlled greenhouse environment in Rheinberg, Germany since June, 2016 has shown that the unique features of this new *Impatiens* plant are stable and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

Plants of the new *Impatiens* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with

**2**

variations in environmental conditions such as temperature, daylight and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Duetamarcherblos’. These characteristics in combination distinguish ‘Duetamarcherblos’ as a new and distinct *Impatiens* plant:

1. Upright, outwardly spreading and mounding plant habit.
2. Moderately vigorous growth habit; medium in size.
3. Freely branching habit.
4. Dark green-colored leaves.
5. Freely and early flowering habit.
6. Large red purple and light red purple bi-colored flowers.
7. Good garden performance.

Plants of the new *Impatiens* can be compared to plants of the female parent selection. Plants of the new *Impatiens* differ primarily from plants of the female parent selection in flower color as plants of the female parent selection have blue and red bi-colored flowers. In addition, plants of the new *Impatiens* have larger flowers than plants of the female parent selection.

Plants of the new *Impatiens* can be compared to plants of the male parent selection. Plants of the new *Impatiens* differ primarily from plants of the male parent selection in flower size as plants of the male parent selection have smaller flowers.

Plants of the new *Impatiens* can be compared to plants of *Impatiens hawkeri* ‘Kicoiba’, not patented. In side-by-side comparisons, plants of the new *Impatiens* differ primarily from plants of ‘Kicoiba’ in the following characteristics:

1. Plants of the new *Impatiens* are larger than plants of ‘Kicoiba’.
2. Plants of the new *Impatiens* are more freely branching than plants of ‘Kicoiba’.

3. Plants of the new *Impatiens* have longer leaves than plants of 'Kicoiba'.
4. Plants of the new *Impatiens* have larger flower than plants of 'Kicoiba'.
5. Plants of the new *Impatiens* and 'Kicoiba' differ in flower color as plants of 'Kicoiba' have light red purple-colored flowers.
6. Plants of the new *Impatiens* have longer peduncles than plants of 'Kicoiba'.

## BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Impatiens* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Impatiens* plant.

The photograph comprises a side perspective view of a typical flowering plant of 'Duetamarcherblös' grown in a container.

## DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown during the summer in 22-cm containers in a glass-covered greenhouse in Rheinberg, Germany and under cultural practices typical of commercial New Guinea *Impatiens* production. During the production of the plants, day and night temperatures averaged 18° C. and light levels averaged 4,500 lux. Plants were 13 weeks old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* 'Duetamarcherblös'.

Parentage:

*Female, or seed, parent.*—Proprietary selection of *Impatiens hawkeri* identified as code number NN09-002148-001, not patented.

*Male, or pollen, parent.*—Proprietary selection of *Impatiens hawkeri* identified as code number NN-1041, not patented.

Propagation:

*Type.*—By terminal cuttings.

*Time to initiate roots, summer.*—About five days at temperatures about 20° C.

*Time to initiate roots, winter.*—About seven days at temperatures about 20° C.

*Time to produce a rooted young plant, summer.*—About three weeks at temperatures about 20° C.

*Time to produce a rooted young plant, winter.*—About four weeks at temperatures about 20° C.

*Root description.*—Fine, fibrous; white in color.

*Rooting habit.*—Freely branching; dense.

Plant description:

*Plant and growth habit.*—Upright to outwardly spreading and mounding plant habit; medium in size; freely branching habit with about 50 lateral branches developing per plant; moderately vigorous growth habit.

*Plant height.*—About 38 cm.

*Plant diameter.*—About 71 cm.

Lateral branch description:

*Length.*—About 20 cm to 35 cm.

*Diameter.*—About 6 mm to 8 mm.

*Internode length.*—About 4.9 cm.

*Strength.*—Strong.

*Aspect.*—Initially upright to outwardly spreading.

*Texture.*—Smooth, glabrous.

*Color, when developing.*—Close to 146C.

*Color, fully developed.*—Close to 146B.

10 Leaf description:

*Arrangement.*—Opposite or in whorls; simple.

*Length.*—About 9.1 cm.

*Width.*—About 3.5 cm.

*Shape.*—Ovate.

*Apex.*—Apiculate.

*Base.*—Obtuse.

*Margin.*—Serrulate with ciliation.

*Texture, upper and lower surfaces.*—Smooth, glabrous; leathery.

*Venation pattern.*—Pinnate; arcuate.

*Color.*—Developing leaves, upper surface: Close to 146A. Developing leaves, lower surface: Close to 59A. Fully expanded leaves, upper surface: Close to 139A; venation, close to 187A. Fully expanded leaves, lower surface: Close to 187A; venation, close to 59A.

*Petioles.*—Length: About 2.3 cm. Diameter: About 3 mm. Texture, upper and lower surfaces: Smooth, glabrous; leathery. Color, upper surface: Close to 185B. Color, lower surface: Close to 185A.

Flower description:

*Flower type and flowering habit.*—Large single rounded and flat axillary flowers; freely flowering habit, typically about five open flowers and flower buds per lateral branch and about 180 to 220 flowers potentially developing per plant; flowers positioned above and beyond the foliar plane, flowers typically face mostly upright to outwardly.

*Flower longevity.*—Flowers typically last about one to two days on the plant under greenhouse conditions; petals self-cleaning, gynoecium persistent.

*Fragrance.*—None detected.

*Natural flowering season.*—Year-round under greenhouse conditions; in the garden, flowering from spring until fall in Germany; early flowering habit, plants typically begin flowering about eight weeks after planting.

*Flower buds.*—Length: About 1.3 cm. Diameter: About 7.8 mm. Shape: Ovate. Texture: Smooth, glabrous. Color: Close to 63A.

*Flower diameter.*—About 5.8 cm.

*Flower depth.*—About 3 cm.

*Petals.*—Quantity and arrangement: Five per flower in a single whorl. Length: About 3.4 cm. Width: About 4.1 cm. Shape: Obovate. Apex: Cordate; rounded. Base: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; satiny. Color: When opening, upper surface: Close to 69A and N66B. When opening, lower surface: Close to 69B and N57A. Fully opened, upper surface: Close to 69B and N57B; color becoming closer to N155C with development. Fully opened, lower surface: Close to 69B and N57A; color does not change with development.

*Sepals.*—Quantity and arrangement: Three in a single whorl; one modified into an elongated spur. Length:

About 1.3 cm. Width: About 7 mm. Shape: Ovate. Apex: Apiculate. Base: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; leathery. Color: When opening, upper surface: Close to 147D. When opening, lower surface: Close to 60B. Fully opened, upper surface: Close to 65B. Fully opened, lower surface: Close to 63D. Spur length: About 5.4 cm. Spur diameter: At the flower, about 2.9 mm. Spur texture: Smooth, glabrous. Spur color: Close to 145C.

*Peduncles*.—Length: About 5.5 cm. Diameter: About 2.2 mm. Angle: Upright to outward. Strength: Moderately strong; flexible. Texture: Smooth, glabrous. Color: Close to 145A.

*Reproductive organs*.—Stamens: Quantity: Five fused at anthers; filaments free. Filament length: About 1.2 mm. Filament color: Close to N155D. Anther length: About 3.2 mm. Anther shape: Oval. Anther color:

Close to N155D and N57B. Pollen amount: Abundant. Pollen color: Close to 155D. Pistils: Quantity per flower: One. Pistil length: About 5 mm. Stigma shape: Crested. Stigma color: Close to 58A. Style length: About 1 mm. Style color: Close to 58A. Ovary color: Close to 149A.

*Seeds and fruits*.—Seed and fruit production has not been observed on plants of the new *Impatiens*.

Disease & pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to *Impatiens* plants.

Garden performance: Plants of the new *Impatiens* have been observed to have good garden performance and tolerate temperatures ranging from about 5° C. to about 40° C.

It is claimed:

1. A new and distinct *Impatiens* plant named ‘Duetamarcherblos’ as illustrated and described.

\* \* \* \* \*

