


1 574 477

(21) Application No. 40732/77 (22) Filed 30 Sep. 1977
 (31) Convention Application No's 51/118343 (32) Filed 1 Oct. 1976
 52/058369 20 May 1977
 52/092803 2 Aug. 1977 in



(33) Japan (JP)
 (44) Complete Specification Published 10 Sep. 1980
 (51) INT. CL.³ C07C 143/78
 A01N 41/06

(52) Index at Acceptance
 C2C 200 202 220 226 227 22Y 30Y 311
 313 314 315 31Y 323 326 32Y 338
 364 36Y 385 510 51X 51Y 52Y 534
 620 650 652 662 669 697 699 805
 80Y AA SC SF SG

(72) Inventors: TETSUO TAKEMATSU
 MAKOTO KONNAI
 HIROYOSHI OMOKAWA

(54) NOVEL BENZENESULFONAMIDE DERIVATIVES
 AND THEIR USE AS HERBICIDES

(71) We, UTSUNOMIYA UNIVERSITY, a Japanese National University, of 350
 Mine-machi, Utsunomiya-shi, Tochigiken, Japan, do hereby declare the invention for
 which we pray that a patent may be granted to us, and the method by which it is to be
 performed, to be particularly described in and by the following statement:-

5 This invention relates to certain novel N,N-disubstituted benzenesulfonamide derivatives, to a process for their preparation, and to their use as herbicides.


Numerous benzenesulfonamide derivatives have been known. For example, Swiss Patent
 10 No. 224,856 discloses that N-methyl-N-benzoyl-3,4-dichlorobenzene-sulfonamide is useful
 as an insecticide. U. S. Patent 3,245,913 states that N-methyl-N-benzoyl-substituted
 15 benzenesulfonamides are effective as bleaching agents. Japanese Patent Publication No.
 7216/74 suggests the utilization of N-methyl-N-benzoyl-3, 5-dibromo-6-acetoxy-
 benzenesulfonamide as a herbicide.

Elsewhere, Zh. Org. Chim., 7, 363 (1971) reports N-halogenated ethyl-N-substituted
 15 benzoyl-4-methylbenzene-sulfonamides as known compounds, and Ukrain. Khim. Zhur.,
 26, 496 (1960) reports N-isopropyl-N-benzoyl-p-chlorobenzene-sulfonamide as a known
 compound.

20 It has now been found in accordance with this invention that a certain group of novel
 N,N-disubstituted benzenesulfonamide derivatives are useful as herbicides, and have
 superior selective herbicidal activity against the emergency of barnyard grasses (especially,
 25 Echinochloa crusgalli Beauv.) without substantially affecting rice plants. This is surprising
 in view of the fact that since barnyard grass, the most hazardous weed in a paddy field, is a
 graminaceous grass, it has been considered as impossible to selectively control barnyard
 grass without causing phytotoxicity to rice.

The present invention provides a compound of the general formula

25



35 wherein R_O represents an alkyl group optionally having a substituent selected from a cyano group, lower alkoxy groups and di-(lower alkyl)amino groups, a lower alkenyl group, or a lower alkynyl group; R₁, R₂ and R₃, independently from each other, represent a hydrogen

5

10

15


20

25


30

35

atom, a halogen atom, a lower alkyl group, or a lower alkoxy group; R_4 , R_5 , R_6 , R_7 and R_8 , independently from each other, represent a hydrogen atom, a halogen atom, a lower alkyl group, or a lower alkoxy group; Y represents a group of the formula



10 in which one of R_9 and R_{10} is a lower alkyl group and the other is a hydrogen atom or a lower alkyl group; with the proviso that when Y is the group



and R_0 represents an unsubstituted alkyl group, R_4 , R_5 , R_6 , R_7 and R_8 do not represent hydrogen atoms at the same time.

20 In the present specification and the appended claims, the term "lower" means that a group or radical modified by this term contains up to 6, preferably 1 to 4, carbon atoms. 20

The term "alkyl group", as used in the present specification and the appended claims, denotes a saturated aliphatic hydrocarbon group which may be of straight chain or branched chain, and contain generally up to 20, preferably up to 15, more preferably up to 12, carbon atoms. Examples of the alkyl group include methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, n- or neo-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, n-undecyl, and n-dodecyl. Wherein R_0 in formula (I) represents an alkyl group, the alkyl group may contain a substituent. The substituent is selected from a cyano group (CN), lower alkoxy groups such as a methoxy or ethoxy group, and di(lower alkyl)amino groups such as a dimethylamino or diethylamino group. When either one of R_1 to R_8 represents a lower alkyl, it is desirably a methyl or ethyl group.

The term "lower alkenyl group" denotes a lower aliphatic hydrocarbon group which contains one carbon-to-carbon double bond, and may be of straight chain or branched chain. Examples are vinyl, allyl and butenyl groups, the allyl group being most preferred.

35 The term "lower alkynyl group" denotes a lower aliphatic hydrocarbon group containing one carbon-to-carbon triple bond. A propargyl group is a typical example of the lower alkynyl group.


The term "lower alkoxy group" means a lower alkyl ether group and includes, for example, methoxy, ethoxy, n- or iso-propoxy, n-, iso-, sec- or tert-butoxy. Of these, methoxy and ethoxy groups are preferred.

The term "halogen atom" denotes fluorine, chlorine, bromine and iodine atoms. Of these, a chlorine atom is most preferred, and next comes a fluorine atom. Thus, preferred substituents R_1 are alkyl groups containing up to 15 carbon atoms which


45 Thus, preferred substituents R_0 are alkyl groups containing up to 15 carbon atoms which may contain a substituent selected from a cyano group, lower alkoxy groups and di-(lower alkyl)amino groups; a lower alkenyl group; or a propargyl group. Of these, the lower alkenyl groups, above all, an allyl group, and the propargyl group are especially preferred.

Preferred species of R_1 , R_2 and R_3 are a hydrogen atom, a halogen atom, a methyl group, an ethyl group, a methoxy group, and an ethoxy group. Preferably, at least one of R_1 , R_2 and R_3 is a hydrogen atom, and moreover, at least one of these groups is desirably located at the 4-position of the benzene ring to which they are bonded.

Preferred atoms or groups represented by R_4 , R_5 , R_6 , R_7 and R_8 are a hydrogen atom, a halogen atom, a methyl group, an ethyl group, a methoxy group, and an ethoxy group, the hydrogen atom or halogen atom being especially preferred. When at least one of R_4 to R_8 represent the aforesaid groups other than hydrogen, one or two of them are desirably located at the 2-position and/or the 4-position on the benzene ring. When Y is a carbonyl group




it is most preferred that at least one, preferably one or two, of groups R₄ to R₈ be halogen and the remainder be hydrogen. If Y is a substituted methylene group



5

10 it is most convenient that all of groups R₄ to R₈ be hydrogen atoms at the same time.
Of the compounds of formula (I), a group of preferred compounds are those expressed by the formula


10



15

20 wherein R₀₁ represents an alkyl group containing up to 15 carbon atoms which may contain a substituent selected from a cyano group, lower alkoxy groups and di-(lower alkyl)amino groups, a lower alkenyl group, or a propargyl group; R₁₁ and R₂₁, independently from each other, represent a hydrogen atom, a halogen atom, a methyl group, an ethyl group, or a methoxy group; R₄₁, R₅₁, R₆₁, R₇₁ and R₈₁, independently from each other, represent a hydrogen atom, a halogen atom, a methyl group or a methoxy group; Y₁ represents the group

20




25

30 in which one of R₉₁ and R₁₀₁ is a methyl or ethyl group, and the other is a hydrogen atom or a methyl or ethyl group; with the proviso that when Y₁ is the group CO and R₀₁ is an unsubstituted alkyl group, R₄₁, R₅₁, R₆₁, R₇₁ and R₈₁ do not represent hydrogen atoms at the same time.

35


35 A group of more preferred compounds of formula (I-a) are those in which R₀₁ represents an allyl group or a propargyl group and Y₁ represents the group



40

45 A combination of the group

45



50



55

60 as Y₁ and an allyl group as R₀₁ are especially advantageous.
Thus, a group of especially preferred compounds of formula (I-a) are those in which R₀₁ represents an allyl or propargyl group; Y₁ represents the group

60




65

5 R_{41} is a halogen atom; and (a) R_{51} , R_{61} , R_{71} and R_{81} independently from each other, represent hydrogen or a methyl group and R_{11} and R_{21} , independently from each other, represent a hydrogen atom or a methyl group or (b) one of R_{51} , R_{61} , R_{71} and R_{81} , especially R_{61} , is a hydrogen or halogen atom and the rest are hydrogen atoms. In case (b), it is preferred that R_{11} and R_{21} , independently from each other, represent a hydrogen atom or a methyl group, especially hydrogen.

5

Another group of especially preferred compounds of formula (I-a) are those in which R_{01} is an allyl group, Y_1 is the group



10

15 R_{11} represents a hydrogen atom, a halogen atom, or a methyl or methoxy group, R_{21} is a hydrogen atom, R_{41} and R_{61} , independently from each other, represent a hydrogen atom, a halogen atom or a methyl group, and R_{51} , R_{71} and R_{81} are all hydrogen atoms.


20 As will be described in detail hereinbelow, the compounds of formula (I) provided by this invention have superior selective herbicidal activity. From the view-point of herbicidal activity, most preferred among the compounds of formula (I) are those of the following formula



25


30 wherein R_{02} is an allyl or propargyl group, and X_1 is a hydrogen or chlorine atom; and those of the following formula

30



35

35 wherein Y_2 is the group



40

and X_2 is a hydrogen or chlorine atom.

45 Typical examples of the compounds of formula (I) provided by the present invention except those specifically shown in Examples to be given hereinbelow are listed below.

45

50 N-allyl-N-(4-chlorobenzoyl)benzenesulfonamide,
N-allyl-N-(3,4-dichlorobenzoyl)benzenesulfonamide,
N-allyl-N-(2,5-dichlorobenzoyl)benzenesulfonamide,
N-allyl-N-(3,5-dichlorobenzoyl)benzenesulfonamide,
N-allyl-N-(pentachlorobenzoyl)benzenesulfonamide,
and
N-allyl-N-benzoylbenzenesulfonamide.

50


55 Specific examples of the most preferred compounds in the present invention are:
N-allyl-N-(2-chlorobenzoyl)benzenesulfonamide,
N-allyl-N-(2,4-dichlorobenzoyl)benzenesulfonamide,
N-propargyl-N-(2-chlorobenzoyl)benzenesulfonamide,
N-allyl-N-(α , α -dimethylbenzyl)benzenesulfonamide,
N-allyl-N-(α -methylbenzyl)benzenesulfonamide,
60 N-allyl-N-(α -methyl-4-chlorobenzyl)benzenesulfonamide, and
N-allyl-N-(α , α -dimethyl-4-chlorobenzyl)benzenesulfonamide.

55

60

According to the present invention, the compound of formula (I) can be produced by
 (a) reacting a benzenesulfonamide derivative of the general formula


5



5

wherein M represents a hydrogen atom or an alkali metal atom, and R₀, R₁, R₂ and R₃ are as defined hereinabove, with a compound of the general formula


10



10

15 wherein X₃ is a halogen atom, and R₄, R₅, R₆, R₇, R₈ and Y are as defined hereinabove, or
 (b) reacting a benzenesulfonyl halide of the general formula


20



20

wherein X₄ is a halogen atom, and R₁, R₂ and R₃ are as defined hereinabove, with a compound of the general formula


25



25

30 wherein R₀, R₄, R₅, R₆, R₇, R₈ and Y are as defined hereinabove, or
 (c) reacting a compound of the general formula

35



35

wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, Y and M are as defined hereinabove, with a compound of the formula

40



40

wherein X₄ is a halogen atom, and R₀ is as defined above.

The reaction of the benzenesulfonamide derivative of formula (II) with the compound of formula (III) in method (a) may be carried out in the absence of a solvent. Generally, however, it is preferably carried out in an inert solvent. Examples of usable solvents are water, ethers such as diethyl ether, dioxane or tetrahydrofuran, aromatic hydrocarbons such as benzene, toluene or xylene, N,N-dimethyl formamide, dimethyl sulfoxide, and pyridine. For certain types of the starting materials, alcohols such as methanol or ethanol, ketones such as acetone or methyl ethyl ketone, or halogenated hydrocarbons such as methylene chloride, chloroform or carbon tetrachloride may also be used as the solvents. These solvents are used singly or as mixtures of two or more.

The reaction temperature is not critical, and can be varied over a wide range according, for example, to the types of the starting materials, and the type of the solvent. Advantageously, the reaction temperature is generally from about 0°C. to the reflux temperature of the reaction mixture, preferably from room temperature to the reflux temperature of the reaction mixture. It is sufficient that the reaction pressure is atmospheric pressure, but if desired, reduced or elevated pressures may be employed. Under these conditions, the reaction can be terminated in about 0.5 to 5 hours.

The ratio between the compound of formula (II) and the compound of formula (III) to be reacted is not critical, and can be varied over a wide range. Advantageously, the compound of formula (III) is used in an amount of generally at least 1 mole, preferably 1.1 to 2 moles, per mole of the compound of formula (II).

If a compound of formula (II) in which M is a hydrogen atom is used as the starting material, the above reaction is advantageously carried out in the presence of an acid binder. Examples of usable acid binders are basic substances, for example alkali metal hydroxides such as sodium hydroxide or potassium hydroxide, and organic bases such as pyridine or

50

55

60

65

triethylamine.


Examples of preferred alkali metal atoms for M in formula (II) are lithium, potassium and sodium. A compound of formula (II) in which M is an alkali metal atom may be prepared in advance from a compound of formula (II) in which M is a hydrogen atom and an alkali metal hydride such as sodium hydride or potassium hydride or an alkali metal such as sodium or potassium, and then reacted with the compound of formula (III). Or it may be formed *in situ* by causing the alkali metal hydride or alkali metal to be present in the reaction system.

Thus, in the reaction between the compound of formula (II) and the compound of formula (III), the use of a reaction medium comprising a combination of sodium hydroxide or potassium hydroxide and water; a combination of pyridine and an ether or an aromatic hydrocarbon; or a combination of an alkali metal hydride and N,N-dimethyl formamide or dimethyl sulfoxide is advantageous.

The amount of each of the alkali metal hydroxide, organic base, alkali metal hydride, etc. may be at least 1 mole, preferably 1.1 to 3 moles, per mole of the compound of formula (II).

In the compound of formula (III), a chlorine atom is especially preferred as the halogen atom represented by X₃.

The method (a) can be especially advantageously applied to the production of compounds of formula (I) in which Y represents



The resulting reaction mixture containing the compound of formula (I) can be recovered by a known procedure. For example, water is added to the reaction mixture, and when the final product precipitates as a solid, it can be separated by filtration, centrifugal separation or the like. When the final product precipitates as an oil, it can be separated by solvent extraction, decantation or the like. In the case of solvent extraction, benzene, ethyl acetate, and chloroform may be suitably used as solvents. The crude final product separated can be purified, as needed, by recrystallization, distillation, chromatography, etc.

According to method (b) in accordance with this invention, the benzenesulfonyl halide of formula (IV) is reacted with the amine of formula (V). This reaction can be performed in the absence of a solvent, but usually, it is preferred to carry it out in an inert solvent. Examples of useful inert solvents are water, aromatic hydrocarbons such as benzene, toluene or xylene, ethers such as diethyl ether, tetrahydrofuran or dioxane, ketones such as acetone or methyl ethyl ketone, and halogenated hydrocarbons such as methylene chloride, chloroform or carbon tetrachloride. Water, benzene, and tetrahydrofuran are especially preferred.

The reaction temperature is not critical, and can be varied over a wide range according, for example, to the type of the starting materials and/or the type of the solvent used. It is advantageous that the reaction temperature is generally from about 0°C. to the reflux temperature of the reaction mixture, preferably from room temperature to about 80°C. The reaction pressure is usually atmospheric pressure, but if desired, elevated or reduced pressures may be employed.

If desired, the reaction in method (b) may be carried out in the presence of a catalyst. Useful catalysts are, for example, alkali metal hydroxides such as sodium hydroxide, and organic bases such as pyridine or triethylamine. The amount of the compound of formula (IV) is advantageously about 1 to 2 moles per mole of the compound of formula (IV).

The ratio between the compound of formula (IV) and the compound of formula (V) is neither critical and can be varied over a wide range. Generally, it is advantageous that the compound of formula (IV) is used in an amount of at least 1 mole, preferably 1.1 to 2 moles, per mole of the compound of formula (V).

Under the reaction conditions described hereinabove, the reaction is generally terminated in about 1 to 6 hours. The separation of the final product from the reaction mixture and its purification can be performed in the same manner as mentioned hereinabove with regard to method (a).

In method (c) of this invention, the compound of formula (VI) is treated with an alkylating, alkenylating or alkynylating agent [the compound of formula (VII)] to introduce the group R₉ into the N-atom of the compound of formula (VI).

Examples of the compound of formula (VII) include alkyl halides such as methyl bromide, ethyl chloride, methyl iodide, n-butyl iodide and n-decyl bromide, substituted alkyl halides such as cyanomethyl bromide, methoxypropyl iodide and dimethylaminopropyl bromide, alkenyl halides such as allyl chloride, and alkynyl halides such as propargyl chloride.

Preferably, the reaction between the compound of formula (VI) and the compound of formula (VII) is generally carried out in an inert solvent. Useful solvents include, for example, ethers such as diethyl ether, tetra-hydrofuran or dioxane, aromatic hydrocarbons such as benzene, toluene or xylene, alcohols such as methanol or ethanol, ketones such as acetone or methyl ethyl ketone, halogenated hydrocarbons such as methylene chloride or chloroform, N,N-dimethyl formamide, dimethyl sulfoxide, pyridine, and water. 5

Advantageously, the reaction of a compound of formula (VI) in which M is hydrogen with the compound of formula (VII) can be carried out in the presence of a basic catalyst, for example an alkali metal hydroxide such as sodium hydroxide or potassium hydroxide, an organic base such as sodium hydride or potassium hydride, or an alkali metal such as metallic sodium or potassium. The catalyst can be used in an amount of 1 to 2 moles per mole of the compound of formula (VI). A compound of formula (VI) in which M is an alkali metal atom can be prepared by the reaction of a compound of formula (VI) in which M is hydrogen with the alkali metal hydride or alkali metal. It may also be produced *in situ* in the reaction system. 10

The reaction temperature is not critical, and can be varied widely according, for example, to the types of the starting materials, or the type of the solvent. Generally, the reaction temperature is from about 0°C. to the reflux temperature of the reaction mixture, preferably from room temperature to the reflux temperature of the reaction mixture. 15

Usually, the reaction pressure is atmospheric pressure, but if desired, reduced or elevated pressures may be used. 20

The ratio between the compound of formula (VI) and the compound of formula (VII) is not critical, and can be varied widely. The compound of formula (VII) is used in an amount of at least 1 mole, usually 1.2 to 3 moles, per mole of the compound of formula (VI). 25

Under the reaction conditions described hereinabove, the reaction can be terminated in about 1 to 10 hours. The final product of formula (I) can be obtained in good yields from the resulting reaction mixture in the same manner as described hereinabove. 30

Some of the compound of formula (VI) used in method (c) in this invention, especially those in which Y is the group 35

$$\begin{array}{c}
 R_9 \\
 | \\
 -C- \\
 | \\
 R_{10}
 \end{array}$$

are novel compounds, and can be prepared, for example, by reacting correspondingly substituted benzylamines with the corresponding benzenesulfonyl halides in the presence of catalysts such as sodium hydroxide. 40

The production of the compound (I) of this invention is illustrated by the following Examples. 45

EXAMPLE 1

N-Methyl-N-(α , α -dimethylbenzyl)benzenesulfonamide

27.5 g (0.1 mole) of N-(α , α -dimethylbenzyl)benzenesulfonamide was added to 5.8 g (0.12 mole) of 50% sodium hydride in 200 ml of dry N,N-dimethyl formamide. The mixture was stirred at room temperature for 30 minutes to react them. To the resulting sodium salt of N-(α , α -dimethylbenzyl)benzenesulfonamide was added 16.1 g (0.17 mole) of methyl bromide. The mixture was heated to 60°C. and stirred for 1 hour. The N,N-dimethyl formamide was distilled off under reduced pressure, and cold water was added. The resulting white solid was recrystallized from methanol to afford N-methyl-N-(α , α -dimethylbenzyl)benzenesulfonamide in a yield of 87%. This compound is designated as compound No. 64 in Table 2 below, and its melting point and elemental analysis values are shown in Table 2. 50


The N-(α , α -dimethylbenzyl)benzenesulfonamide used as a starting compound in the above procedure could be prepared in the following manner. A two-layered mixture consisting of 33.8 g (0.25 mole) of α , α -dimethylbenzylamine and 120 ml of a 10% aqueous solution of sodium hydroxide was stirred at below 40°C., and 44.1 g of benzenesulfonyl chloride was added dropwise. Stirring the mixture for an additional 1.5 hours afforded a white precipitate. The solid precipitate was collected by filtration, and recrystallized from methanol to afford N-(α , α -dimethylbenzyl)benzenesulfonamide having melting point of 114 to 115°C. in a yield of 67%. 55

Starting compounds shown in Table 1 were prepared by the same procedure as above except that α -methyl-benzylamine or the correspondingly substituted α , α -dimethylbenzylamines were used instead of the α , α -dimethylbenzylamine, and the 60

65

correspondingly substituted benzenesulfonyl chlorides were used instead of the benzenesulfonyl chloride.

TABLE 1



10	Compound No.	Q ₁	Q ₂	Q ₃	Q ₄	Melting point or refractive index	10
15	2	2-CH ₃	H	CH ₃	H	120 - 121°C.	15
	3	4-CH ₃	H	CH ₃	H	135 - 137°C.	
	4	4-CH ₃ O	H	CH ₃	H	128 - 130°C.	
20	5	4-Cl	H	CH ₃	H	125 - 126°C.	20
	6	2-CH ₃	5-CH ₃	CH ₃	H	112.0-112.5°C.	
25	7	4-C ₂ H ₅	H	CH ₃	H	103.5-104.0°C	25
	8	H	H	CH ₃	4-Cl	105 - 107°C.	
	9	H	H	CH ₃	CH ₃	72 - 78°C.	
30	10	H	H	H	H	102 - 103°C.	30
	11	4-CH ₃	H	H	H	82 - 83.5°C.	
35	12	4-C ₂ H ₅	H	H	H	$n_D^{26} = 1.5656$	35
	13	4-Cl	H	H	H	74 - 75°C.	
	14	2-CH ₃	4-CH ₃	H	H	104.5-105°C.	
40	15	2-CH ₃	5-CH ₃	H	H	72.5-73°C.	40

Compounds Nos. 65 to 88 and 95 to 102 shown in Table 2 could be produced in the same way as in Example 1 using the starting compounds Nos. 2 to 15.

45 **EXAMPLE 2**
N-Allyl-N-(α -methyl-2-chlorobenzyl)benzenesulfonamide
1.97 g (0.01 mole) of N-allylbenzenesulfonamide was added to 0.58 g (0.012 mole) of 50% sodium hydride in 15 ml of dry N,N-dimethylformamide. The mixture was stirred at room temperature for 30 minutes to react them. To the resulting sodium salt was added 2.10 g (0.012 mole) of α -methyl-2-chlorobenzyl chloride. The mixture was stirred at room temperature for 1 hour. The N,N-dimethyl formamide was distilled off under reduced pressure, and cold water was added. The resulting oil layer was extracted with benzene. The benzene layer was dried over anhydrous magnesium sulfate, concentrated, and purified by silica gel chromatography to afford the final product. The final product is designated as compound No. 89 in Table 2, and its melting point and elemental analysis values are shown in Table 2.

Compounds Nos. 90 to 94 shown in Table 2 could be prepared in the same way as in Example 2.

60 **EXAMPLE 3**
N-methyl-N-(2-chlorobenzoyl)benzenesulfonamide
1.71 g (0.01 mole) of N-methylbenzenesulfonamide was added to 0.58 g (0.012 mole) of 50% sodium hydride in 15 ml of dry N,N-dimethylformamide. The mixture was stirred at room temperature for 30 minutes to react them. To the resulting sodium salt was added 2.10 g (0.012 mole) of 2-chlorobenzoyl chloride, and the mixture was stirred at room

temperature for 1 hour. The N,N-dimethyl formamide was distilled off under reduced pressure. Cold water was added, and the resulting oil layer was extracted with benzene. The benzene layer was dried over anhydrous magnesium sulfate, and concentrated to afford almost pure N-methyl-N-(2-chlorobenzoyl)benzenesulfonamide in a yield of 92%. For further purification, this compound was subjected to chromatography or high vacuum distillation. This compound is designated as compound No. 16 in Table 2, and its melting point and elemental analysis values are shown in Table 2.

Compounds Nos. 17 to 63 in Table 2 could be prepared in the same way as in Example 3.

10 EXAMPLE 4

N-Allyl-N-(2-chlorobenzoyl)benzenesulfonamide

3.0 g (0.01 mole) of N-2-chlorobenzoyl benzenesulfonamide prepared from benzenesulfonamide and 2-chlorobenzoyl chloride was added to 0.58 g (0.012 mole) of 50% sodium hydride in 15 ml of dry tetrahydrofuran. After stirring for 30 minutes at room temperature, 1.48 g (0.012 mole) of allyl bromide was added. Then the mixture was reacted for 2 hours under the same conditions. The tetrahydrofuran was distilled off under reduced pressure, and cold water was added. The resulting oil layer was extracted with benzene. The benzene layer was separated, dried over anhydrous magnesium sulfate, and concentrated to afford a white solid. Recrystallization from methanol afforded compound No. 27 in Table 2 in a yield of 35%.

Compounds Nos. 16 to 26 and 28 to 63 could be produced in the same way as in Example 4.

25 EXAMPLE 5

N-Methyl-N-(α -methylbenzyl)benzenesulfonamide

Benzenesulfonyl chloride (2.1 g; 0.012 mole) was added dropwise at below 40°C. to a two-layered mixture consisting of 1.35 g (0.01 mole) of N-methyl- α -phenethylamine [prepared by the method disclosed in Novzlli, J. Ame. Chem. Soc. 61, 520 (1939)] and 12 ml of a 10% aqueous solution of sodium hydroxide. On continuing the reaction for 2 hours, an oily substance formed in the lower layer. It was extracted with benzene, and dried over anhydrous magnesium sulfate. The benzene was distilled off under reduced pressure to give a colorless oily substance. Purification by silica gel column chromatography afforded pure N-methyl-N-(α -methylbenzyl)benzenesulfonamide in yield of 98%. This compound is designated as compound No. 82 in Table 2, and its refractive index and elemental analysis values are also shown in Table 2.

Compounds Nos. 64 to 72, 74 to 77, 79 to 81, 83 to 88, and 95 to 102 shown in Table 2 could be produced in the same way as in Example 5.

40 EXAMPLE 6

N-Allyl-N-(2,4-dichlorobenzoyl)-4-toluenesulfonamide

4.6 g (0.02 mole) of N-allyl-2,4-dichlorobenzamide was added to 1.2 g (0.024 mole) of 50% sodium hydride in 20 ml of dry tetrahydrofuran. The mixture was stirred at room temperature for 30 minutes to react them. Then, 4.2 g (0.024 mole) of p-toluenesulfonyl chloride was added, and reacted at room temperature for 1 hour. The tetrahydrofuran was distilled off under reduced pressure. After adding cold water to the residual oil, it was extracted with benzene, dried over anhydrous magnesium sulfate, and concentrated to afford an oily substance. It was purified by silica gel chromatography to afford compound No. 42 shown in Table 2 in a yield of 68%. Its refractive index and elemental analysis values are also shown in Table 2.

50 Compounds Nos. 16 to 41 and 43 to 63 in Table 2 could be prepared in the same way as in Example 6.

TABLE 2

Com- ound No.	Structural formula	Elemental analysis values (%) (the upper row: calculated the lower row: found)				X
		C	H	N	S	
16		n_D^{21} 1.5878	54.48	3.90	4.52	10.35 $X=C\ell$ 11.45
17		82-83.5°C	54.34	3.77	4.50	10.53 $X=C\ell$ 11.37
18		n_D^{25} 1.5912	48.85	3.22	4.07	9.32 $X=C\ell$ 20.60
19		53-54°C	48.96	3.41	4.05	9.44 $X=Br$ 22.56
20		48-50°C.	62.35	5.36	4.80	11.19 -

TABLE 2 (continued)

Compound No.	Structural formula	n_D^{25}	Elemental analysis values (%)				
			the upper row: calculated		the lower row: found		X
			C	H	N	S	
21		1.5791	48.14	3.50	3.74	8.57	X=Cf 18.95
22		1.5750	48.03	3.38	3.75	8.66	18.90
23		1.5768	55.64	4.36	4.33	9.90	X=Cf 10.95
24		1.5650	55.51	4.40	4.35	9.78	10.87
25		99-100°C.	51.62	4.06	3.76	8.61	X=Cf 19.05
			51.53	4.18	3.74	8.53	18.95
			56.88	4.77	4.15	9.49	10.50
			56.92	4.73	4.08	9.31	10.47
			56.88	4.77	4.15	9.49	X=Cf 10.50
			56.75	4.70	4.18	9.53	10.43

TABLE 2 (continued)

Compound No.	Structural formula	Elemental analysis values (%)						X	
		Melting point or refractive index		C		H			
		n_D^{25}	n_D^{25}	49.24	3.62	3.59	8.22		
26		1.5598	49.38	3.76	3.52	8.30	$X=C\ell$		
27		65.5-66.0°C.	57.22	4.20	4.17	9.55	$X=C\ell$		
28		78.5-79.0°C.	56.98	4.31	4.13	9.32	$X=C\ell$		
29		39.0-40.0°C.	51.90	3.54	3.78	8.66	$X=C\ell$		
30		51.5-52.5°C.	59.98	4.37	4.43	9.90	$X=F$		
			60.17	4.42	4.39	10.04	5.95	$X=I$	
			44.97	3.30	3.28	7.50	29.71		
			45.31	3.17	3.35	7.63	29.65		

TABLE 2 (continued)

Compound No.	Structural formula	Elemental analysis values (%)					
		Melting point or refractive index		C H		N S X	
31		49.0-49.5°C.	57.22	4.20	4.17	9.55	X=Cf 10.56
32		111.0-112.0°C.	56.96	4.27	4.21	9.42	X=Cf 10.39
33		96.0-98.0°C.	51.90	3.54	3.78	8.66	X=Cf 19.15
34		54.5-55.0°C.	52.04	3.65	3.83	8.54	X=Cf 19.03
35		57.0-57.5°C.	47.48	2.99	3.46	7.92	X=Cf 26.28
			47.57	3.08	3.52	7.74	26.15
			64.74	5.43	4.44	10.17	-
			65.20	5.31	4.52	10.29	-
			64.74	5.43	4.44	10.17	-
			64.80	5.33	4.49	10.26	-

TABLE 2 (continued)

Compound No.	Structural formula	Elemental analysis values (%)						X
		Melting point or refractive index		C		H		
the upper row: calculated		the lower row: found		C	H	N	S	
36		86.0-87.0°C.	61.61	5.17	4.23	9.68	-	
37		56.0-56.5°C.	61.45	5.24	4.35	9.52	-	
38		n_D^{26} 1.5775	61.61	5.17	4.23	9.68	-	
39		n_D^{26} 1.5700	61.50	5.23	4.15	9.75	-	
40		n_D^{26} 1.5735	51.01	3.78	3.50	8.01	17.72	X=Cℓ

TABLE 2 (continued)

Compound No.	Structural formula	Melting point or refractive index	Elemental analysis values (%)				X
			C	H	N	S	
41		n_D^{26} 1.5805	53.13	3.93	3.65	8.34	X=Cℓ 18.45
42		n_D^{25} 1.5649	53.01	3.85	3.58	8.55	18.53
43		n_D^{28} 1.5796	53.13	3.93	3.65	8.34	X=Cℓ 18.45
44		n_D^{26} 1.5953	53.20	3.85	3.50	8.46	18.50
45		81.0-82.0°C.	54.28	4.30	3.52	8.05	X=Cℓ 17.80
			54.34	4.41	3.45	8.12	17.72
			47.48	2.99	3.46	7.92	X=Cℓ 26.28
			47.45	3.05	3.58	7.80	26.35
			51.90	3.54	3.78	8.66	X=Cℓ 19.15
			51.75	3.65	3.71	8.68	19.30

TABLE 2 (continued)

Compound No.	Structural formula	Elemental analysis values (%)						
		Melting point or refractive index		C		H	N	S
(the upper row: calculated)		(the lower row: found)						
46		69.5-70.0°C.	47.48	2.99	3.46	7.92	26.28	X=Cℓ
47		n_D^{18}	47.62	3.03	3.30	7.85	26.36	X=Cℓ
48		57.0-59.5°C.	54.31	3.70	3.96	9.06	10.02	X=Cℓ
49		43.0-45.0°C.	54.34	3.63	4.05	9.22	9.91	X=Cℓ
50		81.0-82.0°C.	55.81	4.41	3.83	8.77	9.69	X=Cℓ
			55.68	4.60	3.75	8.90	9.65	X=Cℓ
			44.20	3.55	3.22	7.38	24.47	
			44.14	3.29	3.28	7.53	24.61	

TABLE 2 (continued)




Compound No.	Structural formula	Melting point or refractive index	Elemental analysis values (%)				X=Cf 10.08
			the upper row: calculated		the lower row: found		
		C	H	N	S		
51		n_D^{24} 1.5575	58.03 58.18	5.16 5.34	3.98 3.95	9.11 9.04	10.13
52		n_D^{24} 1.5550	58.03 58.18	5.16 5.25	3.98 3.98	9.11 9.15	10.08
53		50.0-52.0°C. 57.94	5.16 5.07	3.98 3.84	9.11 9.21	10.08 10.12	X=Cf 10.08
54		n_D^{24} 1.5580	59.09 5.51	5.51 3.83	3.83 8.76	9.69 9.48	X=Cf 9.69
55		62.5-63.5°C. 61.01	6.14 6.16	3.56 3.51	8.14 8.18	8.79 8.79	X=Cf 9.00

TABLE 2 (continued)


Compound No.	Structural formula	Elemental analysis values (%) (the upper row: calculated the lower row: found)					
		C	H	N	S	X	X=Cℓ
56		62.0-63.5°C.	61.82	6.42	3.43	7.86	8.69
57		88.0-89.0°C.	61.78	6.50	3.40	7.91	8.67
58		42.0-43.5°C.	62.62	6.69	3.32	7.60	8.40
59		n_D^{26} 1.5327	63.45	6.89	3.30	7.38	8.15
60		n_D^{24} 1.5584	56.88	5.62	7.30	8.48	9.28

TABLE 2 (continued)

Compound No.	Structural formula	Melting point or refractive index	Elemental analysis values (%)				X	
			(the upper row: calculated the lower row: found)		C	H	N	
			X	X=Cf				
61		n_D^{24} 1.5760	53.81	3.31	8.37	9.58	X=Cf 10.57	
62		n_D^{21} 1.5650	53.73	3.22	8.40	9.53	10.70	
63		n_D^{25} 1.5778	55.50	4.93	3.81	8.72	X=Cf 9.64	
64		93.0°C.	57.57	3.62	4.20	9.61	X=Cf 10.62	
65		65.0-66.0°C.	66.40	6.62	4.84	11.08	-	
			66.25	6.58	4.75	11.00	-	
			67.29	6.98	4.62	10.57	-	
			67.35	7.02	4.58	10.52	-	

TABLE 2 (continued)

Compound No.	Structural formula	Melting point or refractive index	Elemental analysis values (%) (the upper row: calculated) (the lower row: found)					X
			C	H	N	S		
66		79.0-80.0°C.	68.10	7.30	4.41	10.10	-	
67		74.5-75.0°C.	68.05	7.21	4.39	10.15	-	
68		n_D^{21} 1.5750	68.98	6.11	4.47	10.23	-	
69		47.0-48.0°C.	68.85	6.08	4.49	10.20	-	
70		68.0°C.	69.29	7.04	4.25	9.73	-	
			69.09	7.08	4.13	9.82	-	
			68.84	7.60	4.23	9.67	-	
			68.75	7.64	4.18	9.75	-	

TABLE 2 (continued)

Compound No.	Structural formula	Elemental analysis values (%)					
		Melting point or refractive index		(the upper row: calculated the lower row: found)			
		C	H	N	S	X	
71		n_D^{21} 1.5450	70.23 70.05	8.14 8.17	3.90 3.85	8.92 8.99	-
72		n_D^{21} 1.5190	73.09 73.15	9.32 9.35	3.16 3.10	7.23 7.18	-
73		n_D^{21} 1.5530	63.92 63.86	6.63 6.48	4.39 4.50	10.04 10.12	-
74		63.0-66.0°C.		69.27 69.20	7.04 7.08	4.25 4.21	9.73 9.78
75		n_D^{27} 1.5676	69.13 69.13	7.08 7.08	4.21 4.21	9.76 9.76	-

TABLE 2 (continued)

Compound No.	Structural formula	Elemental analysis values (%)					
		the upper row: calculated			the lower row: found		
		C	H	N	S	X	
76		n_D^{27} 1.5710	66.06	6.71	4.06	9.28	-
77		n_D^{27} 1.5770	66.15	6.67	4.10	9.22	-
78		n_D^{26} 1.5672	61.79	5.76	4.00	9.17	$X=C\ell$ 10.13
79		n_D^{26} 1.5672	61.71	5.79	4.06	9.12	10.09
			61.79	5.76	4.00	9.17	$X=C\ell$ 10.13
			61.68	5.71	3.95	9.23	10.10
			69.26	7.04	4.25	9.73	-
			69.17	7.10	4.20	9.78	-
			61.0-63.0°C.				
			(mixture of O-, m-, p-isomers)				

TABLE 2 (continued)

Compound No.	Structural formula	Melting point or refractive index	Elemental analysis values (%)				X
			C	H	N	S	
80		n_D^{26} 1.5653	69.93	7.34	4.08	9.34	-
81		n_D^{26} 1.5630	69.93	7.34	4.08	9.34	-
82		78.0-79.0°C.	69.92	7.40	4.13	9.29	-
83		62.0-62.5°C.	65.42	6.22	5.09	11.65	-
84		50.0-52.0°C.	65.34	6.19	5.15	11.71	-
			66.40	6.62	4.84	11.08	-
			64.48	6.56	4.79	11.16	-
			67.29	6.98	4.62	10.57	-
			67.21	6.92	4.70	10.65	-

TABLE 2 (continued)

Compound No.	Structural formula	Melting point or refractive index	Elemental analysis values (%)					X	
			(the upper row: calculated)		(the lower row: found)				
			C	H	N	S			
85		n_D^{27} 1.5598	67.74	6.35	4.65	10.64	-		
86		n_D^{27} 1.5507	68.10	7.30	4.41	10.10	-		
87		47.0-48.5°C.	68.25	7.25	4.37	10.26	-		
88		n_D^{27} 1.5066	72.68	9.15	3.26	7.46	-		
89		65.0-67.5°C.	72.52	9.18	3.20	7.52	-	$X=C\ell$ 10.56	
			60.79	5.40	4.17	9.55			
			60.65	5.35	4.23	9.61		10.63	

TABLE 2 (continued)

Compound No.	Structural formula	Melting point or refractive index	Elemental analysis values (%)				X
			(the upper row: calculated)		(the lower row: found)		
			C	H	N	S	
90		n_D^{26} 1.5705	60.79	5.40	4.17	9.55	X=Cf 10.56
91		n_D^{26} 1.5720	60.79	5.40	4.17	9.55	X=Cf 10.56
92		n_D^{26} 1.5821	55.14	4.63	3.78	8.66	X=Cf 19.15
93		n_D^{27} 1.5598	55.05	4.69	3.75	8.71	19.03
94		n_D^{26} 1.5670	65.12	6.35	4.31	9.79	-

TABLE 2 (continued)

Compound No.	Structural formula	Melting point or refractive index	Elemental analysis values (%)				
			the upper row: calculated		the lower row: found		
		C	H	N	S	X	
95		n_D^{26} 1.5666	69.27	7.04	4.25	9.73	-
96		n_D^{26} 1.5671	69.25	7.01	4.28	9.69	-
97		n_D^{27} 1.5647	69.30	7.02	4.15	9.75	-
98		n_D^{26} 1.5630	68.54	6.71	4.44	10.17	-
99		n_D^{27} 1.5700	65.15	6.46	4.19	9.76	-

TABLE 2 (continued)

Compound No.	Structural formula	Elemental analysis values (%)				X
		Melting point or refractive index		C	H	
		(the upper row: calculated the lower row: found)				
						X=Cℓ 10.56
100		n_D^{27} 1.5748	60.79 60.65	5.40 5.32	4.17 4.25	9.55 9.63
101		n_D^{27} 1.5891	68.20 68.12	5.72 5.75	4.68 4.61	10.71 10.65
102		n_D^{27} 1.5621	68.54 68.39	6.71 6.75	4.44 4.40	10.17 10.29

The compounds of formula (I) provided by this invention have superior herbicidal activities, and are useful as active ingredients of herbicides for controlling various weeds in agricultural crops. Examples of weeds which can be controlled by the compounds of formula (I) of this invention are various species of barnyard grass (such as *Echinochloa crus-galli* Beauv., *Echinochloa crusgalli* var. *oryzicola* Ohwi, and *Echinochloa crus-galli* Subsp. *genuine* var. *echinata* Honda), spikerush (*Eleocharis pellucida* Presl), sedge sp. (*Cyperus hakoneensis* saiat), umbrella plant (*Cyperus difformis* L.), pipewort (*Eriocaulon sieboldianum* Sieb.), waterwort (*Elatine triandra*), redstem sp. (*Rotala indica* Koehne), bulrush (*Scirpus juncoides* Roxb.), redstem sp. (*Ammannia multiflora* Roxb.), false pimpernel (*Lindernia pyxidaria* L.), and slender spikerush (*Eleocharis acicularis* Roem. et Schult var. *longiseta* Svenson). These examples are not limitative, and it should be understood that the compounds of formula (I) of this invention exhibit herbicidal effects also against other kinds of weeds.

It has been found that the compounds of formula (I) of this invention exhibit marked effects in controlling weeds which occur in fields containing much water, such as a paddy field, rather than those which occur in dry upland fields.

Thus, the compounds of formula (I) exhibit excellent control effects against various species of barnyard grass, especially *Echinochloa crus-galli* Beauv. which is a very hazardous weed in an aquatic paddy and is considered as one of the five greatest weeds in the world. This weed grows in paddy fields, especially submerged paddy fields, throughout the world. The compounds of formula (I) have the ability to inhibit the germination of the barnyard grass strongly and to prevent its growth in paddy fields.

Moreover, the compounds of formula (I) are very characteristic in that they have excellent selective herbicidal activity which ensures substantial freedom from phytotoxicity to useful agricultural crops such as rice.

Many herbicides have heretofore been suggested for application to paddy fields, and some have come into actual use. Almost none of them, however, have selectivity in physiological herbicidal action between barnyard grass and rice plant. The conventional methods for weed killing in paddy fields are directed to the treatment of paddy fields in the rice growing stage (including the transplanting stage) to control the sprouting of barnyard grass. They are based either on the utilization of the differences in resistance to herbicides between barnyard grass and rice plant according to the differences in their growing stages, or on the principle of chemical adsorption in the upper layer of soil ("artificial selectivity") whereby rice plants are transplanted in such a manner that their roots are located below the herbicide-treated layer, and barnyard grass in the upper layer is controlled while protecting the rice plants from the herbicide.

Barnyard grass is a graminaceous weed as rice, and they physiologically resemble each other very well. Hence, controlling of barnyard grass with herbicidal chemicals often causes phytotoxicity to rice plant, and it is extremely difficult to control this weed selectively in paddy fields. Barnyard grass has therefore been considered to be difficult to eradicate in paddy fields, and there has been a strong demand for the advent of herbicides which can selectively control barnyard grass.

The compounds of formula (I) of this invention meet this demand of agriculture. They have the excellent property of acting selectively on the seeds and seedlings of barnyard grass to strongly inhibit their germination, but causing no substantial phytotoxicity to rice plant. This property renders the compounds of formula (I) very suitable as active ingredients of herbicides for application to paddy fields.

The superior herbicidal activity of the compounds (I) of this invention can be demonstrated by the experimental fact that when N-allyl-N-(2-chlorobenzoyl)benzenesulfonamide was applied at a rate of 62.5 g per 10 ares to a paddy field where rice plant and barnyard grass were simultaneously sown, the germination of the barnyard grass was completely inhibited, whereas the rice plant showed normal emergence and growth without any phytotoxicity, and that even when the rate of the compound applied was increased to 1,000 g per 10 ares, the rice plant showed normal germination and growth without any phytotoxicity. Thus, the compound of this invention, when applied in an amount about 20 times as large as the amount required for completely controlling barnyard grass, does not exert any substantial effect on the germination and growth of rice plant.

Such a high selectivity of the compounds of this invention between barnyard grass and rice plant is ascribable presumably to the specific physiological activities of the compounds of this invention against barnyard grass and rice plant. This superior selectivity cannot be expected from the conventional herbicides available for application to paddy fields.

The compound of formula (I) of this invention may be applied directly as a herbicide. Generally, however, it is formulated into a herbicidal composition by mixing it with inert liquid or solid carriers or diluents which are commonly employed in herbicide formulations.

In the present invention, any inert liquid or solid carriers or diluents known in the art can be used. Examples of the inert solid carrier or diluent are kaolin, diatomaceous earth, talc, bentonite, silica, and clay minerals. Examples of the inert liquid carrier or diluent are water, xylene, toluene, benzene, N,N-dimethyl formamide, dimethyl sulfoxide, and liquefied gases such as tetrafluoroethane.

In addition to the inert liquid or solid carrier or diluent, the herbicidal composition may, as needed, contain surface-active agents such as polyoxyethylene monolaurate or polyethylene sorbitol in usual amounts chosen according to the form of the herbicidal composition.

The herbicidal composition may contain the active compound of formula (I) in an amount of at least 0.5% by weight, preferably 1 to 90% by weight, more preferably 2 to 80% by weight, based on the weight of the composition itself.

The herbicidal composition can be in any conventional forms such as a dust, granule, wettable powder, solution, emulsifiable concentrate, or spray according to the method of

application. Any methods of formulation known in the art can be used for this purpose. For example, when making a dust, granule or wettable powder, at least one active compound of formula (I) is mixed with at least one inert solid carrier or diluent. The mixture is pulverized and mixed uniformly with a suitable amount of a surface active agent. The solution or emulsifiable concentrate can be prepared by dissolving or dispersing at least one active compound of formula (I) in at least one inert liquid carrier or diluent, followed, if desired, by adding a surface active agent.

Conveniently, the amount of the active compound of formula (I) is 3 to 20% by weight for the dust and granule, 25 to 75% by weight for the wettable powder, and 20 to 50% by weight for the solution and emulsifiable concentrate, all based on the weight of the resulting composition.

The herbicidal composition may further contain agricultural chemicals commonly used in cultivating agricultural crops, such as fungicides, insecticides, nematocides, and fertilizers. Typical examples of the fungicides are Benomyl [methyl 1-(n-butyloxycarbonyl)-1H-benzimidazol-2-yl-carbamate], Hymexazol (5-methyl-3-isoxazolol), Captan [3a,4,7,7a-tetrahydro-N-(trichloromethanesulphenyl) phthalimide], and Zineb [zinc ethylenebis-(dithiocarbamate)]. Examples of the insecticides are Disulfoton (0,0-diethylS-2-ethylthioethyl phosphorodithioate) and Propoxur (2-isopropoxyphenyl methylcarbamate). Examples of the nematocides are Methomyl[S-methyl N-(methylcarbamoyloxy) thioacetamide] and Alidicarb [2-methyl-2-(methylthio) propionaldehyde 0-methylcarbamoyloxime].

It is also possible to incorporate at least one other herbicidally active compound used heretofore in the art into the herbicidal composition of this invention. This frequently brings about a high herbicidal effect against a broad spectrum of weeds. Examples of the other herbicidally active compounds include MCP (2-methyl-4-chlorophenoxy acetic acid), TOK (2,4-dichlorophenyl-4'-nitrophenyl ether), Benthiocarb (4-chlorobenzyl-N,N-diethylthiocarbamate), Molinate (S-ethyl-N,N-hexamethylene thiocarbamate), Oxadiaxon[2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl-5-oxo-1,3,4-oxadiazone], and Butachlor [2-chloro-2',6'-diethyl-N-(butoxymethyl)acetanilide]. It should be understood that these examples are not limitative, and other active compounds can be equally incorporated in the herbicidal composition of this invention as needed.

The herbicidal compositions of this invention containing these other herbicidally active compounds are especially useful for application to paddy fields in the rice growing stage, for example to a paddy field in which transplantation has ended.

The herbicide containing the compound of formula (I) as an active ingredient can be used to control various weeds in areas where agricultural crops are cultivated. In particular, the herbicide of this invention is effective against weeds in wet paddies rather than dry fields, and exhibits a very strong selective herbicidal effect against barnyard grass which accompanies rice plant in paddy fields, such as Echinochloa crus-galli Beauv.

Herbicides containing as active ingredients 2,4,6-trichlorophenyl-4-nitrophenyl ether (CNP), S-(4-chlorobenzyl)-N,N-diethylthiocarbamate (Benthiocarb), 2-chloro-2',6'-diethyl-N-(butoxymethyl)acetanilide (Butachlor), and S-ethyl-N,N-hexamethyl enethiolcarbamate (Molinate), which now gain widespread acceptance for application to paddy fields, do not show selectivity between barnyard grass and rice plant in the germinating stage, nor are they absolutely safe to transplanted rice plants in the early stage of growth. With these conventional herbicides, the risk of phytotoxicity cannot be avoided in the event of changes in the environmental condition of paddy fields, for example when rice plants are transplanted shallow at the soil surface, the soil is sandy, the water leaks, root growth is abnormal, or the temperature becomes unusually high.

Since the herbicide provided by this invention is based on physiological selectivity, it has the advantage of being applicable to all growing stages of rice plants ranging from the germinating to the growing stage, and being substantially free from phytotoxicity to rice plants by changes in environmental conditions. Thus, its contributes greatly to the cultivation of agricultural crops.

Heretofore, 3',4'-dichloropropionanilide (Propanil) has been used worldwide as an agent having selective activity against barnyard grass in a paddy field. Propanil, however, is an agent suited for foliar application and has no effect of inhibiting germination. In contrast, the herbicide of this invention exhibits far higher selective activity during the emergency of barnyard grass and rice plant than Propanil, and it is no exaggeration to say that the herbicide of this invention is an epoch-making weed killer having no equal among known herbicides of this kind.

As a result of fundamental and applied tests on the herbicides of this invention, it has been found that most effective, and most preferred, herbicides of this invention are those containing N-allyl-N-(2-chlorobenzoyl) benzenesulfonamide, N-allyl-N-(2,4-dichlorobenzoyl)benzenesulfonamide, N-propargyl-N-(2-chlorobenzoyl)benzenesulfonamide, N-

allyl-N-(α , α -dimethylbenzyl)benzenesulfonamide, and N-allyl-N-(α -methylbenzyl)benzene sulfonamide as active ingredients. These active compounds commonly have the property of strongly inhibiting the germination of barnyard grass without affecting rice plants in emergence. There is, however, a slight difference in activity among these compounds. For example, herbicides containing benzoyl-type active compounds have long persistence in soil and can inhibit the growth of weeds for long periods of time. Herbicides containing benzyl-type active compounds, on the other hand, have a very wide range of selectivity between barnyard grass and rice plant immediately after emergence. 5

In use, the herbicide of this invention containing the active ingredient of formula (I) is applied to the locus to be protected from weeds. 10

Thus, according to still another aspect of this invention, there is provided a method for controlling weeds in agricultural crops which comprises applying the compound of general formula (I) described hereinbefore to the locus to be protected from the weeds. 15

The time of application of the compound of formula (I) is not strictly limited, and differs according to the agricultural crops and/or the weeds to be controlled. Generally, in order for the active compound of this invention to exhibit the best herbicidal effect, it is most convenient to apply it just before the weeds to be controlled sprout, or during their germinating stage. It is of course possible to apply it to weeds after emergence, and this brings about some extent of control effect. 20

There is no particular restriction on the locus to which the active compound of this invention can be applied. It can be applied to various types of agricultural land as is the case with conventional herbicides. It can be best applied however to wet paddies, especially aquatic paddies in the submerged state, and when applied to upland farms of low water content, the active compound of this invention tends to have a somewhat decreased herbicidal effect. 25

In order for the herbicide of this invention to exhibit its herbicidal effect most, it is applied to a field in a submerged condition before or during the germination of weeds. 30

The active compound of formula (I) of this invention exhibits herbicidal effects against the various weeds described hereinabove, but have excellent effects of inhibiting germination of various kinds of barnyard grass, especially *Echinochloa crus-galli* Beauv. which accompany rice plants, without any substantial toxicity to rice plant. Thus, the active compounds of formula (I) can be effectively applied to control barnyard grass selectively and protect rice plants therefrom. 35

The rate of application of the active compound of formula (I) is not critical, and can be varied widely according to the type of the active compound, the time of application, the procedure of application, etc. It is advantageous that the rate of application of the active compound of formula (I) is generally at least 25 g, preferably 50 to 1000 g, more preferably 100 to 500 g, per 10 ares. 40

The method of application may be any conventional method. For example, the herbicidal composition of this invention may be sprayed onto the locus to be protected from weeds from above the ground or from the air. Or it may be sprayed together with the seeds of an agricultural crop at the time of seeding the crop. 45

Furthermore, according to the present invention, seeds of a crop may be dipped prior to sowing in an aqueous liquid containing the active compound of this invention to control the germination of weed seeds that may be present mixed. 50

The active compounds of formula (I) of this invention have little toxicity on useful agricultural crops and low mammalian toxicity, and therefore are very suitable as herbicides. 55

The following Examples further illustrate the formulation of the herbicides provided by the present invention, and their selective herbicidal activities. 60

In these Examples, all parts and percentages are by weight. The numbers of the compounds refer to those given in Table 2. 65

Example A (wettable powder)

40 Parts of compound No. 27, 55 parts of a 2:1 mixture of Zeeklite and Kunilite (a trade-mark of a product of Kunimine Kabushiki Kaisha), and Sorpol 800 as a surfactant (a trademark for a product of Toho Kagaku Kogyo K.K.) were mixed and pulverized to form a 40% wettable powder. 55

Example B (emulsifiable concentrate)

25 Parts of compound No. 85, 65 parts of benzene and 10 parts of Sorpol 800 as a surfactant were mixed and dissolved to form a 25% emulsifiable concentrate. 60

Example C (granule)

Five parts of compound No. 67, 50 parts of bentonite, 40 parts of Kunilite and 5 parts of

Sorpol 800 as a surfactant were mixed, and pulverized. Then, 10 parts of water was added, and the mixture was uniformly stirred to form a paste. The paste was extruded through a hole with a diameter of 0.7 mm, dried, and then cut to a length of 1 to 2 mm to form a 5% granule.

5

5

Example D

Pots (1/5000 are) were filled with a paddy soil in the muddy state, and barnyard grass (*Echinochloa crus-galli* Beauv.) and other broad-leaves weeds (waterwort, pipewort, unblella plant, spikerush, sedge sp., red stem, and false pimpernel), bulurusk, and slender spikerush were sown or planted. At the same time, 10 germinated rice seeds (variety: Nihonbare) were sown, and two rice seedlings in the 3-lead stage were planted as one stock.

Two days later, a wettable power containing each of the active compounds of this invention shown in Table 3 was weighed to a predetermined amount, diluted with 10 ml of water, and applied to the irrigated water surface in each pot. Then, the treated pots were allowed to stand in a greenhouse, and the control effect and phytotoxicity on rice plants were examined two weeks later.

The evaluation was made on a scale of 10 grades where 0 represents normal growth and 10 represents complete withering.

The results are shown in Table 3.

10

10

15

15

TABLE 3

Amount of the active compound (g/10 ares)	Phytotoxicity to rice			Control effect		
	Active compound	Trans-plant-ation	Direct sowing	Barn-yard grass	Broad-leaved weeds	Slender spikerush
No. 16	1000	0	3	10	4	4
	500	0	1	8	3	1
	250	0	0	7	1	0
	125	0	0	1	0	0
No. 17	1000	0	6	3	4	0
	500	0	4	1	2	0
	250	0	2	0	0	0
	125	0	1	0	0	0
No. 18	1000	2	10	5	7	3
	500	0	7	4	5	0
	250	0	1	0	3	0
	125	0	0	0	0	0
No. 19	1000	0	10	6	6	3
	500	0	8	4	4	0
	250	0	3	3	2	0
	125	0	2	0	0	0
No. 20	1000	0	6	3	3	0
	500	0	3	2	2	0
	250	0	0	0	0	0
	125	0	0	0	0	0

TABLE 3 (continued)

Active compound	Amount of the active compound (g/10 ares)	Phytotoxicity to rice			Control effect		
		Trans-plant-ation	Direct sowing	Barn-yard grass	Broad-leaved weeds	Bulrush	Slender spikerush
No. 21	1000	0	2	10	2	6	4
	500	0	0	7	2	4	1
	250	0	0	3	1	1	0
	125	0	0	1	0	0	0
No. 22	1000	0	3	10	4	7	3
	500	0	1	5	3	3	1
	250	0	0	4	1	2	0
	125	0	0	2	0	1	0
No. 23	1000	0	2	10	3	6	3
	500	0	1	6	2	3	1
	250	0	0	2	1	0	0
	125	0	0	1	0	0	0
No. 24	1000	0	2	10	4	0	4
	500	0	1	6	1	0	1
	250	0	0	3	0	0	0
	125	0	0	1	0	0	0
No. 25	1000	0	0	6	3	3	2
	500	0	0	3	1	1	0
	250	0	0	1	0	0	0
	125	0	0	0	0	0	0
No. 26	1000	0	0	7	3	4	0
	500	0	0	4	2	2	0
	250	0	0	2	0	0	0
	125			0	0	0	0

TABLE 3 (continued)

Active compound	Amount of the active compound (g/10 ares)	Phytotoxicity to rice						Control effect		
		Trans-plant-ation	Direct sowing	Barn-yard grass	Broad-leaved weeds	Bulrush	Slender spikerush			
No. 27	1000	0	0	10	8	10	6			
	500	0	0	10	3	10	3			
	250	0	0	10	0	10	0			
	125	0	0	10	0	10	0			
No. 28	1000	0	0	10	7	10	4			
	500	0	0	10	4	10	2			
	250	0	0	10	1	10	0			
	125	0	0	10	0	10	0			
No. 29	1000	0	5	10	2	10	2			
	500	0	0	10	1	10	0			
	250	0	0	8	0	8	0			
	125	0	0	4	0	2	0			
No. 30	1000	0	0	10	3	10	3			
	500	0	0	10	1	0	6			
	250	0	0	9	0	0	0			
	125	0	0	1	0	0	0			
No. 31	1000	0	0	10	5	10	5			
	500	0	0	5	0	2	2			
	250	0	0	4	0	4	0			
	125	0	0	4	0	1	0			
No. 32	1000	0	0	10	6	10	6			
	500	0	0	10	9	9	1			
	250	0	0	9	0	3	6			
	125	0	0	8	0	0	1			

TABLE 3 (continued)

Active compound	Amount of the active compound (g/10 ares)	Phytotoxicity to rice				Control effect		
		Trans-plant-ation	Direct sowing	Barn-yard grass	Broad-leaved weeds	Bulrush	Slender spikerush	
No. 33	1000	0	0	10	2	10	2	
	500	0	0	9	0	8	0	
	250	0	0	9	0	6	0	
	125	0	0	8	0	4	0	
No. 34	1000	0	3	8	0	6	0	
	500	0	0	7	0	4	0	
	250	0	0	6	0	3	0	
	125	0	0	3	0	0	0	
No. 35	1000	0	0	10	3	10	2	
	500	0	0	10	1	10	0	
	250	0	0	8	0	8	0	
	125	0	0	4	0	2	0	
No. 36	1000	0	0	10	10	10	8	
	500	0	0	10	9	10	6	
	250	0	0	6	0	8	0	
	125	0	0	0	0	0	0	
No. 37	1000	0	0	8	8	8	0	
	500	0	0	6	6	4	0	
	250	0	0	0	0	0	0	
	125	0	0	0	0	0	0	
No. 38	1000	0	0	0	10	10	7	
	500	0	0	0	10	8	4	
	250	0	0	0	10	7	1	
	125	0	0	0	10	0	0	

TABLE 3 (continued)

		Control effect						
		Phytotoxicity to rice	Trans- plant- ation	Direct sowing	Barn- yard grass	Broad- leaved weeds	Bulrush	Slender spikerush
Amount of the active compound (g/10 acres)	Active compound							
No. 39	1000	0	5	10	9	6	5	0
	500	0	3	9	4	4	4	0
	250	0	1	8	0	0	0	0
	125	0	0	6	0	0	0	0
No. 40	1000	0	4	10	0	6	0	0
	500	0	1	9	0	4	0	0
	250	0	0	9	0	2	0	0
	125	0	0	8	0	0	0	0
No. 41	1000	0	2	10	3	6	3	1
	500	0	0	10	1	5	1	0
	250	0	0	8	0	0	0	0
	125	0	0	6	0	0	0	0
No. 42	1000	0	0	10	6	9	3	0
	500	0	0	10	4	6	0	0
	250	0	0	4	0	2	0	0
	125	0	0	2	0	0	0	0
No. 43	1000	0	0	8	0	4	0	0
	500	0	0	6	0	1	0	0
	250	0	0	2	0	0	0	0
	125	0	0	0	0	0	0	0
No. 44	1000	0	0	10	4	10	0	0
	500	0	0	10	8	8	4	1
	250	0	0	9	0	0	0	0
	125	0	0	8	0	0	0	0

TABLE 3 (continued)

		Phytotoxicity to rice				Control effect		
		Amount of the active compound (g/10 ares)	Trans-plant-ation	Direct sowing	Barn-yard grass	Broad-leaved weeds	Bulrush	Slender spikerush
No. 45	1000	0	3	10	5	6	5	5
	500	0	0	10	2	4	3	3
	250	0	0	10	0	3	0	0
	125	0	0	8	0	2	0	0
No. 46	1000	0	0	10	10	7	0	0
	500	0	0	9	6	4	0	0
	250	0	0	7	6	1	0	0
	125	0	0	4	4	1	0	0
No. 47	1000	0	6	10	4	10	3	3
	500	0	4	10	0	7	1	0
	250	0	2	6	0	4	0	0
	125	0	0	2	0	2	0	0
No. 48	1000	0	0	10	0	10	8	8
	500	0	0	10	0	10	4	4
	250	0	0	10	0	9	0	0
	125	0	0	9	0	8	0	0
No. 49	1000	0	0	10	8	5	0	0
	500	0	0	10	8	4	0	0
	250	0	0	9	0	2	0	0
	125	0	0	4	0	0	0	0
No. 50	1000	0	0	10	8	5	5	5
	500	0	0	10	7	10	4	4
	250	0	0	10	4	10	0	0
	125	0	0	0	0	10	8	8

TABLE 3 (continued)

Active compound	Amount of the active compound (g/10 ares)	Phytotoxicity to rice						Control effect
		Trans-plant-ation	Direct sowing	Barn-yard grass	Broad-leaved weeds	Bulrush	Slender spikerush	
No. 51	1000	0	0	9	6	3	6	
	500	0	0	4	4	1	0	
	250	0	0	2	2	0	0	
	125	0	0	1	0	0	0	
No. 52	1000	0	0	9	4	2	0	
	500	0	0	5	1	1	0	
	250	0	0	3	0	0	0	
	125	0	0	0	0	0	0	
No. 53	1000	0	0	10	3	3	3	
	500	0	0	6	1	0	0	
	250	0	0	3	0	0	0	
	125	0	0	0	0	0	0	
No. 54	1000	0	0	9	2	4	0	
	500	0	0	6	1	2	0	
	250	0	0	3	0	0	0	
	125	0	0	0	0	0	0	
No. 55	1000	0	0	8	4	1	2	
	500	0	0	2	1	0	0	
	250	0	0	0	0	0	0	
	125	0	0	0	0	0	0	
No. 56	1000	0	0	7	2	4	0	
	500	0	0	3	1	0	0	
	250	0	0	0	0	0	0	
	125	0	0	0	0	0	0	

TABLE 3 (continued)

Active compound	Amount of the active compound (g/10 ares)	Phytotoxicity to rice				Control effect
		Trans-plant-ation	Direct sowing	Barn-yard grass	Broad-leaved weeds	
No. 57	1000	0	0	6	3	3
	500	0	0	2	0	1
	250	0	0	1	0	0
	125	0	0	0	0	0
No. 58	1000	0	0	6	0	0
	500	0	0	4	0	0
	250	0	0	2	0	0
	125	0	0	0	0	0
No. 59	1000	0	0	7	0	0
	500	0	0	5	0	0
	250	0	0	3	0	0
	125	0	0	0	0	0
No. 60	1000	0	0	4	2	1
	500	0	0	2	0	0
	250	0	0	0	0	0
	125	0	0	0	0	0
No. 61	1000	0	0	5	0	1
	500	0	0	2	0	0
	250	0	0	0	0	0
	125	0	0	0	0	0
No. 62	1000	0	0	10	1	4
	500	0	0	3	0	1
	250	0	0	0	0	0
	125	0	0	0	0	0

TABLE 3 (cont...)

Active compound	Amount of the active compound (g/10 ares)	Phytotoxicity						Control effect
		tc. rice	Trans-plant-ation	Direct sowing	Barn-yard grass	Broad-leaved weeds	Slender spikerush	
No. 63	1000	0	3	10	10	10	10	7
	500	0	1	10	10	6	3	3
	250	0	0	10	4	8	1	1
	125	0	0	10	3	0	0	0
Known compound A (*)	1000	0	0	1	0	1	1	0
	500	0	0	0	0	0	0	0
	250	0	0	0	0	0	0	0
	125	0	0	0	0	0	0	0
Known compound B (*)	1000	0	0	1	0	1	1	0
	500	0	0	0	0	0	0	0
	250	0	0	0	0	0	0	0
	125	0	0	0	0	0	0	0
Benthio-carb (control)	500	6	10	10	10	10	10	10
	250	2	10	10	7	4	4	10
	125	0	10	10	4	1	1	6
	62.5	0	8	8	0	0	1	1
CNP (control)	500	7	10	10	10	9	4	6
	250	2	10	10	9	4	0	0
	125	0	10	6	1	0	0	0
	62.5	0	10	6	1	0	0	0
Untreated	-	0	0	0	0	0	0	0

(*) Known compound A: N-nethyl-N-benzyl-2-nitrobenzene-sulfonamide
 Known compound B: N-2-bromoethyl-N-benzoyl-4-methylbenzene-sulfonamide

Example E

Two sheets of filter paper were spread in a Petri dish with a diameter of 9 cm, and 6 ml of a wettable powder of each of the active compounds of this invention shown in Table 4 in a predetermined concentration was dropped onto them. Ten seeds each of barnyard grass and rice plant (variety: Nihonbare) were directly sown. The Petri dish was put into a thermostat chamber at 26°C., and 72 hours later, the lengths of young stems and leaves and young roots were measured.

The evaluation was made on a scale of 10 grades in which 10 represents complete inhibition and 0 represents normal growth.

10

5

10

TABLE 4

15	Active compound No.	Concen- tration (ppm)	Barnyard grass		Rice plant		15
			Young stem	Root	Young stem	Root	
20	64	1,000	7	9	0	0	20
		300	5	7	0	0	
		100	4	7	0	0	
		30	2	3	0	0	
		10	1	2	0	0	
25	65	1,000	10	9	0	0	25
		300	7	8	0	0	
		100	4	5	0	0	
		30	3	5	0	0	
		10	1	2	0	0	
30	66	1,000	9	10	0	0	30
		300	9	9	0	0	
		100	7	5	0	0	
		30	2	3	0	0	
		10	1	1	0	0	
35	67	1,000	10	10	3	2	35
		300	10	10	1	0	
		100	10	10	0	0	
		30	8	7	0	0	
		10	5	4	0	0	
40	68	1,000	10	10	0	0	40
		300	9	10	0	0	
		100	5	8	0	0	
		30	3	6	0	0	
		10	2	3	0	0	
45	69	1,000	7	10	0	0	45
		300	5	7	0	0	
		100	3	6	0	0	
		30	1	3	0	0	
		10	0	1	0	0	
50	69	1,000	7	10	0	0	50
		300	5	7	0	0	
		100	3	6	0	0	
		30	1	3	0	0	
		10	0	1	0	0	

TABLE 4 (continued)
Barnyard grass Rice plant

5	Active compound No.	Concen- tration (ppm)	Young stem	Root	Young stem	Root	5
10	70	1,000	10	10	3	1	
		300	8	10	0	0	
		100	5	7	0	0	10
		30	2	2	0	0	
		10	1	2	0	0	
15	71	1,000	7	6	0	0	
		300	6	6	0	0	15
		100	3	5	0	0	
		30	1	2	0	0	
		10	0	0	0	0	
20	72	1,000	6	8	0	0	20
		300	4	3	0	0	
		100	2	1	0	0	
		30	0	0	0	0	
		10	0	0	0	0	
25	73	1,000	8	10	0	0	
		300	6	9	0	0	
		100	4	5	0	0	
		30	1	3	0	0	
		10	0	0	0	0	30
35	74	1,000	10	10	4	1	
		300	9	10	2	0	
		100	7	9	0	0	
		30	3	4	0	0	35
		10	1	2	0	0	
40	75	1,000	10	10	5	4	
		300	10	10	2	1	
		100	10	10	0	0	
		30	9	10	0	0	
		10	9	7	0	0	40

TABLE 4 (continued)
Barnyard grass Rice plant

5	Active compound No.	Concen- tration (ppm)	Young stem	Root	Young stem	Root	5
10	76	1,000	10	10	7	6	
		300	10	10	3	2	
		100	10	10	0	0	10
		30	9	9	0	0	
		10	9	9	0	0	
15	77	1,000	10	10	5	2	
		300	10	10	4	1	15
		100	10	10	1	0	
		30	9	9	0	0	
		10	8	7	0	0	
20	78	1,000	10	10	7	3	20
		300	10	10	4	1	
		100	8	9	0	0	
		30	7	6	0	0	
		10	6	3	0	0	25
25	79	1,000	10	10	7	6	
		300	10	10	5	3	
		100	9	8	1	0	
		30	7	5	0	0	
		10	3	3	0	0	30
30	80	1,000	10	10	6	3	
		300	10	9	4	2	
		100	10	6	0	0	
		30	7	2	0	0	35
		10	2	1	0	0	
35	81	1,000	10	10	8	5	
		300	10	10	2	2	
		100	10	6	0	0	
		30	6	2	0	0	
		10	3	1	0	0	

TABLE 4 (continued)
Barnyard grass Rice plant

5	Active compound No.	Concentration (ppm)	Young stem	Root	Young stem	Root	5
10	82	1,000	8	10	0	0	10
		300	4	7	0	0	
		100	2	6	0	0	
		30	1	3	0	0	
		10	0	1	0	0	
15	83	1,000	8	10	1	0	15
		300	7	9	0	0	
		100	5	3	0	0	
		30	4	0	0	0	
		10	0	0	0	0	
20	84	1,000	8	10	0	0	20
		300	7	9	0	0	
		100	6	4	0	0	
		30	3	1	0	0	
		10	1	0	0	0	
25	85	1,000	10	10	6	7	25
		300	10	10	0	0	
		100	10	10	0	0	
		30	9	10	0	0	
		10	9	9	0	0	
30	86	1,000	10	10	7	4	30
		300	10	3	0	0	
		100	8	0	0	0	
		30	6	0	0	0	
		10	1	0	0	0	
35	87	1,000	5	10	8	9	35
		300	9	7	0	1	
		100	8	3	0	0	
		30	4	2	0	0	
		10	1	1	0	0	
40	87	1,000	5	10	8	9	40
		300	9	7	0	1	
		100	8	3	0	0	
		30	4	2	0	0	
		10	1	1	0	0	

TABLE 4 (continued)
 Barnyard grass Rice plant

5	Active compound No.	Concen- tration (ppm)	Young stem	Root	Young stem	Root	5
10	88	1,000	10	10	5	1	
		300	10	3	2	1	
		100	8	2	0	0	10
		30	6	0	0	0	
		10	2	0	0	0	
15	89	1,000	10	10	8	9	
		300	10	2	7	3	15
		100	10	0	5	1	
		30	10	0	0	0	
		10	7	0	0	0	
20	90	1,000	10	10	10	10	20
		300	10	10	6	4	
		100	10	4	6	0	
		30	10	0	2	0	
		10	9	0	1	0	25
25	91	1,000	10	10	10	10	
		300	10	8	8	4	
		100	10	3	7	0	
		30	10	0	1	0	
		10	9	0	0	0	30
30	92	1,000	10	10	9	9	
		300	10	3	5	2	
		100	10	0	0	0	
		30	6	0	0	0	35
		10	2	0	0	0	
35	93	1,000	10	10	10	10	
		300	10	9	5	1	
		100	10	7	1	0	40
		30	7	6	0	0	
		10	4	1	0	0	

TABLE 4 (continued)
Barnyard grass Rice plant

5	Active compound No.	Concen- tration (ppm)	Young stem	Root	Young stem	Root	5
10	94	1,000	10	10	9	9	
		300	10	7	6	1	
		100	7	4	2	0	10
		30	6	1	0	0	
		10	4	0	0	0	
15	95	1,000	10	10	8	7	
		300	10	10	1	3	15
		100	10	10	0	0	
		30	9	6	0	0	
		10	4	2	0	0	
20	96	1,000	10	10	7	5	20
		300	10	10	4	2	
		100	10	7	1	0	
		30	7	4	0	0	
		10	4	0	0	0	
25	97	1,000	10	10	10	10	25
		300	10	9	10	9	
		100	10	4	10	6	
		30	6	1	7	2	
		10	2	0	3	0	30
35	98	1,000	10	10	8	8	
		300	10	9	6	4	
		100	9	8	4	2	
		30	7	6	2	0	35
		10	4	2	0	0	
40	99	1,000	10	10	7	6	
		300	10	10	6	4	
		100	9	9	2	1	
		30	8	7	0	0	
		10	7	6	0	0	40

TABLE 4 (continued)
Barnyard grass Rice plant

5	Active compound No.	Concen- tration (ppm)	Young stem	Root	Young stem	Root	5
10	100	1,000	10	10	7	6	
		300	9	9	5	3	
		100	8	8	1	1	10
		30	7	6	0	0	
		10	6	4	0	0	
15	101	1,000	10	10	4	3	
		300	5	4	1	1	15
		100	4	2	0	0	
		30	1	0	0	0	
		10	0	0	0	0	
20	102	1,000	10	10	5	2	20
		300	6	5	2	1	
		100	5	4	0	0	
		30	2	1	0	0	
		10	0	0	0	0	25
25	Benthio- carb (con- trol)	1,000	10	10	10	10	
		300	10	9	10	9	
		100	10	4	10	6	
		30	6	1	7	2	30
		10	2	0	3	0	
Untreated		—	0	0	0	0	

35 *Example F*

A paddy soil was put into pots (1/5000 are), and the surface soil to a depth of 1 cm and 30 seeds of barnyard grass were mixed. Ten seeds of rice (variety: Nihonbare) were sown on the surface of the soil, and two seedlings of rice each in the 1-leaf stage, 2-leaf stage, and 3.5-leaf stage respectively were transplanted as one stock. The pots were irrigated to a water depth of 2 cm. One day later, a predetermined amount of each of the active compounds of this invention indicated in Table 5 was diluted with 10 ml of water, and dropped onto the soil surface in each pot. The control effect and phytotoxicity were examined in the same manner as in Example D 20 days after the treatment. The results are shown in Table 5.

35

40

40

TABLE 5

5	Active compound	Amount of the active compound (g/10 ares)	Control effect on barnyard grass	Phytotoxicity to rice			5
				Directly sown	1-leaf stage	2-leaf stage	
10	No. 27	1000	10	0	0	0	0
		500	10	0	0	0	0
		250	10	0	0	0	0
		125	10	0	0	0	0
		62.5	10	0	0	0	0
15	No. 28	1000	10	0	0	0	0
		500	10	0	0	0	0
		250	10	0	0	0	0
		125	10	0	0	0	0
		62.5	10	0	0	0	0
20	No. 67	1000	10	0	0	0	0
		500	10	0	0	0	0
		250	10	0	0	0	0
		125	10	0	0	0	0
		62.5	7	0	0	0	0
25	No. 85	1000	10	0	0	0	0
		500	10	0	0	0	0
		250	10	0	0	0	0
		125	9	0	0	0	0
		62.5	8	0	0	0	0
30	Benthio-carb (control)	1000	10	10	10	9	7
		500	10	10	10	6	3
		250	10	10	5	1	0
		125	10	10	2	0	0
		62.5	9	8	0	0	0
35	CNP (control)	1000	10	10	10	10	9
		500	10	10	8	7	4
		250	10	10	7	4	1
		125	10	10	5	2	0
		62.5	7	7	3	0	0
45	Untreated	—	0	0	0	0	45

Example G

A test on an enlarged scale was conducted under assumed field conditions.

50 Paddy soil was put into a concrete pot having a length of 70 cm, a width of 70 cm and a height of 50 cm and rendered muddy. Sixty seeds of barnyard grass (*Echinochloa crus-galli* Beauv.) were sown, and the pot was irrigated to a water depth of 3 cm. Further, 20 germinated seeds of rice (variety: Nihonbare) were sown on the soil. Three-leaf stage rice seedlings of the same variety of rice were planted in four stocks each consisting of two seedlings. One day later, a granule of each of the active compounds of this invention shown in Table 6 and Benthiocarb as a control was sprayed by hand in each of the amounts indicated in Table 6. The control effect (broad-leaved grasses were those which occurred naturally) and phytotoxicity to rice were examined two weeks after the treatment. The results are shown in Table 6.

50

55

TABLE 6

Active compound	Amount of the active compound (g/10 ares)	Phytotoxicity to rice			Control effect		
		Direct sowing	Trans-plantation	Barnyard grass	Broad-leaved weeds		
No. 27	1000	0	0	10	2		
	500	0	0	10	0		
	250	0	0	10	0		
	125	0	0	7	0		
No. 28	1000	0	0	10	0		
	500	0	0	10	0		
	250	0	0	10	0		
	125	0	0	6	0		
No. 31	1000	0	0	10	0		
	500	0	0	10	0		
	250	0	0	8	0		
	125	0	0	6	0		
No. 32	1000	0	0	10	0		
	500	0	0	10	0		
	250	0	0	6	0		
	125	0	0	3	0		
No. 33	1000	0	0	10	0		
	500	0	0	8	0		
	250	0	0	5	0		
	125	0	0	4	0		
No. 67	1000	0	0	10	0		
	500	0	0	10	0		
	250	0	0	10	8		

TABLE 6 (continued)

Active compound	Amount of the active compound (g/10 ares)	Phytotoxicity to rice			Control effect
		Direct sowing	Trans-plantation	Barnyard grass	
No. 85	1000	0	0	10	0
	500	0	0	10	0
	250	0	0	7	0
	125	0	0	5	0
Benthio-carb (control)	500	10	3	10	8
	250	10	1	10	5
	125	8	0	9	2
	62.5	3	0	4	1
Untreated	—	0	0	0	0

Example H

Silt loam (a clay content of 48.5%) and sandy soil (a clay content of 15.4%) were each placed in pots (1/5000 are), and agitated until a paddy field condition was attained. Thirty seeds of barnyard grass (*Echinochloa crus-galli* Beauv.) were sown per pot. Further, ten germinated seeds of rice were sown, and rice seedlings in the three-leaf stage were planted in two stocks each consisting of two seedlings. The soil surface was submerged to a water depth of 3 cm. One day later, a wettable powder of each of the active compounds of this invention indicated in Table 7 was applied. Two weeks after the treatment, the control effect was examined in the same way as in Example D. The results are shown in Table 7.

Active compound	Amount of the active compound (g/10 ares)	Sandy soil		Silt loam	
		Directly sown rice plant	Transplanted rice plant	Directly sown rice plant	Transplanted rice plant
No. 27	500	10	0	10	0
	250	10	0	10	0
	125	10	0	10	0
	62.5	10	0	10	0
No. 31	500	10	0	10	0
	250	10	0	7	0
	125	10	0	4	0
	62.5	4	0	1	0
No. 32	500	10	0	10	0
	250	10	0	9	0
	125	9	0	8	0
	62.5	7	0	4	0
No. 33	500	10	0	10	0
	250	10	0	9	0
	125	8	0	7	0
	62.5	6	0	4	0
Butachlor (control)	500	10	10	10	10
	250	10	10	10	10
	125	10	10	10	10
	62.5	10	4	7	8
Benthio-carb (control)	500	10	10	10	10
	250	10	10	10	10
Untreated	125	10	10	6	3
	62.5	10	10	3	0
Untreated		—	0	0	0

Example I

Three-leaf stage rice seedlings were machine-transplanted in a paddy field, and four days after, the paddy field was divided into areas of 2 m × 2 m by vinyl resin plates. A wettable powder of each of the active compounds of this invention shown in Table 8 in a predetermined amount was diluted with 400 ml of water per area and sprayed onto the entire water surface. The control effect and phytotoxicity to rice were examined three weeks later.

The evaluation of the control effect was made on a scale of 10 grades. The phytotoxicity to rice was evaluated on the following standards:

- : Normal
- +: slight injury
- ++: minor injury
- +++: moderate injury

The results are shown in Table 8.


TABLE 8

Active compound	Amount of the active compound (g/10 ares)	Control effect				Toxicity to rice plants
		Barnyard grass	Bulrush	Umbrella plant	Broad-leaved weeds	
No. 27	1000	10	8	10	5	-
	500	10	6	9	3	-
	250	7	4	4	1	-
	125	4	2	3	1	-
No. 67	1000	10	7	10	6	-
	500	9	5	9	4	-
	250	7	3	6	1	-
	125	4	1	3	0	-
No. 85	1000	10	7	10	8	+
	500	8	6	9	5	++
	250	5	4	6	1	++
	125	3	1	1	0	++
Mo (control)	500	10	10	9	4	-
	250	8	5	6	1	-
	125	4	2	2	0	-
Untreated		0	0	0	0	-

WHAT WE CLAIM IS:

1. A compound of the general formula

5



5

10

wherein R_0 represents an alkyl group optionally having a substituent selected from a cyano group, lower alkoxy groups and di(lower alkyl)amino groups, a lower alkenyl group, or a lower alkynyl group; R_1 , R_2 and R_3 , independently from each other, represent a hydrogen atom, a halogen atom, a lower alkyl group, or a lower alkoxy group; R_4 , R_5 , R_6 , R_7 and R_8 , independently from each other, represent a hydrogen atom, a halogen atom, a lower alkyl group, or a lower alkoxy group; Y represents a group of the formula

10

15



15

25

in which one of R_9 and R_{10} is a lower alkyl group and the other is a hydrogen atom or a lower alkyl group; with the proviso that when Y is the group

25

30



30

and R_0 represents an unsubstituted alkyl group, R_4 , R_5 , R_6 , R_7 and R_8 do not represent hydrogen atoms at the same time.

35

2. A compound of claim 1 wherein R_0 represents an alkyl group containing up to 15 carbon atoms which may contain a substituent selected from a cyano group, lower alkoxy groups and di-(lower alkyl)amino groups; a lower alkenyl group; or a propargyl group.

35

3. A compound of claim 1 wherein R_0 is an allyl or propargyl group.

4. A compound of claim 1, 2 or 3 wherein Y represents the group

40

40



45


and at least one of R_4 , R_5 , R_6 , R_7 and R_8 is a halogen atom and the rest are hydrogen atoms.

45

5. A compound of claim 1, 2 or 3 wherein Y represents the group

50

50



55


55

and R_4 , R_5 , R_6 , R_7 and R_8 all represent hydrogen atoms.

6. A compound of claim 1 which is a compound of the general formula

60

60




65

65

wherein R_{01} represents an alkyl group containing up to 15 carbon atoms which may contain

5 a substituent selected from a cyano group, lower alkoxy groups and di-(lower alkyl)amino groups, a lower alkenyl group, or a propargyl group; R₁₁ and R₂₁, independently from each other, represent a hydrogen atom, a halogen atom, a methyl group, an ethyl group, or a methoxy group, R₄₁, R₅₁, R₆₁, R₇₁ and R₈₁, independently from each other represent a hydrogen atom, a halogen atom, a methyl group or a methoxy group; Y₁ represents the group

5



10

15 in which one of R₉₁ and R₁₀₁ is a methyl or ethyl group, and the other is a hydrogen atom or a methyl or ethyl group; with the proviso that when Y₁ is the group CO and R₀₁ is an unsubstituted alkyl group, R₄₁, R₅₁, R₆₁, R₇₁ and R₈₁ do not represent hydrogen atoms at the same time.

15

20 7. A compound of claim 6 wherein R₀₁ represents an allyl or propargyl group, and Y₁ represents the group -CO-, -CH(CH₃)- or -C(CH₃)₂-.

20

8. A compound of claim 6 wherein R₀₁ represents an allyl or propargyl group and Y₁ represents the group -CO-.

9. A compound of claim 8 wherein R₄₁ is a halogen atom, and one of R₅₁, R₆₁, R₇₁ and R₈₁ is a hydrogen atom or a halogen atom, and the rest are hydrogen atoms.

25 10. A compound of claim 8 wherein R₄₁ is a halogen atom, R₁₁ and R₂₁, independently from each other, represent a hydrogen atom or a methyl group and R₅₁, R₆₁, R₇₁ and R₈₁, independently from each other, represent a hydrogen atom or a methyl group.

25

11. A compound of claim 9 wherein R₁₁ and R₂₁, independently from each other, represent a hydrogen atom or a methyl group.


30 12. A compound of claim 6 wherein R₀₁ is an allyl group and Y₁ is the group -CH(CH₃)- or -C(CH₃)₂-.

30

13. A compound of claim 12 wherein R₁₁ represents a hydrogen atom, a methyl group or a methoxy group, R₂₁ is a hydrogen atom, R₄₁ and R₆₁, independently from each other, represent a hydrogen atom, a halogen atom, or a methyl group and R₅₁, R₇₁ and R₈₁ are all hydrogen atoms.

35

35 14. A compound of claim 1 which is a compound of the formula



40

wherein R₀₂ is an allyl or propargyl, and X₁ is a hydrogen or chlorine atom.

45 15. The compound of claim 1 which is N-allyl-N-(2-chlorobenzoyl)benzenesulfonamide.


45

16. The compound of claim 1 which is N-allyl-N-(2,4-dichlorobenzoyl)benzenesulfonamide.

50 17. The compound of claim 1 which is N-propargyl-N-(2-chlorobenzoyl)benzenesulfonamide.

50

18. A compound of claim 1 which is a compound of the formula



55

wherein Y₂ is the group



60

65 and X₂ is a hydrogen or chlorine atom.

65

19. The compound of claim 1 which is N-allyl-N-(α , α -dimethylbenzyl)benzenesulfonamide. 5

20. The compound of claim 1 which is N-allyl-N-(α -methylbenzyl)benzenesulfonamide.

21. The compound of claim 1 which is N-allyl-N-(α -methyl-4-chlorobenzyl)benzenesulfonamide. 5

22. The compound of claim 1 which is N-allyl-N-(α , α -dimethyl-4-chlorobenzyl)benzenesulfonamide. 5

23. Any of the compounds according to claim 1 that are hereinbefore described in the Examples, other than the compounds of claims 15-17 and 19-22. 10

24. A process for preparing a compound according to claim 1, which comprises (a) reacting a benzenesulfonamide derivative of the general formula 10

15

wherein M represents a hydrogen atom or an alkali metal atom, and R₀, R₁, R₂ and R₃ are as defined in claim 1, with a compound of the general formula 15

20

25

wherein X₃ is a halogen atom, and R₄, R₅, R₆, R₇, R₈ and Y are as defined in claim 1, or (b) reacting a benzenesulfonyl halide of the general formula 25

30

wherein X₄ is a halogen atom, and R₁, R₂ and R₃ are as defined in claim 1, with a compound of the general formula 30

35

40

45

50

55

60

65

wherein R₀, R₄, R₅, R₆, R₇, R₈ and Y are as defined in claim 1, or (c) reacting a compound of the general formula 40

40

45

50

55

60

65

wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, Y and M are as defined in claim 1 with a compound of the formula 45

R₀ - X₄ (VII) 50

wherein X₄ is a halogen atom, and R₀ is as defined in claim 1. 50

25. A herbicidal composition as an active ingredient a compound of the general formula (I) according to any of claims 1 to 23, in admixture with an inert, liquid or solid diluent or carrier. 55






26. A herbicide of claim 25 wherein the amount of the compound of formula (I) is at least 0.5% by weight. 55

27. A herbicide of claim 26 wherein the amount of the compound of formula (I) is 1 to 90% by weight. 60

28. A herbicide of claim 25, 26 or 27 which is in the form of a dust, granule, wettable powder, solution or emulsifiable concentrate. 60

29. A method for controlling weeds in an agricultural crop which comprises applying a compound of the general formula (I) according to any of claims 1 to 23 to the locus to be protected from the weeds. 65

30. A method of claim 29 wherein the compound of formula (I) is applied before or during the germination of the weeds. 65



31. A method of claim 29, 30 or 31 wherein the agricultural crop is rice.
33. A method of any of claims 29 to 32 wherein the weed is barnyard grass, *Echinochloa crus-galli* Beauv.
34. A method of any of claims 29 to 33 wherein the amount of the compound of formula 5 (I) applied is at least 25 g/10 ares.
35. A herbicide according to claim 25 substantially as hereinbefore described in any of the Examples A to I.
36. A method according to claim 29 substantially as hereinbefore described in any one of the Examples D to I.

10

5

10

15

15

For the Applicants
CARPMAELS & RANSFORD,
Chartered Patent Agents,
43 Bloomsbury Square,
London WC1A 2RA.

Printed for Her Majesty's Stationery Office, by Croydon Printing Company Limited, Croydon, Surrey, 1980.
Published by The Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from
which copies may be obtained.