



US00PP16560P3

(12) **United States Plant Patent**  
**Delabroye**

(10) **Patent No.:** **US PP16,560 P3**

(45) **Date of Patent:** **May 23, 2006**

(54) **HEUCHERA PLANT NAMED 'CARMEL'**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(50) Latin Name: *Heuchera villosa*  
Varietal Denomination: **Caramel**

(52) **U.S. Cl.** ..... **Plt./263**

(58) **Field of Classification Search** ..... **Plt./263**  
See application file for complete search history.

(76) Inventor: **Sandrine Delabroye**, Rue Roger  
Salengro 40, Hantay (FR), F-59496

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 375 days.

*Primary Examiner*—Anne Marie Grunberg  
(74) *Attorney, Agent, or Firm*—Penny J. Aguirre

(57) **ABSTRACT**

A new cultivar of *Heuchera villosa*, 'Caramel', character-  
ized by its uniquely colored foliage of cordate-shaped leaves  
that emerge greyed-orange and mature to golden yellow-  
green, its vigorous growth habit and its resistance to disease.

(21) Appl. No.: **10/935,969**

(22) Filed: **Sep. 7, 2004**

(65) **Prior Publication Data**

US 2006/0053519 P1 Mar. 9, 2006

**2 Drawing Sheets**

**1**

**2**

Botanical classification: *Heuchera villosa*.  
Cultivar designation: 'Caramel'.

4. 'Caramel' is vigorous grower compared to the growth  
rate that is typical for *Heuchera* with colored foliage  
and has been shown to be resistant to diseases under the  
conditions tested.

**BACKGROUND OF THE INVENTION**

**BRIEF DESCRIPTION OF THE DRAWING**

The present invention relates to a new and distinct cultivar  
of *Heuchera villosa*, and is hereinafter referred to by the  
cultivar name 'Caramel'.

The inventor discovered the new cultivar, 'Caramel' as a  
chance seedling in a cultivated nursery bed in Hantay,  
France, CT in 2003. Although the parentage is unknown, the  
characteristics of the new cultivar and the proximity of  
plants of *Heuchera villosa* 'Autumn Pride' (not patented)  
suggests that 'Autumn Pride' is a probable parent.

The new cultivar was selected for its unique foliage that  
emerges grey-red in color and matures to a golden yellow-  
green color. The foliage color of 'Autumn Pride' is green.  
'Caramel' is most similar in foliage coloration to *Heuchera*  
'Amber Waves' (U.S. Plant Pat. No. 13,348), a *Heuchera* of  
hybrid origin. 'Caramel' differs in having foliage that is less  
ruffled, more cordate in shape, and more sunfast.

Asexual reproduction of the new cultivar was first accom-  
plished under direction by the inventor by in vitro propa-  
gation in The Netherlands in the fall of 2003. Asexual  
reproduction of the new cultivar by division and tissue  
culture has shown that the unique features of 'Caramel' are  
stable and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and  
represent the characteristics of the new cultivar. These  
attributes in combination distinguish 'Caramel' as a new and  
unique cultivar of *Heuchera*.

1. 'Caramel' has foliage with the unique coloration of  
emerging greyed-orange and turning to a golden  
yellow-green on maturity.
2. The leaves of 'Caramel' are cordate in shape and fairly  
flat in aspect.
3. 'Caramel' reaches a plant height of 60 cm in bloom.

The accompanying colored photographs illustrate the  
overall appearance and distinct characteristics of the new  
*Heuchera*. The photographs were taken in The Netherlands  
in April of a one year-old plant grown outdoors in a field  
plot.

The photograph on the first sheet provides an overall view  
of foliage characteristics of new cultivar.

The photographs on the second sheet are close-up views  
of the foliage of 'Caramel'; the top photo provides a view of  
the foliage as it emerges and changes with maturity  
(clockwise) and the bottom photograph is of a mature leaf.  
The colors in the photographs are as close as possible with  
the photographic and printing technology utilized. The color  
values cited in the detailed botanical description accurately  
describe the colors of the new *Heuchera*.

**DETAILED BOTANICAL DESCRIPTION OF  
THE PLANT**

The following is a detailed description of a one year-old  
plant of the new cultivar as grown outdoors in The Nether-  
lands. The phenotype of the new cultivar may vary with  
variations in environmental, climatic, and cultural  
conditions, as it has not been tested under all possible  
environmental conditions. The color determination is in  
accordance with the 2001 R.H.S. Colour Chart of The Royal  
Horticultural Society, London, England, except where gen-  
eral color terms of ordinary dictionary significance are used.

Botanical classification: 'Caramel' is a cultivar of *Heuchera  
villosa*.

Parentage: Chance seedling, most probably a seedling of  
*Heuchera villosa* 'Autumn Pride'.

## General description:

*Plant habit.*—Clump-forming herbaceous perennial, mounded foliage.

*Height and spread.*—Reaches about 60 cm in height in bloom and about 28 cm in width. Foliage reaches a height of about 12 cm.

*Hardiness.*—Cold hard to at least USDA Zone 6.

*Culture.*—Full sun to medium shade in moist, well-drained, fertile soils.

*Diseases and pests.*—Disease free in the conditions tested, no susceptibility or resistance to pests has been observed.

*Root description.*—Fibrous roots on woody rootstalks.

*Branching habit.*—Basal rosettes of leaves on petioles.

*Propagation.*—In vitro propagation is the preferred method, division are also possible.

*Root initiation.*—Roots appear in rooting media in 10 day at 20° C. in the laboratory without supplemental lighting.

*Root development.*—Rooted transplants from tissue culture fully develop in a 2.5 in container in about 20 days in a greenhouse with average temperatures of about 20° C. without supplemental lighting in The Netherlands.

*Growth rate.*—Vigorous compared to what is typically observed for *Heuchera* with colored foliage. A 2.5 inch container will fully develop in a one-gallon container in 2 to 2.5 months in the Netherlands.

## Foliage description:

*Leaf shape.*—Cordate.

*Leaf division.*—Simple, but lobed with about 7 lobes per leaf.

*Leaf base.*—Strongly cordate (leaves at base overlap).

*Leaf apex.*—Acute to rounded.

*Leaf venation.*—Palmate, conspicuous but not prominent, vein color on upper surface of mature leaves 152D, vein color on lower surface of mature leaves N170D.

*Leaf margins.*—Crenate, average of 2.5 teeth per cm.

*Leaf attachment.*—Petiolate.

*Leaf arrangement.*—Basal rosettes.

*Leaf orientation.*—Held horizontal to petiole, held nearly flat (not ruffled or curled).

*Leaf surface.*—Upper: pubescent with 1 mm stiff white hairs, lower: glabrous with pubescence on veins.

*Leaf color.*—Young leaves (upper surface): greyed-orange 176A with margin area 177B. Young leaves (lower surface): greyed-red to greyed-purple ranges from 178A to 183C. Mature leaves (upper surface): yellow-green 152A to 152B (color slightly more brownish as if overlaid with 199A). Mature leaves (lower surface): greyed-red 182B to 182C.

*Leaf size.*—About 8 cm in length, up to about 7.7 cm in width.

*Leaf quantity.*—About 11 per basal rosette.

*Petiole size.*—Up to about 9 cm in length, about 2 mm in width.

*Petiole color.*—182C.

*Petiole shape.*—Round.

*Stipules.*—2 leafy stipules at the base of each petiole, oblong in shape, 1 cm in length, 2 mm in width, color 182C (both surfaces), margins sparsely covered with short hairs.

Flower and seed description: Flowering has not been observed on the plants under the conditions tested.

It is claimed:

1. A new and distinct cultivar of *Heuchera* plant named 'Caramel' as herein illustrated and described.

\* \* \* \* \*



