



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2722776 A1 2009/11/12

(21) **2 722 776**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2009/05/11
(87) Date publication PCT/PCT Publication Date: 2009/11/12
(85) Entrée phase nationale/National Entry: 2010/10/27
(86) N° demande PCT/PCT Application No.: US 2009/043502
(87) N° publication PCT/PCT Publication No.: 2009/137843
(30) Priorité/Priority: 2008/05/09 (US61/127,098)

(51) Cl.Int./Int.Cl. *A61K 31/496* (2006.01),
A61K 31/4465 (2006.01), *A61P 25/24* (2006.01)
(71) Demandeur/Applicant:
EMORY UNIVERSITY, US
(72) Inventeurs/Inventors:
DINGLEDINE, RAYMOND J., US;
TRAYNELIS, STEPHEN F., US
(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : ANTAGONISTES DES RECEPTEURS NMDA UTILISABLES POUR LE TRAITEMENT DE TROUBLES
NEUROPSYCHIATRIQUES
(54) Title: NMDA RECEPTOR ANTAGONISTS FOR THE TREATMENT OF NEUROPSYCHIATRIC DISORDERS



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

REVISED VERSION

(19) World Intellectual Property Organization
International Bureau



PCT



(10) International Publication Number
WO 2009/137843 A9

(51) International Patent Classification:
A61K 31/496 (2006.01) *A61P 25/24* (2006.01)
A61K 31/4465 (2006.01)

(21) International Application Number:
PCT/US2009/043502

(22) International Filing Date:
11 May 2009 (11.05.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/127,098 9 May 2008 (09.05.2008) US

(71) Applicant (*for all designated States except US*):
EMORY UNIVERSITY [US/US]; 1599 Clifton Road,
NE, 4th Floor, Atlanta, GA 30322 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): DINGLELINE,
Raymond, J. [US/US]; 2695 Rangewood Dr. NE, At-
lanta, GA 30345 (US). TRAYNELIS, Stephen, F.
[US/US]; 2640 Pangborn Rd, Decatur, GA 30033 (US).

(74) Agents: KAUFMAN, Rebecca et al.; King& Spalding,
1180 Peachtree St., Atlanta, GA 30309 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ,
EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO,
NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG,
SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR),
OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
MR, NE, SN, TD, TG).

Published:
— *with declaration under Article 17(2)(a); without abstract;
title not checked by the International Searching Authority*

(48) Date of publication of this revised version:
11 March 2010

(15) Information about Correction:
see Notice of 11 March 2010



WO 2009/137843 A9

(54) Title: NMDA RECEPTOR ANTAGONISTS FOR THE TREATMENT OF NEUROPSYCHIATRIC DISORDERS

(57) Abstract:

NMDA RECEPTOR ANTAGONISTS FOR THE TREATMENT OF NEUROPSYCHIATRIC DISORDERS

CROSS-REFERENCE TO RELATED APPLICATION

The application claims priority to U.S. Provisional Patent Application No. 61/127,098, filed May 9, 2008.

FIELD OF THE INVENTION

The present invention provides certain NMDA receptor blockers, including pH-sensitive NMDA receptor blockers, in the treatment of neuropsychiatric disorders including depression, anxiety and other related diseases.

BACKGROUND OF THE INVENTION

Glutamate and aspartate play dual roles in the central nervous system as essential amino acids and as the principal excitatory neurotransmitters (hereinafter referred to as excitatory amino acids or EAAs). There are at least four classes of EAA receptors: NMDA, AMPA (2-amino-3-(methyl-3-hydroxyisoxazol-4-yl)propanoic acid), kainate and metabotropic receptors. These EAA receptors mediate a wide range of signaling events that impact all physiological brain functions. For example, it has been reported that NMDA receptor antagonists produce an analgesic effect under certain conditions (Wong, et al. (1995) *Acta Anaesthesiologica. Sinica* 33, 227-232).

The NMDA subtype of glutamate-gated ion channels mediates excitatory synaptic transmission between neurons in the central nervous system (Dingledine et al. (1999), *Pharmacological Reviews* 51:7-61). NMDA receptors participate in a wide range of both physiological and pathological processes in the central nervous system. A high density of NMDA receptors has been found in the cortico-limbic regions of the brain which have been postulated to play a role in emotional functions, anxiety and depression (Tzschentke TM (2002) *Amino Acids* 23:147-152). Extensive studies have demonstrated antidepressant-like effects of various antagonists of the NMDA receptors. The antidepressant-like activity of competitive and non-competitive antagonists and inorganic inhibitors of NMDA receptor (zinc and magnesium) has been reported (see Decollogne, et al. (1997) *Pharmacol Biochem Behav* 58:261-268; Krocza, et al. (2001) *Brain Res Bull* 55:297-300; Krocza, et al. (2000)

Pol J Pharmacol. 52:403–406; Poleszak, et al. (2004) *Pharmacol Biochem Behav* 78:7–12; Poleszak, et al. (2007) *Pharmacol Biochem Behav* 88:158–164; Poleszak, et al. (2007) *Pharmacol Rep* 57:654–658; Przegaliński, et al. (1997) *Neuropharmacology* 36:31–37; Przegaliński, et al. (1998) *Pol J Pharmacol* 50: 349–354; Skolnick P *Eur J Pharmacol* 375:31–40; Skolnick, et al. (2001) *Pharmacol Res* 43:411–423; and Trullas, et al. (1990) *Eur J Pharmacol* 185:1–10). Poleszak, et al. showed that the NMDA receptor binding of certain antagonists, specifically CGP 37849 and L-701,324, are directly related to their antidepressant-like effects (Poleszak, et al. (2007) *Pharm. Reports* 59:595-600).

NMDA receptors are composed of NR1, NR2 (A, B, C, and D), and NR3 (A and B) subunits, which determine the functional properties of native NMDA receptors. Expression of the NR1 subunit alone does not produce a functional receptor. Co-expression of one or more NR2 subunits is required to form functional channels. In addition to glutamate, the NMDA receptor requires the binding of a co-agonist, glycine, to allow the receptor to function. A glycine binding site is found on the NR1 and NR3 subunits, whereas the glutamate binding site is found on NR2 subunits. At resting membrane potentials, NMDA receptors are largely inactive due to a voltage-dependent block of the channel pore by magnesium ions. Depolarization releases this channel block and permits passage of calcium as well as other ions.

The NMDA receptor is modulated by a number of endogenous and exogenous compounds including, sodium, potassium and calcium ions that can not only pass through the NMDA receptor channel but also modulate the activity of receptors. Zinc blocks the channel through NR2A- and NR2B-containing receptors noncompetitive and voltage-independent manner. Polyamines can also either potentiate or inhibit glutamate-mediated responses.

Neuropsychiatric disorders including schizophrenia and bipolar disorder and mood disorders affect more than 60 million Americans each year. Four basic forms of mood disorders are major depression, cyclothymia (a mild form of bipolar disorder), SAD (seasonal affective disorder) and mania (euphoric, hyperactive, over inflated ego, unrealistic optimism.) About 20% of the U.S. population reports at least one depressive symptom in a given month, and 12% report two or more in a year. A survey conducted in 1992 found rates of major depression reaching 5% in the previous 30 days, 17% for a lifetime. Bipolar disorder is less common, occurring at a rate of 1% in the general population, but some believe the diagnosis is often overlooked because manic elation is too rarely reported as an illness.

Depression, formally called major depression, major depressive disorder or clinical depression, is a medical illness that involves the mind and body. Most health professionals

today consider depression a chronic illness that requires long-term treatment, much like diabetes or high blood pressure. Although some people experience only one episode of depression, most have repeated episodes of depression symptoms throughout their life. Depression is also a common feature of mental illness, whatever its nature and origin. A person with a history of any serious psychiatric disorder has almost as high a chance of developing major depression as someone who has had major depression itself in the past. Most people with major depression also show some signs of anxiety, and 15-30% have panic attacks.

Depression is associated with physical illness as well. Some 25% of hospitalized medical patients have noticeable depressive symptoms and about 5% are suffering from major depression. Chronic medical conditions associated with depression include heart disease, cancer, vitamin deficiencies, diabetes, hepatitis, and malaria. Depression also is a common effect of neurological disorders, including Parkinson's and Alzheimer's diseases, multiple sclerosis, strokes, and brain tumors. Even moderate depressive symptoms are associated with a higher than average rate of arteriosclerosis, heart attacks, and high blood pressure. Depression can mimic medical illness and any illness feels worse to someone suffering from depression.

It's not known specifically what causes depression. As with many mental illnesses, it's thought that a variety of biochemical, genetic and environmental factors may cause depression. Despite the many advances that occurred from a better understanding of neuropharmacology, many psychiatric diseases remain untreated or inadequately treated with current pharmaceutical agents. In addition, many of the current agents interact with a number of cellular targets, potentially resulting in side effects that can greatly influence the overall outcome of therapy.

Numerous treatments for depression are available, including dozens of medications. Typical protocols include a selective serotonin reuptake inhibitor (SSRI). SSRIs include fluoxetine (Prozac, Sarafem), paroxetine (Paxil), sertraline (Zoloft), citalopram (Celexa) and escitalopram (Lexapro). Other common first choices for antidepressants include serotonin and norepinephrine reuptake inhibitors (SNRIs), norepinephrine and dopamine reuptake inhibitors (NDRIs), combined reuptake inhibitors and receptor blockers, and tetracyclic antidepressants. Tricyclic antidepressants (TCAs) are also effective, but because TCAs tend to have more numerous and more severe side effects, they are often less prescribed. Monoamine oxidase inhibitors (MAOIs) are often prescribed as a last resort, when other medications haven't worked.

Functional antagonists of the NMDA receptor complex exhibit antidepressant-like activity in the rodent test and models of depression. In 1990, Trullas and Skolnick demonstrated the antidepressant activity of AP-7, MK-801 and ACPC in the mouse forced swim test (FST) and tail suspension test (TST) (Trullas R, Skolnick P (1990) *Eur J Pharmacol* 185:1–10). Since then, a number of reports have confirmed and extended this finding. The NMDA antagonists are active in the FST in mice (Layer , et al. (1995) *Pharmacol Biochem Behav* 52:621–627; Maj et al. (1992) *Pol J Pharmacol* 44:337–346) and rats (Moryl , et al. (1993) *Pharmacol Toxicol* 72:394–397; Przeglasiński, et al (1997) *Neuropharmacology* 36:31–37) and tail suspension test in mice (Layer , et al. (1995) *Pharmacol Biochem Behav* 52:621–627), and in learned helplessness (Meloni, et al. (1993) *Pharmacol Biochem Behav* 46:423–426), chronic unpredicted stress (Ossowska, et al. (1997) *J Physiol Pharmacol* 48:127–135), chronic mild stress (Papp, et al. *Eur J Pharmacol* 263:1–7), and bulbectomy models (Redmond, et al. (1997) *Pharmacol Biochem Behav* 58:355–359). NMDA antagonists also demonstrate efficacy in clinical studies. Ketamine appears effective in major depression (Berman, et al. (2000) *Biol Psychiatry* 47:351–354; Zarate, et al. (2006) *Arch Gen Psychiatry* 63:856–864), although the clinical efficacy of memantine is not as clear (Ferguson, et al. (2007) *Clin Neuropharmacol* 30:136–144; Zarate, et al. (2006) *Am J Psychiatry* 163:153–155). Furthermore, the palliative effect of non-specific NMDA antagonist (amantadine and zinc) supplementation to antidepressant therapy has been suggested. On the other hand, antidepressants induce adaptive changes in the NMDA receptor complex (Skolnick , et al. (1996) *Pharmacopsychiatry* 29:23–26; Skolnick, et al. (2001) *Pharmacol Res* 43:411–423). Alterations in this receptor complex were demonstrated in the animal paradigm used for antidepressant screening (FST), in models of depression (Nowak, et al. (1998) *Pol J Pharmacol* 50:365–369; Nowak, et al. (1995) *J Neurochem* 64:925–927) and suicide victims (Nowak , et al. (1995) *Brain Res* 675:157–164). Thus, depression may be associated with enhanced NMDA signal transduction and the mechanism of antidepressant effect is related to reduction of this transmission.

U.S. Patent No. 7,019,016 to Pfizer provides methods for treating certain disorders including depression which comprise administration of certain NR2B subunit selective NMDA antagonists. The disorders that can be treated by the invention include hearing loss, vision loss, neurodegeneration caused by epileptic seizures, neurotoxin poisoning, Restless Leg Syndrome, multi-system atrophy, non-vascular headache, and depression.

U.S. Patent No. 5,710,168 claims the use of certain compounds having NR2B subunit selectivity for treating a disease or condition which is susceptible to treatment by blocking of

NMDA receptor sites, including traumatic brain injury, spinal cord trauma, pain, psychotic conditions, drug addiction, migraine, hypoglycemia, anxiolytic conditions, urinary incontinence, and ischemic events arising from CNS surgery, open heart surgery or any procedure during which the function of the cardiovascular system is compromised.

U.S. Patent No. 6,479,553 to AstraZeneca provides certain compounds, in particular memantine, budipine, amantidine, 5-aminocarbonyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5,10-imine, dextromethorphan and NPS 1506, and the compounds disclosed in EP 279 937 and EP 633 879, specifically (S)-1-phenyl-2-(2-pyridyl)ethanamine as potentially useful as antidepressant agents. In particular, the compounds were expected to be useful in the treatment of depression associated with neurodegenerative disorders such as Alzheimer's disease.

U.S. Patent No. 6,432,985 to Hoffman La-Roche provides certain neuroprotective substituted piperidine compounds with activity as NMDA NR2B subtype selective antagonists.

PCT Publication No. WO 06/017409 to Merck & Co. provides certain 1,3-disubstituted heteroaryl compounds are N-methyl-D-aspartate receptor antagonists useful for treating neurological condition e.g. pain, Parkinson's disease, Alzheimer's disease, anxiety, epilepsy and stroke.

PCT Publication No. WO 02/072542 to Emory University describes a class of pH-dependent NMDA receptor antagonists that exhibit pH sensitivity tested in vitro using an oocyte assay and in an experimental model of epilepsy.

While NMDA-receptor antagonists might be useful to treat a number of very challenging disorders, to date, dose-limiting side effects have prevented clinical use of NMDA receptor antagonists for these conditions. Thus, despite the potential for glutamate antagonists to treat many serious diseases, the severity of the side effects have caused many to abandon hope that a well-tolerated NMDA receptor antagonist could be developed (Hoyte L. et al (2004) *Curr. Mol. Med.* 4(2): 131-136; Muir, K. W. and Lees, K. R. (1995) *Stroke* 26:503-513; Herrling, P. L., ed. (1997) "Excitatory amino acid clinical results with antagonists" Academic Press; Parsons et al. (1998) *Drug News Perspective* II: 523 569).

There remains a need for improved neuroprotective compounds and methods for the treatment and/or prophylaxis of neuropsychiatric disorders. In particular, there is a need for compounds that have enhanced efficacy in treatment of neuropsychiatric disorders. In addition, there remains a need for effective compounds that exhibit reduced side effects upon administration. In particular there is a need for improved treatments for depression and


anxiety.

It is therefore an object of the present invention to provide new pharmaceutical compositions and methods for the treatment of neuropsychiatric disorders, and in particular for the treatment of depression and anxiety.

SUMMARY OF THE INVENTION


Compounds of Formula I, II, III and IV are provided for the treatment or prophylaxis of neuropsychiatric disorders. In particular, compounds for use in the treatment or prophylaxis of depression or anxiety in a host at risk of or suffering from the disorder are provided. In certain instances, the disorders are specifically known to result from NMDA receptor activation. Certain NMDA receptor antagonists described herein have enhanced activity in brain tissue having lower-than-normal pH due to conditions associated with a mood disorder.

In one particular embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula I or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, optionally in combination with a pharmaceutically acceptable carrier, to a host in need thereof:



Formula I


wherein the substituents are described herein. More typically, the compounds are of Formula A:



Formula A, wherein the substituents are described herein.


In a separate embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula II or a pharmaceutically acceptable salt, ester, prodrug or derivative

thereof, optionally in combination with a pharmaceutically acceptable carrier, to a host in need thereof:



Formula II

wherein the substituents are described herein. More typically, the compounds are of Formula B:



Formula B, wherein the substituents are described herein.

In certain embodiments, the compounds are used for the treatment of neuropsychiatric disorders, and in particular embodiments, neuropsychiatric mood disorders. These disorders include depression, bipolar disorders, seasonal affective disorders (SAD) and manias. In certain embodiments, the compounds are used for the treatment of depression in a host diagnosed with the disorder. In certain other embodiments, the compounds are used for treatment of a bipolar disorder in a host diagnosed with the disorder. The compounds can also be used to prevent or diminish future depressive or manic episodes. The compounds can be provided on a seasonal basis, especially in a host who has been diagnosed or is at risk of SAD or of depression.

In certain other embodiments, the compounds are useful in the treatment or prophylaxis of a neuropsychiatric disorder associated with a physiological insult. The disorder can include depression or bipolar disorder associated with an injury or with aging. The compounds may also be useful in treatment or prophylaxis of schizophrenia.

In certain embodiments, the compounds are administered to a host in need thereof. In certain other embodiments, the compounds are administered in combination or alternation with other compounds, in particular embodiments another compound useful in the treatment or prophylaxis of neuropsychiatric disorders.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is graph of the immobility time (in seconds) of CD1 mice dosed with a test compound in a forced swim test. Structures of test compounds are shown in Table 26

Figure 2 is is graph of the immobility time (in seconds) of CD1 mice dosed with a test compound in a forced swim test.

Figure 3 is a graph of the distance traveled by CD1 mice injected with a test compound in an open field activity test.

Figure 4 is a graph of the motor performance of the CD1 mice on a rotorod after dosing with test compounds.

Figure 5 is a graph of the cell toxicity of the the test compunds as assessed by percent total LDH release.

Figure 6 is a graph of the hERG binding IC_{50} (μM) for selected compounds plotted against the patch clamp IC_{50} (μM).

Figure 7 is a graph of the QT interval (msec) correlated with the log of the concentration of the test compound. Langendorff QT effects are shown for compounds NP10075, NP10239 and NP10076.

Figure 8 is a graph of PCP discrimination test data for NP10031 and NP10097.

DETAILED DESCRIPTION OF THE INVENTION

Certain compounds are provided as useful in the treatment or prophylaxis of neuropsychiatric disorders. Typically, these compounds act as NMDA antagonists. In particular, compounds of Formulas I, II, III and IV are provided for treatment of mood disorders including depression or anxiety. In certain instances, the disorders are specifically known to result from NMDA receptor activation. In certain embodiments, the compounds are allosteric NMDA inhibitors. In one embodiment, the IC_{50} value of the compound is 0.01 to 10 μM , 0.01 to 9 μM , 0.01 to 8 μM , 0.01 to 7 μM , 0.01 to 6 μM , 0.01 to 5 μM , 0.01 to 4 μM , 0.01 to 3 μM , 0.01 to 2 μM , 0.01 to 1 μM , 0.05 to 7 μM , 0.05 to 6 μM , 0.05 to 5 μM , 0.05 to 4 μM , 0.05 to 3 μM , 0.05 to 2 μM , 0.05 to 1 μM , 0.05 to 0.5 μM , 0.1 to 7 μM , 0.1 to 6 μM , 0.1 to 5 μM , 0.1 to 4 μM , 0.1 to 3 μM , 0.1 to 2 μM , 0.1 to 1 μM , 0.1 to 0.5 μM , 0.1 to 0.4 μM , 0.1 to 0.3 μM , or 0.1 to 0.2 μM .

Certain NMDA receptor antagonists described herein have enhanced activity in tissue having lower-than-normal pH. Certain studies have indicated that pH may be altered in brains of individuals suffering from certain neuropsychiatric disorder (see e.g. Karolewicz, et al.

(2004) *J. Neurochem* 91:1057-66. Xing, et al. (2002) *Schizophr Res.* 58:21-30.) A reduced brain pH can be harnessed as a switch to activate the neuroprotective agents described herein. In this way side effects are minimized in unaffected tissue since drug at these sites are less active.

In particular embodiments, the compound is pH sensitive. In specific embodiments, the compound exhibits a potency boost of at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15 or at least 20 when comparing the IC₅₀ at physiological pH versus the IC₅₀ diseased pH (i.e., (IC₅₀ at phys pH/IC₅₀ at diseased pH)).

In one embodiment, the compound has an IC₅₀ value of less than 10 μ M at a pH of about 6 to about 9. In one embodiment, the compound has an IC₅₀ value of less than 10 μ M at a pH of about 6.9. In another embodiment, the compound has an IC₅₀ value of less than 10 μ M at a pH of about 7.6. In one embodiment, the compound has an IC₅₀ value of less than 10 μ M at physiological pH. In one embodiment, the compound has an IC₅₀ value of less than 10 μ M at ischemic pH.

In one embodiment, the IC₅₀ value of the compound is 0.01 to 10 μ M, 0.01 to 9 μ M, 0.01 to 8 μ M, 0.01 to 7 μ M, 0.01 to 6 μ M, 0.01 to 5 μ M, 0.01 to 4 μ M, 0.01 to 3 μ M, 0.01 to 2 μ M, 0.01 to 1 μ M, 0.05 to 7 μ M, 0.05 to 6 μ M, 0.05 to 5 μ M, 0.05 to 4 μ M, 0.05 to 3 μ M, 0.05 to 2 μ M, 0.05 to 1 μ M, 0.05 to 0.5 μ M, 0.1 to 7 μ M, 0.1 to 6 μ M, 0.1 to 5 μ M, 0.1 to 4 μ M, 0.1 to 3 μ M, 0.1 to 2 μ M, 0.1 to 1 μ M, 0.1 to 0.5 μ M, 0.1 to 0.4 μ M, 0.1 to 0.3 μ M, or 0.1 to 0.2 μ M, and the ratio of the IC₅₀ values at pH 7.6 to pH 6.9 for the compound is greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, or 100.

In one embodiment, the IC₅₀ value of the compound is 0.01 to 10 μ M, 0.01 to 9 μ M, 0.01 to 8 μ M, 0.01 to 7 μ M, 0.01 to 6 μ M, 0.01 to 5 μ M, 0.01 to 4 μ M, 0.01 to 3 μ M, 0.01 to 2 μ M, 0.01 to 1 μ M, 0.05 to 7 μ M, 0.05 to 6 μ M, 0.05 to 5 μ M, 0.05 to 4 μ M, 0.05 to 3 μ M, 0.05 to 2 μ M, 0.05 to 1 μ M, 0.05 to 0.5 μ M, 0.1 to 7 μ M, 0.1 to 6 μ M, 0.1 to 5 μ M, 0.1 to 4 μ M, 0.1 to 3 μ M, 0.1 to 2 μ M, 0.1 to 1 μ M, 0.1 to 0.5 μ M, 0.1 to 0.4 μ M, 0.1 to 0.3 μ M, or 0.1 to 0.2 μ M, and the ratio of the IC₅₀ values at pH 7.6 to pH 6.9 for the compound is between 1 and 100, 2 and 100, 3 and 100, 4 and 100, 5 and 100, 6 and 100, 7 and 100, 8 and 100, 9 and 100, 10 and 100, 15 and 100, 20 and 100, 25 and 100, 30 and 100, 40 and 100, 50 and 100, 60 and 100, 70 and 100, 80 and 100, or 90 and 100.

Definitions

Whenever a term in the specification is identified as a range (i.e. C₁₋₄ alkyl), the range independently refers to each element of the range. As a non-limiting example, C₁₋₄ alkyl

means, independently, C₁, C₂, C₃ or C₄ alkyl. Similarly, when one or more substituents are referred to as being “independently selected from” a group, this means that each substituent can be any element of that group, and any combination of these groups can be separated from the group. For example, if R¹ and R² can be independently selected from X, Y and Z, this separately includes the groups R¹ is X and R² is X; R¹ is X and R² is Y; R¹ is X and R² is Z; R¹ is Y and R² is X; R¹ is Y and R² is Y; R¹ is Y and R² is Z; R¹ is Z and R² is X; R¹ is Z and R² is Y; and R¹ is Z and R² is Z.

The term “alkyl” is used herein, unless otherwise specified, refers to a substituted or unsubstituted, saturated, straight, branched, or cyclic (also identified as cycloalkyl), primary, secondary, or tertiary hydrocarbon, including but not limited to those of C₁ to C₆. Illustrative examples of alkyl groups are methyl, ethyl, propyl, *isopropyl*, cyclopropyl, butyl, *secbutyl*, *isobutyl*, *tertbutyl*, cyclobutyl, 1-methylbutyl, 1,1-dimethylpropyl, pentyl, cyclopentyl, *isopentyl*, *neopentyl*, cyclopentyl, hexyl, *isohexyl*, and cyclohexyl. Unless otherwise specified, the alkyl group can be unsubstituted or substituted with one or more moieties selected from the group consisting of alkyl, halo, haloalkyl, hydroxyl, carboxyl, acyl, acyloxy, amino, amido, carboxyl derivatives, alkylamino, dialkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, thio, sulfonyl, ester, carboxylic acid, amide, phosphonyl, phosphinyl, thioether, oxime, or any other viable functional group that does not inhibit the pharmacological activity of this compound, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene, *et al.*, Protective Groups in Organic Synthesis, John Wiley and Sons, Second Edition, 1991. In certain embodiments, alkyl may be optionally substituted by one or more fluoro, chloro, bromo, iodo, hydroxy, heterocyclic, heteroaryl, carboxy, alkoxy, nitro, NH₂, N(alkyl)₂, NH(alkyl), alkoxycarbonyl, -N(H or alkyl)C(O)(H or alkyl), -N(H or alkyl)C(O)N(H or alkyl)₂, -N(H or alkyl)C(O)O(H or alkyl), -OC(O)N(H or alkyl)₂, -S(O)_n-(H or alkyl), -C(O)-N(H or alkyl)₂, cyano, alkenyl, cycloalkyl, acyl, hydroxyalkyl, heterocyclic, heteroaryl, aryl, aminoalkyl, oxo, carboxyalkyl, -C(O)-NH₂, -C(O)-N(H)O(H or alkyl), -S(O)₂-NH₂, -S(O)_n-N(H or alkyl)₂ and/or -S(O)₂-N(H or alkyl)₂.

The term “halo” or “halogen,” refers to chloro, bromo, iodo, or fluoro.

The term “heteroaryl” or “heteroaromatic,” refers to an aromatic that includes at least one sulfur, oxygen, nitrogen or phosphorus in the aromatic ring. The term “heterocyclic” refers to a non-aromatic cyclic group wherein there is at least one heteroatom, such as oxygen, sulfur, nitrogen, or phosphorus in the ring. Nonlimiting examples of heteroaryl and heterocyclic groups include furyl, furanyl, pyridyl, pyrimidyl, thienyl, isothiazolyl,

imidazolyl, tetrazolyl, pyrazinyl, benzofuranyl, benzothiophenyl, quinolyl, isoquinolyl, benzothienyl, isobenzofuryl, pyrazolyl, indolyl, isoindolyl, benzimidazolyl, purinyl, carbazolyl, oxazolyl, thiazolyl, isothiazolyl, 1,2,4-thiadiazolyl, isooxazolyl, pyrrolyl, quinazolinyl, cinnolinyl, phthalazinyl, xanthinyl, hypoxanthinyl, thiophene, furan, pyrrole, isopyrrole, pyrazole, imidazole, 1,2,3-triazole, 1,2,4-triazole, oxazole, isoxazole, thiazole, isothiazole, pyrimidine or pyridazine, pteridinyl, aziridines, thiazole, isothiazole, oxadiazole, thiazine, pyridine, pyrazine, piperazine, piperidine, pyrrolidine, oxaziranes, phenazine, phenothiazine, morpholinyl, pyrazolyl, pyridazinyl, pyrazinyl, quinoxalinyl, xanthinyl, hypoxanthinyl, pteridinyl, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, pyrazolopyrimidinyl, adenine, N⁶-alkylpurines, N⁶-benzylpurine, N⁶-halopurine, N⁶-vinypurine, N⁶-acetylenic purine, N⁶-acyl purine, N⁶-hydroxyalkyl purine, N⁶-thioalkyl purine, thymine, cytosine, 6-azapyrimidine, 2-mercaptopyrimidine, uracil, N⁵-alkylpyrimidines, N⁵-benzylpyrimidines, N⁵-halopyrimidines, N⁵-vinylpyrimidine, N⁵-acetylenic pyrimidine, N⁵-acyl pyrimidine, N⁵-hydroxyalkyl purine, and N⁶-thioalkyl purine, and isoxazolyl. The heteroaromatic or heterocyclic group can be optionally substituted with one or more substituent selected from halogen, haloalkyl, alkyl, alkoxy, hydroxy, carboxyl derivatives, amido, amino, alkylamino, dialkylamino. The heteroaromatic can be partially or totally hydrogenated as desired. Nonlimiting examples include dihydropyridine and tetrahydrobenzimidazole. In some embodiment, the heteroaryl may be optionally substituted by one or more fluoro, chloro, bromo, iodo, hydroxy, heterocyclic, heteroaryl, carboxy, alkoxy, nitro, NH₂, N(alkyl)₂, NH(alkyl), alkoxycarbonyl, -N(H or alkyl)C(O)(H or alkyl), -N(H or alkyl)C(O)N(H or alkyl)₂, -N(H or alkyl)C(O)O(H or alkyl), -OC(O)N(H or alkyl)₂, -S(O)_n-(H or alkyl), -C(O)-N(H or alkyl)₂, cyano, alkenyl, cycloalkyl, acyl, hydroxyalkyl, heterocyclic, heteroaryl, aryl, aminoalkyl, oxo, carboxyalkyl, -C(O)-NH₂, -C(O)-N(H)O(H or alkyl), -S(O)₂-NH₂, -S(O)_n-N(H or alkyl)₂ and/or -S(O)₂-N(H or alkyl)₂. Functional oxygen and nitrogen groups on the heteroaryl group can be protected as necessary or desired. Suitable protecting groups are well known to those skilled in the art, and include trimethylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, and *t*-butyldiphenylsilyl, trityl or substituted trityl, alkyl groups, acyl groups such as acetyl and propionyl, methanesulfonyl, and *p*-tolylsulfonyl.

The term “aryl,” unless otherwise specified, refers to a carbon based aromatic ring, including phenyl, biphenyl, or naphthyl. The aryl group can be optionally substituted with one or more moieties selected from the group consisting of hydroxyl, acyl, amino, halo, alkylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate,

or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in *Greene, et al.*, "Protective Groups in Organic Synthesis," John Wiley and Sons, Second Edition, 1991. In certain embodiments, the aryl group is optionally substituted by one or more fluoro, chloro, bromo, iodo, hydroxy, heterocyclic, heteroaryl, carboxy, alkoxy, nitro, NH_2 , $\text{N}(\text{alkyl})_2$, $\text{NH}(\text{alkyl})$, alkoxycarbonyl, $-\text{N}(\text{H or alkyl})\text{C}(\text{O})(\text{H or alkyl})$, $-\text{N}(\text{H or alkyl})\text{C}(\text{O})\text{N}(\text{H or alkyl})_2$, $-\text{N}(\text{H or alkyl})\text{C}(\text{O})\text{O}(\text{H or alkyl})$, $-\text{OC}(\text{O})\text{N}(\text{H or alkyl})_2$, $-\text{S}(\text{O})_n-\text{H or alkyl}$, $-\text{C}(\text{O})-\text{N}(\text{H or alkyl})_2$, cyano, alkenyl, cycloalkyl, acyl, hydroxyalkyl, heterocyclic, heteroaryl, aryl, aminoalkyl, oxo, carboxyalkyl, $-\text{C}(\text{O})-\text{NH}_2$, $-\text{C}(\text{O})-\text{N}(\text{H})\text{O}(\text{H or alkyl})$, $-\text{S}(\text{O})_2-\text{NH}_2$, $-\text{S}(\text{O})_n-\text{N}(\text{H or alkyl})_2$ and/or $-\text{S}(\text{O})_2-\text{N}(\text{H or alkyl})_2$.

The term "aralkyl," unless otherwise specified, refers to an aryl group as defined above linked to the molecule through an alkyl group as defined above.

The term "alkaryl," unless otherwise specified, refers to an alkyl group as defined above linked to the molecule through an aryl group as defined above. Other groups, such as acyloxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkylaminoalkyl, alkylthioalkyl, amidoalkyl, aminoalkyl, carboxyalkyl, dialkylaminoalkyl, haloalkyl, heteroaralkyl, heterocyclicalkyl, hydroxyalkyl, sulfonamidoalkyl, sulfonylalkyl and thioalkyl are named in a similar manner.

The term "alkoxy," unless otherwise specified, refers to a moiety of the structure $-\text{O}-\text{alkyl}$, wherein alkyl is as defined above.

The term "acyl," refers to a group of the formula $\text{C}(\text{O})\text{R}'$ or "alkyl-oxy", wherein R' is an alkyl, aryl, alkaryl or aralkyl group, or substituted alkyl, aryl, aralkyl or alkaryl.

The term "alkenyl" The term "alkenyl" means a monovalent, unbranched or branched hydrocarbon chain having one or more double bonds therein. The double bond of an alkenyl group can be unconjugated or conjugated to another unsaturated group. Suitable alkenyl groups include, but are not limited to (C_2-C_8) alkenyl groups, such as vinyl, allyl, butenyl, pentenyl, hexenyl, butadienyl, pentadienyl, hexadienyl, 2-ethylhexenyl, 2-propyl-2-butenyl, 4-(2-methyl-3-butene)-pentenyl. An alkenyl group can be unsubstituted or substituted with one or two suitable substituents.

The term "carbonyl" refers to a functional group composed of a carbon atom double-bonded to an oxygen atom : $-\text{C}=\text{O}$. Similarly, $\text{C}(\text{O})$ or $\text{C}(=\text{O})$ refers to a carbonyl group.

The term "amino" refers to $-\text{NH}_2$, $-\text{NH}(\text{alkyl})$ or $-\text{N}(\text{alkyl})_2$.

The term "thio" indicates the presence of a sulfur group. The prefix thio- denotes that there is at least one extra sulfur atom added to the chemical. The prefix 'thio-' can also be placed before the name of a compound to mean that an oxygen atom in the compound has been

replaced by a sulfur atom. Although typically the term "thiol" is used to indicate the presence of -SH, in instances in which the sulfur atom would be have improper valance a radical if the hydrogen is improperly designated, the terms 'thio' and 'thiol' are used interchangeably, unless otherwise indicated.

The term "amido" indicates a group (H or alkyl)-C(O)-NH-.

The term "carboxy" designates the terminal group -C(O)OH.

The term "sulfonyl" indicates an organic radical of the general formula (H or alkyl)-S(=O)₂-(H or alkyl'), where there are two double bonds between the sulfur and oxygen.

The term "pharmaceutically acceptable salt" refers to salts or complexes that retain the desired biological activity of the compounds of the present invention and exhibit minimal undesired toxicological effects. Nonlimiting examples of such salts are (a) acid addition salts formed with inorganic acids (for example, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid, and the like), and salts formed with organic acids such as acetic acid, oxalic acid, tartaric acid, succinic acid, malic acid, ascorbic acid, benzoic acid, tannic acid, pamoic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, naphthalenedisulfonic acid, and polygalcturonic acid; (b) base addition salts formed with metal cations such as zinc, calcium, bismuth, barium, magnesium, aluminum, copper, cobalt, nickel, cadmium, sodium, potassium, and the like, or with a cation formed from ammonia, N,N-dibenzylethylenediamine, D-glucosamine, tetraethylammonium, or ethylenediamine; or (c) combinations of (a) and (b); e.g., a zinc tannate salt or the like. Also included in this definition are pharmaceutically acceptable quaternary salts known by those skilled in the art, which specifically include the quaternary ammonium salt of the formula -NR⁺A⁻, wherein R is H or alkyl and A is a counterion, including chloride, bromide, iodide, -O-alkyl, toluenesulfonate, methylsulfonate, sulfonate, phosphate, or carboxylate (such as benzoate, succinate, acetate, glycolate, maleate, malate, citrate, tartrate, ascorbate, benzoate, cinnamoate, mandeloate, benzyloate, and diphenylacetate).


The term "protected" as used herein and unless otherwise defined refers to a group that is added to an oxygen, nitrogen, or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis.

It should be understood that the various possible stereoisomers of the groups mentioned above and herein are within the meaning of the individual terms and examples, unless otherwise specified. As an illustrative example, "1-methyl-butyl" exists in both (R) and the (S) form, thus, both (R)-1-methyl-butyl and (S)-1-methyl-butyl is covered by the term

"1-methyl-butyl", unless otherwise specified.

Compounds

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety, are provided comprising administering a compound of Formula I or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



FORMULA I

wherein:

each L is independently C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, alkaryl, hydroxy, -O-alkyl, -O-aryl, -SH, -S-alkyl, -S-aryl, fluoro, chloro, bromo, iodo, nitro, or cyano; or two L groups may be taken together with Ar¹ to form: a dioxolane ring or a cyclobutane ring;

k = 0, 1, 2, 3, 4 or 5;

each Ar¹ and Ar² is independently aryl or heteroaryl;

W is a bond, C₁-C₄ alkyl, or C₂-C₄ alkenyl;

X is a bond, NR¹ or O;

each R¹ and R² is independently H, C₁-C₆ alkyl, C₂-C₆ alkenyl or C₆-C₁₂ aralkyl; or


R¹ and R² can be taken together to form a 5-8 membered ring;

each R³ and R⁴ is independently H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR³R⁴ is C=O;

n and p are each independently 1, 2, 3 or 4;


each R⁵ and R⁶ is independently H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR⁵R⁶ is C=O or C=CH₂;

or wherein -NR²- (CR⁵R⁶)_p- can be




Y is a bond, O, S, SO, SO₂, CH₂, NH, N(C₁-C₆ alkyl), or NHC(=O);


Z is OH, NR^6R^7 , $\text{NR}^8\text{SO}_2(\text{C}_1\text{-C}_6 \text{ alkyl})$, $\text{NR}^8\text{C}(\text{O})\text{NR}^6\text{R}^7$, $\text{NR}^8\text{C}(\text{S})\text{NR}^6\text{R}^7$, $\text{NR}^8\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_6 \text{ alkyl})$, NR^8 -dihydrothiazole, or NR^8 -dihydroimidazole; wherein each R^6 , R^7 and R^8 is independently H, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_6\text{-C}_{12}$ aralkyl; or



In one embodiment, when Y is $\text{NHC}(=\text{O})$, Z is not OH or $\text{NR}^8\text{SO}_2(\text{C}_1\text{-C}_6 \text{ alkyl})$. In one subembodiment, when R^1 and R^2 are taken together to form a 5-8 membered ring so that



subembodiment, when R^1 and R^2 are taken together to form a 5-8 membered ring so that -



In one embodiment, X is NR^1 . In another embodiment, X is O. In another embodiment, X is a bond. In a particular subembodiment, X is a bond, n is 1, R^3 and R^4 are both H, and W is C_2 alkenyl.

In particular subembodiment, Ar^1 is phenyl, pyridyl, pyrimidinyl, thiophenyl, imidazolyl, furanyl, indolyl, benzothiophenyl, benzofuranyl, or benzoimidazolyl.

In another particular subembodiment, L is $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}(=\text{O})-(\text{C}_1\text{-C}_4)\text{-alkyl}$, $\text{C}_1\text{-C}_6$ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano. In a further subembodiment, L is methyl, trifluoromethyl, methoxy, nitro, fluoro, chloro or hydroxy. In one further subembodiment, there are one, two or three L groups substituting Ar^1 . In one subembodiment, Ar^1 is substituted with one fluoro group. In one subembodiment, Ar^1 is substituted with two fluoro groups. In one subembodiment, Ar^1 is substituted with one fluoro

group and one chloro group. In one subembodiment, Ar^1 is substituted with one chloro group. In one subembodiment, Ar^1 is substituted with two chloro groups. In one subembodiment, Ar^1 is substituted with one methyl group. In one subembodiment, Ar^1 is substituted with one trifluoromethyl group.


In one subembodiment, Ar^1 is phenyl. In one subembodiment, Ar^1 is phenyl and is substituted with an L group at the 2, 3, or 4 position. In another subembodiment, Ar^1 is phenyl and is substituted with L groups at the 2 and 4 positions. In another subembodiment, Ar^1 is phenyl and is substituted with L groups at the 3 and 4 positions.










In one subembodiment, Ar^1 is pyridyl. In another subembodiment, Ar^1 is 2-pyridyl, 3-pyridyl, or 4-pyridyl.

In one embodiment, Ar^1 is a bicyclic group wherein the W group is attached to the heterocyclic ring.


In one embodiment, W is a bond. In another embodiment, W is CH_2 . In another embodiment, W is $\text{C}_2\text{-C}_4$ alkenyl.

In one embodiment, each R^1 and R^2 is independently H or $\text{C}_1\text{-C}_4$ alkyl, for example, methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl or *tert*-butyl. In one embodiment, R^1 and R^2 are both H. In one embodiment, R^1 and R^2 are both $\text{C}_1\text{-C}_4$ alkyl, for example *n*-butyl. In another embodiment, R^1 and R^2 can be taken together to form a 5-8 membered ring so that


$-\text{NR}^1-(\text{CR}^3\text{R}^4)_n-\text{NR}^2-$ is . In one embodiment, n is 2. In one embodiment, n is 3. In another embodiment, R^1 and R^2 are each CH_2 . In a subembodiment, CR^3R^4 is CH_2 and n is 2. In a subembodiment, CR^3R^4 is CH_2 and n is 3. In a subembodiment, CR^3R^4 is C=O and n is 1.


In one embodiment,  is , ,  or . In one embodiment,  is . In another embodiment,  is .

In one embodiment, each R^5 and R^6 is independently H, C₁-C₄ alkyl, C₁-C₄ alkoxy, C(=O)-(C₁-C₄)-alkyl, C₁-C₄ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano. In one embodiment, CR^5R^6 is C=O or C=CH₂. In one embodiment, p is 2, 3, or 4. In another embodiment, p is 3. In one embodiment, R^5 and R^6 are H. In another embodiment, one of R^5 and R^6 is hydroxy. In another embodiment, CR^5R^6 is C=CH₂. In another embodiment, CR^5R^6 is C=O. In one embodiment, $(CR^5R^6)_p$ is selected from the group consisting of




Compounds of Formula I can include compounds wherein when p is greater than 1, each (CR^5R^6) can be independently selected, for example, in one embodiment p is 2 and one (CR^5R^6) is C=O and the other (CR^5R^6) is CH₂. In one embodiment, R^5 is not fluoro. In another embodiment, R^6 is not fluoro.

In one embodiment, $-NR^2-(CR^5R^6)_p-$ is . In a particular

subembodiment, the compound is . In another

particular subembodiment, the compound is



In one embodiment, Y is a bond, O or CH₂. In one embodiment, Y is O. In another embodiment, Y is CH₂. In one embodiment, Y is not NH. In another embodiment, Y is not NHC(=O).

In one embodiment, Ar² is aryl. In one embodiment, Ar² is aryl, but not phenyl or heteroaryl. In one embodiment Ar² is phenyl. In one subembodiment, Ar² is phenyl and is substituted with a Z group at the 4 position. In one embodiment, Ar² is not heteroaryl. In one embodiment, Ar² is aryl, but not phenyl or heteroaryl.

In one embodiment, Z is OH, NR⁶R⁷, NR⁸SO₂(C₁-C₆ alkyl), NR⁸C(O)NR⁶R⁷, NR⁸C(S)NR⁶R⁷, NR⁸C(O)O(C₁-C₆ alkyl), NR⁸-dihydrothiazole, or NR⁸-dihydroimidazole.

In one embodiment, Ar²—Z is

, or


subembodiment, Ar²—Z is

is

subembodiment, R⁹ and R¹⁰ are each H.

In one embodiment, Z is NR⁸C(O)NR⁶R⁷, for example NHC(O)NH₂ or NHC(O)N(CH₃)₂.

In another embodiment, Z and Ar² are taken together and selected from the group



In one embodiment, the compound is a compound of Formula I, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein:

L is C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, alkaryl, hydroxy, -O-alkyl, -O-aryl, -SH, -S-alkyl, -S-aryl, fluoro, chloro, bromo, iodo, nitro, or cyano; or two L groups may be taken together with Ar¹ to form a dioxolane ring or a cyclobutane ring;

k = 0, 1, 2, 3, 4 or 5;

Ar¹ is phenyl, pyridyl, pyrimidinyl, thiophenyl, imidazolyl, furanyl, indolyl, benzothiophenyl, benzofuranyl, benzoimidazolyl;

Ar² is phenyl;

W is a bond, C₁-C₄ alkyl, or C₂-C₄ alkenyl;

each R¹ and R² is independently H, C₁-C₄ alkyl; or

R¹ and R² can be taken together to form a 5-8 membered ring;


each R³ and R⁴ is independently H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR³R⁴ is C=O;

n = 1, 2, 3 or 4;

each R⁵ and R⁶ is independently H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR⁵R⁶ is C=O, C=CH₂;

Y is a bond, O, S, SO, SO₂, CH₂, NH, N(C₁-C₆ alkyl), NHC(=O);

Z is OH, NH₂, NHSO₂(C₁-C₄ alkyl), NHC(O)NR⁶R⁷, NR⁸C(S)NR⁶R⁷, NHC(O)O(C₁-C₄ alkyl), NH-dihydrothiazole, or NH-dihydroimidazole; wherein each R⁶ and R⁷ is independently H, C₁-C₆ alkyl; or



In one embodiment, the compound is a compound of Formula I, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein:

L is C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, alkaryl, hydroxy, -O-alkyl, -O-aryl, -SH, -S-alkyl, -S-aryl, fluoro, chloro, bromo, iodo, nitro, or cyano; or two L groups may be taken together with Ar¹ to form a dioxolane ring or a cyclobutane ring;

k = 0, 1, 2, 3, 4 or 5;

Ar¹ is phenyl, pyridyl, pyrimidinyl, thiophenyl, imidazolyl, furanyl, indolyl, benzothiophenyl, benzofuranyl, benzoimidazolyl;

Ar² is phenyl;

W is a bond, C₁-C₄ alkyl, or C₂-C₄ alkenyl;

each R¹ and R² is independently H, C₁-C₄ alkyl; or

R¹ and R² can be taken together to form a 5-8 membered ring;


each R³ and R⁴ is independently H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR³R⁴ is C=O;

n = 1, 2, 3 or 4;

each R⁵ and R⁶ is independently H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR⁵R⁶ is C=O, C=CH₂;

Y is a bond, O, S, SO, SO₂, CH₂, NH, N(C₁-C₆ alkyl), NHC(=O);

Z is OH, NH₂, NHSO₂(C₁-C₄ alkyl), NHC(O)NR⁶R⁷, NR⁸C(S)NR⁶R⁷, NHC(O)O(C₁-C₄ alkyl), NH-dihydrothiazole, or NH-dihydroimidazole; wherein each R⁶ and R⁷ is independently H, C₁-C₆ alkyl; or



In one embodiment, the compound is a compound of Formula I, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein:

L is C₁-C₄ alkyl, C₁-C₄ alkoxy, C(=O)-(C₁-C₄)-alkyl, C₁-C₄ haloalkyl, alkaryl, hydroxy, -O-alkyl, -O-aryl, -SH, -S-alkyl, -S-aryl, fluoro, chloro, bromo, iodo, or nitro; or

two L groups may be taken together to form a dioxolane ring with Ar¹;

k = 0, 1, 2, 3, 4 or 5;

Ar¹ is phenyl or pyridyl;

Ar² is phenyl;

W is a bond or C₁-C₄ alkyl;

X is NR¹;

each R¹ and R² is independently H or C₁-C₄ alkyl; or

R¹ and R² can be taken together to form a 5-8 membered ring;


each R³ and R⁴ is independently H or C₁-C₄ alkyl; or CR³R⁴ is C=O;

n = 2 or 3;

each R⁵ and R⁶ is independently H, C₁-C₄ alkyl or OH; or CR⁴R⁵ is C=O or C=CH₂;

Y is O or CH₂;

Z is OH, NH₂, NHSO₂(C₁-C₄ alkyl), NHC(O)NR⁶R⁷, NR⁸C(S)NR⁶R⁷, NHC(O)O(C₁-C₄ alkyl), NH-dihydrothiazole, or NH-dihydroimidazole; wherein each R⁶ and R⁷ is independently H or C₁-C₄ alkyl; or



R⁹ is H or C₁-C₄ alkyl.

In one embodiment, the compound is a compound of Formula I, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein:

L is C₁-C₄ alkyl, C₁-C₄ alkoxy, C(=O)-(C₁-C₄)-alkyl, C₁-C₄ haloalkyl, alkaryl, hydroxy, -O-alkyl, -O-aryl, -SH, -S-alkyl, -S-aryl, fluoro, chloro, bromo, iodo, or nitro; or

two L groups may be taken together to form a dioxolane ring with Ar¹;

k = 0, 1, 2, 3, 4 or 5;

Ar¹ is phenyl or pyridyl;

Ar² is phenyl;

W is a bond or C₁-C₄ alkyl;

X is O;

R² is H or C₁-C₄ alkyl;


each R³ and R⁴ is independently H or C₁-C₄ alkyl; or CR³R⁴ is C=O;

n = 2 or 3;

each R⁵ and R⁶ is independently H, C₁-C₄ alkyl or OH; or CR⁴R⁵ is C=O or C=CH₂;

Y is O or CH₂;

Z is OH, NH₂, NHSO₂(C₁-C₄ alkyl), NHC(O)NR⁶R⁷, NHC(O)O(C₁-C₄ alkyl), NH-dihydrothiazole, or NH-dihydroimidazole; wherein each R⁶ and R⁷ is independently H or C₁-C₄ alkyl; or



R⁹ is H or C₁-C₄ alkyl.

In one embodiment, the compound is a compound of Formula I, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein:

L is C₁-C₄ alkyl, C₁-C₄ alkoxy, C(=O)-(C₁-C₄)-alkyl, C₁-C₄ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, or nitro; or

two L groups may be taken together to form a dioxolane ring with Ar¹;

k = 0, 1, 2, 3, 4 or 5;

Ar¹ is phenyl or pyridyl;

Ar² is phenyl;

W is C₂-C₄ alkenyl;

X is a bond;

R² is H or C₁-C₄ alkyl;


each R³ and R⁴ is independently H or C₁-C₄ alkyl; or CR³R⁴ is C=O;

n = 1, 2 or 3;

each R^5 and R^6 is independently H, C_1 - C_4 alkyl or OH; or CR^4R^5 is $C=O$ or $C=CH_2$;

Y is O or CH_2 ;






Z is OH, NH_2 , $NHSO_2(C_1-C_4 \text{ alkyl})$, $NHC(O)NR^6R^7$, $NR^8C(S)NR^6R^7$, $NHC(O)O(C_1-C_4 \text{ alkyl})$, NH-dihydrothiazole, or NH-dihydroimidazole; wherein each R^6 and R^7 is independently H or C_1 - C_4 alkyl; or



R^9 is H or C_1 - C_4 alkyl.

In one embodiment, the compound is selected from the compounds in Table 1.

Table 1.




Compound	NAME
	N-(4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(4-Chloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-(4-{3-[4-phenyl-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(4-Hydroxy-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(2-Pyridyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide

	N-(4-{3-[4-(4-Pyridyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-{4-[2-(<i>S</i>)-Hydroxy-3-(2-phenylamino-ethylamino)-propoxy]-phenyl}-methanesulfonamide
	N-{4-[2-(<i>S</i>)-Hydroxy-3-(2-(3,4-difluorophenyl)amino-ethylamino)-propoxy]-phenyl}-methanesulfonamide
	N-(4-{3-[3-(3,4-Dichloro-phenyl)-allylamino]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-[4-(3-{Butyl-[3-(3,4-dichloro-phenyl)-allyl]-amino}-2-(<i>S</i>)-hydroxy-propoxy)-phenyl]-methanesulfonamide
	N-(4-{3-[3-(3,4-Difluoro-phenyl)-allylamino]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide

In one embodiment, the compound is selected from the compounds in Table 2.




Table 2.









Compound	NAME
	6-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one









	6-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one
	6-{3-[4-(4-Methyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one
	6-{3-[2-(4-Chloro-phenylamino)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one





In one embodiment, the compound is selected from the compounds in Table 3.

Table 3.

Compound	NAME
	4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-(<i>R</i>)-hydroxy-propoxy}-phenol



	4-{3-[4-(4-Fluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(3,4-Dimethyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(4-Methyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(4-Cyano-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(4-Bromo-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(4-Hydroxy-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(4-Methoxy-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(4-Trifluoromethyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol








	4-{3-[4-(4-Biphenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(2-Fluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(2-Chloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(2-Chloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(2-Cyano-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-Phenyl-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(3-Fluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol








	4-{3-[4-(3-Chloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(3-Methyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-{3-[4-(3-Trifluoromethyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	








In one embodiment, the compound is selected from the compounds in Table 4.







Table 4.








Compound	NAME
	(4-{2-[4-(3,4-Difluoro-phenyl)-piperazin-1-ylmethyl]-allyloxy}-phenyl)-urea
	(4-{3-[4-(4-Fluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-urea







	(4-{3-[4-(4-Chloro-phenyl)-piperazin-1-yl]-2-hydroxy-propoxy}-phenyl)-urea
	1-Ethyl-3-(4-{3-[4-(4-fluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-urea
	(4-{3-[4-(4-Fluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-carbamic acid methyl ester
	(S)-1-(4-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(R)-1-(4-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(4-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea







	(S)-1-(4-(3-(4-(3,4-dimethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4- <i>p</i> -tolylpiperazin-1-yl)propoxy)phenyl)urea
	(S)-1-(4-(3-(4-(4-cyanophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(4-bromophenyl)piperazin-1-yl)-2-(<i>S</i>)-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4-(4-hydroxyphenyl)piperazin-1-yl)propoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4-(4-methoxyphenyl)piperazin-1-yl)propoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4-(4-(trifluoromethyl)phenyl)piperazin-1-yl)propoxy)phenyl)urea





	(S)-1-(4-(3-(4-(biphenyl-4-yl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(2,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(2-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(2-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4-o-tolylpiperazin-1-yl)propoxy)phenyl)urea
	(S)-1-(4-(3-(4-(2-cyanophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4-phenylpiperazin-1-yl)propoxy)phenyl)urea

	(S)-1-(4-(3-(4-(3-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(3-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4- <i>m</i> -tolylpiperazin-1-yl)propoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)propoxy)phenyl)urea
	(S)-1-(4-(3-(4-(4-ethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(4-isopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea

	(S)-1-(4-(3-(4-(4-cyclopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(4-propylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(4-butylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(3-(4-(4-isobutylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(S)-1-(4-(2-hydroxy-3-(4-(4-(prop-1-ynyl)phenyl)piperazin-1-yl)propoxy)phenyl)urea
	(S)-1-(4-(3-(4-(2-naphthyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	1-(4-((S)-3-((R)-4-(4-chlorophenyl)-2-methylpiperazin-1-yl)-2-hydroxypropoxy)phenyl)urea




	1-(4-((<i>S</i>)-3-((<i>S</i>)-4-(4-chlorophenyl)-2-methylpiperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	1-(4-((<i>S</i>)-3-4-(4-chlorophenyl)-cis-2,6-dimethylpiperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	1-(4-((<i>S</i>)-3-(cis-2,6-dimethyl-4-p-tolylpiperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(4-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(4-ethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea

	(<i>R</i>)-1-(4-(3-(4-(4-isopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(4-cyclopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(4-propylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(4-butylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(4-isobutylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(2-hydroxy-3-(4-(4-(prop-1-ynyl)phenyl)piperazin-1-yl)propoxy)phenyl)urea









	(<i>R</i>)-1-(4-(3-(4-(2-naphthyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(4-methylphenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-phenyl-piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea
	(<i>R</i>)-1-(4-(3-(4-(4-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)phenyl)urea

In one embodiment, the compound is selected from the compounds in Table 5.

Table 5.

Compound	Name
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-2-(4-methanesulfonylamino-phenoxy)-acetamide
	N-(4-{2-[2-(3,4-Dichloro-phenylamino)-ethylamino]-ethoxy}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-3-oxo-propyl}-phenyl)-methanesulfonamide








	N-(4-{3-[4-(3,4-DiFluoro-phenyl)-piperazin-1-yl]-3-oxo-propyl}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-propyl}-phenyl)-methanesulfonamide
	N-(4-{2-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-methanesulfonamide
	6-{2-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-oxo-ethoxy}-3H-benzooxazol-2-one
	6-{2-[4-(3,4-DiFluoro-phenyl)-piperazin-1-yl]-2-oxo-ethoxy}-3H-benzooxazol-2-one
	6-{2-[4-(4-Chloro-phenyl)-piperazin-1-yl]-2-oxo-ethoxy}-3H-benzooxazol-2-one
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-2-(2-oxo-2,3-dihydro-benzooxazol-6-yloxy)-acetamide
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-2-(4-hydroxy-phenoxy)-acetamide
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-3-(4-hydroxy-phenyl)-propionamide
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-2-(3-fluoro-4-hydroxy-phenoxy)-acetamide








	N-[3-(3,4-Dichloro-phenyl)-allyl]-2-(4-methanesulfonylamino-phenoxy)-acetamide
	N-[2-(3,4-Dichloro-phenoxy)-ethyl]-2-(4-methanesulfonylamino-phenoxy)-acetamide
	N-[2-(3,4-Dichloro-phenoxy)-ethyl]-2-(4-hydroxy-phenoxy)-acetamide
	N-[2-(3,4-Dichloro-phenoxy)-ethyl]-2-(4-ureido-phenoxy)-acetamide
	(S)-1-(4-chlorophenyl)-3-(2-hydroxy-3-(4-hydroxyphenoxy)propyl)imidazolidin-2-one
	(S)-N-(4-(3-(3-(3,4-dichlorophenyl)-2-oxoimidazolidin-1-yl)-2-hydroxypropoxy)phenyl)methanesulfonamide
	(S)-3-(2-(4-chlorophenylamino)ethyl)-5-((4-hydroxyphenoxy)methyl)oxazolidin-2-one
	(S)-N-(4-((3-(2-(3,4-dichlorophenylamino)ethyl)-2-oxooxazolidin-5-yl)methoxy)phenyl)methanesulfonamide








In one embodiment, the compound is selected from Table 6.








Table 6.








Compound	NAME
----------	------








	(S)-5-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(R)-5-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(3-(4-(4-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(3-(4-(3,4-dimethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-p-tyl)piperazin-1-yl)propoxy)indolin-2-one
	(S)-4-(4-(2-hydroxy-3-(2-oxoindolin-5-yloxy)propyl)piperazin-1-yl)benzonitrile






	(S)-5-(3-(4-(4-bromophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-(4-hydroxyphenyl)piperazin-1-yl)propoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-(4-methoxyphenyl)piperazin-1-yl)propoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-(4-(trifluoromethyl)phenyl)piperazin-1-yl)propoxy)indolin-2-one
	(S)-5-(3-(4-(biphenyl-4-yl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(3-(4-(2,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(3-(4-(2-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one

	(S)-5-(3-(4-(2-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4- <i>o</i> -tolylpiperazin-1-yl)propoxy)indolin-2-one
	(S)-2-(4-(2-hydroxy-3-(2-oxoindolin-5-yloxy)propyl)piperazin-1-yl)benzonitrile
	(S)-5-(2-hydroxy-3-(4-phenylpiperazin-1-yl)propoxy)indolin-2-one
	(S)-5-(3-(4-(3-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(3-(4-(3-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4- <i>m</i> -tolypiperazin-1-yl)propoxy)indolin-2-one

	(S)-5-(2-hydroxy-3-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)propoxy)indolin-2-one
	(S)-5-(3-(4-(4-ethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-(4-isopropylphenyl)piperazin-1-yl)propoxy)indolin-2-one
	(S)-5-(3-(4-(4-cyclopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-(4-propylphenyl)piperazin-1-yl)propoxy)indolin-2-one
	(S)-5-(3-(4-(4-butylphenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-(4-isobutylphenyl)piperazin-1-yl)propoxy)indolin-2-one


	(S)-5-(2-hydroxy-3-(4-(4-(prop-1-ynyl)phenyl)piperazin-1-yl)propoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-(naphthalen-2-yl)piperazin-1-yl)propoxy)indolin-2-one
	5-((S)-3-((R)-4-(4-chlorophenyl)-2-methylpiperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	5-((S)-3-((S)-4-(4-chlorophenyl)-2-methylpiperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	5-((S)-3-((2S,6R)-4-(4-chlorophenyl)-2,6-dimethylpiperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	5-((S)-3-((2S,6R)-2,6-dimethyl-4-p-tolylpiperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(R)-5-(3-(4-(4-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one








	(<i>R</i>)-5-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(<i>R</i>)-5-(3-(4-(4-ethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(<i>R</i>)-5-(2-hydroxy-3-(4-(4-isopropylphenyl)piperazin-1-yl)propoxy)indolin-2-one
	(<i>R</i>)-5-(3-(4-(4-cyclopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(<i>R</i>)-5-(2-hydroxy-3-(4-(4-propylphenyl)piperazin-1-yl)propoxy)indolin-2-one
	(<i>R</i>)-5-(3-(4-(4-butylphenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(<i>R</i>)-5-(2-hydroxy-3-(4-(4-isobutylphenyl)piperazin-1-yl)propoxy)indolin-2-one








	(<i>R</i>)-5-(2-hydroxy-3-(4-(4-(prop-1-ynyl)phenyl)piperazin-1-yl)propoxy)indolin-2-one
	(<i>R</i>)-5-(2-hydroxy-3-(4-(naphthalen-2-yl)piperazin-1-yl)propoxy)indolin-2-one
	(<i>R</i>)-5-(2-hydroxy-3-(4- <i>p</i> -tolylpiperazin-1-yl)propoxy)indolin-2-one
	(<i>R</i>)-5-(2-hydroxy-3-(4-phenylpiperazin-1-yl)propoxy)indolin-2-one
	(<i>R</i>)-5-(3-(4-(4-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)indolin-2-one








In one embodiment, the compound is selected from Table 7.

Table 7.








Compound	NAME
	(<i>S</i>)-6-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one








	(S)-6-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(R)-6-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(4-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(3,4-dimethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4- <i>p</i> -tolylpiperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-4-(4-(2-hydroxy-3-(2-oxo-1,2,3,4-tetrahydroquinolin-6-yloxy)propyl)piperazin-1-yl)benzonitrile
	(S)-6-(3-(4-(4-bromophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one





	(S)-6-(2-hydroxy-3-(4-(4-hydroxyphenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-methoxyphenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-(trifluoromethyl)phenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(biphenyl-4-yl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(2,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(2-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(2-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one

	(S)-6-(2-hydroxy-3-(4- <i>o</i> -tolylpiperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-2-(4-(2-hydroxy-3-(2-oxo-1,2,3,4-tetrahydroquinolin-6-yloxy)propyl)piperazin-1-yl)benzonitrile
	(S)-6-(2-hydroxy-3-(4-phenylpiperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(3-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(3-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4- <i>m</i> -tolylpiperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one

	(S)-6-(3-(4-(4-ethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-isopropylphenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(4-cyclopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-propylphenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(4-butylphenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-isobutylphenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-(prop-1-ynyl)phenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one



	(<i>S</i>)-6-(2-hydroxy-3-(4-(naphthalen-2-yl)-piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(<i>1H</i>)-one
	6-((<i>S</i>)-3-((<i>R</i>)-4-(4-chlorophenyl)-2-methylpiperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(<i>1H</i>)-one
	6-((<i>S</i>)-3-((<i>S</i>)-4-(4-chlorophenyl)-2-methylpiperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(<i>1H</i>)-one
	6-((<i>S</i>)-3-((2 <i>S</i> ,6 <i>R</i>)-4-(4-chlorophenyl)-2,6-dimethylpiperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(<i>1H</i>)-one
	6-((<i>S</i>)-3-((2 <i>S</i> ,6 <i>R</i>)-2,6-dimethyl-4- <i>p</i> -tolylpiperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(<i>1H</i>)-one
	(<i>R</i>)-6-(3-(4-(4-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(<i>1H</i>)-one
	(<i>R</i>)-6-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(<i>1H</i>)-one








	(<i>R</i>)-6-(3-(4-(4-ethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(2-hydroxy-3-(4-(4-isopropylphenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(3-(4-(4-cyclopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(2-hydroxy-3-(4-(4-propylphenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(3-(4-(4-butylphenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(2-hydroxy-3-(4-(4-isobutylphenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(2-hydroxy-3-(4-(4-(prop-1-ynyl)phenyl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one








	(<i>R</i>)-6-(2-hydroxy-3-(4-(naphthalen-2-yl)piperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(2-hydroxy-3-(4- <i>p</i> -tolylpiperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(2-hydroxy-3-(4-phenylpiperazin-1-yl)propoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(3-(4-(4-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1 <i>H</i>)-one








In another embodiment, the compound is selected from Table 8.








Table 8.

Compound	NAME
	(<i>S</i>)-6-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>S</i>)-6-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one








	(<i>R</i>)-6-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>S</i>)-6-(3-(4-(4-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>S</i>)-6-(3-(4-(3,4-dimethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>S</i>)-6-(2-hydroxy-3-(4- <i>p</i> -tolylpiperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(<i>S</i>)-4-(4-(2-hydroxy-3-(2-oxo-1,2-dihydroquinolin-6-yloxy)propyl)piperazin-1-yl)benzonitrile
	(<i>S</i>)-6-(3-(4-(4-bromophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>S</i>)-6-(2-hydroxy-3-(4-(4-hydroxyphenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one







	(S)-6-(2-hydroxy-3-(4-(4-methoxyphenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-(trifluoromethyl)phenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(biphenyl-4-yl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(2,4-difluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(2-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(2-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4- <i>o</i> -tolylpiperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one

	(S)-2-(4-(2-hydroxy-3-(2-oxo-1,2-dihydroquinolin-6-yloxy)propyl)piperazin-1-yl)benzonitrile
	(S)-6-(2-hydroxy-3-(4-phenylpiperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(3-fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(3-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4- <i>m</i> -tolypiperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(4-ethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one

	(S)-6-(2-hydroxy-3-(4-(4-isopropylphenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(4-cyclopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-propylphenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(3-(4-(4-butylphenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-isobutylphenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(4-(prop-1-ynyl)phenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(S)-6-(2-hydroxy-3-(4-(naphthalen-2-yl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one

	6-((<i>S</i>)-3-((<i>R</i>)-4-(4-chlorophenyl)-2-methylpiperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	6-((<i>S</i>)-3-((<i>S</i>)-4-(4-chlorophenyl)-2-methylpiperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	6-((<i>S</i>)-3-((2 <i>S</i> ,6 <i>R</i>)-4-(4-chlorophenyl)-2,6-dimethylpiperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	6-((<i>S</i>)-3-((2 <i>S</i> ,6 <i>R</i>)-2,6-dimethyl-4- <i>p</i> -tolylpiperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(3-(4-(4-chlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(3-(4-(4-ethylphenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one








	(R)-6-(2-hydroxy-3-(4-(4-isopropylphenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(R)-6-(3-(4-(4-cyclopropylphenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(R)-6-(2-hydroxy-3-(4-(4-propylphenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(R)-6-(3-(4-(4-butylphenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(R)-6-(2-hydroxy-3-(4-(4-isobutylphenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(R)-6-(2-hydroxy-3-(4-(4-(prop-1-ynyl)phenyl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(R)-6-(2-hydroxy-3-(4-(naphthalen-2-yl)piperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one

	(<i>R</i>)-6-(2-hydroxy-3-(4- <i>p</i> -tolylpiperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(2-hydroxy-3-(4-phenylpiperazin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(3-(4-(fluorophenyl)piperazin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>S</i>)-6-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>R</i>)-6-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)quinolin-2(1 <i>H</i>)-one
	(<i>S</i>)-6-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)quinolin-2(1 <i>H</i>)-one

In another embodiment, the compound is selected from Table 9.


Table 9.









Compound	Name
----------	------









	N-(4-(2-(3-(3,4-dichlorophenyl)-2-oxoimidazolidin-1-yl)ethoxy)phenyl)methanesulfonamide
	N-(4-(3-(3-(3,4-dichlorophenyl)-2-oxoimidazolidin-1-yl)propyl)phenyl)methanesulfonamide
	(S)-N-(4-(3-(3-(3,4-dichlorophenyl)-2-oxoimidazolidin-1-yl)-2-hydroxypropoxy)phenyl)methanesulfonamide
	(S)-N-(4-(3-(3-(4-chlorophenyl)-2-oxoimidazolidin-1-yl)-2-hydroxypropoxy)phenyl)methanesulfonamide
	(S)-1-(4-chlorophenyl)-3-(2-hydroxy-3-(4-hydroxyphenoxy)propyl)imidazolidin-2-one
	1-(4-chlorophenyl)-3-(3-(4-hydroxyphenoxy)propyl)imidazolidin-2-one
	(S)-1-(4-(3-(3-(4-chlorophenyl)-2-oxoimidazolidin-1-yl)-2-hydroxypropoxy)phenyl)urea









In another embodiment, the compound is selected from Table 10.









Table 10.








Compound	Name
	N-(2-(3,4-difluorophenylamino)ethyl)-2-(4-(methylsulfonamido)phenoxy)acetamide

	N-(2-(3,4-dichlorophenylamino)ethyl)-2-(4-(methylsulfonamido)phenoxy)acetamide
	N-(2-(3,4-dichlorophenylthio)ethyl)-2-(4-(methylsulfonamido)phenoxy)acetamide
	N-(2-(3,4-dichlorophenoxy)ethyl)-2-(4-(methylsulfonamido)phenoxy)acetamide
	(E)-N-(3-(3,4-dichlorophenyl)allyl)-2-(4-(methylsulfonamido)phenoxy)acetamide
	N-(3-(3,4-dichlorophenyl)propyl)-2-(4-(methylsulfonamido)phenoxy)acetamide
	N-(2-(3,4-dichlorophenylamino)ethyl)-3-(4-(methylsulfonamido)phenyl)propanamide
	N-(2-(3,4-dichlorophenylthio)ethyl)-3-(4-(methylsulfonamido)phenyl)propanamide
	N-(2-(3,4-dichlorophenoxy)ethyl)-3-(4-(methylsulfonamido)phenyl)propanamide
	(E)-N-(3-(3,4-dichlorophenyl)allyl)-3-(4-(methylsulfonamido)phenyl)propanamide









	N-(3-(3,4-dichlorophenyl)propyl)-3-(4-(methanesulfonamido)phenyl)propanamide
	N-(4-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-3-oxopropyl)phenyl)methanesulfonamide
	N-(4-(2-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-oxoethoxy)phenyl)methanesulfonamide
	N-(4-(3-(4-(3,4-dichlorophenyl)-2-oxopiperazin-1-yl)propyl)phenyl)methanesulfonamide
	N-(4-(2-(4-(3,4-dichlorophenyl)-2-oxopiperazin-1-yl)ethoxy)phenyl)methanesulfonamide
	(S)-N-(4-(3-(4-(4-chlorophenyl)-2-oxopiperazin-1-yl)-2-hydroxypropoxy)phenyl)methanesulfonamide
	(S)-N-(4-(3-(4-(3,4-difluorophenyl)-2-oxopiperazin-1-yl)-2-hydroxypropoxy)phenyl)methanesulfonamide
	N-(4-(3-(4-(3,4-difluorophenyl)-2-oxopiperazin-1-yl)propoxy)phenyl)methanesulfonamide





	<i>N</i> -(4-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-3-oxopropyl)phenyl)methanesulfonamide
	<i>N</i> -(4-(2-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-oxoethoxy)phenyl)methanesulfonamide
	6-(2-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-oxoethoxy)benzo[<i>d</i>]oxazol-2(3 <i>H</i>)-one
	6-(2-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-oxoethoxy)benzo[<i>d</i>]oxazol-2(3 <i>H</i>)-one
	6-(2-(4-(4-chlorophenyl)piperazin-1-yl)-2-oxoethoxy)benzo[<i>d</i>]oxazol-2(3 <i>H</i>)-one
	5-(2-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-oxoethoxy)indolin-2-one
	5-(2-(4-(4-chlorophenyl)piperazin-1-yl)-2-oxoethoxy)indolin-2-one
	5-(2-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-oxoethoxy)indolin-2-one


	6-(2-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-oxoethoxy)-3,4-dihydroquinolin-2(1H)-one
	6-(2-(4-(4-chlorophenyl)piperazin-1-yl)-2-oxoethoxy)-3,4-dihydroquinolin-2(1H)-one
	6-(2-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-oxoethoxy)-3,4-dihydroquinolin-2(1H)-one
	6-(2-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-oxoethoxy)quinolin-2(1H)-one
	6-(2-(4-(4-chlorophenyl)piperazin-1-yl)-2-oxoethoxy)quinolin-2(1H)-one
	6-(2-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-oxoethoxy)quinolin-2(1H)-one
	6-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-3-oxopropoxy)benzo[d]oxazol-2(3H)-one
	6-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-3-oxopropoxy)benzo[d]oxazol-2(3H)-one


	6-(3-(4-(4-chlorophenyl)piperazin-1-yl)-3-oxopropoxy)benzo[d]oxazol-2(3H)-one
	5-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-3-oxopropoxy)indolin-2-one
	5-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-3-oxopropoxy)indolin-2-one
	5-(3-(4-(4-chlorophenyl)piperazin-1-yl)-3-oxopropoxy)indolin-2-one
	6-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-3-oxopropoxy)-3,4-dihydroquinolin-2(1H)-one
	6-(3-(4-(4-chlorophenyl)piperazin-1-yl)-3-oxopropoxy)-3,4-dihydroquinolin-2(1H)-one
	6-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-3-oxopropoxy)-3,4-dihydroquinolin-2(1H)-one

	6-(3-(4-(3,4-difluorophenyl)piperazin-1-yl)-3-oxopropoxy)quinolin-2(1H)-one
	6-(3-(4-(4-chlorophenyl)piperazin-1-yl)-3-oxopropoxy)quinolin-2(1H)-one
	6-(3-(4-(3,4-dichlorophenyl)piperazin-1-yl)-3-oxopropoxy)quinolin-2(1H)-one
	5-(2-(4-(3,4-dichlorophenyl)-2-oxopiperazin-1-yl)ethoxy)indolin-2-one
	5-(2-(4-(4-chlorophenyl)-2-oxopiperazin-1-yl)ethoxy)indolin-2-one
	5-(2-(4-(3,4-difluorophenyl)-2-oxopiperazin-1-yl)ethoxy)indolin-2-one
	5-(3-(4-(3,4-difluorophenyl)-2-oxopiperazin-1-yl)propoxy)indolin-2-one
	5-(3-(4-(3,4-dichlorophenyl)-2-oxopiperazin-1-yl)propoxy)indolin-2-one

	5-(3-(4-(4-chlorophenyl)-2-oxopiperazin-1-yl)propoxy)indolin-2-one
	6-(2-(4-(3,4-dichlorophenyl)-2-oxopiperazin-1-yl)ethoxy)-3,4-dihydroquinolin-2(1H)-one
	6-(2-(4-(4-chlorophenyl)-2-oxopiperazin-1-yl)ethoxy)-3,4-dihydroquinolin-2(1H)-one
	6-(2-(4-(3,4-difluorophenyl)-2-oxopiperazin-1-yl)ethoxy)-3,4-dihydroquinolin-2(1H)-one
	6-(2-(4-(3,4-dichlorophenyl)piperazin-1-yl)-2-oxoethoxy)quinolin-2(1H)-one
	6-(2-(4-(4-chlorophenyl)piperazin-1-yl)-2-oxoethoxy)quinolin-2(1H)-one
	6-(2-(4-(3,4-difluorophenyl)piperazin-1-yl)-2-oxoethoxy)quinolin-2(1H)-one
	N-(2-(3,4-dichlorophenylamino)ethyl)-2-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yloxy)acetamide



	<i>N</i> -(2-(3,4-difluorophenylamino)ethyl)-2-(2-oxo-2,3-dihydrobenzo[<i>d</i>]oxazol-6-yloxy)acetamide
	<i>N</i> -(2-(3,4-difluorophenylamino)ethyl)-2-(4-hydroxyphenoxy)acetamide
	<i>N</i> -(2-(3,4-difluorophenylamino)ethyl)-3-(4-hydroxyphenyl)propanamide
	<i>N</i> -(2-(3,4-difluorophenylamino)ethyl)-2-(3-fluoro-4-hydroxyphenoxy)acetamide















In one embodiment, the compound is not . In















another embodiment, the compound is not .

















In one embodiment, the compound has an IC₅₀ value of 600 nM or less. In one embodiment, the compound has an IC₅₀ value of 600 nM or less at pH 6.9 or an ischemic pH. In one embodiment, the compound is selected from Table 11.


Table 11.

	
--	---

 <chem>OS(=O)(=O)c1ccc(OCC(O)CN2CCCN(C2)c3cc(Cl)cc(Cl)c3)cc1</chem>	 <chem>OS(=O)(=O)c1ccc(OCC(O)CN2CCCN(C2)c3ccc(Cl)cc3)cc1</chem>
 <chem>OS(=O)(=O)c1ccc(OCC(O)CN2CCCN(C2)c3ccccc3)cc1</chem>	 <chem>OS(=O)(=O)c1ccc(OCC(O)CN2CCCN(C2)CNc3ccccc3)cc1</chem>
 <chem>OS(=O)(=O)c1ccc(OCC(O)CN2CCCN(C2)c3cc(F)cc(F)c3)cc1</chem>	 <chem>OS(=O)(=O)c1ccc(OCC(O)CN2CCCN(C2)/C=C/c3cc(Cl)cc(Cl)c3)cc1</chem>
 <chem>OS(=O)(=O)c1ccc(OCC(O)CN2CCCN(C2)/C=C/c3cc(Cl)cc(Cl)c3)cc1</chem>	 <chem>OS(=O)(=O)c1ccc(OCC(O)CN2CCCN(C2)/C=C/c3cc(F)cc(F)c3)cc1</chem>
 <chem>O=C1OC(=O)N1c2ccc(OCC(O)CN2CCCN(C2)c3ccc(Cl)cc3)cc2</chem>	 <chem>O=C1OC(=O)N1c2ccc(OCC(O)CN2CCCN(C2)c3cc(F)cc(F)c3)cc2</chem>
 <chem>O=C1OC(=O)N1c2ccc(OCC(O)CN2CCCN(C2)c3cc(Cl)cc(Cl)c3)cc2</chem>	 <chem>Oc1ccc(OCC(O)CN2CCCN(C2)c3ccc(Cl)cc3)cc1</chem>
 <chem>Oc1ccc(OCC(O)CN2CCCN(C2)c3cc(Cl)cc(Cl)c3)cc1</chem>	 <chem>Oc1ccc(OCC(O)CN2CCCN(C2)c3cc(F)cc(F)c3)cc1</chem>













 <chem>Oc1ccc(O[C@H](CO)N2CCN(C2)c3cc(F)c(F)cc3)cc1</chem>	 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccc(F)cc3)cc1</chem>
 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccc(C)cc3)cc1</chem>	 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccc(C#N)cc3)cc1</chem>
 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccc(Br)cc3)cc1</chem>	 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccc(O)cc3)cc1</chem>
 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccc(OC)cc3)cc1</chem>	 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccc(C(F)(F)F)cc3)cc1</chem>
 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3cc(F)c(cc3F)F)cc1</chem>	 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccccc3F)cc1</chem>
 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3cc(C)ccccc3)cc1</chem>	 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3ccccc3)cc1</chem>
 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3cccc(F)c3)cc1</chem>	 <chem>Oc1ccc(O[C@@H](CO)N2CCN(C2)c3cccc(Cl)c3)cc1</chem>



















In one embodiment, the compound has an IC₅₀ value of 600 nM or less at pH 7.6 or a physiological pH. In one embodiment, the compound is selected from Table 12.










Table 12.


In one embodiment, the compound has a pH boost of 5 or more. In one embodiment, the compound is selected from Table 13.


Table 13.


	
	
	
	
	

In one embodiment, the compound is selected from the group consisting of:












In one embodiment, the compound is




In one embodiment, the compound is selected from the group consisting of:







In another embodiment, the compound is selected from the group consisting of:




In one embodiment, the compound has an IC_{50} of 600 nM or less and a pH boost of 5 or more. In a particular embodiment, the compound is




. In another embodiment, the compound is




. In another embodiment, the compound is,




. In another embodiment, the compound is,




. In a particular embodiment, the compound is




. In another particular embodiment, the compound is



. In another embodiment, the compound is,






. In another embodiment, the compound is,





.


In one embodiment, the compound is selected from the group consisting of:





















Formula II

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula II or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



FORMULA II


wherein:

each G is independently F, Cl, Br, I, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₆-C₁₂ aralkyl, -O-aryl, -S-aryl, -NH-aryl;

f = 0, 1, 2, 3, 4 or 5;

Ar^a and Ar^b are each independently aryl or heteroaryl;

B is selected from the group consisting of:



wherein R^a, R^b, R^c, R^d, R^e, R^f, R^g, R^h, R^k and R^p are each independently selected from H, C₁-C₆ alkyl, C₁-C₆ alkoxy, OH or halo;

R^j is H, C₁-C₆ alkyl, OH or P(O)(OC₁-C₄ alkyl)₂;


R^m is C₁-C₄ alkyl or C₂-C₄ alkenyl;

Rⁿ is C₁-C₄ alkyl, C₂-C₄ alkenyl, C₆-C₁₂ aralkyl, -CH₂O-, -CH(C₁-C₆ alkyl)O-, -CH(C₂-C₁₂ aralkyl)O-;

t, w, y and z each = 0, 1, 2, or 3;

X and X' are independently selected from a bond, O, S, SO, SO₂, CH₂, NH, N(C₁-C₆ alkyl), and NHC(=O);

M is OH, F, Cl, Br, I, NH₂, NR^qR^r, NO₂, O(C₁-C₆ alkyl), OCF₃, CN, C(O)OH, C(O)O(C₁-C₆ alkyl), C₆-C₁₂ aralkyl, NR^sC(O)CR^t₃, NR^sSO₂(C₁-C₆ alkyl), or NR^uC(O)NR^v₂; wherein each R^q, R^r, R^s, R^u and R^v is each independently H or C₁-C₆ alkyl; and each R^t is independently H, C₁-C₆ alkyl or halo; or two M groups may be taken together with Ar^b to form:



and R^w are independently H, C_1 - C_6 alkyl or C_6 - C_{12} aralkyl; and


$h = 1, 2, 3, 4$ or 5 .

In some embodiments, when B contains a piperidin-4-ol or a pyrrolidin-2-ol moiety, and Ar^a and Ar^b are each phenyl, M is not OH at the para position on Ar^b .

In some embodiments of Formula II, each G is independently F, Cl, Br, I, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_6 - C_{12} aralkyl, -O-aryl, -S-aryl, -NH-aryl;

$f = 0, 1, 2, 3, 4$ or 5 ;

Ar^a and Ar^b are each independently aryl or heteroaryl;



B is

wherein R^{a-h} , R^k and R^p are each independently selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, OH or halo;

R^j is H, C_1 - C_6 alkyl, OH or $P(O)(OC_1-C_4 \text{ alkyl})_2$;


R^m is C_1 - C_4 alkyl or C_2 - C_4 alkenyl;

R^n is C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_6 - C_{12} aralkyl, $-CH_2O-$, $-CH(C_1-C_6 \text{ alkyl})O-$, $-CH(C_2-C_{12} \text{ aralkyl})O-$;

t, w, y and z each = 0, 1, 2, or 3;

X and X' are independently selected from a bond, O, S, SO, SO_2 , CH_2 , NH, $N(C_1-C_6 \text{ alkyl})$, and $NHC(=O)$;

M is OH, F, Cl, Br, I, NH_2 , NR^qR^r , NO_2 , $O(C_1-C_6 \text{ alkyl})$, OCF_3 , CN, $C(O)OH$, $C(O)O(C_1-C_6 \text{ alkyl})$, C_6 - C_{12} aralkyl, $NR^sC(O)CR^t$, $NR^8SO_2(C_1-C_6 \text{ alkyl})$, or $NR^uC(O)NR^v$; wherein each R^q , R^r , R^s , R^u and R^v is each independently H or C_1 - C_6 alkyl; and each R^t is independently H, C_1 - C_6 alkyl or halo; or two M groups may be taken together with Ar^b to form:


alkyl or C₆-C₁₂ aralkyl; and

$h = 1, 2, 3, 4$ or 5 .

In one embodiment, G is F or Cl. In another embodiment, f is 1 or 2.


In one embodiment, Ar^a is phenyl. In another embodiment, Ar^b is phenyl. In another embodiment, Ar^a and Ar^b are each phenyl. In one embodiment, Ar^a is phenyl and is substituted with two G groups. In a subembodiment, both G groups are Cl. In another subembodiment, both G groups are F. In another subembodiment, one G group is Cl and the other G group is F. In one embodiment, G is selected from the group consisting of C₆-C₁₂ aralkyl, -O-aryl, -S-aryl, and -NH-aryl.

In one embodiment, B is




R^j . In a subembodiment, $R^a, R^b, R^c, R^d, R^e, R^g$ and R^h are H; R^j is H, C_1 - C_6 alkyl, OH or $P(O)(OC_1-C_4 \text{ alkyl})_2$; R^f is H, halo or OH; t is 0, 1, 2, or 3; and w, y and z are each 1.

In one embodiment, B is




In a subembodiment, R^a, R^b, R^c, R^d, R^e, R^g and R^h are H; R^f and R^k are independently H, halo or OH; R^m is C₁-C₄ alkyl; t is 1, 2, or 3; and w, y and z are each 1. In specific subembodiments, B is




, and R^f and R^k are independently H or OH.

In certain subembodiments, Ar^b is substituted with one, two or three M groups, wherein the M group is independently selected from OH, F, Cl, Br, I, or $\text{NR}^u\text{C}(\text{O})\text{NR}^v_2$; wherein each R^u and R^v is each independently H or $\text{C}_1\text{-C}_6$ alkyl or two M groups may be taken together with Ar^b to form



. In certain subembodiments, X' is selected from a bond, O, S, CH_2 , NH. In particular subembodiments, f is 1 and G is at a para position of Ar^a .

In one embodiment, B is



. In a subembodiment,

R^a , R^b , R^c , R^d , R^e , R^g and R^h are H; R^f is H, halo or OH; R^p is H, halo or OH; R^n is $-\text{CH}_2\text{O}-$; t is 0, 1, 2, or 3; and w, y and z are each 1.

In one embodiment, the sum of w, y and z does not exceed 6. In one embodiment, the sum of w, y and z is 2, 3, 4, 5 or 6.


In one embodiment, X is a bond, O, S or CH_2 . In another embodiment, X is O. In another embodiment, X is CH_2 .

In one embodiment, X' is a bond, NH, S or CH_2 . In another embodiment, X' is a bond. In another embodiment, X' is S. In another embodiment, X' is NH. In another embodiment, X' is CH_2 .


In one embodiment, M is OH. In another embodiment, M is F or Cl. In another embodiment, M is $\text{O}(\text{C}_1\text{-C}_6 \text{ alkyl})$, for example OCH_3 , OCH_2CH_3 , $\text{O}(\text{CH}_2)_2\text{CH}_3$, $\text{OCH}(\text{CH}_3)_2$ or $\text{OC}(\text{CH}_3)_3$. In another embodiment, M is NH_2 . In another embodiment, M is NR^qR^r . In another embodiment, M is NO_2 . In another embodiment, M is OCF_3 . In one embodiment, M is CN. In one embodiment, M is $\text{C}(\text{O})\text{OH}$. In one embodiment, M is $\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_6 \text{ alkyl})$, for example $\text{C}(\text{O})\text{OCH}_3$, $\text{C}(\text{O})\text{OCH}_2\text{CH}_3$, $\text{C}(\text{O})\text{O}(\text{CH}_2)_2\text{CH}_3$, $\text{C}(\text{O})\text{OCH}(\text{CH}_3)_2$ or

$\text{C(O)OC(CH}_3)_3$. In one embodiment, M is $\text{C}_6\text{-C}_{12}$ aralkyl, for example $\text{CH}_2\text{-phenyl}$. In one embodiment, M is $\text{NR}^s\text{C(O)CR}^t_3$. In a subembodiment, R^s is H. In a subembodiment, R^t is H or Cl. In one embodiment, M is $\text{NR}^u\text{C(O)NR}^v_2$, for example, NHC(O)NH_2 . In a subembodiment, R^u is H and R^v is H or alkyl.

In one embodiment, two M groups may be taken together with Ar^b to form:



subembodiment, two M groups may be taken together with Ar^b to form:



2. In one embodiment, R^u and R^w are both H. In one embodiment, h is 1 or


In one embodiment, the compound is a compound of Formula II, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein:


each G is independently F, Cl, Br or I;

f is 0, 1, 2, 3, 4, or 5;

Ar^a and Ar^b are each independently selected from the group consisting of phenyl, pyridyl, pyrimidinyl, thiophenyl, imidazolyl, furanyl, indolyl, benzothiophenyl, benzofuranyl, benzoimidazolyl;

B is selected from the group consisting of:





wherein R^a , R^b , R^c , R^d , R^e , R^f , R^g , R^h , R^k and R^p are each independently selected from H, C_1 - C_6 alkyl, OH, or halo;

R^j is H, C_1 - C_6 alkyl, C_7 - C_{12} aralkyl, or OH;

R^m is C_1 - C_4 alkyl or C_2 - C_4 alkenyl;


R^n is C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_6 - C_{12} aralkyl, $-\text{CH}_2\text{O}-$, $-\text{CH}(\text{C}_1\text{-C}_6 \text{ alkyl})\text{O}-$, $-\text{CH}(\text{C}_2\text{-C}_{12} \text{ aralkyl})\text{O}-$;

t , w , y and z each = 0, 1, 2, or 3;

X is a bond, CH_2 or O;

X' is a bond, CH_2 , S or NH;

M is OH, F, Cl, Br, I, NH_2 , NR^qR^r , NO_2 , $\text{O}(\text{C}_1\text{-C}_6 \text{ alkyl})$, OCF_3 , CN, $\text{C}(\text{O})\text{OH}$, $\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_6 \text{ alkyl})$, $\text{C}_6\text{-C}_{12}$ aralkyl, $\text{NR}^s\text{C}(\text{O})\text{CR}^t_3$, or $\text{NR}^u\text{C}(\text{O})\text{NR}^v_2$; wherein each R^q , R^r , R^s , R^u and R^v is each independently H or C_1 - C_6 alkyl; and each R^t is independently H, C_1 - C_6 alkyl or halo; or two M groups may be taken together with Ar^b to form:



and R^w are independently H or C_1 - C_4 alkyl; and

$h = 1, 2$ or 3 .


In one embodiment, the compound is a compound of Formula II, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein:

each G is independently F, Cl, Br or I;

$f = 0, 1, 2, 3, 4$ or 5 ;

Ar^a and Ar^b are each phenyl;

B is selected from the group consisting of:



wherein R^{a} , R^{b} , R^{c} , R^{d} , R^{e} , R^{f} , R^{g} , R^{h} , R^{k} and R^{p} are each independently selected from H, C_1 - C_6 alkyl, OH, or halo;

R^{j} is H, C_1 - C_6 alkyl, or OH;

R^{m} is C_1 - C_4 alkyl or C_2 - C_4 alkenyl;


R^{n} is C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_6 - C_{12} aralkyl, $-\text{CH}_2\text{O}-$, $-\text{CH}(\text{C}_1\text{-C}_6 \text{ alkyl})\text{O}-$, $-\text{CH}(\text{C}_2\text{-C}_{12} \text{ aralkyl})\text{O}-$;

t, w, y and z each = 0, 1, 2, or 3;

X is a bond, CH_2 or O;

X' is a bond, CH_2 , S or NH;

M is OH, F, Cl, Br, I, NH_2 , $\text{NR}^{\text{q}}\text{R}^{\text{r}}$, NO_2 , $\text{O}(\text{C}_1\text{-C}_6 \text{ alkyl})$, OCF_3 , CN, $\text{C}(\text{O})\text{OH}$, $\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_6 \text{ alkyl})$, $\text{C}_6\text{-C}_{12} \text{ aralkyl}$, $\text{NR}^{\text{s}}\text{C}(\text{O})\text{CR}^{\text{t}}_3$; wherein each R^{q} , R^{r} , and R^{s} is each independently H or C_1 - C_6 alkyl; and each R^{t} is independently H, C_1 - C_6 alkyl or halo; or two M groups may be taken together with Ar^{b} to form:






; and wherein R^{u} is H or C_1 - C_4 alkyl; and

h = 1, 2 or 3.




In one embodiment, M is $\text{NR}^{\text{u}}\text{C}(\text{O})\text{NR}^{\text{v}}_2$, for example $\text{NHC}(\text{O})\text{NH}_2$ or $\text{NHC}(\text{O})\text{N}(\text{CH}_3)_2$.








In another embodiment, $\text{Ar}^{\text{b}}\text{-M}$ is selected from the group consisting of:














In one embodiment, the compound is selected from the compounds in Table 14.

Table 14.

Compound	NAME
	4-{3-[2-(3,4-Dichloro-phenyl)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-phenol
	4-(3-{Butyl-[2-(3,4-dichloro-phenyl)-ethyl]-amino}-2-(<i>S</i>)-hydroxy-propoxy)-phenol
	4-{3-[2-(3,4-Dichloro-phenyl)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-3-fluoro-phenol



	4-{3-[2-(3,4-Dichloro-phenyl)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-2-fluoro-phenol
	1-[2-(<i>S</i>)-Hydroxy-3-(4-hydroxy-phenoxy)-propyl]-4-phenyl-piperidin-4-ol
	<i>(R)</i> -1-(4-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)phenyl)urea
	<i>(S)</i> -1-(4-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)phenyl)urea
	<i>(S)</i> -1-(4-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)phenyl)urea
	<i>(R)</i> -1-(4-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)phenyl)urea
	1-(4-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)phenyl)urea




	1-(4-(3-(4-(4-chlorobenzyl)-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)phenyl)urea
	1-(4-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)phenyl)urea
	(S)-5-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(R)-5-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)indolin-2-one
	(S)-5-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)indolin-2-one
	(R)-5-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)indolin-2-one
	(R)-6-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1H)-one

	(S)-6-(3-(4-benzyl-4-hydroxypiperidin-1-yl)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1H)-one
	(S)-6-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)-3,4-dihydroquinolin-2(1H)-one
	(R)-6-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)-3,4-dihydroquinolin-2(1H)-one
	(R)-6-(2-hydroxy-3-(4-hydroxy-4-phenylpiperidin-1-yl)propoxy)quinolin-2(1H)-one


In one embodiment, the compound has an IC₅₀ value of 600 nM or less. In one embodiment, the compound has an IC₅₀ value of 600 nM or less at pH 6.9 or an ischemic pH. In one embodiment, the compound is selected from Table 15.

Table 15.


	
---	---

In one embodiment, the compound has a pH boost of 5 or more. In one embodiment,




In one embodiment, the compound is selected from the group consisting of:




In one embodiment, the compound is

In another embodiment, the compound is selected from the group consisting of:



In another embodiment the compound is



In one embodiment, one or more of R^c , R^d , R^e , R^f , R^g , and R^h is an OH group which creates a stereogenic center. In a particular subembodiment, one of R^c , R^d , R^e , R^f , R^g , and R^h is an OH group which creates a stereogenic center. In another subembodiment, the OH group at one of R^c , R^d , R^e , R^f , R^g , and R^h is in the R configuration. In another subembodiment, the OH group at one of R^c , R^d , R^e , R^f , R^g , and R^h is in the S configuration.

In certain embodiments, the binding to both hERG and alpha-1 adrenergic receptors can be modulated by changing the G substituent or G substituents. In particular, for compounds wherein Ar^a is phenyl, the binding to both hERG and alpha-1 adrenergic receptors can be modulated by changing the substitution at the 3 and/or 4 positions. In one embodiment, the Ar^a phenyl is substituted at the 3 and/or 4 position with, for example, fluorine or chlorine. In certain embodiments, substitution at the 3 and/or 4 positions of the Ar^a phenyl can increase potency.


In certain embodiments, both hERG and alpha-1 adrenergic binding can be reduced by substituting N at the R^j position with C_7 - C_{12} aralkyl. In a particular subembodiment, R^j is benzyl.

In certain embodiments, alpha-1 adrenergic binding is reduced when R^j is C_1 - C_6 alkyl.


When Ar^b is phenyl, *para* substitution of the M substituent is particularly preferred. Additional M substituents on the Ar^b phenyl are preferred at one or more *ortho* positions. Additional substitution on the Ar^b phenyl at one or more *meta* positions can reduce potency.

In certain embodiments, the Ar^a phenyl is not substituted by two fluoro groups. In one embodiment, the Ar^a phenyl is not substituted by two methyl groups. In one embodiment, the Ar^a phenyl is not substituted by one halo group. In one embodiment, the Ar^a phenyl is not substituted by one fluoro or alkyl group at the C-2 position. In one embodiment, the Ar^a phenyl is not substituted by a OH or NO₂ group.

In one embodiment, when Ar^a and Ar^b are both phenyl, at least one of f or h is not 0. In one embodiment, when Ar^a and Ar^b are both phenyl, f is not 0. In one embodiment, when Ar^a and Ar^b are both phenyl, h is not 0. In one embodiment, when Ar^a and Ar^b are both phenyl, X is not CH₂. In one embodiment, when Ar^a and Ar^b are both phenyl, X' is not CH₂. In another embodiment, M is not OH. In one embodiment, the compound is not



In one embodiment, M is not aralkoxy. In one embodiment, the compound is not



In one embodiment, B does not contain a piperidinyl moiety. In another embodiment, when B contains a piperidinyl moiety, and Ar^a and Ar^b are both phenyl, M is not OH. In one embodiment, when B contains a piperidinyl moiety, M is NR^uC(O)NR^v₂, for example, NHC(O)NH₂. In a subembodiment, R^u is H and R^v is H or alkyl. In one embodiment, when B contains a piperidinyl moiety, X is not CH₂. In one embodiment, when B contains a piperidinyl moiety, X' is not CH₂. In one embodiment, R^k is not OH. In one embodiment, R^p is not OH.

In one embodiment, when B contains a hydroxy-substituted-piperidinyl moiety, X is not CH₂. In one embodiment, when B contains a hydroxy-substituted-piperidinyl moiety, X' is not CH₂. In one embodiment, B does not contain a hydroxy-substituted-piperidinyl moiety.

In one embodiment, X is not SO₂. In another embodiment, X' is not SO₂. In one embodiment, when B contains a piperidinyl moiety, X is not SO₂. In one embodiment, when B contains a piperidinyl moiety, X' is not SO₂.


In one embodiment, X is not S. In another embodiment, X' is not S. In one embodiment, when B contains a piperidinyl moiety, X is not S. In one embodiment, when B contains a piperidinyl moiety, X' is not S.

In another embodiment, M is not OCH₃ or OCF₃. In another embodiment, M is not NO₂. In one embodiment, when B contains a nitrogen-containing heterocycle, Ar^b-X is not heteroaryl-NH. In another embodiment, when B contains a nitrogen-containing heterocycle, Ar^a-X' is not heteroaryl-NH.

In one embodiment, when B contains a nitrogen-containing heterocycle, X is not NH(C=O). In another embodiment, when B contains a nitrogen-containing heterocycle, X' is not NH(C=O).

Formula III


In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula III or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



FORMULA III

wherein:

Z^* is OH, $\text{NR}^{10*}\text{R}^{11*}$, $\text{NR}^{12*}\text{SO}_2\text{R}^{11*}$, $\text{NR}^{12*}\text{C(O)NR}^{10*}\text{R}^{11*}$, $\text{NR}^{12*}\text{C(O)OR}^{10*}$, NR^{12*} -dihydrothiazole, or NR^{12*} -dihydroimidazole; wherein each R^{10*} , R^{11*} and R^{12*} is independently H, C_1 - C_6 alkyl or C_6 - C_{12} aralkyl; or



Ar^{1*} and Ar^{2*} are each independently aryl or heteroaryl;

R^{1*} , R^{2*} , R^{4*} , R^{5*} , R^{7*} , R^{8*} are independently H, OH or C_1 - C_4 alkyl;

$n^* = 1, 2, 3$ or 4 ;

$p^* = 0, 1, 2$ or 3 ;

$q^* = 0, 1$ or 2 ;


R^{3*} and R^{6*} are each independently H or C_1 - C_4 alkyl;

X^{1*} and X^{2*} are each independently O, S, $\text{N}(\text{C}_1\text{-C}_4 \text{ alkyl})$ or $\text{C}(\text{H or C}_1\text{-C}_4 \text{ alkyl})_2$;


W^* is NR^{9*} or $\text{CR}^{13*}\text{R}^{14*}$; wherein R^{9*} , R^{13*} and R^{14*} are each independently is H or C_1 - C_4 alkyl;

each L^* is independently C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $\text{C(=O)-(C}_1\text{-C}_6\text{)-alkyl}$, C_1 - C_6 haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or two L groups may be taken together with Ar^{2*} to form a dioxolane ring or a cyclobutane ring;

$k^* = 0, 1, 2, 3, 4$ or 5 ;



In one embodiment, Z^* is OH, $NR^{12*}SO_2R^{11*}$; wherein R^{12*} is H or C_1 - C_4 alkyl, and R^{11*} is C_1 - C_4 alkyl or C_7 - C_{10} aralkyl. In one embodiment, Z^* is OH. In another embodiment, Z^* is $NR^{12*}SO_2R^{11*}$, for example, $NHSO_2CH_3$.



In one embodiment, Ar^{1*} and Ar^{2*} are each phenyl.

In one embodiment, R^{1*} , R^{2*} , R^{4*} , R^{5*} , R^{7*} , R^{8*} are H.

In a particular embodiment, n^* is 2.

In one embodiment, p^* is 0, 1 or 2. In another embodiment, p^* is 0. In another embodiment, p^* is 1. In another embodiment, p^* is 2.

In one embodiment, q^* is 0. In another embodiment, q^* is 1. In another embodiment, q^* is 2.

In one embodiment, R^{3*} and R^{6*} are both H. In one embodiment, R^{6*} is C_1 - C_4 alkyl, for example methyl, ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl, or *tert*-butyl.

In one embodiment, X^* is S. In one embodiment, X^* is O.

In one embodiment, W^* is NR^{7*} , for example NH. In another embodiment, W^* is $CR^{13*}R^{14*}$, for example CH_2 .

In one embodiment, each L^* is independently selected from C_1 - C_4 alkyl, F, Cl, Br, I, or C_1 - C_4 haloalkyl, for example, Cl, CH_3 or CF_3 . In one embodiment, k^* is 1. In another embodiment, k^* is 2.

In one embodiment, the compound is a compound of Formula III, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein: Z^* is OH, $NHSO_2CH_3$;

Ar^{1*} is phenyl;

R^{1*} , R^{2*} , R^{4*} , R^{5*} are independently H or $\text{C}_1\text{-C}_4$ alkyl;

$n^* = 2$;

$p^* = 0, 1$ or 2 ;

$q^* = 0, 1$ or 2 ;

R^{3*} and R^{6*} are each independently H or $\text{C}_1\text{-C}_4$ alkyl;


X^* is O or S;

W^* is NR^{7*} or $\text{CR}^{13*}\text{R}^{14*}$; wherein R^{7*} , R^{13*} and R^{14*} are each independently is H or $\text{C}_1\text{-C}_4$ alkyl;


Ar^{2*} is phenyl;


each L^* is independently selected from $\text{C}_1\text{-C}_4$ alkyl, F, Cl, Br, I, $\text{C}_1\text{-C}_4$ haloalkyl;

$k^* = 0, 1, 2, 3, 4$ or 5 ;




In one embodiment, the compound is selected from the group consisting of:





In another embodiment, the compound is selected from the group consisting of:



Formula IV

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a

compound of Formula IV or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



FORMULA IV

wherein:

each L^{**} is independently C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $C(=O)$ -(C_1 - C_6)-alkyl, C_1 - C_6 haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or two L^{**} groups may be taken together with Ar^{1**} to form: a dioxolane ring or a cyclobutane ring;

$k^{**} = 0, 1, 2, 3, 4$ or 5 ;

each Ar^{1**} and Ar^{2**} is independently aryl or heteroaryl;


X^{**} is S, O or NR^3 ; wherein R^3 is H, C_1 - C_6 alkyl, or C_6 - C_{12} aralkyl;

each R^{1**} and R^{2**} is independently H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_6 - C_{12} aralkyl, $C(=O)$ -(C_1 - C_6)-alkyl, C_1 - C_6 haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR^1R^2 can be $C=O$ or $C=CH_2$;

$n^{**} = 1, 2, 3$ or 4 ;

Y^{**} is a bond, O, S, SO, SO_2 , CH_2 , NH, $N(C_1-C_6 \text{ alkyl})$, or $NHC(=O)$;

Z^{**} is OH, $NR^{6**}R^{7**}$, $NR^{8**}SO_2(C_1-C_6 \text{ alkyl})$, $NR^{8**}C(O)NR^{6**}R^{7**}$, $NR^{8**}C(O)O(C_1-C_6 \text{ alkyl})$, NR^{8**} -dihydrothiazole, or NR^{8**} -dihydroimidazole; wherein each R^{6**} , R^{7**} and R^{8**} is independently H, C_1 - C_6 alkyl or C_6 - C_{12} aralkyl; or



In particular subembodiment, Ar^{1**} is phenyl, pyridyl, pyrimidinyl, thiophenyl, imidazolyl, furanyl, indolyl, benzothiophenyl, benzofuranyl, or benzoimidazolyl. In one


embodiment, Ar^{2**} is phenyl. In another embodiment, Ar^{1**} is benzoimidazolyl. In a particular subembodiment, Ar^{2**} is phenyl and Ar^{1**} is a heteroaryl, for example benzoimidazolyl. In one embodiment, Ar^{1**} is a bicyclic group wherein the X^{**} group is attached to the heterocyclic ring.


In one embodiment, X^{**} is S. In one embodiment, X^{**} is O. In one embodiment, X^{**} is NR^{3**} , for example NH.

In another particular subembodiment, L^{**} is C_1 - C_4 alkyl, C_1 - C_4 alkoxy, $\text{C}(=\text{O})$ -(C_1 - C_4)-alkyl, C_1 - C_6 haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano. In a further subembodiment, L^{**} is methyl, trifluoromethyl, methoxy, nitro, fluoro, chloro or hydroxy. In one further subembodiment, there are one, two or three L^{**} groups substituting Ar^{1**} . In one subembodiment, Ar^{1**} is substituted with one fluoro group. In one subembodiment, Ar^{1**} is substituted with two fluoro groups. In one subembodiment, Ar^{1**} is substituted with one fluoro group and one chloro group. In one subembodiment, Ar^{1**} is substituted with one chloro group. In one subembodiment, Ar^{1**} is substituted with two chloro groups. In one subembodiment, Ar^{1**} is substituted with one methyl group. In one subembodiment, Ar^{1**} is substituted with one trifluoromethyl group.

In one embodiment, each R^{1**} and R^{2**} is independently H or C_1 - C_4 alkyl, for example, methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl or *tert*-butyl. In one embodiment, R^{1**} and R^{2**} are both H. In one embodiment, one R^{1**} or R^{2**} is hydroxy. In one embodiment, n^{**} is 2, 3, or 4. In one embodiment, n^{**} is 3.


In one embodiment, one $\text{CR}^{1**}\text{R}^{2**}$ is $\text{C}=\text{O}$ or $\text{C}=\text{CH}_2$. In one embodiment,





In one embodiment, Y^{**} is a bond, O or CH_2 . In one embodiment, Y^{**} is O. In one subembodiment, Ar^{2**} is phenyl and is substituted with a Z^{**} group at the 4 position.

In one embodiment, Z^{**} is OH, $NR^{6**}R^{7**}$, $NR^{8**}SO_2(C_1-C_6 \text{ alkyl})$, $NR^{8**}C(O)NR^{6**}R^{7**}$, $NR^{8**}C(O)O(C_1-C_6 \text{ alkyl})$, NR^{8**} -dihydrothiazole, or NR^{8**} -dihydroimidazole. In one subembodiment, Ar^{2**} is phenyl and is substituted with a Z^{**} group at the 4 position. In one embodiment,



In one embodiment, the compound is a compound of Formula IV, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof, wherein:

L^{**} is C_1-C_4 alkyl, C_1-C_4 alkoxy, $C(=O)-(C_1-C_4)$ -alkyl, C_1-C_4 haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, or nitro;

$k^{**} = 0, 1, 2, 3, 4$ or 5 ;

Ar^{1**} is selected from the group consisting of phenyl, pyridyl, pyrimidinyl, thiophenyl, imidazolyl, furanyl, indolyl, benzothiophenyl, benzofuranyl, or benzoimidazolyl;

Ar^{2**} is phenyl;


X^{**} is S;

each R^{1**} and R^{2**} is independently H, hydroxy or C_1 - C_4 alkyl; or $CR^{1**}R^{2**}$ is $C=O$;

$n^{**} = 2, 3$ or 4 ;


Y^{**} is O;

Z^{**} is OH, NH_2 , $NHSO_2(C_1-C_4 \text{ alkyl})$, $NHC(O)NR^{6**}R^{7**}$, $NHC(O)O(C_1-C_4 \text{ alkyl})$, NH-dihydrothiazole, or NH-dihydroimidazole; wherein each R^{6**} and R^{7**} is independently H or C_1 - C_4 alkyl; or




R^{9**} is H or C_1 - C_4 alkyl.


In one embodiment, the compound is selected from the group consisting of:





In one embodiment, the compound is




In one embodiment, the compound is selected from the group consisting of:






In one embodiment, the compound is

In another embodiment, the compound is selected from the group consisting of:





and



In another embodiment, the compound is selected from Table 16.

Table 16.

Compound	Name
	(<i>R</i>)-4-(3-(5,6-dichloro-1H-benzo[d]imidazol-2-ylthio)-2-hydroxypropoxy)phenol
	(<i>R</i>)-5-(3-(5,6-dichloro-1H-benzo[d]imidazol-2-ylthio)-2-hydroxypropoxy)indolin-2-one
	(<i>R</i>)-6-(3-(5,6-dichloro-1H-benzo[d]imidazol-2-ylthio)-2-hydroxypropoxy)-3,4-dihydroquinolin-2(1H)-one
	(<i>R</i>)-6-(3-(5,6-dichloro-1H-benzo[d]imidazol-2-ylthio)-2-hydroxypropoxy)quinolin-2(1H)-one
	(<i>R</i>)-1-(4-(3-(5,6-dichloro-1H-benzo[d]imidazol-2-ylthio)-2-hydroxypropoxy)phenyl)urea

	(<i>R</i>)-N-(4-(3-(5,6-dichloro-1H-benzo[d]imidazol-2-ylthio)-2-hydroxypropoxy)phenyl)methanesulfonamide
	(<i>R</i>)-N-(4-(2-hydroxy-3-(6-methyl-1H-benzo[d]imidazol-2-ylthio)propoxy)phenyl)methanesulfonamide


Formula V

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula V or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



FORMULA V

wherein B' is selected from the group consisting of:



W' is a bond or C₁-C₄ alkyl;

W'' is C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, C₁-C₄ haloalkyl or C(=O)-C₁-C₄ alkyl;


Y' is selected from a bond, O, S, CH₂ and N;

Ar' is an substituted or unsubstituted aromatic or nonaromatic cycloalkyl which optionally may include 0-3 heteroatoms;


Ar'' is an aromatic or nonaromatic cycloalkyl which optionally may include 0-3 heteroatoms;

Z' is NRC(O)NR₂ wherein each R is independently selected from H, C₁-C₆ alkyl or C₆-C₁₂ aralkyl; or

Ar''-Z' are taken together and selected from the group consisting



In one embodiment, Ar' is substituted by (L')_{k'}, wherein each L' is independently C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, alkaryl, hydroxy, -O-alkyl, -O-aryl, -SH, -S-alkyl, -S-aryl, fluoro, chloro, bromo, iodo, nitro, or cyano; or two L' groups may be taken together with Ar' to form a dioxolane ring or a cyclobutane ring; and k' = 1, 2, 3, 4 or 5.



In one embodiment, W' is a bond. In another embodiment, W' is C₁-C₄ alkyl, for example methylene, ethylene, or propylene. In a particular subembodiment, W' is CH₂.

In one embodiment W'' is C₁-C₄ alkyl, for example methylene, ethylene, propylene, methylpropylene, or butylene. In another embodiment, W'' is C₁-C₄ hydroxyalkyl, for example hydroxymethylene, hydroxyethylene, or hydroxypropylene. In a particular subembodiment, W'' is -CH₂, CH(OH)-CH₂-. In another embodiment, W'' is C₁-C₄ haloalkyl, for example fluoroethylene, fluoropropylene, chloroethylene, or chloropropylene. In another embodiment, W'' is C(=O)-C₁-C₄ alkyl, for example -C(=O)-CH₂- or -C(=O)-CH₂-CH₂-.




In one embodiment, Ar' is an aromatic cycloalkyl, for example phenyl. In another embodiment, Ar' is a nonaromatic cycloalkyl, for example cyclopentyl or cyclohexyl. In another embodiment, Ar' is an aromatic cycloalkyl including 1-3 heteroatoms, for example pyrrole, furan, thiophene, pyridine, pyrimidine, pyrazine, pyridazine. Heteroatoms include


but are not limited to N, S and O. In another embodiment, Ar' is a nonaromatic cycloalkyl including 1-3 heteroatoms, for example pyrrolidine, pyrroline, dihydrofuran, tetrahydrofuran, dihydrothiophene, tetrahydrothiophene, piperidine, tetrahydropyran, pyran, thiane, thiiine, piperazine, oxazine, dithiane, or dioxane. In another embodiment, Ar' is an aromatic or nonaromatic cycloalkyl including 1 heteroatom. In another embodiment, Ar' is an aromatic or nonaromatic cycloalkyl including 2 heteroatoms. In another embodiment, Ar' is an aromatic or nonaromatic cycloalkyl including 3 heteroatoms.

In one embodiment, Ar'' is an aromatic cycloalkyl, for example phenyl. In another embodiment, Ar'' is a nonaromatic cycloalkyl, for example cyclopentyl or cyclohexyl. In another embodiment, Ar'' is an aromatic cycloalkyl including 1-3 heteroatoms, for example pyrrole, furan, thiophene, pyridine, pyrimidine, pyrazine, or pyridazine. In another embodiment, Ar'' is a nonaromatic cycloalkyl including 1-3 heteroatoms, for example pyrrolidine, pyrroline, dihydrofuran, tetrahydrofuran, dihydrothiophene, tetrahydrothiophene, piperidine, tetrahydropyran, pyran, thiane, thiiine, piperazine, oxazine, dithiane, or dioxane. In another embodiment, Ar'' is an aromatic or nonaromatic cycloalkyl including 1 heteroatom. In another embodiment, Ar'' is an aromatic or nonaromatic cycloalkyl including 2 heteroatoms. In another embodiment, Ar'' is an aromatic or nonaromatic cycloalkyl including 3 heteroatoms.


In one embodiment, Z' is NRC(O)NR_2 , for example NHC(O)NH_2 or $\text{NHC(O)N(CH}_3)_2$.

In another embodiment, Z and Ar'' are taken together and selected from the group






. In another subembodiment, Ar''-Z' is




. In another



subembodiment, Ar''-Z' is

. In another subembodiment, Ar''-Z' is




. In a particular subembodiment of any of the above embodiments, R is H.


In a particular subembodiment of any of the above embodiments, Ar'' is phenyl.

In one embodiment, each L' is independently halo, C₁-C₆ alkyl, or C₁-C₆ haloalkyl. In a particular subembodiment Ar' has at least one L'. In a particular subembodiment Ar' is phenyl and is substituted with one or more L' groups wherein one L' is in the para position. In a particular embodiment, at least one L' is halo, for example fluoro, chloro, bromo, or iodo. In a particular subembodiment, at least two L' are halo and may be the same or different. In another embodiment, at least one L' is C₁-C₆ alkyl, for example methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, or hexyl. In another embodiment, at least one L' is C₁-C₆ haloalkyl, for example, trifluoromethyl.

In one embodiment, Ar' is unsubstituted. In another embodiment, k' is 1. In a subembodiment, when k' is 1 and Ar' is phenyl, L' is in the para position. In another embodiment, k' is 2. In a subembodiment, when k' is 2 and Ar' is phenyl, one L' is in the para position and one L' is in a meta position. In another embodiment, k' is 3. In another embodiment, k' is 4. In another embodiment, k' is 5.


In one embodiment, the compound is selected from the group consisting of:





Formula A

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula A or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof::



Formula A

wherein R^1 is H, F, Cl, Br, CF_3 , C_{1-6} alkyl, $C(O)CH_3$, $C(O)CO-(C_{1-6}$ alkyl), CH_2OH , CN, NH_2 , $N(C_{1-6}$ alkyl) $_2$, OH, $O-(C_{1-6}$ alkyl), OCF_3 , $S-(C_{1-6}$ alkyl), $SO_2-(C_{1-6}$ alkyl);

R^2 is H, F, Cl, methyl, CF_3 ;

R^3 is H, F, Cl, CH_3 , CF_3 , CN;

each of R^4 and $R^{4'}$ are independently selected from H or methyl;

each of R^5 and $R^{5'}$ can be H or OH, or R^5 and $R^{5'}$ can be taken together to form $=CH_2$ or $=O$;

R^6 is H or F;


X is H or F;

Y is OH, $NHSO_2R^7$, or $NHC(O)NHR^8$;


R^7 is C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

R^8 is H, C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

or X and Y are taken together to form a heterocycle wherein the moiety



of a compound of Formula A is selected from the group consisting of:



In one embodiment, C_{1-6} alkyl includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, cyclopropyl. C_{1-6} alkyl may also include tert-butyl, pentyl, cyclopentyl, hexyl, or cyclohexyl.

In one embodiment, R^1 is H. In one embodiment, R^1 is F. In one embodiment, R^1 is Cl. In another embodiment, R^1 is C_{1-6} alkyl, for example methyl or isopropyl. In one embodiment, R^1 is OH. In one embodiment, R^1 is CF_3 .

In one embodiment, R^2 is H. In one embodiment, R^2 is F. In one embodiment, R^2 is Cl. In another embodiment, R^2 is C_{1-6} alkyl, for example methyl. In one embodiment, R^2 is CF_3 .

In one embodiment, R^3 is H. In one embodiment, R^3 is F. In one embodiment, R^3 is Cl. In another embodiment, R^3 is C_{1-6} alkyl, for example methyl. In one embodiment, R^3 is CF_3 . In another embodiment, R^3 is CN.

In one embodiment, R^4 is H. In one embodiment, R^4 is methyl. In one embodiment, $R^{4'}$ is H. In one embodiment, $R^{4'}$ is methyl. In a particular embodiment, R^4 and $R^{4'}$ are both H. In another embodiment, one of R^4 and $R^{4'}$ is methyl.

In another embodiment, R^6 is H. In another embodiment, R^6 is F.

In another embodiment, X is H. In another embodiment, X is F.



In one embodiment, Y is OH. In one embodiment, Y is not OH. In one embodiment, Y is $NHSO_2R^7$. In another embodiment, Y is not $NHSO_2R^7$. In one embodiment, Y is $NHC(O)NHR^8$.

In one embodiment, R^7 is C_{1-6} alkyl, for example methyl.


In one embodiment, R^8 is H or C_{1-6} alkyl, for example methyl, ethyl or propyl.

In a particular subembodiment, X and Y are taken together to form a heterocycle


wherein the moiety




In a particular embodiment, the moiety . In




another embodiment, the moiety . In one




embodiment, the moiety . In one embodiment,




the moiety .



In one embodiment, the moiety . In




one embodiment, the moiety or



In one embodiment, the compound of Formula A, is selected from the compounds in Table 26, for example, the compound is selected from the group consisting of: NP10039, NP10165, NP10075, NP10153, NP10150, NP10146, NP10056, NP10122, NP10231, NP10002, NP10030, NP10070, NP10119, and NP10045.

Formula B

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula B or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



Formula B

wherein R¹ is H, F, Cl, Br, CF₃, or C₁₋₆ alkyl;

Z is O, S, NH, CH₂ or a bond;

R² is H or OH;

R⁶ is H or F;


X is H or F;

Y is OH, NHSO₂R⁷ or NHC(O)NHR⁸;



R⁷ is C₁₋₆ alkyl, C₆₋₁₂ aryl, or C₇₋₁₃ aralkyl;

R⁸ is H, C₁₋₆ alkyl, C₆₋₁₂ aryl, or C₇₋₁₃ aralkyl;


or X and Y are taken together to form a heterocycle wherein the moiety



of a compound of Formula B is selected from the group consisting of:



of a compound of Formula B is selected from the group consisting of:



In one embodiment, R^1 is H. In one embodiment, R^1 is not H. In one embodiment, R^1 is Cl. In another embodiment, R^1 is H or Cl. In one embodiment, R^1 is F, Cl or Br. In one embodiment, R^1 is CF_3 . In one embodiment, R^1 is C_{1-6} alkyl.

In one embodiment, Z is O. In another embodiment, Z is S. In another embodiment, Z is NH. In another embodiment, Z is CH_2 . In another embodiment, Z is a bond. In one embodiment, Z is not a bond. In another embodiment, Z is not CH_2 .

In one embodiment, R^2 is OH. In another embodiment, R^2 is H.

In another embodiment, R^6 is H. In another embodiment, R^6 is F.


In one embodiment, X is H. In a particular embodiment X is F.

In one embodiment, Y is OH. In one embodiment, Y is not OH. In one embodiment, Y is $NHSO_2R^7$. In another embodiment, Y is not $NHSO_2R^7$. In one embodiment, Y is $NHC(O)NHR^8$.


In one embodiment, R^7 is C_{1-6} alkyl, for example methyl.

In one embodiment, R^8 is H or C_{1-6} alkyl, for example methyl, ethyl or propyl.



In one embodiment, X and Y are taken together to form a heterocycle wherein the moiety



of a compound of Formula B is selected from the group consisting of:




In a particular embodiment, the moiety




. In


another embodiment, the moiety




is




or




. In one embodiment, the moiety



is not



or




.

In one embodiment, the compound is selected from the compounds in Table 26, for example compounds NP10250 and NP10185.

Formula C

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula C or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



Formula C

wherein each R^1 and R^2 is independently selected from H, F, Cl, Br, CF_3 , or C_{1-6} alkyl;
 R^6 is H or F;


X is H or F;

Y is OH, NHSO_2R^7 or NHC(O)NHR^8 ;


R^7 is C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

R^8 is H, C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

or X and Y are taken together to form a heterocycle wherein the moiety



of a compound of Formula C is selected from the group consisting of:



In one embodiment, R^1 is Cl. In one embodiment, R^1 is F. In one embodiment, R^1 is Br. In one embodiment, R^1 is H. In one embodiment, R^1 is not H. In one embodiment, R^1 is C_{1-6} alkyl, for example methyl.

In one embodiment, R^2 is Cl. In one embodiment, R^2 is F. In one embodiment, R^2 is Br. In one embodiment, R^2 is H. In one embodiment, R^2 is not H.

In one embodiment, R^6 is H. In another embodiment, R^6 is F.


In one embodiment, X is H. In a particular embodiment X is F.

In one embodiment, Y is OH. In one embodiment, Y is not OH. In one embodiment, Y is NHSO_2R^7 . In another embodiment, Y is not NHSO_2R^7 . In one embodiment, Y is NHC(O)NHR^8 .


In one embodiment, R^7 is C_{1-6} alkyl, for example methyl.

In one embodiment, R^8 is H or C_{1-6} alkyl, for example methyl, ethyl or propyl.

In a particular subembodiment, X and Y are taken together to form a heterocycle wherein the moiety



of a compound of Formula C is selected from the group consisting of:




In a particular embodiment, the moiety is . In

another embodiment, the moiety is . In one

embodiment, the moiety is . In one embodiment,


the moiety is .



In one embodiment, the compound is

Formula D-1

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula D-1 or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



Formula D-1

wherein each R^1 and R^2 is independently selected from H, F, Cl, Br, CF_3 , or C_{1-6} alkyl;

R^3 is H or OH;

R^6 is H or F;


X is H or F;

Y is OH, NH_2 , $N(R^8)_2$, $NHSO_2R^7$ or $NHC(O)NHR^8$;


R^7 is C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

each R^8 is independently selected from H, C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

or X and Y are taken together to form a heterocycle wherein the moiety



of a compound of Formula D-1 is selected from the group consisting of:



In one embodiment, R^1 is Cl. In one embodiment, R^1 is F. In one embodiment, R^1 is Br. In one embodiment, R^1 is H. In one embodiment, R^1 is not H. In one embodiment, R^1 is C_{1-6} alkyl, for example methyl.

In one embodiment, R^2 is Cl. In one embodiment, R^2 is F. In one embodiment, R^2 is Br. In one embodiment, R^2 is H. In one embodiment, R^2 is not H.

In one embodiment, one of R^1 and R^2 is Cl. In another embodiment, both of R^1 and R^2 are Cl.

In one embodiment, R^3 is H. In another embodiment, R^3 is OH.

In one embodiment, R^6 is H. In another embodiment, R^6 is F.


In one embodiment, X is H. In a particular embodiment, X is F.

In one embodiment, Y is OH. In one embodiment, Y is not OH. In one embodiment, Y is NH_2 . In one embodiment, Y is $N(R^8)_2$. In one embodiment, Y is $NHSO_2R^7$. In another embodiment, Y is not $NHSO_2R^7$. In one embodiment, Y is $NHC(O)NHR^8$.




In one embodiment, R^7 is C_{1-6} alkyl, for example methyl.

In one embodiment, R^8 is H or C_{1-6} alkyl, for example methyl, ethyl or propyl.

In a particular subembodiment, X and Y are taken together to form a heterocycle wherein the moiety





of a compound of Formula C is selected from the group consisting of:



,

and



.



In a particular embodiment, the moiety

is



. In



another embodiment, the moiety

is



. In one



embodiment, the moiety

is

. In one embodiment,




the moiety

is

.

Formula D-2

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula D-2 or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



Formula D-2

wherein R^1 is H, F, Cl, Br, CF_3 , or C_{1-6} alkyl;

Z^1 and Z^2 are each independently selected from the group consisting of $-CH_2-$ or $-C(=O)-$;

each of R^2 and $R^{2'}$ can be H or OH., or R^2 and $R^{2'}$ can be taken together to form $=CH_2$;

R^6 is H or F;


X is H or F;

Y is OH, NH_2 , $N(R^8)_2$, $NHSO_2R^7$ or $NHC(O)NHR^8$;


R^7 is C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

each R^8 is independently selected from H, C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

or X and Y are taken together to form a heterocycle wherein the moiety



of a compound of Formula D-2 is selected from the group consisting of:



In one embodiment, R^1 is Cl. In one embodiment, R^1 is F. In one embodiment, R^1 is Br. In one embodiment, R^1 is H. In one embodiment, R^1 is not H. In one embodiment, R^1 is C_{1-6} alkyl, for example methyl.

In one embodiment, R^2 is H. In one embodiment, R^2 is OH. In one embodiment, $R^{2'}$ is H. In one embodiment, $R^{2'}$ is OH. In one embodiment, one of R^2 and $R^{2'}$ is OH. In another embodiment, both of R^2 and $R^{2'}$ are H. In another embodiment, R^2 and $R^{2'}$ are taken together to form $=CH_2$.

In one embodiment, R^6 is H. In another embodiment, R^6 is F.


In one embodiment, X is H. In a particular embodiment, X is F.

In one embodiment, Y is OH. In one embodiment, Y is not OH. In one embodiment, Y is NH_2 . In one embodiment, Y is $N(R^8)_2$. In one embodiment, Y is $NHSO_2R^7$. In another embodiment, Y is not $NHSO_2R^7$. In one embodiment, Y is $NHC(O)NHR^8$.


In one embodiment, R^7 is C_{1-6} alkyl, for example methyl.

In one embodiment, R^8 is H or C_{1-6} alkyl, for example methyl, ethyl or propyl.


In a particular subembodiment, X and Y are taken together to form a heterocycle wherein the moiety




of a compound of Formula C is selected from the group consisting of:





,



, and




.




In a particular embodiment, the moiety

is

. In




is




. In one

another embodiment, the moiety




is




. In one embodiment,

embodiment, the moiety



is




the moiety

.

In one embodiment, the compound is selected from the compounds in Table 26, for example compounds NP10076 or NP10226.

Formula F

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of Formula F or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:



Formula F

wherein R^1 is H, F, Cl, Br, CF_3 , C_{1-6} alkyl, $C(O)CH_3$, $C(O)CO-(C_{1-6}$ alkyl), CH_2OH , CN, NH_2 , $N(C_{1-6}$ alkyl) $_2$, OH, O-(C_{1-6} alkyl), OCF_3 , S-(C_{1-6} alkyl), $SO_2-(C_{1-6}$ alkyl);

R^2 is H, F, Cl, methyl, CF_3 ;

R^3 is H, F, Cl, CH_3 , CF_3 , CN;

R^4 is H or methyl;

n is 0, 1 or 2;


R^6 is H or F;

X is H or F;


Y is OH, $NHSO_2R^7$, $NHC(S)NHR^8$ or $NHC(O)NHR^8$;

wherein R^7 or R^8 are each independently C_{1-6} alkyl, C_{6-12} aryl, C_{7-13} aralkyl;

or X and Y are taken together to form a heterocycle wherein the moiety



of a compound of Formula F is selected from the group consisting of:



In one embodiment, C₁₋₆ alkyl includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, cyclopropyl. C₁₋₆ alkyl may also include tert-butyl, pentyl, cyclopentyl, hexyl, or cyclohexyl.

In one embodiment, R¹ is F. In one embodiment, R¹ is Cl. In one embodiment, R¹ is Br. In a particular embodiment, R¹ is CF₃. In a particular embodiment, R¹ is C₁₋₆ alkyl, for example methyl. In one embodiment, R¹ is not H. In one embodiment, R¹ is F, Cl or methyl.

In another embodiment, R² is H. In one embodiment, R² is F. In one embodiment, R² is Cl.

In another embodiment, R³ is H.

In one embodiment, n is 0. In one embodiment, n is 1. In one embodiment, n is 2.


In one embodiment, R⁴ is H. In one embodiment, R⁴ is methyl. In one embodiment, R^{4'} is H. In one embodiment, R^{4'} is methyl. In a particular embodiment, R⁴ and R^{4'} are both H. In another embodiment, one of R⁴ and R^{4'} is methyl.



In another embodiment, R⁶ is H. In another embodiment, R⁶ is F.



In another embodiment, X is H. In another embodiment, X is F.



In one embodiment, Y is OH. In one embodiment, Y is not OH. In one embodiment, Y is NHSO₂R⁷. In another embodiment, Y is not NHSO₂R⁷. In one embodiment, Y is NHC(O)NHR⁸. In one embodiment, Y is NHC(S)NHR⁸.



In a particular subembodiment, X and Y are taken together to form a heterocycle wherein the moiety




In a particular embodiment, the moiety  is . In

another embodiment, the moiety  is . In one

embodiment, the moiety  is . In one embodiment,

the moiety  is .


In one embodiment, the compound is



Other Compound Embodiments

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of described in WO 02/072542 to Emory University, the entire disclosure of which is hereby incorporated by reference, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof.


In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound selected from the group consisting of





or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof.

In another embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound of described in WO 09/006437 to Emory University and NeurOp, Inc., or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof.

In one embodiment, methods of treatment or prophylaxis of neuropsychiatric disorders, in particular depression and anxiety are provided comprising administering a compound selected from the group consisting of







or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof.

Enantiomers

In certain embodiments, the compounds are provided as enantiomers. In one embodiment, the compound is provided as an enantiomer or mixture of enantiomers. In a particular embodiment, the compound is present as a racemic mixture. The enantiomer can be named by the configuration at the chiral center, such as R or S. In certain embodiments, the compound is present as a racemic mixture of R- and S- enantiomers. In certain embodiments, the compound is present as a mixture of two enantiomers. In one embodiment, the mixture has an enantiomeric excess in R. In one embodiment, the mixture has an enantiomeric excess in S. In certain other embodiments, the compound is in an enantiomeric excess of the R- or S- enantiomer. The enantiomeric excess can be 51% or more, such as 51% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, or 99% or more in the single enantiomer. The enantiomeric excess can be 51% or more, such as 51% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, or 99% or more in the R enantiomer. The enantiomeric excess can be 51% or more, such as 51% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, or 99% or more in the S enantiomer.

In other embodiments, the compound is substantially in the form of a single enantiomer. In some embodiments, the compound is present substantially in the form of the R enantiomer. In some embodiments, the compound is present substantially in the form of the

S enantiomer. The phrase “substantially in the form of a single enantiomer” is intended to mean at least 70% or more in the form of a single enantiomer, for example 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, or 99% or more in either the R or S enantiomer.

The enantiomer can be named by the direction in which it rotates the plane of polarized light. If it rotates the light clockwise as seen by the viewer towards whom the light is traveling, the isomer can be labeled (+) and if it rotates the light counterclockwise, the isomer can be labeled (-). In certain embodiments, the compound is present as a racemic mixture of (+) and (-) isomers. In certain embodiments, the compound is present as a mixture of two isomers. In one embodiment, the mixture has an excess in (+). In one embodiment, the mixture has an excess in (-). In certain other embodiments, the compound is in an excess of the (+) or (-) isomer. The isomeric excess can be 51% or more, such as 51% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, or 99% or more in the (+) isomer. The enantiomeric excess can be 51% or more, such as 51% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, or 99% or more in the (-) isomer.

In other embodiments, the compound is substantially in the form of a single optical isomer. In some embodiments, the compound is present substantially in the form of the (+) isomer. In other embodiments, the compound is present substantially in the form of the (-) isomer. The phrase “substantially in the form of a single optical isomer” is intended to mean at least 70% or more in the form of a single isomer, for example 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, or 99% or more of either the (+) or (-) isomer.

Methods of Use

In certain embodiments, the compounds are used for the treatment or prevention of neuropsychiatric disorders. The compounds of the invention can generally be administered to a host at risk of, or suffering from, a neuropsychiatric disorder related to NMDA receptor activation. Representative neuropsychiatric disorders include, without limitation, depression, anxiety, schizophrenia, bipolar disorder, obsessive-compulsive disorder, alcohol and substance abuse, and attention-deficit disorders such as ADH or ADHD. In particular embodiments, the disorders are neuropsychiatric mood disorders, non-limiting examples of which include depression, including major depression, bipolar disorders including cyclothymia (a mild form of bipolar disorder), affective disorders such as SAD (seasonal affective disorder) and mania (euphoric, hyperactive, over inflated ego, unrealistic optimism). In certain embodiments, a method of treatment a neuropsychiatric disorder is provided including administering a compound of the invention, alone or in combination to a host diagnosed with the disorder. Uses of the compounds in the treatment or manufacture of a medicament for such disorders are also provided.

In certain embodiments, the compounds are used for the treatment of depression in a host diagnosed with the disorder. Depression, formally called major depression, major depressive disorder or clinical depression, is a medical illness that involves the mind and body. Most health professionals today consider depression a chronic illness that requires long-term treatment, much like diabetes or high blood pressure. Although some people experience only one episode of depression, most have repeated episodes of depression symptoms throughout their life. Depression is also a common feature of mental illness, whatever its nature and origin. In some instances, the host or patient has a history of a major psychiatric disorder, such as schizophrenia. In other instances, the host does not have a history of a major psychiatric disorder but has been diagnosed with suffering from at least one depressive episode. In other instances, the host has been diagnosed with bipolar disorder. The host may also have been diagnosed as suffering from panic attacks or anxiety.

In one embodiment, the compounds of the present invention are used to diminish the severity of a depressive episode.

In some instances, the host is not suffering from a chronic disorder but is at risk of a depressive episode, anxiety or a panic attack due to environmental circumstances. The compounds may be given prophylactically to prevent onset of such an episode. For instance, in certain instances the compounds can be provided to a host before a plane trip, a public

speech, or other potential stressful even that could lead to an episode. In some embodiments, therefore, a method of prevention of a neuropsychiatric episode is provided, including administering a compound of the invention to a host at risk of suffering from such an episode.

In one embodiment, the compounds of the present invention are used to prevent a future depressive episode.

In certain embodiments, the compounds are administered to a host suffering from or at risk of suffering from age-related depression. The compounds can be administered prophylactically to a host over the age of 60, or over the age of 70, or over the age of 80 to prevent or reduce the severity of depressive episodes.

Depression is associated with physical illness as well. Chronic medical conditions associated with depression include heart disease, cancer, vitamin deficiencies, diabetes, hepatitis, and malaria. Depression also is a common effect of neurological disorders, including Parkinson's and Alzheimer's diseases, multiple sclerosis, strokes, and brain tumors. Even moderate depressive symptoms are associated with a higher than average rate of arteriosclerosis, heart attacks, and high blood pressure. Depression can mimic medical illness and any illness feels worse to someone suffering from depression. In certain other embodiments, the compounds are useful in the treatment or prophylaxis of a neuropsychiatric disorder associated with a medical condition, including but not limited to heart disease, cancer, vitamin deficiencies, diabetes, hepatitis, and malaria by administering the compound to a host suffering from the medical condition. In other instances, the compounds are useful in treatment or prophylaxis of a neuropsychiatric disorder associated with a neurological disorder or physiological insult by administering the compound to a host suffering from a neurological disorder or physiological insult. In non-limiting embodiments, these can include Parkinson's and Alzheimer's diseases, multiple sclerosis, strokes, and brain tumors. In some instances, the compounds are useful for treatment or prophylaxis of disorders such as depression or bipolar disorder associated with an injury or with aging. The compounds may also be useful in treatment or prophylaxis of schizophrenia.

In certain other embodiments, the compounds are used for treatment of a bipolar disorder in a host diagnosed with the disorder. The compounds can also be used to diminish the severity of manic episodes or prevent future episodes.

In certain embodiments, methods of treating seasonal disorders is provided including administering the compound to a host at risk of suffering from a SAD. In particular, the compounds can be provided on a seasonal basis. In some embodiments, the host has been diagnosed or is at risk of SAD.

In certain embodiments, the host is suffering from an attention deficit disorders such as ADH or ADHD.

Certain NMDA receptor antagonists described herein have enhanced activity in tissue having lower-than-normal pH. The tissue can be brain tissue. In certain embodiments, the reduced pH is associated with neuropsychiatric conditions. In some embodiments, the conditions can be associated with a physiological insult. In other embodiments, the conditions are mood disorders.

The compounds provided herein block the NR2B-containing NMDA receptors, have varying activity against receptors containing NR2A or NR2D, and may be selective for other members of the NMDA receptor family (NR2C, NR3A and NR3B). In one embodiment, the compounds are selective NMDA receptor blockers. General blocking of NMDA receptors throughout the brain causes adverse effects such as ataxia, memory deficits, hallucinations and other neurological problems. In one embodiment, the compounds are NMDA receptors antagonists selective for NR2B, NR2A, NR2C, NR2D, NR3A, and/or NR3B that do not interact with other receptors or ion channels at therapeutic concentrations. In one embodiment, the compound is a selective NR1/NR2A NMDA receptor and/or a NR1/NR2B NMDA receptor antagonist. In one particular embodiment, the compounds can bind to the NR2B subunit of the NMDA receptor. In another particular embodiment, the compounds are selective for the NR2B subunit of the NMDA receptor. In one embodiment, the compound is not an NMDA receptor glutamate site antagonist. In another embodiment, the compound is not an NMDA receptor glycine site antagonist.

In one embodiment, the compound does not exhibit substantial toxic side effects, such as, for example, motor impairment or cognitive impairment. In a particular embodiment, the compound has a therapeutic index equal to or greater than at least 2. In another embodiment, the compound is at least 10 times more selective for binding to an NMDA receptor than any other glutamate receptor.

Further, compounds selected according to the methods or processes described herein can be used prophylactically to prevent or protect against such diseases or neurological conditions, such as those described herein. In one embodiment, patients with a predisposition for a neuropsychiatric disorder, in particular a mood disorder, such as a genetic predisposition, can be treated prophylactically with the methods and compounds described herein.

Pharmaceutical Compositions

Mammals, and specifically humans, suffering from or at risk of neuropsychiatric disorders can be treated by either targeted or systemic administration, via oral, inhalation, topical, trans- or sub-mucosal, subcutaneous, parenteral, intramuscular, intravenous or transdermal administration of a composition comprising an effective amount of the compounds described herein or a pharmaceutically acceptable salt, ester or prodrug thereof, optionally in a pharmaceutically acceptable carrier.

The compounds or composition is typically administered by oral administration. Alternatively, compounds can be administered by inhalation. In another embodiment, the compound is administered transdermally (for example via a slow release patch), or topically. In yet another embodiment, the compound is administered subcutaneously, intravenously, intraperitoneally, intramuscularly, parenterally, or submucosally. In any of these embodiments, the compound is administered in an effective dosage range to treat the target condition.

In one embodiment, compounds of the present invention are administered orally. Oral compositions will generally include an inert diluent or an edible carrier. They may be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition.

When the compound is administered orally in the form of a dosage unit such as a tablets, pills, capsules, troches and the like, these can contain any of the following ingredients, or compounds of a similar nature: a binder (such as microcrystalline cellulose, gum tragacanth or gelatin); an excipient (such as starch or lactose), a disintegrating agent (such as alginic acid, Primogel, or corn starch); a lubricant (such as magnesium stearate or Sterotes); a glidant (such as colloidal silicon dioxide); a sweetening agent (such as sucrose or saccharin); and/or a flavoring agent (such as peppermint, methyl salicylate, or orange flavoring). When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier (such as a fatty oil). In addition, dosage unit forms can contain various other materials which modify the physical form of the dosage unit, for example, coatings of sugar, shellac, or other enteric agents.

The compound or its salts can also be administered orally as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the

active compounds, a sweetening agent (such as sucrose, saccharine, etc.) and preservatives, dyes and colorings and flavors.

The compounds of the invention may be also administered in specific, measured amounts in the form of an aqueous suspension by use of a pump spray bottle. The aqueous suspension compositions of the present invention may be prepared by admixing the compounds with water and other pharmaceutically acceptable excipients. The aqueous suspension compositions according to the present invention may contain, inter alia, water, auxiliaries and/or one or more of the excipients, such as: suspending agents, e.g., microcrystalline cellulose, sodium carboxymethylcellulose, hydroxypropyl-methyl cellulose; humectants, e.g. glycerin and propylene glycol; acids, bases or buffer substances for adjusting the pH, e.g., citric acid, sodium citrate, phosphoric acid, sodium phosphate as well as mixtures of citrate and phosphate buffers; surfactants, e.g. Polysorbate 80; and antimicrobial preservatives, e.g., benzalkonium chloride, phenylethyl alcohol and potassium sorbate.

In a separate embodiment, the compounds of the invention are in the form of an inhaled dosage. In this embodiment, the compounds may be in the form of an aerosol suspension, a dry powder or liquid particle form. The compounds may be prepared for delivery as a nasal spray or in an inhaler, such as a metered dose inhaler. Pressurized metered-dose inhalers ("MDI") generally deliver aerosolized particles suspended in chlorofluorocarbon propellants such as CFC-11, CFC-12, or the non-chlorofluorocarbons or alternate propellants such as the fluorocarbons, HFC-134A or HFC-227 with or without surfactants and suitable bridging agents. Dry-powder inhalers can also be used, either breath activated or delivered by air or gas pressure such as the dry-powder inhaler disclosed in the Schering Corporation International Patent Application No. PCT/US92/05225, published 7 Jan. 1993 as well as the Turbuhaler™ (available from Astra Pharmaceutical Products, Inc.) or the Rotahaler™ (available from Allen & Hanburys) which may be used to deliver the aerosolized particles as a finely milled powder in large aggregates either alone or in combination with some pharmaceutically acceptable carrier e.g. lactose; and nebulizers.

Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include at least some of the following components: a sterile diluent (such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents); antibacterial agents (such as benzyl alcohol or methyl parabens); antioxidants (such as ascorbic acid or sodium bisulfite); chelating agents (such as ethylenediaminetetraacetic acid); buffers (such as acetates, citrates or phosphates); and/or agents for the adjustment of tonicity (such as sodium chloride or dextrose). The pH of the

solution or suspension can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide.

A parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Suitable vehicles or carriers for topical application can be prepared by conventional techniques, such as lotions, suspensions, ointments, creams, gels, tinctures, sprays, powders, pastes, slow-release transdermal patches, suppositories for application to rectal, vaginal, nasal or oral mucosa. In addition to the other materials listed above for systemic administration, thickening agents, emollients, and stabilizers can be used to prepare topical compositions. Examples of thickening agents include petrolatum, beeswax, xanthan gum, or polyethylene, humectants such as sorbitol, emollients such as mineral oil, lanolin and its derivatives, or squalene.

If administered intravenously, carriers can be physiological saline, bacteriostatic water, Cremophor ELTM (BASF, Parsippany, NJ) or phosphate buffered saline (PBS).

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) are also preferred as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811 (which is incorporated herein by reference in its entirety). For example, liposome formulations may be prepared by dissolving appropriate lipid(s) (such as stearyl phosphatidyl ethanolamine, stearyl phosphatidyl choline, arachadoyl phosphatidyl choline, and cholesterol) in an inorganic solvent that is then evaporated, leaving behind a thin film of dried lipid on the surface of the container. An aqueous solution of the compound is then introduced into the container. The container is then swirled by hand to free lipid material from the sides of the container and to disperse lipid aggregates, thereby forming the liposomal suspension.

Dosing

The compound is administered for a sufficient time period to alleviate the undesired symptoms and the clinical signs associated with the condition being treated. In one embodiment, the compounds are administered less than three times daily. In one embodiment, the compounds are administered in one or two doses daily. In one embodiment, the compounds are administered once daily. In some embodiments, the compounds are administered in a single oral dosage once a day.

The active compound is included in the pharmaceutically acceptable carrier or diluent in an amount sufficient to deliver to a patient a therapeutic amount of compound *in vivo* in the absence of serious toxic effects. An effective dose can be readily determined by the use of conventional techniques and by observing results obtained under analogous circumstances. In determining the effective dose, a number of factors are considered including, but not limited to: the species of patient; its size, age, and general health; the specific disease involved; the degree of involvement or the severity of the disease; the response of the individual patient; the particular compound administered; the mode of administration; the bioavailability characteristics of the preparation administered; the dose regimen selected; and the use of concomitant medication.

Typical systemic dosages for the herein described conditions are those ranging from 0.01 mg/kg to 1500 mg/kg of body weight per day as a single daily dose or divided daily doses. Preferred dosages for the described conditions range from 0.5-1500 mg per day. A more particularly preferred dosage for the desired conditions ranges from 5-750 mg per day. Typical dosages can also range from 0.01 to 1500, 0.02 to 1000, 0.2 to 500, 0.02 to 200, 0.05 to 100, 0.05 to 50, 0.075 to 50, 0.1 to 50, 0.5 to 50, 1 to 50, 2 to 50, 5 to 50, 10 to 50, 25 to 50, 25 to 75, 25 to 100, 100 to 150, or 150 or more mg/kg/day, as a single daily dose or divided daily doses. In one embodiment, the daily dose is between 10 and 500 mg/day. In another embodiment, the dose is between about 10 and 400 mg/day, or between about 10 and 300 mg/day, or between about 20 and 300 mg/day, or between about 30 and 300 mg/day, or between about 40 and 300 mg/day, or between about 50 and 300 mg/day, or between about 60 and 300 mg/day, or between about 70 and 300 mg/day, or between about 80 and 300 mg/day, or between about 90 and 300 mg/day, or between about 100 and 300 mg/day, or about 200 mg/day. In one embodiment, the compounds are given in doses of between about 1 to about 5, about 5 to about 10, about 10 to about 25 or about 25 to about 50 mg/kg. Typical dosages for topical application are those ranging from 0.001 to 100% by weight of the active compound.

The concentration of active compound in the drug composition will depend on absorption, inactivation, and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition. The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at varying intervals of time.

Combination Treatment

The compound can also be mixed with other active materials which do not impair the desired action, or with materials that supplement the desired action. The active compounds can be administered in conjunction, i.e. combination or alternation, with other medications used in the treatment or prevention neuropsychiatric disorders, such as those in which NMDA receptor activation is involved. In certain embodiments, the combination can be synergistic.

In certain embodiments, the compound is administered in combination or alternation with a compound useful for treatment of neuropsychiatric disorders, such as a selective serotonin reuptake inhibitor (SSRI), a serotonin and norepinephrine reuptake inhibitor (SNRI), norepinephrine and dopamine reuptake inhibitor (NDRI), combined reuptake inhibitor and receptor blocker, tetracyclic antidepressant, tricyclic antidepressants (TCAs) (although TCAs tend to have numerous and severe side effects), or a monoamine oxidase inhibitor (MAOI).

Electroconvulsive therapy (ECT) can also be used to treat depression in conjunction with administration of a compound of the invention. Nontraditional treatment options include vagus nerve stimulation, transcranial magnetic stimulation and deep brain stimulation.

SSRIs include fluoxetine (Prozac, Sarafem), paroxetine (Paxil), sertraline (Zoloft), citalopram (Celexa) and escitalopram (Lexapro). SSRIs that have been approved by the Food and Drug Administration specifically to treat depression are: Citalopram (Celexa), Escitalopram (Lexapro), Fluoxetine (Prozac, Prozac Weekly), Paroxetine (Paxil, Paxil CR) and Sertraline (Zoloft). SNRIs that have been approved by the Food and Drug Administration specifically to treat depression are: Duloxetine (Cymbalta) and Venlafaxine

(Effexor, Effexor XR). The only NDRI that has been approved by the Food and Drug Administration specifically to treat depression is Bupropion (Wellbutrin, Wellbutrin SR, Wellbutrin XL). The only tetracyclic antidepressant that has been approved by the Food and Drug Administration specifically to treat depression is Mirtazapine (Remeron, Remeron SolTab). Other compounds approved for treatment of neuropsychiatric disorders include Anafranil (clomipramine HCl); Aventyl (nortriptyline HCl); Desyrel (trazodone HCl); Elavil (amitriptyline HCl); Limbitrol (chlordiazepoxide/amitriptyline); Ludiomil (Maprotiline HCl); Luvox (fluvoxamine maleate); Marplan (isocarboxazid); Nardil (phenelzine sulfate); Norpramin (desipramine HCl); Pamelor (nortriptyline HCl); Parnate (tranylcypromine sulfate); Pexeva (paroxetine mesylate); Prozac (fluoxetine HCl); Sarafem (fluoxetine HCl); Serzone (nefazodone HCl); Sinequan (doxepin HCl); Surmontil (trimipramine); Symbyax (olanzapine/fluoxetine); Tofranil (imipramine HCl); Tofranil-PM (imipramine pamoate); Triavil (Perphenazine/Amitriptyline); Vivactil (protriptyline HCl); Wellbutrin (bupropion HCl); and Zyban (bupropion HCl). Combined inhibitors and blockers that have been approved by the Food and Drug Administration specifically to treat depression are: Trazodone, Nefazodone and Maprotiline.

Tricyclic antidepressants (TCAs) inhibit the reabsorption (reuptake) of serotonin and norepinephrine. They were among the earliest of antidepressants, hitting the market in the 1960s, and they remained the first line of treatment for depression through the 1980s, before newer antidepressants arrived. TCAs that have been approved by the Food and Drug Administration specifically to treat depression are: Amitriptyline, Amoxapine, Desipramine (Norpramin), Doxepin (Sinequan), Imipramine (Tofranil), Nortriptyline (Pamelor), Protriptyline (Vivactil) and Trimipramine (Surmontil).

MAOIs that have been specifically approved by the Food and Drug Administration to treat depression are: Phenelzine (Nardil), Tranylcypromine (Parnate), Isocarboxazid (Marplan) and Selegiline (Emsam). Emsam is the first skin (transdermal) patch for depression.

Any of the compounds of the invention can be administered in combination with another active agent. In certain embodiments, the second active is one that is effective in treatment of a neuropsychiatric disorder. However, in certain other embodiments, the second active is one that is effective against an underlying disorder that is associated with a neuropsychiatric symptom. Examples of such disorders are heart disease, Alzheimer's disease and Parkinson's diseases. In certain embodiments, the compounds can be administered in combination in a single dosage form or injection, or administered

concurrently. In other embodiments, the compounds are administered in alternation.

Side Effects

In an additional aspect of the methods and processes described herein, the compound does not exhibit substantial toxic an/or psychotic side effects. Toxic side effects include, but are not limited to: agitation, hallucination, confusion, stupor, paranoia, delirium, psychotomimetic-like symptoms, rotarod impairment, amphetamine-like stereotyped behaviors, stereotypy, psychosis memory impairment, motor impairment, anxiolytic-like effects, increased blood pressure, decreased blood pressure, increased pulse, decreased pulse, hematological abnormalities, electrocardiogram (ECG) abnormalities, cardiac toxicity, heart palpitations, motor stimulation, psychomotor performance, mood changes, short-term memory deficits, long-term memory deficits, arousal, sedation, extrapyramidal side-effects, ventricular tachycardia. Lengthening of cardiac repolarisation, ataxia, cognitive deficits and/or schizophrenia-like symptoms.

Further, in another embodiment, the compounds selected or identified according to the processes and methods described herein do not have substantial side effects associated with other classes of NMDA receptor antagonists. In one embodiment, such compounds do not substantially exhibit the side effects associated with NMDA antagonists of the glutamate site, such as selfotel, D-CPPene (SDZ EAA 494) and AR-R15896AR (ARL 15896AR), including, agitation, hallucination, confusion and stupor (Davis et al. (2000) *Stroke* 31(2):347-354; Diener et al. (2002), *J Neurol* 249(5):561-568); paranoia and delirium (Grotta et al. (1995), *J Intern Med* 237:89-94); psychotomimetic-like symptoms (Loscher et al. (1998), *Neurosci Lett* 240(1):33-36); poor therapeutic ratio (Dawson et al. (2001), *Brain Res* 892(2):344-350); amphetamine-like stereotyped behaviors (Potschka et al. (1999), *Eur J Pharmacol* 374(2):175-187). In another embodiment, such compounds do not exhibit the side effects associated with NMDA antagonists of the glycine site, such as HA-966, L-701,324, d-cycloserine, CGP-40116, and ACEA 1021, including significant memory impairment and motor impairment (Wlaz, P (1998), *Brain Res Bull* 46(6):535-540). In a still further embodiment, such compounds do not exhibit the side effects of NMDA high affinity receptor channel blockers, such as MK-801 and ketamine, including, psychosis-like effects (Hoffman, D C (1992), *J Neural Transm Gen Sect* 89:1-10); cognitive deficits (decrements in free recall, recognition memory, and attention; Malhotra et al (1996), *Neuropsychopharmacology* 14:301-307); schizophrenia-like symptoms (Krystal et al (1994), *Arch Gen Psychiatry* 51:199-214; Lahti et al. (2001), *Neuropsychopharmacology* 25:455-467), and hyperactivity

and increased stereotpy (Ford et al (1989) *Physiology and behavior* 46: 755-758.

In a further additional or alternative embodiment, the compound has a therapeutic index equal to or greater than at least 2:1, at least 3:1, at least 4:1, at least 5:1, at least 6:1, at least 7:1, at least 8:1, at least 9:1, at least 10:1, at least 15:1, at least 20:1, at least 25:1, at least 30:1, at least 40:1, at least 50:1, at least 75:1, at least 100:1 or at least 1000:1. The therapeutic index can be defined as the ratio of the dose required to produce toxic or lethal effects to dose required to produce therapeutic responses. It can be the ratio between the the median toxic dose (the dosage at which 50% of the group exhibits the adverse effect of the drug) and the median effective dose (the dosage at which 50% of the population respond to the drug in a specific manner). The higher the therapeutic index, the more safe the drug is considered to be. It simply indicates that it would take a higher dose to invoke a toxic response that it does to cause a beneficial effect.

The side effect profile of compounds can be determined by any method known to those skilled in the art. In one embodiment, motor impairment can be measured by, for example, measuring locomotor activity and/or rotorod performance. Rotorod experiments involve measuring the duration that an animal can remain on an accelerating rod. In another embodiment, memory impairment can be assessed, for example, by using a passive avoidance paradigm; Sternberg memory scanning and paired words for short-term memory, or delayed free recall of pictures for long-term memory. In a further embodiment, anxiolytic-like effects can be measured, for example, in the elevated plus maze task. In other embodiments, cardiac function can be monitored, blood pressure and/or body temperature measured and/or electrocardiograms conducted to test for side effects. In other embodiments, psychomotor functions and arousal can be measured, for example by analyzing critical flicker fusion threshold, choice reaction time, and/or body sway. In other embodiments, mood can be assessed using, for example, self-ratings. In further embodiments, schizophrenic symptoms can be evaluated, for example, using the PANSS, BPRS, and CGI, side-effects were assessed by the HAS and the S/A scale.

EXAMPLES


The following examples are provided to illustrate the present invention and are not intended to limit the scope thereof. Those skilled in the art will readily understand that kown variations of the conditions and processes of the following preparative procedures can be used to manufacture the desired compounds. The materials required for the embodiments and

the examples are known in the literature, readily commercially available, or can be made by known methods from the known starting materials by those skilled in the art.

Synthesis of Compounds

The compounds for use in the methods described herein can be prepared by any methods known in the art, such as in accordance with the methods and general synthetic strategies provided in WO 02/072542 or WO 09/006437, or by the following synthetic methods, or variations of those procedures readily understood to those skilled in the art.

Examples 1 and 2. N-(4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-(*S*)-hydroxy-propoxy}-phenyl)-methanesulfonamide (Compound 1) and N-(4-{3-[2-(3,4-Dichloro-phenylamino)-ethylamino]-2-(*S*)-hydroxy-propoxy}-phenyl)-methanesulfonamide (Compound 2).



Step (i). 3-(4-Nitro-phenoxy)-2-(*S*)-propyleneoxide (**i-1**). 4-Nitrophenol (6.6 mmol) was dissolved in 5 ml anhydrous DMF. Cesium fluoride (19.9 mmol) was added to the reaction. The reaction mixture was stirred for 1 hour at room temperature and (S)-Glycidyl nosylate (6.6 mmol) was added to the reaction mixture. The reaction stirred for 24 hours at room temperature. Water (150 mL) was added and the solution was extracted with ethyl acetate.

The organic phase was dried over MgSO_4 and evaporated. The residue was purified with column chromatograph using ethylacetate: hexane (50:50) solvent system to give the desired product **i-1**. This step can be substituted with (*R*)-Glycidyl nosylate to get the *R* isomer.








Step (ii). 3-(4-Amino-phenoxy)-2-(*S*)-propyleneoxide (**i-2**). (*S*)-Glycidyl-4-nitrophenyl ether (2.6 mmol, **i-1**) and 5% Pd/C(en) [Sajiki et al, Chemistry- a european journal 6(12):2200-2204 (2000).] (10% of the weight of starting material) in 5 ml anhydrous THF was hydrogenated at ambient pressure and temperature for 3 hours. The reaction mixture was filtered by using membrane filter (13, 0.22 μm) and the filtrate was concentrated in vacuum. The compound was afforded as a crude mixture of amino reduction compound **i-2**.





Step (iii). 3-(4-methansufonylamido-phenoxy)-2-(*S*)-propyleneoxide (**i-3**). (*S*)-Glycidyl-4-aminophenyl ether (2.4 mmol, **i-2**) dissolved in 20 ml anhydrous DCM and *N,N*-diisopropyl-*N*-ethylamine (2.6 mmol) was added at 0°C. After stirring 15 minutes methanesulfonyl chloride (2.6 mmol) was added drop wise to the reaction mixture at 0°C. After stirring over night, the reaction extracted with water and washed with brine. Organic phase dried over magnesium sulfate and evaporated. The residue was purified with flash chromatography using Ethyl acetate: DCM (30:70) solvent system to give the desired product **i-3**.

Step (iv). N-(4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-(*S*)-hydroxy-propoxy}-phenyl)-methanesulfonamide (Compound 1). Compound **i-3** (2.00 mmol) and *N*-(3,4-Difluorophenyl)piperazine (2.00 mmol) were heated under reflux conditions in 20 ml ethanol for 8 hours. Then solvent was evaporated and residue was purified with flash chromatography using dichloromethane:methanol (90:10) solvent system to get compound 1. Compound 1 was dissolved in ethanol and bubbled HCl gas to get the HCl salt of the compound 1.


Step (v). N-(4-{3-[2-(3,4-Dichloro-phenylamino)-ethylamino]-2-(*S*)-hydroxy-propoxy}-phenyl)-methanesulfonamide (Compound 2). The epoxide (**i-3**, 1.58 mmol) was dissolved in EtOH (20 ml), and then the 3,4-dicholoro-ethylene diamine (1.58 mmol) (preparation: Isabel Perillo, M. Cristina Caterina, Julieta López, Alejandra Salerno. Synthesis 2004, 6, 851-856) was added and the solution refluxed for 16 hours. The solvent was evaporated and the product purified with column chromatography using 10% MeOH/DCM + 1% NH_4OH to give compound 2.

The following compounds were synthesized according to the procedures provided in examples 1 and 2.

COMPOUND	NAME AND PHYSICAL DATA
	N-(4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide MS: 475.14486
	N-(4-{3-[4-(4-Chloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide MS:440.14017
	N-(4-{3-[4-phenyl-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide MS:406.20183
	N-(4-{3-[4-(4-Hydroxy-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(2-Pyridyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(4-Pyridyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide
	N-{4-[2-(<i>S</i>)-Hydroxy-3-(2-phenylamino-ethylamino)-propoxy]-phenyl}-methanesulfonamide

	N-{4-[2-(<i>S</i>)-Hydroxy-3-(2-(3,4-difluorophenyl)amino-ethylamino)-propoxy]-phenyl}-methanesulfonamide MS:416.15842
	N-(4-{3-[3-(3,4-Dichloro-phenyl)-allylamino]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide MS:446.09479
	N-[4-(3-{Butyl-[3-(3,4-dichloro-phenyl)-allyl]-amino}-2-(<i>S</i>)-hydroxy-propoxy)-phenyl]-methanesulfonamide MS:501.13871
	N-(4-{3-[3-(3,4-Difluoro-phenyl)-allylamino]-2-(<i>S</i>)-hydroxy-propoxy}-phenyl)-methanesulfonamide MS:413.58425

Example 3. 6-{3-[4-(4-Chloro-phenyl)-piperazin-1-yl]-2-(*S*)-hydroxy-propoxy}-3H-benzooxazol-2-one (Compound 3).






Step (i). 6-(2-(*S*)-Oxiranylmethoxy)-3H-benzooxazol-2-one (**ii-1**). 5-hydroxy-benzoxazole (310 mg) and cesium carbonate (780 mg) were combined in 6 mL of N,N-dimethylformamide. The reaction was stirred for room temperature for 1 hour. (*S*)-glycidal nosylate (520 mg) was added, and the reaction stirred at room temperature overnight. The reaction was quenched with NH₄Cl(aq) solution and extracted with ethyl acetate. The organic layer was washed with NH₄Cl(aq) and NaCl(aq) solutions, separated, and dried over



Na₂SO₄(s). Filtration and solvent removal was followed by absorption onto silica gel.

Elution with an ethyl acetate/methanol mixture (4:1) followed by solvent removal gave 445 mg of a yellow, oily solid.


Step (ii). 6-{3-[4-(4-Chloro-phenyl)-piperazin-1-yl]-2-(*S*)-hydroxy-propoxy}-3H-benzooxazol-2-one (Compound 3). To a solution of 300 mg of epoxide (ii-1) in 10 mL of absolute ethanol was added 300 mg of 4-(4-chlorophenyl)-piperazine. The solution was heated to 70°C for 8 hours. The reaction was cooled and the solvent removed under vacuum. The residue was purified by column chromatography on silica gel using ethyl acetate as solvent. Obtained 240 mg of a light brown solid (45% yield). ¹H NMR (d₆-DMSO, 400 MHz): δ 2.37 (dq, 2H, J=6Hz, J=13Hz), 2.51 (m, 4H), 3.02 (m, 4H), 3.68 (q, 1H, J=8Hz), 3.84 (dd, 1H, J=4Hz, J=14Hz), 4.02 (bs, 1H), 5.07 (d, 1H, J=5Hz), 6.61 (dd, 1H, J=2Hz, J=9Hz), 6.73 (d, 1H, J=2Hz), 6.91 (d, 2H, J=9Hz), 7.05 (d, 1H, J=8Hz), 7.21 (d, 2H, J=9Hz), 9.43 (s, 1H); MS (m/z): 404 (M+H), 406 (M+2+H); HRMS Calcd. for C₂₀H₂₃ClN₃O₄: 404.13771; Found: 404.13673.

The following compounds were synthesized according to the procedure in Example 3.

COMPOUND	NAME AND PHYSICAL DATA
	6-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one MS: 406.15664
	6-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one
	6-{3-[4-(4-Methyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one MS: 384.19077

	6-{3-[2-(3,4-Dichloro-phenyl)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one
	6-{3-[2-(4-Chloro-phenylamino)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-3H-benzooxazol-2-one MS: 378.12089

Example 4. 4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(*S*)-hydroxy-propoxy}-phenol (Compound 4).






Step (i). 3-(4-tert-Butyldimethylsilyloxy-phenoxy)-2-(*S*)-propyleneoxide (**iii-1**). 4-(tert-Butyldimethylsilyloxy)phenol (1.45 g, 6.25 mmol) in 5 ml anhydrous THF was added dropwise to the suspension of NaH (0.158 g, 6.25 mmol) in 5 ml THF. After stirring at room temperature for 2 hours glycidyl nosylate (1.30 g, 5 mmol) and then 15-crown-5 (25 mol%) were added to the reaction mixture. After stirring 24 hours reaction was poured to ice-water and extracted with ethyl acetate. Organic phase was washed with water and brine, then dried over sodium sulfate and evaporated. Product was purified by column chromatography using EtOAc: Hexane (1:9) (yield: 1.06 g 76%). ¹H-NMR (400MHz, CDCl₃) δ 0.17 (6H, s), 0.98









(9H, s), 2.75 (1H, dd, J = 2.4, 4.4 Hz), 2.89 (1H, q, J = 4.4Hz), 3.33-3.36 (1H, m), 3.90 (1H, dd, J = 5.6, 10.8 Hz), 4.16 (1H, dd, J = 3.6, 11.2 Hz), 6.69-6.81 (4H, m).









Step (ii). 4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(*S*)-hydroxy-propoxy}-phenoxy-tert-butyldimethyl silane (**iii-2**). Compound **iii-1** (0.280 g, 1 mmol) and 1-(4-chlorophenyl)piperazine (0.200 g, 1 mmol) were dissolved in 5 ml EtOH and refluxed for 90 minutes. Solvent was evaporated and the material was used in the next step without purification.









Step (iii). 4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(*S*)-hydroxy-propoxy}-phenol (Compound **4**). Compound **iii-2** was dissolved in 5 ml THF and 2 ml TBAF in 1.0M THF solution was added, and stirred for 2 hours. Quenched with ammonium chloride solution, extracted with EtOAc. Organic phase was dried over sodium sulfate and evaporated. Product was purified using column chromatography using EtOAc:MeOH (95:5). $^1\text{H-NMR}$ (400MHz, DMSO- d_6) δ 2.36-2.61 (6H, m), 3.11 (4H, t, J = 4.8 Hz), 3.76 (1H, dd, J = 4.0, 6.0 Hz), 3.86(1H, dd, J = 4.4, 10.0 Hz), 3.91-3.95 (1H, m), 4.85 (1H, d, J = 4.8 Hz), 6.66 (1H, dd, J = 2.4, 6.8 Hz), 6.75 (1H, dd, J = 2.4, 6.8 Hz), 6.92 (1H, dd, J = 2.4, 6.8 Hz), 7.21 (1H, dd, J = 2.4, 6.8 Hz), 8.90 (1H, s). HRMS: 362.1397 calculated. 362.14696 found.

The following compounds were synthesized according to Example 4.


COMPOUND	NAME AND PHYSICAL DATA
	4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:397.10811
	4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:365.16672
	4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-(<i>R</i>)-hydroxy-propoxy}-phenol MS: 365.16657

	4-{3-[4-(4-Fluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:347.17602
	4-{3-[4-(3,4-Dimethyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:357.21716
	4-{3-[4-(4-Methyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:343.20093
	4-{3-[4-(4-Cyano-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:354.18073
	4-{3-[4-(4-Bromo-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:407.09663
	4-{3-[4-(4-Hydroxy-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:345.18061
	4-{3-[4-(4-Methoxy-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:359.19608
	4-{3-[4-(4-Trifluoromethyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:397.17327

	4-{3-[4-(4-Biphenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS: 405.21643
	4-{3-[4-(2,4-Difluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:365.16651
	4-{3-[4-(2-Fluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:347.17595
	4-{3-[4-(2-Chloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS: 363.14695
	4-{3-[4-(2-Chloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:343.20108
	4-{3-[4-(2-Cyano-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:354.18070
	4-{3-[4-Phenyl-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:329.18547
	4-{3-[4-(3-Fluoro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:347.17608

	4-{3-[4-(3-Chloro-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:363.14741
	4-{3-[4-(3-Methyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:343.20109
	4-{3-[4-(3-Trifluoromethyl-phenyl)-piperazin-1-yl]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:397.17269
	4-{3-[2-(3,4-Dichloro-phenyl)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-phenol MS:357.10159
	4-(3-{Butyl-[2-(3,4-dichloro-phenyl)-ethyl]-amino}-2-(<i>S</i>)-hydroxy-propoxy)-phenol MS:413.12769
	4-{3-[2-(3,4-Dichloro-phenyl)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-3-fluoro-phenol MS:375.03418
	4-{3-[2-(3,4-Dichloro-phenyl)-ethylamino]-2-(<i>S</i>)-hydroxy-propoxy}-2-fluoro-phenol MS:375.03409
	1-[2-(<i>S</i>)-Hydroxy-3-(4-hydroxy-phenoxy)-propyl]-4-phenyl-piperidin-4-ol MS:344.18509

Example 5. (4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-urea (Compound 5).



Step (i). [4-(3-Bromo-propoxy)-phenyl]-carbamic acid tert-butyl ester (**iv-1**). To a solution of 2.1g of 4-t-butylcarbonylamino-phenol in 20 mL of acetonitrile was added 3.25g of cesium carbonate. The reaction was stirred for one hour, and then 1.5 mL of 1,3-dibromopropane was added and the reaction stirred for 20 hours. The reaction was then quenched with $\text{NH}_4\text{Cl(aq.)}$ solution. The mixture was extracted with ethyl acetate and washed with $\text{NH}_4\text{Cl(aq.)}$ and NaCl(aq.) solutions. The organic layer was separated and dried over $\text{Na}_2\text{SO}_4\text{(s)}$. Filtration and solvent removal gave a light brown oily solid. Hexanes were added and the resulting solids filtered and washed with Hexanes three times. Drying gave 2.4g of an off-white solid.


Step (ii). (4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-carbamic acid tert-butyl ester (**iv-2**). To 305 mg of 4-(3,4Difluoro-phenyl)-piperazine and 335 mg of compound **iv-1** was added 5 mL of acetonitrile. The reaction was heated to 65°C overnight. The reaction was cooled, and then extracted with ethyl acetate. The organic layers were washed with $\text{NaHCO}_3\text{(aq.)}$ twice, and the organic layers separated and dried over $\text{Na}_2\text{SO}_4\text{(s)}$. Filtration and solvent removal gave an light brown solid. Dilution with hexanes, filtration,





and washing with hexanes gave 458 mg of a white solid (**iv-2**). MS (m/z): 430 (M+H); HRMS: Obsd for C₂₄H₃₃FN₃O₃: 430.24951.

Step (iii). Compound **iv-2** (430 mg) was dissolved in 6 mL of dichloromethane. Next, 4 mL of trifluoroacetic acid was added and the reaction was stirred for 6 hours. Then NaHCO₃(s) was added until the bubbling stopped. Then water was added to the reaction mixture and the reaction was extracted with dichloromethane and washed with NaHCO₃(aq.) twice. The organics were dried over Na₂SO₄(s), and then the solution was filtered and the solvent removed under vacuum. The residue was used in the next step without any purification.


Step (iv). (4-{3-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-urea (Compound **5**). The aniline from the previous step was dissolved in 10 mL of N,N-dimethyl formamide. Next, 1 mL of trimethylsilyl isocyanate was added, and the reaction was stirred at room temperature overnight. The reaction was then quenched with NaHCO₃(aq.) solution. The reaction was extracted with ethyl acetate and washed with NaHCO₃(aq.) solution twice. The organic layer was separated and dried over Na₂SO₄(s). Filtration and solvent removal gave a brown solid. Filtration over a plug of silica gel with ethyl acetate/methanol (4:1) was followed by solvent removal. Trituration of the resulting solids with ethyl ether and filtration gave 98 mg of an off-white solid. MS (m/z): 391 (M+H); HRMS: calcd. for C₂₀H₂₅F₂N₄O₂: 391.19456, Found: 391.19184.

The following compounds were synthesized according to the methods and variations of described for Example 5.

COMPOUND	NAME AND PHYSICAL DATA
	(4-{2-[4-(3,4-Difluoro-phenyl)-piperazin-1-ylmethyl]-allyloxy}-phenyl)-urea MS: 403.19326

	(4-{3-[4-(4-Fluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-urea MS: 373.20319
	(4-{3-[4-(4-Chloro-phenyl)-piperazin-1-yl]-2-hydroxy-propoxy}-phenyl)-urea
	1-Ethyl-3-(4-{3-[4-(4-fluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-urea
	(4-{3-[4-(4-Fluoro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-carbamic acid methyl ester MS: 388.20251

Examples 6, 7, and 8. N-[2-(3,4-Dichloro-phenylamino)-ethyl]-3-(4-methanesulfonylamino-phenyl)-propionamide (Compound 6), N-(4-{3-[2-(3,4-Dichloro-phenylamino)-ethylamino]-propyl}-phenyl)-methanesulfonamide (Compound 7), and N-(4-(3-(3-(3,4-dichlorophenyl)-2-oxoimidazolidin-1-yl)propyl)phenyl) methanesulfonamide (Compound 8).



Step (i) Methyl 3-(4-aminophenyl)propanoate (**v-1**). Thionyl chloride (14.6 ml, 200 mmol, 3.3 equiv) was added dropwise to a solution of dry methanol (60 ml, 1453 mmol, 24 equiv) at -10°C . After stirring for 10 minutes, 3-(4-aminophenyl)propanoic acid (10.0 g, 61 mmol) was added to give a yellow suspension. The solution stirred for 1 hour and was slowly warmed to room temperature. The resulting solution was concentrated to give a yellow solid. The solid was suspended in ethyl acetate, and NaHCO_3 (aq.) was added until the salt dissolved fully. Solid sodium bicarbonate was added to give pH 8. The layers were separated and the organics were washed with brine (aq.). The resulting solution was dried over MgSO_4 , filtered, and concentrated to give a yellow solid (10.6 g, 98%). ^1H NMR (300 MHz, CDCl_3) 7.00 (d, $J=8.3$ Hz, 2H), 6.63 (d, $J=8.3$ Hz, 2H), 3.67 (s, 3H), 3.59 (bs, NH_2 , 2H), 2.85 (t,

$J=7.6$ Hz, 2H), 2.58 (t, $J=8.3$ Hz, 2H). ^{13}C NMR (300 MHz, CDCl_3) 173.8, 144.9, 130.7, 129.3, 115.5, 51.8, 36.4, 30.4. M.S. (ESI) $m/z=180.102$ (M+H).

Step (ii). Methyl 3-(4-(methylsulfonamido)phenyl)propanoate (**v-2**). The ester (7.38 g, 41.2 mmol) was dissolved in pyridine (17.0 ml, excess). After cooling to 0°C , methanesulfonyl chloride (4.55 ml, 57.7 mmol, 1.4 equiv) was added dropwise. The reaction was warmed to room temperature and stirred overnight. The reaction was quenched with water and diluted with DCM. The layers were separated and the organics were washed with brine. The resulting solution was concentrated to give a red solid. The crude material was purified using silica gel chromatography (1 EtOAc/1 Hexanes) to give a white solid (87%). ^1H NMR: (CDCl_3 , 400 MHz) 7.20 (d, $J=8.6$ Hz, 2H), 7.15 (d, $J=8.6$ Hz, 2H), 6.45 (bs, NH, 1H), 3.68 (s, 3H), 3.00 (s, 3H), 2.94 (t, $J=7.6$ Hz, 2H), 2.63 (t, $J=7.5$ Hz, 2H). ^{13}C NMR (CDCl_3 , 400 MHz): 173.4, 137.6, 135.2, 129.4, 121.4, 51.7, 38.5, 35.5, 30.1. M.S. (ESI) $m/z=257.56$ (M+H)

Step (iii). 3-(4-(methylsulfonamido)phenyl)propanoic acid (**v-3**). The sulfonamide ester (1.16 g, 4.5 mmol) was dissolved in methanol (50 ml). To this solution, 1.0 N NaOH (17.0 ml, 17.0 mmol, 3.8 equiv) was added. The mixture was stirred at room temperature overnight. TLC indicated the reaction was finished. The pH of the solution was adjusted to 3 with a solution of aqueous HCl. The volume of methanol was reduced by rotary evaporation (40 mbar), upon which the product crashed out of solution. The yellow crystals were filtered off and dried (0.900 g, 82%). ^1H NMR (400 MHz, CD_3OD) 7.21 (d, $J=8.6$ Hz, 2H), 7.17 (d, $J=8.6$ Hz, 2H), 2.91 (s, 3H), 2.89 (t, $J=7.6$ Hz, 2H), 2.59 (t, $J=7.6$ Hz, 2H). ^{13}C NMR (400 MHz, CD_3OD) 176.7, 139.0, 137.7, 130.5, 122.3, 39.1, 36.8, 31.4. M.S. (ESI) $m/z=242.05$ (M-H).

Step (iv). N-(2-(3,4-dichlorophenylamino)ethyl)-3-(4-(methylsulfonamido)phenyl)propanamide Compound **6**). The carboxylic acid (0.700 g, 2.88 mmol) was dissolved in DMF (30.0 ml) and cooled to 0°C . To this solution, DMAP (0.352 g, 2.28 mmol, 1.1 equiv), and EDCI (0.552 g, 2.88 mmol, 1.0 equiv) were added to give a clear suspension. After stirring for 30 minutes, the amine (0.590 g, 2.88 mmol, 1.0 equiv) in THF (5.0 ml) was added dropwise to give a brown solution. The mixture was warmed to room temperature and stirred






overnight. The reaction was monitored by TLC. To quench the reaction, 20 mL of 1.0 N HCl was added and the solution was extracted with 3x30 mL of EtOAc. The organic layer was dried with MgSO_4 , filtered, and concentrated to give red oil. The crude material was purified by taking the residue up in DCM and stirring. Immediately a white powder precipitated out (0.920 g, 74%). ^1H NMR (400 MHz, CD_3OD) 7.14-7.10 (mult, 5H), 6.72 (d, $J=2.9$ Hz, 1H), 6.51 (dd, $J_1=8.9$ Hz, $J_2=2.6$ Hz, 1H), 3.27 (t, 2H), 3.10 (t, $J=6.4$ Hz, 2H), 2.88 (s, 3H), 2.87 (t, $J=6.5$ Hz, 2H), 2.46, (t, $J=6.4$ Hz, 2H). ^{13}C NMR (300 MHz, CDCl_3) 175.8, 150.0, 138.7, 138.3, 131.7, 130.5, 122.3, 114.4, 113.6, 44.0, 39.8, 39.2, 39.0, 32.3. M.S. Calc'd 429.0681 Found (HRMS) 431.08143 (M+H). E.A. Calc'd: C 50.24, H 4.92, N 9.76 Found: C 49.94, H 4.91, N 9.74.










Step (v). N-(4-(3-(2-(3,4-dichlorophenylamino)ethylamino)propyl)phenyl) methanesulfonamide (Compound 7). The sulfonamide amide ((0.500 g, 1.2 mmol) was dissolved in THF (30.0 ml). After cooling to 0°C , a solution of Lithium Aluminum hydride (2.0 M solution in THF, 2.3 ml, 4.6 mmol, 4.0 equiv) was added dropwise. After stirring for 10 minutes at 0°C , the ice bath was removed and the reaction mixed was warmed to room temperature and stirred overnight. The mixture was diluted with DCM and water to give an emulsion. Rochelle's salt (sat'd solution) was added and the mixture stirred for 20 minutes before filtering over a pad of celite. The resulting liquid was separated, and the organics were washed with brine, dried over MgSO_4 and concentrated to give a white foam (0.358 g, 74%). The free base was converted to the HCl salt by bubbling HCl (g) through a solution of substrate dissolved in ethanol. The white powder precipitated out and was filtered off. ^1H NMR (300 MHz, CDCl_3) 7.20-7.11 (mult, 5H), 6.69 (d, $J=2.8$ Hz, 1H), 6.46 (dd, $J_1=8.8$ Hz, $J_2=2.8$ Hz), 4.36 (bs, 1H, NH), 3.156 (mult, 2H), 3.00 (s, 3H), 2.88 (t, $J=6.2$ Hz, 2H), 2.66 (t, $J=7.1$ Hz, 4H), 1.86-1.79 (mult, 3H). ^{13}C NMR (300 MHz, CDCl_3) 148.1, 139.4, 134.8, 132.8, 130.7, 129.7, 121.6, 119.7, 113.9, 112.9, 49.0, 48.2, 43.2, 39.3, 32.9, 31.5. M.S. Calc'd 416.088. Found (HRMS): 416.069.

Step (vi). N-(4-(3-(3-(3,4-dichlorophenyl)-2-oxoimidazolidin-1-yl)propyl)phenyl) methanesulfonamide (Compound 8). The starting material diamine (0.113 g, 0.27 mmol) was dissolved in THF (10.0 ml). To this solution 1,1-carbonyldiimidazole (0.048 g, 0.30 mmol, 1.1 equiv) was added. The mixture stirred at room temperature overnight. After completion,

the solution was evaporated to dryness and the residue was taken up in ethyl acetate, washed with brine (1x) and dried over Na₂SO₄, filtered, and concentrated to give a clear oil. The crude material was purified using silica gel chromatography (100% EtOAc) to give a white foam (0.070 g, 58%). ¹H (400 MHz, CDCl₃) 7.72 (s, 1H), 7.16 (d, *J*=8.6 Hz, 2H), 7.07 (d, *J*=8.6 Hz, 2H), 7.02 (s, 1H), 6.64 (d, *J*=2.9 Hz, 1H), 6.41 (dd, *J*₁=8.5 Hz, *J*₂=2.9 Hz), 3.64 (t, *J*=6.0 Hz, 2H), 3.40-3.36 (mult, 4H), 2.97 (s, 3H), 2.57 (t, *J*=7.3 Hz, 2H), 1.26 (t, *J*=7.3 Hz, 2H). ¹³C (75 MHz, CDCl₃) 152.6, 147.1, 137.3, 136.8, 135.6, 130.9, 129.5, 121.6, 118.0, 113.6, 112.5. M.S. (ESI) Calc'd: 441.0681 Found: 442.07527 (M+H).

Compounds in the following table were synthesized according to variations in methods described for Examples 6, 7, and 8.

COMPOUND	NAME
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-2-(4-methanesulfonylamino-phenoxy)-acetamide
	N-(4-{2-[2-(3,4-Dichloro-phenylamino)-ethylamino]ethoxy}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-3-oxo-propyl}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(3,4-DiFluoro-phenyl)-piperazin-1-yl]-3-oxo-propyl}-phenyl)-methanesulfonamide
	N-(4-{3-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-propyl}-phenyl)-methanesulfonamide

	N-(4-{2-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-methanesulfonamide
	6-{2-[4-(3,4-Dichloro-phenyl)-piperazin-1-yl]-2-oxo-ethoxy}-3H-benzooxazol-2-one
	6-{2-[4-(3,4-Difluoro-phenyl)-piperazin-1-yl]-2-oxo-ethoxy}-3H-benzooxazol-2-one
	6-{2-[4-(4-Chloro-phenyl)-piperazin-1-yl]-2-oxo-ethoxy}-3H-benzooxazol-2-one
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-2-(2-oxo-2,3-dihydro-benzooxazol-6-yloxy)-acetamide
	N-[3-(3,4-Dichloro-phenyl)-allyl]-2-(4-methanesulfonylamino-phenoxy)-acetamide
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-2-(4-hydroxy-phenoxy)-acetamide
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-3-(4-hydroxy-phenyl)-propionamide
	N-[2-(3,4-Dichloro-phenylamino)-ethyl]-2-(3-fluoro-4-hydroxy-phenoxy)-acetamide

Biological Data

Example 9. Expression of glutamate receptors in *Xenopus laevis* oocytes.

cRNA was synthesized from linearized template cDNA for rat glutamate receptor subunits according to manufacturer specifications (Ambion). Quality of synthesized cRNA was assessed by gel electrophoresis, and quantity was estimated by spectroscopy and gel electrophoresis. Stage V and VI oocytes were surgically removed from the ovaries of large, well-fed and healthy *Xenopus laevis* anesthetized with 3-amino-benzoic acid ethyl ester (3 gm/l) as previously described. Clusters of isolated oocytes were incubated with 292 U/ml Worthington (Freehold, NJ) type IV collagenase or 1.3 mg/ml collagenase (Life Technologies, Gaithersburg, MD; 17018-029) for 2 hr in Ca^{2+} -free solution comprised of (in mM) 115 NaCl, 2.5 KCl, and 10 HEPES, pH 7.5, with slow agitation to remove the follicular cell layer. Oocytes were then washed extensively in the same solution supplemented with 1.8 mM CaCl_2 and maintained in Barth's solution comprised of (in mM): 88 NaCl, 1 KCl, 2.4 NaHCO_3 , 10 HEPES, 0.82 MgSO_4 , 0.33 $\text{Ca}(\text{NO}_3)_2$, and 0.91 CaCl_2 and supplemented with 100 $\mu\text{g/ml}$ gentamycin, 10 $\mu\text{g/ml}$ streptomycin, and 10 $\mu\text{g/ml}$ penicillin. Oocytes were manually defolliculated and injected within 24 hrs of isolation with 3-5 ng of NR1 subunit cRNA and 7-10 ng of NR2 cRNA subunit in a 50 nl volume, or 5-10 ng of AMPA or kainate receptor cRNAs in a 50 nl volume, and incubated in Barth's solution at 18°C for 1-7 d. Glass injection pipettes had tip sizes ranging from 10-20 microns, and were backfilled with mineral oil.

Example 10. Two electrode voltage clamp recording from *Xenopus laevis* oocytes

Two electrode voltage-clamp recordings were made 2-7 days post-injection as previously described. Oocytes were placed in a dual-track plexiglass recording chamber with a single perfusion line that splits in a Y-configuration to perfuse two oocytes. Dual recordings were made at room temperature (23°C) using two Warner OC725B two-electrode voltage clamp amplifiers, arranged as recommended by the manufacturer. Glass microelectrodes (1-10 Megaohms) were filled with 300 mM KCl (voltage electrode) or 3 M KCl (current electrode). The bath clamps communicated across silver chloride wires placed into each side of the recording chamber, both of which were assumed to be at a reference potential of 0 mV. Oocytes were perfused with a solution comprised of (in mM) 90 NaCl, 1 KCl, 10 HEPES, and 0.5 BaCl_2 ; pH was adjusted by addition of 1-3 M NaOH or HCl. Oocytes were recorded under voltage clamp at -40 mV. Final concentrations for control application of glutamate (50 μM) plus glycine (30 μM) were achieved by adding appropriate volumes from 100 and 30 mM stock solutions, respectively. In addition, 10 μM final EDTA was obtained by adding a




1:1000 dilution of 10 mM EDTA, in order to chelate contaminant divalent ions such as Zn^{2+} . Concentration-response curves for experimental compounds were obtained by applying in successive fashion maximal glutamate/glycine, followed by glutamate/glycine plus variable concentrations of experimental compounds. Dose response curves consisting of 4 to 8 concentrations were obtained in this manner. The baseline leak current at -40 mV was measured before and after recording, and the full recording linearly corrected for any change in leak current. Oocytes with glutamate-evoked responses smaller than 50 nA were not included in the analysis. The level of inhibition by applied experimental compounds was expressed as a percent of the initial glutamate response, and averaged together across oocytes from a single frog. Each experiment consisted of recordings from 3 to 10 oocytes obtained from a single frog. Results from 3-6 experiments were pooled, and the average percent responses at antagonist concentrations were fitted by the equation,










$$\text{Percent Response} = (100 - \text{minimum}) / (1 + ([\text{conc}] / \text{IC}_{50})^{nH}) + \text{minimum}$$

where *minimum* is the residual percent response in saturating concentration of the experimental compounds, IC_{50} is the concentration of antagonist that causes half of the achievable inhibition, and nH is a slope factor describing steepness of the inhibition curve. *Minimum* was constrained to be greater than or equal to 0.

Assay results for test compounds are reported in Tables 17-21.

Table 17. pH Dependence of NMDA Antagonism

COMPOUND	IC_{50} at pH6.9 (nM)	IC_{50} at pH7.6 (nM)
	31	557
	12	62
	296	1,020

	32	270
	213	1,020
	2,400	28,000
	1,810	12,700
	>30,000	>30,000
	55	133
	108	199
	7	38
	163	2,580


	38	71
---	----	----

Table 18. pH Dependence of NMDA Antagonism































COMPOUND	IC ₅₀ at pH 6.9 (nM)	IC ₅₀ at pH 7.6 (nM)
	79	726
	145	1,700
	185	1,500
	72	644
	185	446
	25	97

Table 19. pH Dependence of NMDA Antagonism

COMPOUND	IC ₅₀ at pH 6.9 (nM)	IC ₅₀ at pH 7.6 (nM)
	80	814
	131	611
	87	980
	50	272
	87	529
	1,500	5,790
	70	260

	460	4,370
	73	204
	576	4,700
	562	9,740
	113	448
	6,950	191,300
	529	3,560
	541	1,730

	978	4,390
	430	3,930
	3,210	8,330
	217	476
	211	598
	257	4,800
	341	1,740
	386	3,790
	197	501









	585	6,500
	116	330
	114	549
	191	298

Table 20. pH Dependence of NMDA Antagonism

COMPOUND	IC ₅₀ at pH 6.9	IC ₅₀ at pH 7.6
	82	696
	557	10,400
	117	625
	46	452




















	19	60
	3,030	15,700
	1,540	16,200

Table 21. pH Dependence of NMDA Antagonism

COMPOUND	IC ₅₀ at pH 6.9	IC ₅₀ at pH 7.6
	30	55
	2	21
	160	809
	13	58
	14	36

	670	5,330
	2,200	8,400
	230	2,730
	40	129
	41	45
	307	467
	189	243
	80	150
	10	70

	280	515
	210	339

Example 11. In vitro binding studies for secondary effects












Compounds were evaluated for binding to the human ether-a-go-go potassium channel (hERG) expressed in HEK293 cells by displacement of $^3\text{[H]}$ -astemizole according to the methods by Finlayson et al. (K. Finlayson., L. Turnbull, C.T. January, J. Sharkey, J.S. Kelly; $^3\text{[H]}$ Dofetilide binding to HERG transfected membranes: a potential high throughput preclinical screen. *Eur. J. Pharmacol.* 2001, 430, 147-148). Compounds were incubated at 1 or 10 μM final concentration, in duplicate, and the amount of displaced $^3\text{[H]}$ -astemizole determined by liquid scintillation spectroscopy. In some cases, a seven concentration (each concentration in duplicate) displacement curve was generated to determine an IC_{50} .

Binding to the rat alpha-1 adrenergic receptor in rat brain membranes was determined by displacement of $^3\text{[H]}$ -prazosin (P. Greengrass and R. Bremner; Binding characteristics of 3H-prazosin to rat brain α -adrenergic receptors. *Eur.J. Pharmacol.* 1979, 55: 323-326). Compounds were incubated at 0.3 or 3 μM final concentration, in duplicate, and the amount of displaced $^3\text{[H]}$ -prazosin determined by liquid scintillation spectroscopy.

Binding IC_{50} values were determined from displacement curves (four-six concentrations, each concentration in duplicate) fit by a non-linear, least squares, regression analysis using MathIQ (ID Business Solutions Ltd., UK). The binding K_i 's were determined from the IC_{50} according to the method of Cheng and Prusoff (Y. Cheng and W.H. Prusoff; Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 percent inhibition (IC_{50}) of an enzymatic reaction. *Biochem. Pharmacol.* 1973, 22: 3099-3108).

Table 22: NMDA antagonism at pH 6.9 and 7.6 and hERG and $\alpha 1$ adrenergic receptor binding

Compound	IC_{50} pH 6.9 (nM)	IC_{50} pH 7.6 (nM)	hERG K_i (nM)	$\alpha 1$ IC_{50} (nM)
----------	------------------------------------	---------------------------------	------------------------------	--

	74	555	39,000	720
	51	447	1,600	620
	32	270	553	350
	46	452	13,000	340
	410	3,830	~1,000	~300
	69	550	~7,500	~350
	14	103	~7,500	~350
	>10,000	>10,000	-	-
	370	29	~10,000	~100
	16	76	~7,500	~150
	12	66	~5,000	~100

Example 12. Metabolic stability

Compounds were incubated with pooled human (from at least 10 donors) or rat liver microsomes, 1.0 mg/ml microsomal protein, and 1 mM NADPH, in buffer at 37°C in a shaking water bath according to the method of Clarke and Jeffrey (S.E. Clarke and P. Jeffrey; *Xenobiotica* 2001. 31: 591-598). At 60 min the samples were extracted and analyzed for the presence of the parent compound by LC-MS/MS. The parent material remaining in the sample at 60 min was compared to that at 0 min and expressed as a percentage. A control compound, testosterone, was run in parallel.






Example 13. Plasma half-life and brain exposure

Rats (n=3 per dose) were administered compounds at a doses of 1-4 mg/kg in a single bolus i.v. infusion (2 ml/kg body weight) via the tail vein formulated in 2% dimethyl acetamide/ 98% 2-hydroxy-propyl cyclodextrin (5%). Animals were fasted overnight prior to dose administration and food returned to the animals two hours after dosing. Following IV dosing, blood samples (*ca* 200 µL) were collected into separate tubes containing anticoagulant (K-EDTA) *via* the orbital plexus at various times post administration. Plasma samples were prepared immediately after collection by centrifugation for ten minutes using a tabletop centrifuge, and stored at -80°C. Brain tissue was weighed, homogenized on ice in 50 mM phosphate buffer (2 ml per brain) and the homogenate stored at -80°C. Plasma and brain homogenate samples were extracted by the addition of 5 volumes of cold acetonitrile, mixed well by vortexing and centrifuged at 4000 rpm for 15 minutes. The supernatant fractions were analyzed by LC-MS/MS operating in multiple reaction monitoring mode (MRM). The amount of parent compound in each sample was calculated by comparing the response of the analyte in the sample to that of a standard curve.

Penetration Classification (Table 24) using the *in vitro* cell permeability assay to predict brain penetration potential: Transwell® wells containing MDR1-MDCK cell monolayers that express the multidrug transporter P-gp were used for measuring the percent recovery of compound after dosing both sides of a cell monolayer with the test article. Monolayers were grown for 7-11 days at which time 5 µM of the test article was made by dilution from DMSO stocks into a Hank's balanced salt solution (pH 7.4), final DMSO not greater than 1%, and added to: a) the apical side for A-B permeability (apical to basal) assessment, or separately b) the basal side for the B-A permeability (basal to apical)

assessment, all at pH 7.4. After a 2 hr incubation (37°C) both the apical and the basal compartments were sampled and the amount of test article present determined by generic LC-MS/MS methods against a ≥ 4 point calibration curve. Experiments were done in duplicate. Apparent permeability (P_{app} units are reported $\times 10^{-6}$ cm/s) are determined for the A-B and the B-A directions as well as the Efflux ratio ($P_{app}B-A/P_{app}A-B$). The blood-brain barrier penetration potential is classified as follows: “High” when $P_{app} A-B \geq 3.0 \times 10^{-6}$ cm/s, and efflux < 3.0 ; “Moderate” when $P_{app} A-B \geq 3.0 \times 10^{-6}$ cm/s, and $10 > \text{efflux} \geq 3.0$; and “Low” when either $P_{app} A-B \geq 3.0 \times 10^{-6}$ cm/s, and efflux ≥ 10 , or when $P_{app} A-B < 3.0 \times 10^{-6}$ cm/s.

Table 23: Plasma Stability Results

Compound	PK-i.v. t _{1/2} (hr)	i.v. C _{max} ng/ml	AUC (0-last h) i.v. XMPK (hr*ng/mL)	Dosing	Formulation
	2.7	252	1031*	(a) 1 MPK, 0-4 hr i.v.;	2% DMA/98% 2-HPBCD (5%in Water)
	2.4	241	452	(b) 1 MPK, 0-6 hr i.v.;	5% DMA/95% 2-HPBCD (5%in Water)
	2.6	1069	2351	(b) 3MPK, 0-4 hr i.v.;	2% DMA/98% 2-HPBCD (5%in Water)
	> 2 i.p.**	13403 (30m), 13057 (2hr),		60 MPK, i.p.	5% DMA/95% 2-HPBCD (5%in Water)
	0.8	464	347	(b) 1 MPK, 0-6 hr i.v.; .	5% DMA/95% 2-HPBCD (5%in Water)


















Compound	PK-i.v. t _{1/2} (hr)	i.v. C _{max} ng/ml	AUC (0-last h) i.v. XMPK (hr*ng/mL)	Dosing	Formulation
	0.6	193	187	(a) 1 MPK, 0-4 hr i.v.;	2% DMA/98% 2-HPBCD (5%in Water)
	0.6	288	217	(a) 1 MPK, 0-4 hr i.v.	2% DMA/98% 2-HPBCD (5%in Water)
	0.6	568	364	(a) 1 MPK, 0-4 hr i.v.;	2% DMA/98% 2-HPBCD (5%in Water)
	1.14	715	787.4	(b) 1MPK, 0-6 hr i.v.;	10% DMA/10% EtOH/30% 2-HPBCD/50% water
	≤ 0.8 hr	377 (30m)		3 MPK i.v.	2% DMA/98% 2-HPBCD (5%in Water)
	1.84	575.7	1096	(b) 1MPK, 0-6 hr i.v.;	10% DMA/10% EtOH/30% 2-HPBCD/50% water
	~0.83 i.p.	960 (15m)		10 MPK i.p.	50% DMSO/50% saline
	≤ 0.8 hr	398 (30m)		3 MPK i.v.	2% DMA/98% 2-HPBCD (5%in Water)
	~0.5 hr	413 (30m)		3 MPK i.v.	2% DMA/98% 2-HPBCD (5%in Water)

Table 24: Brain Penetration

Compound	RATIO: Brain:plasma (Direct)	Dosing	Formulation	BBB Pene Classification
	10.2 (avg); 11 (30m), 9.5 (1hr),	3 MPK i.v.	2% DMA/98% 2-HPBCD (5%in Water)	High
	2.3 (avg); 2.3 (30m), 2.9 (1hr), 1.7 (2hr)	3 MPK i.v.	2% DMA/98% 2-HPBCD (5%in Water)	Moderate
	0.77 (avg); 0.7 (30m), 0.7 (2hr), 0.9 (4hr)	60 MPK, i.p.	5% DMA/95% 2-HPBCD (5%in Water)	Moderate
	0.42 (avg); 0.24 (30m), 0.53 (1hr), 0.49 (6hr)	3 MPK i.v. @ 6hr, 1MPK @ 0.5hr and 2 hr	2% DMA/98% 2-HPBCD (5%in Water)	Low
	0.59 (avg); 0.68 (30m), 0.49 (1hr),	3 MPK i.v.	2% DMA/98% 2-HPBCD (5%in Water)	High
	Below Detection limit (~ 0)	1 MPK i.v.	10% DMA/10% EtOH/30% 2- HPBCD/50% water	Low
	0.33 (avg); 0.25 (15m), 0.49 (30m), 0.30 (1hr), 0.36 (2hr)	10 MPK i.p.	50% DMSO/50% saline	Moderate
	0.07 (avg.); 0.04 (15m), 0.06 (30m), 0.07 (1hr), 0.10 (3hr)	1 MPG i.v.	10% DMA/10% EtOH/30% 2- HPBCD/50% water	Moderate








Compound	RATIO: Brain:plasma (Direct)	Dosing	Formulation	BBB Pene Classification
	0.25 (avg); 0.15 (30m), 0.35 (1hr), BLQ (2hr)	3 MPK i.v.	2% DMA/98% 2-HPBCD (5%in Water)	High
	BLQ (avg); BLQ (30m), BLQ (1hr),	3 MPK i.v.	2% DMA/98% 2-HPBCD (5%in Water)	Moderate

Table 25: Oral Absorption

Compound	% Ab- sorbed (%F)	p.o. Cmax	p.o. Tmax hr	p.o. t1/2 hr	AUC (0-last h) p.o. (hr*ng/ mL)	Dosin g	Formulatio n
	145.7* *	1556.6	2.67	6.46	8451	10 MPK, 0-8 hr p.o.	2% DMA/98% 2-HPBCD (5%in Water)
	82.2	558	0.5	3.04	1856	5 MPK, 0-6 hr p.o.	5% DMA/95% 2-HPBCD (5%in Water)
	31.6	407	0.33	0.85	549	5 MPK, 0-6 hr p.o.	5% DMA/95% 2-HPBCD (5%in Water)
	7.9**	57.8	6.7	nc	284	10 MPK, 0-8 hr p.o.	2% DMA/98% 2-HPBCD (5%in Water)
	6.2**	30.6	8	nc	132	10 MPK, 0-8 hr p.o.	2% DMA/98% 2-HPBCD (5%in Water)































Compound	% Absorbed (%F)	p.o. Cmax	p.o. Tmax hr	p.o. t1/2 hr	AUC (0-last h) p.o. (hr*ng/mL)	Dosin g	Formulation
	3.3**	10.2	5.3	nc	56.4	10 MPK, 0-8 hr p.o.	2% DMA/98% 2-HPBCD (5%in Water)

Table 26 - Structures of compounds referenced in Examples 14-22.

Label	Compound
93-31	
93-97	
93-108	 HCl
NP10001	
NP10002	

NP10030	
NP10031	
NP10039	
NP10045	
NP10056	
NP10066	
NP10068	
NP10070	

NP10075	
NP10076	
NP10097	
NP10099	
NP10119	
NP10122	
NP10146	
NP10150	

NP10153	
NP10165	
NP10185	
NP10226	
NP10231	
NP10239	
NP10250	
NP10272	

Example 14: Forced swim model

CD1 mice were administered a compounds shown in Table 11, desipramine, Ro 25-6981 or a control vehicle and subjected to a forced swim test. All compounds were administered as intraperitoneal injections. Animals were placed into a beaker (15 cm diameter) of water held at 25°C with a depth of 15 cm 30 min after compound administration. Behavior was videotaped for 6 minutes from the side of the beaker and scored subsequently for struggling behavior. Results were analyzed by one-way ANOVA and post-hoc Bonferroni tests. Immobility time date from the forced swim tests is shown in Figures 1 and 2. Total immobility time refers to the time that the animal spends floating or engaged in minimal activity to keep afloat for at least 3 seconds. Subtle movements of feet, tail or head required to maintain the eyes, ears, and nose above the surface of the water were excluded as immobility. The video-tapes were scored by investigators unfamiliar with the treatments of the mice.

For the data in Figure 1, test compounds were tested at a dose of 10 mg/ kg. Desipramine was tested at a dose of 20 mg/kg. Ro 25-6981 was tested at a dose of 5 mg/kg. Number of CD1 mice tested per groups was 8-10. ANOVA: $F(11,98) = 3.638$, $p < 0.01$. * = $p < 0.05$ compared to vehicle. + = $p < 0.05$ compared to desipramine.

For the data in Figure 2, compounds NP10075 and NP10076 were tested at doses of 5 mg/kg, 7.5 mg/kg and 10 mg/kg; desipramine was tested at a dose of 20 mg/kg; and Ro 25-6981 was tested at a dose of 5 mg/kg. Number of CD1 mice tested per group was 8-10. * = $p < 0.05$, ** = $p < 0.01$ ANOVA, post-hoc Bonferroni from control.

Example 15: Open field activity test

Spontaneous activity was evaluated in an automated Omnitech Digiscan apparatus (AccuScan Instruments, Columbus, OH). Animals were given vehicle, imipramine, or a dose of a test compound. All compounds were administered as intraperitoneal injections. Activity was summated at 5 minute intervals over the 90 min period of testing. At 60 minutes, the mice were injected with 10 mg/kg NP10075, 10 mg/kg NP10076 or a vehicle. Locomotion was measured in terms of the total distance traveled (horizontal activity). Results were analyzed by one-way ANOVA and post-hoc Bonferroni tests. Neither NP10075 or NP10076 altered mouse open field activity at a dosage of 10 mg/kg. Data for these tests is shown in Figure 3.

Example 16: Plasma and Brain Exposure Assessment

Mice were administered a dose of test compound (10 mg/kg by i.p.) and blood and brain tissue samples collected at the indicated times post drug administration (n=3-5). Blood samples were collected in K-EDTA tubes and centrifuged for ten minutes immediately after collection, and the plasma was stored at -80°C until analysis. Brains were immediately removed from the skull and the meninges and cerebellum removed, rinsed with ice-cold PBS, weighed and then homogenized at 4°C in 2-3 volumes of 50 mM potassium phosphate buffer (pH 7.4) and stored at -80°C until analysis. Plasma and brain homogenates were extracted by the addition of 5 volumes of cold acetonitrile, mixed well by vortexing and centrifuged at 4000 rpm for 15 minutes. The supernatant fractions were analyzed by LC-MS/MS operating in multiple reaction monitoring mode (MRM) and analyzed for the parent compound to determine the plasma or brain concentration. Internal standards were added to calibrate each sample. An eight point standard curve was prepared similarly in naïve plasma and brain for each compound of interest. Plasma and brain exposure assessment data is provided in Table 27. Based on occupancy studies of other NR2B antagonists in rodents (CP-101,606, Ro25-6981, and Merck20j), plus plasma levels achieved in Preskorn et al., brain exposures for the test compounds were consistent with “anticipated levels” required for efficacy. Results are shown in Table 27.

Table 27. Plasma and Brain Drug Concentration

Depression Result	Label	Rat IC₅₀ (μM)	Plasma* (ng/ml)	Free Plasma** (ng/ml)	Free Plasma (μM)	Brain*** (μM)
active	NP10075	0.108	3137	220	0.542	0.130
active	NP10076	0.072	377	34	0.094	0.064
active	NP10068	0.060	679	61	0.145	0.048

* = Plasma sample analysis by LC-MS/MS, in mice dosed 10 mg/kg, i.p., 30 minute time point. ** = Based on % free fraction measured in protein binding study, by LC-MS/MS or NR2B receptor bioassay. *** = Calculated from the brain:plasma ratio for the compounds determined by LC-MS/MS.

Example 17: Rotorod test (in vivo safety)

The rotorod test is a modification of the procedure described by Rozas and Labandeira-Garcia (1997). The test is initiated by placing mice on a rotating rod (5 rpm) that is 3.8 cm diameter by 8 cm wide and suspended 30 cm from the floor of a chamber. After 10

sec the rotation is accelerated from 5 to 35 rpm over a 5 minute period. The time the mouse falls from the rod (the latency time) is recorded automatically with a light-activated sensor in the bottom of the chamber. Animals were trained four times each day for two days, with a within-day inter-trial interval of 20-25 min and a between-day interval of 24 hrs. On day 3, mice were randomly assigned to groups and injected in a blinded fashion with either vehicle, positive control (0.3 mg/kg (+)MK-801 or 10 mg/kg ifenprodil), or doses of NP compound. All drugs were administered i.p. Results were analyzed by ANOVA and Dunnett's tests. Data is shown in Figure 4.

Example 18: Cell toxicity in cortical neuron culture

Primary cultures of rat cerebral cortex were prepared from Sprague-Dawley rat embryos (E16-E19). Cells were plated into 24 well plates at a density of 3×10^5 per well, in Neurobasal medium supplemented with L-glutamine (2 mM), penicillin (5 U/ml), streptomycin (10 μ g/ml) and B-27. After 14-22 days in culture, cells were treated with test compounds (in triplicate wells) at 10 μ M, final, and incubated for 24 hrs. Cell death was assessed by measuring the amount of lactate dehydrogenase (LDH) released into the culture medium (Tox-7 kit; Sigma Chemical Co, St. Louis, Mo). Released LDH was expressed as the fraction of total LDH present in each well. Maximal cell death was determined by treating separate wells (in triplicate) with saturating concentrations of NMDA (100 μ M) and glycine (10 μ M) for 24 hrs. Results shown are the Mean $\pm$ SEM from a minimum of three separate cultures. Cell toxicity was assessed by % total LDH release after 24 hr incubation of 10 μ M compound in cell culture. For each compound, three cultures were treated with 10 μ M compound. Data is shown in Figure 5.

Example 19: Ames test for genotoxicity

The Ames test determines the ability of a compound to reverse an introduced mutation in two strains of *Salmonella typhimurium* (selected from TA98, TA100, TA15345, TA1537, and TA102). (See for example Maron, D.M. and Ames, B.N., *Mutat. Res.*, 1983, 113, 173–215.) Compounds were tested at eight dose levels 1.5, 5, 15, 50, 150, 500, 1500, and 5000 μ g/plate in both the presence and absence of S-9 microsomal fraction in two bacterial strains (TA98, TA102). After incubation at 37° the number of revertant colonies was compared with the number of spontaneous revertants on negative (vehicle) plates. Positive control plates containing a known mutagen active in each of the strains in the presence of S-9 extract (2-aminoanthracene at 1-5 μ g/plate) were also run. Data is shown in Table 28.

Table 28. Genotoxicity Test Results

Label	Conc #, range (µg/plate)	+/- S9 Activation	Strains	Valid Assay	Result
93-31	8; 1.5-5000	Both	TA98, TA102	Yes	No positive response
NP10056	8; 1.5-5000	Both	TA98, TA102	Yes	No positive response
NP10075	8; 1.5-5000	Both	TA98, TA102	Yes	No positive response
NP10122	8; 1.5-5000	Both	TA98, TA102	Yes	No positive response
NP10226	8; 1.5-5000	Both	TA98, TA102	Yes	No positive response
NP10231	8; 1.5-5000	Both	TA98, TA102	Yes	No positive response

Example 20: hERG Binding

Compounds were evaluated for binding to the human ether-a-go-go potassium channel (hERG) expressed in HEK293 cells by displacement of 3[H]-astemizole. Binding studies were performed either at a single concentration of 10 µM (in duplicate) or binding IC₅₀ values determined from displacement curves (four-six concentrations, each point in duplicate) fit by a non-linear, least squares, regression analysis using MathIQ (ID Business Solutions Ltd., UK). Data is shown in Figure 6. Compounds were evaluated for binding to the human ether-a-go-go potassium channel (hERG) expressed in HEK293 cells by displacement of 3[H]-astemizole. Binding studies were performed either at a single concentration of 10 µM (in duplicate) or binding IC₅₀ values determined from displacement curves (four-six concentrations, each point in duplicate) fit by a non-linear, least squares, regression analysis using MathIQ (ID Business Solutions Ltd., UK). Functional hERG channel block was determined using patch clamp methods with stable hERG channel transfectants in HEK293 cells. All experiments were performed at ambient temperature. Each cell acted as its own control. Three to five concentrations of the test article was applied at 5 minute intervals via micropipette tips to cells expressing hERG (n≥3 cells/concentration). Duration of exposure to each test article concentration was 5 minutes. After vehicle application, the positive control was applied in the same manner to verify sensitivity to hERG blockade. Intracellular solution for whole cell recordings consisted of (composition in mM): potassium aspartate, 130; MgCl₂, 5; EGTA, 5; ATP, 4; HEPES, 10; pH adjusted to 7.2 with KOH. After establishment of a whole-cell configuration, membrane currents were recorded

using QPatch HT® system. Before digitization, the current records were low-pass filtered at one-fifth of the sampling frequency. Onset and block of hERG current was measured using a stimulus voltage pattern consisting of a 200 ms prepulse to -40 mV (leakage subtraction), a 2-second activating pulse to +40 mV, followed by a 2-second test pulse to -40 mV. The pulse pattern was repeated continuously at 10 s intervals, from a holding potential of -80 mV. Peak tail currents were measured during the -40 mV test pulse. Leakage current was calculated from the current amplitude evoked by the prepulse and subtracted from the total membrane current recorded. Data acquisition and analysis was performed using the suite of Assay Software programs (Sophion Bioscience A/S, Denmark). Steady state was defined by the limiting constant rate of change with time (linear time dependence). The steady state before and after test article application was used to calculate the percentage of current inhibited at each concentration. Concentration-response data were fit to the following equation:

$$\% \text{ Block} = \{1 - 1/[1 + ([\text{Test}]/\text{IC}_{50})^N]\} * 100$$

where [Test] is the concentration of test article, IC₅₀ is the concentration of the test article producing half-maximal inhibition, N is the Hill coefficient, and % Block is the percentage of hERG potassium current inhibited at each concentration of the test article. Data were fit by a nonlinear least squares fits with the Solver add-in for Excel 2000 (Microsoft, Redmond, WA). Data is shown in Figure 6.

Example 21: Langendorff Heart Preparation and Measurement of QT Effects

The effects of test compounds on the QT-interval of the electrocardiogram were evaluated in vitro using an isolated retrograde perfused rabbit (New Zealand white female) heart preparation (Langendorff) with an ablated AV node and stimulated at a basic cycle length of 1s. Test article concentrations were prepared by diluting stock solutions in DMSO into Krebs-Henseleit (KH) solution (composition in mM): NaCl, 129; KCl, 3.7; CaCl₂, 1.3; MgSO₄, 0.64; Na-Pyruvate, 2.0; NaHCO₃, 17.8; Glucose, 5. The solution was aerated with a mixture of 95% O₂ and 5% CO₂ (pH 7.3-7.45). All test solutions contained 0.3% DMSO, final. Briefly, rabbits were heparinized and anesthetized with sodium pentothal and hearts rapidly removed via a midsternal thoracotomy and placed in chilled oxygenated (95% O₂ + 5% CO₂) KH solution. The heart was mounted in a Langendorff heart perfusion apparatus and perfused at a constant flow with KH solution (37 °C) in a retrograde fashion through the aorta. The A-V node was ablated to slow the intrinsic heart rate to a ventricular escape less than 60 beats/min. Following immersion of the heart into the bath the volume-conducted

ECG was recorded via bath-mounted electrodes. Three Ag/AgCl pellet electrodes were positioned in the bath chamber to form an equilateral triangle centered on the heart. Each heart was paced by repetitive electrical stimuli (0.1-5 ms, approximately 1.5 x threshold) by a pulse generator. The ECG signals were conditioned by an AC-coupled preamplifier (Grass Model P511) with low-pass filtering to achieve a bandwidth of 10 – 300 Hz. A stabilization period was at least 30 minutes long before obtaining baseline control responses. Test article concentrations were applied sequentially, in ascending order for exposure periods of at least 15 minutes/concentration to allow equilibration with the tissue. The average responses from at least three hearts were analyzed for each test condition. The QT interval was calculated and the Mean $\pm$ SEM values from the last four beats in the equilibration period were measured. Test results are shown in Figure 7.

Example 22: PCP Discrimination Test


Development of N-methyl-D-aspartate (NMDA) antagonists for a variety of disorders has been hindered by their production of phencyclidine (PCP)-like psychological effects and abuse potential. Drug discrimination studies allow direct comparisons to be made among the discriminative stimulus effects of drugs (Balster, 1990; Holtzman, 1990) and are considered to be predictive of subjective effects in humans. Sprague-Dawley rats were trained to discriminate 2mg/kg (i.p.) PCP or saline when administered intraperitoneally 15 min before the session under a double alternation schedule. Rats were placed in the operant chambers and the session initiated, as signaled by illumination of the chamber houselight. Completion of a FR32 on the correct lever resulted in delivery of a 45- mg food pellet (PJ Noyes Company, Inc., Lancaster, New Hampshire, USA). Incorrect responding reset the FR for correct-lever responding. Training was continued until the animals respond reliably and complete the first FR with more than 80% of total responses on the correct lever during a minimum of four consecutive sessions. Subsequent to acquisition of the PCP-saline discrimination, test sessions commenced on when animals met the following criteria on the most recent PCP and saline training sessions: (i) first FR completed on the correct lever, and (ii) greater than 85% correct-lever responding over the entire session. The animals were tested with different doses of test drug (as shown), generally given in an ascending order across test days. Various doses of PCP and test compounds were administered intraperitoneally 15 min before session initiation. To demonstrate the degree of stimulus control, tests with 2 mg/kg PCP and saline were carried out before and after each dose–response curve. In addition vehicle was also tested. Between test sessions, animals continued

to train with PCP and saline injections. Illumination of lights, recording of responses and pellet delivery was controlled by a microcomputer using MEDPC software (Med Associates). For data analysis the mean ($\pm$ SE) percentage PCP-lever responding and response rate (resp/s) effects was evaluated for all test sessions. Full substitution for PCP required greater than 80% PCP-lever responding, partial substitution as producing between 20 and 80% PCP-lever responding, and less than 20% PCP-lever responding will be indicative of a lack of PCP-like discriminative stimulus effects. Additionally, the mean response rate for all animals during each test session was determined to reveal any nonspecific effects on behavior.

Data for 93-31 (NP031) and 93-97 (NP097) compared to PCP are shown in Figure 8.

WE CLAIM

1. A method of treatment or prophylaxis of a neuropsychiatric disorders comprising administering a compound of Formula I or II, or a pharmaceutically acceptable salt, ester, prodrug or derivative thereof to a host in need thereof:

FORMULA I

wherein:

each L is independently C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, alkaryl, hydroxy, -O-alkyl, -O-aryl, -SH, -S-alkyl, -S-aryl, fluoro, chloro, bromo, iodo, nitro, or cyano; or two L groups may be taken together with Ar¹ to form: a dioxolane ring or a cyclobutane ring;

k = 0, 1, 2, 3, 4 or 5;

each Ar¹ and Ar² is independently aryl or heteroaryl;


W is a bond, C₁-C₄ alkyl, or C₂-C₄ alkenyl;

X is a bond, NR¹ or O wherein each R¹ and R² is independently H, C₁-C₆ alkyl, C₂-C₆ alkenyl or C₆-C₁₂ aralkyl; or R¹ and R² can be taken together to form a 5-8 membered ring;

each R³ and R⁴ is independently H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR³R⁴ is C=O;

n and p are independently 1, 2, 3 or 4;


each R⁵ and R⁶ is independently H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C(=O)-(C₁-C₆)-alkyl, C₁-C₆ haloalkyl, hydroxy, fluoro, chloro, bromo, iodo, nitro, or cyano; or CR⁵R⁶ is



C=O or C=CH₂; or wherein -NR²-(CR⁵R⁶)_p- can be

Y is a bond, O, S, SO, SO₂, CH₂, NH, N(C₁-C₆ alkyl), or NHC(=O);

Z is OH, NR^6R^7 , $\text{NR}^8\text{SO}_2(\text{C}_1\text{-C}_6 \text{ alkyl})$, $\text{NR}^8\text{C}(\text{O})\text{NR}^6\text{R}^7$, $\text{NR}^8\text{C}(\text{S})\text{NR}^6\text{R}^7$, $\text{NR}^8\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_6 \text{ alkyl})$, NR^8 -dihydrothiazole, or NR^8 -dihydroimidazole; wherein each R^6 , R^7 and R^8 is independently H, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_6\text{-C}_{12}$ aralkyl; or



FORMULA II


wherein:

each G is independently F, Cl, Br, I, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_6\text{-C}_{12}$ aralkyl, -O-aryl, -S-aryl, -NH-aryl;

$f = 0, 1, 2, 3, 4$ or 5 ;

Ar^a and Ar^b are each independently aryl or heteroaryl;

B is selected from the group consisting of:



wherein $R^a, R^b, R^c, R^d, R^e, R^f, R^g, R^h, R^k$ and R^p are each independently selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, OH or halo;

R^j is H, C_1 - C_6 alkyl, OH or $P(O)(OC_1-C_4 \text{ alkyl})_2$;


R^m is C_1 - C_4 alkyl or C_2 - C_4 alkenyl;

R^n is C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_6 - C_{12} aralkyl, $-CH_2O-$, $-CH(C_1-C_6 \text{ alkyl})O-$, $-CH(C_2-C_{12} \text{ aralkyl})O-$;

t, w, y and z each = 0, 1, 2, or 3;

X and X' are independently selected from a bond, O, S, SO, SO_2 , CH_2 , NH, $N(C_1-C_6 \text{ alkyl})$, and $NHC(=O)$;


M is OH, F, Cl, Br, I, NH_2 , NR^qR^r , NO_2 , $O(C_1-C_6 \text{ alkyl})$, OCF_3 , CN, $C(O)OH$, $C(O)O(C_1-C_6 \text{ alkyl})$, C_6 - C_{12} aralkyl, $NR^sC(O)CR^t_3$, $NR^8SO_2(C_1-C_6 \text{ alkyl})$, or $NR^uC(O)NR^v_2$; wherein each R^q, R^r, R^s, R^u and R^v is each independently H or C_1 - C_6 alkyl; and each R^t is independently H, C_1 - C_6 alkyl or halo; or two M groups may be taken together with Ar^b to form:



; and wherein R^u and R^w are independently H, C_1 - C_6 alkyl or C_6 - C_{12} aralkyl; and

$h = 1, 2, 3, 4$ or 5 .

2. The method of claim 1 wherein the compound is of Formula II and each G is independently F, Cl, Br, I and $f = 0, 1$ or 2 ;




wherein $R^{a-e, g, h, k, p}$ are each from H and R^f is selected from H, OH or halo;

R^m is C_1 - C_4 alkyl or C_2 - C_4 alkenyl;


t, w, y and z each = 0, 1, 2, or 3;

X and X' are independently selected from a bond, O, S, CH₂, and NH;

M is OH, F, Cl, Br, I, NH₂, NR^qR^r, NO₂, O(C₁-C₆ alkyl), OCF₃, CN, C(O)OH, C(O)O(C₁-C₆ alkyl), C₆-C₁₂ aralkyl, NR^sC(O)CR^t₃, NR⁸SO₂(C₁-C₆ alkyl), or NR^uC(O)NR^v₂; wherein each R^q, R^r, R^s, R^u and R^v is each independently H or C₁-C₆ alkyl; and each R^t is independently H, C₁-C₆ alkyl or halo; or two M groups may be taken together with Ar^b to form:



3. The method of claim 1, wherein the compound is a compound of Formula A:



FORMULA A

wherein:

R¹ is H, F, Cl, Br, CF₃, C₁₋₆ alkyl, C(O)CH₃, C(O)CO-(C₁₋₆ alkyl), CH₂OH, CN, NH₂, N(C₁₋₆ alkyl)₂, OH, O-(C₁₋₆ alkyl), OCF₃, S-(C₁₋₆ alkyl), SO₂-(C₁₋₆ alkyl);

R² is H, F, Cl, methyl, CF₃;

R³ is H, F, Cl, CH₃, CF₃, CN;

each of R⁴ and R^{4'} are independently selected from H or methyl;

each of R⁵ and R^{5'} can be H or OH, or R⁵ and R^{5'} can be taken together to form =CH₂;

R^6 is H or F;


X is H or F;

Y is OH, NHSO_2R^7 , or NHC(O)NHR^8 ;


R^7 is C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

R^8 is H, C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;


or X and Y are taken together to form a heterocycle wherein the moiety



is selected from the group consisting of:



4. The method of claim 1, wherein the compound is a compound of Formula B:



FORMULA B

wherein:

R^1 is H, F, Cl, Br, CF_3 , or C_{1-6} alkyl;

Z is O, S, NH, CH_2 or a bond;

R^2 is H or OH;

R^6 is H or F;


X is H or F;

Y is OH, NHSO_2R^7 or NHC(O)NHR^8 ;


R^7 is C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;

R^8 is H, C_{1-6} alkyl, C_{6-12} aryl, or C_{7-13} aralkyl;


or X and Y are taken together to form a heterocycle wherein the moiety



is selected from the group consisting of:



5. The method of claim 1 wherein the disorder is depression.
6. The method of claim 4 wherein the host has been diagnosed with a major depression.
7. The method of claim 1 wherein the compound is administered to a host at risk of suffering from a depressive episode.
8. The method of claim 1 wherein the compound is administered in combination with a pharmaceutically acceptable carrier.
9. The method of claim 1 wherein the compound is administered in combination or alternation with a second active agent.

**Figure 1**

Forced Swim Test




Figure 2

Open Field Activity Test





Figure 3

**Figure 4**




Figure 5




Figure 6




Figure 7




Figure 8