




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A01N 47/34, C07C 275/54</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/19542 (43) International Publication Date: 14 May 1998 (14.05.98)</p>
<p>(21) International Application Number: PCT/US97/18199 (22) International Filing Date: 17 October 1997 (17.10.97) (30) Priority Data: 08/745,387 8 November 1996 (08.11.96) US (71) Applicant: DOW AGROSCIENCES LLC [US/US]; 9330 Zionsville Road, Indianapolis, IN 46268 (US). (72) Inventors: SBRAGIA, Ronald, J.; 3545 Cedar Ravine Road, Placerville, CA 95667 (US). JOHNSON, George, W.; 6715 Knollcreek Drive, Indianapolis, IN 46256 (US). KARR, Laura, L.; 800 N. 200 E., Lebanon, IN 46052 (US). EDWARDS, Jeff, M.; Route 1, Box 47, Torrington, WY 82240 (US). SCHNEIDER, Brian, M.; 75 Red Oak Lane, Carmel, IN 46033 (US). (74) Agent: STUART, Donald, R.; Dow AgroSciences LLC, Law Dept., 9330 Zionsville Road, Indianapolis, IN 46268-1054 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>

(54) Title: NEW BENZOYLPHENYLUREA INSECTICIDES AND METHODS OF USING CERTAIN BENZOYLPHENYLUREAS TO CONTROL COCKROACHES, ANTS, FLEAS, AND TERMITES



(57) Abstract

Compounds of formula (I) wherein R¹ and R² are H, methyl, or ethyl, have unexpected activity against cockroaches, ants, fleas, and termites. Compounds wherein at least one of R¹ and R² is methyl or ethyl are novel.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**NEW BENZOYLPHENYLUREA INSECTICIDES AND
METHODS OF USING CERTAIN BENZOYLPHENYLUREAS TO
CONTROL COCKROACHES, ANTS, FLEAS, AND TERMITES**

5 **Background of the Invention**


This invention provides novel benzoylphenylurea insecticides and novel methods of inhibiting cockroaches, ants, fleas, and termites.

10 A broad class of benzoylphenylurea insecticides is disclosed in U.S. Patent No. 3,748,356. European Patent Application 263438 discloses that certain N-substituted phenyl-N'-substituted benzoyl-N-methylureas are highly safe to beneficial aquatic Crustacea while exhibiting equal or superior insecticidal activities to non-alkylated analogs. Hexaflumuron, a commercially significant benzoylphenylurea, is disclosed
15 in U.S. Patent No. 4,468,405. Use of hexaflumuron in methods of controlling termites is disclosed in WO 93/24011. Use of hexaflumuron to control cockroaches is disclosed in WO 94/03066. The compound N-[3,5-dichloro-2-fluoro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]-N'-(2,6-difluorobenzoyl)urea is disclosed in DE 3827133 and European Patent Application A 243,790, but there was no
20 disclosure of the unexpected activity of the compound against cockroaches, ants, fleas, and termites.

We have discovered that certain benzoylphenylurea compounds, some of which are novel, have substantially greater activity against cockroaches, ants, fleas, and
25 termites than would have been expected based on comparison with the closest prior art, i.e., hexaflumuron. Another significant property of the novel compounds of the invention is their surprisingly low toxicity to *Daphnia*.

Summary of the Invention

The invention provides a method of controlling cockroaches, ants, fleas, or termites which comprises delivering a effective amount of a compound of the formula (I):



wherein R¹ and R² are H, methyl, or ethyl,

to a location where control of cockroaches, ants, fleas, or termites is desired.

More specifically, the invention provides:


10 A method of controlling cockroaches which comprises a compound of the formula (I), in an amount effective to control cockroaches, to a location where control of cockroaches is desired;

A method of controlling ants which comprises delivering a compound of the formula (I), in an amount effective to control ants, to a location where control of ants is desired.

15 A method of controlling fleas which comprises delivering a compound of the formula (I), in an amount effective to control fleas, to a location where control of fleas is desired; and

20 A method of controlling termites which comprises delivering a compound of the formula (I), in an amount effective to control termites, to a location where control of termites is desired.


The invention also provides novel compounds of the formula (II)



25 wherein R^{1'} and R^{2'} are H, methyl, or ethyl, provided that at least one of R^{1'} and R^{2'} is methyl or ethyl.

Detailed Description of the Invention

Intermediate 1: 2,6-difluorobenzoyl isocyanate



5

A mixture of 0.52 g of 2,6-difluorobenzamide and 0.33 ml of oxalyl chloride was stirred under reflux in 15 ml 1,2-dichloroethane overnight. Solvent was removed under vacuum and 10 ml 1,2-dichloroethane was added. Solvent was removed under vacuum to leave the title intermediate, which could be used directly or dissolved in

10 1,2-dichloroethane and stored for future use.

Intermediate 2: 3,5-Dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)aniline




X522838

To a solution of 2.0 g of 4-amino-2,6-dichlorophenol in 40 mL tetrahydrofuran at room temperature was added 0.7 g of 87% potassium hydroxide. The mixture was warmed to 40° C. and stirred for 10 minutes, then chilled to 0° C.

Hexafluoropropene was bubbled in for 5 minutes, and the mixture stirred at room temperature over night. It was then concentrated under vacuum to dryness. The residue was dissolved in 50 mL dichloromethane and washed with 20 mL brine solution. The organic layer was separated and filtered through phase separation filter paper and then concentrated under vacuum to an oil. This was diluted with 50 mL dichloromethane and 50 mL heptane and re concentrated to give 3.05 g of 3,5-dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)aniline as a brown oil. Proton and ¹⁹F nmr spectra were consistent with the proposed structure.

25

Intermediate 3: 3,5-Dichloro-2-fluoro-4-(1,1,2,3,3,3-hexafluoropropoxy)aniline


To a solution of 2.50 g 3,5-dichloro 4-(1,1,2,3,3,3-hexafluoropropoxy)aniline in 60 mL acetonitrile under an atmosphere of nitrogen at room temperature there was

5 added 2.57 g 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2] octane bis (tetrafluoroborate) SELECTFLUOR™ (Air Products) portion wise over a 10 minute period. The mixture was warmed to 70° C. over a one hour period, then cooled to room temperature and poured into 100 mL of saturated sodium bicarbonate solution. The product was extracted with 150 mL ethyl acetate. The organic layer

10 was separated, washed with 50 mL of brine solution, separated, and dried over magnesium sulfate. The magnesium sulfate was removed by filtration, and the filtrate was concentrated under vacuum to give a dark oil 2.52 g. The product was chromatographed using a Michel-Miller low pressure silica gel column eluted with 6:1 heptane/ethyl acetate. Like fractions were pooled and concentrated under

15 vacuum to a brown oil 1.19 g. The proton and ¹⁹F nmr spectra were consistent with the proposed structure. Anal. calcd C₉H₄Cl₂F₇N₁O₁: C, 31.24; H, 1.17; N, 4.05. Found: C, 31.52; H, 1.15; N, 4.02.

20 **Intermediate 4: 3,5-Dichloro-2-fluoro-4-(1,1,2,3,3,3-hexafluoropropoxy)-N-ethyl aniline**




To a solution of 0.28 g 3,5-dichloro-2-fluoro-4-(1,1,2,3,3,3-hexafluoropropoxy) aniline in 8 mL glacial acetic acid at room temperature under an atmosphere of

nitrogen was slowly added 0.31 g sodium borohydride. The addition was carried out over a 1.5 hour period through a solid addition funnel between 25-34° C. with ice water cooling. The mixture was stirred at room temperature over night. Then the reaction mixture was added to 80 mL water and the pH was carefully adjusted to 7
 5 by adding solid sodium carbonate. The product was extracted with 80 mL ethyl acetate. The organic layer was separated, washed with brine solution, separated, and dried over magnesium sulfate. The magnesium sulfate was removed by filtration, and the filtrate was concentrated under vacuum to give a brown oil 0.29 g. The product was chromatographed using a Michel-Miller low pressure silica gel column
 10 eluted with 9:1 heptane/ethyl acetate. Like fractions were pooled and concentrated under vacuum to give 0.19 g of a tan oil. ¹H-NMR d 1.28 (t, 3H), 3.17 (q, 4H), 3.95 (bs, 1H), 4.94-5.21 (md, 1H), 6.59 (d, 1H). Anal. calcd C₁₁H₈Cl₂F₇N: C, 35.31; H, 2.16; N, 3.74. Found: C, 35.32; H, 2.14; N, 3.74.

15

Preparation of Products

Compound 1: N-[3,5-Dichloro-2-fluoro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]-N'-(2,6-difluorobenzoyl)urea




20 To a solution of 1.31 g 3,5-dichloro-2-fluoro-4-(1,1,2,3,3,3-hexafluoropropoxy) aniline in 5 mL 1,2-dichloroethane under an atmosphere of nitrogen at room temperature was added 0.87 g 2,6-difluorobenzoyl isocyanate dissolved in 10 mL dichloroethane dropwise over a 10 minute period. The mixture was stirred, warmed to 40° C. for a one hour period, then concentrated under vacuum to a brown solid 2.0
 25 g. The mixture was chromatographed using a Michel-Miller low pressure silica gel column eluted with 4:1 dichloromethane/heptane. Like fractions were pooled and concentrated under vacuum to give 1.67 g of a light tan solid, mp 156-7° C. Proton

and ^{19}F nmr spectra were consistent with the proposed structure. Anal. calcd $\text{C}_{17}\text{H}_7\text{Cl}_2\text{F}_9\text{N}_2\text{O}_3$: C, 38.58; H, 1.33; N, 5.29. Found: C, 38.64; H, 1.40; N, 5.44.

Compound 2: N-[3,5-Dichloro-2-fluoro-4-(1,1,2,3,3,3-


5 hexafluoropropoxy)phenyl]-N'-(2,6-difluorobenzoyl)-N-ethylurea



To a solution of 1.00 g 3,5-dichloro-2-fluoro-4-(1,1,2,3,3,3-hexafluoropropoxy)-N-ethyl aniline in 2 mL 1,2-dichloroethane under an atmosphere of nitrogen at room temperature there was added 0.54 g 2,6-difluorobenzoyl isocyanate dissolved in 6 mL dichloroethane dropwise over a 10 minute period. The mixture was stirred and warmed to 40° C. for a 1.5 hour period. Analysis by thin layer chromatography silica gel 4:1 heptane/ethyl acetate indicated incomplete reaction. To the mixture was added 0.13 g 2,6-difluorobenzoyl isocyanate in 1.5 mL 1,2-dichloroethane, and the mixture was warmed at 40° C. for 2.5 hour. Analysis by TLC indicated complete reaction. The reaction mixture was concentrated under vacuum to give an oil 1.64 g, which was chromatographed using a Michel-Miller low pressure silica gel column eluted with 4:1 heptane/ethyl acetate. Like fractions were pooled and concentrated under vacuum to give a white solid 0.87 g, mp 106-14° C. Proton nmr and mass spectra were consistent with the proposed structure. Anal. calcd $\text{C}_{19}\text{H}_{11}\text{Cl}_2\text{F}_9\text{N}_2\text{O}_3$: C, 40.95; H, 1.99; N, 5.03. Found: C, 40.89; H, 1.92; N, 5.03.

Compound 3: N-[3,5-Dichloro-2-fluoro-4-(1,1,2,3,3,3-

hexafluoropropoxy)phenyl]-N'-(2,6-difluorobenzoyl)-N'-ethyl urea



Dissolve 1.65 g N-[3,5-dichloro-2-fluoro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]-N'-(2,6-difluorobenzoyl) urea in 10 mL N,N-dimethylformamide and add 0.22 g 87%

potassium hydroxide. Chill to 0° C. and add 0.97 g iodoethane. Stir at 0° C. for a 7 hour period. Store in freezer at 0° C. over night. Analysis by thin layer chromatography silica gel dichloromethane shows product forming with starting urea remaining. The reaction was treated as above at 0° C. for the next two days.

- 5 Additional amounts of iodoethane, 1.94 g, and 0.10 g 87% potassium hydroxide were added. Pour reaction mixture into 50 mL brine solution and extract with 80 mL ethyl acetate. Separate organic layer and dry over magnesium sulfate. Filter drying agent and concentrate the filtrate under vacuum to a red oil 2.63 g. Chromatograph using a Michel-Miller low pressure silica gel column and elute with 2:1
- 10 heptane/dichloromethane. Pool like fractions and concentrate under vacuum to 0.18 g of a pink solid, mp 105-06° C. Proton nmr and mass spectra were consistent with the proposed structure. Anal. calcd C₁₉H₁₁Cl₂F₉N₂O₃: C, 40.95; H, 1.99; N, 5.03. Found: C, 40.99; H, 1.85; N, 4.98.

15

Biological Activity

German cockroach 2nd instars (*Blattella germanica*)

Continuous, low-dose ingestion exposure (treated cornmeal)

Rates: 0.19, 0.78, 3.12, 12.5, 50, 200 ppm

Compound	LC ₅₀ (ppm)	
	21 days	42 days
Compound 1	< 2.2	<0.78
Compound 2	0.35	0.26
Compound 3	8.0	1.5
hexaflumuron	> 200	> 200

- 20 Under continuous exposure, Compounds 1, 2, and 3 were far more active than hexaflumuron.

German cockroach 2nd instars (*Blattella germanica*)

Limited ingestion exposure (48 hr) to treated cornmeal

Rates: 1, 10, 100, 1000, 10000 ppm

Compound	LC ₅₀ (ppm)	
	21 days	42 days
Compound 1	< 22.2	< 9.1
Compound 2	21.1	12.8
hexaflumuron	> 10,000	> 10,000

Under limited exposure, Compounds 1 and 2 were more potent than hexaflumuron at both 21 and 42 days after exposure.

Cat Flea (*Ctenocephalides felis*)

Continuous exposure of larvae to treated media, impact on subsequent adult emergence

Rates: 0.1, 1.0, 10, 100, 1000 ppm

Compound	LC ₅₀ (ppm)	LC ₉₀ (ppm)
	Compound 1	2.8
Compound 2	12.7	18.6
hexaflumuron	65.7	333.5

Compounds 1 and 2 were both far more efficacious than hexaflumuron against cat fleas.

Subterranean Termite (*Reticulitermes flavipes*)

Continuous exposure (56 days) to treated paper

Rates: 0.78, 3.12, 12.5, 50, 200 ppm

Compound	LC ₅₀ (ppm)		LT ₅₀ (days) for 200 ppm trt
	35 days	56 days	
Compound 1	31.2	< 0.78	27.6
Compound 2	< 0.78	< 0.78	not calc.
hexaflumuron	> 200	1.3	33.8

Under continuous exposure, Compounds 1 and 2 were more potent and quicker acting than hexaflumuron.

Subterranean Termite (*Reticulitermes flavipes*)

- 5 Limited exposure (7 days) with mortality determined at 14, 28, 42, and 56 days 33399

Compound	LT ₅₀ (days) for 10000 ppm treatment
Compound 1	23.7
hexaflumuron	32.9

Under limited exposure, a high rate of Compound 1 induced more mortality earlier than did hexaflumuron.

Ant Studies

- Laboratory ant bait studies were carried out with Red Imported Fire Ant (RIFA) (10) (*Solenopsis invicta*) and Pharaoh Ant (*Monomorium pharaonis*). Chitin synthesis inhibitors, such as the compounds of the invention, control ants by killing the molting larvae and/or pupae and potentially preventing the hatching of eggs. Because adult workers are not affected, control is measured by effects on the brood. The studies involved 3-4 day exposure to bait. These limited exposure studies more (15) accurately represent real world bait availability than continuous exposure.

Compound	Concentration tested	Species	Time to Achieve 50% Brood Reduction	Time to Achieve 90% Brood Reduction*
Compound 1	0.1%*	RIFA	2 wks	3 wks
	0.1%	Pharaoh	4 wks	NA (70% @ 13 wks)
Hexaflumuron	0.1%	RIFA	NA	NA
	0.25%	RIFA	4 wks	10 wks
	0.1%	Pharaoh	NA	NA

* Only concentration tested. NA = did not achieve specified percent brood reduction. Percent brood reduction achieved at end of study listed in parenthesis.

Compound 1 is significantly more potent than hexaflumuron based on a short exposure study with RIFA.

Activity against *Daphnia*

5 48 hr exposure to treated water

Compound	LC ₅₀ (ppb) in water
Compound 1	5.0
Compound 2	> 100
Compound 3	> 100
Hexaflumuron	68.1

Compounds 2 and 3, the alkylated derivatives of Compound 1, were much less active against *Daphnia* than were Compound 1 and hexaflumuron; this is surprising since the activities of Compound 1, 2, and 3 in cockroaches are quite similar.

10

Formulations

In order to facilitate the application of the compounds of formula (I) to the desired locus, or to facilitate storage, transport or handling, the compound is normally formulated with a carrier and/or a surface-active agent.

15 A carrier in the present context is any material with which the compound of formula (1)(active ingredient) is formulated to facilitate application to the locus, or storage, transport or handling. A carrier may be a solid or a liquid, including a material which is normally gaseous but which has been compressed to form a liquid. Any of the carriers normally used or known to be usable in formulating insecticidal compositions may be used.

20 Compositions according to the invention contain 0.0001 to 99.9% by weight active ingredient. Preferably, compositions according to the invention contain 0.001 to 10.0% by weight of active ingredient though proportions as low as 0.0001% may be useful in some circumstances.

- Suitable solid carriers include natural and synthetic clays and silicates, for example natural silicas such as diatomaceous earths; magnesium silicates, for example talcs; magnesium aluminium silicates, for example attapulgites and vermiculites; aluminium silicates, for example kaolinites, montmorillonites and micas; calcium carbonate; calcium sulphate; ammonium sulphate; synthetic hydrated silicon oxides and synthetic calcium or aluminium silicates; elements, for example carbon and sulfur; natural and synthetic resins, for example coumarone resins, polyvinyl chloride, and styrene polymers and copolymers; solid polychlorophenols; bitumen; waxes; agar; and solid fertilizers, for example superphosphates. Cellulose based materials, for example wood, sawdust, agar, paper products, cotton linter, or Methocel[®], as well as the other solid carriers that are themselves attractive to or at least non-repellant to termites are particularly suitable and preferable. Mixtures of different solids are often suitable. For example, a mixture of wood flour and agar formulated as a moisture containing solid would be preferable.
- Suitable liquid carriers include water; alcohols, for example isopropanol and glycols; ketones, for example acetone, methyl ethyl ketone, methyl isobutyl ketone, isophorone and cyclohexanone; ethers; aromatic or aliphatic hydrocarbons, for example benzene, toluene and xylene; petroleum fractions, for example kerosene and light mineral oils; chlorinated hydrocarbons, for example carbon tetrachloride, perchloroethylene and trichloroethane; polar organic liquids, such as dimethyl formamide, dimethyl acetamide, dimethyl sulfoxide and N-methylpyrrolidone; oils derived from plants, such as corn oil and peanut oil. Mixtures of different liquids are often suitable, for example a mixture of isophorone with a polar organic solvent such as N-methylpyrrolidone, as are mixtures of solid and liquid carriers.
- Pesticidal compositions are often formulated and transported in a concentrated form which is subsequently diluted by the user before application. The presence of small amounts of a carrier which is a surface-active agent facilitates this process of dilution. Thus it is suitable to use at least one carrier in such a composition which is a surfaceactive agent. For example, the composition may contain at least two carriers, at least one of which is a surface-active agent.

A surface-active agent may be an emulsifying agent, a dispersing agent or a wetting agent; it may be nonionic or ionic. Examples of suitable surface-active agents

include the sodium or calcium salts of polyacrylic acids and lignin sulfonic acids; the condensation of fatty acids or aliphatic amines or amides containing at least 12 carbon atoms in the molecule with ethylene oxide and/or propylene oxide; fatty acid esters of glycerol, sorbitol, sucrose or pentaerythritol; condensates of these with
5 ethylene oxide and/or propylene oxide; condensates of these with ethylene oxide and/or propylene oxide; condensation products of fatty alcohol or alkyl phenols, for example p-octylphenol or p-octylcresol, with ethylene oxide and/or propylene oxide; sulfates or sulfonates of these condensation products; alkali or alkaline earth metal salts, preferably sodium salts, or sulfuric or sulfonic acid esters containing at least 10
10 carbon atoms in the molecule, for example sodium lauryl sulphate, sodium secondary alkyl sulfates, sodium salts of sulfinated castor oil, and sodium alkylaryl sulfonates such as dodecylbenzene sulfonate; and polymers of ethylene oxide and copolymers of ethylene oxide and propylene oxide.

Pesticidal compositions may for example be formulated as wettable powders, dusts,
15 granules, baits, solutions, emulsifiable concentrates, emulsions, suspension concentrates and aerosols.

Wettable powders usually contain 25, 50 or 75% weight of active ingredient and usually contain in addition to solid inert carrier, 3-10% weight of a dispersing agent and, where necessary, 0-10% weight of stabilizer(s) and/or other additives such as
20 penetrants or stickers.

Dusts are usually formulated as a dust concentrate having a similar composition to that of a wettable powder but without a dispersant, and are diluted in the field with further solid carrier to give a composition usually containing 0.5-10% weight of active ingredient.

25 Granules are usually prepared to have a size between 10 and 100 BS mesh (1.676-0.152 mm), and may be manufactured by, for example, agglomeration or impregnation techniques. Generally, granules will contain 0.01-75% weight active ingredient and 0-10% weight of additives such as stabilizers, surfactants, slow release modifiers and binding agents. The so-called "dry flowable powders" consist
30 of relatively small granules having a relatively high concentration of active ingredient. Of particular interest in current practice are the water dispersible granular formulations. These are in the form of dry, hard granules that are essentially

dust-free, and are resistant to attrition on handling, thus minimizing the formation of dust. On contact with water, the granules readily disintegrate to form stable suspensions of the particles of active material. Such formulation contain 90% or more by weight of finely divided active material, 3-7% by weight of a blend of
5 surfactants, which act as wetting dispersing, suspending and binding agents, and 1-3% by weight of a finely divided carrier, which acts as a resuspending agent.

Baits are prepared by, for example, combining a mixture of a suitable food source, such as sawdust for termites or grain or meal for cockroaches, with an amount of active ingredient sufficient to provide the desired result; for example, from about
10 0.001% to about 20% weight active ingredient and forming the mixture into a paste by the addition of about 1% to 5% of a water based binder such as agar. The paste-like mixture may be applied as is or may be packed into a housing such as a hollowed out wooden dowel or a plastic tube or bait station. In other embodiments, sheets of paper or cardboard can be sprayed with or dipped in a diluted formulation
15 containing the active ingredient. Baits are a preferable embodiment of the present invention.

Emulsifiable concentrates usually contain, in addition to a solvent and, when necessary, co-solvent, 10-50% weight per volume active ingredient, 2-20% weight per volume emulsifiers and 0-20% weight per volume of other additives such as
20 stabilizers, penetrants and corrosion inhibitors.

Suspension concentrates are usually compounded so as to obtain a stable, non-sedimenting flowable product and usually contain 10-75% weight active ingredient, 0.5-15% weight of dispersing agents, 0.1-10% weight of suspending agents such as protective colloids and thixotropic agents, 0-10% weight of other
25 additives such as defoamers, corrosion inhibitors, stabilizers, penetrants and stickers, and water or an organic liquid in which the active ingredient is substantially insoluble; certain organic solids or inorganic salts may be present dissolved in the formulation to assist in preventing sedimentation or as anti-freeze agents for water.

Aqueous dispersions and emulsions are compositions which may be obtained by
30 diluting a wettable powder or a concentrate with water. The said emulsions may be of the water-in-oil or of the oil-in-water type, and may have a thick 'mayonnaise'-like consistency.

The method of applying a compound of Formula (I) to combat termites comprises applying the compound, conveniently in a composition comprising the compound of Formula (I) and a carrier as described above, to a locus or area to be treated for the termites, such as soil or timber, already subject to infestation or attack by termites or intended to be protected from infestation by termites. The active ingredient is, of course, applied in an amount sufficient to effect the desired action of combatting termite infestation. This dosage is dependent upon many factors, including the carrier employed, the method and conditions of the application, whether the formulation is present at the locus in the form of a film, or as discrete particles or as a bait, the thickness of film or size of particles, the degree of termite infestation, and the like.

Proper consideration and resolution of these factors to provide the necessary dosage of the active ingredient at the locus to be protected are within the skill of those versed in the art. In general, however, the effective dosage of the compound of the invention at the locus to be protected--i.e., the dosage to which the termite has access--is of the order of 0.001 to 1.0% based on the total weight of the composition, though under some circumstances the effective concentration may be as little as 0.0001% or as much as 2%, on the same basis.

When used to control cockroaches, it is preferred to use the active ingredient in a treated bait or as a surface treatment.

When used to control ants, it is preferred to use the active ingredient in a liquid bait or granular bait.

When used to control termites, it is preferred to use the active ingredient in a cellulose based bait.


When used to control fleas, it is preferred to use the active ingredient on a treated substrate.

30

Suitable formulations include granular, paste, or dust cockroach bait, SP or WP cockroach and/or flea sprayables, cellulose-based termite baits, liquid or granular ant baits, feed-through or topical animal treatment for fleas.


We claim:

1. A method of controlling cockroaches which comprises a compound of the formula (I)



wherein R^1 and R^2 are H, methyl, or ethyl,
in an amount effective to control cockroaches,
to a location where control of cockroaches is desired.


10 2. A method of controlling ants which comprises delivering a compound of the formula (I)



wherein R^1 and R^2 are H, methyl, or ethyl,
in an amount effective to control ants,
to a location where control of ants is desired.


15

3. A method of controlling fleas which comprises delivering a compound of the formula (I)



20 wherein R^1 and R^2 are H, methyl, or ethyl,
in an amount effective to control fleas,
to a location where control of fleas is desired.


4. A method of controlling termites which comprises delivering a compound of the formula (I)



wherein R¹ and R² are H, methyl, or ethyl,

- 5 in an amount effective to control termites,
to a location where control of termites is desired.

5. A compound of the formula (II)



- 10 wherein R^{1'} and R^{2'} are H, methyl, or ethyl, provided that at least one of R^{1'} and R^{2'} is methyl or ethyl.

6. A pesticide composition which comprises an effective amount of a compound of claim 5 in combination with a carrier.

7. A cockroach bait comprising a compound of claim 5 in a conventional
15 bait.

8. An ant bait comprising a compound of claim 5 in combination with a conventional bait.

9. A termite bait comprising a compound of claim 5 in combination with a conventional bait.

INTERNATIONAL SEARCH REPORT

Int. (ional Application No

PCT/US 97/18199

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 A01N47/34 C07C275/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 38 27 133 A (CIBA-GEIGY) 23 February 1989 cited in the application see page 4, line 50 - line 64; claims; example 1.2.1	1-9
A	DE 27 26 684 A (HOECHST) 4 January 1979 see page 3, paragraph 2 - page 4, paragraph 2; examples 6,17,28	1-9
A	PATENT ABSTRACTS OF JAPAN vol. 014, no. 126 (C-0699), 9 March 1990 & JP 02 003659 A (ISHIHARA SANGYO KAISHA LTD), 9 January 1990, see abstract	1-9
	--- -/--	

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 February 1998

Date of mailing of the international search report

02/03/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

Int. l. Application No.
PCT/US 97/18199

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 468 405 A (R.H RIGTERINK, ET AL.) 28 August 1984 cited in the application see column 1; example 3 ---	1-9
A	US 4 170 657 A (R.H. RIGTERINK) 9 October 1979 see column 1 ---	1-9
A	WO 94 03066 A (CIBA-GEIGY) 17 February 1994 cited in the application see page 2 - page 3 ---	1,5-7
A	US 5 556 883 A (E.M. THOMS, ET AL.) 17 September 1996 see claims -----	4-6,9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/18199

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 3827133 A	23-02-89	NONE	
DE 2726684 A	04-01-79	NONE	
US 4468405 A	28-08-84	AU 565944 B AU 5832986 A AU 553953 B AU 8657982 A BR 8204510 A CA 1225401 A CA 1236112 C DK 338782 A EP 0071279 A JP 1493880 C JP 58026858 A JP 63040422 B JP 1581947 C JP 2006749 B JP 62123155 A US 4536341 A,B	01-10-87 20-11-86 31-07-86 03-02-83 26-07-83 11-08-87 03-05-88 31-01-83 09-02-83 20-04-89 17-02-83 11-08-88 11-10-90 13-02-90 04-06-87 20-08-85
US 4170657 A	09-10-79	AU 3482678 A BE 867046 A CA 1122612 A DE 2820696 A DK 212478 A EG 13720 A FR 2390423 A GB 1575799 A IN 146988 A JP 53141242 A NL 7805121 A US RE32418 E ZA 7801892 A	11-10-79 13-11-78 27-04-82 16-11-78 14-11-78 31-12-82 08-12-78 01-10-80 20-10-79 08-12-78 15-11-78 12-05-87 28-03-79
WO 9403066 A	17-02-94	AU 4571493 A	03-03-94
US 5556883 A	17-09-96	US 5691383 A AT 161143 T AU 660821 B	25-11-97 15-01-98 06-07-95

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int: lonal Application No

PCT/US 97/18199

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5556883 A		AU 4248793 A	30-12-93
		CA 2136985 A	09-12-93
		CN 1081318 A	02-02-94
		DE 69315834 D	29-01-98
		EP 0643561 A	22-03-95
		JP 7507307 T	10-08-95
		MX 9303230 A	01-12-93
		NZ 252604 A	25-06-96
		WO 9324011 A	09-12-93
