International Bureau

(43) International Publication Date 18 October 2018 (18.10.2018)





(10) International Publication Number WO 2018/189738 Al

(51) International Patent Classification:

A23J 3/22 (2006.01)

A23L 13/00 (2016.01)

(21) International Application Number:

PCT/IL2018/050398

(22) International Filing Date:

04 April 2018 (04.04.2018)

(25) Filing Language:

English

(26) **Publication Language:** 

English

(30) Priority Data:

62/483,399

09 April 2017 (09.04.2017)

US

- (71) Applicant: SUPERMEAT THE ESSENCE OF MEAT LTD. [IL/IL]; 3 Pinhas Sapir Street, 7403626 Nes Ziona (IL).
- (72) Inventors: SAVIR, Ido; 12 Yotam Street, 4670130 Herzliya (IL). FRIEDMAN, Shir; 7 Nahal Shikma Street, 7175459 Modiin-Maccabim-Reut (IL). BARAK, Koby; 36 Aba-Hilel Street, 5253209 Ramat-Gan (IL).
- (74) Agent: EHRLICH, Gal et al; G. E. EHRLICH (1995) LTD., 11 Menachem Begin Road, 5268104 Ramat Gan (IL).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

#### **Declarations under Rule** 4.17:

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(in))
- f inventorship (Rule 4.17(iv))

#### Published:

with international search report (Art. 21(3))

(54) Title: CULTURED MEAT-CONTAINING HYBRID FOOD

(57) Abstract: A method of producing a hybrid foodstuff is provided. The method comprises combining a plant-originated substance with an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein the animal cells do not form a tissue, and wherein the amount is below 30 % (w/w) of the hybrid foodstuff.

WO 2018/189738 PCT/IL2018/050398

# CULTURED MEAT-CONTAINING HYBRID FOOD

#### FIELD AND BACKGROUND OF THE INVENTION

5

10

15

20

25

30

The present invention, in some embodiments thereof, relates to cultured meat-containing hybrid food.

In recent years, attempts have been made to bring natural taste, odor, appearance and nutritional value of meat to the process of vegetarian food preparation, under the fields of food science, food technology and molecular gastronomy.

Cultured meat, also called synthetic meat, cell-cultured meat, clean meat, and in vitro meat, is meat grown in cell culture instead of inside animal, see Verbeke, Wim, Pierre Sans, and Ellen J. Van Loo. "Challenges and prospects for consumer acceptance of cultured meat." Journal of Integrative AgriculturelA.2 (2015): 285-294, which is incorporated herein as a reference. Several patent documents describe a variety of cultured meat consumable products and related method: US patent app. Nos. 2005010965, 2006121006, and 2006029922 disclose process for production and cultured meat which comprises muscle cells that are grown ex vivo attached to either 2D or 3D support structure and further comprises other cells such as fat cells or cartilage cells, or both, that are grown ex vivo together with the muscle cells. The visual appearance (including color), smell, texture taste and price-per-product of the cultured meat published in the art is not yet satisfying, see Post, Mark J. "Cultured meat from stem cells: Challenges and prospects." Meat Science 92.3 (2012): 297-301; Verbeke, Wim, et al. "Would you eat cultured meat?': Consumers' reactions and attitude formation in Belgium, Portugal and the United Kingdom." Meat science 102 (2015): 49-58; and Forgacs, Gabor, Francoise Marga, and Karoly Robert Jakab. "Engineered comestible meat." U.S. Patent No. 8,703,216. 22 Apr. 2014, all incorporated herein as a reference.

It therefore remains a long felt and unmet need to provide a novel non-meat production method which is more cost-efficient than the current methods of production and non-meat products which are healthier having an improved meaty flavor and appearance.

Additional background art:

US Patent Application Publication No. 20060121006;

U.S. Patent No. 9,332,779;

US Patent Application Publication No. 201 10301249

# SUMMARY OF THE INVENTION

5

10

15

20

25

30

According to an aspect of some embodiments of the present invention there is provided a method of producing a hybrid foodstuff, the method comprising combining a plant-originated substance with an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein the animal cells do not form a tissue, and wherein the amount is below 30 % (w/w) of the hybrid foodstuff.

According to an aspect of some embodiments of the present invention there is provided a method of providing nutrition to a subject in need thereof, the method comprising providing the subject with a hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein the animal cells do not form a tissue, and wherein the amount is below 30 % (w/w) of the hybrid foodstuff.

According to some embodiments of the invention, the subject is at risk of nutritional deficiency.

According to an aspect of some embodiments of the present invention there is provided a method of qualifying a hybrid foodstuff, the method comprising subjecting the hybrid foodstuff to at least one organoleptic test.

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein the animal cells do not form a tissue, and wherein the amount is below 30 % (w/w) of the hybrid foodstuff.

According to some embodiments of the invention, the hybrid foodstuff comprises from about 82% to about 99.5% (w/w) plant-originated substance and/or additives thereof and about 0.5% to about 18% (w/w) cultured animal cells.

According to some embodiments of the invention, the amount is minimal weight percent.

According to some embodiments of the invention, the amount provides the hybrid foodstuff with one 9-PHS point more than a corresponding foodstuff devoid of the animal cells.

According to some embodiments of the invention, the foodstuff comprises about 70 % to about 99.7% (w/w) plant-originated substances and/or additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 0.3% to 30% (w/w), where the improvement is that the oil portion consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured animal cells.

According to some embodiments of the invention, the cultured animal cells are selected from the group consisting of hepatocytes, myoblasts, osteoblasts, fibroblasts, lipoblasts, odontoblasts, adult neuronal progenitor cells, neural stem cells, multipotent stem cells from subventricular forebrain region, ependymal-derived neural stem cells, hematopoietic stem cells, liver-derived hematopoietic stem, marrow-derived stem cell, adipo-fibroblasts, adipose-derived stem cells, islet- cells producing stem cells, pancreatic-derived pluripotent islet-producing stem cells, mesenchymal stem cells, placenta cells, bone marrow stromal cells, muscle side population cells, bone marrow-derived recycling cells, blood-derived mesenchymal precursor cells, bone-marrow derived side population cells, muscle precursor cells, circulating skeleton stem cells, neural progenitor cells, multipotent adult progenitor cells, mesodermal progenitor cells, spinal cord progenitor cells and spore-like cell, and any combinations thereof.

5

10

15

20

25

30

According to some embodiments of the invention, the hybrid foodstuff has the organoleptic properties of a meat product, selected from the group consisting of a chicken, duck, pork and beef.

According to some embodiments of the invention, the hybrid foodstuff comprises less than about 0.5 gr per 100 gr saturated fat, less than about 60 mg per 100 gr cholesterol and less than about 220 mg per 100 gr sodium.

According to some embodiments of the invention, the animal cells are of a single cell type or single cell lineage.

According to some embodiments of the invention, the animal cells are of no more than two cell types or two cell lineages.

According to some embodiments of the invention, the animal cells are of no more than three cell types or three cell lineages.

According to some embodiments of the invention, the foodstuff is free of growth hormones in an amount hazardous for human or animal consumption.

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff comprising from about 82% to about 99.5% (w/w) plant-originated substances and additives thereof and about 0.5% to about 18% (w/w) cultured meat.

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff of claim 17, wherein the portion of the cultured meat in the end product is defined as the minimal weight percent providing the end product with one 9-PHS point more than the corresponding foodstuff being non-hybrid, cultured meat free vegetable-based product.

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff with improved organoleptic and nutritional properties, comprising about 70%

to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 0.3% to 30% (w/w), where the improvement is that the oil portion consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat.

5

10

15

20

25

30

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff with improved organoleptic and nutritional properties, comprising about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 3% to 30% (w/w), where the improvement is that the oil portion consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat; wherein the smallest portion of the cultured meat in the end product is defined as the minimal weight percent providing the end product with at least one 9-PHS point more than a corresponding foodstuff being non-hybrid, cultured meat free vegetable-based product.

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff with improved organoleptic and nutritional properties, characterized by about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 3% to 30% (w/w); the oil portion consists of up to about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat; wherein the role the mixtures in the oil portion (measured as vegetable oil to culture meat (wt/wt) in improving the organoleptic properties (measured as 9-PHS) of the hybrid foodstuff of is provided by any of Formula I, II or III.

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff with improved organoleptic and/or nutritional properties, characterized by about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (e.g., gum, binding, thickening, aroma flavor additives) and/or water immiscible materials (oil portion), ranging from about 3% to 30% (w/w), where the improvement is that the additives thereof (e.g., gum, binding, thickening, aroma flavor additives) and/or water immiscible materials (oil portion)consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat.

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff according to any of claim 17 to 22, wherein at least about 50% of cultured meat portion is cultured meat cells selected from a group consisting of heme-proteins (HPs), hepatocytes, myoblasts, osteoblasts, fibroblasts, lipoblasts, odontoblasts, adult neuronal progenitor cells, neural stem cells, multipotent stem cells from subventricular forebrain region,

ependymal-derived neural stem cells, hematopoietic stem cells, liver-derived hematopoietic stem, marrow-derived stem cell, adipo-fibroblasts, adipose-derived stem cells, islet- cells producing stem cells, pancreatic-derived pluripotent islet-producing stem cells, mesenchymal stem cells, placenta cells, bone marrow stromal cells, muscle side population cells, bone marrow-derived recycling cells, blood-derived mesenchymal precursor cells, bone-marrow derived side population cells, muscle precursor cells, circulating skeleton stem cells, neural progenitor cells, multipotent adult progenitor cells, mesodermal progenitor cells, spinal cord progenitor cells and spore-like cell, mixtures and any combinations thereof.

5

10

15

20

25

30

According to an aspect of some embodiments of the present invention there is provided a hybrid foodstuff according to any of claim 17 to 22, wherein the foodstuff is a consumable, edible item having the final organoleptic properties of a meat product, and especially a product selected from the group consisting of Beef, Beef heart, Beef liver, Beef tongue, Bone soup from allowable meats, Buffalo, Bison, Calf liver, Caribou, Goat, Ham, Horse, Kangaroo, Lamb, Marrow soup, Moose, Mutton, Opossum, Organ Meats, Pork, Bacon, Rabbit, Snake, Squirrel, Sweetbreads, Tripe, Turtle, Veal, Venison, Chicken, Chicken Liver, Cornish Game Hen, Duck, Duck Liver, Emu, Gizzards, Goose, Goose Liver, Grouse, Guinea Hen, Liver, Ostrich, Partridge, Pheasant, Quail, Squab, Turkey, any mixture and a combination thereof.

According to an aspect of some embodiments of the present invention there is provided a meat-flavored hybrid foodstuff-based diet free of harmful substances at harmful quantities; wherein the harmful substances including saturated fat, cholesterol and sodium; wherein harmful quantities are  $5 \, \text{g/d}$ ,  $<\!200 \, \text{mg}$  and  $400 \, \text{mg/d}$ , respectfully; and wherein the each of the hybrid foodstuff comprising from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.

According to an aspect of some embodiments of the present invention there is provided a beef-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein the harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <8 gr and <3.5 gr per 100 gr, respectfully; and wherein the beef-flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.

According to an aspect of some embodiments of the present invention there is provided a pork-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein the harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <5 gr and <1 gr per lOOg, respectfully; and wherein the pork-flavored hybrid

10

15

20

25

30

foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.

According to an aspect of some embodiments of the present invention there is provided a chicken-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein the harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <4 gr and <0.5 gr per IOOgr, respectfully; and wherein the chicken -flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.

According to an aspect of some embodiments of the present invention there is provided a duck-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein the harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <4 gr and <0.5 gr per 10Ogr, respectfully; and wherein the duck -flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.

According to an aspect of some embodiments of the present invention there is provided a method of preparing a hybrid foodstuff comprising the steps of:

providing a plant-originated substance;

providing cultured meat, cells and substances thereof;

interconnecting the plant-originated substance with the cultured meat, cells and substances thereof; and

forming the hybrid foodstuff product into a desired form.

According to some embodiments of the invention, the plant-originated substance is from about 82% to about 99.5% (w/w).

According to some embodiments of the invention, the cultured meat, cells and substances thereof is from about 0.5% to about 18% (w/w).

According to an aspect of some embodiments of the present invention there is provided a method of preparing a hybrid foodstuff comprising the steps of:

providing a plant-originated substance;

providing water immiscible materials;

providing cultured meat, cells and substances thereof;

interconnecting the plant-originated substance with the cultured meat, cells and substances thereof and with the water immiscible materials; and

forming the hybrid foodstuff product into a desired form.

According to some embodiments of the invention, the plant-originated substance is from about 70% to about 99.7% (w/w).

According to some embodiments of the invention, the water immiscible materials is from about 3% to 30% (w/w).

According to some embodiments of the invention, the cultured meat, cells and substances thereof is from about 0.3% to about 30% (w/w).

5

10

15

20

25

30

According to some embodiments of the invention, at least about 50% of cultured meat portion are hepatocytes.

According to some embodiments of the invention, at least about 50% of cultured meat portion is cultured meat cells selected from a group consisting of heme-proteins (HPs), hepatocytes, myoblasts, osteoblasts, fibroblasts, lipoblasts, odontoblasts, adult neuronal progenitor cells, neural stem cells, multipotent stem cells from subventricular forebrain region, ependymal-derived neural stem cells, hematopoietic stem cells, liver-derived hematopoietic stem, marrow-derived stem cell, adipo-fibroblasts, adipose-derived stem cells, islet- cells producing stem cells, pancreatic-derived pluripotent islet-producing stem cells, mesenchymal stem cells, placenta cells, bone marrow stromal cells, muscle side population cells, bone marrow-derived recycling cells, blood-derived mesenchymal precursor cells, bone-marrow derived side population cells, muscle precursor cells, circulating skeleton stem cells, neural progenitor cells, multipotent adult progenitor cells, mesodermal progenitor cells, spinal cord progenitor cells and spore-like cell, mixtures and any combinations thereof.

According to some embodiments of the invention, the foodstuff is a consumable, edible item having the final organoleptic properties of a meat product, and especially a product selected from the group consisting of Beef, Beef heart, Beef liver, Beef tongue, Bone soup from allowable meats, Buffalo, Bison, Calf liver, Caribou, Goat, Ham, Horse, Kangaroo, Lamb, Marrow soup, Moose, Mutton, Opossum, Organ Meats, Pork, Bacon, Rabbit, Snake, Squirrel, Sweetbreads, Tripe, Turtle, Veal, Venison, Chicken, Chicken Liver, Cornish Game Hen, Duck, Duck Liver, Emu, Gizzards, Goose, Goose Liver, Grouse, Guinea Hen, Liver, Ostrich, Partridge, Pheasant, Quail, Squab, Turkey, any mixture and a combination thereof.

According to some embodiments of the invention, the animal-derived protein in the hybrid foodstuff is below 50 % (w/w).

According to some embodiments of the invention, the foodstuff is not a chip and the plant-originated substance is not a plant derived hydrogel.

According to some embodiments of the invention, the hybrid foodstuff comprises the hybrid foodstuff as described herein.

10

15

20

25

30

PCT/IL2018/050398

According to some embodiments of the invention, the foodstuff is a vegetarian foodstuff.

According to some embodiments of the invention, the foodstuff is a vegan foodstuff.

According to some embodiments of the invention, the foodstuff is a meat substitute.

Unless otherwise defined, all technical and/or scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments of the invention, exemplary methods and/or materials are described below. In case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and are not intended to be necessarily limiting.

#### DESCRIPTION OF SPECIFIC EMBODIMENTS OF THE INVENTION

The present invention, in some embodiments thereof, relates to cultured meat-containing hybrid food.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not necessarily limited in its application to the details set forth in the following description or exemplified by the Examples. The invention is capable of other embodiments or of being practiced or carried out in various ways.

Whilst reducing the present invention to practice, the present inventors realized that enriching vegetarian foods with in vitro-cultured isolated animal cells that do not attempt to mimic a tissue structure can enhance organoleptic properties and nutritional properties of foods that are based essentially on plant-originated substances.

As used herein the term "organoleptic properties" refers to the aspects of food, that a consumer experiences via the senses—including taste, sight, smell or touch.

For example, adipocytes may be used to confer a non-meat food with a meaty taste and/or texture (e.g., beef or chicken) when cooked or grilled.

Cardiomyocytes may be used to confer a non-meat food with a texture of meat as well as taste (e.g., bitter/metal) of meat.

Erythrocytes may confer a non-meat food with a grilled meat (e.g., beef or chicken) taste as well as meaty color (e.g., beef).

Hepatocytes may confer a non-meat food with meaty color (e.g., beef) as well as enrich the taste of meat.

The term "meat" is meant to encompass any animal flash that is eaten as food (e.g., beef, pork, poultry, fish as well as additional examples which are provided hereinbelow).

In this case, the organoleptic property is that of meat. For instance, when referring to taste, an organoleptic property may be an umami taste.

As used herein "nutritional properties" refer to the composition of meat that is of valuable to the subject feeding on it.

5

10

15

20

25

For muscle tissue is very high in protein, containing all of the essential amino acids, and in most cases is a good source of zinc, vitamin  $B_{12}$ , selenium, phosphorus, niacin, vitamin  $B_6$ , choline, riboflavin and iron. Several forms of meat are also high in vitamin K. Muscle tissue is very low in carbohydrates and does not contain dietary fiber. Proteins, vitamins, and minerals are available from almost any source of meat and are generally consistent.

The fat content of meat can vary widely depending on the species and breed of animal, the way in which the animal was raised, including what it was fed, the anatomical part of the body, and the methods of butchering and cooking. Wild animals such as deer are typically leaner than farm animals, leading those concerned about fat content to choose game such as venison. The fatty deposits that exist with the muscle fibers in meats soften meat when it is cooked and improve the flavor through chemical changes initiated through heat that allow the protein and fat molecules to interact. The fat, when cooked with meat, also makes the meat seem juicier. However, the nutritional contribution of the fat is mainly calories as opposed to protein. As fat content rises, the meat's contribution to nutrition declines. In addition, there is cholesterol associated with fat surrounding the meat. The cholesterol is a lipid associated with the kind of saturated fat found in meat.

Table A below compares the nutritional content of several types of meat (per 110 gr). While each kind of meat has about the same content of protein and carbohydrates, there is a very wide range of fat content.

Table A

Source	Calories	Protein (gr)	Carbs (gr)	Fat (gr)
Fish	110-140	20-25	0	1-5
Chicken breast	160	28	0	7
Lamb	250	30	0	14
Steak (beef top	210	36	0	7
round)				
Steak (beef T-	450	25	0	35
bone)				

Other major nutritional ingredients and their sources are provided hereinbelow:

Vitamin B12, which is mainly found in fish, meat, poultry and dairy products.

Vitamin D, which is found in oily fish, eggs and dairy.

Docosahexaenoic acid (DHA), which is an essential omega-3 fat found in fatty fish.

Heme-iron, which is predominantly found in meat, especially red meat.

10

15

20

25

30

Zinc, which is mainly found in animal protein sources, such as beef, pork and lamb.

Hence, the improvement/enhancement in nutritional property can be in any of protein, calories, fat, vitamins, minerals such as listed herein.

As used herein "enhancement", "increase", "augmentation" refers to an addition of at least 10 %, 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 %, 55 %, 60 %, 65 %, 70 %, 75 %, 80 %, 85 %, 90 % 95 % 100 %, 200 %, 300 %, 400 %, 500 %, 600 %, 700 %, 800 %, 900 % or 1000 % (e.g., by weight or by score) of the organoleptic or nutritional property as compared to the same food without the cultured meat cells or in which a portion has been substituted with an equivalent amount of cultured meat cells (e.g., lipid portion, hydrolysate, synthetic additives etc.), as described below.

The terms 'hybrid food' and 'products' thereof refer hereinafter to a foodstuff or a portion of a foodstuff comprising at least one first non-meat significantly high portion and at least one second cultured meat low portion, combined (e.g., interconnected) with said at least one first non-meat portion.

According to a specific embodiment, the non-meat is a plant originated substance(s).

According to a specific embodiment, the non-meat is a non-plant originated substances (e.g., minerals, synthetic substance(s)).

According to a specific embodiment, the non-meat is selected from the group consisting of a plant originated substance(s) and non-plant-originated substance(s).

According to a specific embodiment, the foodstuff is a vegetarian foodstuff.

According to a specific embodiment, the foodstuff is a vegan foodstuff.

According to a specific embodiment, the foodstuff comprises a meat substitute or is generally consumed as a meat substitute (plant-based).

According to a specific embodiment, the animal cells are of a single cell type or single cell lineage.

As used herein "cell lineage" refers to any of endoderm, mesoderm and endoderm.

According to a specific embodiment, the animal cells are of no more than two cell types or two cell lineages.

According to a specific embodiment, the animal cells are of no more than three or four cell types.

According to a specific embodiment, the hybrid foodstuff is free of bodily fluids e.g., saliva, serum, plasma, mucus, urine, feces, tears, milk etc. As used herein "animal cells" refer to "non-human cells".

The term 'foodstuff refers hereinafter to any substance with food value; the term further refers to the raw material of food before, within the process and after processing, the food product or portions thereof (e.g., the coating of an edible item such as a schnitzel), by-product(s) and end-product (e.g., sausage, ground meat, schnitzel etc.) thereof. In the present invention, there are provided three families of foodstuffs: plant-originated foodstuffs and substances thereof, and cultured meat, cells and substances thereof, both are known in the art; and the novel hybrid foodstuff which comprises a majority (above 50 % w/w) of said plant-originated substances and a minority (below 50 % w/w) of cultured meat, cells and substances thereof interconnected with said plant-originated substances to form a unified foodstuff with an improved organoleptic and nutritional value.

According to an embodiment of the invention the hybrid foodstuff is an end article of manufacture (product) to be consumed by a human or non-human subject.

Examples of meat substitutes: natural, traditional and commercially made which are contemplated according to some embodiments of the invention:

Alpro and *Provamel*, both usually known for their plant milk range, also offer different vegetarian meat substitutes

Beanfeast

5

10

15

**Beyond Meat** 

Boca Burger

Falafel, a traditional Middle Eastern bean fritter, believed to have been created by ancient Copts as a meat substitute during Lent

Fistulina hepatica, common mushroom known as beefsteakfungus

Ganmodoki, a traditional Japanese tofu based dish similar to veggie burgers

Gardein

25 Gardenburger<sup>[</sup>

Glamorgan sausage

Goshen Alimentos

Green Slice vegetarian, organic and soy free hot dogs and deli slices

Impossible Foods

30 Jackfruit, a fruit whose flesh has a similar texture to pulled pork when cooked

*Koya-dofu*, freeze-dried tofu that has a taste and texture similar to meat when prepared, common in Buddhist vegetarian cuisine

Laetiporus, a mushroom which is also named chicken of the woods

Leaf protein concentrate

WO 2018/189738 PCT/IL2018/050398

LightLife

Linda McCartney Foods

Lyophyllum decastes, mushroom known as fried chicken mushroom

Meat extenders

5 Meatless

Mock duck

Morning star Farms

Muscolo di grano (Wheat's muscle), seitan prepared according to an Italian recipe

Nut roast

10 Oncom

Paneer, for example in such dishes as Paneer tikka

Quorn

Soy protein

Soy pulp, used for veggie burgers and croquettes

15 Tempeh

Textured vegetable protein

Tofu, not traditionally seen as a meat substitute in Asia, but widely used for that purpose in the

Western hemisphere

Tofurkey

20 Tofurky

Turtle Island Foods

Vegetarian bacon

Vegetarian hot dog

Veggie burger

25 Veggie Chicken patty

Veggie Chicken cutlet, breaded (schnitzel)

Veggie Chicken frankfurter (Hot dog)

Viana, one of the largest German vegan food manufacturers, offers a wide range of veggie burgers, croquettes, sausages, minced mock meats, up to vegan doner, vegan gyros and deli

30 slices for sandwiches

Wheat gluten

Any of the above examples is an independent embodiment that is not necessarily associated with a specific vendor.

It will be appreciated that food refers to food or feed (animal feeding).

As used herein "cultured animal cells" refers to any non-human cell (e.g., mammals, avian, insect, fish) that is cultured *in vitro* but is not allowed to form a tissue in vitro.

The cells can be wild-type cells or genetically modified cells (e.g., transgenic, genome edited).

The term "cultured meat" is interchangeably used with "cultured animal cells".

5

10

15

20

25

30

According to a specific embodiment, the cultured animal cells are terminally differentiated.

According to a specific embodiment, the cultured animal cells are partially differentiated (e.g., blood-derived mesenchymal precursor cells, neural progenitor cells, multipotent adult progenitor cells, mesodermal progenitor cells, spinal cord progenitor cells).

According to a specific embodiment, the cultured animal cells are stem cells (e.g., neural stem cells, multipotent stem cells from subventricular forebrain region, ependymal-derived neural stem cells, hematopoietic stem cells, liver-derived hematopoietic stem, marrow-derived stem cell, adipose-derived stem cells, islet- cells producing stem cells, pancreatic-derived pluripotent islet-producing stem cells, mesenchymal stem cells).

According to a specific embodiment, the cells are stem cells selected from the group consisting of mesenchymal stem cells and embryonic stem cells.

The phrase "embryonic stem cells" refers to embryonic cells which are capable of differentiating into cells of all three embryonic germ layers (*i.e.*, endoderm, ectoderm and mesoderm), or remaining in an undifferentiated state. The phrase "embryonic stem cells" may comprise cells which are obtained from the embryonic tissue formed after gestation (e.g., blastocyst) before implantation of the embryo (*i.e.*, a pre-implantation blastocyst), extended blastocyst cells (EBCs) which are obtained from a post-implantation/pre-gastrulation stage blastocyst (see WO2006/040763), embryonic germ (EG) cells which are obtained from the genital tissue of a fetus, and cells originating from an unfertilized ova which are stimulated by parthenogenesis (parthenotes).

Induced pluripotent stem cells (iPS; embryonic-like stem cells), are cells obtained by dedifferentiation of adult somatic cells which are endowed with pluripotency (*i.e.*, being capable of differentiating into the three embryonic germ cell layers, *i.e.*, endoderm, ectoderm and mesoderm). According to some embodiments of the invention, such cells are obtained from a differentiated tissue (e.g., a somatic tissue such as skin) and undergo de-differentiation by genetic manipulation which re-program the cell to aquire embryonic stem cells characteristics.

The phrase "adult stem cells" (also called "tissue stem cells" or a stem cell from a somatic tissue) refers to any stem cell derived from a somatic tissue [of either a postnatal or

prenatal animal (especially the human)]. The adult stem cell is generally thought to be a multipotent stem cell, capable of differentiation into multiple cell types. Adult stem cells can be derived from any adult, neonatal or fetal tissue such as adipose tissue, skin, kidney, liver, prostate, pancreas, intestine, bone marrow and placenta.

Hematopoietic stem cells, which may also referred to as adult tissue stem cells, include stem cells obtained from blood or bone marrow tissue of an individual at any age or from cord blood of a newborn individual.

Placental and cord blood stem cells may also be referred to as "young stem cells".

Mesenchymal stem cells are multipotent stromal cells that can differentiate into a variety of cell types, including osteoblasts (bone cells), chondrocytes (cartilage cells), myocytes (muscle cells) and adipocytes (fat cells which give rise to marrow adipose tissue). The term encompasses multipotent cells derived from the marrow as well as other non-marrow tissues, such as placenta, umbilical cord blood, adipose tissue, adult muscle, corneal stroma or the dental pulp of deciduous baby teeth. The cells do not have the capacity to reconstitute an entire organ.

According to a specific embodiment, the cells are selected from the group consisting of erythrocytes, adipocytes, cardiomyocytes and hepatocytes.

According to a specific embodiment, the cells are erythrocytes.

5

10

15

20

25

30

According to a specific embodiment, the cells are adipocytes.

According to a specific embodiment, the cells are hepatocytes.

According to a specific embodiment, the cells are cardiomyocytes.

According to a specific embodiment, the cells are of domesticated animals, such as listed below (e.g., duck, chicken).

According to a specific embodiment, the cells are produced by in vitro expansion e.g., stem cell expansion.

According to a specific embodiment, the cells are produced by in vitro differentiation.

Methods of cell differentiation are well known in the art.

Generally these methods are based on adherent cultures or suspension cultures, each of which is contemplated herein.

According to a specific embodiment, the cells are produced in a suspension culture.

In vitro meat can be produced by culturing the cells from farm animal species in large quantities starting from a relatively small number. Culturing satellite cells and even more so embryonic stem cells are typically for this purpose since these cells have an almost infinite self-renewal capacity. Recent reports have shown the establishment of stem-cell lines from domesticated ungulate animals e.g., non-human, rat, mouse (Challenges and prospects for the

10

15

20

25

30

15

establishment of embryonic stem cell lines of domesticated ungulates. Anim Reprod Sci. 2007;98(1-2): 147-168. doi: 10.1016/j.anireprosci.2006.10.009., which is hereby incorporated by reference). Bach et al. (Engineering of muscle tissue. Clin Plast Surg. 2003;30(4):589-599. doi: 10.1016/S0094-1298(03)00077-4.) suggested myosatellite cells as the preferred source of primary myoblasts because they recapitulate myogenesis more closely than immortal myogenic cell lines. Myosatellite cells have been isolated and characterized from the skeletal muscle tissue of cattle (Dodson et al. Optimization of bovine satellite cell derived myotube formation in vitro. Tissue Cell. 1987; 19(2): 159-166. doi: 10.1016/0040-8166(87)90001-2.), chicken (Yablonka-Reuveni et al. Dev Biol. 1987;119(1):252-259. doi: 10.1016/0012-1606(87)90226-0.), fish (Powell et al. Cultivation and differentiation of satellite cells from skeletal muscle of the rainbow trout Salmo gairdneri. J Exp Zool. 1989;250(3):333-338), lambs (Dodson et al. Isolation of satellite cells from ovine skeletal muscles. J Tissue Cult Methods. 1986;10(4):233-237. doi: 10.1007/BF01404483), pigs (Blanton Blanton et al. Isolation of two populations of myoblasts from porcine skeletal muscle. Muscle Nerve. 1999;22(1):43-50. doi: 10.1002/(SICI)1097-4598(199901)22:1, Wilschut et al. Isolation and characterization of porcine adult muscle-derived progenitor cells. J Cell Biochem. 2008; 105(5): 1228-1239.), and turkeys (McFarland et al. Proliferation of the turkey myogenic satellite cell in a serum-free medium. Comp Biochem 1991;99(1-2): 163-167. 10.1016/0300-9629(91)90252-8). Physiol. doi: Porcine progenitor cells have the potential for multilineage differentiation into adipogenic, osteogenic and chondrogenic lineages, which may play a role in the development of co-cultures (Wilschut et al. 2008, supra). Advanced technology in tissue engineering and cell biology offer some alternate cell options having practical applications and multilineage potential allowing for co-culture development with suitability for large-scale operations.

Alternatively, as mentioned, adult stem cells from farm animal species can be used. For instance, myosatellite cells are an adult stem-cell type with multilineage potential (Asakura et al. Differentiation. 2001;68(4-5):245-253. doi: 10.1046/j.1432-0436.2001.680412). also have the capacity to differentiate into skeletal muscle cells. A rare population of multipotent cells found in adipose tissue known as adipose tissue-derived adult stem cells (ADSCs) is another relevant cell type for in vitro meat production (Gimble et al. Adipose-derived stem cells regenerative medicine. Circ Res. 2007; 100(9): 1249-1260. for doi: 10.1161/01.RES.0000265074.83288.09) which can be obtained from subcutaneous fat and subsequently transdifferentiated to myogenic, osteogenic, chondrogenic or adipogenic cell lineages (Kim et al. Muscle regeneration by adipose tissue-derived adult stem cells attached to

injectable PLGA spheres. Biochem Biophys Res Commun. 2006;348(2):386-392. doi: 10.1016/j.bbrc.2006.07.063).

Matsumoto et al. (J Cell Physiol. 2007;215(l):210-222.) reported that mature adipocytes can be dedifferentiated in vitro into a multipotent preadipocyte cell line known as dedifferentiated fat (DFAT) cells, reversion of a terminally differentiated cell into a multipotent cell type. These DFAT cells are capable of being transdifferentiated into skeletal myocytes (Kazama et al. Mature adipocyte-derived dedifferentiated fat cells can transdifferentiate into skeletalmyocytes in vitro. Biochem Biophys Res Commun. 2008;377(3):780-785. doi: 10.1016/j.bbrc.2008.10.046) and are an attractive alternative to the use of stem cells.

For the sake of completeness some representative protocols for differentiation of specific cell types are provided herein.

# Chicken adipocytes differentiation from ES <u>Isolation</u> and <u>cultivation</u> of <u>chicken</u> ES cells

5

10

15

20

25

30

Isolation and cultivation of chicken ES cells can be carried out following previously described method (Wu et al., 2008 Wu XS, He XH, Dai MJ, Tian ZQ, Yang HY, Xu Q, Li BC (2008). Isolation and Cultivation of Chicken Embryonic Stem Cells and Production of Single-cell Clone. China Poult., 30: 30-34). In brief, blastoderm cells at stage X are collected by spoon method in tissue culture dishes and rinsed with Ca<sup>2+</sup> and Mg<sup>2+</sup> free PBS to remove the yolks and vitelline membrane. After washing with PBS, ES cells are transferred into fresh tissue culture dishes containing PBS plus 0.25 % trypsin and 0.04 % EDTA. Followed by digestion at 37 0°C for 5 to 8 min, the dissociated cells are collected by centrifugation at 1000 r/min for 8 to 10 min and suspended, then about 3~5xl0 <sup>4</sup> ES cells are maintained in a 5 % C02 humidified atmosphere at 37 0°C with DMEM and chicken embryonic fibroblast (CEF) cell monolayer. The medium is replaced daily with fresh medium. The cell colonies with undifferentiated morphology are rinsed three times with PBS and digested at 37 0°C for 2 to 3 min with 0.25 % trypsin plus 0.04 % EDTA. The dissociated ES clusters are suspended with a pipette and subcultured in a 5 % C02 humidified atmosphere at 37 0°C with gelatin-coated flasks containing a feeder cell layer and DMEM medium. The medium is replaced daily with half fresh medium.

#### **Induction of chicken ES cells into adipocytes**

Chicken ES cells of the third passage are sub- cultured in DMEM medium with 10 % FBS for 48 h. The cells are induced firstly in an adipocyte induction medium with 10 % FBS,  $5.5 \times 10^{-5} \text{mol/l}$   $\beta$ - ME(Sigma-Aldrich, Inc., St. Louis, MO, USA),  $1 \times 10^{-6} \text{mol/L}$  DEX (Maixin),

10 mg/ml Ins (Maixin), and  $5x10^{-4}$  mol/1 IBMX (Sigma) for 3 days and then in DMEM medium supplemented with 10 % FBS,  $5.5x10^{-5}$ mol/L  $\beta$ -ME, 10 mg/ml Ins. After cultivation for an additional day, the medium is changed by induction solutions DMEM with GlutaMAX (Gibco, 31966-021) supplemented with 10 % of Hyclone FBS (Thermo, SH30070.03), 100X Non-Essential Amino Acid solution (Sigma, M7145), 100X Penicillin/Streptomycin (PAA, PI1-010). Thereafter, the earlier step IS repeated three times. 21 days post-induction, the derived adipocytes are identified by oil red O staining as previously described (Yu F, Ge JH, Ni LG, He XH, Xu Q, Li BC (2008). *In vitro* Induction of Differentiation of Chicken Spermatogonial Stem Cells into Adipocytes. J. Agric. Biotechnol. 16: 450-455).

Protocols for differentiation of chicken ES cells can be found in Li B, Zhang Y, Chen X, Shi Q, Fu D, Yin Y, et al. Directional differentiation of chicken embryonic stem cells into osteoblasts, neuron-like cells and adipocytes. African J. Biotechnol. 2011; 10: 7772-7779.

## Erythroblasts differentiation from ES (via embryoid bodies)

Embryonic body (EB) (Li B, Zhang Y, Chen X, Shi Q, Fu D, Yin Y, et al. Directional differentiation of chicken embryonic stem cells into osteoblasts, neuron-like cells and adipocytes. African J. Biotechnol. 2011; 10: 7772-7779) cell expansion:

10<sup>6</sup> cESC cells are thawed in gelatinized plates in cESC proliferation medium containing DMEM with 2 mM L-glutamine, 100 U/mL penicillin, and 100 g/mL streptomycin [all from Gibco/BRL, Carlsbad, CA], 15 % fetal calf serum [FCS; Euroclone, Milano, Italy], 1 % leukemia inhibitory factor [LIFJ—supernatant. 1.5X10<sup>-4</sup> M monothioglycerol [MTG; Sigma, St Louis, MO]) on gelatinized plates. The basic medium is changed from DMEM to Iscove modified Dulbecco medium (IMDM; Gibco/BRL) 2 days prior to in vitro differentiation. Cells are grown without feeder cells. The medium is changed daily using mESC medium. The basic medium was is from DMEM to IMDM 2 days prior to in vitro differentiation.

### Embryonic **Body** (**EB**) Formation

5

10

15

20

25

30

20 LiL drops of the IX10 <sup>4c</sup> cESC cells/mL suspension are placed onto the inner surface of 150 mm plate. They are incubated for 5 days and EBs are collected with 10 mL PBS into a 50 mL conical sterile tube. The EBs are allowed to sediment for 10-20 min at room temperature and PBS is removed. 0.05% trypsin is added to the cell culture and incubated for 2-7 min and Inactivated the trypsin by 2 mL differentiation medium. The EBs obtained from one plate are replated in 150 Place back in the incubator to 1-4 days.

10

15

20

25

30

## **Generation of ES cell-derived erythroid cultures (ES-EPs)**

EB-derived cells at a concentration of 2X106/mL are seeded into serum-free medium (StemPro34 plus nutrient supplement; Gibco/BRL) plus human recombinant erythropoietin (Epo; 2 U/mL; Erypo; Janssen-Cilag AG, Baar, Switzerland), murine recombinant Kit-ligand (KL or stem cell factor [SCF]; 100 ng/mL; R&D Systems, Minneapolis, MN), 106 M dexamethasone (Sigma), and 40 ng/mL insulin-like growth factor 1 (IGF-1; Promega, Madison, WI). At day 1 and day 3 of cultivation, the suspension of cells is collected and replated into a new dish.

Cell density is maintained between 2 and 4X10<sup>6</sup> cells/mL. ES-EPs are washed twice with PBS and cultivated at 2X10<sup>6</sup> cells/mL in serum-free medium (StemPro34 plus nutrient supplement) supplemented with 10 U/mL Epo, insulin (10 ng/mL; Actrapid HM; Novo Nordisk), 3X10<sup>6</sup> M of the glucocorticoid receptor antagonist ZKl 12.993,3 and one mg/mL iron-saturated human transferrin. ES-EPs derived from 6-day-old embryoid bodies are cultivated for 16 days under erythroid conditions, washed twice in PBS, and resuspended in IMDM.

Protocols for erythrocyte differentiation can be found in Carotta, S.; Pilat, S.; Mairhofer, A.; Schmidt, U.; Dolznig, H.; Steinlein, P.; Beug, H. Directed differentiation and mass cultivation of pure erythroid progenitors from mouse embryonic stem cells. *Blood* **2004**, *104*, 1873-1880.

Protocols for hepatocyte differentiation can be found in

Stock et al., *The generation of hepatocytes from mesenchymal stem cells and engraftment into murine liver.* Nature protocols, VOL.5 NO.4, 2010; Protocols for cardiomyocyte differentiation can be found in Breckwoldt et al., Differentiation of cardiomyocytes and generation of human engineered heart tissue Nature protocols, VOL.12 NO.6, 2017.

According to a specific embodiment, the cells are differentiated from mesenchymal stem cells, hematopoietic stem cells or embryonic stem cells.

As used herein "tissue" refers to a structure of more than one cell type that is aggregated (e.g., connected) by natural or synthetic means (e.g., matrix).

According to a specific embodiment, the foodstuff is devoid of animal contaminants such as animal pathogens (that occur in vivo), hazardous concentrations of hormone(s) (e.g., ketosteroids, GH) and antibiotics that typically cause rapid weight gain).

According to a specific embodiment, the cells are grown in a free pathogen, virus, bacteria and nitrate and are distributed in such conditions as well.

Ultroser G is an example of a commercially available serum substitute containing growth factors, binding proteins, adhesin factors, vitamins, hormones, mineral trace elements and has

been designed specially to replace fetal bovine serum for growth of anchorage-dependent cells in vitro (Duque et al. Use of two replacements of serum during bovine embryo culture in vitro. Theriogenology. 2003;59(3-4):889-899. doi: 10.1016/S0093-691X(02)01 134-2). Benjaminson et al. (vitro edible muscle protein production system (MPPS): Stage 1, fish. Acta Astronaut. 2002;51(12):879-889. doi: 10.1016/S0094-5765(02)00033-4) succeeded in using a serum-free medium made from maitake mushroom extract that achieved higher rates of growth than fetal bovine serum and recently it has been shown that lipids such as sphingosine-1-phosphate can replace serum in supporting the growth and differentiation of embryonic tissue explants.

5

10

15

20

25

30

According to a specific embodiment, the foodstuff is free of growth hormone (GH) in an amount hazardous for human or animal consumption (undetectable amounts by HPLC).

According to a specific embodiment, the plant-originated substance augments (by way of reaction) the organoleptic properties of the non-human cultured meat cells within the hybrid foodstuff.

The plant-originated substance is selected such that it supports the development of flavor components when reacting with the cultured biomass during the preparation of the hybrid food, such as cooking, baking, frying, grilling, roasting etc. Thus the hybrid foodstuff mimics the equivalent development of flavors in conventional meat preparations.

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 50 % (w/w).

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 45 % (w/w).

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 40 % (w/w).

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 35 % (w/w).

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 30 % (w/w).

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 50 % (w/w).

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 25 % (w/w).

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 20 % (w/w).

WO 2018/189738 PCT/IL2018/050398

20

According to a specific embodiment, animal-derived protein in the hybrid foodstuff is below 15 % (w/w).

According to a specific embodiment, the animal protein in the hybrid foodstuff is 10-49~% (w/w).

According to a specific embodiment, the foodstuff is not a chip and and/or the plant-originated substance is not a plant derived hydrogel.

5

10

15

20

25

30

As mentioned, the foodstuff preparation is performed by combining a plant-originated substance with an amount of cultured meat cells.

The terms "connection between" and "interconnected with" as defined above refers to those specific cases where at least two different portions are provided *inter alia*, by any suitable means, including (i) where the at least one second portion is comprised, immersed, wetted, solubilized, suspended, doped, glued, attached, aggregated, mixed, contained or otherwise, directly or indirectly, provided in contact within and/or upon or on top of the aforesaid at least one first portion; and (ii), where the at least one second portion is chemically, biologically and/or physically reacts with the aforesaid at least one first portion so that either integral or non-integral non-cultured meat and cultured meat substantially unified phase, matter or composition is obtained.

The combined matter can be subjected to further processing such as by means of rising, kneading, extruding, molding, shaping, cooking, stewing, boiling, broiling, baking, frying and any combination of same.

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 18% to about 1% (w/w) cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 82% plant-originated substances and additives thereof and 18% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 83% plant-originated substances and additives thereof and 17% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 84% plant-originated substances and additives thereof and 16% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 85% plant-originated substances and additives thereof and 15% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 86% plant-originated substances and additives thereof and 14% (w/w) or lower cultured meat (cultured animal cells).

5

10

15

20

25

30

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 87% plant-originated substances and additives thereof and 13% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 88% plant-originated substances and additives thereof and 12% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 89% plant-originated substances and additives thereof and 11% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 90% plant-originated substances and additives thereof and 10% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 91% plant-originated substances and additives thereof and 9% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 92% plant-originated substances and additives thereof and 8% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 93% plant-originated substances and additives thereof and 7% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 94% plant-originated substances and additives thereof and 6% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 95% plant-originated substances and additives thereof and 5% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 96% plant-originated substances and additives thereof and 4% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 97% plant-originated substances and additives thereof and 3% (w/w) or lower cultured meat (cultured animal cells).

5

10

15

20

25

30

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 98% plant-originated substances and additives thereof and 2% (w/w) or lower cultured meat (cultured animal cells).

The present invention discloses a novel hybrid foodstuff technology where the foodstuff comprises, *inter alia*, at least about 99% plant-originated substances and additives thereof and 1% (w/w) or lower cultured meat (cultured animal cells).

According to an aspect of the invention there is provided a hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 30 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 25 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 20 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 15 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 10 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or

meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 5 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 3 % (w/w) of the hybrid foodstuff.

5

10

15

20

25

30

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 3 - 30 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 3 - 25 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 3 - 20 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 3 - 15% (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 3 - 10 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 5 - 15% (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 10 - 15% (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 7.5 - 15 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 3 - 5 % (w/w) of the hybrid foodstuff.

5

10

15

20

25

30

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 7.5 - 10 % (w/w) of the hybrid foodstuff.

According to an embodiment, the hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is 3 - 12 % (w/w) of the hybrid foodstuff.

The present invention also discloses a novel hybrid foodstuff technology wherein a portion of the cultured meat in the end product is defined as the minimal weight percent providing this end product with an enhancement in a meat organoleptic property or meat nutritional property than the corresponding foodstuff being non-hybrid, cultured meat free vegetable-based product.

An aspect of the invention also relates to a method of qualifying a hybrid foodstuff, the method comprising subjecting the hybrid foodstuff to at least one organoleptic test.

According to a specific embodiment, the hybrid foodstuff is any of the hybrid foodstuffs contemplated herein.

Methods of organoleptic assaying are well known in the art, some of which are described infra.

Organoleptic (sensory) evaluation is a common and very useful tool in quality assessment of processed food (e.g., meat) products, but to date not for hybrid foodstuffs. It makes use of the senses to evaluate the general acceptability and quality attributes of the products. The assays typically make use of dedicated panelists and/or artificial means.

All panelists must use proven and identical test methods in order to make their results comparable. Each panelist involved in such tests is given a scoresheet, where they mark their

findings. Scoresheets of the team of panelists are evaluated and a test result for each individual product is produced based on multiple observations.

Common test methods used in sensory evaluation are:

- 1. **Paired comparison test for simple difference** where two coded samples are presented to the panelists for evaluation on simple difference (Table B).
- 2. **Triangle test** where three coded samples are presented at the same time, two are identical and the third is odd and the panelist is asked to identify the odd sample (Table C).
- 3. **Hedonic scale rating test or acceptability test** where samples are tested to determine their acceptability or preference (Table D).

## 10 Table B: Score sheet for Paired-Comparison Test for Simple Difference

Name:		Product:	
Panelist No.:		Date;	
Instructions: You are given a pair of obstween them by placing an write down any comments.		ate if there is a difference opropriate column. Please	
SAMPLE PAIR CODE	THERE IS A DIFFERENCE	THERE IS NO DIFFERENCE	
Comments:	<u>.</u>		

## **Table C: Score sheet for Triangle Test**

Name :	Product:
Panelist No .:	Date :
Instructions: YOU are presented with three: coded samples these samples are identical while the third is o taste each sample, Indicate the code of the o mark across the code, Please write down any com	dd or different. Smell and dd sample by placing an x
SAMPLE CODE	ODD SAMPLE
Comments:	

Table D: Score sheet for hedonic-scale rating test

Name:	Product:	Product:	
Panelist No.:	Date;		
Instructions: Taste the given samples, then publich best describes your feeling.	lace an x mark on the point in the sca	ale	
	SAMPLE CODE		
SCORE*			
(9) Like extremely			
(8) Like very much			
(7) Like moderately			
(6) Like slightly	, i		
(5) Neither like nor dislike			
(4) Dislike slightly			
(3) Dislike moderately			
(2) Dislike very much			
(1) Dislike extremely		- 1	

<sup>\*</sup> Note: Numbers in parentheses are to be assigned during data analysis and are not to appear in the score sheet.

Tests for simple difference and the triangle test are very useful methods for quality control and product development. Newly formulated products can be evaluated by determining if a simple difference exists between the new products developed and the old ones. Similarly, the hedonic scale rating can be used for internal factory testing, and this method is also suitable for market research by determining the consumer's acceptance or preference for certain products. Such a method i.e., the 9-PHS method is further described hereinbelow.

#### Instruments for texture measurement

5

10

15

20

25

Sensory testing (chewing) is normally sufficient to test tenderness/toughness or homogenous/fibrous structure of meat and meat products. If more objective results are desired, special instruments for texture measurement can be employed. Such a device typically measures the shear-force necessary to cut through meat/meat products. Comparative texture measurements are usually taken from same tissues or products which were submitted to different treatments such as ripening, cooking etc.

According to a specific embodiment, organoleptic assaying is effected by a combination of expert panel studies and by consumer difference and preference tests.

The expert panel studies may include two types of tests:

Difference testing - proving statistically the sensory benefits of the hybrid foodstuff as compared to the corresponding product without the cultured animal cells (substituted or not as defined); Quantitative Descriptive Analysis (QDA) - demonstrating how the the hybrid foodstuff differs in specific sensory attributes from the corresponding product without the cultured animal cells (substituted or not as defined), with emphasis on the magnitude of meat related texture and flavor characteristics.

Consumer studies will demonstrated how well the products are liked, which are preferred, and will quantify the effect of the SuperMeat on preferred sensory attributes.

Sensory attributes

5

10

15

20

25

30

The list of relevant sensory attributes includes three main groups, adjusted individually per type of product, as follows:

Appearance: surface color, internal color, texture (coarsness, uniformity), overall rating with relevance to the type of product tested.

Texture: hardness/softness, juiciness/dryness, cohesiveness, chewiness, fatty/oily mouthfeel, overall rating.

Taste and flavor (possible list of positive and negative characteristics of aroma and taste): meaty, cooked chicken, roasted chicken, bouillon-like (brothy), greasy, burned, sweet, bitter, rancid, overall rating.

The detailed sensory list applies to the expert panel studies, mainly the QDA comparisons.

Selected and easily understood attributes in each sensory category (apperance, texture and taste) are used for the consumer acceptance and preference studies.

The present invention also discloses a novel hybrid foodstuff technology wherein a portion of the cultured meat in the end product is defined as the minimal weight percent providing this end product with one point more in a 9-PHS (as will defined below) than the corresponding foodstuff being non-hybrid, cultured meat free vegetable-based product.

The present invention further discloses a novel hybrid foodstuff technology wherein this foodstuff comprises, *inter alia*, about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 3% to 30% (w/w), where the improvement is that said oil portion consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat.

The present invention further discloses a novel hybrid foodstuff technology wherein this foodstuff comprises, *inter alia*, about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (e.g., gum, binding, thickening, aroma flavor additives) and/or water immiscible materials (oil portion), ranging from about 3% to 30% (w/w), where the improvement is that said additives thereof (e.g., gum, binding, thickening, aroma flavor additives) and/or water immiscible materials (oil portion) consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat.

Where the improvement is that it allows for the replacement of meat taste and aroma flavor additives which are mostly synthetic or based on different types of hydrolysates (of plant

WO 2018/189738 PCT/IL2018/050398

or yeast proteins) by incorporating a composition of cultured meat cells such as erythrocytes, hepatocytes and/or adipocytes, allowing for the development of hybrid foodstuff with minimal additives and flavorings ingredients providing authentic meat taste.

28

Where the improvement is that it allows for the replacement gums and binding or thickening additives in meat substitutes by incorporating a composition of cultured meat ceils such as erythrocytes, hepatocytes and/or adipocytes, allowing for the development of hybrid foodstuff with minimal additives ingredients providing authentic meat texture.

5

10

15

20

25

30

Thus, the present teachings contemplate for substitution of various components which are non meat in the foodstuff so as to render them more nutritional and/or with improved organoleptic properties.

The present invention additionally discloses a novel hybrid foodstuff technology wherein this foodstuff comprises, *inter alia*, about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 3% to 30% (w/w), where the improvement is that said oil portion consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat; wherein the smallest portion of the cultured meat in the end product is defined as the minimal weight percent providing said end product with at least one 9-PHS point more than a corresponding foodstuff being non-hybrid, cultured meat free vegetable-based product.

The control is defined as the same plant-originated foodstuff that is devoid of the supplementation or addition of the cultured meat cells.

The present invention further discloses a novel hybrid foodstuff technology wherein this foodstuff comprises, *inter alia*, about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (e.g., gum, binding, thickening, aroma flavor additives) and/or water immiscible materials (oil portion), ranging from about 3% to 30% (w/w), where the improvement is that said additives thereof (e.g., gum, binding, thickening, aroma flavor additives) and/or water immiscible materials (oil portion)consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat, wherein the smallest portion of the cultured meat in the end product is defined as the minimal weight percent providing said end product with at least one 9-PHS point more than a corresponding foodstuff being non-hybrid, cultured meat free vegetable-based product.

It is hence on the scope of the invention wherein a cultured meat containing hybrid foodstuff as defined above comprises:  $X_T$ % by weight of non-cultured meat foodstuff portion, e.g., plant-originated substances, such as roots, leaves, stems, flowers and fruits of plants, where solids of soy protein hydrolysate is an example; Xi% by weight of an  $i^{th}$  at least one ingredient of

said non-cultured meat foodstuff portion, canola oil is an example; and Y % by weight of cultured meat cells.

29

Moreover, the present invention hence discloses a novel hybrid foodstuff technology wherein this foodstuff is characterized by about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 3% to 30% (w/w); said oil portion consists of up to about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat; wherein the role said mixtures in the oil portion (measured as vegetable oil to culture meat (w/w) in improving the organoleptic properties (measured as 9-PHS) of said hybrid foodstuff of is provided by any of **Formula I, II** or **III** as defined below.

# THE ROLE OF VEGETABLE FAT SUBSTITUTION BY CULTURED HEPATOCYTES IN HYBRID FOODSTUFFS

According to an embodiment of the invention,  $X_T$ , Xi & Y values, here % are defined are weight to total dry weight ratio, are provided in **Table 1** below. It is acknowledged that in this patent the term 'about' refers to any value being up to 20% (e.g., 10 %) bigger or 20% smaller (e.g., 10 %) than the defined measure.

**Table 1.** Possible X<sub>T</sub>, Xi & Y values

5

10

15

	X <sub>T</sub> , %	<b>Xi,</b> % (wt/dry	<b>Y</b> , % (wt/dry
	(wt/dry wt)	wt)	wt)
1st hybrid	from about	from about	from about
foodstuff	51% to about 75%	1% to about 5%	49% to about 25%
2 <sup>nd</sup> hybrid	from about	from about	from about
foodstuff	51% to about 75%	6% to about 10%	49% to about 25%
3 <sup>rd</sup> hybrid	from about	from about	from about
foodstuff	51% to about 75%	11% to about 15%	49% to about 25%
4th hybrid	from about	from about	from about
foodstuff	51% to about 75%	16% to about 20%	49% to about 25%
5 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	51% to about 75%	21% to about 25%	49% to about 25%
6 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	51% to about 75%	26% to about 30%	49% to about 25%
7 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	76% to about 90%	1% to about 5%	24% to about 10%

30

		30	
8 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	76% to about 90%	6% to about 10%	24% to about 10%
9 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	76% to about 90%	11% to about 15%	24% to about 10%
10 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	76% to about 90%	16% to about 20%	24% to about 10%
1 1 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	76% to about 90%	21% to about 25%	24% to about 10%
12 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	76% to about 90%	26% to about 30%	24% to about 10%
13 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	91% to about 95%	1% to about 7%	9% to about 5%
14 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	91% to about 95%	8% to about 14%	9% to about 5%
15 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	91% to about 95%	15% to about 21%	9% to about 5%
16st hybrid	from about	from about	from about
foodstuff	91% to about 95%	22% to about 29%	9% to about 5%
17 <sup>nd</sup> hybrid	from about	from about	from about
foodstuff	91% to about 95%	30% to about 37%	9% to about 5%
18 <sup>th</sup> hybrid	from about	from about	from about
foodstuff	91% to about 95%	38% to about 45%	9% to about 5%

In one embodiment of the invention, hepatocytes, *i.e.*, cells of the main parenchymal tissue of the liver mass, are harvested and provided as a substituted canola oil various fractions in both sausage-like, ground beef-like and schnitzel-like novel hybrid foods.

The term **'hepatocytes cells'** used hereinafter concurrently refers to (a) hepatocytes cells; (b) genetically engineered liver tissue ingredients;

5

10

A hedonic rating scale is utilizable to measure food acceptance, see Lim, Juyun. "Hedonic scaling: A review of methods and theory." *Food quality and preference* 22.8 (2011): 733-747, which is incorporated herein as a reference. A 9-point hedonic scale, utilizable herein, is a balanced bipolar scale around neutral at the center with four positive and four negative categories on each side. The categories are labeled with phrases representing various degrees of

affect and those labels are arranged successively to suggest a single continuum of likes and dislikes, as follows:

9-Point Hedonic Scale			
9	Like Extremely		
8	Like Very Much		
7	Like Moderately		
6	Like Slightly		
5	Neither Like nor Dislike		
4	Dislike Slightly		
3	Dislike Moderately		
2	Dislike Very Much		
1	Dislike Extremely		

Table 2 below describes potential results for assessment of chicken sausage-like, ground beef-like and calf schnitzel-like cultured meet-containing hybrid foods, as can be measured by 9-point hedonic scale (9-PHS).

Can be tested on individuals of the following characterization N=25, age 30-65 yrs, about 50% male,

10 **Table 2** 

		X <sub>T</sub> , %	Xi, %	Υ, %	9
		(wt/dry wt)	(wt/dry wt)	(wt/dry wt)	-PHS
		Soy protein hydrolysate- based solids	Canola oil	Cultured hepatocytes	
	sausage-like	100%	30%	0%	5
dish	sausuge mie	10070	2070	0,70	
	sausage-like	95%	25%	5%	6
dish					
	sausage-like	87%	17%	13%	7
dish	sausage-like	80%	10%	20%	7
dish	-				
	sausage-like	73%	3%	27%	7
dish					

		32		
ground beef-	100%	30%	0%	4
like dish				
ground beef-	95%	25%	5%	5
like dish				
ground beef-	87%	17%	13%	6
like dish				
ground beef-	80%	10%	20%	7
like dish				
ground beef-	73%	3%	27%	8
like dish				
schnitzel-like	100%	30%	0%	4
dish				
schnitzel-like	95%	25%	5%	6
dish				
schnitzel-like	87%	17%	13%	8
dish				
schnitzel-like	80%	10%	20%	8
dish				
schnitzel-like	73%	3%	27%	6
dish				

The function of such a replacement is provided useful  $[R^2=0.92]$  by formula I:

**Formula I** 9-points hedonic score =  $a+bX^c$ 

Where: a is ranging from about -2 to about 2; e.g.,  $2*10^{-3}$ ;

b is ranging from about -3 to about 3; e.g., 1;

5

10

c is ranging from about -1 to about 5; e.g., 3.4.

The function of such a replacement is provided useful [R<sup>2</sup>=0.98] by formula II:

**Formula II** 9-points hedonic score =  $a+bX+cX^2$ 

Where: a is ranging from about 0 to about 4; e.g., 3.4

b is ranging from about 1 to about 3; e.g., 1.8; and

c is ranging from about -1 to about 2.5; e.g., 0.2

The function of such a replacement is provided useful [R<sup>2</sup>=0.98] by formula III:

**Formula III** 9-points hedonic score =  $a+bX+cX^2$ 

Where: a is ranging from about 0 to about 4; e.g., -0.4

b is ranging from about 2 to about 6; e.g., 4.8; and c is ranging from about -1 to about 2.5; e.g., -0.7.

5

10

15

20

25

30

#### THE ROLE OF CULTURED HEME-POTEINS IN HYBRID FOODSTUFFS

The two main heme proteins (HPs), hemoglobin and myoglobin, are important factors determining meat quality aspects such as color and hemorrhage, see Kranen, R. W., et al. "Hemoglobin and myoglobin content in muscles of broiler chickens." *Poultry science* 78.3 (1999): 467-476; and see Boles, Jane Ann, and Ronald Pegg. "Meat color." Montana State University and Saskatchewan *Food Product Innovation, Program* University of Saskatchewan (2010), which are both incorporated herein as a reference.

It is well in the scope of the invention where HPs are cultured in various methods, various sources, including, e.g., liver, bone marrow, mesenchymal cells or any other multipotent stromal cells that can differentiate into a variety of cell types, including osteoblasts (bone cells), chondrocytes (cartilage cells), myocytes (muscle cells) and adipocytes (fat cells); yet for convenient and in a non-limiting manner, one set of examples will be hereto disclosed.

Hepatocytes are enriched by those two main HPs and thus provide meat-like foods with improved organoleptic properties including meat-like color, texture, mouth feel, odor, flavor and appearance. Heme derived from the hepatocytes increasing bioavailability of the iron at about 24 % higher than that in food containing iron sulfate, see G. Gonzalez-Rosendo, *et al.*, "Bioavailability of a heme-iron concentrate product added to chocolate biscuit filling in adolescent girls living in a rural area of Mexico," *Journal of Food Science*, vol. 75, pp. H73-H78, 2010 which is incorporated herein as a reference.

In another embodiment of the present invention, six different novel hybrid foodstuffs are disclosed: (a) A chicken-flavored cultured-meat containing foodstuff, namely sausage-like hybrid food (I) and ground meat-like hybrid food (II), where the differentiated hepatocytes cells are cultivated to provide about 0.03% to about 0.07% HPs in the foodstuff; (a) A chicken-flavored cultured-meat containing foodstuff, where the differentiated hepatocytes cells are cultivated to provide about 0.1% to about 0.7% HPs; (b), A beef-flavored cultured-meat containing foodstuff, where the differentiated hepatocytes cells are cultivated to provide about 0.75 to about 2.5% HPs; (c) A chicken-flavored cultured-meat containing foodstuff, where the differentiated hepatocytes cells are cultivated to provide about 0.1% to about 0.3% oxymyoglobin; (d) A chicken-flavored cultured-meat containing foodstuff, where the differentiated hepatocytes cells are cultivated to provide about 0.1% to about 0.3% myoglobin; (e) A chicken-flavored cultured-meat containing foodstuff, where the differentiated hepatocytes cells are cultivated to provide about 0.1% to about 0.3% myoglobin; (e) A chicken-flavored cultured-meat containing foodstuff, where the differentiated hepatocytes

PCT/IL2018/050398 WO 2018/189738

cells are cultivated to provide about 0.1% to about 0.3% metmyoglobin; and (f) a combination (2.5% in total) of the same.

As mentioned, cultured cells can be used in their native form in the hybrid foodstuff or treated or genetically engineered to enrich a certain trait/phenotype e.g., level of expression of HPs.

Methods of genetically modifying cells are well known in the art.

Table 3 below describes potential results for assessment of sausage-like hybrid food (I) and ground meat-like hybrid food (II) cultured meet-containing hybrid foods as can be measured by 9-point hedonic scale (9-PHS).

The assay can be executed on individuals of the following characterization N= 25, age 30-65 yrs, about 50% male,

Table 3

5

10

	Table 5		T		
		Heme-	Heme-	Heme-	9
		containing	containing	containing	-PHS
		myoglobin,	oxymyoglobin,	metmyoglobin,	
		% (wt/dry	% (wt/dry	% (wt/dry	
		wt)	wt)	wt)	
	sausage-like	0.01			
dish					4
	sausage-like	0.20			
1. 1	sudsuge like	0.20			4
dish					4
	sausage-like	0.50			
dish					5
	sausage-like		0.01		
dish	C				6
uisii					U
	sausage-like		0.20		
dish					6
	sausage-like		0.50		
dish					7
GISH	1'1			0.01	,
	sausage-like			0.01	
dish					5
	sausage-like			0.20	
dish					5
	sausaga lika			0.50	
	sausage-like			0.50	6

WO 2018/189738 PCT/IL2018/050398

		35		
dish				
sausage-like	0.01	0.	0.04	
dish				6
sausage-like	0.01	1.5	0.14	
dish				6
sausage-like	0.05	1.9	0.04	
dish				4
sausage-like	0.15	0.6	0.04	
dish				4
sausage-like	0.20	0.1	0.04	
dish				5
ground meat -	0.01			
like dish				6
ground meat -	0.20			_
like dish	0.50			6
ground meat -	0.50			7
like dish		0.01		7
ground meat -		0.01		_
like dish		0.20		5
ground meat -		0.20		_
like dish ground meat -		0.50		5
like dish		0.30		6
ground meat -			0.01	
like dish			0.01	6
ground meat -			0.20	3
like dish			0.20	6
ground meat -			0.50	
like dish			<b></b>	4

**Table 3** above hence demonstrates the potential of *in vitro* produced HPs in cultured meat containing hybrid food and the synergic role of mixtures thereof in some of the foods, where improving food organoleptic features is to be shown.

The hybrid food is characterized by relatively high nutritional value. It provides with useful in administrating an effective dose of nutritional supplement, vitamins, minerals and proteins, e.g., those which *inter alia* selected from the group consisting of vitamins E, thiamine (B-l), riboflavin (B-2), niacin (B-3), pantothenic acid (B-5), B-6, biotin (B-7), folic Acid B-9, B-12, A, C, D, K, magnesium, iron, zinc, omega-3 fatty acids, creatine, carnosine and mixtures of the same.

5

10

15

20

25

30

The hybrid cultured meat and products thereof are enriched to some degree, when required, with additives to protect or modify its flavor or color, to improve its tenderness, juiciness or cohesiveness, or to aid with its preservation. Cultured meat additives hence potentially include, inter alia, salt and other means to impart flavor and inhibits microbial growth, extends the product's shelf life and helps emulsifying finely processed products, such as sausages. Nitrite is utilizable in curing meat to stabilize the meat's color and flavor, and inhibits the growth of spore-forming microorganisms such as C. botulinum. Phosphates used in meat processing are normally alkaline polyphosphates such as sodium tripolyphosphate. Erythorbate or its equivalent ascorbic acid (vitamin C) is utilizable to stabilize the color of cured meat. Sweeteners such as sugar or corn syrup impart a sweet flavor, bind water and assist surface browning during cooking in the Maillard reaction. Seasonings impart or modify flavor. They include spices or oleoresins extracted from them, herbs, vegetables and essential oils. Flavorings such as monosodium glutamate impart or strengthen a particular flavor. Tenderizers break down collagens to make the meat more palatable for consumption. They include proteolytic enzymes, acids, salt and phosphate. Dedicated antimicrobials include lactic, citric and acetic acid, sodium diacetate, acidified sodium chloride or calcium sulfate, cetylpyridinium chloride, activated lactoferrin, sodium or potassium lactate, or bacteriocins such as nisin. Antioxidants include a wide range of chemicals that limit lipid oxidation, which creates an undesirable "off flavor", in precooked meat products. Acidifiers, most often lactic or citric acid, can impart a tangy or tart flavor note, extend shelf-life, tenderize fresh meat or help with protein denaturation and moisture release in dried meat. They substitute for the process of natural fermentation that acidifies some meat products such as hard salami or prosciutto.

It is thus within the scope of the invention wherein the hybrid food additionally comprising Acidity regulators, Acidity regulators, Anticaking agents, Anticaking agents, Antifoaming agents, Antifoaming agents, natural and other Antioxidants, Bulking agents, Food coloring agents, color retention agents, Emulsifiers, Flavors, Flavor enhancers, Flour treatment agents, Glazing agents, Humectants, Tracer gas, Preservatives, Probiotic microorganisms, Stabilizers, Sweeteners, Thickeners and any mixtures thereof.

WO 2018/189738 PCT/IL2018/050398

37

In other embodiment of the present invention, the hybrid food product is a consumable, edible item having the final organoleptic properties of a meat product, and especially product(s) selected from the group consisting of Beef, Beef heart, Beef liver, Beef tongue, Bone soup from allowable meats, Buffalo, Bison, Calf liver, Caribou, Goat, Ham, Horse, Kangaroo, Lamb, Marrow soup, Moose, Mutton, Opossum, Organ Meats, Pork, Bacon, Rabbit, Snake, Squirrel, Sweetbreads, Tripe, Turtle, Veal, Venison, Chicken, Chicken Liver, Cornish Game Hen, Duck, Duck Liver, Emu, Gizzards, Goose, Goose Liver, Grouse, Guinea Hen, Liver, Ostrich, Partridge, Pheasant, Quail, Squab, and Turkey.

In other embodiment of the present invention, the hybrid food comprises *ex vivo* produced cells (cultured animal cells) selected in a non-limiting manner from a group consisting of myoblasts, osteoblasts, fibroblasts, lipoblasts, odontoblasts, adult neuronal progenitor cells, neural stem cells, multipotent stem cells from subventricular forebrain region, ependymal-derived neural stem cells, hematopoietic stem cells, liver-derived hematopoietic stem, marrow-derived stem cell, adipo-fibroblasts, adipose-derived stem cells, islet- cells producing stem cells, pancreatic-derived pluripotent islet-producing stem cells, mesenchymal stem cells, bone marrow stromal cells, muscle side population cells, bone marrow-derived recycling cells, blood-derived mesenchymal precursor cells, bone-marrow derived side population cells, muscle precursor cells, circulating skeleton stem cells, neural progenitor cells, multipotent adult progenitor cells, mesodermal progenitor cells, spinal cord progenitor cells and spore-like cell, mixtures or any combinations thereof.

#### **Surfactant filtration**

5

10

15

20

25

30

In one embodiment of the present invention, the media of the cells is recovered and filtrated in order to extract compounds secreted from the cells. Said compounds are then used in the mix of the foodstuff.

## **Use of scaffolds**

In general, the cell lines are grown preferably without the use of scaffolds. It is also acknowledged that for those specific cell lines that scaffold is required for differentiation purposes, said scaffold will be used. Also, said scaffold will be an edible scaffold. Alternatively, the cells can be removed from the scaffold prior to inclusion in the hybrid food.

10

15

20

25

30

The present invention further provides a **method** of producing a hybrid food product, comprising steps of: a. providing plant-originated substance; b. providing cultured meat, cells and substances thereof; c. interconnecting said plant-originated substance with said cultured meat, cells and substances thereof; d. forming said hybrid food product into a desired form.

More specific, the present invention further provides a **method** of producing a hybrid food product, comprising steps of: a. providing plant-originated substance in a quantity following Table 1; b. providing cultured meat, cells and substances thereof in a quantity following Table 1; c. interconnecting said plant-originated substance with said cultured meat, cells and substances thereof; d. forming said hybrid food product into a desired form.

It is included in any of the methods, known in the art steps of cooking, sterilizing, pasteurizing, packaging and storing the hybrid food product.

Also provided is a method of providing nutrition to a subject in need thereof. The method comprising providing the subject with a hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 50 % (w/w) of the hybrid foodstuff (e.g., as described hereinabove).

According to a specific embodiment, the subject is at risk of nutritional deficiency.

According to a specific embodiment, the subject is a healthy subject (e.g., not suffering from a disease associated with nutrition/absorption).

According to a specific embodiment, the subject suffers from malnutrition.

According to a specific embodiment, the subject suffers from a disease associated with nutrition/absorption e.g., liypocoba!arninemia. iron deficiency anemia, zinc deficiency and vitamin D deficiency, fatty acid deficiency.

The terms "comprises", "comprising", "includes", "including", "having" and their conjugates mean "including but not limited to".

The term "consisting of means "including and limited to".

The term "consisting essentially of" means that the composition, method or structure may include additional ingredients, steps and/or parts, but only if the additional ingredients, steps and/or parts do not materially alter the basic and novel characteristics of the claimed composition, method or structure.

As used herein, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a compound" or "at least one compound" may include a plurality of compounds, including mixtures thereof.

Throughout this application, various embodiments of this invention may be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

5

10

15

20

25

30

Whenever a numerical range is indicated herein, it is meant to include any cited numeral (fractional or integral) within the indicated range. The phrases "ranging/ranges between" a first indicate number and a second indicate number and "ranging/ranges from" a first indicate number "to" a second indicate number are used herein interchangeably and are meant to include the first and second indicated numbers and all the fractional and integral numerals therebetween.

As used herein the term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known manners, means, techniques and procedures by practitioners of the chemical, pharmacological, biological, biochemical and medical arts.

As used herein, the term "treating" includes abrogating, substantially inhibiting, slowing or reversing the progression of a condition, substantially ameliorating clinical or aesthetical symptoms of a condition or substantially preventing the appearance of clinical or aesthetical symptoms of a condition.

# MEAT-FLAVORED HYBRID FOODSTUFF FREE OF HARMFUL SUBSTANCES AT HARMFUL QUANTITIES

[1] Meat is generally regarded by nutritionists as a healthy component of a well-balanced diet if eaten in moderation, although it's clearly not essential, as vegetarians have proven for eons; see Johnathon Andrew "Benefits & Disadvantages of a Meat Diet" in the currently available link //livehealthy(dot)chron(dot)com/benefits-disadvantages-meat-diet-1908(dot)html(dot). Meat is an excellent source of protein and contains many of the vitamins and minerals that people need. On the other hand, meat is relatively high in fat and cholesterol, which are linked to increased risk of cardiovascular disease.

Meat Diet: Meat is typically defined as the flesh, muscle tissue and organs from animals, particularly mammals such as cows, pigs and lamb. Birds such as chickens and turkeys are often classified generically as meat, but sometimes differentiated as poultry. Fish has its own category and is not usually referred to as meat. According to the Harvard School of Public Health, meats should be placed at the very top of any food pyramid, which signifies they should be eaten sparingly and in small portions. In comparison, poultry and fish are usually placed further down on food pyramids, reflecting higher consumption recommendations and greater health value.

5

10

15

20

25

- Main Benefits: Meat is an excellent source of protein in terms of both quantity and quality. For example, a typical 3-ounce portion of meat contains about 21 grams of protein, which is almost half the recommended dietary allowance for women each day. Meat is also a complete protein source because it contains all the essential amino acids your body cannot make. Amino acids are important building blocks for most tissues, such as muscle and skin, as well as enzymes. Meat is also a very good source of certain vitamins and minerals, especially most B vitamins, iron, zinc and phosphorus. Additionally, the density and caloric value of meat usually satisfies hunger better compared to equivalent amounts of vegetable foods.
- Main Disadvantages: The main disadvantages of eating meat are pertaining with ethical considerations, including moral (e.g., killing animals), cultural and religious (Kosher and Halal issues, for example) highly important issues. Other disadvantages are related to the fat, cholesterol and sodium content. People need some dietary fat and cholesterol on a regular basis for energy and to make cellular membranes and certain hormones. However, meat is relatively high in saturated fat and cholesterol, which increase the risk of cardiovascular diseases such as heart attack and atherosclerosis. Sodium is also a serious concern because too much leads to high blood pressure and increased risk of stroke and heart failure. On average, processed meat contains approximately 310 milligrams of sodium per ounce, whereas unprocessed meat has about 77 milligrams. Younger adults should aim for less than 2,300 milligrams of sodium daily, whereas elderly adults and those with risk factors for hypertension should consume less than 1,500 milligrams per day.
- [5] Modest to aggressive changes in diet can improve health and reduce annual national medical expenditures by \$60 billion to \$120 billion, see Doll, Timothy M., et al. "Potential health benefits and medical cost savings from calorie, sodium, and saturated fat reductions in the American diet." *American Journal of Health Promotion* 23.6 (2009): 412-422 and to Karppanen, Heikki, and Eero Mervaala. "Sodium intake and hypertension." *Progress in cardiovascular diseases* 49.2 (2006): 59-75 both are incorporated herein as a reference. They

have estimate that permanent 100-kcal reductions in daily intake would eliminate approximately 71.2 million cases of overweight/obesity and save \$58 billion annually. Long-term **sodium** intake reductions of **400 mg/d** in those with uncontrolled hypertension would eliminate about 1.5 million cases, saving \$2.3 billion annually. Decreasing **5 g/d of saturated fat** intake in those with elevated cholesterol would eliminate 3.9 million cases, saving \$2.0 billion annually. Likewise, according to Kalmijn, Sandra, et al. "Dietary fat intake and the risk of incident dementia in the Rotterdam Study." *Annals of neurology* 42.5 (1997): 776-782 which is incorporated herein as a reference, high intakes of the following nutrients were associated with an increased risk of dementia: total fat (RR = 2.4% intake as percentage of total energy intake), **saturated fat** (RR = 1.9%) and **cholesterol** (RR = 1.7%).

5

10

15

20

25

- [6] Hybrid foods of the present invention comprises significantly less quantities of harmful substances, such as cholesterol, fat and saturated fat, as shown in **Table 4** below.
- [7] Hence, the present invention discloses a beef-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein said harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <8 gr and <3.5 gr per 100 gr, respectfully; and wherein said beef-flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.
- [8] The present invention also discloses a pork-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein said harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <5 gr and <1 gr per 10Ogr, respectfully; and wherein said pork-flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.
- [9] The present invention further discloses a chicken-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein said harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <4 gr and <0.5 gr per lOOgr, respectfully; and wherein said chicken -flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.
- [10] The present invention further discloses a duck-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein said harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <4 gr and <0.5 gr per lOOgr, respectfully; and wherein said chicken -flavored hybrid foodstuff comprises from about 82% to

about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.

**Table 4** Cholesterol, fat and saturated fat content in 100 gr of hybrid foods of the current invention, as compared with 100 gr non-cultured (commercially available) foods, adapted from UCSF Medical Center web site, currently available at www(dot)ucsfhealth(dot)org/education/(dot)

5

10

15

Meats & Protein	Cholesterol (mg)	Total Fat (g)	Saturated Fat (g)	
Beef (ground, lean)	78	18	7	
Beef (short ribs)	94	42	18	
Beef (sirloin)	89	12	5	
Beef Liver	389	5	2	
Veal (top round)	135	5	2	
Lamb (foreshank)	106	14	6	
Ham	53	6	2	
Pork (tenderloin)	79	6	2	
Pork (chop)	85	25	10	
Chicken Liver	631	6	2	
Chicken (no skin)	85	5	1	
Beef-like hybrid food	<60	<8	<3.5	
Pork-like hybrid food	<60	<5	<1	
Chicken-like hybrid food	<60	<4	<0.5	

It is thus in the scope of the invention wherein a meat-flavored hybrid foodstuff based diet is provided useful for having nutritional value of meat, yet is free of harmful substances at harmful quantities; wherein said harmful substances including saturated fat, cholesterol and sodium wherein harmful quantities are 5 g/d, <200 mg and 400 mg/d, respectfully; and wherein said hybrid foodstuff comprising from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 18% to about 1% (w/w) cultured meat.

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in

the context of a single embodiment, may also be provided separately or in any suitable subcombination or as suitable in any other described embodiment of the invention. Certain features described in the context of various embodiments are not to be considered essential features of those embodiments, unless the embodiment is inoperative without those elements.

Various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below find experimental support in the following examples.

#### **EXAMPLES**

Reference is now made to the following examples, which together with the above descriptions illustrate some embodiments of the invention in a non limiting fashion.

#### 10 Example 1 Hybrid chicken-like sausage

## Ingredients:

5

15

20

- A. ErythrocytesSoy protein tissue 12Kg, soy protein isolate 7.25Kg, sodium caseinate 1.25Kg, potato starch 7.5Kg, corn starch 3Kg;
- B. 75% canola oil, 20% hepatocytes cells and 5% of water, gluten, Guar gum, egg, modified corn starch MC, salt.

^Seasoning step is also optional.

- Step 1 Rehydrated tissue protein: Soy protein tissues are soaked in water for four hours until full absorption of Soy protein and softening.
- Step 2 Brushed: After re-organization of the protein by dry water, a chopping machine brushed is used with low speed, blade speed 1,000 rpm for 2 minutes to give the length and thickness of relatively uniform fibers.
- Step 3 Emulsified: Into the chopping machine ingredients A (isolated soy protein, starch, water, spices) and ingredients B were added using high gear speed, blade speed 5,000 rpm chopping five minutes to achieve fully emulsified mixture.
- 25 Step 4 Vacuum mixing: The emulsion prepared in step 2 of fibrous tissue proteins and Step 3 is placed in a vacuum mixer, fully stirred, for 30 minutes.
  - Step 5 Filling: Accomplished by stuffing sausage filling machine is completed, select a breathable plastic casing, smoke treatment may be carried out, casing diameter 18mm.

- Step 6 Thermal processing: The drying temperature is 58° C, for 10 minutes; smoked temperature 57 °C, for 10 minutes; cooking temperature 82° C, 20 minutes to complete the ripening and sterilization process.
- Step 7 Natural cooling to an internal temperature below 10° C.
- Step 8 Packaging & storage. The airtight plastic bag.  $0^{\circ}$  C to about  $4^{\circ}$  C refrigerated or frozen below -18  $^{0}$  C.

#### Example 2 vegetarian beef-like sausage

### Ingredients:

10

- A. Soy protein tissue 13Kg, soy protein isolate 6Kg, sodium caseinate I. 5Kg, potato starch 5.5Kg, corn starch 5Kg.
- B. 75% canola oil, 20% hepatocytes cells and 5% of water, gluten, Guar gum, egg, MC and salt.

Rehydrated tissue protein - Soy protein tissues were soaked in water for six hours. Soy protein was fully absorbed and become soft.

### 15 Example 3 vegetarian pork-like sausage

#### Ingredients:

- A. Soy protein 15Kg, soy protein isolate 6Kg, sodium caseinate 2Kg, potato starch lOKg.
- B. 75% canola oil, 20% hepatocytes cells and 5% of water, gluten, Guar gum, egg, MC, salt.

#### 20 Example 4 Hybrid chicken-like nuggets

#### Ingredients:

- A. Chickpea protein 300 gr, textured wheat protein 100 gr, soy protein isolate 50 gr, water 15 gr, chicken flavor 2 gr, Erythrocytes cells 0.5 gr (-2%), Dijon mustard 3 Tbsp, canola oil 2 Tbsp, corn starch 1 Tsp, dextrose 0.5 Tsp.
- B. Soy milk 25 gr, flavoring 1 Tbsp.
  - C. Bread crumbles 50 gr, nutritional yeast 1 Tbsp, flavoring 1 Tbsp.
- Step 1 Into a chopping machine ingredients A are added using high gear speed, blade speed 5,000 rpm chopping five minutes to achieve fully emulsified mixture.
- Step 2 Preparation of nugget sized flatten pieces from the mixture.
- 30 Step 3 The nugget pieces are dipped in ingredients B liquid mixture.
  - Step 4 The nugget pieces are dipped in ingredients C breading mixture.

- Step 5 Thermal processing: The frying temperature is 190  $^{\circ}$ C for 4 minutes. Cooking temperature 82  $^{\circ}$ C, for 20 minutes to complete the ripening and sterilization process.
- Step 6 Natural cooling to an internal temperature below 10 °C.
- Step 7 Packaging & storage. An airtight plastic bag. 0 °C to about 4 °C refrigerated or frozen below -18 °C.

## Example 5 Hybrid duck-like burger

#### Ingredients:

5

10

20

25

30

- A. Soy protein 350 gr, textured wheat protein 100 gr, soy protein isolate 50 gr, adipocytes cells 20 gr (-4%), water 15 gr, coconut oil 10 gr, duck flavor 2 gr, canola oil 2 Tbsp, corn starch 1 Tsp, dextrose 0.5 Tsp.
- Step 1 Into a chopping machine ingredients A are added using high gear speed, blade speed 5,000 rpm chopping five minutes to achieve fully emulsified mixture.
- Step 2 Preparation of flatten patties from the mixture.
- 15 Step 3 Thermal processing: The frying temperature is 190 °C for 4 minutes. Cooking temperature 82 °C, for 20 minutes to complete the ripening and sterilization process.
  - Step 4 Natural cooling to an internal temperature below 10 °C.
  - Step 5 Packaging & storage. An airtight plastic bag. 0 °C to about 4 °C refrigerated or frozen below -18 °C.
  - Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

All publications, patents and patent applications mentioned in this specification are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference. In addition, citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention. To the extent that section headings are used, they should not be construed as necessarily limiting.

## WHAT IS CLAIMED IS:

1. A method of producing a hybrid foodstuff, the method comprising combining a plant-originated substance with an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 30 % (w/w) of the hybrid foodstuff.

PCT/IL2018/050398

- 2. A method of providing nutrition to a subject in need thereof, the method comprising providing the subject with a hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 30 % (w/w) of the hybrid foodstuff.
- 3. The method of claim 2, wherein the subject is at risk of nutritional deficiency.
- 4. A method of qualifying a hybrid foodstuff, the method comprising subjecting the hybrid foodstuff to at least one organoleptic test.
- 5. A hybrid foodstuff comprising a plant-originated substance and an amount of cultured animal cells so as to enhance a meat organoleptic and/or meat nutritional property in the hybrid foodstuff, wherein said animal cells do not form a tissue, and wherein said amount is below 30 % (w/w) of the hybrid foodstuff.
- 6. The method or hybrid foodstuff of any one of claims 1-5, wherein said hybrid foodstuff comprises from about 82% to about 99.5% (w/w) plant-originated substance and/or additives thereof and about 0.5% to about 18% (w/w) cultured animal cells.
- 7. The method or hybrid foodstuff of any one of claims 1-6, wherein said amount is minimal weight percent.
- 8. The method or hybrid foodstuff of claim 7, wherein said amount provides said hybrid foodstuff with one 9-PHS point more than a corresponding foodstuff devoid of said animal cells.

- 9. The method or hybrid foodstuff of any one of claims 1-8, wherein said foodstuff comprises about 70 % to about 99.7% (w/w) plant-originated substances and/or additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 0.3% to 30% (w/w), where the improvement is that said oil portion consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured animal cells.
- 10. The method or hybrid foodstuff of any one of claims 1-9, wherein said cultured animal cells are selected from the group consisting of hepatocytes, myoblasts, osteoblasts, fibroblasts, lipoblasts, odontoblasts, adult neuronal progenitor cells, neural stem cells, multipotent stem cells from subventricular forebrain region, ependymal-derived neural stem cells, hematopoietic stem cells, liver-derived hematopoietic stem, marrow-derived stem cell, adipo-fibroblasts, adiposederived stem cells, islet- cells producing stem cells, pancreatic-derived pluripotent islet-producing stem cells, mesenchymal stem cells, placenta cells, bone marrow stromal cells, muscle side population cells, bone marrow-derived recycling cells, blood-derived mesenchymal precursor cells, bone-marrow derived side population cells, muscle precursor cells, circulating skeleton stem cells, neural progenitor cells, multipotent adult progenitor cells, mesodermal progenitor cells, spinal cord progenitor cells and spore-like cell, and any combinations thereof.
- 11. The method or hybrid foodstuff of any one of claims 1-10, wherein the hybrid foodstuff has the organoleptic properties of a meat product, selected from the group consisting of a chicken, duck, pork and beef.
- 12. The method or hybrid foodstuff of any one of claims 1-11, wherein said hybrid foodstuff comprises less than about 0.5 gr per 100 gr saturated fat, less than about 60 mg per 100 gr cholesterol and less than about 220 mg per 100 gr sodium.
- 13. The method or hybrid foodstuff of any one of claims 1-12, wherein said animal cells are of a single cell type or single cell lineage.
- 14. The method or hybrid foodstuff of any one of claims 1-12, wherein said animal cells are of no more than two cell types or two cell lineages.
- 15. The method or hybrid foodstuff of any one of claims 1-12, wherein said animal cells are of no more than three cell types or three cell lineages.

- 16. The method or hybrid foodstuff of any one of claims 1-15, wherein said foodstuff is free of growth hormones in an amount hazardous for human or animal consumption.
- 17. Hybrid foodstuff comprising from about 82% to about 99.5% (w/w) plant-originated substances and additives thereof and about 0.5% to about 18% (w/w) cultured meat.
- 18. Hybrid foodstuff of claim 17, wherein the portion of the cultured meat in the end product is defined as the minimal weight percent providing said end product with one 9-PHS point more than the corresponding foodstuff being non-hybrid, cultured meat free vegetable-based product.
- 19. Hybrid foodstuff with improved organoleptic and nutritional properties, comprising about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 0.3% to 30% (w/w), where the improvement is that said oil portion consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat.
- 20. Hybrid foodstuff with improved organoleptic and nutritional properties, comprising about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 3% to 30% (w/w), where the improvement is that said oil portion consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat; wherein the smallest portion of the cultured meat in the end product is defined as the minimal weight percent providing said end product with at least one 9-PHS point more than a corresponding foodstuff being non-hybrid, cultured meat free vegetable-based product.
- 21. Hybrid foodstuff with improved organoleptic and nutritional properties, characterized by about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (vegetable portion) and water immiscible materials (oil portion), ranging from about 3% to 30% (w/w); said oil portion consists of up to about 30% (w/w) or less vegetable oil, and from about 0.3% to about 30% (w/w) cultured meat; wherein the role said mixtures in the oil portion (measured as vegetable oil to culture meat (wt/wt) in improving the organoleptic properties (measured as 9-PHS) of said hybrid foodstuff of is provided by any of **Formula** I, II or III.

22. Hybrid foodstuff with improved organoleptic and/or nutritional properties, characterized by about 70% to about 99.7% (w/w) plant-originated substances and additives thereof (e.g., gum, binding, thickening, aroma flavor additives) and/or water immiscible materials (oil portion), ranging from about 3% to 30% (w/w), where the improvement is that said additives thereof (e.g., gum, binding, thickening, aroma flavor additives) and/or water immiscible materials (oil portion)consists of about 30% (w/w) or less vegetable oil, and from about 0.3% to

about 30% (w/w) cultured meat.

- 23. Hybrid foodstuff according to any of claim 17 to 22, wherein at least about 50% of cultured meat portion is cultured meat cells selected from a group consisting of heme-proteins (HPs), hepatocytes, myoblasts, osteoblasts, fibroblasts, lipoblasts, odontoblasts, adult neuronal progenitor cells, neural stem cells, multipotent stem cells from subventricular forebrain region, ependymal-derived neural stem cells, hematopoietic stem cells, liver-derived hematopoietic stem, marrow-derived stem cell, adipo-fibroblasts, adipose-derived stem cells, islet- cells producing stem cells, pancreatic-derived pluripotent islet-producing stem cells, mesenchymal stem cells, placenta cells, bone marrow stromal cells, muscle side population cells, bone marrow-derived recycling cells, blood-derived mesenchymal precursor cells, bone-marrow derived side population cells, muscle precursor cells, circulating skeleton stem cells, neural progenitor cells, multipotent adult progenitor cells, mesodermal progenitor cells, spinal cord progenitor cells and spore-like cell, mixtures and any combinations thereof.
- 24. Hybrid foodstuff according to any of claim 17 to 22, wherein said foodstuff is a consumable, edible item having the final organoleptic properties of a meat product, and especially a product selected from the group consisting of Beef, Beef heart, Beef liver, Beef tongue, Bone soup from allowable meats, Buffalo, Bison, Calf liver, Caribou, Goat, Ham, Horse, Kangaroo, Lamb, Marrow soup, Moose, Mutton, Opossum, Organ Meats, Pork, Bacon, Rabbit, Snake, Squirrel, Sweetbreads, Tripe, Turtle, Veal, Venison, Chicken, Chicken Liver, Cornish Game Hen, Duck, Duck Liver, Emu, Gizzards, Goose, Goose Liver, Grouse, Guinea Hen, Liver, Ostrich, Partridge, Pheasant, Quail, Squab, Turkey, any mixture and a combination thereof.
- 25. A meat-flavored hybrid foodstuff-based diet free of harmful substances at harmful quantities; wherein said harmful substances including saturated fat, cholesterol and sodium; wherein harmful quantities are 5 g/d, <200 mg and 400 mg/d, respectfully; and wherein said

each of said hybrid foodstuff comprising from about 82% to about 99% (w/w) plant-originated

substances and additives thereof and about 1% to about 18% (w/w) cultured meat.

26. A beef-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein said harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <8 gr and <3.5 gr per 100 gr, respectfully; and wherein said beef-flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.

- 27. A pork-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein said harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <5 gr and <1 gr per 100g, respectfully; and wherein said pork-flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.
- 28. A chicken-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein said harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <4 gr and <0.5 gr per lOOgr, respectfully; and wherein said chicken -flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.
- 29. A duck-flavored hybrid foodstuff free of harmful substances at harmful quantities; wherein said harmful substances including cholesterol, total fat and saturated fat; wherein harmful quantities are <60 mg, <4 gr and <0.5 gr per IOOgr, respectfully; and wherein said duck -flavored hybrid foodstuff comprises from about 82% to about 99% (w/w) plant-originated substances and additives thereof and about 1% to about 18% (w/w) cultured meat.
- 30. A method of preparing a hybrid foodstuff comprising the steps of: providing a plant-originated substance;

providing cultured meat, cells and substances thereof;

interconnecting said plant-originated substance with said cultured meat, cells and substances thereof; and

forming said hybrid foodstuff product into a desired form.

WO 2018/189738 PCT/IL2018/050398

31. The method according to claim 30, wherein said plant-originated substance is from about 82% to about 99.5% (w/w).

- 32. The method according to claim 30 or 31, wherein said cultured meat, cells and substances thereof is from about 0.5% to about 18% (w/w).
- 33. A method of preparing a hybrid foodstuff comprising the steps of:
  providing a plant-originated substance;
  providing water immiscible materials;
  providing cultured meat, cells and substances thereof;
  interconnecting said plant-originated substance with said cultured meat, cells and substances thereof and with said water immiscible materials; and
- 34. The method according to claim 33, wherein said plant-originated substance is from about 70% to about 99.7% (w/w).

forming said hybrid foodstuff product into a desired form.

- 35. The method according to claim 33, wherein said water immiscible materials is from about 3% to 30% (w/w).
- 36. The method according to claim 33, wherein said cultured meat, cells and substances thereof is from about 0.3% to about 30% (w/w).
- 37. The method according to claims 30 and 33, wherein at least about 50% of cultured meat portion are hepatocytes.
- 38. The method according to claims 30 and 33, wherein at least about 50% of cultured meat portion is cultured meat cells selected from a group consisting of heme-proteins (HPs), hepatocytes, myoblasts, osteoblasts, fibroblasts, lipoblasts, odontoblasts, adult neuronal progenitor cells, neural stem cells, multipotent stem cells from subventricular forebrain region, ependymal-derived neural stem cells, hematopoietic stem cells, liver-derived hematopoietic stem, marrow-derived stem cell, adipo-fibroblasts, adipose-derived stem cells, islet- cells producing stem cells, pancreatic-derived pluripotent islet-producing stem cells, mesenchymal stem cells, placenta cells, bone marrow stromal cells, muscle side population cells, bone

marrow-derived recycling cells, blood-derived mesenchymal precursor cells, bone-marrow derived side population cells, muscle precursor cells, circulating skeleton stem cells, neural progenitor cells, multipotent adult progenitor cells, mesodermal progenitor cells, spinal cord progenitor cells and spore-like cell, mixtures and any combinations thereof.

- 39. The method according to claims 30 and 33, wherein said foodstuff is a consumable, edible item having the final organoleptic properties of a meat product, and especially a product selected from the group consisting of Beef, Beef heart, Beef liver, Beef tongue, Bone soup from allowable meats, Buffalo, Bison, Calf liver, Caribou, Goat, Ham, Horse, Kangaroo, Lamb, Marrow soup, Moose, Mutton, Opossum, Organ Meats, Pork, Bacon, Rabbit, Snake, Squirrel, Sweetbreads, Tripe, Turtle, Veal, Venison, Chicken, Chicken Liver, Cornish Game Hen, Duck, Duck Liver, Emu, Gizzards, Goose, Goose Liver, Grouse, Guinea Hen, Liver, Ostrich, Partridge, Pheasant, Quail, Squab, Turkey, any mixture and a combination thereof.
- 40. The method of hybrid foodstuff of any one of claims 17-39, wherein animal-derived protein in the hybrid foodstuff is below 50 % (w/w).
- 41. The method of hybrid foodstuff of any one of claims 17-40, wherein said foodstuff is not a chip and said plant-originated substance is not a plant derived hydrogel.
- 42. The method of claim 4, wherein the hybrid foodstuff comprises the hybrid foodstuff of any one of claims 5-29 and 40-41.
- 43. The method or hybrid foodstuff of any one of claims 17-42, wherein said foodstuff is a vegetarian foodstuff.
- 44. The method or hybrid foodstuff of any one of claims 17-42, wherein said foodstuff is a vegan foodstuff.
- 45. The method or hybrid foodstuff of any one of claims 17-42, wherein said foodstuff is a meat substitute.

#### INTERNATIONAL SEARCH REPORT

International application No PCT/IL2018/050398

a. classification of subject matter INV. A23J3/22 A23L A23L13/00 ADD. According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED searched (classification system followed by classification symbols) A23L C12N A23J Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal , WPI Data C. DOCUMENTS CONSIDERED TO BE RELEVANT Category\* Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. Υ wo 2013/016547 A2 (UNIV MISSOURI [US]; 1-45 FORGACS GABOR [US]; MARGA FRANCOISE [US]; JAKAB KA) 31 January 2013 (2013-01-31) paragraphs [0072] , [0118] Χ W0 2013/010042 AI (SAND HILL FOODS INC 1-45 [US]; BROWN PATRICK [US]; VRLJIC MARIJA [US]; VARA) 17 January 2013 (2013-01-17) Υ exampl es 1-45 wo 2006/041429 A2 (VEIN JON [US]) Α 1-45 20 Apri I 2006 (2006-04-20) abstract Α WO 2015/161099 AI (SAVAGE RIVER INC DBA 1-45 BEYOND MEAT [US]) 22 October 2015 (2015-10-22) claim 1 X See patent family annex. Further documents are listed in the continuation of Box C. \* Special categories of cited documents : ater document published after the international filing date or priority date and not in conflict with the application but cited to understand "A" document defining the general state of the art which is not considered the principle or theory underlying the invention to be of particular relevance "E" earlier application or patent but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive filing date ocumentwhich may throw doubts on priority claim(s) orwhich is cited to establish the publication date of another citation or other step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be special reason (as specified) considered to involve an inventive step when the document combined with one or more other such documents, such combination being obvious to a person skilled in the art "O" document referring to an oral disclosure, use, exhibition or other "P" document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 27/07/2018 18 July 2018 Name and mailing address of the ISA/ Authorized officer European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016 Saunders , Thomas

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/IL2018/050398

Patent document cited in search report	Publication date		Patent family member(s)		Publication date
wo <b>2013016547</b> A	2 31-01-2013	AU	2012286817	Al	13-02-2014
		ΑU	2016204474	ΑI	21-07-2016
		CA	2842837	Αl	31-01-2013
		CN	103747693	Α	23-04-2014
		EP	2736357	A2	04-06-2014
		JР	2014521336	Α	28-08-2014
		KR	20140050064	Α	28-04-2014
		US	2013029008	Αl	31-01-2013
		US	2014093618	Αl	03-04-2014
		Wo	2013016547	A2	31-01-2013
wo 2013010042 AI	d 17-01-2013	AU	2012281069	Al	13 -02 -2014
		ΑU	2016231499	Αl	06-10-2016
			112014000614	A2	11-07 -2017
		CA	2841473	Αl	17-01-2013
		CN	103889243	Α	25 -06 -2014
		CN	105724726	Α	06-07-2016
		EP	2731446	ΑI	21-05 -2014
		JР	2014520554	Α	25 - 08 - 2014
		KR	20140047125	Α	21-04-2014
		RU	2014104812	Α	27-08-2015
		US	2014193547	Αl	10-07-2014
		US	2015289541	ΑI	15-10-2015
		US	2017290363	ΑI	12 <b>-10 -201</b> 7
		WO	2013010042	Al	17-01-2013
wo 2006041429 A2	2 20-04-2006	AT	536750	T	15-12-2011
		ΑU	2004324099	ΑI	20-04-2006
		BR	PI0419055	Α	11-12 -2007
		CA	2580433	ΑI	20-04-2006
		DK	1789063	T3	26-03-2012
		EA	200700660		26-10-2007
		EP	1789063		30 - 05 - 2007
		ES	2378822		18 -04 -2012
		۱L	181859		31-03-2011
		JР	2008513006		01-05-2008
		KR	20070065381		22-06-2007
		WO	2006041429	A2	20-04-2006
wo <b>2015161099</b> Al	d 22-10-2015	AU	2015247529		24-11-2016
		CA	2946125		22-10-2015
		CN	106455622		22-02-2017
		EP	3131406	ΑI	22-02-2017
		JР	2017517273		29 - 06 - 2017
		US	2015296834		22-10-2015
		WO	2015161099	ΑI	22-10-2015