



US 20070054918A1

(19) **United States**(12) **Patent Application Publication****Braje et al.**(10) **Pub. No.: US 2007/0054918 A1**(43) **Pub. Date: Mar. 8, 2007**(54) **N-[(PIPERAZINYL)HETARY]ARYLSULFONAMIDE COMPOUNDS WITH AFFINITY FOR THE DOPAMINE D3 RECEPTOR**(76) Inventors: **Wilfried Braje**, Rinteln (DE); **Andreas Haupt**, Schwetzingen (DE); **Wilfried Lubisch**, Heidelberg (DE); **Roland Grandel**, Dossenheim (DE); **Karla Drescher**, Dossenheim (DE); **Herve Geneste**, Neuhausen (DE); **Lilliane Unger**, Ludwigshafen (DE); **Daryl R. Sauer**, Trevor, WI (US); **Sean C. Turner**, Mannheim (DE)

Correspondence Address:

WOOD, PHILLIPS, KATZ, CLARK & MORTIMER**500 W. MADISON STREET
SUITE 3800
CHICAGO, IL 60661 (US)**(21) Appl. No.: **10/552,842**(22) PCT Filed: **Apr. 13, 2004**(86) PCT No.: **PCT/EP04/03872**


§ 371(c)(1),

(2), (4) Date: **Aug. 22, 2006**(30) **Foreign Application Priority Data**

Apr. 14, 2003 (US)..... 10413233

Publication Classification(51) **Int. Cl.****A61K 31/506** (2007.01)**A61K 31/501** (2007.01)**A61K 31/496** (2007.01)**C07D 403/14** (2007.01)**C07D 403/02** (2007.01)(52) **U.S. Cl.** **514/252.02; 544/238; 544/364; 544/295; 514/252.18; 514/253.01**(57) **ABSTRACT**

The invention relates to N-[(piperazinyl)hetaryl]arylsulfonamide compounds of the general formula (I) in which Q is a bivalent, 6-membered heteroaromatic radical which possesses 1 or 2 N atoms as ring members and which optionally carries one or two substituents R^a which is/are selected, independently of each other, from halogen, CN, NO₂, CO₂R⁴, COR⁵, C₁—C₄-alkyl and C₁—C₄-haloalkyl; Ar is phenyl or a 6-membered heteroaromatic radical which possesses 1 or 2 N atoms as ring members and which optionally carries one or two substituents R^b, which is/are selected from halogen, NO₂, CN, CO₂R⁴, COR⁵, C₁—C₆-alkyl, C₂—C₆-alkenyl, C₂—C₆-alkynyl, C₃—C₆-cycloalkyl, C₃—C₆-cycloalkyl-C₁—C₄-alkyl and C₁—C₄-haloalkyl, with it also being possible for two radicals R^b which are bonded to adjacent C atoms of Ar to be together C₃—C₄-alkylene; R¹ is hydrogen, C₁—C₄-alkyl, C₁—C₄-haloalkyl, C₃—C₆-cycloalkyl, C₃—C₆-cycloalkyl-C₁—C₄-alkyl, C₁—C₄-hydroxyalkyl, C₁—C₄-alkoxy-C₁—C₄-alkyl, C₃—C₄-alkenyl or C₃—C₄-alkynyl; with the radicals n, R¹, R², R³, R⁴ and R⁵ having the meanings given in the patent claims, to the N-oxides and to the physiologically tolerated acid addition salts of these compounds and to pharmaceutical compositions which comprise at least one N-[(piperazinyl)hetaryl]arylsulfonamide compound as claimed in one of claims 1 to 10 and/or at least one physiologically tolerated acid addition salt of I and/or an N-oxide of I, where appropriate together with physiologically acceptable carriers and/or auxiliary substances for treating diseases which respond to influencing by dopamine D₃ receptor antagonists or agonists, in particular for treating diseases of the central nervous system and disturbances of kidney function.



(I)

**N-[(PIPERAZINYL)HETARY]ARYLSULFONAMIDE
COMPOUNDS WITH AFFINITY FOR THE
DOPAMINE D₃ RECEPTOR**

DESCRIPTION

[0001] The present invention relates to novel N-[(piperazinyl)hetaryl]arylsulfonamide compounds. The compounds possess valuable therapeutic properties and are suitable, in particular, for treating diseases which respond to modulation of the dopamine D₃ receptor.

[0002] Neurons obtain their information by way of G protein-coupled receptors, inter alia. A large number of substances exert their effect by way of these receptors. One of them is dopamine. Confirmed findings exist with regard to the presence of dopamine and its physiological function as a neurotransmitter. Disturbances in the dopaminergic transmitter system result in diseases of the central nervous system which include, for example, schizophrenia, depression and Parkinson's disease. These diseases, and others, are treated with drugs which interact with the dopamine receptors.


[0003] Up until 1990, two subtypes of dopamine receptor had been clearly defined pharmacologically, namely the D₁ and D₂ receptors. More recently, a third subtype was found, namely the D₃ receptor which appears to mediate some effects of antipsychotics and antiparkinsonians (J. C. Schwartz et al., The Dopamine D₃ Receptor as a Target for Antipsychotics, in Novel Antipsychotic Drugs, H. Y. Meltzer, Ed. Raven Press, New York 1992, pages 135-144; M. Dooley et al., Drugs and Aging 1998, 12, 495-514, J. N. Joyce, Pharmacology and Therapeutics 2001, 90, pp. 231-59 "The Dopamine D₃ Receptor as a Therapeutic Target for Antipsychotic and Antiparkinsonian Drugs").

[0004] Since then, the dopamine receptors have been divided into two families. On the one hand, there is the D₂ group, consisting of D₂, D₃ and D₄ receptors, and, on the other hand, the D₁ group, consisting of D₁ and D₅ receptors. Whereas D₁ and D₂ receptors are widely distributed, D₃ receptors appear to be expressed regiospecifically. Thus, these receptors are preferentially to be found in the limbic system and the projection regions of the mesolimbic dopamine system, especially in the nucleus accumbens, but also in other regions, such as the amygdala. Because of this comparatively regiospecific expression, D₃ receptors are regarded as being a target having few side-effects and it is assumed that while a selective D₃ ligand would have the properties of known antipsychotics, it would not have their dopamine D₂ receptor-mediated neurological side-effects (P. Sokoloff et al., Localization and Function of the D₃ Dopamine Receptor, *Arzneim. Forsch./Drug Res.* 42(1), 224 (1992); P. Sokoloff et al. Molecular Cloning and Characterization of a Novel Dopamine Receptor (D₃) as a Target for Neuroleptics, *Nature*, 347, 146 (1990)).

[0005] Compounds having an affinity for the dopamine D₃ receptor have been described in the prior art on various occasions, e.g. in WO 96/02519, WO 96/02520, WO 96/02249, WO 96/02246 and DE 10131543 and WO 99/02503. Some of these compounds possess high affinities for the dopamine D₃ receptor. They have therefore been proposed as being suitable for treating diseases of the central nervous system. Some of the compounds described in these publications possess a piperazinylhetaryl structure.

[0006] The invention is based on the object of providing compounds which act as selective dopamine D₃ receptor ligands.

[0007] This object is achieved by means of N-[(piperazinyl)hetaryl]arylsulfonamide compounds of the general formula I



in which

[0008] R is oxygen, a group N—R³ or a group CR^{3a}R^{3b};

[0009] Q is a bivalent, 6-membered heteroaromatic radical which possesses 1 or 2N atoms as ring members and which optionally carries one or two substituents R^a which is/are selected, independently of each other, from halogen, CN, NO₂, CO₂R⁴, COR⁵, C₁—C₄-alkyl, C₁—C₄-alkoxy, C₁—C₄-haloalkyl, NH₂, NHR⁶, NR⁶R⁷ and C₁—C₄-haloalkoxy;

[0010] Ar is phenyl or a 6-membered heteroaromatic radical which possesses 1 or 2N atoms as ring members and which optionally carries one or two substituents R^b, which is/are selected from halogen, NO₂, CN, CO₂R⁴, COR⁵, NH₂, NHR⁶, NR⁶R⁷, C₁—C₆-alkyl, C₁—C₆-haloalkyl, C₁—C₆-alkoxy, C₁—C₆-haloalkoxy, C₂—C₆-alkenyl, C₂—C₆-alkynyl, C₃—C₆-cycloalkyl, C₃—C₆-cycloalkoxy, C₃—C₆-cycloalkyl-C₁—C₄-alkyl and C₁—C₄-haloalkyl, with it also being possible for two radicals R^b which are bonded to adjacent C atoms of Ar to be together C₃—C₄-alkylene;

[0011] n is 0, 1 or 2;

[0012] R¹ is hydrogen, C₁—C₄-alkyl, C₁—C₄-haloalkyl, C₃—C₆-cycloalkyl, C₃—C₆-cycloalkyl-C₁—C₄-alkyl, C₁—C₄-hydroxyalkyl, C₁—C₄-alkoxy-C₁—C₄-alkyl, C₃—C₄-alkenyl or C₃—C₄-alkynyl;

[0013] R² is C₁—C₄-alkyl, C₁—C₄-haloalkyl or, together with R¹, is C₂—C₅-alkylene or, in the case of n=2, the two radicals R² can together be C₁—C₄-alkylene;

[0014] R³ is hydrogen or C₁—C₄-alkyl;

[0015] R^{3a}, R^{3b} are, independently of each other, hydrogen or C₁—C₄-alkyl;

[0016] R⁴ is C₁—C₄-alkyl, C₁—C₄-haloalkyl, C₂—C₄-alkenyl, C₃—C₈-cycloalkyl, C₃—C₆-cycloalkyl-C₁—C₄-alkyl, phenyl or benzyl;

[0017] R⁵ is hydrogen, C₁—C₄-alkyl, C₁—C₄-haloalkyl, C₂—C₄-alkenyl, C₃—C₆-cycloalkyl, C₃—C₆-cycloalkyl-C₁—C₄-alkyl, phenyl or benzyl; and

[0018] R⁶, R⁷ are each independently selected from C₁—C₄-alkyl, C₁—C₄-haloalkyl or together with the nitrogen to which they are bound form a saturated 3-, 4-, 5- or 6-membered heterocycle, which additionally

may comprise an oxygen atom or an additional nitrogen atom as a ring member and which may carry 1, 2, 3 or 4 C_1-C_4 alkyl groups;

the N-oxides thereof and the physiologically tolerated acid addition salts of these compounds.

[0019] These compounds have not previously been described, with the exception of 4-methyl-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide and 4-chloro-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide, which are offered for sale by Ambinter, Paris, as test substances for exploratory libraries.

[0020] The present invention therefore relates to N-[(piperazinyl)hetaryl]arylsulfonamide compounds of the general formula I, to their N-oxides and to their physiologically tolerated acid addition salts, with the exception of the compounds 4-methyl-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide and 4-chloro-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide.

[0021] The present invention also relates to the use of N-[(piperazinyl)hetaryl]arylsulfonamide compounds of the general formula I, of their N-oxides and of their acid addition salts for producing a pharmaceutical composition for treating diseases which respond to the influence of dopamine- D_3 receptor antagonists or agonists.

[0022] The diseases which respond to the influence of dopamine D_3 receptor antagonists or agonists include, in particular, disturbances and diseases of the central nervous system, in particular affective disturbances, neurotic disturbances, stress disturbances and somatoform disturbances and psychoses, especially schizophrenia and depression and, in addition, disturbances of kidney function, in particular kidney function disturbances which are caused by diabetes mellitus (see WO 00/67847).

[0023] According to the invention, at least one compound of the general formula I having the meanings mentioned at the outset is used for treating the abovementioned indications. Provided the compounds of the formula I possess one or more centers of asymmetry, it is also possible to use enantiomeric mixtures, in particular racemates, diastereomeric mixtures and tautomeric mixtures, preferably, however, the respective essentially pure enantiomers, diastereomers and tautomers.

[0024] It is likewise possible to use physiologically tolerated salts of the compounds of the formula I, especially acid addition salts with physiologically tolerated acids. Examples of suitable physiologically tolerated organic and inorganic acids are hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, C_1-C_4 -alkylsulfonic acids, such as methanesulfonic acid, aromatic sulfonic acids, such as benzenesulfonic acid and toluenesulfonic acid, oxalic acid, maleic acid, fumaric acid, lactic acid, tartaric acid, adipic acid and benzoic acid. Other utilizable acids are described in Fortschritte der Arzneimittelforschung [Advances in drug research], Volume 10, pages 224 ff., Birkhauser Verlag, Basel and Stuttgart, 1966.

[0025] It is likewise possible to use N-oxides of the compounds of the formula I. In the N-oxides of the compounds of the formula I, one or more of the N atoms which is/are ring members, and in particular ring members in the aromatic heterocycles Q and/or Ar, are present as an N-oxide

group. Preference is given to those N-oxides of the formula I in which the ring nitrogen atoms in the piperazine ring do not form any N-oxide group. Particularly preferred N-oxides exhibit a N-oxide group on one or two of the ring nitrogen atoms of Ar and/or Q.

[0026] Here and in that which follows, halogen is fluorine, chlorine, bromine or iodine.

[0027] C_n-C_m -Alkyl (in radicals such as alkoxy, alkoxy-alkyl, alkylthio, alkylamino, dialkylamino etc., as well) is a straight-chain or branched alkyl group having from n to m carbon atoms, e.g. from 1 to 4 carbon atoms. Examples of an alkyl group are methyl, ethyl, n-propyl, iso-propyl, n-butyl, 2-butyl, iso-butyl, tert-butyl, n-pentyl, 2-pentyl, neopentyl, n-hexyl and the like.

[0028] C_1-C_4 -Haloalkyl (in radicals such as haloalkoxy, haloalkoxyalkyl, haloalkylthio, etc., as well) is an alkyl group having from 1 to 4 C atoms in which all or some, e.g. 1, 2, 3 or 4 of the hydrogen atoms, is/are replaced by halogen atoms, in particular by chlorine or fluorine. Preferred haloalkyl is C_1-C_2 -fluoroalkyl or C_1-C_2 -fluorochloroalkyl, in particular CF_3 , CHF_2 , CF_2Cl , CH_2F , and CH_2CF_3 .

[0029] C_1-C_4 -Hydroxyalkyl is a C_1-C_4 -alkyl group which possesses an OH group, such as 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 2-hydroxybutyl, 3-hydroxybutyl, 2-methyl-2-hydroxypropyl etc.

[0030] C_1-C_4 -Alkoxy- C_1-C_4 -alkyl is a C_1-C_4 -alkyl group which carries a C_1-C_4 -alkoxy substituent, e.g. methoxymethyl, ethoxymethyl, 2-methoxyethyl, 1-methoxyethyl, 2-ethoxyethyl, 1-ethoxyethyl, n-propoxymethyl, isopropoxymethyl, n-butoxymethyl, (1-methylpropoxy)methyl, (2-methylpropoxy)methyl, $CH_2OC(CH_3)_3$, 2-(methoxy)ethyl, 2-(ethoxy)ethyl, 2-(n-propoxy)ethyl, 2-(1-methylethoxy)ethyl, 2-(n-butoxy)ethyl, 2-(1-methylpropoxy)ethyl, 2-(2-methylpropoxy)ethyl, 2-(1',1-dimethylethoxy)ethyl, 2-(methoxy)propyl, 2-(ethoxy)propyl, 2-(n-propoxy)propyl, 2-(1-methylethoxy)propyl, 2-(n-butoxy)propyl, 2-(1-methylpropoxy)propyl, 2-(2-methylpropoxy)propyl, 2-(1,1-dimethylethoxy)propyl, 3-(methoxy)propyl, 3-(ethoxy)propyl, 3-(n-propoxy)propyl, 3-(1-methylethoxy)propyl, 3-(n-butoxy)propyl, 3-(1-methylpropoxy)propyl, 3-(2-methylpropoxy)propyl, 3-(1,1-dimethylethoxy)propyl, 2-(methoxy)butyl, 2-(ethoxy)butyl, 2-(n-propoxy)butyl, 2-(1-methylethoxy)butyl, 2-(n-butoxy)butyl, 2-(1-methylpropoxy)butyl, 2-(2-methylpropoxy)butyl, 2-(1,1-dimethylethoxy)butyl, 3-(methoxy)butyl, 3-(ethoxy)butyl, 3-(n-propoxy)butyl, 3-(1-methylethoxy)butyl, 3-(n-butoxy)butyl, 3-(1-methylpropoxy)butyl, 3-(2-methylpropoxy)butyl, 3-(1,1-dimethylethoxy)butyl, 4-(methoxy)butyl, 4-(ethoxy)butyl, 4-(n-propoxy)butyl, 4-(1-methylethoxy)butyl, 4-(n-butoxy)butyl, 4-(1-methylpropoxy)butyl, 4-(2-methylpropoxy)butyl or 4-(1,1-dimethylethoxy)butyl, preferably methoxymethyl, ethoxymethyl, 2-methoxyethyl, 2-ethoxyethyl, 2-(methoxy)propyl, 2-(ethoxy)propyl or 3-(methoxy)propyl, or 3-(ethoxy)propyl.

[0031] C_3-C_6 -Cycloalkyl is a cycloaliphatic radical having from 3 to 6 C atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

[0032] C_3-C_6 -Cycloalkyl- C_1-C_4 -alkyl is a C_1-C_4 -alkyl group which carries a C_3-C_6 -cycloalkyl radical, e.g. cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl,

1-cyclopropylethyl, 1-cyclobutylethyl, 1-cyclopentylethyl, 2-cyclopropylethyl, 2-cyclobutylethyl, 2-cyclopentylethyl, 1-cyclopropylpropyl, 1-cyclobutylpropyl, 1-cyclopentylpropyl, 2-cyclopropylpropyl, 2-cyclobutylpropyl, 2-cyclopentylpropyl, 3-cyclopropylpropyl, 3-cyclobutylpropyl, 3-cyclopentylpropyl, 1-cyclopropyl-1-methylethyl, 1-cyclopentyl-1-methylethyl, 1-cyclopentyl-1-methylethyl, 3-cyclohexylpropyl, 1-cyclohexyl-1-methylethyl, 1-cyclohexyl-1-methylethyl or 1-cyclohexyl-1-methylethyl.

[0033] C_2-C_4 -Alkenyl (in radicals such as alkenyloxy as well) is a singly unsaturated hydrocarbon radical having 2, 3, 4, 5 or 6 atoms, e.g. vinyl, allyl(2-propen-1-yl), 1-propen-1-yl, 2-propen-2-yl, methallyl(2-methylprop-2-en-1-yl) and the like. C_3-C_4 -Alkenyl is, in particular, allyl, 1-methylprop-2-en-1-yl, 2-buten-1-yl, 3-buten-1-yl, methallyl, 2-penten-1-yl, 3-penten-1-yl, 4-penten-1-yl, 1-methylbut-2-en-1-yl or 2-ethylprop-2-en-1-yl.

[0034] C_3-C_6 -Alkynyl (in radicals such as alkynyloxy as well) is a hydrocarbon radical having 2, 3, 4, 5 or 6 C atoms which possesses a triple bond, e.g. propargyl (2-propyn-1-yl), 1-methylprop-2-yn-1-yl, 2-butyne-1-yl, 3-butyne-1-yl, 2-pentyne-1-yl, 1-pentyne-3-yl, etc.

[0035] Examples of 6-membered heteroaromatic radicals which possess 1 or 2 nitrogen atoms as ring members are, in particular, 2-, 3- or 4-pyridinyl, 2-, 4- or 5-pyrimidinyl, 2- or 3-pyrazinyl and 3- or 4-pyridazinyl. Examples of bivalent, 6-membered heteroaromatic radicals which possess 1 or 2 nitrogen atoms as ring members are, in particular, pyridin-2,4-diyl, pyridin-2,5-diyl, pyridin-2,6-diyl, pyridin-3,5-diyl, pyrimidin-2,4-diyl, pyrimidin-2,5-diyl, pyrimidin-4,6-diyl, pyrazin-2,5-diyl, pyrazin-2,6-diyl, pyridazin-3,6-diyl and pyridazin-3,5-diyl.

[0036] With regard to using the compounds according to the invention as dopamine D_3 receptor ligands, preference is given to those compounds of formula I in which the piperazin ring is bonded to the heteroaromatic radical Q in the meta position or, in particular, in the para position with respect to the group R.

[0037] In one embodiment the radical R^a is selected from halogen, CN, NO_2 , CO_2R^4 , COR^5 , C_1-C_4 -alkyl and C_1-C_4 -haloalkyl. Preferred meanings of R^4 and R^5 are, independently of each other, H or C_1-C_4 -alkyl. In another embodiment R^a is selected from C_1-C_4 -alkoxy, C_1-C_4 -haloalkoxy, NH_2 , NHR^6 and NR^6R^7 . In this embodiment preferred meanings of R^6 and R^7 are independently of each other methyl or ethyl.

[0038] The heteroaromatic radical Q may be unsubstituted or possess a substituent R^a which is selected from halogen, C_1-C_4 -alkyl and C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy, NH_2 , NHR^6 , NR^6R^7 and C_1-C_4 -haloalkoxy. In one embodiment the radical R^a is selected from halogen, C_1-C_4 -alkyl and C_1-C_4 -haloalkyl. In another preferred embodiment R^a is selected from C_1-C_4 -alkoxy, C_1-C_4 -haloalkoxy, NH_2 , NHR^6 and NR^6R^7 with specific preference given to methoxy, NH_2 , methylamino, dimethylamino, ethylamino, diethylamino and methylethyl amino.


[0039] In a very preferred embodiment, Q is unsubstituted.

[0040] In another very preferred embodiment, Q carries a radical selected from C_1-C_4 -haloalkoxy, C_1-C_4 -alkoxy,

C_1-C_4 -haloalkyl, C_1-C_4 -alkyl, NH_2 , NHR^6 or NR^6R^7 , and especially methoxy or methyl.

[0041] Preference is given to the variables Q, R^1 , R^2 , R^3 and Ar preferably having, independently of each other, the meanings given below:

[0042] Q is preferably a radical of the formula A:



in which A_1 , A_2 and A_3 are, independently of each other, N or CH, and one or two of the variables A_1 , A_2 and A_3 can also be $C-R^1$, with A_1 , A_2 and A_3 not simultaneously being N or being simultaneously selected from CH and $C-R^a$. In formula A, k is 0 or 1 and R^a has the previously mentioned meanings.

[0043] In formula A, R^a is preferably selected from halogen, especially chlorine or fluorine, C_1-C_4 -alkyl, especially methyl, and C_1-C_4 -haloalkyl, especially trifluoromethyl, C_1-C_4 -alkoxy especially methoxy, C_1-C_4 -haloalkoxy, especially difluoromethoxy or trifluoromethoxy, NH_2 , NHR^6 especially methylamino or ethylamino, and NR^6R^7 , especially dimethylamino, diethylamino or methyl-ethylamino. The C atom which is located between the atoms A_1 and A_3 preferably carries the piperazinyl radical.

[0044] In particular, none of the variables A_1 , A_2 and A_3 is $C-R^a$. Preferred radicals Q are those of formula A, in which A_1 and/or A_3 is/are N, the remaining variable A_1 or A_2 is CH or $C-R^a$, A_2 is CH, and the piperazinyl radical is bonded to the C atom which is located between A_1 and A_3 .

[0045] Preference is furthermore given to compounds of the formula I, in which A_1 and A_2 in formula A are N and A_3 is CH or $C-R^a$.

[0046] In a very preferred embodiment k in formula A is 0. In particular, Q is pyridin-2,5-diyl or pyrimidin-2,5-diyl which are unsubstituted or able to possess a substituent R^a which is different from hydrogen. The piperazinyl radical is then preferably arranged in the 2 position.

[0047] In another very preferred embodiment k in formula A is 1, A_3 is N. A_1 and A_2 are, independently of each other, N or CH and R^a is selected from C_1-C_4 -alkoxy, C_1-C_4 -alkyl, NH_2 , NHR^6 , NR^6R^7 and C_1-C_4 -haloalkoxy. In particular R^a is methoxy or methyl. Most preferably the piperazine radical is located in the 2 position. In this embodiment compounds are especially preferred, in which A_1 is N or CH, in particular CH and A_2 is CH.

[0048] Ar is preferably phenyl or pyridyl which, where appropriate, possesses one or two of the abovementioned substituents R^b . With regard to using the compounds according to the invention as dopamine D_3 receptor ligands, preference is given to those compounds of formula I in which Ar carries one substituent R^b in the para position and, where appropriate, a further substituent R^b in the ortho position or in the metaposition, in each case related to the

binding site for the sulfonamide group. The radicals R^b may be identical or different. Preference is given to the radicals R^b in the para position being selected from C_2-C_6 -alkyl, C_2-C_6 -alkenyl, C_2-C_6 -alkynyl, C_3-C_6 -cycloalkyl, NH_2 , NHR^6 and NR^6R^7 .

[0049] In one preferred embodiment the radical R^b in the para position is selected from C_2-C_6 -alkyl, C_2-C_6 -alkenyl, C_2-C_6 -alkynyl, C_3-C_6 -cycloalkyl, in particular, from branched C_3-C_6 -alkyl, especially isopropyl, and C_3-C_6 -cycloalkyl, especially cyclopropyl. Very particular preference is given to the radical R^b which is arranged in the para position of Ar being isopropyl.

[0050] In one preferred embodiment the radical R^b in the para position is selected from NHR^6 and NR^6R^7 . In this embodiment preferred meanings of R^6 and R^7 are independently of each other methyl or ethyl or form together with the N atom a saturated 3-, 4-, 5- or 6-membered heterocycle, which additionally may comprise an oxygen atom or an additional nitrogen atom as a ring member and which may carry 1, 2, 3 or 4 C_1-C_4 alkyl groups, e.g. a radical from the group consisting of aziridinyl, azetidiny, pyrrolidinyl, piperidinyl, morpholinyl etc.

[0051] Preferred radicals R^b in the meta position or ortho position are selected from halogen, especially chlorine and fluorine, C_1-C_4 -alkyl, especially methyl, CN, trifluoromethyl and difluoromethyl.


[0052] With regard to using the compounds according to the invention as dopamine D_3 receptor ligands, preference is given to those compounds of the formula I in which R^1 is different from hydrogen, in particular hydrogen and methyl. In particular, R^1 is C_2-C_3 -alkyl, cyclopropylmethyl or, particularly preferably, ethyl, allyl or n-propyl.

[0053] The variable n is preferably 0 or 1. Provided n is $\neq 0$, R^2 is preferably methyl. When n is $\neq 0$, the group R^2 is preferably bonded to a carbon atom in the piperazine ring which is adjacent to the group R^1-N . In particularly preferred compounds, n=0. Particular preference is also given to compounds of the formula I in which it applies that n=1 and R^2 is a methyl group which is bonded to a carbon atom in the piperazine ring which is adjacent to the group R^1-N . The compounds can then be present as a racemate, as pure enantiomers or as nonracemic mixtures of the enantiomers. Among these, particular preference is given to those compounds in which the C atom which carries the methyl group exhibits the S configuration.

[0054] R^3 is preferably hydrogen or C_1-C_4 -alkyl and, in particular, hydrogen.

[0055] If R is a group $CR^{3a}R^{3b}$, at least one of the radicals R^{3a} or R^{3b} is hydrogen. More preferably both radicals R^{3a} and R^{3b} are hydrogen.

[0056] Among the compounds of the general formula I, preference is given to the compounds I, wherein R is $N-R^3$, wherein R^3 is as defined above and especially H. Amongst these compounds preference is given to the compounds of the general formula Ia




in which n, R^1 , R^2 , R^3 , R^a and R^b have the previously mentioned meanings, in particular the meanings specified as being preferred, and in which A_1 , A_2 and A_3 are, independently of each other, N or CH, and one of the variables A_1 , A_2 and A_3 can also be $C-R^a$, with A_1 , A_2 and A_3 not simultaneously being N or simultaneously being selected from CH and $C-R^a$, and X and Y are selected from CH, $C-R^b$ and N, in which R^b is halogen, methyl, CN, difluoromethyl or trifluoromethyl, with X and Y not simultaneously being N or simultaneously being $C-R^b$, and k is 0 or 1. R^a has the previously mentioned meanings. In particular, R^a is selected from halogen, especially chlorine or fluorine, C_1-C_4 -alkyl, especially methyl, and C_1-C_4 -haloalkyl, especially trifluoromethyl, C_0-C_4 -alkoxy especially methoxy, C_1-C_4 -haloalkoxy, especially difluoromethoxy or trifluoromethoxy, NH_2 , NHR^6 especially methylamino or ethylamino, and NR^6R^7 , especially dimethylamino, diethylamino or methylethylamino.

[0057] In particular, none of the variables A^1 , A^2 and A^3 is $C-R^a$. Preferred compounds Ia are those in which A_1 and/or A_3 is/are N, the remaining variable A_1 or A_2 is CH or $C-R^a$, A_2 is CH.

[0058] In particular preferred embodiment, k=0. Among these, preference is furthermore given to compound I in which A_1 and A_2 are N and A_3 is CH or $C-R^a$. Among these, preference is given to those compounds of the formula Ia in which X or Y is CH or N and, in particular, both are CH.


[0059] In another very preferred embodiment k in formula Ia is 1, A_3 is N, A_1 and A_2 are, independently of each other, N or CH and R^a is selected from C_1-C_4 -alkyl, C_1-C_4 -alkoxy, C_1-C_4 -haloalkyl, NH_2 , NHR^6 , NR^6R^7 and C_1-C_4 -haloalkoxy. In particular R^a is methoxy or methyl. In this embodiment compounds are especially preferred, in which A_1 is N or CH, in particular CH and A_2 is CH.

[0060] Among the compounds of general formula Ia, preference is given to the compounds of general formula Ia.1







in which n, X, Y, R^1 , R^2 , R^3 , R^a and R^b have the previously mentioned meanings, in particular the meanings specified as being preferred, and q is 0, 1 or 2 and in particular 0 or 1. If q is 1, the radical R^a is preferably bound to the carbon atom which is adjacent to the nitrogen atom of the pyridine ring

[0061] Among the compounds of general formula Ia, preference is furthermore given to the compounds of general formula Ia.2








in which n, X, Y, R¹, R², R³, R^a and R^b have the previously mentioned meanings, in particular the meanings specified as being preferred, and q and q' are independently of each other 0 or 1, with q+q' preferably being 0 or 1.

[0062] Examples of compounds of the formula Ia.1 are the compounds of the following general formulae Ia.1a, Ia.1b, Ia.1c, Ia.1d, Ia.1e, Ia.1f, Ia.1g, Ia.1h and Ia.1k:




-continued




in which R¹, R^{2a}, R^{2b}, R^{2c}, R³, X, Y and R^b, have the meanings specified in one line in Table 1.




[0063] Examples of compounds of the formula Ia.2 are the compounds of the following general formulae Ia.2a, Ia.2b, Ia.2c, Ia.2d and Ia.2e:



-continued



-continued



[0064] in which R¹, R^a, R^{2b}, R^{2c}, R³, X, Y and R^b have the meanings specified in one line in Table 1.

TABLE 1

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
1.	H	H	H	H	H	CH	CH	CH(CH ₃) ₂
2.	CH ₃	H	H	H	H	CH	CH	CH(CH ₃) ₂
3.	CH ₂ CH ₃	H	H	H	H	CH	CH	CH(CH ₃) ₂
4.	CH ₂ CH=CH ₂	H	H	H	H	CH	CH	CH(CH ₃) ₂
5.	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	CH	CH(CH ₃) ₂
6.	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	CH	CH(CH ₃) ₂
7.	H	(s)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
8.	CH ₃	(s)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
9.	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
10.	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
11.	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
12.	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
13.	CH ₃	rac-CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
14.	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
15.	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
16.	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
17.	CH ₃	(R)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
18.	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
19.	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
20.	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	CH	CH(CH ₃) ₂
21.	CH ₃	H	CH ₃	H	H	CH	CH	CH(CH ₃) ₂
22.	CH ₂ CH=CH ₂	H	CH ₃	H	H	CH	CH	CH(CH ₃) ₂
23.	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	H	CH	CH	CH(CH ₃) ₂
24.	CH ₂ CH ₂ CH ₃	H	CH ₃	H	H	CH	CH	CH(CH ₃) ₂
25.	CH ₃	CH ₃	H	CH ₃	H	CH	CH	CH(CH ₃) ₂
26.	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	H	CH	CH	CH(CH ₃) ₂
27.	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	H	CH	CH	CH(CH ₃) ₂
28.	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	H	CH	CH	CH(CH ₃) ₂
29.	CH ₃	CH ₃	CH ₃	H	H	CH	CH	CH(CH ₃) ₂
30.	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	H	CH	CH	CH(CH ₃) ₂
31.	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	H	CH	CH	CH(CH ₃) ₂
32.	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	H	CH	CH	CH(CH ₃) ₂
33.	(s)(CH ₂) ₃	H	H	H	H	CH	CH	CH(CH ₃) ₂
34.	(s)(CH ₂) ₄	H	H	H	H	CH	CH	CH(CH ₃) ₂
35.	rac(CH ₂) ₃	H	H	H	H	CH	CH	CH(CH ₃) ₂
36.	rac(CH ₂) ₄	H	H	H	H	CH	CH	CH(CH ₃) ₂
37.	(R)(CH ₂) ₃	H	H	H	H	CH	CH	CH(CH ₃) ₂
38.	(R)(CH ₂) ₄	H	H	H	H	CH	CH	CH(CH ₃) ₂
39.	CH ₂ CH=CH ₂	H	H	H	H	C—Cl	CH	CH(CH ₃) ₂
40.	CH ₂ -c-C ₃ H ₅	H	H	H	H	C—Cl	CH	CH(CH ₃) ₂
41.	CH ₂ CH ₂ CH ₃	H	H	H	H	C—Cl	CH	CH(CH ₃) ₂
42.	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
43.	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
44.	CH ₂ CH ₃	(s)CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
45.	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
46.	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
47.	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
48.	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
49.	CH ₂ CH ₃	rac-CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
50.	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
51.	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
52.	CH ₂ CH ₃	(R)CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
53.	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	C—Cl	CH	CH(CH ₃) ₂
54.	CH ₂ CH=CH ₂	H	CH ₃	H	H	C—Cl	CH	CH(CH ₃) ₂
55.	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	H	C—Cl	CH	CH(CH ₃) ₂
56.	CH ₂ CH ₂ CH ₃	H	CH ₃	H	H	C—Cl	CH	CH(CH ₃) ₂
57.	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	H	C—Cl	CH	CH(CH ₃) ₂
58.	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	H	C—Cl	CH	CH(CH ₃) ₂
59.	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	H	C—Cl	CH	CH(CH ₃) ₂
60.	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	H	C—Cl	CH	CH(CH ₃) ₂
61.	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	H	C—Cl	CH	CH(CH ₃) ₂
62.	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	H	C—Cl	CH	CH(CH ₃) ₂
63.	(CH ₂) ₃	H	H	H	H	C—Cl	CH	CH(CH ₃) ₂
64.	(CH ₂) ₄	H	H	H	H	C—Cl	CH	CH(CH ₃) ₂
65.	CH ₂ CH=CH ₂	H	H	H	H	CH	C—Cl	CH(CH ₃) ₂
66.	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	C—Cl	CH(CH ₃) ₂
67.	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	C—Cl	CH(CH ₃) ₂
68.	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
69.	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
70.	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
71.	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
72.	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
73.	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
74.	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
75.	CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
76.	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
77.	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
78.	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
79.	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	C—Cl	CH(CH ₃) ₂
80.	CH ₂ CH=CH ₂	H	CH ₃	H	H	CH	C—Cl	CH(CH ₃) ₂
81.	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	H	CH	C—Cl	CH(CH ₃) ₂
82.	CH ₂ CH ₂ CH ₃	H	CH ₃	H	H	CH	C—Cl	CH(CH ₃) ₂
83.	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	H	CH	C—Cl	CH(CH ₃) ₂
84.	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	H	CH	C—Cl	CH(CH ₃) ₂
85.	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	H	CH	C—Cl	CH(CH ₃) ₂
86.	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	H	CH	C—Cl	CH(CH ₃) ₂
87.	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	H	CH	C—Cl	CH(CH ₃) ₂
88.	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	H	CH	C—Cl	CH(CH ₃) ₂
89.	(CH ₂) ₃	H	H	H	H	CH	C—Cl	CH(CH ₃) ₂
90.	(CH ₂) ₄	H	H	H	H	CH	C—Cl	CH(CH ₃) ₂
91.	CH ₂ CH=CH ₂	H	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
92.	CH ₂ -c-C ₃ H ₅	H	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
93.	CH ₂ CH ₂ CH ₃	H	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
94.	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
95.	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
96.	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
97.	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
98.	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
99.	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
100.	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
101.	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
102.	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
103.	CH ₂ CH=CH ₂	H	CH ₃	H	H	C—CH ₃	CH	CH(CH ₃) ₂
104.	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	H	C—CH ₃	CH	CH(CH ₃) ₂
105.	CH ₂ CH ₂ CH ₃	H	CH ₃	H	H	C—CH ₃	CH	CH(CH ₃) ₂
106.	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	H	C—CH ₃	CH	CH(CH ₃) ₂
107.	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	H	C—CH ₃	CH	CH(CH ₃) ₂
108.	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	H	C—CH ₃	CH	CH(CH ₃) ₂
109.	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	H	C—CH ₃	CH	CH(CH ₃) ₂
110.	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	H	C—CH ₃	CH	CH(CH ₃) ₂
111.	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	H	C—CH ₃	CH	CH(CH ₃) ₂

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
112	(CH ₂) ₃	H	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
113	(CH ₂) ₄	H	H	H	H	C—CH ₃	CH	CH(CH ₃) ₂
114	CH ₂ CH=CH ₂	H	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
115	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
116	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
117	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
118	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
119	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
120	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
121	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
122	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
123	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
124	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
125	CH ₂ CH ₂ CH ₃	(T)CH ₃	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
126	CH ₂ CH=CH ₂	H	CH ₃	H	H	CH	C—CH ₃	CH(CH ₃) ₂
127	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	H	CH	C—CH ₃	CH(CH ₃) ₂
128	CH ₂ CH ₂ CH ₃	H	CH ₃	H	H	CH	C—CH ₃	CH(CH ₃) ₂
129	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	H	CH	C—CH ₃	CH(CH ₃) ₂
130	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	H	CH	C—CH ₃	CH(CH ₃) ₂
131	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	H	CH	C—CH ₃	CH(CH ₃) ₂
132	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	H	CH	C—CH ₃	CH(CH ₃) ₂
133	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	H	CH	C—CH ₃	CH(CH ₃) ₂
134	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	H	CH	C—CH ₃	CH(CH ₃) ₂
135	(CH ₂) ₃	H	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
136	(CH ₂) ₄	H	H	H	H	CH	C—CH ₃	CH(CH ₃) ₂
137	H	H	H	H	H	CH	CH	c-C ₃ H ₅
138	CH ₃	H	H	H	H	CH	CH	c-C ₃ H ₅
139	CH ₂ CH ₃	H	H	H	H	CH	CH	c-C ₃ H ₅
140	CH ₂ CH=CH ₂	H	H	H	H	CH	CH	c-C ₃ H ₅
141	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	CH	c-C ₃ H ₅
142	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	CH	c-C ₃ H ₅
143	H	(s)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
144	CH ₃	(s)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
145	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
146	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
147	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
148	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
149	CH ₃	rac-CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
150	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
151	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
152	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
153	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
154	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
155	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
156	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	CH	c-C ₃ H ₅
157	CH ₃	H	CH ₃	H	H	CH	CH	c-C ₃ H ₅
158	CH ₂ CH=CH ₂	H	CH ₃	H	H	CH	CH	c-C ₃ H ₅
159	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	H	CH	CH	c-C ₃ H ₅
160	CH ₂ CH ₂ CH ₃	H	CH ₃	H	H	CH	CH	c-C ₃ H ₅
161	CH ₃	CH ₃	H	CH ₃	H	CH	CH	c-C ₃ H ₅
162	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	H	CH	CH	c-C ₃ H ₅
163	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	H	CH	CH	c-C ₃ H ₅
164	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	H	CH	CH	c-C ₃ H ₅
165	CH ₃	CH ₃	CH ₃	H	H	CH	CH	c-C ₃ H ₅
166	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	H	CH	CH	c-C ₃ H ₅
167	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	H	CH	CH	c-C ₃ H ₅
168	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	H	CH	CH	c-C ₃ H ₅
169	(s)(CH ₂) ₃	H	H	H	H	CH	CH	c-C ₃ H ₅
170	(s)(CH ₂) ₄	H	H	H	H	CH	CH	c-C ₃ H ₅
171	rac(CH ₂) ₃	H	H	H	H	CH	CH	c-C ₃ H ₅
172	rac(CH ₂) ₄	H	H	H	H	CH	CH	c-C ₃ H ₅
173	(R)(CH ₂) ₃	H	H	H	H	CH	CH	c-C ₃ H ₅
174	(R)(CH ₂) ₄	H	H	H	H	CH	CH	c-C ₃ H ₅
175	CH ₂ CH=CH ₂	H	H	H	H	CH	C—Cl	c-C ₃ H ₅
176	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	C—Cl	c-C ₃ H ₅
177	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	C—Cl	c-C ₃ H ₅
178	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
179	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
180	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
181	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
182	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
183	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
184	CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
185	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
186	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
187	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
188	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
189	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	C—Cl	c-C ₃ H ₅
190	CH ₂ CH=CH ₂	H	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
191	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
192	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
193	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
194	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
195	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
196	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
197	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
198	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
199	CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
200	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
201	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
202	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
203	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
204	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	C—CH ₃	c-C ₃ H ₅
205	CH ₂ CH=CH ₂	H	H	H	H	C—Cl	CH	c-C ₃ H ₅
206	CH ₂ -c-C ₃ H ₅	H	H	H	H	C—Cl	CH	c-C ₃ H ₅
207	CH ₂ CH ₂ CH ₃	H	H	H	H	C—Cl	CH	c-C ₃ H ₅
208	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	C—Cl	CH	c-C ₃ H ₅
209	CH ₂ -c-C ₃ H ₅	(S)CH ₃	H	H	H	C—Cl	CH	c-C ₃ H ₅
210	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	C—Cl	CH	c-C ₃ H ₅
211	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	C—Cl	CH	c-C ₃ H ₅
212	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	C—Cl	CH	c-C ₃ H ₅
213	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	C—Cl	CH	c-C ₃ H ₅
214	CH ₂ CH=CH ₂	H	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
215	CH ₂ -c-C ₃ H ₅	H	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
216	CH ₂ CH ₂ CH ₃	H	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
217	CH ₂ CH=CH ₂	(S)CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
218	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
219	CH ₂ CH ₃	(s)CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
220	CH ₂ CH ₂ CH ₃	(S)CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
221	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
222	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
223	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
224	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
225	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
226	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	C—CH ₃	CH	c-C ₃ H ₅
227	H	H	H	H	H	CH	CH	C ₂ H ₅
228	CH ₃	H	H	H	H	CH	CH	C ₂ H ₅
229	CH ₂ CH ₃	H	H	H	H	CH	CH	C ₂ H ₅
230	CH ₂ CH=CH ₂	H	H	H	H	CH	CH	C ₂ H ₅
231	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	CH	C ₂ H ₅
232	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	CH	C ₂ H ₅
233	H	(s)CH ₃	H	H	H	CH	CH	C ₂ H ₅
234	CH ₃	(s)CH ₃	H	H	H	CH	CH	C ₂ H ₅
235	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	CH	C ₂ H ₅
236	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	CH	C ₂ H ₅
237	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	CH	C ₂ H ₅
238	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	CH	C ₂ H ₅
239	CH ₃	rac-CH ₃	H	H	H	CH	CH	C ₂ H ₅
240	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	CH	C ₂ H ₅
241	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	H	CH	CH	C ₂ H ₅
242	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	CH	C ₂ H ₅
243	CH ₃	(R)CH ₃	H	H	H	CH	CH	C ₂ H ₅
244	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	CH	C ₂ H ₅
245	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	CH	C ₂ H ₅
246	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	CH	C ₂ H ₅
247	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	CH	C ₂ H ₅
248	CH ₃	H	CH ₃	H	H	CH	CH	C ₂ H ₅
249	CH ₂ CH=CH ₂	H	CH ₃	H	H	CH	CH	C ₂ H ₅
250	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	H	CH	CH	C ₂ H ₅
251	CH ₂ CH ₂ CH ₃	H	CH ₃	H	H	CH	CH	C ₂ H ₅
252	CH ₃	CH ₃	H	CH ₃	H	CH	CH	C ₂ H ₅
253	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	H	CH	CH	C ₂ H ₅
254	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	H	CH	CH	C ₂ H ₅
255	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	H	CH	CH	C ₂ H ₅
256	CH ₃	CH ₃	CH ₃	H	H	CH	CH	C ₂ H ₅
257	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	H	CH	CH	C ₂ H ₅
258	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	H	CH	CH	C ₂ H ₅
259	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	H	CH	CH	C ₂ H ₅

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
260	(s)(CH ₂) ₃	H	H	H	H	CH	CH	C ₂ H ₅
261	(s)(CH ₂) ₄	H	H	H	H	CH	CH	C ₂ H ₅
262	rac(CH ₂) ₃	H	H	H	H	CH	CH	C ₂ H ₅
263	rac(CH ₂) ₄	H	H	H	H	CH	CH	C ₂ H ₅
264	(R)(CH ₂) ₃	H	H	H	H	CH	CH	C ₂ H ₅
265	(R)(CH ₂) ₄	H	H	H	H	CH	CH	C ₂ H ₅
266	H	H	H	H	H	CH	CH	CH ₃
267	CH ₃	H	H	H	H	CH	CH	CH ₃
268	CH ₂ CH ₃	H	H	H	H	CH	CH	CH ₃
269	CH ₂ CH=CH ₂	H	H	H	H	CH	CH	CH ₃
270	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	CH	CH ₃
271	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	CH	CH ₃
272	H	(s)CH ₃	H	H	H	CH	CH	CH ₃
273	CH ₃	(s)CH ₃	H	H	H	CH	CH	CH ₃
274	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	CH	CH ₃
275	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	CH	CH ₃
276	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	CH	CH ₃
277	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	CH	CH ₃
278	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	CH	CH ₃
279	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	CH	CH ₃
280	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	CH	CH ₃
281	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	CH	CH ₃
282	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	CH	CH ₃
283	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	CH	CH ₃
284	CH ₂ CH=CH ₂	H	CH ₃	H	H	CH	CH	CH ₃
285	CH ₂ CH ₂ CH ₃	H	CH ₃	H	H	CH	CH	CH ₃
286	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	H	CH	CH	CH ₃
287	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	H	CH	CH	CH ₃
288	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	H	CH	CH	CH ₃
289	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	H	CH	CH	CH ₃
290	(s)(CH ₂) ₃	H	H	H	H	CH	CH	CH ₃
291	(s)(CH ₂) ₄	H	H	H	H	CH	CH	CH ₃
292	rac(CH ₂) ₃	H	H	H	H	CH	CH	CH ₃
293	rac(CH ₂) ₄	H	H	H	H	CH	CH	CH ₃
294	(R)(CH ₂) ₃	H	H	H	H	CH	CH	CH ₃
295	(R)(CH ₂) ₄	H	H	H	H	CH	CH	CH ₃
296	H	H	H	H	H	N	CH	CH(CH ₃) ₂
297	CH ₃	H	H	H	H	N	CH	CH(CH ₃) ₂
298	CH ₂ CH ₃	H	H	H	H	N	CH	CH(CH ₃) ₂
299	CH ₂ CH=CH ₂	H	H	H	H	N	CH	CH(CH ₃) ₂
300	CH ₂ -c-C ₃ H ₅	H	H	H	H	N	CH	CH(CH ₃) ₂
301	CH ₂ CH ₂ CH ₃	H	H	H	H	N	CH	CH(CH ₃) ₂
302	H	(s)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
303	CH ₃	(s)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
304	CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
305	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
306	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
307	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
308	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
309	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
310	CH ₂ CH ₃	(R)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
311	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
312	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
313	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	N	CH	CH(CH ₃) ₂
314	(s)(CH ₂) ₃	H	H	H	H	N	CH	CH(CH ₃) ₂
315	(s)(CH ₂) ₄	H	H	H	H	N	CH	CH(CH ₃) ₂
316	rac(CH ₂) ₃	H	H	H	H	N	CH	CH(CH ₃) ₂
317	rac(CH ₂) ₄	H	H	H	H	N	CH	CH(CH ₃) ₂
318	(R)(CH ₂) ₃	H	H	H	H	N	CH	CH(CH ₃) ₂
319	(R)(CH ₂) ₄	H	H	H	H	N	CH	CH(CH ₃) ₂
320	H	H	H	H	H	N	CH	CH=CH ₂
321	CH ₃	H	H	H	H	N	CH	CH=CH ₂
322	CH ₂ CH ₃	H	H	H	H	N	CH	CH=CH ₂
323	CH ₂ CH=CH ₂	H	H	H	H	N	CH	CH=CH ₂
324	CH ₂ -c-C ₃ H ₅	H	H	H	H	N	CH	CH=CH ₂
325	CH ₂ CH ₂ CH ₃	H	H	H	H	N	CH	CH=CH ₂
326	H	(s)CH ₃	H	H	H	N	CH	CH=CH ₂
327	CH ₃	(s)CH ₃	H	H	H	N	CH	CH=CH ₂
328	CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	CH=CH ₂
329	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	N	CH	CH=CH ₂
330	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	N	CH	CH=CH ₂
331	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	CH=CH ₂
332	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	N	CH	CH=CH ₂
333	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	N	CH	CH=CH ₂

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
334	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	N	CH	CH=CH ₂
335	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	N	CH	CH=CH ₂
336	H	H	H	H	H	N	CH	c-C ₃ H ₅
337	CH ₃	H	H	H	H	N	CH	c-C ₃ H ₅
338	CH ₂ CH ₃	H	H	H	H	N	CH	c-C ₃ H ₅
339	CH ₂ CH=CH ₂	H	H	H	H	N	CH	c-C ₃ H ₅
340	CH ₂ -c-C ₃ H ₅	H	H	H	H	N	CH	c-C ₃ H ₅
341	CH ₂ CH ₂ CH ₃	H	H	H	H	N	CH	c-C ₃ H ₅
342	H	(s)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
343	CH ₃	(s)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
344	CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
345	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
346	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
347	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
348	CH ₂ CH ₃	(R)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
349	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
350	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	N	CH	c-C ₃ H ₅
351	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	N	CH	c-C ₃ H ₅
352	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	N	CH	c-C ₃ H ₅
353	H	H	H	H	H	N	CH	CH ₃
354	CH ₃	H	H	H	H	N	CH	CH ₃
355	CH ₂ CH ₃	H	H	H	H	N	CH	CH ₃
356	CH ₂ CH=CH ₂	H	H	H	H	N	CH	CH ₃
357	CH ₂ -c-C ₃ H ₅	H	H	H	H	N	CH	CH ₃
358	CH ₂ CH ₂ CH ₃	H	H	H	H	N	CH	CH ₃
359	H	(s)CH ₃	H	H	H	N	CH	CH ₃
360	CH ₃	(s)CH ₃	H	H	H	N	CH	CH ₃
361	CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	CH ₃
362	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	N	CH	CH ₃
363	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	N	CH	CH ₃
364	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	CH ₃
365	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	N	CH	CH ₃
366	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	N	CH	CH ₃
367	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	N	CH	CH ₃
368	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	N	CH	CH ₃
369	H	H	H	H	H	N	CH	CF ₃
370	CH ₃	H	H	H	H	N	CH	CF ₃
371	CH ₂ CH ₃	H	H	H	H	N	CH	CF ₃
372	CH ₂ CH=CH ₂	H	H	H	H	N	CH	CF ₃
373	CH ₂ -c-C ₃ H ₅	H	H	H	H	N	CH	CF ₃
374	CH ₂ CH ₂ CH ₃	H	H	H	H	N	CH	CF ₃
375	H	(s)CH ₃	H	H	H	N	CH	CF ₃
376	CH ₃	(s)CH ₃	H	H	H	N	CH	CF ₃
377	CH ₂ CH ₃	H	H	H	N	CH	CF ₃	
378	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	N	CH	CF ₃
379	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	N	CH	CF ₃
380	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	N	CH	CF ₃
381	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	N	CH	CF ₃
382	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	N	CH	CF ₃
383	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	N	CH	CF ₃
384	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	N	CH	CF ₃
385	H	H	H	H	H	CH	N	CH(CH ₃) ₂
386	CH ₃	H	H	H	H	CH	N	CH(CH ₃) ₂
387	CH ₂ CH ₃	H	H	H	H	CH	N	CH(CH ₃) ₂
388	CH ₂ CH=CH ₂	H	H	H	H	CH	N	CH(CH ₃) ₂
389	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	N	CH(CH ₃) ₂
390	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	N	CH(CH ₃) ₂
391	H	(s)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
392	CH ₃	(s)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
393	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
394	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
395	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
396	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
397	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
398	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
399	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
400	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
401	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
402	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	CH(CH ₃) ₂
403	H	H	H	H	H	CH	N	CH=CH ₂
404	CH ₃	H	H	H	H	CH	N	CH=CH ₂
405	CH ₂ CH ₃	H	H	H	H	CH	N	CH=CH ₂
406	CH ₂ CH=CH ₂	H	H	H	H	CH	N	CH=CH ₂
407	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	N	CH=CH ₂

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
408	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	N	CH=CH ₂
409	H	(s)CH ₃	H	H	H	CH	N	CH=CH ₂
410	CH ₃	(s)CH ₃	H	H	H	CH	N	CH=CH ₂
411	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	CH=CH ₂
412	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	N	CH=CH ₂
413	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	N	CH=CH ₂
414	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	CH=CH ₂
415	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	N	CH=CH ₂
416	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	N	CH=CH ₂
417	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	CH=CH ₂
418	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	N	CH=CH ₂
419	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	N	CH=CH ₂
420	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	CH=CH ₂
421	H	H	H	H	H	CH	N	c-C ₃ H ₅
422	CH ₃	H	H	H	H	CH	N	c-C ₃ H ₅
423	CH ₂ CH ₃	H	H	H	H	CH	N	c-C ₃ H ₅
424	CH ₂ CH=CH ₂	H	H	H	H	CH	N	c-C ₃ H ₅
425	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	N	c-C ₃ H ₅
426	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	N	c-C ₃ H ₅
427	H	(s)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
428	CH ₃	(s)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
429	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
430	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
431	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
432	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
433	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	N	c-C ₃ H ₅
434	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	N	c-C ₃ H ₅
435	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
436	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
437	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
438	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	c-C ₃ H ₅
439	H	H	H	H	H	CH	N	CH ₃
440	CH ₃	H	H	H	H	CH	N	CH ₃
441	CH ₂ CH ₃	H	H	H	H	CH	N	CH ₃
442	CH ₂ CH=CH ₂	H	H	H	H	CH	N	CH ₃
443	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	N	CH ₃
444	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	N	CH ₃
445	H	(s)CH ₃	H	H	H	CH	N	CH ₃
446	CH ₃	(s)CH ₃	H	H	H	CH	N	CH ₃
447	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	CH ₃
448	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	N	CH ₃
449	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	N	CH ₃
450	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	CH ₃
451	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	N	CH ₃
452	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	N	CH ₃
453	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	CH ₃
454	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	N	CH ₃
455	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	N	CH ₃
456	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	CH ₃
457	H	H	H	H	H	CH	N	CF ₃
458	CH ₃	H	H	H	H	CH	N	CF ₃
459	CH ₂ CH ₃	H	H	H	H	CH	N	CF ₃
460	CH ₂ CH=CH ₂	H	H	H	H	CH	N	CF ₃
461	CH ₂ -c-C ₃ H ₅	H	H	H	H	CH	N	CF ₃
462	CH ₂ CH ₂ CH ₃	H	H	H	H	CH	N	CF ₃
463	H	(s)CH ₃	H	H	H	CH	N	CF ₃
464	CH ₃	(s)CH ₃	H	H	H	CH	N	CF ₃
465	CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	CF ₃
466	CH ₂ CH=CH ₂	(s)CH ₃	H	H	H	CH	N	CF ₃
467	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	H	CH	N	CF ₃
468	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	H	CH	N	CF ₃
469	CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	CF ₃
470	CH ₂ CH=CH ₂	(R)CH ₃	H	H	H	CH	N	CF ₃
471	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	H	CH	N	CF ₃
472	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	H	CH	N	CF ₃
473	CH ₂ CH=CH ₂	rac-CH ₃	H	H	H	CH	N	CF ₃
474	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	H	CH	N	CF ₃
475	H	H	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
476	CH ₃	H	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
477	CH ₂ CH ₃	H	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
478	CH ₂ CH=CH ₂	H	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
479	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
480	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
481	H	(s)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
482	CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
483	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
484	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
485	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
486	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
487	CH ₃	rac-CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
488	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
489	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
490	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
491	CH ₃	(R)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
492	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
493	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
494	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	CH	CH	CH(CH ₃) ₂
495	CH ₃	H	CH ₃	H	CH ₃	CH	CH	CH(CH ₃) ₂
496	CH ₂ CH=CH ₂	H	CH ₃	H	CH ₃	CH	CH	CH(CH ₃) ₂
497	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	CH ₃	CH	CH	CH(CH ₃) ₂
498	CH ₂ CH ₂ CH ₃	H	CH ₃	H	CH ₃	CH	CH	CH(CH ₃) ₂
499	CH ₃	CH ₃	H	CH ₃	CH ₃	CH	CH	CH(CH ₃) ₂
500	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	CH ₃	CH	CH	CH(CH ₃) ₂
501	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	CH ₃	CH	CH	CH(CH ₃) ₂
502	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	CH ₃	CH	CH	CH(CH ₃) ₂
503	CH ₃	CH ₃	CH ₃	H	CH ₃	CH	CH	CH(CH ₃) ₂
504	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	CH ₃	CH	CH	CH(CH ₃) ₂
505	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	CH ₃	CH	CH	CH(CH ₃) ₂
506	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	CH ₃	CH	CH	CH(CH ₃) ₂
507	(S)(CH ₂) ₃	H	H	CH ₃	CH	CH	CH	CH(CH ₃) ₂
508	(S)(CH ₂) ₄	H	H	CH ₃	CH	CH	CH	CH(CH ₃) ₂
509	rac(CH ₂) ₃	H	H	CH ₃	CH	CH	CH	CH(CH ₃) ₂
510	rac(CH ₂) ₄	H	H	CH ₃	CH	CH	CH	CH(CH ₃) ₂
511	(R)(CH ₂) ₃	H	H	CH ₃	CH	CH	CH	CH(CH ₃) ₂
512	(R)(CH ₂) ₄	H	H	CH ₃	CH	CH	CH	CH(CH ₃) ₂
513	CH ₂ CH=CH ₂	H	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
514	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
515	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
516	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
517	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
518	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
519	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
520	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
521	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
522	CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
523	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
524	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
525	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
526	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
527	CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
528	CH ₂ CH=CH ₂	H	CH ₃	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
529	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
530	CH ₂ CH ₂ CH ₃	H	CH ₃	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
531	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	CH ₃	C—Cl	CH	CH(CH ₃) ₂
532	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	CH ₃	C—Cl	CH	CH(CH ₃) ₂
533	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	CH ₃	C—Cl	CH	CH(CH ₃) ₂
534	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
535	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
536	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
537	(CH ₂) ₃	H	H	CH ₃	C—Cl	CH	CH	CH(CH ₃) ₂
538	(CH ₂) ₄	H	H	CH ₃	C—Cl	CH	CH	CH(CH ₃) ₂
539	CH ₂ CH=CH ₂	H	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
540	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
541	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
542	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
543	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
544	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
545	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
546	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
547	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
548	CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
549	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
550	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
551	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
552	CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
553	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
554	CH ₂ CH=CH ₂	H	CH ₃	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂
555	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	CH ₃	C—Cl	CH	CH(CH ₃) ₂

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
556	CH ₂ CH ₂ CH ₃	H	CH ₃	H	CH ₃	CH	C—Cl	CH(CH ₃) ₂
557	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	CH ₃	CH	C—Cl	CH(CH ₃) ₂
558	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	CH ₃	CH	C—Cl	CH(CH ₃) ₂
559	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	CH ₃	CH	C—Cl	CH(CH ₃) ₂
560	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	CH ₃	CH	C—Cl	CH(CH ₃) ₂
561	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	CH ₃	CH	C—Cl	CH(CH ₃) ₂
562	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	CH ₃	CH	C—Cl	CH(CH ₃) ₂
563	(CH ₂) ₃	H	H	H	CH ₃	CH	C—Cl	CH(CH ₃) ₂
564	(CH ₂) ₄	H	H	H	CH ₃	CH	C—Cl	CH(CH ₃) ₂
565	CH ₂ CH=CH ₂	H	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
566	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
567	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
568	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
569	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
570	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
571	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
572	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
573	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
574	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
575	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
576	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
577	CH ₂ CH=CH ₂	H	CH ₃	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
578	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
579	CH ₂ CH ₂ CH ₃	H	CH ₃	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
580	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
581	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
582	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
583	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
584	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
585	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
586	(CH ₂) ₃	H	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
587	(CH ₂) ₄	H	H	H	CH ₃	C—CH ₃	CH	CH(CH ₃) ₂
588	CH ₂ CH=CH ₂	H	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
589	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
590	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
591	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
592	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
593	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
594	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
595	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
596	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
597	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
598	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
599	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
600	CH ₂ CH=CH ₂	H	CH ₃	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
601	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
602	CH ₂ CH ₂ CH ₃	H	CH ₃	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
603	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
604	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
605	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
606	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
607	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
608	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
609	(CH ₂) ₃	H	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
610	(CH ₂) ₄	H	H	H	CH ₃	CH	C—CH ₃	CH(CH ₃) ₂
611	H	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
612	CH ₃	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
613	CH ₂ CH ₃	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
614	CH ₂ CH=CH ₂	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
615	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
616	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
617	H	(s)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
618	CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
619	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
620	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
621	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
622	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
623	CH ₃	rac-CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
624	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
625	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
626	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
627	CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
628	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
629	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅

TABLE 1-continued

No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
630	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	CH	CH	c-C ₃ H ₅
631	CH ₃	H	CH ₃	H	CH ₃	CH	CH	c-C ₃ H ₅
632	CH ₂ CH=CH ₂	H	CH ₃	H	CH ₃	CH	CH	c-C ₃ H ₅
633	CH ₂ -c-C ₃ H ₅	H	CH ₃	H	CH ₃	CH	CH	c-C ₃ H ₅
634	CH ₂ CH ₂ CH ₃	H	CH ₃	H	CH ₃	CH	CH	c-C ₃ H ₅
635	CH ₃	CH ₃	H	CH ₃	CH ₃	CH	CH	c-C ₃ H ₅
636	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	CH ₃	CH	CH	c-C ₃ H ₅
637	CH ₂ -c-C ₃ H ₅	CH ₃	H	CH ₃	CH ₃	CH	CH	c-C ₃ H ₅
638	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	CH ₃	CH	CH	c-C ₃ H ₅
639	CH ₃	CH ₃	CH ₃	H	CH ₃	CH	CH	c-C ₃ H ₅
640	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	CH ₃	CH	CH	c-C ₃ H ₅
641	CH ₂ -c-C ₃ H ₅	CH ₃	CH ₃	H	CH ₃	CH	CH	c-C ₃ H ₅
642	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	CH ₃	CH	CH	c-C ₃ H ₅
643	(s)(CH ₂) ₃	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
644	(s)(CH ₂) ₄	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
645	rac(CH ₂) ₃	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
646	rac(CH ₂) ₄	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
647	(R)(CH ₂) ₃	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
648	(R)(CH ₂) ₄	H	H	H	CH ₃	CH	CH	c-C ₃ H ₅
649	CH ₂ CH=CH ₂	H	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
650	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
651	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
652	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
653	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
654	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
655	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
656	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
657	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
658	CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
659	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
660	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
661	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
662	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
663	CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	CH	C—Cl	c-C ₃ H ₅
664	CH ₂ CH=CH ₂	H	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
665	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
666	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
667	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
668	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
669	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
670	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
671	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
672	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
673	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
674	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
675	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	CH	C—CH ₃	c-C ₃ H ₅
676	CH ₂ CH=CH ₂	H	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
677	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
678	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
679	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
680	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
681	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
682	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
683	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
684	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
685	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
686	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
687	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
688	CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
689	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	C—Cl	CH	c-C ₃ H ₅
690	CH ₂ CH=CH ₂	H	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
691	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
692	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
693	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
694	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
695	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
696	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
697	CH ₂ -c-C ₃ H ₅	rac-CH ₃	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
698	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	C—CH ₃	CH	c-C ₃ H ₅
699	CH ₃	H	H	H	CH ₃	CH	CH	C ₂ H ₅
700	CH ₂ CH=CH ₂	H	H	H	CH ₃	CH	CH	C ₂ H ₅
701	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	CH	CH	C ₂ H ₅
702	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	CH	CH	C ₂ H ₅
703	CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅

TABLE 1-continued


No.	R ¹	R ^{2a}	R ^{2b}	R ^{2c}	R ³	X	Y	R ^b
704	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
705	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
706	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
707	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
708	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
709	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
710	CH ₃	(R)CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
711	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
712	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
713	CH ₂ CH=CH ₂	H	CH ₃	H	CH ₃	CH	CH	C ₂ H ₅
714	CH ₂ CH ₂ CH ₃	H	CH ₃	H	CH ₃	CH	CH	C ₂ H ₅
715	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	CH ₃	CH	CH	C ₂ H ₅
716	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	CH ₃	CH	CH	C ₂ H ₅
717	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	CH ₃	CH	CH	C ₂ H ₅
718	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	CH ₃	CH	CH	C ₂ H ₅
719		(s)(CH ₂) ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
720		(s)(CH ₂) ₄	H	H	CH ₃	CH	CH	C ₂ H ₅
721		rac(CH ₂) ₃	H	H	CH ₃	CH	CH	C ₂ H ₅
722		rac(CH ₂) ₄	H	H	CH ₃	CH	CH	C ₂ H ₅
723	H	H	H	H	CH ₃	CH	CH	CH=CH ₂
724	CH ₃	H	H	H	CH ₃	CH	CH	CH=CH ₂
725	CH ₂ CH ₃	H	H	H	CH ₃	CH	CH	CH=CH ₂
726	CH ₂ CH=CH ₂	H	H	H	CH ₃	CH	CH	CH=CH ₂
727	CH ₂ -c-C ₃ H ₅	H	H	H	CH ₃	CH	CH	CH=CH ₂
728	CH ₂ CH ₂ CH ₃	H	H	H	CH ₃	CH	CH	CH=CH ₂
729	H	(s)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
730	CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
731	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
732	CH ₂ CH=CH ₂	(s)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
733	CH ₂ -c-C ₃ H ₅	(s)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
734	CH ₂ CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
735	CH ₂ CH=CH ₂	rac-CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
736	CH ₂ CH ₂ CH ₃	rac-CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
737	CH ₂ CH ₃	(s)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
738	CH ₂ CH=CH ₂	(R)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
739	CH ₂ -c-C ₃ H ₅	(R)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
740	CH ₂ CH ₂ CH ₃	(R)CH ₃	H	H	CH ₃	CH	CH	CH=CH ₂
741	CH ₂ CH=CH ₂	H	CH ₃	H	CH ₃	CH	CH	CH=CH ₂
742	CH ₂ CH ₂ CH ₃	H	CH ₃	H	CH ₃	CH	CH	CH=CH ₂
743	CH ₂ CH=CH ₂	CH ₃	H	CH ₃	CH ₃	CH	CH	CH=CH ₂
744	CH ₂ CH ₂ CH ₃	CH ₃	H	CH ₃	CH ₃	CH	CH	CH=CH ₂
745	CH ₂ CH=CH ₂	CH ₃	CH ₃	H	CH ₃	CH	CH	CH=CH ₂
746	CH ₂ CH ₂ CH ₃	CH ₃	CH ₃	H	CH ₃	CH	CH	CH=CH ₂
747		(CH ₂) ₃	H	H	CH ₃	CH	CH	CH=CH ₂
748		(CH ₂) ₄	H	H	CH ₃	CH	CH	CH=CH ₂

rac: racemate;

(S): S configuration;


(R) R configuration.

[0065] Other examples of compounds according to the invention are the compounds of the general formulae Ia.3, Ib, Ic, Id, Ie and If:




-continued




(Ib)



(Ic)




-continued




in which R^1 , R^{2a} , R^{2b} , R^{2c} , R^3 , X , Y and R^b have the meanings specified in one line in Table 1.

[0066] Among the compounds of the general formula I, preference is also given to the compounds of the general formula Ig





in which n , k , R^1 , R^2 , R^a , R^b , A^1 , A^2 , A^3 , X and Y have the meanings given for formula Ia.

[0067] Among the compounds of the general formula I, preference is also given to the compounds of the general formula Ih






in which n , k , R^1 , R^2 , R^a , R^b , A^1 , A^2 , A^3 , X and Y have the meanings given for formula Ia.

[0068] Among the compounds of general formulae Ig and Ih, preference is given to the compounds of general formula Ig.1, Ig.2, Ih.1 and Ih.2









in which n , q , q' , X , Y , R^1 , R^2 , R^a and R^b have the meanings mentioned previously for formulae Ia.1 and Ia.2 and R is CH_2 (compounds Ig.1 and Ig.2) or O (compounds Ih.1 or Ih.2).

[0069] Examples of compounds of the formula Ig.1 and Ih.1 are the compounds of the following general formulae Ig.1a, Ig.1b, Ig.1c, Ig.1d, Ig.1e, Ig.1f, Ig.1g, Ig.1h, Ig.1k, Ih.1a, Ih.1b, Ih.1c, Ih.1h, Ih.1e, Ih.1f, Ih.1g, Ih.1h and Ih.1k:








-continued

(Ig.1d: R = CH₂)
(Ih.1d: R = O)(Ig.1e: R = CH₂)
(Ih.1e: R = O)(Ig.1f: R = CH₂)
(Ih.1f: R = O)(Ig.1g: R = CH₂)
(Ih.1g: R = O)(Ig.1h: R = CH₂)
(Ih.1h: R = O)(Ig.1k: R = CH₂)
(Ih.1k: R = O)

in which R¹, R^{2a}, R^{2b}, R^{2c}, X, Y and R^b have the meanings specified in one of the lines 1 to 474 in Table 1.

[0070] Examples of compounds of the formula Ig.1 and Ih.1 are the compounds of the following general formulae


Ig.2a, Ig.2b, Ig.2c, Ig.2d, Ig.2e, Ih.2a, Ih.2b, Ih.2c, Ih.2d and Ih.2e:

Ig.2a: R = CH₂
Ih.2a: R = OIg.2b: R = CH₂
Ih.2b: R = OIg.2c: R = CH₂
Ih.2c: R = OIg.2d: R = CH₂
Ih.2d: R = OIg.2e: R = CH₂
Ih.2e: R = O

in which R¹, R^{2a}, R^{2b}, R^{2c}, X, Y and R^b have the meanings specified in one of the lines 1 to 474 in Table 1.

[0071] The compounds I according to the invention are prepared in analogy with methods known from the literature. An important approach to the compounds according to the invention with R being O or N—R³ is offered by the reaction of a hetaryl compound If with an arylsulfonic acid derivative III as depicted in scheme 1.

Scheme 1:





[0072] In scheme 1, n , R^1 , R^2 , R^3 , Ar and Q have the previously mentioned meanings. R is O or $N-R^3$. X is a nucleophilically displaceable leaving group, in particular a halogen atom and, especially, chlorine or bromine. The reaction depicted in scheme 1 takes place under the reaction conditions which are customary for preparing arylsulfonamide compounds or arylsulfonic esters, respectively, and which are described, for example, in J. March, Advanced Organic Chemistry, 3rd edition, John Wiley & Sons, New York, 1985 p 444 and the literature cited therein, European J. Org. Chem. 2002 (13), pp. 2094-2108, Tetrahedron 2001, 57 (27) pp. 5885-5895, Bioorganic and Medicinal Chemistry Letters, 2000, 10(8), pp. 835-838 and Synthesis 2000 (1), pp. 103-108.

[0073] The reaction customarily takes place in an inert solvent, for example in an ether, such as diethyl ether, diisopropyl ether, methyl tert-butyl ether or tetrahydrofuran, a halohydrocarbon, such as dichloromethane, an aliphatic or cycloaliphatic hydrocarbon, such as pentane, hexane or cyclohexane, or an aromatic hydrocarbon, such as toluene, xylene, cumene and the like, or in a mixture of the above-mentioned solvents.

[0074] The reaction of II with III is customarily carried out in the presence of an auxiliary base. Suitable bases are inorganic bases, such as sodium carbonate or potassium carbonate, or sodium hydrogencarbonate or potassium hydrogencarbonate, and organic bases, for example trialkylamines, such as triethylamine, or pyridine compounds, such as pyridine, lutidine and the like. The latter compounds can at the same time serve as solvents. The auxiliary base is customarily employed in at least equimolar quantities, based on the amine compound II.

[0075] The compounds of the general formula II are known per se or, in case R is $N-R^3$, can be prepared in the manner shown in schemes 2.

Scheme 2:



[0076] In scheme 2, n , R^2 and Q have the previously mentioned meanings. R^1 has the meanings different from hydrogen which are specified for R^1 or is a suitable protecting group. Suitable protecting groups are disclosed, for example, in P. Kocienski, Protecting Groups, Thieme-Verlag, Stuttgart 2000, Chapter 6. Y is a nucleophilically displaceable leaving group, in particular a halogen atom, e.g. chlorine or bromine, or an alkylsulfonyl group, e.g. methylsulfonyl.

[0077] The reaction depicted in step a) in scheme 2 takes place under the reaction conditions which are customary for a nucleophilic substitution on an aromatic radical and which are described, for example, in Tetrahedron 1999, 55(33), pp. 10243-10252, J. Med. Chem. 1997, 40(22), pp. 3679-3686 and Synthetic Communications, 1993, 23(5), pp. 591-599. Where appropriate, it can be advantageous to convert a ring nitrogen atom in the Q group into its N-oxide (see, for example, Angew. Chem. Int. Ed. Engl., 2002 41(1), pp. 1937-1940, J. Med. Chem. 1985, 28(2), pp. 248-252 and Tetrahedron Lett. 2002 43(17) pp. 3121-3123). This approach has proved to be of value, in particular, for preparing compounds I in which Q is a pyridin-2,4-diyl group. In connection with the subsequent reduction of the nitro group in VI (step b), the N-oxide group is also reduced. For this, the reduction is carried out, for example, in the presence of indium salts.

[0078] If 5-bromonitropyridine is used as compound V in step a) in accordance with scheme 2, the coupling is also achieved under palladium catalysis in the presence of an auxiliary base, for example an alkali metal carbonate such as cesium carbonate. Particularly suitable palladium catalysts in this connection are palladium(0) compounds or palladium compounds which are able to form a palladium(0) compound under reaction conditions, e.g. palladium dichloride, tetrakis(triphenylphosphine)palladium(0), tris(dibenzylideneacetone)dipalladium(0), advantageously in combination with phosphine ligands, e.g. triarylphosphines, such as triphenylphosphine, trialkylphosphines, such as tributylphosphine, and cycloalkylphosphines, such as tricyclohexylphosphine, and, especially, using phosphine chelate ligands, such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl. The conditions which are required for reactions of this nature are described, for example, in Tetrahedron Lett. 2001, 42(22), p. 3681 and Tetrahedron Lett. 2002, 43(12), pp. 2171-2173.

[0079] In step b), the nitro group in VI is reduced to the NH_2 group in II. Subsequently, in step c), the NH_2 group can


be converted into a $\text{—NR}^3\text{H}$ group, in which R^3 has the meanings different from hydrogen which are specified for R^3 .

[0080] The reaction conditions which are required for step b) correspond to the customary conditions for reducing aromatic nitro groups which have been described extensively in the literature (see, for example, J. March, *Advanced Organic Chemistry*, 3rd ed., J. Wiley & Sons, New-York, 1985, p.1183 and the literature cited in this reference).

[0081] The reduction is achieved, for example, by reacting the nitro compound VII with a metal such as iron, zinc or tin under acidic reaction conditions, i.e. using nascent hydrogen, or using a complex hydride such as lithium aluminum hydride or sodium borohydride, preferably in the presence of transition metal compounds of nickel or cobalt such as $\text{NiCl}_2(\text{P}(\text{phenyl})_3)_2$, or CoCl_2 , (see Ono et al. Chem. Ind. (London), 1983 p.480), or using NaBH_4S_3 (see Lalancette et al. Can. J. Chem. 49, 1971, p. 2990), with it being possible to carry out these reductions, depending on the given reagent, in substance or in a solvent or diluent. Alternatively, the reduction of VI to II can be carried out with hydrogen in the presence of a transition metal catalyst, e.g. using hydrogen in the presence of catalysts based on platinum, palladium, nickel, ruthenium or rhodium. The catalysts can contain the transition metal in elemental form or in the form of a complex compound, of a salt or of an oxide of the transition metal, with it being possible, for the purpose of modifying the activity, to use customary coligands, e.g. organic phosphine compounds, such as triphenylphosphine, tricyclohexylphosphine or tri-*n*-butylphosphines or phosphites. The catalyst is customarily employed in quantities of from 0.001 to 1 mol per mol of compound VI, calculated as catalyst metal. In a preferred variant, the reduction is effected using tin(II) chloride in analogy with the methods described in Bioorganic and, Medicinal Chemistry Letters, 2002, 12(15), pp. 1917-1919 and J. Med. Chem. 2002, 45(21), pp. 4679-4688. The reaction of VI with tin(II) chloride is preferably carried out in an inert organic solvent, preferably an alcohol such as methanol, ethanol, isopropanol or butanol.

[0082] Reducing VI results in compounds II in which R³ is hydrogen. Customary methods can then be used to react these compounds with an alkylating agent R³-X, in which R³ is C₁—C₄-alkyl and X is a nucleophilically displaceable leaving group (e.g. halogen, such as chlorine, bromine or iodine), resulting in a compound II in which R³=alkyl (step c). The reaction conditions which are required for this are disclosed, for example, in WO 02/83652, Tetrahedron 2000, 56(38) pp. 7553-7560 and Synlett. 2000 (4), pp. 475-480.


[0083] The compound I can also be prepared by the route depicted in scheme 3:



[0084] In scheme 3, n, R, R¹, R², Ar and 0 have the previously mentioned meanings. Y is a nucleophilically displaceable leaving group, in particular a halogen atom, e.g. chlorine or bromine, or an alkylsulfonyl group, e.g. meth-

ylsulfonyl. The reaction of VII with VIII, as depicted in scheme 3, takes place under the reaction conditions specified for scheme 2, step a). Compounds of the general formulae VII and VIII are known or can be prepared in analogy with the methods known from the literature. Compounds of the formula V, wherein R is N—R³ or an oxygen atom can be prepared by amidation of the the corresponding aminocompound X-Q-NHR³ (or esterification of the corresponding hydroxycompound X-Q-OH) with a sulfonylhalide, especially with a sulfonylchloride of the formula Z—SO₂—Ar, wherein Z is halogen, especially chlorine or bromine, according to standard methods of organic chemistry (see e.g. J. March, *Advanced Organic Chemistry*, 3rd edition, John Wiley & Sons, New York, 1985 p 444 and the literature cited therein).

[0085] Compounds of general formula I, in which R¹ is an allyl group, can be converted into compounds possessing different R¹ substituents using the synthetic route shown in scheme 4.



[0086] In scheme 4, n, R, R², Ar and Q have the previously mentioned meanings. The elimination of the allyl group, as depicted in step a) in scheme 4, is achieved, for example, by reacting I [R¹=allyl] with an allyl trapping agent, such as mercaptobenzoic acid or 1,3-dimethylbarbituric acid, in the presence of catalytic quantities of palladium (0) compounds or palladium compounds which are able to form a palladium(0) compound under reaction conditions, e.g. palladium dichloride, tetra(is(triphenylphosphine)palladium(0) or tris-(dibenzylideneacetone)dipalladium(0), advantageously in combination with phosphine ligands, e.g. triarylphosphines, such as triphenylphosphine, trialkylphosphines, such as tributylphosphine, and cycloalkylphosphines, such as tricyclohexylphosphine, and especially with phosphine chelate


ligands, such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl or 1,4-bis(diphenylphosphino)butane, using methods known from the literature (with regard to eliminating N-allyl in the presence of mercaptobenzoic acid, see WO 94/24088; with regard to eliminating in the presence of 1,3-dimethylbarbituric acid, see J. Am. Chem. Soc. 2001, 123 (28), pp. 6801-6808 and J. Org. Chem 2002, 67(11) pp. 3718-3723). Alternatively, the elimination of N-allyl, as depicted in scheme 4 step a), can also be effected by reacting in the presence of rhodium compounds, such as tris(triphenylphosphine)chlororhodium(1), using methods known from the literature (see J. Chem. Soc., Perkin Transaction I: Organic and BioOrganic Chemistry 1999 (21) pp. 3089-3104 and Tetrahedron Asymmetry 1997, 8(20), pp. 3387-3391).

[0087] The resulting piperazine compound I [$R^1=H$] can then be reacted, in a known manner, in the sense of an alkylation, with a compound R^1-X . In this compound, R^1 is C_1-C_4 -alkyl, C_3-C_6 -Cycloalkyl, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy- C_1-C_4 -alkyl or C_3-C_6 -cycloalkyl- C_1-C_4 -alkyl and X is a nucleophilically displaceable leaving group, e.g. halogen, trifluoroacetate, alkylsulfonate, arylsulfonate, alkyl sulfate and the like. The reaction conditions which are required for the alkylation in step b) have been adequately disclosed, e.g. in Bioorganic and Medicinal Chemistry Lett. 2002, 12(7), pp. 2443-2446 and also 2002, 12(5), pp.1917-1919.

[0088] The conversion, as depicted in scheme 4, step b), of the piperazine compound I [$R^1=H$] obtained in step a) can also be achieved, in the sense of a reductive amination, by reacting I [$R^1=H$] with a suitable ketone or aldehyde in the presence of a reducing agent, e.g. in the presence of a borohydride such as sodium borohydride, sodium cyanoborohydride or sodium triacetoxyborohydride. The skilled person is familiar with the reaction conditions which are required for a reductive amination, e.g. from Bioorganic and Medicinal Chemistry Lett. 2002, 12(5), pp. 795-798 and 12(7) pp. 1269-1273.

[0089] The conversion, as depicted in scheme 4, step b), of the piperazine compound I [$R^1=H$] obtained in step a) can also be achieved by successive acylation and subsequent reduction of the acylation product, using the method depicted in scheme 4a:

Scheme 4a:




R^x = alkyl, cycloalkyl, cycloalkylalkyl, alkenyl haloalkyl, alkynyl alkoxyalkyl, hydroxyalkyl

[0090] In scheme 4a, n, R, R^2 , Ar and Q have the previously mentioned meanings. The acylation in step a) and the reduction in step b) are effected using standard methods of organic chemistry as are described, for example, in J. March, Advanced Organic Chemistry, 3rd ed. J. Wiley & Sons, New York 1985, p. 370 and 373 (acylation) and p. 1099 f. and in the literature cited in this publication (with regard to acylation, see also Synth. Commun. 1986, 16, p. 267, and with regard to reduction, see also J. Heterocycl. Chem. 1979, 16, p. 1525).

[0091] In compounds of the general formula I which carry a halogen atom, in particular bromine or iodine, on the aromatic radical Ar, the halogen atom can be converted into an alkyl, alkenyl, cycloalkyl, alkynyl or cycloalkylalkyl group using methods which are known per se. The conversion is achieved by coupling the halo compound I to an alkyl-, alkenyl-, alkynyl-, cycloalkyl- or cycloalkylalkyl-boronic acid compound under the conditions of a Suzuki coupling as is described, for example, in Tetrahedron Lett. 2002, 43, pp. 6987-6990; Chem. Rev. 1995, 95, pp. 2457-2483 and J. Org. Chem. 66(21) (2001), pp. 7124-7128.


[0092] Compounds of the formula I, wherein R is CR^aR^b , can be also prepared by the synthetic route illustrated in scheme 5:



[0093] In scheme 5, n, R^1 , R^2 , Q, R^{3a} , R^{3b} and Ar are as defined above. L is a nucleophilically displaceable leaving group, in particular a halogen atom such as chlorine or bromine or a sulfonate group, e.g. a C_1-C_4 -alkylsulfonate such as methanesulfonate or an arylsulfonate such as benzenesulfonate or tosylate (toluene sulfonate). L may also be OH, when R^{3a} , R^{3b} are different from hydrogen. The reaction depicted in the first step of scheme 5 takes place under the reaction conditions which are customary for preparing arylsulfide compounds which are described, for example, in J. March, Advanced Organic Chemistry, 3rd edition, John Wiley & Sons, New York, 1985 p 360 to 362, and in the literature cited therein.

[0094] In the second step of scheme 5 the sulfide X obtained in the first step is oxidized to the corresponding sulfone I ($R=CR^aR^b$) under reaction conditions which are customary for preparing arylsulfone compounds from the corresponding arylsulfides and which are described, for example, in J. March, Advanced Organic Chemistry, 3rd edition, John Wiley & Sons, New York, 1985 p 1089 f. and in the literature cited therein.

[0095] The compounds of the formula IX are known in the art or can be prepared according to the method depicted in scheme 6:



[0096] In scheme 6, n , R^1 , R^2 , Q , R^{3a} , R^{3b} and Ar are as defined above. Hal is halogen, especially chlorine, bromine or iodine. In step (a) of scheme 6 the halogen compound XI is first metallized, i.e. converted into the corresponding lithium compound or magnesium compound by standard methods of organic chemistry and the thus obtained metal compound is reacted with an aldehyde or ketone to obtain the alcohol XII (see e.g. Tetrahedron 2003, 59 (24), 4303-4308). In step (b) the OH group of XII is then converted into a suitable leaving group, e.g. into a halogen atom by reaction with thionylhalide such as SOCl_2 or phosphorylhalide such as POCl_3 , or into an alkylsulfonate or arylsulfonate by reaction with an alkylsulfonylhalide such as methanesulfonylchloride or with an arylsulfonylhalide such as toluenesulfonylchloride (see e.g. J. Med. Chem. 1985, 28 (12), 1790-1796).

[0097] Alternatively the halogen compound XI is converted into the carbonyl compound XIII (step c, see e.g. J. Med. Chem. 1987, 30 (8), 1494-1497), which is then reduced ($R^{3b}=\text{H}$) or reacted with an alkylgrignard $R^{3b}\text{-Mg-X}$ ($\text{X}=\text{halogen}$, e.g. Cl or Br) to obtain the alcohol XII (step d), see e.g. J. Org. Chem. 2003, 68 (9), 3736-3738).

[0098] If not otherwise indicated, the above-described reactions are generally carried out in a solvent at temperatures between room temperature and the boiling temperature of the solvent employed. Alternatively, the activation energy which is required for the reaction can be introduced into the reaction mixture using microwaves, something which has proved to be of value, in particular, in the case of the reactions catalyzed by transition metals (with regard to reactions using microwaves, see Tetrahedron 2001, 57, p. 9199 ff. p. 9225 ff. and also, in a general manner, "Microwaves in Organic Synthesis", Andre Loupy (Ed.), Wiley-VCH 2002).

[0099] Examples of solvents which can be used are ethers, such as diethyl ether, diisopropyl ether, methyl tert-butyl ether or tetrahydrofuran, aprotic polar solvent, such as dimethylformamide, dimethyl sulfoxide, dimethoxyethane, and acetonitrile, aromatic hydrocarbons, such as toluene and

xylene, ketones, such as acetone or methyl ethyl ketone, halohydrocarbons, such as dichloromethane, trichloromethane and dichloroethane, esters, such as ethyl acetate and methyl butyrate, carboxylic acids, such as acetic acid or propionic acid, and alcohols, such as methanol, ethanol, n-propanol, isopropanol and butanol.

[0100] If desired, it is possible for a base to be present in order to neutralize protons which are released in the reactions. Suitable bases include inorganic bases, such as sodium carbonate, potassium carbonate, sodium hydrogen carbonate or potassium hydrogen carbonate, and, in addition, alkoxides, such as sodium methoxide or sodium ethoxide, alkali metal hydrides, such as sodium hydride, and also organometallic compounds, such as butyllithium compounds or alkylmagnesium compounds, or organic nitrogen bases, such as triethylamine or pyridine. The latter compounds can at the same time serve as solvents.

[0101] The crude product is isolated in a customary manner, for example by filtering, distilling off the solvent or extracting from the reaction mixture, etc. The resulting compounds can be purified in a customary manner, for example by means of recrystallizing from a solvent, by means of chromatography or by means of converting into an acid addition salt.

[0102] The acid addition salts are prepared in a customary manner by mixing the free base with a corresponding acid, where appropriate in solution in an organic solvent, for example a lower alcohol, such as methanol, ethanol or propanol, an ether, such as methyl tert-butyl ether or diisopropyl ether, a ketone, such as acetone or methyl ethyl ketone, or an ester, such as ethyl acetate.

[0103] The compounds according to the invention of the formula I are highly selective dopamine D₃ receptor ligands which, because of their low affinity for other receptors such as D₁ receptors, D₄ receptors, $\alpha 1$ -adrenergic and/or $\alpha 2$ -adrenergic receptors, muscarinic receptors, histamine receptors, opiate receptors and, in particular, dopamine D₂ receptors, give rise to fewer side-effects than do the classic neuroleptics, which are D₂ receptor antagonists.

[0104] The high affinity of the compounds according to the invention for D_3 receptors is reflected in very low in-vitro K_i values of as a rule less than 100 nM (nmoVI), in particular less than 50 nM and, in particular, of less than 10 nM. The displacement of [125 I]-iodosulpride can, for example, be used in receptor binding studies for determining binding affinities for D_3 receptors.

[0105] The selectivity $K_i(D_2)/K_i(D_3)$ of the compounds according to the invention is as a rule at least 10, preferably at least 30, even better at least 50 and particularly advantageously at least 100. The displacement of [3 H]SCH23390, [125 I] idosulpride or [125 I] spiperone can be used, for example, for carrying out receptor binding studies on D_1 , D_2 and D_4 receptors.

[0106] Because of their binding profile, the compounds can be used for treating diseases which respond to dopamine D_3 ligands, i.e. they are effective for treating those disturbances or diseases in which exerting an influence on (modulating) the dopamine D_3 receptors leads to an improvement in the clinical picture or to the disease being cured. Examples of these diseases are disturbances or diseases of the central nervous system.

[0107] Disturbances or diseases of the central nervous system are understood as meaning disturbances which affect the spinal chord and, in particular, the brain. Within the meaning of the invention, the term "disturbance" denotes anomalies which are as a rule regarded as being pathological conditions or functions and which can manifest themselves in the form of particular signs, symptoms and/or malfunctions. While the treatment according to the invention can be directed toward individual disturbances, i.e. anomalies or pathological conditions, it is also possible for several anomalies, which may be causatively linked to each other, to be combined into patterns, i.e. syndromes, which can be treated in accordance with the invention.

[0108] The disturbances which can be treated in accordance with the invention are, in particular, psychiatric and neurological disturbances. These disturbances include, in particular, organic disturbances, including symptomatic disturbances, such as psychoses of the acute exogenous reaction type or attendant psychoses of organic or exogenous cause, e.g., in association with metabolic disturbances, infections and endocrinopathies; endogenous psychoses, such as schizophrenia and schizotypic and delusional disturbances; affective disturbances, such as depressions, mania and/or manic depressive conditions; and also mixed forms of the above-described disturbances; neurotic and somatoform disturbances and also disturbances in association with stress; dissociative disturbances, e.g. loss of consciousness, clouding of consciousness, double consciousness and personality disturbances; disturbances in attention and waking/sleeping behavior, such as behavioral disturbances and emotional disturbances whose onset lies in childhood and youth, e.g. hyperactivity in children, intellectual deficits, in particular attention disturbances (attention deficit disorders), memory disturbances and cognitive disturbances, e.g. impaired learning and memory (impaired cognitive function), dementia, narcolepsy and sleep disturbances, e.g. restless legs syndrome; development disturbances; anxiety states, delirium; sexlife disturbances, e.g. impotence in men; eating disturbances, e.g. anorexia or bulimia; addiction; and other unspecified psychiatric disturbances.

[0109] The disturbances which can be treated in accordance with the invention also include Parkinson's disease and epilepsy and, in particular, the affective disturbances connected thereto.

[0110] The addiction diseases include psychic disturbances and behavioral disturbances which are caused by the abuse of psychotropic substances, such as pharmaceuticals or narcotics, and also other addiction diseases, such as addiction to gaming (impulse control disorders not elsewhere classified). Examples of addictive substances are: opioids (e.g. morphine, heroin and codeine), cocaine; nicotine; alcohol; substances which interact with the GABA chloride channel complex, sedatives, hypnotics and tranquilizers, for example benzodiazepines; LSD; cannabinoids; psychomotor stimulants, such as 3,4-methylenedioxy-N-methylamphetamine (ecstasy); amphetamine and amphetamine-like substances such as methylphenidate and other stimulants including caffeine. Addictive substances which come particularly into consideration are opioids, cocaine, amphetamine or amphetamine-like substances, nicotine and alcohol.

[0111] With regard to the treatment of addiction diseases, particular preference is given to those compounds according to the invention of the formula I which themselves do not possess any psychotropic effect. This can also be observed in a test using rats, which, after having been administered compounds which can be used in accordance with the invention, reduce their self administration of psychotropic substances, for example cocaine.

[0112] According to another aspect of the present invention, the compounds according to the invention are suitable for treating disturbances whose causes can at least partially be attributed to an anomalous activity of dopamine D_3 receptors.

[0113] According to another aspect of the present invention, the treatment is directed, in particular, toward those disturbances which can be influenced, within the sense of an expedient medicinal treatment, by the binding of preferably exogenously administered binding partners (ligands) to dopamine D_3 receptors.

[0114] The diseases which can be treated with the compounds according to the invention are frequently characterized by progressive development, i.e. the above-described conditions change over the course of time; as a rule, the severity increases and conditions may possibly merge into each other or other conditions may appear in addition to those which already exist.

[0115] The compounds according to the invention can be used to treat a large number of signs, symptoms and/or malfunctions which are connected with the disturbances of the central nervous system and, in particular, the abovementioned conditions. These signs, symptoms and/or malfunctions include, for example, a disturbed relationship to reality, lack of insight and ability to meet customary social norms or the demands made by life, changes in temperament, changes in individual drives, such as hunger, sleep, thirst, etc., and in mood, disturbances in the ability to observe and combine, changes in personality, in particular emotional lability, hallucinations, ego-disturbances, distractedness, ambivalence, autism, depersonalization and false perceptions, delusional ideas, chanting speech, lack of synkinesia, short-step gait,

flexed posture of trunk and limbs, tremor, poverty of facial expression, monotonous speech, depressions, apathy, impeded spontaneity and decisiveness, impoverished association ability, anxiety, nervous agitation, stammering, social phobia, panic disturbances, withdrawal symptoms in association with dependency, manifold syndromes, states of excitation and confusion, dysphoria, dyskinetic syndromes and tic disturbances, e.g. Huntington's chorea and Gilles-de-la-Tourette's syndrome, vertigo syndromes, e.g. peripheral positional, rotational and oscillatory vertigo, melancholia, hysteria, hypochondria and the like.

[0116] Within the meaning of the invention, a treatment also includes a preventive treatment (prophylaxis), in particular as relapse prophylaxis or phase prophylaxis, as well as the treatment of acute or chronic signs, symptoms and/or malfunctions. The treatment can be orientated symptomatically, for example as the suppression of symptoms. It can be effected over a short period, be orientated over the medium term or can be a long-term treatment, for example within the context of a maintenance therapy.

[0117] The compounds according to the invention are preferentially suitable for treating diseases of the central nervous system, in particular for treating affective disturbances; neurotic disturbances, stress disturbances and somatoform disturbances and psychoses, and, in particular, for treating schizophrenia and depression. Because of their high selectivity with regard to the D₃ receptor, the compounds I according to the invention are also suitable for treating disturbances of kidney function, in particular disturbances of kidney function which are caused by diabetes mellitus (see WO 00/67847) and, especially, diabetic nephropathy.

[0118] Within the context of the treatment, the use according to the invention of the described compounds involves a method. In this method, an effective quantity of one or more compounds, as a rule formulated in accordance with pharmaceutical and veterinary practice, is administered to the individual to be treated, preferably a mammal, in particular a human being, productive animal or domestic animal. Whether such a treatment is indicated, and in which form it is to take place, depends on the individual case and is subject to medical assessment (diagnosis) which takes into consideration signs, symptoms and/or malfunctions which are present, the risks of developing particular signs, symptoms and/or malfunctions, and other factors.

[0119] As a rule, the treatment is effected by means of single or repeated daily administration, where appropriate together, or alternating, with other active compounds or active compound-containing preparations such that a daily dose of preferably from about 0.1 to 1000 mg/kg of bodyweight, in the case of oral administration, or of from about 0.1 to 100 mg/kg of bodyweight, in the case of parenteral administration, is supplied to an individual to be treated.

[0120] The invention also relates to the production of pharmaceutical compositions for treating an individual, preferably a mammal, in particular a human being, productive animal or domestic animal. Thus, the ligands are customarily administered in the form of pharmaceutical compositions which comprise a pharmaceutically acceptable excipient together with at least one ligand according to the invention and, where appropriate, other active compounds. These compositions can, for example, be administered

orally, rectally, transdermally, subcutaneously, intravenously, intramuscularly or intranasally.

[0121] Examples of suitable pharmaceutical formulations are solid medicinal forms, such as powders, granules, tablets, in particular film tablets, lozenges, sachets, cachets, sugar-coated tablets, capsules, such as hard gelatin capsules and soft gelatin capsules, suppositories or vaginal medicinal forms, semisolid medicinal forms, such as ointments, creams, hydrogels, pastes or plasters, and also liquid medicinal forms, such as solutions, emulsions, in particular oil-in-water emulsions, suspensions, for example lotions, injection preparations and infusion preparations, and eyedrops and eardrops. Implanted release devices can also be used for administering inhibitors according to the invention. In addition, it is also possible to use liposomes or microspheres. When producing the compositions, inhibitors according to the invention are usually mixed or diluted with an excipient. Excipients can be solid, semisolid or liquid materials which serve as vehicles, carriers or medium for the active compound.

[0122] Suitable excipients are listed in the specialist medicinal monographs. In addition, the formulations can comprise pharmaceutically acceptable carriers or customary auxiliary substances, such as glidants; wetting agents; emulsifying and suspending agents; preservatives; antioxidants; antirritants; chelating agents; coating auxiliaries; emulsion stabilizers; film formers; gel formers; odor masking agents; taste corrigents; resin; hydrocolloids; solvents; solubilizers; neutralizing agents; diffusion accelerators; pigments; quaternary ammonium compounds; refatting and overfatting agents; raw materials for ointments, creams or oils; silicone derivatives; spreading auxiliaries; stabilizers; sterilants; suppository bases; tablet auxiliaries, such as binders, fillers, glidants, disintegrants or coatings; propellants; drying agents; opacifiers; thickeners; waxes; plasticizers and white mineral oils. A formulation in this regard is based on specialist knowledge as described, for example, in Fiedler, H. P., *Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete* [Encyclopedia of auxiliary substances for pharmacy, cosmetics and related fields], 4th edition, Aulendorf: ECV-Editio-Kantor-Verlag, 1996.

[0123] The following examples serve to explain the invention without limiting it.

[0124] The magnetic nuclear resonance spectral properties (NMR) refer to the chemical shifts (δ) expressed in parts per million (ppm). The relative area of the shifts in the ¹H NMR spectrum corresponds to the number of hydrogen atoms for a particular functional type in the molecule. The nature of the shift, as regards multiplicity, is indicated as singlet (s), broad singlet (s. br.), doublet (d), broad doublet (d br.), triplet (t), broad triplet (t br.), quartet (q), quintet (quint.) and multiplet (m).

PREPARATION EXAMPLES

Example 1

N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide

1.1 1-Allyl-4-(5-nitropyridin-2-yl)piperazine

[0125] 2.0 g (12.61 mmol) of 2-chloro-5-nitropyridine were dissolved in 8 ml of dimethylformamide, and 3.49 g

(25.23 mmol) of potassium carbonate were added. After that, a solution of 1.75 g (13.88 mmol) of N-allylpiperazine in 2 ml of dimethylformamide was added slowly dropwise to the reaction mixture (exothermic reaction). The reaction mixture was then stirred at room temperature for 2 hours. After the solvent had been concentrated down to dryness, the resulting residue was stirred up in 100 ml of heptane. The precipitate which remained was filtered off with suction. The filtrate was concentrated, resulting in 720 mg of the title compound. The precipitate which had been filtered off with suction was treated with 150 ml of water and extracted three times with diethyl ether. The organic phase was washed with a saturated solution of sodium chloride and dried over sodium sulfate. A further 2.24 g of the title compound were isolated after the solvent had been filtered and concentrated down to dryness. The total yield of 1-allyl-4-(5-nitropyridin-2-yl)piperazine was 2.96 g (95% of theory).

[0126] MS [m+1]: 249.

1.2 6-(4-Allylpiperazin-1-yl)pyridine-3-amine

[0127] 2.2 g (8.86 mmol) of 1-allyl-4-(5-nitropyridin-2-yl)piperazine from Example 1.1 were dissolved in 150 ml of methanol after which 18 g (79.75 mmol) of tin(II) chloride dihydrate were added and the mixture was stirred at 70° C. for 4 hours. After the solvent had been evaporated down to dryness, water was added to the residue. The aqueous reaction mixture was made alkaline with dilute sodium hydroxide solution and then extracted with ethyl acetate. The solid which had precipitated out was filtered off. After that, the phases were separated and the aqueous phase was extracted in each case twice with ethyl acetate and dichloromethane. The combined organic phases were dried over sodium sulfate. 1.74 g (90% of theory) of the title compound were obtained after the drying agent had been removed and the solvent had been evaporated down to dryness.

[0128] MS [m+1]: 219.

1.3N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide

[0129] 1.4 g (7.97 mmol) of 6-(4-allylpiperazin-1-yl)pyridin-3-ylamine from Example 1.2 and 1.74 g (7.97 mmol) of 4-isopropylbenzenesulfonyl chloride were dissolved in 30 ml of tetrahydrofuran at room temperature. 3.3 ml (23.91 mmol) of triethylamine were then added to this mixture. After that, the reaction mixture was stirred overnight at room temperature. After the solvent had been evaporated to dryness, water was added to the residue. The aqueous reaction mixture was made acid with 1N hydrochloric acid and extracted twice with diethyl ether. After that, the aqueous phase was made alkaline (pH 9-10) with a 1N aqueous solution of sodium hydroxide and then extracted twice with diethyl ether. After the combined organic phases had been dried over sodium sulfate, the drying agent had been filtered off and the solvent had been evaporated down to dryness, the resulting residue was chromatographed on silica gel using cyclohexane/ethyl acetate (45:55% to 100% ethyl acetate). The filtrate was evaporated down to dryness. The resulting residue was thoroughly stirred in 10 ml of heptane, filtered off in suction and dried, with 1.93 g (61% of theory) of the title compound being obtained.

[0130] ¹H-NMR (500 MHz, CDCl₃): δ[ppm] 7.7 (s, 1H); 7.6 (d, 2H); 7.4 (d, 1H); 7.3 (d, 2H); 6.6 (d, 1H); 6.4 (bs, 1H);

5.9 (m, 1H); 5.2 (m, 2H); 3.5 (m, 4H); 3.1 (m, 2H); 3.0 (m, 1H); 2.5 (m, 4H); 1.2 (d, 6H).

[0131] MS [m+1]: 401.

Example 2

N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-propylbenzenesulfonamide

[0132] 373 mg of the title compound were obtained in an analogous manner to that described in Example 1.3 when starting with 4-n-propylbenzenesulfonyl chloride.

[0133] ¹H-NMR (500 MHz, CDCl₃): δ[ppm] 7.7 (m, 1H); 7.6 (m, 2H); 7.4 (d, 1H); 7.3 (m, 2H); 6.6 (d, 1H); 6.3 (bs, 1H); 5.9 (m, 1H); 5.2 (m, 2H); 3.5 (m, 4H); 3.1 (m, 2H); 2.6 (m, 2H); 2.5 (m, 4H); 1.7 (m, 2H); 0.9 (m, 3H).

[0134] MS [m+1]: 401.

Example 3

N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-butylbenzenesulfonamide

[0135] 405 mg of the title compound were obtained in an analogous manner to that described in Example 1.3 when starting with 4-n-butylbenzenesulfonyl chloride.

[0136] ¹H-NMR (500 MHz, CDCl₃): δ[ppm] 7.7 (m, 1H); 7.6 (m, 2H); 7.4 (d, 1H); 7.3 (m, 2H); 6.6 (d, 1H); 6.2 (bs, 1H); 5.9 (m, 1H); 5.2 (m, 2H); 3.5 (m, 4H); 3.0 (m, 2H);

[0137] 2.7 (m, 2H); 2.5 (m, 4H); 1.6 (m, 2H); 1.4 (m, 2H); 0.9 (m, 3H).

[0138] MS [m+1]: 415.

Example 4

N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-trifluoromethylbenzenesulfonamide

[0139] 500 mg of the title compound were obtained in an analogous manner to that described in Example 1.3 when starting with 4-trifluoromethylbenzenesulfonyl chloride.

[0140] ¹H-NMR (500 MHz, CDCl₃): δ[ppm] 7.9 (d, 2H); 7.8 (m, 3H); 7.3 (d, 1H); 6.6 (d, 1H); 5.9 (m, 1H); 5.2 (m, 2H); 3.5 (m, 4H); 3.1 (m, 2H); 2.5 (m, 4H).

[0141] MS [m+1]: 427.

Example 5

N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-ethylbenzenesulfonamide Hydrochloride

[0142] The Example 1.3 was repeated with 4-ethylbenzenesulfonyl chloride being used instead of 4-isopropylbenzenesulfonyl chloride. The resulting reaction product was converted into the hydrochloride with ethereal hydrochloric acid, with 480 mg (please complete) of the title compound being obtained.

[0143] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.5 (bs, 1H); 10.0 (s, 1H); 7.8 (d, 2H); 7.6 (d, 2H); 7.4 (m, 3H); 6.9 (d, 1H); 6.0 (m, 1H); 5.5 (m, 2H); 4.3 (m, 2H); 3.8 (m, 2H); 3.4 (m, 2H); 3.3 (m, 2H); 3.0 (m, 2H); 2.7 (m, 2H); 1.2 (t, 3H).

[0144] MS [m+1]: 387 (free base).

Example 6

N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-vinylbenzenesulfonamide Hydrochloride

[0145] Example 1.3 was repeated with 4-vinylbenzenesulfonyl chloride being used instead of 4-isopropylbenzenesulfonyl chloride. The resulting reaction product was converted into the hydrochloride with ethereal hydrochloric acid, with 300 mg of the title compound being obtained.

[0146] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.1 (bs, 1H); 10.0 (s, 1H); 7.8 (d, 1H); 7.6 (m, 4H); 7.3 (d, 1H); 6.9 (d, 1H); 6.8 (dd, 1H); 6.0 (m, 2H); 5.5 (m, 3H); 4.3 (m, 2H); 3.8 (m, 2H); 3.4 (m, 2H); 3.2 (m, 2H); 3.0 (m, 2H).

[0147] MS [m+1]: 385 (free base).

Example 7

4-Isopropyl-N-(6-piperazin-1-ylpyridin-3-yl)benzenesulfonamide

[0148] 95 mg (0.1 mmol) of tris-(dibenzylideneacetone)dipalladium(0) and 44 mg (0.1 mmol) of 1,4-bis-(diphenylphosphino)butane were dissolved in 10 ml of tetrahydrofuran under an argon atmosphere. A solution composed of 1.1 g (2.75 mmol) of N-[6-(4-allylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide from Example 1.3 in 3 ml of tetrahydrofuran was then added dropwise to the reaction mixture. After that, a solution of 386 mg (2.5 mmol) of 2-mercaptobenzoic acid in 2 ml of tetrahydrofuran was added dropwise to the reaction mixture and the mixture was stirred at room temperature for 90 minutes. A solution of a further 386 mg (2.5 mmol) of 2-mercaptobenzoic acid in 2 ml of tetrahydrofuran was then added dropwise to the reaction mixture. The reaction mixture was stirred overnight at room temperature and, after that, the solvent was evaporated down to dryness. 150 ml of water were added to the resulting residue, after which the mixture was made acid with 1N aqueous hydrochloric acid and extracted three times with diethyl ether. The aqueous phase was then made alkaline, to pH>11, with a 1N aqueous solution of sodium hydroxide and subsequently extracted three times with dichloromethane. After that, the aqueous phase was adjusted to pH 8-9, saturated with an aqueous solution of sodium chloride and, after that, extracted several times with dichloromethane. 840 mg (82% of theory) of the title compound were obtained after the combined organic phases had been dried over sodium sulfate and the solvent had been filtered and evaporated down to dryness.

[0149] ¹H-NMR (400 MHz, CDCl₃): δ[ppm] 7.7 (d, 1H); 7.6 (d, 2H); 7.4 (dd, 1H); 7.3 (d, 2H); 6.6 (d, 1H); 3.5 (m, 4H); 3.0 (m, 5H); 1.2 (d, 6H).

[0150] MS [m+1]: 361.

Example 8

N-{6-[4-(Cyclohexylmethyl)piperazin-1-yl]pyridin-3-yl}-4-isopropylbenzenesulfonamide Hydrochloride

[0151] 150 mg (0.42 mmol) of 4-isopropyl-N-(6-piperazin-1-ylpyridin-3-yl)-benzenesulfonamide from Example 7 and 51 mg (0.46 mmol) of cyclohexanaldehyde were dissolved in 5 ml of dichloromethane and 40 μl (0.62 mmol)

of glacial acetic acid under a nitrogen atmosphere. 133 mg (0.63 mmol) of sodium trisacetoxyborohydride were then added. The mixture was stirred at room temperature for 90 minutes and, after that, the solvent was evaporated down to dryness. The resulting residue was taken up in water and this mixture was made to pH>11 with a 1N aqueous solution of sodium hydroxide. After that, the aqueous reaction mixture was extracted with diethyl ether. After the organic phase had been dried over sodium sulfate and the solvent had been filtered and evaporated down to dryness, the resulting residue was converted into the hydrochloride with ethereal hydrochloric acid, resulting in 156 mg (76% of theory) of the title compound.

[0152] ¹H-NMR (500 MHz, DMSO-d₆): δ[ppm] 10.4 (bs, 1H); 10.0 (s, 1H); 7.8 (d, 1H);

[0153] 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (d, 1H); 6.9 (d, 1H); 4.2 (m, 2H); 3.5 (m, 2H); 3.4 (m, 2H); 3.0 (m, 5H); 1.8 (m, 3H); 1.7 (m, 3H); 1.2 (m, 9H); 1.0 (m, 2H).

[0154] MS [m+1]: 457 (free base).

[0155] The compounds of Examples 9 to 12 were prepared in an analogous manner.

Example 9

N-[6-(4-Isobutylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide Hydrochloride

[0156] ¹H-NMR (500 MHz, DMSO-d₆): δ[ppm] 10.4 (bs, 1H); 10.0 (s, 1H); 7.8 (m, 1H); 7.6 (d, 2H); 7.5 (d, 2H); 7.4 (m, 1H); 6.9 (d, 1H); 4.2 (d, 2H); 3.5 (d, 2H); 3.4 (m, 2H); 3.0 (m, 5H); 2.1 (m, 1H); 1.2 (d, 6H); 1.0 (d, 6H).

[0157] MS [m+1]: 417 (free base).

Example 10

4-Isopropyl-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide

[0158] ¹H-NMR (500 MHz, CDCl₃): δ[ppm] 7.7 (d, 1H); 7.6 (d, 2H); 7.4 (dd, 1H); 7.3 (d, 2H); 6.6 (d, 1H); 3.5 (m, 4H); 3.0 (m, 1H); 2.5 (m, 4H); 2.3 (s, 3H); 1.2 (d, 6H).

[0159] MS [m+1]: 375.

Example 11

N-[6-(4-Ethylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide Hydrochloride

[0160] ¹H-NMR (500 MHz, DMSO-d₆): δ[ppm] 10.4 (bs, 1H); 10.0 (s, 1H); 7.8 (d, 1H);

[0161] 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (d, 1H); 6.9 (d, 1H); 4.3 (m, 2H); 3.5 (m, 2H); 3.2 (m, 2H); 3.1 (m, 2H); 3.0 (m, 3H); 1.3 (m, 3H); 1.2 (d, 6H).

[0162] MS [m+1]: 389 (free base).

Example 12

N-{6-[4-(Cyclopropylmethyl)piperazin-1-yl]pyridin-3-yl}-4-isopropylbenzenesulfonamide Hydrochloride

[0163] ¹H-NMR (500 MHz, DMSO-d₆): δ[ppm] 10.8 (bs, 1H); 10.0 (s, 1H); 7.8 (d, 1H);

[0164] 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (d, 1H); 6.9 (d, 1H); 4.3 (m, 2H); 3.6 (m, 2H); 3.3 (m, 2H); 3.0 (m, 5H); 1.2 (d, 6H); 1.1 (m, 1H); 0.6 (m, 2H); 0.4 (m, 2H).

[0165] MS [m+1]: 415 (free base)

Example 13

N-[6-(4-Allyl-3-methylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide Hydrochloride

13.1 3-Methyl-1-(5-nitropyridin-2-yl)piperazine

[0166] 872 mg (6.31 mmol) of potassium carbonate were added to a solution of 500 mg (3.15 mmol) of 2-chloro-5-nitropyridine in 7 ml of dimethylformamide. After that, a solution of 350 mg (3.32 mmol) of 2-methylpiperazine in 3 ml of dimethylformamide was slowly added dropwise to the reaction mixture while cooling with ice (exothermic reaction). The reaction mixture was stirred for 1 hour while cooling with ice and then stirred overnight at room temperature. After the solvent had been evaporated to dryness, the residue was taken up in water and this mixture was extracted three times with diethyl ether. The combined organic phases were dried over sodium sulfate, filtered and evaporated to dryness, with 3-methyl-1-(5-nitropyridin-2-yl)piperazine (Yield: 650 mg, 89% of theory) being obtained.

[0167] ¹H-NMR (500 MHz, CDCl₃): δ[ppm] 9.0 (s, 1H); 8.2 (d, 1H); 6.6 (d, 1H); 4.4 (m, 2H); 3.2 (m, 1H); 3.1 (m, 1H); 2.9 (m, 2H); 2.7 (m, 1H); 1.2 (m, 3H).

[0168] ¹³C-NMR (125 MHz, CDCl₃): 160.4 (C); 146.5 (CH); 134.9 (C); 133.0 (C); 104.5 (CH); 52.2 (CH₂); 50.6 (CH); 45.7 (CH₂); 45.4 (CH₂); 19.6 (CH₃).

13.2 1-Allyl-2-methyl-4-(5-nitropyridin-2-yl)piperazine

[0169] 630 mg (2.72 mmol) of 3-methyl-1-(5-nitropyridin-2-yl)piperazine from Example 13.1 and 267 mg (3.09 mmol) of allyl bromide were dissolved in 10 ml of dimethylformamide. 1.2 ml (8.4 mmol) of triethylamine were then added dropwise to the solution. After the mixture had been stirred at room temperature for 1 hour, a further 65 μl (0.75 mmol) of allyl bromide were added dropwise to the reaction mixture, which was then stirred for a further hour. After that, a further 65 μl (0.75 mmol) of allyl bromide and 0.5 ml (3.6 mmol) of triethylamine were added dropwise. The mixture was then stirred overnight at room temperature. After the solvent had been evaporated down to dryness, the resulting residue was taken up in water and this solution was made alkaline using a 1N aqueous solution of sodium hydroxide. After that, the aqueous reaction mixture was extracted three times with diethyl ether. The combined organic phases were dried over sodium sulfate, filtered and evaporated down to dryness, with 707 mg (90% of theory) of the title compound being obtained.

[0170] MS [m+1]: 263.

13.3 6-(4-Allyl-3-methylpiperazin-1-yl)pyridine-3-amine

[0171] 4.975 g (22.05 mmol) of tin(II) chloride dihydrate were added to a solution of 707 mg (2.45 mmol) of 1-allyl-2-methyl-4-(5-nitropyridin-2-yl)piperazine from Example 13.2 in 50 ml of methanol and the resulting mixture was stirred at 70° C. for 90 minutes. After the solvent had been evaporated down to dryness, water was added to the result-

ing residue and the mixture was made alkaline using a dilute aqueous solution of sodium hydroxide. After that, the aqueous reaction mixture was extracted with ethyl acetate. The solid which had precipitated out was filtered off with suction and the phases were separated. The aqueous phase was extracted with dichloromethane. After that, the combined organic phases were dried over sodium sulfate, filtered and evaporated down to dryness. The resulting title compound was used in the next step without any further purification.

[0172] MS [m+1]: 233.

13.4 N-[6-(4-Allyl-3-methylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide Hydrochloride

[0173] 305 mg (1.31 mmol) of 6-(4-allyl-3-methylpiperazin-1-yl)pyridin-3-ylamine from Example 13.3 and 301 mg (1.38 mmol) of 4-isopropylbenzenesulfonyl chloride were dissolved in 10 ml of tetrahydrofuran at room temperature, after which 0.55 ml (3.94 mmol) of triethylamine was added dropwise. After that, the reaction mixture was stirred overnight at room temperature. After the solvent had been evaporated down to dryness, the resulting residue was treated with water and the mixture was made acid with 1N hydrochloric acid and extracted twice with diethylether. The aqueous phase was made alkaline, to pH 9-10, using a 1N aqueous solution of sodium hydroxide and then extracted twice with diethyl ether. After the combined organic phases had been dried over sodium sulfate and the solvent had been filtered and evaporated down to dryness, the resulting residue was purified by column chromatography (cyclohexane/ethylacetate from 50:50 to 20:80). After that, the filtrate was evaporated down to dryness. The resulting residue was converted into the hydrochloride using ethereal hydrochloric acid, with 417 mg (74% of theory) of the title compound being obtained.

[0174] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.3 (bs, 1H); 10.0 (s, 1H); 7.8 (d, 1H); 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (d, 1H); 6.9 (d, 1H); 6.0 (m, 1H); 5.5 (m, 2H); 4.3 (m, 1H); 4.0 (m, 1H); 3.7 (m, 1H); 3.4 (m, 1H); 3.2 (m, 3H); 3.0 (m, 3H); 1.4 (d, 3H); 1.2 (d, 6H).

[0175] MS [m+1]: 415 (free base).

Example 13a

N-{6-[4-Allyl-(3S)-methylpiperazin-1-yl]pyridin-3-yl}-4-isopropylbenzenesulfonamide (S Enantiomer as Free Base)

[0176] The preparation was effected in analogy with the preparation of the racemic compound, with enantiomerically pure (2S)-methylpiperazine being used in step 13.1 instead of racemic 2-methylpiperazine.

[0177] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.3 (bs, 1H); 10.0 (s, 1H); 7.8 (s, 1H);

[0178] 7.6 (d, 2H); 7.4 (d, 1H); 7.3 (d, 1H); 6.9 (d, 1H); 6.0 (m, 1H); 5.5 (m, 2H); 4.3 (m, 2H); 4.0 (m, 1H); 3.7 (m, 1H); 3.4 (m, 1H); 3.2 (m, 2H); 3.1 (m, 1H); 3.0 (m, 2H); 1.4 (d, 3H); 1.2 (d, 6H).

[0179] MS [m+1]: 415 (free base)

Example 14

4-Isopropyl-N-[6-(3-methyl-4-propylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide Hydrochloride

[0180] 100 mg (0.24 mmol) of N-[6-(4-allyl-3-methylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide

hydrochloride from Example 13.4 were dissolved in 10 ml of ethyl acetate, after which 10 mg of palladium on active charcoal (10%) were added and the mixture was stirred overnight at room temperature under a hydrogen atmosphere. After that, the catalyst was filtered off and the filtrate was evaporated down to dryness. After 1 ml of dichloromethane had been added to the resulting residue, diethyl ether was slowly added dropwise until the solution became cloudy. The reaction mixture was stirred for 30 minutes and the precipitate which had formed was filtered off with suction. The filtrate was evaporated down to dryness, after which the residue was dissolved in a 1:1 mixture of dichloromethane and diethyl ether and converted into the hydrochloride by adding ethereal hydrochloric acid. 71 mg (63% of theory) of the title compound were obtained.

[0181] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 10.9 (bs, 1H); 10.0 (s, 1H); 7.8 (d, 1H); 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (d, 1H); 6.9 (d, 1H); 4.2 (m, 2H); 3.6 (m, 1H); 3.4-3.0 (m, 7H); 1.7 (m, 2H); 1.4 (d, 3H); 1.2 (d, 6H); 0.9 (m, 3H).

[0182] MS [m+1]: 417 (free base).

Example 14a

4-Isopropyl-N-{6-[(3S)-methyl-4-propylpiperazin-1-yl]pyridin-3-yl}benzenesulfonamide as Free Base
(S Enantiomer)

[0183] The preparation was effected in analogy with the preparation of the racemic compound, with enantiomerically pure (2S)-methylpiperazine being used instead of racemic 2-methylpiperazine.

[0184] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 9.7 (s, 1H); 7.7 (s, 1H); 7.6 (d, 2H); 7.4 (d, 2H); 7.2 (d, 1H); 6.7 (d, 1H); 3.8 (m, 2H); 2.9 (m, 2H); 2.8 (m, 1H); 2.6 (m, 2H); 2.3 (m, 1H); 2.1 (m, 2H); 1.4 (m, 2H); 1.2 (d, 6H); 1.0 (m, 3H); 0.8 (m, 3H). MS [m+1]: 417 (free base)

Example 15

N-[5-(4-Allylpiperazin-1-yl)pyridin-2-yl]-4-isopropylbenzenesulfonamide Hydrochloride

15.1 1-Allyl-4-(6-nitropyridin-3-yl)piperazine

[0185] 315 mg (2.5 mmol) of N-allylpiperazine were dissolved in 5 ml of toluene under an argon atmosphere. 93 mg (0.1 mmol) of tris-(dibenzylideneacetone)dipalladium(0) (Pd₂dba₃), 126 mg (0.2 mmol) of 2,2'-bis-(diphenylphosphino)-1,1'-binaphthyl (BINAP), 1.14 g (3.5 mmol) of cesium carbonate and 515 mg (2.54 mmol) of 5-bromo-2-nitropyridine were then added and the mixture was stirred at 120° C., in a microwave oven, for 4 hours. After the reaction mixture had cooled down to room temperature, a saturated aqueous solution of ammonium chloride was added. After that, the aqueous reaction mixture was extracted three times with in each case 50 ml of ethyl acetate. After the organic phase had been dried over sodium sulfate, the drying agent had been filtered off and the solvent had been evaporated down to dryness, the residue was chromatographed through silica gel using ethyl acetate/

methanol (4:1), with 304 mg (46% of theory) of the title compound being obtained.

[0186] ¹H-NMR (400 MHz, CDCl₃): δ[ppm] 8.2 (m, 2H); 7.2 (dd, 1H); 5.9 (m, 1H); 5.3 (m, 2H); 3.5 (m, 4H); 3.1 (m, 2H); 2.6 (m, 4H).

[0187] MS [m+1]: 249

15.2 5-(4-Allylpiperazin-1-yl)pyridine-2-amine

[0188] 300 mg (1.21 mmol) of 1-allyl-4-(6-nitropyridin-3-yl)piperazine from Example 15.1 were dissolved in 20 ml of methanol, after which 2.18 g (9.67 mmol) of tin(II) chloride dihydrate were added and the mixture was stirred at 70° C. for 2 hours. After the solvent had been evaporated down to dryness, the resulting residue was treated with water and this mixture was made alkaline using a dilute aqueous solution of sodium hydroxide and extracted with ethyl acetate. The solid which had precipitated out was filtered off with suction. The phases were then separated and the aqueous phase was extracted three times with ethyl acetate. The combined organic phases were dried over sodium sulfate, filtered and evaporated down to dryness, with 183 mg (69% of theory) of the title compound being obtained.

[0189] MS [m+1]: 219.

15.3 N-[5-(4-Allylpiperazin-1-yl)pyridin-2-yl]-4-isopropylbenzenesulfonamide hydrochloride

[0190] 520 mg (2.38 mmol) of 5-(4-allylpiperazin-1-yl)pyridin-2-ylamine and 495 mg (2.26 mmol) of 4-isopropylbenzenesulfonyl chloride were dissolved in 5 ml of tetrahydrofuran at room temperature, after which 1.0 ml (7.15 mmol) of triethylamine was added dropwise and the mixture was stirred at 40-50° C. for 6 hours. After the solvent had been evaporated down to dryness, the resulting residue was treated with water and this mixture was made acid using 1N aqueous hydrochloric acid and extracted twice with diethyl ether. The aqueous phase was made alkaline, to pH 9-10, using a 1N aqueous solution of sodium hydroxide and then extracted twice with ethyl acetate. After the combined organic phases had been dried over sodium sulfate, the drying agent had been filtered off and the solvent had been evaporated down to dryness, the resulting residue was chromatographed on silica gel using ethyl acetate. After the solvent had been removed, the resulting residue was brought into solution using a little diethyl ether in dichloromethane and then converted into the hydrochloride using ethereal hydrochloric acid. 415 mg (44% of theory) of the title compound were obtained.

[0191] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.6 (bs, 1H); 7.9 (d, 1H); 7.8 (d, 2H); 7.5 (dd, 1H); 7.4 (d, 2H); 7.1 (d, 1H); 6.0 (m, 1H); 5.5 (m, 2H); 3.7 (m, 4H); 3.4 (m, 2H); 3.1 (m, 4H); 3.0 (m, 1H); 1.2 (d, 6H).

[0192] ¹³C-NMR (100 MHz, DMSO-d₆): δ[ppm] 153.3 (C); 144.5 (C); 141.6 (C); 138.4 (C); 134.3 (CH); 127.3 (CH); 127.0 (CH); 126.8 (CH); 124.8 (CH₂); 113.8 (CH); 57.3 (CH₂); 49.6 (CH₂); 45.2 (CH₂); 33.3 (CH); 23.4 (CH₃).

[0193] MS [m+1]: 401.

Example 16

N-[2-(4-Allylpiperazin-1-yl)pyrimidin-5-yl]-4-isopropylbenzenesulfonamide

16.1 2-(4-Allylpiperazin-1-yl)-5-nitropyrimidine

[0194] 114 mg (2.38 mmol) of 50% sodium hydride were added, under a nitrogen atmosphere and while cooling with ice, to a solution of 273 mg (2.17 mmol) of N-allylpiperazine in 5 ml of dimethylformamide. After 30 minutes, a solution of 440 mg (2.17 mmol) of 2-(methylsulfone)-5-nitropyrimidine in 5 ml of dimethylformamide was added dropwise to the reaction mixture. After 10 minutes, 70 ml of water were added and the reaction mixture was extracted twice with in each case 50 ml of ethyl acetate. After the combined organic phases had been dried over sodium sulfate, the drying agent had been filtered off and the solvent had been evaporated to dryness, 535 mg (99% of theory) of the title compound were obtained.

[0195] $^1\text{H-NMR}$ (400 MHz, CDCl_3): δ [ppm] 9.0 (s, 2H); 5.8 (m, 1H); 5.2 (m, 2H); 4.0 (m, 4H); 3.1 (m, 2H); 2.5 (m, 4H).

[0196] MS [m+1]: 250.

16.2 2-(4-Allylpiperazin-1-yl)pyrimidine-5-amine

[0197] 3.84 g (17.0 mmol) of tin(II) chloride dihydrate were added to a solution of 530 mg (2.13 mmol) of 2-(4-allylpiperazin-1-yl)-5-nitropyrimidine from Example 16.1 in 20 ml of methanol and, after that, the reaction mixture was heated at reflux for 1 hour. After the solvent had been evaporated to dryness, the residue was treated with saturated aqueous sodium chloride solution and then made alkaline using dilute aqueous sodium hydroxide solution. After that, the aqueous reaction mixture was extracted with ethyl acetate. The solid which had precipitated out was filtered off with suction. The phases were then separated and the aqueous phase was extracted in each case twice with ethyl acetate and dichloromethane. After the combined organic phases had been dried over sodium sulfate, the drying agent had been filtered off and the solvent had been evaporated down to dryness, 220 mg (46% of theory) of the title compound were obtained.

16.3 N-[2-(4-Allylpiperazin-1-yl)pyrimidin-5-yl]-4-isopropylbenzenesulfonamide

[0198] 216 mg (0.98 mmol) of 2-(4-Allylpiperazin-1-yl)pyrimidin-5-ylamine from Example 16.2 and 215 mg (0.98 mmol) of 4-isopropylbenzenesulfonyl chloride were dissolved in 20 ml of tetrahydrofuran at room temperature, after which 0.4 ml (3.0 mmol) of triethylamine was added dropwise and the mixture was stirred at room temperature overnight. After the solvent had been evaporated down to dryness, water was added to the resulting residue. The aqueous reaction mixture was made acid using 1N aqueous hydrochloric acid and extracted twice with diethyl ether. The aqueous phase was made alkaline to pH 9-10, using a 1N solution of sodium hydroxide and then extracted three times with diethyl ether. The combined organic phases were dried over sodium sulfate. The residue which was obtained after filtering off the drying agent and evaporating the solvent down to dryness was thoroughly stirred with a mixture composed of heptane and diethyl ether, filtered off with suction and dried, with 71 mg (18% of theory) of the title compound being obtained.

[0199] $^1\text{H-NMR}$ (500 MHz, CDCl_3): δ [ppm] 8.0 (s, 2H); 7.7 (d, 2H); 7.3 (d, 2H); 6.2 (bs, 1H); 5.9 (m, 1H); 5.2 (m, 2H); 3.8 (m, 4H); 3.1 (m, 2H); 3.0 (m, 1H); 2.5 (m, 4H); 1.3 (d, 6H).

[0200] MS [m+1]: 402.

Example 17

4-Isopropyl-N-[2-(4-propylpiperazin-1-yl)pyrimidin-5-yl]benzenesulfonamide Hydrochloride

[0201] 70 mg (0.17 mmol) of N-[2-(4-allylpiperazin-1-yl)pyrimidin-5-yl]-4-isopropylbenzenesulfonamide from Example 16.3 were dissolved in 30 ml of ethyl acetate, after which 10 mg of palladium on active charcoal (10%) were added and the mixture was stirred at room temperature for 2 hours under a hydrogen atmosphere. The catalyst was then filtered off and the filtrate was concentrated by evaporation. The residue was brought into solution using 25 ml of diethyl ether and converted into the hydrochloride with ethereal hydrochloric acid, resulting in 58 mg (76% of theory) of the title compound being obtained.

[0202] $^1\text{H-NMR}$ (400 MHz, DMSO-d_6): δ [ppm] 11.0 (bs, 1H); 10.0 (s, 1H); 8.1 (s, 2H); 7.7 (d, 2H); 7.5 (d, 2H); 4.6 (m, 2H); 3.5 (m, 2H); 3.4 (m, 2H); 3.0 (m, 5H); 1.7 (m, 2H); 1.3 (d, 6H); 0.9 m, 3H).

[0203] MS [m+1]: 404 (free base).

EXAMPLE 18

N-[6-(4-Allylpiperazin-1-yl)pyrimidin-4-yl]-4-isopropylbenzenesulfonamide

18.1 N-(6-Chloropyrimidin-4-yl)-4-isopropylbenzenesulfonamide

[0204] 996 mg (5.0 mmol) of isopropylbenzenesulfonamide were dissolved in 20 ml of dimethyl sulfoxide, after which 288 mg (6.0 mmol) of 50% sodium hydride were added and the mixture was stirred at room temperature for 30 minutes. 819 mg (5.5 mmol) of 4,6-dichloropyrimidine were then added and the reaction mixture was stirred overnight at room temperature. Subsequently, the mixture was heated at 90° C. for 3 hours and, after that, stirred at 120° C., in a microwave oven, for 30 minutes. After the reaction mixture had cooled down to room temperature, it was diluted with 150 ml of water, neutralized with citric acid and extracted three times with diethyl ether. The residue, which was obtained after drying with sodium sulfate and after removing the solvent, was dissolved in 100 ml of diethyl ether and extracted with an aqueous solution of sodium hydrogen carbonate. The aqueous phase was acidified and extracted with diethyl ether. The organic phase was dried, filtered and evaporated down to dryness, with 440 mg (28% of theory) of the title compound being obtained.

[0205] MS [m+1]: 312.

18.2 N-[6-(4-Allylpiperazin-1-yl)pyrimidin-4-yl]-4-isopropylbenzenesulfonamide

[0206] 430 mg (1.38 mmol) of N-(6-chloropyrimidin-4-yl)-4-isopropylbenzenesulfonamide from Example 18.1

were dissolved in 3 ml of dimethyl sulfoxide, after which 1.74 g (13.79 mmol) of N-allylpiperazine were added and the mixture was stirred overnight. Subsequently, the reaction mixture was stirred at 100° C., in a microwave oven, for 45 minutes. After the reaction mixture had cooled down to room temperature, it was diluted with 50 ml of water. After that, the aqueous reaction mixture was extracted with 50 ml of ethyl acetate and the precipitate was filtered off with suction, with 190 mg (34% of theory) of the title compound being obtained.

[0207] ¹H-NMR (400 MHz, CDCl₃): δ[ppm] 8.4 (s, 1H); 7.8 (d, 2H); 7.3 (d, 2H); 6.1 (s, 1H); 5.9 (m, 1H); 5.2 (m, 2H); 3.6 (m, 4H); 3.0 (m, 3H); 2.5 (m, 4H); 1.3 (d, 6H).

[0208] MS [m+1]: 402.

Example 19

N-[2-(4-Allylpiperazin-1-yl)pyridin-5-yl]-4-bromobenzenesulfonamide hydrochloride

[0209] The preparation was effected in analogy with Example 1.3, with 4-bromobenzenesulfonyl chloride being used instead of 4-isopropylbenzenesulfonyl chloride. The reaction product which was obtained was converted into the hydrochloride using ethereal hydrochloric acid, resulting in 398 mg of the title compound.

[0210] MS [m+1]: 436/438

Example 20

N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-cyclopropylbenzenesulfonamide

[0211] 398 mg (0.84 mmol) of N-[6-(4-allylpiperazin-1-yl)pyridin-3-yl]-4-bromobenzenesulfonamide from Example 19, 101 mg (1.18 mmol) of cyclopropylboronic acid, 676 mg (3.19 mmol) of K₃PO₄ and 26 mg (0.09 mmol) of tricyclohexylphosphine were dissolved in 4 ml of toluene and 0.2 ml of water under a nitrogen atmosphere. 10 mg (0.04 mmol) of palladium(II) acetate were then added and the mixture was stirred at 100° C., in a microwave oven, for one hour. After the solvent had been evaporated down to dryness, the resulting residue was treated with water and the mixture was then extracted with ethyl acetate. Because the phases only separated poorly, the finely divided solid was filtered off. The aqueous phase was extracted twice with ethyl acetate. After the combined organic phases had been dried over sodium sulfate and the solvent had been filtered and evaporated down to dryness, the resulting residue was purified by column chromatography.

[0212] MS [m+1]: 399

[0213] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.1 (bs, 1H); 9.9 (s, 1H); 7.8 (s, 1H); 7.6 (d, 2H); 7.3 (dd, 1H); 7.2 (d, 2H); 6.9 (d, 1H); 6.0 (m, 1H); 5.5 (d, 2H); 4.3 (m, 2H); 3.8 (m, 2H); 3.4 (m, 2H); 3.2 (m, 2H); 3.0 (m, 2H); 2.0 (m, 1H); 1.0 (m, 2H); 0.8 (m, 2H).

[0214] The compounds of the following examples 21 to 40 were prepared in analogous manner:

Example 21

4-Isopropyl-N-[2-(4-propylpiperazin-1-yl)pyridin-3-yl]-benzenesulfonamide Hydrochloride

[0215] MS [m+1]: 403 (free base).

Example 22

4-Isopropyl-N-[2-(3,5-dimethyl-4-propylpiperazin-1-yl)pyridin-3-yl]-benzenesulfonamide trifluoroacetate

[0216] MS [m+1]: 431 (free base).

Example 23

N-[2-(4-Allyl-3-methylpiperazin-1-yl)pyridin-3-yl]-4-trifluoromethylbenzenesulfonamide Hydrochloride

[0217] MS [m+1]: 441 (free base).

Example 24

N[6-(4-Allyl-3,5-dimethylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide Hydrochloride

[0218] MS [m+1]: 429 (free base)

Example 25

N-[6-(4-Allyl-3,5-dimethylpiperazin-1-yl)pyridin-3-yl]-4-trifluoromethylbenzenesulfonamide Hydrochloride

[0219] MS [m+1]: 455 (free base)

Example 26

N[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-trifluoromethylbenzenesulfonamide

[0220] MS [m+1]: 427

Example 27

4-Bromo-N-[6-(4-propylpiperazin-1-yl)pyridin-3-yl]-benzenesulfonamide

[0221] MS [m+1]: 439/441

Example 28

4-Chloro-N-[6-(4-propylpiperazin-1-yl)pyridin-3-yl]-benzenesulfonamide

[0222] MS [m+1]: 395

Example 29

4-Isopropyl-N-[6-(5-propyl-2,5-diazabicyclo[2.2.1]hept-2-yl)pyridin-3-yl]-benzenesulfonamide Hydrochloride

[0223] MS [m+1]: 415 (free base)

Example 30

N-[6-(5-Allyl-2,5-diazabicyclo[2.2.1]hept-2-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide Hydrochloride

[0224] MS [m+1]: 413 (free base)

Example 31

N-[6-(4-Propylpiperazin-1-yl)pyridin-3-yl]-4-vinylbenzenesulfonamide Hydrochloride

[0225] MS [m+1]: 387 (free base)

Example 32

N-{6-[4-(3-Fluoropropyl)piperazin-1-yl]pyridin-3-yl}-4-isopropylbenzenesulfonamide Hydrochloride

[0226] MS [m+1]: 421 (free base)

Example 33

4-Isopropyl-N-[6-(4-prop-2-yn-1-yl)piperazin-1-yl]pyridin-3-yl]-benzenesulfonamide Hydrochloride

[0227] MS [m+1]: 399 (free base)

Example 34

4-Ethyl-N-[6-(4-propylpiperazin-1-yl)pyridin-3-yl]-benzenesulfonamide Hydrochloride

[0228] MS [m+1]: 389 (free base)

Example 35

N-[6-(4-Allylpiperazin-1-yl)pyridin-3-yl]-4-chlorobenzenesulfonamide Hydrochloride

[0229] MS [m+1]: 393 (free base)

Example 36

4-Isopropyl-N-(4-methyl-6-piperazin-1-ylpyridin-3-yl)-benzenesulfonamide Hydrochloride

[0230] MS [m+1]: 375 (free base)

Example 37

N-[6-(4-Allylpiperazin-1-yl)-4-methylpyridin-3-yl]-4-isopropylbenzenesulfonamide Hydrochloride

[0231] MS [m+1]: 415 (free base)

Example 38

4-Isopropyl-N-[4-methyl-6-(4-propylpiperazin-1-yl)pyridin-3-yl]-benzenesulfonamide Hydrochloride

[0232] MS [m+1]: 417 (free base)

Example 39

N-[4-Methyl-6-(4-propylpiperazin-1-yl)pyridin-3-yl]-4-vinylbenzenesulfonamide Hydrochloride

[0233] MS [m+1]: 401 (free base)

Example 40

N-[6-(4-Butylpiperazin-1-yl)pyridin-3-yl]-4-isopropylbenzenesulfonamide Hydrochloride

[0234] MS [m+1]: 417 (free base)

Example 41

N-{6-[(3S)-4-Ethyl-3-methylpiperazin-1-yl]pyridin-3-yl}-4-isopropylbenzenesulfonamide Hydrochloride

[0235] MS [m+1]: 403 (free base)

Example 42

N-[2-(4-Allylpiperazin-1-yl)pyridin-5-yl]-4-(N-pyrrolidinyl)benzenesulfonamide Fumarate

[0236] Reaction of 0.300 g of 6-(4-allylpiperazin-1-yl)-2-methylpyridin-3-amine (1.29 mmol) and 0.282 g of 4-isopropylbenzene-1-sulfonyl chloride (1.29 mmol) in 10 ml of a mixture of pyridine and dichloromethane (1:2) yielded 0.465 g (87%) of the title compound as a white solid. The title compound was converted into the fumarate salt in methanol by addition of fumaric acid.

[0237] MS [m+1]: 415

[0238] ¹H-NMR (400 MHz, CH₃OD): δ[ppm]: 7.6 (d, 2H), 7.5 (d, 2H), 7.3 (m, 1H), 7.2 (m, 1H), 7.1 (d, 1H), 6.5 (d, 1H), 5.9 (m, 1H), 5.4 (m, 2H), 3.5 (m, 5H), 3.0 (m, 4H), 2.8 (m, 2H), 1.9 (s, 3H), 1.2 (m, 6H).

Example 43

4-Isopropyl-[N-[2-(4-allylpiperazin-1-yl)-6-methylpyridin-5-yl]-4-(N-pyrrolidinyl)benzenesulfonamide, Fumarate

[0239] Reaction of 0.300 g of 6-(4-allylpiperazin-1-yl)-2-methylpyridin-3-amine (1.29 mmol) and 0.282 g of 4-isopropylbenzene-1-sulfonyl chloride (1.29 mmol) in 10 ml of a mixture of pyridine/dichloromethane (1:2) yielded 0.465 g (87%) of the title compound as a white solid. The title compound was converted into the fumarate salt in methanol by addition of fumaric acid.

[0240] MS [m+1]: 415

[0241] ¹H-NMR (400 MHz, MeOD): δ (ppm) 7.6 (d, 2H), 7.5 (d, 2H), 7.3 (m, 1H), 7.2 (m, 1H), 7.1 (d, 1H), 6.5 (d, 1H), 5.9 (m, 1H), 5.4 (m, 2H), 3.5 (m, 5H), 3.0 (m, 4H), 2.8 (m, 2H), 1.9 (s, 3H), 1.2 (m, 6H).

Example 44

4-tert-Butyl-[N-[2-(4-allylpiperazin-1-yl)-6-methylpyridin-5-yl]-benzenesulfonamide, Fumarate

[0242] Reaction of 0.300 g of 6-(4-allylpiperazin-1-yl)-2-methylpyridin-3-amine (1.29 mmol) and 0.300 g of 4-tert-butylbenzene-1-sulfonyl chloride (1.29 mmol) in 10 ml of a mixture of pyridine/dichloromethane (1:2) yielded 0.525 g (95%) of the title compound as a white solid. The title compound was converted into the fumarate salt in methanol by addition of fumaric acid.

[0243] MS [m+1]: 429

[0244] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 9.3 (s, 2H), 8.6 (d, 2H), 7.8 (m, 1H), 7.4 (m, 2H), 7.0 (d, 1H), 6.6 (d, 1H), 5.8 (m, 1H), 5.2 (m, 2H), 3.4 (m, 4H), 2.9 (m, 2H), 2.4 (m, 4H), 1.9 (s, 3H), 1.3 (s, 9H).

Example 45

4-tert-pentyl-[N-[2-(4-Allylpiperazin-1-yl)-6-methylpyridin-5-yl]-benzenesulfonamide, Fumarate

[0245] Reaction of 0.300 g of 6-(4-allylpiperazin-1-yl)-2-methylpyridin-3-amine (1.29 mmol) and 0.318 g of 4-tert-pentylbenzene-1-sulfonyl chloride (1.29 mmol) in 10 mL pyridinedichloromethane (1:2) yielded 0.564 g (99%) of the title compound as a white solid. The title compound was converted into the fumarate salt in methanol by addition of fumaric acid.

[0246] MS [m+1]: 443

[0247] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 9.3 (s, 2H), 8.6 (d, 2H), 7.8 (m, 1H), 7.4 (m, 2H), 7.0 (d, 1H), 6.5 (d, 1H), 5.8 (m, 1H), 5.2 (m, 2H), 3.4 (m, 4H), 3.0 (m, 2H), 2.4 (m, 4H), 1.9 (s, 3H), 1.6 (m, 2H), 1.2 (s, 6H), 0.6 (t, 3H).

[0248] The compounds of Examples 46 to were prepared in analogous manner:

Example 46

4-Ethyl-N-[6-((S)-3-methyl-4-propyl-piperazin-1-yl)-pyridin-3-yl]-benzenesulfonamide, Hydrochloride

[0249] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 10.9 (bs, 1H); 10.0 (s, 1H); 7.8 (m, 1H); 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (dd, 1H); 6.9 (d, 1H); 4.3 (m, 2H); 3.5-2.9 (m, 7H); 2.7 (m, 2H); 1.7 (m, 2H); 1.4 (m, 2H); 1.2 (m, 4H); 0.9 (t, 3H).

[0250] MS [m+1]: 403

Example 47

N-[6-((S)-3-methyl-4-propyl-piperazin-1-yl)-pyridin-3-yl]-4-vinylbenzenesulfonamide, Hydrochloride

[0251] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 10.4 (bs, 1H); 10.0 (s, 1H); 7.8 (s, 1H); 7.7 (m, 4H); 7.3 (d, 1H); 6.9 (d, 1H); 6.8 (dd, 1H); 6.0 (d, 1H); 5.4 (d, 1H); 4.3 (m, 2H); 3.5 (m, 1H); 3.3-2.9 (m, 6H); 1.7 (m, 2H); 1.4 (m, 2H); 1.2 (m, 1H); 0.9 (t, 3H).

[0252] MS [m+1]: 401

Example 48

N-[6-((S)-4-Allyl-3-methyl-piperazin-1-yl)-2-methoxy-pyridin-3-yl]-4-isopropylbenzenesulfonamide, Hydrochloride

[0253] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.1 (bs, 1H); 9.2 (s, 1H); 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (m, 1H); 6.4 (m, 1H); 6.0 (m, 1H); 5.5 (m, 2H); 4.3 (m, 2H); 4.0 (m, 1H); 3.7 (m, 2H); 3.4-3.2 (m, 5H); 3.1 (m, 1H); 3.0 (m, 2H); 1.4 (m, 2H); 1.2 (m, 7H).

[0254] MS [m+1]: 445

Example 49

4-Isopropyl-N-[2-methoxy-6-((S)-3-methyl-4-propyl-piperazin-1-yl)-pyridin-3-yl]-benzenesulfonamide, Hydrochloride

[0255] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 10.8 (bs, 1H); 9.3 (s, 1H); 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (m, 1H); 6.4

(m, 1H); 4.2 (m, 2H); 3.5-3.2 (m, 8H); 3.1 (m, 1H); 3.0 (m, 2H); 1.7 (m, 2H); 1.4 (m, 2H); 1.2 (m, 7H); 0.9 (t, 3H).

[0256] MS [m+1]: 447

Example 50

N-[6-((S)-4-Allyl-3-ethyl-piperazin-1-yl)-pyridin-3-yl]-4-isopropylbenzenesulfonamide, Hydrochloride

[0257] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.2 (bs, 1H); 10.0 (s, 1H); 7.8 (d, 1H); 7.6 (d, 2H); 7.4 (d, 2H); 7.3 (m, 1H); 6.9 (d, 1H); 6.0 (m, 1H); 5.5 (m, 2H); 4.2 (m, 2H); 4.0 (m, 2H); 3.4-3.0 (m, 6H); 1.7 (m, 2H); 1.2 (d, 6H); 1.0 (t, 3H).

[0258] MS [m+1]: 429

Example 51

N-[6-((S)-3-Ethyl-4-propyl-piperazin-1-yl)-pyridin-3-yl]-4-isopropylbenzenesulfonamide, Hydrochloride

[0259] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.2 (bs, 1H); 10.2 (s, 1H); 7.8 (d, 1H); 7.7 (d, 2H); 7.4 (m, 3H); 7.0 (d, 1H); 4.2 (m, 2H); 3.6-3.0 (m, 8H); 2.0 (m, 1H); 1.7 (m, 3H); 1.2 (d, 6H); 1.0 (m, 6H).

[0260] MS [m+1]: 431

Example 52

4-Isopropyl-N-(2-piperazin-1-yl-pyrimidin-5-yl)-benzenesulfonamide

[0261] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 8.0 (s, 2H); 7.6 (d, 2H); 7.4 (d, 2H); 3.6 (m, 4H); 3.0 (m, 1H); 2.7 (m, 4H); 1.2 (d, 6H).

[0262] MS [m+1]: 362

Example 53

N-[2-(4-Ethyl-piperazin-1-yl)-pyrimidin-5-yl]-4-isopropylbenzenesulfonamide, Hydrochloride

[0263] ¹H-NMR (400 MHz, D₂O): δ[ppm] 7.7 (s, 2H); 7.3 (d, 2H); 7.1 (d, 2H); 4.3 (m, 2H); 3.3 (m, 2H); 3.0 (m, 4H); 2.7 (m, 3H); 1.0 (t, 3H); 0.9 (d, 6H).

[0264] MS [m+1]: 390

Example 54

N-[2-((S)-4-Ethyl-3-methyl-piperazin-1-yl)-pyrimidin-5-yl]-4-isopropylbenzenesulfonamide, Hydrochloride

[0265] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 10.7 (bs, 1H); 10.0 (s, 1H); 8.1 (m, 2H); 7.6 (d, 2H); 7.4 (d, 2H); 4.6 (m, 2H); 3.4 (m, 4H); 3.2 (m, 1H); 3.0 (m, 3H); 1.4 (m, 2H); 1.2 (m, 10H).

[0266] MS [m+1]: 404

Example 55

N-[2-((S)-4-Allyl-3-methyl-piperazin-1-yl)-pyrimidin-5-yl]-4-isopropylbenzenesulfonamide, Hydrochloride

[0267] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.0 (bs, 1H); 10.0 (s, 1H); 8.1 (s, 2H); 7.6 (d, 2H); 7.4 (d, 2H); 6.0

(m, 1H); 5.5 (m, 2H); 4.6 (m, 2H); 4.0 (m, 2H); 3.7 (m, 2H); 3.3 (m, 3H); 3.0 (m, 1H); 1.4 (m, 2H); 1.2 (m, 7H).

[0268] MS [m+1]: 416

Example 56

4-Isopropyl-N-[2-((S)-3-methyl-4-propyl-piperazin-1-yl)-pyrimidin-5-yl]benzenesulfonamide, Hydrochloride

[0269] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 10.6 (bs, 1H); 10.0 (s, 1H); 8.1 (m, 2H); 7.6 (d, 2H); 7.4 (d, 2H); 4.5 (m, 2H); 3.5 (m, 2H); 3.3 (m, 3H); 3.0 (m, 3H); 1.7 (m, 2H); 1.4 (m, 2H); 1.2 (m, 7H); 0.9 (t, 3H).

[0270] MS [m+1]: 418

Example 57

4-Ethyl-N-[2-((S)-3-methyl-4-propyl-piperazin-1-yl)-pyrimidin-5-yl]benzenesulfonamide, Hydrochloride

[0271] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 10.8 (bs, 1H); 10.0 (s, 1H); 8.1 (m, 2H); 7.6 (d, 2H); 7.4 (d, 2H); 4.5 (m, 2H); 3.5 (m, 2H); 3.3 (m, 3H); 3.0 (m, 2H); 2.7 (m, 2H); 1.7 (m, 2H); 1.4 (m, 2H); 1.2 (m, 4H); 0.9 (t, 3H).

[0272] MS [m+1]: 404

Example 58

N-[2-((S)-3-Methyl-4-propyl-piperazin-1-yl)-pyrimidin-5-yl]-4-Vinylbenzenesulfonamide, Hydrochloride

[0273] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 10.7 (bs, 1H); 10.0 (s, 1H); 8.1 (d, 2H); 7.7 (m, 4H); 6.8 (dd, 1H); 6.0 (d, 1H); 5.5 (d, 1H); 4.5 (m, 2H); 3.5 (m, 2H); 3.3 (m, 3H); 3.0 (m, 2H); 1.7 (m, 2H); 1.4 (m, 2H); 1.2 (m, 1H); 0.9 (t, 3H).

[0274] MS [m+1]: 402

Example 59

4-Isopropyl-benzenesulfonic acid 6-(4-allyl-piperazin-1-yl)-pyridin-3-yl Ester

[0275] 59.1 4-Isopropyl-benzenesulfonic acid 6-chloropyridin-3-yl ester

[0276] A reaction flask containing 500 mg of 6-chloropyridin-3-ol (3.86 mmol) and 844 mg of 4-isopropyl-benzenesulfonylchloride (0.20 mmol) in dry tetrahydrofuran (10 ml) was flushed with N₂. 1.6 ml of triethylamine were added and the reaction mixture was stirred at room temperature for 2 h. Thereby 4-isopropyl-benzenesulfonic acid 6-chloropyridin-3-yl ester was obtained in 98% yield.

[0277] MS [m+1]: 312

[0278] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 8.1 (d, 1H); 7.8 (d, 2H); 7.6 (m, 4H); 3.0 (m, 1H); 1.2 (d, 6H).

[0279] 59.2 4-Isopropyl-benzenesulfonic acid 6-(4-allyl-piperazin-1-yl)-pyridin-3-yl ester

[0280] A flask containing 45 mg of palladium(II) acetate (0.2 mmol), 126 mg of BINAP (0.20 mmol) and 0.233 mg

of sodium tert-butyrate in dry toluene (15 mL) was flushed with nitrogen. 1.275 g of 1-allylpiperazin (10 mmol) were added and the reaction mixture was heated to 50° C. 630 mg of 4-isopropyl-benzenesulfonic acid 6-chloropyridin-3-yl ester (2.0 mmol) were dissolved in toluene and the solution was slowly added over a period of 10 minutes into the reaction mixture with stirring. The mixture was heated to reflux with stirring for 7 h. Thereby 4-isopropyl-benzenesulfonic acid 8-(4-allyl-piperazin-1-yl)-pyridin-3-yl ester was obtained in 21% yield.

[0281] MS [m+1]: 402

[0282] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 7.8 (d, 2H); 7.7 (d, 1H); 7.6 (d, 2H); 7.2 (dd, 1H); 6.8 (d, 1H); 5.8 (m, 1H); 5.2 (m, 2H); 3.5 (m, 4H); 3.0 (m, 3H); 2.5 (m, 4H); 1.2 (d, 6H).

Example 60

4-Isopropyl-benzenesulfonic acid 6-(4-propyl-piperazin-1-yl)-pyridin-3-yl Ester, Hydrochloride

[0283] In a reaction flask 5 mg of palladium on charcoal were added to 50 mg of 4-isopropylbenzenesulfonic acid 6-(4-allyl-piperazin-1-yl)-pyridin-3-yl ester from example 59 (0.12 mmol) in dry ethyl acetate (5 ml). The atmosphere was then charged with hydrogen gas and the reaction mixture was stirred at room temperature for 1 hour. Thereby, 4-isopropyl-benzenesulfonic acid 6-(4-propyl-piperazin-1-yl)-pyridin-3-yl ester were obtained in 58% yield. The compound was converted into its hydrochloride salt by addition of HCl in ether.

[0284] MS [m+1]: 404

[0285] ¹H-NMR (400 MHz, DMSO-d₆): δ[ppm] 11.0 (bs, 1H); 7.8 (m, 3H); 7.6 (d, 2H); 7.3 (dd, 1H); 6.9 (d, 1H); 4.3 (d, 2H); 3.5 (d, 2H); 3.3 (dd, 2H); 3.0 (m, 5H); 1.7 (m, 2H); 1.3 (d, 6H); 0.9 (t, 3H).

[0286] Examples of galenic administration forms

A) Tablets

[0287] Tablets of the following composition are pressed on a tablet press in the customary manner:

[0288] 40 mg of substance from Example 2

[0289] 120 mg of corn starch

[0290] 13.5 mg of gelatin

[0291] 45 mg of lactose

[0292] 2.25 mg of Aerosil® (chemically pure silicic acid in submicroscopically fine dispersion)

[0293] 6.75 mg of potato starch (as a 6% paste)

B) Sugar-coated tablets

[0294] 20 mg of substance from Example 2

[0295] 60 mg of core composition

[0296] 70 mg of saccharification composition

[0297] The core composition consists of 9 parts of corn starch, 3 parts of lactose and 1 part of 60:40 vinylpyrrolidone/vinyl acetate copolymer. The saccharification composition consists of 5 parts of cane sugar, 2 parts of corn starch, 2 parts of calcium carbonate and 1 part of talc. The sugar-

coated tablets which had been prepared in this way are subsequently provided with a gastric juice-resistant coating.

Biological investigations—receptor binding studies:

[0298] The substance to be tested was either dissolved in methanol/Chremophor (BASF-AG) or in dimethyl sulfoxide and then diluted with water to the desired concentration.

[0299] Dopamine D₃ receptor:

[0300] The assay mixture (0.250 ml) was composed of membranes derived from ~10⁶ HEK-293 cells possessing stably expressed human dopamine D₃ receptors, 0.1 nM [¹²⁵I]-iodosulpride and incubation buffer (total binding) or, in addition, test substance (inhibition curve) or 1 μM spiperone (nonspecific binding). Each assay mixture was run in triplicate. The incubation buffer contained 50 mM tris, 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 2 mM MgCl₂ and 0.1% bovine serum albumin, 10 μM quinolone and 0.1% ascorbic acid (prepared fresh daily). The buffer was adjusted to pH 7.4 with HCl.

[0301] Dopamine D_{2L} receptor:

[0302] The assay mixture (1 ml) was composed of membranes from ~10⁶ HEK-293 cells possessing stably expressed human dopamine D_{2L} receptors (long isoform) and 0.01 nM [¹²⁵I] iodospiperone and incubation buffer (total binding) or, in addition, test substance (inhibition curve) or 1 μM haloperidol (nonspecific binding). Each assay mixture was run in triplicate.

[0303] The incubation buffer contained 50 mM tris, 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 2 mM MgCl₂ and 0.1% bovine serum albumin. The buffer was adjusted to pH 7.4 with HCl.

[0304] Measurement and analysis:

[0305] After having been incubated at 25° C. for 60 minutes, the assay mixtures were filtered through a Whatman GF/B glass fiber filter under vacuum using a cell collecting device. The filters were transferred to scintillation vials using a filter transfer system. After 4 ml of Ultima Gold® (Packard) have been added, the samples were shaken for one hour and the radioactivity was then counted in a Beta-Counter (Packard, Tricarb 2000 or 2200CA). The cpm values were converted into dpm using a standard quench series and the program belonging to the instrument.

[0306] The inhibition curves were analyzed by means of iterative nonlinear regression analysis using the Statistical Analysis System (SAS) which is similar to the "LIGAND" program described by Munson and Rodbard.

[0307] In these tests, the compounds according to the invention exhibit very good affinities for the D₃ receptor (<100 nM, frequently <50 nM) and bind selectively to the D₃ receptor. The results of the binding tests are given in Table 1.

TABLE 1


Example	K _i (D ₃) [nM]	Selectivity vs. D _{2L} *
1	3.0	232
2	5.5	25
3	5.9	15
5	11.4	108

TABLE 1-continued

Example	K _i (D ₃) [nM]	Selectivity vs. D _{2L} *
6	9.7	169
7	11.4	68
10	7.5	93
11	6.2	77
13	3.6	131
13a	2.7	96
14	2.5	81
14a	1.5	184
16	3.8	131
17	8.2	148
19	36.9	91
22	21.9	22
24	25.0	47
27	21.4	55
28	25.3	67
29	16.9	31
30	11.1	17
31	14.0	96
32	17.0	74
34	9.6	73
35	26.6	51
36	5.4	50
37	2.7	86
38	17.2	22
39	34.6	30
42	5.3	33
43	1.1	80
44	1.4	41
45	1.4	24
46	7.1	50
47	6.1	118
48	0.8	21
49	0.4	19
50	9.9	57
51	3.1	101
52	14.2	63
53	8.8	149
54	2.3	84
55	2.9	159
56	1.6	399
57	7.3	234
58	6.2	254
59	16.2	75
60	17.4	47

*K_i(D₃)/K_i(D_{2L})

1. An N-[(piperazinyl)hetaryl]arylsulfonamide compound of the general formula I



in which

R is oxygen, a group N—R³ or a group CR^{3a}R^{3b};

Q is a bivalent, 6-membered heteroaromatic radical which possesses 1 or 2 N atoms as ring members and which optionally carries one or two substituents R^a which is/are selected, independently of each other, from halogen, CN, NO₂, CO₂R⁴, COR⁵, C₁—C₄-alkyl, C₁—C₄-alkoxy, C₁—C₄-haloalkyl, NH₂, NHR⁶, NR⁶R⁷ and C₁—C₄-haloalkoxy;

Ar is phenyl or a 6-membered heteroaromatic radical which possesses 1 or 2 N atoms as ring members and which optionally carries one or two substituents R^b , which is/are selected from halogen, NO_2 , CN, CO_2R^4 , COR^5 , NH_2 , NHR^6 , NR^6R^7 , C_1-C_6 -alkyl, C_1-C_6 -haloalkyl, C_1-C_6 -alkoxy, C_1-C_6 -haloalkoxy, C_2-C_6 -alkenyl, C_2-C_6 -alkynyl, C_3-C_6 -cycloalkyl, C_3-C_6 -cycloalkoxy, C_3-C_6 -cycloalkyl- C_1-C_4 -alkyl and C_1-C_4 -haloalkyl, with it also being possible for two radicals R^b which are bonded to adjacent C atoms of Ar to be together C_3-C_4 -alkylene;

n is 0, 1 or 2;

R^1 is hydrogen, C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_3-C_6 -cycloalkyl, C_3-C_6 -cycloalkyl- C_1-C_4 -alkyl, C_1-C_4 -hydroxyalkyl, C_1-C_4 -alkoxy- C_1-C_4 -alkyl, C_3-C_4 -alkenyl or C_3-C_4 -alkynyl;

R^2 is C_1-C_4 -alkyl or, together with R^1 , is C_2-C_5 -alkylene or, in the case of $n=2$, the two radicals R^2 can together be C_1-C_4 -alkylene;

R^3 is hydrogen or C_1-C_4 -alkyl;

R^{3a} , R^{3b} are, independently of each other, hydrogen or C_1-C_4 -alkyl;

R^4 is C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_2-C_4 -alkenyl, C_3-C_6 -cycloalkyl, C_3-C_6 -cycloalkyl- C_1-C_4 -alkyl, phenyl or benzyl; and

R^5 is hydrogen, C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_2-C_4 -alkenyl, C_3-C_6 -cycloalkyl,

C_3-C_6 -cycloalkyl- C_1-C_4 -alkyl, phenyl or benzyl;

R^6 , R^7 are each independently selected from C_1-C_4 -alkyl, C_1-C_4 -haloalkyl or together with the nitrogen to which they are bound form a saturated 3-, 4-, 5- or 6-membered heterocycle, which additionally may comprise an oxygen atom or an additional nitrogen atom as a ring member and which may carry 1, 2, 3 or 4 C_1-C_4 -alkyl groups;

the N-oxides thereof and the physiologically tolerated acid addition salts of these compounds;

with the exception of the compounds: 4-methyl-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide and 4-chloro-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]benzenesulfonamide.

2. The compound as claimed in claim 1, wherein R is $N-R^3$ with R^3 being H or C_1-C_4 -alkyl.

3. The compound as claimed in claim 2, wherein

Q is a bivalent, 6-membered heteroaromatic radical which possesses 1 or 2 N atoms as ring members and which optionally carries one or two substituents


R^a which is/are selected, independently of each other, from halogen, CN, NO_2 , CO_2R^4 , COR^5 , C_1-C_4 -alkyl and C_1-C_4 -haloalkyl and

Ar is phenyl or a 6-membered heteroaromatic radical which possesses 1 or 2N atoms as ring members and which optionally carries one or two substituents R^b , which is/are selected from halogen, NO_2 , CN, CO_2R^4 , COR^5 , C_1-C_6 -alkyl, C_2-C_6 -alkenyl, C_2-C_6 -alkynyl, C_3-C_6 -cycloalkyl, C_3-C_6 -cycloalkyl- C_1-C_4 -alkyl and C_1-C_4 -haloalkyl, with it also

being possible for two radicals R^b which are bonded to adjacent C atoms of Ar to be together C_3-C_4 -alkylene.

4. The compound as claimed in claim 1, in which the piperazine ring is bonded to the heteroaromatic radical Q in the para position in relation to the group $R-SO_2-Ar$.

5. The compound as claimed in claim 1, in which Q is a radical of the formula




in which A_1 , A_2 and A_3 are, independently of each other, N or CH, one or two of the variables A_1 , A_2 and A_3 can also be $C-R^a$, $k=0$ or 1 and R^a is selected from halogen, C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy, NH_2 , NHR^6 , NR^6R^7 , C_1-C_4 -haloalkoxy, with A_1 , A_2 and A_3 not simultaneously being N or simultaneously being selected from CH and $C-R^a$.

6. The compound as claimed in claim 5, in which A_3 is nitrogen, A_2 is CH and A_1 is N or CH and wherein the piperazine radical is located in the 2 position.

7. The compound as claimed in claim 6, in which Q is pyridin-2,5-diyl which carries the piperazine radical in the 2 position.

8. The compound as claimed in claim 6, in which Q is a radical of the formula



in which A_1 and A_2 are, independently of each other, N or CH and R^a is selected from, C_1-C_4 -alkoxy, NH_2 , NHR^6 , NR^6R^7 and C_1-C_4 -haloalkoxy.


9. The compound as claimed in claim 8, in which A_1 is N or CH and A_2 is CH and wherein the piperazine radical is located in the 2 position.

10. The compound as claimed in claim 1, in which the radical Ar carries a substituent R^b in the para position and, where appropriate, a further substituent R^b in the meta position or in the ortho position, in each case based on the binding site of the sulfonamide group.

11. The compound as claimed in claim 1, in which Ar is phenyl or pyridyl, which radicals possess, where appropriate, one or 2 R^b substituents.

12. The compound as claimed in claim 1, in which R^1 is different from hydrogen and methyl.

13. The compound as claimed in claim 1 of the general formula Ia



in which n, R¹, R², R³, R^a and R^b have the meanings given in claim 1 and in which either

A₁, A₂ and A₃ are, independently of each other, N or CH and one or two of the variables A₁, A₂ and A₃ can also be C—R^a, with A₁, A₂ and A₃ not simultaneously being N or simultaneously being selected from CH and C—R^a,

X and Y are selected from CH, C—R^b and N, in which R^b is halogen, methyl, CN, difluoromethyl or trifluoromethyl, with X and Y not simultaneously being N or simultaneously being C—R^b, and

k is 0 or 1.

14. The compound of the formula Ia as claimed in claim 13, in which k=0, with A₁, A₂ and A₃ being, independently of each other, N or CH and A₁, A₂ and A₃ not simultaneously being N or simultaneously being CH.

15. The compound of the formula Ia as claimed in claim 14, in which A₁ is CH or N, A₂ is CH and A₃ is N.

16. The compound of the formula Ia as claimed in claim 13, in which k is 1, A₁ is CH or N, A₂ is CH and A₃ is N, and R^a is selected from, C₁—C₄-alkoxy, NH₂, NHR⁶, NR⁶R⁷ and C₁—C₄-haloalkoxy and R^a is bound to the carbon atom adjacent to A₃.


17. The compound of the formula Ia as claimed in claim 13, in which n is 0 or 1 and, in the case of n=1, R² is bonded to the C atom of the piperazine ring which is adjacent to the group R¹—N and is a methyl group having the S configuration.

18. The compound of the formula Ia as claimed claim 13, in which the radical Ar carries a substituent R^b in the para position and, where appropriate, a further substituent R^b in the meta position or in the ortho position, in each case based on the binding site of the sulfonamide group.

19. The compound of the formula Ia as claimed in claim 13, in which Ar is phenyl or pyridyl, which radicals possess, where appropriate, one or 2 R^b substituents.


20. The compound of the formula Ia as claimed in claim 13, in which R¹ is different from hydrogen and methyl.

21. The compound of the formula Ia as claimed in claim 13, of the general formula Ia.1



in which n, X, Y, R¹, R², R³, R^a and R^b have the meanings given in claim 13 and q is 0, 1 or 2.


22. The compound of the formula Ia as claimed in claim 13 of the general formula Ia.2



in which n, X, Y, R¹, R², R³, R^a and R^b have the meanings given in claim 13 and q is 0 or 1.

23. A pharmaceutical composition which comprises at least one N-[(piperazinyl)hetaryl]arylsulfonamide compound as claimed in claim 1 and/or at least one physiologically tolerated acid addition salt of I and/or an N-oxide of I, where appropriate together with physiologically acceptable carriers and/or auxiliary substances.

24. The use of at least one N-[(piperazinyl)hetaryl]aryl-sulfonamide compound of the formula I




in which Q, Ar, n, R¹, R² and R³ have the previously mentioned meanings, of the N-oxides thereof and of the physiologically tolerated acid addition salts thereof for producing a pharmaceutical composition for treating diseases which respond to influencing by dopamine D₃ receptor antagonists or dopamine D₃ agonists.

25. The use as claimed in claim 24 for treating diseases of the central nervous system.

26. The use as claimed in claim 24 for treating kidney function disturbances.

27. A method for treating a medical disorder susceptible to treatment with a dopamine D₃ receptor antagonist or a dopamine D₃ agonist, said method comprising administering an effective amount of at least one compound of the claim 1



to a subject in need thereof.

28. The method as claimed in claim 27, wherein the medical disorder is a disease of the central nervous system.

29. The method as claimed in claim 27 wherein the medical disorder is a disturbance of kidney function.