




(86) Date de dépôt PCT/PCT Filing Date: 2009/06/24
(87) Date publication PCT/PCT Publication Date: 2010/01/21
(85) Entrée phase nationale/National Entry: 2010/12/21
(86) N° demande PCT/PCT Application No.: US 2009/048504
(87) N° publication PCT/PCT Publication No.: 2010/008894
(30) Priorité/Priority: 2008/06/24 (US61/075,307)

(51) Cl.Int./Int.Cl. *C07C 233/24* (2006.01),
A61K 31/167 (2006.01), *A61K 31/216* (2006.01),
A61K 31/27 (2006.01), *A61P 25/08* (2006.01),
C07C 271/28 (2006.01), *C07C 69/734* (2006.01)
(71) Demandeur/Applicant:
VALEANT PHARMACEUTICALS INTERNATIONAL, US
(72) Inventeurs/Inventors:
WU, JIM ZHEN, CN;
VERNIER, JEAN-MICHEL, US;
CHEN, HUANMING, US;
SONG, JIANLAN, US
(74) Agent: OSLER, HOSKIN & HARCOURT LLP

(54) Titre : DERIVES DE BENZYLOXY ANILIDE UTILES EN TANT QUE MODULATEURS DU CANAL POTASSIQUE
(54) Title: BENZYLOXY ANILIDE DERIVATIVES USEFUL AS POTASSIUM CHANNEL MODULATORS



(57) **Abrégé/Abstract:**

The present invention relates to benzyloxylanilide derivatives having the following structural formula (I): The compounds of the present invention are useful for the treatment and prevention of diseases and disorders which are affected by activation or modulation of potassium ion channels. One such condition is seizure disorders.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
21 January 2010 (21.01.2010)

PCT

(10) International Publication Number
WO 2010/008894 A1

(51) International Patent Classification:

C07D 213/30 (2006.01) *A61K 31/472* (2006.01)

(21) International Application Number:

PCT/US2009/048504

(22) International Filing Date:

24 June 2009 (24.06.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/075,307 24 June 2008 (24.06.2008) US

(71) Applicant (for all designated States except US):
VALEANT PHARMACEUTICALS INTERNATIONAL-AL; One Enterprise, Aliso Viego, CA 92656 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WU, Jim, Zhen** [CN/CN]; 720 Cai Lan Road, Building 5, Pudong, Shanghai, 201203 (CN). **VERNIER, Jean-michel** [FR/US]; 24831 Cutter, Laguna Niguel, CA 92677 (US). **CHEN, Huanming** [CN/US]; 12 Armagosa, Irvine, CA 92602 (US). **SONG, Jianlan** [CN/US]; 13531 Semora Place, Cerritos, CA 90703 (US).(74) Agents: **CADENA, Deborah, L.** et al.; Mcdermott, Will & Emery Llp, 11682 El Camino Real, Suite 400, San Diego, CA 92130-2047 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: BENZYLOXY ANILIDE DERIVATIVES USEFUL AS POTASSIUM CHANNEL MODULATORS



(57) Abstract: The present invention relates to benzyloxyanilide derivatives having the following structural formula (I): The compounds of the present invention are useful for the treatment and prevention of diseases and disorders which are affected by activation or modulation of potassium ion channels. One such condition is seizure disorders.

**BENZYLOXY ANILIDE DERIVATIVES USEFUL
AS POTASSIUM CHANNEL MODULATORS**

Field of the Invention

5 This application claims the benefit of priority of U.S. provisional application Serial No. 61/075,307, filed June 24, 2008, the entire contents of which is incorporated herein by reference.

This invention provides benzyloxy anilide derivatives which activate or otherwise modulate voltage-gated potassium channels. The compounds are useful for the treatment and prevention of diseases and disorders which are affected by modulation of potassium ion channels. One such
10 condition is seizure disorders.

Background of the Invention

Epilepsy is a well-known neurological disease, found in about 3% of the population. Approximately 30% of patients with epilepsy do not respond to currently available therapies. Such unfortunate patients – who number hundreds of thousands of people world-wide – must
15 contend with both uncontrolled seizures and the resulting narrowing of their options in such crucial areas of life as health insurance, employment, and driving.

Retigabine (N-[2-amino-4-(4-fluorobenzylamino)phenyl]carbamic acid, ethyl ester) (U.S. Patent No. 5,384,330) has been found to be an effective treatment of seizure disorders and has also been found useful in treating pain. Retigabine has been found to be particularly potent in models for
20 the drug-refractory types of epilepsy. Bialer, M. *et al.*, *Epilepsy Research* 1999, 34, 1-41; Blackburn-Munro and Jensen, *Eur. J. Pharmacol.* 2003, 460, 109-116; Wickenden, A.D. *et al.*, *Expert Opin. Ther. Patents*, 2004, 14(4).

"Benign familial neonatal convulsions," an inherited form of epilepsy, has been associated with mutations in the KCNQ2/3 channels. Biervert, C. *et al.*, *Science* 1998, 27, 403-06; Singh, N.A.,
25 *et al.*, *Nat. Genet.* 1998, 18, 25-29; Charlier, C. *et al.*, *Nat. Genet.* 1998, 18, 53-55; Rogawski, *Trends in Neurosciences* 2000, 23, 393-398. Subsequent investigations have established that one important site of action of retigabine is the KCNQ2/3 channel. Wickenden, A.D. *et al.*, *Mol. Pharmacol.* 2000, 58, 591-600; Main, M.J. *et al.*, *Mol. Pharmacol.* 2000, 58, 253-62. Retigabine has been shown to increase the conductance of the channels at the resting membrane potential,
30 with a possible mechanism involving binding of the activation gate of the KCNQ 2/3 channel. Wuttke, T.V., *et al.*, *Mol. Pharmacol.* 2005. Additionally, retigabine has been shown to increase


neuronal M currents and to increase the channel open probability of KCNQ 2/3 channels. Delmas, P. and Brown, D.A. *Nat. Revs Neurosci.*, vol. 6, 2005, 850-62; Tatulian, L. and Brown, D.A., *J. Physiol.*, (2003) 549, 57-63.

The seizure type that has been most resistant to therapy is the so-called "complex partial seizure." Retigabine is active in several seizure models, including, as indicated above, models for drug-refractory epilepsy. Because of retigabine's broad spectrum of activity and its unusual molecular mechanism, there is hope that retigabine will be effective in management of several seizure types, including the complex partial seizure, which have been resistant to treatment. Porter, R. J., Nohria, V., and Rundfeldt, C., *Neurotherapeutics*, 2007, vol. 4, 149-154.

The recognition of retigabine as a potassium channel opener has inspired a search among compounds with structural features in common with retigabine for other compounds which can affect the opening of, or otherwise modulate, potassium ion channels.

Brief Description of the Invention

The present invention provides a compound of formula I



I

wherein R_1 and R_2 , are, independently, H, CN, halogen, CH_2CN , OH, NO_2 , CH_2F , CHF_2 , CF_3 , CF_2CF_3 , $\text{C}_1\text{-C}_6$ alkyl, $\text{C}(=\text{O})\text{C}_1\text{-C}_6$ alkyl; NH_2 , $\text{NH-C}_1\text{-C}_6$ alkyl; $\text{N}(\text{C}_1\text{-C}_6 \text{ alkyl})\text{-C}_1\text{-C}_6$ alkyl, $\text{NHC}(=\text{O})\text{C}_1\text{-C}_6$ alkyl, $\text{C}(=\text{O})\text{N}(\text{CH}_3)_2$, $\text{C}(=\text{O})\text{N}(\text{Et})_2$, $\text{C}(=\text{O})\text{NH}_2$, $\text{C}(=\text{O})\text{NH-C}_1\text{-C}_6$ alkyl, SO_2NH_2 , $\text{NHSO}_2\text{-C}_1\text{-C}_6$ alkyl; $\text{C}(=\text{O})\text{OC}_1\text{-C}_6$ alkyl, $\text{OC}(=\text{O})\text{C}_1\text{-C}_6$ alkyl, $\text{OC}_1\text{-C}_6$ alkyl, $\text{SC}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, $(\text{CH}_2)_m\text{C}_3\text{-C}_6$ cycloalkyl, $\text{C}_3\text{-C}_6$ cycloalkenyl, $(\text{CH}_2)_m\text{C}_3\text{-C}_6$ cycloalkenyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, Ar, $(\text{CH}_2)_m$ thienyl, $(\text{CH}_2)_m$ furyl, $(\text{CH}_2)_m$ imidazolyl, $(\text{CH}_2)_m$ pyrazyl, $(\text{CH}_2)_m$ oxazolyl, $(\text{CH}_2)_m$ isoxazolyl, $(\text{CH}_2)_m$ thiazolyl, $(\text{CH}_2)_m$ isothiazolyl, $(\text{CH}_2)_m$ phenyl, $(\text{CH}_2)_m$ pyrrolyl, $(\text{CH}_2)_m$ pyridyl, or $(\text{CH}_2)_m$ pyrimidyl, which cycloalkyl and said cycloalkenyl groups optionally contain one or two heteroatoms selected independently from O, N, and S, and which are optionally substituted as described below; wherein m is zero, 1, or 2, Ar is a 5- to 10-member mono- or bicyclic aromatic group, optionally containing 1 – 4 ring heteroatoms selected

independently from N, O, and S; or R₁ and R₂, together with the ring carbon atoms to which they are attached, form a 5- or 6- member fused ring, which ring may be saturated, unsaturated, or aromatic, which optionally contains one or two heteroatoms selected independently from O, N, and S, and which is optionally substituted as described below; R₃ and R₄ are, independently, H, CN, halogen, CF₃, OCF₃, OC₁-C₃ alkyl, or C₁-C₆ alkyl; X = O or S; Y is O or S; q = 1 or zero; R₅ is C₁-C₆ alkyl, (CHR₆)_wC₃-C₆ cycloalkyl, (CHR₆)_wCH₂C₃-C₆ cycloalkyl, CH₂(CHR₆)_wC₃-C₆ cycloalkyl, CR₆=CH-C₃-C₆ cycloalkyl, CH=CR₆-C₃-C₆ cycloalkyl, (CHR₆)_wC₅-C₆ cycloalkenyl, CH₂(CHR₆)_wC₅-C₆ cycloalkenyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, Ar, (CHR₆)_wAr, CH₂(CHR₆)_wAr, or (CHR₆)_wCH₂Ar, wherein w = zero, 1, 2, or 3, Ar is a 5- to 10- member mono- or bicyclic aromatic group, optionally containing 1 – 4 ring heteroatoms selected independently from N, O, and S; R₆ is H or C₁-C₃ alkyl; wherein all cycloalkyl and cycloalkenyl groups optionally contain one or two ring heteroatoms selected independently from N, O, and S; wherein all alkyl, cycloalkyl, alkenyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, alkynyl, aryl, and heteroaryl groups in R₁, R₂, R₃, R₄, R₅, R₆, and Ar are optionally substituted with one or two substituents selected independently from C₁-C₃ alkyl, OCF₃, halogen, CN, OH, OMe, OEt, CN, CH₂F, and trifluoromethyl; and wherein, additionally, all cycloalkyl and heterocycloalkyl groups are optionally substituted with a carbonyl group, and halogen designates Cl, F, Br or I and the terms alkyl refers to branch or unbranched alkyl groups and pharmaceutically acceptable salts, solvates, and esters thereof. Such compounds are potassium channel activators or modulators.

Essentially all combinations of the several variables in formula I are embraced by this invention.

In another embodiment, this invention provides a composition comprising a pharmaceutically acceptable carrier or diluent and at least one of the following: a pharmaceutically effective amount of a compound of formula I, a pharmaceutically acceptable salt of a compound of formula I, a pharmaceutically acceptable solvate of a compound of formula I, and a pharmaceutically acceptable ester of a compound of formula I.

In yet another embodiment, this invention provides a pediatric pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent, a syrup for pediatric use, and at least one of the following: a pharmaceutically effective amount of a compound of formula I, a pharmaceutically acceptable salt of a compound of formula I, a pharmaceutically acceptable ester of a compound of formula I, and a pharmaceutically acceptable solvate of a compound of formula I.

In yet another embodiment, this invention provides a chewable tablet, suitable for pediatric pharmaceutical use, comprising a pharmaceutically acceptable carrier or diluent, and at least one of the following: a pharmaceutically effective amount of a compound of formula I, a pharmaceutically acceptable salt of a compound of formula I, a pharmaceutically acceptable
5 solvate of a compound of formula I, and a pharmaceutically acceptable ester of a compound of formula I.

In yet another embodiment, this invention provides a method of preventing or treating a disease or disorder which is affected by activation voltage-gated potassium channels, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of
10 formula IA or a salt or ester or solvate thereof.

This invention includes all tautomers and salts of compounds of this invention. This invention also includes all compounds of this invention wherein one or more atoms are replaced by a radioactive isotope thereof.

This invention provides compounds of formula I above wherein the group $\text{NH-C(=X)-(Y)}_q\text{-R}_5$ is each of the following: NHC(=O)R_5 , NHC(=O)OR_5 , NHC(=S)R_5 , NHC(=S)SR_5 , NHC(=S)OR_5 ,
15 and NHC(=O)SR_5 .

Thus, in one embodiment, this invention provides a compound of formula I, wherein $\text{NH-C(=X)-(Y)}_q\text{-R}_5$ is NHC(=O)R_5 .

In another embodiment, this invention provides a compound of formula I, wherein $\text{NH-C(=X)-(Y)}_q\text{-R}_5$ is NHC(=S)R_5 .
20

In another embodiment, this invention provides a compound of formula I, wherein $\text{NH-C(=X)-(Y)}_q\text{-R}_5$ is NHC(=S)SR_5 .


In another embodiment, this invention provides a compound of formula I, wherein $\text{NH-C(=X)-(Y)}_q\text{-R}_5$ is each NHC(=O)OR_5 .

25 In another embodiment, this invention provides a compound of formula I, wherein $\text{NH-C(=X)-(Y)}_q\text{-R}_5$ is NHC(=S)OR_5 .

In another embodiment, this invention provides a compound of formula I, wherein $\text{NH-C(=X)-(Y)}_q\text{-R}_5$ is NHC(=O)SR_5 .


In another generic embodiment, this invention provides a compound of formula I, wherein q is zero and R_5 is C_1 - C_6 alkyl, or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula I wherein R_1 is located as shown below:




5

In another embodiment, R_1 is located as shown below:



In another embodiment, R_1 is located as shown below



10

In another embodiment, this invention provides a compound of formula I, wherein R_2 is H.

In another embodiment, this invention provides a compound of formula I, wherein R_2 is halogen.

In another embodiment, this invention provides a compound of formula I, wherein R_2 is Cl or F.

15 In another embodiment, this invention provides a compound of formula I, wherein R_2 is trifluoromethyl.

In another embodiment, this invention provides a compound of formula I, wherein R_3 and R_4 are, independently, H, Cl, methyl, ethyl, trifluoromethyl, or methoxy.

In another embodiment, this invention provides a compound of formula I, wherein q is zero and R_3 and R_4 are Cl, ethyl, methoxy, or methyl.

- 5 In another embodiment, this invention provides a compound of formula I, wherein q is zero and R_3 and R_4 are both methyl.

In another embodiment, this invention provides a compound of formula I, wherein R_3 and R_4 are, independently, H, Cl, ethyl, methoxy, or methyl.

- 10 In another embodiment, this invention provides a compound of formula I, wherein R_3 and R_4 are, independently, H, Cl, ethyl, methoxy, or methyl.

In another embodiment, this invention provides a compound of formula I, wherein R_3 and R_4 are, independently, H, Cl, ethyl, or methyl.

In another embodiment, this invention provides a compound of formula I, wherein q is zero, and R_5 is C_1 - C_6 alkyl, or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

- 15 In another embodiment, this invention provides a compound of formula I, wherein q is 1; Y is O; and R_5 is C_1 - C_6 alkyl, or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein q is 1; Y is S; and R_5 is C_1 - C_6 alkyl, or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

- 20 In another embodiment, this invention provides a compound of formula I, wherein R_2 is H and R_5 is C_1 - C_6 alkyl, or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein R_2 is H and R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$.

- 25 In another embodiment, this invention provides a compound of formula I, wherein R_2 is H and R_5 is $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula I, wherein R_2 is H and R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl or $CH=CR_6$ - C_3 - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein R₃ and R₄ are H, Cl, ethyl, or methyl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ is Cl or F; and R₃ and R₄ are H, Cl, ethyl, or methyl.

- 5 In another embodiment, this invention provides a compound of formula I, wherein R₁ is Cl or F; R₃ and R₄ are H, Cl, ethyl, or methyl; and R₅ is C₁-C₆ alkyl, or (CHR₆)_wC₃-C₆ cycloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ is phenyl, optionally substituted.

- 10 In another embodiment, this invention provides a compound of formula I, wherein R₁ is phenyl, optionally substituted, and R₅ is C₁-C₆ alkyl, or (CHR₆)_wC₃-C₆ cycloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ is NH-C₁-C₆ alkyl, N(C₁-C₆ alkyl)-C₁-C₆ alkyl, C(=O)NH-C₁-C₆ alkyl, NH-C(=O)C₁-C₆ alkyl; O-C₁-C₆ alkyl, C(=O)-C₁-C₆ alkyl, C(=O)-OC₁-C₆ alkyl, or OC(=O)C₁-C₆ alkyl; and R₅ is C₁-C₆ alkyl, or (CHR₆)_wC₃-C₆ cycloalkyl.

- 15 In another embodiment, this invention provides a compound of formula I, wherein R₁ is NH-C₁-C₆ alkyl, N(C₁-C₆ alkyl)-C₁-C₆ alkyl, C(=O)NH-C₁-C₆ alkyl, or NH-C(=O)C₁-C₆ alkyl.

In yet another embodiment, this invention provides a compound of formula I, wherein R₁ is C(=O)OC₁-C₆ alkyl, OC(=O)C₁-C₆ alkyl, or OC₁-C₆ alkyl.

- 20 In another specific embodiment, this invention provides a compound of formula I, wherein R₁ is H, methyl, methoxy, or halogen, and R₂ is methyl or ethyl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ is H, methyl, methoxy, or halogen, and R₂ is phenyl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ is H, methyl, methoxy, or halogen, and R₂ is F.

- 25 In another embodiment, this invention provides a compound of formula I, wherein R₁ is methoxy, methoxymethyl, ethoxymethyl, or methoxyethyl.

In another embodiment, this invention provides a compound of formula I, wherein R_1 is methoxy, methoxymethyl, ethoxymethyl, or methoxyethyl; R_2 is H, methyl, or halogen; and R_3 is methyl or Cl.

5 In another embodiment, this invention provides a compound of formula I, wherein R_1 is phenyl, optionally substituted, and R_2 is H, methyl, methoxy, or halogen.

In another embodiment, this invention provides a compound of formula I, wherein R_1 is CF_3 or C_1 - C_3 alkyl, and R_2 is H, methyl, methoxy, or halogen.

In another embodiment, this invention provides a compound of formula I, wherein R_1 is methoxy, and R_2 is H, methyl, methoxy, or halogen.

10 In another embodiment, this invention provides a compound of formula I, wherein R_1 is 2-dimethylamino ethyl, and R_2 is H, methyl, methoxy, or halogen.


In another embodiment, this invention provides a compound of formula I, wherein q is zero, R_2 is H, methyl, methoxy, or halogen, R_1 is phenyl, optionally substituted; and R_3 and R_4 are H, Cl, ethyl, or methyl.

15 In another embodiment, this invention provides a compound of formula I, wherein q is zero, R_2 is H, methyl, methoxy, or halogen; R_1 is CF_3 or C_1 - C_3 alkyl; and R_3 and R_4 are H, Cl, ethyl, or methyl.

In another embodiment, this invention provides a compound of formula I, wherein q is zero, R_2 is H, methyl, methoxy, or halogen; R_1 is F; and R_3 and R_4 are H, Cl, ethyl, or methyl.


20 In another embodiment, this invention provides a compound of formula I, wherein q is zero; R_1 is Br; R_2 is H, methyl, methoxy, or halogen; and R_3 and R_4 are H, Cl, ethyl, or methyl.

In another embodiment, the invention provides a compound of formula IA-1 below.



wherein R_1 is selected from the group consisting of H, halogen, CN, CH_2CN , CHF_2 , CF_3 , $\text{C}_1\text{-C}_6$ alkyl, OCH_3 , $(\text{C}=\text{O})\text{OCH}_3$, $\text{O}(\text{C}=\text{O})\text{CH}_3$, OCF_3 , $(\text{CH}_2)_m\text{C}_3\text{-C}_6$ cycloalkyl, phenyl, and pyridyl; R_2 is selected from the group consisting of H, F, OCH_3 , CH_3 , and CF_3 ; R_3 and R_4 are independently, selected from the group consisting of H, F, Cl, CF_3 , OCF_3 , $\text{OC}_1\text{-C}_3$ alkyl, and $\text{C}_1\text{-C}_3$ alkyl; and R_5 is selected from the group consisting of $\text{C}_1\text{-C}_6$ alkyl, $(\text{CHR}_6)_w\text{C}_3\text{-C}_6$ cycloalkyl, $(\text{CHR}_6)_w\text{CH}_2\text{C}_3\text{-C}_6$ cycloalkyl, $\text{CH}_2(\text{CHR}_6)_w\text{C}_3\text{-C}_6$ cycloalkyl, $\text{CR}_6=\text{CH-C}_3\text{-C}_6$ cycloalkyl, $\text{CH}=\text{CR}_6\text{-C}_3\text{-C}_6$ cycloalkyl, $(\text{CHR}_6)_w\text{C}_5\text{-C}_6$ cycloalkenyl, $\text{CH}_2(\text{CHR}_6)_w\text{C}_5\text{-C}_6$ cycloalkenyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, Ar, $(\text{CHR}_6)_w\text{Ar}$, $\text{CH}_2(\text{CHR}_6)_w\text{Ar}$, and $(\text{CHR}_6)_w\text{CH}_2\text{Ar}$, wherein $w = 0 - 3$, Ar is selected from the group consisting of phenyl, furyl, pyrrolyl, oxazolyl, thiazolyl, thienyl, and pyridyl; and R_6 is $\text{C}_1\text{-C}_3$ alkyl; wherein all alkyl, cycloalkyl, aryl, and heteroaryl groups in R_1 , R_2 , R_3 , R_4 , R_5 , and R_6 , and Ar are optionally substituted with one or two substituents selected independently from $\text{C}_1\text{-C}_3$ alkyl, halogen, OCH_3 , OCH_2CH_3 , CN, and CF_3 .


In another embodiment, this invention provides a compound of formula IA-2 below. In another embodiment, this invention provides a compound of formula IA-3 below,



wherein R_1 is selected from the group consisting of H, halogen, CN, CH_2CN , CHF_2 , CF_3 , $\text{C}_1\text{-C}_6$ alkyl, OCH_3 , $(\text{C}=\text{O})\text{OCH}_3$, $\text{O}(\text{C}=\text{O})\text{CH}_3$, OCF_3 , $(\text{CH}_2)_m\text{C}_3\text{-C}_6$ cycloalkyl, phenyl, and pyridyl; R_2 is selected from the group consisting of H, F, OCH_3 , CH_3 , and CF_3 ; R_3 and R_4 are independently, selected from the group consisting of H, F, Cl, CF_3 , OCF_3 , $\text{OC}_1\text{-C}_3$ alkyl, and $\text{C}_1\text{-C}_3$ alkyl; and R_5 is selected from the group consisting of $\text{C}_1\text{-C}_6$ alkyl, $(\text{CHR}_6)_w\text{C}_3\text{-C}_6$ cycloalkyl, $(\text{CHR}_6)_w\text{CH}_2\text{C}_3\text{-C}_6$ cycloalkyl, $\text{CH}_2(\text{CHR}_6)_w\text{C}_3\text{-C}_6$ cycloalkyl, $\text{CR}_6=\text{CH-C}_3\text{-C}_6$ cycloalkyl, $\text{CH}=\text{CR}_6\text{-C}_3\text{-C}_6$ cycloalkyl, $(\text{CHR}_6)_w\text{C}_5\text{-C}_6$ cycloalkenyl, $\text{CH}_2(\text{CHR}_6)_w\text{C}_5\text{-C}_6$ cycloalkenyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, Ar, $(\text{CHR}_6)_w\text{Ar}$, $\text{CH}_2(\text{CHR}_6)_w\text{Ar}$, and $(\text{CHR}_6)_w\text{CH}_2\text{Ar}$, wherein $w = 0 - 3$, Ar is selected from the group consisting of phenyl, furyl, pyrrolyl, oxazolyl, thiazolyl, thienyl, and pyridyl; and R_6 is $\text{C}_1\text{-C}_3$ alkyl; wherein all alkyl, cycloalkyl, aryl, and heteroaryl groups in R_1 , R_2 , R_3 , R_4 , R_5 , and R_6 , and Ar are optionally substituted with one or two substituents selected independently from $\text{C}_1\text{-C}_3$ alkyl, halogen, OCH_3 , OCH_2CH_3 , CN, and CF_3 .


10

In still another embodiment, this invention provides a compound of formula IA-3 below.




In yet another embodiment, this invention provides a compound of formula IB-1 below.

5




In another specific embodiment, this invention provides a compound of formula IB-2 below.



10

In another specific embodiment, this invention provides a compound of formula IB-3 below.



In formulas IA-3, IB-1, IB-2 and IB-3, R₁, R₂, R₃, R₄ and R₅ are defined as in formula I.

In another more specific embodiment, this invention provides a compound of formula IA-1, formula IA-2, or formula IA-3, wherein R_2 is H, alkyl, or halogen; and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

5 In another specific embodiment, this invention provides a compound of formula IA-1, formula IA-2, or formula IA-3, wherein R_1 is $(CH_2)_mC_3$ - C_6 cycloalkyl; R_2 is H, alkyl, or halogen; and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

10 In another specific embodiment, this invention provides a compound of formula IA-1, formula IA-2, or formula IA-3, wherein R_1 is methoxy, methoxymethyl, or methoxyethyl; R_2 is H, alkyl, or halogen; and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

In yet another specific embodiment, this invention provides a compound of formula IA-1, wherein R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

15 In yet another specific embodiment, this invention provides a compound of formula IA-2, wherein R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

In yet another specific embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$.

20 In yet another specific embodiment, this invention provides a compound of formula IA-2, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$.

In yet another embodiment, this invention provides a compound of formula IA-3, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$.

25 In yet another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In yet another embodiment, this invention provides a compound of formula IA-2, wherein R_5 is $CR_6=CH-C_3-C_6$ cycloalkyl, $CH=CR_6-C_3-C_6$ cycloalkyl, $(CHR_6)_wC_5-C_6$ cycloalkenyl, $CH_2(CHR_6)_wC_5-C_6$ cycloalkenyl, C_2-C_6 alkenyl, or C_2-C_6 alkynyl.

5 In yet another embodiment, this invention provides a compound of formula IA-1, wherein R_2 is H; R_3 is methyl; and R_5 is C_1-C_6 alkyl, $(CHR_6)_wC_3-C_6$ cycloalkyl, $(CHR_6)_wCH_2C_3-C_6$ cycloalkyl, or $CH_2(CHR_6)_wC_3-C_6$ cycloalkyl.

In yet another embodiment, this invention provides a compound of formula IA-2, wherein R_2 is H; R_3 is methyl; and R_5 is C_1-C_6 alkyl, $(CHR_6)_wC_3-C_6$ cycloalkyl, $(CHR_6)_wCH_2C_3-C_6$ cycloalkyl, or $CH_2(CHR_6)_wC_3-C_6$ cycloalkyl.

10 In yet another embodiment, this invention provides a compound of formula IA-3, wherein R_2 is H; R_3 is methyl; and R_5 is C_1-C_6 alkyl, $(CHR_6)_wC_3-C_6$ cycloalkyl, $(CHR_6)_wCH_2C_3-C_6$ cycloalkyl, or $CH_2(CHR_6)_wC_3-C_6$ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is H, F, Cl, Br, methoxy, methoxymethyl, ethoxymethyl, methoxyethyl, or trifluoromethyl; R_3 is methyl;
15 and R_5 is C_4-C_6 alkyl, $(CHR_6)_wCH_2C_5-C_6$ cycloalkyl, or $CH_2(CHR_6)_wC_5-C_6$ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R_1 is H, F, Cl, Br, methoxy, methoxymethyl, ethoxymethyl, methoxyethyl, or trifluoromethyl; R_3 is methyl; and R_5 is C_4-C_6 alkyl, $(CHR_6)_wCH_2C_5-C_6$ cycloalkyl, or $CH_2(CHR_6)_wC_5-C_6$ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is C_4-C_6 alkyl, $(CHR_6)_wC_5-C_6$ cycloalkyl, or $CH_2(CHR_6)_wC_5-C_6$ cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, methoxymethyl, ethoxymethyl, methoxyethyl, or trifluoromethyl.
20

In another embodiment, this invention provides a compound of formula IA-2, wherein R_5 is C_4-C_6 alkyl, $(CHR_6)_wC_5-C_6$ cycloalkyl, or $CH_2(CHR_6)_wC_5-C_6$ cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, methoxymethyl, ethoxymethyl, methoxyethyl, or trifluoromethyl.

25 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is H, F, Cl, Br, methoxy, methoxymethyl, ethoxymethyl, methoxyethyl, or trifluoromethyl; R_2 is H, methyl, or F; R_3 is methyl; R_4 is methyl or Cl; and R_5 is C_4-C_6 alkyl, $(CHR_6)_wC_5-C_6$ cycloalkyl, or $CH_2(CHR_6)_wC_5-C_6$ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R_1 is H, F, Cl, Br, methoxy, methoxymethyl, ethoxymethyl, methoxyethyl, or trifluoromethyl; R_2 is H, methyl, or F; R_3 is methyl; R_4 is methyl or Cl; and R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl.

- 5 In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is H, F, Cl, Br, methoxy, methoxymethyl, ethoxymethyl, methoxyethyl, or trifluoromethyl; R_2 is H, methyl, or F; R_3 is methyl; R_4 is methyl or Cl; and R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl.

- 10 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is $(CH_2)_mC_3$ - C_6 cycloalkyl, C_3 - C_6 cycloalkenyl, or $(CH_2)_mC_3$ - C_6 cycloalkenyl; and R_3 is methyl or Cl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is $(CH_2)_mC_3$ - C_6 cycloalkyl, C_3 - C_6 cycloalkenyl, or $(CH_2)_mC_3$ - C_6 cycloalkenyl.

- 15 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is methoxy, methoxymethyl, ethoxymethyl; or methoxyethyl; R_2 is H or F; R_3 is methyl; R_4 is methyl or Cl; and R_5 is $(CHR_6)_wC_5$ - C_6 cycloalkenyl or $(CHR_6)_wAr$.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is phenyl, optionally substituted.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is methyl, halomethyl, ethyl, or haloethyl.

- 20 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is C_1 - C_4 alkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is methyl or ethyl.

- 25 In another embodiment, this invention provides a compound of formula IA-1, wherein R_2 is fluoro, R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R₂ is 4-fluoro, R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

5 In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is (CH₂)_mimidazolyl, (CH₂)_mpyrazyl, (CH₂)_m furyl, (CH₂)_m thienyl, (CH₂)_moxazolyl, (CH₂)_misoxazolyl, (CH₂)_mthiazolyl, (CH₂)_misothiazolyl, (CH₂)_mphenyl, (CH₂)_mpyrrolyl, (CH₂)_mpyridyl, or (CH₂)_mpyrimidyl; and R₂ is H.

10 In another embodiment, this invention provides a compound of formula IA-2, wherein R₁ is (CH₂)_mimidazolyl, (CH₂)_mpyrazyl, (CH₂)_m furyl, (CH₂)_m thienyl, (CH₂)_moxazolyl, (CH₂)_misoxazolyl, (CH₂)_mthiazolyl, (CH₂)_misothiazolyl, (CH₂)_mphenyl, (CH₂)_mpyrrolyl, (CH₂)_mpyridyl, or (CH₂)_mpyrimidyl; R₂ is H and R₅ is pyridyl or phenyl, optionally substituted.

In another embodiment, this invention provides a compound of formula IA-2, wherein R₂ is CF₃ or C₁-C₃ alkyl; R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

15 In another embodiment, this invention provides a compound of formula IA-2, wherein R₃ and R₄ are methyl or trifluoromethyl; R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

20 In another embodiment, this invention provides a compound of formula IA-2, wherein R₂ is methoxy or ethoxy; and R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R₂ is phenyl, optionally substituted; R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

25 In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is (CH₂)_mimidazolyl, (CH₂)_mpyrazyl, (CH₂)_m furyl, (CH₂)_m thienyl, (CH₂)_moxazolyl, (CH₂)_misoxazolyl, (CH₂)_mthiazolyl, (CH₂)_misothiazolyl, (CH₂)_mphenyl, (CH₂)_mpyrrolyl, (CH₂)_mpyridyl, or (CH₂)_mpyrimidyl; R₂ is H; and R₅ is pyridyl or phenyl, optionally substituted.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_2 is CF_3 or C_1 - C_3 alkyl; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

5 In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are methyl or trifluoromethyl; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_2 is methoxy or ethoxy; and R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

10 In another embodiment, this invention provides a compound of formula IA-1, wherein R_2 is phenyl, optionally substituted; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R' is 4-phenyl, optionally substituted; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R' is CF_3 or C_1 - C_3 alkyl; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

20 In another embodiment, this invention provides a compound of formula IA-1, wherein R' is 4-methyl or 4-ethyl; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R' is methoxy or ethoxy, R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

25 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_2 is H, F, or methyl; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R' is H.

- 5 In another embodiment, this invention provides a compound of formula IA-2, wherein R' is halogen.

In another embodiment, this invention provides a compound of formula IA-2, wherein R' is F.

In another embodiment, this invention provides a compound of formula IA-2, wherein R' is methyl or ethyl.

- 10 In another embodiment, this invention provides a compound of formula IA-2, wherein R' is methyl or ethyl; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R' is halogen; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is

- 15 H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R' is H; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

- In another embodiment, this invention provides a compound of formula IA-2, wherein R' is 1-phenyl, optionally substituted.

In another embodiment, this invention provides a compound of formula IA-2, wherein R' is 4-phenyl, optionally substituted.

In another embodiment, this invention provides a compound of formula IA-2, wherein R' is CF_3 or C_1 - C_3 alkyl.

- 25 In another embodiment, this invention provides a compound of formula IA-3, wherein R' is H; R_5 is C_4 - C_6 alkyl, $(CHR_6)_wC_5$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_5$ - C_6 cycloalkyl; and R_1 is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R' is F; R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

5 In another embodiment, this invention provides a compound of formula IA-3, wherein R' is 1-phenyl, optionally substituted; R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R' is 4-phenyl, optionally substituted; R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

10 In another embodiment, this invention provides a compound of formula IA-3, wherein R₂ is CF₃ or C₁-C₃ alkyl; R₅ is C₄-C₆ alkyl, (CHR₆)_wC₅-C₆ cycloalkyl, or CH₂(CHR₆)_wC₅-C₆ cycloalkyl; and R₁ is H, F, Cl, Br, methoxy, or trifluoromethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ and R₂, are, independently, H, CN, F, Cl, Br, CH₂CN, OCH₃, CH₂OCH₃, CH₂CH₂OCH₃, CH₂OCH₂CH₃,
15 CH₂F, CHF₂, CF₃, CF₂CF₃, or C₁-C₆ alkyl and R₅ is C₁-C₆ alkyl or CH₂(CHR₆)_wC₃-C₆ cycloalkyl, wherein w = 0, 1, or 2.

In another embodiment, this invention provides a compound of formula IA-1, R₁ is H, CN, F, Cl, Br, CH₂CN, OCH₃, CH₂OCH₃, CH₂CH₂OCH₃, CH₂OCH₂CH₃, CH₂F, CHF₂, CF₃,

CF₂CF₃, or C₁-C₆ alkyl; R₂ is H, F, Cl, or methyl; R₃ is methyl or chloro; and R₅ is C₁-C₆ alkyl or
20 CH₂(CHR₆)_wC₃-C₆ cycloalkyl, wherein R₆ is H or methyl and w = 1 or 2.

In another embodiment, this invention provides a compound of formula IA, wherein R₅ is Ar, (CHR₆)_wAr, CH₂(CHR₆)_wAr, or (CHR₆)_wCH₂Ar.

In another embodiment, this invention provides a compound of formula IA-2, wherein R₅ is Ar, (CHR₆)_wAr, CH₂(CHR₆)_wAr, or (CHR₆)_wCH₂Ar₁.

25 In another embodiment, this invention provides a compound of formula IA-3, wherein R₅ is Ar, (CHR₆)_wAr, CH₂(CHR₆)_wAr, or (CHR₆)_wCH₂Ar.

In another embodiment, this invention provides a compound of formula I, wherein R_5 is $CR_6=CH-C_3-C_6$ cycloalkyl, $CH=CR_6-C_3-C_6$ cycloalkyl, $(CHR_6)_wC_5-C_6$ cycloalkenyl, $CH_2(CHR_6)_wC_5-C_6$ cycloalkenyl, C_2-C_6 alkenyl, or C_2-C_6 alkynyl.

5 In another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is $CR_6=CH-C_3-C_6$ cycloalkyl, $CH=CR_6-C_3-C_6$ cycloalkyl, $(CHR_6)_wC_5-C_6$ cycloalkenyl, $CH_2(CHR_6)_wC_5-C_6$ cycloalkenyl, C_2-C_6 alkenyl, or C_2-C_6 alkynyl.

In another embodiment, this invention provides a compound of formula I, wherein R_5 is haloalkyl.

10 In another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is haloalkyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R_5 is haloalkyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_5 is haloalkyl.

15 In another embodiment, this invention provides a compound of formula IB-1, wherein R_5 is haloalkyl.

In another embodiment, this invention provides a compound of formula IB-2, wherein R_5 is haloalkyl.

20 In another embodiment, this invention provides a compound of formula IB-3, wherein R_5 is haloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein R_5 is methoxy alkyl.

In another embodiment, this invention provides a compound of formula I, wherein R_5 is cyano alkyl.

25 In another embodiment, the invention provides a compound of formula I, wherein R_5 is CH_2 -cycloalkyl or CH_2CH_2 -cycloalkyl.

In another embodiment, the invention provides a compound of formula IA-1, wherein R₅ is CH₂-cycloalkyl or CH₂CH₂-cycloalkyl.

In another embodiment, the invention provides a compound of formula IA-2, wherein R₅ is CH₂-cycloalkyl or CH₂CH₂-cycloalkyl.

- 5 In another embodiment, the invention provides a compound of formula IA-3, wherein R₅ is CH₂-cycloalkyl or CH₂CH₂-cycloalkyl.

In another embodiment, the invention provides a compound of formula IB-1, wherein R₅ is CH₂-cycloalkyl or CH₂CH₂-cycloalkyl.

- 10 In another embodiment, the invention provides a compound of formula IB-2, wherein R₅ is CH₂-cycloalkyl or CH₂CH₂-cycloalkyl.

In another embodiment, the invention provides a compound of formula IB-3, wherein R₅ is CH₂-cycloalkyl or CH₂CH₂-cycloalkyl.

In another embodiment, the invention provides a compound of formula IA-1, wherein R₅ is CH₂-C₅-C₆ cycloalkyl or CH₂CH₂-C₅-C₆ cycloalkyl.

- 15 In another embodiment, the invention provides a compound of formula IA-2, wherein R₅ is CH₂-C₅-C₆ cycloalkyl or CH₂CH₂-C₅-C₆ cycloalkyl.

In another embodiment, the invention provides a compound of formula IA-3, wherein R₅ is CH₂-C₅-C₆ cycloalkyl or CH₂CH₂-C₅-C₆ cycloalkyl.

- 20 In another embodiment, the invention provides a compound of formula IB-1, wherein R₅ is CH₂-C₅-C₆ cycloalkyl or CH₂CH₂-C₅-C₆ cycloalkyl.

In another embodiment, the invention provides a compound of formula IB-2, wherein R₅ is CH₂-C₅-C₆ cycloalkyl or CH₂CH₂-C₅-C₆ cycloalkyl.

In another embodiment, the invention provides a compound of formula IB-3, wherein R₅ is CH₂-C₅-C₆ cycloalkyl or CH₂CH₂-C₅-C₆ cycloalkyl.

- 25 In another embodiment, the invention provides a compound of formula IA-1, wherein R₅ is C₅-C₆ alkyl or CH₂C₅-C₆ alkyl.

In another embodiment, the invention provides a compound of formula IA-2, wherein R₅ is C₅-C₆ alkyl or CH₂C₅-C₆ alkyl.

In another embodiment, the invention provides a compound of formula IA-3, wherein R₅ is C₅-C₆ alkyl or CH₂C₅-C₆ alkyl.

- 5 In another embodiment, the invention provides a compound of formula IB-1, wherein R₅ is C₅-C₆ alkyl or CH₂C₅-C₆ alkyl.

In another embodiment, the invention provides a compound of formula IB-2, wherein R₅ is C₅-C₆ alkyl or CH₂C₅-C₆ alkyl.

- 10 In another embodiment, the invention provides a compound of formula IB-3, wherein R₅ is C₅-C₆ alkyl or CH₂C₅-C₆ alkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₃ and R₄ are chloro, methoxy, or methyl and R₅ is CH₂-cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₃ and R₄ are chloro, methoxy, or methyl and R₅ is haloalkyl, hydroxyalkyl, or methoxyalkyl.

- 15 In another embodiment, this invention provides a compound of formula IA-1, wherein R₃ and R₄ are methyl and R₅ is C₅-C₆ alkyl or methoxy alkyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R₃ and R₄ are methyl and R₅ is C₅-C₆ alkyl or chloroalkyl.

- 20 In another embodiment, this invention provides a compound of formula IA-2, wherein R₃ and R₄ are trifluoromethyl and R₅ is methoxyalkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₃ and R₄ are both methyl and R₅ is 2-(2-halo cyclopentyl) ethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₃ and R₄ are both methyl and R₅ is 2-(2-furyl) ethyl.

- 25 In another embodiment, this invention provides a compound of formula IA-1, wherein R₃ and R₄ are both methyl and R₅ is 2-(2-tetrahydrofuryl) ethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are both methyl and R_5 is 2-phenyl ethyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are both methyl and R_5 is 3-phenyl propyl.

- 5 In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are both methyl and R_5 is 2-phenyl propyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl; and R_1 is halogen.

- 10 In another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl; R_2 is H or halogen; and R_1 is halogen.

- In another embodiment, this invention provides a compound of formula IA-2, wherein R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl; R_2 is H or halogen; and R_1 is halogen.
- 15

In another embodiment, this invention provides a compound of formula IA-3, wherein R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl; R_2 is H or halogen; and R_1 is halogen.

- In another embodiment, this invention provides a compound of formula IA-2, wherein R_5 is C_1 - C_6 alkyl or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl; R_2 is hydrogen; and R_1 is halogen.
- 20

In another embodiment, this invention provides a compound of formula IA-3, wherein R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl; R_2 is hydrogen; and R_1 is halogen.

- In another embodiment, this invention provides a compound of formula I, wherein R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.
- 25

In another embodiment, this invention provides a compound of formula IA, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar_1$.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is haloalkyl; R_2 is H or F; R_3 and R_4 are Cl, methoxy, or methyl; and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA, wherein R_1 is C_1 - C_3 alkyl, halogen, or haloalkyl; R_2 is H or F; R_3 and R_4 are H, methyl, or Cl; and R_5 is CH_2CR_6 - C_3 - C_6 cycloalkyl, $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_4 - C_6 alkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is C_1 - C_3 alkyl, halogen, or haloalkyl; R_2 is H or F; R_3 and R_4 are H, methyl, or Cl; and R_5 is CH_2CR_6 - C_3 - C_6 cycloalkyl, $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_4 - C_6 alkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is C_1 - C_3 alkyl, halogen, or haloalkyl; R_2 is H or F; R_3 and R_4 are H, methyl, or Cl; and R_5 is CH_2CR_6 - C_3 - C_6 cycloalkyl, $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_4 - C_6 alkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is C_1 - C_3 alkyl, halogen, or haloalkyl; R_2 is H or F; R_3 and R_4 are H, methyl, or Cl; and R_5 is CH_2CR_6 - C_3 - C_6 cycloalkyl, or C_2 - C_6 alkyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is C_1 - C_3 alkyl, halogen, or haloalkyl; R_2 is H or F; R_3 and R_4 are H, methyl, or Cl; and R_5 is CH_2CR_6 - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is halogen or haloalkyl; R_2 is H or F; and R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is halogen or haloalkyl; R_2 is H or F; and R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is halogen or haloalkyl; R_2 is H or F; R_3 and R_4 are Cl, methoxy, or methyl; and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

5 In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is halogen or haloalkyl; R_2 is H or F; and R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

10 In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is methyl, fluoro, or fluoroalkyl; R_2 is H or F; and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is Cl, F, or CF_3 ; R_2 is H or F; R' is H or CH_3 ; and R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

15 In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is Cl, F, or CF_3 ; R_2 is H or F; R' is H or CH_3 ; and R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$.

20 In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are H, methyl, or Cl; and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are H, methyl, or Cl; and R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

25 In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are H, methyl, or Cl; and wherein R_1 and R_2 , on adjacent carbons, form a six-membered ring.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are H, methyl, or Cl; wherein R_5 is C_2 - C_6 alkyl, CH_2 - C_5 - C_6 cycloalkyl, CH_2CH_2 - C_5 - C_6 cycloalkyl, $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, or C_2 - C_6 alkenyl; and wherein R_1 and R_2 , are on adjacent carbons, and are both other than H.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are H, methyl, or Cl; wherein R_5 is C_2 - C_6 alkyl, CH_2 - C_5 - C_6 cycloalkyl, CH_2CH_2 - C_5 - C_6 cycloalkyl, $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, or C_2 - C_6 alkenyl; and wherein R_1 and R_2 , on adjacent carbons, are both halogen.

- 5 In another embodiment, this invention provides a compound of formula IA-1, wherein R_3 and R_4 are H, methyl, or Cl; wherein R_5 is C_2 - C_6 alkyl, CH_2 - C_5 - C_6 cycloalkyl, CH_2CH_2 - C_5 - C_6 cycloalkyl, $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, or C_2 - C_6 alkenyl; and wherein R_1 and R_2 , on adjacent carbons, are both fluorine.

- 10 In an embodiment, this invention provides a compound of formula IA-1, wherein R' is F, methyl, or H; R_3 and R_4 are H, methyl, or Cl; and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R' is F, methyl, or H; R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

- 15 In another embodiment, this invention provides a compound of formula IA-1, wherein R' is halogen and R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar_1$.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 and R_2 are on adjacent carbon atoms and are both other than H.

- 20 In an embodiment, this invention provides a compound of formula IA-1, wherein R_1 and R_2 , on adjacent carbon atoms are, independently trifluoromethyl or halogen; and wherein R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

- 25 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is halogen and R_2 is H, or R_1 and R_2 , on adjacent carbon atoms are, independently trifluoromethyl or halogen; and wherein R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar_1$.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is halogen or trifluoromethyl and R_2 is H, or R_1 and R_2 , on adjacent carbon atoms are, independently trifluoromethyl or halogen; and wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$.

In another embodiment, this invention provides a compound of formula IA, wherein X is S, $q = 1$, Y is O, and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

10 In another embodiment, this invention provides a compound of formula IA, wherein X is S, $q = 1$, Y is O, and R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA, wherein X is S, $q = 1$, Y is O, and R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$.

15 In another embodiment, this invention provides a compound of formula IA, wherein X is S, $q =$ zero, and R_5 is C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, or $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA, wherein X is S, $q =$ zero, and R_5 is $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl.

In another embodiment, this invention provides a compound of formula IA-2 wherein R_5 is C_1 - C_6 alkyl or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_5 is C_1 - C_6 alkyl or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

25 In another embodiment, this invention provides a compound of formula IA-2, wherein R_1 is halogen or trifluoromethyl and R_2 is H or R_1 and R_2 , on adjacent carbon atoms, are, independently, halogen or trifluoromethyl; and R_5 is C_1 - C_6 alkyl or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is halogen or trifluoromethyl and R_2 is H or R_1 and R_2 , on adjacent carbon atoms, are, independently, halogen or trifluoromethyl; and R_5 is C_1 - C_6 alkyl or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

5 In another embodiment, this invention provides a compound of formula IA-2, wherein R_1 and R_2 are, independently, methyl, methoxy, trifluoromethyl, F, Cl, or H; and R_5 is C_1 - C_6 alkyl or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 and R_2 are, independently, methyl, methoxy, trifluoromethyl, F, Cl, or H; R' is H; and R_5 is C_1 - C_6 alkyl or $(CHR_6)_wC_3$ - C_6 cycloalkyl.

10 In another embodiment, this invention provides a compound of formula IA-1 or IA-2 or IA-3, wherein R_1 is halogen, C_1 - C_6 alkyl, mono-halo C_1 - C_6 alkyl, CN, di-halo C_1 - C_6 alkyl, CF_3 , CN, or O- C_1 - C_6 alkyl; R' is methyl or ethyl; and R_5 is C_5 - C_6 alkyl or CH_2 - C_3 - C_6 cycloalkyl.

15 In another embodiment, this invention provides a compound of formula IA-1 or IA-2 or IA-3, wherein R_1 is H, halogen, cyano, CF_3 , or methoxy, R_2 is H, F, or methyl, R' is H, halogen, methyl, ethyl, or methoxy, and R_5 is C_5 - C_6 alkyl or CH_2 - C_3 - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein R_1 is F, Cl, or CF_3 ; R_2 is H; and R' is halogen, methyl, ethyl, or methoxy; R_3 and R_4 are H, methyl, or Cl; and R_5 is C_5 - C_6 alkyl or CH_2 - C_3 - C_6 cycloalkyl.

20 In another embodiment, this invention provides a compound of formula I, wherein R_1 is halogen or CF_3 ; R_2 is H, F, or methyl, R' is phenyl; R_3 and R_4 are H, methyl, or Cl; and R_5 is C_5 - C_6 alkyl or CH_2 - C_5 - C_6 cycloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein R_1 is halogen or CF_3 ; R_2 is H, F, or methyl, R' is halophenyl; R_3 and R_4 are H, methyl, or Cl; and R_5 is C_5 - C_6 alkyl or CH_2 - C_5 - C_6 cycloalkyl.

25 In another embodiment, this invention provides a compound of formula I, wherein R_1 is NH_2 , NH - C_1 - C_6 alkyl; N (C_1 - C_6 alkyl)- C_1 - C_6 alkyl, $NHC(=O)$ - C_1 - C_6 alkyl, $C(=O)N(CH_3)_2$, $C(=O)N(Et)_2$, $C(=O)NH_2$, $C(=O)NH$ - C_1 - C_6 alkyl, SO_2NH_2 , $NHSO_2$ - C_1 - C_6 alkyl.

In another embodiment, this invention provides a compound of formula I wherein R_1 is NH_2 , $NH-C_1-C_6$ alkyl; or $N(C_1-C_6$ alkyl)- C_1-C_6 alkyl; and R_2 is H or halogen.

In another embodiment, this invention provides a compound of formula I wherein R_1 is $NHC(=O)C_1-C_6$ alkyl, $C(=O)N(CH_3)_2$, $C(=O)N(Et)_2$, $C(=O)NH_2$, or $C(=O)NH-C_1-C_6$ alkyl.

- 5 In another embodiment, this invention provides a compound of formula 1 wherein R_1 is $NHC(=O)C_1-C_6$ alkyl, $C(=O)N(CH_3)_2$, $C(=O)N(Et)_2$, $C(=O)NH_2$, or $C(=O)NH-C_1-C_6$ alkyl.

In another embodiment, this invention provides a compound of formula 1 wherein R_1 is SO_2NH_2 or $NHSO_2-C_1-C_6$ alkyl.

- 10 In another embodiment, this invention provides a compound of formula IA-2 wherein R_1 is SO_2NH_2 or $NHSO_2-C_1-C_6$ alkyl.

In another embodiment, this invention provides a compound of formula I, wherein R_1 is $C(=O)OC_1-C_6$ alkyl, $OC(=O)C_1-C_6$ alkyl, OC_1-C_6 alkyl, or SC_1-C_6 alkyl.

In another embodiment, this invention provides a compound of formula I, wherein R_1 is $(CH_2)_mC_3-C_6$ cycloalkenyl, C_2-C_6 alkenyl, or C_2-C_6 alkynyl.

- 15 In another embodiment, this invention provides a compound of formula I, wherein R_1 is CH_2OCH_3 , $CH_2OCH_2CH_3$, OC_1-C_6 alkyl, or SC_1-C_6 alkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is $C(=O)OC_1-C_6$ alkyl, $OC(=O)C_1-C_6$ alkyl, OC_1-C_6 alkyl, or SC_1-C_6 alkyl.

- 20 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is CH_2OCH_3 , $CH_2OCH_2CH_3$, OC_1-C_6 alkyl, or SC_1-C_6 alkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is $C(=O)OC_1-C_6$ alkyl, $OC(=O)C_1-C_6$ alkyl, OC_1-C_6 alkyl, or SC_1-C_6 alkyl; R_2 is H, F, or methyl, R' is halogen or methyl; and R_5 is C_5-C_6 alkyl or $CH_2-C_5-C_6$ cycloalkyl.

- 25 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is NH_2 , $NH-C_1-C_6$ alkyl; or $N(C_1-C_6$ alkyl)- C_1-C_6 alkyl; R_2 is H, F, or methyl, R' is halogen or methyl; and R_5 is C_5-C_6 alkyl or $CH_2-C_5-C_6$ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is $\text{NHC(=O)C}_1\text{-C}_6$ alkyl, $\text{C(=O)N(CH}_3)_2$, C(=O)N(Et)_2 , C(=O)NH_2 , $\text{C(=O)NH-C}_1\text{-C}_6$ alkyl, SO_2NH_2 , or $\text{NHSO}_2\text{-C}_1\text{-C}_6$ alkyl; R_2 is H, F, or methyl, R' is halogen or methyl; and R_5 is $\text{C}_5\text{-C}_6$ alkyl or $\text{CH}_2\text{-C}_5\text{-C}_6$ cycloalkyl.

- 5 In another embodiment, this invention provides a compound of formula I, wherein R_1 is $\text{C}_2\text{-C}_6$ alkynyl, optionally substituted.

In another embodiment, this invention provides a compound of formula I, wherein R_1 and R_2 form a fused, nitrogen-containing ring.

- 10 In another embodiment, this invention provides a compound of formula I, wherein R_1 and R_2 form a fused, oxygen-containing ring.

In another embodiment, this invention provides a compound of formula I, wherein R_1 and R_2 form a fused thiazolo or isothiazolo group.

In another embodiment, this invention provides a compound of formula I, wherein R_1 and R_2 form a fused cyclopentane, optionally substituted.

- 15 In another embodiment, this invention provides a compound of formula I, wherein R_1 and R_2 form a fused cyclohexane, optionally substituted.

In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R_1 and R_2 form a fused, nitrogen-containing ring.

- 20 In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R_1 and R_2 form a fused, oxygen-containing ring.

In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R_1 and R_2 form a fused thiazolo or isothiazolo group.

In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R_1 and R_2 form a fused cyclopentane, optionally substituted.

- 25 In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R_1 and R_2 form a fused cyclohexane, optionally substituted.

In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R₁ and R₂ form a fused, nitrogen-containing ring; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R₁ and R₂ form a fused, oxygen-containing ring; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

- 5 In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R₁ and R₂ form a fused thiazolo or isothiazolo group; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R₁ and R₂ form a fused cyclopentane, optionally substituted; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

- 10 In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R₁ and R₂ form a fused cyclohexane, optionally substituted; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

- In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is halogen; R₂ is H, F, or methyl, R' is halogen or methyl; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.
- 15

In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is halogen; R₂ is H, F, or methyl, R' is 2-(dimethylamino) ethyl; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

- In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is halogen; R₂ is H, halogen, or methyl, R' is H; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.
- 20

In another embodiment, this invention provides a compound of formula IA-2, wherein R₁ is halogen; R₂ is H or methyl, R' is halogen or methyl; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

- In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is Br, Cl, F or methyl; R₂ is H or F and R₅ is *t*-butyl or cyclopentylmethyl.
- 25

In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is trifluoromethyl; R₂ is H or methyl, R' is halogen or methyl; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-2, wherein R₁ wherein R₁ is trifluoromethyl; R₂ is H or methyl, R' is halogen or methyl; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

5 In another embodiment, this invention provides a compound of formula IA-3, wherein R₁ wherein R₁ is trifluoromethyl; R₂ is H or methyl, R' is halogen or methyl; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

In another embodiment, this invention provides a compound of formula IA-1 or IA-2, wherein R₁ wherein R₁ is trifluoromethyl; R₂ is H or methyl, R' is halogen or methyl; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

10 In another embodiment, this invention provides a compound of formula IA-2, wherein R₁ is trifluoromethyl; R₂ is F; R' is halogen or methyl; and R₅ is C₅-C₆ alkyl or CH₂-C₅-C₆ cycloalkyl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ or R₅ is CH₂Ar or CH₂CH₂-Ar, wherein Ar is phenyl, pyridyl, pyrrolyl, imidazolyl, oxazolyl, or thiazolyl.

15 In another embodiment, this invention provides a compound of formula I, wherein R₁ is F.

In another embodiment, this invention provides a compound of formula I, wherein R₁ is Cl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ is Br.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is F.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is Cl.

20 In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is Br.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is F and R₂ is H, OCH₃, or F.

In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is F; R₃ and R₄ are both methyl; and R' is H.

25 In another embodiment, this invention provides a compound of formula IA-1, wherein R₁ is CF₃; R₃ and R₄ are both methyl; and R' is H.

In another embodiment, this invention provides a compound of formula I, wherein R_1 and R_2 are both F.

In another embodiment, this invention provides a compound of formula I, wherein R_1 is mono-, di-, or tri-halomethyl.

- 5 In another embodiment, this invention provides a compound of formula I, wherein R_1 is CH_2F , CHF_2 , or CF_3 .

In another embodiment, this invention provides a compound of formula I, wherein R_1 is CH_2Cl .

In another embodiment, this invention provides a compound of formula I, wherein R_1 is CH_2Br .

- 10 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 and R_2 are both F; R_3 and R_4 are both methyl; and R' is H.

In another embodiment, this invention provides a compound of formula IA-2, wherein R_1 is F.

In another embodiment, this invention provides a compound of formula IA-2, wherein R_1 and R_2 are both F.

In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 is F.

- 15 In another embodiment, this invention provides a compound of formula IA-3, wherein R_1 and R_2 are both F.

In another embodiment, this invention provides a compound of formula I, wherein R_1 or R_5 is CH_2Ar or $\text{CH}_2\text{CH}_2\text{-Ar}$, wherein Ar is isoxazolyl or isothiazolyl.

- 20 In another embodiment, this invention provides a compound of formula I, wherein R_1 or R_5 is CH_2Ar or $\text{CH}_2\text{CH}_2\text{-Ar}$, wherein Ar is quinolyl or isoquinolyl.

In another embodiment, this invention provides a compound of formula I, wherein R_1 or R_5 is CH_2Ar or $\text{CH}_2\text{CH}_2\text{-Ar}$, wherein Ar is pyrimidyl or purinyl.

In another embodiment, this invention provides a compound of formula I, wherein R_1 or R_5 is CH_2Ar or $\text{CH}_2\text{CH}_2\text{-Ar}$, wherein Ar is indolyl, isoindolyl, or benzimidazolyl.

- 25 In an embodiment, this invention provides a compound of formula I, wherein R_1 or R_5 is CH_2Ar or $\text{CH}_2\text{CH}_2\text{-Ar}$, wherein Ar is halo phenyl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ or R₅ is CH₂Ar or CH₂CH₂-Ar, wherein Ar is dihalophenyl or dihalopyridyl.

In another embodiment, invention provides a compound of formula I, wherein R₁ or R₅ is CH₂Ar or CH₂CH₂-Ar, wherein Ar is mono- or di-halothienyl, mono- or di-halofuryl, mono- or di-
5 halobenzothienyl, or mono- or di-halobenzofuryl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ or R₅ is CH₂Ar or CH₂CH₂-Ar, wherein Ar is *o*-, *m*-, or *p*-xylyl or *o*-, *m*-, or *p*-anisyl.

In another embodiment, this invention provides a compound of formula I, wherein R₁ or R₅ is CH₂Ar or CH₂CH₂-Ar, wherein Ar is *m*- or *p*-cyanophenyl or *m*- or *p*-cyanomethyl phenyl.

In another embodiment, this invention provides a compound of formula I, in which R_3 and R_4 are halogen, CF_3 , or C_1 - C_3 alkyl and R_5 is C_1 - C_6 alkyl, wherein the alkyl group is substituted with one or two groups selected, independently, from OH, OMe, OEt, F, CF_3 , Cl, or CN.

5 In another embodiment, this invention provides a compound of formula I, in which R_3 and R_4 are halogen, CF_3 , OCF_3 , C_1 - C_3 alkyl, or OC_1 - C_3 alkyl, and R_5 is $(CH_2)_wC_3$ - C_6 cycloalkyl, wherein w is 1 or 2, wherein the cycloalkyl group is substituted with Me, OH, OMe, OEt, F, CF_3 , Cl, or CN.

In an embodiment, this invention provides a compound of formula IA-1, in which R_3 and R_4 are halogen, CF_3 , or C_1 - C_3 alkyl, and R_5 is $(CH_2)_wC_5$ - C_6 cycloalkyl, optionally substituted, or $(CH_2)_wC_5$ - C_6 heterocycloalkyl, optionally substituted.

10 In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is CH_2 phenyl or CH_2CH_2 -phenyl.

In another embodiment, this invention provides a compound of formula IA-1, wherein R_1 is Ar, CH_2 Ar or CH_2CH_2 -Ar, wherein Ar is 3,5-dichlorophenyl or 3,5-difluorophenyl.

15 In an embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_w$ Ar, $CH_2(CHR_6)_w$ Ar, or $(CHR_6)_wCH_2$ Ar, wherein Ar is phenyl or pyridyl; R_3 and R_4 are H or C_1 - C_6 alkyl, unsubstituted or substituted with one or two groups selected from OH, OMe; and R_6 is CN, CH_2 CN, or halogen.

20 In another embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_w$ Ar, $CH_2(CHR_6)_w$ Ar, or $(CHR_6)_wCH_2$ Ar, wherein Ar is phenyl or pyridyl; and R_1 is F, CH_2 F, CHF_2 , CF_3 , or CF_2CF_3 .

In an embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_w$ Ar, $CH_2(CHR_6)_w$ Ar, or $(CHR_6)_wCH_2$ Ar, wherein Ar is phenyl or pyridyl, and R_1 is OC_1 - C_6 alkyl or $C(=O)C_1$ - C_6 alkyl.

In an embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl, and R_1 is $C(=O)OC_1-C_6$ alkyl or $OC(=O)C_1-C_6$ alkyl.

5 In an embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl, R_1 is C_2-C_6 alkenyl or C_2-C_6 alkynyl, q is 1, and X and Y are both O.

In an embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, Ar is phenyl or pyridyl, and R_1 is SC_1-C_6 alkyl.

10 In an embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl, R_3 and R_4 are H, Cl, methoxy, or C_1-C_3 alkyl, and R_1 is C_1-C_6 alkyl.

In an embodiment, this invention provides a compound of formula IA-2, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl; R_3 and R_4 are H, Cl, methoxy, or C_1-C_2 alkyl, unsubstituted or substituted with one or two groups selected
15 from OH, OMe; and R_1 is CN, CH_2CN , or halogen.

In another embodiment, this invention provides a compound of formula IA-2, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl; and R_1 is F, CH_2F , CHF_2 , CF_3 , or CF_2CF_3 .

20 In an embodiment, this invention provides a compound of formula IA-1, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl, and R_1 is OC_1-C_6 alkyl or $C(=O)C_1-C_6$ alkyl.

In an embodiment, this invention provides a compound of formula IA-2, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl, and R_1 is OC_1-C_6 alkyl or $C(=O)C_1-C_6$ alkyl.

In an embodiment, this invention provides a compound of formula IA-3, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl, and R_1 is OC_1-C_6 alkyl or $C(=O)C_1-C_6$ alkyl.

5 In an embodiment, this invention provides a compound of formula IA-3, wherein R' is phenyl or methoxy, R_2 is H, and R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl, and R_1 is $C(=O)OC_1-C_6$ alkyl or $OC(=O)C_1-C_6$ alkyl.

In an embodiment, this invention provides a compound of formula IA-2, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, Ar is phenyl or pyridyl, and R_1 is SC_1-C_6 alkyl.

10 In an embodiment, this invention provides a compound of formula IA-2, wherein R_5 is Ar, $(CHR_6)_wAr$, $CH_2(CHR_6)_wAr$, or $(CHR_6)_wCH_2Ar$, wherein Ar is phenyl or pyridyl, R_3 and R_4 are H or C_1-C_3 alkyl, and R_1 is C_1-C_6 alkyl.

15 In another embodiment, this invention provides a method of treating or preventing a disease, disorder, or condition that is affected by modulation of potassium ion channels in a patient comprising administration of a compound of formula I in an amount of up to about 2000 mg per day.

In another embodiment, this invention provides a method of treating or preventing a disease, disorder, or condition that is affected by modulation of potassium ion channels in a patient comprising administration of a compound of formula I in an amount of from about 10 mg to about 2000 mg per day.

20 In another embodiment, this invention provides a method of treating or preventing a disease, disorder, or condition that is affected by modulation of potassium ion channels in a patient comprising administration of a compound of formula IA-1 in an amount of up to about 2000 mg per day.

In another embodiment, this invention provides a method of treating or preventing a seizure disorder in a patient comprising administration of a compound of formula I in an amount of up to about 2000 mg per day.

5 In another embodiment, this invention provides a method of treating or preventing a seizure disorder in a patient comprising administration of a compound of formula I in an amount of from about 10 mg per day to about 2000 mg per day.

In another embodiment, this invention provides a method of treating or preventing a seizure disorder in a patient comprising administration of a compound of formula I in an amount of from about 300 mg per day to about 2000 mg per day.

10 In another embodiment, this invention provides a method of treating or preventing a seizure disorder in a patient comprising administration of a compound of formula I in an amount of from about 300 mg per day to about 1200 mg per day.

In another embodiment, this invention provides a method of treating or preventing a seizure disorder in a patient comprising administration of a compound of formula IA-1 in an amount of
15 up to 2000 mg per day.

In another embodiment, this invention provides a method of treating or preventing a seizure disorder in a patient comprising administration of a compound of formula IA-1 in an amount of from about 10 mg per day to about 2000 mg per day.

In another embodiment, this invention provides a method of treating or preventing a seizure
20 disorder in a patient comprising administration of a compound of formula IA-1 in an amount of from about 300 mg per day to about 2000 mg per day.

In another embodiment, this invention provides a method of treating or preventing a seizure disorder in a patient comprising administration of a compound of formula IA-1 in an amount of from about 300 mg per day to about 1200 mg per day.

Detailed Description of Invention

As provided by this invention, compounds of formula IA are designed for oral or intravenous dosing of up to 2000 mg per day. Yet the high activities of many of these compounds indicate that dosing of less than 1200 mg per day – the current anticipated dosing level of retigabine in adults – is possible. Thus, this invention comprises tablets, capsules, solutions, and suspensions of compounds of formula IA which are formulated for oral administration. Similarly, solutions and suspensions suitable for oral pediatric administration, comprising, in addition to compounds of formula IA, a syrup such as sorbitol or propylene glycol, among many other examples, are also contemplated. ally, solutions and suspensions comprising, in addition to compounds of formula IA, a syrup such as sorbitol or propylene glycol, along with colorants and flavorings suitable for oral pediatric administration, are also contemplated. Additionally, both chewable and non-chewable tablets comprising compounds of formula IA, along with pharmaceutically acceptable tableting agents and other pharmaceutically acceptable carriers and excipients, are also contemplated. As used herein, the term pharmaceutically acceptable carrier comprises such excipients, binders, lubricants, tableting agents, disintegrants, preservatives, anti-oxidants, flavours and colourants as are typically used in the art of formulation of pharmaceuticals. Examples of such agents include – but are not limited to – starch, calcium carbonate, dibasic calcium phosphate, dicalcium phosphate, microcrystalline cellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose lactose, polyethylene glycols, polysorbates, glycols, safflower oil, sesame oil, soybean oil, and Povidone. Additionally, disintegrants such as sodium starch glycolate; lubricants such as magnesium stearate, stearic acid, and SiO₂; and solubility enhancers such as cyclodextrins, among a great many other examples for each group, are contemplated. Such materials and the methods of using them are well known in the pharmaceutical art. Additional examples are provided in Kibbe, *Handbook of Pharmaceutical Excipients*, London, Pharmaceutical Press, 2000.

As used herein, the term "pharmaceutically acceptable acid salts" refers to acid addition salts formed from acids which provide non-toxic anions. The pharmaceutically acceptable anions include, but are not limited to, acetate, aspartate, benzoate, bicarbonate, carbonate, bisulfate, sulfate, chloride, bromide, benzene sulfonate, methyl sulfonate, phosphate, acid phosphate, lactate, maleate, malate, malonate, fumarate, lactate, tartrate, borate, camsylate, citrate, edisylate, esylate, formate, fumarate, gluceptate, glucuronate, gluconate oxalate, palmitate, pamoate, saccharate, stearate, succinate, tartrate, tosylate and trifluoroacetate salts, among a great many other examples. Hemi-salts, including but not limited to hemi-sulfate salts, are likewise

contemplated.

For a review on suitable salts, see "Handbook of Pharmaceutical Salts: Properties, Selection, and Use" by Stahl and Wermuth (Wiley-VCH, Weinheim, Germany, 2002).

As is well known, pharmaceutically acceptable salts of compounds of formula I may be prepared by reaction of a compound of formula I with the desired acid; by removal of a protecting group from a suitable precursor of the compound of formula I or by ring-opening a suitable cyclic precursor, for example, a lactone or lactam, using the desired acid or base; and by conversion of one salt of the compound of formula I to another by reaction with an appropriate acid or base or by passage through an appropriate ion-exchange column.

As used herein, the term "pharmaceutically acceptable solvate" refers to describe a molecular complex comprising the compound of the invention and a stoichiometric amount of one or more pharmaceutically acceptable solvent molecules, including but not limited to water and ethanol. Thus, the term solvate includes a hydrate as one example and an ethanolate as another example.

The compounds of the present invention may possess one or more asymmetric carbons. Accordingly, any optical isomers as separated and any mixtures including racemic mixtures are embraced by the scope of the present invention. Resolution of racemic mixtures can be accomplished by methods known to those skilled in the art.


The compounds of the present invention may also exist as geometric isomers and in different tautomeric forms. Those geometric isomers and tautomeric forms are included within the scope of the present invention.

As used herein, modulation of ion channels refers to activating the ion channels, to affecting the kinetics of opening and closing of the ion channels, or to causing any change in the channel open probability of the ion channels.

Preparation of Compounds

5 **The Preparation of compounds of formula I is outlined in Schemes I and II.**

Scheme I



While in the above Scheme I nitrophenol is exemplified, a great many substituted nitrophenols are known and are therefore embraced by Scheme I. Thus, for example, compounds of Formula I wherein R₃ and R₄ are CF₃, OCF₃, OC₁-C₃ alkyl or C₁-C₃ alkyl or C₁-C₃ alkyl can be prepared from the appropriate substituted nitrophenol starting material. Likewise the exemplified fluorobenzylchloride of Scheme I can be replaced by a number of other substituted benzyl chloride. For example, if the compounds of Formula I wherein R₁ and R₂ are independently selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl or OC₁-C₆ alkyl can be prepared from the appropriate substituted benzyl chloride starting material. It is to be understood that while the benzyl chloride of Scheme I is substituted by a single R group, a second R group which would correspond to the R₁ and R₂ groups of Formula I is included.

While Scheme I exemplifies the preparation of a substituted carbamic acid ethyl ester wherein the ethyl group would correspond to the R₅ position of Formula I, it is to be understood that substitution of the appropriate reagent for the diethyl pyrocarbonate of Scheme I would yield the corresponding moiety which is defined by R₅.

1-nitro-4-(4-fluorobenzyl)oxy)-benzene

A mixture of 4-nitrophenol (1.39g, 10mmol), potassium carbonate (1.38g, 10mmol) and 4-fluorobenzyl bromide (1.89g, 10mmol) in 20 ml of anhydrous DMF was stirred at 100 °C for 2 days. After cooling to room temperature, the reaction mixture was poured into 200 ml of ice-water with stirring. The solid was filtered and washed with water and dried to give 2.37g (96%)

of 1-nitro-4-(4-fluorobenzyloxy)-benzene as yellow solids.

The following compounds were synthesized by the same procedure:

- 5
 1-Nitro-2-fluoro-4-(4-fluorobenzyloxy)-benzene, yellow solids, 96%
 1-Nitro-2-methyl-4-(4-fluorobenzyloxy)-benzene, yellow solids, 97%
 1-Nitro-2-trifluoromethyl-4-(4-fluorobenzyloxy)-benzene, yellow solids, 95%
 1-Nitro-3-fluoro-4-(4-fluorobenzyloxy)-benzene, yellow solids, 91%
 1-Nitro-3-chloro-4-(4-fluorobenzyloxy)-benzene, yellow solids, 85%
 1-Nitro-3-methoxy-4-(4-fluorobenzyloxy)-benzene, yellow solids, 97%


4-(4-Fluoro-benzyloxy)-aniline

- 10 1-Nitro-4-(4-fluorobenzyloxy)-benzene (2.37g) was dissolved in 200 ml of methanol and catalytic amount of Raney Ni was added. The mixture was stirred at room temperature under atmospheric pressure in a hydrogen atmosphere for 3 hours. After filtering off Raney Ni over Celite and washing with methanol, the obtained filtrate was concentrated under reduced pressure to give a solid product 4-(4-fluoro-benzyloxy)-aniline, which is pure enough for next step.

- 15 The following compounds were synthesized by the same procedure:

- 20
 2-Fluoro-4-(4-Fluoro-benzyloxy)-aniline
 2-Trifluoromethyl-4-(4-Fluoro-benzyloxy)-aniline
 2-Methyl-4-(4-Fluoro-benzyloxy)-aniline
 3-Fluoro-4-(4-Fluoro-benzyloxy)-aniline
 3 -Chloro-4-(4-Fluoro-benzyloxy)-aniline
 3 -Methoxy-4-(4-Fluoro-benzyloxy)-aniline.

[4-(4-Fluoro-benzyloxy)-phenyl] carbamic acid ethyl ester




- 25 4-(4-Fluoro-benzyloxy)-aniline (0.22g, 1mmol) was dissolved in 8 ml of anhydrous ethanol and diethyl pyrocarbonate (0.20g, 1.2mmol) was added dropwise at room temperature. The resulting mixture was stirred at room temperature for 4 hours, then stored at -20 °C overnight. The crystals was filtered and washed with cold ethanol to give pure product as crystal solids (88mg, 30%). The filtrate was concentrated to dryness under reduced pressure and the residue was purified by silica gel column to give another batch of pure products. ¹H NMR (DMSO-*d*₆, 300MHz): 8.9.38 (brs, 1H, exchangeable with D₂O, NH), 7.46 (dd, J=5.7 and 8.1Hz, 2H), 7.32 (d,
- 30

J=8.7Hz, 2H), 7.18 (t, J=8.7Hz, 2H), 6.90 (d, J=8.7Hz, 2H), 5.00 (s, 2H), 4.07 (q, J=7.2Hz, 2H), 1.20 (t, J=7.2Hz, 3H). MS: 288 (M-1).


The following compounds were synthesized by the same procedure:

[2-Methyl-4-(4-fluoro-benzyloxy)-phenyl]-carbamic acid ethyl ester




¹H NMR (DMSO-*d*₆, 300MHz): δ 8.62 (brs, 1H, exchangeable with D₂O, NH), 7.46 (dd, J=5.7 and 8.1Hz, 2H), 7.19 (t, J=8.7Hz, 2H), 7.12 (d, J=8.7Hz, 1H), 6.83 (d, J=2.7Hz, 1H), 6.76 (dd, J=2.7 and 8.7Hz, 1H), 5.02 (s, 2H), 4.04 (q, J=7.2Hz, 2H), 2.12 (s, 3H), 1.19 (t, J=7.2Hz, 3H). MS: 302 (M-1).

[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-carbamic acid ethyl ester




¹H NMR (DMSO-*d*₆, 300MHz): δ 8.97 (brs, 1H, exchangeable with D₂O, NH), 7.47 (dd, J=5.7 and 8.1Hz, 2H), 7.37 (t, J=8.7Hz, 1H), 7.20 (t, J=8.7Hz, 2H), 6.93 (dd, J=2.7 and 12.3Hz, 1H), 6.78 (dd, J=2.7 and 8.7Hz, 1H), 5.05 (s, 2H), 4.05 (q, J=7.2Hz, 2H), 1.19 (t, J=7.2Hz, 3H). MS: 307 (M-1).


[2-Trifluoromethyl-4-(4-fluoro-benzyloxy)-phenyl] carbamic acid ethyl ester




¹H NMR (DMSO-*d*₆, 300MHz): δ 8.83 (brs, 1H, exchangeable with D₂O, NH), 7.50 (dd, J=5.7 and 8.1Hz, 2H), 7.33 (d, J=8.7Hz, 1H), 7.21 (t, J=8.7Hz, 2H), 7.26 (dd, J=2.7 and 8.7Hz, 1H), 7.24 (d, J=2.7Hz, 1H), 5.15 (s, 2H), 4.03 (q, J=7.2Hz, 2H), 1.16 (t, J=7.2Hz, 3H). MS: 356 (M-1).

[3-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-carbamic acid ethyl ester

¹H NMR (DMSO-d₆, 300MHz): δ 9.61 (brs, 1H, exchangeable with D₂O, NH), 7.46 (dd, J=5.7 and 8.1Hz, 2H), 7.36 (dd, J=1.8 and 13.8Hz, 1H), 7.20 (t, J=8.7Hz, 2H), 7.16 (d, J=9.0Hz, 1H),
 5 7.09 (dd, J=1.8 and 9.0Hz, 1H), 5.06 (s, 2H), 4.08 (q, J=7.2Hz, 2H), 1.21 (t, J=7.2Hz, 3H). MS: 306 (M-1).


[3-Chloro-4-(4-fluoro-benzyloxy)-phenyl]-carbamic acid ethyl ester

¹H NMR (DMSO-d₆, 300MHz): δ 9.59 (brs, 1H, exchangeable with D₂O, NH), 7.55 (d, J=1.8Hz,
 10 1H), 7.48 (dd, J=5.7 and 8.7Hz, 2H), 7.29 (dd, J=1.8 and 8.7Hz, 1H), 7.21 (t, J=8.7Hz, 2H), 7.15 (d, J=8.7Hz, 1H), 5.10 (s, 2H), 4.08 (q, J=7.2Hz, 2H), 1.21 (t, J=7.2Hz, 3H). MS: 322 (M-1).

[3-Methoxy-4-(4-fluoro-benzyloxy)-phenyl]-carbamic acid ethyl ester

¹H NMR (DMSO-d₆, 300MHz): δ 9.40 (brs, 1H, exchangeable with D₂O, NH), 7.45 (dd, J=5.7 and 8.4Hz, 2H), 7.18 (t, J=8.7Hz, 2H), 7.18 (m, 1H), 6.89 (m, 2H), 4.96 (s, 2H), 4.08 (q,
 15 J=7.2Hz, 2H), 1.21 (t, J=7.2Hz, 3H). MS: 318 (M-1).

Scheme II



While in the above scheme II 3, 5-dimethylphenol is exemplified, a great many substituted phenols are known and are embraced by this scheme. Thus for example, compounds of formula I wherein R_3 and R_4 are CF_3 , OCF_3 , OC_1-C_3 alkyl or C_1-C_3 alkyl can be prepared from the appropriate substituted or disubstituted phenol starting material to arrive at the corresponding nitrophenol. Likewise the exemplified (fluoro)-benzyl chloride of the above scheme can be replaced by a number of other substituted benzyl chlorides. For example if compound of formula I wherein R_1 and R_2 are independently selected C_1-C_3 alkyl, C_1-C_3 cycloalkyl or OC_1-C_6 alkyl can be prepared from the appropriate substituted benzyl chloride starting material. It is to be understood that while the benzyl chloride of Scheme I is substituted by a single R group, a second R group which would correspond to the R_1 and R_2 groups of formula I is included.

In the above scheme II, R_1 of scheme II would correspond to R_5 of formula I and R would correspond to R_1 and R_2 of formula I.

15 3,5-Dimethyl-4-nitrophenol

[3,5-Dimethyl-4-nitrophenol was synthesized by the reference procedures (US 4564640)] 750 ml of Concentrated hydrochloric acid was added to a solution of 3,5-dimethylphenol (80.6g) in 750 ml of 95% ethanol. The mixture was cooled to 0 °C in an ice/methanol bath. While maintaining the temperature of the reaction mixture below 5 °C, a solution of $NaNO_2$ (69.0g) in 150 ml of water was added dropwise to the reaction mixture. The mixture was stirred at 0 °C for more than an hour and then poured into 9 liters of water. The aqueous mixture was filtered to give a yellow solid which was recrystallized from hot methanol and filtered to give 71.45g of 3,5-dimethyl-4-nitrosophenol as a yellow solid. Mp. 180-181 °C.

A mixture of 3,5-dimethyl-4-nitrosophenol (70.63g) from above and $(NH_4)_6Mo_7O_{24} \cdot 4H_2O$ (2.83g) in 770 ml of glacial acetic acid was warmed to 100 °C. 30% H_2O_2 (84ml) was added to the mixture in 10 ml portions until an exothermic reaction was observed. The reaction mixture

was then stirred and the remainder of the H₂O₂ solution added in small portions. The reaction mixture was heated and stirred until a clear dark red solution results. A yellow-orange solid precipitated from the solution after stirring for another 20-30 minutes. The reaction mixture was stirred overnight and filtered to give a small amount of a yellow solid and a clear dark red filtrate. The red filtrate was concentrated in vacuo and partitioned between water and ether. The aqueous layer was washed with ether and the combined ether extracts washed with 10% sodium carbonate until the aqueous layer becomes basic. The ether extract was dried over anhydrous sodium sulfate, filtered, concentrated and allowed to cool overnight. After cooling in an ice bath the mixture was filtered to give 49.5g of 3,5-dimethyl-4-nitrophenol as a yellow-green solid. Mp. 106-108°C.

5-(4-Fluoro-benzyloxy)-2-nitro-meta-xylene

A mixture of 3,5-dimethyl-4-nitrophenol (0.97g, 5.79mmol), 4-fluorobenzyl chloride (1.26g, 8.69mmol) and anhydrous potassium carbonate (1.24g, 9.0mmol) in 50 ml of acetone was stirred under reflux for 22 hours. TLC showed this reaction is complete. The reaction mixture was cooled to room temperature and filtered and washed with acetone. The filtrate was evaporated to dryness to give the crude product, which was used for next step without further purification.

The following compounds were synthesized by the same procedure:

5-(4-Chloro-benzyloxy)-2-nitro-meta-xylene
5-(4-Bromo-benzyloxy)-2-nitro-meta-xylene
5-(4-Methyl-benzyloxy)-2-nitro-meta-xylene
5-(4-Trifluoromethyl-benzyloxy)-2-nitro-meta-xylene
5-(2-Trifluoromethyl -benzyloxy)-2-nitro-meta-xylene
5-(3-Trifluoromethyl -benzyloxy)-2-nitro-meta-xylene
5-(3-Chloro-benzyloxy)-2-nitro-meta-xylene
5-(3-Fluoro-benzyloxy)-2-nitro-meta-xylene
5-(2,4-Difluoro-benzyloxy)-2-nitro-meta-xylene
5-(3,4-Difluoro-benzyloxy)-2-nitro-meta-xylene


4-(4-Fluoro-benzyloxy)-2,6-dimethylaniline

1-Nitro-4-(4-fluorobenzyloxy)-benzene (0.5g) was dissolved in 60 ml of methanol and catalytic amount of Raney Ni was added. The mixture was stirred at room temperature under atmospheric pressure in a hydrogen atmosphere for 3 hours. After filtering off Raney Ni over Celite and washing with methanol, the obtained filtrate was concentrated under reduced pressure to give a solid product 4-(4-fluoro-benzyloxy)-2,6-dimethylaniline, which is pure enough for next step.

The following compounds were synthesized by the same procedure:

- 5
10
- 4-(4-Chloro-benzyloxy)-2,6-dimethylaniline
 - 4-(4-Bromo-benzyloxy)-2,6-dimethylaniline
 - 4-(4-Methyl-benzyloxy)-2,6-dimethylaniline
 - 4-(3,4-Difluoro-benzyloxy)-2,6-dimethylaniline
 - 4-(2,4-Difluoro-benzyloxy)-2,6-dimethylaniline
 - 4-(4-Trifluoromethyl-benzyloxy)-2,6-dimethylaniline
 - 4-(3-Trifluoromethyl-benzyloxy)-2,6-dimethylaniline
 - 4-(2-Trifluoromethyl-benzyloxy)-2,6-dimethylaniline
 - 4-(3-Fluoro-benzyloxy)-2,6-dimethylaniline
 - 4-(3-Chloro-benzyloxy)-2,6-dimethylaniline


N-[4-(4-Fluoro-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide



- 15 To a solution of 4-(4-fluoro-benzyloxy)-2,6-dimethylaniline(0.20g, 0.82mmol) from above and triethylamine (125mg, 1.24mmol) in anhydrous methylene chloride (20 ml) was added dropwise *tert*-butylacetyl chloride (135mg, 1 mmol) at 0 °C. The reaction mixture was stirred at room temperature for 18 hours. Water was added to the reaction mixture, and the mixture was washed with saturated brine and dried over sodium sulfate. The solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ISCO, hexane/EtOAc, 0-40%, 40 min) and recrystallized from hexane/EtOAc (5:1) to give 230 mg (82%) of the white solids. ¹H NMR (DMSO-d₆, 400MHz): δ 8.92 (brs, 1H, exchangeable with D₂O, NH), 7.46 (dd, J=4.7 and 6.4Hz, 2H), 7.19 (t, J=7.0Hz, 2H), 6.70 (s, 2H), 5.02 (s, 2H), 2.16 (s, 2H), 2.08 (s, 6H), 1.03 (s, 9H). MS: 344 (M+1).
- 20


The following compounds were synthesized by the same procedure:

N-[4-(4-Chloro-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide




- 5 ^1H NMR (DMSO- d_6 , 400MHz): δ 8.94 (brs, 1H, exchangeable with D_2O , NH), 7.46 (s, 4H), 6.72 (s, 2H), 5.07 (s, 2H), 2.18 (s, 2H), 2.10 (s, 6H), 1.05 (s, 9H). MS: 360 (M+1).

N-[4-(4-Bromo-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide




- 10 ^1H NMR (DMSO- d_6 , 400MHz): δ 8.94 (brs, 1H, exchangeable with D_2O , NH), 7.59 (d, $J=8.0\text{Hz}$, 2H), 7.39 (d, $J=8.0\text{Hz}$, 2H), 6.71 (s, 2H), 5.05 (s, 2H), 2.18 (s, 2H), 2.10 (s, 6H), 1.05 (s, 9H). MS: 404 (M+1).

N-[4-(4-Methyl-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide




15


- ^1H NMR (DMSO- d_6 , 400MHz): δ 8.94 (brs, 1H, exchangeable with D_2O , NH), 7.30 (d, $J=8.0\text{Hz}$, 2H), 7.20 (d, $J=8.0\text{Hz}$, 2H), 6.70 (s, 2H), 5.01 (s, 2H), 2.31 (s, 3H), 2.18 (s, 2H), 2.09 (s, 6H), 1.05 (s, 9H). MS: 340 (M+1).

N-[4-(4-Trifluoromethyl-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide


¹H NMR (DMSO-d₆, 400MHz): δ 8.95 (brs, 1H, exchangeable with D₂O, NH), 7.76 (d, J=8.0Hz, 2H), 7.66 (d, J=8.0Hz, 2H), 6.74 (s, 2H), 5.19 (s, 2H), 2.18 (s, 2H), 2.10 (s, 6H), 1.05 (s, 9H). MS: 394 (M+1).

N-[4-(2-Trifluoromethyl-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide


¹H NMR (DMSO-d₆, 400MHz): δ 8.96 (brs, 1H, exchangeable with D₂O, NH), 7.80 (d, J=8.0Hz, 1H), 7.74 (t, J=8.0Hz, 1H), 7.71 (d, J=8.0Hz, 1H), 7.59 (t, J=8.0Hz, 1H), 6.72 (s, 2H), 5.19 (s, 2H), 2.19 (s, 2H), 2.11 (s, 6H), 1.05 (s, 9H). MS: 394 (M+1).

N-[4-(3-Trifluoromethyl-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide


¹H NMR (DMSO-d₆, 400MHz): δ 8.96 (brs, 1H, exchangeable with D₂O, NH), 7.80 (s, 1H), 7.75 (d, J=8.0Hz, 1H), 7.70 (d, J=8.0Hz, 1H), 7.64 (t, J=8.0Hz, 1H), 6.75 (s, 2H), 5.18 (s, 2H), 2.19 (s, 2H), 2.11 (s, 6H), 1.05 (s, 9H). MS: 394 (M+1).

N[4-(3-Chloro-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide

^1H NMR (DMSO- d_6 , 400MHz): δ 8.94 (brs, 1H, exchangeable with D_2O , NH), 7.50 (s, 1H), 7.41 (m, 3H), 6.73 (s, 2H), 5.09 (s, 2H), 2.18 (s, 2H), 2.10 (s, 6H), 1.05 (s, 9H). MS: 360 (M+1).


5 N-[4-(3-Fluoro-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide

^1H NMR (DMSO- d_6 , 400MHz): δ 8.94 (brs, 1H, exchangeable with D_2O , NH), 7.50 (s, 1H), 7.41 (m, 3H), 6.73 (s, 2H), 5.09 (s, 2H), 2.18 (s, 2H), 2.10 (s, 6H), 1.05 (s, 9H). MS: 360 (M+1).

N-[4-(2,4-Difluoro-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide


10

^1H NMR (DMSO- d_6 , 400MHz): δ 8.96 (brs, 1H, exchangeable with D_2O , NH), 7.61 (dd, $J=8.3$ and 15.3Hz , 1H), 7.30 (dt, $J=2.2$ and 10.1Hz , 1H), 7.13 (dt, $J=2.2$ and 8.3Hz , 1H), 6.74 (s, 2H), 5.06 (s, 2H), 2.19 (s, 2H), 2.11 (s, 6H), 1.05 (s, 9H). MS: 362 (M+1).


N-[4-(3,4-Difluoro-benzyloxy)-2,6-dimethyl-phenyl]-3,3-dimethyl-butyramide

15

^1H NMR (DMSO- d_6 , 400MHz): δ 8.93 (brs, 1H, exchangeable with D_2O , NH), 7.49 (t, $J=7.7\text{Hz}$, 1H), 7.43 (dd, $J=6.9$ and 15.2Hz , 1H), 7.28 (m, 1H), 6.70 (s, 2H), 5.03 (s, 2H), 2.16 (s, 2H), 2.08 (s, 6H), 1.03 (s, 9H). MS: 362 (M+1).

3-Cyclopentyl-N-[4-(3,4-difluoro-benzyloxy)-2,6-dimethyl-phenyl]-propionamide

¹H NMR (DMSO-d₆, 400MHz): δ 8.99 (brs, 1H, exchangeable with D₂O, NH), 7.48 (t, J=7.7Hz, 1H), 7.42 (dd, J=6.9 and 15.2 Hz, 1H), 7.28 (m, 1H), 6.70 (s, 2H), 5.03 (s, 2H), 2.27 (t, J=6.0Hz, 2H), 2.06 (s, 6H), 1.75 (m, 3H), 1.60 (m, 4H), 1.47 (m, 2H), 1.10 (m, 2H). MS: 388 (M+1).

3-Cyclopentyl-N-[4-(4-fluoro-benzyloxy)-2,6-dimethyl-phenyl]-propionamide

¹H NMR (DMSO-d₆, 400MHz): δ 8.98 (brs, 1H, exchangeable with D₂O, NH), 7.46 (dd, J=4.7 and 6.5Hz, 2H), 7.20 (t, J=7.0Hz, 1H), 6.69 (s, 2H), 5.02 (s, 2H), 2.28 (t, J=6.0Hz, 2H), 2.11 (s, 6H), 1.76 (m, 3H), 1.60 (m, 4H), 1.47 (m, 2H), 1.10 (m, 2H). MS: 370 (M+1).

Biological Results

Compounds of this invention formula were evaluated for activity toward potassium channels in a cell-based Rb⁺ efflux assay. This cellular bioassay is believed to faithfully represent the M current channel activities identified with KCNQ2/3 heteromultimers. The most active compounds of this invention have EC₅₀s in the single-digit nM range. Additionally, antiseizure activity *in vivo* was evaluated in a mouse maximal electroshock seizure (MES) model, and neurotoxicities were determined from a rotorod neurocognitive motor impairment model.

Methods:**Rubidium Efflux Test**

PC-12 cells were grown at 37 °C and 5 % CO₂ in DMEM/F12 Medium (Dulbecco's Modified Eagle Medium with Nutrient Mix F-12, available from Invitrogen of Carlsbad, CA), supplemented with 10 % horse serum, 5 % fetal bovine serum, 2 mM glutamine, 100 U/ml penicillin, and 100 U/ml streptomycin. They were plated in poly-D-lysine-coated 96-well cell

culture microplates at a density of 40,000 cells/well and differentiated with 100 ng/ml NGF-7s for 2-5 days. For the assay, the medium was aspirated, and the cells were washed once with 0.2 ml in wash buffer (25 mM HEPES, pH 7.4, 150 mM NaCl, 1 mM MgCl₂, 0.8 mM NaH₂PO₄, 2 mM CaCl₂). The cells were then loaded with 0.2 ml Rb⁺ loading buffer (wash buffer plus 5.4 mM RbCl₂, 5 mM glucose) and incubated at 37 °C for 2 h. Attached cells were quickly washed three times with buffer (same as Rb⁺ loading buffer, but containing 5.4 mM KCl instead of RbCl) to remove extracellular Rb⁺. Immediately following the wash, 0.2 ml of depolarization buffer (wash buffer plus 15 mM KCl) with or without compounds was added to the cells to activate efflux of potassium ion channels. After incubation for 10 min at room temperature, the supernatant was carefully removed and collected. Cells were lysed by the addition of 0.2 ml of lysis buffer (depolarization buffer plus 0.1 % Triton X-100) and the cell lysates were also collected. If collected samples were not immediately analyzed for Rb⁺ contents by atomic absorption spectroscopy (see below), they were stored at 4 °C without any negative effects on subsequent Rb⁺ analysis.

The concentrations of Rb⁺ in the supernatants (Rb⁺_{Sup}) and the cell lysates (Rb⁺_{Lys}) were quantified using an ICR8000 flame atomic absorption spectrometer (Aurora Biomed Inc., Vancouver, B.C.) under conditions defined by the manufacturer. Samples 0.05 ml in volume were processed automatically from microtiter plates by dilution with an equal volume of Rb⁺ sample analysis buffer and injection into an air-acetylene flame. The amount of Rb⁺ in the sample was measured by absorption at 780 nm using a hollow cathode lamp as light source and a PMT detector. A calibration curve covering the range 0–5 mg/L Rb⁺ in sample analysis buffer was generated with each set of plates. The percent Rb⁺ efflux (*F*) was defined by

$$F = [\text{Rb}^+_{\text{Sup}} / (\text{Rb}^+_{\text{Sup}} + \text{Rb}^+_{\text{Lys}})] \times 100 \%$$

wherein the *F*_c is the efflux in the presence of compound in depolarization buffer, *F*_b is the efflux in basal buffer, and *F*_s is the efflux in depolarization buffer, and *F*_c is the efflux in the presence of compound in depolarization buffer. The efflux (*F*) and compound concentration relationship was plotted to calculate an *EC*₅₀ value, a compound's concentration for 50% of maximal Rb⁺ efflux. The results are shown below in Table 1.

30 Maximal Electroshock Seizure (MES) and Acute Toxicity Tests

MES Test

The MES testing protocol is based on procedures established at the National Institute of

Neurological Disorders and Stroke in conjunction with the Anticonvulsant Screening Program (ASP) at the University of Utah (White, H.S., Woodhead, J.H., Wilcox, K.S., Stables, J.P., Kupferberg, H.J and Wolf, H.H. 2002. "General Principles: Discovery and Preclinical Development of Antiepileptic Drugs," in *Antiepileptic Drugs*, 5th Edition, R.H. Levy, ed.; R.H. Mattson, B.S. Meldrum, and E. Perucca. Philadelphia, Lippincott Williams & Wilkins.), The goal of the test rapid identification and characterization of the *in vivo* anticonvulsant activity of any compounds that have been shown active in PC-12 cellular based Rb⁺ efflux assay.

Adult male CF-1 albino mice (18-25 g, Charles River Laboratories) are exclusively used for in-house MES screen of compounds. Male Sprague-Dawley albino rats (100-125g, Charles River Laboratories) are also used to test anticonvulsant compounds. Variability of test outcomes is reduced by using animals of the same sex, age, and weight. Animals are permitted to rest and recover from transit for at least 48 hr prior to experimentation. Animals are used for AED testing only once. In some instances, the animals may be anesthetized prior to blood collection and/or whole brain extraction for pharmacokinetic assay. All animals are maintained and handled as outlined in standard animal care guidelines.

In the experiments, testing compounds are prepared as suspensions in 0.5% methyl cellulose (Sigma, Cat # M0512, Viscosity 4000 cP at 20°C) in water, regardless of solubility. Dry powder compounds are initially ground with a glass rod in a test tube in several drops of methyl cellulose to create a paste and to break down any large chunks. After several minutes of grinding, the volume of the suspension is increased to the final concentration desired. The suspension is then sonicated using a Branson sonicator model 3510 in a water bath at room temperature for 15 minutes. Compound suspensions are further vortexed prior to animal dosing. In some of the cases, DMSO is used to initially solubilize compounds in small volumes and then this solution is added to the 0.5% methyl cellulose solution, in order to create more even and less aggregated compound suspensions. The final concentration of DMSO is 3.75%, an amount with no apparent toxicity or neuroprotective effects in our usual rotarod and MES tests. Methyl cellulose/DMSO compound suspensions are identically prepared for intraperitoneally (i.p.) to mice or orally (p.o.) to rat dosing.

Initially the animals are weighed with an electronic scale and then marked. Data recording sheets are generated for each compound assessment. Mice or rats are dosed with the compound suspension at 0.01 mL/g of body weight. The typical injection volume range is between 180-250 µl for mice. Compounds are dosed by i.p. to mice using a 25 or 22 gauge needle, depending on

the viscosity of the suspension. Rats are p.o. dosed using a flexible feeding tube, typically starting at a compound dose of 5 mg/kg.

A Rodent Electroconvulsive Stimulator (Model 200, Hamit-Darvin-Freesh, Snow Canyon Clinic, Ivins, UT) is used for MES testing. A 60-Hz alternating current (50 mA for mice; 150 mA for rats) is delivered for 0.2 seconds through corneal electrodes to the mice. A drop of 0.5% tetracaine (Sigma, Cat. # T-7508) solution is placed on the eye prior to current delivery. The electrodes are subsequently placed gently onto the eyes of the animal and the electrical shock is initiated by triggering through a foot-pedal activator. The animals are restrained by hand and gently released as the shock is delivered and the seizure commences. Animals are monitored for hind limb tonic extension as the end point for this test. Current delivery is recorded as a measure of overall seizure-induction potential. Electrical current delivery can vary from approximately 30-55 mA (mice) or 90-160 mA (rats) depending on impedance in the animal and quality of the current delivery (i.e., correct placement of the electrodes on the cornea). Seizures will be successfully induced in control animals throughout this current range. Tonic extension is considered abolished if the hind limbs fail to become fully extended at 180° with the plane of the body. Lack of tonic extension suggests that the test compound has prevented the spread of seizure discharge through neural tissue. Although unnecessary in mice, the rats are pre-screened for seizure induction potential using the MES 24hr prior to compound dosing and the subsequent MES test. A success rate of 92-100% has been determined for the rat seizure induction potential. Rats that fail to develop tonic/clonic seizures during the pre-screening are not used for drug testing.

For compound testing, time-to-peak effect studies are initially performed using 0.5, 1, 2, 4, 8 and 24 hr time points, typically using a single 5 or 25 mg/kg dose. The determined time-to-peak effect is used for further titration of a compound's potency (ED₅₀, the dose of a drug that protects 50% of animals from electrical induced seizure) in both mouse and rat models. For titrations, 8 animals are used per concentration and dose (normal 5 concentrations) is varied until a full dose response curve can be obtained. Probit analysis (ASP method) or non-linear regression analysis on Graph Pad (constraining the lower dose/effect value) is used to calculate an ED₅₀ value for the test compound.

30 Rotarod Test

Prior to MES testing, compound dosed mice are scrutinized for abnormal neurologic status as defined by motor impairment on a slowly turning (6 rpm) rotarod apparatus (Model 755, Series

8, IITC Life Sciences, Woodland Hills, CA). The inability of a mouse to maintain its balance on the rotarod over a period of one minute (three falls = failure) signifies motor impairment and hence acute toxicity. These measurements are done at the same time points as the MES assay. Untreated normal mice are able to maintain balance on the rotarod for at least one minute
5 without falling. Median toxicity of a compound (TD_{50} , the dose of a drug that results in motor impairment in 50% of animals) is determined.

Open Field Test






Before the MES test, compound treated rats are visually observed for acute toxicity signs for approximately one minute in the open field test. Here, rats are gently placed into a plexiglass
10 enclosure and are monitored for behavior consistent with toxicity including ataxia, trembling, hypoactivity (including failure to seek the walls), hypersensitivity, lack of exploratory behavior and lack of avoidance of the open area. Typically if the rats exhibits two or more of these abnormal behaviors they are scored as toxic.






TABLE 1






ACTIVITIES OF EXEMPLARY COMPOUNDS



Legend: A: $EC_{50} \leq 1 \text{ nM}$; B: $1 \text{ nM} < EC_{50} \leq 10 \text{ nM}$;
C: $10 \text{ nM} < EC_{50} \leq 50 \text{ nM}$; D: $50 \text{ nM} < EC_{50} \leq 500 \text{ nM}$

5

COMPOUND	ACTIVITY EC_{50}
	D
	D
	D
	D
	B

COMPOUND	ACTIVITY EC ₅₀
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2cc(F)c(F)cc2)cc1C</chem>	C
 <chem>C1CCCCC1CCC(=O)Nc1cc(C)c(OCc2ccc(F)cc2)cc1C</chem>	B
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2ccc(F)cc2)cc1C</chem>	C
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2ccc(C)cc2)cc1C</chem>	C
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2cc(C(F)(F)F)ccc2)cc1C</chem>	C

COMPOUND	ACTIVITY EC ₅₀
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2ccccc2C(F)(F)F)c(C)c1</chem>	D
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2ccc(Cl)cc2)c(C)c1</chem>	C
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2ccc(Br)cc2)c(C)c1</chem>	C
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2ccccc2C(F)(F)F)c(C)c1</chem>	C
 <chem>CC(C)(C)C(=O)Nc1cc(C)c(OCc2cc(F)ccc2F)c(C)c1</chem>	C

COMPOUND	ACTIVITY EC ₅₀
	C
	C

Studies of KCNQ2/3 opening activity and KCNQ subtype selectivity using electrophysiological patch clamp in *Xenopus* oocytes

5 Expression in *Xenopus laevis* oocytes

Female *Xenopus laevis* extracted ovaries can be purchased from eNASCO (LM00935MX, eNASCO Fort Atkinson, WI). Following manual dissection of the oocytes into smaller groups, the oocytes are defolliculated by enzymatic treatment with collagenase type 2 (LS004177, 10 Worthington, Lakewood, NJ) for 1-1/2 hour in the presence of calcium-free Culture Bath solution (88 mM NaCl, 1 mM KCl, 0.82 mM MgSO₄, 2.4 mM NaHCO₃, and 5 mM HEPES, pH 7.5). Oocytes are then kept in supplemented Culture Bath solution (88 mM NaCl, 1 mM KCl, 0.82 mM MgSO₄, 0.9 mM CaCl₂, 2.4 mM NaHCO₃, 1 mM sodium pyruvate, 0.05 mg/ml Geneticin, 100 U/ml penicillin, 0.1 mg/ml streptomycin and 5 mM HEPES, pH 7.5) at 19°C for 15 24 hours before injection of cRNA. Approximately 50 nl cRNA (about 50 ng) is injected for KCNQ1, KCNQ4, and KCNQ5 using a Nanoject microinjector (Drummond, Broomall, PA, USA). For co-expression of KCNQ2 and KCNQ3 and of KCNQ1 and KCNE1, cRNA's are mixed in equal molar ratios before injection of approximately 50 nl. The mixtures contain about 10 + 10 ng and 12.5 + 2.5 ng cRNA, respectively. The smaller amounts are needed because 20 larger currents arise when KCNQ2/KCNQ3 and KCNQ1/KCNE1 are co-expressed. Oocytes are

kept in Culture Barth solution at 19°C which is changed daily and currents are recorded after 3 to 5 days.

Electrophysiology

5 KCNQ channel currents expressed in *Xenopus laevis* oocytes are recorded using a two-electrode voltage-clamp. The recordings are made at room temperature in recording solution (96 mM NaCl, 2 mM KCl, 1 mM MgCl₂, 1.8 mM CaCl₂, and 5 mM HEPES, pH 7.5) using a two-electrode voltage-clamp amplifier (OC-725C, Warner Instrument, Hamden, CT, USA). The oocytes are placed in custom built perfusion chambers connected to a continuous flow system
10 and impaled with a current electrode and a voltage-clamp electrode pulled from borosilicate glass on a Flaming/Brown Micropipette Puller (Sutter Instruments Co, Novato, CA, USA). Recording electrodes are filled with 3 M KCl and had a resistance of 0.5 to 2.5 MΩ.

Compounds

15 All compounds are dissolved in DMSO to obtain concentrated stock solutions. On the day of electrophysiological experiments the stock solutions are thawed and diluted in recording solution to their final concentrations. The final DMSO concentration never exceeds 0.1%. Compound delivery is performed using a custom built multi-barrel apparatus connected to the flow system.

Calculations

20 Data are acquired by means of an Axograph X software (Axograph Scientific, Sydney, AU) and analyzed using Graph Pad Prism (GraphPad Software Inc., CA, USA).

Concentration - response curves are constructed by plotting the increase in steady-state current expressed in percentages as a function of drug concentration. During the course of the
25 experiment, while various concentrations of the drug are being dosed, the resting voltage is held at -90 mV and pulsed to -60 mV, -40 mV, and -50 mV for 5 s for KCNQ2/KCNQ3, KCNQ4 and KCNQ5 channels respectively. The plot was then fitted to a Hill function:

$$\text{Response} = R2 + (R1 - R2) / [1 + (C / EC_{50})^{nH}]$$

30 wherein R1 is the initial response, R2 is the maximum response, C is the drug concentration and nH is the slope (Hill coefficient) of the curve.

The efficacy of compounds of this invention in comparison with Retigabine (as a positive control) are determined by recording the steady current using the above voltage protocol for the

channels in the presence of the EC₇₅ of the drugs. After steady channel current is recorded in the presence of Retigabine at its EC₇₅, recorded oocyte is washed with the recording solution until its steady current returned to its normal level without the presence of any drugs. Then the channel steady current is recorded in the presence of the test compound at its EC₇₅. The percent efficacy is then expressed as:

$$\% \text{ efficacy} = (C2/C1) \times 100 \%$$


wherein C2 is the recorded steady current in the presence of follow-on compound at its EC₇₅ and C1 is the recorded steady current in the presence of Retigabine at its EC₇₅.

10 In view of the above, it will be seen that the several objects of the invention are achieved and other advantageous results attained.

As various changes could be made in the above methods and products without departing from the scope of the invention, it is intended that all matter contained in the above description shall be interpreted as illustrative and not in a limiting sense.

What is claimed is:


1. A compound having the following structural formula I:



- wherein R_1 and R_2 are independently selected from the group consisting of H, CN, halogen, NH_2 , CH_2CN , OH, NO_2 , CH_2F , CHF_2 , CF_3 , CF_2CF_3 , C_1 - C_6 alkyl, $C(=O)C_1$ - C_6 alkyl; NH - C_1 - C_6 alkyl; $N(C_1$ - C_6 alkyl)- C_1 - C_6 alkyl, $NHC(=O)C_1$ - C_6 alkyl, $C(=O)N(CH_3)_2$, $C(=O)N(Et)_2$, $C(=O)NH_2$, $C(=O)NH$ - C_1 - C_6 alkyl, SO_2NH_2 , $NHSO_2$ - C_1 - C_6 alkyl; $C(=O)OC_1$ - C_6 alkyl, $OC(=O)C_1$ - C_6 alkyl, OC_1 - C_6 alkyl, SC_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, $(CH_2)_mC_3$ - C_6 cycloalkyl, C_3 - C_6 cycloalkenyl, $(CH_2)_mC_3$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, Ar, $(CH_2)_m$ thienyl, $(CH_2)_m$ imidazolyl, $(CH_2)_m$ pyrazyl, $(CH_2)_m$ oxazolyl, $(CH_2)_m$ isoxazolyl, $(CH_2)_m$ thiazolyl, $(CH_2)_m$ isothiazolyl, $(CH_2)_m$ phenyl, $(CH_2)_m$ pyrrolyl, $(CH_2)_m$ pyridyl, and $(CH_2)_m$ pyrimidyl, wherein $m =$ zero, 1, or 2, Ar is a 5- to 10- member mono or bicyclic aromatic group, optionally containing 1-4 ring heteroatoms selected independently from N, O, and S; or R_1 and R_2 , together with the ring carbon atoms to which they are attached, form a 5- or 6- member fused ring, which ring may be saturated, unsaturated, or aromatic, which optionally contains one or two heteroatoms selected independently from O, N, and S; R_3 and R_4 are independently selected from the group consisting of H, CN, halogen, CF_3 , OCF_3 , OC_1 - C_3 alkyl, and C_1 - C_3 alkyl; X is O or S; Y is O or S; $q = 1$ or zero; R_5 is selected from the group consisting of C_1 - C_6 alkyl, $(CHR_6)_wC_3$ - C_6 cycloalkyl, $(CHR_6)_wCH_2C_3$ - C_6 cycloalkyl, $CH_2(CHR_6)_wC_3$ - C_6 cycloalkyl, $CR_6=CH$ - C_3 - C_6 cycloalkyl, $CH=CR_6$ - C_3 - C_6 cycloalkyl, $(CHR_6)_wC_5$ - C_6 cycloalkenyl, $CH_2(CHR_6)_wC_5$ - C_6 cycloalkenyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, Ar, $(CHR_6)_w$ Ar, $CH_2(CHR_6)_w$ Ar, and $(CHR_6)_wCH_2$ Ar, wherein $w =$ zero, 1, 2, or 3, Ar is a 5- to 10- member mono or bicyclic aromatic group, optionally containing 1-4 ring heteroatoms selected independently from N, O, and S; R_6 is H or C_1 - C_3 alkyl; wherein all cycloalkyl and cycloalkenyl groups optionally contain one or two ring heteroatoms selected independently from N, O, and S; wherein all alkyl, cycloalkyl, alkenyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, alkynyl, aryl, and heteroaryl groups in R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , and Ar are optionally substituted with one or

two substituents independently selected from the group consisting of C₁-C₃ alkyl, halogen, OH, OEt, OMe, CN, CH₂F, OCF₃, and CF₃; and wherein, additionally, all cycloalkyl and heterocycloalkyl groups are optionally substituted with a carbonyl group, and their pharmaceutically acceptable salts, solvates and esters thereof.

- 5 2. The compound according to claim 1 having the following structural formula:






wherein R₁ is selected from the group consisting of H, halogen, CN, CH₂CN, CHF₂, CF₃, C₁-C₆ alkyl, OCH₃, (C=O)OCH₃, O(C=O)CH₃, OCF₃, (CH₂)_mC₃-C₆ cycloalkyl, phenyl, and pyridyl; R₂ is selected from the group consisting of H, F, OCH₃, CH₃, and CF₃; R₃ and R₄ are independently, selected from the group consisting of H, F, Cl, CF₃, OCF₃, OC₁-C₃ alkyl, and C₁-C₃ alkyl; and R₅ is selected from the group consisting of C₁-C₆ alkyl, (CHR₆)_wC₃-C₆ cycloalkyl, (CHR₆)_wCH₂C₃-C₆ cycloalkyl, CH₂(CHR₆)_wC₃-C₆ cycloalkyl, CR₆=CH-C₃-C₆ cycloalkyl, CH=CR₆-C₃-C₆ cycloalkyl, (CHR₆)_wC₅-C₆ cycloalkenyl, CH₂(CHR₆)_wC₅-C₆ cycloalkenyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, Ar, (CHR₆)_wAr, CH₂(CHR₆)_wAr, and (CHR₆)_wCH₂Ar, wherein w = 0 – 3, Ar is selected from the group consisting of phenyl, furyl, pyrrolyl, oxazolyl, thiazolyl, thienyl, and pyridyl; and R₆ is C₁-C₃ alkyl; wherein all alkyl, cycloalkyl, aryl, and heteroaryl groups in R₁, R₂, R₃, R₄, R₅, and R₆, and Ar are optionally substituted with one or two substituents selected independently from C₁-C₃ alkyl, halogen, OCH₃, OCH₂CH₃, CN, and CF₃.







3. The compound according to claim 2 wherein R₁ is selected from the group consisting of H, F, Cl, Br, CHF₂, CF₃, C₁-C₆ alkyl, OCH₃, CH₂OCH₃, CH₂CH₂OCH₃, CH₂OCH₂CH₃, OCH₂CH₃, and CH₂OCH₃; R₂ is selected from the group consisting of H, CH₃, CH₂CH₃, and halogen; R₃ and R₄ are, independently, selected from the group consisting of H, F, Cl, CF₃, OCF₃, OCH₃, and CH₃; and R₅ is selected from the group consisting of C₁-C₆ alkyl, CH₂C₃-C₆ cycloalkyl, CH₂CH₂C₃-C₆ cycloalkyl, CH=CH-C₃-C₆ cycloalkyl, CH=CH-C₅-C₆ cycloalkenyl, CH₂C₅-C₆ cycloalkenyl, CH₂CH₂C₅-C₆ cycloalkenyl, C₂-C₆ alkenyl, and (CH₂)_wAr, wherein w = 1 or 2; Ar is phenyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furyl, thienyl, pyrrolyl, pyridyl; wherein

all alkyl, cycloalkyl, aryl, and hetroaryl groups in R₁, R₂, R₃, R₄, R₅, R₆ and Ar are optionally substituted with one or two substituents selected independently from the group consisting of CH₃, halogen, OCH₃, OCH₂CH₃, CN, and CF₃.

4. A compound according to claim 3 wherein R₁ is selected from the group consisting of H, F, Cl, Br, CF₃, CHF₂, and C₁-C₆ alkyl; R₂ is selected from the group consisting of H, CH₃, CH₂CH₃ and halogen; R₃ and R₄ are independently selected from the group consisting of H, F, OCH₃ and CH₃; and R₅ is selected from the group consisting of C₁-C₆ alkyl, CH₂C₃-C₆ cycloalkyl or CH₂CH₂C₃-C₆ cycloalkyl.
5. The compound according to claim 4 wherein the compound is selected from the group consisting of:





63





5

10


64



5







6. The compound according to claim 1 having the following structural formula:



- wherein R_1 is selected from the group consisting of H, halogen, CN, CH_2CN , CHF_2 , CF_3 , $\text{C}_1\text{-C}_6$ alkyl, OCH_3 , $(\text{C}=\text{O})\text{OCH}_3$, $\text{O}(\text{C}=\text{O})\text{CH}_3$, OCF_3 , $(\text{CH}_2)_m\text{C}_3\text{-C}_6$ cycloalkyl, phenyl, and pyridyl; R_2 is selected from the group consisting of H, F, OCH_3 , CH_3 , and CF_3 ; R_3 and R_4 are independently, selected from the group consisting of H, F, Cl, CF_3 , OCF_3 , $\text{OC}_1\text{-C}_3$ alkyl, and $\text{C}_1\text{-C}_3$ alkyl; and R_5 is selected from the group consisting of $\text{C}_1\text{-C}_6$ alkyl, $(\text{CHR}_6)_w\text{C}_3\text{-C}_6$ cycloalkyl, $(\text{CHR}_6)_w\text{CH}_2\text{C}_3\text{-C}_6$ cycloalkyl, $\text{CH}_2(\text{CHR}_6)_w\text{C}_3\text{-C}_6$ cycloalkyl, $\text{CR}_6=\text{CH-C}_3\text{-C}_6$ cycloalkyl, $\text{CH}=\text{CR}_6\text{-C}_3\text{-C}_6$ cycloalkyl, $(\text{CHR}_6)_w\text{C}_5\text{-C}_6$ cycloalkenyl, $\text{CH}_2(\text{CHR}_6)_w\text{C}_5\text{-C}_6$ cycloalkenyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, Ar, $(\text{CHR}_6)_w\text{Ar}$, $\text{CH}_2(\text{CHR}_6)_w\text{Ar}$, and $(\text{CHR}_6)_w\text{CH}_2\text{Ar}$, wherein $w = 0 - 3$, Ar is selected from the group consisting of phenyl, furyl, pyrrolyl, oxazolyl, thiazolyl, thienyl, and pyridyl; and R_6 is $\text{C}_1\text{-C}_3$ alkyl; wherein all alkyl, cycloalkyl, aryl, and heteroaryl groups in R_1 , R_2 , R_3 , R_4 , R_5 , and R_6 , and Ar are optionally substituted with one or two substituents selected independently from $\text{C}_1\text{-C}_3$ alkyl, halogen, OCH_3 , OCH_2CH_3 , CN, and CF_3 .
7. The compound according to claim 2 wherein R_1 is selected from the group consisting of H, F, Cl, Br, CHF_2 , CF_3 , $\text{C}_1\text{-C}_6$ alkyl, OCH_3 , CH_2OCH_3 , $\text{CH}_2\text{CH}_2\text{OCH}_3$, $\text{CH}_2\text{OCH}_2\text{CH}_3$, OCH_2CH_3 , and CH_2OCH_3 ; R_2 is selected from the group consisting of H, CH_3 , CH_2CH_3 , and halogen; R_3 and R_4 are, independently, selected from the group consisting of H, F, Cl, CF_3 , OCF_3 , OCH_3 , and CH_3 ; and R_5 is selected from the group consisting of $\text{C}_1\text{-C}_6$ alkyl, $\text{CH}_2\text{C}_3\text{-C}_6$ cycloalkyl, $\text{CH}_2\text{CH}_2\text{C}_3\text{-C}_6$ cycloalkyl, $\text{CH}=\text{CH-C}_3\text{-C}_6$ cycloalkyl, $\text{CH}=\text{CH-C}_5\text{-C}_6$ cycloalkenyl, $\text{CH}_2\text{C}_5\text{-C}_6$ cycloalkenyl, $\text{CH}_2\text{CH}_2\text{C}_5\text{-C}_6$ cycloalkenyl, $\text{C}_2\text{-C}_6$ alkenyl, and $(\text{CH}_2)_w\text{Ar}$, wherein $w = 1$ or 2 ; Ar is phenyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, furyl, thienyl, pyrrolyl, pyridyl; wherein all alkyl, cycloalkyl, aryl, and heteroaryl groups in R_1 , R_2 , R_3 , R_4 , R_5 , R_6 and Ar are optionally substituted with one or two substituents selected independently from the group consisting of CH_3 , halogen, OCH_3 , OCH_2CH_3 , CN, and CF_3 .

8. A compound according to claim 3 wherein R_1 is selected from the group consisting of H, F, Cl, Br, CF_3 , CHF_2 , and C_1 - C_6 alkyl; R_2 is selected from the group consisting of H, CH_3 , CH_2CH_3 and halogen; R_3 and R_4 are independently selected from the group consisting of H, F, OCH_3 and CH_3 ; and R_5 is selected from the group consisting of C_1 - C_6 alkyl, CH_2C_3 - C_6 cycloalkyl or $CH_2CH_2C_3$ - C_6 cycloalkyl.
9. The compound according to claim 8 wherein the compound is selected from the group consisting of:






10. A composition comprising a pharmaceutically acceptable carrier and one or more of the following:

- i. a compound of formula IA-1 or formula IA-2,
- ii. a pharmaceutically acceptable solvate of a compound of formula IA-1 or formula IA-2,
- 5 iii. a pharmaceutically acceptable salt of a compound of formula IA-1 or formula IA-2;
- iv. a pharmaceutically acceptable ester of a compound of formula IA-1 or formula IA-2.




11. The composition according to claim 10 wherein the formula is IA-1.

12. The composition according to claim 10 wherein the formula is IA-2.



13. The composition according to claim 11 wherein the compound of formula IA-1 is selected
10 from the group consisting of:




68







5



10




69



5

10

14. The composition according to claim 12 wherein the compound of formula IA-2 is selected from the group consisting of:








15. A method of preventing or treating a disease or disorder which is affected by activation of voltage-gated potassium channels, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of formula IA-1 or formula IA-2 or a salt or ester or solvate thereof.

16. The method according to claim 15 wherein the formula is IA-1.

17. The method according to claim 15 wherein the formula is IA-2.



18. The method according to claim 16 wherein the compound of formula IA-1 is selected from the group consisting of:






5

10

72






5




10

73




5

19. The method according to claim 17 wherein the compound of formula IA-2 is selected from the group consisting of :





10



and

74





(I)