

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
6 February 2003 (06.02.2003)

PCT

(10) International Publication Number
WO 03/010286 A2

- (51) International Patent Classification⁷: **C12N**
- (21) International Application Number: PCT/US02/23254
- (22) International Filing Date: 23 July 2002 (23.07.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/909,796 23 July 2001 (23.07.2001) US
10/141,647 7 May 2002 (07.05.2002) US
- (71) Applicant: **SENESCO, INC.** [US/US]; 303 George Street, Suite 420, New Brunswick, NJ 08901 (US).
- (72) Inventors: **TAYLOR, Catherine**; 11 Devitt Avenue South, Apt.301, Waterloo, Ontario N2J 1Y6 (CA). **WANG, Tzann-Wei**; 607 Salzburg Drive, Waterloo, Ontario N2V 2L5 (CA). **PETROV, Larry**; 66 Mooregate Crescent, Apt. 1210, Kitchener, Ontario N2M 5E6 (CA). **CARLSON, John, C.**; 126 Woodbend Crescent, Waterloo, Ontario N2T 1G9 (CA). **NARAYANSINGH, Richard**; 119 Lorilee Crescent, Kitchener, Ontario N2N 2J5 (CA). **THOMPSON, John, E.**; 640 Woolwich Street, Waterloo, Ontario N2J 4G8 (CA). **CLICHE, Dominic**; 75 Old Chicopee Drive, Unit #52, Kitchener, Ontario N2A 2G4 (CA). **KAUP, Marianne**; 102 Perth Street, New Hamburg, Ontario N0B 2G0 (CA).
- (74) Agents: **LAVENUE, Teresa, A.** et al.; Kenyon & Kenyon, 1500 K Street N.W., Suite 700, Washington, DC 20005 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: NUCLEIC ACIDS, POLYPEPTIDES, AND METHODS FOR MODULATING APOPTOSIS



(57) Abstract: The present invention relates to isolated and/or purified rat apoptosis-specific eucaryotic initiation Factor-5A (eIF-5A) and deoxyhypusine synthase (DHS) nucleic acids and polypeptides. The present invention also relates to methods of modulating apoptosis using apoptosis-specific eIF-5A and DHS, and antisense oligonucleotides and expression vectors of apoptosis-specific eIF-5A and DHS useful in such methods.

NUCLEIC ACIDS, POLYPEPTIDES, AND METHODS FOR MODULATING APOPTOSIS

FIELD OF THE INVENTION

5

The present invention relates to apoptosis-specific eucaryotic initiation Factor-5A (eIF-5A) and deoxyhypusine synthase (DHS) nucleic acids and polypeptides and methods for modulating apoptosis in cells using apoptosis-specific eIF-5A and DHS.

10

BACKGROUND OF THE INVENTION

Apoptosis is a genetically programmed cellular event that is characterized by well-defined morphological features, such as cell shrinkage, chromatin condensation, nuclear fragmentation, and membrane blebbing. Kerr et al. (1972) *Br. J. Cancer*, **26**, 239-257; Wyllie et al. (1980) *Int. Rev. Cytol.*, **68**, 251-306. It plays an important role in normal tissue development and homeostasis, and defects in the apoptotic program are thought to contribute to a wide range of human disorders ranging from neurodegenerative and autoimmunity disorders to neoplasms. Thompson (1995) *Science*, **267**, 1456-1462; Mullauer et al. (2001) *Mutat. Res*, **488**, 211-231. Although the morphological characteristics of apoptotic cells are well characterized, the molecular pathways that regulate this process have only begun to be elucidated.

One group of proteins that is thought to play a key role in apoptosis is a family of cysteine proteases, termed caspases, which appear to be required for most pathways of apoptosis. Creagh & Martin (2001) *Biochem. Soc. Trans*, **29**, 696-701; Dales et al. (2001) *Leuk. Lymphoma*, **41**, 247-253. Caspases trigger apoptosis in response to apoptotic stimuli by cleaving various cellular proteins, which results in classic manifestations of apoptosis, including cell shrinkage, membrane blebbing and DNA fragmentation. Chang & Yang (2000) *Microbiol. Mol. Biol. Rev.*, **64**, 821-846.

Pro-apoptotic proteins, such as Bax or Bak, also play a key role in the apoptotic pathway by releasing caspase-activating molecules, such as mitochondrial cytochrome c, thereby promoting cell death through apoptosis. Martinou & Green (2001) *Nat. Rev. Mol. Cell. Biol.*, **2**, 63-67; Zou et al. (1997) *Cell*, **90**, 405-413. Anti-apoptotic proteins, such as Bcl-2, promote cell survival by antagonizing the activity of the pro-apoptotic proteins, Bax and Bak. Tsujimoto (1998) *Genes Cells*, **3**, 697-707; Kroemer (1997) *Nature Med.*, **3**, 614-620. The ratio of Bax:Bcl-2 is thought to be one way in which cell fate is determined; an excess of Bax promotes apoptosis and an excess of Bcl-2 promotes cell survival. Salomons et al. (1997) *Int. J. Cancer*, **71**, 959-965; Wallace-Brodeur & Lowe (1999) *Cell Mol. Life Sci.*, **55**, 64-75.

Another key protein involved in apoptosis is that encoded by the tumor suppressor gene p53. This protein is a transcription factor that regulates cell growth and induces apoptosis in cells that are damaged and genetically unstable, presumably through up-regulation of Bax. Bold et al. (1997) *Surgical Oncology*, **6**, 133-142; Ronen et al., 1996; Schuler & Green (2001) *Biochem. Soc. Trans.*, **29**, 684-688; Ryan et al. (2001) *Curr. Opin. Cell Biol.*, **13**, 332-337; Zörnig et al. (2001) *Biochem. Biophys. Acta*, **1551**, F1-F37.

The distinct morphological features that characterize cells undergoing apoptosis have given rise to a number of methods for assessing the onset and progress of apoptosis. One such feature of apoptotic cells that can be exploited for their detection is activation of a flippase, which results in externalization of phosphatidylserine, a phospholipid normally localized to the inner leaflet of the plasma membrane. Fadok et al. (1992) *J. Immunol.*, **149**, 4029-4035. Apoptotic cells bearing externalized phosphatidylserine can be detected by staining with a phosphatidylserine-binding protein, Annexin V, conjugated to a fluorescent dye. The characteristic DNA fragmentation that occurs during the apoptotic process can be detected by labeling the exposed 3'-OH ends of the DNA fragments with fluorescein-labeled deoxynucleotides. Fluorescent dyes that bind nucleic acids, such as Hoescht 33258, can be used to detect chromatin condensation and nuclear

fragmentation in apoptotic cells. The degree of apoptosis in a cell population can also be inferred from the extent of caspase proteolytic activity present in cellular extracts.

As a genetically defined process, apoptosis, like any other developmental
5 program, can be disrupted by mutation. Alterations in the apoptotic pathways are believed to play a key role in a number of disease processes, including cancer. Wyllie et al. (1980) *Int. Rev. Cytol.*, **68**, 251-306; Thompson (1995) *Science*, **267**, 1456-1462; Sen & D'Incalci (1992) *FEBS Letters*, **307**, 122-127; McDonnell et al. (1995) *Seminars in Cancer and Biology*, **6**, 53-60. Investigations into cancer
10 development and progression have traditionally been focused on cellular proliferation. However, the important role that apoptosis plays in tumorigenesis has recently become apparent. In fact, much of what is now known about apoptosis has been learned using tumor models, since the control of apoptosis is invariably altered in some way in tumor cells. Bold et al. (1997) *Surgical*
15 *Oncology*, **6**, 133-142.

Apoptosis can be triggered during tumor development by a variety of signals. Extracellular signals include growth or survival factor depletion, hypoxia and ionizing radiation. Internal signals that can trigger apoptosis include DNA damage, shortening telomeres, and oncogenic mutations that produce
20 inappropriate proliferative signals. Lowe & Lin (2000) *Carcinogenesis*, **21**, 485-495. Ionizing radiation and nearly all cytotoxic chemotherapy agents used to treat malignancies are thought to act by triggering endogenous apoptotic mechanisms to induce cell death. Rowan & Fisher (1997) *Leukemia*, **11**, 457-465; Kerr et al. (1994) *Cancer*, **73**, 2013-2026; Martin & Schwartz (1997) *Oncology Research*,
25 **9**, 1-5.

Evidence would suggest that early in the progression of cancer, tumor cells are sensitive to agents (such as ionizing radiation or chemotherapeutic drugs) that induce apoptosis. However, as the tumor progresses, the cells develop resistance to apoptotic stimuli. Naik et al. (1996) *Genes and Development*, **10**,
30 2105-2116. This may explain why early cancers respond better to treatment than

more advanced lesions. The ability of late-stage cancers to develop resistance to chemotherapy and radiation therapy appears to be linked to alterations in the apoptotic pathway that limit the ability of tumor cells to respond to apoptotic stimuli. Reed et al. (1996) *Journal of Cellular Biology*, **60**, 23-32; Meyn et al. 5 (1996) *Cancer Metastasis Reviews*, **15**, 119-131; Hannun (1997) *Blood*, **89**, 1845-1853; Reed (1995) *Toxicology Letters*, **82-83**, 155-158; Hickman (1996) *European Journal of Cancer*, **32A**, 921-926. Resistance to chemotherapy has been correlated to overexpression of the anti-apoptotic gene bcl-2 and deletion or mutation of the pro-apoptotic bax gene in chronic lymphocytic leukemia and 10 colon cancer, respectively.

The ability of tumor cells to successfully establish disseminated metastases also appears to involve alterations in the apoptotic pathway. Bold et al. (1997) *Surgical Oncology*, **6**, 133-142. For example, mutations in the tumor suppressor gene p53 are thought to occur in 70 % of tumors. Evan et al. (1995) 15 *Curr. Opin. Cell Biol.*, **7**, 825-834. Mutations that inactivate p53 limit the ability of cells to induce apoptosis in response to DNA damage, leaving the cell vulnerable to further mutations. Ko & Prives (1996) *Genes and Development*, **10**, 1054-1072.

Therefore, apoptosis is intimately involved in the development and 20 progression of neoplastic transformation and metastases, and a better understanding of the apoptotic pathways involved may lead to new potential targets for the treatment of cancer by the modulation of apoptotic pathways through gene therapy approaches. Bold et al. (1997) *Surgical Oncology*, **6**, 133-142.

25 Deoxyhypusine synthase (DHS) and hypusine-containing eucaryotic translation initiation Factor-5A (eIF-5A) are known to play important roles in many cellular processes including cell growth and differentiation. Hypusine, a unique amino acid, is found in all examined eucaryotes and archaebacteria, but not in eubacteria, and eIF-5A is the only known hypusine-containing protein. 30 Park (1988) *J. Biol. Chem.*, **263**, 7447-7449; Schümann & Klink (1989) *System.*

Appl. Microbiol., **11**, 103-107; Bartig et al. (1990) *System. Appl. Microbiol.*, **13**, 112-116; Gordon et al. (1987a) *J. Biol. Chem.*, **262**, 16585-16589. Active eIF-5A is formed in two post-translational steps: the first step is the formation of a deoxyhypusine residue by the transfer of the 4-aminobutyl moiety of spermidine
5 to the α -amino group of a specific lysine of the precursor eIF-5A catalyzed by deoxyhypusine synthase; the second step involves the hydroxylation of this 4-aminobutyl moiety by deoxyhypusine hydroxylase to form hypusine.

The amino acid sequence of eIF-5A is well conserved between species, and there is strict conservation of the amino acid sequence surrounding the
10 hypusine residue in eIF-5A, which suggests that this modification may be important for survival. Park et al. (1993) *Biofactors*, **4**, 95-104. This assumption is further supported by the observation that inactivation of both isoforms of eIF-5A found to date in yeast, or inactivation of the DHS gene, which catalyzes the first step in their activation, blocks cell division. Schnier et al. (1991) *Mol. Cell.*
15 *Biol.*, **11**, 3105-3114; Sasaki et al. (1996) *FEBS Lett.*, **384**, 151-154; Park et al. (1998) *J. Biol. Chem.*, **273**, 1677-1683. However, depletion of eIF-5A protein in yeast resulted in only a small decrease in total protein synthesis suggesting that eIF-5A may be required for the translation of specific subsets of mRNA's rather than for protein global synthesis. Kang et al. (1993), "Effect of initiation factor
20 eIF-5A depletion on cell proliferation and protein synthesis," in Tuite, M. (ed.), *Protein Synthesis and Targeting in Yeast*, NATO Series H. The recent finding that ligands that bind eIF-5A share highly conserved motifs also supports the importance of eIF-5A. Xu & Chen (2001) *J. Biol. Chem.*, **276**, 2555-2561. In addition, the hypusine residue of modified eIF-5A was found to be essential for
25 sequence-specific binding to RNA, and binding did not provide protection from ribonucleases.

The first cDNA for eIF-5A was cloned from human in 1989 by Smit-McBride et al., and since then cDNAs or genes for eIF-5A have been cloned from various eukaryotes including yeast, rat, chick embryo, alfalfa, and tomato. Smit-
30 McBride et al. (1989a) *J. Biol. Chem.*, **264**, 1578-1583; Schnier et al. (1991)

(yeast); Sano, A. (1995) in Imahori, M. et al. (eds), *Polyamines, Basic and Clinical Aspects*, VNU Science Press, The Netherlands, 81-88 (rat); Rinaudo & Park (1992) *FASEB J.*, **6**, A453 (chick embryo); Pay et al. (1991) *Plant Mol. Biol.*, **17**, 927-929 (alfalfa); Wang et al. (2001) *J. Biol. Chem.*, **276**, 17541-17549
5 (tomato).

In addition, intracellular depletion of eIF-5A resulted in a significant accumulation of specific mRNAs in the nucleus, indicating that eIF-5A may be responsible for shuttling specific classes of mRNAs from the nucleus to the cytoplasm. Liu & Tartakoff (1997) Supplement to *Molecular Biology of the Cell*,
10 **8**, 426a. Abstract No. 2476, 37th American Society for Cell Biology Annual Meeting. The accumulation of eIF-5A at nuclear pore-associated intranuclear filaments and its interaction with a general nuclear export receptor further suggest that eIF-5A is a nucleocytoplasmic shuttle protein, rather than a component of polysomes. Rosorius et al. (1999) *J. Cell Science*, **112**, 2369-2380.

15 Expression of eIF-5A mRNA has been explored in various human tissues and mammalian cell lines. For example, changes in eIF-5A expression have been observed in human fibroblast cells after addition of serum following serum deprivation. Pang & Chen (1994) *J. Cell Physiol.*, **160**, 531-538. Age-related decreases in deoxyhypusine synthase activity and abundance of precursor eIF-5A
20 have also been observed in senescing fibroblast cells, although the possibility that this reflects averaging of differential changes in isoforms was not determined. Chen & Chen (1997b) *J. Cell Physiol.*, **170**, 248-254.

Studies have shown that eIF-5A may be the cellular target of viral proteins such as the human immunodeficiency virus type 1 Rev protein and human T cell
25 leukemia virus type 1 Rex protein. Ruhl et al. (1993) *J. Cell Biol.*, **123**, 1309-1320; Katahira et al. (1995) *J. Virol.*, **69**, 3125-3133. Preliminary studies indicate that eIF-5A may target RNA by interacting with other RNA-binding proteins such as Rev, suggesting that these viral proteins may recruit eIF-5A for viral RNA processing. Liu et al. (1997) *Biol. Signals*, **6**, 166-174.

Deoxyhypusine synthase and eIF-5A are known to play important roles in key cellular processes including cell growth and senescence. For example, antisense reduction of deoxyhypusine synthase expression in plants results in delayed senescence of leaves and fruits, indicating that there is a senescence-inducing isoform of eIF-5A in plants. See WO 01/02592; PCT/US01/44505; U.S. Application No. 09/909,796. Inactivation of the genes for deoxyhypusine synthase or eIF-5A in yeast results in inhibition of cell division. Schnier et al. (1991) *Mol. Cell. Biol.*, **11**, 3105-3114; Sasaki et al. (1996) *FEBS Lett.*, **384**, 151-154; Park et al. (1998) *J. Biol. Chem.*, **273**, 1677-1683.

Spermidine analogs have been successfully used to inhibit deoxyhypusine synthase *in vitro*, as well as to inhibit the formation of hypusine *in vivo*, which is accompanied by an inhibition of protein synthesis and cell growth. Jakus et al. (1993) *J. Biol. Chem.*, **268**, 13151-13159; Park et al. (1994) *J. Biol. Chem.*, **269**, 27827-27832. Polyamines themselves, in particular putrescine and spermidine, also appear to play important roles in cellular proliferation and differentiation. Tabor & Tabor (1984) *Annu. Rev. Biochem.*, **53**, 749-790; Pegg (1988) *Cancer Res.*, **48**, 759-774. For example, yeast mutants in which the polyamine biosynthesis pathway has been blocked are unable to grow unless provided with exogenous polyamines. Cohn et al. (1980) *J. Bacteriol.*, **134**, 208-213.

Polyamines have also been shown to protect cells from the induction of apoptosis. For example, apoptosis of thymocytes has been blocked by exposure to spermidine and spermine, the mechanism of which appears to be the prevention of endonuclease activation. Desiderio et al. (1995) *Cell Growth Differ.*, **6**, 505-513; Brune et al. (1991) *Exp. Cell Res.*, **195**, 323-329. In addition, exogenous polyamines have been shown to repress B cell receptor-mediated apoptosis as well as apoptosis in the unicellular parasite, *Trypanosoma cruzi*. Nitta et al. (2001) *Exptl. Cell Res.*, **265**, 174-183; Piacenza et al. (2001) *Proc. Natl. Acad. Sci., USA*, **98**, 7301-7306. Low concentrations of spermine and spermidine have also been observed to reduce the number of nerve cells lost during normal development of newborn rats, as well as protect the brain from

neuronal damage during cerebral ischaemia. Gilad et al. (1985) *Brain Res.*, **348**, 363-366; Gilad & Gilad (1991) *Exp. Neurol.*, **111**, 349-355. Polyamines also inhibit senescence, a form of programmed cell death, of plant tissues. Spermidine and putrescine have been shown to delay post-harvest senescence of carnation
5 flowers and detached radish leaves. Wang & Baker (1980) *HortScience*, **15**, 805-806 (carnation flowers); Altman (1982) *Physiol. Plant.*, **54**, 189-193 (detached radish leaves).

In other studies, however, induction of apoptosis has been observed in response to exogenous polyamines. For example, human breast cancer cell lines
10 responded to a polyamine analogue by inducing apoptosis, and excess putrescine has been shown to induce apoptosis in DH23A cells. McCloskey et al. (1995) *Cancer Res.*, **55**, 3233-3236; Tome et al. (1997) *Biochem. J.*, **328**, 847-854.

The results from these experiments with polyamines collectively suggest that existence of specific isoforms of eIF-5A play a role in induction of apoptosis.
15 Specifically, the data are consistent with the view that there is an apoptosis-specific isoform of eIF-5A, which is activated by DHS. The fact that this DHS reaction requires spermidine is consistent with the finding that polyamines have been shown to elicit activation of caspase, a key executor of apoptosis-related proteolysis. Stefanelli et al. (2000) *Biochem. J.*, **347**, 875-880; Stefanelli et al.
20 (1999) *FEBS Lett.*, **451**, 95-98. In a similar vein, inhibitors of polyamine synthesis can delay apoptosis. Das et al. (1997) *Oncol. Res.*, **9**, 565-572; Monti et al. (1998) *Life Sci.*, **72**, 799-806; Ray et al. (2000) *Am. J. Physiol.*, **278**, C480-C489; Packham & Cleveland (1994) *Mol. Cell Biol.*, **14**, 5741-5747.

The finding that exogenous polyamines both inhibit and promote
25 apoptosis can be explained by the fact that, depending upon the levels applied, they can either inhibit the DHS reaction leading to activation of eIF-5A and hence impede apoptosis, or induce apoptosis by reason of being toxic. The finding that low concentrations of exogenous polyamines block apoptosis in plant and animal systems is consistent with the fact that low concentrations of polyamines and their
30 analogues act as competitive inhibitors of the DHS reaction. Indeed, even

exogenous spermidine, which is a substrate for the DHS reaction, will impede the reaction through substrate inhibition. Jakus et al. (1993) *J. Biol. Chem.*, **268**, 13153-13159.

However, all polyamines, and their analogues, are toxic at high concentrations and are able to induce apoptosis. This occurs despite their ability to inhibit activation of the putative apoptosis-specific isoform of eIF-5A for two reasons. First, activated eIF-5A has a long half-life. Torrelio et al. (1987) *Biochem. Biophys. Res. Commun.*, **145**, 1335-1341; Dou & Chen (1990) *Biochim. Biophys. Acta.*, **1036**, 128-137. Accordingly, depletion of activated apoptosis-specific eIF-5A arising from inhibition of deoxyhypusine synthase activity may not occur in time to block apoptosis caused by the toxic effects of spermidine. Second, polyamines are competitive inhibitors of the deoxyhypusine reaction and hence not likely to completely block the reaction even at concentrations that are toxic.

The present invention relates to cloning of an eIF-5A cDNA that is up regulated immediately before the induction of apoptosis. This apoptosis-specific eIF-5A is likely to be a suitable target for intervention in apoptosis-causing disease states since it appears to act at the level of post-transcriptional regulation of downstream effectors and transcription factors involved in the apoptotic pathway. Specifically, the apoptosis-specific eIF-5A appears to selectively facilitate the translocation of mRNAs encoding downstream effectors and transcription factors of apoptosis from the nucleus to the cytoplasm, where they are subsequently translated. The ultimate decision to initiate apoptosis appears to stem from a complex interaction between internal and external pro- and anti-apoptotic signals. Lowe & Lin (2000) *Carcinogenesis*, **21**, 485-495. Through its ability to facilitate the translation of downstream apoptosis effectors and transcription factors, the apoptosis-related eIF-5A appears to tip the balance between these signals in favor of apoptosis.

As described previously, it is well established that anticancer agents induce apoptosis and that alterations in the apoptotic pathways can attenuate

drug-induced cell death. Schmitt & Lowe (1999) *J. Pathol.*, **187**, 127-137. For example, many anticancer drugs upregulate p53, and tumor cells that have lost p53 develop resistance to these drugs. However, nearly all chemotherapy agents can induce apoptosis independently of p53 if the dose is sufficient, indicating that
5 even in drug-resistant tumors, the pathways to apoptosis are not completely blocked. Wallace-Brodeur & Lowe (1999) *Cell Mol. Life Sci.*, **55**, 64-75. This suggests that induction of apoptosis eIF-5A, even though it may not correct the mutated gene, may be able to circumvent the p53-dependent pathway and induce apoptosis by promoting alternative pathways.

10 Induction of apoptosis-related eIF-5A has the potential to selectively target cancer cells while having little or no effect on normal neighboring cells. This arises because mitogenic oncogenes expressed in tumor cells provide an apoptotic signal in the form of specific species of mRNA that are not present in normal cells. Lowe et al. (1993) *Cell*, **74**, 954-967; Lowe & Lin (2000)
15 *Carcinogenesis*, **21**, 485-495. For example, restoration of wild-type p53 in p53-mutant tumor cells can directly induce apoptosis as well as increase drug sensitivity in tumor cell lines and xenographs. (Spitz et al., 1996; Badie et al., 1998).

The selectivity of apoptosis-eIF-5A arises from the fact that it selectively
20 facilitates translation of mRNAs for downstream apoptosis effectors and transcription factors by mediating their translocation from the nucleus into the cytoplasm. Thus, for apoptosis eIF-5A to have an effect, mRNAs for these effectors and transcription factors have to be transcribed. Inasmuch as these mRNAs would be transcribed in cancer cells, but not in neighboring normal cells,
25 it is to be expected that apoptosis eIF-5A would promote apoptosis in cancer cells but have minimal, if any, effect on normal cells. Thus, restoration of apoptotic potential in tumor cells with apoptosis-related eIF-5A may decrease the toxicity and side effects experienced by cancer patients due to selective targeting of tumor cells. Induction of apoptotic eIF-5A also has the potential to potentiate the
30 response of tumor cells to anti-cancer drugs and thereby improve the

effectiveness of these agents against drug-resistant tumors. This in turn could result in lower doses of anti-cancer drugs for efficacy and reduced toxicity to the patient.

5

SUMMARY OF INVENTION

The present invention provides isolated and/or purified rat apoptosis-specific eIF-5A and DHS nucleic acids and polypeptides and antisense oligonucleotides and expression vectors of apoptosis-specific eIF-5A and DHS. The present invention also provides isolated and/or purified human eIF-5A2 (also referred to herein as proliferating eIF-5A. The present invention also provides methods of modulating apoptosis using apoptosis-specific eIF-5A and DHS.

10

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the nucleotide sequence and derived amino acid sequence of the 3' end of rat apoptosis-specific eIF-5A.

15

Figure 2 depicts the nucleotide sequence and derived amino acid sequence of the 5' end of rat apoptosis-specific eIF-5A cDNA.

Figure 3 depicts the nucleotide sequence of rat corpus luteum apoptosis-specific eIF-5A full-length cDNA.

20

Figure 4 depicts the nucleotide sequence and derived amino acid sequence of the 3' end of rat apoptosis-specific DHS cDNA.

Figure 5 is an alignment of the full-length nucleotide sequence of rat corpus luteum apoptosis-specific eIF-5A cDNA with the nucleotide sequence of human eIF-5A (Accession number BC000751 or NM_001970, SEQ ID NO:3).

25

Figure 6 is an alignment of the full-length nucleotide sequence of rat corpus luteum apoptosis-specific eIF-5A cDNA with the nucleotide sequence of human eIF-5A (Accession number NM-020390, SEQ ID NO:4).

Figure 7 is an alignment of the full-length nucleotide sequence of rat corpus luteum apoptosis-specific eIF-5A cDNA with the nucleotide sequence of

mouse eIF-5A (Accession number BC003889). Mouse nucleotide sequence (Accession number BC003889) is SEQ ID NO:5.

Figure 8 is an alignment of the derived full-length amino acid sequence of rat corpus luteum apoptosis-specific eIF-5A with the derived amino acid
5 sequence of human eIF-5A (Accession number BC000751 or NM_001970).

Figure 9 is an alignment of the derived full-length amino acid sequence of rat corpus luteum apoptosis-specific eIF-5A with the derived amino acid sequence of human eIF-5A (Accession number NM_020390).

Figure 10 is an alignment of the derived full-length amino acid sequence
10 of rat corpus luteum apoptosis-specific eIF-5A with the derived amino acid sequence of mouse eIF-5A (Accession number BC003889).

Figure 11 is an alignment of the partial-length nucleotide sequence of rat corpus luteum apoptosis-specific DHS cDNA with the nucleotide sequence of human DHS (Accession number BC000333, SEQ ID NO:8).

15 Figure 12 is a restriction map of rat corpus luteum apoptosis-specific eIF-5A cDNA.

Figure 13 is a restriction map of the partial-length rat apoptosis-specific DHS cDNA.

Figure 14 is a Northern blot (Figure 14A) and an ethidium bromide
20 stained gel (Figure 14B) of total RNA probed with the ³²P-dCTP-labeled 3'-end of rat corpus luteum apoptosis-specific eIF-5A cDNA.

Figure 15 is a Northern blot (Figure 15A) and an ethidium bromide stained gel (Figure 15B) of total RNA probed with the ³²P-dCTP-labeled 3'-end of rat corpus luteum apoptosis-specific DHS cDNA.

25 Figure 16 depicts a DNA laddering experiment in which the degree of apoptosis in superovulated rat corpus lutea was examined after injection with PGF-2 α .

Figure 17 is an agarose gel of genomic DNA isolated from apoptosing rat corpus luteum showing DNA laddering after treatment of rats with PGF F-2 α .

Figure 18 depicts a DNA laddering experiment in which the degree of apoptosis in dispersed cells of superovulated rat corpora lutea was examined in rats treated with spermidine prior to exposure to prostaglandin F-2 α (PGF-2 α).

Figure 19 depicts a DNA laddering experiment in which the degree of apoptosis in superovulated rat corpus lutea was examined in rats treated with spermidine and/or PGF-2 α .

Figure 20 is a Southern blot of rat genomic DNA probed with ^{32} P-dCTP-labeled partial-length rat corpus luteum apoptosis-specific eIF-5A cDNA.

Figure 21 depicts pHM6, a mammalian epitope tag expression vector (Roche Molecular Biochemicals).

Figure 22 is a Northern blot (Figure 23A) and ethidium bromide stained gel (Figure 23B) of total RNA isolated from COS-7 cells after induction of apoptosis by withdrawal of serum probed with the ^{32} P-dCTP-labeled 3'-untranslated region of rat corpus luteum apoptosis-specific DHS cDNA.

Figure 23 is a flow chart illustrating the procedure for transient transfection of COS-7 cells.

Figure 24 is a Western blot of transient expression of foreign proteins in COS-7 cells following transfection with pHM6.

Figure 25 illustrates enhanced apoptosis as reflected by increased caspase activity when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation.

Figure 26 illustrates enhanced apoptosis as reflected by increased DNA fragmentation when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation.

Figure 27 illustrates detection of apoptosis as reflected by increased nuclear fragmentation when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation.

Figure 28 illustrates enhanced apoptosis as reflected by increased nuclear fragmentation when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation.

Figure 29 illustrates detection of apoptosis as reflected by phosphatidylserine exposure when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation.

5 Figure 30 illustrates enhanced apoptosis as reflected by increased phosphatidylserine exposure when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation.

10 Figure 31 illustrates enhanced apoptosis as reflected by increased nuclear fragmentation when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation.

Figure 32 illustrates enhanced apoptosis when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation.

15 Figure 33 illustrates down-regulation of Bcl-2 when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation. Figure 31A is the Coomassie-blue-stained protein blot; Figure 31B is the corresponding Western blot.

20 Figure 34 is a Coomassie-blue-stained protein blot and the corresponding Western blot of COS-7 cells transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the antisense orientation using Bcl-2 as a probe.

25 Figure 35 is a Coomassie-blue-stained protein blot and the corresponding Western blot of COS-7 cells transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation using c-Myc as a probe.

30 Figure 36 is a Coomassie-blue-stained protein blot and the corresponding Western blot of COS-7 cells transiently transfected with pHM6 containing full-length rat apoptosis-specific eIF-5A in the sense orientation when p53 is used as a probe.

Figure 37 is a Coomassie-blue-stained protein blot and the corresponding Western blot of expression of pHM6-full-length rat apoptosis-specific eIF-5A in COS-7 cells using an anti-[HA]-peroxidase probe and a Coomassie-blue-stained protein blot and the corresponding Western blot of expression of pHM6-full-length rat apoptosis-specific eIF-5A in COS-7 cells when a p53 probe is used.

Figure 38 is an alignment of human eIF5A2 isolated from RKO cells with the sequence of human eIF5A2 (Genbank accession number XM_113401).

Figure 39 is a graph depicting the percentage of apoptosis occurring in RKO and RKO-E6 cells following transient transfection. RKO and RKO-E6 cells were transiently transfected with pHM6-LacZ or pHM6-eIF5A1. RKO cells treated with Actinomycin D and transfected with pHM6-eIF5A1 showed a 240% increase in apoptosis relative to cells transfected with pHM6-LacZ that were not treated with Actinomycin D. RKO-E6 cells treated with Actinomycin D and transfected with pHM6-eIF5A1 showed a 105% increase in apoptosis relative to cells transfected with pHM6-LacZ that were not treated with Actinomycin D.

Figure 40 is a graph depicting the percentage of apoptosis occurring in RKO cells following transient transfection. RKO cells were transiently transfected with pHM6-LacZ, pHM6-eIF5A1, pHM6-eIF5A2, or pHM6-truncated eIF5A1. Cells transfected with pHM6-eIF5A1 showed a 25% increase in apoptosis relative to control cells transfected with pHM6-LacZ. This increase was not apparent for cells transfected with pHM6-eIF5A2 or pHM6-truncated eIF5A1.

Figure 41 is a graph depicting the percentage of apoptosis occurring in RKO cells following transient transfection. RKO cells were either left untransfected or were transiently transfected with pHM6-LacZ or pHM6-eIF5A1. After correction for transfection efficiency, 60 % of the cells transfected with pHM6-eIF5A1 were apoptotic.

Figure 42 provides the results of a flow cytometry analysis of RKO cell apoptosis following transient transfection. RKO cells were either left untransfected or were transiently transfected with pHM6-LacZ, pHM6-eIF5A1, pHM6-eIF5A2, or pHM6-truncated eIF5A1. The table depicts the percentage of cells undergoing

apoptosis calculated based on the area under the peak of each gate. After correction for background apoptosis in untransfected cells and for transfection efficiency, 80% of cells transfected with pHM6-eIF5A1 exhibited apoptosis. Cells transfected with pHM6-LacZ, pHM6-eIF5A2 or pHM6-truncated eIF5A1 exhibited only background levels of apoptosis.

Figure 43 provides Western blots of protein extracted from RKO cells treated with 0.25 µg/ml Actinomycin D for 0, 3, 7, 24, and 48 hours. The top panel depicts a Western blot using anti-p53 as the primary antibody. The middle panel depicts a Western blot using anti-eIF5A1 as the primary antibody. The bottom panel depicts the membrane used for the anti-eIF5A1 blot stained with Coomassie blue following chemiluminescent detection to demonstrate equal loading. p53 and eIF5A1 are both upregulated by treatment with Actinomycin D.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based, in part, on the discovery and characterization of a full-length cDNA encoding an eIF-5A isolated from rat corpus luteum, which is involved in apoptosis (apoptosis-specific). Therefore, in one embodiment, the present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding a rat apoptosis-specific eIF-5A polypeptide. Also provided by the present invention is a purified polypeptide comprising an amino acid sequence of a rat apoptosis-specific eIF-5A polypeptide. Rat apoptosis-specific eIF-5A polypeptide means any polypeptide specific to rats that is differentially expressed in apoptosing cells and that results from formation of a deoxyhypusine residue by the transfer of the 4-aminobutyl moiety of spermidine to the α-amino group of a specific conserved lysine of a precursor eIF-5A catalyzed by deoxyhypusine synthase and hydroxylation of this 4-aminobutyl moiety by deoxyhypusine hydroxylase to form hypusine, thereby activating eIF-5A.

In addition, the nucleic acid and polypeptide rat apoptosis-specific eIF-5A sequences of the present invention can be used to isolate apoptosis-specific

nucleic acids and polypeptides from other cells, tissues, organs, or animals using guidance provided herein and techniques well known to those skilled in the art. The present invention also provides nucleic acid molecules that are suitable as primers or hybridization probes for the detection of nucleic acids encoding a rat
5 apoptosis-specific eIF-5A polypeptide of the invention.

The nucleic acids of the present invention can be DNA, RNA, DNA/RNA duplexes, protein-nucleic acid (PNA), or derivatives thereof. As used herein, a nucleic acid or polypeptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other
10 chemicals. It should be appreciated that the term isolated or purified does not refer to a library-type preparation containing a myriad of other sequence fragments. The nucleic acid or polypeptide of the present invention can be purified to homogeneity or other degrees of purity. The level of purification will be based on the intended use. The critical feature is that the preparation allows
15 for the desired function of the nucleic acid or polypeptide, even if in the presence of considerable amounts of other components.

The isolated polypeptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. For example recombinant
20 production of proteins involves cloning a nucleic acid molecule encoding either the apoptosis inducing eIF-5A or DHS into an expression vector. The expression vector is introduced into a host cell and the protein is expressed in the host cell. The protein can then be isolated from the cells by any appropriate purification scheme using standard protein purification techniques.

25 Preferably, the isolated nucleic acid encoding a rat apoptosis-specific eIF-5A polypeptide of the present invention has a nucleotide sequence of SEQ ID NO:1 and the purified polypeptide of the present invention has an amino acid sequence of SEQ ID NO:2. The present inventive rat apoptosis-specific eIF-5A nucleic acids and polypeptides also encompass sequences that have substantial

sequence identity or homology to SEQ ID NO:1 and SEQ ID NO:2, respectively, as well as functional derivatives and variants thereof.

As used herein, the term "substantial sequence identity" or "substantial homology" is used to indicate that a sequence exhibits substantial structural or functional equivalence with another sequence. Any structural or functional differences between sequences having substantial sequence identity or substantial homology will be *de minimus*; that is, they will not affect the ability of the sequence to function as indicated in the desired application. Differences may be due to inherent variations in codon usage among different species, for example.

Structural differences are considered *de minimus* if there is a significant amount of sequence overlap or similarity between two or more different sequences or if the different sequences exhibit similar physical characteristics even if the sequences differ in length or structure. Such characteristics include, for example, the ability to hybridize under defined conditions, or in the case of proteins, immunological crossreactivity, similar enzymatic activity, etc. The skilled practitioner can readily determine each of these characteristics by art known methods.

Additionally, two nucleotide sequences are "substantially complementary" if the sequences have at least about 70 percent or greater, more preferably 80 percent or greater, even more preferably about 90 percent or greater, and most preferably about 95 percent or greater sequence similarity between them. Two amino acid sequences are substantially homologous if they have at least 50%, preferably at least 70%, more preferably at least 80%, even more preferably at least 90%, and most preferably at least 95% similarity between the active, or functionally relevant, portions of the polypeptides.

To determine the percent identity of two sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, at least 30%, 40%, 50%, 60%, 70%, 80%,

or 90% or more of the length of a reference sequence is aligned for comparison purposes. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the
5 corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap,
10 which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome
15 Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991).

20 The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (1990) *J. Mol Biol.* 215:403-10. BLAST
25 nucleotide searches can be performed with the NBLAST program. BLAST protein searches can be performed with the XBLAST program to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing

BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

The term "functional derivative" of a nucleic acid is used herein to mean a homolog or analog of the gene or nucleotide sequence. A functional derivative
5 may retain at least a portion of the function of the given gene, which permits its utility in accordance with the invention. "Functional derivatives" of the apoptosis-specific eIF-5A polypeptide as described herein are fragments, variants, analogs, or chemical derivatives of apoptosis-specific eIF-5A that retain at least a portion of the apoptosis-specific eIF-5A activity or immunological cross
10 reactivity with an antibody specific for apoptosis-specific eIF-5A. A fragment of the apoptosis-specific eIF-5A polypeptide refers to any subset of the molecule.

Functional variants can also contain substitutions of similar amino acids that result in no change or an insignificant change in function. Amino acids that are essential for function can be identified by methods known in the art, such as
15 site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham et al. (1989) Science 244:1081-1085). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as kinase activity or in assays such as an in vitro proliferative activity. Sites that are critical for binding partner/substrate
20 binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith et al. (1992) J. Mol. Biol. 224:899-904; de Vos et al. (1992) Science 255:306-312).

A "variant" refers to a molecule substantially similar to either the entire gene or a fragment thereof, such as a nucleotide substitution variant having one or
25 more substituted nucleotides, but which maintains the ability to hybridize with the particular gene or to encode mRNA transcript which hybridizes with the native DNA. A "homolog" refers to a fragment or variant sequence from a different animal genus or species. An "analog" refers to a non-natural molecule substantially similar to or functioning in relation to the entire molecule, a variant
30 or a fragment thereof.

Variant peptides include naturally occurring variants as well as those manufactured by methods well known in the art. Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other
5 proteins based on sequence and/or structural homology to the eIF-5A or DHS proteins of the present invention. The degree of homology/identity present will be based primarily on whether the protein is a functional variant or non-functional variant, the amount of divergence present in the paralog family and the evolutionary distance between the orthologs.

10 Non-naturally occurring variants of the eIF-5A or DHS proteins of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the proteins. For example, one class of substitutions are conserved amino acid substitution. Such substitutions are those that substitute a
15 given amino acid in a protein by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln; exchange of the basic
20 residues Lys and Arg; and replacements among the aromatic residues Phe and Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie et al., Science 247:1306-1310 (1990).

Alternatively, but also preferably, the nucleic acid encoding a rat apoptosis-specific eIF-5A polypeptide of the present invention hybridizes under
25 highly stringent conditions with a nucleotide sequence that is complementary to that of SEQ ID NO:1. The term "hybridization" as used herein is generally used to mean hybridization of nucleic acids at appropriate conditions of stringency as would be readily evident to those skilled in the art depending upon the nature of the probe sequence and target sequences. Conditions of hybridization and
30 washing are well known in the art, and the adjustment of conditions depending

upon the desired stringency by varying incubation time, temperature and/or ionic strength of the solution are readily accomplished. *See, e.g.* Sambrook, J. et al., Molecular Cloning: A Laboratory Manual, 2nd edition, Cold Spring Harbour Press, Cold Spring Harbor, New York, 1989.

5 The choice of conditions is dictated by the length of the sequences being hybridized, in particular, the length of the probe sequence, the relative G-C content of the nucleic acids and the amount of mismatches to be permitted. Low stringency conditions are preferred when partial hybridization between strands that have lesser degrees of complementarity is desired. When perfect or near
10 perfect complementarity is desired, high stringency conditions are preferred. For typical high stringency conditions, the hybridization solution contains 6X S.S.C., 0.01 M EDTA, 1X Denhardt's solution and 0.5% SDS. Hybridization is carried out at about 68°C for about 3 to 4 hours for fragments of cloned DNA and for about 12 to 16 hours for total eucaryotic DNA. For lower stringencies, the
15 temperature of hybridization is reduced to about 42° C below the melting temperature (T_m) of the duplex. The T_m is known to be a function of the G-C content and duplex length as well as the ionic strength of the solution.

As used herein, the phrase "hybridizes to a corresponding portion" of a DNA or RNA molecule means that the molecule that hybridizes, e.g.,
20 oligonucleotide, polynucleotide, or any nucleotide sequence (in sense or antisense orientation) recognizes and hybridizes to a sequence in another nucleic acid molecule that is of approximately the same size and has enough sequence similarity thereto to effect hybridization under appropriate conditions. For example, a 100 nucleotide long sense molecule will recognize and hybridize to an
25 approximately 100 nucleotide portion of a nucleotide sequence, so long as there is about 70% or more sequence similarity between the two sequences. It is to be understood that the size of the "corresponding portion" will allow for some mismatches in hybridization such that the "corresponding portion" may be smaller or larger than the molecule which hybridizes to it, for example 20-30% larger or
30 smaller, preferably no more than about 12-15% larger or smaller.

In addition, functional variants of polypeptides can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham et al., Science 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity or in assays.

For example, an analog of apoptosis-specific eIF-5A refers to a non-natural protein or peptidomimetic substantially similar to either the entire protein or a fragment thereof. Chemical derivatives of apoptosis-specific eIF-5A contain additional chemical moieties not normally a part of the peptide or peptide fragment. Modifications can be introduced into peptide or fragment thereof by reacting targeted amino acid residues of the peptide with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues.

The initial discovery and characterization of a full-length cDNA encoding an apoptosis-specific eIF-5A isolated from rat corpus luteum led to the discovery and characterization of a partial-length cDNA clone encoding a DHS, which is also isolated from rat corpus luteum and involved in apoptosis. Accordingly, in an additional embodiment, the present invention provides an isolated nucleic acid comprising a nucleotide sequence encoding a rat apoptosis-specific DHS polypeptide. Also provided is a purified polypeptide comprising an amino acid sequence of a rat apoptosis-specific DHS polypeptide. Rat apoptosis-specific DHS polypeptide means any suitable polypeptide specific to rats that is differentially expressed in apoptosing cells and that catalyzes formation of a deoxyhypusine residue by the transfer of the 4-aminobutyl moiety of spermidine to the α -amino group of a specific conserved lysine of inactive eIF-5A to form deoxyhypusine, thereby activating eIF-5A.

Preferably, the isolated nucleic acid encoding a rat apoptosis-specific DHS polypeptide of the present invention has a nucleotide sequence of SEQ ID

NO:22 and the purified polypeptide of the present invention has an amino acid sequence of SEQ ID NO:23. The present inventive rat apoptosis-specific DHS nucleic acids and polypeptides also encompass sequences that have substantial sequence identity or homology to SEQ ID NO:22 and SEQ ID NO:23,

5 respectively, as well as functional derivatives and variants thereof, which have been described previously. Alternatively, and also preferably, the isolated nucleic acid of the present invention has a nucleotide sequence that hybridizes under highly stringent conditions with the complement of SEQ ID NO:22, which also has been described previously.

10 As is the case with the nucleic acids and polypeptides of the rat apoptosis-specific eIF-5A sequences described herein, the nucleic acids and polypeptides of the rat apoptosis-specific DHS sequences of the present invention can be used to isolate apoptosis-specific DHS nucleic acids and polypeptides from other animals, including humans. Isolation of such DHS sequences from animals and
15 human can be achieved using art known methods and guidance provided herein, based on sequence similarities of at least 80% across species. The present invention also provides nucleic acid molecules that are suitable as primers or hybridization probes for the detection of nucleic acids encoding a rat apoptosis-specific DHS polypeptide of the invention.

20 Apoptosis-specific eIF-5A and DHS are suitable targets for regulation of apoptosis, including apoptosis underlying disease processes, since it likely acts in the post-transcriptional regulation of downstream effectors and transcription factors involved in the apoptotic pathway. Thus, the present invention also provides methods of modulating apoptosis in a cell by administering to the cell an
25 agent that modulates apoptosis-specific eIF-5A and/or DHS function. It should be appreciated by one of skill in the art that the agent can be one that modulates only apoptosis-specific eIF-5A function, only apoptosis-specific DHS function alone, or both apoptosis-specific eIF-5A and DHS function.

Apoptosis can be modulated by any suitable alteration in the normal level
30 of apoptosis-specific eIF-5A and/or DHS function in the cell. As intended herein,

modification or alteration can be complete or partial and can include a change in transcriptional or translational control or other change altering apoptosis-specific eIF-5A and/or DHS function in the cell. Apoptosis-specific eIF-5A or DHS function means any activity relating to formation of a deoxyhypusine residue by the transfer of the 4-aminobutyl moiety of spermidine to the α -amino group of a specific conserved lysine of a precursor eIF-5A, which is catalyzed by DHS, and hydroxylation of this 4-aminobutyl moiety by deoxyhypusine hydroxylase to form hypusine, thereby activating eIF-5A.

In one embodiment of the present invention, the agent can inhibit apoptosis-specific eIF-5A and/or DHS function, thereby inhibiting apoptosis. Inhibiting apoptosis means any decrease, in intensity and/or number, and/or delay in onset of any or all of the well-defined morphological features characteristic of apoptosis, such as, for example, cell shrinkage, chromatin condensation, nuclear fragmentation, and membrane blebbing.

One agent that can inhibit apoptosis-specific eIF-5A and/or DHS function is an antisense oligonucleotide. Preferably, the antisense oligonucleotide has a nucleotide sequence encoding a portion of an apoptosis-specific eIF-5A polypeptide and/or an apoptosis-specific DHS polypeptide. Many suitable nucleic acid sequences encoding an apoptosis-specific eIF-5A polypeptide and/or DHS polypeptide are known in the art. For example, SEQ ID NOS:1, 3, 4, 5, 11, 15, 19, 20, and 21 (apoptosis-specific eIF-5A nucleic acid sequences), SEQ ID NOS:6 and 8 (apoptosis-specific DHS nucleic acid sequences), SEQ ID NOS:12 and 16 eIF-5A (apoptosis-specific polypeptide sequences), and SEQ ID NO:7 (apoptosis-specific DHS polypeptide sequences), or portions thereof, provide suitable sequences. Others suitable sequences can be found using the known sequences as probes according to the methods described herein.

Accordingly, the present invention also provides antisense oligonucleotides encoding a portion of an apoptosis-specific eIF-5A polypeptide and/or an apoptosis-specific DHS polypeptide, or a complement thereof. The antisense oligonucleotides of the present invention can be in the form of RNA or

DNA, e.g., cDNA, genomic DNA, or synthetic RNA or DNA. The DNA can be double-stranded or single stranded, and if single stranded can be the coding strand or non-coding strand. The specific hybridization of an oligomeric compound with its target nucleic acid, resulting in interference with the normal function of the nucleic acid, is generally referred to as "antisense." The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which can be engaged in or facilitated by the RNA. The overall effect of such antisense oligonucleotide is inhibiting of expression of apoptosis-specific eIF-5A and/or DHS and/or the amount of activated apoptosis-specific eIF-5A produced.

Alternatively, the activation of apoptosis-specific eIF-5A by apoptosis-specific DHS can be inhibited by administering chemical agents that inhibit the DHS enzymatic reaction. For example, the onset of DNA laddering reflecting apoptosis is delayed in rat corpus luteum when the animals are treated with spermidine, an inhibitor of the DHS reaction after induction of apoptosis by injection of PGF-2 α (FIGS. 18-19). Jakus et al., (1993) *J. Biol. Chem.* 268: 13151-13159.

Apoptosis also can be inhibited or substantially decreased by adding agents that degrade apoptosis-specific eIF-5A DNA, RNA, or protein, or that degrade apoptosis-specific DHS DNA, RNA, or protein, thereby preventing the activation of apoptosis-specific eIF-5A by apoptosis-specific DHS. In another embodiment of the invention, inhibition of expression of endogenous mammalian apoptosis-specific DHS, apoptosis-specific eIF-5A, or both, are affected through the use of ribozymes. Examples of suitable drugs include those that inhibit the activation of apoptosis-specific eIF-5A by apoptosis-specific DHS, those that inhibit the activation of apoptosis-specific eIF-5A by deoxyhypusine hydroxylase, those that inhibit transcription and/or translation of apoptosis-specific DHS, those that inhibit transcription and/or translation of apoptosis-specific deoxyhypusine

hydroxylase, and those that inhibit transcription or translation of apoptosis-specific eIF-5A. Examples of drugs that inhibit the activation of eIF-5A by apoptosis-specific DHS are spermidine, 1,3-Diamino-propane, 1,4-Diamino-butane (putrescine), 1,7-Diamino-heptane, or 1,8-Diamino-octane.

5 It is also possible to inhibit apoptosis-specific eIF-5A by inactivating the gene coding for apoptosis-specific eIF-5A in a cell. Such inactivation can occur by deleting the gene in the cell or by introducing a deletion or mutation into the gene and thereby inactivating the gene. The gene can also be inactivated by inserting into the gene another DNA fragment such that expression of the
10 endogenous apoptosis-specific eIF-5A protein does not occur. Likewise, it is possible to inhibit activation of apoptosis-specific eIF-5A by inactivating the gene coding for apoptosis-specific DHS in a cell. Methods for introducing mutations, such as deletions and insertions, into genes in eukaryotic cells are known in the art, e.g., U.S. Patent No. 5,464,764. Oligonucleotides and expression vectors
15 useful for mutation of genes in cells can be made according to methods known in the art and guidance provided herein; for example, methods useful for making and expressing antisense oligonucleotides can be used to make oligonucleotides and expression vectors useful for mutating genes in cells.

 It is also possible to inhibit expression of apoptosis-specific eIF-5A by
20 suppressing expression of the gene coding for apoptosis-specific eIF-5A in a cell. Such inactivation can be accomplished via cosuppression, e.g., by introducing nucleotide sequence(s) coding for apoptosis-specific eIF-5A into a cell such that cosuppression occurs. Likewise, it is possible to inhibit activation of apoptosis-specific eIF-5A by suppressing the expression of the gene coding for apoptosis-specific DHS in a cell via cosuppression. Oligonucleotides and expression
25 vectors useful for cosuppression can be made according to methods known in the art and guidance provided herein; for example, methods useful for making and expressing antisense oligonucleotides can be used to make oligonucleotides and expression vectors useful for cosuppression. Methods for cosuppression are
30 known in the art, e.g., U.S. Patent No. 5,686,649.

One result of the inhibition (through, e.g., antisense, mutation, or cosuppression) is a reduction in the amount of endogenous translatable apoptosis-specific eIF-5A or DHS-encoding mRNA. Consequently, the amount of apoptosis-specific DHS protein produced is reduced, thereby reducing the amount of activated eIF-5A, which in turn reduces translation of apoptosis-specific proteins. Apoptosis is thus inhibited or delayed, since *de novo* protein synthesis is required for the onset of apoptosis.

In another embodiment of the present invention, the agent can induce apoptosis-specific eIF-5A or DHS function, thereby inducing apoptosis. Inducing apoptosis means any increase, in intensity or number, or acceleration in onset of any or all of the well-defined morphological features characteristic of apoptosis, such as, for example, cell shrinkage, chromatin condensation, nuclear fragmentation, and membrane blebbing.

Any suitable agent that induces apoptosis-specific eIF-5A and/or DHS function can be used. It is appreciated by one of skill in the art that both the inactive and active forms of apoptosis-specific eIF-5A can be administered. If the inactive form, or hypusine-unmodified form, is administered, native apoptosis-specific DHS will activate the eIF-5A. Many suitable nucleic acid sequences encoding an apoptosis-specific eIF-5A polypeptide and/or DHS polypeptide are known in the art. For example, SEQ ID NOS:1, 3, 4, 5, 11, 15, 19, 20, and 21 (apoptosis-specific eIF-5A nucleic acid sequences), SEQ ID NOS:6 and 8 (apoptosis-specific DHS nucleic acid sequences), SEQ ID NOS:12 and 16 eIF-5A (apoptosis-specific polypeptide sequences), and SEQ ID NO:7 (apoptosis-specific DHS polypeptide sequences), or portions thereof, provide suitable sequences. Others suitable sequences can be found using the known sequences as probes according to the methods described herein.

For example, naked nucleic acids (naked DNA vectors such as oligonucleotides or plasmids), or polypeptides, including recombinantly produced polypeptides, can be administered to a cell. Recombinantly produced polypeptides means that the DNA sequences encoding the eIF-5A or the DHS

proteins are placed into a suitable expression vector, which is described in detail below. The host is transfected with the expression vector and thereafter produces the desired polypeptides. The polypeptides are then isolated from the host cell. Recombinant apoptosis-inducing eIF-5A protein can be made, for example, in
5 Chinese Hamster Ovary (CHO) cells and activated using recombinant DHS by those skilled in the art. Wang et al. (2001) *J. Biol. Chem.*, 276, 17541-17549; Eriksson et al., (2001) *Semin. Hematol.*, 38, 24-31. The polypeptides can also be synthetic, which are synthesized using known protein synthesis methods.

Polypeptide uptake can be facilitated using ligands, for example, a ligand
10 derived from anthrax that mediates uptake into a broad range of cells. Liu et al. (2001) *J. Biol. Chem.*, 276, 46326-46332. Recombinant protein can also be administered to target cells, tissues, and organs of mammals using liposomes. Liposomes occluding the protein are administered intravenously. Targeting can be achieved by incorporating ligands to specific cell receptors into the liposomes.
15 See, e.g., Kaneda, *Adv Drug Delivery Rev* 43: 197-205 (2000).

One preferred agent that can induce induce apoptosis-specific eIF-5A or DHS function is an expression vector. Accordingly, the present invention provides expression vectors having a promoter operably linked to a nucleic acid encoding an apoptosis-specific eIF-5A polypeptide and/or DHS polypeptide. The
20 expression vectors of the present invention can be in the form of RNA or DNA, e.g., cDNA, genomic DNA, or synthetic RNA or DNA. The DNA can be double-stranded or single stranded, and if single stranded can be the coding strand or non-coding strand. Any appropriate expression vector (see, e.g., Pouwels et al., *Cloning Vectors: A Laboratory Manual* (Elsevier, N.Y.: 1985)) can be used.
25 Preferably, the expression vector has a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific (related) eIF-5A polypeptide and/or apoptosis-specific (related) DHS polypeptide.

Within the expression vector, the desired nucleic acid and the promoter are operably linked such that the promoter is able to drive the expression of the
30 nucleic acid. Any suitable promoter can be used provided that the nucleic acid is

expressed. Examples of such suitable promoters include various viral promoters, eucaryotic promoters, and constitutively active promoters. As long as this operable linkage is maintained, the expression vector can include more than one nucleic acid (e.g., nucleic acids encoding both apoptosis-specific eIF-5A and/or
5 DHS). The expression vector can optionally include other elements, such as polyadenylation sequences, ribosome entry sites, transcriptional regulatory elements (e.g., enhancers, silencers, etc.), other sequences for enhancing the stability of the vector or transcript or the translation or processing of the desired transcript within the cells (e.g., secretion signals, leaders, etc.), or any other
10 suitable element.

Expression vector can be derived from viruses such as adenovirus, adeno-associated virus, herpesvirus, retrovirus or lentivirus. The expression vector of the present invention can be transfected into host cells, which include, but are not limited to, bacterial species, mammalian or insect host cell systems including
15 baculovirus systems (see, e.g., Luckow et al., Bio/Technology, 6, 47 (1988)), and established cell lines such 293, COS-7, C127, 3T3, CHO, HeLa, BHK, etc.

Adenoviral vectors are preferred because, unlike plasmids and other viral vectors (e.g., herpes simplex virus), adenoviral vectors achieve gene transfer in both dividing and nondividing cells, with high levels of protein expression in
20 cardiovascular relevant sites such as myocardium, vascular endothelium, and skeletal muscle. Furthermore, the gene transferred by an adenoviral vector functions in an epichromosomal position and thus carries little risk of inappropriately inserting the transferred gene into a critical site of the host genome. The adenoviral vector also desirably is deficient in at least one gene
25 function required for viral replication. Preferably, the adenoviral vector is deficient in at least one essential gene function of the E1, E2, and/or E4 regions of the adenoviral genome. More preferably, the vector additionally is deficient in at least part of the E3 region of the adenoviral genome (e.g., an XbaI deletion of the E3 region).

Recombinant adenovirus can be delivered to cultured cells by simply adding the virus to the culture media. Infection of host animals/humans can be achieved by directly injecting the viral particles into the bloodstream or into the desired tissue. The half-life of the virus in serum can be extended by complexing
5 the virus with liposomes (e.g. Lipofectin, Life Technologies) or polyethylene glycol. The adenovirus vector normally enters the cell through an interaction between the knob domain of the viral fiber protein and the coxsackievirus and adenovirus receptor, CAR. The viral vector can be directed to specific cells, or to cells which do not express the CAR, by genetically engineering the virus to
10 express a ligand specific to a certain cell receptor.

In an alternate embodiment, apoptosis can be initiated or enhanced by chemically upregulating the transcription of endogenous apoptosis-specific eIF-5A, or apoptosis-specific DHS, or both, with chemicals, or by chemically enhancing the activation of apoptosis-specific eIF-5A. In one such embodiment,
15 PGF-2 α is administered to the cancer cells or tumor of the animal/human to upregulate the transcription of DHS and eIF-5A.

Apoptosis-specific eIF-5A is a suitable target for regulation of apoptosis, including apoptosis underlying disease processes, since it likely acts in the post-transcriptional regulation of downstream effectors and transcription factors
20 involved in the apoptotic pathway. The present inventive methods of modulating apoptosis-specific eIF-5A and apoptosis-specific DHS, either alone or in combination, can be accomplished in cells of animals resulting in induction or enhancement of apoptosis and giving rise to novel methods and compositions for the treatment and prevention of diseases caused by, causing, or otherwise having
25 an etiology associated with an inability of cells to undergo apoptosis.

Many important human diseases are caused by abnormalities in the control of apoptosis. These abnormalities can result in either a pathological increase in cell number (e.g. cancer) or a damaging loss of cells (e.g. degenerative diseases). As non-limiting examples, the methods and compositions of the
30 present invention can be used to prevent or treat the following apoptosis-

associated diseases and disorders: neurological/ neurodegenerative disorders (e.g., Alzheimer's, Parkinson's, Huntington's, Amyotrophic Lateral Sclerosis (Lou Gehrig's Disease), autoimmune disorders (e.g., rheumatoid arthritis, systemic lupus erythematosus (SLE), multiple sclerosis), Duchenne Muscular Dystrophy
5 (DMD), motor neuron disorders, ischemia, chronic heart failure, stroke, infantile spinal muscular atrophy, cardiac arrest, renal failure, atopic dermatitis, sepsis and septic shock, AIDS, hepatitis, glaucoma, diabetes (type 1 and type 2), asthma, retinitis pigmentosa, osteoporosis, xenograft rejection, and burn injury.

The present inventive methods can be used for therapeutic treatments to
10 an animal having a cancerous cell or suffering from a tumor in an amount sufficient to kill the cancerous cell or inhibit the progression of the tumor, respectively. An amount adequate to accomplish this is defined as a therapeutically effective dose. Amounts effective for this use will depend upon the severity of the disease and the general state of the animal's own immune
15 system.

Inhibition of tumor growth means prevention or reduction of the progression of the tumor, e.g, the growth, invasiveness, metastases and/or recurrence of the tumor. The present inventive methods can be used to treat any suitable tumor, including, for example, tumors of the breast, heart, lung, small
20 intestine, colon, spleen, kidney, bladder, head and neck, ovary, prostate, brain, pancreas, skin, bone, bone marrow, blood, thymus, uterus, testicles, cervix or liver. Animals, preferably mammals, and more preferably humans can be treated using the compositions and methods of the present invention. The present inventive methods can thus be carried out *in vitro*, *ex vivo*, or *in vivo*.

25 Dosing schedules will also vary with the disease state and status of the animal, and will typically range from a single bolus dosage or continuous infusion to multiple administrations per day (e.g., every 4-6 hours), or as indicated by the treatment and the animals's condition. It should be noted, however, that the present invention is not limited to any particular dose.

In the present invention, any suitable method or route can be used for administration, for example, oral, intravenous, intraperitoneal, subcutaneous, or intramuscular administration. The dose of antagonist administered depends on numerous factors, including, for example, the type of molecule administered, the type and severity tumor being treated and the route of administration. It should be emphasized, however, that the present invention is not limited to any particular method or route of administration.

In one alternative embodiment, the present inventive methods can be used in combination with one or more traditional therapies. For example, a suitable antineoplastic agent can be used, such as a chemotherapeutic agent or radiation. In an additional alternative embodiment, the present inventive methods can be used in combination with one or more suitable adjuvants, such as, for example, cytokines (IL-10 and IL-13, for example) or other immune stimulators.

In another alternative embodiment, diagnosis of an apoptosis-related disorder can be made using apoptosis-specific eIF-5A and proliferating eIF-5A, which differs from the apoptosis-specific eIF-5A in that they are transcribed from different locations by different promoters; although the two are structurally homologous, with differences in the carboxy terminus. The method of diagnosis of the present invention involves comparing the amount of proliferating eIF-5A present in a given cell with the amount of apoptosis-specific eIF-5A present in the same cell. During normal functioning, a cell will have a greater amount of proliferating eIF-5A (also referred to herein as eIF-5A2) than apoptosis-specific eIF-5A (also referred to herein as eIF-5A1). However, in some cancer cells, the normal regulatory mechanisms go awry and the amount of apoptosis-specific eIF-5A relative to the amount of proliferating eIF-5A is altered. This potentially allows for diagnosis of a cell as cancerous prior to any phenotypic changes in the cell.

In yet another embodiment, the ratio of proliferating eIF-5A to apoptosis-specific eIF-5A can be used in drug screening. Such a method also involves comparing the amount of proliferating eIF-5A present in a given cell with the

amount of apoptosis-specific eIF-5A present in the same cell. The normal ratio of proliferating eIF-5A to apoptosis-specific eIF-5A would be compared to the ratio of proliferating eIF-5A to apoptosis-specific eIF-5A after contacting the cell with the drug candidate. Alterations in the ratio of proliferating eIF-5A to apoptosis-specific eIF-5A after contact allows for identification of those candidates that have apoptosis-modulating activity. Candidates having apoptosis-modulating activity can be useful in treating diseases associated with apoptosis, either through inhibition or induction of apoptosis. In addition, alterations in the ratio of proliferating eIF-5A to apoptosis-specific eIF-5A can be used to modulate apoptosis, which may also be useful to treat any of the conditions described herein as relating to abnormal apoptosis.

Using this method a large number of potential candidates, i.e., a library can be effectively screened to identify members of the library that modulate apoptosis. Any candidate or library of candidates can be screened using this method. For example, biological response modifiers that have shown promise as apoptosis modulators, including monoclonal antibodies that alter signal transduction pathways, cytokines such as TRAIL (Apo2 ligand), ligands for retinoid/steroid family nuclear receptors, and small-molecule compounds that bind and inhibit protein kinases, can be screened for definite apoptosis-modulating activity using the present methods.

One suitable candidate is a protein kinase C-alpha antisense oligonucleotide, ISIS 3521 (ISIS Pharmaceuticals, Inc., Carlsbad, CA), which has anti-tumor activity. Other specific candidates include caspases (Idun Pharmaceuticals, San Diego, CA), which are known to play a crucial role in the triggering and execution of apoptosis in a variety of cell types leading to cancer and neurodegenerative diseases; GENASENSE™ (Genta, Inc., Berkeley Heights, NJ), which is an antisense drug that blocks the production of Bcl-2; INGN 241 (Introgen Therapeutics, Inc., Houston, TX), which is a gene therapy targeting P53; rituximab (IDEC Corporation, Osaka, Japan), which is an anti-CD20

monoclonal antibody; and general apoptosis driven therapies for cardiovascular disease and cancer (Ægera Therapeutics Inc., Quebec, Canada).

It is understood that the nucleic acids and polypeptides of the present invention, where used in an animal for the purpose of prophylaxis or treatment, will be administered in the form of a composition additionally comprising a pharmaceutically acceptable carrier. Suitable pharmaceutically acceptable carriers include, for example, one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. Pharmaceutically acceptable carriers can further comprise minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the binding proteins. The compositions of the injection can, as is well known in the art, be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the mammal.

The compositions of this invention can be in a variety of forms. These include, for example, solid, semi-solid and liquid dosage forms, such as tablets, pills, powders, liquid solutions, dispersions or suspensions, liposomes, suppositories, injectable and infusible solutions. The preferred form depends on the intended mode of administration and therapeutic application.

Such compositions can be prepared in a manner well known in the pharmaceutical art. In making the composition the active ingredient will usually be mixed with a carrier, or diluted by a carrier, and/or enclosed within a carrier which can, for example, be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, excipient or medium for the active ingredient. Thus, the composition can be in the form of tablets, lozenges, sachets, cachets, elixirs, suspensions, aerosols (as a solid or in a liquid medium), ointments containing for example up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, injection solutions, suspensions, sterile packaged powders and as a topical patch.

Having now generally described the invention, the same will be more readily understood through reference to the following examples, which are provided by way of illustration. The Examples are set forth to aid in understanding the invention but are not intended to, and should not be construed to, limit its scope in any way. The examples do not include detailed descriptions of conventional methods. Such methods are well known to those of ordinary skill in the art and are described in numerous publications. Detailed descriptions of conventional methods, such as those employed in the construction of vectors and plasmids, the insertion of nucleic acids encoding polypeptides into such vectors and plasmids, the introduction of plasmids into host cells, and the expression and determination thereof of genes and gene products can be obtained from numerous publications, including Sambrook, J. et al., (1989) *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press. All references mentioned herein are incorporated in their entirety.

15

EXAMPLES

Example 1

The present example demonstrates isolation and characterization of a full-length cDNA encoding a rat eIF-5A nucleic acid exhibiting apoptosis-specific expression.

20

Superovulation and Induction of Apoptosis in Rat Corpus Luteum

Immature (21-30 day old) female rats were superovulated by subcutaneous injection with 50 IU of PMSG (Pregant Mare Serum Gonadotropin) and 60 to 65 hours later with 50 IU of HCG (Human Chorionic Gonadotropin). Seven days after the treatment with HCG, corpus luteum apoptosis was induced by subcutaneous injection with 500 mg of PGF-2 α . Rats were sacrificed at various times (e.g., 1, 8, and 24 hours) after PGF-2 α treatment, and the corpora lutea were

25

removed and placed in liquid nitrogen. Control corpus luteum tissue was obtained by sacrificing rats immediately before PGF-2 α treatment.

Dispersion of Rat Ovarian Corpus Luteum Cells

- 5 Six to nine days after superovulation, rats were treated by multisite subcutaneous injection with 500 mg PGF-2 α . Fifteen to thirty minutes later, the ovaries were removed from the superovulated rats, placed in EBSS (Gibco) on ice, blotted dry and weighed. Connective tissue was trimmed away, and the ovaries were minced finely with a razor blade and washed twice with EBSS 2X.
- 10 Collagenase solution was prepared by vortexing 6.5 mg of collagenase (Sigma, Catalogue # C 5138) in 5ml of EBSS. Minced tissue from 8 ovaries was added to 5 ml of collagenase in EBSS in a 50 ml Erlenmeyer flask and agitated gently by withdrawing several times into a Diamed pipette. The flask with minced tissue was then placed in a water bath at 37° C for 20 minutes with gentle shaking
- 15 (Position 45 on GFL incubator) under 95% air, 5% CO₂.

- Following this incubation, the flask was placed on ice, and the suspension of cells was transferred with a plastic transfer pipet onto a Nitex filter fitted with Swiss Nitex Nylon Monofilament (75 m). The filtrate was collected into a 15 ml Falcon test tube. A second aliquot (2.5 ml) of collagenase solution (6.5 mg
- 20 collagenase/5ml EBSS) was added to the minced tissue remaining in the 50 ml Erlenmeyer flask, agitated gently using a pipette, incubated for 10 minutes and filtered as above. The two filtrates were combined and centrifuged in a clinical centrifuge (~200g) for 5 minutes at room temperature. All but ~2ml of the supernatant were removed with a pipet and discarded, and the sedimented cells
- 25 were resuspended in the remaining 2 ml of supernatant.

 The cells were washed twice by adding 5 ml of MEM and centrifuging and resuspending as above. The washed cells were resuspended in 30 mls of MEM containing 10 mm glutamine in a 50 ml Erlenmeyer flask and incubated for 1 hour without shaking at 37° C under 95% air, 5% CO₂. The cells were then

sedimented by centrifugation as above and resuspended in MEM containing 10 mM glutamine.

The concentration of dispersed cells was determined using a hemocytometer, and viability was assessed using trypan blue dye. Aliquots of 2-5
5 $\times 10^5$ cells were placed in 12x75 mm test tubes and incubated without shaking for 2-5 hours at 37° C under 95 % air, 5 % CO₂. The progress of apoptosis during this period was monitored by assessing the degree of DNA laddering.

Visualization of Apoptosis in Rat Corpus Luteum by DNA Laddering

10 The degree of apoptosis was determined by DNA laddering. Genomic DNA was isolated from dispersed corpus luteal cells or from excised corpus luteum tissue using the QIAamp DNA Blood Kit (Qiagen) according to the manufacturer's instructions. Corpus luteum tissue was excised before the induction of apoptosis by treatment with PGF-2 α , 1 and 24 hours after induction
15 of apoptosis. The isolated DNA was end-labeled by incubating 500 ng of DNA with 0.2 μ Ci [α -³²P]dCTP, 1 mM Tris, 0.5 mM EDTA, 3 units of Klenow enzyme, and 0.2 pM each of dATP, dGTP, and dTTP at room temperature for 30 minutes. Unincorporated nucleotides were removed by passing the sample through a 1 ml Sepadex G-50 column according to Sambrook et al. The samples were then
20 resolved by Tris-acetate-EDTA (1.8 %) gel electrophoresis. The gel was dried for 30 minutes at room temperature under vacuum and exposed to x-ray film at - 80° C for 24 hours.

In one experiment, the degree of apoptosis in superovulated rat corpus lutea was examined either 0, 1, or 24 hours after injection with PGF-2 α . In the 0
25 hour control, the ovaries were removed without PGF-2 α injection. Laddering of low molecular weight DNA fragments reflecting nuclease activity associated with apoptosis is not evident in control corpus luteum tissue excised before treatment with PGF-2 α , but is discernible within 1 hour after induction of apoptosis and is pronounced by 24 hours after induction of apoptosis, which is shown in FIG. 16.
30 In this figure, the top panel is an autoradiograph of the Northern blot probed with

the ^{32}P -dCTP-labeled 3'-untranslated region of rat corpus luteum apoptosis-specific DHS cDNA. The lower panel is the ethidium bromide stained gel of total RNA. Each lane contains 10 μg RNA. The data indicate that there is down-regulation of eIF-5A transcript following serum withdrawal.

5 In another experiment, the corresponding control animals were treated with saline instead of PGF-2 α . Fifteen minutes after treatment with saline or PGF-2 α , corpora lutea were removed from the animals. Genomic DNA was isolated from the corpora lutea at 3 hours and 6 hours after removal of the tissue from the animals. DNA laddering and increased end labeling of genomic DNA
10 are evident 6 hours after removal of the tissue from the PGF-2 α -treated animals, but not at 3 hours after removal of the tissue. *See* FIG. 17. DNA laddering reflecting apoptosis is also evident when corpora lutea are excised 15 minutes after treatment with PGF-2 α and maintained for 6 hours under *in vitro* conditions in EBSS (Gibco). Nuclease activity associated with apoptosis is also evident
15 from more extensive end labeling of genomic DNA.

 In another experiment, superovulation was induced by subcutaneous injection with 500 μg of PGF-2 α . Control rats were treated with an equivalent volume of saline solution. Fifteen to thirty minutes later, the ovaries were removed and minced with collagenase. The dispersed cells from rats treated with
20 PGF-2 α were incubated in 10 mm glutamine + 10 mm spermidine for 1 hour and for a further 5 hours in 10 mm glutamine without spermidine (lane 2) or in 10 mm glutamine + 10 mm spermidine for 1 hour and for a further 5 hours in 10 mm glutamine + 1 mm spermidine (lane 3). Control cells from rats treated with saline were dispersed with collagenase and incubated for 1 hour and a further 5 hours in
25 glutamine only (lane 1). Five hundred nanograms of DNA from each sample was labeled with [α - ^{32}P]-dCTP using klenow enzyme, separated on a 1.8 % agarose gel, and exposed to film for 24 hours. Results are shown in FIG. 18.

 In yet another experiment, superovulated rats were injected subcutaneously with 1 mg/100 g body weight of spermidine, delivered in three
30 equal doses of 0.333 mg/100 g body weight, 24, 12, and 2 hours prior to a

subcutaneous injection with 500 μ g PGF-2 α . Control rats were divided into three sets: no injections, three injections of spermidine but no PGF-2 α ; and three injections with an equivalent volume of saline prior to PGF-2 α treatment. Ovaries were removed from the rats either 1 hour and 35 minutes or 3 hours and 45 minutes after prostaglandin treatment and used for the isolation of DNA. Five hundred nanograms of DNA from each sample was labeled with [α -³²P]-dCTP using Klenow enzyme, separated on a 1.8 % agarose gel, and exposed to film for 24 hours: lane 1, no injections (animals were sacrificed at the same time as for lanes 3-5); lane 2, three injections with spermidine (animals were sacrificed at the same time as for lanes 3-5); lane 3, three injections with saline followed by injection with PGF-2 α (animals were sacrificed 1 h and 35 min after treatment with PGF-2 α); lane 4, three injections with spermidine followed by injection with PGF-2 α (animals were sacrificed 1 h and 35 min after treatment with PGF-2 α); lane 5, three injections with spermidine followed by injection with PGF-2 α (animals were sacrificed 1 h and 35 min after treatment with PGF-2 α); lane 6, three injections with spermidine followed by injection with PGF-2 α (animals were sacrificed 3 h and 45 min after treatment with PGF-2 α); lane 7, three injections with spermidine followed by injection with PGF-2 α (animals were sacrificed 3 h and 45 min after treatment with PGF-2 α). Results are shown in FIG. 19.

RNA Isolation

Total RNA was isolated from corpus luteum tissue removed from rats at various times after PGF-2 α induction of apoptosis. Briefly, the tissue (5 g) was ground in liquid nitrogen. The ground powder was mixed with 30 ml guanidinium buffer (4 M guanidinium isothiocyanate, 2.5 mM NaOAc pH 8.5, 0.8% β -mercaptoethanol). The mixture was filtered through four layers of Miracloth and centrifuged at 10,000g at 4° C for 30 minutes. The supernatant was then subjected to cesium chloride density gradient centrifugation at 11,200g for 20 hours. The pelleted RNA was rinsed with 75% ethanol, resuspended in

600 ml DEPC-treated water and the RNA precipitated at -70° C with 1.5 ml 95% ethanol and 60 ml of 3M NaOAc.

Genomic DNA Isolation and Laddering

5 Genomic DNA was isolated from extracted corpus luteum tissue or dispersed corpus luteal cells using the QIAamp DNA Blood Kit (Qiagen) according to the manufacturer's instructions. The DNA was end-labeled by incubating 500 ng of DNA with 0.2 μ Ci [α -³²P]dCTP, 1 mM Tris, 0.5 mM EDTA, 3 units of Klenow enzyme, and 0.2 pM each of dATP, dGTP, and dTTP, 10 at room temperature for 30 minutes. Unincorporated nucleotides were removed by passing the sample through a 1-ml Sephadex G-50 column according to the method described by Maniatis et al. The samples were then resolved by Tris-acetate-EDTA (2 %) gel electrophoresis. The gel was dried for 30 minutes at room temperature under vacuum and exposed to x-ray film at - 80° C for 24 15 hours.

Plasmid DNA Isolation, DNA Sequencing

The alkaline lysis method described by Sambrook et al., *supra*, was used to isolate plasmid DNA. The full-length positive cDNA clone was sequenced 20 using the dideoxy sequencing method. Sanger et al., Proc. Natl. Acad. Sci. USA, 74:5463-5467. The open reading frame was compiled and analyzed using BLAST search (GenBank, Bethesda, MD) and sequence alignment was achieved using a BCM Search Launcher: Multiple Sequence Alignments Pattern-Induced Multiple Alignment Method (see F. Corpet, Nuc. Acids Res., 16:10881-10890, 25 (1987). Sequences and sequence alignments are shown in FIGS. 5-11.

Northern Blot Hybridization of Rat Corpus Luteum RNA

Twenty milligrams of total RNA isolated from rat corpus luteum at various stages of apoptosis were separated on 1% denatured formaldehyde 30 agarose gels and immobilized on nylon membranes. The full-length rat

apoptosis-specific eIF-5A cDNA (SEQ ID NO:1) labeled with ^{32}P -dCTP using a random primer kit (Boehringer) was used to probe the membranes 7×10^7 . Alternatively, full length rat apoptosis-specific DHS cDNA (SEQ ID NO:6) labeled with ^{32}P -dCTP using a random primer kit (Boehringer) was used to probe the membranes (7×10^7 cpm). The membranes were washed once with $1\times$ SSC, 0.1% SDS at room temperature and three times with $0.2\times$ SSC, 0.1% SDS at 65°C . The membranes were dried and exposed to X-ray film overnight at -70°C .

As can be seen, eIF-5A and DHS are both upregulated in apoptosing corpus luteum tissue. Expression of apoptosis-specific eIF-5A is significantly enhanced after induction of apoptosis by treatment with PGF- 2α – low at time zero, increased substantially within 1 hour of treatment, increased still more within 8 hours of treatment and increased slightly within 24 hours of treatment (FIG. 14). Expression of DHS was low at time zero, increased substantially within 1 hour of treatment, increased still more within 8 hours of treatment and increased again slightly within 24 hours of treatment (FIG. 15).

*Generation of an Apoptosing Rat Corpus Luteum
RT-PCR Product Using Primers Based on Yeast,
Fungal and Human eIF-5A Sequences*

A partial-length apoptosis-specific eIF-5A sequence (SEQ ID NO:11) corresponding to the 3' end of the gene was generated from apoptosing rat corpus luteum RNA template by RT-PCR using a pair of oligonucleotide primers designed from yeast, fungal and human eIF-5A sequences. The upstream primer used to isolate the 3'end of the rat eIF-5A gene is a 20 nucleotide degenerate primer: 5' TCSAARACHGGNAAGCAYGG 3' (SEQ ID NO:9), wherein S is selected from C and G; R is selected from A and G; H is selected from A, T, and C; Y is selected from C and T; and N is any nucleic acid. The downstream primer used to isolate the 3'end of the rat eIF-5A gene contains 42 nucleotides: 5' GCGAAGCTTCCATGG CTCGAGTTTTTTTTTTTTTTTTTTTTTTT 3' (SEQ ID NO:10). A reverse transcriptase polymerase chain reaction (RT-PCR) was

carried out. Briefly, using 5 mg of the downstream primer, a first strand of cDNA was synthesized. The first strand was then used as a template in a RT-PCR using both the upstream and downstream primers.

5 Separation of the RT-PCR products on an agarose gel revealed the presence a 900 bp fragment, which was subcloned into pBluescriptTM (Stratagene Cloning Systems, LaJolla, CA) using blunt end ligation and sequenced (SEQ ID NO:11). The cDNA sequence of the 3' end is SEQ ID NO:11 and the amino acid sequence of the 3' end is SEQ ID NO:12. *See* FIGS. 1-2.

A partial-length apoptosis-specific eIF-5A sequence (SEQ ID NO:15) corresponding to the 5' end of the gene and overlapping with the 3' end was generated from apoptosing rat corpus luteum RNA template by RT-PCR. The 5' primer is a 24-mer having the sequence, 5' CAGGTCTAGAGTTGGAATCGAAGC 3' (SEQ ID NO:13), that was designed from human eIF-5A sequences. The 3' primer is a 30-mer having the sequence, 5' ATATCTCGAGCCTT GATTGCAACAGCTGCC 3' (SEQ ID NO:14) that was designed according to the 3' end RT-PCR fragment. A reverse transcriptase-polymerase chain reaction (RT-PCR) was carried out. Briefly, using 5 mg of the downstream primer, a first strand of cDNA was synthesized. The first strand was then used as a template in a RT-PCR using both the upstream and downstream primers.

20 Separation of the RT-PCR products on an agarose gel revealed the presence a 500 bp fragment, which was subcloned into pBluescriptTM (Stratagene Cloning Systems, LaJolla, CA) using XbaI and XhoI cloning sites present in the upstream and downstream primers, respectively, and sequenced (SEQ ID NO:15). The cDNA sequence of the 5' end is SEQ ID NO:15, and the amino acid sequence of the 5' end is SEQ ID NO:16. *See* FIG. 2.

The sequences of the 3' and 5' ends of the rat apoptosis-specific eIF-5A (SEQ ID NO:11 and SEQ ID NO:15, respectively) overlapped and gave rise to the full-length cDNA sequence (SEQ ID NO:1). This full-length sequence was aligned and compared with sequences in the GeneBank data base. *See* FIGS. 1-2.

The cDNA clone encodes a 154 amino acid polypeptide (SEQ ID NO:2) having a calculated molecular mass of 16.8 KDa. The nucleotide sequence, SEQ ID NO:1, for the full length cDNA of the rat apoptosis-specific corpus luteum eIF-5A gene obtained by RT-PCR is depicted in FIG. 3 and the corresponding derived amino acid sequence is SEQ ID NO:9. The derived full-length amino acid sequence of eIF-5A was aligned with human and mouse eIF-5a sequences. See FIG. 7-9.

Generation of an Apoptosing Rat Corpus Luteum

RT-PCR Product Using Primers Based on a

Human DHS Sequence

A partial-length apoptosis-specific DHS sequence (SEQ ID NO:6) corresponding to the 3' end of the gene was generated from apoptosing rat corpus luteum RNA template by RT-PCR using a pair of oligonucleotide primers designed from a human DHS sequence. The 5' primer is a 20-mer having the sequence, 5' GTCTGTGTATTATTGGGCCC 3' (SEQ ID NO. 17).; the 3' primer is a 42-mer having the sequence, 5' GCGAAGCTTCCATGGC TCGAGTTTTTTTTTTTTTTTTTTTTTTT 3' (SEQ ID NO:18). A reverse transcriptase polymerase chain reaction (RT-PCR) was carried out. Briefly, using 5 mg of the downstream primer, a first strand of cDNA was synthesized. The first strand was then used as a template in a RT-PCR using both the upstream and downstream primers.

Separation of the RT-PCR products on an agarose gel revealed the presence a 606 bp fragment, which was subcloned into pBluescriptTM (Stratagene Cloning Systems, LaJolla, CA) using blunt end ligation and sequenced (SEQ ID NO:6). The nucleotide sequence (SEQ ID NO:6) for the partial length cDNA of the rat apoptosis-specific corpus luteum DHS gene obtained by RT-PCR is depicted in FIG. 4 and the corresponding derived amino acid sequence is SEQ ID NO.7.

Isolation of Genomic DNA and Southern Analysis

Genomic DNA for southern blotting was isolated from excised rat ovaries. Approximately 100 mg of ovary tissue was divided into small pieces and placed into a 15 ml tube. The tissue was washed twice with 1 ml of PBS by gently shaking the tissue suspension and then removing the PBS using a pipette. The tissue was resuspended in 2.06 ml of DNA-buffer (0.2 M Tris-HCl pH 8.0 and 0.1 mM EDTA) and 240 μ l of 10 % SDS and 100 μ l of proteinase K (Boehringer Mannheim; 10 mg/ml) was added. The tissue was placed in a shaking water bath at 45° C overnight. The following day another 100 μ l of proteinase K (10 mg/ml) was added and the tissue suspension was incubated in a waterbath at 45 °C for an additional 4 hours. After the incubation the tissue suspension was extracted once with an equal volume of phenol:chloroform:iso-amyl alcohol (25:24:1) and once with an equal volume of chloroform:iso-amyl alcohol (24:1). Following the extractions 1/10th volume of 3M sodium acetate (pH 5.2) and 2 volumes of ethanol were added. A glass pipette sealed and formed into a hook using a Bunsen burner was used to pull the DNA threads out of solution and to transfer the DNA into a clean microcentrifuge tube. The DNA was washed once in 70 % ethanol and air-dried for 10 minutes. The DNA pellet was dissolved in 500 μ l of 10 mM Tris-HCl (pH 8.0), 10 μ l of RNase A (10 mg/ml) was added, and the DNA was incubated for 1 hour at 37 °C. The DNA was extracted once with phenol:chloroform:iso-amyl alcohol (25:24:1) and the DNA was precipitated by adding 1/10th volume of 3 M sodium acetate (pH 5.2) and 2 volumes of ethanol. The DNA was pelleted by centrifugation for 10 minutes at 13,000 x g at 4° C. The DNA pellet was washed once in 70 % ethanol and dissolved in 200 μ l 10 mM Tris-HCl (pH 8.0) by rotating the DNA at 4 °C overnight.

For Southern blot analysis, genomic DNA isolated from rat ovaries was digested with various restriction enzymes that either do not cut in the endogenous gene or cut only once. To achieve this, 10 μ g genomic DNA, 20 μ l 10X reaction buffer and 100 U restriction enzyme were reacted for five to six hours in a total reaction volume of 200 μ l. Digested DNA was loaded onto a 0.7 % agarose gel

and subjected to electrophoresis for 6 hours at 40 volts or overnight at 15 volts. After electrophoresis, the gel was depurinated for 10 minutes in 0.2 N HCl followed by two 15-minute washes in denaturing solution (0.5 M NaOH, 1.5 M NaCl) and two 15 minute washes in neutralizing buffer (1.5 M NaCl, 0.5 M Tris-HCl pH 7.4). The DNA was transferred to a nylon membrane, and the membrane was prehybridized in hybridization solution (40 % formamide, 6 X SSC, 5 X Denhart's, solution (1 X Denhart's solution is 0.02 % Ficoll, 0.02 % PVP, and 0.02 % BSA), 0.5 % SDS, and 1.5 mg of denatured salmon sperm DNA). A 700 bp PCR fragment of the 3' UTR of rat eIF-5A cDNA (650 bp of 3' UTR and 50 bp of coding) was labeled with [α - ^{32} P]-dCTP by random priming and added to the membrane at 1 X 10⁶ cpm/ml.

Similarly, a 606 bp PCR fragment of the rat DHS cDNA (450 bp coding and 156 bp 3' UTR) was random prime labeled with [α - ^{32} P]-dCTP and added at 1 X 10⁶ cpm/ml to a second identical membrane. The blots were hybridized overnight at 42° C and then washed twice with 2 X SSC and 0.1 % SDS at 42° C and twice with 1 X SSC and 0.1 % SDS at 42° C. The blots were then exposed to film for 3-10 days.

Rat corpus genomic DNA was cut with restriction enzymes as indicated on FIG. 20 and probed with ^{32} P-dCTP-labeled full-length eIF-5A cDNA. Hybridization under high stringency conditions revealed hybridization of the full-length cDNA probe to several restriction fragments for each restriction enzyme digested DNA sample, indicating the presence of several isoforms of eIF-5A. Of particular note, when rat genomic DNA was digested with EcoRV, which has a restriction site within the open reading frame of apoptosis-specific eIF-5A, two restriction fragments of the apoptosis-specific isoform of eIF-5A were detectable in the Southern blot. The two fragments are indicated with double arrows in FIG. 20. The restriction fragment corresponding to the apoptosis-specific isoform of eIF-5A is indicated by a single arrow in the lanes labeled EcoR1 and BamH1, restriction enzymes for which there are no cut sites within the open reading frame. These results suggest that the apoptosis-specific eIF-5A is a single copy

gene in rat. As shown in FIGS. 5 through 13, the eIF-5A gene is highly conserved across species, and so it would be expected that there is a significant amount of conservation between isoforms within any species.

Figure 21 shows a Southern blot of rat genomic DNA probed with ³²P-dCTP-labeled partial-length rat corpus luteum apoptosis-specific DHS cDNA. The genomic DNA was cut with EcoRV, a restriction enzyme that does not cut the partial-length cDNA used as a probe. Two restriction fragments are evident indicating that there are two copies of the gene or that the gene contains an intron with an EcoRV site.

10

Example 2

The present example demonstrates modulation of apoptosis with apoptosis-specific eIF-5A and DHS.

15 *Culturing of COS-7 Cells and Isolation of RNA*

COS-7, an African green monkey kidney fibroblast-like cell line transformed with a mutant of SV40 that codes for wild-type T antigen, was used for all transfection-based experiments. COS-7 cells were cultured in Dulbecco's Modified Eagle's medium (DMEM) with 0.584 grams per liter of L-glutamine, 4.5 g of glucose per liter, and 0.37 % sodium bicarbonate. The culture media was supplemented with 10 % fetal bovine serum (FBS) and 100 units of penicillin/streptomycin. The cells were grown at 37° C in a humidified environment of 5 % CO₂ and 95 % air. The cells were subcultured every 3 to 4 days by detaching the adherent cells with a solution of 0.25 % trypsin and 1 mM EDTA. The detached cells were dispensed at a split ratio of 1:10 in a new culture dish with fresh media.

COS-7 cells to be used for isolation of RNA were grown in 150-mm tissue culture treated dishes (Corning). The cells were harvested by detaching them with a solution of trypsin-EDTA. The detached cells were collected in a centrifuge tube, and the cells were pelleted by centrifugation at 3000 rpm for 5

30

minutes. The supernatant was removed, and the cell pellet was flash-frozen in liquid nitrogen. RNA was isolated from the frozen cells using the GenElute Mammalian Total RNA Miniprep kit (Sigma) according to the manufacturer's instructions.

5

Construction of Recombinant Plasmids and Transfection of COS-7 Cells

Recombinant plasmids carrying the full-length coding sequence of rat apoptosis eIF-5A in the sense orientation and the 3' untranslated region (UTR) of rat apoptosis eIF-5A in the antisense orientation were constructed using the mammalian epitope tag expression vector, pHM6 (Roche Molecular Biochemicals), which is illustrated in FIG. 21. The vector contains the following: CMV promoter - human cytomegalovirus immediate-early promoter/enhancer; HA - nonapeptide epitope tag from influenza hemagglutinin; BGH pA - Bovine growth hormone polyadenylation signal; fl ori - fl origin; SV40 ori - SV40 early promoter and origin; Neomycin - Neomycin resistance (G418) gene; SV40 pA - SV40 polyadenylation signal; Col E1 - ColE1 origin; Ampicillin- Ampicillin resistance gene. The full-length coding sequence of rat apoptosis eIF-5A and the 3' UTR of rat apoptosis eIF-5A were amplified by PCR from the original rat eIF-5A RT-PCR fragment in pBluescript (SEQ ID NO:1). To amplify the full-length eIF-5A the primers used were as follows: Forward 5' GCCAAGCTTAATGGCAGATGATTT GG 3' (Hind3) and Reverse 5' CTGAATTCCAGT TATTTTGCCATGG 3' (EcoR1). To amplify the 3' UTR rat eIF-5A the primers used were as follows: forward 5' AATGAATTCCGCCATGACAGAGGAGGC 3' (EcoR1) and reverse 5' GCGAAGCTTCCATGGCTCGAGTTTTTTTTTTTTTTTTTTTTTTT 3' (Hind3).

The full-length rat eIF-5A PCR product isolated after agarose gel electrophoresis was 430 bp in length while the 3' UTR rat eIF-5A PCR product was 697 bp in length. Both PCR products were subcloned into the Hind 3 and EcoR1 sites of pHM6 to create pHM6-full-length eIF-5A and pHM6-antisense 3'UTReIF-5A. The full-length rat eIF-5A PCR product was subcloned in frame

with the nonapeptide epitope tag from influenza hemagglutinin (HA) present upstream of the multiple cloning site to allow for detection of the recombinant protein using an anti-[HA]-peroxidase antibody. Expression is driven by the human cytomegalovirus immediate-early promoter/enhancer to ensure high level
5 expression in mammalian cell lines. The plasmid also features a neomycin-resistance (G418) gene, which allows for selection of stable transfectants, and a SV40 early promoter and origin, which allows episomal replication in cells expressing SV40 large T antigen, such as COS-7.

COS-7 cells to be used in transfection experiments were cultured in either
10 24 well cell culture plates (Corning) for cells to be used for protein extraction, or 4 chamber culture slides (Falcon) for cells to be used for staining. The cells were grown in DMEM media supplemented with 10 % FBS, but lacking penicillin/streptomycin, to 50 to 70 % confluency. Transfection medium sufficient for one well of a 24-well plate or culture slide was prepared by diluting
15 0.32 µg of plasmid DNA in 42.5 µl of serum-free DMEM and incubating the mixture at room temperature for 15 minutes. 1.6 µl of the transfection reagent, LipofectAMINE (Gibco, BRL), was diluted in 42.5 µl of serum-free DMEM and incubated for 5 minutes at room temperature. After 5 minutes the LipofectAMINE mixture was added to the DNA mixture and incubated together
20 at room temperature for 30 to 60 minutes. The cells to be transfected were washed once with serum-free DMEM before overlaying the transfection medium and the cells were placed back in the growth chamber for 4 hours.

After the incubation, 0.17 ml of DMEM + 20 % FBS was added to the cells. The cells were the cultured for a further 40 hours before either being
25 induced to undergo apoptosis prior to staining or harvested for Western blot analysis. As a control, mock transfections were also performed in which the plasmid DNA was omitted from the transfection medium.

Protein Extraction and Western Blotting

Protein was isolated for Western blotting from transfected cells by washing the cells twice in PBS (8 g/L NaCl, 0.2 g/L KCl, 1.44 g/L Na₂HPO₄, and 0.24 g/L KH₂PO₄) and then adding 150 µl of hot SDS gel-loading buffer (50 mM
5 Tris-HCl pH 6.8, 100 mM dithiothreitol, 2 % SDS, 0.1 % bromophenol blue, and 10 % glycerol). The cell lysate was collected in a microcentrifuge tube, heated at 95° C for 10 minutes, and then centrifuged at 13,000 x g for 10 minutes. The supernatant was transferred to a fresh microcentrifuge tube and stored at -20° C until ready for use.

10 For Western blotting, 2.5 or 5 µg of total protein was separated on a 12 % SDS-polyacrylamide gel. The separated proteins were transferred to a polyvinylidene difluoride membrane. The membrane was then incubated for one hour in blocking solution (5 % skim milk powder, 0.02 % sodium azide in PBS) and washed three times for 15 minutes in PBS-T (PBS + 0.05 % Tween-20). The
15 membrane was stored overnight in PBS-T at 4 °C. After being warmed to room temperature the next day, the membrane was blocked for 30 seconds in 1 µg/ml polyvinyl alcohol. The membrane was rinsed 5 times in deionized water and then blocked for 30 minutes in a solution of 5 % milk in PBS. The primary antibody was preincubated for 30 minutes in a solution of 5 % milk in PBS prior to
20 incubation with the membrane.

Several primary antibodies were used. An anti-[HA]-peroxidase antibody (Roche Molecular Biochemicals) was used at a dilution of 1:5000 to detect expression of the recombinant proteins. Since this antibody is conjugated to peroxidase, no secondary antibody was necessary, and the blot was washed and
25 developed by chemiluminescence. The other primary antibodies that were used are monoclonal antibodies from Oncogene that recognize p53 (Ab-6), Bcl-2 (Ab-1), and c-Myc (Ab-2). The monoclonal antibody to p53 was used at a dilution of 0.1 µg/ml, and the monoclonal antibodies to Bcl-2 and c-Myc were both used at a dilution of 0.83 µg/ml. After incubation with primary antibody for 60 to 90
30 minutes, the membrane was washed 3 times for 15 minutes in PBS-T. Secondary

antibody was then diluted in 1 % milk in PBS and incubated with the membrane for 60 to 90 minutes. When p53 (Ab-6) was used as the primary antibody, the secondary antibody used was a goat anti-mouse IgG conjugated to alkaline phosphatase (Rockland) at a dilution of 1:1000. When Bcl-2 (Ab-1) and c-Myc (Ab-2) were used as the primary antibody, a rabbit anti-mouse IgG conjugated to peroxidase (Sigma) was used at a dilution of 1:5000. After incubation with the secondary antibody, the membrane was washed 3 times in PBS-T.

Two detection methods were used to develop the blots, a colorimetric method and a chemiluminescent method. The colorimetric method was used only when p53 (Ab-6) was used as the primary antibody in conjunction with the alkaline phosphatase-conjugated secondary antibody. Bound antibody was visualized by incubating the blot in the dark in a solution of 0.33 mg/mL nitro blue tetrazolium, 0.165 mg/mL 5-bromo-4-chloro-3-indolyl phosphate, 100 mM NaCl, 5 mM MgCl₂, and 100 mM Tris-HCl (pH 9.5). The color reaction was stopped by incubating the blot in 2 mM EDTA in PBS. A chemiluminescent detection method was used for all other primary antibodies, including anti-[HA]-peroxidase, Bcl-2 (Ab-1), and c-Myc (Ab-2). The ECL Plus Western blotting detection kit (Amersham Pharmacia Biotech) was used to detect peroxidase-conjugated bound antibodies. In brief, the membrane was lightly blotted dry and then incubated in the dark with a 40:1 mix of reagent A and reagent B for 5 minutes. The membrane was blotted dry, placed between sheets of acetate, and exposed to X-ray film for time periods varying from 10 seconds to 10 minutes.

Induction of Apoptosis in COS 7 Cells

Two methods were used to induce apoptosis in transfected COS-7 cells, serum deprivation and treatment with Actinomycin D, *streptomyces* sp (Calbiochem). For both treatments, the medium was removed 40 hours post-transfection. For serum starvation experiments, the media was replaced with serum- and antibiotic-free DMEM. Cells grown in antibiotic-free DMEM supplemented with 10 % FBS were used as a control. For Actinomycin D

induction of apoptosis, the media was replaced with antibiotic-free DMEM supplemented with 10 % FBS and 1 µg/ml Actinomycin D dissolved in methanol. Control cells were grown in antibiotic-free DMEM supplemented with 10 % FBS and an equivalent volume of methanol. For both methods, the percentage of apoptotic cells was determined 48 hours later by staining with either Hoescht or Annexin V-Cy3. Induction of apoptosis was also confirmed by Northern blot analyses, as shown in FIG. 22.

Hoescht Staining

10 The nuclear stain, Hoescht, was used to label the nuclei of transfected COS-7 cells in order to identify apoptotic cells based on morphological features such as nuclear fragmentation and condensation. A fixative, consisting of a 3:1 mixture of absolute methanol and glacial acetic acid, was prepared immediately before use. An equal volume of fixative was added to the media of COS-7 cells growing on a culture slide and incubated for 2 minutes. The media/fixative mixture was removed from the cells and discarded, and 1 ml of fixative was added to the cells. After 5 minutes the fixative was discarded, and 1 ml of fresh fixative was added to the cells and incubated for 5 minutes. The fixative was discarded, and the cells were air-dried for 4 minutes before adding 1 ml of Hoescht stain (0.5 µg/ml Hoescht 33258 in PBS). After a 10-minute incubation in the dark, the staining solution was discarded and the slide was washed 3 times for 1 minute with deionized water. After washing, 1 ml of McIlvaine's buffer (0.021 M citric acid, 0.058 M Na₂HPO₄·7H₂O; pH 5.6) was added to the cells, and they were incubated in the dark for 20 minutes. The buffer was discarded, the cells were air-dried for 5 minutes in the dark and the chambers separating the wells of the culture slide were removed. A few drops of Vectashield mounting media for fluorescence (Vector Laboratories) was added to the slide and overlaid with a coverslip. The stained cells were viewed under a fluorescence microscope using a UV filter. Cells with brightly stained or fragmented nuclei were scored as apoptotic.

Annexin V-Cy3 Staining

An Annexin V-Cy3 apoptosis detection kit (Sigma) was used to fluorescently label externalized phosphatidylserine on apoptotic cells. The kit
5 was used according to the manufacturer's protocol with the following modifications. In brief, transfected COS-7 cells growing on four chamber culture slides were washed twice with PBS and three times with 1 X Binding Buffer. 150 μ l of staining solution (1 μ g/ml AnnCy3 in 1 X Binding Buffer) was added, and the cells were incubated in the dark for 10 minutes. The staining solution was
10 then removed, and the cells were washed 5 times with 1 X Binding Buffer. The chamber walls were removed from the culture slide, and several drops of 1 X Binding Buffer were placed on the cells and overlaid with a coverslip. The stained cells were analyzed by fluorescence microscopy using a green filter to visualize the red fluorescence of positively stained (apoptotic) cells. The total
15 cell population was determined by counting the cell number under visible light.

Example 3

The present example demonstrates modulation of apoptosis with apoptosis-specific eIF-5A and DHS.

20 Using the general procedures and methods described in the previous examples, FIG. 23 is a flow chart illustrating the procedure for transient transfection of COS-7 cells, in which cells in serum-free medium were incubated in plasmid DNA in lipofectAMINE for 4 hours, serum was added, and the cells were incubated for a further 40 hours. The cells were then either incubated in
25 regular medium containing serum for a further 48 hours before analysis (i.e. no further treatment), deprived of serum for 48 hours to induce apoptosis before analysis, or treated with actinomycin D for 48 hours to induce apoptosis before analysis.

FIG. 22 is a Western blot illustrating transient expression of foreign
30 proteins in COS-7 cells following transfection with pHM6. Protein was isolated

from COS-7 cells 48 hours after either mock transfection, or transfection with pHM6-LacZ, pHM6-Antisense 3' rF5A (pHM6-Antisense 3' UTR rat apoptosis eIF-5A), or pHM6-Sense rF5A (pHM6-Full length rat apoptosis eIF-5A). Five μ g of protein from each sample was fractionated by SDS-PAGE, transferred to a
5 PVDF membrane, and Western blotted with anti-[HA]-peroxidase. The bound antibody was detected by chemiluminescence and exposed to x-ray film for 30 seconds. Expression of LacZ (lane 2) and of sense rat apoptosis eIF-5A (lane 4) is clearly visible.

As described above, COS-7 cells were either mock transfected or
10 transfected with pHM6-Sense rF5A (pHM6-Full length rat eIF-5A). Forty hours after transfection, the cells were induced to undergo apoptosis by withdrawal of serum for 48 hours. The caspase proteolytic activity in the transfected cell extract was measured using a fluorometric homogenous caspase assay kit (Roche Diagnostics). DNA fragmentation was also measured using the FragEL DNA
15 Fragmentation Apoptosis Detection kit (Oncogene) which labels the exposed 3'-OH ends of DNA fragments with fluorescein-labeled deoxynucleotides.

Additional COS-7 cells were either mock transfected or transfected with pHM6-Sense rF5A (pHM6-Full length rat eIF-5A). Forty hours after transfection, the cells were either grown for an additional 48 hours in regular
20 medium containing serum (no further treatment), induced to undergo apoptosis by withdrawal of serum for 48 hours or induced to undergo apoptosis by treatment with 0.5 μ g/ml of Actinomycin D for 48 hours. The cells were either stained with Hoescht 33258, which depicts nuclear fragmentation accompanying apoptosis, or stained with Annexin V-Cy3, which depicts phosphatidylserine exposure
25 accompanying apoptosis. Stained cells were also viewed by fluorescence microscopy using a green filter and counted to determine the percentage of cells undergoing apoptosis. The total cell population was counted under visible light.

FIG. 25 illustrates enhanced apoptosis as reflected by increased caspase activity when COS-7 cells were transiently transfected with pHM6 containing

full-length rat apoptosis-induced eIF-5A in the sense orientation. Expression of rat apoptosis-induced eIF-5A resulted in a 60% increase in caspase activity.

FIG. 26 illustrates enhanced apoptosis as reflected by increased DNA fragmentation when COS-7 cells were transiently transfected with pHM6

5 containing full-length rat apoptosis-induced eIF-5A in the sense orientation.

Expression of rat apoptosis-induced eIF-5A resulted in a 273% increase in DNA fragmentation. FIG. 27 illustrates detection of apoptosis as reflected by increased nuclear fragmentation when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense orientation.

10 There is a greater incidence of fragmented nuclei in cells expressing rat apoptosis-induced eIF-5A. FIG. 28 illustrates enhanced apoptosis as reflected by increased nuclear fragmentation when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense orientation. Expression of rat apoptosis-induced eIF-5A resulted in a 27 % and
15 63 % increase in nuclear fragmentation over control in non-serum starved and serum starved samples, respectively.

FIG. 29 illustrates detection of apoptosis as reflected by phosphatidylserine exposure when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense

20 orientation. FIG. 30 illustrates enhanced apoptosis as reflected by increased phosphatidylserine exposure when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense orientation. Expression of rat apoptosis-induced eIF-5A resulted in a 140 % and 198 % increase in phosphatidylserine exposure over control, in non-serum starved
25 and serum starved samples, respectively.

FIG. 31 illustrates enhanced apoptosis as reflected by increased nuclear fragmentation when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense orientation.

Expression of rat apoptosis-induced eIF-5A resulted in a 115 % and 62 %
30 increase in nuclear fragmentation over control in untreated and treated samples,

respectively. FIG. 32 illustrates a comparison of enhanced apoptosis under conditions in which COS-7 cells transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense orientation were either given no further treatment or treatment to induce apoptosis.

5

Example 4

The present example demonstrates modulation of apoptotic activity following administration of apoptosis-specific eIF-5A and DHS.

Moreover, COS-7 cells were either mock transfected, transfected with
10 pHM6-LacZ or transfected with pHM6-Sense rF5A (pHM6-Full length rat eIF-5A) and incubated for 40 hours. Five µg samples of protein extract from each sample were fractionated by SDS-PAGE, transferred to a PVDF membrane, and Western blotted with a monoclonal antibody that recognizes Bcl-2. Rabbit anti-mouse IgG conjugated to peroxidase was used as a secondary antibody, and
15 bound antibody was detected by chemiluminescence and exposure to x-ray film. Results are shown in FIG. 32. Less Bcl-2 is detectable in cells transfected with pHM6-Sense rF5A than in those transfected with pHM6-LacZ; therefore, Bcl-2 is down-regulated.

Additional COS-7 cells were either mock transfected, transfected with
20 pHM6-antisense 3' rF5A (pHM6-antisense 3' UTR of rat apoptosis-specific eIF-5A) or transfected with pHM6-Sense rF5A (pHM6-Full length rat apoptosis-specific eIF-5A). Forty hours after transfection, the cells were induced to undergo apoptosis by withdrawal of serum for 48 hours. Five µg samples of protein extract from each sample were fractionated by SDS-PAGE, transferred to
25 a PVDF membrane, and Western blotted with a monoclonal antibody that recognizes Bcl-2. Rabbit anti-mouse IgG conjugated to peroxidase was used as a secondary antibody, and bound antibody was detected by chemiluminescence and exposure to x-ray film.

Also additionally, COS-7 cells were either mock transfected, transfected
30 with pHM6-LacZ or transfected with pHM6-Sense rF5A (pHM6-Full length rat

eIF-5A) and incubated for 40 hours. Five µg samples of protein extract from each sample were fractionated by SDS-PAGE, transferred to a PVDF membrane, and Western blotted with a monoclonal antibody that recognizes p53. Goat anti-mouse IgG conjugated to alkaline phosphatase was used as a secondary antibody,
5 and bound antibody was detected a colorimetrically.

Finally, COS-7 cells were either mock transfected, transfected with pHM6-LacZ or transfected with pHM6-Sense rF5A (pHM6-Full length rat eIF-5A) and incubated for 40 hours. Five µg samples of protein extract from each sample were fractionated by SDS-PAGE, transferred to a PVDF membrane, and
10 probed with a monoclonal antibody that recognizes p53. Corresponding protein blots were probed with with anti-[HA]-peroxidase to determine the level of rat apoptosis-specific eIF-5A expression. Goat anti-mouse IgG conjugated to alkaline phosphatase was used as a secondary antibody, and bound antibody was detected by chemiluminescence.

15 FIG. 33 illustrates downregulation of Bcl-2 when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense orientation. The upper panel illustrates the Coomassie-blue-stained protein blot; the lower panel illustrates the corresponding Western blot. Less Bcl-2 is detectable in cells transfected with pHM6-Sense rF5A than in those
20 transfected with pHM6-LacZ.

FIG. 34 illustrates upregulation of Bcl-2 when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the antisense orientation. The upper panel illustrates the Coomassie-blue-stained protein blot; the lower panel illustrates the corresponding Western
25 blot. More Bcl-2 is detectable in cells transfected with pHM6-antisense 3' rF5A than in those mock transfected or transfected with pHM6-Sense rF5A.

FIG. 35 illustrates upregulation of c-Myc when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense orientation. The upper panel illustrates the Coomassie-blue-stained protein blot; the lower panel illustrates the corresponding Western blot.
30

More c-Myc is detectable in cells transfected with pHM6-Sense rF5A than in those transfected with pHM6-LacZ or the mock control.

FIG. 36 illustrates upregulation of p53 when COS-7 cells were transiently transfected with pHM6 containing full-length rat apoptosis-induced eIF-5A in the sense orientation. The upper panel illustrates the Coomassie-blue-stained protein blot; the lower panel illustrates the corresponding Western blot. More p53 is detectable in cells transfected with pHM6-Sense rF5A than in those transfected with pHM6-LacZ or the mock control.

FIG. 37 illustrates the dependence of p53 upregulation upon the expression of pHM6-full length rat apoptosis-induced eIF-5A in COS-7 cells. In the Western blot probed with anti-[HA]-peroxidase, the upper panel illustrates the Coomassie-blue-stained protein blot and the lower panel illustrates the corresponding Western blot. More rat apoptosis-induced eIF-5A is detectable in the first transfection than in the second transfection. In the Western blot probed with anti-p53, the upper panel in A illustrates a corresponding Coomassie-blue-stained protein blot and the lower panel illustrates the Western blot with p53. For the first transfection, more p53 is detectable in cells transfected with pHM6-Sense rF5A than in those transfected with pHM6-LacZ or the mock control. For the second transfection in which there was less expression of rat apoptosis-induced eIF-5A, there was no detectable difference in levels of p53 between cells transfected with pHM6-Sense rF5A, pHM6-LacZ or the mock control.

Example 5

The present example demonstrates that apoptosis-specific eIF-5A can induce apoptosis in cells with active p53 (RKO cells) and in cells without active p53 (RKO-E6 cells), indicating that apoptosis-specific eIF-5A can initiate apoptosis through pathway(s) other than the p53 pathway. This also supports our contention that it is acting upstream and likely able to kill a wide range of different types of cancers.

The present example also indicates that the active site of eIF-5A1 is the

carboxy terminus of the protein (i.e. see experiments with truncated eIF-5A1), which most likely contains the RNA binding domain.

Further, the present example also demonstrates that human eIF-5A2 is most likely a proliferating eIF-5A as it is unable to induce apoptosis. Thus, it is
5 believed that of the two eIF-5A genes in the human data bank, apoptosis-specific eIF-5A1 is the apoptosis gene, and eIF-5A2 is the proliferation gene.

Culture of RKO and RKO-E6 Cells

RKO (American Type Culture Collection CRL-2577), a human colon
10 carcinoma cell line expressing wild-type p53, and RKO-E6 (American Type Culture Collection CRL-2578), a cell line derived from RKO that contains a stably integrated human papilloma virus E6 oncogene that causes a decrease in normal p53 level and function, were used for transfection-based experiments. RKO and RKO-E6 cells were cultured in Minimum Essential Medium Eagle
15 (MEM) with non-essential amino acids, Earle's salts, and L-glutamine. The culture media was supplemented with 10 % fetal bovine serum (FBS) and 100 units of penicillin/streptomycin. The cells were grown at 37°C in a humidified environment of 5 % CO₂ and 95 % air. The cells were subcultured every 3 to 4 days by detaching the adherent cells with a solution of 0.25 % trypsin and 1 mM
20 EDTA. The detached cells were dispensed at a split ratio of 1:10 to 1:12 into a new culture dish with fresh media.

Cloning of human eIF5A2

Human eIF5A2 was isolated by RT-PCR from RNA isolated from RKO
25 cells using primers designed against the sequence of human eIF5A2 available from GenBank (ACCESSION XM_113401). Figure 38 provides an alignment of human eIF-5A isolated from RKO cells with the sequence of human eIF-5A2. RNA was isolated from RKO cells using the GenElute Mammalian Total RNA Miniprep Kit (Sigma). The forward primer used to amplify eIF5A2 had the
30 sequence 5' AAACCTACCATCTCCCCTGCC 3' and the reverse primer had the

sequence 5' TGCCCTACACAGGCTGAAAG 3'. The resulting 936 bp PCR product was subcloned into the pGEM-T Easy Vector (Promega) and sequenced.

The pGEM-T Easy-eIF5A2 construct was then used as template to generate a eIF5A2 PCR fragment to be subcloned in frame into the mammalian expression vector pHM6 (Roche). The forward primer used to amplify human eIF5A2 was 5' ATCAAGCTTGCCACCATGGCAGACG 3' and the reverse primer was 5' AACGAATTCCATGCCTGATGTTTCCG 3'. The resulting 505 bp PCR product was digested with Hind 3 and EcoR 1 and subcloned into the Hind 3 and EcoR1 sites of pHM6.

10

Construction of pHM6-truncated eIF5A1

In order to determine if the carboxy-terminal region of eIF5A1 is important for its apoptosis-inducing activity, a carboxy-terminal deleted eIF5A1 was constructed. The truncated eIF5A1, coding for amino acids 1 to 127, was generated by PCR using pBS-rat eIF5A1 as template. The forward PCR primer was 5' GCCAAGCTTAATGGCAGATGATTTGG 3' and the reverse primer was 5' TCCGAATTCGTACTTCTGCTCAATC 3'. The resulting 390 bp PCR product was digested with EcoR 1 and Hind 3 and subcloned into the EcoR 1 and Hind 3 sites of pHM6.

20

Transfection

RKO or RKO-E6 cells to be used in transfection experiments were cultured in 8 well chamber culture slides (Falcon) for cells to be used for Hoescht staining or 6 well plates for cells to be analyzed by flow cytometry. The cells were grown in MEM media supplemented with 10 % FBS but lacking penicillin/streptomycin to 70 to 80 % confluency. Transfection medium sufficient for one well of an 8 well culture slide was prepared by diluting 0.425 µg of plasmid DNA in 22 µl of serum-free MEM and incubating the mixture at room temperature for 15 minutes. 0.85 µl of the transfection reagent, LipofectAMINE (Gibco, BRL), was diluted in 22 µl of serum-free MEM and

30

incubated for 5 minutes at room temperature. After 5 minutes the LipofectAMINE mixture was added to the DNA mixture and incubated at room temperature for 30 to 60 minutes. The cells to be transfected were washed once with serum-free MEM before adding 44 μ l of MEM to the transfection medium and overlaying it over the cells. The cells were placed back in the growth chamber for 4 hours. After the incubation, 88 μ l of MEM + 20 % FBS was added to the cells. The cells were then cultured for a further 44 hours and then stained with Hoescht 33258 as previously described. In another set of experiments, RKO or RKO-E6 cells in 8-well culture slides were treated with 0.25 μ g/ml Actinomycin D 24 hours after transfection and stained with Hoescht 20 hours later. Transfections carried out in 6-well plates were performed in the same manner except that all the reagents were increased by 4.81 times. RKO cells transfected in 6 well plates were harvested 48 hours after transfection and fixed for analysis by flow cytometry as described below.

15

Determination of Transfection Efficiency

The efficiency of transfection was determined by staining pHM6-LacZ-transfected cells with 5-bromo-4-chloro-3-indolyl-beta-D-galactopyranoside (X-GAL). Blue-stained cells are LacZ-expressing transfected cells, and transfection efficiency was calculated as the number of blue stained cells divided by the total number of cells. Transfected cells were stained 48 hours following transfection. The cells were washed twice with PBS and then fixed for 10 minutes at room temperature in 0.5 % glutaraldehyde/PBS. The cells were washed three times with 1 mM MgCl_2 /PBS and then incubated with staining solution [5 mM $\text{K}_4\text{Fe}(\text{CN})_6 \cdot 3\text{H}_2\text{O}$, 5 mM $\text{K}_3\text{Fe}(\text{CN})_6$, 1 mM MgCl_2 , 0.1 % X-GAL in PBS] until blue-stained cells appeared.

25

Hoescht Staining

The nuclear stain, Hoescht, was used to label the nuclei of transfected RKO and RKO-E6 cells in order to identify apoptotic cells based on nuclear

30

fragmentation and condensation. A fixative, consisting of a 3:1 mixture of absolute methanol and glacial acetic acid, was prepared immediately before use. An equal volume of fixative was added to the media of cells growing on a culture slide and incubated for 2 minutes. The media/fixative mixture was removed from the cells and discarded, and 1 ml of fixative was added to the cells. After 5 minutes the fixative was discarded and 1 ml of fresh fixative was added to the cells and incubated for 5 minutes. The fixative was discarded and the cells were air-dried for 4 minutes before adding 1 ml of Hoescht stain (0.5 µg/ml Hoescht 33258 in PBS). After a 10 minute incubation in the dark, the staining solution was discarded and the slide was washed 3 times for 1 minute with deionized water. After washing, 1 ml of McIlvaine's buffer (0.021 M citric acid, 0.058 M Na₂HPO₄·7H₂O; pH 5.6) was added to the cells and incubated in the dark for 20 minutes. The buffer was discarded and the cells were air-dried for 5 minutes in the dark and the chambers separating the wells of the culture slide were removed. A few drops of Vectashield mounting media for fluorescence (Vector Laboratories) was added to the slide and overlaid with a coverslip. The stained cells were viewed under a fluorescent microscope using a UV filter. Cells with brightly stained or fragmented nuclei were scored as apoptotic.

DNA Fragmentation Detection by Flow Cytometry

DNA fragments generated during apoptosis were labeled with fluorescein-labeled deoxynucleotides using the Fluorescein-FragEL™ DNA Fragmentation Detection Kit (Oncogene Research Products). Cells transfected with various constructs in 6 well culture plates were harvested by trypsinization 48 hours after transfection and fixed and labeled according to the manufacturer's instructions. Briefly, the cells were pelleted at 1000 x g for 5 minutes at 4°C and washed once in PBS (8 g/L NaCl, 0.2 g/L KCl, 1.44 g/L Na₂HPO₄, and 0.24 g/L KH₂PO₄). The cells were resuspended in 4% formaldehyde/PBS and incubated at room temperature for 10 minutes. The cells were pelleted again, resuspended in 1 ml of 80 % ethanol, and stored at 4°C. On the day of analysis, 1 ml of fixed cells (at 1

X 10⁶ cells/ml) was transferred to a microfuge tube and the cells pelleted by centrifugation at 1000 x g for 5 minutes. The pelleted cells were resuspended in 200 µl of 1 X TBS (20 mM Tris pH 7.6, 140 mM NaCl) and incubated 10 to 15 minutes at room temperature. The cells were then pelleted again and resuspended
5 in 100 µl of 20 µg/ml proteinase K and incubated for 5 minutes at room temperature. The cells were pelleted and resuspended in 100 µl of 1 X TdT Equilibration buffer and incubated at room temperature for 10 to 30 minutes. The cells were then pelleted by centrifugation and resuspended in 60 µl of TdT Labeling Reaction Mixture and incubated for 1 to 1.5 hours in the dark. After the
10 incubation, the cells were pelleted by centrifugation and washed twice in 200 µl of 1 X TBS. The cells were resuspended in a final volume of 0.5 ml 1 X TBS and analyzed on a flow cytometer equipped with a 488 nm argon ion laser source.

Protein Extraction and Western Blotting

15 Protein was isolated for Western blotting from transfected cells by washing the cells twice in PBS and then adding 150 µl of hot SDS gel-loading buffer (50 mM Tris-HCl pH 6.8, 100 mM dithiothreitol, 2 % SDS, 0.1 % bromophenol blue, and 10 % glycerol). The cell lysate was collected in a microcentrifuge tube, heated at 95 °C for 10 minutes, and then centrifuged at
20 13,000 X g for 10 minutes. The supernatant was transferred to a fresh microcentrifuge tube and stored at – 20 °C until ready for use.

For Western blotting, 5 µg or 10 µg of total protein was separated on a 12 % SDS-polyacrylamide gel. The separated proteins were transferred to a polyvinylidene difluoride membrane. The membrane was then incubated for one
25 hour in blocking solution (5 % skim milk powder, 0.02 % sodium azide in PBS) and washed three times for 15 minutes in PBS-T (PBS + 0.05 % Tween-20). The membrane was stored overnight in PBS-T at 4 °C. After being warmed to room temperature the next day, the membrane was blocked for 30 seconds in 1 µg/ml polyvinyl alcohol. The membrane was rinsed 5 times in deionized water and then
30 blocked for 30 minutes in a solution of 5 % milk in PBS. The primary antibody

was preincubated for 30 minutes in a solution of 5 % milk in PBS/0.025% Tween-20 prior to incubation with the membrane.

The membranes were blotted with either a monoclonal antibody from Oncogene which recognizes p53 (Ab-6), or a polyclonal antibody directed against a synthetic peptide (amino-CRLPEGDLGKEIEQKYD-carboxy) homologous to the c-terminal end of human eIF5A1 that was raised in chickens. The monoclonal antibody to p53 was used at a dilution of 0.1 µg/ml and the antibody against eIF5A1 was used at a dilution of 1:1000. After incubation with primary antibody for 60 to 90 minutes, the membrane was washed 3 times for 15 minutes in PBS-T. Secondary antibody was then diluted in 1 % milk in PBS/0.025 % Tween-20 and incubated with the membrane for 60 to 90 minutes. When p53 (Ab-6) was used as the primary antibody, the secondary antibody used was a goat anti-mouse IgG conjugated to alkaline phosphatase (Rockland) at a dilution of 1:1000. When anti-eIF5A1 was used as the primary antibody, a rabbit anti-chicken IgY conjugated to peroxidase (Gallus Immunotech) was used at a dilution of 1:10000. After incubation with the secondary antibody, the membrane was washed 3 times in PBS-T.

Two detection methods were used to develop the blots, a colourimetric method and a chemiluminescent method. The colourimetric method was used only when p53 (Ab-6) was used as the primary antibody in conjunction with the alkaline phosphatase-conjugated secondary antibody. Bound antibody was visualized by incubating the blot in the dark in a solution of 0.33 mg/mL nitro blue tetrazolium, 0.165 mg/mL 5-bromo-4-chloro-3-indolyl phosphate, 100 mM NaCl, 5 mM MgCl₂, and 100 mM Tris-HCl (pH 9.5). The color reaction was stopped by incubating the blot in 2 mM EDTA in PBS. A chemiluminescent detection method was used for all other primary antibodies, including anti-[HA]-peroxidase and anti-eIF5A1. The ECL Plus Western blotting detection kit (Amersham Pharmacia Biotech) was used to detect peroxidase-conjugated bound antibodies. In brief, the membrane was lightly blotted dry and then incubated in the dark with a 40:1 mix of reagent A and reagent B for 5 minutes. The

membrane was blotted dry, placed between sheets of acetate, and exposed to X-ray film for time periods varying from 10 seconds to 30 minutes.

Figure 39 shows a graph depicting the percentage of apoptosis occurring in RKO and RKO-E6 cells following transient transfection. RKO and RKO-E6 cells were transiently transfected with pHM6-LacZ or pHM6-eIF5A1. 24 hours later, the cells were treated either with 0.25 μ g/ml Actinomycin D or an equivalent volume of methanol (control). The cells were stained with Hoescht 20 hours later and were viewed under a fluorescent microscope using a UV filter. Cells that stained brightly due to condensed chromatin were scored as apoptotic. The experiments above reveal that RKO cells treated with Actinomycin D and transfected with pHM6-eIF5A1 showed a 240% increase in apoptosis relative to cells transfected with pHM6-LacZ that were not treated with Actinomycin D. RKO-E6 cells treated with Actinomycin D and transfected with pHM6-eIF5A1 showed a 105% increase in apoptosis relative to cells transfected with pHM6-LacZ that were not treated with Actinomycin D.

Figure 40 provides a graph depicting the percentage of apoptosis occurring in RKO cells following transient transfection. RKO cells were transiently transfected with pHM6-LacZ, pHM6-eIF5A1, pHM6-eIF5A2, or pHM6-truncated eIF5A1. The cells were stained with Hoescht 44 hours later and were viewed under a fluorescent microscope using a UV filter. Cells that stained brightly due to condensed chromatin were scored as apoptotic. Cells transfected with pHM6-eIF5A1 showed a 25% increase in apoptosis relative to control cells transfected with pHM6-LacZ. This increase was not apparent for cells transfected with pHM6-eIF5A2 or pHM6-truncated eIF5A1.

Figure 41 provides a graph depicting the percentage of apoptosis occurring in RKO cells following transient transfection. RKO cells were either left untransfected or were transiently transfected with pHM6-LacZ or pHM6-eIF5A1. The cells were stained with Hoescht 44 hours later and were viewed under a fluorescent microscope using a UV filter. Cells that stained brightly due to condensed chromatin were scored as apoptotic. After correction for transfection

efficiency, 60 % of the cells transfected with pHM6-eIF5A1 were apoptotic.

Figure 42 provides flow cytometry analysis of RKO cell apoptosis following transient transfection. RKO cells were either left untransfected or were transiently transfected with pHM6-LacZ, pHM6-eIF5A1, pHM6-eIF5A2, or
5 pHM6-truncated eIF5A1. 48 hours later the cells were harvested and fixed. Fragmented DNA reflecting apoptosis was labeled with fluorescein-conjugated deoxynucleotides and analyzed on a flow cytometer equipped with a 488 nm argon ion laser source. Fluorescence occurring under gate E is from non-apoptotic cells, and fluorescence occurring under gate F is from cells undergoing
10 apoptosis. The table depicts the percentage of cells undergoing apoptosis calculated based on the area under the peak of each gate. After correction for background apoptosis in untransfected cells and for transfection efficiency, 80% of cells transfected with pHM6-eIF5A1 exhibited apoptosis. Cells transfected with pHM6-LacZ, pHM6-eIF5A2 or pHM6-truncated eIF5A1 exhibited only
15 background levels of apoptosis.

Figure 43 provides Western blots of protein extracted from RKO cells treated with 0.25 µg/ml Actinomycin D for 0, 3, 7, 24, and 48 hours. 5 µg (for anti-eIF5A1) or 10 µg (for anti-p53) of total protein was separated on a 12 % SDS-polyacrylamide gel and transferred to a polyvinylidene difluoride
20 membrane. The top panel depicts a Western blot using anti-p53 as the primary antibody. The middle panel depicts a Western blot using anti-eIF5A1 as the primary antibody. The bottom panel depicts the membrane used for the anti-eIF5A1 blot stained with Coomassie blue following chemiluminescent detection to demonstrate equal loading. p53 and eIF5A1 are both upregulated by treatment
25 with Actinomycin D.

We Claim:

1. An isolated nucleic acid comprising a nucleotide sequence encoding a rat apoptosis-specific eIF-5A polypeptide.
2. The isolated nucleic acid of claim 1, wherein the nucleotide sequence comprises SEQ ID NO:1.
3. The isolated nucleic acid of claim 1, wherein the nucleotide sequence hybridizes under highly stringent conditions with the complement of SEQ ID NO:1.
4. The isolated nucleic acid of claim 1, wherein the nucleotide sequence encodes SEQ ID NO:2.
5. A purified polypeptide comprising an amino acid sequence of a rat apoptosis-specific eIF-5A polypeptide.
6. The purified polypeptide of claim 5, wherein the amino acid sequence comprises SEQ ID NO:2.
7. An isolated nucleic acid comprising a nucleotide sequence encoding a rat apoptosis-specific DHS polypeptide.
8. The isolated nucleic acid of claim 7, wherein the nucleotide sequence comprises SEQ ID NO:22.
9. The isolated nucleic acid of claim 7, wherein the nucleotide sequence hybridizes under highly stringent conditions with the complement of SEQ ID NO:22.
10. The isolated nucleic acid of claim 7, wherein the nucleotide sequence encodes SEQ ID NO:23.
11. A purified polypeptide comprising an amino acid sequence of a rat apoptosis-specific DHS polypeptide.
12. The purified polypeptide of claim 11, wherein the amino acid sequence comprises SEQ ID NO:23.
13. A method of modulating apoptosis in a cell comprising administering to the cell an agent that modulates apoptosis-specific eIF-5A function.

14. The method of claim 13, wherein the agent inhibits apoptosis-specific eIF-5A function, thereby inhibiting apoptosis.
15. The method of claim 14, wherein the agent is an antisense oligonucleotide.
16. The method of claim 15, wherein the antisense oligonucleotide comprises a nucleotide sequence encoding a portion of an apoptosis-specific eIF-5A polypeptide.
17. The method of claim 16, wherein the antisense oligonucleotide comprises SEQ ID NO:19, 20, or 21.
18. The method of claim 15, wherein the antisense oligonucleotide comprises a nucleotide sequence encoding a portion of an apoptosis-specific DHS polypeptide.
19. The method of claim 15, wherein the agent is a chemical or drug that inhibits apoptosis-specific eIF-5A function.
20. The method of claim 13, wherein the agent inhibits apoptosis-specific DHS function, thereby inhibiting apoptosis.
21. The method of claim 20, wherein the agent is an antisense oligonucleotide.
22. The method of claim 21, wherein the antisense oligonucleotide comprises a nucleotide sequence encoding a portion of an apoptosis-specific eIF-5A polypeptide.
23. The method of claim 22, wherein the nucleotide sequence comprising SEQ ID NO:19, 20, or 21.
24. The method of claim 21, wherein the antisense oligonucleotide comprises a nucleotide sequence encoding a portion of an apoptosis-specific DHS polypeptide.
25. The method of claim 20, wherein the agent is a chemical or drug that inhibits apoptosis-specific DHS function.
26. The method of claim 13, wherein the agent increases apoptosis-specific eIF-5A function, thereby inducing apoptosis in the cell.

27. The method of claim 26, wherein the agent comprises an expression vector.

28. The method of claim 27, wherein the expression vector comprises a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific eIF-5A polypeptide.

29. The method of claim 27, wherein the expression vector comprises a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific DHS polypeptide.

30. The method of claim 13, wherein the agent increases apoptosis-specific DHS function, thereby inducing apoptosis in the cell.

31. The method of claim 30, wherein the agent comprises an expression vector.

32. The method of claim 31, wherein the expression vector comprises a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific eIF-5A polypeptide.

33. The method of claim 31, wherein the expression vector comprises a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific DHS polypeptide.

34. The method of claim 13, wherein the cell is comprised within an animal.

35. The method of claim 34, wherein the agent is administered to the animal.

36. A method of modulating apoptosis in a cell comprising administering to the cell an agent that modulates apoptosis-specific DHS function.

37. The method of claim 36, wherein the agent inhibits apoptosis-specific DHS function, thereby inhibiting apoptosis.

38. The method of claim 37, wherein the agent is an antisense oligonucleotide.

39. The method of claim 38, wherein the antisense oligonucleotide comprises a nucleotide sequence encoding a portion of an apoptosis-specific eIF-5A polypeptide.

40. The method of claim 39, wherein the antisense oligonucleotide comprises SEQ ID NO:19, 20, or 21.

41. The method of claim 38, wherein the antisense oligonucleotide comprises a nucleotide sequence encoding a portion of an apoptosis-specific DHS polypeptide.

42. The method of claim 37, wherein the agent is a chemical or drug that inhibits apoptosis-specific eIF-5A function.

43. The method of claim 36, wherein the agent inhibits apoptosis-specific DHS function, thereby inhibiting apoptosis.

44. The method of claim 44, wherein the agent is an antisense oligonucleotide.

45. The method of claim 45, wherein the antisense oligonucleotide comprises a nucleotide sequence encoding a portion of an apoptosis-specific eIF-5A polypeptide.

46. The method of claim 45, wherein the antisense oligonucleotide comprises a nucleotide sequence encoding a portion of an apoptosis-specific DHS polypeptide.

47. The method of claim 44, wherein the agent is a chemical or drug that inhibits apoptosis-specific DHS function.

48. The method of claim 36, wherein the agent increases apoptosis-specific eIF-5A function, thereby inducing apoptosis in the cell.

49. The method of claim 48, wherein the agent comprises an expression vector.

50. The method of claim 49, wherein the expression vector comprises a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific eIF-5A polypeptide.

51. The method of claim 49, wherein the expression vector comprises a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific DHS polypeptide.

52. The method of claim 36, wherein the agent increases apoptosis-specific DHS function, thereby inducing apoptosis in the cell.

53. The method of claim 52, wherein the agent comprises an expression vector.

54. The method of claim 53, wherein the expression vector comprises a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific eIF-5A polypeptide.

55. The method of claim 53, wherein the expression vector comprises a promoter sequence operably linked to a nucleotide sequence encoding an apoptosis-specific DHS polypeptide.

56. The method of claim 36, wherein the cell is comprised within an animal.

57. The method of claim 56, wherein the agent is administered to the animal.

58. An antisense oligonucleotide comprising a nucleotide sequence encoding a portion of an apoptosis-specific eIF-5A polypeptide.

59. The antisense oligonucleotide of claim 58, wherein the nucleotide sequence comprises SEQ ID NO:19, 20, or 21.

60. An antisense oligonucleotide comprising a nucleotide sequence encoding a portion of an apoptosis-specific DHS polypeptide.

61. An expression vector comprising a promoter operably linked to a nucleotide sequence encoding an apoptosis-specific eIF-5A polypeptide.

62. An expression vector comprising a promoter operably linked to a nucleotide sequence encoding an apoptosis-specific DHS polypeptide.

63. A method of identifying modulation of apoptosis comprising:

- (i) providing a test cell;
- (ii) contacting the test cell with a drug candidate; and

(iii) determining whether a ratio of proliferating eIF-5A to apoptosis-specific eIF-5A in the test cell in the presence of the drug candidate has been altered,

wherein alterations in the ratio of proliferating eIF-5A to apoptosis-specific eIF-5A is indicative of modulation of apoptosis.

T CGAAGACCGGTAAGCACGGCCATGCCAAGGTCCATCTGGTGGTATTGATATTTTACTGGGAAGAAATAT
S K T G K H G H A K V H L V G I D I F T G K K Y
GAAGATATCTGCCCCGTCGACTCATAACATGGATGTCCCCAACATCAAAAGGAATGATTTCCAGCTGATTGGC
E D I C P S T H N M D V P N I K R N D F Q L I G
ATCCAGGATGGGTACCTATCCCTGCTCCAGGACAGTGGGGAGGTACGAGAGGACCTTCGTCTGCCTGAGGGA
I Q D G Y L S L L Q D S G E V R E D L R L P E G
GACCTTGGCAAGGAGATTGAGCAGAAGTATGACTGTGGAGAAGAGATCCTGATCACAGTGTGTCCGCCATG
D L G K E I E Q K Y D C G E E I L I T V L S A M
ACAGAGGAGGCAGCTGTTGCAATCAAGGCCATGGCAAAATAACTGGCTTCCAGGGTGGCGGTGGTGGCAGCA
T E E A A V A I K A M A K
GTGATCCATGAGCCTACAGAGGCCCCCTCCCCAGCTCTGGCTGGGCCCCITGGCTGGACTCCTATCCAATTTA
TTTGACGTTTTATTTTGGTTTTCTCACCCCTTCAAACCTGTCCGGGAGACCCTGCCCTTCACCTAGCTCCCT
TGGCCAGGCATGAGGGAGCCATGGCCTTGGTGAAGCTACCTGCCCTCTTCTCTCGCAGCCCTGATGGGGGAAA
GGGAGTGGGTACTGCCCTGTGGTTTAGGTTCCCTCTCCCTTTTCTTTTAAATTCATTTGGAATCAGAAAG
CTGTGGATTCTGGCAAATGGTCTTGTGTCTTTTATCCCACTCAAACCCATCTGGTCCCCTGTTCTCCATAGT
CCTTCACCCCCAAGCACCCTGACAGACTGGGGACCAGCCCCCTTCCCTGCCCTGTGTCTTTCCTCCAAACCCC
TCTATAGGGGTGACAAGAAGAGGAGGGGGGGAGGGGACACGATCCCTCCTCAGGCATCTGGGAAGGCCTTGC
CCCCATGGGCTTTACCTTTCTCTGTGGGCTTTCTCCCTGACACATTTGTTAAAAATCAAACCTGAATAAAAC
TACAAGTTTAAATATGAAAAAAAAAAAAAAAAAAAAA
(972 NT, 109 aa)

Figure 1

CAGGTCTAGAGTTGGAATCGAAGCCTCTTAAAAATGGCAGATGATTGGACTTCGAGACAGGAGATGCAGGGG
M A D D L D F E T G D A G
CCTCAGCCACCTTCCCAATGCAGTGCTCAGCATTACGTAAGAATGGTTTTGTGGTGCTCAAGGGCCGGCCAT
A S A T F P M Q C S A L R K N G P V V L K G R P
GTAAGATCGTCGAGATGTCTACTTCGAAGACTGGCAAGCATGGCCATGCCAAGGTCCATCTGGTTGGTATTG
C K I V E M S T S K T G K H G H A K V H L V G I
ATATTTTTACTGGGAAGAAATATGAAGATATCTGCCCGTCGACTCATAACATGGATGTCCCCAACATCAAAA
D I F T G K K Y E D I C P S T H N M D V P N I K
GGAATGATTTCCAGCTGATTGGCATCCAGGATGGGTACCTATCCCTGCTCCAGGACAGTGGGGAGGTACGAG
R N D F Q L I G I Q D G Y L S L L Q D S G E V R
AGGACCTTCGTCTGCCTGAGGGAGACCTTGSCAAGGAGATTGAGCAGAAGTATGACTGTGGAGAAGAGATCC
E D L R L P E G D L G K E I E Q K Y D C G E E I
TGATCACAGTGCTGTCCGCCATGACAGAGGAGGCAGCTGTTGCAATCAAGGCTCGAG
L I T V L S A M T E E A A V A I K A

(488 NT, 151 aa)

Figure 2

```

CAGGTCTAGAGTTTGGGAATCGAAGCCTCTTAAAATGGCAGATGATTGGAAGCTTCGAGACAGGAGATGCAGGGG
      M A D D L D F E T G D A G      13
CCTCAGCCACCTTCCCAATGCAGTGCTCAGCATTACGTAAGAATGGTTTTGTGGTGCTCAAGGGCCGGCCAT      144
A S A T F P M Q C S A L R K N G F V V L K G R P
GTAAGATCGTTCGAGATGCTCTACTTCGAAGACTGGCAAGCATGGCCATGCCAAGGTCCATCTGGTTGGTATTG
C K I V E M S T S K T G K H G H A K V H L V G I      61
ATATTTTTTACTGGGAAGAAATATGAAGATATCTGCCCGTCGACTCATAACATGGATGTCCCCAACATCAGAAA      288
D I F T G K K Y E D I C P S T H N M D V P N I K
GGAATGATTTCAGCTGATTGGCATCCAGGATGGGTACCTATCCCTGCTCCAGGACAGTGGGGAGGTACGAG
R N D F Q L I G I Q D G Y L S L L Q D S G E V R      109
AGGACCTTCGTCTGCCTGAGGGAGACCTTGGCAAGGAGATTGAGCAGAAGTATGACTGTGGAGAAGAGATCC      432
E D L R L P E G D L G K E I E Q K Y D C G E E I
TGATCACAGTGCTGTCCGCCATGACAGAGGAGGCAGCTGTTGCAATCAAGGCCATGGCAAAATAAAGTGGCTT
L I T V L S A M T E E A A V A I K A M A K *      154
CCAGGGTGGCGGTGGTGGCAGCAGTGATCCATGAGCCTACAGAGGCCCTCCCCAGCTCTGGCTGGGCCCTT      576
TGGCTGGACTCCTATCCAATTTATTTGACGTTTTATTTTGGTTTTCTCACCCTTCAAAGTGTGGGGGAGA
CCCTGCCCCCTTACCTAGCTCCCTTGGCCAGGCATGAGGGAGCCATGGCCTTGGTGAAGCTACCTGCCTCTTC      720
TCTCGCAGCCCTGATGGGGGAAAGGGAGTGGGTACTGCCCTGTGGTTTAGGTTCCCTCTCCCTTTTTCTTTT
TAATTCAATTTGGAATCAGAAAGCTGTGGATTCTGGCAATGGTCTTGTGTCTTTATCCCACTCAAACCCA      864
TCTGGTCCCCTGTTCTCCATAGTCCTTCACCCCAAGCACCCTGACAGACTGGGGACCAGCCCCCTTCCCT
GCCTGTGTCTCTTCCCAAACCCCTCTATAGGGGTGACAAGAAGAGGAGGGGGGAGGGGACACGATCCCTCC      1008
TCAGGCATCTGGGAAGGCCTTGCCCCCATGGGCTTTACCCTTTCTGTGGGCTTTCTCCCTGACACATTGT
TAAAAATCAAACCTGAATAAAACTACAAGTTTAATATGAAAAAAAAAAAAAAAAAAAAA      1139

```

... (1139 NT, 154 aa)

Figure 3

GCTGTGTATTATTGGGCCCATAAGAACCACATACCTGTGCTGAGTCCTGCACTCACAGACGGCTCACTGGGT
A V Y Y W A H K N H I P V L S P A L T D G S L G
GACATGATCTTTTTCCATTCCCTATAAAAACCCAGGCTTGGTCCTGGACATCGTTGAAGACCTGCGGCTCATC
D M I F F H S Y K N P G L V L D I V E D L R L I
AACATGCAGGCCATTTTCGCCAAGCGCACTGGGATGATCATCCTGGGTGGAGGCGTGGTCAAGCACCCACATC
N M Q A I F A K R T G M I I L G G G V V K H H I
GCCAATGCTAACCTCATGCGGAATGGAGCTGACTACGCTGTTTATATCAACACAGCCCAGGAGTTTGATGGC
A N A N L M R N G A D Y A V Y I N T A Q E F D G
TCAGACTCAGGAGCCCGGCCAGATGAGGCTGTCTCCTGGGGCAAGATCCGGATGGATGCACAGCCAGTAAAG
S D S G A R P D E A V S W G K I R M D A Q P V K
GTCTATGCTGATGCATCTCTGGTTTTCCCTTGCTGGTGGCTGAGACATTGCCCCAAAAGGCAGATGCCTTC
V Y A D A S L V F P L L V A E T F A Q K A D A F
AGAGCTGAGAAGAATGAGGACTGAGCAGATGGGTAAAGACGGAGGCTTCTGCCACACCTTTATTTATTATT
R A E K N E D
GCATACCAACCCCTCCTGGGCCCTCTCCTTGGTCAGCAGCATCTTGAGAATAAATGGCCTTTTTGTTGGTTT
CTGTAAAAAAAGGACTTTAAAAA

(606 NT, 151 aa)

Figure 4

Figure 5

Figure 5

rat vs. human(NM_020390) 72.5% identity (coding)

```

      10      20      30      40      50      60
rat   ATGGCAGATGATTTCGACTTCGAGACAGGAGATGCAGGGGCCTCAGCCACCTTCCCAATG
      :::::  ::  :  ::  ::  ::  :::::  :::::  ::  :::  :  :::  :::
human ATGGCAGACGAAATTGATTTCACACTACTGGAGATGCCGGGGCTTCCAGCACTTACCCCTATG
      10      20      30      40      50      60

      70      80      90     100     110     120
rat   CAGTGCTCAGCATTACGTAAGAATGGTTTTGTGGTGCTCAAGGGCCGGCCATGTAAGATC
      :::::  ::  ::  ::  ::  ::  ::  :::::  ::  ::  :::  ::  ::  ::
human CAGTGCTCGGCCTTGCGCAAAACGGCTTCGTGGTGCTCAAAGGACGACCATGCNAAATA
      70      80      90     100     110     120

      130     140     150     160     170     180
rat   GTCGAGATGTCTACTTCGAAGACTGGCAAGCATGGCCATGCCAAGGTCCATCTGGTTGGT
      ::  :::::  ::  ::  ::  ::  :::::  :::::  ::  ::  :::  ::  :::
human GTGGAGATGTCAACTTCCAAACCTGGAAAGCATGGTCATGCCAAGGTTACCTTGTGGA
      130     140     150     160     170     180

      190     200     210     220     230     240
rat   ATTGATATTTTACTGGGAAGAAATATGAAGATATCTGCCCGTCGACTCATAACATGGAT
      :::::  ::  ::  ::  :::::  :::::  ::  ::  ::  ::  ::  :::::
human ATTGATATTTTCACGGGCNAAAAATATGAAGATATTTGTCCTTCTACTCACAACATGGAT
      190     200     210     220     230     240

      250     260     270     280     290     300
rat   GTCCCCAACATCAAAGGAATGATTTCCAGCTGATTGGCATCCAGGATGGGTACCTATCC
      ::  ::  ::  ::  ::  :::::  ::  :::::  ::  ::  ::  ::  :::::
human GTTCCAAATATTAAGAGAAATGATTATCAACTGATATGCATTCAAGATGGTTACCTTTCC
      250     260     270     280     290     300

      310     320     330     340     350     360
rat   CTGCTCCAGGACAGTGGGGAGGTACGAGAGGACCTTCGTCTGCCTGAGGGAGACCTTGGC
      :::::  ::  :  ::  ::  ::  :::::  ::  :::::  ::  ::  ::  ::  :::
human CTGCTGACAGAACTGGTGAAATTCTGTGAGGATCTTAAACTGCCAGAAGGTGAAGTAGGC
      310     320     330     340     350     360

      370     380     390     400     410     420
rat   AAGGAGATTGAGCAGAAGTATGACTGTGGAGAAGAGATCCTGATCACAGTGCTGTCCGCC
      ::  ::  ::  ::  ::  ::  ::  ::  ::  ::  ::  ::  ::  ::  ::  ::
human AAAGAAATAGAGGGAAAATACAATGCAGGTGAAGATGTACAGGTGTCTGTCTATGTGTGCA
      370     380     390     400     410     420

      430     440     450     460
rat   ATGACAGAGGAGGAGCTGTTGCAATCAAGGCCATGGCAAAA
      :::  ::  ::  :::::  ::  ::  ::  ::  ::  :::
human ATGAGTGAAGAATATGCTGTAGCCATAAAACCCT--GCAAT
      430     440     450     460

```

Figure 6

rat vs. mouse (BC003889) 98.3% identity (coding)

```

      10      20      30      40      50      60
rat   ATGGCAGATGATTTGGACTTCGAGACAGGAGATGCAGGGGCCTCAGCCACCTTCCCAATG
      .....
mouse ATGGCAGATGATTTGGACTTCGAGACAGGAGATGCAGGGGCCTCAGCCACCTTCCCAATG
      10      20      30      40      50      60

      70      80      90     100     110     120
rat   CAGTGCTCAGCATTACGTAAGAATGGTTTTGTGGTGCTCAAGGGCCGGCCATGTAAGATC
      .....
mouse CAGTGCTCAGCATTACGTAAGAATGGTTTTGTGGTGCTCAAGGGCCGGCCATGTAAGATC
      70      80      90     100     110     120

      130     140     150     160     170     180
rat   GTCGAGATGTCTACTTCGAAGACTGGCAAGCATGGCCATGCCAAGGTCCATCTGGTTGGT
      .....
mouse GTCGAGATGTCTACTTCGAAGACTGGCAAGCATGGCCATGCCAAGGTCCATCTGGTTGGC
      130     140     150     160     170     180

      190     200     210     220     230     240
rat   ATTGATATTTTTACTGGGAAGAAATATGAAGATATCTGCCCGTCGACTCATAACATGGAT
      .....
mouse ATTGACATTTTTACTGGGAAGAAATATGAAGATATCTGCCCGTCGACTCATAATATGGAT
      190     200     210     220     230     240

      250     260     270     280     290     300
rat   GTCCCCAACATCAAAAGGAATGATTTCCAGCTGATTGGCATCCAGGATGGGTACCTATCC
      .....
mouse GTCCCCAACATCAAAAGGAATGATTTCCAGCTGATTGGCATCCAGGATGGGTACCTATCC
      250     260     270     280     290     300

      310     320     330     340     350     360
rat   CTGCTCCAGGACAGTGGGGAGGTACGAGAGGACCTTCGTCTGCCTGAGGGAGACCTTGGC
      .....
mouse CTGCTCCAGGACAGTGGGGAGGTACGAGAGGACCTTCGTCTGCCTGAAGGAGACCTTGGC
      310     320     330     340     350     360

      370     380     390     400     410     420
rat   AAGGAGATTGAGCAGAAGTATGACTGTGGAGAAGAGATCCTGATCACAGTGCTGTCCGCC
      .....
mouse AAGGAGATTGAGCAGAAGTATGACTGTGGAGAAGAGATCCTGATCACAGTGCTGTCTGCC
      370     380     390     400     410     420

      430     440     450     460
rat   ATGACAGAGGAGGCAGCTGTTGCAATCAAGGCCATGGCAAAA
      .....
mouse ATGACAGAGGAGGCAGCTGTTGCAATCAAGGCCATGGCAAAA
      430     440     450     460

```

Figure 7

```
rat vs. human(BC000751 or NM_001970) 100.0% identity

      10      20      30      40      50      60
rat   MADDLDFETGDAGASATFPMQCSALRKNGFVVLKGRPCKIVEMSTSKTGKHGHAKVHLVG
      : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human MADDLDFETGDAGASATFPMQCSALRKNGFVVLKGRPCKIVEMSTSKTGKHGHAKVHLVG
      10      20      30      40      50      60

      70      80      90     100     110     120
rat   IDIFTGKKYEDICPSTHNMDVPNIKRNDFQLIGIQDGYLSLLQDSGEVREDLRLPEGDLG
      : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human IDIFTGKKYEDICPSTHNMDVPNIKRNDFQLIGIQDGYLSLLQDSGEVREDLRLPEGDLG
      70      80      90     100     110     120

      130     140     150
rat   KEIEQKYDCGEEILITVLSAMTEEA A VAIKAMAK
      : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human KEIEQKYDCGEEILITVLSAMTEEA A VAIKAMAK
      130     140     150
```

Figure 8

rat vs. human(NM_020390) 82.5% identity

	10	20	30	40	50	60
rat	MADDLDFETGDAGASATFFPMQCSALRKNGFVVLKGRPCKIVEMSTSKTGKHGHAKVHLVG					
						
human	MADEIDFTTGDAGASSTYPMQCSALRKNGFVVLKGRPCKIVEMSTSKTGKHGHAKVHLVG					
	10	20	30	40	50	60
	70	80	90	100	110	120
rat	IDIFTGKKYEDICPSTHNMDVPNIKRNDYQLICIQDGYLSLLTETGEVREDLRLPEGDLG					
						
human	IDIFTGKKYEDICPSTHNMDVPNIKRNDYQLICIQDGYLSLLTETGEVREDLKLPEGLG					
	70	80	90	100	110	120
	130	140	150			
rat	KEIEQKYDCGEEILITVLSAMTEEA AVAIKAMAK					
						
human	KEIEGKYNAGEDVQVSVMCAMSE EYAVAIKP-CK					
	130	140	150			

Figure 9

rat vs. mouse (BC003889) 100.0% identity

	10	20	30	40	50	60
rat	MADDLDFETGDAGASATFPMQCSALRKNGFVVLKGRPCKIVEMSTSKTGKEGHAKVHLVG					
	::					
mouse	MADDLDFETGDAGASATFPMQCSALRKNGFVVLKGRPCKIVEMSTSKTGKEGHAKVHLVG					
	10	20	30	40	50	60
	70	80	90	100	110	120
rat	IDIFTGKKYEDICPSTHNMDVPNIKRNDFQLIGIQDGYLSLLQDSGEVREDLRLPEGDLG					
	::					
mouse	IDIFTGKKYEDICPSTHNMDVPNIKRNDFQLIGIQDGYLSLLQDSGEVREDLRLPEGDLG					
	70	80	90	100	110	120
	130	140	150			
rat	KEIEQKYDCGEEILITVLSAMTEEA A VAIKAMAK					
	::					
mouse	KEIEQKYDCGEEILITVLSAMTEEA A VAIKAMAK					
	130	140	150			

Figure 10

rat vs. human (BC000333) 87.4% identity (coding)

```

      10      20      30      40      50      60
rat   GCTGTGTATTATTGGGCCCCATAAGAACCACATACCTGTGCTGAGTCCTGCACTCACAGAC
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human TCCGTGTATTACTGGGCCCCAGAAGAACCACATCCCTGTGTTAGTCCCGCACTTACAGAC
      10      20      30      40      50      60

      70      80      90     100     110     120
rat   GGCTCACTGGGTGACATGATCTTTTCCATTCCCTATAAAAACCCAGGCTTGGTCCTGGAC
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human GGCTCGCTGGGCGACATGATCTTCTTCCATTCCCTACAAGAACCAGGCTTGGTCCTGGAC
      70      80      90     100     110     120

      130     140     150     160     170     180
rat   ATCGTTGAAGACCTGCGGCTCATCAACATGCAGGCCATTTTCGCCAAGCGCACTGGGATG
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human ATCGTTGAGGACCTGAGGCTCATCAACACACAGGCCATCTTTGCCAAGTGCCTGGGATG
      130     140     150     160     170     180

      190     200     210     220     230     240
rat   ATCATCCTGGGTGGAGGCGTGGTCAAGCACCACATCGCCAATGCTAACCTCATGCGGAAT
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human ATCAITCTGGGCGGGGCGTGGTCAAGCACCACATTGCCAATGCCAACCTCATGCGGAAC
      190     200     210     220     230     240

      250     260     270     280     290     300
rat   GGAGCTGACTACGCTGTTTATATCAACACAGCCCAGGAGTTTGATGGCTCAGACTCAGGA
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human GGGGCGGACTACGCTGTTTACATCAACACAGCCCAGGAGTTTGATGGCTCTGACTCAGGT
      250     260     270     280     290     300


      310     320     330     340     350     360
rat   GCCCGGCCAGATGAGGCTGTCTCCTGGGGCAAGATCCGGATGGATGCACAGCCAGTAAAG
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human GCCCGACCAGACGAGGCTGTCTCCTGGGGCAAGATCCGGGTGGATGCACAGCCCGTCAAG
      310     320     330     340     350     360


      370     380     390     400     410     420
rat   GTCTATGCTGATGCATCTCTGGTTTTCCCTTGCTGGTGGCTGAGACATTGCCCCAAAAG
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human GTCTATGCTGACGCTCCCTGGTCTTCCCCCTGCTTGTGGCTGAAACCTTTGCCCAGAAG
      370     380     390     400     410     420

      430     440     450
rat   GCAGATGCCCTTCAGAGCTGAGAAGAATGAGGAC
      : : : : : : : : : : : : : : : : : : : : : : : : : : : :
human ATGGATGCCCTTCATGCATGAGAAGAACGAGGAC
      430     440     450

```

Figure 11

**Figure 12**

**Figure 13**




Figure 14


**Figure 15**

FIG. 16

Hours After PGF-2 α Treatment

0

1

24




**Figure 17**

FIG. 18






FIG. 19



Southern Blot of Rat Genomic DNA



EcoR1 EcoRV BamH1

**Full-length
rat eIF-5A
cDNA probe**




FIG. 20




Figure 21




Figure 22

Transient Transfection





Figure 23

Mock Transfected
pHM6-LacZ
pHM6-Antisense 3' rF5A
pHM6-Sense rF5A



Anti-[HA]-
Peroxidase
(1:5000)

Figure 24

Enhancement of Apoptosis in COS-7 cells
expressing sense rat eIF-5A (Caspase
Activity)




Figure 25






Figure 26

**Hoescht Staining of Transformed
COS-7 Cells Deprived of Serum**



Mock Transformed



pHM6-Sense rat F5A

Figure 27





Figure 28



Sense rF5A 20 X



Mock Transfected 20 X




Figure 29




Figure 30




Figure 31




Figure 32






Figure 33

**Bcl-2 Expression Levels in Transiently Transfected COS-7
Cells Serum-Starved for 48 Hours**



**Bcl-2
(0.5 µg/ul)**

Figure 34




Figure 35




Figure 36




Figure 37

Figure 38

Sequence and Alignment of human eIF5A2 isolated from RKO cells with
sequence of human eIF5A2 in Genbank (ACCESSION XM_113401)

XM_113401	MADEIDFTTG	DAGASSTYPM	QCSALRKNGF	VVLKGRPCKI	VEMSTSKTGK
PCR	MADEIDFTTG	DAGASSTYPM	QCSALRKNGF	VVLKGRPCKI	VEMSTSKTGK
Consensus	MADEIDFTTG	DAGASSTYPM	QCSALRKNGF	VVLKGRPCKI	VEMSTSKTGK
	51				100
XM_113401	HGHAKVHLVG	IDIFTGKKYE	DICPSTHNMD	VPNIKRNDYQ	LICIQDGYLS
PCR	HGHAKVHLVG	IDIFTGKKYE	DICPSTHNMD	VPNIKRNDYQ	LICIQDGCLS
Consensus	HGHAKVHLVG	IDIFTGKKYE	DICPSTHNMD	VPNIKRNDYQ	LICIQDGcLS
	101				150
XM_113401	LLTETGEVRE	DLKLPEGELG	KEIEGKYNAG	EDVQVSVMCA	MSEYAVAIK
PCR	LLTETGEVRE	DLKLPEGELG	KEIEGKYNAG	EDVQVSVMCA	MSEYAVAIK
Consensus	LLTETGEVRE	DLKLPEGELG	KEIEGKYNAG	EDVQVSVMCA	MSEYAVAIK
	151				
XM_113401	PCK				
PCR	PCK				
Consensus	PCK				

Figure 39




Figure 40






Figure 41



Transfection efficiency = 30 %


Figure 42



Transfected Plasmid	Percent Apoptosis
Untransfected	21.2 %
pHM6-Lac Z	21.7 %
pHM6-eIF5A1	60.7 % (80%)*
pHM6-eIF5A2	20.5 %
pHM6-truncated eIF5A1	24.1 %

*Corrected for background apoptosis in untransfected cells and for transfection efficiency

Figure 43



SEQUENCE LISTING

<110> SENESCO, Inc.

<120> NUCLEIC ACIDS, POLYPEPTIDES, AND METHODS FOR MODULATING APOPTOSIS

<130> 10799/40

<140>

<141>

<150> 09/909,796

<151> 2001-07-23

<150> 10/141,647

<151> 2002-05-07

<160> 21

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1139

<212> DNA

<213> Rodent

<220>

<221> CDS

<222> (33)...(497)

<400> 1

caggtctaga gttggaatcg aagcctctta aa atg gca gat gat ttg gac ttc	53
Met Ala Asp Asp Leu Asp Phe	
1 5	

gag aca gga gat gca ggg gcc tca gcc acc ttc cca atg cag tgc tca	101
Glu Thr Gly Asp Ala Gly Ala Ser Ala Thr Phe Pro Met Gln Cys Ser	
10 15 20	

gca tta cgt aag aat ggt ttt gtg gtg ctc aag ggc cgg cca tgt aag	149
Ala Leu Arg Lys Asn Gly Phe Val Val Leu Lys Gly Arg Pro Cys Lys	
25 30 35	

atc gtc gag atg tct act tcg aag act ggc aag cat ggc cat gcc aag	197
Ile Val Glu Met Ser Thr Ser Lys Thr Gly Lys His Gly His Ala Lys	
40 45 50 55	

```

gtc cat ctg gtt ggt att gat att ttt act ggg aag aaa tat gaa gat 245
Val His Leu Val Gly Ile Asp Ile Phe Thr Gly Lys Lys Tyr Glu Asp
        60                65                70

atc tgc ccg tcg act cat aac atg gat gtc ccc aac atc aaa agg aat 293
Ile Cys Pro Ser Thr His Asn Met Asp Val Pro Asn Ile Lys Arg Asn
        75                80                85

gat ttc cag ctg att ggc atc cag gat ggg tac cta tcc ctg ctc cag 341
Asp Phe Gln Leu Ile Gly Ile Gln Asp Gly Tyr Leu Ser Leu Leu Gln
        90                95                100

gac agt ggg gag gta cga gag gac ctt cgt ctg cct gag gga gac ctt 389
Asp Ser Gly Glu Val Arg Glu Asp Leu Arg Leu Pro Glu Gly Asp Leu
       105                110                115

ggc aag gag att gag cag aag tat gac tgt gga gaa gag atc ctg atc 437
Gly Lys Glu Ile Glu Gln Lys Tyr Asp Cys Gly Glu Glu Ile Leu Ile
      120                125                130                135

aca gtg ctg tcc gcc atg aca gag gag gca gct gtt gca atc aag gcc 485
Thr Val Leu Ser Ala Met Thr Glu Glu Ala Ala Val Ala Ile Lys Ala
        140                145                150

atg gca aaa taa ctggcttcca gggtggcggt ggtggcagca gtgatccatg 537
Met Ala Lys *

```

```

agcctacaga ggcccctccc ccagctctgg ctggggccctt ggctggactc ctatccaatt 597
tatttgacgt tttattttgg ttttcctcac cccttcaaac tgtcgggggag accctgccct 657
tcacctagct cccttgGCCa ggcatgaggg agccatggcc ttgggtgaagc tacctgcctc 717
ttctctcgca gccctgatgg gggaaagggg gtgggtactg cctgtgggtt aggttcccct 777
ctcccttttt ctttttaatt caatttggaa tcagaaagct gtggattctg gcaaattggc 837
ttgtgtcctt tatcccactc aaacccatct ggtcccctgt tctccatagt ccttcacccc 897
caagcaccac tgacagactg gggaccagcc cccttcctg cctgtgtctc ttcccaaacc 957
cctctatagg ggtgacaaga agaggagggg gggaggggac acgatccctc ctcaggcatc 1017
tggaaggcc ttgcccccat gggctttacc ctttcctgtg ggcttttctc ctgacacatt 1077
tgttaaaaat caaacctgaa taaaactaca agtttaatat gaaaaaaaaa aaaaaaaaaa 1137
aa 1139

```

<210> 2
 <211> 154
 <212> PRT
 <213> Rodent

<400> 2
 Met Ala Asp Asp Leu Asp Phe Glu Thr Gly Asp Ala Gly Ala Ser Ala
 1 5 10 15
 Thr Phe Pro Met Gln Cys Ser Ala Leu Arg Lys Asn Gly Phe Val Val

			20					25					30				
Leu	Lys	Gly	Arg	Pro	Cys	Lys	Ile	Val	Glu	Met	Ser	Thr	Ser	Lys	Thr		
		35					40					45					
Gly	Lys	His	Gly	His	Ala	Lys	Val	His	Leu	Val	Gly	Ile	Asp	Ile	Phe		
	50					55					60						
Thr	Gly	Lys	Lys	Tyr	Glu	Asp	Ile	Cys	Pro	Ser	Thr	His	Asn	Met	Asp		
65					70					75				80			
Val	Pro	Asn	Ile	Lys	Arg	Asn	Asp	Phe	Gln	Leu	Ile	Gly	Ile	Gln	Asp		
			85					90					95				
Gly	Tyr	Leu	Ser	Leu	Leu	Gln	Asp	Ser	Gly	Glu	Val	Arg	Glu	Asp	Leu		
		100						105					110				
Arg	Leu	Pro	Glu	Gly	Asp	Leu	Gly	Lys	Glu	Ile	Glu	Gln	Lys	Tyr	Asp		
		115					120					125					
Cys	Gly	Glu	Glu	Ile	Leu	Ile	Thr	Val	Leu	Ser	Ala	Met	Thr	Glu	Glu		
	130					135					140						
Ala	Ala	Val	Ala	Ile	Lys	Ala	Met	Ala	Lys								
145					150												

<210> 3
 <211> 462
 <212> DNA
 <213> Rodent

<400> 3
 atggcagatg acttggactt cgagacagga gatgcagggg cctcagccac cttcccaatg 60
 cagtgtctcag cattacgtaa gaatggcttt gtggtgctca aaggccggcc atgtaagatc 120
 gtcgagatgt ctacttcgaa gactggcaag cacggccacg ccaagggtcca tctggttggt 180
 attgacatct ttactgggaa gaaatatgaa gatattctgcc cgtcaactca taatatggat 240
 gtccccaaca tcaaaaggaa tgacttccag ctgattggca tccaggatgg gtacctatca 300
 ctgctccagg acagcgggga ggtacgagag gaccttcgct tccctgaggg agaccttggt 360
 aaggagattg agcagaagta cgactgtgga gaagagatcc tgatcacggt gctgtctgcc 420
 atgacagagg aggcagctgt tgcaatcaag gccatggcaa aa 462

<210> 4
 <211> 462
 <212> DNA
 <213> Rodent

<220>
 <221> misc_feature
 <222> (1)...(462)
 <223> n = A,T,C or G

<400> 4
 atggcagacg aaattgatatt cactactgga gatgccgggg cttccagcac ttaccctatg 60
 cagtgtctcgg ccttgcgcaa aaacggcttc gtggtgctca aaggacgacc atgcaaaaata 120
 gtggagatgt caacttccaa aactggaaag catggatcatg ccaagggttca ccttggttgga 180
 attgatattt tcacgggcaa aaaatatgaa gatatttgct cttctactca caacatggat 240

```

gttcctaaata ttaagagaaa tgattatcaa ctgatatgca ttcaagatgg ttacctttcc 300
ctgctgacag aaactgggtga agttcgtgag gatcttaaac tgccagaagg tgaactaggc 360
aaagaaatag agggaaaata caatgcaggt gaagatgtac aggtgtctgt catgtgtgca 420
atgagtgaag aatatgctgt agccataaaa ccctnngcaa at 462

```

<210> 5
 <211> 462
 <212> DNA
 <213> Rodent

```

<400> 5
atggcagatg atttggactt cgagacagga gatgcagggg cctcagccac cttcccaatg 60
cagtgtcag cattacgtaa gaatggtttt gtgggtgctca aaggccggcc atgtaagatc 120
gtcgagatgt ctacttcgaa gactggcaag catggccatg ccaagggtcca tctggttggc 180
attgacattht ttactgggaa gaaatatgaa gatattctgcc cgtcgactca taatatggat 240
gtccccaaca tcaaacggaa tgacttccag ctgattggca tccaggatgg gtacctatcc 300
ctgctccagg acagtgggga ggtacgagag gaccttcgtc tgcctgaagg agaccttggc 360
aaggagattg agcagaagta tgactgtgga gaagagatcc tgatcacagt gctgtctgcc 420
atgacagagg aggcagctgt tgcaatcaag gccatggcaa aa 462

```

<210> 6
 <211> 606
 <212> DNA
 <213> Rodent

<220>
 <221> CDS
 <222> (1)...(456)

```

<400> 6
gct gtg tat tat tgg gcc cat aag aac cac ata cct gtg ctg agt cct 48
Ala Val Tyr Tyr Trp Ala His Lys Asn His Ile Pro Val Leu Ser Pro
1 5 10 15

gca ctc aca gac ggc tca ctg ggt gac atg atc ttt ttc cat tcc tat 96
Ala Leu Thr Asp Gly Ser Leu Gly Asp Met Ile Phe Phe His Ser Tyr
20 25 30

aaa aac cca ggc ttg gtc ctg gac atc gtt gaa gac ctg cgg ctc atc 144
Lys Asn Pro Gly Leu Val Leu Asp Ile Val Glu Asp Leu Arg Leu Ile
35 40 45

aac atg cag gcc att ttc gcc aag cgc act ggg atg atc atc ctg ggt 192
Asn Met Gln Ala Ile Phe Ala Lys Arg Thr Gly Met Ile Ile Leu Gly
50 55 60

gga ggc gtg gtc aag cac cac atc gcc aat gct aac ctc atg cgg aat 240
Gly Gly Val Val Lys His His Ile Ala Asn Ala Asn Leu Met Arg Asn
65 70 75 80

```

gga gct gac tac gct gtt tat atc aac aca gcc cag gag ttt gat ggc 288
 Gly Ala Asp Tyr Ala Val Tyr Ile Asn Thr Ala Gln Glu Phe Asp Gly
 85 90 95

tca gac tca gga gcc cgg cca gat gag gct gtc tcc tgg ggc aag atc 336
 Ser Asp Ser Gly Ala Arg Pro Asp Glu Ala Val Ser Trp Gly Lys Ile
 100 105 110

cgg atg gat gca cag cca gta aag gtc tat gct gat gca tct ctg gtt 384
 Arg Met Asp Ala Gln Pro Val Lys Val Tyr Ala Asp Ala Ser Leu Val
 115 120 125

ttc ccc ttg ctg gtg gct gag aca ttc gcc caa aag gca gat gcc ttc 432
 Phe Pro Leu Leu Val Ala Glu Thr Phe Ala Gln Lys Ala Asp Ala Phe
 130 135 140

aga gct gag aag aat gag gac tga gcagatgggt aaagacggag gcttctgcc 486
 Arg Ala Glu Lys Asn Glu Asp *
 145 150

cacctttatt tattatttgc ataccaaccc ctcttgggcc ctctccttgg tcagcagcat 546
 cttgagaata aatggccttt ttgttggttt ctgtaaaaaa aggactttta aaaaaaaaaa 606

<210> 7
 <211> 151
 <212> PRT
 <213> Rodent

<400> 7
 Ala Val Tyr Tyr Trp Ala His Lys Asn His Ile Pro Val Leu Ser Pro
 1 5 10 15
 Ala Leu Thr Asp Gly Ser Leu Gly Asp Met Ile Phe Phe His Ser Tyr
 20 25 30
 Lys Asn Pro Gly Leu Val Leu Asp Ile Val Glu Asp Leu Arg Leu Ile
 35 40 45
 Asn Met Gln Ala Ile Phe Ala Lys Arg Thr Gly Met Ile Ile Leu Gly
 50 55 60
 Gly Gly Val Val Lys His His Ile Ala Asn Ala Asn Leu Met Arg Asn
 65 70 75 80
 Gly Ala Asp Tyr Ala Val Tyr Ile Asn Thr Ala Gln Glu Phe Asp Gly
 85 90 95
 Ser Asp Ser Gly Ala Arg Pro Asp Glu Ala Val Ser Trp Gly Lys Ile
 100 105 110
 Arg Met Asp Ala Gln Pro Val Lys Val Tyr Ala Asp Ala Ser Leu Val
 115 120 125
 Phe Pro Leu Leu Val Ala Glu Thr Phe Ala Gln Lys Ala Asp Ala Phe
 130 135 140

Arg Ala Glu Lys Asn Glu Asp
145 150

<210> 8
<211> 453
<212> DNA
<213> Rodent

<400> 8
tccgtgtatt actgggcccga gaagaaccac atccctgtgt ttagtcccgc acttacagac 60
ggctcgctgg gcgacatgat cttcttccat tcctacaaga acccgggcct ggtcctggac 120
atcgttgagg acctgaggct catcaacaca caggccatct ttgccaagtg cactgggatg 180
atcattcttg gcgggggcgt ggtcaagcac cacattgcca atgccaacct catgcggaac 240
ggggccgact acgctgttta catcaacaca gcccgaggag ttgatggctc tgactcaggt 300
gcccgaccag acgaggctgt ctctgggggc aagatccggg tggatgcaca gcccgtaag 360
gtctatgctg acgcctccct ggtcttcccc ctgcttgtgg ctgaaacctt tgcccagaag 420
atggatgcct tcatgcatga gaagaacgag gac 453

<210> 9
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<221> misc_feature
<222> (1)...(20)
<223> n = A,T,C or G

<400> 9
tcsaarachg gnaagcaygg 20

<210> 10
<211> 42
<212> DNA
<213> Rodent

<220>
<223> Primer

<400> 10
gcgaagcttc catggctcga gttttttttt tttttttttt tt 42

<210> 11
<211> 972
<212> DNA
<213> Rodent

<220>

<221> CDS

<222> (1)...(330)

<400> 11

tcg aag acc ggt aag cac ggc cat gcc aag gtc cat ctg gtt ggt att	48
Ser Lys Thr Gly Lys His Gly His Ala Lys Val His Leu Val Gly Ile	
1 5 10 15	
gat att ttt act ggg aag aaa tat gaa gat atc tgc ccg tcg act cat	96
Asp Ile Phe Thr Gly Lys Lys Tyr Glu Asp Ile Cys Pro Ser Thr His	
20 25 30	
aac atg gat gtc ccc aac atc aaa agg aat gat ttc cag ctg att ggc	144
Asn Met Asp Val Pro Asn Ile Lys Arg Asn Asp Phe Gln Leu Ile Gly	
35 40 45	
atc cag gat ggg tac cta tcc ctg ctc cag gac agt ggg gag gta cga	192
Ile Gln Asp Gly Tyr Leu Ser Leu Leu Gln Asp Ser Gly Glu Val Arg	
50 55 60	
gag gac ctt cgt ctg cct gag gga gac ctt ggc aag gag att gag cag	240
Glu Asp Leu Arg Leu Pro Glu Gly Asp Leu Gly Lys Glu Ile Glu Gln	
65 70 75 80	
aag tat gac tgt gga gaa gag atc ctg atc aca gtg ctg tcc gcc atg	288
Lys Tyr Asp Cys Gly Glu Glu Ile Leu Ile Thr Val Leu Ser Ala Met	
85 90 95	
aca gag gag gca gct gtt gca atc aag gcc atg gca aaa taa	330
Thr Glu Glu Ala Ala Val Ala Ile Lys Ala Met Ala Lys *	
100 105	
ctggccttcca ggggtggcgggt ggtggcagca gtgatccatg agcctacaga ggcccctccc	390
ccagctctgg ctgggccctt ggctggactc ctatccaatt tatttgacgt tttatatttg	450
ttttcctcac cccttcaaac tgtcggggag accctgccct tcacctagct cccttggcca	510
ggcatgaggg agccatggcc ttggtgaagc tacctgcctc ttctctcgca gccctgatgg	570
gggaaaggga gtgggtactg cctgtggttt aggttcccct ctcccttttt ctttttaatt	630
caatttgga tcagaaagct gtggattctg gcaaattggc ttgtgtcctt tatcccactc	690
aaacccatct ggtcccctgt tctccatagt ccttcacccc caagcaccac tgacagactg	750
gggaccagcc cccttccctg cctgtgtctc ttcccaaacc cctctatagg ggtgacaaga	810
agaggagggg gggaggggac acgatccctc ctcaaggcatc tgggaaggcc ttgcccccat	870
gggctttacc ctttcctgtg ggcttttctc ctgacacatt tgttaaaaaat caaacctgaa	930
taaaactaca agttttaatat gaaaaaaaaa aaaaaaaaaa aa	972

<210> 12

<211> 109

<212> PRT

<213> Rodent

<400> 12

Ser	Lys	Thr	Gly	Lys	His	Gly	His	Ala	Lys	Val	His	Leu	Val	Gly	Ile
1				5					10					15	
Asp	Ile	Phe	Thr	Gly	Lys	Lys	Tyr	Glu	Asp	Ile	Cys	Pro	Ser	Thr	His
			20					25					30		
Asn	Met	Asp	Val	Pro	Asn	Ile	Lys	Arg	Asn	Asp	Phe	Gln	Leu	Ile	Gly
		35					40					45			
Ile	Gln	Asp	Gly	Tyr	Leu	Ser	Leu	Leu	Gln	Asp	Ser	Gly	Glu	Val	Arg
	50					55					60				
Glu	Asp	Leu	Arg	Leu	Pro	Glu	Gly	Asp	Leu	Gly	Lys	Glu	Ile	Glu	Gln
65					70					75					80
Lys	Tyr	Asp	Cys	Gly	Glu	Glu	Ile	Leu	Ile	Thr	Val	Leu	Ser	Ala	Met
			85						90					95	
Thr	Glu	Glu	Ala	Ala	Val	Ala	Ile	Lys	Ala	Met	Ala	Lys			
			100					105							

<210> 13

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 13

caggtctaga gttggaatcg aagc

24

<210> 14

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 14

atatctcgag ccttgattgc aacagctgcc

30

<210> 15

<211> 489

<212> DNA

<213> Rodent

<220>

<221> CDS

<222> (33)...(485)

<400> 15

caggtctaga gttggaatcg aagcctctta aa	atg gca gat gat ttg gac ttc	53
	Met Ala Asp Asp Leu Asp Phe	
	1 5	
gag aca gga gat gca ggg gcc tca gcc acc ttc cca atg cag tgc tca	101	
Glu Thr Gly Asp Ala Gly Ala Ser Ala Thr Phe Pro Met Gln Cys Ser		
	10 15 20	
gca tta cgt aag aat ggt ttt gtg gtg ctc aag ggc cgg cca tgt aag	149	
Ala Leu Arg Lys Asn Gly Phe Val Val Leu Lys Gly Arg Pro Cys Lys		
	25 30 35	
atc gtc gag atg tct act tcg aag act ggc aag cat ggc cat gcc aag	197	
Ile Val Glu Met Ser Thr Ser Lys Thr Gly Lys His Gly His Ala Lys		
	40 45 50 55	
gtc cat ctg gtt ggt att gat att ttt act ggg aag aaa tat gaa gat	245	
Val His Leu Val Gly Ile Asp Ile Phe Thr Gly Lys Lys Tyr Glu Asp		
	60 65 70	
atc tgc ccg tcg act cat aac atg gat gtc ccc aac atc aaa agg aat	293	
Ile Cys Pro Ser Thr His Asn Met Asp Val Pro Asn Ile Lys Arg Asn		
	75 80 85	
gat ttc cag ctg att ggc atc cag gat ggg tac cta tcc ctg ctc cag	341	
Asp Phe Gln Leu Ile Gly Ile Gln Asp Gly Tyr Leu Ser Leu Leu Gln		
	90 95 100	
gac agt ggg gag gta cga gag gac ctt cgt ctg cct gag gga gac ctt	389	
Asp Ser Gly Glu Val Arg Glu Asp Leu Arg Leu Pro Glu Gly Asp Leu		
	105 110 115	
ggc aag gag att gag cag aag tat gac tgt gga gaa gag atc ctg atc	437	
Gly Lys Glu Ile Glu Gln Lys Tyr Asp Cys Gly Glu Glu Ile Leu Ile		
	120 125 130 135	
aca gtg ctg tcc gcc atg aca gag gag gca gct gtt gca atc aag gct	485	
Thr Val Leu Ser Ala Met Thr Glu Glu Ala Ala Val Ala Ile Lys Ala		
	140 145 150	
cgag	489	

<210> 16

<211> 151

<212> PRT

<213> Rodent

<400> 16

Met	Ala	Asp	Asp	Leu	Asp	Phe	Glu	Thr	Gly	Asp	Ala	Gly	Ala	Ser	Ala
1				5					10					15	
Thr	Phe	Pro	Met	Gln	Cys	Ser	Ala	Leu	Arg	Lys	Asn	Gly	Phe	Val	Val
			20					25					30		
Leu	Lys	Gly	Arg	Pro	Cys	Lys	Ile	Val	Glu	Met	Ser	Thr	Ser	Lys	Thr
		35					40					45			
Gly	Lys	His	Gly	His	Ala	Lys	Val	His	Leu	Val	Gly	Ile	Asp	Ile	Phe
	50					55					60				
Thr	Gly	Lys	Lys	Tyr	Glu	Asp	Ile	Cys	Pro	Ser	Thr	His	Asn	Met	Asp
65					70				75					80	
Val	Pro	Asn	Ile	Lys	Arg	Asn	Asp	Phe	Gln	Leu	Ile	Gly	Ile	Gln	Asp
				85					90					95	
Gly	Tyr	Leu	Ser	Leu	Leu	Gln	Asp	Ser	Gly	Glu	Val	Arg	Glu	Asp	Leu
			100					105					110		
Arg	Leu	Pro	Glu	Gly	Asp	Leu	Gly	Lys	Glu	Ile	Glu	Gln	Lys	Tyr	Asp
		115					120					125			
Cys	Gly	Glu	Glu	Ile	Leu	Ile	Thr	Val	Leu	Ser	Ala	Met	Thr	Glu	Glu
	130					135					140				
Ala	Ala	Val	Ala	Ile	Lys	Ala									
145					150										

<210> 17

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 17

gtctgtgtat tattgggccc

20

<210> 18

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 18

gcgaagcttc catggctcga gttttttttt tttttttttt tt

42

<210> 19

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 19

ttgaaggggt gaggaaaa

18

<210> 20

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 20

ttgagtggga taaag

15

<210> 21

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 21

aatcatctgc cattttaa

18