



(86) Date de dépôt PCT/PCT Filing Date: 2012/03/29
(87) Date publication PCT/PCT Publication Date: 2012/10/11
(85) Entrée phase nationale/National Entry: 2013/10/02
(86) N° demande PCT/PCT Application No.: EP 2012/055595
(87) N° publication PCT/PCT Publication No.: 2012/136549
(30) Priorité/Priority: 2011/04/05 (EP11161142.2)

(51) Cl.Int./Int.Cl. *A61K 31/517* (2006.01),
A61P 35/00 (2006.01), *G01N 33/48* (2006.01)
(71) Demandeur/Applicant:
BAYER INTELLECTUAL PROPERTY GMBH, DE
(72) Inventeurs/Inventors:
LIU, NINGSHU, DE;
SCHNEIDER, CLAUDIA, DE
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : UTILISATION DE 2,3-DIHYDROIMIDAZO[1,2-C]QUINAZOLINES SUBSTITUEES
(54) Title: USE OF SUBSTITUTED 2,3-DIHYDROIMIDAZO[1,2-C]QUINAZOLINES

(57) **Abrégé/Abstract:**

The present invention relates to: - use of a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or of a pharmaceutical composition containing same, as a sole active agent, or of a combination of a) said compound or a pharmaceutical composition containing said compound and b) one or more further active agents, for the preparation of a medicament for the treatment or prophylaxis of cancer; - combinations of a) said compound and b) one or more further active agents; - a pharmaceutical composition comprising said compound as a sole active agent for the treatment of breast cancer; - a pharmaceutical composition comprising a combination of a) said compound and b) one or more further active agents; - use of biomarkers involved in the modification of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN for predicting the sensitivity and /or resistance of a cancer patient to said compound and providing a rationale-based synergistic combination as defined herein to increase sensitivity and/or to overcome resistance; and - a method of determining the level of a component of one or more of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 October 2012 (11.10.2012)

WIPO | PCT

(10) International Publication Number
WO 2012/136549 A1

(51) International Patent Classification:

A61K 31/517 (2006.01) *A61P 35/00* (2006.01)
G01N 33/48 (2006.01)

(21) International Application Number:

PCT/EP2012/055595

(22) International Filing Date:

29 March 2012 (29.03.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

11161142.2 5 April 2011 (05.04.2011) EP

(71) **Applicant** (for all designated States except US): **BAYER PHARMA AKTIENGESELLSCHAFT** [DE/DE]; Müller Strasse 178, 13353 Berlin (DE).

(72) **Inventors; and**

(75) **Inventors/Applicants** (for US only): **LIU, Ningshu** [DE/DE]; Limonenstr. 28, 12203 Berlin (DE).
SCHNEIDER, Claudia [DE/DE]; Bastianstr. 10, 13357 Berlin (DE).

(74) **Agent: BAYER PHARMA AKTIENGESELLSCHAFT**; Law and Patents, Patents and Licensing, Müller Strasse 178, 13353 Berlin (DE).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))

(54) **Title:** USE OF SUBSTITUTED 2,3-DIHYDROIMIDAZO[1,2-C]QUINAZOLINES

(57) **Abstract:** The present invention relates to: - use of a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or of a pharmaceutical composition containing same, as a sole active agent, or of a combination of a) said compound or a pharmaceutical composition containing said compound and b) one or more further active agents, for the preparation of a medicament for the treatment or prophylaxis of cancer; - combinations of a) said compound and b) one or more further active agents; - a pharmaceutical composition comprising said compound as a sole active agent for the treatment of breast cancer; - a pharmaceutical composition comprising a combination of a) said compound and b) one or more further active agents; - use of biomarkers involved in the modification of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN for predicting the sensitivity and / or resistance of a cancer patient to said compound and providing a rationale-based synergistic combination as defined herein to increase sensitivity and/or to overcome resistance; and - a method of determining the level of a component of one or more of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN.



WO 2012/136549 A1

USE OF SUBSTITUTED 2,3-DIHYDROIMIDAZO[1,2-C]QUINAZOLINES

The present invention relates to :

- 5 - use of a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or of a pharmaceutical composition containing same, as a sole active agent, or of a combination of a) said compound or a pharmaceutical composition containing said compound and b) one or more further active agents, for the preparation of a medicament for the treatment or prophylaxis of
10 cancer, e.g. breast cancer, which is classified into several subtypes in the clinic, such as hormone receptor positive breast cancer, Her2 receptor positive breast cancer, triple negative breast cancer and inflammatory breast cancer, as a single agent or in combination with one or more other active agents ;
 - 15 - combinations of a) said compound and b) one or more further active agents ;
 - a pharmaceutical composition comprising said compound as a sole
20 active agent for the treatment of cancer ;
 - a pharmaceutical composition comprising a combination of a) said compound and b) one or more further active agents ;
 - 25 - use of biomarkers involved in the modification of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN for predicting the sensitivity and/or resistance of a cancer patient to said compound and providing a rationale-based synergistic combination as defined herein to increase sensitivity and/or to
30 overcome resistance ;
- and

- a method of determining the level of a component of one or more of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN.

5 BACKGROUND OF THE INVENTION

In recent decades the concept of developing anti-cancer medications which target abnormally active protein kinases has led to a number of successes. In addition to the actions of protein kinases, lipid kinases
10 also play an important role in generating critical regulatory second messengers. The PI3K family of lipid kinases generates 3'-phosphoinositides that bind to and activate a variety of cellular targets, initiating a wide range of signal transduction cascades (Vanhaesebroeck *et al.*, 2001; Toker, 2002; Pendaries *et al.*, 2003; Downes *et al.*, 2005).
15 These cascades ultimately induce changes in multiple cellular processes, including cell proliferation, cell survival, differentiation, vesicle trafficking, migration, and chemotaxis.

PI3Ks can be divided into three distinct classes based upon differences in
20 both structure, and substrate preference. While members of the Class II family of PI3Ks have been implicated in the regulation of tumor growth (Brown and Shepard, 2001; Traer *et al.*, 2006), the bulk of research has focused on the Class I enzymes and their role in cancer (Vivanco and Sawyers, 2002; Workman, 2004, Chen *et al.*, 2005; Hennessey *et al.*,
25 2005; Stauffer *et al.*, 2005; Stephens *et al.*, 2005; Cully *et al.*, 2006).

Class I PI3Ks have traditionally been divided into two distinct sub-classes based upon differences in protein subunit composition. The Class I_A PI3Ks are comprised of a catalytic p110 catalytic subunit (p110 α , p110 β or
30 p110 γ) heterodimerized with a member of the p85 regulatory subunit family. In contrast, the Class I_B PI3K catalytic subunit (p110 γ) heterodimerizes with a distinct p101 regulatory subunit (reviewed by

Vanhaesebroeck and Waterfield, 1999; Funaki *et al.*, 2000; Katso *et al.*, 2001). The C-terminal region of these proteins contains a catalytic domain that possesses distant homology to protein kinases. The PI3K γ structure is similar to Class I α p110s, but lacks the N-terminal p85 binding site (Domin and Waterfield, 1997). Though similar in overall structure, the homology between catalytic p110 subunits is low to moderate. The highest homology between the PI3K isoforms is in the kinase pocket of the kinase domain.

10 The Class I PI3K isoforms associate with activated receptor tyrosine kinases (RTKs) (including PDGFR, EGFR, VEGFR, IGF1-R, c-KIT, CSF-R and Met), cytokine receptors, GPCRs, integrins, or with tyrosine phosphorylated adapter proteins (such as Grb2, Cbl, IRS-1 or Gab1), via their p85 regulatory subunits resulting in stimulation of the lipid kinase activity. Activation of the lipid kinase activity of the p110 β and p110 γ isoforms has been shown to occur in response to binding to activated forms of the ras Oncogene (Kodaki *et al.*, 1994). In fact, the oncogenic activity of these isoforms may require binding to ras (Kang *et al.*, 2006). In contrast, the p110 α and p110 δ isoforms exhibit oncogenic activity independent of ras binding, through constitutive activation of Akt.

Class I PI3Ks catalyze the conversion of PI(4,5)P₂ [PIP₂] to PI(3,4,5)P₃ [PIP₃]. The production of PIP₃ by PI3K affects multiple signaling processes that regulate and coordinate the biological end points of cell proliferation, cell survival, differentiation and cell migration. PIP₃ is bound by Pleckstrin-Homology (PH) domain-containing proteins, including the phosphoinositide-dependent kinase, PDK1 and the Akt proto-oncogene product, localizing these proteins in regions of active signal transduction and also contributing directly to their activation (Klippel *et al.*, 1997; Fleming *et al.*, 2000; Itoh and Takenawa, 2002; Lemmon, 2003). This co-localization of PDK1 with Akt facilitates the phosphorylation and activation of Akt. Carboxy-terminal phosphorylation

of Akt on Ser⁴⁷³ promotes phosphorylation of Thr³⁰⁸ in the Akt activation loop (Chan and Tsichlis, 2001; Hodgekinson *et al.*, 2002; Scheid *et al.*, 2002; Hresko *et al.*, 2003). Once active, Akt phosphorylates and regulates multiple regulatory kinases of pathways that directly influence
5 cell cycle progression and cell survival.

Many of the effects of Akt activation are mediated via its negative regulation of pathways which impact cell survival and which are commonly dysregulated in cancer. Akt promotes tumor cell survival by
10 regulating components of the apoptotic and cell cycle machinery. Akt is one of several kinases that phosphorylate and inactivate pro-apoptotic BAD proteins (del Paso *et al.*, 1997; Pastorino *et al.*, 1999). Akt may also promote cell survival through blocking cytochrome C-dependent caspase activation by phosphorylating Caspase 9 on Ser¹⁹⁶ (Cardone *et al.*, 1998).

15 Akt impacts gene transcription on several levels. The Akt-mediated phosphorylation of the MDM2 E3 ubiquitin ligase on Ser¹⁶⁶ and Ser¹⁸⁶ facilitates the nuclear import of MDM2 and the formation and activation of the ubiquitin ligase complex. Nuclear MDM2 targets the p53 tumor suppressor for degradation, a process that can be blocked by LY294002
20 (Yap *et al.*, 2000; Ogarawa *et al.*, 2002). Downregulation of p53 by MDM2 negatively impacts the transcription of p53-regulated pro-apoptotic genes (e.g. Bax, Fas, PUMA and DR5), the cell cycle inhibitor, p21^{Cip1}, and the PTEN tumor suppressor (Momand *et al.*, 2000; Hupp *et al.*, 2000; Mayo *et al.*, 2002; Su *et al.*, 2003). Similarly, the Akt-mediated
25 phosphorylation of the Forkhead transcription factors FKHR, FKHL and AFX (Kops *et al.*, 1999; Tang *et al.*, 1999), facilitates their binding to 14-3-3 proteins and export from the cell nucleus to the cytosol (Brunet *et al.*, 1999). This functional inactivation of Forkhead activity also impacts
30 pro-apoptotic and pro-angiogenic gene transcription including the transcription of Fas ligand (Ciechomska *et al.*, 2003) Bim, a pro-apoptotic Bcl-2 family member (Dijkers *et al.*, 2000), and the

Angiopoietin-1 (Ang-1) antagonist, Ang-2 (Daly *et al.*, 2004). Forkhead transcription factors regulate the expression of the cyclin-dependent kinase (Cdk) inhibitor p27^{Kip1}. Indeed, PI3K inhibitors have been demonstrated to induce p27^{Kip1} expression resulting in Cdk1 inhibition,
5 cell cycle arrest and apoptosis (Dijkers *et al.*, 2000). Akt is also reported to phosphorylate p21^{Cip1} on Thr¹⁴⁵ and p27^{Kip1} on Thr¹⁵⁷ facilitating their association with 14-3-3 proteins, resulting in nuclear export and cytoplasmic retention, preventing their inhibition of nuclear Cdks (Zhou *et al.*, 2001; Motti *et al.*, 2004; Sekimoto *et al.*, 2004). In addition to
10 these effects, Akt phosphorylates IKK (Romashkova and Makarov, 1999), leading to the phosphorylation and degradation of I κ B and subsequent nuclear translocation of NF κ B, resulting in the expression of survival genes such as IAP and Bcl-X_L.

15 The PI3K/Akt pathway is also linked to the suppression of apoptosis through the JNK and p38^{MAPK} MAP Kinases that are associated with the induction of apoptosis. Akt is postulated to suppress JNK and p38^{MAPK} signaling through the phosphorylation and inhibition of two JNK/p38 regulatory kinases, Apoptosis Signal-regulating Kinase 1 (ASK1) (Kim *et al.*, 2001; Liao and Hung, 2003; Yuan *et al.*, 2003), and Mixed Lineage
20 Kinase 3 (MLK3) (Lopez-Illasaca *et al.*, 1997; Barthwal *et al.*, 2003; Figueroa *et al.*, 2003;). The induction of p38^{MAPK} activity is observed in tumors treated with cytotoxic agents and is required for those agents to induce cell death (reviewed by Olson and Hallahan, 2004). Thus,
25 inhibitors of the PI3K pathway may promote the activities of co-administered cytotoxic drugs.

An additional role for PI3K/Akt signaling involves the regulation of cell cycle progression through modulation of Glycogen Synthase Kinase 3
30 (GSK3) activity. GSK3 activity is elevated in quiescent cells, where it phosphorylates cyclin D₁ on Ser²⁸⁶, targeting the protein for ubiquitination and degradation (Diehl *et al.*, 1998) and blocking entry

into S-phase. Akt inhibits GSK3 activity through phosphorylation on Ser⁹ (Cross *et al.*, 1995). This results in the elevation of Cyclin D₁ levels which promotes cell cycle progression. Inhibition of GSK3 activity also impacts cell proliferation through activation of the wnt/beta-catenin signaling pathway (Abbosh and Nephew, 2005; Naito *et al.*, 2005; Wilker *et al.*, 2005; Kim *et al.*, 2006; Segrelles *et al.*, 2006). Akt mediated phosphorylation of GSK3 results in stabilization and nuclear localization of the beta-catenin protein, which in turn leads to increased expression of c-myc and cyclin D1, targets of the beta-catenin/Tcf pathway.

10

Although PI3K signaling is utilized by many of the signal transduction networks associated with both oncogenes and tumor suppressors, PI3K and its activity have been linked directly to cancer. Overexpression of both the p110 α and p110 β isoforms has been observed in bladder and colon tumors and cell lines, and overexpression generally correlates with increased PI3K activity (Bénistant *et al.*, 2000). Overexpression of p110 α has also been reported in ovarian and cervical tumors and tumor cell lines, as well as in squamous cell lung carcinomas. The overexpression of p110 α in cervical and ovarian tumor lines is associated with increased PI3K activity (Shayesteh *et al.*, 1999; Ma *et al.*, 2000). Elevated PI3K activity has been observed in colorectal carcinomas (Phillips *et al.*, 1998) and increased expression has been observed in breast carcinomas (Gershtein *et al.*, 1999).

Over the last few years, somatic mutations in the gene encoding p110 α (PIK3CA) have been identified in numerous cancers. The data collected to date suggests that PIK3CA is mutated in approximately 32% of colorectal cancers (Samuels *et al.*, 2004; Ikenoue *et al.*, 2005), 18-40% of breast cancers (Bachman *et al.*, 2004; Campbell *et al.*, 2004; Levine *et al.*, 2005; Saal *et al.*, 2005; Wu *et al.*, 2005), 27% of glioblastomas (Samuels *et al.*, 2004; Hartmann *et al.*, 2005, Gallia *et al.*, 2006), 25% of gastric cancers (Byun *et al.*, 2003; Samuels *et al.*, 2004; Li *et al.*, 2005),

30

36% of hepatocellular carcinomas (Lee *et al.*, 2005), 4-12% of ovarian cancers (Levine *et al.*, 2005; Wang *et al.*, 2005), 4% of lung cancers (Samuels *et al.*, 2004; Whyte and Holbeck, 2006), and up to 40% of endometrial cancers (Oda *et al.*, 2005). PIK3CA mutations have been reported in oligodendroma, astrocytoma, medulloblastoma, and thyroid tumors as well (Broderick *et al.*, 2004; Garcia-Rostan *et al.*, 2005). Based upon the observed high frequency of mutation, PIK3CA is one of the two most frequently mutated genes associated with cancer, the other being K-ras. More than 80% of the PIK3CA mutations cluster within two regions of the protein, the helical (E545K) and catalytic (H1047R) domains. Biochemical analysis and protein expression studies have demonstrated that both mutations lead to increased constitutive p110 α catalytic activity and are in fact, oncogenic (Bader *et al.*, 2006; Kang *et al.*, 2005; Samuels *et al.*, 2005; Samuels and Ericson, 2006). Recently, it has been reported that PIK3CA knockout mouse embryo fibroblasts are deficient in signaling downstream from various growth factor receptors (IGF-1, Insulin, PDGF, EGF), and are resistant to transformation by a variety of oncogenic RTKs (IGFR, wild-type EGFR and somatic activating mutants of EGFR, Her2/Neu)(Zhao *et al.*, 2006).

20

Functional studies of PI3K *in vivo* have demonstrated that siRNA-mediated downregulation of p110 β inhibits both Akt phosphorylation and HeLa cell tumor growth in nude mice (Czauderna *et al.*, 2003). In similar experiments, siRNA-mediated downregulation of p110 β was also shown to inhibit the growth of malignant glioma cells *in vitro* and *in vivo* (Pu *et al.*, 2006). Inhibition of PI3K function by dominant-negative p85 regulatory subunits can block mitogenesis and cell transformation (Huang *et al.*, 1996; Rahimi *et al.*, 1996). Several somatic mutations in the genes encoding the p85 α and p85 β regulatory subunits of PI3K that result in elevated lipid kinase activity have been identified in a number of cancer cells as well (Janssen *et al.*, 1998; Jimenez *et al.*, 1998; Philp *et al.*, 2001; Jucker *et al.*, 2002; Shekar *et al.*, 2005). Neutralizing PI3K

30

antibodies also block mitogenesis and can induce apoptosis *in vitro* (Roche *et al.*, 1994; Roche *et al.*, 1998; Bénistant *et al.*, 2000). *In vivo* proof-of-principle studies using the PI3K inhibitors LY294002 and wortmannin, demonstrate that inhibition of PI3K signaling slows tumor growth *in vivo* (Powis *et al.*, 1994; Shultz *et al.*, 1995; Semba *et al.*, 2002; Ihle *et al.*, 2004).

Overexpression of Class I PI3K activity, or stimulation of their lipid kinase activities, is associated with resistance to both targeted (such as imatinib and trastuzumab) and cytotoxic chemotherapeutic approaches, as well as radiation therapy (West *et al.*, 2002; Gupta *et al.*, 2003; Osaki *et al.*, 2004; Nagata *et al.*, 2004; Gottschalk *et al.*, 2005; Kim *et al.*, 2005). Activation of PI3K has also been shown to lead to expression of multidrug resistant protein-1 (MRP-1) in prostate cancer cells and the subsequent induction of resistance to chemotherapy (Lee *et al.*, 2004).

The importance of PI3K signaling in tumorigenesis is further underscored by the findings that the PTEN tumor suppressor, a PI(3)P phosphatase, is among the most commonly inactivated genes in human cancers (Li *et al.*, 1997, Steck *et al.*, 1997; Ali *et al.*, 1999; Ishii *et al.*, 1999). PTEN dephosphorylates PI(3,4,5)P₃ to PI(4,5)P₂ thereby antagonizing PI3K-dependent signaling. Cells containing functionally inactive PTEN have elevated levels of PIP₃, high levels of activity of PI3K signaling (Haas-Kogan *et al.*, 1998; Myers *et al.*, 1998; Taylor *et al.*, 2000), increased proliferative potential, and decreased sensitivity to pro-apoptotic stimuli (Stambolic *et al.*, 1998). Reconstitution of a functional PTEN suppresses PI3K signaling (Taylor *et al.*, 2000), inhibits cell growth and re-sensitizes cells to pro-apoptotic stimuli (Myers *et al.*, 1998; Zhao *et al.*, 2004). Similarly, restoration of PTEN function in tumors lacking functional PTEN inhibits tumor growth *in vivo* (Stahl *et al.*, 2003; Su *et al.*, 2003; Tanaka and Grossman, 2003) and sensitizes cells to cytotoxic agents (Tanaka and Grossman, 2003).

The class I family of PI3Ks clearly plays an important role in the regulation of multiple signal transduction pathways that promote cell survival and cell proliferation, and activation of their lipid kinase activity contributes significantly to the development of human malignancies. Furthermore, inhibition of PI3K may potentially circumvent the cellular mechanisms that underlie resistance to chemotherapeutic agents. A potent inhibitor of Class I PI3K activities would therefore have the potential not only to inhibit tumor growth but to also sensitize tumor cells to pro-apoptotic stimuli *in vivo*.

Signal transduction pathways originating from chemoattractant receptors are considered to be important targets in controlling leukocyte motility in inflammatory diseases. Leukocyte trafficking is controlled by chemoattractant factors that activate heterotrimeric GPCRs and thereby trigger a variety of downstream intracellular events. Signal transduction along one of these pathways that results in mobilization of free Ca^{2+} , cytoskeletal reorganization, and directional movement depends on lipid-derived second messengers produced by PI3K activity (Wymann *et al.*, 2000; Stein and Waterfield, 2000).

PI3K γ modulates baseline cAMP levels and controls contractility in cells. Recent research indicates that alterations in baseline cAMP levels contribute to the increased contractility in mutant mice. This research, therefore, shows that PI3K γ inhibitors afford potential treatments for congestive heart failure, ischemia, pulmonary hypertension, renal failure, cardiac hypertrophy, atherosclerosis, thromboembolism, and diabetes.

PI3K inhibitors would be expected to block signal transduction from GPCRs and block the activation of various immune cells, leading to a broad anti-inflammatory profile with potential for the treatment of

inflammatory and immunoregulatory diseases, including asthma, atopic dermatitis, rhinitis, allergic diseases, chronic obstructive pulmonary disease (COPD), septic shock, joint diseases, autoimmune pathologies such as rheumatoid arthritis and Graves' disease, diabetes, cancer, myocardial contractility disorders, thromboembolism, and atherosclerosis.

Breast cancer is a world health problem, and in the United States this disease is the second most common cause of cancer death in women. About 1 in 8 women in the United States (12%) will develop invasive breast cancer over the course of her lifetime. In 2010, an estimated 207,090 new cases of invasive breast cancer were expected to be diagnosed, along with 54,010 new cases of non-invasive breast cancer. About 39,840 women were expected to die in 2010 from breast cancer. The classification and treatment options are usually based on the receptor status. The three most important in the present classification are estrogen receptor (ER), progesterone receptor (PR), and HER2/neu. Cells with or without these receptors are called ER positive (ER+), ER negative (ER-), PR positive (PR+), PR negative (PR-), HER2 positive (HER2+), and HER2 negative (HER2-). Cells with none of these receptors are called basal-like or triple negative. Recently, DNA-based classification is also used in the clinic. As specific DNA mutations or gene expression profiles are identified in the cancer cells, this classification may guide the selection of treatments, either by targeting these changes, or by predicting from the DNA profile which non-targeted therapies are most effective.

The PI3K/PTEN/AKT pathway has been found to be frequently activated and/or mutated in human breast cancer, which contributes to the development and progression of breast cancer, as well as drug resistance. As genetic alterations of *PIK3CA* and *PTEN*, as well as PI3K pathway activation are observed in almost all breast cancer subtypes,

such as HER2 positive, hormone receptor positive, or triple negative breast cancers, it is important to define the strategy for the development of PI3K pathway inhibitors in breast cancer. The present invention is thus to identify molecular markers predicting the sensitivity
5 and/or resistance of the cancer patients toward the PI3K inhibitors described herein. Furthermore, the present invention also relates to the identification of resistance mechanisms and therefore provides a rationale-based synergistic combination to overcome the resistance.

10 To the Applicant's knowledge, no specific disclosure in the prior art is known that 2,3-dihydroimidazo[1,2-c]quinazoline compounds would be effective in the treatment or prophylaxis of inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

15 It has been found, and this is the basis of the present invention, that 2,3-dihydroimidazo[1,2-c]quinazoline compounds, as described and defined herein, show a beneficial effect in the treatment or prophylaxis of breast cancer, in particular inflammatory breast cancer, triple
20 negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

Thus, in accordance with a first aspect, the present invention relates to the use of 2,3-dihydroimidazo[1,2-c]quinazoline compounds, or a
25 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, as a sole active agent, or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in
30 particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In accordance with a second aspect, the present invention relates to combinations of :

5 a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative,
10 antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- 15 - a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- 20 - a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

30

In accordance with a third aspect, the present invention relates to pharmaceutical compositions comprising a 2,3-dihydroimidazo[1,2-

c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, as a sole active agent, for the treatment of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive
5 breast cancer, hormone receptor positive breast cancer.

In accordance with a fourth aspect, the present invention relates to pharmaceutical compositions comprising a combination of :

10 a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative,
15 antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

20 - a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;

25 - a Bcl binding peptide ;

- a Bcl siRNA, such as PNT-2258 ;

- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001,
30 Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014,

AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

In accordance with a fifth aspect, the present invention relates to the
5 use of combinations of :

a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;
or of a pharmaceutical composition containing such a compound or a
10 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

and

15 b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from
20 the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL
25 (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- 30 - an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779,

Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

5 for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

10 In accordance with a sixth aspect, the present invention relates to use of biomarkers involved in the modification of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN for predicting the sensitivity and/or resistance of a patient with cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative
15 breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer, to a 2,3-dihydroimidazo[1,2-c]quinazoline compound as defined herein, thus providing a rationale-based synergistic combination as defined herein to overcome the resistance (patient stratification).

20

In accordance with a seventh aspect, the present invention relates to a method of determining the level of a component of one or more of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN, wherein :

- 25
- in said Bcl expression, said component is Bcl, for example,
 - in said HER family expression and/or activation, PIK3CA signaling, said component is EGF-R, for example, and
 - in said loss of PTEN, said component is PTEN, for example.

30 In accordance a particular embodiment of any of the above aspects of the present invention, said breast cancer is inflammatory breast cancer.


In accordance a particular embodiment of any of the above aspects of the present invention, said breast cancer is triple negative breast cancer.

- 5 In accordance a particular embodiment of any of the above aspects of the present invention, said breast cancer is Her2 receptor positive breast cancer.

- 10 In accordance a particular embodiment of any of the above aspects of the present invention, said breast cancer is hormone receptor positive breast cancer.

Detailed description of the Invention

- 15 A first aspect of the present invention relates to the use of a compound of general formula (A) :



(A)

- 20 in which :

X represents CR⁵R⁶ or NH;

Y¹ represents CR³ or N;

- the chemical bond between Y²-----Y³ represents a single bond or double bond,

with the proviso that when the Y²-----Y³ represents a double bond, Y² and Y³ independently represent CR⁴ or N, and

when $Y^2 \text{-----} Y^3$ represents a single bond, Y^2 and Y^3 independently represent CR^3R^4 or NR^4 ;

Z^1 , Z^2 , Z^3 and Z^4 independently represent CH, CR^2 or N;

5

R^1 represents aryl optionally having 1 to 3 substituents selected from R^{11} , C_{3-8} cycloalkyl optionally having 1 to 3 substituents selected from R^{11} ,

10

C_{1-6} alkyl optionally substituted by aryl, heteroaryl, C_{1-6} alkoxyaryl, aryloxy, heteroaryloxy or one or more halogen, C_{1-6} alkoxy optionally substituted by carboxy, aryl, heteroaryl, C_{1-6} alkoxyaryl, aryloxy, heteroaryloxy or one or more halogen,

or

15

a 3 to 15 membered mono- or bi-cyclic heterocyclic ring that is saturated or unsaturated, optionally having 1 to 3 substituents selected from R^{11} , and contains 1 to 3 heteroatoms selected from the group consisting of N, O and S,

20

wherein

25

R^{11} represents halogen, nitro, hydroxy, cyano, carboxy, amino, N-(C_{1-6} alkyl)amino, N-(hydroxy C_{1-6} alkyl)amino, N,N-di(C_{1-6} alkyl)amino, N-(C_{1-6} acyl)amino, N-(formyl)-N-(C_{1-6} alkyl)amino, N-(C_{1-6} alkanesulfonyl) amino, N-(carboxy C_{1-6} alkyl)-N-(C_{1-6} alkyl)amino, N-(C_{1-6} alkoxycarbonyl)amino,

30

N-[N,N-di(C_{1-6} alkyl)amino methylene]amino, N-[N,N-di(C_{1-6} alkyl)amino (C_{1-6} alkyl)methylene]amino, N-[N,N-di(C_{1-6} alkyl)amino C_{2-6} alkenyl]amino, aminocarbonyl, N-(C_{1-6} alkyl)aminocarbonyl, N,N-di(C_{1-6} alkyl)aminocarbonyl, C_{3-8} cycloalkyl, C_{1-6} alkylthio, C_{1-6} alkanesulfonyl, sulfamoyl, C_{1-6} alkoxycarbonyl,

5 N-arylamino wherein said aryl moiety is optionally having 1 to 3 substituents selected from R¹⁰¹, N-(aryl C₁₋₆alkyl)amino wherein said aryl moiety is optionally having 1 to 3 substituents selected from R¹⁰¹, aryl C₁₋₆alkoxycarbonyl wherein said aryl moiety is optionally having 1 to 3 substituents selected from R¹⁰¹,

10 C₁₋₆alkyl optionally substituted by mono-, di- or tri-halogen, amino, N-(C₁₋₆alkyl)amino or N,N-di(C₁₋₆alkyl)amino,

C₁₋₆alkoxy optionally substituted by mono-, di- or tri-halogen, N-(C₁₋₆alkyl)sulfonamide, or N-(aryl)sulfonamide, or

15 a 5 to 7 membered saturated or unsaturated ring having 1 to 3 heteroatoms selected from the group consisting of O, S and N, and optionally having 1 to 3 substituents selected from R¹⁰¹

20 wherein

R¹⁰¹ represents halogen, carboxy, amino, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, aminocarbonyl, N-(C₁₋₆alkyl)aminocarbonyl, N,N-di(C₁₋₆alkyl)aminocarbonyl, pyridyl,

25

C₁₋₆ alkyl optionally substituted by cyano or mono-di- or tri- halogen, and

30 C₁₋₆alkoxy optionally substituted by cyano, carboxy, amino, N-(C₁₋₆ alkyl)amino, N,N-di(C₁₋₆alkyl)amino, aminocarbonyl, N-(C₁₋₆alkyl)aminocarbonyl, N,N-

di(C₁₋₆alkyl)aminocarbonyl or mono-, di- or tri-halogen;

5 R² represents hydroxy, halogen, nitro, cyano, amino, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, N-(hydroxyC₁₋₆alkyl)amino, N-(hydroxyC₁₋₆alkyl)-N-(C₁₋₆alkyl)amino, C₁₋₆acyloxy, aminoC₁₋₆acyloxy, C₂₋₆alkenyl, aryl,

10 a 5-7 membered saturated or unsaturated heterocyclic ring having 1 to 3 heteroatoms selected from the group consisting O, S and N, and optionally substituted by

15 hydroxy, C₁₋₆alkyl, C₁₋₆alkoxy, oxo, amino, amino C₁₋₆alkyl, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, N-(C₁₋₆acyl)amino, N-(C₁₋₆alkyl)carbonylamino, phenyl, phenyl C₁₋₆alkyl, carboxy, C₁₋₆alkoxycarbonyl, aminocarbonyl, N-(C₁₋₆alkyl)aminocarbonyl, or N,N-di(C₁₋₆alkyl)amino, -C(O)-R²⁰ wherein

20 R²⁰ represents C₁₋₆alkyl, C₁₋₆alkoxy, amino, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, N-(C₁₋₆acyl)amino, or a 5-7 membered saturated or unsaturated heterocyclic ring having 1 to 3 heteroatoms selected from the group consisting O, S and N, and optionally substituted by C₁₋₆alkyl, C₁₋₆alkoxy, oxo, amino, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, N-(C₁₋₆acyl)amino, phenyl, or benzyl,

25

C₁₋₆alkyl optionally substituted by R²¹,
or

30 C₁₋₆alkoxy optionally substituted by R²¹,
wherein

5 R²¹ represents cyano, mono-, di or tri- halogen, hydroxy, amino, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, N- (hydroxyC₁₋₆ alkyl) amino, N- (halophenylC₁₋₆ alkyl) amino, amino C₂₋₆ alkylenyl, C₁₋₆ alkoxy, hydroxyC₁₋₆ alkoxy, - C(O)- R²⁰¹, -NHC(O)- R²⁰¹, C₃₋₈cycloalkyl, isoindolino, phthalimidyl, 2-oxo-1,3-oxazolidinyl, aryl or a 5 or 6 membered saturated or unsaturated heterocyclic ring
 10 having 1 to 4 heteroatoms selected from the group consisting O, S and N , and optionally substituted by hydroxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxycarbonyl, hydroxyC₁₋₆ alkoxy, oxo, amino, aminoC₁₋₆alkyl, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, N-(C₁₋₆acyl)amino, or benzyl,
 15

wherein

20 R²⁰¹ represents hydroxy, amino, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, N- (halophenylC₁₋₆ alkyl) amino, C₁₋₆alkyl, aminoC₁₋₆ alkyl, aminoC₂₋₆ alkylenyl, C₁₋₆ alkoxy, a 5 or 6 membered saturated or unsaturated heterocyclic ring having 1
 25 to 4 heteroatoms selected from the group consisting O, S and N, and optionally substituted by hydroxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxycarbonyl, hydroxyC₁₋₆ alkoxy, oxo, amino, N-(C₁₋₆alkyl)amino, N,N-di(C₁₋₆alkyl)amino, N-(C₁₋₆acyl)amino or benzyl;
 30

R³ represents hydrogen, halogen, aminocarbonyl, or C₁₋₆ alkyl optionally substituted by aryl C₁₋₆ alkoxy or mono-, di- or tri- halogen;

5

R⁴ represents hydrogen or C₁₋₆ alkyl;

R⁵ represents hydrogen or C₁₋₆ alkyl; and

10 R⁶ represents halogen, hydrogen or C₁₋₆ alkyl,
or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
as a sole active agent,
or of combinations of :

15 a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a
physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent
selected from an anti-angiogenesis, anti-hyper-proliferative,
20 antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic,
anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral
agent, more particularly one or more further active agents selected from
the group consisting of :

25 - a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25,
YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;

30 - a Bcl binding peptide ;

- a Bcl siRNA, such as PNT-2258 ;

- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of
5 mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a
10 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations,

for the preparation of a medicament for the treatment or prophylaxis of
15 cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In a particular embodiment of the above-mentioned first aspect, the
20 present invention relates to the use of a compound selected from the following list,

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

25 or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent
30 selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral

agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- 10 - a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel),
- 15 Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;
- or of pharmaceutical compositions containing such compounds or a
- 20 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
- or of pharmaceutical compositions containing such combinations

for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer :


N-(7,8-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;

- 2-(7, 8-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1-pyridin-3-ylethylenol;
 N-(7, 8-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1H-benzimidazole-5-carboxamide;
- 5 6-(acetamido)-N-(7,8-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
 N-{5-[2-(7,8-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1-hydroxyvinyl]pyridin-2-yl}acetamide;
- 10 2-({5-[2-hydroxy-2-pyridin-3-ylvinyl]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-8-yl}oxy)-N,N-dimethylacetamide;
 2-[7-methoxy-8-(tetrahydro-2H-pyran-2-ylmethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1-pyridin-3-ylethylenol;
 2-[8-(2-hydroxyethoxy)-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1-pyridin-3-ylethylenol;
- 15 ({5-[2-hydroxy-2-pyridin-3-ylvinyl]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-8-yl}oxy)acetic acid;
 4-({5-[2-hydroxy-2-pyridin-3-ylvinyl]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-8-yl}oxy)butanoic acid;
 ({5-[2-hydroxy-2-pyridin-3-ylvinyl]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-8-yl}oxy)acetonitrile;
- 20 2-[7-methoxy-8-(2H-tetrazol-5-ylmethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1-pyridin-3-ylethylenol;
 2-[7-methoxy-8-(4-morpholin-4-yl-4-oxobutoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1-pyridin-3-ylethylenol;
- 25 5-[1-hydroxy-2-(8-morpholin-4-yl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)vinyl]pyridin-3-ol ;
 N-(2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-5-hydroxynicotinamide;
 6-(acetamido)-N-(7,9-dimethoxy-8-methyl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- 30 N-(8,9-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-5-hydroxynicotinamide;

- 5-hydroxy-N-(7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- N-(7,8-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-5-[(4-methoxybenzyl)oxy]nicotinamide;
- 5 N-(7,8-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-5-hydroxynicotinamide;
- 5-hydroxy-N-[8-(trifluoromethyl)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-{8-[3-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)propoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 10 N-(7-bromo-8-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- 6-amino-N-(8-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- 15 1-(1H-benzimidazol-5-yl)-2-(8,9-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)ethylenol;
- 2-(8,9-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1-(2,4-dimethyl-1,3-thiazol-5-yl)ethylenol;
- N-(9-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1H-benzimidazole-5-carboxamide;
- 20 N-(8-bromo-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- N-(8-bromo-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1H-benzimidazole-5-carboxamide;
- N-(8-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1H-benzimidazole-5-carboxamide;
- 25 N-(8-methyl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1H-benzimidazole-5-carboxamide;
- N-[8-(trifluoromethyl)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1H-benzimidazole-5-carboxamide;
- 30 N-(7-fluoro-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1H-benzimidazole-5-carboxamide;
- N-(7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;

- N-(8-chloro-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1H-benzimidazole-5-carboxamide;
- 6-(acetamido)-N-(8-morpholin-4-yl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- 5 1-(1H-benzimidazol-5-yl)-2-(8-morpholin-4-yl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)ethylenol;
- N-{5-[1-hydroxy-2-(8-morpholin-4-yl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)vinyl]pyridin-2-yl}acetamide;
- 6-methyl-N-(8-morpholin-4-yl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- 10 1-(1H-benzimidazol-5-yl)-2-[8-(4-methylpiperazin-1-yl)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]ethylenol;
- N-(2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-3H-imidazo[4,5-b]pyridine-6-carboxamide;
- 15 N-(7,8-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-3H-imidazo[4,5-b]pyridine-6-carboxamide;
- N-[7-(trifluoromethyl)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1H-benzimidazole-5-carboxamide;
- N-(7,9-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1H-benzimidazole-5-carboxamide;
- 20 N-{5-[2-(7,9-dimethoxy-8-methyl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1-hydroxyvinyl]pyridin-2-yl}acetamide;
- N-{5-[2-(7-bromo-9-methyl-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1-hydroxyvinyl]pyridin-2-yl}acetamide; and
- 25 2-(8,9-dimethoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-1-pyridin-3-ylethylenol;

Another embodiment of the present invention encompasses the use of a compound having the formula (I) :



(I)

5 or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, in which :

R¹ represents $-(CH_2)_n-(CHR^4)-(CH_2)_m-N(R^5)(R^{5'})$;

R² represents a heteroaryl optionally substituted with 1, 2 or 3 R⁶
10 groups ;

R³ represents alkyl or cycloalkyl ;

R⁴ represents hydrogen, hydroxy or alkoxy ; and

R⁵ and R^{5'} may be the same or different and represent independently,
hydrogen, alkyl, cycloalkylalkyl, or alkoxyalkyl or R⁵ and R^{5'} may
15 be taken together with the nitrogen atom to which they are bound
to form a 3-7 membered nitrogen containing heterocyclic ring
optionally containing at least one additional heteroatom selected
from oxygen, nitrogen or sulfur and which may be optionally
substituted with 1 or more R^{6'} groups, or R⁴ and R⁵ may be taken
20 together with the atoms to which they are bound to form a 5-6
membered nitrogen containing heterocyclic ring optionally
containing 1 or more nitrogen, oxygen or sulfur atoms and which
may be optionally substituted with 1 or more R^{6'} groups ;

each occurrence of R^6 may be the same or different and is independently halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclic ring, heterocyclalkyl, alkyl-OR⁷, alkyl-SR⁷, alkyl-N(R⁷)(R^{7'}), alkyl-COR⁷, -CN, -
5 COOR⁷, -CON(R⁷)(R^{7'}), -OR⁷, -SR⁷, -N(R⁷)(R^{7'}), or -NR⁷COR⁷ each of which may be optionally substituted with 1 or more R⁸ groups ;

each occurrence of $R^{6'}$ may be the same or different and is independently alkyl, cycloalkylalkyl, or alkyl-OR⁷;

10

each occurrence of R^7 and $R^{7'}$ may be the same or different and is independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heterocyclic ring, heterocyclalkyl, or heteroarylalkyl ;

15

each occurrence of R^8 is independently nitro, hydroxy, cyano, formyl, acetyl, halogen, amino, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heterocyclic ring, heterocyclalkyl, or heteroarylalkyl ;

20

n is an integer from 1-4 and m is an integer from 0-4 with the proviso that when R^4 and R^5 are taken together with the atoms to which they are bound to form a 5-6 membered nitrogen containing ring, $n + m \leq 4$;

25

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

or of combinations of :

30

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In a preferred embodiment, the invention encompasses the use of a compound of Formula (I), wherein R^2 is a nitrogen containing heteroaryl optionally substituted with 1, 2 or 3 R^6 groups,

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from a n anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;

- a Bcl binding peptide ;

- a Bcl siRNA, such as PNT-2258 ;

- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;or of pharmaceutical

compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
or of pharmaceutical compositions containing such combinations,
for the preparation of a medicament for the treatment or prophylaxis of
5 cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In another preferred embodiment, the invention encompasses the use of
10 a compound of Formula (I), wherein R^5 and $R^{5'}$ are independently alkyl, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
as a sole active agent,
or of combinations of :

15 a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative,
20 antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- 25 - a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- 30 - a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of
5 mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a
10 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations,

for the preparation of a medicament for the treatment or prophylaxis of
15 cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In still another preferred embodiment, the invention encompasses the
20 use of a compound of Formula (I), wherein R⁵ and R^{5'} are taken together with the nitrogen atom to which they are bound to form a 5-6 membered nitrogen containing heterocyclic ring containing at least one additional heteroatom selected from oxygen, nitrogen or sulfur and which may be optionally substituted with 1 or more R^{6'} groups,

25 or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a
30 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In yet another preferred embodiment, the invention encompasses the use of a compound of Formula (I), wherein R⁴ is hydroxyl, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

5 as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

10 b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from
15 the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL
20 (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- 25 - an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014,
30 AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations,

5

for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

10

In another preferred embodiment, the invention encompasses the use of a compound of Formula (I), wherein R^4 and R^5 are taken together with the atoms to which they are bound to form a 5-6 membered nitrogen containing heterocyclic ring optionally containing 1 or more nitrogen, oxygen or sulfur atoms and which may be optionally substituted with 1

15

or more R^6 groups,

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

20

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

25

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

30

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-

-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
 - 5 - a Bcl siRNA, such as PNT-2258 ;
 - an antisense therapy oligonucleotide, such as BclKlex ; and
 - an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779,
 - 10 Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;
- 15 or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
- or of pharmaceutical compositions containing such combinations,
- for the preparation of a medicament for the treatment or prophylaxis of
- 20 cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.
- In yet another preferred embodiment, the invention encompasses the
- 25 use of a compound of Formula (I), wherein R³ is methyl,
- or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
- as a sole active agent,
- or of combinations of :
- 30 a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In still another preferred embodiment, the invention encompasses the use of a compound of Formula (I), wherein R² is pyridine, pyridazine, pyrimidine, pyrazine, pyrrole, oxazole, thiazole, furan or thiophene, optionally substituted with 1, 2 or 3 R⁶ groups; more preferably pyridine,
5 pyridazine, pyrimidine, pyrazine, pyrrole, oxazole or thiazole, optionally substituted with 1, 2 or 3 R⁶ groups,

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

10 or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent
15 selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

20

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-
25 3139 (Genasense), Evotec, PIB-1402, EU-517 ;

- a Bcl binding peptide ;

- a Bcl siRNA, such as PNT-2258 ;


- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin
30 analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR

kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;or of pharmaceutical compositions containing such compounds or a physiologically acceptable
 5 salt, solvate, hydrate or stereoisomer thereof,
 or of pharmaceutical compositions containing such combinations,

for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer,
 10 triple negative breast cancer, Her2 receptor positive breast cancer,
 hormone receptor positive breast cancer.

In a distinct embodiment, the invention encompasses the use of a compound of formula (Ia)



15

(Ia)


or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, wherein R^2 is as defined above,
 20 or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
 as a sole active agent,
 or of combinations of :
 25 a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ; or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
- or of pharmaceutical compositions containing such combinations, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

30

In another distinct embodiment, the invention encompasses the use of a compound of formula (Ib) :



(Ib)

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, wherein R^2 is as defined above,

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;

- a Bcl binding peptide ;

- a Bcl siRNA, such as PNT-2258 ;


- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of
5 mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a
10 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations,
for the preparation of a medicament for the treatment or prophylaxis of
cancer, e.g. breast cancer, in particular inflammatory breast cancer,
15 triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In still another distinct embodiment, the invention encompasses the use of a compound of formula (Ic) :



20

(Ic)

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, wherein R² is as defined above,
25 or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

5 b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from
10 the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL
15 (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- 20 - an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014,
25 AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;


or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer
30 thereof,

or of pharmaceutical compositions containing such combinations,

for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

5

In yet another distinct embodiment, the invention encompasses the use of a compound of the formula (Id):



10

(Id)

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, wherein R^2 and R^4 are as defined above,

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

15

as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

20

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from

25


the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- 10 - an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, 15 AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

20 or of pharmaceutical compositions containing such combinations, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, 25 hormone receptor positive breast cancer.

In yet another distinct embodiment, the invention encompasses the use of a compound of the formula (Ie) :



(le)

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, wherein R² and R⁴ are as defined above,
 or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
 as a sole active agent,
 or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;

- an antisense therapy oligonucleotide, such as BclKlex ; and
 - an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;
- 10 or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
- or of pharmaceutical compositions containing such combinations,
- for the preparation of a medicament for the treatment or prophylaxis of
- 15 cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In a preferred embodiment, the invention encompasses the use of a

20 compound of formula (I) - (V), wherein R^2 is pyridine, pyridazine, pyrimidine, pyrazine, pyrole, oxazole, thiazole, furan or thiophene, optionally substituted with 1, 2 or 3 R^6 groups; more preferably wherein R^2 is pyridine, pyridazine, pyrimidine, pyrazine, pyrole, oxazole or thiazole, optionally substituted with 1, 2 or 3 R^6 groups,

25 or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a

30 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ; or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
- or of pharmaceutical compositions containing such combinations, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

30

In still another preferred embodiment, the invention encompasses the use of a compound having the formula :

- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;
- 5 N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide ;
- N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-2,4-dimethyl-1,3-thiazole-5-carboxamide;
- 10 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1,3-thiazole-5-carboxamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]isonicotinamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-4-methyl-1,3-thiazole-5-
- 15 carboxamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-4-propylpyrimidine-5-carboxamide;
- 20 N-{8-[2-(4-ethylmorpholin-2-yl)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}pyrimidine-5-carboxamide;
- 25 N-(8-{3-[2-(hydroxymethyl)morpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- N-(8-{3-[2-(hydroxymethyl)morpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide 1-oxide;
- 30 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-(2-pyrrolidin-1-ylethyl)nicotinamide;
6-(cyclopentylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
5 N-[8-(2-hydroxy-3-morpholin-4-ylpropoxy)-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
N-{7-methoxy-8-[3-(3-methylmorpholin-4-yl)propoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
10 N-(8-{3-[2-(hydroxymethyl)morpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
N-(8-{2-[4-(cyclobutylmethyl)morpholin-2-yl]ethoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
N-(7-methoxy-8-{2-[4-(2-methoxyethyl)morpholin-2-yl]ethoxy}-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
15 N-{8-[(4-ethylmorpholin-2-yl)methoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
N-(7-methoxy-8-{[4-(2-methoxyethyl)morpholin-2-yl]methoxy}-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
20 N-{7-methoxy-8-[(4-methylmorpholin-2-yl)methoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-4-carboxamide;
2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-4-carboxamide;
25 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1-methyl-1H-imidazole-4-carboxamide;
rel-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)pyrimidine-5-carboxamide;
30

rel-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-6-methylnicotinamide;

5 rel-6-acetamido-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1-methyl-1H-imidazole-5-carboxamide;

10 6-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-methylnicotinamide;
2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-4-methylpyrimidine-5-carboxamide;

15 6-amino-5-bromo-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1,3-oxazole-5-carboxamide;
N-[7-methoxy-8-(morpholin-2-ylmethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;

20 2-{[2-(dimethylamino)ethyl]amino}-N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}pyrimidine-5-carboxamide;
2-amino-N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}-1,3-thiazole-5-carboxamide;
25 rel-2-amino-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)pyrimidine-5-carboxamide;

rel-6-amino-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;

30

2-[(2-hydroxyethyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

5 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-[(3-methoxypropyl)amino]pyrimidine-5-carboxamide;

2-amino-N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}pyrimidine-5-carboxamide;

10 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-[(3-morpholin-4-ylpropyl)amino]pyrimidine-5-carboxamide;

2-[(2-methoxyethyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

15 2-{[2-(dimethylamino)ethyl]amino}-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

6-amino-N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;

20 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-pyrrolidin-1-ylpyrimidine-5-carboxamide;

25 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-(4-methylpiperazin-1-yl)pyrimidine-5-carboxamide;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-morpholin-4-ylpyrimidine-5-carboxamide;

30 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-piperazin-1-ylnicotinamide hydrochloride;

- 6-[(3S)-3-aminopyrrolidin-1-yl]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide hydrochloride hydrate;
- 5 6-[(3R)-3-aminopyrrolidin-1-yl]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide hydrochloride;
- 6-[(4-fluorobenzyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 10 6-[(2-furylmethyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 6-[(2-methoxyethyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-(1H-pyrrol-1-yl)nicotinamide;
- 15 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-morpholin-4-ylnicotinamide;
- N-{7-methoxy-8-[3-(methylamino)propoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 20 6-[(2,2-dimethylpropanoyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 6-[(cyclopropylcarbonyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide
- 25 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-(2,2,2-trifluoroethoxy)nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-
- 30 (trifluoromethyl)nicotinamide;

- 6-(isobutyrylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-{7-methoxy-8-[3-(4-methylpiperazin-1-yl)propoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 5 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-
{[(methylamino)carbonyl]amino}-1,3-thiazole-4-carboxamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-
10 {[(methylamino)carbonyl]amino}nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-(methylamino)-1,3-thiazole-4-carboxamide;
- N-[7-methoxy-8-(2-morpholin-4-ylethoxy)-2,3-dihydroimidazo[1,2-
15 c]quinazolin-5-yl]nicotinamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}-2,4-dimethyl-1,3-thiazole-5-carboxamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-
20 dihydroimidazo[1,2-c]quinazolin-5-yl}-6-methylnicotinamide;
- 6-{[(isopropylamino)carbonyl]amino}-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-
25 dihydroimidazo[1,2-c]quinazolin-5-yl]-6-pyrrolidin-1-ylnicotinamide;
- 6-(dimethylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-piperidin-1-ylpropoxy)-2,3-dihydroimidazo[1,2-
30 c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(2-pyrrolidin-1-ylethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;

- N-[7-methoxy-8-(2-piperidin-1-ylethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 6-{[(ethylamino)carbonyl]amino}-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 5 6-fluoro-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1,3-oxazole-4-carboxamide;
- 2-(ethylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1,3-thiazole-4-carboxamide;
- 10 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrazine-2-carboxamide;
- N-[8-(2-aminoethoxy)-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 15 6-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]isonicotinamide;
- N-{8-[3-(diethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 20 N-{8-[2-(diisopropylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- N-{8-[2-(diethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 25 N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 30 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-(methylamino)pyrimidine-5-carboxamide;

- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-(methylthio)pyrimidine-5-carboxamide;
- 5 N-[8-(3-aminopropoxy)-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide trifluoroacetate;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]thiophene-2-carboxamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2,4-dimethyl-1,3-thiazole-5-
- 10 carboxamide;
- 2-methoxy-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-3-furamide;
- 15 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]thiophene-3-carboxamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-methyl-1,3-thiazole-4-carboxamide;
- 20 6-methoxy-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 5-methoxy-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-methylnicotinamide;
- 25 6-(acetylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 30 or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,
- as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

5 b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from
10 the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL
15 (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- 20 - an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014,
25 AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a physiologically acceptable salt, solvate, hydrate or stereoisomer
30 thereof,

or of pharmaceutical compositions containing such combinations,

for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

5

In a preferred embodiment, the invention encompasses the use of a compound having the formula:

- 10 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-methylnicotinamide;
- 5-methoxy-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 15 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2,4-dimethyl-1,3-thiazole-5-carboxamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 20 N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 6-{[(isopropylamino)carbonyl]amino}-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}-2,4-dimethyl-1,3-thiazole-5-carboxamide;
- 25 N-[7-methoxy-8-(2-morpholin-4-ylethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- rel-6-amino-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- 30

rel-2-amino-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)pyrimidine-5-carboxamide;

2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;
 N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}pyrimidine-5-carboxamide;
 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-

c]quinazolin-5-yl]pyrimidine-5-carboxamide;

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

as a sole active agent,

or of combinations of :

a) such a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;

- a Bcl binding peptide ;

- a Bcl siRNA, such as PNT-2258 ;

- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of
5 mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such compounds or a
10 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations,
for the preparation of a medicament for the treatment or prophylaxis of
cancer, e.g. breast cancer, in particular inflammatory breast cancer,
15 triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In a preferred embodiment, the invention encompasses the use of a
compound having the formula :

20

2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-
dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a
physiologically acceptable salt, solvate, hydrate or stereoisomer
thereof ;

25

as a sole active agent,

or of pharmaceutical compositions containing such a compound or a
physiologically acceptable salt, solvate, hydrate or stereoisomer
30 thereof,

for the preparation of a medicament for the treatment or prophylaxis of
cancer, e.g. breast cancer, in particular inflammatory breast cancer,

triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In a preferred embodiment, the invention encompasses the use of combinations of :

- a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and
- b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :
- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
 - a Bcl binding peptide ;
 - a Bcl siRNA, such as PNT-2258 ;
 - an antisense therapy oligonucleotide, such as BclKlex ; and
 - an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014,

AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such a compound or a
5 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations,
for the preparation of a medicament for the treatment or prophylaxis of
cancer, e.g. breast cancer, in particular inflammatory breast cancer,
10 triple negative breast cancer, Her2 receptor positive breast cancer,
hormone receptor positive breast cancer.

In a preferred embodiment, the invention encompasses the use of
combinations of :

15

a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-
dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a
physiologically acceptable salt, solvate, hydrate or stereoisomer
thereof ; and

20

b) one or more further active agents selected from the group consisting
of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25,
25 YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- 30 - a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of
5 mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such a compound or a
10 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations,
for the preparation of a medicament for the treatment or prophylaxis of
cancer, e.g. breast cancer, in particular inflammatory breast cancer,
15 triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In a preferred embodiment, the invention encompasses the use of combinations of :

20

a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

25

b) one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25,
30 YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL

(Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;

- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- 5 - an antisense therapy oligonucleotide, such as BclKlex ;

or of pharmaceutical compositions containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

- 10 or of pharmaceutical compositions containing such combinations, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

15

In a preferred embodiment, the invention encompasses the use of combinations of :

- a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a
20 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

- b) one or more further active agents, which is ABT-737 ;

25

or of pharmaceutical compositions containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

- or of pharmaceutical compositions containing such combinations,
30 for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer,

triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In a preferred embodiment, the invention encompasses the use of combinations of :

a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents selected from the group consisting of :

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or of pharmaceutical compositions containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations, for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In a preferred embodiment, the invention encompasses the use of combinations of :

5 a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and

b) one or more further active agents, which is Rapamycin (Sirolimus) ;

10

or of pharmaceutical compositions containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

or of pharmaceutical compositions containing such combinations,

15

for the preparation of a medicament for the treatment or prophylaxis of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

20

Where there is a discrepancy between the chemical name and the chemical structure depicted, the chemical structure depicted takes precedence over the chemical name given.

25 Without being bound by theory or mechanism, the compounds of the present invention display surprising activity for the inhibition of phosphatidylinositol-3-kinase and chemical and structural stability over those compounds of the prior art. It is believed that this surprising activity is based on the chemical structure of the compounds, in particular the basicity of the compounds as a result of R¹ being amino
30 optionally substituted with R⁵ and R^{5'}. Further, the appropriate choice

of R^3 and R^2 provide the necessary activity against the appropriate isoforms to allow for activity in vivo.

5 In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is inflammatory breast cancer.

10 In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is triple negative breast cancer.

In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is Her2 receptor positive breast cancer.

15 In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is hormone receptor positive breast cancer.

20

Definitions

25 The term 'alkyl' refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing solely of carbon and hydrogen atoms, containing no unsaturation, having from one to eight carbon atoms, and which is attached to the rest of the molecule by a single bond, such as illustratively, methyl, ethyl, n-propyl 1-methylethyl (isopropyl), n-butyl, n-pentyl, and 1,1-dimethylethyl (t-butyl).

30

The term "alkenyl" refers to an aliphatic hydrocarbon group containing a carbon-carbon double bond and which may be a straight or branched or

branched chain having about 2 to about 10 carbon atoms, e.g., ethenyl, 1-propenyl, 2-propenyl (allyl), iso-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-and butenyl.

- 5 The term "alkynyl" refers to a straight or branched chain hydrocarbonyl radicals having at least one carbon-carbon triple bond, and having in the range of about 2 up to 12 carbon atoms (with radicals having in the range of about 2 up to 10 carbon atoms presently being preferred) e.g., ethynyl.

10

The term "alkoxy" denotes an alkyl group as defined herein attached via oxygen linkage to the rest of the molecule. Representative examples of those groups are methoxy and ethoxy.

- 15 The term "alkoxyakyl" denotes an alkoxy group as defined herein attached via oxygen linkage to an alkyl group which is then attached to the main structure at any carbon from alkyl group that results in the creation of a stable structure the rest of the molecule. Representative examples of those groups are $-\text{CH}_2\text{OCH}_3$, $-\text{CH}_2\text{OC}_2\text{H}_5$.

20

The term "cycloalkyl" denotes a non-aromatic mono or multicyclic ring system of about 3 to 12 carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and examples of multicyclic cycloalkyl groups include perhydronaphthyl, adamantyl and norbornyl groups bridged
25 cyclic group or spirobicyclic groups e.g spiro (4,4) non-2-yl.

- The term "cycloalkylalkyl" refers to cyclic ring-containing radicals containing in the range of about about 3 up to 8 carbon atoms directly attached to alkyl group which is then also attached to the main structure
30 at any carbon from the alkyl group that results in the creation of a stable structure such as cyclopropylmethyl, cyclobuylethyl, cyclopentylethyl.

The term "aryl" refers to aromatic radicals having in the range of 6 up to 14 carbon atoms such as phenyl, naphthyl, tetrahydronaphthyl, indanyl, biphenyl .

5

The term "arylalkyl" refers to an aryl group as defined herein directly bonded to an alkyl group as defined herein which is then attached to the main structure at any carbon from alkyl group that results in the creation of a stable structure the rest of the molecule. e.g., --CH₂C₆H₅, -C₂H₅C₆H₅ .

10

The term "heterocyclic ring" refers to a stable 3- to 15 membered ring radical which consists of carbon atoms and from one to five heteroatoms selected from the group consisting of nitrogen, phosphorus, oxygen and sulfur. For purposes of this invention, the heterocyclic ring radical may be a monocyclic, bicyclic or tricyclic ring system, which may include fused, bridged or spiro ring systems, and the nitrogen, phosphorus, carbon, oxygen or sulfur atoms in the heterocyclic ring radical may be optionally oxidized to various oxidation states. In addition, the nitrogen atom may be optionally quaternized; and the ring radical may be partially or fully saturated (i.e., heteroaromatic or heteroaryl aromatic). Examples of such heterocyclic ring radicals include, but are not limited to, azetidiny, acridiny, benzodioxoly, benzodioxany, benzofurnyl, carbazoly, cinnoliny, dioxolany, indoliziny, naphthyridiny, perhydroazepiny, phenaziny, phenothiaziny, phenoxaziny, phthalazil, pyridyl, pteridiny, puriny, quinazoliny, quinoxaliny, quinoliny, isoquinoliny, tetrazoyl, imidazolyl tetrahydroisouinolyl, piperidiny, piperaziny, 2-oxopiperaziny, 2-oxopiperidiny, 2-oxopyrrolidiny, 2-oxoazepiny, azepiny, pyrroly, 4-piperidony, pyrrolidiny, pyraziny, pyrimidiny, pyridaziny, oxazolyl oxazoliny oxasolidiny, triazolyl, indanyl, isoxazolyl, isoxasolidiny, morpholiny, thiazolyl, thiazoliny, thiazolidiny, isothiazolyl, quinuclidiny, isothiazolidiny, indoly,

15

20

25

30

isoindolyl, indolyl, isoindolyl, octahydroindolyl, octahydroisoindolyl
quinolyl, isoquinolyl, decahydroisoquinolyl, benzimidazolyl, thiadiazolyl,
benzopyranyl, benzothiazolyl, benzooxazolyl, furyl, tetrahydrofurtyl,
tetrahydropyranyl, thienyl, benzothienyl, thiamorpholinyl,
5 thiamorpholinyl sulfoxide thiamorpholinyl sulfone, dioxaphospholanyl,
oxadiazolyl, chromanyl, isochromanyl .

The term "heteroaryl" refers to heterocyclic ring radical as defined
herein which are aromatic. The heteroaryl ring radical may be attached
10 to the main structure at any heteroatom or carbon atom that results in
the creation of a stable structure.

The heterocyclic ring radical may be attached to the main structure at
any heteroatom or carbon atom that results in the creation of a stable
15 structure.

The term "heteroarylalkyl" refers to heteroaryl ring radical as defined
herein directly bonded to alkyl group. The heteroarylalkyl radical may
be attached to the main structure at any carbon atom from alkyl group
20 that results in the creation of a stable structure.

The term "heterocyclyl" refers to a heterocyclic ring radical as defined
herein. The heterocyclyl ring radical may be attached to the main
structure at any heteroatom or carbon atom that results in the creation
25 of a stable structure.

The term "heterocyclylalkyl" refers to a heterocyclic ring radical as
defined herein directly bonded to alkyl group. The heterocyclylalkyl
radical may be attached to the main structure at carbon atom in the
30 alkyl group that results in the creation of a stable structure.

The term "carbonyl" refers to an oxygen atom bound to a carbon atom of the molecule by a double bond.

5 The term "halogen" refers to radicals of fluorine, chlorine, bromine and iodine.

Where the plural form of the word compounds, salts, polymorphs, hydrates, solvates and the like, is used herein, this is taken to mean also a single compound, salt, polymorph, isomer, hydrate, solvate or the like.

10

The compounds of this invention may contain one or more asymmetric centers, depending upon the location and nature of the various substituents desired. Asymmetric carbon atoms may be present in the (*R*) or (*S*) configuration, resulting in racemic mixtures in the case of a
15 single asymmetric center, and diastereomeric mixtures in the case of multiple asymmetric centers. In certain instances, asymmetry may also be present due to restricted rotation about a given bond, for example, the central bond adjoining two substituted aromatic rings of the specified compounds. Substituents on a ring may also be present in
20 either cis or trans form. It is intended that all such configurations (including enantiomers and diastereomers), are included within the scope of the present invention. Preferred compounds are those, which produce the more desirable biological activity. Separated, pure or partially purified isomers and stereoisomers or racemic or
25 diastereomeric mixtures of the compounds of this invention are also included within the scope of the present invention. The purification and the separation of such materials can be accomplished by standard techniques known in the art.

30 The present invention also relates to useful forms of the compounds as disclosed herein, such as pharmaceutically acceptable salts, co-precipitates, metabolites, hydrates, solvates and prodrugs of all the

compounds of examples. The term "pharmaceutically acceptable salt" refers to a relatively non-toxic, inorganic or organic acid addition salt of a compound of the present invention. For example, see S. M. Berge, *et al.* "Pharmaceutical Salts," *J. Pharm. Sci.* 1977, 66, 1-19.

5 Pharmaceutically acceptable salts include those obtained by reacting the main compound, functioning as a base, with an inorganic or organic acid to form a salt, for example, salts of hydrochloric acid, sulfuric acid, phosphoric acid, methane sulfonic acid, camphor sulfonic acid, oxalic acid, maleic acid, succinic acid and citric acid. Pharmaceutically
10 acceptable salts also include those in which the main compound functions as an acid and is reacted with an appropriate base to form, e.g., sodium, potassium, calcium, magnesium, ammonium, and chlorine salts. Those skilled in the art will further recognize that acid addition salts of the claimed compounds may be prepared by reaction of the
15 compounds with the appropriate inorganic or organic acid via any of a number of known methods. Alternatively, alkali and alkaline earth metal salts of acidic compounds of the invention are prepared by reacting the compounds of the invention with the appropriate base via a variety of known methods.

20

Representative salts of the compounds of this invention include the conventional non-toxic salts and the quaternary ammonium salts which are formed, for example, from inorganic or organic acids or bases by means well known in the art. For example, such acid addition salts
25 include acetate, adipate, alginate, ascorbate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cinnamate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, chloride, bromide, iodide, 2-hydroxyethanesulfonate, itaconate, lactate, maleate,
30 mandelate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate,

picrate, pivalate, propionate, succinate, sulfonate, sulfate, tartrate, thiocyanate, tosylate, and undecanoate.

Base salts include alkali metal salts such as potassium and sodium salts, alkaline earth metal salts such as calcium and magnesium salts, and ammonium salts with organic bases such as dicyclohexylamine and N-methyl-D-glucamine. Additionally, basic nitrogen containing groups may be quaternized with such agents as lower alkyl halides such as methyl, ethyl, propyl, or butyl chlorides, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl sulfate, or diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others.

A solvate for the purpose of this invention is a complex of a solvent and a compound of the invention in the solid state. Exemplary solvates would include, but are not limited to, complexes of a compound of the invention with ethanol or methanol. Hydrates are a specific form of solvate wherein the solvent is water.

20

The synthesis of the compounds listed above is described in International Patent Application No. PCT/EP2003/010377, published as WO 2004/029055 A1, and in International Patent Application No. PCT/US2007/024985, published as WO 2008/070150, both of which are hereby incorporated herein in their entirety by reference.

In accordance with another embodiment, the present invention relates to a 2,3-dihydroimidazo[1,2-c]quinazoline compound as defined herein, in particular 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, as a sole agent, for the treatment of cancer, e.g. breast cancer,

30

in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

5 In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is inflammatory breast cancer.

10 In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is triple negative breast cancer.

15 In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is Her2 receptor positive breast cancer.

In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is hormone receptor positive breast cancer.

20

Combination therapies

As mentioned *supra*, the present invention relates to combinations of :

25 a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound as defined *supra*, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or pharmaceutical compositions containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

30

and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

In a preferred embodiment, the invention encompasses combinations of :

- a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or pharmaceutical compositions containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

and

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :

- 10 - a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- 15 - a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor),
- 20 Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

25

In a preferred embodiment, the invention encompasses combinations of :

- a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or pharmaceutical compositions containing such a compound or
- 30

a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

and

5

b) one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, 10 YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- 15 - a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, 20 Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

25 In a preferred embodiment, the invention encompasses combinations of :

- a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a 30 physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or pharmaceutical compositions containing such a compound or

a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

and

5

b) one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, 10 YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- a Bcl binding peptide ;
- 15 - a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex.

In a preferred embodiment, the invention encompasses the use of combinations of :

20

- a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or pharmaceutical compositions containing such a compound or 25 a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

and

30

- b) one or more further active agents, which is ABT-737.

In a preferred embodiment, the invention encompasses combinations of :

- a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or pharmaceutical compositions containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

10

and

- b) one or more further active agents selected from the group consisting of :

15

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

20

In a preferred embodiment, the invention encompasses combinations of :

25

- a) 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or pharmaceutical compositions containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

30

and

b) one or more further active agents, which is Rapamycin (Sirolimus).

5

The compounds of this invention can be administered as the sole pharmaceutical agent or in combination with one or more other pharmaceutical agents (or “further active agents”) where the combination causes no unacceptable adverse effects. For example, the
10 compounds of this invention can be combined with known anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agents, and the like, as well as with admixtures and combinations thereof.

15

The additional pharmaceutical agent or agents (or “further active agent”) can be, but are not limited to aldesleukin, alendronic acid, alfaferone, alitretinoin, allopurinol, aloprim, aloxi, altretamine, aminoglutethimide, amifostine, amrubicin, amsacrine, anastrozole,
20 anzmet, aranesp, arglabin, arsenic trioxide, aromasin, 5-azacytidine, azathioprine, BCG or tice BCG, bestatin, betamethasone acetate, betamethasone sodium phosphate, bexarotene, bleomycin sulfate, broxuridine, bortezomib, busulfan, calcitonin, campath, capecitabine, carboplatin, casodex, cefesone, celmoleukin, cerubidine, chlorambucil, cisplatin, cladribine, cladribine, clodronic acid, cyclophosphamide,
25 cytarabine, dacarbazine, dactinomycin, DaunoXome, decadron, decadron phosphate, delestrogen, denileukin diftitox, depo-medrol, deslorelin, dexomethasone, dexrazoxane, diethylstilbestrol, diflucan, docetaxel, doxifluridine, doxorubicin, dronabinol, DW-166HC, eligard, elitek, ellence, emend, epirubicin, epoetin alfa, epogen, eptaplatin,
30 ergamisol, estrace, estradiol, estramustine phosphate sodium, ethinyl estradiol, ethyol, etidronic acid, etopophos, etoposide, fadrozole,

farston, filgrastim, finasteride, fligrastrim, floxuridine, fluconazole, fludarabine, 5-fluorodeoxyuridine monophosphate, 5-fluorouracil (5-FU), fluoxymesterone, flutamide, formestane, fosteabine, fotemustine, fulvestrant, gammagard, gemcitabine, gemtuzumab, gleevec, gliadel, 5 goserelin, granisetron HCl, herceptin, histrelin, hycamtin, hydrocortone, eyrthro-hydroxynonyladenine, hydroxyurea, ibritumomab tiuxetan, idarubicin, ifosfamide, interferon alpha, interferon-alpha 2, interferon alfa-2A, interferon alfa-2B, interferon alfa-n1, interferon alfa-n3, interferon beta, interferon gamma-1a, interleukin-2, intron A, iressa, 10 irinotecan, kytril, lapatinib, lentinan sulphate, letrozole, leucovorin, leuprolide, leuprolide acetate, lenalidomide, levamisole, levofolinic acid calcium salt, levothroid, levoxyl, lomustine, lonidamine, marinol, mechlorethamine, mecobalamin, medroxyprogesterone acetate, megestrol acetate, melphalan, menest, 6-mercaptopurine, Mesna, 15 methotrexate, metvix, miltefosine, minocycline, mitomycin C, mitotane, mitoxantrone, Modrenal, Myocet, nedaplatin, neulasta, neumega, neupogen, nilutamide, nolvadex, NSC-631570, OCT-43, octreotide, ondansetron HCl, orapred, oxaliplatin, paclitaxel, pediaped, pegaspargase, Pegasys, pentostatin, picibanil, pilocarpine HCl, 20 pirarubicin, plicamycin, porfimer sodium, prednimustine, prednisolone, prednisone, premarin, procarbazine, procrit, RDEA 119, raltitrexed, rebif, rhenium-186 etidronate, rituximab, roferon-A, romurtide, salagen, sandostatin, sargramostim, semustine, sizofiran, sobuzoxane, solu-medrol, sparfosic acid, stem-cell therapy, streptozocin, strontium-89 25 chloride, sunitinib, synthroid, tamoxifen, tamsulosin, tasonermin, tastolactone, taxotere, teceleukin, temozolomide, teniposide, testosterone propionate, testred, thioguanine, thiotepa, thyrotropin, tiludronic acid, topotecan, toremifene, tositumomab, trastuzumab, treosulfan, tretinoin, trexall, trimethylmelamine, trimetrexate, 30 triptorelin acetate, triptorelin pamoate, UFT, uridine, valrubicin, vesnarinone, vinblastine, vincristine, vindesine, vinorelbine, virulizin, zinecard, zinostatin stimalamer, zofran, ABI-007, acolbifene,

actimmune, affinitak, aminopterin, arzoxifene, asoprisnil, atamestane, atrasentan, BAY 43-9006 (sorafenib), avastin, CCI-779, CDC-501, celebrex, cetuximab, crisnatol, cyproterone acetate, decitabine, DN-101, doxorubicin-MTC, dSLIM, dutasteride, edotecarin, eflornithine, 5 exatecan, fenretinide, histamine dihydrochloride, histrelin hydrogel implant, holmium-166 DOTMP, ibandronic acid, interferon gamma, intron-PEG, ixabepilone, keyhole limpet hemocyanin, L-651582, lanreotide, lasofoxifene, libra, lonafarnib, miproxifene, minodronate, MS-209, liposomal MTP-PE, MX-6, nafarelin, nemorubicin, neovastat, 10 nolatrexed, oblimersen, onco-TCS, osidem, paclitaxel polyglutamate, pamidronate disodium, PN-401, QS-21, quazepam, R-1549, raloxifene, ranpirnase, 13-cis -retinoic acid, satraplatin, seocalcitol, T-138067, tarceva, taxoprexin, thalidomide, thymosin alpha 1, tiazofurine, tipifarnib, tirapazamine, TLK-286, toremifene, TransMID-107R, 15 valspodar, vaporeotide, vatalanib, verteporfin, vinflunine, Z-100, zoledronic acid or combinations thereof.

In accordance with an embodiment, the additional pharmaceutical agent or agents (or “further active agent”) is selected from the group 20 consisting of : 131I-chTNT, abarelix, abiraterone, aclarubicin, aldesleukin, alemtuzumab, alitretinoin, altretamine, aminoglutethimide, amrubicin, amsacrine, anastrozole, arglabin, arsenic trioxide, asparaginase, azacitidine, basiliximab, BAY 80-6946, BAY 1000394, BAY 86-9766 (RDEA 119), belotecan, bendamustine, bevacizumab, 25 bexarotene, bicalutamide, bisantrene, bleomycin, bortezomib, buserelin, busulfan, cabazitaxel, calcium folinate, calcium levofolate, capecitabine, carboplatin, carmofur, carmustine, catumaxomab, celecoxib, celmoleukin, cetuximab, chlorambucil, chlormadinone, chlormethine, cisplatin, cladribine, clodronic acid, clofarabine, 30 crisantaspase, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, darbepoetin alfa, dasatinib, daunorubicin, decitabine, degarelix, denileukin diftitox, denosumab, deslorelin, dibrospidium

chloride, docetaxel, doxifluridine, doxorubicin, doxorubicin + estrone, eculizumab, edrecolomab, elliptinium acetate, eltrombopag, endostatin, enocitabine, epirubicin, epitiostanol, epoetin alfa, epoetin beta, eptaplatin, eribulin, erlotinib, estradiol, estramustine, etoposide, everolimus, exemestane, fadrozole, filgrastim, fludarabine, fluorouracil, flutamide, formestane, fotemustine, fulvestrant, gallium nitrate, ganirelix, gefitinib, gemcitabine, gemtuzumab, glutoxim, goserelin, histamine dihydrochloride, histrelin, hydroxycarbamide, I-125 seeds, ibandronic acid, ibritumomab tiuxetan, idarubicin, ifosfamide, imatinib, imiquimod, improsulfan, interferon alfa, interferon beta, interferon gamma, ipilimumab, irinotecan, ixabepilone, lanreotide, lapatinib, lenalidomide, lenograstim, lentinan, letrozole, leuprorelin, levamisole, lisuride, lobaplatin, lomustine, lonidamine, masoprocol, medroxyprogesterone, megestrol, melphalan, mepitiostane, mercaptopurine, methotrexate, methoxsalen, Methyl aminolevulinate, methyltestosterone, mifamurtide, miltefosine, miriplatin, mitobronitol, mitoguazone, mitolactol, mitomycin, mitotane, mitoxantrone, nedaplatin, nelarabine, nilotinib, nilutamide, nimotuzumab, nimustine, nitracrine, ofatumumab, omeprazole, oprelvekin, oxaliplatin, p53 gene therapy, paclitaxel, palifermin, palladium-103 seed, pamidronic acid, panitumumab, pazopanib, pegaspargase, PEG-epoetin beta (methoxy PEG-epoetin beta), pegfilgrastim, peginterferon alfa-2b, pemetrexed, pentazocine, pentostatin, peplomycin, perfosfamide, picibanil, pirarubicin, plerixafor, plicamycin, poliglusam, polyestradiol phosphate, polysaccharide-K, porfimer sodium, pralatrexate, prednimustine, procarbazine, quinagolide, raloxifene, raltitrexed, ranimustine, razoxane, regorafenib, risedronic acid, rituximab, romidepsin, romiplostim, sargramostim, sipuleucel-T, sizofiran, sobuzoxane, sodium glycididazole, sorafenib, streptozocin, sunitinib, talaporfin, tamibarotene, tamoxifen, tasonermin, teceleukin, tegafur, tegafur + gimeracil + oteracil, temoporfin, temozolomide, temsirolimus, teniposide, testosterone, tetrofosmin, thalidomide, thiotepa,

thymalfasin, tioguanine, tocilizumab, topotecan, toremifene, tositumomab, trabectedin, trastuzumab, treosulfan, tretinoin, trilostane, triptorelin, trofosfamide, tryptophan, ubenimex, valrubicin, vandetanib, vapreotide, vemurafenib, vinblastine, vincristine, vindesine,
5 vinflunine, vinorelbine, vorinostat, vorozole, yttrium-90 glass microspheres, zinostatin, zinostatin stimalamer, zoledronic acid, zorubicin.

10 The additional pharmaceutical agent can also be gemcitabine, paclitaxel, cisplatin, carboplatin, sodium butyrate, 5-FU, doxorubicin, tamoxifen, etoposide, trastumazab, gefitinib, intron A, rapamycin, 17-AAG, U0126, insulin, an insulin derivative, a PPAR ligand, a sulfonylurea drug, an α -glucosidase inhibitor, a biguanide, a PTP-1B inhibitor, a DPP-
15 IV inhibitor, a 11-beta-HSD inhibitor, GLP-1, a GLP-1 derivative, GIP, a GIP derivative, PACAP, a PACAP derivative, secretin or a secretin derivative.

Optional anti-hyper-proliferative agents which can be added to the
20 composition include but are not limited to compounds listed on the cancer chemotherapy drug regimens in the 11th Edition of the *Merck Index*, (1996), which is hereby incorporated by reference, such as asparaginase, bleomycin, carboplatin, carmustine, chlorambucil, cisplatin, colaspase, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, daunorubicin, doxorubicin (adriamycine), epirubicin,
25 etoposide, 5-fluorouracil, hexamethylmelamine, hydroxyurea, ifosfamide, irinotecan, leucovorin, lomustine, mechlorethamine, 6-mercaptopurine, mesna, methotrexate, mitomycin C, mitoxantrone, prednisolone, prednisone, procarbazine, raloxifen, streptozocin, tamoxifen, thioguanine, topotecan, vinblastine, vincristine, and
30 vindesine.

Other anti-hyper-proliferative agents suitable for use with the composition of the invention include but are not limited to those compounds acknowledged to be used in the treatment of neoplastic diseases in *Goodman and Gilman's The Pharmacological Basis of Therapeutics* (Ninth Edition), editor Molinoff et al., publ. by McGraw-Hill, pages 1225-1287, (1996), which is hereby incorporated by reference, such as aminoglutethimide, L-asparaginase, azathioprine, 5-azacytidine, cladribine, busulfan, diethylstilbestrol, 2',2'-difluorodeoxycytidine, docetaxel, erythrohydroxynonyl adenine, ethinyl estradiol, 5-fluorodeoxyuridine, 5-fluorodeoxyuridine monophosphate, fludarabine phosphate, fluoxymesterone, flutamide, hydroxyprogesterone caproate, idarubicin, interferon, medroxyprogesterone acetate, megestrol acetate, melphalan, mitotane, paclitaxel, pentostatin, N-phosphonoacetyl-L-aspartate (PALA), plicamycin, semustine, teniposide, testosterone propionate, thiotepa, trimethylmelamine, uridine, and vinorelbine.

Other anti-hyper-proliferative agents suitable for use with the composition of the invention include but are not limited to other anti-cancer agents such as epothilone and its derivatives, irinotecan, raloxifen and topotecan.

Generally, the use of cytotoxic and/or cytostatic agents in combination with a compound or composition of the present invention will serve to:

25

(1) yield better efficacy in reducing the growth of a tumor or even eliminate the tumor as compared to administration of either agent alone,

30 (2) provide for the administration of lesser amounts of the administered chemotherapeutic agents,

(3) provide for a chemotherapeutic treatment that is better tolerated in the patient with fewer deleterious pharmacological complications than observed with single agent chemotherapies and certain other combined therapies,

5

(4) provide for treating a broader spectrum of different cancer types in mammals, especially humans,

(5) provide for a higher response rate among treated patients,

10

(6) provide for a longer survival time among treated patients compared to standard chemotherapy treatments,

(7) provide a longer time for tumor progression, and/or

15

(8) yield efficacy and tolerability results at least as good as those of the agents used alone, compared to known instances where other cancer agent combinations produce antagonistic effects.

20 In accordance with an embodiment, the invention relates to combinations wherein said 2,3-dihydroimidazo[1,2-c]quinazoline compound is 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide.

25 In accordance with an embodiment, the invention relates to combinations wherein said further active agent is rapamycin or ABT-737.

In accordance with an embodiment, the invention relates to combinations wherein said 2,3-dihydroimidazo[1,2-c]quinazoline
30 compound is 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide and said further active agent is rapamycin or ABT-737.

Pharmaceutical compositions of the compounds of the invention

5

As mentioned *supra*, the present invention relates to pharmaceutical compositions :

- comprising a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, as a sole active agent, for the treatment of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer, and
- comprising a pharmaceutical composition which comprises a combination of :
 - a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; and
 - b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyperproliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral agent, more particularly one or more further active agents selected from the group consisting of :
 - a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
 - a Bcl binding peptide ;

30

- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus),
5 Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-
10 01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

In accordance with another embodiment, the present invention relates to pharmaceutical compositions which comprise a 2,3-dihydroimidazo[1,2-c]quinazoline compound as defined herein, in
15 particular 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, as a sole agent, for the treatment of cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer,
20 Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is
25 inflammatory breast cancer.

In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is triple negative breast cancer.

In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is Her2 receptor positive breast cancer.

- 5 In accordance a particular embodiment of any of the above aspects, or embodiments thereof, of the present invention, said breast cancer is hormone receptor positive breast cancer.

- Said pharmaceutical compositions contain one or more compounds.
- 10 These compositions can be utilized to achieve the desired pharmacological effect by administration to a patient in need thereof. A patient, for the purpose of this invention, is a mammal, including a human, in need of treatment for the particular condition or disease. Therefore, the present invention includes pharmaceutical compositions
- 15 that are comprised of a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound, or salt thereof, of the present invention. A pharmaceutically acceptable carrier is preferably a carrier that is relatively non-toxic and innocuous to a patient at concentrations consistent with effective activity of the active
- 20 agent so that any side effects ascribable to the carrier do not vitiate the beneficial effects of the active agent. A pharmaceutically effective amount of compound is preferably that amount which produces a result or exerts an influence on the particular condition being treated. The compounds of the present invention can be administered with
- 25 pharmaceutically-acceptable carriers well known in the art using any effective conventional dosage unit forms, including immediate, slow and timed release preparations, orally, parenterally, topically, nasally, ophthalmically, optically, sublingually, rectally, vaginally, and the like.
- 30 For oral administration, the compounds can be formulated into solid or liquid preparations such as capsules, pills, tablets, troches, lozenges, melts, powders, solutions, suspensions, or emulsions, and may be

prepared according to methods known to the art for the manufacture of pharmaceutical compositions. The solid unit dosage forms can be a capsule that can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers such as
5 lactose, sucrose, calcium phosphate, and corn starch.

In another embodiment, the compounds of this invention may be tableted with conventional tablet bases such as lactose, sucrose and cornstarch in combination with binders such as acacia, corn starch or
10 gelatin, disintegrating agents intended to assist the break-up and dissolution of the tablet following administration such as potato starch, alginic acid, corn starch, and guar gum, gum tragacanth, acacia, lubricants intended to improve the flow of tablet granulation and to prevent the adhesion of tablet material to the surfaces of the tablet dies
15 and punches, for example talc, stearic acid, or magnesium, calcium or zinc stearate, dyes, coloring agents, and flavoring agents such as peppermint, oil of wintergreen, or cherry flavoring, intended to enhance the aesthetic qualities of the tablets and make them more acceptable to the patient. Suitable excipients for use in oral liquid dosage forms
20 include dicalcium phosphate and diluents such as water and alcohols, for example, ethanol, benzyl alcohol, and polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent or emulsifying agent. Various other materials may be present as coatings or to otherwise modify the physical form of the
25 dosage unit. For instance tablets, pills or capsules may be coated with shellac, sugar or both.

Dispersible powders and granules are suitable for the preparation of an aqueous suspension. They provide the active agent in admixture with a
30 dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional

excipients, for example those sweetening, flavoring and coloring agents described above, may also be present.

The pharmaceutical compositions of this invention may also be in the
5 form of oil-in-water emulsions. The oily phase may be a vegetable oil such as liquid paraffin or a mixture of vegetable oils. Suitable emulsifying agents may be (1) naturally occurring gums such as gum acacia and gum tragacanth, (2) naturally occurring phosphatides such as soy bean and lecithin, (3) esters or partial esters derived from fatty
10 acids and hexitol anhydrides, for example, sorbitan monooleate, (4) condensation products of said partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

15 Oily suspensions may be formulated by suspending the active agent in a vegetable oil such as, for example, arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent such as, for example, beeswax, hard paraffin, or cetyl alcohol. The suspensions may also
20 contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate; one or more coloring agents; one or more flavoring agents; and one or more sweetening agents such as sucrose or saccharin.

Syrups and elixirs may be formulated with sweetening agents such as, for
25 example, glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, and preservative, such as methyl and propyl parabens and flavoring and coloring agents.

The compounds of this invention may also be administered parenterally,
30 that is, subcutaneously, intravenously, intraocularly, intrasynovially, intramuscularly, or interperitoneally, as injectable dosages of the compound in preferably a physiologically acceptable diluent with a

pharmaceutical carrier which can be a sterile liquid or mixture of liquids such as water, saline, aqueous dextrose and related sugar solutions, an alcohol such as ethanol, isopropanol, or hexadecyl alcohol, glycols such as propylene glycol or polyethylene glycol, glycerol ketals such as 2,2-
5 dimethyl-1,1-dioxolane-4-methanol, ethers such as poly(ethylene glycol) 400, an oil, a fatty acid, a fatty acid ester or, a fatty acid glyceride, or an acetylated fatty acid glyceride, with or without the addition of a pharmaceutically acceptable surfactant such as a soap or a detergent, suspending agent such as pectin, carbomers, methycellulose,
10 hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agent and other pharmaceutical adjuvants.

Illustrative of oils which can be used in the parenteral formulations of this invention are those of petroleum, animal, vegetable, or synthetic
15 origin, for example, peanut oil, soybean oil, sesame oil, cottonseed oil, corn oil, olive oil, petrolatum and mineral oil. Suitable fatty acids include oleic acid, stearic acid, isostearic acid and myristic acid. Suitable fatty acid esters are, for example, ethyl oleate and isopropyl myristate. Suitable soaps include fatty acid alkali metal, ammonium,
20 and triethanolamine salts and suitable detergents include cationic detergents, for example dimethyl dialkyl ammonium halides, alkyl pyridinium halides, and alkylamine acetates; anionic detergents, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates; non-ionic detergents, for
25 example, fatty amine oxides, fatty acid alkanolamides, and poly(oxyethylene-oxypropylene)s or ethylene oxide or propylene oxide copolymers; and amphoteric detergents, for example, alkyl-beta-aminopropionates, and 2-alkylimidazoline quarternary ammonium salts, as well as mixtures.

30

The parenteral compositions of this invention will typically contain from about 0.5% to about 25% by weight of the active agent in solution.

Preservatives and buffers may also be used advantageously. In order to minimize or eliminate irritation at the site of injection, such compositions may contain a non-ionic surfactant having a hydrophile-lipophile balance (HLB) preferably of from about 12 to about 17. The quantity of surfactant in such formulation preferably ranges from about 5% to about 15% by weight. The surfactant can be a single component having the above HLB or can be a mixture of two or more components having the desired HLB.

Illustrative of surfactants used in parenteral formulations are the class of polyethylene sorbitan fatty acid esters, for example, sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol.

The pharmaceutical compositions may be in the form of sterile injectable aqueous suspensions. Such suspensions may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents such as, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents which may be a naturally occurring phosphatide such as lecithin, a condensation product of an alkylene oxide with a fatty acid, for example, polyoxyethylene stearate, a condensation product of ethylene oxide with a long chain aliphatic alcohol, for example, heptadeca-ethyleneoxycetanol, a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol such as polyoxyethylene sorbitol monooleate, or a condensation product of an ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride, for example polyoxyethylene sorbitan monooleate.

The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent. Diluents and solvents that may be employed are, for example, water, Ringer's solution, isotonic sodium chloride solutions and isotonic
5 glucose solutions. In addition, sterile fixed oils are conventionally employed as solvents or suspending media. For this purpose, any bland, fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid can be used in the preparation of injectables.

10

A composition of the invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritation excipient which is solid at ordinary temperatures but liquid at the rectal
15 temperature and will therefore melt in the rectum to release the drug. Such materials are, for example, cocoa butter and polyethylene glycol.

Another formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal
20 patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art (see, e.g., US Patent No. 5,023,252, issued June 11, 1991, incorporated herein by reference).
25 Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Controlled release formulations for parenteral administration include liposomal, polymeric microsphere and polymeric gel formulations that
30 are known in the art.

It may be desirable or necessary to introduce the pharmaceutical composition to the patient via a mechanical delivery device. The construction and use of mechanical delivery devices for the delivery of pharmaceutical agents is well known in the art. Direct techniques for, for example, administering a drug directly to the brain usually involve placement of a drug delivery catheter into the patient's ventricular system to bypass the blood-brain barrier. One such implantable delivery system, used for the transport of agents to specific anatomical regions of the body, is described in US Patent No. 5,011,472, issued April 30, 1991.

The compositions of the invention can also contain other conventional pharmaceutically acceptable compounding ingredients, generally referred to as carriers or diluents, as necessary or desired. Conventional procedures for preparing such compositions in appropriate dosage forms can be utilized. Such ingredients and procedures include those described in the following references, each of which is incorporated herein by reference: Powell, M.F. *et al*, "Compendium of Excipients for Parenteral Formulations" *PDA Journal of Pharmaceutical Science & Technology* 1998, 52(5), 238-311; Strickley, R.G. "Parenteral Formulations of Small Molecule Therapeutics Marketed in the United States (1999)-Part-1" *PDA Journal of Pharmaceutical Science & Technology* 1999, 53(6), 324-349; and Nema, S. *et al*, "Excipients and Their Use in Injectable Products" *PDA Journal of Pharmaceutical Science & Technology* 1997, 51(4), 166-171.

Commonly used pharmaceutical ingredients that can be used as appropriate to formulate the composition for its intended route of administration include:

acidifying agents (examples include but are not limited to acetic acid, citric acid, fumaric acid, hydrochloric acid, nitric acid);

alkalinizing agents (examples include but are not limited to ammonia solution, ammonium carbonate, diethanolamine, monoethanolamine, potassium hydroxide, sodium borate, sodium carbonate, sodium hydroxide, triethanolamine, triamine);

5

adsorbents (examples include but are not limited to powdered cellulose and activated charcoal);

aerosol propellants (examples include but are not limited to carbon dioxide, CCl_2F_2 , $\text{F}_2\text{ClC-CClF}_2$ and CClF_3)

10

air displacement agents (examples include but are not limited to nitrogen and argon);

antifungal preservatives (examples include but are not limited to benzoic acid, butylparaben, ethylparaben, methylparaben, propylparaben, sodium benzoate);

15

antimicrobial preservatives (examples include but are not limited to benzalkonium chloride, benzethonium chloride, benzyl alcohol, cetylpyridinium chloride, chlorobutanol, phenol, phenylethyl alcohol, phenylmercuric nitrate and thimerosal);

20

antioxidants (examples include but are not limited to ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, hypophosphorus acid, monothioglycerol, propyl gallate, sodium ascorbate, sodium bisulfite, sodium formaldehyde sulfoxylate, sodium metabisulfite);

25

binding materials (examples include but are not limited to block polymers, natural and synthetic rubber, polyacrylates, polyurethanes, silicones, polysiloxanes and styrene-butadiene copolymers);

30

buffering agents (examples include but are not limited to potassium metaphosphate, dipotassium phosphate, sodium acetate, sodium citrate anhydrous and sodium citrate dihydrate)

5

carrying agents (examples include but are not limited to acacia syrup, aromatic syrup, aromatic elixir, cherry syrup, cocoa syrup, orange syrup, syrup, corn oil, mineral oil, peanut oil, sesame oil, bacteriostatic sodium chloride injection and bacteriostatic water for injection)

10

chelating agents (examples include but are not limited to edetate disodium and edetic acid)

colorants (examples include but are not limited to FD&C Red No. 3, FD&C Red No. 20, FD&C Yellow No. 6, FD&C Blue No. 2, D&C Green No. 5, D&C Orange No. 5, D&C Red No. 8, caramel and ferric oxide red);

15

clarifying agents (examples include but are not limited to bentonite);

emulsifying agents (examples include but are not limited to acacia, cetomacrogol, cetyl alcohol, glyceryl monostearate, lecithin, sorbitan monooleate, polyoxyethylene 50 monostearate);

20

encapsulating agents (examples include but are not limited to gelatin and cellulose acetate phthalate)

25

flavorants (examples include but are not limited to anise oil, cinnamon oil, cocoa, menthol, orange oil, peppermint oil and vanillin);

humectants (examples include but are not limited to glycerol, propylene glycol and sorbitol);

30

levigating agents (examples include but are not limited to mineral oil and glycerin);

oils (examples include but are not limited to arachis oil, mineral oil,
5 olive oil, peanut oil, sesame oil and vegetable oil);

ointment bases (examples include but are not limited to lanolin, hydrophilic ointment, polyethylene glycol ointment, petrolatum,
10 hydrophilic petrolatum, white ointment, yellow ointment, and rose water ointment);

penetration enhancers (transdermal delivery) (examples include but are not limited to monohydroxy or polyhydroxy alcohols, mono-or
15 polyvalent alcohols, saturated or unsaturated fatty alcohols, saturated or unsaturated fatty esters, saturated or unsaturated dicarboxylic acids, essential oils, phosphatidyl derivatives, cephalin, terpenes, amides, ethers, ketones and ureas)

20 **plasticizers** (examples include but are not limited to diethyl phthalate and glycerol);

solvents (examples include but are not limited to ethanol, corn oil, cottonseed oil, glycerol, isopropanol, mineral oil, oleic acid, peanut oil,
25 purified water, water for injection, sterile water for injection and sterile water for irrigation);

stiffening agents (examples include but are not limited to cetyl alcohol, cetyl esters wax, microcrystalline wax, paraffin, stearyl alcohol, white
30 wax and yellow wax);

suppository bases (examples include but are not limited to cocoa butter and polyethylene glycols (mixtures));

5 **surfactants** (examples include but are not limited to benzalkonium chloride, nonoxynol 10, oxtoxynol 9, polysorbate 80, sodium lauryl sulfate and sorbitan mono-palmitate);

10 **suspending agents** (examples include but are not limited to agar, bentonite, carbomers, carboxymethylcellulose sodium, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, kaolin, methylcellulose, tragacanth and veegum);

15 **sweetening agents** (examples include but are not limited to aspartame, dextrose, glycerol, mannitol, propylene glycol, saccharin sodium, sorbitol and sucrose);

tablet anti-adherents (examples include but are not limited to magnesium stearate and talc);

20 **tablet binders** (examples include but are not limited to acacia, alginic acid, carboxymethylcellulose sodium, compressible sugar, ethylcellulose, gelatin, liquid glucose, methylcellulose, non-crosslinked polyvinyl pyrrolidone, and pregelatinized starch);

25 **tablet and capsule diluents** (examples include but are not limited to dibasic calcium phosphate, kaolin, lactose, mannitol, microcrystalline cellulose, powdered cellulose, precipitated calcium carbonate, sodium carbonate, sodium phosphate, sorbitol and starch);

30 **tablet coating agents** (examples include but are not limited to liquid glucose, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl

methylcellulose, methylcellulose, ethylcellulose, cellulose acetate phthalate and shellac);

5 **tablet direct compression excipients** (examples include but are not limited to dibasic calcium phosphate);

tablet disintegrants (examples include but are not limited to alginic acid, carboxymethylcellulose calcium, microcrystalline cellulose, polacrillin potassium, cross-linked polyvinylpyrrolidone, sodium alginate, 10 sodium starch glycollate and starch);

tablet glidants (examples include but are not limited to colloidal silica, corn starch and talc);

15 **tablet lubricants** (examples include but are not limited to calcium stearate, magnesium stearate, mineral oil, stearic acid and zinc stearate);

tablet/capsule opaquants (examples include but are not limited to 20 titanium dioxide);

tablet polishing agents (examples include but are not limited to carnuba wax and white wax);

25 **thickening agents** (examples include but are not limited to beeswax, cetyl alcohol and paraffin);

tonicity agents (examples include but are not limited to dextrose and sodium chloride);

30

viscosity increasing agents (examples include but are not limited to alginic acid, bentonite, carbomers, carboxymethylcellulose sodium,

methylcellulose, polyvinyl pyrrolidone, sodium alginate and tragacanth);
and

wetting agents (examples include but are not limited to
5 heptadecaethylene oxycetanol, lecithins, sorbitol monooleate,
polyoxyethylene sorbitol monooleate, and polyoxyethylene stearate).

Pharmaceutical compositions according to the present invention can be
illustrated as follows:

10

Sterile IV Solution: A 5 mg/mL solution of the desired compound of this
invention can be made using sterile, injectable water, and the pH is
adjusted if necessary. The solution is diluted for administration to 1 - 2
mg/mL with sterile 5% dextrose and is administered as an IV infusion
15 over about 60 minutes.

Lyophilized powder for IV administration: A sterile preparation can be
prepared with (i) 100 - 1000 mg of the desired compound of this
invention as a lyophilized powder, (ii) 32- 327 mg/mL sodium citrate, and
20 (iii) 300 - 3000 mg Dextran 40. The formulation is reconstituted with
sterile, injectable saline or dextrose 5% to a concentration of 10 to 20
mg/mL, which is further diluted with saline or dextrose 5% to 0.2 - 0.4
mg/mL, and is administered either IV bolus or by IV infusion over 15 - 60
minutes.

25

Intramuscular suspension: The following solution or suspension can be
prepared, for intramuscular injection:

50 mg/mL of the desired, water-insoluble compound of this invention

5 mg/mL sodium carboxymethylcellulose

30 4 mg/mL TWEEN 80

9 mg/mL sodium chloride

9 mg/mL benzyl alcohol

Hard Shell Capsules: A large number of unit capsules are prepared by filling standard two-piece hard galantine capsules each with 100 mg of powdered active agent, 150 mg of lactose, 50 mg of cellulose and 6 mg of magnesium stearate.

Soft Gelatin Capsules: A mixture of active agent in a digestible oil such as soybean oil, cottonseed oil or olive oil is prepared and injected by means of a positive displacement pump into molten gelatin to form soft gelatin capsules containing 100 mg of the active agent. The capsules are washed and dried. The active agent can be dissolved in a mixture of polyethylene glycol, glycerin and sorbitol to prepare a water miscible medicine mix.

Tablets: A large number of tablets are prepared by conventional procedures so that the dosage unit is 100 mg of active agent, 0.2 mg. of colloidal silicon dioxide, 5 mg of magnesium stearate, 275 mg of microcrystalline cellulose, 11 mg. of starch, and 98.8 mg of lactose. Appropriate aqueous and non-aqueous coatings may be applied to increase palatability, improve elegance and stability or delay absorption.

Immediate Release Tablets/Capsules: These are solid oral dosage forms made by conventional and novel processes. These units are taken orally without water for immediate dissolution and delivery of the medication. The active agent is mixed in a liquid containing ingredient such as sugar, gelatin, pectin and sweeteners. These liquids are solidified into solid tablets or caplets by freeze drying and solid state extraction techniques. The drug compounds may be compressed with viscoelastic and thermoelastic sugars and polymers or effervescent components to produce porous matrices intended for immediate release, without the need of water.

Method of treating breast cancer

The present invention also relates to a method of treating or prophylaxis
5 of cancer, e.g. breast cancer, which is classified into several subtypes in
the clinic, such as hormone receptor positive breast cancer, Her2
receptor positive breast cancer, triple negative breast cancer and
inflammatory breast cancer, in a mammal, said method comprising
administering a 2,3-dihydroimidazo[1,2-c]quinazoline compound as
10 defined herein, or a pharmaceutical composition containing same, as a
sole active agent, or administering a combination of a) said compound or
a pharmaceutical composition containing said compound and b) one or
more further active agents as defined herein.

15 In accordance a particular embodiment of any of the above aspects, or
embodiments thereof, of the present invention, said breast cancer is
inflammatory breast cancer.

In accordance a particular embodiment of any of the above aspects, or
20 embodiments thereof, of the present invention, said breast cancer is
triple negative breast cancer.

In accordance a particular embodiment of any of the above aspects, or
embodiments thereof, of the present invention, said breast cancer is
25 Her2 receptor positive breast cancer.

In accordance a particular embodiment of any of the above aspects, or
embodiments thereof, of the present invention, said breast cancer is
hormone receptor positive breast cancer.

30

The embodiments of the methods of treating or prophylaxis of cancer,
e.g. breast cancer, as defined supra, are as described in the

embodiments of the use of the compounds/combinations, as described supra.

5 The present invention relates to a method for using the compounds of the present invention and compositions thereof, to treat mammalian breast cancer. Compounds can be utilized to inhibit, block, reduce, decrease, etc., cell proliferation and/or cell division, and/or produce apoptosis, in the treatment or prophylaxis of breast cancer. This method comprises administering to a mammal in need thereof, including
10 a human, an amount of a compound or combination of this invention, or a pharmaceutically acceptable salt, isomer, polymorph, metabolite, hydrate, solvate or ester thereof; etc. which is effective for the treatment or prophylaxis of breast cancer.

15 This disorder has been well characterized in humans, but also exists with a similar etiology in other mammals, and they can be treated by administering pharmaceutical compositions of the present invention.

The term “treating” or “treatment” as stated throughout this document
20 is used conventionally, e.g., the management or care of a subject for the purpose of combating, alleviating, reducing, relieving, improving the condition of, etc., of a disease or disorder, such as a carcinoma.

Dose and administration

25

Based upon standard laboratory techniques known to evaluate compounds useful for the treatment or prophylaxis of breast cancer, by standard toxicity tests and by standard pharmacological assays for the determination of treatment of the conditions identified above in
30 mammals, and by comparison of these results with the results of known medicaments that are used to treat these conditions, the effective dosage of the compounds of this invention can readily be determined for

treatment of the indication. The amount of the active agent to be administered in the treatment of the condition can vary widely according to such considerations as the particular compound and dosage unit employed, the mode of administration, the period of treatment, the
5 age and sex of the patient treated, and the nature and extent of the condition treated.

The total amount of the active agent to be administered will generally range from about 0.001 mg/kg to about 200 mg/kg body weight per day,
10 and preferably from about 0.01 mg/kg to about 20 mg/kg body weight per day. Clinically useful dosing schedules will range from one to three times a day dosing to once every four weeks dosing. In addition, "drug holidays" in which a patient is not dosed with a drug for a certain period of time, may be beneficial to the overall balance between
15 pharmacological effect and tolerability. A unit dosage may contain from about 0.5 mg to about 1,500 mg of active agent, and can be administered one or more times per day or less than once a day. The average daily dosage for administration by injection, including intravenous, intramuscular, subcutaneous and parenteral injections, and
20 use of infusion techniques will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily rectal dosage regimen will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily vaginal dosage regimen will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily topical dosage regimen will
25 preferably be from 0.1 to 200 mg administered between one to four times daily. The transdermal concentration will preferably be that required to maintain a daily dose of from 0.01 to 200 mg/kg. The average daily inhalation dosage regimen will preferably be from 0.01 to 100 mg/kg of total body weight.

30

Of course the specific initial and continuing dosage regimen for each patient will vary according to the nature and severity of the condition as

determined by the attending diagnostician, the activity of the specific compound employed, the age and general condition of the patient, time of administration, route of administration, rate of excretion of the drug, drug combinations, and the like. The desired mode of treatment and
5 number of doses of a compound of the present invention or a pharmaceutically acceptable salt or ester or composition thereof can be ascertained by those skilled in the art using conventional treatment tests.

10

Biomarkers :

15 Biomarkers used for patient stratification are e.g. Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN for predicting the sensitivity and/or resistance of a cancer patient to said compound, thus providing rationale-based synergistic combination as defined herein to overcome the resistance.

20

Examples:

The invention is demonstrated in the following examples which are not
25 meant to limit the invention in any way:

Example 1

In accordance with the invention, the compound were assessed in a cell-
30 based assay that measures the capacity of the compounds to inhibit tumor cell proliferation following a 72-hour drug exposure. Cell viability was conducted by Invitrogen (breast Oncopanel, Invitrogen, USA), or was determined using CellTiter-Glo® Luminescent Cell Viability Assay

(Promega) as follows. Cells were plated at 300 - 5000 cells/well depending on the cell lines or microtiter plates (96- or 384-well) in 100 μ L growth medium in 96-well and 20 μ L growth medium in 384-well microtiterplate. For each cell line assayed, cells were plated onto a
5 separate plate for determination of luminescence at $t = 0$ hour and $t = 72$ hour time points. Following overnight incubation at 37 °C, luminescence values for the $t = 0$ samples were determined. Dose plates for the $t = 72$ hour time points were treated with compounds diluted into growth medium. Cells were then incubated for 72 hours at 37 °C.
10 Luminescence values for the $t = 72$ hour samples were determined. For data analysis, briefly, $t = 0$ values were subtracted from those determined for the $t = 72$ hour time points, for both the treated and untreated samples. Percent differences in luminescence between drug-treated and control values were used to determine the percent
15 inhibition of growth.

Tumor cells were seeded in 384-well plates at a concentration of 1000-2500 cells/well/25 μ L depending on cell lines. After overnight incubation, cells were treated with compounds (5 μ L). Then, Caspase
20 3/7 activities were determined at 24 to 48 hours post-treatment using Caspase-Glo3/7 assay kits (Promega Cat # G8212).

The combination effects of PI3K inhibitors and other agent were evaluated using combination index isobologram analysis (Chou et al.
25 Pharmacology Reviews 2006). The efficacy parameters were the median effect in a 72-hour cell proliferation assay and the 24- or 48-hour Caspase 3/7 activity assay described above. Briefly, cells were plated in 384-well plate with 25 μ L medium. After 24 hours, 5 μ L of experimental media containing either drug 1 (D1), or drug 2 (D2), or the combination
30 of D1 plus D2 at different ratios (0.8xD1+0.2xD2, 0.6xD1+0.4xD2, 0.4xD1+0.6xD2, 0.2xD1+0.8xD2, 0.1xD1+0.9xD2) were used to make serial three-fold dilutions to generate 7 dose curves. Experiments were

conducted in triplicate. For determination of EC₅₀s and IC₅₀s, a linear regression analysis are used to determine drug concentration which results in a 50% inhibition or efficacy. The corresponding component doses of D1 and D2 at the E(I)C₅₀/ E(I)C₉₀ were calculated and used for plotting isobolograms. The multiple drug effect was analyzed as described by Chou et al. (Pharmacology Reviews 2006) and the combination index was calculated using the formula :

$$\text{Combination Index} = [D1x] / D1' + [D2x] / D2'$$


10

where D1x and D2x refer to the Drug 1 and Drug 2 concentration at EC₅₀/IC₅₀ or EC₉₀/IC₉₀, respectively, in combination; and D1' and D2' refer to the EC₅₀/IC₅₀ or EC₉₀/IC₉₀ values of D1 and D2, respectively, as single agents. In this analysis, values less than 1.0 indicate synergistic interactions, values greater than 1.0 indicate antagonistic interactions, and values around 1.0 indicate additive interactions.

15

In the following, "compound of formula I" refers to 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide, of structure :

20




(I)

25

or a solvate, hydrate or stereoisomer thereof.

In the following, "compound A" refers to 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide dihydrochloride, of structure :



10

or a solvate, hydrate or stereoisomer thereof.

The synthesis of compound A is described in European patent application number EP 11 161 111.7, which is hereby incorporated herein in its entirety by reference.

15

Synthesis of compound A :

To a suspension of the compound of formula I (400 g) in water (1,1 L) at room temperature was added a 32% aqueous 32% (aqueous) hydrochloric acid solution iswith stirring dosed at room temperature to a suspension of 400 g of the compound of formula (I) in 1.1 L water until a pH of 3-4 is was reached. Additional 90 mL water (90 mL) and 32% hydrochloric acid are were added until a pH of 1.8 to 2.0 is was attained. E160 mL ethanol (160 mL) are dosed into was added to the mixture, followed by seed crystals. After stirring for 30 minutes, 1740 gadditional ethanol (2,2 L) are dosed within 5 hwas added into the mixture over 5 h, which isand the resulting mixture was subsequently stirred for 1 h. The suspension is filtered and the residue is washed first with a mixture of 130 g water and 215 g ethanol, secondly with a mixture of 80 g water and 255 g

20

25

30

ethanol and then with 320 g pure ethanol. The filter cake is dried at 40 °C under vacuum to yield 457 g product (99% of theory).

Characterization of compound A:

- 5 The chemical structure of compound A has been confirmed using the described methods of structural analysis.

IR and Raman spectroscopy

Apparatus and measuring conditions

FT-IR / FT-Raman-Spectrometer	Bruker IFS 66v	/	Bruker RFS 100
Spectral resolution	2 cm ⁻¹	/	2 cm ⁻¹
Number of interferograms	32	/	64
Wave number range	4000 - 500 cm ⁻¹	/	3500 - 100 cm ⁻¹
Laser power	-	/	350mW
Sample preparation	KBr pellet	/	solid in test tube

Assignment of the characteristic bands

- 10 Table: Assignment of the characteristic active vibrations to the spectrum with ν $\equiv$ stretching vibrations; δ $\equiv$ bending vibrations; o.o.p. $\equiv$ out of plane.

Assigned Structure	IR Band position [cm ⁻¹]	Raman Band position [cm ⁻¹]
ν N-H	3336	-
ν =C-H	3176	3090
ν C-H	2942	2990 - 2963
ν NH ⁺	2687 - 2474	-
ν Amide I	1669	1664
ν C=C, ν C=N, δ N-H, Amide II	1618 - 1477	1619 - 1476
ν C-O	1285	1291
δ =C-H o.o.p.	812	-

- 15 ν $\equiv$ stretching vibrations; δ $\equiv$ bending vibrations; o.o.p. $\equiv$ out of plane

The IR spectrum is given in Figure 7.

The Raman spectrum is given in Figure 8.

5

UV/VIS spectroscopy

Apparatus and measuring conditions

UV/VIS spectrometer	Varian Cary 4
Cuvette	Quartz, 1 cm
Wave number range	200-800 nm
Sample preparation	4.67 mg / 500 mL water
Bands	309 nm

10 The UV/vis spectrum is given in Figure 9.

NMR spectroscopy

¹H-NMR-spectroscopy


15

Equipment and experimental parameters:

NMR spectrometer	Bruker, model Avance
Working frequency	500.13 MHz
Solvent	Dimethylsulfoxide (DMSO-d ₆)
Internal reference compound	Tetramethylsilane (TMS)
Concentration	3.08 mg/mL solution
Diameter of sample tube	5 mm
Temperature	approx. 25 °C
Technique	Fourier transform mode
Spectral width	20.65ppm

Digital resolution	0.079 Hz/Pt
Pulse length	4.5 μ sec, 30° Pulse flip angle
Acquisition time	6.34sec
Relaxation time	0.5sec
No. of free induction decays	32

Structural Formula for the assignment of NMR signals



5

Chemical shift, signal multiplicity, relative number of nuclei :

H-atoms(a)	Chemical shift δ (ppm)	Multiplicity and coupling constants (b)	no. of nuclei H/molecule
------------	----------------------------------	---	-----------------------------

H-26	2.32	M	2
H-29; H-33	3.11; 3.48	M; M	2; 2
H-30; H-32	3.83; 3.98	M; M	2; 2
H-27	3.29	M	2
-OCH ₃	4.00	S	3
H-25	4.37	T	2
H-2; H-3	4.47; 4.19	T; T	2; 2
H-9	7.39	D	1
NH ₂	7.54	S	2
H-10	8.21	D	1
H-16; H-20	8.97	S	1; 1
HCl	11.1; 12.6	bS; bS	1; 1
H-12	13.4	bS	1

a) Numbering refers to the structural formula for the assignment of NMR-signals.

5 b) S = Singlet bS = broad Singlet D = Doublet
 T = Triplet M = Multiplet

The ¹H-NMR Spectrum of compound A is given in Figure 10.

10

¹³C-NMR-spectroscopy

15

Equipment and experimental parameters

NMR spectrometer	Bruker, model Avance
Working frequency	125.76 MHz
Solvent	Dimethylsulfoxide-d ₆ (DMSO)
Internal reference compound	Tetramethylsilane (TMS)
Concentration	37.2 mg/mL solution
Diameter of sample tube	5 mm
Temperature	approx. 27° C
Technique	Fourier transform mode
Spectral width	240.95 ppm
Digital resolution	0.4624 Hz/Pt
Pulse length	11.0 µsec, 90° Pulse flip angle
Acquisition time	1.08 sec
Relaxation time	4 sec
No. of free induction decays	256

Chemical shift, signal multiplicity, rel. no. of nuclei :

C-atoms(a)	Chemical shift δ (ppm)	Multiplicity and coupling constants (b)	no. of nuclei C/molecule
C-26	22.73	T	1
C-2; C-3	44.96; 45.65	T; T	1; 1
C-29; C-33	50.84	T	1; 1
C-27	53.01	T	1
OCH ₃	61.24	Q	1
C-30; C-32	63.03	T	1; 1
C-25	66.81	T	1
C-10a	100.79	S	1
C-9	112.17	D	1
C-15	118.16	S	1
C-10	123.86	D	1
C-6a	132.43	S	1
C-7	133.95	S	1
C-5	148.58	S	1
C-11	156.29	S	1
C-8	156.89	S	1
C-16; C-20	160.20	D	1; 1
C-18	164.61	S	1
C=O	175.65	S	1

5 a) Numbering refers to the structural formula for the assignment of NMR-signals.

b) S = Single (C) D = Doublet (CH) T = Triplet (CH₂) Q = Quadruplet (CH₃)

The ¹³C-NMR Spectra of compound A are given in Figures 11 and 12.

Mass Spectrometry
Instrumental Parameters

5	
Mass spectrometer	Waters ZQ
Ionization mode	ESI (Electrospray-Ionization)
Solvent	CH ₃ CN/H ₂ O

Interpretation of the Spectrum

<u>Mass value (m/z)</u>	<u>Rel. Intensity (%)</u>	<u>Ion Formation</u>
481.2	46	(M + H) +
354.1	5	(C16 H16 N7 O3) ⁺
261.7	26	(M + 2H + CH ₃ CN) ⁺²
241.2	100	(M + 2H) ⁺²

10

The Mass Spectrum of compound A is given in Figure 13. Refer to the spectrum
15 for relative peak intensities.

Elemental Analysis

Elemental analysis was conducted by Bayer Industry Services, Leverkusen, Germany.

5 **Results**

Element	Measured [%]	Calculated [%]	Calculated including 7.0 % water [%]	Difference
C	47.5	49.9	46.4	1.1
H	5.7	5.5	5.9	0.2
N	19.1	20.3	18.8	0.3
O	18.1	11.6	17.0	1.1
Cl	11.9	12.8	11.9	0.0
Sum	102.3	100.1	100.0	-

The elemental analysis is consistent with compound A with 7% water.

Further method of preparation of compound "A"

10

To a suspension of 366 g of compound of formula (I) in 1015 g water, 183 g of an aqueous hydrochloric acid solution (32%) were added while maintaining the temperature at 20 °C (+-2 °) until a pH of 3 to 4 was reached. The resulting mixture was stirred at room temperature for more than 10 min. filtered and the filtercake washed with additional 82 g of water. The filtrate was adjusted to pH 1.8 to 2.0 using aqueous hydrochloric acid solution (32%). The mixture was stirred for 10 min. at room temperature, 146g of ethanol (100%) were added and stirred for another 10 min.. 1 g of seed crystals were added, followed by 1592 g ethanol within 5 h. The resulting substance was removed by filtration,

20

washed with a water-ethanol mixture and dried in vacuo to give 410 g (97%) of compound A of a purity >99% according to HPLC.

Molecular profiling of Compound A in 24 breast tumor cell lines indicated
5 that tumor cells with either *PIK3CA* mutation and/or HER2 expression
were extremely sensitive to Compound A with an average IC₅₀ value of 17
(n=7) and 19 nM (n=8), respectively. In contrast, wild type *PIK3CA* and
HER2 negative breast cancer cells (n=12) were relatively insensitive or
resistant to Compound A with an average IC₅₀ value of 773 nM (~40-fold
10 higher). Using a set of breast cancer cell lines representing differential
receptor expression status and genetic alterations of *PIK3CA* and *PTEN*,
the anti-proliferative and apoptotic effects of Compound A were
investigated. With regard to apoptosis induction, Compound A
demonstrated strong activity in *PIK3CA* mutant and or HER2 positive
15 breast tumor cells (such as BT20, BT474).

Next, the molecular features responsible for the sensitivity and/or
resistance to Compound A mediated induction of apoptosis were
identified. It was found that the expression of Bcl-2, but not Mcl-1 nor
20 surviving, determined sensitivity to apoptosis. Thus, tumor cells that
lack Bcl-2 expression (such as KPL-4, BT-474, and BT-20), even in the
presence of high levels of EGFR and phospho-ERK in BT-20 cells,
immediately underwent apoptosis after exposure to Compound A. On the
other hand, cells expressing high level of Bcl-2, such as T47D
25 (*PIK3CA*^{mut}), were resistant to apoptosis induced by Compound A, despite
the potent anti-proliferative activity of Compound A (single-digit nM
IC₅₀). Combining Compound A with ABT-737, an inhibitor of Bcl-2,
sensitized T47D cells to apoptosis, further supporting the hypothesis that
simultaneous inhibiting PI3K and Bcl-2 could synergistically enhance the
30 tumor killing effect in Bcl-2 positive breast cancer.

On the other hand, PTEN^{null}, EGFR^{high} and Bcl-2 negative MDA-MB-468 tumor cells, are not only insensitive to inhibition of proliferation, but also resistant to induction of apoptosis by Compound A. Inhibition of mTORC1 by rapamycin and/or inhibition of mTORC2 by the knockdown of Rictor, sensitizes MDA-MB468 cells to Compound A with the respect to inhibition of proliferation and induction of apoptosis.

In conclusion, Compound A was extremely effective at inducing apoptosis in breast cancer cells expressing HER2 and/or *PIK3CA* mutation in the absence of Bcl-2. Compound A in combination with anti-Bcl agents or mTOR inhibitors might be the promising approach to achieve tumor responses in Bcl-2 positive tumors or EGFR expressing/PTEN^{null} breast tumors, respectively. These findings provide a rationale for developing personalized therapies for the treatment of molecular subtypes of breast cancer.

Figure 1 shows the correlation of IC₅₀s of Compound A in proliferation assays and genetic status of breast cancer cell lines.

Figure 2 shows kinetics of cell growth inhibition by Compound A. The Xcelligence technique (Roche, Germany) was used to monitor effects on tumor cell growth after treatment by Compound A. Cell index (CI), which correlates with the number of cells attached to the bottom of the plate (Giordano C, Masi A, Pizzini A, Sansone A, Consalvi V, Chiaraluce R, Lucente G. Synthesis and activity of fibrillogenesis peptide inhibitors related to the 17-21 beta-amyloid sequence. Eur J Med Chem. 2009;44:179-189), was detected every 15 minutes for 80 hours treatment period. CI was normalized to the time point of treatment (at 24h CI =1).

Figure 3 shows induction of PARP cleavage by Compound A in BT-20, BT-474, ZR-75-1, T47D and MDA-MB468 cells investigated at the indicated

timepoint by western blot analysis using GAPDH as an internal protein control.

Figure 4 shows pro-survival molecules of the Bcl family investigated in untreated cells (A). Cells were lysed and western blot analysis was performed to detect Bcl-2, Mcl-1, and survivin. GAPDH was used as an internal protein control.

Figure 5 shows quantitative analysis of the combination effect of compound A and ABT-737 on Caspase 3 and 7 induction in T47D cells (C). Apoptosis induction by compound A and ABT-737 as a single agent and in combination was investigated by Caspase-Glo[®] 3/7 (Promega, USA) and analyzed by software Analyze 5 (Bayer HealthCare, Germany). The combination effects are depicted by isobologram and combination index based on the formula described by Chou (Chou et al., 2006). Combination index < 1, =1, and > 1 indicate synergistic, additive, and antagonistic effects, respectively.

Figure 6 shows MDA-MB-468 cells transfected with Lentivirus expressing shRNA against Rictor. The cellular expression of the mTORC2 component Rictor was assessed by western blot analysis (A). Induction of PARP cleavage by compound A treatment in combination with inhibition of mTORC1 or mTORC2 (B). shControl and shRictor MDA-MB468 cells were treated with indicated agents for 48 hours, and cleavage of PARP was analyzed. GAPDH was used as an internal protein control.

CONTROL SUBSTANCES

Rapamycin obtained from Sigma (St Louis, MO, USA) was used as a reference inhibitor. ABT-737 was obtained from Selleck Chemicals (Houston, TX, USA)

In conclusion, Compound A was extremely effective at inducing apoptosis in breast cancer cells expressing HER2 and/or a *PIK3CA* mutation in the absence of Bcl-2. Compound A in combination with anti-
5 Bcl agents or mTOR inhibitors should provide a promising approach to achieve tumor responses in Bcl-2 positive tumors or EGFR expressing/PTEN^{null} breast tumors, respectively. These findings provide a rationale to develop personalized therapies for the treatment of molecular subtypes of breast cancer.

10

Hence, as mentioned *supra*, the present invention relates to the use of biomarkers involved in the modification of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN for predicting the sensitivity and/or resistance of a patient with cancer, e.g.
15 breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer, to a 2,3-dihydroimidazo[1,2-c]quinazoline compound as defined herein, thus providing rationale-based synergistic combination as defined herein to overcome the resistance.

20

In accordance with an embodiment, the present invention relates to the use of biomarkers involved in the modification of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN for predicting the sensitivity and/or resistance of a patient with
25 cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer, to a 2,3-dihydroimidazo[1,2-c]quinazoline compound as defined herein, thus providing a rationale-based synergistic combination as defined herein to overcome the
30 resistance (patient stratification).

In accordance with an embodiment, the present invention relates to a method of determining the level of a component of one or more of Bcl expression, HER family expression and/or activation, PIK3CA signaling and / or loss of PTEN, wherein :

- 5 - in said Bcl expression, said component is Bcl, for example,
- in said HER family expression and/or activation, PIK3CA signaling, said component is EGF-R, for example, and
- in said loss of PTEN, said component is PTEN, for example.

10 Further, as mentioned *supra*, the present invention thus relates to combinations of :

a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound as defined *supra*, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or pharmaceutical compositions containing such a compound or
15 a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

and

20

b) one or more further active agents, in particular an active agent selected from an anti-angiogenesis, anti-hyper-proliferative, antiinflammatory, analgesic, immunoregulatory, diuretic, antiarrhythmic, anti-hypercholesterolemia, anti-dyslipidemia, anti-diabetic or antiviral
25 agent, more particularly one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- 30

- a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin
5 analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01,
10 PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235,

as defined *supra*.

In accordance a particular embodiment of any of the above aspects, or
15 embodiments thereof, of the present invention, said breast cancer is inflammatory breast cancer.

In accordance a particular embodiment of any of the above aspects, or
embodiments thereof, of the present invention, said breast cancer is
20 triple negative breast cancer.

In accordance a particular embodiment of any of the above aspects, or
embodiments thereof, of the present invention, said breast cancer is
Her2 receptor positive breast cancer.

25

In accordance a particular embodiment of any of the above aspects, or
embodiments thereof, of the present invention, said breast cancer is
hormone receptor positive breast cancer.

30

References:

1. Abbosh, P. H.; Nephew, K. P. Multiple signaling pathways converge on b-catenin in thyroid cancer. *Thyroid* 2005, 15, 551-561.
- 5 2. Ali, I. U.; Schriml, L. M.; Dean, M. Mutational spectra of PTEN/MMAC1 gene: a tumor suppressor with lipid phosphatase activity. *J. Natl. Cancer Inst.* 1999, 91, 1922-1932.
3. Bachman, K. E.; Argani, P.; Samuels, Y.; Silliman, N.; Ptak, J.; Szabo, S.; Konishi, H.; Karakas, B.; Blair, B. G.; Lin, C.; Peters, B. A.;
10 Velculescu, V. E.; Park, B. H. The PIK3CA gene is mutated with high frequency in human breast cancers. *Cancer Biol. Therap.* 2004, 3, 772-775.
4. Bader, A. G.; Kang, S.; Vogt, P. K. Cancer-specific mutations in PIK3CA are oncogenic in vivo. *Proc. Natl. Acad. Sci. U. S. A.* 2006, 103,
15 1475-1479.
5. Barthwal, M. K.; Sathyanarayana, P.; Kundu, C. N.; Rana, B.; Pradeep, A.; Sharma, C.; Woodgett, J. R.; Rana, A. Negative Regulation of Mixed Lineage Kinase 3 by Protein Kinase B/AKT Leads to Cell Survival. *J. Biol. Chem.* 2003, 278, 3897-3902.
- 20 6. Bénistant, C.; Chapuis, H.; Roche, S. A specific function for phosphatidylinositol 3-kinase a (p85a-p110a) in cell survival and for phosphatidylinositol 3-kinase b (p85a-p110b) in de novo DNA synthesis of human colon carcinoma cells. *Oncogene* 2000, 19, 5083-5090.
7. Broderick, D. K.; Di, C.; Parrett, T. J.; Samuels, Y. R.; Cummins, J. M.; McLendon, R. E.; Fults, D. W.; Velculescu, V. E.; Bigner, D. D.;
25 Yan, H. Mutations of PIK3CA in anaplastic oligodendrogliomas, high-grade astrocytomas, and medulloblastomas. *Cancer Res.* 2004, 64, 5048-5050.
8. Brown, R. A.; Shepherd, P. R. Growth factor regulation of the novel class II phosphoinositide 3-kinases. *Biochem. Soc. Trans.* 2001, 29,
30 535-537.
9. Brunet, A.; Bonni, A.; Zigmond, M. J.; Lin, M. Z.; Juo, P.; Hu, L. S.; Anderson, M. J.; Arden, K. C.; Blenis, J.; Greenberg, M. E. Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* 1999, 96, 857-868.
- 35 10. Byun, D.-S.; Cho, K.; Ryu, B.-K.; Lee, M.-G.; Park, J.-I.; Chae, K.-S.; Kim, H.-J.; Chi, S.-G. Frequent monoallelic deletion of PTEN and its reciprocal association with PIK3CA amplification in gastric carcinoma. *Int. J. Cancer* 2003, 104, 318-327.

11. Campbell, I. G.; Russell, S. E.; Choong, D. Y. H.; Montgomery, K. G.; Ciavarella, M. L.; Hooi, C. S. F.; Cristiano, B. E.; Pearson, R. B.; Phillips, W. A. Mutation of the PIK3CA gene in ovarian and breast cancer. *Cancer Res.* 2004, 64, 7678-7681.
- 5 12. Cardone, M. H.; Roy, N.; Stennicke, H. R.; Salvesen, G. S.; Franke, T. F.; Stanbridge, E.; Frisch, S.; Reed, J. C. Regulation of cell death protease caspase-9 by phosphorylation. *Science* 1998, 282, 1318-1321.
- 10 13. Chen, Y. L.; Law, P.-Y.; Loh, H. H. Inhibition of PI3K/Akt signaling: An emerging paradigm for targeted cancer therapy. *Curr. Med. Chem. Anticancer Agents* 2005, 5, 575-589.
- 15 14. Ciechomska, I.; Pyrzynska, B.; Kazmierczak, P.; Kaminska, B. Inhibition of Akt kinase signaling and activation of Forkhead are indispensable for up-regulation of FasL expression in apoptosis of glioma cells. *Oncogene* 2003, 22, 7617-7627.
- 15 15. Cross, D. A. E.; Alessi, D. R.; Cohen, P.; Andjelkovich, M.; Hemmings, B. A. Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B. *Nature* 1995, 378, 785-9.
- 20 16. Cully, M.; You, H.; Levine, A. J.; Mak, T. W. Beyond PTEN mutations: the PI3K pathway as an integrator of multiple inputs during tumorigenesis. *Nat. Rev. Cancer* 2006, 6, 184-192.
- 25 17. Czauderna, F.; Fechtner, M.; Aygun, H.; Arnold, W.; Klippel, A.; Giese, K.; Kaufmann, J. Functional studies of the PI(3)-kinase signaling pathway employing synthetic and expressed siRNA. *Nucleic Acids Res.* 2003, 31, 670-682.
18. del Peso, L.; González-Garcia, M.; Page, C.; Herrera, R.; Nunez, G. Interleukin-3-induced phosphorylation of BAD through the protein kinase Akt. *Science* 1997, 278, 687-689.
- 30 19. Diehl, J. A.; Cheng, M.; Roussel, M. F.; Sherr, C. J. Glycogen synthase kinase-3b regulates cyclin D1 proteolysis and subcellular localization. *Genes Dev.* 1998, 12, 3499-3511.
- 35 20. Dijkers, P. F.; Medema, R. H.; Lammers, J.-W. J.; Koenderman, L.; Coffey, P. J. Expression of the pro-apoptotic Bcl-2 family member Bim is regulated by the Forkhead transcription factor FKHR-L1. *Curr. Biol.* 2000, 10, 1201-1204.
21. Domin, J.; Waterfield, M. D. Using structure to define the function of phosphoinositide 3-kinase family members. *FEBS Lett.* 1997, 410, 91-95.

22. Downes, C. P.; Gray, A.; Lucocq, J. M. Probing phosphoinositide functions in signaling and membrane trafficking. *Trends Cell Biol.* 2005, 15, 259-268.
23. Figueroa, C.; Tarras, S.; Taylor, J.; Vojtek, A. B. Akt2 negatively
5 regulates assembly of the POSH-MLK-JNK signaling complex. *J. Biol. Chem.* 2003, 278, 47922-47927.
24. Fleming, I. N.; Gray, A.; Downes, C. P. Regulation of the Rac1-specific exchange factor tiam1 involves both phosphoinositide 3-kinase-dependent and -independent components. *Biochem. J.* 2000, 351, 173-
10 182.
25. Funaki, M.; Katagiri, H.; Inukai, K.; Kikuchi, M.; Asano, T. Structure and function of phosphatidylinositol-3,4 kinase. *Cell. Signal.* 2000, 12, 135-142.
26. Gallia, G. L.; Rand, V.; Siu, I. M.; Eberhart, C. G.; James, C. D.;
15 Marie, S. K. N.; Oba-Shinjo, S. M.; Carlotti, C. G.; Caballero, O. L.; Simpson, A. J. G.; Brock, M. V.; Massion, P. P.; Carson, B. S., Sr.; Riggins, G. J. PIK3CA gene mutations in pediatric and adult glioblastoma multiforme. *Mol. Cancer Res.* 2006, 4, 709-714.
27. Gershtein, E. S.; Shatskaya, V. A.; Ermilova, V. D.; Kushlinsky, N. E.; Krasil'nikov, M. A. Phosphatidylinositol 3-kinase expression in human
20 breast cancer. *Clin. Chim. Acta* 1999, 287, 59-67.
28. Gottschalk, A. R.; Doan, A.; Nakamura, J. L.; Stokoe, D.; Haas-Kogan, D. A. Inhibition of phosphatidylinositol-3-kinase causes increased sensitivity to radiation through a PKB-dependent mechanism. *Int. J. Radiat. Oncol. Biol. Phys.* 2005, 63, 1221-1227.
25
29. Gupta, A. K.; Cerniglia, G. J.; Mick, R.; Ahmed, M. S.; Bakanauskas, V. J.; Muschel, R. J.; McKenna, W. G. Radiation sensitization of human cancer cells in vivo by inhibiting the activity of PI3K using LY294002. *Int. J. Radiat. Oncol. Biol. Phys.* 2003, 56, 846-853.
30. Haas-Kogan, D.; Shalev, N.; Wong, M.; Mills, G.; Yount, G.; Stokoe, D. Protein kinase B (PKB/Akt) activity is elevated in glioblastoma cells due to mutation of the tumor suppressor PTEN/MMAC. *Curr. Biol.* 1998, 8, 1195-1198.
31. Hartmann, C.; Bartels, G.; Gehlhaar, C.; Holtkamp, N.; von
35 Deimling, A. PIK3CA mutations in glioblastoma multiforme. *Acta Neuropathol.* 2005, 109, 639-642.
32. Hennesy, B. T.; Smith, D. L.; Ram, P. T.; Lu, Y.; Mills, G. B. Exploiting the PI3K/AKT Pathway for Cancer Drug Discovery. *Nat. Rev. Drug Disc.* 2005, 4, 988-1004.

33. Hodgkinson, C. P.; Sale, E. M.; Sale, G. J. Characterization of PDK2 activity against Protein Kinase B gamma. *Biochemistry* 2002, 41, 10351-10359.
34. Hresko, R. C.; Murata, H.; Mueckler, M. Phosphoinositide-dependent Kinase-2 is a distinct protein kinase enriched in a novel cytoskeletal fraction associated with adipocyte plasma membranes. *J. Biol. Chem.* 2003, 278, 21615-21622.
35. Huang, C.; Ma, W.-Y.; Dong, Z. Requirement for phosphatidylinositol 3-kinase in epidermal growth factor-induced AP-1 transactivation and transformation in JB6 P+ cells. *Mol. Cell. Biol.* 1996, 16, 6427-6435.
36. Hupp, T. R.; Lane, D. P.; Ball, K. L. Strategies for manipulating the p53 pathway in the treatment of human cancer. *Biochem. J.* 2000, 352, 1-17.
37. Ihle, N. T.; Williams, R.; Chow, S.; Chew, W.; Berggren, M. I.; Paine-Murrieta, G.; Minion, D. J.; Halter, R. J.; Wipf, P.; Abraham, R.; Kirkpatrick, L.; Powis, G. Molecular pharmacology and antitumor activity of PX-866, a novel inhibitor of phosphoinositide-3-kinase signaling. *Mol. Cancer Therap.* 2004, 3, 763-772.
38. Ikenoue, T.; Kanai, F.; Hikiba, Y.; Obata, T.; Tanaka, Y.; Imamura, J.; Ohta, M.; Jazag, A.; Guleng, B.; Tateishi, K.; Asaoka, Y.; Matsumura, M.; Kawabe, T.; Omata, M. Functional analysis of PIK3CA gene mutations in human colorectal cancer. *Cancer Res.* 2005, 65, 4562-4567.
39. Ishii, N.; Maier, D.; Merlo, A.; Tada, M.; Sawamura, Y.; Diserens, A.-C.; Van Meir, E. G. Frequent co-alterations of TP53, p16/CDKN2A, p14ARF, PTEN tumor suppressor genes in human glioma cell lines. *Brain Pathol.* 1999, 9, 469-479.
40. Itoh, T.; Takenawa, T. Phosphoinositide-binding domains. Functional units for temporal and spatial regulation of intracellular signaling. *Cell. Signal.* 2002, 14, 733-743.
41. Janssen, J. W. G.; Schleithoff, L.; Bartram, C. R.; Schulz, A. S. An oncogenic fusion product of the phosphatidylinositol 3-kinase p85b subunit and HUMORF8, a putative deubiquitinating enzyme. *Oncogene* 1998, 16, 1767-1772.
42. Jimenez, C.; Jones, D. R.; Rodriguez-Viciana, P.; Gonzalez-Garcia, A.; Leonardo, E.; Wennstrom, S.; Von Kobbe, C.; Toran, J. L.; R.-Borlado, L.; Calvo, V.; Copin, S. G.; Albar, J. P.; Gaspar, M. L.; Diez, E.; Marcos, M. A. R.; Downward, J.; Martinez-A, C.; Merida, I.; Carrera, A. C. Identification and characterization of a new oncogene derived from

the regulatory subunit of phosphoinositide 3-kinase. *EMBO J.* 1998, 17, 743-753.

43. Jucker, M.; Sudel, K.; Horn, S.; Sickel, M.; Wegner, W.; Fiedler, W.; Feldman, R. A. Expression of a mutated form of the p85a regulatory subunit of phosphatidylinositol 3-kinase in a Hodgkin's lymphoma-derived cell line (CO). *Leukemia* 2002, 16, 894-901.
44. Kang, S.; Bader, A. G.; Vogt, P. K. Phosphatidylinositol 3-kinase mutations identified in human cancer are oncogenic. *Proc. Natl. Acad. Sci. U. S. A.* 2005, 102, 802-807.
45. Kang, S.; Denley, A.; Vanhaesebroeck, B.; Vogt, P. K. Oncogenic transformation induced by the p110b, -g, and -d isoforms of class I phosphoinositide 3-kinase. *Proc. Natl. Acad. Sci. U. S. A.* 2006, 103, 1289-1294.
46. Katso, R.; Okkenhaug, K.; Ahmadi, K.; White, S.; Timms, J.; Waterfield, M. D. Cellular function of phosphoinositide 3-kinases: implications for development, immunity, homeostasis, and cancer. *Annu. Rev. Cell Dev. Biol.* 2001, 17, 615-675.
47. Kim, A. H.; Khursigara, G.; Sun, X.; Franke, T. F.; Chao, M. V. Akt phosphorylates and negatively regulates apoptosis signal-regulating kinase 1. *Mol. Cell. Biol.* 2001, 21, 893-901.
48. Kim, D.; Dan, H. C.; Park, S.; Yang, L.; Liu, Q.; Kaneko, S.; Ning, J.; He, L.; Yang, H.; Sun, M.; Nicosia, S. V.; Cheng, J. Q. AKT/PKB signaling mechanisms in cancer and chemoresistance. *Front. Biosci.* 2005, 10, 975-987.
49. Klippel, A.; Kavanaugh, W. M.; Pot, D.; Williams, L. T. A specific product of phosphatidylinositol 3-kinase directly activates the protein kinase Akt through its pleckstrin homology domain. *Mol. Cell. Biol.* 1997, 17, 338-44.
50. Kodaki, T.; Woscholski, R.; Hallberg, B.; Rodriguez-Viciano, P.; Downward, J.; Parker, P. J. The activation of phosphatidylinositol 3-kinase by Ras. *Curr. Biol.* 1994, 4, 798-806.
51. Kops, G. J. P. L.; De Ruiter, N. D.; De Vries-Smits, A. M. M.; Powell, D. R.; Bos, J. L.; Burgering, B. M. T. Direct control of the Forkhead transcription factor AFX by protein kinase B. *Nature* 1999, 398, 630-634.
52. Lee, J. T., Jr.; Steelman, L. S.; McCubrey, J. A. Phosphatidylinositol 3'-Kinase Activation Leads to Multidrug Resistance Protein-1 Expression and Subsequent Chemoresistance in Advanced Prostate Cancer Cells. *Cancer Res.* 2004, 64, 8397-8404.

53. Lee, J. W.; Soung, Y. H.; Kim, S. Y.; Lee, H. W.; Park, W. S.; Nam, S. W.; Kim, S. H.; Lee, J. Y.; Yoo, N. J.; Lee, S. H. PIK3CA gene is frequently mutated in breast carcinomas and hepatocellular carcinomas. *Oncogene* 2005, 24, 1477-1480.
- 5 54. Lemmon, M. A. Phosphoinositide recognition domains. *Traffic* 2003, 4, 201-213.
55. Levine, D. A.; Bogomolnii, F.; Yee, C. J.; Lash, A.; Barakat, R. R.; Borgen, P. I.; Boyd, J. Frequent Mutation of the PIK3CA Gene in Ovarian and Breast Cancers. *Clin. Cancer Res.* 2005, 11, 2875-2878.
- 10 56. Li, J.; Yen, C.; Liaw, D.; Podsypanina, K.; Bose, S.; Wang, S. I.; Puc, J.; Miliarensis, C.; Rodgers, L.; McCombie, R.; Bigner, S. H.; Giovanella, B. C.; Ittmann, M.; Tycko, B.; Hibshoosh, H.; Wigler, M. H.; Parsons, R. PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* 1997, 275, 1943-1947.
- 15 57. Li, V. S. W.; Wong, C. W.; Chan, T. L.; Chan, A. S. W.; Zhao, W.; Chu, K.-M.; So, S.; Chen, X.; Yuen, S. T.; Leung, S. Y. Mutations of PIK3CA in gastric adenocarcinoma. *BMC Cancer* 2005, 5, 29.
58. Liao, Y.; Hung, M.-C. Regulation of the activity of p38 mitogen-activated protein kinase by Akt in cancer and adenoviral protein E1A-mediated sensitization to apoptosis. *Mol. Cell. Biol.* 2003, 23, 6836-6848.
- 20 59. Lopez-Illasaca, M.; Li, W.; Uren, A.; Yu, J.-c.; Kazlauskas, A.; Gutkind, J. S.; Heidaran, M. A. Requirement of phosphatidylinositol-3 kinase for activation of JNK/SAPKs by PDGF. *Biochem. Biophys. Res. Commun.* 1997, 232, 273-277.
- 25 60. Ma, Y.-Y.; Wei, S.-J.; Lin, Y.-C.; Lung, J.-C.; Chang, T.-C.; Whang-Peng, J.; Liu, J. M.; Yang, D.-M.; Yang, W. K.; Shen, C.-Y. PIK3CA as an oncogene in cervical cancer. *Oncogene* 2000, 19, 2739-2744.
61. Mayo, L. D.; Dixon, J. E.; Durden, D. L.; Tonks, N. K.; Donner, D. B. PTEN protects p53 from Mdm2 and sensitizes cancer cells to chemotherapy. *J. Biol. Chem.* 2002, 277, 5484-5489.
- 30 62. Momand, J.; Wu, H.-H.; Dasgupta, G. MDM2 - master regulator of the p53 tumor suppressor protein. *Gene* 2000, 242, 15-29.
63. Motti, M. L.; De Marco, C.; Califano, D.; Fusco, A.; Viglietto, G. Akt-dependent T198 phosphorylation of cyclin-dependent kinase inhibitor p27kip1 in breast cancer. *Cell Cycle* 2004, 3, 1074-1080.
- 35 64. Myers, M. P.; Pass, I.; Batty, I. H.; Van Der Kaay, J.; Stolarov, J. P.; Hemmings, B. A.; Wigler, M. H.; Downes, C. P.; Tonks, N. K. The lipid

phosphatase activity of PTEN is critical for its tumor suppressor function. Proc. Natl. Acad. Sci. U. S. A. 1998, 95, 13513-13518.

65. Nagata, Y.; Lan, K.-H.; Zhou, X.; Tan, M.; Esteva, F. J.; Sahin, A. A.; Klos, K. S.; Li, P.; Monia, B. P.; Nguyen, N. T.; Hortobagyi, G. N.; Hung, M.-C.; Yu, D. PTEN activation contributes to tumor inhibition by trastuzumab, and loss of PTEN predicts trastuzumab resistance in patients. Cancer Cell 2004, 6, 117-127.
66. Naito, A. T.; Akazawa, H.; Takano, H.; Minamino, T.; Nagai, T.; Aburatani, H.; Komuro, I. Phosphatidylinositol 3-Kinase-Akt Pathway Plays a Critical Role in Early Cardiomyogenesis by Regulating Canonical Wnt Signaling. Circ. Res. 2005, 97, 144-151.
67. Oda, K.; Stokoe, D.; Taketani, Y.; McCormick, F. High Frequency of Coexistent Mutations of PIK3CA and PTEN Genes in Endometrial Carcinoma. Cancer Res. 2005, 65, 10669-10673.
68. Ogawara, Y.; Kishishita, S.; Obata, T.; Isazawa, Y.; Suzuki, T.; Tanaka, K.; Masuyama, N.; Gotoh, Y. Akt enhances Mdm2-mediated ubiquitination and degradation of p53. J. Biol. Chem. 2002, 277, 21843-21850.
69. Olson, J. M.; Hallahan, A. R. p38 MAP kinase: a convergence point in cancer therapy. Trends Mol. Med. 2004, 10, 125-129.
70. Osaki, M.; Oshimura, M.; Ito, H. PI3K-Akt pathway: Its functions and alterations in human cancer. Apoptosis 2004, 9, 667-676.
71. Pastorino, J. G.; Tafani, M.; Farber, J. L. Tumor necrosis factor induces phosphorylation and translocation of BAD through a phosphatidylinositide-3-OH kinase-dependent pathway. J. Biol. Chem. 1999, 274, 19411-19416.
72. Pendaries, C.; Tronchere, H.; Plantavid, M.; Payrastre, B. Phosphoinositide signaling disorders in human diseases. FEBS Lett. 2003, 546, 25-31.
73. Phillips, W. A.; St. Clair, F.; Munday, A. D.; Thomas, R. J. S.; Mitchell, C. A. Increased levels of phosphatidylinositol 3-kinase activity in colorectal tumors. Cancer 1998, 83, 41-47.
74. Philp, A. J.; Campbell, I. G.; Leet, C.; Vincan, E.; Rockman, S. P.; Whitehead, R. H.; Thomas, R. J. S.; Phillips, W. A. The phosphatidylinositol 3'-kinase p85a gene is an oncogene in human ovarian and colon tumors. Cancer Res. 2001, 61, 7426-7429.
75. Powis, G.; Bonjouklian, R.; Berggren, M. M.; Gallegos, A.; Abraham, R.; Ashendel, C.; Zalkow, L.; Matter, W. F.; Dodge, J.

Wortmannin, a potent and selective inhibitor of phosphatidylinositol-3-kinase. *Cancer Res.* 1994, 54, 2419-23.

76. Pu, P.; Kang, C.; Zhang, Z.; Liu, X.; Jiang, H. Downregulation of PIK3CB by siRNA suppresses malignant glioma cell growth in vitro and in vivo. *Technol. Cancer Res. Treat.* 2006, 5, 271-280.

77. Rahimi, N.; Tremblay, E.; Elliott, B. Phosphatidylinositol 3-kinase activity is required for hepatocyte growth factor-induced mitogenic signals in epithelial cells. *J. Biol. Chem.* 1996, 271, 24850-24855.

78. Roche, S.; Downward, J.; Raynal, P.; Courtneidge, S. A. A function for phosphatidylinositol 3-kinase b (p85a-p110b) in fibroblasts during mitogenesis: requirement for insulin- and lysophosphatidic acid-mediated signal transduction. *Mol. Cell. Biol.* 1998, 18, 7119-7129.

79. Roche, S.; Koegl, M.; Courtneidge, S. A. The phosphatidylinositol 3-kinase a is required for DNA synthesis induced by some, but not all, growth factors. *Proc. Natl. Acad. Sci. U. S. A.* 1994, 91, 9185-9.

80. Romashkova, J. A.; Makarov, S. S. Nf-kB is a target of Akt in anti-apoptotic PDGF signaling. *Nature* 1999, 401, 86-90.

81. Saal, L. H.; Holm, K.; Maurer, M.; Memeo, L.; Su, T.; Wang, X.; Yu, J. S.; Malmstroem, P.-O.; Mansukhani, M.; Enoksson, J.; Hibshoosh, H.; Borg, A.; Parsons, R. PIK3CA mutations correlate with hormone receptors, node metastasis, and ERBB2, and are mutually exclusive with PTEN loss in human breast carcinoma. *Cancer Res.* 2005, 65, 2554-2559.

82. Samuels, Y.; Diaz, L. A., Jr.; Schmidt-Kittler, O.; Cummins, J. M.; DeLong, L.; Cheong, I.; Rago, C.; Huso, D. L.; Lengauer, C.; Kinzler, K. W.; Vogelstein, B.; Velculescu, V. E. Mutant PIK3CA promotes cell growth and invasion of human cancer cells. *Cancer Cell* 2005, 7, 561-573.

83. Samuels, Y.; Ericson, K. Oncogenic PI3K and its role in cancer. *Curr. Opin. Oncol.* 2006, 18, 77-82.

84. Samuels, Y.; Wang, Z.; Bardelli, A.; Silliman, N.; Ptak, J.; Szabo, S.; Yan, H.; Gazdar, A.; Powell, S. M.; Riggins, G. J.; Willson, J. K. V.; Markowitz, S.; Kinzler, K. W.; Vogelstein, B.; Velculescu, V. E. Brevia: High frequency of mutations of the PIK3Ca gene in human cancers. *Science* 2004, 304, 554.

85. Scheid, M. P.; Marignani, P. A.; Woodgett, J. R. Multiple phosphoinositide 3-kinase-dependent steps in activation of protein kinase B. *Mol. Cell. Biol.* 2002, 22, 6247-6260.


86. Schultz, R. M.; Merriman, R. L.; Andis, S. L.; Bonjouklian, R.; Grindey, G. B.; Rutherford, P. G.; Gallegos, A.; Massey, K.; Powis, G. In vitro and in vivo antitumor activity of the phosphatidylinositol-3-kinase inhibitor, wortmannin. *Anticancer Res.* 1995, 15, 1135-9.
- 5 87. Segrelles, C.; Moral, M.; Lara, M. F.; Ruiz, S.; Santos, M.; Leis, H.; Garcia-Escudero, R.; Martinez-Cruz, A. B.; Martinez-Palacio, J.; Hernandez, P.; Ballestin, C.; Paramio, J. M. Molecular determinants of Akt-induced keratinocyte transformation. *Oncogene* 2006, 25, 1174-1185.
- 10 88. Sekimoto, T.; Fukumoto, M.; Yoneda, Y. 14-3-3 suppresses the nuclear localization of threonine 157-phosphorylated p27Kip1. *EMBO J.* 2004, 23, 1934-1942.
- 15 89. Semba, S.; Itoh, N.; Ito, M.; Youssef, E. M.; Harada, M.; Moriya, T.; Kimura, W.; Yamakawa, M. Down-regulation of PIK3CG catalytic subunit of phosphatidylinositol 3-OH kinase by CpG hypermethylation in human colorectal carcinoma. *Clin. Cancer Res.* 2002, 8, 3824-3831.
- 20 90. Shayesteh, L.; Lu, Y.; Kuo, W.-L.; Baldocchi, R.; Godfrey, T.; Collins, C.; Pinkel, D.; Powell, B.; Mills, G. B.; Gray, J. W. PIK3CA is implicated as an oncogene in ovarian cancer. *Nat. Genet.* 1999, 21, 99-102.
91. Shekar, S. C.; Wu, H.; Fu, Z.; Yip, S.-C.; Nagajyothi; Cahill, S. M.; Girvin, M. E.; Backer, J. M. Mechanism of Constitutive Phosphoinositide 3-Kinase Activation by Oncogenic Mutants of the p85 Regulatory Subunit. *J. Biol. Chem.* 2005, 280, 27850-27855.
- 25 92. Stahl, J. M.; Cheung, M.; Sharma, A.; Trivedi, N. R.; Shanmugam, S.; Robertson, G. P. Loss of PTEN Promotes Tumor Development in Malignant Melanoma. *Cancer Res.* 2003, 63, 2881-2890.
93. Stambolic, V.; Suzuki, A.; De La Pompa, J. L.; Brothers, G. M.; Mirtsos, C.; Sasaki, T.; Ruland, J.; Penninger, J. M.; Siderovski, D. P.; Mak, T. W. Negative regulation of PKB/Akt-Dependent cell survival by the tumor suppressor PTEN. *Cell* 1998, 95, 29-39.
- 30 94. Stauffer, F.; Holzer, P.; Garcia-Echeverria, C. Blocking the PI3K/PKB pathway in tumor cells. *Curr. Med. Chem. Anticancer Agents* 2005, 5, 449-462.
- 35 95. Steck, P. A.; Pershouse, M. A.; Jasser, S. A.; Yung, W. K. A.; Lin, H.; Ligon, A. H.; Langford, L. A.; Baumgard, M. L.; Hattier, T.; Davis, T.; Frye, C.; Hu, R.; Swedlund, B.; Teng, D. H. F.; Tavtigian, S. V. Identification of a candidate tumor suppressor gene, MMAC1, at chromosome 10q23.3 that is mutated in multiple advanced cancers. *Nat. Genet.* 1997, 15, 356-362.
- 40

96. Stein, R. C.; Waterfield, M. D. PI3-kinase inhibition: a target for drug development? *Mol. Med. Today* 2000, 6, 347-358.
97. Stephens, L.; Williams, R.; Hawkins, P. Phosphoinositide 3-kinases as drug targets in cancer. *Curr. Opin. Pharmacol.* 2005, 5, 357-365.
- 5 98. Su, J. D.; Mayo, L. D.; Donner, D. B.; Durden, D. L. PTEN and Phosphatidylinositol 3'-Kinase Inhibitors Up-Regulate p53 and Block Tumor-induced Angiogenesis: Evidence for an Effect on the Tumor and Endothelial Compartment. *Cancer Res.* 2003, 63, 3585-3592.
- 10 99. Tanaka, M.; Grossman, H. B. In vivo gene therapy of human bladder cancer with PTEN suppresses tumor growth, downregulates phosphorylated Akt, and increases sensitivity to doxorubicin. *Gene Ther.* 2003, 10, 1636-1642.
- 15 100. Tang, E. D.; Nunez, G.; Barr, F. G.; Guan, K.-L. Negative regulation of the forkhead transcription factor FKHR by Akt. *J. Biol. Chem.* 1999, 274, 16741-16746.
101. Taylor, V.; Wong, M.; Brandts, C.; Reilly, L.; Dean, N. M.; Cowser, L. M.; Moodie, S.; Stokoe, D. 5' Phospholipid phosphatase SHIP-2 causes protein kinase B inactivation and cell cycle arrest in glioblastoma cells. *Mol. Cell. Biol.* 2000, 20, 6860-6871.
- 20 102. Toker, A. Phosphoinositides and signal transduction. *Cell. Mol. Life Sci.* 2002, 59, 761-779.
103. Traer, C. J.; Foster, F. M.; Abraham, S. M.; Fry, M. J. Are class II phosphoinositide 3-kinases potential targets for anticancer therapies? *Bull. Cancer (Paris)*. 2006, 93, E53-8.
- 25 104. Vanhaesebroeck, B.; Leivers, S. J.; Ahmadi, K.; Timms, J.; Katso, R.; Driscoll, P. C.; Woscholski, R.; Parker, P. J.; Waterfield, M. D. Synthesis and function of 3-phosphorylated inositol lipids. *Annu. Rev. Biochem.* 2001, 70, 535-602.
105. Vanhaesebroeck, B.; Waterfield, M. D. Signaling by Distinct Classes of Phosphoinositide 3-Kinases. *Exp. Cell Res.* 1999, 253, 239-254.
- 30 106. Vivanco, I.; Sawyers, C. L. The phosphatidylinositol 3-Kinase-AKT pathway in human cancer. *Nat. Rev. Cancer* 2002, 2, 489-501.
107. Wang, Y.; Helland, A.; Holm, R.; Kristensen Gunnar, B.; Borresen-Dale, A.-L. PIK3CA mutations in advanced ovarian carcinomas. *Hum. Mutat.* 2005, 25, 322.
- 35

108. West, K. A.; Castillo, S. S.; Dennis, P. A. Activation of the PI3K/Akt pathway and chemotherapeutic resistance. *Drug Resist. Update.* 2002, 5, 234-48.
109. Whyte, D. B.; Holbeck, S. L. Correlation of PIK3Ca mutations with
5 gene expression and drug sensitivity in NCI-60 cell lines. *Biochem. Biophys. Res. Commun.* 2006, 340, 469-475.
110. Wilker, E.; Lu, J.; Rho, O.; Carbajal, S.; Beltran, L.; DiGiovanni, J. Role of PI3K/Akt signaling in insulin-like growth factor-1 (IGF-1) skin tumor promotion. *Mol. Carcinog.* 2005, 44, 137-145.
- 10 111. Workman, P. Inhibiting the phosphoinositide 3-kinase pathway for cancer treatment. *Biochem. Soc. Trans.* 2004, 32, 393-396.
112. Wu, G.; Xing, M.; Mambo, E.; Huang, X.; Liu, J.; Guo, Z.; Chatterjee, A.; Goldenberg, D.; Gollin, S. M.; Sukumar, S.; Trink, B.; Sidransky, D. Somatic mutation and gain of copy number of PIK3CA in
15 human breast cancer. *Breast Cancer Res.* 2005, 7, R609-R616.
113. Wymann, M. P.; Sozzani, S.; Altruda, F.; Mantovani, A.; Hirsch, E. Lipids on the move: phosphoinositide 3-kinases in leukocyte function. *Immunol. Today* 2000, 21, 260-264.
114. Yap, D. B.; Hsieh, J. K.; Lu, X. Mdm2 inhibits the apoptotic
20 function of p53 mainly by targeting it for degradation. *J. Biol. Chem.* 2000, 275, 37296-302.
115. Yuan, Z.-q.; Feldman, R. I.; Sussman, G. E.; Coppola, D.; Nicosia, S. V.; Cheng, J. Q. AKT2 Inhibition of Cisplatin-induced JNK/p38 and Bax Activation by Phosphorylation of ASK1: Implication of AKT2 in
25 Chemoresistance. *J. Biol. Chem.* 2003, 278, 23432-23440.
116. Zhao, H.; Dupont, J.; Yakar, S.; Karas, M.; LeRoith, D. PTEN inhibits cell proliferation and induces apoptosis by downregulating cell surface IGF-IR expression in prostate cancer cells. *Oncogene* 2004, 23, 786-794.
- 30 117. Zhao, J. J.; Cheng, H.; Jia, S.; Wang, L.; Gjoerup, O. V.; Mikami, A.; Roberts, T. M. The p110a isoform of PI3K is essential for proper growth factor signaling and oncogenic transformation. *Proc. Natl. Acad. Sci. U. S. A.* 2006, 103, 16296-300.
118. Zhou, B. P.; Liao, Y.; Xia, W.; Spohn, B.; Lee, M.-H.; Hung, M.-C. Cytoplasmic localization of p21Cip1/WAF1 by Akt-induced
35 phosphorylation in HER-2/neu-overexpressing cells. *Nat. Cell Biol.* 2001, 3, 245-252.

CLAIMS

1. Use of a 2,3-dihydroimidazo[1,2-c]quinazoline compound of general
5 formula :



(I)

10

or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof, in which :

R¹ represents $-(CH_2)_n-(CHR^4)-(CH_2)_m-N(R^5)(R^{5'})$;

R² represents a heteroaryl optionally substituted with 1, 2 or 3 R⁶
15 groups ;

R³ represents alkyl or cycloalkyl ;

R⁴ represents hydrogen, hydroxy or alkoxy ;

R⁵ and R^{5'} may be the same or different and are independently,
hydrogen, alkyl, cycloalkylalkyl, or alkoxyalkyl or R⁵ and R^{5'} may
20 be taken together with the nitrogen atom to which they are bound
to form a 3-7 membered nitrogen containing heterocyclic ring
optionally containing at least one additional heteroatom selected
from oxygen, nitrogen or sulfur and which may be optionally
substituted with 1 or more R^{6'} groups, or R⁴ and R⁵ may be taken
25 together with the atoms to which they are bound to form a 5-6
membered nitrogen containing heterocyclic ring optionally

containing 1 or more nitrogen, oxygen or sulfur atoms and which may be optionally substituted with 1 or more $R^{6'}$ groups;

each occurrence of R^6 may be the same or different and is
5 independently halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclic ring, heterocyclalkyl, alkyl-OR⁷, alkyl-SR⁷, alkyl-N(R⁷)(R^{7'}), alkyl-COR⁷, -CN, -COOR⁷, -CON(R⁷)(R^{7'}), -OR⁷, -SR⁷, -N(R⁷)(R^{7'}), or -NR⁷COR⁷ each of which may be optionally substituted with 1 or more R^8 groups;

10

each occurrence of $R^{6'}$ may be the same or different and is independently alkyl, cycloalkylalkyl, or alkyl-OR⁷;

each occurrence of R^7 and $R^{7'}$ may be the same or different and is
15 independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heterocyclic ring, heterocyclalkyl, or heteroarylalkyl;

each occurrence of R^8 is independently nitro, hydroxy, cyano, formyl, acetyl, halogen, amino, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heterocyclic ring, heterocyclalkyl, or heteroarylalkyl;

n is an integer from 1-4 and m is an integer from 0-4 with the proviso that when R^4 and R^5 are taken together with the atoms to which they are bound to form a 3-7 membered nitrogen containing ring,
25 $n + m \leq 4$;

as a sole active agent,

30 or of a pharmaceutical composition containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,


for the preparation of a medicament for the treatment or prophylaxis of breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor
5 positive breast cancer.

2. The use according to claim 1, wherein in said compound of formula (I), R^4 is hydroxy.

10 3. The use according to claim 1, wherein in said compound of formula (I), R^4 and R^5 are taken together with the atoms to which they are bound to form a 5-6 membered nitrogen containing heterocyclic ring optionally containing 1 or more nitrogen, oxygen or sulfur atoms and which may be optionally substituted with 1 or more $R^{6'}$ groups.

15 4. The use according to claim 1, wherein in said compound of formula (I), R^2 is pyridine, pyridazine, pyrimidine, pyrazine, pyrrole, oxazole, thiazole, furan or thiophene, optionally substituted with 1, 2 or 3 R^6 groups.

20 5. The use according to claim 1, wherein said compound of formula (I) has the formula :



6. The use according to claim 5, wherein in said compound, R^2 is pyridine, pyridazine, pyrimidine, pyrazine, pyrrole, oxazole, thiazole,
25 furan or thiophene, optionally substituted with 1, 2 or 3 R^6 groups.

7. The use according to claim 1, wherein said compound is, namely:

- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;
- 5 N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide ;
- N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-2,4-dimethyl-1,3-thiazole-5-carboxamide;
- 10 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1,3-thiazole-5-carboxamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]isonicotinamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-4-methyl-1,3-thiazole-5-
- 15 carboxamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-4-propylpyrimidine-5-carboxamide;
- 20 N-{8-[2-(4-ethylmorpholin-2-yl)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}pyrimidine-5-carboxamide;
- N-(8-{3-[2-(hydroxymethyl)morpholin-4-yl]propoxy}-7-methoxy-
- 25 2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- N-(8-{3-[2-(hydroxymethyl)morpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-
- 30 c]quinazolin-5-yl}nicotinamide 1-oxide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-(2-pyrrolidin-1-ylethyl)nicotinamide;
6-(cyclopentylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
5 N-[8-(2-hydroxy-3-morpholin-4-ylpropoxy)-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
N-{7-methoxy-8-[3-(3-methylmorpholin-4-yl)propoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
10 N-(8-{3-[2-(hydroxymethyl)morpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
N-(8-{2-[4-(cyclobutylmethyl)morpholin-2-yl]ethoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
N-(7-methoxy-8-{2-[4-(2-methoxyethyl)morpholin-2-yl]ethoxy}-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
15 N-{8-[(4-ethylmorpholin-2-yl)methoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
N-(7-methoxy-8-{[4-(2-methoxyethyl)morpholin-2-yl]methoxy}-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
20 N-{7-methoxy-8-[(4-methylmorpholin-2-yl)methoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-4-carboxamide;
2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-4-carboxamide;
25 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1-methyl-1H-imidazole-4-carboxamide;
rel-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)pyrimidine-5-carboxamide;
30

rel-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)-6-methylnicotinamide;

5 rel-6-acetamido-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1-methyl-1H-imidazole-5-carboxamide;

10 6-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-methylnicotinamide;
2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-4-methylpyrimidine-5-carboxamide;

15 6-amino-5-bromo-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1,3-oxazole-5-carboxamide;
N-[7-methoxy-8-(morpholin-2-ylmethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;

20 2-{[2-(dimethylamino)ethyl]amino}-N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}pyrimidine-5-carboxamide;
2-amino-N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}-1,3-thiazole-5-carboxamide;
25 rel-2-amino-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)pyrimidine-5-carboxamide;

rel-6-amino-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;

30

2-[(2-hydroxyethyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

5 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-[(3-methoxypropyl)amino]pyrimidine-5-carboxamide;

2-amino-N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}pyrimidine-5-carboxamide;

10 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-[(3-morpholin-4-ylpropyl)amino]pyrimidine-5-carboxamide;

2-[(2-methoxyethyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

15 2-{[2-(dimethylamino)ethyl]amino}-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

6-amino-N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;

20 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-pyrrolidin-1-ylpyrimidine-5-carboxamide;

25 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-(4-methylpiperazin-1-yl)pyrimidine-5-carboxamide;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-morpholin-4-ylpyrimidine-5-carboxamide;

30 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-piperazin-1-ylnicotinamide hydrochloride;

- 6-[(3S)-3-aminopyrrolidin-1-yl]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide hydrochloride hydrate;
- 5 6-[(3R)-3-aminopyrrolidin-1-yl]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide hydrochloride;
- 6-[(4-fluorobenzyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 10 6-[(2-furylmethyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 6-[(2-methoxyethyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 15 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-(1H-pyrrol-1-yl)nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-morpholin-4-ylnicotinamide;
- 20 N-{7-methoxy-8-[3-(methylamino)propoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 6-[(2,2-dimethylpropanoyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 25 6-[(cyclopropylcarbonyl)amino]-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-(2,2,2-trifluoroethoxy)nicotinamide;
- 30 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-(trifluoromethyl)nicotinamide;

- 6-(isobutyrylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-{7-methoxy-8-[3-(4-methylpiperazin-1-yl)propoxy]-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 5 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-
{[(methylamino)carbonyl]amino}-1,3-thiazole-4-carboxamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-
- 10 {[(methylamino)carbonyl]amino}nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-(methylamino)-1,3-thiazole-4-carboxamide;
- N-[7-methoxy-8-(2-morpholin-4-ylethoxy)-2,3-dihydroimidazo[1,2-
- 15 c]quinazolin-5-yl]nicotinamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}-2,4-dimethyl-1,3-thiazole-5-carboxamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-
- 20 dihydroimidazo[1,2-c]quinazolin-5-yl}-6-methylnicotinamide;
- 6-{[(isopropylamino)carbonyl]amino}-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-
- 25 dihydroimidazo[1,2-c]quinazolin-5-yl]-6-pyrrolidin-1-ylnicotinamide;
- 6-(dimethylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-piperidin-1-ylpropoxy)-2,3-dihydroimidazo[1,2-
- 30 c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(2-pyrrolidin-1-ylethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;

- N-[7-methoxy-8-(2-piperidin-1-ylethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 6-{[(ethylamino)carbonyl]amino}-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 5 6-fluoro-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1,3-oxazole-4-carboxamide;
- 2-(ethylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-1,3-thiazole-4-carboxamide;
- 10 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrazine-2-carboxamide;
- N-[8-(2-aminoethoxy)-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- 15 6-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]isonicotinamide;
- N-{8-[3-(diethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 20 N-{8-[2-(diisopropylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- N-{8-[2-(diethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 25 N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 30 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-(methylamino)pyrimidine-5-carboxamide;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-(methylthio)pyrimidine-5-carboxamide;

5 N-[8-(3-aminopropoxy)-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide trifluoroacetate;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]thiophene-2-carboxamide;

10 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2,4-dimethyl-1,3-thiazole-5-carboxamide;

2-methoxy-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-3-furamide;

15 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]thiophene-3-carboxamide;
N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2-methyl-1,3-thiazole-4-carboxamide;

20 6-methoxy-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;

5-methoxy-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;

25 N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-methylnicotinamide;

6-(acetylamino)-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide ;

30

preferably,

- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-6-methylnicotinamide;
- 5 5-methoxy-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]-2,4-dimethyl-1,3-thiazole-5-carboxamide;
- 10 N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- N-{8-[3-(dimethylamino)propoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}nicotinamide;
- 15 6-{[(isopropylamino)carbonyl]amino}-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}-2,4-dimethyl-1,3-thiazole-5-carboxamide;
- 20 N-[7-methoxy-8-(2-morpholin-4-ylethoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]nicotinamide;
- rel-6-amino-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)nicotinamide;
- 25 rel-2-amino-N-(8-{3-[(2R,6S)-2,6-dimethylmorpholin-4-yl]propoxy}-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl)pyrimidine-5-carboxamide;
- 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;
- 30 N-{8-[2-(dimethylamino)ethoxy]-7-methoxy-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl}pyrimidine-5-carboxamide;

N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide;

5 or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof.

8. A combination of :

10 a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound according to any one of claims 1 to 7, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ; or a pharmaceutical composition containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

15 and

b) one or more further active agents selected from the group consisting of :

- 20 - a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;
- 25 - a Bcl binding peptide ;
- a Bcl siRNA, such as PNT-2258 ;
- an antisense therapy oligonucleotide, such as BclKlex ; and
- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of
- 30 mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014,

AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

9. The combination according to claim 8, wherein said 2,3-
5 dihydroimidazo[1,2-c]quinazoline compound is 2-amino-N-[7-methoxy-8-(3-morpholin-4-ylpropoxy)-2,3-dihydroimidazo[1,2-c]quinazolin-5-yl]pyrimidine-5-carboxamide.

10 10. The combination according to claim 8 or 9, wherein said further active agent is rapamycin.

11. The combination according to claim 8 or 9, wherein said further active agent is ABT-737.

15

12. A pharmaceutical composition which comprises a combination of :

a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound according to any one of claims 1 to 7, or a physiologically acceptable salt, solvate,
20 hydrate or stereoisomer thereof ; and

b) one or more further active agents selected from the group consisting of :

25 - a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;

30 - a Bcl binding peptide ;

- a Bcl siRNA, such as PNT-2258 ;

- an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001, Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01, PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235.

13. Use of a combination of :

10

a) a 2,3-dihydroimidazo[1,2-c]quinazoline compound according to any one of claims 1 to 7, or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof ;

or of a pharmaceutical composition containing such a compound or a physiologically acceptable salt, solvate, hydrate or stereoisomer thereof,

and

20 b) one or more further active agents selected from the group consisting of :

- a Bcl inhibitor, such as ABT-737, ABT-263 (Navitoclax), EM20-25, YC137, GX-015-070 (Obatoclax), Tetrocarcin A, UCB-1350883, AT-101 ((-)-Gossypol), SPC-2004 (Beclanorsen), IG-105, WL-276, BI-97C1, I-VRL (Immunovivorelbine), DATS (Allitridin), CNDO-103 (Apogossypol), D-G-3139 (Genasense), Evotec, PIB-1402, EU-517 ;

- a Bcl binding peptide ;

- a Bcl siRNA, such as PNT-2258 ;

30 - an antisense therapy oligonucleotide, such as BclKlex ; and

- an inhibitor of the mTOR pathway, such as rapamycin or a rapamycin analogue, such as Rapamycin (Sirolimus), Everolimus (RAD-001,

Afinitor), Zotarolimus (ABT-578, Endeavor), Temisirolimus (CCI-779, Torisel), Ridaforolimus (AP-23576, MK-8669), TAFA-93, or an inhibitor of mTOR kinase, such as WYE-132, OSI-027, INK-128, OSI-027, AZD-2014, AZD-8055, CC-223, ABI-009, EXEL-3885, EXEL-4451, NV-128, OXA-01,
5 PKI-402, SB-2015, WYE-354, KU-0063794, X-387, BEZ-235 ;

or a pharmaceutical composition containing such a combination,

for the preparation of a medicament for the treatment or prophylaxis of
10 cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer.

14. Use of a biomarker such as Bcl expression, HER family expression
15 and/or activation, PIK3CA signaling and / or loss of PTEN for predicting the sensitivity and/or resistance of a patient with cancer, e.g. breast cancer, in particular inflammatory breast cancer, triple negative breast cancer, Her2 receptor positive breast cancer, hormone receptor positive breast cancer, to a compound according to any one of claims 1 to 7, thus
20 providing rationale-based synergistic combination according to any one of claims 8 to 10 to overcome the resistance (patient stratification).

15. A method of determining the level of a component of one or more of Bcl expression, HER family expression and/or activation, PIK3CA
25 signaling and / or loss of PTEN,
wherein in said Bcl expression, said component is Bcl, for example,
wherein in said HER family expression and/or activation, PIK3CA signaling, said component is EGF-R, for example, and
wherein said loss of PTEN, said component is PTEN, for example.

Application number / numéro de demande: 2832120

Page 5 Fig:5

Unscannable items
received with this application
(Request original documents in File Prep. Section on the 10th floor)

Documents reçu avec cette demande ne pouvant être balayés
(Commander les documents originaux dans la section de préparation des dossiers au
10^{ème} étage)

Fig. 1

5

10

15

20




Fig. 2




Fig. 3




Fig. 4






Fig. 6

5

A**B**

1 - DMSO

2 - 0.5 μ M Rapamycin3 - 2 μ M compound A4- 2 μ M compound A + 0.5 μ M Rapamycin

FIGURE 7

IR spectrum of the compound A




FIGURE 8

Raman spectrum of the compound A




FIGURE 9

UV/VIS spectra of the compound A

5




FIGURE 10


 ^1H -NMR Spectrum of the compound A

FIGURE 11


 ^{13}C -NMR Spectrum of the compound A

FIGURE 12


 ^{13}C -NMR Spectra of the compound A

FIGURE 13
Mass Spectrum of the compound A

