



US 20120028872A1

(19) **United States**

(12) **Patent Application Publication**
Perez-Prat Vinuesa et al.

(10) **Pub. No.: US 2012/0028872 A1**

(43) **Pub. Date: Feb. 2, 2012**

(54) **LIQUID DETERGENT COMPOSITION**

Related U.S. Application Data

(76) Inventors: **Eva Maria Perez-Prat Vinuesa**,
Brussel (BE); **Anna Asmanidou**,
Strombeek-Bever (BE); **Karl**
Shiqing Wei, Mason, OH (US);
Mark Francois Theophile Evers,
Strombeek-Bever (BE); **Robby**
Renilde François Keuleers,
Lippelo (BE); **Jean-Luc Philippe**
Bettiol, Brussel (BE)

(60) Provisional application No. 61/368,744, filed on Jul.
29, 2010.

Publication Classification

(51) **Int. Cl.**
C11D 3/60 (2006.01)

(52) **U.S. Cl.** **510/236**; 510/235; 510/237

(57) **ABSTRACT**

A method of cleaning dishware with a liquid detergent composition having a hydrophobic emollient and a crystalline structurant to provide improved hand skin care benefits and superior grease cleaning and/or suds mileage.

(21) Appl. No.: **13/191,489**

(22) Filed: **Jul. 27, 2011**

LIQUID DETERGENT COMPOSITION

CROSS REFERENCE TO RELATED APPLICATION

[0001] This application claims the benefit of U.S. Provisional Application No. 61/368,744, filed Jul. 29, 2010.

FIELD OF INVENTION

[0002] The present invention relates to a method of cleaning dishware with a liquid detergent composition comprising a hydrophobic emollient and a crystalline structurant, to provide improved hand skin care benefits and superior grease cleaning and/or suds mileage.

BACKGROUND OF THE INVENTION

[0003] During the manual dishwashing process, hands of users are exposed to dishwashing detergents. Such compositions comprise surfactants and other ingredients which can damage the skin and/or cause skin irritation and dryness. Some users will therefore feel the need to apply soothing or moisturizing cream in order to restore moisturization, after the dishwashing process.

[0004] EPO410567 and WO2008046778 illustrate liquid dishwashing detergent compositions comprising low levels of emollients to give skin care benefit.

[0005] However, there remains an unmet need for a shelf stable liquid dishwashing composition, that prevents skin damage and even provides superior moisture maintenance, improved skin feel and skin appearance, while not affecting the cleaning and sudsing performance of said liquid dishwashing composition. Indeed, skin care benefits are often obtained at the expense of effective cleaning and/or sudsing thus making it difficult to formulate a liquid dishwashing composition with such benefits.

[0006] The present invention is directed to the delivery of superior skin care to the user's hands, such requiring high levels of emollients. It has been found that a crystalline structurant improves the stability of the liquid dishwashing detergent composition such that higher levels of hydrophobic emollient can be formulated without negatively impacting cleaning.

[0007] The hand dishwashing method of the present invention has been found to provide the right equilibrium in providing improvement in skin care and stability of the composition whilst providing superior grease cleaning, shine and suds on dishware. Indeed, the washing of dishes requires the difficult technical combination of tough grease cleaning whilst at the same time delighting the consumer with improved skin care and suds.

[0008] A further advantage of this invention is that the skin care benefit can be delivered under all conditions typically found during usage of the liquid dishwashing composition. These conditions vary according to the various different methods of washing dishes used by consumers, i.e. from neat application to dilute conditions. The liquid dishwashing detergent composition of the present invention can be used to deliver improved skin feel and skin appearance in the context of manual dish washing operations, wherein the product undergoes a broad range of concentration levels.

SUMMARY OF THE INVENTION

[0009] The present application relates to a method of cleaning dishware with a liquid detergent composition comprising a hydrophobic emollient and a crystalline structurant.

[0010] The present invention further encompasses the use of liquid detergent composition comprising a hydrophobic emollient and a crystalline structurant to improve hand skin care benefits such as hand skin appearance and skin feel.

DETAILED DESCRIPTION OF THE INVENTION

[0011] As used herein "grease" means materials comprising at least in part (i.e., at least about 0.5 wt % by weight of the grease) saturated and unsaturated fats and oils, preferably oils and fats derived from animal sources, such as beef and/or chicken, and/or vegetable sources.

[0012] As used herein "shelf stable" means a neat hand dish liquid cleansing composition that under ambient conditions does not phase separate for at least two weeks, preferably for at least six months, and more preferably never.

[0013] As used herein "hydration" means optimization of the water level in the skin through importing water from outside into the skin.

[0014] As used herein "moisturization" means optimization of the water level in the skin through hydration and/or through minimization of water loss from the skin via water binding, occlusion and/or the improvement of the skin barrier condition.

[0015] As used herein "dishware" refers to a hard surface such as dishes, glasses, pots, pans, baking dishes and flatware made from ceramic, china, metal, glass, plastic (polyethylene, polypropylene, polystyrene, etc.) and wood.

[0016] As used herein "liquid dishwashing detergent composition" refers to those compositions that are employed in manual (i.e. hand) dishwashing. Such compositions are generally high sudsing or foaming in nature and are shelf stable.

[0017] A used herein "hand skin care benefit" means any benefit relating to hand skin appearance (such as smoothness, elasticity and absence of redness), skin feel (such as softness and suppleness), and skin moisture level

[0018] As used herein "suds profile" means amount of sudsing (high or low) and the persistence of sudsing (sustained or prevention) throughout the washing process resulting from the use of the liquid detergent composition of the present composition. Liquid dishwashing detergent compositions require high sudsing and sustained suds. This is particularly important with respect to liquid dishwashing detergent compositions as the consumer uses high sudsing as an indicator of the performance of the detergent composition. Moreover, the consumer in a liquid dishwashing detergent composition also uses the sudsing profile as an indicator that the wash solution still contains active detergent ingredients. The consumer usually renews the wash solution when the sudsing subsides. Thus, a low sudsing liquid dishwashing detergent composition formulation will tend to be replaced by the consumer more frequently than is necessary because of the low sudsing level.

The Process of Cleaning/Treating a Dishware

[0019] The method of cleaning dishware of the present invention surprisingly provides improved skin care benefits, especially sensory benefits such as skin smoothness, softness, suppleness, and improved skin appearance, while maintaining adequate dishware cleaning performance and sudsing profile and the necessary product stability.

[0020] The present invention is directed to a process of cleaning a dishware with a liquid composition comprising the crystalline structurant and hydrophobic emollient described herein. Said process comprises the step of applying said composition onto the dishware surface, typically in diluted or neat form and rinsing or leaving said composition to dry on said surface without rinsing said surface.

[0021] By “in its neat form”, it is meant herein that said liquid composition is applied directly onto the surface to be treated and/or onto a cleaning device or implement such as a dish cloth, a sponge or a dish brush without undergoing any dilution by the user (immediately) prior to the application. By “diluted form”, it is meant herein that said liquid composition is diluted by the user with an appropriate solvent, typically water. By “rinsing”, it is meant herein contacting the dishware cleaned with the process according to the present invention with substantial quantities of appropriate solvent, typically water, after the step of applying the liquid composition herein onto said dishware. By “substantial quantities”, it is meant usually about 5 to about 20 liters.

[0022] In one embodiment of the present invention, the composition herein can be applied in its diluted form. Soiled dishes are contacted with an effective amount, typically from about 0.5 ml to about 20 ml (per about 25 dishes being treated), preferably from about 3 ml to about 10 ml, of the liquid detergent composition of the present invention diluted in water. The actual amount of liquid detergent composition used will be based on the judgment of user, and will typically depend upon factors such as the particular product formulation of the composition, including the concentration of active ingredients in the composition, the number of soiled dishes to be cleaned, the degree of soiling on the dishes, and the like. Generally, from about 0.01 ml to about 150 ml, preferably from about 3 ml to about 40 ml of a liquid detergent composition of the invention is combined with from about 2000 ml to about 20000 ml, more typically from about 5000 ml to about 15000 ml of water in a sink having a volumetric capacity in the range of from about 1000 ml to about 20000 ml, more typically from about 5000 ml to about 15000 ml. The soiled dishes are immersed in the sink containing the diluted compositions then obtained, where contacting the soiled surface of the dish with a cloth, sponge, or similar article cleans them. The cloth, sponge, or similar article may be immersed in the detergent composition and water mixture prior to being contacted with the dish surface, and is typically contacted with the dish surface for a period of time ranged from about 1 to about 10 seconds, although the actual time will vary with each application and user. The contacting of cloth, sponge, or similar article to the dish surface is preferably accompanied by a concurrent scrubbing of the dish surface.

[0023] Another method of the present invention will comprise immersing the soiled dishes into a water bath or held under running water without any liquid dishwashing detergent. A device for absorbing liquid dishwashing detergent, such as a sponge, is placed directly into a separate quantity of undiluted liquid dishwashing composition for a period of time typically ranging from about 1 to about 5 seconds. The

absorbing device, and consequently the undiluted liquid dishwashing composition, is then contacted individually to the surface of each of the soiled dishes to remove said soiling. The absorbing device is typically contacted with each dish surface for a period of time range from about 1 to about 10 seconds, although the actual time of application will be dependent upon factors such as the degree of soiling of the dish. The contacting of the absorbing device to the dish surface is preferably accompanied by concurrent scrubbing.

[0024] Alternatively, the device may be immersed in a mixture of the hand dishwashing composition and water prior to being contacted with the dish surface, said concentrated solution is made by diluting the hand dishwashing composition with water in a small container that can accommodate the cleaning device at ratios ranging from about 95:5 to about 5:95, preferably about 80:20 to about 20:80 and more preferably about 70:30 to about 30:70 of hand dishwashing liquid: water respectively depending upon the user habits and the cleaning task.

Dependent on the geographical use of the composition, the water used in the method of the present invention can have a hardness level of about 2-30 gpg (“gpg” is a measure of water hardness that is well known to those skilled in the art, and it stands for “grains per gallon”).

Liquid Composition

[0025] The composition used in the method of the present invention is formulated as a liquid dishwashing detergent composition comprising a hydrophobic emollient and a crystalline structurant.

[0026] The liquid detergent compositions herein may further contain from about 30% to about 90% by weight of an aqueous liquid carrier in which the other essential and optional composition components are dissolved, dispersed or suspended. Preferably the aqueous liquid carrier will comprise from about 45% to about 70%, more preferably from about 45% to about 65% of the compositions herein described.

[0027] One preferred component of the aqueous liquid carrier is water. The aqueous liquid carrier, however, may contain other materials which are liquid, or which dissolve in the liquid carrier, at room temperature (about 20° C.-about 25° C.) and which may also serve some other function besides that of an inert filler. Such materials can include, for example, hydrotropes and solvents, discussed in more detail below.

[0028] The liquid detergent composition may have any suitable pH. Preferably the pH of the composition is adjusted to between about 4 and about 14. More preferably the composition has pH of between 6 and 13, most preferably between about 6 and about 10. The pH of the composition can be adjusted using pH modifying ingredients known in the art.

The Hydrophobic Emollient

[0029] The liquid detergent composition for the method of the present invention requires one or more hydrophobic emollients. Hydrophobic emollients are ingredients that soften or soothe the skin by slowing the evaporation of water. Hydrophobic emollients form an oily layer on the surface of the skin that slows water loss increasing skin moisture content and skin water holding capacity. Hydrophobic emollients lubricate the skin and enhance skin barrier function improving skin elasticity and appearance.

[0030] Preferably, the liquid detergent comprises high levels of hydrophobic emollient, typically up to about 10% by weight. The hydrophobic emollient is preferably present from about 0.25% to about 10%, more preferably from about 0.3% to about 8%, most preferably from about 0.5% to about 6% by weight of the total composition.

[0031] Hydrophobic emollients suitable for use in the compositions herein are hydrocarbon oils and waxes; silicones; fatty acid derivatives; glyceride esters, di and tri-glycerides, acetoglyceride esters; alkyl and alkenyl esters; cholesterol and cholesterol derivatives; vegetable oils, vegetable oil derivatives, liquid nondigestible oils, or blends of liquid digestible or nondigestible oils with solid polyol polyesters; natural waxes such as lanolin and its derivatives, beeswax and its derivatives, spermaceti, candelilla, and carnauba waxes; phospholipids such as lecithin and its derivatives; sphingolipids such as ceramide; and homologs thereof and mixtures thereof.

[0032] Examples of suitable Hydrocarbon Oils and Waxes include: petrolatum, mineral oil, micro-crystalline waxes, polyalkenes (e.g. hydrogenated and nonhydrogenated polybutene and polydecene), paratrans, cerasin, ozokerite, polyethylene and perhydroqualene. Preferred hydrocarbon oils are petrolatum and/or blends of petrolatum and mineral oil.

[0033] Examples of suitable Silicone Oils include: dimethicone copolyol, dimethylpolysiloxane, diethylpolysiloxane, high molecular weight dimethicone, mixed C₁₋₃₀alkyl polysiloxane, phenyl dimethicone, dimethiconol, and mixtures thereof. More preferred are non-volatile silicones selected from dimethicone, dimethiconol, mixed C₁₋₃₀alkyl polysiloxane, and mixtures thereof.

[0034] Examples of suitable glyceride esters include: castor oil, soy bean oil, derivatized soybean oils such as maleated soy bean oil, safflower oil, cotton seed oil, corn oil, walnut oil, peanut oil, olive oil, cod liver oil, almond oil, avocado oil, vegetable oils and vegetable oil derivatives; coconut oil and derivatized coconut oil, cottonseed oil and derivatized cottonseed oil, jojoba oil, cocoa butter, and the like. Preferred glyceride is castor oil.

[0035] Acetoglyceride esters may also be used, an example being acetylated monoglycerides.

[0036] Preferred hydrophobic emollients are petrolatum, mineral oil and/or blends of petrolatum and mineral oil; triglycerides such as the ones derived from vegetable oils; oily sugar derivatives; beeswax; lanolin and its derivatives including but not restricted to lanolin oil, lanolin wax, lanolin alcohols, lanolin fatty acids, isopropyl lanolate, cetylated lanolin, acetylated lanolin alcohols, lanolin alcohol linoleate, lanolin alcohol riconoleate; ethoxylated lanolin.

[0037] More preferred hydrophobic emollients are petrolatum; blends of petrolatum and mineral oil wherein the ratio petrolatum:mineral oil ranks from about 90:10 to about 50:50, and preferably is about 70:30; vegetable oils and vegetable waxes such as castor oil, and carnauba wax; blends of petrolatum and vegetable oils such as castor oil; oily sugar derivatives such as the ones taught in WO98/16538. WO98/16538 describes cyclic polyol derivatives or reduced saccharide derivatives resulting from about 35% to about 100% of the hydroxyl group of the cyclic polyol or reduced saccharide being esterified and/or etherified and in which at least two or more ester or ether groups are independently attached to a C8 to C22 alkyl or alkenyl chain, that may be linear or branched. In the context of the present invention, the term cyclic polyol encompasses all forms of saccharides. Especially preferred

are monosaccharides and disaccharides. Examples of monosaccharides include xylose, arabinose, galactose, fructose, and glucose. Example of reduced saccharide is sorbitan. Examples of disaccharides are sucrose, lactose, maltose and cellobiose. Sucrose is especially preferred. Particularly preferred are sucrose esters with 4 or more ester groups. These are commercially available under the trade name Sefose® from Procter & Gamble Chemicals, Cincinnati Ohio.

[0038] Even more preferred hydrophobic emollients are petrolatum, mineral oil, Castor oil, natural waxes such as beeswax, carnauba, spermaceti, lanolin and lanolin derivatives such as liquid lanolin or lanolin oil sold by Croda International under the trade name of Fluilan, and lanolin derivatives such as ethoxylated lanolin sold by Croda International under the trade name of Solan E (PEG-75 lanolin).

[0039] Most preferred hydrophobic emollients are petrolatum, mineral oil, Castor oil, and mixtures thereof.


The Crystalline Structurant

[0040] The liquid detergent composition for the method of the present invention requires one or more crystalline structurants. Crystalline structurants are materials that form a thread-like structuring system and/or an insoluble particle network throughout the matrix of the composition. The crystalline structurants may be crystallized in situ within the aqueous liquid matrix of the composition herein or within a pre-mix which is used to form such an aqueous liquid matrix. It has been found that the network generated by the crystalline structurant prevents the hydrophobic emollient droplets from coalescing and phase splitting in the product, thereby providing excellent stability of a hand dishwashing liquid composition. Such stability allows formulating higher levels of hydrophobic emollient without the need to over emulsify the emollient, process that can result in poor release of the hydrophobic emollient upon product usage which prevents the emollient from delivering the desired hand skin care benefit.

[0041] The crystalline structurant will typically be comprised at a level of from about 0.02% to about 5%, preferably about 0.025% to about 3%, more preferably from about 0.05% to about 2%, most preferably from about 0.1% to about 1.5% by weight of the total composition. Preferred crystalline structurants are: Hydroxyl-containing crystalline structuring agents such as a hydroxyl-containing fatty acid, fatty ester or fatty soap wax-like materials or the like such as the ones described in U.S. Pat. No. 6,080,707. Said crystalline hydroxyl-containing structuring agent is insoluble in water under ambient to near ambient conditions.

[0042] The preferred crystalline hydroxyl-containing structuring agent is selected from the group consisting of a structuring agent with formula (I), (II) or mixtures thereof.


Formula (I)




Wherein R¹ is the chemical moiety described below

R¹ is

[0043]




R^2 is R^1 or H; R^3 is R^1 or H; R^4 is independently C_{10} - C_{22} alkyl or alkenyl comprising at least one hydroxyl group;



Formula (II)

wherein: R^7 is R^4 as defined above in (I) and M is Na^+ , K^+ , Mg^{++} or Al^{3+} , or H.

[0044] Some preferred hydroxyl-containing stabilizers include 12-hydroxystearic acid, 9,10-dihydroxystearic acid, tri-9,10-dihydroxystearin and tri-12-hydroxystearin. Tri-12-hydroxystearin is most preferred for use in the liquid compositions herein.



[0045] Castor wax or hydrogenated castor oil is produced by the hydrogenation (saturation of triglyceride fatty acids) of pure castor oil and is mainly composed of tri-12-hydroxystearin. Commercially available, castor oil-based, crystalline, hydroxyl-containing stabilizers include THIXCIN® from Rheox, Inc. (now Elementis).

[0046] It has been found that the crystalline stabilizing thread-like network formed by these stabilizers provides the composition of the present invention with a pseudoplastic or shear thinning rheology profile and with time-dependent recovery of viscosity after shearing (thixotropy).

[0047] Other suitable crystalline structurants are C_{10-22} ethylene glycol fatty acid esters. C_{10-22} ethylene glycol fatty acid esters can be used alone or in combination with other crystalline structurant such as hydrogenated castor oil. Typical examples are monoesters and/or diesters of ethylene glycol, propylene glycol, diethylene glycol, dipropylene glycol, triethylene glycol or tetraethylene glycol with fatty acids containing from about 6 to about 22, preferably from about 12 to about 18 carbon atoms, such as caproic acid, caprylic acid, 2-ethylhexanoic acid, capric acid, lauric acid, isotridecanoic acid, myristic acid, palmitic acid, palmitoleic acid, stearic acid, isostearic acid, oleic acid, elaidic acid, petroselic acid, linoleic acid, linolenic acid, arachic acid, gadoleic acid, behenic acid, erucic acid, and mixtures thereof.

[0048] The ester is preferably a diester, more preferably a C_{14-18} diester, most preferably ethylene glycol distearate. There are several commercial sources for these materials: e.g. PEG6000MS® available from Stepan, Empilan EGDS/A® available from Albright & Wilson; and Euperlan PK900 Benz-W, Euperlan PK 3000 AM, and Euperlan PK 711 produced by Cognis Corp.

[0049] While not being bound by theory, it is believed that the ester forms an insoluble particle network, preferably platelet crystals, that prevents the coalescence of hydrophobic emollient particles, thus preventing phase separation of the product.

[0050] It has surprisingly been found that the stability of compositions used in the method according to the present invention can be further enhanced by using a combination of two different crystalline structurants to provide the insoluble particle network and thread-like network. Hence in a preferred embodiment, the composition will comprise the combination of (a) typically about 0.02%-5%, preferably about 0.02%-3.5%, more preferably about 0.02%-0.8% by weight of the composition of hydrogenated castor oil, and (b) typically about 0.00001%-1.5%, preferably about 0.0001-1%, more preferably about 0.0001-0.5% by weight of the composition of ethylene glycol distearate.

Blends of Hydrophobic Emollient and Crystalline Structurant

[0051] In a preferred embodiment, the liquid detergent composition according to the present invention comprises a blend of specific hydrophobic emollient and of a specific crystalline structurant. Preferably the hydrophobic emollient is a vegetable oil such as castor oil and the crystalline structurant is hydrogenated castor oil. A preferred blend of castor oil and hydrogenated castor oil is commercially available from Vertellus Specialties Inc: Castorlatum®.

Surfactants

[0052] A preferred further ingredient of the composition of the present invention is a surfactant selected from nonionic, anionic, cationic surfactants, amphoteric, zwitterionic, semi-polar nonionic surfactants, and mixtures thereof. Surfactants may be comprised at a level of from about 1.0% to about 50% by weight, preferably from about 5% to about 40% by weight, more preferably about 10% to about 30% by weight and even more preferably from about 5% to about 20% by weight of the liquid detergent composition. Non-limiting examples of suitable surfactants are discussed below.

[0053] In a preferred embodiment, an efficient but mild to hands surfactant system will typically comprise about 4% to about 40%, preferably about 6% to about 32%, more preferably about 11% to about 25%, and most preferably about 11% to about 18% by weight of the total composition of an anionic surfactant and so preferably with no more than about 15%, preferably no more than about 10%, more preferably no more than about 5% by weight of the total composition, of a sulfonate surfactant.

[0054] Suitable anionic surfactants to be used in the compositions and methods of the present invention are sulfate, sulfosuccinates, sulfonate, and/or sulfoacetate; preferably alkyl sulfate and/or alkyl ethoxy sulfates; more preferably a combination of alkyl sulfates and/or alkyl ethoxy sulfates with a combined ethoxylation degree less than about 5, preferably less than about 3, more preferably less than about 2.

[0055] In an alternative embodiment, the surfactant system could be based on high levels of nonionic surfactant (Such as about 10% to about 45%, preferably about 15 to about 40%, more preferably about 20 to about 35% by weight of the total composition), preferably combined with an amphoteric surfactant, and more preferably with a low level of anionic sur-

factant (such as less than 20%, preferably less than 10%, more preferably less than about 5% by weight of the total composition).

Sulphate Surfactants

[0056] Suitable sulphate surfactants for use in the compositions herein include water-soluble salts or acids of C₁₀-C₁₄ alkyl or hydroxyalkyl, sulphate and/or ether sulfate. Suitable counterions include hydrogen, alkali metal cation or ammonium or substituted ammonium, but preferably sodium.

[0057] Where the hydrocarbyl chain is branched, it preferably comprises C₁₋₄ alkyl branching units. The average percentage branching of the sulphate surfactant is preferably greater than about 30%, more preferably from about 35% to about 80% and most preferably from about 40% to about 60% of the total hydrocarbyl chains.

[0058] The sulphate surfactants may be selected from C₈-C₂₀ primary, branched-chain and random alkyl sulphates (AS); C₁₀-C₁₈ secondary (2,3) alkyl sulphates; C₁₀-C₁₈ alkyl alkoxy sulphates (AE_xS) wherein preferably x is from 1-30; C₁₀-C₁₈ alkyl alkoxy carboxylates preferably comprising 1-5 ethoxy units; mid-chain branched alkyl sulphates as discussed in U.S. Pat. No. 6,020,303 and U.S. Pat. No. 6,060,443; mid-chain branched alkyl alkoxy sulphates as discussed in U.S. Pat. No. 6,008,181 and U.S. Pat. No. 6,020,303.

Alkyl Sulfosuccinates—Sulfoacetate

[0059] Other suitable anionic surfactants are alkyl, preferably dialkyl, sulfosuccinates and/or sulfoacetate. The dialkyl sulfosuccinates may be a C₆₋₁₅ linear or branched dialkyl sulfosuccinate. The alkyl moieties may be symmetrical (i.e., the same alkyl moieties) or asymmetrical (i.e., different alkyl moiety.es). Preferably, the alkyl moiety is symmetrical.

Sulfonate Surfactants

[0060] The compositions of the present invention will preferably comprise no more than about 15% by weight, preferably no more than about 10%, even more preferably no more than about 5% by weight of the total composition, of a sulfonate surfactant. These include water-soluble salts or acids of C₁₀-C₁₄ alkyl or hydroxyalkyl, sulfonates; C₁₁-C₁₈ alkyl benzene sulfonates (LAS), modified alkylbenzene sulfonate (MLAS) as discussed in WO 99/05243, WO 99/05242, WO 99/05244, WO 99/05082, WO 99/05084, WO 99/05241, WO 99/07656, WO 00/23549, and WO 00/23548; methyl ester sulfonate (MES); and alpha-olefin sulfonate (AOS). These also include the paraffin sulfonates may be monosulfonates and/or disulfonates, obtained by sulphonating paraffins of 10 to 20 carbon atoms. The sulfonate surfactants also include the alkyl glyceryl sulfonate surfactants.

Amphoteric and Zwitterionic Surfactants


[0061] The amphoteric and zwitterionic surfactant may be comprised at a level of from about 0.01% to about 20%, preferably from about 0.2% to about 15%, more preferably about 0.5% to about 12% by weight of the liquid detergent composition. Suitable amphoteric and zwitterionic surfactants are amine oxides and betaines.

[0062] Most preferred are amine oxides, especially coco dimethyl amine oxide or coco amido propyl dimethyl amine oxide. Amine oxide may have a linear or mid-branched alkyl moiety. Typical linear amine oxides include water-soluble amine oxides of formula R¹-N(R²)(R³)→O, wherein R¹ is a C₈₋₁₈ alkyl moiety; R² and R³ are independently selected from

the group consisting of C₁₋₃ alkyl groups and C₁₋₃ hydroxyalkyl groups and preferably include methyl, ethyl, propyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl and 3-hydroxypropyl. The linear amine oxide surfactants in particular may include linear C₁₀-C₁₈ alkyl dimethyl amine oxides and linear C₈-C₁₂ alkoxy ethyl dihydroxy ethyl amine oxides. Preferred amine oxides include linear C₁₀, linear C₁₀-C₁₂, and linear C₁₂-C₁₄ alkyl dimethyl amine oxides. As used herein "mid-branched" means that the amine oxide has one alkyl moiety having n₁ carbon atoms with one alkyl branch on the alkyl moiety having n₂ carbon-atoms. The alkyl branch is located on the α carbon from the nitrogen on the alkyl moiety. This type of branching for the amine oxide is also known in the art as an internal amine oxide. The total sum of n₁ and n₂ is from 10 to 24 carbon atoms, preferably from 12 to 20, and more preferably from 10 to 16. The number of carbon atoms for the one alkyl moiety (n₁) should be approximately the same number of carbon atoms as the one alkyl branch (n₂) such that the one alkyl moiety and the one alkyl branch are symmetric. As used herein "symmetric" means that |n₁-n₂| is less than or equal to 5, preferably 4, most preferably from 0 to 4 carbon atoms in at least about 50 wt %, more preferably at least about 75 wt % to about 100 wt % of the mid-branched amine oxides for use herein.

[0063] The amine oxide further comprises two moieties, independently selected from a C₁₋₃ alkyl, a C₁₋₃ hydroxyalkyl group, or a polyethylene oxide group containing an average of from about 1 to about 3 ethylene oxide groups. Preferably the two moieties are selected from a C₁₋₃ alkyl, more preferably both are selected as a C₁ alkyl.

[0064] Other suitable surfactants include betaines such alkyl betaines, alkylamidobetaine, amidazoliniumbetaine, sulfobetaine (INCI Sultaines) as well as the Phosphobetaine and preferably meets formula I:



[0065] R¹ is a saturated or unsaturated C₆₋₂₂ alkyl residue, preferably C₈₋₁₈ alkyl residue, in particular a saturated C₁₀₋₁₆ alkyl residue, for example a saturated C₁₂₋₁₄ alkyl residue;

[0066] X is NH, NR⁴ with C₁₋₄ Alkyl residue R⁴, O or S,

[0067] n a number from 1 to 10, preferably 2 to 5, in particular 3,

[0068] x 0 or 1, preferably 1,



[0069] R², R³ are independently a C₁₋₄ alkyl residue, potentially hydroxy substituted such as a hydroxyethyl, preferably a methyl.

[0070] m a number from 1 to 4, in particular 1, 2 or 3,

[0071] y 0 or 1 and

[0072] Y is COO, SO₃, OPO(OR⁵)O or P(O)(OR⁵)O, whereby R⁵ is a hydrogen atom H or a C1-4 alkyl residue.

[0073] Preferred betaines are the alkyl betaines of the formula (Ia), the alkyl amido betaine of the formula (Ib), the sulfo betaines of the formula (Ic) and the amido sulfobetaine of the formula (Id);



in which R¹ has the same meaning as in formula I. Particularly preferred betaines are the Carbobetaine [wherein Y⁻=COO⁻], in particular the carbobetaine of the formula (Ia) and (Ib), more preferred are the alkylamidobetaine of the formula (Ib).

[0074] Examples of suitable betaines and sulfobetaine are the following [designated in accordance with INCI]: Almondamidopropyl of betaines, Apricotamidopropyl betaines, Avocamidopropyl of betaines, Babassuamidopropyl of betaines, Behenamidopropyl betaines, Behenyl of betaines, betaines, Canolamidopropyl betaines, Capryl/Capramidopropyl betaines, Carnitine, Cetyl of betaines, Cocamidoethyl of betaines, Cocamidopropyl betaines, Cocamidopropyl Hydroxysultaine, Coco betaines, Coco Hydroxysultaine, Coco/Oleamidopropyl betaines, Coco Sultaine, Decyl of betaines, Dihydroxyethyl Oleyl Glycinate, Dihydroxyethyl Soy Glycinate, Dihydroxyethyl Stearyl Glycinate, Dihydroxyethyl Tallow Glycinate, Dimethicone Propyl of PG-betaines, Erucamidopropyl Hydroxysultaine, Hydrogenated Tallow of betaines, Isostearamidopropyl betaines, Lauramidopropyl betaines, Lauryl of betaines, Lauryl Hydroxysultaine, Lauryl Sultaine, Milkamidopropyl betaines, Minkamidopropyl of betaines, Myristamidopropyl betaines, Myristyl of betaines, Oleamidopropyl betaines, Oleamidopropyl Hydroxysultaine, Oleyl of betaines, Olivamidopropyl of betaines, Palmamidopropyl betaines, Palm itam idopropyl betaines, Palmitoyl Carnitine, Palm Kernelamidopropyl betaines, Polytetrafluoroethylene Acetoxypopyl of betaines, Ricinoleamidopropyl betaines, Sesam idopropyl betaines, Soyamidopropyl betaines, Stearamidopropyl betaines, Stearyl of betaines, Tallowamidopropyl betaines, Tallowamidopropyl Hydroxysultaine, Tallow of betaines, Tallow Dihydroxyethyl of betaines, Undecylenamidopropyl betaines and Wheat Germamidopropyl betaines.

[0075] A preferred betaine is, for example, Cocoamidopropyl betaine (Cocoamidopropyl betaine).

[0076] A preferred surfactant system is a mixture of anionic surfactant and amphoteric or zwitterionic surfactants in a ratio within the range of about 1:1 to about 5:1, preferably from about 1:1 to about 3.5:1.

[0077] It has been found that such surfactant system will provide the excellent cleaning and suds profile required from a hand dishwashing liquid composition while being mild to the hands.

Nonionic Surfactants

[0078] Nonionic surfactant, when present as co-surfactant, is comprised in a typical amount of from about 0.1% to about 20%, preferably about 0.5% to about 15%, more preferably from about 0.5% to about 10% by weight of the liquid detergent composition. When present as main surfactant, it is comprised in a typical amount of from about 10 to about 45%, preferably about 15 to about 40%, more preferably about 20 to about 35% by weight of the total composition. Suitable nonionic surfactants include the condensation products of aliphatic alcohols with from about 1 to 25 moles of ethylene oxide. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from 8 to 22 carbon atoms. Particularly preferred are the condensation products of alcohols having an alkyl group containing from 10 to 18 carbon atoms, preferably from 10 to 15 carbon atoms with from 2 to 18 moles, preferably 2 to 15, more preferably 5-12 moles of ethylene oxide per mole of alcohol.

[0079] Also suitable are alkylpolyglycosides having the formula R²O(C_nH_{2n}O)_t(glycosyl)_x (formula (III)), wherein R² of formula (III) is selected from the group consisting of alkyl, alkyl-phenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups' contain from 10 to 18, preferably from 12 to 14, carbon atoms; n of formula (III) is 2 or 3, preferably 2; t of formula (III) is from 0 to 10, preferably 0; and x of formula (III) is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. Also suitable are alkylglycerol ethers and sorbitan esters.


[0080] Also suitable are fatty acid amide surfactants having the formula (IV):



wherein R⁶ of formula (IV) is an alkyl group containing from 7 to 21, preferably from 9 to 17, carbon atoms and each R⁷ of formula (IV) is selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, and -(C₂H₄O)_xH where x of formula (IV) varies from 1 to 3. Preferred amides are C₈-C₂₀ ammonia amides, monoethanolamides, diethanolamides, and isopropanolamides.

Cationic Surfactants

[0081] Cationic surfactants, when present in the composition, are present in an effective amount, more preferably from 0.1% to 20%, by weight of the liquid detergent composition. Suitable cationic surfactants are quaternary ammonium surfactants. Suitable quaternary ammonium surfactants are selected from the group consisting of mono C₆-C₁₆, preferably C₆-C₁₀ N-alkyl or alkenyl ammonium surfactants, wherein the remaining N positions are substituted by methyl, hydroxyethyl or hydroxypropyl groups. Another preferred cationic surfactant is an C₆-C₁₈ alkyl or alkenyl ester of a quaternary ammonium alcohol, such as quaternary chlorine esters. More preferably, the cationic surfactants have the formula (V):



wherein R¹ of formula (V) is C₈-C₁₈ hydrocarbyl and mixtures thereof, preferably, C₈₋₁₄ alkyl, more preferably, C₈, C₁₀ or C₁₂ alkyl, and X⁻ of formula (V) is an anion, preferably, chloride or bromide.

The Cationic Polymer

[0082] In a preferred embodiment, the compositions herein may comprise at least one cationic polymer for further enhanced skin care benefits. Furthermore, it is believed that the interaction of the cationic polymer with the anionic surfactant results in a coacervation which aids hydrophobic emollient deposition and enhances the deposition of the cationic polymer on the skin.

[0083] The cationic polymer will typically be present at a level of from about 0.001% to about 10%, preferably from about 0.01% to about 5%, more preferably from about 0.05% to about 1% by weight of the total composition.

[0084] Suitable cationic polymers for use in current invention contain cationic nitrogen containing moieties such as quaternary ammonium or cationic protonated amino moieties. The average molecular weight of the cationic polymer is between about 5000 to about 10 million, preferably at least about 100000, more preferably at least about 200000, but preferably not more than about 3000000. The polymers also have a cationic charge density ranging from about 0.1 meq/g to about 5 meq/g, preferably at least about 0.2 meq/g, more preferably at least about 0.3 meq/g, at the pH of intended use of the dishwashing liquid formulation. As used herein the “charge density” of the cationic polymers is defined as the number of cationic sites per polymer gram atomic weight (molecular weight), and can be expressed in terms of meq/gram of cationic charge. In general, adjustments of the proportions of amine or quaternary ammonium moieties in the polymer in function of the pH of the liquid dishwashing liquid in the case of amines, will affect the charge density. Any anionic counterions can be used in association with cationic deposition polymers, so long as the polymer remains soluble in water and in the liquid hand dishwashing liquid matrix, and so long that the counterion is physically and chemically stable with the essential components of this liquid hand dishwashing liquid, or do not unduly impair product performance, stability nor aesthetics. Non-limiting examples of such counterions include halides (e.g. chlorine, fluorine, bromine, iodine), sulphate and methylsulfate.

[0085] Specific examples of the water soluble cationized polymer include cationic polysaccharides such as cationized cellulose derivatives, cationized starch and cationized guar gum derivatives. Also included are synthetically derived copolymers such as homopolymers of diallyl quaternary ammonium salts, diallyl quaternary ammonium salt/acrylamide copolymers, quaternized polyvinylpyrrolidone derivatives, polyglycol polyamine condensates, vinylimidazolium trichloride/vinylpyrrolidone copolymers, dimethyldiallylammonium chloride copolymers, vinylpyrrolidone/quaternized dimethylaminoethyl methacrylate copolymers, polyvinylpyrrolidone/alkylamino acrylate copolymers, polyvinylpyrrolidone/alkylamino acrylate/vinylcaprolactam copolymers, vinylpyrrolidone/methacrylamidopropyl trimethylammonium chloride copolymers, alkylacrylamide acrylate alkylaminoalkylacrylamide/polyethylene glycol methacrylate copolymers, adipic acid/dimethylaminohydroxypropyl ethylenetriamine copolymer (“Cartaretin”—product of Sandoz/USA), and optionally quaternized/protonated condensation polymers having at least one heterocyclic end group connected to the polymer backbone through a unit derived from an alkylamide, the connection comprising an optionally substituted ethylene group (as described in WO 2007 098889, pages 2-19)

[0086] Specific non-limiting examples of commercial water soluble cationized polymers described generally above include: “Merquat 550” (a copolymer of acrylamide and diallyl dimethyl ammonium salt—CTFA name: Polyquaternium-7, product of ONDEO-NALCO), “Luviquat FC370” (a copolymer of 1-vinyl-2-pyrrolidone and 1-vinyl-3-methylimidazolium salt—CTFA name: Polyquaternium-16, product of BASF), “Gafquat 755N” (a copolymer of 1-vinyl-2-pyrrolidone and dimethylaminoethyl methacrylate—CTFA name: Polyquaternium-11, product ex ISP), “Polymer KG,” “Polymer JR series” and “Polymer LR series” (salt of a reac-

tion product between trimethyl ammonium substituted epoxide and hydroxyethyl cellulose—CTFA name: Polyquaternium-10, product of Amerchol) and “Jaguar series” (guar hydroxypropyl trimonium chloride, product of Rhodia) or “N-hance series” (guar hydroxypropyl trimonium chloride, product of Aqualon)

[0087] Preferred cationic polymers are cationic polysaccharides, more preferably cationic cellulose derivatives such as the salts of hydroxyethyl cellulose reacted with trimethyl ammonium substituted epoxide, referred to in the industry (CTFA) as Polyquaternium-10, such as the UCARE LR400, or UCARE JR-400 ex Dow Amerchol, even more preferred are cationic guar gum derivatives such as guar hydroxypropyltrimonium chloride, such as the Jaguar series ex Rhodia and N-Hance polymer series available from Aqualon.

Humectant

[0088] A further preferred ingredients are humectants. It has been found that such composition comprising a humectant will provide additional hand skin mildness benefits.

[0089] When present, the humectant will typically be present in the composition of the present invention at a level of from about 0.1% to about 50%, preferably from about 1% to about 20%, more preferably from about 1% to about 10%, even more preferably from about 1% to about 6%, and most preferably from about 2% to about 5% by weight of the total composition.

[0090] Humectants that can be used according to this invention include those substances that exhibit an affinity for water and help enhance the absorption of water onto a substrate, preferably skin. Specific non-limiting examples of particularly suitable humectants include glycerol, diglycerol, polyethyleneglycol (PEG-4), propylene glycol, hexylene glycol, butylene glycol, (di)propylene glycol, glyceryl triacetate, polyalkyleneglycols, and mixtures thereof. Others can be polyethylene glycol ether of methyl glucose, pyrrolidone carboxylic acid (PCA) and its salts, pidolic acid and salts such as sodium pidolate, polyols like sorbitol, xylitol and maltitol, or polymeric polyols like polydextrose or natural extracts like quillaia, or lactic acid or urea. Also included are alkyl polyglycosides, polybetaine polysiloxanes, and mixtures thereof. Additional suitable humectants are polymeric humectants of the family of water soluble and/or swellable polysaccharides such as hyaluronic acid, chitosan and/or a fructose rich polysaccharide which is e.g. available as Fucogel®1000 (CAS-Nr 178463-23-5) by SOLABIA S.

[0091] Humectants containing oxygen atoms are preferred over those containing nitrogen or sulphur atoms. More preferred humectants are polyols or are carboxyl containing such as glycerol, diglycerol, sorbitol, Propylene glycol, Polyethylene Glycol, Butylene glycol; and/or pidolic acid and salts thereof, and most preferred are humectants selected from the group consisting of glycerol (sourced from Procter & Gamble chemicals), sorbitol, sodium lactate, and urea, or mixtures thereof.

Enzymes

[0092] In a preferred embodiment the composition of the present invention may comprise an enzyme such as an amylase, a protease, a cellulase, a mannanase, a pectinase, a xyloglucanase and/or a lipase; preferably an amylase, protease and/or lipase, more preferably a protease.

[0093] It has been found that the combination of a protease and high levels of hydrophobic emollient does provide the desired superior level of cleaning while providing superior hand feel and mildness to the hands, as well as superior moisturization. Indeed, it is believed that protease promotes exfoliation therefore revealing a relatively more moisturized surface for the deposition of the hydrophobic emollient.

[0094] Protease of microbial origin is preferred. Chemically or genetically modified mutants are included. The protease may be a serine protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of neutral or alkaline proteases include:

[0095] (a) subtilisins (EC 3.4.21.62), especially those derived from *Bacillus*, such as *Bacillus lentus*, *B. alkalophilus*, *B. subtilis*, *B. amyloliquefaciens*, *Bacillus licheniformis*, *Bacillus pumilus* and *Bacillus gibsonii*, and *Cellulomonas* described in U.S. Pat. No. 6,312,936 B1, U.S. Pat. No. 5,679,630, U.S. Pat. No. 4,760,025, U.S. Pat. No. 5,030,378, WO 05/052146, DEA6022216A1 and DEA 6022224A1.

[0096] (b) trypsin-like proteases are trypsin (e.g., of porcine or bovine origin) and the *Fusarium* protease described in WO 89/06270.

[0097] (c) metalloproteases, especially those derived from *Bacillus amyloliquefaciens* described in WO 07/044,993A2.

[0098] Preferred proteases for use herein include polypeptides demonstrating at least about 90%, preferably at least about 95%, more preferably at least about 98%, even more preferably at least about 99% and especially about 100% identity with the wild-type enzyme from *Bacillus lentus* or the wild-type enzyme from *Bacillus amyloliquefaciens*, comprising mutations in one or more of the following positions, using the BPN' numbering system and amino acid abbreviations as illustrated in WO00/37627, which is incorporated herein by reference: 3, 4, 68, 76, 87, 99, 101, 103, 104, 118, 128, 129, 130, 159, 160, 167, 170, 194, 199, 205, 217, 222, 232, 236, 245, 248, 252, 256 & 259.

[0099] More preferred proteases are those derived from the BPN' and Carlsberg families, especially the subtilisin BPN' protease derived from *Bacillus amyloliquefaciens*. In one embodiment the protease is that derived from *Bacillus amyloliquefaciens*, comprising the Y217L mutation whose sequence is described in EP342177A2 (pages 4, 5, 21 and 22).

[0100] Preferred commercially available protease enzymes include those sold under the trade names Alcalase®, Savinase®, Primase®, Durazym®, Polarzyme®, Kannase®, Liquanase®, Ovozyme®, Neutrase®, Everlase® and Esperase® by Novozymes A/S (Denmark), those sold under the tradename Maxatase®, Maxacal®, Maxapem®, Properase®, Purafect®, Purafect Prime®, Purafect Ox®, FN3®, FN4®, Excellase® and Purafect OXP® by Genencor International, and those sold under the tradename Opticlean® and Optimase® by Solvay Enzymes. In one aspect, the preferred protease is a subtilisin BPN' protease derived from *Bacillus amyloliquefaciens*, preferably comprising the Y217L mutation, sold under the tradename Purafect Prime®, supplied by Genencor International.

[0101] Enzymes may be incorporated at a level of typically from about 0.00001% to about 1%, preferably at a level of from about 0.0001% to about 0.5%, more preferably at a level of from about 0.0001% to about 0.1% of enzyme protein by weight of the total composition.

[0102] The aforementioned enzymes can be provided in the form of a stabilized liquid or as a protected liquid or encapsulated enzyme. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid or a

protease stabilizer such as 4-formyl phenyl boronic acid according to established methods. Protected liquid enzymes or encapsulated enzymes may be prepared according to the methods disclosed in U.S. Pat. No. 4,906,396, U.S. Pat. No. 6,221,829, U.S. Pat. No. 6,359,031 and U.S. Pat. No. 6,242,405.

Pearlescent Agent

[0103] The composition herein may comprise as an optional ingredient one or more pearlescent agents. Suitable agents are crystalline or glassy solids, transparent or translucent compounds capable of reflecting and refracting light to produce a pearlescent effect. The composition of the present invention can typically comprise either an organic and/or an inorganic pearlescent agent.

[0104] When the composition of the present invention comprises an organic pearlescent agent, it is typically comprised at an active level of from about 0.05% to about 2.0%, preferably from about 0.1% to about 1.0% by weight of the total composition of the 100% active organic pearlescent agents. When the composition of the present invention comprise an inorganic pearlescent agent, it is typically comprised at an active level of from about 0.001% to about 1.0%, preferably from about 0.001% to about 0.3%, and more preferably from about 0.01% to about 0.2% by weight of the composition of the 100% active inorganic pearlescent agents. Inorganic pearlescent agents include aluminosilicates and/or borosilicates. Preferred are the aluminosilicates and/or borosilicates which have been treated to have a very high refractive index, preferably silica, metal oxides, oxychloride coated aluminosilicate and/or borosilicates. More preferably inorganic pearlescent agent is mica, even more preferred titanium dioxide treated mica such as BASF Mearlin Superfine.

[0105] Other commercially available suitable inorganic pearlescent agents are available from Merck under the tradenames Iriodin, Biron, Xirona, Timiron Colorona, Dichrona, Candurin and Ronastar. Other commercially available inorganic pearlescent agent are available from BASF (Engelhard, Mearl) under tradenames Biju, Bi-Lite, Chroma-Lite, Pearl-Glo, Mearlite and from Eckart under the tradenames Prestige Soft Silver and Prestige Silk Silver Star.

[0106] Particle size (measured across the largest diameter of the sphere) of the pearlescent agent is typically below about 200 microns, preferably below about 100 microns, more preferably below about 50 microns.

Cleaning Polymer

[0107] The composition herein may optionally further comprise one or more alkoxylated polyethyleneimine polymer. The composition may comprise from about 0.01% to about 10%, preferably from about 0.01% to about 2%, more preferably from about 0.1% to about 1.5%, even more preferably from about 0.2% to about 1.5% by weight of the total composition of an alkoxylated polyethyleneimine polymer as described on page 2, line 33 to page 5, line 5 and exemplified in examples 1 to 4 at pages 5 to 7 of WO2007/135645 The Procter & Gamble Company.

[0108] The alkoxylated polyethyleneimine polymer of the present composition has a polyethyleneimine backbone having from about 400 to about 10000 weight average molecular weight, preferably from about 400 to about 7000 weight average molecular weight, alternatively from about 3000 to about 7000 weight average molecular weight.

[0109] The alkoxylation of the polyethyleneimine backbone includes: (1) one or two alkoxylation modifications per nitrogen atom, dependent on whether the modification occurs at a internal nitrogen atom or at an terminal nitrogen atom, in the polyethyleneimine backbone, the alkoxylation modification consisting of the replacement of a hydrogen atom by a polyalkoxylene chain having an average of about 1 to about 40 alkoxy moieties per modification, wherein the terminal alkoxy moiety of the alkoxylation modification is capped with hydrogen, a C₁-C₄ alkyl or mixtures thereof; (2) a substitution of one C₁-C₄ alkyl moiety or benzyl moiety and one or two alkoxylation modifications per nitrogen atom, dependent on whether the substitution occurs at a internal nitrogen atom or at an terminal nitrogen atom, in the polyethyleneimine backbone, the alkoxylation modification consisting of the replacement of a hydrogen atom by a polyalkoxylene chain having an average of about 1 to about 40 alkoxy moieties per modification wherein the terminal alkoxy moiety is capped with hydrogen, a C₁-C₄ alkyl or mixtures thereof; or (3) a combination thereof.

[0110] The composition may further comprise the amphiphilic graft polymers based on water soluble polyalkylene oxides (A) as a graft base and sides chains formed by polymerization of a vinyl ester component (B), said polymers having an average of ≤ 1 graft site per 50 alkylene oxide units and mean molar mass Mw of from about 3,000 to about 100,000 described in BASF patent application WO2007/138053 on pages 2 line 14 to page 10, line 34 and exemplified on pages 15-18.

Magnesium Ions

[0111] The optional presence of magnesium ions may be utilized in the detergent composition when the compositions are used in softened water that contains few divalent ions. When utilized, the magnesium ions preferably are added as a hydroxide, chloride, acetate, sulphate, formate, oxide or nitrate salt to the compositions of the present invention. When included, the magnesium ions are present at an active level of from about 0.01% to about 1.5%, preferably from about 0.015% to about 1%, more preferably from about 0.025% to about 0.5%, by weight of the liquid detergent composition.

Solvent

[0112] The present compositions may optionally comprise a solvent. Suitable solvents include C₄₋₁₄ ethers and diethers, glycols, alkoxyated glycols, C₆-C₁₆ glycol ethers, alkoxyated aromatic alcohols, aromatic alcohols, aliphatic branched alcohols, alkoxyated aliphatic branched alcohols, alkoxyated linear C₁-C₅ alcohols, linear C₁-C₅ alcohols, amines, C₈-C₁₄ alkyl and cycloalkyl hydrocarbons and halo-hydrocarbons, and mixtures thereof. When present, the liquid detergent composition will contain from about 0.01% to about 20%, preferably from about 0.5% to about 20%, more preferably from about 1% to about 10% by weight of the liquid detergent composition of a solvent. These solvents may be used in conjunction with an aqueous liquid carrier, such as water, or they may be used without any aqueous liquid carrier being present.

Hydrotrope

[0113] The liquid detergent compositions of the invention may optionally comprise a hydrotrope in an effective amount so that the liquid detergent compositions are appropriately

compatible in water. Suitable hydrotropes for use herein include anionic-type hydrotropes, particularly sodium, potassium, and ammonium xylene sulfonate, sodium, potassium and ammonium toluene sulfonate, sodium potassium and ammonium cumene sulfonate, and mixtures thereof, and related compounds, as disclosed in U.S. Pat. No. 3,915,903. The liquid detergent compositions of the present invention typically comprise from about 0% to about 15% by weight of the total liquid detergent composition of a hydrotrope, or mixtures thereof, preferably from about 1% to about 10%, most preferably from about 3% to about 10% by weight of the total liquid hand dishwashing composition.

Polymeric Suds Stabilizer

[0114] The compositions of the present invention may optionally contain a polymeric suds stabilizer. These polymeric suds stabilizers provide extended suds volume and suds duration of the liquid detergent compositions. These polymeric suds stabilizers may be selected from homopolymers of (N,N-dialkylamino) alkyl esters and (N,N-dialkylamino) alkyl acrylate esters. The weight average molecular weight of the polymeric suds boosters, determined via conventional gel permeation chromatography, is from about 1,000 to about 2,000,000, preferably from about 5,000 to about 1,000,000, more preferably from about 10,000 to about 750,000, more preferably from about 20,000 to about 500,000, even more preferably from about 35,000 to about 200,000. The polymeric suds stabilizer can optionally be present in the form of a salt, either an inorganic or organic salt.

[0115] One preferred polymeric suds stabilizer is (N,N-dimethylamino)alkyl acrylate esters. Other preferred suds boosting polymers are copolymers of hydroxypropylacrylate/dimethyl aminoethylmethacrylate (copolymer of HPA/DMAM).

[0116] When present in the compositions, the polymeric suds booster/stabilizer may be present from about 0.01% to about 15%, preferably from about 0.05% to about 10%, more preferably from about 0.1% to about 5%, by weight of the liquid detergent composition.

[0117] Another preferred class of polymeric suds booster polymers is hydrophobically modified cellulosic polymers having a number average molecular weight (Mw) below about 45,000; preferably between about 10,000 and about 40,000; more preferably between about 13,000 and about 25,000. The hydrophobically modified cellulosic polymers include water soluble cellulose ether derivatives, such as non-ionic and cationic cellulose derivatives. Preferred cellulose derivatives include methylcellulose, hydroxypropyl methylcellulose, hydroxyethyl methylcellulose, and mixtures thereof.

Diamines

[0118] Another optional ingredient of the compositions according to the present invention is a diamine. Since the habits and practices of the users of liquid-detergent compositions show considerable variation, the composition will preferably contain about 0% to about 15%, preferably about 0.1% to about 15%, preferably about 0.2% to about 10%, more preferably about 0.25% to about 6%, more preferably about 0.5% to about 1.5% by weight of said composition of at least one diamine.

[0119] Preferred organic diamines are those in which pK1 and pK2 are in the range of about 8.0 to about 11.5, preferably in the range of about 8.4 to about 11, even more preferably from about 8.6 to about 10.75. Preferred materials include 1,3-bis(methylamine)-cyclohexane (pKa=10 to 10.5), 1,3 propane diamine (pK1=10.5; pK2=8.8), 1,6 hexane diamine (pK1=11; pK2=10), 1,3 pentane diamine (DYTEK EP®) (pK1=10.5; pK2=8.9), 2-methyl 1,5 pentane diamine (DYTEK A®) (pK1=11.2; pK2=10.0). Other preferred materials include primary/primary diamines with alkylene spacers ranging from C₄ to C₈.

Carboxylic Acid

[0120] The liquid detergent compositions according to the present invention may comprise a linear or cyclic carboxylic acid or salt thereof to improve the rinse feel of the composition. The presence of anionic surfactants, especially when present in higher amounts in the region of about 15-35% about by weight of the composition, results in the composition imparting a slippery feel to the hands of the user and the dishware. This feeling of slipperiness is reduced when using the carboxylic acids as defined herein i.e. the rinse feel becomes draggy.

[0121] Carboxylic acids useful herein include C₁₋₆ linear or at least about 3 carbon containing cyclic acids. The linear or cyclic carbon-containing chain of the carboxylic acid or salt thereof may be substituted with a substituent group selected from the group consisting of hydroxyl, ester, ether, aliphatic groups having from 1 to 6, more preferably 1 to 4 carbon atoms, and mixtures thereof.

[0122] Preferred carboxylic acids are those selected from the group consisting of salicylic acid, maleic acid, acetyl salicylic acid, 3 methyl salicylic acid, 4 hydroxy isophthalic acid, dihydroxyfumaric acid, 1,2,4 benzene tricarboxylic acid, pentanoic acid and salts thereof and mixtures thereof. Where the carboxylic acid exists in the salt form, the cation of the salt is preferably selected from alkali metal, alkaline earth metal, monoethanolamine, diethanolamine or triethanolamine and mixtures thereof.

[0123] The carboxylic acid or salt thereof, when present, is preferably present at the level of from about 0.1% to about 5%, more preferably from about 0.2% to about 1% and most preferably from about 0.25% to about 0.5%, by weight of the total composition.

Other Optional Components:

[0124] The liquid detergent compositions herein can further comprise a number of other optional ingredients suitable for use in liquid detergent compositions such as perfume, dyes, opacifiers, enzymes, chelants, pH buffering means and rheology modifiers including those of the polyacrylate, polysaccharide or polysaccharide derivative type and/or a combination of a solvent and a polycarboxylate polymer. A further discussion of acceptable optional ingredients suitable for use in light-duty liquid detergent composition may be found in U.S. Pat. No. 5,798,505.

Thickness of the Composition

[0125] The liquid hand dishwashing compositions herein are preferably thickened and have preferably a viscosity from about 50 to about 5000 centipoises (50-5000 mPa*s), more preferably from about 100 to about 4000 centipoises (100-4000 mPa*s), even more preferably from about 200-3500 centipoises (200-3500 mPa*s), and most preferably from about 400 to about 3000 centipoises (400-3000 mPa*s) at 20 s⁻¹ and 20° C. Viscosity can be determined by conventional methods. Viscosity according to the present invention is measured using an AR 550 rheometer from TA instruments using a plate steel spindle at 40 mm diameter and a gap size of 500 µm. The high shear viscosity at 20 s⁻¹ and low shear viscosity at 0.05 s⁻¹ can be obtained from a logarithmic shear rate sweep from 0.1 s⁻¹ to 25 s⁻¹ in 3 minutes time at 20° C.

[0126] The preferred rheology described therein may be achieved using internal existing structuring with detergent ingredients or by employing an external rheology modifier and/or a crystalline structurant, which provides the composition with a pseudoplastic or shear thinning rheology profile and with time-dependent recovery of viscosity after shearing (thixotropy).

Examples

Liquid Dishwashing Detergent Compositions

[0127]

	Ex. 1	Ex. 2	Ex. 3	Ex. 4	Ex. 5
Alkyl C ₁₀₋₁₆ Ethoxy _{0.2-4} Sulfate	18%	15%	17%	18%	—
Coco amido propyl Betaine	—	—	9%	5%	—
Ethoxylated alkyl alcohol	—	—	1%	—	33%
Di-methyl Coco Amine Oxide	6%	5.5%	—	4%	2%
Alkylpolyglucoside	—	—	—	4%	—
Ethanol	—	—	5%	7%	9%
Polypropyleneglycol	1%	0.8%	—	—	—
Citrate	—	—	0.3%	0.6%	—
NaCl	1.2%	1.0%	—	0.5%	—
Sodium cumene sulfonate	—	—	0.8%	—	3%
Petrolatum	2.1%	1%	7%	0.7%	3%
White mineral oil	0.9%	—	3%	0.3%	1%
Glycerol	—	1%	2%	—	—
cationic polymer (1)	0.2%	0.15%	—	0.3%	—
Purafect Prime™ Protease (ppm) - Genencor	—	50	70	30	—
Ethylene glycol distearate	0.5%	—	—	1%	0.05%
Hydrogenated Castor Oil	0.28%	0.6%	1.5%	—	0.35%
Mica (BASF Mearlin superfine)	—	0.05	0.1	—	0.02
Minors*	Balance to 100% with water				
pH	9	9	6	6	7

-continued

	Ex. 6	Ex. 7	Ex. 8	Ex. 9	Ex. 10
Linear Alkylbenzene Sulfonate	—	—	12%	7%	2%
Alkyl C ₁₀₋₁₆ Ethoxy _{0.2-4} Sulfate	9%	25%	11%	—	—
Paraffin Sulfonate	20%	—	—	—	—
Coco amido propyl Betaine	4%	1.5%	—	—	2%
Ethoxylated alkyl alcohol	6%	0.4%	0.6%	2%	32%
Di-methyl Coco Amine Oxide	—	—	5%	0.5%	—
Ethanol	3%	—	4%	—	8%
Polypropyleneglycol	—	—	—	0.5%	0.3%
Citrate	0.1%	0.5%	0.3%	0.8%	—
NaCl	0.3%	0.6%	0.2%	—	—
Sodium cumene sulfonate	—	—	2%	—	3.5%
Sucrose esters having esterification of 6-8 with C ₁₆ -C ₁₈ fatty acids	3%	0.25%	10%	—	—
PEG-75 lanolin (such as Solan E ® from Croda International	—	2%	—	5%	7%
cationic polymer (2)	0.2%	0.25%	—	0.25%	0.1%
Protease (ppm) Purafect Prime TM from Genencor	—	30	65	100	70
Ethylene glycol distearate	1.2%	—	0.01%	—	0.8%
Hydrogenated Castor Oil	—	0.14%	0.3%	0.1%	—
Mica (BASF Mearlin superfine)	—	0.1%	—	0.05%	—
Minors*	Balance to 100% with water				
pH	7	5.5	7	6	6.5
	Ex. 11	Ex. 12	Ex. 13	Ex. 14	Ex. 15
Linear Alkylbenzene Sulfonate	13%	—	—	—	—
Alkyl C ₁₀₋₁₆ Ethoxy _{0.2-4} Sulfate	5%	18%	17%	4%	3%
Paraffin Sulfonate	—	—	3%	10%	—
Coco amido propyl Betaine	—	—	5%	1%	—
Ethoxylated alkyl alcohol	1.5%	—	1%	0.5%	29%
Di-methyl Coco Amine Oxide	0.5%	6%	2%	1.5%	3%
Ethanol	3%	—	2%	3%	9.5%
Polypropyleneglycol	0.5%	—	1%	—	—
Citrate	0.6%	0.5%	1.5%	—	—
NaCl	0.5%	0.5%	—	1%	—
Sodium cumene sulfonate	—	—	—	—	2.9%
Mixture of hydrogenated castor oil and castor oil (Castorlatum ® from Vertellus)	5%	2.7%	1%	2.25%	3%
Petrolatum	—	—	—	0.75%	—
cationic polymer (3)	0.1%	0.15%	—	0.05%	0.2%
Protease (ppm) Purafect Prime TM from Genencor	50	—	—	90	—
Ethylene glycol distearate	0.4%	0.05%	—	—	1%
Hydrogenated Castor Oil	—	—	—	0.15%	—
Mica (BASF Mearlin superfine)	—	0.025%	—	0.2%	—
Minors*	Balance to 100% with water				
pH	5	9	7.5	7.7	7.5

*Minors: dyes, opacifier, perfumes, preservatives, hydrotropes, processing aids, rheology modifiers and/or stabilizers.

(1) Cationically modified hydroxyethyl cellulose (Polyquaternium-10).

(2) A copolymer of acrylamide and diallyl dimethyl ammonium salt - CTFA name: Polyquaternium-7

(3) Guar hydroxypropyl trimmonium chloride

[0128] The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as “40 mm” is intended to mean “about 40 mm.”

[0129] Every document cited herein, including any cross referenced or related patent or application, is hereby incorporated herein by reference in its entirety unless expressly excluded or otherwise limited. The citation of any document is not an admission that it is prior art with respect to any invention disclosed or claimed herein or that it alone, or in any

combination with any other reference or references, teaches, suggests or discloses any such invention. Further, to the extent that any meaning or definition of a term in this document conflicts with any meaning or definition of the same term in a document incorporated by reference, the meaning or definition assigned to that term in this document shall govern.

[0130] While particular embodiments of the present invention have been illustrated and described, it would be obvious to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the invention. It is therefore intended to cover in the appended claims all such changes and modifications that are within the scope of this invention.

What is claimed is:

1. A method of cleaning dishware comprising the step of applying a liquid detergent composition comprising a hydrophobic emollient and a crystalline structurant, onto said dishware.

2. A method according to claim 1, wherein said crystalline structurant is selected from the group consisting of hydroxyl-containing fatty acid, fatty ester or fatty soap wax-like materials, ethylene glycol fatty acid esters, and mixtures thereof.

3. A method according to claim 1, wherein said hydrophobic emollient is selected from the group consisting of hydrocarbon oils and waxes; silicones; fatty acid derivatives; glyceride esters, di and tri-glycerides, acetoglyceride esters; alkyl and alkenyl esters; cholesterol and cholesterol derivatives; vegetable oils, vegetable oil derivatives, liquid nondigestible oils, or blends of liquid digestible or nondigestible oils with solid polyol polyesters; natural waxes, beeswax and its derivatives, spermaceti, candelilla, and carnauba waxes; phospholipids such as lecithin and its derivatives; sphingolipids such as ceramide; and mixtures thereof.

4. A method according to claim 1, wherein said hydrophobic emollient is selected from the group consisting of hydrocarbon oils and waxes, vegetable oils, natural waxes and mixtures thereof.

5. A method according to claim 1 wherein the crystalline structurant is hydrogenated castor oil and the hydrophobic emollient is selected from the group consisting of petrolatum, castor oil, mineral oil, and mixtures thereof.

6. A method according to claim 1 wherein said hydrophobic emollient is comprised at a level of from about 0.25% to about 10% by weight of the composition.

7. A method according to claim 1 wherein said hydrophobic emollient is comprised at a level of from about 0.3% to about 6% by weight of the composition.

8. A method according to claim 1 wherein said crystalline structurant is comprised at a level of from about 0.02% to about 5% by weight of the composition.

9. A method according to claim 1 wherein the crystalline structurant is a blend of from about 0.02% to about 5% by weight of the composition of hydrogenated castor oil and

from about 0.00001% to about 1.5% by weight of the composition of ethylene glycol distearate.

10. A method according to claim 1 wherein said composition further comprises from about 4% to about 40% by weight of the composition of an anionic surfactant and no more than about 15%, by weight of the composition of a sulfonate surfactant.

11. A method according to claim 1 wherein said composition further comprises from about 0.01% to about 20% by weight of the composition of a surfactant selected from the group consisting of an amphoteric surfactant, a zwitterionic surfactant, and mixtures thereof.

12. A method according to claim 1 wherein said composition comprises a coco dimethyl amine oxide.

13. A method according to claim 1 wherein said composition further comprises from about 0.1% to about 45% by weight of the composition of a nonionic surfactant selected from the group consisting of C₈-C₂₂ aliphatic alcohols with about 1 to about 25 moles of ethylene oxide, alkylpolyglycosides, fatty acid amide surfactants, and mixtures thereof.

14. A method according to claim 1 wherein said composition further comprises an enzyme selected from the group consisting of amylase, protease, lipase and mixtures thereof.

15. A method according to claim 1 wherein said composition further comprises a humectant selected from the group consisting of glycerol, urea and mixtures thereof.

16. A method according to claim 1 wherein said composition further comprises a cationic polymer.

17. A method according to claim 1 wherein said composition further comprises a pearlescent agent.

18. The use of a hydrophobic emollient and a crystalline structurant in a liquid hand dishwashing detergent composition for improved hand skin care benefits, wherein said crystalline structurant is selected from the group consisting of hydroxyl-containing fatty acids, fatty esters, fatty soap wax-like materials, and mixtures thereof.

* * * * *