

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 April 2003 (10.04.2003)

PCT

(10) International Publication Number
WO 03/029459 A2

(51) International Patent Classification⁷: **C12N 15/11**,
A61K 48/00

37083 Göttingen (DE). **RAUHUT, Reinhard** [DE/DE];
Merkelstrasse 16, 37085 Göttingen (DE).

(21) International Application Number: PCT/EP02/10881

(74) Agents: **WEICKMANN, Franz, Albert et al.**; Weickmann & Weickmann, Postfach 860 820, 81635 München (DE).

(22) International Filing Date:
27 September 2002 (27.09.2002)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(26) Publication Language: English

Published:
— without international search report and to be republished upon receipt of that report

(30) Priority Data:
01123453.1 28 September 2001 (28.09.2001) EP
02006712.0 22 March 2002 (22.03.2002) EP
02016772.2 26 July 2002 (26.07.2002) EP

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicant (*for all designated States except US*): **MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V.** [DE/DE]; Hofgartenstrasse 8, 80539 München (DE).

(72) Inventors; and
(75) Inventors/Applicants (*for US only*): **TUSCHL, Thomas** [DE/DE]; Keplerstrasse 9, 37085 Göttingen (DE). **LAGOS-QUINTANA, Mariana** [MX/DE]; Kurze Geismarstrasse 13, 37073 Göttingen (DE). **LENDECKEL, Winfried** [DE/DE]; Blinde Gasse 52, 37318 Hohengandern (DE). **MEYER, Jutta** [DE/DE]; Sternstrasse 16,

WO 03/029459 A2

(54) Title: MICRORNA MOLECULES

(57) Abstract: In *Caenorhabditis elegans*, lin-4 and let-7 encode 22- and 21-nucleotide RNAs, respectively, that function as key regulators of developmental timing. Because the appearance of these short RNAs is regulated during development, they are also referred to as "small temporal RNAs" (stRNAs). We show that many more 21- and 22-nt expressed RNAs, termed microRNAs, (miRNAs), exist in invertebrates and vertebrates, and that some of these novel RNAs, similar to let-7 stRNA, are also highly conserved. This suggests that sequence-specific post-transcriptional regulatory mechanisms mediated by small RNAs are more general than previously appreciated.

- 1 -

MicroRNA molecules

Description

5

The present invention relates to novel small expressed (micro)RNA molecules associated with physiological regulatory mechanisms, particularly in developmental control.

10 In *Caenorhabditis elegans*, *lin-4* and *let-7* encode 22- and 21-nucleotide RNAs, respectively (1, 2), that function as key regulators of developmental timing (3-5). Because the appearance of these short RNAs is regulated during development, they are also referred to as "microRNAs" (miRNAs) or small temporal RNAs (stRNAs) (6). *lin-4* and *let-21* are the only known
15 miRNAs to date.

Two distinct pathways exist in animals and plants in which 21- to 23-nucleotide RNAs function as post-transcriptional regulators of gene expression. Small interfering RNAs (siRNAs) act as mediators of sequence-specific mRNA degradation in RNA interference (RNAi) (7-11) whereas
20 miRNAs regulate developmental timing by mediating sequence-specific repression of mRNA translation (3-5). siRNAs and miRNAs are excised from double-stranded RNA (dsRNA) precursors by Dicer (12, 13, 29), a multidomain RNase III protein, thus producing RNA species of similar size.
25 However, siRNAs are believed to be double-stranded (8, 11, 12), while miRNAs are single-stranded (6).

We show that many more short, particularly 21- and 22-nt expressed RNAs, termed microRNAs (miRNAs), exist in invertebrates and vertebrates,
30 and that some of these novel RNAs, similar to *let-7* RNA (6), are also highly conserved. This suggests that sequence-specific post-transcriptional

- 2 -

regulatory mechanisms mediated by small RNAs are more general than previously appreciated.

The present invention relates to an isolated nucleic acid molecule comprising:

- 5 (a) a nucleotide sequence as shown in Table 1, Table 2, Table 3 or Table 4
- 10 (b) a nucleotide sequence which is the complement of (a),
- (c) a nucleotide sequence which has an identity of at least 80%, preferably of at least 90% and more preferably of at least 99%, to a sequence of (a) or (b) and/or
- 15 (d) a nucleotide sequence which hybridizes under stringent conditions to a sequence of (a), (b) and/or (c).

In a preferred embodiment the invention relates to miRNA molecules and analogs thereof, to miRNA precursor molecules and to DNA molecules encoding miRNA or miRNA precursor molecules.

Preferably the identity of sequence (c) to a sequence of (a) or (b) is at least 90%, more preferably at least 95%. The determination of identity (percent) may be carried out as follows:

$$I = n : L$$

wherein I is the identity in percent, n is the number of identical nucleotides between a given sequence and a comparative sequence as shown in Table 1, Table 2, Table 3 or Table 4 and L is the length of the comparative sequence. It should be noted that the nucleotides A, C, G and U as depicted in Tables 1, 2, 3 and 4 may denote ribonucleotides,

- 3 -

deoxyribonucleotides and/or other nucleotide analogs, e.g. synthetic non-naturally occurring nucleotide analogs. Further nucleobases may be substituted by corresponding nucleobases capable of forming analogous H-bonds to a complementary nucleic acid sequence, e.g. U may be
5 substituted by T.

Further, the invention encompasses nucleotide sequences which hybridize under stringent conditions with the nucleotide sequence as shown in Table 1, Table 2, Table 3 or Table 4, a complementary sequence thereof or a
10 highly identical sequence. Stringent hybridization conditions comprise washing for 1 h in 1 x SSC and 0.1% SDS at 45°C, preferably at 48°C and more preferably at 50°C, particularly for 1 h in 0.2 x SSC and 0.1% SDS.

The isolated nucleic acid molecules of the invention preferably have a
15 length of from 18 to 100 nucleotides, and more preferably from 18 to 80 nucleotides. It should be noted that mature miRNAs usually have a length of 19-24 nucleotides, particularly 21, 22 or 23 nucleotides. The miRNAs, however, may be also provided as a precursor which usually has a length
20 of 50-90 nucleotides, particularly 60-80 nucleotides. It should be noted that the precursor may be produced by processing of a primary transcript which may have a length of >100 nucleotides.

The nucleic acid molecules may be present in single-stranded or double-stranded form. The miRNA as such is usually a single-stranded molecule,
25 while the mi-precursor is usually an at least partially self-complementary molecule capable of forming double-stranded portions, e.g. stem- and loop-structures. DNA molecules encoding the miRNA and miRNA precursor molecules. The nucleic acids may be selected from RNA, DNA or nucleic acid analog molecules, such as sugar- or backbone-modified ribonucleotides or deoxyribonucleotides. It should be noted, however, that other nucleic analogs, such as peptide nucleic acids (PNA) or locked nucleic acids (LNA), are also suitable.
30

- 4 -

In an embodiment of the invention the nucleic acid molecule is an RNA- or DNA molecule, which contains at least one modified nucleotide analog, i.e. a naturally occurring ribonucleotide or deoxyribonucleotide is substituted by a non-naturally occurring nucleotide. The modified nucleotide analog may be located for example at the 5'-end and/or the 3'-end of the nucleic acid molecule.

Preferred nucleotide analogs are selected from sugar- or backbone-modified ribonucleotides. It should be noted, however, that also nucleobase-modified ribonucleotides, i.e. ribonucleotides, containing a non-naturally occurring nucleobase instead of a naturally occurring nucleobase such as uridines or cytidines modified at the 5-position, e.g. 5-(2-amino)propyl uridine, 5-bromo uridine; adenosines and guanosines modified at the 8-position, e.g. 8-bromo guanosine; deaza nucleotides, e.g. 7-deazaadenosine; O- and N-alkylated nucleotides, e.g. N6-methyl adenosine are suitable. In preferred sugar-modified ribonucleotides the 2'-OH-group is replaced by a group selected from H, OR, R, halo, SH, SR, NH₂, NHR, NR₂ or CN, wherein R is C₁-C₆ alkyl, alkenyl or alkynyl and halo is F, Cl, Br or I. In preferred backbone-modified ribonucleotides the phosphoester group connecting to adjacent ribonucleotides is replaced by a modified group, e.g. of phosphothioate group. It should be noted that the above modifications may be combined.

The nucleic acid molecules of the invention may be obtained by chemical synthesis methods or by recombinant methods, e.g. by enzymatic transcription from synthetic DNA-templates or from DNA-plasmids isolated from recombinant organisms. Typically phage RNA-polymerases are used for transcription, such as T7, T3 or SP6 RNA-polymerases.

The invention also relates to a recombinant expression vector comprising a recombinant nucleic acid operatively linked to an expression control sequence, wherein expression, i.e. transcription and optionally further

- 5 -

processing results in a miRNA-molecule or miRNA precursor molecule as described above. The vector is preferably a DNA-vector, e.g. a viral vector or a plasmid, particularly an expression vector suitable for nucleic acid expression in eukaryotic, more particularly mammalian cells. The 5 recombinant nucleic acid contained in said vector may be a sequence which results in the transcription of the miRNA-molecule as such, a precursor or a primary transcript thereof, which may be further processed to give the miRNA-molecule.

10 Further, the invention relates to diagnostic or therapeutic applications of the claimed nucleic acid molecules. For example, miRNAs may be detected in biological samples, e.g. in tissue sections, in order to determine and classify certain cell types or tissue types or miRNA-associated pathogenic disorders which are characterized by differential expression of miRNA- molecules or miRNA-molecule patterns. Further, the developmental stage 15 of cells may be classified by determining temporarily expressed miRNA- molecules.

Further, the claimed nucleic acid molecules are suitable for therapeutic 20 applications. For example, the nucleic acid molecules may be used as modulators or targets of developmental processes or disorders associated with developmental dysfunctions, such as cancer. For example, miR-15 and miR-16 probably function as tumor-suppressors and thus expression or delivery of these RNAs or analogs or precursors thereof to tumor cells may 25 provide therapeutic efficacy, particularly against leukemias, such as B-cell chronic lymphocytic leukemia (B-CLL). Further, miR-10 is a possible regulator of the translation of Hox Genes, particularly Hox 3 and Hox 4 (or Scr and Dfd in Drosophila).

30 In general, the claimed nucleic acid molecules may be used as a modulator of the expression of genes which are at least partially complementary to said nucleic acid. Further, miRNA molecules may act as target for

- 6 -

therapeutic screening procedures, e.g. inhibition or activation of miRNA molecules might modulate a cellular differentiation process, e.g. apoptosis.

Furthermore, existing miRNA molecules may be used as starting materials
5 for the manufacture of sequence-modified miRNA molecules, in order to modify the target-specificity thereof, e.g. an oncogene, a multidrug-resistance gene or another therapeutic target gene. The novel engineered miRNA molecules preferably have an identity of at least 80% to the starting miRNA, e.g. as depicted in Tables 1, 2, 3 and 4. Further, miRNA molecules can be modified, in order that they are symmetrically processed
10 and then generated as double-stranded siRNAs which are again directed against therapeutically relevant targets.

Furthermore, miRNA molecules may be used for tissue reprogramming
15 procedures, e.g. a differentiated cell line might be transformed by expression of miRNA molecules into a different cell type or a stem cell.

For diagnostic or therapeutic applications, the claimed RNA molecules are
20 preferably provided as a pharmaceutical composition. This pharmaceutical composition comprises as an active agent at least one nucleic acid molecule as described above and optionally a pharmaceutically acceptable carrier.

The administration of the pharmaceutical composition may be carried out
25 by known methods, wherein a nucleic acid is introduced into a desired target cell in vitro or in vivo.

Commonly used gene transfer techniques include calcium phosphate,
DEAE-dextran, electroporation and microinjection and viral methods [30,
30 31, 32, 33, 34]. A recent addition to this arsenal of techniques for the introduction of DNA into cells is the use of cationic liposomes [35].

- 7 -

Commercially available cationic lipid formulations are e.g. Tfx 50 (Promega) or Lipofectamin 2000 (Life Technologies).

The composition may be in form of a solution, e.g. an injectable solution, a cream, ointment, tablet, suspension or the like. The composition may be administered in any suitable way, e.g. by injection, by oral, topical, nasal, rectal application etc. The carrier may be any suitable pharmaceutical carrier. Preferably, a carrier is used, which is capable of increasing the efficacy of the RNA molecules to enter the target-cells. Suitable examples of such carriers are liposomes, particularly cationic liposomes.

Further, the invention relates to a method of identifying novel microRNA-molecules and precursors thereof, in eukaryotes, particularly in vertebrates and more particularly in mammals, such as humans or mice. This method comprises: ligating 5'- and 3'-adapter-molecules to the end of a size-fractionated RNA-population, reverse transcribing said adapter-ligated RNA-population, and characterizing said reverse transcribed RNA-molecules, e.g. by amplification, concatamerization, cloning and sequencing.

A method as described above already has been described in (8), however, for the identification of siRNA molecules. Surprisingly, it was found now that the method is also suitable for identifying the miRNA molecules or precursors thereof as claimed in the present application.

Further, it should be noted that as 3'-adaptor for derivatization of the 3'-OH group not only 4-hydroxymethylbenzyl but other types of derivatization groups, such as alkyl, alkyl amino, ethylene glycol or 3'-deoxy groups are suitable.

Further, the invention shall be explained in more detail by the following Figures and Examples:

- 8 -

Figure Legends

Fig. 1A. Expression of *D. melanogaster* miRNAs. Northern blots of total RNA isolated from staged populations of *D. melanogaster* were probed for the indicated miRNAs. The position of 76-nt val-tRNA is also indicated on the blots. 5S rRNA serves as loading control. E, embryo; L, larval stage; P, pupae; A, adult; S2, Schneider-2 cells. It should be pointed out, that S2 cells are polyclonal, derived from an unknown subset of embryonic tissues, and may have also lost some features of their tissue of origin while maintained in culture. miR-3 to miR-6 RNAs were not detectable in S2 cells (data not shown). miR-14 was not detected by Northern blotting and may be very weakly expressed, which is consistent with its cloning frequency. Similar miRNA sequences are difficult to distinguish by Northern blotting because of potential cross-hybridization of probes.

15

Fig. 1B. Expression of vertebrate miRNAs. Northern blots of total RNA isolated from HeLa cells, mouse kidneys, adult zebrafish, frog ovaries, and S2 cells were probed for the indicated miRNAs. The position of 76-nt val-tRNA is also indicated on the blots. 5S rRNA from the preparations of total RNA from the indicated species is also shown. The gels used for probing of miR-18, miR-19a, miR-30, and miR-31 were not run as far as the other gels (see tRNA marker position). miR-32 and miR-33 were not detected by Northern blotting, which is consistent with their low cloning frequency. Oligodeoxynucleotides used as Northern probes were:

25

let-7a, 5' TACTATACAACCTACTACCTCAATTGCC (SEQ ID NO:1);
let-7d, 5' ACTATGCAACCTACTACCTCT (SEQ ID NO:2);
let-7e, 5' ACTATACAACCTCCTACCTCA (SEQ ID NO:3);
D. melanogaster val-tRNA, 5' TGGTGTTCGCCCCGGAA (SEQ ID NO:4);
miR-1, 5' TGGAATGTAAAGAAGTATGGAG (SEQ ID NO:5);
30 miR-2b, 5' GCTCCTCAAAGCTGGCTGTGATA (SEQ ID NO:6);
miR-3, 5' TGAGACACACTTGCCCAGTGA (SEQ ID NO:7);
miR-4, 5' TCAATGGTTGTCTAGCTTTAT (SEQ ID NO:8);

- 9 -

miR-5, 5' CATATCACAAACGATCGTCCCTT (SEQ ID NO:9);
miR-6, 5' AAAAAGAACAGCCACTGTGATA (SEQ ID NO:10);
miR-7, 5' TGGAAGACTAGTGATTTGTTGT (SEQ ID NO:11);
miR-8, 5' GACATCTTACCTGACAGTATT (SEQ ID NO:12);
5 miR-9, 5' TCATACAGCTAGATAACCAAAGA (SEQ ID NO:13);
miR-10, 5' ACAAAATTCGGATCTACAGGGT (SEQ ID NO:14);
miR-11, 5' GCAAGAACTCAGACTGTGATG (SEQ ID NO:15);
miR-12, 5' ACCAGTACCTGATGTAATACTCA (SEQ ID NO:16);
miR-13a, 5' ACTCGTAAAATGGCTGTGATA (SEQ ID NO:17);
10 miR-14, 5' TAGGAGAGAGAAAAAGACTGA (SEQ ID NO:18);
miR-15, 5' TAGCAGCACATAATGGTTGT (SEQ ID NO:19);
miR-16, 5' GCCAATATTACGTGCTGCTA (SEQ ID NO:20);
miR-17, 5' TACAAGTGCCTCACTGCAGTA (SEQ ID NO:21);
miR-18, 5' TATCTGCACTAGATGCACCTTA (SEQ ID NO:22);
15 miR-19a, 5' TCAGTTTGCATAGATTCGCACA (SEQ ID NO:23);
miR-20, 5' TACCTGCACTATAAGCACTTTA (SEQ ID NO:24);
miR-21, 5' TCAACATCAGTCTGATAAGCTA (SEQ ID NO:25);
miR-22, 5' ACAGTTCTCAACTGGCAGCTT (SEQ ID NO:26);
miR-23, 5' GGAAATCCCTGGCAATGTGAT (SEQ ID NO:27);
20 miR-24, 5' CTGTTCTGCTGAACTGAGCCA (SEQ ID NO:28);
miR-25, 5' TCAGACCGAGACAAGTGCATG (SEQ ID NO:29);
miR-26a, 5' AGCCTATCCTGGATTACTTGAA (SEQ ID NO:30);
miR-27; 5' AGCGGAACCTAGCCACTGTGAA (SEQ ID NO:31);
miR-28, 5' CTCAATAGACTGTGAGCTCCTT (SEQ ID NO:32);
25 miR-29, 5' AACCGATTCAGATGGTGCTAG (SEQ ID NO:33);
miR-30, 5' GCTGCAAACATCCGACTGAAAG (SEQ ID NO:34);
miR-31, 5' CAGCTATGCCAGCATCTTGCCT (SEQ ID NO:35);
miR-32, 5' GCAACTTAGTAATGTGCAATA (SEQ ID NO:36);
miR-33, 5' TGCAATGCAACTACAATGCACC (SEQ ID NO:37).

30

Fig. 2. Genomic organization of miRNA gene clusters. The precursor structure is indicated as box and the location of the miRNA within the

- 10 -

precursor is shown in gray; the chromosomal location is also indicated to the right. (A) *D. melanogaster* miRNA gene clusters. (B) Human miRNA gene clusters. The cluster of let-7a-1 and let-7f-1 is separated by 26500 nt from a copy of let-7d on chromosome 9 and 17. A cluster of let-7a-3 and let-7b, separated by 938 nt on chromosome 22, is not illustrated.

Fig. 3. Predicted precursor structures of *D. melanogaster* miRNAs. RNA secondary structure prediction was performed using mfold version 3.1 [28] and manually refined to accommodate G/U wobble base pairs in the helical segments. The miRNA sequence is underlined. The actual size of the stem-loop structure is not known experimentally and may be slightly shorter or longer than represented. Multicopy miRNAs and their corresponding precursor structures are also shown.

Fig. 4. Predicted precursor structures of human miRNAs. For legend, see Fig. 3.

Fig. 5. Expression of novel mouse miRNAs. Northern blot analysis of novel mouse miRNAs. Total RNA from different mouse tissues was blotted and probed with a 5'-radiolabeled oligodeoxynucleotide complementary to the indicated miRNA. Equal loading of total RNA on the gel was verified by ethidium bromide staining prior to transfer; the band representing tRNAs is shown. The fold-back precursors are indicated with capital L. Mouse brains were dissected into midbrain, mb, cortex, cx, cerebellum, cb. The rest of the brain, rb, was also used. Other tissues were heart, ht, lung, lg, liver, lv, colon, co, small intestine, si, pancreas, pc, spleen, sp, kidney, kd, skeletal muscle, sm, stomach, st, H, human Hela SS3 cells. Oligodeoxynucleotides used as Northern probes were:

miR-1a, CTCCATACTTCTTACATTCCA (SEQ ID NO:38);
miR-30b, GCTGAGTGTAGGATGTTACA (SEQ ID NO:39);
miR-30a-s, GCTTCCAGTCGAGGATGTTACA (SEQ ID NO:40);
miR-99b, CGCAAGGTCGGTTCTACGGGTG (SEQ ID NO:41);

- 11 -

miR-101, TCAGTTATCACAGTACTGTA (SEQ ID NO:42);
miR-122a, ACAAACACCATTGTCACACTCCA (SEQ ID NO:43);
miR-124a, TGGCATTCACCGCGTGCCTTA (SEQ ID NO:44);
miR-125a, CACAGGTTAAAGGGTCTCAGGGA (SEQ ID NO:45);
5 miR-125b, TCACAAGTTAGGGTCTCAGGGA (SEQ ID NO:46);
miR-127, AGCCAAGCTCAGACGGATCCGA (SEQ ID NO:47);
miR-128, AAAAGAGACCGGTTCACTCTGA (SEQ ID NO:48);
miR-129, GCAAGCCCAGACCGAAAAAAG (SEQ ID NO:49);
miR-130, GCCCTTTAACATTGCACTC (SEQ ID NO:50);
10 miR-131, ACTTCGGTTATCTAGCTTTA (SEQ ID NO:51);
miR-132, ACGACCATGGCTGTAGACTGTTA (SEQ ID NO:52);
miR-143, TGAGCTACAGTGCTTCATCTCA (SEQ ID NO:53).

15 Fig.6. Potential orthologs of lin-4 stRNA. (A) Sequence alignment of *C. elegans* lin-4 stRNA with mouse miR-125a and miR-125b and the *D. melanogaster* miR-125. Differences are highlighted by gray boxes. (B)
20 Northern blot of total RNA isolated from staged populations of *D. melanogaster*, probed for miR-125. E, embryo; L, larval stage; P, pupae; A, adult; S2, Schneider-2 cells.

Fig. 7. Predicted precursor structures of miRNAs, sequence accession numbers and homology information. RNA secondary structure prediction was performed using mfold version 3.1 and manually refined to accommodate G/U wobble base pairs in the helical segments. Dashes were inserted into the secondary structure presentation when asymmetrically bulged nucleotides had to be accommodated. The excised miRNA sequence is underlined. The actual size of the stem-loop structure is not known experimentally and may be slightly shorter or longer than represented. Multicopy miRNAs and their corresponding precursor structures are also shown. In cases where no mouse precursors were yet deposited in the database, the human orthologs are indicated. miRNAs
25
30

- 12 -

which correspond to *D. melanogaster* or human sequences are included. Published *C. elegans* miRNAs [36, 37] are also included in the table. A recent set of new HeLa cell miRNAs is also indicated [46]. If several ESTs were retrieved for one organism in the database, only those with different precursor sequences are listed. miRNA homologs found in other species are indicated. Chromosomal location and sequence accession numbers, and clusters of miRNA genes are indicated. Sequences from cloned miRNAs were searched against mouse and human in GenBank (including trace data), and against *Fugu rubripes* and *Danio rerio* at www.jgi.doe.gov and www.sanger.ac.uk, respectively.

EXAMPLE 1: MicroRNAs from *D. melanogaster* and human.

We previously developed a directional cloning procedure to isolate siRNAs after processing of long dsRNAs in *Drosophila melanogaster* embryo lysate (8). Briefly, 5' and 3' adapter molecules were ligated to the ends of a size-fractionated RNA population, followed by reverse transcription, PCR amplification, concatamerization, cloning and sequencing. This method, originally intended to isolate siRNAs, led to the simultaneous identification of 14 novel 20- to 23-nt short RNAs which are encoded in the *D. melanogaster* genome and which are expressed in 0 to 2 h embryos (Table 1). The method was adapted to clone RNAs in a similar size range from HeLa cell total RNA (14), which led to the identification of 19 novel human stRNAs (Table 2), thus providing further evidence for the existence of a large class of small RNAs with potential regulatory roles. According to their small size, we refer to these novel RNAs as microRNAs or miRNAs. The miRNAs are abbreviated as miR-1 to miR-33, and the genes encoding miRNAs are named mir-1 to mir-33. Highly homologous miRNAs are classified by adding a lowercase letter, followed by a dash and a number for designating multiple genomic copies of a mir gene.

- 13 -

The expression and size of the cloned, endogenous short RNAs was also examined by Northern blotting (Fig. 1, Table 1 and 2). Total RNA isolation was performed by acid guanidinium thiocyanate-phenol-chloroform extraction [45]. Northern analysis was performed as described [1], except that the total RNA was resolved on a 15% denaturing polyacrylamide gel, transferred onto Hybond-N+ membrane (Amersham Pharmacia Biotech), and the hybridization and wash steps were performed at 50°C. Oligodeoxynucleotides used as Northern probes were 5'-32P-phosphorylated, complementary to the miRNA sequence and 20 to 25 nt in length.

5S rRNA was detected by ethidium staining of polyacrylamide gels prior to transfer. Blots were stripped by boiling in 0.1% aqueous sodium dodecylsulfate/0.1x SSC (15 mM sodium chloride, 1.5 mM sodium citrate, pH 7.0) for 10 min, and were re-probed up to 4 times until the 21-nt signals became too weak for detection. Finally, blots were probed for val-tRNA as size marker.

For analysis of *D. melanogaster* RNAs, total RNA was prepared from different developmental stages, as well as cultured Schneider-2 (S2) cells, which originally derive from 20-24 h *D. melanogaster* embryos [15] (Fig. 1, Table 1). miR-3 to miR-7 are expressed only during embryogenesis and not at later developmental stages. The temporal expression of miR-1, miR-2 and miR-8 to miR-13 was less restricted. These miRNAs were observed at all developmental stages though significant variations in the expression levels were sometimes observed. Interestingly, miR-1, miR-3 to miR-6, and miR-8 to miR-11 were completely absent from cultured Schneider-2 (S2) cells, which were originally derived from 20-24 h *D. melanogaster* embryos [15], while miR-2, miR-7, miR-12, and miR-13 were present in S2 cells, therefore indicating cell type-specific miRNA expression. miR-1, miR-8, and miR-12 expression patterns are similar to those of lin-4 stRNA in *C. elegans*, as their expression is strongly upregulated in larvae and sustained

- 14 -

to adulthood [16]. miR-9 and miR-11 are present at all stages but are strongly reduced in the adult which may reflect a maternal contribution from germ cells or expression in one sex only.

5 The mir-3 to mir-6 genes are clustered (Fig. 2A), and mir-6 is present as triple repeat with slight variations in the mir-6 precursor sequence but not in the miRNA sequence itself. The expression profiles of miR-3 to miR-6 are highly similar (Table 1), which suggests that a single embryo-specific precursor transcript may give rise to the different miRNAs, or that the
10 same enhancer regulates miRNA-specific promoters. Several other fly miRNAs are also found in gene clusters (Fig. 2A).

15 The expression of HeLa cell miR-15 to miR-33 was examined by Northern blotting using HeLa cell total RNA, in addition to total RNA prepared from mouse kidneys, adult zebrafish, *Xenopus laevis* ovary, and *D. melanogaster* S2 cells (Fig. 1B, Table 2). miR-15 and miR-16 are encoded in a gene cluster (Fig. 2B) and are detected in mouse kidney, fish, and very weakly in frog ovary, which may result from miRNA expression in somatic ovary tissue rather than oocytes. miR-17 to miR-20 are also clustered (Fig. 2B),
20 and are expressed in HeLa cells and fish, but undetectable in mouse kidney and frog ovary (Fig. 1, Table 2), and therefore represent a likely case of tissue-specific miRNA expression.

25 The majority of vertebrate and invertebrate miRNAs identified in this study are not related by sequence, but a few exceptions, similar to the highly conserved let-7 RNA [6], do exist. Sequence analysis of the *D. melanogaster* miRNAs revealed four such examples of sequence conservation between invertebrates and vertebrates. miR-1 homologs are encoded in the genomes of *C. elegans*, *C. briggsae*, and humans, and are
30 found in cDNAs from zebrafish, mouse, cow and human. The expression of miR-1 was detected by Northern blotting in total RNA from adult zebrafish and *C. elegans*, but not in total RNA from HeLa cells or mouse kidney

- 15 -

(Table 2 and data not shown). Interestingly, while mir-1 and let-7 are expressed both in adult flies (Fig. 1A) [6] and are both undetected in S2 cells, miR-1 is, in contrast to let-7, undetectable in HeLa cells. This represents another case of tissue-specific expression of a miRNA, and
5 indicates that miRNAs may not only play a regulatory role in developmental timing, but also in tissue specification. miR-7 homologs were found by database searches in mouse and human genomic and expressed sequence tag sequences (ESTs). Two mammalian miR-7 variants are predicted by sequence analysis in mouse and human, and were detected by Northern blotting in HeLa cells and fish, but not in mouse kidney (Table 2). Similarly,
10 we identified mouse and human miR-9 and miR-10 homologs by database searches but only detected mir-10 expression in mouse kidney.

The identification of evolutionary related miRNAs, which have already
15 acquired multiple sequence mutations, was not possible by standard bioinformatic searches. Direct comparison of the *D. melanogaster* miRNAs with the human miRNAs identified an 11-nt segment shared between *D. melanogaster* miR-6 and HeLa miR-27, but no further relationships were detected. One may speculate that most miRNAs only act on a single target
20 and therefore allow for rapid evolution by covariation, and that highly conserved miRNAs act on more than one target sequence, and therefore have a reduced probability for evolutionary drift by covariation [6]. An alternative interpretation is that the sets of miRNAs from *D. melanogaster* and humans are fairly incomplete and that many more miRNAs remain to
25 be discovered, which will provide the missing evolutionary links.

lin-4 and let-7 stRNAs were predicted to be excised from longer transcripts that contain approximately 30 base-pair stem-loop structures [1, 6]. Database searches for newly identified miRNAs revealed that all miRNAs
30 are flanked by sequences that have the potential to form stable stem-loop structures (Fig. 3 and 4). In many cases, we were able to detect the predicted, approximately 70-nt precursors by Northern blotting (Fig. 1).

- 16 -

Some miRNA precursor sequences were also identified in mammalian cDNA (EST) databases [27], indicating that primary transcripts longer than 70-nt stem-loop precursors do also exist. We never cloned a 22-nt RNA complementary to any of the newly identified miRNAs, and it is as yet unknown how the cellular processing machinery distinguishes between the miRNA and its complementary strand. Comparative analysis of the precursor stem-loop structures indicates that the loops adjacent to the base-paired miRNA segment can be located on either side of the miRNA sequence (Fig. 3 and 4), suggesting that the 5' or 3' location of the stem-closing loop is not the determinant of miRNA excision. It is also unlikely that the structure, length or stability of the precursor stem is the critical determinant as the base-paired structures are frequently imperfect and interspersed by less stable, non-Watson-Crick base pairs such as G/A, U/U, C/U, A/A, and G/U wobbles. Therefore, a sequence-specific recognition process is a likely determinant for miRNA excision, perhaps mediated by members of the Argonaute (rde-1/ago1/piwi) protein family. Two members of this family, alg-1 and alg-2, have recently been shown to be critical for stRNA processing in *C. elegans* [13]. Members of the Argonaute protein family are also involved in RNAi and PTGS. In *D. melanogaster*, these include argonaute2, a component of the siRNA-endonuclease complex (RISC) [17], and its relative aubergine, which is important for silencing of repeat genes [18]. In other species, these include rde-1, argonaute1, and qde-2, in *C. elegans* [19], *Arabidopsis thaliana* [20], and *Neurospora crassa* [21], respectively. The Argonaute protein family therefore represents, besides the RNase III Dicer [12, 13], another evolutionary link between RNAi and miRNA maturation.

Despite advanced genome projects, computer-assisted detection of genes encoding functional RNAs remains problematic [22]. Cloning of expressed, short functional RNAs, similar to EST approaches (RNomics), is a powerful alternative and probably the most efficient method for identification of such novel gene products [23-26]. The number of functional RNAs has been

- 17 -

widely underestimated and is expected to grow rapidly because of the development of new functional RNA cloning methodologies.

The challenge for the future is to define the function and the potential targets of these novel miRNAs by using bioinformatics as well as genetics, and to establish a complete catalogue of time- and tissue-specific distribution of the already identified and yet to be uncovered miRNAs. lin-4 and let-7 stRNAs negatively regulate the expression of proteins encoded by mRNAs whose 3' untranslated regions contain sites of complementarity to the stRNA [3-5].

Thus, a series of 33 novel genes, coding for 19- to 23-nucleotide microRNAs (miRNAs), has been cloned from fly embryos and human cells. Some of these miRNAs are highly conserved between vertebrates and invertebrates and are developmentally or tissue-specifically expressed. Two of the characterized human miRNAs may function as tumor suppressors in B-cell chronic lymphocytic leukemia. miRNAs are related to a small class of previously described 21- and 22-nt RNAs (lin-4 and let-7 RNAs), so-called small temporal RNAs (stRNAs), and regulate developmental timing in *C. elegans* and other species. Similar to stRNAs, miRNAs are presumed to regulate translation of specific target mRNAs by binding to partially complementary sites, which are present in their 3'-untranslated regions.

Deregulation of miRNA expression may be a cause of human disease, and detection of expression of miRNAs may become useful as a diagnostic. Regulated expression of miRNAs in cells or tissue devoid of particular miRNAs may be useful for tissue engineering, and delivery or transgenic expression of miRNAs may be useful for therapeutic intervention. miRNAs may also represent valuable drug targets itself. Finally, miRNAs and their precursor sequences may be engineered to recognize therapeutic valuable targets.

- 18 -

EXAMPLE 2: miRNAs from mouse.

To gain more detailed insights into the distribution and function of miRNAs in mammals, we investigated the tissue-specific distribution of miRNAs in adult mouse. Cloning of miRNAs from specific tissues was preferred over whole organism-based cloning because low-abundance miRNAs that normally go undetected by Northern blot analysis are identified clonally.

Also, *in situ* hybridization techniques for detecting 21-nt RNAs have not yet been developed. Therefore, 19- to 25-nucleotide RNAs were cloned and sequenced from total RNA, which was isolated from 18.5 weeks old

BL6 mice. Cloning of miRNAs was performed as follows: 0.2 to 1 mg of total RNA was separated on a 15% denaturing polyacrylamide gel and RNA of 19- to 25-nt size was recovered. A 5'-phosphorylated 3'-adapter oligonucleotide (5'-pUUUaaccgcgaattccagx: uppercase, RNA; lowercase,

DNA; p, phosphate; x, 3'-Amino-Modifier C-7, ChemGenes, Ashland, Ma, USA, Cat. No. NSS-1004; SEQ ID NO:54) and a 5'-adapter oligonucleotide (5'-acggaattcctcaactAAA: uppercase, RNA; lowercase, DNA; SEQ ID NO:55) were ligated to the short RNAs. RT/PCR was performed with 3'-

primer (5'-GACTAGCTGGAATTCGCGGTTAAA; SEQ ID NO:56) and 5'- primer (5'-CAGCCAACCGAACCCGAAATTCCCTCACTAAA; SEQ ID NO:57). In order to introduce Ban I restriction sites, a second PCR was performed using the primer pair 5'-CAGCCAACCGAACCCGAAATTCCCTCACTAAA (SEQ ID NO:57) and 5'-GACTAGCTTGGTGCGAACATTGCGCGGTTAAA (SEQ ID NO:56), followed by concatamerization after Ban I digestion and T4 DNA ligation. Concatamers of 400 to 600 basepairs were cut out from 1.5%

agarose gels and recovered by Biotrap (Schleicher & Schuell) electroelution (1x TAE buffer) and by ethanol precipitation. Subsequently, the 3' ends of the concatamers were filled in by incubating for 15 min at 72°C with Taq polymerase in standard PCR reaction mixture. This solution was diluted 3-fold with water and directly used for ligation into pCR2.1 TOPO vectors. Clones were screened for inserts by PCR and 30 to 50 samples were subjected to sequencing. Because RNA was prepared from combining

- 19 -

tissues of several mice, minor sequence variations that were detected multiple times in multiple clones may reflect polymorphisms rather than RT/PCR mutations. Public database searching was used to identify the genomic sequences encoding the approx. 21-nt RNAs. The occurrence of
5 a 20 to 30 basepair fold-back structure involving the immediate upstream or downstream flanking sequences was used to assign miRNAs [36-38].

We examined 9 different mouse tissues and identified 34 novel miRNAs, some of which are highly tissue-specifically expressed (Table 3 and Figure
10 5). Furthermore, we identified 33 new miRNAs from different mouse tissues and also from human Soas-2 osteosarcoma cells (Table 4). miR-1 was previously shown by Northern analysis to be strongly expressed in adult heart, but not in brain, liver, kidney, lung or colon [37]. Here we show that miR-1 accounts for 45% of all mouse miRNAs found in heart,
15 yet miR-1 was still expressed at a low level in liver and midbrain even though it remained undetectable by Northern analysis. Three copies or polymorphic alleles of miR-1 were found in mice. The conservation of tissue-specific miR-1 expression between mouse and human provides additional evidence for a conserved regulatory role of this miRNA. In liver,
20 variants of miR-122 account for 72% of all cloned miRNAs and miR-122 was undetected in all other tissues analyzed. In spleen, miR-143 appeared to be most abundant, at a frequency of approx. 30%. In colon, miR-142-as, was cloned several times and also appeared at a frequency of 30%. In small intestine, too few miRNA sequences were obtained to permit
25 statistical analysis. This was due to strong RNase activity in this tissue, which caused significant breakdown of abundant non-coding RNAs, e.g. rRNA, so that the fraction of miRNA in the cloned sequences was very low. For the same reason, no miRNA sequences were obtained from pancreas.

30

To gain insights in neural tissue miRNA distribution, we analyzed cortex, cerebellum and midbrain. Similar to heart, liver and small intestine, variants

- 20 -

of a particular miRNA, miR-124, dominated and accounted for 25 to 48% of all brain miRNAs. miR-101, -127, -128, -131, and -132, also cloned from brain tissues, were further analyzed by Northern blotting and shown to be predominantly brain-specific. Northern blot analysis was performed as
5 described in Example 1. tRNAs and 5S rRNA were detected by ethidium staining of polyacrylamide gels prior to transfer to verify equal loading. Blots were stripped by boiling in deionized water for 5 min, and reprobed up to 4 times until the 21-nt signals became too weak for detection.

10 miR-125a and miR-125b are very similar to the sequence of *C. elegans* lin-4 stRNA and may represent its orthologs (Fig. 6A). This is of great interest because, unlike let-7 that was readily detected in other species, lin-4 has acquired a few mutations in the central region and thus escaped bioinformatic database searches. Using the mouse sequence miR-125b, we
15 could readily identify its ortholog in the *D. melanogaster* genome. miR-125a and miR-125b differ only by a central diuridine insertion and a U to C change. miR-125b is very similar to lin-4 stRNA with the differences located only in the central region, which is presumed to be bulged out during target mRNA recognition [41]. miR-125a and miR-125b were cloned
20 from brain tissue, but expression was also detected by Northern analysis in other tissues, consistent with the role for lin-4 in regulating neuronal remodeling by controlling lin-14 expression [43]. Unfortunately, orthologs to *C. elegans* lin-14 have not been described and miR-125 targets remain to be identified in *D. melanogaster* or mammals. Finally, miR-125b
25 expression is also developmentally regulated and only detectable in pupae and adult but not in embryo or larvae of *D. melanogaster* (Fig. 6B).

Sequence comparison of mouse miRNAs with previously described miRNA reveals that miR-99b and miR-99a are similar to *D. melanogaster*, mouse
30 and human miR-10 as well as *C. elegans* miR-51 [36], miR-141 is similar to *D. melanogaster* miR-8 , miR-29b is similar to *C. elegans* miR-83 , and miR-131 and miR-142-s are similar to *D. melanogaster* miR-4 and *C.*

- 21 -

5 *elegans* miR-79 [36]. miR-124a is conserved between invertebrates and vertebrates. In this respect it should be noted that for almost every miRNA cloned from mouse was also encoded in the human genome, and frequently detected in other vertebrates, such as the pufferfish, *Fugu rubripes*, and the zebrafish, *Danio rerio*. Sequence conservation may point to conservation in function of these miRNAs. Comprehensive information about orthologous sequences is listed in Fig. 7.

10 In two cases both strands of miRNA precursors were cloned (Table 3), which was previously observed once for a *C. elegans* miRNA [36]. It is thought that the most frequently cloned strand of a miRNA precursor represents the functional miRNA, which is miR-30c-s and miR-142-as, s and as indicating the 5' or 3' side of the fold-back structure, respectively.

15 The mir-142 gene is located on chromosome 17, but was also found at the breakpoint junction of a t(8;17) translocation, which causes an aggressive B-cell leukemia due to strong up-regulation of a translocated MYC gene [44]. The translocated MYC gene, which was also truncated at the first exon, was located only 4-nt downstream of the 3'-end of the miR-142 precursor. This suggests that translocated MYC was under the control of the upstream miR-142 promoter. Alignment of mouse and human miR-142 containing EST sequences indicate an approximately 20 nt conserved sequence element downstream of the mir-142 hairpin. This element was lost in the translocation. It is conceivable that the absence of the 20 conserved downstream sequence element in the putative miR-142/mRNA fusion prevented the recognition of the transcript as a miRNA precursor and therefore may have caused accumulation of fusion transcripts and overexpression of MYC.

30 miR-155, which was cloned from colon, is excised from the known noncoding BIC RNA [47]. BIC was originally identified as a gene transcriptionally activated by promoter insertion at a common retroviral

- 22 -

integration site in B cell lymphomas induced by avian leukosis virus. Comparison of BIC cDNAs from human, mouse and chicken revealed 78% identity over 138 nucleotides [47]. The identity region covers the miR-155 fold-back precursor and a few conserved boxes downstream of the 5 fold-back sequence. The relatively high level of expression of BIC in lymphoid organs and cells in human, mouse and chicken implies an evolutionary conserved function, but BIC RNA has also been detected at low levels in non-hematopoietic tissues [47].

10 Another interesting observation was that segments of perfect complementarity to miRNAs are not observed in mRNA sequences or in genomic sequences outside the miRNA inverted repeat. Although this could be fortuitous, based on the link between RNAi and miRNA processing [11, 13, 43] it may be speculated that miRNAs retain the potential to cleave 15 perfectly complementary target RNAs. Because translational control without target degradation could provide more flexibility it may be preferred over mRNA degradation.

In summary, 63 novel miRNAs were identified from mouse and 4 novel 20 miRNAs were identified from human Soas-2 osteosarcoma cells (Table 3 and Table 4), which are conserved in human and often also in other non-mammalian vertebrates. A few of these miRNAs appear to be extremely tissue-specific, suggesting a critical role for some miRNAs in tissue-specification and cell lineage decisions. We may have also identified 25 the fruitfly and mammalian ortholog of *C. elegans* lin-4 stRNA. The establishment of a comprehensive list of miRNA sequences will be instrumental for bioinformatic approaches that make use of completed genomes and the power of phylogenetic comparison in order to identify miRNA-regulated target mRNAs.

- 23 -

References and Notes

1. R. C. Lee, R. L. Feinbaum, V. Ambros, *Cell* 75, 843 (1993).
2. B. J. Reinhart et al., *Nature* 403, 901 (2000).
- 5 3. V. Ambros, *Curr. Opin. Genet. Dev.* 10, 428 (2000).
4. E. G. Moss, *Curr. Biol.* 10, R436 (2000).
5. F. Slack, G. Ruvkun, *Annu. Rev. Genet.* 31, 611 (1997).
6. A. E. Pasquinelli et al., *Nature* 408, 86 (2000).
7. S. M. Elbashir et al., *Nature* 411, 494 (2001).
- 10 8. S. M. Elbashir, W. Lendeckel, T. Tuschl, *Genes & Dev.* 15, 188 (2001).
9. A. J. Hamilton, D. C. Baulcombe, *Science* 286, 950 (1999).
10. S. M. Hammond, E. Bernstein, D. Beach, G. J. Hannon, *Nature* 404, 293 (2000).
- 15 11. P. D. Zamore, T. Tuschl, P. A. Sharp, D. P. Bartel, *Cell* 101, 25 (2000).
12. G. Hutvágner, J. McLachlan, É. Bálint, T. Tuschl, P. D. Zamore, *Science* 93, 834 (2001).
13. A. Grishok et al., *Cell* 106, 23 (2001).
- 20 14. Cloning of 19- to 24-nt RNAs from *D. melanogaster* 0-2 h embryo lysate was performed as described (8). For cloning of HeLa miRNAs, 1 mg of HeLa total RNA was separated on a 15% denaturing polyacrylamide gel and RNA of 19- to 25-nt size was recovered. A 5' phosphorylated 3' adapter oligonucleotide (5' pUUU-aaccgcgaattccagx: uppercase, RNA; lowercase, DNA; p, phosphate; x, 4-hydroxymethylbenzyl; SEQ ID NO:54) and a 5' adapter oligonucleotide (5' acggaattcctcactAAA: uppercase, RNA; lowercase, DNA; SEQ ID NO:55) were ligated to the short HeLa cell RNAs. RT/PCR was performed with 3' primer (5' GACTAGCTGGAATTCGCGGTTAAA; SEQ ID NO:56) and 5' primer (5' CAGCCAACGGAATTCCCTCACTAAA; SEQ ID NO:57), and followed by concatamerization after Eco RI digestion and T4 DNA
- 25
- 30

- 24 -

ligation (8). After ligation of concatamers into pCR2.1 TOPO vectors, about 100 clones were selected and subjected to sequencing.

15. I. Schneider, J Embryol Exp Morphol 27, 353 (1972).
- 5 16. R. Feinbaum, V. Ambros, Dev. Biol. 210, 87 (1999).
17. S. M. Hammond, S. Boettcher, A. A. Caudy, R. Kobayashi, G. J. Hannon, Science 293, 1146 (2001).
18. A. A. Aravin et al., Curr. Biol. 11, 1017 (2001).
19. H. Tabara et al., Cell 99, 123 (1999).
- 10 20. M. Fagard, S. Boutet, J. B. Morel, C. Bellini, H. Vaucheret, Proc. Natl. Acad. Sci. USA 97, 11650 (2000).
21. C. Catalanotto, G. Azzalin, G. Macino, C. Cogoni, Nature 404, 245 (2000).
22. S. R. Eddy, Curr. Opin. Genet. Dev. 9, 695 (1999).
- 15 23. J. Cavaille et al., Proc. Natl. Acad. Sci. USA 97, 14311 (2000).
24. A. Hüttnerhofer et al., EMBO J. 20, 2943 (2001).
25. L. Argaman et al., Curr. Biol. 11, 941 (2001).
26. K. M. Wassarman, F. Repoila, C. Rosenow, G. Storz, S. Gottesman, Genes & Dev. 15, 1637 (2001).
- 20 27. Supplementary Web material is available on Science Online at www.sciencemag.org/cgi/content/full/xxx
28. D. H. Mathews, J. Sabina, M. Zuker, D. H. Turner, J. Mol. Biol. 288, 911 (1999).
29. E. Bernstein, A. A. Caudy, S. M. Hammond, G. J. Hannon, Nature 409, 363 (2001).
- 25 30. Graham, F.L. and van der Eb, A.J., (1973), Virol. 52, 456.
31. McCutchan, J.H. and Pagano, J.S., (1968), J. Natl. Cancer Inst. 41, 351.
32. Chu, G. et al., (1987), Nucl. Acids Res. 15, 1311.
- 30 33. Fraley, R. et al., (1980), J. Biol. Chem. 255, 10431.
34. Capecchi, M.R., (1980), Cell 22, 479.
35. Felgner, P.L. et al., (1987), Proc. Natl. Acad. Sci USA 84, 7413.

- 25 -

36. Lau N.C., Lim L.P., Weinstein E.G., Bartel D.P., (2001), *Science* 294, 858-862.
37. Lee R.C., Ambros V., (2001), *Science* 294, 862-864.
38. Ambros V., (2001), *Cell* 107, 823-826.
- 5 39. Ambros V., Horvitz H.R., (1984), *Science* 226, 409-416.
40. Wightman B., Ha I., Ruvkun G., (1993), *Cell* 75, 855-862.
41. Rougvie A.E., (2001), *Nat. Rev. Genet.* 2, 690-701.
42. Ketting R.F., Fischer S.E., Bernstein E., Sijen T., Hannon G.J., Plasterk R.H., (2001), *Genes & Dev.* 15, 2654-2659.
- 10 43. Hallam S.J., Jin Y., (1998), *Nature* 395, 78-82.
44. Gauwerky C.E., Huebner K., Isobe M., Nowell P.C., Croce C.M., (1989), *Proc. Natl. Acad. Sci. USA* 86, 8867-8871.
45. P. Chomczynski, N. Sacchi, *Anal Biochem* 162, 156, (1987).
46. Mourelatos Z., Dostie J., Paushkin S., Sharma A., Charroux B., Abel 15 L., J.R., Mann M., Dreyfuss G., (2002), *Genes & Dev.*, in press.
47. Tam W., (2001), *Gene* 274, 157-167.

- 26 -

Table 1

D. melanogaster miRNAs. The sequences given represent the most abundant, and typically longest miRNA sequence identified by cloning; miRNAs frequently vary in length by one or two nucleotides at their 3' termini. From 222 short RNAs sequenced, 69 (31%) corresponded to miRNAs, 103 (46%) to already characterized functional RNAs (rRNA, 7SL RNA, tRNAs), 30 (14%) to transposon RNA fragments, and 20 (10%) sequences with no database entry. The frequency (freq.) for cloning a particular miRNA relative to all identified miRNAs is indicated in percent.

5 Results of Northern blotting of total RNA isolated from staged populations of D. melanogaster are summarized. E, embryo; L, larval stage; P, pupae; A, adult; S2, Schneider-2 cells. The strength of the signal within each blot is represented from strongest (++) to undetected (-). let-7 stRNA was probed as control. Genbank accession numbers and homologs of miRNAs

10 identified by database searching in other species are provided as supplementary material.

15

miRNA	sequence (5' to 3')	freq. (%)	E 0-3 h	E 0-6 h	L1+ L2	L3	P	A	S2
miR-1	UGGAUUGUAAGAAGUAUGGAG (SEQ ID NO:58)	32	+	+	++ +	++ +	++	++ +	-
20 miR-2a*	UAUCACAGCCAGCUUUGAUGAGC (SEQ ID NO:59)	3							
miR-2b*	UAUCACAGCCAGCUUUGAGGAGC (SEQ ID NO:60)	3	++	++	++ +	++ +	++	+ +	++ +
25 miR-3	UCACUGGGCAAAGUGUGUCUCA#	9	+++	+++	-	-	-	-	-
miR-4	AUAAAGCUAGACAACCAUUGA (SEQ ID NO:62)	6	+++	+++	-	-	-	-	-
miR-5	AAAGGAACGAUCGUUGUGAU AUG (SEQ ID NO:63)	1	+++	+++	+/-	+/-	-	-	-
miR-6	UAUCACAGUGGCUGUUUCUUUU	13	+++	+++	+/-	+/-	-	-	-
miR-7	UGGAAGACUAGUGAUUUUGUUGU (SEQ ID NO:65)	4	+++	++	+/-	+/-	+/-	+/-	+/-
miR-8	UAAUACUGUCAGGUAAAGAUGUC (SEQ ID NO:66)	3	+/-	+/-	++ +	++ +	+	++ +	-

- 27 -

	miR-9	UCUUJUGGUUAUCUAGCUGUAUGA (SEQ ID NO:67)	7	+++	++	++	++	++	+/-	-
	miR-10	ACCCUGUAGAUCCGAAUUUGU (SEQ ID NO:68)	1	+	+	++	++	+/-	+	-
	miR-11	CAUCACAGUCUGAGUUUCUUGC (SEQ ID NO:69)	7	+++	+++	++	++	++	+	-
	miR-12	UGAGUAUUACAUCAUCAGGUACUGGU (SEQ ID NO:70)	7	+	+	++	++	+	++	+/-
5	miR-13a*	UAUCACAGCCAUUUJUGACGAGU (SEQ ID NO:71)	1	+++	+++	++	++	+	++	++
	miR-13b*	UAUCACAGCCAUUUJUGAUGAGU (SEQ ID NO:72)	0	-	-	-	-	-	-	-
	miR-14	UCAGUCUUUUUCUCUCUCCUA (SEQ ID NO:73)	1	-	-	-	-	-	-	-
	let-7	UGAGGUAGUAGGUUGUAUAGUU (SEQ ID NO:74)	0	-	-	-	-	++	++	-

10 # = (SEQ ID NO:61)

*Similar miRNA sequences are difficult to distinguish by Northern blotting because of potential cross-hybridization of probes.

- 28 -

Table 2

Human miRNAs. From 220 short RNAs sequenced, 100 (45%) corresponded to miRNAs, 53 (24%) to already characterized functional RNAs (rRNA, snRNAs, tRNAs), and 67 (30%) sequences with no database entry. Results of Northern blotting of total RNA isolated from different vertebrate species and S2 cells are indicated. For legend, see Table 1.

miRNA	sequence (5' to 3')	freq. (%)	HeLa cells	mouse kidney	adult fish	frog ovary	S2
let-7a*	UGAGGUAGUAGGUUGGUUAUAGUU#	10	+++	+++	+++	-	-
let-7b*	UGAGGUAGUAGGUUGGUUGGUU (SEQ ID NO:76)	13					
let-7c*	UGAGGUAGUAGGUUGGUUAUGGUU (SEQ ID NO:77)	3					
let-7d*	AGAGGUAGUAGGUUGCAUAGU (SEQ ID NO:78)	2	+++	+++	+++	-	-
let-7e*	UGAGGUAGGAGGUUGGUUAUAGU (SEQ ID NO:79)	2	+++	+++	+++	-	-
let-7f*	UGAGGUAGUAGAUUGGUUAUAGUU (SEQ ID NO:80)	1					
15 miR-15	UAGCAGCACAUAAUGGUUUGUG (SEQ ID NO:81)	3	+++	++	+	+/-	-
miR-16	UAGCAGCACGUAAAUAUUGGCG (SEQ ID NO:82)	10	+++	+	+/-	+/-	-
miR-17	ACUGCAGUGAAGGCACUUGU (SEQ ID NO:83)	1	+++	-	-	-	-
miR-18	UAAGGUGCAUCUAGUGCAGAUA (SEQ ID NO:84)	2	+++	-	-	-	-
miR-19a*	UGUGCAAAUCUAUGCAAAACUGA (SEQ ID NO:85)	1	+++	-	+/-	-	-
20 miR-19b*	UGUGCAAAUCUAUGCAAAACUGA (SEQ ID NO:86)	3					
miR-20	UAAAGUGCUUAUAGUGCAGGU (SEQ ID NO:87)	4	+++	-	+	-	-
miR-21	UAGCUUAUCAGACUGAUGUJGA (SEQ ID NO:88)	10	+++	+	++	-	-
miR-22	AAGCUGCAGUUGAAGAACUGU (SEQ ID NO:89)	10	+++	+++	+	+/-	-
miR-23	AUCACAUUGCAGGGAUUCC (SEQ ID NO:90)	2	+++	+++	+++	+	-

- 29 -

	miR-24	UGGCUCAGUUCAGCAGGAACAG (SEQ ID NO:91)	4	++	+++	++	-	-
	miR-25	CAUUGCACUUJGUCUCGGUCUGA (SEQ ID NO:92)	3	+++	+	++	-	-
	miR-26a*	UUCAAGUAUUCAGGAUAGGU (SEQ ID NO:93)	2	+	++	+++	-	-
	miR-26b*	UUCAAGUAUUCAGGAUAGGU (SEQ ID NO:94)	1					-
5	miR-27	UUCACAGUGGCUAAGUUCCGCU (SEQ ID NO:95)	2	+++	+++	++	-	-
	miR-28	AAGGAGCUCACAGUCUAUJGAG (SEQ ID NO:96)	2	+++	+++	-	-	-
	miR-29	CUAGCACCAUCUGAAAUCGGUU (SEQ ID NO:97)	2	+	+++	+/-	-	-
	miR-30	CUUUCAGUCGGAUGUUUUGCAGC (SEQ ID NO:98)	2	+++	+++	+++	-	-
	miR-31	GGCAAGAUGCGUGGCAUAGCUG (SEQ ID NO:99)	2	+++	-	-	-	-
10	miR-32	UAUUGCACAUUACUAAGUUGC (SEQ ID NO:100)	1	-	-	-	-	-
	miR-33	GUGCAUUGUAGUUGCAUUG (SEQ ID NO:101)	1	-	-	-	-	-
	miR-1	UGGA AUGUAAGAAGUAUGGAG (SEQ ID NO:102)	0	-	-	+	-	-
	miR-7	UGGAAGACUAGUGAUUUJGUUGU (SEQ ID NO:103)	0	+	-	+/-	-	+/-
	miR-9	UCUUUGGUUAUCUAGCUGUAUGA (SEQ ID NO:104)	0	-	-	-	-	-
15	miR-10	ACCCUGUAGAUC CGAAUJGU (SEQ ID NO:105)	0	-	+	-	-	-

= (SEQ ID NO:75)

*Similar miRNA sequences are difficult to distinguish by Northern

20 blotting because of potential cross-hybridization of probes.

- 30 -

Table 3

Mouse miRNAs. The sequences indicated represent the longest miRNA sequences identified by cloning. The 3'-terminus of miRNAs is often truncated by one or two nucleotides. miRNAs that are more than 85% identical in sequence (i.e. share 18 out of 21 nucleotides) or contain 1- or 2-nucleotide internal deletions are referred to by the same gene number followed by a lowercase letter. Minor sequence variations between related miRNAs are generally found near the ends of the miRNA sequence and are thought to not compromise target RNA recognition. Minor sequence variations may also represent A to G and C to U changes, which are accommodated as G-U wobble base pairs during target recognition. miRNAs with the suffix -s or -as indicate RNAs derived from either the 5'-half or the 3'-half of a miRNA precursor. Mouse brains were dissected into midbrain, mb, cortex, cx, cerebellum, cb. The tissues analyzed were heart, ht; liver, lv; small intestine, si; colon, co; cortex, ct; cerebellum, cb; midbrain, mb.

	miRNA	sequence (5' to 3')	Number of clones							
			ht	lv	sp	si	co	cx	cb	mb
20	let-7a	UGAGGUAGUAGGUUGUAUAGUU (SEQ ID NO:106)		3			1	1		7
	let-7b	UGAGGUAGUAGGUUGUGUGGUU (SEQ ID NO:107)		1	1				2	5
	let-7c	UGAGGUAGUAGGUUGUAUGGUU (SEQ ID NO:108)		2			2	5	19	
	let-7d	AGAGGUAGUAGGUUGCAUAGU (SEQ ID NO:109)	2			2	2			2
25	let-7e	UGAGGUAGGAGGUUGUAUAGU (SEQ ID NO:110)				1				2
	let-7f	UGAGGUAGUAGAUUGUAUAGUU (SEQ ID NO:111)			2			3	3	
	let-7g	UGAGGUAGUAGUUUGUACAGUA (SEQ ID NO:112)					1	1		2
	let-7h	UGAGGUAGUAGUGUGUACAGUU (SEQ ID NO:113)					1	1		

- 31 -

	let-7i	UGAGGUAGUAGUUUGUGCU (SEQ ID NO:114)			1	1		
	miR-1b	UGGAAUGUAAGAAGUAUGUAA (SEQ ID NO:115)	4	2				1
	miR-1c	UGGAAUGUAAGAAGUAUGUAC (SEQ ID NO:116)	7					
	miR-1d	UGGAAUGUAAGAAGUAUGUAUU (SEQ ID NO:117)	16					1
5	miR-9	UCUUUGGUUAUCUAGCUGUAUGA (SEQ ID NO:118)			3	4	4	
	miR-15a	UAGCAGCACAUAAUGGUUJUG (SEQ ID NO:119)	1					2
	miR-15b	UAGCAGCACAUCAUGGUUUACA (SEQ ID NO:120)	1					
	miR-16	UAGCAGCACGUAAAUAUUGGCG (SEQ ID NO:121)	1		1	2	1	2
	miR-18	UAAGGUGCAUCUAGUGCAGAUA (SEQ ID NO:122)		1				
10	miR-19b	UGUGCAAAUCCAUGCAAAACUGA (SEQ ID NO:123)		1				
	miR-20	UAAAGUGCUUAUAGUGCAGGUAG (SEQ ID NO:124)			1			
	miR-21	UAGCUUAUCAGACUGAUGUUGA (SEQ ID NO:125)	1		1	2	1	
	miR-22	AAGCUGCCAGUUGAAGAACUGU (SEQ ID NO:126)	2	1		1		1
	miR-23a	AUCACAUUGCAGGGAUUUCC (SEQ ID NO:127)	1					2
15	miR-23b	AUCACAUUGCAGGGAUUACCAC (SEQ ID NO:128)			1			
	miR-24	UGGCUCAGUUCAGCAGGAACAG (SEQ ID NO:129)	1		1	1		1
	miR-26a	UUCAAGUAAUCCAGGAUAGGCCU (SEQ ID NO:130)					3	2
	miR-26b	UUCAAGUAAUUCAGGAUAGGUU (SEQ ID NO:131)		2		4	1	
	miR-27a	UUCACAGUGGCUAAGUUCCGCU (SEQ ID NO:132)	1	2	1	1	2	1
20	miR-27b	UUCACAGUGGCUAAGUUCUG (SEQ ID NO:133)						1
	miR-29a	CUAGCACCAUCUGAAAUCGGUU (SEQ ID NO:134)	1		1		1	
	miR-29b/miR-102	UAGCACCAUJUGAAAUCAGUGUU (SEQ ID NO:135)	1		1	5		3
	miR-29c/	UAGCACCAUJUGAAAUCGGUUA (SEQ ID NO:136)	1		3			1

- 32 -

	miR-30a-s/miR-97	UGUAAAACAUCUCGACUGGAAGC (SEQ ID NO:137)	1	1	1
	miR-30a-as ^a	CUUUCAGUCGGAUGUUUGCAGC (SEQ ID NO:138)			1
	miR-30b	UGUAAAACAUCUACACUCAGC (SEQ ID NO:139)	1		2
	miR-30c	UGUAAAACAUCUACACUCAGC (SEQ ID NO:140)	2		1 1
5	miR-30d	UGUAAAACAUCCCGACUGGAAG (SEQ ID NO:141)	1		
	miR-99a/miR-99	ACCCGUAGAUCCGAUCUUGU (SEQ ID NO:142)		1	
	miR-99b	CACCGUAGAACCGACCUGCG (SEQ ID NO:143)			1
	miR-101	UACAGUACUGUGAUACUGA (SEQ ID NO:144)		2	1 1
	miR-122a	UGGAGUGUGACAAUGGUGUUJGU (SEQ ID NO:145)	3		
10	miR-122b	UGGAGUGUGACAAUGGUGUUJGA (SEQ ID NO:146)	11		
	miR-122a,b	UGGAGUGUGACAAUGGUGUUJUG (SEQ ID NO:147)	23		
	miR-123	CAUUAUUACUUUJGGUACCGG (SEQ ID NO:148)	1 2		
	miR-124a ^b	UUAAGGCACGCCG-UGAAUGC (SEQ ID NO:149)		1	37 41 24
	miR-124b	UUAAGGCACGCCGGUGAAUGC (SEQ ID NO:150)		1	3
15	miR-125a	UCCCUGAGACCUUUACCUGUG (SEQ ID NO:151)		1	1
	miR-125b	UCCCUGAGACCCU--AACUUGUGA (SEQ ID NO:152)		1	
	miR-126	UCGUACCGUGAGUAUAAAUGC (SEQ ID NO:153)	4		1
	miR-127	UCGGAUCCGUCUGAGCUUGGCU (SEQ ID NO:154)			1
	miR-128	UCACAGUGAACCGGUCUCUUU (SEQ ID NO:155)		2 2	2
20	miR-129	CUUUUUUCGGUCUGGGCUUGC (SEQ ID NO:156)			1
	miR-130	CAGUGCAAUGUAAAAGGGC (SEQ ID NO:157)			1
	miR-131	UAAAGCUAGUAACCGAAAGU (SEQ ID NO:158)		1 1	1
	miR-132	UAACAGUCUACAGCCAUGGUCGU (SEQ ID NO:159)			1

- 33 -

	miR-133	UUGGUCCCCUUCAACCAGCUGU (SEQ ID NO:160)	4		1	
	miR-134	UGUGACUGGUUGACCAGAGGGA (SEQ ID NO:161)			1	
	miR-135	UAUGGCCUUUUAUUCCUAUGUGAA (SEQ ID NO:162)			1	
	miR-136	ACUCCAUUJUGUUUUUGAUGAUGGA (SEQ ID NO:163)			1	
5	miR-137	UAUJGCUUAAGAAUACGCGUAG (SEQ ID NO:164)			1	1
	miR-138	AGCUGGGUGUJUGUGAAUC (SEQ ID NO:165)			1	
	miR-139	UCUACAGUGCACGUGUCU (SEQ ID NO:166)			1	1
	miR-140	AGUGGUUUUACCCUAUGGUAG (SEQ ID NO:167)		1		
	miR-141	AACACUGUCUGGUAAAAGAUGG (SEQ ID NO:168)		1	1	1
10	miR-142-s	CAUAAAGUAGAAAGCACUAC (SEQ ID NO:169)			1	1
	miR-142-as ^b	UGUAGUGUUUCCUACUUUAUGG (SEQ ID NO:170)		1	1	6
	miR-143	UGAGAUGAAGCACUGUAGCUCA (SEQ ID NO:171)	3	7		2
	miR-144	UACAGUAUAGAUGAUGUACUAG (SEQ ID NO:172)	2			1
	miR-145	GUCCAGUUUCCAGGAAUCCUU (SEQ ID NO:173)	1			
15	miR-146	UGAGAACUGAAUCCAUGGGUUU (SEQ ID NO:174)	1			
	miR-147	GUGUGUGAAAUGCUCUGCC (SEQ ID NO:175)		1		
	miR-148	UCAGUGCACUACAGAACUUUGU (SEQ ID NO:176)		1		
	miR-149	UCUGGCUCCGUGUCUUCACUCC (SEQ ID NO:177)	1			
	miR-150	UCUCCCAACCCUUGUACCAGUGU (SEQ ID NO:178)			1	
20	miR-151	CUAGACUGAGGCUCCUUGAGGU (SEQ ID NO:179)			1	
	miR-152	UCAGUGCAUGACAGAACUJUGG (SEQ ID NO:180)			1	
	miR-153	UUGCAUAGUCACAAAAGUGA (SEQ ID NO:181)				1
	miR-154	UAGGUUAUCCGUGUUGCCUUCG (SEQ ID NO.182)				1

- 34 -

miR-155

UUAAUGCUAAUUGUGAUAGGG
(SEQ ID NO:183)

1

- 5 ^aThe originally described miR-30 was renamed to miR-30a-as in order to distinguish it from the miRNA derived from the opposite strand of the precursor encoded by the mir-30a gene. miR-30a-s is equivalent to miR-97 [46].
- ^bA 1-nt length heterogeneity is found on both 5' and 3' end. The 22-nt miR sequence is shown, but only 21-nt miRNAs were cloned.

- 35 -

Table 4

Mouse and human miRNAs. The sequences indicated represent the longest miRNA sequences identified by cloning. The 3' terminus of miRNAs is often truncated by one or two nucleotides. miRNAs that are more than 85% identical in sequence (i.e. share 18 out of 21 nucleotides) or contain 1- or 2-nucleotide internal deletions are referred to by the same gene number followed by a lowercase letter. Minor sequence variations between related miRNAs are generally found near the ends of the miRNA sequence and are thought to not compromise target RNA recognition. Minor sequence variations may also represent A to G and C to U changes, which are accommodated as G-U wobble base pairs during target recognition. Mouse brains were dissected into midbrain, mb, cortex, cx, cerebellum, cb. The tissues analyzed were lung, ln; liver, lv; spleen, sp; kidney, kd; skin, sk; testis, ts; ovary, ov; thymus, thy; eye, ey; cortex, ct; cerebellum, cb; midbrain, mb. The human osteosarcoma cells SAOS-2 cells contained an inducible p53 gene (p53-, uninduced p53; p53+, induced p53); the differences in miRNAs identified from induced and uninduced SAOS cells were not statistically significant.

5	miRNA	Sequence (5' to 3')	number of clones										
			mouse tissues					human SAOS-2 cells					
			In	lv	sp	kd	sk	ts	ov	thy	ey	p53-	p53+
10	miR-C1	AACAUUCAACGGCUGUCGGUGAGU	1				1				2		(SEQ ID NO.184)
	miR-C2	UUUGGCAAUGGUAGAACUCACA									1		(SEQ ID NO.185)
	miR-C3	UAUGGCACUGGGUAGAAUUCACUG									1		(SEQ ID NO.186)
	miR-C4	CUDUUUGGGCUCUGGGCUUGUU					1				1		(SEQ ID NO.187)
	miR-C5	UGGACGGAGAACUGAUAGGGU									2		(SEQ ID NO.188)
	miR-C6	UGGAGAGAAAAGGCAGUTUC									1		(SEQ ID NO.189)
	miR-C7	CAAAGAAUUCUCCUUUGGGCUU									1		(SEQ ID NO.190)
	miR-C8	UCCGGUCUUGGGUGCGAGCCGG									1		(SEQ ID NO.191)
	miR-C9	UAAACACUGUCUGGGUAAACGAUG									1		(SEQ ID NO.192)
	miR-C10	CAUCCCCUUGCAUGGGGGGGU									1		(SEQ ID NO.193)
	miR-C11	GUGCCUACUGAGCUGACAUCAUCAGU									1		(SEQ ID NO.194)
	miR-C12	UGGAAUAGUUUUGAUAAUATJAGGU									2		(SEQ ID NO.195)
20	miR-C13	CAACGGAAUCCAAAAGCAGCU									2	1	(SEQ ID NO.196)
	miR-C14	CUGACCUAUGGAAUUGACAA									2	1	(SEQ ID NO.197)

		(SEQ ID NO.198)
		(SEQ ID NO.199)
		(SEQ ID NO.200)
		(SEQ ID NO.201)
		(SEQ ID NO.202)
		(SEQ ID NO.203)
		(SEQ ID NO.204)
		(SEQ ID NO.205)
		(SEQ ID NO.206)
		(SEQ ID NO.207)
		(SEQ ID NO.208)
		(SEQ ID NO.209)
		(SEQ ID NO.210)
		(SEQ ID NO.211)
		(SEQ ID NO.212)
		(SEQ ID NO.213)
		(SEQ ID NO.214)
		(SEQ ID NO.215)
		(SEQ ID NO.216)
		(SEQ ID NO.217)
5	miR-C15	UACCAACAGGGUAGAACCAACCGGA 1
	miR-C16	AACUGGCCUACAAAGUCCCCAG 1
	miR-C17	UGUAACAGGCAACUCCAUUGGGA 1
	miR-C18	UAGCAGCACAGAAAUAUUGGC 2
	miR-C19	UAGGUAGUUUCAUGUUGUUG 1
	miR-C20	UUACACCACCUUCUCCACCCAGC 1
	miR-C21	GUCCAGAGGGAGAUAGG 1
	miR-C22	CCAGUGUUCAGACUACCGUU 1
	miR-C23	UAAUACUGCCUGGUAAUGAUGAC 2
	miR-C24	UACUCAGUAAGGCCAUUGUU 1
	miR-C25	AGAGGUAUACGCCAUGGAAAGA 1
	miR-C26	UGAAAUGUUUAGGACCAUCAG 1
	miR-C27	UUCCCUUUGUGCAUCCUACGCCUG 1
	miR-C28	UCCUUCAUUCACCGGAGUCUG 1
	miR-C29	GUGAA AUGUUUAGGACCAUCAGA 2
	miR-C30	UGGA AUGUAAAGGAAGUGUGGG 2
	miR-C31	UACAGUAGUUCUGCACAUUGGU 1
	miR-C32	CCUGUAGAACCGAAAUUGUGU 1
	miR-C33	AACCCGUAGAUCCGAACUUGUGAA 1
	miR-C34	GCUCUCCUGGCCUCCUCCUC 1
20		

- 38 -

Table 5

D. melanogaster miRNA sequences and genomic location. The sequences given represent the most abundant, and typically longest miRNA sequences identified by cloning. It was frequently observed that miRNAs vary in length by one or two nucleotides at their 3'-terminus. From 222 short RNAs sequenced, 69 (31%) corresponded to miRNAs, 103 (46%) to already characterized functional RNAs (rRNA, 7SL RNA, tRNAs), 30 (14%) to transposon RNA fragments, and 20 (10%) sequences with no database entry. RNA sequences with a 5'-guanosine are likely to be underrepresented due to the cloning procedure (8). miRNA homologs found in other species are indicated. Chromosomal location (chr.) and GenBank accession numbers (acc. nb.) are indicated. No ESTs matching miR-1 to miR-14 were detectable by database searching.

miRNA	sequence (5' to 3')	chr., acc. nb.	remarks
15			
miR-1	UGGAAUGUAAGAAGUAUGGAG (SEQ ID NO:58)	2L, AE003667	homologs: <i>C. briggsae</i> , G20U, AC87074; <i>C.elegans</i> G20U, U97405; mouse, G20U, G22U, AC020867; human, chr. 20, G20U, G22U, AL449263; ESTs: zebrafish, G20U, G22U, BF157-601; cow, G20U, G22U, BE722-224; human, G20U, G22U, AI220268
20	miR-2a	UAUCACAGCCAGCUUUGAUGAGC (SEQ ID NO:59)	2L, AE003663 2 precursor variants clustered with a copy of <i>mir-2b</i>
20	miR-2b	UAUCACAGCCAGCUUUGAGGAGC (SEQ ID NO:60)	2L, AE003620 2L, AE003663 2 precursor variants
25	miR-3	UCACUGGGCAAAGUGUGUCUCA (SEQ ID NO:61)	2R, AE003795 in cluster <i>mir-3</i> to <i>mir-6</i>
	miR-4	AUAAAGCUAGACAACCAUUGA (SEQ ID NO:62)	2R, AE003795 in cluster <i>mir-3</i> to <i>mir-6</i>

- 39 -

	miR-5	AAAGGAACGAUCGUUGUGUAUAG (SEQ ID NO: 63)	2R, AE003795	in cluster <i>mir-3</i> to <i>mir-6</i>
	miR-6	UAUCACAGUGGCUGUUUCUUUUU (SEQ ID NO: 64)	2R, AE003795	in cluster <i>mir-3</i> to <i>mir-6</i> with 3 variants
5	miR-7	UGGAAGACUAGUGAUUUUGUUGU (SEQ ID NO: 65)	2R, AE003791	homologs: human, chr. 19 AC006537, EST BF373391; mouse chr. 17 AC026385, EST AA881786
	miR-8	UAAAUCUGUCAGGUAAAAGAUGUC (SEQ ID NO: 66)	2R, AE003805	
10	miR-9	UCUUUGGUUAUCUAGCUGUAUGA (SEQ ID NO: 67)	3L, AE003516	homologs: mouse, chr. 19, AF155142; human, chr. 5, AC026701, chr. 15, AC005316
	miR-10	ACCCUGUAGAUCCGAAUUUUGU (SEQ ID NO: 68)	AE001574	homologs: mouse, chr 11, AC011194; human, chr. 17, AF287967
	miR-11	CAUCACAGUCUGAGUUUCUUGC (SEQ ID NO: 69)	3R, AE003735	intronic location
15	miR-12	UGAGUAUUACAUCAGGUACUGGU (SEQ ID NO: 70)	X, AE003499	intronic location
	miR-13a	UAUCACAGCCAUUUUGACGAGU (SEQ ID NO: 71)	3R, AE003708 X, AE003446	<i>mir-13a</i> clustered with <i>mir-13b</i> on chr. 3R
20	miR-13b	UAUCACAGCCAUUUUGAUGAGU (SEQ ID NO: 72)	3R, AE003708	<i>mir-13a</i> clustered with <i>mir-13b</i> on chr. 3R
	miR-14	UCAGUCUUUUUCUCUCUCCUA (SEQ ID NO: 73)	2R, AE003833	no signal by Northern analysis

- 40 -

Table 6

Human miRNA sequences and genomic location. From 220 short RNAs sequenced, 100 (45%) corresponded to miRNAs, 53 (24%) to already characterized functional RNAs (rRNA, snRNAs, tRNAs), and 67 (30%) sequences with no database entry. For legend, see Table 1.

	miRNA	sequence (5' to 3')	chr. or EST, acc. nb.	remarks*
10	let-7a	UGAGGUAGUAGGUUGUAUAGUU (SEQ ID NO:75)	9, AC007924, 11, AP001359, 17, AC087784, 22, AL049853	sequences of chr 9 and 17 identical and clustered with <i>let-7f</i> , homologs: <i>C. elegans</i> , AF274345; <i>C. briggsae</i> , AF210771, <i>D. melanogaster</i> , AE003659
	let-7b	UGAGGUAGUAGGUUGUGUGGUU (SEQ ID NO:76)	22, AL049853†, ESTs, AI382133, AW028822	homologs: mouse, EST AI481799; rat, EST, BE120662
15	let-7c	UGAGGUAGUAGGUUGUAUGGUU (SEQ ID NO:77)	21, AP001667	Homologs: mouse, EST, AA575575
	let-7d	AGAGGUAGUAGGUUGCAUAGU (SEQ ID NO:78)	17, AC087784, 9, AC007924	identical precursor sequences
	let-7e	UGAGGUAGGAGGUUGUAUAGU (SEQ ID NO:79)	19, AC018755	
20	let-7f	UGAGGUAGUAGAUUGUAUAGUU (SEQ ID NO:80)	9, AC007924, 17, AC087784, X, AL592046	sequences of chr 9 and 17 identical and clustered with <i>let-7a</i>
	miR-15	UAGCAGCACAUAAUGGUUUGUG (SEQ ID NO:81)	13, AC069475	in cluster with <i>mir-16</i> homolog
	miR-16	UAGCAGCACGUAAAUAUUGGCG (SEQ ID NO:82)	13, AC069475	in cluster with <i>mir-15</i> homolog

- 41 -

	miR-17	ACUGGCAGUGAAGGCACUUGU (SEQ ID NO:83)	13, AL138714	in cluster with <i>mir-17</i> to <i>mir-20</i>
	miR-18	UAAGGGUGCAUCUAGUGCAGAUA (SEQ ID NO:84)	13, AL138714	in cluster with <i>mir-17</i> to <i>mir-20</i>
5	miR-19a	UGUGCAAAUCUAUGCAAAACUG A (SEQ ID NO:85)	13, AL138714	in cluster with <i>mir-17</i> to <i>mir-20</i>
	miR-19b	UGUGCAAAUCCAUGCAAAACUG A (SEQ ID NO:86)	13, AL138714, X, AC002407	in cluster with <i>mir-17</i> to <i>mir-20</i>
10	miR-20	UAAAAGUGCUUAUAGUGCAGGUA (SEQ ID NO:87)	13, AL138714	in cluster with <i>mir-17</i> to <i>mir-20</i>
	miR-21	UAGCUUAUCAGACUGAUGUUGA (SEQ ID NO:88)	17, AC004686, EST, BF326048	homologs: mouse, EST, AA209594
	miR-22	AAGCUGCCAGUUGAAGAACUGU (SEQ ID NO:89)	ESTs, AW961681†, AA456477, AI752503, BF030303, HS1242049	human ESTs highly similar; homologs: mouse, ESTs, e.g. AA823029; rat, ESTs, e.g. BF543690
15	miR-23	AUCACAUUGCAGGGAUUUCC (SEQ ID NO:90)	19, AC020916	homologs: mouse, EST, AW124037; rat, EST, BF402515
	miR-24	UGGCUCAGUUUCAGCAGGAACAG (SEQ ID NO:91)	9, AF043896, 19, AC020916	homologs: mouse, ESTs, AA111466, AI286629; pig, EST, BE030976
20	miR-25	CAUUGCACUUGUCUCGGUCUGA (SEQ ID NO:92)	7, AC073842, EST, BE077684	human chr 7 and EST identical; highly similar precursors in mouse ESTs (e.g. AI595464); fish precursor different STS: G46757
	miR-26a	UUCAAGUAAUCCAGGAUAGGCU (SEQ ID NO:93)	3, AP000497	

- 42 -

miR-26b	UUCAAGUAAUUCAGGAUAGGUU (SEQ ID NO:94)	2, AC021016	
miR-27	UUCACAGUGGCUAAGUUCCGUU (SEQ ID NO:95)	19, AC20916	U22C mutation in human genomic sequence
5	miR-28	AAGGAGCUCACAGUCUAUUGAG (SEQ ID NO:96)	3, AC063932
10	miR-29	CUAGCACCAUCUGAAAUCGGUU (SEQ ID NO:97)	7, AF017104
15	miR-30	CUUUCAGUCGGGAUGUUUGCAGC (SEQ ID NO:98)	6, AL035467
20	miR-31	GGCAAGAUGCUGGCAUAGCUG (SEQ ID NO:99)	9, AL353732
	miR-32	UAUUGCACAUUACUAAGUUGC (SEQ ID NO:100)	9, AL354797 not detected by Northern blotting
25	miR-33	GUGCAUUGUAGUUGCAUUG (SEQ ID NO:101)	22, Z99716 not detected by Northern blotting

*If several ESTs were retrieved for one organism in the database, only those with different precursor sequences are listed.

20 †precursor structure shown in Fig. 4.

Claims

1. Isolated nucleic acid molecule comprising

5

(a) a nucleotide sequence as shown in Table 1, Table 2, Table 3 or Table 4 or a precursor thereof as shown in Figure 3, Figure 4 or Figure 7.

10

(b) a nucleotide sequence which is the complement of (a),

(c) a nucleotide sequence which has an identity of at least 80% to a sequence of (a) or (b) and/or

15

(d) a nucleotide sequence which hybridizes under stringent conditions to a sequence of (a), (b) and/or (c).

2. The nucleic acid molecule of claim 1, wherein the identity of sequence (c) is at least 90%.

20

3. The nucleic acid molecule of claim 1, wherein the identity of sequence (c) is at least 95%.

25

4. The nucleic acid molecule of any one of claims 1-3, which is selected from miR 1-14 as shown in Table 1 or miR 15-33 as shown in Table 2 or miR 1-155 as shown in Table 3 or miR-C1-34 as shown in Table 4 or a complement thereof.

30

5. The nucleic acid molecule of any one of claims 1-3, which is selected from mir 1-14 as shown in Figure 3 or let 7a-7f or mir 15-33, as shown in Figure 4 or let 7a-i or mir 1-155 or mir-c1-34, as shown in Figure 7 or a complement thereof.

- 44 -

6. The nucleic acid molecule of any one of claims 1-4 which is a miRNA molecule or an analog thereof having a length of from 18-25 nucleotides.

5 7. The nucleic acid molecule of any one of claims 1-3 or 5, which is a miRNA precursor molecule having a length of 60-80 nucleotides or a DNA molecule coding therefor.

10 8. The nucleic acid molecule of any one of claims 1-7, which is single-stranded.

9. The nucleic acid molecule of any one of claims 1-7, which is at least partially double-stranded.

15 10. The nucleic acid molecule of any one of claims 1-9, which is selected from RNA, DNA or nucleic acid analog molecules.

11. The nucleic acid molecule of claim 10, which is a molecule containing at least one modified nucleotide analog.

20 12. The nucleic molecule of claim 10 which is a recombinant expression vector.

13. A pharmaceutical composition containinig as an active agent at least one nucleic acid molecule of any one of claims 1-12 and optionally a pharmaceutically acceptable carrier.

25 14. The composition of claim 13 for diagnostic applications.

15. The composition of claim 13 for therapeutic applications.

30 16. The composition of any one of claims 13-15 as a marker or a modulator for developmental or pathogenic processes.

- 45 -

17. The composition of claim 13 as a marker or modulator of developmental disorders, particularly cancer, such a B-cell chronic leukemia.
18. The composition of any one of claims 13-15 as a marker or modulator of gene expression.
19. The composition of claim 18 as a marker or modulator of the expression of a gene, which is at least partially complementary to said nucleic acid molecule.
- 10 20. A method of identifying microRNA molecules or precursor molecules thereof comprising ligating 5'- and 3'-adapter molecules to the ends of a size-fractionated RNA population, reverse transcribing said adapter-containing RNA population and characterizing the reverse transcription products.

15

Fig. 1 A




Fig. 1 B




Fig. 2






Fig. 3




. Fig. 4

*Fig. 5***miR-1a miR-122a**

ht kd lv pc sp



ht kd lv pc sp




— L

— 21-nt

miR-124abrain

rbmbcx cb ht lg lv co si pc sp kd sm st H



— L

— 21-nt




— tRNAs

Fig. 5 (cont.)

brain

rbmb cx cb ht lg lv co si pc sp kd sm st H


miR-30a-s



brain

rbmb cx cb ht lg lv co si pc sp kd sm st H

miR-101



— miR-L


— miR-S

— tRNAs

brain

rbmb cx cb ht lg lv co si pc sp kd sm st H


miR-125a



brain

rbmb cx cb ht lg lv co si pc sp kd sm st H

miR-125b



— miR-L

— miR-S






Fig. 5 (cont.)

Fig. 5 (cont.)



*Fig. 6***A**

- C. elegans* lin-4
D. melanogaster miR-125
M. musculus/H. sapiens miR-125b
M. musculus/H. sapiens miR-125a

UCCCUGAGACCUC--AAG-UGUGA
UCCCUGAGACCCU--AACUUGUGA
UCCCUGAGACCCU--AACUUGUGA
UCCCUGAGACCCUUUAACCUGUGA


B

Fig. 7

name	sequence	structure
let-7a-1	UGAGGUAGGUUGGUUAUAGUU	UG UGGGA GAGGUAGGUUGGUUAUAGUU CAC UGGGA GAGGUAGGUUGGUUAUAGUU GUG AUCCU UUCGUCAUCUACAUUCAA CA -
let-7a-2	UGAGGUAGGUUGGUUAUAGUU	UU G U AGG GAG UAG AGGUAGGUUGGUUAUAGUU UCC UUC AUC UCCGACAUGUCAA U- G C
let-7a-3	UGAGGUAGGUUGGUUAUAGUU	U GGG GAGGUAGGUUGGUUAUAGUU UCC UUCGUCAUCUACAUUCAA U
let-7b	UGAGGUAGGUUGGUUGGUUGGUU	GG CGGG GAGGUAGGUUGGUUGGUU UC GUCCC UCCGUCAUCCAACAUUCAA AG --- -
let-7c	UGAGGUAGGUUGGUUAUAGGU	A UU G U GC UCCGGG GAG UAG AGGUAGGUUGGUU CG AGGUUC UUCGUCCAGC UAUCAA - CU G U
let-7d	AGAGGUAGGUUGGUUAUAGU	A C UUA CCUAGGA GAGGUAGGUUGGUUAUAGU GGAUUCU UUCGUCCAGC UAUCAA - A
let-7e	UGAGGUAGGAGGUUGGUUAUAGU	C CU G U CC GGG GAG UAGGAGGUUGGUUAUAGU GG CCC UUCGUCCAGC UAUCAA A CU G

Fig. 7 (cont.)

let-7f-1	UGAGGUAGUAGAUUGUAAGUU	AGU UCAG AGUC CC-	GAGGUAGUAGAUUGUAAGUU UUCGUUAUCUAAACAUCAAUAA GAGGACUUG	GGGGUAG UCCAUU UU	UG
let-7f-2	UGAGGUAGUAGAUUGUAAGUU	<u>U</u> CUGUGGGA GGCACCCU	GAGGUAGUAGAUUGUAAGUU UUCGUCAUCUGACAUCAAUAA UAGA	UUAGGG GGUUCU ACCC	A C AC
let-7g	UGAGGUAGUAGUUGUACAGUA	A <u>U</u> CC GG CG A -	UGAGG GAGGUAGU GUU UCCGUCA CGGACAU U C	GUCU UG UAGA AC GG -	A- A A A C
let-7h	UGAGGUAGUAGUAGUACAGUU				
let-7i	UGAGGUAGUAGUUGUGCU	<u>U</u> CUGGC GAUCG	GAGGUAGUAGUUGUGCU UCCGUCAUCGCG CAA	UU U UAGGAGGUG	U GGGGU UC UUAC
miR-1	UGGAAGUAAAGGAAGUAGGAG	A UUGAGA UUC GAG CGG - UCUAAG	UUGAGA GUCCAGGUUC UUGCAUUC CGAGGUAGGAAG AAUGUAAG	C A A G	AUA GUU U ACU
miR-1b	UGGAAGUAAAGGAAGUAGUAA	A ACAU ACUCU A -	ACAU UUGAAGGAAAGUA GGUAU A	AC UGG AUC CGA GU	AC C C U AL449263.5

Fig. 7 (cont.)

miR-1c	UGGAUGUAAAGAGUAUGUAC		
miR-1d	UGGAUGUAAAGAGUAUGUAUU	GC C GCUUGGGA ACAUACUUUUAU CGGACU A-	UGAACCC U GG G CGAACU
miR-2a-1	UAUCACAGCCAGGUUUGAGGC	- - GGUGGCC UCAAAG CGAUUC A U G	A AUUUC UU CGC GCG U CG
miR-2a-2	UAUCACAGCCAGGUUUGAGGC	A C AUCU AGC UAGG UCG A U	GAUAC UGGUUGGUAUAG ACCGACACUAUAC CG GCAAC
miR-2b-1	UAUCACAGCCAGGUUUGAGGC	U CU CAAC GUUG AGGGAGUUC C CG	A C UG UCUUCAAG UGGC GUGA ACCG CACU A AUACU A
miR-2b-2	UAUCACAGCCAGGUUUGAGGC	A U C G	UUU--- A U GGUGUC UUCUCAAG UGGUUGGUGA AUG AGGGCAG GAGGAGUUC C G
miR-3	UCACUGGGCAAGUGUGUCUCA	C A G	UUCA A UG GU AC A CUCU

Fig. 7 (cont.)

miR-4	AUAAAGCUAGACACCAUUGA	U UU C C C GG UU UGGCAAU AGUUUC UGGU GUC AGC UUA UGAUU \/ GGGUUG UGAAAG ACCA CAG UCG AAU ACUGG U C UU A A A --- CC
miR-5	AAAGGAAACGAUCGGUUGUGAAUAG	UA--- C AGUUGU GC AAAGGAA GAUCGUUGUGAU AUG CG UUUCCUU UUAGUGACACUAUAC U C AAUA - AAUCCU
miR-6-1	UAUCACAGUGGGUGUUCUUUU	A- C AG UAAUA UUUA UGUAGAGGGAAUAGUUUGUGUGUG UGUA U \/ AAAU AUGUUUUUCUUUGGGUGUGACAC AUAU A U CC UU CU UACCA
miR-6-2	UAUCACAGUGGGUGUUCUUUU	C UU UG C UU - G UAACC AAGGGAAC C CUG UGAUUA UA UU A GUUGG UUUUCUUG G GAC ACUUAU AU AA A U UC GU - C C A
miR-6-3	UAUCACAGUGGGUGUUCUUUU	A A U AAAC CAA AGAAGGGAACGGGUUGUGUG UGAUGUAG UUG \/ GUUU UUUUUCUUUGGGUGAC ACUUAUU AAC U G - U ACUC
miR-7	UGGAAGACAUAGUGAUUUGUGU	U U U U --- UGGUC GAGUGCAU CGGUAGAGC AG GAUUU UGUUGUU \/ UUUACGUG GCGAU UCUCUUG UC CUAAA ACAUAUA U C - U C UA UGGUU
miR-8	UAAUACUGUCAGGUAAAGAUUC	CUGUUC - G C UCCUU AGGGACAU ACAUCUU ACC GGCAG AUUAGA \/ UCCUGUG UGUAGAA UGG CUGUC UAAUCU U CCUGG- A A A CAAUAU

Fig. 7 (cont.)

miR-9	UCUUGGUUAUCUAGCUGUAUGA	- UAU <u>U</u> G - GAU GCUA UGUUG CUUUGGU CUAGCU UAUCA GU A CGAU AUAAA GAAGCCA GAUCGA AUACU CA A U UUC A G AUA
miR-10	ACCCUGUAGAUCCGAAUUUGU	CU - G U AUACU CCACGU ACC CU UAGA CCGAAUUVGGUUU A GGUGUG UGG GA AUUC GGCUUAACAGGA G UU A G U AUUC
miR-11	CAUCACAGUCUGAGGUUCUUGC	U UCU CCC U ACU GCACUUG CAAGAACUU CUGUGA GCG GU U CGUGAGU GUUCUUGAG GACACU CGC CG A C UCU A--- AAA
miR-12	UGAGUAUUAUACUAGGUACUGGU	UG U <u>C</u> - GCCUU UACGGU AGUAU ACAU AGGUACUGGU GU A GUGCCG UCAUA UGUA UUCAUGACCA CA A CA C - ACCUA
miR-13a	UAUCACAGCCAUUUUGAUGAGU	U C - A UC-- CU UACG AACUC UCAAAG GGUGUGA AUG GA A GUGC UUGAG AGUUUU CCGACACU UAC CU U U U A A UCAU AU
miR-13b-1	UAUCACAGCCAUUUUGACGEGAU	UG- U ACU UAUU CCA UGUAAAAG UUGUGA UAUG C GGU AGCAGUUUAC GACACU AUAC A UUG U --- UAAC
miR-13b-2	UAUCACAGCCAUUUUGACGAGU	UAUU G A GCUA UU AAC CGUCAAAAG CUGUGA UGUGGA U UUG GCAGUUUAC GACACU AUACUU G GU-- A C ---- CA

Fig. 7 (cont.)

miR-14	UCAGUCUUUUCUCUCCUA	C C C GCUU UGGGGAG GAGA GGGGACU ACUGU \/ AUUCCUC CUCU UUUCUGA UGAUA A U U C AAUU
miR-15a	UAGCAGCACAUAAUGGUUG	GAGUAAGUA UA GA U CCUUG GCAGCACA AUGGUUUGUG UUU \ GGAAC CGUCGUGU UACGGACGU AAA G AUAAAACUC UA GG A
miR-15b	UAGCAGCACAUCAUGGUUACA	U C C A ACA CUG AGCAGCA AU AUGGUU CAU CU \/ GAU UCGUCGU UA UACUAAG GUA GA G C U U C - ACU
miR-16	UAGCAGCACGUAAAUUUGCG	AG C - A CGUUA UCUA GUAGC UGC UUAGCAGCAC GU AAUAUUGG AGAU \/ CAGUUG AUG AGUCGUCGUG CA UUAUGACC UCUA A GA A U A ----- UUAA
miR-16	only different precursor	UC CU UA C AG AU GU CACU AGCAGCACG AAUAUUGG GU UGA A CA GUGA UCGUCGUG UUUAACC CA AUU U GU UU CA A A----- AUAA
miR-17	ACUGGCAGUGAACGCCACUUG	GA CA- A G G - AU GUCA AUAAUGU AGUGGUU CA UGCAG UAG UG \ CAGU UAUUACG UUCACGGA GU ACCUC AUC AC U GG AUG A G - U GUG
miR-18	UAAGGGUGCAUCUAGGUAGUA	CU U C U A UGAA AG UGUU AAGG GCAG UAG GCAG UAG GU A ACGG UUCC CGUG AUC CGUC AUC CG U UC U A C - UA-- AU

Hg. 7 (cont.)

miR-19a	UGUGCAAAUCUAUGCAAAACUGA	U U --- --- AGA CCAG CC CUGUAGUUUUGCAUAG UGGCAC UACA \\\nCGUC GG GGUAGUCAAAACGUAC AACGUG AUGU A C U UA UUG AAG
miR-19b-1	UGUGCAAAUCCAUGCAAAACUGA	UU CACUG UUAGGUAGUUUUGCA GG UUUGCA CAGC \\ GUGAU GGUGUCAGUCHAAACGU CC AAACGU GUUG A --- A U --- UCUUAU
miR-19b-2	UGUGCAAAUCCAUGCAAAACUGA	CUAC UUACAAUUAUGUUUUGCA GG UUUGCAU GCGUAUA A ACAUUG UGUAAU AGUGUUAUGICA AAAACGU CC AAACGUG UGUUAU U --- A U UCGG G
miR-20	UAAAGUGCUUAUGUGGAGGUAG	C A- G - UU GUAG ACU AAGUGCUUAUGUGGAG UAG UG U CGUC UGA UUCAGGAGUAUACGUC AUC AU A A AA - U UG
miR-21	UAGCUUUAUCAGACUGAUUGA	A A A A U AA UGUCGGGUAGCUUAUC GACUG UGGUG CUGU G \\ ACAGUCUGUGGGUAG CUGAC ACAAC GGUAC U A - C - UC
miR-22	AAGCUGCCAGUUGAGAACUGU	U CC A U CCUG GGC GAG GCAGUAGUUCUUCAG UGGCA GCUUUA GU \\ CGC CUC CGUUGUCAAGAGGU ACCGU CGAAAU CG A U C - G - ACCC
miR-23a	AUCACAUUGCAGGGAUUUC	C C G G C UUC GG CGG UGGGG UUCCUGG GAUG GAUUG C CC GCC ACCUU AGGGACC UAC CUAAAC U A A U G A ACUG

Fig. 7 (cont.)

miR-23b	AUCACAUUCCAGGGAUUACAC	C U --- GG UGC UGG GUUCCUGGCA V G UGAUUU CC ACG ACC UAGGGACC <u>GU</u> AC ACUAAA A C AU U - AUUAGA	- C GUGACU U GUGACU G G A UA UCUCAU CUCC GU CCU CUGAGCU <u>GA</u> UCAGU GAGG CA GGA GACUUGACU GGUC <u>CA</u> A A C C - CACAUU
miR-24-1	UGGCUCAGGUUCAGCAGGAACAG	CC CG CU- AA-- UU CUCUG UCC UGC ACUGAGGUG ACACAG GGGAC AGG ACG <u>UGACUCGGU</u> UGUGUU G A- --- ACU CACA UG	
miR-24-2	UGGCUCAGGUUCAGCAGGAACAG	A AG UU G UG ACG GGCC GUGUG AGGC GAGAC G GCAAU CUGG C CCGG CGGAC <u>UCUG</u> CUCUG C GUUA GGUC U C AG G <u>UU</u> A CG CG	
miR-25	CAUUGCACUUUGUCUCGGCU <u>GA</u>	- G U U GAG AGGCC GUG CCUCGU CAAGUAA CCAGGAUAGGCGU G UCCGG CGC GGGCA GUUCAUU GGUUCUAUCCGGUA U G A C - ACCC	
miR-26a	UUCAAGUAUCCAGGAUAGGU	GA - U U GUG CCC CCC AGU CAAGUAA AGGAUAGGUUG GGCC GGG UCG GUCAUA UCUGUCCGAC C AG C - CC CUGU	
miR-26b	UUCAAGUAUUCAGGAUAGGU	A A A U G UCCAC CUG GG GC GGGCUUAGCUGCU GUGAGCA GG GAC CC CG CUUGAACGGUGA CACUUGU CU A C C C - G GAACC	
miR-27a	UUUCACAGGGCUAAGUUCGGCU		

Fig. 7 (cont.)

miR-27b	UUCACAGUGGUAAAGUUCUG	AGGUGGAGGCUUAGCUG UCCACGCUUAGAUCGGU	AUUG GA--	GUGAACAG CACIUGUU	UGAU GCC U	U V
miR-28	AAGGAGCUCACAGCUAAUUGAG	GGU CUGGCCUC AGGAGCUCACAGCUA UCA GGACGGGAG UCCUCGAGGUUAGAU	C A C G	U <u>GGGUU AGAG</u> ACCACGA U <u>CUU</u>	U <u>GUUA</u> C C	CC U
miR-29a	CURGCACCAUCAUCUGAAUUCGGUU	AUGACUGAUUUC U <u>AUUGGCCUAAAG</u>	U <u>UU</u> <u>UCU</u>	UGGUGGUU ACCACGA	U <u>CAAU</u> U <u>AAAU</u>	CC A
miR-29b	UAGGCACCAUUGAAAUCAGUGUU	AGGA GCUGGUUUC UCUU U <u>GACUAAAGU</u>	A G	U <u>GGUG</u> U <u>ACCAC</u>	U <u>UAGAU</u> GAUCUG	U A
miR-29c	UAGGCACCAUUGAAAUCGGuuu					
miR-30a-s	UGUAAAACAUCUCCGACUGGAAGC	GGC CUGUAAACAAUCC CGU GACGUUUGUAGG	A C	U <u>C</u> ---	GACUGGGAGCU CUGACUUUCGG	----- GUAAA C
miR-30a-as	CUUUCAGUCGGGAUGGUUUGGAGC	GCG CUGUAAAACAUC CGU GACGUUUGUAGG	A C	U <u>C</u> ---	GACUGGAAGCU CUGACUUUCGG	----- GUAGA C

Fig. 7 (cont.)

miR-30b	UGUAAACAUCCUACACUCAGC	U U - UCAUA AUGUAAA <u>ACA</u> <u>UCC</u> ACA CUCAGCUG C UGCAU <u>UUG</u> UAGG UGU GGGU <u>CGG</u> U A - A UGC <u>GU</u>
miR-30c	UGUAAACAUCCUACACUCAGC	UACU U U - AGA GUAA <u>ACA</u> CCU CUC <u>U</u> <u>GG</u> AA UCU CAU <u>UUG</u> U GGA GAGG <u>GU</u> GA G - A-- AAGAAU human
miR-30d	UGUAAACAUCCCGACUGGAG	U U CCC GUAA <u>ACAU</u> GACUGGG <u>AG</u> CU C GU GU CG CGUUG <u>UAG</u> CUGACUU <u>U</u> <u>CG</u> A U U A--- AUGCAC chr8 human
miR-31	GGCAAGAUGCUGGCAUAGCUG	GA G G GAA GGAGAG GGC <u>AA</u> AUG UGGCAU <u>AGC</u> U- GAA CCUUUC CGUU UAC ACCGU <u>AUCG</u> CAA C UA A A UC GGG
miR-32	UAUUGCACAUUACUAAGUUGC	U - UU C GGAGAU <u>UUG</u> CACAU ACUAAGU <u>U</u> <u>GG</u> CAU G GU A CUUUAUGUGUGUG UGAU <u>U</u> <u>AACG</u> UA C CG C - A UC G
miR-33	GUGCAUUGUAGUUGCAUUG	A UU UUCU UG CUGUGGUG <u>CAU</u> GU G GCAU <u>U</u> <u>GG</u> CAUG GG GACACU <u>ACG</u> U <u>AC</u> C UGUAA <u>AC</u> UAC C C G C UU ---- AU
miR-99a	ACCCGUAGAUCCGAUCUUG	A U G G A G C CAUA ACC <u>CG</u> U <u>AGA</u> CGA CU <u>U</u> <u>GG</u> U G GUG UGG <u>GU</u> AU <u>U</u> GCU GA <u>AC</u> GC GC G C UU C - CAG

Fig. 7 (cont.)

miR-99b	CACCCGUAGAACCGACCUUGGG	<u>C</u> GGCAC CUGUG CC	<u>A</u> ACCGUAGA UGGUGUCU GU	<u>C</u> CGA GCU C	<u>---</u> UGGGG ACGCC ACAC	<u>C</u> GG CU G	<u>-</u> C	
miR-101	UACAGUACUGUGAUAAACUGA				UCAGUUAUCACAGUGGUG AGUCAAUAGUGCAUGAC	<u>A</u> UGGU -	GUCCA UGCU AUGG AAAUC	<u>A</u> U U -
miR-122a	UGGAGUGUGACAUAUGGUUUGU				AGCUGU UCGAUA AA	<u>G</u> AGUGUGA UCACACU A	<u>C</u> AUGGUUUUG UUAACCGCAAA UAUCA	<u>Y</u> GUCC A A woodchuck
miR-122b	UGGAGUGUGACAUAUGGUUUGA							
miR-122a,b	UGGAGUGUGACAUAUGGUUUG							
miR-123	CAUUAUACUUUUGGUACGGG				UGAC ACUG G	<u>A</u> GC CG C	<u>U</u> CAUAAA GUAAA U	<u>CG</u> GUAC GCCAUGC UCAA-
miR-124a*	UUAGGGCACGGGGAAUGCCA					<u>-</u> GAGA A	<u>C</u> G C - G	<u>A</u> GA GCG GG AC UAAUG UU C CAUAU

Fig. 7 (cont.)

miR-124b	UUAAGGCACGGGGUGAAUGC	CC A GA UAAUG CUCU GUGUUAC GCG CCUUUGAUU GAGA CGUAAGUG CGC GGAAUAAA U AC G AC CAUAC AC021518
miR-125a	UCCCCUGAGACCCUUUACCUUGUG potential lin-4 ortholog	C C C UA --- A CUGGGU CCUGAGA CCUU ACCUGUGA --- GG C GGUCCG GGGUUCU GGAG UGGACACU CC G A U --- GGGAA U
miR-125b	UCCCCUGAGACCCUAACUUGUGA potential lin-4 ortholog	UC C C A GG- U GCUUAG CCUGAGA CCU ACUUGUGA UAU U CGGAUC GGGUUCU GGA UGAACACU AUG U CA U C ACA A
miR-126	UCGUACCGUGAGUAAAUGC	A U CGCUG C GC CAUUAUUACUU UGGUACG UGA A CG GUAAAUGC AGCUGA CC AGGCUA AG A C U UCAA- U
miR-127	UCGGGAUCCGUCUGAGCUUGGU	A U G C --- AG CC GCC GCU AAGCUCAGA GG UCUGAU UC \ GG UGG CGG UUCGAGUCU CC AGGCUA AG A C U - G U CU AA
miR-128	UCACAGUGAACCGUCUUUUU	UUC UAG CU U GUUGCA GGGCCG CACUGU GAGGGU U CGACUU UUU CUCUGGC GUGACA CUCUUA A CAA --- C
miR-129	CUUUUUUCGGUCUGGGCUUGC	- C CU G UUCCU C GGAU CUUUUUG GGU GGGCUU CUG CU A UCUA GAAAAAC CCA CCCGAA GAC GA A U C U G UGAU- C human

Fig. 7 (cont.)

miR-130	CAGUGCA AUGUUAAAAGGGC	- C GCUUUU ACAUUGGGCU CU A GCUAAC CU CGGGAAAA UGUAACGUGA GA G A U C GCCAUGU
miR-131	UAAAGCUUAGAUAAACCAGAAAGU	G C G U A GUU UUAU UUUGGUUAUCUAGCU UAUGAG GU U CAA AUG AGCCAAUAGAUCGA AUACUU UG U A A A C G
miR-132	UAACAGUCUACAGCCAUGGU	A G G G C GGGC ACCGUGGCC CCCG UGGUACCGA A UUC G- G GUU GGUUACU UGG G- GUACAAUGG GCC A A C AU AG A
miR-133	UUGGUCCCCUUCAACCAGCUGU	A A A G G C GCUA AGCUGGU AA GG ACCAAAC CGAU UCGACCA UU CC UGGGUUAG G AC C C GUAC GU GGCUC
miR-134	UGUGACUGGUUGACCAAGGGAA	GU AGGGU UCCCA GUGACUGG UG CCA AGGG CACUGAUC AC GGU UCCC AC C CG G ACU- UC G GGU AC GU GCGU AC
miR-135	UAUGGCCUUUUUAUCCUAUGGAA	UU UUCUAU CUAUGGCCUUU AUCCUAUGUGA GGUGCCGAGG UAGGGAUUAACU AC U- CGCUCC G GGU AC GU GCGU AC
miR-136	ACUCCAUUUUUUGAUGGAA	C UUU UUCU GUAGACUC AUUUG UGAUGAUGGA CUUCUGAG UAAAC GCUACUACCU AC UCU CGAA G GGU AC GU GCGU AC

Fig. 7 (cont.)

miR-137	UAUUGCUUAAGAAUACGGUAG	G G A - GA CUUCGGU ACG GUAUUCUUGGGUGG UAAA CG \ GGAGCG UGC CAUAGAUAUCCGU AUUGU GC U A G - - U AU
miR-138	ACCUUGGUUGGUAGAACU	-- UCA AC- C CG CAGGU GGUGUUGUGAA GGC CG GAG AG C GUUGG CCACAGCACUU UCGGC UUC UC A GA UA- CCA - CU
miR-139	UCUACAGUGGCACGGUUCU	G U AAUUCUA CAG GC CGUGUCUCCAGU CA AUGAGGU GUC CG GCGAGGGUUC GUUGG C - GAGGC human
miR-140	AGUGGGUUUACCCUUAUGGUAG	- A CCUG CC GUGGCCCCU UGGUAGG ACG A GGAC GG CACCAAGAUGGG ACCAUCU UGU U A - C - -- CG
miR-141	AACACUGUCUGGUAAAGAUGG	U --- U AU GAAG GGG CCAUCUU CCAG GCAGUGUUGG GGUU \\ CCC GGAGAA GGUC UGUCACAAUC UCGA U - AU - C- AGUA
miR-142s	CAUAAAGUAGAAAGCACUAC	AC- A CCAUAAGUAG AAGCACUAC CA C GUAAUUCUAC UUUGUGAUG GU A GUA C UGGGAG C
miR-142as*	UGUAGUGUUUCCUACUUUAUGG	AC- A CCAUAAGUAG AAGCACUAC CA C GUAAUUCUAC UUUGUGAUG GU A GUA C UGGGAG C

Fig. 7 (cont.)

new	AUAAGCAGCAAAAGCUUGU	G GGGGAGCUUUU CG CG UUUAAC UG \ ACUG UGUUCGAAAG GCG AGC AAAUAG AC G G A AG C UC AL049829.4
miR-143	UGAGAUGAACGACUGUAGCUAG UUAUGAAGCACUGUAG	G G G U - AG CCUGAG UGCAGUGCU CAUCUC GG UC U GGACUC AUGUCACGA GUAGAG CU AG U G G G AC008681.7
miR-144	UACAGUAUAGAUGAUGUACUAG	G A A - GU GGCUUGG AUAUCAUC UAUACUGUA GUUU G CUGAUC UGUAGUAG AU AUGACAU CAGA A A - CA GU
miR-145	GUCCAGUUUCCAGGAAUCCUU	C UC U C UGG AUG CUCA GG CAGU UU CCAGGAAUCCU C GAGU UC GUCA AA GGUCUUAGGG C - UU U A UAGAU
miR-146	UGAGAACUGAAUCCAUUGGUUU	CY C AUUC AGCU GAGAACUGAAUU CAUGGGUU A UCGA UUCUUGACUUA GUGUCCAG A C- A ACUGU
miR-147	GUGUGUGAAUCCUUCUGCC	A- CAA ACA --- GA AAUCUA AGA CAUUCUGCACAC CCA \ UUAGAU UCU GUAAAGGUGUGUG GGU C CG UC- ACCGAA AU human
miR-148	UCAGUGCACUACAGAACUUUGU	- A- CC - AGU GAGGCCAAGGUUCUG AG CACU GACU CUG \ CUCUGUUUCAAGAC UC GUGA CUGA GAU A A AC --- A AGU human

Fig. 7 (cont.)

miR-149	UCUGGUCCGGUGUCUACUCC	<u>G</u> <u>C</u> <u>G</u> <u>A</u> <u>GUG</u> <u>G</u> <u>GGCUCUG</u> <u>CUC</u> <u>GU</u> <u>UCUUC</u> <u>CUC</u> <u>UUU</u> <u>U</u> <u>UCGGGGC</u> <u>GAG</u> <u>CA</u> <u>GGAGG</u> <u>GAGG</u> <u>GAG</u> <u>C</u> G A G - AG- C
miR-150	UCUCCCCAACCUUAGGU	<u>AC</u> <u>U</u> <u>UG-</u> <u>UG</u> <u>CCCUGUCUCCA</u> <u>CCU</u> <u>GUACCAG</u> <u>CUG</u> \ <u>GGGAUAGGGGU</u> <u>GGA</u> <u>CAUGGUC</u> <u>GAC</u> <u>C</u> CC - CCA UC
miR-151	CUAGACUGAGGCCUCUTGAGGU	<u>C</u> <u>CA</u> <u>UGUCU</u> <u>CCUG</u> <u>CCUCGAGGAGCU</u> <u>CAGCUUAGUA</u> \ <u>GGAC</u> <u>GGAGGUUCCUCGG</u> <u>GUCAGAUCAU</u> C A- CCCUC
miR-152	UCAGUGCAUGACAGAACUUGG	<u>G</u> <u>A</u> <u>CC</u> <u>CGG</u> <u>C</u> <u>CGGGCCUAGGUUCUGU</u> <u>AU</u> <u>CACU</u> <u>GACU</u> <u>GCU</u> <u>U</u> <u>GCCCCGGGUUCAAGACA</u> <u>UA</u> <u>GUGA</u> <u>CUGA</u> <u>CGA</u> <u>G</u> G C -- --- G
miR-153	UUGCAUAGUCACAAAGUGA	<u>-</u> <u>GU</u> <u>A-</u> <u>AAU</u> <u>CAGUG</u> <u>UCAUUUUUGUAI</u> <u>UGCAGCU</u> <u>GU</u> \ <u>GUUAC</u> <u>AGUGAAAACACUG</u> <u>ACGUUGA</u> <u>CG</u> <u>A</u> U AU CC AGU
miR-154	UAGGUUAUCCGGGUUCCUUCG	<u>U</u> <u>-</u> <u>CCU</u> <u>--</u> <u>UUU</u> <u>GAAGAUAGGUUA</u> <u>CCGUGU</u> <u>UG</u> <u>UCGC</u> \ <u>UUUUUAUCCAGU</u> <u>GGCACCA</u> <u>AC</u> <u>AGUG</u> A U U UAAGC UUU
miR-155 [BIC-RNA]	UUAAUGGUAAUUGUGAUAGGG	<u>U</u> <u>U</u> <u>A</u> <u>UUGGCC</u> <u>CUGUUAAUGGUAAU</u> <u>G</u> <u>G</u> <u>UAGGGGUU</u> \ <u>GCAAUUAGGAIUG</u> <u>U</u> <u>C</u> <u>AUCUCAG</u> <u>U</u> - C - UCAGUC

Fig. 7 (cont.)

name	sequence	structure
miR-C1	AACAUUCAACGCGUGUGAGU	U <u>A</u> U <u>C</u> A <u>G</u> GGGAUUCA CCA GG ACA UCAACG GUCCGGUG GUUT GGU CC UGU AGUUGC CAGCCAC CAAA A U A C --- - AAAACAAA
miR-C2	UUUGGCCAAUGGUAGAACUCACA	U <u>U</u> <u>GG</u> <u>UCA</u> UAAGGU ACCAU <u>UUGGCAA</u> <u>UAGAAC</u> <u>CACCGG</u> A UGGUA AACCGUU AUCUTUG GUGGCC A UC CAG --- CAGGGU
miR-C3	UAUGGCCACUGGUAGAACUCACUG	G <u>AC</u> - <u>GA</u> --- AC CUGU <u>UAUGGC</u> UGGUA <u>AUUCACUG</u> UGA A GACA AUACCG GCCAU <u>UAAGUGAC</u> ACU G A GGAA --- UG CU
miR-C4	CUUUUGGGGUCUGGGCUUUGU	- <u>C</u> <u>CU</u> <u>G</u> UUUU C UGGAU <u>CUUUUGG</u> <u>GGCUU</u> <u>CUG</u> CU G AUCUA GAAAAC CCA CCCGAA GAC GA A U C UU G UGAU C
miR-C5	UGGACGGAGAACUGAUAGGGU	U <u>C</u> <u>AG</u> --- UG CCU UCCUUAUCA UUUUC C CAGC UUUG A GGA <u>GGGAAUAGU</u> <u>AAGAGG</u> <u>GGUUG</u> GAAU C U <u>C</u> <u>CA</u> U CU
miR-C6	UGGAGAGAAAGGCAGTUC	AGGGAUUGGAG <u>A</u> <u>G</u> AU UC TUCCUGGGUCU <u>C</u> <u>CA</u> G GUGGCCUG GG C - G --- UC

Fig 7 (cont.)

name	sequence	structure
miR-C7	CAAAGAAUUCUCCUUUUGGGCUU	<pre> U UU \ _ ACUUTUCCAAAGGAUUC CCUU GGGCUU UGAGGGUUUUUAAG GGAA CCCGAA U U- UUUUAU </pre>
miR-C8	UCGUGUCUUGGUUGUGCAGCGG	<pre> A A C CGCUGC UC GGCU CAACACAGGAC CGGG U GG CCGA GUUGGUUGUCUG GCUC C - C _ CCCAGU </pre>
miR-C9	UAACACUGUCUGGUAAAGAUGU	<pre> - C UU UUUG GGGCAUC UTACCGGACAGUG UGGA UC \ CUUGUAG AAUGGUUCUGUAC AUCU AG G - C- _ UUC </pre>
miR-C10	CAUCCCUGCAUGGGAGGGU	<pre> CA UC GU UGAGCUC CA CCUUGCAUG GGAGGG U AGG GU GGGACGUAC CUCUCC C AC UU AC CAAAAGU </pre>
miR-C11	GUGCCUACUGAGCUGACAUAGU	<pre> G G A UA UCUCAU CUCC GU CCU CUGAGCUGA UCAGU \ GAGG CA GA GACUTUGACU GGUCA U A A C _ CACACU </pre>
miR-C12	UGAUUAUGUUUGAUAAUAGGU	<pre> U- UA--- UU CUGUG GAUAUGUUUGAUAAU _ GACAU UUUAUCGAACUAAUA CUAU A CC UCAAC UU </pre>

Fig. 7 (cont.)

name	sequence	structure
miR-C13	CAACGGAAUCCAAAAGCAGCU	C AGCGGG <u>AACGGAAUUC</u> C <u>AA</u> GCAGGUG GU CU C UCGUCC UUGCUUAGG UU CGUCGAC UA GA A C - CA CU C G
miR-C14	CUGACCUAUGAAUUGACA	C UGACCUAUG <u>AAUUG</u> CAGCCAG ACUGGAUAC UUAAC GUCCGUC U - C C UCCCCUC
miR-C15	UACCACAGGGUAGAACCGGA	- G A UU UC UCCUG CCG UGGUUUACCU UGGUAGG ACG A AGGAC <u>GGC</u> ACCAAGAUGGGA ACCAUUC UGU U A - C - CG
miR-C16	AACUGGCCUACAAAGUCCAG	A U C A AGU GAG GCUGGG CUUUG GGGC AG UGAG G CUC UGACCC <u>GAAAC</u> UCCG UC ACUU U C - U A G A GAC
miR-C17	UGUAACAGCAAUCUCAUGGGA	U AUCGG GUAAACAGCA CUCCAU <u>UGGA</u> CUG G UAGUCU CAUUGUCGU GAGGUG ACCU GGC C U C - UA U
miR-C18	UAGGAGGCACAGAAAUUGGC	U <u>AGCAGCACAG</u> <u>A-</u> AAUAUUGGCA GG G UCGUCCGUGUC UUAUAACCGU CU U GG -- GAG

Fig 7 (cont.)

name	sequence	structure
miR-C19	UAGGUAGUUCAUGUUGGG	<pre> A A C GGCCUGGG GGU GUU AUGUUGUUG CACUAG CCAAA UACAACAC C C U ACAAGUCU </pre>
miR-C20	UUCACCACCUUCACCCAGC	<pre> C A CA GA - A GGCUGUGC GGGU GAGAGGG GUGG GGU AAG G CCCGUACG CCCA CUCUCC CACU CCA UUC C A C AC UC C U </pre>
miR-C21	GGUCCAGAGGGAGAUAGG	<pre> G - C G U UCCUG UCAAU G UC A AGGGAGA AGG AGUAA U AG U UCUCUUCU UCC A A A A - UUUUA </pre>
miR-C22	CCCAGUGUAGACUACUGUU	<pre> AAC U C U G--- G CCAGUGU CAGACUAC UGU CA GAG \ CGG GGUTACA GUCUGAUG ACA GU CUC C AUU C - U GUAA U </pre>
miR-C23	AAAUAACUGCCUGGUAAUGAUGAC	<pre> GGC - C TAGUG GCCGU CAUC UUACUGGGCAG AUUGGA U CGGCA GUAG AAUGGUCCGUC UAAUCU C --- U A CUAGU </pre>
miR-C24	UACUCAGUAAGGCAUTGGUUCU	<pre> U U U UUC A UACCUUAC CAG AAGGCAUUGUUC UAU U AUGGGAUG GUC UCCGUGACAAAG AUA U U U A A </pre>

Fig.7 (cont.)

name	sequence	structure
miR-C25	AGAGGUUAUGGCAUGGGAAAGA	<pre> U A- UG C GUCC UUUCCUAUGC UAUACUTUCU UGGAU \ CGAGG AGAAGGGUACG AUUAGGAGAA AUCUG U U CG -- G </pre>
miR-C26	UGAAAUGUUUAGGACCACUAG	<pre> C U G A C U GGUC AGUGGUUCU GACA UUCA CAGUU UG \ CCAG UCACCAGGA UUGU AAGU GUUAA AC A A U A - C G </pre>
miR-C27	UUCCCCUUUGCUAUCCUAGGCCUG	<pre> U A U GAGAAUA UGGAC UCCCUUUGUC UCCUA GCCU \ ACUTG AGGGAAACGG AGGGU CGGA C A - GGAAGUA </pre>
miR-C28	UCCUCAUUCACCGGAGUCUG	<pre> UC C UCUUA CUCUTUG CUCAUTUCAC GGAGUCUG U GAGGAC GAAGUGAGGUG CUTUAGAC G UC A - CAACC </pre>
miR-C29	GUGAAAUGUUUAGGACCACUAGA	<pre> U C U G A C U GCC GGUC AGUGGUUCU GACA UUCA CAGUU UG \ CGG CCAG UCACCAGGA UUGU AAGU GUUAA AC A C A U A - C G </pre>
miR-C30	UGGAUGUAAGGAAGUGUGGG	<pre> - C U AUUC CCAGG CCACAUUGCUUUAUAU C CAUAG \ GGUUU GGUGUGUGAAGGAUUGUA G GUUAC U A - ACCAC </pre>

Fig 7 (cont.)

name	sequence	structure
miR-C31	UACAGUAGUCUGGACAUUUGGU	AUC U C G GCC CCAGUGU CAGACUAC UGU UCAG A CGG GGUUACA GUCUGAUG ACA GGUC G AUU <u>C</u> - <u>UGUACAG</u> G
miR-C32	CCCUUGUAGAACCGAAUUUGUGU a miR-10 variant	A G C UG- AC UAUAU CCCU UAGAA CGAAUUUGUG GU C AUAAA GGGG AUCCU GCUUAGACAC UA C - A UGA CA
miR-C33	AACCCGUAGAUCCGAACUUGUGA A a miR-99a variant	A ACC C AU GUGU GUAGAU CGA CTUGUG UG U - A UACUG GUU GAACAC AC C - C - GU
miR-C34	GCUUUCUCCUGGGCUCUCCUCUC	C U UUG - GGAG AAGG AGGGG GAGGGG CGGGAGGAGC G UUCC UCUCC CUCCUC GUCCUCUUCG GUUCG C - UCG C GCGU

Fig. 7 (cont)

name	human	C. elegans	mouse						Drosophila	fugu	fish	zebrafish
			liver	small intes	colon	cerebellum	ortex	midbrain				
let-7a-1	AC007924 chr9 AC08784 chr17 identical precursor		num.hits in trace data, 3 families of similar precursors	found			nearly identical precursor	found				
let-7a-2	AF001359 chr11						nearly identical precursor					
let-7a-3	AL049853 chr22 AF274345 chrX with diff. precursor											
let-7b	AL049853 chr22 nearly identical precursor			nearly ident precursor trace#48311003								
let-7c	AP001667 chr21 AC007924.3 chr9 AC08784 chr17 identical		identical and diff. precursors		numerous genomic hits	ident precursor; diff precursor -> EST AI-614897	numerous genomic hits	found				
let-7d				found	trace#83587042	trace#83587042 found	2 nearly ident prec	found				
let-7e	AC018755 chr19							found				
let-7f-1	AC007924 chr9 AC007704 chr17					Ident Precursor genomic DNA		found				
let-7f-2	AI592046 chrX					Ident. precursor in multitrace 187113911						
let-7g	precursor ident. to mouse in AC092045.2 chr3					genomic hits,no EST		found				
let-7h						found in cortex,no db hit						

Fig. 7 (cont.)

		found, supported by EST BB661268	found				
let-7i	precursor ident. to mouse (AL117383.19); also AC018341.22						
mir-1							2L, AE003667
mir-1b	AL419263.5 chr20 nt1-21	U97405.1 nt 1-21 (22G)	no mouse hit (only nt1-21)				
mir-1c				found	found		
mir-1d	AL419263.5 chr20 nt1-22 (23G)				found, but no db hit		
mir-2a-1							
mir-2a-2						2L, AE003663	
mir-2b-1						2L, AE003620	
mir-2b-2						2L, AE003663	
mir-3						2R, AE003795	
mir-4						2R, AE003795	

35/46

Fig. 7 (cont.)

Fig. 7 (cont.)

miR-13b-1						3R, AE003708		
miR-13b-2						X, AE003446		
miR-14						2R, AE003833		
miR-15a	13, AC069475				found	trace#72 137197 prec sig diff		
miR-15b						trace#79 105069		
miR-16	13, AC069475 interesting leukemia locus			genomic hits with 2 slightly diff precursor trace#502 93866, 783168660	found			AL606727 diff precurs
miR-16	3, NT_005740.6		several found	trace#7910506 g murine ident prec. as in human				
miR-17	13, AL138714							
miR-18	13, AL138714							
miR-19a	13, AL138714							
miR-19b-1	13, AL138714				found			G46757 with a U9C

Fig. 7 (cont.)

37/46

mir-19b-2	X, AC002407							
mir-20	13, AL138714		found					
mir-21	17, AC004686	AL604063 found chr11, near ly ident precursor					found	
mir-22	several highly similar ESTs; AW961681 shown	CDNAS from var. tissues, ide ntical precursor	AK008813 (cDNA), prec ident to human precursor				found	
mir-23a	19, AC020916						found trace 62 540691 prec sli diff	Three hits in db
mir-23b	XN_072557..1 chr9, also human ESTs, prec nearly ident to mouse						EST AW124037 AR84865 cerebellum	
mir-24-1	9, AF043896		found				found EST H1286529 (thymus); nearly ident. to mir-24-1; EST AB11466 (whole embryo) different precursor	G46757 similar precursor
mir-24-2	19, AC020916							
mir-25	7, AC073842 second ident copy found in chr7							
mir-26a	3, AP000497						AC055318..9 trf found ace#88411973 precursor diff. from human	Scaffold_ 4097 different precursor
mir-26b	2, AC021016	found					found, trace 6986 6494, slight diff precursor	

Fig. 7 (cont.)

	19, AC020916	found	found, hit	found, but no db hit for mouse	found, but no db hit for mouse	found	found
miR-27a	XM_099943.1 chr9 identical precursor						
miR-27b					found maps to chr 13 MGSC mmtrace 44611617		
miR-28	3, AC063932						
miR-29a	7, AF017104 second ident. copy found in chr7 CLUSTER, this cluster also conserved in mouse; AC02913.32	found, AC024913.3 2	found, AC024913.3 mmtrace#2346734 AC024913.32	nearly ident precursor trace#2346733 4, EST AC024913.32	trace, EST, nearly ident prec		
miR-29b	AL032209.1 chr11 CLUSTER of miR-29-b and 29-c; miRNA similar to miR-83	found		AC024913.32; d iff precursor in EST BG342396 (retina)	FOUND		Scaffold 17610.(A- third copy)
miR-29c				found	found, support by ESTs		Scaffold 17610 has two copies of this RNA
miR-30a-s	nearly ident fold in AL034467.23 chr6	found ESTs 'trace6002 3889 all with 22G	found	found	found		
miR-30a-s	6, AL03467			found with diff. precursor in trace #85261735			
miR-30b	human AF151227.6 chr8 different precursor			trace#72329251	found		Scaffold 3483, diff precursor
miR-30c	AL1316164.8 chr.6 supported by ESTs (BF94736.1)			found, but no db hit for mouse	found	found	

Fig. 7 (cont.)

	A#159227.5 chr8						Found, but no mouse db hit	Scaffold 3463, diff fold
mir-30d	9 , AL353732							
mir-31	9 , AL354797							
mir-32	22 , 299716							
mir-33	AP000962.2 chr21;ident to mouse; similar to mir-10 and mir-51]					trace#4891071 4		G44780 with diff.precu risor
mir-99a	AC018755.3 chr.19; [similar to miR- 10 and miR-51]					mntrace #92340982		
mir-99b	AL358147.17 chr9 diff precursor					AK021366.1 cDNA eyeball	Found	U53213.1 <i>T.fluviat</i> <i>lilis</i>
mir-101								
mir-122a						abundant but no db hit,except woodchuck X13234		
mir-122b								
mir-122a,b								
mir-123						genomic hits (trace#6108 147), no EST		Scaffold_ 3295

Fig. 7 (cont.)

		found in 772504.1 chrIV intron diff precursor	found	most abundant in centab., genomic hits (trace#21097008, 11137241). found, but no db found	most abundant; seve ral trace hits;precurs= centabulum	slightly diff precursor AC009251 chr2L
mir-124a*	nearly ident. precursor in chr8 [AC021518] chr20 [AL096828]					
mir-124b	AC021518 chr8, nearly ident chr20 AL096828.29					
mir-125a	Ident precursor in AC018755.3 chr 19					
mir-125b	AP001359.4 chr11 AP001667.1 chr21 (chr21 like mouse)					
mir-126						
mir-127	human AL17150.6 chr.14 same precurs as in mouse					
mir-128	Ident in AC016743.10 chr 2:diff prec in AC016943.7 chr.3					
mir-129	human AC018662.3 chr7					
mir-130						
mir-131	AC005317.2 chr 15 slight diff precursor, but 5 ident					
mir-132	AL137038.5 chr17 prec slight, diff from mouse					

Fig. 7 (cont.)

		found, trace#	found		
mir-133	AL391221.15 chr6 diff. precursor (ident to rat L33722.1)	AC093440.1 Scaffold_ 1049;prec u nearly like mouse	62407955		
mir-134	AL332709.5 chr14 similar precursor		trace#6462031 1		
mir-135	AC092045.2 chr3 AC016639.35 chr12 (ident or simil to mouse)	trace#7149523 5 ESTB#80195 .1 (kidn., splic en) (=chr3huma n)	found		
mir-136	AL117190.6 chr14 ident to mouse	trace#8607175 3			
mir-137	AC027691.1 chr1 ident to mouse, nearly ident fish	trace#8977454 3,EST (hypothal)A18 52436.1.ident			Scaffold_ 18244 nearly ident to mouse/mam
mir-138	AC006058.1 chr3 precursor diff	mouse EST BB528620.2			
mir-139	AP003065.2 chr11	found, but no mouse hit			
mir-140	AC026468.8 chr.16,precurso r nearly ident,	several trace hits; trace#1053 0393			
mir-141	AC000512.12 chr12,Precursor slightli diff	AC002397 chr6			found
mir-142s	AC0004687.1 chr17 BC11/myq translocation locus,like mouse	found	found		
mir-142as*		several EST AI153235	found		found

Fig. 7 (cont.)

new	AL049829.4 chr14					found but no db hit		
miR-143	AC008681.7 chr5				found	found, but no db hit	found	found
miR-144	XM_064366.1 precursor nearly ident						AJ290206 .1 trace 2143909	
miR-145	AC008681.7 chr5 GG>GA precursor nearly like mouse, see 2 positions above AC008318.7 chr5 diff precursor				found		EST BF163348 .1 lung	Scaffold 934 similar
miR-146	AL592549.7						trace#34 639321	
miR-147	AC010719.4					found		
miR-148							found, no db hit	
miR-149							trace#05 955550	
miR-150								
miR-151	human chr 17 AC04477.1, nearly identical				trace#8472 1065,10352 B01			
					trace#8845 6669			
						found in colon, supported by trace#83700445; close match MGSC in chr18 (additional 14C unlikely, not supported by trace and		

Fig. 7 (cont.)

miR-153	AC003372.2 Chr7 ident.precursor						found sever. mmtrace 87010874
miR-154	AL12709.5 chr4 nearly identical precursor						found sever. mmtrace 86715639
miR-155 [BIC-RNA]	human BIC RNA RP402776.1 (has U12C)				found; chr 16 mouse		

Fig. 7 (cont.)

name	human	mouse	eye	kidney	testes	lung	thymus	skin	Drosophila	fugu fish	zebrafish
	spleen	mouse	trace #76647842		Found				scaffold_1819		
miR-C1	with different precursors in chr9 AL15805.11, chr1 AL136321.5										
miR-C2	chr7 AC084864.2 similar precursor	mouse trace #88841093							scaffold_967	AL590150.2	
miR-C3	chr7 AC084864.2 ident.precursor	trace #86029980							scaffold_967	AL590150.2	
miR-C4	similar precurs.in chr7 AC018662.3	trace #13885686	found								
miR-C5	chr15 AC069082.9	trace #87318220							found	scaffold_3571	
miR-C6	chr22 AC005664.2 ident.precursor	Chr16 AC012526.32									
miR-C7	chr1 AL512443.7 similar prec.	trace #86694995									
miR-C8				found, trace #51673384							
miR-C9					found, trace #78964803						
miR-C10	chrX AF222686.1 nearly ident. precursor				found, trace #6192892						
miR-C11	chr9 XM_098943.1 has C170;prec.nearly identical to mouse				found cDNA AL286659.1, has C170						
miR-C12					found trace#760450				scaffold_2294		
miR-C13		found			found trace #88722637						

Fig. 7 (cont.)

name	human		mouse		Drosophila		fugu fish		zebrafish	
	spleen	eye	kidney	testes	lung	thymus	skin			
miR-C14	chr11 AC000159.6		found, but no db hit							
miR-C15	chr16 AC026468.6 nearly ident.precursor		EST BI687377.1, several trace						scaffold_ 2083	
miR-C16	chr17 AC003101.1, similar precursor		found, trace#95 55103						scaffold_ 246	
miR-C17	chr11 AC000159.6, chr1 AC013590.2; diff.prec.		found, trace #87796602						scaffold_ 152	
miR-C18			found, trace #47833768 (close to miR- 16)	found	found	found	found			
miR-C19	chr17 AC009789.21 cloned from human cell line only								scaffold_ 18334	
miR-C20	chr1 AL355310.19 cloned from human cell line only									
miR-C21	chr3 AC053952.15 cloned from human cell line only									
miR-C22	chr19 AC007229.1; chr1 AL137157.7 similar precursor; cloned from human cell line only								scaffold_ 8399	
miR-C23				trace #72257777	found				scaffold_ 2210	
miR-C24								trace #69879879		
miR-C25								trace #49754566		
miR-C26	All3601 ident. precursor						trace #11977216			

Fig. 7 (cont.)

name	human	mouse						Drosophila	fugu fish	zebrafish
		spleen	eye	kidney	testes	lung	thymus			
miR-C27	chr9 AL159990.12 identical precursor		trace #91503159						scaffold_725	
miR-C28	XN_036612.4, precursor very similar								scaffold_13664	
miR-C29	chr14 AL156001.6 nearly identical precursor								XM_149012.1	
miR-C30	chr6 AL391221.15 similar precursor								trace #18453604	
miR-C31	chr9 AC006312.8								trace #84055510	
miR-C32									scaffold_5830	
miR-C33									trace #83079710	
miR-C34									UTT364_1, intronic location Hoxd4 gene trace #84780544	scaffold_82
									trace #72109322	scaffold_15612

SEQUENCE LISTING

<110> Max-Planck-Gesellschaft zur Förderung der Wissensc

<120> Identification of novel genes encoding for small
temporal RNA

<130> 26250PWO_DR

<140>

<141>

<150> EP 01 123 453.1

<151> 2001-09-28

<150> EP 02 006 712.0

<151> 2002-03-22

<150> EP 02 016 772.2

<151> 2002-07-26

<160> 217

<170> PatentIn Ver. 2.1

<210> 1

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 1

tactatacaa cctactacacct caatggcc

29

<210> 2

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 2

actatgcaac ctactacctc t

21

<210> 3
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 3
actataacaac ctcctacctc a

21

<210> 4
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 4
tggtgtttcc gccccggaa

19

<210> 5
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 5
tggaaatgtaa agaagtatgg ag

22

<210> 6
<211> 23
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 6
gctcctcaaa gctggctgtg ata

23

<210> 7
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 7
tgagacacac tttgccagt ga

22

<210> 8
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 8
tcaatggttg tctagcttta t

21

<210> 9
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 9
catatatcacaa cgatcggtcc ttt

23

<210> 10
<211> 22

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 10
aaaaagaaca gccactgtga ta

22

<210> 11
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 11
tggaagacta gtgattttgt tgt

23

<210> 12
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 12
gacatcttta cctgacagta tta

23

<210> 13
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 13
tcatacagct agataaccaa aga

23

<210> 14
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 14
acaaatttcgg atctacaggg t

21

<210> 15
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 15
gcaagaactc agactgtgat g

21

<210> 16
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 16
accagtacct gatgtaatac tca

23

<210> 17
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:

Oligonucleotide

<400> 17
actcgtaaaa atggctgtga ta 22

<210> 18
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 18
taggagagag aaaaagactg a 21

<210> 19
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 19
tagcagcacaca taatggtttg t 21

<210> 20
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 20
gccaaatattt acgtgctgct a 21

<210> 21
<211> 22
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 21

tacaagtgcc ttcactgcag ta

22

<210> 22

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 22

tatctgcact agatgcacct ta

22

<210> 23

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 23

ttagtttgc atagatttgc aca

23

<210> 24

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 24

tacctgcact ataaggactt ta

22

<210> 25
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 25
tcaacatca g tctgataa gc ta

22

<210> 26
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 26
acagttcttc aactggcagc tt

22

<210> 27
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 27
ggaaatccct ggcaatgtga t

21

<210> 28
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 28

ctgttcctgc tgaactgagc ca

22

<210> 29

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 29

tcaagaccgag acaaggtaaa tg

22

<210> 30

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 30

agcctatcct ggattacttg aa

22

<210> 31

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 31

agcgaaacctt agccactgtg aa

22

<210> 32

<211> 22

<212> DNA

<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 32
ctcaatagac tgtgagctcc tt 22

<210> 33
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 33
aacccgatttc agatggtgct ag 22

<210> 34
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 34
gctgcaaaca tccgactgaa ag 22

<210> 35
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 35
cagctatgcc agcatcttgc ct 22

<210> 36
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 36
gcaacttagt aatgtgcaat a 21

<210> 37
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 37
tgcaatgcaa ctacaatgca cc 22

<210> 38
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 38
ctccatactt ctttacattc ca 22

<210> 39
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 39
gctgagtgta ggatgttac a 21

<210> 40
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 40
gcttccagtc gaggatgtt aca 23

<210> 41
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 41
cgcaaggtcg gttctacggg tg 22

<210> 42
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 42
tcagttatca cagtactgta 20

<210> 43
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 43
acaaaacacca ttgtcacact cca

23

<210> 44
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 44
tggcattcac cgctgcctt a

21

<210> 45
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 45
cacaggttaa aggtctcag gga

23

<210> 46
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 46
tcacaagtta ggtctcagg ga

22

<210> 47

<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 47
agccaaaggctc agacggatcc ga

22

<210> 48
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 48
aaaagagacc ggttcactct ga

22

<210> 49
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 49
gcaagcccaag accgaaaaaaaa g

21

<210> 50
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 50

gcccttttaa cattgcactc

20

<210> 51
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 51
actttcggtt atctagctt a

21

<210> 52
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 52
acgaccatgg ctgttagactg tta

23

<210> 53
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Combined DNA/RNA Molecule:
Oligonucleotide

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 53
tgagctacag tgcttcatct ca

22

<210> 54
<211> 18

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 54
uuuaaccgcg aattccag

18

<210> 55
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 55
acggaattcc tcactaaa

18

<210> 56
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 56
cacaggttaa agggtctcag gga

23

<210> 57
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
Oligonucleotide

<400> 57
cagccaacgg aattcctcac taaa

24

<210> 58
<211> 22
<212> RNA
<213> D. melanogaster

<400> 58
uggaauguaa agaaguaugg ag

22

<210> 59
<211> 23
<212> RNA
<213> D. melanogaster

<400> 59
uaucacagcc agcuuugaug agc

23

<210> 60
<211> 23
<212> RNA
<213> D. melanogaster

<400> 60
uaucacagcc agcuuugagg agc

23

<210> 61
<211> 22
<212> RNA
<213> D. melanogaster

<400> 61
ucacugggca aagugugucu ca

22

<210> 62
<211> 21
<212> RNA
<213> D. melanogaster

<400> 62
auaaagcuag acaaccauug a

21

<210> 63

<211> 23
<212> RNA
<213> D. melanogaster

<400> 63
aaaggaacga ucguugugau aug

23

<210> 64
<211> 22
<212> RNA
<213> D. melanogaster

<400> 64
uaucacagug gcuguucuuu uu

22

<210> 65
<211> 23
<212> RNA
<213> D. melanogaster

<400> 65
uggaagacua gugauuuugu ugu

23

<210> 66
<211> 23
<212> RNA
<213> D. melanogaster

<400> 66
uaauuacuguc agguaaagau guc

23

<210> 67
<211> 23
<212> RNA
<213> D. melanogaster

<400> 67
ucuuuggua ucuagcugua uga

23

<210> 68
<211> 21
<212> RNA
<213> D. melanogaster

<400> 68
acccuguaga uccgaaauuug u 21

<210> 69
<211> 21
<212> RNA
<213> D. melanogaster

<400> 69
caucacaguc ugaguucuug c 21

<210> 70
<211> 23
<212> RNA
<213> D. melanogaster

<400> 70
ugaguauuac aucagguacu ggu 23

<210> 71
<211> 22
<212> RNA
<213> D. melanogaster

<400> 71
uaucacagcc auuuugacga gu 22

<210> 72
<211> 22
<212> RNA
<213> D. melanogaster

<400> 72
uaucacagcc auuuugauga gu 22

<210> 73
<211> 21
<212> RNA
<213> D. melanogaster

<400> 73
ucagucuuuu ucucucuccu a 21

<210> 74
<211> 22
<212> RNA
<213> D. melanogaster

<400> 74
ugagguagua gguuguauag uu

22

<210> 75
<211> 22
<212> RNA
<213> Human

<400> 75
ugagguagua gguuguauag uu

22

<210> 76
<211> 22
<212> RNA
<213> Human

<400> 76
ugagguagua gguugugugg uu

22

<210> 77
<211> 22
<212> RNA
<213> Human

<400> 77
ugagguagua gguuguaugg uu

22

<210> 78
<211> 21
<212> RNA
<213> Human

<400> 78
agagguagua gguugcauag u

21

<210> 79

<211> 21
<212> RNA
<213> Human

<400> 79
ugagguagga gguuguauag u 21

<210> 80
<211> 22
<212> RNA
<213> Human

<400> 80
ugagguagua gauuguauag uu 22

<210> 81
<211> 22
<212> RNA
<213> Human

<400> 81
uagcagcaca uaaugguuug ug 22

<210> 82
<211> 22
<212> RNA
<213> Human

<400> 82
uagcagcacg uaaaauauugg cg 22

<210> 83
<211> 20
<212> RNA
<213> Human

<400> 83
acugcaguga aggcacuugu 20

<210> 84
<211> 22
<212> RNA
<213> Human

<400> 84
uaaggugcau cuagugcaga ua 22

<210> 85
<211> 23
<212> RNA
<213> Human

<400> 85
ugugcaaauc uaugcaaaac uga 23

<210> 86
<211> 23
<212> RNA
<213> Human

<400> 86
ugugcaaauc caugcaaaac uga 23

<210> 87
<211> 22
<212> RNA
<213> Human

<400> 87
uaaagugcuu auagugcagg ua 22

<210> 88
<211> 22
<212> RNA
<213> Human

<400> 88
uagcuaauca gacugauguu ga 22

<210> 89
<211> 22
<212> RNA
<213> Human

<400> 89
aagcugccag uugaagaacu gu 22

<210> 90
<211> 21
<212> RNA
<213> Human

<400> 90
aucacauugc cagggauuuc c 21

<210> 91
<211> 22
<212> RNA
<213> Human

<400> 91
uggcucaguu cagcaggaac ag 22

<210> 92
<211> 22
<212> RNA
<213> Human

<400> 92
cauugcacuu gucucggucu ga 22

<210> 93
<211> 22
<212> RNA
<213> Human

<400> 93
uucaaguaau ccaggauagg cu 22

<210> 94
<211> 22
<212> RNA
<213> Human

<400> 94
uucaaguaau ucaggauagg uu 22

<210> 95

<211> 22
<212> RNA
<213> Human

<400> 95
uucacagugg cuaaguuccg cu 22

<210> 96
<211> 22
<212> RNA
<213> Human

<400> 96
aaggagcuca cagucuauug ag 22

<210> 97
<211> 22
<212> RNA
<213> Human

<400> 97
cuagcaccau cugaaaucgg uu 22

<210> 98
<211> 22
<212> RNA
<213> Human

<400> 98
cuuucagucg gauguuugca gc 22

<210> 99
<211> 21
<212> RNA
<213> Human

<400> 99
ggcaagaugc uggcauagcu g 21

<210> 100
<211> 21
<212> RNA
<213> Human

<400> 100
uauugcacau uacuaaguug c

21

<210> 101
<211> 19
<212> RNA
<213> Human

<400> 101
gugcauugua guugcauug

19

<210> 102
<211> 22
<212> RNA
<213> Human

<400> 102
uggaaauguaa agaaguaugg ag

22

<210> 103
<211> 23
<212> RNA
<213> Human

<400> 103
uggaagacua gugauuuugu ugu

23

<210> 104
<211> 23
<212> RNA
<213> Human

<400> 104
ucuuugguua ucuagcugua uga

23

<210> 105
<211> 21
<212> RNA
<213> Human

<400> 105
acccuguaaga uccgaauuug u

21

<210> 106
<211> 22
<212> RNA
<213> Mouse

<400> 106
ugagguagua gguuguauag uu

22

<210> 107
<211> 22
<212> RNA
<213> Mouse

<400> 107
ugagguagua gguugugugg uu

22

<210> 108
<211> 22
<212> RNA
<213> Mouse

<400> 108
ugagguagua gguuguaagg uu

22

<210> 109
<211> 21
<212> RNA
<213> Mouse

<400> 109
agagguagua gguugcauag u

21

<210> 110
<211> 21
<212> RNA
<213> Mouse

<400> 110
ugagguaggga gguuguauag u

21

<210> 111

<211> 22
<212> RNA
<213> Mouse

<400> 111
ugagguagua gauuguaauag uu

22

<210> 112
<211> 22
<212> RNA
<213> Mouse

<400> 112
ugagguagua guuuguacag ua

22

<210> 113
<211> 22
<212> RNA
<213> Mouse

<400> 113
ugagguagua guguguacag uu

22

<210> 114
<211> 19
<212> RNA
<213> Mouse

<400> 114
ugagguagua guuugugcu

19

<210> 115
<211> 22
<212> RNA
<213> Mouse

<400> 115
uggaaauguaa agaaguaugu aa

22

<210> 116
<211> 22
<212> RNA
<213> Mouse

<400> 116
uggaauguaa agaaguaugu ac 22

<210> 117
<211> 23
<212> RNA
<213> Mouse

<400> 117
uggaauguaa agaaguaugu auu 23

<210> 118
<211> 23
<212> RNA
<213> Mouse

<400> 118
ucuuuggua ucuagcugua uga 23

<210> 119
<211> 22
<212> RNA
<213> Mouse

<400> 119
uagcagcac a uaugguuug ug 22

<210> 120
<211> 22
<212> RNA
<213> Mouse

<400> 120
uagcagcac a ucaugguuua ca 22

<210> 121
<211> 22
<212> RNA
<213> Mouse

<400> 121
uagcagcacg uaaaauauugg cg 22

<210> 122
<211> 22
<212> RNA
<213> Mouse

<400> 122
uaaggugcau cuagugcaga ua

22

<210> 123
<211> 23
<212> RNA
<213> Mouse

<400> 123
uggugcaaauc caugcaaaac uga

23

<210> 124
<211> 23
<212> RNA
<213> Mouse

<400> 124
uaaagugcuu auagugcagg uag

23

<210> 125
<211> 22
<212> RNA
<213> Mouse

<400> 125
uagcuuauca gacugauguu ga

22

<210> 126
<211> 22
<212> RNA
<213> Mouse

<400> 126
aagcugccag uugaagaacu gu

22

<210> 127

<211> 21
<212> RNA
<213> Mouse

<400> 127
aucacauugc cagggauuuc c 21

<210> 128
<211> 23
<212> RNA
<213> Mouse

<400> 128
aucacauugc cagggauuac cac 23

<210> 129
<211> 22
<212> RNA
<213> Mouse

<400> 129
uggcucaguu cagcaggaac ag 22

<210> 130
<211> 22
<212> RNA
<213> Mouse

<400> 130
...uuacaguaau ccaggauagg cu 22

<210> 131
<211> 22
<212> RNA
<213> Mouse

<400> 131
uucaaguaaau ucaggauagg uu 22

<210> 132
<211> 22
<212> RNA
<213> Mouse

<400> 132
uucacagugg cuaaguuccg cu 22

<210> 133
<211> 20
<212> RNA
<213> Mouse

<400> 133
uucacagugg cuaaguucug 20

<210> 134
<211> 22
<212> RNA
<213> Mouse

<400> 134
cuagcaccau cugaaaucgg uu 22

<210> 135
<211> 23
<212> RNA
<213> Mouse

<400> 135
uagcaccauu ugaaaucagu guu 23

<210> 136
<211> 22
<212> RNA
<213> Mouse

<400> 136
uagcaccauu ugaaaucggu ua 22

<210> 137
<211> 23
<212> RNA
<213> Mouse

<400> 137
uguaaaacauc cucgacugga agc 23

<210> 138
<211> 22
<212> RNA
<213> Mouse

<400> 138
cuuucagucg gauguuugca gc

22

<210> 139
<211> 21
<212> RNA
<213> Mouse

<400> 139
uguaaacauc cuacacucag c

21

<210> 140
<211> 23
<212> RNA
<213> Mouse.

<400> 140
uguaaacauc cuacacucuc agc

23

<210> 141
<211> 22
<212> RNA
<213> Mouse

<400> 141
uguaaacauc cccgacugga ag

22

<210> 142
<211> 20
<212> RNA
<213> Mouse

<400> 142
acccguagau ccgaucuugu

20

<210> 143

<211> 22
<212> RNA
<213> Mouse

<400> 143
cacccguaga accgaccuug cg

22

<210> 144
<211> 20
<212> RNA
<213> Mouse

<400> 144
uacaguacug ugauuaacuga

20

<210> 145
<211> 23
<212> RNA
<213> Mouse

<400> 145
uggaguguga caaugguguu ugu

23

<210> 146
<211> 23
<212> RNA
<213> Mouse

<400> 146
uggaguguga caaugguguu uga

23

<210> 147
<211> 22
<212> RNA
<213> Mouse

<400> 147
uggaguguga caaugguguu ug

22

<210> 148
<211> 21
<212> RNA
<213> Mouse

<400> 148
cauuauuacu uuuggguacgc g

21

<210> 149
<211> 22
<212> RNA
<213> Mouse

<400> 149
uuaaggcacg cggugaaugc ca

22

<210> 150
<211> 21
<212> RNA
<213> Mouse

<400> 150
uuaaggcacg cgggugaaug c

21

<210> 151
<211> 23
<212> RNA
<213> Mouse

<400> 151
ucccugagac ccuuuaaccu gug

23

<210> 152
<211> 22
<212> RNA
<213> Mouse

<400> 152
ucccugagac ccuaacuugu ga

22

<210> 153
<211> 21
<212> RNA
<213> Mouse

<400> 153
ucguaccgug aguaauaaug c

21

<210> 154
<211> 22
<212> RNA
<213> Mouse

<400> 154
ucggauccgu cugagcuugg cu

22

<210> 155
<211> 22
<212> RNA
<213> Mouse

<400> 155
ucacagugaa ccggucucuu uu

22

<210> 156
<211> 21
<212> RNA
<213> Mouse

<400> 156
cuuuuuucgg ucugggcuug c

21

<210> 157
<211> 20
<212> RNA
<213> Mouse

<400> 157
cagugcaaug uuaaaaaggc

20

<210> 158
<211> 21
<212> RNA
<213> Mouse

<400> 158
uaaagcuaga uaaccgaaag u

21

<210> 159

<211> 23
<212> RNA
<213> Mouse

<400> 159
uaaacagucua cagccauuggu cgu

23

<210> 160
<211> 22
<212> RNA
<213> Mouse

<400> 160
uugguccccu ucaaccagcu gu

22

<210> 161
<211> 22
<212> RNA
<213> Mouse

<400> 161
ugugacuggu ugaccagagg ga

22

<210> 162
<211> 24
<212> RNA
<213> Mouse

<400> 162
uauggcuuuu uauuccuaug ugaa

24

<210> 163
<211> 23
<212> RNA
<213> Mouse

<400> 163
acuccauuuug uuuugaugau gga

23

<210> 164
<211> 22
<212> RNA
<213> Mouse

<400> 164
uauugcuuaa gaauacgcgu ag 22

<210> 165
<211> 17
<212> RNA
<213> Mouse

<400> 165
agcugguguu gugaauc 17

<210> 166
<211> 18
<212> RNA
<213> Mouse

<400> 166
ucuacagugc acgugucu 18

<210> 167
<211> 21
<212> RNA
<213> Mouse

<400> 167
agugguuuuua cccuauggua g 21

<210> 168
<211> 21
<212> RNA
<213> Mouse

<400> 168
aacacugucu gguaaaagaug g 21

<210> 169
<211> 20
<212> RNA
<213> Mouse

<400> 169
cauaaaguag aaaggcacuac 20

<210> 170
<211> 22
<212> RNA
<213> Mouse

<400> 170
uguaguguuu ccuacuuuau gg

22

<210> 171
<211> 22
<212> RNA
<213> Mouse

<400> 171
ugagaugaag cacuguagcu ca

22

<210> 172
<211> 22
<212> RNA
<213> Mouse

<400> 172
uacaguauag augauguacu ag

22

<210> 173
<211> 24
<212> RNA
<213> Mouse

<400> 173
guccaguuuu cccagggaauc ccuu

24

<210> 174
<211> 23
<212> RNA
<213> Mouse

<400> 174
ugagaacuga auuccauggg uuu

23

<210> 175

<211> 21
<212> RNA
<213> Mouse

<400> 175
guguguggaa augcuucugc c

21

<210> 176
<211> 22
<212> RNA
<213> Mouse

<400> 176
ucagugcacu acagaacuuu gu

22

<210> 177
<211> 22
<212> RNA
<213> Mouse

<400> 177
ucuggccuccg ugucuucacu cc

22

<210> 178
<211> 23
<212> RNA
<213> Mouse

<400> 178
ucucccaacc cuuguaccag ugu

23

<210> 179
<211> 22
<212> RNA
<213> Mouse

<400> 179
cuagacugag gcuccuugag gu

22

<210> 180
<211> 21
<212> RNA
<213> Mouse

<400> 180
ucagugcaug acagaacuug g 21

<210> 181
<211> 20
<212> RNA
<213> Mouse

<400> 181
uugcauaguc aaaaaaguga 20

<210> 182
<211> 22
<212> RNA
<213> Mouse

<400> 182
uagguuaucc guguugccuu cg 22

<210> 183
<211> 22
<212> RNA
<213> Mouse

<400> 183
uuaaugcuaa uugugauagg gg 22

<210> 184
<211> 23
<212> RNA
<213> Human/Mouse

<400> 184
aacauucaac gcugucggug agu 23

<210> 185
<211> 22
<212> RNA
<213> Human/Mouse

<400> 185
uuuggcaaung guagaacuca ca 22

<210> 186

<211> 23

<212> RNA

<213> Human/Mouse

<400> 186

uauggcacug guagaauuca cug

23

<210> 187

<211> 22

<212> RNA

<213> Human/Mouse

<400> 187

cuuuuuugcgg ucuggggcuug uu

22

<210> 188

<211> 22

<212> RNA

<213> Human/Mouse

<400> 188

uggacggaga acugauaagg gu

22

<210> 189

<211> 18

<212> RNA

<213> Human/Mouse

<400> 189

uggagagaaa ggcaguuc

18

<210> 190

<211> 23

<212> RNA

<213> Human/Mouse

<400> 190

caaagaauuc uccuuuuggg cuu

23

<210> 191

<211> 22
<212> RNA
<213> Human/Mouse

<400> 191
ucgugucuuug uguugcagcc gg

22

<210> 192
<211> 21
<212> RNA
<213> Human/Mouse

<400> 192
uaacacuguc ugguaacgau g

21

<210> 193
<211> 22
<212> RNA
<213> Human/Mouse

<400> 193
cauccccuugc augguggagg gu

22

<210> 194
<211> 23
<212> RNA
<213> Human/Mouse

<400> 194
gugccuacug agcugacauc agu

23

<210> 195
<211> 22
<212> RNA
<213> Human/Mouse

<400> 195
ugauauguuu gauauauuag gu

22

<210> 196
<211> 22
<212> RNA
<213> Human/Mouse

<400> 196
caacggaauc ccaaaaagcag cu 22

<210> 197
<211> 18
<212> RNA
<213> Human/Mouse

<400> 197
cugaccuaug aauugaca 18

<210> 198
<211> 22
<212> RNA
<213> Human/Mouse

<400> 198
uaccacaggg uagaaccacg ga 22

<210> 199
<211> 21
<212> RNA
<213> Human/Mouse

<400> 199
aacuggccua caaaguccca g 21

<210> 200
<211> 22
<212> RNA
<213> Human/Mouse

<400> 200
uguaacagca acuccaughug ga 22

<210> 201
<211> 21
<212> RNA
<213> Human/Mouse

<400> 201
uagcagcaca gaaaauauugg c 21

<210> 202
<211> 21
<212> RNA
<213> Human/Mouse

<400> 202
uagguaguuu cauguuguug g

21

<210> 203
<211> 22
<212> RNA
<213> Human/Mouse

<400> 203
uucaccacccu ucuccaccca gc

22

<210> 204
<211> 19
<212> RNA
<213> Human/Mouse

<400> 204
gguccagagg ggagauagg

19

<210> 205
<211> 22
<212> RNA
<213> Human/Mouse

<400> 205
cccaguguuc agacuaccug uu

22

<210> 206
<211> 23
<212> RNA
<213> Human/Mouse

<400> 206
uaauacugcc ugguaaugau gac

23

<210> 207

<211> 21

<212> RNA

<213> Human/Mouse

<400> 207

uacucaguuaa ggcauuguuc u

21

<210> 208

<211> 22

<212> RNA

<213> Human/Mouse

<400> 208

agagguauaag cgcaugggaa ga

22

<210> 209

<211> 21

<212> RNA

<213> Human/Mouse

<400> 209

ugaaauguuu aggaccacua g

21

<210> 210

<211> 23

<212> RNA

<213> Human/Mouse

<400> 210

uucccuuugu cauccuaugc cug

23

<210> 211

<211> 22

<212> RNA

<213> Human/Mouse

<400> 211

uccuucauuc caccggaguc ug

22

<210> 212

<211> 23

<212> RNA

<213> Human/Mouse

<400> 212
gugaaauguu uaggaccacu aga

23

<210> 213
<211> 22
<212> RNA
<213> Human/Mouse

<400> 213
uggaauguaa ggaagugugu gg

22

<210> 214
<211> 22
<212> RNA
<213> Human/Mouse

<400> 214
uacaguaguc ugcacauugg uu

22

<210> 215
<211> 22
<212> RNA
<213> Human/Mouse

<400> 215
cccuguagaa ccgaauuugu gu

22

<210> 216
<211> 24
<212> RNA
<213> Human/Mouse

<400> 216
aacccguaga uccgaacuug ugaa

24

<210> 217
<211> 23
<212> RNA
<213> Human/Mouse

<400> 217
gcuucuccug gcucuccucc cuc

23