




(86) Date de dépôt PCT/PCT Filing Date: 2015/06/05
(87) Date publication PCT/PCT Publication Date: 2015/12/10
(45) Date de délivrance/Issue Date: 2023/09/19
(85) Entrée phase nationale/National Entry: 2016/11/24
(86) N° demande PCT/PCT Application No.: US 2015/034372
(87) N° publication PCT/PCT Publication No.: 2015/188046
(30) Priorités/Priorities: 2014/06/06 (US62/009,145);
2014/06/30 (US62/019,289)

(51) Cl.Int./Int.Cl. *A61K 31/202* (2006.01),
A61P 9/00 (2006.01)
(72) Inventeurs/Inventors:
SANDNES, OLAV, US;
MILLER, BRUCE A., US;
DYERBERG, JORN, DK;
STENDER, STEEN, DK
(73) Propriétaire/Owner:
MARINE INGREDIENTS, LLC, US
(74) Agent: CPST INTELLECTUAL PROPERTY INC.

(54) Titre : COMPOSITIONS D'OMEGA -3, FORMES POSOLOGIQUES ET PROCEDES D'UTILISATION
(54) Title: OMEGA-3 COMPOSITIONS, DOSAGE FORMS, AND METHODS OF USE



(57) **Abrégé/Abstract:**

Omega-3 compositions, dosage forms, and methods of use are disclosed herein. The omega-3 compositions and dosage forms disclosed herein may comprise DHA and EPA at a ratio of about 5:2. The omega-3 compositions and dosage forms disclosed herein may comprise re- esterified triglycerides. Methods of treatment using the compositions and dosage forms for reducing the resting heart rate of subject are also disclosed.

ABSTRACT

Omega-3 compositions, dosage forms, and methods of use are disclosed herein. The omega-3 compositions and dosage forms disclosed herein may comprise DHA and EPA at a ratio of about 5:2. The omega-3 compositions and dosage forms disclosed herein may comprise re-esterified triglycerides. Methods of treatment using the compositions and dosage forms for reducing the resting heart rate of subject are also disclosed.

OMEGA-3 COMPOSITIONS, DOSAGE FORMS, AND METHODS OF USE

RELATED APPLICATIONS

[0001] This application claims priority to both United States Provisional Application No. 62/009,145 filed on June 6, 2014 and titled "OMEGA-3 COMPOSITIONS, DOSAGE FORMS, AND METHODS OF USE" and United States Provisional Application No. 62/019,289 filed on June 30, 2014 and titled "OMEGA-3 COMPOSITIONS, DOSAGE FORMS, AND METHODS OF USE."

TECHNICAL FIELD

[0002] Embodiments of the invention relate generally to medicinal compositions, and more particularly to omega-3 compositions, dosage forms, and methods of use.

BACKGROUND

[0003] Mammals have only a limited ability to synthesize omega-3 fatty acids, and thus generally rely on other sources to obtain 5Z,8Z,11Z,14Z,17Z-eicosapentaenoic acid (EPA) and 4Z,7Z,10Z,13Z,16Z,19Z-docosahexaenoic acid (DHA).

[0004] Omega-3 fatty acids are associated with numerous health benefits, including protection against heart disease. Some studies have identified potential benefits for other conditions as well, including autoimmune diseases, inflammatory bowel disease, and cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

[0005] FIG. 1 is a box-and-whisker plot showing changes in blood serum triglyceride levels during the study disclosed in Example 1.

[0006] FIG. 2 is a bar graph of median non-fasting triglyceride levels as measured at the beginning and at the end of the study disclosed in Example 1.

[0007] FIG. 3 is a bar graph depicting changes in median non-fasting triglyceride levels as measured at the beginning and at the end of the study disclosed in Example 1.

[0008] FIG. 4 is a bar graph of median corrected non-fasting triglyceride levels as measured at the beginning and at the end of the study disclosed in Example 1.

[0009] FIG. 5 is a bar graph depicting changes in median corrected non-fasting triglyceride levels as measured at the beginning and at the end of the study disclosed in Example 1.

[0010] FIG. 6 is a bar graph providing hazard ratios for all-cause mortality in each experimental arm of the study disclosed in Example 1.

[0011] FIG. 7 is a box-and-whisker plot of changes in omega-3 index over the course of the study disclosed in Example 1.

[0012] FIG. 8 is a bar graph depicting the mean omega-3 index for each arm of the study disclosed in Example 1, both before and after treatment.

[0013] FIG. 9 is a bar graph depicting changes in omega-3 index over the course of the study disclosed in Example 1.

[0014] FIG. 10 is the bar graph depicting the mean dose-corrected omega-3 index for each arm of the study disclosed in Example 1, both before and after treatment.

[0015] FIG. 11 is a bar graph depicting dose-corrected changes in omega-3 index over the course of the study disclosed in Example 1.

[0016] FIG. 12 is a scatter plot of the change in triglyceride levels over the course of the study disclosed in Example 1, and the change in omega-3 index observed over the course of the same study.

[0017] FIG. 13 is a bar graph depicting mean EPA levels in red blood cells both at the beginning and at the conclusion of the study disclosed in Example 1.

[0018] FIG. 14 is a bar graph depicting mean DHA levels in red blood cells at the beginning and at the conclusion of the study disclosed in Example 1.

[0019] FIG. 15 is a box-and-whisker plot of the change in heart rate observed over the course of the study disclosed in Example 1.

[0020] FIG. 16 is a bar graph depicting mean changes in heart rate for each arm of the study disclosed in Example 1.

[0021] FIG. 17 is a box-and-whisker plot depicting changes in high-density lipoprotein-cholesterol concentration over the course of the study disclosed in Example 1.

[0022] FIG. 18 is a bar graph depicting mean high-density lipoprotein-cholesterol concentration at the beginning and at the conclusion of the study disclosed in Example 1.

[0023] FIG. 19 is a bar graph depicting changes in mean high-density lipoprotein-cholesterol concentration over the course of the study disclosed in Example 1.

[0024] FIG. 20 is a bar graph of mean non-high-density lipoprotein-cholesterol concentration for each arm of the study disclosed in Example 1.

[0025] FIG. 21 is a bar graph depicting mean total cholesterol levels for each arm of the study disclosed in Example 1.

[0026] FIG. 22 is a bar graph depicting mean apolipoprotein B concentrations for each arm of the study disclosed in Example 1.

[0027] FIG. 23 is a bar graph depicting mean changes in apolipoprotein B concentrations over the course of the study disclosed in Example 1.

[0028] FIG. 24 is a bar graph depicting the ratio of apolipoprotein B to apolipoprotein A for each arm of the study disclosed in Example 1.

[0029] FIG. 25 is a bar graph depicting the changes in the apolipoprotein B/apolipoprotein A ratio across the course of the study disclosed in Example 1.

[0030] FIG. 26 is a bar graph depicting the systolic blood pressure in each arm of the study disclosed in Example 1.

[0031] FIG. 27 is a bar graph depicting the diastolic blood pressure in each arm of the study disclosed in Example 1.

[0032] FIG. 28 is a bar graph depicting the mean concentration of C-reactive protein in each arm of the study disclosed in Example 1.

[0033] FIG. 29 is a bar graph depicting the percent change of various measurements and values determined over the course of the study disclosed in Example 1.

[0034] FIG. 30 is a chromatogram of oil used in the re-esterified triglyceride arm of the study disclosed in Example 1.

DETAILED DESCRIPTION

[0035] The following detailed description of various embodiments is not intended to limit the scope of the present disclosure, but is merely representative of various embodiments. The present disclosure relates to medicinal compositions, and more particularly, to omega-3 compositions, dosage forms, and methods of use.

[0036] The term “about,” when used with reference to ratios, amounts, or percentages of one or more elements of a composition, dosage form, or portion thereof, encompasses both the actual ratios, amounts, and percentages of the elements (as measured) and the ratios, amounts, and percentages after correction using standard correction methods (e.g., to compensate for the flame ionization detection response for each component). For example, a marine oil comprising fatty acids, including omega-3 polyunsaturated fatty acids (PUFAs), wherein at least about 50% of the fatty acids are triglycerides (TGs), encompasses marine oils where at least about 50% of the fatty acids, on either a corrected or uncorrected basis, are TGs.

[0037] Some compositions disclosed herein comprise a marine oil that comprises fatty acids, including omega-3 polyunsaturated fatty acids (PUFAs). In these compositions, 4Z,7Z,10Z,13Z,16Z,19Z-docosahexaenoic acid (DHA) and 5Z,8Z,11Z,14Z,17Z-eicosapentaenoic acid (EPA) comprise at least a portion of the PUFAs, and the ratio of DHA to EPA may be about 3:2 to about 4:1. For example, in some compositions, the ratio of DHA to EPA is about 5:2.

[0038] In some compositions, at least about 50% (more particularly about 55–70%) of the fatty acids are triglycerides (TGs). In some embodiments, at least about 55% of the fatty acids are TGs.

[0039] In some compositions, about 25% to about 45% of the fatty acids are diglycerides (DGs). In some compositions, about 0% to about 5% of the fatty acids are monoglycerides (MGs). In some compositions, about 5% or less (or, more particularly, about 1% or less) of the fatty acids are ethyl esters. In some compositions, about 60% of the fatty acids are PUFAs. In some compositions, at least about 55% of the fatty acids are omega-3 PUFAs. In some compositions, at least about 50% of the fatty acids are either DHA or EPA.

[0040] In some compositions, about 0% to about 35% of the fatty acids are mono-unsaturated fatty acids. In some compositions, about 0% to about 5% of the fatty acids are saturated fatty acids. In some compositions, at least about 80% of the PUFAs are DHA and EPA. In some compositions at least about 15% of the marine oil is derived from cephalopod oil. In some compositions, about 10% to about 25% of the marine oil is derived from cephalopod oil and the remainder is derived from fish oil. In some compositions about 25% to about 50% of the marine oil is derived from cephalopod oil and the remainder is derived from fish oil. In some compositions, at least about 50% of the marine oil is derived from cephalopod oil.

[0041] In some compositions, the ratio of DHA to EPA is about 5:2, about 55% to about 70% of the fatty acids are re-esterified triglycerides (rTGs), and about 25% to about 45% of the fatty acids are re-esterified diglycerides (rDGs).

[0042] In the embodiments described herein, a portion or all of the triglycerides, diglycerides, and/or monoglycerides may be re-esterified. For example, in some compositions, at least about 50% of the fatty acids are rTGs.

[0043] Some dosage forms disclosed herein comprise a marine oil comprising fatty acids, including omega-3 PUFAs, wherein DHA and EPA comprise at least a

portion of the PUFAs. In such dosage forms, the amount of DHA may be at least about 450 mg and the amount of EPA may be at least about 150 mg.

[0044] In some dosage forms, the ratio of DHA to EPA may be about 3:2 to about 4:1 (e.g., about 5:2).

[0045] In some dosage forms, at least about 50% of the fatty acids are triglycerides.

[0046] In some dosage forms, at least about 50% of the fatty acids are re-esterified triglycerides.

[0047] In some dosage forms, the amount of DHA is about 450 mg to about 700 mg. In some dosage forms, the amount of EPA is about 150 mg to about 300 mg.

[0048] In some embodiments, the dosage form is a capsule, such as a liquid-filled capsule, where the liquid is a marine oil disclosed herein.

[0049] In some embodiments, the dosage form is a liquid, such as delivered in a 150 milliliter bottle. In such embodiments, the marine oil may be dispersed in a flavored carrier.

[0050] In some embodiments, the mass of the dosage form is about 1200 mg to about 1600 mg.

[0051] In some dosage forms, at least about 55% of the fatty acids are TGs. In some dosage forms, about 55% to about 70% of the fatty acids are TGs.

[0052] In some dosage forms, about 25% to about 45% of the PUFAs are DGs.

[0053] In some dosage forms, about 0% to about 5% of the fatty acids are MGs.

[0054] In some dosage forms, about 5% or less (e.g., 1% or less) of the fatty acids are ethyl esters. In some dosage forms, less than about 1% of the fatty acids are ethyl esters.

[0055] In some dosage forms, at least about 60% of the fatty acids are PUFAs.

[0056] In some dosage forms, at least about 55% of the fatty acids are omega-3 PUFAs.

[0057] In some dosage forms, at least about 50% of the fatty acids are omega-3 PUFAs.

[0058] In some dosage forms, about 0% to about 35% of the fatty acids are mono-unsaturated fatty acids.

[0059] In some dosage forms, about 0% to about 5% of the fatty acids are saturated fatty acids.

[0060] In some dosage forms, at least about 80% of the PUFAs are DHA and EPA.

[0061] In some dosage forms, at least about 15% (e.g., at least about 50%) of the marine oil is derived from cephalopod oil. In some dosage forms, about 10% to about 25% of the marine oil is derived from cephalopod oil and the remainder is derived from fish oil. In some dosage forms, about 25% to about 50% of the marine oil is derived from cephalopod oil and the remainder is derived from fish oil.

[0062] The foregoing compositions and dosage forms (and other compositions and dosage forms disclosed herein) may be used in methods of treatment. For example, the present disclosure encompasses a method of reducing the risk of mortality in a subject comprising administering a composition or dosage form disclosed herein.

[0063] The present disclosure encompasses a method of extending the life of a subject comprising administering a composition or dosage form disclosed herein.

[0064] The present disclosure encompasses a method of reducing the risk of coronary heart disease (CHD) in a subject comprising administering a composition or dosage form disclosed herein.

[0065] The present disclosure encompasses a method of reducing the risk of sudden cardiac death (SCD) in a subject comprising administering a composition or dosage form disclosed herein.

[0066] The present disclosure encompasses a method of reducing the risk of cardiac arrest in a subject comprising administering a composition or dosage form disclosed herein.

[0067] The present disclosure encompasses a method of reducing non-fasting triglyceride levels in a subject's blood comprising administering a composition or dosage form disclosed herein.

[0068] The present disclosure encompasses a method of reducing fasting triglyceride levels in a subject's blood comprising administering a composition or dosage form disclosed herein.

[0069] The present disclosure encompasses a method of increasing the omega-3 index in red blood cells of a subject comprising administering a composition or dosage form disclosed herein. Some of such embodiments further comprise identifying a subject with less than an average of about 8% EPA and/or DHA in the red blood cells of the subject.

[0070] The present disclosure encompasses a method of reducing the resting heart rate of a subject comprising administering a composition or dosage form disclosed herein.

[0071] The present disclosure encompasses a method of increasing high-density lipoprotein cholesterol (HDL-c) in a subject comprising administering a composition or dosage form disclosed herein.

[0072] The present disclosure encompasses a method of reducing non-high-density-lipoprotein cholesterol (non-HDL-c) in a subject comprising administering a composition or dosage form disclosed herein.

[0073] The present disclosure encompasses a method of reducing total cholesterol in a subject comprising administering a composition or dosage form disclosed herein.

[0074] The present disclosure encompasses a method of reducing apolipoprotein B (Apo-B) in a subject's blood comprising administering a composition or dosage form disclosed herein.

[0075] The present disclosure encompasses a method of reducing systolic blood pressure in a subject comprising administering a composition or dosage form disclosed herein.

[0076] The present disclosure encompasses a method of reducing diastolic blood pressure in a subject comprising administering a composition or dosage form disclosed herein.

[0077] The present disclosure encompasses a method of treating hypertriglyceridemia in a subject comprising administering a composition or dosage form disclosed herein. More particularly, some methods of treating hypertriglyceridemia using a composition or dosage form disclosed herein comprise identifying a subject with hypertriglyceridemia.

[0078] In some embodiments of each of the methods of treatment disclosed herein, the dose of DHA is at least about 1800 mg/day (e.g., about 1800 mg/day to about 2400 mg/day). In some such embodiments, the dose of DHA is at least about 1800 mg/day and the dose of EPA is about 700 mg/day to about 1000 mg/day. In some of such embodiments, the dose of DHA is about 900 mg to about 1200 mg, twice a day, and the dose of EPA is about 350 mg to about 500 mg, twice a day.

[0079] Some embodiments of each of the methods or treatment disclosed herein may reduce apolipoprotein B (Apo-B) in a subject's blood and increase apolipoprotein A-1 (Apo-A-1) in a subject's blood, thereby decreasing the ratio of Apo-B to Apo-A-1.

[0080] Compositions and precursors of compositions disclosed herein may be manufactured by any suitable method. For example, some compositions may be manufactured, at least in part, from crude fish oil or crude cephalopod (e.g., squid) oil.

The following methods and/or steps for processing fish and/or squid oil are illustrative, and are not meant to limit the scope of this disclosure.

[0081] In some methods, crude squid or fish oil may be deacidified by treatment with a one or more bases such as NaOH, thereby reducing oil acidity. More particularly, in some methods, aqueous NaOH is added to the crude fish or squid oil, and the crude oil is then isolated by one or more of aqueous/organic phase separation and distillation. In some methods, after addition of a base, volatile material (e.g., free fatty acid) may be stripped by means of a stripping gas (e.g., steam at a high temperature and low pressure).

[0082] In some methods, ethyl esters may be produced from the fish or squid oil by chemical reaction with the deacidified fish or squid oil. For example, the deacidified fish or squid oil may be reacted with ethanol (e.g., through a sodium ethoxide-catalyzed transesterification reaction) to form ethyl esters. Additionally or alternatively, in some embodiments, microbial lipases may be used for such a transesterification reaction.

[0083] In some methods, the resulting ethyl esters may then be concentrated by molecular distillation. In some methods, the ethyl esters may additionally or alternatively be concentrated through urea complexation. In some urea complexation procedures, ethyl esters are mixed with an ethanolic solution of urea with moderate heating. The mixture is then allowed to cool, causing the urea to crystallize. Because urea crystallizes into a hexagonal structure with channels of appropriate size to accommodate straight-chain saturated fatty acids, fatty acids with a low degree of unsaturation remain complexed with the urea, while fatty acids with a higher degree of saturation (e.g., DHA and EPA) are separated in the solution and may be isolated by filtration.

[0084] A method of processing crude fish or squid oil may comprise re-esterification of ethyl ester fatty acids to produce a re-esterified triglyceride. For example, ethyl esters may be enzymatically converted to triglycerides.

[0085] In some methods, fish or squid oil (either in ethyl ester form or as a re-esterified triglyceride) may, at some point, undergo a bleaching process. For example, the oil may be heated (e.g., to approximately 80–85 °C) and mixed with activated bleaching clay and/or activated carbon. The bleaching clay, activated carbon, or other bleaching agent may adsorb soaps, sulfur-containing compounds, trace metals, pigments, and/or other components.

[0086] In some embodiments, fish or squid oil may be deodorized to remove, *inter alia*, free fatty acids, aldehydes, and ketones or other compounds or materials. In the absence of such deodorization, the fish oil may have objectionable flavor and/or smell characteristics. In some embodiments, volatile material is stripped by means of a stripping gas (e.g., steam at high temperature and low pressure).

[0087] Some compositions disclosed herein comprise a mixture of oils from distinct sources. For example, some compositions or dosage forms may comprise cephalopod oil and/or fish oil. In some embodiments, cephalopod oil may be combined with fish oil. In some circumstances, the oils may be combined in large containers and that will be used to fill smaller containers (e.g., nitrogen-flushed bottles for individual sale to consumers). Further, in some embodiments, antioxidants, flavorings, tocopherol, or other additives may be added prior to sealing the containers for distribution to end users.

[0088] It should be understood that for some embodiments only a portion of the processing steps disclosed above may be performed. Additionally, some compositions or dosage forms may not be processed by the methods described above, but by alternative processes known in the art. Also, some compositions or dosage forms may be processed by blending already-purified oils.

[0089] Some embodiments of the compositions and dosage forms disclosed herein comprise re-esterified triglycerides (rTGs), including re-esterified EPA and DHA. rTGs may have advantages in therapeutic settings relative to other fatty acids, such as the ethyl ester forms of EPA and DHA. For example, as shown in Example 1 below, the triglyceride levels of a group receiving a composition with re-esterified triglycerides were reduced to a greater extent than the triglyceride levels of a group that received ethyl ester forms of EPA and DHA. Further, the group receiving a composition with re-esterified EPA and DHA had a lower heart rate, a lower hazard ratio, lower systolic and diastolic blood pressure, a higher omega-3 index, increased HDL-c levels, lower non-HDL-c levels, and lower cholesterol levels relative to the group that received ethyl ester forms of EPA and DHA.

[0090] Compositions and dosage forms comprising a relatively high ratio of DHA to EPA may have one or more advantages relative to compositions that have a lower ratio of DHA to EPA. For example, as shown in Example 1 below, patients who received about 5:2 DHA:EPA as compared to patients who received about 4:5

DHA:EPA (at comparable total omega-3 PUFA amounts) had reduced triglyceride levels, lower heart rate, decreased blood pressure, and an increased omega-3 index.

[0091] Example 1

[0092] One hundred and nineteen individuals who had been diagnosed with moderate fasting hypertriglyceridemia (i.e., 150 mg triglyceride per dL of blood plasma (1.7 mM) to 500 mg triglyceride per dL of blood plasma (5.65 mM)) were recruited to investigate the response of such individuals to three different lipophilic combinations over an eight-week period.

[0093] The 119 individuals (median age of 64) were randomly assigned to three separate groups in a 1:1:1 ratio. The first group, which is referred to herein as the re-esterified triglyceride (rTG) group, was prescribed 5.5 g of LIPOMAR™ per day (two 1.375 g capsules taken twice per day). The 5.5 g of prescribed LIPOMAR™ included 767 mg EPA and 1930 mg DHA (i.e., 2696 mg of combined EPA and DHA). The oil of the LIPOMAR™ capsules was a blend of refined and re-esterified squid and fish oils. The second group, which is referred to herein as the ethyl ester (EE) group, was prescribed 4.0 g of LOVAZA™ (two 1.0 mg capsules taken twice per day). The 4.0 g of prescribed LOVAZA™ included 1702 mg EPA and 1382 mg DHA in ethyl ester form (i.e., 3085 mg of combined EPA and DHA). The third group, referred to herein as the placebo group, received 4.0 g of olive oil (OO) per day (two 1.0 mg capsules taken twice per day).

[0094] At the beginning of the study, each individual's resting heart rate and blood pressure were measured. Additionally, one or more blood samples of each patient was taken and used to assess the concentration and/or ratio of blood components. For example, the patient's blood was tested to determine non-fasting triglyceride levels, cholesterol levels, the omega-3 index of red blood cell membranes, and the concentrations and/or ratios of various proteins and lipoproteins found in the blood. Other tests on the blood plasma were also conducted. Table 1 provides a summary of the characteristics of those in each experimental arm of the study.

	Re-esterified Triglyceride	Ethyl ester	Olive oil (placebo)
General characteristics			
Number of participants/group	N = 39	N = 40	N = 40
Average age	63.3	60.4	63.6

Number of male participants	36 (92.3) ¹	26 (65.0)	32 (80.0)
BMI (mean)	29	28	29
Average weight of female (kg)	82.1	71.1	81.8
Range of female weight (kg)	68–98	54–86	55–96
Average weight of male (kg)	93.1	89.8	90.4
Range of male weight (kg)	75–120	71–117	72–118
History (self-declared)			
Cardiovascular disease	30 (76.9)	30 (75.0)	32 (80.0)
Dyslipidæmi	30 (76.9)	35 (87.5)	30 (75.0)
Hypertension	22 (56.4)	19 (47.5)	20 (50.0)
Diabetes Mellitus Type II	5 (12.8)	6 (15.0)	8 (20.0)
Psychiatric disorder	1 (2.6)	2 (5.0)	1 (2.5)
Therapy			
Statin therapy users	30 (76.9)	29 (72.5)	29 (72.5)
Ezetemibe users	3 (7.7)	5 (12.5)	2 (5.0)
Hypertension therapy users	27 (69.2)	26 (65)	30 (75.0)
Psychopharmaca	3 (7.7)	5 (12.5)	2 (5.0)
Blood pressure (average)			
Systolic (mmHg)	145	141	145
Diastolic (mmHg)	84	83	85
Heart rate (average # of beats/min)	67	63	65
Lipid values (average)			
Cholesterol (mg/dL)	247	240	218
HDL (mg/dL)	40.2	42.9	44.4
Non-HDL (mg/dL)	141.3	146.7	145.9
Triglyceride (mg/dL)	247	240	218
Omega-3 index (%)	6.7	6.4	6.2

Table 1

[0095] After four and eight weeks of treatment, the individuals in each group returned, and the measurements and tests performed at the beginning of the study

¹ Numbers enclosed within parentheses denote the percentage of participants in each group that fall within the relevant category.

were repeated. A summary of these results is set forth in the text below, and the tables and figures referenced therein.

[0096] FIGS. 1–3 provide graphical depictions of data related to changes in triglyceride levels in each arm of the study. More particularly, FIG. 1 provides a box-and-whisker plot depicting, within each group, changes in blood serum triglyceride levels between baseline levels measured at the beginning of the study and blood serum triglyceride levels measured at the conclusion of the study. The box-and-whisker plot shows the median, upper and lower quartile, and maximum and minimum values (excluding outliers). Outlier data are depicted with circles or asterisks.

[0097] FIG. 2 provides a bar graph of median non-fasting triglyceride levels (mg/dL) as measured at the beginning of the study (i.e., baseline levels) and at the conclusion of the study. FIG. 3 depicts changes in median non-fasting triglyceride levels (mg/dL) as measured at the beginning of the study relative to median non-fasting triglyceride levels as measured at the conclusion of the study. As depicted in these figures, the median non-fasting triglyceride levels for the rTG arm and the EE arm of the study were lowered from baseline levels (from 2.79 ± 1.12 mM to 1.81 ± 0.82 mM for the rTG arm and from 2.70 ± 1.39 mM to 2.10 ± 1.23 mM for the EE arm), while the median non-fasting triglyceride level for the placebo arm did not decrease (changing from 2.46 ± 1.38 mM to 2.69 ± 1.62 mM).

[0098] The decrease in triglyceride levels for the rTG and EE arms relative to baseline levels was statistically significant (p-values < 0.001). The decrease in triglyceride levels in both the rTG and EE arms was also statistically significant when compared with the placebo group receiving olive oil (p-values < 0.001). Changes in triglyceride levels within the placebo group were not statistically significant (p-value = 0.52).

[0099] The total amount (by weight) of combined EPA and DHA differs in the two experimental arms of the study. More particularly, the ethyl ester treatment arm (i.e., LOVAZA) receives 14% more combined EPA and DHA by weight than the group receiving re-esterified triglyceride. FIGS. 4 and 5 are analogous to FIGS. 2 and 3, but correct for this difference in total amount of supplied EPA and DHA. As shown in these figures, the decrease in median triglyceride levels for the rTG arm of the study is even more pronounced when the data are corrected to account for the difference in the total amount of combined EPA and DHA by weight.

[00100] FIG. 6 is a bar graph providing hazard ratios for all-cause mortality in each experimental arm of the study. The hazard ratios are determined and defined as set forth in Thomsen et al., *Low Nonfasting Triglycerides and Reduced All-Cause Mortality: A Mendelian Randomization Study*, 60 Clin. Chem 737–46 (2014). The hazard ratio remained constant throughout the study in both the EE group and the placebo group. However, the hazard ratio for those in the rTG group was lowered at the conclusion of the study relative to the hazard ratio at the beginning of the study.

[00101] FIGS. 7, 8, and 9 provide graphical depictions of data related to the changes in the omega-3 index of members of the study population. The omega-3 index is defined as the ratio of the combined number of EPA and DHA fatty acids in the measured erythrocyte membranes relative to the total number of all fatty acids (including EPA and DHA) in the measured erythrocyte membranes. The ratio is expressed either as a ratio or a percentage. FIG. 7 provides a box-and-whisker plot of the changes in omega-3 index between baseline levels measured at the beginning of the study and levels measured at the conclusion of the study. The box-and-whisker plot shows the median, upper and lower quartile, and maximum and minimum values (excluding outliers). There were no outliers in the rTG and ethyl ester treatment arms. The four outliers of the control group are depicted with circles and an asterisk in the corresponding box plot. As shown in FIG. 7, the omega-3 index of those in the rTG and EE treatment arms generally increased, while the omega-3 index of those in the placebo (olive oil) group, generally did not increase.

[00102] FIG. 8 depicts the mean omega-3 index for each arm of the study both before treatment and at the conclusion of the treatment period. FIG. 9 depicts changes in the mean omega-3 index across the course of the study. As shown in these figures, the mean omega-3 index in the rTG and EE treatment arms increased relative to both the placebo group and to baseline levels within each group. The increase in omega-3 index values in both the rTG and EE treatment arms was statistically significant relative to both the placebo group and baseline values (p -values < 0.001). The increase in omega-3 index values for the rTG group relative to the EE group, as depicted in these figures, was not statistically significant. Additionally, changes in the omega-3 index in the placebo group relative to baseline values were not statistically significant.

[00103] FIGS. 10 and 11 are analogous to FIGS. 8 and 9, but corrected for the total amount of EPA and DHA. As shown in these figures, the increase in mean omega-3 index values relative to baseline values is even more pronounced when the data are

corrected to account for the difference in the total amount of combined EPA and DHA (by weight) in treatment arms.

[00104] FIG. 12 provides a scatter plot depicting the difference between triglyceride levels (mg/dL) measured at the end of the study from triglyceride levels measured at the beginning of the study (i.e., baseline levels) on the y-axis, and the difference in omega-3 index (i.e., omega-3 index levels at the conclusion of the study minus baseline omega-3 index levels measured at the beginning of the study) on the x-axis. As is evidenced by the negative slope of the best-fit line, this plot shows a negative correlation between these two variables (R^2 value = 0.233)

[00105] FIG. 13 provides a bar graph depicting mean EPA levels (mg/dL) in red blood cells at the beginning of the study and at the conclusion of the study. As depicted in FIG. 13, mean EPA concentrations increased in both the rTG and EE treatment arms across the course of the study, while the mean EPA concentration of the placebo group did not increase. The increase in EPA concentrations for both treatment groups was statistically significant relative to baseline levels (p-value < 0.001). The increase in EPA concentration in the EE arm was significantly greater than the increase in EPA concentrations in the rTG group (p-value < 0.043). Changes in the placebo group were not statistically significant.

[00106] FIG. 14 provides a bar graph depicting mean DHA levels (mg/dL) in red blood cells before and after treatment. As depicted in FIG. 14, the mean DHA concentrations increased in both the rTG and EE treatment arms across the course of the study, while the mean DHA concentration in the control group did not increase. The increase in DHA concentrations for both treatment groups was statistically significant relative to baseline levels (p-values < 0.001). The increase in DHA concentration in the rTG arm was significantly greater than the increase DHA concentration in the EE arm (p-value < 0.001). Changes in the olive oil placebo group were not statistically significant.

[00107] FIGS. 15 and 16 provide graphical representations of heart rate data collected during the study. More particularly, FIG. 15 provides a box-and-whiskers plot of the changes in heart rate (beats/minute) between a baseline heart rate measured at the beginning of the study and a heart rate measured at the conclusion of the study. The box-and-whisker plot shows the median, upper and lower quartile, and maximum and minimum values (excluding outliers). Outliers are shown as circles or asterisks.

As shown in FIG. 15, heart rate levels generally decreased in the rTG group, while heart rates of those in the EE and placebo (olive oil) groups did not decrease.

[00108] FIG. 16 is a bar graph depicting mean changes in heart rate (expressed in beats/minute) for each group of the study. The mean heart rate for those in the rTG arm decreased more than 2.5 beats per minute, while no decrease was observed in the EE and placebo groups. The decrease in heart rate in the rTG group was significant relative to both the placebo group (p -value = 0.045) and the baseline values prior to treatment (p -value = 0.038). Changes in heart rate in the EE and placebo groups relative to baseline values were not statistically significant.

[00109] FIGS. 17, 18, and 19 provide graphical representations of high-density lipoprotein-cholesterol (HDL-c) concentration levels as measured during the study. More particularly, FIG. 17 provides a box-and-whisker plot that depicts, within each group, changes in blood plasma HDL-c levels (mM) between baseline levels measured at the beginning of the study and levels measured at the conclusion of the study. The box-and-whisker plot shows the median, upper and lower quartile, and maximum and minimum values (excluding outliers). Outliers are depicted with circles or asterisks.

[00110] As shown in FIG. 17, HDL-c concentrations in the blood plasma of those in the rTG and ethyl ester group generally increased, while the HDL-c concentrations of those in the placebo group generally did not increase.

[00111] FIG. 18 provides a bar graph of mean HDL-c levels for each arm of the study. FIG. 19 is a bar graph depicting the changes in mean HDL-c concentrations within each group across the course of the study. These bar graphs show an increase in mean HDL-c concentrations for those in the rTG and EE treatment arms, while the mean HDL-c concentration for those in the placebo group decreased. The increase in HDL-c concentration within the rTG group was statistically significant relative to both the placebo group (p -value < 0.001) and baseline values prior to treatment (p -value < 0.001). The increase in HDL-c concentration within the EE group was also statistically significant relative to both the placebo group (p = 0.025) and baseline values prior to treatment (p -value = 0.026).

[00112] FIG. 20 is a bar graph of mean non-HDL-c concentrations (mg/dL) for each arm of the study. This bar graph shows a decrease in mean non-HDL-c concentrations for those in the rTG and EE treatment arms relative to baseline (i.e., pre-treatment) values. The decrease in mean non-HDL-c levels relative to baseline values is statistically significant for both the rTG and EE groups (p -values \leq 0.001). Additionally,

the decrease in mean HDL-c concentrations for the rTG group was significant relative to the placebo group ($p = 0.027$), while the decrease in mean HDL-c concentrations for the EE group was not statistically significant relative to the placebo group (p -value = 0.064). Changes in non-HDL-c concentration within the placebo group were not statistically significant.

[00113] FIG. 21 is a bar graph depicting mean total cholesterol levels (mg/dL) for each arm of the study both at the beginning of the study (i.e., baseline values) and at the conclusion of the study. This bar graphs shows a decrease a mean total cholesterol levels for those in the rTG and EE treatment arms relative to baseline (i.e., pre-treatment) values. The decrease in total cholesterol levels in these groups relative to baseline levels was statistically significant (p -values < 0.05). However, neither the decrease in total cholesterol levels between the rTG and EE groups versus the placebo control (olive oil) nor the decrease in cholesterol levels within the placebo group relative to baseline levels was statistically significant. Likewise, the difference in the change in total cholesterol levels within the rTG group relative to the EE group was not statistically significant.

[00114] FIGS. 22 and 23 provide graphical depictions of data related to blood plasma apolipoprotein B (Apo-B) concentration levels (mg/dL) in each arm of the study. More particularly, FIG. 22 is a bar graph depicting mean Apo-B concentrations for each arm of at the beginning of the study (i.e., baseline levels) and at the conclusion of the study. FIG. 23 is a bar graph depicting the change in mean Apo-B concentrations in each arm across the study. The decrease in mean Apo-B concentrations shown in these figures was significant in both the rTG and EE groups relative to baseline (i.e., pre-treatment) levels (p -value < 0.01). The change in mean Apo-B concentrations found in the rTG and EE groups relative to the olive oil placebo group was not statistically significant. The decrease in Apo-B concentrations found in the olive oil placebo group relative to baseline (i.e., pre-treatment) values was also not statistically significant.

[00115] FIGS. 24 and 25 provide graphical depictions of data relating to the relative concentrations of Apo-B and apolipoprotein A (Apo-A) in the blood plasma of those in each arm of the study. More particularly, FIG. 24 is a bar graph depicting the mean ratio of Apo-B to Apo-A in each group, both at the beginning and at the conclusion of the study. FIG. 25 is a bar graph that depicts changes in mean Apo-B/Apo-A concentration across the course of the study. These figures show a decrease in the

mean Apo-B/Apo-A ratio for both the rTG and EE groups. The decrease in the mean Apo-B/Apo-A ratio relative to baseline ratios in both the rTG and EE groups was statistically significant, with p-values of 0.003 and 0.008 respectively. The decrease in mean Apo-B/Apo-A ratios for the rTG groups was also statistically significant relative to the placebo group (p-value = 0.027). The decrease in mean Apo-B/Apo-A ratios for the EE group was not statistically significant relative to the placebo group (p-value = 0.182). Changes to the Apo-B/Apo-A ratio within the placebo group relative to baseline values were not statistically significant.

[00116] FIG. 26 is a bar graph depicting the mean systolic blood pressure in each group both at the beginning and at the conclusion of the study. FIG. 27 is a similar bar graph depicting the mean diastolic blood pressure in each group both at the beginning and at the conclusion of the study. The decrease in systolic and diastolic blood pressure within the rTG and EE groups relative to baseline levels was statistically significant (p-values < 0.05). The changes in systolic and diastolic blood pressure in the rTG and EE groups relative to the placebo group (olive oil) were not statistically significant.

[00117] FIG. 28 is a bar graph depicting the mean concentrations of C-reactive protein (mg/L) in each arm of the study both at the beginning and at the conclusion of the study. The mean values of C-reactive protein were lowered in both the rTG group and the olive oil placebo group, but increased in the EE group. None of these changes in the rTG and EE groups was significant relative to baseline values or the placebo group.

[00118] FIG. 29 is a bar graph depicting the percent change of the following measurements or values, relative to baseline (i.e., pre-treatment) levels: triglyceride concentration, HDL-c concentration, non-HDL-c concentration, the omega-3 index, heart rate, and the ratio of Apo-B to Apo-A.

[00119] Table 2 provides statistical data for between-group analyses (i.e., ANOVA-derived p-values) and within-group analyses (i.e., p-values derived from a paired t-test). In addition to the measurements, values, and indexes noted above, the table provides statistical data for changes in levels of alanine transaminase (ALAT), alkaline phosphatase, bilirubin, and HgbA1c. The table also provides statistical data for changes in the clotting tendency of the blood (as measured using the INR test) and the body mass index (BMI) of those in the study.

	Group difference, ANOVA, p-values two-tailed			Individual change, Paired t-test (post-pre)		
	rTG vs. EE	rTG vs. OO	EE vs. OO	rTG	EE	OO
Triglyceride	0.785	<0.001	<0.001	<0.001	<0.001	0.138
Total cholesterol	0.996	0.213	0.180	0.010	<0.001	0.323
HDL cholesterol	0.088	<0.001	0.025	<0.001	0.026	0.131
Non-HDL cholesterol	0.930	0.027	0.064	0.001	<0.001	0.521
Apo-B/A ratio	0.676	0.027	0.182	0.003	0.008	0.917
Apo-A	0.478	0.478	1.000	0.682	0.131	0.277
Apo-B	0.992	0.337	0.274	0.010	0.002	0.395
Heart rate	0.233	0.045	0.717	0.038	0.868	0.070
Diastolic BP	0.947	0.996	0.972	0.006	0.010	0.043
Systolic BP	0.805	0.807	0.420	0.050	0.006	0.258
ALAT	0.379	0.070	0.638	0.009	0.600	0.492
Alkaline phosphatase	0.837	0.110	0.311	0.003	0.026	0.863
Bilirubin	0.964	0.969	0.872	0.919	0.585	0.818
CRP	0.410	0.469	0.994	0.243	0.651	0.739
INR	0.068	0.031	0.999	0.021	0.604	0.239
HgbA1c	0.993	0.563	0.631	0.041	0.227	0.842
BMI	0.827	0.534	0.878	0.303	0.686	0.738
DHA	<0.001	<0.001	<0.001	<0.001	<0.001	0.316
EPA	0.043	<0.001	<0.001	<0.001	<0.001	0.690
O3 index	0.788	<0.001	<0.001	<0.001	<0.001	0.173

Table 2

[00120] FIG. 30 and Table 3 provide information regarding the fatty acid composition of the oil prescribed to those in the rTG arm of the study. The oil was analyzed by gas-liquid chromatography to generate a fatty acid composition profile and to determine the relative concentrations of fatty acids in the oil. The resulting chromatogram is shown in FIG. 30, and data corresponding to the chromatogram are shown in Table 3. The percentages reported in Table 3 are relative to the total amount of detected fatty

acids, and do not necessarily correspond with the total mass of the oil composition. As shown in this figure, the oil comprises a plurality of fatty acid compositions of varying length. The two most prevalent fatty acids in the composition are DHA (peak 29) and EPA (peak 22).

Peak #	Time	Component Name	Area [%]	Corrected Area [%]	Omega-3 Area [%]	Saturated FAC Area [%]	Mono-unsaturated FAC Area [%]	Poly-unsaturated FAC Area [%]
1	6.035	C14:0	0.13	0.1263	0.0000	0.1263	0.0000	0.0000
2	7.336		0.07					
3	7.523	C16:0	0.50	0.4839	0.0000	0.4839	0.0000	0.0000
4	7.781	C16:1 n-7	0.31	0.3082	0.0000	0.0000	0.3082	0.0000
5	8.401		0.06					
6	8.534		0.08					
7	9.918	C18:0	1.19	1.1554	0.0000	1.1554	0.0000	0.0000
8	10.213	C18:1 n-9	3.26	3.1903	0.0000	0.0000	3.1903	0.0000
9	10.323	C18:1 n-7	1.06	1.0641	0.0000	0.0000	1.0641	0.0000
10	10.530		0.11					
11	10.949	C18:2 n-6	0.40	0.3998	0.0000	0.0000	0.0000	0.3998
12	12.102	C18:3 n-3	0.20	0.1960	0.1960	0.0000	0.0000	0.1960
13	12.707	C18:4 n-3	0.39	0.3899	0.3899	0.0000	0.0000	0.3899
14	13.554	C20:0	0.35	0.3468	0.0000	0.3468	0.0000	0.0000
15	13.867	C20:1 n-11	6.93	6.9333	0.0000	0.0000	6.9333	0.0000
16	13.957	C20:1 n-9	5.61	5.6066	0.0000	0.0000	5.6066	0.0000
17	14.150		0.65					
18	15.014		0.74					
19	16.179	C20:4 n-6	1.63	1.6317	0.0000	0.0000	0.0000	1.6317
20	16.590		0.34					
21	17.271	C20:4 n-3	1.08	1.0842	1.0842	0.0000	0.0000	1.0842
22	17.871	C20:5 n-3	16.58	16.5818	16.5818	0.0000	0.0000	16.5818
23	18.794	C22:1 n-11 + 13	10.97	11.0829	0.0000	0.0000	11.0829	0.0000

24	18.947	C22:1 n-9	1.50	1.4992	0.0000	0.0000	1.4992	0.0000
25	19.202		0.42					
26	20.810	C21:5 n-3	0.90	0.8992	0.8992	0.0000	0.0000	0.8992
27	22.545	C22:5 n-6	0.85	0.8484	0.0000	0.0000	0.0000	0.8484
28	23.666	C22:5 n-3	2.77	2.9399	2.9399	0.0000	0.0000	2.9399
29	24.587	C22:6 n-3	39.63	40.0274	40.0274	0.0000	0.0000	40.0274
30	24.821	C24:1	1.30	1.3788	0.0000	0.0000	1.3788	0.0000

Table 3

[00121] Any methods disclosed herein may include one or more steps or actions for performing the described method. The method steps and/or actions may be interchanged with one another. In other words, unless a specific order of steps or actions is required for proper operation of the embodiment, the order and/or use of specific steps and/or actions may be modified. Moreover, sub routines or only a portion of a method described herein may be a separate method within the scope of this disclosure. Stated otherwise, some methods may include only a portion of the steps described in a more detailed method.


[00122] It will be apparent to those having skill in the art that changes may be made to the details of the above-described embodiments without departing from the underlying principles of the present disclosure.

We Claim:

1. A dosage form comprising:
a marine oil comprising fatty acids, including omega-3 polyunsaturated fatty acids (PUFAs), including re-esterified 4Z,7Z,10Z,13Z,16Z,19Z-docosahexaenoic acid (DHA) and re-esterified 5Z,8Z,11Z,14Z,17Z-eicosapentaenoic acid (EPA), and
wherein an amount of DHA is 450 mg to 700 mg and an amount of EPA is 150 mg to 300 mg, wherein at least 55 wt.% of the fatty acids are omega-3 PUFAs, and wherein at least 50 wt.% of the fatty acids are re-esterified triglycerides (rTGs).
2. The dosage form of claim 1, wherein a ratio of DHA to EPA is about 3:2 to about 4:1.
3. The dosage form of claim 1 or 2, wherein a ratio of DHA to EPA is about 5:2.
4. The dosage form of any one of claims 1-3, wherein the dosage form is a capsule.
5. The dosage form of any one of claims 1-3, wherein the dosage form is a liquid-filled capsule.
6. The dosage form of any one of claims 1-3, wherein the dosage form is a liquid.
7. The dosage form of any one of claims 1-6, wherein a mass of the dosage form is about 1200 mg to about 1600 mg.
8. The dosage form of any one of claims 1-7, wherein at least 55 wt.% of the fatty acids are TGs.
9. The dosage form of any one of claims 1-8, wherein about 55 wt.% to about 70 wt.% of the fatty acids are TGs.
10. The dosage form of any one of claims 1-9, wherein about 25 wt.% to about 45 wt.% of the PUFAs are diglycerides (DGs).
11. The dosage form of any one of claims 1-11, wherein 0 wt.% to about 5 wt.% of the fatty acids are monoglycerides (MGs).
12. The dosage form of any one of claims 1-12, wherein about 5 wt% or less of the fatty acids are ethyl esters.

13. The dosage form of any one of claims 1-12, wherein less than 1 wt.% of the fatty acids are ethyl esters.
14. The dosage form of any one of claims 1-13, wherein at least 60 wt.% of the fatty acids are PUFAs.
15. The dosage form of any one of claims 1-14, wherein at least 50 wt.% of the fatty acids are DHA and EPA.
16. The dosage form of any one of claims 1-15, wherein 0 wt.% to about 35 wt.% of the fatty acids are mono-unsaturated fatty acids.
17. The dosage form of any one of claims 1-16, wherein 0 wt.% to about 5 wt.% of the fatty acids are saturated fatty acids.
18. The dosage form of any one of claims 1-17, wherein at least 80 wt.% of the PUFAs are DHA and EPA.
19. The dosage form of any one of claims 1-18, wherein at least 15 wt.% of the marine oil is derived from cephalopod oil.
20. The dosage form of any one of claims 1-19, wherein about 10 wt.% to about 25 wt.% of the marine oil is derived from cephalopod oil and the remainder is derived from fish oil.
21. The dosage form of any one of claims 1-19, wherein about 25 wt.% to about 50 wt.% of the marine oil is derived from cephalopod oil and the remainder is derived from fish oil.
22. The dosage form of any one of claims 1-19, wherein at least 50 wt.% of the marine oil is derived from cephalopod oil.
23. A dosage form for use in reducing the resting heart rate of a subject, the dosage form comprising the dosage form of any one of claims 1-22.

1/30

**FIG. 1**

2/30




FIG. 2

3/30





FIG. 3

4/30

**FIG. 4**

5/30




FIG. 5

6/30





FIG. 6

7/30

**FIG. 7**

8/30




FIG. 8

9/30




FIG. 9

10/30




FIG. 10

11/30





FIG. 11

12/30

**FIG. 12**

13/30





FIG. 13

14/30

**FIG. 14**

15/30




FIG. 15

16/30





FIG. 16

17/30

**FIG. 17**

18/30






FIG. 18

19/30**FIG. 19**

20/30**FIG. 20**

21/30




FIG. 21

22/30




FIG. 22

23/30




FIG. 23

24/30




FIG. 24

25/30




FIG. 25

26/30




FIG. 26

27/30




FIG. 27

28/30



FIG. 28

29/30




FIG. 29

WO 2015/188046

PCT/US2015/034372

30/30

FIG. 30

