



(11) (21) (C) **2,157,879**
(86) 1994/01/13
(87) 1994/09/15
(45) 2000/08/29

(72) Labroo, Virender M., US

(72) Piggott, James R., US

(72) Bain, Steven D., US

(73) ZYMOGENETICS, INC., US

(51) Int.Cl.⁶ A61K 31/35, A61K 31/40

(30) 1993/03/11 (08/029,729) US

(54) **METHODES UTILISANT LE 3,4-DIARYLCHROMANE POUR
EMPECHER LA PERTE DE SUBSTANCE OSSEUSE**

(54) **METHODS FOR INHIBITING BONE LOSS WITH 3,4-
DIARYLCHROMAN**

(57) Procédés et préparations pharmaceutiques permettant de réduire les pertes osseuses. Les 3,4-diarylchromanes et leur sels pharmacocompatibles sont formulés pour servir de médicaments traitant les pertes osseuses résultant de l'ostéoporose et d'autres états. Un exemple de 3,4-diarylchromane est représenté par le centchromane (3,4-trans-2,2-diméthyl-3-phényl-4-[p-(bêta-pyrrolidinoéthoxy)phényl]-7-méthoxy-chromane). Les formulations décrites se présentent sous forme de comprimés ou autres présentations à administration orale, et d'implants sous-cutanés à diffusion contrôlée.

(57) Methods and pharmaceutical compositions for reducing bone loss are disclosed. 3,4-diarylchromans and their pharmaceutically acceptable salts are formulated into medicaments for the treatment of bone loss due to osteoporosis or other conditions. An exemplary 3,4-diarylchroman is centchroman (3,4-trans-2,2-dimethyl-3phenyl-4-(p-(beta-pyrrolidinoethoxy) phenyl)-7-methoxy-chroman). Formulations include tablets and other forms suitable for administration and controlled-release subdermal implants.






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/40, 31/35		A1	(11) International Publication Number: WO 94/20098
			(43) International Publication Date: 15 September 1994 (15.09.94)
(21) International Application Number: PCT/US94/00633		(81) Designated States: AU, BR, BY, CA, CZ, FI, HU, JP, KR, KZ, NO, NZ, PL, RO, RU, SK, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 13 January 1994 (13.01.94)			
(30) Priority Data: 08/029,729 11 March 1993 (11.03.93) US		2157879 Published With international search report.	
(71) Applicant: ZYMOGENETICS, INC. [US/US]; 4225 Roosevelt Way N.E., Seattle, WA 98105 (US).			
(72) Inventors: LABROO, Virender, M.; 2814 163rd Place S.E., Mill Creek, WA 98012 (US). PIGGOTT, James, R.; 23612 Meridian Avenue West, Bothell, WA 98021 (US). BAIN, Steven, D.; 123 NW 180th, Seattle, WA 98177 (US).			
(74) Agent: PARKER, Gary, E.; Zymogenetics, Inc., 4225 Roosevelt Way N.E., Seattle, WA 98105 (US).			

(54) Title: METHODS FOR INHIBITING BONE LOSS WITH 3,4-DIARYLCHROMAN



a=significantly decreased vs ovx/E2 and ovx/C groups; $P < .05$.
b=significantly decreased vs sham and ovx/E2 groups; $P < .01$.

(57) Abstract

Methods and pharmaceutical compositions for reducing bone loss are disclosed. 3,4-diarylchromans and their pharmaceutically acceptable salts are formulated into medicaments for the treatment of bone loss due to osteoporosis or other conditions. An exemplary 3,4-diarylchroman is centchroman (3,4-*trans*-2,2-dimethyl-3-phenyl-4-[*p*-(beta-pyrrolidinoethoxy) phenyl]-7-methoxy-chroman). Formulations include tablets and other forms suitable for oral administration and controlled-release subdermal implants.

METHODS FOR INHIBITING BONE LOSS WITH
3,4-DIARYLCHROMAN

Background of the Invention

5 Bone remodeling is the dynamic process whereby
skeletal mass and architecture are renewed and maintained.
This renewal and maintenance is a balance between bone
resorption and bone formation, with the osteoclast and the
osteoblast considered the two key participants in the
10 remodeling process. The osteoclast initiates the
remodeling cycle by resorbing a cavity in the bone which
is subsequently refilled when the osteoblast synthesizes
and deposits new bone matrix into the excavation. The
activities of osteoclast and osteoblast are regulated by
15 complex interactions between systemic hormones and the
local production of growth factors and cytokines at active
remodeling sites.

 Imbalances in bone remodeling are associated
with such conditions as osteoporosis, Paget's disease, and
20 hyperparathyroidism. Osteoporosis, characterized by a
decrease in the skeletal mass, is one of the most common
diseases of postmenopausal women and is often the cause of
debilitating and painful fractures of the spine, hip and
wrist.

25 Approximately 25% of all postmenopausal women
suffer from osteoporosis, and it is generally accepted
that the etiology of the disease involves the reduction of
circulating estrogens (Komm et al., Science 241:81-84,
1988). Komm et al. further report that the proportion of
30 caucasian women in the United States who are at risk for a
hip fracture is 15%, or 247,000 hip fractures per year in
women over the age of 45.

 The costs of osteoporosis, both personal and
financial, are enormous. In 1984, 145,000 in-patient
35 fracture reductions and 107,000 hip arthroplasties and
replacements were performed on American women over 65

2157879

- 2 -

years of age. Among patients who lived alone prior to hip fracture, 15% to 20% required long-term care as a result of the fracture and one year after the fracture had still not regained their independence. The total financial cost of osteoporosis treatment, including fractures, in the United States in 1986 was 7-10 billion dollars (Peck et al., Am. J. Med. 84:275-282, 1988).

Bone loss associated with osteoporosis has been arrested by the administration of exogenous estrogens. To be effective, estrogen therapy must begin within a few years of the onset of menopause, and should continue for 10 to 15 years, according to Thorneycroft (Am. J. Obstet. Gynecol. 160:1306-1310, 1989). While there are several different types of estrogens, 17- β -estradiol is the primary estrogen found naturally occurring in premenopausal women and is often the compound of choice for therapeutic use. At the recommended dose, however, there are significant side effects, the most disturbing being the well-established correlation of estrogen therapy with endometrial and breast cancers. The incidence of carcinoma is both dose-dependent and duration-dependent.

Avoidance of the cancer risk has been achieved by the concomitant use of a progestogen with estrogen. This combination, however, causes menses to return, which many women find unacceptable. A further disadvantage is that the long-term effects of the progestogen have not been fully determined. Thus, a large population of women require alternatives to hormone replacement therapies that can safely prevent the rapid bone loss that accompanies the menopause.

Centchroman is a non-steroidal compound known to have antiestrogenic activity. It is in use in India as an oral contraceptive (see, for example, Salman et al., U.S. Patent No. 4,447,622; Singh et al., Acta Endocrinol (Copenh) 126:444-450, 1992; Grubb, Curr. Opin. Obstet. Gynecol. 3:491-495, 1991; Sankaran et al., Contraception

- 3 -

9:279-289, 1974; Indian Patent No. 129187). Centchroman has also been investigated as an anti-cancer agent for treatment of advanced breast cancer (Misra et al., Int. J. Cancer 43:781-783, 1989), but has not previously been
5 shown to have an effect on bone resorption.

There remains a need in the art for compositions and methods useful in reducing bone loss, in particular bone loss associated with osteoporosis. There is a further need for such compositions that lack the
10 undersirable side effects of estrogens. The present invention provides such compositions and methods and also provides other, related advantages.

Brief Description of the Drawings

15 Figure 1 illustrates the effects of centchroman on bone loss in ovariectomized mice.

Figure 2 illustrates the effects of centchroman on cancellous bone volumes in the proximal tibiae of ovariectomized rats.

20 Figure 3 illustrates the effects of centchroman on bone resorption (left) and uterine weights (right) in ovariectomized rats.

Detailed Description of the Invention


25 The present invention is based in part on the discovery that a representative 3,4-diarylchroman, centchroman (3,4-trans-2,2-dimethyl-3-phenyl-4-[p-(beta-pyrrolidinoethoxy)phenyl]-7-methoxy-chroman), is an effective inhibitor of bone loss in ovariectomized mice
30 and rats. These animal models mimic the post-menopausal condition and are generally recognized models of osteoporosis. These data thus indicate that the 3,4-diarylchromans are useful as therapeutic agents for reducing bone loss in mammals, including primates such as
35 humans.

2157879

- 4 -

Within the present invention, compounds of formula (I) or their pharmaceutically acceptable salts are used for reducing bone loss in a patient.

5



(I)


10

Within formula (I), R1, R4 and R5 are individually hydrogen, halo, trifluoromethyl, lower alkyl, lower alkoxy or tertiary amino lower alkoxy; and R2 and R3 are individually H or a lower alkyl. As used herein, the term "lower alkyl" includes straight and branched chain alkyl radicals containing from 1 to 6 carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, n-amyl, sec-amyl, n-hexyl, 2-ethylbutyl, 2,3-dimethylbutyl and the like. The term "lower alkoxy" includes straight and branched chain alkoxy radicals containing from 1 to 6 carbon atoms, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, tert-butoxy, n-amyl, sec-amyl, n-hexyloxy, 2-ethyl-butoxy, 2,3-dimethylbutoxy and the like. "Halo" includes chloro, fluoro, bromo and iodo. The tertiary amino radical may be a dialkylamine such as a dimethyl, diethyl, dipropyl, dibutyl or a polymethyleneimine, e.g. piperidine, pyrrolidine, N-methyl piperazine or morpholine. Preferred compounds include those in which R1 is lower alkoxy; R2 and R3 are lower alkyl, especially methyl; R4 is H; and R5 is tertiary amino lower alkoxy of the polymethyleneimine

type. Within particularly preferred embodiments, R1 is in the 7-position and is lower alkoxy, particularly methoxy; each of R2 and R3 is methyl, R4 is H and R5 is in the 4-position and is a tertiary amino lower alkoxy radical such as pyrrolidinoethoxy.

It is preferred to use the compounds of structure (I) in the *trans* configuration. These compounds may be used as racemic mixtures, or the isolated *d*- or *l*-enantiomers may be used.

A particularly preferred compound for use within the present invention is centchroman (II):



15

Although only one enantiomer is shown, it will be understood that the structure II is used herein to designate the *trans* configuration of the 3- and 4-phenyl groups and that both the *d*- and *l*-enantiomers, as well as the racemic mixture, are included.

3,4-diarylchromans are prepared according to known methods, such as those disclosed in U.S. Patent No. 3,340,276 to Carney et al., U.S. Patent No. 3,822,287 to Bolger, and Ray et al., J. Med. Chem. 19:276-279, 1976.

Conversion of the *cis* isomer to the *trans* configuration by means of an

organometallic base-catalyzed rearrangement is disclosed in U.S. Patent No. 3,822,287. The optically active *d*- and *l*- enantiomers may be prepared as disclosed by Salman et al. in U.S. Patent No. 4,447,622

5 by forming an optically active acid salt which is subjected to alkaline hydrolysis to produce the desired enantiomer.

Within the present invention, 3,4-diarylchromans may be prepared in the form of pharmaceutically acceptable
10 salts, especially acid-addition salts, including salts of organic acids and mineral acids. Examples of such salts include salts of organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, succinic acid, malic acid, tartaric
15 acid, citric acid, benzoic acid, salicylic acid and the like. Suitable inorganic acid-addition salts include salts of hydrochloric, hydrobromic, sulfuric and phosphoric acids and the like. The acid addition salts may be obtained as the direct products of compound
20 synthesis. In the alternative, the free base may be dissolved in a suitable solvent containing the appropriate acid; and the salt isolated by evaporating the solvent or otherwise separating the salt and solvent.

3,4-diarylchromans and their salts are useful
25 within human and veterinary medicine for the regulation of bone metabolism. These compounds may be used, for example, in the treatment of patients suffering from bone loss due to osteoporosis (including post-menopausal osteoporosis and glucocorticoid-related osteoporosis),
30 Paget's disease, hyperparathyroidism, hypercalcemia of malignancy and other conditions characterized by excessive rates of bone resorption and/or decreased rates of bone formation.

For use within the present invention, 3,4-
35 diarylchromans and their pharmaceutically acceptable salts are formulated with a pharmaceutically acceptable carrier

to provide a medicament for parenteral, oral, nasal, rectal, subdermal or transdermal administration according to conventional methods. Formulations may further include one or more diluents, fillers, emulsifiers, preservatives, buffers, excipients, etc. and may be provided in such forms as liquids, powders, emulsions, suppositories, liposomes, transdermal patches, controlled release subdermal implants, tablets, etc. One skilled in this art may formulate the compounds in an appropriate manner, and in accordance with accepted practices, such as those disclosed in Remington's Pharmaceutical Sciences, Gennaro, ed., Mack Publishing Co., Easton, PA, 1990.

Oral administration is preferred. Thus, the active compound is prepared in a form suitable for oral administration, such as a tablet or capsule. Typically, a pharmaceutically acceptable salt of the compound is combined with a carrier and molded into a tablet. Suitable carriers in this regard include starch, sugars, dicalcium phosphate, calcium stearate, magnesium stearate and the like. Such compositions may further include one or more auxiliary substances, such as wetting agents, emulsifiers, preservatives, stabilizers, coloring, etc.

Pharmaceutical compositions are administered at daily to weekly intervals. An "effective amount" of such a pharmaceutical composition is the amount that provides a clinically significant inhibition of bone loss. Such amounts will depend, in part, on the particular condition to be treated, age, weight, and general health of the patient, and other factors evident to those skilled in the art. In general, inhibition of bone loss is manifested as a statistically significant difference in cancellous bone volume between treatment and control groups. This can be seen as, for example, a 5-10% or more difference in spinal bone mass or bone mineral content over two years. Data from accepted animal models, such as the ovariectomized

mouse or rat models of osteoporosis, are generally predictive of doses in humans to within one order of magnitude. For example, therapeutic doses for the treatment of osteoporosis will generally range from 0.01-50 mg/kg/day, preferably 0.05-10 mg/kg/day, most preferably 0.1-5.0 mg/kg/day. The use of *cis*-isomers or racemic mixtures may necessitate doses in the higher end of the stated range.

The pharmaceutical compositions may be administered in unit dosage form on a daily to weekly basis. In the alternative, they may be provided as controlled release formulations suitable for subdermal implantation. Implants are formulated to provide release of active compound over the desired period of time, which can be up to several years. Controlled-release formulations are disclosed by, for example, Sanders et al., J. Pharm. Sci. 73: 1294-1297, 1984; U.S. Patent No. 4,489,056; and U.S. Patent No. 4,210,644.

The following examples are offered by way of illustration, not limitation.

Example 1

The ability of centchroman to prevent osteopenia induced by estrogen deficiency was evaluated in the ovariectomized mouse model. Twenty-four female Swiss-Webster mice (8 weeks old) received either an ovariectomy or sham surgery prior to the initiation of a 4 week treatment protocol. For the ovariectomy, a flank incision through the skin, muscle and abdominal peritoneum was made on each side, the ovaries were located and dissected free of adherent fat and connective tissue, and excised. In the sham procedure the ovaries were exteriorized and replaced. In all animals the peritoneum and muscle were

2157879

- 9 -

sutured together and the skin incisions were closed with wound clips.

Centchroman was dissolved in a minimal amount of dimethylsulfoxide and diluted in oil vehicle to a concentration of 50 $\mu\text{g}/100 \mu\text{l}$. The mice were treated twice per week for 4 weeks with a subcutaneous injection of centchroman or oil vehicle according to the following outline: Sham/oil vehicle (SV); OVX/oil vehicle; OVX/50 μg centchroman two times per week. There were 8 animals in each group.

At the conclusion of the 4-week centchroman treatment, the mice were anesthetized with ether and sacrificed by cervical dislocation. Immediately after sacrifice, the femurs were removed and fixed in 70% ethyl alcohol (EtOH) and dehydrated in a series of increasing alcohol concentrations: 95% EtOH for 24 hours followed by three changes in 100% EtOH of 24 hours each. After the final 100% EtOH the femurs were cleared in two changes of xylene and then processed undecalcified and embedded in methacrylate plastic according to previously described methods (Bain et al., Stain Technology 65: 159-163, 1990). Frontal sections of the distal metaphyses of the femur 5 μm thick were cut on a Reichert-Jung 2050 rotary microtome equipped with a tungsten-carbide knife. The 5 μm sections were mounted on glass slides and stained with Goldner's trichrome stain.

Histomorphometric measurements of the distal metaphyses were determined using the Bioquant Bone Morphometry Program (Biometrics, Inc., Nashville, TN) interfaced via a camera lucida with an Olympus BH-2 light/epifluorescent microscope (Scientific Instruments, Inc., Redmond, WA). Morphometric measurements of cancellous bone volume (BV/TV) were performed in the tissue space greater than 0.25 mm from the growth plate-metaphyseal junction to exclude primary spongiosa.

2157879

- 10 -

Inc., Redmond, WA). Morphometric measurements of cancellous bone volume (BV/TV) were performed in the tissue space greater than 0.25 mm from the growth plate-metaphyseal junction to exclude primary spongiosa.

5 Data, shown in Figure 1, are expressed as the mean $\pm$ SD for each group. Comparison of cancellous bone volumes of the distal femur were based on analysis of variance using Statview® statistical programs (Abacus Concepts, Inc., Berkeley, CA). Treatment differences
10 indicated by the ANOVA were compared using Dunnett's multiple comparison procedure. A P value of less than .05 was considered significant.

In ovariectomized mice treated with oil vehicle a 50% decrease in the cancellous bone volume of the distal
15 femur compared to sham animals treated with vehicle was observed. In the ovariectomized animals treated with 50 μ g of centchroman twice per week this bone loss was completely prevented.

20 Example 2

To evaluate centchroman effects on bone resorption and bone mass, the skeletons of 54 female Sprague-Dawley rats were pre-labeled over four consecutive weeks with tritiated tetracycline ($^3\text{H-T}$; obtained from
25 Dupont NEN Research Products, Boston, MA). Animals were given 12-15 injections of 15 μCi each for a total of approximately 200 μCi per animal. Three days after the final $^3\text{H-T}$ injection, eight animals were sacrificed as baseline controls, and the remaining animals were
30 randomized for estrogen (E2) or centchroman (C) treatments according to the following outline: sham/placebo; ovariectomy(ovx)/placebo; ovariectomy/E2 (0.05 mg/kg/day); ovariectomy/C (0.05 mg/kg/day); ovariectomy/C (0.5 mg/kg/day); and ovariectomy/C (5.0 mg/kg/day). The
35 hormone treatments were delivered via subcutaneous pellet implants containing a matrix of cholesterol, lactose,

2157879

- 11 -

femurs and vertebrae were also determined to document changes in bone physical properties, and quantitative histomorphometry was used to compare changes in cancellous bone volumes of the proximal tibiae.

5 Sixty days after the initiation of the treatment protocols, the animals were anesthetized with ether and sacrificed by cervical dislocation. Immediately after sacrifice, the uteri were removed and weights recorded; both femurs and three thoracic vertebrae (T11-T13) were
10 excised for the bone resorption assay; one tibia and the first lumbar vertebra were collected for determination of bone physical properties; and the second tibia was excised and processed for bone histomorphometry. All tissues were fixed initially in 70% ethyl alcohol (EtOH) and dehydrated
15 in an ascending series of EtOH to 100%. After the final change of 100% EtOH, the specimens were processed according to the assay protocols outlined below.

 The assay of whole bone resorption was based on the levels of $^3\text{H-T}$ retained in the pre-labeled femurs and
20 vertebrae essentially as disclosed by Klein and Jackman (Calcified Tissue Research 20: 275-290, 1976). Briefly, the samples were defatted in three changes of chloroform of 24 hours each and dried at 100°C for 24 hours, and the weights were recorded. To extract the $^3\text{H-T}$, the femurs
25 and vertebrae were demineralized in 15 ml of 0.5 N hydrochloric acid (HCl), and the supernatants were decanted and reserved. To quantify tritium levels, 625 μl aliquots were pipetted into glass scintillation vials containing 10 ml of Optiflor scintillation fluid (Packard
30 Instruments, Meriden, CT), and the $^3\text{H-T}$ levels were counted on a liquid scintillation spectrometer (Beckman LS 1800).

 Following the EtOH dehydration, the samples for measuring bone mass were defatted in three changes of
35 chloroform of 24 hours each and dried in a 60°C oven

2157879

- 12 -

overnight. Bone mass was expressed as mg of dry weight per gram of body weight.

After the final 100% EtOH treatment, the tibiae were cleared in two changes of xylene, processed undecalcified and embedded in methacrylate plastic essentially as disclosed by Bain et al. (Stain Technology 65: 159-163, 1990). Frontal sections of the proximal tibiae 5 μ m thick were cut on a Reichert-Jung 2050 rotary microtome (Leica Instruments, Nusslock, Germany) equipped with a tungsten carbide knife. The 5 μ m sections were stained with Goldner's trichrome stain.

Histomorphometric measurements of the proximal tibiae were determined using a Bioquant Bone Morphometry program (Biometrics, Inc., Nashville, TN) interfaced via a camera lucida with an Olympus BH-2 light/epifluorescent microscope (Scientific Instruments, Inc., Redmond, WA). Morphometric measurements of cancellous bone volume (cn.BV/TV) were performed in a 3.0 mm² tissue space 1.5 millimeters from the growth plate-metaphyseal junction to exclude measurements of primary spongiosa. A minimum of four separate sections were measured from each animal.

Analyses of uterine weights, bone resorption assays, bone physical properties and bone histomorphometry were based on analysis of variance (ANOVA) using Statview® statistical programs (Abacus Concepts, Inc., Berkeley, CA). When significance was indicated by the ANOVA, control and treatment means were compared using Dunnett's multiple comparison procedure. P values of less than 0.05 were considered significant.

Compared to sham-vehicle treated animals, ovariectomy led to significant decreases in uterine wet weights. Estrogen replacement restored uterine weights to sham values, but centchroman treatment had no statistically significant effect on uterine weight, even at the highest dose of 5.0 mg per day. Indicative of increased bone resorption, ovariectomy reduced the

- 13 -

skeletal retention of $^3\text{H-T}$ in femurs and vertebrae (Tables 1 and 2, respectively). As expected, estrogen treatment increased the skeletal retention of $^3\text{H-T}$. Centchroman mimicked the effect of estrogen on bone resorption by causing a dose-dependent increase in the skeletal retention of $^3\text{H-T}$ at both skeletal sites (r^2 values equal 0.96 and 0.92 for femora and vertebrae, respectively).

Table 1

Retention of $^3\text{H-T}$ in Femurs

<u>Group</u>	<u>Count</u>	<u>cpm (thousands)</u>	
		<u>Mean</u>	<u>Std. Dev.</u>
sham/placebo	8	3832.25	928.975
ovx/placebo	8	3184.375	680.598
ovx/0.05 mg E2	10	4782	1501.813
ovx/0.05 mg C	9	3373.556	733.896
ovx/0.5 mg C	10	3405.1	1107.792
ovx/5.0 mg C	9	4192.222	1107.3

Table 2

Retention of $^3\text{H-T}$ in Vertebrae

<u>Group</u>	<u>Count</u>	<u>cpm (thousands)</u>	
		<u>Mean</u>	<u>Std. Dev.</u>
sham/placebo	8	1876.75	481.552
ovx/placebo	8	1531.625	416.205
ovx/0.05 mg E2	9	2467.556	733.064
ovx/0.05 mg C	9	1523.333	406.635
ovx/0.5 mg C	10	1616	509.954
ovx/5.0 mg C	10	2156.7	495.741

The ability of centchroman to inhibit bone resorption and prevent bone loss was confirmed by measurements of cancellous bone volume in the tibiae and bone mass determinations of femora and vertebrae. Compared to ovariectomized rats treated with vehicle, centchroman caused a dose-dependent increase in cancellous bone volume of the proximal tibiae (Figure 2; $r^2 = 0.99$). Similarly, centchroman had dose-dependent effects on bone mass in femora and vertebrae.

In summary, these data indicate that the ability of centchroman to prevent bone loss in the ovariectomized


2157879

rat is independent of any apparent uterotrophic activity.
This is clearly demonstrated in Figure 3 by combining
uterine weight data with the bone resorption data from the
femur to show the independent effects of centchroman on
5 these two tissues.

Although the foregoing invention has been
described in some detail by way of illustration and
example for purposes of clarity of understanding, it will
10 be evident that certain changes and modifications may be
practiced within the scope of the appended claims.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS.

1. Use of a compound of the formula



for the preparation of a pharmaceutical composition for reducing bone loss due to osteoporosis, Paget's disease or hyperparathyroidism, wherein:

R1, R4 and R5 are individually hydrogen, hydroxy, halo, trifluoromethyl, lower alkyl, lower alkoxy or tertiary amino lower alkoxy; and

R2 and R3 are individually hydrogen or lower alkyl.

2. The use according to claim 1 wherein said compound is an isolated *l*-enantiomer.

3. The use according to claim 1 or 2 wherein R1 is lower alkoxy, R2 and R3 are lower alkyl, R4 is hydrogen and R5 is tertiary amino lower alkoxy.


4. The use according to claim 1 or 2 wherein R1 is methoxy.

2157879

5. The use according to claim 1 or 2 wherein R2 and R3 are methyl.


6. The use according to claim 1 or 2 wherein R4 is hydrogen.

7. The use according to claim 1 or 2 wherein R5 is




8. The use according to claim 1 wherein said compound is an isolated *d*- or *l*-enantiomer.

9. The use according to claim 1 wherein said compound is



2157879

10. The use according to claim 1 wherein said compound is



11. The use according to claim 10 wherein said compound is an isolated *d*- or *l*-enantiomer.


12. The use according to claim 10 wherein said compound is an isolated *l*-enantiomer.

13. The use according to claim 1 or 2 wherein said osteoporosis, Paget's disease or hyperparathyroidism is present in a post-menopausal female.

14. The use according to claim 1 or 2 wherein said composition is in a form suitable for oral administration.

15. The use according to claim 1 or 2 wherein said composition is in suitable form for subdermal implant.

16. Use of a composition comprising an isolated *d*- or *l*-enantiomer of a compound of the formula



or a pharmaceutically acceptable salt thereof, for the preparation of a pharmaceutical composition for reducing bone loss due to osteoporosis, wherein:

R1, R4 and R5 are individually hydrogen, hydroxy, halo, trifluoromethyl, lower alkyl, lower alkoxy or tertiary amino lower alkoxy; and

R2 and R3 are individually hydrogen or lower alkyl;
in an amount sufficient to inhibit bone resorption.

17. The use of a composition of claim 16 wherein said isolated compound is the *l*-enantiomer.


18. The use according to claim 16 or 17 wherein R1 is lower alkoxy, R2 and R3 are lower alkyl, R4 is hydrogen and R5 is tertiary amino lower alkoxy.

19. The use according to claim 16 or 17 wherein R1 is methoxy.


20. The use according to claim 16 or 17 wherein R2 and R3 are methyl.

21. The use according to claim 16 or 17 wherein R4 is hydrogen.

22. The use according to claim 16 or 17 wherein R5 is




23. The use according to claim 16 or 17 wherein said compound is




2157879

24. The use according to claim 16 or 17 wherein said compound is



25. A pharmaceutical composition suitable for reducing bone loss in a human, which comprises as an active ingredient a bone loss inhibiting compound of the formula




wherein R1, R4 and R5 are individually selected from the group consisting of hydrogen, hydroxy, halo, trifluoromethyl, lower alkyl, lower alkoxy and tertiary

2157879

amino lower alkoxy and R2 and R3 are individually hydrogen or lower alkyl; or a pharmaceutically acceptable salt thereof, in an amount effective to inhibit bone loss associated with osteoporosis, Paget's disease or hyperparathyroidism, in admixture with a pharmaceutically acceptable diluent or carrier, or other biologically compatible vehicle.

(10696)

26. The use of a composition comprising a bone loss inhibiting compound of the formula



or a pharmaceutically acceptable salt thereof, wherein R1, R4 and R5 are individually hydrogen, hydroxy, halo, trifluoromethyl, lower alkyl, lower alkoxy or tertiary amino lower alkoxy; and R2 and R3 are individually hydrogen or lower alkyl, in combination with a pharmaceutically acceptable carrier for reducing or preventing bone loss in a patient.


27. The use according to claim 26 wherein R1 is lower alkoxy, R2 and R3 are lower alkyl, R4 is hydrogen and R5 is tertiary amino lower alkoxy.

28. The use according to claim 26 wherein R1 is methoxy.

29. The use according to claim 26 wherein R2 and R3 are methyl.


30. The use according to claim 26 wherein R4 is hydrogen.

31. The use according to claim 26 wherein R5 is:




32. The use according to claim 26 wherein said compound is an isolated d- or l-enantiomer.

5 33. The use according to claim 26 wherein said compound is:



34. The use according to claim 26 wherein said compound is:



30 35. The use according to claim 26 wherein said compound is an isolated d- or l-enantiomer.

35 36. The use according to claim 26 wherein said patient is a post-menopausal female.


37. The use according to claim 26 wherein said composition is in a form suitable for oral administration.

38. The use according to claim 26 wherein said compound is used at a dose of 0.1 - 5.0 mg/kg patient weight/day.

5 39. The use according to claim 26 wherein said composition is administered at daily to weekly intervals.

40. The use according to claim 26 wherein said composition is in the form of a subdermal implant.

10 41. The use according to claim 26 wherein the compound is an isolated l-enantiomer.

1/3

*P<.05 decreased compared to sham animals treated with vehicle and ovx mice treated with centchroman; n = 8 mice/group.

Figure 1

2/3





Figure 2

2157879

3/3



a=significantly decreased vs ovx/E2 and ovx/C groups; $P < .05$.
 b=significantly decreased vs sham and ovx/E2 groups; $P < .01$.

Figure 3