

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2013217114 B2

(54) Title
CD47 antibodies and methods of use thereof

(51) International Patent Classification(s)
C12P 21/08 (2006.01) **A61K 39/395** (2006.01)

(21) Application No: **2013217114** (22) Date of Filing: **2013.02.06**

(87) WIPO No: **WO13/119714**

(30) Priority Data

(31) Number (32) Date (33) Country
61/595,216 **2012.02.06** **US**
61/659,752 **2012.06.14** **US**

(43) Publication Date: **2013.08.15**
(44) Accepted Journal Date: **2017.03.30**

(71) Applicant(s)
Inhibrx LP

(72) Inventor(s)
Eckelman, Brendan;Timmer, John;Razai, Amir;Deveraux, Quinn;Jones, Kyle;Nguy, Peter L.

(74) Agent / Attorney
Griffith Hack, GPO Box 1285, Melbourne, VIC, 3001, AU

(56) Related Art
US 2011/0014119

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2013/119714 A1

(43) International Publication Date
15 August 2013 (15.08.2013)

WIPO | PCT

(51) International Patent Classification:
CI2P 21/08 (2006.01) *A61K 39/395* (2006.01)

(74) Agents: **ELRIFI, Ivor, R.** et al.; Mintz Levin Cohn Ferris Glovsky, And Popeo, P.c., One Financial Center, Boston, MA 02111 (US).

(21) International Application Number:

PCT/US2013/024995

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
6 February 2013 (06.02.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/595,216 6 February 2012 (06.02.2012) US
61/659,752 14 June 2012 (14.06.2012) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US 61/659,752 (CIP)
Filed on 14 June 2012 (14.06.2012)
US 61/595,216 (CIP)
Filed on 6 February 2012 (06.02.2012)

(71) Applicant: **INHIBRX LLC** [US/US]; 11099 N. Torrey Pines Road, Suite 130, La Jolla, CA 92037 (US).

(72) Inventors: **ECKELMAN, Brendan**; 11099 N. Torrey Pines Road, Suite 130, La Jolla, CA 92037 (US). **TIM-MER, John**; 11099 N. Torrey Pines Road, Suite 130, La Jolla, CA 92037 (US). **RAZAI, Amir**; 11099 N. Torrey Pines Road, Suite 130, La Jolla, CA 92037 (US). **DE-VERAUX, Quinn**; 11099 N. Torrey Pines Road, Suite 130, La Jolla, CA 92037 (US). **JONES, Kyle**; 11099 N. Torrey Pines Road, Suite 130, La Jolla, CA 92037 (US). **NGUY, Peter, L.**; 11099 N. Torrey Pines Road, Suite 130, La Jolla, CA 92037 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

WO 2013/119714 A1

(54) Title: CD47 ANTIBODIES AND METHODS OF USE THEREOF

(57) Abstract: This invention relates generally to monoclonal antibodies that recognize CD47, more specifically to CD47 antibodies that do not cause a significant level of agglutination of cells, to methods of generating these antibodies, and to methods of using these monoclonal antibodies as therapeutics.

CD47 Antibodies and Methods of Use Thereof

RELATED APPLICATIONS

[0001] This application claims the benefit of priority to U.S. Provisional Application No: 61/595,216, filed February 6, 2012 and to U.S. Provisional Application No. 61/659,752, filed June 14, 2012. Each of these applications is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0002] This invention relates generally to monoclonal antibodies that recognize CD47, more specifically to CD47 antibodies that do not cause a significant level of hemagglutination of human red blood cells, to methods of generating these antibodies, and to methods of using these monoclonal antibodies as therapeutics.

BACKGROUND OF THE INVENTION

[0003] CD47, also known as integrin-associated protein (IAP), ovarian cancer antigen OA3, Rh-related antigen and MER6, is a multi-spanning transmembrane receptor belonging to the immunoglobulin superfamily. CD47 expression and/or activity have been implicated in a number of diseases and disorders. Accordingly, there exists a need for therapies that target CD47.

[0003a] It is to be understood that if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art in Australia or any other country.

SUMMARY OF THE INVENTION

[0004] The present invention provides monoclonal antibodies that recognize and bind to CD47, particularly human CD47. The antibodies of the invention are capable of modulating, *e.g.*, blocking, inhibiting, reducing, antagonizing, neutralizing or otherwise interfering with CD47 expression, activity and/or signaling, and these antibodies do not cause a significant level of hemagglutination of human red blood cells, also referred to herein as erythrocytes. However, the ability of the antibodies of the present invention to bind CD47 on the cell surface and not cause a cellular clumping phenomenon is not limited to red blood

cells. The antibodies of the present invention uniquely bind CD47 in a manner that does not promote clumping of CD47 positive cells. The antibodies of the invention and derivatives thereof are capable of modulating, *e.g.*, blocking, inhibiting, reducing, antagonizing, neutralizing or otherwise interfering with the interaction between CD47 and SIRP (signal-regulatory-protein), and these antibodies do not cause a significant level of hemagglutination of human red blood cells. The antibodies provided herein are referred to collectively as “CD47 antibodies.” The CD47 antibodies of the invention are a significant improvement over existing CD47 antibodies that cause hemagglutination of human red blood cells (*See, e.g.*, Kikuchi Y, Uno S, Yoshimura Y et al. A bivalent single-chain Fv fragment against CD47 induces apoptosis for leukemic cells. *Biochem Biophys Res Commun* 2004; 315: 912–8). For example, the CD47 antibodies of the invention are a significant improvement over the existing CD47 antibodies B6H12, BRC126, and CC2C6, each of which block SIRP , but cause hemagglutination of RBCs, as described in detail below. The full IgG CD47 antibodies of the present invention (*e.g.*, 2A1 and its humanized derivatives including those provided in Table 1) do not agglutinate cells at a significant level. For example, the CD47 antibodies of the invention do not hemagglutinate red blood cells (RBCs). Described herein are the first CD47 antibodies in a full IgG format that block SIRP and do not cause a significant level of agglutination.

[0004a] A first aspect provides an isolated monoclonal antibody or immunologically active fragment thereof that binds CD47, wherein the antibody or immunologically active fragment thereof comprises a VH complementarity determining region 1 (CDR1) sequence set forth in SEQ ID NO: 50, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 64, SEQ ID NO: 65, or SEQ ID NO: 66, a VH CDR2 sequence set forth in SEQ ID NO: 51, SEQ ID NO: 72, SEQ ID NO: 73, SEQ ID NO: 74, SEQ ID NO: 75, or SEQ ID NO: 76, a VH CDR3 sequence set forth in SEQ ID NO: 52 or SEQ ID NO: 77, a VL CDR1 sequence set forth in SEQ ID NO: 53, SEQ ID NO: 67, or SEQ ID NO: 68, a VL CDR2 sequence set forth in SEQ ID NO: 54, SEQ ID NO: 69, SEQ ID NO: 70, or SEQ ID NO: 71 and a VL CDR3 sequence set forth in SEQ ID NO: 55.

[0004b] A second aspect provides an isolated monoclonal antibody or immunologically active fragment thereof that binds to human CD47, said antibody or immunologically active

fragment thereof comprising:

a VH CDR1, a VH CDR2, and a VH CDR3, respectively comprising the amino acid sequences of a CDR1, a CDR2, and a CDR3 within a VH chain region having the sequence set forth in any of SEQ ID NOs: 5-30; and

a VL CDR1, a VL CDR2, and a VL CDR3, respectively comprising the amino acid sequences of a CDR1, a CDR2, and a CDR3 within a VL chain region having the sequence set forth in any of SEQ ID NOs: 31-47.

[0004c] A third aspect provides a pharmaceutical composition comprising the antibody or immunologically active fragment thereof of the first or second aspect and a carrier.

[0004d] A fourth aspect provides a method of alleviating a symptom of a cancer or other neoplastic condition, the method comprising administering the antibody or immunologically active fragment thereof of the first or second aspect to a subject in need thereof in an amount sufficient to alleviate the symptom of the cancer or other neoplastic condition in the subject.

[0004e] A fifth aspect provides isolated nucleic acid(s) encoding the antibody or immunologically active fragment thereof of the first or second aspect.

[0004f] A sixth aspect provides an isolated host cell comprising nucleic acid(s) encoding the antibody or immunologically active fragment thereof of the first or second aspect.

[0004g] A seventh aspect provides use of the antibody or immunologically active fragment thereof of the first or second aspect in the manufacture of a medicament for alleviating a symptom of a cancer or other neoplastic condition.

[0005] The CD47 antibodies of the invention exhibit numerous desirable characteristics, such as, by way of non-limiting example, potent blocking of the interaction between CD47 and its ligand SIRP_α, without causing a significant level of hemagglutination

of erythrocytes, as well as potent anti-tumor activity. For example, the CD47 antibodies of the invention block at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 95%, or at least 99% of the interaction between CD47 and SIRP as compared to the level of interaction between CD47 and SIRP in the absence of the CD47 antibody described herein.

[0006] The CD47 antibodies of the invention do not cause a significant level of agglutination of cells, *e.g.*, the CD47 antibodies of the invention do not cause a significant level of hemagglutination of red blood cells. In some cases, a significant level of agglutination of cells refers to the level of agglutination in the presence of existing CD47 antibodies. In one embodiment, the level of agglutination in the presence of the CD47 antibodies of the invention is reduced by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 99% compared to the level of agglutination in the presence existing CD47 antibodies. In some embodiments, the CD47 antibodies of the invention do not cause a significant level of agglutination if the level of agglutination in the presence of the CD47 antibodies of the

invention is reduced by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 99% compared to the level of agglutination in the presence of existing CD47 antibodies. In other embodiments, the CD47 antibodies of the invention do not cause a significant level of agglutination if the level of agglutination in the presence of the CD47 antibodies of the invention is reduced by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 99% compared to the level of agglutination in the presence of CD47 antibody, 1B4, which comprises a variable heavy and variable light chain sequence provided in SEQ ID NO: 80 and SEQ ID NO: 81, respectively. Preferably, the CD47 antibodies of the invention do not cause a significant level of agglutination of cells at an antibody concentration of between 10 pM and 10 μ M, *e.g.*, at an antibody concentration of 50 pM, 100 pM, 1 nM, 10 nM, 50 nM, 100 nM, 1 μ M, or 5 μ M.

[0007] The antibodies of the present invention are also significantly more potent in tumor models compared to antibodies known in the art. For example, the ability of macrophages to phagocytose tumor cells in the presence of CD47 antibodies of the invention is increased by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 99% compared to the ability of macrophages to phagocytose tumor cells in the presence of existing CD47 antibodies.

[0008] Those skilled in the art will recognize that it is possible to quantitate, without undue experimentation, the level of agglutination, *e.g.*, the level of hemagglutination of RBCs. For example, those skilled in the art will recognize that the level of hemagglutination is ascertained by measuring the area of an RBC dot after performing a hemagglutination assay in the presence of the CD47 antibodies of the invention, as described in the Examples below. In some cases, the area of the RBC dot in the presence of the CD47 antibody of the invention is compared to the area of the RBC dot in the absence of a CD47 antibody, *i.e.*, in the presence of zero hemagglutination. In this manner, hemagglutination is quantified relative to a baseline control. A larger RBC dot area corresponds to a higher level of hemagglutination. Alternatively, densitometry of the RBC dot may also be utilized to quantitate hemagglutination.

[0009] The CD47 antibodies described herein are useful in treating, delaying the progression of, preventing relapse of or alleviating a symptom of a cancer or other neoplastic condition. For example, the CD47 antibodies described herein are useful in

treating hematological malignancies and/or tumors, *e.g.*, hematological malignancies and/or tumors. For example, the CD47 antibodies described herein are useful in treating CD47+ tumors. By way of non-limiting example, the CD47 antibodies described herein are useful in treating non-Hodgkin's lymphoma (NHL), acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), multiple myeloma (MM), breast cancer, ovarian cancer, head and neck cancer, bladder cancer, melanoma, colorectal cancer, pancreatic cancer, lung cancer, leiomyoma, leiomyosarcoma, glioma, glioblastoma, and so on. Solid tumors include, *e.g.*, breast tumors, ovarian tumors, lung tumors, pancreatic tumors, prostate tumors, melanoma tumors, colorectal tumors, lung tumors, head and neck tumors, bladder tumors, esophageal tumors, liver tumors, and kidney tumors.

[0010] As used herein, "hematological cancer" refers to a cancer of the blood, and includes leukemia, lymphoma and myeloma among others. "Leukemia" refers to a cancer of the blood in which too many white blood cells that are ineffective in fighting infection are made, thus crowding out the other parts that make up the blood, such as platelets and red blood cells. It is understood that cases of leukemia are classified as acute or chronic. Certain forms of leukemia include, by way of non-limiting example, acute lymphocytic leukemia (ALL); acute myeloid leukemia (AML); chronic lymphocytic leukemia (CLL); chronic myelogenous leukemia (CML); Myeloproliferative disorder/neoplasm (MPDS); and myelodysplasia syndrome. "Lymphoma" may refer to a Hodgkin's lymphoma, both indolent and aggressive non-Hodgkin's lymphoma, Burkitt's lymphoma, and follicular lymphoma (small cell and large cell), among others. Myeloma may refer to multiple myeloma (MM), giant cell myeloma, heavy-chain myeloma, and light chain or Bence-Jones myeloma.

[0011] Exemplary monoclonal antibodies of the invention include, for example, the antibodies described herein. Exemplary antibodies include antibodies having a variable heavy chain selected from SEQ ID NOs: 5-30 and a variable light chain selected from SEQ ID NOs: 31-47. The antibodies also include antibodies having a variable heavy chain that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the sequence set forth in at least one of SEQ ID NOs: 5-30 and a variable light chain that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the sequence set forth in at least one of SEQ ID NOs: 31-47. Preferably, the antibodies recognize and bind to human CD47 and do not cause a significant level of hemagglutination of human red blood cells. These antibodies are respectively referred to herein as CD47

antibodies. CD47 antibodies include fully human monoclonal antibodies, as well as humanized monoclonal antibodies and chimeric antibodies. These antibodies show specificity for human CD47, and they have been shown to modulate, *e.g.*, block, inhibit, reduce, antagonize, neutralize or otherwise interfere with CD47 expression, activity and/or signaling without causing a significant level of hemagglutination of red blood cells.

[0012] The CD47 antibodies provided herein exhibit inhibitory activity, for example by inhibiting CD47 expression (*e.g.*, inhibiting cell surface expression of CD47), activity, and/or signaling, or by interfering with the interaction between CD47 and SIRPa. The antibodies provided herein completely or partially reduce or otherwise modulate CD47 expression or activity upon binding to, or otherwise interacting with, CD47, *e.g.*, a human CD47. The reduction or modulation of a biological function of CD47 is complete, significant, or partial upon interaction between the antibodies and the human CD47 polypeptide and/or peptide. The antibodies are considered to completely inhibit CD47 expression or activity when the level of CD47 expression or activity in the presence of the antibody is decreased by at least 95%, *e.g.*, by 96%, 97%, 98%, 99% or 100% as compared to the level of CD47 expression or activity in the absence of interaction, *e.g.*, binding, with the antibody described herein. The CD47 antibodies are considered to significantly inhibit CD47 expression or activity when the level of CD47 expression or activity in the presence of the CD47 antibody is decreased by at least 50%, *e.g.*, 55%, 60%, 75%, 80%, 85% or 90% as compared to the level of CD47 expression or activity in the absence of binding with a CD47 antibody described herein. The antibodies are considered to partially inhibit CD47 expression or activity when the level of CD47 expression or activity in the presence of the antibody is decreased by less than 95%, *e.g.*, 10%, 20%, 25%, 30%, 40%, 50%, 60%, 75%, 80%, 85% or 90% as compared to the level of CD47 expression or activity in the absence of interaction, *e.g.*, binding, with an antibody described herein.

[0013] Antibodies of the invention also include monoclonal antibodies that specifically bind CD47, wherein the antibody does not cause a significant level of agglutination, *e.g.*, red blood cell hemagglutination (“RBC hemagglutination”). The antibodies of the present invention uniquely bind CD47 in a manner that does not promote clumping of CD47 positive cells; however, the ability of the antibodies of the present invention to bind CD47 on the cell surface and not cause a cellular clumping phenomenon is not limited to red blood cells.

[0014] Pharmaceutical compositions according to the invention can include an antibody of the invention and a carrier. These pharmaceutical compositions can be included in kits, such as, for example, diagnostic kits.

[0015] The invention provides monoclonal antibodies that bind to CD47 or an immunologically active fragment thereof, wherein the antibody does not cause a significant level of agglutination of cells after administration, e.g., the antibody does not cause a significant level of hemagglutination of red blood cells after administration. In some embodiments, the antibody is chimeric, humanized, or fully human. In some embodiments, the antibodies bind to human CD47. In some embodiments, the antibody or immunologically active fragment thereof prevents CD47 from interacting with SIRP α . The antibodies are considered to completely inhibit the interaction of CD47 and SIRP α when the level of CD47/SIRP α interaction in the presence of the antibody is decreased by at least 95%, e.g., by 96%, 97%, 98%, 99% or 100% as compared to the level of CD47/SIRP α interaction in the absence of interaction with the antibody, e.g., binding with the antibody. The antibodies are considered to partially inhibit CD47/SIRP α interaction when the level of CD47/SIRP α interaction in the presence of the antibody is decreased by less than 95%, e.g., 10%, 20%, 25%, 30%, 40%, 50%, 60%, 75%, 80%, 85% or 90% as compared to the level of CD47/SIRP α interaction in the absence of interaction with the antibody, e.g., binding with the antibody.

[0016] The amount of antibody sufficient to treat or prevent cancer in the subject is, for example, an amount that is sufficient to reduce CD47 signaling (See, e.g., Yamauchi et al., 2013 Blood, Jan 4. [Epub ahead of print]; Soto-Pantoja et al., 2013 Expert Opin Ther Targets, 17: 89-103; Irandoust et al., 2013 PLoS One, Epub Jan 8; Chao et al., 2012 Curr Opin Immunol, 24:225-32; Theocharides et al., 2012 J Exp Med, 209(10): 1883-99; Csányi et al., 2012 Arterioscler Thromb Vasc Biol, 32: 2966-73; Maxhimer et al., 2009 Sci Transl Med, 1: 3ra7; Sarfati et al., 2008 Curr Drug Targets, 9: 842-850; Miyashita et al., 2004 Mol Biol Cell, 15: 3950-3963; E.J. Brown and W.A. Frazier, 2001 Trends Cell Biol, 11: 130-135; Oldenborg et al., 2001 J Exp Med, 193: 855-862; Blazar et al., 2001 J Exp Med, 194: 541-549; Oldenborg et al., 2000 Science, 288: 2051-2054; and Gao et al., 1996 J Biol Chem, 271: 21-24). For example, the amount of antibody sufficient to treat or prevent cancer in the subject is an amount that is sufficient to reduce the phagocytic inhibitory signal in macrophages generated by CD47/SIRP α interaction in the CD47/SIRP α signaling axis, i.e., the antibody of the invention promotes macrophage-mediated phagocytosis of a CD47-expressing cell. As used herein, the term “reduced” refers to a decreased CD47

signaling in the presence of the antibody of the invention. CD47 mediated signaling is decreased when the level of CD47 signaling in the presence of a CD47 antibody of the invention is greater than or equal to 5%, 10%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 75%, 80%, 90%, 95%, 99%, or 100% lower than a control level of CD47 signaling (*i.e.*, the level of CD47 signaling in the absence of the antibody). Level of CD47 signaling is measured using any of a variety of standard techniques, such as, by way of non-limiting example, measurement of down-stream gene activation, and/or luciferase reporter assays responsive to CD47 activation. Those skilled in the art will appreciate that the level of CD47 signaling can be measured using a variety of assays, including, for example, commercially available kits.

[0017] In some embodiments, the antibody or immunologically active fragment thereof is an IgG isotype. In some embodiments, the constant region of the antibody is of human IgG1 isotype, having an amino acid sequence:

ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV
HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDKKVEP
KSCDKTHTCP PCPAPE~~LL~~GG PSVFLFPPKP KDTLMISRTP EVTCVVVDVS
HEDPEVKFNW YVDGVEVHNA KTKPREEQY~~N~~ STYRVVSVLT VLHQDWLNGK
EYKCKVSNKA LPAPIEKTI S KAKGQPREPQ VYTLPPSRDE LTKNQVSLTC
LVKGFYPSDI AVEWESNGQP ENNYKTPPPV LDSDGSFFLY SKLTVDKSRW
QQGNVFSCSV MHEALHNHYT QKSLSLSPGK (SEQ ID NO: 1)

[0018] In some embodiments, the human IgG1 constant region is modified at amino acid Asn297 (Boxed, Kabat Numbering) to prevent glycosylation of the antibody, for example Asn297Ala (N297A). In some embodiments, the constant region of the antibody is modified at amino acid Leu235 (Kabat Numbering) to alter Fc receptor interactions, for example Leu235Glu (L235E) or Leu235Ala (L235A). In some embodiments, the constant region of the antibody is modified at amino acid Leu234 (Kabat Numbering) to alter Fc receptor interactions, *e.g.*, Leu234Ala (L234A). In some embodiments, the constant region of the antibody is altered at both amino acid 234 and 235, for example Leu234Ala and Leu235Ala (L234A/L235A) (EU index of *Kabat et al 1991 Sequences of Proteins of Immunological Interest*).

[0019] In some embodiments, the constant region of the antibody is of human IgG2 isotype, having an amino acid sequence:

ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV
HTFPAVLQSS GLYSLSSVVT VPSSNFGTQT YTCNVDHKPS NTKVDKTVER
KCCVECPPCP APPVAGPSVF LFPPKPDKTL MISRTPEVTC VVVDVSHEDP
EVQFNWYVDG VEVHNAKTKP REEQF~~N~~STFR VVSVLTVVHQ DWLNGKEYKC
KVSNKGLPAP IEKTISKTG QPREPQVYTL PPSREEMTKN QVSLTCLVKKG

FYPSDISVEW ESNGQPENNY KTPPPMLDSD GSFFFLYSKLT VDKSRWQQGN
VFSCSVMHEA LHNHYTQKSL SLSPGK (SEQ ID NO: 2)

[0020] In some embodiments, the human IgG2 constant region is modified at amino acid Asn297 (Boxed, Kabat Numbering) to prevent to glycosylation of the antibody, *e.g.*, Asn297Ala (N297A).

[0021] In some embodiments, the constant region of the antibody is of human IgG3 isotype, having an amino acid sequence:

ASTKGPSVFP LAPCSRSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV
HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT YTCNVNHKPS NTKVDKRVEL
KTPLGDTTHT CPRCPEPKSC DTTPPCPRCP EPKSCDTPPP CPRCPEPKSC
DTTPPCPRCP APELLGGPSV FLFPPPKPKDT LMISRTPEVT CVVVDVSCHED
PEVQFKWYVD GVEVHNNAKTK PREEQY~~N~~STF RVVSVLTVLH QDWLNGKEYK
CKVSNKALPA PIEKTISKTK GQPREPQVYT LPPSREEMTK NQVSLTCLVK
GFYPSDIAVE WESSGQPENN YNTTPPMLDS DGSSFLYSKLT VDKSRWQQG
NIFSCSVMHE ALHN~~R~~FTQKS LSLSPGK (SEQ ID NO: 3)

[0022] In some embodiments, the human IgG3 constant region is modified at amino acid Asn297 (Boxed, Kabat Numbering) to prevent to glycosylation of the antibody, *e.g.*, Asn297Ala (N297A). In some embodiments, the human IgG3 constant region is modified at amino acid 435 to extend the half-life, *e.g.*, Arg435His (R435H) (EU index of *Kabat et al 1991 Sequences of Proteins of Immunological Interest*).

[0023] In some embodiments, the constant region of the antibody is of human IgG4 isotype, having an amino acid sequence:

ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV
HTFPAVLQSS GLYSLSSVVT VPSSSLGTKT YTCNVDHKPS NTKVDKRVES
KYGPPCP~~S~~CP APEF~~L~~GGPSV FLFPPPKPKDT LMISRTPEVT CVVVDVSQED
PEVQFNWYVD GVEVHNNAKTK PREEQF~~N~~STY RVVSVLTVLH QDWLNGKEYK
CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK NQVSLTCLVK
GFYPSDIAVE WESNGQPENN YKTTPPVLDS DGSSFLYSRLL VDKSRWQEG
NVFSCSVMHE ALHNHYTQKS LSLSLGK (SEQ ID NO: 4)

[0024] In some embodiments, the human IgG4 constant region is modified within the hinge region to prevent or reduce strand exchange, *e.g.*, Ser228Pro (S228P). In other embodiments, the human IgG4 constant region is modified at amino acid 235 to alter Fc receptor interactions, *e.g.*, Leu235Glu (L235E). In some embodiments, the human IgG4 constant region is modified within the hinge and at amino acid 235, *e.g.*, Ser228Pro and Leu235Glu (S228P/L235E). In some embodiments, the human IgG4 constant region is modified at amino acid Asn297 (Kabat Numbering) to prevent to glycosylation of the

antibody, *e.g.*, Asn297Ala (N297A). (EU index of *Kabat et al* 1991 *Sequences of Proteins of Immunological Interest*).

[0025] In some embodiments, the human IgG constant region is modified to enhance FcRn binding. Examples of Fc mutations that enhance binding to FcRn are Met252Tyr, Ser254Thr, Thr256Glu (M252Y, S254T, T256E, respectively) (Kabat numbering, Dall'Acqua *et al* 2006, *J. Biol Chem* Vol 281(33) 23514–23524), or Met428Leu and Asn434Ser (M428L, N434S) (Zalevsky *et al* 2010 *Nature Biotech*, Vol 28(2) 157-159). (EU index of *Kabat et al* 1991 *Sequences of Proteins of Immunological Interest*). In some embodiments, the human IgG constant region is modified to alter antibody-dependent cellular cytotoxicity (ADCC) and/or complement-dependent cytotoxicity (CDC), *e.g.*, the amino acid modifications described in Natsume *et al.*, 2008 *Cancer Res*, 68(10): 3863-72; Idusogie *et al.*, 2001 *J Immunol*, 166(4): 2571-5; Moore *et al.*, 2010 *mAbs*, 2(2): 181-189; Lazar *et al.*, 2006 *PNAS*, 103(11): 4005–4010, Shields *et al.*, 2001 *JBC*, 276(9): 6591–6604; Stavenhagen *et al.*, 2007 *Cancer Res*, 67(18): 8882-8890; Stavenhagen *et al.*, 2008 *Advan. Enzyme Regul.*, 48: 152-164; Alegre *et al*, 1992 *J Immunol*, 148: 3461-3468; Reviewed in Kaneko and Niwa, 2011 *Biodrugs*, 25(1):1-11.

[0026] In some embodiments, the human IgG constant region is modified to induce heterodimerization. For example, having an amino acid modification within the CH3 domain at Thr366, which when replaced with a more bulky amino acid, *e.g.*, Try (T366W), is able to preferentially pair with a second CH3 domain having amino acid modifications to less bulky amino acids at positions Thr366, Leu368, and Tyr407, *e.g.*, Ser, Ala and Val, respectively (T366S/L368A/Y407V). Heterodimerization via CH3 modifications can be further stabilized by the introduction of a disulfide bond, for example by changing Ser354 to Cys (S354C) and Y349 to Cys (Y349C) on opposite CH3 domains (Reviewed in Carter, 2001 *Journal of Immunological Methods*, 248: 7–15).

[0027] The invention also provides pharmaceutical compositions that include one or more monoclonal antibodies that bind to CD47 or an immunologically active fragment thereof, wherein the antibody does not cause a significant level of hemagglutination of red blood cells after administration.

[0028] Hemagglutination is an example of a homotypic interaction, wherein two CD47 expressing cells are caused to aggregate or clump when treated with a bivalent CD47 binding entity. The ability of the antibodies of the present invention to bind CD47 on the cell surface and not cause a cellular clumping phenomenon is not limited to red blood cells.

The antibodies of the present invention have been observed to uniquely bind CD47 in a manner that does not promote clumping of CD47 positive cell lines, *e.g.*, Daudi cells.

[0029] In some cases, the antibody comprises a variable heavy (VH) chain region selected from the group consisting of SEQ ID NOs: 5-30. The antibody optionally comprises a variable light (VL) chain region selected from the group consisting of SEQ ID NOs: 31-47. In some cases, the antibody comprises a VH chain region selected from the group consisting of SEQ ID NOs: 5-30 and a VL chain region selected from the group consisting of SEQ ID NOs: 31-47. The antibodies of the invention also include antibodies having a variable heavy chain that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the sequence set forth in at least one of SEQ ID NOs: 5-30 and a variable light chain that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identical to the sequence set forth in at least one of SEQ ID NOs: 31-47. In other embodiments, the antibody comprises a VH region provided in any one of SEQ ID NOs: 5, 7, 8, 11, 15-17, 20-22, and 27-30 paired with a VL region provided in any one of SEQ ID NOs: 31-39, 42, 43, 44, and 47. In another embodiment, the antibody comprises a VH region provided in any one of SEQ ID NOs: 5, 7, 8, 11, 12, 15-17, 20-22, and 27-30 paired with a VL region provided in any one of SEQ ID NOs: 31, 32, 35, 40, 41, 42, 43, 44, and 47. In yet another embodiment, the antibody comprises a combination of a VH chain region and a VL chain region selected from the combinations listed in Table 1.

[0030] In some embodiments, the CD47 antibody or immunologically active fragment thereof comprises a VH complementarity determining region 1 (CDR1) sequence set forth in SEQ ID NO: 50, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 64, SEQ ID NO: 65, or SEQ ID NO: 66, a VH CDR2 sequence set forth in SEQ ID NO: 51, SEQ ID NO: 72, SEQ ID NO: 73, SEQ ID NO: 74, SEQ ID NO: 75, or SEQ ID NO: 76, a VH CDR3 sequence set forth in SEQ ID NO: 52 or SEQ ID NO: 77, a VL CDR1 sequence set forth in SEQ ID NO: 53, SEQ ID NO: 67, or SEQ ID NO: 68, a VL CDR2 sequence set forth in SEQ ID NO: 54, SEQ ID NO: 69, SEQ ID NO: 70, or SEQ ID NO: 71 and a VL CDR3 sequence set forth in SEQ ID NO: 55. For example, the antibody or immunologically active fragment thereof comprises a VH CDR1 sequence set forth in SEQ ID NO: 50, a VH CDR2 sequence set forth in SEQ ID NO: 51, a VH CDR3 sequence set forth in SEQ ID NO: 52, a VL CDR1 sequence set forth in

2013217114 21 Feb 2017

SEQ ID NO: 53, a VL CDR2 sequence set forth in SEQ ID NO: 54, and a VL CDR3 sequence set forth in SEQ ID NO: 55. In another example, the antibody or immunologically active fragment thereof comprises a VH CDR1 sequence set

forth in SEQ ID NO: 50, a VH CDR2 sequence set forth in SEQ ID NO: 72, a VH CDR3 sequence set forth in SEQ ID NO: 52, a VL CDR1 set forth in SEQ ID NO: 53, a VL CDR2 sequence set forth in SEQ ID NO: 71, and a VL CDR3 sequence set forth in SEQ ID NO: 55.

[0031] In one embodiment, the antibodies of the present invention bind to CD47 in a head to side orientation that positions the heavy chain near the membrane of CD47 expressing cell, while the light chain occludes the SIRP α binding site on CD47. In another embodiment, the antibodies of the present invention bind to CD47 in a head to side orientation that positions the light chain near the membrane of CD47 expressing cell, while the heavy chain occludes the SIRP α binding site on CD47.

[0032] The CD47 antibodies bind to an epitope that includes any one of amino acid residues 1-116 of CD47 when numbered in accordance with SEQ ID NO: 147 (*i.e.*, SEQ ID NO: 48 excluding the signal sequence (amino acids 1-18)). For example, the antibodies of the present invention bind to an epitope that includes one or more of amino acid residues Q31, N32, T33, T34, E35, V36, Y37, V38, K39, W40, K41, F42, K43, G44, R45, D46, I47, Y48, T49, F50, D51, G52, A53, L54, N55, K56, S57, T58, V59, P60, T61, D62, F63, S64, S65, A66, K67, I68, E69, V70, S71, Q72, L73, L74, K75, G76, D77, A78, S79, L80, K81, M82, D83, K84, S85, D86, A87, V88, S89, H90, T91, G92, N93, Y94, T95, C96, E97, V98, T99, E100, L101, T102, R103, E104, G105, E106, T107, I108, I109, and E110 of CD47 when numbered in accordance with SEQ ID NO: 147.

[0033] In some cases, the antibodies of the present invention bind to a discontinuous epitope that includes one or more of amino acid residues Y37, V38, K39, W40, K41, F42, K43, G44, R45, D46, I47, Y48, T49, F50, and D51 of CD47 when numbered in accordance with SEQ ID NO: 147. For example, the antibodies of the present invention bind to a discontinuous epitope comprising amino acids residues Y37, K39, K41, K43, G44, R45, D46, D51, H90, N93, E97, T99, E104, or E106 of CD47 when numbered in accordance with SEQ ID NO: 147. For example, the antibodies of the present invention bind to a discontinuous epitope that includes at least residues of the KGRD (SEQ ID NO: 56) loop (residues 43-46) of CD47 when numbered in accordance with SEQ ID NO: 147. For example, the antibodies of the present invention bind to a discontinuous epitope that includes at least residues Y37, K39, K41, the KGRD (SEQ ID NO: 56) loop (residues 43-46), D51, H90, N93, E97, T99, E104, and E106 of CD47 when numbered in accordance with SEQ ID NO: 147. For example, the antibodies of the present invention bind to a discontinuous epitope that includes residues Y37, K39, K41, the KGRD (SEQ ID NO: 56)

loop (residues 43-46), D51, H90, N93, E97, T99, E104, and E106 of CD47 when numbered in accordance with SEQ ID NO: 147.

[0034] The VH region of the CD47 antibodies described herein is primarily involved in binding to the KGRD (SEQ ID NO: 56) loop of CD47. Thus, the unique epitope to which antibodies of the present invention bind is on the side of CD47. In contrast to existing CD47 antibodies known in the art, the orientation of the VH domain of the CD47 antibodies described herein in a membrane proximal position is a critical feature of these antibodies that prevents cellular clumping, *e.g.*, red blood cell hemagglutination, by constraining the antibodies such that they cannot bridge to CD47 molecules on adjacent cells. Additionally, because the VK domain of the CD47 antibodies described herein interacts with apical residues such as Y37, T102, and E104, which are involved in SIRP α binding, it is primarily the VK domain that physically precludes SIRP α binding to CD47.

[0035] Also provided is an isolated antibody or an immunologically active fragment thereof which competes with the CD47 antibodies described herein for preventing CD47 from interacting with SIRP α .

[0036] The invention provides a polypeptide comprising amino acids residues Y37, K39, K41, K43, G44, R45, D46, D51, H90, N93, E97, T99, E104, and E106 of CD47 when numbered in accordance with SEQ ID NO: 147. Also provided is a polypeptide comprising any one of amino acid residues 1-116 of CD47 when numbered in accordance with SEQ ID NO: 147. For example, the polypeptide comprises one or more of amino acid residues Q31, N32, T33, T34, E35, V36, Y37, V38, K39, W40, K41, F42, K43, G44, R45, D46, I47, Y48, T49, F50, D51, G52, A53, L54, N55, K56, S57, T58, V59, P60, T61, D62, F63, S64, S65, A66, K67, I68, E69, V70, S71, Q72, L73, L74, K75, G76, D77, A78, S79, L80, K81, M82, D83, K84, S85, D86, A87, V88, S89, H90, T91, G92, N93, Y94, T95, C96, E97, V98, T99, E100, L101, T102, R103, E104, G105, E106, T107, I108, I109, and E110 of CD47 when numbered in accordance with SEQ ID NO: 147. Also provided are methods of using this polypeptide as an antigen, *e.g.*, an antigen which binds a CD47 antibody.

[0037] The invention also provides methods of alleviating a symptom of a cancer or other neoplastic condition by administering to a subject in need thereof one or more monoclonal antibodies that bind to CD47 or an immunologically active fragment thereof, wherein the antibody does not cause a significant level of hemagglutination of red blood cells after administration. The antibody is administered in an amount sufficient to alleviate the symptom of the cancer or other neoplastic condition in the subject. In some embodiments, the subject is a human. In some embodiments, the antibody is chimeric,

humanized, or fully human. In some embodiments, the antibody binds to human CD47. In some embodiments, the antibody or immunologically active fragment thereof prevents CD47 from interacting with SIRP⁺. In some embodiments, the antibody or immunologically active fragment thereof is an IgG isotype selected from the group consisting of IgG1 isotype, IgG2 isotype, IgG3 isotype, and IgG4 isotype. In some embodiments, the antibody or immunologically active fragment thereof is an IgG isotype selected from IgG4P and IgG4PE.

[0038] In some embodiments, the CD47 antibodies described herein are used in conjunction with one or more additional agents or a combination of additional agents. Suitable additional agents include current pharmaceutical and/or surgical therapies for an intended application, such as, for example, cancer. For example, the CD47 antibodies can be used in conjunction with one or more additional chemotherapeutic or anti-neoplastic agents. Alternatively, the additional chemotherapeutic agent is radiotherapy. In some embodiments, the chemotherapeutic agent is a cell death-inducing agent. In some embodiments, the chemotherapeutic agent induces a loss of phospholipid asymmetry across the plasma membrane, for example causes cell surface exposure of phosphatidylserine (PS). In some embodiments, the chemotherapeutic agent induces endoplasmic reticulum (ER) stress. In some embodiments, the chemotherapeutic agent is a proteasome inhibitor. In some embodiments, the chemotherapeutic agent induces the translocation of ER proteins to the cell surface. In some embodiments, the chemotherapeutic agent induces the translocation and cell surface exposure of calreticulin.

[0039] In some embodiments, the CD47 antibody and additional agent are formulated into a single therapeutic composition, and the CD47 antibody and additional agent are administered simultaneously. Alternatively, the CD47 antibody and additional agent are separate from each other, *e.g.*, each is formulated into a separate therapeutic composition, and the CD47 antibody and the additional agent are administered simultaneously, or the CD47 antibody and the additional agent are administered at different times during a treatment regimen. For example, the CD47 antibody is administered prior to the administration of the additional agent, the CD47 antibody is administered subsequent to the administration of the additional agent, or the CD47 antibody and the additional agent are administered in an alternating fashion. As described herein, the CD47 antibody and additional agent are administered in single doses or in multiple doses.

[0040] One skilled in the art will appreciate that the antibodies of the invention have a variety of uses. For example, the antibodies of the invention are used as therapeutic

agents, as reagents in diagnostic kits or as diagnostic tools, or as reagents in competition assays to generate therapeutic reagents.

BRIEF DESCRIPTION OF THE DRAWINGS

[0041] Figure 1A is a graph depicting the binding of CD47 on Daudi cells by antibodies in hybridoma supernatants as assessed by flow cytometry. Figure 1B is a graph showing the ability of some of the CD47 antibodies within the hybridoma supernatant to block the binding of recombinant human SIRP α to recombinant human CD47, as determined by an ELISA.

[0042] Figure 2 is a series of graphs depicting (A) the binding of purified murine CD47 antibodies to Raji cells, a cultured line of lymphoblastoid cells derived from a Burkitt lymphoma, and (B) CCRF-CEM cells, a CD47 positive human T cell lymphoblast-like cell line, as analyzed by flow cytometry. This experiment compares the binding of the murine antibodies of the present invention to the commercially available CD47 antibodies, B6H12 and 2D3.

[0043] Figure 3 is a series of graphs depicting the capacity of CD47 antibodies to block SIRP α binding using (A) an ELISA with recombinant human protein, or (B) by flow cytometry using CCRF-CEM cells and recombinant human SIRP α protein.

[0044] Figure 4 is a series of photographs, graphs, and a table that show RBC hemagglutination by CD47 antibodies. RBC hemagglutination is evidenced by a haze appearance in the well, whereas non-agglutinated RBC appear as punctate. Figure 4A shows that the 2A1 antibody displays no hemagglutination at all concentrations tested. Hemagglutination index is depicted in the graph. Figure 4B shows that 2A1 is rare amongst many CD47 antibodies in its inability to agglutinate RBCs. Also shown is the lack of agglutinating activity by the human chimeric version of 2A1 (2A1-xi). Figure 4C shows that CD47 monoclonal antibody 2D3, which does not block SIRP α , does not cause hemagglutination. Figure 4D shows a high concentration range of the CD47 antibodies in the hemagglutination assay and demonstrates the pro-zone effect. Hemagglutination index is depicted in the graph. Figure 4E demonstrates the more narrow concentration range for hemagglutination by the CD47 antibody, 1B4, whereas this effect is absent upon 2A1 binding. Figure 4F shows that 2A1, chimeric 2A1 (2A1-xi), and humanized variants do not cause hemagglutination. In most experiments the 9E4 antibody and the commercial B6H12 antibody were used as positive controls for hemagglutination. Other commercially

available antibodies used in these assays were the SIRP blocking antibodies, BRC126 and CC2C6, and the non-SIRP blocking antibody, 2D3.

[0045] Figure 5 is a graph showing 2A1 and B6H12 binding to cynomolgus (cyno) monkey B-cells and Raji, as assessed by flow cytometry. 2A1 binds human and cyno CD47 with equivalent affinity, as does B6H12, albeit with lower affinity to both human and cyno CD47 than 2A1.

[0046] Figure 6 is a graph depicting the binding of 2A1, 2A1-xi, and B6H12 to Raji cells, as assessed by flow cytometry. Importantly, the graph shows that the variable region sequences of the heavy (VH) and light (VL) chains were correctly elucidated in the chimeric version of 2A1.

[0047] Figure 7A-7J is a series of graphs showing binding of humanized variants of 2A1 to Raji cells. 2A1-xi was used as an internal control on most graphs. Numerous combinations of heavy and light chains were tested as described in Example 8.

[0048] Figure 8A is image of the trace from size exclusion chromatography using an AKTA FLPC with a superdex200 column. Shown are the IgG1, IgG4P, and IgG4PE variants of the AB6.12 antibody. All three variant are over 97% monomeric. Figure 8B is a photograph of a coomassie blue stained SDS-PAGE gel of numerous humanized variants of 2A1 under reducing (R) and non-reducing (NR) conditions.

[0049] Figure 9 is a series of graphs depicting the ability of CD47 antibodies to promote phagocytosis of human tumor cell lines by human monocyte derived macrophages (MDM). Figure 9A is a graph showing the phagocytic index, wherein antibodies used were the commercial antibody B6H12, the murine 2A1 antibody, a humanized variant AB2.05 antibody, and the non-blocking commercial antibody 2D3. Figure 9B is a graph showing the phagocytic index, wherein the antibodies used were the commercial antibody B6H12, humanized antibody AB2.05 (human IgG1), and the IgG1, IgG4P, and IgG4PE variants of the humanized antibody AB6.12. CCRF-CEM cells were used as the CD47 target cell line in these experiments.

[0050] Figure 10 is a series of graphs showing the anti-tumor effects of CD47 antibodies in a Raji tumor model. Figure 10A is a graph that shows the efficacy of the murine antibodies 9E4, 1B4, 2A1, and the commercial antibody B6H12. Figure 10B is a graph that shows the efficacy of the IgG1, IgG4P, and IgG4PE isotypes of the humanized antibody AB6.12, along with the murine 2A1 antibody. In both models, mice were treated with 200 μ g antibody doses three times a week.

[0051] Figure 11 is a graphic representation of the co-crystal complexes of CD47-IgV with SIRP α -IgV domain (A, (Protein Data Bank (PDB) Reference No. 2JJS), B6H12 (B), and 2A1 (C). 2A1 and B6H12 bind in very different orientations and distinct epitopes on CD47, both of which overlap with the SIRP α binding site. The 2A1 antibody binds in a head to side orientation on the CD47 protein.

DETAILED DESCRIPTION

[0052] The present invention provides monoclonal antibodies that specifically bind CD47, including human CD47. These antibodies are collectively referred to herein as CD47 antibodies.

[0053] CD47, a multi-spanning transmembrane receptor belonging to the immunoglobulin superfamily, interacts with SIRP α (signal-regulatory-protein α) on macrophages and thereby dampens phagocytosis. Cancer cells that co-opt this pathway evade phagocytosis. As described in detail below, this is a new mechanism of tumor immune avoidance, and therapeutically targeting CD47 has widespread application in numerous cancers.

[0054] The expression of CD47 correlates with worse clinical outcomes in many distinct malignancies including Non-Hodgkin Lymphoma (NHL), Acute Lymphocytic Leukemia (ALL), Acute Myelogenous Leukemia (AML), ovarian cancer, glioma, glioblastoma, etc. In addition, CD47 has been identified as a cancer stem cell marker in both leukemias and solid tumors (Jaiswal et al., 2009 Cell, 138(2): 271-85; Chan et al., 2009 Proc Natl Acad Sci USA, 106(33): 14016-21; Chan et al., 2010 Curr Opin Urol, 20(5): 393-7; Majeti R et al., 2011 Oncogene, 30(9): 1009-19).

[0055] CD47 blocking antibodies have demonstrated anti-tumor activity multiple *in vivo* tumor models. Furthermore, these antibodies have been shown to synergize with other therapeutic antibodies including Rituxan \circledR and Herceptin \circledR in tumor models. Blocking the interaction of CD47 with SIRP α is capable of promoting phagocytosis of CD47 expressing cells by macrophages (reviewed in Chao et al., 2012 Curr Opin Immunol, 24(2): 225-32). Mice lacking CD47 are markedly resistant to radiation therapy, suggesting a role for targeting CD47 in combination with radiotherapy (Isenberg et al., 2008 Am J Pathol, 173(4): 1100-1112; Maxhimer et al., 2009 Sci Transl Med, 1(3): 3ra7). Furthermore, syngeneic tumor models in these mice display decreased bone metastasis compared wild-type mice (Uluçkan et al., 2009 Cancer Res, 69(7): 3196-204).

[0056] Importantly, most CD47 antibodies have been reported to cause hemagglutination of human erythrocytes. Hemagglutination is an example of a homotypic interaction, wherein two CD47 expressing cells are caused to aggregate or clump when treated with a bivalent CD47 binding entity. For example, the CD47 antibody, MABL, as a full IgG or F(ab')₂, has been reported to cause hemagglutination of erythrocytes, and, only when MABL was altered into an scFv or bivalent scFv, was this effect mitigated. (See e.g., Uno S, Kinoshita Y, Azuma Y et al. Antitumor activity of a monoclonal antibody against CD47 in xenograft models of human leukemia. *Oncol Rep* 2007; 17: 1189–94; Kikuchi Y, Uno S, Yoshimura Y et al. A bivalent single-chain Fv fragment against CD47 induces apoptosis for leukemic cells. *Biochem Biophys Res Commun* 2004; 315: 912–8). Other known CD47 antibodies including B6H12, BRC126, and CC2C6 also cause hemagglutination of RBCs, as described in detail below. Thus, the aggregation of cells represents a major limitation of therapeutically targeting CD47 with existing full IgG antibodies.

[0057] Moreover, an important characteristic of CD47 antibodies is the ability to block the interaction of CD47 and SIRP α in order to promote the phagocytosis of CD47 expressing cells by macrophages. Many existing CD47 antibodies block SIRP α ; however, prior to the invention described herein, existing antibodies that blocked SIRP α caused the side effect of hemagglutination, which, as described above, is undesirable. Other existing antibodies, such as 2D3, do not cause hemagglutination; however, these antibodies also do not block SIRP α , rendering them ineffective in the promotion of phagocytosis. Thus, prior to the invention described herein, there was a pressing need to identify CD47 antibodies that blocked SIRP α without causing cellular clumping.

[0058] The CD47 antibodies of the present invention avoid the undesirable effect of hemagglutination, thereby increasing the efficacy of therapeutically targeting CD47, and maintain the ability to block the interaction of CD47 with SIRP α , thereby promoting phagocytosis of CD47 expressing cells. Specifically, the full IgG CD47 antibodies of the present invention (e.g., 2A1 and its humanized derivatives including those provided in Table 1) do not agglutinate cells at a significant level. For example, the CD47 antibodies of the invention do not hemagglutinate RBCs at a significant level. Described herein are the first CD47 antibodies in a full IgG format that block SIRP α and do not cause a significant level of hemagglutination. Taken together, the antibodies of the invention (e.g., the 2A1 antibody and its humanized derivatives) are unique among existing CD47 antibodies in their ability to block SIRP α , but not cause a significant level of hemagglutination.

[0059] The CD47 antibodies of the invention exhibit numerous desirable characteristics, such as, by way of non-limiting example, potent blocking of the interaction between CD47 and its ligand SIRP α , without causing a significant level of or otherwise modulating hemagglutination of erythrocytes, as well as potent anti-tumor activity. For example, the CD47 antibodies of the invention block at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 95%, or at least 99% of the interaction between CD47 and SIRP α as compared to the level of interaction between CD47 and SIRP α in the absence of the CD47 antibody described herein. The CD47 antibodies of the invention do not cause a significant level of agglutination of cells, *e.g.*, the CD47 antibodies of the invention do not cause a significant level of hemagglutination of red blood cells. For example, the level of agglutination in the presence of the CD47 antibodies of the invention is reduced by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 99% compared to the level of agglutination in the presence existing CD47 antibodies. In some embodiments, the CD47 antibodies of the invention do not cause a significant level of agglutination if the level of agglutination in the presence of the CD47 antibodies of the invention is reduced by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 99% compared to the level of agglutination in the presence of CD47 antibody, 1B4, which comprises a variable heavy and variable light chain sequence provided in SEQ ID NO: 80 and SEQ ID NO: 81, respectively. The antibodies of the present invention are also significantly more potent in tumor models compared to antibodies known in the art. For example, the ability of macrophages to phagocytose tumor cells in the presence of CD47 antibodies of the invention is increased by at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 99% compared to the ability of macrophages to phagocytose tumor cells in the presence of existing CD47 antibodies.

[0060] Those skilled in the art will recognize that it is possible to quantitate, without undue experimentation, the level of agglutination, *e.g.*, the level of hemagglutination of RBCs. For example, those skilled in the art will recognize that the level of hemagglutination is ascertained by measuring the area of an RBC dot after performing a hemagglutination assay in the presence of the CD47 antibodies of the invention, as described in the Examples below. In some cases, the area of the RBC dot in the presence of the CD47 antibody of the invention is compared to the area of the RBC dot in the absence

of a CD47 antibody, *i.e.*, in the presence of zero hemagglutination. In this manner, hemagglutination is quantified relative to a baseline control. A larger RBC dot area corresponds to a higher level of hemagglutination. Alternatively, densitometry of the RBC dot may also be utilized to quantitate hemagglutination.

[0061] The CD47 antibodies of the invention bind to human CD47 and block its interaction with SIRP α (Figures 1B, 3, and 7J). These antibodies do not cause a significant level of hemagglutination of human erythrocytes (Figure 4). These antibodies are capable of promoting phagocytosis of tumor cells by macrophages (Figure 9). Furthermore, the CD47 antibodies display potent anti-tumor activity in a mouse model of human lymphoma (Figure 10). Thus, the CD47 antibodies of the invention circumvent a major limiting factor for the therapeutic targeting CD47. Accordingly, the CD47 antibodies of the invention stand to be of great importance in treatment a multitude of cancers.

[0062] Antibodies of the invention that specifically bind human CD47, block, inhibit, disrupt or otherwise modulate the interaction between human CD47 and human SIRP α , without causing a significant level of or otherwise modulating hemagglutination of erythrocytes.

[0063] The antibodies of the present invention bind to a CD47 epitope with an equilibrium binding constant (K_d) of $\leq 1 \mu\text{M}$, *e.g.*, $\leq 100 \text{ nM}$, preferably $\leq 10 \text{ nM}$, and more preferably $\leq 1 \text{ nM}$. For example, the CD47 antibodies provided herein exhibit a K_d in the range approximately between $\leq 1 \text{ nM}$ to about 1 pM .

[0064] The CD47 antibodies of the invention serve to modulate, block, inhibit, reduce, antagonize, neutralize or otherwise interfere with the functional activity of the widely distributed CD47. Functional activities of CD47 include for example, signaling via the interaction with SIRP α , modulating, *e.g.*, increasing, intracellular calcium concentration upon cell adhesion to extracellular matrix, interacting with the C-terminal cell binding domain of thrombospondin, interacting with fibrinogen, and interacting with various integrins. For example, the CD47 antibodies completely or partially inhibit CD47 functional activity by partially or completely modulating, blocking, inhibiting, reducing, antagonizing, neutralizing, or otherwise interfering with the binding of CD47 to SIRP α .

[0065] The CD47 antibodies are considered to completely modulate, block, inhibit, reduce, antagonize, neutralize or otherwise interfere with CD47 functional activity when the level of CD47 functional activity in the presence of CD47 antibody is decreased by at least 95%, *e.g.*, by 96%, 97%, 98%, 99% or 100% as compared to the level of CD47 functional

activity in the absence of binding with a CD47 antibody described herein. The CD47 antibodies are considered to significantly block, inhibit, reduce, antagonize, neutralize or otherwise interfere with CD47 functional activity when the level of CD47 activity in the presence of the CD47 antibody is decreased by at least 50%, *e.g.*, 55%, 60%, 75%, 80%, 85% or 90% as compared to the level of CD47 activity in the absence of binding with a CD47 antibody described herein. The CD47 antibodies are considered to partially modulate, block, inhibit, reduce, antagonize, neutralize or otherwise interfere with CD47 functional activity when the level of CD47 activity in the presence of the CD47 antibody is decreased by less than 95%, *e.g.*, 10%, 20%, 25%, 30%, 40%, 50%, 60%, 75%, 80%, 85% or 90% as compared to the level of CD47 activity in the absence of binding with a CD47 antibody described herein.

Definitions

[0066] Unless otherwise defined, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures utilized in connection with, and techniques of, cell and tissue culture, molecular biology, and protein and oligo- or polynucleotide chemistry and hybridization described herein are those well known and commonly used in the art. Standard techniques are used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (*e.g.*, electroporation, lipofection). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and procedures are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. *See e.g.*, Sambrook *et al.* Molecular Cloning: A Laboratory Manual (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). The nomenclatures utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[0067] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[0067a] In the claims which follow and in the description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

[0068] As used herein, the terms CD47, integrin-associated protein (IAP), ovarian cancer antigen OA3, Rh-related antigen and MER6 are synonymous and may be used interchangeably.

[0069] The terms red blood cell(s) and erythrocyte(s) are synonymous and used interchangeably herein.

[0070] The term agglutination refers to cellular clumping, while the term hemagglutination refers to clumping of a specific subset of cells, *i.e.*, red blood cells. Thus, hemagglutination is a type of agglutination.

[0071] As used herein, the term “antibody” refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, *i.e.*, molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. By “specifically bind” or “immunoreacts with” “or directed against” is meant that the antibody reacts with one or more antigenic determinants of the desired antigen and does not react with other polypeptides or binds at much lower affinity ($K_d > 10^{-6}$). Antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, dAb (domain antibody), single chain, Fab, Fab⁻ and F(ab')₂ fragments, F_v, scFvs, and an Fab expression library.

[0072] The basic antibody structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kDa) and one “heavy” chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. In general, antibody molecules obtained from humans relate to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain.

[0073] The term “monoclonal antibody” (MAb) or “monoclonal antibody composition”, as used herein, refers to a population of antibody molecules that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs contain an antigen binding site capable of immunoreacting with a particular epitope of the antigen characterized by a unique binding affinity for it.

[0074] In general, antibody molecules obtained from humans relate to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain.

[0075] The term “antigen-binding site” or “binding portion” refers to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable (“V”) regions of the heavy (“H”) and light (“L”) chains. Three highly divergent stretches within the V regions of the heavy and light chains, referred to as “hypervariable regions,” are interposed between more conserved flanking stretches known as “framework regions,” or “FRs”. Thus, the term “FR” refers to amino acid sequences which are naturally found between, and adjacent to, hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three-dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as “complementarity-determining regions,” or “CDRs.” The assignment of amino acids to each domain is in accordance with the definitions of Kabat Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987 and 1991)), or Chothia & Lesk J. Mol. Biol. 196:901-917 (1987), Chothia *et al.* Nature 342:878-883 (1989).

[0076] As used herein, the term “epitope” includes any protein determinant capable of specific binding to an immunoglobulin or fragment thereof, or a T-cell receptor. The term “epitope” includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. An antibody is said to specifically bind an antigen when the dissociation constant is $\leq 1 \mu\text{M}$; *e.g.*, $\leq 100 \text{ nM}$, preferably $\leq 10 \text{ nM}$ and more preferably $\leq 1 \text{ nM}$.

[0077] As used herein, the terms “immunological binding,” and “immunological binding properties” refer to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The

strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and geometric parameters that equally influence the rate in both directions. Thus, both the “on rate constant” (k_{on}) and the “off rate constant” (k_{off}) can be determined by calculation of the concentrations and the actual rates of association and dissociation. (See *Nature* 361:186-87 (1993)). The ratio of k_{off}/k_{on} enables the cancellation of all parameters not related to affinity, and is equal to the dissociation constant K_d . (See, generally, Davies et al. (1990) *Annual Rev Biochem* 59:439-473). An antibody of the present invention is said to specifically bind to CD47, when the equilibrium binding constant (K_d) is $\leq 1 \mu\text{M}$, preferably $\leq 100 \text{ nM}$, more preferably $\leq 10 \text{ nM}$, and most preferably $\leq 100 \text{ pM}$ to about 1 pM, as measured by assays such as radioligand binding assays, surface plasmon resonance (SPR), flow cytometry binding assay, or similar assays known to those skilled in the art.

[0078] The term “isolated polynucleotide” as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the “isolated polynucleotide” (1) is not associated with all or a portion of a polynucleotide in which the “isolated polynucleotide” is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence.

[0079] The term “isolated protein” referred to herein means a protein of cDNA, recombinant RNA, or synthetic origin or some combination thereof, which by virtue of its origin, or source of derivation, the “isolated protein” (1) is not associated with proteins found in nature, (2) is free of other proteins from the same source, *e.g.*, free of marine proteins, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

[0080] The term “polypeptide” is used herein as a generic term to refer to native protein, fragments, or analogs of a polypeptide sequence. Hence, native protein fragments, and analogs are species of the polypeptide genus.

[0081] The term “naturally-occurring” as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be

isolated from a source in nature and which has not been intentionally modified by man in the laboratory or otherwise is naturally-occurring.

[0082] The term “operably linked” as used herein refers to positions of components so described are in a relationship permitting them to function in their intended manner. A control sequence “operably linked” to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

[0083] The term “control sequence” as used herein refers to polynucleotide sequences which are necessary to effect the expression and processing of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term “control sequences” is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences. The term “polynucleotide,” as referred to herein, refers to a polymeric boron of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

[0084] The term “oligonucleotide” referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. Preferably oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, *e.g.*, for probes, although oligonucleotides may be double stranded, *e.g.*, for use in the construction of a gene mutant. Oligonucleotides of the invention are either sense or antisense oligonucleotides.

[0085] The term “naturally occurring nucleotides” referred to herein includes deoxyribonucleotides and ribonucleotides. The term “modified nucleotides” referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term “oligonucleotide linkages” referred to herein includes Oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselerloate, phosphorodiselenoate, phosphoroanilothioate, phosphoranylilate, phosphoromimidate, and the like. *See e.g.*, LaPlanche *et al.* Nucl. Acids Res. 14:9081 (1986); Stec *et al.* J. Am. Chem. Soc. 106:6077

(1984), Stein *et al.* Nucl. Acids Res. 16:3209 (1988), Zon *et al.* Anti Cancer Drug Design 6:539 (1991); Zon *et al.* Oligonucleotides and Analogues: A Practical Approach, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England (1991)); Stec *et al.* U.S. Patent No. 5,151,510; Uhlmann and Peyman Chemical Reviews 90:543 (1990). An oligonucleotide can include a label for detection, if desired.

[0086] The term “selectively hybridize” referred to herein means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments thereof in accordance with the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. High stringency conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. Generally, the nucleic acid sequence homology between the polynucleotides, oligonucleotides, and fragments of the invention and a nucleic acid sequence of interest will be at least 80%, and more typically with preferably increasing homologies of at least 85%, 90%, 95%, 99%, and 100%. Two amino acid sequences are homologous if there is a partial or complete identity between their sequences. For example, 85% homology means that 85% of the amino acids are identical when the two sequences are aligned for maximum matching. Gaps (in either of the two sequences being matched) are allowed in maximizing matching gap lengths of 5 or less are preferred with 2 or less being more preferred. Alternatively and preferably, two protein sequences (or polypeptide sequences derived from them of at least 30 amino acids in length) are homologous, as this term is used herein, if they have an alignment score of at more than 5 (in standard deviation units) using the program ALIGN with the mutation data matrix and a gap penalty of 6 or greater. *See* Dayhoff, M.O., in *Atlas of Protein Sequence and Structure*, pp. 101-110 (Volume 5, National Biomedical Research Foundation (1972)) and Supplement 2 to this volume, pp. 1-10. The two sequences or parts thereof are more preferably homologous if their amino acids are greater than or equal to 50% identical when optimally aligned using the ALIGN program. The term “corresponds to” is used herein to mean that a polynucleotide sequence is homologous (*i.e.*, is identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide sequence, or that a polypeptide sequence is identical to a reference polypeptide sequence. In contradistinction, the term “complementary to” is used herein to mean that the complementary sequence is homologous to all or a portion of a reference polynucleotide sequence. For illustration, the nucleotide sequence “TATAC” corresponds to a reference sequence “TATAC” and is complementary to a reference sequence “GTATA”.

[0087] The following terms are used to describe the sequence relationships between two or more polynucleotide or amino acid sequences: “reference sequence”, “comparison window”, “sequence identity”, “percentage of sequence identity”, and “substantial identity”. A “reference sequence” is a defined sequence used as a basis for a sequence comparison a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing or may comprise a complete cDNA or gene sequence. Generally, a reference sequence is at least 18 nucleotides or 6 amino acids in length, frequently at least 24 nucleotides or 8 amino acids in length, and often at least 48 nucleotides or 16 amino acids in length. Since two polynucleotides or amino acid sequences may each (1) comprise a sequence (*i.e.*, a portion of the complete polynucleotide or amino acid sequence) that is similar between the two molecules, and (2) may further comprise a sequence that is divergent between the two polynucleotides or amino acid sequences, sequence comparisons between two (or more) molecules are typically performed by comparing sequences of the two molecules over a “comparison window” to identify and compare local regions of sequence similarity. A “comparison window”, as used herein, refers to a conceptual segment of at least 18 contiguous nucleotide positions or 6 amino acids wherein a polynucleotide sequence or amino acid sequence may be compared to a reference sequence of at least 18 contiguous nucleotides or 6 amino acid sequences and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions, deletions, substitutions, and the like (*i.e.*, gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, (Genetics Computer Group, 575 Science Dr., Madison, Wis.), Geneworks, or MacVector software packages), or by inspection, and the best alignment (*i.e.*, resulting in the highest percentage of homology over the comparison window) generated by the various methods is selected.

[0088] The term “sequence identity” means that two polynucleotide or amino acid sequences are identical (*i.e.*, on a nucleotide-by-nucleotide or residue-by-residue basis) over the comparison window. The term “percentage of sequence identity” is calculated by

comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (*e.g.*, A, T, C, G, U or I) or residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the comparison window (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms “substantial identity” as used herein denotes a characteristic of a polynucleotide or amino acid sequence, wherein the polynucleotide or amino acid comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 18 nucleotide (6 amino acid) positions, frequently over a window of at least 24-48 nucleotide (8-16 amino acid) positions, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the comparison window. The reference sequence may be a subset of a larger sequence.

[0089] As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. *See Immunology - A Synthesis* (2nd Edition, E.S. Golub and D.R. Gren, Eds., Sinauer Associates, Sunderland7 Mass. (1991)). Stereoisomers (*e.g.*, D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4 hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N- acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, σ -N-methylarginine, and other similar amino acids and imino acids (*e.g.*, 4- hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

[0090] Similarly, unless specified otherwise, the left-hand end of single- stranded polynucleotide sequences is the 5' end the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction sequence regions on the DNA strand having the same sequence as the RNA and which are 5' to the 5' end of the RNA transcript are referred to as “upstream sequences”, sequence regions on the DNA

strand having the same sequence as the RNA and which are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences".

[0091] As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, preferably at least 90 percent sequence identity, more preferably at least 95 percent sequence identity, and most preferably at least 99 percent sequence identity.

[0092] Preferably, residue positions which are not identical differ by conservative amino acid substitutions.

[0093] Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide- containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur- containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine valine, glutamic- aspartic, and asparagine-glutamine.

[0094] As discussed herein, minor variations in the amino acid sequences of antibodies or immunoglobulin molecules are contemplated as being encompassed by the present invention, providing that the variations in the amino acid sequence maintain at least 75%, more preferably at least 80%, 90%, 95%, and most preferably 99%. In particular, conservative amino acid replacements are contemplated. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. Genetically encoded amino acids are generally divided into families: (1) acidic amino acids are aspartate, glutamate; (2) basic amino acids are lysine, arginine, histidine; (3) non-polar amino acids are alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan, and (4) uncharged polar amino acids are glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. The hydrophilic amino acids include arginine, asparagine, aspartate, glutamine, glutamate, histidine, lysine, serine, and threonine. The hydrophobic amino acids include alanine, cysteine, isoleucine, leucine, methionine, phenylalanine, proline, tryptophan, tyrosine and valine. Other families of amino acids include (i) serine and threonine, which are the aliphatic-hydroxy family; (ii) asparagine and

glutamine, which are the amide containing family; (iii) alanine, valine, leucine and isoleucine, which are the aliphatic family; and (iv) phenylalanine, tryptophan, and tyrosine, which are the aromatic family. For example, it is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the binding or properties of the resulting molecule, especially if the replacement does not involve an amino acid within a framework site. Whether an amino acid change results in a functional peptide can readily be determined by assaying the specific activity of the polypeptide derivative. Assays are described in detail herein. Fragments or analogs of antibodies or immunoglobulin molecules can be readily prepared by those of ordinary skill in the art. Preferred amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains. Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. Preferably, computerized comparison methods are used to identify sequence motifs or predicted protein conformation domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. Bowie *et al.* Science 253:164 (1991). Thus, the foregoing examples demonstrate that those of skill in the art can recognize sequence motifs and structural conformations that may be used to define structural and functional domains in accordance with the invention.

[0095] Preferred amino acid substitutions are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and (4) confer or modify other physicochemical or functional properties of such analogs. Analogs can include various mutants of a sequence other than the naturally-occurring peptide sequence. For example, single or multiple amino acid substitutions (preferably conservative amino acid substitutions) may be made in the naturally- occurring sequence (preferably in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. A conservative amino acid substitution should not substantially change the structural characteristics of the parent sequence (*e.g.*, a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in Proteins, Structures and Molecular Principles (Creighton, Ed., W. H.

Freeman and Company, New York (1984)); Introduction to Protein Structure (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al. *Nature* 354:105 (1991).

[0096] The term “polypeptide fragment” as used herein refers to a polypeptide that has an amino terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the naturally-occurring sequence deduced, for example, from a full length cDNA sequence. Fragments typically are at least 5, 6, 8 or 10 amino acids long, preferably at least 14 amino acids long, more preferably at least 20 amino acids long, usually at least 50 amino acids long, and even more preferably at least 70 amino acids long. The term “analog” as used herein refers to polypeptides which are comprised of a segment of at least 25 amino acids that has substantial identity to a portion of a deduced amino acid sequence and which has specific binding to CD47, under suitable binding conditions. Typically, polypeptide analogs comprise a conservative amino acid substitution (or addition or deletion) with respect to the naturally- occurring sequence. Analogs typically are at least 20 amino acids long, preferably at least 50 amino acids long or longer, and can often be as long as a full-length naturally-occurring polypeptide.

[0097] Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed “peptide mimetics” or “peptidomimetics”. Fauchere, J. *Adv. Drug Res.* 15:29 (1986), Veber and Freidinger *TINS* p.392 (1985); and Evans *et al.* *J. Med. Chem.* 30:1229 (1987). Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (*i.e.*, a polypeptide that has a biochemical property or pharmacological activity), such as human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from the group consisting of: --CH₂NH--, --CH₂S--, --CH₂CH₂--, --CH=CH--(cis and trans), --COCH₂--, CH(OH)CH₂--, and -CH₂SO--, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (*e.g.*, D-lysine in place of L-lysine) may be used to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Giersch *Ann. Rev. Biochem.* 61:387

(1992)); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

[0098] The term “agent” is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials.

[0099] As used herein, the terms “label” or “labeled” refers to incorporation of a detectable marker, *e.g.*, by incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (*e.g.*, streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods). In certain situations, the label or marker can also be therapeutic. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (*e.g.*, ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (*e.g.*, FITC, rhodamine, lanthanide phosphors), enzymatic labels (*e.g.*, horseradish peroxidase, p-galactosidase, luciferase, alkaline phosphatase), chemiluminescent, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (*e.g.*, leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance. The term “pharmaceutical agent or drug” as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient.

[00100] The term “antineoplastic agent” is used herein to refer to agents that have the functional property of inhibiting a development or progression of a neoplasm in a human, particularly a malignant (cancerous) lesion, such as a carcinoma, sarcoma, lymphoma, or leukemia. Inhibition of metastasis is frequently a property of antineoplastic agents.

[00101] Other chemistry terms herein are used according to conventional usage in the art, as exemplified by The McGraw-Hill Dictionary of Chemical Terms (Parker, S., Ed., McGraw-Hill, San Francisco (1985)).

[00102] As used herein, “substantially pure” means an object species is the predominant species present (*i.e.*, on a molar basis it is more abundant than any other individual species in the composition), and preferably a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present.

[00103] Generally, a substantially pure composition will comprise more than about 80 percent of all macromolecular species present in the composition, more preferably more than about 85%, 90%, 95%, and 99%. Most preferably, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

CD47 Antibodies

[00104] Monoclonal antibodies of the invention have the ability to bind CD47, to inhibit the binding of SIRP α to CD47, decrease CD47-SIRP α -mediated signaling, promote phagocytosis, and to inhibit tumor growth and/or migration. Inhibition is determined, for example, using the cellular assay described herein in the Examples.

[00105] Exemplary antibodies of the invention include the 2A1 antibody, the chimeric version of 2A1, and humanized variants of 2A1. Exemplary antibodies of the invention include an antibody having a variable heavy (VH) chain selected from SEQ ID NOS: 5-30, and having a variable light (VL) chain selected from SEQ ID NOS: 31-47. Specifically, exemplary antibodies include those provided in Table 1.

Table 1.

Antibody	Variable heavy (VH) chain	Variable light (VL) chain
2A1	SEQ ID NO: 5	SEQ ID NO: 31
2A1-xi	SEQ ID NO: 5	SEQ ID NO: 32
AB2.03	SEQ ID NO: 7	SEQ ID NO: 33
AB2.04	SEQ ID NO: 7	SEQ ID NO: 34
AB2.05	SEQ ID NO: 7	SEQ ID NO: 35
AB2.06	SEQ ID NO: 7	SEQ ID NO: 36
AB2.07	SEQ ID NO: 7	SEQ ID NO: 37
AB2.08	SEQ ID NO: 7	SEQ ID NO: 38
AB2.09	SEQ ID NO: 7	SEQ ID NO: 39
AB2.13	SEQ ID NO: 7	SEQ ID NO: 43
AB3.09	SEQ ID NO: 8	SEQ ID NO: 39
AB6.12	SEQ ID NO: 11	SEQ ID NO: 42
AB6.13	SEQ ID NO: 11	SEQ ID NO: 43
AB6.14	SEQ ID NO: 11	SEQ ID NO: 44

AB6.17	SEQ ID NO: 11	SEQ ID NO: 47
AB10.13	SEQ ID NO: 15	SEQ ID NO: 43
AB10.14	SEQ ID NO: 15	SEQ ID NO: 44
AB11.05	SEQ ID NO: 16	SEQ ID NO: 35
AB12.05	SEQ ID NO: 17	SEQ ID NO: 35
AB15.05	SEQ ID NO: 20	SEQ ID NO: 35
AB16.05	SEQ ID NO: 21	SEQ ID NO: 35
AB17.05	SEQ ID NO: 22	SEQ ID NO: 35
AB22.05	SEQ ID NO: 27	SEQ ID NO: 35
AB23.05	SEQ ID NO: 28	SEQ ID NO: 35
AB24.05	SEQ ID NO: 29	SEQ ID NO: 35
AB25.05	SEQ ID NO: 30	SEQ ID NO: 35

[00106] Also included in the invention are antibodies that bind to the same epitope as the CD47 antibodies described herein. For example, antibodies of the invention specifically bind to an epitope that includes one or more amino acid residues on human CD47 (*see e.g.*, GenBank Accession No. Q08722.1).

[00107] The amino acid sequence of an exemplary human CD47 is provided below (GenBank Accession No. Q08722.1 (GI:1171879), incorporated herein by reference). The signal sequence (amino acids 1-18) is underlined.

```

1 mwplvaalll gsaccgsaql lfnktksvef tfcnndtvvip cfvtnmeaqn
ttevyvkwkf
       61 kgrdiytfdg alnkstvptd fssakievsq llkgdaslkm dksdavshtg
nytcevtelt
       121 regetiielk yrvvswfspd enilivifpi faillfwgqf giktlkyrsg
gmdektiall
       181 vaglvitviv ivgailfvpg eyslknatgl glivtstgil illhyyvfst
aigltsfvia
       241 ilviqviayi lavvglslci aacipmhgpl lisglslal aqlglvymk
fvasnqktiq
       301 pprkaveepl nafkeskgmm nde (SEQ ID NO: 48)

```

[00108] For clarity, the amino acid sequence of an exemplary human CD47 excluding the signal sequence is provided below.

```

1 qllfnktksv eftfcndtvv ipcfvtnmea qnttevyvkw kfkgrdiytf
dgalnkstvp
       61 tdfssakiev sql1kgdasl kmdksdavsh tgnytcevte ltregetiie
lkyrvvswfs
       121 pnenilivif pifaillfwg qfgiktlkyr sggmdektia llvaglvitv
ivivgailfv
       181 pgeyslknat glglivtstg ilillhyyvf staigltsfv iailviqvia
yilavvglsl

```

241 ciaacipmhg pllisglsil alaqlglv y mkfvasnqkt iqpprkavee
 plnafkeskg
 301 mmnde (SEQ ID NO: 147)

[00109] The amino acid sequence of an exemplary human CD47-IgV domain is provided below:

19 qlfnktksv eftfcndtvv ipcfvtnmea qnttevyvkw kfkgrdiytf
 dgalnkstvp
 79 tdfssakiev sqllkgdasl kmdksdavsh tgnytcevte ltregetiie lkyrvv
 (SEQ ID NO: 49)

[00110] Exemplary monoclonal antibodies of the invention include, for example, humanized antibodies having a variable heavy chain region (VH) and/or variable light (VL) chain region shown in the sequences below.

[00111] Variable heavy (VH) chain regions of the CD47 antibodies are provided below. The complementarity determining regions (CDRs) of the VH chain of the CD47 antibodies are highlighted below. In some embodiments, the amino acid sequence of VH CDR1 is GFNIKDYYLH (SEQ ID NO: 50), GYTFTYYYLH (SEQ ID NO: 57), GFTFTYYYLH (SEQ ID NO: 58), GYNFTYYYLH (SEQ ID NO: 59), GYTITYYYYLH (SEQ ID NO: 60), GYTFKYYYLH (SEQ ID NO: 61), GYTFTDYYLH (SEQ ID NO: 62), GFTFTDYYLH (SEQ ID NO: 63), GFTITDYYLH (SEQ ID NO: 64), GYTFKDYYLH (SEQ ID NO: 65), or GTFKDYYLH (SEQ ID NO: 66). In some embodiments, the amino acid sequence of VH CDR2 is WIDPDNGDTE (SEQ ID NO: 51), WIDPDQGDTE (SEQ ID NO: 72), WIDPDYGDTE (SEQ ID NO: 73), WIDPDSGDTE (SEQ ID NO: 74), WIDPDNADTE (SEQ ID NO: 75), or WIDPDNTDTE (SEQ ID NO: 76). In some embodiments, the amino acid sequence of VH CDR3 is NAAYGSSSYPPMDY (SEQ ID NO: 52) or NAAYGSSPYPPMDY (SEQ ID NO: 77).

EVQLQQSGAELVRSRSGASVQLSCTASGFNIKDYYLHWVKQRPEQGLEWIGWIDPDNGDTEFAPKFQGKATMTADT
 SSNTAYLQLSSLTSEDTAVYYCNAAYGSSSYPPMDYWGQGTSVTV (SEQ ID NO: 5)

EVQLVQSGAEVKKPGATVKISCKVSGFNIKDYYLHWVQQAPGKGLEWMGWIDPDNGDTEYAEKFQGRVTITADT
 STDTAYMELSSLRSEDTAVYYCNAAYGSSSYPPMDYWGQGTTVTV (SEQ ID NO: 6)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGWIDPDNGDTEYAQKFQDRVTITRDR
 SMSTAYMELSSLRSEDTAMYYCNAAYGSSSYPPMDYWGQGTTVTV (SEQ ID NO: 7)

EVQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVQRQAPGQALEWMGWIDPDNGDTEYAQKFQDRVTITRDR
 SMSTAYMELSSLRSEDTAMYYCNAAYGSSSYPPMDYWGQGTTVTV (SEQ ID NO: 8)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGWIDPDNGDTEYAQKFQGRVTMTADT
 SSNTAYMELSSLRSEDTAMYYCNAAYGSSSYPPMDYWGQGTTVTV (SEQ ID NO: 9)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQGRVTMTEDT
STDTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 10)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDQGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 11)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDYGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 12)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDSGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 13)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDNADTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 14)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDNTDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 15)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSPYPMDYWGQGTTVTV (SEQ ID NO: 16)

QMQLVQSGAEVKKTGSSVKVSCKASGYTFTYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 17)

QMQLVQSGAEVKKTGSSVKVSCKASGFTFTYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 18)

QMQLVQSGAEVKKTGSSVKVSCKASGYNFYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 19)

QMQLVQSGAEVKKTGSSVKVSCKASGYTITYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 20)

QMQLVQSGAEVKKTGSSVKVSCKASGYTFKYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 21)

QMQLVQSGAEVKKTGSSVKVSCKASGYTFDYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 22)

QMQLVQSGAEVKKTGSSVKVSCKASGFTFTDYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 23)

QMQLVQSGAEVKKTGSSVKVSCKASGFTITDYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 24)

QMQLVQSGAEVKKTGSSVKVSCKASGYTFKDYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 25)

QMQLVQSGAEVKKTGSSVKVSCKASGFTFKDYYYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 26)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYLQLSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 27)

QMQLVQSGAEVKKTGSSVKVSCKASGFNIKDYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLTSEDTAVYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 28)

EVQLVQSGAEVKKPGATVKISCKVSGFNIKDYYLHWVRQAPGQALEWMGVIDPDNGDTEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYYCNAAYGSSSSYPMDYWGQGTTVTV (SEQ ID NO: 29)

EVQLVQSGAEVKKPGATVKISCKVSGFNIKDYYLHWVQQAPGKGLEWMGWIDPDNGDEYAQKFQDRVTITRDR
SMSTAYMELSSLRSEDTAMYCNAAYGSSSYPMDWGQGTTVTV (SEQ ID NO: 30)

[00112] Variable light (VL) chain regions of the CD47 antibodies are provided below. The CDRs of the VL chain of the CD47 antibodies are highlighted below. In some embodiments, the amino acid sequence of VL CDR1 is KASQDIHYRQLS (SEQ ID NO: 53), RASQDIHYRQLA (SEQ ID NO: 67), or RARQGIHYRQLS (SEQ ID NO: 68). In some embodiments, the amino acid sequence of VL CDR2 is RANRLVD (SEQ ID NO: 54), RANRLQS (SEQ ID NO: 69), RANRRAT (SEQ ID NO: 70), or RANRLVS (SEQ ID NO: 71). In some embodiments, the amino acid sequence of VL CDR3 is LQYDEFPYT (SEQ ID NO: 55).

DIKMTQSPSSLYASLGERVTITCKASQDIHYRQLSWFQQKPGKSPKILLYRANRLVDGVPSRFSGSGSGQDYSLT
ISSLEYEDMGIYYCLQYDEFPYTFGGGTKLEMK (SEQ ID NO: 31)

DIKMTQSPSSLYASLGERVTITCKASQDIHYRQLSWFQQKPGKSPKILLYRANRLVDGVPSRFSGSGSGQDYSLT
ISSLEYEDMGIYYCLQYDEFPYTFGGGTKLEIK (SEQ ID NO: 32)

DIQMTQSPSSLSASVGDRVTITCKASQDIHYRQLSWYQQKPGKAPKLLIYRANRLVDGVPSRFSGSGSGTDFTFT
ISSLQPEDIATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 33)

DIQMTQSPSSLSASVGDRVTITCKASQDIHYRQLSWFQQKPGKAPKSLIYRANRLVDGVPSRFSGSGSGTDFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 34)

NIQMTQSPSAMSASVGDRVTITCKASQDIHYRQLWFQQKPGKVKHLLIYRANRLVDGVPSRFSGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 35)

DIQMTQSPSSLSASVGDRVTITCKASQDIHYRQLSWYQQKPGKAPKRLIYRANRLVDGVPSRFSGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 36)

DIQMTQSPSSLSASVGDRVTITCRASQDIHYRQLAWYQQKPGKVKLLIYRANRLQSGVPSRFSGSGSGTDFTLT
ISSLQPEDVATYYCLQYDEFPYTFGQGKVEIK (SEQ ID NO: 37)

EIVLTQSPATLSLSPGERATLSCRASQDIHYRQLAWYQQKPGQAPRLLIYRANRRATGIPARFSGSGSGTDFTLT
ISSLQPEDFAVYYCLQYDEFPYTGFQGTRLEIK (SEQ ID NO: 38)

DIQMTQSPSAMSASVGDRVTITCKASQDIHYRQLSWFQQKPGKVKHLLIYRANRLVDGVPSRFSGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 39)

NIQMTQSPSAMSASVGDRVTITCRARQGIHYRQLWFQQKPGKVKHLLIYRANRLVDGVPSRFSGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 40)

NIQMTQSPSAMSASVGDRVTITCKASQDIHYRQLSWFQQKPGKVKLLIYRANRLVDGVPSRFSGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 41)

NIQMTQSPSAMSASVGDRVTITCKASQDIHYRQLSWFQQKPGKVKHLLIYRANRLVSGVPSRFSGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 42)

NIQMTQSPSAMSASVGDRVTITCRARQGIHYRQLWFQQKPGKVKLLIYRANRLVDGVPSRFSGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 43)

NIQMTQSPSAMSASVGDRVTITCRARQGIHYRQLSWFQQKPGKVKHLLIYRANRLVSGVPSRFSGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 44)

NIQMTQSPSAMSASVGDRVTITCKASQDIHRYLSWFQQKPGKVPKLLIYRANRLVDGVPSRFGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 45)

NIQMTQSPSAMSASVGDRVTITCKASQDIHRYLSWFQQKPGKVPKLLIYRANRLVSGVPSRFGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 46)

NIQMTQSPSAMSASVGDRVTITCRARQGITHRYLSWFQQKPGKVPKLLIYRANRLVSGVPSRFGSGSGTEFTLT
ISSLQPEDFATYYCLQYDEFPYTFGGGTKVEIK (SEQ ID NO: 47)

[00113] In some cases, the CD47 antibodies described herein comprise a variable heavy chain region selected from SEQ ID NOs: 5-30 and a variable light chain region selected from SEQ ID NOs: 31-47. An exemplary CD47 antibody comprises a variable heavy chain region set forth in SEQ ID NO: 5 and a variable light chain region set forth in SEQ ID NO: 31; a variable heavy chain region set forth in SEQ ID NO: 7 and a variable light chain region set forth in SEQ ID NO: 35; a variable heavy chain region set forth in SEQ ID NO: 11 and a variable light chain region set forth in SEQ ID NO: 42, a variable heavy chain region set forth in SEQ ID NO: 5 and a variable light chain region set forth in SEQ ID NO: 32, a variable heavy chain region set forth in SEQ ID NO: 7 and a variable light chain region set forth in SEQ ID NO: 33, a variable heavy chain region set forth in SEQ ID NO: 7 and a variable light chain region set forth in SEQ ID NO: 34, a variable heavy chain region set forth in SEQ ID NO: 7 and a variable light chain region set forth in SEQ ID NO: 36, a variable heavy chain region set forth in SEQ ID NO: 7 and a variable light chain region set forth in SEQ ID NO: 37, a variable heavy chain region set forth in SEQ ID NO: 7 and a variable light chain region set forth in SEQ ID NO: 38, a variable heavy chain region set forth in SEQ ID NO: 29 and a variable light chain region set forth in SEQ ID NO: 35, a variable heavy chain region set forth in SEQ ID NO: 30 and a variable light chain region set forth in SEQ ID NO: 35, a variable heavy chain region set forth in SEQ ID NO: 7 and a variable light chain region set forth in SEQ ID NO: 43, a variable heavy chain region set forth in SEQ ID NO: 11 and a variable light chain region set forth in SEQ ID NO: 43, a variable heavy chain region set forth in SEQ ID NO: 11 and a variable light chain region set forth in SEQ ID NO: 47, a variable heavy chain region set forth in SEQ ID NO: 43, a variable heavy chain region set forth in SEQ ID NO: 15 and a variable light chain region set forth in SEQ ID NO: 43, a variable heavy chain region set forth in SEQ ID NO: 15 and a variable light chain region set forth in SEQ ID NO: 44, a variable heavy chain region set forth in SEQ ID NO: 11 and a variable light chain region set forth in SEQ ID NO: 44, a variable heavy chain region set forth in SEQ ID NO: 22 and a variable light chain region set forth in SEQ ID NO: 35, a variable heavy chain region set forth in SEQ ID NO: 7 and a variable light chain region set forth in

SEQ ID NO: 39, a variable heavy chain region set forth in SEQ ID NO: 8 and a variable light chain region set forth in SEQ ID NO: 39, a variable heavy chain region set forth in SEQ ID NO: 16 and a variable light chain region set forth in SEQ ID NO: 35, a variable heavy chain region set forth in SEQ ID NO: 20 and a variable light chain region set forth in SEQ ID NO: 35, a variable heavy chain region set forth in SEQ ID NO: 21 and a variable light chain region set forth in SEQ ID NO: 35, a variable heavy chain region set forth in SEQ ID NO: 17 and a variable light chain region set forth in SEQ ID NO: 35, a variable heavy chain region set forth in SEQ ID NO: 28 and a variable light chain region set forth in SEQ ID NO: 35, or a variable heavy chain region set forth in SEQ ID NO: 27 and a variable light chain region set forth in SEQ ID NO: 35.

[00114] The CD47 antibodies described herein comprise any one of the VH regions provided in SEQ ID NOS: 5-30 paired with any one of the VL regions provided in SEQ ID NOS: 31-47. Specifically, the CD47 antibodies described herein comprise any one of the VH regions provided in SEQ ID NOS: 5, 7, 8, 11, 15-17, 20-22, and 27-30 paired with any one of the VL regions provided in SEQ ID NOS: 31-39, 42, 43, 44, and 47.

[00115] The CD47 antibodies described herein comprise any one of the VH CDR1 regions provided in SEQ ID NO: 50, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 64, SEQ ID NO: 65, and SEQ ID NO: 66, any one of the VH CDR2 regions provided in SEQ ID NO: 51, SEQ ID NO: 72, SEQ ID NO: 73, SEQ ID NO: 74, SEQ ID NO: 75, and SEQ ID NO: 76, any one of the VH CDR3 regions provided in SEQ ID NO: 52 and SEQ ID NO: 77, any one of the VL CDR1 regions provided in SEQ ID NO: 53, SEQ ID NO: 67, and SEQ ID NO: 68, any one of the VL CDR2 regions provided in SEQ ID NO: 54, SEQ ID NO: 69, SEQ ID NO: 70, and SEQ ID NO: 71, and the VL CDR3 region provided in SEQ ID NO: 55.

[00116] Those skilled in the art will recognize that it is possible to determine, without undue experimentation, if a monoclonal antibody has the same specificity as a monoclonal antibody of the invention (*e.g.*, the 2A1 antibody, or an antibody having a variable heavy chain selected from SEQ ID NOS: 5-31, and a variable light chain selected from SEQ ID NOS: 31-47) by ascertaining whether the former prevents the latter from binding to CD47. If the monoclonal antibody being tested competes with the monoclonal antibody of the invention, as shown by a decrease in binding by the monoclonal antibody of the invention, then the two monoclonal antibodies bind to the same, or a closely related, epitope.

[00117] An alternative method for determining whether a monoclonal antibody has the specificity of monoclonal antibody of the invention is to pre-incubate the monoclonal antibody of the invention with soluble CD47 protein (with which it is normally reactive), and then add the monoclonal antibody being tested to determine if the monoclonal antibody being tested is inhibited in its ability to bind CD47. If the monoclonal antibody being tested is inhibited then, in all likelihood, it has the same, or functionally equivalent, epitopic specificity as the monoclonal antibody of the invention.

Antibodies of the Present Invention

[00118] Screening of monoclonal antibodies of the invention, can be also carried out, *e.g.*, by measuring CD47- and/or CD47/SIRP α -mediated signaling, and determining whether the test monoclonal antibody is able to modulate, block, inhibit, reduce, antagonize, neutralize or otherwise interfere with CD47- and/or CD47/SIRP α -mediated signaling. These assays can include competitive binding assays. Additionally, these assays can measure a biologic readout, for example the ability to promote phagocytosis of a CD47 expressing cell by a macrophage, as is described in Example 9 (Figure 9).

[00119] Various procedures known within the art may be used for the production of monoclonal antibodies directed against CD47, or against derivatives, fragments, analogs homologs or orthologs thereof. (*See, for example*, Antibodies: A Laboratory Manual, Harlow E, and Lane D, 1988, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference). Fully human antibodies are antibody molecules in which the entire sequence of both the light chain and the heavy chain, including the CDRs, arise from human genes. Such antibodies are termed “human antibodies” or “fully human antibodies” herein. Human monoclonal antibodies are prepared, for example, using the procedures described in the Examples provided below. Human monoclonal antibodies can be also prepared by using the trioma technique; the human B-cell hybridoma technique (*see* Kozbor, et al., 1983 *Immunol Today* 4: 72); and the EBV hybridoma technique to produce human monoclonal antibodies (*see* Cole, et al., 1985 *In: MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized and may be produced by using human hybridomas (*see* Cote, et al., 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus *in vitro* (*see* Cole, et al., 1985 *In: MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96).

[00120] Antibodies are purified by well-known techniques, such as affinity chromatography using protein A or protein G, which provide primarily the IgG fraction of

immune serum. Subsequently, or alternatively, the specific antigen which is the target of the immunoglobulin sought, or an epitope thereof, may be immobilized on a column to purify the immune specific antibody by immunoaffinity chromatography. Purification of immunoglobulins is discussed, for example, by D. Wilkinson (The Scientist, published by The Scientist, Inc., Philadelphia PA, Vol. 14, No. 8 (April 17, 2000), pp. 25-28).

[00121] The CD47 antibodies of the invention are monoclonal antibodies. Monoclonal antibodies that modulate, block, inhibit, reduce, antagonize, neutralize or otherwise interfere with CD47- and/or CD47/SIRPa-mediated cell signaling are generated, *e.g.*, by immunizing an animal with membrane bound and/or soluble CD47, such as, for example, human CD47 or an immunogenic fragment, derivative or variant thereof. Alternatively, the animal is immunized with cells transfected with a vector containing a nucleic acid molecule encoding CD47 such that CD47 is expressed and associated with the surface of the transfected cells. Alternatively, the antibodies are obtained by screening a library that contains antibody or antigen binding domain sequences for binding to CD47. This library is prepared, *e.g.*, in bacteriophage as protein or peptide fusions to a bacteriophage coat protein that is expressed on the surface of assembled phage particles and the encoding DNA sequences contained within the phage particles (*i.e.*, “phage displayed library”). Hybridomas resulting from myeloma/B cell fusions are then screened for reactivity to CD47.

[00122] Monoclonal antibodies are prepared, for example, using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes can be immunized *in vitro*.

[00123] The immunizing agent will typically include the protein antigen, a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, (1986) pp. 59-103). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture

medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (“HAT medium”), which substances prevent the growth of HGPRT-deficient cells.

[00124] Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of monoclonal antibodies. (See Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63)).

[00125] The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980). Moreover, in therapeutic applications of monoclonal antibodies, it is important to identify antibodies having a high degree of specificity and a high binding affinity for the target antigen.

[00126] After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods. (See Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103). Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells can be grown *in vivo* as ascites in a mammal.

[00127] The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

[00128] Monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as Chinese hamster ovary (CHO) cells, Human Embryonic Kidney (HEK) 293 cells, simian COS cells, PER.C6®, NS0 cells, SP2/0, YB2/0, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (*see* U.S. Patent No. 4,816,567; Morrison, *Nature* 368, 812-13 (1994)) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

Human Antibodies and Humanization of Antibodies

[00129] Monoclonal antibodies of the invention include fully human antibodies or humanized antibodies. These antibodies are suitable for administration to humans without engendering an immune response by the human against the administered immunoglobulin.

[00130] A CD47 antibody is generated, for example, using the procedures described in the Examples provided below. For example, CD47 antibodies of the invention are identified using a modified RIMMS (Repetitive Immunization Multiple Sites) immunization strategy in mice and subsequent hybridoma generation.

[00131] In other, alternative methods, a CD47 antibody is developed, for example, using phage-display methods using antibodies containing only human sequences. Such approaches are well-known in the art, *e.g.*, in WO92/01047 and U.S. Pat. No. 6,521,404, which are hereby incorporated by reference. In this approach, a combinatorial library of phage carrying random pairs of light and heavy chains are screened using natural or recombinant source of cd47 or fragments thereof. In another approach, a CD47 antibody can be produced by a process wherein at least one step of the process includes immunizing a

transgenic, non-human animal with human CD47 protein. In this approach, some of the endogenous heavy and/or kappa light chain loci of this xenogenic non-human animal have been disabled and are incapable of the rearrangement required to generate genes encoding immunoglobulins in response to an antigen. In addition, at least one human heavy chain locus and at least one human light chain locus have been stably transfected into the animal. Thus, in response to an administered antigen, the human loci rearrange to provide genes encoding human variable regions immunospecific for the antigen. Upon immunization, therefore, the xenomouse produces B-cells that secrete fully human immunoglobulins.

[00132] A variety of techniques are well-known in the art for producing xenogenic non-human animals. For example, see U.S. Pat. No. 6,075,181 and No. 6,150,584, which is hereby incorporated by reference in its entirety. This general strategy was demonstrated in connection with generation of the first XenoMouse™ strains as published in 1994. *See* Green *et al.* *Nature Genetics* 7:13-21 (1994), which is hereby incorporated by reference in its entirety. *See also*, U.S. Patent Nos. 6,162,963; 6,150,584; 6,114,598; 6,075,181; and 5,939,598 and Japanese Patent Nos. 3 068 180 B2, 3 068 506 B2, and 3 068 507 B2 and European Patent No., EP 0 463 151 B1 and International Patent Applications No. WO 94/02602, WO 96/34096, WO 98/24893, WO 00/76310 and related family members.

[00133] In an alternative approach, others have utilized a “minilocus” approach in which an exogenous Ig locus is mimicked through the inclusion of pieces (individual genes) from the Ig locus. Thus, one or more VH genes, one or more DH genes, one or more JH genes, a mu constant region, and a second constant region (preferably a gamma constant region) are formed into a construct for insertion into an animal. *See e.g.*, U.S. Patent Nos. 5,545,806; 5,545,807; 5,591,669; 5,612,205; 5,625,825; 5,625,126; 5,633,425; 5,643,763; 5,661,016; 5,721,367; 5,770,429; 5,789,215; 5,789,650; 5,814,318; 5,877; 397; 5,874,299; 6,023,010; and 6,255,458; and European Patent No. 0 546 073 B1; and International Patent Application Nos. WO 92/03918, WO 92/22645, WO 92/22647, WO 92/22670, WO 93/12227, WO 94/00569, WO 94/25585, WO 96/14436, WO 97/13852, and WO 98/24884 and related family members.

[00134] Generation of human antibodies from mice in which, through microcell fusion, large pieces of chromosomes, or entire chromosomes, have been introduced, has also been demonstrated. *See* European Patent Application Nos. 773 288 and 843 961.

[00135] Human anti-mouse antibody (HAMA) responses have led the industry to prepare chimeric or otherwise humanized antibodies. While chimeric antibodies have a human constant region and an immune variable region, it is expected that certain human

anti-chimeric antibody (HACA) responses will be observed, particularly in chronic or multi-dose utilizations of the antibody. Thus, it would be desirable to provide fully human antibodies against CD47 in order to vitiate or otherwise mitigate concerns and/or effects of HAMA or HACA response.

[00136] The production of antibodies with reduced immunogenicity is also accomplished via humanization, chimerization and display techniques using appropriate libraries. It will be appreciated that murine antibodies or antibodies from other species can be humanized or primatized using techniques well known in the art. *See e.g.*, Winter and Harris Immunol Today 14:43 46 (1993) and Wright *et al.* Crit, Reviews in Immunol. 12:125-168 (1992). The antibody of interest may be engineered by recombinant DNA techniques to substitute the CH1, CH2, CH3, hinge domains, and/or the framework domain with the corresponding human sequence (*See* WO 92102190 and U.S. Patent Nos. 5,530,101; 5,585,089; 5,693,761; 5,693,792; 5,714,350; and 5,777,085). Also, the use of Ig cDNA for construction of chimeric immunoglobulin genes is known in the art (Liu *et al.* P.N.A.S. 84:3439 (1987) and J. Immunol. 139:3521 (1987)). mRNA is isolated from a hybridoma or other cell producing the antibody and used to produce cDNA. The cDNA of interest may be amplified by the polymerase chain reaction using specific primers (U.S. Pat. Nos. 4,683,195 and 4,683,202). Alternatively, a library is made and screened to isolate the sequence of interest. The DNA sequence encoding the variable region of the antibody is then fused to human constant region sequences. The sequences of human constant regions genes may be found in Kabat *et al.* (1991) Sequences of Proteins of immunological Interest, N.I.H. publication no. 91-3242. Human C region genes are readily available from known clones. The choice of isotype will be guided by the desired effector functions, such as complement fixation, or activity in antibody-dependent cellular cytotoxicity. Preferred isotypes are IgG1, IgG2, IgG3, and IgG4. Either of the human light chain constant regions, kappa or lambda, may be used. The chimeric, humanized antibody is then expressed by conventional methods.

[00137] Antibody fragments, such as Fv, F(ab')₂ and Fab may be prepared by cleavage of the intact protein, *e.g.*, by protease or chemical cleavage. Alternatively, a truncated gene is designed. For example, a chimeric gene encoding a portion of the F(ab')₂ fragment would include DNA sequences encoding the CH1 domain and hinge region of the H chain, followed by a translational stop codon to yield the truncated molecule.

[00138] Consensus sequences of H and L J regions may be used to design oligonucleotides for use as primers to introduce useful restriction sites into the J region for

subsequent linkage of V region segments to human C region segments. C region cDNA can be modified by site directed mutagenesis to place a restriction site at the analogous position in the human sequence.

[00139] Expression vectors include plasmids, retroviruses, YACs, EBV derived episomes, and the like. A convenient vector is one that encodes a functionally complete human CH or CL immunoglobulin sequence, with appropriate restriction sites engineered so that any VH or VL sequence can be easily inserted and expressed. In such vectors, splicing usually occurs between the splice donor site in the inserted J region and the splice acceptor site preceding the human C region, and also at the splice regions that occur within the human CH exons. Polyadenylation and transcription termination occur at native chromosomal sites downstream of the coding regions. The resulting chimeric antibody may be joined to any strong promoter, including retroviral LTRs, *e.g.*, SV-40 early promoter, (Okayama *et al.* Mol. Cell. Bio. 3:280 (1983)), Rous sarcoma virus LTR (Gorman *et al.* P.N.A.S. 79:6777 (1982)), and moloney murine leukemia virus LTR (Grosschedl *et al.* Cell 41:885 (1985)). Also, as will be appreciated, native Ig promoters and the like may be used.

[00140] Further, human antibodies or antibodies from other species can be generated through display type technologies, including, without limitation, phage display, retroviral display, ribosomal display, and other techniques, using techniques well known in the art and the resulting molecules can be subjected to additional maturation, such as affinity maturation, as such techniques are well known in the art. Wright *et al.* Crit. Reviews in Immunol. 12:125-168 (1992), Hanes and Plückthun PNAS USA 94:4937-4942 (1997) (ribosomal display), Parmley and Smith Gene 73:305-318 (1988) (phage display), Scott, TIBS, vol. 17:241-245 (1992), Cwirla *et al.* PNAS USA 87:6378-6382 (1990), Russel *et al.* Nucl. Acids Research 21:1081-1085 (1993), Hoganboom *et al.* Immunol. Reviews 130:43-68 (1992), Chiswell and McCafferty TIBTECH; 10:80-8A (1992), and U.S. Patent No. 5,733,743. If display technologies are utilized to produce antibodies that are not human, such antibodies can be humanized as described above.

[00141] Using these techniques, antibodies can be generated to CD47 expressing cells, soluble forms of CD47, epitopes or peptides thereof, and expression libraries thereto (See *e.g.*, U.S. Patent No. 5,703,057) which can thereafter be screened as described above for the activities described herein.

[00142] The CD47 antibodies of the invention can be expressed by a vector containing a DNA segment encoding the single chain antibody described above.

[00143] These can include vectors, liposomes, naked DNA, adjuvant-assisted DNA, gene gun, catheters, *etc.* Vectors include chemical conjugates such as described in WO 93/64701, which has targeting moiety (*e.g.* a ligand to a cellular surface receptor), and a nucleic acid binding moiety (*e.g.* polylysine), viral vector (*e.g.* a DNA or RNA viral vector), fusion proteins such as described in PCT/US95/02140 (WO 95/22618) which is a fusion protein containing a target moiety (*e.g.* an antibody specific for a target cell) and a nucleic acid binding moiety (*e.g.* a protamine), plasmids, phage, *etc.* The vectors can be chromosomal, non-chromosomal or synthetic.

[00144] Preferred vectors include viral vectors, fusion proteins and chemical conjugates. Retroviral vectors include moloney murine leukemia viruses. DNA viral vectors are preferred. These vectors include pox vectors such as orthopox or avipox vectors, herpesvirus vectors such as a herpes simplex I virus (HSV) vector (*see* Geller, A. I. et al., *J. Neurochem.* 64:487 (1995); Lim, F., et al., in *DNA Cloning: Mammalian Systems*, D. Glover, Ed. (Oxford Univ. Press, Oxford England) (1995); Geller, A. I. et al., *Proc Natl. Acad. Sci.: U.S.A.* 90:7603 (1993); Geller, A. I., et al., *Proc Natl. Acad. Sci USA* 87:1149 (1990), Adenovirus Vectors (*see* LeGal LaSalle et al., *Science*, 259:988 (1993); Davidson, et al., *Nat. Genet.* 3:219 (1993); Yang, et al., *J. Virol.* 69:2004 (1995) and Adeno-associated Virus Vectors (*see* Kaplitt, M. G., et al., *Nat. Genet.* 8:148 (1994)).

[00145] Pox viral vectors introduce the gene into the cells cytoplasm. Avipox virus vectors result in only a short term expression of the nucleic acid. Adenovirus vectors, adeno-associated virus vectors and herpes simplex virus (HSV) vectors are preferred for introducing the nucleic acid into neural cells. The adenovirus vector results in a shorter term expression (about 2 months) than adeno-associated virus (about 4 months), which in turn is shorter than HSV vectors. The particular vector chosen will depend upon the target cell and the condition being treated. The introduction can be by standard techniques, *e.g.* infection, transfection, transduction or transformation. Examples of modes of gene transfer include *e.g.*, naked DNA, CaPO₄ precipitation, DEAE dextran, electroporation, protoplast fusion, lipofection, cell microinjection, and viral vectors.

[00146] The vector can be employed to target essentially any desired target cell. For example, stereotaxic injection can be used to direct the vectors (*e.g.* adenovirus, HSV) to a desired location. Additionally, the particles can be delivered by intracerebroventricular (icv) infusion using a minipump infusion system, such as a SynchroMed Infusion System. A method based on bulk flow, termed convection, has also proven effective at delivering large molecules to extended areas of the brain and may be useful in delivering the vector to the

target cell. (See Bobo et al., Proc. Natl. Acad. Sci. USA 91:2076-2080 (1994); Morrison et al., Am. J. Physiol. 266:292-305 (1994)). Other methods that can be used include catheters, intravenous, parenteral, intraperitoneal and subcutaneous injection, and oral or other known routes of administration.

[00147] These vectors can be used to express large quantities of antibodies that can be used in a variety of ways. For example, to detect the presence of CD47 in a sample. The antibody can also be used to try to bind to and disrupt CD47- and/or the CD47/SIRP α interaction and CD47/SIRP α -mediated signaling.

[00148] Techniques can be adapted for the production of single-chain antibodies specific to an antigenic protein of the invention (see e.g., U.S. Patent No. 4,946,778). In addition, methods can be adapted for the construction of Fab expression libraries (see e.g., Huse, et al., 1989 Science 246: 1275-1281) to allow rapid and effective identification of monoclonal Fab fragments with the desired specificity for a protein or derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the idiotypes to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F(ab')₂ fragment produced by pepsin digestion of an antibody molecule; (ii) an Fab fragment generated by reducing the disulfide bridges of an F_(ab')₂ fragment; (iii) an Fab fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F_v fragments.

[00149] The invention also includes F_v, Fab, Fab' and F(ab')₂ CD47 fragments, single chain CD47 antibodies, single domain antibodies (e.g., nanobodies or VHJs), bispecific CD47 antibodies, and heteroconjugate CD47 antibodies.

[00150] Bispecific antibodies are antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for CD47. The second binding target is any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

[00151] Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, Nature, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed

in WO 93/08829, published 13 May 1993, and in Traunecker et al., EMBO J., 10:3655-3659 (1991).

[00152] Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

[00153] According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (*e.g.* tyrosine or tryptophan). Compensatory “cavities” of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (*e.g.* alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[00154] Bispecific antibodies can be prepared as full length antibodies or antibody fragments (*e.g.* F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., Science 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

[00155] Additionally, Fab' fragments can be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

[00156] Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., *J. Immunol.* 152:5368 (1994).

[00157] Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

[00158] Exemplary bispecific antibodies can bind to two different epitopes, at least one of which originates in the protein antigen of the invention. Alternatively, an anti-antigenic arm of an immunoglobulin molecule can be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (*e.g.* CD2, CD3, CD28, or B7), or Fc receptors for IgG (Fc γ R), such as Fc γ RI (CD64), Fc γ RII (CD32) and Fc γ RIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular antigen. Bispecific antibodies can also be used to direct cytotoxic agents to cells which express a particular antigen. These antibodies possess an antigen-binding arm and an

arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the protein antigen described herein and further binds tissue factor (TF).

[00159] Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (see U.S. Patent No. 4,676,980), and for treatment of HIV infection (see WO 91/00360; WO 92/200373; EP 03089). It is contemplated that the antibodies can be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptopbutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

[00160] It can be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, *e.g.*, the effectiveness of the antibody in treating diseases and disorders associated with aberrant CD47 signaling. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). (See Caron et al., *J. Exp Med.*, 176: 1191-1195 (1992) and Shope, *J. Immunol.*, 148: 2918-2922 (1992)). Alternatively, an antibody can be engineered that has dual Fc regions and can thereby have enhanced complement lysis and ADCC capabilities. (See Stevenson et al., *Anti-Cancer Drug Design*, 3: 219-230 (1989)).

[00161] The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).

[00162] Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), *momordica charantia* inhibitor, curcin, crotin, *sapaonaria officinalis* inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the trichothecenes. A variety

of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

[00163] Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., *Science* 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. (See WO94/11026).

[00164] Those of ordinary skill in the art will recognize that a large variety of possible moieties can be coupled to the resultant antibodies of the invention. (See, for example, "Conjugate Vaccines", Contributions to Microbiology and Immunology, J. M. Cruse and R. E. Lewis, Jr (eds), Carger Press, New York, (1989), the entire contents of which are incorporated herein by reference).

[00165] Coupling may be accomplished by any chemical reaction that will bind the two molecules so long as the antibody and the other moiety retain their respective activities. This linkage can include many chemical mechanisms, for instance covalent binding, affinity binding, intercalation, coordinate binding and complexation. The preferred binding is, however, covalent binding. Covalent binding can be achieved either by direct condensation of existing side chains or by the incorporation of external bridging molecules. Many bivalent or polyvalent linking agents are useful in coupling protein molecules, such as the antibodies of the present invention, to other molecules. For example, representative coupling agents can include organic compounds such as thioesters, carbodiimides, succinimide esters, diisocyanates, glutaraldehyde, diazobenzenes and hexamethylene diamines. This listing is not intended to be exhaustive of the various classes of coupling agents known in the art but, rather, is exemplary of the more common coupling agents. (See Killen and Lindstrom, *Jour. Immun.* 133:1335-2549 (1984); Jansen et al., *Immunological Reviews* 62:185-216 (1982); and Vitetta et al., *Science* 238:1098 (1987)).

[00166] Preferred linkers are described in the literature. (See, for example, Ramakrishnan, S. et al., *Cancer Res.* 44:201-208 (1984) describing use of MBS (M-

maleimidobenzoyl-N-hydroxysuccinimide ester). *See also*, U.S. Patent No. 5,030,719, describing use of halogenated acetyl hydrazide derivative coupled to an antibody by way of an oligopeptide linker. Particularly preferred linkers include: (i) EDC (1-ethyl-3-(3-dimethylamino-propyl) carbodiimide hydrochloride; (ii) SMPT (4-succinimidylsuccinimido-carbonyl-alpha-methyl-alpha-(2-pridyl-dithio)-toluene (Pierce Chem. Co., Cat. (21558G); (iii) SPDP (succinimidyl-6 [3-(2-pyridyl)dithio] propionamido]hexanoate (Pierce Chem. Co., Cat #21651G); (iv) Sulfo-LC-SPDP (sulfosuccinimidyl 6 [3-(2-pyridyl)dithio]-propianamide] hexanoate (Pierce Chem. Co. Cat. #2165-G); and (v) sulfo-NHS (N-hydroxysulfo-succinimide: Pierce Chem. Co., Cat. #24510) conjugated to EDC.

[00167] The linkers described above contain components that have different attributes, thus leading to conjugates with differing physio-chemical properties. For example, sulfo-NHS esters of alkyl carboxylates are more stable than sulfo-NHS esters of aromatic carboxylates. NHS-ester containing linkers are less soluble than sulfo-NHS esters. Further, the linker SMPT contains a sterically hindered disulfide bond, and can form conjugates with increased stability. Disulfide linkages, are in general, less stable than other linkages because the disulfide linkage is cleaved *in vitro*, resulting in less conjugate available. Sulfo-NHS, in particular, can enhance the stability of carbodiimide couplings. Carbodiimide couplings (such as EDC) when used in conjunction with sulfo-NHS, forms esters that are more resistant to hydrolysis than the carbodiimide coupling reaction alone.

[00168] The antibodies disclosed herein can also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

[00169] Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction.

Use of antibodies against CD47

[00170] It will be appreciated that administration of therapeutic entities in accordance with the invention will be administered with suitable carriers, excipients, and other agents that are incorporated into formulations to provide improved transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences (15th ed, Mack Publishing Company, Easton, PA (1975)), particularly Chapter 87 by Blaug, Seymour, therein. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as LipofectinTM), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. Any of the foregoing mixtures may be appropriate in treatments and therapies in accordance with the present invention, provided that the active ingredient in the formulation is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. *See also* Baldrick P. "Pharmaceutical excipient development: the need for preclinical guidance." *Regul. Toxicol Pharmacol.* 32(2):210-8 (2000), Wang W. "Lyophilization and development of solid protein pharmaceuticals." *Int. J. Pharm.* 203(1-2):1-60 (2000), Charman WN "Lipids, lipophilic drugs, and oral drug delivery-some emerging concepts." *J Pharm Sci.* 89(8):967-78 (2000), Powell *et al.* "Compendium of excipients for parenteral formulations" *PDA J Pharm Sci Technol.* 52:238-311 (1998) and the citations therein for additional information related to formulations, excipients and carriers well known to pharmaceutical chemists.

[00171] In one embodiment, antibodies of the invention, which include a monoclonal antibody of the invention, may be used as therapeutic agents. Such agents will generally be employed to diagnose, prognose, monitor, treat, alleviate, and/or prevent a disease or pathology associated with aberrant CD47 expression, activity and/or signaling in a subject. A therapeutic regimen is carried out by identifying a subject, *e.g.*, a human patient suffering from (or at risk of developing) a disease or disorder associated with aberrant CD47 expression, activity and/or signaling, *e.g.*, a cancer or other neoplastic disorder, using standard methods. An antibody preparation, preferably one having high specificity and high affinity for its target antigen, is administered to the subject and will generally have an effect due to its binding with the target. Administration of the antibody may abrogate or inhibit or

interfere with the expression, activity and/or signaling function of the target (*e.g.*, CD47). Administration of the antibody may abrogate or inhibit or interfere with the binding of the target (*e.g.*, CD47) with an endogenous ligand (*e.g.*, SIRP α) to which it naturally binds. For example, the antibody binds to the target and modulates, blocks, inhibits, reduces, antagonizes, neutralizes, or otherwise interferes with CD47 expression, activity and/or signaling.

[00172] Diseases or disorders related to aberrant CD47 expression, activity and/or signaling include, by way of non-limiting example, hematological cancer and/or solid tumors. Hematological cancers include, *e.g.*, leukemia, lymphoma and myeloma. Certain forms of leukemia include, by way of non-limiting example, acute lymphocytic leukemia (ALL); acute myeloid leukemia (AML); chronic lymphocytic leukemia (CLL); chronic myelogenous leukemia (CML); Myeloproliferative disorder/neoplasm (MPDS); and myelodysplasia syndrome. Certain forms of lymphoma include, by way of non-limiting example, Hodgkin's lymphoma, both indolent and aggressive non-Hodgkin's lymphoma, Burkitt's lymphoma, and follicular lymphoma (small cell and large cell). Certain forms of myeloma include, by way of non-limiting example, multiple myeloma (MM), giant cell myeloma, heavy-chain myeloma, and light chain or Bence-Jones myeloma. Solid tumors include, *e.g.*, breast tumors, ovarian tumors, lung tumors, pancreatic tumors, prostate tumors, melanoma tumors, colorectal tumors, lung tumors, head and neck tumors, bladder tumors, esophageal tumors, liver tumors, and kidney tumors.

[00173] Symptoms associated with cancers and other neoplastic disorders include, for example, inflammation, fever, general malaise, fever, pain, often localized to the inflamed area, loss of appetite, weight loss, edema, headache, fatigue, rash, anemia, muscle weakness, muscle fatigue and abdominal symptoms such as, for example, abdominal pain, diarrhea or constipation.

[00174] A therapeutically effective amount of an antibody of the invention relates generally to the amount needed to achieve a therapeutic objective. As noted above, this may be a binding interaction between the antibody and its target antigen that, in certain cases, interferes with the functioning of the target. The amount required to be administered will furthermore depend on the binding affinity of the antibody for its specific antigen, and will also depend on the rate at which an administered antibody is depleted from the free volume other subject to which it is administered. Common ranges for therapeutically effective dosing of an antibody or antibody fragment of the invention may be, by way of

nonlimiting example, from about 0.1 mg/kg body weight to about 100 mg/kg body weight. Common dosing frequencies may range, for example, from twice daily to once a week.

[00175] Efficaciousness of treatment is determined in association with any known method for diagnosing or treating the particular inflammatory-related disorder. Alleviation of one or more symptoms of the inflammatory-related disorder indicates that the antibody confers a clinical benefit.

[00176] Methods for the screening of antibodies that possess the desired specificity include, but are not limited to, enzyme linked immunosorbent assay (ELISA) and other immunologically mediated techniques known within the art.

[00177] In another embodiment, antibodies directed against CD47 may be used in methods known within the art relating to the localization and/or quantitation of CD47 (*e.g.*, for use in measuring levels of CD47 and/or both CD47 and SIRP α within appropriate physiological samples, for use in diagnostic methods, for use in imaging the protein, and the like). In a given embodiment, antibodies specific to CD47, or derivative, fragment, analog or homolog thereof, that contain the antibody derived antigen binding domain, are utilized as pharmacologically active compounds (referred to hereinafter as “Therapeutics”).

[00178] In another embodiment, an antibody specific for CD47 can be used to isolate a CD47 polypeptide, by standard techniques, such as immunoaffinity, chromatography or immunoprecipitation. Antibodies directed against the CD47 protein (or a fragment thereof) can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

[00179] In yet another embodiment, an antibody according to the invention can be used as an agent for detecting the presence of CD47 and/or both CD47 and SIRPa protein (or a protein fragment thereof) in a sample. In some embodiments, the antibody contains a detectable label. Antibodies are polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab, scFv, or F(ab')₂) is used. The term “labeled”, with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin. The term “biological sample” is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. Included within the usage of the term “biological sample”, therefore, is blood and a fraction or component of blood including blood serum, blood plasma, or lymph. That is, the detection method of the invention can be used to detect an analyte mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of an analyte mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of an analyte protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, and immunofluorescence. *In vitro* techniques for detection of an analyte genomic DNA include Southern hybridizations. Procedures for conducting immunoassays are described, for example in “ELISA: Theory and Practice: Methods in Molecular Biology”, Vol. 42, J. R. Crowther (Ed.) Human Press, Totowa, NJ, 1995; “Immunoassay”, E. Diamandis and T. Christopoulos, Academic Press, Inc., San Diego, CA, 1996; and “Practice and Theory of Enzyme Immunoassays”, P. Tijssen, Elsevier Science Publishers, Amsterdam, 1985. Furthermore, *in vivo* techniques for detection of an analyte protein include introducing into a subject a labeled anti-analyte protein antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

Therapeutic Administration and Formulations of CD47 antibodies

[00180] The antibodies of the invention (also referred to herein as “active compounds”), and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Principles and considerations involved in preparing such compositions, as well as guidance in the choice of

components are provided, for example, in Remington's Pharmaceutical Sciences: The Science And Practice Of Pharmacy 19th ed. (Alfonso R. Gennaro, et al., editors) Mack Pub. Co., Easton, Pa.: 1995; Drug Absorption Enhancement: Concepts, Possibilities, Limitations, And Trends, Harwood Academic Publishers, Langhorne, Pa., 1994; and Peptide And Protein Drug Delivery (Advances In Parenteral Sciences, Vol. 4), 1991, M. Dekker, New York.

[00181] Such compositions typically comprise the antibody and a pharmaceutically acceptable carrier. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. (See, e.g., Marasco et al., Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993)).

[00182] As used herein, the term "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, ringer's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated.

[00183] The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

[00184] A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (*i.e.*, topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or

sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[00185] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

[00186] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[00187] Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[00188] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

[00189] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[00190] The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[00191] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as sustained/controlled release formulations, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art.

[00192] For example, the active ingredients can be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions.

[00193] Sustained-release preparations can be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT TM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

[00194] The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) and can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

[00195] It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

[00196] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

[00197] The formulation can also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[00198] In one embodiment, the active compounds are administered in combination therapy, *i.e.*, combined with other agents, *e.g.*, therapeutic agents, that are useful for treating pathological conditions or disorders, such as various forms of cancer, autoimmune disorders and inflammatory diseases. The term “in combination” in this context means that the agents are given substantially contemporaneously, either simultaneously or sequentially. If given sequentially, at the onset of administration of the second compound, the first of the two compounds is preferably still detectable at effective concentrations at the site of treatment.

[00199] For example, the combination therapy can include one or more antibodies of the invention coformulated with, and/or coadministered with, one or more additional therapeutic agents, *e.g.*, one or more cytokine and growth factor inhibitors, immunosuppressants, anti-inflammatory agents, metabolic inhibitors, enzyme inhibitors, and/or cytotoxic or cytostatic agents, as described in more detail below. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

[00200] Preferred therapeutic agents used in combination with an antibody of the invention are those agents that interfere at different stages in an inflammatory response. In one embodiment, one or more antibodies described herein may be coformulated with, and/or coadministered with, one or more additional agents such as other cytokine or growth factor antagonists (*e.g.*, soluble receptors, peptide inhibitors, small molecules, ligand fusions); or antibodies or antigen binding fragments thereof that bind to other targets (*e.g.*, antibodies that bind to other cytokines or growth factors, their receptors, or other cell surface molecules); and anti-inflammatory cytokines or agonists thereof.

[00201] In other embodiments, the antibodies of the invention are used as vaccine adjuvants against autoimmune disorders, inflammatory diseases, etc. The combination of adjuvants for treatment of these types of disorders are suitable for use in combination with a wide variety of antigens from targeted self-antigens, *i.e.*, autoantigens, involved in

autoimmunity, *e.g.*, myelin basic protein; inflammatory self-antigens, *e.g.*, amyloid peptide protein, or transplant antigens, *e.g.*, alloantigens. The antigen may comprise peptides or polypeptides derived from proteins, as well as fragments of any of the following: saccharides, proteins, polynucleotides or oligonucleotides, autoantigens, amyloid peptide protein, transplant antigens, allergens, or other macromolecular components. In some instances, more than one antigen is included in the antigenic composition.

Design and Generation of Other Therapeutics

[00202] In accordance with the present invention and based on the activity of the antibodies that are produced and characterized herein with respect to CD47, the design of other therapeutic modalities beyond antibody moieties is facilitated. Such modalities include, without limitation, advanced antibody therapeutics, such as bispecific antibodies, immunotoxins, and radiolabeled therapeutics, generation of peptide therapeutics, gene therapies, particularly intrabodies, antisense therapeutics, and small molecules.

[00203] For example, in connection with bispecific antibodies, bispecific antibodies can be generated that comprise (i) two antibodies- one with a specificity to CD47 and another to a second molecule that are conjugated together, (ii) a single antibody that has one chain specific to CD47 and a second chain specific to a second molecule, or (iii) a single chain antibody that has specificity to CD47 and a second molecule. Such bispecific antibodies are generated using techniques that are well known for example, in connection with (i) and (ii) *See e.g.*, Fanger *et al.* Immunol Methods 4:72-81 (1994) and Wright *et al.* Crit, Reviews in Immunol. 12:125-168 (1992), and in connection with (iii) *See e.g.*, Traunecker *et al.* Int. J. Cancer (Suppl.) 7:51-52 (1992).

[00204] In connection with immunotoxins, antibodies can be modified to act as immunotoxins utilizing techniques that are well known in the art. *See e.g.*, Vitetta Immunol Today 14:252 (1993). *See also* U.S. Patent No. 5,194,594. In connection with the preparation of radiolabeled antibodies, such modified antibodies can also be readily prepared utilizing techniques that are well known in the art. *See e.g.*, Junghans *et al.* in Cancer Chemotherapy and Biotherapy 655-686 (2d edition, Chafner and Longo, eds., Lippincott Raven (1996)). *See also* U.S. Patent Nos. 4,681,581, 4,735,210, 5,101,827, 5,102,990 (RE 35,500), 5,648,471, and 5,697,902. Each of immunotoxins and radiolabeled molecules would be likely to kill cells expressing CD47.

[00205] In connection with the generation of therapeutic peptides, through the utilization of structural information related to CD47 and antibodies thereto, such as the

antibodies of the invention or screening of peptide libraries, therapeutic peptides can be generated that are directed against CD47. Design and screening of peptide therapeutics is discussed in connection with Houghten *et al.* *Biotechniques* 13:412-421 (1992), Houghten *PNAS USA* 82:5131-5135 (1985), Pinalla *et al.* *Biotechniques* 13:901-905 (1992), Blake and Litzi-Davis *BioConjugate Chem.* 3:510-513 (1992). Immunotoxins and radiolabeled molecules can also be prepared, and in a similar manner, in connection with peptidic moieties as discussed above in connection with antibodies. Assuming that the CD47 molecule (or a form, such as a splice variant or alternate form) is functionally active in a disease process, it will also be possible to design gene and antisense therapeutics thereto through conventional techniques. Such modalities can be utilized for modulating the function of CD47. In connection therewith the antibodies of the present invention facilitate design and use of functional assays related thereto. A design and strategy for antisense therapeutics is discussed in detail in International Patent Application No. WO 94/29444. Design and strategies for gene therapy are well known. However, in particular, the use of gene therapeutic techniques involving intrabodies could prove to be particularly advantageous. *See e.g.*, Chen *et al.* *Human Gene Therapy* 5:595-601 (1994) and Marasco *Gene Therapy* 4:11-15 (1997). General design of and considerations related to gene therapeutics is also discussed in International Patent Application No. WO 97/38137.

[00206] Knowledge gleaned from the structure of the CD47 molecule and its interactions with other molecules in accordance with the present invention, such as SIRP α and/or the antibodies of the invention, and others can be utilized to rationally design additional therapeutic modalities. In this regard, rational drug design techniques such as X-ray crystallography, computer-aided (or assisted) molecular modeling (CAMM), quantitative or qualitative structure-activity relationship (QSAR), and similar technologies can be utilized to focus drug discovery efforts. Rational design allows prediction of protein or synthetic structures which can interact with the molecule or specific forms thereof which can be used to modify or modulate the activity of IL-6R α . Such structures can be synthesized chemically or expressed in biological systems. This approach has been reviewed in Capsey *et al.* *Genetically Engineered Human Therapeutic Drugs* (Stockton Press, NY (1988)). Further, combinatorial libraries can be designed and synthesized and used in screening programs, such as high throughput screening efforts.

Screening Methods

[00207] The invention provides methods (also referred to herein as “screening assays”) for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*,

peptides, peptidomimetics, small molecules or other drugs) that modulate or otherwise interfere with the binding of CD47 to SIRP α , or candidate or test compounds or agents that modulate or otherwise interfere with the signaling function of CD47 and/or CD47-SIRP α . Also provided are methods of identifying compounds useful to treat disorders associated with aberrant CD47 and/or CD47-SIRP α expression, activity and/or signaling. The screening methods can include those known or used in the art or those described herein. For example, CD47 can be immobilized on a microtiter plate and incubated with a candidate or test compound, *e.g.*, a CD47 antibody, in the presence of SIRP α . Subsequently, bound SIRP α can be detected using a secondary antibody, and absorbance can be detected on a plate reader.

[00208] Methods of identifying compounds capable of promoting phagocytosis of tumor cells by macrophages are also provided. These methods can include those known or used in the art or those described herein. For example, macrophages are incubated with labeled tumor cells in the presence of a candidate compound, *e.g.*, a CD47 antibody. After a period of time, the macrophages can be observed for internalization of the tumor label to identify phagocytosis. Additional details regarding these methods, *e.g.*, SIRP α blocking assays and phagocytosis assays, are provided in the Examples.

[00209] The invention also includes compounds identified in the screening assays described herein.

[00210] In one embodiment, the invention provides assays for screening candidate or test compounds which modulate the signaling function of CD47. The test compounds of the invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the “one-bead one-compound” library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds. (*See, e.g.*, Lam, 1997. *Anticancer Drug Design* 12: 145).

[00211] A “small molecule” as used herein, is meant to refer to a composition that has a molecular weight of less than about 5 kD and most preferably less than about 4 kD. Small molecules can be, *e.g.*, nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids or other organic or inorganic molecules. Libraries of chemical and/or biological mixtures, such as fungal, bacterial, or algal extracts, are known in the art and can be screened with any of the assays of the invention.

[00212] Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt, et al., 1993. Proc. Natl. Acad. Sci. U.S.A. 90: 6909; Erb, et al., 1994. Proc. Natl. Acad. Sci. U.S.A. 91: 11422; Zuckermann, et al., 1994. J. Med. Chem. 37: 2678; Cho, et al., 1993. Science 261: 1303; Carrell, et al., 1994. Angew. Chem. Int. Ed. Engl. 33: 2059; Carell, et al., 1994. Angew. Chem. Int. Ed. Engl. 33: 2061; and Gallop, et al., 1994. J. Med. Chem. 37: 1233.

[00213] Libraries of compounds may be presented in solution (*see e.g.*, Houghten, 1992. Biotechniques 13: 412-421), or on beads (*see* Lam, 1991. Nature 354: 82-84), on chips (*see* Fodor, 1993. Nature 364: 555-556), bacteria (*see* U.S. Patent No. 5,223,409), spores (*see* U.S. Patent 5,233,409), plasmids (*see* Cull, et al., 1992. Proc. Natl. Acad. Sci. USA 89: 1865-1869) or on phage (*see* Scott and Smith, 1990. Science 249: 386-390; Devlin, 1990. Science 249: 404-406; Cwirla, et al., 1990. Proc. Natl. Acad. Sci. U.S.A. 87: 6378-6382; Felici, 1991. J. Mol. Biol. 222: 301-310; and U.S. Patent No. 5,233,409.).

[00214] In one embodiment, a candidate compound is introduced to an antibody-antigen complex and determining whether the candidate compound disrupts the antibody-antigen complex, wherein a disruption of this complex indicates that the candidate compound modulates the signaling function of CD47 and/or the interaction between CD47 and SIRP α . In another embodiment, a soluble CD47 and/or both CD47 and SIRP α protein of the invention is provided and exposed to at least one neutralizing monoclonal antibody. Formation of an antibody-antigen complex is detected, and one or more candidate compounds are introduced to the complex. If the antibody-antigen complex is disrupted following introduction of the one or more candidate compounds, the candidate compounds is useful to treat disorders associated with aberrant CD47 and/or CD47-SIRP α signaling.

[00215] Determining the ability of the test compound to interfere with or disrupt the antibody-antigen complex can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the antigen or biologically-active portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically-labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

[00216] In one embodiment, the assay comprises contacting an antibody-antigen complex with a test compound, and determining the ability of the test compound to interact with the antigen or otherwise disrupt the existing antibody-antigen complex. In this embodiment, determining the ability of the test compound to interact with the antigen and/or disrupt the antibody-antigen complex comprises determining the ability of the test compound to preferentially bind to the antigen or a biologically-active portion thereof, as compared to the antibody.

[00217] In another embodiment, the assay comprises contacting an antibody-antigen complex with a test compound and determining the ability of the test compound to modulate the antibody-antigen complex. Determining the ability of the test compound to modulate the antibody-antigen complex can be accomplished, for example, by determining the ability of the antigen to bind to or interact with the antibody, in the presence of the test compound.

[00218] Those skilled in the art will recognize that, in any of the screening methods disclosed herein, the antibody may be a neutralizing antibody, which modulates or otherwise interferes with CD47 activity and/or signaling.

[00219] The screening methods disclosed herein may be performed as a cell-based assay or as a cell-free assay. The cell-free assays of the invention are amenable to use of either the soluble form or the membrane-bound form of CD47 and fragments thereof. In the case of cell-free assays comprising the membrane-bound form of CD47, it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the proteins are maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton® X-100, Triton® X-114, Thesit®, Isotridecypoly(ethylene glycol ether)_n, N-dodecyl-N,N-dimethyl-3-ammonio-1-propane sulfonate, 3-(3-cholamidopropyl) dimethylamminiol-1-propane sulfonate (CHAPS), or 3-(3-cholamidopropyl)dimethylamminiol-2-hydroxy-1-propane sulfonate (CHAPSO).

[00220] In more than one embodiment, it may be desirable to immobilize either the antibody or the antigen to facilitate separation of complexed from uncomplexed forms of one or both following introduction of the candidate compound, as well as to accommodate automation of the assay. Observation of the antibody-antigen complex in the presence and absence of a candidate compound can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein that adds a domain that allows one or both of the proteins to be bound to a matrix can be provided. For example, GST-antibody

fusion proteins or GST-antigen fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates, that are then combined with the test compound, and the mixture is incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly. Alternatively, the complexes can be dissociated from the matrix, and the level of antibody-antigen complex formation can be determined using standard techniques.

[00221] Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either the antibody (*e.g.*, the 2A1 antibody, or an antibody having a variable heavy chain selected from SEQ ID NOS: 5-30 and a variable light chain selected from SEQ ID NOS: 31-47) or the antigen (*e.g.* CD47 protein) can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated antibody or antigen molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well-known within the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, other antibodies reactive with the antibody or antigen of interest, but which do not interfere with the formation of the antibody-antigen complex of interest, can be derivatized to the wells of the plate, and unbound antibody or antigen trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using such other antibodies reactive with the antibody or antigen.

[00222] The invention further pertains to novel agents identified by any of the aforementioned screening assays and uses thereof for treatments as described herein.

Diagnostic and Prophylactic Formulations

[00223] The CD47 MAbs of the invention are used in diagnostic and prophylactic formulations. In one embodiment, a CD47 MAb of the invention is administered to patients that are at risk of developing one or more of the aforementioned diseases, such as for example, without limitation, cancer or other neoplastic condition. A patient's or organ's predisposition to one or more of the aforementioned cancers or other neoplastic conditions can be determined using genotypic, serological or biochemical markers.

[00224] In another embodiment of the invention, the CD47 antibody is administered to human individuals diagnosed with a clinical indication associated with one or more of the aforementioned diseases, such as for example, without limitation, cancer or other neoplastic condition. Upon diagnosis, the CD47 antibody is administered to mitigate or reverse the effects of the clinical indication associated with one or more of the aforementioned diseases.

[00225] Antibodies of the invention are also useful in the detection of CD47 and/or SIRP α in patient samples and accordingly are useful as diagnostics. For example, the CD47 antibodies of the invention are used in *in vitro* assays, *e.g.*, ELISA, to detect CD47 and/or SIRP α levels in a patient sample.

[00226] In one embodiment, a CD47 antibody of the invention is immobilized on a solid support (*e.g.*, the well(s) of a microtiter plate). The immobilized antibody serves as a capture antibody for any CD47 and/or SIRP α that may be present in a test sample. Prior to contacting the immobilized antibody with a patient sample, the solid support is rinsed and treated with a blocking agent such as milk protein or albumin to prevent nonspecific adsorption of the analyte.

[00227] Subsequently the wells are treated with a test sample suspected of containing the antigen, or with a solution containing a standard amount of the antigen. Such a sample is, *e.g.*, a serum sample from a subject suspected of having levels of circulating antigen considered to be diagnostic of a pathology. After rinsing away the test sample or standard, the solid support is treated with a second antibody that is detectably labeled. The labeled second antibody serves as a detecting antibody. The level of detectable label is measured, and the concentration of CD47 and/or SIRP α in the test sample is determined by comparison with a standard curve developed from the standard samples.

[00228] It will be appreciated that based on the results obtained using the CD47 antibodies of the invention in an *in vitro* diagnostic assay, it is possible to stage a disease (*e.g.*, a clinical indication associated with ischemia, an autoimmune or inflammatory disorder) in a subject based on expression levels of CD47 and/or SIRP α . For a given disease, samples of blood are taken from subjects diagnosed as being at various stages in the progression of the disease, and/or at various points in the therapeutic treatment of the disease. Using a population of samples that provides statistically significant results for each stage of progression or therapy, a range of concentrations of the antigen that may be considered characteristic of each stage is designated.

[00229] All publications and patent documents cited herein are incorporated herein by reference as if each such publication or document was specifically and individually

indicated to be incorporated herein by reference. Citation of publications and patent documents is not intended as an admission that any is pertinent prior art, nor does it constitute any admission as to the contents or date of the same. The invention having now been described by way of written description, those of skill in the art will recognize that the invention can be practiced in a variety of embodiments and that the foregoing description and examples below are for purposes of illustration and not limitation of the claims that follow.

EXAMPLES

[00230] The following examples, including the experiments conducted and results achieved are provided for illustrative purposes only and are not to be construed as limiting upon the present invention.

EXAMPLE 1: Generation and Selection of CD47 Antibodies

[00231] CD47 antibodies were generated by immunizing mice with a recombinant protein representing CD47-IgV (immunoglobulin-like variable-type), implementing a modified rapid immunization strategy in multiple sites (Kilpatrick *et al.* (1997) Rapid development of affinity matured monoclonal antibodies using RIMMS. Hybridoma 16, 381-389). In addition, half of the mice in the immunized group received a single injection of the anti-mouse GITR agonist antibody, DTA-1. Following the immunization schedule, lymph nodes from all mice (DTA-1 treated and untreated) were harvested and dissociated, thereby enabling B-cell isolation and subsequent fusion to a mouse myeloma cell line. Hybridoma supernatants were screened for binding to CD47 by ELISA and by flow cytometry on Daudi (ATCC# CCL-213) cells (Figure 1A). Hybridoma supernatants were also analyzed for the ability to block the CD47-SIRP α interaction (Figure 1B). Recombinant CD47 was immobilized on a Medisorp (NUNC) microtiter plate and subsequently incubated with the hybridoma supernatants in the presence of recombinant human SIRP α -ECD fused to a human IgG Fc domain. Bound SIRP α was detected using an HRP conjugated anti-human IgG Fc specific secondary antibody (Jackson Immuno Research), and absorbance at 650nm detected in plate reader.

EXAMPLE 2: Characterization of CD47 Antibodies

[00232] Exemplary murine CD47 antibodies of the present invention are shown in Figure 2. Affinity ranking of SIRP α blocking CD47 antibodies was conducted by flow

cytometry on Raji (ATCC# CCL-86) (Figure 2A) and CCRF-CEM (ATCC# CCL-119) cells (Figure 2B). Bound CD47 antibodies were detected using a FITC conjugated anti-mouse IgG secondary antibody (Jackson ImmunoResearch). The CD47 antibody known in the art, B6H12, was included as a positive control (See, *e.g.*, U.S. Patent 5,057,604). In Figure 2B, both the B6H12 and the 2D3, a commercially available non-SIRP α blocking antibody, were compared to antibodies generated herein. The antibodies of the present invention display higher affinity toward the endogenous (cell surface) form of CD47 compared to the B6H12 and 2D3 antibodies.

EXAMPLE 3: SIRP α Blocking Activity of CD47 antibodies

[00233] The potency of SIRP α blocking by CD47 antibodies was measured by an ELISA wherein recombinant His-tagged-CD47-IgV was immobilized on a Medisorp microtiter plate. Binding of recombinant SIRP α fused to an Fc domain of human IgG was monitored in the presence of increasing amounts of the CD47 antibodies. Bound SIRP α was determined using an HRP conjugated anti-human IgG (Fc specific) secondary antibody (Jackson ImmunoResearch). The antibodies of the present invention display enhanced potency of SIRP α blocking compared to the B6H12 antibody. Figure 3A shows representative data of the ELISA based SIRP α blocking assay.

[00234] CD47 antibodies were analyzed by flow cytometry for their ability to block recombinant SIRP α binding to cell surface CD47. CCRF-CEM (ATCC# CCL-119) cells were used as the source of CD47 in the assay and binding of recombinant SIRP α fused to an Fc domain of human IgG was monitored in the presence of increasing amounts of the CD47 antibodies. Bound SIRP α was determined using an APC conjugated anti-human IgG (Fc specific) secondary antibody (Jackson ImmunoResearch) (Figure 3B). B6H12 and commercially available non-SIRP α blocking CD47 antibody 2D3 where used a positive and negative controls respectively.

EXAMPLE 4: CD47 Antibody-mediated Homotypic interactions

[00235] SIRP α blocking CD47 antibodies were analyzed for their ability to induce cellular clustering, as known as homotypic interactions, between CD47 positive cells. Daudi and Raji cells were used as candidate CD47 expressing cells lines. Among the antibodies examined, the 2A1 antibody of the present invention was the only SIRP α blocking antibody that did not promote homotypic interactions of CD47 expressing cells.

EXAMPLE 5: Hemagglutination Activity of CD47 Antibodies

[00236] One example of a homotypic interaction is hemagglutination, as evidenced by RBC aggregation. CD47 antibodies were screened for RBC agglutination, as observed

by the ability of an antibody to prevent the settling of human RBCs. Unexpectedly, the 2A1 antibody was found to be unique among other CD47 antibodies for its inability to promote hemagglutination, while having high affinity and the ability to block SIRP α . Other antibodies that displayed reduced hemagglutination did not block SIRP α binding to CD47.

[00237] To evaluate the hemagglutinating capacity of CD47 antibodies, human RBCs were diluted to 10% in PBS and incubated at 37°C for 2-6 hours with a titration of CD47 antibodies in a round bottom 96 well plate. Evidence of hemagglutination is demonstrated by the presence of non-settled RBCs, appearing as a haze compared to a punctuate red dot of non-hemagglutinated RBCs. Unexpectedly, as shown in Figure 4A, CD47 antibodies of the invention, particularly the antibody referred to herein as 2A1, did not exhibit hemagglutinating activity. The graph shows the quantitation of the hemagglutination assay, denoted “hemagglutination index” determined by quantitating the area of the RBC pellet in the presence of the antibody, normalized to that in the absence of the antibody.

[00238] The murine 9E4 antibody caused the most profound hemagglutination at all concentrations tested. Thus, The 9E4 antibody binds CD47 and blocks CD47 interaction with SIRP α ; however, the 9E4 antibody causes profound hemagglutination.

[00239] The VH chain region of the 9E4 antibody is provided below.

EVQLRQSGPELVKPGASVKISCKASGYSFTDYYMYWVKQSRVRS LAWIGRINPYTGATGYDQNFKDKASLIVDK
SSSTAYMELRSLTSEDSAVYYCARGRNR YDGWFAYWGQGTLVTV (SEQ ID NO: 78)

[00240] The VL chain region of the 9E4 antibody is provided below.

EIQMTQTSSL SASLGDRVT ISCRASQD ISNYLNWYQQKPDGTVKLLI YYTSRLHSGVPSRFSGSGSGTDYSLT
ISNLQEDIATYFCQQGNALPPTFGGGTNLEIK (SEQ ID NO: 79)

[00241] The control antibody B6H12 caused hemagglutination as is expected for SIRP α blocking CD47 antibodies.

[00242] In order to investigate the uniqueness of the non-hemagglutinating activity of the 2A1 antibody, numerous other CD47 antibodies were screened in the RBC hemagglutination assay (Figure 4B). Included in this assay was the chimeric version of the 2A1 antibody (2A1-xi), which consists of the murine variable heavy chain region of 2A1, the murine variable light chain region of 2A1 modified at amino acid 106 (i.e., M106I), and the constant regions of human IgG1 and human Igkappa. The VH and VL region sequences of 2A1 antibody and 2A1-xi antibody are provided in Table 1. Antibodies were tested at 12.5, 25, 50, and 100nM. Unexpectedly, 2A1 is rare amongst the CD47 antibodies examined in Figure 4B, in that it was the only antibody in Figure 4B with absent or reduced

hemagglutinating activities. Figure 4E shows that 2A1, chimeric 2A1 (2A1-xi), and humanized variants do not cause hemagglutination.

[00243] Figure 4C shows the results of screening additional CD47 antibodies in the RBC hemagglutination assay. As shown in Figure 4C, the commercially available CD47 monoclonal antibody 2D3, which does not block SIRP α , did not cause hemagglutination. However, other commercially available CD47 antibodies (*e.g.*, CC2C6, BRC126, and B6H12) which block SIRP α caused hemagglutination (Figure 4C). Thus, prior to the invention described herein, existing antibodies that blocked SIRP α caused hemagglutination, while existing antibodies, such as 2D3, that did not block SIRP α did not cause hemagglutination. Taken together, the antibodies of the invention (*e.g.*, the 2A1 antibody and its humanized derivatives) are unique among existing CD47 antibodies in their ability to block SIRP α , but not cause hemagglutination.

[00244] A high concentration range of select CD47 antibodies was retested in the hemagglutination assay (Figure 4D). This assay revealed a pro-zone effect of hemagglutination by B6H12 and 9E4, wherein hemagglutination was reduced at high and low ends of the concentration range tested. The graphical representation of the hemagglutination index also highlights the pro-zone effect. The pro-zone effect was also evident in Figures 4C and 4E. Importantly, the mouse 2A1 and chimeric 2A1 CD47 antibodies were devoid of hemagglutinating activity at all concentrations.

[00245] As shown in Figure 4E, the murine 1B4 antibody displayed a narrow range of hemagglutination.

[00246] The VH chain region of the 1B4 antibody is provided below.

QIQLQQSGPELVKPGASVKISCKASGYTFTDYYIHWKQRPGQGLEWIGWIYPGSGNTKYNERFKGKATLTVAT
SSSTAYMQLSSLTSEDTAVYFCARREEDYFDYWGQQGLTV (SEQ ID NO: 80)

[00247] The VL chain region of the 1B4 antibody is provided below.

DIVMSQSPSSLAWSVGEKVTMSCKSSQSLYSSNQKNYLTWYQQKPGQSPKLLIYWASTRESGVPDFTGSGSG
TDFTLTISSVKAEDLAVYYCQQYYSPLTGAGTKLEIK (SEQ ID NO: 81)

[00248] The hemagglutinating capacity of humanized antibodies derived from the murine 2A1 was tested as above. Importantly, the representative humanized antibody AB6.12 in numerous human IgG isotypes (IgG1, IgG4-S228P, and IgG4-S228P/L235E) did not cause any RBC hemagglutination. 2A1 and 2A1-xi were included as controls for non-hemagglutinating antibodies, whereas B6H12 and 9E4 were included as positive controls for hemagglutination (Figure 4F).

EXAMPLE 6: Binding to Cynomolgus Monkey CD47

[00249] The ability of murine 2A1 to bind to cynomolgus (cyno) monkey CD47 was assessed. The B6H12 antibody has previously been reported to be cross-reactive with cyno CD47 and was used as a positive control for the presence on cyno CD47 in the assay. The experiment to measure binding of 2A1 to cynomolgus monkey CD47 was designed to compare binding of 2A1 to CD47 on cynomolgus monkey B-cells and human cells, wherein the Raji cell line was used as a human CD47 positive cell. Cynomolgus peripheral blood mononuclear cells (PBMCs) were isolated from cynomolgus whole blood by ficoll-paque gradient centrifugation. Cynomolgus and human B-cells (Raji) were labeled with the human CD20 antibody ofatumumab (Arzerra) at 10 µg/ml, and reacted with a dilution series of murine CD47 antibody 2A1 or B6H12. B-cells labeled with human CD20 antibody were detected with polyclonal anti-human antibody conjugated to DyLite 649, while the CD47 murine antibodies were detected with polyclonal anti-mouse antibody conjugated to DyLite 488. Cells were analyzed by flow cytometry, first gated on live cells by FSC and SSC, then on cells positive for FL4 (CD20 positive), and lastly the median FL1 (CD47 positive) was measured. The data were normalized by dividing the signal at each concentration by the maximum signal for each antibody on each cell population. The normalized results shown in Figure 5 reveal that 2A1 does cross react with cyno CD47 and has identical affinity as compared to human CD47. Consistent with the results presented above, B6H12 had lower affinity for cell surface CD47 on both Raji and cynomolgus B-cells compared to antibodies of the present invention.

EXAMPLE 7: Chimeric Antibody Generation

[00250] In order to identify the sequences of the variable regions of the heavy (VH) and light (VL) chains of the murine 2A1 antibody, ribonucleic acid (RNA) was isolated from the hybridoma and utilized in reverse transcription polymerase chain reaction (RT-PCR) (Phusion RT-PCR Kit Thermo Scientific) to generate first strand cDNA. A degenerative primer set that covers the complete repertoire of murine of antibody leader sequences of both VH and VL was used in a PCR wherein the first strand cDNA served as the template.

[00251] The forward primers (murine IgG leader) are provided below.

Name	Sequence
VH1-1	CACTGCAGGTRTCCACTCC (SEQ ID NO: 82)
VH1-2	CATAGCAGGTGTCCACTCC (SEQ ID NO: 83)
VH1-3	CRCTACAGGTGTCCACTCC (SEQ ID NO: 84)
VH1-4	GCYACAGMTGTCCACTCC (SEQ ID NO: 85)
VH1-5	CACTGCAGGTGTCCWMTCC (SEQ ID NO: 86)
VH1-6	CRCTRCAGGTGTCACACTCC (SEQ ID NO: 87)

VH1-7	GCTAWMGGTGTCCACTCC (SEQ ID NO: 88)
VH1-8	CCTCAGGTGTCCACTCC (SEQ ID NO: 89)
VH1-9	GCTACAGGTGCTCACTCC (SEQ ID NO: 90)
VH1-10	CACTGCAGGTGTCCCTCT (SEQ ID NO: 91)
VH1-11	CAYTGCAGGTGTCCAYTGC (SEQ ID NO: 92)
VH1-12	GCTAMMGGTGTCCACTTC (SEQ ID NO: 93)
VH1-13	CTCCTGTCAKTAACTKCAGGT (SEQ ID NO: 94)
VH1-14	CAACTGCAGGTGTCTCTCT (SEQ ID NO: 95)
VH1-15	CRCTRCAAGGYGTCCACTCT (SEQ ID NO: 96)
VH2-1	CCAAGCTGTATCCTTCC (SEQ ID NO: 97)
VH2-2	CCAAGCTGTGTCCTRCC (SEQ ID NO: 98)
VH3-1	CTTGACAGYCVTCCCKGGT (SEQ ID NO: 99)
VH3-2	CTTCACAGCCTTCCTGGT (SEQ ID NO: 100)
VH4	CTTAAAAGGGTCCAGTGT (SEQ ID NO: 101)
VH5-1	CAYTTAAAARGTGTCAAGTGT (SEQ ID NO: 102)
VH5-2	GTTTTAAAAGGTGTCTGTG (SEQ ID NO: 103)
VH6	CTYTTAAAAGGKGTCAGWG (SEQ ID NO: 104)
VH7-1	CYTTTAMATGGTATCCAGTGT (SEQ ID NO: 105)
VH7-2	CTTTTACATGGTTCAAGTGT (SEQ ID NO: 106)
VH8	GTCCTGCATATGTCYT (SEQ ID NO: 107)
VH9	GATGGCAGCWGCYCAAAG (SEQ ID NO: 108)
VH10	CTATCAAGGTGTGCATTGT (SEQ ID NO: 109)
VH11	CTTTTAAAAGWTGTCCAGKGT (SEQ ID NO: 110)
VH12	GTGACAGTCCTCCTGGTAG (SEQ ID NO: 111)
VH14	CTTCCTGATGGCAGTGGTT (SEQ ID NO: 112)
VH15	GCTACAGGTATCCAATCC (SEQ ID NO: 113)

[00252] The reverse primer (murine IgG constant) is provided below.

Name	Sequence
HC-Rev	GCGTCTAGAACAYCTCCACACACAGGRRCCAGTGGATAGAC (SEQ ID NO: 114)

[00253] The forward primers (murine IgKappa leader) are provided below.

Name	Sequence
VK1-1	CTGWTGTTCTGGATTCCCTG (SEQ ID NO: 115)
VK1-2	GGTCAGACAGTCAGCAGT (SEQ ID NO: 116)
VK2	GTGCTCTGGATTGGGAA (SEQ ID NO: 117)
VK4/5-1	CAGCTTCYTGCTAACAGTG (SEQ ID NO: 118)
VK4/5-2	CTAACAGTGCTTCAGGA (SEQ ID NO: 119)
VK8-1	GTGGGTATCTGGTRCSTGTG (SEQ ID NO: 120)
VK8-2	GGAAATTAAAAGTACCTGTGGG (SEQ ID NO: 121)
VK9A/9B-1	GGTTTCMAGGTRCCAGATGT (SEQ ID NO: 122)
VK9A/9B-2	CTCTGGTTYCCAGGTATC (SEQ ID NO: 123)
VK10	CTGTTTCAAGGTRCCAGATGT (SEQ ID NO: 124)
VK11	GTTGTAATGTCCAGAGGA (SEQ ID NO: 125)
VK12/13-1	CTTACAGGTGCCAGATGT (SEQ ID NO: 126)
VK12/13-2	CTCAATTGTAGRTGCCAGATGT (SEQ ID NO: 127)
VK12/13-3	CACAGTAGGTGTCAGATGT (SEQ ID NO: 128)
VK12/13-4	GTCGTAGTTGTCAGATGT (SEQ ID NO: 129)
VK12/13-5	CCTCCTTCTGGCCAAGA (SEQ ID NO: 130)
VK19/28-1	CTTATATGGAGCTGATGGG (SEQ ID NO: 131)

VK19/28-2	GTGTCTGGTGCTCATGGG (SEQ ID NO: 132)
VK19/28-3	CTSTGGTTGTCTGGTGTGA (SEQ ID NO: 133)
VK20	GTCTCTGATTCTAGGGCA (SEQ ID NO: 134)
VK21-1	CTKCKCTGGGTTCCAG (SEQ ID NO: 135)
VK21-2	GCAGGTGTTGACGGA (SEQ ID NO: 136)
VK22-1	CAGGTGCCTCGTGCAC (SEQ ID NO: 137)
VK22-2	CTCTGGTGCGCTGTGCA (SEQ ID NO: 138)
VK23	CTGGAYTYCAGCCTCCAGA (SEQ ID NO: 139)
VK24/25-1	GWTCTCTRGAGTCAGTGGG (SEQ ID NO: 140)
VK24/25-2	CTGGATCCCTGGAKCYACT (SEQ ID NO: 141)
VK32	GTTCTGCTTTTAGGTGTG (SEQ ID NO: 142)
VK33/34	GATCCCAGGCATGATATGT (SEQ ID NO: 143)
VK31/38C	CTTCATGGTGCTCAGTGT (SEQ ID NO: 144)
VKRF	CCATATCAGGTGCCAGTGT (SEQ ID NO: 145)

[00254] The reverse primer (murine IgKappa constant) is provided below.

Name	Sequence
LC -rev	GCGTCTAGAACTGGATGGTGGGAAGATGG (SEQ ID NO: 146)

[00255] Amplified VH and VL were subsequently cloned in-frame into vectors containing appropriate antibody secretion sequences and human IgG1 and Igkappa constants regions, respectively, to generate murine:human chimeric DNA constructs. These constructs were co-transfected into 293Freestyle cells (Life Technologies) and the resultant antibody was purified from the cell culture supernatant by Protein-A chromatography. To determine that the correct VH and VL sequences had been identified, the chimeric 2A1 (denoted 2A1-xi) was compared to the murine parental 2A1 antibody and CD47 binding assay by flow cytometry on Raji cells (Figure 6). B6H12 was also included as a positive control in this assay. Bound 2A1-xi was detected using a FITC-conjugated anti-human IgG secondary antibody. Bound 2A1 and B6H12 were detected using a FITC-conjugated anti-mouse IgG secondary antibody. Apparent affinities were determined by non-linear fits (Prism Graphpad Software) of the median fluorescence intensities at various antibody concentrations (Table 2). The 2A1-xi antibody has a similar binding affinity as the murine 2A1 antibody toward cell surface CD47, demonstrating that the VH and VL sequences had been properly identified.

Table 2.

	KD(apparent) (pM)	Std Error	R ²
2A1-mIgG1	93.6	±10.1	0.9977
2A1-xi	78	±14.9	0.9922
B6H12	3786	±310	0.9998

EXAMPLE 8: Antibody Humanization

[00256] The murine 2A1 CD47 antibody was humanized to reduce the potential of immunogenicity when administered to human patient. The sequences of the VH and VL region of 2A1 were compared to human antibody sequences in the IMGT databank. Subsequently, a structural model was generated of the 2A1 VH and VL regions using the known structures of the most closely related humanized and human antibodies in the Protein Data Bank (PDB). The 3 complementary determining regions (CDR) in both the heavy and light chains of the 2A1 antibody were fixed and the murine frameworks were replaced with numerous human frameworks that had the highest possibility of maintaining the proper orientation of the CDRs. Constructs corresponding to each the humanized 2A1 variants were generated by gene synthesis and cloned in frame into vectors containing an appropriate secretion sequence and human IgG1 and Igkappa constant regions. Various combinations of humanized heavy and light chains were co-transfected in to 293Freestyle cells (Life Technologies), and resultant antibodies were purified from the cell culture supernatant by Protein-A chromatography.

[00257] Humanized antibodies were tested for their ability to bind Raji cells by flow cytometry (Figure 7). The 2A1-xi antibody was used as a control in most of these assays to set the benchmark for binding affinity. Humanized antibodies were further optimized to enhance expression and reduce problematic sites including potential isomerization and deamidation sites. An example of an optimized humanized antibody derived from the murine 2A1 antibody is denoted as AB6.12 antibody, which displays very similar binding affinity as the 2A1-xi antibody (Figure 7H; Table 3). Apparent affinities were determined by non-linear fits (Prism Graphpad Software) of the median fluorescence intensities at various antibody concentrations.

Table 3.

	KD(apparent) (pM)	Std Error	R ²
2A1-xi	36.4	±8.54	0.9908
AB6.12	39.9	±5.54	0.9964

[00258] The AB6.12 antibody was subsequently converted from an IgG1 to other human IgG isotypes by replacing the constant domain of the heavy chain. As shown in Figure 7I, changing the IgG isotype to a hinge stabilized version of IgG4 (IgG4P: S228P),

and reduced Fc receptor binding variant of the hinge stabilized IgG4 (IgG4PE: S228P/L235E) did not alter binding affinity of the humanized antibody toward cells surface CD47 (Figure 7I; Table 4). Apparent affinities were determined by non-linear fits (Prism Graphpad Software) of the median fluorescence intensities at various antibody concentrations.

Table 4.

	KD(apparent) (pM)	Std Error	R ²
AB6.12-IgG1	38.6	±10.5	0.9798
AB6.12-IgG4P	35.7	±8.4	0.9841
AB6.12-IgG4PE	34.6	±10.9	0.9727

[00259] Throughout the humanization process, the CD47 antibodies were tested to ensure the SIRP α blocking functionality was intact. As shown in Figure 7J, multiple IgG isotypes of the humanized antibody, AB6.12, blocked the SIRP α :CD47 interaction, using the flow cytometry-based method described above in Example 3. Exemplary CD47 antibodies and their corresponding VH region and VL region include those provided in Table 1.

[00260] During the humanization process, it was determined that in some embodiments, an amino acid sequence motif, “NA,” at the beginning of VH CDR3 (SEQ ID NO: 52 or SEQ ID NO: 77) is important for binding of the CD47 antibodies described herein. In some embodiments, in the absence of amino acid residues “NA” at the beginning of VH CDR3 (SEQ ID NO: 52 or SEQ ID NO: 77), the CD47 antibodies of the invention do not bind to their target or bind to their target with lower affinity than they would in the presence of amino acid residues “NA.” For example, when the “NA” motif was changed to more canonical motifs of “AR” or “AT,” binding was substantially reduced (i.e., greater than ten-fold). In other embodiments, in the absence of amino acid residues “NA” at the beginning of VH CDR3 (SEQ ID NO: 52 or SEQ ID NO: 77), the CD47 antibodies of the invention bind to their target with equivalent affinity compared to binding in the presence of amino acid residues “NA.”

[00261] Those skilled in the art will recognize that it is possible to determine, without undue experimentation, if an amino acid substitution in the sequences of the CD47 antibodies of the invention will result in an antibody with substantially the same function, *e.g.*, a CD47 antibody with the ability to block SIRP α and not cause a significant level of hemagglutination.

[00262] An image of the trace from size exclusion chromatography using an AKTA FLPC with a superdex200 column is shown in Figure 8A. The IgG1, IgG4P, and IgG4PE variants of the AB6.12 antibody are shown. All three variant are over 98% monomeric. Figure 8B is a photograph of a coomassie blue stained SDS-PAGE gel of numerous humanized variants of 2A1 under reducing (R) and non-reducing (NR) conditions.

EXAMPLE 9: CD47 Antibodies Promote Phagocytosis of Tumor Cell Lines

[00263] CD47 is a cell surface receptor that is upregulated on tumor cells and is also thought to contribute to immune evasion through its interaction with its natural ligand SIRP α . Ligation of CD47 to SIRP α on macrophages results in decreased phagocytic activity. As described in detail below, it was determined if the CD47 binding and SIRP α blocking activity of the 2A1 antibody, and variations thereof, promote tumor cell phagocytosis in the presence of human macrophages.

[00264] PBMCs were isolated from human blood, and the monocytes were differentiated into macrophages by incubating them in AIM-V media (Life Technologies) for 7 days. These monocyte derived macrophages (MDMs) become adherent allowing other cells to be washed away. MDMs were scraped and re-plated in 12-well dishes and allowed to adhere for 24 hours. The human tumor cell line CCRF-CEM was chosen as a target cell type because of its high CD47 expression. CCRF-CEM cells were labeled with 0.3 μ M CFSE at 37°C for 15 minutes, then washed and added to MDMs at a ratio of 4:1 tumor cells per phagocyte, and CD47 antibody was added at various concentrations. Phagocytosis of target cells was allowed for 3 hours. Subsequently, non-phagocytosed target cells were washed away with PBS. The remaining phagocytes were scraped off, stained with an antibody to the macrophage marker CD14 conjugated to DyLite 649 (Biolegend), and analyzed by flow cytometry. Phagocytosis was measured by gating on live cells that were FL4 positive (CD14+), and then assessing the percent of FL1 (CFSE+) positive cells.

[00265] Figure 9 shows that the CD47 antibody 2A1 and its humanized variants demonstrated a dose-dependent increase in phagocytosis of tumor cells by MDMs. Antibody 2A1 and the humanized variant AB2.05 were unique in their ability to induce phagocytosis of tumor cells at 66.7 pM, whereas B6H12 had no activity at that concentration (Figure 9A). Figure 9B shows how 2A1, and the humanized variants AB2.05, AB6.12-IgG1, AB6.12-IgG4P, and AB6.12-IgG4PE all induce maximal phagocytosis at 0.3 μ g/ml or 2 nM, while B6H12 requires higher concentrations. This data demonstrates that the CD47 antibody, 2A1 (and humanized variants derived from it), induce macrophage-

mediated phagocytosis of CD47 positive tumor cells. In this example, CCFR-CEM cells were utilized as the CD47 positive target cell.

Example 10: Antitumor Activity of CD47 Antibodies

[00266] The anti-tumor activity of the murine CD47 antibodies was evaluated in a Raji model of lymphoma. Raji cells were implanted subcutaneously in NOD/SCID mice, and randomized into 5 groups (10 mice per group, day 0). Group 1: Vehicle (buffer only); Group 2: B6H12 (positive control); Group 3: 1B4; Group 4: 2A1; and Group 5: 9E4. Treatment with each antibody or vehicle (buffer only) began when tumors were palpable (50 mm³, day 13) and mice were euthanized when their tumor volumes reached ~1500 mm³. Tumor volumes were measured 3 times per week. Antibodies were dosed intravenously (IV) with 200 μ g 3 times per week for 3 weeks (9 total doses per mouse). Treatment started on day 13 and ended on day 32.

[00267] As shown in Figure 10A, CD47 antibodies of the invention, particularly the 2A1 antibody, demonstrated anti-tumor activity in this animal model of lymphoma. To reach a tumor volume of 1500 mm³, Group 1 (vehicle only) required ~25 days; Group 2 (B6H12.2) required ~45 days; Group 3 (1B4) required ~37 days; Group 4 (2A1) required ~85 days; and Group 5 (9E4) required ~40 days to reach a tumor volume ~1500 mm³. These data indicate that antibody 2A1 was significantly more potent than all CD47 binding antibodies tested, including B6H12 that was known to bind CD47, block CD47 interaction with SIRP α , and suppress tumor formation in mouse models of human cancer.

Unexpectedly, tumor suppression activity of these CD47 antibodies did not correlate with their potency of binding CD47 or blocking CD47 interaction with SIRP α , which would be expected based upon published data.

[00268] As described in Examples 2 and 3, 2A1, 1B4, and 9E4 had similar affinity for CD47 and similar potency for blocking CD47 interaction with SIRP α . In addition, the enhanced efficacy of 2A1 cannot be explained by differences in the Fc domain of the antibodies described since all antibodies used in this study were comprised of identical mouse IgG1 domains. Thus, in addition to unique composition of matter, the 2A1 antibody possesses unexpected and unique characteristics including the inability to induce homotypic interactions between CD47 expressing cells, *e.g.*, red blood cells, and enhanced tumor suppression activity that cannot be explained by enhanced binding to CD47 or an enhanced ability to block CD47 interaction with SIRP α .

[00269] To confirm that the humanized 2A1 antibodies maintained their anti-tumor activity, a similar Raji tumor study was conducted. The study design was the same as described above. Raji cells were implanted subcutaneously in NOD/SCID mice and randomized into 5 groups (10 mice per group, day 0). In this study, antibodies were dosed intraperitoneal (IP) with 200 μ g 3 times per week for 3 weeks (9 total doses per mouse), and tumor volumes were measured 3 times per week. However, for this study, the mouse IgG1 2A1 antibody (group 2) was compared to a humanized derivative, AB6.12. For this study, AB6.12 was constructed (as described in EXAMPLE 8) into human IgG1 (Group 3), human IgG4P (Group 4) and human IgG4PE (Group 4). Thus, this experiment was designed to address the influence of 2A1 humanization on its tumor suppression activity and the potential role of Fc domain effector function that is known in the art to contribute to the antitumor activities of many antibodies. It has been well documented that human IgG1 possesses significantly more effector function compared to human IgG4P. IgG4PE was developed to further reduce effector function. As can be seen in Figure 10B, humanization of 2A1 did not diminish the antitumor activity of 2A1, and in fact may have enhanced it. AB6.12-hIgG1, AB6.12-hIgG4P, and AB6.12-hIgG4PE all showed similar anti-tumor activity that appears significantly greater than mouse 2A1 (2A1mIgG). This result is unexpected since 2A1mIgG1, AB6.12-hIgG1, AB6.12-hIgG4P and AB6.12-hIgG4PE have similar CD47 binding and SIRP α blocking activities. In addition, since AB6.12-hIgG1, AB6.12-hIgG4P and AB6.12-hIgG4PE have similar anti-tumor activities, it appears that effector function does play a role in the efficacy of the humanized 2A1 antibody AB6.12.

Example 11: Co-crystallization of CD47 antibodies with CD47

[00270] CD47 is 5 pass transmembrane protein with a single extracellular IgV (immunoglobulin-like variable-type) domain that is highly glycosylated at 6 sites. The structure of the CD47-IgV domain has been solved in complex with the IgV domain of SIRP α , its natural ligand (Protein Data Bank (PDB) Reference No. 2JJS; Hatherley et al., 2008 Mol Cell, 25;31(2): 266-77 (Figure 11A)). The structure shows SIRP α -IgV binding to CD47-IgV on an apical epitope including the N-terminal pyroglutamate of CD47. This structure sufficiently explains how both cell surface transmembrane proteins can productively interact from adjacent cells in a head to head orientation. The X-ray crystallographic structure of CD47-IgV in complex with the B6H12 Fab is presented in Figure 11B. For clarity, the constant regions of the Fab (CH1 and CL) were omitted in the Figure, and only the Fv (VH and VL) is presented. This revealed an apical binding site,

positioning this antibody on a surface extremely distal from the cell membrane (Figure 11B). The mechanism of SIRP α blocking by B6H12 is apparent from this structure. The orientation purposes relative location of the cell membrane is depicted as a dashed line in Figure 11.

[00271] In order to determine the target epitope of the antibodies of the present invention, the X-ray crystallographic structure of the co-complex of CD47-IgV domain and the Fab of 2A1-xi (chimeric antibody with human CH1 and CL domains) was determined (Figure 11C). For clarity, the constant regions of the Fab (CH1 and CL) were omitted in the Figure, and only the Fv (VH and VL) is presented. Unlike the previously determined structure of CD47 binding SIRP α in a head to head orientation (Figure 11A), and the B6H12 antibody being positioned apically away from the membrane (Figure 11B), the structure of 2A1 in complex with CD47 revealed binding of the antibody to CD47 near the membrane in an unexpected and unique head to side orientation (Figure 11C). The 2A1 epitope on CD47 is discontinuous, and includes residues Y37, K39, K41, the KGRD (SEQ ID NO: 56) loop (residues 43-46), D51, H90, N93, E97, T99, E104, and E106 of CD47 when numbered in accordance with SEQ ID NO: 147 (*i.e.*, SEQ ID NO: 48 excluding the signal sequence (amino acids 1-18)). The structure of 2A1 bound to CD47 also reveals that the VH is primarily involved in binding to the KGRD (SEQ ID NO: 56) loop of CD47, while the VK domain interacts with apical residues including Y37, T102, and E104, which are involved in SIRP α binding. Therefore, it is primarily the VK domain that physically precludes SIRP α binding to CD47. These structural studies suggest that the unique epitope which 2A1 binds to is on the side of CD47. In contrast to CD47 antibodies known in the art, the orientation of the 2A1 VH region in a membrane proximal position are critical features of this antibody that prevent a significant level of red blood cell hemagglutination by constraining the antibody such that it cannot bridge to CD47 molecules on adjacent cells.

Claims

1. An isolated monoclonal antibody or immunologically active fragment thereof that binds CD47, wherein the antibody or immunologically active fragment thereof comprises a VH complementarity determining region 1 (CDR1) sequence set forth in SEQ ID NO: 50, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 64, SEQ ID NO: 65, or SEQ ID NO: 66, a VH CDR2 sequence set forth in SEQ ID NO: 51, SEQ ID NO: 72, SEQ ID NO: 73, SEQ ID NO: 74, SEQ ID NO: 75, or SEQ ID NO: 76, a VH CDR3 sequence set forth in SEQ ID NO: 52 or SEQ ID NO: 77, a VL CDR1 sequence set forth in SEQ ID NO: 53, SEQ ID NO: 67, or SEQ ID NO: 68, a VL CDR2 sequence set forth in SEQ ID NO: 54, SEQ ID NO: 69, SEQ ID NO: 70, or SEQ ID NO: 71 and a VL CDR3 sequence set forth in SEQ ID NO: 55.
2. The antibody or immunologically active fragment thereof of claim 1, wherein the antibody or immunologically active fragment thereof comprises a VH CDR1 sequence set forth in SEQ ID NO: 50, a VH CDR2 sequence set forth in SEQ ID NO: 51, a VH CDR3 sequence set forth in SEQ ID NO: 52, a VL CDR1 sequence set forth in SEQ ID NO: 53, a VL CDR2 sequence set forth in SEQ ID NO: 54, and a VL CDR3 sequence set forth in SEQ ID NO: 55.
3. The antibody or immunologically active fragment thereof of claim 1, wherein the antibody or immunologically active fragment thereof comprises a VH CDR1 sequence set forth in SEQ ID NO: 50, a VH CDR2 sequence set forth in SEQ ID NO: 72, a VH CDR3 sequence set forth in SEQ ID NO: 52, a VL CDR1 set forth in SEQ ID NO: 53, a VL CDR2 sequence set forth in SEQ ID NO: 71, and a VL CDR3 sequence set forth in SEQ ID NO: 55.
4. An isolated monoclonal antibody or immunologically active fragment thereof that binds to human CD47, said antibody or immunologically active fragment thereof comprising:
 - a VH CDR1, a VH CDR2, and a VH CDR3, respectively comprising the amino acid sequences of a CDR1, a CDR2, and a CDR3 within a VH chain region having the sequence set forth in any of SEQ ID NOs: 5-30; and

a VL CDR1, a VL CDR2, and a VL CDR3, respectively comprising the amino acid sequences of a CDR1, a CDR2, and a CDR3 within a VL chain region having the sequence set forth in any of SEQ ID NOs: 31-47.

5. The antibody or immunologically active fragment thereof of any one of claims 1-4, wherein the antibody or immunologically active fragment thereof comprises a variable heavy (VH) chain region selected from the group consisting of SEQ ID NOs: 5-30.

6. The antibody or immunologically active fragment thereof of any one of claims 1-5, wherein the antibody or immunologically active fragment thereof comprises a variable light (VL) chain region selected from the group consisting of SEQ ID NOs: 31-47.

7. The antibody or immunologically active fragment thereof of any one of claims 1-6, wherein the antibody or immunologically active fragment thereof comprises a VH region provided in any one of SEQ ID NOs: 5-30 and a VL region provided in any one of SEQ ID NOs: 31-47.

8. The antibody or immunologically active fragment thereof of claim 8, wherein the antibody or immunologically active fragment thereof comprises a VH region provided in any one of SEQ ID NOs: 5, 7, 8, 11, 12, 15-17, 20-22, and 27-30 paired with a VL region provided in any one of SEQ ID NOs: 31, 32, 35, 40, 41, 42, 43, 44, and 47.

9. The antibody or immunologically active fragment thereof of claim 8, wherein the antibody comprises a combination of a VH chain region and a VL chain region selected from the combinations listed in Table 1:

Antibody	Variable heavy (VH) chain	Variable light (VL) chain
2A1	SEQ ID NO: 5	SEQ ID NO: 31
2A1-xi	SEQ ID NO: 5	SEQ ID NO: 32
AB2.03	SEQ ID NO: 7	SEQ ID NO: 33
AB2.04	SEQ ID NO: 7	SEQ ID NO: 34
AB2.05	SEQ ID NO: 7	SEQ ID NO: 35
AB2.06	SEQ ID NO: 7	SEQ ID NO: 36

AB2.07	SEQ ID NO: 7	SEQ ID NO: 37
AB2.08	SEQ ID NO: 7	SEQ ID NO: 38
AB2.09	SEQ ID NO: 7	SEQ ID NO: 39
AB2.13	SEQ ID NO: 7	SEQ ID NO: 43
AB3.09	SEQ ID NO: 8	SEQ ID NO: 39
AB6.12	SEQ ID NO: 11	SEQ ID NO: 42
AB6.13	SEQ ID NO: 11	SEQ ID NO: 43
AB6.14	SEQ ID NO: 11	SEQ ID NO: 44
AB6.17	SEQ ID NO: 11	SEQ ID NO: 47
AB10.13	SEQ ID NO: 15	SEQ ID NO: 43
AB10.14	SEQ ID NO: 15	SEQ ID NO: 44
AB11.05	SEQ ID NO: 16	SEQ ID NO: 35
AB12.05	SEQ ID NO: 17	SEQ ID NO: 35
AB15.05	SEQ ID NO: 20	SEQ ID NO: 35
AB16.05	SEQ ID NO: 21	SEQ ID NO: 35
AB17.05	SEQ ID NO: 22	SEQ ID NO: 35
AB22.05	SEQ ID NO: 27	SEQ ID NO: 35
AB23.05	SEQ ID NO: 28	SEQ ID NO: 35
AB24.05	SEQ ID NO: 29	SEQ ID NO: 35
AB25.05	SEQ ID NO: 30	SEQ ID NO: 35

10. The isolated monoclonal antibody or immunologically active fragment thereof of any one of claims 1-9, wherein the antibody does not cause a significant level of agglutination of cells after administration.


11. The antibody or immunologically active fragment thereof of any one of claims 1-10, wherein the antibody is chimeric, humanized, or fully human.

12. The antibody or immunologically active fragment thereof of any one of claims 1-11, wherein the CD47 is human CD47.

13. The antibody or immunologically active fragment thereof of any one of claims 1-12, wherein the antibody or immunologically active fragment thereof prevents CD47 from interacting with signal-regulatory-protein (SIRP).
14. The antibody or immunologically active fragment thereof of any one of claims 1-13, wherein the antibody or immunologically active fragment thereof promotes macrophage-mediated phagocytosis of a CD47-expressing cell.
15. The antibody or immunologically active fragment thereof of any one of claims 1-14, wherein the antibody or immunologically active fragment thereof is an IgG isotype selected from the group consisting of IgG1 isotype, IgG2 isotype, IgG3 isotype, and IgG4 isotype.
16. The antibody or immunologically active fragment thereof of any one of claims 1-15, wherein the antibody binds to CD47 in a head to side orientation, wherein a VH chain of the antibody is positioned near the membrane of a CD47 expressing cell, and wherein a VL chain of the antibody occludes a SIRP binding site on CD47.
17. The antibody or immunologically active fragment thereof of any one of claims 1-15, wherein the antibody binds to CD47 in a head to side orientation, wherein a VL chain of the antibody is positioned near the membrane of a CD47 expressing cell, and wherein a VH chain of the antibody occludes a SIRP binding site on CD47.
18. The antibody or immunologically active fragment thereof of any one of claims 1-17, wherein the antibody or immunologically active fragment thereof binds to a discontinuous epitope on CD47.
19. The antibody or immunologically active fragment thereof of claim 18, wherein the antibody or immunologically active fragment thereof binds to a CD47 loop comprising SEQ ID NO: 56.

20. The antibody or immunologically active fragment thereof of claim 18, wherein the discontinuous epitope comprises amino acids residues Y37, K39, K41, K43, G44, R45, D46, D51, H90, N93, E97, T99, E104, or E106 of CD47 when numbered in accordance with SEQ ID NO: 147.
21. The antibody or immunologically active fragment thereof of any one of claims 1-20, wherein the antibody or immunologically active fragment thereof does not cause a significant level of hemagglutination of red blood cells after administration.
22. The antibody or immunologically active fragment thereof of any one of claims 1-21, wherein the antibody or immunologically active fragment thereof is an IgG isotype selected from IgG4P and IgG4PE.
23. A pharmaceutical composition comprising the antibody or immunologically active fragment thereof of any one of claims 1-22 and a carrier.
24. A method of alleviating a symptom of a cancer or other neoplastic condition, the method comprising administering the antibody or immunologically active fragment thereof of any one of claims 1-22 to a subject in need thereof in an amount sufficient to alleviate the symptom of the cancer or other neoplastic condition in the subject.
25. The method of claim 24, wherein the subject is a human.
26. The method of claim 24 or claim 25, further comprising administering chemotherapy.
27. The method of claim 26, wherein said chemotherapy is radiotherapy.
28. Isolated nucleic acid(s) encoding the antibody or immunologically active fragment thereof of any one of claims 1-22.
29. An isolated host cell comprising nucleic acid(s) encoding the antibody or immunologically active fragment thereof of any one of claims 1-22.

30. Use of the antibody or immunologically active fragment thereof of any one of claims 1-22 in the manufacture of a medicament for alleviating a symptom of a cancer or other neoplastic condition.






FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B






FIG. 4A

[Ab] (nM)	100	50	25	12.5	100	50	25	12.5	100	50	25	12.5
A	9E4	9E4	9E4	9E4	11B9	11B9	11B9	11B9	15B9	15B9	15B9	15B9
B	Blank	Blank	Blank	Blank	10F4	10F4	10F4	10F4	18F11	18F11	18F11	18F11
C	B6H12	B6H12	B6H12	B6H12	Blank	Blank	Blank	Blank	18B5	18B5	18B5	18B5
D	2A1	2A1	2A1	2A1	3B10	3B10	3B10	3B10	17F11	17F11	17F11	17F11
E	2A1-XI	2A1-XI	2A1-XI	2A1-XI	15B7	15B7	15B7	15B7	Blank	Blank	Blank	Blank
F	6B2	6B2	6B2	6B2	2B11	2B11	2B11	2B11	Blank	Blank	Blank	Blank
G	14E10	14E10	14E10	14E10	5B11	5B11	5B11	5B11	Blank	Blank	Blank	Blank
H	Blank	Blank	Blank	Blank	Blank	Blank	Blank	Blank	Blank	Blank	Blank	Blank
	1	2	3	4	5	6	7	8	9	10	11	12



2A1 →
2A1-XI →

FIG. 4B




FIG. 4C





FIG. 4D




FIG. 4E




FIG. 4F




FIG. 5




FIG. 6

FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D






FIG. 7E




FIG. 7F




FIG. 7G





FIG. 7H



12/19




FIG. 7I





FIG. 7J

FIG. 8A






FIG. 8B







FIG. 9A

FIG. 9B
human MDM phagocytosis of CCRF-CEM cells

17/19

FIG. 10A
Raji Tumor Model: Murine CD47 Antibodies








FIG. 11A

FIG. 11B

FIG. 11C

31 Jul 2015

2013217114

5641102_1
SEQUENCE LISTING

<110> InhibRx LLC

<120> CD47 Antibodies and Methods of Use Thereof

<130> 42967-506001WO

<140> PCT/US2013/024995

<141> 2013-02-06

<150> US 61/595,216

<151> 2012-02-06

<150> US 61/659,752

<151> 2012-06-14

<160> 147

<170> PatentIn version 3.5

<210> 1

<211> 330

<212> PRT

<213> Homo sapiens

<400> 1

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys

1

5

10

15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

20

25

30

31 Jul 2015

2013217114

5641102_1

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100 105 110

Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
115 120 125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130 135 140

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
145 150 155 160

2013217114 31 Jul 2015

5641102_1
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165 170 175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180 185 190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
195 200 205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
210 215 220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu
225 230 235 240

Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245 250 255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
260 265 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290 295 300

31 Jul 2015

5641102_1

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305 310 315 320

2013217114

Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
325 330

<210> 2
<211> 326
<212> PRT
<213> Homo sapiens

<400> 2

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15

Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr

31 Jul 2015

2013217114

5641102_1

65 70 75 80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys

85 90 95

Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro

100 105 110

Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp

115 120 125

Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp

130 135 140

Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly

145 150 155 160

Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn

165 170 175

Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp

180 185 190

Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro

195 200 205

2013217114 31 Jul 2015

5641102_1

Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu
210 215 220

Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
225 230 235 240

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
245 250 255

Ser Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
260 265 270

Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
275 280 285

Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
290 295 300

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
305 310 315 320

Ser Leu Ser Pro Gly Lys
325

<210> 3

31 Jul 2015

5641102_1

<211> 377

<212> PRT

<213> Homo sapiens

<400> 3

2013217114

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg

1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser

35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr

65 70 75 80

Tyr Thr Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

85 90 95

Arg Val Glu Leu Lys Thr Pro Leu Gly Asp Thr Thr His Thr Cys Pro

100 105 110

2013217114 31 Jul 2015

5641102_1
Arg Cys Pro Glu Pro Lys Ser Cys Asp Thr Pro Pro Pro Cys Pro Arg
115 120 125

Cys Pro Glu Pro Lys Ser Cys Asp Thr Pro Pro Pro Cys Pro Arg Cys
130 135 140

Pro Glu Pro Lys Ser Cys Asp Thr Pro Pro Pro Cys Pro Arg Cys Pro
145 150 155 160

Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
165 170 175

Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
180 185 190

Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Lys Trp Tyr
195 200 205

Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
210 215 220

Gln Tyr Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Leu His
225 230 235 240

Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
245 250 255

31 Jul 2015

2013217114

5641102_1

Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln
260 265 270

Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
275 280 285

Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
290 295 300

Ser Asp Ile Ala Val Glu Trp Glu Ser Ser Gly Gln Pro Glu Asn Asn
305 310 315 320

Tyr Asn Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu
325 330 335

Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Ile
340 345 350

Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln
355 360 365

Lys Ser Leu Ser Leu Ser Pro Gly Lys
370 375

2013217114 31 Jul 2015

5641102_1

<210> 4
<211> 327
<212> PRT
<213> Homo sapiens

<400> 4

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg

1 5 10 15

Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser

35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr

65 70 75 80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys

85 90 95

Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Ser Cys Pro Ala Pro

100 105 110

2013217114 31 Jul 2015

5641102_1

Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
115 120 125

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
130 135 140

Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
145 150 155 160

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
165 170 175

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
180 185 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
195 200 205

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
210 215 220

Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
225 230 235 240

Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
250 255 260 265

31 Jul 2015

2013217114

5641102_1
245 250 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
260 265 270

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
275 280 285

Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
290 295 300

Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
305 310 315 320

Leu Ser Leu Ser Leu Gly Lys
325

<210> 5

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 5

Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Ser Gly Ala

31 Jul 2015
2013217114

5641102_1

1

5

10

15

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Tyr

20

25

30

Tyr Leu His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile

35

40

45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Phe Ala Pro Lys Phe

50

55

60

Gln Gly Lys Ala Thr Met Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr

65

70

75

80

Leu Gln Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys

85

90

95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100

105

110

Gly Thr Ser Val Thr Val

115

<210> 6

<211> 118

<212> PRT

2013217114 31 Jul 2015

5641102_1

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 6

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala

1

5

10

15

Thr Val Lys Ile Ser Cys Lys Val Ser Gly Phe Asn Ile Lys Asp Tyr

20

25

30

Tyr Leu His Trp Val Gln Gln Ala Pro Gly Lys Gly Leu Glu Trp Met

35

40

45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Glu Lys Phe

50

55

60

Gln Gly Arg Val Thr Ile Thr Ala Asp Thr Ser Thr Asp Thr Ala Tyr

65

70

75

80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys

85

90

95

Asn Ala Ala Tyr Gly Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100

105

110

2013217114 31 Jul 2015

5641102_1

Gly Thr Thr Val Thr Val

115

<210> 7

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 7

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1

5

10

15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr

20

25

30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35

40

45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50

55

60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65

70

75

80

2013217114 31 Jul 2015

5641102_1

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 8
<211> 118
<212> PRT
<213> Artificial Sequence

<220>
<223> variable heavy chain region of a humanized CD47 antibody

<400> 8

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

2013217114 31 Jul 2015

5641102_1

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 9

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 9

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

2013217114 31 Jul 2015

5641102_1

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

Gln Gly Arg Val Thr Met Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 10
<211> 118
<212> PRT
<213> Artificial Sequence

2013217114 31 Jul 2015

5641102_1

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 10

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Gly Arg Val Thr Met Thr Glu Asp Thr Ser Thr Asp Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

2013217114 31 Jul 2015

5641102_1

115

<210> 11
<211> 118
<212> PRT
<213> Artificial Sequence

<220>
<223> Variable heavy chain region of a humanized CD47 antibody

<400> 11

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

Gly Trp Ile Asp Pro Asp Gln Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

31 Jul 2015

2013217114

5641102_1
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

<210> 12

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 12

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Tyr Gly Asp Thr Glu Tyr Ala Gln Lys Phe

2013217114
31 Jul 2015

50 55 5641102_1
60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

<210> 13

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 13

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr

31 Jul 2015
2013217114

20 25 5641102_1 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Ser Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

<210> 14

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

2013217114
31 Jul 2015

5641102_1

<400> 14

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Asn Ala Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

31 Jul 2015

5641102_1

<210> 15
<211> 118
<212> PRT
<213> Artificial Sequence

2013217114

<220>
<223> variable heavy chain region of a humanized CD47 antibody
<400> 15

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

Gly Trp Ile Asp Pro Asp Asn Thr Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

2013217114 31 Jul 2015

5641102_1

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 16
<211> 118
<212> PRT
<213> Artificial Sequence

<220>
<223> Variable heavy chain region of a humanized CD47 antibody

<400> 16

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

2013217114 31 Jul 2015

5641102_1

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Pro Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 17

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 17

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Tyr Tyr
20 25 30

2013217114 31 Jul 2015

5641102_1

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 18

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 18

2013217114 31 Jul 2015

5641102_1

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Tyr Tyr
20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 19

2013217114 31 Jul 2015

5641102_1

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 19

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Asn Phe Thr Tyr Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

31 Jul 2015

2013217114

100 105 5641102_1 110

Gly Thr Thr Val Thr Val

115

<210> 20

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 20

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Ile Thr Tyr Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

31 Jul 2015

2013217114

5641102_1
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

<210> 21

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 21

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Lys Tyr Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

31 Jul 2015
2013217114

5641102_1
35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

<210> 22

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 22

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

31 Jul 2015
2013217114

5641102_1
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30
Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45
Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60
Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95
Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110
Gly Thr Thr Val Thr Val
115
<210> 23
<211> 118
<212> PRT

2013217114 31 Jul 2015

5641102_1

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 23

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Asp Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

2013217114 31 Jul 2015

5641102_1

Gly Thr Thr Val Thr Val

115

<210> 24

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 24

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1

5

10

15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Ile Thr Asp Tyr

20

25

30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35

40

45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50

55

60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65

70

75

80

2013217114 31 Jul 2015

5641102_1

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 25
<211> 118
<212> PRT
<213> Artificial Sequence

<220>
<223> variable heavy chain region of a humanized CD47 antibody

<400> 25

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Lys Asp Tyr
20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

2013217114 31 Jul 2015

5641102_1

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 26

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 26

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser
1 5 10 15

2013217114 31 Jul 2015

5641102_1

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Lys Asp Tyr
20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met
35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe
50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val
115

<210> 27
<211> 118
<212> PRT
<213> Artificial Sequence

2013217114 31 Jul 2015

5641102_1

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 27

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Leu Gln Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

2013217114 31 Jul 2015

5641102_1

115

<210> 28

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable heavy chain region of a humanized CD47 antibody

<400> 28

Gln Met Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Thr Gly Ser

1

5

10

15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Asn Ile Lys Asp Tyr

20

25

30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35

40

45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50

55

60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65

70

75

80

Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys

31 Jul 2015

2013217114

5641102_1
85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

<210> 29

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 29

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala

1 5 10 15

Thr Val Lys Ile Ser Cys Lys Val Ser Gly Phe Asn Ile Lys Asp Tyr

20 25 30

Tyr Leu His Trp Val Arg Gln Ala Pro Gly Gln Ala Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

2013217114
31 Jul 2015

50

55

5641102_1
60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

<210> 30

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> variable heavy chain region of a humanized CD47 antibody

<400> 30

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala

1 5 10 15

Thr Val Lys Ile Ser Cys Lys Val Ser Gly Phe Asn Ile Lys Asp Tyr

31 Jul 2015
2013217114

20 25 5641102_1 30

Tyr Leu His Trp Val Gln Gln Ala Pro Gly Lys Gly Leu Glu Trp Met

35 40 45

Gly Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Arg Val Thr Ile Thr Arg Asp Arg Ser Met Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr Trp Gly Gln

100 105 110

Gly Thr Thr Val Thr Val

115

<210> 31

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

31 Jul 2015
2013217114

5641102_1

<400> 31

Asp Ile Lys Met Thr Gln Ser Pro Ser Ser Leu Tyr Ala Ser Leu Gly

1 5 10

15

Glu Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr

20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Ser Pro Lys Ile Leu Ile

35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Gln Asp Tyr Ser Leu Thr Ile Ser Ser Leu Glu Tyr

65 70 75 80

Glu Asp Met Gly Ile Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

85 90 95

Thr Phe Gly Gly Thr Lys Leu Glu Met Lys

100 105

<210> 32

<211> 107

<212> PRT

2013217114 31 Jul 2015

5641102_1

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 32

Asp Ile Lys Met Thr Gln Ser Pro Ser Ser Leu Tyr Ala Ser Leu Gly

1 5 10 15

Glu Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr

20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Ser Pro Lys Ile Leu Ile

35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Gln Asp Tyr Ser Leu Thr Ile Ser Ser Leu Glu Tyr

65 70 75 80

Glu Asp Met Gly Ile Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

85 90 95

Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys

100 105

31 Jul 2015

5641102_1

<210> 33
<211> 107
<212> PRT
<213> Artificial Sequence

2013217114

<220>
<223> variable light chain region of a humanized CD47 antibody
<400> 33

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr
20 25 30

Leu Ser Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr
85 90 95

2013217114 31 Jul 2015

5641102_1

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 34

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 34

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr
20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Ala Pro Lys Ser Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

2013217114 31 Jul 2015

5641102_1

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr
85 90 95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 35

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable light chain region of a humanized CD47 antibody

<400> 35

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr
20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys His Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

2013217114 31 Jul 2015

5641102_1

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr
85 90 95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 36

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 36

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr
20 25 30

Leu Ser Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile
35 40 45

2013217114 31 Jul 2015

5641102_1

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr
85 90 95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 37

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable light chain region of a humanized CD47 antibody

<400> 37

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile His Arg Tyr
20 25 30

2013217114 31 Jul 2015

5641102_1

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Val Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr
85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
100 105

<210> 38

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable light chain region of a humanized CD47 antibody

<400> 38

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

2013217114 31 Jul 2015

5641102_1

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Asp Ile His Arg Tyr
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr
85 90 95

Thr Gly Phe Gln Gly Thr Arg Leu Glu Ile Lys
100 105

<210> 39

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 39

2013217114 31 Jul 2015

5641102_1

Asp Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr
20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys His Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr
85 90 95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 40
<211> 107
<212> PRT
<213> Artificial Sequence

2013217114 31 Jul 2015

5641102_1

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 40

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Arg Gln Gly Ile His Arg Tyr

20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys His Leu Ile

35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

85 90 95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys

100 105

<210> 41

2013217114 31 Jul 2015

5641102_1

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 41

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr

20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys Ile Leu Ile

35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

85 90 95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys

2013217114 31 Jul 2015

100 105 5641102_1

<210> 42
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> Variable light chain region of a humanized CD47 antibody

<400> 42

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr
20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys His Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Leu Val Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

31 Jul 2015

2013217114

5641102_1
85 90 95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys

100 105

<210> 43

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable light chain region of a humanized CD47 antibody

<400> 43

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Arg Gln Gly Ile His Arg Tyr

20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys Ile Leu Ile

35 40 45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

31 Jul 2015

2013217114

65 70 75 80

5641102_1
Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr
85 90 95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 44

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 44

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Arg Gln Gly Ile His Arg Tyr
20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys His Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Leu Val Ser Gly Val Pro Ser Arg Phe Ser Gly

2013217114
31 Jul 2015

5641102_1
50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

85 90 95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys

100 105

<210> 45

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Variable light chain region of a humanized CD47 antibody

<400> 45

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr

20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Ile

31 Jul 2015

2013217114

5641102_1

35

40

45

Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly

50

55

60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65

70

75

80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

85

90

95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys

100

105

<210> 46

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 46

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly

1

5

10

15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile His Arg Tyr

31 Jul 2015
2013217114

20 25 5641102_1 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Ile

35 40 45

Tyr Arg Ala Asn Arg Leu Val Ser Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

85 90 95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys

100 105

<210> 47

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> variable light chain region of a humanized CD47 antibody

<400> 47

Asn Ile Gln Met Thr Gln Ser Pro Ser Ala Met Ser Ala Ser Val Gly

31 Jul 2015
2013217114

5641102_1

1

5

10

15

Asp Arg Val Thr Ile Thr Cys Arg Ala Arg Gln Gly Ile His Arg Tyr

20

25

30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Ile

35

40

45

Tyr Arg Ala Asn Arg Leu Val Ser Gly Val Pro Ser Arg Phe Ser Gly

50

55

60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65

70

75

80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Tyr

85

90

95

Thr Phe Gly Gly Thr Lys Val Glu Ile Lys

100

105

<210> 48

<211> 323

<212> PRT

<213> Homo sapiens

<400> 48

2013217114
31 Jul 2015

5641102_1
Met Trp Pro Leu Val Ala Ala Leu Leu Leu Gly Ser Ala Cys Cys Gly
1 5 10 15

Ser Ala Gln Leu Leu Phe Asn Lys Thr Lys Ser Val Glu Phe Thr Phe
20 25 30

Cys Asn Asp Thr Val Val Ile Pro Cys Phe Val Thr Asn Met Glu Ala
35 40 45

Gln Asn Thr Thr Glu Val Tyr Val Lys Trp Lys Phe Lys Gly Arg Asp
50 55 60

Ile Tyr Thr Phe Asp Gly Ala Leu Asn Lys Ser Thr Val Pro Thr Asp
65 70 75 80

Phe Ser Ser Ala Lys Ile Glu Val Ser Gln Leu Leu Lys Gly Asp Ala
85 90 95

Ser Leu Lys Met Asp Lys Ser Asp Ala Val Ser His Thr Gly Asn Tyr
100 105 110

Thr Cys Glu Val Thr Glu Leu Thr Arg Glu Gly Glu Thr Ile Ile Glu
115 120 125

Leu Lys Tyr Arg Val Val Ser Trp Phe Ser Pro Asn Glu Asn Ile Leu
130 135 140

31 Jul 2015

2013217114

5641102_1

Ile Val Ile Phe Pro Ile Phe Ala Ile Leu Leu Phe Trp Gly Gln Phe
145 150 155 160

Gly Ile Lys Thr Leu Lys Tyr Arg Ser Gly Gly Met Asp Glu Lys Thr
165 170 175

Ile Ala Leu Leu Val Ala Gly Leu Val Ile Thr Val Ile Val Ile Val
180 185 190

Gly Ala Ile Leu Phe Val Pro Gly Glu Tyr Ser Leu Lys Asn Ala Thr
195 200 205

Gly Leu Gly Leu Ile Val Thr Ser Thr Gly Ile Leu Ile Leu Leu His
210 215 220

Tyr Tyr Val Phe Ser Thr Ala Ile Gly Leu Thr Ser Phe Val Ile Ala
225 230 235 240

Ile Leu Val Ile Gln Val Ile Ala Tyr Ile Leu Ala Val Val Gly Leu
245 250 255

Ser Leu Cys Ile Ala Ala Cys Ile Pro Met His Gly Pro Leu Leu Ile
260 265 270

2013217114 31 Jul 2015

5641102_1
Ser Gly Leu Ser Ile Leu Ala Leu Ala Gln Leu Leu Gly Leu Val Tyr
275 280 285

Met Lys Phe Val Ala Ser Asn Gln Lys Thr Ile Gln Pro Pro Arg Lys
290 295 300

Ala Val Glu Glu Pro Leu Asn Ala Phe Lys Glu Ser Lys Gly Met Met
305 310 315 320

Asn Asp Glu

<210> 49

<211> 116

<212> PRT

<213> Homo sapiens

<400> 49

Gln Leu Leu Phe Asn Lys Thr Lys Ser Val Glu Phe Thr Phe Cys Asn
1 5 10 15

Asp Thr Val Val Ile Pro Cys Phe Val Thr Asn Met Glu Ala Gln Asn
20 25 30

Thr Thr Glu Val Tyr Val Lys Trp Lys Phe Lys Gly Arg Asp Ile Tyr
35 40 45

2013217114 31 Jul 2015

5641102_1

Thr Phe Asp Gly Ala Leu Asn Lys Ser Thr Val Pro Thr Asp Phe Ser
50 55 60

Ser Ala Lys Ile Glu Val Ser Gln Leu Leu Lys Gly Asp Ala Ser Leu
65 70 75 80

Lys Met Asp Lys Ser Asp Ala Val Ser His Thr Gly Asn Tyr Thr Cys
85 90 95

Glu Val Thr Glu Leu Thr Arg Glu Gly Glu Thr Ile Ile Glu Leu Lys
100 105 110

Tyr Arg Val Val
115

<210> 50

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 50

Gly Phe Asn Ile Lys Asp Tyr Tyr Leu His
1 5 10

31 Jul 2015

5641102_1

<210> 51
<211> 10
<212> PRT
<213> Artificial Sequence

2013217114

<220>
<223> amino acid sequence of VH CDR2

<400> 51

Trp Ile Asp Pro Asp Asn Gly Asp Thr Glu
1 5 10

<210> 52
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> amino acid sequence of VH CDR3

<400> 52

Asn Ala Ala Tyr Gly Ser Ser Ser Tyr Pro Met Asp Tyr
1 5 10

<210> 53
<211> 11
<212> PRT
<213> Artificial Sequence

2013217114 31 Jul 2015

5641102_1

<220>

<223> amino acid sequence of VL CDR1

<400> 53

Lys Ala Ser Gln Asp Ile His Arg Tyr Leu Ser

1 5 10

<210> 54

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VL CDR2

<400> 54

Arg Ala Asn Arg Leu Val Asp

1 5

<210> 55

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VL CDR3

<400> 55

Leu Gln Tyr Asp Glu Phe Pro Tyr Thr

2013217114 31 Jul 2015

5641102_1

1 5

<210> 56
<211> 4
<212> PRT
<213> Homo sapiens

<400> 56

Lys Gly Arg Asp

1

<210> 57
<211> 10
<212> PRT
<213> Artificial sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 57

Gly Tyr Thr Phe Thr Tyr Tyr Tyr Leu His

1 5 10

<210> 58
<211> 10
<212> PRT
<213> Artificial sequence

<220>

2013217114 31 Jul 2015

5641102_1

<223> amino acid sequence of VH CDR1

<400> 58

Gly Phe Thr Phe Thr Tyr Tyr Tyr Leu His

1 5 10

<210> 59

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 59

Gly Tyr Asn Phe Thr Tyr Tyr Tyr Leu His

1 5 10

<210> 60

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 60

Gly Tyr Thr Ile Thr Tyr Tyr Tyr Leu His

1 5 10

2013217114 31 Jul 2015

5641102_1

<210> 61
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> amino acid sequence of VH CDR1

<400> 61

Gly Tyr Thr Phe Lys Tyr Tyr Tyr Leu His
1 5 10

<210> 62
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> amino acid sequence of VH CDR1

<400> 62

Gly Tyr Thr Phe Thr Asp Tyr Tyr Tyr Leu His
1 5 10

<210> 63
<211> 10
<212> PRT
<213> Artificial Sequence

31 Jul 2015

5641102_1

<220>

<223> amino acid sequence of VH CDR1

<400> 63

2013217114

Gly Phe Thr Phe Thr Asp Tyr Tyr Leu His

1 5 10

<210> 64

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 64

Gly Phe Thr Ile Thr Asp Tyr Tyr Leu His

1 5 10

<210> 65

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 65

2013217114 31 Jul 2015

5641102_1

Gly Tyr Thr Phe Lys Asp Tyr Tyr Leu His

1 5 10

<210> 66

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 66

Gly Phe Thr Phe Lys Asp Tyr Tyr Leu His

1 5 10

<210> 67

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VL CDR1

<400> 67

Arg Ala Ser Gln Asp Ile His Arg Tyr Leu Ala

1 5 10

<210> 68

<211> 11

31 Jul 2015

5641102_1

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VL CDR1

2013217114

<400> 68

Arg Ala Arg Gln Gly Ile His Arg Tyr Leu Ser

1 5 10

<210> 69

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VL CDR2

<400> 69

Arg Ala Asn Arg Leu Gln Ser

1 5

<210> 70

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VL CDR2

31 Jul 2015

5641102_1

<400> 70

Arg Ala Asn Arg Arg Ala Thr

1 5

2013217114

<210> 71

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VL CDR2

<400> 71

Arg Ala Asn Arg Leu Val Ser

1 5

<210> 72

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 72

Trp Ile Asp Pro Asp Gln Gly Asp Thr Glu

1 5 10

2013217114 31 Jul 2015

5641102_1

<210> 73
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> amino acid sequence of VH CDR1

<400> 73

Trp Ile Asp Pro Asp Tyr Gly Asp Thr Glu
1 5 10

<210> 74
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> amino acid sequence of VH CDR1

<400> 74

Trp Ile Asp Pro Asp Ser Gly Asp Thr Glu
1 5 10

<210> 75
<211> 10
<212> PRT
<213> Artificial Sequence

<220>

2013217114 31 Jul 2015

<223> amino acid sequence of VH CDR1 5641102_1

<400> 75

Trp Ile Asp Pro Asp Asn Ala Asp Thr Glu

1 5 10

<210> 76

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR1

<400> 76

Trp Ile Asp Pro Asp Asn Thr Asp Thr Glu

1 5 10

<210> 77

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> amino acid sequence of VH CDR3

<400> 77

Asn Ala Ala Tyr Gly Ser Ser Pro Tyr Pro Met Asp Tyr

1 5 10

2013217114 31 Jul 2015

5641102_1

<210> 78
<211> 118
<212> PRT
<213> Artificial Sequence

<220>
<223> VH chain region of the 9E4 antibody

<400> 78

Glu Val Gln Leu Arg Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
20 25 30

Tyr Met Tyr Trp Val Lys Gln Ser Arg Val Arg Ser Leu Ala Trp Ile
35 40 45

Gly Arg Ile Asn Pro Tyr Thr Gly Ala Thr Gly Tyr Asp Gln Asn Phe
50 55 60

Lys Asp Lys Ala Ser Leu Ile Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

31 Jul 2015

2013217114

5641102_1

Ala Arg Gly Arg Asn Arg Tyr Asp Gly Trp Phe Ala Tyr Trp Gly Gln
100 105 110

Gly Thr Leu Val Thr Val
115

<210> 79
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> VL chain region of the 9E4 antibody

<400> 79

Glu Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly
1 5 10 15

Asp Arg Val Thr Ile Ser Cys Arg Ala Ser Gln Asp Ile Ser Asn Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile
35 40 45

Tyr Tyr Thr Ser Arg Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

31 Jul 2015

5641102_1

Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Asp Gln
65 70 75 80

2013217114

Glu Asp Ile Ala Thr Tyr Phe Cys Gln Gln Gly Asn Ala Leu Pro Pro
85 90 95

Thr Phe Gly Gly Thr Asn Leu Glu Ile Lys
100 105

<210> 80

<211> 115

<212> PRT

<213> Artificial sequence

<220>

<223> VH chain region of the 1B4 antibody

<400> 80

Gln Ile Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Tyr Ile His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

31 Jul 2015

2013217114

5641102_1

Gly Trp Ile Tyr Pro Gly Ser Gly Asn Thr Lys Tyr Asn Glu Arg Phe
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Ala Thr Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Phe Cys
85 90 95

Ala Arg Arg Glu Glu Asp Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val
115

<210> 81
<211> 113
<212> PRT
<213> Artificial Sequence

<220>
<223> VL chain region of the 1B4 antibody

<400> 81

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

31 Jul 2015

2013217114

5641102_1

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Thr Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Ile
100 105 110

Lys

<210> 82

<211> 19

<212> DNA

<213> Artificial Sequence

2013217114 31 Jul 2015

5641102_1

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 82

cactgcaggt rtccactcc

19

<210> 83

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 83

catagcaggt gtccactcc

19

<210> 84

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 84

crctacaggt gtccactcc

19

<210> 85

<211> 18

31 Jul 2015

5641102_1

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

2013217114

<400> 85

gcyacagmtg tccactcc

18

<210> 86

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 86

cactgcaggt gtccwmtcc

19

<210> 87

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 87

crctrcaagg gtckactcc

19

2013217114 31 Jul 2015

5641102_1

<210> 88
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 88
gctawmrgtg tccactcc 18

<210> 89
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 89
cctcaggtgt ccactcc 17

<210> 90
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 90
gctacaggtg ctcactcc 18

2013217114 31 Jul 2015

5641102_1

<210> 91
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 91
cactgcaggt gtcctctct

19

<210> 92
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 92
caytgcaggt gtccaytgc

19

<210> 93
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

2013217114 31 Jul 2015

5641102_1

<400> 93

gcttammggtg tccacttc

18

<210> 94

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 94

ctcctgtcak taactkcagg t

21

<210> 95

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 95

caactgcagg tgtctctct

19

<210> 96

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

31 Jul 2015

<223> 5641102_1
Chemically synthesized oligonucleotide primer
<400> 96
crctrccaggy gtccactct

19

2013217114

<210> 97
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 97
ccaagctgtc tcctttcc

18

<210> 98
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 98
ccaagctgtc tcctttcc

18

<210> 99
<211> 19
<212> DNA
<213> Artificial Sequence

2013217114 31 Jul 2015

5641102_1

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 99

cttgacagyc vttcckggt

19

<210> 100

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 100

cttcacagcc tttcctggt

19

<210> 101

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 101

cttaaaaggg gtccagtgt

19

<210> 102

<211> 22

31 Jul 2015

5641102_1

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

2013217114

<400> 102

caytttaaaa rgtgtc magt gt

22

<210> 103

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 103

gtttttaaaag gtgtcctgtg

20

<210> 104

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 104

ctyttaaaag gkgtccagwg

20

2013217114 31 Jul 2015

5641102_1

<210> 105
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 105
cytttamatg gtatccagtg t

21

<210> 106
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 106
cttttacatg gtttcaagtg t

21

<210> 107
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 107
gtccctgcat atgtcyt

17

2013217114 31 Jul 2015

5641102_1

<210> 108

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 108

gatggcagcw gcycaaag

18

<210> 109

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 109

ctatcaaggt gtgcattgt

19

<210> 110

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

2013217114 31 Jul 2015

5641102_1

<400> 110

cttttaaaag wtgtccagkg t

21

<210> 111

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 111

gtgacagtcc ttcctggtag

20

<210> 112

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 112

tttcctgatg gcagtggtt

19

<210> 113

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

31 Jul 2015

5641102_1
<223> Chemically synthesized oligonucleotide primer
<400> 113
gctacaggta tccaatcc

18

2013217114

<210> 114
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 114
gcgtctagaa yctccacaca caggrrccag tggatagac

39

<210> 115
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 115
ctgwtgttct ggattcctg

19

<210> 116
<211> 18
<212> DNA
<213> Artificial Sequence

201321714 31 Jul 2015

5641102_1

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 116

ggtcagacag tcagcagt

18

<210> 117

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 117

tgctctgga ttcgggaa

18

<210> 118

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 118

cagtttcytg ctaatcagtg

20

<210> 119

<211> 18

31 Jul 2015

5641102_1

<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

2013217114

<400> 119
ctaatcagtg cttcagga 18

<210> 120
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 120
gtgggtatct ggtrcstgtg 20

<210> 121
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 121
ggaaatcaa aagtacctgt ggg 23

2013217114 31 Jul 2015

5641102_1

<210> 122
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 122
ggtttc~~m~~agg trccagatgt 20

<210> 123
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 123
ctctggttc caggtatc 18

<210> 124
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 124
ctgtttcaa ggtrccagat gt 22

2013217114 31 Jul 2015

5641102_1

<210> 125

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 125

gttgtaatgt ccagagga

18

<210> 126

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 126

cttacaggtg ccagatgt

18

<210> 127

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

2013217114 31 Jul 2015

5641102_1

<400> 127
ctcaattgta grtgccagat gt 22

<210> 128
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 128
cacagtaggt gtcagatgt 19

<210> 129
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 129
gtcgttagttg tcagatgt 18

<210> 130
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

31 Jul 2015

<223> 5641102_1
chemically synthesized oligonucleotide primer
<400> 130
cctccttctt ggccaaga

18

2013217114

<210> 131
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> 131
chemically synthesized oligonucleotide primer

<400> 131
cttatatgga gctgatggg

19

<210> 132
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> 132
chemically synthesized oligonucleotide primer

<400> 132
gtgtctggtg ctcatggg

18

<210> 133
<211> 20
<212> DNA
<213> Artificial Sequence

2013217114 31 Jul 2015

5641102_1

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 133

ctstggtgtt ctgggttgta

20

<210> 134

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 134

gtctctgatt ctagggca

18

<210> 135

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 135

ctkckctggg ttccag

16

<210> 136

<211> 15

31 Jul 2015

5641102_1

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

2013217114

<400> 136

gcaggtgttg acgga

15

<210> 137

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 137

caggtgcctc gtgcac

16

<210> 138

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 138

ctctggtgcc tgtgca

16

2013217114 31 Jul 2015

5641102_1

<210> 139
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 139
ctggaytyca gcctccaga 19

<210> 140
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 140
gwtctctrga gtcagtggg 19

<210> 141
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 141
ctggatccct ggakcyact 19

2013217114 31 Jul 2015

5641102_1

<210> 142

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 142

gttctgcttt ttaggttg

19

<210> 143

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

<400> 143

gatcccaggc atgatatgt

19

<210> 144

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Chemically synthesized oligonucleotide primer

2013217114 31 Jul 2015

5641102_1

<400> 144
cttcatggtg ctcagtgt 18

<210> 145
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 145
ccatatcagg tgcccagtgt 20

<210> 146
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Chemically synthesized oligonucleotide primer

<400> 146
gcgtctagaa ctggatggtg ggaagatgg 29

<210> 147
<211> 305
<212> PRT
<213> Homo sapiens

<400> 147

2013217114 31 Jul 2015

5641102_1

Gln Leu Leu Phe Asn Lys Thr Lys Ser Val Glu Phe Thr Phe Cys Asn
1 5 10 15

Asp Thr Val Val Ile Pro Cys Phe Val Thr Asn Met Glu Ala Gln Asn
20 25 30

Thr Thr Glu Val Tyr Val Lys Trp Lys Phe Lys Gly Arg Asp Ile Tyr
35 40 45

Thr Phe Asp Gly Ala Leu Asn Lys Ser Thr Val Pro Thr Asp Phe Ser
50 55 60

Ser Ala Lys Ile Glu Val Ser Gln Leu Leu Lys Gly Asp Ala Ser Leu
65 70 75 80

Lys Met Asp Lys Ser Asp Ala Val Ser His Thr Gly Asn Tyr Thr Cys
85 90 95

Glu Val Thr Glu Leu Thr Arg Glu Gly Glu Thr Ile Ile Glu Leu Lys
100 105 110

Tyr Arg Val Val Ser Trp Phe Ser Pro Asn Glu Asn Ile Leu Ile Val
115 120 125

Ile Phe Pro Ile Phe Ala Ile Leu Leu Phe Trp Gly Gln Phe Gly Ile

2013217114
31 Jul 2015

5641102_1
130 135 140
Lys Thr Leu Lys Tyr Arg Ser Gly Gly Met Asp Glu Lys Thr Ile Ala
145 150 155 160
Leu Leu Val Ala Gly Leu Val Ile Thr Val Ile Val Ile Val Gly Ala
165 170 175
Ile Leu Phe Val Pro Gly Glu Tyr Ser Leu Lys Asn Ala Thr Gly Leu
180 185 190
Gly Leu Ile Val Thr Ser Thr Gly Ile Leu Ile Leu Leu His Tyr Tyr
195 200 205
Val Phe Ser Thr Ala Ile Gly Leu Thr Ser Phe Val Ile Ala Ile Leu
210 215 220
Val Ile Gln Val Ile Ala Tyr Ile Leu Ala Val Val Gly Leu Ser Leu
225 230 235 240
Cys Ile Ala Ala Cys Ile Pro Met His Gly Pro Leu Leu Ile Ser Gly
245 250 255
Leu Ser Ile Leu Ala Leu Ala Gln Leu Leu Gly Leu Val Tyr Met Lys
260 265 270

201321714 31 Jul 2015

5641102_1

Phe Val Ala Ser Asn Gln Lys Thr Ile Gln Pro Pro Arg Lys Ala Val
275 280 285

Glu Glu Pro Leu Asn Ala Phe Lys Glu Ser Lys Gly Met Met Asn Asp
290 295 300

Glu
305