



(86) Date de dépôt PCT/PCT Filing Date: 2011/09/16	(51) Cl.Int./Int.Cl. <i>C12N 15/82</i> (2006.01), <i>C07K 14/415</i> (2006.01), <i>C12N 15/29</i> (2006.01)
(87) Date publication PCT/PCT Publication Date: 2012/07/12	
(45) Date de délivrance/Issue Date: 2018/04/10	(72) Inventeurs/Inventors: KIM, SANGWOO, KR; LEE, YOUNGSOOK, KR; NISHIDA, IKUO, JP; YAMAOKA, YASUYO, JP; ONO, HIROFUMI, JP
(85) Entrée phase nationale/National Entry: 2013/07/05	
(86) N° demande PCT/PCT Application No.: KR 2011/006826	(73) Propriétaire/Owner: POSTECH ACADEMY-INDUSTRY FOUNDATION, KR
(87) N° publication PCT/PCT Publication No.: 2012/093764	
(30) Priorité/Priority: 2011/01/06 (KR10-2011-0001531)	(74) Agent: BROUILLETTE LEGAL INC.

- (54) Titre : COMPOSITION CONTENANT UN GENE CODANT POUR PROTEINES DE TRANSPORTEUR ABC POUR AUGMENTER LA TAILLE DE GRAINES DE PLANTES ET LA TENEUR EN GRAISSE STOCKEE DANS LES GRAINES
- (54) Title: COMPOSITION CONTAINING GENE ENCODING ABC TRANSPORTER PROTEINS FOR INCREASING SIZE OF PLANT SEED AND CONTENT OF FAT STORED WITHIN SEED

(57) **Abrégé/Abstract:**

The present invention relates to a composition for increasing the size of a plant seed and the content of fat store within the seed, and more specifically to a composition containing a gene encoding ABC transporter proteins for increasing the size of a plant seed and the content of fat store within the seed, and a method for increasing the size of a plant seed and the content of fat store within the seed comprising a step of introducing said gene and a promoter for over expressing said gene to a plant.

[ABSTRACT]

The present invention relates to a composition for increasing the size of a plant seed and the content of fat store within the seed, and more specifically to a composition containing a gene encoding ABC transporter proteins for increasing the size of a plant seed and the content of fat store within the seed, and a method for increasing the size of a plant seed and the content of fat store within the seed comprising a step of introducing said gene and a promoter for overexpressing said gene to a plant.

[DESCRIPTION]**[Invention Title]**

COMPOSITION CONTAINING GENE ENCODING ABC TRANSPORTER
5 PROTEINS FOR INCREASING SIZE OF PLANT SEED AND CONTENT OF FAT
STORED WITHIN SEED

[Technical Field]

The present invention relates to a composition for
10 increasing plant seeds in size or in content of fat store.
More particularly, the present invention relates to a
composition for increasing plant seeds in size or in content
of fat store, comprising a gene coding for an ABC (ATP-binding
cassette) transporter protein, and the method for increasing
15 plant seeds in size or in content of fat store, comprising the
step of introducing the gene, and a promoter for
overexpressing the gene into a plant body.

[Background Art]

20 Most plants store oil and fat in their seeds, with
triacylglycerol being more predominant than other oil and fat
forms. In plants, the first energy product is in the form of
carbohydrate which are later converted and stored into
triacylglycerol, a more dense form. Upon germination, 1 g of
25 the stored oil is converted into 2.7 g of carbohydrate, which

is directly used as an energy source during germination.

Vegetable oils are more abundant in unsaturated fatty acids compared to animal oils, and plants contain 18:2 and 18:3 fatty acids, and polyunsaturated derivatives thereof, 5 which cannot be synthesized in animals. These fatty acids are called essential fatty acids, since they cannot be synthesized in vivo, and thus could be only ingested through food. In addition to the importance as food, vegetable oils are emerging as an alternative energy source, with extensive 10 attention being paid as biodiesel due to the recent exhaustion of fossil fuel.

The production of oilseeds in the world amounted to 3.19 billion metric tons in 2001. While some are used directly as food, most of them are used for oil extraction. On average, 15 the content of oil in oilseed accounts for roughly 26 % of its total weight. Globally, the consumption of animal oil has been decreased or maintained since in 1970, while the consumption of vegetable oil has more than doubled between 1980 and 2000 (ca. 670 million tons), with a rapid increase in consumption 20 tendency (refer to USDA food disappearance records, U.S. PUFA Consumption, 1909-2005).

As can be seen, the demand for vegetable oils tends to increase as days go by, while supply can't catch up the demand. To date, the output of oils from oilseeds has been 25 maximized thanks to the continuous improvement in breeding and

crossbreeding, and it is anticipated that with the current breeding and crossbreeding, it is difficult to catch up the demand in a limited cultivation area. As a solution to this limitation, genetically modified organisms have come to the fore. For the production of vegetable oil which is expected for tremendous worldwide demand, it is anticipated that developing genetically modified organisms with high oil output is essential. In this regard, many researchers around the globe are conducting research on increasing the oil content per unit weight of seed and increasing the total amount of oil, but many genes are complicatedly involved in triacylglycerol synthesis in the seed and the control process which take up most of the stored vegetable oil, and thus, the goal isn't being accomplished easily.

In order to improve the stored oil of plant seed, big multinational corporations, such as Monsanto and DuPont are conducting a research. But yet, accomplishments haven't been reported since although a fat content of a seed increases, there are lots of cases when the overall productivity decreases due to the reduction of overall plant growth and the number of silique. There is therefore a need to develop a gene that functions to allow an increase in oil content in a seed that mostly stores the vegetable oil, without decreasing the size of seeds or the number of siliques, or more ideally, increases both the size of seeds and the number of siliques.

If this technology is developed, it can create higher values when applied in conjunction with other technologies developed in regards of plant fats. For example, the technology can be used to produce a greater amount of useful fats when it is
5 applied to a plant producing omega-3 fatty acids or tocopherol or a certain useful fat / fatty acids, together with a technique of increasing the size of seeds and the number of siliques.

10 **[Disclosure]**

[Technical Problem]

Accordingly, the present invention has been invented to solve the above problems occurring in the prior art, and an object of the present invention is to provide a composition
15 for increasing plant seeds in size and in content of fat store, comprising more than one selected from the group consisting of an ABC (ATP-binding cassette) transporter protein and a gene encoding the ABC transporter protein.

Another object of the present invention is to a method
20 for increasing plant seeds in size or in content of fat store, comprising the step of introducing a gene coding for ABC (ATP-binding cassette) transporter protein) and a promoter for overexpressing the gene into a plant.

25 **[Technical Solution]**

To fulfill the above purpose, the present inventors confirmed through research that when a gene coding for ABC (ATP-binding cassette) transporter protein is inserted and the gene is overexpressed, not only the size of the seeds of the transgenic plant increases, but also the fat store content itself inside the seeds increases, which eventually improves the stored fat store, thereby greatly increasing the productivity of seeds and fat store within seeds, and completed the invention.

10 In accordance with one aspect thereof, the present invention provides a composition for increasing plant seeds in size and in fat store content, comprising more than one selected from the group consisting of a plant ABC (ATP-binding cassette) transporter protein and a gene coding for the ABC
15 transporter protein.

In accordance with another aspect thereof, the present invention provides a method for increasing plant seeds in size or in fat store content, comprising the step of introducing a gene coding for a plant (ATP-binding cassette) transporter
20 protein, and a promoter for overexpression the gene into a plant.

In accordance with a further aspect thereof, the present invention provides a transgenic plant which bears seeds improved in size and in fat store content according to the
25 method.

In accordance with another aspect thereof, the present invention further provides a composition for increasing seed size of a plant or content of fat stored within the seeds, wherein seed size and content of fat stored within the seeds are increased in
5 comparison to the seeds of the same plant that has not been subjected to said composition, the composition comprising a transformation vector comprising a gene encoding a plant ATP-binding cassette A9 (ABC A9) transporter protein; and a promoter for overexpressing the gene; wherein the ABC A9 transporter protein
10 is a polypeptide having an amino acid sequence of SEQ ID NO: 1.

In accordance with another aspect thereof, the present invention also provides a method for increasing seed size of a plant or content of fat stored within the seeds, wherein seed size and content fat stored within the seeds are increased in comparison
15 to the seeds of the same plant that has not been subjected to the step outlined below, the method comprising the steps of inserting a gene encoding the plant ATP-binding cassette A9 (ABC A9) transporter protein and a promoter for overexpressing the gene into a transformation vector; and inserting the transformation vector
20 into the plant;
wherein the ABC A9 transporter protein is a polypeptide having an amino acid sequence of SEQ ID NO: 1.

In accordance with still another aspect thereof, the present invention provides a seed which is improved in size and fat store content according to the method.

A detailed description will be given of the present
5 invention, below.

In accordance with an aspect thereof, the present invention provides a composition for increasing plant seeds in size and in fat store content, comprising more than one selected from the group consisting of a plant ABC (ATP-binding cassette) transporter
10 protein and a gene coding for the ABC transporter protein.

As used herein, the term "ABC (ATP-binding cassette) transporter protein" refers to a protein, composed of six biomembrane-spanning domains and one ATP-binding domain and positioned after passing through a biomembrane consisted of
15 lipidic bilayer, and is a transporter that carried substances using the energy from ATP disintegration, and also plays a role of absorbing nutrition into the cell or transporting poisonous substance outside of the cell, and especially refers to a protein that directly or indirectly involves in transporting precursor for
20 the synthesis of a vegetable fat (or oil).

The term "overexpression," as used herein, refers to

expressing a gene encoding the ABC (ATP-binding cassette) transporter protein of the present invention at a higher level of that in a wild-type plant, and the method of overexpression may be performed using various well-known techniques, without
5 particular limitations loaded thereto. For example, overexpression may be achieved by mutation in a ribosomal binding site, a promoter, or a regulatory region upstream of a structural gene or introduced to increase the number of appropriate gene copies, or an upstream from a structural gene
10 introduced expression cassette can be worked in the same manner. Also, expression may be increased by introducing a gene inducible promoter encoding the ABC (ATP-binding cassette) transporter protein of the present invention or by extending the life of mRNA. A medium composition and/or
15 changing the culturing technique can also overexpress the above gene.

The term "transformation" in the present invention means a molecular biological technique that changes the genetic trait of a cell by a DNA chain fragment or plasmid that
20 possess a different type of foreign gene from that of the original cell have, penetrate among the cells and combine with the DNA that existed in the original cell. In the context of the present invention, transformation means that the above genes along with the overexpression promoter are inserted into
25 the plant body.

The term "plant body," as used herein, may be selected from the group consisting of a plant, plant tissue, cell, and seed.

As used herein, the term "transformants" refers to the
5 above plant body whose genetic trait has been altered through the above transformation.

The "plant cell" can be any cell of a plant. The plant cell may be a cultured cell, a cultured tissue, a cultured organ or a whole plant, preferably a cultured cell, a cultured
10 tissue or a cultured organ, and more preferably any form of cultured cell.

The "plant tissue" is intended to encompass differentiated or undifferentiated plant tissues including, but not limited to, roots, stems, leaves, pollens, seeds,
15 cancer tissue and various type of cells used in culture, that is, single cell, protoplast, sprout, and calluses. The plant tissue may be in planta, or may be in the status of organ culture, tissue culture or cell culture.

The expression "increase of seed in size," as used
20 herein, is intended to preferably include, but not limited to the increase of plant seed and seed's weight, volume, number and size of silique, and the increase of yield and productivity thereof. The expression "increase of fat store content in a seed" refers to increasing the content of fat (or
25 oil) stored in plant seed, and the increase of fat store

content in the seed may preferably expand in meaning to, but not limited to, the increase of fat store content itself, but also the increase of fat store content in the seed and the productivity increase of vegetable fat (or oil) as a result, 5 through the increase of seed in size.

The typical example of fat store in the above seed is triacylglycerols. A triacylglycerol (triacylglyceride; triglyceride) has a structure of one glycerol skeletal structure bound with three fatty acids, and is a principle 10 component for both vegetable oil and animal fat. This triacylglycerols is transformed as energy source and used in animal upon scarce carbohydrate, and in plants, triacylglycerols is a representative fat store which is mainly stored in seed and used as nutrient upon seed germination.

15 The composition for increasing plant seeds in size or fat store content in accordance with the present invention is characterized by comprising an ABC transport protein, and this ABC transport protein may be preferably a polypeptide having the amino acid sequence of SEQ ID NO: 1 20 (AtABCA9/AtATH11), but is not limited thereto.

Preferably, it may be a polypeptide composed of a homologous sequence with amino acid sequence of SEQ ID NO: 1 (AtABCA9/AtATH11), and the homologous sequence may preferably share a homology of from 65 % to 99 %, more preferably from 25 80 % to 99 %, far more preferably from 90 % to 99 %, and most

preferably from 90 % to 95 % or from 95 % to 99 % with the amino acid sequence of SEQ ID NO: 1.

Moreover, the gene encoding the polypeptide having the amino acid sequence of SEQ ID NO: 1, could preferably be, but
5 not limited to the gene of SEQ ID NO: 2 (AtABCA9/AtATH11) consisted of base sequence.

In addition to the gene, the composition may comprise the gene, a transformation vector inserted with a promoter for overexpressing the gene, or a microorganism transformed by the
10 above transformation vector.

The term "transformation vector," as used herein, refers to a recombinant DNA molecule comprising the proper nucleic acid sequence which is essential for expressing coding sequence of interest and a coding sequence that is connected
15 only to operate in specific host organism. The proper nucleic acid sequence may be a promoter, and may further comprise an enhancer, a transcriptional terminator, and a polyadenylation signal. Promoters, enhancers, transcriptional terminators, and polyadenylation signals which are available in eukaryotic
20 cells are well known.

Examples of the promoter is a promoter for plant expression which can use CaMV (Cauliflower Mosaic Virus) 35S promoter, NOS (nopaline synthase) promoter of Agrobacterium tumefaciens Ti plasmid, OCS (octopine synthase) promoter and
25 MAS (mannopine synthase) promoter, but are not limited

thereto.

The transcriptional terminator may use typical one examples of which include nopaline synthase (NOS), rice α -amylase RAmyl A terminator, phaseoline terminator, and a
5 terminator for optopine gene of *Agrobacterium tumefaciens*, but are not limited thereto.

Preferably, the transformation vector may further comprise a marker for identifying the gene expression or selecting a transformants. The marker is a nucleotide
10 sequence typically having a property based on which it can be selected by a common chemical method, and any gene that is available to differentiate transformed cells from non-transformed cells can be a marker. Marker gene may include a gene resistant to an antibiotic such as kanamycin,
15 spectinomycin, or a gene encoding GUS (β -glucuronidase), but are not limited thereto. The marker is delivered to a plant together with the vector, and is cultivated in a medium which contains a specific antibiotic and allows selecting transformants.

20 The transformation vector may be a plant expression vector in which a base sequence of the gene is inserted and be directly introduced into a plant cell or be introduced into a microorganism that trigger infection to a plant.

A preferred embodiment of the plant expression vector is
25 a Ti-plasmid vector, which can transfer its segment, known as

T-region, to plant cell when existing in a proper host such as *Agrobacterium tumefaciens*. There are many strains of *Agrobacterium* that can be used in this manipulation and are known in the art. Another type of T-plasmid vector is used as
5 transferring hybrid DNA sequence, and is a protoplast which can produce a new plant that properly inserts the present plant cell or a hybrid DNA sequence into a plant's genome.

Another preferred type of the Ti-plasmid vector is a binary vector disclosed in EP 0120516 B1 and U.S. Pat. No.
10 4,940,838. Another proper vector which can be used to introduce the gene according to the present invention into a plant host may be selected from among virus vectors that can be derived from double stranded plant viruses including CaMV, single stranded viruses, and geminiviruses, for example, a
15 non-complete plant virus vector. Use of these vectors may be especially advantageous when it is difficult to properly transform a plant host. Most preferred may be a pCAMBIA1302 vector with a cauliflower mosaic virus(CaMV) 35S promoter.

More particularly, the vector may be
20 pCAMBIA1302::AtABCA9/AtATH11 which may comprise a 35S promoter, an AtABCA9/AtATH11 gene, and a nopaline synthase transcription terminator.

So long as it causes an infection in plants, any microorganism may be used without limitation in the present
25 invention. For example, various viruses or bacteria may be

employed, preferably, the soil bacterium, *Agrobacterium tumefaciens*, which causes crown gall disease in dicotyledonous plant.

The plants may be, for example, herbaceous plants and/or
5 woody plants and may be flowering plants, garden plants, rice, wheat, barley, corn, soybean, red bean, oat, sorghum, onion, carrot, cucumber, olive, sweet potato, potato, Chinese cabbage, radish, lettuce, broccoli, tobacco, petunia, sunflower, leaf mustard, turf, *Arabidopsis*, *Brassica*
10 *campestris*, *B. napus*, *Betula platyphylla*, poplar, hybrid poplar, and *Betula schmidtii*.

According to another exemplary embodiment of the present invention, the present invention provides a method for
15 increasing plant seeds in size or fat store content, comprising the step of introducing a gene coding for a plant ABC (ATP-binding cassette) transporter protein, and a promoter for overexpressing the gene into a plant body.

The ABC transport protein may be preferably a polypeptide
20 having the amino acid sequence of SEQ ID NO: 1 (*AtABCA9/AtATH11*), but is not limited thereto.

As for the gene encoding the polypeptide having the amino acid sequence of SEQ ID NO: 1, could preferably be but limited to, SEQ ID NO: 2 (*AtABCA9/AtATH11*) gene which is consisted of
25 base sequence.

In order to introduce the gene into a plant body, an Agrobacterium tumefaciens-mediated gene transfer method may be preferably used, and more preferably, may use recombinant Agrobacterium dipping method which is produced through
5 selecting the method among electroporation, micro-particle injection and gene gun, but is not limited thereto.,

According to further another exemplary embodiment of the present invention, the present invention provides a plant body
10 which is increased in seed size and fat store content in the seed according to the method of increasing the seed size of the plant or the fat store content of the seed.

The plant body may be preferably selected from the group consisting of a whole plant, a plant tissue, a plant cell, and
15 a plant seed.

According to still another exemplary embodiment of the present invention, the present invention provides a seed which is increased in size or fat store content according to the method of increasing the seed size of the plant or the fat
20 store content of the seed.

[Advantageous Effects]

The composition for increasing the plant seed size and fat store content in the seed and the method for increasing
25 plant seeds in size or fat store content according to the

present invention can be used to increase the size of the plant seed of course, and the weight, and volume of the seed, the number and size of siliques, and thereby improving seed yield and productivity. Moreover, increases not only the
 5 content of stored fat (or oil) in seeds itself, but also more increases the fat store content in the seed through increasing the size of the seed which is a main storage organ of the plant oil as stated, and thus making a significant contribution to the supply of biodiesel and edible plant fats
 10 (or oils) in response to the rapidly increasing demand.

[Description of Drawings]

FIG. 1 shows three different mutants according to Example 1, and Fig. 1a shows three different alleles (*abca9-1/ath11-1*.
 15 *abca9-2/ath11-2*, *abca9-3/ath11-3*) with T-DNA to produce AtABCA9/AtATH11 mutant *Arabidopsis thaliana*, Fig. 1b shows the result of running gene-specific PCR (a) and T-DNA-specific PCR (b) according to Example 1 to examine the knockout due to the insertion of T-DNA. (WT: wild-type; 1st: *abca9-1*, 2nd: *abca9-2*,
 20 3rd: *abca9-3*).

FIG. 2 shows the result of measuring the germination rates of the three different mutants that can't express AtABCA9/AtATH11 and the wild species in accordance with Example 2-1.

25 FIG. 3 shows the result of comparing the growth of the

three different mutants that can't express AtABCA9/AtATH11 and the wild species in accordance with Example 2-2.

FIG. 4a shows the result of comparing wild species and the seeds of three different mutants that can't express AtABCA9/AtATH11 in accordance with Example 3, and FIG. 4b shows the result of observing the inside of the siliques of the wild species and the three different mutants that can't express AtABCA9/AtATH11 (1st: *abca9-1*, 2nd: *abca9-2*, 3rd: *abca9-3*) as observed every 4 days after flowering (4, 8, 12, 16, and 20 DAF (Day After Flowering)) in accordance with Example 3,.

FIG. 5a is a graph showing the result confirmed through quantitative RT-PCR by classifying the expression of AtABCA9/AtATH11 in organizations (WS: whole seedling), L: leaf, S: stem, F: flower, S1: silique 4 ~ 6 DAF, S2: silique 10 ~ 12 DAF, and S3: silique 16 ~ 18, DAF: (Day After Flowering)), in accordance with Example 5, and FIG. 5b is a graph showing the expression levels of the overexpression transformants through quantitative RT-PCR (WT: wild-type, T1~T12 : overexpression transformants) in accordance with Example 5.

FIGS. 6a and 6b show the result of measuring fat store in a seed in accordance with Example 6.

FIG. 7 shows the result of observing the plant growth of wild species and transformants that overexpress AtABCA9/AtATH11 in accordance with Example 7 (WT: wild-type,

T5, T7, T9, T11 and T12 : overexpression transformants).

FIG. 8 shows the result of observing the number of siliques and the number of seeds in the silique of wild species and transformants that overexpress AtABCA9/AtATH11 in accordance with Example 7 (WT: wild-type, T5, T7, T9, T11 and T12 : overexpression transformants).

FIGS. 9a to 9c each shows the result of measuring siliques (Fig.9a), the weight of seeds (Fig.9b), and the content of TAG of the seed(Fig.9c) inside the silique of wild species and overexpressing transformants in accordance with Example 7 (WT: wild-type, T1~T12 : overexpression transformants).

FIG. 10 shows the amino acid of sequence ID No:1 consisting the ABC transporter protein of the present invention.

FIG. 11 shows the base sequence of sequence ID No:2 consisting the gene encoding ABC transporter protein of the present invention.

20 [Best Mode]

Hereinafter, preferred embodiments of the present invention will be presented for a better understanding of the present invention. However, the description proposed herein is just an example for a better understanding, not intended to limit the scope of the invention.

REFERENCE EXAMPLE 1: Isolation of Genomic DNA (gDNA)

Genomic DNA was extracted using a DNA extraction buffer (50 mM Tris-HCl, 10 mM EDTA, 100 mM NaCl, 1%(v/v) SDS, 10 mM β -mercaptoethanol) from leaves of *Arabidopsis thaliana* (wild-type (Columbia-0, SALK institute) and mutant) cultured for 2 weeks in a 1/2 MS medium (Murashige and Skoog, 1962),.

In detail, leaves of *Arabidopsis thaliana* were homogenously pulverized using liquid nitrogen, and mixed with 750 μ l of the DNA extraction buffer per 1 g of the pulverized sample by vortexing, followed by incubation at 65°C for 30 min. Subsequently, 250 μ l of 5 M potassium acetate was added to the sample and was shaken, cooled for 20 minutes in ice, and centrifugation was processed at 12000 rpm in 4°C for 10 min. The supernatant was transferred to a fresh tube, and 700 μ l of isopropanol was added and shaken slightly. The pellet formed after removing supernatant through centrifugation at 12000 rpm, in 4°C for 10 min, was washed with 80 % (v/v) ethanol and then dried. It was dissolved in 50 μ l of a TE buffer and separated genomic DNA and was used in PCR, etc.

EXAMPLE 1: Isolation of *Arabidopsis thaliana* that can't express AtABCA9/AtATH11

To search for a mutant incapable of expressing

AtABCA9/AtATH11 due to the insertion of T-DNA into the AtABCA9/AtATH11 gene, a mutant with T-DNA inserted in three different sites was ordered from SALK Institute (<http://signal.salk.edu/>) (FIG. 1a) The three mutants ordered
 5 were SALK_058070: abca9-1, SALK_084342:abca9-2, and SALK_023744:abca9-3, and the three mutants and wild-type (Columbia-0, SALK institute) *Arabidopsis thaliana* were cultured for 2 weeks in a 1/2 MS medium (Murashige and Skoog, 1962) containing kanamycin and extracted genomic DNA of each and
 10 identified the knockout status of genes.

In detail, the genomic DNA extracted from the Example 1 method was the template and performed the gene (AtABCA9/AtATH11)-specific PCR and T-DNA-specific PCR response, and for the PCR response, the following primers were
 15 used.

<List of Primers used >

1. pROKLBb1 (SEQ ID NO: 3: 5'-GCGTGGAACCGCTTGCTGCAACT-3')
2. SALK_058070LP (SEQ ID NO: 4: 5'-CTACATATGGCTCGTGGGAAC-
 3')
- 20 3. SALK_058070RP (SEQ ID NO: 5: 5'-AAAGAGGTGGAGGTGCTCTTC-
 3')
4. SALK_084342LP (SEQ ID NO: 6: 5'-ATGACTCTGCGAGAAGGCTT-
 3')
5. SALK_084342RP (SEQ ID NO: 7: 5'-GAAAGAGACCAAACACACC-
 25 3'F)

In this list, the pROKLBb1 primer of 1 was specific for T-DNA while all the primers of 2 to 5 were specific for the gene AtABCA9/AtATH11.

In SALK_058070: for the gene-specific PCR of abca9-1, no. 2 of the list was used as a forward primer and no. 3 as a reverse primer, and for T-DNA-specific PCR, no.1 was used as a forward primer and no. 3 as a reverse primer.

In SALK_084342: for the gene-specific PCR of abca9-2, no. 4 of the list was used as a forward primer and no. 5 as a reverse primer, and for T-DNA-specific PCR, no.1 was used as a forward primer and no. 5 as a reverse primer.

In SALK_023744: for the gene-specific PCR of abca9-3, no. 2 of the list was used as a forward primer and no. 3 as a reverse primer, and for T-DNA-specific PCR, no.2 was used as a forward primer and no. 1 as a reverse primer.

By performing PCR(35 cycles of 95°C 30 sec, 56°C 30 sec, and 72°C 30 sec) using each primers as described above, a mutant in which T-DNA was inserted as a homozygote into the AtABCA9/AtATH11 gene was excavated (FIGS. 1a and 1b).

In detail, as a result of performing gene specific PCR using primer pairs specific for the gene (AtABCA9/AtATH11), band that becomes PCR only in wild-type(WT) was identified((a) of FIG. 1b), and as a result of performing T-DNA specific PCR using RP or LP primers specific for each gene (abca9-1, abca9-2: RP, abca9-3: LP) and primers specific for T-DNA, band that

only becomes PCR in knockout lanes was identified and excavated mutants in which T-DNA was inserted as a homozygote.

**EXAMPLE 2: Effect of AtABCA9/AtATH11 Gene on Germination
5 and Growth of Plant**

**EXAMPLE 2-1. Effectiveness Measurement on Germination
Rate**

To measure germination rates of *Arabidopsis thaliana* mutants that can't express AtABCA9/AtATH11 (abca9-1, abca9-2,
10 abca9-3) identified in Example 1 and the wild-type (Columbia-0, SALK institute), each seeds were planted in a 1/2 MS medium (Murashige and Skoog, 1962), and was monitored after 48 hrs at intervals of 12 hrs by optical microscopy. Germination was determined on the basis of the emergence of radicles.

15 The observation result, as shown in FIG. 2, while the wild-type was 98 % germinated after 48 hrs since the seed was planted in the medium, the mutant that can't express AtABCA9/AtATH11 showed late germination comparing to wild-type due to small and crumpled seed, and even if it was germinated,
20 only 20 ~ 40 % were germinated, and roughly 20 % of the mutant did not germinate to the end.

EXAMPLE 2-2 Effectiveness Measurement on Growth

To measure growth rates of *Arabidopsis thaliana* mutants
25 that can't express AtABCA9/AtATH11 (abca9-1, abca9-2, abca9-3)

identified in Example 1 and the wild-type (Columbia-0, SALK institute), each seeds were planted in rich media containing sucrose and a sucrose-free media of 1/2 MS media (Murashige et al., 1962, Duchefa), and after cultivated for 3 weeks, the growth was observed and measured the effect on the growth of *Arabidopsis thaliana*

As a result shown in FIG. 3, there were no differences between the wild-type and the three different mutants that can't express AtABCA9/AtATH11 in the rich media containing 1 % of sucrose, while when there isn't a sucrose and should germinate and grow fully using the nutrients in the seed, the mutants that can't express AtABCA9/AtATH11 showed slow growth and couldn't grow well compared to wild-type.

EXAMPLE 3: Effect Measurement on the seed of AtABCA9/AtATH11 Gene

As a result of observing the seed of *Arabidopsis thaliana* mutants(abca9-1, abca9-2, abca9-3) that can't express the AtABCA9/AtATH11 identified in Example 1, it was identified that it produced a far greater number of small and rumped seeds compared to the wild-type (Columbia-0, SALK institute) (FIG. 4a), and when observing the seed inside the siliques of the wild-type and three different mutants that can't express AtABCA9/AtATH11 at intervals of 4 days after flowering (4, 8, 12, 16, and 20 DAF (Day After Flowering)), three different mutants that can't express AtABCA9/AtATH11 produced small,

pale colored seeds from earlier times compared to the wild-type and eventually produced more small and rumped seeds which indicated that AtABCA9/AtATH11 gene has an effect from the early time of the formation of siliques that actually
 5 forms seeds (FIG. 4b).

EXAMPLE 4: Preparation of Arabidopsis thaliana Transformants that Overexpress AtABCA9/AtATH11 Gene

In reference of Example 1, serving the gDNA as a template
 10 that was extracted from the wild-type Arabidopsis thaliana (Columbia-0, SALK institute), AtABCA9/AtATH11 gene was amplified by PCR (25 cycles of 95°C 30 sec, 56°C 30 sec, 72°C 2.5 min) using AtATH11-ATG (SEQ ID NO: 8: 5'-ACTAGTATGACTCTGCGAGAAGGCTT-3') and AtATH11-Stop (SEQ ID NO: 9:
 15 5'-ACTAGTTTAATTGTTAGATTCATAATCA-3') primers, and T-blunt vector (Solgent, <http://www.solgent.co.kr/>) was added, and was recombined in pCAMBIA1302 binary vector (Cambia, <http://www.cambia.org/daisy/cambia/585.html> using SpeI (Roche, <http://www.roche.co.kr/portal/kr>). pCAMBIA1302 vector was used
 20 as plant expression vector to produce Arabidopsis thaliana transformants and this contains a CaMV 35S promoter, a multi-cloning site, and a nopaline synthetase terminator.

Using the floral dipping method (Clough and Bent, 1988), Agrobacterium (GV3101) (obtained from Cauliflower mosaic virus
 25 (CaMV)) that contains pCAMBIA1302:AtABCA9/AtATH11 gene was

used to transform into *Arabidopsis thaliana*. The transformed seed was selected in 1/2 MS (Murashige and Skoog, 1962) medium containing 50 µg/L kanamycin antibiotics, and the seeds were taken from the plants survived. Homozygote seeds in the 3rd generation were used for phenotype analysis.

EXAMPLE 5: Quantitative RT-PCR

EXAMPLE 5-1: RNA Isolation and cDNA Synthesis

<Preparation of *Arabidopsis thaliana* sample>

1) Samples to measure the expression of AtABCA9/AtATH11 by tissues

To measure the expression of AtABCA9/AtATH11 according to tissues, samplings of each tissues (WS: whole seedling grown for 2 weeks in a medium, L: leaves grown for 2 weeks in a medium and for 2 weeks in soil, S: stems grown for 2 weeks a medium and for 3 weeks in soil, F: flowers grown for 2 weeks in a medium and for 3 weeks in soil, S1: siliques 4 ~ 6 DAF, S2: siliques 10 ~ 12 DAF, and S3: siliques 16 ~ 18 DAF, DAF: Day After Flowering) of wild-type *Arabidopsis thaliana*(Columbia-0, SALK institute) were done, and were used for RNA isolation and cDNA synthesis.

2) Samples for measuring expression of overexpressing-transformants

To measure the expression of overexpressing-transformants, the wild-type *Arabidopsis thaliana* and the

Arabidopsis thaliana transformants which was produced according to the Example 4 were cultured for 6 weeks, and samplings were done on each siliques, and were used for total RNA isolation and cDNA synthesis.

5

< RNA Isolation and cDNA Synthesis >

After 100 mg of Arabidopsis thaliana samples of 1) and 2) were evenly pulverized using liquid nitrogen, the pulverized samples were placed in 2 ml tubes, and were well mixed with
10 800 µl of an RNA extraction buffer (p-aminosalicylic acid 2.7 g, triisopropyl naphthalene sulfonic acid 0.45 g, 1.25M Tris HCl pH 9.0, 1.25 M NaCl, 0.25 M EDTA, β-mercaptoethanol 1 ml, acidic phenol 1.2 ml, DEPC water 36 ml) and 400 µl of acidic phenol. After inserting 400µl of chloroform and well blended,
15 place it in ice for 10 min. After centrifugation at 6000 rpm in 4°C for 10 min, the supernatant was transferred to a fresh tube, and 200 µl of isopropanol was added. After centrifugation at 6000 rpm in 4°C for 15 min, the supernatant was removed and the pellet was dried.

20 After the dried pellet was treated with DNase (Roche, <http://www.roche.co.kr/portal/kr>), 50 µl of RNase-free water was inserted to the tube, and the pellet was dissolved to earn RNA.

To the 2 µg of RNA obtained from above, cDNA was
25 synthesized using Powerscript RT (reverse transcription)-kit

(BD Bioscience Clontech) and oligo dT primer according to the manufacturer's instruction.

EXAMPLE 5-2 : Quantitative RT- PCR

To know the expression pattern by tissues of
 5 AtABCA9/AtATH11 gene and the level of expression of the
 overexpressing transformants, each cDNA of wild-type and
 transformants through Example 5-1 were served as template, and
 quantitative RT-PCR was performed (45 cycles of 94°C 30 sec,
 56°C 15 sec, 15°C sec) using the AtABCA9/AtATH11 F1 primer
 10 (SEQ ID NO: 10: 5'-ATGACTCTGCGAGAAGGCTT-3') and
 AtABCA9/AtATH11 R1 primer (SEQ ID NO: 11: 5'-
 CATGGAAGAATCGGAAGAGA-3').

As a results of measuring the expression patterns
 according to tissues of AtABCA9/AtATH11 (WS, whole seedling;
 15 L, leave; S, stem; F, flower; S1, siliques 4~6DAF; S2,
 siliques 10~12 DAF; and S3, siliques 16~18 DAF, DAF; Day After
 Flowering) AtABCA9/AtATH11 showed a higher expression level in
 silique compared to other tissues(FIG. 5a), and according to
 the level of expression of overexpressing transformants,
 20 expression of AtABCA9/AtATH11 gene at overexpressing
 transformants (T1~T12 in FIG. 5b) is significantly higher
 compared to the wild-type (WT in FIG. 5b). ubiquitin 1
 (AY059080) was used as a control plot.

EXAMPLE 6: Measurement of Fat store in Seeds

To measure the fat store in the seeds of mutants that can't express AtABCA9/AtATH11 identified in Example 1 and wild-type (Columbia-0, SALK institute), stored oils were extracted from 300 seeds of each. 300 seeds were put in 1 mL of 2-propanol and were boiled at 80°C for 5 min, and were chilled, and then 2 mL of chloroform was added and were grinded finely with Polytron Homogenizer. And then, after it was centrifuged at 3,000 rpm in room temperature for 10 min, the supernatant was collected, and 0.9 % (v/v) of KCl was added and shaken, followed by centrifugation at 3,000 rpm in room temperature for 10 min.

When the solvent was evaporated after the lower layer was collected, only the oil remained, and by weighing this the total weight of oil could be measured (FIG. 6a), and then was dissolved by chloroform, and 1 mg of the oil was loaded to a TLC plate (Merck 1.05721). In order to observe the triacylglycerol, a main neutral fat of a seed, 90 ml of hexane, 7.5 ml of diethylether, and 1 ml of acetic acid were mixed in a glass tank as a solvent to separate neutral fat, and then deployed inserting TLC plate that loaded samples. After deployment, the TLC plate was stained with primuline and triacylglycerol was observed under UV light. Triacylglycerol was scraped off from the TLC plate and after it was dissolved in 3 mL of hexane, the quantity of triacylglycerol which is a main oil constituent was measured using gas chromatography

(FIG. 6b).

As can be seen in FIG. 6a and 6b, for mutants that can't express AtABCA9/AtATH11, the total weight of seed and total fat (or oil) decreased but didn't show much difference in protein content compared to wild-type (FIG. 6a), and for triacylglycerol, which is a main fat store (or oil) of seeds, decreased about 35 % in mutants compared to wild-type (FIG. 6b). These results indicate that AtABCA9/AtATH11 plays an important role in the fat store of seed.

10

EXAMPLE 7: Productivity Measurement of Overexpressing Transformants

To compare the productivity of the transformants that overexpresses the AtABCA9/AtATH11 gene produced in Example 4, the seeds from wild-type *Arabidopsis thaliana* (Columbia-0, SALK institute) and the transformed *Arabidopsis thaliana* were stored for 2 days after the surfaces were sterilized with ethanol and chlororox, at 4°C in a dark place, and then was cultured in 1/2 MS medium (Murashige and Skoog 1962). After the medium was cultured horizontally for 2 weeks (16h light/8h dark, 22 °C /18 °C), they were transferred to soil and were grown for 4 weeks and the growth rate, the number of siliques, the number of seeds per silique, the size and weight of seeds, and the TAG content in seeds were measured and compared from wild-type and transformants. The TAG measurement was done by

25

extracting TAG using seeds from wild-type and overexpressing transformants according to the method disclosed in Example 6.

As a result, the growth of plants were similar between wild type (WT) and overexpressing transformants (T5, T7, T9, T11 and T12) or transformants showed slightly higher growth (FIG. 7, and when the number of siliques and the number of seeds in the silique were compared, the number of siliques was increased by 6.2 % on average in the overexpressing transformants, while there were no difference in the number of seeds per silique therebetween (FIG. 8).

In addition, for the overexpressing transformants (T5 and T8), the seeds in silique became much bigger in size (FIG. 9a) compared to wild type, and especially the transformants T8 showed increase up to 123 % (FIG. 9b) in seed weight, compared to the wild-type. Also, the transformants, particularly T9, contained triactylglycerols (TAG), which is the main fat store of seed, in an amount 132 % higher than that of the wild-type (FIG. 9c).

Consequently, the transformants that overexpress AtABCA9/AtATH11 gene of the present invention showed significant increase in the number of silique, the size of seeds in siliques, and the content of TAG, although there hasn't been much difference in the number of seeds per silique, and thus identified a great improvement in view of total productivity and fat store.

[Industrial Applicability]

The present invention can significantly improve the productivity of vegetable fat (or oil) in a limited space since it could more increase the fat store content in seeds through the increase of content itself of the fat store (or oil) stored in plant seed and the increase of seed in size which is a main storage organ of vegetable oil. And thus can make a significant contribution to the supply of biodiesel and edible plant fats (or oils) in response to the rapidly increasing demand therefor.

[Sequence List text]

15 SEQ. ID. NO. 1 : amino acid sequence of AtABCA9/AtATH11

Met Thr Leu Arg Glu Gly Leu Pro Leu Phe His Gln Gln Phe
 Thr Ala Leu Phe Lys Lys Asn Leu Leu Leu Ser Trp Arg Asn Lys
 Arg Ala Thr Cys Leu His Leu Phe Ser Ser Phe Phe Phe Ile Leu
 Leu Ile Phe Ser Ile Glu Glu Ser Ser Lys Ala Ser Asp Leu Thr
 20 Ser Thr Arg His Lys Asn Val Thr Asp Pro Lys Ala Leu Val Ser
 Leu Pro Ile Leu Pro Cys Glu Asp Lys Phe Phe Val Arg Leu Pro
 Cys Phe Asp Phe Val Trp Ser Gly Asn Gln Ser Arg Arg Val Thr
 Asp Ile Val Ser Ala Ile Met Ala Asn Asn Pro Gly Arg Pro Ile
 Pro Thr Asn Lys Val Gln Ser Phe Thr Lys Pro Glu Glu Val Asp
 25 Ala Trp Phe Met Ser His Pro Ser Gln Val Thr Gly Ala Leu His

Phe Val Glu Lys Asn Ala Thr Val Ile Ser Tyr Gly Ile Gln Thr
 Asn Ser Ser Ser Glu Lys Lys Arg Gly Arg Arg Glu Asp Pro Thr
 Phe Lys Phe Leu Val Pro Leu Gln Ile Ala Ala Glu Arg Glu Ile
 Ala Arg Ser Leu Ile Gly Asp Pro Lys Phe Ser Trp Asp Phe Gly
 5 Phe Lys Glu Phe Ala Arg Pro Ala Ile Gly Gly Glu Val Ile Ile
 Ser Ala Phe Tyr Leu Met Gly Pro Val Phe Phe Leu Ala Phe Ser
 Met Phe Gly Phe Val Leu Gln Leu Gly Ser Val Val Thr Glu Lys
 Glu Leu Lys Leu Arg Glu Ala Met Thr Thr Met Gly Val Tyr Glu
 Ser Ala Tyr Trp Leu Ser Trp Leu Ile Trp Glu Gly Ile Leu Thr
 10 Phe Val Ser Ser Leu Phe Leu Val Leu Phe Gly Met Met Phe Gln
 Phe Glu Phe Phe Leu Lys Asn Ser Phe Val Leu Val Phe Leu Leu
 Phe Phe Leu Phe Gln Phe Asn Met Ile Gly Leu Ala Phe Ala Leu
 Ser Ser Ile Ile Ser Lys Ser Ser Ser Ala Thr Thr Val Gly Phe
 Leu Val Phe Leu Val Gly Phe Ile Thr Gln Ile Val Thr Thr Ala
 15 Gly Phe Pro Tyr Ser Ser Ala Tyr Ser Ile Gly Ser Arg Val Ile
 Trp Ser Leu Phe Pro Pro Asn Thr Phe Ser Ala Gly Leu Gln Leu
 Leu Leu Glu Ala Thr Ser Ser Pro Gly Asp Ser Gly Ile Ser Trp
 Ser Glu Arg Ala Ile Cys Ala Gly Gly Glu Ser Thr Cys Val Ile
 Thr Thr Asn Lys Ile Tyr Ile Trp Leu Val Gly Thr Phe Phe Phe
 20 Trp Phe Val Leu Ala Leu Tyr Phe Asp Asn Ile Ile Pro Asn Ala
 Ser Gly Val Arg Lys Ser Ile Phe Tyr Phe Leu Lys Pro Ser Tyr
 Trp Thr Gly Lys Glu Gly Asn Lys Val Glu Glu Gly Ser Ile Cys
 Ser Cys Ile Gly Ser Val Pro Pro Val Glu His Ile Thr Pro Glu
 Asp Glu Asp Val Leu Glu Glu Glu Ile Leu Val Lys Gln Gln Ala
 25 Met Asp Gly Arg Val Asp Pro Asn Ile Ala Val Gln Ile His Gly

Leu Ala Lys Thr Tyr Pro Gly Thr Thr Lys Leu Gly Cys Cys Lys
 Cys Thr Lys Thr Ser Pro Phe His Ala Val Lys Gly Leu Trp Met
 Asn Ile Ala Lys Asp Gln Leu Phe Cys Leu Leu Gly Pro Asn Gly
 Ala Gly Lys Thr Thr Thr Ile Ser Cys Leu Thr Gly Ile Asn Pro
 5 Val Thr Gly Gly Asp Ala Lys Ile Tyr Gly Asn Ser Ile Arg Ser
 Ser Val Gly Met Ser Asn Ile Arg Lys Met Ile Gly Val Cys Pro
 Gln Phe Asp Ile Leu Trp Asp Ala Leu Ser Ser Glu Glu His Leu
 His Leu Phe Ala Ser Ile Lys Gly Leu Pro Pro Ser Ser Ile Lys
 Ser Ile Ala Glu Lys Leu Leu Val Asp Val Lys Leu Thr Gly Ser
 10 Ala Lys Ile Arg Ala Gly Ser Tyr Ser Gly Gly Met Lys Arg Arg
 Leu Ser Val Ala Ile Ala Leu Ile Gly Asp Pro Lys Leu Val Phe
 Leu Asp Glu Pro Thr Thr Gly Met Asp Pro Ile Thr Arg Arg His
 Val Trp Asp Ile Ile Gln Glu Ser Lys Lys Gly Arg Ala Ile Ile
 Leu Thr Thr His Ser Met Glu Glu Ala Asp Ile Leu Ser Asp Arg
 15 Ile Gly Ile Met Ala Lys Gly Arg Leu Arg Cys Ile Gly Thr Ser
 Ile Arg Leu Lys Ser Arg Phe Gly Thr Gly Phe Val Ala Thr Val
 Ser Phe Ile Glu Asn Lys Lys Asp Gly Ala Pro Glu Pro Leu Lys
 Arg Phe Phe Lys Glu Arg Leu Lys Val Glu Pro Thr Glu Glu Asn
 Lys Ala Phe Met Thr Phe Val Ile Pro His Asp Lys Glu Gln Leu
 20 Leu Lys Gly Phe Phe Ala Glu Leu Gln Asp Arg Glu Ser Glu Phe
 Gly Ile Ala Asp Ile Gln Leu Gly Leu Ala Thr Leu Glu Glu Val
 Phe Leu Asn Ile Ala Arg Arg Ala Glu Leu Glu Ser Ala Thr Val
 Glu Gly Thr Met Val Thr Leu Glu Leu Glu Ser Gly Ile Ala Val
 Glu Ile Pro Val Gly Ala Arg Phe Val Gly Ile Pro Gly Thr Glu
 25 Asn Ala Glu Asn Pro Arg Gly Leu Met Val Glu Val Tyr Trp Gln

Gln Asp Gly Ser Gly Ser Met Cys Ile Ser Gly His Ser Ala Glu
 Met Arg Ile Pro Glu Asn Val Ser Val Ile Tyr Glu Pro Ser Ser
 Gln Val Leu Gly His Gly Gln Arg Arg Val Arg Gly Ile Val Ile
 Asp Tyr Glu Ser Asn Asn

5 SEQ. ID. NO. 2 : AtABCA9/AtATH11 gene

```

      atgactctgc gagaaggctt gccgcttttt catcagcaat tcacagcttt
gttcaagaag aatctactgc tttcatggag gaacaagaga gccacgtgtc
ttcatctctt ctcttcggtc ttcttcatcc ttctcatctt ctctatagag
gaatcttcca aagcgagcga cttaacctcg actaggcaca agaatgtcac
10 agatcctaaa gcattagtct ctcttccgat tcttccatgt gaggataagt
ttttcgtgag acttccatgt tttgacttcg tgtggagtgg caaccagagc
cgtcgtgtca ctgacattgt ctctgcaatt atggctaaca atcctggacg
accaattcca accaataagg ttcaatcatt taaaagcct gaggaagtag
atgcatggtt tatgtcacat ccatcgcaag taacaggggc tttgcatttt
15 gtggaaaaga atgctacagt gatcagctat ggaattcaaa caaactcttc
atcggagaaa aaacgtggtc ggcgtaaga tcctacgttt aagttccttg
ttcctcttca aattgctgca gagcgtgaaa tcgcaaggtc ttttaattgga
gatccaaagt ttagttggga ttttggattt aaggaatttg cgcgtccagc
gattggtggt gaagtgatca tttctgcatt ttatcttatg ggaccagtgt
20 tctttcttgc tttctccatg tttggttttg ttctccaact cgggtctgtg
gttaccgaga aagagctaaa acttcgcgag gcaatgacaa cgatgggtgt
ttatgaatct gcatattggt tgatcatggc catatgggaa ggaatcctta
cctttgtctc ctactcttc ttggtcctct ttggaatgat gttccagttt
gagtttttct tgaagaacag ttttgttctt gtcttcctac ttttctttct
25 ttttcagttt aatatgattg gcctagcatt cgcgctatca tctatcatta

```

	gcaaatacatc	ttcggcaaca	actggttggt	tccttggtgt	tctggttggt
	tttataacac	agattgtaac	aactgctgga	ttcccttatt	caagcgcata
	ttcgattggt	agccgtgtca	tttggtcact	ctttccaccg	aatacctttt
	ctgcgggtct	gcagctgctt	cttgaagcga	catcatctcc	cggagactct
5	ggaatcagtt	ggagtgaaag	agcaatatgt	gcagggggcg	agagcacttg
	cgttattaca	actaataaaa	tctacatatg	gctcgtggga	acgttctttt
	tctggtttgt	attggctctc	tactttgaca	acatcatccc	caatgcatcc
	ggtgtgagaa	aatcgatctt	ctactttcta	aaacctagtt	attggactgg
	caaagaaggc	aacaaagtgg	aagaaggag	catctgtagc	tgtattggtt
10	cagttccacc	agtagagcat	attacaccag	aggacgaaga	tgtgcttgaa
	gaggagattt	tagttaaaca	acaagcaatg	gatggaagag	ttgatcctaa
	cattgcagtt	cagatacatg	gtcttgcaaa	gacatatcct	ggaacaacaa
	agcttggaatg	ctgcaaatgc	acaaaaactt	cgccttttca	tgctgtaaag
	ggtttggtga	tgaatattgc	caaagatcag	ttgttttgtc	ttctcggacc
15	taatggcgca	gggaaaacaa	ctactattag	ttgtttgact	ggcataaatc
	cagtcactgg	tggggatgca	aaaatctatg	gaaattccat	aagaagctct
	gttggtatgt	ccaacattcg	taaaatgata	ggagtttgtc	ctcagtttga
	tattcttttg	gatgctttgt	ctagtgaaga	gcacctccac	ctctttgcta
	gcatcaaagg	gttgccacca	tcatcgatca	aatcgattgc	agagaagtta
20	ctggtagatg	tgaagctaac	aggatcggcg	aaaattagag	caggaagtta
	cagtgggtga	atgaaacgtc	ggctgagtgt	tgcaatagca	ctcattggtg
	atcccaagct	ggtttttcta	gatgaaccga	ctactggcat	ggaccctatc
	acgaggagac	atgtgtggga	cattatacaa	gagtcaaaga	aaggtcgtgc
	catcatacta	acgacgcatt	ctatggagga	agctgatatt	ttaagtgatc
25	gaatagggat	catggctaaa	ggcaggctcc	gctgcattgg	aacctcaatc

```

    aggttaaaat   ctcgcttttg   cacgggattt   gttgctaccg   ttagcttcat
    cgaaaacaaa   aaagacggtg   cacccgagcc   attgaaaaga   ttctttaagg
    agcgtctaaa   agttgagcca   acagaagaaa   acaaagcttt   catgactttt
    gtaatcccac   acgacaaaga   gcaacttttg   aaggggtttt   tcgcggagct
5   acaagataga   gaatctgaat   ttggtatcgc   agacattcag   ctcggtcttg
    ccactcttga   agaagtgttt   ttgaacatcg   ctagacgtgc   tgaactagaa
    agcgcaactg   ttgaaggaaac   tatggtaact   ctcgagttag   aatcaggcat
    cgcagtcgag   atacctgtgg   gagcaagatt   tgtaggtatc   cctggaacag
    aaaacgcaga   gaatccaaga   ggactaatgg   tggaagtgtg   ctggcaacaa
10  gacgggtcag   gatcgatgtg   catttctgga   cactcggcgg   agatgcggat
    cccagagaat   gtatcgggtg   tatatgaacc   atcatcacao   gtattaggac
    atggacagcg acgagttcgg ggtattgtga ttgattatga atctaacaat taa

```

SEQ. ID. NO. 3 : pROKLBb1

gcgtggaacc gcttgctgca act

15 SEQ. ID. NO. 4 : SALK_058070LP

ctacatatgg ctcggtggaa c

SEQ. ID. NO. 5 : SALK_058070RP

aaagaggtgg aggtgctctt c

SEQ. ID. NO. 6 : SALK_084342LP

20 atgactctgc gagaaggctt

SEQ. ID. NO. 7 : SALK_084342RP

gaaagagacc aaaccacacc

SEQ. ID. NO. 8 : AtATH11-ATG primer

actagtatga ctctgcgaga aggctt

25 SEQ. ID. NO. 9 : AtATH11-Stop primer

actagtttaa ttgtagatt cataatca

SEQ. ID. NO. 10 : AtABCA9/AtATH11 F1 primer

atgactctgc gagaaggctt

SEQ. ID. NO. 11 : AtABCA9/AtATH11 R1 primer

5 catggaagaa tcggaagaga

WHAT IS CLAIMED IS:

1. A composition for increasing seed size of a plant or content of fat stored within the seeds, wherein seed size and content of fat stored within the seeds are increased in comparison to seeds of the same plant that has not been subjected to said composition, the composition comprising a transformation vector comprising:

a gene encoding a plant ATP-binding cassette A9 (ABC A9) transporter protein;
and

a promoter for overexpressing said gene;

wherein the ABC A9 transporter protein is a polypeptide having an amino acid sequence of SEQ ID NO: 1.

2. The composition of claim 1, wherein the gene has a base sequence of SEQ ID NO: 2.

3. The composition of claim 1 or 2, wherein the fat stored in the seeds is a triacylglycerol.

4. The composition according to any one of claims 1 to 3, wherein the composition further comprises a microorganism transformed by said transformation vector.

5. The composition of any one of claims 1 to 4, wherein the plant is rice, wheat, barley, corn, soybean, red bean, oat, sorghum, onion, carrot, cucumber, olive, sweet

potato, potato, Chinese cabbage, radish, lettuce, broccoli, tobacco, petunia, sunflower, leaf mustard, turf, *Arabidopsis thaliana*, *Brassica campestris*, *B. napus*, *Betula platyphylla*, poplar, hybrid poplar, or *Betula schmidtii*.

5 6. A method for increasing seed size of a plant or content of fat stored within the seeds, wherein seed size and content of a fat stored within the seeds are increased in comparison to seeds of the same plant that has not been subjected to the step outlined below, the method comprising the steps of:

 inserting a gene encoding the plant ATP-binding cassette A9 (ABC A9)
10 transporter protein and a promoter for overexpressing the gene into a transformation vector; and

 inserting the transformation vector into the plant;

 wherein the ABC A9 transporter protein is a polypeptide having an amino acid sequence of SEQ ID NO: 1.

15

 7. The method of claim 6, wherein the gene has a nucleotide sequence of SEQ ID NO: 2.

 8. The method of claim 6 or 7, wherein the fat stored in the seeds is a
20 triacylglycerol.

 9. The method of any one of claims 6 to 8, wherein the plant is rice, wheat, barley, corn, soybean, red bean, oat, sorghum, onion, carrot, cucumber, olive, sweet potato,

potato, Chinese cabbage, radish, lettuce, broccoli, tobacco, petunia, sunflower, leaf
mustard, turf, *Arabidopsis thaliana*, *Brassica campestris*, *B. napus*, *Betula platyphylla*,
poplar, hybrid poplar, or *Betula schmidtii*.

5

*


*

*

FIGURE
1/11

FIG. 1

a



b




FIGURE
2/11

FIG. 2





FIGURE
3/11

FIG. 3

1% sucrose



no sucrose





FIGURE
4/11

FIG. 4

a



b





FIGURE
5/11

FIG. 5

a



b





FIGURE
6/11

FIG. 6

a



b




FIGURE
7/11

FIG. 7




FIGURE
8/11

FIG. 8






FIGURE
9/11

FIG. 9

a



b



c




FIGURE
10/11

FIG. 10

MTLREGLPLFHQQFTALFKKNLLLSWRNKRATCLHLFSSFFFILLIFSIBESSKASDL
TSTRHKNVTDPKALVSLPILPCEDKFFVRLPCFDFVWSGNQSRRTDIVSAIMANNP
GRPIPTNKVQSFTKPEEVDWFMSSHPSQVTGALHFVEKNA TVISYGIQTNSSEKK
RGRREDPTFKFLVPLQIAAREIARSLIGDPKFSWDFGFKEFARPAIGGEVIISAFYLM
GPVFFLAFSMFGFVLQLGSVVTEKELKLREAMTTMGVYESAYWLSWLIWBGILTFV
SSLFLVLRGMMFQFBFFLKNSFVLVFLFFLFOFNMI GLAFALSSISKSSSATTVGFL
VFLVGFTQIVTTAGFPYSSAYSIGSRVIWSLFPNTFSAGLQLLLEATSSPGDSGISW
SERAICAGGESTCVITTNKIYIWLVGTFRRFWFVLALYFDNIIPNASGVRKSIFYFLKPS
YWTGKEGNKVEEGSICSCIGSVPPVEHITPEDEDVLEEEILVKQQAMDGRVDPNIAVQ
IHGLAKTYPGTTKLGCKCTKTSPFHAVKGLWMNIAKDQLFCLLGPNGAGKTTTIS
CLTGINPVTGGDAKIYGNSSIRSSVGMSNIRKMIGVCPQFDILWDALSSEEHLHLFASIK
GLPPSSIKSIAEKLLVDVKLTGSAKIRAGSYSGGMKRRLSVAIALIGDPKLVFLDEPTT
GMDPI TRRHVWDIIQESKKGRAILLTTHSMEEADILSDRIGIMAKGRLRCIGTSIRLKSR
FGTGfVATVSFIENKKDGAPEPLKRFFKERLKVEPTENKAFMTFVIPHDKEQLLKG
FFAELQDRESEFGIADIQLGLATLEEVFLNIARRAELESA TVEGTMVTLELESGIAVEI
PVGARFVGIPGTENAENPRGLMVEVYWQQDGSGSMCISGHS AEMRIPENVSVIYBPS
SQVLGHGQRRVRGIVIDYESNN

FIGURE
11/11

FIG. 11

ATGACTCTGCGAGAAGGCTTGCCGCTTTTTTCATCAGCAATTCACAGCTTTGTTCAAGAA
GAATCTACTGCTTTTCATGGAGGAACAAGAGAGCCAGGTGCTTCAATCTCTCTCTTCGT
TCTTCTTCATCCTTCTCATCTTCTCTATAGAGGAATCTTCCAAAGCGAGCGACTTAACC
TCGACTAGGCACAAGAATGTCACAGATCTTAAAGCATTAGTCTCTCTTCCGATTCTTCC
ATGTGAGGATAAGTTTTTCGTGAGACTTCCATGTTTTGACTTCGTGTGGAGTGGCAACC
AGAGCCGTCTGTCACTGACATTGTCTCTGCAATTAATGGCTAACAAATCCTGGACGACCA
ATTCCAACCAATAAGGTTCAATCATTACAAAGCCTGAGGAAGTAGATGCAATGGTTTA
TGTCACATCCATCGCAAGTAACAGGGGCTTTGCAATTTGTGAAAAGAATGCTACAGTG
ATCAGCTATGGAATTCAAACAAACTCTTCATCGGAGAAAAACGTGGTCCGCGTGAAGA
TCCTACGTTTAAAGTTCTTGTTCCTCTTCAATTTGCTGCAGAGCGTGAAATCCGAAGGT
CTTTAATTGGAGATCCAAAGTTTAGTTGGGATTTTGGATTTAAGGAATTTGCGGTCC
AGCGATTGGTGGTGAAGTGATCAATTTCTGCAATTTATCTTATGGGACCAGTGTCTTT
CTTGCTTTCTCCAATGTTTGGTTTGTCTCTCAACTCGGGTCTGTGGTTACCGAGAAAGA
GCTAAACCTTCGCGAGGCAATGACAACGATGGGTGTTTATGAATCTGCATATTGGTTGT
CATGGCTCATATGGGAAGGAATCCTTACCTTTGCTCTCTCACTCTTCTTGGTCTCTTT
GGAATGATGTTCCAGTTTGAGTTTTCTTGAAGAACAGTTTTGTTCTTGTCTTCTCTAC
TTTTCTTTCTTTTTTCAGTTTAAATATGATTGGCCTAGCATTCGCGCTATCATCTATCAT
TAGCAAAATCATCTTCGCAACAACCTGTTGGTTTTCTTGTGTTTCTGGTTGGTTTTATA
ACACAGATTGTAACAACCTGCTGGAATCCCTTATTCGAAGCGCATATTCGATTGGTAGCCG
TGTCATTTGGTCACTCTTTCCACCGAATACCTTTTCTGCGGCTCTGCAGCTGCTTCTTG
AAGCGACATCATCTCCCGGAGACTCTGCAATCAGTTGGAGTGAAAGAGCAATATGTGCA
GGGGCGAGAGCACTTGCGTTATTACAACATAAATCTACATATGGCTCGTGGGAAC
GTTCTTTTCTGGTTTGTATTGGCTCTCTACTTTGACAACATCATCCCAATGCATCCG
GTGTGAGAAAAATCGATCTTCTACTTTCTAAAACCTAGTTATTGGACTGGCAAGAAGG
CAACAAAGTGGAAGAAGGGAGCATCTGTAGCTGTATTGGTTCACTTCCACCAGTAGAGC
ATATTACACAGAGGACGAAGATGTGCTTGAAGAGGAGATTTTACTTAAACAACAAGC
AATGGATGGAAGAGTTGATCCTAACATTGCAAGTTACAGATACATGGTCTTCCAAAGACA
TATCCTGGAACAACAAGCTTGGATGCTGCAAAATGCACCAAACTTCGCTTTTTCATGC
TGTAAGGGTTTGTGGAATGAATATTGCCAAAGATCAGTTGTTTTGTCTTCTCGGACCT
AATGGCGCAGGGAAAAACAACCTACTATTAGTTGTTTGGACTGGCATAAATCCAGTCACTGG
TGGGGATGCAAAAATCTATGGAAATTCATAAGAAGCTCTGTTGGTATGTCCAACATT
CGTAAATGATAGGAGTTTGTCTCAGTTTGATATTCTTTGGGATGCTTTGTCTAGTG
AAGAGCACCTCCACCTCTTTGCTAGCATCAAAGGGTTGCCACCATCATCGATCAAAATCG
ATTGCAGAGAAGTTACTGCTAGATGTGAAGCTAACAGGATCGGCGAAAAATTAGAGCAG
GAAGTTACAGTGGTGGAAATGAAACGTGGGCTGAGTGTGCAATAGCACTCATTTGGTGAT
CCCAAGCTGGTTTTTCTAGATGAACCGACTACTGGCAATGGACCTATCACGAGGAGACA
TGTTGGGACATTATACAAGAGTCAAAGAAAGGTCGTGCCATCATACTAACGACGCATT
CTATGGAGGAAGCTGATATTTTAAGTGATCGAATAGGGATCATGGCTAAAGGCAGGCTC
CGCTGCATTGGAACCTCAATCAGGTTAAATCTCGCTTTGGCACGGGATTTGTTGCTAC
CGTTAGCTTCATCGAAAAACAAAAAGACGGTGCACCCGAGCCATTGAAAAGATTCTTTA
AGGAGCGTCTAAAAGTTGAGCCAAACAGAAAGAAAGCTTTCAATGACTTTTCTAAT
CCCACACGACAAAGAGCAACTTTTGAAGGGGTTTTTCGGGAGCTACAAGATAGAGAAT
CTGAATTTGGTATCGCAGACATTGAGCTCGGTCTTGGCACTCTTGAAGAAGTGTTTTTG
AACATCGCTAGACGTGCTGAATAGAAAGCGCAACTGTTGAAGGAACATGGTAACCTCT
CGAGTTAGAAATCAGGCATCGCAGTCGAGATACCTGTGGGAGCAAGATTTGTAGGTATCC
CTGGAACAGAAAAACCGCAGAGAAATCCAAGAGGACTAATGGTGGAGGTGTAAGTGGCAACA
GACGGGTGAGGATCGATGTGCAATTTCTGGACACTCGGCGGAGATCGGATCCCAAGAGAA
TGTATCGGTGATATATGAACCATCATCAACAAGTATTAGGACATGGACAGCGACGAGTTC
GGGGTATTGTGATTGATTAATGAATCTAACAAATTA