



(86) Date de dépôt PCT/PCT Filing Date: 2011/07/01

(87) Date publication PCT/PCT Publication Date: 2012/01/05

(45) Date de délivrance/Issue Date: 2021/01/19

(85) Entrée phase nationale/National Entry: 2012/12/28

(86) N° demande PCT/PCT Application No.: US 2011/042834

(87) N° publication PCT/PCT Publication No.: 2012/003468

(30) Priorité/Priority: 2010/07/02 (US61/361,151)

(51) Cl.Int./Int.Cl. *C07D 489/08* (2006.01)


(72) Inventeurs/Inventors:
KESKENY, ERNO M., US;
MENCEL, JAMES J., US;
DUNG, JEN-SEN, US

(73) Propriétaire/Owner:
JOHNSON MATTHEY PUBLIC LIMITED COMPANY, GB

(74) Agent: SMART & BIGGAR LLP

(54) Titre : PROCEDE DE SYNTHESE ET DE PURIFICATION D'OXYCODONE

(54) Title: PROCESS FOR THE SYNTHESIS AND PURIFICATION OF OXYCODONE



(57) Abrégé/Abstract:

A method of preparing oxycodone includes forming 14-hydroxycodeine by reduction of 14-hydroxycodeinone and rearrangement of the 14-hydroxycodeine to form the oxycodone. During the reduction step, the ketone group of an undesirable contaminant precursor, 8, 14-dihydroxy-7,8-dihydrocodeinone, is reduced to a hydroxyl group thus forming a triol. This triol is substantially inert with respect to reforming 14-hydroxycodeinone and can be readily separated from oxycodone.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
5 January 2012 (05.01.2012)

PCT

(10) International Publication Number
WO 2012/003468 A1(51) International Patent Classification:
C07D 489/08 (2006.01)(21) International Application Number:
PCT/US2011/042834(22) International Filing Date:
1 July 2011 (01.07.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/361,151 2 July 2010 (02.07.2010) US(71) Applicant (for all designated States except US): **JOHNSON MATTHEY PUBLIC LIMITED COMPANY**
[GB/GB]; 5th Floor, 25 Farringdon Street, London EC4A 4AB (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KESKENY, Erno, M.** [US/US]; 1021 Trenton Place, Wilmington, DE 19801 (US). **MENCEL, James, J.** [US/US]; 103 Usher Lane, North Wales, PA 19454 (US). **DUNG, Jen-Sen** [US/US]; 2105 Sunnyside Lane, Boothwyn, PA 19061 (US).(74) Agents: **LEWIS, Christopher, R.** et al.; RatnerPrestia,
P.O. Box. 980, Valley Forge, PA 19482 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: PROCESS FOR THE SYNTHESIS AND PURIFICATION OF OXYCODONE

(57) Abstract: A method of preparing oxycodone includes forming 14-hydroxycodone by reduction of 14-hydroxycodone and rearrangement of the 14-hydroxycodone to form the oxycodone. During the reduction step, the ketone group of an undesirable contaminant precursor, 8, 14-dihydroxy-7,8-dihydrocodeinone, is reduced to a hydroxyl group thus forming a triol. This triol is substantially inert with respect to reforming 14-hydroxycodone and can be readily separated from oxycodone.



WO 2012/003468 A1

- 1 -

PROCESS FOR THE SYNTHESIS AND PURIFICATION OF OXYCODONE**BACKGROUND OF THE INVENTION**

Various reports have identified the role of two isomers of 8,14-dihydroxy-7,8-dihydrocodeinone in the formation of 14-hydroxycodeinone - an α,β -unsaturated ketone ("ABUK") and a purportedly genotoxic impurity in oxycodone hydrochloride. For example, U.S. Patent No. 7,683,072 of Chapman et al. notes that "[d]uring salt formation reactions known in the art, the 8,14-dihydroxy-7,8-dihydrocodeinone component is converted to 14-hydroxycodeinone by acid-catalyzed dehydration. See also Weiss, J. Org. Chem., 22(11): 1505-08 (1957). Thus, 14-hydroxycodeinone is increased in the final product." (col. 8, lines 7-12). Similarly, Cox (WO 2008/070656 and WO 2008/070658) notes that "DHDHC [8,14-dihydroxy-7,8-dihydrocodeinone] is easily converted to 14-hydroxycodeinone. This conversion occurs during the conversion of oxycodone base (a.k.a. oxycodone free base) to oxycodone hydrochloride, thus 14-hydroxycodeinone is present in the final oxycodone hydrochloride." (p. 3, lines 5-7). Thus, the isomeric 8,14-dihydroxy-7,8-dihydrocodeinone diols are understood to be unstable such that, during conversion of the oxycodone free base to oxycodone hydrochloride, they undergo acid-catalyzed dehydration to form the 14-hydroxycodeinone ABUK impurity. Weiss teaches that 8,14-dihydroxy-7,8-dihydrocodeinone can be recrystallized from hot 2N HCl if the treatment is rapid, but also reports that treatment in "dilute acid (1:1) in a boiling water bath for 20 minutes" converts the dihydroxy species to 14-hydroxycodeinone. While Weiss does not indicate the mixture of dihydroxy isomers handled during the 2N recrystallization and subsequent hydrolysis to 14-hydroxycodeinone, it implies that both isomers of 8,14-dihydroxy-7,8-dihydrocodeinone will dehydrate to form 14-hydroxycodeinone. Recent reports, including the Chapman '072 patent, and Baldwin (see Baldwin Declaration in support of U.S. Pat. Appln. No. 11/729,741) have pointed to the alpha isomer as the apparent more hydrolytically unstable diol of 8,14-dihydroxy-7,8-dihydrocodeinone. Baldwin cites Example 3 of Chapman's U.S. Patent No. 7,674,800 as demonstrating that dehydration occurs under conditions of aqueous HCl as low as 0.2N at 75°C. Baldwin contrasts this with the Weiss report to assert that one of the two isomers is much more prone to acid catalyzed dehydration.

- 2 -

The present inventors prepared and characterized the 8 β ,14-dihydroxy-7,8-dihydrocodeinone and used this to develop analytical methodology to measure the alpha and beta isomers by LC/MS to a detection limit of 2 ppm. Two HPLC peaks at relative retention times (RRT) 0.82 and 0.91 versus oxycodone were observed as having a mass
5 corresponding to 8,14-dihydroxy-7,8-dihydrocodeinone upon LC/MS analysis. The peak at RRT 0.91 was identified as the 8 β diol. Exposure of a mixture of the two species to acid showed a correlation between the loss of the RRT 0.82 peak with the growth of 14-hydroxycodeinone over a 20 hour period. The peak at RRT 0.91 degraded little under the same conditions over the 20 hour period. On the basis of this study and in view of
10 Baldwin's assertions, the peak at RRT 0.82 was assigned as the 8 α diol for the purposes of developing the invention.

A variety of approaches to producing oxycodone hydrochloride with low levels of 14-hydroxycodeinone have been published. The Chapman '072 patent, as well as other patents in the Chapman patent family, and U.S. Patent Application Publication No.
15 2007/0149559 to Shafer et al., for example, report methods wherein the diols, as contaminants in oxycodone, are dehydrated using aqueous acid or organic acids in organic solvents to form 14-hydroxycodeinone. Chapman describes converting the 14-hydroxycodeinone to oxycodone by hydrogenation as the dehydration occurs. This prevents reversal by rehydration of the 7,8 double bond and minimizes the possibility of having 8,14-
20 dihydroxy-7,8-dihydrocodeinone as an ABUG precursor in the final oxycodone hydrochloride product. Shafer traps the 14-hydroxycodeinone with a thiol nucleophile, forming a water-extractable adduct that can be readily separated from oxycodone.

International Patent Application Publication No. WO 2007/103105 of Buehler et al. likewise describes conditions that may dehydrate the diols to an extent and, in
25 a fashion similar to Shafer, treats the 14-hydroxycodeinone with a sulfur nucleophile or a sulfur-containing inorganic acid or salt thereof to form a water-soluble adduct. The Buehler procedure uses species akin to *meta*-bisulfite, which react with oxycodone in a 1,2 fashion and with 14-hydroxycodeinone in a 1,4 and 1,2 fashion to form soluble adducts of each. Upon a pH adjustment to ca. pH 9, the 1,2 sulfite adducts hydrolyze to the ketone; however,
30 the 1,4 adduct derived from 14-hydroxycodeinone remains intact and confers water solubility. At pH 9, the oxycodone precipitates or may be preferentially extracted into organic solvent and the adducted 14-hydroxycodeinone remains in the aqueous mixture.

The Shafer approach differs in emphasizing the formation of the 1,4 adduct of 14-hydroxycodone and does not explicitly rely upon selective 1,2 hydrolysis to recover and separate the oxycodone from the water-soluble 1,4 adduct.

5 The derivatization of 14-hydroxycodone with a sulfur-based reagent, either as a free species or resin-bound, is common to Shafer, Cox and Buehler. At least Chapman and Shafer find commonality in: (1) forcing the dehydration of the diols in the presence of oxycodone; and (2) subsequently consuming or removing the resulting 14-hydroxycodone either as oxycodone or as a derivative separable from oxycodone.

SUMMARY OF THE INVENTION

10 The inventors have found that the alpha diol of 8,14-dihydroxy-7,8- dihydrocodeinone can be present in up to 1,000 ppm in oxycodone base, and that the beta diol can be present at even higher levels. In essence, the approaches outlined above have the potential to actively create and contaminate oxycodone hydrochloride with over 1,000 ppm of the undesired, purportedly genotoxic, impurity then require measures to rigorously remove the impurity.

15 There is also the chance that measures to destroy diol precursors may not proceed exhaustively as noted, for example, in Example 3 of U.S. Patent No. 7,129,248 of Chapman et al., where a cautionary note appears that purified oxycodone salt must be handled with care to avoid dehydrating remaining diols to reform 14-hydroxycodone and contaminate the product.

The above approaches include the conversion of the diols to 14-hydroxycodone in the presence of oxycodone free base. A variety of chemical approaches are used to remove the

20 14-hydroxycodone, especially during conversion of the oxycodone free base to oxycodone hydrochloride. Thus, the above approaches include the formation of 14-hydroxycodone - the actual ABUG impurity one is trying to eliminate - during the formation and in the presence of oxycodone, and in some cases, oxycodone hydrochloride. Accordingly, new methods of producing oxycodone and its salts with low levels of impurities, including ABUG, would be

25 beneficial.

A novel process is now disclosed for producing oxycodone and its salts, especially, oxycodone hydrochloride, with low levels of impurities, including ABUG, that diverges from

the approaches described above.

In one broad aspect, the present specification discloses a method of preparing oxycodone, comprising contacting 14-hydroxycodeine with a catalyst under conditions sufficient to cause rearrangement of the 14-hydroxycodeine to form oxycodone free base.

5 In another broad aspect, the present specification discloses an oxycodone free base comprising less than 2 ppm of 8 α ,14-dihydroxy-7,8-dihydrocodeinone and less than 5 ppm 14-hydroxycodeinone.

 In yet another broad aspect, the present specification discloses an oxycodone hydrochloride having less than 25 ppm 14-hydroxycodeinone and less than 2 ppm of 8 α ,14-
10 dihydroxy-7,8-dihydrocodeinone.

In brief, the inventors have found, among other things, that reduction of the ketone group of 14-hydroxycodeinone to afford 14-hydroxycodeine also reduces the ketone groups of 8,14-dihydroxy-7,8-dihydrocodeinone, thereby converting the diols to 8,14-dihydroxy-7,8-dihydrocodein-6-ol. An exemplary sequence of steps according to preferred embodiments of the invention is shown in Schemes 1 and 2. As shown, after oxidation and reduction steps, a rearrangement step for producing oxycodone free base is used to significantly reduce or eliminate impurities, including especially ABUK, prior to converting the oxycodone free base to its salt. Also described are several optional purification steps, and a salt formation step.

10 BRIEF DESCRIPTION OF THE DRAWINGS

A detailed description of specific exemplary embodiments is provided herein below with reference to the accompanying drawings in which:

FIG. 1 shows a typical chromatogram using 0.2% TFA/H₂O as Blank;

FIG. 2 shows a typical chromatogram using a Resolution Solution (10 PPM ABUK Spiked);

15 FIG. 3 shows a typical chromatogram using a Resolution Solution (10 PPM .beta.-Diol Spiked);

FIG. 4 shows a typical chromatogram using a Sample Solution (Containing about 3 PPM of 14-Hydroxy Codeinone);

20 FIG. 5 shows a typical chromatogram using a Sample Solution (Containing about 30 PPM of beta-Diol).

In the figures, exemplary embodiments are illustrated by way of example. It is to be expressly understood that the description and figures are only for the purpose of illustrating certain embodiments and are an aid for understanding. They are not intended to be a definition of the limits of the invention.


DETAILED DESCRIPTION OF THE INVENTION

It is known that oxidation reactions of commercially available varieties of thebaine will produce 14-hydroxycodeinone. This is described, for example, by Chapman and Cox as well
5 as in U.S. Patent No. 7,153,966 of Casner. During oxidation, especially under aqueous acidic conditions, the two isomers of 8,14-dihydroxy-7,8-dihydrocodeinone are also formed as impurities. See, e.g., Proska, Arch. Pharm. Pharm. Med. Chem., 332, 369-70 (1999). Next, the ketone of 14-hydroxycodeinone is reduced to form 14-hydroxycodeine and, as a consequence of the reduction, the ketones of the two isomers of 8,14-dihydroxy-7,8-dihydrocodeinone are
10 also reduced, creating related 8,14-dihydroxy-7,8-dihydrocodein-6-ol isomers, or triols. Only traces of the two isomers of the diols remain and these are purged when the oxycodone base is purified prior to salt formation.


In a novel metal-catalyzed process, 14-hydroxycodeine is rearranged to oxycodone in a reaction whereby the 6-hydroxyl group is oxidized and the 7,8 double bond is reduced.
15 Because the two isomers of 8,14-dihydroxy-7,8-dihydrocodein-6-ol do not have the allylic C6, C7, C8 topography contained within 14-hydroxycodeine, they do not participate in the rearrangement. The triols are observed in the resulting oxycodone. The triols also do not appear to interfere and therefore cannot convert to 14-hydroxycodeinone. In this sense, they are inert with regard to conversion to either 14-hydroxycodeinone or oxycodone. Moreover,
20 the triols are far more polar than oxycodone and can potentially be separated from oxycodone by precipitating the oxycodone from a polar solvent or solvent mixture, or by extracting oxycodone into a suitable organic solvent. Notwithstanding, even if some of the triols were to remain with oxycodone, these species lack the purportedly genotoxic alpha, beta unsaturated ketone, cannot generate it, and do not fall under FDA alert

compound status. When the resulting oxycodone base is converted to the hydrochloride salt, the active, alpha diol is not present and any traces of the triols are incapable of generating 14-hydroxycodeinone, so the salt can be produced without risk of regenerating that impurity.

5 **Scheme 1. Synthetic Route To Oxycodone Base and Oxycodone Hydrochloride**



Scheme 2: Alternative Routes from Crude Oxycodone Base to Low ABUG Oxycodone HCl




5

Oxidation of thebaine to 14-hydroxycodeinone may be performed by any means known in the art, for example as described in U.S. Patent Application Publication No. 2006/0111383 to Casner et al. Reduction of 14-hydroxycodeinone to 14-hydroxycodine may involve any method known in the art for reducing ketone groups of

alpha, beta unsaturated ketones, with sodium borohydride being one example as described below.

Suitable catalysts for Step 3 of Scheme 1 include any of a variety of ligand-complexed metal catalyst, including such catalysts wherein the ligand is a phosphine and the metal is rhodium or ruthenium. Wilkinson's catalyst and analogs thereof are generally
5 suitable. Specific examples of suitable catalysts are disclosed in U.S. Patent No. 7,323,565 and U.S. Patent No. 7,321,038 to Wang et al. One suitable catalyst is shown below, the preparation and use of which is detailed in the Examples herein.



10 The new route to oxycodone enjoys several advantages over the aforementioned routes as per Chapman, Shafer, Buehler, and Cox. The process converts known precursors of the purportedly genotoxic 14-hydroxycodeinone to triol species that are incapable of giving rise to this impurity. This conversion may be performed concurrently with the conversion of 14-hydroxycodeinone to 14-hydroxycodeine (i.e.,
15 before converting the 14-hydroxycodeine to oxycodone). The 14-hydroxycodeine may be isolated as a solid or extracted into a suitable organic solvent, either approach allowing the opportunity to purge these triols a full chemical stage before oxycodone is formed. In some examples trace levels of 8β,14-dihydroxy-7,8-dihydrocodeinone are observed by LC/MS in the isolated 14-hydroxycodeine; however, 8α,14-dihydroxy-7,8-dihydrocodeinone is not
20 detectable at this stage of the process. In a further embodiment of the invention, the 14-hydroxycodeine may be treated with sodium *meta*-bisulfite as per Rappoport et al., J. Amer. Chem. Soc. 89:8, 1967, 1942-1947,1 or similar species as a means to remove any remaining traces of 14-hydroxycodeinone or of the ketone-containing diols. In still a further embodiment, the 14-hydroxycodeine may be treated with resin-supported
25 borohydride to remove any trace amounts of 14-hydroxycodeinone. In yet another embodiment of the invention, residual 14-hydroxycodeine may be treated with organic thiol species to selectively form water-soluble or resin bound derivatives of any remains 14-

- 8 -

hydroxycodone such that 14-hydroxycodone may be produced free of 14-hydroxycodone. By means of the reducing step and, optionally, the incorporation of treatments using *meta*-bisulfite-related or thiol-type species, 14-hydroxycodone substantially free of residual 14-hydroxycodone and of the two diols can be produced. In
5 another embodiment, any trace amounts of 14-hydroxycodone in 14-hydroxycodone are carried through the rearrangement step and removed from the oxycodone base by catalytic hydrogenation in an organic solvent.

As will be understood from the above explanation, diols that are potential precursors to 14-hydroxycodone are not converted to that compound, but instead are
10 converted to species that are incapable of giving rise to 14-hydroxycodone. In this fashion, the drug entity is not actively contaminated with the purportedly genotoxic impurity one is also trying to remove.

It is noteworthy that 14-hydroxycodone is both the undesired, purportedly genotoxic species and the penultimate intermediate common to most traditional syntheses of
15 oxycodone. Conventional synthesis routes rely upon hydrogenation of 14-hydroxycodone to form oxycodone, and the hydrogenation may often be left incomplete such that large amounts of 14-hydroxycodone – 1,000-5,000 ppm – are carried forward directly into the isolated oxycodone. See, e.g. U.S. Patent No. 7,674,800, Examples 2 and 3). The aforementioned approaches may thus be burdened by the need to overcome this
20 reservoir of 14-hydroxycodone carried forward along with that formed by forced dehydration of the diols. In contrast, the new process converts 14-hydroxycodone to a new, non-ketone intermediate (i.e., 14-hydroxycodone) with zero or low single digit ppm levels of remaining 14-hydroxycodone. Should non-zero levels of 14-hydroxycodone remain, these can be removed as described above using *meta*-bisulfite or organic thiol
25 agents. Alternatively, 14-hydroxycodone can be removed from oxycodone base by catalytic hydrogenation in an organic solvent or a mixture of solvents.

The new route employs what is believed to be a novel rearrangement of a C-14 oxidized codeine-like entity to oxycodone. There are many published reports for the rearrangements of 6-hydroxy 7,8 double bond allylic systems such as found in codeine to
30 form, for example, hydrocodone. However, the inventors are not aware of any reports of a similar rearrangement of related species bearing the 14-hydroxy group. Indeed, 14-hydroxycodone, the intermediate in the new process disclosed here, is a “bisallylic” entity

- 9 -

to the extent that the 7,8 double bond and 14-hydroxylic moiety constitute a second allylic function. The lack of reports of metal-catalyzed rearrangements of C-14 hydroxyl analogues of codeine to form the 6-ketone function suggests that those skilled in the art would not expect this to be a particularly productive, or even probable, reaction.

5 The reduction of 14-hydroxycodeinone to 14-hydroxycodeine is expected to entirely convert the diols to the triols. 14-hydroxycodeine containing ca. 0-50 ppm of 14-hydroxycodeinone will undergo a metal catalyzed rearrangement to form oxycodone base with about 45-350 ppm of 14-hydroxycodeinone. LC/MS analysis indicates that the presumed active diol, 8 α ,14-dihydroxy-7,8-dihydrocodeinone, may be present at up to 16
10 ppm while 8 β ,14-dihydroxy-7,8-dihydrocodeinone may be present, with two other species having molecular weights matching the triols, at up to about 110 ppm. The minor increase in 14-hydroxycodeinone is thought to result from very slow oxidation of oxycodone by the metal rearrangement catalyst. This is supported by the fact that the inventors have demonstrated that oxycodone with undetectable (or, at most, single digit ppm) levels of
15 either isomer of 8 α ,14-dihydroxy-7,8-dihydrocodeinone will slowly give rise to 14-hydroxycodeinone upon exposure to the metal rearrangement catalyst under conditions of the rearrangement to levels of over 60 ppm, far higher than levels obtainable from the quantity of diols present. The very low levels of 14-hydroxycodeinone observed in oxycodone formed by the current invention are, however, in sharp contrast to the high levels
20 found in oxycodone produced by the aforementioned processes, as noted above, which may contain 1,000-5,000 ppm of 14-hydroxycodeinone, *and* the potential for more due to the 1000 ppm or more of each of the diol precursors to 14-hydroxycodeinone, i.e., 8 α ,14-dihydroxy-7,8-dihydrocodeinone and 8 β ,14-dihydroxy-7,8-dihydrocodeinone. The traces of the 8 α ,14-dihydroxy-7,8-dihydrocodeinone and 8 β ,14-dihydroxy-7,8-dihydrocodeinone
25 observed after the rhodium catalyzed rearrangement are purged in the next stage of the process while purifying the oxycodone base.

Oxycodone base prepared by the new process, containing ca. 0-350 ppm of 14-hydroxycodeinone, can be rendered virtually free of this impurity by following a purification process such as the sodium *meta*-bisulfite procedure described by Rappoport et
30 al., J. Amer. Chem. Soc. 89:8, 1967, 1942-1947. This affords an oxycodone free base having undetectable (or, at most, very low) levels of 14-hydroxycodeinone. Alternatively, one may reduce the residual 14-hydroxycodeinone to levels under 5 ppm by catalytic

- 10 -

hydrogenation using a palladium on carbon catalyst in an organic solvent or solvent mixture. The hydrogenation may be run at about 13-50 psi and at temperatures from about room temperature (e.g., 18°C) to about 40°C. Suitable catalysts include, but are not limited to, 5% and 10% palladium on carbon. Suitable solvents include alcohols, chlorinated
5 solvents, or mixtures thereof. The isolation of the hydrogenated oxycodone as a solid purges most of any residual trace amounts of 8 α ,14-dihydroxy-7,8-dihydrocodeinone and 8 β ,14-dihydroxy-7,8-dihydrocodeinone.

The exposure of oxycodone to hydrogenation conditions has been found to produce low levels of 6-hydroxy oxycodol. This is removed using an ethanol slurry,
10 preferably at an elevated temperature, with the added benefit of also removing any further trace amounts of 14-hydroxycodeinone, 8 α ,14-dihydroxy-7,8-dihydrocodeinone, and 8 β ,14-dihydroxy-7,8-dihydrocodeinone in the unslurried oxycodone base. Purified oxycodone base produced via a rhodium-catalyzed rearrangement reaction of 14-hydroxycodeine, subsequent hydrogenation and an ethanol slurry contains no detectable amounts of 8 α ,14-
15 dihydroxy-7,8-dihydrocodeinone, ca. 10 ppm of 8 β ,14-dihydroxy-7,8-dihydrocodeinone, and 1-5 ppm of 14-hydroxycodeinone.


In a further embodiment, crude oxycodone base containing traces of 14-hydroxycodeine may be purified by hydrogenation as described above, and used as a solution after catalyst filtration to form and isolate oxycodone HCl salt having less than 5
20 ppm of residual 14-hydroxycodeinone.

Importantly, unlike the Buehler, Cox, Chapman, or Shafer processes, the new process does not require breaking down the diol precursors to form 14-hydroxycodeinone in the presence of oxycodone. There is also no threat that unconverted 8 α ,14-dihydroxy-7,8-dihydrocodeinone will be carried into the oxycodone product and pose a risk of regenerating
25 14-hydroxycodeinone during salt formation, stability testing or upon storage. Again, this is in contrast to aforementioned processes. For example, as noted by Chapman in Example 3, care is required when handling the HCl salt from the base in the route described there due to the possible presence of unconverted diol precursors. These precursors can convert to 14-hydroxycodeinone during salt formation or on storage or stability testing. Because 8 α ,14-
30 dihydroxy-7,8-dihydrocodeinone is undetected in oxycodone base prepared by the new process, no possibility exists to regenerate 14-hydroxycodeinone from it while making the HCl salt. The conversion of oxycodone base produced by the process above, and containing

- 11 -

less than 5 ppm 14-hydroxycodeinone and no detectable 8 α ,14-dihydroxy-7,8-dihydrocodeinone, showed no change in the level of 14-hydroxycodeinone level in the isolated oxycodone HCl salt. Furthermore, treatment of the oxycodone base or oxycodone HCl salt made by the above process in 65°C aq. hydrochloride acid at pH <1 resulted in no or minimal growth in 14-hydroxycodeinone after 24 hours. Exposure of oxycodone base and oxycodone HCl salt made by the above process to conditions of dry heat (greater than 60°C) resulted in no growth of 14-hydroxycodeinone after 14 days. In view of assertions noted above by Baldwin regarding the ease of dehydration of 8 α ,14-dihydroxy-7,8-dihydrocodeinone in weak acid (i.e., 0.2N; US Patent No. 7,674,800, Example 3), the slow, trace growth observed in hot acid is attributable to trace levels of residual 8 β ,14-dihydroxy-7,8-dihydrocodeinone as per Weiss, or to other unknown precursors to 14-hydroxycodeinone. This stability provides wider process latitude during oxycodone salt formation on an industrial scale and suggests a highly stable solid product with regard to growth of 14-hydroxycodeinone.

EXAMPLES

Example 1 - Reduction of 14-hydroxycodeinone to 14-hydroxycodeine


A portion of 14-hydroxycodeinone (26.0 g wet, 16.9 g dry weight, 54.0 mmol) was combined with 135 ml methylene chloride and 15 ml methanol under nitrogen in a round bottomed flask equipped with a thermometer. The slurry was cooled to 0-5 °C in an ice bath. Sodium borohydride (2.54 g, 66.8 mmol) was added to the slurry in one portion. Excess hydrogen was released through a bubbler, and the reaction mixture was allowed to warm up to ambient temperature and was stirred for 48 hours.

After the starting material was consumed, the excess sodium borohydride was quenched with 70 ml 10 wt % hydrochloric acid. The aqueous layer was at pH=1. The two phase mixture was stirred for one hour. The layers were separated and the upper aqueous layer was separated from the lower organic layer. The organic layer was washed

- 12 -

with distilled water (2x20 ml). The aqueous layers were combined and the pH was adjusted to 9 with the addition of 45 ml of 5 M sodium hydroxide solution. An oily precipitate formed and the aqueous slurry was extracted with dichloromethane (3X50 ml). The combined dichloromethane phase was washed with 20 % sodium chloride solution, dried over sodium sulfate, filtered and evaporated yielding 15.1 g (89 %) brown oil which solidified upon standing. Trituration of the solidified oil with cold 2-propanol afforded 12.5 g (74 %) white crystalline 14-hydroxycodeine.

Example 2 - Rearrangement of 14-hydroxycodeine to Oxycodone



A 100 ml 4 necked round bottom flask, equipped with condenser and a gas sparging tube, gas outlet and a thermometer, was flushed with nitrogen and charged with methanol (40 ml). A portion of 14-Hydroxycodeine (5.0 g, 15.9 mmol) was added at room temperature and the mixture was stirred until a homogenous solution formed. The solution was deoxygenated by sparging with nitrogen gas while stirring.

A 20 ml Schlenk flask, equipped with a gas sparging tube and gas outlet, was flushed with nitrogen and charged with methanol (10 ml). The solvent was deoxygenated by sparging nitrogen gas through it while stirring. Bis(norbornadiene)rhodium (I) tetrafluoroborate (70 mg, 0.19 mmol) and 1,4-bis(diphenylphosphino)butane (80 mg, 0.19 mmol) were added under nitrogen and the orange solution was stirred at room temperature for 30 minutes. The solution was then sparged with hydrogen gas for 30 minutes. The color of the solution changed from orange to dark reddish orange. Excess hydrogen was removed by sparging the solution with nitrogen for 10 minutes. The solution of the catalyst was transferred to the solution of 14-hydroxycodeine via cannula. The combined solution was heated at 50 °C for 90 minutes. The product, oxycodone free base, precipitated from the reaction mixture and a thick slurry formed. A small sample was taken for an in-process

- 13 -

test to confirm the reaction completion. The reaction mixture was cooled to 0-5 °C for 2 hours, and the product was filtered off and washed with 10 ml cold 2-propanol.

The product, oxycodone free base, was dried in a vacuum oven at 50 °C.
Yield: 4.1 g, 71.9 %.

5 Example 3 - Removal of 14-hydroxycodeinone from 14-hydroxycodeine

A 100 ml round bottom flask, equipped with gas outlet and a thermometer, was flushed with nitrogen and charged with methanol (40 ml). 14-Hydroxycodeine (5.0 g, 15.9 mmol) containing a small amount of 14-hydroxycodeinone as an impurity was added at room temperature and the mixture was stirred until a homogenous solution formed.


10 Resin supported borohydride (0.15g) was added to the solution and it was stirred for 24 h at 55 °C. A small sample was taken for the in process test to confirm the removal of 14-hydroxycodeinone. The mixture was then filtered through 45 µm filter under nitrogen blanket into a 100 ml 4 necked round bottom flask, equipped with condenser and a gas sparging tube, gas outlet, and a thermometer. The solution was deoxygenated by
15 sparging nitrogen gas through it while stirring, for use in Example 4.

Example 4 - Rearrangement of 14-hydroxycodeine to Oxycodone

A 20 ml Schlenk flask, equipped with a gas sparging tube and gas outlet, was flushed with nitrogen and charged with methanol (10 ml). The solvent was deoxygenated by sparging nitrogen gas through it while stirring. Bis(norbornadiene)rhodium (I)
20 tetrafluoroborate (70 mg, 0.19 mmol) and 1,4-bis(diphenylphosphino)butane (80 mg, 0.19 mmol) were added under nitrogen and the orange solution was stirred at room temperature for 30 minutes. The solution was then sparged with hydrogen gas for 30 minutes. The color of the solution changed from orange to dark reddish orange. Excess hydrogen was removed by sparging the solution with nitrogen for 10 minutes. The solution of the catalyst
25 was transferred to the solution of 14-hydroxycodeine from Example 3 via cannula. The combined solution was heated at 50 °C for 90 minutes. Oxycodone free base precipitated from the reaction mixture and a thick slurry formed. A small sample was taken for an in-process test to confirm the reaction completion. The reaction mixture was cooled to 0-5 °C for 2 hours and the product was filtered off, washed with 10 ml cold 2-propanol.

30 The product, oxycodone free base, was dried in a vacuum oven at 50 °C.
Yield: 3.43 g, 68.6 %.

- 14 -


Example 5 - Reduction of 14-Hydroxycodeinone to 14-Hydroxycodeine

5 14-Hydroxycodeinone (25.0g wet, 22.225g dry weight, 70.93mmol) was combined with 178mL methylene chloride and 13mL methanol under nitrogen in a round bottomed flask equipped with a thermometer and a solution resulted. The batch was cooled to 0-5 °C in an ice bath. Sodium borohydride (3.44g, 90.93mmol) was added in one portion. The mixture was stirred at 0-5°C for 7 hours and then allowed to warm up to ambient

10 temperature and stirred for 17 hours. After the starting material was consumed, the excess sodium borohydride was quenched with 110mL of 2.4N hydrochloric acid at 0-5°C. The aqueous layer was at pH=1. The two phase mixture was stirred at 0-5°C for 30 min. The layers were settled and separated. The upper aqueous layer was separated from the lower organic layer. The organic layer was washed with 15mL of 2.4N hydrochloric acid. The

15 aqueous layers were combined and the pH was adjusted to 9.5 with the addition of 30 mL of 25% sodium hydroxide solution at < 10°C. There was no precipitate formed in the beginning at 10°C. The ice/water bath was then removed and the batch was stirred at ambient temperature. Solids precipitated at the overnight agitation. The batch was cooled to 0-5°C and stirred for 1.5h. Product was filtered, rinsed with cold water (25 mL x 2), and

20 dried. 8.99g (40.2%) of off-white solid was obtained as Lot 2377-085 (4 ppm 14-hydroxycodeinone).

Example 6 - Rearrangement of 14-Hydroxycodeine to Oxycodone Base

- 15 -


An oven-dried 100 mL round bottom flask was charged with 5 g 14-hydroxycodone (15.9 mmol, Lot 2377-085) and 40mL of deoxygenated methanol. The mixture was stirred under nitrogen until homogenous solution formed.

An oven-dried 25mL round bottom flask was charged with 10mL of deoxygenated methanol and placed under nitrogen. 75mg of bis(norbornadiene)rhodium (I) tetrafluoroborate (0.2 mmol) and 86mg of 1,4-bis(diphenylphosphino)butane (0.2 mmol) were added under nitrogen and the orange solution was stirred at RT for 10 minutes. The solution was then sparged with hydrogen gas for 45 minutes. The color of the solution changed from orange to reddish orange. Excess hydrogen was removed by sparging the solution with nitrogen for 10 minutes. The solution of the catalyst was transferred to the solution of 14-hydroxycodone via cannula. The combined solution was heated at 50 °C for 9 hours. The product, Oxycodone free base, precipitated from the reaction mixture and a thick slurry formed. A small sample of the supernatant was taken for the in process test to confirm the reaction completion.

The reaction mixture was cooled to 0-5 °C for 2 hours and the product was filtered off, and the filter cake washed with cold ethanol (10 mL x 2).

The product was dried in a vacuum oven at 55 °C. 1.71g (34.2% yield, as Lot 2377-095) of oxycodone base was obtained. (45 ppm 14-hydroxycodone, undetected 8 α ,14-dihydroxy-7,8-dihydrocodeinone by LC/MS and 5 ppm 8 β ,14-dihydroxy-7,8-dihydrocodeinone by LC/MS; 99.41A% pure by HPLC).

Example 7 – Purification of Oxycodone Base by Sodium Metabisulfite



To a 25mL flask is charged 1.5g (4.756mmol, Lot 2377-095, prepared via Rh catalyst rearrangement) of oxycodone base and 7.5mL of water. The mixture is stirred at room temperature (RT, 20 $\pm$ 5°C). pH is 8.45, which is then adjusted to 7.02 at RT by adding 0.056g of conc. HCl.


- 16 -

Preparation of pH 7 $\text{Na}_2\text{S}_2\text{O}_5/\text{Na}_2\text{SO}_3$ buffered solution: To a small flask is charged 0.362g of sodium *meta*-bisulfite (1.904mmol, 0.4eq) and 1mL of water. Stirred at RT and a clear solution resulted within 4 min (pH~4-5). To another small flask, 1.43g (11.35mmol) of sodium sulfite and 6mL water are charged. The mixture is stirred at RT and a clear solution resulted within 4 min (pH~9-10). The sodium *meta*-bisulfite solution is transferred to sodium sulfite solution at RT and stirred. The pH is 6.90, which is then adjusted to 6.97 at RT by adding 1.43g of saturated sodium sulfite solution.

The pH 7 buffered solution of sodium *meta*-bisulfite/sodium sulfite prepared above is transferred to the batch at RT. The pH went up to 7.98, which is then adjusted to 7.01 at RT by adding 0.777g of conc. HCl. A solution results. The mixture is stirred at RT for 3-24h, or until the reaction is complete. (In-process test for reaction completion by LC/MS: $\leq$ 1ppm 14-hydroxycodeinone.) pH is adjusted to 9.18 by adding 0.95g (1.1mL) of conc. ammonium hydroxide. The mixture (a white slurry) is stirred at RT for 2h, filtered, washed with water (6mL x2) and dried until transferable.

The wet cake is transferred to a 25mL flask, slurried in 12mL water for 2h, filtered, washed with water (6mL x2), and dried until weight is constant. 1.34g (89.3% yield) of low ABUG oxycodone base is obtained, as Lot 2377-103 (~1 ppm 14-hydroxycodeinone by LC/MS, undetected $8\alpha,14$ -dihydroxy-7,8-dihydrocodeinone by LC/MS and 4 ppm $8\beta,14$ -dihydroxy-7,8-dihydrocodeinone by LC/MS; 99.48A% pure by HPLC).

Example 8A – Purification of Oxycodone Base by Hydrogenation



To a 250mL hydrogenation bottle was charged 4.05g (12.84mmol, prepared via the Rh-catalyzed rearrangement, 267 ppm 14-hydroxycodeinone; 16 ppm $8\alpha,14$ -dihydroxy-7,8-dihydrocodeinone, 111 ppm of $8\beta,14$ -dihydroxy-7,8-dihydrocodeinone) of oxycodone base, 36mL of dichloromethane (DCM) and 4mL of methanol. The mixture was agitate until

- 17 -

a clear solution was formed, and 0.4g of 5% Pd/C was added. The mixture was hydrogenated with agitation at 21-22°C (room temperature), 13-17 psi H₂ and was processed in portions to isolated oxycodone base after one and two days of hydrogenation.

After one day under hydrogenation conditions, about 16mL of the mixture was removed from hydrogenation bottle, catalyst was filtered was removed by filtration, and the filtered solution was concentrated by distilling off DCM/methanol. After most of the DCM/methanol had been removed, isopropanol (IPA) was added and the mixture was distilled at 65-70°C for 10min. The mixture was cooled to room temperature and then to 0-5°C. The product was isolated by filtration, rinsed and dried. 1.359g of low ABUG oxycodone base was obtained. (2.4 ppm 14-hydroxycodone by LC/MS; undetected 8 α ,14-dihydroxy-7,8-dihydrocodeinone by LC/MS, 28 ppm 8 β ,14-dihydroxy-7,8-dihydrocodeinone by LC/MS; 97.61A% pure by HPLC).

After the second day under hydrogenation conditions, the rest of the mixture was filtered to remove catalyst was and the DCM/methanol was removed by distillation. After most DCM/methanol had been removed, IPA and continued the mixture was distilled at 75-76°C for 10min. The mixture was cooled to room temperature and then to 0-5°C. The product was isolated by filtration, rinsed and dried. 2.442g of low ABUG oxycodone base was obtained (2.7 ppm 14-hydroxycodone by LC/MS, 3 ppm 8 α ,14-dihydroxy-7,8-dihydrocodeinone by LC/MS, 41 ppm 8 β ,14-dihydroxy-7,8-dihydrocodeinone by LC/MS; 97.67A% pure by HPLC).

Example 8B – Purification of Crude Oxycodone Base by Hydrogenation at 35°C


To a 250mL hydrogenation bottle was charged 5.0g (15.85 mmol, prepared via the Rh-catalyzed rearrangement , 267 ppm 14-hydroxycodone; 16 ppm 8 α , 11 ppm of 8 beta) of oxycodone base, 30mL of dichloromethane (DCM) and 10mL of isopropanol (IPA). The mixture was agitated until a clear solution was formed, and 0.5g of 5% Pd/C was added. The mixture was hydrogenated with agitation at 35°C , ca. 25 psi H₂ for one day. The catalyst was removed by filtration, and the filtered solution was concentrated by distilling off DCM/IPA. After most of the DCM/IPA had been removed, 3 mL of isopropanol (IPA) was added and the mixture was distilled at 65-70°C for 5 min. The mixture was cooled to room temperature and then to 0-5°C. The product was isolated by filtration, rinsed with cold IPA, and dried. The low ABUG oxycodone base obtained showed 1.5 ppm 14-hydroxycodone by LC/MS; undetected 8 α ,14-dihydroxy-7,8-

Example 11 - Conversion of Purified, Low ABUG Oxycodone Base to Oxycodone Hydrochloride Salt

- 19 -

To a 25mL flask was charged 2.0g (6.34mmol, from Example 9) of low ABUK oxycodone base, 6.4mL of ethanol and 1.3mL of water. The mixture was stirred at RT. The batch was heated to 41-46°C, added 0.643g (6.527mmol, 1.03 molar equivalents) of 37% hydrochloric acid, and a solution was then resulted. The mixture was heated to 55°C and stirred at 55°C for 10 hours to simulate the production salt formation timeframe. The mixture was cooled to room temperature and then to 0-5°C. The mixture was stirred at 0-5°C for 1.5 hours, filtered, and the product on the filter was rinsed with cold ethanol (2.5mL x 2) and dried. 1.899g of oxycodone HCl (85.12% yield, as Lot 2418-123) was obtained (2.5 ppm 14-hydroxycodeinone by LC/MS, undetected 8 α ,14-dihydroxy-7,8-dihydrocodeinone by LC/MS and 8 β ,14-dihydroxy-7,8-dihydrocodeinone by LC/MS; 0.01 HPLC A% 6-oxycodol; 99.93 HPLC A% oxycodone HCl).

Example 12 – Telescoped Conversion Of Crude Oxycodone Base to Low ABUK Oxycodone HCl Salt



To a 250mL hydrogenation bottle was charged 1.5g (4.75mmol, prepared via Rh catalyst rearrangement, 424 ppm 14-hydroxycodeinone) of oxycodone base, 6mL of dichloromethane (DCM) and 2mL of isopropanol (IPA). The mixture was agitated until a solution was obtained, then 0.15g of 10% Pd/C was added. The mixture was hydrogenated with agitation at 35°C, 25 psi H₂ and processed to isolated oxycodone HCl in portions after one and two days of hydrogenation.

After one day of hydrogenation, about 1.5mL of the mixture was removed from hydrogenation bottle and filtered to remove catalyst. The filtered mixture was distilled to minimize the solvents (DCM/IPA). The batch was diluted with 1.2mL of IPA, 0.4mL of water and 0.105g of 37% hydrochloric acid (1.066 mmol), and stirred at ambient temperature. The mixture was then heated to 55-60°C and stirred at 55-60°C for 10 min. The batch was cooled to room temperature and then to 0-5°C. The mixture was stirred at 0-5°C for 1.5 hours, filtered, and the solid product was rinsed with cold IPA (1mL) and dried.

- 20 -

0.369g of oxycodone HCl (as Lot 2418-047) was obtained (3 ppm 14-hydroxycodeinone by LC/MS; 99.08A% pure by HPLC).

After two days of hydrogenation, the rest of the mixture was filtered to remove catalyst. The filtered mixture was distilled to minimize the solvents (DCM/IPA). The mixture was diluted with 1.5mL of IPA, 0.5mL of water and 0.13g of 37% hydrochloric acid (1.32mmol), and stirred at ambient temperature. The mixture was then heated to 55-60°C and stirred at 55-60°C for 10 min. The mixture was cooled to room temperature and then to 0-5°C. The mixture was stirred at 0-5°C for 1.5 hours, filtered, and the solid product was rinsed with cold IPA (1.5mL x 2) and dried. 0.461g of oxycodone HCl (as Lot 2418-051) was obtained (2 ppm 14-hydroxycodeinone by LC/MS; 98.66A% pure by HPLC).

Example 13 - Hot Acid Stress Test of Oxycodone Hydrochloride from Example 11

To a 10mL flask are charged 0.232 g (0.659) of low ABUK oxycodone HCl (T-zero: 2.5 ppm 14-hydroxycodeinone) and 5mL of water. The mixture is stirred at room temperature to form a solution. 0.26g of 3N aq. HCl (0.659 mmole, 1 molar equivalent) is then added. The batch is heated to 65°C and monitored over 24 hours with LC/MS analysis. LC/MS analysis showed 4.2 ppm 14-hydroxycodeinone after 5h and 6.1ppm after 24h. The slow growth over 24 was ascribed to degradation of unknown species other than 8 α ,14-dihydroxy-7,8-dihydrocodeinone.

Entry	Hours in Hot Acid	ABUK (ppm)
1	0	2.5
2	2	3.6
3	5	4.2
4	24	6.1

Example 14 - Dry Heat Stress Test of Oxycodone Hydrochloride from Example 11

A 0.0225 g (0.597 sample of low ABUK oxycodone HCl (T-zero: 3.9 ppm 14-hydroxycodeinone) was placed in a standard lab drying oven under air at 62°C and the level of 14-hydroxycodeinone was monitored by LC/MS for 14 days. No growth in the level of 14-hydroxycodeinone was observed.

Entry	Days (at 62°C)	ABUK (ppm)
1	T-zero	3.9 ppm
2	4	3.7 ppm

- 21 -

3	7	4.0 ppm
4	14	3.7 ppm

Example 15: Hot Acid Stress Test of Oxycodone Base from Example 9

To a 10mL flask are charged 0.20 g (0.634) of low ABUK oxycodone base (T-zero: 2.5 ppm 14-hydroxycodeinone by LC/MS) and 5mL of water. The mixture is stirred at room temperature to form a solution. 0.26g of 3N aq. HCl (0.659 mmole) is then added. The batch is heated to 65°C and monitored over 24 hours with LC/MS analysis. LC/MS analysis showed 3.3 ppm 14-hydroxycodeinone after 5h and 6.4 ppm after 24h. The slow growth after 24h was ascribed to degradation of unknown species other than 8 α ,14-dihydroxy-7,8-dihydrocodeinone.

10

Entry	Hours in Hot Acid	ABUK (ppm)
1	0 (r.t.)	2.5
2	2	2.5
3	5	3.3
4	24	6.4

Example 16 - Dry Heat Stress Test of Oxycodone Base from Example 9

A 0.0225 g (0.597 sample of low ABUK oxycodone (T zero: 3.3 ppm 14-hydroxycodeinone by LC/MS) was placed in a standard lab drying oven under air at 62°C and the level of 14-hydroxycodeinone was monitored by LC/MS for 14 days. No growth in the level of 14-hydroxycodeinone was observed

Entry	Days (at 62°C)	ABUK (ppm)
1	T-zero	3.3 ppm
2	4	3.4 ppm
3	7	3.3 ppm
4	14	3.6 ppm

20 Analytical Method for PPM Level 14-Hydroxycodeinone and for 8 α ,14-dihydroxy-7,8-dihydrocodeinone and 8 β ,14-dihydroxy-7,8-dihydrocodeinone (HPLC/MS-SIM)

1.1 Reagents and Materials: (Equivalent reagents and materials may be substituted)

Methanol (MeOH)	Fisher Scientific, HPLC Grade
Acetonitrile (ACN)	Fisher Scientific, HPLC Grade
Ammonium Acetate (NH ₄ OAc)	J.T. Baker, HPLC Grade

- 22 -

Purified Water (H ₂ O)	MilliQ, Model A10 Gradient Water System
14-Hydroxy Codeinone	JM Qualified Reference Standard
7,8-Dihydro-8 β -14-Dihydroxycodeinone (Inert β -Diol)	Retention Time Marker (not qualified)

1.2 Instrumentation: (Equivalent instrumentation can be used)

HPLC	Waters Acquity UPLC System
UV Detector	Waters Acquity PDA Detector
Mass Spec	Waters Q-ToF Premier
Data System	Waters MassLynx 4.1
Balance	Mettler-Toledo, Model AT261 or PG503-S, Delta Range

5 1.3 Mobile Phase Preparation: (For 1 L each)

Mobile Phase A: Weigh ~ 0.77 (± 0.03) g of Ammonium Acetate into a suitable mobile phase bottle, dissolve with 950 mL of deionized water, add 25 mL of Acetonitrile and 25 mL of MeOH to the container. Mix well and degas.

10 *Mobile Phase B:* Weigh ~ 0.77 (± 0.03) g of Ammonium Acetate into a suitable mobile phase bottle, dissolve with 100 mL of deionized water, add 450 mL of Acetonitrile and 450 mL of MeOH to the container. Mix well and degas.

The diluent: MeOH (free base) or 0.2% (v/v) TFA/H₂O (HCl salt).

1.4 Operating Conditions:

Column	Waters, XBridge, Phenyl, 5 μ m , 4.6 x 150 mm
Col. Temperature	Ambient
Sample Temp	Ambient
Injection Volume	10 μ L
Detection	UV at 220 nm
Flow Rate	0.8 mL/min, with a splitter to lead ~0.2 mL/min to the mass spec.
Analysis Time	30 min
Run Time	35 min

Linear Gradient (Mixing) Conditions:	Time (min)	% MP A	% MP B	Curve
	initial	100	0	6
	22	73	27	6
	25	10	90	6
	30	10	90	6
	31	100	0	6
	40	100	0	6

15

Mass Spec Parameters:		
Source	Instrument	Acquisition

- 23 -

Capillary	0.5 (diols) 1.5 (ABUK)	LM Resol	4.7	Source	ESI
Spl Cone	30	HM Resol	15.0		
Extra Cone	5.0	Ion Energy	0.0	Source	ESI
Ion Guide	3.5	Pre-filter	5.0	Polarity	+
Source Temp	120	Collision Energy	5.0	Analyzer Mode	V
Cone Gas	0	Cell Entrance	2.0	Sensitivity	Maximum at 332 a.u. (Diols) Maximum at 314 a.u. (ABUK)
De-solvation Temp	450	Cell Exit	-10.0	Scan	0.3
				Scan Delay	0.02
Detector Voltage	1920	Collision Cell	0.4	Data Format	Continuum
De-solvation	700	Ion Guide	0.0	Mass Range	332.0 – 332.3 a.u. (Diols) 313.9 - 314.3 a.u. (ABUK)

1.5 Approximate Retention Times of Known Analytes:

Analyte	Approximate Retention Time* (min)	RRT
α -Diol	15.8	0.82
β -Diol	17.7	0.91
Oxycodone	19.5**	1.00
14-Hydroxy Codeinone	23.0	1.18

* The retention time is extremely sensitive to the mobile phase.

** The peak is saturated.

5

1.6 Impurity Working Standard Solution Preparation

- Weigh 15 mg (± 20 %, accurate to the second digit passed the decimal point) each of 14-Hydroxy Codeinone reference standard and β -Diol into a 100 mL volumetric flask. Dissolve and then dilute to the volume with MeOH. Sonicate for 30 sec and mix well. This is the impurity stock solution.
- Transfer 2.0 mL of the impurity stock solution into a 100 mL volumetric flask, dilute to volume with 0.2% TFA/H₂O, and mix well. This is the impurity stock solution-2.

10

- 24 -

- Transfer 5.0 mL of the impurity stock solution-2 into a 100 mL volumetric flask, dilute to volume with 0.2% TFA/H₂O, and mix well. This is the impurity working standard solution.

1.7 Resolution Solution Preparation:

5 Accurately weigh 150 mg ($\pm$ 10 mg) of the Oxycodone material, which has the lowest possible level of β -Diol, into a 10 mL volumetric flask. Dissolve and dilute to volume with the impurity stock solution-2 (Sonication may be necessary).

1.8 Sample Solution Preparation:

10 In duplicate, accurately weigh 150 mg ($\pm$ 10 mg) of the sample into a 10 mL volumetric flask. Dissolve the sample and dilute to volume with the diluent (Sonication may be necessary).

1.9 System Equilibration and Conditioning:

15 Pump Mobile Phase A through the column for at least 20 minutes followed by pump Mobile Phase B for at least another 20 minutes at a flow rate of 0.8 mL/min. Switch to Initial assay conditions and pump for at least 20 minutes.

1.10 Procedure:

- Separately inject the diluent as a blank.
- Inject the resolution solution once.
- Inject the impurity working standard three times.
- 20 • Inject each sample solution under the full gradient.
- Inject the diluent at the end.
- Quantify the level of 14-Hydroxy Codeinone in the sample by comparing to the averaged corresponding peak specific response (SR) of the standard solution.
- Report the level of 14-Hydroxy Codeinone found in the sample to the nearest 0.0001%.
- 25 • Report the peak area of the diols for information only.

1.11 System Suitability:

The RRT of the inert diol (MS signal) relevant to the saturated Oxycodone (UV signal) is NMT 0.95.

1.12 Calculations:

30 1.12.1 ABUK %w/w (in free base form):

$$\%w/w = \frac{(\text{ABUK in Sample}^{\text{Avg PA}})(100)(\text{ABUK Std}^{\text{Conc. mg/mL}})(\text{ABUK Std Purity (in decimal)})}{(\text{ABUK Std}^{\text{Avg PA}})(\text{Sample}^{\text{Conc. mg/mL}}) \times \text{CF}}$$

Where: PA = Peak Area Std = Standard

35

Conversion Factor (CF) = MW of the base form


MW of HCl Salt form

$$\text{PPM} = \%w/w \times 1000000$$

What is claimed:

1. A method of preparing oxycodone, comprising contacting 14-hydroxycodeine with a catalyst under conditions sufficient to cause rearrangement of the 14-hydroxycodeine to form oxycodone free base.
2. The method of claim 1, wherein the 14-hydroxycodeine is formed by reduction of 14-hydroxycodeinone.
3. The method of claim 2, wherein the 14-hydroxycodeinone is derived from thebaine.
4. The method of claim 1, further comprising purifying the oxycodone free base.
5. The method of claim 1, further comprising converting the oxycodone free base to an oxycodone salt.
6. The method of claim 3, wherein the 14-hydroxycodeinone contains at least one isomer of 8,14-dihydroxy-7,8-dihydrocodeinone as an impurity and wherein, during the reduction of 14-hydroxycodeinone to form 14-hydroxycodeine, the 8,14-dihydroxy-7,8-dihydrocodeinone is reduced to at least one isomer of 8,14-dihydroxy-7,8-dihydrocodein-6-ol.
7. The method of claim 6, wherein at least one isomer of 8,14-dihydroxy-7,8-dihydrocodeinone is formed during an oxidation of the thebaine.
8. The method of claim 6, wherein the at least one isomer of 8,14-dihydroxy-7,8-dihydrocodeinone is an alpha-isomer, a beta-isomer or both the alpha-isomer and the beta-isomer at the C-8 hydroxyl group.
9. The method of claim 2, wherein the reduction comprises use of a hydride reagent.
10. The method of claim 9, wherein the hydride reagent is selected from L-selectride, N-selectride and sodium borohydride.
11. The method of claim 2, wherein the reduction is performed in a solvent or a mixture of solvents.

12. The method of claim 11, where in the mixture of solvents comprises at least one alcohol and at least one aprotic organic solvent.
13. The method of claim 12, wherein the alcohol is selected from the group consisting of methanol, ethanol, isopropanol, n-propanol, isobutanol, n-butanol and 2-butanol.
14. The method of claim 12, wherein the aprotic organic solvent is selected from the group consisting of methylene chloride, 1,2-dichloroethane, ethyl acetate, isopropyl acetate, tetrahydrofuran, toluene, xylene and methyl tert-butyl ether.
15. The method of claim 2, wherein less than 100 ppm of 14-hydroxycodeinone is present after the reduction.
16. The method of claim 2, wherein less than 50 ppm of 14-hydroxycodeinone is present after the reduction.
17. The method of claim 8, wherein less than 2 ppm of 8 α ,14-dihydroxy-7,8-dihydrocodeinone is present after the 14-hydroxycodeinone is reduced.
18. The method of claim 1, wherein the catalyst is a ligand-complexed metal catalyst.
19. The method of claim 18, wherein the ligand is a phosphine.
20. The method of claim 18, wherein the metal is rhodium or ruthenium.
21. The method of claim 20, wherein the catalyst is



22. The method of claim 1, wherein the contacting is performed in a solvent or a mixture of solvents.
23. The method of claim 22, wherein the contacting is performed in said mixture of solvents and wherein the mixture of solvents comprises at least one alcohol and at least one aprotic organic solvent.
24. The method of claim 23, wherein the alcohol is selected from the group consisting of methanol, ethanol, isopropanol, n-propanol, isobutanol, n-butanol and 2-butanol.
25. The method of claim 23, wherein the aprotic organic solvent is selected from the group consisting of methylene chloride, 1,2-dichloroethane, ethyl acetate, isopropyl acetate, tetrahydrofuran, toluene, xylene and methyl tert-butyl ether.
26. The method of claim 1, wherein the contacting is performed at a temperature in a range from about 25°C to about 100° C.
27. The method of claim 4, wherein the purifying comprises contacting the oxycodone free base with a solution comprising sodium meta-bisulfate.
28. The method of claim 27, wherein the solution is a $\text{Na}_2\text{S}_2\text{O}_5/\text{Na}_2\text{SO}_3$ buffered solution.
29. The method of claim 5, wherein the oxycodone salt is oxycodone hydrochloride.
30. An oxycodone free base having less than 25 ppm 14-hydroxycodeinone and less than 2 ppm of 8 α ,14-dihydroxy-7,8-dihydrocodeinone, wherein the oxycodone free base is made by the method of claim 4.
31. The oxycodone free base of claim 30, wherein less than 10 ppm of 14-hydroxycodeinone is present.
32. The oxycodone free base of claim 30, wherein less than 5 ppm of 14-hydroxycodeinone is present.

33. An oxycodone free base having less than 25 ppm 14-hydroxycodeinone and less than 2 ppm of 8 α , 14-dihydroxy-7,8-dihydrocodeinone, wherein the oxycodone free base is made by the method of claim 27.
34. The method of claim 4, wherein the purifying comprises contacting the oxycodone free base with a mixture comprising an alcohol, a palladium on carbon catalyst, and hydrogen, and optionally a chlorinated solvent.
35. The method of claim 34, wherein the alcohol comprises at least one of methanol, isopropanol, ethanol, and butanol.
36. The method of claim 34, wherein the chlorinated solvent is dichloromethane, dichloroethane, chloroform, or chlorobenzene.
37. The method of claim 34, wherein the palladium on carbon catalyst is 5% or 10% by weight of palladium on carbon.
38. The method of claim 34, wherein the hydrogen is pressurized from about 15 psi to about 50 psi.
39. The method of claim 34, wherein the contacting is performed at a temperature from about 18°C to about 35°C.
40. The method of claim 34, wherein the oxycodone free base is isolated as a solid.
41. The method of claim 40, wherein the solid contains less than 10 ppm 8 α ,14-dihydroxy-7,8-dihydrocodeinone.
42. The method of claim 41, wherein less than 2 ppm of 8 α ,14-dihydroxy-7,8-dihydrocodeinone is present.
43. The method of claim 40, wherein the solid contains less than 10 ppm 14-hydroxycodeinone.
44. The method of claim 40, wherein the solid contains less than 2 ppm 14-hydroxycodeinone.

45. The method of claim 34, wherein the oxycodone free base is converted to oxycodone hydrochloride salt and the oxycodone hydrochloride salt is isolated as a solid.
46. The method of claim 45, wherein the oxycodone free base is not isolated as a solid prior to said converting.
47. The method of claim 45, wherein the oxycodone hydrochloride salt contains less than 5 ppm 14-hydroxycodeinone.
48. The method of claim 40, wherein the solid oxycodone free base is contacted with ethanol.
49. The method of claim 48, wherein the contacting is performed at an elevated temperature.
50. The method of claim 48, wherein the oxycodone free base is re-isolated after the contacting with ethanol.
51. The method of claim 50, wherein the re-isolated oxycodone free base contains less than 5 ppm 8.alpha.,14-dihydroxy-7,8-dihydrocodeinone.
52. The method of claim 51, wherein less than 2 ppm of 8.alpha.,14-dihydroxy-7,8-dihydrocodeinone is present.
53. The method of claim 50, wherein the re-isolated oxycodone free base contains less than 5 ppm 14-hydroxycodeinone.
54. An oxycodone free base comprising less than 2 ppm of 8.alpha.,14-dihydroxy-7,8-dihydrocodeinone and less than 5 ppm 14-hydroxycodeinone.
55. An oxycodone hydrochloride having less than 25 ppm 14-hydroxycodeinone and less than 2 ppm of 8.alpha.,14-dihydroxy-7,8-dihydrocodeinone.
56. The oxycodone hydrochloride of claim 55 having less than 10 ppm 14-hydroxycodeinone.

57. The oxycodone hydrochloride of claim 55 having less than 5 ppm 14-hydroxycodeinone.
58. The oxycodone hydrochloride of claim 55, wherein less than 2 ppm of 14-hydroxycodeinone is present.
59. An oxycodone free base made by the method of claim 34 comprising less than 2 ppm of 8 α ,14-dihydroxy-7,8-dihydrocodeinone and less than 5 ppm 14-hydroxycodeinone.
60. An oxycodone free base made by the method of claim 48 comprising less than 2 ppm of 8 α ,14-dihydroxy-7,8-dihydrocodeinone and less than 5 ppm 14-hydroxycodeinone.
61. An oxycodone hydrochloride made by the method of claim 5 having less than 25 ppm 14-hydroxycodeinone and less than 2 ppm of 8 α ,14-dihydroxy-7,8-dihydrocodeinone.
62. The oxycodone hydrochloride of claim 61 having less than 10 ppm 14-hydroxycodeinone.
63. The oxycodone hydrochloride of claim 61 having less than 5 ppm 14-hydroxycodeinone.
64. The oxycodone hydrochloride of claim 61, wherein less than 2 ppm of 14-hydroxycodeinone is present.

1/5





FIG. 1

2/5



SUBSTITUTE SHEET (RULE 26)

3/5




FIG. 3

4/5




FIG. 4

SUBSTITUTE SHEET (RULE 26)

5/5









FIG. 5



Low ABUK Base


Undetected (<2 ppm) α-diol
<0.1% oxycodone
< 5 ppm 14-hydroxycodeinone



Low ABUK Oxycodone HCl

< 5 ppm 14-hydroxycodeinone

Base Purification by hydrogenation
with direct formation of salt



Low ABUK Oxycodone HCl

< 5 ppm 14-hydroxycodeinone

Base Purification by hydrogenation
with ethanol slurry option.