

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199662305 B2
(10) Patent No. 720892

(54) Title
Animal feed

(51)⁷ International Patent Classification(s)
A23K 001/14 A23K 001/18
A23K 001/16 A61K 035/78

(21) Application No: **199662305**

(22) Application Date: **1996.06.14**

(87) WIPO No: **WO97/00018**

(30) Priority Data

(31) Number	(32) Date	(33) Country
9512106	1995.06.14	GB

(43) Publication Date : **1997.01.15**

(43) Publication Journal Date : **1997.03.06**

(44) Accepted Journal Date : **2000.06.15**

(71) Applicant(s)
Matforsk Norwegian Food Research Institute

(72) Inventor(s)
Erik Slinde; Stefan Sahlstrom; Leif Arne Fardal; Askild Holck; Grete Skrede

(74) Agent/Attorney
CALLINAN LAWRIE, Private Bag 7, KEW VIC 3101

(56) Related Art
EP 162805
FR 2695619

OPI DATE 15/01/97 APPLN. ID 62305/96
AOPP DATE 06/03/97 PCT NUMBER PCT/GB96/01414



AU9662305

CT)

(51) International Patent Classification ⁶ : A23K 1/14, 1/16, 1/18, A61K 35/78		A1	(11) International Publication Number: WO 97/00018
			(43) International Publication Date: 3 January 1997 (03.01.97)
(21) International Application Number: PCT/GB96/01414		(74) Agents: COCKBAIN, Julian, Roderick, Michaelson et al.; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB).	
(22) International Filing Date: 14 June 1996 (14.06.96)			
(30) Priority Data: 9512106.7 14 June 1995 (14.06.95) GB		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): MATFORSK NORWEGIAN FOOD RESEARCH INSTITUTE [NO/NO]; Osloveien 1, N-1430 Ås (NO).			
(71) Applicant (for GB only): COCKBAIN, Julian, Roderick, Michaelson [GB/GB]; 27 Ladbroke Road, London W11 3PD (GB).			
(72) Inventors; and (75) Inventors/Applicants (for US only): SLINDE, Erik [NO/NO]; Natlandsfjellet 44, N-5030 Landås (NO). SAHLSTRØM, Stefan [NO/NO]; Rydningen 12, N-1430 Ås (NO). FARDAL, Leif, Arne [NO/NO]; Lindtrupbakken 12D, N-1440 Drøbak (NO). HOLCK, Askild [NO/NO]; Askeveien 34, N-1430 Ås (NO). SKREDE, Grete [NO/NO]; Arneborgveien 10, N-1430 Ås (NO).		Published <i>With international search report.</i>	
(54) Title: ANIMAL FEED			
(57) Abstract The present invention relates to animal feed with improved quality, improved resistance to undesirable bacterial strains and improved digestibility and to a process for preparing animal feed containing cereal involving cereal fermentation to produce a feed.			

ANIMAL FEED

The present invention relates to animal feed and to a process for preparing animal feed which yields improved feed quality and digestibility.

Carbohydrates and starch are used as feedstuffs in the farming industry and in this regard cereals, such as for example oats, used as a component of fish feed in the fish farming industry are a well-known carbohydrate source for farmed salmonids. Studies have shown that salmonids are capable of digesting starch from cereals and that cereals can be mixed in fish feed without adverse effects on digestibility or on the performance of the fish. However, in general, it is observed that the higher the level of cereal present in the feed, the lower the level of digestibility. As a typical example, an increase from 10% to 30% by weight of oats in fish feed typically reduces starch digestibility from about 61% to about 33%. Thus there is generally a need to increase starch digestibility in fish feed.

However, to date only small quantities of cereals have been used as a component of fish feed because of difficulty in extruding feed mixtures into pellets. Conventional feed pellets are also prone to contamination by bacterial strains such as Listeria monocytogenes. This poses a significant problem in the fish farming industry.

We have now found that pre-fermenting of cereal facilitates formation of firm feed pellets and produces feed which is more digestible than conventional feed containing non-fermented cereal, and moreover that where pre-fermentation is effected using a bacteriocin producing strain of lactic acid bacteria (lactobacilli), the growth of undesirable bacterial strains such as food-borne pathogens or food spoilage bacteria, (for

example species of Listeria, eg. Listeria monocytogenes) during the production and storage of the feed is suppressed. Furthermore, fish feed containing prefermented cereal has also been found to exhibit an immunostimulating effect.

Bacteriocins are peptides or proteins released by bacteria which show bactericidal or bacteriostatic activity towards both the producing strain and/or other bacteria. Generally, only bacteria closely related to the producing strain are inhibited although in some cases other, non related bacteria, including certain pathogens may also be affected.

Due to their potential use as antibacterial agents, bacteriocins have been the subject of intensive research. In recent years there has in particular been considerable interest in bacteriocins isolated from lactic acid bacteria (LAB), in view of their potential utility in the food and brewing industries, in particular in the preservation of food products.

It is known that short chain (1-3) β -glucans have an immunostimulating effect in salmon. For example, salmon treated with short chain (1-3) β -glucans have been shown to have increased resistance to different types of bacteria including Yersina ruckeri (Enteric redmouth disease), Vibro anguillarum (Classical vibriose) and Vibro salmonicidia (Hitra disease) (see Robertsen B, Rørstad G, Engstad R. and Raa J, Journal of fish diseases 13, 1990, 391-400).

Thus viewed from one aspect the present invention provides an animal feed, comprising a cereal fermented with lactic acid bacteria, wherein said cereal is selected from the group consisting of wheat, oats, rye and barley, together with at least one animal foodstuff component, said feed has been subjected to expansion, compression, extrusion and/or drying.

Preferably the animal feed is a fish (e.g. salmon) feed or bird (e.g. chicken) feed.



Viewed from a further aspect the present invention provides a process for producing an animal feed, said process comprising the steps of:

(a) fermenting a cereal with lactic acid bacteria, said cereal being selected from wheat, barley, rye and oats;

(b) admixing the fermented cereal with at least one animal foodstuff; and, optionally

(c) processing said admixture by expansion, compression, extrusion and/or drying.

Especially preferably a cereal is fermented with a bacteriocin-producing lactobacillus strain, particularly especially preferably a Sakacin A or Sakacin P producing lactobacillus strain.

Viewed from a still further aspect the invention provides the use of fermented cereal, preferably a bacteriocin containing fermented cereal, as a component in processed animal feed, preferably in fish or bird feed.

Viewed from a yet still further aspect the present invention provides a method of achieving an immunostimulatory response in an animal body by administering thereto an animal feed as defined hereinbefore and a method of reducing the level of cholesterol in an animal body by administering thereto an animal feed as defined hereinbefore together with a digestible amount of fibre capable of reducing the cholesterol level in said animal.

Viewed from an even further aspect the present invention provides the use of a fermented cereal in the manufacture of an animal feed for use in reducing the level of cholesterol in the animal to which the feed is fed or for use in a method of achieving an immunostimulatory response in an animal body.

Cereal fermentation has been found to have the advantage of changing the level of mono- and disaccharides present in the cereal and the solubility



- 4 -


of polysaccharides in the cereal, allowing the cereal grains to swell and partly dissolve. The admixture may then more easily undergo extrusion and thereby become more digestible for the fish.

Moreover, fermentation yields a feed with a lower pH than
5 identical feed containing unfermented cereal. This is particularly significant in view of the fact that the main pigments used in salmonid feed, astaxanthins, are known to have increased stability under acidic conditions. The pH stabilisation of pigments (carotenoids) in pelletised fish feeds is, therefore, an important
10 advantage of the invention, particularly in view of the high costs of pigments in feed.

Furthermore, cereal fermentation has been found to improve firmness and to lower water solubility of the feed overall, both of which being highly advantageous properties when feeding fish kept in
15 sea cages.

A further advantage of the process according to the invention is that it produces animal feed which may exhibit an increased immunostimulating effect in the animal to which it is fed. In this regard, it has been found that fish feed containing fermented barley
20 has an immunostimulating effect in the fish to which it is fed. Without wishing to be bound by any theoretical considerations, it is thought that this effect may be due to the formation of short chain (1-3) β -glucan during cereal fermentation by degradation of polysaccharides such as β -glucan.


Another advantage of animal feed prepared in accordance with the process according to the invention is that fermentation of the cereal provides a fish feed with improved digestibility, particularly with regard to starch digestibility. This in turn allows a greater fibre content to be added to the feed than would
30 otherwise be possible. This is important because fibre is an important dietary requirement for animals and is known to have a cholesterol lowering effect. Typically



- 4a -

up to 10% by weight of fibre will be digestible in fish feed according to the invention compared to an upper limit of 2.5% in conventional feed. Preferably the fibre content will be in the range 4-10% by weight.

5



- 5 -

The feed produced according to the invention may be processed to any conventional form, eg. by expansion, extrusion, heating, drying, or pelleting. By pellets, is meant any form of self-supporting feed unit, eg. tablets, cakes, bricks etc. Pelleting can be effected by compression but most conveniently will be by extrusion. While feeds in pellet form are preferred, the feeds according to the invention may be in other forms, eg. flakes or powders.

The cereal used in the present invention may be any feedstuff cereal. However for fish feed barley, wheat and oats are preferred and oats and barley are especially preferred. The digestibility in salmonids of cereal starch present in fish feed prepared in accordance with the process according to the invention is improved and it has been shown that for barley the increase in digestibility is significant (nearly 100%) and for wheat it is 20%.

The fermented cereal conveniently makes up from 1 to 95%, preferably 10 to 40% by weight of the animal feed. The remaining components may be conventional animal feedstuffs and feedstuff additives, such as unfermented cereals, fish meal, bacteriocins, fats, vitamins, minerals, water, colorants, flavour-enhancers, therapeutic agents, stabilizers, antioxidants and pH regulators. Cereal components may be whole grain, grain fragment, crushed grain or milled grain (eg. flour).

In the case of fish feed, the total cereal content of the feed is preferably no more than about 40% by weight and of this it is preferred that the major part be pre-fermented. Especially preferably substantially all the cereal component is pre-fermented.

The bacteriocins present as components of the fermented cereal may be produced during fermentation as mentioned hereinbefore but may also be produced separately and added pure or in a crude or partly purified form prior to processing.

The bacteriocins used may have one or several inhibition spectra. Preferably bacteriocin having

- 6 -

inhibitative effects against pathogen-producing bacteria (such as *Listeria*) found in feedstuffs will be used.

The cereal fermentation is preferably effected using a lactic acid bacteria and in particular a bacteriocin producing *Lactobacillus* strain. Many *Lactobacilli* are known and are commercially available, for example for use in bread baking, eg. from Chr Hanson, Denmark. Preferred fermentation agents include the lactic acid bacteria containing sour dough cultivars.

To inhibit bacterial strains such as *Listeria monocytogenes* in the animal feed, a bacteriocin producing *Lactobacillus* strain may be used. Several such strains are known, for example L. Sake strain Lb674 and Lb706 producing Sakacin P and Sakacin A respectively (L. Sake Lb674 is available from Dr. Lothar Kröckel, Federal Institute of Meat Research, Kulmach, Germany and L. Sake Lb706 from MATFORSK, The Norwegian Food Research Institute, Osloveien, Norway).

Lactic acid bacteria bacteriocins are usually small peptides, seldom containing more than 60 amino acids. Many bacteriocins from lactic acid bacteria have now been described (for a review see Nettles et al., 1993) and include, most notably, the well-studied nisin, (see for example Gross, et al., J. Am. Chem. Soc. 93: 4634-4635, 1971 and Hurst, Adv. Appl. Microbiol. 27: 85-123, 1981).

Preferably a bacteriocin producing strain should simultaneously carry out lactic acid fermentation and produce bacteriocin. As bacteriocins are peptides, they may be decomposed and inactivated by proteinases in the flour. In such cases it may be preferable to add bacteriocin prior to processing or to pre-heat the flour to inactivate any such proteinases. If the bacteriocin is unable to survive processing (eg. extrusion) then further bacteriocin or bacteriocin producing strains may be added after processing.

The fermentation step in the process according to the invention is conveniently effected in an aqueous

- 7 -

medium, generally water, containing 40-60% by weight cereal, preferably in the form of finely ground whole grain flour, for a period of 10 to 48 hours, preferably about 24-48 hours, until an acid pH is obtained, for example a pH of 1-6, especially 3-5, and at ambient or above ambient temperature, for example 25 to 35°C, especially about 30°C.

In the fermentation step, a starter culture may be used or alternatively one may use a spontaneous culture obtained from the cereal used, optionally one obtained in advance. Using a starter culture the fermentation period would typically be 10 to 16 hours, while with spontaneous fermentation a longer period of 24 to 48 hours might be used. The fermentation will not generally be effected using an initially acid medium. The acid pH develops as a result of acid fermentation during fermentation.

The following Examples are intended to illustrate the invention in a non-limiting manner:

EXAMPLE 1

The raw materials used to carry out the following experiments were:

- (a) Wheat - whole grain wheat flour;
- (b) Oats - finely ground whole grain oat flour;
- (c) Fish meal - Norseamink, Norwegian fish meal with antioxidant, Egersund Sildoljefabrikk, Norway.

Four mixtures were prepared by admixing fish meal with the following untreated or pre-treated cereals:

- (1) Untreated wheat (15%)
- (2) Untreated oats (15%)
- (3) Fermented oats (sour-dough) - (based on 15% flour weight)
- (4) Scalded oats (based on 15% flour weight)

Fish feed mixtures containing cereals (1) and (2) were prepared by manually mixing the untreated wheat or oat flour (15%) with fish meal. The water content for both mixtures was 8.1%.

For the pre-treatment of the fish feed mixture containing (3), fermentation was carried out by adding equal amounts of oats flour and water to an equivalent amount of lactic acid bacteria (sour dough) culture. The mixture (20°C, pH 4.6) was fermented for 24 hours at 30°C to produce a preculture. The preculture was refreshed regularly with oat flour and water prior to extrusion a week later. The pH of the final fermentation mixture was 4.0 and when admixed with fish meal yielded a mixture with a water content of 19.9%.

For the pre-treatment of the fish feed mixture containing (4), boiling water was added to oats flour in a 1:1 ratio. Shortly after addition, the temperature was 76°C and cooling to 30°C led to a weight loss of 3%. The scalded flour/water mixture was added to fish meal yielding a mixture with a water content of 19.9%.

Extrusion was carried out on a Werner & Pleider, Continua 37 extruder. The temperature of the extruder was in the range 130-150°C and water was added at a rate of between 2 and 7.5 l/hour. Further variants included

- 9 -

the rotating screw (280-320 rpm), the flour feeding rate (1.2-2.4 g/min) and the feeding temperature (130-145°C). During the extrusion process, the pressure varied between 10 and 25 bar.

The pellets were allowed to cool and dry at ambient temperature and then packed in closed drums. For each experiment at least two fractions were sampled and the results therefrom are compiled in Tables 1 and 2 below.

The % dry matter for each sample was determined shortly after production and again after 24 hours in air at ambient temperature. The pH was also measured. Breakage and crushing strength were measured in a Kramer cell in an Instron Universal testing machine. The breakage strength was measured perpendicular to the length of the pellet and averaged over 12 pellets. The width and length of each pellet were measured together with their water solubility and rate of sinking. As used herein, the term solubility refers to the % weight of pellets dissolved in water during a certain period of soaking in water.

Results of the tests carried out on the 4 types of fish feed mixture are presented in Tables 1 and 2 below.

Table 1

Dry matter (%) and pH of pellets from wheat and oats

Treatment	Dry matter (%) at production	after 24 h	pH
Untreated wheat	81.9	89.9	6.1
Untreated oats	77.1	87.3	6.0
Fermented oats	75.6	87.7	5.8
Scalded oats	76.4	86.6	6.0

Table 2
Breakage and water solubility of pellets
from wheat and oats

Treatment	Breakage N	Crushing N	Water solubility (%)	
			2 min	20 h
Untreated wheat	16.4	213	8	62
Untreated oats	16.6	103	15	38
Fermented oats	16.4	127	9	32
Scalded oats	16.3	134	19	100

Notably from the results in Table 1, drying for 24 hours at ambient temperature led to an 8 to 12.1% increase in the dry matter content with the most extensive drying observed in the fermented oats. All of the pellets proved to be stable toward mould growth or any other visible deterioration over a 12 month period. In addition, pellets containing fermented oats had a lower pH than the other pellet-types.

Whilst breakage strength showed no significant variation amongst the pellet-types, a greater force was required to crush the wheat-containing pellets than any of the oat-containing pellets.

Water solubility was determined from the time taken for pellets to sink in water. All pellet types took several hours to sink but when finally wetted, solubility rose.

EXAMPLE 2

Experiments in Atlantic salmon fed with feed containing different amounts of fermented and unfermented wheat or barley flour

The following describes the utility of animal feed produced by the process according to the invention. Feed composition is stated alongside digestibility



- 11 -

results, carbohydrate analysis and immunostimulating effects in salmon. In all experiments, the salmon performed well.

The raw materials used to carry out the following experiments were:

- (a) Wheat - whole grain wheat flour, finely ground;
- (b) Barley - whole grain barley flour, coarsely ground.

Eight mixtures were prepared by admixing fish meal with the following pre-treated (fermented) or untreated (unfermented) cereals:

- (1) Pretreated wheat (12%) (HF 12)
- (2) Pretreated wheat (24%) (HF 24)
- (3) Pretreated wheat (36%) (HF 36)
- (4) Untreated wheat (24%) (HU 24)
- (5) Pretreated barley (12%) (BF 12)
- (6) Pretreated barley (24%) (BF 24)
- (7) Pretreated barley (36%) (BF 36)
- (8) Untreated barley (24%) (BU 24)

The feed compositions were chosen to give equal levels of energy and are shown in Table 3. Extrusion conditions were as described above in Example 1.

- 12 -

Table 3
Feed compositions (kg/100 kg feed)

Type of feed ingredient	Fermented			Unfermented
	1	2	3	4
Coding	H(B) F 12	H(B) F 24	H(B) F 36	H(B) U24
Wholemeal wheat/barley	12	24	36	24
Fish meal	66	57	48	57
Fish oil	22	19	16	19
Sum	100	100	100	100
Indicator (Yttriumoxide)	100 mg/kg feed			
Mineral mix	100 mg/kg feed			
Astaxanthin	75 ppm (937.5 mg Carophyll Pink/kg feed)			

The feed was fed *ad libitum* to Atlantic salmon weighing 0.5 kg. After an adaption period of 9 days, the experimental period lasted for 3 weeks. Faeces were collected on day 8 and day 16 according to the method of Austreng, E. (see Aquaculture, 13: 265-272, 1978). Faeces were freeze-dried prior to analysis.

Chemical composition of feed

Chemical composition of feed as revealed by analysis is given in Table 4.

- 13 -

Table 4
Chemical composition of feed (g/100g feed)

Type of feed	Dry matter	Protein	Fat	Starch	beta-glucan	sugars (mono- and di)
Wheat						
HF 12	93.7	47.4	28.5	5.6	0.05	0.21
HF 24	93.7	43.0	25.8	11.5	0.08	0.48
HF 36	93.1	39.0	21.8	17.2	0.07	0.67
HU 24	94.0	42.2	26.2	11.1	0.08	0.20
Barley						
BF 12	94.4	47.3	28.9	4.5	0.28	0.11
BF 24	94.1	42.5	24.8	10.2	0.53	0.38
BF 36	93.6	38.0	22.0	15.3	0.79	0.59
BU 24	94.2	42.8	26.2	8.2	0.40	0.00

Effects of fermentation on digestibility

Digestibility (as %) of protein, fat, starch and energy were calculated from chemical analysis of feed and faeces collected from the fish. The results are given in Table 5.

Table 5

Digestibility (%) of fish feed fed to Atlantic salmon

Type of feed	Protein	Fat	Starch	Energy
Wheat				
HF 12	83.3	88.7	70.9	83.6
HF 24	84.3	89.6	57.2	81.6
HF 36	85.5	84.5	38.4	75.8
HU 24	84.3	86.5	47.9	79.8
Barley				
BF 12	84.4	85.4	68.9	81.7
BF 24	85.7	86.9	63.2	81.1
BF 36	85.8	84.4	48.0	75.4
BU 24	84.6	87.3	33.2	80.2

For protein, there was a tendency for increased digestibility with increasing amounts of fermented wheat or barley flour in the fish feed. For feed containing barley, there was a slightly higher protein digestibility when the flour was fermented (85.7) compared with non-fermented feed (84.6). These feeds were both formulated with 24 % barley flour.

For fats, no systematic variation in % digestibility due to carbohydrate level or treatment were seen.

For starch digestibility, significant positive effects of fermentation were obtained for both wheat- and barley-containing feeds. In feeds containing 24% fermented wheat flour, starch digestibility was 57.2% compared with 47.9% for unfermented feed. This corresponds to an increase in starch digestibility of 19%.

For barley the effect of starch digestibility due to fermentation was even more pronounced; non-fermented feed with 24% flour had a digestibility of only 33.2% while that of the fermented feed was 63.2%. This is an



- 15 -

increase in starch digestibility of more than 90%.

As is known from previous research, starch digestibility decreases with increased levels of cereals. This was also seen in the present experiments. However, for barley the fermentation process compensated for this so that the starch digestibility of feed containing 36% fermented barley (48.0%) was higher than the digestibility of feed containing the lower level of 24% unfermented barley (33.2%).

The increased digestibility of starch due to fermentation was supported by the results for energy digestibility. The increased digestibility of fermented samples caused higher energy digestibility than the unfermented samples. For wheat containing 24% flour the digestibilities were 81.6% and 79.8% for fermented and unfermented feed respectively. For barley, the corresponding figures were 81.1% and 80.2%.

Effects of fermentation on fish performance

The salmon performed well during the experimental period with no problems with appetite in the fish. Faeces were of normal consistency, even at the higher carbohydrate levels. This is reflected in normal dry matter content of the faeces, with no sign of diarrhoea (dry matter below 12%).

Type of feed	Dry matter in faeces (g/100g)
Wheat	
HF 12	15.0
HF 24	14.4
HF 36	15.7
HU 24	14.3
Barley	
BF 12	14.9
BF 24	13.6
BF 36	17.2
BU 24	15.3

Effects of fermentation on carbohydrate composition

The solubility of carbohydrates in the feed was influenced by the fermentation process. Water soluble carbohydrate contents of wheat and barley flour prior to and after fermentation are given in Table 6.

Table 6
Carbohydrates (g/100g feed) in wheat and barley flour
prior to and after fermentation

Sample	Glucose (G)	Maltose (M)	Water soluble polysaccharide	Total G+M	Total
Unfermented					
Wheat	0.72	5.72	2.05	6.44	8.49
Barley	1.05	4.81	5.61	5.86	11.47
Fermented					
Wheat	2.04	1.16	1.24	3.20	4.44
Barley	2.00	0.06	2.76	2.06	4.82

- 17 -

As these results show, the content of maltose decreased considerably, whilst the amount of glucose increased during fermentation. The amount of mono- and disaccharides was nearly half that of the unfermented flour for both wheat and barley. Also the amount of water soluble polysaccharides decreased during the fermentation process. Approximately half as much water soluble polymeric compounds was detected after fermentation. The lowering in soluble polysaccharides was most extensive in barley flour where the initial level of 11.5% was lowered to 4.8%, a reduction of 59%.

Immunostimulating effects

Experimental fish feed was fed to salmon for 16 days and blood samples were tested for serum lysozyme activity and used to evaluate the immunostimulating effect in the blood. The results are shown in Table 7.

Table 7

Serum lysozyme activity in salmon fed feed containing fermented and unfermented barley

Type of feed	Lysozyme activity (%)
Barley	
BF 12	28.445 ^{abc} *
BF 24	32.852 ^a
BF 36	28.480 ^{abc}
BU 24	23.004 ^c

* figures denoted with the same letter are not significantly different ($P < 0.5$)

Serum lysozyme activity was determined by the *Micrococcus lysoplate* assay. In this method the sample is placed in a small well stamped out in an agarose gel containing the bacteria *Micrococcus lysodeicticus*. After incubation, the agarose gels are stained and

- 18 -

destained and the diameter of the lysed zones are measured. The results are compared with the diameter of a reference serum (100%) known to contain lysozyme activity and given as a percent of the diameter of the reference. Actual levels of lysozymes may vary between groups of fish due to health status, age etc. The effect in a special trial is evaluated as the relative lysozyme activities of fish within that trial.

Lysozyme is an enzyme which hydrolyses cell wall components of bacteria. By degradation of the cell walls, the growth of the bacteria is inhibited. High lysozyme activity may therefore give an increased resistance toward pathogen bacteria and thereby enhanced health to the animal.

The results clearly demonstrate increased levels of lysozyme activity in salmon fed with fermented barley compared with salmon fed with unfermented feed. Lysozyme activity was significantly higher in salmon fed with 24% fermented barley (32.85%) compared with salmon fed with the same amount of unfermented barley (23.00%).

- 19 -

The claims defining the invention are as follows:-

1. An animal feed comprising a cereal fermented by lactic acid bacteria together with at least one animal feedstuff component, said
5 cereal being selected from the group consisting of wheat, oats, rye and barley, wherein said cereal is grain fragment, crushed grain, milled grain, wholemeal wheat, wholemeal barley or finely ground whole grain flour or coarsely ground barley, wherein said feed has been subjected to expansion, compression, extrusion and/or drying.
- 10 2. A feed as claimed in claim 1 wherein said cereal is oats or barley.
3. A feed as claimed in any preceding claim having an improved
15 digestibility over its non-fermented cereal-containing animal feed analogue.
4. A feed as claimed in claim 3 having improved starch
20 digestibility over its non-fermented cereal-containing animal feed analogue.
5. A feed as claimed in any preceding claim wherein the animal
feedstuff component is selected from unfermented cereals, bacteriocin, fish meal and pigment.
- 25 6. A feed as claimed in any preceding claim comprising one or more bacteriocins.
7. A feed as claimed in claim 6 wherein the bacteriocin has an
30 inhibitory effect against pathogen producing bacteria.
8. A feed as claimed in claim 7 wherein said bacteria is a species of listeria.
- 35 9. A feed as claimed in any preceding claim comprising an amount of short chain (1-3) β -glucan capable of achieving an immunostimulatory response.



- 20 -

10. A feed as claimed in any preceding claim comprising an amount of fibre capable of having a cholesterol reducing effect.

5 11. An animal feed as claimed in any preceding claim wherein the feed is in the form of tablets, cakes, bricks, pellets, flakes or powders.

10 12. An animal feed as claimed in any preceding claim wherein the fermentation is carried out in an aqueous medium containing 40 to 60% by wheat cereal.

15 13. An animal feed as claimed in any preceding claim wherein fermentation is carried out at a temperature of 25 to 35°C.

14. An animal feed as claimed in any preceding claim wherein fermentation is carried out for 10 to 48 hours.

20 15. A process for producing an animal feed comprising the steps of:

(a) fermenting a cereal with a lactic acid bacteria, said cereal being selected from wheat, barley, rye and oats;

(b) admixing the fermented cereal with at least one animal feedstuff; and optionally,

25 (c) processing said admixture by expansion, compression, extrusion and/or drying.

30 16. A process as claimed in claim 15 wherein the feed is in the form of tablets, cakes, bricks, pellets, flakes or powders.

17. A process as claimed in claim 15 or claim 16 wherein said lactic acid bacteria is a bacteriocin-producing strain of lactobacillus bacteria.

35 18. A process as claimed in claim 17 wherein said bacteriocin-producing lactobacillus strain is a Sakacin A or Sakacin P producing lactobacillus strain.



- 21 -

19. A process as claimed in any of claims 15 to 18 wherein bacteriocins or bacteriocin-producing bacterial strains are added in a further step.

5

20. A process as claimed in claim 19 wherein bacteriocins are added to the admixture prior to or after step (b).

10

21. A process as claimed in any of claims 17 to 20 wherein the bacteriocin has an inhibitory effect against pathogen producing bacteria.

15

22. A process as claimed in claim 21 wherein said bacteria is a species of listeria.

23. A process as claimed in any one of claims 15 to 22 wherein the fermentation is carried out in an aqueous medium containing 40 to 60% by wheat cereal.

20

24. A process as claimed in any one of claims 16 to 23 wherein fermentation is carried out at a temperature of 25 to 35°C.

25. A process as claimed in any one of claims 16 to 24 wherein fermentation is carried out for 10 to 48 hours.


25

26. Use of a cereal fermented by lactic acid bacteria as defined in any one of claims 1 to 14 as a component of fish feed.

30

27. Use as claimed in claim 23 of a bacteriocin containing cereal fermented by lactic acid bacteria as a component of fish feed.

28. A method of reducing the level of cholesterol in a non-human animal body by administering thereto an animal feed as defined in claim 1 together with an effective amount of fibre.



29. A method of achieving an immunostimulatory response in a non-human animal body by administering thereto an animal feed as defined in claim 1.

5 30. Use of a cereal fermented by lactic acid bacteria in the
manufacture of an animal feed for use in reducing the level of
cholesterol in the animal to which the feed is fed.

31. Use of a cereal fermented by lactic acid bacteria in the
10 manufacture of an animal feed for use in a method of achieving an
immunostimulatory response in an animal body.

32. A feed as defined in claim 1, a process as defined in claim 15, use as defined in any one of claims 26, 27 or 28 substantially as hereinbefore defined with reference to the Examples.

DATED this 19th day of April, 2000.

20 MATFORSK, NORWEGIAN FOOD RESEARCH INSTITUTE
By their Patent Attorneys:
CALLINAN LAWRIE

R. E. Nelson

