





(86) Date de dépôt PCT/PCT Filing Date: 2016/03/04
(87) Date publication PCT/PCT Publication Date: 2016/09/09
(85) Entrée phase nationale/National Entry: 2017/08/22
(86) N° demande PCT/PCT Application No.: US 2016/020802
(87) N° publication PCT/PCT Publication No.: 2016/141258
(30) Priorités/Priorities: 2015/03/04 (IN615/DEL/2015);
2015/03/09 (IN640/DEL/2015)

(51) Cl.Int./Int.Cl. *C07D 409/14* (2006.01),
A61K 31/381 (2006.01), *A61K 31/395* (2006.01),
A61K 31/404 (2006.01), *A61K 31/4178* (2006.01),
A61K 31/433 (2006.01), *A61K 31/437* (2006.01),
A61K 31/443 (2006.01), *A61K 31/4436* (2006.01),
A61K 31/4439 (2006.01), *A61K 31/454* (2006.01),
A61K 31/4545 (2006.01), ...

(71) Demandeur/Applicant:
MEDIVATION TECHNOLOGIES, INC., US

(72) Inventeurs/Inventors:
PUJALA, BRAHMAM, IN;
JANGIR, RAMNIWAS, IN;
GUGULOTH, RAMBABU, IN; ...

(54) Titre : INHIBITEURS DES PROTEINES DE LIAISON A L'ELEMENT DE REGULATION DES STEROLS (SREBP)
(54) Title: STEROL REGULATORY ELEMENT-BINDING PROTEINS (SREBPS) INHIBITORS



(57) **Abrégé/Abstract:**

This disclosure provides compounds (Ia) and (Ib) and those compounds for use to treat metabolic disorders and hyperproliferative disorders, including administration of the compounds in conjunction with hormone receptor antagonists.



(51) Cl.Int./Int.Cl. (suite/continued) *A61K 31/4985* (2006.01), *A61K 31/506* (2006.01), *A61P 3/00* (2006.01), *A61P 35/00* (2006.01), *C07D 409/04* (2006.01), *C07D 413/14* (2006.01), *C07D 417/04* (2006.01), *C07D 417/14* (2006.01), *C07D 471/04* (2006.01), *C07D 487/04* (2006.01), *C07D 491/107* (2006.01)

(72) Inventeurs(suite)/Inventors(continued): SHINDE, BHARAT UTTAM, IN; RAI, ROOPA, US; PHAM, SON MINH, US; BERNALES, SEBASTIAN, US; LINDQUIST, JEFFREY, US; GUHA, MAUSUMEE, US; KALLEM, SATYANARAYANA, IN; BHATT, BHAWANA, IN; BHAGWAT, VIKAS RAMDAS, IN

(74) Agent: SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

9 September 2016 (09.09.2016)

WIPO | PCT



(10) International Publication Number

WO 2016/141258 A1

(51) International Patent Classification:

C07D 401/14 (2006.01) A61K 31/42 (2006.01)
 C07D 413/14 (2006.01) A61K 31/427 (2006.01)
 C07D 409/04 (2006.01) A61K 31/381 (2006.01)
 C07D 409/14 (2006.01) A61K 31/4436 (2006.01)
 C07D 417/04 (2006.01) A61P 3/00 (2006.01)
 C07D 417/14 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2016/020802

(22) International Filing Date:

4 March 2016 (04.03.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

615/DEL/2015 4 March 2015 (04.03.2015) IN
 640/DEL/2015 9 March 2015 (09.03.2015) IN

(71) Applicant: MEDIVATION TECHNOLOGIES, INC.

[US/US]; 525 Market Street, 36th Floor, San Francisco,
 California 94105 (US).

(72) Inventors: PUJALA, Brahmam; c/o Integral Biosciences Pvt Ltd., C-64 Hosiery Complex Phase-II Extension, Noida 201306 (IN). JANGIR, Ramniwas; c/o Integral Biosciences Pvt Ltd., C-64 Hosiery Complex Phase-II Extension, Noida 201306 (IN). GUGULOTH, Rambabu; c/o Integral Biosciences Pvt Ltd., C-64 Hosiery Complex Phase-II Extension, Noida 201306 (IN). SHINDE, Bharat Uttam; c/o Integral Biosciences Pvt Ltd., C-64 Hosiery Complex Phase-II Extension, Noida 201306 (IN). RAI, Roopa; c/o Medivation Technologies, Inc., 525 Market Street, 36th Floor, San Francisco, California 94105 (US). PHAM, Son Minh; c/o Medivation Technologies, Inc., 525 Market Street, 36th Floor, San Francisco, California 94105 (US). BERNALES, Sebastian; c/o Medivation Technologies, Inc., 525 Market Street, 36th Floor, San Francisco, CA 94105 (US). LINDQUIST, Jeffrey; c/o Medivation Technologies, Inc., 525 Market Street, 36th

Floor, San Francisco, California 94105 (US). GUHA, Mausumee; c/o Medivation Technologies, Inc., 525 Market Street, 36th Floor, San Francisco, California 94105 (US). KALLEM, Satyanarayana; c/o Integral Biosciences Pvt Ltd., C-64 Hosiery Complex Phase-II Extension, Noida 201306 (IN). BHATT, Bhawana; c/o Integral Biosciences Pvt Ltd., C-64 Hosiery Complex Phase-II Extension, Noida 201306 (IN). BHAGWAT, Vikas Ramdas; c/o Integral Biosciences Pvt Ltd., C-64 Hosiery Complex Phase-II Extension, Noida 201306 (IN).

(74) Agent: HEMMENDINGER, Lisa M.; Banner & Witcoff, Ltd., 1100 13th St NW, Suite 1200, Washington, District of Columbia 20005 (US).



(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: STEROL REGULATORY ELEMENT-BINDING PROTEINS (SREBPS) INHIBITORS



(57) Abstract: This disclosure provides compounds (1a) and (1b) and those compounds for use to treat metabolic disorders and hyperproliferative disorders, including administration of the compounds in conjunction with hormone receptor antagonists.

WO 2016/141258 A1

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 180

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 180

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

STEROL REGULATORY ELEMENT-BINDING PROTEINS (SREBPS) INHIBITORS

[001] This application claims priority to and incorporates by reference Indian applications 615/DEL/2015 filed on March 4, 2015 and 640/DEL/2015 filed on March 9, 2015. Each reference cited in this disclosure is incorporated herein by reference in its entirety.

TECHNICAL FIELD

[002] This disclosure relates generally to therapeutics for treatment of hyperproliferative disorders, metabolic disorders, and pancreatitis.

BACKGROUND


[003] Sterol regulatory element-binding proteins (SREBPs) are major transcription factors regulating the biosynthesis of cholesterol, fatty acid, and triglyceride. They control the expression of crucial genes involved in lipogenesis and uptake. Inhibition of the SREBP pathway can reduce lipid biosynthesis and thus can be a strategy to treat metabolic diseases, such as type II diabetes, insulin resistance, fatty liver and atherosclerosis [Xiao et al. *Acta Biochim. Biophys. Sin* (2013) 45:1, pp 2-10]. In mammals, three SREBP isoforms are known, designated SREBP-1a, SREBP-1c, and SREBP-2. SREBP-1a controls a broad range of SREBP targets including production of fatty acids, triglycerides, phospholipids and cholesterol. SREBP-1c preferentially activates genes of fatty acid and triglyceride metabolism, whereas SREBP-2 preferentially activates genes of cholesterol metabolism, both of which have been studied in human and mice models [Horton et al. *J. Clin. Invest.* (2002) 109:9, pp 1125-1131], as well as *Drosophila* [Rawson. *Nature Rev. Mol. Cell Biol.* (2003) 4:8, pp 631-640].

[004] Recent studies have also presented a link between upregulation of lipid synthesis and prostate cancer [Suburu et al. *Prostaglandins Other Lipid Mediat.* (2012) 98:0, pp 1-10]. The metabolic shift from catabolic to anabolic metabolism is a hallmark of cancer cells. Many cancers require synthesis of fatty acids, and other lipids such as cholesterol and androgens are implicated in prostate cancer. SREBP-1c is the major transcriptional regulator of enzymes in the fatty acid synthesis pathway, and its expression can be stimulated by androgens and epidermal growth factor (EGF) in prostate cancer cells. Overexpression of SREBP-1c is sufficient to cause tumorigenicity and invasion of prostate cancer cells. SREBP-1 can also increase expression of NOX5, a prominent producer of reactive oxygen species (ROS) and regulator of prostate cancer cell growth [Brar et al. *Am. J. Physiol. Cell Physiol.* (2003) 285:2, pp C353-369; Huang et al. *Mol. Cancer Res.* (2012) 10:1, pp 133-142; Huang et al. *Cancer Research* (2012) 72:8, SUPPL. 1; Huang et al. *Mol. Cancer Res.* (2014) 13:4, pp 855-866].


[005] SREBP-2, a regulator of androgen synthesis, is also itself regulated by androgens, demonstrating a direct feedback circuit for regulation of androgen production. SREBP-2 expression increases during disease progression and is significantly higher after castration. This transcription factor also lacks its feedback inhibition in prostate cancer cells, implicating a role for cholesterol and androgen synthesis in prostate cancer [Eberle et al. *Biochimie* (2004) 86:11, pp 839-848; Ettinger et al. *Cancer Res.* (2004), 64:6, pp2212-2221; Chen et al. *Int. J. Cancer* (2001), 91:1, pp 41-45].

[006] Blocking SREBP functions linked to disease states therefore represents an important therapeutic approach for limiting lipid/cholesterol synthesis in membrane production which occurs in metabolic diseases and in cancer progression, as well as in viral pathogenesis [Naar et al. *Clin. Lipidol.* (2012) 7:1, pp 27-36]. Small molecule therapeutics affecting metabolic regulators such as mTOR, AMPK or SIRT1, including Rapamycin, Metformin, or Resveratrol, respectively, may impinge on the transcriptional activity of SREBPs. Recently, two non-sterol small molecules, fatostatin and betulin have been found to inhibit SREBP processing [Kamisuki et al. *Chem. Biol.* (2009) 16:8, pp 882-892; Tang et al. *Cell. Metab.* (2011) 13:1, pp 44-56]. Methods for the treatment of cancers having a p53 mutation, such as breast cancer cells, using SREBP inhibitors have been presented [Freed-Pastor et al. *PCT. Publication* WO2013-110007A1].

[007] Fatostatin analogs have recently been described as potential therapeutics for the treatment of metabolic disorders [Uesugi et al. *U.S. Patent No.* 8,207,196]. Key compounds presented therein are based around Formula X:



Fatostatin




Formula X

wherein R is H, F, Cl, Br, OBz, OH, OCH₃, OCH₂CO₂Me, OCH₂CO₂H, NH₂, NH_iPr, NHCOCH₃, NHSO₂Me, NH[benzyl], NH[cyclopropyl], NH[tertbutyloxycarbonyl], NH[cyclohexyl], NH[tosyl], NH[quinolin-8-yl], and NH[thiophen-2-yl]. In particular, one compound (FGH10019), the methanesulfonamide derivative of fatostatin above wherein R is NHSO₂Me, has been described as a lead candidate [Kamisuki et al. *J. Med. Chem.* (2011) 54:13, pp 4923-4927]. Further examples of Fatostatin analogs have been presented [Chakravarty et al. *PCT Publication* WO2015/031650A1].

BRIEF SUMMARY

[008] This disclosure relates generally to therapeutics for treatment of hyperproliferative disorders, metabolic disorders, and pancreatitis. This disclosure provides compounds and methods of using those compounds to treat benign and malignant hyperproliferative disorders, as well as metabolic disorders and pancreatitis, as described below. The disclosed compounds may also find use in treating cancer.

[009] Compounds disclosed herein fall within Formulae (Ia) or (Ib):



or a pharmaceutically acceptable salt thereof, wherein:

A is either:

- i. a heteroaryl having only one ring, substituted with one or more **R1** or **R2**; or
- ii. an aryl or heteroaryl, each having more than one ring, optionally substituted with one or more **R1** or **R3**;

R1 is C1-C6 linear or branched alkyl, C2-C6 linear or branched alkenyl, C2-C6 linear or branched alkynyl, C3-C6 cycloalkyl, C3-C6 cycloalkylmethyl, C3-C6 cycloalkenyl, aryl, or heteroaryl; each optionally substituted with one or more **R3**;

R2 is halogen, -CN, -OH, -O(Alkyl), -NO₂, -SH, -S(Alkyl), -S(O)(Alkyl), -S(O)₂(Alkyl), -CH₂OCH₃, -OBn, -CO₂H, -CO₂(Alkyl), -NR₁₀R₁₁, -CONR₁₀R₁₁, and -S(O)₂NR₁₀R₁₁;

R3 is halogen, -CN, -OH, -O(Alkyl), =O, -NO₂, -SH, -S(Alkyl), -S(O)(Alkyl), -S(O)₂(Alkyl), -CH₂OCH₃, -OBn, -CO₂H, -CO₂(Alkyl), -NR₁₀R₁₁, -CONR₁₀R₁₁, and -S(O)₂NR₁₀R₁₁;

Each **R_{C4}**, **R_{C5}**, and **R_{C6}**, is independently hydrogen, **R1**, or **R2**; or **R_{C4}** is taken with **R_{C7}** to form a heterocyclyl optionally substituted with one or more **R3**;

Each **R_{C7}** and **R_{C8}** is independently hydrogen, **R1**, or -SO₂**R_{C12}**; or **R_{C7}** and **R_{C8}** are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more **R1** or **R3**; or **R_{C7}** is taken with **R_{C4}** to form a heterocyclyl optionally substituted with one or more **R3**;

R10 and **R11** are independently hydrogen, C1-C6 linear or branched alkyl, $-\text{C}(\text{O})\text{R}_{\text{C12}}$, $-\text{C}(\text{O})_2\text{R}_{\text{C12}}$, $-\text{C}(\text{O})\text{N}(\text{R}_{\text{C12}})_2$, $-\text{SO}_2\text{R}_{\text{C12}}$, or are taken together with the N to which they are attached to form a C3-C8 heterocyclyl;

R_{C12} is hydrogen, a linear or branched C1-C6 alkyl, a linear or branched C2-C6 alkenyl, a linear or branched C2-C6 alkynyl, or aryl; wherein each alkyl, alkenyl, alkynyl, or aryl group is optionally substituted with one or more halogen atoms, one or more $-\text{NH}_2$ groups, or one or more $-\text{OH}$ groups;

V is the moiety $-\text{C}(=\text{O})-$ or $-\text{S}(=\text{O})_2-$; and

The ring comprising **W**, **X**, and **Y** is a 5-membered heteroaromatic, wherein **W** and **Y** are independently selected from the group consisting of **CR_A**, N, **NR_B**, O and S; **X** is selected from the group consisting of **NR_B**, O and S; wherein **R_A** is hydrogen, **R1** or **R2**; and **R_B** is hydrogen or **R1**;

With the proviso that when **W** is N, **X** is S, and **Y** is C, then one of **R_{C4}**, **R_{C5}** and **R_{C6}** is other than H.

[010] Examples of some of the disclosed compounds are described in Table 1, such as a compound selected from the group consisting of Compound Nos. 1 to 104; or a pharmaceutically acceptable salt, solvate, prodrug, or N-oxide thereof.

[011] Examples of other of the disclosed compounds are described in Table 2, such as a compound selected from the group consisting of Compound Nos. 2.1 to 2.152; or a pharmaceutically acceptable salt, solvate, prodrug, or N-oxide thereof.

[012] Further provided is a pharmaceutical composition, comprising a compound of Formulae (Ia) or (Ib), or any variations described herein, or a salt thereof, and a pharmaceutically acceptable carrier or excipient.

[013] Further provided is a kit, comprising a compound of Formulae (Ia) or (Ib), or any variations described herein, or a salt thereof, and instructions for use.

[014] Further provided are methods of treating one or more of the following: hyperproliferative disorders, metabolic disorders and/or pancreatitis in individuals in need thereof, such as humans, comprising administering to an individual in need thereof a therapeutically effective amount of a compound of Formulae (Ia) or (Ib), or any variations described herein, or a pharmaceutically acceptable salt thereof.

[015] Further provided are methods of treating any diseases or conditions for which the modulation of SREBP is believed to be or is beneficial, comprising administering to an

individual in need thereof a therapeutically effective amount of a compound of Formulae (Ia) or (Ib), or any variations described herein, or a pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION

Definitions

[016] For use herein, unless clearly indicated otherwise, use of the terms “a,” “an” and the like refers to one or more.

[017] As used herein, reference to “about” a value or parameter herein includes (and describes) embodiments that are directed to that value or parameter per se. For example, description referring to “about X” includes description of “X.”

[018] “Alkyl” refers to and includes saturated linear or branched univalent hydrocarbon structures and combinations thereof. Particular alkyl groups are those having 1 to 6 carbon atoms (a “C1-C6 alkyl”). Unless otherwise defined, “C1-C6 linear or branched alkyl” means methyl, ethyl, and C1-C6, C1-C5, C1-C4, C1-C3, C1-C2, C2-C6, C2-C5, C2-C4, C2-C3, C3-C6, C3-C5, C3-C4, C4-C6, C4-C5, C5-C6, C3, C4, C5, and C6 linear or branched alkyl. When an alkyl residue having a specific number of carbons is named, unless otherwise defined, all geometric isomers having that number of carbons are intended to be encompassed and described; thus, for example, “butyl” is meant to include *n*-butyl, *sec*-butyl, *iso*-butyl, and *tert*-butyl; “propyl” includes *n*-propyl and *iso*-propyl. Examples of alkyl include methyl, *t*-butyl, *n*-heptyl, octyl, and the like.

[019] “Alkenyl” refers to an unsaturated hydrocarbon group having at least one site of olefinic unsaturation (*i.e.*, having at least one moiety of the formula C=C). Unless otherwise defined, “C2-C6 linear or branched alkenyl” means ethenyl and C2-C6, C2-C5, C2-C4, C2-C3, C3-C6, C3-C5, C3-C4, C4-C6, C4-C5, C5-C6, C3, C4, C5, and C6 linear or branched alkenyl. Examples of alkenyl include but are not limited to ethenyl “-CH=CH₂,” -CH₂-CH=CH-CH₃ and -CH=CH-CH=CH₂.

[020] “Alkynyl” refers to an unsaturated hydrocarbon group having at least one site of acetylenic unsaturation (*i.e.*, having at least one moiety of the formula C≡C). Unless otherwise defined, “C2-C6 linear or branched alkynyl” means ethynyl and C2-C6, C2-C5, C2-C4, C2-C3, C3-C6, C3-C5, C3-C4, C4-C6, C4-C5, C5-C6, C3, C4, C5, and C6 linear or branched alkynyl. Examples of alkynyl include but are not limited to ethynyl “-C≡CH,” -CH₂-C≡C-CH₃ and -C≡C-C≡CH.

[021] “Cycloalkyl” refers to and includes cyclic univalent hydrocarbon structures. Cycloalkyl can consist of one ring, such as cyclohexyl, or multiple rings, such as adamantyl. A cycloalkyl comprising more than one ring may be fused, spiro or bridged, or combinations thereof. A preferred cycloalkyl is a saturated cyclic hydrocarbon having from 3 to 13 annular carbon atoms. A more preferred cycloalkyl is a saturated cyclic hydrocarbon having from 3 to 6 annular carbon atoms (a “C3-C6 cycloalkyl”). Examples of cycloalkyl groups include adamantyl, decahydronaphthalenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like. Unless otherwise defined, “C3-C6 cycloalkyl” means C3-C6, C3-C5, C3-C4, C4-C6, C4-C5, C5-C6, C3, C4, C5, and C6 cycloalkyl.

[022] “Cycloalkenyl” refers to an unsaturated hydrocarbon group within a cycloalkyl having at least one site of olefinic unsaturation (*i.e.*, having at least one moiety of the formula C=C). Cycloalkenyl can consist of one ring, such as cyclohexyl, or multiple rings, such as norbornenyl. A more preferred cycloalkenyl is an unsaturated cyclic hydrocarbon having from 3 to 8 annular carbon atoms (*e.g.*, “C3-C8 cycloalkenyl,” “C3-C6 cycloalkenyl”). Unless otherwise defined, “C3-C6 cycloalkenyl” means C3-C6, C3-C5, C3-C4, C4-C6, C4-C5, C5-C6, C3, C4, C5, and C6 cycloalkenyl. Examples of cycloalkenyl groups include cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, and the like.

[023] “Heterocycle,” “heterocyclic,” or “heterocyclyl” refers to a saturated or an unsaturated non-aromatic group having a single ring or multiple condensed rings, and having from 1 to 10 annular carbon atoms and from 1 to 4 annular heteroatoms, such as nitrogen, sulfur or oxygen, and the like. A heterocycle comprising more than one ring may be fused, spiro or bridged, or any combination thereof. In fused ring systems, one or more of the rings can be aryl or heteroaryl. A heterocycle having more than one ring where at least one ring is aromatic may be connected to the parent structure at either a non-aromatic ring position or at an aromatic ring position. In one variation, a heterocycle having more than one ring where at least one ring is aromatic is connected to the parent structure at a non-aromatic ring position. Unless otherwise defined, “C3-C8 heterocycle” (or “heterocyclyl” or “heterocyclic”) means C3-C8, C3-C7, C3-C6, C3-C4, C3-C5, C4-C8, C4-C7, C4-C6, C4-C5, C5-C8, C5-C7, C5-C6, C6-C8, C6-C7, C7-C8, C3, C4, C5, C6, C7, and C8 heterocycle (or heterocyclyl or heterocyclic).

[024] “Aryl” as used herein refers to an unsaturated aromatic carbocyclic group having a single ring (*e.g.*, phenyl) or multiple condensed rings (*e.g.*, naphthyl or anthryl) which condensed rings may or may not be aromatic. The aryl group may be optionally substituted independently with one or more substituents described herein. Particular aryl groups are those having from 6 to 14 annular (*i.e.*, ring) carbon atoms (a “C6-C14 aryl”). An aryl group having more than one ring

where at least one ring is non-aromatic may be connected to the parent structure at either an aromatic ring position or at a non-aromatic ring position. In one variation, an aryl group having more than one ring where at least one ring is non-aromatic is connected to the parent structure at an aromatic ring position.

[025] “Heteroaryl” as used herein refers to an unsaturated aromatic cyclic group having from 1 to 14 annular (i.e., ring) carbon atoms and at least one annular heteroatom, including but not limited to heteroatoms such as nitrogen, oxygen and sulfur. A heteroaryl group may have a single ring (e.g., pyridyl, furyl) or multiple condensed rings (e.g., indolizinyl, benzothienyl) which condensed rings may or may not be aromatic. The heteroaryl group may be optionally substituted independently with one or more substituents described herein. Particular heteroaryl groups are 5- to 14-membered rings having 1 to 12 annular (i.e., ring) carbon atoms and 1 to 6 annular (i.e., ring) heteroatoms independently selected from nitrogen, oxygen and sulfur; 5- to 10-membered rings having 1 to 8 annular carbon atoms and 1 to 4 annular heteroatoms independently selected from nitrogen, oxygen and sulfur; and 5-, 6- or 7-membered rings having 1 to 5 annular carbon atoms and 1 to 4 annular heteroatoms independently selected from nitrogen, oxygen and sulfur. In one variation, heteroaryl includes monocyclic aromatic 5-, 6- or 7-membered rings having from 1 to 6 annular carbon atoms and 1 to 4 annular heteroatoms independently selected from nitrogen, oxygen and sulfur. In another variation, heteroaryl includes polycyclic aromatic rings having from 1 to 12 annular carbon atoms and 1 to 6 annular heteroatoms independently selected from nitrogen, oxygen and sulfur. A heteroaryl group having more than one ring where at least one ring is non-aromatic may be connected to the parent structure at either an aromatic ring position or at a non-aromatic ring position. In one variation, a heteroaryl group having more than one ring where at least one ring is non-aromatic is connected to the parent structure at an aromatic ring position.

[026] “Alkoxy” refers to the group alkyl-O-, which includes, by way of example, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, tert-butoxy, sec-butoxy, n-pentoxy, n-hexoxy, 1,2-dimethylbutoxy, and the like.

[027] “Hydroxyalkyl” refers to the group alkyl-OH, which includes, by way of example, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 1-hydroxyprop-1-yl, 2-hydroxyprop-2-yl, 3-hydroxypropyl, 4-hydroxybutyl, 5-hydroxypentyl, 6-hydroxyhexyl, and the like.


[028] “Halogen” refers to elements of the Group 17 series having atomic number 9 to 85. Preferred halo groups include the radicals of fluorine, chlorine, bromine and iodine. Where a residue is substituted with more than one halogen, it may be referred to by using a prefix corresponding to the number of halogen moieties attached, e.g., dihaloaryl, dihaloalkyl,

trihaloaryl etc. refer to aryl and alkyl substituted with two (“di”) or three (“tri”) halo groups, which may be but are not necessarily the same halogen; thus 4-chloro-3-fluorophenyl is within the scope of dihaloaryl. An alkyl group in which each H is replaced with a halo group is referred to as a “perhaloalkyl.” An alkenyl group in which each H is replaced with a halo group is referred to as a “perhaloalkenyl.” An alkynyl group in which each H is replaced with a halo group is referred to as a “perhaloalkynyl.” A preferred perhaloalkyl group is trifluoroalkyl (-CF₃). Similarly, “perhaloalkoxy” refers to an alkoxy group in which a halogen takes the place of each H in the hydrocarbon making up the alkyl moiety of the alkoxy group. An example of a perhaloalkoxy group is trifluoromethoxy (-OCF₃).

[029] Any and all stereoisomers, including geometric isomers (*cis/trans* or *E/Z* isomers), tautomers, salts, N-oxides, and solvates of the compounds described herein can be used in the disclosed methods. This disclosure also provides methods of making such compounds.

Compounds

[030] Compounds disclosed herein fall within Formulae (Ia) or (Ib):



or a pharmaceutically acceptable salt thereof, wherein:

A is either:

- i. a heteroaryl having only one ring, substituted with one or more **R1** or **R2**; or
- ii. an aryl or heteroaryl, each having more than one ring, optionally substituted with one or more **R1** or **R3**;

R1 is C1-C6 linear or branched alkyl, C2-C6 linear or branched alkenyl, C2-C6 linear or branched alkynyl, C3-C6 cycloalkyl, C3-C6 cycloalkylmethyl, C3-C6 cycloalkenyl, aryl, or heteroaryl; each optionally substituted with one or more **R3**;

R2 is halogen, -CN, -OH, -O(Alkyl), -NO₂, -SH, -S(Alkyl), -S(O)(Alkyl), -S(O)₂(Alkyl), -CH₂OCH₃, -OBn, -CO₂H, -CO₂(Alkyl), -NR₁₀R₁₁, -CONR₁₀R₁₁, and -S(O)₂NR₁₀R₁₁;

R3 is halogen, -CN, -OH, -O(Alkyl), =O, -NO₂, -SH, -S(Alkyl), -S(O)(Alkyl), -S(O)₂(Alkyl), -CH₂OCH₃, -OBn, -CO₂H, -CO₂(Alkyl), -NR₁₀R₁₁, -CONR₁₀R₁₁, and -S(O)₂NR₁₀R₁₁;

Each R_{C4} , R_{C5} , and R_{C6} , is independently hydrogen, $R1$, or $R2$; or R_{C4} is taken with R_{C7} to form a heterocyclyl optionally substituted with one or more $R3$;

Each R_{C7} and R_{C8} is independently hydrogen, $R1$, or $-SO_2R_{C12}$; or R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more $R1$ or $R3$; or R_{C7} is taken with R_{C4} to form a heterocyclyl optionally substituted with one or more $R3$;

$R10$ and $R11$ are independently hydrogen, C1-C6 linear or branched alkyl, $-C(O)R_{C12}$, $-C(O)_2R_{C12}$, $-C(O)N(R_{C12})_2$, $-SO_2R_{C12}$, or are taken together with the N to which they are attached to form a C3-C8 heterocyclyl;

R_{C12} is hydrogen, a linear or branched C1-C6 alkyl, a linear or branched C2-C6 alkenyl, a linear or branched C2-C6 alkynyl, or aryl; wherein each alkyl, alkenyl, alkynyl, or aryl group is optionally substituted with one or more halogen atoms, one or more $-NH_2$ groups, or one or more $-OH$ group;

V is the moiety $-C(=O)-$ or $-S(=O)_2-$; and

The ring comprising W , X , and Y is a 5-membered heteroaromatic, wherein W and Y are independently selected from the group consisting of CR_A , N, NR_B , O and S; X is selected from the group consisting of NR_B , O and S; wherein R_A is hydrogen, $R1$ or $R2$; and R_B is hydrogen or $R1$;

With the proviso that when W is N, X is S, and Y is C, then one of R_{C4} , R_{C5} and R_{C6} is other than H.

Variations in which A is a heteroaryl having only one ring

[031] In some variations of Formulae (Ia) or (Ib), A is a heteroaryl having only one ring, substituted with one $R1$ (*i.e.*, there is a first $R1$). In some variations, A is a heteroaryl having only one ring, substituted with two $R1$ (*i.e.*, there is a first $R1$ and a second $R1$). In some variations, A is a heteroaryl having only one ring, substituted with three $R1$ (*i.e.*, there is a first $R1$, a second $R1$, and a third $R1$).

[032] In some variations, the first $R1$ is unsubstituted C1-C6 linear or branched alkyl. In some variations, the first $R1$ is unsubstituted C2-C6 linear or branched alkenyl. In some variations, the first $R1$ is unsubstituted C2-C6 linear or branched alkynyl. In some variations, the first $R1$ is unsubstituted C3-C6 cycloalkyl. In some variations, the first $R1$ is unsubstituted C3-C6 cycloalkylmethyl. In some variations, the first $R1$ is unsubstituted C3-C6 cycloalkenyl. In some variations, the first $R1$ is unsubstituted aryl. In some variations, the first $R1$ is unsubstituted

heteroaryl. In some variations, the first **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some variations, the first **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some variations, the first **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some variations, the first **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some variations, the first **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some variations, the first **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the first **R1** is aryl substituted with one or more **R3**. In some variations, the first **R1** is heteroaryl substituted with one or more **R3**.

[033] In some variations, the second **R1** is unsubstituted C1-C6 linear or branched alkyl. In some variations, the second **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some variations, the second **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some variations, the second **R1** is unsubstituted C3-C6 cycloalkyl. In some variations, the second **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some variations, the second **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the second **R1** is unsubstituted aryl. In some variations, the second **R1** is unsubstituted heteroaryl. In some variations, the second **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some variations, the second **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some variations, the second **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some variations, the second **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some variations, the second **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some variations, the second **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the second **R1** is aryl substituted with one or more **R3**. In some variations, the second **R1** is heteroaryl substituted with one or more **R3**.

[034] In some variations, the third **R1** is unsubstituted C1-C6 linear or branched alkyl. In some variations, the third **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some variations, the third **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some variations, the third **R1** is unsubstituted C3-C6 cycloalkyl. In some variations, the third **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some variations, the third **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the third **R1** is unsubstituted aryl. In some variations, the third **R1** is unsubstituted heteroaryl. In some variations, the third **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some variations, the third **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some variations, the third **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some variations, the third **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some variations, the third **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some variations, the third **R1** is C3-C6 cycloalkenyl substituted with

one or more **R3**. In some variations, the third **R1** is aryl substituted with one or more **R3**. In some variations, the third **R1** is heteroaryl substituted with one or more **R3**.

[035] In some variations, at least one **R3** is halogen. In some variations, at least one **R3** is -CN. In some variations, at least one **R3** is -O(**R12**). In some variations, at least one **R3** is =O. In some variations, at least one **R3** is -NO₂. In some variations, at least one **R3** is -S(**R12**). In some variations, at least one **R3** is -S(O)(**R12**). In some variations, at least one **R3** is -S(O)₂(**R12**). In some variations, at least one **R3** is -CH₂OCH₃. In some variations, at least one **R3** is -OBn. In some variations, at least one **R3** is -CO₂H. In some variations, at least one **R3** is -CO₂(**R12**). In some variations, at least one **R3** is -NR10R11. In some variations, at least one **R3** is -CONR10R11. In some variations, at least one **R3** is -S(O)₂NR10R11.

[036] In some variations of Formulae (Ia) or (Ib), **A** is a heteroaryl having only one ring, substituted with one **R2** (a first **R2**). In some variations, **A** is a heteroaryl having only one ring, substituted with two **R2** (a first **R2** and a second **R2**). In some variations, **A** is a heteroaryl having only one ring, substituted with three **R2** (a first **R2**, a second **R2**, and a third **R2**). In some of these variations, the first **R2** is halogen. In some of these variations, the first **R2** is -CN. In some of these variations, the first **R2** is -OH. In some of these variations, the first **R2** is -O(Alkyl). In some of these variations, the first **R2** is -NO₂. In some of these variations, the first **R2** is -SH. In some of these variations, the first **R2** is -S(Alkyl). In some of these variations, the first **R2** is -S(O)(Alkyl). In some of these variations, the first **R2** is -S(O)₂(Alkyl). In some of these variations, the first **R2** is -CH₂OCH₃. In some of these variations, the first **R2** is -OBn. In some of these variations, the first **R2** is -CO₂H. In some of these variations, the first **R2** is -CO₂(Alkyl). In some of these variations, the first **R2** is -NR10R11. In some of these variations, the first **R2** is -CONR10R11. In some of these variations, the first **R2** is -S(O)₂NR10R11. In some of these variations, the second **R2** is halogen. In some of these variations, the second **R2** is -CN. In some of these variations, the second **R2** is -OH. In some of these variations, the second **R2** is -O(Alkyl). In some of these variations, the second **R2** is -NO₂. In some of these variations, the second **R2** is -SH. In some of these variations, the second **R2** is -S(Alkyl). In some of these variations, the second **R2** is -S(O)(Alkyl). In some of these variations, the second **R2** is -S(O)₂(Alkyl). In some of these variations, the second **R2** is -CH₂OCH₃. In some of these variations, the second **R2** is -OBn. In some of these variations, the second **R2** is -CO₂H. In some of these variations, the second **R2** is -CO₂(Alkyl). In some of these variations, the second **R2** is -NR10R11. In some of these variations, the second **R2** is -CONR10R11. In some of these variations, the second **R2** is -S(O)₂NR10R11. In some of these variations, the third **R2** is halogen. In some of these variations, the third **R2** is -CN. In some of these variations, the third

R2 is –OH. In some of these variations, the third **R2** is –O(Alkyl). In some of these variations, the third **R2** is –NO₂. In some of these variations, the third **R2** is –SH. In some of these variations, the third **R2** is –S(Alkyl). In some of these variations, the third **R2** is –S(O)(Alkyl). In some of these variations, the third **R2** is –S(O)₂(Alkyl). In some of these variations, the third **R2** is –CH₂OCH₃. In some of these variations, the third **R2** is –OBn. In some of these variations, the third **R2** is –CO₂H. In some of these variations, the third **R2** is –CO₂(Alkyl). In some of these variations, the third **R2** is –NR₁₀R₁₁. In some of these variations, the third **R2** is –CONR₁₀R₁₁. In some of these variations, the third **R2** is –S(O)₂NR₁₀R₁₁.

Variations in which A is an aryl or heteroaryl, each having more than one ring

[037] In some variations of Formulae (Ia) or (Ib), **A** is an aryl or heteroaryl, each having more than one ring.

Variations in which A is an aryl having more than one ring

[038] In some variations of Formulae (Ia) or (Ib), **A** is an aryl having more than one ring, substituted with one **R1** (a first **R1**). In some variations, **A** is an aryl having more than one ring, substituted with two **R1** (a first **R1** and a second **R1**). In some variations, **A** is an aryl having more than one ring, substituted with three **R1** (a first **R1**, a second **R1**, and a third **R1**). In some of these variations, the first **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the second **R1** is unsubstituted aryl. In some variations, the second **R1** is unsubstituted heteroaryl. In some of these variations, the first **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the first **R1** is aryl substituted with one or more **R3**. In some variations, the first **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, the second **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the

second **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the second **R1** is unsubstituted aryl. In some variations, the second **R1** is unsubstituted heteroaryl. In some of these variations, the second **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the second **R1** is aryl substituted with one or more **R3**. In some variations, the second **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, the third **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the third **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the third **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the third **R1** is unsubstituted aryl. In some variations, the third **R1** is unsubstituted heteroaryl. In some of these variations, the third **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the third **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the third **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the third **R1** is aryl substituted with one or more **R3**. In some variations, the third **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, at least one **R3** is halogen. In some of these variations, at least one **R3** is -CN. In some of these variations, at least one **R3** is -OH. In some of these variations, at least one **R3** is -O(Alkyl). In some of these variations, at least one **R3** is =O. In some of these variations, at least one **R3** is -NO₂. In some of these variations, at least one **R3** is -SH. In some of these variations, at least one **R3** is -S(Alkyl). In some of these variations, at least one **R3** is -S(O)(Alkyl). In some of these variations, at least one **R3** is -S(O)₂(Alkyl). In some of these variations, at least one **R3** is -CH₂OCH₃. In some of these variations, at least one **R3** is -OBn. In some of these variations, at least one **R3** is -CO₂H. In some of these variations, at

least one **R3** is -CO₂(Alkyl). In some of these variations, at least one **R3** is -NR₁₀R₁₁. In some of these variations, at least one **R3** is -CONR₁₀R₁₁. In some of these variations, at least one **R3** is -S(O)₂NR₁₀R₁₁.

[039] In some variations of Formulae (Ia) or (Ib), **A** is an aryl having more than one ring, substituted with one **R3** (a first **R3**). In some variations, **A** is an aryl having more than one ring, substituted with two **R3** (a first **R3** and a second **R3**). In some variations, **A** is an aryl having more than one ring, substituted with three **R3** (a first **R3**, a second **R3**, and a third **R3**). In some of these variations, the first **R3** is halogen. In some of these variations, the first **R3** is -CN. In some of these variations, the first **R3** is -OH. In some of these variations, the first **R3** is -O(Alkyl). In some of these variations, the first **R3** is =O. In some of these variations, the first **R3** is -NO₂. In some of these variations, the first **R3** is -SH. In some of these variations, the first **R3** is -S(Alkyl). In some of these variations, the first **R3** is -S(O)(Alkyl). In some of these variations, the first **R3** is -S(O)₂(Alkyl). In some of these variations, the first **R3** is -CH₂OCH₃. In some of these variations, the first **R3** is -OBn. In some of these variations, the first **R3** is -CO₂H. In some of these variations, the first **R3** is -CO₂(Alkyl). In some of these variations, the first **R3** is -NR₁₀R₁₁. In some of these variations, the first **R3** is -CONR₁₀R₁₁. In some of these variations, the first **R3** is -S(O)₂NR₁₀R₁₁. In some of these variations, the second **R3** is halogen. In some of these variations, the second **R3** is -CN. In some of these variations, the second **R3** is -OH. In some of these variations, the second **R3** is -O(Alkyl). In some of these variations, the second **R3** is =O. In some of these variations, the second **R3** is -NO₂. In some of these variations, the second **R3** is -SH. In some of these variations, the second **R3** is -S(Alkyl). In some of these variations, the second **R3** is -S(O)(Alkyl). In some of these variations, the second **R3** is -S(O)₂(Alkyl). In some of these variations, the second **R3** is -CH₂OCH₃. In some of these variations, the second **R3** is -OBn. In some of these variations, the second **R3** is -CO₂H. In some of these variations, the second **R3** is -CO₂(Alkyl). In some of these variations, the second **R3** is -NR₁₀R₁₁. In some of these variations, the second **R3** is -CONR₁₀R₁₁. In some of these variations, the second **R3** is -S(O)₂NR₁₀R₁₁. In some of these variations, the third **R3** is halogen. In some of these variations, the third **R3** is -CN. In some of these variations, the third **R3** is -OH. In some of these variations, the third **R3** is -O(Alkyl). In some of these variations, the third **R3** is =O. In some of these variations, the third **R3** is -NO₂. In some of these variations, the third **R3** is -SH. In some of these variations, the third **R3** is -S(Alkyl). In some of these variations, the third **R3** is -S(O)(Alkyl). In some of these variations, the third **R3** is -S(O)₂(Alkyl). In some of these variations, the third **R3** is -CH₂OCH₃. In some of these variations, the third **R3** is -OBn. In some of these variations, the third **R3** is -CO₂H. In some of these variations, the third **R3** is -CO₂(Alkyl). In some of these variations, the third **R3** is -

NR10R11. In some of these variations, the third **R3** is -CONR10R11. In some of these variations, the third **R3** is -S(O)₂NR10R11.

Variations in which A is a heteroaryl having more than one ring


[040] In some variations of Formulae (Ia) or (Ib), **A** is a heteroaryl having more than one ring, substituted with one **R1** (a first **R1**). In some variations, **A** is a heteroaryl having more than one ring, substituted with two **R1** (a first **R1** and a second **R1**). In some variations, **A** is a heteroaryl having more than one ring, substituted with three **R1** (a first **R1**, a second **R1**, and a third **R1**). In some of these variations, the first **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the first **R1** is unsubstituted aryl. In some variations, the first **R1** is unsubstituted heteroaryl. In some of these variations, the first **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the first **R1** is aryl substituted with one or more **R3**. In some variations, the first **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, the second **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the second **R1** is unsubstituted aryl. In some variations, the second **R1** is unsubstituted heteroaryl. In some of these variations, the second **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkylmethyl substituted with one or


more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the second **R1** is aryl substituted with one or more **R3**. In some variations, the second **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, the third **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the third **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the third **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the third **R1** is unsubstituted aryl. In some variations, the third **R1** is unsubstituted heteroaryl. In some of these variations, the third **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the third **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the third **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the third **R1** is aryl substituted with one or more **R3**. In some variations, the third **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, at least one **R3** is halogen. In some of these variations, at least one **R3** is -CN. In some of these variations, at least one **R3** is -OH. In some of these variations, at least one **R3** is -O(Alkyl). In some of these variations, at least one **R3** is =O. In some of these variations, at least one **R3** is -NO₂. In some of these variations, at least one **R3** is -SH. In some of these variations, at least one **R3** is -S(Alkyl). In some of these variations, at least one **R3** is -S(O)(Alkyl). In some of these variations, at least one **R3** is -S(O)₂(Alkyl). In some of these variations, at least one **R3** is -CH₂OCH₃. In some of these variations, at least one **R3** is -OBn. In some of these variations, at least one **R3** is -CO₂H. In some of these variations, at least one **R3** is -CO₂(Alkyl). In some of these variations, at least one **R3** is -NR₁₀R₁₁. In some of these variations, at least one **R3** is -CONR₁₀R₁₁. In some of these variations, at least one **R3** is -S(O)₂NR₁₀R₁₁.


[041] In some variations of Formulae (Ia) or (Ib), **A** is a heteroaryl having more than one ring, substituted with one **R3** (a first **R3**). In some variations, **A** is a heteroaryl having more than one ring, substituted with two **R3** (a first **R3** and a second **R3**). In some variations, **A** is a heteroaryl having more than one ring, substituted with three **R3** (a first **R3**, a second **R3**, and a third **R3**). In some of these variations, the first **R3** is halogen. In some of these variations, the first **R3** is -CN. In some of these variations, the first **R3** is -OH. In some of these variations, the first **R3** is -O(Alkyl). In some of these variations, the first **R3** is =O. In some of these variations, the first **R3**

is -NO₂. In some of these variations, the first **R3** is -SH. In some of these variations, the first **R3** is -S(Alkyl). In some of these variations, the first **R3** is -S(O)(Alkyl). In some of these variations, the first **R3** is -S(O)₂(Alkyl). In some of these variations, the first **R3** is -CH₂OCH₃. In some of these variations, the first **R3** is -OBn. In some of these variations, the first **R3** is -CO₂H. In some of these variations, the first **R3** is -CO₂(Alkyl). In some of these variations, the first **R3** is -NR₁₀R₁₁. In some of these variations, the first **R3** is -CONR₁₀R₁₁. In some of these variations, the first **R3** is -S(O)₂NR₁₀R₁₁. In some of these variations, the second **R3** is halogen. In some of these variations, the second **R3** is -CN. In some of these variations, the second **R3** is -OH. In some of these variations, the second **R3** is -O(Alkyl). In some of these variations, the second **R3** is =O. In some of these variations, the second **R3** is -NO₂. In some of these variations, the second **R3** is -SH. In some of these variations, the second **R3** is -S(Alkyl). In some of these variations, the second **R3** is -S(O)(Alkyl). In some of these variations, the second **R3** is -S(O)₂(Alkyl). In some of these variations, the second **R3** is -CH₂OCH₃. In some of these variations, the second **R3** is -OBn. In some of these variations, the second **R3** is -CO₂H. In some of these variations, the second **R3** is -CO₂(Alkyl). In some of these variations, the second **R3** is -NR₁₀R₁₁. In some of these variations, the second **R3** is -CONR₁₀R₁₁. In some of these variations, the second **R3** is -S(O)₂NR₁₀R₁₁. In some of these variations, the third **R3** is halogen. In some of these variations, the third **R3** is -CN. In some of these variations, the third **R3** is -OH. In some of these variations, the third **R3** is -O(Alkyl). In some of these variations, the third **R3** is =O. In some of these variations, the third **R3** is -NO₂. In some of these variations, the third **R3** is -SH. In some of these variations, the third **R3** is -S(Alkyl). In some of these variations, the third **R3** is -S(O)(Alkyl). In some of these variations, the third **R3** is -S(O)₂(Alkyl). In some of these variations, the third **R3** is -CH₂OCH₃. In some of these variations, the third **R3** is -OBn. In some of these variations, the third **R3** is -CO₂H. In some of these variations, the third **R3** is -CO₂(Alkyl). In some of these variations, the third **R3** is -NR₁₀R₁₁. In some of these variations, the third **R3** is -CONR₁₀R₁₁. In some of these variations, the third **R3** is -S(O)₂NR₁₀R₁₁.


[042] In some variations of Formulae (Ia) or (Ib), **A** is a moiety selected from the group consisting of:







[047] In some variations of Formulae (Ia) or (Ib), the ring comprising **W**, **X**, and **Y**, and the carbon atoms to which they are attached, is a 5-membered heteroaromatic group selected from the group consisting of:




wherein **RA** is hydrogen, **R1**, or **R2**; and **RB** is hydrogen or **R1**. In some variations, the 5-

membered group is . In particular variations, the 5-


membered group is . In some of these embodiments, **RA** is hydrogen and **RB** is hydrogen. In some of these embodiments, **RA** is hydrogen and **RB** is a first **R1**. In some of these embodiments, **RA** is a first **R1** and **RB** is hydrogen. In some of these embodiments, **RA** is a first **R1** and **RB** is a second **R1**. In some of these embodiments, **RA** is **R2** and **RB** is hydrogen. In some of these embodiments, **RA** is a **R2** and **RB** is a first **R1**. In some of these variations, the first **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the first **R1** is unsubstituted or C3-C6 cycloalkenyl. In some of these variations, the first **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the

first **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the first **R1** is or C3-C6 cycloalkenyl substituted with one or more **R3**. In some of these variations, the second **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the second **R1** is unsubstituted or C3-C6 cycloalkenyl. In some of these variations, the second **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the second **R1** is or C3-C6 cycloalkenyl substituted with one or more **R3**. In some of these variations, at least one **R3** is halogen. In some of these variations, at least one **R3** is -CN. In some of these variations, at least one **R3** is -OH. In some of these variations, at least one **R3** is -O(Alkyl). In some of these variations, at least one **R3** is =O. In some of these variations, at least one **R3** is -NO₂. In some of these variations, at least one **R3** is -SH. In some of these variations, at least one **R3** is -S(Alkyl). In some of these variations, at least one **R3** is -S(O)(Alkyl). In some of these variations, at least one **R3** is -S(O)₂(Alkyl). In some of these variations, at least one **R3** is -CH₂OCH₃. In some of these variations, at least one **R3** is -OBn. In some of these variations, at least one **R3** is -CO₂H. In some of these variations, at least one **R3** is -CO₂(Alkyl). In some of these variations, at least one **R3** is -NR₁₀R₁₁. In some of these variations, at least one **R3** is -CONR₁₀R₁₁. In some of these variations, at least one **R3** is -S(O)₂NR₁₀R₁₁. In some of these variations, **R2** is halogen. In some of these variations, **R2** is -CN. In some of these variations, **R2** is -OH. In some of these variations, **R2** is -O(Alkyl). In some of these variations, **R2** is -NO₂. In some of these variations, **R2** is -SH. In some of these variations, **R2** is -S(Alkyl). In some of these variations, **R2** is -S(O)(Alkyl). In some of these variations, **R2** is -S(O)₂(Alkyl). In some of these variations, **R2** is -CH₂OCH₃. In some of these variations, **R2** is -OBn. In some of these variations, **R2** is -CO₂H. In some of these variations, **R2** is -CO₂(Alkyl). In some of these variations, **R2** is -NR₁₀R₁₁. In some of these variations, **R2** is -CONR₁₀R₁₁. In some of these variations, **R2** is -S(O)₂NR₁₀R₁₁.

[048] In some variations of Formulae (Ia) or (Ib), the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is a moiety selected from the group consisting of:



[049] In particular variations of Formulae (Ia) or (Ib), the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is a moiety selected from:



[050] In some variations of Formulae (Ia) or (Ib), one of R_{C4} , R_{C5} and R_{C6} is hydrogen. In some variations, two of R_{C4} , R_{C5} and R_{C6} are hydrogen. In some variations, each R_{C4} , R_{C5} and R_{C6} is hydrogen.

[051] In some variations of Formulae (Ia) or (Ib), one of R_{C4} , R_{C5} and R_{C6} is R_1 . In some of these variations, R_{C4} is R_1 . In some of these variations, R_{C5} is R_1 . In some of these variations, R_{C6} is R_1 . In some of these variations, R_1 is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, R_1 is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, R_1 is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, R_1 is unsubstituted C3-C6 cycloalkyl. In some of these variations, R_1 is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, R_1 is unsubstituted C3-C6 cycloalkenyl. In some variations, R_1 is unsubstituted aryl. In some variations, R_1 is unsubstituted heteroaryl. In some of these variations, R_1 is C1-C6 linear or branched alkyl substituted with one or more R_3 . In some of these variations, R_1 is C2-C6 linear or branched alkenyl substituted with one or more R_3 . In some of these variations, R_1 is C2-C6 linear or branched alkynyl substituted with one or more R_3 . In some of these variations, R_1 is C3-C6 cycloalkyl substituted with one or more R_3 .

In some of these variations, **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, **R1** is aryl substituted with one or more **R3**. In some variations **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, at least one **R3** is halogen. In some of these variations, at least one **R3** is -CN. In some of these variations, at least one **R3** is -OH. In some of these variations, at least one **R3** is -O(Alkyl). In some of these variations, at least one **R3** is =O. In some of these variations, at least one **R3** is -NO₂. In some of these variations, at least one **R3** is -SH. In some of these variations, at least one **R3** is -S(Alkyl). In some of these variations, at least one **R3** is -S(O)(Alkyl). In some of these variations, at least one **R3** is -S(O)₂(Alkyl). In some of these variations, at least one **R3** is -CH₂OCH₃. In some of these variations, at least one **R3** is -OBn. In some of these variations, at least one **R3** is -CO₂H. In some of these variations, at least one **R3** is -CO₂(Alkyl). In some of these variations, at least one **R3** is -NR₁₀R₁₁. In some of these variations, at least one **R3** is -CONR₁₀R₁₁. In some of these variations, at least one **R3** is -S(O)₂NR₁₀R₁₁.

[052] In some variations of Formulae (Ia) or (Ib), two of **R_{C4}**, **R_{C5}** and **R_{C6}** are **R1**. In some of these variations, **R_{C4}** is a first **R1** and **R_{C5}** is a second **R1**. In some of these variations, **R_{C4}** is a first **R1** and **R_{C6}** is a second **R1**. In some of these variations, **R_{C5}** is a first **R1** and **R_{C6}** is a second **R1**. In some of these variations, the first **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the first **R1** is unsubstituted aryl. In some variations, the first **R1** is unsubstituted heteroaryl. In some of these variations, the first **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the first **R1** is aryl substituted with one or more **R3**. In some variations, the first **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, the second **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the

second **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the second **R1** is unsubstituted aryl. In some variations, the second **R1** is unsubstituted heteroaryl. In some of these variations, the second **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the second **R1** is aryl substituted with one or more **R3**. In some variations, the second **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, at least one **R3** is halogen. In some of these variations, at least one **R3** is -CN. In some of these variations, at least one **R3** is -OH. In some of these variations, at least one **R3** is -O(Alkyl). In some of these variations, at least one **R3** is =O. In some of these variations, at least one **R3** is -NO₂. In some of these variations, at least one **R3** is -SH. In some of these variations, at least one **R3** is -S(Alkyl). In some of these variations, at least one **R3** is -S(O)(Alkyl). In some of these variations, at least one **R3** is -S(O)₂(Alkyl). In some of these variations, at least one **R3** is -CH₂OCH₃. In some of these variations, at least one **R3** is -OBn. In some of these variations, at least one **R3** is -CO₂H. In some of these variations, at least one **R3** is -CO₂(Alkyl). In some of these variations, at least one **R3** is -NR₁₀R₁₁. In some of these variations, at least one **R3** is -CONR₁₀R₁₁. In some of these variations, at least one **R3** is -S(O)₂NR₁₀R₁₁.

[053] In some variations of Formulae (Ia) or (Ib), each **R_{C4}**, **R_{C5}** and **R_{C6}** is **R1** (*i.e.*, **R_{C4}** is a first **R1**, **R_{C5}** is a second **R1**, and **R_{C6}** is a third **R1**). In some of these variations, the first **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the first **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the first **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the first **R1** is unsubstituted aryl. In some variations, the first **R1** is unsubstituted heteroaryl. In some of these variations, the first **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the first **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations,

the first **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the first **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the first **R1** is aryl substituted with one or more **R3**. In some variations, the first **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, the second **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the second **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the second **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the second **R1** is unsubstituted aryl. In some variations, the second **R1** is unsubstituted heteroaryl. In some of these variations, the second **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the second **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the second **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the second **R1** is aryl substituted with one or more **R3**. In some variations, the second **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, the third **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, the third **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, the third **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, the third **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the third **R1** is unsubstituted aryl. In some variations, the third **R1** is unsubstituted heteroaryl. In some of these variations, the third **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, the third **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, the third **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, the third **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the third **R1** is aryl substituted with one or more **R3**. In some variations, the third **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, at least one **R3** is

halogen. In some of these variations, at least one **R3** is $-\text{CN}$. In some of these variations, at least one **R3** is $-\text{OH}$. In some of these variations, at least one **R3** is $-\text{O}(\text{Alkyl})$. In some of these variations, at least one **R3** is $=\text{O}$. In some of these variations, at least one **R3** is $-\text{NO}_2$. In some of these variations, at least one **R3** is $-\text{SH}$. In some of these variations, at least one **R3** is $-\text{S}(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{S}(\text{O})(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{CH}_2\text{OCH}_3$. In some of these variations, at least one **R3** is $-\text{OBn}$. In some of these variations, at least one **R3** is $-\text{CO}_2\text{H}$. In some of these variations, at least one **R3** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{NR}_{10}\text{R}_{11}$. In some of these variations, at least one **R3** is $-\text{CONR}_{10}\text{R}_{11}$. In some of these variations, at least one **R3** is $-\text{S}(\text{O})_2\text{NR}_{10}\text{R}_{11}$.

[054] In some variations of Formulae (Ia) or (Ib), one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is C3-C6 cycloalkyl substituted with one or more **R3**. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some of these variations, at least one **R3** is halogen. In some of these variations, at least one **R3** is $-\text{CN}$. In some of these variations, at least one **R3** is $-\text{OH}$. In some of these variations, at least one **R3** is $-\text{O}(\text{Alkyl})$. In some of these variations, at least one **R3** is $=\text{O}$. In some of these variations, at least one **R3** is $-\text{NO}_2$. In some of these variations, at least one **R3** is $-\text{SH}$. In some of these variations, at least one **R3** is $-\text{S}(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{S}(\text{O})(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{CH}_2\text{OCH}_3$. In some of these variations, at least one **R3** is $-\text{OBn}$. In some of these variations, at least one **R3** is $-\text{CO}_2\text{H}$. In some of these variations, at least one **R3** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{NR}_{10}\text{R}_{11}$. In some of these variations, at least one **R3** is $-\text{CONR}_{10}\text{R}_{11}$.

[055] In some variations of Formulae (Ia) or (Ib), one of **R_{C4}**, **R_{C5}** and **R_{C6}** is **R2**. In some of these variations, **R2** is halogen. In some of these variations, **R2** is $-\text{CN}$. In some of these variations, **R2** is $-\text{OH}$. In some of these variations, **R2** is $-\text{O}(\text{Alkyl})$. In some of these variations, **R2** is $-\text{NO}_2$. In some of these variations, **R2** is $-\text{SH}$. In some of these variations, **R2** is $-\text{S}(\text{Alkyl})$. In some of these variations, **R2** is $-\text{S}(\text{O})(\text{Alkyl})$. In some of these variations, **R2** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some of these variations, **R2** is $-\text{CH}_2\text{OCH}_3$. In some of these variations, **R2** is $-\text{OBn}$. In some

of these variations, **R2** is $-\text{CO}_2\text{H}$. In some of these variations, **R2** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, **R2** is $-\text{NR}_{10}\text{R}_{11}$. In some of these variations, **R2** is $-\text{CONR}_{10}\text{R}_{11}$. In some of these variations, **R2** is $-\text{S}(\text{O})_2\text{NR}_{10}\text{R}_{11}$.

[056] In some variations of Formulae (Ia) or (Ib), two of **R_{C4}**, **R_{C5}** and **R_{C6}** are **R2**. In some variations, **R_{C4}** is a first **R2** and **R_{C5}** is a second **R2**. In some variations, **R_{C4}** is a first **R2** and **R_{C6}** is a second **R2**. In some variations, **R_{C5}** is a first **R2** and **R_{C6}** is a second **R2**. In some of these variations, the first **R2** is halogen. In some of these variations, the first **R2** is $-\text{CN}$. In some of these variations, the first **R2** is $-\text{OH}$. In some of these variations, the first **R2** is $-\text{O}(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{NO}_2$. In some of these variations, the first **R2** is $-\text{SH}$. In some of these variations, the first **R2** is $-\text{S}(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{S}(\text{O})(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{CH}_2\text{OCH}_3$. In some of these variations, the first **R2** is $-\text{OBn}$. In some of these variations, the first **R2** is $-\text{CO}_2\text{H}$. In some of these variations, the first **R2** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{NR}_{10}\text{R}_{11}$. In some of these variations, the first **R2** is $-\text{CONR}_{10}\text{R}_{11}$. In some of these variations, the first **R2** is $-\text{S}(\text{O})_2\text{NR}_{10}\text{R}_{11}$. In some of these variations, the second **R2** is halogen. In some of these variations, the second **R2** is $-\text{CN}$. In some of these variations, the second **R2** is $-\text{OH}$. In some of these variations, the second **R2** is $-\text{O}(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{NO}_2$. In some of these variations, the second **R2** is $-\text{SH}$. In some of these variations, the second **R2** is $-\text{S}(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{S}(\text{O})(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{CH}_2\text{OCH}_3$. In some of these variations, the second **R2** is $-\text{OBn}$. In some of these variations, the second **R2** is $-\text{CO}_2\text{H}$. In some of these variations, the second **R2** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{NR}_{10}\text{R}_{11}$. In some of these variations, the second **R2** is $-\text{CONR}_{10}\text{R}_{11}$.

[057] In some variations of Formulae (Ia) or (Ib), each **R_{C4}**, **R_{C5}** and **R_{C6}** is **R2** (*i.e.*, there is a first **R2**, a second **R2**, and a third **R2**). In some of these variations, the first **R2** is halogen. In some of these variations, the first **R2** is $-\text{CN}$. In some of these variations, the first **R2** is $-\text{OH}$. In some of these variations, the first **R2** is $-\text{O}(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{NO}_2$. In some of these variations, the first **R2** is $-\text{SH}$. In some of these variations, the first **R2** is $-\text{S}(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{S}(\text{O})(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{CH}_2\text{OCH}_3$. In some of these variations, the first **R2** is $-\text{OBn}$. In some of these variations, the first **R2** is $-\text{CO}_2\text{H}$. In some of these variations, the first **R2** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, the first **R2** is $-\text{NR}_{10}\text{R}_{11}$. In some of these variations, the first **R2** is $-\text{CONR}_{10}\text{R}_{11}$. In some of these


variations, the first **R2** is $-\text{S}(\text{O})_2\text{NR10R11}$. In some of these variations, the second **R2** is halogen. In some of these variations, the second **R2** is $-\text{CN}$. In some of these variations, the second **R2** is $-\text{OH}$. In some of these variations, the second **R2** is $-\text{O}(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{NO}_2$. In some of these variations, the second **R2** is $-\text{SH}$. In some of these variations, the second **R2** is $-\text{S}(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{S}(\text{O})(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{CH}_2\text{OCH}_3$. In some of these variations, the second **R2** is $-\text{OBn}$. In some of these variations, the second **R2** is $-\text{CO}_2\text{H}$. In some of these variations, the second **R2** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, the second **R2** is $-\text{NR10R11}$. In some of these variations, the second **R2** is $-\text{CONR10R11}$. In some of these variations, the second **R2** is $-\text{S}(\text{O})_2\text{NR10R11}$. In some of these variations, the third **R2** is halogen. In some of these variations, the third **R2** is $-\text{CN}$. In some of these variations, the third **R2** is $-\text{OH}$. In some of these variations, the third **R2** is $-\text{O}(\text{Alkyl})$. In some of these variations, the third **R2** is $-\text{NO}_2$. In some of these variations, the third **R2** is $-\text{SH}$. In some of these variations, the third **R2** is $-\text{S}(\text{Alkyl})$. In some of these variations, the third **R2** is $-\text{S}(\text{O})(\text{Alkyl})$. In some of these variations, the third **R2** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some of these variations, the third **R2** is $-\text{CH}_2\text{OCH}_3$. In some of these variations, the third **R2** is $-\text{OBn}$. In some of these variations, the third **R2** is $-\text{CO}_2\text{H}$. In some of these variations, the third **R2** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, the third **R2** is $-\text{NR10R11}$. In some of these variations, the third **R2** is $-\text{CONR10R11}$. In some of these variations, the third **R2** is $-\text{S}(\text{O})_2\text{NR10R11}$.

[058] In some variations of Formulae (Ia) or (Ib), one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is halogen. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{CN}$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{OH}$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{O}(\text{Alkyl})$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{NO}_2$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{SH}$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{S}(\text{Alkyl})$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{S}(\text{O})(\text{Alkyl})$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{S}(\text{O})_2(\text{Alkyl})$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{CH}_2\text{OCH}_3$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{OBn}$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{CO}_2\text{H}$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{CO}_2(\text{Alkyl})$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{NR10R11}$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{CONR10R11}$. In some variations, one or more of **R_{C4}**, **R_{C5}** and **R_{C6}** is $-\text{S}(\text{O})_2\text{NR10R11}$.

[059] In some variations of Formulae (Ia) or (Ib), one of **R_{C4}**, **R_{C5}** and **R_{C6}** is halogen. In some variations, two of **R_{C4}**, **R_{C5}** and **R_{C6}** are halogen. In some variations each **R_{C4}**, **R_{C5}** and **R_{C6}** is

halogen. In some variations, one of R_{C4} , R_{C5} and R_{C6} is hydrogen, and the remaining two of R_{C4} , R_{C5} and R_{C6} are halogen. In some variations, two of R_{C4} , R_{C5} and R_{C6} are hydrogen, and the remaining one of R_{C4} , R_{C5} and R_{C6} is halogen. In some variations, R_{C4} , R_{C5} and R_{C6} are each halogen. In some embodiments, one of R_{C4} , R_{C5} and R_{C6} is fluoro, chloro or bromo. In some embodiments, two of R_{C4} , R_{C5} and R_{C6} are fluoro, chloro or bromo. In some embodiments, one of R_{C4} , R_{C5} and R_{C6} is chloro. In some embodiments, one of R_{C4} , R_{C5} and R_{C6} is chloro and one of R_{C4} , R_{C5} and R_{C6} is bromo. In some embodiments, R_{C4} is chloro. In some embodiments, R_{C5} is chloro. In some embodiments, R_{C4} is bromo. In some embodiments, R_{C5} is bromo.

[060] In some variations of Formulae (Ia) or (Ib), R_{C4} is taken together with R_{C7} to form a heterocyclyl optionally substituted with one or more R_3 . In some variations, R_{C4} is taken together with R_{C7} to form a moiety selected from:



[061] In some variations of Formulae (Ia) or (Ib), R_{C7} is hydrogen. In some variations, R_{C7} is R_1 . In some of these variations, R_1 is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, R_1 is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, R_1 is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, R_1 is unsubstituted C3-C6 cycloalkyl. In some of these variations, R_1 is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, R_1 is unsubstituted C3-C6 cycloalkenyl. In some variations, R_1 is unsubstituted aryl. In some variations, R_1 is unsubstituted heteroaryl. In some of these variations, R_1 is C1-C6 linear or branched alkyl substituted with one or more R_3 . In some of these variations, R_1 is C2-C6 linear or branched alkenyl substituted with one or more R_3 . In some of these variations, R_1 is C2-C6 linear or branched alkynyl substituted with one or more R_3 . In some of these variations, R_1 is C3-C6 cycloalkyl substituted with one or more R_3 . In some of these variations, R_1 is C3-C6 cycloalkylmethyl substituted with one or more R_3 . In some of these variations, R_1 is C3-C6 cycloalkenyl substituted with one or more R_3 . In some variations, R_1 is aryl substituted with one or more R_3 . In some variations, R_1 is heteroaryl substituted with one or more R_3 . In some of these variations, at least one R_3 is halogen. In some of these variations, at least one R_3 is $-CN$. In some of these variations, at least one R_3 is $-OH$. In some of these variations, at least one R_3 is $-O(Alkyl)$. In some of these variations, at least one R_3 is $=O$. In some of these variations, at least one R_3 is $-NO_2$. In some of these variations, at least one R_3 is $-SH$. In some of these variations, at least one R_3 is $-S(Alkyl)$. In some of these variations, at least one R_3 is $-S(O)(Alkyl)$. In some of these variations, at least one R_3 is

-S(O)₂(Alkyl). In some of these variations, at least one **R3** is -CH₂OCH₃. In some of these variations, at least one **R3** is -OBn. In some of these variations, at least one **R3** is -CO₂H. In some of these variations, at least one **R3** is -CO₂(Alkyl). In some of these variations, at least one **R3** is -NR₁₀R₁₁. In some of these variations, at least one **R3** is -CONR₁₀R₁₁. In some of these variations, at least one **R3** is -S(O)₂NR₁₀R₁₁.

[062] In some variations of Formulae (Ia) or (Ib), **R_{C7}** is -SO₂**R_{C12}**. In some of these variations, **R_{C12}** is unsubstituted linear or branched C1-C6 alkyl. In some of these variations, **R_{C12}** is unsubstituted linear or branched C2-C6 alkenyl. In some of these variations, **R_{C12}** is unsubstituted linear or branched C2-C6 alkynyl. In some of these variations, **R_{C12}** is aryl. In some of these variations, **R_{C12}** is linear or branched C1-C6 alkyl substituted with one or more halogen atoms. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkenyl substituted with one or more halogen atoms. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkynyl substituted with one or more halogen atoms. In some of these variations, **R_{C12}** is aryl substituted with one or more halogen atoms. In some of these variations, **R_{C12}** is linear or branched C1-C6 alkyl substituted with one or more one or more -NH₂ groups. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkenyl substituted with one or more one or more -NH₂ groups. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkynyl substituted with one or more one or more -NH₂ groups. In some of these variations, **R_{C12}** is aryl substituted with one or more one or more -NH₂ groups. In some of these variations, **R_{C12}** is linear or branched C1-C6 alkyl substituted with one or more one or more -OH groups. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkenyl substituted with one or more one or more -OH groups. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkynyl substituted with one or more one or more -OH groups. In some of these variations, **R_{C12}** is aryl substituted with one or more one or more -OH groups.

[063] In some variations of Formulae (Ia) or (Ib), **R_{C8}** is hydrogen. In some variations, **R_{C8}** is **R1**. In some of these variations, **R1** is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, **R1** is unsubstituted C3-C6 cycloalkyl. In some of these variations, **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, **R1** is unsubstituted aryl. In some variations, **R1** is unsubstituted heteroaryl. In some of these variations, **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some of these variations, **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some of these variations, **R1** is C2-C6 linear or branched alkynyl substituted with one or

more **R3**. In some of these variations, **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some of these variations, **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some of these variations, **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, **R1** is aryl substituted with one or more **R3**. In some variations, **R1** is heteroaryl substituted with one or more **R3**. In some of these variations, at least one **R3** is halogen. In some of these variations, at least one **R3** is -CN. In some of these variations, at least one **R3** is -OH. In some of these variations, at least one **R3** is -O(Alkyl). In some of these variations, at least one **R3** is =O. In some of these variations, at least one **R3** is -NO₂. In some of these variations, at least one **R3** is -SH. In some of these variations, at least one **R3** is -S(Alkyl). In some of these variations, at least one **R3** is -S(O)(Alkyl). In some of these variations, at least one **R3** is -S(O)₂(Alkyl). In some of these variations, at least one **R3** is -CH₂OCH₃. In some of these variations, at least one **R3** is -OBn. In some of these variations, at least one **R3** is -CO₂H. In some of these variations, at least one **R3** is -CO₂(Alkyl). In some of these variations, at least one **R3** is -NR₁₀R₁₁. In some of these variations, at least one **R3** is -CONR₁₀R₁₁. In some of these variations, at least one **R3** is -S(O)₂NR₁₀R₁₁.

[064] In some variations of Formulae (Ia) or (Ib), **R_{C8}** is -SO₂**R_{C12}**. In some of these variations, **R_{C12}** is unsubstituted linear or branched C1-C6 alkyl. In some of these variations, **R_{C12}** is unsubstituted linear or branched C2-C6 alkenyl. In some of these variations, **R_{C12}** is unsubstituted linear or branched C2-C6 alkynyl. In some of these variations, **R_{C12}** is aryl. In some of these variations, **R_{C12}** is linear or branched C1-C6 alkyl substituted with one or more halogen atoms. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkenyl substituted with one or more halogen atoms. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkynyl substituted with one or more halogen atoms. In some of these variations, **R_{C12}** is aryl substituted with one or more halogen atoms. In some of these variations, **R_{C12}** is linear or branched C1-C6 alkyl substituted with one or more one or more -NH₂ groups. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkenyl substituted with one or more one or more -NH₂ groups. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkynyl substituted with one or more one or more -NH₂ groups. In some of these variations, **R_{C12}** is aryl substituted with one or more one or more -NH₂ groups. In some of these variations, **R_{C12}** is linear or branched C1-C6 alkyl substituted with one or more one or more -OH groups. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkenyl substituted with one or more one or more -OH groups. In some of these variations, **R_{C12}** is linear or branched C2-C6 alkynyl substituted with one or more one or more -OH groups. In some of these variations, **R_{C12}** is aryl substituted with one or more one or more -OH groups.

[065] In some variations of Formulae (Ia) or (Ib), R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more $R1$. In some of these variations, $R1$ is unsubstituted C1-C6 linear or branched alkyl. In some of these variations, $R1$ is unsubstituted C2-C6 linear or branched alkenyl. In some of these variations, $R1$ is unsubstituted C2-C6 linear or branched alkynyl. In some of these variations, $R1$ is unsubstituted C3-C6 cycloalkyl. In some of these variations, $R1$ is unsubstituted C3-C6 cycloalkylmethyl. In some of these variations, $R1$ is unsubstituted C3-C6 cycloalkenyl. In some variations, $R1$ is unsubstituted aryl. In some variations, $R1$ is unsubstituted heteroaryl. In some of these variations, $R1$ is C1-C6 linear or branched alkyl substituted with one or more $R3$. In some of these variations, $R1$ is C2-C6 linear or branched alkenyl substituted with one or more $R3$. In some of these variations, $R1$ is C2-C6 linear or branched alkynyl substituted with one or more $R3$. In some of these variations, $R1$ is C3-C6 cycloalkyl substituted with one or more $R3$. In some of these variations, $R1$ is C3-C6 cycloalkylmethyl substituted with one or more $R3$. In some of these variations, $R1$ is C3-C6 cycloalkenyl substituted with one or more $R3$. In some variations, $R1$ is aryl substituted with one or more $R3$. In some variations, $R1$ is heteroaryl substituted with one or more $R3$. In some of these variations, at least one $R3$ is halogen. In some of these variations, at least one $R3$ is $-CN$. In some of these variations, at least one $R3$ is $-OH$. In some of these variations, at least one $R3$ is $-O(Alkyl)$. In some of these variations, at least one $R3$ is $=O$. In some of these variations, at least one $R3$ is $-NO_2$. In some of these variations, at least one $R3$ is $-SH$. In some of these variations, at least one $R3$ is $-S(Alkyl)$. In some of these variations, at least one $R3$ is $-S(O)(Alkyl)$. In some of these variations, at least one $R3$ is $-S(O)_2(Alkyl)$. In some of these variations, at least one $R3$ is $-CH_2OCH_3$. In some of these variations, at least one $R3$ is $-OBn$. In some of these variations, at least one $R3$ is $-CO_2H$. In some of these variations, at least one $R3$ is $-CO_2(Alkyl)$. In some of these variations, at least one $R3$ is $-NR10R11$. In some of these variations, at least one $R3$ is $-CONR10R11$. In some of these variations, at least one $R3$ is $-S(O)_2NR10R11$.

[066] In some variations of Formulae (Ia) or (Ib), R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more $R3$. In some of these variations, at least one $R3$ is halogen. In some of these variations, at least one $R3$ is $-CN$. In some of these variations, at least one $R3$ is $-OH$. In some of these variations, at least one $R3$ is $-O(Alkyl)$. In some of these variations, at least one $R3$ is $=O$. In some of these variations, at least one $R3$ is $-NO_2$. In some of these variations, at least one $R3$ is $-SH$. In some of these variations, at least one $R3$ is $-S(Alkyl)$. In some of these variations, at least one $R3$ is $-S(O)(Alkyl)$. In some of these variations, at least one $R3$ is $-S(O)_2(Alkyl)$. In some of these variations, at least one $R3$ is $-CH_2OCH_3$. In some of these variations, at least one $R3$ is $-OBn$. In


some of these variations, at least one **R3** is $-\text{CO}_2\text{H}$. In some of these variations, at least one **R3** is $-\text{CO}_2(\text{Alkyl})$. In some of these variations, at least one **R3** is $-\text{NR}_{10}\text{R}_{11}$. In some of these variations, at least one **R3** is $-\text{CONR}_{10}\text{R}_{11}$. In some of these variations, at least one **R3** is $-\text{S}(\text{O})_2\text{NR}_{10}\text{R}_{11}$.

[067] In some variations of Formulae (Ia) or (Ib), one or both of **R10** and **R11** are hydrogen. In some variations, one or both of **R10** and **R11** are C1-C6 linear or branched alkyl. In some variations, one or both of **R10** and **R11** are $-\text{C}(\text{O})\text{R}_{\text{C12}}$. In some variations, one or both of **R10** and **R11** are $-\text{C}(\text{O})_2\text{R}_{\text{C12}}$. In some variations, one or both of **R10** and **R11** are $-\text{C}(\text{O})\text{N}(\text{R}_{\text{C12}})_2$. In some variations, one or both of **R10** and **R11** are $-\text{SO}_2\text{R}_{\text{C12}}$. In some variations, **R10** and **R11** are taken together with the N to which they are attached to form a C3-C8 heterocycle.


[068] In some variations of Formulae (Ia) or (Ib), **R_{C12}** is hydrogen. In some variations, **R_{C12}** is a linear or branched C1-C6 alkyl. In some variations, **R_{C12}** is a linear or branched C2-C6 alkenyl. In some variations, **R_{C12}** is a linear or branched C2-C6 alkynyl. In some variations, **R_{C12}** is aryl.

[069] In some variations of Formulae (Ia) or (Ib), **R_{C12}** is a linear or branched C1-C6 alkyl having at least two halogen atoms. In some variations, **R_{C12}** is a linear or branched C2-C6 alkenyl having at least two halogen atoms. In some variations, **R_{C12}** is a linear or branched C2-C6 alkynyl having at least two halogen atoms.


[070] In some variations of Formulae (Ia) or (Ib), **R_{C12}** is selected from:




[071] In some variations, **R_{C12}** is selected from:




[072] In some variations of Formulae (Ia) or (Ib), A is a moiety selected from




the 5-membered heteroaromatic group comprising **W**, **X** and **Y** is selected from:




and the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is a moiety selected from:




[073] In some variations of Formulae (Ia) or (Ib), **A** is a moiety selected from




the 5-membered heteroaromatic group comprising **W**, **X** and **Y** is selected from:




and the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is a moiety selected from:




[074] In some variations of Formulae (Ia) or (Ib), A is




the 5-membered heteroaromatic group comprising W, X and Y is selected from:




and the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is:




[075] In some variations, the compound is of Formulae (Ia), wherein A is a moiety selected from




the 5-membered heteroaromatic group comprising W, X and Y is selected from:




and the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is a moiety selected from:




[076] In some variations, the compound is of Formulae (Ia), wherein A is



the 5-membered heteroaromatic group comprising W, X and Y is selected from:




and the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is:



. In some of these variations, R_{C4} , R_{C5} , R_{C6} , are each independently H or halogen. In some of these variations, R_{C4} and R_{C6} are each H, and R_{C5} is halogen.


[077] In some variations, the compound of Formula (Ia) is a compound of Formula (IIa):



wherein R_{C5} is halogen, and each R_{C7} and R_{C8} is independently hydrogen, R_1 , or $-SO_2R_{C12}$; or R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more R_1 or R_3 . In some variations, R_{C5} is chloro. In some variations, R_{C7} and R_{C8} are taken together with the N to which they are attached to form

a pyrrolidinyl, piperidinyl, piperazinyl, or morpholinyl, each optionally substituted with one or more **R1** or **R3**.

[078] In some variations, the compound of Formula (Ib) is a compound of Formula (IIb):



wherein **R_{C5}** is halogen, and each **R_{C7}** and **R_{C8}** is independently hydrogen, **R1**, or -SO₂**R_{C12}**; or **R_{C7}** and **R_{C8}** are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more **R1** or **R3**. In some variations, **R_{C5}** is chloro. In some variations, **R_{C7}** and **R_{C8}** are taken together with the N to which they are attached to form a pyrrolidinyl, piperidinyl, piperazinyl, or morpholinyl, each optionally substituted with one or more **R1** or **R3**.

Variations of Formula (IIa) or (IIb) in which **R_{C7}** and **R_{C8}** are taken together with the N to which they are attached to form an optionally substituted C3-C8 heterocycle

[079] In some variations of Formula (IIa) or (IIb), **R_{C7}** and **R_{C8}** are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more **R1**. In some of these variations, the C3-C8 heterocycle is unsubstituted. In some of these variations, the C3-C8 heterocycle is substituted with a first **R1**. In some of these variations, the C3-C8 heterocycle is substituted with a first **R1** and a second **R1**. In some of these variations, the C3-C8 heterocycle is substituted with a first **R1**, a second **R1**, and a third **R1**. In some of these variations, the C3-C8 heterocycle is substituted with a first **R31**. In some of these variations, the C3-C8 heterocycle is substituted with a first **R3** and a second **R3**. In some of these variations, the C3-C8 heterocycle is substituted with a first **R3**, a second **R3**, and a third **R3**. In some of these variations, the C3-C8 heterocycle is substituted with a first **R1** and a first **R3**. In some of these variations, the C3-C8 heterocycle is substituted with a first **R1**, a second **R1**, and a first **R3**. In some of these variations, the C3-C8 heterocycle is substituted with a first **R1**, a first **R3**, and a second **R3**.

[080] In some variations, the first **R1** is unsubstituted C1-C6 linear or branched alkyl. In some variations, the first **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some variations, the first **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some variations, the first **R1** is unsubstituted C3-C6 cycloalkyl. In some variations, the first **R1** is unsubstituted C3-C6

cycloalkylmethyl. In some variations, the first **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the first **R1** is unsubstituted aryl. In some variations, the first **R1** is unsubstituted heteroaryl. In some variations, the first **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some variations, the first **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some variations, the first **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some variations, the first **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some variations, the first **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some variations, the first **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the first **R1** is aryl substituted with one or more **R3**. In some variations, the first **R1** is heteroaryl substituted with one or more **R3**. In some variations, the first **R1** is substituted with at least one halogen. In some variations, the first **R1** is substituted with at least one -CN. In some variations, the first **R1** is substituted with at least one -OH. In some variations, the first **R1** is substituted with at least one -O(Alkyl). In some variations, the first **R1** is substituted with at least one =O. In some variations, the first **R1** is substituted with at least one -NO₂. In some variations, the first **R1** is substituted with at least one -SH. In some variations, the first **R1** is substituted with at least one -S(Alkyl). In some variations, the first **R1** is substituted with at least one -S(O)(Alkyl). In some variations, the first **R1** is substituted with at least one -S(O)₂(Alkyl). In some variations, the first **R1** is substituted with at least one -CH₂OCH₃. In some variations, the first **R1** is substituted with at least one -OBn. In some variations, the first **R1** is substituted with at least one -CO₂H. In some variations, the first **R1** is substituted with at least one -CO₂(Alkyl). In some variations, the first **R1** is substituted with at least one -NR₁₀R₁₁. In some variations, the first **R1** is substituted with at least one -CONR₁₀R₁₁. In some variations, the first **R1** is substituted with at least one -S(O)₂NR₁₀R₁₁.

[081] In some variations, the second **R1** is unsubstituted C1-C6 linear or branched alkyl. In some variations, the second **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some variations, the second **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some variations, the second **R1** is unsubstituted C3-C6 cycloalkyl. In some variations, the second **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some variations, the second **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the second **R1** is unsubstituted aryl. In some variations, the second **R1** is unsubstituted heteroaryl. In some variations, the second **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some variations, the second **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some variations, the second **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some variations, the second **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some variations, the second **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some variations, the second

R1 is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the second **R1** is aryl substituted with one or more **R3**. In some variations, the second **R1** is heteroaryl substituted with one or more **R3**. In some variations, the second **R1** is substituted with at least one halogen. In some variations, the second **R1** is substituted with at least one -CN. In some variations, the second **R1** is substituted with at least one -OH. In some variations, the second **R1** is substituted with at least one -O(Alkyl). In some variations, the second **R1** is substituted with at least one =O. In some variations, the second **R1** is substituted with at least one -NO₂. In some variations, the second **R1** is substituted with at least one -SH. In some variations, the second **R1** is substituted with at least one -S(Alkyl). In some variations, the second **R1** is substituted with at least one -S(O)(Alkyl). In some variations, the second **R1** is substituted with at least one -S(O)₂(Alkyl). In some variations, the second **R1** is substituted with at least one -CH₂OCH₃. In some variations, the second **R1** is substituted with at least one -OBn. In some variations, the second **R1** is substituted with at least one -CO₂H. In some variations, the second **R1** is substituted with at least one -CO₂(Alkyl). In some variations, the second **R1** is substituted with at least one -NR₁₀R₁₁. In some variations, the second **R1** is substituted with at least one -CONR₁₀R₁₁. In some variations, the second **R1** is substituted with at least one -S(O)₂NR₁₀R₁₁.

[082] In some variations, the third **R1** is unsubstituted C1-C6 linear or branched alkyl. In some variations, the third **R1** is unsubstituted C2-C6 linear or branched alkenyl. In some variations, the third **R1** is unsubstituted C2-C6 linear or branched alkynyl. In some variations, the third **R1** is unsubstituted C3-C6 cycloalkyl. In some variations, the third **R1** is unsubstituted C3-C6 cycloalkylmethyl. In some variations, the third **R1** is unsubstituted C3-C6 cycloalkenyl. In some variations, the third **R1** is unsubstituted aryl. In some variations, the third **R1** is unsubstituted heteroaryl. In some variations, the third **R1** is C1-C6 linear or branched alkyl substituted with one or more **R3**. In some variations, the third **R1** is C2-C6 linear or branched alkenyl substituted with one or more **R3**. In some variations, the third **R1** is C2-C6 linear or branched alkynyl substituted with one or more **R3**. In some variations, the third **R1** is C3-C6 cycloalkyl substituted with one or more **R3**. In some variations, the third **R1** is C3-C6 cycloalkylmethyl substituted with one or more **R3**. In some variations, the third **R1** is C3-C6 cycloalkenyl substituted with one or more **R3**. In some variations, the third **R1** is aryl substituted with one or more **R3**. In some variations, the third **R1** is heteroaryl substituted with one or more **R3**. In some variations, the third **R1** is substituted with at least one halogen. In some variations, the third **R1** is substituted with at least one -CN. In some variations, the third **R1** is substituted with at least one -OH. In some variations, the third **R1** is substituted with at least one -O(Alkyl). In some variations, the third **R1** is substituted with at least one =O. In some variations, the third **R1** is substituted with at least one -NO₂. In some variations, the third **R1** is substituted with at least one

–SH. In some variations, the third **R1** is substituted with at least one –S(Alkyl). In some variations, the third **R1** is substituted with at least one –S(O)(Alkyl). In some variations, the third **R1** is substituted with at least one –S(O)₂(Alkyl). In some variations, the third **R1** is substituted with at least one –CH₂OCH₃. In some variations, the third **R1** is substituted with at least one –OBn. In some variations, the third **R1** is substituted with at least one –CO₂H. In some variations, the third **R1** is substituted with at least one –CO₂(Alkyl). In some variations, the third **R1** is substituted with at least one –NR₁₀R₁₁. In some variations, the third **R1** is substituted with at least one –CONR₁₀R₁₁. In some variations, the third **R1** is substituted with at least one –S(O)₂NR₁₀R₁₁.

[083] In some of these variations, the first **R3** is halogen. In some of these variations, the first **R3** is –CN. In some of these variations, the first **R3** is –OH. In some of these variations, the first **R3** is –O(Alkyl). In some of these variations, the first **R3** is =O. In some of these variations, the first **R3** is –NO₂. In some of these variations, the first **R3** is –SH. In some of these variations, the first **R3** is –S(Alkyl). In some of these variations, the first **R3** is –S(O)(Alkyl). In some of these variations, the first **R3** is –S(O)₂(Alkyl). In some of these variations, the first **R3** is –CH₂OCH₃. In some of these variations, the first **R3** is –OBn. In some of these variations, the first **R3** is –CO₂H. In some of these variations, the first **R3** is –CO₂(Alkyl). In some of these variations, the first **R3** is –NR₁₀R₁₁. In some of these variations, the first **R3** is –CONR₁₀R₁₁. In some of these variations, the first **R3** is –S(O)₂NR₁₀R₁₁.

[084] In some of these variations, the second **R3** is halogen. In some of these variations, the second **R3** is –CN. In some of these variations, the second **R3** is –OH. In some of these variations, the second **R3** is –O(Alkyl). In some of these variations, the second **R3** is =O. In some of these variations, the second **R3** is –NO₂. In some of these variations, the second **R3** is –SH. In some of these variations, the second **R3** is –S(Alkyl). In some of these variations, the second **R3** is –S(O)(Alkyl). In some of these variations, the second **R3** is –S(O)₂(Alkyl). In some of these variations, the second **R3** is –CH₂OCH₃. In some of these variations, the second **R3** is –OBn. In some of these variations, the second **R3** is –CO₂H. In some of these variations, the second **R3** is –CO₂(Alkyl). In some of these variations, the second **R3** is –NR₁₀R₁₁. In some of these variations, the second **R3** is –CONR₁₀R₁₁. In some of these variations, the second **R3** is –S(O)₂NR₁₀R₁₁.

[085] In some of these variations, the third **R3** is halogen. In some of these variations, the third **R3** is –CN. In some of these variations, the third **R3** is –OH. In some of these variations, the third **R3** is –O(Alkyl). In some of these variations, the third **R3** is =O. In some of these variations, the third **R3** is –NO₂. In some of these variations, the third **R3** is –SH. In some of

these variations, the third **R3** is -S(Alkyl). In some of these variations, the third **R3** is -S(O)(Alkyl). In some of these variations, the third **R3** is -S(O)₂(Alkyl). In some of these variations, the third **R3** is -CH₂OCH₃. In some of these variations, the third **R3** is -OBn. In some of these variations, the third **R3** is -CO₂H. In some of these variations, the third **R3** is -CO₂(Alkyl). In some of these variations, the third **R3** is -NR₁₀R₁₁. In some of these variations, the third **R3** is -CONR₁₀R₁₁. In some of these variations, the third **R3** is -S(O)₂NR₁₀R₁₁.

[086] It is intended and understood that each and every variation of **A**, **R1**, **R2**, **R3**, **R_{C4}**, **R_{C5}**, **R_{C6}**, **R_{C7}**, **R_{C8}**, **R₁₀**, **R₁₁**, **R_{C12}**, **W**, **X**, **Y**, **R_A**, and **R_B**, where present, described for formulae (Ia) or (Ib) may be combined with each and every variation of **A**, **R1**, **R2**, **R3**, **R_{C4}**, **R_{C5}**, **R_{C6}**, **R_{C7}**, **R_{C8}**, **R₁₀**, **R₁₁**, **R_{C12}**, **W**, **X**, **Y**, **R_A**, and **R_B**, as if each and every combination is individually described.

[087] In some embodiments, the compound is selected from the group consisting of:

N-methyl-4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide;

N,N-dimethyl-4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide;

4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide;

4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1,3-thiazol-2-yl}-1H-indole;

2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1,3-thiazol-2-yl}pyridine;

3-chloro-4-[2-(1H-indol-4-yl)-1,3-thiazol-4-yl]-N,N-dimethylbenzamide;

4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-2-one;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-cyclobutylbenzamide;

3-chloro-N,N-dimethyl-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzamide;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-4-methylpiperazine;

2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]thiophen-2-yl}pyridine;

2-tert-butyl-4-{4-[2-chloro-4-(3,3-difluoropyrrolidine-1-carbonyl)phenyl]thiophen-2-yl}pyridine;

2-tert-butyl-4-{4-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]thiophen-2-yl}pyridine;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-cyclopropylbenzamide;

N-tert-butyl-4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamide;

2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-5-methylthiophen-2-yl}pyridine;

4-(3-chloro-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzoyl)morpholine;

3-chloro-N-(2-hydroxypropyl)-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzamide;

tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)carbamate;

tert-butyl N-[(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

tert-butyl N-[(3S)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-1 λ ⁶,4-thiomorpholine-1,1-dione;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-amine;

(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-amine;

(3S)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-amine;

4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}morpholine;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-phenylbenzamide;

2-tert-butyl-4-{4-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]-1H-imidazol-2-yl}pyridine;

2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1H-imidazol-2-yl}pyridine;

6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-2-methyl-1,2,3,4-tetrahydroisoquinolin-1-one;

tert-butyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazine-1-carboxylate;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazine;

1-(4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-1-yl)ethan-1-one;

- 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-4-methanesulfonylpiperazine;
- N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)methanesulfonamide;
- N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)acetamide;
- 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N,N-bis(propan-2-yl)benzamide;
- 1-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)pyrrolidin-2-one;
- 6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-2-(propan-2-yl)-1,2,3,4-tetrahydroisoquinolin-1-one;
- 2-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-1 λ ⁶,2-thiazolidine-1,1-dione;
- tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)carbamate;
- N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)acetamide;
- 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-1-ethylpiperazin-2-one;
- 2-tert-butyl-4-{5-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]thiophen-3-yl}pyridine;
- 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-methyl-N-phenylbenzamide;
- 4-{4-[4-(azetidine-1-carbonyl)-2-chlorophenyl]thiophen-2-yl}-2-tert-butylpyridine;
- 6-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2-oxa-6-azaspiro[3.3]heptane;
- ethyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)carbamate;
- ethyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazine-1-carboxylate;

1-(4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-1-yl)-2,2-dimethylpropan-1-one;

N-[2-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorophenyl}-N-ethylformamido)ethyl]acetamide;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-ol;

(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-ol;

(3S)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-ol;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(3-hydroxyphenyl)benzamide;

(2S,6R)-4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2,6-dimethylmorpholine;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-1,2,3,4-tetrahydro-1,8-naphthyridine;

6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-7-chloro-2-(propan-2-yl)-1,2,3,4-tetrahydroisoquinolin-1-one;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-[(4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl]benzamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(4-fluorophenyl)benzamide

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-2,2-dimethylpropanamide;

1-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-3-cyclopropylurea;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-ol;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(pyridin-3-yl)benzamide;

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)methanesulfonamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-methyl-N-(1,3-oxazol-2-yl)benzamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-[2-(morpholin-4-yl)-2-oxoethyl]benzamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-{8-methyl-8-azabicyclo[3.2.1]octan-3-yl}benzamide;

ethyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)carbamate;

tert-butyl N-[1-(3-chloro-4-{5-[2-(morpholin-4-yl)pyridin-4-yl]thiophen-3-yl}benzoyl)piperidin-3-yl]carbamate;

1-(3-chloro-4-{5-[2-(morpholin-4-yl)pyridin-4-yl]thiophen-3-yl}benzoyl)piperidin-3-amine;

1-{4-[5-(2-tert-butylpyridin-4-yl)-2-methylthiophen-3-yl]-3-chlorobenzoyl}piperidin-3-amine;

4-[5-(2-tert-butylpyridin-4-yl)-2-methylthiophen-3-yl]-3-chloro-N-phenylbenzamide;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-ol;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-ol;

tert-butyl N-[(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-N-methylcarbamate;

tert-butyl 3-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamido}piperidine-1-carboxylate;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-amine;

6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-5-chloro-N-phenylpyridine-3-carboxamide;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-methylpiperidin-3-amine;

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)-2,2-dimethylpropanamide;

4-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperazin-2-one;

tert-butyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamido}piperidine-1-carboxylate;

tert-butyl N-[(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

tert-butyl N-[(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-[(4-chlorophenyl)methyl]piperidin-3-amine;

(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-[(4-chlorophenyl)methyl]piperidin-3-amine;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-N-[(4-chlorophenyl)methyl]piperidin-3-amine;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-[(3-chlorophenyl)methyl]piperidin-3-amine;

(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}pyrrolidin-3-ol;

7-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2,7-diazaspiro[4.5]decan-1-one;

2-tert-butyl-4-[4-(2-chloro-4-{octahydropyrrolo[1,2-a]piperazine-2-carbonyl}phenyl)thiophen-2-yl]pyridine;

1-{4-[2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl]-3-chlorobenzoyl}piperidin-3-amine;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-4-ol;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidine-4-carboxylic acid;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-ol;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-ol;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-ol;

tert-butyl N-[(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}pyrrolidin-3-yl]carbamate;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidine-3-carboxylic acid;

2-tert-butyl-4-[5-(2-chloro-4-{3H,4H,5H,6H,7H-imidazo[4,5-c]pyridine-5-carbonyl}phenyl)thiophen-3-yl]pyridine;

4-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperazin-2-one;

1-[(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]imidazolidin-2-one;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-amine;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-amine;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-amine;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-amine;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-amine;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-1,2,3,4-tetrahydro-1,5-naphthyridine;

1-{4-[2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl]-3-chlorobenzoyl}piperidin-4-ol;

2-tert-butyl-4-{5-[2-chloro-4-(3-methoxypiperidine-1-carbonyl)phenyl]thiophen-3-yl}pyridine;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-4-methylpiperidin-4-ol;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-ol;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-ol;
 4-(3-chloro-4-{5-[1-(3-methylbutyl)-1H-pyrazol-5-yl]thiophen-3-yl}benzoyl)morpholine;
 1-(3-chloro-4-{5-[1-(3-methylbutyl)-1H-pyrazol-5-yl]thiophen-3-yl}benzoyl)-4,4-difluoropiperidine;
 1-(3-chloro-4-{4-[1-(3-methylbutyl)-1H-pyrazol-5-yl]thiophen-2-yl}benzoyl)piperidin-3-ol; and
 3-(1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorophenyl}-N-(4-fluorophenyl)formamido)propanoic acid.

[088] In some embodiments, the compound is selected from the group consisting of:

piperidin-1-yl(4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)methanone;
 piperidin-1-yl(3-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)methanone;
 (4-(5-(1-isobutyl-1H-pyrazol-5-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;
 (3-chloro-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;
 (4-(4-(2-tert-butylpyridin-4-yl)thiophen-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
 (3-chloro-4-(5-(1,2-dimethyl-1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)phenyl)
 (piperidin-1-yl)methanone;
 (3-chloro-4-(5-(1-propyl-1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;
 (3-chloro-4-(5-(2,3-dimethyl-3H-imidazo[4,5-b]pyridin-7-yl)thiophen-3-yl)phenyl)
 (piperidin-1-yl)methanone;
 (3-chloro-4-(5-(3-propyl-3H-imidazo[4,5-b]pyridin-7-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;
 (4-(5-(3H-imidazo[4,5-b]pyridin-7-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
 (3-chloro-4-(5-(1,2-dipropyl-1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)phenyl)
 (piperidin-1-yl)methanone;
 (4-(5-(6-(butylamino)-2-isobutylpyrimidin-4-yl)thiophen-3-yl)-3-chlorophenyl)
 (piperidin-1-yl)methanone;

(3-chloro-4-(5-(6-(dibutylamino)-2-isobutylpyrimidin-4-yl)thiophen-3-yl)phenyl)
(piperidin-1-yl)methanone;

(4-(5-(6-amino-2-isobutylpyrimidin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(2-aminopyrimidin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(propylamino)pyrimidin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-(butyl(propyl)amino)pyrimidin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-propylthiazol-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(5-propyl-1,2,4-thiadiazol-3-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(5-propyl-1,2,4-oxadiazol-3-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-propylthiazol-5-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(5-propyl-1,3,4-thiadiazol-2-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(5-propyl-1,3,4-oxadiazol-2-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

5-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)-1-isobutylpyridin-2(1H)-one;

(3-chloro-4-(5-(2-morpholinopyridin-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(2-bromo-3-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3,5-dichloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-(trifluoromethyl)phenyl)(piperidin-1-yl)methanone;

(2-cyclopropyl-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-methylphenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-isopropylphenyl)(piperidin-1-yl) methanone;

(2-ethynyl-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-3-ethynyl-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(2-bromo-6-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(2-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-6-methylphenyl)(piperidin-1-yl) methanone;

(4-(4-(2-tert-butylpyridin-4-yl)-1-methyl-1H-imidazol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(2-(2-isobutylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-3-chloro-4-(2-(2-isobutylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-5-chloro-4-(2-(2-isobutylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)oxazol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(2-(2-tert-butylpyridin-4-yl)oxazol-4-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(dimethylamino)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(3,5-dichloro-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

N-(4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)pyridin-2-yl) methanesulfonamide;

(3-chloro-4-(5-(2-(3,3,3-trifluoropropyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(1-methyl-2-(2-(3,3,3-trifluoropropyl)pyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-hydroxy-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-2-hydroxyphenyl)(piperidin-1-yl)methanone;

(2-methoxy-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-methoxyphenyl)(piperidin-1-yl)methanone;

(4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-2-methoxyphenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-3-(trifluoromethyl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-neopentylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-5-(trifluoromethyl)phenyl)(piperidin-1-yl)methanone;

(2,3-dichloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(2-(2-isobutylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)-2-methoxyphenyl)(piperidin-1-yl)methanone;

(2-methoxy-4-(1-methyl-2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(4-(2-(2-tert-butylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)-2-methoxyphenyl)(piperidin-1-yl)methanone;

(4-(5-(2-benzylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(2-aminopyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

N-(4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)-1-methyl-1H-imidazol-2-yl)pyridin-2-yl)methanesulfonamide;

(2-hydroxy-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(5-(2-(3,3,3-trifluoropropyl)pyridin-4-yl)furan-3-yl)phenyl)(piperidin-1-yl)
methanone;

(2-methoxy-4-(5-(2-neopentylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

N-(4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)pyridin-2-yl)-1,1,1-
trifluoromethanesulfonamide;

1-(4-(4-(4-(2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl)thiophen-2-yl)pyridin-
2-yl)piperazin-1-yl)ethanone;

(3-chloro-4-(5-(1-isopentyl-1H-pyrazol-5-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(5-(1-isopentyl-1H-pyrazol-5-yl)furan-3-yl)phenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(5-(2-(cyclohexylmethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-
yl)methanone;

(3-chloro-4-(5-(naphthalen-1-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(1H-indol-6-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(1H-indol-4-yl)thiophen-3-yl)-3-chlorophenyl)(4,4-difluoropiperidin-1-yl)
methanone;

(4-(5-(1H-benzo[d]imidazol-4-yl)thiophen-3-yl)-3-chlorophenyl)(4,4-difluoropiperidin-
1-yl)methanone;

(3-chloro-4-(5-(isoquinolin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(4-methylpiperazin-1-yl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-
1-yl)methanone;

(3-chloro-4-(5-(3,5-dimethylisoxazol-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(5-(1-isopentyl-1H-pyrazol-5-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-
1-yl)methanone;

(3-chloro-4-(5-(1-ethyl-1H-indol-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl) methanone;

(4-(5-(1H-indol-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(1-ethyl-1H-benzo[d]imidazol-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(5-(2,3-dihydrobenzofuran-5-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(2-(hydroxymethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(3-(hydroxymethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(2-(piperidin-1-yl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(3-methylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(2-(piperidin-1-yl)pyridin-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(5-(5-fluoro-1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(benzo[b]thiophen-3-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(3-(pyrrolidin-1-yl)methyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(cyclopentylamino)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(4-(5-(1H-indol-3-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(quinolin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-chloropyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl) methanone;

(2-bromo-4-(2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)
methanone;

(2-bromo-4-(5-(2-tert-butylpyridin-4-yl)furan-3-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(2-bromo-4-(5-(2-tert-butylpyridin-4-yl)-1H-pyrrol-3-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl)-3-fluorophenyl)(piperidin-1-yl)
methanone;

(4-(4-(2-tert-butylpyridin-4-yl)-1H-imidazol-2-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(2-bromo-4-(2-(2-(piperidin-1-yl)pyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)
methanone;

(4-(2-(2-tert-butylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(5-(2-tert-butylpyridin-4-yl)-1H-pyrrol-3-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(4-(2-tert-butylpyridin-4-yl)furan-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)-1H-pyrrol-2-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(4-(2-tert-butylpyridin-4-yl)-1-methyl-1H-pyrrol-2-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(4-(2-tert-butylpyridin-4-yl)thiazol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(2-tert-butylpyridin-4-yl)isothiazol-3-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(5-(2-tert-butylpyridin-4-yl)isoxazol-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(2-tert-butylpyridin-4-yl)-1H-pyrazol-3-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(3-(2-tert-butylpyridin-4-yl)isothiazol-5-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(3-(2-tert-butylpyridin-4-yl)isoxazol-5-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(3-(2-tert-butylpyridin-4-yl)-1H-pyrazol-5-yl)-3-chlorophenyl)(piperidin-1-yl) methanone;

(4-(3-(2-tert-butylpyridin-4-yl)-1-methyl-1H-pyrazol-5-yl)-3-chlorophenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(2-isobutylpyridin-4-yl)-1H-pyrrol-3-yl)phenyl)(piperidin-1-yl) methanone;

(3-fluoro-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(2-(2-isobutylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(2-(2-isobutylpyridin-4-yl)oxazol-4-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(2-(2-isobutylpyridin-4-yl)thiazol-4-yl)phenyl)(piperidin-1-yl) methanone;

(3-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(2-bromo-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(2-bromo-4-(2-(2-isobutylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl) methanone;

(4-(5-(2-butylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(2-(methoxymethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(4,6-dipropylpyridin-2-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

piperidin-1-yl(4-(5-(4-propylpyridin-2-yl)thiophen-3-yl)phenyl) methanone;

(4-(5-(2,6-dipropylpyridin-3-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(4-(5-(2-(cyclohexylmethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-3-methoxyphenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(2-((trifluoromethylsulfonyl)methyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(1-methyl-2-(1H-pyrrolo[2,3-b]pyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl) methanone;

(3-chloro-4-(5-(2-propyl-3H-imidazo[4,5-b]pyridin-7-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)-N-cyclopropyl picolinamide;

4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)-N-cyclohexyl picolinamide;

(4-(2-(1H-indol-4-yl)-1-methyl-1H-imidazol-4-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(4-(2-(piperidin-1-yl)pyridin-4-yl)thiophen-2-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(4-(2-(piperidin-1-yl)pyridin-4-yl)thiophen-2-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-chloro-4-(2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-5-chloro-4-(2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(5-(2-cyclopropylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

piperidin-1-yl(4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)-3-(trifluoromethyl)phenyl)methanone;

(3-chloro-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(3,3-difluoropyrrolidin-1-yl)methanone;

(3-chloro-4-(2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

5-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-(piperidine-1-carbonyl)benzonitrile;
 piperidin-1-yl(4-(5-(6-propylpyridin-3-yl)thiophen-3-yl)phenyl)methanone;
 (4-(5-(2-pentylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone; and
 (2-bromo-4-(1-methyl-2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)
 methanone.

[089] In some embodiments, the compound is selected from the group consisting of:

(4-(2-(1H-indol-4-yl)thiazol-4-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
 (4-(2-(2-tert-butylpyridin-4-yl)thiazol-4-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
 4-(2-(1H-indol-4-yl)thiazol-4-yl)-3-chloro-N,N-dimethylbenzamide;
 3-chloro-N,N-dimethyl-4-(2-(2-(piperidin-1-yl)pyridin-4-yl)thiazol-4-yl)benzamide;
 (4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
 (4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(3,3-difluoropyrrolidin-1-yl)methanone;
 (4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(4,4-difluoropiperidin-1-yl)methanone;
 4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-3-chloro-N-cyclopropylbenzamide; and
 N-tert-butyl-4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-3-chlorobenzamide.

[090] The compounds described below are not intended to be limiting; rather, these embodiments and variations are intended to provide examples of compounds within the scope of Formulae (Ia) or (Ib).

[091] Representative compounds are presented in Table 1 and Table 2.







Table 1		
 1	 2	 3
 4	 5	 6

Table 1

























 7	 8	 9
 10	 11	 12
 13	 14	 15
 16	 17	 18 18a, 18b
 19 19a, 19b	 20	 21 21, 21b
 22	 23	 24
 25	 26	 27
 28	 29	 30

























Table 1		
 31 31a, 31b	 32 32a, 32b	 33
 34 34a, 34b	 35	 36 36a, 36b
 37	 38	 39
 40	 41	 42
 43	 44 44a, 44b	 45
 46	 47	 48 48a, 48b
 49	 50 50a, 50b, 50c	 51
 52	 53	 54

























Table 1		
 55 55a, 55b	 56 56a, 56b	 57
 58	 59	 60
 61	 62	 63
 64 64a, 64b	 65 65a, 65b	 66 66a, 66b
 67	 68 68a, 68b	 69 69a, 69b
 70 70a, 70b	 71 71a, 71b	 72
 73	 74 74a, 74b	 75 75a, 75b
 76	 77	 78

























Table 1		
 79 79a, 79b	 80 80a, 80b	 81 81a, 81b
 82 82a, 82b	 83 83a, 83b	 84 84a, 84b
 85 85a, 85b	 86	 87 87a, 87b
 88	 89 89a, 89b	 90 90a, 90b
 91 91a, 91b	 92	 93
 94 94a, 94b	 95 95a, 95b	 96 96a, 96b
 97	 98	 99 99a, 99b
 100	 101 101a, 101b	 102 102a, 102b

Table 1



 103	 104 104, 104b	
--	---	--

Table 2






















 2.1	 2.2	 2.3
 2.4	 2.5	 2.6
 2.7	 2.8	 2.9
 2.10	 2.11	 2.12
 2.13	 2.14	 2.15
 2.16	 2.17	 2.18
 2.19	 2.20	 2.21

Table 2




























 2.22	 2.23	 2.24
 2.25	 2.26	 2.27
 2.28	 2.29	 2.30
 2.31	 2.32	 2.33
 2.34	 2.35	 2.36
 2.37	 2.38	 2.39
 2.40	 2.41	 2.42
 2.43	 2.44	 2.45
 2.46	 2.47	 2.48

Table 2




























 2.49	 2.50	 2.51
 2.52	 2.53	 2.54
 2.55	 2.56	 2.57
 2.58	 2.59	 2.60
 2.61	 2.62	 2.63
 2.64	 2.65	 2.66
 2.67	 2.68	 2.69
 2.70	 2.71	 2.72
 2.73	 2.74	 2.75

Table 2































 2.76	 2.77	 2.78
 2.79	 2.80	 2.81
 2.82	 2.83	 2.84
 2.85	 2.86	 2.87
 2.88	 2.89	 2.90
 2.91	 2.92	 2.93
 2.94	 2.95	 2.96
 2.97	 2.98	 2.99
 2.100	 2.101	 2.102
 2.103	 2.104	 2.105
















































Table 2		
 2.106	 2.107	 2.108
 2.109	 2.110	 2.111
 2.112	 2.113	 2.114
 2.115	 2.116	 2.117
 2.118	 2.119	 2.120
 2.121	 2.122	 2.123
 2.124	 2.125	 2.126
 2.127	 2.128	 2.129
 2.130	 2.131	 2.132

Table 2		
 2.133	 2.134	 2.135
 2.136	 2.137	 2.138
 2.139	 2.140	 2.141
 2.142	 2.143	 2.144
 2.145	 2.146	 2.147
 2.148	 2.149	 2.150
 2.151	 2.152	

[092] In some embodiments, the compound is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, or 104, or stereoisomers thereof.

[093] In some embodiments, the compound is 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 2.10, 2.11, 2.12, 2.13, 2.14, 2.15, 2.16, 2.17, 2.18, 2.19, 2.20, 2.21, 2.22, 2.23, 2.24, 2.25, 2.26, 2.27, 2.28, 2.29, 2.30, 2.31, 2.32, 2.33, 2.34, 2.35, 2.36, 2.37, 2.38, 2.39, 2.40, 2.41, 2.42, 2.43, 2.44, 2.45, 2.46, 2.47, 2.48, 2.49, 2.50, 2.51, 2.52, 2.53, 2.54, 2.55, 2.56, 2.57, 2.58, 2.59, 2.60, 2.61, 2.62, 2.63, 2.64, 2.65, 2.66, 2.67, 2.68, 2.69, 2.70, 2.71, 2.72, 2.73, 2.74, 2.75, 2.76, 2.77, 2.78,

2.79, 2.80, 2.81, 2.82, 2.83, 2.84, 2.85, 2.86, 2.87, 2.88, 2.89, 2.90, 2.91, 2.92, 2.93, 2.94, 2.95, 2.96, 2.97, 2.98, 2.99, 2.100, 2.101, 2.102, 2.103, 2.104, 2.105, 2.106, 2.107, 2.108, 2.109, 2.110, 2.111, 2.112, 2.113, 2.114, 2.115, 2.116, 2.117, 2.118, 2.119, 2.120, 2.121, 2.122, 2.123, 2.124, 2.125, 2.126, 2.127, 2.128, 2.129, 2.130, 2.131, 2.132, 2.133, 2.134, 2.135, 2.136, 2.137, 2.138, 2.139, 2.140, 2.141, 2.142, 2.143, 2.144, 2.145, 2.146, 2.147, 2.148, 2.149, 2.150, 2.151, or 2.152.

[094] Representative examples of compounds detailed herein, including intermediates and final compounds, are depicted in Table 1 and elsewhere herein. It is understood that in one aspect, any of the compounds may be used in the methods detailed herein, including, where applicable, intermediate compounds that may be isolated and administered to an individual.

[095] This disclosure also includes any or all of the stereochemical forms, including any enantiomeric or diastereomeric forms of the compounds described. The structure or name is intended to embrace all possible stereoisomers of a compound depicted, and each unique stereoisomer has a compound number bearing a suffix “a,” “b,” *etc.* All forms of the compounds are also embraced by this disclosure, such as crystalline or non-crystalline forms of the compounds. Compositions comprising one or more disclosed compounds are also intended, such as a composition of substantially pure compound, including a specific stereochemical form thereof, or a composition comprising mixtures of compounds in any ratio, including two or more stereochemical forms, such as in a racemic or non-racemic mixture.

[096] The compounds depicted herein may be present as salts even if salts are not depicted, and it is understood that the compositions and methods provided herein embrace all salts and solvates of the compounds depicted here, as well as the non-salt and non-solvate form of the compound, as is well understood by the skilled artisan. In some embodiments, the salts of the compounds provided herein are pharmaceutically acceptable salts.

[097] In one embodiment, the compound is a pharmaceutically acceptable salt of a compound of Formulae (Ia) or (Ib), or any variation presented herein, or a pharmaceutically acceptable salt thereof.

[098] The disclosed compounds may include isotopically-labeled and/or isotopically-enriched forms of the compounds. The compounds herein may contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. In some embodiments, the compound is isotopically-labeled, such as an isotopically-labeled compound of the Formulae (Ia) or (Ib), or variations thereof described herein, where a fraction of one or more atoms are replaced by an isotope of the same element. Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus,

sulfur, chlorine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}O , ^{17}O , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl . Certain isotope labeled compounds (e.g. ^3H and ^{14}C) are useful in compound or substrate tissue distribution study. Incorporation of heavier isotopes such as deuterium (^2H) can afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased in vivo half-life, or reduced dosage requirements, hence may be preferred in some instances.

[099] Isotopically-labeled compounds can generally be prepared by standard methods and techniques known to those skilled in the art or by procedures similar to those described in the accompanying Examples substituting appropriate isotopically-labeled reagents in place of the corresponding non-labeled reagent.

[0100] This disclosure also includes any or all metabolites of any of the compounds described. The metabolites may include any chemical species generated by a biotransformation of any of the compounds described, such as intermediates and products of metabolism of the compound.

[0101] In some embodiments, certain compounds presented herein are considered to be “prodrug” forms of other compounds herein. Prodrugs are precursor derivatives that, upon administration to a patient, undergo metabolism *in-vivo* such as, for example, hydrolysis to release the active form of the compound – the ‘parent’ compound. The prodrug form itself is either inactive or less active than the parent. Prodrugs are designed to improve bioavailability or to improve selective administration to particular organs, such as the liver [see, for example, Erion et al. *PNAS* (2007) 104:39, pp 15490-15495; Erion et al. *J. Pharmacol. Exp. Ther.* (2005) 312:2, pp 554-560; Meyer et al. *Patent Publication US 2006-0281695A1*]. In some embodiments, prodrug forms of compounds presented herein are provided. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in “*Design of Prodrugs*,” ed. H. Bundgaard, Elsevier, 1985; Beaumont, K. et al. *Curr. Drug Metab.* (2003) 4, pp 461-485; Mizen, L et al. *Pharm. Biotechnol.* (1998) 11, pp 345-365. In addition to prodrugs, this disclosure provides salts, esters, amides, and other protected or derivatized forms of the described compounds.

Pharmaceutically acceptable salts

[0102] “Pharmaceutically acceptable salts” are those salts which retain at least some of the biological activity of the free (non-salt) compound and which can be administered as drugs or pharmaceuticals to an individual. Such salts, for example, include: (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, oxalic acid,

propionic acid, succinic acid, maleic acid, tartaric acid and the like; (2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, *e.g.*, an alkali metal ion, an alkaline earth metal ion, or an aluminum ion; or coordinates with an organic base. Acceptable organic bases include ethanolamine, diethanolamine, triethanolamine and the like. Acceptable inorganic bases include aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, sodium hydroxide, and the like. Further examples of pharmaceutically acceptable salts include those listed in Berge *et al.*, *Pharmaceutical Salts*, *J. Pharm. Sci.* 1977 Jan; 66(1):1-19. Pharmaceutically acceptable salts can be prepared *in situ* in the manufacturing process, or by separately reacting a purified compound disclosed herein in its free acid or base form with a suitable organic or inorganic base or acid, respectively, and isolating the salt thus formed during subsequent purification. In one embodiment, a pharmaceutically acceptable salt includes the solvent addition forms or crystal forms thereof, particularly solvates or polymorphs. Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and are often formed during the process of crystallization. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Polymorphs include the different crystal packing arrangements of the same elemental composition of a compound. Polymorphs usually have different X-ray diffraction patterns, infrared spectra, melting points, density, hardness, crystal shape, optical and electrical properties, stability, and solubility. Various factors such as the recrystallization solvent, rate of crystallization, and storage temperature may cause a single crystal form to dominate.

Pharmaceutical Compositions

[0103] A compound of Formulae (Ia) or (Ib), or any variation presented herein, typically is provided in a pharmaceutical composition comprising a pharmaceutically acceptable carrier and/or excipient. A “pharmaceutically acceptable” carrier or excipient is a material that is not biologically or otherwise undesirable, *e.g.*, the material may be incorporated into a pharmaceutical composition administered to an individual without causing significant undesirable biological effects or interacting in a deleterious manner with any of the other components of the composition, wherein it is contained. Pharmaceutically acceptable carriers or excipients meet the required standards of toxicological and manufacturing testing and/or are included on the Inactive Ingredient Guide prepared by the U.S. Food and Drug administration.

[0104] A pharmaceutical composition can comprise one or more compounds of Formulae (Ia) or (Ib), or any variation presented herein. In some embodiments, a pharmaceutical composition further comprises chemotherapeutic agent, as described below.

[0105] Preferably a compound of Formulae (Ia) or (Ib), or any variation presented herein, is bioavailable orally. However, the compounds may also be formulated for parenteral (*e.g.*, intravenous) administration.

[0106] A compound of Formulae (Ia) or (Ib), or any variation presented herein, can be used in the preparation of a medicament by combining the compound or compounds as an active ingredient with a pharmacologically acceptable carrier, which is known in the art. Depending on the therapeutic form of the medication, the carrier may be in various forms. In one variation, the manufacture of a medicament is for use in any of the methods disclosed herein.

[0107] Methods as provided herein may comprise administering to an individual a pharmacological composition that contains an effective amount of a compound of Formulae (Ia) or (Ib), or any variation presented herein, and a pharmaceutically acceptable carrier. The effective amount of the compound may in one aspect be a dose of between about 0.01 and about 100 mg.

[0108] The compound may be formulated for any available delivery route, including an oral, mucosal (*e.g.*, nasal, sublingual, vaginal, buccal or rectal), parenteral (*e.g.*, intramuscular, subcutaneous or intravenous), topical or transdermal delivery form. A compound may be formulated with suitable carriers to provide delivery forms that include, but are not limited to, tablets, caplets, capsules (such as hard gelatin capsules or soft elastic gelatin capsules), cachets, troches, lozenges, gums, dispersions, suppositories, ointments, cataplasms (poultices), pastes, powders, dressings, creams, solutions, patches, aerosols (*e.g.*, nasal spray or inhalers), gels, suspensions (*e.g.*, aqueous or non-aqueous liquid suspensions, oil-in-water emulsions or water-in-oil liquid emulsions), solutions and elixirs.

[0109] A compound of Formulae (Ia) or (Ib), or any variation presented herein, can be used in the preparation of a formulation, such as a pharmaceutical formulation, by combining the compound or compounds as an active ingredient with a pharmaceutically acceptable carrier, such as those mentioned above. Depending on the therapeutic form of the system (*e.g.*, transdermal patch vs. oral tablet), the carrier may be in various forms. In addition, pharmaceutical formulations may contain preservatives, solubilizers, stabilizers, re-wetting agents, emulgators, sweeteners, dyes, adjusters, salts for the adjustment of osmotic pressure, buffers, coating agents or antioxidants. Formulations comprising the compound may also contain other substances which have valuable therapeutic properties. Pharmaceutical formulations may be prepared by known pharmaceutical methods. Suitable formulations can be found, *e.g.*, in *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Philadelphia, PA, 20th ed. (2000), which is incorporated herein by reference.

[0110] A compound of Formulae (Ia) or (Ib), or any variation presented herein, may be administered to individuals in a form of generally accepted oral compositions, such as tablets, coated tablets, gel capsules in a hard or in soft shell, emulsions or suspensions. Examples of carriers, which may be used for the preparation of such compositions, are lactose, corn starch or its derivatives, talc, stearate or its salts, *etc.* Acceptable carriers for gel capsules with soft shell are, for instance, plant oils, wax, fats, semisolid and liquid poly-ols, and so on. In addition, pharmaceutical formulations may contain preservatives, solubilizers, stabilizers, re-wetting agents, emulgators, sweeteners, dyes, adjusters, salts for the adjustment of osmotic pressure, buffers, coating agents or antioxidants.

[0111] A compound of Formulae (Ia) or (Ib), or any variation presented herein, can be formulated in a tablet in any dosage form described, for example, a compound as described herein or a pharmaceutically acceptable salt thereof can be formulated as a 10 mg tablet.

[0112] A compound of Formulae (Ia) or (Ib), or any variation presented herein, may be administered to an individual in accordance with an effective dosing regimen for a desired period of time or duration, such as at least about one month, at least about 2 months, at least about 3 months, at least about 6 months, or at least about 12 months or longer, which in some variations may be for the duration of the individual's life. In one variation, a compound of Formulae (Ia) or (Ib), or any variation presented herein, is administered on a daily or intermittent schedule. The compound can be administered to an individual continuously (for example, at least once daily) over a period of time. The dosing frequency can also be less than once daily, *e.g.*, about a once weekly dosing. The dosing frequency can be more than once daily, *e.g.*, twice or three times daily. The dosing frequency can also be intermittent (*e.g.*, once daily dosing for 7 days followed by no doses for 7 days, repeated for any 14 day time period, such as about 2 months, about 4 months, about 6 months or more). Any of the dosing frequencies can employ any of the compounds described herein together with any of the dosages described herein.

[0113] In some embodiments, a pharmaceutical composition is provided as a unit dosage form, such as a tablet, capsule, or individually packaged container (*e.g.*, an ampoule, syringe, or vial).

[0114] In some embodiments, the unit dosage form contains a daily dose of a compound of Formulae (Ia) or (Ib), or any variation presented herein. In some embodiments, the unit dosage form contains a daily sub-dose of the compound.

[0115] In some embodiments, the unit dosage form contains a daily dose of each of two or more compounds of Formulae (Ia) or (Ib), or any variation presented herein. In some embodiments, the unit dosage form contains a daily sub-dose of each of two or more compounds.

[0116] In some embodiments, the unit dosage form contains a daily dose of compound of Formulae (Ia) or (Ib), or any variation presented herein, and a daily dose of each of one or more chemotherapeutic agents. In some embodiments, the unit dosage form contains a daily sub-dose of the compound and a daily sub-dose of each of one or more chemotherapeutic agents.

[0117] In some embodiments, the unit dosage form contains a daily dose of each of two or more compounds of Formulae (Ia) or (Ib), or any variation presented herein, and a daily dose of each of one or more chemotherapeutic agents. In some embodiments, the unit dosage form contains a daily sub-dose of each of two or more compounds and a daily dose of each of one or more chemotherapeutic agents.

Kits and Articles of Manufacture

[0118] This disclosure also provides kits and articles of manufacture comprising one or more compounds of Formulae (Ia) or (Ib), or any variation presented herein, or a pharmacological composition comprising a compound of Formulae (Ia) or (Ib), or any variation presented herein,. The kits may employ any of the compounds disclosed herein. In one variation, the kit employs a compound described herein or a pharmaceutically acceptable salt thereof.

[0119] Kits generally comprise suitable packaging. The kits may comprise one or more containers comprising any compound described herein. Each component (if there is more than one component) can be packaged in separate containers or some components can be combined in one container where cross-reactivity and shelf life permit.

[0120] The kits may be in unit dosage forms, bulk packages (*e.g.*, multi-dose packages) or sub-unit doses. For example, kits may be provided that contain sufficient dosages of a compound as disclosed herein and/or a second pharmaceutically active compound useful for a disease detailed herein to provide effective treatment of an individual for an extended period, such as any of a week, 2 weeks, 3 weeks, 4 weeks, 6 weeks, 8 weeks, 3 months, 4 months, 5 months, 7 months, 8 months, 9 months, or more. Kits may also include multiple unit doses of the compounds and instructions for use and be packaged in quantities sufficient for storage and use in pharmacies (*e.g.*, hospital pharmacies and compounding pharmacies).

[0121] The kits may optionally include a set of instructions, generally written instructions, although electronic storage media (*e.g.*, magnetic diskette or optical disk) containing instructions are also acceptable, relating to the use of component(s) of the disclosed methods. The instructions included with the kit generally include information as to the components and their administration to an individual.

Therapeutic Uses

[0122] Unless otherwise defined, “treat,” as used herein, refers to the reduction of one or more symptoms associated with a disorder or to slowing the progression of one or more such symptoms.

[0123] Compounds of Formulae (Ia) or (Ib), or any variation presented herein, can be used to treat hyperproliferative disorders. A “hyperproliferative disorder” is a disorder associated with some degree of abnormal cell proliferation. A hyperproliferative disorder can be benign (including pre-cancerous disorders) or malignant.

[0124] In some embodiments, the hyperproliferative disorder is benign, such as benign prostatic hyperplasia, neurofibromatosis, actinic keratosis, hypertrophic actinic keratosis, Bowenoid actinic keratosis, arsenical keratosis, hydrocarbon keratosis, thermal keratosis, radiation keratosis, chronic scar keratosis, viral keratosis, actinic cheilitis, Bowen’s disease, erythroplakia of queyrat, oral erythroplakia, leukoplakia, intraepidermal epithelioma, psoriasis, polyps, Barrett’s esophagus, atrophic gastritis, cervical dysplasia, benign meningioma, and benign ovarian epithelial tumors (*e.g.*, serous adenomas, mucinous adenomas, Brenner tumors).

[0125] In some embodiments, the hyperproliferative disorder is malignant, *e.g.*, adenocarcinoma, bladder cancer, breast cancer, cervical cancer, cholangiocarcinoma, CNS cancer (*e.g.*, astrocytoma, dendroma, ependymoma, glioma, malignant meningioma, medulloblastoma, neuroblastoma, neuroglioma, oligodendroglioma), gastrointestinal cancer (*e.g.*, gastrointestinal stromal carcinoma, colorectal cancer), kidney cancer, leukemia (*e.g.*, acute lymphocytic leukemia; acute myelogenous leukemia; chronic lymphocytic leukemia; chronic myelogenous leukemia), liver cancer (*e.g.*, hepatic cancer, hepatocellular carcinoma), lung cancer (*e.g.*, lung squamous carcinoma, small-cell lung carcinoma, non-small-cell lung carcinoma, mesothelioma), lymphoma (*e.g.*, Hodgkin lymphoma and non-Hodgkin lymphoma), melanoma, myeloma (*e.g.*, multiple myeloma, plasmacytoma), ovarian cancer, pancreatic cancer, prostate cancer, renal cancer, thyroid cancer, and uterine cancer.

[0126] In some embodiments, the breast cancer is AR+, ER+, and Her2+. In some embodiments, the breast cancer is AR+, ER+, and PR+. In some embodiments, the breast cancer is AR+, ER+, Her2+, and PR+. In some embodiments, the breast cancer is AR-, ER+, and Her2+. In some embodiments, the breast cancer is AR-, ER+, and PR+. In some embodiments, the breast cancer is AR-, ER+, Her2+, and PR+.

[0127] In some embodiments, the breast cancer is ductal carcinoma *in situ*. In some embodiments, the breast cancer is invasive ductal carcinoma. In some embodiments, the breast

cancer is triple negative breast cancer (*e.g.*, basal-like type 1 (BL1), basal-like type 2 (BL2), immunomodulatory (IM), mesenchymal (M), mesenchymal stem-like (MSL), and luminal androgen receptor (LAR) subtypes). In some embodiments, the breast cancer is inflammatory breast cancer. In some embodiments, the breast cancer is BRCA1-related breast cancer. In some embodiments, the breast cancer is medullary breast cancer, metaplastic breast cancer. In some embodiments, the breast cancer is special histologic type of breast cancer. In some embodiments, the breast cancer is resistant to endocrine therapy.

[0128] In some embodiments, the prostate cancer is hormone-sensitive prostate cancer. In some embodiments, the prostate cancer is castration-resistant prostate cancer.

[0129] In some embodiments, the ovarian cancer is an epithelial carcinoma. In some embodiments, the ovarian cancer is a germ cell tumor. In some embodiments, the ovarian cancer is an ovarian stromal tumor (*e.g.*, granulosa-theca tumors and Sertoli-Leydig cell tumors).

[0130] Compounds disclosed herein also can be used to treat a variety of metabolic disorders, including metabolic disorders that are mediated by genetic factors (*e.g.*, Niemann-Pick disease, Fabry disease, Gaucher disease, Forbe's disease, Tangier disease) and environmental factors (*e.g.*, diets rich in fat and/or sugar). Compounds also may be useful in the treatment of complications of metabolic diseases, such as cardiovascular disease, non-alcoholic hepatic steatosis, hyperlipemia, and obesity.

[0131] Compounds disclosed herein also can be used to treat pancreatitis.

[0132] Compounds can be used to treat liver fibrosis, elevated cholesterol levels, and insulin resistance.

a. Liver Fibrosis

[0133] Compounds disclosed herein can be administered to an individual to treat liver fibrosis. In some embodiments, the liver fibrosis is secondary to chronic hepatitis C virus infection. In some embodiments, the liver fibrosis is secondary to alcohol abuse. In some embodiments, the liver fibrosis is secondary to nonalcoholic steatohepatitis (NASH).

[0134] In some embodiments in which the liver fibrosis is secondary to NASH, the individual has been diagnosed with NASH following a liver biopsy in which one or more of steatosis, hepatocyte ballooning, lobular inflammation, Mallory hyaline bodies, mixed inflammatory infiltrate, pericellular fibrosis, and perisinusoidal fibrosis is detected.

[0135] In other embodiments in which the liver fibrosis is secondary to NASH, the individual is suspected of having NASH based on one or more symptoms such as such as elevated serum

levels of liver enzymes (*e.g.*, alanine aminotransferase (ALT), aspartate aminotransferase (AST), γ -glutamyltransferase, alkaline phosphatase); focal or diffuse accumulation of lipid as detected by imaging techniques such as magnetic resonance spectroscopy, ultrasonography, computed tomography; abdominal discomfort, acanthosis nigricans, bowel dysmotility, coma, constipation, disseminated intravascular coagulopathy, epigastric pain, fatigue, fluid retention, hepatomegaly, hypoglycemia, intestinal bleeding, jaundice, lipomatosis, lipoatrophy, lipodystrophy, muscle wasting, nausea, neurological defects, Palmer erythema, panniculitis, periumbilical pain, small bowel bacterial overgrowth, spider angiomas, splenomegaly, subacute liver failure, or vomiting.

[0136] In some embodiments, the individual is overweight or obese. In some embodiments, the individual has symptoms of insulin resistance, as described below; *i.e.*, the individual is pre-diabetic or has type II diabetes.

[0137] In some embodiments, administration of one or more disclosed compounds prevents or slows the histologic progression of liver fibrosis and the clinical progression to cirrhosis in patients with NASH.

[0138] Treatment can be assessed by one or more of the following:

- reduction in average liver fat concentration (measured, *e.g.*, by NMRS or MRI);
- reduction of serum ALT;
- reduction of serum AST;
- reduction of serum γ -glutamyltransferase;
- reduction of serum alkaline phosphatase;
- increased in plasma concentrations of the cholesterol synthesis intermediate lathosterol;
- improved NASH activity index or NAFLD activity score (NAS) (Kleiner *et al.*, Hepatology 41, 1313-21, 2005);
- improved SAF score (Bedossa *et al.*, Hepatology 56, 1751-59, 2012);
- changes in insulin resistance (measured, *e.g.*, by Homeostasis Model Assessment of Insulin Resistance (HOMA-IR);
- reduced hemoglobin A1c levels;
- adiponectin level;
- leptin:adiponectin ratio (LAR);
- reduction in one or more markers of inflammation or fibrosis such as fibrinogen, CK-18, C-reactive protein (CRP), TNF α , IL-6;
- reduction in body weight; or

reduced histological features such as cholestasis, fat cysts, fibrosis, granular iron, hepatocellular ballooning, increased numbers of eosinophils, inflammation, lobular disarray, lobular inflammation, macrovesicular steatosis, Mallory bodies, megamitochondria, necrosis, periodic acid-Schiff stained globulins, portal inflammation, microvesicular steatosis, or steatosis.

b. Elevated Cholesterol Levels

[0139] Compounds disclosed herein can be administered to an individual to treat elevated cholesterol levels, *e.g.*, cholesterol levels above 200 mg/dL (5.2 mmol/L), such as 200-239 mg/dL (5.2-6.2 mmol/L) or 240 mg/dL (6.2 mmol/L) and above. In some embodiments, the individual has LDL cholesterol levels in the range of 130-159 mg/dL (3.4-4.1 mmol/L). In some embodiments, the individual has LDL cholesterol levels in the range of 160-189 mg/dL (4.1-4.9 mmol/L). In some embodiments, the individual has LDL cholesterol levels in the range of 190 mg/dL (4.9 mmol/L) and above. In some embodiments, the individual has HDL cholesterol levels below 40 mg/dL (1 mmol/L; men) or below 50 mg/dL (1.3 mmol/L; women). In some embodiments, the individual has HDL cholesterol levels in the range of 50-59 mg/dL (1.3-1.5 mmol/L). In some embodiments, the individual has triglyceride levels in the range of 150-199 mg/dL (1.7-2.2 mmol/L). In some embodiments, the individual has triglyceride levels in the range of 200-499 mg/dL (2.3-5.6 mmol/L). In some embodiments, the individual has triglyceride levels in the range of 500 mg/dL (5.6 mmol/L) and above.

[0140] In some embodiments, the individual has NASH or is suspected of having NASH, as described above.

[0141] In some embodiments, the individual has symptoms of insulin resistance, as described below.

c. Insulin Resistance

[0142] Compounds disclosed herein can be administered to an individual to treat insulin resistance. In some embodiments, the individual has pre-diabetes, *e.g.*, the individual has a hemoglobin A1C level between 5.7 and 6.4 percent and/or a fasting blood sugar level from 100-125 mg/dL (5.6-6.9 mmol/L). In some embodiments, the individual has type II diabetes *e.g.*, the individual has a hemoglobin A1C level of 6.5 percent or higher and/or a fasting blood sugar level of 126 mg/dL (7 mmol/L) or higher.

[0143] In some embodiments, the individual has one or more of the following symptoms, which may improve upon treatment with one or more compounds disclosed herein: increased thirst and

frequent urination, increased hunger; weight loss; fatigue; blurred vision; slow-healing sores or frequent infections; and acanthosis nigricans.

[0144] In some embodiments, the individual has NASH or is suspected of having NASH, as described above.

d. Combination Therapies

[0145] Compounds disclosed herein can be administered in combination with other therapeutic interventions for treating elevated cholesterol levels, liver fibrosis, or insulin resistance. Unless otherwise defined, “in combination” includes any coordinated administration of such therapeutic interventions with one or more therapeutic compounds disclosed herein, including sequential administration, alternating administration, and substantially simultaneous administration.

[0146] Therapeutic interventions for treating elevated cholesterol levels include, but are not limited to, statins, such as atorvastatin (*e.g.*, LIPITOR®), fluvastatin (*e.g.*, LESCOL®), lovastatin (*e.g.*, ALTOPREV®, MEVACOR®), pitavastatin (*E.G.*, LIVALO®), pravastatin (*e.g.*, PRAVACHOL®), rosuvastatin (*e.g.*, CRESTOR®), and SIMVASTATIN (*e.g.*, ZOCOR®); bile acid binding resins, such as cholestyramine (*e.g.*, PREVALITE®), colestevlam (*e.g.*, WELCHOL®), and colestipol (*e.g.*, COLESTID®); and cholesterol absorption inhibitors such as ezetimibe (*e.g.*, ZETIA®).

[0147] Therapeutic interventions for treating liver fibrosis include, but are not limited to, angiotensin inhibitors, colchicine, corticosteroids, endothelin inhibitors, interferon- α , interleukin 10, pentoxifylline or oxpentifylline (*e.g.*, TRENTAL®), phosphatidylcholine, PPAR antagonists, S-adenosyl-methionine, TGF- β inhibitors, and tocopherol.

[0148] Therapeutic interventions for treating insulin resistance include, but are not limited to, insulin-sensitizing agents such as metformin (*e.g.*, GLUCOPHAGE®), thiazolidinediones such as pioglitazone (*e.g.*, ACTOS®) and rosiglitazone (*e.g.*, AVANDIA®); and leptin); α -glucosidase inhibitors such as miglitol (*e.g.*, GIYSET®); insulin; meglitinides such as repaglinide (*e.g.*, PRANDIN®) and nateglinide (*e.g.*, STARLIX®); sulfonylureas such as glyburide (*e.g.*, ORINASE®, TOLINASE®, MICRONASE®, GLYNASE®, DIABETA®, AMARYL®) and chlorpropamide (*e.g.*, DIABINASE®, GLUTROL®, GLUCOTROL XL®); and combinations such as AVANDAMET® (metformin and rosiglitazone).

Treatment Regimens

[0149] Compounds can be administered alone or in conjunction with other therapeutic interventions. The disclosed compounds decrease the synthesis of cholesterol and fatty acids, which are essential components of cell membranes and cell division; accordingly, administration

of a compound should decrease the rate of cell division. These effects, coupled with alterations in lipid-mediated cell signaling pathways, induce cell death.

[0150] Administration of a compound of Formulae (Ia) or (Ib), or any variation presented herein, “in conjunction with” another therapeutic intervention may include any of the following regimens.

Dosing and Method of Administration

[0151] The dose of a compound administered to an individual (such as a human) may vary with the particular compound or salt thereof, the method of administration, and the particular stage of hyperproliferative or metabolic disorder being treated. For therapeutic purposes, “effective dose” or “effective amount” refers to that amount of each active compound or pharmaceutical agent, alone or in combination, that elicits the biological or medicinal response in a tissue system, animal, or human that is being sought by a researcher, veterinarian, medical doctor, or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated. For prophylactic purposes (i.e., preventing or inhibiting the onset or progression of a disorder), the term “effective dose” or “effective amount” refers to that amount of each active compound or pharmaceutical agent, alone or in combination, that inhibits in a subject the onset or progression of a disorder as being sought by a researcher, veterinarian, medical doctor, or other clinician, the delaying of which disorder is mediated, at least in part, by blocking of SREBP function. The amount should be sufficient to produce a desirable response, such as a therapeutic or prophylactic response against the disorder. In some embodiments, the amount of the compound or salt thereof is a therapeutically effective amount. In some embodiments, the amount of the compound or salt thereof is a prophylactically effective amount. In some embodiments, the amount of compound or salt thereof is below the level that induces a toxicological effect (e.g., an effect above a clinically acceptable level of toxicity) or is at a level where a potential side effect can be controlled or tolerated when the composition is administered to the individual.

[0152] In some embodiments, the amount of compound or salt thereof is an amount sufficient to inhibit cancer cell growth and/or proliferation or increase apoptosis of cancer cells.

[0153] The effective amount of the compound may in one aspect be a dose of between about 0.01 and about 100 mg/kg.

[0154] Any of the methods provided herein may in one aspect comprise administering to an individual a pharmaceutical composition that contains an effective amount of a compound provided herein or a salt thereof and a pharmaceutically acceptable excipient.

[0155] A compound or composition provided herein may be administered to an individual in accordance with an effective dosing regimen for a desired period of time or duration, such as at least about one month, at least about 2 months, at least about 3 months, at least about 6 months, or at least about 12 months or longer, which in some variations may be for the duration of the individual's life. In one variation, the compound is administered on a daily or intermittent schedule. The compound can be administered to an individual continuously (for example, at least once daily) over a period of time. The dosing frequency can also be less than once daily, *e.g.*, about a once weekly dosing. The dosing frequency can be more than once daily, *e.g.*, twice or three times daily. The dosing frequency can also be intermittent (*e.g.*, once daily dosing for 7 days followed by no doses for 7 days, repeated for any 14 day time period, such as about 2 months, about 4 months, about 6 months or more). Any of the dosing frequencies can employ any of the compounds described herein together with any of the dosages described herein.

[0156] The compounds provided herein or a salt thereof may be administered to an individual via various routes, including, *e.g.*, intravenous, intramuscular, subcutaneous, oral and transdermal.

[0157] In one aspect, provided is a method of treating cancer in an individual by parenterally administering to the individual (*e.g.*, a human) an effective amount of a compound or salt thereof. In some embodiments, the route of administration is intravenous, intra-arterial, intramuscular, or subcutaneous. In some embodiments, the route of administration is oral. In still other embodiments, the route of administration is transdermal. In one aspect are provided compositions (including pharmaceutical compositions) as described herein for the use in treating a hyperproliferative or metabolic disorder, such as cancer (*e.g.*, prostate cancer).

[0158] Also provided are compositions (including pharmaceutical compositions) as described herein for the use in treating, preventing, and/or delaying the onset and/or development of cancer and other methods described herein. In certain embodiments, the composition comprises a pharmaceutical formulation which is present in a unit dosage form.

[0159] Also provided are articles of manufacture, comprising a compound provided herein or a salt thereof, composition, and unit dosages described herein in suitable packaging for use in the methods described herein. Suitable packaging is known in the art and includes, for example, vials, vessels, ampules, bottles, jars, flexible packaging and the like. An article of manufacture may further be sterilized and/or sealed.

Anti-Proliferative Agents

[0160] An “anti-proliferative agent” is an intervention that increases apoptosis of hyperproliferating cells. In some embodiments, a compound of Formulae (Ia) or (Ib), or any variation presented herein, is used in conjunction with an anti-proliferative agent which is a chemotherapeutic agent. Chemotherapeutic agents include any pharmacological agent which is currently approved by the FDA in the U.S. (or elsewhere by any other regulatory body) for use as pharmacological treatment for hyperproliferative disorders, including cancer, or which is currently being used experimentally as part of a clinical trial program.

General Synthetic Methods

[0161] The compounds may be prepared by a number of processes as generally described below in the General Synthetic Schemes and more specifically in the Examples hereinafter. In the following process descriptions, the symbols when used in the formulae depicted are to be understood to represent those groups described above in relation to the formulae herein.


[0162] Chromatography, recrystallization and other conventional separation procedures may also be used with intermediates or final products where it is desired to obtain a particular isomer of a compound or to otherwise purify a product of a reaction.

[0163] The following abbreviations are used herein: thin layer chromatography (TLC); h (h); minute (min); second (sec); ethanol (EtOH); dimethylsulfoxide (DMSO); *N,N*-dimethylformamide (DMF); trifluoroacetic acid (TFA); tetrahydrofuran (THF); Normal(N); aqueous (aq.); methanol (MeOH); dichloromethane (DCM); ethyl acetate (EtOAc); Retention factor (Rf); room temperature (RT).

[0164] The following General Synthetic Schemes and Examples are provided to illustrate but not to limit the scope of this disclosure. Those skilled in the art will be familiar with many of the reaction steps described. See also WO2015/031650, which is incorporated herein by reference in its entirety.

General Synthetic Scheme 1

[0165] General Synthetic Scheme 1 provides methods to prepare compounds with a thiazole or imidazole B-ring as presented herein. Substituents R₁-R₆ are as exemplified in the Examples below. Syntheses of tricyclic substituted thiazoles, oxazoles and imidazoles, such as those presented herein, will be familiar to those skilled in the art. An example to illustrate a synthesis of a substituted thiazole is presented below. Complete details for syntheses of the compounds presented herein are provided in the Examples.




A = aromatic or heteroaromatic ring
Y = O, S, NH

[0166] General procedure: The corresponding substituted pyridine-4-carbothioamide or isonicotinamide and the corresponding substituted 2-bromoacetylbenzene are dissolved in EtOH. The resultant reaction mixture is stirred at between RT and 70 °C for between 30 min and 2h. The progress of the reaction is monitored by TLC and LCMS. The reaction mixture is cooled to RT, basified with aq. sodium bicarbonate solution and the mixture extracted with EtOAc. The organic layer is dried over sodium sulfate and concentrated to obtain the crude product, which is purified by silica gel (100-200 mesh) column chromatography or by HPLC to obtain the desired product.

EXAMPLES

Example 1. Preparation of Compound No. 1

Synthesis of N-methyl-4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide




[0167] To DCM (2 mL) was added oxalyl chloride (0.04 mL, 0.462 mmol) and DMF (0.01 mL) dropwise respectively at 0 °C. The resultant mixture was stirred at 0 °C for 10 min. To the reaction mixture was added 4-(2-(2-propylpyridin-4-yl)thiazol-4-yl)benzoic acid (50 mg, 0.154 mmol), and the reaction mixture was allowed to come to RT. The reaction mixture was stirred at RT for 30 min. To the reaction mixture was added methyl amine (2M in THF) (0.23 mL, 0.462 mmol) dropwise at 0 °C and the mixture stirred for 10 min. To this reaction mixture was added water (2 mL) and then the reaction mixture was diluted with DCM (10 mL). The DCM layer was separated and the aqueous layer was extracted with DCM (2x5 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated to obtain the crude product. The crude product was purified by HPLC to obtain N-methyl-4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide (10 mg).

[0168] ^1H NMR (methanol- d_4): δ (ppm): 8.75 (d, $J=6.1$ Hz, 1H), 8.42 (m, 3H), 8.20 (d, $J=8.0$ Hz, 2H), 7.95 (d, $J=8.0$ Hz, 2H), 3.02 (t, 2H), 2.95 (s, 3H), 1.91 (h, 2H), 1.09 (t, 3H).

Example 2. Preparation of Compound No. 2

Synthesis of N,N-dimethyl-4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide




[0169] To DCM (2 mL) was added oxalyl chloride (0.04 mL, 0.462 mmol) and DMF (0.01 mL) dropwise respectively at 0 °C. The resultant mixture was stirred at 0 °C for 10 min. To this reaction mixture was added 4-(2-(2-propylpyridin-4-yl)thiazol-4-yl)benzoic acid (50 mg, 0.154 mmol) and the reaction mixture was allowed to come to RT. The reaction mixture was then stirred at RT for 30 min. To this reaction mixture was added liquid ammonia (till pH became basic) dropwise at 0 °C and the mixture stirred for 10 min at the same temperature. Then, to the reaction mixture was added water (2 mL) and the reaction mixture was diluted with DCM (10 mL). The DCM layer was separated and the aqueous layer was extracted with DCM (2x5 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated to obtain the crude product, as a mixture with Compound No. 3. The crude product was purified by HPLC to get N,N-dimethyl-4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide (11 mg).

[0170] ^1H NMR (methanol- d_4): δ (ppm): 8.78 (d, $J=6.3$ Hz, 1H), 8.55 (s, 1H), 8.47 (m, 2H), 8.21 (d, $J=8.0$, 2H), 7.57 (d, $J=7.8$ Hz, 2H), 3.13-3.08 (m, 8H), 1.91 (m, 2H), 1.10 (t, $J=7.3$ Hz 3H).

Example 3. Preparation of Compound No. 3

Synthesis of 4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide




[0171] To DCM (2 mL) was added oxalyl chloride (0.04 mL, 0.462 mmol) and DMF (0.01 mL) dropwise respectively at 0 °C. The resultant mixture was stirred at the same temperature for 10 min. To this stirred reaction mixture was added 4-(2-(2-propylpyridin-4-yl)thiazol-4-yl)benzoic acid (50 mg, 0.154 mmol) and the reaction mixture was allowed to come to RT. The mixture was stirred at RT for 30 min. To this reaction mixture was added liquid ammonia (till pH became

basic) dropwise at 0 °C and the mixture stirred for 10 min. Water (2 mL) was added to the reaction mixture and then the reaction mixture was diluted with DCM (10 mL). The DCM layer was separated and the aqueous layer was extracted with DCM (2x5 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated to obtain the crude product, as a mixture with Compound No. 2. The crude product was purified by HPLC to obtain get 4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide (12 mg).

[0172] ^1H NMR (methanol- d_4): δ (ppm): 8.78 (d, $J=6.0$ Hz, 1H), 8.53 (s, 1H), 8.47 (d, $J=4.8$, 1H), 8.22 (d, $J=8.0$ Hz, 2H), 8.01 (d, $J=7.9$ Hz, 2H), 3.18-2.77 (m, 3H), 1.92 (m, 2H), 1.10 (t, $J=7.3$, 3H).

Example 4. Preparation of Compound No. 4

Synthesis of 4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1,3-thiazol-2-yl}-1H-indole



Step-1: Synthesis of [2-chloro-4-(piperidine-1-carbonyl)phenyl]boronic acid

[0173] In a 250 mL flask was placed 4-bromo-3-chlorobenzoic acid (1.0 g, 5 mmol, 1eq.) in DMF (25 mL), followed by addition of DIPEA (3.6 mL, 20 mmol, 4 eq.) and the mixture was stirred for 5 min at RT. Then HATU (3.8 g, 10 mmol, 2 eq.) and piperidine (1.7 mL, 17.5 mmol, 3.5 eq.) were added and the mixture stirred at RT overnight. After completion of reaction (monitored by TLC and ^1H -NMR/LCMS), the mixture was diluted with 1N aq. HCl (100 mL) and extracted with EtOAc (2x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by silica gel (60-120) column chromatography using acetone: hexane (0-35%) to elute pure compound as a semisolid which solidified in a refrigerator overnight (1.0g).

Step 2: Synthesis of 4-bromo-2-(1H-indol-4-yl)thiazole

[0174] In a 100 mL glass bottle was placed 2,4 dibromothiazole (1 g, 4.11 mmol, 1eq.) along with 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (1 g, 4.11 mmol, 1eq.) and potassium phosphate (2.18 g, 2.5 eq.) in (30 mL) of THF. Then it was purged with nitrogen for 15 min then Xantphos (167 mg, 0.07 eq.) and tetrakis (333 mg, 0.07 eq.) were added; repurged for 5 min more and the mixture stirred at 60 °C overnight. After completion of reaction (monitored by TLC & LCMS), the mixture was allowed to cool to RT and then diluted with water (100 mL) and extracted with EtOAc (3x150 mL). The combined organic layer was washed with (50 mL) brine then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-bromo-2-(1H-indol-4-yl)thiazole pure compound as a yellow colored solid (950 mg).


Step-3: Synthesis of [3-chloro-4-[2-(1H-indol-4-yl) thiazol-4-yl] phenyl]-(1-piperidyl)methanone

[0175] In a 25 mL glass bottle were placed 4-bromo-2-(1H-indol-4-yl)thiazole (250 mg, 0.89 mmol, 1eq.) and [2-chloro-4-(piperidine-1-carbonyl)phenyl]boronic acid (313 mg, 1.16 mmol, 1.3 eq.), and sodium carbonate (238 mg, 2.5 eq. dissolved in water (1.0 mL) in DMF (7 mL) and purged with nitrogen gas for 5 min. After adding tetrakis (103 mg, 0.089 mmol, 0.1 eq.), the mixture was repurged for 2 min and heated to 70 °C for 4h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc (3x100 mL). The combined organic layer was washed with water (4x50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude viscous compound, which was purified by reverse phase HPLC to afford [3-chloro-4-[2-(1H-indol-4-yl)thiazol-4-yl]phenyl]-(1-piperidyl)methanon (53 mg) as a white solid, the free base.

[0176] ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 11.49 (s, 1H), 8.24 (s, 1H), 8.13 (d, J = 7.9 Hz, 1H), 7.75 – 7.61 (m, 1H), 7.60 (d, J = 5.1Hz, 1H), 7.60 – 7.47 (m, 3H), 7.24 (s, 2H), 3.61 (s, 2H), 1.63 (s, 4H), 1.53 (s, 4H). LCMS: (M+1) 422.1.

Example 5. Preparation of Compound No. 5

Synthesis of 2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1,3-thiazol-2-yl} pyridine



Step-1: Synthesis of [2-chloro-4-(piperidine-1-carbonyl) phenyl] boronic acid

[0177] In a 250 mL flask was placed 4-borono-3-chloro-benzoic acid (1.0 g, 5 mmol, 1 eq.) dissolved in DMF (25 mL), followed by addition of DIPEA (3.6 mL, 20 mmol, 4 eq.) and the mixture was stirred for 5 min. Then HATU (3.8 g, 10 mmol, 2 eq.) and piperidine (1.7 mL, 17.5 mmol, 3.5 eq.) were added and the mixture stirred at RT overnight. After completion of reaction (monitored by TLC and ¹H-NMR/LCMS), the mixture was diluted with 1N aq. HCl (100 mL) and extracted with EtOAc (2x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by silica gel (60-120) column chromatography using acetone: hexane (0-35%) to elute pure compound as a semisolid which solidified in a refrigerator overnight (1.0g).

Step-2: Synthesis of 4-bromo-2-(2-tert-butyl-4-pyridyl)thiazole

[0178] In a 100 mL glass bottle were placed 2,4 dibromothiazole (200 mg, 0.82 mmol, 1 eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (215, 0.82 mmol, 1 eq.) and potassium phosphate (349 mg, 2.5 eq.) dissolved in 10 mL of THF. The mixture was purged with nitrogen for 15 min, then Xantphos (47 mg, 0.1 eq.) and tetrakis (95 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. When the reaction was completed (monitored by TLC & LCMS), the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x150 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-15%) as eluent system to afford 4-bromo-2-(1H-indol-4-yl)thiazole pure compound as a yellow colored solid (200 mg).


Step-3: Synthesis of [4-[2-(2-tert-butyl-4-pyridyl)thiazol-4-yl]-3-chloro-phenyl]-(1-piperidyl)methanone

[0179] In a 25mL glass bottle were placed 4-bromo-2-(2-tert-butyl-4-pyridyl)thiazole (200 mg, 0.67 mmol, 1eq.), [2-chloro-4-(piperidine-1-carbonyl)phenyl]boronic acid (234 mg, 0.87 mmol, 1.3 eq.), and sodium carbonate (180 mg, 2.5 eq. dissolved in water (1.0 mL), in DMF (7 mL). The mixture was purged with nitrogen gas for 5 min. After adding tetrakis (78 mg, 0.067 mmol, 0.1 eq.), the mixture was repurged for 2 min and then heated to 70 °C for 4h. The reaction was monitored by TLC and LCMS. The reaction mixture was allowed to come to RT, diluted with water (50 mL) and extracted with EtOAc (3x100 mL). The combined organic layer was washed with water (4x50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to afford [4-[2-(2-tert-butyl-4-pyridyl)thiazol-4-yl]-3-chloro-phenyl]-(1-piperidyl)methanone (35 mg) as an off-white solid, the free base.

[0180] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.62 (d, J = 5.2 Hz, 1H), 8.22 (s, 1H), 8.11 (d, J = 8.0 Hz, 1H), 8.04 (s, 1H), 7.79 (d, J = 5.1 Hz, 1H), 7.60 (s, 1H), 7.47 (d, J = 8.0 Hz, 1H), 3.73 (m, 2H), 3.43 (m, 2H), 1.72 (s, 4H), 1.59 (m, 2H), 1.44 (s, 9H). LCMS: (M+1) 440.2.

Example 6. Preparation of Compound No. 6

Synthesis of 3-chloro-4-[2-(1H-indol-4-yl)-1,3-thiazol-4-yl]-N,N-dimethylbenzamide



Step-1: Synthesis of [2-chloro-4-(dimethylcarbamoyl)phenyl]boronic acid

[0181] In a 250 mL flask was placed 4-borono-3-chlorobenzoic acid (1.5 g, 7.5mmol, 1eq.) was dissolved in DMF (25 mL), followed by addition of DIPEA (6.9mL, 37.5 mmol, 5 eq.) and resulting mixture was stirred for 5 min then at RT. Then HATU (5.7 g, 15, mmol, 2 eq.) and

dimethyl amine hydrochloride (2.43 g, 30 mmol, 4 eq.) were added and the mixture stirred at RT overnight. After completion of reaction (monitored by TLC and ¹H-NMR/LCMS), the mixture was diluted with 1N aq. HCl (100 mL) and extracted with EtOAc (2x100 mL), then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by silica gel (60-120) column chromatography using acetone: hexane (0-35%) to elute pure compound as a semisolid which solidified in a refrigerator overnight (1.05g).

Step-2: Synthesis of 4-bromo-2-(1H-indol-4-yl)thiazole

[0182] In a 100 mL glass bottle were placed 2,4 dibromothiazole (1 g, 4.11 mmol, 1eq.) along with 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (1 g, 4.11 mmol, 1eq.) and potassium phosphate (2.18 g, 2.5 eq.) in 30 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (167 mg, .07 eq.) and tetrakis (333 mg, .07 eq.) were added to the mixture and repurged for 5 min. Then the reaction mixture was stirred at 60 °C overnight. After completion of reaction (monitored by TLC & LCMS), the mixture was allowed to cool to RT followed by addition of water (100 mL) and the mixture was extracted with EtOAc (3x150 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-bromo-2-(1H-indol-4-yl)thiazole pure compound as a yellow colored solid (950 mg).


Step-3: Synthesis of 3-chloro-4-[2-(1H-indol-4-yl) thiazol-4-yl]-N, N-dimethyl-benzamide

[0183] In a 25 mL glass bottle were placed 4-bromo-2-(1H-indol-4-yl)thiazole (250 mg, 0.89 mmol, 1eq.) and [2-chloro-4-(dimethylcarbamoyl)phenyl]boronic acid (265 mg, 1.16 mmol, 1.3 eq.), sodium carbonate (258 mg, 2.5 eq. dissolved in water (1.0 mL), in DMF (7 mL), and purged with nitrogen gas for 5 min. After adding tetrakis (104 mg, 0.089 mmol, 0.1 eq.), the mixture was repurged for 2 min and heated to 70 °C overnight. The reaction was monitored by TLC and LCMS. The reaction mixture was allowed to come to RT; water (50 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with water (4x50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to afford 3-chloro-4-[2-(1H-indol-4-yl)thiazol-4-yl]-N,N-dimethyl-benzamide (53 mg-as the freebase) as an off yellow solid.

[0184] ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 11.48 (s, 1H), 8.24 (s, 1H), 8.13 (d, J = 8.0 Hz, 1H), 7.72 (d, J = 7.4 Hz, 1H), 7.64 (s, 1H), 7.62 – 7.51 (m, 3H), 7.24 (s, 2H), 3.00 (m, J = 13.7 Hz, 6H). LCMS: (M+1) 382.1.

Example 7. Preparation of Compound No. 7

Synthesis of 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-2-one



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0185] In a 100 mL glass bottle was placed 2,4-dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0186] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chlorobenzoic acid (397 mg, 1.98 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic

layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazin-2-one

[0187] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (125 mg, 0.34 mmol, 1 eq.) was dissolved in DMF (5 mL), followed by addition of DIPEA (0.25 mL, 1.34 mmol, 4 eq.) and HATU (255 mg, 0.68 mmol, 2 equiv), and the resulting mixture was stirred for 10 min at RT and piperazine-2-one (168 mg, 1.68 mmol, 5 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-2-one (25 mg) as an off-white solid, the free base.

[0188] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.49 (d, J = 5.2 Hz, 1H), 7.92 (s, 1H), 7.8 (s, 1H), 7.75 – 7.64 (m, 3H), 7.58 – 7.41 (m, 2H), 4.31 (s, 1H), 4.18 (m, 1H) 3.95 (s, 1H), 3.72 (s, 1H), 3.41 (s, 2H), 1.42 (s, 9H). LCMS: (M+1) 454.1.

Example 8. Preparation of Compound No. 8

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-cyclobutylbenzamide



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butylpyridine

[0189] In a 100 mL glass bottle was placed 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.)

and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0190] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-cyclobutyl-benzamide

[0191] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (200 mg, 0.54 mmol, 1 eq.) was dissolved in DMF (7 mL), followed by addition of DIPEA (0.4 mL, 2.15 mmol, 4 eq.) and HATU (409 mg, 1.07 mmol, 2 equiv), and the resulting mixture was stirred for 10 min at RT and cyclobutyl amine (191 mg, 2.69 mmol, 5 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-cyclobutylbenzamide (13 mg) as an off-white solid, the free base.

[0192] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.3$ Hz, 1H), 8.02 (s, 1H), 7.94 – 7.89 (m, 1H), 7.87 – 7.77 (m, 2H), 7.72 – 7.62 (m, 2H), 7.52 (d, $J = 4.7$ Hz, 1H), 4.61 – 4.47 (m, 1H), 2.37 (s, 2H), 2.14 (p, $J = 9.6$ Hz, 2H), 1.86 – 1.73 (m, 2H), 1.42 (s, 9H). LCMS: (M+1) 425.2.

Example 9. Preparation of Compound No. 9

Synthesis of 3-chloro-N,N-dimethyl-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzamide



Step-1: Synthesis of [2-chloro-4-(dimethylcarbamoyl)phenyl]boronic acid

[0193] In a 250 mL flask was placed 4-borono-3-chlorobenzoic acid (1.5 g, 7.5 mmol, 1eq.) in (25 mL) of DMF, then at RT, DIPEA (6.9 mL, 37.5 mmol, 5 eq.) was added, stirred for 5 min then at RT, HATU (5.7 g, 15 mmol, 2 eq.) and dimethyl amine hydrochloride (2.43 g, 30 mmol, 4 eq.) were added and the mixture stirred at RT overnight. When the reaction was completed (monitored by TLC and ^1H -NMR/LCMS), 1N aq. HCl (100 mL) was added and the mixture extracted with EtOAc (2x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by silica gel (60-120) column chromatography using acetone: hexane (0-35%) to elute pure compound as a semisolid which solidified in a refrigerator overnight (1.05g).

Step-2: Synthesis of 4-bromo-2-[3-(1-piperidinyl)phenyl]thiazole

[0194] In a 100 mL glass bottle were placed 2,4 dibromothiazole (400 mg, 1.64 mmol, 1eq.) along with [2-(1-piperidinyl)-4-pyridyl]boronic acid (340 mg, 1.64 mmol, 1eq.) and potassium phosphate (697 mg, 3.29 mmol, 2.5 eq.) dissolved in 15 mL of THF. Then the mixture was

purged with nitrogen for 15 min then Xantphos (95 mg, 0.1 eq.) and tetrakis (190 mg, 0.1 eq.) were added then repurged nitrogen for 5 min and the mixture stirred at 60 °C overnight. After completion of reaction (monitored by TLC & LCMS), the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL), the combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-bromo-2-(1H-indol-4-yl)thiazole pure compound as a yellow colored solid (325 mg).


Step-3: Synthesis of 3-chloro-N,N-dimethyl-4-[2-[2-(1-piperidyl)-4-pyridyl]thiazol-4-yl]benzamide

[0195] In a 25 mL glass bottle were placed 4-bromo-2-[3-(1-piperidyl)phenyl]thiazole (300 mg, 0.92 mmol, 1eq.) and [2-chloro-4-(dimethylcarbamoyl)phenyl]boronic acid (295 mg, 1.3 mmol, 1.4 eq.), sodium carbonate (246 mg, 2.32 mmol, 2.5 eq. dissolved in water (1.5 mL), in DMF (7 mL), and purged with nitrogen gas for 5 min. After adding tetrakis (107 mg, 0.062 mmol, 0.1 eq.), the mixture was repurged for 2 min and heated to 80 °C overnight. The reaction was monitored by TLC and LCMS. The reaction mixture was allowed to come to RT; water (50 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with water (4x50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to afford 3-chloro-N,N-dimethyl-4-[2-[2-(1-piperidyl)-4-pyridyl]thiazol-4-yl]benzamide (100 mg) as a yellow solid, the free base.

[0196] ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 8.34 (s, 1H), 8.23 (d, J = 5.1 Hz, 1H), 8.05 (d, J = 7.9 Hz, 1H), 7.63 (s, 1H), 7.51 (d, J = 7.9 Hz, 1H), 7.28 (s, 1H), 7.12 (d, J = 5.1 Hz, 1H), 3.60 (t, J = 5.1 Hz, 2H), 2.98 (d, J = 19.1 Hz, 6H), 1.9 (m, 2H), 1.6 (t, J = 11.3 Hz, 6H). LCMS: (M+1) 427.2.

Example 10. Preparation of Compound No. 10

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-4-methylpiperazine



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0197] In a 100 mL glass bottle was placed 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0198] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(4-methylpiperazin-1-yl)methanone

[0199] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (200 mg, 0.54 mmol, 1 eq.) was dissolved in DMF (7 mL), followed by addition of DIPEA (0.4 mL, 2.15 mmol, 4 eq.) and HATU (409 mg, 1.07 mmol, 2 equiv), and the resulting mixture was stirred for 10 min at RT and 4-methyl piperidine (269 mg, 2.69 mmol, 5 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-4-methylpiperazine (20 mg) as an off-white solid, the free base.

[0200] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.49 (d, $J = 5.3$ Hz, 1H), 7.91 (d, $J = 1.4$ Hz, 1H), 7.78 (d, $J = 1.5$ Hz, 1H), 7.71 – 7.58 (m, 3H), 7.52 (dd, $J = 5.3, 1.8$ Hz, 1H), 7.43 (dd, $J = 7.9, 1.7$ Hz, 1H), 3.80 (s, 2H), 3.54 (s, 2H), 2.53 (s, 4H), 2.37 (s, 3H), 1.42 (s, 9H). LCMS: (M+1) 454.1.

Example 11. Preparation of Compound No. 11

Synthesis of 2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]thiophen-2-yl}pyridine



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0201] In a 100 mL glass bottle was placed 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.), 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound that was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0202] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL), and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10) (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.

Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(1-piperidyl)methanone


[0203] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (150 mg, 0.40 mmol, 1 eq.) dissolved in DMF (10 mL), followed by addition of DIPEA (0.37 mL, 1.61 mmol, 5 eq.) and HATU (307 mg, 0.80 mmol, 2 equiv), and the resulting mixture was stirred for 5 min at RT. Piperidine (137 mg, 1.61 mmol, 4 eq.) was added and the mixture was stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced

pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(1-piperidyl)methanone (40 mg) as an off-white solid, the free base.

[0204] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.49 (d, $J = 5.2$ Hz, 1H), 7.91 (s, 1H), 7.77 (s, 1H), 7.71 – 7.61 (m, 2H), 7.59 – 7.45 (m, 2H), 7.41 (d, $J = 8.0$ Hz, 1H), 3.73 (m, 2H), 3.44 (m, 2H), 1.73 (m, 4H), 1.59 (m, 2H), 1.42 (s, 9H). LCMS: (M+1) 439.2.

Example 12. Preparation of Compound No. 12

Synthesis of 2-tert-butyl-4-{4-[2-chloro-4-(3,3-difluoropyrrolidine-1-carbonyl)phenyl] thiophen-2-yl}pyridine



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butylpyridine

[0205] In a 100 mL glass bottle was placed 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butylpyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0206] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.), in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(3,3-difluoropyrrolidin-1-yl)methanone

[0207] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (150 mg, 0.40 mmol, 1 eq.) in DMF (10 mL), followed by addition of DIPEA (0.37 mL, 1.61 mmol, 5 eq.) and HATU (307 mg, 0.80 mmol, 2 equiv), and the resulting mixture was stirred for 5 min at RT and 3,3-difluoropyrrolidine (173 mg, 1.61 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(3,3-difluoropyrrolidin-1-yl)methanone (33 mg) as an off-white solid, the free base.

[0208] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.49 (d, J = 5.3 Hz, 1H), 7.91 (s, 1H), 7.79 (s, 1H), 7.73 (s, 1H), 7.68 (d, J = 10.2 Hz, 2H), 7.54 (dd, J = 16.3, 6.5 Hz, 2H), 3.97 (t, J = 14.1 Hz, 2H), 3.81 (d, J = 7.4 Hz, 2H), 2.49 (s, 2H), 1.42 (s, 9H). LCMS: (M+1) 461.1.

Example 13. Preparation of Compound No. 13

Synthesis of 2-tert-butyl-4-{4-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl] thiophen-2-yl}pyridine



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0209] In a 100 mL glass bottle was placed 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0210] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chlorobenzoic acid (397 mg, 1.98 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic

layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(4,4-difluoro-1-piperidyl)methanone

[0211] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (150 mg, 0.40 mmol, 1 eq.) in DMF (10 mL), followed by addition of DIPEA (0.37 mL, 1.61 mmol, 5 eq.) and HATU (307 mg, 0.80 mmol, 2 equiv), and the resulting mixture was stirred for 5 min at RT and 3,3-difluoropiperidine (196 mg, 1.61 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(4,4-difluoro-1-piperidyl)methanone (50 mg) as an off-white solid, the free base.

[0212] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.49 (d, $J = 5.2$ Hz, 1H), 7.91 (s, 1H), 7.78 (s, 1H), 7.71 – 7.61 (m, 3H), 7.52 (d, $J = 5.2$ Hz, 1H), 7.47 (d, $J = 7.9$ Hz, 1H), 3.87 (t, 2H), 3.62 (t, 2H), 2.08 (t, 4H), 1.42 (s, 9H). LCMS: (M+1) 475.2.

Example 14. Preparation of Compound No. 14

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-cyclopropylbenzamide



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0213] In a 100 mL glass bottle was placed 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0214] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.

Step-3: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-cyclopropyl-benzamide


[0215] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (160 mg, 0.43 mmol, 1 eq.) in DMF (10 mL), followed by addition of DIPEA (0.40 mL, 2.15 mmol, 5 eq.) and HATU (327 mg, 0.86 mmol, 2 equiv), and the resulting mixture was stirred for 5 min at RT and cyclopropyl amine (98 mg, 1.72 mmol, 5 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to

afford 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-cyclopropyl-benzamide (18 mg) as an off-white solid, the free base.

[0216] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.3$ Hz, 1H), 7.97 (s, 1H), 7.91 (s, 1H), 7.81 (d, $J = 9.7$ Hz, 2H), 7.71 – 7.61 (m, 2H), 7.55 – 7.49 (m, 1H), 2.91 – 2.84 (m, 1H), 1.42 (s, 9H), 0.83 (t, $J = 6.7$ Hz, 2H), 0.67 (t, 2H). LCMS: (M+1) 411.1.

Example 15. Preparation of Compound No. 15

Synthesis of N-tert-butyl-4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamide



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butylpyridine

[0217] In a 100 mL glass bottle was placed 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butylpyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0218] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of N-tert-butyl-4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzamide

[0219] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (160 mg, 0.43 mmol, 1 eq.) was dissolved in DMF (10 mL), followed by addition of DIPEA (0.40 mL, 2.15 mmol, 5 eq.) and HATU (327 mg, 0.86 mmol, 2 equiv), and the resulting mixture was stirred for 5 min at RT and isobutyl amine (127 mg, 1.72 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford N-tert-butyl-4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzamide (16 mg) as an off-white solid, the free base.

[0220] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.48 (d, J = 5.2 Hz, 1H), 7.91 (m, 2H), 7.82 – 7.73 (m, 2H), 7.72 – 7.67 (m, 1H), 7.63 (d, J = 8.0 Hz, 1H), 7.55 – 7.47 (m, 1H), 1.47 (s, 9H), 1.42 (s, 9H). LCMS: (M+1) 427.2.

Example 16. Preparation of Compound No. 16

Synthesis of 2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-5-methylthiophen-2-yl}pyridine



Step-1: Synthesis of 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butylpyridine

[0221] In a 100 mL glass bottle was placed 3,5-dibromo-2-methylthiophene (500 mg, 1.95 mmol, 1 eq.) along with (2-tert-butyl-4-pyridyl)boronic acid (350 mg, 1 eq.) and potassium phosphate (1.03 g, 2.5 eq.) in 15 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (90 mg, 0.08 eq.) and tetrakis (180 mg, 0.08 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butylpyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chlorobenzoic acid

[0222] In a 25 mL glass bottle was placed 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butylpyridine (350 mg, 1.13 mmol, 1 eq.) and 4-borono-3-chlorobenzoic acid (292 mg, 1.46 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (298 mg, 2.5 equiv dissolved in water (1.5 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (130 mg, 0.113 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chlorobenzoic acid (350 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-phenyl]-(1-piperidyl)methanone

[0223] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chlorobenzoic acid (200 mg, 0.52 mmol, 1 eq.) was dissolved in DMF (15 mL), followed by addition of DIPEA (0.38 mL, 2.07 mmol, 4 eq.) and HATU (394 mg, 1.03 mmol, 2 equiv), and the resulting mixture was stirred for 10 min at RT and piperidine (221 mg, 2.59 mmol, 5 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford 2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-5-methylthiophen-2-yl}pyridine (34 mg) as an off-white solid, the free base.

[0224] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.49 (d, J = 5.2 Hz, 1H), 7.76 (s, 1H), 7.59 (dd, J = 5.2, 1.6 Hz, 2H), 7.49 (d, J = 7.8 Hz, 1H), 7.45 – 7.35 (m, 2H), 3.60 (s, 2H), 2.34 (s, 3H), 1.71-1.22 (m, 8H), 1.34 (s, 9H). LCMS: (M+1) 453.2.

Example 17. Preparation of Compound No. 17

Synthesis of 4-(3-chloro-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzoyl)morpholine




[0225] To a stirred solution of 3-chloro-4-[2-[2-(1-piperidyl)-4-pyridyl]thiazol-4-yl]benzoic acid (230 mg, 0.576 mmol) in DMF (6 mL) was added DIPEA (0.4 mL, 2.305 mmol) followed by addition of HATU (394 mg, 1.037 mmol) and the mixture stirred at RT for 30 min. Then morpholine (125 mg, 1.441 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain the crude compound which was purified by HPLC to obtain [3-chloro-4-

[2-[2-(1-piperidyl)-4-pyridyl]thiazol-4-yl]phenyl]-morpholino-methanone (70 mg) freebase as a solid.

[0226] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.21 – 8.07 (m, 3H), 7.59 (s, 1H), 7.49 (d, J = 8.0 Hz, 1H), 7.37 (s, 1H), 7.15 (d, J = 5.2 Hz, 1H), 3.77 (m, 4H), 3.71 – 3.59 (m, 4H), 3.51 (m, 2H), 3.34 – 3.28 (m, 4H), 1.69 (m, 4H). LCMS:- 469 (M+1).

Example 18. Preparation of Compound Nos. 18, 18a and 18b

Synthesis of 3-chloro-N-(2-hydroxypropyl)-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzamide




[0227] To a stirred solution of 3-chloro-4-[2-[2-(1-piperidyl)-4-pyridyl]thiazol-4-yl]benzoic acid (230 mg, 0.576 mmol) in DMF (6 mL), was added DIPEA (0.4 mL, 2.305 mmol) followed by addition of HATU (394 mg, 1.037 mmol) and the mixture stirred at RT for 30 min. Then (S)-1-aminopropan-2-ol (108 mg, 1.441 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain the crude compound which was purified by HPLC to obtain 3-chloro-N-(2-hydroxypropyl)-4-[2-[2-(1-piperidyl)-4-pyridyl]thiazol-4-yl]benzamide (34 mg) freebase as a solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0228] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.18 (d, J = 5.3 Hz, 2H), 8.11 (d, J = 8.1 Hz, 1H), 8.04 (s, 1H), 7.89 (d, J = 8.2 Hz, 1H), 7.37 (s, 1H), 7.15 (d, J = 5.3 Hz, 1H), 3.99 (q, J = 6.2 Hz, 1H), 3.62 (t, J = 4.8 Hz, 4H), 3.45 (dd, J = 13.6, 4.7 Hz, 1H), 3.39 – 3.31 (m, 1H), 1.69 (m, 6H), 1.22 (d, J = 6.3 Hz, 3H). LCMS:- 457 (M+1).

Example 19. Preparation of Compound Nos. 19, 19a, and 19b

Synthesis of tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)carbamate




[0229] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chlorobenzoic acid (100 mg, 0.268 mmol) in DMF (5 mL) was added DIPEA (0.18 mL, 1.072 mmol) followed by addition of HATU (183 mg, 0.482 mmol) and the mixture stirred at RT for 30 min. Then tert-butyl N-(3-piperidyl)carbamate (134 mg, 0.670 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain the crude compound which was purified by HPLC to obtain tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)carbamate (15 mg) freebase as a solid. Each of the enantiomers of this racemate were prepared using both the (R) and (S) forms of the carbamate reagent, to give Compound 19a [tert-butyl (R)-(1-(4-(5-(2-(tert-butyl)pyridin-4-yl)thiophen-3-yl)-3-chlorobenzoyl)piperidin-3-yl)carbamate] and Compound 19b [tert-butyl (S)-(1-(4-(5-(2-(tert-butyl)pyridin-4-yl)thiophen-3-yl)-3-chlorobenzoyl)piperidin-3-yl)carbamate].

[0230] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.2$ Hz, 1H), 7.88 (s, 1H), 7.75 (s, 1H), 7.70 – 7.57 (m, 3H), 7.51 (dd, $J = 5.3, 1.8$ Hz, 1H), 7.43 (dd, $J = 7.9, 1.7$ Hz, 1H), 3.87 (m, 1H), 3.56 (m, 3H), 2.01 (d, $J = 19.4$ Hz, 2H), 1.87 (m, 1H), 1.76 (m, 1H), 1.47 (m, 1H), 1.41 (s, 18H). LCMS: - 554 (M+1).

Example 20. Preparation of Compound No. 20

Synthesis of 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-1 λ^6 ,4-thiomorpholine-1,1-dione




[0231] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chlorobenzoic acid (100 mg, 0.268 mmol) in DMF (5 mL), was added DIPEA (0.18 mL, 1.072 mmol) followed by addition of HATU (183 mg, 0.482 mmol) and the mixture stirred at RT for 30 min. Then 1,4-thiazinane 1,1-dioxide (91 mg, 0.670 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL). The organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain the crude compound which was purified by HPLC to obtain 4-[4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl]-1,4-thiomorpholine-1,1-dione (11.5 mg) freebase as a solid.

[0232] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.49 (d, $J = 5.2$ Hz, 1H), 7.90 (s, 1H), 7.79 (d, $J = 1.2$ Hz, 1H), 7.69 (t, $J = 10.2$ Hz, 3H), 7.52 (d, $J = 5.8$ Hz, 2H), 4.08 (m, 4H), 3.25 (m, 4H), 1.42 (s, 9H). LCMS:- 489 (M+1).

Example 21. Preparation of Compound Nos. 21, 21a, and 21b

Synthesis of 1-[4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl]piperidin-3-amine




[0233] A stirred solution of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chlorobenzoyl]-3-piperidyl]carbamate (10 mg, 0.0180 mmol) in ethanolic HCl (3 mL) was stirred at RT for 2h. The reaction was monitored by LCMS. After completion of reaction, the mixture was concentrated under reduced pressure to obtain 1-[4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl]piperidin-3-amine HCl (8 mg) as a solid. The enantiomers were prepared from

chiral HPLC resolution of the racemate to give Compound 21a [(R)-(3-aminopiperidin-1-yl)(4-(5-(2-(tert-butyl)pyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)methanone] and Compound 21b [(S)-(3-aminopiperidin-1-yl)(4-(5-(2-(tert-butyl)pyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)methanone].

[0234] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.60 (d, $J = 6.3$ Hz, 1H), 8.36 (s, 1H), 8.15 (d, $J = 15.3$ Hz, 3H), 7.74 – 7.66 (m, 2H), 7.51 (dd, $J = 7.8, 1.7$ Hz, 1H), 4.43 (m, 2H), 3.64 (m, 1H), 3.39 (m, 2H), 3.34 (m, 2H), 2.18 (m, 1H), 1.74 (d, $J = 15.5$ Hz, 1H), 1.56 (s, 9H). LCMS: - 454(M+1).

Example 22. Preparation of Compound No. 22

Synthesis of 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}morpholine



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0235] In a 100 mL glass bottle was placed 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) in 25 mL of THF. Then the mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. The progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg).

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0236] In a 25 mL glass bottle was placed 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.) in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis triphenylphosphine palladium(0) (176 mg, 0.152 mmol, 0.1 eq.), the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-morpholino-methanone

[0237] In a 100 mL flask was placed 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (125 mg, 0.34 mmol, 1 eq.) was dissolved in DMF (5 mL), followed by addition of DIPEA (0.25 mL, 1.34 mmol, 4 eq.) and HATU (255 mg, 0.68 mmol, 2 equiv), and the resulting mixture was stirred for 10 min at RT and piperidine (146 mg, 1.68 mmol, 5 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}morpholine (50 mg) as the formate salt as a yellow solid.

[0238] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.49 (d, J = 5.2 Hz, 1H), 7.90 (d, J = 1.5 Hz, 1H), 7.78 (d, J = 1.4 Hz, 1H), 7.71 – 7.64 (m, 2H), 7.62 (d, J = 1.6 Hz, 1H), 7.52 (dd, J = 5.2, 1.7 Hz, 1H), 7.44 (dd, J = 7.8, 1.7 Hz, 1H), 3.76 (s, 5H), 3.52 (s, 3H), 1.42 (s, 9H). LCMS: (M+1) 441.0.

Example 23. Preparation of Compound No. 23

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-phenylbenzamide



[0239] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (100 mg, 0.269 mmol) in DMF (5 mL), was added DIPEA (0.2 mL, 1.076 mmol) followed by addition of HATU (204 mg, 0.538 mmol) and stirred at RT for 30 min. Then aniline (125 mg, 1.347 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x50 mL). The organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude compound, which was purified by Reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-phenyl-benzamide (45 mg) freebase, as a solid.

[0240] ^1H NMR (400 MHz, Dms o-d_6) δ (ppm): 10.4 (s, 1H), 8.58 (d, $J = 5.2$ Hz, 1H), 8.20 (s, 1H), 8.10 (s, 1H), 8.00 (m, 2H), 7.8 (m, 3H), 7.65 (s, 1H), 7.5 (s, 1H), 7.38 (m, 2H), 7.10 (m, 1H), 1.41 (s, 9H). LCMS=446.9 ($\text{M}+1$).

Example 24. Preparation of Compound No. 24

Synthesis of 2-tert-butyl-4-{4-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]-1H-imidazol-2-yl}pyridine

[0241] To a solution of (4-(2-(2-tert-butyl-4-pyridyl)-3-(methoxymethyl)-1H-imidazole-5-yl)-3-chloro-phenyl)-(-4,4-difluoro-1-piperidyl) methanone (360 mg, 0.716 mmol) in THF (20 mL) was added 3M HCl solution (10 mL) and heated at 80 °C for 2h. The reaction was monitored by LCMS and when complete, the reaction mixture was basified with saturated sodium bicarbonate solution (30mL) (pH~7) and extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (50mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product, which was purified by reverse phase HPLC to afford (40mg) of 2-tert-butyl-4-{4-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]-1H-imidazol-2-yl}pyridine.

[0242] ^1H NMR (400 MHz, DMSO- d_6): 8.55 (d, $J = 8.5$ Hz, 1H), 8.10 (s, 1H), 7.98 (s, 2H), 7.71 (d, $J = 8.2$ Hz, 1H), 7.45 (s, 1H), 7.38 (d, $J = 3.5$ Hz, 1H), 3.55 (s, 2H), 3.25 (s, 2H), 1.58 (s, 4H), 1.40 (s, 2H), 1.30 (s, 9H). LCMS – ($\text{M}+1$) 459.3.

Example 25. Preparation of Compound No. 25


Synthesis of 2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1H-imidazol-2-yl}pyridine

[0243] To a solution of (4-(2-(2-tert-butyl-4-pyridyl)-3-(methoxymethyl)-1H-imidazole-5-yl)-3-chloro-phenyl)-(1-piperidyl) methanone (400 mg, 0.854 mmol) and THF (20 mL) was added 3M HCl solution (10 mL) and the mixture heated at 80 °C for 2h. The reaction was monitored by LCMS and when complete, the reaction mixture was basified with saturated sodium bicarbonate solution (30mL) (pH~7) and extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (50mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product, which was purified by reverse phase HPLC to afford (30mg) of 2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1H-imidazol-2-yl}pyridine.

[0244] ¹H NMR (400 MHz, DMSO-d₆): 8.55 (d, J = 8.5 Hz, 1H), 8.10(s, 1H), 7.98(s, 2H), 7.71 (d, J = 8.2 Hz, 1H), 7.45(s, 1H), 7.38 (d, J = 3.5 Hz, 1H), 3.55 (s, 2H), 3.25(s, 2H), 1.58(s, 4H), 1.40(s, 2H), 1.30 (s, 9H). LCMS=423.2 (M+1).

Example 26. Preparation of Compound No. 26

Synthesis of 6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-2-methyl-1,2,3,4-tetrahydroisoquinolin-1-one



Step-1: Synthesis of 6-bromo-3,4-dihydro-2H-isoquinolin-1-one

[0245] To a stirred solution of 5-bromoindan-1-one (4 g, 0.0189 mol) in DCM (30 mL) was added methanesulfonic acid (20 mL) and stirred for 2 min at 0 °C under nitrogen atmosphere. Then sodium azide (2.46 g, 0.0379 mol) was added portionwise and stirred at the same temperature for 2 h. The reaction was monitored by TLC and LCMS. After completion of

reaction, the mixture was quenched with 20 % NaOH solution and extracted with DCM (2x100 mL). The combined organic layer was washed with water (2x50 mL) and brine (50 mL). The organic layer dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product, which was purified by chromatography using eluent 40% EtOAc in hexane to obtain 6-bromo-3,4-dihydro-2H-isoquinolin-1-one (3.59 g) as an off-white solid.

Step-2: Synthesis of 6-bromo-2-methyl-3,4-dihydroisoquinolin-1-one

[0246] To a stirred solution of 6-bromo-3,4-dihydro-2H-isoquinolin-1-one (3.5 g, 0.01548 mol) in DMF (20 mL) was added NaH (60% mineral oil) (1.2 g, 0.0309 mol) at 0 °C under nitrogen atmosphere and stirred for 30 min at RT. Then methyl iodide (1.5 mL, 0.023 mol) was added and the reaction mixture was stirred for 2h. The reaction was monitored by TLC and HNMR. After completion of reaction, the mixture was quenched with ice cold water (50 mL) and extracted with EtOAc (2x100 mL). The combined organic layer was washed with water (2x50 mL) and brine (50 mL). The organic layer dried over anhydrous sodium sulfate and concentrated under reduced pressure to 6-bromo-2-methyl-3,4-dihydroisoquinolin-1-one (3.3 gm) as an off-white solid.

Step-3: Synthesis of 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroisoquinolin-1-one

[0247] A mixture of 6-bromo-2-methyl-3,4-dihydroisoquinolin-1-one (1 g, 0.00418 mol), bis(pinacolato)diboron (1.59 g, 0.00627 mol) and potassium acetate (1.2 g, 0.0125 mol) were dissolved in DMF (15 mL) and the mixture was degassed with nitrogen for 15 min. Then, a catalytic amount of PdCl₂(dppf).CH₂Cl₂ (0.17 g, 0.0002 mol) was added and purging with nitrogen continued for 5 min. Then the mixture was heated at 80 °C for 2h. The reaction was monitored by TLC and ¹HNMR. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x100 mL). The combined organic layer was washed with water (2x50 mL) and brine (50 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product, which was purified by chromatography using eluent 40 % EtOAc in hexane to obtain 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroisoquinolin-1-one (890 mg) as a light yellow oil.

Step-4: Synthesis of 6-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-2-methyl-3,4-dihydroisoquinolin-1-one


[0248] To a stirred solution of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (200 mg, 0.675 mmol) and 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroisoquinolin-1-one (291 mg, 1.0135 mmol) in DMF (4 mL) was added a solution of sodium carbonate (179 mg,

1.689 mmol) in water (4 mL). The mixture was degassed with nitrogen for 20 min. Then tetrakis (62.4 mg, 0.054 mmol) was added and purging with nitrogen was continued for 5 min. The reaction mixture was heated at 80 °C overnight. The reaction was monitored by TLC and LCMS. After the completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL). The organic layer washed with brine (20 mL), dried over sodium sulfate and concentrated under reduced pressure to obtain a crude product, which was purified by chromatography using eluent 50 % EtOAc in hexane to obtain the product, which was triturated with n-pentane (10 mL) and dried to obtain 6-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-2-methyl-3,4-dihydroisoquinolin-1-one (95 mg) as the freebase, an off-white solid.

[0249] ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 8.54 (d, J = 5.1 Hz, 1H), 8.38 (s, 1H), 8.15 (s, 1H), 7.91 (d, J = 8.1 Hz, 1H), 7.85 – 7.71 (m, 3H), 7.51 (dd, J = 5.2, 1.7 Hz, 1H), 3.59 (t, J = 6.6 Hz, 2H), 3.05 (d, J = 13.6 Hz, 2H), 3.04 (s, 3H), 1.38 (s, 9H). LCMS=377 (M+1).

Example 27. Preparation of Compound No. 27

Synthesis of tert-butyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl} piperazine-1-carboxylate




[0250] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (700 mg, 1.88 mmol) in DMF (10 mL), was added DIPEA (1.3 mL, 7.52 mmol) followed by addition of HATU (1.286 g, 3.384 mmol) and the mixture stirred at RT for 30 min. Then tert-butyl piperazine-1-carboxylate (1.75 g, 9.411 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x100 mL). The organic layer was washed with water (2x50 mL) and brine (50 mL) and dried over anhydrous sodium sulfate to obtain a crude compound (820 mg) and 80 mg of the crude compound was purified by reverse phase HPLC to obtain tert-butyl 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazine-1-carboxylate (15 mg) freebase as a solid.

[0251] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.3$ Hz, 1H), 7.89 (d, $J = 1.5$ Hz, 1H), 7.70 – 7.59 (m, 4H), 7.54 – 7.40 (m, 2H), 3.73 (m, 2H), 3.48 (m, 6H), 1.47 (s, 9H), 1.41 (s, 9H). LCMS:- 540 (M+1).

Example 28. Preparation of Compound No. 28

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazine




[0252] A solution of tert-butyl 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazine-1-carboxylate (700 mg, 1.296 mmol) in ethanolic HCl (30 mL) was stirred at RT for 2h. Progress of the reaction was monitored by LCMS. After completion of reaction, the mixture was concentrated under reduced pressure to obtain 650 mg of crude compound and 100 mg crude compound was purified by reverse phase HPLC to obtain [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-piperazin-1-yl-methanone (28 mg) as a solid.

[0253] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.49 (d, $J = 5.2$ Hz, 1H), 7.90 (d, $J = 1.4$ Hz, 1H), 7.71 – 7.60 (m, 4H), 7.55 – 7.41 (m, 2H), 3.77 (m, 2H), 3.61 (m, 2H), 3.00 (m, 4H), 1.42 (s, 9H). LCMS:- 440(M+1).

Example 29. Preparation of Compound No. 29

Synthesis of 1-(4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-1-yl)ethan-1-one




[0254] To a stirred solution of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-piperazin-1-yl-methanone (150 mg, 0.3409 mmol) in DMF (5 mL) was added trimethyl amine (0.237 mL, 1.704 mmol) at 0 °C under nitrogen atmosphere. Then, acetyl bromide (76 mg, 0.613 mmol) was added at the same temperature. The reaction mixture was stirred at RT for 1h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL). The combined organic layer

was washed with water (2x25 mL) and brine (20 mL) and dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude compound which was purified by reverse phase HPLC to obtain 1-[4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazin-1-yl]ethanone (46 mg) as the freebase, a white solid.

[0255] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.2$ Hz, 1H), 7.90 (d, $J = 1.4$ Hz, 1H), 7.77 (s, 1H), 7.70 – 7.61 (m, 3H), 7.56 – 7.42 (m, 2H), 3.64 (m, 8H), 2.14 (s, 3H), 1.41 (s, 9H). LCMS :- 482 (M+1).

Example 30. Preparation of Compound No. 30


Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-4-methane sulfonylpiperazine



[0256] To a stirred solution of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-piperazin-1-yl-methanone 150 mg, 0.3409 mmol in DMF (5 mL) was added triethylamine (0.237 mL, 1.704 mmol) at 0 °C under nitrogen atmosphere. Then methane sulphonyl chloride (69 mg, 0.613 mmol) was added at the same temperature. The reaction mixture was stirred at RT for 1h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL). The combined organic layer was washed with water (2x25 mL) and brine (20 mL) and dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude compound which was purified by reverse phase HPLC to obtain [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(4-methylsulfonylpiperazin-1-yl)methanone (46 mg) as the freebase, a white solid.

Example 31. Preparation of Compound Nos. 31, 31a, and 31b

Synthesis of N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)methanesulfonamide



Step 1. Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butylpyridine:

[0257] In a 100 mL glass bottle, 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in THF (25 mL). The mixture was purged with nitrogen for 15 min and then xantphos (145 mg, 0.1 eq.) and tetrakis ((289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the resultant solution was stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butylpyridine as a colorless semi-solid (500 mg) in pure form.

Step 2. Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid:

[0258] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butylpyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.), were charged in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL)

and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (176 mg, 0.152 mmol, 0.1 equiv) the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come at RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.

Step 3. Synthesis of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate:

[0259] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (700 mg, 1.88 mmol, 1 eq.) was dissolved in DMF (10 mL), followed by addition of DIPEA (1.4 mL, 7.54 mmol, 4 eq.) and HATU (1.4 mg, 3.77 mmol, 2 equiv), and the resulting mixture was stirred for 10 min at RT and tert-butyl N-(3-piperidyl)carbamate (1.5 g, 7.54 mmol, 4 eq.) was added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (700 mg) as a crude viscous compound which was used as such for the next step of synthesis.

Step 4. Synthesis of (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone:

[0260] In a 100 mL flask, tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (700 mg, 1.26 mmol, 1 eq.) was charged in DCM (15 mL), the reaction mixture was maintained at 0 °C and trifluoroacetic acid (5 mL) was added dropwise and the mixture stirred at RT for 2.5 h. Progress of reaction was monitored by TLC/LCMS. After completion of reaction, the DCM was evaporated under reduced pressure, the residue basified with saturated aqueous sodium bicarbonate solution and extracted with EtOAc (2x100 mL). The organic layer was washed with brine (100 mL) and dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure to afford (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone as a crude viscous compound (570 mg) which was used as such for the next step of synthesis.


Step 5. Synthesis of N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]methanesulfonamide:


[0261] In a 25 mL flask, (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone, (150 mg, 0.33 mmol, 1 eq.) was suspended in DCM (5 mL), then pyridine (1 mL) was added and the mixture stirred for 10 min at RT. The mixture was maintained at 0 °C condition and mesyl chloride (63 mg, 0.43 mmol, 1.3 eq.) was added dropwise and the mixture stirred for 1h at the same temperature. Progress of the reaction was monitored by TLC & ¹H-NMR/LCMS. After maximum conversion of the starting material into product, the mixture was diluted with (100 mL) water and extracted with EtOAc (3x100 mL), the combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous desired compound which was submitted for reverse phase HPLC for purification to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl] methanesulfonamide as the freebase (10 mg) as an off-white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0262] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.48 (d, J = 5.2 Hz, 1H), 7.91 (s, 1H), 7.78 (s, 1H), 7.72 – 7.60 (m, 3H), 7.52 (d, J = 5.3 Hz, 1H), 7.45 (d, J = 7.9 Hz, 1H), 3.82 (m, 1H), 3.45-3.43 (m, 2H), 3.1-3.04 (m, 1H), 2.92 (s, 3H), 2.10 (m, 1H), 1.92 (m, 2H), 1.63 (m, 2H), 1.42 (s, 9H). LCMS – (M+1) 532.1.

Example 32. Preparation of Compound Nos. 32, 32a, and 32b

Synthesis of N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl] acetamide





Steps 1-4: see Example 31.

[0263] Step 5: In a 25 mL flask, (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone (150 mg, 0.33 mmol, 1 eq.) was suspended in DCM (5 mL), then pyridine (1 mL) was added and the mixture stirred for 10 min at RT. The mixture was maintained at 0 °C condition and acetyl chloride (26 mg, 0.33 mmol, 1 eq.) was added dropwise and the mixture stirred for 1h at the same temperature. Progress of the reaction was monitored by TLC & ¹H-NMR/LCMS. After maximum conversion of the starting material into product, the mixture was diluted with water (100 mL) and extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous desired compound which was submitted for reverse phase HPLC for purification to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]acetamide as the freebase (50 mg) as an off-white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0264] ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 8.50 (d, J = 5.2 Hz, 1H), 8.04 – 7.99 (m, 1H), 7.89 (s, 1H), 7.64 (m, 2H), 7.54 – 7.45 (m, 2H), 7.39 (s, 1H), 4.18 (d, 1H), 3.41 (m, 2H), 3.08 (m, 2H), 2.88 (m, 1H), 1.80 – 1.70 (m, 3H), 1.48 (m, 2H), 1.33 (s, 9H), 1.20 (m, 2H). LCMS – (M+1) 496.2.

Example 33. Preparation of Compound No. 33

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N,N-bis(propan-2-yl)benzamide


[0265] In a 25 mL flask, 4-(5-(2-tert-butyl-4-pyridyl)-3-thienyl)-3-chloro-benzoic acid (100 mg, 0.26 mmol, 1 eq.) was suspended in DMF (5 mL) then to it was added diisopropylethylamine (139.8 mg, 1.076 mmol) and HATU (204 mg, 0.53 mmol) and the mixture stirred for 10 min at RT. Then added N-isopropylpropan-2-amine (136 mg, 1.347 mmol) and the mixture stirred for 12h at RT. Progress of the reaction was monitored by TLC & ¹H-NMR/LCMS, the reaction mixture was diluted with water (100 mL) and extracted with EtOAc (3x100 mL), the combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous desired compound which was submitted

for reverse phase HPLC for purification to afford 4-(5-(2-tert-butyl-4-pyridyl)-3-thienyl)-3-chloro-N,N-diisopropyl-benzamide as the freebase (12 mg) as an off-white solid.

[0266] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.50 (d, J = 5.2 Hz, 1H), 8.04 – 7.99 (m, 1H), 7.89 (s, 1H), 7.64 (m, 2H), 7.54 – 7.45 (m, 2H), 7.39 (s, 1H), 3.61 (m, 2H), 1.35 (s, 9H), 1.21 (s, 12H). LCMS – (M+1) 455.1.

Example 34. Preparation of Compound Nos. 34, 34a, and 34b

Synthesis of 1-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)pyrrolidin-2-one




[0267] To a stirred solution of (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone (100 mg, 0.204 mmol) in DMF (5 mL) was added NaH (60%) (16 mg, 0.408 mmol) at 0 °C under nitrogen atmosphere and the mixture stirred for 30 min at RT for 30 min. Then 4-chlorobutanoyl chloride (43 mg, 0.306 mmol) was added and the reaction mixture was heated at 70 °C for 2h. The reaction was monitored by LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by reverse phase HPLC to obtain 1-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]pyrrolidin-2-one (22 mg) freebase, an off-white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0268] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, J = 5.3 Hz, 1H), 7.90 (d, J = 1.5 Hz, 1H), 7.77 (d, J = 1.4 Hz, 1H), 7.71 – 7.57 (m, 3H), 7.51 (dd, J = 5.3, 1.7 Hz, 1H), 7.45 (s, 1H), 4.57 (m, 1H), 4.00 (m, 1H), 3.70 (m, 1H), 3.52 (m, 2H), 3.07 (m, 1H), 2.84 (m, 1H), 2.40 (m, 1H), 2.33 (m, 1H), 2.01 (m, 3H), 1.95 – 1.79 (m, 2H), 1.67 (m, 1H), 1.41 (s, 9H). LCMS:- 522 (M+1).

Example 35. Preparation of Compound No. 35

Synthesis of 6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-2-(propan-2-yl)-1,2,3,4-tetrahydroisoquinolin-1-one



Step-1. Synthesis of 6-bromo-3,4-dihydro-2H-isoquinolin-1-one:

[0269] To a stirred solution of 5-bromoindan-1-one (4 g, 0.0189 mol) in DCM (30 mL) was added methane sulfonic acid (20 mL) and the mixture stirred for 2 min at 0 °C under nitrogen atmosphere. Then sodium azide (2.46 g, 0.0379 mol) was added portionwise and the mixture stirred at the same temperature for 2h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was quenched with 20 % NaOH solution and extracted with DCM (2x100 mL). The combined organic layer was washed with water (2x50 mL) and brine (50 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by chromatography using eluent 40 % EtOAc in hexane to obtain 6-bromo-3,4-dihydro-2H-isoquinolin-1-one (3.59 g) as an off-white solid.

Step-2. Synthesis of 6-bromo-2-isopropyl-3,4-dihydroisoquinolin-1-one:

[0270] To a stirred solution of 6-bromo-3,4-dihydro-2H-isoquinolin-1-one (1 g, 0.0044 mol) in DMF (20 mL) was added NaH (60% mineral oil) (354 mg, 0.0088 mol) at 0 °C under nitrogen atmosphere and the mixture stirred for 30 min at RT. Then isopropyl iodide (1.12 g, 0.0066 mol) was added and The reaction mixture was stirred at 70 °C for 2h. The reaction was monitored by TLC and HNMR. After completion of reaction, the mixture was quenched with ice cold water (50 mL) and extracted with EtOAc (2x100 mL). The combined organic layer was washed with water (2x50 mL) and brine (50 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to 66-bromo-2-isopropyl-3,4-dihydroisoquinolin-1-one (1.2 g) as a light brown solid.

Step-3. Synthesis of 2-isopropyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroisoquinolin-1-one:

[0271] A mixture of 6-bromo-2-methyl-3,4-dihydroisoquinolin-1-one (1.2 g, 0.0044 mol), bis(pinacolato)diboron (1.704 g, 0.0067 mol) and potassium acetate (1.293 g, 0.0132 mol) were dissolved in DMF (15 mL) and the mixture was degassed with nitrogen for 15 min. Then, a catalytic amount of PdCl₂(dppf).CH₂Cl₂ (0.179 g, 0.00022 mol) was added and the mixture purged further with nitrogen for 5 min. Then the mixture was heated at 80 °C for 2h. The reaction was monitored by TLC and HNMR. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x100 mL). The combined organic layer was washed with water (2x50 mL) and brine (50 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by chromatography using 40 % EtOAc in hexane to obtain 2-isopropyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroisoquinolin-1-one (600 mg) as a light yellow oil.


Step-4. Synthesis of 6-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-2-isopropyl-3,4-dihydroisoquinolin-1-one:

[0272] To a stirred solution of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (281 mg, 0.996 mmol) and 2-isopropyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroisoquinolin-1-one (200 mg, 0.644 mmol) in DMF (5 mL) was added a solution of sodium carbonate (140 mg, 1.228 mmol) in water (5 mL). The mixture was degassed nitrogen for 20 min. Then tetrakis (62.4 mg, 0.053 mmol) was added and purging with nitrogen was continued for 5 min. The reaction mixture was heated at 80 °C overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL) and organic layer washed with brine (20 mL) and dried over sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by chromatography using 50 % EtOAc in hexane to obtain the product, which was triturated n-pentane (10 mL) to obtain 6-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-2-isopropyl-3,4-dihydroisoquinolin-1-one (20 mg) as the freebase, an off-white solid.

[0273] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.47 (d, J = 5.3 Hz, 1H), 8.10 (s, 1H), 7.98 (d, J = 7.9 Hz, 1H), 7.88 (s, 1H), 7.72 (d, J = 8.8 Hz, 2H), 7.64 (s, 1H), 7.52 (d, J = 5.3 Hz, 1H), 4.98 (p, J = 6.7, 6.3 Hz, 1H), 3.52 (t, J = 6.5 Hz, 2H), 3.02 (t, J = 6.5 Hz, 2H), 1.42 (s, 9H), 1.24 (d, J = 6.9 Hz, 6H). LCMS:- 405 (M+1).

Example 36. Preparation of Compound Nos. 36, 36a, and 36b

Synthesis of 2-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-1λ⁶,2-thiazolidine-1,1-dione




[0274] To a stirred solution of (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone (100 mg, 0.204 mmol) in DMF (5 mL) was added NaH (60%) (16 mg, 0.408 mmol) at 0 °C under nitrogen atmosphere and the mixture stirred for 30 min at RT for 30 min. Then addition of 3-chloropropane-1-sulfonyl chloride (54 mg, 0.306 mmol) and the reaction mixture was heated at 70 °C for 2h. The reaction was monitored by LCMS. After completion of reaction, the reaction mixture was diluted with water (10 mL) and extracted with EtOAc (2x100 mL) combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by reverse phase HPLC to obtain [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-(1,1-dioxo-1,2-thiazolidin-2-yl)-1-piperidyl]methanone (12 mg) freebase, an off-white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0275] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.48 (d, J = 5.3 Hz, 1H), 7.91 (d, J = 1.4 Hz, 1H), 7.78 (d, J = 1.4 Hz, 1H), 7.72 – 7.57 (m, 3H), 7.52 (dd, J = 5.2, 1.7 Hz, 1H), 7.44 (d, J = 7.8 Hz, 1H), 4.57 (m, 1H), 3.64 (m, 3H), 3.25 (d, J = 12.7 Hz, 3H), 2.84 (m, 2H), 2.09 (d, J = 12.5 Hz, 2H), 1.93 (m, 2H), 1.62 (m, 2H), 1.42 (s, 9H). LCMS:- 558 (M+1).

Example 37. Preparation of Compound No. 37

Synthesis of tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)carbamate




[0276] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoic acid (200 mg, 0.538 mmol) in DMF (10 mL), was added DIPEA (0.36 mL, 2.152 mmol) followed by addition of HATU (408 mg, 1.076 mmol) and the mixture stirred at RT for 30 min. Then tert-

butyl N-(4-piperidyl) carbamate (409 mg, 2.04 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-4-piperidyl]carbamate (15 mg) freebase as a solid.

[0277] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.2$ Hz, 1H), 7.88 (s, 1H), 7.75 (s, 1H), 7.70 – 7.57 (m, 3H), 7.51 (dd, $J = 5.3, 1.8$ Hz, 1H), 7.43 (dd, $J = 7.9, 1.7$ Hz, 1H), 3.87 (m, 1H), 3.56 (m, 3H), 2.01 (d, $J = 19.4$ Hz, 2H), 1.87 (m, 1H), 1.76 (m, 1H), 1.47 (m, 1H), 1.41 (s, 18H). LCMS:- 554 (M+1).

Example 38. Preparation of Compound No. 38

Synthesis of N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl) acetamide




[0278] To a solution of (4-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chlorophenyl] methanone (100 mg, 0.2208 mmol) in DCM (5 mL), was added pyridine (1 mL) and the mixture stirred for 15 min at 0 °C. Acetyl chloride (17.2 mg, 0.2208 mmol) was added and the mixture stirred for 1h at RT. The reaction mixture was monitored by TLC and LCMS. The reaction mixture was diluted with water (20 mL) and extracted with EtOAc (3x50 mL). The combined organic layers were concentrated under reduced pressure to obtain a crude product which was purified by reverse phase HPLC to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-4-piperidyl]acetamide (24 mg) as a white solid.

[0279] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.54 (d, $J = 5.1$ Hz, 1H), 8.07 (d, $J = 1.5$ Hz, 1H), 7.84 (d, $J = 7.5$ Hz, 1H), 7.71 – 7.64 (m, 2H), 7.57 (d, $J = 1.6$ Hz, 1H), 7.52 – 7.38 (m, 2H), 4.30 (s, 1H), 3.82 (s, 3H), 3.56 (s, 1H), 3.19 (s, 2H), 3.02 (s, 2H), 1.80 (s, 3H), 1.36 (s, 9H). LCMS – (M+1) 495.

Example 39. Preparation of Compound No. 39

Synthesis of 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-1-ethylpiperazin-2-one



Step-1. Synthesis of 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazin-2-one:

[0280] 4-[5-(2-tert-Butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (200 mg, 0.53 mmol) was taken in DMF (8 mL), followed by addition of DIPEA (0.37 mL, 2.15 mmol), HATU (409 mg, 1.078 mmol), DMF (8 mL), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5 h, followed by addition of piperazin-2-one (215 mg, 2.15 mmol). The reaction mixture was stirred at RT for 16h. After completion of reaction, water (50 mL) was added and the mixture extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl] piperazin-2-one crude (500 mg) as a brown oil.


Step-2. Synthesis of 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-1-ethyl-piperazin-2-one:

[0281] 4-[4-[5-(2-tert-Butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl] piperazin-2-one (300 mg, 0.66 mmol) was taken in dry DMF (5 mL) under nitrogen atmosphere and to it was added NaH (60%) (52 mg, 1.32 mmol) at 0 °C, then the reaction mixture was stirred at RT for 15 min. Ethyl iodide (0.05 mL, 0.66 mmol) was added at 0 °C, then the reaction mixture was stirred at RT for 2h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was quenched with ice cold water (10 mL) and extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (2x25 mL) and finally brine (2x25 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to obtain 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-1-ethyl-piperazin-2-one (37 mg) as the freebase, an off-white solid.

[0282] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.54 (d, $J = 5.1$ Hz, 1H), 8.08 (s, 1H), 7.97 (d, $J = 1.5$ Hz, 1H), 7.70 (d, $J = 8.0$ Hz, 1H), 7.68 (s, 2H), 7.55 – 7.46 (m, 2H), 4.14 (m, 2H), 3.62 (m, 2H), 3.42 – 3.32 (m, 4H), 1.36 (s, 9H), 1.06 (t, $J = 7.1$ Hz, 3H). LCMS:- (M+1) 482.1.

Example 40. Preparation of Compound No. 40

Synthesis of 2-tert-butyl-4-{5-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]thiophen-3-yl}pyridine



Step-1. Synthesis of (4-(4-bromothiophen-2-yl)-3-chlorophenyl)(4,4-difluoropiperidin-1-yl)methanone:

[0283] In a 25 mL glass bottle [2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]boronic acid (200 mg, 0.6601 mmol), 2,4-dibromothiophene (189.2 mg, 0.7921 mmol), and sodium carbonate (173 mg 1.65 mmol), DMF (7 mL), water (2 mL) were charged and purged with nitrogen gas for 5 min. After adding tetrakis (76 mg 0.066 mmol) and Xantphos (38.2 mg, 0.066 mmol), the reaction mixture was heated to 100 °C. The reaction mixture was stirred overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain (4-(4-bromothiophen-2-yl)-3-chlorophenyl)(4,4-difluoropiperidin-1-yl)methanone (300 mg) as a brown colored crude product.

Step-2. Synthesis of [4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-phenyl]-(4,4-difluoro-1-piperidyl)methanone:


[0284] (4-(4-Bromothiophen-2-yl)-3-chlorophenyl)(4,4-difluoropiperidin-1-yl)methanone (300 mg, 0.7160 mmol), (2-tert-butyl-4-pyridyl)boronic acid (153.9 mg, 0.8592 mmol), and sodium carbonate (188 mg 1.79 mmol), DMF (10 mL), water (1 mL) were charged in a 25 mL glass bottle and purged with nitrogen gas for 5 min. After adding tetrakis (82.7 mg 0.0716 mmol), the reaction mixture was heated to 100 °C. The reaction mixture was stirred overnight. The reaction

was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain the crude product which was purified by reverse phase HPLC to afford (4-(4-bromothiophen-2-yl)-3-chlorophenyl)(4,4-difluoropiperidin-1-yl)methanone (20 mg) as a white solid.

[0285] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.51 (d, J = 5.1 Hz, 1H), 7.99 (s, 1H), 7.88 (s, 1H), 7.65 (s, 1H), 7.45 (m, 2H), 7.34 (t, J = 7.7 Hz, 1H), 7.25 (d, J = 7.3 Hz, 1H), 3.88 (m, 4H), 1.34 (s, 9H), 1.12 (m, 4H). LCMS=475.1 (M+1).

Example 41. Preparation of Compound No. 41

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-methyl-N-phenyl benzamide




[0286] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chlorobenzoic acid (200 mg, 0.538 mmol) in DMF (10 mL), was added DIPEA (0.36 mL, 2.152 mmol) followed by addition of HATU (408 mg, 1.076 mmol) and the mixture stirred at RT for 30 min. Then N-methyl aniline (219 mg, 2.04 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-methyl-N-phenylbenzamide (10 mg) freebase as a solid.

[0287] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.48 (d, J = 5.2 Hz, 1H), 8.0(d, J = 5.2 Hz, 1H), 7.88 (s, 1H), 7.75 (m, 1H), 7.47 (m, 3H), 7.43 (dd, J = 7.9, 1.7 Hz, 1H), 7.20-7.42 (m, 5H), 3.41 (s, 3H), 1.41 (s, 9H). LCMS=461.1 (M+1).

Example 42. Preparation of Compound No. 42

Synthesis of 4-{4-[4-(azetidine-1-carbonyl)-2-chlorophenyl]thiophen-2-yl}-2-tert-butylpyridine



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0288] In a 100 mL glass bottle, 2,4 dibromothiophene (600 mg, 2.5 mmol, 1 eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1 eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in 25 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0289] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (1 g, 2.69 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (700 mg, 3.5 mmol, 1.3 eq.), were charged in DMF (13 mL) followed by addition of sodium carbonate (898 mg, 2.5 equiv dissolved in water (2.0 mL)) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (311 mg, 0.269 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 80 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (95:05), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a solid compound, 3-chloro-4-

[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (700 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of azetidin-1-yl-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone

[0290] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (200 mg, 0.54 mmol, 1 eq.) was dissolved in DMF (7 mL), followed by addition of DIPEA (0.5 mL, 2.76 mmol, 5 eq.) and HATU (410 mg, 1.08 mmol, 2 eq.), and the resulting mixture was stirred for 10 min at RT and azetidine hydrochloride (200 mg, 2.15 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure which was submitted to reverse phase HPLC process for its purification to afford azetidin-1-yl-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone (64 mg-as freebase) as an off-white solid.

[0291] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.3$ Hz, 1H), 7.91 (d, $J = 1.5$ Hz, 1H), 7.82 – 7.76 (m, 2H), 7.71 – 7.59 (m, 3H), 7.52 (dd, $J = 5.3, 1.8$ Hz, 1H), 4.44 (t, $J = 7.8$ Hz, 2H), 4.23 (t, $J = 7.9$ Hz, 2H), 2.39 (tt, $J = 14.3, 7.7$ Hz, 2H), 1.42 (s, 9H). LCMS=411.1 (M+1).

Example 43. Preparation of Compound No. 43

Synthesis of 6-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2-oxa-6-azaspiro[3.3]heptane



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butylpyridine

[0292] In a 100 mL glass bottle 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in 25 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0293] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (1 g, 2.69 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (700 mg, 3.5 mmol, 1.3 eq.), were charged in DMF (13 mL) followed by addition of sodium carbonate (898 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (311 mg, 0.269 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 80 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (95:05), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (700 mg) which was used as such for the next step of synthesis.

Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(6-oxa-2-azaspiro[3.3]heptan-2-yl)methanone


[0294] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (150 mg, 0.40 mmol, 1 eq.) was dissolved in DMF (7 mL), followed by addition of DIPEA (0.4 mL, 2.01 mmol, 5 eq.) and HATU (307 mg, 0.80 mmol, 2 eq.), and the resulting mixture was stirred for 10 min at RT and 6-oxa-2-azaspiro[3.3]heptane oxalate (305 mg, 1.61 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated

under reduced pressure which was submitted to reverse phase HPLC process for its purification to afford [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(6-oxa-2-azaspiro[3.3]heptan-2-yl)methanone (54 mg-as freebase) as an off-white solid.

[0295] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.3$ Hz, 1H), 7.90 (s, 1H), 7.79 (d, $J = 4.3$ Hz, 2H), 7.71 – 7.59 (m, 3H), 7.52 (dt, $J = 5.3, 1.3$ Hz, 1H), 4.83 (d, $J = 2.8$ Hz, 4H), 4.59 (d, 2H), 4.36 (d, 2H), 1.42 (s, 9H). LCMS:- 453.1.

Example 44. Preparation of Compound Nos. 44, 44a, and 44b

Synthesis of ethyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)carbamate




[0296] (3-Amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone (200 mg, 0.451 mmol) was taken in DCM (2 mL) under nitrogen atmosphere. Pyridine (1.0 mL) was added at 0 °C followed by addition of ethylcarbonochloride (71 mg, 0.622 mmol) then The reaction mixture was stirred at RT for 4h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was quenched with water (10 mL) and extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (2x25 mL) dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to obtain ethyl -N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl] carbamate (20 mg) as the freebase, an off-white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0297] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.60 (d, $J = 5.5$ Hz, 1H), 8.24 (s, 1H), 8.07 (s, 1H), 7.84 (s, 1H), 7.74 – 7.54 (m, 2H), 7.47 (d, $J = 9.0$ Hz, 1H), 7.29 (s, 1H), 4.18-3.81 (m, 6H), 3.09-2.98 (m, 4H), 1.41 (s, 9H), 1.23 (t, 3H). LCMS:- (M+1) 526.2.

Example 45. Preparation of Compound No. 45

Synthesis of ethyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazine-1-carboxylate




[0298] To a stirred solution of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-piperazin-1-yl-methanone (150 mg, 0.3409 mmol) in DCM (4mL) and DMF (2 mL), was added triethylamine (68.9 mg, 0.6818 mmol) at 0°C under nitrogen atmosphere. Then ethylchloroformate (55.49 mg, 0.5113 mmol) was added at the same temperature. The reaction mixture was stirred at RT for 2h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (25 mL) and extracted with DCM (2x80 mL). The organic layer was washed with brine solution (50 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by reverse phase chromatography to obtain ethyl 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazine-1-carboxylate (38.87 mg). This was dissolved in ethanolic HCl and the mixture stirred at for 30 min at RT then concentrated under reduced pressure to obtain ethyl 4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazine-1-carboxylate HCl (40 mg), an off-white solid.

[0299] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.61 (d, J = 6.4 Hz, 1H), 8.41 (d, J = 1.4 Hz, 1H), 8.25 – 8.14 (m, 3H), 7.73 – 7.64 (m, 2H), 7.49 (dd, J = 7.8, 1.7 Hz, 1H), 4.16 (q, J = 7.1 Hz, 2H), 3.77 (m, 2H), 3.60 (m, 2H), 3.51 (m, 4H), 1.58 (s, 10H), 1.27 (dd, J = 8.9, 5.1 Hz, 3H). LCMS=512 (M+1).

Example 46. Preparation of Compound No. 46

Synthesis of 1-(4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-1-yl)-2,2-dimethylpropan-1-one




[0300] To a stirred solution of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-piperazin-1-yl-methanone (150 mg, 0.3409 mmol) in DCM (4 mL) and DMF (2 mL). Triethylamine (68.9 mg, 0.6818 mmol) was added at 0°C under nitrogen atmosphere. Then pivaloyl chloride (61.65 mg, 0.5113 mmol) was added at the same temperature. The reaction mixture was stirred at RT for 2h. The reaction was monitored by TLC and LCMS. After

completion of reaction, the mixture was diluted with water (50 mL) and extracted with DCM (2x100 mL). The organic layer was washed with brine solution (50 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by reverse phase chromatography to obtain 1-[4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazin-1-yl]-2,2-dimethyl-propan-1-one (36 mg). This was dissolved in ethanolic HCl and the mixture stirred at for 30 min at RT then concentrated under reduced pressure to obtain 1-[4-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]piperazin-1-yl]-2,2-dimethyl-propan-1-one HCl (38 mg), an off-white solid.

[0301] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.58 (d, $J = 6.2$ Hz, 1H), 8.31 (s, 1H), 8.14 – 8.03 (m, 3H), 7.75 – 7.65 (m, 2H), 7.49 (dd, $J = 7.9, 1.7$ Hz, 1H), 3.80 (m, 4H), 3.71 (m, 2H), 3.53 (m, 2H), 1.54 (s, 9H), 1.30 (s, 9H). LCMS=524(M+1).

Example 47. Preparation of Compound No. 47

Synthesis of N-[2-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorophenyl}-N-ethylformamido)ethyl]acetamide



Step-1: Synthesis of N-[2-(ethyl amino) ethyl] acetamide

[0302] N-(2-Aminoethyl) acetamide (500 mg, 4.90 mmol) and dry DMF (5 mL) were charged in a 50 mL 2 neck flask under nitrogen atmosphere, and to it was added NaH (60%) (392 mg, 9.80 mmol) at 0°C then the reaction mixture was stirred at RT for 15 min. Ethyl iodide (0.29 mL, 4.90 mmol) was added at 0°C then the reaction mixture was stirred at RT for 2h. The reaction was monitored by TLC. After completion of reaction, the mixture was quenched with ice cold water (10 mL) and extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (2x25 mL) and brine (2x25 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain crude -N-[2-(ethyl amino)ethyl]acetamide (500 mg) as a brown oil.


Step-2: N-(2-acetamidoethyl)-4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-ethyl-benzamide

[0303] 4-[5-(2-tert-Butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (150 mg, 0.40 mmol) was taken in DMF (5 mL), followed by addition of DIPEA (0.28 mL, 1.61 mmol), HATU (307 mg, 0.80 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5 h, followed by addition of N-[2-(ethyl amino)ethyl]acetamide (210 mg, 1.61 mmol). The reaction mixture was stirred at RT for 16h. After completion of reaction, water (50 mL) was added and the mixture extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to obtain N-(2-acetamidoethyl)-4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-ethyl-benzamide (12 mg) as the freebase, an off-white solid.

[0304] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.54 (d, J = 5.2 Hz, 1H), 8.06 (d, J = 24.5 Hz, 1H), 7.97 (s, 1H), 7.77 – 7.63 (m, 2H), 7.52(s, 2H), 7.43 (d, J = 8.6 Hz, 1H), 3.66-3.44 (m, 2H), 3.33-3.22 (m, 2H), 3.25-3.12 (m, 2H), 1.84(s, 3H), 1.44 (s, 9 H), 1.19 (t, 3H). LCMS:- (M+1) 485.1.

Example 48. Preparation of Compound Nos. 48, 48a, and 48b

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-ol



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0305] In a 100 mL glass bottle 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.), 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in 25 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT,

water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0306] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (1 g, 2.69 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (700 mg, 3.5 mmol, 1.3 eq.), were charged in DMF (13 mL) followed by addition of sodium carbonate (898 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (311 mg, 0.269 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 80 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (95:05), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (700 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(3-hydroxy-1-piperidyl)methanone

[0307] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (150 mg, 0.40 mmol, 1 eq.) was dissolved in DMF (7 mL), followed by addition of DIPEA (0.4 mL, 2.01 mmol, 5 eq.) and HATU (307 mg, 0.80 mmol, 2 eq.), and the resulting mixture was stirred for 10 min at RT and 3-hydroxypiperidine (163 mg, 1.61 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure which was submitted to reverse phase HPLC process for its purification to afford [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(3-hydroxy-1-piperidyl)methanone (54 mg- as freebase) as an off-white solid then it was treated with ethanolic HCl (1.25 M) to obtain its HCl salt (55 mg) as an off-white solid. The enantiomers were prepared from chiral HPLC resolution of the racemate to give Compound 48a [(R)-(4-(5-(2-(tert-butyl)pyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(3-hydroxypiperidin-1-yl)methanone] and Compound 48b [(S)-(4-(5-(2-(tert-butyl)pyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(3-hydroxypiperidin-1-yl)methanone].

[0308] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.61 (d, $J = 6.3$ Hz, 1H), 8.40 (d, $J = 1.5$ Hz, 1H), 8.24 – 8.13 (m, 3H), 7.66 (t, $J = 11.3$ Hz, 2H), 7.49 (d, $J = 8.8$ Hz, 1H), 4.11 (d, $J = 12.5$ Hz, 1H), 3.79 – 3.73 (s, 2H), 3.64–3.4 (m, 2H), 3.56 (d, $J = 14.1$ Hz, 1H), 1.99 – 1.92 (m, 2H), 1.81–1.61 (m, 1H), 1.58 (s, 9H). LCMS=455.1 (M+1).

Example 49. Preparation of Compound No. 49

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(3-hydroxyphenyl) benzamide



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0309] In a 100 mL glass bottle 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in 25 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0310] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (1 g, 2.69 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (700 mg, 3.5 mmol, 1.3 eq.), were charged in DMF (13

mL) followed by addition of sodium carbonate (898 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (311 mg, 0.269 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 80 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (95:05), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (700 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-(3-hydroxyphenyl)benzamide

[0311] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (250 mg, 0.67 mmol, 1 eq.) was dissolved in DMF (7 mL), followed by addition of DIPEA (0.5 mL, 2.69 mmol, 4 eq.) and HATU (513 mg, 1.34 mmol, 2 eq.), and the resulting mixture was stirred for 10 min at RT and 3-hydroxyaniline (295 mg, 2.69 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure which was submitted to reverse phase HPLC process for its purification to afford 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-(3-hydroxyphenyl)benzamide (110 mg freebase) as an off-white solid then it was treated with ethanolic HCl (1.25 M) to obtain its HCl salt (118 mg) as an off-white solid.

[0312] ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 10.29 (s, 1H), 9.46 (s, 1H), 8.66 (d, J = 5.5 Hz, 1H), 8.44 (s, 1H), 8.25 (d, J = 9.1 Hz, 1H), 8.18 (d, J = 1.8 Hz, 1H), 8.06 – 7.98 (m, 2H), 7.96 (s, 1H), 7.79 (d, J = 8.0 Hz, 1H), 7.37 (t, J = 2.2 Hz, 1H), 7.16 (dt, J = 15.9, 8.0 Hz, 2H), 6.58 – 6.50 (m, 1H), 1.46 (d, J = 5.0 Hz, 9H). LCMS=463.2 (M+1).

Example 50. Preparation of Compound Nos. 50, 50a, 50b, and 50c

Synthesis of (2R,6S)-4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2,6-dimethylmorpholine




[0313] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chlorobenzoic acid (100 mg, 0.269 mmol) in DMF 5 mL, was added DIPEA (34.8 mg, 0.2695 mmol) followed by addition of HATU (204 mg, 0.539 mmol) and the mixture stirred at RT for 30 min. Then (2R, 6S)-2,6-dimethyl morpholine (124 mg, 1.07 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chlorophenyl-((2R,6S)-2,6-dimethylmorpholin-4-yl) methanone (25 mg) as the HCl salt. The other diastereomers can be prepared using appropriate stereoisomers of the 2,6-dimethylmorpholine reagent.

[0314] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.45 (d, $J = 5.2$ Hz, 1H), 7.95(s, 1H), 7.78(s, 1H), 7.65(m, 2H), 7.61(s, 1H), 7.51(d, $J = 5.2$ Hz, 1H), 7.40 (d, $J = 5.2$ Hz, 1H), 4.5 (s, 1H), 3.60 (s, 3H), 2.96 (s, 1H), 2.60 (s, 1H), 1.41 (s, 9H), 1.20 (s, 3H), 1.08 (s, 3H). LCMS=469.1 ($M+1$).

Example 51. Preparation of Compound No. 51

Synthesis of 1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-1,2,3,4-tetrahydro-1,8-naphthyridine




[0315] To a stirred solution of 4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chlorobenzoic acid (300 mg, 0.80 mmol) in DMF (5 mL), were added DIPEA (0.6 mL, 3.22 mmol) and HATU (612 mg, 1.61 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5h, followed by addition of 1,2,3,4-tetrahydro-1,8-naphthyridine (324 mg, 1.61 mmol). The reaction mixture was stirred at RT for 16h. After completion of reaction, water (100 mL) was added and the mixture extracted with EtOAc (3x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to

obtain a residue, which was purified by Reverse phase HPLC to obtain [4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-phenyl]-(3,4-dihydro-2H-1,8-naphthyridin-1-yl)methanone (12 mg) as the HCl salt.

[0316] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.68 (s, 1H), 8.63 (d, $J = 6.4$ Hz, 1H), 8.32 (s, 1H), 8.25 (d, $J = 6.1$ Hz, 1H), 8.16 (s, 1H), 8.03 (d, $J = 5.4$ Hz, 1H), 7.93 (d, $J = 7.5$ Hz, 1H), 7.73 (d, $J = 8.1$ Hz, 1H), 7.66 (s, 1H), 7.43 (d, $J = 8.1$ Hz, 1H), 7.33 – 7.25 (m, 1H), 4.02 (t, $J = 6.1$ Hz, 2H), 3.02 (t, $J = 6.5$ Hz, 2H), 2.12 (q, $J = 6.3$ Hz, 2H), 1.58 (s, 9H). LCMS = (M+1) 488.2.

Example 52. Preparation of Compound No. 52

Synthesis of 6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-7-chloro-2-(propan-2-yl)-1,2,3,4-tetrahydroisoquinolin-1-one



Step-1: Synthesis of 5-bromo-6-nitro-indan-1-one

[0317] Fuming nitric acid (25 mL) was cooled to -15°C and 5-bromoindan-1-one (3.5 g, 0.0165 mol) was then added portionwise. The reaction mixture was stirred at -10°C under nitrogen atmosphere for 4h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was poured into ice cold water (100 mL) and extracted with EtOAc (2x200 mL), the organic layer was washed with aq. sodium bicarbonate solution (2x100 mL) and finally brine solution washed (50 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude compound that was dissolved in DCM (15

mL) and n-pentane (150 mL) was added. The precipitated compound was filtered and dried under vacuum to obtain 5-bromo-6-nitro-indan-1-one (2.8 g) as a light brown solid.

Step-2: Synthesis of 6-amino-5-bromo-indan-1-one

[0318] A mixture of 5-bromo-6-nitro-indan-1-one (2.8 g, 0.0109 mol), iron (1.42 g, 0.0546 mol) and NH_4Cl (0.877 g, 0.0164) in EtOH (20 mL) and water (5 mL) was heated to reflux for 90 min. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was filtered through a celite bed, the celite bed washed with hot ethanol (2x25 mL) and the filtrate concentrated under reduced pressure to obtain a crude compound that was diluted with water (50 mL) and extracted with EtOAc (2x200 mL). The extracts were dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain 6-amino-5-bromo-indan-1-one (1.7 g) as a brown solid.

Step-3: Synthesis of 5-bromo-6-chloro-indan-1-one

[0319] To a stirred solution of 6-amino-5-bromo-indan-1-one (500 mg, 2.21 mmol) in conc. HCl (5 mL) at 0 °C under nitrogen atmosphere, was added portionwise sodium nitrite (168 mg, 2.43 mmol) at the same temperature for 30 min, then the reaction mixture was brought to RT and the mixture stirred for 30 min. A solution of copper (I) chloride (307 mg, 3.09 mmol) in conc. HCl (3 mL) was added slowly. The reaction mixture was heated at 60 °C for 50 min. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was cooled to RT and diluted with ice cold water (50 mL) and extracted with EtOAc (2x100 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain 5-bromo-6-chloro-indan-1-one (370 mg).

Step-4: Synthesis of 6-bromo-7-chloro-3,4-dihydro-2H-isoquinolin-1-one

[0320] To a stirred suspension of 5-bromo-6-chloro-indan-1-one (370 mg, 1.507 mmol) in DCM (10 mL) was added methanesulfonic acid (3 mL) at 0 °C under nitrogen atmosphere. Sodium azide (196 mg, 3.014 mmol) was added portionwise and the reaction mixture was stirred at the same temperature for 30 min. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was basified with using 20 % NaOH solution and extracted with EtOAc (12x50 mL). The organic layer was washed with water (2x50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by combi-flash chromatography using EtOAc in hexane to obtain 6-bromo-7-chloro-3,4-dihydro-2H-isoquinolin-1-one (200 mg).

Step-5: Synthesis of 6-bromo-7-chloro-2-isopropyl-3,4-dihydroisoquinolin-1-one

[0321] To a stirred solution of 6-bromo-7-chloro-3,4-dihydro-2H-isoquinolin-1-one (400 mg, 1.538 mmol) in DMF (10 mL) was added NaH (60%) (123 mg, 3.076 mmol) at 0 °C under nitrogen atmosphere and the reaction mixture was stirred at RT for 30 min. Then isopropyl iodide (392 mg, 2.307 mmol) at RT. The reaction mixture was heated at 70 °C for 2h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was cooled to RT and quenched with ammonium chloride solution and extracted with EtOAc (2x100 mL). The organic layer was washed with water (2x50 mL) and brine (50 mL) and dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain 6-bromo-7-chloro-2-isopropyl-3,4-dihydroisoquinolin-1-one (300 mg).

Step-6: Synthesis of 2-tert-butyl-4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-thienyl]pyridine

[0322] A stirred solution of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (1 g, 3.37 mmol), boron bis-pinacolato ester (1.28 g, 5.06 mmol) and potassium tert-butoxide (995 mg, 10.13 mmol) in anhydrous 1,4-dioxane (20 mL), was purged with nitrogen for 15 min and Bis(triphenyl phosphine)palladium(II) dichloride (118 mg, 0.16 mmol) was added and purging continued with nitrogen for 5 min. The reaction mixture was stirred at RT for 10 min and then stirred at 70 °C for 4 h. The reaction was monitored by TLC/¹H-NMR & LCMS. After completion of reaction, the mixture was cooled to RT and diluted with water (100 mL) and extracted with EtOAc (3x100 mL) then washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by silica gel (60-120) column chromatography using EtOAc/ Hexane (0-20%) to elute to obtain 2-tert-butyl-4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-thienyl]pyridine pure compound as a light yellow highly viscous (900 mg) oil.

Step-7: Synthesis of 6-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-7-chloro-2-isopropyl-3,4-dihydroisoquinolin-1-one

[0323] To a solution of 6-bromo-7-chloro-2-isopropyl-3,4-dihydroisoquinolin-1-one (300 mg, 0.991 mmol) and 2-tert-butyl-4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-thienyl]pyridine (408 mg, 1.189 mmol) in DMF (5mL) was added a solution of sodium carbonate (263 mg, 2.478 mmol) in water (5 mL), and purged with nitrogen for 20 min. tetrakis(triphenyl phosphine)palladium (91.6 mg, 0.079 mmol) was added and purging continued with nitrogen for 5 min. The reaction mixture was heated in a reagent bottle at 80 °C overnight. The reaction was monitored by LCMS. After completion of reaction, the mixture was acidified with 6M HCl and extracted with diethyl ether (2x20 mL). The aq layer was basified with saturated sodium carbonate and extracted with EtOAc (2x100 mL). The organic layer was dried

over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by chromatography to obtain pure compound (110 mg). This was dissolved in ethanolic HCl (10 mL) and concentrated under reduced pressure to obtain 6-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-7-chloro-2-isopropyl-3,4-dihydroisoquinolin-1-one HCl salt (120 mg) as a light brown solid.

[0324] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.60 (d, $J = 6.4$ Hz, 1H), 8.41 (d, $J = 1.5$ Hz, 1H), 8.28 – 8.13 (m, 3H), 8.05 (s, 1H), 7.53 (s, 1H), 4.97 (p, $J = 6.8$ Hz, 1H), 3.56 (t, $J = 6.5$ Hz, 2H), 3.03 (t, $J = 6.5$ Hz, 2H), 1.57 (s, 9H), 1.26 (d, $J = 6.8$ Hz, 6H). LCMS=439 (M+1).

Example 53. Preparation of Compound No. 53


Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-[(4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl]benzamide

[0325] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (300 mg, 0.808 mmol) in DMF 10 mL, was added DIPEA (0.6 mL, 3.23 mmol) followed by addition of HATU (614 mg, 1.616 mmol) and stirred at RT for 30 min. Then 3-(amino methyl)-4,6-dimethyl-1H-pyridin-2-one (607 mg, 3.23 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude compound. This was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-[(4,6-dimethyl-2-oxo-1H-pyridin-3-yl)methyl]benzamide (90 mg) freebase as a solid.

[0326] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.48 (d, $J = 5.2$ Hz, 1H), 8.08 (m, 2H), 7.95 (s, 1H), 7.82 (d, $J = 7.9$, 1H), 7.62 (m, 2H), 7.42 (d, $J = 5.3$, 1H), 5.90 (s, 1H), 4.25 (s, 2H), 2.10 (s, 3H), 2.08 (s, 3H), 1.40 (s, 9H). LCMS=506.0 (M+1).

Example 54. Preparation of Compound No. 54

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(4-fluorophenyl) benzamide




[0327] 4-[5-(2-tert-Butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (300 mg, 0.80 mmol) was taken in DMF (10 mL), followed by addition of DIPEA (0.57mL, 3.22 mmol), HATU (600 mg, 1.61 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5 h, followed by addition of 4-fluoroaniline (358 mg, 3.22 mmol). The reaction mixture was stirred at RT for 16h. After completion of reaction, water (150 mL) was added and the mixture extracted with EtOAc (2x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-(4-fluorophenyl)benzamide (90 mg) as the HCl salt.

[0328] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 10.52 (s, 1H), 8.68 (d, $J = 6.0$ Hz, 1H), 8.49 (s, 1H), 8.29 (s, 1H), 8.22 (d, $J = 1.8$ Hz, 1H), 8.09 – 8.02 (m, 2H), 7.88 – 7.77 (m, 3H), 7.22 (t, $J = 8.9$ Hz, 2H), 1.48 (s, 9H). LCMS=465.2 (M+1).

Example 55. Preparation of Compound Nos. 55, 55a, and 55b

Synthesis of N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-2,2-dimethylpropanamide



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0329] In a 100 mL glass bottle 2,4 dibromothiophene (600 mg, 2.5 mmol, 1 eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1 eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in 25 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0330] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.), were charged in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (176 mg, 0.152 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.

Step-3: Synthesis of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate

[0331] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (700 mg, 1.88 mmol, 1 eq.) was dissolved in DMF (10 mL), followed by addition of DIPEA (1.4 mL, 7.54 mmol, 4 eq.) and HATU (1.4 mg, 3.77 mmol, 2 eq.), and the resulting mixture was stirred for 10 min at RT and tert-butyl N-(3-piperidyl)carbamate (1.5 g, 7.54 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then

dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (700 mg) as a crude viscous compound which was used as such for the next step of synthesis.

Step-4: Synthesis of (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone

[0332] In a 100 mL flask, tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (700 mg, 1.26 mmol, 1 eq.) was charged in DCM (15 mL). The reaction mixture was maintained at 0 °C and trifluoroacetic acid (5 mL) was added dropwise and the mixture stirred at RT for 2.5 h. Progress of reaction was monitored by TLC/LCMS. After completion of reaction, the DCM was evaporated under reduced pressure, and the residue basified with saturated aqueous sodium bicarbonate solution and extracted with EtOAc (2x100 mL). The organic layer was washed with brine (100 mL) and dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure to afford (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone as a crude viscous compound (570 mg) which was used as such for the next step of synthesis.

Step-5: Synthesis of N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]-2,2-dimethyl-propanamide


[0333] In a 25 mL flask, (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone, (150 mg, 0.33 mmol, 1 eq.) was suspended in 5 mL of DCM then TEA (2 eq.) added and the mixture stirred for 10 min at RT. The mixture was maintained at 0 °C condition and pivaloyl chloride (59.4 mg, 0.49 mmol, 1.5 eq.) was added dropwise and the mixture stirred for 1h at the same temperature. Progress of the reaction was monitored by TLC & ¹H-NMR/LCMS. After maximum conversion of the starting material into product, the mixture was diluted with (100 mL) water and extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous desired compound which was submitted for reverse phase HPLC for purification to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]-2,2-dimethyl-propanamide as the freebase, which was treated with ethanolic HCl to afford its (100 mg-HCl Salt) as an off-white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0334] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.60 (d, J = 6.4 Hz, 1H), 8.41 (s, 1H), 8.25 – 8.14 (m, 3H), 7.72 – 7.60 (m, 2H), 7.48 (s, 1H), 4.39 (m, 1H), 3.82 (s, 1H), 3.63 (s, 1H), 3.06 (t,

J = 11.5 Hz, 2H), 1.991.91 (m, 2H), 1.68 (m, 2H), 1.57 (s, 9H), 1.21 (s, 4H), 1.13 (s, 5H). LCMS – (M+1) 538.1.

Example 56. Preparation of Compound Nos. 56, 56a, and 56b

Synthesis of 1-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-3-cyclopropylurea




[0335] To a stirred solution of (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl] methanone TFA salt (230 mg, 0.404 mmol) in DCM: DMF (9:1) (10 mL) was added triethylamine (0.22 mL, 1.6197 mmol) at 0 °C under nitrogen atmosphere and the mixture stirred for 5 min at the same temperature. Then isocyanatocyclopropane (50.4 mg, 0.607 mmol) and the mixture stirred for 30 min at RT. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with DCM (12x50 mL), washed with brine solution (2x50 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product which was purified by trituration with using n-pentane (2x10 mL) and diethyl ether (10 mL) to obtain 1-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]-3-cyclopropyl-urea 120 mg as the freebase. 40 mg of pure compound was dissolved in ethanolic HCl (10 mL) and concentrated under reduced pressure to obtain 1-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]-3-cyclopropyl-urea hydrochloride salt (45 mg) as an off-white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0336] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.58 (d, J = 6.2 Hz, 1H), 8.32 (s, 1H), 8.15 – 8.05 (m, 3H), 7.66 (d, J = 7.8 Hz, 1H), 7.60 (s, 1H), 7.45 (d, J = 8.0 Hz, 1H), 3.82 (m, 1H), 3.69 (m, 2H), 2.42 (m, 1H), 2.00 (m, 1H), 1.85 (m, 1H), 1.65 (m, 3H), 1.55 (s, 9H), 1.36 – 1.27 (m, 2H), 0.66 (m, 2H), 0.46 (m, 1H), 0.40 (m, 1H). LCMS=537 (M+1).

Example 57. Preparation of Compound No. 57

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-ol



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0337] In a 100 mL glass bottle 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in 25 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0338] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (1 g, 2.69 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (700 mg, 3.5 mmol, 1.3 eq.), were charged in DMF (13 mL) followed by addition of sodium carbonate (898 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (311 mg, 0.269 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 80 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (95:05), (3x100 mL). The combined organic layer was dried over anhydrous

sodium sulfate and concentrated under reduced pressure to afford a solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (700 mg) which was used as such for the next step of synthesis.


Step-3: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(4-hydroxy-1-piperidyl)methanone

[0339] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (150 mg, 0.40 mmol, 1 eq.) was dissolved in DMF (7 mL), followed by addition of DIPEA (0.4 mL, 2.01 mmol, 5 eq.) and HATU (307 mg, 0.80 mmol, 2 eq.), and the resulting mixture was stirred for 10 min at RT and 3-hydroxypiperidine (163 mg, 1.61 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure which was submitted to reverse phase HPLC process for its purification to afford [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(4-hydroxy-1-piperidyl)methanone (130 mg-as freebase) as an off-white solid then it was treated with ethanolic HCl (1.25 M) to obtain its HCl salt (140 mg) as an off-white solid.

[0340] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.61 (d, $J = 6.4$ Hz, 1H), 8.41 (d, $J = 1.4$ Hz, 1H), 8.25 – 8.14 (m, 3H), 7.68 (d, $J = 7.9$ Hz, 1H), 7.62 (d, $J = 1.6$ Hz, 1H), 7.46 (dd, $J = 7.9, 1.7$ Hz, 1H), 4.18 (s, 1H), 3.93 (dt, $J = 8.4, 4.4$ Hz, 1H), 3.67 (s, 1H), 3.40 (s, 2H), 1.97 (s, 1H), 1.86 (s, 1H), 1.58 (s, 9H), 1.50 (s, 2H). LCMS=455.1 (M+1).

Example 58. Preparation of Compound No. 58

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(pyridin-3-yl)benzamide




[0341] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (100 mg, 0.269 mmol) in DMF 5 mL, was added DIPEA (34.8 mg, 0.2695 mmol) followed by addition of HATU (204 mg, 0.539 mmol) and the mixture stirred at RT for 30 min. Then pyridine-3-amine (101 mg, 1.07 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion

of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-(3-pyridyl) benzamide (34 mg) as the HCl salt.

[0342] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.91 (d, J = 5.2 Hz, 1H), 8.51 (d, J = 5.2 Hz, 1H), 8.35 (d, J = 5.8 Hz, 1H), 8.31 (d, J = 5.2 Hz, 1H), 8.19 (s, 1H), 7.98 (d, J = 5.2 Hz, 1H), 7.90 (s, 1H), 7.82 (s, 1H), 7.78 (d, J = 5.2 Hz, 1H), 7.70 (s, 1H), 7.59 (d, J = 7.9 Hz, 1H), 7.45 (d, J = 7.9 Hz, 1H), 1.41 (s, 9H). LCMS=448.1 ($M+1$).

Example 59. Preparation of Compound No. 59

Synthesis of N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)methanesulfonamide




[0343] To a solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (200 mg, 0.5391 mmol) in DMF (7 mL), was added DIPEA (278.7 mg, 2.6954 mmol), followed by the addition of HATU (410 mg, 1.078 mmol). The reaction mixture was stirred for 30 min. N-(4-Piperidinyl)methanesulfonamide (480.1 mg, 2.6954 mmol) was added and the reaction mixture was stirred overnight at RT. The reaction was monitored by TLC and LCMS. On completion, the reaction mixture was diluted with water (25 mL) and extracted with EtOAc (3x75 mL). The combined organic layer was washed with brine (50 mL) and water (50 mL), and dried over anhydrous sodium sulfate to obtain a crude product. This compound was purified by reverse phase HPLC to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-4-piperidinyl]methanesulfonamide (25 mg) as a white solid.

[0344] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.54 (d, 1H), 8.08 (s, 1H), 7.96 (s, 1H), 7.68 (dd, 1H), 7.67 (s, 1H), 7.58 (d, 1H), 7.52 – 7.39 (d, 1H), 7.18 (d, 1H), 4.28 (s, 1H), 3.03 (s, 1H), 3.18 (s, 1H), 3.33 (s, 1H), 3.56 (s, 1H), 2.95 (s, 3H), 1.85 (s, 2H), 1.36 (s, 1H), 1.33 (s, 9H), 1.23 (s, 1H). LCMS=532 ($M+1$).

Example 60. Preparation of Compound No. 60

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-methyl-N-(1,3-oxazol-2-yl)benzamide



Step-1: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-oxazol-2-yl-benzamide

[0345] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (200 mg, 0.538 mmol) in DMF (10 mL) was added DIPEA (0.4 mL, 2.152 mmol) followed by addition of HATU (408 mg, 1.076 mmol), and the mixture stirred at RT for 30 min. Then oxazol-2-amine (171.1 mg, 2.044 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x50 mL). The organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product, which was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-oxazol-2-yl-benzamide (100 mg) freebase as a solid.


Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-methyl-N-oxazol-2-yl-benzamide

[0346] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-oxazol-2-yl-benzamide (200 mg, 0.456 mmol) in DMF (10 mL), was added sodium hydride (35 mg, 0.913 mmol) at 0 °C, and stirred for 30min. Methyl iodide (96 mg, 0.6849 mmol) was added slowly dropwise and the mixture stirred for 3h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x50 mL). The organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product, which was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-methyl-N-oxazol-2-yl-benzamide (20 mg) freebase as a solid.

[0347] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.58 (d, $J = 5.2$ Hz, 1H), 8.30(s, 1H), 8.18(s, 1H), 7.98(s, 1H), 7.95 (d, $J = 5.2$ Hz, 1H), 7.80(s, 1H), 7.58 (d, $J = 7.9$, 1H), 7.48 (s, 1H), 7.38 (d, $J = 5.3$, 1H), 7.06(s, 1H), 3.38(s, 3H), 1.40 (s, 9H). LCMS=452.0 (M+1).

Example 61. Preparation of Compound No. 61

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-[2-(morpholin-4-yl)-2-oxoethyl]benzamide




[0348] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (100 mg, 0.269 mmol) in DMF 5 mL, was added DIPEA (34.8 mg, 0.2695 mmol) followed by addition of HATU (204 mg, 0.539 mmol) and the mixture stirred at RT for 30 min. Then 2-amino-1-morpholino-ethanone (101 mg, 1.07 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-(3-pyridyl) benzamide (36 mg) as the HCl salt.

[0349] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.45 (d, $J = 5.2$ Hz, 1H), 8.06 (d, $J = 5.2$ Hz, 1H), 7.96(s, 1H), 7.90(d, $J = 5.2$ Hz, 1H), 7.80 (s, 1H), 7.68 (m, 2H), 7.51 (d, $J = 5.2$ Hz, 1H), 4.25 (s, 2H), 3.65(m, 4H), 3.58 (s, 4H), 1.41 (s, 9H). LCMS=498.1 (M+1).

Example 62. Preparation of Compound No. 62

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-{8-methyl-8-azabicyclo[3.2.1]octan-3-yl}benzamide




[0350] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (200 mg, 0.537 mmol) in DMF (5 mL), was added DIPEA (0.37 mL, 2.15 mmol) followed by addition of HATU (327 mg, 0.860 mmol) and the mixture stirred at RT for 30 min. Then 8-methyl-8-azabicyclo[3.2.1]octan-3-amine (188 mg, 1.34 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x100 mL). The organic layer was washed with water (2x50 mL) and brine (50 mL) and dried over anhydrous sodium sulfate to obtain a crude product (820 mg). 80 mg of the crude was purified by reverse phase HPLC to obtain 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-N-(8-methyl-8-azabicyclo[3.2.1]octan-3-yl)benzamide (20 mg) freebase as a solid.

[0351] ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 8.54 (d, J = 5.1 Hz, 1H), 8.20 (s, 1H), 8.07 (s, 1H), 7.97 (d, J = 10.2 Hz, 1H), 7.82 (d, J = 7.9 Hz, 1H), 7.77 – 7.65 (m, 2H), 7.53 – 7.47 (m, 1H), 2.31 (m, 4H), 2.11 (d, J = 14.0 Hz, 3H), 2.03 (s, 3H), 1.92 (m, 2H), 1.36 (s, 9H).
LCMS=494 (M+1).

Example 63. Preparation of Compound No. 63

Synthesis of ethyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)carbamate




[0352] 4-[5-(2-tert-Butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (100mg, 0.26 mmol) was taken in DMF (5.0 mL), followed by addition of DIPEA (0.2mL, 1.07 mmol), HATU (204 mg, 0.053 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5 h, followed by addition of ethyl N-(4-piperidyl)carbamate (184 mg, 0.10 mmol). The reaction mixture was stirred at RT for 16h. After completion of reaction, water (150 mL) was added and the mixture extracted with EtOAc (2x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to obtain ethyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-4-piperidyl]carbamate (30 mg) as a white solid.

[0353] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.54 (d, $J = 5.1$ Hz, 1H), 8.08 (d, $J = 1.4$ Hz, 1H), 7.68 (dd, $J = 4.6, 3.2$ Hz, 2H), 7.56 (d, $J = 1.6$ Hz, 1H), 7.49 (dd, $J = 5.2, 1.7$ Hz, 1H), 7.41 (dd, $J = 7.9, 1.6$ Hz, 1H), 7.19 (d, $J = 7.9$ Hz, 1H), 4.30 (s, 1H), 3.98 (q, $J = 7.1$ Hz, 2H), 3.58 (s, 2H), 3.18 (s, 2H), 3.00 (s, 2H), 1.83 (s, 1H), 1.76 (s, 1H), 1.36 (s, 9H), 1.16 (t, $J = 7.1$ Hz, 3H). LCMS:- (M+1) 525.9.

Example 64. Preparation of Compound Nos. 64, 64a, and 64b

Synthesis of tert-butyl N-[1-(3-chloro-4-{5-[2-(morpholin-4-yl)pyridin-4-yl]thiophen-3-yl}benzoyl)piperidin-3-yl]carbamate




[0354] 4-[4-(4-Bromo-2-thienyl)-2-pyridyl]morpholine (500 mg, 1.53 mmol), [4-[3-(tert-butoxycarbonylamino)piperidine-1-carbonyl]-2-chloro-phenyl]boronic acid (705 mg, 1.84 mmol), sodium carbonate (407 mg, 3.82 mmol), DMF (10 mL), and water (3 mL) were charged in a 25 mL glass bottle and purged with nitrogen gas for 5 min. After adding tetrakis (176 mg 0.153 mmol), the reaction mixture was heated at 100 °C for 16h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (100 mL). The organic layer was washed with brine (50 mL) and water (50 mL) and dried over anhydrous sodium sulfate to obtain a crude product, which was purified by reverse phase chromatography to afford tert-butyl N-[1-[3-chloro-4-[5-(2-morpholino-4-pyridyl)-3-thienyl]benzoyl]-3-piperidyl]carbamate (56 mg) freebase as a white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0355] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.15 (d, $J = 5.2$ Hz, 1H), 8.03 (s, 1H), 7.91 (s, 1H), 7.64 (d, $J = 8.3$ Hz, 1H), 7.44 (d, $J = 7.7$ Hz, 1H), 7.10 (s, 1H), 7.03 (s, 1H), 7.00 – 6.94 (m, 1H), 3.71 (t, $J = 4.8$ Hz, 4H), 3.52 (t, $J = 4.8$ Hz, 4H), 3.05 (s, 2H), 2.78 (s, 2H), 1.84 (s, 1H), 1.32 (s, 9H), 1.21 (s, 2H), 1.12 (s, 2H). LCMS:- (M+1) 583.0.

Example 65. Preparation of Compound Nos. 65, 65a, and 65b

Synthesis of 1-(3-chloro-4-{5-[2-(morpholin-4-yl)pyridin-4-yl]thiophen-3-yl}benzoyl)piperidin-3-amine





[0356] tert-Butyl-N-[1-[3-chloro-4-[5-(2-morpholino-4-pyridyl)-3-thienyl]benzoyl]-3-piperidyl]carbamate (60 mg, 0.10mmol) was taken in EtOH (4.0 mL) at 0 °C, followed by dropwise addition of a 1.25M HCl in EtOH under nitrogen atmosphere. The mixture was stirred at RT for 16h. Completion of reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was concentrated to get (3-amino-1-piperidyl)-[3-chloro-4-[5-(2-morpholino-4-pyridyl)-3-thienyl]phenyl]methanone as the HCl salt, a light yellow solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0357] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.40 (s, 1H), 8.33 (s, 1H), 8.27 (s, 1H), 8.15 (s, 1H), 8.06 (s, 1H), 7.68 (d, $J = 8.6$ Hz, 1H), 7.52 (s, 1H), 7.29 – 7.23 (m, 1H), 4.19 (s, 1H), 3.78 (s, 8H), 3.35 (s, 1H), 3.24 (s, 2H), 3.15 (s, 1H), 2.03 (s, 1H), 1.75 (s, 2H), 1.53 (s, 1H). LCMS:- (M+1) 483.1.

Example 66. Preparation of Compound Nos. 66, 66a, and 66b

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)-2-methylthiophen-3-yl]-3-chlorobenzoyl} piperidin-3-amine





Step-1: Synthesis of 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butyl-pyridine

[0358] In a 100 mL glass bottle 3,5-dibromo-2-methyl-thiophene (500 mg, 1.95 mmol, 1eq.) along with (2-tert-butyl-4-pyridyl)boronic acid (350 mg, 1eq.) and potassium phosphate (1.03 g, 2.5 eq.) were charged in 15 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (90 mg, 0.08 eq.) and tetrakis (180 mg, 0.08 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoic acid

[0359] In a 25 mL glass bottle, 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butyl-pyridine (350 mg, 1.13 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (292 mg, 1.46 mmol, 1.3 eq.), were charged in DMF (7 mL) followed by addition of sodium carbonate (298 mg, 2.5 equiv dissolved in water (1.5 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (130 mg, 0.113 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoic acid (350 mg) which was used as such for the next step of synthesis.

Step-3: Synthesis of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate

[0360] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoic acid 180 mg, 0.46 mmol, 1 eq.) was dissolved in DMF (8 mL), followed by addition of DIPEA (0.34 mL, 1.87 mmol, 4 eq.) and HATU (355 mg, 0.92 mmol, 2 eq.), and the resulting mixture

was stirred for 10 min at RT and tert-butyl N-(3-piperidyl)carbamate (373 mg, 1.87 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, to afford tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (180 mg) as the freebase.


Step-4: Synthesis of (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-phenyl]methanone


[0361] In a 100 mL flask, tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (150 mg, 0.31 mmol, 1 eq.) was charged in DCM (10 mL), the reaction mixture was maintained at 0 °C and trifluoroacetic acid (4 mL) was added dropwise and the mixture stirred at RT for 2.5 h. Progress of reaction was monitored by TLC/LCMS. After completion of reaction, the DCM was evaporated under reduced pressure, and the residue basified with saturated aqueous sodium bicarbonate solution and extracted with EtOAc (2x100 mL). The organic layer was washed with brine (100 mL) and dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure and submitted to reverse phase HPLC process for its purification to afford (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-phenyl]methanone as a greyish solid freebase (56 mg), which was treated with ethanolic HCl to give its respective HCl salt as a yellow-greyish solid (62 mg). The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0362] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.55 (d, J = 6.3 Hz, 1H), 8.08 (d, J = 6.4 Hz, 3H), 7.71 (s, 1H), 7.53 (s, 2H), 4.44 (m, 1H), 4.10 (m, 1H), 3.62 (d, J = 7.2 Hz, 1H), 3.41 (m, 2H), 2.47 (s, 3H), 2.20 (m, 1H), 1.73 (m, 3H), 1.55 (s, 9H). LCMS:- 468.2(M+1).

Example 67. Preparation of Compound No. 67

Synthesis of 4-[5-(2-tert-butylpyridin-4-yl)-2-methylthiophen-3-yl]-3-chloro-N-phenyl benzamide





Step-1: Synthesis of 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butyl-pyridine

[0363] In a 100 mL glass bottle, 3,5-dibromo-2-methyl-thiophene (500 mg, 1.95 mmol, 1 eq.) along with (2-tert-butyl-4-pyridyl)boronic acid (350 mg, 1 eq.) and potassium phosphate (1.03 g, 2.5 eq.) were charged in 15 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (90 mg, 0.08 eq.) and tetrakis (180 mg, 0.08 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoic acid

[0364] In a 25 mL glass bottle, 4-(4-bromo-5-methyl-2-thienyl)-2-tert-butyl-pyridine (350 mg, 1.13 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (292 mg, 1.46 mmol, 1.3 eq.), were charged in DMF (7 mL) followed by addition of sodium carbonate (298 mg, 2.5 equiv dissolved in water (1.5 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (130 mg, 0.113 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 90 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a semi-solid compound, 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoic acid (350 mg) which was used as such for the next step of synthesis.

Step-3: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-N-phenylbenzamide


[0365] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-benzoic acid (180 mg, 0.46 mmol, 1 eq.) was dissolved in DMF (8 mL), followed by addition of DIPEA (0.34 mL, 1.87 mmol, 4 eq.) and HATU (355 mg, 0.92 mmol, 2 eq.), and the resulting mixture

was stirred for 10 min at RT and aniline (174 mg, 1.87 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to give a viscous compound, which was purified by reverse phase HPLC process to afford 4-[5-(2-tert-butyl-4-pyridyl)-2-methyl-3-thienyl]-3-chloro-N-phenyl-benzamide (56 mg) as the freebase as an off-white solid which was treated with ethanolic HCl to give its respective HCl salt as an off-white solid (61 mg).

[0366] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.54 (d, $J = 6.4$ Hz, 1H), 8.17 (d, $J = 1.8$ Hz, 1H), 8.14 – 8.10 (m, 2H), 8.10 – 8.05 (m, 1H), 8.03 – 7.96 (m, 1H), 7.71 (d, $J = 8.0$ Hz, 2H), 7.57 (d, $J = 7.9$ Hz, 1H), 7.39 (t, $J = 7.8$ Hz, 2H), 7.18 (t, $J = 7.5$ Hz, 1H), 2.48 (s, 3H), 1.55 (s, 9H). LCMS:- (M+1) 461.2.

Example 68. Preparation of Compound Nos. 68a, and 68b

Synthesis of (3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-ol




[0367] 4-[4-(2-tert-Butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoic acid (0.5 g, 1.34 mmol) was taken in DMF (10 mL), followed by addition of DIPEA (0.95 mL, 5.37 mmol), and HATU (1.08 g, 2.68 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5 h, followed by addition of (3S)-pyrrolidin-3-ol (0.467 g, 5.37 mmol). The mixture was stirred at RT for 16h. After completion of reaction, water (100 mL) was added and the mixture extracted with EtOAc (3x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a residue, which was purified by reverse phase HPLC to obtain Compound 68a [4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-phenyl]-[(3S)-3-hydroxypyrrolidin-1-yl]methanone (230 mg) as the HCl salt. Compound 68b, the (3R)-enantiomer, was prepared by using the (3S)-pyrrolidin-3-ol reagent.

[0368] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.62 (d, $J = 5.7$ Hz, 2H), 8.26 (s, 1H), 8.20 – 8.10 (m, 2H), 7.85 – 7.73 (m, 2H), 7.59 (t, $J = 6.8$ Hz, 1H), 4.51 (s, 1H), 4.41 (s, 1H), 3.82 – 3.66

(m, 2H), 3.65 – 3.51 (m, 1H), 3.42 – 3.32 (m, 1H), 2.19 (s, 1H), 2.05 (s, 1H), 1.57 (s, 9H).
LCMS=441.2 (M+1).

Example 69. Preparation of Compound Nos. 69, 69a, and 69b

Synthesis of tert-butyl N-[(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-3-chloro-benzoic acid

[0369] 2,4-Dibromothiophene (10.0 g, 40.84 mmol), 4-borono-3-chloro-benzoic acid (10 mg, 50.2 mmol), sodium carbonate (10.9 g, 104.1 mmol) in water (30 mL), and DMF (100 mL) were charged in a 200 mL glass bottle and purged with nitrogen gas for 10 min. After adding tetrakis (4.8 g, 4.16 mmol), the reaction mixture was heated at 100 °C for 16h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL), filtered and dried with washings of ether to afford a white solid 4-(4-bromo-2-thienyl)-3-chloro-benzoic acid (5.0 g) as the freebase.

Step-2: Synthesis of tert-butyl N-[(3R)-1-[4-(4-bromo-2-thienyl)-3-chloro-benzoyl]-3-piperidyl]carbamate

[0370] 4-(4-Bromo-2-thienyl)-3-chloro-benzoic acid (0.5 g, 1.58 mmol) was taken in DMF (10 mL), followed by addition of DIPEA (3.0 mL, 15.8 mmol), and HATU (2.4 g, 6.3 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5 h, followed by addition of tert-butyl N-[(3R)-3-piperidyl]carbamate (0.63 g, 3.1 mmol). The reaction mixture was stirred at RT for 16h. After completion, water (100 mL) was added and the mixture extracted with EtOAc (3x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain crude tert-butyl N-[(3R)-1-[4-(4-bromo-2-thienyl)-3-chloro-benzoyl]-3-piperidyl]carbamate (400 mg) as the freebase.


Step-3: Synthesis of tert-butyl N-[(3R)-1-[4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-3-piperidyl] carbamate

[0371] tert-Butyl N-[(3R)-1-[4-(4-bromo-2-thienyl)-3-chloro-benzoyl]-3-piperidyl]carbamate (300 mg, 0.70 mmol), 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (200 mg, 0.77 mmol), sodium carbonate (183 mg, 1.75 mmol) in water (2.0 mL), and DMF (8.0 mL) were charged in a 25 mL glass bottle and purged with nitrogen gas for 10 min. After adding tetrakis (80 mg, 0.07 mmol), the reaction mixture was heated at 100 °C for 16h. The reaction was monitored by TLC and LCMS. After completion, water (100 mL) was added and the mixture extracted with EtOAc (3x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a residue that was purified by reverse phase HPLC to obtain Compound 69a [tert-butyl N-[(3R)-1-[4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate] as the freebase, an off-white solid. Compound 69b, the (S)-enantiomer, can be prepared from using the (S)-carbamate reagent in Step-2.

[0372] ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 8.54 (d, J = 5.1 Hz, 1H), 8.37 (d, J = 1.5 Hz, 1H), 8.07 (s, 1H), 7.84 (d, J = 8.0 Hz, 1H), 7.76 (s, 1H), 7.64 (s, 1H), 7.57 (dd, J = 5.2, 1.7 Hz, 1H), 7.46 (d, J = 7.9 Hz, 1H), 7.00 (s, 1H), 4.22 (s, 1H), 3.42 (s, 2H), 3.14 (s, 2H), 2.81 (s, 1H), 1.96 (s, 1H), 1.87 (s, 1H), 1.68 (s, 17H), 1.23 (s, 9H). LCMS:- (M+1) 554.0.

Example 70. Preparation of Compound Nos. 70, 70a, and 70b

Synthesis of tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl} piperidin-3-yl)-N-methylcarbamate




[0373] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (500 mg, 1.347 mmol) in DMF (10 mL), was added DIPEA (0.92 mL, 5.390 mmol) followed by addition of HATU (1.0 g, 2.69 mmol), and the mixture stirred at RT for 30 min. Then tert-butyl N-methyl- N-(3-piperidyl) carbamate (1.1 g, 5.39 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL). The organic layer was washed with water (2x20 mL) and brine (20 mL), and

dried over anhydrous sodium sulfate to obtain a crude product, which was purified by reverse phase HPLC to obtain tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]-N-methyl-carbamate (74 mg) freebase as a solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0374] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.48 (d, $J = 5.2$ Hz, 1H), 7.88 (s, 1H), 7.75 (s, 1H), 7.70 – 7.57 (m, 3H), 7.51 (dd, $J = 5.3, 1.8$ Hz, 1H), 7.43 (dd, $J = 7.9, 1.7$ Hz, 1H), 3.87 (m, 1H), 3.56 (m, 3H), 2.01 (d, $J = 19.4$ Hz, 3H), 1.87 (m, 2H), 1.76 (m, 2H), 1.47 (m, 1H), 1.41 (s, 18H). LCMS=554 (M+1).

Example 71. Preparation of Compound Nos. 71, 71a, and 71b

Synthesis of tert-butyl 3-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamido} piperidine-1-carboxylate




[0375] 4-[5-(2-tert-Butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (300 mg, 0.80 mmol) was taken in DMF (10 mL), followed by addition of DIPEA (0.62 mL, 3.22 mmol), and HATU (611 mg, 1.62 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5 h, followed by addition of tert-butyl 3-aminopiperidine-1-carboxylate (640 mg, 3.22 mmol). The reaction mixture was stirred at RT for 16h. After completion, water (150 mL) was added and the mixture extracted with EtOAc (100 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product, which was purified by reverse phase HPLC to obtain tert-butyl 3-[[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]amino]piperidine-1-carboxylate (31 mg) freebase as a white solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0376] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.54 (d, $J = 5.1$ Hz, 1H), 8.46 (d, $J = 7.3$ Hz, 1H), 8.11 – 8.03 (m, 2H), 7.90 (d, $J = 7.8$ Hz, 1H), 7.76 – 7.65 (m, 2H), 7.51 (dd, $J = 5.1, 1.7$ Hz, 1H), 3.89 (s, 2H), 2.84 (s, 2H), 1.86 (s, 2H), 1.78 (s, 2H), 1.5 (t, $J = 12$ Hz, 1H), 1.36 (s, 9H), 1.26 (s, 9H). LCMS:- (M+1) 554.3.

Example 72. Preparation of Compound No. 72

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-amine



Step-1: Synthesis of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-4-piperidyl]carbamate

[0377] To a solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (300 mg, 0.8086 mmol) in DMF (7 mL), was added DIPEA (418 mg, 3.2344 mmol), followed by HATU (615 mg, 1.6172 mmol), and the reaction mixture was stirred for 30 min at RT. tert-Butyl N-(4-piperidyl)carbamate (809 mg, 4.043 mmol) was added and the reaction mixture was stirred for 16h at RT. The reaction was monitored by TLC and LCMS. On completion, the reaction mixture was diluted with water (25 mL) and extracted with EtOAc (3x75 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL) and dried over anhydrous sodium sulfate to obtain 500 mg of crude product. The crude compound was purified by reverse phase chromatography to afford tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-4-piperidyl]carbamate (250 mg) as a white solid.


Step-2: Synthesis of (4-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone

[0378] To a solution of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-4-piperidyl]carbamate (250 mg, 0.2711 mmol) in EtOH (5 mL), was added 1.2 M ethanolic HCl (5 mL) at 0 °C, and the reaction mixture stirred for 16 h at RT. The reaction was monitored by TLC and LCMS. The mixture was concentrated under reduced pressure to obtain (4-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone (170 mg) as a white solid.

[0379] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.55 (d, J = 6.0 Hz, 1H), 8.33 (s, 1H), 8.23 (s, 1H), 8.03 – 7.95 (m, 2H), 7.67 (d, J = 7.9 Hz, 1H), 7.57 (s, 1H), 7.42 (d, J = 7.8 Hz, 1H), 4.45 (s, 1H), 3.60 (s, 1H), 3.30 (m, 1H), 3.17 (s, 1H), 2.87 (s, 1H), 2.00 (s, 2H), 1.88 (s, 2H), 1.42 (s, 9H). LCMS = (M+1) 454.2.

Example 73. Preparation of Compound No. 73

Synthesis of 3-(N-[4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-4-fluoro-anilino)propanoic acid



Step-1: Synthesis of ethyl 3-(4-fluoroanilino)propanoate

[0380] 4-Fluoroaniline (5.0 g, 44.64 mmol) and ethyl prop-2-enoate (4.8 g, 44.64 mmol) were taken in acetic acid (5.0 mL) in a 25 mL microwave vial, then the reaction was stirred at 200 °C for 20 min in a microwave. Completion of reaction was monitored by TLC and LCMS. After completion of reaction, water (200 mL) was added and the mixture extracted with EtOAc (300 mL). The combined organic layer was washed with brine (100 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product, which was purified by chromatography to obtain ethyl 3-(4-fluoroanilino)propanoate compound (2.5 g), as a brown liquid. LCMS:- (M+1) 212.

Step-2: Synthesis of ethyl 3-(N-[4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-4-fluoro-anilino) propanoate

[0381] 4-[4-(2-tert-Butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoic acid (300 mg, 0.806 mmol) was taken in DCM (5 mL) followed by oxalyl chloride (0.2 mL) at 0 °C and a catalytic amount of DMF. The reaction was stirred at RT for 1 h. After completion of reaction, the mixture was completely evaporated to remove all oxalyl chloride. Separately, ethyl 3-(4-fluoroanilino) propionate prop-2-enoate (188 mg, 0.88 mmol) was taken in DCM (5.0 mL) under nitrogen. Dropwise addition of triethyl amine (0.23 mL, 0.96 mmol) was followed by addition of the above solution of acid chloride in DCM at 0 °C. The reaction mixture was stirred RT for 16 h and the reaction was monitored by TLC and LCMS. After completion of reaction, water (100 mL) was added and the mixture extracted with DCM (150 mL). The combined organic layer was washed with brine (25 mL), dried over anhydrous sodium sulfate and concentrated under reduced

pressure to obtain crude ethyl 3-(N-[4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-4-fluoro-anilino)propanoate (150 mg), a yellow solid. LCMS:- (M+1) 565.3.


Step-3: Synthesis of 3-(N-[4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-4-fluoro-anilino)propanoic acid.

[0382] Ethyl 3-(N-[4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-4-fluoro-anilino)propanoate (150 mg, 0.265 mmol) was taken in THF:Water (4:2 mL), followed by addition of lithium hydroxide (44 mg, 1.06 mmol) under nitrogen atmosphere. The reaction mixture was stirred at RT for 16 h. After completion of reaction, the mixture was completely concentrated and water (30 mL) was added and washed with DCM (2x50 mL). The aqueous layer was separated and cooled to 0°C then acidified slowly with 1N HCl to make up to pH=2 and then extracted with EtOAc (3x50 mL). The organic layer was separated and dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain a crude product, which was purified by HPLC to obtain 3-(N-[4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoyl]-4-fluoro-anilino)propanoic acid (10 mg) as the freebase. LCMS:- (M+1) 537.0.

[0383] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.46 (d, J = 5.3 Hz, 1H), 8.06 (s, 1H), 7.81 (s, 1H), 7.72 (s, 1H), 7.51 (d, J = 4.8 Hz, 3H), 7.27 (s, 3H), 7.08 (t, J = 8.4 Hz, 2H), 4.18 (s, 2H), 2.63 (s, 2H), 1.41 (s, 9H).

Example 74. Preparation of Compound Nos. 74, 74a and 74b

Synthesis of (R)-(3-aminopiperidin-1-yl)(4-(4-(2-(tert-butyl)pyridin-4-yl)thiophen-2-yl)-3-chlorophenyl)methanone




[0384] tert-Butyl (R)-(1-(4-(4-(2-(tert-butyl)pyridin-4-yl)thiophen-2-yl)-3-chlorobenzoyl)piperidin-3-yl)carbamate (135 mg, 0.244mmol) was taken in EtOH (3.0 mL), followed by dropwise addition of 1.25M HCl in EtOH (3.0 mL) under nitrogen atmosphere at 0°C. The reaction mixture was stirred at RT for 16h. Completion of reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was concentrated to get (R)-(3-aminopiperidin-1-yl)(4-(4-(2-(tert-butyl)pyridin-4-yl)thiophen-2-yl)-3-chlorophenyl)methanone (120mg) (Compound No. 74a) as the HCl salt light yellow solid. Compound No. 74b, the (S)-enantiomer was prepared using the opposite enantiomeric starting material.

[0385] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.94 (s, 1H), 8.76 (d, $J = 6.0$ Hz, 1H), 8.44 (s, 1H), 8.30 (s, 2H), 7.91 (d, $J = 7.9$ Hz, 1H), 7.74 (s, 1H), 7.55 (d, $J = 7.8$ Hz, 1H), 4.82 (s, 1H), 4.19 (s, 1H), 3.81 (s, 1H), 3.39 (s, 1H), 3.19 (s, 2H), 2.96 (s, 1H), 2.12 (s, 1H), 1.71 (s, 1H), 1.54 (s, 9H). LCMS:- (M+1) 453.9.

Example 75. Preparation of Compound Nos. 75, 75a, and 75b

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-methylpiperidin-3-amine




[0386] A solution of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]-N-methyl-carbamate (50 mg, 0.0880 mmol) in ethanolic HCl (3 mL) was stirred at RT overnight. The reaction was monitored by LCMS. After completion of reaction, the mixture was concentrated under reduced pressure to obtain [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-(methylamino)-1-piperidyl] methanone (34 mg) as a solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0387] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.60 (d, $J = 6.3$ Hz, 1H), 8.36 (s, 1H), 8.15 (d, $J = 15.3$ Hz, 3H), 7.74 – 7.66 (m, 2H), 7.51 (dd, $J = 7.8, 1.7$ Hz, 1H), 4.3 (s, 1H), 3.64 (m, 2H), 3.38 (m, 2H), 2.80 (s, 3H), 2.22 (s, 1H), 1.80 (m, 2H), 1.74 (s, 1H), 1.56 (s, 9H). LCMS:- 468(M+1).

Example 76. Preparation of Compound No. 76

Synthesis of N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)-2,2-dimethylpropanamide




[0388] To a solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (300 mg, 0.8086 mmol) in DMF (10 mL), was added DIPEA (418.1 mg, 3.234 mmol), followed by the addition of HATU (615 mg, 1.6172 mmol), the reaction mixture was stirred for 30 min. 2,2-Dimethyl-N-(4-piperidyl)propanamide (744.7 mg, 4.0431 mmol) was added and the mixture

stirred for 16h at RT. The reaction was monitored by TLC and LCMS. On completion, the reaction mixture was diluted with water (25 mL) and extracted with EtOAc (3x75 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL) and dried over anhydrous sodium sulfate to obtain a crude compound. This crude compound was purified by reverse phase chromatography to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-4-piperidyl]-2,2-dimethyl-propanamide (50 mg) as a white solid.

[0389] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.54 (d, $J = 5.0$ Hz, 1H), 8.08 (s, 1H), 7.96 (s, 1H), 7.69 (d, $J = 7.4$ Hz, 1H), 7.55 (s, 1H), 7.49 (d, $J = 5.1$ Hz, 1H), 7.41 (d, $J = 7.8$ Hz, 1H), 7.21 (d, $J = 7.7$ Hz, 1H), 4.41 (s, 1H), 3.86 (s, 1H), 3.57 (s, 1H), 3.17 (s, 1H), 2.89 (s, 1H), 1.76 (s, 2H), 1.71 (s, 2H), 1.36 (s, 9H), 1.08 (s, 9H). LCMS:- (M+1) 538.2.

Example 77. Preparation of Compound No. 77

Synthesis of 4-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl} piperazin-2-one



Step-1. Synthesis of 4-(4-bromo-5-methyl-2-thienyl)-3-chloro-benzoic acid

[0390] 3,5-Dibromo-2-methyl-thiophene (10 g, 39.2 mmol), 4-borono-3-chloro-benzoic acid (9.4 g, 47.0 mmol), and sodium carbonate (10.2 g, 98.0 mmol) in water (10 mL) and DMF (100.0 mL) were charged in a 250 mL glass bottle and purged with nitrogen gas for 20 min. After adding tetrakis (4.5g, 3.9 mmol), the reaction mixture was heated at 100 °C for 16h. The reaction was monitored by TLC and LCMS. After completion of reaction, water (200 mL) was added and filtered with washings of ether and hexane obtain 4-(4-bromo-5-methyl-2-thienyl)-3-chloro-benzoic acid (10gm) as the freebase, a yellow solid. LCMS:- (M+1) 331.0.

Step-2. Synthesis of 4-[4-(2-tert-butyl-4-pyridyl)-5-methyl-2-thienyl]-3-chloro-benzoic acid:

[0391] 4-(4-Bromo-5-methyl-2-thienyl)-3-chloro-benzoic acid (5.0 g, 15.1 mmol), 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (4.3 g, 16.1 mmol), sodium carbonate (3.9 g, 37.9 mmol) in water (5.0 mL), and DMF (50.0 mL) were charged in a 250 mL glass bottle and purged with nitrogen gas for 15 min. After adding tetrakis (1.7 g, 1.5 mmol), the reaction mixture was heated at 100 °C for 16h. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was cooled to 0 °C and 3N HCl was added dropwise. A yellow precipitate was obtained which was filtered to obtain 4-[4-(2-tert-butyl-4-pyridyl)-5-methyl-2-thienyl]-3-chloro-benzoic acid (4.0 g) as a yellow solid. LCMS:- (M+1) 386.1.


Step-3. Synthesis of 4-[4-[4-(2-tert-butyl-4-pyridyl)-5-methyl-2-thienyl]-3-chloro-benzoyl]piperazin-2-one:

[0392] 4-[4-(2-tert-Butyl-4-pyridyl)-5-methyl-2-thienyl]-3-chloro-benzoic acid (500 mg, 1.29 mmol) was taken in DMF (10 mL), followed by addition of DIPEA (0.9mL, 5.18 mmol), HATU (980 mg, 2.58 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5h, followed by addition of piperazin-2-one (510 mg, 5.18 mmol). The reaction mixture was stirred at RT for 16h. After completion of reaction, water (150 mL) was added and the mixture extracted with EtOAc (2x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to obtain 4-[4-[4-(2-tert-butyl-4-pyridyl)-5-methyl-2-thienyl]-3-chloro-benzoyl]piperazin-2-one (54mg) as the HCl salt.

[0393] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.69 (d, J = 6.3 Hz, 1H), 8.16 – 8.04 (m, 2H), 7.85 – 7.75 (m, 2H), 7.73 – 7.62 (m, 1H), 7.58– 7.47 (m, 1H), 4.31 (s, 1H), 4.13 (s, 1H), 3.95 (s, 1H), 3.70 (s, 1H), 3.41 (s, 2H), 2.77 (s, 3H), 1.58 (s, 9H). LCMS:- (M+1) 468.0.

Example 78. Preparation of Compound No. 78

Synthesis of tert-butyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamido} piperidine-1-carboxylate




[0394] To a solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (300 mg, 0.8086 mmol) in DMF (10 mL), was added DIPEA (696.8 mg, 5.3908 mmol), followed by the addition of HATU (1025 mg, 2.6954 mmol), the reaction mixture was stirred for 30 min. tert-

Butyl 4-aminopiperidine-1-carboxylate (645 mg, 3.22 mmol) was added to the reaction mixture and the mixture stirred the reaction mixture overnight at R.T. The reaction was monitored by TLC and LCMS. On completion, the reaction mixture was diluted with water (30 mL) and extracted with EtOAc (3x50 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL) and dried over anhydrous sodium sulfate to obtain a crude compound. The crude compound was purified by reverse phase HPLC to afford tert-butyl 4-[[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro benzoyl]amino]piperidine-1-carboxylate (45 mg) as a white solid.

[0395] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.52 (d, $J = 8.3$ Hz 1H), 8.50 (d, $J = 8.5$ Hz 1H), 8.03 (d, $J = 9.1$ Hz, 2H), 7.87 (s, $J = 8.0$ Hz, 1H), 7.85 (d, $J = 9.1$ Hz 1H), 7.71 (d, 1H), 7.65 (s, 1H), 7.49 (d, $J = 5.1$ Hz, 1H), 3.94 (d, $J = 14.1$ Hz, 3H), 2.83 (s, 2H), 1.79 (d, $J = 11.9$ Hz, 3H), 1.39 (s, 10H), 1.34 (s, 9H). LCMS – (M+1) 554.2.

Example 79. Preparation of Compound Nos. 79, 79a and 79b

Synthesis of tert-butyl N-[(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate




[0396] To a stirred solution 4-[4-(2-tert-butyl-4-pyridyl)-5-methyl-2-thienyl]-3-chloro-benzoic acid (500 mg, 1.347 mmol) in DMF (10 mL), was added DIPEA (0.92 mL, 5.390 mmol) followed by addition of HATU (1.0 g, 2.69 mmol) and the mixture stirred at RT for 30 min. Then (*R*)-tert-butyl N-methyl- N-(3-piperidyl) carbamate (1.1 g, 5.39 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (10 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain Compound 79a [tert-butyl-(*R*)-N-[1-[4-[4-(2-tert-butyl-4-pyridyl)-5-methyl-2-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate] (174 mg) freebase as a solid. Compound 79b, the (*S*)-enantiomer, was also prepared using the (*S*)-tert-butyl N-methyl- N-(3-piperidyl) carbamate reagent.

[0397] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.58 (d, $J = 5.2$ Hz, 1H), 7.88 (s, 1H), 7.62 (s, 2H), 7.52 (s, 1H), 7.42 (d, $J = 5.3$, 1H), 7.36 (d, $J = 7.9$, 1H), 6.98 (s, 1H), 4.2 (s, 1H), 3.42 (s,

2H), 3.08 (s, 2H), 2.75 (s, 1H), 2.58 (s, 3H) 1.80 (s, 3H), 1.41-1.20 (m, 18H). LCMS:- 568.1 (M+1).

Example 80. Preparation of Compound Nos. 80, 80a, and 80b

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-[(4-chlorophenyl)methyl]piperidin-3-amine



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0398] In a 100 mL glass bottle 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in 25 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by

silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2 Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0399] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.), were charged in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (176 mg, 0.152 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 80 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.

Step-3: Synthesis of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate

[0400] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (500 mg, 1.34 mmol, 1 eq.) was dissolved in DMF (13 mL), followed by addition of DIPEA (1.0 mL, 5.4 mmol, 4 eq.) and HATU (1.07 g, 2.69 mmol, 2 eq.), and the resulting mixture was stirred for 10 min at RT and tert-butyl N-(3-piperidyl)carbamate (1.07 g, 5.4 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (700 mg) as a crude viscous compound which was used as such for the next step of synthesis.

Step-4: Synthesis of (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone

[0401] In a 100 mL flask, tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (700 mg, 1.26 mmol, 1 eq.) was charged in DCM (15 mL), the reaction mixture was maintained at 0 °C and trifluoroacetic acid (5 mL) was added dropwise and

the mixture stirred at RT for 2.5 h. Progress of reaction was monitored by TLC/LCMS. After completion of reaction, the DCM was evaporated under reduced pressure, and the residue basified with saturated aqueous sodium bicarbonate solution and extracted with EtOAc (2x100 mL). The organic layer was washed with brine (100 mL) and dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure to afford (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone as a crude viscous compound (570 mg) which was used as such for the next step of synthesis.

Step-5: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-[(4-chlorophenyl)methylamino]-1-piperidyl]methanone

[0402] In a 100 mL flask, tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (230 mg, 0.5 mmol, 1 eq.) was charged in (10 mL) of methanol and the mixture stirred at RT with 4-chlorobenzaldehyde (64 mg, 0.45 mmol, 1 eq.) for 2h. Then at RT, 3 drops of acetic acid and NaCNBH₃ (64 mg, 1.01 mmol, 2 eq.) were added and the mixture stirred for 30 min at RT. Progress of reaction was monitored by TLC/LCMS. After completion of reaction, methanol was evaporated under reduced pressure, and the residue basified with saturated aqueous sodium bicarbonate solution and extracted with EtOAc (2x100 mL). The organic layer was washed with brine (100 mL) and dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure, to give a viscous compound that was purified by reverse phase chromatography to afford [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-[(4-chlorophenyl)methylamino]-1-piperidyl] methanone as a solid free base compound (94 mg).


Step 6: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-[(4-chlorophenyl)methylamino]-1-piperidyl]methanone hydrochloride

[0403] In a 250 mL flask, [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-[(4-chlorophenyl)methylamino]-1-piperidyl]methanone (94 mg, 0.163 mmol, 1 eq.) was dissolved in 3 mL of EtOH and at 0 °C treated with 2 mL of ethanolic HCl, then evaporated under reduced pressure and directly lyophilized to get a free flowing solid (100 mg, off-white colored) as its HCl salt. The enantiomers were prepared from chiral HPLC resolution of the racemate, to give Compound 80a [(R)-(4-(5-(2-(tert-butyl)pyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(3-((4-chlorobenzyl)amino)piperidin-1-yl)methanone] and Compound 80b [(S)-(4-(5-(2-(tert-butyl)pyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(3-((4-chlorobenzyl)amino)piperidin-1-yl)methanone].

[0404] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.61 (d, $J = 6.3$ Hz, 1H), 8.41 (s, 1H), 8.23 – 8.14 (m, 3H), 7.75 – 7.67 (m, 2H), 7.52 (m, $J = 7.7$ Hz, 5H), 4.67 (s, 1H), 4.37 (m, 2H), 3.70 (s, 1H), 3.42 (s, 3H), 2.37 (m, 1H), 1.85 (m, 2H), 1.68 (m, 1H), 1.57 (s, 9H). LCMS=578 (M+1).

Example 81. Preparation of Compound Nos. 81, 81a, and 81b

Synthesis of 1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-[(3-chlorophenyl)methyl]piperidin-3-amine



Step-1: Synthesis of 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine

[0405] In a 100 mL glass bottle, 2,4 dibromothiophene (600 mg, 2.5 mmol, 1eq.) along with 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (652 mg, 1eq.) and potassium phosphate (1.325 g, 2.5 eq.) were charged in 25 mL of THF. The mixture was purged with nitrogen for 15 min then Xantphos (145 mg, 0.1 eq.) and tetrakis (289 mg, 0.1 eq.) were added, and the mixture repurged for 5 min and the mixture stirred at 60 °C overnight. Progress of the reaction was monitored by TLC & LCMS. When the reaction was completed, the mixture was cooled to RT, water (100 mL) was added and the mixture extracted with EtOAc (3x100 mL). The combined organic layer was washed with brine (50 mL) then dried over anhydrous sodium sulfate, concentrated under reduced pressure to give a viscous compound which was purified by

silica gel (60-120) column chromatography using EtOAc/Hexane (0-20%) as eluent system to afford 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine as a colorless semi-solid (500 mg) in pure form.

Step-2: Synthesis of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid

[0406] In a 25 mL glass bottle, 4-(4-bromo-2-thienyl)-2-tert-butyl-pyridine (450 mg, 1.52 mmol, 1 eq.) and 4-borono-3-chloro-benzoic acid (397 mg, 1.98 mmol, 1.3 eq.), were charged in DMF (7 mL) followed by addition of sodium carbonate (404 mg, 2.5 equiv dissolved in water (2.0 mL) and the reaction mixture was purged with nitrogen gas for 5 min. After adding tetrakis (176 mg, 0.152 mmol, 0.1 eq.) the mixture was repurged for 2 min and was stirred at 80 °C overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was allowed to come to RT and diluted with water (50 mL) and extracted with EtOAc:MeOH (90:10), (3x100 mL). The combined organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford a solid compound, 3-chloro-4-[5-[3-(1-piperidyl)phenyl]-3-thienyl]benzoic acid (500 mg) which was used as such for the next step of synthesis.

Step-3: Synthesis of tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate

[0407] In a 100 mL flask, 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (500 mg, 1.34 mmol, 1 eq.) was dissolved in DMF (13 mL), followed by addition of DIPEA (1.0 mL, 5.4 mmol, 4 eq.) and HATU (1.07 g, 2.69 mmol, 2 eq.), and the resulting mixture was stirred for 10 min at RT and tert-butyl N-(3-piperidyl)carbamate (1.07 g, 5.4 mmol, 4 eq.) were added and the mixture stirred at RT under nitrogen atmosphere overnight. The progress of the reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (2x100 mL), washed with water (4x100 mL) then dried over anhydrous sodium sulfate and the combined organic layer was concentrated under reduced pressure to afford N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (700 mg) as a crude viscous compound which was used as such for the next step of synthesis.

Step-4: Synthesis of (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone

[0408] In a 100 mL flask, tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (700 mg, 1.26 mmol, 1 eq.) was charged in DCM (15 mL), the reaction mixture was maintained at 0 °C and trifluoroacetic acid (5 mL) was added dropwise and

the mixture stirred at RT for 2.5 h. Progress of reaction was monitored by TLC/LCMS. After completion of reaction, the DCM was evaporated under reduced pressure, and the residue basified with saturated aqueous sodium bicarbonate solution and extracted with EtOAc (2x100 mL). The organic layer was washed with brine (100 mL) and dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure to afford (3-amino-1-piperidyl)-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone as a crude viscous compound (570 mg) which was used as such for the next step of synthesis.

Step-5: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-[(3-chlorophenyl)methylamino]-1-piperidyl]methanone

[0409] In a 100 mL flask, tert-butyl N-[1-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3-piperidyl]carbamate (250 mg, 0.55 mmol, 1 eq.) was charged in (8 mL) of DMF and the mixture stirred with potassium carbonate (228 mg, 1.65 mmol, 3 eq.) at RT for 30 min then 3-chlorobenzylbromide (113 mg, 1 eq.), was added and the mixture stirred at RT for 2h. Progress of reaction was monitored by TLC/LCMS. After maximum conversion of starting material into product, extracted with EtOAc (2x100 mL). Then washed with water (4x100 mL). The organic layer was washed with brine (100 mL) and dried over anhydrous sodium sulfate. The solution was concentrated under reduced pressure, viscous compound and purified by reverse phase chromatography to afford [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-[(3-chlorophenyl)methylamino]-1-piperidyl]methanone as a solid free base compound (47 mg).


Step-6: Synthesis of [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-[(3-chlorophenyl)methylamino]-1-piperidyl]methanone hydrochloride

[0410] In 250 mL flask, [4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-[3-[(3-chlorophenyl)methylamino]-1-piperidyl]methanone (47 mg, 0.081 mmol, 1 eq.) was dissolved in 3 mL of EtOH and at 0 °C treated with 2 mL of ethanolic HCl, then evaporated the EtOH under reduced pressure and directly lyophilized to get free flow solid (50 mg, off-white colored) as its respective HCl salt. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0411] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.61 (d, J = 6.4 Hz, 1H), 8.41 (d, J = 1.4 Hz, 1H), 8.24 – 8.14 (m, 3H), 7.75 – 7.67 (m, 2H), 7.64 (s, 1H), 7.51 (d, J = 8.8 Hz, 4H), 4.68 (m, 1H), 4.37 (m, 2H), 3.5 (m, 2H), 2.37 (m, 2H), 1.85 (m, 2H), 1.69 (m, 2H), 1.57 (s, 9H).
LCMS=578 (M+1).

Example 82. Preparation of Compound Nos. 82, 82a and 82b

Synthesis of (3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}pyrrolidin-3-ol




[0412] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (500 mg, 1.34 mmol) in DMF (10 mL), was added DIPEA (0.9 mL, 5.390 mmol) followed by addition of HATU (1.0 g, 2.694 mmol) and the mixture stirred at RT for 30 min. Then (R)-pyrrolidin-3-ol (470 mg, 5.390 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain Compound 82a {[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]-(3-hydroxypyrrolidin-1-yl)methanone} (132 mg) as a solid. Compound 82b, the (S)-enantiomer, can be prepared by using the (S)-pyrrolidin-3-ol.

[0413] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.7 (d, $J = 5.2$ Hz, 1H), 8.50 (s, 1H), 8.28 (s, 1H), 8.18 – 8.00 (m, 2H), 7.70 (m, 2H), 7.59 (m, 1H), 4.38-4.20 (m, 1H), 3.6 (m, 2H), 3.4 (m, 1H), 3.2 (m, 1H), 1.95 (m, 1H), 1.80 (m, 1H), 1.41 (s, 9H). LCMS=441.0 (M+1). LCMS=441.0 (M+1).

Example 83. Preparation of Compound Nos. 83, 83a, and 83b

Synthesis of 7-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2,7-diazaspiro[4.5]decan-1-one




[0414] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (500 mg, 1.34 mmol) in DMF (10 mL), was added DIPEA (0.9 mL, 5.390 mmol) followed by addition of HATU (1.0 g, 2.694 mmol) and the mixture stirred at RT for 30 min. Then 3,9-diazaspiro[4.5] decan-4-one (1.0 g, 5.390 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain 9-[4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoyl]-3,9-diazaspiro[4.5]decan-4-one (140 mg) as a solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0415] ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.72 (d, $J = 5.2$ Hz, 1H), 8.48 (s, 1H), 8.24 (s, 1H), 8.18 – 8.00 (m, 2H), 7.70 (dd, $J = 5.3, 1.8$ Hz, 1H), 7.60 (s, 1H), 7.42 (dd, $J = 7.9, 1.7$ Hz, 1H), 4.20-4.00 (m, 2H), 3.80 (m, 2H), 3.20 (m, 3H), 3.0 (m, 1H), 2.00- 1.87 (m, 2H), 1.70 (m, 2H), 1.41 (s, 9H). LCMS=508.1 (M+1).

Example 84. Preparation of Compound Nos. 84, 84a, and 84b

Synthesis of 2-tert-butyl-4-[4-(2-chloro-4-{octahydropyrrolo[1,2-a]piperazine-2-carbonyl} phenyl)thiophen-2-yl]pyridine




[0416] To a stirred solution of 4-[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-benzoic acid (500 mg, 1.34 mmol) in DMF (10 mL), was added DIPEA (0.9 mL, 5.390 mmol) followed by addition of HATU (1.0 g, 2.694 mmol) and the mixture stirred at RT for 30 min. Then 1,2,3,4,6,7,8,8a-octahydropyrrolo[1,2-a]pyrazine (680 mg, 5.390 mmol) was added at the same temperature. The reaction mixture was stirred at RT overnight. The reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (50 mL) and extracted with EtOAc (2x50 mL), the organic layer was washed with water (2x20 mL) and brine (20 mL) and dried over anhydrous sodium sulfate to obtain a crude product which was purified by reverse phase HPLC to obtain 3, 4, 6, 7, 8, 8a-hexahydro-1H-pyrrolo [1,2-a]pyrazin-2-yl-[4-

[5-(2-tert-butyl-4-pyridyl)-3-thienyl]-3-chloro-phenyl]methanone (130 mg) as a solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0417] ^1H NMR (400 MHz, Dms o-d_6) δ (ppm): 8.58 (d, $J = 5.2$ Hz, 1H), 8.04(s, 1H), 7.98 (s, 1H), 7.70-7.50 (s, 3H), 7.52 – 7.48 (m, 2H), 3.87 (m, 2H), 3.60 (m, 2H), 3.42(s, 1H), 3.25(m, 2H), 3.10 (s, 1H), 2.90 (s, 1H), 1.98 (m, 4H), 1.38 (s, 9H). LCMS=480.2 (M+1).

Example 85. Preparation of Compound Nos. 85, 85a, and 85b

Synthesis of 1-{4-[2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl]-3-chlorobenzoyl}piperidin-3-amine



Step-1: Synthesis of 2,4-dibromo-1-(methoxymethyl)imidazole

[0418] 2,4-Dibromo-1H-imidazole (3.0 g, 13.3 mmol) was taken in THF (25.0 mL) followed by addition of NaH (0.98 g, 20.0mmol) at 0 °C. The reaction mixture was stirred at RT for 15 min followed by addition of chloro(methoxy)methane (1.2 g, 14.6 mmol). The reaction mixture was stirred to RT for 1h and completion of reaction was monitored by TLC and LCMS. After completion of reaction, ice cooled water (200 mL) was added and the mixture extracted with EtOAc (300 mL). The organic layer was separated, dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain 2,4-dibromo-1-(methoxymethyl)imidazole (4.0 g) as a crude product.

Step-2: Synthesis of 4-[4-bromo-1-(methoxymethyl) imidazol-2-yl]-2-tert-butyl-pyridine

[0419] 2,4-Dibromo-1-(methoxymethyl)imidazole (2.0 g, 7.50 mmol), 2-tert-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (1.6 g, 6.00 mmol), sodium carbonate (2.0 g, 18.8 mmol) in water (5.0 mL), and dioxane (20.0 mL) were charged in a 100 mL glass bottle and purged with nitrogen gas for 15 min. Tetrakis (0.866 g, 0.80 mmol) and Xantphos (0.463 g, 0.86 mmol) were added then the reaction mixture was heated at 80 °C for 16h. Completion of reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (150 mL) and extracted with EtOAc (200 mL). The organic layer was separated and washed with brine (100 mL) and dried over anhydrous sodium sulfate to obtain crude 4-[4-bromo-1-(methoxymethyl)imidazol-2-yl]-2-tert-butyl-pyridine (3.0 g) as the freebase.

Step-3: Synthesis of tert-butyl N-[1-[4-[2-(2-tert-butyl-4-pyridyl)-1-(methoxymethyl)imidazol-4-yl]-3-chloro-benzoyl]-3-piperidyl] carbamate

[0420] 4-[4-Bromo-1-(methoxymethyl)imidazol-2-yl]-2-tert-butyl-pyridine (500 mg, 1.54 mmol), [4-[3-(tert-butoxycarbonylamino)piperidine-1-carbonyl]-2-chloro-phenyl]boronic acid (1.18 g, 3.08 mmol), and sodium carbonate (405 mg, 3.85 mmol) in water (1.0 mL), DMF (5.0mL) were charged in a 25.0 mL glass bottle and purged with nitrogen gas for 15 min. After addition of tetrakis (178 mg, 0.015 mmol), the reaction mixture was heated at 100 °C for 16h. Completion of reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was diluted with water (100 mL) and extracted with EtOAc (150 mL). The organic layer was separated and washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude product was purified by reverse phase HPLC to obtain tert-butyl N-[1-[4-[2-(2-tert-butyl-4-pyridyl)-1-(methoxymethyl)imidazol-4-yl]-3-chloro-benzoyl]-3-piperidyl] carbamate (200 mg) as freebase.

Step-4: Synthesis of (3-amino-1-piperidyl)-[4-[2-(2-tert-butyl-4-pyridyl)-1H-imidazol-4-yl]-3-chloro-phenyl]methanone


[0421] tert-Butyl N-[1-[4-[2-(2-tert-butyl-4-pyridyl)-1-(methoxymethyl)imidazol-4-yl]-3-chloro-benzoyl]-3-piperidyl]carbamate (200 mg, 0.34mmol) was taken in 1.25M HCl in EtOH (5.0 mL) under nitrogen atmosphere. The reaction mixture was stirred at 50 °C for 2h. Completion of reaction was monitored by TLC and LCMS. After completion of reaction, the mixture was concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC method to obtain (3-amino-1-piperidyl)-[4-[2-(2-tert-butyl-4-pyridyl)-1H-imidazol-4-yl]-3-chloro-phenyl]methanone (15 mg) as the HCl salt, a yellow solid. The enantiomers can be prepared from chiral HPLC resolution of the racemate.

[0422] ¹H NMR (400 MHz, Methanol-d₄) δ (ppm): 8.74 (d, J = 6.3 Hz, 1H), 8.57 (s, 1H), 8.43 (d, J = 6.4 Hz, 1H), 8.27 – 8.17 (m, 2H), 7.68 (d, J = 1.7 Hz, 1H), 7.53 (d, J = 8.4 Hz, 1H), 4.41

(s, 1H), 3.64 (s, 1H), 3.39 (s, 1H), 2.18 (s, 1H), 1.84 (s, 1H), 1.73 (d, $J = 15.5$ Hz, 2H), 1.60 (s, 9H), 1.29 (s, 2H).

Example 86. Preparation of Compound No. 86

Synthesis of 1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-4-ol




[0423] 4-[4-(2-tert-Butyl-4-pyridyl)-2-thienyl]-3-chloro-benzoic acid (300 mg, 0.80 mmol) was taken in DMF (6.0 mL), followed by addition of DIPEA (0.60 mL, 3.22 mmol), HATU (612 mg, 1.61 mmol), under nitrogen atmosphere. The reaction mixture was stirred at RT for 0.5 h, followed by addition of piperidin-4-ol (322 mg, 3.22 mmol). The reaction mixture was stirred at RT for 16h. After completion of reaction, water (50 mL) was added and the mixture extracted with EtOAc (2x50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to obtain the crude product which was purified by reverse phase HPLC to obtain [4-[4-(2-tert-butyl-4-pyridyl)-2-thienyl]-3-chloro-phenyl]-(4-hydroxy-1-piperidyl)methanone (40 mg) as the freebase.

[0424] ^1H NMR (400 MHz, Methanol- d_4) δ (ppm): 8.70 – 8.60 (m, 2H), 8.31 (s, 1H), 8.24 (d, $J = 6.4$ Hz, 1H), 8.14 (d, $J = 1.6$ Hz, 1H), 7.82 (d, $J = 8.0$ Hz, 1H), 7.64 (d, $J = 1.7$ Hz, 1H), 7.47 (dd, $J = 7.9, 1.7$ Hz, 1H), 4.18 (s, 1H), 3.93 (dt, $J = 8.4, 4.5$ Hz, 1H), 3.67 (s, 1H), 3.40 (s, 2H), 1.97 (s, 1H), 1.86 (s, 1H), 1.58 (s, 9H), 1.51 (s, 2H). LCMS:- ($M+1$) 454.0.

Example 87. Preparation of Compound Nos. 87, 87a and 87b

Synthesis of (3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine



[0425] tert-Butyl N-[(1S)-3-[4-[4-(2-tert-butyl-4-pyridyl)-5-methyl-2-thienyl]-3-chlorobenzoyl]cyclohexyl]carbamate (40 mg, 0.074mmol) was taken in EtOH (3.0 mL) at 0 °C, followed by dropwise addition of 1.25M HCl in EtOH (3.0 mL) under nitrogen atmosphere. The

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 180

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 180


NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

CLAIMS

1. A compound of Formulae (Ia) or (Ib):



or a pharmaceutically acceptable salt thereof, wherein:

A is either:

- iii. a heteroaryl having only one ring, substituted with one or more **R1** or **R2**; or
- iv. an aryl or heteroaryl, each having more than one ring, optionally substituted with one or more **R1** or **R3**;

R1 is C1-C6 linear or branched alkyl, C2-C6 linear or branched alkenyl, C2-C6 linear or branched alkynyl, C3-C6 cycloalkyl, C3-C6 cycloalkylmethyl, C3-C6 cycloalkenyl, aryl, or heteroaryl; each optionally substituted with one or more **R3**;

R2 is halogen, -CN, -OH, -O(Alkyl), -NO₂, -SH, -S(Alkyl), -S(O)(Alkyl), -S(O)₂(Alkyl), -CH₂OCH₃, -OBn, -CO₂H, -CO₂(Alkyl), -NR₁₀R₁₁, -CONR₁₀R₁₁, and -S(O)₂NR₁₀R₁₁;

R3 is halogen, -CN, -OH, -O(Alkyl), =O, -NO₂, -SH, -S(Alkyl), -S(O)(Alkyl), -S(O)₂(Alkyl), -CH₂OCH₃, -OBn, -CO₂H, -CO₂(Alkyl), -NR₁₀R₁₁, -CONR₁₀R₁₁, and -S(O)₂NR₁₀R₁₁;

Each **R_{C4}**, **R_{C5}**, and **R_{C6}**, is independently hydrogen, **R1**, or **R2**; or **R_{C4}** is taken with **R_{C7}** to form a heterocyclyl optionally substituted with one or more **R3**;

Each **R_{C7}** and **R_{C8}** is independently hydrogen, **R1**, or -SO₂**R_{C12}**; or **R_{C7}** and **R_{C8}** are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more **R1** or **R3**; or **R_{C7}** is taken with **R_{C4}** to form a heterocyclyl optionally substituted with one or more **R3**;

R10 and **R11** are independently hydrogen, C1-C6 linear or branched alkyl, -C(O)**R_{C12}**, -C(O)₂**R_{C12}**, -C(O)N(**R_{C12}**)₂, -SO₂**R_{C12}**, or are taken together with the N to which they are attached to form a C3-C8 heterocyclyl;

R_{C12} is hydrogen, a linear or branched C1-C6 alkyl, a linear or branched C2-C6 alkenyl, a linear or branched C2-C6 alkynyl, or aryl; wherein each alkyl, alkenyl, alkynyl, or aryl


group is optionally substituted with one or more halogen atoms, one or more $-NH_2$ groups, or one or more $-OH$ groups;


V is the moiety $-C(=O)-$ or $-S(=O)_2-$; and

The ring comprising **W**, **X**, and **Y** is a 5-membered heteroaromatic, wherein **W** and **Y** are independently selected from the group consisting of CR_A , N, NR_B , O and S; **X** is selected from the group consisting of NR_B , O and S; wherein **R_A** is hydrogen, **R1** or **R2**; and **R_B** is hydrogen or **R1**;


With the proviso that when **W** is N, **X** is S, and **Y** is C, then one of **R_{C4}**, **R_{C5}** and **R_{C6}** is other than H.

2. The compound of claim 1, wherein **A** is a moiety selected from the group consisting of:







3. The compound of claim 2, wherein A is a moiety selected from the group consisting of:




4. The compound of claim 3, wherein A is a moiety selected from the group consisting of:



5. The compound of any of claims 1-4, wherein the ring comprising **W**, **X**, and **Y**, and the carbon atoms to which they are attached, is a 5-membered heteroaromatic group selected from the group consisting of:




6. The compound of claim 5, wherein the ring comprising **W**, **X**, and **Y**, and the carbon atoms to which they are attached, is a 5-membered heteroaromatic group selected from the group consisting of:




7. The compound of claim 6, wherein **R_B** is a linear C1-C6 alkyl selected from methyl, ethyl, n-propyl, n-butyl, n-pentyl or n-hexyl.

8. The compound of claim 6, wherein **R_B** is a branched C1-C6 alkyl selected from iso-propyl, iso-pentyl, and tert-butyl.
9. The compound of any of claims 1-8, wherein the phenyl ring containing the groups **R_{C4}**, **R_{C5}**, **R_{C6}**, and **-C(=O)NR_{C7}R_{C8}**, is a moiety selected from:




10. The compound of claim 9, wherein the phenyl ring containing the groups **R_{C4}**, **R_{C5}**, **R_{C6}**, and **-C(=O)NR_{C7}R_{C8}**, is a moiety selected from:




11. The compound of any of claims 1-10, wherein each **R_{C4}**, **R_{C5}** and **R_{C6}**, is independently hydrogen, **R₁**, or **R₂**.
12. The compound of claim 11, wherein at least one of **R_{C4}**, **R_{C5}** and **R_{C6}** is hydrogen.
13. The compound of claim 11, wherein two of **R_{C4}**, **R_{C5}** and **R_{C6}** are hydrogen.
14. The compound of claim 11, wherein each **R_{C4}**, **R_{C5}** and **R_{C6}** is hydrogen.
15. The compound of any of claims 1-14, wherein at least one of **R_{C4}**, **R_{C5}** and **R_{C6}** is halogen.
16. The compound of claim 15, wherein two of **R_{C4}**, **R_{C5}** and **R_{C6}** are halogen.
17. The compound of claim 15, wherein each **R_{C4}**, **R_{C5}** and **R_{C6}** is halogen.

18. The compound of any of claims 1-11, wherein one of R_{C4} , R_{C5} and R_{C6} is hydrogen, and the remaining two of R_{C4} , R_{C5} and R_{C6} are halogen.
19. The compound of any of claims 1-11, wherein two of R_{C4} , R_{C5} and R_{C6} are hydrogen, and the remaining one of R_{C4} , R_{C5} and R_{C6} is halogen.
20. The compound of any of claims 1-11, wherein R_{C4} , R_{C5} and R_{C6} are each halogen.
21. The compound of any of claims 1-10, wherein R_{C4} is taken with R_{C7} to form a heterocyclyl optionally substituted with one or more R_3 .
22. The compound of claim 21, wherein R_{C4} is taken together with R_{C7} to form a moiety selected from:




23. The compound of any of claims 1-21, wherein:


A is a moiety selected from



the 5-membered heteroaromatic group comprising W , X and Y is selected from:




the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is a moiety selected from:




24. The compound of any of claims 1-21, wherein:


A is a moiety selected from



the 5-membered heteroaromatic group comprising **W**, **X** and **Y** is selected from:




the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is a moiety selected from:




25. The compound of any of claims 1-21, wherein:


A is




the 5-membered heteroaromatic group comprising **W**, **X** and **Y** is selected from:




the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is:




26. The compound of any of claims 1-25, wherein R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more **R3**.
27. The compound of any of claims 1-21, wherein:




the 5-membered heteroaromatic group comprising **W**, **X** and **Y** is selected from



the phenyl ring containing the groups R_{C4} , R_{C5} , R_{C6} , and $-C(=O)NR_{C7}R_{C8}$, is




28. The compound of claim 1, which is a compound of Formula (IIa):



wherein R_{C5} is halogen, and each R_{C7} and R_{C8} is independently hydrogen, $R1$, or - SO_2R_{C12} ; or R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more $R1$ or $R3$.

29. The compound of claim 28, wherein R_{C5} is chloro.
30. The compound of claim 28 or 29 wherein R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle, wherein the C3-C8 heterocycle is selected from the group consisting of pyrrolidinyl, piperidinyl, piperazinyl, and morpholinyl, each optionally substituted with one or more $R1$ or $R3$.
31. The compound of claim 1, which is a compound of Formula (IIb):



wherein R_{C5} is halogen, and each R_{C7} and R_{C8} is independently hydrogen, $R1$, or - SO_2R_{C12} ; or R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle optionally substituted with one or more $R1$ or $R3$.

32. The compound of claim 31, wherein R_{C5} is chloro.
33. The compound of claim 31 or 32, wherein R_{C7} and R_{C8} are taken together with the N to which they are attached to form a C3-C8 heterocycle, wherein the C3-C8 heterocycle is selected from the group consisting of pyrrolidinyl, piperidinyl, piperazinyl, and morpholinyl, each optionally substituted with one or more $R1$ or $R3$.
34. A compound selected from the group consisting of:
 N-methyl-4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide;
 N,N-dimethyl-4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide;
 4-[2-(2-propylpyridin-4-yl)-1,3-thiazol-4-yl]benzamide;
 4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1,3-thiazol-2-yl}-1H-indole;
 2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1,3-thiazol-2-yl}pyridine;
 3-chloro-4-[2-(1H-indol-4-yl)-1,3-thiazol-4-yl]-N,N-dimethylbenzamide;
 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-2-one;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-cyclobutylbenzamide;

3-chloro-N,N-dimethyl-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzamide;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-4-methylpiperazine;

2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]thiophen-2-yl}pyridine;

2-tert-butyl-4-{4-[2-chloro-4-(3,3-difluoropyrrolidine-1-carbonyl)phenyl]thiophen-2-yl}pyridine;

2-tert-butyl-4-{4-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]thiophen-2-yl}pyridine;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-cyclopropylbenzamide;

N-tert-butyl-4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamide;

2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-5-methylthiophen-2-yl}pyridine;

4-(3-chloro-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzoyl)morpholine;

3-chloro-N-(2-hydroxypropyl)-4-{2-[2-(piperidin-1-yl)pyridin-4-yl]-1,3-thiazol-4-yl}benzamide;

tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)carbamate;

tert-butyl N-[(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

tert-butyl N-[(3S)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-1 λ ⁶,4-thiomorpholine-1,1-dione;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-amine;

(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-amine;

(3S)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-amine;

4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}morpholine;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-phenylbenzamide;

2-tert-butyl-4-{4-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]-1H-imidazol-2-yl}pyridine;

2-tert-butyl-4-{4-[2-chloro-4-(piperidine-1-carbonyl)phenyl]-1H-imidazol-2-yl}pyridine;

6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-2-methyl-1,2,3,4-tetrahydroisoquinolin-1-one;

tert-butyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazine-1-carboxylate;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazine;

1-(4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-1-yl)ethan-1-one;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-4-methanesulfonylpiperazine;

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)methanesulfonamide;

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)acetamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N,N-bis(propan-2-yl)benzamide;

1-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)pyrrolidin-2-one;

6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-2-(propan-2-yl)-1,2,3,4-tetrahydroisoquinolin-1-one;

2-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-1λ⁶,2-thiazolidine-1,1-dione;

tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)carbamate;

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)acetamide;

4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-1-ethylpiperazin-2-one;

2-tert-butyl-4-{5-[2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl]thiophen-3-yl}pyridine;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-methyl-N-phenylbenzamide;

4-{4-[4-(azetidine-1-carbonyl)-2-chlorophenyl]thiophen-2-yl}-2-tert-butylpyridine;

6-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2-oxa-6-azaspiro[3.3]heptane;

ethyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)carbamate;

ethyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazine-1-carboxylate;

1-(4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperazin-1-yl)-2,2-dimethylpropan-1-one;

N-[2-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorophenyl}-N-ethylformamido)ethyl]acetamide;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-ol;

(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-ol;

(3S)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-ol;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(3-hydroxyphenyl)benzamide;

(2S,6R)-4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2,6-dimethylmorpholine;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-1,2,3,4-tetrahydro-1,8-naphthyridine;

6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-7-chloro-2-(propan-2-yl)-1,2,3,4-tetrahydroisoquinolin-1-one;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-[(4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl]benzamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(4-fluorophenyl)benzamide

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-2,2-dimethylpropanamide;

1-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-3-cyclopropylurea;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-ol;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-(pyridin-3-yl)benzamide;

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)methanesulfonamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-methyl-N-(1,3-oxazol-2-yl)benzamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-[2-(morpholin-4-yl)-2-oxoethyl]benzamide;

4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chloro-N-{8-methyl-8-azabicyclo[3.2.1]octan-3-yl}benzamide;

ethyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)carbamate;

tert-butyl N-[1-(3-chloro-4-{5-[2-(morpholin-4-yl)pyridin-4-yl]thiophen-3-yl}benzoyl)piperidin-3-yl]carbamate;

1-(3-chloro-4-{5-[2-(morpholin-4-yl)pyridin-4-yl]thiophen-3-yl}benzoyl)piperidin-3-amine;

1-{4-[5-(2-tert-butylpyridin-4-yl)-2-methylthiophen-3-yl]-3-chlorobenzoyl}piperidin-3-amine;

4-[5-(2-tert-butylpyridin-4-yl)-2-methylthiophen-3-yl]-3-chloro-N-phenylbenzamide;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-ol;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-ol;

tert-butyl N-[(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

tert-butyl N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-3-yl)-N-methylcarbamate;

tert-butyl 3-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamido}piperidine-1-carboxylate;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-amine;

6-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-5-chloro-N-phenylpyridine-3-carboxamide;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-methylpiperidin-3-amine;

N-(1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}piperidin-4-yl)-2,2-dimethylpropanamide;

4-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperazin-2-one;

tert-butyl 4-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzamido}piperidine-1-carboxylate;

tert-butyl N-[(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

tert-butyl N-[(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]carbamate;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-[(4-chlorophenyl)methyl]piperidin-3-amine;

(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-[(4-chlorophenyl)methyl]piperidin-3-amine;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-N-[(4-chlorophenyl)methyl]piperidin-3-amine;

1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-N-[(3-chlorophenyl)methyl]piperidin-3-amine;

(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}pyrrolidin-3-ol;

7-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}-2,7-diazaspiro[4.5]decan-1-one;

2-tert-butyl-4-[4-(2-chloro-4-{octahydropyrrolo[1,2-a]piperazine-2-carbonyl}phenyl)thiophen-2-yl]pyridine;

- 1-{4-[2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl]-3-chlorobenzoyl}piperidin-3-amine;
- 1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-4-ol;
- (3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine;
- (3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)-5-methylthiophen-2-yl]-3-chlorobenzoyl}piperidin-3-amine;
- 1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidine-4-carboxylic acid;
- 1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-ol;
- (3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-ol;
- (3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-ol;
- tert-butyl N-[(3R)-1-{4-[5-(2-tert-butylpyridin-4-yl)thiophen-3-yl]-3-chlorobenzoyl}pyrrolidin-3-yl]carbamate;
- (3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidine-3-carboxylic acid;
- 2-tert-butyl-4-[5-(2-chloro-4-{3H,4H,5H,6H,7H-imidazo[4,5-c]pyridine-5-carbonyl}phenyl)thiophen-3-yl]pyridine;
- 4-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperazin-2-one;
- 1-[(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}piperidin-3-yl]imidazolidin-2-one;
- 1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-amine;
- (3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-amine;
- (3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-amine;
- (3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-amine;
- (3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}pyrrolidin-3-amine;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-1,2,3,4-tetrahydro-1,5-naphthyridine;

1-{4-[2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl]-3-chlorobenzoyl}piperidin-4-ol;

2-tert-butyl-4-{5-[2-chloro-4-(3-methoxypiperidine-1-carbonyl)phenyl]thiophen-3-yl}pyridine;

1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-4-methylpiperidin-4-ol;

(3R)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-ol;

(3S)-1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorobenzoyl}-3-methylpiperidin-3-ol;

4-(3-chloro-4-{5-[1-(3-methylbutyl)-1H-pyrazol-5-yl]thiophen-3-yl}benzoyl)morpholine;

1-(3-chloro-4-{5-[1-(3-methylbutyl)-1H-pyrazol-5-yl]thiophen-3-yl}benzoyl)-4,4-difluoropiperidine;

1-(3-chloro-4-{4-[1-(3-methylbutyl)-1H-pyrazol-5-yl]thiophen-2-yl}benzoyl)piperidin-3-ol; and

3-(1-{4-[4-(2-tert-butylpyridin-4-yl)thiophen-2-yl]-3-chlorophenyl}-N-(4-fluorophenyl)formamido)propanoic acid.

35. A compound selected from the group consisting of:

piperidin-1-yl(4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)methanone;

piperidin-1-yl(3-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)methanone;

(4-(5-(1-isobutyl-1H-pyrazol-5-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)thiophen-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(1,2-dimethyl-1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(1-propyl-1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

- (3-chloro-4-(5-(2,3-dimethyl-3H-imidazo[4,5-b]pyridin-7-yl)thiophen-3-yl)phenyl)
(piperidin-1-yl)methanone;
- (3-chloro-4-(5-(3-propyl-3H-imidazo[4,5-b]pyridin-7-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;
- (4-(5-(3H-imidazo[4,5-b]pyridin-7-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
- (3-chloro-4-(5-(1,2-dipropyl-1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)phenyl)
(piperidin-1-yl)methanone;
- (4-(5-(6-(butylamino)-2-isobutylpyrimidin-4-yl)thiophen-3-yl)-3-chlorophenyl)
(piperidin-1-yl)methanone;
- (3-chloro-4-(5-(6-(dibutylamino)-2-isobutylpyrimidin-4-yl)thiophen-3-yl)phenyl)
(piperidin-1-yl)methanone;
- (4-(5-(6-amino-2-isobutylpyrimidin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
- (4-(5-(2-aminopyrimidin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
- (3-chloro-4-(5-(2-(propylamino)pyrimidin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;
- (4-(5-(2-(butyl(propyl)amino)pyrimidin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;
- (3-chloro-4-(5-(2-propylthiazol-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;
- (3-chloro-4-(5-(5-propyl-1,2,4-thiadiazol-3-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;
- (3-chloro-4-(5-(5-propyl-1,2,4-oxadiazol-3-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;
- (3-chloro-4-(5-(2-propylthiazol-5-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;
- (3-chloro-4-(5-(5-propyl-1,3,4-thiadiazol-2-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;
- (3-chloro-4-(5-(5-propyl-1,3,4-oxadiazol-2-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

5-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)-1-isobutylpyridin-2(1H)-one;

(3-chloro-4-(5-(2-morpholinopyridin-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(2-bromo-3-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3,5-dichloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-(trifluoromethyl)phenyl)(piperidin-1-yl)methanone;

(2-cyclopropyl-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-methylphenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-isopropylphenyl)(piperidin-1-yl)methanone;

(2-ethynyl-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-3-ethynyl-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-6-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-6-methylphenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)-1-methyl-1H-imidazol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(2-(2-isobutylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-3-chloro-4-(2-(2-isobutylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-5-chloro-4-(2-(2-isobutylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)phenyl)
(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)oxazol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(2-(2-tert-butylpyridin-4-yl)oxazol-4-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(dimethylamino)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

(3,5-dichloro-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

N-(4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)pyridin-2-yl)
methanesulfonamide;

(3-chloro-4-(5-(2-(3,3,3-trifluoropropyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-
yl)methanone;

(3-chloro-4-(1-methyl-2-(2-(3,3,3-trifluoropropyl)pyridin-4-yl)-1H-imidazol-4-yl)phenyl)
(piperidin-1-yl)methanone;

(2-hydroxy-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-2-hydroxyphenyl)(piperidin-1-yl)
methanone;

(2-methoxy-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-methoxyphenyl)(piperidin-1-yl)
methanone;

(4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-2-methoxyphenyl)(piperidin-1-yl)
methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-3-(trifluoromethyl)phenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(5-(2-neopentylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-5-(trifluoromethyl)phenyl)
(piperidin-1-yl)methanone;

(2,3-dichloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

(4-(2-(2-isobutylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)-2-methoxyphenyl)(piperidin-1-yl)methanone;

(2-methoxy-4-(1-methyl-2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(4-(2-(2-tert-butylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)-2-methoxyphenyl)(piperidin-1-yl)methanone;

(4-(5-(2-benzylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(2-aminopyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

N-(4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)-1-methyl-1H-imidazol-2-yl)pyridin-2-yl)methanesulfonamide;

(2-hydroxy-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(3,3,3-trifluoropropyl)pyridin-4-yl)furan-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-methoxy-4-(5-(2-neopentylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

N-(4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)pyridin-2-yl)-1,1,1-trifluoromethanesulfonamide;

1-(4-(4-(4-(2-chloro-4-(4,4-difluoropiperidine-1-carbonyl)phenyl)thiophen-2-yl)pyridin-2-yl)piperazin-1-yl)ethanone;

(3-chloro-4-(5-(1-isopentyl-1H-pyrazol-5-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(1-isopentyl-1H-pyrazol-5-yl)furan-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(cyclohexylmethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(naphthalen-1-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(1H-indol-6-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(1H-indol-4-yl)thiophen-3-yl)-3-chlorophenyl)(4,4-difluoropiperidin-1-yl)methanone;

(4-(5-(1H-benzo[d]imidazol-4-yl)thiophen-3-yl)-3-chlorophenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(5-(isoquinolin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(4-methylpiperazin-1-yl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(3,5-dimethylisoxazol-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(1-isopentyl-1H-pyrazol-5-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(5-(1-ethyl-1H-indol-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(4-(5-(1H-indol-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(1-ethyl-1H-benzo[d]imidazol-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(5-(2,3-dihydrobenzofuran-5-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(hydroxymethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(3-(hydroxymethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(piperidin-1-yl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(3-methylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(piperidin-1-yl)pyridin-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(5-(5-fluoro-1H-pyrrolo[2,3-b]pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(benzo[b]thiophen-3-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(3-(pyrrolidin-1-ylmethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(cyclopentylamino)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(1H-indol-3-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(quinolin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-chloropyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(5-(2-tert-butylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(5-(2-tert-butylpyridin-4-yl)furan-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(5-(2-tert-butylpyridin-4-yl)-1H-pyrrol-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl)-3-fluorophenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)-1H-imidazol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(2-(2-(piperidin-1-yl)pyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(4-(2-(2-tert-butylpyridin-4-yl)-1-methyl-1H-imidazol-4-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(2-tert-butylpyridin-4-yl)-1H-pyrrol-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)furan-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)-1H-pyrrol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)-1-methyl-1H-pyrrol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(4-(2-tert-butylpyridin-4-yl)thiazol-2-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(2-tert-butylpyridin-4-yl)isothiazol-3-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(5-(2-tert-butylpyridin-4-yl)isoxazol-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(5-(2-tert-butylpyridin-4-yl)-1H-pyrazol-3-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(3-(2-tert-butylpyridin-4-yl)isothiazol-5-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(3-(2-tert-butylpyridin-4-yl)isoxazol-5-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(4-(3-(2-tert-butylpyridin-4-yl)-1H-pyrazol-5-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(4-(3-(2-tert-butylpyridin-4-yl)-1-methyl-1H-pyrazol-5-yl)-3-chlorophenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(5-(2-isobutylpyridin-4-yl)-1H-pyrrol-3-yl)phenyl)(piperidin-1-yl)
methanone;

(3-fluoro-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(2-(2-isobutylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(2-(2-isobutylpyridin-4-yl)oxazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(2-(2-isobutylpyridin-4-yl)thiazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(3-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(2-(2-isobutylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)
methanone;

(4-(5-(2-butylpyridin-4-yl)thiophen-3-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-(methoxymethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

(3-chloro-4-(5-(4,6-dipropylpyridin-2-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)
methanone;

piperidin-1-yl(4-(5-(4-propylpyridin-2-yl)thiophen-3-yl)phenyl)methanone;

(4-(5-(2,6-dipropylpyridin-3-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-(cyclohexylmethyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-3-methoxyphenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-((trifluoromethylsulfonyl)methyl)pyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(1-methyl-2-(1H-pyrrolo[2,3-b]pyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-propyl-3H-imidazo[4,5-b]pyridin-7-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)-N-cyclopropylpicolinamide;

4-(4-(2-chloro-4-(piperidine-1-carbonyl)phenyl)thiophen-2-yl)-N-cyclohexylpicolinamide;

(4-(2-(1H-indol-4-yl)-1-methyl-1H-imidazol-4-yl)-3-chlorophenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(4-(2-(piperidin-1-yl)pyridin-4-yl)thiophen-2-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(4-(2-(piperidin-1-yl)pyridin-4-yl)thiophen-2-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-4-(2-(2-tert-butylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-chloro-4-(2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(2-bromo-5-chloro-4-(2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)methanone;

(3-chloro-4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(4,4-difluoropiperidin-1-yl)methanone;

(3-chloro-4-(5-(2-cyclopropylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

piperidin-1-yl(4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)-3-(trifluoromethyl)phenyl)
methanone;

(3-chloro-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(3,3-difluoropyrrolidin-1-yl)
methanone;

(3-chloro-4-(2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)
methanone;

(2-bromo-4-(5-(2-propylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

(4-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone;

5-(5-(2-isobutylpyridin-4-yl)thiophen-3-yl)-2-(piperidine-1-carbonyl)benzonitrile;

piperidin-1-yl(4-(5-(6-propylpyridin-3-yl)thiophen-3-yl)phenyl)methanone;

(4-(5-(2-pentylpyridin-4-yl)thiophen-3-yl)phenyl)(piperidin-1-yl)methanone; and

(2-bromo-4-(1-methyl-2-(2-propylpyridin-4-yl)-1H-imidazol-4-yl)phenyl)(piperidin-1-yl)
methanone.

36. A pharmaceutical composition comprising a compound of any of claims 1-35 and a pharmaceutically acceptable vehicle.
37. A method of treating a metabolic disorder, comprising administering to an individual in need thereof a therapeutically effective amount of a compound of any of claims 1-35.
38. The method of claim 37, wherein the individual is diagnosed with or suspected of having nonalcoholic steatohepatitis (NASH).
39. A method of treating a hyperproliferative disorder, comprising administering to an individual in need thereof a therapeutically effective amount of a compound of any of claims 1-35.
40. The method of claim 39, wherein the hyperproliferative disorder is cancer.
41. The method of claim 40, wherein the cancer is breast cancer, prostate cancer, or ovarian cancer.
42. Use of a compound of any of claims 1-35 in the manufacture of a medicament for treating a metabolic disorder or a hyperproliferative disorder.
43. Use of a compound of any of claims 1-35 to treat a metabolic disorder or a hyperproliferative disorder.

