



(51) International Patent Classification:

C07D 487/04 (2006.01) A61P 35/00 (2006.01)

A61K 31/519 (2006.01)

(21) International Application Number:

PCT/US2016/048698

(22) International Filing Date:

25 August 2016 (25.08.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/210,264 26 August 2015 (26.08.2015) US

(71) Applicant: BLUEPRINT MEDICINES CORPORATION [US/US]; 38 Sidney Street, Suite 200, Cambridge, MA 02139 (US).

(72) Inventors: WENGLOWSKY, Steven, Mark; 266 Harvard Street, Apt 6, Cambridge, MA 02139 (US). BROOIJMANS, Natasja; 43 Concord Square, Apt 2, Boston, MA 02118 (US). MIDUTURU, Chandrasekhar, V.; 348 Har-

vard Street, Apt 2, Cambridge, MA 02138 (US). BIFULCO, Neil; 15 Lillian Ave, Sudbury, MA 01776 (US).

(74) Agent: MCCARTY, Catherine, M.; Lando & Anastasi, LLP, Riverfront Office Park, One Main Street, Suite 1100, Cambridge, MA 02142 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,

[Continued on next page]

(54) Title: COMPOUNDS AND COMPOSITIONS USEFUL FOR TREATING DISORDERS RELATED TO NTRK

FIGURE 1

#	Structure	NMR; LCMS
1		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.13-7.02 (m, 3H), 4.66 (s, 2H), 4.17 (d, 2H, J = 3.2 Hz), 2.32-2.29 (m, 1H), 2.11-2.08 (m, 1H), 1.91-1.87 (m, 2H), 1.86-1.74 (m, 2H); LCMS: 389.1
2		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.319-7.15 (m, 1H), 7.02-7.01 (m, 1H), 6.99-6.88 (m, 1H), 4.61 (s, 2H), 4.38-4.31 (m, 1H), 4.13-4.11 (m, 1H), 4.03-4.02 (m, 1H), 2.44-2.41 (m, 1H), 2.34 (s, 3H), 2.16-2.13 (m, 1H), 1.79-1.70 (m, 2H); LCMS: 401.1
3		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.21 (s, 1H), 7.15-7.09 (m, 2H), 7.08-7.01 (m, 1H), 4.65 (s, 2H), 4.41-4.38 (m, 1H), 4.03-3.96 (m, 1H), 2.41-2.36 (m, 1H), 1.90-1.89 (m, 1H), 1.58-1.56 (m, 1H), 1.44-1.41 (m, 1H), 0.76-0.73 (m, 1H), 0.50-0.47 (m, 1H); LCMS: 401.1
4		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.12 (s, 1H), 7.06-6.99 (m, 2H), 6.93-6.92 (m, 1H), 4.56 (s, 2H), 4.32-4.29 (m, 1H), 3.93-3.87 (m, 1H), 2.32-2.25 (m, 1H), 1.81-1.80 (m, 1H), 1.49-1.47 (m, 1H), 1.35-1.33 (m, 1H), 0.67-0.64 (m, 1H), 0.41-0.40 (m, 1H); LCMS: 401.1

(57) Abstract: This invention relates to inhibitors of NTRK that are active against wild-type NTRK and its resistant mutants.



DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT,
LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE,
SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

— *as to the applicant's entitlement to claim the priority of
the earlier application (Rule 4.17(iii))*

Published:

— *with international search report (Art. 21(3))*

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii))*

COMPOUNDS AND COMPOSITIONS USEFUL FOR TREATING DISORDERS RELATED TO NTRK

CLAIM OF PRIORITY

This application claims priority from U.S.S.N. 62/210,264 filed August 26, 2015, which is incorporated herein by reference in its entirety.


BACKGROUND

Neurotrophic Tyrosine Receptor Kinase (NTRK) 1, 2 and 3 are receptor tyrosine kinases (RTKs) that activate multiple downstream pathways involved in cell proliferation and survival. Various genetic fusions, arising from aberrant chromosomal translocations of the genes coding for these RTKs, are implicated in the etiology of multiple cancers including high and low grade glioma, cholangiocarcinoma, papillary thyroid carcinoma, colon cancer and non-small cell lung cancer. A genomics analysis on the landscape of kinase fusions identified NTRK fusions in a wide array of additional cancer types including head and neck squamous cell carcinoma, pancreatic adenocarcinoma, sarcoma and melanoma, thereby providing further therapeutic rationale for deploying inhibitors of these kinases to treat multiple oncologic indications.

The identification of NTRK fusions as the underlying cause of certain cancers prompted the discovery and clinical development of several NTRK kinase inhibitors to treat tumors that harbor an NTRK fusion protein. Early clinical data support the viability of this approach in providing benefit to patients with specific human malignancies. Ultimately however, despite clear signs of clinical activity, most patients' cancers will become resistant to kinase inhibitor therapy leading to relapse and progression of the disease. Kinase reactivation via an intrinsic mutation is a frequent mechanism of resistance. When resistance occurs, the patient's treatment options are often very limited. There is thus a need for compounds that inhibit NTRK, as well as its resistant mutants.

SUMMARY OF THE INVENTION

The invention features compounds and pharmaceutical compositions comprising compounds of Formula (I) or pharmaceutically acceptable salts thereof, wherein:



Formula (I)

Rings A and B are each independently selected from aryl, heteroaryl, cycloalkyl and heterocyclyl;

each L¹ and L² is independently selected from a bond, -C(O)-, -N(R¹)-, -N(R¹)-C(O)-, -C(O)-N(R¹)-, -(C₁-C₆ alkylene)-N(R¹)-, -N(R¹)-(C₁-C₆ alkylene)-, -N(R¹)-C(O)-(C₁-C₆ alkylene)-, and -C(O)-N(R¹)-(C₁-C₆ alkylene)-; wherein each alkylene, is independently substituted with 0-5 occurrences of R⁷;

each R^A and R^B is independently selected from hydroxyl, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, halo, C₁-C₆ heteroalkyl, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, C₁-C₆ hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, heterocyclylalkyl, nitro, cyano, -C(O)R¹, -OC(O)R¹, -C(O)OR¹, -(C₁-C₆ alkylene)-C(O)R¹, -SR¹, -S(O)₂R¹, -S(O)₂-N(R¹)(R¹), -(C₁-C₆ alkylene)-S(O)₂R¹, -(C₁-C₆ alkylene)-S(O)₂-N(R¹)(R¹), -N(R¹)(R¹), -C(O)-N(R¹)(R¹), -N(R¹)-C(O)R¹, -N(R¹)-C(O)OR¹, -(C₁-C₆ alkylene)-N(R¹)-C(O)R¹, -N(R¹)S(O)₂R¹, and -P(O)(R¹)(R¹); wherein each of alkyl, alkenyl, alkynyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, and heterocyclylalkyl is independently substituted with 0-5 occurrences of R^a; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^a;

each R¹ is independently selected from hydrogen, hydroxyl, halo, thiol, C₁-C₆ alkyl, C₁-C₆ thioalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, C₁-C₆ hydroxyalkyl, C₁-C₆ heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, wherein each of alkyl, thioalkyl, alkoxy, haloalkyl, hydroxyalkyl, heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl is independently substituted with 0-5 occurrences of R^b, or 2 R¹ together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^b;

each R^a and R^b is independently selected from C₁-C₆ alkyl, halo, hydroxyl, C₁-C₆ haloalkyl,

C₁-C₆ heteroalkyl, C₁-C₆ hydroxyalkyl, C₁-C₆ alkoxy, cycloalkyl, heterocyclyl, and cyano, wherein each of alkyl, haloalkyl, heteroalkyl, hydroxyalkyl, alkoxy, cycloalkyl and heterocyclyl is independently substituted with 0-5 occurrences of R';

each R' is independently selected from C₁-C₆ alkyl, C₁-C₆ heteroalkyl, halo, hydroxyl, C₁-C₆ haloalkyl, C₁-C₆ hydroxyalkyl, cycloalkyl and cyano; or 2 R' together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring;

p is 0, 1, 2, 3, 4, or 5; and

q is 0, 1, 2, 3, or 4.

Any of the compounds disclosed herein may be used, alone or in combination with another therapeutic agent, to treat any of the diseases disclosed herein.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the structure of various exemplary compounds of the invention, as well as their NMR peaks and mass as determined by LC-MS.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used herein, the terms a "patient," "subject," "individual," and "host" refer to either a human or a non-human animal suffering from or suspected of suffering from a disease or disorder associated with aberrant NTRK expression (*i.e.*, increased NTRK activity caused by signaling through NTRK) or biological activity.

"Treat" and "treating" such a disease or disorder refers to ameliorating at least one symptom of the disease or disorder. These terms, when used in connection with a condition such as a cancer, refer to one or more of: impeding growth of the cancer, causing the cancer to shrink by weight or volume, extending the expected survival time of the patient, inhibiting tumor growth, reducing tumor mass, reducing size or number of metastatic lesions, inhibiting the development of new metastatic lesions, prolonging survival, prolonging progression-free survival, prolonging time to progression, and/or enhancing quality of life.

The term "preventing" when used in relation to a condition or disease such as cancer, refers to a reduction in the frequency of, or delay in the onset of, symptoms of the condition or disease. Thus, prevention of cancer includes, for example, reducing the number of detectable cancerous

growths in a population of patients receiving a prophylactic treatment relative to an untreated control population, and/or delaying the appearance of detectable cancerous growths in a treated population versus an untreated control population, *e.g.*, by a statistically and/or clinically significant amount.

The term “therapeutic effect” refers to a beneficial local or systemic effect in animals, particularly mammals, and more particularly humans, caused by administration of a compound or composition of the invention. The phrase “therapeutically-effective amount” means that amount of a compound or composition of the invention that is effective to treat a disease or condition caused by over expression of NTRK or aberrant NTRK biological activity at a reasonable benefit/risk ratio. The therapeutically effective amount of such substance will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of skill in the art.

As used herein, “developing resistance” means that when a drug is first administered to the patient, the patient’s symptoms improve, whether measured by decrease in tumor volume, a decrease in the number of new lesions, or some other means that a physician uses to judge disease progression; however, those symptoms stop improving, or even worsen at some point. At that time, the patient is said to have developed resistance to the drug.

“Aliphatic group” means a straight-chain, branched-chain, or cyclic hydrocarbon group and includes saturated and unsaturated groups, such as an alkyl group, an alkenyl group, and an alkynyl group.

“Alkylene” refers to a divalent radical of an alkyl group, *e.g.*, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, and $\text{CH}_2\text{CH}_2\text{CH}_2-$.

“Alkenyl” means an aliphatic group containing at least one double bond.

“Alkoxy” or “alkoxy” means an alkyl group having an oxygen radical attached thereto. Representative alkoxy groups include methoxy, ethoxy, propyloxy, tert-butoxy and the like. The term “haloalkoxy” refers to an alkoxy in which one or more hydrogen atoms are replaced by halo, and includes alkoxy moieties in which all hydrogens have been replaced by halo (*e.g.*, perfluoroalkoxy).

“Alkyl” refers to a monovalent radical of a saturated straight or branched hydrocarbon, such

as a straight or branched group of 1-12, 1-10, or 1-6 carbon atoms, referred to herein as C₁-C₁₂ alkyl, C₁-C₁₀ alkyl, and C₁-C₆ alkyl, respectively. Exemplary alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl, etc.

“Alkenylene” refers to an alkenyl group having two connecting points. For example, “ethenylene” represents the group -CH=CH-. Alkenylene groups can also be in an unsubstituted form or substituted form with one or more substituents.

“Alkynyl” refers to a straight or branched hydrocarbon chain containing 2-12 carbon atoms and characterized in having one or more triple bonds. Examples of alkynyl groups include, but are not limited to, ethynyl, propargyl, and 3-hexynyl. One of the triple bond carbons may optionally be the point of attachment of the alkynyl substituent.

“Alkynylene” refers to an alkynyl having two connecting points. For example, “ethynylene” represents the group -C≡C-. Alkynylene groups can also be in an unsubstituted form or substituted form with one or more substituents.

“Hydroxyalkylene” or “hydroxyalkyl” refers to an alkylene or alkyl moiety in which an alkylene or alkyl hydrogen atom is replaced by a hydroxyl group. Hydroxyalkylene or hydroxyalkyl includes groups in which more than one hydrogen atom has been replaced by a hydroxyl group.

“Aromatic ring system” is art-recognized and refers to a monocyclic, bicyclic or polycyclic hydrocarbon ring system, wherein at least one ring is aromatic.

“Aryl” refers to a monovalent radical of an aromatic ring system. Representative aryl groups include fully aromatic ring systems, such as phenyl, naphthyl, and anthracenyl, and ring systems where an aromatic carbon ring is fused to one or more non-aromatic carbon rings, such as indanyl, phthalimidyl, naphthimidyl, or tetrahydronaphthyl, and the like.

“Arylalkyl” or “aralkyl” refers to an alkyl moiety in which an alkyl hydrogen atom is replaced by an aryl group. Aralkyl includes groups in which more than one hydrogen atom has been replaced by an aryl group. Examples of “arylalkyl” or “aralkyl” include benzyl, 2-phenylethyl,

3-phenylpropyl, 9-fluorenyl, benzhydryl, and trityl groups.

“Aryloxy” refers to -O-(aryl), wherein the heteroaryl moiety is as defined herein.

“Halo” refers to a radical of any halogen, *e.g.*, -F, -Cl, -Br, or -I.

“Haloalkyl” and “haloalkoxy” refers to alkyl and alkoxy structures that are substituted with one or more halo groups or with combinations thereof. For example, the terms “fluoroalkyl” and “fluoroalkoxy” include haloalkyl and haloalkoxy groups, respectively, in which the halo is fluorine.

“Haloalkylene” refers to a divalent alkyl, *e.g.*, -CH₂-, -CH₂CH₂-, and -CH₂CH₂CH₂-, in which one or more hydrogen atoms are replaced by halo, and includes alkyl moieties in which all hydrogens have been replaced by halo.

“Heteroalkyl” refers to an optionally substituted alkyl, which has one or more skeletal chain atoms selected from an atom other than carbon, *e.g.*, oxygen, nitrogen, sulfur, phosphorus or combinations thereof. A numerical range may be given, *e.g.* C₁-C₆ heteroalkyl which refers to the number of carbons in the chain, which in this example includes 1 to 6 carbon atoms. For example, a -CH₂OCH₂CH₃ radical is referred to as a “C₃” heteroalkyl. Connection to the rest of the molecule may be through either a heteroatom or a carbon in the heteroalkyl chain. “Heteroalkylene” refers to a divalent optionally substituted alkyl, which has one or more skeletal chain atoms selected from an atom other than carbon, *e.g.*, oxygen, nitrogen, sulfur, phosphorus or combinations thereof.

“Carbocyclic ring system” refers to a monocyclic, bicyclic or polycyclic hydrocarbon ring system, wherein each ring is either completely saturated or contains one or more units of unsaturation, but where no ring is aromatic.

“Carbocyclyl” refers to a monovalent radical of a carbocyclic ring system. Representative carbocyclyl groups include cycloalkyl groups (*e.g.*, cyclopentyl, cyclobutyl, cyclopentyl, cyclohexyl and the like), and cycloalkenyl groups (*e.g.*, cyclopentenyl, cyclohexenyl, cyclopentadienyl, and the like).

“Cycloalkyl” refers to a cyclic, bicyclic, tricyclic, or polycyclic non-aromatic hydrocarbon groups having 3 to 12 carbons. Any substitutable ring atom can be substituted (*e.g.*, by one or more substituents). The cycloalkyl groups can contain fused or spiro rings. Fused rings are rings that share a common carbon atom. Examples of cycloalkyl moieties include, but are not limited to, cyclopropyl, cyclohexyl, methylcyclohexyl, adamantyl, and norbornyl. In some embodiments, the cycloalkyl is bicyclo[3.1.0]hexanyl.

“Cycloalkylalkyl” refers to a –(cycloalkyl)-alkyl radical where cycloalkyl and alkyl are as disclosed herein. The “cycloalkylalkyl” is bonded to the parent molecular structure through the cycloalkyl group.

“Heteroaromatic ring system” is art-recognized and refers to monocyclic, bicyclic or polycyclic ring system wherein at least one ring is both aromatic and comprises at least one heteroatom (*e.g.*, N, O or S); and wherein no other rings are heterocyclyl (as defined below). In certain instances, a ring which is aromatic and comprises a heteroatom contains 1, 2, 3, or 4 ring heteroatoms in such ring.

“Heteroaryl” refers to a monovalent radical of a heteroaromatic ring system. Representative heteroaryl groups include ring systems where (i) each ring comprises a heteroatom and is aromatic, *e.g.*, imidazolyl, oxazolyl, thiazolyl, triazolyl, pyrrolyl, furanyl, thiophenyl pyrazolyl, pyridinyl, pyrazinyl, pyridazinyl, pyrimidinyl, indoliziny, purinyl, naphthyridinyl, and pteridinyl; (ii) each ring is aromatic or carbocyclyl, at least one aromatic ring comprises a heteroatom and at least one other ring is a hydrocarbon ring or *e.g.*, indolyl, isoindolyl, benzothienyl, benzofuranyl, dibenzofuranyl, indazolyl, benzimidazolyl, benzthiazolyl, quinolyl, isoquinolyl, cinnoliny, phthalazinyl, quinazolinyl, quinoxalinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, pyrido[2,3-*b*]-1,4-oxazin-3-(4H)-one, 5,6,7,8-tetrahydroquinoliny and 5,6,7,8-tetrahydroisoquinoliny; and (iii) each ring is aromatic or carbocyclyl, and at least one aromatic ring shares a bridgehead heteroatom with another aromatic ring, *e.g.*, 4H-quinoliziny.

“Heterocyclic ring system” refers to monocyclic, bicyclic and polycyclic ring systems where at least one ring is saturated or partially unsaturated (but not aromatic) and comprises at least one heteroatom. A heterocyclic ring system can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure and any of the ring atoms can be optionally substituted.

“Heterocyclyl” refers to a monovalent radical of a heterocyclic ring system. Representative heterocyclyls include ring systems in which (i) every ring is non-aromatic and at least one ring comprises a heteroatom, *e.g.*, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothienyl, pyrrolidinyl, pyrrolidonyl, piperidinyl, pyrrolinyl, decahydroquinoliny, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, diazepinyl, oxazepinyl, thiazepinyl, morpholiny, and quinuclidinyl; (ii) at least one ring is non-aromatic and comprises a heteroatom and at least one other ring is an aromatic carbon ring, *e.g.*, 1,2,3,4-tetrahydroquinoliny, 1,2,3,4-tetrahydroisoquinoliny; and (iii) at least one ring is

non-aromatic and comprises a heteroatom and at least one other ring is aromatic and comprises a heteroatom, *e.g.*, 3,4-dihydro-1H-pyrano[4,3-c]pyridine, and 1,2,3,4-tetrahydro-2,6-naphthyridine.

“Heterocyclalkyl” refers to an alkyl group substituted with a heterocycl group.

“Cyano” refers to a –CN radical.

“Nitro” refers to –NO₂.

“Hydroxy” or “hydroxyl” refers to –OH.

“Hydroxyalkylene” refers to a divalent alkyl, *e.g.*, –CH₂–, –CH₂CH₂–, and –CH₂CH₂CH₂–, in which one or more hydrogen atoms are replaced by a hydroxy, and includes alkyl moieties in which all hydrogens have been replaced by hydroxy.

“Substituted”, whether preceded by the term “optionally” or not, means that one or more hydrogens of the designated moiety are replaced with a suitable substituent. Unless otherwise indicated, an “optionally substituted” group may have a suitable substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at each position. Combinations of substituents envisioned under this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term “stable”, as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, in certain embodiments, their recovery, purification, and use for one or more of the purposes disclosed herein.

As used herein, the definition of each expression, *e.g.*, alkyl, m, n, etc., when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

Certain compounds of the present invention may exist in particular geometric or stereoisomeric forms. The present invention contemplates all such compounds, including *cis*- and *trans*-isomers, *R*- and *S*-enantiomers, diastereomers, (*D*)-isomers, (*L*)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

If, for instance, a particular enantiomer of compound of the present invention is desired, it may be prepared by asymmetric synthesis, or by derivation with a chiral auxiliary, where the

resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

Unless otherwise indicated, when a disclosed compound is named or depicted by a structure without specifying the stereochemistry and has one or more chiral centers, it is understood to represent all possible stereoisomers of the compound, as well as enantiomeric mixtures thereof. The “enantiomeric excess” or “% enantiomeric excess” of a composition can be calculated using the equation shown below. In the example shown below a composition contains 90% of one enantiomer, *e.g.*, the S enantiomer, and 10% of the other enantiomer, *i.e.*, the R enantiomer.

$$ee = (90-10)/100 = 80\%.$$

Thus, a composition containing 90% of one enantiomer and 10% of the other enantiomer is said to have an enantiomeric excess of 80%.

The compounds or compositions described herein may contain an enantiomeric excess of at least 50%, 75%, 90%, 95%, or 99% of one form of the compound, *e.g.*, the S-enantiomer. In other words such compounds or compositions contain an enantiomeric excess of the S enantiomer over the R enantiomer.


The compounds described herein may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example deuterium (^2H), tritium (^3H), carbon-13 (^{13}C), or carbon-14 (^{14}C). All isotopic variations of the compounds disclosed herein, whether radioactive or not, are intended to be encompassed within the scope of the present invention. In addition, all tautomeric forms of the compounds described herein are intended to be within the scope of the invention.

The compound can be useful as the free base or as a salt. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts

and the like. (See, for example, Berge et al. (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19.)

Compounds

The invention features compounds of Formula (I), or a stereoisomer, enantiomer, tautomer, or isotopically labeled form thereof, or a pharmaceutically acceptable salt of any of the foregoing, wherein:



Formula (I)

Rings A and B are each independently selected from aryl, heteroaryl, cycloalkyl and heterocyclyl;

each L^1 and L^2 is independently selected from a bond, $-C(O)-$, $-N(R^1)-$, $-N(R^1)-C(O)-$, $-C(O)-N(R^1)-$, $-(C_1-C_6 \text{ alkylene})-N(R^1)-$, $-N(R^1)-(C_1-C_6 \text{ alkylene})-$, $-N(R^1)-C(O)-(C_1-C_6 \text{ alkylene})-$, and $-C(O)-N(R^1)-(C_1-C_6 \text{ alkylene})-$; wherein each alkylene, is independently substituted with 0-5 occurrences of R^1 ;

each R^A and R^B is independently selected from hydroxyl, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_1-C_6 alkoxy, halo, C_1-C_6 heteroalkyl, C_1-C_6 haloalkyl, C_1-C_6 haloalkoxy, C_1-C_6 hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, heterocyclylalkyl, nitro, cyano, $-C(O)R^1$, $-OC(O)R^1$, $-C(O)OR^1$, $-(C_1-C_6 \text{ alkylene})-C(O)R^1$, $-SR^1$, $-S(O)_2R^1$, $-S(O)_2-N(R^1)(R^1)$, $-(C_1-C_6 \text{ alkylene})-S(O)_2R^1$, $-(C_1-C_6 \text{ alkylene})-S(O)_2-N(R^1)(R^1)$, $-N(R^1)(R^1)$, $-C(O)-N(R^1)(R^1)$, $-N(R^1)-C(O)R^1$, $-N(R^1)-C(O)OR^1$, $-(C_1-C_6 \text{ alkylene})-N(R^1)-C(O)R^1$, $-N(R^1)S(O)_2R^1$, and $-P(O)(R^1)(R^1)$; wherein each of alkyl, alkenyl, alkynyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, and heterocyclylalkyl is independently substituted with 0-5 occurrences of R^a ; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^a ;

each R^1 is independently selected from hydrogen, hydroxyl, halo, thiol, C_1-C_6 alkyl, C_1-C_6 thioalkyl, C_1-C_6 alkoxy, C_1-C_6 haloalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, wherein each of alkyl,

thioalkyl, alkoxyl, haloalkyl, hydroxyalkyl, heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl is independently substituted with 0-5 occurrences of R^b , or 2 R^1 together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^b ;

each R^a and R^b is independently selected from C_1 - C_6 alkyl, halo, hydroxyl, C_1 - C_6 haloalkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 alkoxyl, cycloalkyl, heterocyclyl, and cyano, wherein each of alkyl, haloalkyl, heteroalkyl, hydroxyalkyl, alkoxyl, cycloalkyl and heterocyclyl is independently substituted with 0-5 occurrences of R' ;

each R' is independently selected from C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, halo, hydroxyl, C_1 - C_6 haloalkyl, C_1 - C_6 hydroxyalkyl, cycloalkyl and cyano; or 2 R' together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring; and


p is 0, 1, 2, 3, 4, or 5; and

q is 0, 1, 2, 3, or 4.

In some embodiments, Ring A is cycloalkyl. In some embodiments, Ring A is a 5-membered or 6-membered cycloalkyl ring. In some embodiments, Ring A is cyclopentyl or cyclohexyl. In some embodiments, Ring A is heterocyclyl. In some embodiments, Ring A is a 5-membered or 6-membered heterocyclyl. In some embodiments, Ring A is tetrahydropyranyl, tetrahydrofuranyl, or pyrrolidinyl. In some embodiments, Ring A is a cycloalkenyl ring. In some embodiments, Ring A is cyclopentenyl.

In some embodiments, Ring B is aryl. In some embodiments, Ring B is phenyl. In some embodiments, Ring B is heteroaryl. In some embodiments, Ring B is pyridyl. In some embodiments, Ring B is heterocyclyl. In some embodiments, Ring B is pyrrolidinyl.

In some embodiments, L^1 is a bond, $-C(O)-$, or $-N(R^1)-$; and L^2 is $-N(R^1)-C(O)-(C_1-C_6$ alkylene)- or $-C(O)-N(R^1)-(C_1-C_6$ alkylene)-. In some embodiments, L^1 is $-NH-$ and L^2 is $-C(O)-NH-CH(CH_2OH)-*$, $-C(O)-N(CH_3)-CH_2-*$, $-C(O)-N(CH_3)-CH(CH_3)-*$, $-C(O)N(CH_2CH_3)CH_2-*$, -

$C(O)NHCH(CH_3)-*$, $-C(O)N(CD_3)CH_2-*$, $-C(O)NHCH(CF_3)-*$, and , wherein "*" represents a portion of L^2 bound to ring B. In some embodiments, L^1 is $-NH-$, L^2 is $-C(O)-$ and ring B is pyrrolidinyl. In some embodiments, L^1 is $-NH-$, L^2 is $-C(O)-$ and ring B is pyrrolidin-1-yl.

In some embodiments, each R^1 is independently selected from hydrogen and C_1 - C_6 alkyl

substituted with 0-5 occurrences of R^b . In some embodiments, each R^1 is independently selected from hydrogen and $-CH_3$.

In some embodiments, each R^A and R^B is independently selected from hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halo, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, C_1 - C_6 hydroxyalkyl, cycloalkyl, aryl, heteroaryl, nitro, cyano, $-C(O)R^1$, $-OC(O)R^1$, $-C(O)OR^1$, $-SR^1$, $-S(O)_2R^1$, $-S(O)_2-N(R^1)(R^1)$, $-N(R^1)(R^1)$, $-C(O)-N(R^1)(R^1)$, $-N(R^1)-C(O)R^1$, $-N(R^1)-C(O)OR^1$, and $-N(R^1)S(O)_2R^1$; wherein each of alkyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, and heteroaryl, is independently substituted with 0-5 occurrences of R^a ; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^a .

In some embodiments, each R^A is independently selected from hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halo, $-C(O)-N(R^1)(R^1)$, $-C(O)OR^1$, $-S(O)_2R^1$, and C_1 - C_6 haloalkyl. In some embodiments, each R^A is additionally and independently selected from $-CN$, oxetanyl, and C_1 - C_6 hydroxyalkyl, or two R^A bound to adjacent ring carbon atoms on ring A are taken together to form a C_3 - C_6 cycloalkyl fused to ring A. In some embodiments each R^A is independently selected from hydroxyl, fluoro, oxetan-3-yl, $-CHF_2$, $-CH_2CH_3$, $-C(CH_3)_2OH$, $-OCH_3$, $-C(O)N(CH_3)_2$, $-C(O)OCH_3$, $-S(O)_2CH_3$; or two R^A bound to adjacent ring carbon atoms on ring A are taken together to form a cyclopropyl fused to ring A.

In some embodiments, each R^B is independently selected from halo, C_1 - C_6 alkyl, cyano, C_1 - C_6 alkoxy, aryl, heteroaryl, and C_1 - C_6 haloalkoxy. In some embodiments, each R^B is additionally selected from oxo.


In some embodiments, Ring B is pyrrolidinyl and at least one R^B is optionally substituted aryl or heteroaryl. In some embodiments, Ring B is pyrrolidinyl and at least one R^B is optionally substituted phenyl or pyridyl.

In some embodiments, Ring B is pyrrolidinyl, and at least one R^B is selected from 2,3,5-trifluorophenyl, 2,3-difluorophenyl, 2,5-difluorophenyl, 2-chloro-5-fluorophenyl, 2-chloro-5-fluoropyridin-3-yl, 2-cyano-5-fluorophenyl, 2-fluoro-5-chlorophenyl, 2-methoxy-3,5-difluorophenyl, 2-methoxy-5-fluoropyridin-3-yl, 2-trifluoromethoxy-5-fluorophenyl, 3,5-difluorophenyl, 3-chloro-5-fluorophenyl, 3-cyano-5-fluorophenyl, 3-difluoromethoxy-5-fluorophenyl, 3-fluorophenyl, 5-fluoropyridin-3-yl, and phenyl.

In some embodiments, Ring B is pyrrolidinyl and one additional R^B , if present, is fluoro.

In some embodiments, Ring B is other than pyrrolidinyl, and each R^B is independently selected from chloro, fluoro, oxo, $-CH_3$, $-CF_3$, $-CN$, $-OCH_3$, $-OCF_3$, and $-OCHF_2$.

In another aspect, the invention features compounds of Formula (Ia):



(Ia), or a stereoisomer, enantiomer, tautomer, or isotopically labeled form thereof, or a pharmaceutically acceptable salt of any of the foregoing, wherein:

Ring A is selected from aryl, heteroaryl, cycloalkyl and heterocyclyl;

L^1 is selected from a bond, $-C(O)-$, $-N(R^1)-$, $-N(R^1)-C(O)-$, $-C(O)-N(R^1)-$, $-(C_1-C_6 \text{ alkylene})-N(R^1)-$, $-N(R^1)-(C_1-C_6 \text{ alkylene})-$, $-N(R^1)-C(O)-(C_1-C_6 \text{ alkylene})-$, and $-C(O)-N(R^1)-(C_1-C_6 \text{ alkylene})-$; wherein each alkylene, is independently substituted with 0-5 occurrences of R^1 ;

each R^A and R^B is independently selected from hydroxyl, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_1-C_6 alkoxy, halo, C_1-C_6 heteroalkyl, C_1-C_6 haloalkyl, C_1-C_6 haloalkoxy, C_1-C_6 hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, heterocyclylalkyl, nitro, cyano, $-C(O)R^1$, $-OC(O)R^1$, $-C(O)OR^1$, $-(C_1-C_6 \text{ alkylene})-C(O)R^1$, $-SR^1$, $-S(O)_2R^1$, $-S(O)_2-N(R^1)(R^1)$, $-(C_1-C_6 \text{ alkylene})-S(O)_2R^1$, $-(C_1-C_6 \text{ alkylene})-S(O)_2-N(R^1)(R^1)$, $-N(R^1)(R^1)$, $-C(O)-N(R^1)(R^1)$, $-N(R^1)-C(O)R^1$, $-N(R^1)-C(O)OR^1$, $-(C_1-C_6 \text{ alkylene})-N(R^1)-C(O)R^1$, $-N(R^1)S(O)_2R^1$, and $-P(O)(R^1)(R^1)$; wherein each of alkyl, alkenyl, alkynyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, and heterocyclylalkyl is independently substituted with 0-5 occurrences of R^A ; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^A ;

each R^1 is independently selected from hydrogen, hydroxyl, halo, thiol, C_1-C_6 alkyl, C_1-C_6 thioalkyl, C_1-C_6 alkoxy, C_1-C_6 haloalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, wherein each of alkyl, thioalkyl, alkoxy, haloalkyl, hydroxyalkyl, heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl is independently substituted with 0-5 occurrences of R^B , or 2 R^1 together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring

independently substituted with 0-5 occurrences of R^b;

each R^a and R^b is independently selected from C₁-C₆ alkyl, halo, hydroxyl, C₁-C₆ haloalkyl, C₁-C₆ heteroalkyl, C₁-C₆ hydroxyalkyl, C₁-C₆ alkoxy, cycloalkyl, heterocyclyl, and cyano, wherein each of alkyl, haloalkyl, heteroalkyl, hydroxyalkyl, alkoxy, cycloalkyl and heterocyclyl is independently substituted with 0-5 occurrences of R['];

each R['] is independently selected from C₁-C₆ alkyl, C₁-C₆ heteroalkyl, halo, hydroxyl, C₁-C₆ haloalkyl, C₁-C₆ hydroxyalkyl, cycloalkyl and cyano; or 2 R['] together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring; and

p is 0, 1, 2, 3, 4, or 5; and

q is 0, 1, 2, 3, or 4.

In some embodiments, Ring A is cycloalkyl. In some embodiments, Ring A is a 5-membered or 6-membered cycloalkyl ring. In some embodiments, Ring A is cyclopentyl or cyclohexyl. In some embodiments, Ring A is heterocyclyl. In some embodiments, Ring A is a 5-membered or 6-membered heterocyclyl. In some embodiments, Ring A is tetrahydropyran, tetrahydrofuran, or pyrrolidinyl. In some embodiments, Ring A is a cycloalkenyl ring. In some embodiments, Ring A is cyclopentenyl.

In some embodiments, L¹ is a bond, -C(O)-, or -N(R¹)-. In some embodiments, L¹ is -NH-.

In some embodiments, each R¹ is independently selected from hydrogen and C₁-C₆ alkyl substituted with 0-5 occurrences of R^b. In some embodiments, each R¹ is independently selected from hydrogen and -CH₃.

In some embodiments, each R^A and R^B is independently selected from hydroxyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halo, C₁-C₆ heteroalkyl, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, C₁-C₆ hydroxyalkyl, cycloalkyl, aryl, heteroaryl, nitro, cyano, -C(O)R¹, -OC(O)R¹, -C(O)OR¹, -SR¹, -S(O)₂R¹, -S(O)₂-N(R¹)(R¹), -N(R¹)(R¹), -C(O)-N(R¹)(R¹), -N(R¹)-C(O)R¹, -N(R¹)-C(O)OR¹, and -N(R¹)S(O)₂R¹; wherein each of alkyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, and heteroaryl, is independently substituted with 0-5 occurrences of R^a; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^a.

In some embodiments, each R^A is independently selected from hydroxyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halo, -C(O)-N(R¹)(R¹), -C(O)OR¹, -S(O)₂R¹, and C₁-C₆ haloalkyl. In some embodiments,

each R^A is additionally and independently selected from -CN, oxetanyl, and C_1 - C_6 hydroxyalkyl, or two R^A bound to adjacent ring carbon atoms on ring A are taken together to form a C_3 - C_6 cycloalkyl fused to ring A. In some embodiments each R^A is independently selected from hydroxyl, fluoro, oxetan-3-yl, $-\text{CHF}_2$, $-\text{CH}_2\text{CH}_3$, $-\text{C}(\text{CH}_3)_2\text{OH}$, $-\text{OCH}_3$, $-\text{C}(\text{O})\text{N}(\text{CH}_3)_2$, $-\text{C}(\text{O})\text{OCH}_3$, $-\text{S}(\text{O})_2\text{CH}_3$; or two R^A bound to adjacent ring carbon atoms on ring A are taken together to form a cyclopropyl fused to ring A.

In some embodiments, each R^B is independently selected from halo, C_1 - C_6 alkyl, cyano, C_1 - C_6 alkoxy, aryl, heteroaryl, and C_1 - C_6 haloalkoxy. In some embodiments, each R^B is additionally selected from oxo.


In some embodiments, when ring B is pyrrolidinyl, at least one R^B is selected from 2,3,5-trifluorophenyl, 2,3-difluorophenyl, 2,5-difluorophenyl, 2-chloro-5-fluorophenyl, 2-chloro-5-fluoropyridin-3-yl, 2-cyano-5-fluorophenyl, 2-fluoro-5-chlorophenyl, 2-methoxy-3,5-difluorophenyl, 2-methoxy-5-fluoropyridin-3-yl, 2-trifluoromethoxy-5-fluorophenyl, 3,5-difluorophenyl, 3-chloro-5-fluorophenyl, 3-cyano-5-fluorophenyl, 3-difluoromethoxy-5-fluorophenyl, 3-fluorophenyl, 5-fluoropyridin-3-yl, and phenyl.

In some embodiments, when ring B is pyrrolidinyl one additional R^B , if present, is fluoro.

In some embodiments, p is 0, 1 or 2.

In some embodiments, q is 1, 2 or 3.

In another aspect, the invention features compounds of Formula (II):



(II), or a stereoisomer, enantiomer, tautomer, or

isotopically labeled form thereof, or a pharmaceutically acceptable salt of any of the foregoing, wherein:

Rings A and B are each independently selected from aryl, heteroaryl, cycloalkyl and heterocyclyl;

each R^A and R^B is independently selected from hydroxyl, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_6 alkoxy, halo, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, C_1 - C_6 hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, heterocyclylalkyl, nitro,

cyano, $-C(O)R^1$, $-OC(O)R^1$, $-C(O)OR^1$, $-(C_1-C_6 \text{ alkylene})-C(O)R^1$, $-SR^1$, $-S(O)_2R^1$, $-S(O)_2-N(R^1)(R^1)$, $-(C_1-C_6 \text{ alkylene})-S(O)_2R^1$, $-(C_1-C_6 \text{ alkylene})-S(O)_2-N(R^1)(R^1)$, $-N(R^1)(R^1)$, $-C(O)-N(R^1)(R^1)$, $-N(R^1)-C(O)R^1$, $-N(R^1)-C(O)OR^1$, $-(C_1-C_6 \text{ alkylene})-N(R^1)-C(O)R^1$, $-N(R^1)S(O)_2R^1$, and $-P(O)(R^1)(R^1)$; wherein each of alkyl, alkenyl, alkynyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, and heterocyclalkyl is independently substituted with 0-5 occurrences of R^a ; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocycl ring independently substituted with 0-5 occurrences of R^a ;

each R^1 is independently selected from hydrogen, hydroxyl, halo, thiol, C_1-C_6 alkyl, C_1-C_6 thioalkyl, C_1-C_6 alkoxy, C_1-C_6 haloalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocycl, and heterocyclalkyl, wherein each of alkyl, thioalkyl, alkoxy, haloalkyl, hydroxyalkyl, heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocycl, and heterocyclalkyl is independently substituted with 0-5 occurrences of R^b , or 2 R^1 together with the atom(s) to which they are attached form a cycloalkyl or heterocycl ring independently substituted with 0-5 occurrences of R^b ;

R^{1a} is selected from hydrogen, C_1-C_6 alkyl, and deuterated C_1-C_6 alkyl;

R^{1b} is selected from hydrogen and C_1-C_6 alkyl;

each R^a and R^b is independently selected from C_1-C_6 alkyl, halo, hydroxyl, C_1-C_6 haloalkyl, C_1-C_6 heteroalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 alkoxy, cycloalkyl, heterocycl, and cyano, wherein each of alkyl, haloalkyl, heteroalkyl, hydroxyalkyl, alkoxy, cycloalkyl and heterocycl is independently substituted with 0-5 occurrences of R' ;

each R' is independently selected from C_1-C_6 alkyl, C_1-C_6 heteroalkyl, halo, hydroxyl, C_1-C_6 haloalkyl, C_1-C_6 hydroxyalkyl, cycloalkyl and cyano; or 2 R' together with the atom(s) to which they are attached form a cycloalkyl or heterocycl ring; and

p is 0, 1, 2, 3, 4, or 5; and

q is 0, 1, 2, 3, or 4.

In some embodiments, Ring A is cycloalkyl. In some embodiments, Ring A is a 5-membered or 6-membered cycloalkyl ring. In some embodiments, Ring A is cyclopentyl or cyclohexyl. In some embodiments, Ring A is heterocycl. In some embodiments, Ring A is a 5-membered or 6-membered heterocycl. In some embodiments, Ring A is tetrahydropyran,

tetrahydrofuran, or pyrrolidinyl. In some embodiments, Ring A is a cycloalkenyl ring. In some embodiments, Ring A is cyclopentenyl.

In some embodiments, Ring B is aryl. In some embodiments, Ring B is phenyl. In some embodiments, Ring B is heteroaryl. In some embodiments, Ring B is pyridyl.

In some embodiments, each R^1 is independently selected from hydrogen and C_1 - C_6 alkyl substituted with 0-5 occurrences of R^b .

In some embodiments, R^{1a} is hydrogen, $-CH_3$, $-CD_3$, or $-CH_2CH_3$.

In some embodiments, R^{1b} is hydrogen.

In some embodiments, each R^A and R^B is independently selected from hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halo, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, C_1 - C_6 hydroxyalkyl, cycloalkyl, aryl, heteroaryl, nitro, cyano, $-C(O)R^1$, $-OC(O)R^1$, $-C(O)OR^1$, $-SR^1$, $-S(O)_2R^1$, $-S(O)_2N(R^1)(R^1)$, $-N(R^1)(R^1)$, $-C(O)-N(R^1)(R^1)$, $-N(R^1)-C(O)R^1$, $-N(R^1)-C(O)OR^1$, and $-N(R^1)S(O)_2R^1$; wherein each of alkyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, and heteroaryl, is independently substituted with 0-5 occurrences of R^a ; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^a .

In some embodiments, each R^A is independently selected from hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halo, $-C(O)-N(R^1)(R^1)$, $-C(O)OR^1$, $-S(O)_2R^1$, and C_1 - C_6 haloalkyl. In some embodiments, each R^A is additionally and independently selected from $-CN$, oxetanyl, and C_1 - C_6 hydroxyalkyl, or two R^A bound to adjacent ring carbon atoms on ring A are taken together to form a C_3 - C_6 cycloalkyl fused to ring A. In some embodiments each R^A is independently selected from hydroxyl, fluoro, oxetan-3-yl, $-CHF_2$, $-CH_2CH_3$, $-C(CH_3)_2OH$, $-OCH_3$, $-C(O)N(CH_3)_2$, $-C(O)OCH_3$, $-S(O)_2CH_3$; or two R^A bound to adjacent ring carbon atoms on ring A are taken together to form a cyclopropyl fused to ring A.

In some embodiments, each R^B is independently selected from halo, C_1 - C_6 alkyl, cyano, C_1 - C_6 alkoxy, aryl, heteroaryl, and C_1 - C_6 haloalkoxy. In some embodiments, each R^B is additionally selected from oxo. In some embodiments, each R^B is independently selected from chloro, fluoro, oxo, $-CH_3$, $-CF_3$, $-CN$, $-OCH_3$, $-OCF_3$, and $-OCHF_2$.

In some embodiments, each R' is independently selected from C_1 - C_6 alkyl, C_1 - C_6 haloalkyl and C_1 - C_6 hydroxyalkyl; or 2 R' together with the atom(s) to which they are attached form a

cycloalkyl or heterocyclcyl ring. In some embodiments one R' is hydrogen, and the other R' is selected from hydrogen, C₁-C₆ alkyl, C₁-C₆ haloalkyl and C₁-C₆ hydroxyalkyl; or 2 R' together with the atom(s) to which they are attached form a cycloalkyl ring. In some embodiments one R' is hydrogen, and the other R' is selected from hydrogen, -CH₂OH, -CH₃, or -CF₃, or 2 R' together with the atom(s) to which they are attached form a cycloprop-1,1-diyl ring.

In some embodiments, p is 0, 1 or 2.

In some embodiments, q is 0, 1, 2 or 3.

Although, as indicated above, various embodiments and aspects thereof for a variable in Formula (I), (Ia), or (II), may be selected from a group of chemical moieties, the invention also encompasses as further embodiments and aspects thereof situations where such variable is: a) selected from any subset of chemical moieties in such a group; and b) any single member of such a group.

Although various embodiments and aspects thereof are set forth (or implied, as discussed in the preceding paragraph) individually for each variable in Formula (I), (Ia) and (II), the invention encompasses all possible combinations of the different embodiments and aspects for each of the variables in Formula (I), (Ia), and (II).

The structures, as well as the NMR and LCMS data of exemplary compounds of the invention are shown in Figure 1. In certain embodiments, the compound of the invention is selected from the group consisting of any one of the compounds in Figure 1 and pharmaceutically acceptable salts, solvates, hydrates, tautomers, stereoisomers, and isotopically labeled derivatives thereof.

The invention also features pharmaceutical compositions containing a pharmaceutically acceptable carrier and any compound of Formulas (I), (Ia) and (II).

Pharmaceutically acceptable salts of these compounds are also contemplated for the uses described herein.

"Pharmaceutically acceptable salt" refers to any salt of a compound of the invention which retains its biological properties and which is not toxic or otherwise undesirable for pharmaceutical use. Pharmaceutically acceptable salts may be derived from a variety of organic and inorganic counter-ions well known in the art and include. Such salts include: (1) acid addition salts formed with organic or inorganic acids such as hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, sulfamic, acetic, trifluoroacetic, trichloroacetic, propionic, hexanoic, cyclopentylpropionic, glycolic,

glutaric, pyruvic, lactic, malonic, succinic, sorbic, ascorbic, malic, maleic, fumaric, tartaric, citric, benzoic, 3-(4-hydroxybenzoyl)benzoic, picric, cinnamic, mandelic, phthalic, lauric, methanesulfonic, ethanesulfonic, 1,2-ethane-disulfonic, 2-hydroxyethanesulfonic, benzenesulfonic, 4-chlorobenzenesulfonic, 2-naphthalenesulfonic, 4-toluenesulfonic, camphoric, camphorsulfonic, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic, glucoheptonic, 3-phenylpropionic, trimethylacetic, tert-butylacetic, lauryl sulfuric, gluconic, benzoic, glutamic, hydroxynaphthoic, salicylic, stearic, cyclohexylsulfamic, quinic, muconic acid and the like acids; or (2) salts formed when an acidic proton present in the parent compound either (a) is replaced by a metal ion, *e.g.*, an alkali metal ion, an alkaline earth ion or an aluminum ion, or alkali metal or alkaline earth metal hydroxides, such as sodium, potassium, calcium, magnesium, aluminum, lithium, zinc, and barium hydroxide, ammonia or (b) coordinates with an organic base, such as aliphatic, alicyclic, or aromatic organic amines, such as ammonia, methylamine, dimethylamine, diethylamine, picoline, ethanolamine, diethanolamine, triethanolamine, ethylenediamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylene-diamine, chlorprocaine, diethanolamine, procaine, N-benzylphenethylamine, N-methylglucamine piperazine, tris(hydroxymethyl)-aminomethane, tetramethylammonium hydroxide, and the like. Pharmaceutically acceptable salts further include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium and the like, and when the compound contains a basic functionality, salts of non-toxic organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, besylate, acetate, maleate, oxalate and the like.

Pharmaceutical Compositions

Pharmaceutical compositions of the invention comprise one or more compounds of the invention and one or more physiologically or pharmaceutically acceptable carrier. The term “pharmaceutically acceptable carrier” refers to a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting any subject composition or component thereof. Each carrier must be “acceptable” in the sense of being compatible with the subject composition and its components and not injurious to the patient. Some examples of materials which may serve as pharmaceutically acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt;

(6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

The compositions of the invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. In some embodiments, the compositions of the invention are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tween, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

The pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

Alternatively, the pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

The pharmaceutically acceptable compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs. Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

For topical applications, the pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutically acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The pharmaceutically acceptable compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in

the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

The amount of the compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the compositions should be formulated so that a dosage of between 0.01-100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

Dosages

Toxicity and therapeutic efficacy of compounds of the invention, including pharmaceutically acceptable salts and deuterated variants, can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The LD₅₀ is the dose lethal to 50% of the population. The ED₅₀ is the dose therapeutically effective in 50% of the population. The dose ratio between toxic and therapeutic effects (LD₅₀/ED₅₀) is the therapeutic index. Compounds that exhibit large therapeutic indexes are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

Data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds may lie within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (i.e., the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion,

drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present invention in the composition will also depend upon the particular compound in the composition.

Treatment

NTRK fusions have been implicated in several types of cancers. These fusions harbor an intact NTRK kinase domain that is identical to the native or wild-type form of the receptor; therefore, as used herein, any NTRK protein (NTRK1, 2 or 3) with the same kinase domain as wild-type NTRK will be referred to as “wild-type NTRK.” Mutations can occur in the NTRK kinase domain, leading to mutants that are resistant to kinase inhibitor therapy. These resistance mutations can be predicted using structural biology and computational analyses, as well as by examining codon sequences in which a sequence change gives rise to a codon for a different amino acid. Alternatively, resistance mutations for a given inhibitor can be identified experimentally by administration of that inhibitor (*e.g.*, a known NTRK wild-type inhibitor) and exposing cells to a mutation-promoting agent, such as ENU. The cells are washed, then plated with increasing concentrations (2-100X proliferation IC_{50}) of the compound of choice. The wells with cellular outgrowth are then collected after 3-4 weeks. In particular, a mutation at amino acid position 595 within the NTRK fusion (NTRK1 wt numbering), effecting a change from a glycine to an arginine residue (heretofore designated ‘G595R’) was identified via both methods. This mutation was subsequently demonstrated to confer significant resistance to two NTRK inhibitors that are being clinically evaluated (shown in the table below). As shown in the table, the compounds are active against the wild-type NTRK, but are markedly less active against the G595R mutant form of the NTRK fusion.

Compound	NTRK wt Enzyme Assay IC_{50} (nM)	NTRK wt Cellular GI_{50} (nM)	NTRK G595R Cellular GI_{50} (nM)
Entrectinib	0.6	2	2700
TSR-011	2.3	32	12000
Crizotinib	9.3	87	9000

Accordingly, in another aspect the invention provides a method for treating a subject suffering from a condition mediated by aberrant neurotrophic tyrosine receptor kinase (NTRK)

activity, comprising administering to the subject a therapeutically effective amount of a compound or pharmaceutical composition of a compound described herein.

The invention provides compounds that inhibit both wild-type NTRK and resistant G595R mutants of NTRK.

In another aspect, the invention provides a method for treating a subject who has developed resistance to a cancer treatment, comprising administering to the subject a therapeutically effective amount of a compound or pharmaceutical composition of a compound described herein.

Furthermore, the inhibitors can be selective for wild-type NTRK, over other kinases, thus leading to reduced toxicities associated with inhibiting other kinases. Because of their activity against wild-type and mutant NTRK, the compounds described herein can be used to treat a patient with a condition associated with aberrant NTRK activity. They can also be used to treat various cancers. In some embodiments, the cancer is selected from non-small cell lung cancer, breast cancer, melanoma, low and high grade glioma, glioblastoma, pediatric astrocytoma, colorectal cancer, papillary thyroid carcinoma, pancreatic adenocarcinoma, head and neck cancer, cholangiocarcinoma, acute myelogenous leukemia, secretory breast cancer, salivary cancer and spitzoid neoplasms.

The compounds can also be used to treat a patient who has developed resistance to a wild-type NTRK inhibitor, or a patient with a mutant form of NTRK, such as the G595R mutant. The method includes the step of administering a compound or composition of the invention that is active against the NTRK resistant mutant. By “active” is meant that a compound has an IC_{50} of less than 1 μ M, 500 nM, 250 nM, 100 nM, 75 nM, 50 nM, 25 nM, 10 nM, or 5 nM when measured in a biochemical assay, against at least one resistant mutant.

The compounds and compositions described herein can be administered alone or in combination with other compounds, including other NTRK-modulating compounds, or other therapeutic agents. In some embodiments, the compound or composition of the invention may be administered in combination with one or more compounds selected from Cabozantinib (COMETRIQ), Vandetanib (CALPRESA), Sorafenib (NEXAVAR), Sunitinib (SUTENT), Regorafenib (STAVARGA), Ponatinib (ICLUSIG), Bevacizumab (AVASTIN), Crizotinib (XALKORI), or Gefitinib (IRESSA). The compound or composition of the invention may be administered simultaneously or sequentially with the other therapeutic agent by the same or

different routes of administration. The compound of the invention may be included in a single formulation with the other therapeutic agent or in separate formulations.

Synthesis

Compounds of the invention, including salts and N -oxides thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes, such as those in the Schemes below. The reactions for preparing compounds of the invention can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, *e.g.*, temperatures which can range from the solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected by the skilled artisan.

Preparation of compounds of the invention can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of appropriate protecting groups, can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in Wuts and Greene, *Protective Groups in Organic Synthesis*, 4th ed., John Wiley & Sons: New Jersey, (2006), which is incorporated herein by reference in its entirety.

Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance (NMR) spectroscopy (*e.g.*, ^1H or ^{13}C), infrared (IR) spectroscopy, spectrophotometry (*e.g.*, UV-visible), mass spectrometry (MS), or by chromatographic methods such as high performance liquid chromatography (HPLC) or thin layer chromatography (TLC). Analytical instruments and methods for compound characterization:

LC-MS: Unless otherwise indicated, all liquid chromatography-mass spectrometry (LC-MS) data (sample analyzed for purity and identity) were obtained with an Agilent model-1260 LC system using an Agilent model 6120 mass spectrometer utilizing ES-API ionization fitted with an Agilent Poroshel 120 (EC-C18, 2.7 μm particle size, 3.0 x 50mm dimensions) reverse-phase column at 22.4 degrees Celsius. The mobile phase consisted of a mixture of solvent 0.1% formic acid in

water and 0.1% formic acid in acetonitrile. A constant gradient from 95% aqueous/5% organic to 5% aqueous/95% organic mobile phase over the course of 4 minutes was utilized. The flow rate was constant at 1mL/min.

Prep LC-MS: Preparative HPLC was performed on a Shimadzu Discovery VP® Preparative system fitted with a Luna 5u C18(2) 100A, AXIA packed, 250 x 21.2 mm reverse-phase column at 22.4 degrees Celsius. The mobile phase consisted of a mixture of solvent 0.1% formic acid in water and 0.1% formic acid in acetonitrile. A constant gradient from 95% aqueous/5% organic to 5% aqueous/95% organic mobile phase over the course of 25 minutes was utilized. The flow rate was constant at 20 mL/min. Reactions carried out in a microwave were done so in a Biotage Initiator microwave unit.

Chiral HPLC: Preparative HPLC to resolve chiral mixtures was performed on a Thar SFC Pre-80 instrument fitted with a Chiralpak AS-H column (5 mm, 3.0 cm id x 25 cm L). The mobile phases consisted of SFC CO₂ (A) and MeOH/0.1% NH₄OH (B). A constant gradient from 67% to 33% (B) was maintained at a flow rate of 65 g/min, with a system back pressure of 100 bar. The separation progress was monitored by UV detection at a wavelength of 220 nm.

Silica gel chromatography: Silica gel chromatography was performed on either a Teledyne Isco CombiFlash® Rf unit or a Biotage® Isolera Four unit.


Proton NMR: Unless otherwise indicated, all ¹H NMR spectra were obtained with a Varian 400MHz Unity Inova 400 MHz NMR instrument (acquisition time = 3.5 seconds with a 1 second delay; 16 to 64 scans). Where characterized, all protons were reported in DMSO-d₆ solvent as parts-per million (ppm) with respect to residual DMSO (2.50 ppm).

EXAMPLES

The following examples are intended to be illustrative, and are not meant in any way to be limiting.


The below Schemes are meant to provide general guidance in connection with preparing the compounds of the invention. One skilled in the art would understand that the preparations shown in the Schemes can be modified or optimized using general knowledge of organic chemistry to prepare various compounds of the invention.

General Synthesis 1:




For certain compounds, the general synthesis begins with appropriate nitrogen-protection (P) of a di-halide substituted 1H-pyrazolo[3,4-d]pyrimidine. The nitrogen-protected bicycle can be substituted at the halide on the pyrimidine ring with an appropriately substituted Ring A under appropriate conditions, for example, nucleophilic aromatic substitution reaction conditions, using a base, such as diisopropylethylamine (DIPEA), in a polar solvent such as dioxane to provide the bicycle substituted with Ring A. The halide of the pyrazole ring can be substituted under Palladium-mediated carbonyl insertion reaction conditions followed by hydrolysis to provide the resultant carboxylic acid. The carboxylic acid can be reacted with Ring B under appropriate coupling conditions, for example amide coupling reaction conditions, to afford the nitrogen-protected compound substituted with Rings A and B. The removal of the protecting group can afford compounds of Formula I.


Synthetic Protocol 1:



A slightly more specific version of General Synthesis scheme 1 is shown above in Synthetic Protocol 1. The synthetic protocol begins with SEM-group protection of 3-bromo-4-chloro-1H-pyrazolo[3,4-d]pyrimidine **1**. The SEM-protected heterocycle **2** can be substituted with an amino alcohol under nucleophilic aromatic substitution reaction conditions using a base such as diisopropylethylamine (DIPEA) in a polar solvent such as dioxane to provide the amine-substituted heterocycle **3**. The 3-bromo pyrazolo pyrimidine **3** is subjected to a palladium-mediated carbonyl insertion reaction in a DMF-MeOH solvent mixture to afford the methyl ester **4**. Following the hydrolysis of the ester with NaOH treatment, the carboxylic acid is **5** reacted with a benzyl amine or a pyrrolidine under amide coupling reaction conditions to afford the SEM-protected compound **6**. The SEM-protecting group can be removed using TBAF or under acidic conditions to afford the final compound **7**. The compounds described below were prepared using General Synthesis 1, 2 or 3, as further detailed in Synthetic Protocol 1, 2, or 3, respectively.


General Synthesis 2





For certain compounds, the general synthesis begins with appropriate nitrogen-protection (P) of 4-oxo-4,5-dihydro-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid. The nitrogen-protected bicycle can be chlorinated and coupled with an amine in the presence of a chlorinating reagent such as thionyl chloride. The resulting compound can be substituted at the halide on the pyrimidine ring with an appropriately substituted Ring A under appropriate conditions, for example, nucleophilic aromatic substitution reaction conditions, using a base, such as diisopropylethylamine (DIPEA), in a polar solvent such as dioxane to provide the bicycle substituted with Ring A. The removal of the protecting group can afford compounds of Formula I. Compounds described below can be prepared using this general synthesis. Further, chiral HPLC can be employed to resolve chiral mixtures of compounds of Formula I, (Ia), (Ia-1), (Ia-2), (Ib), (Ib-1), (Ib-2), II, (IIa), (IIb), (IIc).


Synthetic Protocol 2



A slightly more specific version of General Synthesis scheme 2 is shown above in Synthetic Protocol 2. The synthetic protocol begins with SEM-protected 4-oxo-4,5-dihydro-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid **1** which can be chlorinate with thionyl chloride/DMF and then coupled with a benzyl amine or a pyrrolidine under mild heat to afford the SEM-protected


compound **2**. The SEM-protected heterocycle **2** can be substituted with an amino alcohol under nucleophilic aromatic substitution reaction conditions using a base such as diisopropylethylamine (DIPEA) in a polar solvent such as dioxane to provide the amine-substituted heterocycle **3**. The SEM-protecting group can be removed using TBAF or under acidic conditions to afford the final compound **4**.

General Synthesis 3





For certain compounds, the general synthesis begins with appropriate nitrogen-protection (P) of 4-oxo-4,5-dihydro-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid. The carboxylic acid can be coupled to an amine using amide coupling conditions. The resulting compound can be chlorinated using thionyl chloride followed by substitution at the chloride on the pyrimidine ring with an appropriately substituted Ring A under appropriate conditions, for example, nucleophilic aromatic substitution reaction conditions, using a base, such as diisopropylethylamine (DIPEA), in a polar solvent such as dioxane to provide the bicyclic substituted with Ring A. The removal of the protecting group can afford compounds of Formula I. Compounds described below can be prepared using this general synthesis. Further, chiral HPLC can be employed to resolve chiral mixtures of compounds of Formula I, (Ia), (Ia-1), (Ia-2), (Ib), (Ib-1), (Ib-2), II, (IIa), (IIb), (IIc).

Synthetic Protocol 3




A slightly more specific version of General Synthesis scheme 3 is shown above in Synthetic Protocol 3. The synthetic protocol begins with SEM-protected 4-oxo-4,5-dihydro-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid **1** which can be coupled with a benzyl amine or a pyrrolidine under amide coupling conditions. The SEM-protected heterocycle **2** can be chlorinated with thionyl chloride/DMF and then substituted with an amino alcohol under nucleophilic aromatic substitution reaction conditions using a base such as diisopropylethylamine (DIPEA) in a polar solvent such as dioxane to provide the amine-substituted heterocycle **3**. The SEM-protecting group can be removed using TBAF or under acidic conditions to afford the final compound **4**.

All of the compounds set forth in Figure 1, as well as other compounds of the invention were prepared using one of three general synthesis schemes and protocols depicted above. Further, chiral HPLC can be employed to resolve chiral mixtures of compounds of Formula I, (Ia), (Ia-1), (Ia-2), (Ib), (Ib-1), (Ib-2), II, (IIa), (IIb), (IIc). Certain specific examples of synthesis are set forth in the Examples.

Example 1. Synthesis of Compound 45**Step 1: Synthesis of 3-bromo-4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine**


To a solution of 3-bromo-4-chloro-1H-pyrazolo[3,4-d]pyrimidine (10.00 g, 42.84 mmol) in DMF (50.00 mL) was added NaH (2.57 g, 64.25 mmol) in portions at 0 °C. After stirring for 0.5 hr, SEM-Cl (8.57 g, 51.40 mmol) was added dropwise to the reaction at 0 °C over 0.5 hr. The reaction was slowly warmed to 25 °C and stirred for 16 hrs. After TLC (PE:EtOAc=1:1, R_f =0.88) showed the reaction was complete, the reaction was slowly quenched by 50 mL of H₂O. The mixture was extracted with EtOAc (50 mL*3), and the organic layer was washed with brine (20 mL*3), dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography on silica gel (PE:EtOAc=20:1) to give 3-bromo-4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine (6.20 g, yield: 39.80%) as a colorless oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 8.83 (s, 1H), 5.82 (s, 2H), 3.70 (t, 2H, J = 8.4 Hz), 0.97 (t, 2H, J = 8.4 Hz), 0.00 (s, 9H).

Step 2: Synthesis of (1R,2R)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)cyclopentan-1-ol



To a mixture of 3-bromo-4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine (2.50 g, 6.87 mmol) and (1R, 2R)-2-aminocyclopentanol hydrochloride (945.38 mg, 6.87 mmol) in dioxane (15 mL) was added DIPEA (1.78 g, 13.74 mmol), the reaction mixture was allowed to stir at 70 °C for 16 hrs. Once TLC (PE:EtOAc = 5:1) showed the starting material was consumed completely, the mixture was concentrated in vacuum and purified by column chromatography on silica gel (PE:EtOAc = 30:1~10:1) to provide (1R,2R)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)cyclopentan-1-ol (2.10 g, 4.41 mmol, yield: 64.22%) as a yellow oil. ¹H-NMR (400 MHz, CD₃OD) δ ppm 8.41 (s, 1H), 6.26 (s, 1H), 5.70 (s, 2H), 4.13-4.12 (m, 2H), 3.69-3.65 (m, 2H), 2.40-2.39 (m, 1H), 2.20-2.18 (m, 1H), 1.95-1.84 (m, 2H), 1.72-1.61 (m, 2H), 0.97 (d, 2H, *J* = 4.0 Hz), 0.00 (s, 9H).


Step 3: Synthesis of methyl 4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate



To a mixture of (1R,2R)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)cyclopentan-1-ol (2.10 g, 4.90 mmol) in DMF (10 mL) and MeOH (15 mL) was added Pd(dppf)Cl₂ (717.07 mg, 980.00 μmol) and Et₃N (1.49 g, 14.70 mmol) in one portion, and the reaction mixture was allowed to stir at 70 °C for 30 hrs under a CO (50 psi) atmosphere. Once TLC (PE:EtOAc = 1:1) and LCMS showed the starting material was consumed completely, the mixture was filtered and the filtrate was concentrated in vacuum to get methyl 4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate (2.70 g, crude) as a yellow oil, which was used directly without further purification. ¹H-NMR (400 MHz, CDCl₃) δ ppm 8.85 (s, 1H), 8.45 (s, 1H), 5.83 (s, 2H), 4.14 (br.s,


2H), 4.10 (s, 3H), 3.70 (t, 2H, $J = 8.4$ Hz), 2.40-2.38 (m, 1H), 2.20-2.17 (m, 1H), 1.93-1.79 (m, 4H), 0.98 (t, 2H, $J = 8.4$ Hz), 0.00 (s, 9H).

Step 4: Synthesis of 4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid




To a mixture of methyl 4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate (2.70 g, 6.63 mmol) in MeOH (10 mL), H₂O (10 mL) was added, followed by NaOH (530.01 mg, 13.25 mmol) in one portion. The reaction mixture was then allowed to stir at 26 °C for 16 hrs. Once LCMS and TLC (PE:EtOAc =1:1) showed the starting material was consumed completely, MeOH was removed by concentration in vacuum and the residue was washed with EtOAc (8 mL*2). Aqueous HCl (1 N) was then added until the pH < 7 and the formation of white precipitate was observed. The solid was collected by filtration and dried in vacuo to provide 4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (1.20 g, yield: 37.89%) as a white solid. ¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40 (s, 1H), 5.80 (s, 2H), 4.37-4.32 (m, 1H), 4.22-4.19 (m, 1H), 3.75 (t, 2H, $J = 8.0$ Hz), 2.39-2.36 (m, 1H), 2.11-2.06 (m, 1H), 1.96-1.91 (m, 2H), 1.78-1.73 (m, 2H), 0.95 (t, 2H, $J = 8.0$ Hz), 0.00 (s, 9H).

Step 5: Synthesis of ((R)-2-(2,5-difluorophenyl)pyrrolidin-1-yl)(4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone




To a mixture of 4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (100.00 mg, 254.12 μmol) and (2R)-2-(2,5-difluorophenyl)pyrrolidine (55.87 mg, 304.95 μmol) in DMF (2 mL) was added HATU (144.94 mg, 381.18 μmol) and Et_3N (128.57 mg, 1.27 mmol) at 20 °C, and the reaction was stirred at 20 °C for 16 hrs. After LCMS showed the reaction was complete, H_2O (5 mL) was added to the mixture, and the reaction was extracted with EtOAc (10 mL*3) and washed with brine (5 mL*3). The organic layer was then dried over Na_2SO_4 and concentrated. The residue was purified by preparative TLC (PE:EtOAc=1:1, R_f =0.5) to give ((R)-2-(2,5-difluorophenyl)pyrrolidin-1-yl)(4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone (20.00 mg, yield: 14.09%) as a colorless oil.


Step 6: Synthesis of ((R)-2-(2,5-difluorophenyl)pyrrolidin-1-yl)(4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone



To a solution of ((R)-2-(2,5-difluorophenyl)pyrrolidin-1-yl)(4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone (20.00 mg, 35.80 μmol) in dioxane (20 mL) was added TBAF (80.61 mg, 358.00 μmol) at 20 °C, and the reaction was heated at 80 °C for 16 hrs. After TLC (EtOAc, R_f = 0.1) showed the reaction was complete, the solution was concentrated. And 10 mL of H_2O was added to the residue. The solution was extracted with EtOAc (10 mL*3), and the organic layer was dried over Na_2SO_4 and concentrated. The residue was purified by acidic preparative HPLC (MeOH/ H_2O /TFA solvent system) to give ((R)-2-(2,5-difluorophenyl)pyrrolidin-1-yl)(4-(((1R,2R)-2-hydroxycyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone (11.10 mg, yield: 72.37%) as a brown solid.

Example 2. Synthesis of Compound 97 and Compound 98


Step 1: Synthesis of (1R,2R,4R)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-(methylsulfonyl)cyclopentan-1-ol



To a mixture of 3-bromo-4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine (300.00 mg, 824.83 μ mol) in dioxane (10.00 mL) was added DIPEA (319.80 mg, 2.47 mmol) and (1R,2R,4R)-2-amino-4-(methylsulfonyl)cyclopentan-1-ol (195.71 mg, 907.31 μ mol),


and the mixture was stirred at 90 °C for 32 hrs. After LCMS showed the reaction was complete, the mixture was concentrated to remove 1,4-dioxane, and the residue was dissolved in DCM (20 mL). The organic layer was washed with water (10 mL*4), dried over Na₂SO₄, and concentrated to give (1R,2R,4R)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-(methylsulfonyl)cyclopentan-1-ol (350.00 mg, yield: 83.78%) as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 8.43 (s, 1H), 6.23 (br.s, 1H), 5.71 (s, 2H), 4.50 (br.s, 1H), 4.26-4.24 (m, 1H), 3.70-3.66 (m, 3H), 2.97 (br.s, 4H), 2.65 (m, 1H), 2.37-2.20 (m, 2H), 0.97 (t, 2H, *J* = 8.4 Hz), 0.00 (s, 9H).

Step 2: Synthesis of methyl 4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate




To a mixture of (1R,2R,4R)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-(methylsulfonyl)cyclopentan-1-ol (350.00 mg, 691.03 μmol) in MeOH (10.00 mL)/DMF (2.00 mL) was added Et₃N (139.85 mg, 1.38 mmol) and Pd(dppf)Cl₂ (25.28 mg, 34.55 μmol). After addition, the mixture was stirred at 75 °C for 16 hrs under CO (50 Psi). Once LCMS showed the reaction was complete, the mixture was concentrated to give the crude product, which was purified by preparative TLC (PE:EtOAc=0:1) to give methyl 4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate (250.00 mg, yield: 74.50%) as a red solid.

Step 3: Synthesis of 4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid



To a mixture of methyl 4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate (250.00 mg, 514.80 μmol) in MeOH (10.00 mL)/H₂O (5.00 mL) was added NaOH (41.18 mg, 1.03 mmol), which was stirred at 20 °C for 16 hrs. Once LCMS showed the reaction was complete, the mixture was concentrated to remove MeOH. The water layer was washed with EtOAc (3 mL*2) and acidified by HCl (1 M) until the pH = 4, after which the mixture was filtered and the filter cake was dried under vacuum to afford 4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (180.00 mg, yield: 74.14%) as a black/brown solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 8.47 (s, 1H), 5.77 (s, 2H), 4.43-4.40 (m, 1H), 4.17-4.12 (m, 1H), 3.85-8.82 (m, 1H), 3.69 (t, 2H, J = 8.4 Hz), 3.03 (s, 3H), 2.45-2.41 (m, 2H), 1.99-1.91 (m, 2H), 0.92 (t, 2H, J = 8.4 Hz), 0.00 (s, 9H).


Step 4: Synthesis of N-(2,5-difluorobenzyl)-4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide



To a mixture of 4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (60.00 mg, 127.23 μmol) in DMF (2.00 mL) was added DIPEA (16.44 mg, 127.23 μmol), (2,5-difluorophenyl)methanamine (36.42 mg, 254.46 μmol) and T₃P (40.48 mg, 127.23 μmol). After addition, the mixture was stirred at 20 °C for 1 hr, wherein LCMS showed the reaction was complete. The mixture was added to water (4 mL) and extracted with EtOAc (5 mL*3), and the


organic layer was dried over Na_2SO_4 and concentrated to give N-(2,5-difluorobenzyl)-4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (50.00 mg, crude) as a red oil.

Step 5: Synthesis of N-(2,5-difluorobenzyl)-4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide




N-(2,5-difluorobenzyl)-4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (50.00 mg, 83.79 μmol) in DCM (5.00 mL) was stirred in a mixture of TFA (5.00 mL) at 20 °C for 16 hrs, after which LCMS showed the reaction was complete. The mixture was concentrated to give the crude product, which was purified by preparative HPLC (TFA) and chiral HPLC (retention times of the resolved isomers were 7.46 min and 9.20 min, respectively). N-(2,5-difluorobenzyl)-4-(((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (2.80 mg, yield: 7.16%) and N-(2,5-difluorobenzyl)-4-(((1S,2S,4S)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (4.00 mg, yield: 10.23%) were obtained as white solids. LC-MS conditions for these compounds were as follows: flow rate = 0.8 mL $\cdot$ min⁻¹, mobile phase: from 99% [water + 0.375% v/v TFA] and 1% [CH_3CN + 0.188% v/v TFA], under this condition for 0.4 min, then changed to 10% [water + 0.375% v/v TFA] and 90% [CH_3CN + 0.188% v/v TFA] in 3.0 min, then changed to 100% [CH_3CN + 0.188% v/v TFA] in 0.45 min, finally changed to 99% [water + 0.375% v/v TFA] and 1% [CH_3CN + 0.188% v/v TFA] in 0.01 min, then under this condition for 0.64 min; 98.887% purity and 96.551%, respectively.

Example 3. Synthesis of Compound 20 and Compound 21




Step 1: (1S,2R,5S)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-5-fluorocyclopentan-1-ol




To a mixture of 3-bromo-4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine (600.00 mg, 1.65 mmol) and 2-amino-5-fluoro-cyclopentanol (196.58 mg, 1.65 mmol) in dioxane (15 mL) was added DIPEA (426.49 mg, 3.30 mmol). The mixture was stirred at 110 °C for 16 hrs, after which TLC (PE/EtOAc =1:1) showed the reaction was complete. The mixture was cooled to 25 °C and concentrated in reduced pressure at 50 °C. To the residue was added EtOAc (50 mL), and the organic phase was washed with H₂O (20 mL*3), dried over Na₂SO₄, filtered and concentrated in vacuum to afford (1S,2R,5S)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-5-fluorocyclopentan-1-ol (600.00 mg, crude). The residue was used directly in next step without further purification.

Step 2: Methyl 4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate




To a solution of (1S,2R,5S)-2-((3-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-5-fluorocyclopentan-1-ol (600.00 mg, 1.34 mmol) in MeOH/DMF (20 mL, v:v = 2/1) was added Pd(dppf)Cl₂ (49.17 mg, 67.21 μ mol) and Et₃N (408.03 mg, 4.03 mmol) under N₂. The suspension was degassed under vacuum and purged with CO several times. The mixture was stirred under CO (50 psi) at 70 °C for 16 hrs, after which TLC (PE/EtOAc = 1:1) showed the starting material was consumed completely. The reaction mixture was filtered and the filtrate was concentrated to afford methyl 4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate (700.00 mg, crude). The crude product was used directly without purification.

Step 3: 4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid




To a solution of methyl 4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylate (700.00 mg, 1.65 mmol) in MeOH/H₂O (15 mL, v/v = 2/1) was added NaOH (132.00 mg, 3.30 mmol) in one portion, which was stirred at 25 °C for 2 hrs. After LCMS showed the reaction was complete, the mixture was concentrated in reduced pressure at 40 °C. The aqueous phase was adjusted to pH = 4 and filtered to afford 4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (700.00 mg, crude) as a white solid.

Step 4: N-(2,5-difluorobenzyl)-4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide



To a mixture of 4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (100.00 mg, 243.01 μmol) and T₃P (231.97 mg, 729.04 μmol) was added Et₃N (49.18 mg, 486.03 μmol) in DMF (2.00 mL) at 25 °C, followed by the addition of (2,5-difluorophenyl)methanamine (69.57 mg, 486.03 μmol) in one portion after 10 min. The mixture was stirred at 25 °C for 16 hrs. After LCMS showed the reaction was complete, the mixture was concentrated under reduced pressure at 60 °C to afford N-(2,5-difluorobenzyl)-4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (200 mg, crude). The residue was not purified and used directly.

Step 5: N-(2,5-difluorobenzyl)-4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide and N-(2,5-difluorobenzyl)-4-(((1S,2R,3R)-3-fluoro-2-hydroxycyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide




A mixture of N-(2,5-difluorobenzyl)-4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (200.00 mg, 372.70 μmol) in TFA/DCM (15.00 mL, v/v = 1/1) was stirred at 25 °C for 3 hrs, then concentrated under reduced pressure at 30 °C. To the residue was added MeOH (20 mL) and KOAc (100 mg), and the mixture was stirred for 16 hrs at 25 °C. Once LCMS showed the reaction was complete, the mixture was concentrated under reduced pressure at 30 °C. The residue was purified by acidic preparative HPLC followed by chiral preparative HPLC to afford N-(2,5-difluorobenzyl)-4-(((1R,2S,3S)-3-fluoro-2-hydroxycyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (25.00 mg, yield: 16.51%) as a white solid and N-(2,5-difluorobenzyl)-4-(((1S,2R,3R)-3-fluoro-2-hydroxycyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (30.00 mg, yield: 19.81%) as a grey solid. LC-MS conditions for these compounds were as follows: flow rate = 0.8 mL·min⁻¹, mobile phase: from 95% [water + 10mM NH₄HCO₃] and 5% CH₃CN, under this condition for 0.4 min, then changed to 10% [water + 10mM NH₄HCO₃] and 90% CH₃CN in 2.6

min, then changed to 100% CH₃CN in 0.85 min, finally changed to 95% [water + 10mM NH₄HCO₃] and 5% CH₃CN in 0.01 min, then under this condition for 0.64 min. 97.125% purity and 97.690% purity, respectively.

Synthesis of Amine Intermediates

Example 4: Synthesis of (1R,2S,3R)-3-aminocyclopentane-1,2-diol



Step 1: (3aS,4S,6aR)-2,2-Dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-ol

To a solution of (3aR,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-one (92.00 g, 596.78 mmol, 1.00 *eq*) in MeOH (2.00 L) was added Pd-C (10%, 12 g). The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (30 psi) at 20 °C for 4 hrs, at which point TLC (PE:EtOAc=3:1) showed the starting material was consumed completely. The reaction mixture was filtered, and to the filtrate was added NaBH₄ (34.09 g, 901.14 mmol, 1.51 *eq*) in portions at 0 °C, and the resulting mixture was stirred at 20 °C for 0.5 hr. The mixture was then concentrated, and to the residue was added H₂O (500 mL). The mixture was extracted with EtOAc (500 mL*3), dried over Na₂SO₄, and concentrated to give (3aS,4S,6aR)-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-ol (84.00 g, yield: 88.98%) as a yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 4.60 (t, 1H, *J* = 5.2 Hz), 4.39 (t, 1H, *J* = 5.6 Hz), 3.83 (br.s, 1H), 2.37-2.35 (m, 1H), 1.88-1.76 (m, 2H), 1.65-1.56 (m, 1H), 1.48 (s, 3H), 1.45-1.36 (m, 1H), 1.33 (s, 3H).

Step 2: 2-((3aS,4R,6aR)-2,2-Dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione

To a stirred mixture of (3aS,4S,6aR)-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-ol (70.00 g, 442.51 mmol, 1.00 *eq*), isoindoline-1,3-dione (80.00 g, 543.74 mmol, 1.23 *eq*) and PPh₃ (175.00 g, 667.20 mmol, 1.51 *eq*) in dry toluene (1.00 L) under N₂ was added DIAD (135.00 g, 667.62 mmol, 1.51 *eq*) dropwise. The resulting mixture was stirred at 80 °C for 20 hrs under N₂. After TLC (PE:EtOAc=3:1) showed the starting material was consumed completely, the mixture was concentrated, and the residue was purified by column chromatography on silica gel (PE:EtOAc = 80:1/50:1/20:1/10:1) to give 2-((3aS,4R,6aR)-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (90.00 g, yield: 70.79%) as a white solid. ¹H-

NMR (400 MHz, CDCl₃) δ ppm 7.82-7.80 (m, 2H), 7.72-7.70 (m, 2H), 5.03-4.96 (m, 2H), 4.61-4.60 (m, 1H), 2.28-2.20 (m, 2H), 1.94-1.85 (m, 2H), 1.50 (s, 3H), 1.31 (s, 3H)


Step 3: (3aS,4R,6aR)-2,2-Dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-amine

A mixture of 2-((3aS,4R,6aR)-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (90.00 g, 313.25 mmol, 1.00 *eq*) and NH₂NH₂·H₂O (32.00 g, 626.50 mmol, 2.00 *eq*) in EtOH (600.00 mL) was stirred at 80 °C for 16 hrs. After TLC (PE:EtOAc = 3:1) showed the starting material was consumed completely, the mixture was filtered and concentrated, and EtOH (500 mL) was added to the residue. After concentration to remove the solvent, PE (1000 mL) was added and the mixture filtered and concentrated to give (3aS,4R,6aR)-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-amine (41.00 g, crude) as a yellow oil, which was solidified by standing as a yellow crystal. ¹H-NMR (400 MHz, CDCl₃) δ ppm 4.72 (t, 1H, *J* = 5.2 Hz), 4.18 (d, 1H, *J* = 5.6 Hz), 3.39 (d, 1H, *J* = 4.0 Hz), 2.01-1.93 (m, 2H), 1.78-1.77 (m, 1H), 1.40 (s, 3H), 1.38-1.35 (m, 1H), 1.26 (s, 3H), 1.10 (br.s, 2H).

Step 4: (1R,2S,3R)-3-Aminocyclopentane-1,2-diol hydrochloride

A mixture of (3aS,4R,6aR)-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-amine (10.00 g, 63.61 mmol, 1.00 *eq*) in H₂O (55.00 mL) and HCl (5.00 mL, 12 M) was stirred at 20 °C for 2 hrs. TLC (EtOAc:MeOH = 10:1) showed the starting material was consumed completely. The mixture was concentrated to give (1R,2S,3R)-3-aminocyclopentane-1,2-diol hydrochloride (9.20 g, yield: 94.15%) as a yellow solid. ¹H-NMR (400 MHz, CD₃OD) δ ppm 4.03 (br.s, 1H), 3.90 (dd, 1H, *J* = 8.4, 4.4 Hz), 3.48-3.41 (m, 1H), 2.25-2.19 (m, 1H), 2.05-2.02 (m, 1H), 1.75-1.65 (m, 1H), 1.58-1.56 (m, 1H).

Example 5: Synthesis of (1R,2R,3S,4R,5S)-4-Aminobicyclo[3.1.0]hexane-2,3-diol



Step 1: (3aS,4S,6aR)-2,2-Dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol

To a 0 °C stirred mixture of (3aR,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-one (90.00 g, 583.81 mmol, 1.00 *eq*) and CeCl₃·7H₂O (240.00 g, 644.16 mmol, 1.10 *eq*) in MeOH (2.00 L) was added NaBH₄ (44.00 g, 1.16 mol, 1.99 *eq*) in portions over 0.5 hr. After addition, the mixture was stirred at 18 °C for 0.5 hr. TLC (PE:EtOAc = 3:1) showed

the starting material was consumed completely. The mixture was concentrated, and to the residue was added EtOAc (2000 mL) and the solution stirred at 18 °C for 0.5 hr. The mixture was then filtered, the filtrate was dried over Na₂SO₄ and concentrated to give the crude product (3aS,4S,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol (79.00 g, crude) as a light yellow oil which was used directly to the next step without further purification. ¹H-NMR (400 MHz, CDCl₃) δ ppm 5.88 (s, 2 H), 5.01 (d, 1H, *J* = 5.6 Hz), 4.74 (t, 1H, *J* = 5.6 Hz), 4.55 (dd, 1H, *J* = 9.6, 5.6 Hz), 2.76 (d, 1H, *J* = 9.6 Hz), 1.43 (s, 3H), 1.40 (s, 3H).

Step 2: 3aR,3bR,4aS,5S,5aS)-2,2-Dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-ol

To a 0°C stirred mixture of (3aS,4S,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol (40.00 g, 256.11 mmol, 1.00 *eq*) in DCM (50.00 mL) was added dropwise ZnEt₂ (1 M, 1.00 L, 3.90 *eq*). After 15 min, CH₂I₂ (550.00 g, 2.05 mol, 8.02 *eq*) was added into the mixture, which was stirred at 20 °C for 16 hrs. TLC (PE:EtOAc = 1:1) showed the starting material was consumed. The mixture was quenched by saturated NH₄Cl solution (200 mL), followed by addition of H₂O (500 mL). The mixture was extracted with DCM (500 mL*5), and the combined organic layers were dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified by column chromatography on silica gel (PE:EtOAc = 0:1/100:1/80:1/50:1/20:1/10:1/5:1) to give (3aR,3bR,4aS,5S,5aS)-2,2-dimethylhexahydrocyclopropa[3,4]-cyclopenta[1,2-d][1,3]dioxol-5-ol (18.00 g, yield: 41.29%) as a light yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 4.87 (t, 1H, *J* = 6.0 Hz), 4.53-4.45 (m, 2H), 2.34 (br.s, 1H), 1.85-1.82 (m, 1H), 1.64-1.62 (m, 1H), 1.54 (s, 3H), 1.28 (s, 3H), 0.98-0.94 (m, 1H), 0.63-0.60 (m, 1H).

Step 3: 2-((3aR,3bR,4aS,5R,5aS)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-yl)isoindoline-1,3-dione

To a stirred mixture of (3aR,3bR,4aS,5S,5aS)-2,2-dimethylhexahydrocyclopropa[3,4]-cyclopenta[1,2-d][1,3]dioxol-5-ol (10.00 g, 58.75 mmol, 1.00 *eq*), isoindoline-1,3-dione (12.00 g, 81.56 mmol, 1.39 *eq*) and PPh₃ (24.00 g, 91.50 mmol, 1.56 *eq*) in dry toluene (500.00 mL) under N₂ was added DIAD (20.00 g, 98.91 mmol, 1.68 *eq*) dropwise. The resulting mixture was stirred at 80 °C for 20 hrs under N₂. TLC (PE:EtOAc = 3:1) showed the starting material was consumed completely. The mixture was concentrated, and the residue was purified by column chromatography on silica gel (PE:EtOAc = 80:1/50:1/30:1/20:1/10:1) to give 2-

((3aR,3bR,4aS,5R,5aS)-2,2-dimethylhexahydrocyclopropa[3,4]-cyclopenta[1,2-d][1,3]dioxol-5-yl)isoindoline-1,3-dione (14.00 g, yield: 79.61%) as a light yellow oil. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 7.86-7.83 (m, 2H), 7.74-7.72 (m, 2H), 5.37-5.34 (m, 1H), 4.78-4.76 (m, 1H), 4.73 (s, 1H), 2.01-1.95 (m, 1H), 1.51 (s, 3H), 1.47-1.42 (m, 1H), 1.24 (s, 3H), 0.85-0.79 (m, 2H).


Step 4: (3aR,3bR,4aS,5R,5aS)-2,2-Dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-amine


A mixture of 2-((3aR,3bR,4aS,5R,5aS)-2,2-dimethylhexahydrocyclopropa[3,4]-cyclopenta[1,2-d][1,3]dioxol-5-yl)isoindoline-1,3-dione (14.00 g, 46.77 mmol, 1.00 *eq*) and $\text{NH}_2\text{NH}_2\cdot\text{H}_2\text{O}$ (4.78 g, 93.54 mmol, 2.00 *eq*) in EtOH (200.00 mL) was stirred at 70 °C for 16 hrs. TLC (PE:EtOAc=3:1) showed the starting material was consumed completely. The mixture was filtered, the filtrate was concentrated, and to the residue was added EtOAc (20 mL). The mixture was then filtered, and concentrated to give (3aR,3bR,4aS,5R,5aS)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-amine (7.00 g, yield: 88.45%) as a yellow oil. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 5.06-5.03 (m, 1H), 4.30 (d, 1H, $J = 6.8$ Hz), 3.45 (s, 1H), 1.73-1.71 (m, 1H), 1.48 (s, 3H), 1.43-1.39 (m, 1H), 1.25 (s, 3H), 0.74-0.67 (m, 2H).

Step 5: (1R,2R,3S,4R,5S)-4-aminobicyclo[3.1.0]hexane-2,3-diol*HCl


A mixture of (3aR,3bR,4aS,5R,5aS)-2,2-dimethylhexahydrocyclopropa[3,4]cyclopenta[1,2-d][1,3]dioxol-5-amine (1.00 g, 63.61 mmol, 1.00 *eq*) in H_2O (5.500 mL) and HCl (0.5 mL, 12 M) was stirred at 15 °C for 2 hrs. TLC (EtOAc:MeOH = 10:1) showed the starting material was consumed completely. The mixture was concentrated to give (1R,2R,3S,4R,5S)-4-aminobicyclo[3.1.0]hexane-2,3-diol*HCl (780 mg g, yield: 79.9%) as a yellow solid.

Example 6: Synthesis of (1S,2S,4S)-2-amino-4-fluorocyclopentan-1-ol (relative stereochemistry)






Step 1: ((Cyclopent-3-en-1-yloxy)methyl)benzene



To a mixture of cyclopent-3-en-1-ol (60.00 g, 713.27 mmol) in THF (600.00 mL) was added NaH (37.09 g, 927.25 mmol) in portions at 0 °C. After effervescence had ceased, bromomethylbenzene (158.59 g, 927.25 mmol) was added dropwise at 0 °C over 45 min period, then warmed to 25 °C and stirred for 16 hrs. TLC (PE/EtOAc = 50/1) showed the reaction was complete. Excess NaH was quenched with MeOH (120 mL) at a temperature below 5 °C. The mixture was warmed to 25 °C, diluted with H₂O (600.00 mL), and the two layers were separated. The aqueous phase was extracted with ethyl acetate (200 mL*3). The combined organic phase was washed with brine (200 mL), dried over Na₂SO₄, filtered and concentrated in vacuum. The residue was purified by silica gel chromatography (PE/EtOAc = 1/0) to afford ((cyclopent-3-en-1-yloxy)methyl)benzene (120.00 g, yield: 96.56%) as yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.38-7.28 (m, 5H), 5.72 (s, 2H), 4.52 (s, 2H), 4.35-4.30 (m, 1H), 2.64-2.59 (m, 2H), 2.50-2.46 (m, 2H).


Step 2: (1R,3S,5S)-3-(benzyloxy)-6-oxabicyclo[3.1.0]hexane and (1R,3r,5S)-3-(benzyloxy)-6-oxabicyclo[3.1.0]hexane



To a mixture of ((cyclopent-3-en-1-yloxy)methyl)benzene (120.00 g, 688.71 mmol) in DCM (600.00 mL), was added *m*-CPBA (297.68 g, 1.38 mol) in one portion at 0 °C. The mixture was

stirred at 25 °C for 16 hrs. TLC (PE:EtOAc=20:1) showed the reaction was complete. The mixture was filtered and excess *m*-CPBA was reduced by the addition of saturated *aq.* Na₂SO₃ until a negative starch iodide test was observed. The mixture was filtered and concentrated in vacuum. The residue was purified by silica gel chromatography (PE:EtOAc=1:0, 20:1) to afford (1R,3R,5S)-3-(benzyloxy)-6-oxabicyclo[3.1.0]hexane (56.00 g, yield: 42.74%) and (1R,3S,5S)-3-(benzyloxy)-6-oxabicyclo[3.1.0]hexane (37.00 g, yield: 28.24%) as yellow oil. Spectra analysis of (1R,3S,5S)-3-(benzyloxy)-6-oxabicyclo[3.1.0]hexane. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.31-7.20 (m, 5H), 4.36 (s, 2H), 3.84-3.74 (m, 1H), 3.43 (s, 2H), 2.51-2.35 (m, 2H), 1.66-1.57 (m, 2H). Spectra analysis of (1R,3R,5S)-3-(benzyloxy)-6-oxabicyclo[3.1.0]hexane ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.31-7.15 (m, 5H), 4.36 (s, 2H), 3.85-3.74 (m, 1H), 3.43 (s, 2H), 2.51-2.35 (m, 2H), 1.66-1.60 (m, 2H).

Step 3: (1S,2S,4R)-2-Azido-4-(benzyloxy)cyclopentan-1-ol




To a mixture of (1R,3S,5S)-3-(benzyloxy)-6-oxabicyclo[3.1.0]hexane (20.00 g, 105.13 mmol) in EtOH (760.00 mL) and H₂O (230.00 mL) was added NH₄Cl (20.98 g, 392.13 mmol), NaN₃ (24.00 g, 369.17 mmol) in one portion at 25 °C. The mixture was heated to 80 °C and stirred for 16 hrs. TLC (PE:EtOAc=10:1) showed the reaction was complete. The mixture was cooled to 25°C and EtOH was removed by N₂, and the aqueous phase was extracted with DCM (100 mL*3). The combined organic phase was washed with H₂O (30 mL*3), dried over Na₂SO₄, filtered and concentrated in vacuum to afford (1S,2S,4R)-2-azido-4-(benzyloxy)cyclopentan-1-ol (23.00 g, yield: 93.79%) as a yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.43-7.28 (m, 5H), 4.58-4.47 (m, 2H), 4.27-4.25 (m, 1H), 4.11-4.08 (m, 1H), 3.66-3.61 (m, 1H), 2.49-2.44 (m, 2H), 2.16-2.13 (m, 1H), 1.89 (br.s, 1H), 1.87-1.80 (m, 2H).

Step 4: (1S,2S,4R)-2-Azido-4-(benzyloxy)cyclopentyl acetate




To a solution of (1S,2S,4R)-2-azido-4-(benzyloxy)cyclopentan-1-ol (22.90 g, 98.17 mmol), Et₃N (59.60 g, 589.02 mmol) in DCM (550 mL) was added a solution of acetylchloride (38.53 g, 490.85 mmol) in DCM (50 mL) dropwise at 0 °C over a period of 30 mins under N₂, during which the temperature was maintained below 5 °C. The reaction mixture was then warmed to 25 °C and stirred for 16 hrs. TLC (PE:tOAc=10:1) showed the starting material was consumed completely. The reaction was quenched by the slow addition of H₂O(100 mL). The organic phase was washed with saturated brine (50 mL), dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by column chromatography on silica gel (PE: EtOAc=100:1, 50:1) to afford (1S,2S,4R)-2-azido-4-(benzyloxy)cyclopentyl acetate (17.00 g, yield: 62.90%) as yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.37-7.28 (m, 5H), 5.24-5.12 (m, 1H), 4.51 (s, 2H), 4.14-4.11 (m, 1H), 3.88-3.85 (m, 1H), 2.45-2.40 (m, 1H), 2.36-2.32 (m, 1H), 2.07 (s, 3H), 1.95-1.88 (m, 2H).

Step 5: (1S,2S,4R)-2-((tert-butoxycarbonyl)amino)-4-Hydroxycyclopentyl acetate




To a solution of (1S,2S,4R)-2-azido-4-(benzyloxy)cyclopentyl acetate (8.80 g, 31.97 mmol) in EtOH (100.00 mL) was added Pd(OH)₂ (4.42 g, 31.97 mmol) under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (50 psi) at 70 °C for 32 hrs. TLC (PE: EtOAc = 2:1) showed the starting material was consumed completely. The reaction mixture was filtered and the filtrate was concentrated. The crude product was purified by silica gel chromatography (PE:EtOAc=10:1, 2:1) to give (1S,2S,4R)-2-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentyl acetate (3.80 g, yield: 45.84%) as a yellow solid.

Step 6: (1S,2S,4S)-2-((tert-butoxycarbonyl)amino)-4-Fluorocyclopentyl acetate



To a mixture of (1S,2S,4R)-2-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentyl acetate (2.57 g, 9.91 mmol) in DCM (150.00 mL) was added DAST (2.40 g, 14.87 mmol) dropwise at -70 °C under N₂. The mixture was stirred at -70 °C for 30 min. TLC (PE:EtOAc=2:1) showed the reaction was complete. The mixture was cooled to 0 °C and *aq.* NaHCO₃ (5 mL, 10%) was added and allowed to stir for 10 min. The aqueous phase was extracted with EtOAc (15 mL*2), and the combined organic phase was washed with brine (10 mL), dried over Na₂SO₄, filtered, and concentrated in vacuum. The crude product was purified by silica gel chromatography (PE:EtOAc=20:1, 10:1) to give (1S,2S,4S)-2-((tert-butoxycarbonyl)amino)-4-fluorocyclopentyl acetate (700.00 mg, yield: 27.03%) as a yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 5.14 (s, 0.5H), 5.00 (br.s, 1H), 4.71 (s, 0.5H), 4.14-4.13 (m, 1H), 2.49-2.47 (m, 2H), 2.07-1.94 (m, 3H), 1.81-1.74 (m, 2H), 1.43-1.41 (m, 9H).

Step 7: tert-butyl ((1S,2S,4S)-4-Fluoro-2-hydroxycyclopentyl)carbamate



To a mixture of (1S,2S,4S)-2-((tert-butoxycarbonyl)amino)-4-fluorocyclopentyl acetate (700.00 mg, 2.68 mmol) in MeOH (20.00 mL) was added NaOH (160.80 mg, 4.02 mmol) in one portion. The mixture was stirred at 25 °C for 1 hr. TLC (PE:EtOAc=3:1) showed the reaction was complete. The mixture was concentrated under reduced pressure at 30 °C to afford tert-butyl ((1S,2S,4S)-4-fluoro-2-hydroxycyclopentyl)carbamate (650.00 mg, crude) as a white solid.


Step 8: (1S,2S,4S)-2-Amino-4-fluorocyclopentan-1-ol (relative stereochemistry)




The mixture of tert-butyl ((1S,2S,4S)-4-fluoro-2-hydroxycyclopentyl)carbamate (650.00 mg, 2.96 mmol) in MeOH/HCl (20.00 mL, 4 M) was stirred for 1 hr at 25 °C. TLC (PE:EtOAc = 2:1)

showed the reaction was complete. The mixture was concentrated under reduced pressure at 30 °C to afford (1S,2S,4S)-2-amino-4-fluorocyclopentan-1-ol (relative stereochemistry) (400.00 mg, yield: 86.85%) as a white solid. ¹H-NMR (400 MHz, DMSO-*d*₆) δ ppm 8.38 (br.s, 3H), 5.09 (d, 1H, *J* = 53.6 Hz), 4.09-4.03 (m, 1H), 3.34-3.30 (m, 1H), 2.44-2.39 (m, 1H), 2.21-2.16 (m, 1H), 1.95-1.87 (m, 1H), 1.75-1.66 (m, 1H).

Example 7: Synthesis (1R,2R,4R)-2-amino-4-(methylsulfonyl)cyclopentan-1-ol (relative stereochemistry)




Step 1: Preparation of (1R,2R,4S)-2-((tert-butoxycarbonyl)amino)-4-((methylsulfonyl)oxy)cyclopentyl acetate




To a mixture of (1R,2R,4S)-2-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentyl acetate (3.00 g, 11.57 mmol) and Et₃N (4.68 g, 46.28 mmol) in DCM (50.00 mL) was added dropwise MsCl (3.98 g, 34.71 mmol) at 0 °C, then the mixture was stirred at 20 °C for 3 hrs. TLC (PE:EtOAc=1:1) showed the reaction was complete. The mixture was washed with water (100 mL*3), then the organic layer was dried over Na₂SO₄ and concentrated to give (1R,2R,4S)-2-((tert-butoxycarbonyl)amino)-4-((methylsulfonyl)oxy)cyclopentyl acetate (3.5 g, crude: 100%) as a yellow solid.

Step 2: (1R,2R,4R)-2-((tert-butoxycarbonyl)amino)-4-(Methylthio)cyclopentyl acetate




To a mixture of (1R,2R,4S)-2-((tert-butoxycarbonyl)amino)-4-((methylsulfonyl)oxy)cyclopentyl acetate (3.50 g, 10.37 mmol) in DMF (30.00 mL) was added NaSMe (4.36 g, 12.45 mmol). The mixture was then stirred at 90 °C for 2 hrs, and TLC (PE:EtOAc = 2:1) showed the reaction was complete. The mixture was concentrated to give the crude (1R,2R,4R)-2-((tert-butoxycarbonyl)amino)-4-(methylthio)cyclopentyl acetate (3.00 g, crude 100%) as a yellow solid.

Step 3: tert-butyl ((1R,2R,4R)-2-Hydroxy-4-(methylthio)cyclopentyl)carbamate



To a mixture of (1R,2R,4R)-2-((tert-butoxycarbonyl)amino)-4-(methylthio)cyclopentyl acetate (3.00 g, 10.37 mmol) in MeOH (100.00 mL) was added K₂CO₃ (2.87 g, 20.74 mmol). The mixture was stirred at 25 °C for 16 hrs. TLC (PE:EtOAc=2:1) showed the reaction was complete. The mixture was concentrated and purified by column chromatography on silica gel (PE:EtOAc = 5:1-1:1) to give tert-butyl ((1R,2R,4R)-2-hydroxy-4-(methylthio)-cyclopentyl)carbamate (1.60 g, yield: 62.38%) as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 4.10-3.99 (m, 2H), 3.85 (br.s, 1H), 3.14-3.11 (m, 1H), 2.50-2.46 (m, 1H), 2.17-2.10 (m, 4H), 1.86-1.82 (m, 1H), 1.81-1.67 (m, 1H), 1.45 (s, 9H).


Step 4: tert-butyl ((1R,2R,4R)-2-Hydroxy-4-(methylsulfonyl)cyclopentyl)carbamate



To a mixture of tert-butyl ((1R,2R,4R)-2-hydroxy-4-(methylthio)cyclopentyl)carbamate (1.60 g, 6.47 mmol) in DCM (100.00 mL) was added *m*-CPBA (3.49 g, 16.18 mmol). The mixture was stirred at 25 °C for 16 hrs. After TLC (PE:EtOAc = 2:1) showed the starting material was consumed completely, the mixture was washed with saturated Na₂SO₃ (aq. 20 mL)


and saturated NaHCO_3 (aq. 20 mL*3). The organic layer was then dried over Na_2SO_4 , concentrated, washed with PE (10 mL), filtered, and the filter cake was dried under vacuum to give tert-butyl ((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)carbamate (1.70 g, yield: 94.06%) as a white solid. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 4.62 (br.s, 1H), 4.07-4.04 (m, 1H), 3.81-3.79 (m, 1H), 3.48-3.45 (m, 1H), 2.82 (s, 3H), 2.57-2.40 (m, 2H), 2.11-2.07 (m, 1H), 1.90-1.87 (m, 2H), 1.38 (s, 9H).

Step 5: (1R,2R,4R)-2-Amino-4-(methylsulfonyl)cyclopentan-1-ol (relative stereochemistry)




The mixture of tert-butyl ((1R,2R,4R)-2-hydroxy-4-(methylsulfonyl)cyclopentyl)carbamate (1.2 g, 4.30 mmol) in HCl/MeOH (10.00 mL) was stirred at 25 °C for 16 hrs, after which LCMS showed the reaction was complete, and the mixture was concentrated to give (1R,2R,4R)-2-amino-4-(methylsulfonyl)cyclopentan-1-ol (1.0 g, crude: 100%) as a yellow solid. $^1\text{H-NMR}$ (400 MHz, CD_3OD) δ ppm 4.11-4.07 (m, 2H), 3.76 (br.s, 1H), 2.94 (s, 3H), 2.53 (br.s, 2H), 1.99 (br.s, 2H).

Example 8: Synthesis of (1S,2R,5R)-2-Amino-5-fluorocyclopentan-1-ol hydrochloride (relative stereochemistry)




Step 1: Cyclopent-2-en-1-ol



To a mixture of $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (24.00 g, 64.42 mmol) in MeOH (120.00 mL) was added cyclopent-2-en-1-one (5.00 g, 60.90 mmol) at 15 °C. After 5 min, NaBH_4 (4.61 g, 121.80


mmol) was added into the mixture in portions at 0 °C. The resulting mixture was stirred at 25 °C for 1 hr, after which TLC (PE:EtOAc=5:1) showed several spots were generated and a part of the starting material was remained. The reaction was quenched by H₂O (100 mL) and the organic solvent was concentrated in vacuum. To the residue was added H₂O (300 mL), followed by extraction with MTBE (200 mL*3). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum to give the crude product cyclopent-2-en-1-ol (3.00 g, crude) as a brown oil. It was used directly to the next step without further purification. ¹H-NMR (400 MHz, CDCl₃) δ ppm 5.91 (d, 1H, *J* = 4.8 Hz), 5.77-5.76 (m, 1H), 4.79 (d, 1H, *J* = 3.6 Hz), 2.47-2.42 (m, 1H), 2.21-2.15 (m, 2H), 1.64-1.59 (m, 1H).

Step 2: ((Cyclopent-2-en-1-yloxy)methyl)benzene



To a mixture of cyclopent-2-en-1-ol (9.00 g, 106.99 mmol) in THF (200.00 mL) was added NaH (6.80 g, 170.11 mmol) in portions at 0 °C. After addition, the mixture was stirred at 20 °C for 0.5 hr, then BnBr (20.00 g, 116.94 mmol) was added into the mixture dropwise at 0 °C. The resulting mixture was stirred at 20 °C for 2 hrs. TLC (PE:EtOAc = 20:1) showed formation of a new species (*R*_f = 0.6, 254 nm). At this point, H₂O (20 mL) was added, followed by extraction with EtOAc (20 mL*3). The combined organics were dried over Na₂SO₄ and concentrated to give the crude product, which was purified by column chromatography on silica gel (PE:EtOAc = 1:0/100:1/80:1) to give ((cyclopent-2-en-1-yloxy)methyl)benzene (8.00 g, yield: 42.91%) as a yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.34-7.25 (m, 5H), 6.02 (br.s, 1H), 5.88 (br.s, 1H), 4.66 (br.s, 1H), 4.55-4.47 (m, 2H), 2.52-4.48 (m, 1H), 2.27-2.24 (m, 1H), 2.16-2.13 (m, 1H), 1.87-1.84 (m, 1H).


Step 3: (1S,2S,5S)-2-(Benzyloxy)-6-oxabicyclo[3.1.0]hexane



To a mixture of ((cyclopent-2-en-1-yloxy)methyl)benzene (8.50 g, 29.27 mmol) in DCM (50.00 mL) was added *m*-CPBA (13.50 g, 58.67 mmol) in portions at 0 °C. The mixture was stirred at 25 °C for 4 hrs. Once TLC (PE:EtOAc=10:1) showed the starting material was consumed


completely, the mixture was filtered, the filtrate was concentrated and purified by column chromatography on silica gel (PE:EtOAc=1:0/100:1/80:1/50:1) to give the crude product. DCM (20 mL) was then added, the mixture filtered, and H₂O (20 mL) and Na₂CO₃ (500 mg) were added to the filtrate, followed by stirring of the mixture at 25 °C for 0.5 hr. The mixture was then extracted with DCM (20 mL*3), dried over Na₂SO₄, and concentrated to give (1S,2S,5S)-2-(benzyloxy)-6-oxabicyclo[3.1.0]hexane (2.40 g, yield: 43.10%) as a colorless oil, which was confirmed by NOE. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.37-7.26 (m, 5H), 4.61-4.51 (m, 2H), 4.09 (d, 1H, *J* = 5.2 Hz), 3.55 (br.s, 1H), 3.49 (br.s, 1H), 1.99-1.95 (m, 1H), 1.87-1.75 (m, 2H), 1.54-1.52 (m, 1H).

Step 4: (1S,2R,5S)-2-Azido-5-(benzyloxy)cyclopentan-1-ol



To a mixture of (1S,2S,5S)-2-(benzyloxy)-6-oxabicyclo[3.1.0]hexane (2.40 g, 12.62 mmol) and NH₄Cl (1.55 g, 29.03 mmol) in H₂O (3.00 mL) and MeOH (24.00 mL) was added NaN₃ (4.10 g, 63.10 mmol), which was stirred at 80 °C for 16 hrs. After TLC (PE:EtOAc=10:1) showed the starting material was consumed, the organic solvent was dried by N₂ and the residue was diluted with H₂O (20 mL), extracted with DCM (20 mL*3). The combined organic phases were washed with H₂O (10 mL*3), dried over Na₂SO₄, and concentrated to give (1S,2R,5S)-2-azido-5-(benzyloxy)cyclopentan-1-ol (2.60 g, yield: 88.32%) as a brown oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.42-7.31 (m, 5H), 4.64-4.55 (m, 2H), 4.02-3.99 (m, 1H), 3.82-3.80 (m, 1H), 3.66-3.63 (m, 1H), 2.25 (br.s, 1H), 2.07-2.01 (m, 2H), 1.80-1.77 (m, 2H).


Step 5: (((1S,2R,5S)-2-Azido-5-(benzyloxy)cyclopentyl)oxy)(tert-butyl)dimethylsilane



A mixture of (1S,2R,5S)-2-azido-5-(benzyloxy)cyclopentan-1-ol (2.50 g, 10.72 mmol), imidazole (1.61 g, 23.69 mmol) and TBDMSCl (2.42 g, 16.08 mmol) in CHCl₃ (5.00 mL) was stirred at 80 °C for 16 hrs. Once TLC (PE:EtOAc=10:1) showed the starting material was consumed completely, the mixture was concentrated and purified by column chromatography on


silica gel (PE:EtOAc=1:0/100:1/80:1) to give (((1S,2R,5S)-2-azido-5-(benzyloxy)cyclopentyl)oxy)(tert-butyl)dimethylsilane (3.00 g, yield: 80.52%) as a colorless oil. ^1H -NMR (400 MHz, CDCl_3) δ ppm 7.24-7.15 (m, 5H), 4.40 (d, 2H, $J = 2.4$ Hz), 3.88-3.86 (m, 1H), 3.63-3.61 (m, 1H), 3.47-3.43 (m, 1H), 1.94-1.82 (m, 2H), 1.70-1.65 (m, 2H), 0.79 (s, 9H), 0.03 (s, 3H), 0.00 (s, 3H).

Step 6: tert-butyl ((1R,2S,3S)-2-((tert-butyldimethylsilyl)oxy)-3-Hydroxycyclopentyl)carbamate



To a mixture of (((1S,2R,5S)-2-azido-5-(benzyloxy)cyclopentyl)oxy)(tert-butyl)dimethylsilane (2.90 g, 8.34 mmol) and Boc_2O (2.20 g, 10.10 mmol) in MeOH (50.00 mL) was added $\text{Pd}(\text{OH})_2$ (1.50 g, 5.42 mmol), which was stirred at 50 °C under H_2 (50 psi) for 20 hrs. After TLC (PE:EtOAc = 3:1) showed the starting material was consumed completely, the mixture was filtered, and the filtrate was concentrated and purified by column chromatography on silica gel (PE:EtOAc=10:1/8:1/5:1) to give tert-butyl ((1R,2S,3S)-2-((tert-butyldimethylsilyl)oxy)-3-hydroxycyclopentyl)carbamate (2.10 g, yield: 75.95%) as a white solid. ^1H -NMR (400 MHz, CDCl_3) δ ppm 4.81 (br.s, 1H), 3.89-3.88 (m, 1H), 3.70-3.67 (m, 2H), 2.06-1.89 (m, 2H), 1.58-1.56 (m, 2H), 1.35 (s, 9H), 0.79 (s, 9H), 0.02 (s, 3H), 0.00 (s, 3H).


Step 7: tert-butyl ((1R,2S,3R)-2-((tert-butyldimethylsilyl)oxy)-3-Fluorocyclopentyl)carbamate



To a mixture of tert-butyl ((1R,2S,3S)-2-((tert-butyldimethylsilyl)oxy)-3-hydroxycyclopentyl)carbamate (1.10 g, 3.32 mmol) in DCM (50.00 mL) was added DAST (1.61 g, 9.96 mmol) at -70 °C. The reaction mixture was stirred at -70 °C for 1 hr and 25 °C for 1 hr. After TLC (PE:EtOAc=5:1, $R_f = 0.6$) showed the reaction was complete, ice water (5 mL) was added to the reaction. The solution was extracted with DCM (20 mL*3) and washed with brine (30 mL). The organic layer was dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography on silica gel (PE:EtOAc = 100:1) to give tert-butyl ((1R,2S,3R)-2-


((tert-butyldimethylsilyl)oxy)-3-fluorocyclopentyl)carbamate (100.00 mg, yield: 9.03%) as a white solid. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 4.64 (d, 1H, $J = 54.4$ Hz), 4.26 (br.s, 1H), 3.90-3.86 (m, 1H), 3.76-3.68 (m, 1H), 2.11-1.79 (m, 4H), 1.34 (s, 9H), 0.81 (s, 9H), 0.00 (s, 6H).

Step 8: (1S,2R,5R)-2-Amino-5-fluorocyclopentan-1-ol hydrochloride(relative stereochemistry)




A solution of tert-butyl ((1R,2S,3R)-2-((tert-butyldimethylsilyl)oxy)-3-fluorocyclopentyl)carbamate (100.00 mg, 299.84 μmol) in HCl/MeOH (20.00 mL, 4 M) was stirred at 25 °C for 16 hrs. TLC (PE:EtOAc=5:1, $R_f = 0$) showed the reaction was complete. The solution was dried by N_2 and (1S,2R,5R)-2-amino-5-fluorocyclopentan-1-ol hydrochloride (relative stereochemistry) (45.00 mg, yield: 96.45%) was obtained as a yellow solid. $^1\text{H-NMR}$ (400 MHz, CD_3OD) δ ppm 4.93-4.89 (m, 1H), 3.98-3.88 (m, 1H), 3.52-3.47 (m, 1H), 2.30-2.00 (m, 3H), 1.66-1.62 (m, 1H).

Example 9: Synthesis of (1S,2R,5S)-2-amino-5-fluorocyclopentan-1-ol (relative stereochemistry)




Step 1: (1R,2R,5S)-N,N-Dibenzyl-6-oxabicyclo[3.1.0]hexan-2-amine



Cl_3CCOOH (154.72 g, 949.20 mmol) was added to a stirring solution of N,N-dibenzylcyclopent-2-en-1-amine (50.00 g, 189.84 mmol) in DCM (640 mL), and the resulting mixture was stirred at 20 °C for 0.1 hr. *m*-CPBA (43.00 g, 199.33 mmol) was added in one portion


and the reaction mixture was allowed to continue to stir at 20 °C for 16 hrs. After TLC (PE:EtOAc=10:1) showed the reaction was complete, the mixture was diluted with DCM (500 mL) and *sat. aq.* Na₂SO₃ was added until starch-iodide paper indicated no remaining *m*-CPBA. *Sat. aq.* NaHCO₃ (500 mL) was added and the layers were separated. The organic layer was washed with *aq.* NaHCO₃ (200 mL*2) then dried, concentrated, and purified by column chromatography on silica gel (PE:EtOAc=100:1-50:1) to give (1R,2R,5S)-N,N-dibenzyl-6-oxabicyclo[3.1.0]hexan-2-amine (30.00 g, yield: 56.56%) as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.41 (d, 4H, *J* = 7.2 Hz), 7.31 (t, 4H, *J* = 7.6 Hz), 7.25-7.22 (m, 2H), 3.86-3.70 (m, 4H), 3.44 (s, 1H), 3.32 (s, 1H), 3.28-3.24 (m, 1H), 2.04-2.01 (m, 1H), 1.54-1.45 (m, 3H).

Step 2: (1R,2R,3R)-3-(Dibenzylamino)-2-hydroxycyclopentyl acetate



The solution of (1R,2R,5S)-N,N-dibenzyl-6-oxabicyclo[3.1.0]hexan-2-amine (30.00 g, 107.38 mmol, 1.00 Eq) in AcOH (200 mL) was stirred at 50 °C for 16 hrs. After TLC (PE:EtOAc = 10:1) showed the reaction was complete, the mixture was concentrated to remove AcOH, the residue was dissolved in DCM (100 mL), and the organic layer was washed with *aq.* NaHCO₃ (100 mL*3), and dried over Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography (PE:EtOAc = 100:1-50:1) to give (1R,2R,3R)-3-(dibenzylamino)-2-hydroxycyclopentyl acetate (20.00 g, yield: 54.87%) as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.36-7.28 (m, 10H), 5.05-5.02 (m, 1H), 4.05 (d, 1H, *J* = 4.0 Hz), 3.81-3.69 (m, 4H), 3.27-3.24 (m, 1H), 2.40-2.36 (m, 1H), 2.07 (s, 3H), 1.97-1.94 (m, 1H), 1.78-1.73 (m, 1H), 1.60-1.54 (m, 1H).


Step 3: (1R,2R,3R)-3-(Dibenzylamino)-2-((methylsulfonyl)oxy)cyclopentyl acetate



MsCl (8.10 g, 70.71 mmol) was added dropwise to a mixture of (1R,2R,3R)-3-(dibenzylamino)-2-hydroxycyclopentyl acetate (20.00 g, 58.92 mmol), Et₃N (18.48 g, 182.66 mmol) and DMAP (719.83 mg, 5.89 mmol) in DCM (200 mL). After addition, the mixture was

stirred at 20 °C for 16 hrs. After TLC (PE:EtOAc=10:1) showed the reaction was complete, the mixture was washed with water (100 mL*2), the organic layer was dried over Na₂SO₄, and purified by column chromatography on silica gel (PE:EtOAc = 80:1-60:1) to give (1R,2R,3R)-3-(dibenzylamino)-2-((methylsulfonyl)oxy)cyclopentyl acetate (15.00 g, yield: 60.97%) as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.41 (d, 4H, *J* = 7.6 Hz), 7.33 (t, 4H, *J* = 6.8 Hz), 7.31-7.25 (m, 2H), 5.16-5.13 (m, 1H), 5.01-5.00 (m, 1H), 3.92-3.81 (m, 4H), 3.37-3.32 (m, 1H), 3.15 (s, 3H), 2.33-2.30 (m, 1H), 2.01 (s, 3H), 1.98-1.91 (m, 2H), 1.56-1.53 (m, 1H).

Step 4: (1S,2R,5R)-N,N-Dibenzyl-6-oxabicyclo[3.1.0]hexan-2-amine



K₂CO₃ (5.96 g, 43.11 mmol) was added to a mixture of (1R,2R,3R)-3-(dibenzylamino)-2-((methylsulfonyl)oxy)cyclopentyl acetate (15.00 g, 35.93 mmol) in MeOH (70 mL)/THF (30 mL). The mixture was stirred at 20 °C for 16 hrs. After TLC (PE:EtOAc = 10:1) showed the reaction was complete, the mixture was concentrated to remove MeOH and THF. The mixture was then dissolved in DCM (20 mL), the organic layer was washed with water (10 mL*2), dried over Na₂SO₄, and concentrated to give (1S,2R,5R)-N,N-dibenzyl-6-oxabicyclo[3.1.0]hexan-2-amine (10.00 g, yield: 99.62%) as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.37 (d, 4H, *J* = 7.6 Hz), 7.32 (d, 4H, *J* = 7.2 Hz), 7.29-7.23 (m, 2H), 3.73-3.69 (m, 2H), 3.53-3.41 (m, 5H), 2.06-2.00 (m, 1H), 1.91-1.90 (m, 1H), 1.87-1.76 (m, 1H), 1.51-1.48 (m, 1H).


Step 5: (1S,2R,5S)-2-(Dibenzylamino)-5-fluorocyclopentan-1-ol



To a mixture of (1S,2R,5R)-N,N-dibenzyl-6-oxabicyclo[3.1.0]hexan-2-amine (6.50 g, 23.27 mmol) in DCM (200 mL) was added HBF₄/Et₂O (7.54 g, 46.53 mmol), and the mixture was allowed to stir at 30 °C for 0.2 hr. After TLC showed the reaction was complete, the mixture was added to Na₂CO₃ (100 mL) and extracted with DCM (200 mL*2). The organic layer was dried over Na₂SO₄, concentrated, and purified by column chromatography on silica gel (PE:EtOAc = 60:1 to 20:1) to give (1S,2R,5S)-2-(dibenzylamino)-5-fluorocyclopentan-1-ol (1.80


g, yield: 25.84%) as a white solid. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 7.38-7.22 (m, 10H), 4.81-4.67 (m, 1H), 4.23-4.15 (m, 1H), 3.83-3.55 (m, 4H), 3.04-2.98 (m, 1H), 1.93-1.79 (m, 4H).

Step 6: (1S,2R,5S)-2-Amino-5-fluorocyclopentan-1-ol(relative stereochemistry)




To a mixture of (1S,2R,5S)-2-(dibenzylamino)-5-fluorocyclopentan-1-ol (1.60 g, 5.34 mmol) in MeOH (10 mL) was added $\text{Pd}(\text{OH})_2$ (500.00 mg, 3.61 mmol), and the mixture was allowed to stir at 30 °C for 16 hrs under H_2 (30 psi). After TLC showed the reaction was complete, the mixture was filtered by celite and the filtrate was concentrated to give (1S,2R,5S)-2-amino-5-fluorocyclopentan-1-ol (600.00 mg, yield: 94.31%) as a white solid. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 4.93-4.77 (m, 1H), 3.92-3.84 (m, 1H), 3.10-3.04 (m, 1H), 2.07-1.98 (m, 3H), 1.64-1.61 (m, 1H).

Example 10: Synthesis of 3-Fluoro-5-((2R,4S)-4-fluoropyrrolidin-2-yl)pyridine




Step 1: (R)-4-((tert-butyldimethylsilyl)oxy)pyrrolidin-2-one




To the mixture of (R)-4-hydroxypyrrolidin-2-one (9.0 g, 89.1 mmol) in DMF (50 mL) was added imidazole (9.09 g, 134 mmol) and TBDMSCl (14.1 g, 93.6 mmol) in one portion at 0°C. The reaction mixture was stirred at 25 °C for 3 hrs. TLC (DCM/MeOH = 10/1, R_f = 0.8) showed the reaction was complete, then water (200 mL) was added the resulting precipitate was collected by filtration and dried in vacuo to give (R)-4-((tert-butyldimethylsilyl)oxy)pyrrolidin-2-one (15.5 g, yield: 80.7%) as a white solid. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 5.90 (br.s, 1H), 4.55-4.53 (m, 1H), 3.60-3.56 (m, 1H), 3.24-3.21 (m, 1H), 2.56-2.50 (m, 1H), 2.28-2.23 (m, 1H), 0.87-0.85 (m, 9H), 0.06-0.00 (m, 6H).

Step 2: Preparation of tert-butyl (R)-4-((tert-butyldimethylsilyl)oxy)-2-oxopyrrolidine-1-carboxylate



To the mixture of (R)-4-((tert-butyldimethylsilyl)oxy)pyrrolidin-2-one (15.5 g, 72.0 mmol) in CH_3CN (150 mL) was added Et_3N (8.72 g, 86.4 mmol), DMAP (4.39 g, 36 mmol), and Boc_2O (20.4 g, 93.7 mmol) in one portion at 0 °C. The reaction mixture was stirred at 25 °C for 10 hrs. TLC (PE/EtOAc = 3/1) showed the reaction was complete, then water (600 mL) was added, the resulting precipitate was collected by filtration and dried in vacuo to give tert-butyl (R)-4-((tert-butyldimethylsilyl)oxy)-2-oxopyrrolidine-1-carboxylate (19.2 g, yield: 84.6%) as a pink solid. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 4.33-4.30 (m, 1H), 3.81-3.77 (m, 1H), 3.56-3.54 (m, 1H), 2.67-2.61 (m, 1H), 2.41-2.37 (m, 1H), 1.46 (s, 9H), 0.80 (s, 9H), 0.00 (s, 6H).


Step 3: tert-butyl ((2R)-2-((tert-butyldimethylsilyl)oxy)-4-(5-fluoropyridin-3-yl)-4-hydroxybutyl)carbamate



To the mixture of 3-bromo-5-fluoro-pyridine (3.35 g, 19.02 mmol, 1.20 eq) in THF (40.00

mL) was added *i*-PrMgCl-LiCl (1.3 M, 17.56 mL, 1.44 *eq*) dropwise at 0 °C over 30 mins (exothermic). After addition, the temperature was raised to 25 °C over 1 hr and stirred at 25 °C for 30 mins. TLC (PE/EtOAc = 10/1) showed a new spot was generated indicating that the Mg reagent was prepared successfully. Tert-butyl (R)-4-((tert-butyldimethylsilyl)oxy)-2-oxopyrrolidine-1-carboxylate (5.00 g, 15.85 mmol, 1.00 *eq*) in THF (50 mL) was then added dropwise to the solution at -78 °C over 30 mins. The mixture was allowed to warm to 25 °C over 1 hr, then stirred at 25 °C for 16 hrs. TLC (PE/EtOAc = 3/1) showed the starting material was consumed completely and the desired product tert-butyl (R)-2-((tert-butyldimethylsilyl)oxy)-4-(5-fluoropyridin-3-yl)-4-oxobutylcarbamate was detected. The reaction was quenched by addition of MeOH (50 mL) at 0 °C. NaBH₄ (1.20 g, 31.70 mmol, 2.00 *eq*) was added at 0 °C, then the mixture was stirred at 25 °C for 4 hrs. TLC (PE/EtOAc = 2/1) and LCMS showed the reaction was complete. The combined reaction mixture (4 parallel reactions) was quenched by aqueous NH₄Cl (400 mL) and extracted with EtOAc (600 mL*3). The combined organics were dried over Na₂SO₄ and concentrated in vacuo, and the residue was purified by HPLC to give tert-butyl ((2R)-2-((tert-butyldimethylsilyl)oxy)-4-(5-fluoropyridin-3-yl)-4-hydroxybutyl)carbamate (1.24 g, yield: 18.91%) as a yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 8.26-8.22 (m, 2H), 7.37 (d, 1H, *J* = 8.8 Hz), 4.95-4.88 (m, 2H), 4.69 (br.s, 1H), 4.00-3.98 (m, 2H), 3.23-3.10 (m, 2H), 1.73 (br.s, 2H), 1.32 (s, 9H), 0.80-0.79 (m, 9H), 0.00 (s, 6H).


Step 4: tert-butyl (4R)-4-((tert-butyldimethylsilyl)oxy)-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate



To the mixture of tert-butyl ((2R)-2-((tert-butyldimethylsilyl)oxy)-4-(5-fluoropyridin-3-yl)-4-hydroxybutyl)carbamate (8.70 g, 20.98 mmol, 1.00 *eq*) and Et₃N (31.84 g, 314.70 mmol, 15.00 *eq*) in DCM (500.00 mL) was added dropwise MsCl (31.24 g, 272.74 mmol, 13.00 *eq*) at -60 °C over 0.5 hr. The mixture was then stirred at -60 °C for 1 hr, and the reaction mixture was allowed to warm to 25 °C and stirred for 18 hrs. LCMS showed the starting material was consumed completely. The mixture was then washed with H₂O (200 mL*3), and the aqueous phase was


extracted with DCM (200 mL*4). The combined organic layers were dried over Na₂SO₄ and concentrated in vacuo to give crude product tert-butyl (4R)-4-((tert-butyldimethylsilyl)oxy)-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (8.30 g, crude) as a black/ brown oil, which was used directly without purification.

Step 5: tert-butyl (4R)-2-(5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate



To the mixture of tert-butyl (4R)-4-((tert-butyldimethylsilyl)oxy)-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (8.30 g, 20.93 mmol, 1.00 *eq*) in THF (250.00 mL) was added TBAF (9.43 g, 41.86 mmol, 2.00 *eq*) at 25 °C. The mixture was stirred at 25 °C for 16 hrs. After TLC (PE/EtOAc = 1/1) showed the reaction was complete, the mixture was concentrated and the residue was dissolved in EtOAc (600 mL), washed with water (200 mL*5), dried over Na₂SO₄, and concentrated. The crude product was purified by PLC to give tert-butyl (4R)-2-(5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate (4.70 g, 16.65 mmol, yield: 79.54%) as a brown black oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 8.37-8.33 (m, 2H), 7.48 (br.s, 1H), 5.09-4.89 (m, 1H), 4.56-4.54 (m, 1H), 3.80-3.65 (m, 2H), 2.63-2.43 (m, 1H), 2.03-1.96 (m, 1H), 1.56-1.20 (m, 9H).


Step 6: tert-butyl (2R,4S)-4-fluoro-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate



To the mixture of tert-butyl (4R)-2-(5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate (4.70 g, 16.65 mmol, 1.00 *eq*) in DCM (150.00 mL) was added DAST dropwise (29.52 g, 183.15 mmol, 11.00 *eq*) at -78 °C over 0.5 hr. The reaction mixture was stirred at -78 °C for 2 hrs, then allowed to warm to 25 °C and stirred for 20 hrs. After TLC (PE/EtOAc = 0/1) showed the starting material was consumed completely, the mixture was cooled to 0 °C and quenched by saturated NaHCO₃ solution (100 mL) dropwise. The organic phase was separated and dried over Na₂SO₄, concentrated to give the residue, then purified by column chromatography on silica gel


(PE:EtOAc from 10:1, 8:1 to 5:1, then 3:1) to give tert-butyl (2R,4S)-4-fluoro-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (1.38 g, 4.85 mmol, yield: 29.15%, $R_f = 0.53$) as a white solid and tert-butyl (2S,4S)-4-fluoro-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (1.36 g, 4.78 mmol, yield: 28.73%, $R_f = 0.43$) as a yellow oil. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 8.31-8.27 (m, 2H), 7.20-7.18 (m, 1H), 5.18 (d, 1H, $J = 51.6$ Hz), 4.97-4.88 (m, 1H), 4.04-4.00 (m, 1H), 3.64 (dd, 1H, $J = 38.8, 12.8$ Hz), 2.67 (dd, 1H, $J = 15.6, 6.8$ Hz), 1.97-1.67 (m, 1H), 1.56-1.12 (m, 9H).

Step 7: 3-Fluoro-5-((2R,4S)-4-fluoropyrrolidin-2-yl)pyridine



To the mixture of tert-butyl (2R,4S)-4-fluoro-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (1.38 g, 4.85 mmol, 1.00 *eq*) in EtOAc (10 mL) was added dropwise HCl/EtOAc (40.00 mL, 4 M) at 0 °C. The mixture was allowed to warm to 25 °C and stirred 3 hrs. After TLC (PE:EtOAc = 1:1) showed the reaction was complete, the solvent was evaporated to give 3-fluoro-5-((2R,4S)-4-fluoropyrrolidin-2-yl)pyridine (1.25 g, 4.86 mmol, yield: 100.00%) as a brown solid. $^1\text{H-NMR}$ (400 MHz, CD_3OD) δ ppm 8.84-8.81 (m, 2H), 8.31 (d, 1H, $J = 9.2$ Hz), 5.62 (dt, 1H, $J = 52.0, 2.4$ Hz), 5.23-5.18 (m, 1H), 4.00-3.95 (m, 1H), 3.88-3.71 (m, 1H), 2.67 (td, 1H, $J = 16.0, 6.0$ Hz), 1.69-1.59 (m, 1H).

Example 11: Synthesis of (3R,4S,5R)-5-aminotetrahydro-2H-pyran-3,4-diol



Step 1: (S)-3,6-dihydro-2H-pyran-3-yl acetate

(3R,4S)-3,4-dihydro-2H-pyran-3,4-diyl diacetate (2.9 g, 14.49 mmol) was taken up in DCM

(15 ml), and stirred under N₂ at room temperature. Triethylsilane (2.55 ml, 15.93 mmol) was added and stirred for 5 minutes. BF₃.OEt₂ (1.836 ml, 14.49 mmol) was added dropwise and stirring was continued for 30 minutes. The reaction mixture was quenched with 30ml of saturated bicarbonate and the layers were separated. The combined organic layers were dried over sodium sulfate and the solvent was removed. The residue was purified via flash chromatography (0-30% Hex/EtOAc). (S)-3,6-dihydro-2H-pyran-3-yl acetate (1.9g, 92% yield) was recovered as clear oil. ¹H NMR (400 MHz, DMSO-d₆) δ 6.10 (dddt, J = 10.2, 3.2, 2.1, 1.0 Hz, 1H), 5.84 (ddt, J = 10.1, 4.3, 2.1 Hz, 1H), 4.96 (dtd, J = 4.3, 2.7, 1.5 Hz, 1H), 4.16 - 3.89 (m, 2H), 3.73 (t, J = 2.9 Hz, 2H), 2.01 (d, J = 0.9 Hz, 3H).

Step 2: (S)-3,6-dihydro-2H-pyran-3-ol

(S)-3,6-dihydro-2H-pyran-3-yl acetate (1.9 g, 13.37 mmol) was taken up in MeOH (30 ml) and Water (20 ml). Triethylamine (7 ml, 50.2 mmol) was added and stirred at room temperature for 30 min. The solvent was removed under reduced pressure. The residual water was then extracted with EtOAc three times. The organic layers were combined, dried over sodium sulfate and the solvent was removed. (S)-3,6-dihydro-2H-pyran-3-ol (1.1 g, 10.99 mmol, 82 % yield) was recovered as a clear oil. The crude product was carried on without further purification.

Step3: (1S,5R,6R)-3,7-dioxabicyclo[4.1.0]heptan-5-ol

(S)-3,6-dihydro-2H-pyran-3-ol (1.1 g, 10.99 mmol) was taken up in CH₂Cl₂ (20 ml) and cooled to 0 °C. mCPBA (4.55 g, 13.18 mmol) was added portion wise. The reaction mixture was stirred while warming to room temperature, overnight. The white precipitate of the reaction mixture was filtered off, the elutant was retained, the solvent was removed and triturated with diethyl ether. This step was repeated. The residue, (1S,5R,6R)-3,7-dioxabicyclo[4.1.0]heptan-5-ol (1.2g, 100% yield) was carried on without further purification.

Step 4: (3R,4S,5R)-5-(((R)-1-phenylethyl)amino)tetrahydro-2H-pyran-3,4-diol

(1S,5R,6R)-3,7-dioxabicyclo[4.1.0]heptan-5-ol (1.26 g, 10.85 mmol), (R)-1-phenylethanamine (1.658 ml, 13.02 mmol) were taken up in 2-BuOH (15ml). The reaction mixture was heated to 100 °C for 18 hours. The reaction mixture was cooled to room temperature, the solvent was removed, and the residue was then purified on ISCO 0-100% EtOAc. The fractions were combined, the solvent removed, and then the residue was treated with MTBE and stirred overnight. The white precipitate of the organic mixture was filtered off. Recovered (3R,4S,5R)-5-


((*(R)*-1-phenylethyl)amino)tetrahydro-2H-pyran-3,4-diol (0.350g, 14% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.38 – 7.24 (m, 4H), 7.22 – 7.12 (m, 1H), 4.61 (d, J = 5.6 Hz, 1H), 4.46 (d, J = 4.8 Hz, 1H), 3.88 (q, J = 6.5 Hz, 1H), 3.64 (tt, J = 5.0, 2.8 Hz, 1H), 3.47 (dd, J = 11.4, 4.7 Hz, 1H), 3.39 (ddd, J = 8.4, 5.6, 3.1 Hz, 1H), 3.31 (s, 2H), 3.29 (t, J = 3.0 Hz, 1H), 3.25 (d, J = 2.5 Hz, 0H), 2.79 (dd, J = 11.1, 7.8 Hz, 1H), 2.57 (td, J = 7.8, 3.9 Hz, 1H), 1.86 (s, 1H), 1.21 (d, J = 6.6 Hz, 3H).

Step 5: (3*R*,4*S*,5*R*)-5-aminotetrahydro-2H-pyran-3,4-diol

(3*R*,4*S*,5*R*)-5-(((*(R)*-1-phenylethyl)amino)tetrahydro-2H-pyran-3,4-diol (0.350 g, 1.475 mmol) was taken up in EtOH (3 ml) and Pd-C (0.031 g, 0.295 mmol) was added. The reaction mixture was stirred under H_2 balloon overnight. The reaction mixture was filtered through Celite and the solvent was removed to give (3*R*,4*S*,5*R*)-5-aminotetrahydro-2H-pyran-3,4-diol (0.190 g, 1.427 mmol, 97 % yield) as an off white solid. The crude product was carried on without further purification. LCMS ($\text{M}+\text{H}$) 134.

Example 12. (*(R)*-3-(4,4-difluoropyrrolidin-2-yl)-5-fluoropyridine


Step 1: tert-butyl (2*R*,4*R*)-4-((tert-butyldimethylsilyl)oxy)-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate



To a mixture of tert-butyl ((2*R*)-2-((tert-butyldimethylsilyl)oxy)-4-(5-fluoropyridin-3-yl)-4-hydroxybutyl)carbamate (6.80 g, 16.40 mmol) and Et_3N (24.89 g, 246.00 mmol) in DCM (500.00 mL) was added MsCl (24.42 g, 213.20 mmol) dropwise at -60°C over 30 minutes. The mixture was stirred at -60°C for 1 hr. The reaction mixture was allowed to warm to 25°C and stirred for an additional 18 hrs. The mixture was washed with H_2O (200 mL*3). The aqueous phase was extracted with DCM (200 mL*4). The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuo*. The residue was purified by silica gel chromatography (PE:EtOAc = 50/1, 20/1, 10/1) to afford tert-butyl (2*S*,4*R*)-4-((tert-butyldimethylsilyl)oxy)-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (2.70 g, yield: 41.52%) and tert-butyl (2*R*,4*R*)-4-((tert-butyldimethylsilyl)oxy)-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (2.40 g, yield: 36.89%) as brown oil. ^1H -NMR (400 MHz, CDCl_3) δ ppm 8.40 (br.s, 2H), 7.56-7.45 (m, 1H), 5.11-4.94 (m, 2H), 4.53 (br.s, 1H), 3.85-3.79 (m, 1H), 3.66-3.53 (m, 1H), 2.62-2.58 (m, 1H), 2.04-2.01 (m, 1H),


1.56 (s, 3H), 1.32 (s, 6H), 0.99-0.88 (m, 9H), 0.18-0.00 (m, 6H).

Step 2: tert-butyl (2R,4R)-2-(5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate




To a mixture of tert-butyl (2R,4R)-4-((tert-butyldimethylsilyl)oxy)-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (2.40 g, 6.05 mmol) in THF (60.00 mL) was added TBAF (3.16 g, 12.10 mmol) in one portion at 25°C. The mixture was concentrated under reduced pressure at 50°C. The residue was added to water (20 mL). The aqueous phase was extracted with ethyl acetate (30 mL*3). The combined organic phase was washed with saturated brine (20 mL*2), dried over Na₂SO₄, filtered and concentrated *in vacuo*. The residue was purified by silica gel chromatography (PE:EtOAc = 20/1, 10/1, 1/3) to afford tert-butyl (2R,4R)-2-(5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate (1.30 g, yield: 76.11%) as a yellow solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 8.26 (d, 2H, J = 12.8 Hz), 7.39 (br.s, 1H), 4.95-4.81 (m, 1H), 4.48-4.47 (m, 1H), 3.73 (br.s, 1H), 3.56-3.53 (m, 1H), 2.55 (br.s, 1H), 1.97-1.98 (m, 1H), 1.65-1.16 (m, 9H).


Step 3: tert-butyl (R)-2-(5-fluoropyridin-3-yl)-4-oxopyrrolidine-1-carboxylate



To a mixture of tert-butyl (2R,4R)-2-(5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate (1.30 g, 4.60 mmol) and trichloroisocyanuric acid (1.10 g, 4.60 mmol) was added TEMPO (72.41 mg, 460.49 μmol) at -10°C. The mixture was stirred at -10°C for 15 min, then warmed to 25 °C and stirred for 1 hr. TLC (EtOAc) showed the reaction was complete. The organic phase was washed with NaHCO₃ (20 mL*2), dried over Na₂SO₄, filtered and concentrated in vacuum. The residue was purified by silica gel chromatography (petroleum ether/ethyl acetate = 50/1, 10/1) to afford tert-butyl (R)-2-(5-fluoropyridin-3-yl)-4-oxopyrrolidine-1-carboxylate (1.10 g, yield: 85.32%) as a brown oil.


Step 4: tert-butyl (R)-4,4-difluoro-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate

To a mixture of tert-butyl (R)-2-(5-fluoropyridin-3-yl)-4-oxopyrrolidine-1-carboxylate (1.00 g, 3.57 mmol) in DCM (100.00 mL) was added DAST (14.39 g, 89.25 mmol) dropwise at -70°C under N_2 . The mixture was stirred at -70°C for 30 min. Then the mixture was stirred at 25°C for 16 hrs. The reaction mixture was quenched by saturated aq. NaHCO_3 slowly at 0°C and the aqueous phase was extracted with DCM (50 mL*4). The combined organic phase was washed with saturated brine (30 mL), dried over Na_2SO_4 , filtered and concentrated *in vacuo*. The residue was purified by silica gel chromatography (petroleum ether/ethyl acetate = 100/1, 30/1) to afford tert-butyl (R)-4,4-difluoro-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (1.00 g, yield: 92.66%) as a brown oil. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 8.40 (s, 1H), 8.34 (s, 1H), 7.30-7.21 (m, 1H), 5.06 (br.s, 1H), 4.14-3.85 (m, 2H), 2.91-2.84 (m, 1H), 2.39-2.32 (m, 1H), 1.43-1.14 (m, 9H).

Step 5: (R)-3-(4,4-difluoropyrrolidin-2-yl)-5-fluoropyridine


A mixture of tert-butyl (R)-4,4-difluoro-2-(5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (1.00 g, 3.31 mmol) in HCl/EtOAc (50.00 mL, 4 M) was stirred for 2 hrs at 25°C . The mixture was concentrated under reduced pressure at 30°C to afford (R)-3-(4,4-difluoropyrrolidin-2-yl)-5-fluoropyridine (840.00 mg, yield: 92.25%) as a white solid as bis HCl salt. $^1\text{H-NMR}$ (400 MHz, MeOD) δ ppm 8.68-8.63 (m, 1H), 7.97 (d, 1H, $J = 9.2$ Hz), 5.26-5.21 (m, 1H), 4.03-3.90 (m, 2H), 3.13-2.92 (m, 2H).

Example 13. (3S,5R)-5-(2,5-difluorophenyl)pyrrolidine-3-carbonitrileStep 1: tert-butyl ((2R)-2-((tert-butyldimethylsilyl)oxy)-4-(2,5-difluorophenyl)-4-hydroxybutyl)carbamate



To a solution of 2-bromo-1,4-difluoro-benzene (3.01 g, 15.60 mmol, 1.20 Eq) in THF (15 mL) was added isopropylmagnesium chloride complex (2.27 g, 15.60 mmol, 1.20 Eq) at 0°C dropwise under N₂. The reaction was stirred at 15°C for 1 hr to prepare (2, 5-difluorophenyl) magnesium bromide (23 mL). To a solution of tert-butyl (R)-4-((tert-butyldimethylsilyl)oxy)-2-oxopyrrolidine-1-carboxylate (4.10 g, 13.00 mmol, 1.00 Eq) in THF (50 mL) was added (2,5-difluorophenyl) magnesium bromide (23 mL) dropwise at 0°C over 30 mins. The reaction mixture was stirred at 0°C for 1 hr. Methanol (20 mL) was added to the mixture followed by NaBH₄ (738 mg, 19.50 mmol, 1.50 Eq) at 0°C. The mixture was stirred at 0°C for 1 hr then poured into 10% aqueous NH₄Cl. The mixture was extracted with EtOAc (20 mL*2), the combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude product was purified by medium pressure liquid chromatography (MPLC) to give tert-butyl ((2R)-2-((tert-butyldimethylsilyl)oxy)-4-(2,5-difluorophenyl)-4-hydroxybutyl)carbamate (2.22 g, 5.14 mmol, 39.6% yield). ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.17-7.15 (m, 1H), 6.86-6.79 (m, 2H), 5.11-5.06 (m, 1H), 4.70 (br.s, 1H), 4.02-3.98 (m, 1H), 3.69 (br.s, 0.5H), 3.46 (br.s, 0.5H), 3.33-3.14 (m, 2H), 1.80-1.69 (m, 2H), 1.35 (s, 9H), 0.84-0.82 (9H, m), 0.04-0.03 (6H, m).


Step 2: tert-butyl (4R)-4-((tert-butyldimethylsilyl)oxy)-2-(2,5-difluorophenyl)pyrrolidine-1-carboxylate



To a solution of tert-butyl ((2R)-2-((tert-butyldimethylsilyl)oxy)-4-(2,5-difluorophenyl)-4-hydroxybutyl)carbamate (13.40 g, 31.05 mmol, 1.00 Eq) and Et₃N (9.43 g, 93.14 mmol, 3.00 Eq) in DCM (50 mL) was added dropwise methanesulfonyl chloride (5.33 g, 46.57 mmol, 1.50 Eq) at -60°C by under N₂. The mixture was stirred at -60°C for 2 hrs and 15°C for 16 hrs. LCMS showed the starting material was consumed completely. The reaction mixture was extracted with DCM (30


mL*2) and the combined organics were washed with brine (50 mL), dried over Na_2SO_4 and filtered, concentrated to give tert-butyl (4R)-4-((tert-butyldimethylsilyl)oxy)-2-(2,5-difluorophenyl)pyrrolidine-1-carboxylate (12.00 g, 26.11 mmol, yield: 84.10%, 90% purity) which was used directly without further purification.

Step 3: tert-butyl (2R,4R)-2-(2,5-difluorophenyl)-4-hydroxypyrrolidine-1-carboxylate



To a solution of tert-butyl (4R)-4-((tert-butyldimethylsilyl)oxy)-2-(2,5-difluorophenyl)pyrrolidine-1-carboxylate (4.50 g, 10.88 mmol, 1.00 Eq) in THF (30 mL) was added TBAF/THF (1 M, 14.15 mL, 1.30 Eq) at 15°C. The mixture was stirred at 15°C for 16 hrs. TLC (PE:EtOAc = 3:1) showed the starting material was consumed completely. The reaction mixture was quenched by H_2O (50 mL), extracted with EtOAc (30 mL*2) and the combined organics were washed with brine (10 mL), dried over Na_2SO_4 , filtered and concentrated *in vacuo*. The residue was purified by neutral *prep*-HPLC to afford tert-butyl (2R,4R)-2-(2,5-difluorophenyl)-4-hydroxypyrrolidine-1-carboxylate (1.00 g, 3.34 mmol, yield: 30.70%) as a white solid. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 7.04-6.80 (m, 3H), 5.10-5.00 (m, 1H), 4.43 (s, 1H), 3.75 (br.s, 1H), 3.53-3.49 (m, 1H), 2.53 (br.s, 1H), 1.93-1.90 (m, 1H), 1.40-1.16 (m, 9H).


Step 4: tert-butyl (2R,4R)-2-(2,5-difluorophenyl)-4-((methylsulfonyl)oxy)pyrrolidine-1-carboxylate



To a mixture of tert-butyl (2R,4R)-2-(2,5-difluorophenyl)-4-hydroxypyrrolidine-1-carboxylate (3.00 g, 10.02 mmol, 1.00 eq) and Et_3N (2.03 g, 20.04 mmol, 2.00 eq) in DCM (80.00 mL) was added MsCl (1.61 g, 14.03 mmol, 1.40 eq) dropwise at 0°C. The mixture was stirred at 18°C for 2 hrs. The mixture was quenched by H_2O (30 mL). The aqueous phase was extracted by DCM (50 mL*3). The combined organic layer was dried over Na_2SO_4 and concentrated under reduced pressure. tert-Butyl (2R,4R)-2-(2,5-difluorophenyl)-4-((methylsulfonyl)oxy)pyrrolidine-1-


carboxylate (3.60 g, 9.54 mmol, yield: 95.20%) was obtained as a brown solid.

Step 5: tert-butyl (2R,4S)-4-cyano-2-(2,5-difluorophenyl)pyrrolidine-1-carboxylate




To a mixture of tert-Butyl (2R,4R)-2-(2,5-difluorophenyl)-4-((methylsulfonyl)oxy)pyrrolidine-1-carboxylate (3.60 g, 9.54 mmol, 1.00 eq) in DMSO (20.00 mL) was added KCN (745.49 mg, 11.45 mmol, 1.20 eq) in one portion. The mixture was stirred at 90°C for 3 hrs. 80 mL of H₂O was added to the mixture, and the mixture was extracted by EtOAc (80 mL*4). The combined organic layer was concentrated under reduced pressure. The residue was purified by silica gel chromatography (PE/EtOAc = 40:1, 30:1, 10:1). tert-Butyl (2R,4S)-4-cyano-2-(2,5-difluorophenyl)pyrrolidine-1-carboxylate (1.60 g, 5.19 mmol, yield: 54.40%) was obtained as light green liquid.

Step 6: (3S,5R)-5-(2,5-difluorophenyl)pyrrolidine-3-carbonitrile



A mixture of tert-Butyl (2R,4S)-4-cyano-2-(2,5-difluorophenyl)pyrrolidine-1-carboxylate (800.00 mg, 2.59 mmol, 1.00 eq) in TFA (4.00 mL)/DCM (20.00 mL) was stirred at 18°C for 3 hrs. The mixture was dried under N₂. (3S,5R)-5-(2,5-difluorophenyl)pyrrolidine-3-carbonitrile (780.00 mg, 2.42 mmol, yield: 93.44%) was obtained as a light yellow solid.


Example 14. 3-fluoro-5-((2R,4S)-4-fluoropyrrolidin-2-yl)benzamide



3-Fluoro-5-((2R,4S)-4-fluoropyrrolidin-2-yl)benzonitrile (0.050 g, 0.240 mmol) (Prepared as in WO 2012/034095) was taken up in TFA (0.800 ml, 10.38 mmol) and H_2SO_4 (0.200 ml, 3.75 mmol) and stirred overnight at room temperature. The reaction mixture was diluted with ice water (3ml) and the solid was isolated by filtration, and used directly.


Example 15. 2-chloro-5-fluoro-3-((2R,4S)-4-fluoropyrrolidin-2-yl)pyridine

Step 1: (S,Z)-N-((2-chloro-5-fluoropyridin-3-yl)methylene)-2-methylpropane-2-sulfinamide




2-chloro-5-fluoronicotinaldehyde (20 g, 125 mmol) was taken up in THF (150 ml) at 0 °C. (R)-2-Methylpropane-2-sulfinamide (16.71 g, 138 mmol) was added followed by dropwise addition of titaniumtetraethanolate (22.88 ml, 150 mmol). The reaction mixture was stirred while warming to RT. After 3 hours the reaction mixture was cooled to 0 °C, and 150ml of brine was added and stirred for 20 minutes. The mixture was filtered through Celite. The aqueous layer was separated and discarded. The organic layer with dried over Na_2SO_4 and the solvent was removed to give (S,Z)-N-((2-chloro-5-fluoropyridin-3-yl)methylene)-2-methylpropane-2-sulfinamide (32 g, 122 mmol, 97 % yield), which was carried on without further purification. LCMS: 263 M+H.

Step 2: (R)-N-((R)-1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-yl)-2-methylpropane-2-sulfinamide




(R,E)-N-((2-chloro-5-fluoropyridin-3-yl)methylene)-2-methylpropane-2-sulfinamide (32.9 g, 125 mmol) was dissolved in HMPA (100 ml) and cooled to 0 °C. Zinc (16.37 g, 250 mmol), allyl bromide (21.67 ml, 250 mmol) and water (2.256 ml, 125 mmol) were added at 0 °C and the reaction mixture was allowed to warm to RT overnight. LCMS showed complete conversion to desired product. 100 ml of water was added at RT and stirred for 30 minutes. 30 ml of MBTE was added followed by 60 ml of 10% citric acid and the reaction mixture was stirred for 30 minutes. The mixture was filtered through Celite and washed with MTBE. The organic layer was washed with 10% citric acid, water and brine. The solvent was removed under vacuum to give (R)-N-((R)-1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-yl)-2-methylpropane-2-sulfinamide (14.5 g, 47.6 mmol, 38.0 % yield) as an orange oil. LCMS: 305 M+H.

Step 3: (R)-1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-amine, HCl



(R)-N-((R)-1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-yl)-2-methylpropane-2-sulfinamide (7.5 g, 24.61 mmol) was taken up in 10 ml MeOH. HCl (4M in dioxane) (30.8 ml, 123 mmol) was added and stirred at RT for 1h. The solvent was removed under vacuum and the residue was diluted in DCM and washed with saturated aqueous NaHCO₃. The layers were separated and the organic layer was dried with Na₂SO₄ and the solvent was removed under vacuum. Recovered (R)-1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-amine, HCl (5.83 g, 24.59 mmol, 100 % yield) as a solid. LCMS: 201 M+H.


Step 4: (R)-N-(1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-yl)acetamide



To (R)-1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-amine•HCl (5.83 g, 24.59 mmol) in DCM (70.3 ml) at 0 °C was added TEA (4.11 ml, 29.5 mmol) and acetic anhydride (2.320 ml, 24.59 mmol). The mixture was stirred for 2 hours. The reaction mixture was poured into saturated


aqueous NaHCO_3 and extracted with DCM. The organic layer was washed with brine, dried over MgSO_4 , and evaporated under reduced pressure. Recovered (R)-N-(1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-yl)acetamide (5.97 g, 24.60 mmol, 100 % yield) and was carried on without further purification. LCMS: 243 M+H.

Step 5: (5R)-5-(2-chloro-5-fluoropyridin-3-yl)pyrrolidin-3-yl acetate




(R)-N-(1-(2-chloro-5-fluoropyridin-3-yl)but-3-en-1-yl)acetamide (5.97 g, 24.60 mmol) was taken up in THF (56.2 ml) and water (14.06 ml), followed by addition of I_2 (18.73 g, 73.8 mmol) and stirred overnight at RT. The crude reaction was diluted with saturated NaHCO_3 and $\text{Na}_2\text{S}_2\text{O}_3$ solutions and extracted twice with EtOAc. Aqueous layer was basified with saturated aqueous NaHCO_3 and extracted with EtOAc to obtain (5R)-5-(2-chloro-5-fluoropyridin-3-yl)pyrrolidin-3-yl acetate (5.9 g, 22.81 mmol, 93 % yield) as a light yellow oil. LCMS: 259 M+H.

Step 6: (2R)-tert-butyl 4-acetoxy-2-(2-chloro-5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate




To a solution of (5R)-5-(2-chloro-5-fluoropyridin-3-yl)pyrrolidin-3-yl acetate (5.9 g, 22.81 mmol) in dioxane (76 ml) and water (76 ml) was added BOC-anhydride (7.94 ml, 34.2 mmol) followed by careful addition of 2N NaOH (7ml) to achieve pH ~9. The reaction mixture was stirred for 1 hour at RT. The reaction mixture was diluted with water and extracted with EtOAc three times. The organic layer was dried over Na_2SO_4 and the solvent was removed under vacuum to give (2R)-tert-butyl 4-acetoxy-2-(2-chloro-5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (3.5 g, 9.75 mmol, 42.8 % yield), which was carried on without further purification. LCMS: 359 M+H.

Step 7: (2R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate




(2R)-tert-butyl 4-acetoxy-2-(2-chloro-5-fluoropyridin-3-yl)pyrrolidine-1-carboxylate (3.5 g, 9.75 mmol) was taken up in MeOH (48.8 ml) followed by addition of 2M NaOH (5.37 ml, 10.73 mmol) and the reaction mixture was stirred at RT for 2 hours. The solvent was removed under vacuum and the aqueous layer was neutralized with 1N HCl, and extracted with EtOAc three times. The combined organic layers were dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified via silica gel chromatography (0-70% Hex/EtOAc) to give (2R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate (2.1 g, 6.63 mmol, 68.0 % yield). LCMS: 317 M+H.

Step 8: (R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-oxopyrrolidine-1-carboxylate




(2R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate (2.1 g, 6.63 mmol) was taken up in DCM (66.3 ml) and NaHCO₃ (0.557 g, 6.63 mmol) was added followed by Dess-Martin periodinane (8.44 g, 19.89 mmol). The reaction mixture was stirred overnight. Water was added (0.119 ml, 6.63 mmol) followed by Dess-Martin periodinane (8.44 g, 19.89 mmol) and stirred for 18 hours. The pH was adjusted to ~7 with saturated aqueous NaHCO₃ and extracted with DCM x3. The organic layers were combined, dried over Na₂SO₄ and the solvent was removed under vacuum. The residue was purified via flash chromatography (0-70% Hex/EtOAc) to give (R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-oxopyrrolidine-1-carboxylate (1.6 g, 5.08 mmol, 77 % yield). LCMS: 315 M+H.

Step 9: (2R,4R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate




(R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-oxopyrrolidine-1-carboxylate (1.6 g, 5.08 mmol) was suspended in ethanol (33.9 ml) and cooled to 0°C. NaBH₄ was added portionwise (0.096 g, 2.54 mmol) and stirred for 45 minutes at 0°C. The reaction was quenched slowly with saturated NH₄Cl and allowed to warm to RT, and the solution was extracted with DCM x3. The organic layers were combined and dried over Na₂SO₄. The residue was purified via flash chromatography (0-70% Hex/EtOAc) to give (2R,4R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate (1.446 g, 4.57 mmol, 90 % yield). LCMS: 317 M+H.

Step 10: (2R,4S)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-fluoropyrrolidine-1-carboxylate




(2R,4R)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-hydroxypyrrolidine-1-carboxylate (1.0 g, 3.16 mmol) was taken up in DCM (25 ml) and cooled to -78°C. TEA-HF (1.098 ml, 9.47 mmol) was added and stirred for 10 minutes. XtalFluor-E (1.446 g, 6.31 mmol) was added and after 10 minutes the reaction mixture was transferred to an ice bath and allowed to warm to 0°C. After 2 hours the reaction mixture was diluted with DCM and quenched with saturated aqueous NaHCO₃. The organic layers were separated, and the solvent was removed under vacuum. The residue was purified via ISCO (0-50% Hex/EtOAc; 12g column) to give (2R,4S)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-fluoropyrrolidine-1-carboxylate (0.805 g, 2.53 mmol, 80 % yield) as a white solid. LCMS: 319 M+H.

Step 11: 2-chloro-5-fluoro-3-((2R,4S)-4-fluoropyrrolidin-2-yl)pyridine, HCl



(2R,4S)-tert-butyl 2-(2-chloro-5-fluoropyridin-3-yl)-4-fluoropyrrolidine-1-carboxylate (0.805 g, 2.53 mmol, 80 % yield) was taken up in EtOAc (5ml) and 4N HCl/dioxane (3ml) was added. The reaction mixture was stirred at RT for 1 hour. The precipitate was filtered off, washed with ether, and dried under high vacuum overnight to give 2-chloro-5-fluoro-3-((2R,4S)-4-fluoropyrrolidin-2-yl)pyridine, HCl (0.612 g, 2.399 mmol, 76 % yield) as an off white solid. LCMS: 219 M+H.


Example 16. 5-fluoro-3-((2R,4S)-4-fluoropyrrolidin-2-yl)-2-methoxypyridine



5-Fluoro-3-((2R,4S)-4-fluoropyrrolidin-2-yl)-2-methoxypyridine was prepared in the same way as 3-fluoro-5-((2R,4S)-4-fluoropyrrolidin-2-yl)benzamide, substituting for 5-fluoro-2-methoxynicotinaldehyde for 2-chloro-5-fluoronicotinaldehyde.


Example 17. Methyl (1R,3R,4R)-3-amino-4-hydroxycyclopentane-1-carboxylate

Step 1: (1S,2R,4S,5R)-3-oxa-6-azatricyclo[3.2.1.0^{2,4}]octan-7-one




To a solution of (1R,4S)-2-azabicyclo[2.2.1]hept-5-en-3-one (30.00 g, 274.90 mmol, 1.00 eq) in NaH₂PO₄ (395.00 mL, 0.2M) and Na₂HPO₄ (55.00 mL, 0.2 M) was added H₂O (450.00 mL) and oxone (669.31 g, 4.40 mol, 16.00 eq) at 0°C portion-wise over 5 hrs, and maintaining the pH = 6 by addition of aq. NaOH (12 M) and keeping the temperature at 0°C. After addition, the mixture was stirred at 0°C for further 2 hrs, TLC (PE:EtOAc = 1:1) showed the starting material was consumed completely, the mixture was filtered and aqueous phase was extracted with DCM (400 mL*5), the combined organic layers were dried over Na₂SO₄, concentrated in vacuum to get (1S,2R,4S,5R)-3-oxa-6-azatricyclo[3.2.1.0^{2,4}]octan-7-one (9.00 g, 71.93 mmol, yield: 26.16%) as a yellow solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 5.96 (br.s, 1H), 3.86 (s, 1H), 3.62 (1H, d, J = 3.2 Hz), 3.53 (1H, d, J = 2.8 Hz), 2.86 (s, 1H), 1.82 (d, 1H, J = 9.6 Hz), 1.64 (d, 1H, J = 10.0 Hz).

Step 2: tert-butyl (1S,2R,4S,5R)-7-oxo-3-oxa-6-azatricyclo[3.2.1.0^{2,4}]octane-6-carboxylate




To a solution of (1S,2R,4S,5R)-3-oxa-6-azatricyclo[3.2.1.0.2,4]octan-7-one (9.00 g, 71.93 mmol, 1.00 *eq*) in DCM (100.00 mL) was added Boc₂O (17.27 g, 79.12 mmol, 1.10 *eq*), Et₃N (8.73 g, 86.32 mmol, 1.20 *eq*) and DMAP (878.71 mg, 7.19 mmol, 0.10 *eq*), the mixture was stirred at 25°C for 16 hrs, LCMS showed the starting material was consumed completely, the mixture was washed with NH₄Cl *aq.* (100 mL*3), the combined organic layers were dried over Na₂SO₄, concentrated in vacuum, the crude product was purified with column chromatography on silica gel (PE:EtOAc = 5:1~1:1) to get tert-butyl (1S,2R,4S,5R)-7-oxo-3-oxa-6-azatricyclo[3.2.1.0.2,4]octane-6-carboxylate (12.00 g, 53.28 mmol, yield: 74.07%) as a yellow solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 4.56 (s, 1H), 3.71 (d, 1H, *J* = 2.8 Hz), 3.54 (d, 1H, *J* = 2.8 Hz), 3.00 (s, 1H), 1.75 (d, 1H, *J* = 10.0 Hz), 1.57 (d, 1H, *J* = 10.8 Hz), 1.46 (s, 9H).

Step 3: methyl (3R,4R)-4-((tert-butoxycarbonyl)amino)-3-hydroxycyclopent-1-ene-1-carboxylate




Na (3.37 mg, 146.50 μmol, 0.01 *eq*) was added to MeOH (10.00 mL) at 0°C, then the solution was stirred at 0°C for 0.5 hr, the solution was added to tert-butyl (1S,2R,4S,5R)-7-oxo-3-oxa-6-azatricyclo[3.2.1.0.2,4]octane-6-carboxylate (3.30 g, 14.65 mmol, 1.00 *eq*) in MeOH (30.00 mL), and then mixture was stirred at 16°C for 13.5 hrs, LCMS showed the starting material was consumed completely, the reaction was quenched with acetic acid (5 mL), and then washed with NaHCO₃ (20 mL*3), the organic layer was dried over Na₂SO₄, and concentrated in vacuum, the crude product was washed with PE (20 mL) to get methyl (3R,4R)-4-((tert-butoxycarbonyl)amino)-3-hydroxycyclopent-1-ene-1-carboxylate (2.10 g, 8.16 mmol, yield: 55.72%) as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 6.65 (s, 1H), 4.98 (s, 1H), 4.81 (d, 1H, *J* = 2.8 Hz), 4.46 (s, 1H), 3.98-3.92 (m, 1H), 3.75 (s, 3H), 3.07-3.01 (m, 1H), 2.36-2.29 (m, 1H), 1.45 (s, 9H).

Step 4: methyl (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentane-1-carboxylate



A mixture of methyl (3R,4R)-4-((tert-butoxycarbonyl)amino)-3-hydroxycyclopent-1-ene-1-carboxylate (2.00 g, 7.77 mmol, 1.00 *eq*), (1Z,5Z)-cycloocta-1,5-diene;(2S,5S)-1-[2-[(2S,5S)-2,5-dimethylphospholan-1-yl]ethyl]-2,5-dimethyl-phospholane;rhodium(1+);trifluoromethanesulfonate (48.08 mg, 77.74 μ mol, 0.01 *eq*) in MeOH (50.00 mL) was degassed and purged with H₂ for 3 times, and then the mixture was stirred at 55°C for 16 hrs under H₂ (40 psi) atmosphere, TLC (PE:EtOAc = 1:1) showed the starting material was consumed, the mixture was concentrated in vacuum, and then dissolved in EtOAc (5 mL), then to the mixture was added PE (20 mL), white solid was formed, the precipitate was collected, dried in vacuum, the solid was dissolved in MeOH (7 mL), and purified with acidic *prep*-HPLC (HCl) to get methyl (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentane-1-carboxylate (750.00 mg, 2.89 mmol, yield: 37.23%) as a yellow oil, the structure was confirmed by chiral HPLC and ¹H-NMR. ¹H-NMR (400 MHz, CDCl₃) δ ppm 4.04-3.99 (m, 1H), 3.80-3.77 (m, 1H), 3.70 (s, 3H), 2.93-2.91 (m, 1H), 2.48-2.44 (m, 1H), 2.43-2.35 (m, 1H), 1.92-1.89 (m, 1H), 1.87-1.69 (m, 1H), 1.45 (s, 9H).

Example 18. Synthesis of methyl (1R,3R,4R)-3-amino-4-hydroxycyclopentane-1-carboxylate




Methyl (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentane-1-carboxylate (700.00 mg, 2.70 mmol, 1.00 *eq*) in HCl/dioxane (10.00 mL, 4 M) was stirred at 19°C for 5 hrs, LCMS showed the starting material was consumed, the mixture was concentrated in vacuum to get methyl (1R,3R,4R)-3-amino-4-hydroxycyclopentane-1-carboxylate (500.00 mg, 2.56 mmol, yield: 94.66%) as yellow oil. ¹H-NMR (400 MHz, CD₃OD) δ ppm 4.09-4.05 (m, 1H), 3.75-3.66 (m, 3H), 3.37-3.33 (m, 1H), 3.06-3.04 (m, 1H), 2.45-2.37 (m, 2H), 1.87-1.81 (m, 2H). LC-MS (mobile phase: from 95% [water + 0.375% v/v TFA] and 5% [CH₃CN + 0.188% v/v TFA], under this condition for 0.25 min, then changed to 15% [CH₃CN + 0.188% v/v TFA] in 10.0 min, under this condition for 5 min, finally changed to 95% [water + 0.375% v/v TFA] and 5% [CH₃CN

+ 0.188% v/v TFA] in 0.01 min, then under this condition for 5 min. The flow is 1.0 mL·min⁻¹ all along) purity is 98.803%, Rt = 0.893 min, MS Calcd.: 159.2, MS Found: 160.1 ([M+1]⁺).


Example 19. (3aS,4R,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-amine

Step 1: (3aS,4S,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol



(3aR,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-one (50.00 g, 324.34 mmol, 1.00 *eq*) was taken up to MeOH (1.00 L), then CeCl₃·7H₂O (120.84 g, 324.34 mmol, 30.83 mL, 1.00 *eq*) was added. The mixture was cooled to 0°C. Then NaBH₄ (24.54 g, 648.68 mmol, 2.00 *eq*) was added portion-wise at 0°C among 1.5 hrs. After addition, the reaction was complete checked by TLC (PE/EtOAc = 5/1). The reaction was quenched by saturated NH₄Cl (1000 mL), extracted with DCM (500 mL*5). The combined organic layers were dried over Na₂SO₄ and concentrated in vacuo at 45°C. (3aS,4S,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol (50.66 g, 324.37 mmol, yield: 100.00%) was obtained as a yellow liquid, which was used directly without purification.


Step 2: 2-((3aS,4R,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione



To the mixture of (3aS,4S,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-ol (8.00 g, 51.22 mmol, 1.00 *eq*) and isoindoline-1,3-dione (9.04 g, 61.46 mmol, 1.20 *eq*) in toluene (250.00 mL) was added PPh₃ (20.15 g, 76.83 mmol, 1.50 *eq*) at 20°C. Then DIAD (15.54 g, 76.83 mmol, 1.50 *eq*) was added dropwise to the mixture at 0°C. After addition, the mixture was allowed to 80°C and stirred for 16 hrs. TLC (PE/EtOAc = 5/1) showed the reaction was complete. The mixture was concentrated. The residue was purified by column chromatography on silica gel (PE/EtOAc = 25/1 to 15/1). The obtained product was crude as yellow oil with some polar spots on TLC. So 80 mL of MeOH was added and the white precipitate was generated and collected by filtration. 2-((3aS,4R,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-

yl)isoindoline-1,3-dione (9.60 g, 33.65 mmol, yield: 65.70%) was obtained as a white solid.


Step 3: (3aS,4R,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-amine



To the mixture of 2-((3aS,4R,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (9.52 g, 33.37 mmol, 1.00 *eq*) in EtOH (300.00 mL) was added ethane-1,2-diamine (4.01 g, 66.74 mmol, 2.00 *eq*). The resulting mixture was stirred at 80°C for 16 hrs. Lots of white precipitate was generated. TLC (PE/EtOAc = 5/1) showed the starting material was consumed completely. The precipitate was filtered. To the filtrate was added 300 mL of NaOH (0.5 M). The mixture was extracted with DCM (200 mL*5), dried over Na₂SO₄ and concentrated. (3aS,4R,6aR)-2,2-dimethyl-3a,6a-dihydro-4H-cyclopenta[d][1,3]dioxol-4-amine (4.90 g, 31.57 mmol, yield: 94.62%) was obtained as a yellow oil.


Example 20. (1R,3R,4R)-3-amino-4-hydroxycyclopentane-1-carbonitrile

Step 1: (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentane-1-carboxylic acid




A mixture of methyl (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentane-1-carboxylate (5.00 g, 19.28 mmol, 1.00 *eq*), LiOH.H₂O (2.43 g, 57.85 mmol, 3.00 *eq*) in MeOH (10.00 mL) and H₂O (10.00 mL) was stirred at 15°C for 16 hrs, TLC (PE:EtOAc = 1:1) showed the reaction was complete, to the mixture was added diluted HCl (1 M) until pH = 6, and concentrated in vacuum, then the mixture was the dissolved in DCM (15 mL) and EtOAc (5 mL), the mixture was filtered, and the filtrate was concentrated in vacuum to get the product (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentane-1-carboxylic acid (6.50 g, crude) as a white solid. ¹H-NMR (400 MHz, CD₃OD) δ ppm 3.95-3.90 (m, 1H), 3.74-3.69 (m, 1H), 2.91-2.87 (m, 1H), 2.32-2.22 (m, 2H), 1.82-1.72 (m, 2H), 1.45 (s, 9H).

Step 2: tert-butyl ((1R,2R,4R)-4-carbamoyl-2-hydroxycyclopentyl)carbamate



To a mixture of (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentane-1-carboxylic acid (3.00 g, 12.23 mmol, 1.00 *eq*) in DMF (40.00 mL) was added HATU (6.05 g, 15.90 mmol, 1.30 *eq*), DIPEA (4.74 g, 36.69 mmol, 3.00 *eq*) and NH_4Cl (1.96 g, 36.69 mmol, 3.00 *eq*), and then the mixture was stirred at 15°C for 32 hrs, LCMS showed the reaction was complete, the mixture was concentrated in vacuum to get tert-butyl ((1R,2R,4R)-4-carbamoyl-2-hydroxycyclopentyl)carbamate (11 g, crude) (2 batches were set up and purified together). ^1H -NMR (400 MHz, CD_3OD) δ ppm 4.58 (br.s, 1H), 3.96-3.94 (m, 1H), 3.77-3.75 (m, 1H), 2.25-2.20 (m, 2H), 1.79-1.75 (m, 2H), 1.44 (s, 9H).

Step 3: tert-butyl ((1R,2R,4R)-4-cyano-2-hydroxycyclopentyl)carbamate



To a mixture of tert-butyl ((1R,2R,4R)-4-carbamoyl-2-hydroxycyclopentyl)carbamate (2.00 g, 8.19 mmol, 1.00 *eq*), pyridine (1.94 g, 24.56 mmol, 3.00 *eq*) in THF (3.00 mL) was added TFAA (2.58 g, 12.28 mmol, 1.50 *eq*) dropwise at 0°C, then the mixture was stirred at 0°C for 0.5 hr, then to the mixture was added Et_3N (2.49 g, 24.56 mmol, 3.00 *eq*) at 15°C, and the mixture was stirred at 15°C for 0.5 hr, and to the mixture was added TFAA (2.58 g, 12.28 mmol, 1.50 *eq*), and the mixture was stirred at 15°C for 0.5 hr, LCMS showed the reaction was complete, the mixture was concentrated in vacuum, purified by *prep*-HPLC (TFA, MS) to get tert-butyl ((1R,2R,4R)-4-cyano-2-hydroxycyclopentyl)carbamate (380.00 mg, 1.68 mmol, yield: 20.51%) as a colorless oil. ^1H -NMR (400 MHz, CD_3OD) δ ppm 4.07-3.97 (m, 1H), 3.79-3.78 (m, 0.5H), 3.51-3.49 (m, 0.5H), 3.19-3.05 (m, 1H), 2.44-2.35 (m, 0.5H), 2.33-2.28 (m, 1.5H), 1.97-1.94 (m, 1H), 1.92-1.82 (m, 1H), 1.41 (s, 9H).


Step 4: (1R,3R,4R)-3-amino-4-hydroxycyclopentane-1-carbonitrile



A mixture of tert-butyl ((1R,2R,4R)-4-cyano-2-hydroxycyclopentyl)carbamate (800.00 mg, 3.54 mmol, 1.00 *eq*) in TFA (5.00 mL) and DCM (5.00 mL) was stirred at 20°C for 2 hrs, LCMS showed the reaction was complete, the mixture was concentrated in vacuum to get (1R,3R,4R)-3-amino-4-hydroxycyclopentane-1-carbonitrile (545.00 mg, 2.27 mmol, yield: 64.10%) as a colorless oil. ¹H-NMR (400 MHz, CD₃OD) δ ppm 4.12-4.06 (m, 1H), 3.51-3.47 (m, 1H), 3.23-3.20 (m, 1H), 2.48-2.42 (m, 2H), 2.11-2.10 (m, 1H), 1.94-1.90 (m, 1H).


Example 21. (3aS,4R,6aS)-6,6-difluoro-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-amine

Step 1: 2-((3aS,4R,6S,6aR)-6-hydroxy-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione




A mixture of (3aR,4S,6R,6aS)-6-amino-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-ol (0.43 g, 2.46 mmol, 1.00 *eq*), Phthalic anhydride (0.36 g, 2.46 mmol, 1 *eq*) and DIEA (0.65 mL, 3.7 mmol, 1.5 *eq*) in Toluene (6.2 mL) was stirred at 100 °C for 9 hrs. LCMS showed the reaction was complete. EtOAc was added to the reaction mixture and then washed with aqueous saturated sodium bicarbonate solution (15 mL). The combined organic layers were washed with saturated brine solution, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica gel (Hexanes/EtOAc) to get the product 2-((3aS,4R,6S,6aR)-6-hydroxy-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (0.62 g, 83%) as a white solid.

Step 2: 2-((3aS,4R,6aS)-2,2-dimethyl-6-oxotetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione



To a solution of 2-((3aS,4R,6S,6aR)-6-hydroxy-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (0.20 g, 0.68 mmol, 1.00 *eq*) in DCM (4.5 mL) was added PCC (0.29 g, 1.35 mmol, 2 *eq*) and the solution was stirred at 23 °C for 16 hrs. Another aliquot of PCC (0.15g, 0.67 mmol) was added and the reaction continued for another 16 hours. LCMS showed the reaction was complete. EtOAc was added to the reaction mixture and then filtered through a celite pad. The residue was concentrated and then purified by column chromatography on silica gel (Hexanes/EtOAc) to get the product 2-((3aS,4R,6aS)-2,2-dimethyl-6-oxotetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (0.19 g, 94%) as an off-white solid.


Step 3: 2-((3aS,4R,6aS)-6,6-difluoro-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione



To a solution of 2-((3aS,4R,6aS)-2,2-dimethyl-6-oxotetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (0.16 g, 0.53 mmol, 1.00 *eq*) in DCM (3.5 mL) was added DAST (0.42 g, 2.64 mmol, 5 *eq*) and the solution was stirred at reflux for 16 hrs. Another aliquot of DAST (0.42 g, 2.64 mmol, 5 *eq*) was added and the reaction continued for another 16 hours at 23 °C. The reaction mixture was diluted with DCM and then washed with aqueous saturated sodium bicarbonate solution. The combined organic layers were washed with saturated brine solution, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica gel (Hexanes/EtOAc) to get the product 2-((3aS,4R,6aS)-6,6-difluoro-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (0.065 g,

38%).


Step 4: (3aS,4R,6aS)-6,6-difluoro-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-amine



To a solution of 2-((3aS,4R,6aS)-6,6-difluoro-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-yl)isoindoline-1,3-dione (0.065 g, 0.2 mmol, 1.00 *eq*) in Ethanol (1.8 mL) was added Hydrazine monohydrate (0.015 mL, 0.3 mmol, 1.5 *eq*) and the solution was stirred at 50 °C for 2 hrs and then at 70 °C for another 2 hours. The heterogeneous reaction mixture was filtered using minimum volume of Ethanol. The filtrate was then concentrated and the isolated crude product (3aS,4R,6aS)-6,6-difluoro-2,2-dimethyltetrahydro-4H-cyclopenta[d][1,3]dioxol-4-amine was used without further purification in the next step.


Example 22. Synthesis of (1R,3R,4R)-4-aminocyclohexane-1,3-diol

Step 1: (1r,4r)-4-(benzyloxy)cyclohexanol and (1s,4s)-4-(benzyloxy)cyclohexanol




To an ice-bath cooled solution of 4-(benzyloxy)cyclohexanone (31.0 g, 152 mmol) in 500 mL methanol, sodium borohydride (5.78 g, 153 mmol) was added in several portions during a period of 10 min, then the solution was stirred at 20 °C for 2 h. Then the mixture was quenched by saturated aqueous solution of ammonium chloride (50 mL), concentrated and the residue was dissolved in 200 mL water and extracted with ethyl acetate (200 mL x 3), the combined organic phase was dried over sodium sulfate, then concentrated under vacuo to give title product (1r,4r)-4-(benzyloxy)cyclohexanol and (1s,4s)-4-(benzyloxy)cyclohexanol as a pale yellow oil (31.0 g, crude) which was used to next step directly without further purification, . MS (ES+) $C_{13}H_{18}O_2$ requires: 206, found: 207[M+H]⁺.

Step 2: ((Cyclohex-3-enyloxy)methyl)benzene




To an ice-bath cooled solution of (1R,4R)-4-(benzyloxy)cyclohexanol and (1S,4S)-4-(benzyloxy)cyclohexanol (30.0 g, 145 mmol) and N,N-Diisopropylethylamine (28.1 g, 218 mmol) in 1200 mL dichloromethane, trifluoromethanesulfonic anhydride (30.7 g, 109 mmol) was added dropwise during a period of 30 min, then the solution was stirred at 25 °C for 18 h. Then the mixture was concentrated under vacuo and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate = 12:1 to give the title compound (28.0 g, yield 100%) as a yellow oil. MS (ES+) $\text{C}_{13}\text{H}_{16}\text{O}$ requires: 188, found: 189 $[\text{M}+\text{H}]^+$.

Step 3: (1R,3R,6S)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane




A solution of ((cyclohex-3-enyloxy)methyl)benzene (12.0 g, 63.7 mmol) in dichloromethane (200 mL) was treated at 0 °C with *meta*-chloroperoxybenzoic acid (21.9 g, 127 mmol). The reaction mixture was stirred 2 h at 0 °C and then 15 min at room temperature. Evaporation of the washed (10% aqueous solution of sodium sulfite, 5% aqueous sodium hydroxide solution and then water) organic solution afforded a liquid residue, which was separated with silica gel column chromatography, eluting with hexane:isopropyl ether:ethyl acetate = 65:28:7 to give the title compound (4.18 g, yield 32%) as a yellow oil. The *trans*-(1S,3R,6R)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane showed a little less polarity on TLC and eluted firstly. The *cis*-(1R,3R,6S)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane eluted secondly. MS (ES+) $\text{C}_{13}\text{H}_{16}\text{O}_2$ requires: 204, found: 205 $[\text{M}+\text{H}]^+$.

Step 4: (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol



Lithium perchlorate (7.27 g, 68.4 mmol) was added to an ice-bath cooled stirred solution of (1R,3R,6S)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane (7.0 g, 34.2 mmol) in 120 mL 4A-MS dried acetonitrile, the bath was removed and (S)-1-phenylethanamine (5.58 g, 46.1 mmol) was added dropwise during a period of 15 min, then the solution was stirred at 25 °C for 18 h. Then the mixture was diluted in 200 mL water and extracted with ethyl acetate (200 mL x 3), the combined organic phase was dried over sodium sulfate, then concentrated and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate:triethylamine = 98:0:2 ~ 49:49:2 to give the title compound (3.5 g, yield 31%) as a yellow oil. The (1S,2S,5S)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol showed a little less polarity on TLC, and eluted firstly. The (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol eluted secondly. MS (ES+) $\text{C}_{21}\text{H}_{27}\text{NO}_2$ requires: 325, found: 326[M + H]⁺.


Step 5: (1R,2R,4R)-4-(benzyloxy)-2-(tert-butyldimethylsilyloxy)-N-((S)-1-phenylethyl)cyclohexanamine



tert-Butyldimethylsilyl trifluoromethanesulfonate (13.0 g, 49.5 mmol) was added to an ice-bath cooled, stirred solution of (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol (5.4 g, 16.5 mmol) and triethylamine (5.0 g, 49.5 mmol) in 100 mL dried dichloromethane. After 30


min, this was washed with a saturated aqueous solution of sodium bicarbonate, dried over sodium sulfate. Removal of the solvent, and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate = 100:0 ~ 70:30 to give the title compound (5.4 g, yield 71%) as a yellow oil. MS (ES+) $C_{27}H_{41}NO_2Si$ requires: 439, found: 440 $[M+H]^+$.

Step 7: (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol



Tetrabutylammonium fluoride (2.66 g, 10.2 mmol) was added to a stirred solution of (1R,2R,4R)-4-(benzyloxy)-2-(tert-butyldimethylsilyloxy)-N-((S)-1-phenylethyl)cyclohexanamine (1.5 g, 3.41 mmol) in 50 mL dried oxolane at room temperature. Then this solution was stirred at 65 °C for 2 h. Then the mixture was concentrated under vacuo and the residue was diluted in 200 mL water and extracted with ethyl acetate (200 mL x 3), the combined organic phase was washed with water and saturated aqueous solution of sodium chloride, dried over sodium sulfate. Removal of the solvent, and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate = 100:0 ~ 70:30 to give the title compound (0.75 g, yield 68%) as a colorless oil. MS (ES+) $C_{21}H_{27}NO_2$ requires: 325, found: 326 $[M + H]^+$.

Step 8: (1R,3R,4R)-4-aminocyclohexane-1,3-diol




10% Palladium hydroxide in activated carbon (697 mg, catalyst) was added to a solution of (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol (650 mg, 1.99 mmol) in 15 mL ethanol at room temperature. Then this solution was stirred at 50 °C for 20 h under hydrogen. Then the mixture was cooled and filtered through celite, the filter-cake was washed with

methanol:dichloromethane=1:10, the filtrate was concentrated under vacuo and the residue was diluted in 20 mL methanol:dichloromethane=1:10 solution and concentrated, dried under high-vacuo, then cooled at -20 °C to give the title compound (240 mg, yield 92%) as a white crystal. MS (ES+) $C_6H_{13}NO_2$ requires: 131, found: 132[M+H]⁺. ¹H-NMR (400 MHz, 6d-DMSO) δ ppm 4.62-4.49 (m, 2H), 3.42-3.33 (m, 2H, $J = 3.2$ Hz), 2.93-2.86 (m, 1H), 2.25-2.18 (m, 1H), 1.98-1.92 (m, 1H), 1.72-1.59 (m, 3H), 1.13-1.03 (m, 2H), 0.97-0.90 (m, 1H).


Example 23. Synthesis of tert-butyl (1R,2R)-2-(tert-butyldimethylsilyloxy)-4-oxocyclohexylcarbamate

Step 1: tert-butyl (1R,2R,4R)-2-(tert-butyldimethylsilyloxy)-4-hydroxycyclohexylcarbamate



10% Palladium hydroxide in activated carbon (1.9 g, catalyst) was added to a solution of (1R,2R,4R)-4-(benzyloxy)-2-(tert-butyldimethylsilyloxy)-N-((S)-1-phenylethyl)cyclohexanamine from the previous example (2.0 g, 4.54 mmol) and di-tert-butyl dicarbonate (3.95 g, 18.1 mmol) in 60 mL ethanol at room temperature. Then this solution was stirred at 50 °C for 20 h under hydrogen. Then the mixture was cooled and filtered through celite, the filter-cake was washed with methanol:dichloromethane=1:10, the filtrate was concentrated under vacuo and the residue was diluted in 20 mL methanol:dichloromethane = 1:10 solution and concentrated, dried under high-vacuo to give the title compound (1.2 g, yield 77%) as a colorless oil. MS (ES+) $C_{17}H_{35}NO_4Si$ requires: 345, found: 346 [M+H]⁺.

Step 2: tert-butyl (1R,2R)-2-(tert-butyldimethylsilyloxy)-4-oxocyclohexylcarbamate




1,1,1-Triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one (Dess-Martin periodinane, 4.11 g, 9.71 mmol) was added to a solution of tert-butyl (1R,2R,4R)-2-(tert-butyldimethylsilyloxy)-4-hydroxycyclohexylcarbamate (1.4 g, 4.05 mmol) in 50 mL dichloromethane at room temperature.

Then this solution was stirred at 40 °C for 3 h under nitrogen. Then the mixture concentrated under vacuo and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate = 100:0 ~ 95:5 to give the title compound (1.2 g, yield 86%) as a yellow oil. MS (ES+) $C_{17}H_{33}NO_4Si$ requires: 343, found: 344[M+H]⁺. ¹H-NMR (400 MHz, CDCl₃) δ ppm 4.70-4.49 (br. , 1H), 4.10-3.90 (br. s, 1H), 3.79-3.65 (br. , 1H), 2.64 (dd, 1H, *J* = 14.4, 4.0 Hz), 2.42-2.28 (m, 4H), 1.46 (s, 9H), 0.87 (s, 9H), 0.08 (d, 6H, *J* = 6.8 Hz).


Example 23. Synthesis of tert-butyl (1R,2R)-2-(tert-butyldimethylsilyloxy)-4-oxocyclohexylcarbamate and (1R,3R,4R)-4-aminocyclohexane-1,3-diol

Step 1: (1R,4R)-4-(benzyloxy)cyclohexanol and (1S,4S)-4-(benzyloxy)cyclohexanol:



To an ice-bath cooled solution of 4-(benzyloxy)cyclohexanone (31.0 g, 152 mmol) in 500 mL methanol, sodium borohydride (5.78 g, 153 mmol) was added in several portions during a period of 10 min, then the solution was stirred at 20 °C for 2 h. Then the mixture was quenched by saturated aqueous solution of ammonium chloride (50 mL), concentrated and the residue was dissolved in 200 mL water and extracted with ethyl acetate (200 mL x 3), the combined organic phase was dried over sodium sulfate, then concentrated under vacuo to give title product (1R,4R)-4-(benzyloxy)cyclohexanol and (1S,4S)-4-(benzyloxy)cyclohexanol as a pale yellow oil (31.0 g, crude) which was used to next step directly without further purification. MS (ES+) $C_{13}H_{18}O_2$ requires: 206, found: 207[M+H]⁺.


Step 2: ((Cyclohex-3-enyloxy)methyl)benzene:



To an ice-bath cooled solution of (1R,4R)-4-(benzyloxy)cyclohexanol and (1S,4S)-4-(benzyloxy)cyclohexanol (30.0 g, 145 mmol) and N,N-Diisopropylethylamine (28.1 g, 218 mmol)

in 1200 mL dichloromethane, trifluoromethanesulfonic anhydride (30.7 g, 109 mmol) was added dropwise during a period of 30 min, then the solution was stirred at 25 °C for 18 h. Then the mixture was concentrated under vacuo and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate = 12:1 to give the title compound (28.0 g, yield 100%) as a yellow oil. MS (ES+) $C_{13}H_{16}O$ requires: 188, found: 189 $[M+H]^+$.

Step 3: (1R,3R,6S)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane:




A solution of ((cyclohex-3-enyloxy)methyl)benzene (12.0 g, 63.7 mmol) in dichloromethane (200 mL) was treated at 0 °C with *meta*-chloroperoxybenzoic acid (21.9 g, 127 mmol). The reaction mixture was stirred 2 h at 0 °C and then 15 min at room temperature. Evaporation of the washed (10% aqueous solution of sodium sulfite, 5% aqueous sodium hydroxide solution and then water) organic solution afforded a liquid residue, which was separated with silica gel column chromatography, eluting with hexane:isopropyl ether:ethyl acetate = 65:28:7 to give the title compound (4.18 g, yield 32%) as a yellow oil. The *trans*-(1S,3R,6R)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane showed a little less polarity on TLC and eluted firstly. The *cis*-(1R,3R,6S)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane eluted secondly. MS (ES+) $C_{13}H_{16}O_2$ requires: 204, found: 205 $[M+H]^+$.

***cis*-(1R,3R,6S)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane:** 1H -NMR (400 MHz, $CDCl_3$) δ ppm 7.35-7.27 (m, 5H), 4.56-4.45 (m, 2H), 3.35-3.29 (m, 1H), 3.12-3.09 (m, 2H), 2.37-2.32 (m, 1H), 2.25-2.20 (m, 1H), 1.89-1.68 (m, 3H), 1.49-1.44 (m, 1H).

***Trans*-(1S,3R,6R)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane:** 1H -NMR (400 MHz, $CDCl_3$) δ ppm 7.35-7.27 (m, 5H), 4.48 (dd, 2H, $J = 28.0, 12.4$ Hz), 3.56-3.52 (m, 1H), 3.19-3.17 (m, 2H), 2.24-2.18 (m, 1H), 2.15-2.07 (m, 1H), 2.00-1.91 (m, 2H), 1.64-1.53 (m, 2H).

Step 3: (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol:




Lithium perchlorate (7.27 g, 68.4 mmol) was added to an ice-bath cooled stirred solution of (1R,3R,6S)-3-(benzyloxy)-7-oxa-bicyclo[4.1.0]heptane (7.0 g, 34.2 mmol) in 120 mL 4A-MS dried acetonitrile, the bath was removed and (S)-1-phenylethanamine (5.58 g, 46.1 mmol) was added dropwise during a period of 15 min, then the solution was stirred at 25 °C for 18 h. Then the mixture was diluted in 200 mL water and extracted with ethyl acetate (200 mL x 3), the combined organic phase was dried over sodium sulfate, then concentrated and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate:triethylamine = 98:0:2 ~ 49:49:2 to give the title compound (3.5 g, yield 31%) as a yellow oil. The (1S,2S,5S)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol showed a little less polarity on TLC, and eluted firstly. The (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol eluted secondly. MS (ES+) $C_{21}H_{27}NO_2$ requires: 325, found: 326[M + H]⁺.

(1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol: ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.35-7.24 (m, 10H), 4.52 (d, 2H, *J* = 2.0 Hz), 3.97 (q, 1H, *J* = 6.8 Hz), 3.42-3.34 (m, 1H), 3.19-3.12 (m, 1H), 2.39 (dd, 1H, *J* = 12.0, 2.4 Hz), 2.16 (dd, 1H, *J* = 12.0, 3.6 Hz), 2.09-2.00 (m, 2H), 1.65-1.49 (m, 1H), 1.35 (d, 3H, *J* = 6.4 Hz), 1.28-1.15 (m, 2H), 0.90 (qd, 1H, *J* = 13.2, 3.6 Hz).

(1S,2S,5S)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol: ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.36-7.22 (m, 10H), 4.54 (d, 2H, *J* = 3.2 Hz), 3.90 (q, 1H, *J* = 6.4 Hz), 3.44-3.35 (m, 1H), 3.15-3.09 (m, 1H), 2.51-2.45 (m, 1H), 2.43-2.36 (m, 1H), 2.04-1.99 (m, 1H), 1.95-1.90 (m, 1H), 1.47-1.29 (m, 3H), 1.34 (d, 3H, *J* = 6.4 Hz), 0.82 (qd, 1H, *J* = 13.2, 3.2 Hz).


Step 4: Synthesis of (1R,2R,4R)-4-(benzyloxy)-2-(tert-butyldimethylsilyloxy)-N-((S)-1-

phenylethyl)cyclohexanamine:




tert-Butyldimethylsilyl trifluoromethanesulfonate (13.0 g, 49.5 mmol) was added to an ice-bath cooled, stirred solution of (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol (5.4 g, 16.5 mmol) and triethylamine (5.0 g, 49.5 mmol) in 100 mL dried dichloromethane. After 30 min, this was washed with a saturated aqueous solution of sodium bicarbonate, dried over sodium sulfate. Removal of the solvent, and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate = 100:0 ~ 70:30 to give the title compound (5.4 g, yield 71%) as a yellow oil. MS (ES+) $C_{27}H_{41}NO_2Si$ requires: 439, found: 440 $[M+H]^+$.

Step 5: (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol:



Tetrabutylammonium fluoride (2.66 g, 10.2 mmol) was added to a stirred solution of (1R,2R,4R)-4-(benzyloxy)-2-(*tert*-butyldimethylsilyloxy)-*N*-((S)-1-phenylethyl)cyclohexanamine (1.5 g, 3.41 mmol) in 50 mL dried oxolane at room temperature. Then this solution was stirred at 65 °C for 2 h. Then the mixture was concentrated under vacuo and the residue was diluted in 200 mL water and extracted with ethyl acetate (200 mL x 3), the combined organic phase was washed with water and saturated aqueous solution of sodium chloride, dried over sodium sulfate. Removal of the solvent, and the residue was purified with silica gel column chromatography, eluting with petroleum ether:ethyl acetate = 100:0 ~ 70:30 to give the title compound (0.75 g, yield 68%) as a colorless oil. MS (ES+) $C_{21}H_{27}NO_2$ requires: 325, found: 326 $[M+H]^+$.


Step 6: (1R,3R,4R)-4-aminocyclohexane-1,3-diol:



10% Palladium hydroxide in activated carbon (697 mg, catalyst) was added to a solution of (1R,2R,5R)-5-(benzyloxy)-2-((S)-1-phenylethylamino)cyclohexanol (650 mg, 1.99 mmol) in 15 mL ethanol at room temperature. Then this solution was stirred at 50 °C for 20 h under hydrogen. Then the mixture was cooled and filtered through celite, the filter-cake was washed with methanol:dichloromethane=1:10, the filtrate was concentrated under vacuo and the residue was diluted in 20 mL methanol:dichloromethane=1:10 solution and concentrated, dried under high-vacuo, then cooled at -20 °C to give the title compound (240 mg, yield 92%) as a white crystal. MS (ES+) $C_6H_{13}NO_2$ requires: 131, found: 132[M+H]⁺.


(1R,3R,4R)-4-aminocyclohexane-1,3-diol: ¹H-NMR (400 MHz, 6d-DMSO) δ ppm 4.62-4.49 (m, 2H), 3.42-3.33 (m, 2H, $J = 3.2$ Hz), 2.93-2.86 (m, 1H), 2.25-2.18 (m, 1H), 1.98-1.92 (m, 1H), 1.72-1.59 (m, 3H), 1.13-1.03 (m, 2H), 0.97-0.90 (m, 1H).

Example 24: Synthesis of Compound 232:




Step 1: (4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-

2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone



To a solution of 4-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-4,5-dihydro-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (1.00 g, 3.22 mmol, 1.00 *eq*) in SOCl₂ (164.00 g, 1.38 mol, 428.11 *eq*) was added DMF (235.49 mg, 3.22 mmol, 1.00 *eq*) at 15°C. The reaction was heated at 50°C for 16 hrs. TLC (PE:EtOAc = 3:1, R_f = 0.8 and 0.7) showed the reaction was complete. The mixture was concentrated. The residue was cooled to -10°C and dissolved in DCM (25.00 mL). Et₃N (1.63 g, 16.10 mmol, 5.00 *eq*) and (2R,4S)-2-(2,5-difluorophenyl)-4-fluoro-pyrrolidine (497.40 mg, 2.09 mmol, 0.65 *eq*, HCl) was added to the reaction. The reaction was stirred at 0°C for 0.2 hr. TLC (PE:EtOAc = 3:1, R_f = 0.38) showed the reaction was complete. The solution was washed with brine (5 mL), dried over Na₂SO₄ and concentrated. The residue was purified by *prep*-TLC (PE:EtOAc = 10:1) to give (4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone (400.00 mg, yield: 24.26%) as a yellow solid.


Step 2: methyl (1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-carbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxycyclopentane-1-carboxylate



To a solution of (4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone (200.00 mg, 390.63 μmol, 1.00 *eq*) and methyl (1R,3R,4R)-3-amino-4-hydroxycyclopentanecarboxylate (80.24


mg, 410.16 μmol , 1.05 *eq*, HCl) in dioxane (10.00 mL) was added DIPEA (151.46 mg, 1.17 mmol, 3.00 *eq*). The reaction was heated at 90°C for 5 hrs. LCMS showed the reaction was complete. The solution was concentrated. The residue was purified by *prep*-TLC (PE:EtOAc = 2:1) to give methyl (1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxycyclopentane-1-carboxylate (120.00 mg, yield: 48.40%) as a yellow solid.

Step 3: methyl (1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxycyclopentane-1-carboxylate



To a solution of methyl (1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxycyclopentane-1-carboxylate (120.00 mg, 189.06 μmol , 1.00 *eq*) in DCM (3.00 mL) was added TFA (3.00 mL) at 15°C. The reaction was stirred at 15°C for 16 hrs. LCMS showed the starting material was consumed. Little of (1R,3R,4R)-methyl 3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1-(hydroxymethyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxycyclopentanecarboxylate was detected. The reaction was concentrated. The residue was dissolved in MeOH (20.00 mL). KOAc (185.54 mg, 1.89 mmol, 10.00 *eq*) was added to the reaction. The reaction was heated at 50°C for 16 hrs. LCMS showed the reaction was complete. The solution was concentrated. The residue was dissolved in EtOAc (20 mL) and washed with brine (10 mL), dried over Na_2SO_4 and concentrated to give methyl (1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxycyclopentane-1-carboxylate (80.00 mg, crude) as a red solid which was used in the next step without purification.


Step 4: ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((1R,2R,4R)-2-hydroxy-4-(2-hydroxypropan-2-yl)cyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone




To a solution of methyl ((1R,3R,4R)-3-(((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)-4-hydroxycyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)carboxylate (80.00 mg, 158.59 μmol , 1.00 *eq*) in THF (20.00 mL) was added MeMgBr (3 M, 1.59 mL, 30.00 *eq*) at -70°C . The reaction was slowly warmed to 15°C and stirred for 2 hrs. TLC (EtOAc, $R_f = 0.24$) and LCMS showed the reaction was complete. The solution was neutralized with 1N *aq.* HCl to pH = 7. The reaction mixture was concentrated. The residue was purified by neutral *prep*-HPLC. to give ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((1R,2R,4R)-2-hydroxy-4-(2-hydroxypropan-2-yl)cyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone (17.40 mg, yield: 21.75%) as a yellow solid.


For this compound and compound 156, below, LC-MS conditions were as follows: (mobile phase: from 99% [water + 0.375% *v/v* TFA] and 1% [CH_3CN + 0.188% *v/v* TFA], under this condition for 0.4 min, then changed to 10% [water + 0.375% *v/v* TFA] and 90% [CH_3CN + 0.188% *v/v* TFA] in 3.0 min, then changed to 100% [CH_3CN + 0.188% *v/v* TFA] in 0.45 min, finally changed to 99% [water + 0.375% *v/v* TFA] and 1% [CH_3CN + 0.188% *v/v* TFA] in 0.01 min, then under this condition for 0.64 min. The flow is $0.8\text{ mL}\cdot\text{min}^{-1}$ all along.) Purity is 99.870%

Example 25. Synthesis of Compound 229:






Step 1: tert-butyl (3R,4R)-3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxypyrrolidine-1-carboxylate




A solution of 3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one (0.085 g, 0.17 mmol, 1 eq) was stirred with thionyl chloride (0.031 mL, 0.43 mmol, 2.5 eq) and few drops of DMF in DCM (0.7 mL) at 50 °C for 3 hours. LCMS indicated complete consumption of SM to the chloro-heterocycle intermediate. The reaction mixture was cooled on ice and added Dioxane (0.7 mL) followed by DIEA (0.21 mL, 1.21 mmol, 7 eq) and tert-butyl (3R,4R)-3-amino-4-hydroxypyrrolidine-1-carboxylate (0.05g, 0.26 mmol, 1.5 eq). The reaction mixture was then stirred at 70 °C for 3 hours. LCMS indicated reaction was complete. The reaction mixture was then diluted with DCM and washed with aqueous saturated sodium bicarbonate solution. The combined organic layers were washed with saturated brine solution, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica gel (Hexanes/EtOAc) to get the product tert-butyl (3R,4R)-3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxypyrrolidine-1-carboxylate (0.084g, 72%).

Step 2: ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((3R,4R)-4-hydroxypyrrolidin-3-yl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone



A solution of (3R,4R)-3-((3-((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidine-1-carbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-4-hydroxypyrrolidine-1-carboxylate (0.084g, 0.12 mmol, 1 eq) in EtOAc (1.25 mL) was treated with HCl in Dioxane (4M, 0.9 mL, 3.72 mmol, 30 eq). After stirring at 23 °C for 4 hours, LCMS indicated reaction was complete. The reaction mixture was diluted with EtOAc and washed with aqueous saturated sodium bicarbonate solution. The combined organic layers were washed with saturated brine solution, dried over Na₂SO₄ and concentrated in vacuo. The crude product ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((3R,4R)-4-hydroxypyrrolidin-3-yl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone was used without further purification in the next step.


Step 3: ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((3R,4R)-4-hydroxy-1-(oxetan-3-yl)pyrrolidin-3-yl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone



To a solution of ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((3R,4R)-4-hydroxypyrrolidin-3-yl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone (0.036g, 0.062 mmol, 1 eq) in DCE (0.5 mL) was added few drops of Acetic acid (2 μ L, 0.031 mmol, 0.5 eq) followed by oxetan-3-one (0.015 mL, 0.21 mmol, 3.3 eq) and the reaction mixture was heated at 60 °C for 2 hours. Added sodium triacetoxyborohydride (0.033 g, 0.16 mmol, 2.5 eq) and the solution was stirred at 23 °C for 24 hours. LCMS indicated the reaction was complete. The reaction mixture was diluted with EtOAc and washed with aqueous saturated sodium


bicarbonate solution. The combined organic layers were washed with saturated brine solution, dried over Na_2SO_4 and concentrated in vacuo. The residue was then purified by column chromatography on silica gel (DCM/MeOH) to isolate the product ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((3R,4R)-4-hydroxy-1-(oxetan-3-yl)pyrrolidin-3-yl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone (0.030 g, 75%).

Step 4: ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((3R,4R)-4-hydroxy-1-(oxetan-3-yl)pyrrolidin-3-yl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone




A solution of ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((3R,4R)-4-hydroxy-1-(oxetan-3-yl)pyrrolidin-3-yl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone (0.060 g, 0.095 mmol, 1 eq) in DCM (1 mL) was treated with TFA (0.73 mL, 9.5 mmol, 100 eq) for 16 hours. The reaction mixture was diluted with DCM and washed with aqueous saturated sodium bicarbonate solution. The combined organic layers were washed with saturated brine solution, dried over Na_2SO_4 and concentrated in vacuo. To the intermediate in DCM/MeOH (1/1, 1 mL) was added sodium acetate (0.016 g, 0.19 mmol, 2 eq) and the reaction was stirred at 23 °C for 2 hours. The reaction mixture was diluted with DCM and then washed with aqueous saturated sodium bicarbonate solution. The combined organic layers were washed with saturated brine solution, dried over Na_2SO_4 and concentrated in vacuo. The residue was then purified first by column chromatography on silica gel (DCM/MeOH containing 10% NH_4OH) and then by preparative-TLC to isolate the product ((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)(4-(((3R,4R)-4-hydroxy-1-(oxetan-3-yl)pyrrolidin-3-yl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)methanone (0.034g, 70%).

Example 26. Synthesis of Compound 156




Step 1: 1-bromo-2-(bromomethyl)-4-chlorobenzene



A solution of 1-bromo-4-chloro-2-methylbenzene (3.00 g, 14.60 mmol, 1.00 *eq*), NBS (2.34 g, 13.14 mmol, 0.90 *eq*) and AIBN (239.75 mg, 1.46 mmol, 0.10 *eq*) in CCl_4 (20.00 mL) was stirred at 90°C for 12 hrs. The solution was concentrated under vacuum to give a crude product 1-bromo-2-(bromomethyl)-4-chlorobenzene (5.40 g, crude) as a yellow solid which was used directly in the next step without further purification.


Step 2: 1-(2-bromo-5-chlorophenyl)-N-methylmethanamine



The solution of 1-bromo-2-(bromomethyl)-4-chlorobenzene (5.40 g, 18.99 mmol, 1.00 *eq*) in $\text{MeNH}_2/\text{H}_2\text{O}$ (30.00 mL) was stirred at 25°C for 15 hrs. When the reaction was complete, the product was extracted with EtOAc (50 mL * 3), the combined organic layers were concentrated under vacuum to give a crude product. The crude product was purified by column chromatography on silica gel (PE:EtOAc = 5:1 to 1:1) to obtain 1-(2-bromo-5-chlorophenyl)-N-methylmethanamine (1.00 g, yield: 22.45%) as a off-white solid. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 7.45 (d, 1H, $J =$


8.8 Hz), 7.40-7.39 (m, 1H), 7.10 (dd, 1H, $J = 8.4, 2.4$ Hz), 3.79 (s, 2H), 2.46 (s, 3H).

Step 3: N-(2-bromo-5-chlorobenzyl)-4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide




To the solution of 1-(2-bromo-5-chlorophenyl)-N-methylmethanamine (500.00 mg, 1.22 mmol, 1.00 *eq*) and 4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (314.98 mg, 1.34 mmol, 1.10 *eq*) in DMF (5.00 mL) was added DIPEA (315.35 mg, 2.44 mmol, 2.00 *eq*) and HATU (556.66 mg, 1.46 mmol, 1.20 *eq*), the resulting mixture was stirred at 25°C for 15 hrs. When the reaction was complete, H₂O (20 mL) was added, the product was extracted by EtOAc (20 mL * 3), the combined organic layers were concentrated under vacuum to give a crude product. The crude product was purified by *prep*-TLC (EtOAc) to obtain N-(2-bromo-5-chlorobenzyl)-4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (520.00 mg, yield: 81.20%) as a red oil. ¹H-NMR (400 MHz, CDCl₃) δ ppm 9.77 (br.s, 0.5H), 9.49 (br.s, 0.5H), 8.45 (d, 1H, $J = 11.4$ Hz), 7.63 (dd, 1H, $J = 8.4, 4.4$ Hz), 7.33-7.31 (m, 1H), 7.26-7.25 (m, 1H), 5.87 (s, 1H), 5.67 (s, 1H), 5.46 (s, 1H), 4.98 (s, 1H), 4.45-4.44 (m, 1H), 4.31-4.29 (m, 1H), 4.03-4.02 (m, 1H), 3.79 (t, 1H, $J = 8.0$ Hz), 3.71 (s, 1.5H), 3.55 (t, 1H, $J = 8.0$ Hz), 3.29 (s, 1.5H), 3.16 (d, 1H, $J = 6.0$ Hz), 2.65-2.61 (m, 1H), 2.05-1.83 (m, 2H), 1.05 (t, 1H, $J = 8.4$ Hz), 0.90 (t, 1H, $J = 8.0$ Hz), 0.07 (s, 4.5H), 0.00 (s, 4.5H).

Step 4: N-(5-chloro-2-cyanobenzyl)-4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide



The mixture of N-(2-bromo-5-chlorobenzyl)-4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (420.00 mg, 670.91 μmol , 1.00 *eq*), Zn(CN)_2 (630.22 mg, 5.37 mmol, 340.66 μL , 8.00 *eq*) and $\text{Pd(PPh}_3)_4$ (77.53 mg, 67.09 μmol , 0.10 *eq*) was dissolved in DMF (3.00 mL) in a sealed tube, it was radiated with microwave at 140°C for 3 hrs. After 3 hrs, LCMS showed the starting material was not consumed completely, so more Zn(CN)_2 (2 *eq*) and $\text{Pd(PPh}_3)_4$ (0.1 *eq*) was added, it was radiated with microwave at 150°C for another 2 hrs. When the reaction was complete, H_2O (15 mL) was added, the product was extracted by EtOAc (20 mL * 3). The combined organic layers were dried over anhydrous Na_2SO_4 , concentrated to give a crude product. The crude product was purified by *prep*-TLC (EtOAc) to obtain N-(5-chloro-2-cyanobenzyl)-4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (520.00 mg, crude, including PPh_3O) was obtained as a yellow oil. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 9.72 (br.s, 0.5H), 9.44 (br.s, 0.5H), 8.45 (d, 1H, $J = 6.4$ Hz), 7.79-7.54 (m, 3H), 5.86 (s, 1H), 5.70 (s, 1H), 5.67 (s, 1H), 5.11 (s, 1H), 4.45-4.44 (m, 1H), 4.30-4.29 (m, 1H), 4.03-4.02 (m, 1H), 3.81-3.76 (m, 2.5H), 3.59 (t, 1H, $J = 8.0$ Hz), 3.32 (s, 1.5H), 3.18-3.16 (m, 1H), 2.19-1.86 (m, 3H), 1.04 (t, 1H, $J = 8.0$ Hz), 0.92 (t, 1H, $J = 8.0$ Hz), 0.06 (s, 4.5H), 0.00 (s, 4.5H).


Step 5: N-(5-chloro-2-cyanobenzyl)-4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide



To the mixed solvents TFA (10.00 mL) and DCM (10.00 mL) was added N-(5-chloro-2-cyanobenzyl)-4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (520.00 mg, 908.88 μ mol, 1.00 *eq*), the resulting mixture was stirred at 25°C for 1 hr. The solvent was evaporated by N₂ to give a crude product. The crude product was dissolved in MeOH (15.00 mL), adjusted to pH = 7-8 by NaHCO₃, and KOAc (178.39 mg, 1.82 mmol, 2.00 *eq*) was added, it was stirred at 50°C for 2 hrs. When the reaction was complete, the mixture was concentrated under vacuum to give a crude product which was dissolved in EtOAc (50 mL), washed by H₂O (15 mL * 3). The organic layer was concentrated under vacuum to give a crude product which was purified by acidic *prep*-HPLC (TFA) to obtain N-(5-chloro-2-cyanobenzyl)-4-(((1R,2S,3R)-2,3-dihydroxycyclopentyl)amino)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (171.00 mg, yield: 33.85%, TFA) as a white solid.


Example 27. Synthesis of Compound 226

Step 1: Methyl (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-((tert-butyldiphenylsilyl)oxy)cyclopentane-1-carboxylate:




To a solution of methyl (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-hydroxycyclopentane-1-carboxylate (1.50 g, 5.78 mmol, 1.00 *eq*) and imidazole (590.74 mg, 8.67 mmol, 1.50 *eq*) in DMF (10.00 mL) was added TBDPSCI (1.67 g, 6.07 mmol, 1.05 *eq*) at 0°C. The reaction was stirred at 15°C for 16 hrs. TLC (PE:EtOAc = 5:1, R_f = 0.43) showed the reaction was complete. The solution was poured into water (20 mL) and extracted with EtOAc (10 mL * 3). The organic layer was dried over Na₂SO₄ and concentrated. The residue was recrystallized from PE (1 mL) to give methyl (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-((tert-butyldiphenylsilyl)oxy)cyclopentane-1-carboxylate (2.80 g, yield: 97.40%) as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.69-7.64 (m, 4H), 7.43-7.37 (m, 6H), 4.10 (br.s, 1H), 3.92-3.97 (m, 2H), 3.66 (s, 3H), 2.75-2.71 (m, 1H), 2.46-2.43 (m, 1H), 2.01-1.95 (m, 2H), 1.65-1.61 (m, 1H), 1.40 (s, 9H), 1.05 (s, 9H).

Step 2: tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-(hydroxymethyl)cyclopentyl)carbamate:



To a solution of methyl (1R,3R,4R)-3-((tert-butoxycarbonyl)amino)-4-((tert-butyldiphenylsilyl)oxy)cyclopentane-1-carboxylate (2.80 g, 5.63 mmol, 1.00 *eq*) in THF (30.00 mL) was added LiAlH₄ (427.32 mg, 11.26 mmol, 2.00 *eq*) at -30°C. The reaction was slowly warmed to 15°C and stirred for 2 hrs. TLC (PE:EtOAc = 3:1, R_f = 0.24) showed the reaction was complete. The reaction was quenched with 0.43 mL of H₂O and 0.43 mL of 10% *aq.* NaOH at 0°C. The mixture was filtered and the filtrate was concentrated. The residue was washed with PE (5 mL). The solid was collected. tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-(hydroxymethyl)cyclopentyl)carbamate (2.30 g, yield: 86.98%) as a white solid was obtained. ¹H-NMR (400 MHz, CDCl₃) δ ppm 7.69-7.65 (m, 4H), 7.43-7.38 (m, 6H), 4.11-4.10 (m, 1H), 3.89 (br.s, 2H), 3.54 (br.s, 2H), 2.14-2.10 (m, 1H), 1.97-1.89 (m, 2H), 1.62-1.58 (m, 1H), 1.39 (s, 9H), 1.06 (s, 9H).

Step 3: tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-formylcyclopentyl)carbamate:



To a solution of tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-(hydroxymethyl)cyclopentyl)carbamate (2.30 g, 4.90 mmol, 1.00 *eq*) in DCM (50.00 mL) was added Dess-Martin periodinane reagent (3.12 g, 7.35 mmol, 1.50 *eq*) at 0°C. The reaction was stirred at 15°C for 16 hrs. TLC (PE:EtOAc = 3:1, R_f = 0.7) showed the reaction was complete. The reaction was quenched with sat'd *aq.* NaHCO₃ (30 mL) at 0°C and extracted with DCM (30 mL). The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by column chromatography on silica gel (PE:EtOAc = 30:1). tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-formylcyclopentyl)carbamate (900.00 mg, yield: 39.27%) as a yellow oil


was obtained. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 9.55 (s, 1H), 7.61-7.56 (m, 4H), 7.37-7.31 (m, 6H), 4.08 (d, 1H, $J = 7.2$ Hz), 3.93-3.89 (m, 1H), 3.79 (br.s, 1H), 2.62-2.55 (m, 1H), 2.44-2.40 (m, 1H), 1.86-1.83 (m, 2H), 1.57-1.52 (m, 1H), 1.34 (s, 9H), 0.97 (s, 9H).

Step 4: tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-(difluoromethyl)cyclopentyl)carbamate:



To a solution of tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-formylcyclopentyl)carbamate (900.00 mg, 1.92 mmol, 1.00 *eq*) in DCM (50.00 mL) was added DAST (928.45 mg, 5.76 mmol, 3.00 *eq*) at 0°C. The reaction was stirred at 15°C for 5 hrs. TLC (PE:EtOAc = 3:1, $R_f = 0.6$) showed the reaction was complete. The solution was quenched with sat'd aq. NaHCO_3 (20 mL) and extracted with DCM (20 mL). The organic layer was dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography on silica gel (PE:EtOAc = 30:1~20:1). tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-(difluoromethyl)cyclopentyl)carbamate (250.00 mg, yield: 26.59%) as a yellow oil was obtained. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 7.62-7.56 (m, 4H), 7.34-7.31 (m, 6H), 5.76-5.43 (m, 1H), 3.88-3.82 (m, 2H), 2.30-2.24 (m, 1H), 2.07-2.04 (m, 1H), 1.80-1.75 (m, 1H), 1.65-1.57 (m, 1H), 1.33 (s, 9H), 0.99 (s, 9H).


Step 5: (1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-(difluoromethyl)cyclopentan-1-amine:



To a solution of tert-butyl ((1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-(difluoromethyl)cyclopentyl)carbamate (50.00 mg, 102.11 μmol , 1.00 *eq*) in EtOAc (2.00 mL) was added HCl/EtOAc (10.00 mL, 4 M) at 15°C. The reaction was stirred at 15°C for 1 hr. TLC (PE:EtOAc = 3:1, $R_f = 0.05$) showed the reaction was complete. The solvent was blown to dryness by N_2 . The residue wasn't further purified. (1R,2R,4R)-2-((tert-butyldiphenylsilyl)oxy)-4-


(difluoromethyl)cyclopentan-1-amine (40.00 mg, yield: 91.95%, HCl) as a yellow oil was obtained. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ ppm 7.70-7.68 (m, 4H), 7.49-7.43 (m, 6H), 5.81 (td, 1H, $J = 57.2$, 4.8 Hz), 4.25 (dd, 1H, $J = 12.0$, 5.6 Hz), 3.50 (dd, 1H, $J = 13.2$, 6.4 Hz), 2.22-2.19 (m, 1H), 1.80-1.74 (m, 2H), 1.66-1.63 (m, 1H), 1.08 (s, 9H).

Step 6: (4-(((1R,2R,4R)-2-((tert-butyl)phenylsilyl)oxy)-4-(difluoromethyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone:



To a solution of (1R,2R,4R)-2-((tert-butyl)phenylsilyl)oxy)-4-(difluoromethyl)cyclopentan-1-amine (40.00 mg, 93.89 μmol , 1.00 *eq*, HCl) and (4-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone (48.07 mg, 93.89 μmol , 1.00 *eq*) in dioxane (10.00 mL) was added DIPEA (60.67 mg, 469.45 μmol , 5.00 *eq*). The reaction was heated at 90°C for 0.5 hr. LCMS showed the reaction was complete. The solution was concentrated. The residue was purified by *prep*-TLC (PE:EtOAc = 3:1). (4-(((1R,2R,4R)-2-((tert-butyl)phenylsilyl)oxy)-4-(difluoromethyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone (38.00 mg, yield: 46.78%) as a yellow oil was obtained.

Step 7: (4-(((1R,2R,4R)-4-(difluoromethyl)-2-hydroxycyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone (Compound 226):






















A solution of (4-(((1R,2R,4R)-2-((tert-butyl(diphenyl)silyloxy)-4-(difluoromethyl)cyclopentyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone (38.00 mg, 43.93 μmol , 1.00 *eq*) in TBAF/THF (5.00 mL) was heated at 50°C for 2 hrs. LCMS showed (4-(((1R,2R,4R)-4-(difluoromethyl)-2-hydroxycyclopentyl)amino)-1-(hydroxymethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone was remained. The reaction mixture was concentrated. The residue was dissolved in EtOAc (20 mL) and washed with brine (10 mL*2). The organic layer was concentrated and dissolved in MeOH (20.00 mL). KOAc (21.56 mg, 219.65 μmol , 5.00 *eq*) was added to the reaction. The reaction was heated at 50°C for 16 hrs. LCMS showed the reaction was complete. The solution was concentrated. The residue was purified by *prep*-HPLC (MeOH/TFA system). (13.50 mg, yield: 50.34%, TFA) of (4-(((1R,2R,4R)-4-(difluoromethyl)-2-hydroxycyclopentyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)((2R,4S)-2-(2,5-difluorophenyl)-4-fluoropyrrolidin-1-yl)methanone as a yellow solid was obtained.















Example 28. Synthesis of Other Compounds













Additional compounds of the invention were synthesized using similar techniques to those set forth in the above examples. The table below indicates the specific example ("Example") upon which the synthesis of each compound ("Cmpd") was based, as well as the appropriate amino alcohol and amine that were used to synthesize each specific compound.

Table 1. Protocol and Intermediates Used for Synthesizing Exemplary Compounds.

Cmpd	Amino Alcohol	Amine	Example
1			1
2			1
3			1
4			1
5			1
6			1










Cmpd	Amino Alcohol	Amine	Example
7			1
8			1
9			2
10			2
11			1
12			2












Cmpd	Amino Alcohol	Amine	Example
19			3
20			3
21			3
22			1
23			1
24			1
25			1












Cmpd	Amino Alcohol	Amine	Example
13			2
14			3
15			3
16			3
17			3
18			3










Cmpd	Amino Alcohol	Amine	Example
32			1
33			1
34			2
35			2
36			1
37			1
38			1











Cmpd	Amino Alcohol	Amine	Example
26			1
27			1
28			1
29			1
30			1
31			1













Cmpd	Amino Alcohol	Amine	Example
45			1
46			1
47			1
48			1
49			1
50			1











Cmpd	Amino Alcohol	Amine	Example
39			1
40			1
41			1
42			1
43			1
44			1







Cmpd	Amino Alcohol	Amine	Example
57			1
58			1
59			1
60			1
61			3
62			3
63			1











Cmpd	Amino Alcohol	Amine	Example
51			23
52			1
53			1
54			1
55			1
56			1











Cmpd	Amino Alcohol	Amine	Example
70			2
71			2
72			1
73			1
74			1








Cmpd	Amino Alcohol	Amine	Example
64			1
65			1
66			2
67			2
68			1
69			1













Cmpd	Amino Alcohol	Amine	Example
80			1
81			1
82			1
83			1
84			22
85			1

Cmpd	Amino Alcohol	Amine	Example
75			3
76			3
77			1
78			1
79			1

Cmpd	Amino Alcohol	Amine	Example
91			2
92			1
93			1
94			1
95			3









Cmpd	Amino Alcohol	Amine	Example
86			2
87			2
88			1
89			1
90			2











Cmpd	Amino Alcohol	Amine	Example
102			23
103			1
104			1
105			1
106			1














Cmpd	Amino Alcohol	Amine	Example
96			3
97			2
98			2
99			1
100			1
101			1











Cmpd	Amino Alcohol	Amine	Example
113			23
114			1
115			2
116			2
117			1
118			1













Cmpd	Amino Alcohol	Amine	Example
107			1
108			1
109			23
110			1
111			23
112			1













Cmpd	Amino Alcohol	Amine	Example
124			2
125			2
126			1
127			1
128			1













Cmpd	Amino Alcohol	Amine	Example
119			1
120			1
121			1
122			1
123			3













Cmpd	Amino Alcohol	Amine	Example
135			24
136			1
137			1
138			1
139			1
140			24
141			24

Cmpd	Amino Alcohol	Amine	Example
129			1
130			1
131			1
132			1
133			1
134			1

Cmpd	Amino Alcohol	Amine	Example
148			1
149			1
150			1
151			3
152			24
153			1











Cmpd	Amino Alcohol	Amine	Example
142			1
143			24
144			1
145			1
146			24
147			1










Cmpd	Amino Alcohol	Amine	Example
160			I
161			I
162			I
163			I
164			I
165			24











Cmpd	Amino Alcohol	Amine	Example
154			24
155			I
156			24
157			I
158			I
159			I











Cmpd	Amino Alcohol	Amine	Example
171			1
172			1
173			1
174			1
175			1











Cmpd	Amino Alcohol	Amine	Example
166			1
167			24
168			1
169			1
170			1











Cmpd	Amino Alcohol	Amine	Example
181			1
182			1
183			24
184			1
185			1







Cmpd	Amino Alcohol	Amine	Example
176			1
177			1
178			3
179			1
180			1











Cmpd	Amino Alcohol	Amine	Example
191			1
192			1
193			1
194			1
195			1











Cmpd	Amino Alcohol	Amine	Example
186			3
187			3
188			1
189			1
190			1











Cmpd	Amino Alcohol	Amine	Example
201			1
202			1
203			1
204			1
205			1

Cmpd	Amino Alcohol	Amine	Example
196			1
197			1
198			1
199			1
200			1

Cmpd	Amino Alcohol	Amine	Example
211			1
212			1
213			1
214			1
215			1

Cmpd	Amino Alcohol	Amine	Example
206			1
207			3
208			3
209			1
210			22

Cmpd	Amino Alcohol	Amine	Example
221			1
222			1
223			1
224			1
225			1

Cmpd	Amino Alcohol	Amine	Example
216			1
217			1
218			1
219			1
220			1

Cmpd	Amino Alcohol	Amine	Example
231			1
232			22
233			1
234			1
235			1

Cmpd	Amino Alcohol	Amine	Example
226			27
227			1
228			1
229			23
230			1

Cmpd	Amino Alcohol	Amine	Example
241			1
242			1
243			1
244			1

Cmpd	Amino Alcohol	Amine	Example
236			1
237			1
238			1
239			1
240			1

The NMR and LC MS data obtained for compounds disclosed herein are shown in Figure 1.

Example 29. Assays

NTRK1 Wild Type Assay at 1 mM ATP

In each well of a 384-well plate, 1 nM – 1.5 nM of wild type NTRK1 enzyme (BPS Bioscience; 40280) was incubated in a total of 12.5 μ L of buffer (100 mM HEPES pH 7.5, 0.015% Brij 35, 10 mM $MgCl_2$, 1mM DTT) with 1-2 μ M CSKtide (Tuft's University or Anaspec; FITC-AHA-KKKKD DIYFFFG-NH₂) and 1 mM ATP at 25°C for 60 minutes in the presence or absence of a dosed concentration series of compound (1% DMSO final concentration). The reaction was stopped by the addition of 70 μ L of Stop buffer (100 mM HEPES pH 7.5, 0.015% Brij 35, 35 mM EDTA and 0.2% of Coating Reagent 3 (Caliper Lifesciences)). The plate was then read on a Caliper EZReader 2 (protocol settings: -1.7 psi, upstream voltage -500, downstream voltage -3000, post sample sip 35s). Data was normalized to 0% and 100% inhibition controls and the IC₅₀ calculated using a 4-parameter fit in the CORE LIMS.

NTRK Wild Type and G595R Mutant Cellular Assays Protocol

KM12 wild type colon carcinoma cell line harboring the TPM3-NTRK1 fusion protein was obtained from the National Cancer Institute (NCI). This line has been previously shown to be dependent upon the NTRK activity derived from the NTRK fusion protein for growth and survival. The KM12 Cliff (G595R) cell line was generated by mutagenizing the wild type KM12 line with a DNA methylating agent and subsequently selecting for clones that were resistant to chronic exposure to high concentration of a known NTRK inhibitor (Crizotinib). Cells were first plated in 384-well plates at 1000 cells/well in complete media (10% FBS and 1% pen/strep) and incubated overnight at 37°C. Cells were then dosed with test articles at varying concentrations using the Bravo liquid handling system. Concentrations ranged from 25 μ M down to 9.5pM (4-fold dilutions, 10 concentrations total). Each compound was run in duplicate per plate. DMSO and staurosporine (25 μ M) were included on each plate as negative and positive controls for growth inhibition. 72hr after dosing, assay plates were developed using CellTiter-Glo (Promega) and resultant luminescence was read on the Envision plate reader. IC₅₀ determinations were calculated using a 4-parameter curve fitting algorithm

The table below summarizes the results from the biological assays described above. The following designations are used to indicate IC₅₀ in each assay: A < 10.00 nM; B = 10.01-100.0 nM;

C = 100.01-1000.0 nM; and D > 1000.1 nM; "ND" = not determined.

Compound Number	Enz NTRK1	KM12 (WT)	KM12 (G595R)
1	B	B	B
2	C	C	C
3	D	D	D
4	C	D	D
5	C	C	C
6	A	B	B
7	B	B	B
8	B	B	C
9	B	B	C
10	C	C	D
11	C	C	C
12	C	C	D
13	B	B	C
14	C	C	C
15	C	C	D
16	C	D	D
17	B	B	C
18	C	C	D
19	D	D	D
20	B	B	C
21	C	C	D
22	C	D	D
23	A	B	B
24	B	C	C
25	B	B	C
26	B	B	C
27	A	B	B
28	A	B	B
29	A	B	B
30	C	B	C
31	C	C	C
32	B	C	C
33	B	B	B
34	B	B	C
35	C	C	D

Compound Number	Enz NTRK1	KM12 (WT)	KM12 (G595R)
36	B	C	C
37	B	C	C
38	C	C	C
39	A	B	B
40	B	C	C
41	B	C	C
42	C	D	D
43	B	B	C
44	B	B	C
45	A	A	A
46	A	A	A
47	C	C	C
48	B	B	C
49	A	B	B
50	B	B	C
51	C	C	C
52	B	B	B
53	B	B	C
54	B	C	C
55	B	B	B
56	B	C	C
57	A	A	B
58	A	B	B
59	C	C	C
60	B	B	B
61	C	C	D
62	B	B	C
63	A	B	B
64	C	C	C
65	C	ND	ND
66	D	D	D
67	D	D	D
68	A	A	A
69	A	B	B
70	A	B	B

Compound Number	Enz NTRK1	KM12 (WT)	KM12 (G595R)
71	B	B	C
72	A	B	B
73	A	A	B
74	A	B	B
75	B	C	D
76	A	B	B
77	A	B	B
78	A	A	B
79	A	A	B
80	B	C	C
81	B	C	C
82	B	C	C
83	B	C	C
84	A	B	B
85	A	B	B
86	A	A	A
87	A	B	B
88	A	A	A
89	A	B	A
90	D	D	D
91	B	B	C
92	A	A	A
93	A	A	A
94	C	C	D
95	A	A	A
96	B	B	C
97	A	B	C
98	C	D	D
99	A	A	A
100	B	B	C
101	C	C	C
102	A	B	B
103	D	D	D
104	A	A	A
105	A	B	B
106	A	A	A
107	A	B	B
108	B	C	C

Compound Number	Enz NTRK1	KM12 (WT)	KM12 (G595R)
109	A	B	C
110	A	B	B
111	A	B	B
112	B	B	C
113	A	B	B
114	A	A	A
115	A	B	B
116	A	A	A
117	C	C	D
118	A	A	A
119	A	A	A
120	B	B	B
121	A	A	A
122	A	A	A
123	A	B	B
124	A	ND	ND
125	A	ND	ND
126	A	A	A
127	A	A	A
128	A	A	A
129	A	B	B
130	A	A	A
131	C	C	C
132	B	B	C
133	C	C	C
134	C	C	C
135	A	A	A
136	A	B	B
137	B	B	B
138	C	C	C
139	B	B	B
140	A	B	A
141	A	B	A
142	B	B	C
143	B	B	B
144	B	B	C
145	C	B	C
146	A	B	B

Compound Number	Enz NTRK1	KM12 (WT)	KM12 (G595R)
147	B	C	C
148	A	A	A
149	A	B	B
150	B	B	B
151	B	B	B
152	A	B	A
153	B	B	B
154	A	B	B
155	B	B	B
156	B	B	A
157	A	B	B
158	A	B	B
159	A	B	B
160	A	B	B
161	B	B	B
162	A	B	B
163	A	B	B
164	B	B	B
165	A	A	A
166	A	C	C
167	A	A	A
168	A	A	A
169	B	B	B
170	A	A	A
171	B	B	B
172	A	B	B
173	A	A	A
174	A	A	A
175	A	B	B
176	A	B	B
177	A	A	A
178	A	B	B
179	B	B	C
180	A	A	A
181	A	B	B
182	A	A	B
183	A	A	A
184	A	A	A

Compound Number	Enz NTRK1	KM12 (WT)	KM12 (G595R)
185	A	A	A
186	A	A	A
187	A	B	B
188	A	A	A
189	A	A	B
190	A	B	B
191	A	B	B
192	A	B	B
193	A	B	B
194	B	B	B
195	A	A	A
196	A	A	A
197	A	B	B
198	A	A	A
199	B	A	A
200	A	B	B
201	A	A	A
202	A	B	B
203	A	A	A
204	A	A	A
205	A	A	A
206	A	A	A
207	B	B	B
208	A	A	A
209	A	B	B
210	A	A	A
211	A	A	A
212	A	A	A
213	A	A	A
214	A	A	A
215	A	A	A
216	A	A	A
217	A	A	A
218	A	A	A
219	A	A	A
220	A	A	A
221	A	A	A
222	A	B	B

Compound Number	Enz NTRK1	KM12 (WT)	KM12 (G595R)
223	A	B	B
224	A	A	A
225	A	B	B
226	A	A	A
227	A	A	A
228	B	B	B
229	A	A	A
230	A	B	B
231	B	B	C
232	A	A	A
233	A	A	A

Compound Number	Enz NTRK1	KM12 (WT)	KM12 (G595R)
234	A	A	A
235	A	B	B
236	A	A	B
237	A	B	B
238	ND	B	B
239	A	A	B
240	A	A	A
241	A	A	A
242	A	A	A
243	A	A	A
244	A	A	A

Incorporation by Reference


All publications and patents mentioned herein are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually indicated to be incorporated by reference.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

We claim:

1. A compound of Formula (I):



(I), or a pharmaceutically acceptable salt thereof, wherein:

Rings A and B are each independently selected from aryl, heteroaryl, cycloalkyl and heterocyclyl;

each L^1 and L^2 is independently selected from a bond, $-C(O)-$, $-N(R^1)-$, $-N(R^1)-C(O)-$, $-C(O)-N(R^1)-$, $-(C_1-C_6 \text{ alkylene})-N(R^1)-$, $-N(R^1)-(C_1-C_6 \text{ alkylene})-$, $-N(R^1)-C(O)-(C_1-C_6 \text{ alkylene})-$, and $-C(O)-N(R^1)-(C_1-C_6 \text{ alkylene})-$; wherein each alkylene portion of L^1 and L^2 , is independently substituted with 0-5 occurrences of R' ;

each R^A and R^B is independently selected from hydroxyl, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_1-C_6 alkoxy, halo, C_1-C_6 heteroalkyl, C_1-C_6 haloalkyl, C_1-C_6 haloalkoxy, C_1-C_6 hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, heterocyclylalkyl, nitro, cyano, $-C(O)R^1$, $-OC(O)R^1$, $-C(O)OR^1$, $-(C_1-C_6 \text{ alkylene})-C(O)R^1$, $-SR^1$, $-S(O)_2R^1$, $-S(O)_2-N(R^1)(R^1)$, $-(C_1-C_6 \text{ alkylene})-S(O)_2R^1$, $-(C_1-C_6 \text{ alkylene})-S(O)_2-N(R^1)(R^1)$, $-N(R^1)(R^1)$, $-C(O)-N(R^1)(R^1)$, $-N(R^1)-C(O)R^1$, $-N(R^1)-C(O)OR^1$, $-(C_1-C_6 \text{ alkylene})-N(R^1)-C(O)R^1$, $-N(R^1)S(O)_2R^1$, and $-P(O)(R^1)(R^1)$; wherein each alkyl, alkenyl, alkynyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, and heterocyclylalkyl portion of each of R^A and R^B is independently substituted with 0-5 occurrences of R^a ; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^a ;

each R^1 is independently selected from hydrogen, hydroxyl, halo, thiol, C_1-C_6 alkyl, C_1-C_6 thioalkyl, C_1-C_6 alkoxy, C_1-C_6 haloalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, wherein each alkyl, thioalkyl, alkoxy, haloalkyl, hydroxyalkyl, heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl portion of R^1 is independently substituted with 0-5 occurrences of R^b , or 2 R^1 together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^b ;

each R^a and R^b is independently selected from C₁-C₆ alkyl, halo, hydroxyl, C₁-C₆ haloalkyl, C₁-C₆ heteroalkyl, C₁-C₆ hydroxyalkyl, C₁-C₆ alkoxy, cycloalkyl, heterocyclyl, and cyano, wherein each alkyl, haloalkyl, heteroalkyl, hydroxyalkyl, alkoxy, cycloalkyl and heterocyclyl portion of R^a and R^b is independently substituted with 0-5 occurrences of R';

each R' is independently selected from C₁-C₆ alkyl, C₁-C₆ heteroalkyl, halo, hydroxyl, C₁-C₆ haloalkyl, C₁-C₆ hydroxyalkyl, cycloalkyl and cyano; or 2 R' together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring; and

p is 0, 1, 2, 3, 4, or 5; and

q is 0, 1, 2, 3, or 4.

2. The compound of claim 1, wherein Ring A is selected from cycloalkyl, heterocyclyl, and cycloalkenyl.

3. The compound of claim 2, wherein Ring A is selected from cyclopentyl, cyclohexyl, tetrahydropyranyl, tetrahydrofuranyl, pyrrolidinyl, and cyclopentenyl.

4. The compound of any one of claims 1 to 3, wherein Ring B is selected from aryl, heteroaryl, or heterocyclyl.


5. The compound of claim 4, wherein ring B is selected from phenyl, pyridyl, or pyrrolidinyl.

6. The compound of any one of claims 1 to 3, wherein each L¹ is a bond, -C(O)-, or -N(R¹)-; and L² is -N(R¹)-C(O)-(C₁-C₆ alkylene)- or -C(O)-N(R¹)-(C₁-C₆ alkylene)-.

7. The compound of claim 6, wherein:

L¹ is -NH-; and

L² is selected from -C(O)-NH-CH(CH₂OH)-*, -C(O)-N(CH₃)-CH₂-, -C(O)-N(CH₃)-CH(CH₃)-*, -C(O)N(CH₂CH₃)CH₂-, -C(O)NHCH(CH₃)-*, -C(O)N(CD₃)CH₂-, -



wherein "*" represents a portion of L^2 bound to ring B.

8. The compound of claim 7, wherein L^1 is -NH- , L^2 is -C(O)- and ring B is pyrrolidin-1-yl.

9. The compound of any one of claims 1 to 8, wherein each R^1 is independently selected from hydrogen and $\text{C}_1\text{-C}_6$ alkyl substituted with 0-5 occurrences of R^b .

10. The compound of claim 9, wherein each R^1 is independently selected from hydrogen and -CH_3 .

11. The compound of any one of claims 1 to 10, wherein each R^A and R^B is independently selected from hydroxyl, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, halo, $\text{C}_1\text{-C}_6$ heteroalkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ haloalkoxy, $\text{C}_1\text{-C}_6$ hydroxyalkyl, cycloalkyl, aryl, heteroaryl, nitro, cyano, -C(O)R^1 , -OC(O)R^1 , -C(O)OR^1 , -SR^1 , $\text{-S(O)}_2\text{R}^1$, $\text{-S(O)}_2\text{-N(R}^1\text{)(R}^1\text{)}$, $\text{-N(R}^1\text{)(R}^1\text{)}$, $\text{-C(O)-N(R}^1\text{)(R}^1\text{)}$, $\text{-N(R}^1\text{)-C(O)R}^1$, $\text{-N(R}^1\text{)-C(O)OR}^1$, and $\text{-N(R}^1\text{)S(O)}_2\text{R}^1$; wherein each alkyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, and heteroaryl portion of each R^a and each R^b is independently substituted with 0-5 occurrences of R^a ; or 2 R^A or 2 R^B together with the carbon atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of R^a .

12. The compound of any one of claims 1 to 6, wherein each R^A is independently selected from hydroxyl, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, halo, $\text{-C(O)-N(R}^1\text{)(R}^1\text{)}$, -C(O)OR^1 , $\text{-S(O)}_2\text{R}^1$, $\text{C}_1\text{-C}_6$ haloalkyl, -CN , oxetanyl, and $\text{C}_1\text{-C}_6$ hydroxyalkyl, or two R^A bound to adjacent ring carbon atoms on ring A are taken together to form a $\text{C}_3\text{-C}_6$ cycloalkyl fused to ring A.

13. The compound of claim 12, wherein each R^A is independently selected from hydroxyl, fluoro, oxetan-3-yl, -CHF_2 , $\text{-CH}_2\text{CH}_3$, $\text{-C(CH}_3\text{)}_2\text{OH}$, -OCH_3 , $\text{-C(O)N(CH}_3\text{)}_2$, -C(O)OCH_3 ,

-S(O)₂CH₃; or two R^A bound to adjacent ring carbon atoms on ring A are taken together to form a cyclopropyl fused to ring A.


14. The compound of any one of claims 1 to 13, wherein each R^B is independently selected from halo, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, aryl, heteroaryl, oxo, and C₁-C₆ haloalkoxy.

15. The compound of any one of claims 1-14, wherein Ring B is pyrrolidinyl and at least one R^B is optionally substituted aryl or heteroaryl.

16. The compound of claim 15, wherein Ring B is pyrrolidinyl, and at least one R^B is selected from 2,3,5-trifluorophenyl, 2,3-difluorophenyl, 2,5-difluorophenyl, 2-chloro-5-fluorophenyl, 2-chloro-5-fluoropyridin-3-yl, 2-cyano-5-fluorophenyl, 2-fluoro-5-chlorophenyl, 2-methoxy-3,5-difluorophenyl, 2-methoxy-5-fluoropyridin-3-yl, 2-trifluoromethoxy-5-fluorophenyl, 3,5-difluorophenyl, 3-chloro-5-fluorophenyl, 3-cyano-5-fluorophenyl, 3-difluoromethoxy-5-fluorophenyl, 3-fluorophenyl, 5-fluoropyridin-3-yl, and phenyl.


17. The compound of any one of claims 1-14, wherein Ring B is other than pyrrolidinyl, and each R^B is independently selected from chloro, fluoro, oxo, -CH₃, -CF₃, -CN, -OCH₃, -OCF₃, and -OCHF₂.

18. The compound of claim 1, wherein the compound is a compound of Formula (Ia):



(Ia), or a pharmaceutically acceptable salt thereof, wherein each of Ring A, R^A, R^B, p and q are as defined in claim 1.

19. The compound of claim 1, wherein the compound is a compound of Formula (II):



(II) or a pharmaceutically acceptable salt thereof,

wherein:

R^{1a} is selected from hydrogen, C_1 - C_6 alkyl, and deuterated C_1 - C_6 alkyl;

R^{1b} is selected from hydrogen and C_1 - C_6 alkyl; and

Ring A, Ring B, R' , R^A , R^B , p and q are as defined in claim 1.

20. A compound selected from any one of the compounds set forth in Table 1, or a pharmaceutically acceptable salt thereof.

21. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of any one of claims 1 to 20.

22. The composition of claim 21 for use in treating a subject suffering from a condition mediated by aberrant neurotrophic tyrosine receptor kinase (NTRK) activity.

23. The composition of claim 21 for use in treating a subject who has developed resistance to a cancer treatment.

24. The composition of claim 21 for use in treating a condition selected from the group consisting of non-small cell lung cancer, breast cancer, melanoma, low and high grade glioma, glioblastoma, pediatric astrocytoma, colorectal cancer, papillary thyroid carcinoma, pancreatic adenocarcinoma, head and neck cancer, cholangiocarcinoma, acute myelogenous leukemia, secretory breast cancer, salivary cancer, and spitzoid neoplasms,.

FIGURE 1

#	Structure	NMR; LCMS
1		1H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.13-7.02 (m, 3H), 4.66 (s, 2H), 4.17 (d, 2H, J = 3.2 Hz), 2.32-2.29 (m, 1H), 2.11-2.08 (m, 1H), 1.91-1.87 (m, 2H), 1.86-1.74 (m, 2H).; LCMS: 389.1
2		1H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.319-7.15 (m, 1H), 7.02-7.01 (m, 1H), 6.99-6.88 (m, 1H), 4.61 (s, 2H), 4.38-4.31 (m, 1H), 4.13-4.11 (m, 1H), 4.03-4.02 (m, 1H), 2.44-2.41 (m, 1H), 2.34 (s, 3H), 2.16-2.13 (m, 1H), 1.79-1.70 (m, 2H).; LCMS: 401.1
3		1H-NMR (400 MHz, CD ₃ OD) δ ppm 8.21 (s, 1H), 7.15-7.09 (m, 2H), 7.08-7.01 (m, 1H), 4.65 (s, 2H), 4.41-4.38 (m, 1H), 4.03-3.96 (m, 1H), 2.41-2.36 (m, 1H), 1.90-1.89 (m, 1H), 1.58-1.56 (m, 1H), 1.44-1.41 (m, 1H), 0.76-0.73 (m, 1H), 0.50-0.47 (m, 1H).; LCMS: 401.1
4		1H-NMR (400 MHz, CD ₃ OD) δ ppm 8.12 (s, 1H), 7.06-6.99 (m, 2H), 6.93-6.92 (m, 1H), 4.56 (s, 2H), 4.32-4.29 (m, 1H), 3.93-3.87 (m, 1H), 2.32-2.25 (m, 1H), 1.81-1.80 (m, 1H), 1.49-1.47 (m, 1H), 1.35-1.33 (m, 1H), 0.67-0.64 (m, 1H), 0.41-0.40 (m, 1H).; LCMS: 401.1

FIGURE 1





5		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.51-8.47 (m, 2H), 8.38 (d, 1H, J = 2.0 Hz), 7.74 (d, 1H, J = 9.2 Hz), 5.50 (s, 1H), 4.94 (s, 1H), 4.39-4.34 (m, 1H), 4.15-4.13 (m, 1H), 4.07-4.01 (m, 1H), 3.70 (s, 1.5H), 3.20 (s, 1.5H), 2.47-2.44 (m, 1H), 2.19-2.16 (m, 1H), 1.84-1.71 (m, 2H).; LCMS: 402.1
6		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40 (d, 1H, J = 4.0 Hz), 7.22-7.07 (m, 3H), 5.51 (s, 1H), 4.26-4.17 (m, 2H), 3.70 (s, 1.5H), 3.21 (s, 1.5H), 2.38-2.36 (m, 1H), 2.17-2.06 (m, 1H), 1.96-1.76 (m, 4H).; LCMS: 403.1
7		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 7.22-7.21 (m, 1H), 7.13-7.11 (m, 1H), 7.04-7.02 (m, 1H), 5.56-5.51 (m, 1H), 4.16-4.13 (m, 2H), 2.32-2.29 (m, 1H), 2.09-2.08 (m, 1H), 1.90-1.70 (m, 4H), 1.59 (d, 3H, J = 7.2 Hz).; LCMS: 403.1
8		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32 (s, 1H), 7.15-7.09 (m, 2H), 7.05-7.03 (m, 1H), 4.67 (s, 2H), 4.15-4.11 (m, 1H), 4.05-4.04 (m, 1H), 3.98-3.95 (m, 1H), 3.85-3.84 (m, 1H), 3.56 (t, 1H, J = 10.8 Hz), 3.43 (t, 1H, J = 10.8 Hz), 2.10-2.07 (m, 1H), 1.76-1.68 (m, 1H).; LCMS: 405.0

FIGURE 1





9		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35 (s, 1H), 7.16-7.03 (m, 3H), 4.68 (s, 2H), 4.37 (q, 1H, J = 8.4 Hz), 4.12 (d, 1H, J = 4.4 Hz), 4.02 (dd, 1H, J = 8.4, 4.4 Hz), 2.45-2.42 (m, 1H), 2.17-2.14 (m, 1H), 1.82-1.71 (m, 2H).; LCMS: 405.1
10		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35 (s, 1H), 7.16-7.03 (m, 3H), 4.68 (s, 2H), 4.40-4.36 (m, 1H), 4.12 (d, 1H, J = 4.0 Hz), 4.02 (dd, 1H, J = 8.4, 4.4 Hz), 2.45-2.42 (m, 1H), 2.17-2.13 (m, 1H), 1.81-1.70 (m, 2H).; LCMS: 405.1
11		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 7.16-7.10 (m, 2H), 7.04-7.03 (m, 1H), 4.68 (s, 2H), 4.07-4.01 (m, 1H), 4.00-3.97 (m, 2H), 3.67-3.65 (m, 1H), 3.52-3.51 (m, 1H), 3.27-3.24 (m, 1H), 2.19-2.15 (m, 1H), 1.83-1.79 (m, 1H).; LCMS: 405.1
12		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.17 (s, 1H), 7.05-7.00 (m, 2H), 6.93-6.92 (m, 1H), 4.60-4.56 (m, 2H), 4.42-4.38 (m, 1H), 4.28-4.26 (m, 1H), 4.03-3.97 (m, 1H), 2.41-2.36 (m, 1H), 2.17-2.14 (m, 1H), 1.90-1.85 (m, 1H), 1.60-1.57 (m, 1H).; LCMS: 405.1

FIGURE 1





13		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.15-7.09 (m, 2H), 7.03-7.02 (m, 1H), 4.65 (s, 2H), 4.53-4.49 (m, 1H), 4.39-4.35 (m, 1H), 4.12-4.07 (m, 1H), 2.50-2.45 (m, 1H), 2.26-2.25 (m, 1H), 1.99-1.92 (m, 1H), 1.70-1.66 (m, 1H).; LCMS: 405.1
14		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.16-7.09 (m, 2H), 7.02-7.01 (m, 1H), 5.27-5.13 (m, 1H), 4.65 (s, 2H), 4.38-4.33 (m, 2H), 2.79-2.70 (m, 1H), 2.37-2.35 (m, 1H), 1.99-1.88 (m, 2H).; LCMS: 407.1
15		¹H-NMR (400 MHz, CDCl₃) δ ppm 9.76 (s, 1H), 8.35 (s, 1H), 7.57 (t, 1H, J = 5.6 Hz), 7.05-6.90 (m, 3H), 5.16 (d, 1H, J = 52.4 Hz), 4.68-4.62 (m, 2H), 4.39-4.34 (m, 1H), 4.15-4.12 (m, 1H), 2.78-2.70 (m, 1H), 2.48-2.40 (m, 1H), 2.18-2.08 (m, 1H), 1.93-1.81 (m, 1H).; LCMS: 407.1
16		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.12-7.07 (m, 2H), 7.02-7.00 (m, 1H), 5.15 (d, 1H, J = 53.6 Hz), 4.64 (s, 2H), 4.58-4.52 (m, 1H), 4.17 (dd, 1H, J = 12.8, 6.4 Hz), 2.59-2.51 (m, 2H), 1.99-1.87 (m, 2H).; LCMS: 407.1

FIGURE 1





17		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.14-7.08 (m, 2H), 7.02-7.00 (m, 1H), 5.13 (d, 1H, J = 53.2 Hz), 4.64 (s, 2H), 4.57-4.52 (m, 1H), 4.19-4.16 (m, 1H), 2.59-2.51 (m, 2H), 1.99-1.90 (m, 2H).; LCMS: 407.1
18		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.14 (s, 1H), 7.04-6.97 (m, 2H), 6.91-6.90 (m, 1H), 4.89-4.88 (m, 1H), 4.55 (s, 2H), 4.53-4.46 (m, 1H), 3.98-3.89 (m, 1H), 2.38-2.3 (m, 1H), 2.18-2.00 (m, 1H), 1.90-1.83 (m, 1H), 1.56-1.52 (m, 1H).; LCMS: 407.1
19		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.13 (s, 1H), 7.05-6.98 (m, 2H), 6.91-6.90 (m, 1H), 4.89-4.87 (m, 1H), 4.55 (s, 2H), 4.50-4.46 (m, 1H), 3.98-3.89 (m, 1H), 2.37-2.32 (m, 1H), 2.19-2.00 (m, 1H), 1.90-1.83 (m, 1H), 1.56-1.51 (m, 1H).; LCMS: 407.1
20		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.15-7.01 (m, 3H), 4.94-4.92 (m, 1H), 4.66 (s, 2H), 4.32-4.28 (m, 1H), 4.17-4.10 (m, 1H), 2.34-1.88 (m, 4H).; LCMS: 407.2

FIGURE 1





21		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.17 (s, 1H), 7.05-6.92 (m, 3H), 4.86-4.83 (m, 1H), 4.56 (s, 2H), 4.23-4.21 (m, 1H), 4.08-4.02 (m, 1H), 2.24-1.78 (m, 4H).; LCMS: 407.2
22		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (s, 1H), 7.58 (s, 1H), 7.47 (s, 2H), 4.67 (s, 2H), 4.39-4.37 (m, 1H), 4.13 (s, 1H), 4.02 (s, 1H), 2.44 (br.s, 1H), 2.16-2.15 (m, 1H), 1.79-1.69 (m, 2H).; LCMS: 412.1
23		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37, 8.34 (s, 1H), 7.25-7.18 (m, 1H), 6.93-6.82 (m, 2H), 5.40 (s, 1H), 4.42-4.35 (m, 1H), 4.16-3.99 (m, 2H), 3.61 (s, 1.5H), 3.22 (s, 1.5H), 2.47-2.43 (m, 1H), 2.31 (s, 1.5H), 2.28 (s, 1.5H), 2.18-2.12 (m, 1H), 1.83-1.75 (m, 2H).; LCMS: 415.1
24		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.07 (s, 1H), 9.45 (d, <i>J</i> = 7.7 Hz, 1H), 9.26 (t, <i>J</i> = 6.4 Hz, 1H), 8.27 (s, 1H), 7.11 – 6.81 (m, 3H), 6.51 (s, 0H), 5.10 (d, <i>J</i> = 4.8 Hz, 1H), 4.47 (d, <i>J</i> = 6.3 Hz, 2H), 4.04 (dd, <i>J</i> = 7.8, 4.0 Hz, 1H), 3.96 (dd, <i>J</i> = 11.2, 3.9 Hz, 1H), 3.82 (s, 3H), 3.81 – 3.71 (m, 1H), 3.65 (s, 1H), 3.43 (t, <i>J</i> = 9.3 Hz, 1H), 3.18 (dd, <i>J</i> = 11.2, 7.6 Hz, 1H), 1.93 – 1.85 (m, 1H), 1.53 – 1.43 (m, 1H); LCMS: 417

FIGURE 1





25		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.09 (s, 1H), 9.48 – 9.31 (m, 2H), 8.30 (s, 1H), 7.25 (td, <i>J</i> = 9.4, 4.5 Hz, 1H), 7.15 (td, <i>J</i> = 7.5, 6.7, 4.1 Hz, 2H), 4.81 (d, <i>J</i> = 4.1 Hz, 1H), 4.55 (d, <i>J</i> = 6.2 Hz, 2H), 4.40 (d, <i>J</i> = 7.1 Hz, 1H), 4.35 (d, <i>J</i> = 7.7 Hz, 2H), 3.64 (t, <i>J</i> = 5.1 Hz, 1H), 1.56 (dq, <i>J</i> = 8.7, 4.2 Hz, 1H), 1.32 (p, <i>J</i> = 4.6 Hz, 1H), 1.07 (q, <i>J</i> = 4.1 Hz, 1H), 0.43 (td, <i>J</i> = 8.3, 4.6 Hz, 1H); LCMS: 417
26		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37-8.36 (m, 1H), 7.17-7.07 (m, 3H), 4.81 (s, 2H), 4.16-4.13 (m, 2H), 3.66-3.60 (m, 2H), 2.40-2.27 (m, 1H), 2.20-2.03 (m, 1H), 1.91-1.72 (m, 4H), 1.40-1.24 (m, 3H); LCMS: 417.1
27		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.06 (s, 1H), 9.24 (d, <i>J</i> = 7.2 Hz, 1H), 9.11 (d, <i>J</i> = 8.6 Hz, 1H), 8.20 (s, 1H), 7.35 (ddd, <i>J</i> = 9.1, 5.7, 3.2 Hz, 1H), 7.20 (td, <i>J</i> = 9.3, 4.6 Hz, 1H), 7.10 (ddt, <i>J</i> = 9.0, 7.3, 3.6 Hz, 1H), 5.44 – 5.32 (m, 1H), 5.11 (t, <i>J</i> = 5.9 Hz, 1H), 5.02 – 4.91 (m, 1H), 4.16 (p, <i>J</i> = 6.6 Hz, 1H), 3.86 (ddt, <i>J</i> = 9.1, 6.2, 3.6 Hz, 1H), 3.77 – 3.60 (m, 2H), 2.15 – 2.02 (m, 1H), 1.78 (m, 1H), 1.64 (m, 2H), 1.42 (m, 2H); LCMS: 419
28		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.08 (br s, 1H), 9.30 (m, 1H), 8.28 (s, 1H), 7.31 (m, 1H), 7.20 (m, 2H), 5.36 (d, <i>J</i> = 2.4 Hz, 1H), 4.90 – 4.76 (m, 1H), 4.40 (q, <i>J</i> = 7.5 Hz, 1H), 3.94 (dtd, <i>J</i> = 11.7, 5.6, 5.1, 3.4 Hz, 1H), 3.72 (ddd, <i>J</i> = 22.6, 7.5, 4.5 Hz, 2H), 3.60 (br s, 1H), 3.07 (s, 3H), 2.32 – 2.19 (m, 1H), 1.94 (m, 1H), 1.58 (m, 1H), 1.41 – 1.26 (m, 1H); LCMS: 419

FIGURE 1





29		¹H-NMR (400 MHz, CDCl₃) δ ppm 10.62-10.49 (m, 1H), 8.39 (d, 1H, J = 5.6 Hz), 7.10-6.97 (m, 3H), 5.47-5.34 (m, 1H), 4.83 (s, 1H), 4.26-4.16 (m, 2H), 4.02-3.89 (m, 2H), 3.60-3.45 (m, 3.5H), 3.13 (s, 1.5H), 2.17-2.13 (m, 1H), 1.81 (br.s, 1H).; LCMS: 419.1
30		¹H-NMR (400 MHz, CDCl₃) δ ppm 13.68 (br.s, 1H), 9.40 (d, 1H, J = 6.8 Hz), 8.33 (s, 1H), 7.61 (d, 1H, J = 9.2 Hz), 7.03-6.88 (m, 3H), 5.40 (t, 1H, J = 7.6 Hz), 4.13-4.09 (m, 2H), 3.95-3.92 (m, 1H), 3.78-3.76 (m, 1H), 3.42-3.34 (m, 2H), 2.04 (dd, 1H, J = 10.8, 2.0 Hz), 1.73-1.70 (m, 1H), 1.57 (d, 3H, J = 7.2 Hz).; LCMS: 419.1
31		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 7.23-7.21 (m, 1H), 7.13-7.10 (m, 1H), 7.04-7.01 (m, 1H), 5.58-5.52 (m, 1H), 4.03-3.94 (m, 3H), 3.62-3.61 (m, 1H), 3.52-3.49 (m, 1H), 3.27-3.22 (m, 1H), 2.17-2.13 (m, 1H), 1.86-1.80 (m, 1H), 1.60 (d, 3H, J = 7.2 Hz).; LCMS: 419.1
32		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.21 (s, 1H), 7.14-7.10 (m, 2H), 7.03-7.01 (m, 1H), 4.66 (s, 2H), 4.51-4.45 (m, 1H), 4.00 (s, 1H), 3.70-3.63 (m, 1H), 2.09-2.07 (m, 1H), 1.85-1.78 (m, 2H), 1.61-1.56 (m, 2H), 1.45-1.36 (m, 1H).; LCMS: 419.1

FIGURE 1





33		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (d, 1H, J = 5.6 Hz), 6.99-6.96 (m, 2H), 6.92-6.88 (m, 1H), 5.43 (s, 1H), 4.90 (s, 1H), 4.43-4.37 (m, 1H), 4.15-4.13 (m, 1H), 4.05-4.01 (m, 1H), 3.64 (s, 1.5H), 3.16 (s, 1.5H), 2.46-2.43 (m, 1H), 2.19-2.15 (m, 1H), 1.82-1.65 (m, 2H).; LCMS: 419.1
34		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.21 (br.s, 1H), 7.11-7.09 (m, 1H), 7.01 (br.s, 1H), 5.54 (d, 1H, J = 7.2 Hz), 4.39-4.32 (m, 1H), 4.10 (br.s, 1H), 3.99-3.96 (m, 1H), 2.43-2.40 (m, 1H), 2.14-2.12 (m, 1H), 1.78-1.74 (m, 1H), 1.69 (br.s, 1H), 1.59 (d, 3H, J = 6.8 Hz).; LCMS: 419.2
35		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.21 (br.s, 1H), 7.13-7.09 (m, 1H), 7.03-7.01 (m, 1H), 5.55-5.51 (m, 1H), 4.37-4.31 (m, 1H), 4.11 (s, 1H), 4.02-3.99 (m, 1H), 2.42-2.38 (m, 1H), 2.15-2.11 (m, 1H), 1.78-1.76 (m, 1H), 1.67-1.64 (m, 1H), 1.59 (d, 3H, J = 6.8 Hz).; LCMS: 419.2
36		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.10 (s, 1H), 9.49 (m, 1H), 9.37 (d, J = 7.3 Hz, 1H), 8.27 (s, 1H), 7.53 (dd, J = 8.8, 5.1 Hz, 1H), 7.18 (td, J = 8.4, 2.9 Hz, 1H), 7.12 (dd, J = 9.4, 3.5 Hz, 1H), 5.68 (d, J = 7.4 Hz, 1H), 4.87 (m, 1H), 4.56 (m, 1H), 4.48 (m, 1H), 4.39 (p, J = 7.6 Hz, 2H), 3.90 (m, 1H), 3.70 (m, 1H), 2.22 (m, 1H), 1.91 (m, 1H), 1.60 – 1.47 (m, 1H), 1.31 (m, 1H); LCMS: 421

FIGURE 1





37		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 13.80 (s, 1H), 9.43 (t, <i>J</i> = 6.2 Hz, 1H), 9.31 (d, <i>J</i> = 7.8 Hz, 1H), 8.22 (s, 1H), 7.47 (dd, <i>J</i> = 8.8, 5.1 Hz, 1H), 7.13 (td, <i>J</i> = 8.4, 3.1 Hz, 1H), 7.06 (dd, <i>J</i> = 9.6, 3.1 Hz, 1H), 5.06 (d, <i>J</i> = 4.8 Hz, 1H), 4.51 (d, <i>J</i> = 6.2 Hz, 2H), 3.99 (qd, <i>J</i> = 7.7, 3.8 Hz, 1H), 3.91 (dd, <i>J</i> = 11.1, 4.0 Hz, 1H), 3.72 (dt, <i>J</i> = 11.3, 4.6 Hz, 1H), 3.59 (tt, <i>J</i> = 8.5, 4.5 Hz, 1H), 3.38 (ddd, <i>J</i> = 11.7, 9.0, 3.1 Hz, 1H), 3.12 (dd, <i>J</i> = 11.1, 7.7 Hz, 1H), 1.84 (ddt, <i>J</i> = 9.2, 7.6, 3.6 Hz, 1H), 1.43 (dtd, <i>J</i> = 13.0, 8.7, 4.0 Hz, 1H); LCMS: 421
38		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.24 (s, 1H), 7.23 (s, 1H), 7.08 (d, 2H, <i>J</i> = 8.8 Hz), 4.60 (s, 2H), 4.51-4.46 (m, 1H), 4.13 (br.s, 1H), 3.94-3.90 (m, 1H), 2.46-2.40 (m, 1H), 2.11-2.06 (m, 1H), 1.78-1.74 (m, 1H), 1.62-1.58 (m, 1H); LCMS: 421.2
39		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 9.19 (d, <i>J</i> = 145.0 Hz, 1H), 8.08 (d, <i>J</i> = 31.1 Hz, 1H), 7.37 – 7.02 (m, 3H), 5.58 (s, 1H), 5.18 (s, 1H), 4.78 (s, 1H), 4.08 – 3.90 (m, 2H), 3.78 (s, 1H), 3.63 (s, 1H), 3.43 (d, <i>J</i> = 11.5 Hz, 1H), 3.17 (dd, <i>J</i> = 11.2, 7.4 Hz, 1H), 1.92 (s, 2H), 1.48 (s, 1H); LCMS: 422
40		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.19 (s, 1H), 9.51 (t, <i>J</i> = 6.3 Hz, 2H), 7.57 – 7.35 (m, 1H), 7.11 – 6.94 (m, 1H), 4.60 (d, <i>J</i> = 6.1 Hz, 2H), 4.04 (dt, <i>J</i> = 11.7, 5.8 Hz, 1H), 3.97 (dd, <i>J</i> = 11.1, 4.0 Hz, 1H), 3.78 (dd, <i>J</i> = 11.3, 5.1 Hz, 1H), 3.60 (tt, <i>J</i> = 6.6, 3.4 Hz, 1H), 3.48 – 3.40 (m, 1H), 3.19 (dd, <i>J</i> = 11.1, 7.7 Hz, 1H), 3.12 (qd, <i>J</i> = 7.3, 4.2 Hz, 2H), 1.91 (d, <i>J</i> = 13.1 Hz, 1H), 1.50 (dq, <i>J</i> = 8.7, 4.6 Hz, 1H); LCMS: 423

FIGURE 1





41		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37 (s, 1H), 7.09-6.98 (m, 2H), 4.71 (s, 2H), 4.41-4.34 (m, 1H), 4.13 (br.s, 1H), 4.04-4.02 (m, 1H), 2.45-2.43 (m, 1H), 2.17-2.13 (m, 1H), 1.82-1.69 (m, 2H).; LCMS: 423.1
42		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33 (s, 1H), 7.14-6.99 (m, 3H), 4.65 (s, 2H), 4.49-4.43 (m, 1H), 4.35-4.32 (m, 1H), 2.85-2.80 (m, 1H), 2.67-2.58 (m, 1H), 2.26-2.16 (m, 2H).; LCMS: 425.1
43		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.28 (s, 1H), 7.13-6.99 (m, 3H), 4.65 (s, 2H), 4.52-4.46 (m, 1H), 4.33-4.28 (m, 1H), 2.87-2.79 (m, 1H), 2.64-2.59 (m, 1H), 2.22-2.14 (m, 2H).; LCMS: 425.1
44		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33 (d, 1H, J = 6.0 Hz), 7.60 (d, 1H, J = 9.2 Hz), 7.52-7.49 (m, 2H), 5.43 (d, 1H, J = 3.6 Hz), 4.90 (d, 1H, J = 3.2 Hz), 4.43-4.38 (m, 1H), 4.13 (br.s, 1H), 4.03-3.96 (m, 1H), 3.65 (s, 1.5H), 3.17 (s, 1.5H), 2.48-2.44 (m, 1H), 2.17-2.13 (m, 1H), 1.83-1.66 (m, 2H).; LCMS: 426.1

FIGURE 1





45		¹H-NMR (400 MHz, CDCl₃) δ ppm 11.58-11.21 (m, 1H), 8.47 (br.s, 0.5H), 6.98-6.55 (m, 3.5H), 6.33 (d, 0.5H, J = 8.0 Hz), 5.51-5.49 (m, 0.5H), 4.38-3.83 (m, 4H), 2.37-2.32 (m, 2H), 2.05-1.64 (m, 8H).; LCMS: 429.1
46		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.50 (br.s, 1H), 8.40 (br.s, 1H), 8.38 (s, 1H), 7.72 (d, 1H, J = 9.2 Hz), 5.54-5.41 (m, 2H), 4.97-4.92 (m, 1H), 4.59-4.45 ((m, 1H), 4.09-4.04 (m, 2H), 2.91-2.84 (m, 1H), 2.29-2.26 (m, 2H), 1.79-1.66 (m, 5H).; LCMS: 430.1
47		¹H-NMR (400 MHz, CD₃OD) δ ppm 9.57 (s, 1H), 8.34 (s, 1H), 7.38-7.37 (m, 1H), 7.06-6.96 (m, 2H), 4.13-4.09 (m, 1H), 4.00-3.96 (m, 2H), 3.87-3.76 (m, 1H), 3.58-3.47(m, 1H), 3.45-3.44 (m, 1H), 2.11-2.07 (m, 1H), 1.75-1.66 (m, 1H), 1.37 (s, 4H).; LCMS: 431.1
48		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (d, 1H, J = 4.8 Hz), 6.73 (d, 1H, J = 12.0 Hz), 6.69-6.63 (m, 2H), 5.40 (s, 1H), 4.82 (s, 1H), 4.39-4.36 (m, 1H), 4.15-4.12 (m, 1H), 4.06-4.02 (m, 1H), 3.78 (d, 3H, J = 6.4 Hz), 3.62 (s, 1.5H), 3.14 (s, 1.5H), 2.47-2.44 (m, 1H), 2.18-2.15 (m, 1H), 1.82-1.70 (m, 2H).; LCMS: 431.1

FIGURE 1





49		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35 (d, 1H, J = 3.6 Hz), 7.01-6.93 (m, 1H), 5.44-5.33 (m, 1H), 4.83 (s, 1H), 4.40-4.35 (m, 1H), 4.14-4.12 (m, 1H), 4.05-4.01 (m, 1H), 3.86 (s, 1.5H), 3.75 (s, 1.5H), 3.63 (s, 1.5H), 3.12 (s, 1.5H), 2.46-2.43 (m, 1H), 2.17-2.15 (m, 1H), 1.82-1.63 (m, 2H).; LCMS: 431.1
50		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.60 (br.s, 1H), 8.50 (br.s, 1H), 8.43 (s, 1H), 7.89 (d, 1H, J = 9.2 Hz), 5.57-5.35 (m, 2H), 4.58-4.38 (m, 3H), 4.25-4.13 (m, 2H), 4.00-3.93 (m, 1H), 3.84-3.71 (m, 2H), 2.90-2.82 (m, 1H), 2.34-2.21 (m, 1H).; LCMS: 432.1
51		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (s, 1H), 7.22-7.19 (m, 1H), 7.12-7.11 (m, 1H), 7.02-7.01 (m, 1H), 5.52-5.50 (m, 1H), 5.04-5.02 (m, 1H), 4.78-4.76 (m, 2H), 4.30-3.89 (m, 1H), 3.80-3.75 (m, 2H), 3.36-3.34 (m, 2H), 1.59 (d, 3H, J = 6.8 Hz), 1.50-1.34 (m, 3H).; LCMS: 432.2
52		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35 (d, 1H, J = 6.0 Hz), 7.94 (dd, 1H, J = 9.2, 2.4 Hz), 7.47-7.41 (m, 1H), 5.31 (s, 1H), 4.78 (s, 1H), 4.41-4.35 (m, 1H), 4.14-4.11 (m, 1H), 4.04-4.02 (m, 1H), 3.97 (s, 1.5H), 3.89 (s, 1.5H), 3.70 (s, 1.5H), 3.17 (s, 1.5H), 2.47-2.40 (m, 1H), 2.18-2.14 (m, 1H), 1.81-1.64 (m, 2H).; LCMS: 432.2

FIGURE 1





53		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.38 (d, 1H, <i>J</i> = 5.2 Hz), 7.20-7.05 (m, 3H), 4.87 (s, 2H), 4.17-4.11 (m, 2H), 4.00-3.97 (m, 3H), 3.64-3.44 (m, 3H), 2.13-2.06 (m, 1H), 1.76-1.70 (m, 1H), 1.39-1.18 (m, 3H).; LCMS: 433.1
54		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.19 (s, 1H), 9.51 (t, <i>J</i> = 6.3 Hz, 2H), 7.57 – 7.35 (m, 1H), 7.11 – 6.94 (m, 1H), 4.60 (d, <i>J</i> = 6.1 Hz, 2H), 4.04 (dt, <i>J</i> = 11.7, 5.8 Hz, 1H), 3.97 (dd, <i>J</i> = 11.1, 4.0 Hz, 1H), 3.78 (dd, <i>J</i> = 11.3, 5.1 Hz, 1H), 3.60 (tt, <i>J</i> = 6.6, 3.4 Hz, 1H), 3.48 – 3.40 (m, 1H), 3.19 (dd, <i>J</i> = 11.1, 7.7 Hz, 1H), 3.12 (qd, <i>J</i> = 7.3, 4.2 Hz, 2H), 1.91 (d, <i>J</i> = 13.1 Hz, 1H), 1.50 (dq, <i>J</i> = 8.7, 4.6 Hz, 1H); LCMS: 435
55		LCMS: 435
56		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.23-7.21 (m, 1H), 7.13-7.12 (m, 1H), 7.11-7.03 (m, 1H), 5.57-5.54 (m, 1H), 4.14-4.01 (m, 1H), 4.00-3.92 (m, 4H), 3.91-3.90 (m, 1H), 3.61-3.44 (m, 2H), 2.05-2.02 (m, 1H), 1.72-1.68 (m, 1H).; LCMS: 435.1

FIGURE 1





57		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (d, 1H, J = 14.8 Hz), 7.50-7.45 (m, 1H), 7.09-7.04 (m, 2H), 5.46 (s, 1H), 4.96 (s, 1H), 4.40-4.35 (m, 1H), 4.16-3.97 (m, 2H), 3.67 (s, 1.5H), 3.22 (s, 1.5H), 2.47-2.45 (m, 1H), 2.20-2.16 (m, 1H), 1.87-1.77 (m, 2H).; LCMS: 435.1
58		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (d, 1H, J = 5.2 Hz), 7.23 (d, 1H, J = 12.0 Hz), 7.16-7.09 (m, 2H), 5.42 (s, 1H), 4.85 (s, 1H), 4.40-4.36 (m, 1H), 4.15-4.13 (m, 1H), 4.06-4.02 (m, 1H), 3.64 (s, 1.5H), 3.16 (s, 1.5H), 2.47-2.44 (m, 1H), 2.19-2.15 (m, 1H), 1.84-1.72 (m, 2H).; LCMS: 435.1
59		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.24 (s, 1H), 6.94-6.89 (m, 2H), 4.64 (s, 2H), 4.52-4.48 (m, 1H), 4.14-4.12 (m, 1H), 3.95 (s, 3H), 3.93-3.92 (m, 1H), 2.43-2.40 (m, 1H), 2.10-2.08 (m, 1H), 1.78-1.75 (m, 1H), 1.61-1.57 (m, 1H).; LCMS: 435.2
60		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26-8.22 (m, 2H), 7.64-7.59 (m, 1H), 5.34 (s, 1H), 4.89 (s, 1H), 4.51-4.47 (m, 1H), 4.15-4.09 (m, 1H), 3.95-3.86 (m, 1H), 3.68 (s, 1.2H), 3.22 (s, 1.8H), 2.46-2.40 (m, 1H), 2.11-2.09 (m, 1H), 1.80-1.49 (m, 2H).; LCMS: 436.2

FIGURE 1





61		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.30 (s, 1H), 7.25-7.15 (m, 2H), 7.08-7.07 (m, 1H), 5.55-5.52 (m, 1H), 5.15 (d, 1H, J = 53.6 Hz), 4.60-4.55 (m, 1H), 4.21-4.18 (m, 1H), 3.95-3.88 (m, 2H), 2.62-2.51 (m, 2H), 2.00-1.89 (m, 2H).; LCMS: 436.4
62		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.30 (s, 1H), 7.25-7.15 (m, 2H), 7.08-7.07 (m, 1H), 5.55-5.52 (m, 1H), 5.16 (d, 1H, J = 53.6 Hz), 4.60-4.54 (m, 1H), 4.18-4.14 (m, 1H), 3.94-3.89 (m, 2H), 2.60-2.51 (m, 2H), 1.99-1.90 (m, 2H).; LCMS: 436.4
63		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.09 (br s, 1H), 9.25 (Abqd, 1H), 8.33 (d, J = 8.2 Hz, 1H), 7.67 – 7.47 (m, 1H), 7.14 (m, 1H), 5.46 (m, 1H), 4.96 (m, 1H), 4.59 (dd, J = 9.6, 4.0 Hz, 1H), 4.47 (q, J = 7.8 Hz, 1H), 4.00 (m, 1H), 3.84 – 3.71 (m, 1H), 3.58 (s, 1H), 3.40 (s, 3H), 2.42 – 2.26 (m, 1H), 2.08 – 1.92 (m, 1H), 1.72 – 1.57 (m, 1H), 1.37 (m, 1H); LCMS: 437
64		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.12 (s, 1H), 9.58 – 9.35 (m, 2H), 8.29 (s, 1H), 7.37 – 7.12 (m, 3H), 5.72 (d, J = 6.2 Hz, 1H), 4.67 – 4.49 (m, 2H), 4.29 (s, 1H), 3.76 (m, 1H), 2.04 (m, 2H), 1.99 – 1.79 (m, 1H), 1.71 (s, 1H), 1.47 (m, 2H); LCMS: 439

FIGURE 1





65		LCMS: 439
66		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.17 (s, 1H), 7.38-7.34 (m, 3H), 7.10-7.06 (m, 1H), 6.01-5.95 (m, 1H), 4.17-4.12 (m, 1H), 4.07-4.03 (m, 1H), 2.19-2.17 (m, 1H), 1.96-1.94 (m, 1H), 1.76-1.73 (m, 2H), 1.60-1.57 (m, 2H); LCMS: 439.2
67		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.15 (s, 1H), 7.38-7.35 (m, 3H), 7.10-7.05 (m, 1H), 6.02-5.96 (m, 1H), 4.24-4.14 (m, 1H), 4.03-3.99 (m, 1H), 2.24-2.19 (m, 1H), 1.94-1.93 (m, 1H), 1.77-1.74 (m, 2H), 1.60-1.55 (m, 2H); LCMS: 439.2
68		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.09 (s, 1H), 9.01 (d, <i>J</i> = 7.1 Hz, 1H), 8.19 (s, 1H), 7.31 (m, 1H), 7.16 – 7.10 (m, 2H), 7.00 (m, 1H), 5.43-5.30 (m, 2H), 4.79 (d, <i>J</i> = 5.6 Hz, 1H), 4.60 (m, 1H), 4.48 – 4.33 (m, 2H), 4.33 – 4.22 (m, 1H), 3.78 (m, 1H), 3.56 (m, 1H), 2.77 – 2.59 (m, 1H), 2.10 (m, 2H), 1.99 – 1.76 (m, 2H), 1.46 (m, 1H), 1.27 – 1.15 (m, 1H); LCMS: 445

FIGURE 1





69		¹H-NMR (400 MHz, CDCl₃) δ ppm 11.87-11.48 (m, 1H), 8.50-8.46 (m, 1H), 7.06-6.65 (m, 3H), 6.40-6.39 (m, 0.5H), 5.62-5.59 (m, 0.5H), 4.46-3.95 (m, 6H), 3.55-3.51 (m, 2H), 2.42-2.11 (m, 1H), 2.04-1.76 (m, 5H).; LCMS: 445.1
70		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35, 8.28 (s, 1H), 7.14-6.80 (m, 3H), 6.43 (d, 0.5H, J = 7.6 Hz), 5.60 (dd, 0.5H, J = 7.6, 4.0 Hz), 4.52-4.31 (m, 2H), 4.05-3.89 (m, 3H), 2.48-2.41 (m, 2H), 2.12-2.06 (m, 4H), 1.94-1.61 (m, 2H).; LCMS: 445.1
71		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33, 8.27 (s, 1H), 7.14-6.92 (m, 3H), 6.43 (d, 0.5H, J = 8.0 Hz), 5.58 (dd, 0.5H, J = 8.0, 4.4 Hz), 4.39-4.31 (m, 2H), 4.15-4.08 (m, 2H), 3.94-3.91 (m, 1H), 2.48-2.43 (m, 2H), 2.12-2.06 (m, 3H), 1.76-1.51 (m, 3H).; LCMS: 445.1
72		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.42 (br.s, 1H), 8.31 (br.s, 1H), 8.26 (s, 1H), 7.63-7.61 (m, 1H), 5.46-5.26 (m, 2H), 4.50-4.37 (m, 1H), 4.26-4.21 (m, 1H), 3.97 (br.s, 2H), 3.80-3.77 (m, 1H), 2.80-2.76 (m, 1H), 2.33-2.30 (m, 1H), 2.29-2.27 (m, 1H), 2.07-2.00 (m, 1H), 1.69-1.67 (m, 1H), 1.52-1.51 (m, 1H).; LCMS: 446.1

FIGURE 1





73		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.52 (s, 1H), 8.41 (s, 1H), 8.35-8.29 (m, 1H), 7.76 (d, 1H, J = 9.2 Hz), 5.57-5.36 (m, 2H), 4.96-4.94 (m, 1H), 4.60-4.51 (m, 1H), 3.98-3.89 (m, 3H), 3.50-3.47 (m, 2H), 3.24-3.21 (m, 1H), 2.88-2.86 (m, 1H), 2.18-2.09 (m, 2H), 1.75-1.71 (m, 1H).; LCMS: 446.1
74		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.60 (br.s, 2H), 8.35 (s, 1H), 7.75 (d, 1H, J = 9.2 Hz), 5.57-5.35 (m, 2H), 4.59-4.45 (m, 1H), 4.09-4.05 (m, 1H), 3.95-3.91 (m, 2H), 3.65-3.75 (m, 1H), 3.53-3.50 (m, 1H), 3.40-3.35 (m, 1H), 2.88-2.86 (m, 1H), 2.29-2.27 (m, 2H), 1.97-1.93 (m, 1H), 1.66-1.63 (m, 1H).; LCMS: 446.2
75		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.89-8.87 (m, 0.5H), 8.52-8.43 (m, 1H), 8.39 (s, 1H), 8.13-8.10 (m, 0.5H), 7.77 (d, 1H, J = 9.2 Hz), 5.55-5.40 (m, 2H), 5.16-5.01 (m, 2H), 4.60-4.37 (m, 2H), 4.17-4.15 (m, 1H), 2.89-2.86 (m, 1H), 2.66-2.56 (m, 1H), 2.44-2.42 (m, 1H), 2.30-2.20 (m, 1H), 1.98-1.88 (m, 2H).; LCMS: 448.1
76		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.87-8.86 (m, 0.5H), 8.67-8.50 (m, 1H), 8.46 (s, 1H), 8.12-8.10 (m, 0.5H), 7.75-7.73 (m, 1H), 5.54-5.24 (m, 2H), 5.16-5.03 (m, 2H), 4.58-4.37 (m, 2H), 4.11-4.07 (m, 1H), 2.87-2.85 (m, 1H), 2.55-2.47 (m, 2H), 2.26-2.01 (m, 1H), 1.95-1.84 (m, 2H).; LCMS: 448.1

FIGURE 1





77		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.43 (d, 1H, J = 5.2 Hz), 7.06-6.99 (m, 1H), 6.96-6.87 (m, 1H), 5.57-5.48 (m, 1H), 5.00 (s, 1H), 4.45-4.41 (m, 1H), 4.20-4.19 (m, 1H), 4.12-4.07 (m, 1H), 3.98 (s, 1.5H), 3.89 (s, 1.5H), 3.71 (s, 1.5H), 3.22 (s, 1.5H), 2.53-2.49 (m, 1H), 2.25-2.21 (m, 1H), 1.90-1.79 (m, 2H).; LCMS: 449.1
78		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.14-7.00 (m, 3H), 5.59-5.55 (m, 1H), 5.48-5.32 (m, 1H), 4.46-4.33 (m, 3H), 4.23-4.12 (m, 2H), 3.97-3.94 (m, 1H), 3.69-3.68 (m, 2H), 2.85-2.75 (m, 1H), 2.16-2.09 (m, 1H).; LCMS: 449.1
79		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.41 (s, 1H), 8.35 (s, 1H), 7.20-7.16 (m, 1H), 7.07-7.01 (m, 2H), 5.76-5.73 (m, 1H), 4.84-4.75 (m, 1H), 4.25-4.08 (m, 3H), 3.64-3.61 (m, 1H), 2.81-2.78 (m, 1H), 2.41-2.38 (m, 2H), 2.00-1.68 (m, 5H).; LCMS: 453.5
80		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.35 (s, 1H), 9.60 (q, J = 7.8, 7.0 Hz, 1H), 8.33 (s, 1H), 7.84 (dd, J = 8.7, 5.4 Hz, 1H), 7.39 – 7.16 (m, 2H), 4.72 (d, J = 6.1 Hz, 2H), 4.04 (qd, J = 7.9, 4.0 Hz, 1H), 3.96 (dd, J = 11.1, 4.0 Hz, 1H), 3.78 (dt, J = 11.5, 4.5 Hz, 1H), 3.65 (td, J = 8.2, 4.3 Hz, 1H), 3.48 – 3.38 (m, 1H), 3.23 – 3.15 (m, 1H), 1.89 (dq, J = 12.7, 4.1 Hz, 1H), 1.49 (dtd, J = 13.2, 8.9, 4.0 Hz, 1H), 1.25 (dt, J = 13.3, 7.9 Hz, 1H); LCMS: 455

FIGURE 1





81		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37 (s, 1H), 7.80-7.71 (m, 1H), 7.29-7.27 (m, 1H), 7.21-7.17 (m, 1H), 4.74-4.67 (m, 2H), 4.44-4.30 (m, 1H), 4.11 (s, 1H), 4.02-4.01 (m, 1H), 2.50-2.40 (m, 1H), 2.15-2.12 (m, 1H), 1.78-1.77 (m, 1H), 1.70-1.67 (m, 1H).; LCMS: 455.1
82		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.39 (s, 1H), 8.33 (s, 1H), 7.22-6.90 (m, 3H), 5.74 (br.s, 1H), 4.83-4.73 (m, 1H), 4.51-4.42 (m, 2H), 4.26-4.17 (m, 2H), 3.98-3.95 (m, 1H), 3.82-3.71 (m, 3H), 2.87-2.75 (m, 1H), 2.53-2.50 (m, 1H).; LCMS: 455.4
83		¹H NMR (500 MHz, DMSO-<i>d</i>₆) δ 14.12 (s, 1H), 9.51 (t, <i>J</i> = 6.2 Hz, 1H), 9.33 (d, <i>J</i> = 7.7 Hz, 1H), 8.28 (s, 1H), 7.64 (dd, <i>J</i> = 8.2, 3.0 Hz, 1H), 7.15 (dd, <i>J</i> = 9.2, 3.1 Hz, 1H), 5.09 (d, <i>J</i> = 4.9 Hz, 1H), 4.60 (d, <i>J</i> = 6.1 Hz, 2H), 4.05 (dd, <i>J</i> = 7.7, 4.1 Hz, 1H), 3.96 (dd, <i>J</i> = 11.2, 4.1 Hz, 1H), 3.82 – 3.75 (m, 1H), 3.65 (d, <i>J</i> = 4.9 Hz, 1H), 3.43 (t, <i>J</i> = 9.4 Hz, 1H), 3.17 (dd, <i>J</i> = 11.2, 7.6 Hz, 1H), 1.89 (d, <i>J</i> = 12.9 Hz, 1H), 1.52 – 1.44 (m, 1H); LCMS: 456
84		¹H NMR (400 MHz, DMSO-<i>d</i>₆) δ 14.11 (s, 1H), 9.60 – 9.39 (m, 2H), 8.28 (s, 1H), 7.32 – 7.24 (m, 1H), 7.17 (m, 1H), 5.21 (d, <i>J</i> = 5.2 Hz, 1H), 4.56 (d, <i>J</i> = 6.1 Hz, 2H), 4.29 (m, 1H), 3.96 (d, <i>J</i> = 6.7 Hz, 1H), 3.29 – 3.18 (m, 1H), 2.97 (s, 3H), 2.84 (s, 3H), 2.30 (m, 1H), 2.14 (m, 1H), 1.78 – 1.61 (m, 2H); LCMS: 460

FIGURE 1





85		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.51 (s, 1H), 8.41 (s, 1H), 8.32 (s, 1H), 7.76 (d, 1H, J = 9.2 Hz), 5.56-5.40 (m, 2H), 4.87-4.82 (m, 1H), 4.59-4.50 (m, 1H), 4.11-3.97 (m, 2H), 3.48-3.45 (m, 1H), 2.88-2.86 (m, 1H), 2.27-1.80 (m, 2H), 1.78-1.67 (m, 2H), 1.59-1.46 (m, 3H).; LCMS: 460.1
86		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.25-8.19 (m, 1H), 7.14-6.85 (m, 3H), 5.63-5.59 (m, 1H), 5.42 (d, 1H, J = 52.4 Hz), 4.83 (br.s, 1H), 4.48-4.31 (m, 3H), 4.12-3.99 (m, 1H), 2.88-2.80 (m, 1H), 2.42-2.39 (m, 1H), 2.25-2.20 (m, 2H), 1.92-1.89 (m, 1H), 1.69-1.64 (m, 1H).; LCMS: 462.4
87		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26-8.19 (m, 1H), 7.15-6.86 (m, 3H), 5.63-5.59 (m, 1H), 5.42 (d, 1H, J = 52.0 Hz), 4.85-4.80 (m, 1H), 4.47-4.31 (m, 3H), 4.11-4.03 (m, 1H), 2.83-2.78 (m, 1H), 2.49-2.43 (m, 1H), 2.23-2.20 (m, 2H), 1.87-1.83 (m, 1H), 1.70-1.65 (m, 1H).; LCMS: 462.4
88		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.25 (s, 1H), 7.17-7.03 (m, 3H), 5.62 (t, 1H, J = 8.8 Hz), 5.43 (d, 1H, J = 52.0 Hz), 4.86-4.81 (m, 1H), 4.50-4.40 (m, 2H), 4.12-4.08 (m, 1H), 3.93-3.84 (m, 1H), 2.89-2.79 (m, 1H), 2.41-2.38 (m, 1H), 2.12-2.05 (m, 2H), 1.76-1.75 (m, 1H), 1.56-1.55 (m, 1H); LCMS: 463

FIGURE 1





89		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.02 (s, 1H), 9.23 (d, <i>J</i> = 7.0 Hz, 1H), 8.24 (s, 1H), 7.41 (qd, <i>J</i> = 9.4, 4.9 Hz, 1H), 7.12 (ddt, <i>J</i> = 11.4, 7.6, 2.0 Hz, 1H), 5.48 (m, 1H), 4.90 (m, 1H), 4.46 (m, 1H), 4.36 (m, 2H), 4.09 (m, 1H), 3.89 – 3.83 (m, 1H), 3.59 (m, 1H), 2.49 – 2.38 (m, 1H), 2.30 – 2.10 (m, 2H), 1.97 – 1.79 (m, 3H), 1.54 (m, 2H), 1.37 – 1.24 (m, 2H); LCMS: 463
90		LCMS: 463
91		LCMS: 463
92		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.14-6.86 (m, 3H), 5.63-5.59 (m, 1H), 5.51-5.35 (m, 1H), 4.50-4.41 (m, 1H), 4.10-4.06 (m, 1H), 3.98-3.92 (m, 2H), 3.75-3.70 (m, 1H), 3.52-3.41 (m, 1H), 3.39-3.36 (m, 1H), 2.84-2.81 (m, 1H), 2.25-2.00 (m, 1H), 2.00-1.96 (m, 1H), 1.67-1.64 (m, 1H), 1.38-1.28 (m, 1H).; LCMS: 463.1

FIGURE 1





93		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.14-8.04 (m, 1H), 7.02-6.74 (m, 3H), 5.73-5.48 (m, 1H), 5.37-5.24 (m, 1H), 4.36-4.10 (m, 2H), 3.87-3.76 (m, 2H), 3.44-3.38 (m, 2H), 3.18-3.10 (m, 1H), 2.76-2.72 (m, 1H), 2.10-1.88 (m, 2H), 1.52 (br.s, 1H), 1.33-1.17 (m, 1H).; LCMS: 463.1
94		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.53 (s, 1H), 8.46-8.43 (m, 1H), 8.29-8.24 (m, 1H), 7.81-7.66 (m, 1H), 5.81-5.78 (m, 1H), 4.89-4.85 (m, 1H), 4.57-4.52 (m, 2H), 4.50-4.48 (m, 1H), 4.13-4.11 (m, 1H), 3.88-3.85 (m, 1H), 3.12-3.11 (m, 1H), 2.62-2.59 (m, 1H), 2.44-2.42 (m, 1H), 2.08-2.05 (m, 1H), 1.82-1.77 (m, 1H), 1.55-1.53 (m, 1H).; LCMS: 464.1
95		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.15-8.09 (m, 1H), 7.03-6.74 (m, 3H), 5.52-5.47 (m, 1H), 5.31 (d, 1H, J = 52.0 Hz), 5.06-4.92 (m, 2H), 4.44-4.38 (m, 2H), 3.99-3.96 (m, 1H), 2.72-2.70 (m, 1H), 2.45-2.32 (m, 2H), 1.84-1.75 (m, 3H).; LCMS: 465.1
96		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.22-8.10 (m, 1H), 7.05-6.75 (m, 3H), 5.51-5.47 (m, 1H), 5.31 (d, 1H, J = 52.0 Hz), 4.99 (d, 1H, J = 54.0 Hz), 4.44-4.29 (m, 2H), 4.02-3.99 (m, 1H), 2.90-2.72 (m, 1H), 2.47-2.35 (m, 2H), 2.12-1.64 (m, 3H).; LCMS: 465.1

FIGURE 1





97		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.28 (s, 1H), 7.15-7.11 (m, 2H), 7.04-7.02 (m, 1H), 4.66 (s, 2H), 4.51-4.45 (m, 1H), 4.28-4.23 (m, 1H), 3.84-3.80 (m, 1H), 2.96 (s, 3H), 2.75-2.70 (m, 1H), 2.53-2.50 (m, 1H), 2.15-2.06 (m, 2H).; LCMS: 467.1
98		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.28 (s, 1H), 7.15-7.11 (m, 2H), 7.09-7.02 (m, 1H), 4.66 (s, 2H), 4.51-4.45 (m, 1H), 4.28-4.25 (m, 1H), 3.84-3.80 (m, 1H), 2.96 (s, 3H), 2.75-2.70 (m, 1H), 2.54-2.52 (m, 1H), 2.13-2.06 (m, 2H).; LCMS: 467.1
99		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40, 8.36 (d, 1H, J = 11.6 Hz), 7.85-7.80 (m, 1H), 7.25-7.17 (m, 2H), 5.60 (s, 1H), 5.57 (s, 1H), 4.40-4.35 (m, 1H), 4.16-3.98 (m, 2H), 3.69 (s, 1.5H), 3.23 (s, 1.5H), 2.49-2.43 (m, 1H), 2.21-2.20 (m, 1H), 1.85-1.64 (m, 2H).; LCMS: 469.1
100		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.28-8.22 (m, 1H), 7.19-6.90 (m, 3H), 5.78-5.76 (m, 1H), 4.85-4.80 (m, 1H), 4.71-4.69 (m, 1H), 4.19-4.17 (m, 2H), 4.09-4.08 (m, 2H), 3.75-3.74 (m, 1H), 3.59-3.48 (m, 2H), 2.86-2.76 (m, 1H), 2.51-2.35 (m, 1H), 2.13-1.96 (m, 1H), 1.71-1.60 (m, 1H).; LCMS: 469.4

FIGURE 1





101		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.35 (br.s, 1H), 7.20 (d, 1H, J = 6.8 Hz), 7.13 (t, 1H, J = 8.0 Hz), 4.71 (s, 2H), 4.41-4.35 (m, 1H), 4.12 (br.s, 1H), 4.01-3.99 (m, 1H), 2.45-2.40 (m, 1H), 2.15-2.11 (m, 1H), 1.80-1.68 (m, 2H).; LCMS: 471.1
102		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 11.11 (d, 1H, J = 6.0 Hz), 8.67 (s, 1H), 7.84-7.82 (m, 1H), 7.04-6.92 (m, 3H), 4.64 (d, 2H, J = 6.4 Hz), 4.41 (br.s, 1H), 4.29 (br.s, 1H), 3.42-3.38 (m, 2H), 3.25-3.22 (m, 2H), 3.09-3.07 (m, 1H), 2.79-2.77 (m, 1H).; LCMS: 472.2
103		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.32 (s, 1H), 10.13 (d, J = 9.9 Hz, 1H), 8.99 (d, J = 7.6 Hz, 1H), 8.30 (s, 1H), 8.14 – 8.03 (m, 1H), 7.41 (m, 2H), 6.40 (m, 1H), 5.11 (s, 1H), 4.03 (m, 3H), 3.80 (m, 1H), 3.67 (m, 1H), 3.54 – 3.42 (m, 1H), 3.21 (m, 1H), 1.90 (dq, J = 12.7, 4.3 Hz, 1H), 1.50 (dq, J = 13.7, 5.1 Hz, 1H); LCMS: 473
104		LCMS: 475

FIGURE 1





105		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32-8.25 (m, 1H), 7.15-6.87 (m, 3H), 5.62-5.58 (m, 1H), 5.51-5.38 (m, 1H), 4.94-4.90 (m, 1H), 4.51-4.47 (m, 1H), 4.14-4.07 (m, 1H), 4.02 (br.s, 1H), 3.52-3.49 (m, 1H), 2.83-2.79 (m, 1H), 2.27-2.24 (m, 1H), 2.14-2.05 (m, 1H), 1.88-1.80 (m, 2H), 1.60-1.48 (m, 3H).; LCMS: 477.1
106		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32-8.25 (m, 1H), 7.15-6.87 (m, 3H), 5.62-5.58 (m, 1H), 5.51-5.38 (m, 1H), 4.94-4.90 (m, 1H), 4.51-4.47 (m, 1H), 4.14-4.07 (m, 1H), 4.02 (br.s, 1H), 3.52-3.49 (m, 1H), 2.83-2.79 (m, 1H), 2.27-2.24 (m, 1H), 2.14-2.05 (m, 1H), 1.88-1.80 (m, 2H), 1.60-1.48 (m, 3H).; LCMS: 477.1
107		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35-8.27 (m, 1H), 7.16-7.01 (m, 3H), 5.63-5.59 (m, 1H), 5.44 (d, 1H, J = 52.0 Hz), 4.51-4.32 (m, 2H), 4.04-3.75 (m, 5H), 3.60-3.57 (m, 1H), 3.39-3.38 (m, 1H), 2.86-2.80 (m, 1H), 2.27-2.13 (m, 1H); LCMS: 479
108		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.22-8.16 (m, 1H), 7.15-7.11 (m, 1H), 7.05-7.01 (m, 1H), 6.94-6.92 (m, 1H), 5.79-5.75 (m, 1H), 4.82-4.75 (m, 1H), 4.46-4.41 (m, 1H), 4.04-4.02 (m, 1H), 3.82-3.79 (m, 1H), 2.98-2.94 (m, 1H), 2.52-2.36 (m, 2H), 2.01-1.92 (m, 1H), 1.71-1.60 (m, 1H), 1.32-1.29 (m, 1H), 1.23-1.19 (m, 1H).; LCMS: 481.1

FIGURE 1





109		¹H-NMR (400 MHz, CDCl₃) δ ppm 11.26 (d, 1H, J = 6.0 Hz), 8.67 (s, 1H), 7.94-7.91 (m, 1H), 7.00-6.76 (m, 3H), 4.55 (d, 2H, J = 6.0 Hz), 4.39 (br.s, 1H), 4.28 (br.s, 1H), 3.82 (s, 3H), 3.39-3.35 (m, 2H), 3.25-3.22 (m, 2H), 3.09-3.08 (m, 1H), 2.79-2.78 (m, 1H).; LCMS: 484.2
110		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40, 8.35 (s, 1H), 7.40 (br.s, 1H), 7.20-7.14 (m, 2H), 5.51 (s, 1H), 4.94 (d, 1H, J = 9.6 Hz), 4.41-4.36 (m, 1H), 4.15-3.97 (m, 2H), 3.65 (s, 1.5H), 3.18 (s, 1.5H), 2.48-2.42 (m, 1H), 2.19-2.14 (m, 1H), 1.82-1.65 (m, 2H).; LCMS: 485.1
111		¹H-NMR (400 MHz, CDCl₃) δ ppm 8.77 (br.s, 1H), 7.83 (d, 1H, J = 8.4 Hz), 7.10-6.97 (m, 3H), 5.45-5.41 (m, 1H), 4.44 (br.s, 1H), 4.35 (br.s, 1H), 3.51-3.43 (m, 2H), 3.36-3.33 (m, 2H), 3.21 (d, 1H, J = 8.0 Hz), 2.85 (d, 1H, J = 2.0 Hz), 1.67 (d, 1H, J = 6.8 Hz).; LCMS: 486.2
112		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.38-8.32 (m, 1H), 7.17-6.67 (m, 3H), 5.77 (dd, 1H, J = 9.2, 6.0 Hz), 4.83-4.77 (m, 1H), 4.40-4.37 (m, 1H), 4.30-4.26 (m, 1H), 3.94-3.76 (m, 1H), 3.05-2.99 (m, 1H), 2.57-2.50 (m, 1H), 1.70-1.65 (m, 1H), 1.48-1.46 (m, 1H), 1.25-1.23 (m, 1H), 0.64-0.61 (m, 1H).; LCMS: 493.1

FIGURE 1





113		¹H-NMR (400 MHz, CDCl₃) δ ppm 11.10 (d, 1H, J = 6.0 Hz), 8.69 (s, 1H), 7.80 (s, 1H), 7.28-7.26 (m, 1H), 6.92-6.87 (m, 2H), 4.33 (br.s, 1H), 4.26 (br.s, 1H), 3.46-3.44 (m, 1H), 3.31-3.25 (m, 1H), 3.12-3.10 (m, 1H), 2.73-2.72 (m, 1H), 1.32-1.24 (m, 4H).; LCMS: 498.2
114		LCMS: 513
115		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.15-7.01 (m, 3H), 5.62-5.57 (m, 1H), 5.49-5.36 (m, 1H), 4.44-4.41 (m, 2H), 4.17-4.12 (m, 1H), 3.75-3.68 (m, 1H), 2.96 (s, 3H), 2.80-2.70 (m, 1H), 2.68-2.64 (m, 1H), 2.41-2.39 (m, 1H), 2.10-2.00 (m, 4H).; LCMS: 525.1
116		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.15-7.01 (m, 3H), 5.61-5.57 (m, 1H), 5.49-5.36 (m, 1H), 4.44-4.41 (m, 2H), 4.17-4.12 (m, 1H), 3.75-3.68 (m, 1H), 2.96 (s, 3H), 2.80-2.70 (m, 1H), 2.68-2.64 (m, 1H), 2.41-2.39 (m, 1H), 2.10-2.00 (m, 4H).; LCMS: 525.1

FIGURE 1





117		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (s, 1H), 7.01-6.96 (m, 3H), 4.62 (s, 2H), 4.37-4.31 (m, 1H), 4.13 (br.s, 1H), 4.04-4.01 (m, 1H), 3.88 (s, 3H), 2.26-2.24 (m, 1H), 2.46-2.39 (m, 1H), 2.16-2.14 (m, 1H), 1.82-1.72 (m, 2H).; LCMS:ND
118		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.01 (d, <i>J</i> = 41.7 Hz, 1H), 9.31 (d, <i>J</i> = 7.6 Hz, 1H), 8.98 (d, <i>J</i> = 7.5 Hz, 1H), 8.29 (d, <i>J</i> = 9.0 Hz, 1H), 7.36 – 7.07 (m, 3H), 5.33 (s, 1H), 4.81 (dd, <i>J</i> = 9.2, 5.0 Hz, 2H), 4.48 – 4.23 (m, 3H), 3.72 – 3.55 (m, 1H), 3.49 (s, 2H), 3.05 (s, 1H), 1.56 (dt, <i>J</i> = 29.6, 6.3 Hz, 1H), 1.38 – 1.20 (m, 1H), 1.08 (dq, <i>J</i> = 8.5, 4.1 Hz, 1H)
119		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.18-7.04 (m, 3H), 5.61 (t, 1H, <i>J</i> = 8.4 Hz), 5.47 (d, 1H, <i>J</i> = 51.6 Hz), 4.51-4.38 (m, 1H), 4.14-4.08 (m, 2H), 2.89-2.81 (m, 1H), 2.33-1.69 (m, 8H); LCMS: 447
120		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.28 (s, 0.2H), 8.26 (s, 0.8H), 7.24-7.20 (m, 1H), 7.11-7.09 (m, 1H), 5.77-5.67 (m, 1H), 5.04-4.95 (m, 1H), 4.33-4.27 (m, 1H), 4.05-4.04 (m, 1H), 3.93-3.89 (m, 1H), 3.43 (s, 1.8H), 2.92 (s, 1.2H), 2.40-2.35 (m, 1H), 2.08-2.07 (m, 1H), 1.72-1.71 (m, 1H), 1.62-1.59 (m, 1H); LCMS: 453

FIGURE 1





121		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.14 (s, 1H), 9.01 (d, <i>J</i> = 7.2 Hz, 1H), 8.38 – 8.05 (m, 1H), 7.78 (t, <i>J</i> = 1.5 Hz, 1H), 7.74 – 7.62 (m, 2H), 5.44 – 5.31 (m, 2H), 4.92 (d, <i>J</i> = 4.0 Hz, 1H), 4.77 (dd, <i>J</i> = 21.4, 14.1 Hz, 1H), 4.46 (dd, <i>J</i> = 39.9, 14.0 Hz, 1H), 4.18 (d, <i>J</i> = 4.8 Hz, 1H), 3.79 (t, <i>J</i> = 4.7 Hz, 1H), 2.73 (dt, <i>J</i> = 16.8, 8.4 Hz, 1H), 2.20 – 1.99 (m, 2H), 1.78 – 1.60 (m, 3H), 1.56 – 1.32 (m, 2H)
122		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.23-8.15 (m, 1H), 7.36-7.31 (m, 1H), 7.15-7.14 (m, 1H), 7.06 (d, 1H, <i>J</i> = 9.6 Hz), 6.96-6.94 (m, 1H), 5.41 (br.s, 1H), 5.37 (d, 1H, <i>J</i> = 50.4 Hz), 4.60-4.33 (m, 4H), 4.81-4.70 (m, 1H), 2.83-2.74 (m, 1H), 2.20-2.15 (m, 1H), 1.75-1.63 (m, 1H), 1.46-1.40 (m, 1H), 1.38-1.17 (m, 2H), 0.54-0.51 (m, 1H); LCMS: 457
123		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26 (s, 1H), 7.60 (s, 1H), 7.48-7.43 (m, 2H), 5.48-5.45 (m, 1H), 5.43-5.33 (m, 1H), 5.16-5.02 (m, 1H), 4.80-4.75 (m, 1H), 4.55-4.48 (m, 2H), 4.06-4.03 (m, 1H), 2.83-2.81 (m, 1H), 2.55-2.43 (m, 2H), 1.93-1.85 (m, 3H); LCMS: 472.1
124		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.24-8.17 (m, 1H), 7.15-7.01 (m, 3H), 5.61-5.57 (m, 1H), 5.41 (d, 1H, <i>J</i> = 51.6 Hz), 4.88-4.86 (m, 1H), 4.43-4.37 (m, 2H), 4.09-4.00 (m, 1H), 3.87-3.85 (m, 1H), 3.28 (s, 3H), 2.82-2.80 (m, 1H), 2.45-2.24 (m, 3H), 1.78-1.69 (m, 2H); LCMS: 477

FIGURE 1





125		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.24-8.17 (m, 1H), 7.14-6.84 (m, 3H), 5.61-5.57 (m, 1H), 5.41 (d, 1H, J = 51.6 Hz), 4.89-4.86 (m, 1H), 4.46-4.36 (m, 2H), 4.00-3.96 (m, 1H), 3.88-3.86 (m, 1H), 3.29 (s, 3H), 2.82-2.75 (m, 1H), 2.38-2.25 (m, 3H), 1.86-1.80 (m, 1H), 1.67-1.64 (m, 1H); LCMS:477
126		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.19 (s, 1H), 6.96-6.94 (m, 2H), 6.80-6.77 (m, 1H), 5.44-5.30 (m, 2H), 4.79-4.76 (m, 1H), 4.48-4.38 (m, 2H), 3.87 (br.s, 1H), 3.65-3.53 (m, 1H), 2.84-2.74 (m, 1H), 2.05-2.00 (m, 2H), 1.78-1.72 (m, 2H), 1.53-1.35 (m, 3H).; LCMS: 477.1
127		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.23-8.16 (m, 1H), 7.21 (s, 1H), 7.08-7.05 (m, 2H), 5.44-5.31 (m, 2H), 4.80-4.79 (m, 1H), 4.50-4.41 (m, 2H), 4.06-4.04 (m, 1H), 3.85-3.82 (m, 1H), 2.83-2.77 (m, 1H), 2.38-2.36 (m, 1H), 2.07-2.00 (m, 2H), 1.74-1.71 (m, 1H), 1.52-1.49 (m, 1H).; LCMS: 479
128		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.23-8.20 (m, 2H), 7.59 (dd, 1H, J = 8.8, 3.2 Hz), 5.67-5.62 (m, 1H), 5.40 (d, 1H, J = 52.8 Hz), 4.63-4.40 (m, 3H), 4.03-4.01 (m, 1H), 3.80 (dd, 1H, J = 7.2, 4.4 Hz), 2.95-2.87 (m, 1H), 2.37-2.35 (m, 1H), 2.06-1.99 (m, 2H), 1.71-1.68 (m, 1H), 1.49-1.46 (m, 1H). ; LCMS: 480

FIGURE 1





129		ND
130		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.19-8.12 (m, 1H), 7.20 (s, 1H), 7.05 (d, 2H, J = 8.8 Hz), 5.43-5.30 (m, 2H), 4.77-4.76 (m, 1H), 4.49-4.39 (m, 2H), 3.87 (br.s, 1H), 3.56-3.53 (m, 1H), 2.81-2.75 (m, 1H), 2.15-2.13 (m, 1H), 1.81-1.73 (m, 2H), 1.58-1.35 (m, 4H).; LCMS: 493.1
131		¹H-NMR (400 MHz, CDCl₃) δ ppm 8.38 (br.s, 1H), 7.32-7.22 (m, 5H), 5.42-5.28 (m, 1H), 4.84-4.69 (m, 1H), 4.55 (br.s, 1H), 4.22 (br.s, 1H), 4.04 (br.s, 1H), 3.51 (s, 1.5H), 3.07 (s, 1.5H), 2.42 (br.s, 1H), 2.13 (br.s, 1H), 1.87 (br.s, 1H), 1.71 (br.s, 1H).; LCMS: 383.2
132		¹H-NMR (400 MHz, CDCl₃) δ ppm 9.80 (s, 0.5H), 9.64 (s, 0.5H), 8.38 (d, 1H, J = 5.2 Hz), 7.35-7.27 (m, 1H), 7.13-7.02 (m, 3H), 5.40 (s, 1H), 4.82 (d, 1H, J = 4.8 Hz), 4.38 (br.s, 1H), 4.22 (s, 1H), 3.95 (s, 1H), 3.57 (s, 1.5H), 3.09 (s, 1.5H), 2.52 (br.s, 1H), 2.10-2.05 (m, 1H), 1.96 (m, 1H), 1.89-1.82 (m, 1H).; LCMS: 401.1

FIGURE 1





133		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35-8.33 (m, 1H), 8.16-8.12 (m, 1H), 7.97-7.88 (m, 1H), 7.34-7.30 (m, 1H), 5.47 (s, 1H), 4.85 (s, 1H), 4.38 (br.s, 1H), 4.15-4.12 (m, 1H), 4.03-3.98 (m, 1H), 3.70 (s, 1.5H), 3.35-3.25 (m, 1H), 3.22 (s, 1.5H), 2.48-2.40 (m, 1H), 2.14 (br.s, 1H), 1.81-1.75 (m, 1H).; LCMS: 402.1
134		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (d, 1H, J = 4.8 Hz), 7.76-7.67 (m, 3H), 7.58-7.56 (m, 1H), 5.47 (d, 1H, J = 2.4 Hz), 4.91 (s, 1H), 4.42-4.37 (m, 1H), 4.15-4.14 (m, 1H), 4.04-3.99 (m, 1H), 3.65 (s, 1.5H), 3.16 (s, 1.5H), 2.48-2.45 (m, 1H), 1.84-1.81 (m, 1H), 1.71-1.66 (m, 2H).; LCMS: 408.1
135		¹H NMR (400 MHz, DMSO-d₆) δ 13.98 (s, 1H), 9.26 (m, 1H), 8.26 (d, J = 11.5 Hz, 1H), 7.99 (m, 1H), 7.45 – 7.28 (m, 2H), 5.47 (m, 1H), 5.02 (m, 1H), 4.25 (m, 1H), 3.94 (m, 1H), 3.56 (s, 1H), 3.30 (s, 3H), 3.09 (s, 1H), 2.22 – 2.03 (m, 1H), 1.91 (m, 1H), 1.84 – 1.64 (m, 2H), 1.64 – 1.34 (m, 2H).; LCMS: 410.2
136		¹H NMR (400 MHz, DMSO-d₆) δ 13.89 (s, 1H), 9.10 (ABqd, 1H), 8.21 (d, J = 10.6 Hz, 1H), 7.87 (ddd, J = 8.3, 4.7, 3.3 Hz, 1H), 7.33 (ddd, J = 9.4, 7.9, 3.2 Hz, 1H), 4.97 (m, 2H), 4.46 (s, 1H), 4.27 – 4.12 (m, 1H), 3.99 – 3.81 (m, 1H), 3.39 (d, J = 6.3 Hz, 3H), 3.29 (s, 3H), 2.18 – 2.01 (m, 1H), 1.86 – 1.35 (m, 5H).; LCMS: 416.1

FIGURE 1





137		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (s, 1H), 7.52-7.41 (m, 4H), 5.49 (s, 1H), 4.91 (s, 1H), 4.63-4.60 (m, 1H), 4.27 (br.s, 1H), 4.04 (b.s, 1H), 3.67 (s, 1.7H), 3.21 (s, 1.3H), 2.56-2.55 (m, 1H), 2.23 (br.s, 1H), 1.90 (br.s, 1H), 1.71 (br.s, 1H).; LCMS: 417.1
138		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.30 (m, 2H), 7.79-7.74 (m, 1H), 7.42-7.38 (m, 1H), 5.44 (s, 1H), 4.96 (s, 1H), 4.41-4.34 (m, 1H), 4.15-3.97 (m, 2H), 3.97 (s, 1H), 3.25 (s, 2H), 2.47-2.42 (m, 1H), 2.17-2.15 (m, 1H), 1.82-1.74 (m, 2H).; LCMS: 418.0
139		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.25 (d, 1H, J = 5.2 Hz), 7.23-7.14 (m, 3H), 5.51 (s, 1H), 4.94 (s, 1H), 4.51-4.47 (m, 1H), 4.15-4.13 (m, 1H), 3.94-3.89 (m, 1H), 3.89 (s, 1.5H), 3.15 (s, 1.5H), 2.45-2.41 (m, 1H), 2.12-2.04 (m, 1H), 1.80-1.77 (m, 1H), 1.60-1.58 (m, 1H), 1.38-1.29 (m, 1H).; LCMS: 419.1
140		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40 (d, 1H, J = 8.8 Hz), 7.85 (dd, 1H, J = 7.6, 5.6 Hz), 7.31-7.22 (m, 2H), 6.22-6.17 (m, 1H), 6.09-6.04 (m, 1H), 5.61 (s, 1H), 5.05 (s, 1H), 5.02-5.01 (m, 1H), 4.61 (dd, 1H, J = 14.8, 2.8 Hz), 4.12 (td, 1H, J = 18.4, 5.6 Hz), 3.69 (s, 1.5H), 3.21 (s, 1.5H).; LCMS: 424.1

FIGURE 1





141		¹H-NMR (400 MHz, CDCl₃) δ ppm 8.38 (br.s, 1H), 7.68-7.65 (m, 1H), 7.10-7.03 (m, 2H), 5.55-5.46 (m, 1H), 4.94-4.83 (m, 1H), 4.56 (br.s, 1H), 4.21 (br.s, 1H), 4.09-4.02 (m, 1H), 3.58 (s, 1.5H), 3.11 (s, 1.5H), 2.42 (br.s, 1H), 2.10 (br.s, 1H), 1.85 (br.s, 1H), 1.69 (s, 1H).; LCMS: 426.1
142		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35 (d, 1H, J = 5.6 Hz), 7.75-7.65 (m, 2H), 7.37-7.34 (m, 1H), 5.53 (s, 1H), 4.95 (s, 1H), 4.38-4.33 (m, 1H), 4.13-4.11 (m, 1H), 4.04-3.96 (m, 1H), 3.68 (s, 1.5H), 3.20 (s, 1.5H), 2.45-2.42 (m, 1H), 2.17-2.13 (m, 1H), 1.80-1.69 (m, 2H).; LCMS: 426.1
143		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32 (d, 1H, J = 8.4 Hz), 7.87-7.82 (m, 1H), 7.33-7.23 (m, 2H), 5.66-5.55 (m, 1H), 5.11-5.02 (m, 1H), 4.06-3.95 (m, 3H), 3.68 (s, 1.5H), 3.67-3.58 (m, 1H), 3.52-3.49 (m, 1H), 3.25-3.23 (m, 1H), 3.22 (s, 1.5H), 2.20-2.11 (m, 1H), 1.82-1.74 (m, 1H).; LCMS: 426.1
144		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 7.32-7.25 (m, 3H), 7.24-7.19 (m, 2H), 5.47-5.35 (m, 2H), 4.84-4.83 (m, 1H), 4.53-4.29 (m, 2H), 4.05-3.97 (m, 1H), 3.90-3.87 (m, 1H), 2.89-2.78 (m, 1H), 2.38-2.35 (m, 1H), 2.18-2.06 (m, 2H), 1.77-1.61 (m, 2H).; LCMS: 427.1

FIGURE 1





145		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33 (s, 1H), 7.34-7.19 (m, 5H), 5.48-5.36 (m, 2H), 4.93 (br.s, 1H), 4.53-4.44 (m, 1H), 3.98-3.91 (m, 3H), 3.50-3.46 (m, 2H), 3.23-3.20 (m, 1H), 2.80-2.79 (m, 1H), 2.13-2.07 (m, 2H), 1.77-1.74 (m, 1H).; LCMS: 427.2
146		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.17 (d, 1H, J = 8.4 Hz), 7.78-7.72 (m, 1H), 7.21-7.11 (m, 2H), 5.52 (s, 1H), 5.48-4.98 (m, 1H), 4.95 (s, 1H), 4.50-4.44 (m, 1H), 4.10-4.03 (m, 1H), 3.56 (s, 1.5H), 3.10 (s, 1.5H), 2.52-2.41 (m, 2H), 1.88-1.81 (m, 2H).; LCMS: 428.1
147		¹ H NMR (400 MHz, DMSO-d ₆) δ 13.92 (br d, 1H), 9.19 (ABqd, 1H), 9.05 (d, J = 7.2 Hz, 0H), 8.20 (d, J = 9.9 Hz, 1H), 7.87 (m, 1H), 7.33 (m, 1H), 4.97 (d, J = 3.8 Hz, 1H), 4.87 (dd, J = 14.6, 5.5 Hz, 1H), 4.51 – 4.42 (m, 2H), 4.34 (m, 1H), 3.88 (m, 1H), 3.65 (ddt, J = 26.8, 7.3, 5.1 Hz, 1H), 3.40 (d, J = 6.3 Hz, 3H), 3.29 (s, 3H), 2.20 (m, 1H), 1.87 (m, 1H), 1.52 (m, 1H), 1.36 – 1.15 (m, 1H).; LCMS: 432.1
148		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35 (d, 1H, J = 4.4 Hz), 7.42 (d, 1H, J = 2.4 Hz), 7.41-7.32 (m, 1H), 7.18-7.14 (m, 1H), 5.48 (s, 1H), 4.91-4.89 (m, 1H), 4.40-4.36 (m, 1H), 4.15-4.13 (m, 1H), 4.05-3.99 (m, 1H), 3.66 (s, 1.5H), 3.17 (s, 1.5H), 3.47-2.44 (m, 1H), 2.18-2.14 (m, 1H), 1.82-1.80 (m, 1H), 1.71-1.65 (m, 1H).; LCMS: 435.1

FIGURE 1





149		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35 (d, 1H, J = 12.4 Hz), 8.25 (d, 1H, J = 8.8 Hz), 7.64-7.59 (m, 1H), 5.38 (s, 1H), 4.92 (s, 1H), 4.16-3.88 (m, 4H), 3.72 (s, 1H), 3.59-3.39 (m, 2H), 3.26 (s, 2H), 2.13-2.00 (m, 1H), 1.78-1.68 (m, 1H).; LCMS: 436.1
150		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (dd, 1H, J = 12.8, 2.0 Hz), 8.25 (dd, 1H, J = 9.2, 2.0 Hz), 7.64-7.59 (m, 1H), 5.38 (s, 1H), 4.02-3.94 (m, 3H), 3.72 (s, 1H), 3.68-3.48 (m, 2H), 3.24 (s, 2H), 3.23-3.22 (m, 1H), 2.19-2.10 (m, 1H), 1.88-1.75 (m, 1H).; LCMS: 436.1
151		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.27-8.23 (m, 2H), 7.61-7.57 (m, 1H), 5.32 (s, 1H), 5.24-5.10 (m, 1H), 4.94 (s, 1H), 4.58-4.49 (m, 1H), 4.18-4.08 (m, 1H), 3.67 (s, 1H), 3.20 (s, 2H), 2.65-2.48 (m, 2H), 1.99-1.82 (m, 2H).; LCMS: 438.1
152		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40 (d, 1H, J = 6.0 Hz), 7.88-7.84 (m, 1H), 7.36-7.24 (m, 2H), 5.62 (d, 1H, J = 4.0 Hz), 5.07 (d, 1H, J = 5.6 Hz), 4.61 (dt, 1H, J = 32.0, 5.6 Hz), 4.30 (d, 1H, J = 10.4 Hz), 3.94 (dd, 1H, J = 20.8, 6.0 Hz), 3.72 (s, 1.5H), 3.23 (s, 1.5H), 1.84-1.78 (m, 1H), 1.59-1.53 (m, 1H), 1.32-1.28 (m, 1H), 0.71-0.66 (m, 1H).; LCMS: 438.1

FIGURE 1





153		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33-8.32 (m, 1H), 7.32-7.18 (m, 6H), 5.48-5.35 (m, 2H), 4.92-4.90 (m, 1H), 4.49-4.45 (m, 1H), 4.39-4.36 (m, 1H), 4.24-4.20 (m, 1H), 3.74-3.73 (m, 1H), 2.83-2.75 (m, 1H), 2.24-2.10 (m, 1H), 1.70-1.67 (m, 1H), 1.48-1.46 (m, 1H), 1.28-1.20 (m, 1H), 0.65-0.59 (m, 1H).; LCMS: 439.1
154		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.13 (br.s, 1H), 7.81 (br.s, 1H), 7.29 (br.s, 1H), 7.20 (br.s, 1H), 5.61 (br.s, 1H), 5.02-4.99 (m, 1H), 4.42 (br.s, 1H), 4.02-3.97 (m, 1H), 3.62 (s, 2H), 3.14 (s, 1H), 2.06 (s, 1H), 1.82-1.40 (m, 5H). ; LCMS: 440.3
155		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.31 (s, 1H), 7.32-7.25 (m, 3H), 7.24-7.20 (m, 2H), 5.47-5.35 (m, 2H), 4.53-4.43 (m, 1H), 4.10-3.97 (m, 3H), 3.49-3.46 (m, 1H), 2.81-2.79 (m, 1H), 2.17-2.03 (m, 2H), 1.88-1.83 (m, 2H), 1.59-1.49 (m, 3H).; LCMS: 441.1
156		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33 (d, 1H, J = 8.0 Hz), 7.80-7.77 (m, 1H), 7.58-7.50 (m, 2H), 5.61-5.60 (m, 1H), 5.05 (d, 1H, J = 4.8 Hz), 4.42-4.37 (m, 1H), 4.15-4.11 (m, 1H), 4.04-3.97 (m, 1H), 3.69 (s, 1.5H), 3.22 (s, 1.5H), 2.47-2.41 (m, 1H), 2.16-2.13 (m, 1H), 1.81-1.69 (m, 2H).; LCMS: 442.1

FIGURE 1





157		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32-8.24 (m, 1H), 7.35-7.31 (m, 1H), 7.14-6.93 (m, 3H), 6.10-5.96 (m, 2H), 5.47-5.03 (m, 2H), 4.83-4.87 (m, 1H), 4.52-4.38 (m, 2H), 4.07-3.94 (m, 1H), 2.83-2.67 (m, 1H), 2.17-2.01 (m, 1H).; LCMS: 443.1
158		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.39-8.32 (m, 1H), 7.39-6.97 (m, 4H), 5.51-5.35 (m, 2H), 4.96-4.91 (m, 1H), 4.49-4.46 (m, 1H), 4.06-3.92 (m, 3H), 3.55-3.49 (m, 2H), 3.26-3.24 (m, 1H), 2.85-2.83 (m, 1H), 2.21-2.11 (m, 2H), 1.78-1.72 (m, 1H).; LCMS: 445.2
159		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40 (s, 0.5H), 8.37 (s, 0.5H), 8.23 (dd, 1H, J = 8.8, 2.8 Hz), 7.64-7.59 (m, 1H), 5.36 (s, 1H), 4.89 (s, 1H), 4.57 (td, 1H, J = 49.2, 4.2 Hz), 4.29 (s, 0.5H), 4.26 (s, 0.5H), 3.90 (dd, 1H, J = 26.4, 6.0 Hz), 3.70 (s, 1.5H), 3.25 (s, 1.5H), 1.82-1.81 (m, 1H), 1.56-1.48 (m, 1H), 1.30-1.27 (m, 1H), 1.68-1.62 (m, 1H).; LCMS: 448.1
160		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37 (d, 1H, J = 4.4 Hz), 7.31-7.29 (m, 1H), 7.13-7.06 (m, 2H), 6.24-6.19 (m, 1H), 4.37-4.35 (m, 1H), 4.12-4.09 (m, 1H), 3.95 (s, 1H), 3.90-3.87 (m, 2H), 3.67-3.65 (m, 1H), 3.47-3.44 (m, 1H), 3.33 (s, 1.5H), 2.86 (s, 1.5H), 1.79 (d, 1.5H, J = 6.8 Hz), 1.66 (d, 1.5H, J = 6.4 Hz).; LCMS: 449.1

FIGURE 1





161		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.25-8.19 (m, 2H), 7.16 (s, 1H), 5.33 (s, 1H), 4.94-4.88 (m, 1H), 4.54-4.46 (m, 1H), 4.03-3.95 (m, 1H), 3.66 (s, 1.5H), 3.60-3.58 (m, 1H), 3.21 (s, 1.5H), 2.20-2.00 (m, 1H), 1.85-1.81 (m, 2H), 1.60-1.35 (m, 3H).; LCMS: 450.1
162		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40-8.37 (m, 1H), 7.49-7.44 (m, 1H), 7.10-7.03 (m, 2H), 5.46 (s, 2H), 4.36-4.35 (m, 1H), 4.11-4.10 (m, 1H), 3.97-3.85 (m, 3H), 3.67-3.65 (m, 2.5H), 3.49-3.40 (m, 1H), 3.22 (s, 1.5H).; LCMS: 451
163		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (d, 1H, J = 4.0 Hz), 7.43-7.30 (m, 2H), 7.17-7.12 (m, 1H), 5.45 (s, 1H), 4.88-4.86 (m, 1H), 4.41-4.37 (m, 1H), 4.16-4.11 (m, 1H), 3.97-3.95 (m, 1H), 3.89-3.86 (m, 2H), 3.69-3.65 (m, 1H), 3.64 (s, 1.5H), 3.47-3.45 (m, 1H), 3.16 (s, 1.5H).; LCMS: 451.1
164		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.30 (d, 1H, J = 5.6 Hz), 7.11-6.96 (m, 2H), 5.48 (s, 1H), 4.93 (s, 1H), 4.46 (d, 1H, J = 7.6 Hz), 4.12-4.09 (m, 1H), 3.95-3.80 (m, 3H), 3.66-3.64 (m, 2.5H), 3.44-3.41 (m, 1H), 3.18 (s, 1.5H).; LCMS: 453.1

FIGURE 1





165		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33 (d, 1H, J = 6.4 Hz), 7.79-7.76 (m, 1H), 7.60-7.53 (m, 2H), 5.58 (d, 1H, J = 5.2 Hz), 5.03 (d, 1H, J = 5.2 Hz), 4.64-4.57 (m, 1H), 4.36-4.33 (m, 1H), 3.93-3.87 (dd, 1H, J = 17.2, 6.4 Hz), 3.69 (s, 1.5H), 3.02 (s, 1.5H), 1.80-1.76 (m, 1H), 1.55-1.51 (m, 1H), 1.29-1.25 (m, 1H), 0.67-0.62 (m, 1H). ; LCMS: 454.1
166		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.47 (s, 1H), 8.38-8.23 (m, 2H), 7.70 (d, 1H, J = 9.6 Hz), 5.52-5.38 (m, 2H), 4.93-4.89 (m, 1H), 4.56-4.43 (m, 1H), 4.34-4.32 (m, 1H), 4.11-4.07 (m, 1H), 3.18-3.14 (m, 1H), 2.86-2.82 (m, 1H), 2.56-2.52 (m, 1H), 2.51-2.38 (m, 1H), 2.19-2.16 (m, 2H), 1.94-1.91 (m, 1H).; LCMS: 455.1
167		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32 (s, 0.5H), 8.30 (s, 0.5H), 7.80-7.77 (m, 1H), 7.58-7.52 (m, 2H), 5.66-5.55 (m, 1H), 5.10-5.00 (m, 1H), 4.20-4.07 (m, 2H), 3.70 (s, 1.5 H), 3.66-3.59 (m, 1H), 3.22 (s, 1.5H), 2.14-2.07 (m, 1H), 1.90-1.82 (m, 2H), 1.64-1.55 (m, 3H). ; LCMS: 456.1
168		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.31-8.23 (m, 1H), 7.37-7.33 (m, 1H), 7.15 (d, 1H, J = 8.0 Hz), 7.06 (d, 1H, J = 10.0 Hz), 7.00-6.96 (m, 1H), 5.48-5.35 (m, 2H), 4.54-4.44 (m, 1H), 4.13-4.01 (m, 2H), 3.51-3.48 (m, 1H), 2.84-2.80 (m, 1H), 2.32-2.04 (m, 2H), 1.84-1.79 (m, 2H), 1.59-1.50 (m, 3H).; LCMS: 459.2

FIGURE 1





169		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37-8.29 (m, 1H), 7.44-6.96 (m, 4H), 5.46 (br.s, 1H), 5.40 (d, 1H, J = 52.0 Hz), 4.87-4.82 (m, 1H), 4.47 (dd, 1H, J = 38.4, 14.8 Hz), 4.30 (br.s, 1H), 4.05-3.83 (m, 2H), 3.75-3.58 (m, 2H), 3.55-3.36 (m, 2H), 3.09-2.81 (m, 1H), 2.21-2.08 (m, 1H).; LCMS: 461.1
170		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.29 (s, 1H), 7.16-7.02 (m, 3H), 6.08-6.00 (m, 2H), 5.62-5.58 (m, 1H), 5.42 (d, 1H, J = 52.4 Hz), 5.16 (br.s, 1H), 4.66-4.58 (m, 2H), 4.49-4.36 (m, 1H), 4.07-3.94 (m, 1H), 2.88-2.78 (m, 1H), 2.25-2.09 (m, 1H).; LCMS: 461.1
171		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35-8.28 (m, 1H), 6.96-6.94 (m, 1H), 6.85-6.78 (m, 2H), 6.23-6.04 (m, 2H), 5.46-5.33 (m, 2H), 5.04 (br.s, 1H), 4.56-4.40 (m, 2H), 4.11-3.97 (m, 1H), 2.86-2.76 (m, 1H), 2.19-2.03 (m, 1H).; LCMS: 461.1
172		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32-8.23 (m, 1H), 7.16-7.09 (m, 3H), 6.10-5.98 (m, 2H), 5.66-5.61 (m, 1H), 5.43 (d, 1H, J = 52.0 Hz), 5.06-5.05 (m, 1H), 4.55-4.39 (m, 3H), 3.96-3.93 (m, 1H), 2.87-2.77 (m, 1H), 2.26-2.12 (m, 1H). ; LCMS: 461.1

FIGURE 1





173		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 6.95 (d, 2H, J = 6.8 Hz), 6.87-6.67 (m, 1H), 5.46-5.33 (m, 2H), 4.91-4.89 (m, 1H), 4.53-4.50 (m, 1H), 4.34-4.30 (m, 1H), 4.05-4.04 (m, 1H), 3.91-3.88 (m, 1H), 2.84-2.78 (m, 1H), 2.38-2.37 (m, 1H), 2.10-2.06 (m, 2H), 1.74-1.61 (m, 2H).; LCMS: 463.1
174		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33-8.26 (m, 1H), 6.98 (d, 1H, J = 6.4 Hz), 6.88-6.70 (m, 2H), 5.48-5.35 (m, 2H), 4.55-4.52 (m, 1H), 4.01-3.90 (m, 3H), 3.55-3.48 (m, 2H), 3.25-3.19 (m, 1H), 2.84-2.78 (m, 1H), 2.18-2.09 (m, 2H), 1.76-1.66 (m, 2H).; LCMS: 463.1
175		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32-8.24 (m, 1H), 7.17-6.96 (m, 3H), 5.69-5.65 (m, 1H), 5.46 (d, 1H, J = 52.0 Hz), 4.94-4.89 (m, 1H), 4.57-4.37 (m, 1H), 4.03-3.88 (m, 3H), 3.52-3.48 (m, 2H), 3.24-3.22 (m, 1H), 2.87-2.81 (m, 1H), 2.16-2.10 (m, 2H), 1.74-1.71 (m, 1H). ; LCMS: 463.1
176		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.24-8.16 (m, 1H), 7.16-6.92 (m, 3H), 5.68-5.64 (m, 1H), 5.42 (d, 1H, J = 52.0 Hz), 4.91-4.89 (m, 1H), 4.48-4.41 (m, 2H), 4.06-4.04 (m, 1H), 3.84-3.81 (m, 1H), 2.86-2.80 (m, 1H), 2.40-2.35 (m, 1H), 2.09-2.01 (m, 2H), 1.74-1.72 (m, 1H), 1.53-1.51 (m, 1H). ; LCMS: 463.1

FIGURE 1





177		¹H NMR (400 MHz, DMSO-d₆) δ 14.17 (s, 1H), 8.99 (d, J = 7.2 Hz, 1H), 8.38 (d, J = 2.9 Hz, 1H), 8.26 (s, 1H), 7.87 (dd, J = 8.9, 3.0 Hz, 1H), 5.67 – 5.34 (m, 3H), 4.92 (d, J = 3.9 Hz, 1H), 4.80 (dd, J = 21.4, 14.0 Hz, 1H), 4.59 – 4.34 (m, 1H), 4.18 (s, 1H), 3.85 – 3.75 (m, 1H), 2.87 – 2.71 (m, 1H), 2.26 – 2.00 (m, 2H), 1.66 (dq, J = 13.1, 5.5 Hz, 2H), 1.58 – 1.34 (m, 2H).; LCMS: 464
178		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37 (s, 1H), 6.97-6.95 (m, 2H), 6.84-6.79 (m, 1H), 5.47-5.41 (m, 1H), 5.10 (d, 1H, J = 53.6 Hz), 4.82 (s, 1H), 4.54-4.41 (m, 2H), 4.14-4.11 (m, 1H), 2.83-2.60 (m, 1H), 2.54-2.47 (m, 2H), 2.07-1.84 (m, 3H).; LCMS: 465.1
179		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.42 (br.s, 1H), 7.13-7.03 (m, 2H), 6.97-6.74 (m, 2H), 5.48 (d, 1H, J = 1.6 Hz), 4.91 (s, 1H), 4.47-4.43 (m, 1H), 4.20-4.19 (m, 1H), 4.09-4.03 (m, 1H), 3.69 (s, 1.5H), 3.21 (s, 1.5H), 2.53-2.49 (m, 1H), 2.23-2.20 (m, 1H), 1.87-1.69 (m, 2H).; LCMS: 467.1
180		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 7.22-7.04 (m, 3H), 5.85 (td, 1H, J = 57.2, 4.4 Hz), 5.54-5.51 (m, 1H), 4.25-4.16 (m, 2H), 3.90-3.83 (m, 2H), 3.35-3.33 (m, 1H), 2.31-2.19 (m, 2H), 1.92-1.85 (m, 1H), 1.69-1.66 (m, 1H).; LCMS: 469.1

FIGURE 1





181		¹ H NMR (400 MHz, DMSO-d ₆) δ 14.18 (s, 1H), 9.10 (d, J = 7.3 Hz, 1H), 8.25 (s, 1H), 7.78 (s, 1H), 7.74 – 7.59 (m, 2H), 5.57 – 5.35 (m, 2H), 5.06 (d, J = 5.4 Hz, 1H), 4.76 – 4.63 (m, 1H), 4.50 (dd, J = 39.8, 14.4 Hz, 1H), 4.09 (s, 2H), 3.69 (dd, J = 11.3, 4.5 Hz, 2H), 3.42 – 3.37 (m, 1H), 3.09 (dd, J = 11.4, 8.7 Hz, 1H), 2.73 (dd, J = 33.7, 20.2 Hz, 1H), 2.09 (q, J = 13.8, 11.9 Hz, 2H), 1.50 – 1.33 (m, 1H).; LCMS: 470
182		¹ H NMR (400 MHz, DMSO-d ₆) δ 14.17 (s, 1H), 9.11 (d, J = 7.8 Hz, 1H), 8.26 (s, 1H), 7.88 – 7.55 (m, 3H), 5.52 (s, 1H), 5.45 – 5.35 (m, 1H), 5.02 (d, J = 4.7 Hz, 1H), 4.73 (dd, J = 21.9, 13.9 Hz, 1H), 4.48 (dd, J = 39.7, 13.0 Hz, 1H), 4.03 (dd, J = 7.5, 3.7 Hz, 1H), 3.92 (dd, J = 11.1, 3.7 Hz, 1H), 3.73 (dd, J = 11.0, 5.7 Hz, 1H), 3.56 – 3.46 (m, 1H), 3.43 – 3.34 (m, 1H), 3.18 (dd, J = 11.3, 6.5 Hz, 1H), 2.83 – 2.63 (m, 1H), 2.19 – 1.95 (m, 1H), 1.81 – 1.65 (m, 1H), 1.42 (d, J = 4.4 Hz, 1H).; LCMS: 470
183		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33-8.26 (m, 1H), 7.84-7.81 (m, 1H), 7.36 (dd, 1H, J = 9.2, 2.0 Hz), 7.24-7.19 (m, 1H), 5.66 (dd, 1H, J = 10.4, 8.0 Hz), 5.49 (d, 1H, J = 52.0 Hz), 4.95-4.89 (m, 1H), 4.65-4.56 (m, 1H), 3.97-3.89 (m, 3H), 3.51-3.48 (m, 2H), 3.25-3.22 (m, 1H), 2.90-2.86 (m, 1H), 2.28-2.11 (m, 2H), 1.74-1.70 (m, 1H).; LCMS: 470.1
184		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33-8.26 (m, 1H), 7.13-7.07 (m, 3H), 5.60-5.55 (m, 1H), 5.43 (d, 1H, J = 52.0 Hz), 4.47-4.33 (m, 2H), 4.13-4.09 (m, 1H), 3.20-3.14 (m, 2H), 2.83-2.78 (m, 1H), 2.53-2.51 (m, 1H), 2.49-2.38 (m, 1H), 2.21-2.15 (m, 2H), 1.92-1.89 (m, 1H).; LCMS: 472.1

FIGURE 1

185		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.27 (m, 1H), 6.97 (d, 2H, J = 6.0 Hz), 6.85-6.80 (m, 1H), 5.48-5.34 (m, 2H), 4.92-4.86 (m, 1H), 4.72-4.61 (m, 1H), 4.42-4.36 (m, 2H), 4.13-4.12 (m, 1H), 3.17-3.13 (m, 1H), 2.85-2.80 (m, 1H), 2.54-2.51 (m, 1H), 2.44-2.40 (m, 1H), 2.21-2.16 (m, 1H), 1.94-1.90 (m, 1H).; LCMS: 472.1
186		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.24 (s, 1H), 7.82-7.79 (m, 1H), 7.34-7.31 (m, 1H), 7.21-7.03 (m, 1H), 5.65-5.61 (m, 1H), 5.45 (d, 1H, J = 52.0 Hz), 5.10 (d, 1H, J = 53.6 Hz), 4.89 (s, 1H), 4.61-4.47 (m, 2H), 4.06-4.01 (m, 1H), 2.87-2.85 (m, 1H), 2.53-2.46 (m, 2H), 2.44-2.41 (m, 1H), 1.92-1.84 (m, 2H).; LCMS: 472.1
187		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.25 (s, 1H), 7.83-7.80 (m, 1H), 7.34-7.31 (m, 1H), 7.21-7.13 (m, 1H), 5.65-5.61 (m, 1H), 5.45 (d, 1H, J = 51.6 Hz), 5.08 (d, 1H, J = 54.0 Hz), 4.78 (s, 1H), 4.61-4.47 (m, 2H), 4.15-4.07 (m, 1H), 3.06-2.85 (m, 1H), 2.55-2.42 (m, 2H), 2.25-2.00 (m, 1H), 1.94-1.82 (m, 2H).; LCMS: 472.1
188		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.27 (m, 1H), 6.98 (d, 1H, J = 6.4 Hz), 6.85-6.69 (m, 2H), 5.48-5.35 (m, 2H), 4.64-4.42 (m, 3H), 4.29-4.25 (m, 1H), 3.78 (d, 1H, J = 5.6 Hz), 2.82 (br.s, 1H), 2.19-2.08 (m, 1H), 1.70-1.64 (m, 1H), 1.55-1.47 (m, 1H), 1.23 (s, 1H), 0.66-0.63 (m, 1H).; LCMS: 475.1

FIGURE 1





189		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.31-8.23 (m, 1H), 7.18-6.95 (m, 3H), 5.66-5.62 (m, 1H), 5.45 (d, 1H, J = 52.0 Hz), 4.97-4.95 (m, 1H), 4.46-4.28 (m, 3H), 3.89-3.72 (m, 1H), 2.88-2.79 (m, 1H), 2.30-2.16 (m, 1H), 1.70-1.67 (m, 1H), 1.49-1.46 (m, 1H), 1.27-1.20 (m, 1H), 0.65-0.58 (m, 1H).; LCMS: 475.1
190		¹H NMR (400 MHz, DMSO-d₆) δ 14.16 (s, 1H), 9.15 (d, J = 7.4 Hz, 1H), 8.25 (s, 1H), 8.05 (d, J = 2.9 Hz, 1H), 7.54 (dd, J = 8.5, 3.1 Hz, 1H), 5.56 – 5.32 (m, 2H), 5.07 (d, J = 5.4 Hz, 1H), 4.79 – 4.62 (m, 1H), 4.53 – 4.27 (m, 1H), 4.09 (m, 1H), 3.91 (s, 3H), 3.71 (m, 2H), 3.48 – 3.32 (m, 2H), 3.11 (m, 1H), 2.84 – 2.61 (m, 1H), 2.23 – 1.97 (m, 2H), 1.53 – 1.32 (m, 1H).; LCMS: 476.1
191		¹H NMR (400 MHz, DMSO-d₆) δ 14.11 (s, 1H), 8.90 (d, J = 7.4 Hz, 1H), 8.23 (s, 1H), 7.34 – 7.07 (m, 3H), 5.59 – 5.43 (m, 2H), 4.76 – 4.58 (m, 3H), 4.50 – 4.29 (m, 1H), 3.91 – 3.76 (m, 1H), 3.58 – 3.39 (m, 1H), 2.82 – 2.63 (m, 1H), 2.23 – 1.91 (m, 3H), 1.74 (d, J = 12.7 Hz, 1H), 1.32 – 1.05 (m, 2H).; LCMS: 477.1
192		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35-8.28 (m, 1H), 7.20 (s, 1H), 7.07-6.93 (m, 2H), 6.22-5.98 (m, 2H), 5.46-5.33 (m, 2H), 5.04-5.03 (m, 1H), 4.64-4.43 (m, 2H), 4.00-3.97 (m, 1H), 2.85-2.76 (m, 1H), 2.19-2.03 (m, 1H).; LCMS: 477.1

FIGURE 1





193		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.30-8.22 (m, 1H), 7.17-6.95 (m, 3H), 5.67-5.63 (m, 1H), 5.44 (d, 1H, J = 52.0 Hz), 4.93-4.87 (m, 1H), 4.50-4.37 (m, 1H), 4.11-4.08 (m, 1H), 3.99 (s, 1H), 3.49-3.46 (m, 1H), 2.83-2.82 (m, 1H), 2.24-2.03 (m, 2H), 1.86-1.78 (m, 2H), 1.59-1.47 (m, 3H).; LCMS: 477.1
194		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32-8.25 (m, 1H), 8.21-8.09 (m, 1H), 7.61-7.48 (m, 1H), 6.10-6.08 (m, 1H), 5.98-5.96 (m, 1H), 5.65-5.61 (m, 1H), 5.48-5.35 (m, 1H), 5.04-5.03 (m, 1H), 4.55-4.51 (m, 2H), 4.08-3.92 (m, 1H), 2.96-2.87 (m, 1H), 2.19-2.04 (m, 1H).; LCMS: 478.1
195		¹H-NMR (400 MHz, CD₃OD) δ ppm 9.28 (br.s, 1H), 8.30-8.21 (m, 1H), 7.51 (br.s, 1H), 7.18-7.00 (m, 2H), 5.58-5.39 (m, 2H), 4.65-4.31 (m, 3H), 3.84-3.62 (m, 2H), 2.65 (br.s, 4H), 2.19-1.28 (m, 2H).; LCMS: 479.1
196		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.28 (m, 1H), 7.46-7.42 (m, 1H), 7.09-6.91 (m, 2H), 5.75 (t, 1H, J = 8.8 Hz), 5.50-5.33 (m, 1H), 4.61-4.53 (m, 1H), 4.03-3.89 (m, 4H), 3.52-3.47 (m, 2H), 3.24-3.19 (m, 1H), 2.93-2.91 (m, 1H), 2.17-2.10 (m, 2H), 1.75-1.71 (m, 1H).; LCMS: 479.1

FIGURE 1

197		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35-8.28 (m, 1H), 6.97 (d, 1H, J = 6.4 Hz), 6.87-6.80 (m, 2H), 5.48-5.34 (m, 2H), 4.54-4.33 (m, 1H), 4.32-4.07 (m, 1H), 4.05-4.03 (m, 2H), 3.85-3.84 (m, 2H), 3.77-3.75 (m, 1H), 3.59-3.57 (m, 1H), 3.41-3.36 (m, 1H), 2.86-2.79 (m, 1H), 2.20-2.04 (m, 1H).; LCMS: 479.1
198		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 7.22 (s, 1H), 7.08 (d, 2H, J = 8.8 Hz), 5.47-5.34 (m, 2H), 4.86-4.82 (m, 1H), 4.55-4.50 (m, 1H), 3.97-3.90 (m, 3H), 3.54-3.46 (m, 2H), 3.23-3.21 (m, 1H), 2.83-2.81 (m, 1H), 2.17-2.07 (m, 2H), 1.76-1.73 (m, 1H).; LCMS: 479.1
199		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.30-8.23 (m, 1H), 7.07-7.03 (m, 1H), 6.92-6.91 (m, 1H), 6.13-6.06 (m, 1H), 6.04-5.97 (m, 1H), 5.62-5.58 (m, 1H), 5.41 (d, 1H, J = 51.6 Hz), 5.08-5.07 (m, 1H), 4.55-4.35 (m, 2H), 4.05-3.93 (m, 1H), 2.86-2.76 (m, 1H), 2.24-2.08 (m, 1H).; LCMS: 479.1
200		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.30-8.23 (m, 1H), 7.35-7.26 (m, 3H), 5.61-5.57 (m, 1H), 5.40 (d, 1H, J = 52.4 Hz), 4.52-4.36 (m, 1H), 4.03-3.87 (m, 4H), 3.53-3.47 (m, 2H), 3.24-3.21 (m, 1H), 2.80-2.80 (m, 1H), 2.24-2.10 (m, 2H), 1.69-1.67 (m, 1H).; LCMS: 479.1

FIGURE 1

201		¹H-NMR (400 MHz, CDCl₃) δ ppm 8.37-8.29 (m, 1H), 7.35-7.10 (m, 3H), 5.60 (dd, 1H, J = 10.0, 8.4 Hz), 5.45 (d, 1H, J = 52.0 Hz), 4.87-4.85 (m, 1H), 4.55-4.32 (m, 2H), 4.07-3.91 (m, 2H), 2.88-2.82 (m, 1H), 2.41-2.11 (m, 3H), 1.79-1.63 (m, 2H).; LCMS: 479.1
202		¹H NMR (400 MHz, DMSO-d₆) δ 14.20 (s, 1H), 9.07 (d, J = 7.4 Hz, 1H), 8.39 (d, J = 3.0 Hz, 1H), 8.25 (d, J = 6.0 Hz, 1H), 7.84 (dd, J = 8.8, 3.0 Hz, 1H), 5.60 – 5.42 (m, 2H), 5.05 (d, J = 5.3 Hz, 1H), 4.75 (dd, J = 21.3, 14.1 Hz, 1H), 4.51 (dd, J = 39.5, 15.0 Hz, 1H), 4.09 (s, 2H), 3.71 – 3.62 (m, 2H), 3.13 – 3.02 (m, 1H), 2.08 (s, 2H), 1.44 (dd, J = 27.0, 9.2 Hz, 2H), 1.21 (d, J = 13.5 Hz, 1H).; LCMS: 480
203		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.49 (br.s, 1H), 8.38 (br.s, 1H), 8.33 (br.s, 1H), 7.69 (d, 1H, J = 9.2 Hz), 5.97-5.77 (m, 1H), 5.64-5.39 (m, 2H), 4.57-4.44 (m, 1H), 4.24-4.21 (m, 2H), 4.07 (d, 1H, J = 6.4 Hz), 2.89-2.85 (m, 1H), 2.46-2.44 (m, 1H), 2.27-2.25 (m, 1H), 2.15-2.12 (m, 1H), 1.92-1.81 (m, 2H), 1.66-1.62 (m, 1H).; LCMS: 480.1
204		¹H NMR (400 MHz, DMSO-d₆) δ 14.19 (s, 1H), 9.13 (d, J = 7.8 Hz, 1H), 8.26 (s, 1H), 7.47 – 7.34 (m, 1H), 7.14 (d, J = 8.1 Hz, 1H), 5.59 – 5.48 (m, 2H), 4.80 – 4.60 (m, 1H), 4.41 (dd, J = 39.5, 14.0 Hz, 1H), 4.09 – 3.96 (m, 1H), 3.92 (dd, J = 11.1, 3.7 Hz, 1H), 3.75 (dd, J = 12.0, 5.6 Hz, 1H), 3.47 (dtd, J = 59.1, 8.2, 7.8, 4.5 Hz, 2H), 3.19 (dd, J = 11.2, 6.9 Hz, 1H), 2.85 – 2.60 (m, 1H), 2.33 – 2.04 (m, 1H), 1.79 (dd, J = 11.3, 4.7 Hz, 1H), 1.53 – 1.33 (m, 1H).; LCMS: 481

FIGURE 1

205		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37-9.30 (m, 1H), 7.08-6.98 (m, 2H), 5.68-5.63 (m, 1H), 5.53-5.36 (m, 1H), 4.53-4.40 (m, 1H), 4.02-3.91 (m, 4H), 3.55-3.46 (m, 2H), 3.26-3.21 (m, 1H), 2.86-2.82 (m, 1H), 2.30-2.12 (m, 2H), 1.78-1.70 (m, 1H).; LCMS: 481.1
206		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (s, 1H), 7.10-6.95 (m, 2H), 5.72-5.62 (m, 1H), 5.52-5.39 (m, 1H), 4.93-4.92 (m, 1H), 4.52-4.08 (m, 2H), 4.07-5.06 (m, 1H), 3.93-3.90 (m, 1H), 2.88-2.82 (m, 1H), 2.41-2.10 (m, 3H), 1.78-1.63 (m, 2H).; LCMS: 481.1
207		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.25-8.20 (m, 2H), 7.61-7.58 (m, 1H), 5.66-5.62 (m, 1H), 5.25 (d, 1H, J = 52.0 Hz), 5.14-5.00 (m, 1H), 4.91-4.87 (m, 1H), 4.53-4.44 (m, 2H), 4.06-4.03 (m, 1H), 2.94-2.91 (m, 1H), 2.54-2.43 (m, 2H), 1.91-1.83 (m, 3H).; LCMS: 482.1
208		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26-8.20 (m, 1H), 7.08-7.02 (m, 1H), 6.94-6.92 (m, 1H), 5.64-5.60 (m, 1H), 5.42 (d, 1H, J = 52.0 Hz), 5.10 (dt, 1H, J = 52.0, 6.0 Hz), 4.80 (s, 1H), 4.55-4.44 (m, 2H), 4.09-4.04 (m, 1H), 2.92-2.54 (m, 1H), 2.52-2.43 (m, 2H), 1.93-1.89 (m, 1H), 1.86-1.83 (m, 2H).; LCMS: 483.1

FIGURE 1





209		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.38-8.35 (m, 1H), 7.84-7.80 (m, 1H), 7.24-7.15 (m, 2H), 5.58 (s, 1H), 5.07 (s, 1H), 4.40-4.39 (m, 1H), 4.11-4.10 (m, 1H), 3.98-3.90 (m, 3H), 3.68-3.61 (m, 2.5H), 3.48-3.41 (m, 1H), 3.22 (s, 1.5H).; LCMS: 485.1
210		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.36-7.32 (m, 1H), 7.15 (d, 1H, J = 7.6 Hz), 7.07 (d, 1H, J = 9.6 Hz), 6.97-6.93 (m, 1H), 5.45-5.41 (m, 2H), 4.49-4.36 (m, 1H), 4.13-4.09 (m, 1H), 4.01-3.97 (m, 1H), 3.13-3.09 (m, 1H), 2.84-2.75 (m, 1H), 2.26-2.16 (m, 3H), 1.68-1.59 (m, 2H), 1.50-1.37 (m, 1H), 1.20-1.14 (m, 6H).; LCMS: 487.1
211		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.27 (m, 1H), 7.22 (s, 1H), 7.07 (d, 2H, J = 6.8 Hz), 5.47-5.34 (m, 2H), 4.88-4.87 (m, 1H), 4.79-4.77 (m, 1H), 4.42-4.37 (m, 2H), 4.13-4.12 (m, 1H), 3.17-3.13 (m, 1H), 2.83-2.80 (m, 1H), 2.54-2.51 (m, 1H), 2.44-2.40 (m, 1H), 2.21-2.17 (m, 1H), 1.94-1.91 (m, 1H).; LCMS: 488.1
212		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.30 (m, 1H), 7.09-6.95 (m, 2H), 5.65-5.62 (m, 1H), 5.61-5.39 (m, 1H), 4.95-4.91 (m, 1H), 4.50-4.36 (m, 2H), 4.14-4.11 (m, 1H), 3.21-3.16 (m, 1H), 2.82-2.75 (m, 1H), 2.57-2.54 (m, 1H), 2.52-2.42 (m, 1H), 2.23-2.18 (m, 2H), 2.00-1.94 (m, 1H).; LCMS: 490.0

FIGURE 1





213		¹H-NMR (400 MHz, CD₃OD) δ ppm 7.49-7.46 (m, 1H), 7.12-7.06 (m, 2H), 5.76 (t, 1H, J = 8.4 Hz), 5.47 (d, 1H, J = 52.4 Hz), 4.46-4.28 (m, 3H), 4.00 (s, 1H), 3.79 (d, 1H, J = 6.4 Hz), 2.20-2.04 (m, 1H), 1.71 (br.s, 1H), 1.50 (br.s, 1H), 1.31-1.25 (m, 2H), 0.67-0.64 (m, 1H).; LCMS: 491.1
214		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33-8.27 (m, 1H), 7.22 (s, 1H), 7.10-7.07 (m, 2H), 5.47-5.34 (m, 2H), 4.51-4.38 (m, 2H), 4.27-4.23 (m, 1H), 3.91-3.77 (m, 1H), 3.35-3.33 (m, 1H), 2.84-2.78 (m, 1H), 2.19-2.18 (m, 1H), 1.72-1.68 (m, 1H), 1.48-1.45 (m, 1H), 1.24-1.21 (m, 1H), 0.67-0.60 (m, 1H).; LCMS: 491.1
215		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.41-8.28 (m, 1H), 7.36-6.33 (m, 3H), 5.58-5.54 (m, 1H), 5.51-5.38 (m, 1H), 4.98-4.96 (m, 1H), 4.52-3.76 (m, 4H), 2.97-2.74 (m, 1H), 2.30-2.15 (m, 1H), 1.73-1.48 (m, 2H), 1.26-1.22 (m, 1H), 0.66-0.61 (m, 1H).; LCMS: 491.1
216		¹H NMR (400 MHz, DMSO-d₆) δ 14.19 (s, 1H), 9.02 (d, J = 7.6 Hz, 1H), 8.39 (d, J = 2.8 Hz, 1H), 8.29 (s, 1H), 7.89 (dd, J = 9.0, 3.0 Hz, 1H), 5.64 – 5.37 (m, 3H), 4.92 – 4.71 (m, 2H), 4.43 (d, J = 9.1 Hz, 1H), 4.30 (dd, J = 13.2, 7.4 Hz, 2H), 4.14 (q, J = 6.0 Hz, 1H), 3.48 (t, J = 5.3 Hz, 1H), 2.79 (td, J = 16.6, 16.1, 7.2 Hz, 1H), 1.43 (dd, J = 8.9, 4.7 Hz, 1H), 1.24 (d, J = 16.9 Hz, 1H), 1.02 (q, J = 4.2 Hz, 1H), 0.48 – 0.33 (m, 1H).; LCMS: 492

FIGURE 1





217		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.39-8.30 (m, 1H), 7.51-7.48 (m, 1H), 7.11-6.95 (m, 2H), 5.79 (t, 1H, J = 9.2 Hz), 5.55-5.38 (m, 1H), 4.89 (br.s, 1H), 4.62-4.49 (m, 1H), 4.17-4.07 (m, 2H), 3.57-3.55 (m, 1H), 3.04-2.91 (m, 1H), 2.17-2.10 (m, 2H), 1.93-1.85 (m, 2H), 1.67-1.54 (m, 3H).; LCMS: 493.1
218		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.28 (m, 1H), 6.95-6.86 (m, 2H), 5.66 (t, 1H, J = 8.4 Hz), 5.50-5.37 (m, 1H), 4.52-4.43 (m, 1H), 4.03-3.91 (m, 7H), 3.54-3.49 (m, 2H), 3.26-3.21 (m, 1H), 2.86-2.75 (m, 1H), 2.24-2.11 (m, 2H), 1.76-1.71 (m, 1H).; LCMS: 493.1
219		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.30 (m, 1H), 6.93-6.85 (m, 2H), 5.66 (t, 1H, J = 8.8 Hz), 5.50-5.37 (m, 1H), 4.52-4.31 (m, 2H), 4.07-4.04 (m, 2H), 3.98 (s, 3H), 3.93-3.90 (m, 1H), 2.79-2.75 (m, 1H), 2.43-2.39 (m, 1H), 2.20-2.07 (m, 2H), 1.79-1.77 (m, 1H), 1.64-1.62 (m, 1H).; LCMS: 493.1
220		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37 (s, 1H), 7.10-6.98 (m, 2H), 5.63 (t, 1H, J = 8.4 Hz), 5.54-5.36 (m, 1H), 4.98-4.95 (m, 1H), 4.50-4.37 (m, 2H), 4.29-4.25 (m, 1H), 3.80-3.78 (m, 1H), 2.89-2.79 (m, 1H), 2.31-2.15 (m, 1H), 1.78-1.72 (m, 1H), 1.56-1.50 (m, 1H), 1.30-1.25 (m, 1H), 0.68-0.62 (m, 1H).; LCMS: 493.1

FIGURE 1





221		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.30-8.22 (m, 1H), 7.33-6.35 (m, 3H), 5.85-5.56 (m, 1H), 5.49-5.36 (m, 1H), 4.91-4.88 (m, 1H), 4.50-4.40 (m, 1H), 4.15-4.08 (m, 1H), 3.99 (br.s, 1H), 3.52-3.49 (m, 1H), 2.82-2.80 (m, 1H), 2.24-2.03 (m, 2H), 1.86-1.78 (m, 2H), 1.58-1.47 (m, 3H).; LCMS: 493.1
222		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.38 (s, 1H), 7.49-7.45 (m, 1H), 7.10-7.03 (m, 2H), 5.77 (d, 1H, J = 8.4 Hz), 5.44 (d, 1H, J = 51.6 Hz), 4.59-4.49 (m, 1H), 4.38-4.36 (m, 1H), 4.09-3.61 (m, 6H), 3.42-3.39 (m, 1H), 2.99-2.90 (m, 0.5H), 2.16-2.01 (m, 0.5H).; LCMS: 495.1
223		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.28 (m, 1H), 7.21 (s, 1H), 7.11-7.06 (m, 2H), 5.46-5.33 (m, 2H), 4.84-4.80 (m, 1H), 4.55-4.51 (m, 1H), 4.28-4.27 (m, 1H), 4.05-4.02 (m, 1H), 3.86-3.83 (m, 2H), 3.76-3.73 (m, 1H), 3.58-3.56 (m, 1H), 3.40-3.38 (m, 1H), 2.84-2.78 (m, 1H), 2.19-2.05 (m, 1H).; LCMS: 495.1
224		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33-8.32 (m, 1H), 7.09-7.03 (m, 1H), 6.95-6.93 (m, 1H), 5.65-5.61 (m, 1H), 5.51-5.38 (m, 1H), 4.52-4.42 (m, 1H), 4.10-4.02 (m, 3H), 3.51-3.48 (m, 1H), 2.84-2.82 (m, 1H), 2.24-2.04 (m, 2H), 1.87-1.79 (m, 2H), 1.62-1.48 (m, 3H). ; LCMS: 495.2

FIGURE 1





225		N/A; LCMS: 497.1
226		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34-8.27 (m, 1H), 7.15-7.02 (m, 3H), 5.99-5.62 (m, 1H), 5.60-5.58 (m, 1H), 5.51-5.38 (m, 1H), 4.96-4.90 (m, 1H), 4.49-4.40 (m, 1H), 4.23-4.18 (m, 1H), 4.12-4.09 (m, 1H), 2.87-2.79 (m, 1H), 2.58-2.50 (m, 1H), 2.30-2.23 (m, 2H), 2.15-2.11 (m, 1H), 1.87-1.77 (m, 1H), 1.67-1.64 (m, 1H).; LCMS: 497.1
227		N/A; LCMS: 499.1
228		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (d, 1H, J = 8.4 Hz), 7.40 (br.s, 1H), 7.20-7.14 (m, 2H), 5.50 (s, 1H), 4.94 (d, 1H, J = 7.2 Hz), 4.40-4.38 (m, 1H), 4.11-4.10 (m, 1H), 3.97-3.85 (m, 3H), 3.68-3.61 (m, 2.5H), 3.47-3.40 (m, 1H), 3.17 (s, 1.5H).; LCMS: 501.1

FIGURE 1





229		N/A; LCMS: 504.1
230		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.19 (s, 1H), 7.59 (s, 1H), 7.48-7.41 (m, 2H), 5.47-5.30 (m, 2H), 4.78-4.75 (m, 1H), 4.53-4.43 (m, 1H), 4.34-4.30 (m, 1H), 3.75-3.70 (m, 1H), 2.84-2.79 (m, 1H), 2.18-2.07 (m, 3H), 1.75-1.49 (m, 4H).; LCMS: 504.1
231		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.45 (s, 1H), 8.33 (d, 1H, J = 2.8 Hz), 8.21 (s, 1H), 7.64 (d, 1H, J = 9.6 Hz), 5.51-5.34 (m, 2H), 4.82-4.79 (m, 1H), 4.53-4.40 (m, 1H), 4.33 (br.s, 1H), 3.74-3.66 (m, 1H), 2.84-2.80 (m, 1H), 2.21-2.08 (m, 3H), 1.78-1.71 (m, 2H), 1.61-1.46 (m, 2H).; LCMS: 504.1
232		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.22-8.15 (m, 1H), 7.13-6.83 (m, 3H), 5.61-5.56 (m, 1H), 5.41 (d, 1H, J = 52.0 Hz), 4.91-4.89 (m, 1H), 4.46-4.36 (m, 1H), 4.21-4.20 (m, 1H), 4.10-3.97 (m, 1H), 2.85-2.79 (m, 1H), 2.29-2.00 (m, 4H), 1.63-1.58 (m, 2H), 1.22-1.18 (m, 6H); LCMS: 505.2

FIGURE 1

233		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.30-8.24 (m, 1H), 6.94-6.83 (m, 2H), 5.64 (t, 1H, J = 8.8 Hz), 5.41 (d, 1H, J = 52.0 Hz), 4.48-4.29 (m, 4H), 4.00 (s, 3H), 3.74 (d, 1H, J = 6.0 Hz), 2.79-2.71 (m, 1H), 2.20-2.06 (m, 2H), 1.65-1.60 (m, 1H), 1.47-1.45 (m, 1H), 1.23-1.20 (m, 1H), 0.61-0.55 (m, 1H).; LCMS: 505.3
234		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33-8.27 (m, 1H), 6.95-6.84 (m, 2H), 5.65 (t, 1H, J = 8.8 Hz), 5.50-5.31 (m, 1H), 4.52-4.39 (m, 1H), 4.11-4.08 (m, 1H), 4.03 (s, 1H), 3.98 (s, 3H), 3.73-3.50 (m, 2H), 2.84-2.74 (m, 1H), 2.20-2.06 (m, 2H), 1.86-1.81 (m, 2H), 1.61-1.47 (m, 3H).; LCMS: 507.1
235		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37-8.30 (m, 1H), 7.04-6.95 (m, 2H), 6.85-6.67 (m, 2H), 6.15-6.13 (m, 1H), 5.99-5.97 (m, 1H), 5.46-5.33 (m, 2H), 4.98 (d, 1H, J = 4.4 Hz), 4.80 (s, 1H), 4.53-4.42 (m, 2H), 4.00-3.97 (m, 1H), 2.85-2.75 (m, 1H), 2.19-2.02 (m, 1H).; LCMS: 509
236		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.30 (m, 1H), 6.94-6.84 (m, 2H), 5.67 (t, 1H, J = 8.4 Hz), 5.50-5.31 (m, 1H), 4.52-4.42 (m, 1H), 4.31-4.30 (m, 1H), 4.07-3.89 (m, 5H), 3.83-3.76 (m, 3H), 3.58-3.40 (m, 2H), 2.81-2.75 (m, 1H), 2.22-2.03 (m, 1H).; LCMS: 509.1

FIGURE 1

237		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.27 (m, 1H), 7.05-6.96 (m, 2H), 6.86-6.68 (m, 2H), 5.47-5.34 (m, 2H), 4.81 (s, 1H), 4.54-4.49 (m, 1H), 4.31-4.29 (m, 1H), 4.06-4.04 (m, 1H), 3.91-3.88 (m, 1H), 2.86-2.83 (m, 1H), 2.40-2.36 (m, 1H), 2.17-2.06 (m, 2H), 1.76-1.62 (m, 2H).; LCMS: 511
238		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32 (s, 1H), 7.06-6.97 (m, 2H), 6.87-6.69 (m, 2H), 5.47-5.34 (m, 2H), 4.54-4.50 (m, 1H), 4.01-3.88 (m, 3H), 3.53-3.46 (m, 3H), 3.23-3.18 (m, 1H), 2.83-2.81 (m, 1H), 2.17-2.08 (m, 2H), 1.74-1.65 (m, 1H).; LCMS: 511.1
239		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.27 (m, 1H), 7.05-6.97 (m, 2H), 6.87-6.68 (m, 2H), 5.47-5.34 (m, 2H), 4.89 (s, 1H), 4.62-4.47 (m, 1H), 4.40-4.37 (m, 1H), 4.27-4.23 (m, 1H), 3.77 (d, 1H, J = 6.0 Hz), 2.86-2.80 (m, 1H), 2.22-2.18 (m, 1H), 1.71-1.68 (m, 1H), 1.48-1.46 (m, 1H), 1.27-1.20 (m, 1H), 0.65-0.59 (m, 1H).; LCMS: 523
240		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.23 (s, 1H), 7.36 (br.s, 1H), 7.16-7.08 (m, 2H), 5.65-5.61 (m, 1H), 5.41 (d, 1H, J = 52.0 Hz), 4.80 (s, 1H), 4.57-4.38 (m, 2H), 4.14-4.05 (m, 1H), 3.82-3.79 (m, 1H), 2.80-2.76 (m, 1H), 2.38-2.34 (m, 1H), 2.15-2.01 (m, 2H), 1.74-1.71 (m, 1H), 1.52-1.49 (m, 1H).; LCMS: 529.1

FIGURE 1

241		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.22 (s, 1H), 7.36 (br.s, 1H), 7.17-7.15 (m, 1H), 7.11-7.08 (m, 1H), 5.67-5.62 (m, 1H), 5.41 (d, 1H, J = 52.0 Hz), 4.91 (s, 1H), 4.51-4.38 (m, 1H), 4.20 (br.s, 1H), 3.86-3.83 (m, 2H), 3.54-3.49 (m, 2H), 3.27-3.22 (m, 1H), 2.82-2.73 (m, 1H), 2.19-2.03 (m, 2H), 1.63-1.56 (m, 1H).; LCMS: 529.1
242		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 7.38 (br.s, 1H), 7.18-7.10 (m, 2H), 5.65-5.61 (m, 1H), 5.46 (d, 1H, J = 52.0 Hz), 4.98-4.92 (m, 1H), 4.64-4.53 (m, 1H), 4.49-4.40 (m, 1H), 4.37-4.26 (m, 1H), 3.73 (d, 1H, J = 5.6 Hz), 2.82-2.73 (m, 1H), 2.24-2.07 (m, 1H), 1.67 (br.s, 1H), 1.47 (br.s, 1H), 1.29-1.23 (m, 1H), 0.68-0.59 (m, 1H).; LCMS: 541.2
243		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.19 (s, 1H), 7.35-7.34 (m, 1H), 7.16-7.07 (m, 2H), 5.63-5.59 (m, 1H), 5.40 (d, 1H, J = 52.4 Hz), 4.86-4.80 (m, 1H), 4.49-4.41 (m, 2H), 3.81 (br.s, 1H), 3.56-3.54 (m, 1H), 2.81-2.75 (m, 1H), 2.15-1.97 (m, 2H), 1.77-1.71 (m, 2H), 1.52-1.47 (m, 2H), 1.47-1.27 (m, 1H).; LCMS: 543.1
244		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.23 (s, 1H), 7.35 (br.s, 1H), 7.16-7.07 (m, 2H), 5.67-5.62 (m, 1H), 5.40 (d, 1H, J = 52.0 Hz), 4.77 (s, 1H), 4.54-4.40 (m, 2H), 4.01-3.86 (m, 2H), 3.79-3.72 (m, 3H), 3.57-3.54 (m, 1H), 2.82-2.72 (m, 1H), 2.22-2.12 (m, 1H).; LCMS: 545.1

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/048698A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D487/04 A61K31/519 A61P35/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TRAXLER P ET AL: "USE OF A PHARMACOPHORE MODEL FOR THE DESIGN OF EGF-R TYROSINE KINASE INHIBITORS: 4-(PHENYLAMINO)PYRAZOLO not 3,4-D 3/4 PYRIMIDINES", JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, US, vol. 40, no. 22, 1 January 1997 (1997-01-01), pages 3601-3616, XP001009913, ISSN: 0022-2623, DOI: 10.1021/JM970124V	1,4-6, 9-12,14, 17,21-24
A	page 3604; table 1; compounds 7-32 ----- -/--	2,3,7,8, 13,15, 16,18-20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

7 November 2016

Date of mailing of the international search report

18/11/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Bedel, Christian

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2016/048698

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	GB 2 515 785 A (REDX PHARMA LTD [GB]) 7 January 2015 (2015-01-07) page 48 - page 52; claim 22; example 4; table 3	1,2,4-7, 9-14,21 3,8, 15-20, 22-24
X A	----- WO 2005/117909 A2 (EXELIXIS INC [US]; ANAND NEEL K [US]; BLAZEY CHARLES M) 15 December 2005 (2005-12-15) page 196 - page 197; compounds 356-359,373,391,394,397,398,100,401	1,2,4-6, 11-14,21 3,7-10, 15-20, 22-24
X A	----- WO 2014/118226 A1 (BAYER PHARMA AG [DE]) 7 August 2014 (2014-08-07) examples	1,4-14, 21 2,3, 15-20
X A A	----- WO 2011/139273 A1 (VERTEX PHARMA [US]; JIMENEZ JUAN-MIGUEL [GB]; MILLER ANDREW [GB]) 10 November 2011 (2011-11-10) page 40; table 1; compound 1 page 46; example 1 ----- WO 2014/036387 A2 (UNIV COLORADO [US]) 6 March 2014 (2014-03-06) page 2 - page 4 page 7 - page 8; figures 23,25 page 12, line 13 - line 27; claims 12,26 ----- WO 2009/032694 A1 (DANA FARBER CANCER INST INC [US]; GRAY NATHANAEL [US]; SIM TAEBO [US];) 12 March 2009 (2009-03-12) page 86 - page 87; table 4 -----	1,21 2-20, 22-24 1-24 1-24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/048698

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2515785	A	07-01-2015	NONE
WO 2005117909	A2	15-12-2005	AU 2005249380 A1 15-12-2005
		CA 2563699 A1 15-12-2005	
		EP 1750727 A2 14-02-2007	
		JP 5213229 B2 19-06-2013	
		JP 2007534687 A 29-11-2007	
		US 2008076774 A1 27-03-2008	
		WO 2005117909 A2 15-12-2005	
WO 2014118226	A1	07-08-2014	CA 2899662 A1 07-08-2014
		CN 105377845 A 02-03-2016	
		EP 2951175 A1 09-12-2015	
		JP 2016506944 A 07-03-2016	
		US 2016002245 A1 07-01-2016	
		WO 2014118226 A1 07-08-2014	
WO 2011139273	A1	10-11-2011	AU 2010352577 A1 19-01-2012
		CA 2761074 A1 10-11-2011	
		CN 102471342 A 23-05-2012	
		EP 2566866 A1 13-03-2013	
		JP 2013525481 A 20-06-2013	
		WO 2011139273 A1 10-11-2011	
WO 2014036387	A2	06-03-2014	CA 2882759 A1 06-03-2014
		EP 2890815 A2 08-07-2015	
		JP 2015528295 A 28-09-2015	
		US 2015218652 A1 06-08-2015	
		WO 2014036387 A2 06-03-2014	
WO 2009032694	A1	12-03-2009	AU 2008296479 A1 12-03-2009
		WO 2009032694 A1 12-03-2009	
		WO 2009032703 A1 12-03-2009	



(12)发明专利申请

(10)申请公布号 CN 108137607 A

(43)申请公布日 2018.06.08

(21)申请号 201680060799.0

尼尔·比富尔科

(22)申请日 2016.08.25

(74)专利代理机构 永新专利商标代理有限公司

72002

(30)优先权数据

62/210,264 2015.08.26 US

代理人 过晓东

(85)PCT国际申请进入国家阶段日

2018.04.17

(51)Int.Cl.

C07D 487/04(2006.01)

A61K 31/519(2006.01)

A61P 35/00(2006.01)

(86)PCT国际申请的申请数据

PCT/US2016/048698 2016.08.25

(87)PCT国际申请的公布数据

W02017/035354 EN 2017.03.02

(71)申请人 蓝图药品公司

地址 美国马萨诸塞州

(72)发明人 史蒂文·M·翁劳斯基

娜塔莎·布鲁杰孟

昌德拉塞卡·V·曼德拉

权利要求书3页 说明书106页 附图61页

(54)发明名称


适用于治疗与NTRK相关的病症的化合物和组合物

(57)摘要

本发明涉及对野生型NTRK及其抗性突变体具有活性的NTRK抑制剂。

序	结构	NMR, LC/MS
1		¹ H-NMR (500 MHz, CDCl ₃) δ ppm: 8.54 (s, 1H), 7.03-7.02 (m, 3H), 6.96 (s, 2H), 4.87 (d, 2H, J = 7.2 Hz), 3.92-3.20 (m, 4H), 3.11-3.08 (m, 1H), 1.50-1.87 (m, 2H), 1.36-1.74 (m, 3H); LC/MS: 380.1
2		¹ H-NMR (500 MHz, CDCl ₃) δ ppm: 8.23 (s, 1H), 7.13-7.08 (m, 3H), 7.06-7.01 (m, 3H), 6.99-6.88 (m, 2H), 4.55 (s, 2H), 4.26-4.21 (m, 2H), 4.13-4.11 (m, 1H), 4.09-4.06 (m, 1H), 2.44-2.41 (m, 1H), 2.35 (s, 3H), 2.16-2.13 (m, 1H), 1.25-1.20 (m, 2H); LC/MS: 401.1
3		¹ H-NMR (500 MHz, CDCl ₃) δ ppm: 8.23 (s, 1H), 7.13-7.08 (m, 3H), 7.06-7.01 (m, 3H), 6.99-6.88 (m, 2H), 4.41-4.32 (m, 1H), 4.05-3.96 (m, 2H), 2.41-2.36 (m, 1H), 1.99-1.89 (m, 1H), 1.86-1.86 (m, 1H), 1.44-1.41 (m, 1H), 0.26-0.23 (m, 1H), 0.20-0.17 (m, 1H); LC/MS: 401.1
4		¹ H-NMR (500 MHz, CDCl ₃) δ ppm: 8.12 (s, 1H), 7.08-7.05 (m, 3H), 7.05-6.92 (m, 3H), 6.28 (s, 2H), 4.32-4.29 (m, 1H), 3.93-3.87 (m, 1H), 2.33-2.28 (m, 1H), 1.51-1.50 (m, 1H), 1.49-1.47 (m, 1H), 1.35-1.33 (m, 1H), 0.87-0.84 (m, 1H), 0.41-0.40 (m, 1H); LC/MS: 401.1

1. 一种式 (I) 化合物:



或其药学上可接受的盐, 其中:

环A和B各自独立地选自芳基、杂芳基、环烷基和杂环基;

每个 L^1 和 L^2 独立地选自键、 $-C(O)-$ 、 $-N(R^1)-$ 、 $-N(R^1)-C(O)-$ 、 $-C(O)-N(R^1)-$ 、 $-(C_1-C_6\text{亚烷基})-N(R^1)-$ 、 $-N(R^1)-(C_1-C_6\text{亚烷基})-$ 、 $-N(R^1)-C(O)-(C_1-C_6\text{亚烷基})-$ 以及 $-C(O)-N(R^1)-(C_1-C_6\text{亚烷基})-$; 其中 L^1 和 L^2 的每个亚烷基部分独立地以0-5次出现的 R' 取代;

每个 R^A 和 R^B 独立地选自羟基、 C_1-C_6 烷基、 C_2-C_6 烯基、 C_2-C_6 炔基、 C_1-C_6 烷氧基、卤基、 C_1-C_6 杂烷基、 C_1-C_6 卤烷基、 C_1-C_6 卤烷氧基、 C_1-C_6 羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基、杂环烷基、硝基、氰基、 $-C(O)R^1$ 、 $-OC(O)R^1$ 、 $-C(O)OR^1$ 、 $-(C_1-C_6\text{亚烷基})-C(O)R^1$ 、 $-SR^1$ 、 $-S(O)_2R^1$ 、 $-S(O)_2-N(R^1)(R^1)$ 、 $-(C_1-C_6\text{亚烷基})-S(O)_2R^1$ 、 $-(C_1-C_6\text{亚烷基})-S(O)_2-N(R^1)(R^1)$ 、 $-N(R^1)(R^1)$ 、 $-C(O)-N(R^1)(R^1)$ 、 $-N(R^1)-C(O)R^1$ 、 $-N(R^1)-C(O)OR^1$ 、 $-(C_1-C_6\text{亚烷基})-N(R^1)-C(O)R^1$ 、 $-N(R^1)S(O)_2R^1$ 以及 $-P(O)(R^1)(R^1)$; 其中 R^A 和 R^B 中的每一者的每个烷基、烯基、炔基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基以及杂环烷基部分独立地以0-5次出现的 R^a 取代; 或者2个 R^A 或2个 R^B 与其所连接的碳原子一起形成以0-5次出现的 R^a 取代的环烷基或杂环基环;

每个 R^1 独立地选自氢、羟基、卤基、硫醇、 C_1-C_6 烷基、 C_1-C_6 硫烷基、 C_1-C_6 烷氧基、 C_1-C_6 卤烷基、 C_1-C_6 羟烷基、 C_1-C_6 杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基, 其中 R^1 的每个烷基、硫烷基、烷氧基、卤烷基、羟烷基、杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基部分独立地以0-5次出现的 R^b 取代, 或者2个 R^1 与其所连接的原子一起形成独立地以0-5次出现的 R^b 取代的环烷基或杂环基环;

每个 R^a 和 R^b 独立地选自 C_1-C_6 烷基、卤基、羟基、 C_1-C_6 卤烷基、 C_1-C_6 杂烷基、 C_1-C_6 羟烷基、 C_1-C_6 烷氧基、环烷基、杂环基以及氰基, 其中 R^a 和 R^b 的每个烷基、卤烷基、杂烷基、羟烷基、烷氧基、环烷基以及杂环基部分独立地以0-5次出现的 R' 取代;

每个 R' 独立地选自 C_1-C_6 烷基、 C_1-C_6 杂烷基、卤基、羟基、 C_1-C_6 卤烷基、 C_1-C_6 羟烷基、环烷基以及氰基; 或者2个 R' 与其所连接的原子一起形成环烷基或杂环基环; 并且

p 为0、1、2、3、4或5; 并且

q 为0、1、2、3或4。

2. 如权利要求1所述的化合物, 其中环A选自环烷基、杂环基和环烯基。

3. 如权利要求2所述的化合物, 其中环A选自环戊基、环己基、四氢吡喃基、四氢呋喃基、吡咯烷基以及环戊烯基。

4. 如权利要求1至3中任一项所述的化合物, 其中环B选自芳基、杂芳基或杂环基。


5. 如权利要求4所述的化合物, 其中环B选自苯基、吡啶基或吡咯烷基。

6. 如权利要求1至3中任一项所述的化合物, 其中每个 L^1 为键、 $-C(O)-$ 或 $-N(R^1)-$; 并且 L^2

为-N(R¹)-C(O)-(C₁-C₆亚烷基)-或-C(O)-N(R¹)-(C₁-C₆亚烷基)-。

7. 如权利要求6所述的化合物,其中:

L¹为-NH-;并且

L²选自-C(O)-NH-CH(CH₂OH)-*, -C(O)-N(CH₃)-CH₂*, -C(O)-N(CH₃)-CH(CH₃)-*, -C(O)N(CH₂CH₃)CH₂*, -C(O)NHCH(CH₃)-*, -C(O)N(CD₃)CH₂*, -C(O)NHCH(CF₃)-*以及,

其中“*”表示结合至环B的L²的一部分。

8. 如权利要求7所述的化合物,其中L¹为-NH-, L²为-C(O)-并且环B为吡咯烷-1-基。

9. 如权利要求1至8中任一项所述的化合物,其中每个R¹独立地选自氢和以0-5次出现的R^b取代的C₁-C₆烷基。

10. 如权利要求9所述的化合物,其中每个R¹独立地选自氢和-CH₃。

11. 如权利要求1至10中任一项所述的化合物,其中每个R^A和R^B独立地选自羟基、C₁-C₆烷基、C₁-C₆烷氧基、卤基、C₁-C₆杂烷基、C₁-C₆卤烷基、C₁-C₆卤烷氧基、C₁-C₆羟烷基、环烷基、芳基、杂芳基、硝基、氰基、-C(O)R¹、-OC(O)R¹、-C(O)OR¹、-SR¹、-S(O)₂R¹、-S(O)₂-N(R¹)(R¹)、-N(R¹)(R¹)、-C(O)-N(R¹)(R¹)、-N(R¹)-C(O)R¹、-N(R¹)-C(O)OR¹以及-N(R¹)S(O)₂R¹;其中每个R^a和每个R^b的每个烷基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基以及杂芳基部分独立地以0-5次出现的R^a取代;或者2个R^A或2个R^B与其所连接的碳原子一起形成独立地以0-5次出现的R^a取代的环烷基或杂环基环。

12. 如权利要求1至6中任一项所述的化合物,其中每个R^A独立地选自羟基、C₁-C₆烷基、C₁-C₆烷氧基、卤基、-C(O)-N(R¹)(R¹)、-C(O)OR¹、-S(O)₂R¹、C₁-C₆卤烷基、-CN、氧杂环丁烷基以及C₁-C₆羟烷基,或者结合至环A上的相邻环碳原子的两个R^A结合在一起以形成稠合至环A的C₃-C₆环烷基。

13. 如权利要求12所述的化合物,其中每个R^A独立地选自羟基、氟代、氧杂环丁烷-3-基、-CHF₂、-CH₂CH₃、-C(CH₃)₂OH、-OCH₃、-C(O)N(CH₃)₂、-C(O)OCH₃、-S(O)₂CH₃;或者结合至环A上的相邻环碳原子的两个R^A结合在一起以形成稠合至环A的环丙基。


14. 如权利要求1至13中任一项所述的化合物,其中每个R^B独立地选自卤基、C₁-C₆烷基、氰基、C₁-C₆烷氧基、芳基、杂芳基、氧基以及C₁-C₆卤烷氧基。

15. 如权利要求1至14中任一项所述的化合物,其中环B为吡咯烷基并且至少一个R^B为任选取代的芳基或杂芳基。

16. 如权利要求15所述的化合物,其中环B为吡咯烷基,并且至少一个R^B选自2,3,5-三氟苯基、2,3-二氟苯基、2,5-二氟苯基、2-氯-5-氟苯基、2-氯-5-氟吡啶-3-基、2-氰基-5-氟苯基、2-氟-5-氯苯基、2-甲氧基-3,5-二氟苯基、2-甲氧基-5-氟吡啶-3-基、2-三氟甲氧基-5-氟苯基、3,5-二氟苯基、3-氯-5-氟苯基、3-氰基-5-氟苯基、3-二氟甲氧基-5-氟苯基、3-氟苯基、5-氟吡啶-3-基以及苯基。

17. 如权利要求1至14中任一项所述的化合物,其中环B不是吡咯烷基,并且每个R^B独立地选自氯代、氟代、氧基、-CH₃、-CF₃、-CN、-OCH₃、-OCF₃以及-OCHF₂。

18. 如权利要求1所述的化合物,其中所述化合物为式(Ia)化合物:




或其药学上可接受的盐,其中环A、 R^A 、 R^B 、p和q

(Ia),

中的每一者如权利要求1中定义。

19. 如权利要求1所述的化合物,其中所述化合物为式(II)化合物:



或其药学上可接受的盐,其中:

(II)

R^{1a} 选自氢、 C_1 - C_6 烷基和氘化 C_1 - C_6 烷基;

R^{1b} 选自氢和 C_1 - C_6 烷基;并且

环A、环B、 R' 、 R^A 、 R^B 、p和q为如权利要求1所定义的。

20. 一种选自表1中列出的化合物中的任一种的化合物,或其药学上可接受的盐。

21. 一种药物组合物,其包含药学上可接受的载剂和如权利要求1至20中任一项所述的化合物。

22. 如权利要求21所述的组合物,其用于治疗患有由异常神经营养酪氨酸受体激酶(NTRK)活性介导的病状的受试者。

23. 如权利要求21所述的组合物,其用于治疗已对癌症治疗发展抗性的受试者。

24. 如权利要求21所述的组合物,其用于治疗选自由以下组成的组的病状:非小细胞肺癌、乳腺癌、黑色素瘤、低级和高级神经胶质瘤、胶质母细胞瘤、小儿科星细胞瘤、结直肠癌、乳突甲状腺癌、胰腺癌、头颈部癌、胆管癌、急性髓性白血病、分泌性乳腺癌、唾液癌以及spitz样肿瘤。

适用于治疗与NTRK相关的病症的化合物和组合物

[0001] 优先权要求

[0002] 本申请要求2015年8月26日提交的U.S.S.N.62/210,264的优先权,所述专利以引用的方式整体并入本文。


[0003] 先前技术

[0004] 神经营养酪氨酸受体激酶(NTRK)1、2和3为活化参与细胞增殖和存活的多个下游途径的受体酪氨酸激酶(RTK)。由编码这些RTK的基因的异常染色体易位引起的各种基因融合涉及多种癌症的病因,包括高级别和低级别神经胶质瘤、胆管癌、乳突甲状腺癌、结肠癌和非小细胞肺癌。对于激酶融合概貌的基因组学分析识别广泛多种额外癌症类型中的NTRK融合物,包括头颈部鳞状细胞癌、胰腺癌、肉瘤和黑素瘤,从而提供用于部署这些激酶的抑制剂以治疗多个肿瘤学适应症的进一步治疗理论根据。

[0005] 将NTRK融合物识别为某些癌症的潜在病因推进多种NTRK激酶抑制剂的发现和临床研发,以便治疗具有NTRK融合蛋白质的肿瘤。早期临床数据支持此方法在为患有特定人类恶性肿瘤的患者提供益处方面的可行性。然而,最终,尽管存在临床活性的明显迹象,大多数患者的癌症将变得对于激酶抑制剂疗法有抵抗力,导致疾病的复发和进展。通过内在突变的激酶再活化为抗性的常见机制。当发生抗性时,患者的治疗选项往往非常有限。因而需要抑制NTRK,以及其抗性突变体的化合物。

发明内容

[0006] 本发明提供包含式(I)化合物或其药学上可接受的盐的化合物和药物组合物,其中:



[0007]

式(I)

[0008] 环A和B各自独立地选自芳基、杂芳基、环烷基和杂环基;

[0009] 每个L¹和L²独立地选自键、-C(O)-、-N(R¹)-、-N(R¹)-C(O)-、-C(O)-N(R¹)-、-(C₁-C₆亚烷基)-N(R¹)-、-N(R¹)-(C₁-C₆亚烷基)-、-N(R¹)-C(O)-(C₁-C₆亚烷基)-以及-C(O)-N(R¹)-(C₁-C₆亚烷基)-;其中每个亚烷基独立地以0-5次出现的R'取代;

[0010] 每个R^A和R^B独立地选自羟基、C₁-C₆烷基、C₂-C₆烯基、C₂-C₆炔基、C₁-C₆烷氧基、卤基、C₁-C₆杂烷基、C₁-C₆卤烷基、C₁-C₆卤烷氧基、C₁-C₆羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基、杂环烷基、硝基、氰基、-C(O)R¹、-OC(O)R¹、-C(O)OR¹、-(C₁-C₆亚烷基)-C(O)R¹、-SR¹、-S(O)₂R¹、-S(O)₂-N(R¹)(R¹)、-(C₁-C₆亚烷基)-S(O)₂R¹、-(C₁-C₆亚烷基)-S(O)₂-N(R¹)(R¹)、-N(R¹)(R¹)、-C(O)-N(R¹)(R¹)、-N(R¹)-C(O)R¹、-N(R¹)-C(O)OR¹、-(C₁-C₆亚烷基)-N

(R¹)-C(O)R¹、-N(R¹)S(O)₂R¹以及-P(O)(R¹)(R¹)；其中烷基、烯基、炔基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的R^a取代；或2个R^a或2个R^b与其所连接的碳原子一起形成独立地被0-5次出现的R^a取代的环烷基或杂环基环；

[0011] 每个R¹独立地选自氢、羟基、卤基、硫醇、C₁-C₆烷基、C₁-C₆硫烷基、C₁-C₆烷氧基、C₁-C₆卤烷基、C₁-C₆羟烷基、C₁-C₆杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基，其中烷基、硫烷基、烷氧基、卤烷基、羟烷基、杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的R^b取代，或者2个R¹与其所连接的原子一起形成独立地以0-5次出现的R^b取代的环烷基或杂环基环；

[0012] 每个R^a和R^b独立地选自C₁-C₆烷基、卤基、羟基、C₁-C₆卤烷基、C₁-C₆杂烷基、C₁-C₆羟烷基、C₁-C₆烷氧基、环烷基、杂环基以及氰基，其中烷基、卤烷基、杂烷基、羟烷基、烷氧基、环烷基以及杂环基中的每一者独立地以0-5次出现的R'取代；

[0013] 每个R'独立地选自C₁-C₆烷基、C₁-C₆杂烷基、卤基、羟基、C₁-C₆卤烷基、C₁-C₆羟烷基、环烷基以及氰基；或者2个R'与其所连接的原子一起形成环烷基或杂环基环；

[0014] p为0、1、2、3、4或5；并且

[0015] q为0、1、2、3或4。

[0016] 本文公开的任何化合物可单独或与另一个治疗剂组合用于治疗本文公开的任何疾病。

[0017] 附图简述

[0018] 图1描绘本发明的各种示例性化合物的结构以及其如通过LC-MS确定的NMR峰和质量。

[0019] 发明详述

[0020] 定义

[0021] 如本文使用，术语“患者”、“受试者”、“个体”和“宿主”是指患有或疑似患有与异常NTRK表现(即，通过经由NTRK的信号传导所导致的NTRK活性增加)或生物活性相关的疾病或病症的人类或非人类动物。

[0022] “治疗(Treat/treating)”此疾病或病症是指改善疾病或病症的至少一个症状。这些术语在结合病状诸如癌症来使用时是指以下一种或多种：阻碍癌症生长、导致癌症重量或体积缩小、延长患者的预期存活时间、抑制肿瘤生长、减少肿瘤质量、减少转移性病灶的大小或数目、抑制新转移性病灶的发展、延长存活、延长无进展存活、延长进展时间和/或增强生活品质。

[0023] 术语“预防”在结合病状或疾病诸如癌症使用时是指降低病状或疾病的症状的频率或延迟其发作。因此，预防癌症包括例如相对于未治疗的对照群体减少接受预防性治疗的患者群体中的可检测癌性生长的数目和/或相比于未治疗的对照群体延迟所治疗群体中的可检测癌性生长的出现，例如，以在统计上和/或临床上显著的量。

[0024] 术语“治疗效果”是指通过施用本发明的化合物或组合物在动物(尤其是哺乳动物)并且更具体而言在人类中所引起的有利的局部或全身效果。短语“治疗有效量”意指本发明的化合物或组合物以合理风险比有效于治疗由NTRK的过表达或异常NTRK生物活性所造成的疾病或病状的量。此材料的治疗有效量将根据正在治疗的受试者和疾病病状、所述

受试者的体重和年龄、所述疾病病状的严重性、施用方式及其类似情况来改变,这可由本领域技术人元容易地确定。

[0025] 如本文使用,“发展抗性”意谓在药物首次施用于患者时,患者的症状改善,不论通过肿瘤体积的减少、新病灶的数目减少或医师用于判断疾病进展的一些其他手段来测量;然而,那些症状在某一时刻停止改善或甚至恶化。在此时,患者被认为对于药物显现抗性。

[0026] “脂族基团”是指直链、支链或环烷基团并且包括饱和和不饱和基团,诸如烷基、烯基和炔基。

[0027] “亚烷基”是指烷基的二价基团,例如 $-\text{CH}_2-$ 、 $-\text{CH}_2\text{CH}_2-$ 和 $\text{CH}_2\text{CH}_2\text{CH}_2-$ 。

[0028] “烯基”意指含有至少一个双键的脂族基团。

[0029] “烷氧基”意指具有与其所连接的氧基的烷基。代表性烷氧基包括甲氧基、乙氧基、丙氧基、叔丁氧基及其类似基团。术语“卤烷氧基”是指一个或多个氢原子被卤基替代的烷氧基,并包括所有氢已被卤基替代的烷氧基部分(例如,全氟烷氧基)。

[0030] “烷基”是指饱和直链或支链烃的单价基团,诸如1-12、1-10或1-6个碳原子的直链或支链基团,在本文中分别称为 C_1 - C_{12} 烷基、 C_1 - C_{10} 烷基和 C_1 - C_6 烷基。示例性烷基包括但不限于甲基、乙基、丙基、异丙基、2-甲基-1-丙基、2-甲基-2-丙基、2-甲基-1-丁基、3-甲基-1-丁基、2-甲基-3-丁基、2,2-二甲基-1-丙基、2-甲基-1-戊基、3-甲基-1-戊基、4-甲基-1-戊基、2-甲基-2-戊基、3-甲基-2-戊基、4-甲基-2-戊基、2,2-二甲基-1-丁基、3,3-二甲基-1-丁基、2-乙基-1-丁基、丁基、异丁基、叔丁基、戊基、异戊基、新戊基、己基、庚基、辛基等。

[0031] “亚烯基”是指具有两个连接点的烯基。举例而言,“亚乙烯基”表示基团 $-\text{CH}=\text{CH}-$ 。亚烯基也可为具有一个或多个取代基的未取代形式或取代形式。

[0032] “炔基”是指含有2-12个碳原子并且特征在于具有一个或多个三键的直链或支链烃链。炔基的实例包括但不限于乙炔基、炔丙基和3-己炔基。一个三键碳可任选地为炔基取代基的连接点。

[0033] “亚炔基”是指具有两个连接点的炔基。举例而言,“亚乙炔基”表示基团 $-\text{C}\equiv\text{C}-$ 。亚炔基也可为具有一个或多个取代基的未取代形式或取代形式。

[0034] “羟亚烷基”或“羟烷基”是指其中亚烷基或烷基氢原子被羟基置换的亚烷基或烷基部分。羟亚烷基或羟烷基包括其中超过一个氢原子已被羟基置换的基团。

[0035] “芳族环系统”为本领域公认的并且是指单环、双环或多环烃环系统,其中至少一个环为芳族。

[0036] “芳基”是指芳族环系统的单价基团。代表性芳基包括完全芳族环系统,诸如苯基、萘基和蒽基,以及其中芳族碳环稠合至一个或多个非芳族碳环的环系统,诸如二氢茚基、邻苯二甲酰亚胺基、萘二甲酰亚胺基或四氢萘基,及其类似基团。

[0037] “芳基烷基”或“芳烷基”是指烷基氢原子被芳基置换的烷基部分。芳烷基包括超过一个氢原子已被芳基置换的基团。“芳基烷基”或“芳烷基”的实例包括苯甲基、2-苯基乙基、3-苯基丙基、9-苊基、二苯甲基和三苯甲基。

[0038] “芳氧基”是指 $-\text{O}-$ (芳基),其中杂芳基部分为如本文定义的。

[0039] “卤基”是指任何卤素的基团,例如, $-\text{F}$ 、 $-\text{Cl}$ 、 $-\text{Br}$ 或 $-\text{I}$ 。

[0040] “卤烷基”和“卤烷氧基”是指被一个或多个卤基或其组合取代的烷基和烷氧基结构。举例而言,术语“氟烷基”和“氟烷氧基”分别包括其中卤基为氟的卤烷基和卤烷氧基。

[0041] “卤亚烷基”是指二价烷基,例如, $-\text{CH}_2-$ 、 $-\text{CH}_2\text{CH}_2-$ 和 $-\text{CH}_2\text{CH}_2\text{CH}_2-$,其中一个或多个氢原子被卤基置换,并且包括其中所有氢被卤基置换的烷基部分。

[0042] “杂烷基”意指任选地取代的烷基,其具有的一个或多个骨架链原子选自不为碳的原子,例如氧、氮、硫、磷或其组合。可给定数值范围,例如 C_1 - C_6 杂烷基是指链中的碳的数目,其在此实例中包括1至6个碳原子。举例而言, $-\text{CH}_2\text{OCH}_2\text{CH}_3$ 基团被称为“ C_3 ”杂烷基。与分子结构的其余部分的连接可通过杂烷基链中的杂原子或碳达成。“杂亚烷基”意指二价的任选地取代的烷基,其具有的一个或多个骨架链原子选自不为碳的原子,例如氧、氮、硫、磷或其组合。

[0043] “碳环环系统”是指单环、双环或多环烃环系统,其中每个环为完全饱和的或含有一个或多个不饱和单位,但是其中没有环是芳族环。

[0044] “碳环基”是指碳环环系统的单价基团。代表性碳环基团包括环烷基(例如,环戊基、环丁基、环戊基、环己基及其类似基团)以及环烯基(例如,环戊烯基、环己烯基、环戊二烯基及其类似基团)。

[0045] “环烷基”是指具有3至12个碳的环状、双环、三环或多环非芳族烃基团。任何可取代环原子皆可被取代(例如被一个或多个取代基取代)。环烷基可含有稠环或螺环。稠环为共享共同碳原子的环。环烷基部分的实例包括但不限于环丙基、环己基、甲基环己基、金刚烷基和降冰片基。在一些实施方案中,环烷基为双环[3.1.0]己基。

[0046] “环烷基烷基”是指-(环烷基)-烷基,其中环烷基和烷基为如本文公开的。“环烷基烷基”通过环烷基与母体分子结构键合。

[0047] “杂芳族环系统”为本领域公认的并且是指单环、双环或多环环系统,其中至少一个环为芳族并且包含至少一个杂原子(例如,N、O或S);并且其中没有其他环为杂环基(如以下定义)。在某些情况下,作为芳族并包含杂原子的环在此环中含有1、2、3或4个环杂原子。

[0048] “杂芳基”是指杂芳族环系统的单价基团。代表性杂芳基包括环系统,其中(i)每个环包含杂原子并且为芳族,例如,咪唑基、噁唑基、噻唑基、三唑基、吡咯基、呋喃基、噻吩基、吡唑基、吡啶基、吡嗪基、哒嗪基、嘧啶基、吡嗪基、嘌呤基、嘧啶基和蝶啶基;(ii)每个环为芳族或碳环基,至少一个芳族环包含杂原子并且至少一个其他环为烃环,或者例如吡啶基、异吡啶基、苯并噻吩基、苯并呋喃基、二苯并呋喃基、吡唑基、苯并咪唑基、苯并噻唑基、喹啉基、异喹啉基、噌啉基、酞嗪基、喹啉基、喹啉基、喹啉基、喹啉基、喹啉基、吡啶并[2,3-b]-1,4-噁嗪-3-(4H)-酮、5,6,7,8-四氢喹啉基以及5,6,7,8-四氢异喹啉基;并且(iii)每个环为芳族或碳环基,并且至少一个芳族环与另一个芳族环共享桥头杂原子,例如,4H-喹啉基。

[0049] “杂环环系统”是指单环、双环和多环系统,其中至少一个环为饱和或部分不饱和的(但并非芳族)并且包含至少一个杂原子。杂环环系统可在任何杂原子或碳原子处连接至其侧基以得到稳定结构并且任一环原子均可任选地被取代。

[0050] “杂环基”是指杂环环系统的单价基团。代表性杂环基包括环系统,其中(i)每个环为非芳族并且至少一个环包含杂原子,例如,四氢呋喃基、四氢吡喃基、四氢噻吩基、吡咯烷基、吡咯烷酮基、哌啶基、吡咯啉基、十氢喹啉基、噁唑烷基、哌嗪基、二氧杂环己烷、二氧杂环戊烷、二氮杂卓基、氧氮杂卓基、硫氮杂卓基、吗啉基和奎宁环基;(ii)至少一个环为非芳族并且包含杂原子并且至少一个其他环为芳族碳环,例如,1,2,3,4-四氢喹啉基、1,2,3,4-

四氢异喹啉基;并且(iii)至少一个环为非芳族并且包含杂原子并且至少一个其他环为芳族并且包含杂原子,例如,3,4-二氢-1H-吡喃并[4,3-c]吡啶和1,2,3,4-四氢-2,6-萘啶。

[0051] “杂环基烷基”是指被杂环基团取代的烷基。

[0052] “氰基”是指-CN基团。

[0053] “硝基”是指基团-NO₂。

[0054] “羟基(hydroxy)”或“羟基(hydroxyl)”是指基团-OH。

[0055] “羟亚烷基”是指二价烷基,例如,-CH₂-、-CH₂CH₂-和-CH₂CH₂CH₂-,其中一个或多个氢原子被羟基置换,并且包括其中所有氢被羟基置换的烷基部分。

[0056] “取代的”无论是否前置有术语“任选地”都意指指定部分的一个或多个氢被适合取代基置换。除非另外指示,否则“任选取代的”基团可在基团的每个可取代位置处具有适合取代基,并且当任何给定结构中的超过一个位置可被超过一个选自指定组的取代基取代时,在每个位置的取代基可以相同或不同。在本发明下设想的取代基的组合优选地为使得形成稳定或化学可行化合物的组合。如本文所用的术语“稳定”是指化合物在经受允许其产生、检测以及在某些实施方案中允许其回收、纯化和用于本文公开的一种或多种目的的条件时基本上不会改变。

[0057] 如本文所用,当例如烷基、m、n等的各表述在任何结构中出现超过一次时,其定义旨在独立于其在相同结构中的其他位置的定义。

[0058] 本发明的某些化合物可以特定几何或立体异构形式存在。本发明涵盖所有这些化合物,包括属于本发明范围内的顺式-和反式-异构物、R-和S-对映异构物、非对映异构物、(D)-异构物、(L)-异构物、其外消旋混合物及其他混合物。额外不对称碳原子可存在于取代基诸如烷基中。所有这些异构物以及其混合物旨在包括于本发明中。

[0059] 如果例如需要本发明的化合物的特定对映异构物,其可通过不对称合成或通过使用手性助剂衍生化来制备,其中将所得非对映异构混合物分离并且辅助基团裂解以提供纯的所需对映异构物。或者,当分子含有碱性官能基(诸如氨基)或酸性官能基(诸如羧基)时,使用合适光学活性酸或碱来形成非对映异构盐,随后将由此形成的非对映异构物通过在本领域中熟知的分级结晶或色谱方法来拆分,并且随后回收纯对映异构物。

[0060] 除非另外指示,否则当公开的化合物由未规定立体化学的结构命名或描绘并且具有一个或多个手性中心时,应理解其表示所述化合物的所有可能的立体异构物以及其对映异构混合物。可使用以下所示的等式计算组合物的“对映异构过量”或“对映异构过量%”。在以下所示的实例中,组合物含有90%的一种对映异构物(例如S-对映异构物)和10%的另一种对映异构物(即R-对映异构物)。

[0061] $ee = (90 - 10) / 100 = 80\%$ 。

[0062] 因此,认为含有90%的一种对映异构物和10%的另一种对映异构物的组合物具有80%的对映异构过量。

[0063] 本文所述的化合物或组合物可含有至少50%、75%、90%、95%或99%对映异构过量的一种形式的化合物,例如,S-对映异构物。换言之,此类组合物含有相对于R对映异构物而言对映异构过量的S对映异构物。


[0064] 本文所述的化合物也可在构成这些化合物的一个或多个原子处含有非天然比例的原子同位素。举例而言,化合物可用放射性同位素(例如像氘(²H)、氚(³H)、碘-13(¹³C)或

碳-14 (^{14}C) 进行放射性标记。本文公开的化合物的所有同位素变化形式, 无论是否具有放射性, 都旨在涵盖在本发明的范围内。另外, 本文描述的化合物的所有互变异构形式旨在处于本发明的范围内。

[0065] 化合物可呈游离碱形式呈盐形式而适用。代表性盐包括氢溴酸盐、盐酸盐、硫酸盐、硫酸氢盐、磷酸盐、硝酸盐、乙酸盐、戊酸盐、油酸盐、棕榈酸盐、硬脂酸盐、月桂酸盐、苯甲酸盐、乳酸盐、磷酸盐、甲苯磺酸盐、柠檬酸盐、马来酸盐、富马酸盐、琥珀酸盐、酒石酸盐、萘甲酸盐、甲磺酸盐、葡庚糖酸盐、乳糖醛酸盐和月桂基磺酸盐及其类似物。(参见例如 Berge 等人 (1977) "Pharmaceutical Salts", J. Pharm. Sci. 66:1-19。)

[0066] 化合物

[0067] 本发明提供式 (I) 化合物或其立体异构物、对映异构物、互变异构物或同位素标记形式, 或任何前述材料的药学上可接受的盐, 其中:



[0069]

式(I)

[0070] 环A和B各自独立地选自芳基、杂芳基、环烷基和杂环基;

[0071] 每个L¹和L²独立地选自键、-C(O)-、-N(R¹)-、-N(R¹)-C(O)-、-C(O)-N(R¹)-、-(C₁-C₆亚烷基)-N(R¹)-、-N(R¹)-(C₁-C₆亚烷基)-、-N(R¹)-C(O)-(C₁-C₆亚烷基)-以及-C(O)-N(R¹)-(C₁-C₆亚烷基)-; 其中每个亚烷基独立地以0-5次出现的R¹取代;

[0072] 每个R^A和R^B独立地选自羟基、C₁-C₆烷基、C₂-C₆烯基、C₂-C₆炔基、C₁-C₆烷氧基、卤基、C₁-C₆杂烷基、C₁-C₆卤烷基、C₁-C₆卤烷氧基、C₁-C₆羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基、杂环烷基、硝基、氰基、-C(O)R¹、-OC(O)R¹、-C(O)OR¹、-(C₁-C₆亚烷基)-C(O)R¹、-SR¹、-S(O)₂R¹、-S(O)₂-N(R¹)(R¹)、-(C₁-C₆亚烷基)-S(O)₂R¹、-(C₁-C₆亚烷基)-S(O)₂-N(R¹)(R¹)、-N(R¹)(R¹)、-C(O)-N(R¹)(R¹)、-N(R¹)-C(O)R¹、-N(R¹)-C(O)OR¹、-(C₁-C₆亚烷基)-N(R¹)-C(O)R¹、-N(R¹)S(O)₂R¹以及-P(O)(R¹)(R¹); 其中烷基、烯基、炔基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的R^a取代; 或者2个R^A或2个R^B与其所连接的碳原子一起形成独立地以0-5次出现的R^a取代的环烷基或杂环基环;

[0073] 每个R¹独立地选自氢、羟基、卤基、硫醇、C₁-C₆烷基、C₁-C₆硫烷基、C₁-C₆烷氧基、C₁-C₆卤烷基、C₁-C₆羟烷基、C₁-C₆杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基, 其中烷基、硫烷基、烷氧基、卤烷基、羟烷基、杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的R^b取代, 或者2个R¹与其所连接的原子一起形成独立地以0-5次出现的R^b取代的环烷基或杂环基环;

[0074] 每个R^a和R^b独立地选自C₁-C₆烷基、卤基、羟基、C₁-C₆卤烷基、C₁-C₆杂烷基、C₁-C₆羟烷基、C₁-C₆烷氧基、环烷基、杂环基以及氰基, 其中烷基、卤烷基、杂烷基、羟烷基、烷氧基、环烷基以及杂环基中的每一者独立地以0-5次出现的R¹取代;

[0075] 每个R¹独立地选自C₁-C₆烷基、C₁-C₆杂烷基、卤基、羟基、C₁-C₆卤烷基、C₁-C₆羟烷基、

环烷基以及氰基;或者2个R'与其所连接的原子一起形成环烷基或杂环基环;并且

[0076] p为0、1、2、3、4或5;并且

[0077] q为0、1、2、3或4。

[0078] 在一些实施方案中,环A为环烷基。在一些实施方案中,环A为5元或6元环烷基环。在一些实施方案中,环A为环戊基或环己基。在一些实施方案中,环A为杂环基。在一些实施方案中,环A为5元或6元杂环基。在一些实施方案中,环A为四氢吡喃基、四氢呋喃基或吡咯烷基。在一些实施方案中,环A为环烯基环。在一些实施方案中,环A为环戊烯基。

[0079] 在一些实施方案中,环B为芳基。在一些实施方案中,环B为苯基。在一些实施方案中,环B为杂芳基。在一些实施方案中,环B为吡啶基。在一些实施方案中,环B为杂环基。在一些实施方案中,环B为吡咯烷基。

[0080] 在一些实施方案中, L^1 为键、 $-C(O)-$ 或 $-N(R^1)-$;并且 L^2 为 $-N(R^1)-C(O)-(C_1-C_6$ 亚烷基) $-$ 或 $-C(O)-N(R^1)-(C_1-C_6$ 亚烷基) $-$ 。在一些实施方案中, L^1 为 $-NH-$ 并且 L^2 为 $-C(O)-NH-CH(CH_2OH)-*$ 、 $-C(O)-N(CH_3)-CH_2-*$ 、 $-C(O)-N(CH_3)-CH(CH_3)-*$ 、 $-C(O)N(CH_2CH_3)CH_2-*$ 、 $-C(O)$

$NHCH(CH_3)-*$ 、 $-C(O)N(CD_3)CH_2-*$ 、 $-C(O)NHCH(CF_3)-*$ 以及,其中“*”表示结合至环

B的 L^2 的一部分。在一些实施方案中, L^1 为 $-NH-$, L^2 为 $-C(O)-$ 并且环B为吡咯烷基。在一些实施方案中, L^1 为 $-NH-$, L^2 为 $-C(O)-$ 并且环B为吡咯烷-1-基。

[0081] 在一些实施方案中,每个 R^1 独立地选自氢和以0-5次出现的 R^b 取代的 C_1-C_6 烷基。在一些实施方案中,每个 R^1 独立地选自氢和 $-CH_3$ 。

[0082] 在一些实施方案中,每个 R^A 和 R^B 独立地选自羟基、 C_1-C_6 烷基、 C_1-C_6 烷氧基、卤基、 C_1-C_6 杂烷基、 C_1-C_6 卤烷基、 C_1-C_6 卤烷氧基、 C_1-C_6 羟烷基、环烷基、芳基、杂芳基、硝基、氰基、 $-C(O)R^1$ 、 $-OC(O)R^1$ 、 $-C(O)OR^1$ 、 $-SR^1$ 、 $-S(O)_2R^1$ 、 $-S(O)_2N(R^1)(R^1)$ 、 $-N(R^1)(R^1)$ 、 $-C(O)-N(R^1)(R^1)$ 、 $-N(R^1)-C(O)R^1$ 、 $-N(R^1)-C(O)OR^1$ 以及 $-N(R^1)S(O)_2R^1$;其中烷基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基以及杂芳基中的每一者独立地以0-5次出现的 R^a 取代;或者2个 R^A 或2个 R^B 与其所连接的碳原子一起形成独立地以0-5次出现的 R^a 取代的环烷基或杂环基环。

[0083] 在一些实施方案中,每个 R^A 独立地选自羟基、 C_1-C_6 烷基、 C_1-C_6 烷氧基、卤基、 $-C(O)-N(R^1)(R^1)$ 、 $-C(O)OR^1$ 、 $-S(O)_2R^1$ 以及 C_1-C_6 卤烷基。在一些实施方案中,每个 R^A 另外且独立地选自 $-CN$ 、氧杂环丁烷基和 C_1-C_6 羟烷基,或结合至环A上的相邻环碳原子的两个 R^A 结合在一起以形成稠合至环A的 C_3-C_6 环烷基。在一些实施方案中,每个 R^A 独立地选自羟基、氟代、氧杂环丁烷-3-基、 $-CHF_2$ 、 $-CH_2CH_3$ 、 $-C(CH_3)_2OH$ 、 $-OCH_3$ 、 $-C(O)N(CH_3)_2$ 、 $-C(O)OCH_3$ 、 $-S(O)_2CH_3$;或者结合至环A上的相邻环碳原子的两个 R^A 结合在一起以形成稠合至环A的环丙基。

[0084] 在一些实施方案中,每个 R^B 独立地选自卤基、 C_1-C_6 烷基、氰基、 C_1-C_6 烷氧基、芳基、杂芳基以及 C_1-C_6 卤烷氧基。在一些实施方案中,每个 R^B 另外选自氧基。

[0085] 在一些实施方案中,环B为吡咯烷基并且至少一个 R^B 为任选地取代的芳基或杂芳基。在一些实施方案中,环B为吡咯烷基并且至少一个 R^B 为任选地取代的苯基或吡啶基。


[0086] 在一些实施方案中,环B为吡咯烷基,并且至少一个 R^B 选自2,3,5-三氟苯基、2,3-二氟苯基、2,5-二氟苯基、2-氯-5-氟苯基、2-氯-5-氟吡啶-3-基、2-氰基-5-氟苯基、2-氟-5-氯苯基、2-甲氧基-3,5-二氟苯基、2-甲氧基-5-氟吡啶-3-基、2-三氟甲氧基-5-氟苯基、

3,5-二氟苯基、3-氯-5-氟苯基、3-氰基-5-氟苯基、3-二氟甲氧基-5-氟苯基、3-氟苯基、5-氟吡啶-3-基以及苯基。

[0087] 在一些实施方案中,环B为吡咯烷基并且一个额外 R^B ,如果存在,则为氟代。

[0088] 在一些实施方案中,环B不同于吡咯烷基,并且每个 R^B 独立地选自氯代、氟代、氧基、 $-CH_3$ 、 $-CF_3$ 、 $-CN$ 、 $-OCH_3$ 、 $-OCF_3$ 以及 $-OCHF_2$ 。

[0089] 在另一方面中,本发明提供式(Ia)化合物:



物或同位素标记形式,或任何前述材料的药学上可接受的盐,其中:

[0091] 环A选自芳基、杂芳基、环烷基以及杂环基;

[0092] L^1 独立地选自键、 $-C(O)-$ 、 $-N(R^1)-$ 、 $-N(R^1)-C(O)-$ 、 $-C(O)-N(R^1)-$ 、 $-(C_1-C_6\text{亚烷基})-N(R^1)-$ 、 $-N(R^1)-(C_1-C_6\text{亚烷基})-$ 、 $-N(R^1)-C(O)-(C_1-C_6\text{亚烷基})-$ 以及 $-C(O)-N(R^1)-(C_1-C_6\text{亚烷基})-$;其中每个亚烷基独立地以0-5次出现的 R' 取代;

[0093] 每个 R^A 和 R^B 独立地选自羟基、 C_1-C_6 烷基、 C_2-C_6 烯基、 C_2-C_6 炔基、 C_1-C_6 烷氧基、卤基、 C_1-C_6 杂烷基、 C_1-C_6 卤烷基、 C_1-C_6 卤烷氧基、 C_1-C_6 羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基、杂环烷基、硝基、氰基、 $-C(O)R^1$ 、 $-OC(O)R^1$ 、 $-C(O)OR^1$ 、 $-(C_1-C_6\text{亚烷基})-C(O)R^1$ 、 $-SR^1$ 、 $-S(O)_2R^1$ 、 $-S(O)_2-N(R^1)(R^1)$ 、 $-(C_1-C_6\text{亚烷基})-S(O)_2R^1$ 、 $-(C_1-C_6\text{亚烷基})-S(O)_2-N(R^1)(R^1)$ 、 $-N(R^1)(R^1)$ 、 $-C(O)-N(R^1)(R^1)$ 、 $-N(R^1)-C(O)R^1$ 、 $-N(R^1)-C(O)OR^1$ 、 $-(C_1-C_6\text{亚烷基})-N(R^1)-C(O)R^1$ 、 $-N(R^1)S(O)_2R^1$ 以及 $-P(O)(R^1)(R^1)$;其中烷基、烯基、炔基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的 R^a 取代;或者2个 R^A 或2个 R^B 与其所连接的碳原子一起形成独立地以0-5次出现的 R^a 取代的环烷基或杂环基环;

[0094] 每个 R^1 独立地选自氢、羟基、卤基、硫醇、 C_1-C_6 烷基、 C_1-C_6 硫烷基、 C_1-C_6 烷氧基、 C_1-C_6 卤烷基、 C_1-C_6 羟烷基、 C_1-C_6 杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基,其中烷基、硫烷基、烷氧基、卤烷基、羟烷基、杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的 R^b 取代,或者2个 R^1 与其所连接的原子一起形成独立地以0-5次出现的 R^b 取代的环烷基或杂环基环;

[0095] 每个 R^a 和 R^b 独立地选自 C_1-C_6 烷基、卤基、羟基、 C_1-C_6 卤烷基、 C_1-C_6 杂烷基、 C_1-C_6 羟烷基、 C_1-C_6 烷氧基、环烷基、杂环基以及氰基,其中烷基、卤烷基、杂烷基、羟烷基、烷氧基、环烷基以及杂环基中的每一者独立地以0-5次出现的 R' 取代;

[0096] 每个 R' 独立地选自 C_1-C_6 烷基、 C_1-C_6 杂烷基、卤基、羟基、 C_1-C_6 卤烷基、 C_1-C_6 羟烷基、环烷基以及氰基;或者2个 R' 与其所连接的原子一起形成环烷基或杂环基环;并且

[0097] p 为0、1、2、3、4或5;并且

[0098] q 为0、1、2、3或4。

[0099] 在一些实施方案中,环A为环烷基。在一些实施方案中,环A为5元或6元环烷基环。在一些实施方案中,环A为环戊基或环己基。在一些实施方案中,环A为杂环基。在一些实施

方案中,环A为5元或6元杂环基。在一些实施方案中,环A为四氢吡喃、四氢呋喃或吡咯烷基。在一些实施方案中,环A为环烯基环。在一些实施方案中,环A为环戊烯基。

[0100] 在一些实施方案中, L^1 为一键、 $-C(O)-$ 或 $-N(R^1)-$ 。在一些实施方案中, L^1 为 $-NH-$ 。

[0101] 在一些实施方案中,每个 R^1 独立地选自氢和以0-5次出现的 R^b 取代的 C_1-C_6 烷基。在一些实施方案中,每个 R^1 独立地选自氢和 $-CH_3$ 。

[0102] 在一些实施方案中,每个 R^A 和 R^B 独立地选自羟基、 C_1-C_6 烷基、 C_1-C_6 烷氧基、卤基、 C_1-C_6 杂烷基、 C_1-C_6 卤烷基、 C_1-C_6 卤烷氧基、 C_1-C_6 羟烷基、环烷基、芳基、杂芳基、硝基、氰基、 $-C(O)R^1$ 、 $-OC(O)R^1$ 、 $-C(O)OR^1$ 、 $-SR^1$ 、 $-S(O)_2R^1$ 、 $-S(O)_2-N(R^1)(R^1)$ 、 $-N(R^1)(R^1)$ 、 $-C(O)-N(R^1)(R^1)$ 、 $-N(R^1)-C(O)R^1$ 、 $-N(R^1)-C(O)OR^1$ 以及 $-N(R^1)S(O)_2R^1$;其中烷基、烯基、炔基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的 R^a 取代;或2个 R^A 或2个 R^B 与其连接的碳原子一起形成独立地以0-5次出现的 R^a 取代的环烷基或杂环基环。

[0103] 在一些实施方案中,每个 R^A 独立地选自羟基、 C_1-C_6 烷基、 C_1-C_6 烷氧基、卤基、 $-C(O)-N(R^1)(R^1)$ 、 $-C(O)OR^1$ 、 $-S(O)_2R^1$ 和 C_1-C_6 卤烷基。在一些实施方案中,每个 R^A 另外且独立地选自 $-CN$ 、氧杂环丁烷基和 C_1-C_6 羟烷基,或者结合至环A上的相邻环碳原子的两个 R^A 结合在一起以形成稠合至环A的 C_3-C_6 环烷基。在一些实施方案中,每个 R^A 独立地选自羟基、氟代、氧杂环丁烷-3-基、 $-CHF_2$ 、 $-CH_2CH_3$ 、 $-C(CH_3)_2OH$ 、 $-OCH_3$ 、 $-C(O)N(CH_3)_2$ 、 $-C(O)OCH_3$ 、 $-S(O)_2CH_3$;或者结合至环A上的相邻环碳原子的两个 R^A 结合在一起以形成稠合至环A的环丙基。

[0104] 在一些实施方案中,每个 R^B 独立地选自卤基、 C_1-C_6 烷基、氰基、 C_1-C_6 烷氧基、芳基、杂芳基以及 C_1-C_6 卤烷氧基。在一些实施方案中,每个 R^B 另外选自氧基。


[0105] 在一些实施方案中,在环B为吡咯烷基时,至少一个 R^B 选自2,3,5-三氟苯基、2,3-二氟苯基、2,5-二氟苯基、2-氯-5-氟苯基、2-氯-5-氟吡啶-3-基、2-氰基-5-氟苯基、2-氟-5-氯苯基、2-甲氧基-3,5-二氟苯基、2-甲氧基-5-氟吡啶-3-基、2-三氟甲氧基-5-氟苯基、3,5-二氟苯基、3-氯-5-氟苯基、3-氰基-5-氟苯基、3-二氟甲氧基-5-氟苯基、3-氟苯基、5-氟吡啶-3-基以及苯基。

[0106] 在一些实施方案中,当环B为吡咯烷基时,一个额外 R^B ,如果存在,为氟代。

[0107] 在一些实施方案中,p为0、1或2。

[0108] 在一些实施方案中,q为1、2或3。

[0109] 在另一方面中,本发明提供式(II)化合物:



[0111] 环A和B各自独立地选自芳基、杂芳基、环烷基和杂环基;

[0112] 每个 R^A 和 R^B 独立地选自羟基、 C_1-C_6 烷基、 C_2-C_6 烯基、 C_2-C_6 炔基、 C_1-C_6 烷氧基、卤基、 C_1-C_6 杂烷基、 C_1-C_6 卤烷基、 C_1-C_6 卤烷氧基、 C_1-C_6 羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基、杂环烷基、硝基、氰基、 $-C(O)R^1$ 、 $-OC(O)R^1$ 、 $-C(O)OR^1$ 、 $-(C_1-C_6$ 亚烷基) $-C(O)R^1$ 、 $-$

SR^1 、 $-\text{S}(\text{O})_2\text{R}^1$ 、 $-\text{S}(\text{O})_2-\text{N}(\text{R}^1)(\text{R}^1)$ 、 $-(\text{C}_1-\text{C}_6\text{亚烷基})-\text{S}(\text{O})_2\text{R}^1$ 、 $-(\text{C}_1-\text{C}_6\text{亚烷基})-\text{S}(\text{O})_2-\text{N}(\text{R}^1)(\text{R}^1)$ 、 $-\text{N}(\text{R}^1)(\text{R}^1)$ 、 $-\text{C}(\text{O})-\text{N}(\text{R}^1)(\text{R}^1)$ 、 $-\text{N}(\text{R}^1)-\text{C}(\text{O})\text{R}^1$ 、 $-\text{N}(\text{R}^1)-\text{C}(\text{O})\text{OR}^1$ 、 $-(\text{C}_1-\text{C}_6\text{亚烷基})-\text{N}(\text{R}^1)-\text{C}(\text{O})\text{R}^1$ 、 $-\text{N}(\text{R}^1)\text{S}(\text{O})_2\text{R}^1$ 以及 $-\text{P}(\text{O})(\text{R}^1)(\text{R}^1)$ ；其中烷基、烯基、炔基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基、杂芳基、芳氧基、芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的 R^a 取代；或者2个 R^a 或2个 R^b 与其所连接的碳原子一起形成独立地以0-5次出现的 R^a 取代的环烷基或杂环基环；

[0113] 每个 R^1 独立地选自氢、羟基、卤基、硫醇、 C_1-C_6 烷基、 C_1-C_6 硫烷基、 C_1-C_6 烷氧基、 C_1-C_6 卤烷基、 C_1-C_6 羟烷基、 C_1-C_6 杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基，其中烷基、硫烷基、烷氧基、卤烷基、羟烷基、杂烷基、环烷基、环烷基烷基、杂芳烷基、杂环基以及杂环烷基中的每一者独立地以0-5次出现的 R^b 取代，或者2个 R^1 与其所连接的原子一起形成独立地以0-5次出现的 R^b 取代的环烷基或杂环基环；

[0114] R^{1a} 选自氢、 C_1-C_6 烷基和氧化 C_1-C_6 烷基；

[0115] R^{1b} 选自氢和 C_1-C_6 烷基；

[0116] 每个 R^a 和 R^b 独立地选自 C_1-C_6 烷基、卤基、羟基、 C_1-C_6 卤烷基、 C_1-C_6 杂烷基、 C_1-C_6 羟烷基、 C_1-C_6 烷氧基、环烷基、杂环基以及氰基，其中烷基、卤烷基、杂烷基、羟烷基、烷氧基、环烷基以及杂环基中的每一者独立地以0-5次出现的 R' 取代；

[0117] 每个 R' 独立地选自 C_1-C_6 烷基、 C_1-C_6 杂烷基、卤基、羟基、 C_1-C_6 卤烷基、 C_1-C_6 羟烷基、环烷基以及氰基；或者2个 R' 与其所连接的原子一起形成环烷基或杂环基环；并且

[0118] p 为0、1、2、3、4或5；并且

[0119] q 为0、1、2、3或4。

[0120] 在一些实施方案中，环A为环烷基。在一些实施方案中，环A为5元或6元环烷基环。在一些实施方案中，环A为环戊基或环己基。在一些实施方案中，环A为杂环基。在一些实施方案中，环A为5元或6元杂环基。在一些实施方案中，环A为四氢吡喃、四氢呋喃或吡咯烷基。在一些实施方案中，环A为环烯基环。在一些实施方案中，环A为环戊烯基。

[0121] 在一些实施方案中，环B为芳基。在一些实施方案中，环B为苯基。在一些实施方案中，B为杂芳基。在一些实施方案中，环B为吡啶基。

[0122] 在一些实施方案中，每个 R^1 独立地选自氢和以0-5次出现的 R^b 取代的 C_1-C_6 烷基。

[0123] 在一些实施方案中， R^{1a} 为氢、 $-\text{CH}_3$ 、 $-\text{CD}_3$ 或 $-\text{CH}_2\text{CH}_3$ 。

[0124] 在一些实施方案中， R^{1b} 为氢。

[0125] 在一些实施方案中，每个 R^a 和 R^b 独立地选自羟基、 C_1-C_6 烷基、 C_1-C_6 烷氧基、卤基、 C_1-C_6 杂烷基、 C_1-C_6 卤烷基、 C_1-C_6 卤烷氧基、 C_1-C_6 羟烷基、环烷基、芳基、杂芳基、硝基、氰基、 $-\text{C}(\text{O})\text{R}^1$ 、 $-\text{OC}(\text{O})\text{R}^1$ 、 $-\text{C}(\text{O})\text{OR}^1$ 、 $-\text{SR}^1$ 、 $-\text{S}(\text{O})_2\text{R}^1$ 、 $-\text{S}(\text{O})_2-\text{N}(\text{R}^1)(\text{R}^1)$ 、 $-\text{N}(\text{R}^1)(\text{R}^1)$ 、 $-\text{C}(\text{O})-\text{N}(\text{R}^1)(\text{R}^1)$ 、 $-\text{N}(\text{R}^1)-\text{C}(\text{O})\text{R}^1$ 、 $-\text{N}(\text{R}^1)-\text{C}(\text{O})\text{OR}^1$ 以及 $-\text{N}(\text{R}^1)\text{S}(\text{O})_2\text{R}^1$ ；其中烷基、烷氧基、杂烷基、卤烷基、卤烷氧基、羟烷基、环烷基、芳基以及杂芳基中的每一者独立地以0-5次出现的 R^a 取代；或者2个 R^a 或2个 R^b 与其所连接的碳原子一起形成独立地以0-5次出现的 R^a 取代的环烷基或杂环基环。

[0126] 在一些实施方案中，每个 R^a 独立地选自羟基、 C_1-C_6 烷基、 C_1-C_6 烷氧基、卤基、 $-\text{C}(\text{O})-\text{N}(\text{R}^1)(\text{R}^1)$ 、 $-\text{C}(\text{O})\text{OR}^1$ 、 $-\text{S}(\text{O})_2\text{R}^1$ 以及 C_1-C_6 卤烷基。在一些实施方案中，每个 R^a 另外且独立地选自 $-\text{CN}$ 、氧杂环丁烷基和 C_1-C_6 羟烷基，或者结合至环A上的相邻环碳原子的两个 R^a 结合在一

起以形成稠合至环A的C₃-C₆环烷基。在一些实施方案中,每个R^A独立地选自羟基、氟代、氧杂环丁烷-3-基、-CHF₂、-CH₂CH₃、-C(CH₃)₂OH、-OCH₃、-C(O)N(CH₃)₂、-C(O)OCH₃、-S(O)₂CH₃;或者结合至环A上的相邻环碳原子的两个R^A结合在一起以形成稠合至环A的环丙基。

[0127] 在一些实施方案中,每个R^B独立地选自卤基、C₁-C₆烷基、氰基、C₁-C₆烷氧基、芳基、杂芳基以及C₁-C₆卤烷氧基。在一些实施方案中,每个R^B另外选自氧基。在一些实施方案中,每个R^B独立地选自氯代、氟代、氧基、-CH₃、-CF₃、-CN、-OCH₃、-OCF₃以及-OCHF₂。

[0128] 在一些实施方案中,每个R'独立地选自C₁-C₆烷基、C₁-C₆卤烷基和C₁-C₆羟烷基;或者2个R'与其所连接的原子一起形成环烷基或杂环基环。在一些实施方案中,一个R'为氢,并且另一个R'选自氢、C₁-C₆烷基、C₁-C₆卤烷基和C₁-C₆羟烷基;或者2个R'与其所连接的原子一起形成环烷基环。在一些实施方案中,一个R'为氢,并且另一个R'选自氢、-CH₂OH、-CH₃或-CF₃,或者2个R'与其所连接的原子一起形成环丙-1,1-二基环。

[0129] 在一些实施方案中,p为0、1或2。

[0130] 在一些实施方案中,q为0、1、2或3。

[0131] 虽然,如上所述,式(I)、(Ia)或(II)中的变量的各种实施方案及其方面可自选一组化学部分,但是本发明还包括以下情况作为其另外的实施方案和方面,其中此变量:a)选自此组中的化学部分的任何子集;并且为b)此组的任何单一成员。

[0132] 虽然对于式(I)、(Ia)和(II)中的每个变量个别地阐明(或暗示,如先前段落中论述)其各种实施方案和方面,但是本发明涵盖式(I)、(Ia)和(II)中的每种变量的不同实施方案和方面的所有可能组合。

[0133] 本发明的示例性化合物的结构以及NMR和LCMS数据示出在图1中。在某些实施方案中,本发明的化合物选自由以下组成的组:图1中的任一种化合物及其药学上可接受的盐、溶剂化物、水合物、互变异构物、立体异构物以及同位素标记的衍生物。

[0134] 本发明还提供含有药学上可接受的载剂和任何式(I)、(Ia)和(II)化合物的药物组合物。

[0135] 这些化合物的药学上可接受的盐也预期用于本文所述的用途。

[0136] “药学上可接受的盐”是指保持其生物性质并且没有毒性或在其他方面不合药物用途需要的本发明的化合物的任何盐。药学上可接受的盐可从在本领域中熟知的各种有机和无机抗衡离子衍生并且包括。这些盐包括:(1)与有机酸或无机酸所形成的酸加成盐,诸如盐酸、氢溴酸、硫酸、硝酸、磷酸、氨基磺酸、乙酸、三氟乙酸、三氯乙酸、丙酸、己酸、环戊基丙酸、乙醇酸、戊二酸、丙酮酸、乳酸、丙二酸、琥珀酸、山梨酸、抗坏血酸、苹果酸、马来酸、富马酸、酒石酸、柠檬酸、苯甲酸、3-(4-羟基苯甲酰基)苯甲酸、苦味酸、肉桂酸、扁桃酸、邻苯二甲酸、月桂酸、甲磺酸、乙磺酸、1,2-乙烷-二磺酸、2-羟基乙磺酸、苯磺酸、4-氯苯磺酸、2-萘磺酸、4-甲苯磺酸、樟脑酸、樟脑磺酸、4-甲基双环[2.2.2]-辛-2-烯-1-羧酸、葡庚糖酸、3-苯基丙酸、三甲基乙酸、叔丁基乙酸、月桂基硫酸、葡萄糖酸、苯甲酸、谷氨酸、羟基萘甲酸、水杨酸、硬脂酸、环己基氨基磺酸、奎尼酸、黏康酸及其类似酸;或者(2)当存在于母体化合物中的酸性质子(a)置换为金属离子(例如碱金属离子、碱土金属离子或铝离子)或碱金属或碱土金属氢氧化物(诸如氢氧化钠、氢氧化钾、氢氧化钙、氢氧化镁、氢氧化铝、氢氧化锂、氢氧化锌和氢氧化钡、氨)或者(b)与有机碱配位时所形成的盐,所述有机碱诸如脂族、脂环族或芳族有机胺,诸如氨、甲胺、二甲胺、二乙胺、甲基吡啶、乙醇胺、二乙醇胺、三乙醇

胺、乙二胺、赖氨酸、精氨酸、鸟氨酸、胆碱、N,N'-二苄基乙烯-二胺、氯普鲁卡因、二乙醇胺、普鲁卡因、N-苄基苯乙胺、N-甲基葡萄糖胺、三(羟甲基)-氨基甲烷、四甲基氢氧化铵及其类似物。药学上可接受的盐仅举例而言还包括钠、钾、钙、镁、铵、四烷基铵及其类似物,并且当化合物含有碱性官能度时,包括无毒有机酸或无机酸的盐,诸如盐酸盐、氢溴酸盐、酒石酸盐、甲磺酸盐、苯磺酸盐、乙酸盐、马来酸盐、草酸盐等。

[0137] 药物组合物

[0138] 本发明的药物组合物包含一种或多种本发明的化合物和一种或多种生理上或药学上可接受的载剂。术语“药学上可接受的载剂”是指参与携带或运输任何主题组合物或其组分的药学上可接受的材料、组合物或媒介物,诸如液体或固体填充剂、稀释剂、赋形剂、溶剂或封装材料。每种载剂必须在与主题组合物及其组分相容并且不伤害患者的意义上是“可接受的”。可充当药学上可接受的载剂的材料的一些实例包括:(1)糖,诸如乳糖、葡萄糖和蔗糖;(2)淀粉,诸如玉米淀粉和马铃薯淀粉;(3)纤维素及其衍生物,诸如羧甲基纤维素钠、乙基纤维素和乙酸纤维素;(4)粉末状黄耆胶;(5)麦芽;(6)明胶;(7)滑石;(8)赋形剂,诸如可可脂和栓剂蜡;(9)油,诸如花生油、棉籽油、红花油、芝麻油、橄榄油、玉米油和大豆油;(10)二醇,诸如丙二醇;(11)多元醇,诸如甘油、山梨糖醇、甘露糖醇和聚乙二醇;(12)酯,诸如油酸乙酯和月桂酸乙酯;(13)琼脂;(14)缓冲剂,诸如氢氧化镁和氢氧化铝;(15)海藻酸;(16)无热原水;(17)等张生理盐水;(18)林格氏溶液(Ringer's solution);(19)乙醇;(20)磷酸盐缓冲溶液;以及(21)用于药物制剂中的其他无毒相容材料。

[0139] 本发明的组合物可经口、胃肠外、通过吸入喷雾剂、局部、经直肠、经鼻、经颊、经阴道或通过植入储存器施用。如本文所用的术语“胃肠外”包括皮下、静脉内、肌肉内、关节内、滑膜内、胸骨内、鞘内、肝内、病灶内和颅内注射或输注技术。在一些实施方案中,本发明的组合物经口、腹膜内或静脉内施用。本发明的组合物的无菌可注射形式可为水性或油性悬浮液。这些悬浮液可根据本领域中已知的技术使用适合分散剂或湿润剂和悬浮剂来配制。无菌可注射制剂也可用于无毒胃肠外可接受的稀释剂或溶剂中的无菌可注射溶液或悬浮液,例如呈于1,3-丁二醇中的溶液形式。可采用的可接受的媒介物和溶剂尤其为水、林格氏溶液和等张氯化钠溶液。此外,无菌非挥发性油通常用作溶剂或悬浮介质。

[0140] 出于此目的,可采用任何温和的不挥发性油,包括合成的单甘油酯或二甘油酯。脂肪酸(诸如油酸)及其甘油酯衍生物适用于制备可注射剂,天然药学上可接受的油诸如橄榄油或蓖麻油亦如此,尤其呈其聚氧乙基化形式。这些油溶液或悬浮液也可含有长链醇稀释剂或分散剂,诸如羧甲基纤维素或通常用于配制药学上可接受的剂型(包括乳液和悬浮液)的类似分散剂。通常用于制造药学上可接受的固体、液体或其他剂型的其他常用表面活性剂(诸如吐温(Tween)、斯盘(Span))和其他乳化剂或生物可用性增强剂也可用于配制目的。

[0141] 本发明的药学上可接受的组合物可以任何口腔可接受的剂型经口施用,所述剂型包括但不限于胶囊、片剂、水性悬浮液或溶液。在供经口使用的片剂的情况下,通常使用的载剂包括乳糖和玉米淀粉。还通常添加润滑剂诸如硬脂酸镁。对于以胶囊形式进行的经口施用,适合的稀释剂包括乳糖和干燥玉米淀粉。当水性悬浮液为经口使用所需时,使活性成分与乳化剂和悬浮剂组合。必要时,还可添加某些甜味剂、调味剂或着色剂。

[0142] 或者,本发明的药学上可接受的组合物可以用于经直肠施用的栓剂形式施用。这些组合物可通过将药剂与在室温下为固体但在直肠温度下为液体的适合非刺激性赋形剂

混合来制备并且因此将在直肠中熔融以释放药物。这些材料包括可可脂、蜂蜡和聚乙二醇。

[0143] 本发明的药学上可接受的组合物也可以局部施用,尤其当治疗目标包括通过局部应用可易于接近的区域或器官时,包括眼、皮肤或下肠道疾病。易于制备适合的局部制剂以用于这些区域或器官中的每一个。用于下肠道的局部应用可以直肠栓剂制剂(参见上文)形式或以适合灌肠剂制剂形式实现。还可使用局部经皮贴片。

[0144] 对于局部应用,药学上可接受的组合物可以配制为含有悬浮或溶解于一种或多种载剂中的活性组分的适合软膏剂形式。用于局部施用本发明的化合物的载剂包括但不限于矿物油、液体凡士林、白凡士林、丙二醇、聚氧化乙烯、聚氧化丙烯化合物、乳化蜡以及水。或者,药学上可接受的组合物可以配制为含有悬浮或溶解于一种或多种药学上可接受的载剂中的活性组分的适合洗剂或乳膏剂。适合载剂包括但不限于矿物油、脱水山梨醇单硬脂酸酯、聚山梨醇酯60、十六烷基酯蜡、鲸蜡硬脂醇(cetearyl alcohol)、2-辛基十二烷醇、苯甲醇以及水。

[0145] 本发明的药学上可接受的组合物也可通过经鼻气雾剂或吸入剂施用。这些组合物根据药物制剂领域中熟知的技术制备并且可采用苯甲醇或其他适合防腐剂、增强生物可用性的吸收促进剂、氟碳化合物和/或其他常规增溶剂或分散剂制备成盐水中的溶液。

[0146] 本发明的化合物可与载剂材料组合以产生呈单一剂型的组合物的量将根据所治疗宿主和特定施用模式而变化。优选地,应配制所述组合物以使得可向接受这些组合物的患者施用0.01-100mg/kg体重/天之间的剂量的抑制剂。

[0147] 剂量

[0148] 本发明的化合物(包括药学上可接受的盐和氘化变体)的毒性和治疗功效变体可通过在细胞培养物或实验动物中的标准药物程序来确定。 LD_{50} 为对于50%群体致死的剂量。 ED_{50} 为在50%群体中治疗有效的剂量。毒性作用与治疗作用之间的剂量比率(LD_{50}/ED_{50})为治疗指数。优选展现较大治疗指数的化合物。当可使用显示毒副作用的化合物时,应谨慎设计将此类化合物靶向受影响组织的位置的递送系统以便使对未感染细胞的潜在损害最小化从而减少副作用。

[0149] 从细胞培养测定和/或动物研究获得的数据可用于配制用于人类的剂量范围。这些化合物的剂量优选地在循环浓度范围内,包括具有极低或没有毒性的 ED_{50} 。所述剂量可根据所使用的剂型和所使用的施用途径而在此范围内变化。对于任何化合物,治疗有效剂量可最初从细胞培养测定估计。剂量可在动物模型中配制以实现循环血浆浓度范围,包括如在细胞培养物中所确定的 IC_{50} (即,实现症状的一半最大抑制的测试化合物的浓度)。此信息可用于更准确地确定在人类中的有用剂量。血浆中的水平可例如通过高效液相色谱来测量。

[0150] 还应理解用于任何特定患者的特定剂量和治疗方案将取决于多种因素,包括所用的特定化合物的活性、年龄、体重、一般健康状况、性别、膳食、施用时间、排泄速率、药物组合以及治疗医师的判断和所治疗特定疾病的严重性。本发明的化合物在组合物中的量也将取决于组合物中的特定化合物。

[0151] 治疗

[0152] NTRK融合物涉及多种类型的癌症。这些融合物具有与受体的天然或野生型形式的受体相同的完整NTRK激酶结构域;因此,如本文使用,具有与野生型NTRK相同的激酶结构域

的任何NTRK蛋白质(NTRK1、2或3)被称为“野生型NTRK”。突变可在NTRK激酶结构域中发生,从而导致突变体对于激酶抑制剂疗法具有抗性。这些抗性突变可使用结构生物学和计算分析,以及通过检查其中序列变化产生不同氨基酸的密码子的密码子序列来预测。或者,给定抑制剂的抗性突变可在实验上通过施用该抑制剂(例如,已知NTRK野生型抑制剂)并且使细胞暴露于突变促进剂(诸如ENU)来识别。将细胞洗涤,然后用递增浓度(2-100X增殖IC₅₀)的选定化合物来铺板。然后,在3-4周后收集具有细胞向外生长的孔。具体而言,通过两种方法来识别NTRK融合物内的氨基酸位置595(NTRK1 wt编号)处的突变,其实现从甘氨酸至精氨酸残基的变化(迄今称为‘G595R’)。此突变随后证明对于临床上评估的两种NTRK抑制剂赋予显著抗性(在下表中示出)。如表中示出,化合物对于野生型NTRK具有活性,但是对于NTRK融合物的G595R突变形式的活性显著地较小。

化合物	NTRK wt 酶分析 IC ₅₀ (nM)	NTRK wt 细胞 GI ₅₀ (nM)	NTRK G595R 细胞 GI ₅₀ (nM)
	IC ₅₀ (nM)	GI ₅₀ (nM)	GI ₅₀ (nM)
厄洛替尼	0.6	2	2700
TSR-011	2.3	32	12000
克唑替尼	9.3	87	9000

[0154] 因此,在另一方面,本发明提供一种治疗患有由异常神经营养酪氨酸受体激酶(NTRK)活性介导的病状的受试者的方法,其包括向受试者施用治疗有效量的本文所述的化合物或化合物的药物组合物。

[0155] 本发明提供抑制野生型NTRK和NTRK的抗性G595R突变体的化合物。

[0156] 在另一方面中,本发明提供治疗对于癌症治疗显现抗性的受试者的方法,包含向受试者施用治疗有效量的化合物或本文所述化合物的药物组合物。

[0157] 此外,相对于其他激酶,抑制剂可对野生型NTRK具有选择性,由此导致与抑制其他激酶相关的毒性减少。因为其对于野生型和突变体NTRK的活性,本文描述的化合物可用于治疗患有与异常NTRK活性相关的病状的患者。所述化合物也可用于治疗各种癌症。在一些实施方案中,癌症选自非小细胞肺癌、乳腺癌、黑素瘤、低级和高级神经胶质瘤、胶质母细胞瘤、小儿科星细胞瘤、结直肠癌、乳突甲状腺癌、胰腺癌、头颈部癌、胆管癌、急性髓性白血病、分泌乳腺癌、唾液癌症以及spitz样肿瘤。

[0158] 化合物也可用于治疗对于野生型NTRK抑制剂发展抗性的患者,或具有突变形式的NTRK(诸如G595R突变体)的患者。方法包括施用对于NTRK抗性突变体具有活性的本发明的化合物或组合物的步骤。“活性”意指在生物化学测定中对于至少一种抗性突变体测量时,化合物具有小于1 μ M、500nM、250nM、100nM、75nM、50nM、25nM、10nM或5nM的IC₅₀。

[0159] 本文所述的化合物和组合物可单独与其他化合物(包括其他NTRK调控化合物或其他治疗剂)组合来施用。在一些实施方案中,本发明的化合物或组合物可与一种或多种选自以下的化合物组合施用:卡博替尼(COMETRIQ)、凡德他尼(CALPRESA)、索拉非尼(NEXAVAR)、舒尼替尼(SUTENT)、瑞格非尼(STAVARGA)、帕纳替尼(ICLUSIG)、贝伐单抗(AVASTIN)、克唑替尼(XALKORI)或吉非替尼(IRESSA)。本发明的化合物或组合物可通过相同或不同施用途径与其他治疗剂同时或依次施用。本发明的化合物可与其他治疗剂一起包含在单一制剂中或在单独的制剂中。

[0160] 合成

[0161] 本发明的化合物(包括其盐和N-氧化物)可使用已知的有机合成技术制备并且可根据众多可能的合成途径(诸如以下流程中的合成途径)中的任一种来合成。用于制备本发明的化合物的反应可在适合溶剂中进行,所述溶剂可由有机合成领域技术人员容易选择。在进行反应的温度下,例如在可范围为溶剂冷冻温度至溶剂沸腾温度的温度下,适当的溶剂可大体上不与起始材料(反应物)、中间体或产物反应。给定反应可在一种溶剂或超过一种溶剂的混合物中进行。根据特定反应步骤,适于特定反应步骤的溶剂可由技术人员选择。

[0162] 制备本发明的化合物可涉及各种化学基团的保护和脱保护。保护和脱保护的需要和合适保护基团的选择可通过本领域技术人员容易地确定。保护基团的化学可发现于例如Wuts和Greene, *Protective Groups in Organic Synthesis*, 第4版, John Wiley & Sons: New Jersey, (2006) 中, 其以引用方式整体并入本文。

[0163] 反应可根据在本领域中已知的任何合适方法进行监测。举例而言, 产物形成可通过光谱手段进行监测, 诸如核磁共振(NMR) 光谱学(例如, ^1H 或 ^{13}C)、红外线(IR) 光谱学、分光光度测定法(例如, UV-可见光)、质谱法(MS) 或通过色谱方法诸如高效液相色谱(HPLC) 或薄层色谱(TLC)。用于化合物表征的分析仪器和方法:

[0164] **LC-MS:**除非另外指示, 否则所有液相色谱-质谱(LC-MS) 数据(分析样品的纯度和身份)都用Agilent 1260型LC系统获得, 所述系统使用利用ES-API电离的Agilent 6120型质谱仪, 配备有在摄氏22.4°C下的Agilent Poroshel 120 (EC-C18, 2.7 μm 粒度, 3.0 $\times$ 50mm 尺寸) 反相柱。流动相由含0.1%甲酸的水和含0.1%甲酸的乙腈的溶剂混合物组成。利用在4分钟过程内从95%水性/5%有机至5%水性/95%有机流动相的恒定梯度。流速在1mL/min 下恒定。

[0165] **制备型LC-MS:**在配备有在摄氏22.4°C下的Luna 5u C18 (2) 100A, AXIA填充的250 $\times$ 21.2mm反相柱的Shimadzu Discovery VP®制备型系统上进行制备型HPLC。流动相由含0.1%甲酸的水和含0.1%甲酸的乙腈的溶剂混合物组成。利用在25分钟过程中从95%水性/5%有机至5%水性/95%有机流动相的恒定梯度。流速在20mL/min 下恒定。在微波中执行的反应在Biotage启动器微波单元中以此方式进行。

[0166] **手性HPLC:**拆分手性混合物的制备HPLC在配备有Chiralpak AS-H柱(5mm, 3.0cm id $\times$ 25cm L) 的Thar SFC预-80仪器中执行。流动相由SFC CO₂ (A) 和MeOH/0.1%NH₄OH (B) 组成。67%至33% (B) 的恒定梯度在65g/min的流速下使用100巴的系统背压来保持。分离进展通过220nm的波长下的UV检测来监测。

[0167] **硅胶色谱:**硅胶色谱在Teledyne Isco CombiFlash®Rf单元或Biotage®Isolera Four单元上执行。

[0168] **质子NMR:**除非另外指示, 否则所有 ^1H NMR光谱都用Varian 400MHz Unity Inova 400MHz NMR仪器(采集时间=3.5秒, 伴有1秒延迟; 16至64次扫描) 获得。当表征时, 所有质子均在DMSO-d₆溶剂中以关于残余DMSO (2.50ppm) 的百万分率(ppm) 形式加以报导。


实施例

[0169] 以下实施例旨在为示例性的, 并且不意图以任何方式具有限制性。


[0170] 以下流程图提供关于制备本发明的化合物的一般性指导。本领域技术人员将理

解可使用制备各种本发明的化合物的有机化学的一般知识对所述流程中所示的制剂进行修改或优化。

[0171] 一般合成1:




[0172]



[0173] 对于某些化合物,一般合成开始于二卤化物取代的1H-吡唑并[3,4-d]嘧啶的合适氮保护(P)。氮保护的双环可在合适条件下,例如在亲核芳族取代反应条件下,使用碱(诸如二异丙基乙胺(DIPEA)),在极性溶剂诸如二噁烷中,在嘧啶环上的卤化物处使用适当取代的环A取代以提供被环A取代的双环。吡唑环的卤化物可在钯介导的羰基插入反应条件下取代,随后水解以提供所得羧酸。羧酸可在合适偶合条件下,例如在酰胺偶合反应条件下与环B反应以提供被环A和B取代的氮保护化合物。移除保护基可提供式I化合物。


[0174] 合成方案1:



[0176] 稍微更具具体形式的一般合成流程1在以上合成方案1中示出。合成方案开始于3-溴-4-氯-1H-吡唑并[3,4-d]嘧啶1的SEM基团保护。SEM保护的杂环2可在亲核芳族取代反应条件下在极性溶剂诸如二噁烷中使用碱诸如二异丙基乙胺(DIPEA),使用氨基醇取代以提

供胺取代的杂环3。3-溴吡唑并嘧啶3在DMF-MeOH溶剂混合物中经受钯介导的羰基插入反应以提供甲酯4。在通过NaOH处理来水解酯之后,羧酸5在酰胺偶合反应条件下与苄胺或吡咯烷反应以提供SEM保护的化合物6。SEM保护基可使用TBAF或在酸性条件下移除以提供最终化合物7。以下描述的化合物使用一般合成1、2或3来制备,如分别在合成方案1、2或3中进一步详述。

[0177] 一般合成2




[0178]




[0179] 对于某些化合物,一般合成开始于4-氧基-4,5-二氢-1H-吡唑并[3,4-d]嘧啶-3-羧酸的合适氮保护(P)。氮保护的双环可在氯化试剂诸如亚硫酰氯存在下氯化并与胺偶合。所得化合物可在合适条件下,例如在亲核芳族取代反应条件下,使用碱诸如二异丙基乙胺(DIPEA),在极性溶剂诸如二噁烷中,在嘧啶环上的卤化物处使用适当取代的环A取代以提供用环A取代的双环。移除保护基可提供式I化合物。以下描述的化合物可使用此一般合成来制备。此外,手性HPLC可用于拆分式I、(Ia)、(Ia-1)、(Ia-2)、(Ib)、(Ib-1)、(Ib-2)、II、(IIa)、(IIb)、(IIc)化合物的手性混合物。

[0180] 合成方案2




[0181]




[0182] 稍微更具具体形式的一般合成流程2在以上合成方案2中示出。合成方案开始于SEM保护的4-氧基-4,5-二氢-1H-吡唑并[3,4-d]嘧啶-3-羧酸1,其可使用亚硫酰氯/DMF氯化,并且然后在轻微加热下与苄胺或吡咯烷偶合以提供SEM保护的化合物2。SEM保护的杂环2可在亲核芳族取代反应条件下,在极性溶剂诸如二噁烷中使用碱诸如二异丙基乙胺(DIPEA),使用氨基醇取代以提供胺取代的杂环3。SEM保护基可使用TBAF或在酸性条件下移除以提供最终化合物4。

[0183] 一般合成3




[0184]




[0185] 对于某些化合物,一般合成开始于4-氧基-4,5-二氢-1H-吡唑并[3,4-d]嘧啶-3-羧酸的合适氮保护(P)。羧酸可使用酰胺偶合条件来偶合至胺。所得化合物可使用亚硫酰氯来氯化,随后在合适条件下,例如在亲核芳族取代反应条件下,使用碱诸如二异丙基乙胺(DIPEA),在极性溶剂诸如二噁烷中,在嘧啶环上的卤化物处使用适当取代的环A取代以提供被环A取代的双环。移除保护基可提供式I化合物。以下描述的化合物可使用此一般合成来制备。此外,手性HPLC可用于拆分式I、(Ia)、(Ia-1)、(Ia-2)、(Ib)、(Ib-1)、(Ib-2)、II、(IIa)、(IIb)、(IIc)化合物的手性混合物。

[0186] 合成方案3




[0187]




[0188] 稍微更具体形式的一般合成流程3在以上合成方案3中示出。合成方案开始于SEM保护的4-氧基-4,5-二氢-1H-吡唑并[3,4-d]嘧啶-3-羧酸1,其可在酰胺偶合条件下与苄胺或吡咯烷偶合。SEM保护的杂环2可使用亚硫酰氯/DMF来氯化,并且然后在亲核芳族取代反应条件下在极性溶剂诸如二噁烷中使用碱诸如二异丙基乙胺(DIPEA),使用氨基醇取代以提供胺取代的杂环3。SEM保护基可使用TBAF或在酸性条件下移除以提供最终化合物4。

[0189] 图1列出的所有化合物,以及本发明的其他化合物使用三种一般合成流程之一和以上描绘的方案来制备。此外,手性HPLC可用于拆分式I、(Ia)、(Ia-1)、(Ia-2)、(Ib)、(Ib-1)、(Ib-2)、II、(IIa)、(IIb)、(IIc)化合物的手性混合物。某些特定合成实施例在实施例中阐明。


[0190] 实施例1.合成化合物45



[0191]




[0192] 步骤1:合成3-溴-4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶




[0194] 在0℃下向3-溴-4-氯-1H-吡唑并[3,4-d]嘧啶(10.00g,42.84mmol)于DMF(50.00mL)中的溶液分批添加NaH(2.57g,64.25mmol)。在搅拌0.5h之后,在0℃下将SEM-Cl(8.57g,51.40mmol)逐滴添加至反应物历时0.5h。将反应物缓慢加温至25℃并且搅拌16h。在TLC(PE:EtOAc=1:1,R_f=0.88)显示反应完成之后,通过50mL的H₂O来缓慢淬灭反应。将混合物用EtOAc(50mL*3)萃取,并且将有机层用盐水(20mL*3)洗涤,在Na₂SO₄上干燥并浓缩。通过硅胶柱色谱(PE:EtOAc=20:1)纯化残余物以得到呈无色油状的3-溴-4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶(6.20g,产率:39.80%)。¹H-NMR(400MHz,CDCl₃) δ_{ppm} 8.83(s,1H),5.82(s,2H),3.70(t,2H,J=8.4Hz),0.97(t,2H,J=8.4Hz),0.00(s,9H)。

[0195] 步骤2:合成(1R,2R)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)环戊-1-醇




[0197] 向3-溴-4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶(2.50g, 6.87mmol)和(1R,2R)-2-氨基环戊醇盐酸盐(945.38mg, 6.87mmol)于二噁烷(15mL)中的混合物中添加DIPEA(1.78g, 13.74mmol), 将反应混合物在70℃下搅拌16h。一旦TLC(PE:EtOAc=5:1)显示起始材料完全消耗, 就将混合物真空浓缩并且通过硅胶柱色谱(PE:EtOAc=30:1~10:1)纯化以提供呈黄色油状的(1R,2R)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)环戊-1-醇(2.10g, 4.41mmol, 产率: 64.22%)。¹H-NMR(400MHz, CD₃OD) δppm 8.41(s, 1H), 6.26(s, 1H), 5.70(s, 2H), 4.13-4.12(m, 2H), 3.69-3.65(m, 2H), 2.40-2.39(m, 1H), 2.20-2.18(m, 1H), 1.95-1.84(m, 2H), 1.72-1.61(m, 2H), 0.97(d, 2H, J=4.0Hz), 0.00(s, 9H)。

[0198] 步骤3: 合成4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸甲酯




[0200] 向(1R,2R)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)环戊-1-醇(2.10g, 4.90mmol)于DMF(10mL)和MeOH(15mL)中的混合物中一次性添加Pd(dppf)Cl₂(717.07mg, 980.00μmol)和Et₃N(1.49g, 14.70mmol), 并且将反应混合物在70℃下在CO(50psi)气氛下搅拌30h。一旦TLC(PE:EtOAc=1:1)和LCMS显示起始材料完全消耗, 将混合物过滤并且将滤液真空浓缩以获得呈黄色油状的4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸甲酯(2.70g, 粗产物), 其未进一步纯化即直接使用。¹H-NMR(400MHz, CDCl₃) δppm 8.85(s, 1H), 8.45(s, 1H), 5.83(s, 2H), 4.14(br. s, 2H), 4.10(s, 3H), 3.70(t, 2H, J=8.4Hz), 2.40-2.38(m, 1H), 2.20-2.17(m, 1H), 1.93-1.79(m, 4H), 0.98(t, 2H, J=8.4Hz), 0.00(s, 9H)。

[0201] 步骤4: 合成4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸




[0203] 向甲基4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸(2.70g,6.63mmol)于MeOH(10mL)中的混合物中添加H₂O(10mL),随后一次性添加NaOH(530.01mg,13.25mmol)。然后,将反应混合物在26℃下搅拌16h。一旦LCMS和TLC(PE:EtOAc=1:1)显示起始材料完全消耗,就通过在真空中浓缩来移除MeOH并且将残余物用EtOAc(8mL*2)洗涤。然后,添加HCl水溶液(1N)直至pH<7并且观察到白色沉淀物的形成。通过过滤来收集固体并且在真空中干燥以提供呈白色固体状的4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸(1.20g,产率:37.89%)。¹H-NMR(400MHz,CD₃OD)δppm 8.40(s,1H),5.80(s,2H),4.37-4.32(m,1H),4.22-4.19(m,1H),3.75(t,2H,J=8.0Hz),2.39-2.36(m,1H),2.11-2.06(m,1H),1.96-1.91(m,2H),1.78-1.73(m,2H),0.95(t,2H,J=8.0Hz),0.00(s,9H)。

[0204] 步骤5:合成((R)-2-(2,5-二氟苯基)吡咯烷-1-基)(4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮




[0206] 在20℃下向4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸(100.00mg,254.12μmol)和(2R)-2-(2,5-二氟苯基)吡咯烷(55.87mg,304.95μmol)于DMF(2mL)中的混合物中添加HATU(144.94mg,381.18μmol)和Et₃N(128.57mg,1.27mmol),并且在20℃下将反应物搅拌16h。在LCMS显示反应完成之后,将H₂O(5mL)添加至混合物,并且将反应物用EtOAc(10mL*3)萃取并且用盐水(5mL*3)洗涤。然后,将有机层在Na₂SO₄上干燥并且浓缩。通过制备型TLC(PE:EtOAc=1:1,R_f=0.5)纯化残余物以得到呈无色油状的((R)-2-(2,5-二氟苯基)吡咯烷-1-基)(4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮(20.00mg,产率:14.09%)。

[0207] 步骤6:合成((R)-2-(2,5-二氟苯基)吡咯烷-1-基)(4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮





[0209] 在20℃下向((R)-2-(2,5-二氟苯基)吡咯烷-1-基)(4-(((1R,2R)-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮(20.00mg, 35.80 μ mol)于二噁烷(20mL)中的溶液中添加TBAF(80.61mg, 358.00 μ mol),并且在80℃下将反应物加热16h。在TLC(EtOAc, R_f =0.1)显示反应完成之后,将溶液浓缩并且将10mL的H₂O添加至残余物。将溶液用EtOAc(10mL*3)萃取,并且将有机层在Na₂SO₄上干燥并且浓缩。通过酸性制备型HPLC(MeOH/H₂O/TFA溶剂系统)纯化残余物以得到呈棕色固体状的((R)-2-(2,5-二氟苯基)吡咯烷-1-基)(4-(((1R,2R)-2-羟环戊基)氨基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮(11.10mg, 产率:72.37%)。

[0210] 实施例2.合成化合物97和化合物98




[0211]




[0212] 步骤1:合成(1R,2R,4R)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-(甲磺酰基)环戊-1-醇

[0213]




[0214] 向3-溴-4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶(300.00mg, 824.83 μ mol)于二噁烷(10.00mL)中的混合物中添加DIPEA(319.80mg, 2.47mmol)和(1R,2R,4R)-2-氨基-4-(甲磺酰基)环戊-1-醇(195.71mg, 907.31 μ mol),并且将混合物在90℃下搅拌32h。在LCMS显示反应完成之后,将混合物浓缩以移除1,4-二噁烷,并且将残余物溶解于DCM(20mL)中。将有机层用水(10mL*4)洗涤,在Na₂SO₄上干燥,并且浓缩以得到呈白色固体状的(1R,2R,4R)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-(甲磺酰基)环戊-1-醇(350.00mg, 产率:83.78%)。¹H-NMR(400MHz, CDCl₃) δ ppm 8.43(s, 1H), 6.23(br. s, 1H), 5.71(s, 2H), 4.50(br. s, 1H), 4.26-4.24(m, 1H), 3.70-3.66(m, 3H), 2.97(br. s, 4H), 2.65(m, 1H), 2.37-2.20(m, 2H), 0.97(t, 2H, J=8.4Hz), 0.00(s, 9H)。

[0215] 步骤2:合成4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸甲酯



[0217] 向(1R,2R,4R)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-(甲磺酰基)环戊-1-醇(350.00mg, 691.03 μ mol)于MeOH(10.00mL)/DMF(2.00mL)中的混合物中添加Et₃N(139.85mg, 1.38mmol)和Pd(dppf)Cb(25.28mg, 34.55 μ mol)。在添加之后,将混合物在75℃下在CO(50Psi)下搅拌16h。一旦LCMS显示反应完成,将混合物浓缩以得到粗产物,通过制备型TLC(PE:EtOAc=0:1)纯化所述粗产物以得到呈红色固体状的4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸甲酯(250.00mg, 产率:74.50%)。


[0218] 步骤3:合成4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸



[0220] 向4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸甲酯(250.00mg, 514.80 μ mol)于MeOH(10.00mL)/H₂O(5.00mL)中的混合物中添加NaOH(41.18mg, 1.03mmol),将其在20℃下搅拌16h。一旦LCMS显示反应完成,就将混合物浓缩以移除MeOH。将水层用EtOAc(3mL*2)洗涤并且通过HCl(1M)酸化直至pH=4为止,然后将混合物过滤并且将滤饼在真空下干燥以提供呈


黑色/棕色固体状的4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸(180.00mg,产率:74.14%)。¹H-NMR(400MHz,CDCl₃) δppm 8.47(s,1H),5.77(s,2H),4.43-4.40(m,1H),4.17-4.12(m,1H),3.85-8.82(m,1H),3.69(t,2H,J=8.4Hz),3.03(s,3H),2.45-2.41(m,2H),1.99-1.91(m,2H),0.92(t,2H,J=8.4Hz),0.00(s,9H)。

[0221] 步骤4:合成N-(2,5-二氟苄基)-4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺



[0223] 向4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸(60.00mg,127.23umol)于DMF(2.00mL)中的混合物中添加DIPEA(16.44mg,127.23umol)、(2,5-二氟苄基)胺(36.42mg,254.46umol)和T₃P(40.48mg,127.23umol)。在添加之后,将混合物在20℃下搅拌1h,其中LCMS显示反应完成。将混合物添加至水(4mL)中并且用EtOAc(5mL*3)萃取,并且将有机层在Na₂SO₄上干燥并浓缩以得到呈红色油状的N-(2,5-二氟苄基)-4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺(50.00mg,粗产物)。


[0224] 步骤5:合成N-(2,5-二氟苄基)-4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺




[0226] 在20℃下,将DCM(5.00mL)中的N-(2,5-二氟苄基)-4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺(50.00mg,83.79umol)在TFA(5.00mL)的混合物中搅拌16h,然后LCMS显示反应完成。将混合物浓缩以得到粗产物,通过制备型HPLC(TFA)和手性HPLC来纯化所述粗产物(拆分异构物的保留时间分别为7.46min和9.20min)。获得呈白色固体状的N-(2,5-二氟苄基)-4-(((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺(2.80mg,产率:7.16%)和N-(2,5-二氟苄基)-4-(((1S,2S,4S)-2-羟基-4-(甲磺酰基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺(4.00mg,产率:10.23%)。这些化合物的LC-MS条件如下:流速=0.8mL·min⁻¹,流动相:从99%[水+0.375%v/v TFA]和1%[CH₃CN+

0.188%v/v TFA],在此条件下历时0.4min,然后在3.0min内变化至10% [水+0.375%v/v TFA]和90% [CH₃CN+0.188%v/v TFA],然后在0.45min内变化至100% [CH₃CN+0.188%v/v TFA],最后在0.01min内变化至99% [水+0.375%v/v TFA]和1% [CH₃CN+0.188%v/v TFA],然后在此条件下历时0.64min;纯度分别为98.887%和96.551%。


[0227] 实施例3.合成化合物20和化合物21



[0228]




[0229] 步骤1: (1S,2R,5S)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-5-氟环戊-1-醇



[0231] 向3-溴-4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶(600.00mg,1.65mmol)和2-氨基-5-氟-环戊醇(196.58mg,1.65mmol)于二噁烷(15mL)中的混合物中添加DIPEA(426.49mg,3.30mmol)。将混合物在110℃下搅拌16h,然后TLC(PE/EtOAc=1:1)显示反应完成。将混合物冷却至25℃并且在减压下在50℃下浓缩。向残余物中添加EtOAc(50mL),并且将有机相用H₂O(20mL*3)洗涤,在Na₂SO₄上干燥,过滤并且真空浓缩以提供(1S,2R,5S)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-5-氟环戊-1-醇(600.00mg,粗产物)。残余物未进一步纯化即直接用于下一步骤中。


[0232] 步骤2: 4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸甲酯



[0234] 在N₂下,向(1S,2R,5S)-2-((3-溴-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-5-氟环戊-1-醇(600.00mg,1.34mmol)于MeOH/DMF(20mL,v:


$v=2/1$ 中的溶液中添加 Pd(dppf)Cl₂ (49.17mg, 67.21 μ mol) 和 Et₃N (408.03mg, 4.03mmol)。在真空下将悬浮液脱气并且用 C0 吹扫几次。在 70℃ 下将混合物在 C0 (50psi) 下搅拌 16h, 然后 TLC (PE/EtOAc=1:1) 显示起始材料完全消耗。将反应混合物过滤并且将滤液浓缩以提供 4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸甲酯 (700.00mg, 粗产物)。所述粗产物未进一步纯化即使用。

[0235] 步骤3: 4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸



[0237] 向 4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸甲酯 (700.00mg, 1.65mmol) 于 MeOH/H₂O (15mL, $v/v=2/1$) 中的溶液中一次性添加 NaOH (132.00mg, 3.30mmol), 将其在 25℃ 下搅拌 2h。在 LCMS 显示反应完成之后, 将混合物在 40℃ 下减压浓缩。将水相调整至 pH=4 并且过滤以提供呈白色固体状的 4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸 (700.00mg, 粗产物)。

[0238] 步骤4: N-(2,5-二氟苄基)-4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺




[0240] 在 25℃ 下向 4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-羧酸 (100.00mg, 243.01 μ mol) 和 T₃P (231.97mg, 729.04 μ mol) 的混合物中添加 DMF (2.00mL) 中的 Et₃N (49.18mg, 486.03 μ mol), 随后在 10min 之后一次性添加 (2,5-二氟苄基)甲胺 (69.57mg, 486.03 μ mol)。在 25℃ 下将混合物搅拌 16h。在 LCMS 显示反应完成之后, 将混合物在 60℃ 下在减压下浓缩以提供 N-(2,5-二氟苄基)-4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺 (200mg, 粗产物)。残余物未纯化并且直接使用。

[0241] 步骤5: N-(2,5-二氟苄基)-4-(((1R,2S,3S)-3-氟-2-羟环戊基)氨基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺和 N-(2,5-二氟苄基)-4-(((1S,2R,3R)-3-氟-2-羟环戊基)氨基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺



[0244] 合成胺中间物

[0246]



[0250] 在N₂下,向(3aS,4S,6aR)-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-醇(70.00g,442.51mmol,1.00当量)、异吲哚啉-1,3-二酮(80.00g,543.74mmol,1.23当量)和PPh₃(175.00g,667.20mmol,1.51当量)于干燥甲苯(1.00L)中的搅拌混合物中逐滴添加

DIAD (135.00g, 667.62mmol, 1.51当量)。将所得混合物在80℃下在N₂下搅拌20h。在TLC (PE:EtOAc=3:1) 显示起始材料完全消耗之后,将混合物浓缩,并且通过硅胶柱色谱 (PE:EtOAc=80:1/50:1/20:1/10:1) 纯化残余物以得到呈白色固体状的2-((3aS,4R,6aR)-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基) 异吡啶-1,3-二酮 (90.00g, 产率: 70.79%)。¹H-NMR (400MHz, CDCl₃) δppm 7.82-7.80 (m, 2H), 7.72-7.70 (m, 2H), 5.03-4.96 (m, 2H), 4.61-4.60 (m, 1H), 2.28-2.20 (m, 2H), 1.94-1.85 (m, 2H), 1.50 (s, 3H), 1.31 (s, 3H)


[0251] 步骤3: 3aS,4R,6aR)-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-胺

[0252] 将2-((3aS,4R,6aR)-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基) 异吡啶-1,3-二酮 (90.00g, 313.25mmol, 1.00当量) 和NH₂NH₂·H₂O (32.00g, 626.50mmol, 2.00当量) 于EtOH (600.00mL) 中的混合物在80℃下搅拌16h。在TLC (PE:EtOAc=3:1) 显示起始材料完全消耗之后,将混合物过滤并浓缩,并且将EtOH (500mL) 添加至残余物。在浓缩以移除溶剂之后,添加PE (1000mL) 并且将混合物过滤并浓缩以得到呈黄色油状的 (3aS,4R,6aR)-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-胺 (41.00g, 粗产物), 其在静置时固化为黄色晶体。¹H-NMR (400MHz, CDCl₃) δppm 4.72 (t, 1H, J=5.2Hz), 4.18 (d, 1H, J=5.6Hz), 3.39 (d, 1H, J=4.0Hz), 2.01-1.93 (m, 2H), 1.78-1.77 (m, 1H), 1.40 (s, 3H), 1.38-1.35 (m, 1H), 1.26 (s, 3H), 1.10 (br.s, 2H)。

[0253] 步骤4: (1R,2S,3R)-3-氨基环戊烷-1,2-二醇盐酸盐

[0254] 将 (3aS,4R,6aR)-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-胺 (10.00g, 63.61mmol, 1.00当量) 于H₂O (55.00mL) 和HCl (5.00mL, 12M) 中的混合物在20℃下搅拌2h。TLC (EtOAc:MeOH=10:1) 显示起始材料完全消耗。将混合物浓缩以得到呈黄色固体状的 (1R,2S,3R)-3-氨基环戊烷-1,2-二醇盐酸盐 (9.20g, 产率: 94.15%)。¹H-NMR (400MHz, CD₃OD) δppm 4.03 (br.s, 1H), 3.90 (dd, 1H, J=8.4, 4.4Hz), 3.48-3.41 (m, 1H), 2.25-2.19 (m, 1H), 2.05-2.02 (m, 1H), 1.75-1.65 (m, 1H), 1.58-1.56 (m, 1H)。

[0255] 实施例5: 合成 (1R,2R,3S,4R,5S)-4-氨基二环[3.1.0]己烷-2,3-二醇



[0257] 步骤1: (3aS,4S,6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-醇

[0258] 向 (3aR,6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-酮 (90.00g, 583.81mmol, 1.00当量) 和CeCl₃·7H₂O (240.00g, 644.16mmol, 1.10当量) 于MeOH (2.00L) 中的0℃搅拌混合物中分批添加NaBH₄ (44.00g, 1.16mol, 1.99当量) 历时0.5h。在添加之后,在18℃下将混合物搅拌0.5h。TLC (PE:EtOAc=3:1) 显示起始材料完全消耗。将混合物浓缩,并且向残余物中添加EtOAc (2000mL) 并且将溶液在18℃下搅拌0.5h。然后将混合物过滤,将滤液在Na₂SO₄上干燥并且浓缩以得到呈淡黄色油状的粗产物 (3aS,4S,6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-醇 (79.00g, 粗产物), 其未进一步纯化即直接用于下一步骤中。¹H-NMR (400MHz, CDCl₃) δppm 5.88 (s, 2H), 5.01 (d, 1H, J=5.6Hz), 4.74 (t, 1H, J=5.6Hz), 4.55 (dd, 1H, J=9.6, 5.6Hz), 2.76 (d, 1H, J=9.6Hz), 1.43

(s, 3H), 1.40 (s, 3H)。

[0259] 步骤2: 3aR, 3bR, 4aS, 5S, 5aS)-2, 2-二甲基六氢环丙[3, 4]环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-醇

[0260] 向 (3aS, 4S, 6aR)-2, 2-二甲基-3a, 6a-二氢-4H-环戊[d][1, 3]间二氧杂环戊烯-4-醇 (40.00g, 256.11mmol, 1.00当量) 于DCM (50.00mL) 中的0℃搅拌混合物中逐滴添加ZnEt₂ (1M, 1.00L, 3.90当量)。在15min之后, 将CH₂I₂ (550.00g, 2.05mol, 8.02当量) 添加至混合物中, 将其在20℃下搅拌16h。TLC (PE:EtOAc=1:1) 显示起始材料消耗。通过饱和NH₄Cl溶液 (200mL) 淬灭混合物, 随后添加H₂O (500mL)。将混合物用DCM (500mL*5) 萃取, 并且将合并的有机层在Na₂SO₄中干燥并且浓缩以得到粗产物。通过硅胶柱色谱 (PE:EtOAc=0:1/100:1/80:1/50:1/20:1/10:1/5:1) 纯化粗产物以得到呈淡黄色油状的 (3aR, 3bR, 4aS, 5S, 5aS)-2, 2-二甲基六氢环丙[3, 4]-环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-醇 (18.00g, 产率: 41.29%)。¹H-NMR (400MHz, CDCl₃) δppm 4.87 (t, 1H, J=6.0Hz), 4.53-4.45 (m, 2H), 2.34 (br. s, 1H), 1.85-1.82 (m, 1H), 1.64-1.62 (m, 1H), 1.54 (s, 3H), 1.28 (s, 3H), 0.98-0.94 (m, 1H), 0.63-0.60 (m, 1H)。

[0261] 步骤3: 2-((3aR, 3bR, 4aS, 5R, 5aS)-2, 2-二甲基六氢环丙[3, 4]环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-基) 异吲哚啉-1, 3-二酮

[0262] 在N₂下向 (3aR, 3bR, 4aS, 5S, 5aS)-2, 2-二甲基六氢环丙[3, 4]-环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-醇 (10.00g, 58.75mmol, 1.00当量)、异吲哚啉-1, 3-二酮 (12.00g, 81.56mmol, 1.39当量) 和PPh₃ (24.00g, 91.50mmol, 1.56当量) 于干燥甲苯 (500.00mL) 中的搅拌混合物中逐滴添加DIAD (20.00g, 98.91mmol, 1.68当量)。将所得混合物在80℃下在N₂下搅拌20h。TLC (PE:EtOAc=3:1) 显示起始材料完全消耗。将混合物浓缩, 并且通过硅胶柱色谱 (PE:EtOAc=80:1/50:1/30:1/20:1/10:1) 纯化残余物以得到呈淡黄色油状的 2-((3aR, 3bR, 4aS, 5R, 5aS)-2, 2-二甲基六氢环丙[3, 4]-环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-基) 异吲哚啉-1, 3-二酮 (14.00g, 产率: 79.61%)。¹H-NMR (400MHz, CDCl₃) δppm 7.86-7.83 (m, 2H), 7.74-7.72 (m, 2H), 5.37-5.34 (m, 1H), 4.78-4.76 (m, 1H), 4.73 (s, 1H), 2.01-1.95 (m, 1H), 1.51 (s, 3H), 1.47-1.42 (m, 1H), 1.24 (s, 3H), 0.85-0.79 (m, 2H)。

[0263] 步骤4: (3aR, 3bR, 4aS, 5R, 5aS)-2, 2-二甲基六氢环丙[3, 4]环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-胺


[0264] 将 2-((3aR, 3bR, 4aS, 5R, 5aS)-2, 2-二甲基六氢环丙[3, 4]-环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-基) 异吲哚啉-1, 3-二酮 (14.00g, 46.77mmol, 1.00当量) 和NH₂NH₂·H₂O (4.78g, 93.54mmol, 2.00当量) 于EtOH (200.00mL) 中的混合物在70℃下搅拌16h。TLC (PE:EtOAc=3:1) 显示起始材料完全消耗。过滤混合物, 将滤液浓缩, 并且向残余物中添加EtOAc (20mL)。然后过滤混合物, 并且浓缩以得到呈黄色油状的 (3aR, 3bR, 4aS, 5R, 5aS)-2, 2-二甲基六氢环丙[3, 4]环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-胺 (7.00g, 产率: 88.45%)。¹H-NMR (400MHz, CDCl₃) δppm 5.06-5.03 (m, 1H), 4.30 (d, 1H, J=6.8Hz), 3.45 (s, 1H), 1.73-1.71 (m, 1H), 1.48 (s, 3H), 1.43-1.39 (m, 1H), 1.25 (s, 3H), 0.74-0.67 (m, 2H)。

[0265] 步骤5: (1R, 2R, 3S, 4R, 5S)-4-氨基二环[3.1.0]己烷-2, 3-二醇*HCl


[0266] 将 (3aR, 3bR, 4aS, 5R, 5aS)-2, 2-二甲基六氢环丙[3, 4]环戊[1, 2-d][1, 3]间二氧杂环戊烯-5-胺 (1.00g, 63.61mmol, 1.00当量) 于H₂O (5.500mL) 和HCl (0.5mL, 12M) 中的混合

物在15℃下搅拌2h。TLC (EtOAc:MeOH=10:1) 显示起始材料完全消耗。将混合物浓缩以得到呈黄色固体状的 (1R,2R,3S,4R,5S)-4-氨基二环[3.1.0]己烷-2,3-二醇*HCl (780mg g, 产率:79.9%)。

[0267] 实施例6:合成 (1S,2S,4S)-2-氨基-4-氟环戊-1-醇 (相对立体化学)




[0269] 步骤1: ((环戊-3-烯-1-基氧基) 甲基) 苯



[0271] 在0℃下向环戊-3-烯-1-醇 (60.00g, 713.27mmol) 于THF (600.00mL) 中的混合物中分批添加NaH (37.09g, 927.25mmol)。在起泡停止之后,在0℃下在45min时间段内逐滴添加溴甲基苯 (158.59g, 927.25mmol), 然后加温至25℃并且搅拌16h。TLC (PE/EtOAc=50/1) 显示反应完成。在低于5℃的温度下,用MeOH (120mL) 淬灭过量NaH。将混合物加温至25℃,用H₂O (600.00mL) 稀释,并且分离两个层。将水层用乙酸乙酯 (200mL*3) 萃取。将合并的有机相用盐水 (200mL) 洗涤,在Na₂SO₄上干燥,过滤并且真空浓缩。通过硅胶色谱 (PE/EtOAc=1/0) 纯化残余物以提供呈黄色油状的 ((环戊-3-烯-1-基氧基) 甲基) 苯 (120.00g, 产率: 96.56%)。¹H-NMR (400MHz, CDCl₃) δppm 7.38-7.28 (m, 5H), 5.72 (s, 2H), 4.52 (s, 2H), 4.35-4.30 (m, 1H), 2.64-2.59 (m, 2H), 2.50-2.46 (m, 2H)。


[0272] 步骤2: (1R,3S,5S)-3-(苄氧基)-6-氧杂双环[3.1.0]己烷和 (1R,3r,5S)-3-(苄氧基)-6-氧杂双环[3.1.0]己烷



[0274] 在0℃下向 ((环戊-3-烯-1-基氧基) 甲基) 苯 (120.00g, 688.71mmol) 于DCM (600.00mL) 中的混合物中一次性添加m-CPBA (297.68g, 1.38mol)。在25℃下将混合物搅拌


16h。TLC (PE:EtOAc=20:1) 显示反应完成。将混合物过滤并且通过添加饱和 Na_2SO_3 水溶液来减少过量 m -CPBA直至观察到阴性碘化淀粉测试为止。将混合物过滤并且真空浓缩。通过硅胶色谱 (PE:EtOAc=1:0, 20:1) 纯化残余物以提供呈黄色油状的 (1R,3R,5S)-3-(苄氧基)-6-氧杂双环[3.1.0]己烷 (56.00g, 产率:42.74%) 和 (1R,3S,5S)-3-(苄氧基)-6-氧杂双环[3.1.0]己烷 (37.00g, 产率:28.24%)。对 (1R,3S,5S)-3-(苄氧基)-6-氧杂双环[3.1.0]己烷进行光谱分析。 $^1\text{H-NMR}$ (400MHz, CDCl_3) δ_{ppm} 7.31-7.20 (m, 5H), 4.36 (s, 2H), 3.84-3.74 (m, 1H), 3.43 (s, 2H), 2.51-2.35 (m, 2H), 1.66-1.57 (m, 2H)。对 (1R,3R,5S)-3-(苄氧基)-6-氧杂双环[3.1.0]己烷进行光谱分析。 $^1\text{H-NMR}$ (400MHz, CDCl_3) δ_{ppm} 7.31-7.15 (m, 5H), 4.36 (s, 2H), 3.85-3.74 (m, 1H), 3.43 (s, 2H), 2.51-2.35 (m, 2H), 1.66-1.60 (m, 2H)。

[0275] 步骤3: (1S,2S,4R)-2-叠氮基-4-(苄氧基)环戊-1-醇




[0277] 在25℃下向 (1R,3S,5S)-3-(苄氧基)-6-氧杂双环[3.1.0]己烷 (20.00g, 105.13mmol) 于EtOH (760.00mL) 和 H_2O (230.00mL) 中的混合物中一次性添加 NH_4Cl (20.98g, 392.13mmol)、 NaN_3 (24.00g, 369.17mmol)。将混合物加热至80℃并且搅拌16小时。TLC (PE:EtOAc=10:1) 显示反应完成。将混合物冷却至25℃并且通过 N_2 来移除EtOH, 并且使用DCM (100mL*3) 萃取水相。将合并的有机相用 H_2O (30mL*3) 洗涤, 在 Na_2SO_4 上干燥, 过滤并且真空浓缩以提供呈黄色油状的 (1S,2S,4R)-2-叠氮基-4-(苄氧基)环戊-1-醇 (23.00g, 产率:93.79%)。 $^1\text{H-NMR}$ (400MHz, CDCl_3) δ_{ppm} 7.43-7.28 (m, 5H), 4.58-4.47 (m, 2H), 4.27-4.25 (m, 1H), 4.11-4.08 (m, 1H), 3.66-3.61 (m, 1H), 2.49-2.44 (m, 2H), 2.16-2.13 (m, 1H), 1.89 (br.s, 1H), 1.87-1.80 (m, 2H)。

[0278] 步骤4: (1S,2S,4R)-2-叠氮基-4-(苄氧基)环戊基乙酸酯




[0280] 在0℃下在30分钟时间段内在 N_2 下, 向 (1S,2S,4R)-2-叠氮基-4-(苄氧基)环戊-1-醇 (22.90g, 98.17mmol)、 Et_3N (59.60g, 589.02mmol) 于DCM (550mL) 中的溶液中逐滴添加乙酰氯 (38.53g, 490.85mmol) 于DCM (50mL) 中的溶液, 在此期间将温度保持在5℃以下。然后将反应混合物加温至25℃并且搅拌16h。TLC (PE:tOAc=10:1) 显示起始材料完全消耗。通过缓慢添加 H_2O (100mL) 来淬灭反应物。将有机相用饱和盐水 (50mL) 洗涤, 在 Na_2SO_4 上干燥, 过滤并且在真空中浓缩。通过硅胶柱色谱 (PE:EtOAc=100:1, 50:1) 纯化残余物以提供呈黄色油状的 (1S,2S,4R)-2-叠氮基-4-(苄氧基)环戊基乙酸酯 (17.00g, 产率:62.90%)。 $^1\text{H-NMR}$ (400MHz, CDCl_3) δ_{ppm} 7.37-7.28 (m, 5H), 5.24-5.12 (m, 1H), 4.51 (s, 2H), 4.14-4.11 (m, 1H), 3.88-3.85 (m, 1H), 2.45-2.40 (m, 1H), 2.36-2.32 (m, 1H), 2.07 (s, 3H), 1.95-1.88 (m, 2H)。

[0281] 步骤5: (1S,2S,4R)-2-((叔丁氧羰基)氨基)-4-羟环戊基乙酸酯




[0283] 在 N_2 下向 (1S,2S,4R)-2-叠氮基-4-(苄氧基)环戊基乙酸酯 (8.80g, 31.97mmol) 于 EtOH (100.00mL) 中的溶液中添加 Pd(OH)₂ (4.42g, 31.97mmol)。在真空下将悬浮液脱气并且用 H_2 吹扫几次。将混合物在 H_2 (50psi) 下在 70℃ 下搅拌 32h。TLC (PE:EtOAc=2:1) 显示起始材料完全消耗。过滤反应混合物并且浓缩滤液。通过硅胶色谱 (PE:EtOAc=10:1, 2:1) 纯化粗产物以得到呈黄色固体状的 (1S,2S,4R)-2-((叔丁氧羰基)氨基)-4-羟环戊基乙酸酯 (3.80g, 产率: 45.84%)。

[0284] 步骤6: (1S,2S,4S)-2-((叔丁氧羰基)氨基)-4-氟环戊基乙酸酯




[0286] 在 -70℃ 下在 N_2 下向 (1S,2S,4R)-2-((叔丁氧羰基)氨基)-4-羟环戊基乙酸酯 (2.57g, 9.91mmol) 于 DCM (150.00mL) 中的混合物中逐滴添加 DAST (2.40g, 14.87mmol)。在 -70℃ 下将混合物搅拌 30min。TLC (PE:EtOAc=2:1) 显示反应完成。将混合物冷却至 0℃ 并且添加 NaHCO₃ (5mL, 10%) 水溶液并且将其搅拌 10min。将水相用 EtOAc (15mL*2) 萃取, 并且将合并的有机相用盐水 (10mL) 洗涤, 在 Na₂SO₄ 上干燥, 过滤, 并且真空浓缩。通过硅胶色谱 (PE:EtOAc=20:1, 10:1) 纯化粗产物以得到呈黄色油状的 (1S,2S,4S)-2-((叔丁氧羰基)氨基)-4-氟环戊基乙酸酯 (700.00mg, 产率: 27.03%)。¹H-NMR (400MHz, CDCl₃) δ ppm 5.14 (s, 0.5H), 5.00 (br. s, 1H), 4.71 (s, 0.5H), 4.14-4.13 (m, 1H), 2.49-2.47 (m, 2H), 2.07-1.94 (m, 3H), 1.81-1.74 (m, 2H), 1.43-1.41 (m, 9H)。

[0287] 步骤7: ((1S,2S,4S)-4-氟-2-羟环戊基)氨基甲酸叔丁酯



[0289] 向 (1S,2S,4S)-2-((叔丁氧羰基)氨基)-4-氟环戊基乙酸酯 (700.00mg, 2.68mmol) 于 MeOH (20.00mL) 中的混合物中一次性添加 NaOH (160.80mg, 4.02mmol)。在 25℃ 下将混合物搅拌 1h。TLC (PE:EtOAc=3:1) 显示反应完成。将混合物在 30℃ 下在减压下浓缩以提供呈白色固体状的 ((1S,2S,4S)-4-氟-2-羟环戊基)氨基甲酸叔丁酯 (650.00mg, 粗产物)。


[0290] 步骤8: (1S,2S,4S)-2-氨基-4-氟环戊-1-醇 (相对立体化学)




[0292] 将 ((1S,2S,4S)-4-氟-2-羟环戊基)氨基甲酸叔丁酯 (650.00mg, 2.96mmol) 于 MeOH/HCl (20.00mL, 4M) 中的混合物在 25℃ 下搅拌 1h。TLC (PE:EtOAc=2:1) 显示反应完成。将混合物在 30℃ 下在减压下浓缩以提供呈白色固体状的 (1S,2S,4S)-2-氨基-4-氟环戊-1-醇 (相对立体化学) (400.00mg, 产率: 86.85%)。¹H-NMR (400MHz, DMSO-d₆) δ ppm 8.38 (br. s,

3H), 5.09 (d, 1H, $J=53.6\text{Hz}$), 4.09–4.03 (m, 1H), 3.34–3.30 (m, 1H), 2.44–2.39 (m, 1H), 2.21–2.16 (m, 1H), 1.95–1.87 (m, 1H), 1.75–1.66 (m, 1H)。


[0293] 实施例7:合成 (1R,2R,4R) -2-氨基-4-(甲磺酰基) 环戊-1-醇 (相对立体化学)



[0294]




[0295] 步骤1:制备 (1R,2R,4S) -2-((叔丁氧羰基) 氨基) -4-((甲磺酰基) 氧基) 环戊基乙酸酯




[0297] 在0℃下向 (1R,2R,4S) -2-((叔丁氧羰基) 氨基) -4-羟环戊基乙酸酯 (3.00g, 11.57mmol) 和Et₃N (4.68g, 46.28mmol) 于DCM (50.00mL) 中的混合物中逐滴添加MsCl (3.98g, 34.71mmol), 然后在20℃下将混合物搅拌3h。TLC (PE:EtOAc=1:1) 显示反应完成。将混合物用水 (100mL*3) 洗涤, 然后将有机层在Na₂SO₄上干燥并且浓缩以得到呈黄色固体状的 (1R,2R,4S) -2-((叔丁氧羰基) 氨基) -4-((甲磺酰基) 氧基) 环戊基乙酸酯 (3.5g, 粗产物: 100%)。

[0298] 步骤2: (1R,2R,4R) -2-((叔丁氧羰基) 氨基) -4-(甲硫) 环戊基乙酸酯




[0300] 向 (1R,2R,4S) -2-((叔丁氧羰基) 氨基) -4-((甲磺酰基) 氧基) 环戊基乙酸酯 (3.50g, 10.37mmol) 于DMF (30.00mL) 中的混合物中添加NaSMe (4.36g, 12.45mmol)。然后, 将混合物在90℃下搅拌2h, 并且TLC (PE:EtOAc=2:1) 显示反应完成。将混合物浓缩以得到呈黄色固体状的粗 (1R,2R,4R) -2-((叔丁氧基羰基) 氨基) -4-(甲硫) 环戊基乙酸酯 (3.00g, 粗产物100%)。

[0301] 步骤3: ((1R,2R,4R) -2-羟基-4-(甲硫) 环戊基) 氨基甲酸叔丁酯




[0303] 向 (1R,2R,4R)-2-((叔丁氧羰基)氨基)-4-(甲硫)环戊基乙酸酯 (3.00g, 10.37mmol) 于 MeOH (100.00mL) 中的混合物中添加 K_2CO_3 (2.87g, 20.74mmol)。在 25℃ 下将混合物搅拌 16h。TLC (PE:EtOAc=2:1) 显示反应完成。将混合物浓缩并且通过硅胶柱色谱 (PE:EtOAc=5:1-1:1) 纯化以得到呈白色固体状的 ((1R,2R,4R)-2-羟基-4-(甲硫)-环戊基) 氨基甲酸叔丁酯 (1.60g, 产率:62.38%)。 1H -NMR (400MHz, $CDCl_3$) δ ppm 4.10-3.99 (m, 2H), 3.85 (br.s, 1H), 3.14-3.11 (m, 1H), 2.50-2.46 (m, 1H), 2.17-2.10 (m, 4H), 1.86-1.82 (m, 1H), 1.81-1.67 (m, 1H), 1.45 (s, 9H)。

[0304] 步骤4: ((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基甲酸叔丁酯




[0306] 向 ((1R,2R,4R)-2-羟基-4-(甲硫)环戊基)氨基甲酸叔丁酯 (1.60g, 6.47mmol) 于 DCM (100.00mL) 中的混合物中添加 m-CPBA (3.49g, 16.18mmol)。在 25℃ 下将混合物搅拌 16h。在 TLC (PE:EtOAc=2:1) 显示起始材料完全消耗之后, 将混合物用饱和 Na_2SO_3 (aq. 20mL) 和饱和 $NaHCO_3$ (aq. 20mL*3) 洗涤。然后将有机层在 Na_2SO_4 上干燥, 浓缩, 用 PE (10mL) 洗涤, 过滤, 并且将滤饼在真空下干燥以得到呈白色固体状的 ((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基甲酸叔丁酯 (1.70g, 产率:94.06%)。 1H -NMR (400MHz, $CDCl_3$) δ ppm 4.62 (br.s, 1H), 4.07-4.04 (m, 1H), 3.81-3.79 (m, 1H), 3.48-3.45 (m, 1H), 2.82 (s, 3H), 2.57-2.40 (m, 2H), 2.11-2.07 (m, 1H), 1.90-1.87 (m, 2H), 1.38 (s, 9H)。

[0307] 步骤5: (1R,2R,4R)-2-氨基-4-(甲磺酰基)环戊-1-醇 (相对立体化学)




[0309] 将 ((1R,2R,4R)-2-羟基-4-(甲磺酰基)环戊基)氨基甲酸叔丁酯 (1.2g, 4.30mmol) 于 HCl/MeOH (10.00mL) 中的混合物在 25℃ 下搅拌 16h, 然后 LCMS 显示反应完成, 并且将混合物浓缩以得到呈黄色固体状的 (1R,2R,4R)-2-氨基-4-(甲磺酰基)环戊-1-醇 (1.0g, 粗产物:100%)。 1H -NMR (400MHz, CD_3OD) δ ppm 4.11-4.07 (m, 2H), 3.76 (br.s, 1H), 2.94 (s, 3H), 2.53 (br.s, 2H), 1.99 (br.s, 2H)。

[0310] 实施例8: 合成 (1S,2R,5R)-2-氨基-5-氟环戊-1-醇盐酸盐 (相对立体化学)




[0312] 步骤1: 环戊-2-烯-1-醇




[0314] 在15℃下向CeCl₃·7H₂O (24.00g, 64.42mmol) 于MeOH (120.00mL) 中的混合物中添加环戊-2-烯-1-酮 (5.00g, 60.90mmol)。在5min之后, 在0℃下将NaBH₄ (4.61g, 121.80mmol) 分批添加至混合物。将所得混合物在25℃下搅拌1h, 然后TLC (PE:EtOAc=5:1) 显示生成多个点并且部分起始材料保留。通过H₂O (100mL) 淬灭反应物并且将有机溶剂在真空中浓缩。向残余物中添加H₂O (300mL), 随后用MTBE (200mL*3) 萃取。将合并的有机层在Na₂SO₄上干燥并且在真空下浓缩以得到呈棕色油状的粗产物环戊-2-烯-1-醇 (3.00g, 粗产物)。其未进一步纯化即直接用于下一步骤中。¹H-NMR (400MHz, CDCl₃) δ_{ppm} 5.91 (d, 1H, J=4.8Hz), 5.77-5.76 (m, 1H), 4.79 (d, 1H, J=3.6Hz), 2.47-2.42 (m, 1H), 2.21-2.15 (m, 2H), 1.64-1.59 (m, 1H)。

[0315] 步骤2: ((环戊-2-烯-1-基氧基) 甲基) 苯




[0317] 在0℃下向环戊-2-烯-1-醇 (9.00g, 106.99mmol) 于THF (200.00mL) 中的混合物中分批添加NaH (6.80g, 170.11mmol)。在添加之后, 将混合物在20℃下搅拌0.5h, 然后在0℃下将BnBr (20.00g, 116.94mmol) 逐滴添加至混合物中。在20℃下将所得混合物搅拌2h。TLC (PE:EtOAc=20:1) 显示形成新材料 (R_f=0.6, 254nm)。在此时点, 添加H₂O (20mL), 随后用EtOAc (20mL*3) 萃取。将合并的有机物在Na₂SO₄上干燥并且浓缩以得到粗产物, 通过硅胶柱色谱 (PE:EtOAc=1:0/100:1/80:1) 纯化所述粗产物以得到呈黄色油状的((环戊-2-烯-1-基氧基) 甲基) 苯 (8.00g, 产率: 42.91%)。¹H-NMR (400MHz, CDCl₃) δ_{ppm} 7.34-7.25 (m, 5H), 6.02 (br.s, 1H), 5.88 (br.s, 1H), 4.66 (br.s, 1H), 4.55-4.47 (m, 2H), 2.52-4.48 (m, 1H), 2.27-2.24 (m, 1H), 2.16-2.13 (m, 1H), 1.87-1.84 (m, 1H)。

[0318] 步骤3: (1S, 2S, 5S) -2-(苄氧基) -6-氧杂双环[3.1.0]己烷



[0320] 在0℃下向((环戊-2-烯-1-基氧基) 甲基) 苯 (8.50g, 29.27mmol) 于DCM (50.00mL) 中的混合物中分批添加m-CPBA (13.50g, 58.67mmol)。在25℃下将混合物搅拌4h。一旦TLC (PE:EtOAc=10:1) 显示起始材料完全消耗, 就过滤混合物, 将滤液浓缩并且通过硅胶柱色谱 (PE:EtOAc=1:0/100:1/80:1/50:1) 纯化以得到粗产物。然后添加DCM (20mL), 将混合物过滤, 并且将H₂O (20mL) 和Na₂CO₃ (500mg) 添加至滤液, 随后在25℃下将混合物搅拌0.5h。然后将混合物用DCM (20mL*3) 萃取, 在Na₂SO₄上干燥, 并且浓缩以得到呈无色油状的(1S, 2S, 5S) -2-(苄氧基) -6-氧杂双环[3.1.0]己烷 (2.40g, 产率: 43.10%), 其通过NOE证实。¹H-NMR (400MHz, CDCl₃) δ_{ppm} 7.37-7.26 (m, 5H), 4.61-4.51 (m, 2H), 4.09 (d, 1H, J=5.2Hz), 3.55 (br.s, 1H), 3.49 (br.s, 1H), 1.99-1.95 (m, 1H), 1.87-1.75 (m, 2H), 1.54-1.52 (m, 1H)。

[0321] 步骤4: (1S, 2R, 5S) -2-叠氮基-5-(苄氧基) 环戊-1-醇




[0323] 向(1S,2S,5S)-2-(苄氧基)-6-氧杂双环[3.1.0]己烷(2.40g,12.62mmol)和NH₄Cl(1.55g,29.03mmol)于H₂O(3.00mL)和MeOH(24.00mL)中的混合物中添加NaN₃(4.10g,63.10mmol),将其在80℃下搅拌16h。在TLC(PE:EtOAc=10:1)显示起始材料消耗之后,通过N₂干燥有机溶剂并且将残余物用H₂O(20mL)稀释,用DCM(20mL*3)萃取。将合并的有机相用H₂O(10mL*3)洗涤,在Na₂SO₄上干燥,并且浓缩以得到呈棕色油状的(1S,2R,5S)-2-叠氮基-5-(苄氧基)环戊-1-醇(2.60g,产率:88.32%)。¹H-NMR(400MHz,CDCl₃) δppm 7.42-7.31(m,5H),4.64-4.55(m,2H),4.02-3.99(m,1H),3.82-3.80(m,1H),3.66-3.63(m,1H),2.25(br.s,1H),2.07-2.01(m,2H),1.80-1.77(m,2H)。

[0324] 步骤5:(((1S,2R,5S)-2-叠氮基-5-(苄氧基)环戊基)氧基)(叔丁基)二甲基硅烷



[0326] 将(1S,2R,5S)-2-叠氮基-5-(苄氧基)环戊-1-醇(2.50g,10.72mmol)、咪唑(1.61g,23.69mmol)和TBDMSCl(2.42g,16.08mmol)于CHCl₃(5.00mL)中的混合物在80℃下搅拌16h。一旦TLC(PE:EtOAc=10:1)显示起始材料完全消耗,将混合物浓缩并通过硅胶柱色谱(PE:EtOAc=1:0/100:1/80:1)纯化以得到呈无色油状的(((1S,2R,5S)-2-叠氮基-5-(苄氧基)环戊基)氧基)(叔丁基)二甲基硅烷(3.00g,产率:80.52%)。¹H-NMR(400MHz,CDCl₃) δppm 7.24-7.15(m,5H),4.40(d,2H,J=2.4Hz),3.88-3.86(m,1H),3.63-3.61(m,1H),3.47-3.43(m,1H),1.94-1.82(m,2H),1.70-1.65(m,2H),0.79(s,9H),0.03(s,3H),0.00(s,3H)。


[0327] 步骤6:((1R,2S,3S)-2-((叔丁基二甲基甲硅烷基)氧基)-3-羟环戊基)氨基甲酸叔丁酯



[0329] 向(((1S,2R,5S)-2-叠氮基-5-(苄氧基)环戊基)氧基)(叔丁基)二甲基硅烷(2.90g,8.34mmol)和Boc₂O(2.20g,10.10mmol)于MeOH(50.00mL)中的混合物中添加Pd(OH)₂(1.50g,5.42mmol),将其在50℃下在H₂(50psi)下搅拌20h。在TLC(PE:EtOAc=3:1)显示起始材料完全消耗之后,过滤混合物,并将滤液浓缩并且通过硅胶柱色谱(PE:EtOAc=10:1/8:1/5:1)纯化滤液以得到呈白色固体状的((1R,2S,3S)-2-((叔丁基二甲基甲硅烷基)氧基)-3-羟环戊基)氨基甲酸叔丁酯(2.10g,产率:75.95%)。¹H-NMR(400MHz,CDCl₃) δppm 4.81(br.s,1H),3.89-3.88(m,1H),3.70-3.67(m,2H),2.06-1.89(m,2H),1.58-1.56(m,2H),1.35(s,9H),0.79(s,9H),0.02(s,3H),0.00(s,3H)。


[0330] 步骤7:((1R,2S,3R)-2-((叔丁基二甲基甲硅烷基)氧基)-3-氟环戊基)氨基甲酸叔丁酯

叔丁酯



[0332] 在 -70°C 下向((1R,2S,3S)-2-((叔丁基二甲基甲硅烷基)氧基)-3-羟环戊基)氨基甲酸叔丁酯(1.10g,3.32mmol)于DCM(50.00mL)中的混合物添加DAST(1.61g,9.96mmol)。将反应混合物在 -70°C 下搅拌1h并且在 25°C 下搅拌1小时。在TLC(PE:EtOAc=5:1, $R_f=0.6$)显示反应完成之后,将冰水(5mL)添加至反应物中。将溶液用DCM(20mL*3)萃取并且用盐水(30mL)洗涤。将有机层在 Na_2SO_4 上干燥并且浓缩。通过硅胶柱色谱(PE:EtOAc=100:1)纯化残余物以得到呈白色固体状的((1R,2S,3R)-2-((叔丁基二甲基甲硅烷基)氧基)-3-氟环戊基)氨基甲酸叔丁酯(100.00mg,产率:9.03%)。 $^1\text{H-NMR}$ (400MHz, CDCl_3) δ_{ppm} 4.64(d,1H, $J=54.4\text{Hz}$),4.26(br.s,1H),3.90-3.86(m,1H),3.76-3.68(m,1H),2.11-1.79(m,4H),1.34(s,9H),0.81(s,9H),0.00(s,6H)。

[0333] 步骤8: (1S,2R,5R)-2-氨基-5-氟环戊-1-醇盐酸盐(相对立体化学)




[0335] 将((1R,2S,3R)-2-((叔丁基二甲基甲硅烷基)氧基)-3-氟环戊基)氨基甲酸叔丁酯(100.00mg,299.84 μmol)于HCl/MeOH(20.00mL,4M)中的溶液在 25°C 下搅拌16h。TLC(PE:EtOAc=5:1, $R_f=0$)显示反应完成。通过 N_2 干燥溶液,并且获得呈黄色固体状的(1S,2R,5R)-2-氨基-5-氟环戊-1-醇盐酸盐(相对立体化学)(45.00mg,产率:96.45%)。 $^1\text{H-NMR}$ (400MHz, CD_3OD) δ_{ppm} 4.93-4.89(m,1H),3.98-3.88(m,1H),3.52-3.47(m,1H),2.30-2.00(m,3H),1.66-1.62(m,1H)。


[0336] 实施例9:合成(1S,2R,5S)-2-氨基-5-氟环戊-1-醇(相对立体化学)



[0337]




[0338] 步骤1: (1R,2R,5S)-N,N-联苄基-6-氧杂双环[3.1.0]己-2-胺



[0340] 将 Cl_3CCOOH (154.72g,949.20mmol)添加至N,N-联苄基环戊-2-烯-1-胺(50.00g,


189.84mmol) 于DCM (640mL) 中的搅拌溶液中并且将所得混合物在20℃下搅拌0.1h。一次性添加m-CPBA (43.00g, 199.33mmol) 并且将反应混合物在20℃下持续搅拌16h。在TLC (PE: EtOAc=10:1) 显示反应完成之后, 将混合物用DCM (500mL) 稀释并且添加Na₂SO₃饱和水溶液直至碘化淀粉纸指示没有留下m-CPBA为止。添加NaHCO₃ (500mL) 饱和水溶液并且将各层分离。将有机层用NaHCO₃ (200mL*2) 水溶液洗涤, 然后干燥, 浓缩, 并且通过硅胶柱色谱 (PE: EtOAc=100:1-50:1) 纯化以得到呈白色固体状的 (1R, 2R, 5S) -N,N-联苄基-6-氧杂双环 [3.1.0] 己-2-胺 (30.00g, 产率: 56.56%)。¹H-NMR (400MHz, CDCl₃) δppm 7.41 (d, 4H, J=7.2Hz), 7.31 (t, 4H, J=7.6Hz), 7.25-7.22 (m, 2H), 3.86-3.70 (m, 4H), 3.44 (s, 1H), 3.32 (s, 1H), 3.28-3.24 (m, 1H), 2.04-2.01 (m, 1H), 1.54-1.45 (m, 3H)。

[0341] 步骤2: (1R, 2R, 3R) -3-(联苄基氨基) -2-羟环戊基乙酸酯




[0343] 将 (1R, 2R, 5S) -N,N-联苄基-6-氧杂双环 [3.1.0] 己-2-胺 (30.00g, 107.38mmol, 1.00当量) 于AcOH (200mL) 中的溶液在50℃下搅拌16h。在TLC (PE: EtOAc=10:1) 显示反应完成之后, 将混合物浓缩以移除AcOH, 将残余物溶解于DCM (100mL) 中, 并且将有机层用NaHCO₃ (100mL*3) 水溶液洗涤, 并且在Na₂SO₄上干燥并浓缩。通过硅胶柱色谱 (PE: EtOAc=100:1-50:1) 纯化残余物以得到呈白色固体状的 (1R, 2R, 3R) -3-(联苄基氨基) -2-羟环戊基乙酸酯 (20.00g, 产率: 54.87%)。¹H-NMR (400MHz, CDCl₃) δppm 7.36-7.28 (m, 10H), 5.05-5.02 (m, 1H), 4.05 (d, 1H, J=4.0Hz), 3.81-3.69 (m, 4H), 3.27-3.24 (m, 1H), 2.40-2.36 (m, 1H), 2.07 (s, 3H), 1.97-1.94 (m, 1H), 1.78-1.73 (m, 1H), 1.60-1.54 (m, 1H)。

[0344] 步骤3: (1R, 2R, 3R) -3-(联苄基氨基) -2-((甲磺酰基) 氧基) 环戊基乙酸酯



[0346] 将MsCl (8.10g, 70.71mmol) 逐滴添加至 (1R, 2R, 3R) -3-(联苄基氨基) -2-羟环戊基乙酸酯 (20.00g, 58.92mmol)、Et₃N (18.48g, 182.66mmol) 和DMAP (719.83mg, 5.89mmol) 于DCM (200mL) 中的混合物中。在添加之后, 将混合物在20℃下搅拌16h。在TLC (PE: EtOAc=10:1) 显示反应完成之后, 将混合物用水 (100mL*2) 洗涤, 将有机层在Na₂SO₄上干燥, 并且通过硅胶柱色谱 (PE: EtOAc=80:1-60:1) 纯化以得到呈白色固体状的 (1R, 2R, 3R) -3-(联苄基氨基) -2-((甲磺酰基) 氧基) 环戊基乙酸酯 (15.00g, 产率: 60.97%)。¹H-NMR (400MHz, CDCl₃) δppm 7.41 (d, 4H, J=7.6Hz), 7.33 (t, 4H, J=6.8Hz), 7.31-7.25 (m, 2H), 5.16-5.13 (m, 1H), 5.01-5.00 (m, 1H), 3.92-3.81 (m, 4H), 3.37-3.32 (m, 1H), 3.15 (s, 3H), 2.33-2.30 (m, 1H), 2.01 (s, 3H), 1.98-1.91 (m, 2H), 1.56-1.53 (m, 1H)。

[0347] 步骤4: (1S, 2R, 5R) -N,N-联苄基-6-氧杂双环 [3.1.0] 己-2-胺




[0349] 将 K_2CO_3 (5.96g, 43.11mmol) 添加至 (1R, 2R, 3R) -3-(联苄基氨基)-2-((甲磺酰基)氧基)环戊基乙酸酯 (15.00g, 35.93mmol) 于 MeOH (70mL) / THF (30mL) 中的混合物中。在 20℃ 下将混合物搅拌 16h。在 TLC (PE:EtOAc = 10:1) 显示反应完成之后, 将混合物浓缩以移除 MeOH 和 THF。然后将混合物溶解于 DCM (20mL) 中, 将有机层用水 (10mL*2) 洗涤, 在 Na_2SO_4 上干燥, 并且浓缩以得到呈白色固体状的 (1S, 2R, 5R) -N,N-联苄基-6-氧杂双环[3.1.0]己-2-胺 (10.00g, 产率: 99.62%)。 1H -NMR (400MHz, $CDCl_3$) δ ppm 7.37 (d, 4H, $J=7.6$ Hz), 7.32 (d, 4H, $J=7.2$ Hz), 7.29-7.23 (m, 2H), 3.73-3.69 (m, 2H), 3.53-3.41 (m, 5H), 2.06-2.00 (m, 1H), 1.91-1.90 (m, 1H), 1.87-1.76 (m, 1H), 1.51-1.48 (m, 1H)。

[0350] 步骤5: (1S, 2R, 5S) -2-(联苄基氨基)-5-氟环戊-1-醇



[0352] 向 (1S, 2R, 5R) -N,N-联苄基-6-氧杂双环[3.1.0]己-2-胺 (6.50g, 23.27mmol) 于 DCM (200mL) 中的混合物中添加 $HBF_4 \cdot Et_2O$ (7.54g, 46.53mmol), 并且将混合物在 30℃ 下搅拌 0.2h。在 TLC 显示反应完成之后, 将混合物添加至 Na_2CO_3 (100mL) 并且用 DCM (200mL*2) 萃取。将有机层在 Na_2SO_4 上干燥, 浓缩, 并且通过硅胶柱色谱 (PE:EtOAc = 60:1 至 20:1) 纯化以得到呈白色固体状的 (1S, 2R, 5S) -2-(联苄基氨基)-5-氟环戊-1-醇 (1.80g, 产率: 25.84%)。 1H -NMR (400MHz, $CDCl_3$) δ ppm 7.38-7.22 (m, 10H), 4.81-4.67 (m, 1H), 4.23-4.15 (m, 1H), 3.83-3.55 (m, 4H), 3.04-2.98 (m, 1H), 1.93-1.79 (m, 4H)。


[0353] 步骤6: (1S, 2R, 5S) -2-氨基-5-氟环戊-1-醇 (相对立体化学)




(相对立体化学)

[0355] 向 (1S, 2R, 5S) -2-(联苄基氨基)-5-氟环戊-1-醇 (1.60g, 5.34mmol) 于 MeOH (10mL) 中的混合物中添加 $Pd(OH)_2$ (500.00mg, 3.61mmol), 并且将混合物在 30℃ 下在 H_2 (30psi) 下搅拌 16h。在 TLC 显示反应完成之后, 通过硅藻土过滤混合物并且将滤液浓缩以得到呈白色固体状的 (1S, 2R, 5S) -2-氨基-5-氟环戊-1-醇 (600.00mg, 产率: 94.31%)。 1H -NMR (400MHz, $CDCl_3$) δ ppm 4.93-4.77 (m, 1H), 3.92-3.84 (m, 1H), 3.10-3.04 (m, 1H), 2.07-1.98 (m, 3H), 1.64-1.61 (m, 1H)。

[0356] 实施例10: 合成3-氟-5-((2R, 4S) -4-氟吡咯烷-2-基)吡啶




[0358] 步骤1: (R)-4-((叔丁基二甲基甲硅烷基)氧基)吡咯烷-2-酮



[0360] 在0℃下向(R)-4-羟吡咯烷-2-酮(9.0g, 89.1mmol)于DMF(50mL)中的混合物中一次性添加咪唑(9.09g, 134mmol)和TBDMSCl(14.1g, 93.6mmol)。在25℃下将反应混合物搅拌3h。TLC(DCM/MeOH=10/1, R_f =0.8)显示反应完成, 然后添加水(200mL), 通过过滤收集所得沉淀物并且在真空中干燥以得到呈白色固体状的(R)-4-((叔丁基二甲基甲硅烷基)氧基)吡咯烷-2-酮(15.5g, 产率: 80.7%)。 $^1\text{H-NMR}$ (400MHz, CDCl_3) δ ppm 5.90 (br. s, 1H), 4.55-4.53 (m, 1H), 3.60-3.56 (m, 1H), 3.24-3.21 (m, 1H), 2.56-2.50 (m, 1H), 2.28-2.23 (m, 1H), 0.87-0.85 (m, 9H), 0.06-0.00 (m, 6H)。


[0361] 步骤2: 制备(R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-氧基吡咯烷-1-羧酸叔丁酯



[0363] 在0℃下向(R)-4-((叔丁基二甲基甲硅烷基)氧基)吡咯烷-2-酮(15.5g, 72.0mmol)于 CH_3CN (150mL)中的混合物中一次性添加 Et_3N (8.72g, 86.4mmol)、DMAP(4.39g, 36mmol)和 Boc_2O (20.4g, 93.7mmol)。将反应混合物在25℃下搅拌10h。TLC(PE/EtOAc=3/1)显示反应完成, 然后添加水(600mL), 通过过滤收集所得沉淀物并且在真空中干燥以得到呈粉红色固体状的(R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-氧基吡咯烷-1-羧酸叔丁酯(19.2g, 产率: 84.6%)。 $^1\text{H-NMR}$ (400MHz, CDCl_3) δ ppm 4.33-4.30 (m, 1H), 3.81-3.77 (m, 1H), 3.56-3.54 (m, 1H), 2.67-2.61 (m, 1H), 2.41-2.37 (m, 1H), 1.46 (s, 9H), 0.80 (s, 9H), 0.00 (s, 6H)。


[0364] 步骤3: ((2R)-2-((叔丁基二甲基甲硅烷基)氧基)-4-(5-氟吡啶-3-基)-4-羟基丁基)氨基甲酸叔丁酯

[0365]




[0366] 在0℃下在30分钟内向3-溴-5-氟-吡啶(3.35g, 19.02mmol, 1.20当量)于THF(40.00mL)中的混合物中逐滴添加*i*-PrMgCl-LiCl(1.3M, 17.56mL, 1.44当量)(放热)。在添加之后,在1h内温度升高至25℃并且在25℃下搅拌30分钟。TLC(PE/EtOAc=10/1)显示生成新的点,这指示Mg试剂成功地制备。然后在-78℃下在30分钟内将于THF(50mL)中的(R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-氧基吡咯烷-1-羧酸叔丁酯(5.00g, 15.85mmol, 1.00当量)逐滴添加至溶液。使混合物在1h内加温至25℃,然后在25℃下搅拌16h。TLC(PE/EtOAc=3/1)显示起始材料完全消耗并且检测到所需产物(R)-(2-((叔丁基二甲基甲硅烷基)氧基)-4-(5-氟吡啶-3-基)-4-氧代丁基)氨基甲酸叔丁酯。通过在0℃下添加MeOH(50mL)来淬灭反应物。在0℃下添加NaBH₄(1.20g, 31.70mmol, 2.00当量),然后将混合物在25℃下搅拌4h。TLC(PE/EtOAc=2/1)和LCMS显示反应完成。将合并的反应混合物(4个平行反应)通过NH₄Cl(400mL)水溶液淬灭并且用EtOAc(600mL*3)萃取。将合并的有机物在Na₂SO₄上干燥并且在真空中浓缩,并且通过HPLC纯化残余物以得到呈黄色油状的((2R)-2-((叔丁基二甲基甲硅烷基)氧基)-4-(5-氟吡啶-3-基)-4-羟基丁基)氨基甲酸叔丁酯(1.24g, 产率: 18.91%)。¹H-NMR(400MHz, CDCl₃) δppm 8.26-8.22(m, 2H), 7.37(d, 1H, J=8.8Hz), 4.95-4.88(m, 2H), 4.69(br. s, 1H), 4.00-3.98(m, 2H), 3.23-3.10(m, 2H), 1.73(br. s, 2H), 1.32(s, 9H), 0.80-0.79(m, 9H), 0.00(s, 6H)。

[0367] 步骤4: (4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯




[0369] 在-60℃下在0.5h内向((2R)-2-((叔丁基二甲基甲硅烷基)氧基)-4-(5-氟吡啶-3-基)-4-羟基丁基)氨基甲酸叔丁酯(8.70g, 20.98mmol, 1.00当量)和Et₃N(31.84g, 314.70mmol, 15.00当量)于DCM(500.00mL)中的混合物中逐滴添加MsCl(31.24g, 272.74mmol, 13.00当量)。然后将混合物在-60℃下搅拌1h,并且将反应混合物加温至25℃并且搅拌18h。LCMS显示起始材料完全消耗。然后将混合物用H₂O(200mL*3)洗涤,并且将水相用DCM(200mL*4)萃取。将合并的有机层在Na₂SO₄上干燥并且在真空中浓缩以得到呈黑色/棕色油状的粗产物(4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(8.30g, 粗产物),其未纯化即直接使用。

[0370] 步骤5: (4R)-2-(5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯




[0372] 在25℃下向(4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(8.30g,20.93mmol,1.00当量)于THF(250.00mL)中的混合物中添加TBAF(9.43g,41.86mmol,2.00当量)。在25℃下将混合物搅拌16h。在TLC(PE/EtOAc=1/1)显示反应完成之后,将混合物浓缩并且将残余物溶解于EtOAc(600mL)中,用水(200mL*5)洗涤,在Na₂SO₄上干燥,并浓缩。通过PLC纯化粗产物以得到呈棕色黑色油状的(4R)-2-(5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯(4.70g,16.65mmol,产率:79.54%)。¹H-NMR(400MHz,CDCl₃) δ ppm 8.37-8.33(m,2H),7.48(br.s,1H),5.09-4.89(m,1H),4.56-4.54(m,1H),3.80-3.65(m,2H),2.63-2.43(m,1H),2.03-1.96(m,1H),1.56-1.20(m,9H)。

[0373] 步骤6:(2R,4S)-4-氟-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯



[0375] 在-78℃下在0.5h内向(4R)-2-(5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯(4.70g,16.65mmol,1.00当量)于DCM(150.00mL)中的混合物中逐滴添加DAST(29.52g,183.15mmol,11.00当量)。将反应混合物在-78℃下搅拌2h,然后将其加温至25℃并且搅拌20h。在TLC(PE/EtOAc=0/1)显示起始材料完全消耗之后,将混合物冷却至0℃并且通过饱和NaHCO₃溶液(100mL)来逐滴淬灭。将有机相分离并且在Na₂SO₄上干燥,浓缩以得到残余物,然后通过硅胶柱色谱(PE:EtOAc从10:1、8:1至5:1,然后3:1)纯化以得到呈白色固体状的(2R,4S)-4-氟-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(1.38g,4.85mmol,产率:29.15%,R_f=0.53)和呈黄色油状的(2S,4S)-4-氟-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(1.36g,4.78mmol,产率:28.73%,R_f=0.43)。¹H-NMR(400MHz,CDCl₃) δ ppm 8.31-8.27(m,2H),7.20-7.18(m,1H),5.18(d,1H,J=51.6Hz),4.97-4.88(m,1H),4.04-4.00(m,1H),3.64(dd,1H,J=38.8,12.8Hz),2.67(dd,1H,J=15.6,6.8Hz),1.97-1.67(m,1H),1.56-1.12(m,9H)。


[0376] 步骤7:3-氟-5-((2R,4S)-4-氟吡咯烷-2-基)吡啶




[0378] 在0℃下向(2R,4S)-4-氟-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(1.38g,4.85mmol,1.00当量)于EtOAc(10mL)中的混合物中逐滴添加HCl/EtOAc(40.00mL,4M)。将混合物加温至25℃并且搅拌3h。在TLC(PE:EtOAc=1:1)显示反应完成之后,将溶剂蒸发以得到呈棕色固体状的3-氟-5-((2R,4S)-4-氟吡咯烷-2-基)吡啶(1.25g,4.86mmol,产率:

100.00%)。¹H-NMR (400MHz, CD₃OD) δppm 8.84-8.81 (m, 2H), 8.31 (d, 1H, J=9.2Hz), 5.62 (dt, 1H, J=52.0, 2.4Hz), 5.23-5.18 (m, 1H), 4.00-3.95 (m, 1H), 3.88-3.71 (m, 1H), 2.67 (td, 1H, J=16.0, 6.0Hz), 1.69-1.59 (m, 1H)。

[0379] 实施例11:合成 (3R,4S,5R)-5-氨基四氢-2H-吡喃-3,4-二醇



[0380]



[0381] 步骤1: (S)-3,6-二氢-2H-吡喃-3-基乙酸酯

[0382] 将 (3R,4S)-3,4-二氢-2H-吡喃-3,4-二基二乙酸酯 (2.9g, 14.49mmol) 溶解于 DCM (15ml) 中, 并且在 N₂ 下在室温下搅拌。添加三乙基硅烷 (2.55ml, 15.93mmol) 并且搅拌 5 分钟。逐滴添加 BF₃·OEt₂ (1.836ml, 14.49mmol) 并且继续搅拌 30 分钟。将反应混合物用 30ml 饱和碳酸氢盐淬灭, 并且分离各层。将合并的有机层在硫酸钠上干燥并且将溶剂移除。通过快速色谱 (0-30% Hex/EtOAc) 纯化残余物。回收呈透明油状的 (S)-3,6-二氢-2H-吡喃-3-基乙酸酯 (1.9g, 92% 产率)。¹H NMR (400MHz, DMSO-d₆) δ 6.10 (dddt, J=10.2, 3.2, 2.1, 1.0Hz, 1H), 5.84 (ddt, J=10.1, 4.3, 2.1Hz, 1H), 4.96 (dtd, J=4.3, 2.7, 1.5Hz, 1H), 4.16-3.89 (m, 2H), 3.73 (t, J=2.9Hz, 2H), 2.01 (d, J=0.9Hz, 3H)。

[0383] 步骤2: (S)-3,6-二氢-2H-吡喃-3-醇

[0384] 将 (S)-3,6-二氢-2H-吡喃-3-基乙酸酯 (1.9g, 13.37mmol) 溶解于 MeOH (30ml) 和水 (20ml) 中。添加三乙胺 (7ml, 50.2mmol) 并且在室温下搅拌 30min。将溶剂在减压下移除。然后将残余水用 EtOAc 萃取三次。将有机层合并, 在硫酸钠上干燥并且将溶剂移除。回收呈透明油状的 (S)-3,6-二氢-2H-吡喃-3-醇 (1.1g, 10.99mmol, 82% 产率)。所述粗产物未进一步纯化而继续使用。

[0385] 步骤3: (1S,5R,6R)-3,7-二氧杂双环[4.1.0]庚-5-醇

[0386] 将 (S)-3,6-二氢-2H-吡喃-3-醇 (1.1g, 10.99mmol) 溶解于 CH₂Cl₂ (20ml) 中并且将其冷却至 0°C。分批添加 mCPBA (4.55g, 13.18mmol)。在加温至室温的同时将反应混合物搅拌过夜。将反应混合物的白色沉淀物过滤掉, 保留洗脱剂, 将溶剂移除并且用乙醚研磨。重复此步骤。残余物 (1S,5R,6R)-3,7-二氧杂双环[4.1.0]庚-5-醇 (1.2g, 100% 产率) 未进一步纯化而继续使用。

[0387] 步骤4: (3R,4S,5R)-5-(((R)-1-苯基乙基)氨基)四氢-2H-吡喃-3,4-二醇

[0388] 将 (1S,5R,6R)-3,7-二氧杂双环[4.1.0]庚-5-醇 (1.26g, 10.85mmol)、(R)-1-苯基乙胺 (1.658ml, 13.02mmol) 溶解于 2-BuOH (15ml) 中。将反应混合物加热至 100°C, 持续 18 小时。将反应混合物冷却至室温, 移除溶剂, 并且然后在 ISCO 0-100% EtOAc 上纯化残余物。将级分合并, 将溶剂移除, 并且然后将残余物用 MTBE 处理并搅拌过夜。将有机混合物的白色沉


淀物过滤掉。回收 (3R,4S,5R)-5-((R)-1-苯基乙基)氨基)四氢-2H-吡喃-3,4-二醇 (0.350g, 14% 产率)。¹H NMR (400MHz, DMSO-d₆) δ 7.38-7.24 (m, 4H), 7.22-7.12 (m, 1H), 4.61 (d, J=5.6Hz, 1H), 4.46 (d, J=4.8Hz, 1H), 3.88 (q, J=6.5Hz, 1H), 3.64 (tt, J=5.0, 2.8Hz, 1H), 3.47 (dd, J=11.4, 4.7Hz, 1H), 3.39 (ddd, J=8.4, 5.6, 3.1Hz, 1H), 3.31 (s, 2H), 3.29 (t, J=3.0Hz, 1H), 3.25 (d, J=2.5Hz, 0H), 2.79 (dd, J=11.1, 7.8Hz, 1H), 2.57 (td, J=7.8, 3.9Hz, 1H), 1.86 (s, 1H), 1.21 (d, J=6.6Hz, 3H)。

[0389] 步骤5: (3R,4S,5R)-5-氨基四氢-2H-吡喃-3,4-二醇

[0390] 将 (3R,4S,5R)-5-((R)-1-苯基乙基)氨基)四氢-2H-吡喃-3,4-二醇 (0.350g, 1.475mmol) 溶解于 EtOH (3ml) 中并且添加 Pd-C (0.031g, 0.295mmol)。将反应混合物在 H₂ 气球下搅拌过夜。通过硅藻土过滤反应混合物并且将溶剂移除以得到呈米黄色固体状的 (3R,4S,5R)-5-氨基四氢-2H-吡喃-3,4-二醇 (0.190g, 1.427mmol, 97% 产率)。所述粗产物未进一步纯化即继续使用。LCMS (M+H) 134。


[0391] 实施例12. (R)-3-(4,4-二氟吡咯烷-2-基)-5-氟吡啶

[0392] 步骤1: (2R,4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯



[0394] 在-60℃下在30分钟内向 ((2R)-2-((叔丁基二甲基甲硅烷基)氧基)-4-(5-氟吡啶-3-基)-4-羟基丁基)氨基甲酸叔丁酯 (6.80g, 16.40mmol) 和 Et₃N (24.89g, 246.00mmol) 于 DCM (500.00mL) 中的混合物中逐滴添加 MsCl (24.42g, 213.20mmol)。在-60℃下将混合物搅拌 1h。将反应混合物加温至 25℃ 并且再搅拌 18h。将混合物用 H₂O (200mL*3) 洗涤。将水相用 DCM (200mL*4) 萃取。将合并的有机层在 Na₂SO₄ 上干燥并且在真空中浓缩。通过硅胶色谱 (PE:EtOAc=50/1, 20/1, 10/1) 纯化残余物以提供呈棕色油状的 (2S,4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯 (2.70g, 产率:41.52%) 和 (2R,4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯 (2.40g, 产率:36.89%)。¹H-NMR (400MHz, CDCl₃) δppm 8.40 (br. s, 2H), 7.56-7.45 (m, 1H), 5.11-4.94 (m, 2H), 4.53 (br. s, 1H), 3.85-3.79 (m, 1H), 3.66-3.53 (m, 1H), 2.62-2.58 (m, 1H), 2.04-2.01 (m, 1H), 1.56 (s, 3H), 1.32 (s, 6H), 0.99-0.88 (m, 9H), 0.18-0.00 (m, 6H)。


[0395] 步骤2: (2R,4R)-2-(5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯



[0397] 在 25℃ 下向 (2R,4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯 (2.40g, 6.05mmol) 于 THF (60.00mL) 中的混合物中一次性添加 TBAF (3.16g, 12.10mmol)。将混合物在 50℃ 下在减压下浓缩。将残余物添加至水 (20mL) 中。将水


相用乙酸乙酯(30mL*3)萃取。将合并的有机相用饱和盐水(20mL*2)洗涤,在Na₂SO₄上干燥,过滤并且在真空中浓缩。通过硅胶色谱(PE:EtOAc=20/1,10/1,1/3)纯化残余物以提供呈黄色固体状的(2R,4R)-2-(5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯(1.30g,产率:76.11%)。¹H-NMR(400MHz,CDC13) δ_{ppm} 8.26(d,2H,J=12.8Hz),7.39(br.s,1H),4.95-4.81(m,1H),4.48-4.47(m,1H),3.73(br.s,1H),3.56-3.53(m,1H),2.55(br.s,1H),1.97-1.98(m,1H),1.65-1.16(m,9H)。

[0398] 步骤3: (R)-2-(5-氟吡啶-3-基)-4-氧基吡咯烷-1-羧酸叔丁酯




[0400] 在-10℃下向(2R,4R)-2-(5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯(1.30g,4.60mmol)和三氯异氰尿酸(1.10g,4.60mmol)的混合物中添加TEMPO(72.41mg,460.49μmol)。将混合物在-10℃下搅拌15min,然后加温至25℃并且搅拌1h。TLC(EtOAc)显示反应完成。将有机相用NaHCO₃(20mL*2)洗涤,在Na₂SO₄上干燥,过滤并且在真空中浓缩。通过硅胶色谱(石油醚/乙酸乙酯=50/1,10/1)纯化残余物以提供呈棕色油状的(R)-2-(5-氟吡啶-3-基)-4-氧基吡咯烷-1-羧酸叔丁酯(1.10g,产率:85.32%)。

[0401] 步骤4: (R)-4,4-二氟-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯



[0403] 在-70℃下在N₂下向(R)-2-(5-氟吡啶-3-基)-4-氧基吡咯烷-1-羧酸叔丁酯(1.00g,3.57mmol)于DCM(100.00mL)中的混合物中逐滴添加DAST(14.39g,89.25mmol)。在-70℃下将混合物搅拌30min。然后在25℃下将混合物搅拌16h。在0℃下通过NaHCO₃饱和水溶液来缓慢淬灭反应混合物并且将水相用DCM(50mL*4)萃取。将合并的有机相用饱和盐水(30mL)洗涤,在Na₂SO₄上干燥,过滤并且在真空中浓缩。通过硅胶色谱(石油醚/乙酸乙酯=100/1,30/1)纯化残余物以提供呈棕色油状的(R)-4,4-二氟-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(1.00g,产率:92.66%)。¹H-NMR(400MHz,CDC13) δ_{ppm} 8.40(s,1H),8.34(s,1H),7.30-7.21(m,1H),5.06(br.s,1H),4.14-3.85(m,2H),2.91-2.84(m,1H),2.39-2.32(m,1H),1.43-1.14(m,9H)。


[0404] 步骤5: (R)-3-(4,4-二氟吡咯烷-2-基)-5-氟吡啶



[0406] 在25℃下将(R)-4,4-二氟-2-(5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(1.00g, 3.31mmol)于HCl/EtOAc(50.00mL, 4M)中的混合物搅拌2h。将混合物在30℃下在减压下浓缩以提供呈白色固体状的呈双HCl盐形式的(R)-3-(4,4-二氟吡咯烷-2-基)-5-氟吡啶(840.00mg, 产率:92.25%)。¹H-NMR(400MHz, MeOD) δ ppm 8.68-8.63 (m, 1H), 7.97 (d, 1H, J=9.2Hz), 5.26-5.21 (m, 1H), 4.03-3.90 (m, 2H), 3.13-2.92 (m, 2H)。


[0407] 实施例13. (3S,5R)-5-(2,5-二氟苯基)吡咯烷-3-腈

[0408] 步骤1: ((2R)-2-((叔丁基二甲基甲硅烷基)氧基)-4-(2,5-二氟苯基)-4-羟基丁基)氨基甲酸叔丁酯




[0410] 在0℃下在N₂下,向2-溴-1,4-二氟苯(3.01g, 15.60mmol, 1.20当量)于THF(15mL)中的溶液中逐滴添加异丙基氯化镁复合物(2.27g, 15.60mmol, 1.20当量)。将反应物在15℃下搅拌1h以制备(2,5-二氟苯基)溴化镁(23mL)。在0℃下在30分钟内向(R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-氧基吡咯烷-1-羧酸叔丁酯(4.10g, 13.00mmol, 1.00当量)于THF(50mL)中的溶液中逐滴添加(2,5-二氟苯基)溴化镁(23mL)。在0℃下将反应混合物搅拌1h。在0℃下将甲醇(20mL)添加至混合物,随后添加NaBH₄(738mg, 19.50mmol, 1.50当量)。将混合物在0℃下搅拌1h,然后倾倒至10%NH₄Cl水溶液。将混合物用EtOAc(20mL*2)萃取,将合并的有机层用盐水洗涤,在Na₂SO₄上干燥,过滤并浓缩。通过中压液相色谱(MPLC)纯化粗产物以得到((2R)-2-((叔丁基二甲基甲硅烷基)氧基)-4-(2,5-二氟苯基)-4-羟基丁基)氨基甲酸叔丁酯(2.22g, 5.14mmol, 39.6%产率)。¹H-NMR(400MHz, CDCl₃) δ ppm 7.17-7.15 (m, 1H), 6.86-6.79 (m, 2H), 5.11-5.06 (m, 1H), 4.70 (br. s, 1H), 4.02-3.98 (m, 1H), 3.69 (br. s, 0.5H), 3.46 (br. s, 0.5H), 3.33-3.14 (m, 2H), 1.80-1.69 (m, 2H), 1.35 (s, 9H), 0.84-0.82 (9H, m), 0.04-0.03 (6H, m)。

[0411] 步骤2: (4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(2,5-二氟苯基)吡咯烷-1-羧酸叔丁酯




[0413] 在-60℃下在N₂下,向((2R)-2-((叔丁基二甲基甲硅烷基)氧基)-4-(2,5-二氟苯基)-4-羟基丁基)氨基甲酸叔丁酯(13.40g, 31.05mmol, 1.00当量)和Et₃N(9.43g, 93.14mmol, 3.00当量)于DCM(50mL)中的溶液中逐滴添加甲磺酰氯(5.33g, 46.57mmol, 1.50当量)。将混合物在-60℃下搅拌2h并且在15℃下搅拌16h。LCMS显示起始材料完全消耗。将反应混合物用DCM(30mL*2)萃取并且将合并的有机物用盐水(50mL)洗涤,在Na₂SO₄上干燥并过滤,浓缩以得到(4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(2,5-二氟苯基)吡咯烷-1-羧酸叔丁酯(12.00g, 26.11mmol, 产率:84.10%, 90%纯度),其未进一步纯化即直接使用。

[0414] 步骤3: (2R,4R)-2-(2,5-二氟苯基)-4-羟基吡咯烷-1-羧酸叔丁酯




[0416] 在15℃下向 (4R)-4-((叔丁基二甲基甲硅烷基)氧基)-2-(2,5-二氟苯基)吡咯烷-1-羧酸叔丁酯 (4.50g, 10.88mmol, 1.00当量) 于THF (30mL) 中的溶液中添加TBAF/THF (1M, 14.15mL, 1.30当量)。在15℃下将混合物搅拌16h。TLC (PE:EtOAc=3:1) 显示起始材料完全消耗。通过H₂O (50mL) 淬灭反应混合物, 将其用EtOAc (30mL*2) 萃取并且将合并的有机物用盐水 (10mL) 洗涤, 在Na₂SO₄上干燥, 过滤并在真空中浓缩。通过中性制备型HPLC纯化残余物以提供呈白色固体状的 (2R,4R)-2-(2,5-二氟苯基)-4-羟基吡咯烷-1-羧酸叔丁酯 (1.00g, 3.34mmol, 产率: 30.70%)。¹H-NMR (400MHz, CDCl₃) δppm 7.04-6.80 (m, 3H), 5.10-5.00 (m, 1H), 4.43 (s, 1H), 3.75 (br. s, 1H), 3.53-3.49 (m, 1H), 2.53 (br. s, 1H), 1.93-1.90 (m, 1H), 1.40-1.16 (m, 9H)。

[0417] 步骤4: (2R,4R)-2-(2,5-二氟苯基)-4-((甲磺酰基)氧基)吡咯烷-1-羧酸叔丁酯




[0419] 在0℃下向 (2R,4R)-2-(2,5-二氟苯基)-4-羟基吡咯烷-1-羧酸叔丁酯 (3.00g, 10.02mmol, 1.00当量) 和Et₃N (2.03g, 20.04mmol, 2.00当量) 于DCM (80.00mL) 中的混合物中逐滴添加MsCl (1.61g, 14.03mmol, 1.40当量)。在18℃下将混合物搅拌2h。通过H₂O (30mL) 淬灭混合物。通过DCM (50mL*3) 萃取水相。将合并的有机层在Na₂SO₄上干燥并且在减压下浓缩。获得呈棕色固体状的 (2R,4R)-2-(2,5-二氟苯基)-4-((甲磺酰基)氧基)吡咯烷-1-羧酸叔丁酯 (3.60g, 9.54mmol, 产率: 95.20%)。

[0420] 步骤5: (2R,4S)-4-氰基-2-(2,5-二氟苯基)吡咯烷-1-羧酸叔丁酯




[0422] 向 (2R,4R)-2-(2,5-二氟苯基)-4-((甲磺酰基)氧基)吡咯烷-1-羧酸叔丁酯 (3.60g, 9.54mmol, 1.00当量) 于DMSO (20.00mL) 中的混合物中一次性添加KCN (745.49mg, 11.45mmol, 1.20当量)。在90℃下将混合物搅拌3h。将80mL的H₂O添加至混合物, 并且通过EtOAc (80mL*4) 萃取混合物。在减压下浓缩合并的有机层。通过硅胶色谱 (PE/EtOAc=40:1, 30:1, 10:1) 纯化残余物。获得呈浅绿色液体状的 (2R,4S)-4-氰基-2-(2,5-二氟苯基)吡咯烷-1-羧酸叔丁酯 (1.60g, 5.19mmol, 产率: 54.40%)。

[0423] 步骤6: (3S,5R)-5-(2,5-二氟苯基)吡咯烷-3-腈



[0425] 将(2R,4S)-4-氰基-2-(2,5-二氟苯基)吡咯烷-1-羧酸叔丁酯(800.00mg, 2.59mmol, 1.00当量)于TFA(4.00mL)/DCM(20.00mL)中的混合物在18℃下搅拌3h。将混合物在N₂下干燥。获得呈淡黄色固体状的(3S,5R)-5-(2,5-二氟苯基)吡咯烷-3-腈(780.00mg, 2.42mmol, 产率:93.44%)。


[0426] 实施例14.3-氟-5-((2R,4S)-4-氟吡咯烷-2-基)苯甲酰胺



[0428] 将3-氟-5-((2R,4S)-4-氟吡咯烷-2-基)苄腈(0.050g, 0.240mmol) (如WO 2012/034095中制备)溶解于TFA(0.800mL, 10.38mmol)和H₂SO₄(0.200mL, 3.75mmol)中并且在室温下搅拌过夜。将反应混合物用冰水(3mL)稀释并且通过过滤分离固体,并且直接使用所述固体。


[0429] 实施例15.2-氯-5-氟-3-((2R,4S)-4-氟吡咯烷-2-基)吡啶

[0430] 步骤1: (S,Z)-N-((2-氯-5-氟吡啶-3-基)亚甲基)-2-甲基丙烷-2-亚磺酰胺




[0432] 在0℃下将2-氯-5-氟烟碱醛(20g, 125mmol)溶解于THF(150mL)中。添加(R)-2-甲基丙烷-2-亚磺酰胺(16.71g, 138mmol),随后逐滴添加四乙醇钛(22.88mL, 150mmol)。搅拌反应混合物,同时加温至RT。在3小时之后,将反应混合物冷却至0℃,并且添加150mL盐水并且搅拌20分钟。通过硅藻土过滤混合物。将水层分离并丢弃。将有机层在Na₂SO₄上干燥并且将溶剂移除以得到(S,Z)-N-((2-氯-5-氟吡啶-3-基)亚甲基)-2-甲基丙烷-2-亚磺酰胺(32g, 122mmol, 97%产率),其未进一步纯化即继续使用。LCMS: 263M+H。

[0433] 步骤2: (R)-N-((R)-1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-基)-2-甲基丙烷-2-亚磺酰胺




[0435] 将(R,E)-N-((2-氯-5-氟吡啶-3-基)亚甲基)-2-甲基丙烷-2-亚磺酰胺(32.9g, 125mmol)溶解于HPMA(100ml)中并且冷却至0℃。在0℃下添加锌(16.37g, 250mmol)、烯丙基溴(21.67ml, 250mmol)和水(2.256ml, 125mmol)并且将反应混合物加温至RT过夜。LCMS显示完全转化成所需产物。在RT下添加100ml水并且搅拌30分钟。添加30ml的MBTE, 随后添加60ml的10%柠檬酸并且将反应混合物搅拌30分钟。通过硅藻土过滤混合物并用MTBE洗涤。将有机层用10%柠檬酸、水和盐水洗涤。将溶剂在真空中移除以得到呈橙色油状的(R)-N-((R)-1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-基)-2-甲基丙烷-2-亚磺酰胺(14.5g, 47.6mmol, 38.0%产率)。LCMS: 305M+H。

[0436] 步骤3: (R)-1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-胺, HCl




[0438] 将(R)-N-((R)-1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-基)-2-甲基丙烷-2-亚磺酰胺(7.5g, 24.61mmol)溶解于10ml的MeOH中。添加HCl(二噁烷中的4M)(30.8ml, 123mmol)并且在RT下搅拌1h。在真空中移除溶剂并且将残余物用DCM稀释并用NaHCO₃饱和水溶液洗涤。将各层分离并且将有机层用Na₂SO₄干燥并在真空中移除溶剂。回收呈固体状的(R)-1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-胺, HCl(5.83g, 24.59mmol, 100%产率)。LCMS: 201M+H。

[0439] 步骤4: (R)-N-(1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-基)乙酰胺



[0441] 在0℃下向DCM(70.3ml)中的(R)-1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-胺·HCl(5.83g, 24.59mmol)中添加TEA(4.11ml, 29.5mmol)和乙酸酐(2.320ml, 24.59mmol)。将混合物搅拌2小时。将反应混合物倾倒至饱和NaHCO₃水溶液中并且用DCM萃取。将有机层用盐水洗涤、在MgSO₄上干燥并且在减压下浓缩。回收(R)-N-(1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-基)乙酰胺(5.97g, 24.60mmol, 100%产率)并且未进一步纯化即继续使用。LCMS: 243M+H。

[0442] 步骤5: (5R)-5-(2-氯-5-氟吡啶-3-基)吡咯烷-3-基乙酸酯




[0444] 将(R)-N-(1-(2-氯-5-氟吡啶-3-基)丁-3-烯-1-基)乙酰胺(5.97g, 24.60mmol)溶解于THF(56.2ml)和水(14.06ml)中,随后添加I₂(18.73g, 73.8mmol)并且在RT下搅拌过夜。将粗反应物用饱和NaHCO₃和Na₂S₂O₃溶液稀释并且用EtOAc萃取两次。将水层用饱和NaHCO₃水溶液碱化并且用EtOAc萃取以获得呈淡黄色油状的(5R)-5-(2-氯-5-氟吡啶-3-基)吡咯烷-3-基乙酸酯(5.9g, 22.81mmol, 93%产率)。LCMS: 259M+H。

[0445] 步骤6: (2R)-4-乙酰氧基-2-(2-氯-5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯




[0447] 向(5R)-5-(2-氯-5-氟吡啶-3-基)吡咯烷-3-基乙酸酯(5.9g, 22.81mmol)于二噁烷(76ml)和水(76ml)中的溶液中添加BOC-酐(7.94ml, 34.2mmol),随后谨慎添加2N NaOH(7ml)以达到pH~9。在RT下将反应混合物搅拌1小时。将反应混合物用水稀释并且用EtOAc萃取三次。将有机层在Na₂SO₄上干燥并且将溶剂在真空下移除以得到(2R)-4-乙酰氧基-2-(2-氯-5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(3.5g, 9.75mmol, 42.8%产率),其未进一步纯化即继续使用。LCMS: 359M+H。

[0448] 步骤7: (2R)-2-(2-氯-5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯




[0450] 将(2R)-4-乙酰氧基-2-(2-氯-5-氟吡啶-3-基)吡咯烷-1-羧酸叔丁酯(3.5g, 9.75mmol)溶解于MeOH(48.8ml)中,随后添加2M NaOH(5.37ml, 10.73mmol)并且将反应混合物在RT下搅拌2小时。将溶剂在真空下移除并且将水层用1N HCl中和,并且用EtOAc萃取三次。将合并的有机层在Na₂SO₄上干燥。将溶剂在真空下移除并且通过硅胶色谱(0-70%Hex/EtOAc)纯化残余物以得到(2R)-2-(2-氯-5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯(2.1g, 6.63mmol, 68.0%产率)。LCMS: 317M+H。

[0451] 步骤8: (R)-2-(2-氯-5-氟吡啶-3-基)-4-氧基吡咯烷-1-羧酸叔丁酯




[0453] 将 (2R)-2-(2-氯-5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯 (2.1g, 6.63mmol) 溶解于DCM (66.3ml) 中并且添加NaHCO₃ (0.557g, 6.63mmol), 随后添加戴斯-马丁过碘烷 (8.44g, 19.89mmol)。将反应混合物搅拌过夜。添加水 (0.119ml, 6.63mmol), 随后添加戴斯-马丁过碘烷 (8.44g, 19.89mmol) 并且搅拌18小时。用饱和NaHCO₃水溶液将pH调整至~7并且用DCM x3萃取。将有机层合并, 在Na₂SO₄上干燥并且将溶剂在真空下移除。通过快速色谱 (0-70%Hex/EtOAc) 纯化残余物以得到 (R)-2-(2-氯-5-氟吡啶-3-基)-4-氧基吡咯烷-1-羧酸叔丁酯 (1.6g, 5.08mmol, 77%产率)。LCMS: 315M+H。

[0454] 步骤9: (2R,4R)-2-(2-氯-5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯




[0456] 将 (R)-2-(2-氯-5-氟吡啶-3-基)-4-氧基吡咯烷-1-羧酸叔丁酯 (1.6g, 5.08mmol) 悬浮于乙醇 (33.9ml) 中并且冷却至0℃。分批添加NaBH₄ (0.096g, 2.54mmol) 并且在0℃下搅拌45分钟。用饱和NH₄Cl缓慢淬灭反应物并且将其加温至RT, 并且将溶液用DCM x3萃取。将有机层合并并且在Na₂SO₄上干燥。通过快速色谱 (0-70%Hex/EtOAc) 纯化残余物以得到 (2R,4R)-2-(2-氯-5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯 (1.446g, 4.57mmol, 90%产率)。LCMS: 317M+H。

[0457] 步骤10: (2R,4S)-2-(2-氯-5-氟吡啶-3-基)-4-氟吡咯烷-1-羧酸叔丁酯



[0459] 将 (2R,4R)-2-(2-氯-5-氟吡啶-3-基)-4-羟基吡咯烷-1-羧酸叔丁酯 (1.0g, 3.16mmol) 溶解于DCM (25ml) 中并且冷却至-78℃。添加TEA·HF (1.098ml, 9.47mmol) 并且搅拌10分钟。添加XtalFluor-E (1.446g, 6.31mmol) 并且在10分钟之后将反应混合物转移至冰浴并且将其加温至0℃。在2小时之后, 将反应混合物用DCM稀释并且用饱和NaHCO₃水溶液淬灭。将有机层分离, 并且将溶剂在真空下移除。通过ISCO (0-50%Hex/EtOAc; 12g柱) 纯化残余物以得到呈白色固体状的 (2R,4S)-2-(2-氯-5-氟吡啶-3-基)-4-氟吡咯烷-1-羧酸叔丁酯 (0.805g, 2.53mmol, 80%产率)。LCMS: 319M+H。


[0460] 步骤11: 2-氯-5-氟-3-((2R,4S)-4-氟吡咯烷-2-基)吡啶, HCl



[0462] 将 (2R,4S)-2-(2-氯-5-氟吡啶-3-基)-4-氟吡咯烷-1-羧酸叔丁酯 (0.805g, 2.53mmol, 80%产率) 溶解于EtOAc (5ml) 中并且添加4N HCl/二噁烷 (3ml)。在RT下将反应混合物搅拌1小时。将沉淀物过滤掉, 用醚洗涤, 并且在高真空下干燥过夜以得到呈米黄色固

体状的2-氯-5-氟-3-((2R,4S)-4-氟吡咯烷-2-基)吡啶, HCl (0.612g, 2.399mmol, 76%产率)。LCMS: 219M+H。


[0463] 实施例16. 5-氟-3-((2R,4S)-4-氟吡咯烷-2-基)-2-甲氧基吡啶



[0465] 将5-氟-3-((2R,4S)-4-氟吡咯烷-2-基)-2-甲氧基吡啶以与3-氟-5-((2R,4S)-4-氟吡咯烷-2-基)苯甲酰胺相同的方式来制备, 用5-氟-2-甲氧基烟碱醛替换2-氯-5-氟烟碱醛。


[0466] 实施例17. (1R,3R,4R)-3-氨基-4-羟基环戊烷-1-羧酸甲酯

[0467] 步骤1: (1S,2R,4S,5R)-3-氧杂-6-氮杂三环[3.2.1.0^{2,4}]辛-7-酮



[0469] 在0℃下在5h内向(1R,4S)-2-氮杂双环[2.2.1]庚-5-烯-3-酮(30.00g, 274.90mmol, 1.00当量)于NaH₂PO₄(395.00mL, 0.2M)和Na₂HPO₄(55.00mL, 0.2M)中的溶液中分批添加H₂O(450.00mL)和过硫酸氢钾复合盐(669.31g, 4.40mol, 16.00当量), 并且通过添加NaOH水溶液(12M)来保持pH=6并且将温度保持于0℃。在添加之后, 将混合物在0℃下再搅拌2h, TLC(PE:EtOAc=1:1)显示起始材料完全消耗, 将混合物过滤并且将水相用DCM(400mL*5)萃取, 将合并的有机层在Na₂SO₄上干燥, 真空浓缩以获得呈黄色固体状的(1S,2R,4S,5R)-3-氧杂-6-氮杂三环[3.2.1.0^{2,4}]辛-7-酮(9.00g, 71.93mmol, 产率: 26.16%)。¹H-NMR(400MHz, CDCl₃) δppm 5.96(br.s, 1H), 3.86(s, 1H), 3.62(1H, d, J=3.2Hz), 3.53(1H, d, J=2.8Hz), 2.86(s, 1H), 1.82(d, 1H, J=9.6Hz), 1.64(d, 1H, J=10.0Hz)。


[0470] 步骤2: (1S,2R,4S,5R)-7-氧基-3-氧杂-6-氮杂三环[3.2.1.0^{2,4}]辛烷-6-羧酸叔丁酯



[0472] 向(1S,2R,4S,5R)-3-氧杂-6-氮杂三环[3.2.1.0^{2,4}]辛-7-酮(9.00g, 71.93mmol, 1.00当量)于DCM(100.00mL)中的溶液中添加Boc₂O(17.27g, 79.12mmol, 1.10当量)、Et₃N(8.73g, 86.32mmol, 1.20当量)和DMAP(878.71mg, 7.19mmol, 0.10当量), 将混合物在25℃下搅拌16h, LCMS显示起始材料完全消耗, 将混合物用NH₄Cl水溶液(100mL*3)洗涤, 将合并的有机层在Na₂SO₄上干燥, 在真空中浓缩, 使用硅胶柱色谱(PE:EtOAc=5:1~1:1)纯化粗产物以获得呈黄色固体状的(1S,2R,4S,5R)-7-氧基-3-氧杂-6-氮杂三环[3.2.1.0^{2,4}]辛烷-6-羧酸叔丁酯(12.00g, 53.28mmol, 产率: 74.07%)。¹H-NMR(400MHz, CDCl₃) δppm 4.56(s, 1H), 3.71(d, 1H, J=2.8Hz), 3.54(d, 1H, J=2.8Hz), 3.00(s, 1H), 1.75(d, 1H, J=10.0Hz), 1.57

(d, 1H, J=10.8Hz), 1.46 (s, 9H)。

[0473] 步骤3: (3R,4R)-4-((叔丁氧羰基)氨基)-3-羟基环戊-1-烯-1-羧酸甲酯




[0475] 在0℃下将Na (3.37mg, 146.50μmol, 0.01当量)添加至MeOH (10.00mL),然后将溶液在0℃下搅拌0.5h,将溶液添加至MeOH (30.00mL)中的(1S,2R,4S,5R)-7-氧杂-3-氧杂-6-氮杂三环[3.2.1.0^{2,4}]辛烷-6-羧酸叔丁酯 (3.30g, 14.65mmol, 1.00当量),并且然后将混合物在16℃下搅拌13.5h,LCMS显示起始材料完全消耗,用乙酸 (5mL)淬灭反应物,并且然后用NaHCO₃ (20mL*3)洗涤,将有机层在Na₂SO₄上干燥,并且在真空中浓缩,用PE (20mL)洗涤粗产物以获得呈白色固体状的(3R,4R)-4-((叔丁氧羰基)氨基)-3-羟基环戊-1-烯-1-羧酸甲酯 (2.10g, 8.16mmol, 产率:55.72%)。¹H-NMR (400MHz, CDCl₃) δ_{ppm} 6.65 (s, 1H), 4.98 (s, 1H), 4.81 (d, 1H, J=2.8Hz), 4.46 (s, 1H), 3.98-3.92 (m, 1H), 3.75 (s, 3H), 3.07-3.01 (m, 1H), 2.36-2.29 (m, 1H), 1.45 (s, 9H)。

[0476] 步骤4: (1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-羟基环戊烷-1-羧酸甲酯



[0478] 将(3R,4R)-4-((叔丁氧羰基)氨基)-3-羟基环戊-1-烯-1-羧酸甲酯 (2.00g, 7.77mmol, 1.00当量)、(1Z,5Z)-环辛-1,5-二烯; (2S,5S)-1-[2-[(2S,5S)-2,5-二甲基磷杂环戊烷-1-基]乙基]-2,5-二甲基-磷杂环戊烷; 铯 (1+); 三氟甲磺酸酯 (48.08mg, 77.74μmol, 0.01当量)于MeOH (50.00mL)中的混合物脱气并且用H₂吹扫3次,并且然后将混合物在55℃下在H₂ (40psi)气氛下搅拌16h,TLC (PE:EtOAc=1:1)显示起始材料消耗,将混合物真空浓缩,并且然后溶解于EtOAc (5mL)中,然后向混合物中添加PE (20mL),形成白色固体,将沉淀物收集,在真空中干燥,将固体溶解于MeOH (7mL)中,并且用酸性制备型-HPLC (HC1)纯化以获得呈黄色油的(1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-羟基环戊烷-1-羧酸甲酯 (750.00mg, 2.89mmol, 产率:37.23%),所述结构通过手性HPLC和¹H-NMR来证实。¹H-NMR (400MHz, CDCl₃) δ_{ppm} 4.04-3.99 (m, 1H), 3.80-3.77 (m, 1H), 3.70 (s, 3H), 2.93-2.91 (m, 1H), 2.48-2.44 (m, 1H), 2.43-2.35 (m, 1H), 1.92-1.89 (m, 1H), 1.87-1.69 (m, 1H), 1.45 (s, 9H)。

[0479] 实施例18.合成(1R,3R,4R)-3-氨基-4-羟基环戊烷-1-羧酸甲酯




[0481] 在19℃下将HCl/二噁烷 (10.00mL, 4M)中的(1R,3R,4R)-3-((叔丁氧羰基)氨基)-

4-羟基环戊烷-1-羧酸甲酯(700.00mg, 2.70mmol, 1.00当量)搅拌5h, LCMS显示起始材料消耗, 将混合物在真空中浓缩以获得呈黄色油状的(1R, 3R, 4R)-3-氨基-4-羟基环戊烷-1-羧酸甲酯(500.00mg, 2.56mmol, 产率: 94.66%)。¹H-NMR(400MHz, CD₃OD) δ_{ppm} 4.09-4.05 (m, 1H), 3.75-3.66 (m, 3H), 3.37-3.33 (m, 1H), 3.06-3.04 (m, 1H), 2.45-2.37 (m, 2H), 1.87-1.81 (m, 2H)。

[0482] LC-MS(流动相: 从95% [水+0.375%v/v TFA]和5% [CH₃CN+0.188%v/v TFA], 在此条件下历时0.25min, 然后在10.0min内变化至15% [CH₃CN+0.188%v/v TFA], 在此条件下历时5min, 最后在0.01min内变化至95% [水+0.375%v/v TFA]和5% [CH₃CN+0.188%v/v TFA], 然后在此条件下历时5min。流量自始至终为1.0mL · min⁻¹) 纯度为98.803%, Rt = 0.893min, MS计算值: 159.2, MS实验值: 160.1 ([M+1]⁺)。


[0483] 实施例19. (3aS, 4R, 6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-胺

[0484] 步骤1: (3aS, 4S, 6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-醇



[0486] 将(3aR, 6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-酮(50.00g, 324.34mmol, 1.00当量)溶解于MeOH(1.00L)中, 然后添加CeCl₃·7H₂O(120.84g, 324.34mmol, 30.83mL, 1.00当量)。将混合物冷却至0℃。然后在0℃下在1.5h内分批添加NaBH₄(24.54g, 648.68mmol, 2.00当量)。在添加之后, 通过TLC(PE/EtOAc = 5/1)检查反应完成。通过饱和NH₄Cl(1000mL)淬灭反应物, 将其用DCM(500mL*5)萃取。将合并的有机层在Na₂SO₄上干燥并且在45℃下在真空中浓缩。获得呈黄色液体状的(3aS, 4S, 6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-醇(50.66g, 324.37mmol, 产率: 100.00%), 其未纯化即直接使用。


[0487] 步骤2: 2-((3aS, 4R, 6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮



[0489] 在20℃下向(3aS, 4S, 6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-醇(8.00g, 51.22mmol, 1.00当量)和异吲哚啉-1,3-二酮(9.04g, 61.46mmol, 1.20当量)于甲苯(250.00mL)中的混合物中添加PPh₃(20.15g, 76.83mmol, 1.50当量)。然后在0℃下将DIAD(15.54g, 76.83mmol, 1.50当量)逐滴添加至混合物。在添加之后, 使混合物达到80℃并且搅拌16h。TLC(PE/EtOAc = 5/1)显示反应完成。浓缩混合物。通过硅胶柱色谱(PE/EtOAc = 25/1至15/1)纯化残余物。所获得产物为呈黄色油状的粗产物, 在TLC上有一些极性点。因此, 添加80mL的MeOH并且产生白色沉淀物并通过过滤收集。获得呈白色固体状的2-

((3aS,4R,6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮(9.60g,33.65mmol,产率:65.70%)。


[0490] 步骤3: ((3aS,4R,6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)胺



[0492] 向2-((3aS,4R,6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮(9.52g,33.37mmol,1.00当量)于EtOH(300.00mL)中的混合物中添加乙烷-1,2-二胺(4.01g,66.74mmol,2.00当量)。在80℃下将反应混合物搅拌16h。产生许多白色沉淀物。TLC(PE/EtOAc=5/1)显示起始材料完全消耗。将沉淀物过滤。向滤液中添加300mL的NaOH(0.5M)。将混合物用DCM(200mL*5)萃取,在Na2SO4上干燥并浓缩。获得呈黄色油状的((3aS,4R,6aR)-2,2-二甲基-3a,6a-二氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)胺(4.90g,31.57mmol,产率:94.62%)。

[0493] 实施例20. (1R,3R,4R)-3-氨基-4-羟基环戊烷-1-腈

[0494] 步骤1: (1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-羟基环戊烷-1-羧酸



[0496] 在15℃下将(1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-羟基环戊烷-1-羧酸甲酯(5.00g,19.28mmol,1.00当量)、LiOH·H2O(2.43g,57.85mmol,3.00当量)于MeOH(10.00mL)和H2O(10.00mL)中的混合物搅拌16h,TLC(PE:EtOAc=1:1)显示反应完成,向混合物中添加稀释的HCl(1M)直至pH=6为止,并且真空浓缩,然后将混合物溶解于DCM(15mL)和EtOAc(5mL)中,将混合物过滤,并且将滤液真空浓缩以获得呈白色固体状的产物(1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-羟基环戊烷-1-羧酸(6.50g,粗产物)。¹H-NMR(400MHz,CD₃OD) δppm 3.95-3.90(m,1H),3.74-3.69(m,1H),2.91-2.87(m,1H),2.32-2.22(m,2H),1.82-1.72(m,2H),1.45(s,9H)。


[0497] 步骤2: ((1R,2R,4R)-4-氨甲酰基-2-羟环戊基)氨基甲酸叔丁酯



[0499] 向(1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-羟基环戊烷-1-羧酸(3.00g,12.23mmol,1.00当量)于DMF(40.00mL)中的混合物中添加HATU(6.05g,15.90mmol,1.30当量)、DIPEA(4.74g,36.69mmol,3.00当量)和NH₄Cl(1.96g,36.69mmol,3.00当量),然后将混合物在15℃下搅拌32h,LCMS显示反应完成,将混合物真空浓缩以获得((1R,2R,4R)-4-氨甲


酰基-2-羟环戊基)氨基甲酸叔丁酯(11g,粗产物)(建立2个批次并且在一起纯化)。¹H-NMR (400MHz, CD₃OD) δ ppm 4.58 (br. s, 1H), 3.96-3.94 (m, 1H), 3.77-3.75 (m, 1H), 2.25-2.20 (m, 2H), 1.79-1.75 (m, 2H), 1.44 (s, 9H)。

[0500] 步骤3: ((1R, 2R, 4R)-4-氰基-2-羟环戊基)氨基甲酸叔丁酯



[0502] 在0℃下向((1R, 2R, 4R)-4-氨基-2-羟环戊基)氨基甲酸叔丁酯(2.00g, 8.19mmol, 1.00当量)、吡啶(1.94g, 24.56mmol, 3.00当量)于THF(3.00mL)中的混合物中逐滴添加TFAA(2.58g, 12.28mmol, 1.50当量),然后混合物在0℃下搅拌0.5h,然后在15℃下向混合物中添加Et₃N(2.49g, 24.56mmol, 3.00当量),并且将混合物在15℃下搅拌0.5h,并且向混合物中添加TFAA(2.58g, 12.28mmol, 1.50当量),并且将混合物在15℃下搅拌0.5h, LCMS显示反应完成,将混合物真空浓缩,通过制备型HPLC(TFA, MS)纯化以获得呈无色油状的((1R, 2R, 4R)-4-氰基-2-羟环戊基)氨基甲酸叔丁酯(380.00mg, 1.68mmol, 产率: 20.51%)。¹H-NMR (400MHz, CD₃OD) δ ppm 4.07-3.97 (m, 1H), 3.79-3.78 (m, 0.5H), 3.51-3.49 (m, 0.5H), 3.19-3.05 (m, 1H), 2.44-2.35 (m, 0.5H), 2.33-2.28 (m, 1.5H), 1.97-1.94 (m, 1H), 1.92-1.82 (m, 1H), 1.41 (s, 9H)。

[0503] 步骤4: (1R, 3R, 4R)-3-氨基-4-羟基环戊烷-1-腈




[0505] 将((1R, 2R, 4R)-4-氰基-2-羟环戊基)氨基甲酸叔丁酯(800.00mg, 3.54mmol, 1.00当量)于TFA(5.00mL)和DCM(5.00mL)中的混合物在20℃下搅拌2h, LCMS显示反应完成,将混合物真空浓缩以获得呈无色油状的(1R, 3R, 4R)-3-氨基-4-羟基环戊烷-1-腈(545.00mg, 2.27mmol, 产率: 64.10%)。¹H-NMR (400MHz, CD₃OD) δ ppm 4.12-4.06 (m, 1H), 3.51-3.47 (m, 1H), 3.23-3.20 (m, 1H), 2.48-2.42 (m, 2H), 2.11-2.10 (m, 1H), 1.94-1.90 (m, 1H)。

[0506] 实施例21. (3aS, 4R, 6aS)-6,6-二氟-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-胺

[0507] 步骤1: 2-((3aS, 4R, 6S, 6aR)-6-羟基-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮


[0508]



[0509] 将(3aR,4S,6R,6aS)-6-氨基-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-醇(0.43g,2.46mmol,1.00当量)、邻苯二甲酸酐(0.36g,2.46mmol,1当量)和DIEA(0.65mL,3.7mmol,1.5当量)于甲苯(6.2mL)中的混合物在100℃下搅拌9h。LCMS显示反应完成。将EtOAc添加至反应混合物,并且然后用饱和碳酸氢钠水溶液(15mL)洗涤。将合并的有机层用饱和盐水溶液洗涤,在Na₂SO₄上干燥,在真空中浓缩。通过硅胶柱色谱(己烷/EtOAc)纯化残余物以获得呈白色固体状的产物2-((3aS,4R,6S,6aR)-6-羟基-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮(0.62g,83%)。

[0510] 步骤2:2-((3aS,4R,6aS)-2,2-二甲基-6-氧基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮


[0511]



[0512] 向2-((3aS,4R,6S,6aR)-6-羟基-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮(0.20g,0.68mmol,1.00当量)于DCM(4.5mL)中的溶液中添加PCC(0.29g,1.35mmol,2当量)并且将溶液在23℃下搅拌16h。添加另一个等分试样的PCC(0.15g,0.67mmol)并且将反应再持续16小时。LCMS显示反应完成。将EtOAc添加至反应混合物,并且然后通过硅藻土垫过滤。将残余物浓缩,然后通过硅胶柱色谱(己烷/EtOAc)纯化以获得呈灰白色固体状的产物2-((3aS,4R,6aS)-2,2-二甲基-6-氧基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮(0.19g,94%)。

[0513] 步骤3:2-((3aS,4R,6aS)-6,6-二氟-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮


[0514]



[0515] 向2-((3aS,4R,6aS)-2,2-二甲基-6-氧基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮(0.16g,0.53mmol,1.00当量)于DCM(3.5mL)中的溶液中添加DAST(0.42g,2.64mmol,5当量)并且将溶液在回流下搅拌16h。添加另一个等分试样的DAST(0.42g,2.64mmol,5当量)并且将反应在23℃下再持续16小时。将反应混合物用DCM稀释并

且用饱和碳酸氢钠溶液洗涤。将合并的有机层用饱和盐水溶液洗涤,在 Na_2SO_4 上干燥,在真空中浓缩。通过硅胶柱色谱(己烷/ EtOAc)纯化残余物以获得产物2-((3aS,4R,6aS)-6,6-二氟-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮(0.065g,38%)。


[0516] 步骤4: (3aS,4R,6aS)-6,6-二氟-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-胺



[0518] 向2-((3aS,4R,6aS)-6,6-二氟-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-基)异吲哚啉-1,3-二酮(0.065g,0.2mmol,1.00当量)于乙醇(1.8mL)中的溶液中添加一水合肼(0.015mL,0.3mmol,1.5当量)并且将溶液在50℃下搅拌2h,然后在70℃下再搅拌2小时。使用最小体积乙醇来过滤非均相反应混合物。然后将滤液浓缩并且分离的粗产物(3aS,4R,6aS)-6,6-二氟-2,2-二甲基四氢-4H-环戊[d][1,3]间二氧杂环戊烯-4-胺未进一步纯化即用于下一个步骤中。


[0519] 实施例22. 合成(1R,3R,4R)-4-氨基环己烷-1,3-二醇

[0520] 步骤1: (1r,4r)-4-(苄氧基)环己醇和(1s,4s)-4-(苄氧基)环己醇



[0522] 在10min时间段期间,向4-(苄氧基)环己酮(31.0g,152mmol)于500mL甲醇中的冰浴冷却溶液中,分多批添加硼氢化钠(5.78g,153mmol),然后将溶液在20℃下搅拌2h。然后通过饱和氯化铵水溶液(50mL)淬灭混合物,将其浓缩并且将残余物溶解于200mL水中并且用乙酸乙酯(200mL x 3)萃取,将合并的有机相在硫酸钠上干燥,然后在真空中浓缩以得到呈淡黄色油状的标题产物(1r,4r)-4-(苄氧基)环己醇和(1s,4s)-4-(苄氧基)环己醇(31.0g,粗产物),其未进一步纯化即直接用于下一个步骤。MS (ES+) $\text{C}_{13}\text{H}_{18}\text{O}_2$ 需要值:206,实验值:207 $[\text{M}+\text{H}]^+$ 。


[0523] 步骤2: ((环己-3-烯基氧基)甲基)苯



[0525] 在30min时间段期间,向(1r,4r)-4-(苄氧基)环己醇和(1s,4s)-4-(苄氧基)环己


醇 (30.0g, 145mmol) 以及 N,N-二异丙基乙胺 (28.1g, 218mmol) 于 1200mL 二氯甲烷中的冰浴冷却溶液中逐滴添加三氟甲磺酸酐 (30.7g, 109mmol), 然后将溶液在 25℃ 下搅拌 18h。然后将混合物在真空中浓缩并且使用硅胶柱色谱以石油醚:乙酸乙酯=12:1 洗脱来纯化残余物以得到呈黄色油状的标题化合物 (28.0g, 产率 100%)。MS (ES+) $C_{13}H_{16}O$ 需要值: 188, 实验值: 189 $[M+H]^+$ 。

[0526] 步骤3: (1R, 3R, 6S) -3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷



[0528] 在 0℃ 下使用间氯过氧苯甲酸 (21.9g, 127mmol) 处理 ((环己-3-烯基氧基) 甲基) 苯 (12.0g, 63.7mmol) 于二氯甲烷 (200mL) 中的溶液。将反应混合物在 0℃ 下搅拌 2h, 然后在室温下搅拌 15min。将洗涤的 (10% 亚硫酸钠水溶液、5% 氢氧化钠水溶液、然后为水) 有机溶液蒸发, 提供液体残余物, 使用硅胶柱色谱以己烷:异丙醚:乙酸乙酯=65:28:7 洗脱来分离所述残余物以得到呈黄色油状的标题化合物 (4.18g, 产率 32%)。反式-(1S, 3R, 6R)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷在 TLC 上显示更小极性并且首先洗脱。顺式-(1R, 3R, 6S)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷其次洗脱。MS (ES+) $C_{13}H_{16}O_2$ 需要值: 204, 实验值: 205 $[M+H]^+$ 。


[0529] 步骤4: (1R, 2R, 5R) -5-(苄氧基)-2-((S)-1-苯基乙氨基) 环己醇



[0531] 将高氯酸锂 (7.27g, 68.4mmol) 添加至 (1R, 3R, 6S)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷 (7.0g, 34.2mmol) 于 120mL 4A-MS 干燥乙腈中的冰浴冷却搅拌溶液中, 将浴移除并且在 15min 时间段期间逐滴添加 (S)-1-苯基乙胺 (5.58g, 46.1mmol), 然后将溶液在 25℃ 下搅拌 18h。然后将混合物在 200mL 水中稀释并且用乙酸乙酯 (200mL x 3) 萃取, 将合并的有机相在硫酸钠上干燥, 然后浓缩并且使用硅胶柱色谱以石油醚:乙酸乙酯:三乙胺=98:0:2~49:49:2 洗脱来纯化残余物以得到呈黄色油状的标题化合物 (3.5g, 产率 31%)。 (1S,


2S,5S)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇在TLC上显示更小极性,并且首先洗脱。(1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇其次洗脱。MS (ES+) $C_{21}H_{27}NO_2$ 需要值:325,实验值:326 $[M+H]^+$ 。

[0532] 步骤5: (1R,2R,4R)-4-(苄氧基)-2-(叔丁基二甲基甲硅烷基氧基)-N-((S)-1-苯基乙基)环己胺




[0534] 将三氟甲磺酸叔丁基二甲基甲硅烷基酯(13.0g,49.5mmol)添加至(1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇(5.4g,16.5mmol)和三乙胺(5.0g,49.5mmol)于100mL干燥二氯甲烷中的冰浴冷却的搅拌溶液中。在30min之后,将此用碳酸氢钠的饱和水溶液洗涤,在硫酸钠上干燥。移除溶剂,并且使用硅胶柱色谱以石油醚:乙酸乙酯=100:0~70:30洗脱来纯化残余物以得到呈黄色油状的标题化合物(5.4g,产率71%)。MS (ES+) $C_{27}H_{41}NO_2Si$ 需要值:439,实验值:440 $[M+H]^+$ 。

[0535] 步骤7: (1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇



[0537] 在室温下,将四丁基氟化铵(2.66g,10.2mmol)添加至(1R,2R,4R)-4-(苄氧基)-2-(叔丁基二甲基甲硅烷基氧基)-N-((S)-1-苯基乙基)环己胺(1.5g,3.41mmol)于50mL干燥氧杂环戊烷中的搅拌溶液中。然后在65°C下将此溶液搅拌2h。然后将混合物在真空中浓缩并且将残余物在200mL水中稀释并且用乙酸乙酯(200mL x 3)萃取,将合并的有机相用水和饱和氯化钠水溶液洗涤,在硫酸钠上干燥。移除溶剂,并且使用硅胶柱色谱以石油醚:乙酸乙酯=100:0~70:30洗脱来纯化残余物以得到呈无色油状的标题化合物(0.75g,产率68%)。MS (ES+) $C_{21}H_{27}NO_2$ 需要值:325,实验值:326 $[M+H]^+$ 。

[0538] 步骤8: (1R,3R,4R)-4-氨基环己烷-1,3-二醇




[0540] 在室温下,将活性碳上的10%氢氧化钡(697mg,催化剂)添加至(1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇(650mg,1.99mmol)于15mL乙醇中的溶液。然后在50°C

下在氢气下将此溶液搅拌20h。然后将混合物冷却并且通过硅藻土过滤,将滤饼用甲醇:二氯甲烷=1:10洗涤,将滤液在真空下浓缩并且将残余物在20mL甲醇:二氯甲烷=1:10溶液中稀释并且浓缩,在高真空下干燥,然后在-20℃下冷却以得到呈白色晶体状的标题化合物(240mg,产率92%)。MS (ES+) $C_6H_{13}NO_2$ 需要值:131,实验值:132 $[M+H]^+$ 。 1H -NMR (400MHz, 6d-DMSO) δ ppm 4.62-4.49 (m, 2H), 3.42-3.33 (m, 2H, $J=3.2$ Hz), 2.93-2.86 (m, 1H), 2.25-2.18 (m, 1H), 1.98-1.92 (m, 1H), 1.72-1.59 (m, 3H), 1.13-1.03 (m, 2H), 0.97-0.90 (m, 1H)。


[0541] 实施例23. 合成 (1R,2R)-2-(叔丁基二甲基甲硅烷基氧基)-4-氧环己基氨基甲酸叔丁酯

[0542] 步骤1: (1R,2R,4R)-2-(叔丁基二甲基甲硅烷基氧基)-4-羟基环己基氨基甲酸叔丁酯



[0544] 在室温下,将活性碳上的10%氢氧化钾(1.9g,催化剂)添加至来自先前实施例的(1R,2R,4R)-4-(苄氧基)-2-(叔丁基二甲基甲硅烷基氧基)-N-((S)-1-苯基乙基)环己胺(2.0g,4.54mmol)和二碳酸二-叔丁酯(3.95g,18.1mmol)于60mL乙醇中的溶液中。然后在50℃下在氢气下将此溶液搅拌20h。然后将混合物冷却并且通过硅藻土过滤,将滤饼用甲醇:二氯甲烷=1:10洗涤,将滤液在真空中浓缩并且将残余物在20mL甲醇:二氯甲烷=1:10溶液中稀释并且浓缩,在高真空下干燥,以得到呈无色油状的标题化合物(1.2g,产率77%)。MS (ES+) $C_{17}H_{35}NO_4Si$ 需要值:345,实验值:346 $[M+H]^+$ 。


[0545] 步骤2: (1R,2R)-2-(叔丁基二甲基甲硅烷基氧基)-4-氧环己基氨基甲酸叔丁酯



[0547] 在室温下,将1,1,1-三乙氧基-1,1-二氢-1,2-苯碘酰基-3(1H)-酮(戴斯-马丁过碘烷,4.11g,9.71mmol)添加至(1R,2R,4R)-2-(叔丁基二甲基甲硅烷基氧基)-4-羟基环己基氨基甲酸叔丁酯(1.4g,4.05mmol)于50mL二氯甲烷中的溶液中。然后在40℃下在氮气下将此溶液搅拌3h。然后将混合物在真空下浓缩并且使用硅胶柱色谱以石油醚:乙酸乙酯=100:0~95:5洗脱来纯化残余物以得到呈黄色油状的标题化合物(1.2g,产率86%)。MS (ES+) $C_{17}H_{33}NO_4Si$ 需要值:343,实验值:344 $[M+H]^+$ 。 1H -NMR (400MHz, $CDCl_3$) δ ppm 4.70-4.49 (br., 1H), 4.10-3.90 (br. s, 1H), 3.79-3.65 (br., 1H), 2.64 (dd, 1H, $J=14.4, 4.0$ Hz), 2.42-2.28 (m, 4H), 1.46 (s, 9H), 0.87 (s, 9H), 0.08 (d, 6H, $J=6.8$ Hz)。


[0548] 实施例23. 合成 (1R,2R)-2-(叔丁基二甲基甲硅烷基氧基)-4-氧环己基氨基甲酸叔丁酯和(1R,3R,4R)-4-氨基环己烷-1,3-二醇

[0549] 步骤1: (1R,4R)-4-(苄氧基)环己醇和(1S,4S)-4-(苄氧基)环己醇:




[0551] 在10min时间段期间,向4-(苄氧基)环己酮(31.0g,152mmol)于500mL甲醇中的冰浴冷却溶液中,分多批添加硼氢化钠(5.78g,153mmol),然后将溶液在20℃下搅拌2h。然后通过饱和氯化铵水溶液(50mL)淬灭混合物,将其浓缩并且将残余物溶解于200mL水中并且用乙酸乙酯(200mL x 3)萃取,将合并的有机相在硫酸钠上干燥,然后在真空下浓缩以得到呈淡黄色油状的标题产物(1R,4R)-4-(苄氧基)环己醇和(1S,4S)-4-(苄氧基)环己醇(31.0g,粗产物),其未进一步纯化即直接用于下一个步骤。MS (ES+) $C_{13}H_{18}O_2$ 需要值:206,实验值:207 $[M+H]^+$ 。

[0552] 步骤2: ((环己-3-烯基氧基)甲基)苯:



[0554] 在30min时间段期间,向(1R,4R)-4-(苄氧基)环己醇和(1S,4S)-4-(苄氧基)环己醇(30.0g,145mmol)和N,N-二异丙基乙胺(28.1g,218mmol)于1200mL二氯甲烷中的冰浴冷却溶液中逐滴添加三氟甲磺酸酐(30.7g,109mmol),然后将溶液在25℃下搅拌18h。然后将混合物在真空中浓缩并且使用硅胶柱色谱以石油醚:乙酸乙酯=12:1洗脱来纯化残余物以得到呈黄色油状的标题化合物(28.0g,产率100%)。MS (ES+) $C_{13}H_{16}O$ 需要值:188,实验值:189 $[M+H]^+$ 。

[0555] 步骤3: (1R,3R,6S)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷:




[0557] 在0℃下使用间氯过氧苯甲酸(21.9g,127mmol)处理((环己-3-烯基氧基)甲基)苯(12.0g,63.7mmol)于二氯甲烷(200mL)中的溶液。将反应混合物在0℃下搅拌2h,并且然后在室温下搅拌15min。将洗涤的(10%亚硫酸钠水溶液、5%氢氧化钠水溶液、然后为水)有机溶液蒸发,提供液体残余物,使用硅胶柱色谱以己烷:异丙醚:乙酸乙酯=65:28:7洗脱来分离所述残余物以得到呈黄色油状的标题化合物(4.18g,产率32%)。反式-(1S,3R,6R)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷在TLC上显示更小极性并且首先洗脱。顺式-(1R,3R,6S)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷其次洗脱。MS (ES+) $C_{13}H_{16}O_2$ 需要值:204,实验值:205 $[M+H]^+$ 。

[0558] 顺式-(1R,3R,6S)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷:¹H-NMR (400MHz, CDCl₃) δppm 7.35-7.27 (m, 5H), 4.56-4.45 (m, 2H), 3.35-3.29 (m, 1H), 3.12-3.09 (m, 2H), 2.37-2.32 (m, 1H), 2.25-2.20 (m, 1H), 1.89-1.68 (m, 3H), 1.49-1.44 (m, 1H)。

[0559] 反式-(1S,3R,6R)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷:¹H-NMR (400MHz, CDCl₃) δppm 7.35-7.27 (m, 5H), 4.48 (dd, 2H, J=28.0, 12.4Hz), 3.56-3.52 (m, 1H), 3.19-3.17 (m, 2H), 2.24-2.18 (m, 1H), 2.15-2.07 (m, 1H), 2.00-1.91 (m, 2H), 1.64-1.53 (m, 2H)。

[0560] 步骤3: (1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇:




[0562] 将高氯酸锂 (7.27g, 68.4mmol) 添加至 (1R,3R,6S)-3-(苄氧基)-7-氧杂-双环[4.1.0]庚烷 (7.0g, 34.2mmol) 于120mL 4A-MS干燥乙腈中的冰浴冷却搅拌溶液中, 将浴移除并且在15min时间段期间逐滴添加 (S)-1-苯基乙胺 (5.58g, 46.1mmol), 然后将溶液在25℃下搅拌18h。然后将混合物在200mL水中稀释并且用乙酸乙酯 (200mL x 3) 萃取, 将合并的有机相在硫酸钠上干燥, 然后浓缩并且使用硅胶柱色谱以石油醚:乙酸乙酯:三乙胺=98:0:2~49:49:2洗脱来纯化残余物以得到呈黄色油状的标题化合物 (3.5g, 产率31%)。(1S,2S,5S)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇在TLC上显示更小极性, 并且首先洗脱。(1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇其次洗脱。MS (ES+) C₂₁H₂₇N₂O₂需要值: 325, 实验值: 326 [M+H]⁺。

[0563] (1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇:¹H-NMR (400MHz, CDCl₃) δ ppm 7.35-7.24 (m, 10H), 4.52 (d, 2H, J=2.0Hz), 3.97 (q, 1H, J=6.8Hz), 3.42-3.34 (m, 1H), 3.19-3.12 (m, 1H), 2.39 (dd, 1H, J=12.0, 2.4Hz), 2.16 (dd, 1H, J=12.0, 3.6Hz), 2.09-2.00 (m, 2H), 1.65-1.49 (m, 1H), 1.35 (d, 3H, J=6.4Hz), 1.28-1.15 (m, 2H), 0.90 (qd, 1H, J=13.2, 3.6Hz)。


[0564] (1S,2S,5S)-5-(苄氧基)-2-((S)-1-苯基乙氨基)环己醇:¹H-NMR (400MHz, CDCl₃) δ ppm 7.36-7.22 (m, 10H), 4.54 (d, 2H, J=3.2Hz), 3.90 (q, 1H, J=6.4Hz), 3.44-3.35 (m, 1H), 3.15-3.09 (m, 1H), 2.51-2.45 (m, 1H), 2.43-2.36 (m, 1H), 2.04-1.99 (m, 1H), 1.95-1.90 (m, 1H), 1.47-1.29 (m, 3H), 1.34 (d, 3H, J=6.4Hz), 0.82 (qd, 1H, J=13.2, 3.2Hz)。

[0565] 步骤4: 合成 (1R,2R,4R)-4-(苄氧基)-2-(叔丁基二甲基甲硅烷基氧基)-N-((S)-1-苯基乙基)环己胺:




[0567] 将三氟甲磺酸叔丁基二甲基甲硅烷基酯 (13.0g, 49.5mmol) 添加至 (1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基) 环己醇 (5.4g, 16.5mmol) 和三乙胺 (5.0g, 49.5mmol) 于 100mL 干燥二氯甲烷中的冰浴冷却的搅拌溶液中。在 30min 之后, 将此用碳酸氢钠的饱和水溶液洗涤, 在硫酸钠上干燥。移除溶剂, 并且使用硅胶柱色谱以石油醚: 乙酸乙酯 = 100:0 ~ 70:30 洗脱来纯化残余物以得到呈黄色油状的标题化合物 (5.4g, 产率 71%)。MS (ES+) $C_{27}H_{41}NO_2Si$ 需要值: 439, 实验值: 440 $[M+H]^+$ 。

[0568] 步骤 5: (1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基) 环己醇:



[0570] 在室温下, 将四丁基氟化铵 (2.66g, 10.2mmol) 添加至 (1R,2R,4R)-4-(苄氧基)-2-(叔丁基二甲基甲硅烷基氧基)-N-((S)-1-苯基乙基) 环己胺 (1.5g, 3.41mmol) 于 50mL 干燥氧杂环戊烷中的搅拌溶液中。然后在 65 °C 下将此溶液搅拌 2h。然后将混合物在真空中浓缩并且将残余物在 200mL 水中稀释并用乙酸乙酯 (200mL x 3) 萃取, 将合并的有机相用水和饱和氯化钠水溶液洗涤, 在硫酸钠上干燥。移除溶剂, 并且使用硅胶柱色谱以石油醚: 乙酸乙酯 = 100:0 ~ 70:30 洗脱来纯化残余物以得到呈无色油状的标题化合物 (0.75g, 产率 68%)。MS (ES+) $C_{21}H_{27}NO_2$ 需要值: 325, 实验值: 326 $[M+H]^+$ 。

[0571] 步骤 6: (1R,3R,4R)-4-氨基环己烷-1,3-二醇:




[0573] 在室温下, 将活性碳上的 10% 氢氧化钡 (697mg, 催化剂) 添加至 (1R,2R,5R)-5-(苄氧基)-2-((S)-1-苯基乙氨基) 环己醇 (650mg, 1.99mmol) 于 15mL 乙醇中的溶液中。然后在 50 °C 下在氢气下将此溶液搅拌 20h。然后将混合物冷却并且通过硅藻土过滤, 将滤饼用甲醇: 二氯甲烷 = 1:10 洗涤, 将滤液在真空下浓缩并且将残余物在 20mL 甲醇: 二氯甲烷 = 1:10 溶液中稀释并且浓缩, 在高真空下干燥, 然后在 -20 °C 下冷却以得到呈白色晶体状的标题化合物 (240mg, 产率 92%)。MS (ES+) $C_6H_{13}NO_2$ 需要值: 131, 实验值: 132 $[M+H]^+$ 。


[0574] (1R,3R,4R)-4-氨基环己烷-1,3-二醇: 1H -NMR (400MHz, 6J-DMSO) δ ppm 4.62-4.49

(m, 2H), 3.42-3.33 (m, 2H, J=3.2Hz), 2.93-2.86 (m, 1H), 2.25-2.18 (m, 1H), 1.98-1.92 (m, 1H), 1.72-1.59 (m, 3H), 1.13-1.03 (m, 2H), 0.97-0.90 (m, 1H)。


[0575] 实施例24:合成化合物232:



[0576]




[0577] 步骤1: (4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮



[0578]


[0579] 在15℃下向4-氧基-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-4,5-二氢-1H-吡唑并[3,4-d]嘧啶-3-羧酸(1.00g, 3.22mmol, 1.00当量)于SOCl₂(164.00g, 1.38mol, 428.11当量)中的溶液中添加DMF(235.49mg, 3.22mmol, 1.00当量)。将反应物在50℃下加热16h。TLC(PE:EtOAc=3:1, R_f=0.8和0.7)显示反应完成。浓缩混合物。将残余物冷却至-10℃并且溶解于DCM(25.00mL)中。将Et₃N(1.63g, 16.10mmol, 5.00当量)和(2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷(497.40mg, 2.09mmol, 0.65当量, HCl)添加至反应。在0℃下将反应物搅拌0.2h。TLC(PE:EtOAc=3:1, R_f=0.38)显示反应完成。将溶液用盐水(5mL)洗涤, 在Na₂SO₄上干燥并且浓缩。通过制备型-TLC(PE:EtOAc=10:1)纯化残余物以得到呈黄色固体状的(4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮(400.00mg, 产率:24.26%)。

[0580] 步骤2: (1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基环戊烷-1-羧酸甲酯




[0582] 向(4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮(200.00mg, 390.63 μ mol, 1.00当量)和(1R,3R,4R)-3-氨基-4-羟基-环戊烷羧酸甲酯(80.24mg, 410.16 μ mol, 1.05当量, HCl)于二噁烷(10.00mL)中的溶液中添加DIPEA(151.46mg, 1.17mmol, 3.00当量)。将反应物在90℃下加热5h。LCMS显示反应完成。将溶液浓缩。通过制备型-TLC(PE:EtOAc=2:1)纯化残余物以得到呈黄色固体状的(1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基环戊烷-1-羧酸甲酯(120.00mg, 产率:48.40%)。

[0583] 步骤3: (1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基环戊烷-1-羧酸甲酯



[0585] 在15℃下,向(1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基环戊烷-1-羧酸甲酯(120.00mg, 189.06 μ mol, 1.00当量)于DCM(3.00mL)中的溶液中添加TFA(3.00mL)。在15℃下将反应物搅拌16h。LCMS显示起始材料消耗。检测到极少(1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基环戊烷羧酸甲酯。将反应物浓缩。将残余物溶解于MeOH(20.00mL)中。将KOAc(185.54mg, 1.89mmol, 10.00当量)添加至反应物中。将反应物在50℃下加热16h。LCMS显示反应完成。将溶液浓缩。将残余物溶解于EtOAc(20mL)中并且用盐水(10mL)洗涤,在Na₂SO₄上干燥并且浓缩以得到呈红色固体状的(1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基环戊烷-1-羧酸甲酯(80.00mg, 粗产物),其未纯化即用于下一个步骤中。


[0586] 步骤4: ((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((1R,2R,4R)-2-羟基-4-(2-羟丙-2-基)环戊基)氨基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮




[0588] 在 -70°C 下,向(1R,3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基环戊烷-1-羧酸甲酯(80.00mg, 158.59 μmol , 1.00当量)于THF(20.00mL)中的溶液中添加MeMgBr(3M, 1.59mL, 30.00当量)。将反应物缓慢加温至 15°C 并且搅拌2h。TLC(EtOAc, $R_f=0.24$)和LCMS显示反应完成。将溶液用1N HCl水溶液中和至 $\text{pH}=7$ 。将反应混合物浓缩。通过中性制备型-HPLC纯化残余物以得到呈黄色固体状的((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((1R,2R,4R)-2-羟基-4-(2-羟丙-2-基)环戊基)氨基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮(17.40mg, 产率: 21.75%)。

[0589] 对于此化合物和以下化合物156, LC-MS条件如下: (流动相: 从99% [水+0.375%v/v TFA] 和1% [CH_3CN +0.188%v/v TFA], 在此条件下历时0.4min, 然后在3.0min内变化至10% [水+0.375%v/v TFA] 和90% [CH_3CN +0.188%v/v TFA], 然后在0.45min内变化至100% [CH_3CN +0.188%v/v TFA], 最后在0.01min内变化至99% [水+0.375%v/v TFA] 和1% [CH_3CN +0.188%v/v TFA], 然后在此条件下历时0.64min。流量自始至终为 $0.8\text{mL} \cdot \text{min}^{-1}$ 。) 纯度为99.870%

[0590] 实施例25. 合成化合物229:




[0593] 步骤1: (3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基吡咯烷-1-羧酸叔丁酯




[0595] 在 50°C 下,将3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1-((2-(三甲基硅烷基)乙氧基)甲基)-1,5-二氢-4H-吡唑并[3,4-d]嘧啶-4-酮(0.085g,0.17mmol,1当量)的溶液与亚硫酸氯(0.031mL,0.43mmol,2.5当量)和DCM(0.7mL)中的几滴DMF一起搅拌3小时。LCMS指示SM完全消耗成氯代杂环中间物。将反应混合物在冰上冷却并且添加二噁烷(0.7mL),随后添加DIEA(0.21mL,1.21mmol,7当量)和(3R,4R)-3-氨基-4-羟基吡咯烷-1-羧酸叔丁酯(0.05g,0.26mmol,1.5当量)。接着在 70°C 下将反应混合物搅拌3小时。LCMS指示反应完成。然后将反应混合物用DCM稀释并且用饱和碳酸氢钠溶液洗涤。将合并的有机层用饱和盐水溶液洗涤,在 Na_2SO_4 上干燥并且在真空中浓缩。通过硅胶柱色谱(己烷/EtOAc)纯化残余物以获得产物(3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1-((2-(三甲基硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基吡咯烷-1-羧酸叔丁酯(0.084g,72%)。

[0596] 步骤2: ((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((3R,4R)-4-羟基吡咯烷-3-基)氨基)-1-((2-(三甲基硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮




[0598] 使用二噁烷(4M,0.9mL,3.72mmol,30当量)中的HCl处理(3R,4R)-3-((3-((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-羰基)-1-((2-(三甲基硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-4-基)氨基)-4-羟基吡咯烷-1-羧酸酯(0.084g,0.12mmol,1当量)于EtOAc(1.25mL)中的溶液。在 23°C 下搅拌4小时之后,LCMS指示反应完成。将反应混合物用EtOAc稀释并且用饱和碳酸氢钠溶液洗涤。将合并的有机层用饱和盐水溶液洗涤,在 Na_2SO_4 上干燥,在真空中浓缩。粗产物((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((3R,4R)-4-羟基吡咯烷-3-基)氨基)-1-((2-(三甲基硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮未进一步纯化即用于下一个步骤中。

[0599] 步骤3: ((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((3R,4R)-4-羟基-1-(氧杂环丁烷-3-基)吡咯烷-3-基)氨基)-1-((2-(三甲基硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮




[0601] 向((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((3R,4R)-4-羟吡咯烷-3-基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮(0.036g,0.062mmol,1当量)于DCE(0.5mL)中的溶液中添加几滴乙酸(2 μL ,0.031mmol,0.5当量),随后添加氧杂环丁烷-3-酮(0.015mL,0.21mmol,3.3当量)并且将反应混合物在60℃下加热2小时。添加三乙酰氧基硼氢化钠(0.033g,0.16mmol,2.5当量)并且将溶液在23℃下搅拌24小时。LCMS指示反应完成。将反应混合物用EtOAc稀释并且用饱和碳酸氢钠溶液洗涤。将合并的有机层用饱和盐水溶液洗涤,在 Na_2SO_4 上干燥,在真空中浓缩。然后,通过硅胶柱色谱(DCM/MeOH)纯化残余物以分离产物((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((3R,4R)-4-羟基-1-(氧杂环丁烷-3-基)吡咯烷-3-基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮(0.030g,75%)。

[0602] 步骤4: ((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((3R,4R)-4-羟基-1-(氧杂环丁烷-3-基)吡咯烷-3-基)氨基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮





[0604] 将((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((3R,4R)-4-羟基-1-(氧杂环丁烷-3-基)吡咯烷-3-基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮(0.060g,0.095mmol,1当量)于DCM(1mL)中的溶液用TFA(0.73mL,9.5mmol,100当量)处理16小时。将反应混合物用DCM稀释并且用饱和碳酸氢钠水溶液洗涤。将合并的有机层用饱和盐水溶液洗涤,在 Na_2SO_4 上干燥,并且在真空中浓缩。向DCM/MeOH(1/1,1mL)中的中间物中添加乙酸钠(0.016g,0.19mmol,2当量)并且将反应物在23℃下搅拌2小时。将反应混合物用DCM稀释并且用饱和碳酸氢钠溶液洗涤。将合并的有机层用饱和盐水溶液洗涤,在 Na_2SO_4 上干燥,在真空中浓缩。然后,首先通过硅胶柱色谱(含有10% NH_4OH 的DCM/MeOH)然后通过制备-TLC来纯化残余物以分离产物((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)(4-(((3R,4R)-4-羟基-1-(氧杂环丁烷-3-基)吡咯烷-3-基)氨基)-1H-吡唑并[3,4-d]嘧啶-3-基)甲酮(0.034g,70%)。


[0605] 实施例26.合成化合物156



[0606]

[0607] 步骤1:1-溴-2-(溴甲基)-4-氯苯


[0609] 在90℃下,将1-溴-4-氯-2-甲苯(3.00g,14.60mmol,1.00当量)、NBS(2.34g,13.14mmol,0.90当量)和AIBN(239.75mg,1.46mmol,0.10当量)于CCl₄(20.00mL)中的溶液搅拌12h。将溶液在真空中浓缩以得到呈黄色固体状的粗产物1-溴-2-(溴甲基)-4-氯苯(5.40g,粗产物),其未进一步纯化即用于下一个步骤中。

[0610] 步骤2:1-(2-溴-5-氯苯基)-N-甲基甲胺

[0612] 在25℃下,将1-溴-2-(溴甲基)-4-氯苯(5.40g,18.99mmol,1.00当量)于MeNH₂/H₂O(30.00mL)中的溶液搅拌15h。当反应完成时,用EtOAc(50mL*3)萃取产物,将合并的有机层在真空中浓缩以得到粗产物。通过硅胶柱色谱(PE:EtOAc=5:1至1:1)纯化粗产物以获得呈灰白色固体状的1-(2-溴-5-氯苯基)-N-甲基甲胺(1.00g,产率:22.45%)。¹H-NMR(400MHz,CDCl₃) δppm 7.45(d,1H,J=8.8Hz),7.40-7.39(m,1H),7.10(dd,1H,J=8.4,2.4Hz),3.79(s,2H),2.46(s,3H)。

[0613] 步骤3:N-(2-溴-5-氯苯基)-4-(((1R,2S,3R)-2,3-二羟基环戊基)氨基)-N-甲基-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺


[0614]



[0615] 向1-(2-溴-5-氯苄基)-N-甲基甲胺(500.00mg, 1.22mmol, 1.00当量)和4-(((1R, 2S, 3R)-2,3-二羟基环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3, 4-d]嘧啶-3-羧酸(314.98mg, 1.34mmol, 1.10当量)于DMF(5.00mL)中的溶液中添加DIPEA(315.35mg, 2.44mmol, 2.00当量)和HATU(556.66mg, 1.46mmol, 1.20当量),将所得混合物在25℃下搅拌15h。当反应完成时,添加H₂O(20mL),通过EtOAc(20mL*3)萃取产物,将合并的有机层在真空中浓缩以得到粗产物。通过制备型-TLC(EtOAc)纯化粗产物以获得呈红色油状的N-(2-溴-5-氯苄基)-4-(((1R, 2S, 3R)-2,3-二羟基环戊基)氨基)-N-甲基-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3, 4-d]嘧啶-3-甲酰胺(520.00mg, 产率:81.20%)。¹H-NMR(400MHz, CDCl₃) δppm 9.77(br.s, 0.5H), 9.49(br.s, 0.5H), 8.45(d, 1H, J=11.4Hz), 7.63(dd, 1H, J=8.4, 4.4Hz), 7.33-7.31(m, 1H), 7.26-7.25(m, 1H), 5.87(s, 1H), 5.67(s, 1H), 5.46(s, 1H), 4.98(s, 1H), 4.45-4.44(m, 1H), 4.31-4.29(m, 1H), 4.03-4.02(m, 1H), 3.79(t, 1H, J=8.0Hz), 3.71(s, 1.5H), 3.55(t, 1H, J=8.0Hz), 3.29(s, 1.5H), 3.16(d, 1H, J=6.0Hz), 2.65-2.61(m, 1H), 2.05-1.83(m, 2H), 1.05(t, 1H, J=8.4Hz), 0.90(t, 1H, J=8.0Hz), 0.07(s, 4.5H), 0.00(s, 4.5H)。

[0616] 步骤4:N-(5-氯-2-氰基苄基)-4-(((1R, 2S, 3R)-2,3-二羟基环戊基)氨基)-N-甲基-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3, 4-d]嘧啶-3-甲酰胺


[0617]



[0618] 将N-(2-溴-5-氯苄基)-4-(((1R, 2S, 3R)-2,3-二羟基环戊基)氨基)-N-甲基-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3, 4-d]嘧啶-3-甲酰胺(420.00mg, 670.91μmol, 1.00当量)、Zn(CN)₂(630.22mg, 5.37mmol, 340.66μL, 8.00当量)和Pd(PPh₃)₄(77.53mg, 67.09μmol, 0.10当量)的混合物溶解于密封管中的DMF(3.00mL)中,将其在140℃下用微波辐射3h。在3h之后,LCMS显示起始材料未完全消耗,因此添加更多Zn(CN)₂(2当量)和Pd(PPh₃)₄(0.1当量),将其在150℃下再用微波辐射2h。当反应完成时,添加H₂O(15mL),通过EtOAc(20mL*3)萃取产物。在无水Na₂SO₄干燥上合并的有机层并且将其浓缩得到粗产物。通过制备型-TLC(EtOAc)纯化粗产物以获得呈黄色油状的N-(5-氯-2-氰基苄基)-4-(((1R, 2S, 3R)-2,3-二羟基环戊基)氨基)-N-甲基-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡

唑并[3,4-d]嘧啶-3-甲酰胺 (520.00mg, 粗产物, 包括PPh₃O)。¹H-NMR (400MHz, CDCl₃) δppm 9.72 (br.s, 0.5H), 9.44 (br.s, 0.5H), 8.45 (d, 1H, J=6.4Hz), 7.79-7.54 (m, 3H), 5.86 (s, 1H), 5.70 (s, 1H), 5.67 (s, 1H), 5.11 (s, 1H), 4.45-4.44 (m, 1H), 4.30-4.29 (m, 1H), 4.03-4.02 (m, 1H), 3.81-3.76 (m, 2.5H), 3.59 (t, 1H, J=8.0Hz), 3.32 (s, 1.5H), 3.18-3.16 (m, 1H), 2.19-1.86 (m, 3H), 1.04 (t, 1H, J=8.0Hz), 0.92 (t, 1H, J=8.0Hz), 0.06 (s, 4.5H), 0.00 (s, 4.5H)。


[0619] 步骤5: N-(5-氯-2-氰基苄基)-4-(((1R,2S,3R)-2,3-二羟基环戊基)氨基)-N-甲基-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺



[0621] 向混合溶剂TFA (10.00mL) 和DCM (10.00mL) 中添加N-(5-氯-2-氰基苄基)-4-(((1R,2S,3R)-2,3-二羟基环戊基)氨基)-N-甲基-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺 (520.00mg, 908.88μmol, 1.00当量), 将所得混合物在25℃下搅拌1h。通过N₂蒸发溶剂以得到粗产物。将粗产物溶解于MeOH (15.00mL) 中, 通过NaHCO₃调整至pH=7-8, 并且添加KOAc (178.39mg, 1.82mmol, 2.00当量), 将其在50℃下搅拌2h。当反应完成时, 将混合物在真空中浓缩以得到粗产物, 将其溶解于EtOAc (50mL) 中, 通过H₂O (15mL*3) 洗涤。将有机层在真空中浓缩以得到粗产物, 通过酸性制备型-HPLC (TFA) 纯化所述粗产物以获得呈白色固体状的N-(5-氯-2-氰基苄基)-4-(((1R,2S,3R)-2,3-二羟基环戊基)氨基)-N-甲基-1H-吡唑并[3,4-d]嘧啶-3-甲酰胺 (171.00mg, 产率: 33.85%, TFA)。

[0622] 实施例27. 合成化合物226


[0623] 步骤1: (1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-((叔丁基二苯基甲硅烷基)氧基)环戊烷-1-羧酸甲酯:



[0625] 在0℃下, 向 (1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-羟基环戊烷-1-羧酸甲酯 (1.50g, 5.78mmol, 1.00当量) 和咪唑 (590.74mg, 8.67mmol, 1.50当量) 于DMF (10.00mL) 中的溶液中添加TBDPSCI (1.67g, 6.07mmol, 1.05当量)。在15℃下将反应物搅拌16h。TLC (PE: EtOAc=5:1, R_f=0.43) 显示反应完成。将溶液倾倒至水 (20mL) 中并且用EtOAc (10mL*3) 萃取。将有机层在Na₂SO₄上干燥并且浓缩。将残余物从PE (1mL) 重结晶以得到呈白色固体状的 (1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-((叔丁基二苯基甲硅烷基)氧基)环戊烷-1-羧酸甲酯 (2.80g, 产率: 97.40%)。¹H-NMR (400MHz, CDCl₃) δppm 7.69-7.64 (m, 4H), 7.43-7.37 (m,


6H), 4.10 (br.s, 1H), 3.92-3.97 (m, 2H), 3.66 (s, 3H), 2.75-2.71 (m, 1H), 2.46-2.43 (m, 1H), 2.01-1.95 (m, 2H), 1.65-1.61 (m, 1H), 1.40 (s, 9H), 1.05 (s, 9H)。

[0626] 步骤2: ((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(羟甲基)环戊基)氨基甲酸叔丁酯:



[0628] 在-30℃下,向(1R,3R,4R)-3-((叔丁氧羰基)氨基)-4-((叔丁基二苯基甲硅烷基)氧基)环戊烷-1-羧酸甲酯(2.80g, 5.63mmol, 1.00当量)于THF(30.00mL)中的溶液中添加LiAlH₄(427.32mg, 11.26mmol, 2.00当量)。将反应物缓慢加温至15℃并且搅拌2h。TLC(PE:EtOAc=3:1, R_f=0.24)显示反应完成。在0℃下,将反应物用0.43mL的H₂O和0.43mL的10% NaOH水溶液淬灭。将混合物过滤并且将滤液浓缩。将残余物用PE(5mL)洗涤。将固体收集。获得呈白色固体状的((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(羟甲基)环戊基)氨基甲酸叔丁酯(2.30g, 产率:86.98%)。¹H-NMR(400MHz, CDCl₃) δ_{ppm} 7.69-7.65 (m, 4H), 7.43-7.38 (m, 6H), 4.11-4.10 (m, 1H), 3.89 (br.s, 2H), 3.54 (br.s, 2H), 2.14-2.10 (m, 1H), 1.97-1.89 (m, 2H), 1.62-1.58 (m, 1H), 1.39 (s, 9H), 1.06 (s, 9H)。

[0629] 步骤3: ((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-甲酰基环戊基)氨基甲酸叔丁酯:




[0631] 在0℃下向((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(羟甲基)环戊基)氨基甲酸叔丁酯(2.30g, 4.90mmol, 1.00当量)于DCM(50.00mL)中的溶液中添加戴斯-马丁过碘烷试剂(3.12g, 7.35mmol, 1.50当量)。在15℃下将反应物搅拌16h。TLC(PE:EtOAc=3:1, R_f=0.7)显示反应完成。在0℃下,将反应物用饱和NaHCO₃(30mL)水溶液淬灭并且用DCM(30mL)萃取。将有机层在Na₂SO₄上干燥并且浓缩。通过硅胶柱色谱(PE:EtOAc=30:1)纯化残余物。获得呈黄色油状的((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-甲酰基环戊基)氨基甲酸叔丁酯(900.00mg, 产率:39.27%)。¹H-NMR(400MHz, CDCl₃) δ_{ppm} 9.55 (s, 1H), 7.61-7.56 (m, 4H), 7.37-7.31 (m, 6H), 4.08 (d, 1H, J=7.2Hz), 3.93-3.89 (m, 1H), 3.79 (br.s, 1H), 2.62-2.55 (m, 1H), 2.44-2.40 (m, 1H), 1.86-1.83 (m, 2H), 1.57-1.52 (m, 1H), 1.34 (s, 9H), 0.97 (s, 9H)。

[0632] 步骤4: ((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊基)氨基甲酸叔丁酯:




[0634] 在0℃下,向((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-甲酰基环戊基)氨基甲酸叔丁酯(900.00mg,1.92mmol,1.00当量)于DCM(50.00mL)中的溶液中添加DAST(928.45mg,5.76mmol,3.00当量)。在15℃下将反应物搅拌5h。TLC(PE:EtOAc=3:1, R_f =0.6)显示反应完成。将溶液用饱和NaHCO₃水溶液(20mL)淬灭并且用DCM(20mL)萃取。将有机层在Na₂SO₄上干燥并且浓缩。通过硅胶柱色谱(PE:EtOAc=30:1~20:1)纯化残余物。获得呈黄色油状的((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊基)氨基甲酸叔丁酯(250.00mg,产率:26.59%)。¹H-NMR(400MHz,CDCl₃) δ ppm 7.62-7.56(m,4H), 7.34-7.31(m,6H), 5.76-5.43(m,1H), 3.88-3.82(m,2H), 2.30-2.24(m,1H), 2.07-2.04(m,1H), 1.80-1.75(m,1H), 1.65-1.57(m,1H), 1.33(s,9H), 0.99(s,9H)。

[0635] 步骤5: ((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊-1-胺:




[0637] 在15℃下,向((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊基)氨基甲酸叔丁酯(50.00mg,102.11 μ mol,1.00当量)于EtOAc(2.00mL)中的溶液中添加HCl/EtOAc(10.00mL,4M)。在15℃下将反应物搅拌1h。TLC(PE:EtOAc=3:1, R_f =0.05)显示反应完成。通过N₂将溶剂吹干。残余物未进一步纯化。获得呈黄色油状的((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊-1-胺(40.00mg,产率:91.95%,HCl)。¹H-NMR(400MHz,CDCl₃) δ ppm 7.70-7.68(m,4H), 7.49-7.43(m,6H), 5.81(td,1H, J =57.2, 4.8Hz), 4.25(dd,1H, J =12.0, 5.6Hz), 3.50(dd,1H, J =13.2, 6.4Hz), 2.22-2.19(m,1H), 1.80-1.74(m,2H), 1.66-1.63(m,1H), 1.08(s,9H)。

[0638] 步骤6: (4-(((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮:



[0640] 向 (1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊-1-胺 (40.00mg, 93.89 μ mol, 1.00当量, HCl) 和 (4-氯-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮 (48.07mg, 93.89 μ mol, 1.00当量) 于二噁烷 (10.00mL) 中的溶液中添加 DIPEA (60.67mg, 469.45 μ mol, 5.00当量)。将反应物在 90℃ 下加热 0.5h。LCMS 显示反应完成。将溶液浓缩。通过制备型-TLC (PE:EtOAc=3:1) 纯化残余物。获得呈黄色油状的 (4-(((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮 (38.00mg, 产率: 46.78%)。

[0641] 步骤7: (4-(((1R,2R,4R)-4-(二氟甲基)-2-羟环戊基)氨基)-1H-吡唑并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮 (化合物226):












[0643] 在 50℃ 下将 (4-(((1R,2R,4R)-2-((叔丁基二苯基甲硅烷基)氧基)-4-(二氟甲基)环戊基)氨基)-1-((2-(三甲基甲硅烷基)乙氧基)甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮 (38.00mg, 43.93 μ mol, 1.00当量) 于 TBAF/THF (5.00mL) 中的溶液加热 2h。LCMS 显示留下 (4-(((1R,2R,4R)-4-(二氟甲基)-2-羟环戊基)氨基)-1-(羟甲基)-1H-吡唑并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮。将反应混合物浓缩。将残余物溶解于 EtOAc (20mL) 中并且用盐水 (10mL*2) 洗涤。将有机层浓缩并且溶解于 MeOH (20.00mL) 中。将 KOAc (21.56mg, 219.65 μ mol, 5.00当量) 添加至反应物中。将反应物在 50℃ 下加热 16h。LCMS 显示反应完成。将溶液浓缩。通过制备型-HPLC (MeOH/TFA 系统) 纯化残余物。获得呈黄色固体状的 (13.50mg, 产率:

50.34%，TFA)的(4-(((1R,2R,4R)-4-(二氟甲基)-2-羟环戊基)氨基)-1H-吡啶并[3,4-d]嘧啶-3-基)((2R,4S)-2-(2,5-二氟苯基)-4-氟吡咯烷-1-基)甲酮。











[0644] 实施例28.合成其他化合物









[0645] 本发明的额外化合物使用与以上实施例阐明的技术类似的技术来合成。下表指示基于每种化合物(“Cmpd”)的合成的特定实施例(“实施例”)以及用于合成每种特定化合物的合适氨基醇和胺。

[0646] 表1.用于合成示例性化合物的方案和中间物。










实施例	胺	氨基醇	Cmpd
1			7
1			8
2			9
2			10
1			11













[0647]

实施例	胺	氨基醇	Cmpd
1			1
1			2
1			3
1			4
1			5
1			6













Cmpd	氨基醇	胺	实施例
17			3
18			3
19			3
20			3
21			3

[0648]

Cmpd	氨基醇	胺	实施例
12			2
13			2
14			3
15			3
16			3

Cmpd	氨基醇	胺	实施例
28			1
29			1
30			1
31			1
32			1
33			1

[0649]

Cmpd	氨基醇	胺	实施例
22			1
23			1
24			1
25			1
26			1
27			1

Cmpd	氨基醇	胺	实施例
40			1
41			1
42			1
43			1
44			1

[0650]

Cmpd	氨基醇	胺	实施例
34			2
35			2
36			1
37			1
38			1
39			1

Cmpd	氨基醇	胺	实施例
50			1
51			23
52			1
53			1
54			1

[0651]

Cmpd	氨基醇	胺	实施例
45			1
46			1
47			1
48			1
49			1

Cmpd	氨基醇	胺	实施例
61			3
62			3
63			1
64			1
65			1
66			2










[0652]

Cmpd	氨基醇	胺	实施例
55			1
56			1
57			1
58			1
59			1
60			1











Cmpd	氨基醇	胺	实施例
72			1
73			1
74			1
75			3









[0653]

Cmpd	氨基醇	胺	实施例
67			2
68			1
69			1
70			2
71			2







Cmpd	氨基醇	胺	实施例
81			1
82			1
83			1
84			22
85			1

[0654]

Cmpd	氨基醇	胺	实施例
76			3
77			1
78			1
79			1
80			1

Cmpd	氨基醇	胺	实施例
90			2
91			2
92			1
93			1

[0655]

Cmpd	氨基醇	胺	实施例
86			2
87			2
88			1
89			1

Cmpd	氨基醇	胺	实施例
98			2
99			1
100			1
101			1
102			23











[0656]

Cmpd	氨基醇	胺	实施例
94			1
95			3
96			3
97			2









Cmpd	氨基醇	胺	实施例
107			1
108			1
109			23
110			1
111			23








[0657]

Cmpd	氨基醇	胺	实施例
103			1
104			1
105			1
106			1









Cmpd	氨基醇	胺	实施例
116			2
117			1
118			1
119			1
120			1












[0658]

Cmpd	氨基醇	胺	实施例
112			1
113			23
114			1
115			2











Cmpd	氨基醇	胺	实施例
125			2
126			1
127			1
128			1

[0659]

Cmpd	氨基醇	胺	实施例
121			1
122			1
123			3
124			2

Cmpd	氨基醇	胺	实施例
134			1
135			24
136			1
137			1
138			1
139			1

[0660]

Cmpd	氨基醇	胺	实施例
129			1
130			1
131			1
132			1
133			1

Cmpd	氨基醇	胺	实施例
145			1
146			24
147			1
148			1
149			1











[0661]

Cmpd	氨基醇	胺	实施例
140			24
141			24
142			1
143			24
144			1











Cmpd	氨基醇	胺	实施例
155			1
156			24
157			1
158			1
159			1








[0662]

Cmpd	氨基醇	胺	实施例
150			1
151			3
152			24
153			1
154			24









Cmpd	氨基醇	胺	实施例
165			24
166			1
167			24
168			1
169			1

[0663]

Cmpd	氨基醇	胺	实施例
160			1
161			1
162			1
163			1
164			1

Cmpd	氨基醇	胺	实施例
174			1
175			1
176			1
177			1









[0664]

Cmpd	氨基醇	胺	实施例
170			1
171			1
172			1
173			1








Cmpd	氨基醇	胺	实施例
182			1
183			24
184			1
185			1







[0665]

Cmpd	氨基醇	胺	实施例
178			3
179			1
180			1
181			1








Cmpd	氨基醇	胺	实施例
190			1
191			1
192			1
193			1









[0666]

Cmpd	氨基醇	胺	实施例
186			3
187			3
188			1
189			1








Cmpd	氨基醇	胺	实施例
198			1
199			1
200			1
201			1

[0667]

Cmpd	氨基醇	胺	实施例
194			1
195			1
196			1
197			1

Cmpd	氨基醇	胺	实施例
206			1
207			3
208			3
209			1









[0668]

Cmpd	氨基醇	胺	实施例
202			1
203			1
204			1
205			1









Cmpd	氨基醇	胺	实施例
214			1
215			1
216			1
217			1









[0669]

Cmpd	氨基醇	胺	实施例
210			22
211			1
212			1
213			1









Cmpd	氨基醇	胺	实施例
222			1
223			1
224			1
225			1








[0670]

Cmpd	氨基醇	胺	实施例
218			1
219			1
220			1
221			1









Cmpd	氨基醇	胺	实施例
230			1
231			1
232			22
233			1

[0671]







Cmpd	氨基醇	胺	实施例
226			27
227			1
228			1
229			23

Cmpd	氨基醇	胺	实施例
238			1
239			1
240			1
241			1

[0672]

Cmpd	氨基醇	胺	实施例
234			1
235			1
236			1
237			1

[0673]

实施例	1	1	1
胺			
氨基酸			
Cmpd	242	243	244

[0674] 对于本文公开的化合物所获得的NMR和LC MS数据示出在图1中。

[0675] 实施例29. 测定

[0676] 1mM ATP处的NTRK1野生型测定

[0677] 在384孔板的每个孔中,在25℃下,在一系列剂量浓度的化合物(1%DMSO最终浓度)存在或不存在下,将1nM-1.5nM野生型NTRK1酶(BPS Bioscience;40280)在总共12.5μL缓冲液(100mM HEPES pH 7.5,0.015%Brij 35,10mM MgCl₂,1mM DTT)中与1-2μM CSKtide (Tuft's University or Anaspec;FITC-AHA-KKKKD DIYFFFG-NH₂)和1mM ATP一起孵育60分钟。通过添加70μL终止缓冲液(100mM HEPES pH 7.5,0.015%Brij 35,35mM EDTA和0.2%涂布试剂3(Caliper Lifesciences))来终止反应。然后,将板在Caliper EZReader 2(方案设置:-1.7psi,上游电压-500,下游电压-3000,采样后sip 35s)上读取。数据归一化至0%和100%抑制对照并且IC₅₀使用CORE LIMS中的4-参数拟合计算。

[0678] NTRK野生型和G595R突变型细胞测定方案

[0679] 具有TPM3-NTRK1融合蛋白质的KM12野生型结肠癌细胞系从国家癌症学会(NCI)获得。此细胞系先前证明依赖于来自NTRK融合蛋白质的NTRK活性以便生长和存活。KM12 Cliff(G595R)细胞系通过用DNA甲基化剂诱变野生型KM12细胞系并且随后选择对于长期暴露于高浓度已知NTRK抑制剂(克唑替尼)有抗性的克隆来产生。将细胞首先在384孔板中以1000个细胞/孔涂覆于完全培养基(10%FBS和1%pen/strep)中并且在37℃下孵育过夜。然后,使用Bravo液体处置系统给予细胞不同浓度的测试样品。浓度范围为25uM直至9.5pM(4倍稀释,总共10个浓度)。每种化合物以每个板重复运行两次。DMSO和十字孢碱(25uM)作为

生长抑制的阴性和阳性对照包括于每个板上。在给药之后72h,测定板使用CellTiter-Glo (Promega) 显影并且所得荧光在Envision板阅读器上读取。IC₅₀确定值使用4-参数曲线拟合算法来计算。

[0680] 下表总结如上所述生物测定的结果。以下表示法用于指示每个测定中的IC₅₀:A<10.00nM;B=10.01-100.0nM;C=100.01-1000.0nM;和D>1000.1nM;“ND”=未确定。

[0681]

化合物编号	Enz NTRK1	KM12 (WT)	KM12 (G595R)
1	B	B	B
2	C	C	C
3	D	D	D
4	C	D	D
5	C	C	C
6	A	B	B
7	B	B	B
8	B	B	C
9	B	B	C
10	C	C	D
11	C	C	C
12	C	C	D
13	B	B	C
14	C	C	C
15	C	C	D
16	C	D	D
17	B	B	C
18	C	C	D
19	D	D	D
20	B	B	C
21	C	C	D
22	C	D	D
23	A	B	B
24	B	C	C
25	B	B	C
26	B	B	C
27	A	B	B
28	A	B	B
29	A	B	B
30	C	B	C
31	C	C	C
32	B	C	C
33	B	B	B
34	B	B	C
35	C	C	D
36	B	C	C
37	B	C	C
38	C	C	C
39	A	B	B
40	B	C	C
41	B	C	C
42	C	D	D
43	B	B	C

化合物编号	Enz NTRK1	KM12 (WT)	KM12 (G595R)
44	B	B	C
45	A	A	A
46	A	A	A
47	C	C	C
48	B	B	C
49	A	B	B
50	B	B	C
51	C	C	C
52	B	B	B
53	B	B	C
54	B	C	C
55	B	B	B
56	B	C	C
57	A	A	B
58	A	B	B
59	C	C	C
60	B	B	B
61	C	C	D
62	B	B	C
63	A	B	B
64	C	C	C
65	C	ND	ND
66	D	D	D
67	D	D	D
68	A	A	A
69	A	B	B
70	A	B	B
71	B	B	C
72	A	B	B
73	A	A	B
74	A	B	B
75	B	C	D
76	A	B	B
77	A	B	B
78	A	A	B
79	A	A	B
80	B	C	C
81	B	C	C
82	B	C	C
83	B	C	C
84	A	B	B
85	A	B	B
86	A	A	A

[0682]

化合物编号	Enz NTRK1	KM12 (WT)	KM12 (G595R)
87	A	B	B
88	A	A	A
89	A	B	A
90	D	D	D
91	B	B	C
92	A	A	A
93	A	A	A
94	C	C	D
95	A	A	A
96	B	B	C
97	A	B	C
98	C	D	D
99	A	A	A
100	B	B	C
101	C	C	C
102	A	B	B
103	D	D	D
104	A	A	A
105	A	B	B
106	A	A	A
107	A	B	B
108	B	C	C
109	A	B	C
110	A	B	B
111	A	B	B
112	B	B	C
113	A	B	B
114	A	A	A
115	A	B	B
116	A	A	A
117	C	C	D
118	A	A	A
119	A	A	A
120	B	B	B
121	A	A	A
122	A	A	A
123	A	B	B
124	A	ND	ND
125	A	ND	ND
126	A	A	A
127	A	A	A
128	A	A	A
129	A	B	B

化合物编号	Enz NTRK1	KM12 (WT)	KM12 (G595R)
130	A	A	A
131	C	C	C
132	B	B	C
133	C	C	C
134	C	C	C
135	A	A	A
136	A	B	B
137	B	B	B
138	C	C	C
139	B	B	B
140	A	B	A
141	A	B	A
142	B	B	C
143	B	B	B
144	B	B	C
145	C	B	C
146	A	B	B
147	B	C	C
148	A	A	A
149	A	B	B
150	B	B	B
151	B	B	B
152	A	B	A
153	B	B	B
154	A	B	B
155	B	B	B
156	B	B	A
157	A	B	B
158	A	B	B
159	A	B	B
160	A	B	B
161	B	B	B
162	A	B	B
163	A	B	B
164	B	B	B
165	A	A	A
166	A	C	C
167	A	A	A
168	A	A	A
169	B	B	B
170	A	A	A
171	B	B	B
172	A	B	B

化合物编号	Enz NTRK1	KM12 (WT)	KM12 (G595R)
173	A	A	A
174	A	A	A
175	A	B	B
176	A	B	B
177	A	A	A
178	A	B	B
179	B	B	C
180	A	A	A
181	A	B	B
182	A	A	B
183	A	A	A
184	A	A	A
185	A	A	A
186	A	A	A
187	A	B	B
188	A	A	A
189	A	A	B
190	A	B	B
191	A	B	B
192	A	B	B
193	A	B	B
194	B	B	B
195	A	A	A
196	A	A	A
197	A	B	B
198	A	A	A
199	B	A	A
200	A	B	B
201	A	A	A
202	A	B	B
203	A	A	A
204	A	A	A
205	A	A	A
206	A	A	A
207	B	B	B
208	A	A	A
209	A	B	B
210	A	A	A
211	A	A	A
212	A	A	A
213	A	A	A
214	A	A	A
215	A	A	A

[0683]

化合物编号	Enz NTRK1	KM12 (WT)	KM12 (G595R)
216	A	A	A
217	A	A	A
218	A	A	A
219	A	A	A
220	A	A	A
221	A	A	A
222	A	B	B
223	A	B	B
224	A	A	A
225	A	B	B
226	A	A	A
227	A	A	A
228	B	B	B
229	A	A	A
230	A	B	B
231	B	B	C
232	A	A	A
233	A	A	A
234	A	A	A
235	A	B	B
236	A	A	B
237	A	B	B
238	ND	B	B
239	A	A	B
240	A	A	A
241	A	A	A
242	A	A	A
243	A	A	A
244	A	A	A





[0684] 以引用的方式并入





[0685] 本文提及的所有公布和专利均据此以全文引用的方式并入本文中,其引用程度就





如同已明确并且个别地指示将各个别公布或专利以引用的方式并入本文中一般。





[0686] 等效物





[0687] 本领域技术人员将仅使用不超过常规实验即会认识到或能够确定本文所述的本发明的特定实施方案的许多等效物。这些等效物旨在由以下权利要求书涵盖。





#	结构	NMR; LCMS
1		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.13-7.02 (m, 3H), 4.66 (s, 2H), 4.17 (d, 2H, J = 3.2 Hz), 2.32-2.29 (m, 1H), 2.11-2.08 (m, 1H), 1.91-1.87 (m, 2H), 1.86-1.74 (m, 2H); LCMS: 389.1
2		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.319-7.15 (m, 1H), 7.02-7.01 (m, 1H), 6.99-6.88 (m, 1H), 4.61 (s, 2H), 4.38-4.31 (m, 1H), 4.13-4.11 (m, 1H), 4.03-4.02 (m, 1H), 2.44-2.41 (m, 1H), 2.34 (s, 3H), 2.16-2.13 (m, 1H), 1.79-1.70 (m, 2H); LCMS: 401.1
3		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.21 (s, 1H), 7.15-7.09 (m, 2H), 7.08-7.01 (m, 1H), 4.65 (s, 2H), 4.41-4.38 (m, 1H), 4.03-3.96 (m, 1H), 2.41-2.36 (m, 1H), 1.90-1.89 (m, 1H), 1.58-1.56 (m, 1H), 1.44-1.41 (m, 1H), 0.76-0.73 (m, 1H), 0.50-0.47 (m, 1H); LCMS: 401.1
4		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.12 (s, 1H), 7.06-6.99 (m, 2H), 6.93-6.92 (m, 1H), 4.56 (s, 2H), 4.32-4.29 (m, 1H), 3.93-3.87 (m, 1H), 2.32-2.25 (m, 1H), 1.81-1.80 (m, 1H), 1.49-1.47 (m, 1H), 1.35-1.33 (m, 1H), 0.67-0.64 (m, 1H), 0.41-0.40 (m, 1H); LCMS: 401.1





5		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.51-8.47 (m, 2H), 8.38 (d, 1H, J = 2.0 Hz), 7.74 (d, 1H, J = 9.2 Hz), 5.50 (s, 1H), 4.94 (s, 1H), 4.39-4.34 (m, 1H), 4.15-4.13 (m, 1H), 4.07-4.01 (m, 1H), 3.70 (s, 1.5H), 3.20 (s, 1.5H), 2.47-2.44 (m, 1H), 2.19-2.16 (m, 1H), 1.84-1.71 (m, 2H); LCMS: 402.1
6		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40 (d, 1H, J = 4.0 Hz), 7.22-7.07 (m, 3H), 5.51 (s, 1H), 4.26-4.17 (m, 2H), 3.70 (s, 1.5H), 3.21 (s, 1.5H), 2.38-2.36 (m, 1H), 2.17-2.06 (m, 1H), 1.96-1.76 (m, 4H); LCMS: 403.1
7		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34 (s, 1H), 7.22-7.21 (m, 1H), 7.13-7.11 (m, 1H), 7.04-7.02 (m, 1H), 5.56-5.51 (m, 1H), 4.16-4.13 (m, 2H), 2.32-2.29 (m, 1H), 2.09-2.08 (m, 1H), 1.90-1.70 (m, 4H), 1.59 (d, 3H, J = 7.2 Hz); LCMS: 403.1
8		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.32 (s, 1H), 7.15-7.09 (m, 2H), 7.05-7.03 (m, 1H), 4.67 (s, 2H), 4.15-4.11 (m, 1H), 4.05-4.04 (m, 1H), 3.98-3.95 (m, 1H), 3.85-3.84 (m, 1H), 3.56 (t, 1H, J = 10.8 Hz), 3.43 (t, 1H, J = 10.8 Hz), 2.10-2.07 (m, 1H), 1.76-1.68 (m, 1H); LCMS: 405.0





9		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35 (s, 1H), 7.16-7.03 (m, 3H), 4.68 (s, 2H), 4.37 (q, 1H, J = 8.4 Hz), 4.12 (d, 1H, J = 4.4 Hz), 4.02 (dd, 1H, J = 8.4, 4.4 Hz), 2.45-2.42 (m, 1H), 2.17-2.14 (m, 1H), 1.82-1.71 (m, 2H); LCMS: 405.1
10		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35 (s, 1H), 7.16-7.03 (m, 3H), 4.68 (s, 2H), 4.40-4.36 (m, 1H), 4.12 (d, 1H, J = 4.0 Hz), 4.02 (dd, 1H, J = 8.4, 4.4 Hz), 2.45-2.42 (m, 1H), 2.17-2.13 (m, 1H), 1.81-1.70 (m, 2H); LCMS: 405.1
11		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.16-7.10 (m, 2H), 7.04-7.03 (m, 1H), 4.68 (s, 2H), 4.07-4.01 (m, 1H), 4.00-3.97 (m, 2H), 3.67-3.65 (m, 1H), 3.52-3.51 (m, 1H), 3.27-3.24 (m, 1H), 2.19-2.15 (m, 1H), 1.83-1.79 (m, 1H); LCMS: 405.1
12		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.17 (s, 1H), 7.05-7.00 (m, 2H), 6.93-6.92 (m, 1H), 4.60-4.56 (m, 2H), 4.42-4.38 (m, 1H), 4.28-4.26 (m, 1H), 4.03-3.97 (m, 1H), 2.41-2.36 (m, 1H), 2.17-2.14 (m, 1H), 1.90-1.85 (m, 1H), 1.60-1.57 (m, 1H); LCMS: 405.1





13		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.15-7.09 (m, 2H), 7.03-7.02 (m, 1H), 4.65 (s, 2H), 4.53-4.49 (m, 1H), 4.39-4.35 (m, 1H), 4.12-4.07 (m, 1H), 2.50-2.45 (m, 1H), 2.26-2.25 (m, 1H), 1.99-1.92 (m, 1H), 1.70-1.66 (m, 1H); LCMS: 405.1
14		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.16-7.09 (m, 2H), 7.02-7.01 (m, 1H), 5.27-5.13 (m, 1H), 4.65 (s, 2H), 4.38-4.33 (m, 2H), 2.79-2.70 (m, 1H), 2.37-2.35 (m, 1H), 1.99-1.88 (m, 2H); LCMS: 407.1
15		¹H-NMR (400 MHz, CDCl₃) δ ppm 9.76 (s, 1H), 8.35 (s, 1H), 7.57 (t, 1H, J = 5.6 Hz), 7.05-6.90 (m, 3H), 5.16 (d, 1H, J = 52.4 Hz), 4.68-4.62 (m, 2H), 4.39-4.34 (m, 1H), 4.15-4.12 (m, 1H), 2.78-2.70 (m, 1H), 2.48-2.40 (m, 1H), 2.18-2.08 (m, 1H), 1.93-1.81 (m, 1H); LCMS: 407.1
16		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.26 (s, 1H), 7.12-7.07 (m, 2H), 7.02-7.00 (m, 1H), 5.15 (d, 1H, J = 53.6 Hz), 4.64 (s, 2H), 4.58-4.52 (m, 1H), 4.17 (dd, 1H, J = 12.8, 6.4 Hz), 2.59-2.51 (m, 2H), 1.99-1.87 (m, 2H); LCMS: 407.1





17		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26 (s, 1H), 7.14-7.08 (m, 2H), 7.02-7.00 (m, 1H), 5.13 (d, 1H, J = 53.2 Hz), 4.64 (s, 2H), 4.57-4.52 (m, 1H), 4.19-4.16 (m, 1H), 2.59-2.51 (m, 2H), 1.99-1.90 (m, 2H); LCMS: 407.1
18		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.14 (s, 1H), 7.04-6.97 (m, 2H), 6.91-6.90 (m, 1H), 4.89-4.88 (m, 1H), 4.55 (s, 2H), 4.53-4.46 (m, 1H), 3.98-3.89 (m, 1H), 2.38-2.3 (m, 1H), 2.18-2.00 (m, 1H), 1.90-1.83 (m, 1H), 1.56-1.52 (m, 1H); LCMS: 407.1
19		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.13 (s, 1H), 7.05-6.98 (m, 2H), 6.91-6.90 (m, 1H), 4.89-4.87 (m, 1H), 4.55 (s, 2H), 4.50-4.46 (m, 1H), 3.98-3.89 (m, 1H), 2.37-2.32 (m, 1H), 2.19-2.00 (m, 1H), 1.90-1.83 (m, 1H), 1.56-1.51 (m, 1H); LCMS: 407.1
20		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26 (s, 1H), 7.15-7.01 (m, 3H), 4.94-4.92 (m, 1H), 4.66 (s, 2H), 4.32-4.28 (m, 1H), 4.17-4.10 (m, 1H), 2.34-1.88 (m, 4H); LCMS: 407.2





21		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.17 (s, 1H), 7.05-6.92 (m, 3H), 4.86-4.83 (m, 1H), 4.56 (s, 2H), 4.23-4.21 (m, 1H), 4.08-4.02 (m, 1H), 2.24-1.78 (m, 4H); LCMS: 407.2
22		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (s, 1H), 7.58 (s, 1H), 7.47 (s, 2H), 4.67 (s, 2H), 4.39-4.37 (m, 1H), 4.13 (s, 1H), 4.02 (s, 1H), 2.44 (br.s, 1H), 2.16-2.15 (m, 1H), 1.79-1.69 (m, 2H); LCMS: 412.1
23		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37, 8.34 (s, 1H), 7.25-7.18 (m, 1H), 6.93-6.82 (m, 2H), 5.40 (s, 1H), 4.42-4.35 (m, 1H), 4.16-3.99 (m, 2H), 3.61 (s, 1.5H), 3.22 (s, 1.5H), 2.47-2.43 (m, 1H), 2.31 (s, 1.5H), 2.28 (s, 1.5H), 2.18-2.12 (m, 1H), 1.83-1.75 (m, 2H); LCMS: 415.1
24		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.07 (s, 1H), 9.45 (d, <i>J</i> = 7.7 Hz, 1H), 9.26 (t, <i>J</i> = 6.4 Hz, 1H), 8.27 (s, 1H), 7.11-6.81 (m, 3H), 6.51 (s, 0H), 5.10 (d, <i>J</i> = 4.8 Hz, 1H), 4.47 (d, <i>J</i> = 6.3 Hz, 2H), 4.04 (dd, <i>J</i> = 7.8, 4.0 Hz, 1H), 3.96 (dd, <i>J</i> = 11.2, 3.9 Hz, 1H), 3.82 (s, 3H), 3.81-3.71 (m, 1H), 3.65 (s, 1H), 3.43 (t, <i>J</i> = 9.3 Hz, 1H), 3.18 (dd, <i>J</i> = 11.2, 7.6 Hz, 1H), 1.93-1.85 (m, 1H), 1.53-1.43 (m, 1H); LCMS: 417





25		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.09 (s, 1H), 9.48 – 9.31 (m, 2H), 8.30 (s, 1H), 7.25 (td, <i>J</i> = 9.4, 4.5 Hz, 1H), 7.15 (td, <i>J</i> = 7.5, 6.7, 4.1 Hz, 2H), 4.81 (d, <i>J</i> = 4.1 Hz, 1H), 4.55 (d, <i>J</i> = 6.2 Hz, 2H), 4.40 (d, <i>J</i> = 7.1 Hz, 1H), 4.35 (d, <i>J</i> = 7.7 Hz, 2H), 3.64 (t, <i>J</i> = 5.1 Hz, 1H), 1.56 (dq, <i>J</i> = 8.7, 4.2 Hz, 1H), 1.32 (p, <i>J</i> = 4.6 Hz, 1H), 1.07 (q, <i>J</i> = 4.1 Hz, 1H), 0.43 (td, <i>J</i> = 8.3, 4.6 Hz, 1H); LCMS: 417
26		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37-8.36 (m, 1H), 7.17-7.07 (m, 3H), 4.81 (s, 2H), 4.16-4.13 (m, 2H), 3.66-3.60 (m, 2H), 2.40-2.27 (m, 1H), 2.20-2.03 (m, 1H), 1.91-1.72 (m, 4H), 1.40-1.24 (m, 3H); LCMS: 417.1
27		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.06 (s, 1H), 9.24 (d, <i>J</i> = 7.2 Hz, 1H), 9.11 (d, <i>J</i> = 8.6 Hz, 1H), 8.20 (s, 1H), 7.35 (ddd, <i>J</i> = 9.1, 5.7, 3.2 Hz, 1H), 7.20 (td, <i>J</i> = 9.3, 4.6 Hz, 1H), 7.10 (ddt, <i>J</i> = 9.0, 7.3, 3.6 Hz, 1H), 5.44 – 5.32 (m, 1H), 5.11 (t, <i>J</i> = 5.9 Hz, 1H), 5.02 – 4.91 (m, 1H), 4.16 (p, <i>J</i> = 6.6 Hz, 1H), 3.86 (ddt, <i>J</i> = 9.1, 6.2, 3.6 Hz, 1H), 3.77 – 3.60 (m, 2H), 2.15 – 2.02 (m, 1H), 1.78 (m, 1H), 1.64 (m, 2H), 1.42 (m, 2H); LCMS: 419
28		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.08 (br s, 1H), 9.30 (m, 1H), 8.28 (s, 1H), 7.31 (m, 1H), 7.20 (m, 2H), 5.36 (d, <i>J</i> = 2.4 Hz, 1H), 4.90 – 4.76 (m, 1H), 4.40 (q, <i>J</i> = 7.5 Hz, 1H), 3.94 (ddd, <i>J</i> = 11.7, 5.6, 5.1, 3.4 Hz, 1H), 3.72 (ddd, <i>J</i> = 22.6, 7.5, 4.5 Hz, 2H), 3.60 (br s, 1H), 3.07 (s, 3H), 2.32 – 2.19 (m, 1H), 1.94 (m, 1H), 1.58 (m, 1H), 1.41 – 1.26 (m, 1H); LCMS: 419





29		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 10.62-10.49 (m, 1H), 8.39 (d, 1H, J = 5.6 Hz), 7.10-6.97 (m, 3H), 5.47-5.34 (m, 1H), 4.83 (s, 1H), 4.26-4.16 (m, 2H), 4.02-3.89 (m, 2H), 3.60-3.45 (m, 3.5H), 3.13 (s, 1.5H), 2.17-2.13 (m, 1H), 1.81 (br.s, 1H); LCMS: 419.1
30		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 13.68 (br.s, 1H), 9.40 (d, 1H, J = 6.8 Hz), 8.33 (s, 1H), 7.61 (d, 1H, J = 9.2 Hz), 7.03-6.88 (m, 3H), 5.40 (t, 1H, J = 7.6 Hz), 4.13-4.09 (m, 2H), 3.95-3.92 (m, 1H), 3.78-3.76 (m, 1H), 3.42-3.34 (m, 2H), 2.04 (dd, 1H, J = 10.8, 2.0 Hz), 1.73-1.70 (m, 1H), 1.57 (d, 3H, J = 7.2 Hz); LCMS: 419.1
31		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.23-7.21 (m, 1H), 7.13-7.10 (m, 1H), 7.04-7.01 (m, 1H), 5.58-5.52 (m, 1H), 4.03-3.94 (m, 3H), 3.62-3.61 (m, 1H), 3.52-3.49 (m, 1H), 3.27-3.22 (m, 1H), 2.17-2.13 (m, 1H), 1.86-1.80 (m, 1H), 1.60 (d, 3H, J = 7.2 Hz); LCMS: 419.1
32		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.21 (s, 1H), 7.14-7.10 (m, 2H), 7.03-7.01 (m, 1H), 4.66 (s, 2H), 4.51-4.45 (m, 1H), 4.00 (s, 1H), 3.70-3.63 (m, 1H), 2.09-2.07 (m, 1H), 1.85-1.78 (m, 2H), 1.61-1.56 (m, 2H), 1.45-1.36 (m, 1H); LCMS: 419.1





33		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (d, 1H, J = 5.6 Hz), 6.99-6.96 (m, 2H), 6.92-6.88 (m, 1H), 5.43 (s, 1H), 4.90 (s, 1H), 4.43-4.37 (m, 1H), 4.15-4.13 (m, 1H), 4.05-4.01 (m, 1H), 3.64 (s, 1.5H), 3.16 (s, 1.5H), 2.46-2.43 (m, 1H), 2.19-2.15 (m, 1H), 1.82-1.65 (m, 2H); LCMS: 419.1
34		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.21 (br.s, 1H), 7.11-7.09 (m, 1H), 7.01 (br.s, 1H), 5.54 (d, 1H, J = 7.2 Hz), 4.39-4.32 (m, 1H), 4.10 (br.s, 1H), 3.99-3.96 (m, 1H), 2.43-2.40 (m, 1H), 2.14-2.12 (m, 1H), 1.78-1.74 (m, 1H), 1.69 (br.s, 1H), 1.59 (d, 3H, J = 6.8 Hz); LCMS: 419.2
35		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.21 (br.s, 1H), 7.13-7.09 (m, 1H), 7.03-7.01 (m, 1H), 5.55-5.51 (m, 1H), 4.37-4.31 (m, 1H), 4.11 (s, 1H), 4.02-3.99 (m, 1H), 2.42-2.38 (m, 1H), 2.15-2.11 (m, 1H), 1.78-1.76 (m, 1H), 1.67-1.64 (m, 1H), 1.59 (d, 3H, J = 6.8 Hz); LCMS: 419.2
36		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.10 (s, 1H), 9.49 (m, 1H), 9.37 (d, J = 7.3 Hz, 1H), 8.27 (s, 1H), 7.53 (dd, J = 8.8, 5.1 Hz, 1H), 7.18 (td, J = 8.4, 2.9 Hz, 1H), 7.12 (dd, J = 9.4, 3.5 Hz, 1H), 5.68 (d, J = 7.4 Hz, 1H), 4.87 (m, 1H), 4.56 (m, 1H), 4.48 (m, 1H), 4.39 (p, J = 7.6 Hz, 2H), 3.90 (m, 1H), 3.70 (m, 1H), 2.22 (m, 1H), 1.91 (m, 1H), 1.60-1.47 (m, 1H), 1.31 (m, 1H); LCMS: 421





37		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 13.80 (s, 1H), 9.43 (t, <i>J</i> = 6.2 Hz, 1H), 9.31 (d, <i>J</i> = 7.8 Hz, 1H), 8.22 (s, 1H), 7.47 (dd, <i>J</i> = 8.8, 5.1 Hz, 1H), 7.13 (td, <i>J</i> = 8.4, 3.1 Hz, 1H), 7.06 (dd, <i>J</i> = 9.6, 3.1 Hz, 1H), 5.06 (d, <i>J</i> = 4.8 Hz, 1H), 4.51 (d, <i>J</i> = 6.2 Hz, 2H), 3.99 (qd, <i>J</i> = 7.7, 3.8 Hz, 1H), 3.91 (dd, <i>J</i> = 11.1, 4.0 Hz, 1H), 3.72 (dt, <i>J</i> = 11.3, 4.6 Hz, 1H), 3.59 (tt, <i>J</i> = 8.5, 4.5 Hz, 1H), 3.38 (ddd, <i>J</i> = 11.7, 9.0, 3.1 Hz, 1H), 3.12 (dd, <i>J</i> = 11.1, 7.7 Hz, 1H), 1.84 (ddt, <i>J</i> = 9.2, 7.6, 3.6 Hz, 1H), 1.43 (dtd, <i>J</i> = 13.0, 8.7, 4.0 Hz, 1H); LCMS: 421
38		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.24 (s, 1H), 7.23 (s, 1H), 7.08 (d, 2H, <i>J</i> = 8.8 Hz), 4.60 (s, 2H), 4.51-4.46 (m, 1H), 4.13 (br.s, 1H), 3.94-3.90 (m, 1H), 2.46-2.40 (m, 1H), 2.11-2.06 (m, 1H), 1.78-1.74 (m, 1H), 1.62-1.58 (m, 1H); LCMS: 421.2
39		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 9.19 (d, <i>J</i> = 145.0 Hz, 1H), 8.08 (d, <i>J</i> = 31.1 Hz, 1H), 7.37 – 7.02 (m, 3H), 5.58 (s, 1H), 5.18 (s, 1H), 4.78 (s, 1H), 4.08 – 3.90 (m, 2H), 3.78 (s, 1H), 3.63 (s, 1H), 3.43 (d, <i>J</i> = 11.5 Hz, 1H), 3.17 (dd, <i>J</i> = 11.2, 7.4 Hz, 1H), 1.92 (s, 2H), 1.48 (s, 1H); LCMS: 422
40		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.19 (s, 1H), 9.51 (t, <i>J</i> = 6.3 Hz, 2H), 7.57 – 7.35 (m, 1H), 7.11 – 6.94 (m, 1H), 4.60 (d, <i>J</i> = 6.1 Hz, 2H), 4.04 (dt, <i>J</i> = 11.7, 5.8 Hz, 1H), 3.97 (dd, <i>J</i> = 11.1, 4.0 Hz, 1H), 3.78 (dd, <i>J</i> = 11.3, 5.1 Hz, 1H), 3.60 (tt, <i>J</i> = 6.6, 3.4 Hz, 1H), 3.48 – 3.40 (m, 1H), 3.19 (dd, <i>J</i> = 11.1, 7.7 Hz, 1H), 3.12 (qd, <i>J</i> = 7.3, 4.2 Hz, 2H), 1.91 (d, <i>J</i> = 13.1 Hz, 1H), 1.50 (dq, <i>J</i> = 8.7, 4.6 Hz, 1H); LCMS: 423





41		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.09-6.98 (m, 2H), 4.71 (s, 2H), 4.41-4.34 (m, 1H), 4.13 (br.s, 1H), 4.04-4.02 (m, 1H), 2.45-2.43 (m, 1H), 2.17-2.13 (m, 1H), 1.82-1.69 (m, 2H); LCMS: 423.1
42		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33 (s, 1H), 7.14-6.99 (m, 3H), 4.65 (s, 2H), 4.49-4.43 (m, 1H), 4.35-4.32 (m, 1H), 2.85-2.80 (m, 1H), 2.67-2.58 (m, 1H), 2.26-2.16 (m, 2H); LCMS: 425.1
43		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.28 (s, 1H), 7.13-6.99 (m, 3H), 4.65 (s, 2H), 4.52-4.46 (m, 1H), 4.33-4.28 (m, 1H), 2.87-2.79 (m, 1H), 2.64-2.59 (m, 1H), 2.22-2.14 (m, 3H); LCMS: 425.1
44		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33 (d, 1H, J = 6.0 Hz), 7.60 (d, 1H, J = 9.2 Hz), 7.52-7.49 (m, 2H), 5.43 (d, 1H, J = 3.6 Hz), 4.90 (d, 1H, J = 3.2 Hz), 4.43-4.38 (m, 1H), 4.13 (br.s, 1H), 4.03-3.96 (m, 1H), 3.65 (s, 1.5H), 3.17 (s, 1.5H), 2.48-2.44 (m, 1H), 2.17-2.13 (m, 1H), 1.83-1.66 (m, 2H); LCMS: 426.1





45		¹H-NMR (400 MHz, CDCl₃) δ ppm 11.58-11.21 (m, 1H), 8.47 (br.s, 0.5H), 6.98-6.55 (m, 3.5H), 6.33 (d, 0.5H, J = 8.0 Hz), 5.51-5.49 (m, 0.5H), 4.38-3.83 (m, 4H), 2.37-2.32 (m, 2H), 2.05-1.64 (m, 8H); LCMS: 429.1
46		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.50 (br.s, 1H), 8.40 (br.s, 1H), 8.38 (s, 1H), 7.72 (d, 1H, J = 9.2 Hz), 5.54-5.41 (m, 2H), 4.97-4.92 (m, 1H), 4.59-4.45 (m, 1H), 4.09-4.04 (m, 2H), 2.91-2.84 (m, 1H), 2.29-2.26 (m, 2H), 1.79-1.66 (m, 5H); LCMS: 430.1
47		¹H-NMR (400 MHz, CD₃OD) δ ppm 9.57 (s, 1H), 8.34 (s, 1H), 7.38-7.37 (m, 1H), 7.06-6.96 (m, 2H), 4.13-4.09 (m, 1H), 4.00-3.96 (m, 2H), 3.87-3.76 (m, 1H), 3.58-3.47 (m, 1H), 3.45-3.44 (m, 1H), 2.11-2.07 (m, 1H), 1.75-1.66 (m, 1H), 1.37 (s, 4H); LCMS: 431.1
48		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36 (d, 1H, J = 4.8 Hz), 6.73 (d, 1H, J = 12.0 Hz), 6.69-6.63 (m, 2H), 5.40 (s, 1H), 4.82 (s, 1H), 4.39-4.36 (m, 1H), 4.15-4.12 (m, 1H), 4.06-4.02 (m, 1H), 3.78 (d, 3H, J = 6.4 Hz), 3.62 (s, 1.5H), 3.14 (s, 1.5H), 2.47-2.44 (m, 1H), 2.18-2.15 (m, 1H), 1.82-1.70 (m, 2H); LCMS: 431.1





49		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35 (d, 1H, J = 3.6 Hz), 7.01-6.93 (m, 1H), 5.44-5.33 (m, 1H), 4.83 (s, 1H), 4.40-4.35 (m, 1H), 4.14-4.12 (m, 1H), 4.05-4.01 (m, 1H), 3.86 (s, 1.5H), 3.75 (s, 1.5H), 3.63 (s, 1.5H), 3.12 (s, 1.5H), 2.46-2.43 (m, 1H), 2.17-2.15 (m, 1H), 1.82-1.63 (m, 2H); LCMS: 431.1
50		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.60 (br.s, 1H), 8.50 (br.s, 1H), 8.43 (s, 1H), 7.89 (d, 1H, J = 9.2 Hz), 5.57-5.35 (m, 2H), 4.58-4.38 (m, 3H), 4.25-4.13 (m, 2H), 4.00-3.93 (m, 1H), 3.84-3.71 (m, 2H), 2.90-2.82 (m, 1H), 2.34-2.21 (m, 1H); LCMS: 432.1
51		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (s, 1H), 7.22-7.19 (m, 1H), 7.12-7.11 (m, 1H), 7.02-7.01 (m, 1H), 5.52-5.50 (m, 1H), 5.04-5.02 (m, 1H), 4.78-4.76 (m, 2H), 4.30-3.89 (m, 1H), 3.80-3.75 (m, 2H), 3.36-3.34 (m, 2H), 1.59 (d, 3H, J = 6.8 Hz), 1.50-1.34 (m, 3H); LCMS: 432.2
52		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35 (d, 1H, J = 6.0 Hz), 7.94 (dd, 1H, J = 9.2, 2.4 Hz), 7.47-7.41 (m, 1H), 5.31 (s, 1H), 4.78 (s, 1H), 4.41-4.35 (m, 1H), 4.14-4.11 (m, 1H), 4.04-4.02 (m, 1H), 3.97 (s, 1.5H), 3.89 (s, 1.5H), 3.70 (s, 1.5H), 3.17 (s, 1.5H), 2.47-2.40 (m, 1H), 2.18-2.14 (m, 1H), 1.81-1.64 (m, 2H); LCMS: 432.2





53		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.38 (d, 1H, <i>J</i> = 5.2 Hz), 7.20-7.05 (m, 3H), 4.87 (s, 2H), 4.17-4.11 (m, 2H), 4.00-3.97 (m, 3H), 3.64-3.44 (m, 3H), 2.13-2.06 (m, 1H), 1.76-1.70 (m, 1H), 1.39-1.18 (m, 3H); LCMS: 433.1
54		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.19 (s, 1H), 9.51 (t, <i>J</i> = 6.3 Hz, 2H), 7.57 – 7.35 (m, 1H), 7.11 – 6.94 (m, 1H), 4.60 (d, <i>J</i> = 6.1 Hz, 2H), 4.04 (dt, <i>J</i> = 11.7, 5.8 Hz, 1H), 3.97 (dd, <i>J</i> = 11.1, 4.0 Hz, 1H), 3.78 (dd, <i>J</i> = 11.3, 5.1 Hz, 1H), 3.60 (tt, <i>J</i> = 6.6, 3.4 Hz, 1H), 3.48 – 3.40 (m, 1H), 3.19 (dd, <i>J</i> = 11.1, 7.7 Hz, 1H), 3.12 (qd, <i>J</i> = 7.3, 4.2 Hz, 2H), 1.91 (d, <i>J</i> = 13.1 Hz, 1H), 1.50 (dq, <i>J</i> = 8.7, 4.6 Hz, 1H); LCMS: 435
55		LCMS: 435
56		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.23-7.21 (m, 1H), 7.13-7.12 (m, 1H), 7.11-7.03 (m, 1H), 5.57-5.54 (m, 1H), 4.14-4.01 (m, 1H), 4.00-3.92 (m, 4H), 3.91-3.90 (m, 1H), 3.61-3.44 (m, 2H), 2.05-2.02 (m, 1H), 1.72-1.68 (m, 1H); LCMS: 435.1





57		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (d, 1H, J = 14.8 Hz), 7.50-7.45 (m, 1H), 7.09-7.04 (m, 2H), 5.46 (s, 1H), 4.96 (s, 1H), 4.40-4.35 (m, 1H), 4.16-3.97 (m, 2H), 3.67 (s, 1.5H), 3.22 (s, 1.5H), 2.47-2.45 (m, 1H), 2.20-2.16 (m, 1H), 1.87-1.77 (m, 2H); LCMS: 435.1
58		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (d, 1H, J = 5.2 Hz), 7.23 (d, 1H, J = 12.0 Hz), 7.16-7.09 (m, 2H), 5.42 (s, 1H), 4.85 (s, 1H), 4.40-4.36 (m, 1H), 4.15-4.13 (m, 1H), 4.06-4.02 (m, 1H), 3.64 (s, 1.5H), 3.16 (s, 1.5H), 2.47-2.44 (m, 1H), 2.19-2.15 (m, 1H), 1.84-1.72 (m, 2H); LCMS: 435.1
59		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.24 (s, 1H), 6.94-6.89 (m, 2H), 4.64 (s, 2H), 4.52-4.48 (m, 1H), 4.14-4.12 (m, 1H), 3.95 (s, 3H), 3.93-3.92 (m, 1H), 2.43-2.40 (m, 1H), 2.10-2.08 (m, 1H), 1.78-1.75 (m, 1H), 1.61-1.57 (m, 1H); LCMS: 435.2
60		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26-8.22 (m, 2H), 7.64-7.59 (m, 1H), 5.34 (s, 1H), 4.89 (s, 1H), 4.51-4.47 (m, 1H), 4.15-4.09 (m, 1H), 3.95-3.86 (m, 1H), 3.68 (s, 1.2H), 3.22 (s, 1.8H), 2.46-2.40 (m, 1H), 2.11-2.09 (m, 1H), 1.80-1.49 (m, 2H); LCMS: 436.2





61		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.30 (s, 1H), 7.25-7.15 (m, 2H), 7.08-7.07 (m, 1H), 5.55-5.52 (m, 1H), 5.15 (d, 1H, J = 53.6 Hz), 4.60-4.55 (m, 1H), 4.21-4.18 (m, 1H), 3.95-3.88 (m, 2H), 2.62-2.51 (m, 2H), 2.00-1.89 (m, 2H); LCMS: 436.4
62		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.30 (s, 1H), 7.25-7.15 (m, 2H), 7.08-7.07 (m, 1H), 5.55-5.52 (m, 1H), 5.16 (d, 1H, J = 53.6 Hz), 4.60-4.54 (m, 1H), 4.18-4.14 (m, 1H), 3.94-3.89 (m, 2H), 2.60-2.51 (m, 2H), 1.99-1.90 (m, 2H); LCMS: 436.4
63		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.09 (br s, 1H), 9.25 (Abqd, 1H), 8.33 (d, J = 8.2 Hz, 1H), 7.67-7.47 (m, 1H), 7.14 (m, 1H), 5.46 (m, 1H), 4.96 (m, 1H), 4.59 (dd, J = 9.6, 4.0 Hz, 1H), 4.47 (q, J = 7.8 Hz, 1H), 4.00 (m, 1H), 3.84-3.71 (m, 1H), 3.58 (s, 1H), 3.40 (s, 3H), 2.42-2.26 (m, 1H), 2.08-1.92 (m, 1H), 1.72-1.57 (m, 1H), 1.37 (m, 1H); LCMS: 437
64		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.12 (s, 1H), 9.58-9.35 (m, 2H), 8.29 (s, 1H), 7.37-7.12 (m, 3H), 5.72 (d, J = 6.2 Hz, 1H), 4.67-4.49 (m, 2H), 4.29 (s, 1H), 3.76 (m, 1H), 2.04 (m, 2H), 1.99-1.79 (m, 1H), 1.71 (s, 1H), 1.47 (m, 2H); LCMS: 439





65		LCMS: 439
66		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.17 (s, 1H), 7.38-7.34 (m, 3H), 7.10-7.06 (m, 1H), 6.01-5.95 (m, 1H), 4.17-4.12 (m, 1H), 4.07-4.03 (m, 1H), 2.19-2.17 (m, 1H), 1.96-1.94 (m, 1H), 1.76-1.73 (m, 2H), 1.60-1.57 (m, 2H); LCMS: 439.2
67		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.15 (s, 1H), 7.38-7.35 (m, 3H), 7.10-7.05 (m, 1H), 6.02-5.96 (m, 1H), 4.24-4.14 (m, 1H), 4.03-3.99 (m, 1H), 2.24-2.19 (m, 1H), 1.94-1.93 (m, 1H), 1.77-1.74 (m, 2H), 1.60-1.55 (m, 2H); LCMS: 439.2
68		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.09 (s, 1H), 9.01 (d, <i>J</i> = 7.1 Hz, 1H), 8.19 (s, 1H), 7.31 (m, 1H), 7.16-7.10 (m, 2H), 7.00 (m, 1H), 5.43-5.30 (m, 2H), 4.79 (d, <i>J</i> = 5.6 Hz, 1H), 4.60 (m, 1H), 4.48-4.33 (m, 2H), 4.33-4.22 (m, 1H), 3.78 (m, 1H), 3.56 (m, 1H), 2.77-2.59 (m, 1H), 2.10 (m, 2H), 1.99-1.76 (m, 2H), 1.46 (m, 1H), 1.27-1.15 (m, 1H); LCMS: 445





69		¹H-NMR (400 MHz, CDCl₃) δ ppm 11.87-11.48 (m, 1H), 8.50-8.46 (m, 1H), 7.06-6.65 (m, 3H), 6.40-6.39 (m, 0.5H), 5.62-5.59 (m, 0.5H), 4.46-3.95 (m, 6H), 3.55-3.51 (m, 2H), 2.42-2.11 (m, 1H), 2.04-1.76 (m, 5H); LCMS: 445.1
70		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.35, 8.28 (s, 1H), 7.14-6.80 (m, 3H), 6.43 (d, 0.5H, J = 7.6 Hz), 5.60 (dd, 0.5H, J = 7.6, 4.0 Hz), 4.52-4.31 (m, 2H), 4.05-3.89 (m, 3H), 2.48-2.41 (m, 2H), 2.12-2.06 (m, 4H), 1.94-1.61 (m, 2H); LCMS: 445.1
71		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33, 8.27 (s, 1H), 7.14-6.92 (m, 3H), 6.43 (d, 0.5H, J = 8.0 Hz), 5.58 (dd, 0.5H, J = 8.0, 4.4 Hz), 4.39-4.31 (m, 2H), 4.15-4.08 (m, 2H), 3.94-3.91 (m, 1H), 2.48-2.43 (m, 2H), 2.12-2.06 (m, 3H), 1.76-1.51 (m, 3H); LCMS: 445.1
72		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.42 (br.s, 1H), 8.31 (br.s, 1H), 8.26 (s, 1H), 7.63-7.61 (m, 1H), 5.46-5.26 (m, 2H), 4.50-4.37 (m, 1H), 4.26-4.21 (m, 1H), 3.97 (br.s, 2H), 3.80-3.77 (m, 1H), 2.80-2.76 (m, 1H), 2.33-2.30 (m, 1H), 2.29-2.27 (m, 1H), 2.07-2.00 (m, 1H), 1.69-1.67 (m, 1H), 1.52-1.51 (m, 1H); LCMS: 446.1





73		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.52 (s, 1H), 8.41 (s, 1H), 8.35-8.29 (m, 1H), 7.76 (d, 1H, J = 9.2 Hz), 5.57-5.36 (m, 2H), 4.96-4.94 (m, 1H), 4.60-4.51 (m, 1H), 3.98-3.89 (m, 3H), 3.50-3.47 (m, 2H), 3.24-3.21 (m, 1H), 2.88-2.86 (m, 1H), 2.18-2.09 (m, 2H), 1.75-1.71 (m, 1H); LCMS: 446.1
74		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.60 (br.s, 2H), 8.35 (s, 1H), 7.75 (d, 1H, J = 9.2 Hz), 5.57-5.35 (m, 2H), 4.59-4.45 (m, 1H), 4.09-4.05 (m, 1H), 3.95-3.91 (m, 2H), 3.65-3.75 (m, 1H), 3.53-3.50 (m, 1H), 3.40-3.35 (m, 1H), 2.88-2.86 (m, 1H), 2.29-2.27 (m, 2H), 1.97-1.93 (m, 1H), 1.66-1.63 (m, 1H); LCMS: 446.2
75		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.89-8.87 (m, 0.5H), 8.52-8.43 (m, 1H), 8.39 (s, 1H), 8.13-8.10 (m, 0.5H), 7.77 (d, 1H, J = 9.2 Hz), 5.55-5.40 (m, 2H), 5.16-5.01 (m, 2H), 4.60-4.37 (m, 2H), 4.17-4.15 (m, 1H), 2.89-2.86 (m, 1H), 2.66-2.56 (m, 1H), 2.44-2.42 (m, 1H), 2.30-2.20 (m, 1H), 1.98-1.88 (m, 2H); LCMS: 448.1
76		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.87-8.86 (m, 0.5H), 8.67-8.50 (m, 1H), 8.46 (s, 1H), 8.12-8.10 (m, 0.5H), 7.75-7.73 (m, 1H), 5.54-5.24 (m, 2H), 5.16-5.03 (m, 2H), 4.58-4.37 (m, 2H), 4.11-4.07 (m, 1H), 2.87-2.85 (m, 1H), 2.55-2.47 (m, 2H), 2.26-2.01 (m, 1H), 1.95-1.84 (m, 2H); LCMS: 448.1





77		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.43 (d, 1H, J = 5.2 Hz), 7.06-6.99 (m, 1H), 6.96-6.87 (m, 1H), 5.57-5.48 (m, 1H), 5.00 (s, 1H), 4.45-4.41 (m, 1H), 4.20-4.19 (m, 1H), 4.12-4.07 (m, 1H), 3.98 (s, 1.5H), 3.89 (s, 1.5H), 3.71 (s, 1.5H), 3.22 (s, 1.5H), 2.53-2.49 (m, 1H), 2.25-2.21 (m, 1H), 1.90-1.79 (m, 2H); LCMS: 449.1
78		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.14-7.00 (m, 3H), 5.59-5.55 (m, 1H), 5.48-5.32 (m, 1H), 4.46-4.33 (m, 3H), 4.23-4.12 (m, 2H), 3.97-3.94 (m, 1H), 3.69-3.68 (m, 2H), 2.85-2.75 (m, 1H), 2.16-2.09 (m, 1H); LCMS: 449.1
79		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.41 (s, 1H), 8.35 (s, 1H), 7.20-7.16 (m, 1H), 7.07-7.01 (m, 2H), 5.76-5.73 (m, 1H), 4.84-4.75 (m, 1H), 4.25-4.08 (m, 3H), 3.64-3.61 (m, 1H), 2.81-2.78 (m, 1H), 2.41-2.38 (m, 2H), 2.00-1.68 (m, 5H); LCMS: 453.5
80		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.35 (s, 1H), 9.60 (q, J = 7.8, 7.0 Hz, 1H), 8.33 (s, 1H), 7.84 (dd, J = 8.7, 5.4 Hz, 1H), 7.39 - 7.16 (m, 2H), 4.72 (d, J = 6.1 Hz, 2H), 4.04 (qd, J = 7.9, 4.0 Hz, 1H), 3.96 (dd, J = 11.1, 4.0 Hz, 1H), 3.78 (dt, J = 11.5, 4.5 Hz, 1H), 3.65 (td, J = 8.2, 4.3 Hz, 1H), 3.48 - 3.38 (m, 1H), 3.23 - 3.15 (m, 1H), 1.89 (dq, J = 12.7, 4.1 Hz, 1H), 1.49 (dd, J = 13.2, 8.9, 4.0 Hz, 1H), 1.25 (dt, J = 13.3, 7.9 Hz, 1H); LCMS: 455





81		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.80-7.71 (m, 1H), 7.29-7.27 (m, 1H), 7.21-7.17 (m, 1H), 4.74-4.67 (m, 2H), 4.44-4.30 (m, 1H), 4.11 (s, 1H), 4.02-4.01 (m, 1H), 2.50-2.40 (m, 1H), 2.15-2.12 (m, 1H), 1.78-1.77 (m, 1H), 1.70-1.67 (m, 1H); LCMS: 455.1
82		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.39 (s, 1H), 8.33 (s, 1H), 7.22-6.90 (m, 3H), 5.74 (br.s, 1H), 4.83-4.73 (m, 1H), 4.51-4.42 (m, 2H), 4.26-4.17 (m, 2H), 3.98-3.95 (m, 1H), 3.82-3.71 (m, 3H), 2.87-2.75 (m, 1H), 2.53-2.50 (m, 1H); LCMS: 455.4
83		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.12 (s, 1H), 9.51 (t, <i>J</i> = 6.2 Hz, 1H), 9.33 (d, <i>J</i> = 7.7 Hz, 1H), 8.28 (s, 1H), 7.64 (dd, <i>J</i> = 8.2, 3.0 Hz, 1H), 7.15 (dd, <i>J</i> = 9.2, 3.1 Hz, 1H), 5.09 (d, <i>J</i> = 4.9 Hz, 1H), 4.60 (d, <i>J</i> = 6.1 Hz, 2H), 4.05 (dd, <i>J</i> = 7.7, 4.1 Hz, 1H), 3.96 (dd, <i>J</i> = 11.2, 4.1 Hz, 1H), 3.82-3.75 (m, 1H), 3.65 (d, <i>J</i> = 4.9 Hz, 1H), 3.43 (t, <i>J</i> = 9.4 Hz, 1H), 3.17 (dd, <i>J</i> = 11.2, 7.6 Hz, 1H), 1.89 (d, <i>J</i> = 12.9 Hz, 1H), 1.52-1.44 (m, 1H); LCMS: 456
84		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.11 (s, 1H), 9.60-9.39 (m, 2H), 8.28 (s, 1H), 7.32-7.24 (m, 1H), 7.17 (m, 1H), 5.21 (d, <i>J</i> = 5.2 Hz, 1H), 4.56 (d, <i>J</i> = 6.1 Hz, 2H), 4.29 (m, 1H), 3.96 (d, <i>J</i> = 6.7 Hz, 1H), 3.29-3.18 (m, 1H), 2.97 (s, 3H), 2.84 (s, 3H), 2.30 (m, 1H), 2.14 (m, 1H), 1.78-1.61 (m, 2H); LCMS: 460





85		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.51 (s, 1H), 8.41 (s, 1H), 8.32 (s, 1H), 7.76 (d, 1H, J = 9.2 Hz), 5.56-5.40 (m, 2H), 4.87-4.82 (m, 1H), 4.59-4.50 (m, 1H), 4.11-3.97 (m, 2H), 3.48-3.45 (m, 1H), 2.88-2.86 (m, 1H), 2.27-1.80 (m, 2H), 1.78-1.67 (m, 2H), 1.59-1.46 (m, 3H); LCMS: 460.1
86		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.25-8.19 (m, 1H), 7.14-6.85 (m, 3H), 5.63-5.59 (m, 1H), 5.42 (d, 1H, J = 52.4 Hz), 4.83 (br.s, 1H), 4.48-4.31 (m, 3H), 4.12-3.99 (m, 1H), 2.88-2.80 (m, 1H), 2.42-2.39 (m, 1H), 2.25-2.20 (m, 2H), 1.92-1.89 (m, 1H), 1.69-1.64 (m, 1H); LCMS: 462.4
87		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26-8.19 (m, 1H), 7.15-6.86 (m, 3H), 5.63-5.59 (m, 1H), 5.42 (d, 1H, J = 52.0 Hz), 4.85-4.80 (m, 1H), 4.47-4.31 (m, 3H), 4.11-4.03 (m, 1H), 2.83-2.78 (m, 1H), 2.49-2.43 (m, 1H), 2.23-2.20 (m, 2H), 1.87-1.83 (m, 1H), 1.70-1.65 (m, 1H); LCMS: 462.4
88		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.25 (s, 1H), 7.17-7.03 (m, 3H), 5.62 (t, 1H, J = 8.8 Hz), 5.43 (d, 1H, J = 52.0 Hz), 4.86-4.81 (m, 1H), 4.50-4.40 (m, 2H), 4.12-4.08 (m, 1H), 3.93-3.84 (m, 1H), 2.89-2.79 (m, 1H), 2.41-2.38 (m, 1H), 2.12-2.05 (m, 2H), 1.76-1.75 (m, 1H), 1.56-1.55 (m, 1H); LCMS: 463





89		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.02 (s, 1H), 9.23 (d, <i>J</i> = 7.0 Hz, 1H), 8.24 (s, 1H), 7.41 (qd, <i>J</i> = 9.4, 4.9 Hz, 1H), 7.12 (ddt, <i>J</i> = 11.4, 7.6, 2.0 Hz, 1H), 5.48 (m, 1H), 4.90 (m, 1H), 4.46 (m, 1H), 4.36 (m, 2H), 4.09 (m, 1H), 3.89–3.83 (m, 1H), 3.59 (m, 1H), 2.49–2.38 (m, 1H), 2.30–2.10 (m, 2H), 1.97–1.79 (m, 3H), 1.54 (m, 2H), 1.37–1.24 (m, 2H); LCMS: 463
90		LCMS: 463
91		LCMS: 463
92		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.14–6.86 (m, 3H), 5.63–5.59 (m, 1H), 5.51–5.35 (m, 1H), 4.50–4.41 (m, 1H), 4.10–4.06 (m, 1H), 3.98–3.92 (m, 2H), 3.75–3.70 (m, 1H), 3.52–3.41 (m, 1H), 3.39–3.36 (m, 1H), 2.84–2.81 (m, 1H), 2.25–2.00 (m, 1H), 2.00–1.96 (m, 1H), 1.67–1.64 (m, 1H), 1.38–1.28 (m, 1H); LCMS: 463.1





93		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.14-8.04 (m, 1H), 7.02-6.74 (m, 3H), 5.73-5.48 (m, 1H), 5.37-5.24 (m, 1H), 4.36-4.10 (m, 2H), 3.87-3.76 (m, 2H), 3.44-3.38 (m, 2H), 3.18-3.10 (m, 1H), 2.76-2.72 (m, 1H), 2.10-1.88 (m, 2H), 1.52 (br.s, 1H), 1.33-1.17 (m, 1H); LCMS: 463.1
94		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.53 (s, 1H), 8.46-8.43 (m, 1H), 8.29-8.24 (m, 1H), 7.81-7.66 (m, 1H), 5.81-5.78 (m, 1H), 4.89-4.85 (m, 1H), 4.57-4.52 (m, 2H), 4.50-4.48 (m, 1H), 4.13-4.11 (m, 1H), 3.88-3.85 (m, 1H), 3.12-3.11 (m, 1H), 2.62-2.59 (m, 1H), 2.44-2.42 (m, 1H), 2.08-2.05 (m, 1H), 1.82-1.77 (m, 1H), 1.55-1.53 (m, 1H); LCMS: 464.1
95		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.15-8.09 (m, 1H), 7.03-6.74 (m, 3H), 5.52-5.47 (m, 1H), 5.31 (d, 1H, J = 52.0 Hz), 5.06-4.92 (m, 2H), 4.44-4.38 (m, 2H), 3.99-3.96 (m, 1H), 2.72-2.70 (m, 1H), 2.45-2.32 (m, 2H), 1.84-1.75 (m, 3H); LCMS: 465.1
96		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.22-8.10 (m, 1H), 7.05-6.75 (m, 3H), 5.51-5.47 (m, 1H), 5.31 (d, 1H, J = 52.0 Hz), 4.99 (d, 1H, J = 54.0 Hz), 4.44-4.29 (m, 2H), 4.02-3.99 (m, 1H), 2.90-2.72 (m, 1H), 2.47-2.35 (m, 2H), 2.12-1.64 (m, 3H); LCMS: 465.1





97		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.28 (s, 1H), 7.15-7.11 (m, 2H), 7.04-7.02 (m, 1H), 4.66 (s, 2H), 4.51-4.45 (m, 1H), 4.28-4.23 (m, 1H), 3.84-3.80 (m, 1H), 2.96 (s, 3H), 2.75-2.70 (m, 1H), 2.53-2.50 (m, 1H), 2.15-2.06 (m, 2H); LCMS: 467.1
98		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.28 (s, 1H), 7.15-7.11 (m, 2H), 7.09-7.02 (m, 1H), 4.66 (s, 2H), 4.51-4.45 (m, 1H), 4.28-4.25 (m, 1H), 3.84-3.80 (m, 1H), 2.96 (s, 3H), 2.75-2.70 (m, 1H), 2.54-2.52 (m, 1H), 2.13-2.06 (m, 2H); LCMS: 467.1
99		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.40, 8.36 (d, 1H, J = 11.6 Hz), 7.85-7.80 (m, 1H), 7.25-7.17 (m, 2H), 5.60 (s, 1H), 5.57 (s, 1H), 4.40-4.35 (m, 1H), 4.16-3.98 (m, 2H), 3.69 (s, 1.5H), 3.23 (s, 1.5H), 2.49-2.43 (m, 1H), 2.21-2.20 (m, 1H), 1.85-1.64 (m, 2H); LCMS: 469.1
100		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.28-8.22 (m, 1H), 7.19-6.90 (m, 3H), 5.78-5.76 (m, 1H), 4.85-4.80 (m, 1H), 4.71-4.69 (m, 1H), 4.19-4.17 (m, 2H), 4.09-4.08 (m, 2H), 3.75-3.74 (m, 1H), 3.59-3.48 (m, 2H), 2.86-2.76 (m, 1H), 2.51-2.35 (m, 1H), 2.13-1.96 (m, 1H), 1.71-1.60 (m, 1H); LCMS: 469.4





101		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.35 (br.s, 1H), 7.20 (d, 1H, J = 6.8 Hz), 7.13 (t, 1H, J = 8.0 Hz), 4.71 (s, 2H), 4.41-4.35 (m, 1H), 4.12 (br.s, 1H), 4.01-3.99 (m, 1H), 2.45-2.40 (m, 1H), 2.15-2.11 (m, 1H), 1.80-1.68 (m, 2H); LCMS: 471.1
102		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 11.11 (d, 1H, J = 6.0 Hz), 8.67 (s, 1H), 7.84-7.82 (m, 1H), 7.04-6.92 (m, 3H), 4.64 (d, 2H, J = 6.4 Hz), 4.41 (br.s, 1H), 4.29 (br.s, 1H), 3.42-3.38 (m, 2H), 3.25-3.22 (m, 2H), 3.09-3.07 (m, 1H), 2.79-2.77 (m, 1H); LCMS: 472.2
103		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 14.32 (s, 1H), 10.13 (d, J = 9.9 Hz, 1H), 8.99 (d, J = 7.6 Hz, 1H), 8.30 (s, 1H), 8.14 - 8.03 (m, 1H), 7.41 (m, 2H), 6.40 (m, 1H), 5.11 (s, 1H), 4.03 (m, 3H), 3.80 (m, 1H), 3.67 (m, 1H), 3.54 - 3.42 (m, 1H), 3.21 (m, 1H), 1.90 (dq, J = 12.7, 4.3 Hz, 1H), 1.50 (dq, J = 13.7, 5.1 Hz, 1H); LCMS: 473
104		LCMS: 475





105		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32-8.25 (m, 1H), 7.15-6.87 (m, 3H), 5.62-5.58 (m, 1H), 5.51-5.38 (m, 1H), 4.94-4.90 (m, 1H), 4.51-4.47 (m, 1H), 4.14-4.07 (m, 1H), 4.02 (br.s, 1H), 3.52-3.49 (m, 1H), 2.83-2.79 (m, 1H), 2.27-2.24 (m, 1H), 2.14-2.05 (m, 1H), 1.88-1.80 (m, 2H), 1.60-1.48 (m, 3H); LCMS: 477.1
106		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32-8.25 (m, 1H), 7.15-6.87 (m, 3H), 5.62-5.58 (m, 1H), 5.51-5.38 (m, 1H), 4.94-4.90 (m, 1H), 4.51-4.47 (m, 1H), 4.14-4.07 (m, 1H), 4.02 (br.s, 1H), 3.52-3.49 (m, 1H), 2.83-2.79 (m, 1H), 2.27-2.24 (m, 1H), 2.14-2.05 (m, 1H), 1.88-1.80 (m, 2H), 1.60-1.48 (m, 3H); LCMS: 477.1
107		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35-8.27 (m, 1H), 7.16-7.01 (m, 3H), 5.63-5.59 (m, 1H), 5.44 (d, 1H, J = 52.0 Hz), 4.51-4.32 (m, 2H), 4.04-3.75 (m, 5H), 3.60-3.57 (m, 1H), 3.39-3.38 (m, 1H), 2.86-2.80 (m, 1H), 2.27-2.13 (m, 1H); LCMS: 479
108		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.22-8.16 (m, 1H), 7.15-7.11 (m, 1H), 7.05-7.01 (m, 1H), 6.94-6.92 (m, 1H), 5.79-5.75 (m, 1H), 4.82-4.75 (m, 1H), 4.46-4.41 (m, 1H), 4.04-4.02 (m, 1H), 3.82-3.79 (m, 1H), 2.98-2.94 (m, 1H), 2.52-2.36 (m, 2H), 2.01-1.92 (m, 1H), 1.71-1.60 (m, 1H), 1.32-1.29 (m, 1H), 1.23-1.19 (m, 1H); LCMS: 481.1





109		¹H-NMR (400 MHz, CDCl₃) δ ppm 11.26 (d, 1H, J = 6.0 Hz), 8.67 (s, 1H), 7.94-7.91 (m, 1H), 7.00-6.76 (m, 3H), 4.55 (d, 2H, J = 6.0 Hz), 4.39 (br.s, 1H), 4.28 (br.s, 1H), 3.82 (s, 3H), 3.39-3.35 (m, 2H), 3.25-3.22 (m, 2H), 3.09-3.08 (m, 1H), 2.79-2.78 (m, 1H); LCMS: 484.2
110		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.40, 8.35 (s, 1H), 7.40 (br.s, 1H), 7.20-7.14 (m, 2H), 5.51 (s, 1H), 4.94 (d, 1H, J = 9.6 Hz), 4.41-4.36 (m, 1H), 4.15-3.97 (m, 2H), 3.65 (s, 1.5H), 3.18 (s, 1.5H), 2.48-2.42 (m, 1H), 2.19-2.14 (m, 1H), 1.82-1.65 (m, 2H); LCMS: 485.1
111		¹H-NMR (400 MHz, CDCl₃) δ ppm 8.77 (br.s, 1H), 7.83 (d, 1H, J = 8.4 Hz), 7.10-6.97 (m, 3H), 5.45-5.41 (m, 1H), 4.44 (br.s, 1H), 4.35 (br.s, 1H), 3.51-3.43 (m, 2H), 3.36-3.33 (m, 2H), 3.21 (d, 1H, J = 8.0 Hz), 2.85 (d, 1H, J = 2.0 Hz), 1.67 (d, 1H, J = 6.8 Hz); LCMS: 486.2
112		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.38-8.32 (m, 1H), 7.17-6.67 (m, 3H), 5.77 (dd, 1H, J = 9.2, 6.0 Hz), 4.83-4.77 (m, 1H), 4.40-4.37 (m, 1H), 4.30-4.26 (m, 1H), 3.94-3.76 (m, 1H), 3.05-2.99 (m, 1H), 2.57-2.50 (m, 1H), 1.70-1.65 (m, 1H), 1.48-1.46 (m, 1H), 1.25-1.23 (m, 1H), 0.64-0.61 (m, 1H); LCMS: 493.1





113		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 11.10 (d, 1H, J = 6.0 Hz), 8.69 (s, 1H), 7.80 (s, 1H), 7.28-7.26 (m, 1H), 6.92-6.87 (m, 2H), 4.33 (br.s, 1H), 4.26 (br.s, 1H), 3.46-3.44 (m, 1H), 3.31-3.25 (m, 1H), 3.12-3.10 (m, 1H), 2.73-2.72 (m, 1H), 1.32-1.24 (m, 4H).; LCMS: 498.2
114		LCMS: 513
115		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26 (s, 1H), 7.15-7.01 (m, 3H), 5.62-5.57 (m, 1H), 5.49-5.36 (m, 1H), 4.44-4.41 (m, 2H), 4.17-4.12 (m, 1H), 3.75-3.68 (m, 1H), 2.96 (s, 3H), 2.80-2.70 (m, 1H), 2.68-2.64 (m, 1H), 2.41-2.39 (m, 1H), 2.10-2.00 (m, 4H).; LCMS: 525.1
116		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26 (s, 1H), 7.15-7.01 (m, 3H), 5.61-5.57 (m, 1H), 5.49-5.36 (m, 1H), 4.44-4.41 (m, 2H), 4.17-4.12 (m, 1H), 3.75-3.68 (m, 1H), 2.96 (s, 3H), 2.80-2.70 (m, 1H), 2.68-2.64 (m, 1H), 2.41-2.39 (m, 1H), 2.10-2.00 (m, 4H).; LCMS: 525.1





117		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (s, 1H), 7.01-6.96 (m, 3H), 4.62 (s, 2H), 4.37-4.31 (m, 1H), 4.13 (br.s, 1H), 4.04-4.01 (m, 1H), 3.88 (s, 3H), 2.26-2.24 (m, 1H), 2.46-2.39 (m, 1H), 2.16-2.14 (m, 1H), 1.82-1.72 (m, 2H); LCMS: ND
118		¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 14.01 (d, <i>J</i> = 41.7 Hz, 1H), 9.31 (d, <i>J</i> = 7.6 Hz, 1H), 8.98 (d, <i>J</i> = 7.5 Hz, 1H), 8.29 (d, <i>J</i> = 9.0 Hz, 1H), 7.36-7.07 (m, 3H), 5.33 (s, 1H), 4.81 (dd, <i>J</i> = 9.2, 5.0 Hz, 2H), 4.48-4.23 (m, 3H), 3.72-3.55 (m, 1H), 3.49 (s, 2H), 3.05 (s, 1H), 1.56 (dt, <i>J</i> = 29.6, 6.3 Hz, 1H), 1.38-1.20 (m, 1H), 1.08 (dq, <i>J</i> = 8.5, 4.1 Hz, 1H)
119		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 7.18-7.04 (m, 3H), 5.61 (t, 1H, <i>J</i> = 8.4 Hz), 5.47 (d, 1H, <i>J</i> = 51.6 Hz), 4.51-4.38 (m, 1H), 4.14-4.08 (m, 2H), 2.89-2.81 (m, 1H), 2.33-1.69 (m, 8H); LCMS: 447
120		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.28 (s, 0.2H), 8.26 (s, 0.8H), 7.24-7.20 (m, 1H), 7.11-7.09 (m, 1H), 5.77-5.67 (m, 1H), 5.04-4.95 (m, 1H), 4.33-4.27 (m, 1H), 4.05-4.04 (m, 1H), 3.93-3.89 (m, 1H), 3.43 (s, 1.8H), 2.92 (s, 1.2H), 2.40-2.35 (m, 1H), 2.08-2.07 (m, 1H), 1.72-1.71 (m, 1H), 1.62-1.59 (m, 1H); LCMS: 453





125		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.24-8.17 (m, 1H), 7.14-6.84 (m, 3H), 5.61-5.57 (m, 1H), 5.41 (d, 1H, J = 51.6 Hz), 4.89-4.86 (m, 1H), 4.46-4.36 (m, 2H), 4.00-3.96 (m, 1H), 3.88-3.86 (m, 1H), 3.29 (s, 3H), 2.82-2.75 (m, 1H), 2.38-2.25 (m, 3H), 1.86-1.80 (m, 1H), 1.67-1.64 (m, 1H); LCMS: 477
126		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.19 (s, 1H), 6.96-6.94 (m, 2H), 6.80-6.77 (m, 1H), 5.44-5.30 (m, 2H), 4.79-4.76 (m, 1H), 4.48-4.38 (m, 2H), 3.87 (br.s, 1H), 3.65-3.53 (m, 1H), 2.84-2.74 (m, 1H), 2.05-2.00 (m, 2H), 1.78-1.72 (m, 2H), 1.53-1.35 (m, 3H); LCMS: 477.1
127		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.23-8.16 (m, 1H), 7.21 (s, 1H), 7.08-7.05 (m, 2H), 5.44-5.31 (m, 2H), 4.80-4.79 (m, 1H), 4.50-4.41 (m, 2H), 4.06-4.04 (m, 1H), 3.85-3.82 (m, 1H), 2.83-2.77 (m, 1H), 2.38-2.36 (m, 1H), 2.07-2.00 (m, 2H), 1.74-1.71 (m, 1H), 1.52-1.49 (m, 1H); LCMS: 479
128		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.23-8.20 (m, 2H), 7.59 (dd, 1H, J = 8.8, 3.2 Hz), 5.67-5.62 (m, 1H), 5.40 (d, 1H, J = 52.8 Hz), 4.63-4.40 (m, 3H), 4.03-4.01 (m, 1H), 3.80 (dd, 1H, J = 7.2, 4.4 Hz), 2.95-2.87 (m, 1H), 2.37-2.35 (m, 1H), 2.06-1.99 (m, 2H), 1.71-1.68 (m, 1H), 1.49-1.46 (m, 1H); LCMS: 480





129		ND
130		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.19-8.12 (m, 1H), 7.20 (s, 1H), 7.05 (d, 2H, J = 8.8 Hz), 5.43-5.30 (m, 2H), 4.77-4.76 (m, 1H), 4.49-4.39 (m, 2H), 3.87 (br.s, 1H), 3.56-3.53 (m, 1H), 2.81-2.75 (m, 1H), 2.15-2.13 (m, 1H), 1.81-1.73 (m, 2H), 1.58-1.35 (m, 4H); LCMS: 493.1
131		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.38 (br.s, 1H), 7.32-7.22 (m, 5H), 5.42-5.28 (m, 1H), 4.84-4.69 (m, 1H), 4.55 (br.s, 1H), 4.22 (br.s, 1H), 4.04 (br.s, 1H), 3.51 (s, 1.5H), 3.07 (s, 1.5H), 2.42 (br.s, 1H), 2.13 (br.s, 1H), 1.87 (br.s, 1H), 1.71 (br.s, 1H); LCMS: 383.2
132		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 9.80 (s, 0.5H), 9.64 (s, 0.5H), 8.38 (d, 1H, J = 5.2 Hz), 7.35-7.27 (m, 1H), 7.13-7.02 (m, 3H), 5.40 (s, 1H), 4.82 (d, 1H, J = 4.8 Hz), 4.38 (br.s, 1H), 4.22 (s, 1H), 3.95 (s, 1H), 3.57 (s, 1.5H), 3.09 (s, 1.5H), 2.52 (br.s, 1H), 2.10-2.05 (m, 1H), 1.96 (m, 1H), 1.89-1.82 (m, 1H); LCMS: 401.1





133		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35-8.33 (m, 1H), 8.16-8.12 (m, 1H), 7.97-7.88 (m, 1H), 7.34-7.30 (m, 1H), 5.47 (s, 1H), 4.85 (s, 1H), 4.38 (br.s, 1H), 4.15-4.12 (m, 1H), 4.03-3.98 (m, 1H), 3.70 (s, 1.5H), 3.35-3.25 (m, 1H), 3.22 (s, 1.5H), 2.48-2.40 (m, 1H), 2.14 (br.s, 1H), 1.81-1.75 (m, 1H); LCMS: 402.1
134		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (d, 1H, J = 4.8 Hz), 7.76-7.67 (m, 3H), 7.58-7.56 (m, 1H), 5.47 (d, 1H, J = 2.4 Hz), 4.91 (s, 1H), 4.42-4.37 (m, 1H), 4.15-4.14 (m, 1H), 4.04-3.99 (m, 1H), 3.65 (s, 1.5H), 3.16 (s, 1.5H), 2.48-2.45 (m, 1H), 1.84-1.81 (m, 1H), 1.71-1.66 (m, 2H); LCMS: 408.1
135		¹ H NMR (400 MHz, DMSO-d ₆) δ 13.98 (s, 1H), 9.26 (m, 1H), 8.26 (d, J = 11.5 Hz, 1H), 7.99 (m, 1H), 7.45 - 7.28 (m, 2H), 5.47 (m, 1H), 5.02 (m, 1H), 4.25 (m, 1H), 3.94 (m, 1H), 3.56 (s, 1H), 3.30 (s, 3H), 3.09 (s, 1H), 2.22 - 2.03 (m, 1H), 1.91 (m, 1H), 1.84 - 1.64 (m, 2H), 1.64 - 1.34 (m, 2H); LCMS: 410.2
136		¹ H NMR (400 MHz, DMSO-d ₆) δ 13.89 (s, 1H), 9.10 (ABqd, 1H), 8.21 (d, J = 10.6 Hz, 1H), 7.87 (ddd, J = 8.3, 4.7, 3.3 Hz, 1H), 7.33 (ddd, J = 9.4, 7.9, 3.2 Hz, 1H), 4.97 (m, 2H), 4.46 (s, 1H), 4.27 - 4.12 (m, 1H), 3.99 - 3.81 (m, 1H), 3.39 (d, J = 6.3 Hz, 3H), 3.29 (s, 3H), 2.18 - 2.01 (m, 1H), 1.86 - 1.35 (m, 5H); LCMS: 416.1





137		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (s, 1H), 7.52-7.41 (m, 4H), 5.49 (s, 1H), 4.91 (s, 1H), 4.63-4.60 (m, 1H), 4.27 (br.s, 1H), 4.04 (b.s, 1H), 3.67 (s, 1.7H), 3.21 (s, 1.3H), 2.56-2.55 (m, 1H), 2.23 (br.s, 1H), 1.90 (br.s, 1H), 1.71 (br.s, 1H); LCMS: 417.1
138		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36-8.30 (m, 2H), 7.79-7.74 (m, 1H), 7.42-7.38 (m, 1H), 5.44 (s, 1H), 4.96 (s, 1H), 4.41-4.34 (m, 1H), 4.15-3.97 (m, 2H), 3.97 (s, 1H), 3.25 (s, 2H), 2.47-2.42 (m, 1H), 2.17-2.15 (m, 1H), 1.82-1.74 (m, 2H); LCMS: 418.0
139		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.25 (d, 1H, J = 5.2 Hz), 7.23-7.14 (m, 3H), 5.51 (s, 1H), 4.94 (s, 1H), 4.51-4.47 (m, 1H), 4.15-4.13 (m, 1H), 3.94-3.89 (m, 1H), 3.89 (s, 1.5H), 3.15 (s, 1.5H), 2.45-2.41 (m, 1H), 2.12-2.04 (m, 1H), 1.80-1.77 (m, 1H), 1.60-1.58 (m, 1H), 1.38-1.29 (m, 1H); LCMS: 419.1
140		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.40 (d, 1H, J = 8.8 Hz), 7.85 (dd, 1H, J = 7.6, 5.6 Hz), 7.31-7.22 (m, 2H), 6.22-6.17 (m, 1H), 6.09-6.04 (m, 1H), 5.61 (s, 1H), 5.05 (s, 1H), 5.02-5.01 (m, 1H), 4.61 (dd, 1H, J = 14.8, 2.8 Hz), 4.12 (td, 1H, J = 18.4, 5.6 Hz), 3.69 (s, 1.5H), 3.21 (s, 1.5H); LCMS: 424.1





141		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.38 (br.s, 1H), 7.68-7.65 (m, 1H), 7.10-7.03 (m, 2H), 5.55-5.46 (m, 1H), 4.94-4.83 (m, 1H), 4.56 (br.s, 1H), 4.21 (br.s, 1H), 4.09-4.02 (m, 1H), 3.58 (s, 1.5H), 3.11 (s, 1.5H), 2.42 (br.s, 1H), 2.10 (br.s, 1H), 1.85 (br.s, 1H), 1.69 (s, 1H); LCMS: 426.1
142		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35 (d, 1H, J = 5.6 Hz), 7.75-7.65 (m, 2H), 7.37-7.34 (m, 1H), 5.53 (s, 1H), 4.95 (s, 1H), 4.38-4.33 (m, 1H), 4.13-4.11 (m, 1H), 4.04-3.96 (m, 1H), 3.68 (s, 1.5H), 3.20 (s, 1.5H), 2.45-2.42 (m, 1H), 2.17-2.13 (m, 1H), 1.80-1.69 (m, 2H); LCMS: 426.1
143		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32 (d, 1H, J = 8.4 Hz), 7.87-7.82 (m, 1H), 7.33-7.23 (m, 2H), 5.66-5.55 (m, 1H), 5.11-5.02 (m, 1H), 4.06-3.95 (m, 3H), 3.68 (s, 1.5H), 3.67-3.58 (m, 1H), 3.52-3.49 (m, 1H), 3.25-3.23 (m, 1H), 3.22 (s, 1.5H), 2.20-2.11 (m, 1H), 1.82-1.74 (m, 1H); LCMS: 426.1
144		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.32-7.25 (m, 3H), 7.24-7.19 (m, 2H), 5.47-5.35 (m, 2H), 4.84-4.83 (m, 1H), 4.53-4.29 (m, 2H), 4.05-3.97 (m, 1H), 3.90-3.87 (m, 1H), 2.89-2.78 (m, 1H), 2.38-2.35 (m, 1H), 2.18-2.06 (m, 2H), 1.77-1.61 (m, 2H); LCMS: 427.1

145		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33 (s, 1H), 7.34-7.19 (m, 5H), 5.48-5.36 (m, 2H), 4.93 (br.s, 1H), 4.53-4.44 (m, 1H), 3.98-3.91 (m, 3H), 3.50-3.46 (m, 2H), 3.23-3.20 (m, 1H), 2.80-2.79 (m, 1H), 2.13-2.07 (m, 2H), 1.77-1.74 (m, 1H); LCMS: 427.2
146		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.17 (d, 1H, J = 8.4 Hz), 7.78-7.72 (m, 1H), 7.21-7.11 (m, 2H), 5.52 (s, 1H), 5.48-4.98 (m, 1H), 4.95 (s, 1H), 4.50-4.44 (m, 1H), 4.10-4.03 (m, 1H), 3.56 (s, 1.5H), 3.10 (s, 1.5H), 2.52-2.41 (m, 2H), 1.88-1.81 (m, 2H); LCMS: 428.1
147		¹ H NMR (400 MHz, DMSO-d ₆) δ 13.92 (br d, 1H), 9.19 (ABqd, 1H), 9.05 (d, J = 7.2 Hz, 1H), 8.20 (d, J = 9.9 Hz, 1H), 7.87 (m, 1H), 7.33 (m, 1H), 4.97 (d, J = 3.8 Hz, 1H), 4.87 (dd, J = 14.6, 5.5 Hz, 1H), 4.51-4.42 (m, 2H), 4.34 (m, 1H), 3.88 (m, 1H), 3.65 (ddt, J = 26.8, 7.3, 5.1 Hz, 1H), 3.40 (d, J = 6.3 Hz, 3H), 3.29 (s, 3H), 2.20 (m, 1H), 1.87 (m, 1H), 1.52 (m, 1H), 1.36-1.15 (m, 1H); LCMS: 432.1
148		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35 (d, 1H, J = 4.4 Hz), 7.42 (d, 1H, J = 2.4 Hz), 7.41-7.32 (m, 1H), 7.18-7.14 (m, 1H), 5.48 (s, 1H), 4.91-4.89 (m, 1H), 4.40-4.36 (m, 1H), 4.15-4.13 (m, 1H), 4.05-3.99 (m, 1H), 3.66 (s, 1.5H), 3.17 (s, 1.5H), 3.47-2.44 (m, 1H), 2.18-2.14 (m, 1H), 1.82-1.80 (m, 1H), 1.71-1.65 (m, 1H); LCMS: 435.1





149		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35 (d, 1H, J = 12.4 Hz), 8.25 (d, 1H, J = 8.8 Hz), 7.64-7.59 (m, 1H), 5.38 (s, 1H), 4.92 (s, 1H), 4.16-3.88 (m, 4H), 3.72 (s, 1H), 3.59-3.39 (m, 2H), 3.26 (s, 2H), 2.13-2.00 (m, 1H), 1.78-1.68 (m, 1H); LCMS: 436.1
150		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (dd, 1H, J = 12.8, 2.0 Hz), 8.25 (dd, 1H, J = 9.2, 2.0 Hz), 7.64-7.59 (m, 1H), 5.38 (s, 1H), 4.02-3.94 (m, 3H), 3.72 (s, 1H), 3.68-3.48 (m, 2H), 3.24 (s, 2H), 3.23-3.22 (m, 1H), 2.19-2.10 (m, 1H), 1.88-1.75 (m, 1H); LCMS: 436.1
151		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.27-8.23 (m, 2H), 7.61-7.57 (m, 1H), 5.32 (s, 1H), 5.24-5.10 (m, 1H), 4.94 (s, 1H), 4.58-4.49 (m, 1H), 4.18-4.08 (m, 1H), 3.67 (s, 1H), 3.20 (s, 2H), 2.65-2.48 (m, 2H), 1.99-1.82 (m, 2H); LCMS: 438.1
152		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.40 (d, 1H, J = 6.0 Hz), 7.88-7.84 (m, 1H), 7.36-7.24 (m, 2H), 5.62 (d, 1H, J = 4.0 Hz), 5.07 (d, 1H, J = 5.6 Hz), 4.61 (dt, 1H, J = 32.0, 5.6 Hz), 4.30 (d, 1H, J = 10.4 Hz), 3.94 (dd, 1H, J = 20.8, 6.0 Hz), 3.72 (s, 1.5H), 3.23 (s, 1.5H), 1.84-1.78 (m, 1H), 1.59-1.53 (m, 1H), 1.32-1.28 (m, 1H), 0.71-0.66 (m, 1H); LCMS: 438.1





153		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33-8.32 (m, 1H), 7.32-7.18 (m, 6H), 5.48-5.35 (m, 2H), 4.92-4.90 (m, 1H), 4.49-4.45 (m, 1H), 4.39-4.36 (m, 1H), 4.24-4.20 (m, 1H), 3.74-3.73 (m, 1H), 2.83-2.75 (m, 1H), 2.24-2.10 (m, 1H), 1.70-1.67 (m, 1H), 1.48-1.46 (m, 1H), 1.28-1.20 (m, 1H), 0.65-0.59 (m, 1H); LCMS: 439.1
154		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.13 (br.s, 1H), 7.81 (br.s, 1H), 7.29 (br.s, 1H), 7.20 (br.s, 1H), 5.61 (br.s, 1H), 5.02-4.99 (m, 1H), 4.42 (br.s, 1H), 4.02-3.97 (m, 1H), 3.62 (s, 2H), 3.14 (s, 1H), 2.06 (s, 1H), 1.82-1.40 (m, 5H); LCMS: 440.3
155		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.31 (s, 1H), 7.32-7.25 (m, 3H), 7.24-7.20 (m, 2H), 5.47-5.35 (m, 2H), 4.53-4.43 (m, 1H), 4.10-3.97 (m, 3H), 3.49-3.46 (m, 1H), 2.81-2.79 (m, 1H), 2.17-2.03 (m, 2H), 1.88-1.83 (m, 2H), 1.59-1.49 (m, 3H); LCMS: 441.1
156		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33 (d, 1H, J = 8.0 Hz), 7.80-7.77 (m, 1H), 7.58-7.50 (m, 2H), 5.61-5.60 (m, 1H), 5.05 (d, 1H, J = 4.8 Hz), 4.42-4.37 (m, 1H), 4.15-4.11 (m, 1H), 4.04-3.97 (m, 1H), 3.69 (s, 1.5H), 3.22 (s, 1.5H), 2.47-2.41 (m, 1H), 2.16-2.13 (m, 1H), 1.81-1.69 (m, 2H); LCMS: 442.1





157		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32-8.24 (m, 1H), 7.35-7.31 (m, 1H), 7.14-6.93 (m, 3H), 6.10-5.96 (m, 2H), 5.47-5.03 (m, 2H), 4.83-4.87 (m, 1H), 4.52-4.38 (m, 2H), 4.07-3.94 (m, 1H), 2.83-2.67 (m, 1H), 2.17-2.01 (m, 1H); LCMS: 443.1
158		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.39-8.32 (m, 1H), 7.39-6.97 (m, 4H), 5.51-5.35 (m, 2H), 4.96-4.91 (m, 1H), 4.49-4.46 (m, 1H), 4.06-3.92 (m, 3H), 3.55-3.49 (m, 2H), 3.26-3.24 (m, 1H), 2.85-2.83 (m, 1H), 2.21-2.11 (m, 2H), 1.78-1.72 (m, 1H); LCMS: 445.2
159		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.40 (s, 0.5H), 8.37 (s, 0.5H), 8.23 (dd, 1H, J = 8.8, 2.8 Hz), 7.64-7.59 (m, 1H), 5.36 (s, 1H), 4.89 (s, 1H), 4.57 (td, 1H, J = 49.2, 4.2 Hz), 4.29 (s, 0.5H), 4.26 (s, 0.5H), 3.90 (dd, 1H, J = 26.4, 6.0 Hz), 3.70 (s, 1.5H), 3.25 (s, 1.5H), 1.82-1.81 (m, 1H), 1.56-1.48 (m, 1H), 1.30-1.27 (m, 1H), 1.68-1.62 (m, 1H); LCMS: 448.1
160		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (d, 1H, J = 4.4 Hz), 7.31-7.29 (m, 1H), 7.13-7.06 (m, 2H), 6.24-6.19 (m, 1H), 4.37-4.35 (m, 1H), 4.12-4.09 (m, 1H), 3.95 (s, 1H), 3.90-3.87 (m, 2H), 3.67-3.65 (m, 1H), 3.47-3.44 (m, 1H), 3.33 (s, 1.5H), 2.86 (s, 1.5H), 1.79 (d, 1.5H, J = 6.8 Hz), 1.66 (d, 1.5H, J = 6.4 Hz); LCMS: 449.1





161		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.25-8.19 (m, 2H), 7.16 (s, 1H), 5.33 (s, 1H), 4.94-4.88 (m, 1H), 4.54-4.46 (m, 1H), 4.03-3.95 (m, 1H), 3.66 (s, 1.5H), 3.60-3.58 (m, 1H), 3.21 (s, 1.5H), 2.20-2.00 (m, 1H), 1.85-1.81 (m, 2H), 1.60-1.35 (m, 3H); LCMS: 450.1
162		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.40-8.37 (m, 1H), 7.49-7.44 (m, 1H), 7.10-7.03 (m, 2H), 5.46 (s, 2H), 4.36-4.35 (m, 1H), 4.11-4.10 (m, 1H), 3.97-3.85 (m, 3H), 3.67-3.65 (m, 2.5H), 3.49-3.40 (m, 1H), 3.22 (s, 1.5H); LCMS: 451
163		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (d, 1H, J = 4.0 Hz), 7.43-7.30 (m, 2H), 7.17-7.12 (m, 1H), 5.45 (s, 1H), 4.88-4.86 (m, 1H), 4.41-4.37 (m, 1H), 4.16-4.11 (m, 1H), 3.97-3.95 (m, 1H), 3.89-3.86 (m, 2H), 3.69-3.65 (m, 1H), 3.64 (s, 1.5H), 3.47-3.45 (m, 1H), 3.16 (s, 1.5H); LCMS: 451.1
164		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.30 (d, 1H, J = 5.6 Hz), 7.11-6.96 (m, 2H), 5.48 (s, 1H), 4.93 (s, 1H), 4.46 (d, 1H, J = 7.6 Hz), 4.12-4.09 (m, 1H), 3.95-3.80 (m, 3H), 3.66-3.64 (m, 2.5H), 3.44-3.41 (m, 1H), 3.18 (s, 1.5H); LCMS: 453.1





165		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33 (d, 1H, J = 6.4 Hz), 7.79-7.76 (m, 1H), 7.60-7.53 (m, 2H), 5.58 (d, 1H, J = 5.2 Hz), 5.03 (d, 1H, J = 5.2 Hz), 4.64-4.57 (m, 1H), 4.36-4.33 (m, 1H), 3.93-3.87 (dd, 1H, J = 17.2, 6.4 Hz), 3.69 (s, 1.5H), 3.02 (s, 1.5H), 1.80-1.76 (m, 1H), 1.55-1.51 (m, 1H), 1.29-1.25 (m, 1H), 0.67-0.62 (m, 1H); LCMS: 454.1
166		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.47 (s, 1H), 8.38-8.23 (m, 2H), 7.70 (d, 1H, J = 9.6 Hz), 5.52-5.38 (m, 2H), 4.93-4.89 (m, 1H), 4.56-4.43 (m, 1H), 4.34-4.32 (m, 1H), 4.11-4.07 (m, 1H), 3.18-3.14 (m, 1H), 2.86-2.82 (m, 1H), 2.56-2.52 (m, 1H), 2.51-2.38 (m, 1H), 2.19-2.16 (m, 2H), 1.94-1.91 (m, 1H); LCMS: 455.1
167		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32 (s, 0.5H), 8.30 (s, 0.5H), 7.80-7.77 (m, 1H), 7.58-7.52 (m, 2H), 5.66-5.55 (m, 1H), 5.10-5.00 (m, 1H), 4.20-4.07 (m, 2H), 3.70 (s, 1.5 H), 3.66-3.59 (m, 1H), 3.22 (s, 1.5H), 2.14-2.07 (m, 1H), 1.90-1.82 (m, 2H), 1.64-1.55 (m, 3H); LCMS: 456.1
168		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.31-8.23 (m, 1H), 7.37-7.33 (m, 1H), 7.15 (d, 1H, J = 8.0 Hz), 7.06 (d, 1H, J = 10.0 Hz), 7.00-6.96 (m, 1H), 5.48-5.35 (m, 2H), 4.54-4.44 (m, 1H), 4.13-4.01 (m, 2H), 3.51-3.48 (m, 1H), 2.84-2.80 (m, 1H), 2.32-2.04 (m, 2H), 1.84-1.79 (m, 2H), 1.59-1.50 (m, 3H); LCMS: 459.2





169		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37-8.29 (m, 1H), 7.44-6.96 (m, 4H), 5.46 (br.s, 1H), 5.40 (d, 1H, J = 52.0 Hz), 4.87-4.82 (m, 1H), 4.47 (dd, 1H, J = 38.4, 14.8 Hz), 4.30 (br.s, 1H), 4.05-3.83 (m, 2H), 3.75-3.58 (m, 2H), 3.55-3.36 (m, 2H), 3.09-2.81 (m, 1H), 2.21-2.08 (m, 1H); LCMS: 461.1
170		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.29 (s, 1H), 7.16-7.02 (m, 3H), 6.08-6.00 (m, 2H), 5.62-5.58 (m, 1H), 5.42 (d, 1H, J = 52.4 Hz), 5.16 (br.s, 1H), 4.66-4.58 (m, 2H), 4.49-4.36 (m, 1H), 4.07-3.94 (m, 1H), 2.88-2.78 (m, 1H), 2.25-2.09 (m, 1H); LCMS: 461.1
171		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35-8.28 (m, 1H), 6.96-6.94 (m, 1H), 6.85-6.78 (m, 2H), 6.23-6.04 (m, 2H), 5.46-5.33 (m, 2H), 5.04 (br.s, 1H), 4.56-4.40 (m, 2H), 4.11-3.97 (m, 1H), 2.86-2.76 (m, 1H), 2.19-2.03 (m, 1H); LCMS: 461.1
172		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32-8.23 (m, 1H), 7.16-7.09 (m, 3H), 6.10-5.98 (m, 2H), 5.66-5.61 (m, 1H), 5.43 (d, 1H, J = 52.0 Hz), 5.06-5.05 (m, 1H), 4.55-4.39 (m, 3H), 3.96-3.93 (m, 1H), 2.87-2.77 (m, 1H), 2.26-2.12 (m, 1H); LCMS: 461.1





173		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 6.95 (d, 2H, J = 6.8 Hz), 6.87-6.67 (m, 1H), 5.46-5.33 (m, 2H), 4.91-4.89 (m, 1H), 4.53-4.50 (m, 1H), 4.34-4.30 (m, 1H), 4.05-4.04 (m, 1H), 3.91-3.88 (m, 1H), 2.84-2.78 (m, 1H), 2.38-2.37 (m, 1H), 2.10-2.06 (m, 2H), 1.74-1.61 (m, 2H); LCMS: 463.1
174		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33-8.26 (m, 1H), 6.98 (d, 1H, J = 6.4 Hz), 6.88-6.70 (m, 2H), 5.48-5.35 (m, 2H), 4.55-4.52 (m, 1H), 4.01-3.90 (m, 3H), 3.55-3.48 (m, 2H), 3.25-3.19 (m, 1H), 2.84-2.78 (m, 1H), 2.18-2.09 (m, 2H), 1.76-1.66 (m, 2H); LCMS: 463.1
175		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32-8.24 (m, 1H), 7.17-6.96 (m, 3H), 5.69-5.65 (m, 1H), 5.46 (d, 1H, J = 52.0 Hz), 4.94-4.89 (m, 1H), 4.57-4.37 (m, 1H), 4.03-3.88 (m, 3H), 3.52-3.48 (m, 2H), 3.24-3.22 (m, 1H), 2.87-2.81 (m, 1H), 2.16-2.10 (m, 2H), 1.74-1.71 (m, 1H); LCMS: 463.1
176		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.24-8.16 (m, 1H), 7.16-6.92 (m, 3H), 5.68-5.64 (m, 1H), 5.42 (d, 1H, J = 52.0 Hz), 4.91-4.89 (m, 1H), 4.48-4.41 (m, 2H), 4.06-4.04 (m, 1H), 3.84-3.81 (m, 1H), 2.86-2.80 (m, 1H), 2.40-2.35 (m, 1H), 2.09-2.01 (m, 2H), 1.74-1.72 (m, 1H), 1.53-1.51 (m, 1H); LCMS: 463.1





177		¹ H NMR (400 MHz, DMSO-d ₆) δ 14.17 (s, 1H), 8.99 (d, J = 7.2 Hz, 1H), 8.38 (d, J = 2.9 Hz, 1H), 8.26 (s, 1H), 7.87 (dd, J = 8.9, 3.0 Hz, 1H), 5.67 – 5.34 (m, 3H), 4.92 (d, J = 3.9 Hz, 1H), 4.80 (dd, J = 21.4, 14.0 Hz, 1H), 4.59 – 4.34 (m, 1H), 4.18 (s, 1H), 3.85 – 3.75 (m, 1H), 2.87 – 2.71 (m, 1H), 2.26 – 2.00 (m, 2H), 1.66 (dq, J = 13.1, 5.5 Hz, 2H), 1.58 – 1.34 (m, 2H); LCMS: 464
178		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37 (s, 1H), 6.97-6.95 (m, 2H), 6.84-6.79 (m, 1H), 5.47-5.41 (m, 1H), 5.10 (d, 1H, J = 53.6 Hz), 4.82 (s, 1H), 4.54-4.41 (m, 2H), 4.14-4.11 (m, 1H), 2.83-2.60 (m, 1H), 2.54-2.47 (m, 2H), 2.07-1.84 (m, 3H); LCMS: 465.1
179		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.42 (br.s, 1H), 7.13-7.03 (m, 2H), 6.97-6.74 (m, 2H), 5.48 (d, 1H, J = 1.6 Hz), 4.91 (s, 1H), 4.47-4.43 (m, 1H), 4.20-4.19 (m, 1H), 4.09-4.03 (m, 1H), 3.69 (s, 1.5H), 3.21 (s, 1.5H), 2.53-2.49 (m, 1H), 2.23-2.20 (m, 1H), 1.87-1.69 (m, 2H); LCMS: 467.1
180		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.22-7.04 (m, 3H), 5.85 (td, 1H, J = 57.2, 4.4 Hz), 5.54-5.51 (m, 1H), 4.25-4.16 (m, 2H), 3.90-3.83 (m, 2H), 3.35-3.33 (m, 1H), 2.31-2.19 (m, 2H), 1.92-1.85 (m, 1H), 1.69-1.66 (m, 1H); LCMS: 469.1





181		¹ H NMR (400 MHz, DMSO-d ₆) δ 14.18 (s, 1H), 9.10 (d, J = 7.3 Hz, 1H), 8.25 (s, 1H), 7.78 (s, 1H), 7.74 – 7.59 (m, 2H), 5.57 – 5.35 (m, 2H), 5.06 (d, J = 5.4 Hz, 1H), 4.76 – 4.63 (m, 1H), 4.50 (dd, J = 39.8, 14.4 Hz, 1H), 4.09 (s, 2H), 3.69 (dd, J = 11.3, 4.5 Hz, 2H), 3.42 – 3.37 (m, 1H), 3.09 (dd, J = 11.4, 8.7 Hz, 1H), 2.73 (dd, J = 33.7, 20.2 Hz, 1H), 2.09 (q, J = 13.8, 11.9 Hz, 2H), 1.50 – 1.33 (m, 1H); LCMS: 470
182		¹ H NMR (400 MHz, DMSO-d ₆) δ 14.17 (s, 1H), 9.11 (d, J = 7.8 Hz, 1H), 8.26 (s, 1H), 7.88 – 7.55 (m, 3H), 5.52 (s, 1H), 5.45 – 5.35 (m, 1H), 5.02 (d, J = 4.7 Hz, 1H), 4.73 (dd, J = 21.9, 13.9 Hz, 1H), 4.48 (dd, J = 39.7, 13.0 Hz, 1H), 4.03 (dd, J = 7.5, 3.7 Hz, 1H), 3.92 (dd, J = 11.1, 3.7 Hz, 1H), 3.73 (dd, J = 11.0, 5.7 Hz, 1H), 3.56 – 3.46 (m, 1H), 3.43 – 3.34 (m, 1H), 3.18 (dd, J = 11.3, 6.5 Hz, 1H), 2.83 – 2.63 (m, 1H), 2.19 – 1.95 (m, 1H), 1.81 – 1.65 (m, 1H), 1.42 (d, J = 4.4 Hz, 1H); LCMS: 470
183		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33-8.26 (m, 1H), 7.84-7.81 (m, 1H), 7.36 (dd, 1H, J = 9.2, 2.0 Hz), 7.24-7.19 (m, 1H), 5.66 (dd, 1H, J = 10.4, 8.0 Hz), 5.49 (d, 1H, J = 52.0 Hz), 4.95-4.89 (m, 1H), 4.65-4.56 (m, 1H), 3.97-3.89 (m, 3H), 3.51-3.48 (m, 2H), 3.25-3.22 (m, 1H), 2.90-2.86 (m, 1H), 2.28-2.11 (m, 2H), 1.74-1.70 (m, 1H); LCMS: 470.1
184		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33-8.26 (m, 1H), 7.13-7.07 (m, 3H), 5.60-5.55 (m, 1H), 5.43 (d, 1H, J = 52.0 Hz), 4.47-4.33 (m, 2H), 4.13-4.09 (m, 1H), 3.20-3.14 (m, 2H), 2.83-2.78 (m, 1H), 2.53-2.51 (m, 1H), 2.49-2.38 (m, 1H), 2.21-2.15 (m, 2H), 1.92-1.89 (m, 1H); LCMS: 472.1





185		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.27 (m, 1H), 6.97 (d, 2H, J = 6.0 Hz), 6.85-6.80 (m, 1H), 5.48-5.34 (m, 2H), 4.92-4.86 (m, 1H), 4.72-4.61 (m, 1H), 4.42-4.36 (m, 2H), 4.13-4.12 (m, 1H), 3.17-3.13 (m, 1H), 2.85-2.80 (m, 1H), 2.54-2.51 (m, 1H), 2.44-2.40 (m, 1H), 2.21-2.16 (m, 1H), 1.94-1.90 (m, 1H); LCMS: 472.1
186		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.24 (s, 1H), 7.82-7.79 (m, 1H), 7.34-7.31 (m, 1H), 7.21-7.03 (m, 1H), 5.65-5.61 (m, 1H), 5.45 (d, 1H, J = 52.0 Hz), 5.10 (d, 1H, J = 53.6 Hz), 4.89 (s, 1H), 4.61-4.47 (m, 2H), 4.06-4.01 (m, 1H), 2.87-2.85 (m, 1H), 2.53-2.46 (m, 2H), 2.44-2.41 (m, 1H), 1.92-1.84 (m, 2H); LCMS: 472.1
187		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.25 (s, 1H), 7.83-7.80 (m, 1H), 7.34-7.31 (m, 1H), 7.21-7.13 (m, 1H), 5.65-5.61 (m, 1H), 5.45 (d, 1H, J = 51.6 Hz), 5.08 (d, 1H, J = 54.0 Hz), 4.78 (s, 1H), 4.61-4.47 (m, 2H), 4.15-4.07 (m, 1H), 3.06-2.85 (m, 1H), 2.55-2.42 (m, 2H), 2.25-2.00 (m, 1H), 1.94-1.82 (m, 2H); LCMS: 472.1
188		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.27 (m, 1H), 6.98 (d, 1H, J = 6.4 Hz), 6.85-6.69 (m, 2H), 5.48-5.35 (m, 2H), 4.64-4.42 (m, 3H), 4.29-4.25 (m, 1H), 3.78 (d, 1H, J = 5.6 Hz), 2.82 (br.s, 1H), 2.19-2.08 (m, 1H), 1.70-1.64 (m, 1H), 1.55-1.47 (m, 1H), 1.23 (s, 1H), 0.66-0.63 (m, 1H); LCMS: 475.1





189		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ ppm 8.31-8.23 (m, 1H), 7.18-6.95 (m, 3H), 5.66-5.62 (m, 1H), 5.45 (d, 1H, $J = 52.0$ Hz), 4.97-4.95 (m, 1H), 4.46-4.28 (m, 3H), 3.89-3.72 (m, 1H), 2.88-2.79 (m, 1H), 2.30-2.16 (m, 1H), 1.70-1.67 (m, 1H), 1.49-1.46 (m, 1H), 1.27-1.20 (m, 1H), 0.65-0.58 (m, 1H); LCMS: 475.1
190		$^1\text{H NMR}$ (400 MHz, DMSO-d_6) δ 14.16 (s, 1H), 9.15 (d, $J = 7.4$ Hz, 1H), 8.25 (s, 1H), 8.05 (d, $J = 2.9$ Hz, 1H), 7.54 (dd, $J = 8.5$, 3.1 Hz, 1H), 5.56-5.32 (m, 2H), 5.07 (d, $J = 5.4$ Hz, 1H), 4.79-4.62 (m, 1H), 4.53-4.27 (m, 1H), 4.09 (m, 1H), 3.91 (s, 3H), 3.71 (m, 2H), 3.48-3.32 (m, 2H), 3.11 (m, 1H), 2.84-2.61 (m, 1H), 2.23-1.97 (m, 2H), 1.53-1.32 (m, 1H); LCMS: 476.1
191		$^1\text{H NMR}$ (400 MHz, DMSO-d_6) δ 14.11 (s, 1H), 8.90 (d, $J = 7.4$ Hz, 1H), 8.23 (s, 1H), 7.34-7.07 (m, 3H), 5.59-5.43 (m, 2H), 4.76-4.58 (m, 3H), 4.50-4.29 (m, 1H), 3.91-3.76 (m, 1H), 3.58-3.39 (m, 1H), 2.82-2.63 (m, 1H), 2.23-1.91 (m, 3H), 1.74 (d, $J = 12.7$ Hz, 1H), 1.32-1.05 (m, 2H); LCMS: 477.1
192		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ ppm 8.35-8.28 (m, 1H), 7.20 (s, 1H), 7.07-6.93 (m, 2H), 6.22-5.98 (m, 2H), 5.46-5.33 (m, 2H), 5.04-5.03 (m, 1H), 4.64-4.43 (m, 2H), 4.00-3.97 (m, 1H), 2.85-2.76 (m, 1H), 2.19-2.03 (m, 1H); LCMS: 477.1





193		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.30-8.22 (m, 1H), 7.17-6.95 (m, 3H), 5.67-5.63 (m, 1H), 5.44 (d, 1H, J = 52.0 Hz), 4.93-4.87 (m, 1H), 4.50-4.37 (m, 1H), 4.11-4.08 (m, 1H), 3.99 (s, 1H), 3.49-3.46 (m, 1H), 2.83-2.82 (m, 1H), 2.24-2.03 (m, 2H), 1.86-1.78 (m, 2H), 1.59-1.47 (m, 3H); LCMS: 477.1
194		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32-8.25 (m, 1H), 8.21-8.09 (m, 1H), 7.61-7.48 (m, 1H), 6.10-6.08 (m, 1H), 5.98-5.96 (m, 1H), 5.65-5.61 (m, 1H), 5.48-5.35 (m, 1H), 5.04-5.03 (m, 1H), 4.55-4.51 (m, 2H), 4.08-3.92 (m, 1H), 2.96-2.87 (m, 1H), 2.19-2.04 (m, 1H); LCMS: 478.1
195		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 9.28 (br.s, 1H), 8.30-8.21 (m, 1H), 7.51 (br.s, 1H), 7.18-7.00 (m, 2H), 5.58-5.39 (m, 2H), 4.65-4.31 (m, 3H), 3.84-3.62 (m, 2H), 2.65 (br.s, 4H), 2.19-1.28 (m, 2H); LCMS: 479.1
196		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36-8.28 (m, 1H), 7.46-7.42 (m, 1H), 7.09-6.91 (m, 2H), 5.75 (t, 1H, J = 8.8 Hz), 5.50-5.33 (m, 1H), 4.61-4.53 (m, 1H), 4.03-3.89 (m, 4H), 3.52-3.47 (m, 2H), 3.24-3.19 (m, 1H), 2.93-2.91 (m, 1H), 2.17-2.10 (m, 2H), 1.75-1.71 (m, 1H); LCMS: 479.1





197		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.35-8.28 (m, 1H), 6.97 (d, 1H, J = 6.4 Hz), 6.87-6.80 (m, 2H), 5.48-5.34 (m, 2H), 4.54-4.33 (m, 1H), 4.32-4.07 (m, 1H), 4.05-4.03 (m, 2H), 3.85-3.84 (m, 2H), 3.77-3.75 (m, 1H), 3.59-3.57 (m, 1H), 3.41-3.36 (m, 1H), 2.86-2.79 (m, 1H), 2.20-2.04 (m, 1H); LCMS: 479.1
198		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.22 (s, 1H), 7.08 (d, 2H, J = 8.8 Hz), 5.47-5.34 (m, 2H), 4.86-4.82 (m, 1H), 4.55-4.50 (m, 1H), 3.97-3.90 (m, 3H), 3.54-3.46 (m, 2H), 3.23-3.21 (m, 1H), 2.83-2.81 (m, 1H), 2.17-2.07 (m, 2H), 1.76-1.73 (m, 1H); LCMS: 479.1
199		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.30-8.23 (m, 1H), 7.07-7.03 (m, 1H), 6.92-6.91 (m, 1H), 6.13-6.06 (m, 1H), 6.04-5.97 (m, 1H), 5.62-5.58 (m, 1H), 5.41 (d, 1H, J = 51.6 Hz), 5.08-5.07 (m, 1H), 4.55-4.35 (m, 2H), 4.05-3.93 (m, 1H), 2.86-2.76 (m, 1H), 2.24-2.08 (m, 1H); LCMS: 479.1
200		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.30-8.23 (m, 1H), 7.35-7.26 (m, 3H), 5.61-5.57 (m, 1H), 5.40 (d, 1H, J = 52.4 Hz), 4.52-4.36 (m, 1H), 4.03-3.87 (m, 4H), 3.53-3.47 (m, 2H), 3.24-3.21 (m, 1H), 2.80-2.80 (m, 1H), 2.24-2.10 (m, 2H), 1.69-1.67 (m, 1H); LCMS: 479.1





201		¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.37-8.29 (m, 1H), 7.35-7.10 (m, 3H), 5.60 (dd, 1H, J = 10.0, 8.4 Hz), 5.45 (d, 1H, J = 52.0 Hz), 4.87-4.85 (m, 1H), 4.55-4.32 (m, 2H), 4.07-3.91 (m, 2H), 2.88-2.82 (m, 1H), 2.41-2.11 (m, 3H), 1.79-1.63 (m, 2H); LCMS: 479.1
202		¹ H NMR (400 MHz, DMSO-d ₆) δ 14.20 (s, 1H), 9.07 (d, J = 7.4 Hz, 1H), 8.39 (d, J = 3.0 Hz, 1H), 8.25 (d, J = 6.0 Hz, 1H), 7.84 (dd, J = 8.8, 3.0 Hz, 1H), 5.60-5.42 (m, 2H), 5.05 (d, J = 5.3 Hz, 1H), 4.75 (dd, J = 21.3, 14.1 Hz, 1H), 4.51 (dd, J = 39.5, 15.0 Hz, 1H), 4.09 (s, 2H), 3.71-3.62 (m, 2H), 3.13-3.02 (m, 1H), 2.08 (s, 2H), 1.44 (dd, J = 27.0, 9.2 Hz, 2H), 1.21 (d, J = 13.5 Hz, 1H); LCMS: 480
203		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.49 (br.s, 1H), 8.38 (br.s, 1H), 8.33 (br.s, 1H), 7.69 (d, 1H, J = 9.2 Hz), 5.97-5.77 (m, 1H), 5.64-5.39 (m, 2H), 4.57-4.44 (m, 1H), 4.24-4.21 (m, 2H), 4.07 (d, 1H, J = 6.4 Hz), 2.89-2.85 (m, 1H), 2.46-2.44 (m, 1H), 2.27-2.25 (m, 1H), 2.15-2.12 (m, 1H), 1.92-1.81 (m, 2H), 1.66-1.62 (m, 1H); LCMS: 480.1
204		¹ H NMR (400 MHz, DMSO-d ₆) δ 14.19 (s, 1H), 9.13 (d, J = 7.8 Hz, 1H), 8.26 (s, 1H), 7.47-7.34 (m, 1H), 7.14 (d, J = 8.1 Hz, 1H), 5.59-5.48 (m, 2H), 4.80-4.60 (m, 1H), 4.41 (dd, J = 39.5, 14.0 Hz, 1H), 4.09-3.96 (m, 1H), 3.92 (dd, J = 11.1, 3.7 Hz, 1H), 3.75 (dd, J = 12.0, 5.6 Hz, 1H), 3.47 (dtd, J = 59.1, 8.2, 7.8, 4.5 Hz, 2H), 3.19 (dd, J = 11.2, 6.9 Hz, 1H), 2.85-2.60 (m, 1H), 2.33-2.04 (m, 1H), 1.79 (dd, J = 11.3, 4.7 Hz, 1H), 1.53-1.33 (m, 1H); LCMS: 481





205		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.37-9.30 (m, 1H), 7.08-6.98 (m, 2H), 5.68-5.63 (m, 1H), 5.53-5.36 (m, 1H), 4.53-4.40 (m, 1H), 4.02-3.91 (m, 4H), 3.55-3.46 (m, 2H), 3.26-3.21 (m, 1H), 2.86-2.82 (m, 1H), 2.30-2.12 (m, 2H), 1.78-1.70 (m, 1H); LCMS: 481.1
206		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (s, 1H), 7.10-6.95 (m, 2H), 5.72-5.62 (m, 1H), 5.52-5.39 (m, 1H), 4.93-4.92 (m, 1H), 4.52-4.08 (m, 2H), 4.07-5.06 (m, 1H), 3.93-3.90 (m, 1H), 2.88-2.82 (m, 1H), 2.41-2.10 (m, 3H), 1.78-1.63 (m, 2H); LCMS: 481.1
207		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.25-8.20 (m, 2H), 7.61-7.58 (m, 1H), 5.66-5.62 (m, 1H), 5.25 (d, 1H, J = 52.0 Hz), 5.14-5.00 (m, 1H), 4.91-4.87 (m, 1H), 4.53-4.44 (m, 2H), 4.06-4.03 (m, 1H), 2.94-2.91 (m, 1H), 2.54-2.43 (m, 2H), 1.91-1.83 (m, 3H); LCMS: 482.1
208		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26-8.20 (m, 1H), 7.08-7.02 (m, 1H), 6.94-6.92 (m, 1H), 5.64-5.60 (m, 1H), 5.42 (d, 1H, J = 52.0 Hz), 5.10 (dt, 1H, J = 52.0, 6.0 Hz), 4.80 (s, 1H), 4.55-4.44 (m, 2H), 4.09-4.04 (m, 1H), 2.92-2.54 (m, 1H), 2.52-2.43 (m, 2H), 1.93-1.89 (m, 1H), 1.86-1.83 (m, 2H); LCMS: 483.1





209		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.38-8.35 (m, 1H), 7.84-7.80 (m, 1H), 7.24-7.15 (m, 2H), 5.58 (s, 1H), 5.07 (s, 1H), 4.40-4.39 (m, 1H), 4.11-4.10 (m, 1H), 3.98-3.90 (m, 3H), 3.68-3.61 (m, 2.5H), 3.48-3.41 (m, 1H), 3.22 (s, 1.5H); LCMS: 485.1
210		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.26 (s, 1H), 7.36-7.32 (m, 1H), 7.15 (d, 1H, J = 7.6 Hz), 7.07 (d, 1H, J = 9.6 Hz), 6.97-6.93 (m, 1H), 5.45-5.41 (m, 2H), 4.49-4.36 (m, 1H), 4.13-4.09 (m, 1H), 4.01-3.97 (m, 1H), 3.13-3.09 (m, 1H), 2.84-2.75 (m, 1H), 2.26-2.16 (m, 3H), 1.68-1.59 (m, 2H), 1.50-1.37 (m, 1H), 1.20-1.14 (m, 6H); LCMS: 487.1
211		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34-8.27 (m, 1H), 7.22 (s, 1H), 7.07 (d, 2H, J = 6.8 Hz), 5.47-5.34 (m, 2H), 4.88-4.87 (m, 1H), 4.79-4.77 (m, 1H), 4.42-4.37 (m, 2H), 4.13-4.12 (m, 1H), 3.17-3.13 (m, 1H), 2.83-2.80 (m, 1H), 2.54-2.51 (m, 1H), 2.44-2.40 (m, 1H), 2.21-2.17 (m, 1H), 1.94-1.91 (m, 1H); LCMS: 488.1
212		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36-8.30 (m, 1H), 7.09-6.95 (m, 2H), 5.65-5.62 (m, 1H), 5.61-5.39 (m, 1H), 4.95-4.91 (m, 1H), 4.50-4.36 (m, 2H), 4.14-4.11 (m, 1H), 3.21-3.16 (m, 1H), 2.82-2.75 (m, 1H), 2.57-2.54 (m, 1H), 2.52-2.42 (m, 1H), 2.23-2.18 (m, 2H), 2.00-1.94 (m, 1H); LCMS: 490.0





213		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 7.49-7.46 (m, 1H), 7.12-7.06 (m, 2H), 5.76 (t, 1H, J = 8.4 Hz), 5.47 (d, 1H, J = 52.4 Hz), 4.46-4.28 (m, 3H), 4.00 (s, 1H), 3.79 (d, 1H, J = 6.4 Hz), 2.20-2.04 (m, 1H), 1.71 (br.s, 1H), 1.50 (br.s, 1H), 1.31-1.25 (m, 2H), 0.67-0.64 (m, 1H); LCMS: 491.1
214		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.33-8.27 (m, 1H), 7.22 (s, 1H), 7.10-7.07 (m, 2H), 5.47-5.34 (m, 2H), 4.51-4.38 (m, 2H), 4.27-4.23 (m, 1H), 3.91-3.77 (m, 1H), 3.35-3.33 (m, 1H), 2.84-2.78 (m, 1H), 2.19-2.18 (m, 1H), 1.72-1.68 (m, 1H), 1.48-1.45 (m, 1H), 1.24-1.21 (m, 1H), 0.67-0.60 (m, 1H); LCMS: 491.1
215		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.41-8.28 (m, 1H), 7.36-6.33 (m, 3H), 5.58-5.54 (m, 1H), 5.51-5.38 (m, 1H), 4.98-4.96 (m, 1H), 4.52-3.76 (m, 4H), 2.97-2.74 (m, 1H), 2.30-2.15 (m, 1H), 1.73-1.48 (m, 2H), 1.26-1.22 (m, 1H), 0.66-0.61 (m, 1H); LCMS: 491.1
216		¹ H NMR (400 MHz, DMSO-d ₆) δ 14.19 (s, 1H), 9.02 (d, J = 7.6 Hz, 1H), 8.39 (d, J = 2.8 Hz, 1H), 8.29 (s, 1H), 7.89 (dd, J = 9.0, 3.0 Hz, 1H), 5.64 - 5.37 (m, 3H), 4.92 - 4.71 (m, 2H), 4.43 (d, J = 9.1 Hz, 1H), 4.30 (dd, J = 13.2, 7.4 Hz, 2H), 4.14 (q, J = 6.0 Hz, 1H), 3.48 (t, J = 5.3 Hz, 1H), 2.79 (td, J = 16.6, 16.1, 7.2 Hz, 1H), 1.43 (dd, J = 8.9, 4.7 Hz, 1H), 1.24 (d, J = 16.9 Hz, 1H), 1.02 (q, J = 4.2 Hz, 1H), 0.48 - 0.33 (m, 1H); LCMS: 492





217		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.39-8.30 (m, 1H), 7.51-7.48 (m, 1H), 7.11-6.95 (m, 2H), 5.79 (t, 1H, J = 9.2 Hz), 5.55-5.38 (m, 1H), 4.89 (br.s, 1H), 4.62-4.49 (m, 1H), 4.17-4.07 (m, 2H), 3.57-3.55 (m, 1H), 3.04-2.91 (m, 1H), 2.17-2.10 (m, 2H), 1.93-1.85 (m, 2H), 1.67-1.54 (m, 3H); LCMS: 493.1
218		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.34-8.28 (m, 1H), 6.95-6.86 (m, 2H), 5.66 (t, 1H, J = 8.4 Hz), 5.50-5.37 (m, 1H), 4.52-4.43 (m, 1H), 4.03-3.91 (m, 7H), 3.54-3.49 (m, 2H), 3.26-3.21 (m, 1H), 2.86-2.75 (m, 1H), 2.24-2.11 (m, 2H), 1.76-1.71 (m, 1H); LCMS: 493.1
219		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.30 (m, 1H), 6.93-6.85 (m, 2H), 5.66 (t, 1H, J = 8.8 Hz), 5.50-5.37 (m, 1H), 4.52-4.31 (m, 2H), 4.07-4.04 (m, 2H), 3.98 (s, 3H), 3.93-3.90 (m, 1H), 2.79-2.75 (m, 1H), 2.43-2.39 (m, 1H), 2.20-2.07 (m, 2H), 1.79-1.77 (m, 1H), 1.64-1.62 (m, 1H); LCMS: 493.1
220		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37 (s, 1H), 7.10-6.98 (m, 2H), 5.63 (t, 1H, J = 8.4 Hz), 5.54-5.36 (m, 1H), 4.98-4.95 (m, 1H), 4.50-4.37 (m, 2H), 4.29-4.25 (m, 1H), 3.80-3.78 (m, 1H), 2.89-2.79 (m, 1H), 2.31-2.15 (m, 1H), 1.78-1.72 (m, 1H), 1.56-1.50 (m, 1H), 1.30-1.25 (m, 1H), 0.68-0.62 (m, 1H); LCMS: 493.1

221		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.30-8.22 (m, 1H), 7.33-6.35 (m, 3H), 5.85-5.56 (m, 1H), 5.49-5.36 (m, 1H), 4.91-4.88 (m, 1H), 4.50-4.40 (m, 1H), 4.15-4.08 (m, 1H), 3.99 (br.s, 1H), 3.52-3.49 (m, 1H), 2.82-2.80 (m, 1H), 2.24-2.03 (m, 2H), 1.86-1.78 (m, 2H), 1.58-1.47 (m, 3H); LCMS: 493.1
222		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.38 (s, 1H), 7.49-7.45 (m, 1H), 7.10-7.03 (m, 2H), 5.77 (d, 1H, J = 8.4 Hz), 5.44 (d, 1H, J = 51.6 Hz), 4.59-4.49 (m, 1H), 4.38-4.36 (m, 1H), 4.09-3.61 (m, 6H), 3.42-3.39 (m, 1H), 2.99-2.90 (m, 0.5H), 2.16-2.01 (m, 0.5H); LCMS: 495.1
223		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.28 (m, 1H), 7.21 (s, 1H), 7.11-7.06 (m, 2H), 5.46-5.33 (m, 2H), 4.84-4.80 (m, 1H), 4.55-4.51 (m, 1H), 4.28-4.27 (m, 1H), 4.05-4.02 (m, 1H), 3.86-3.83 (m, 2H), 3.76-3.73 (m, 1H), 3.58-3.56 (m, 1H), 3.40-3.38 (m, 1H), 2.84-2.78 (m, 1H), 2.19-2.05 (m, 1H); LCMS: 495.1
224		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33-8.32 (m, 1H), 7.09-7.03 (m, 1H), 6.95-6.93 (m, 1H), 5.65-5.61 (m, 1H), 5.51-5.38 (m, 1H), 4.52-4.42 (m, 1H), 4.10-4.02 (m, 3H), 3.51-3.48 (m, 1H), 2.84-2.82 (m, 1H), 2.24-2.04 (m, 2H), 1.87-1.79 (m, 2H), 1.62-1.48 (m, 3H); LCMS: 495.2

225		N/A; LCMS: 497.1
226		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34-8.27 (m, 1H), 7.15-7.02 (m, 3H), 5.99-5.62 (m, 1H), 5.60-5.58 (m, 1H), 5.51-5.38 (m, 1H), 4.96-4.90 (m, 1H), 4.49-4.40 (m, 1H), 4.23-4.18 (m, 1H), 4.12-4.09 (m, 1H), 2.87-2.79 (m, 1H), 2.58-2.50 (m, 1H), 2.30-2.23 (m, 2H), 2.15-2.11 (m, 1H), 1.87-1.77 (m, 1H), 1.67-1.64 (m, 1H); LCMS: 497.1
227		N/A; LCMS: 499.1
228		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.36 (d, 1H, J = 8.4 Hz), 7.40 (br.s, 1H), 7.20-7.14 (m, 2H), 5.50 (s, 1H), 4.94 (d, 1H, J = 7.2 Hz), 4.40-4.38 (m, 1H), 4.11-4.10 (m, 1H), 3.97-3.85 (m, 3H), 3.68-3.61 (m, 2.5H), 3.47-3.40 (m, 1H), 3.17 (s, 1.5H); LCMS: 501.1

229		N/A; LCMS: 504.1
230		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.19 (s, 1H), 7.59 (s, 1H), 7.48-7.41 (m, 2H), 5.47-5.30 (m, 2H), 4.78-4.75 (m, 1H), 4.53-4.43 (m, 1H), 4.34-4.30 (m, 1H), 3.75-3.70 (m, 1H), 2.84-2.79 (m, 1H), 2.18-2.07 (m, 3H), 1.75-1.49 (m, 4H); LCMS: 504.1
231		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.45 (s, 1H), 8.33 (d, 1H, J = 2.8 Hz), 8.21 (s, 1H), 7.64 (d, 1H, J = 9.6 Hz), 5.51-5.34 (m, 2H), 4.82-4.79 (m, 1H), 4.53-4.40 (m, 1H), 4.33 (br.s, 1H), 3.74-3.66 (m, 1H), 2.84-2.80 (m, 1H), 2.21-2.08 (m, 3H), 1.78-1.71 (m, 2H), 1.61-1.46 (m, 2H); LCMS: 504.1
232		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.22-8.15 (m, 1H), 7.13-6.83 (m, 3H), 5.61-5.56 (m, 1H), 5.41 (d, 1H, J = 52.0 Hz), 4.91-4.89 (m, 1H), 4.46-4.36 (m, 1H), 4.21-4.20 (m, 1H), 4.10-3.97 (m, 1H), 2.85-2.79 (m, 1H), 2.29-2.00 (m, 4H), 1.63-1.58 (m, 2H), 1.22-1.18 (m, 6H); LCMS: 505.2

233		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.30-8.24 (m, 1H), 6.94-6.83 (m, 2H), 5.64 (t, 1H, J = 8.8 Hz), 5.41 (d, 1H, J = 52.0 Hz), 4.48-4.29 (m, 4H), 4.00 (s, 3H), 3.74 (d, 1H, J = 6.0 Hz), 2.79-2.71 (m, 1H), 2.20-2.06 (m, 2H), 1.65-1.60 (m, 1H), 1.47-1.45 (m, 1H), 1.23-1.20 (m, 1H), 0.61-0.55 (m, 1H); LCMS: 505.3
234		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.33-8.27 (m, 1H), 6.95-6.84 (m, 2H), 5.65 (t, 1H, J = 8.8 Hz), 5.50-5.31 (m, 1H), 4.52-4.39 (m, 1H), 4.11-4.08 (m, 1H), 4.03 (s, 1H), 3.98 (s, 3H), 3.73-3.50 (m, 2H), 2.84-2.74 (m, 1H), 2.20-2.06 (m, 2H), 1.86-1.81 (m, 2H), 1.61-1.47 (m, 3H); LCMS: 507.1
235		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.37-8.30 (m, 1H), 7.04-6.95 (m, 2H), 6.85-6.67 (m, 2H), 6.15-6.13 (m, 1H), 5.99-5.97 (m, 1H), 5.46-5.33 (m, 2H), 4.98 (d, 1H, J = 4.4 Hz), 4.80 (s, 1H), 4.53-4.42 (m, 2H), 4.00-3.97 (m, 1H), 2.85-2.75 (m, 1H), 2.19-2.02 (m, 1H); LCMS: 509
236		¹H-NMR (400 MHz, CD₃OD) δ ppm 8.36-8.30 (m, 1H), 6.94-6.84 (m, 2H), 5.67 (t, 1H, J = 8.4 Hz), 5.50-5.31 (m, 1H), 4.52-4.42 (m, 1H), 4.31-4.30 (m, 1H), 4.07-3.89 (m, 5H), 3.83-3.76 (m, 3H), 3.58-3.40 (m, 2H), 2.81-2.75 (m, 1H), 2.22-2.03 (m, 1H); LCMS: 509.1

237		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34-8.27 (m, 1H), 7.05-6.96 (m, 2H), 6.86-6.68 (m, 2H), 5.47-5.34 (m, 2H), 4.81 (s, 1H), 4.54-4.49 (m, 1H), 4.31-4.29 (m, 1H), 4.06-4.04 (m, 1H), 3.91-3.88 (m, 1H), 2.86-2.83 (m, 1H), 2.40-2.36 (m, 1H), 2.17-2.06 (m, 2H), 1.76-1.62 (m, 2H); LCMS: 511
238		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.32 (s, 1H), 7.06-6.97 (m, 2H), 6.87-6.69 (m, 2H), 5.47-5.34 (m, 2H), 4.54-4.50 (m, 1H), 4.01-3.88 (m, 3H), 3.53-3.46 (m, 3H), 3.23-3.18 (m, 1H), 2.83-2.81 (m, 1H), 2.17-2.08 (m, 2H), 1.74-1.65 (m, 1H); LCMS: 511.1
239		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34-8.27 (m, 1H), 7.05-6.97 (m, 2H), 6.87-6.68 (m, 2H), 5.47-5.34 (m, 2H), 4.89 (s, 1H), 4.62-4.47 (m, 1H), 4.40-4.37 (m, 1H), 4.27-4.23 (m, 1H), 3.77 (d, 1H, J = 6.0 Hz), 2.86-2.80 (m, 1H), 2.22-2.18 (m, 1H), 1.71-1.68 (m, 1H), 1.48-1.46 (m, 1H), 1.27-1.20 (m, 1H), 0.65-0.59 (m, 1H); LCMS: 523
240		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.23 (s, 1H), 7.36 (br.s, 1H), 7.16-7.08 (m, 2H), 5.65-5.61 (m, 1H), 5.41 (d, 1H, J = 52.0 Hz), 4.80 (s, 1H), 4.57-4.38 (m, 2H), 4.14-4.05 (m, 1H), 3.82-3.79 (m, 1H), 2.80-2.76 (m, 1H), 2.38-2.34 (m, 1H), 2.15-2.01 (m, 2H), 1.74-1.71 (m, 1H), 1.52-1.49 (m, 1H); LCMS: 529.1





241		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.22 (s, 1H), 7.36 (br.s, 1H), 7.17-7.15 (m, 1H), 7.11-7.08 (m, 1H), 5.67-5.62 (m, 1H), 5.41 (d, 1H, J = 52.0 Hz), 4.91 (s, 1H), 4.51-4.38 (m, 1H), 4.20 (br.s, 1H), 3.86-3.83 (m, 2H), 3.54-3.49 (m, 2H), 3.27-3.22 (m, 1H), 2.82-2.73 (m, 1H), 2.19-2.03 (m, 2H), 1.63-1.56 (m, 1H); LCMS: 529.1
242		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.34 (s, 1H), 7.38 (br.s, 1H), 7.18-7.10 (m, 2H), 5.65-5.61 (m, 1H), 5.46 (d, 1H, J = 52.0 Hz), 4.98-4.92 (m, 1H), 4.64-4.53 (m, 1H), 4.49-4.40 (m, 1H), 4.37-4.26 (m, 1H), 3.73 (d, 1H, J = 5.6 Hz), 2.82-2.73 (m, 1H), 2.24-2.07 (m, 1H), 1.67 (br.s, 1H), 1.47 (br.s, 1H), 1.29-1.23 (m, 1H), 0.68-0.59 (m, 1H); LCMS: 541.2
243		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.19 (s, 1H), 7.35-7.34 (m, 1H), 7.16-7.07 (m, 2H), 5.63-5.59 (m, 1H), 5.40 (d, 1H, J = 52.4 Hz), 4.86-4.80 (m, 1H), 4.49-4.41 (m, 2H), 3.81 (br.s, 1H), 3.56-3.54 (m, 1H), 2.81-2.75 (m, 1H), 2.15-1.97 (m, 2H), 1.77-1.71 (m, 2H), 1.52-1.47 (m, 2H), 1.47-1.27 (m, 1H); LCMS: 543.1
244		¹ H-NMR (400 MHz, CD ₃ OD) δ ppm 8.23 (s, 1H), 7.35 (br.s, 1H), 7.16-7.07 (m, 2H), 5.67-5.62 (m, 1H), 5.40 (d, 1H, J = 52.0 Hz), 4.77 (s, 1H), 4.54-4.40 (m, 2H), 4.01-3.86 (m, 2H), 3.79-3.72 (m, 3H), 3.57-3.54 (m, 1H), 2.82-2.72 (m, 1H), 2.22-2.12 (m, 1H); LCMS: 545.1

图1