



(21) (A1) **2,256,647**
(86) 1997/05/22
(87) 1997/12/11

(72) PODSZUN, Wolfgang, DE

(72) PRIESNITZ, Uwe, DE

(72) HÖLTERS, Jürgen, DE

(72) REHBOLD, Bodo, DE

(72) ISRAELS, Rafel, DE

(71) BAYER AKTIENGESELLSCHAFT, DE

(51) Int.Cl.⁶ A01N 25/34, A01N 43/72, A01N 47/40, A01N 25/10

(30) 1996/06/04 (196 22 355.5) DE

(54) **CORPS MOULES LIBÉRANT DES PRINCIPES ACTIFS
AGROCHIMIQUES**

(54) **SHAPED BODIES WHICH RELEASE AGROCHEMICAL
ACTIVE SUBSTANCES**

(57) L'invention concerne des corps moulés libérant des principes actifs agrochimiques servant à traiter des plantes ligneuses, qui s'utilisent dans un creux préformé dans la plante ligneuse; dont le volume remplit jusqu'à 10 à 95 % du volume du creux; dont le volume n'augmente pas pendant la période d'utilisation jusqu'à ce que le creux préformé soit entièrement rempli; qui contiennent a) des principes actifs agrochimiques, b) des polymères non solubles dans l'eau, pouvant subir un traitement thermoplastique, c) des polymères solubles dans l'eau et éventuellement d'autres auxiliaires.

(57) The invention concerns shaped bodies which release agrochemical active substances for treating woody plants. The active substances are introduced into a cavity previously formed in the woody plant, their volume filling the volume of the cavity by between 10 and 95 %. Their volume does not increase during the application time to such an extent that the previously formed cavity is completely filled. The shaped bodies contain: a) agrochemical active substances; b) water-insoluble polymers which can be treated thermoplastically; c) water-soluble polymers; and d) optionally further auxiliary substances.

Le A 31 810-Foreign countries

- 45 -

Shaped articles which release agrochemical active compounds

Abstract

The present invention relates to shaped articles which release agrochemical active compounds, are used for treatment of woody plants and

- are employed in a hollow preformed in the woody plant,
- are of a volume which fills the volume of the hollow to the extent of 10 to 95%,
- are of a volume which does not increase during the period of use to the extent that the preformed hollow space is filled completely,
- comprise
 - a) agrochemical active compounds,
 - b) water-insoluble thermoplastically processable polymers,
 - c) water-soluble polymers and
 - d) if appropriate further auxiliaries.

Le A 31 810-Foreign countries / Rt/Ke/SP

- 1 -

FILED IN THIS AMENDED
TO AMPLIFICATION

Shaped articles which release agrochemical active compounds

The invention relates to shaped articles which release agrochemical active compounds, are used for treatment of woody plants and comprise a water-insoluble thermoplastically processable polymer, a water-soluble polymer and one or more agrochemical active compounds.

The incorporation of agrochemical active compounds into polymers is known, for example from DE-A 4 432 126, WO 91/4732, WO 94/8455, EP-A 524 831, WO 91/3940, EP-A 404 727, EP-A 254 196, JP-A 58/39 602.

Polymer mixtures of a water-containing hydrophilic polymer and water-insoluble thermoplastic polymer are known from EP-A 0 344 118 as a carrier material for active compounds, the shaped articles obtained from this material having an improved dimensional stability in damp air. In water, these materials swell very severely and as a result increase their volume several-fold.

A method for plant treatment in which shaped articles of agrochemical active compounds and a solid carrier are introduced into the sap flow of plants is known from EP-A 564 945.

The present invention relates to shaped articles which release agrochemical active compounds, are used for treatment of woody plants and

- are employed in a hollow preformed in the woody plant,
- are of a volume which fills the volume of the hollow to the extent of 10 to 95%,
- are of a volume which does not increase during the

Le A 31 810-Foreign countries

- 2 -

period of use to the extent that the preformed hollow space is filled completely,

- comprise

a) agrochemical active compounds,

5 b) water-insoluble thermoplastically processable polymers,

c) water-soluble polymers and

d) if appropriate further auxiliaries.

10 The shaped articles according to the invention are suitable for treatment of all types of woody plants, in particular trees and shrubs. Ornamental and crop woody plants are included. Use on trees is particularly advantageous. The trunk circumference of the trees can vary here within a wide range, for example from 6 cm to 200 cm.

15 The size of the hollow form depends on the circumference of the shoot axes (trunk, bough, branch) in which the implant is to be employed. For trunks with a circumference from 20 cm upwards, cylindrical hollow forms with a diameter of 0.2 to 1.0 cm and a length of 2.0 to 4.0 cm are particularly
20 suitable. Several hollow forms are expediently located on thicker trunks. The hollow form can be produced by drilling, stamping, milling or sawing.

The volume of the shaped article fills the hollow form to the extent of 10 to 95%, preferably to the extent of 50 to
25 90%. The shape of the body is not critical for the activity, and it can be rod-shaped, cylindrical or conical. Conical bodies with a base area which corresponds to the diameter of the hollow form can be employed such that the hollow form is closed off tightly. However, it is also possible to
30 establish the tight closure with a closing cap or a closing paste, for example of a commercially available grafting wax.

Agrochemical active compounds which may be mentioned are insecticides, fungicides and herbicides.

Le A 31 810-Foreign countries

- 3 -

Insecticides which may preferably be mentioned are organic phosphorous compounds, such as phosphoric acid esters, carbamates, pyrethroids, urea derivatives, such as benzoyl ureas, triazines, agonists or antagonists of the nicotinic acetylcholine receptors of insects. Juvenile hormones and juvenoid synthetic compounds we, for example, pyriproxyfen, methoprene, hydroprene, are also to be mentioned.

The pyrethroids include:

Allethrin = 2-methyl-4-oxo-3-(2-propenyl)-2-cyclopenten-1-yl
2,2-dimethyl-3-(2-methyl-1-propenyl)-cyclopropane-carboxylate.

Barthrin = (6-chloro-1,3-benzodioxol-5-yl)-methyl
2,2-dimethyl-3-(2-methyl-1-propenyl)-cyclopropane-carboxylate.

Bioresmethrin = [5-(phenyl-methyl)-3-furanyl]-methyl
2,2,3-(2-methyl-1-propenyl)-carboxylate.

Bromethrin = (5-benzyl-3-furyl)methyl 2-(2,2-dibromovinyl)-
3,3-dimethylcyclopropane-carboxylate

Cycloethrin = 3-(2-cyclopenten-1-yl)-2-methyl-4-oxo-2-
cyclopenten-1-yl 2,2-dimethyl-3-(2-methyl-propenyl)-
cyclopropane-carboxylate.

Dimethrin = 2,4-dimethylbenzyl 2,2,3-(2-methyl-1-propenyl)-
carboxylate

Pyresmethrin = 3-[(5-benzyl-3-furyl)-methyl]methyl trans-
(+)-3-carboxy- α ,2,2-trimethylcyclopropane-acrylate.

Resmethrin = (5-benzyl-3-furyl)-methyl 2,2,3-(2-methyl-1-
propenyl)-carboxylate.

Le A 31 810-Foreign countries

- 4 -

- Tetramethrin = (1,3,4,5,6,7-hexahydro-1,3-di-oxo-2H-isoindol-2-yl)-methyl 2,2,3-(2-methyl-1-propenyl)-carboxylate.
- 5 K-othrin = (5-benzyl-3-furyl)-methyl-trans-(+)-3-cyclopentylidene-methyl 2,2-dimethylcyclopropane-carboxylate.
- Permethrin (FMC 33297) (NRDC 143) = m-Phenoxybenzyl-cis-trans-(+)-3-(2,2-dichlorovinyl) 2,2-dimethylcyclopropane-carboxylate.
- 10 Cinerin I = 2-(2-butenyl)-4-hydroxy-3-methyl-2-cyclopenten-1-one 2,2,3-(2-methyl-1-propenyl)-carboxylate.
- Pyrethrin I = 4-hydroxy-3-methyl-2-(2,4-pentadienyl)-2-cyclopenten-1-one 2,2,3-(2-methyl-1-propenyl)-carboxylate.
- 15 Cinerin II = 2-(2-butenyl)-4-hydroxy-3-methyl-2-cyclopenten-1-one 2,2,3-(2-methyl-1-propenyl)-carboxylate.
- Pyrethrin II = 4-hydroxy-3-methyl-2-(2,4-pentadienyl)-2-cyclopenten-1-one 2,2,3-(2-methyl-1-propenyl)-carboxylate.
- Jasmolin I = 4',5'-dihydropyrethrin I.
- Jasmolin II = 4',5'-dihydropyrethrin II.
- 20 Bioethanometrin = (5-benzyl-3-furyl)methyl 2,2-dimethyl-3-(2-cyclopentylvinyl)-cyclopropane-carboxylate.
- Bioethanomethrin = (3-diphenyl ether)methyl 2-(2,2-dichlorovinyl)-3,3-dimethyl-cyclopropane-carboxylate.
- 25 Cypermethrin = (3-diphenyl ether)-cycnomethyl 2-(2,2-dichlorovinyl)-3,3-dimethyl-cyclopropane-carboxylate.

Le A 31 810-Foreign countries

- 5 -

Decamethrin = (3-diphenyl ether)-cyanomethyl 2-(2,2-dibromovinyl)-3,3-dimethyl-cyclopropane-carboxylate.

ES-56 = 2,3-dihydrofuran 2,2,3-(2-methyl-1-propenyl)-carboxylate.

5 Fenpropanate (S-3206) = (3-diphenyl ether) cyanomethyl 2,2-dimethyl-3,3-dimethyl-cyclopropane-carboxylate.

Fenvalerate (S-5602) = (3-diphenyl ether)-cyanomethyl [(p-chlorophenyl)-(isopropyl)]-acetate.

10 (S-5439) = (3-diphenyl ether)methyl [(p-chlorophenyl)-(isopropyl)]-acetate.

Cis-methrin = 5-benzyl-3-furylmethyl 2,2,3-(2-methyl-1-propenyl)carboxylate.

15 Phenothrin = (3-phenoxybenzyl)methyl 2,2,3-(2-methyl-1-propenyl)carboxylate.

Cyfluthrin = 4-fluor-3-diphenyl ether-cyanomethylol 2-(2,2-dichlorovinyl)-3,3-dimethylcyclopropane-carboxylate.

The carbamates include:

20 Aldicarb = 2-methyl-2-(methylthio)-propanol O-[(methylamino)carbonyl]oxime.

Aldoxycarb = 2-methyl-2-(methylsulfonyl)propanol O-[(methylamino)carbonyl]-oxime.

Aminocarb = 4-dimethylamino-3-methylphenyl methylcarbamate.

25 Bendiocarb = 2,2-dimethyl-benzo-1,3-dioxol-4-yl N-methylcarbamate.

Le A 31 810-Foreign countries

- 6 -

Bufencarb = 3-(1-methylbutyl)phenyl methylcarbamate and 3-(1-ethylpropyl)-phenyl methylcarbamate (3:1).

Butacarb = 3,5-bis-(1,1-dimethylethyl)phenyl methylcarbamate.

5 Butocarboxim = 3-methylthio-2-butane O-[(methylamino)carbonyl]oxime.

Butoxycarboxim = 3-methylthio-2-butanone O-[(methylamino)carbonyl]oxime.

10 2-sec.-Butylphenyl methylcarbamate = 2-(1-methylpropyl)-phenyl methylcarbamate.

Carbanolate = 2-chloro-4,5-dimethylphenyl methylcarbamate.

Carbaryl = 1-naphthalenyl methylcarbamate.

Carbofuran = 2,3-dihydro-2,2-dimethyl-7-benzofuranyl methylcarbamate.

15 Cartap = S,S'-[2-(dimethylamino)-1,3-propanediyl] carbamothicate.

Decarbofuran = 2,3-dihydro-2-methylbenzofuran-7-yl methylcarbamate.

20 Dimetilan = -[(dimethylamino)-carbonyl]-5-methyl-1H-pyrazol-3-yl dimethylcarbamate.

Dioxocarb = 2-(1,3-dioxolan-2-yl)-phenyl methylcarbamate.

Ethiofencarb = 2-ethylthiomethylphenyl methylcarbamate.

Fenethacarb = 3,5-diethylphenyl methylcarbamate.

Le A 31 810-Foreign countries

- 7 -

Formetanate = 3-dimethylaminoethyleneaminophenyl methylcarbamate.

Formparanate = 3-Methyl-4-dimethylamino-methyleneamino-phenyl methylcarbamate.

5 Isoprocarb = 2-Isopropylphenyl methylcarbamate.

Methiocarb = 3,5-dimethyl-4-methylthiophenyl methylcarbamate.

Methomyl = Methyl N-[[(methylamino) carbonyl] oxy]-ethane-imidothioate.

10 Mexacarbate = 4-dimethylamino-3,5-dimethylphenyl methylcarbamate.

Nabam = disodium 1,2-ethanediyibis(carbamodithioate)

Nitrilacarb = to Cl₂, (4,4-dimethyl-5-methylamino-carbonyloximino)pentanenitrile.

15 Oxamil = methyl 2-(dimethylamino)-N-[[(methylamino)-carbonyl]oxy]-2-oxoethaneiminiothioate.

Pirimicarb = 2-(dimethylamino)-5,6-dimethyl-4-pyrimidinyl dimethylcarbamate.

20 Promecarb = 3-methyl-5-(1-methylethyl)phenyl methylcarbamate.

Propoxur = 2-(1-methylethoxy)phenyl methylcarbamate.

Thiofanox = 3,3-dimethyl-(methylthio)-2-butanone O-[(methylamino) carbonyl]-oxime.

Thiocarboxime = 1-(2-cyanoethylthio)-ethyleneaminoethyl

Le A 31 810-Foreign countries

- 8 -

carbamate.

Thiram = Tetramethylthoperoxy-dicarbonic acid diamides.

Trimethylphenyl methylcarbamate = 3,4,5-trimethylphenyl methylcarbamate.

5 3,4-Xylyl methylcarbamate = 3,4-dimethylphenyl methylcarbamate.

3,5-Xylyl methylcarbamate = 3,5-dimethylphenyl methylcarbamate.

The organophosphorus compounds include:

10 Acephate = O,S-dimethyl acetylphosphoroaminothioate.

Amidithion = S-(N-2-methoxyethylcarbamoylmethyl) dimethyl phosphorodithioate.

Amitone = S-[2-(diethylamino)ethyl] diethyl phosphorothioate.

15 Athidation = O,O-Diethyl S-5-methoxy-2-oxo-1,3,4-thiadiazol-3-yl-methyl phosphorodithioate.

Azinphos-ethyl = O,O-Diethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate.

20 Azinphos-methyl = O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate.

Azothioate = O,O-Dimethyl O-[p-(p-chlorophenylazo)-phenyl] phosphorothioate.

Bromophos = -(4-bromo-2,5-dichlorophenyl) O,O-dimethyl phosphorothioate.

Le A 31 810-Foreign countries

- 9 -

Bromophos-ethyl = -(4-bromo-2,5-dichlorophenyl) O,O-diethyl phosphorothioate.

Butonate = O,O-Dimethyl 1-butyryl-1-butyryloxy.

5 Carbophenothion = S-[[(4-Chlorophenyl)thio]methyl] O,O-diethyl phosphorodithioate.

Chlorfenvinphos = Diethyl 2-chloro-1-(2,4-dichlorophenyl)-ethenyl-phosphate.

Chlormephos = S-Chloromethyl O,O-diethyl phosphorodithioate.

10 Chlorphoxim = 7-(2-Chlorophenyl)-4-ethoxy-3,5-dioxa-6-aza-4-phosphaoct-6-en-8-nitrile-4-sulphur.

Chlorprazophos = O,O-Diethyl O-3-chloro-7-methyl-pyrazolo[1,5-a]pyrimidin-2-yl phosphorothioate.

Chlorpyrifos = O,O-Diethyl 0,3,5,6,-trichloro-2-pyridyl phosphorothioate.

15 Chlorpyrifos-methyl = O,O-Dimethyl 0,3,5,6-trichloro-2-pyridyl phosphorothioate.

Chlorthiophos = O-2,5-Dichloro-4-(methylthio)-phenyl O,O-diethyl phosphorothioate.

20 Coumaphos = O-3-Chloro-4-methylcoumarin-7-yl O,O-diethyl phosphorothioate.

Coumithoat = O,O-Diethyl O-(7,8,9,10-tetrahydro-6-oxo-6H-dibenzo[b,d]pyran-3-yl) phosphorothioate.

Cortoxyphos = 1-Phenylethyl (E)-3-[(dimethoxyphosphonyl)oxy]-2-butenate.

Le A 31 810-Foreign countries

- 10 -

Cruformate = 2-Chlor-4-(1,1-dimethylethyl)phenyl methyl methylphosphoramidate.

Cyanofenphos = O-4-Cyanophenyl O-ethyl phenylphosphonothioate.

5 Cyanophos = O-4-Cyanophenyl O,O-dimethyl phosphorothioate.

Cyanthoate = O,O-Diethyl S-[N-(1-cyano-1-methylethyl)]carbamoylmethyl phosphorothioate.

Demephion = O,O-Dimethyl O-2-methylthioethyl phosphorothioate and of OkO-dimethyl S-2-methylthioethyl phosphorothioate.
10

Demeton = O,O-Diethyl O-2-ethylthioethyl phosphorothioate and O,O-diethyl S-2-ethylthioethyl phosphorothioate.

Demeton-S-methyl = O,O-Dimethyl S-2-ethylthioethyl phosphorothioate.

15 Demeton-S-methyl-sulfon = S-2-ethylsulfonyl ethyl O,O-dimethyl phosphorothioate.

Demeton-S = O,O-Diethyl S-[2-(ethylthio)ethyl] phosphorothioate.

Demeton-O = O,O-Diethyl-O-[2-(ethylthio)ethyl] phosphorothioate.
20

Demeton-O-methyl = O,O-Dimethyl O-[2-(ethylthio)ethyl] phosphorothioate.

Dialifos = S-[2-Chloro-1-(1,3-dihydro-1,3-dioxy-2H-isoindol-2-yl)ethyl] O,O-diethyl phosphorodithioate.

25 Diazinon = O,O-Diethyl O-[6-methyl-2-(1-methylethyl)-4-

Le A 31 810-Foreign countries

- 11 -

pyrimidinyl] phosphorothioate.

Dichlorfenthion = O,O-Diethyl O-(2,4-dichlorophenyl) phosphorothioate.

O-2,4-Dichlorophenyl O-ethylphenyl phosphonothioate.

5 Dichlorvos = Dimethyl 2,2-dichloroethenyl phosphate.

Dicrotophos = Dimethyl 3-(dimethylamino)-1-methyl-3-oxo-1-propenyl phosphate.

Dimefox = Bis(dimethylamino)fluorophosphine oxide.

Dimefox = Bis(dimethylamino)fluorophosphine oxide.

10 Dimethoate = O,O-Dimethyl S-[2-(methylamino)-2-oxo-ethyl] phosphorodithioate.

1,3-Di-(methoxycarbonyl)-1-propen-2-yl dimethyl phosphate = dimethyl 3-[(dimethoxyphosphinyl)oxy]-2-pentenedioate.

15 Dioxathion = S,S'-1,4-dioxan-2,3-diyl O,O',O'-tetraethyl di-(phosphorodithioate).

Disulfoton = O,O-Diethyl S-2-ethylthioethyl phosphorodithioate.

EPN = O-Ethyl O-4-nitrophenyl phenylphosphonothioate.

20 Endothion = O,O-Dimethyl S-(5-methoxy-4-pyron-2-yl-methyl) phosphorothioate.

Ethion = O,O,O'',O''-Tetraethyl S,S'-methylene di(phosphorodithioate).

S-Ethylsulfinylmethyl O,O-diisopropyl phosphorodithioate.

Le A 31 810-Foreign countries

- 12 -

Ethoat-methyl = O,O-Dimethyl S-(N-ethylcarbamoyl-methyl) phosphorodithioate.

Ethoprophos = O-Ethyl S,S-dipropyl phosphorodithioate.

5 Etrimfos = O-(6-Ethoxy-2-ethyl-4-pyrimidinyl) O,O-dimethylphosphorothioate.

Famphur = O,O-Dimethyl O-p-(dimethylsulfamoyl)-phenyl phosphorothioate.

Fenchlorphos = O,O-Dimethyl O-(2,4,5-trichlorophenyl) phosphorothioate.

10 Fensulfothion = O,O-Diethyl O-4-(methylsulfinyl)phenyl phosphorothioate.

Fenthion = O,O-Dimethyl O-[3-methyl-4-(methylthio)-phenyl] phosphorothioate.

Fonophos = O-Ethyl S-phenyl ethylphosphonodithioate.

15 Formothion = S-[2-(formylmethylamino)-2-oxoethyl] O,O-dimethyl phosphorodithioate.

Fospirate = Dimethyl 3,5,6-trichloro-2-pyridyl phosphate.

Fosthietan = Diethyl 1,3-dithietan-2-yl-idene phosphoramidate.

20 Heptenophos = 7-Chlorobicyclo[3,2,0]-hepta-2,6-dien-6-yl dimethyl phosphate.

Iodofenphos = O-2,5-Dichloro-4-iodophenyl O,O-dimethyl phosphorothioate.

Isofenphos = 1-Methylethyl 2-[[ethoxy]-(1-

Le A 31 810-Foreign countries

- 13 -

methylethyl)amino[phosphinothioyl]-oxy]benzoate.

Leptophos = O-4-Bromo-2,5-dichlorophenyl O-methylphenyl phosphonothioate.

5 Lythidathion = O,O-Dimethyl S-(5-ethoxy-2,3-dihydro-2-oxo-1,3,4-thiadiazol-3-yl-methyl) phosphotodithioate.

Malathion = Diethyl (dimethoxyphosphinothioyl)thiobutenedioate.

Mazidox = N,N,N',N'-Tetramethylphosphorodiamidic acid.

10 Mecarbam = Methyl ethyl[[(diethoxyphosphinothioyl)thio]acetal]carbamate.

Mecarphon = N-Methylcarbonyl-N-methyl-carbamoyl-methyl O-methyl methylphosphonodithioate.

Menazon = S-[(4,6-Diamino-1,3,5-triazin-2-yl)methyl] O,O-dimethyl phosphorodithioate.

15 Mephosfolan = Diethyl 4-methyl-1,3-dithiolan-2-yl-dinen phosphoroamidate.

Methamidophos = O,S-Dimethyl phosphoramidothioate.

Methidation = S-[[5-Methoxy-2-oxo-1,3,4-thiadiazol-3-(2H)-yl]methyl] O,O-dimethyl phosphorodithioate.

20 Methocrotophos = Dimethyl cis-2-(N-methoxy-N-methyl)-carbamoyl)-1-methyl-vinyl phosphate.

2-Sulfur of 2-methoxy-4H-benzo-1,3,2-dioxaphosphorin.

Methyl-carbophenotion = S-[[4-Chlorophenyl)thio]-methyl] O,O-dimethyl phosphorodithioate.

Le A 31 810-Foreign countries

- 14 -

Mevinphos = Methyl 3-[(dimethoxyphosphinyl)oxy]-2-buenoate.

Monocrotophos = Dimethyl 1-methyl-3-(methylamino)-3-oxo-1-propenyl phosphate.

5 Morphotion = O,O-Dimethyl S-(morpholino-carbonylmethyl) phosphorodithioate.

Naled = Dimethyl 1,2-dibromo-2,2-dichloroethyl phosphate.

Omethoate = O,O-Dimethyl S-[2-(methylamino)-2-oxoethyl] phosphorothioate.

10 Oxydimeton-methyl = S-[2-(Ethylsulfinyl)ethyl] O,O-dimethyl phosphorothioate.

Oxydisulfoton = O,O-Diethyl S-[2-(ethyl-sulfinyl)-ethyl] phosphorodithioate.

Parathion = O,O-Diethyl O-4-nitrophenyl phosphorothioate.

15 Parathion-methyl = O,O-Dimethyl O-4-nitrophenyl phosphorothioate.

Phenkapton = O,O-Diethyl S-(2,5-dichloro-phenylthiomethyl) phosphorodithioate.

Phenthioate = Ethyl α [(dimethoxyphosphinothioyl)thio]benzene-acetate.

20 Phorate = O,O-Diethyl S-ethylthiomethyl phosphorodithioate.

Phosalone = S-[[(6-Chloro-2-oxo-3) (2H)-benzoxazolyl]-(methyl)] O-diethyl phosphorodithioate.

Phosfolan = Diethyl 1,3-dithiolan-2-ylidene phosphoramidate.

Le A 31 810-Foreign countries

- 15 -

Phosmet = S-[(1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)methyl]
O,O-dimethyl phosphorodithioate.

Phosnichlor = O,O-Dimethyl O-4-chloro-3-nitrophenyl
phosphorothioate.

5 Phosphamidon = Dimethyl 2-Chloro-3-(diethylamino)-1-methyl-
3-oxo-1-propenyl phosphate.

Phoxim = -[[Diethoxyphosphinothioyl)oxy]imino)-
benzeneacetonitrile.

10 Pirimiphos-ethyl = O-[2-(Diethylamino)-6-methyl-4-
pyrimidinyl)] O,O-diethyl phosphorothioate.

Pirimiphos-methyl = O-[2-(Diethylamino)-6-methyl-4-
pyrimidinyl)] O,O-dimethyl phosphorothioate.

Profenofos = O-(4-bromo-2-chlorophenyl) O-ethyl S-propyl
phosphorothioate.

15 Propetamphos = (E)-1-Methylethyl 3-[[(ethylamino)-
methoxyphosphinothioyl)oxy]-2-butenate.

Prothidathion = O,O-Diethyl S-(2,3-dihydro-5-isopropyl-2-
oxo-1,3,4-thiadiazol-3-yl-methyl) phosphorodithioate.

20 Prothoate = O,O-Diethyl S-[2-(1-methylethyl)amino-2-
oxoethyl] phosphorodithioate.

Quinalphos = O,O-Diethyl O-2-quinoxalinyll phosphorothioate.

Quinothion = O,O-Diethyl 2-methylquinolin-4-yl
phosphorothioate.

Quintiofos = O-Ethyl O-8-quinolyphenyl phosphorothioate.

Le A 31 810-Foreign countries

- 16 -

Sophamide = O,O-Dimethyl S-(N-methoxy-methyl)-carbamoyl-methyl phosphorodithioate.

Sulfotepp = Tetraethyl thiodiphosphate.

5 Sulfprofos = O-Ethyl O-(4-methylthiophenyl) S-propyl phosphorodithioate.

Temephos = O,O'-(Thiodi-4,1-phenylene) O,O,O',O'-tetramethyl-di(phosphorodithioate)

Tepp = Tetraethyl diphosphates.

10 Terbufos = S-[(1,1-Dimethylethyl)thiomethyl] O,O-diethyl phosphorodithioate.

Tetrachlorvinphos = Dimethyl trans-2-Chloro-1-(2,4,5-trichlorophenyl)vinyl phosphates.

O,O,O',O'-Tetrapropyl dithiopyrophosphate = Tetrapropyl thiodiphosphate.

15 Thiometon = O,O-Dimethyl S-[2-(ethylthio)ethyl] phosphorodithioate.

Thionazin = O,O-Diethyl O-pyrazinyl phosphorothioate.

Triazophos = O,O-Diethyl O-(phenyl-1H-1,2,4-triazol-3-yl) phosphorothioate.

20 Trichloronate = O-Ethyl O-2,4,5-trichlorophenyl ethylphosphonothioate.

Trichlorphon = Dimethyl (1-hydroxy-2,2,2-trichloro-ethyl)-phosphonate.


Vamidotion = O,O-Dimethyl S-[2-(1-methylcarbamoyl)-

Le A 31 810-Foreign countries

- 17 -

ethylenethyl] phosphorothioate.

The benzoylureas include compounds of the formula (V):



wherein


5 R¹ represents halogen,

R² represents hydrogen or halogen,

R³ represents hydrogen, halogen or C₁₋₄-alkyl,

10 R⁴ for halogen, 1-5-halogeno-C₁₋₄-alkyl, C₁₋₄-alkoxy, 1-5-halogeno-C₁₋₄-alkoxy, C₁₋₄-Alkylthio, 1-5-halogeno-C₁₋₄-alkylthio, Phenoxy or pyridyloxy, which can optionally be substituted by halogen, C₁₋₄-alkyl, 1-5-halogeno-C₁₋₄-alkyl, C₁₋₄-alkoxy, 1-5-halogeno-C₁₋₄-alkoxy, C₁₋₄-alkylthio, 1-5-halogeno-C₁₋₄-alkylthio.

15 Benzoylureas which may be mentioned in particular are those of the formula:






Le A 31 810-Foreign countries


- 18 -

5

10

R ¹	R ²	R ⁴
H	Cl	CF ₃
Cl	Cl	CF ₃
F	F	CF ₃
H	F	CF ₃
H	Cl	SCF ₃
F	F	SCF ₃
H	F	SCF ₃
H	Cl	OCF ₃
F	F	OCF ₃
H	F	OCF ₃
F	F	
F	F	
F	F	

15 The triazines include compounds of the formula



Le A 31 810-Foreign countries

- 19 -

	R ₁	R ₂	R ₃
	Cyclopropyl	H	H
	Cyclopropyl	H	CH ₃
	Cyclopropyl	H	C ₂ H ₅
5	Cyclopropyl	H	C ₃ H ₇ -n
	Cyclopropyl	H	C ₄ H ₉ -n
	Cyclopropyl	H	C ₅ H ₁₁ -n
	Cyclopropyl	H	C ₆ H ₁₃ -n
	Cyclopropyl	H	C ₇ H ₁₅ -n
10	Cyclopropyl	H	C ₁₂ -H ₂₅ -n
	Cyclopropyl	H	CH ₂ -C ₄ H ₉ -t
	Cyclopropyl	H	CH ₂ CH(CH ₃)C ₂ H ₅
	Cyclopropyl	H	CH ₂ CH=CH ₂
	Cyclopropyl	Cl	C ₂ H ₅
15	Cyclopropyl	Cl	C ₆ H ₁₃ -n
	Cyclopropyl	Cl	C ₈ H ₁₇ -n
	Cyclopropyl	Cl	C ₁₂ H ₂₅ -n
	Cyclopropyl	H	Cyclopropyl
	Cyclopropyl	H	COCH ₃
20	Cyclopropyl	H	COCH ₃ .HCl
	Cyclopropyl	H	COC ₂ H ₅ .HCl
	Cyclopropyl	H	COC ₂ H ₅
	Cyclopropyl	H	COC ₃ H ₇ -n
	Cyclopropyl	H	COC ₃ H ₇ -i
25	Cyclopropyl	H	COC ₄ H ₉ -t.HCl

Le A 31 810-Foreign countries

- 20 -

	R ₁	R ₂	R ₃
	Cyclopropyl	H	COC ₄ H ₉ -n
	Cyclopropyl	H	COC ₆ H ₁₃ -n
	Cyclopropyl	H	COC ₁₁ -H ₂₃ -n
	Cyclopropyl	COCH ₃	COC ₂ H ₅
5	Cyclopropyl	COC ₃ H ₇ -n	COC ₆ H ₁₃ -n
	Cyclopropyl	COCH ₃	COC ₃ H ₇ -n
	Cyclopropyl	COC ₂ H ₅	COC ₃ H ₇ -n
	Cyclopropyl	H	COCyclopropyl
	Cyclopropyl	COCyclopropyl	COCyclopropyl
10	Cyclopropyl	COCH ₃	COCH ₃
	Isopropyl	H	H
	Isopropyl	H	COCH ₃
	Isopropyl	H	COC ₃ H ₇ -n
	Cyclopropyl	H	CONHCH ₃
15	Cyclopropyl	H	CONHC ₃ H ₇ -i
	Cyclopropyl	CONHCH ₃	CONHCH ₃
	Cyclopropyl	H	CSNHCH ₃
	Cyclopropyl	H	CONHCH ₂ CH=CH ₂
	Cyclopropyl	CONHCH ₂ CH=CH ₂	CONHCH ₂ CH=CH ₂
20	Cyclopropyl	CSNHCH ₃	CSNHCH ₃

The agonists or antagonists of nicotinic acetylcholin receptors of insects include the known compounds from, for example, European Offenlegungsschriften (European Published Specifications) No. 464 830, 428 941, 425 978, 386 565, 25 383 091, 375 907, 364 844, 315 826, 259 738, 254 859,

Le A 31 810-Foreign countries

- 21 -

235 725, 212 600, 192 060, 163 855, 154 178, 136 636,
 303 570, 302 833, 306 696, 189 972, 455 000, 135 956,
 471 372, 302 389; German Offenlegungsschriften (German
 5 Published Specifications) No. 3 639 877, 3 712 307; Japanese
 Offenlegungsschriften (Japanese Published Specifications)
 No. 3 639 877, 3 712 307; Japanese Offenlegungsschriften
 (Japanese Published Specifications) No. 03 220 176,
 02 207 083, 63 307 857, 63 287 764, 03 246 283, 04 9371,
 03 279 359, 03 255 072; US Patent Specifications No.
 10 5 034 524, 4 948 798, 4 918 086, 5 039 686, 5 034 404; PCT
 Applications No. WO 91/17 659, 91/4965; French Application
 No. 2 611 114; Brazilian Application No. 88 03 621.

These compounds can preferably be represented by the general
 formula (I)

15



in which

- R represents hydrogen, optionally substituted radicals
 from the group consisting of acyl, alkyl, aryl,
 aralkyl, heteroaryl or heteroarylalkyl;
- 20 A represents a monofunctional group from the series
 consisting of hydrogen, acyl, alkyl, aryl, or
 represents a bifunctional group which is linked to the
 radical Z;
- E represents an electron-withdrawing radical, such as NO₂


Le A 31 810-Foreign countries

- 22 -

or CN;

X represents the radicals $-\text{CH}=\text{}$ or $=\text{N}-$, wherein the radical $-\text{CH}=\text{}$ can be linked with the radical Z at the position of the H atom;

5 Z represents a monofunctional group from the series consisting of alkyl, $-\text{O}-\text{R}$, $-\text{S}-\text{R}$,



or represents a bifunctional group which is linked to the radical A or the radical X.

10 Particularly preferred compounds of the formula (I) are those in which the radicals have the following meaning:

R represents hydrogen, and represents optionally substituted radicals from the series consisting of acyl, alkyl, aryl, aralkyl, heteroaryl, 15 heteroarylalkyl.

Acyl radicals which may be mentioned are formyl, alkylcarbonyl, arylcarbonyl, alkylsulphonyl, arylsulphonyl, (alkyl)-(aryl)-phosphoryl, which can be substituted in their turn.

20 Alkyls which may be mentioned are C_{1-10} -alkyl, in particular C_{1-4} -alkyl, specifically methyl, ethyl, isopropyl, sec.- or t-butyl, which can be substituted in their turn.

Aryls which may be mentioned are phenyl, naphthyl, in

Le A 31 810-Foreign countries

- 23 -

particular phenyl.

Aralkyls which may be mentioned are phenylmethyl, phenethyl.

5 Heteroaryls which may be mentioned are heteroaryl having up to 10 ring atoms and N, O, S, in particular N, as heteroatoms. Thienyl, furyl, thiazolyl, imidazolyl, pyridyl, benzothiazolyl may be mentioned specifically.

10 Heteroarylalkyls which may be mentioned are heteroarylmethyl, heteroarylethyl having up to 6 ring atoms, and N, O, S, in particular N, as heteroatoms.

Substituents which may be mentioned as examples and as preferred are:

15 Alkyls having preferably 1 to 4, in particular 1 or 2, carbon atoms, such as methyl, ethyl, n- and i-propyl and n-, i- and t-butyl; alkoxy having preferably 1 to 4, in particular 1 or 2, carbon atoms, such as methoxy, ethoxy, n- and i-propyloxy and n-, i- and t-butyloxy; alkylthio having preferably 1 to 4, in particular 1 or 20 2, carbon atoms, such as methylthio, ethylthio, n- and i-propylthio and n-, i- and t-butylthio; halogenoalkyl having preferably 1 to 4, in particular 1 or 2, carbon atoms and preferably 1 to 5, in particular 1 to 3, halogen atoms, it being possible for the halogen atoms to be identical or different, and for the halogen atoms 25 to be, preferably, fluorine, chlorine or bromine, in particular fluorine, such as trifluoromethyl; hydroxyl; halogen, preferably fluorine, chlorine, bromine and iodine, in particular fluorine, chlorine and bromine; 30 cyano; nitro; amino; monoalkyl- and dialkylamino having preferably 1 to 4, in particular 1 or 2, carbon atoms

Le A 31 810-Foreign countries

- 24 -

per alkyl group, such as methylamino, methyl-ethyl-
 amino, n- and i-propylamino and methyl-n-butylamino;
 carboxyl; carbalkoxy having preferably 2 to 4, in
 particular 2 or 3, carbon atoms, such as carbomethoxy
 5 and carboethoxy; sulpho ($-SO_3H$); alkylsulphonyl having
 preferably 1 to 4, in particular 1 or 2, carbon atoms,
 such as methylsulphonyl and ethylsulphonyl;
 arylsulphonyl having preferably 6 or 10 aryl carbon
 atoms, such as phenylsulphonyl, and heteroarylamino and
 10 heteroarylalkylamino, such as chloropyridylamino and
 chloropyridylmethylamino.

A particularly preferably represents hydrogen, and
 represents optionally substituted radicals from the
 series consisting of acyl, alkyl, aryl, which
 15 preferably have the meanings given in the case of R. A
 furthermore represents a bifunctional group. Optionally
 substituted alkylene having 1-4, in particular 1-2, C
 atoms may be mentioned, substituents which may be
 mentioned being the substituents listed above and it
 20 being possible for the alkylene groups to be
 interrupted by heteroatoms from the series consisting
 of N, O, S.

A and Z, together with the atoms to which they are bonded,
 can form a saturated or unsaturated heterocyclic ring.
 25 The heterocyclic ring can contain a further 1 or 2
 identical or different heteroatoms and/or hetero
 groups. Heteroatoms are preferably oxygen, sulphur or
 nitrogen, and hetero groups are N-alkyl, in which alkyl
 of the N-alkyl group preferably contains 1 to 4, in
 30 particular 1 or 2, carbon atoms. Alkyls which may be
 mentioned are methyl, ethyl, n- and i-propyl and n-, i-
 and t-butyl. The heterocyclic ring contains 5 to 7,
 preferably 5 or 6, ring members.

Le A 31 810-Foreign countries

- 25 -



- 5 Examples of the heterocyclic ring which may be mentioned are pyrrolidine, piperidine, piperazine, hexamethyleneimine, hexahydro-1,3,5-triazine, morpholine, which can optionally be substituted, preferably by methyl.
- E represents an electron-withdrawing radical, NO_2 , CN , halogenoalkylcarbonyl, such as 1,5-halogeno- C_{1-4} -carbonyl, in particular COCF_3 , being mentioned in particular.
- 10 X represents $-\text{CH}=\text{}$ or $-\text{N}=\text{}$
- Z represents optionally substituted radicals alkyl, $-\text{OR}$, $-\text{SR}$, $-\text{NRR}$, wherein R and the substituents preferably have the abovementioned meaning.
- 15 Z can form, in addition to the abovementioned ring, together with the atom to which it is bonded and the radical $\begin{array}{c} | \\ =\text{C}- \end{array}$, at the position of X, a saturated or unsaturated heterocyclic ring. The heterocyclic ring can contain a further 1 or 2 identical or different heteroatoms and/or hetero groups. Heteroatoms are preferably oxygen, sulphur or nitrogen, and hetero groups are N-alkyl, wherein the alkyl or N-alkyl group preferably contains 1 to 4, in particular 1 or 2, carbon atoms. Alkyls which may be mentioned are methyl, ethyl, n- and i-propyl and n-, i- and t-butyl. The heterocyclic ring contains 5 to 7, preferably 5 or 6, ring members.
- 20
- 25
- 30 Examples of the heterocyclic ring which may be mentioned are pyrrolidine, piperidine, piperazine, hexamethyleneimine, morpholine and N-methylpiperazine.

Le A 31 810-Foreign countries

- 26 -

As compounds from the group of agonists and antagonists of nicotinic acetylcholine receptors of insects which can especially preferably be used there may be mentioned compounds of the general formulae (II) and (III):

5





in which

n represents 1 or 2,

10 Subst. represents one of the substituents mentioned above in the preferred or particularly preferred meanings, in particular represents halogen, and especially represents chlorine,









A, Z, X and E have the meanings given above in those as preferred or particularly preferred,

15 The following compounds may be mentioned specifically:











Le A 31 810-Foreign countries

- 27 -










Le A 31 810-Foreign countries

- 28 -



Le A 31 810-Foreign countries

- 29 -



- 5 Fungicides which may be mentioned are, preferably:
 Sulphenamides, such as dichlorfluand (Euparen),
 tolylfuand (Methyleuparen), folpet, fluorfolpet;

Le A 31 810-Foreign countries

- 30 -

Benzimidazoles, such as carbendazim (MBC), benomyl, fuberidazole, thiabendazole or salts thereof;

Thiocyanates, such as thiocyanatomethylthiobenzothiazole (TCMTB), methylene bithiocyanate (MBT);

- 5 quaternary ammonium compounds, such as benzyldimethyltetradecylammonium chloride, benzoyl-dimethyl-dodecyl-ammonium chloride, dodecyl-dimethyl-ammonium chloride; morpholine derivatives, such as C₁₁-C₁₄-4-alkyl-2,6-dimethyl-morpholine homologs (tridemorph), (+)-cis-4-[3-
10 tert-butylphenyl)-2-methylpropyl]-2,6-dimethylmorpholine (fenpropimorph), falimorph;

Phenols, such as o-phenylphenol, tribromophenol, tetrachlorophenol, pentachlorophenol, 3-methyl-4-chlorophenol, dichlorophen, chlorophen or salts thereof;

- 15 Azoles, such as tridimefon, triadimenol, bitertanol, tebuconazole, propiconazole, azaconazole, hexaconazole, prochloraz, cyproconazole, 1-(2-chlorophenyl)-2-(1-chlorocyclopropyl)-3-(1,2,4-triazol-1-yl)-propan-2-ol, 1-(2-chlorophenyl)-2-(1,2,4-triazol-1-yl-methyl)-3,3-dimethyl-
20 butan-2-ol.

Iodopropargyl derivatives, such as iodopropargyl butylcarbamate (IPBC), chlorophenylformal, phenylcarbamate, hexylcarbamate, cyclohexylcarbamate, iodopropargyl oxyethylphenylcarbamate;

- 25 Iodine derivatives, such as diiodomethyl p-aryl sulphones, for example diiodomethyl p-tolyl sulphone;

Bromine derivatives, such as bromopol;

Le A 31 810-Foreign countries

- 31 -

Isothiazolines, such as N-methylisothioazolin-3-one, 5-chloro-N-methylisothiazolin-3-one, 4,5-dichloro-N-octylisothiazoline-3-one, N-octylisothiazolin-3-one (octilinone);

- 5 Benzisothiazolinones, cyclopenteneisothiazolines;

Pyridines, such as 1-hydroxy-2-pyridinethione, tetrachloro-4-methylsulphonylpyridine;

- 10 Nitriles, such as 2,4,5,6-tetrachloroisophthalonitrile (chlorthalonil) and other microbicides having an activated halogen group, such as CI-Ac, MCA, tectamer, bromopol, bromidox;

Benzothiazoles, such as 2-mercaptobenzothiazoles; see above dazomet;

Quinolines, such as 8-hydroxyquinoline.

- 15 Insecticides which may be mentioned as particularly preferred are:

- 20 Phosphoric acid esters, such as azinphos-ethyl, azinphos-methyl, 1-(4-chlorophenyl)-4-(O-ethyl, S-propyl)phosphoryloxypyrazole (TIA-230), chlorpyrifos, coumaphos, demeton, demeton-S-methyl, diazinon, dichlorvos, dimethoate, ethoprophos, etrimfos, fenitrothion, fention, heptenophos, parathion, parathion-methyl, phosalone, phoxion, pirimiphos-ethyl, pirimiphos-methyl, profenofos, prothiofos, sulprofos, triazophos and trichlorphon.

- 25 Carbamates, such as aldicarb, bendiocarb, BPMC (2-(1-methylpropyl)phenyl methylcarbamate), butocarboxim, butoxycarboxim, carbaryl, carbofuran, carbosulfan,

Le A 31 810-Foreign countries

- 32 -

cloethocarb, isoprocarb, methomyl, oxamyl, pirimicarb, promecarb, propoxur and thiodicarb.

Pyrethroids, such as allethrin, alphamethrin, bioresmethrin, byfenthrin (FMC 54 800), cycloprothrin, cyfluthrin, 5 decamethrion, cyhalothrin, cypermethrin, deltamethrin, alpha-cyano-3-phenyl-2-methylbenzyl 2,2-dimethyl-3-(2-chloro-2-trifluoromethylvinyl)-cyclopropanecarboxylate, fenpropathrin, fenfluthrin, fenvalerate, flucythrinate, flumethrin, fluvalinate, permethrin and resmethrin; 10 nitroimino and nitroimides, such as 1-[(6-chloro-3-pyridinyl)-methyl]-4,5-dihydro-N-nitro-1H-imidazole-2-amine (imidacloprid).

Examples of herbicides which may be mentioned are anilides, such as, for example, diflufenican and propanil; 15 arylcarboxylic acids, such as, for example, dichloropicolinic acid, dicamba and picloram; aryloxyalkanoic acids, such as, for example, 2,4-D, 2,4-DB, 2,4-DP, fluroxypyr, MCPA, MCPP and triclopyr; aryloxy-phenoxy-alkanoic acid esters, such as, for example, 20 diclofop-methyl, fenoxaprop-ethyl, fluazifop-butyl, haloxyfop-methyl and quizalofop-ethyl; azinones, such as, for example, chloridazon and norflurazon; carbamates, such as, for example, chlorpropham, desmedipham, phenmedipham and propham; chloroacetanilides, such as, for example, alachlor, 25 acetochlor, butachlor, metazachlor, metolachlor, pretilachlor and propachlor; dinitroanilines, such as, for example, oryzalin, pendimethalin and trifluralin; diphenyl ethers, such as, for example, acifluorfen, bifenox, fluoroglycofen, fomesafen, halosafen, lactofen and 30 oxyfluorfen; ureas, such as, for example, chlortoluron, diuron, flometuron, isoproturon, linuron and methabenzthiazuron; hydroxylamines, such as, for example, alloxydim, clethodim, cycloxydim, sethoxydim and

Le A 31 810-Foreign countries

- 33 -

tralkoxydim; imidazolinones, such as, for example, imazethapyr, imazamethabenz, imazapyr and imazaquin; nitriles, such as, for example, bromoxynil, dichlorbenil and ioxynil; oxacetamides, such as, for example, mefenacet;

5 sulphonylureas, such as, for example, amidosulfuron, bensulfuron-methyl, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, metsulfuron-methyl, nicosulfuron, primisulfuron, pyrazosulfuron-ethyl, thifensulfuron-methyl, triasulfuron and tribenuron-methyl; thiocarbamates, such as,

10 for example, butylate, cycloate, diallate, EPTC, esprocarb, molinate, prosulfocarb, thiobvencarb and triallate; triazines, such as, for example, atrazin, cyanazin, simazin, simetryne, terbutryne and terbutylazin; triazinones, such as, for example, hexazinon, met amitron and metribuzin;

15 others, such as, for example, aminotriazole, benfuresate, bentazone, cinmethylin, clomazone, clopyralid, difenzoquat, dithiopyr, ethofumesate, fluoro-chloride, glufosinate, glyphosate, isoxaben, pyridate, quinchlorac, quinmerac, sulphosate and tridiphane.

20 The shaped articles according to the invention comprise 0.1 to 70% by weight of active compound, preferably 1 to 65% by weight of active compound, particularly preferably 5 to 60% by weight of active compound.

25 Water-insoluble thermoplastically processable polymers are those having a processing temperature of 60 to 210°C, preferably 80 to 150°C, and a solubility of less than 100 mg in one litre of water at 20°C.

30 Polymers which may be mentioned are polyolefins, such as polyethylene, polypropylene and polyisobutylene; vinyl polymers, such as polyvinyl chloride, polyvinyl acetate, polystyrene, polyacrylonitrile, polyacrylates, polymethacrylates, polyacetals; polyesters, such as poly-

Le A 31 810-Foreign countries

- 34 -

hydroxybutyric acid, polyhydroxyvaleric acid, polyalkylene terephthalate; polyamides; polyurethanes; polyethers and polycarbonates.

5 Copolymers, which as overall are distinguished by particularly favourable processing properties, may furthermore be mentioned, such as, for example, copolymers of ethylene and (meth)acrylates, styrene/acrylonitrile copolymers, acrylonitrile/butadiene/styrene polymers, styrene/butadiene copolymers and olefin/maleic anhydride
10 polymers.

Other water-insoluble thermoplastically processable polymers are starch derivatives and cellulose derivatives, such as cellulose esters, cellulose ethers and cellulose nitrate.

15 Vinyl acetate polymers are particularly suitable for production of the shaped articles according to the invention. Vinyl acetate polymers which may be mentioned are polyvinyl acetate and vinyl acetate copolymers, if these are water-insoluble.

20 Copolymers with vinyl chloride, vinylidene chloride and ethylene may be mentioned as particularly preferred. Ethylene/vinyl acetate copolymers are preferred.

Ethylene/vinyl acetate copolymers having a vinyl acetate content of 30 to 80% are particularly preferred. It is also possible to employ ethylene/vinyl acetate copolymers in
25 which all or some of the acetate groups are hydrolysed to alcohol groups.

Water-soluble polymers which may be mentioned are cellulose derivatives, such as, for example, methylcellulose, hydroxyethylcellulose, hydroxypropylcellulose and

Le A 31 810-Foreign countries

- 35 -

carboxyethylcellulose. There may also be mentioned polyacrylic acids and salts thereof, polyacrylamide, polyvinylpyrrolidone, vinylpyrrolidone copolymers and polyalkylene oxides, such as polyethylene oxide, and
5 copolymers of ethylene oxide and propylene oxide. Polyvinyl alcohol may furthermore be mentioned, it being possible to add a hydrophilic plasticizer in this case to improve the processability. Suitable hydrophilic plasticizers for polyvinyl alcohol are glycerol and triethylene glycol, these
10 preferably being added in proportions of 5 to 25% by weight, based on the polyvinyl alcohol. Preferred water-soluble polymers are polyvinylpyrrolidone and copolymers of vinylpyrrolidone, comonomers which may be mentioned as examples being acrylic acid, methacrylic acid, itaconic
15 acid, maleic acid, hydroxyalkyl methacrylate, hydroxyalkyl acrylate and vinyl acetate. Copolymers of polyvinylpyrrolidone and vinyl acetate are preferred.

The weight ratio of water-insoluble thermoplastically processable polymer to water-soluble polymer is 15:85 to
20 16:40, preferably 30:70 to 60:40.

The shaped articles according to the invention can comprise customary additives, such as fillers, plasticizers, stabilizers and lubricants. Examples of fillers are calcium carbonate, magnesium carbonate, silicon dioxide, aluminas,
25 silicates, talc. Particularly suitable plasticizers are esters of phthalic acid and sebacic acid as well as oligomeric ethylene oxides.

The shaped articles according to the invention can be produced from the corresponding blends by extrusion or
30 injection moulding. The blends can be prepared from the individual components by known mixing techniques, for example by kneading in the melt, by extrusion or by

Le A 31 810-Foreign countries

- 36 -

dissolving in a common solvent, from which the blends can be obtained by precipitation with a precipitant or by evaporating of the solvent. Processing by extrusion is preferred.

- 5 The shaped articles according to the invention show a high rate of release of the active compound and very low swelling in water. In general, the swelling is less than 15%. The swelling is defined as the relative increase in weight after storage in water. For determination of this swelling,
10 cylindrical test specimens having a length of 16 mm and a diameter of 4 mm are immersed in water for different times (24 hours, 72 hours, 120 hours), and the weight is determined, after removal of the water adhering to the surface.
- 15 It is particularly important for the volume of the shaped articles to be constant if these shaped articles are to be employed as an implant for the treatment of individual plants. In particular, biological tests have shown that implants which swell significantly in water are poorly
20 tolerated by the plant. Furthermore, the release of active compound from swellable implants into the sap stream of a plant is unfavourable. An increase in volume of not more than the free volume between the preformed hollow space and implant is in general not critical.
- 25 The agrochemical active compounds can be incorporated into the polymer components, a homogeneous distribution of the active compound being achieved in a simple manner, that is to say even under low sheer forces and during short mixing times. The blends containing active compounds are
30 distinguished by an easy processability, in particular by injection moulding.

Le A 31 810-Foreign countries

- 37 -

The blends containing active compound can be employed in the form of films, nets, tiles, woven fabrics or tapes. It is also possible to produce plant vessels, for example pots for growing, from this material.

- 5 The blends according to the invention can particularly preferably be employed in the form of shaped articles for treatment of individual plants, such as, for example, trees. For this, they are preferably incorporated into the sap stream of the plant in the form of suitable shaped articles,
- 10 such as rods, pins, needles, rivets, pegs, clamps, wires, beads, tablets or sheets. A hollow is preferably first produced, into which the shaped article is incorporated as an implant.

Le A 31 810-Foreign countries

- 38 -

Example 1

Production of a blend and shaped article according to the invention

- a) Polymer mixture
- 5 A compound of 50 parts of polyvinyl acetate (Moviol 50), 50 parts of poly(pyrrolidone-covinyl acetate) (Luviskol VA64) and 1 part of highly dispersed silicon dioxide (HDK H2000) a polymer strand is produced in a twin-screw extruder at a speed of rotation of 80 rpm, a melt temperature of 150°C and a nozzle discharge temperature of 175°C and is comminuted with a granulator.
- 10
- b) Blend containing active compound
- 15 75 parts of the granules from 1a) and 25 parts of imidachloride were mixed in a twin-screw extruder at a speed of rotation of 80 rpm, a melt temperature of 145°C and a nozzle discharge temperature of 165°C and then granulated.
- c) Shaped article containing active compound
- 20 Cylindrical shaped articles (length: 16 mm, diameter: 4 mm) containing active compound were produced from the blend 1b) in an injection moulding machine at a melt temperature of 150°C, a nozzle temperature of 155°C and a mould temperature.

Le A 31 810-Foreign countries

- 39 -

Example 2

Production of blends and shaped articles according to the invention

- a) Polymer mixture
- 5 50 parts of dried poly(vinyl acetate-co-ethylene) (Levapren 450) and 50 parts of dried polyvinylpyrrolidone (Luviskol K30) were mixed and processed to granules in accordance with the procedure described under 1a).
- 10 b) Blend containing active compound
- 50 parts of the granules from 2a) and 50 parts of imidachloride were mixed in a twin-screw extruder at a speed of rotation of 60 rpm, a melt temperature of 140°C and a nozzle discharge temperature of 165°C and
- 15 then granulated.
- c) Implant containing active compound
- Shaped articles containing active compound and having the dimensions described above were produced from blend 2b) in an injection moulding machine at a melt
- 20 temperature of 150°C, a nozzle temperature of 155°C and a mould temperature of 50°C.

Example 3-5

Production of blends and shaped articles according to the invention

- 25 a) Polymer mixture
- 40 parts of dried poly(vinyl acetate-co-ethylene) (Levapren 700) and 60 parts of dried poly(vinylpyrrolidone-co-vinyl acetate) (Luviskol VA64)

Le A 31 810-Foreign countries

- 40 -

were mixed and processed to granules in accordance with the procedure described under 1a).

b) Blend containing active compound

5 Blends containing active compound, of the following composition, were prepared in a twin-screw extruder at a speed of rotation of 100 rpm, a melt temperature of 140°C and a nozzle discharge temperature of 165°C.

Example 3b: 75.0 parts of granules from 3a)
25 parts of imidachloride

10 Example 4b: 66.7 parts of granules from 3a)
33.3 parts of imidachloride

Example 5b: 50.0 parts of granules from 3a)
50 parts of imidachloride

c) Shaped article containing active compound

15 Shaped articles containing active compound and having the dimensions described above were produced from blends 3b), 4b) and 5b) in an injection moulding machine at a melt temperature of 135°C, a nozzle temperature of 147°C and a mould temperature of 44°C.

20 Example A, Comparison experiment**Preparation of a polymer mixture according to EP 0 344 118**

a) Polymer mixture

25 A blend was prepared from 90 parts of gelatin and 10 parts of polystyrene in accordance with the instructions from Example 1, No. 5.

b) Blend containing active compound

75 parts of the granules from 6a) and 25 parts of imidachloride were mixed in a twin-screw extruder at a

Le A 31 810-Foreign countries

- 41 -

speed of rotation of 60 rpm, a melt temperature of 155°C and a nozzle discharge temperature of 165°C and then granulated.

- 5 c) Shaped article containing active compound
Shaped articles having the dimensions described under 1 were produced from blend 6b) in an injection moulding machine at a melt temperature of 155°C, a nozzle temperature of 165°C and a mould temperature of 45°C.
- 10 d) Shaped article containing no active compound
For further comparison, the granules 6a) containing no active compound were also processed to shaped articles under the conditions described in 6c).

Determination of the swelling

15 In each case 9 shaped articles from Examples 1 to 5 and A, the individual dry weight of which had been determined accurately beforehand, were stored in a 5 l water container with an overflow, 1 l/hour of fresh water being passed into the container. After 24 hours, 78 hours and 120 hours, in
20 each case 3 test specimens were removed and freed from adhering water on the surface by wiping with a cotton pad, and the weight m_{wet} was determined. The parameter $(m_{\text{wet}} - m_0) \cdot 100 / m_0$ was defined as the swelling value.

Shaped article	24 hours	72 hours	120 hours
1c	10.4%	8.5%	2.5%
2c	5.9%	1.2%	-3.1%
3c	3.6%	-1.6%	-8.1%
4c	3.3%	-1.0%	-7.2%

Le A 31 810-Foreign countries

- 42 -

Shaped article	24 hours	72 hours	120 hours
5c	8.5%	0.6%	-4.0%
Ac	461%	2115%	2133%
Ad	595%	1870%	2895%

Le A 31 810-Foreign countries

- 43 -

Patent claims

1. Shaped articles which release agrochemical active compounds, are used for treatment of woody plants and
 - are employed in a hollow preformed in the woody plant,
 - 5 - are of a volume which fills the volume of the hollow to the extent of 10 to 95%,
 - are of a volume which does not increase during the period of use to the extent that the preformed hollow space is filled completely,
 - 10 - comprise
 - a) agrochemical active compounds,
 - b) water-insoluble thermoplastically processable polymers,
 - c) water-soluble polymers and
 - 15 d) if appropriate further auxiliaries.
2. Shaped articles according to Claim 1 which release agrochemical active compounds, characterized in that the water-insoluble thermoplastically processable polymer is a vinyl acetate polymer.
- 20 3. Shaped articles according to Claim 1 which release agrochemical active compounds, characterized in that the weight ratio of water-insoluble thermoplastically processable polymer to water-soluble polymer is 15:85 to 60:40.
- 25 4. Shaped articles according to Claim 1 which release agrochemical active compounds, characterized in that they comprise 0.1 to 70% by weight of active compound.
5. Polymer blends comprising

Le A 31 810-Foreign countries

- 44 -

- a) agrochemical active compounds,
 - b) water-insoluble thermoplastically processable polymers,
 - c) water-soluble polymers
 - 5 d) if appropriate further auxiliaries.
6. Process for the production of shaped articles which release agrochemical active compounds, characterized in that blends according to Claim 5 are extruded or processed by injection moulding in a manner known per se.
- 10

**Fetherstonhaugh & Co.
Ottawa, Canada
Patent Agents**