



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2488261 C 2012/10/02

(11)(21) **2 488 261**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2003/05/28
(87) Date publication PCT/PCT Publication Date: 2003/12/11
(45) Date de délivrance/Issue Date: 2012/10/02
(85) Entrée phase nationale/National Entry: 2004/12/01
(86) N° demande PCT/PCT Application No.: EP 2003/005725
(87) N° publication PCT/PCT Publication No.: 2003/101213
(30) Priorité/Priority: 2002/06/03 (EP02100662.2)

(51) Cl.Int./Int.Cl. *A23C 19/11* (2006.01),
A23C 19/086 (2006.01), *A23L 1/00* (2006.01)
(72) Inventeurs/Inventors:
WARMERDAM, MARTINUS JOHANNES MARIA, NL;
HAAN, DE, BEN RUDOLF, NL
(73) Propriétaire/Owner:
DSM IP ASSETS B.V., NL
(74) Agent: SMART & BIGGAR

(54) Titre : AMELIORATION APPOREE AU TRAITEMENT DE FROMAGE RAPE AU MOYEN D'UN COMPOSE
POLYENE ANTIFONGIQUE

(54) Title: IMPROVED METHOD FOR THE TREATMENT OF SHREDDED CHEESE WITH A POLYENE ANTIFUNGAL
COMPOUND

(57) **Abrégé/Abstract:**

A method for the treatment of shredded cheese with an aqueous composition whereby the aqueous composing comprises a polyene antifungal compound e.g. notamycine and a thickening agent, e.g. xanthan. Use of a thickening agent to prevent nozzle clogging during the treatment of food, feed and agricultural products with an aqueous composition.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 December 2003 (11.12.2003)

PCT

(10) International Publication Number
WO 03/101213 A1

(51) International Patent Classification⁷: **A23C 19/11**,
19/086, A23L 1/00

(21) International Application Number: PCT/EP03/05725

(22) International Filing Date: 28 May 2003 (28.05.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02100662.2 3 June 2002 (03.06.2002) EP

(71) Applicant (*for all designated States except US*): **DSM IP ASSETS B.V.** [NL/NL]; Het Overloon 1, NL-6411 TE HEERLEN (NL).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants and

(72) Inventors: **WARMERDAM, Martinus, Johannes, Maria** [NL/NL]; Pauwhof 179, NL-2289 BM RIJSWIJK (ZH) (NL). **HAAN, DE, Ben, Rudolf** [NL/NL]; Watertorenlaan 17, NL-2275 AW VOORBURG (NL).

(74) Agent: **MATULEWICZ, Emil, Rudolf, Antonius**; DSM Intellectual Property, Delft Office (600-0240), P.O. Box 1, NL-2600 MA DELFT (NL).


Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: IMPROVED METHOD FOR THE TREATMENT OF SHREDDED CHEESE WITH A POLYENE ANTIFUNGAL COMPOUND

(57) Abstract: A method for the treatment of shredded cheese with an aqueous composition whereby the aqueous composing comprises a polyene antifungal compound e.g. notamycine and a thickening agent, e.g. xanthan. Use of a thickening agent to prevent nozzle clogging during the treatment of food, feed and agricultural products with an aqueous composition.



WO 03/101213 A1

IMPROVED METHOD FOR THE TREATMENT OF SHREDDED CHEESE WITH A POLYENE ANTIFUNGAL COMPOUND

5

Field of the invention

The present invention relates to a method for the treatment of shredded cheese with a polyene antifungal compound.

10

Background of the invention

The prevention of mould growth is an important topic to the food, feed and agricultural industry, especially in the food industry. Fungal spoilage can lead to serious economic losses. Some foods can be considered as a good substrate for fungal growth.

15 Cheese is an example of such a product. Apart from the negative appearance of fungal growth on cheese, fungal spoilage is also a health risk. Some mould species produce mycotoxins, which may penetrate into the product (J.C. Frisvad & U. Thane; "Mycotoxin production by food-borne fungi" in Introduction to food-borne fungi, 4th edn. (ed. R.A. Samson et al.), 1995, 251-260. Therefore, superficial removal of moulds gives no
20 guarantee of safety to the consumer.

For more than 30 years, natamycin has been used to prevent growth of moulds and yeasts on cheeses and sausages. Natamycin is on the market under the brand name of Delvacid®, a powder composition containing 50% (w/w) of natamycin.

Food products can be treated with natamycin in different ways. Natamycin can
25 for example be added to the polymer dispersion that is applied to the cheese rind as a coating (C.B.G. Daamen & G. van den Berg "Prevention of mould growth on cheese by means of natamycin" Voedingsmiddelentechnologie, 1985, 18 (2), 26-29)

Alternatively, food products such as cheeses and sausages can be treated by dipping or spraying with a suspension of natamycin in water (C.B.G. Daamen & G. van
30 den Berg "Prevention of mould growth on cheese by means of natamycin" Voedingsmiddelentechnologie, 1985, 18 (2), 26-29; H.A. Morris & H.B. Castberg "Control of surface growth on blue cheese using pimarinic" Cultured Dairy Products Journal, 1980, 15 (2), 21-23; P. Baldini, F. Palmia, R.G. Raczynski, M. Campanini, "Use of pimarinic for

preventing mould growth on Italian cured meat products", Industria Conserve, 1979, 54 (4), 305-307; R.A. Holley, "Prevention of surface mould growth on Italian dry sausage by natamycin and potassium sorbate", Appl. Environ. Microbiol., 1981, 41 (2).

5 Spraying of a suspension of natamycin can be applied for the treatment of shredded cheese, a cheese product which is very sensitive to fungal spoilage. The production of shredded cheese is as follows: cheese, e.g. Mozzarella, pizza cheese, or Cheddar is shredded and conveyed to a revolving tumbler. An anti-caking agent, for example cellulose (microcrystalline or powdered), starch or modified starch is metered
10 onto the cheese in the first part of the tumbler. Cellulose forms a film around the cheese particles which prevents caking of the shredded cheese. At the end of the tumbler a suspension of natamycin (e.g. Delvolid®) is sprayed onto the shredded cheese. The spray nozzles deliver a fine spray or mist of the fungicide onto the cheese. Finally, the cheese empties onto a conveyor to be transported to the filling equipment.

15 The natamycin suspension is prepared by mixing Delvolid® and water, mostly at the ratio from 2 to 10 grams of Delvolid® to 1 litre of water. The suspension is stored in a liquid container. To prevent sedimentation of the suspension a small amount of the suspension, which is pumped under pressure to spray nozzles, is recirculated to the liquid container via a jet agitator. Alternatively, a stirring device can be used to prevent
20 sedimentation. Mostly approximately 6 to 8 litres of Delvolid® suspension is sprayed onto 1000 kg of cheese. Mostly this treatment is effective to prevent fungal spoilage.

 The spraying or showering of aqueous suspensions on cheeses using nozzles is an effective way of coating the cheeses with the antifungal compounds. In general a less fine spray is used for spraying of complete cheeses. This spray or shower will be
25 focused or directed onto the cheese. However for spraying of shredded cheese, the suspension should be added as a fine spray or mist. This spraying method often gives rise to problems such as the clogging of the spraying nozzles.

70500-110

3

Description of the invention

It now has been found that protection against the development of moulds and yeasts shredded cheese is obtained in a method for the treatment shredded cheese with a composition comprising a polyene antifungal compound, whereby the composition comprises a polyene antifungal agent, a thickening agent and water. Surprisingly it has been found that the thickening agent prevents the clogging of the nozzles. Moreover the aqueous composition can be pumped under pressure to and through the nozzles without problems which would have been expected due to the expected increase of viscosity of the aqueous composition of the invention compared to an aqueous composition without thickening agent. Especially xanthan gives very good results.

Therefore the present invention relates to a method for the treatment of shredded cheese with an aqueous composition whereby the aqueous composition comprises a polyene antifungal compound and a thickening agent.

The invention further relates to a method for preventing nozzle clogging during the treatment of shredded cheese by spraying with an aqueous composition comprising natamycin, wherein the aqueous composition comprises a thickening agent.

The invention further relates to a method for preparing shredded cheese treated with an aqueous composition comprising natamycin, the method comprising the steps of: (i) shredding cheese, (ii) conveying the shredded cheese to a revolving tumbler, (iii) adding an anticaking agent to the shredded cheese in the tumbler, (iv) spraying an aqueous composition comprising natamycin and a thickening agent onto the shredded cheese in the tumbler, the presence of the thickening agent preventing clogging of nozzles during spraying, and (v) transporting the shredded cheese to a filling equipment.

In one aspect, the invention relates to use of a thickening agent for preventing nozzle clogging during the treatment of shredded cheese, wherein the shredded cheese is sprayed with an aqueous solution comprising natamycin and the thickening agent.

70500-110

3a

In another aspect, the invention relates to use of a thickening agent for preventing nozzle clogging during the preparation of shredded cheese, wherein the use comprises: (i) shredding cheese, (ii) conveying the shredded cheese to a revolving tumbler, (iii) adding an anticaking agent to the shredded cheese in the
5 tumbler, (iv) spraying an aqueous composition comprising natamycin and the thickening agent onto the shredded cheese in the tumbler, the presence of the thickening agent preventing clogging of nozzles during spraying, and (v) transporting the shredded cheese to a filling equipment.

The present invention also relates to products produced with the
10 present method. Shredded cheese coated with a polyene antifungal compound and thickening agent forms part of this invention. A thickening agent is used to prevent the nozzle clogging during the treatment of food, feed and agricultural products with an aqueous composition.

The polyene antifungal compounds used in the composition of the
15 invention are preferably one or more of natamycin, nystatin, lucensomycin or amphotericin B. The preferred polyene compound is natamycin. In the preparation of compositions of the invention, one or more polyene antifungal compounds may be used or pre-prepared compositions containing such antifungal components may be used. An example of such a preparation is commercially
20 available powder composition sold under the trade mark Delvolid® which contains 50% (w/w) natamycin.

The concentration of polyene fungicide in the aqueous composition will generally be from 0.1 to 20 g/l, more preferably from 0.5 to 8 g/l and most preferably from 1 to 5 g/l.

25 In compositions of the invention the preferred thickening agents include all thickening agents known in the art for use in food products, preferably the thickening agents are gum, more preferably xanthan gum and/or gellan gum.

The one or more thickening agents in the aqueous composition are preferably present in an amount of 0.5 to 50 g/l, more preferably from 0.5 to 20 g/l and most preferably from 1 to 5 g/l.

5 In some embodiments it has been found that no buffer is needed for the suspension. In general the pH of the suspension will be between 3-10, preferably the pH will be between 6-9, more preferably between 6.2-8.5 for the optimal natamycin addition.

10 Preferably also an anticaking agent is added to the shredded cheese. The anticaking agent preferably is a dry powder composition, which is added separately from the aqueous composition. In general 0.2 to 3% w/w of anticaking agent (on shredded cheese) preferably between 1.5 and 2.5%w/w.

The steps of adding the dry powder composition and the aqueous composition may be performed in any order. Preferably, the dry powder composition is added before the aqueous composition is added.

15 The amount of the aqueous composition added to the shredded cheese may be between 0.01% and 5% (v/w), preferably said amount is at least 0.5% or preferably lower than 2% (v/w).

20 In a preferred embodiment, the aqueous composition used in the method according to the present invention is an aqueous solution or an aqueous dispersion such as an emulsion or suspension.

25 The optimal final concentration of natamycin added according to the method of the present invention, may vary with the type of cheese used as shredded cheese or the kind of shredded used. To prevent fungal growth on shredded cheese, the optimal final concentration may be between 2 and 40 ppm natamycin, preferably the concentration is at least 3 ppm natamycin, more preferably at least 4 ppm natamycin. The concentration is preferably lower than 20 ppm, more preferably lower than 10 ppm.

30 Any anti-caking agent may be used. Examples of suitable anti-caking agents are microcrystalline or powdered cellulose, starch, modified starch, sodium-, magnesium-, potassium-, and zinc silicate, silicium-dioxide, kaolin, talc, potassium and magnesium carbonate, phosphates (di, tri and polyphosphates including sodium, potassium and calcium salts). Also commercial products containing cellulose, such as Keycel or

Floam® may be used as anti-caking agent. Further, any combination of anti-caking agents may be used.

The aqueous composition comprising an antifungal compound, e.g. natamycin, may be prepared by mixing a dry formulation, e.g. Delvacid®, in water using well known methods. Preferably a natamycin suspension for spraying is prepared by using a stock suspension of natamycin as described in EP 0 678 241. This stock suspension is a chemically, physically and microbially stable concentrated suspension of natamycin, which provides a convenient stock for the easy and reproducible dosage of a fungicide to food, feed and agricultural products. Apart from natamycin, the suspension comprises a thickening agent, for example xanthan. The concentration of natamycin in the stock suspension may be as high as 40% (w/w), however also stock suspensions with lower concentrations of natamycin can be used. An example of such a stock suspension is the commercial product Delvacid®-Sol. This product comprises at least 50% (w/w) of natamycin; for example 50 to 54% (w/w) of natamycin, 2% (w/w) of xanthan and 44 to 48% (w/w) of lactose. In the case of shredded cheese the stock suspension is diluted with water to the final concentration required for the treatment. This dilution step can be best executed just before adding the natamycin suspension to the shredded cheese.

In cases where the polyene antifungal compound is natamycin, complexes of natamycin (e.g. with proteins) and salts of natamycin (e.g. the calcium salt) may also be used in the method according to the present invention. Further, the dry and aqueous compositions may also contain components, which are already present in a natamycin preparation. For example, when Delvacid-Instant® or Natamax® is used, lactose will also be present in the final composition.

The following Examples are for illustrative purposes only and are not to be construed as being limiting to the invention.

Examples

Cheddar cheese was shredded using well known methods and conveyed to a revolving tumbler. Microcrystalline cellulose was metered onto the shredded cheese in the first part of the tumbler and thoroughly mixed into the shredded cheese. At the other side of the tumbler, extending about 30 – 60

cm into the tumbler, spray nozzles are situated (S.S. Fogger Tip Assembly #22624-2-80: 80° angle, 2 gal. (8 liters) per hour at 80 PSI (5.3 bar, purchased from Spraying Systems Co. PO Box 95584 Chicago IL 60694). The nozzles are able to deliver a fine spray of mist of one of the following natamycin formulations.

5 Two experiments were carried out using either formulation A (reference) or formulation B (present invention).

A) A suspension of Delvacid®, available by DSM Food Specialties USA, by mixing 10 grams of Delvacid® with 4 liters of tapwater.

10 B) A suspension of Delvacid® and xanthan by mixing 10 grams of Delvacid® with 4 liters of tapwater together with 3 grams Keltrol® RD, available by NutraSweet Kelco Company.

Both suspensions were sprayed on shredded cheese with a flow of 10 liters per hour.

15 The spray characteristics were followed in time during a 2 shift operation, 16 hours per day.

During the tests it was shown that the nozzles could be used with suspension A for a 1 – 6 hour operation. After this period the nozzles were completely blocked by clogging. The nozzles needed to be cleaned completely before further operation was possible.

20 Using suspension B resulted in a operation of 16 hours without any problems such as clogging.

70500-110

7

CLAIMS:

1. Use of a thickening agent for preventing nozzle clogging during the treatment of shredded cheese, wherein the shredded cheese is sprayed with an aqueous solution comprising natamycin and the thickening agent.
- 5 2. The use according to claim 1, wherein the thickening agent is a gum.
3. The use according to claim 1 or 2, wherein the thickening agent is a xanthan gum, gellan gum or a combination thereof.
4. The use according to any one of claims 1-3, wherein the concentration of natamycin in the aqueous composition is from 0.1 to 20 g/l.
- 10 5. The use according to any one of claims 1-4, wherein the concentration of the thickening agent in the aqueous composition is from 0.5 to 50 g/l.
6. The use according to any one of claims 1-5, wherein the method further comprises adding a dry powder composition comprising at least one anticaking agent to the shredded cheese.
- 15 7. The use according to claim 6, wherein the dry powder composition is added before treatment of the shredded cheese by spraying with the aqueous composition.
8. The use according to claim 6 or 7, wherein the anticaking agent is selected from the group consisting of cellulose, starch, modified starch, sodium
20 silicate, magnesium silicate, potassium silicate, zinc silicate, silicium dioxide, kaolin, talc, potassium carbonate, magnesium carbonate and phosphates.
9. The use according to any one of claims 1-8, wherein the aqueous composition is sprayed onto the shredded cheese in an amount of between 0.01 and 5% (v/w).

70500-110

8

10. The use according to any one of claims 1-9, wherein natamycin is sprayed onto the shredded cheese in a final concentration of between 2 and 40 ppm.

11. Use of a thickening agent for preventing nozzle clogging during the preparation of shredded cheese, wherein the use comprises:

5 (i) shredding cheese,

(ii) conveying the shredded cheese to a revolving tumbler,

(iii) adding an anticaking agent to the shredded cheese in the tumbler,

(iv) spraying an aqueous composition comprising natamycin and the thickening agent onto the shredded cheese in the tumbler, the presence of the
10 thickening agent preventing clogging of nozzles during spraying, and

(v) transporting the shredded cheese to a filling equipment.