



(22) Date de dépôt/Filing Date: 2010/04/23
(41) Mise à la disp. pub./Open to Public Insp.: 2010/10/28
(45) Date de délivrance/Issue Date: 2020/02/25
(62) Demande originale/Original Application: 2 758 627
(30) Priorité/Priority: 2009/04/23 (GB0907003.8)

(51) Cl.Int./Int.Cl. *C07C 43/15* (2006.01),
A01N 25/30 (2006.01)
(72) Inventeur/Inventor:
BELL, GORDON ALASTAIR, GB
(73) Propriétaire/Owner:
SYNGENTA PARTICIPATIONS AG, CH
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : ALCOXYLATES D'ALCOOL COMME ADJUVANTS POUR FORMULATIONS AGROCHIMIQUES
(54) Title: ALCOHOL ALKOXYLATES AS ADJUVANTS FOR AGROCHEMICAL FORMULATIONS

(57) **Abrégé/Abstract:**

A bioperformance enhancing adjuvant of Formula (I) $R_1O[BO]_n[AO]_mR_2$ where BO is butylene oxide; and each AO is independently propylene oxide or ethylene oxide; n is from 1 to 12; m is from 0 to 20; R_1 is optionally substituted C_{4-20} alkyl or optionally substituted C_{4-20} alkenyl; and R_2 is hydrogen or optionally substituted C_{1-3} alky.

ABSTRACT

A bioperformance enhancing adjuvant of Formula (I) $R_1O[BO]_n[AO]_mR_2$ where BO is butylene oxide; and each AO is independently propylene oxide or ethylene oxide; n is from 1 to 12; m is from 0 to 20; R_1 is optionally substituted C_{4-20} alkyl or optionally substituted C_{4-20} alkenyl; and R_2 is hydrogen or optionally substituted C_{1-3} alkyl.

ALCOHOL ALKOXYLATES AS ADJUVANTS FOR AGROCHEMICAL FORMULATIONS

This invention relates to bioperformance enhancing adjuvants and to use of such adjuvants.

Numerous adjuvants which enhance the bioperformance of agrochemicals are known. A broad mention of alcohol alkoxylates as penetration aids is made in WO2008/037375.

The present invention provides a bioperformance enhancing adjuvant of formula (I):



where BO is butylene oxide; and each AO is independently propylene oxide or ethylene oxide; n is from 1 to 12; m is from 0 to 20; R₁ is optionally substituted C₄₋₂₀ alkyl or optionally substituted C₄₋₂₀ alkenyl; and R₂ is hydrogen or optionally substituted C₁₋₄ alkyl; provided that R₁ comprises more carbon atoms than R₂.

Each alkyl chain is, independently, straight or branched.

Optional substituents on the alkyl and alkenyl groups are, independently, hydroxy and epoxy groups.

Certain butylene oxide-ethylene oxide block copolymers are disclosed by J.Chlebicki in J.Colloid and Interface Science 206, 77-82 (1998).

Therefore in another aspect, the present invention provides a compound of formula (I) as defined above provided that it is not a compound of formula (Ia):



when R₃ is *n*-butyl, *n*-hexyl, *n*-octyl or *n*-decyl; a is 1, 2, 3 or 4; and b is from 9.7 to 10.2.

When R₁ is optionally substituted alkyl, suitably R₁ is optionally substituted C₆₋₁₈ alkyl; more suitably R₁ is optionally substituted C₁₀₋₁₆ alkyl; even more suitably R₁ is optionally substituted C₁₃₋₁₅ alkyl.

When R₁ is optionally substituted alkenyl, suitably R₁ is optionally substituted C₆₋₁₈ alkenyl; more suitably R₁ is optionally substituted oleyl; even more suitably R₁ is oleyl.

Suitably R₂ is hydrogen or optionally substituted C₁₋₃ alkyl; more suitably R₂ is hydrogen or optionally substituted C₁₋₂ alkyl; even more suitably R₂ is hydrogen or optionally substituted methyl; most suitably R₂ is hydrogen.

Suitably R₁ is non-substituted alkyl.


Suitably R₂ is hydrogen or non-substituted alkyl.

BO [butylene oxide] has the empirical formula C₄H₈O. Every BO unit has the formula CH(R₄)CH(R₅)O but each BO unit is independently selected from the following

options: R_4 is methyl and R_5 is methyl; or R_4 is ethyl and R_5 is hydrogen; or R_4 is hydrogen and R_5 is ethyl.

Propylene oxide [PO] has the empirical formula C_3H_6O . Every PO unit has the formula $CH(R_6)CH(R_7)O$ but each PO unit is independently selected from the following options: R_6 is methyl and R_7 is hydrogen; or R_6 is hydrogen and R_7 is methyl.

In one aspect, the block $[AO]_m$ is a PO block followed by an EO block, where the PO block is bonded to the BO block, such that the adjuvant is of formula (Ib):



where $(m' + m'' = m)$ and R_1 , R_2 , n and m are as defined for compounds of formula (I).

Suitably AO is ethylene oxide.

Suitably n is from 2 to 8; more suitably from 3 to 6; even more suitably it is 4.

Suitably m is from 5 to 15; more suitably it is from 8 to 12; even more suitably it is 10.

In one aspect, $m=0$. Adjuvants for which $m=0$ may display very low levels of phytotoxicity [i.e. little damage to plants].

The bioperformance enhancing adjuvants of the present invention may be used effectively at much lower concentrations than the effective concentrations for conventional adjuvants.

The bioperformance enhancing adjuvants of the present invention may be used synergistically with other bioperformance enhancing adjuvants of the present invention or with conventional adjuvants.

The values of n and m represent values both for individual species and for averages taken over a distribution of compounds. This will be well understood by the skilled person.

Suitably the bioperformance enhancing adjuvants of the present invention are used to enhance the bioperformance of a pesticide. Pesticides suitable for use with the present invention include insecticides, fungicides, herbicides, acaricides, nematocides and biocides suitable for controlling pests, diseases or weeds that are a problem in agriculture. Many such pesticides are known and are described in The Pesticide Manual 14th edition published by the British Crop Protection Council in 2006.

The invention is illustrated by the following non-limiting Examples in which all parts and percentages are by weight unless otherwise stated.

EXAMPLE 1

This example illustrates compounds of the present invention. Twenty eight butylene oxide based compounds were synthesised using standard techniques familiar to a skilled person [for example, see EP0681865A]. Each sample consists of a hydrocarbon tail connected to a butylene oxide section which in turn is connected to an ethylene oxide section; the samples are compounds of formula (I) in which AO is ethylene oxide and R₂ is hydrogen; and R₁, n and m are as defined in Table 1. Samples 1 to 3 and 24 to 27 were prepared using branched alcohols containing an average of 13 carbon atoms. Samples 4 to 7 were prepared with 2-ethylhexyl alcohol. Samples 8 to 11 were prepared using butanol as the starting solvent. Samples 12 to 14 and sample 28 used a fraction of alcohols ranging from 12 to 15 carbons in length. Samples 15 to 23 used an alcohol range from C₁₂ to C₁₆. The values of n and m are average values.

Table 1

Sample	R ₁ alkyl chain	n	m
1	C ₁₃ [average]	2	8
2	C ₁₃ [average]	4	8
3	C ₁₃ [average]	6	12
4	2-ethylhexyl	2	6
5	2-ethylhexyl	2	3
6	2-ethylhexyl	4	6
7	2-ethylhexyl	6	10
8	C ₄	2	8
9	C ₄	4	10
10	C ₄	4	8
11	C ₄	6	8
12	C ₁₂₋₁₅	2	9
13	C ₁₂₋₁₅	4	10
14	C ₁₂₋₁₅	6	12
15	C ₁₂₋₁₆	1	3
16	C ₁₂₋₁₆	1	5
17	C ₁₂₋₁₆	1	8
18	C ₁₂₋₁₆	2	5
19	C ₁₂₋₁₆	2	8

20	C ₁₂₋₁₆	3	5
21	C ₁₂₋₁₆	3	8
22	C ₁₂₋₁₆	4	3
23	C ₁₂₋₁₆	4	8
24	C ₁₃ [average]	2	3
25	C ₁₃ [average]	2	8
26	C ₁₃ [average]	4	3
27	C ₁₃ [average]	4	8
28	C ₁₂₋₁₅	4	0
29	C ₁₈ (oleyl)	4	0
30	C ₁₈ (oleyl)	4	20

EXAMPLE 2

This example shows that samples 15, 17, 24 and 25 from Table 1 behave as adjuvants for fungicides used against brown rust (*puccinia recondita*). Wheat was grown outside in field plots. Each plot was sprayed with water at a rate of 15 litres per hectare, the water containing either difenoconazole or cyproconazole, at a concentration which enabled a pesticide application rate of 0.1, 1, 3, 10 or 30 grams per hectare. Adjuvants were added at the standard rate of 0.2% v/v of the spray volume used. The known adjuvants BrijTM 96 and TEHP (tris 2-ethylhexyl phosphate) were also tested for comparison. Each experiment was replicated four times and the results were averaged at each rate. Plants were examined for both protective action and curative effects. A standard mathematical analytical technique was used: Plots of efficacy against pesticide concentration for each adjuvant were 'logit' transformed and used to estimate the concentration required for 90% effect (ED90). For each sample, its ED90 value was compared to that of the known adjuvant BrijTM96 or TEHP in order to generate a relative potency; the relative potency is the ratio of the ED90 values. Relative potencies are given in Table 2 [relative potency results compared to BrijTM96 for difenoconazole] and Table 3 [relative potency results compared to TEHP for cyproconazole]. Adjuvants which performed better than the standard adjuvants have a relative potency value greater than 1.

Table 2

Sample	Protective	Curative
--------	------------	----------

15	0.99	1.50
17	1.47	1.69
24	1.46	2.84
25	1.22	1.07

Table 3

Sample	Protective	Curative
15	1.80	3.81
17	0.97	2.73
24	1.68	2.97
25	1.22	4.29

EXAMPLE 3

5 In this example Sample 13 from Table 1 was compared to the oil adjuvant blend TurbochargeTM. The herbicide fomesafen was applied to the weed species xanthium strumarium (XANST), setaria viridis (SETVI), abutilon theophrasti (ABUTH), and chenopodium album (CHEAL) at rates of 60 or 120 grams per hectare, using a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to
10 the weeds was assessed visually at time periods of 7, 14 and 21 days after application. Each adjuvant was applied at a rate of 0.5% of the volume of the spray water. Four effects were evaluated: pesticide rate (2 levels), adjuvant type (two levels), weed species (four levels) and days after application (three levels). A standard simple linear model was constructed to judge the significance of these effects. These effects were found to be
15 significant at a 5% level. A model was fitted using multiple linear regression using the statistics package JMP (SAS group). The effect of each adjuvant was pulled out from the model and the least significant differences evaluated using a Student's t method. In further examples, where a greater number than two adjuvants were compared, Tukey's HSD method was used.

20

Table 4 shows the mean efficacy of the butylene oxide adjuvant number 13 with the pesticide fomesafen compared to the oil adjuvant Turbocharge across four weed species and compared to single weed species. The mean values and a letter code denoting significant difference (samples with the same letter are not significantly different from

each other at the 5% level.) In addition the mean values split according to individual weed species are also shown. As can be seen, Sample 13 was more efficacious than the standard adjuvant TurbochargeTM across the weed species, and was as good or better on individual weeds.

5

Table 4

Sample	All weeds	XANST	CHEAL	SETVI	ABUTH
Sample 13	61.8; A	89.7; A	89.4; A	27.5; A	40.55; A
Turbocharge	58.8; B	83.9; B	88.6; A	24.4; B	38.33; A

EXAMPLE 4

In this example Sample 13 from Table 1 was compared to the adjuvant BrijTM96V. The herbicide fomesafen was applied to the weed species xanthium strumarium (XANST), setaria viridis (SETVI), abutilon theophrasti (ABUTH), and chenopodium album (CHEAL) at rates of 60 or 120 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 7, 14 and 21 days after application. Each adjuvant was applied at a rate of 0.2% of the volume of the spray water.

15 The same statistical methodology that was applied in Example 3 was used here.

Table 5 shows the mean efficacy of the butylene oxide adjuvant number 13 with the pesticide fomesafen compared to the adjuvant Brij 96 across four weed species and compared to single weed species. The mean values and a letter denoting which group each adjuvant belonged to are shown. In addition the mean values split according to individual weed species are also shown. As can be seen the butylene oxide adjuvant was more efficacious than the standard adjuvant across the weed species, and was as good or better on individual weeds.

Table 5

Sample	All weeds	ABUTH	CHEAL	SETVI	XANST
Sample 13	57.15; A	32.5; A	85.27; A	24.17; A	86.66; A
Brij 96V	53.33; B	32.2; A	82.5; A	26.66; A	71.94; B

EXAMPLE 5

In this example the rate response of the butylene oxide adjuvant number 13 of Table 1 was measured to display the excellent performance of this adjuvant at very low rates. compared to the adjuvant BrijTM96V. The herbicide fomesafen was applied to the weed species xanthium strumarium (XANST), setaria viridis (SETVI), abutilon theophrasti (ABUTH), and chenopodium album (CHEAL) at rates of 60 or 120 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 7, 14 and 21 days after application. The adjuvants were each applied at a rate of 0.2% of the volume of the spray water.

The same statistical methodology that was applied in Example 3 was used here. Where a sample has more than one letter it is not significantly different to any other sample with one of those letters.

Table 6 shows the mean efficacy of the butylene oxide adjuvant Sample 13 at several addition rates with the pesticide fomesafen compared to the adjuvant Brij 96V across four weed species, the mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen the butylene oxide adjuvant was more efficacious than the standard adjuvant across the weed species. In addition it was as effective at half the rate of the standard Brij96V (Sample 13 at 0.1% cf. Brij 96V at 0.2%). The level of addition of the butylene oxide adjuvant that was statistically better than no adjuvant was 0.025%. This is a very low level of adjuvant addition, indicating the remarkable efficacy of this adjuvant.

Table 6

Sample	Adjuvant rate (%)	Rank	Mean % kill
Sample 13	0.5	A	61.805
Sample 13	0.2	B	57.15
Sample 13	0.1	BC	54.23
Brij 96	0.2	C	53.33
Sample 13	0.05	D	43.95
Sample 13	0.025	E	38.12
No adjuvant	0	F	27.57

EXAMPLE 6

In this example the butylene oxide adjuvant Sample 13 of Table 1 was compared to the adjuvant TweenTM 20. The herbicide mesotrione was applied at rates of 45 and 90 grams per hectare by a laboratory track sprayer to the weed species *brachiaria platyphyla* (BRAPL), *digitaria sanguinalis* (DIGSA), *polygonum convolvulus* (POLCO) and *amaranthus tuberculatus* (AMATU). In each case the experiments were replicated four times and the percentage damage to the weeds was assessed visually at time periods of 7, 14 and 21 days after application. The adjuvants was applied at a rate of 0.5% of the volume of the spray water.

The same statistical methodology that was applied in Example 3 was used here.

Table 7 shows the mean efficacy of the butylene oxide adjuvant Sample 13 at a rate of 0.5 % v/v with the pesticide mesotrione compared to the adjuvant Tween 20 across four weed species and compared to single weed species. The mean values and a letter denoting which group each adjuvant belonged to are shown. In addition the mean values split according to individual weed species are also shown. As can be seen the butylene oxide adjuvant was more efficacious than the standard adjuvant across the weed species, and was as good or better on individual weeds.

Table 7

Sample	All weeds	BRAPL	DIGSA	POLCO	AMATU
Sample 13	69.986; A	54.72; A	56.94; A	97.167; A	71.11; A
Tween 20	63.861; B	44.72; B	48.33; B	95.44; A	66.94; AB
No adjuvant	44.986; C	20.27; C	19.44; C	77.16; B	63.05; B

EXAMPLE 7

In this example the butylene oxide adjuvant Sample 13 of Table 1 was compared to the adjuvant BrijTM96V. The herbicide mesotrione was applied to the weed species brachiaria platyphyla (BRAPL), digitaria sanguinalis (DIGSA), polygonum convolvulus (POLCO) and amaranthus tuberculatus (AMATU) at rates of 45 or 90 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 7, 14 and 21 days after application. The adjuvants were applied at a rate of 0.2% of the volume of the spray water.

The same statistical methodology that was applied in Example 3 was used here.

Table 8 shows the mean efficacy of the butylene oxide adjuvant Sample 13 at a rate of 0.2% v/v with the pesticide mesotrione compared to the adjuvant Brij 96V across four weed species, and compared to single weed species. The mean values and a letter denoting which group each adjuvant belonged to are shown. In addition the mean values split according to individual weed species are also shown. As can be seen the butylene oxide adjuvant was more efficacious than the standard adjuvant across the weed species, and was as good or better on individual weeds.

Table 8

Sample	All weeds	BRAPL	DIGSA	POLCO	AMATU
Sample 13	68.44; A	51.66; A	54.72; A	97.11; A	70.28; A
Brij 96V	61.4; B	41.94; B	44.72; B	93.66; B	65.28; B
No adjuvant	44.99; C	20.27; C	19.44; C	77.16; C	63.05; B

EXAMPLE 8

In this example the rate response of the butylene oxide adjuvant Sample 13 of Table 1 was measured to display the excellent performance of this adjuvant at very low rates compared to the adjuvant BrijTM96V. The herbicide mesotrione was applied to the weed species brachiaria platyphyla (BRAPL), digitaria sanguinalis (DIGSA), polygonum convolvulus (POLCO) and amaranthus tuberculatus (AMATU) at rates of 45 or 90 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 7, 14 and 21 days after application. The butylene oxide adjuvant was applied at rates of 0.025, 0.05,

0.1, 0.2 and 0.5% w/v of the spray water. For comparison Brij96V was added at a rate of 0.2% w/v.

The same statistical methodology that was applied in Example 3 was used here.

- Table 9 shows the mean efficacy of the butylene oxide adjuvant Sample 13 at several addition rates with the pesticide mesotrione compared to the adjuvant Brij 96V across four weed species. The mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen half the rate of the butylene oxide adjuvant was more efficacious than the standard adjuvant (0.1 % vs 0.2 %). In addition it was as effective at one quarter of the rate of the standard Brij96V (0.05 % vs 0.2%).
- The level of addition of the butylene oxide adjuvant that was statistically better than no adjuvant was 0.025 %. This is a very low level of adjuvant addition indicating the remarkable efficacy of this adjuvant.

Table 9

Sample	Adjuvant rate %	Rank	Mean % kill	Standard error
Sample 13	0.5	A	70.0	1.8737
Sample 13	0.2	AB	68.4	
Sample 13	0.1	BC	66.2	
Sample 13	0.05	CD	63.0	
Brij 96	0.2	D	61.4	
Sample 13	0.025	D	59.5	
No adjuvant	0	E	45.0	

15

EXAMPLE 9

- In this example the rate response of the butylene oxide adjuvant Sample 13 of Table 1 was measured to display the excellent performance of this adjuvant at very low rates. The herbicide pinoxaden was applied to the weed species *lolium perenne* (LOLPE), *alopecurus myosuroides* (ALOMY), *setaria viridis* (SETVI) and *avena fatua* (AVEFA) at rates of 7.5 or 15 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application. The butylene oxide adjuvant was applied at rates of 0.025, 0.05, 0.1, 0.2 and 0.5% w/v of the spray water.

The same statistical methodology that was applied in Example 3 was used here.

Table 10 shows the mean efficacy of the butylene oxide adjuvant Sample 13 at several addition rates with the pesticide pinoxaden. Results are meaned across four weed species. The mean values and a letter denoting which group each adjuvant belonged to are shown. The results show that the butylene oxide adjuvant is efficacious at a very low level (0.025%) and that there is a strong rate response to the added adjuvant.

Table 10

Sample	Adjuvant rate %	Rank	Mean % kill	Standard error
Sample 13	0.5	A	76.89	1.441
Sample 13	0.2	B	73.89	
Sample 13	0.1	B	73.45	
Sample 13	0.05	C	67.53	
Sample 13	0.025	D	60.76	
No adjuvant	0	E	4.72	

EXAMPLE 10

In this example the butylene oxide adjuvant Sample 13 of Table 1 was compared to the commercial oil adjuvant blend Atplus™ 411F. The herbicide nicosulfuron was applied to the weed species chenopodium album (CHEAL), digitaria sanguinalis (DIGSA), setaria viridis (SETVI) and abutilon theophrasti (ABUTH) at rates of 30 or 60 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application. The adjuvants were applied at a rate of 0.5% of the volume of the spray water.

The same statistical methodology that was applied in Example 3 was used here.

Table 11 shows the mean efficacy of the butylene oxide adjuvant Sample 13 at a rate of 0.5%v/v with the pesticide nicosulfuron compared to the oil adjuvant Atplus™ 411F across four weed species and compared to single weed species. The mean values and a letter denoting which group each adjuvant belonged to are shown. In addition the mean values split according to individual weed species are also shown. As can be seen the butylene oxide adjuvant was more efficacious than the standard adjuvant across the weed species, and was as good or better on individual weeds.

Table 11

Sample	All weeds	CHEAL	SETVI	DIGSA	ABUTH
Sample 13	76.15; A	84.58; A	90; A	81.25; A	48.75; A
Atplus 411F	66.75; B	74.17; B	81.58; B	78.75; A	32.5; B
No adjuvant	20.94; C	12.08; C	41.67; C	8.75; B	21.25; C

EXAMPLE 11

In this example the butylene oxide adjuvant Sample 13 of Table 1 was compared to the commercial adjuvant tris 2-ethylhexyl phosphate (TEHP). The herbicide nicosulfuron was applied to the weed species chenopodium album (CHEAL), digitaria sanguinalis (DIGSA), setaria viridis (SETVI) and abutilon theophrasti (ABUTH) at rates of 30 or 60 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application. The adjuvants were applied at a rate of 0.2% of the volume of the spray water.

The same statistical methodology that was applied in Example 3 was used here.

Table 12 shows the mean efficacy of the butylene oxide adjuvant Sample 13 at a rate of 0.2% v/v with the pesticide nicosulfuron compared to the adjuvant TEHP the same rate across four weed species, and compared to single weed species. The mean values and a letter denoting which group each adjuvant belonged to are shown along with the standard error. In addition the mean values split according to individual weed species are also shown. As can be seen the butylene oxide adjuvant was more efficacious than the standard adjuvant across the weed species, and was as good or better on individual weeds.

Table 12

Sample	All weeds	CHEAL	SETVI	DIGSA	ABUTH
Sample 13	68.23; A	76.67; A	80; A	73.33; A	42.92; A
TEHP	58.12; B	68.33; B	77.08; A	69.17; A	17.92; B
No adjuvant	20.94; C	12.08; C	41.67; B	8.75; B	21.25; B

EXAMPLE 12

In this example the rate response of the butylene oxide adjuvant Sample 13 of Table 1 was measured and compared to the adjuvant TEHP. The herbicide nicosulfuron was applied to the weed species chenopodium album (CHEAL), digitaria sanguinalis (DIGSA), setaria viridis (SETVI) and abutilon theophrasti (ABUTH) at rates of 30 to 60 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application. The butylene oxide adjuvant was applied at rates of 0.025, 0.05, 0.1, 0.2 and 0.5% w/v of the spray water. For comparison TEHP was added at a rate of 0.2% w/v.

The same statistical methodology that was applied in Example 3 was used here.

Table 13 shows the mean efficacy of the butylene oxide adjuvant Sample 13 at several addition rates with the pesticide nicosulfuron. Results are meaned across four weed species. As a comparison the adjuvant TEHP was added at 0.2% v/v. The mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen half the rate of the butylene oxide adjuvant was more efficacious than the standard adjuvant (Sample 13 at 0.1% compared to TEHP at 0.2%). In addition it was as effective at one quarter of the rate of the standard TEHP (0.05% compared to 0.2%). The level of addition of the butylene oxide adjuvant that was statistically better than no adjuvant was 0.025%. This is a very low level of adjuvant addition, indicating the remarkable efficacy of this adjuvant.

Table 13

Sample	Adjuvant rate %	Rank	Mean % kill	Standard error
Sample 13	0.5	A	76.14	1.803
Sample 13	0.2	B	68.23	
Sample 13	0.1	C	63.23	
TEHP	0.2	D	58.12	
Sample 13	0.05	D	56.35	
Sample 13	0.025	E	51.87	
No adjuvant	0	F	20.93	

EXAMPLE 13

In this example the responses of the butylene oxide adjuvants 13, 17, 19, 21, 22 and 23 of Table 1 were measured at an application rate of 0.2% of the volume of the spray solution used. They were compared to the commercial tank mix adjuvant Atplus™411F, which was applied at the recommended rate of 0.5% by volume. The herbicide nicosulfuron was applied to the weed species chenopodium album (CHEAL), digitaria sanguinalis (DIGSA), setaria viridis (SETVI) and abutilon theophrasti (ABUTH) at rates of 30 or 60 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 14 shows the mean efficacy of six butylene oxide adjuvants at an addition rate of 0.2% with the pesticide nicosulfuron. Results are meaned across two pesticide rates and each sample was replicated four times. As a comparison the adjuvant Atplus™411F was added at 0.5% v/v. The mean values and a letter denoting which group each adjuvant belonged to are shown. The data shows that the butylene oxide adjuvants were at least as good as the standard which was used at a higher rate than the butylene oxide adjuvants (0.2% compared to 0.5%).

Table 14

Sample	All weeds	ABUTH	CHEAL	DIGSA	SETVI
Sample 13	78.44; A	65.42; A	85; A	67.5; A	95.83; A
Sample 21	76.45; AB	66.25; A	85; A	60; AB	94.58; AB
Sample 22	76.35; AB	61.67; A	85.83; A	62.5; AB	95.42; A
Sample 19	75.81; AB	65; A	84.17; A	58.33; BC	95.75; A
Sample 23	74.16; AB	62.92; A	82.08; A	57.92; BC	93.75; AB
Atplus 411F	70.73; B	60.83; A	74.17; B	56.67; BC	91.25; B
Sample 17	69.37; B	58.75; A	76.67; B	50.83; C	91.25; B
No adjuvant	37.5; C	25.83; B	24.16; C	18.33; D	81.67; C

20

EXAMPLE 14

In this example the responses of the adjuvants 13, 17, 19, 21, 22 and 23 were measured at an application rate of 0.2% of the volume of the spray solution used. They were compared to the adjuvant tris 2-ethylhexyl phosphate (TEHP), which was applied at the rate of 0.5% by volume. The herbicide pinoxaden was applied at rates of 7.5 or 15

grams per hectare by a laboratory track sprayer to the weed species *loium perenne* (LOLPE), *alopecurius myosuroides* (ALOMY), *setaria viridis* (SETVI) and *avena fatua* (AVEFA). In each case the experiments were replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 15 shows the mean efficacy of six butylene oxide adjuvants at an addition rate of 0.2% with the pesticide pinoxaden. Results are meaned across two pesticide rates and each sample was replicated four times. As a comparison the adjuvant tris 2 ethylhexyl phosphate was added at 0.5% v/v. The mean values and a letter denoting which group each adjuvant belonged to are shown along with the standard error. The standard TEHP was used at a higher rate than the butylene oxide adjuvants (0.2% compared to 0.5%).

Table 15

Sample	All weeds	SETVI	LOLPE	AVEFA	ALOMY
Sample 23	76.98; A	78.75; ABC	78.33; AB	78.33; AB	72.5; A
Sample 13	76.98; A	83.33; A	77.5; AB	80.42; A	66.67; B
Sample 19	76.25; A	80; AB	77.08; AB	77.08; AB	70.83; AB
TEHP	75.63; A	77.5; ABC	82.5; A	73.75; AB	68.75; AB
Sample 22	73.96; AB	78.33; ABC	72.5; BC	75.83; AB	69.17; AB
Sample 21	72.19; AB	71.25; C	76.67; AB	71.66; B	69.17; AB
Sample 17	68.02; B	72.92; BC	68.75; C	73.33; AB	57.08; C
No adjuvant	15.62; C	17.08; D	7.08; D	23.33; C	15; D

15

EXAMPLE 15

In this example the rate response of the butylene oxide sample 27 of Table 1 was measured, displaying the excellent performance of this adjuvant compared to the commercial adjuvant Turbocharge™, used at a rate of 0.5% by volume. The herbicide fomesafen was applied to the weed species *xanthium strumarium* (XANST), *setaria viridis* (SETVI), *abutilon theophrasti* (ABUTH), and *chenopodium album* (CHEAL) at rates of 60 or 120 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 7, 14 and 21 days after application. Sample 27 was applied at a rate of 0.2% of the volume of the spray water.

The same statistical methodology that was applied in Example 3 was used here. Table 16 shows the mean efficacy of the butylene oxide adjuvant 27 at a rate of 0.2% v/v with the pesticide fomesafen compared to the adjuvant Turbocharge at a rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown along with the standard error. As can be seen the butylene oxide adjuvant was as effective as Turbocharge even although it was applied at 0.2% as opposed to 0.5% for the commercial adjuvant.

Table 16

Sample	All weeds	XANST	SETVI	CHEAL	ABUTH
Sample 27	51.01; A	73.44; A	33.44; A	74.3; A	22.86; B
Turbocharge	52.89; A	72.33; A	28.72; B	75.69; A	34.81; A
No adjuvant	22.94; B	31.55; B	13.78; C	30.69; B	15.75; C

EXAMPLE 16

In this example the rate response of the butylene oxide adjuvant 27 of Table 1 was measured to display the excellent performance of this adjuvant compared to the commercial adjuvant TweenTM20, used at a rate of 0.5% by volume. The herbicide mesotrione was applied to the weed species brachiaria platyphyla (BRAPL), digitaria sanguinalis (DIGSA), polygonum convolvulus (POLCO) and amaranthus tuberculatus (AMATU) at rates of 45 or 90 grams per hectare by a laboratory track sprayer. Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 7, 14 and 21 days after application. Sample 27 was applied at a rate of 0.2% of the volume of the spray water.

The same statistical methodology that was applied in Example 3 was used here.

Table 17 shows the mean efficacy of the butylene oxide adjuvant 27 at a rate of 0.2% v/v with the pesticide mesotrione compared to the adjuvant Tween 20 at a rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen the butylene oxide adjuvant was as effective as Tween 20 even although it was applied at 0.2% as opposed to 0.5% for the commercial adjuvant.

Table 17

Sample	All weeds	POLCO	DIGSA	BRAPL	AMATU
Sample 27	57.26; A	75.44; A	44.99; A	36.94; A	71.64; A
Tween 20	54.54; A	73.61; A	41.67; A	33.06; B	69.86; A
No adjuvant	34.84; B	55.5; B	20.55; B	19.17; C	44.13; B

EXAMPLE 17

In this example the response of butylene oxide sample 27 of Table 1 was measured at an application rate of 0.2% of the volume of the spray solution used. It was compared to the adjuvant GenopolTMX080, which was also applied at a rate of 0.2% by volume. This adjuvant has the same alkyl chain and ethylene oxide head group as Sample 27 however it does not contain the butylene oxide moiety. The herbicide pinoxaden was applied at rates of 7.5 or 15 grams per hectare by a laboratory track sprayer to the weed species *loium perenne* (LOLPE), *alopecurus myosuroides* (ALOMY), *setaria viridis* (SETVI) and *avena fatua* (AVEFA). Each experiment was replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 18 shows the mean efficacy of the butylene oxide adjuvant 27 at a rate of 0.2% v/v with the pesticide pinoxaden compared to the adjuvant Genapol X080 at the same rate. The mean values and a letter denoting which group each adjuvant belonged to are shown. The data shows that the butylene oxide adjuvants was significantly more efficacious than Genapol X080 across the range of weed species tested.

Table 18

Sample	All weeds	SETVI	LOLPE	AVEFA	ALOMY
Sample 27	71.06; A	84.4; A	59.22; A	82.3; A	58.31; A
Genopol X080	52.08; B	70.67; B	20.97; B	75.47; B	41.23; B
No adjuvant	7.44; C	4.75; C	6.53; C	5.74; C	12.73; C

EXAMPLE 18

In this example the response of butylene oxide sample 27 of Table 1 was measured at an application rate of 0.2% by volume of the spray solution used. It was

compared to the commercial tank mix adjuvant Atplus™411F, which was applied at the recommended rate of 0.5% by volume, and to the adjuvant Genapol™X080 which was applied at 0.2%. This adjuvant has the same alkyl chain and ethylene oxide head group as sample 27 however it does not contain the butylene oxide moiety. The herbicide nicosulfuron was applied at rates of 30 or 60 grams per hectare by a laboratory track sprayer to the weed species chenopodium album (CHEAL), digitaria sanguinalis (DIGSA), setaria viridis (SETVI) and abutilon theophrasti (ABUTH). In each case the experiments were replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 19 shows the mean efficacy of the butylene oxide adjuvant 27 at a rate of 0.2 % v/v with the pesticide nicosulfuron compared to the adjuvant Genapol X080 at the same rate, and to Atplus 411F applied at the higher rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. The data shows that the butylene oxide adjuvants was as good as Atplus 411F which was used at a higher rate than the butylene oxide adjuvants (0.2% compared to 0.5%). It was more efficacious than Genapol X080.

Table 19

Sample	All weeds	SETVI	DIGSA	CHEAL	ABUTH
Sample 27	70.26; A	45.13; A	79.55; A	84.58; A	71.76; A
Atplus 411F	65.74; A	25.96; B	76.05; A	85.01; A	75.93; A
Genapol X080	46.47; B	19.72; C	42.71; B	70.42; B	53.01; B
No adjuvant	21.39; C	14.17; D	22.59; C	15.2; C	33.6; C

20

EXAMPLE 19

In this example the rate response of the butylene oxide adjuvant 13 of Table 1 was measured to display the excellent performance of this adjuvant compared to the commercial adjuvant Turbocharge™, used at a rate of 0.5% by volume. The herbicide fomesafen was applied at rates of 60, 90 or 120 grams per hectare by a laboratory track sprayer to the weed species xanthium strumarium (XANST), abutilon theophrasti (ABUTH), and chenopodium album (CHEAL). In each case the experiments were replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application. Sample 13 was applied at a rate of 0.2% of the volume of the spray water.

The same statistical methodology that was applied in Example 3 was used here.

Table 21 shows the mean efficacy of the butylene oxide adjuvant 13 at a rate of 0.2% v/v with the pesticide fomesafen compared to the adjuvant Turbocharge applied at the higher rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen the butylene oxide adjuvant was more effective than Turbocharge even although it was applied at 0.2% as opposed to 0.5% for the commercial adjuvant.

Table 20

Sample	All weeds	XANST	CHEAL	ABUTH
Sample 13	51.24; A	92.56; A	62.5; A	41.39; A
Turbocharge	39.43; B	77.44; B	40; B	35; B
No adjuvant	23.61; C	45; C	24.44; C	25; C

10

EXAMPLE 20

In this example the rate response of the butylene oxide adjuvant 13 of Table 1 was measured to display the excellent performance of this adjuvant compared to the commercial adjuvant TweenTM20, used at a rate of 0.5% by volume. The herbicide mesotrione was applied at rates of 30, 60 or 90 grams per hectare by a laboratory track sprayer to the weed species brachiaria platyphyla (BRAPL), digitaria sanguinalis (DIGSA), and polygonum convolvulus (POLCO). Sample 13 was applied at a rate of 0.2% of the volume of the spray water. In each case the experiments were replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

20 The same statistical methodology that was applied in Example 3 was used here.

Table 21 shows the mean efficacy of the butylene oxide adjuvant 13 at a rate of 0.2% v/v with the pesticide mesotrione compared to the adjuvant Tween 20 applied at the higher rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen the butylene oxide adjuvant was more effective than Tween 20 even though it was applied at 0.2% as opposed to 0.5 % for the commercial adjuvant.

Table 21

Sample	All weeds	POLCO	DIGSA	BRAPL
Sample 13	28.89; A	36.11; A	47.5; A	39.14; A
Tween 20	26.11; B	32.77; B	45.55; A	26.11; B
No adjuvant	13.61; C	27.78; C	17.5; B	9.17; C

EXAMPLE 21

In this example the response of the butylene oxide adjuvant 13 of Table 1 was measured at an application rate of 0.2% of the volume of the spray solution used. It was compared to the commercial tank mix adjuvant Atplus™411F, which was applied at the recommended rate of 0.5% by volume. The herbicide nicosufuron was applied at rates of 30, 45 or 60 grams per hectare by a laboratory track sprayer to the weed species chenopodium album (CHEAL), digitaria sanguinalis (DIGSA), and abutilon theophrasti (ABUTH). In each case the experiments were replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 22 shows the mean efficacy of the butylene oxide adjuvant 13 at a rate of 0.2% v/v with the pesticide nicosulfuron compared to the adjuvant Atplus 411F applied at the higher rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. The data shows that the butylene oxide adjuvant was as good as Atplus 411F which was used at a higher rate than the butylene oxide adjuvant (0.5% compared to 0.2%).

Table 22

Sample	All weeds	DIGSA	CHEAL	ABUTH
Sample 13	67.31; A	78.61; A	88.33; A	35; A
Atplus 411F	63.61; A	75.55; B	88.06; A	27.22; B
No adjuvant	8.11; B	4.05; C	0.56; B	19.72; C

EXAMPLE 22

In this example the response of the butylene oxide adjuvant 13 of Table 1 was measured at an application rate of 0.2% of the volume of the spray solution used. It was compared to the adjuvant tris 2-ethylhexyl phosphate [TEHP], which was applied at the

higher rate of 0.5% by volume. The herbicide pinoxaden was applied at rates of 7.5, 11.25 or 15 grams per hectare by a laboratory track sprayer to the weed species *lolium perenne* (LOLPE), *alopecurus myosuroides* (ALOMY), and *avena fatua* (AVEFA). In each case the experiments were replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 23 shows the mean efficacy of butylene oxide adjuvant 13 at a rate of 0.2% v/v with the pesticide pinoxaden compared to the adjuvant tris 2-ethylhexyl phosphate applied at the higher rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. The data shows that the butylene oxide adjuvant was as effective as TEHP across the range of weed species tested.

Table 23

Sample	All weeds	LOLPE	AVEFA	ALOMY
Sample 13	58.05; A	53.89; B	76.94; A	43.33; A
TEHP	59.91; A	58.06; A	78.33; A	43.33; A
No adjuvant	3.43; B	2.22; C	6.39; B	1.67; B

15

EXAMPLE 23

In this example the rate response of 22 butylene oxide adjuvants of Table 1 were measured at an adjuvant rate of 0.2% by volume, displaying excellent performance of this adjuvant compared to the commercial adjuvant Turbocharge, which was used at a rate of 0.5% by volume. The herbicide mesotrione was applied at rates of 60 or 120 grams per hectare by a laboratory track sprayer to the weed species *brachiaria platyphylla* (BRAPP), *digitaria sanguinalis* (DIGSA), *abutilon theophrasti* (ABUTH) and *amaranthus retroflexus* (AMARE). In each case the experiments were replicated four times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application. All the butylene oxide adjuvants were applied at a rate of 0.2% volume of the spray water whereas Turbocharge was used at 0.5%.

The same statistical methodology that was applied in Example 3 was used here.

Table 23 shows the mean efficacy of 22 butylene oxide adjuvants used at a rate of 0.2% v/v with the pesticide mesotrione compared to the adjuvant Turbocharge applied at the higher rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen most of the butylene oxide adjuvants

were more effective than Turbocharge even although they were applied at 0.2% as opposed to 0.5% for the commercial adjuvant. All of the butylene adjuvants were as good as Turbocharge.

Table 23

Sample	Group	Mean kill across weeds %
3	A	84.69
27	AB	84.27
26	ABC	83.85
21	ABCD	83.65
13	ABCDE	83.44
20	ABCDE	83.44
23	ABCDE	83.44
19	ABCDEF	83.13
7	ABCDEF	83.02
2	ABCDEF	82.60
6	ABCDEF	82.60
1	BCDEFG	81.98
22	CDEFGH	81.88
18	CDEFGH	81.77
25	CDEFGH	81.67
10	DEFGHI	81.35
17	EFGHI	81.25
11	FGHI	80.94
8	GHIJ	80.21
24	HIJ	79.58
16	IJ	79.27
Turbocharge	J	78.44
5	J	78.13
No adjuvant	K	69.48

EXAMPLE 24

In this example the responses of the butylene oxide adjuvants 7, 14, 24 and 25 of Table 1 were measured at an application rate of 0.2% of the volume of the spray solution used. They were compared to the adjuvant BrijTM96V, which was applied at the same rate. The herbicide pinoxaden was applied to the weed species *lolium perenne* (LOLPE), *alopecurus myosuroides* (ALOMY), *setaria viridis* (SETVI) and *avena fatua* (AVEFA) at rates of 7.5 or 15 grams per hectare by a laboratory track sprayer. Each experiment was replicated three times and the percentage damage to the weeds was assessed visually 13 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 25 shows the mean efficacy of 4 butylene oxide adjuvants used at a rate of 0.2% v/v with the pesticide pinoxaden compared to the adjuvant Brij 96V applied at the same rate. The mean values and a letter denoting which group each adjuvant belonged to are shown. The data shows that two of the butylene oxide adjuvants were more effective than Brij 96V and two were as good as Brij 96V across the range of weed species tested.

Table 25

Sample	Group	Mean kill % all weeds
7	A	72.3
14	A	67.3
25	B	57.3
Brij 96V	B	55.0
24	B	52.9
No adjuvant	C	16.3

EXAMPLE 25

In this example the rate responses of four butylene oxide adjuvants of Table 1, applied at 0.2%, displayed excellent performance when compared to the commercial adjuvant TurbochargeTM, used at a rate of 0.5%; and to tris 2-ethylhexyl phosphate (TEHP) also used at 0.5%. The herbicide fomesafen was applied to the weed species *xanthium strumarium* (XANST), *abutilon theophrasti* (ABUTH), *setaria viridis* (SETVI) and *chenopodium album* (CHEAL) at rates of 60 or 120 grams per hectare by a laboratory track sprayer. Each experiment was replicated six times and the percentage damage to the weeds was assessed visually at time periods of 7 and 13 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 26 shows the mean efficacy of 4 butylene oxide adjuvants used at a rate of 0.2% v/v with the pesticide fomesafen compared to the adjuvants tris 2-ethylhexyl phosphate (TEHP) and Turbocharge. The latter two adjuvants were applied at the higher rate of 0.5% v/v. The mean values and a letter denoting which group each adjuvant belonged to are shown along with the standard error. As can be seen the performance of all four of the butylene oxide adjuvants was at least as good as the standard Turbocharge and most of the adjuvants were as good as or better than TEHP.

10

Table 26

Sample	Group	Mean kill across weeds (%)
24	A	78.2
7	B	73.6
25	B	73.1
TEHP	B	72.7
Turbocharge	C	68.7
14	C	66.1
None	D	47.0

EXAMPLE 26

In this example the rate response of sample 28 of Table 1, a non-ethoxylated adjuvant, applied at 0.2% by volume was measured to display the performance of the adjuvant compared to the commercial adjuvant Turbocharge, used at a rate of 0.5%. The herbicide fomesafen was applied to the weed species xanthium strumarium (XANST), abutilon theophrasti (ABUTH), setaria viridis (SETVI) and chenopodium album (CHEAL) at rates of 60 or 120 grams per hectare by a laboratory track sprayer. Each experiment was replicated three times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 27 shows the mean efficacy of a non-ethoxylated butylene oxide adjuvant used at a rate of 0.2% v/v with the pesticide fomesafen compared to the adjuvant Turbocharge. Turbocharge was applied at the higher rate of 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen the performance of the butylene oxide adjuvant was as good as the standard Turbocharge.

Table 27

Sample	Group	Mean kill (%)
Turbocharge	A	67.6
Sample 28	A	64.7
No adjuvant	B	37.7

EXAMPLE 27

5 In this example the responses of sample 28 of table 1, a non-ethoxylated butylene oxide adjuvant was compared to tris 2-ethylhexyl phosphate (TEHP). The butylene oxide adjuvant was applied at an application rate of 0.2% by volume of the spray solution used whereas TEHP was applied at 0.5%. The herbicide pinoxaden was applied to the weed species *lolium perenne* (LOLPE), *alopecurus myosuroides* (ALOMY), *setaria* 10 *viridis* (SETVI) and *avena fatua* (AVEFA) at rates of 7.5 or 15 grams per hectare by a laboratory track sprayer. Each experiment was replicated three times and the percentage damage to the weeds was assessed visually 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 28 shows the mean efficacy of a non-ethoxylated butylene oxide adjuvant 15 compared to TEHP with the pesticide pinoxaden. The rate of the former was 0.2% whereas the latter was applied at 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. The data shows that the butylene oxide adjuvant was as effective as TEHP across the range of weed species tested.

Table 28

Sample	Group	Mean kill (%)
TEHP	A	92.6
Sample 28	A	91.0
No adjuvant	B	17.4

EXAMPLE 28

25 In this example the rate response of a non-ethoxylated butylene oxide adjuvant was measured at an adjuvant rate of 0.2% by volume in order to display the excellent

performance of this adjuvant compared to the commercial adjuvant' Tween 20, which was used at a rate of 0.5% by volume. The herbicide mesotrione was applied at rates of 45 or 90 grams per hectare by a laboratory track sprayer to the weed species brachiaria–decumbens (BRADE), digitaria sanguinalis (DIGSA), polygonum convolvulus (POLCO) and amaranthus retroflexus (AMARE). In each case the experiments were replicated three times and the percentage damage to the weeds was assessed visually at time periods of 7, 14 and 21 days after application. The butylene oxide adjuvant was applied at a rate of 0.2% of the volume of the spray water whereas Tween 20 was used at 0.5%.

The same statistical methodology that was applied in Example 3 was used here. Table 29 shows the mean efficacy of an unethoxylated butylene oxide adjuvant compared to Tween 20 with the pesticide mesotrione. The rate of the former was 0.2% whereas the latter was applied at 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown. As can be seen the butylene oxide adjuvant was as effective as Tween 20 even though it was applied at 0.2% as opposed to 0.5% for the commercial adjuvant.

Table 29

Sample	Group	Mean kill %
Sample 28	A	68.8
Tween 20	A	65.8
No adjuvant	B	47.2

EXAMPLE 29

In this example the response of sample 28 of Table 1, a non-ethoxylated butylene oxide adjuvant was measured at an application rate of 0.2% of the volume of the spray solution used. It was compared to the commercial tank mix adjuvant Atplus™411F, which was applied at the recommended rate of 0.5% by volume. The herbicide nicosufuron was applied at rates of 30 or 60 grams per hectare by a laboratory track sprayer to the weed species chenopodium album (CHEAL), digitaria sanguinalis (DIGSA), setaria viridis (SETVI) and abutilon theophrasti (ABUTH). In each case the experiments were replicated three times and the percentage damage to the weeds was assessed visually at time periods of 14 and 21 days after application.

The same statistical methodology that was applied in Example 3 was used here.

Table 30 shows the mean efficacy of a non-ethoxylated butylene oxide adjuvant compared to Atplus 411F with the pesticide nicosulfuron. The rate of the former was 0.2% whereas the latter was applied at 0.5%. The mean values and a letter denoting which group each adjuvant belonged to are shown along with the standard error. The data shows that the butylene oxide adjuvant was as good as Atplus 411F which was used at a higher rate than the butylene oxide adjuvants (0.2% compared to 0.5%).

Table 30

Sample	Group	Mean kill %
ATPLUS 411 F	A	90.4
Sample 28	A	89.2
No adjuvant	B	75.0

10

EXAMPLE 30

This example shows that samples 29 and 30 from Table 1 behave as adjuvants for fungicides used against the fungus septoria tritici. Wheat was sprayed with water at a rate of 200 litres per hectare, the water containing either isopyrazam or epoxyconazole, at a concentration which enabled a pesticide application rate of 0.6, 2, 6 or 20 grams per hectare. Sample 29 was added at a rate of 0.2% v/v of the spray volume used and sample 30 was added at a rate of 0.1 % v/v. As a comparison to these adjuvants the same formulation was tested without the adjuvant, as a standard. Each experiment was replicated 12 times and the results were averaged at each rate. Plants were examined for curative effects. The percentage disease on each sample was assessed visually and averaged across the replicates at each rate. This was converted to a percentage control by comparison to the disease level on plants which were sprayed using a blank spray application where the pesticide was not included.

Table 31 shows the percentage septoria control for the two adjuvants used with the four levels of isopyrazam as well as the blank formulation. Table 32 shows the percentage septoria control for the two adjuvants used with the four levels of epoxyconazole as well as the blank formulation. In each case it can be seen that the adjuvants have improved the performance of the fungicide.

Table 31

Isopyrazam			
g/ha	Control	Sample 29	Sample 30
20	21	100	99
6	35	85	85
2	21	53	26
0.6	18	12	0

Table 32

Epoxyconazole		Sample	Sample
g/ha	Control	29	30
20	29	99	100
6	11	99	99
2	22	88	90
0.6	4	45	54

EXAMPLE 31

- 5 This is an insecticide example: it shows that sample 28 from Table 1 behaves as an adjuvant for the insecticide thiamethoxam against *Aphis craccivora*. The lower sides of French bean leaves were infested with an aphid population *Aphis craccivora* of mixed ages contained in clip cages. The upper sides of the leaves were sprayed with the test solutions, 1 day after aphid infestation. French bean was sprayed with water at a rate of
- 10 200 litres per hectare, the water containing 3, 6, 12.5 and 25 ppm thiamethoxam. Sample 28 was added at a rate of 0.1% v/v of the spray volume used. As a comparison to this adjuvant the same formulation was tested without the adjuvant, as a standard. 5 days after spray application, the aphids were checked visually for mortality. Each experiment
- 15 was replicated twice and the results were averaged at each rate. In the control experiment the beans were sprayed with water and no mortality was observed.

Table 33

Treatment	3ppm thiamethoxam % mortality	6ppm thiamethoxam % mortality	12.5 ppm thiamethoxam % mortality	25 ppm thiamethoxam % mortality
Actara WG25	0	70	99	100
Actara WG25 + 0.1% v/v sample 28	50	90	97.5	100

ActaraTM WG25 is a commercial product containing thiamethoxam

5

EXAMPLE 32

This is a Phytotoxicity example: it shows that sample 28 from Table 1 is not phytotoxic to soybean, French bean and Chinese cabbage. The plants were sprayed with water at a rate of approximately 500 litres per hectare, the water containing 0.1% v/v or 0.2% v/v adjuvant. The plants were assessed for phytotoxicity 7 days after spray application.

- 10 Each experiment was replicated twice and the results averaged. In the control experiment the plants were sprayed with water and no phytotoxicity was observed. The results show that the adjuvant sample 28 is safer to the crops than the alcohol ethoxylate adjuvant Genapol O100.

Table 34


	Soybean % phytotoxicity	French bean % phytotoxicity	Chinese cabbage % phytotoxicity
0.1%v/v Sample 28	0	0	0
0.2%v/v Sample 28	1	0	0
0.1%v/v Genapol O100	3.5	5	2
0.2%v/v Genapol O100	15	10	10

15

GenapolTM O100 is a commercial surfactant from Clariant, an oleyl ethoxylate with 10 moles ethylene oxide.

WE CLAIM:

1. Use of a compound of formula (I) as a bioperformance enhancing adjuvant of a
 5 pesticide



where BO is CH(R₄)CHR₅O; n is from 1 to 12; R₁ is C₄₋₂₀ alkyl or C₄₋₂₀ alkenyl;
 10 R₂ is hydrogen or C₁₋₃ alkyl; R₄ is methyl and R₅ is methyl or R₄ is ethyl and R₅ is
 hydrogen or R₄ is hydrogen and R₅ is ethyl.

2. The use as claimed in claim 1 where R₁ is C₆₋₁₈ alkyl or C₆₋₁₈ alkenyl.
- 15 3. The use as claimed in claim 2 where R₁ is oleyl.
4. The use as claimed in claim 2 or 3 where R₂ is hydrogen or C₁₋₂ alkyl.
5. The use as claimed in any one of claims 2 to 4 where n is from 2 to 8.
- 20 6. The use as claimed in claim 2 where R₁ is C₁₂₋₁₅ alkyl; n is 4; and R₂ is hydrogen.
7. The use as claimed in claim 2 where R₁ is oleyl; n is 4; and R₂ is hydrogen.

25