

(43) International Publication Date  
23 September 2010 (23.09.2010)(10) International Publication Number  
**WO 2010/107733 A2**

## (51) International Patent Classification:

C12N 15/63 (2006.01) A61K 48/00 (2006.01)  
C12N 15/113 (2010.01) C12Q 1/68 (2006.01)  
C07H 21/00 (2006.01)

## (21) International Application Number:

PCT/US2010/027394

## (22) International Filing Date:

16 March 2010 (16.03.2010)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

61/160,369 16 March 2009 (16.03.2009) US  
61/181,062 26 May 2009 (26.05.2009) US

(71) Applicant (for all designated States except US): **CUR-NA, INC.** [US/US]; 555 Heritage Drive, Suite 110, Jupiter, FL 33458 (US).

## (72) Inventors; and

(75) Inventors/Applicants (for US only): **COLLARD, Joseph** [US/US]; 1004 Brooks Lane, Delray Beach, FL 33483 (US). **KHORKOVA SHERMAN, Olga** [US/US]; 18288 SE Heritage Drive, Tequesta, FL 33469 (US). **COITO, Carlos** [PT/US]; 4518 Corniche Circle, #26, West Palm Beach, FL 33417 (US).

(74) Agents: **ELAMRANI, Samir** et al.; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, CA 94304-1050 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: TREATMENT OF NUCLEAR FACTOR (ERYTHROID-DERIVED 2)-LIKE 2 (NRF2) RELATED DISEASES BY INHIBITION OF NATURAL ANTISENSE TRANSCRIPT TO NRF2

FIGURE 1A



(57) Abstract: The present invention relates to antisense oligonucleotides that modulate the expression of and/or function of Nuclear factor (erythroid-derived 2)-like 2 (NRF2), in particular, by targeting natural antisense polynucleotides of Nuclear factor (erythroid-derived 2)-like 2 (NRF2). The invention also relates to the identification of these antisense oligonucleotides and their use in treating diseases and disorders associated with the expression of NRF2.

**TREATMENT OF NUCLEAR FACTOR (ERYTHROID-DERIVED 2)-LIKE 2 (NRF2)  
RELATED DISEASES BY INHIBITION OF NATURAL ANTISENSE TRANSCRIPT  
TO NRF2**

**REFERENCE TO PRIOR APPLICATIONS**

[0001] The present application claims the priority of U.S. provisional patent application No. 61/160,369, entitled "Antisense Compounds for Nuclear Factors," filed March 16, 2009 and U.S. provisional patent application No. 61/181,062, entitled "Nuclear Factor (Erythroid-Derived 2)-Like 2 Antisense Oligonucleotides," filed May 26, 2009, both of which are  
5 incorporated herein by reference in their entirety.

**FIELD OF THE INVENTION**

[0002] Embodiments of the invention comprise oligonucleotides modulating expression and/or function of NRF2 and associated molecules.

**BACKGROUND**

10 [0003] DNA-RNA and RNA-RNA hybridization are important to many aspects of nucleic acid function including DNA replication, transcription, and translation. Hybridization is also central to a variety of technologies that either detect a particular nucleic acid or alter its expression. Antisense nucleotides, for example, disrupt gene expression by hybridizing to target RNA, thereby interfering with RNA splicing, transcription, translation, and replication.  
15 Antisense DNA has the added feature that DNA-RNA hybrids serve as a substrate for digestion by ribonuclease H, an activity that is present in most cell types. Antisense molecules can be delivered into cells, as is the case for oligodeoxynucleotides (ODNs), or they can be expressed from endogenous genes as RNA molecules. The FDA recently approved an antisense drug, VITRAVENE™ (for treatment of cytomegalovirus retinitis), reflecting that  
20 antisense has therapeutic utility.

**SUMMARY**

[0004] This Summary is provided to present a summary of the invention to briefly indicate the nature and substance of the invention. It is submitted with the understanding that it will not be used to interpret or limit the scope or meaning of the claims.

25 [0005] In one embodiment, the invention provides methods for inhibiting the action of a natural antisense transcript by using antisense oligonucleotide(s) targeted to any region of the

natural antisense transcript resulting in up-regulation of the corresponding sense gene. It is also contemplated herein that inhibition of the natural antisense transcript can be achieved by siRNA, ribozymes and small molecules, which are considered to be within the scope of the present invention.

5 [0006] One embodiment provides a method of modulating function and/or expression of an NRF2 polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising contacting said cells or tissues with an antisense oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to a reverse complement of a polynucleotide comprising 5 to 30 consecutive nucleotides within nucleotides 1 to 5808 of  
10 SEQ ID NO: 3 (Figure 3) thereby modulating function and/or expression of the NRF2 polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0007] In another preferred embodiment, an oligonucleotide targets a natural antisense sequence of NRF2 polynucleotides, for example, nucleotides set forth in SEQ ID NO: 3, and any variants, alleles, homologs, mutants, derivatives, fragments and complementary  
15 sequences thereto. Examples of antisense oligonucleotides are set forth as SEQ ID NOS: 4 to 6 (Figure 4).

[0008] Another embodiment provides a method of modulating function and/or expression of an NRF2 polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising contacting said cells or tissues with an antisense oligonucleotide 5 to 30 nucleotides in length wherein  
20 said oligonucleotide has at least 50% sequence identity to a reverse complement of the an antisense of the NRF2 polynucleotide; thereby modulating function and/or expression of the NRF2 polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0009] Another embodiment provides a method of modulating function and/or expression of an NRF2 polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising contacting  
25 said cells or tissues with an antisense oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to an antisense oligonucleotide to an NRF2 antisense polynucleotide; thereby modulating function and/or expression of the NRF2 polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0010] In a preferred embodiment, a composition comprises one or more antisense  
30 oligonucleotides which bind to sense and/or antisense NRF2 polynucleotides.

[0011] In another preferred embodiment, the oligonucleotides comprise one or more modified or substituted nucleotides.

[0012] In another preferred embodiment, the oligonucleotides comprise one or more modified bonds.

[0013] In yet another embodiment, the modified nucleotides comprise modified bases comprising phosphorothioate, methylphosphonate, peptide nucleic acids, 2'-O-methyl, fluoro- or carbon, methylene or other locked nucleic acid (LNA) molecules. Preferably, the modified nucleotides are locked nucleic acid molecules, including  $\alpha$ -L-LNA.

[0014] In another preferred embodiment, the oligonucleotides are administered to a patient subcutaneously, intramuscularly, intravenously or intraperitoneally.

[0015] In another preferred embodiment, the oligonucleotides are administered in a pharmaceutical composition. A treatment regimen comprises administering the antisense compounds at least once to patient; however, this treatment can be modified to include multiple doses over a period of time. The treatment can be combined with one or more other types of therapies.

[0016] In another preferred embodiment, the oligonucleotides are encapsulated in a liposome or attached to a carrier molecule (e.g. cholesterol, TAT peptide).

[0017] Other aspects are described *infra*.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0018] Figure 1:

Figure 1A is a graph of real time PCR results showing the fold change + standard deviation in NRF2 mRNA after treatment of HepG2 cells with phosphorothioate oligonucleotides introduced using Lipofectamine 2000, as compared to control. Real time PCR results show that the levels of NRF2 mRNA in HepG2 cells are significantly increased 48 h after treatment with two of the oligonucleotides designed to NRF2 antisense BU656954.1. Bars denoted as CUR-0330 and CUR-0332 correspond to samples treated with SEQ ID NOS: 4 and 5 respectively.

Figure 1B is a graph of real time PCR results showing the fold change + standard deviation in NRF2 mRNA after treatment of CHP212 cells with phosphorothioate oligonucleotides introduced using Lipofectamine 2000, as compared to control. Real time PCR results show that the levels of NRF2 mRNA in CHP212 cells are significantly increased 48 h after treatment with two of the oligonucleotides designed to NRF2 antisense BU656954.1. Bars

denoted as CUR-0330, CUR-0332 and CUR-0509 correspond to samples treated with SEQ ID NOS: 4, 5 and 6 respectively.

[0019] Figure 2 shows

SEQ ID NO: 1: Homo sapiens nuclear factor (erythroid-derived 2)-like 2 (NFE2L2),  
5 transcript variant 1, mRNA. (NCBI Accession No.: NM\_006164)

SEQ ID NO: 2: Genomic sequence of NRF2 (exons are shown in capital letters, introns in small).

[0020] Figure 3 shows SEQ ID NO: 3: Natural NRF2 antisense sequence (BU656954.1).

[0021] Figure 4 shows the antisense oligonucleotides, SEQ ID NOs: 4 to 6. \* indicates  
10 phosphothioate bond.

[0022] Figure 5 shows the sense oligonucleotides, SEQ ID NOs: 7 and 8. The sense oligonucleotides SEQ ID NO: 7 and 8 are the reverse complements of the antisense oligonucleotides SEQ ID NO: 4 and 5 respectively.

#### DETAILED DESCRIPTION

[0023] Several aspects of the invention are described below with reference to example  
15 applications for illustration. It should be understood that numerous specific details, relationships, and methods are set forth to provide a full understanding of the invention. One having ordinary skill in the relevant art, however, will readily recognize that the invention can be practiced without one or more of the specific details or with other methods. The present  
20 invention is not limited by the ordering of acts or events, as some acts may occur in different orders and/or concurrently with other acts or events. Furthermore, not all illustrated acts or events are required to implement a methodology in accordance with the present invention.

[0024] Oxygen is essential for humans but reactive oxygen species generated in the body as its by-product is highly toxic. It is known that cellular damages due to oxidative stress caused  
25 by reactive oxygen species are deeply involved in onset or progress of various age-related chronic diseases (e.g., arteriosclerosis, hypertension, diabetes, cerebral nerve degenerative disease, skin disease, eye disease, asthma, and cancer). Enhancing the capability in protecting from oxidative stress is useful in preventing these diseases or delaying their progress (*Biomed Pharmacother.* 57,251-60 (2003)).

[0025] A series of proteins called phase II enzymes collaborate in undertaking protection of  
30 the body from oxidative stress or detoxification of xenobiotic substances. Examples of known

phase II enzymes include glutathione S-transferase (GST), NAD(P)H: quinine oxidoreductase 1 (NQO1), glutaryl-cysteinyl ligase (GCL), heme oxygenase 1 (HO1), and thioredoxin reductase 1 (TXNRD1). A common sequence called antioxidant responsive element (ARE) is present in a promoter of each gene of these phase II enzymes, and its expression is induced by the transcription factor NRF2 (NF-E2 related factor 2) (*Annu Rev Pharmacol Toxicol*, 43:233-60 (2003)). It is generally believed that NRF2 is present in the cytoplasm as a Keap1-NRF2 complex and when an activator acts therein, it moves into the nucleus and forms a heterodimer with a small Maf protein to bind ARE, thereby augmenting the expression of phase II enzymes.

[0026] Since the brain has a relatively high metabolic rate and a low cell regenerative capacity, it is an organ vulnerable to damages due to oxidative stress as compared to the other organs. Further, since the amount of reactive oxidative species in the brain increases with age, it is suggested that the capability in protecting the brain with oxidative stress may decrease with age. The increase in oxidative stress is believed to be closely related to the onset of cerebral nerve diseases such as Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis. For e.g., accumulation of 4-hydroxynonenal or malondialdehyde, which is marker for lipid peroxides, has been observed in the cerebral cortex or hippocampus of Alzheimer's disease patients, the substantia nigra of Parkinson's disease patients, and the spinal fluid of amyotrophic lateral sclerosis patients. Further, in Alzheimer's disease,  $\beta$ -amyloid, which is believed to be a causative substance, increases the amount of intracellular hydrogen peroxide and that in Parkinson's disease, increasing dopamine is known to increase the amount of multiple reactive oxygen species molecular species including hydrogen peroxide. In Parkinson's disease, it has also been reported that the amount of glutathione in the *substantia nigra* makes a biochemical index for the disease and the amount of glutathione decreases with the aggravation of the disease (*Nat Med.* 10 Suppl. S18-25 (2004)). Accordingly, augmentation of the capability in the protection from reactive oxygen species, for e.g., in preventing cellular lesions due to hydrogen peroxide and in increasing the amount of intracellular glutathione is considered to be effective in preventing or ameliorating these cerebral neurodegenerative diseases. It has also been reported that dimethyl fumarate and tert-butylhydroquinone which induce phase II enzymes suppress cell death due to dopamine and hydrogen peroxide (*J.Biol Chem.* 277, 388-94 (2002); *J Neurochem.* 71, 69-77 (1998); *Neurosci Lett.*; 273, 109-12 (1999)).

[0027] All genes, gene names, and gene products disclosed herein are intended to correspond to homologs from any species for which the compositions and methods disclosed herein are applicable. Thus, the terms include, but are not limited to genes and gene products from humans and mice. It is understood that when a gene or gene product from a particular species is disclosed, this disclosure is intended to be exemplary only, and is not to be interpreted as a limitation unless the context in which it appears clearly indicates. Thus, for example, for the genes disclosed herein, which in some embodiments relate to mammalian nucleic acid and amino acid sequences are intended to encompass homologous and/or orthologous genes and gene products from other animals including, but not limited to other mammals, fish, amphibians, reptiles, and birds. In preferred embodiments, the genes or nucleic acid sequences are human.

### *Definitions*

[0028] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention. As used herein, the singular forms "a", "an" and "the" are intended to include the plural forms as well, unless the context clearly indicates otherwise. Furthermore, to the extent that the terms "including", "includes", "having", "has", "with", or variants thereof are used in either the detailed description and/or the claims, such terms are intended to be inclusive in a manner similar to the term "comprising."

[0029] The term "about" or "approximately" means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. For example, "about" can mean within 1 or more than 1 standard deviation, per the practice in the art. Alternatively, "about" can mean a range of up to 20%, preferably up to 10%, more preferably up to 5%, and more preferably still up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value. Where particular values are described in the application and claims, unless otherwise stated the term "about" meaning within an acceptable error range for the particular value should be assumed.

[0030] As used herein, the term "mRNA" means the presently known mRNA transcript(s) of a targeted gene, and any further transcripts which may be elucidated.

[0031] By "antisense oligonucleotides" or "antisense compound" is meant an RNA or DNA molecule that binds to another RNA or DNA (target RNA, DNA). For example, if it is an RNA oligonucleotide it binds to another RNA target by means of RNA-RNA interactions and alters the activity of the target RNA (Eguchi *et al.*, (1991) *Ann. Rev. Biochem.* 60, 631-652).

5 An antisense oligonucleotide can upregulate or downregulate expression and/or function of a particular polynucleotide. The definition is meant to include any foreign RNA or DNA molecule which is useful from a therapeutic, diagnostic, or other viewpoint. Such molecules include, for example, antisense RNA or DNA molecules, interference RNA (RNAi), micro RNA, decoy RNA molecules, siRNA, enzymatic RNA, therapeutic editing RNA and agonist  
10 and antagonist RNA, antisense oligomeric compounds, antisense oligonucleotides, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other oligomeric compounds that hybridize to at least a portion of the target nucleic acid. As such, these compounds may be introduced in the form of single-stranded, double-stranded, partially single-stranded, or circular oligomeric compounds.

15 [0032] In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. The term "oligonucleotide", also includes linear or circular oligomers of natural and/or modified monomers or linkages, including deoxyribonucleosides, ribonucleosides, substituted and alpha-anomeric forms thereof, peptide nucleic acids (PNA), locked nucleic acids (LNA),  
20 phosphorothioate, methylphosphonate, and the like. Oligonucleotides are capable of specifically binding to a target polynucleotide by way of a regular pattern of monomer-to-monomer interactions, such as Watson-Crick type of base pairing, Hoögsteen or reverse Hoögsteen types of base pairing, or the like.

[0033] The oligonucleotide may be "chimeric", that is, composed of different regions. In the  
25 context of this invention "chimeric" compounds are oligonucleotides, which contain two or more chemical regions, for example, DNA region(s), RNA region(s), PNA region(s) etc. Each chemical region is made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotides compound. These oligonucleotides typically comprise at least one region wherein the oligonucleotide is modified in order to exhibit one or more desired properties.  
30 The desired properties of the oligonucleotide include, but are not limited, for example, to increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. Different regions of the oligonucleotide may therefore have different properties. The chimeric oligonucleotides of the present invention can



be formed as mixed structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide analogs as described above.

[0034] The oligonucleotide can be composed of regions that can be linked in "register", that is, when the monomers are linked consecutively, as in native DNA, or linked via spacers. The spacers are intended to constitute a covalent "bridge" between the regions and have in preferred cases a length not exceeding about 100 carbon atoms. The spacers may carry different functionalities, for example, having positive or negative charge, carry special nucleic acid binding properties (intercalators, groove binders, toxins, fluorophors etc.), being lipophilic, inducing special secondary structures like, for example, alanine containing peptides that induce alpha-helices.

[0035] As used herein "NRF2" and "Nuclear factor (erythroid-derived 2)-like 2" are inclusive of all family members, mutants, alleles, fragments, species, coding and noncoding sequences, sense and antisense polynucleotide strands, etc.

[0036] As used herein, the words 'Nuclear factor (erythroid-derived 2)-like 2', Nuclear factor erythroid 2-related factor 2, NFE2-related factor 2, NF-E2-related factor 2, HEBP1, NRF2, Nrf2, NFE2L2, are used interchangeably in the present application.

[0037] As used herein, the term "oligonucleotide specific for" or "oligonucleotide which targets" refers to an oligonucleotide having a sequence (i) capable of forming a stable complex with a portion of the targeted gene, or (ii) capable of forming a stable duplex with a portion of a mRNA transcript of the targeted gene. Stability of the complexes and duplexes can be determined by theoretical calculations and/or in vitro assays. Exemplary assays for determining stability of hybridization complexes and duplexes are described in the Examples below.

[0038] As used herein, the term "target nucleic acid" encompasses DNA, RNA (comprising premRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA, coding, noncoding sequences, sense or antisense polynucleotides. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds, which specifically hybridize to it, is generally referred to as "antisense". The functions of DNA to be interfered include, for example, replication and transcription. The functions of RNA to be interfered, include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one

or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of an encoded product or oligonucleotides.

[0039] RNA interference "RNAi" is mediated by double stranded RNA (dsRNA) molecules that have sequence-specific homology to their "target" nucleic acid sequences (Caplen, N. J., *et al.* (2001) *Proc. Natl. Acad. Sci. USA* 98:9742-9747). In certain embodiments of the present invention, the mediators are 5-25 nucleotide "small interfering" RNA duplexes (siRNAs). The siRNAs are derived from the processing of dsRNA by an RNase enzyme known as Dicer (Bernstein, E., *et al.* (2001) *Nature* 409:363-366). siRNA duplex products are recruited into a multi-protein siRNA complex termed RISC (RNA Induced Silencing Complex). Without wishing to be bound by any particular theory, a RISC is then believed to be guided to a target nucleic acid (suitably mRNA), where the siRNA duplex interacts in a sequence-specific way to mediate cleavage in a catalytic fashion (Bernstein, E., *et al.* (2001) *Nature* 409:363-366; Boutla, A., *et al.* (2001) *Curr. Biol.* 11:1776-1780). Small interfering RNAs that can be used in accordance with the present invention can be synthesized and used according to procedures that are well known in the art and that will be familiar to the ordinarily skilled artisan. Small interfering RNAs for use in the methods of the present invention suitably comprise between about 1 to about 50 nucleotides (nt). In examples of non limiting embodiments, siRNAs can comprise about 5 to about 40 nt, about 5 to about 30 nt, about 10 to about 30 nt, about 15 to about 25 nt, or about 20-25 nucleotides.

[0040] Selection of appropriate oligonucleotides is facilitated by using computer programs that automatically align nucleic acid sequences and indicate regions of identity or homology. Such programs are used to compare nucleic acid sequences obtained, for example, by searching databases such as GenBank or by sequencing PCR products. Comparison of nucleic acid sequences from a range of species allows the selection of nucleic acid sequences that display an appropriate degree of identity between species. In the case of genes that have not been sequenced, Southern blots are performed to allow a determination of the degree of identity between genes in target species and other species. By performing Southern blots at varying degrees of stringency, as is well known in the art, it is possible to obtain an approximate measure of identity. These procedures allow the selection of oligonucleotides that exhibit a high degree of complementarity to target nucleic acid sequences in a subject to be controlled and a lower degree of complementarity to corresponding nucleic acid sequences

in other species. One skilled in the art will realize that there is considerable latitude in selecting appropriate regions of genes for use in the present invention.

[0041] By "enzymatic RNA" is meant an RNA molecule with enzymatic activity (Cech, (1988) *J. American. Med. Assoc.* 260, 3030-3035). Enzymatic nucleic acids (ribozymes) act by first binding to a target RNA. Such binding occurs through the target binding portion of an enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through base pairing, and once bound to the correct site, acts enzymatically to cut the target RNA.

[0042] By "decoy RNA" is meant an RNA molecule that mimics the natural binding domain for a ligand. The decoy RNA therefore competes with natural binding target for the binding of a specific ligand. For example, it has been shown that over-expression of HIV trans-activation response (TAR) RNA can act as a "decoy" and efficiently binds HIV tat protein, thereby preventing it from binding to TAR sequences encoded in the HIV RNA (Sullenger *et al.* (1990) *Cell*, 63, 601- 608). This is meant to be a specific example. Those in the art will recognize that this is but one example, and other embodiments can be readily generated using techniques generally known in the art.

[0043] As used herein, the term "monomers" typically indicates monomers linked by phosphodiester bonds or analogs thereof to form oligonucleotides ranging in size from a few monomeric units, e.g., from about 3-4, to about several hundreds of monomeric units. Analogs of phosphodiester linkages include: phosphorothioate, phosphorodithioate, methylphosphonates, phosphoroselenoate, phosphoramidate, and the like, as more fully described below.

[0044] The term "nucleotide" covers naturally occurring nucleotides as well as nonnaturally occurring nucleotides. It should be clear to the person skilled in the art that various nucleotides which previously have been considered "non-naturally occurring" have subsequently been found in nature. Thus, "nucleotides" includes not only the known purine and pyrimidine heterocycles-containing molecules, but also heterocyclic analogues and tautomers thereof. Illustrative examples of other types of nucleotides are molecules containing adenine, guanine, thymine, cytosine, uracil, purine, xanthine, diaminopurine, 8-oxo- N6-methyladenine, 7-deazaxanthine, 7-deazaguanine, N4,N4-ethanocytosin, N6,N6-ethano-2,6-diaminopurine, 5-methylcytosine, 5-(C3-C6)-alkynylcytosine, 5-fluorouracil, 5-bromouracil,

pseudoisocytosine, 2-hydroxy-5-methyl-4-triazolopyridin, isocytosine, isoguanin, inosine and the "non-naturally occurring" nucleotides described in Benner *et al.*, U.S. Pat No. 5,432,272. The term "nucleotide" is intended to cover every and all of these examples as well as analogues and tautomers thereof. Especially interesting nucleotides are those containing  
5 adenine, guanine, thymine, cytosine, and uracil, which are considered as the naturally occurring nucleotides in relation to therapeutic and diagnostic application in humans. Nucleotides include the natural 2'-deoxy and 2'-hydroxyl sugars, e.g., as described in Kornberg and Baker, DNA Replication, 2nd Ed. (Freeman, San Francisco, 1992) as well as their analogs.

10 [0045] "Analog" in reference to nucleotides includes synthetic nucleotides having modified base moieties and/or modified sugar moieties (see e.g., described generally by Scheit, Nucleotide Analogs, John Wiley, New York, 1980; Freier & Altmann, (1997) *Nucl. Acid. Res.*, 25(22), 4429- 4443, Toulmé, J.J., (2001) *Nature Biotechnology* 19:17-18; Manoharan M., (1999) *Biochemica et Biophysica Acta* 1489:117-139; Freier S. M., (1997) *Nucleic Acid*  
15 *Research*, 25:4429-4443, Uhlman, E., (2000) *Drug Discovery & Development*, 3: 203-213, Herdewin P., (2000) *Antisense & Nucleic Acid Drug Dev.*, 10:297-310); 2'-O, 3'-C-linked [3.2.0] bicycloarabinonucleosides (see e.g. N.K Christensen., *et al*, (1998) *J. Am. Chem. Soc.*, 120: 5458-5463; Prakash TP, Bhat B. (2007) *Curr Top Med Chem.* 7(7):641-9; Cho EJ, *et al.* (2009) *Annual Review of Analytical Chemistry*, 2, 241-264). Such analogs include synthetic  
20 nucleotides designed to enhance binding properties, e.g., duplex or triplex stability, specificity, or the like.

[0046] As used herein, "hybridization" means the pairing of substantially complementary strands of oligomeric compounds. One mechanism of pairing involves hydrogen bonding, which may be Watson-Crick, Hoögsteeen or reversed Hoögsteeen hydrogen bonding, between  
25 complementary nucleoside or nucleotide bases (nucleotides) of the strands of oligomeric compounds. For example, adenine and thymine are complementary nucleotides which pair through the formation of hydrogen bonds. Hybridization can occur under varying circumstances.

[0047] An antisense compound is "specifically hybridizable" when binding of the compound  
30 to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a modulation of function and/or activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences

under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and under conditions in which assays are performed in the case of in vitro assays.

[0048] As used herein, the phrase "stringent hybridization conditions" or "stringent conditions" refers to conditions under which a compound of the invention will hybridize to its target sequence, but to a minimal number of other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances and in the context of this invention, "stringent conditions" under which oligomeric compounds hybridize to a target sequence are determined by the nature and composition of the oligomeric compounds and the assays in which they are being investigated. In general, stringent hybridization conditions comprise low concentrations ( $<0.15\text{M}$ ) of salts with inorganic cations such as  $\text{Na}^{++}$  or  $\text{K}^{++}$  (i.e., low ionic strength), temperature higher than  $20^{\circ}\text{C}$  -  $25^{\circ}\text{C}$ . below the  $T_m$  of the oligomeric compound:target sequence complex, and the presence of denaturants such as formamide, dimethylformamide, dimethyl sulfoxide, or the detergent sodium dodecyl sulfate (SDS). For example, the hybridization rate decreases 1.1% for each 1% formamide. An example of a high stringency hybridization condition is 0.1X sodium chloride-sodium citrate buffer (SSC)/0.1% (w/v) SDS at  $60^{\circ}\text{C}$ . for 30 minutes.

[0049] "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides on one or two oligomeric strands. For example, if a nucleobase at a certain position of an antisense compound is capable of hydrogen bonding with a nucleobase at a certain position of a target nucleic acid, said target nucleic acid being a DNA, RNA, or oligonucleotide molecule, then the position of hydrogen bonding between the oligonucleotide and the target nucleic acid is considered to be a complementary position. The oligomeric compound and the further DNA, RNA, or oligonucleotide molecule are complementary to each other when a sufficient number of complementary positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of precise pairing or complementarity over a sufficient number of nucleotides such that stable and specific binding occurs between the oligomeric compound and a target nucleic acid.

[0050] It is understood in the art that the sequence of an oligomeric compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. Moreover, an oligonucleotide may hybridize over one or more segments such that intervening

or adjacent segments are not involved in the hybridization event (e.g., a loop structure, mismatch or hairpin structure). The oligomeric compounds of the present invention comprise at least about 70%, or at least about 75%, or at least about 80%, or at least about 85%, or at least about 90%, or at least about 95%, or at least about 99% sequence complementarity to a target region within the target nucleic acid sequence to which they are targeted. For example, an antisense compound in which 18 of 20 nucleotides of the antisense compound are complementary to a target region, and would therefore specifically hybridize, would represent 90 percent complementarity. In this example, the remaining noncomplementary nucleotides may be clustered or interspersed with complementary nucleotides and need not be contiguous to each other or to complementary nucleotides. As such, an antisense compound which is 18 nucleotides in length having 4 (four) noncomplementary nucleotides which are flanked by two regions of complete complementarity with the target nucleic acid would have 77.8% overall complementarity with the target nucleic acid and would thus fall within the scope of the present invention. Percent complementarity of an antisense compound with a region of a target nucleic acid can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul *et al.*, (1990) *J. Mol. Biol.*, 215, 403-410; Zhang and Madden, (1997) *Genome Res.*, 7, 649-656). Percent homology, sequence identity or complementarity, can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), using default settings, which uses the algorithm of Smith and Waterman (*Adv. Appl. Math.*, (1981) 2, 482-489).

[0051] As used herein, the term "Thermal Melting Point (T<sub>m</sub>)" refers to the temperature, under defined ionic strength, pH, and nucleic acid concentration, at which 50% of the oligonucleotides complementary to the target sequence hybridize to the target sequence at equilibrium. Typically, stringent conditions will be those in which the salt concentration is at least about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short oligonucleotides (e.g., 10 to 50 nucleotide). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

[0052] As used herein, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene.

[0053] The term "variant," when used in the context of a polynucleotide sequence, may encompass a polynucleotide sequence related to a wild type gene. This definition may also include, for example, "allelic," "splice," "species," or "polymorphic" variants. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or an absence of domains. Species variants are polynucleotide sequences that vary from one species to another. Of particular utility in the invention are variants of wild type gene products. Variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes that give rise to variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

[0054] The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs,) or single base mutations in which the polynucleotide sequence varies by one base. The presence of SNPs may be indicative of, for example, a certain population with a propensity for a disease state, that is susceptibility versus resistance.

[0055] Derivative polynucleotides include nucleic acids subjected to chemical modification, for example, replacement of hydrogen by an alkyl, acyl, or amino group. Derivatives, e.g., derivative oligonucleotides, may comprise non-naturally-occurring portions, such as altered sugar moieties or inter-sugar linkages. Exemplary among these are phosphorothioate and other sulfur containing species which are known in the art. Derivative nucleic acids may also contain labels, including radionucleotides, enzymes, fluorescent agents, chemiluminescent agents, chromogenic agents, substrates, cofactors, inhibitors, magnetic particles, and the like.

[0056] A "derivative" polypeptide or peptide is one that is modified, for example, by glycosylation, pegylation, phosphorylation, sulfation, reduction/alkylation, acylation, chemical coupling, or mild formalin treatment. A derivative may also be modified to contain a

detectable label, either directly or indirectly, including, but not limited to, a radioisotope, fluorescent, and enzyme label.

[0057] As used herein, the term "animal" or "patient" is meant to include, for example, humans, sheep, elks, deer, mule deer, minks, mammals, monkeys, horses, cattle, pigs, goats, dogs, cats, rats, mice, birds, chicken, reptiles, fish, insects and arachnids.

[0058] "Mammal" covers warm blooded mammals that are typically under medical care (e.g., humans and domesticated animals). Examples include feline, canine, equine, bovine, and human, as well as just human.

[0059] "Treating" or "treatment" covers the treatment of a disease-state in a mammal, and includes: (a) preventing the disease-state from occurring in a mammal, in particular, when such mammal is predisposed to the disease-state but has not yet been diagnosed as having it; (b) inhibiting the disease-state, e.g., arresting its development; and/or (c) relieving the disease-state, e.g., causing regression of the disease state until a desired endpoint is reached. Treating also includes the amelioration of a symptom of a disease (e.g., lessen the pain or discomfort), wherein such amelioration may or may not be directly affecting the disease (e.g., cause, transmission, expression, etc.).

[0060] As used herein, the term "cancer" refers to any malignant tumor, particularly arising in the lung, kidney, or thyroid. The cancer manifests itself as a "tumor" or tissue comprising malignant cells of the cancer. Examples of tumors include sarcomas and carcinomas such as, but not limited to: fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma. As used herein, the terms "cancer," "neoplasm," and "tumor," are used interchangeably and in either the singular or



plural form, refer to cells that have undergone a malignant transformation that makes them pathological to the host organism. Primary cancer cells (that is, cells obtained from near the site of malignant transformation) can be readily distinguished from non-cancerous cells by well-established techniques, particularly histological examination. The definition of a cancer cell, as used herein, includes not only a primary cancer cell, but any cell derived from a cancer cell ancestor. This includes metastasized cancer cells, and in vitro cultures and cell lines derived from cancer cells. When referring to a type of cancer that normally manifests as a solid tumor, a "clinically detectable" tumor is one that is detectable on the basis of tumor mass; e.g., by procedures such as CAT scan, MR imaging, X-ray, ultrasound or palpation, and/or which is detectable because of the expression of one or more cancer-specific antigens in a sample obtainable from a patient.

[0061] "Neurological disease or disorder" refers to any disease or disorder of the nervous system and/or visual system. "Neurological disease or disorder" include disease or disorders that involve the central nervous system (brain, brainstem and cerebellum), the peripheral nervous system (including cranial nerves), and the autonomic nervous system (parts of which are located in both central and peripheral nervous system). A Neurological disease or disorder includes but is not limited to acquired epileptiform aphasia; acute disseminated encephalomyelitis; adrenoleukodystrophy; age-related macular degeneration; agenesis of the corpus callosum; agnosia; Aicardi syndrome; Alexander disease; Alpers' disease; alternating hemiplegia; Alzheimer's disease; Vascular dementia; amyotrophic lateral sclerosis; anencephaly; Angelman syndrome; angiomatosis; anoxia; aphasia; apraxia; arachnoid cysts; arachnoiditis; Anronl-Chiari malformation; arteriovenous malformation; Asperger syndrome; ataxia telegiectasia; attention deficit hyperactivity disorder; autism; autonomic dysfunction; back pain; Batten disease; Behcet's disease; Bell's palsy; benign essential blepharospasm; benign focal; amyotrophy; benign intracranial hypertension; Binswanger's disease; blepharospasm; Bloch Sulzberger syndrome; brachial plexus injury; brain abscess; brain injury; brain tumors (including glioblastoma multiforme); spinal tumor; Brown-Sequard syndrome; Canavan disease; carpal tunnel syndrome; causalgia; central pain syndrome; central pontine myelinolysis; cephalic disorder; cerebral aneurysm; cerebral arteriosclerosis; cerebral atrophy; cerebral gigantism; cerebral palsy; Charcot-Marie-Tooth disease; chemotherapy-induced neuropathy and neuropathic pain; Chiari malformation; chorea; chronic inflammatory demyelinating polyneuropathy; chronic pain; chronic regional pain syndrome; Coffin Lowry syndrome; coma, including persistent vegetative state; congenital

facial diplegia; corticobasal degeneration; cranial arteritis; craniosynostosis; Creutzfeldt-Jakob disease; cumulative trauma disorders; Cushing's syndrome; cytomegalic inclusion body disease; cytomegalovirus infection; dancing eyes-dancing feet syndrome; DandyWalker syndrome; Dawson disease; De Morsier's syndrome; Dejerine-Klumke palsy; dementia;
 5 dermatomyositis; diabetic neuropathy; diffuse sclerosis; dysautonomia; dysgraphia; dyslexia; dystonias; early infantile epileptic encephalopathy; empty sella syndrome; encephalitis; encephaloceles; encephalotrigeminal angiomatosis; epilepsy; Erb's palsy; essential tremor; Fabry's disease; Fahr's syndrome; fainting; familial spastic paralysis; febrile seizures; Fisher syndrome; Friedreich's ataxia; fronto-temporal dementia and other "tauopathies"; Gaucher's
 10 disease; Gerstmann's syndrome; giant cell arteritis; giant cell inclusion disease; globoid cell leukodystrophy; Guillain-Barre syndrome; HTLV-1-associated myelopathy; Hallervorden-Spatz disease; head injury; headache; hemifacial spasm; hereditary spastic paraplegia; heredopathia atactica polyneuritiformis; herpes zoster oticus; herpes zoster; Hirayama syndrome; HIV-associated dementia and neuropathy (also neurological manifestations of
 15 AIDS); holoprosencephaly; Huntington's disease and other polyglutamine repeat diseases; hydranencephaly; hydrocephalus; hypercortisolism; hypoxia; immune-mediated encephalomyelitis; inclusion body myositis; incontinentia pigmenti; infantile phytanic acid storage disease; infantile Refsum disease; infantile spasms; inflammatory myopathy; intracranial cyst; intracranial hypertension; Joubert syndrome; Keams-Sayre syndrome;
 20 Kennedy disease; Kinsbourne syndrome; Klippel Feil syndrome; Krabbe disease; Kugelberg-Welander disease; kuru; Lafora disease; Lambert-Eaton myasthenic syndrome; Landau-Kleffner syndrome; lateral medullary (Wallenberg) syndrome; learning disabilities; Leigh's disease; Lennox-Gustaut syndrome; Lesch-Nyhan syndrome; leukodystrophy; Lewy body dementia; Lissencephaly; locked-in syndrome; Lou Gehrig's disease (i.e., motor neuron
 25 disease or amyotrophic lateral sclerosis); lumbar disc disease; Lyme disease--neurological sequelae; Machado-Joseph disease; macrencephaly; megalencephaly; Melkersson-Rosenthal syndrome; Meniere's disease; meningitis; Menkes disease; metachromatic leukodystrophy; microcephaly; migraine; Miller Fisher syndrome; mini-strokes; mitochondrial myopathies; Mobius syndrome; monomelic amyotrophy; motor neuron disease; Moyamoya disease;
 30 mucopolysaccharidoses; multi-infarct dementia; multifocal motor neuropathy; multiple sclerosis and other demyelinating disorders; multiple system atrophy with postural hypotension; muscular dystrophy; myasthenia gravis; myelinoclastic diffuse sclerosis; myoclonic encephalopathy of infants; myoclonus; myopathy; myotonia congenital;

narcolepsy; neurofibromatosis; neuroleptic malignant syndrome; neurological manifestations  
 of AIDS; neurological sequelae of lupus; neuromyotonia; neuronal ceroid lipofuscinosis;  
 neuronal migration disorders; Niemann-Pick disease; O'Sullivan-McLeod syndrome; occipital  
 neuralgia; occult spinal dysraphism sequence; Ohtahara syndrome; olivopontocerebellar  
 5 atrophy; opsoclonus myoclonus; optic neuritis; orthostatic hypotension; overuse syndrome;  
 paresthesia; a neurodegenerative disease or disorder (Parkinson's disease, Huntington's  
 disease, Alzheimer's disease, amyotrophic lateral sclerosis (ALS), dementia, multiple  
 sclerosis and other diseases and disorders associated with neuronal cell death); paramyotonia  
 congenital; paraneoplastic diseases; paroxysmal attacks; Parry Romberg syndrome; Pelizaeus-  
 10 Merzbacher disease; periodic paralyses; peripheral neuropathy; painful neuropathy and  
 neuropathic pain; persistent vegetative state; pervasive developmental disorders; photic  
 sneeze reflex; phytanic acid storage disease; Pick's disease; pinched nerve; pituitary tumors;  
 polymyositis; porencephaly; post-polio syndrome; postherpetic neuralgia; postinfectious  
 encephalomyelitis; postural hypotension; Prader- Willi syndrome; primary lateral sclerosis;  
 15 prion diseases; progressive hemifacial atrophy; progressive multifocal leukoencephalopathy;  
 progressive sclerosing poliomyelitis; progressive supranuclear palsy; pseudotumor cerebri;  
 Ramsay-Hunt syndrome (types I and II); Rasmussen's encephalitis; reflex sympathetic  
 dystrophy syndrome; Refsum disease; repetitive motion disorders; repetitive stress injuries;  
 restless legs syndrome; retrovirus-associated myelopathy; Rett syndrome; Reye's syndrome;  
 20 Saint Vitus dance; Sandhoff disease; Schilder's disease; schizencephaly; septo-optic  
 dysplasia; shaken baby syndrome; shingles; Shy-Drager syndrome; Sjogren's syndrome; sleep  
 apnea; Soto's syndrome; spasticity; spina bifida; spinal cord injury; spinal cord tumors; spinal  
 muscular atrophy; Stiff-Person syndrome; stroke; Sturge-Weber syndrome; subacute  
 sclerosing panencephalitis; subcortical arteriosclerotic encephalopathy; Sydenham chorea;  
 25 syncope; syringomyelia; tardive dyskinesia; Tay-Sachs disease; temporal arteritis; tethered  
 spinal cord syndrome; Thomsen disease; thoracic outlet syndrome; Tic Douloureux; Todd's  
 paralysis; Tourette syndrome; transient ischemic attack; transmissible spongiform  
 encephalopathies; transverse myelitis; traumatic brain injury; tremor; trigeminal neuralgia;  
 tropical spastic paraparesis; tuberous sclerosis; vascular dementia (multi-infarct dementia);  
 30 vasculitis including temporal arteritis; Von Hippel-Lindau disease; Wallenberg's syndrome;  
 Werdnig-Hoffman disease; West syndrome; whiplash; Williams syndrome; Wildon's disease;  
 and Zellweger syndrome.

*Polynucleotide and Oligonucleotide Compositions and Molecules*

[0062] *Targets:* In one embodiment, the targets comprise nucleic acid sequences of Nuclear factor (erythroid-derived 2)-like 2 (NRF2), including without limitation sense and/or antisense noncoding and/or coding sequences associated with NRF2.

[0063] Transcription factor NF-E2-related factor 2 (NRF2) belongs to the basic region-leucine zipper family and is activated in response to electrophiles and reactive oxygen species. NRF2 coordinately regulates the constitutive and inducible transcription of a wide array of genes involved in drug metabolism, detoxification, and antioxidant defenses. During periods of oxidative stress, NRF2 is released from sequestration in the cytoplasm and translocates to the nucleus. NRF2 binds antioxidant response elements (AREs) in the regulatory regions of the target genes and activates transcription.

[0064] Chronic obstructive pulmonary disease (COPD), which comprises emphysema and chronic bronchitis resulting from prolonged exposure to cigarette smoke (CS), is a major public health burden with no effective treatment. Emphysema is also associated with pulmonary hypertension, which can progress to right ventricular failure, an important cause of morbidity and mortality among patients with COPD. Recently, it has been shown that patients with advanced COPD have a decline in expression of the Nrf2 pathway in lungs, indicating that loss of this antioxidative protective response is a key factor in the pathophysiological progression of emphysema. Furthermore, genetic disruption of Nrf2 in mice causes early-onset and severe emphysema.

[0065] The liver is frequently exposed to insults, including toxic chemicals and alcohol, viral infection or metabolic overload. Although it can fully regenerate after acute injury, chronic liver damage causes liver fibrosis and cirrhosis, which can result in complete liver failure.

[0066] In preferred embodiments, antisense oligonucleotides are used to prevent or treat diseases or disorders associated with abnormal nuclear factor (erythroid-derived 2)-like 2 (NRF2) expression and/or function. This includes all forms of nuclear factor (erythroid-derived 2)-like 2 (NRF2) molecules, including mutants and aberrant expression or function of normal or abnormal nuclear factor (erythroid-derived 2)-like 2 (NRF2) molecules.

[0067] In a preferred embodiment, modulation of NF-E2-related factor 2 (Nrf2) transcription factor protects the liver from acute and chronic toxin-mediated damage.

[0068] In another preferred embodiment, up-regulation or activation of Nrf2 prevents or ameliorates toxin-induced liver injury and fibrosis.

[0069] In another preferred embodiment, modulation of NF-E2-related factor 2 (Nrf2) transcription factor prevents or treats gastrointestinal diseases or disorders in patients.

[0070] Brain tumors are associated with genetic alterations of oncogenes and tumor suppressor genes. Accumulation of reactive oxygen species (ROS) in cells leads to oxidative stress-induced damage, resulting in tumorigenesis.

[0071] Nuclear factor erythroid 2-related factor 2 (Nrf2) is a redox-sensitive transcription factor that up-regulates a battery of antioxidative genes and cytoprotective enzymes that constitute the defense against oxidative stress. Modulation of this polynucleotide is important in treatment of diseases or disorders that are associated with oxidative stress. Exemplary Nuclear factor (erythroid-derived 2)-like 2 (NRF2) mediated diseases and disorders which can be treated with cell/tissues regenerated from stem cells obtained using the antisense compounds comprise: chronic obstructive pulmonary disease (COPD), multiple sclerosis, a hepatic disease or disorder, a gastrointestinal disease or disorder, diabetes, autoimmunity, an immune related disease or disorder, an immunodeficiency (e.g., AIDS), a neurological disease or disorder, a neurodegenerative disease or disorder, a disease or disorder or condition associated with oxidative stress, an eye disease (e.g., age-related macular degeneration, cataracts, light retinopathy, retinopathy of prematurity etc.), a skin disease, asthma, arteriosclerosis, a chronic inflammatory diseases or condition (e.g., vasculitis, pulmonary bronchitis, rheumatoid arthritis, osteoarthritis, hepatitis, pancreatitis, dermatitis, esophagitis, ulcerative colitis, Crohn's disease, conjunctivitis etc.), nerve repair and paralysis, neuroendocrine differentiation, an inflammatory disease, a muscular disease or disorder, diseases or disorders associated with infectious organisms, senile plaques, cerebral amyloid angiopathy, atherosclerosis, glioblastoma, amyloid deposition, neurofibrillary tangles, dementia, choriocarcinoma, astrocytoma, amyloidosis, hyperlipidemia, neoplastic transformation, an atherosclerotic plaque, an atherosclerotic obstruction, metastasis, myocardial infarction, pulmonary fibrosis, necrosis, shock, melanoma, colorectal carcinoma, genetic susceptibility, psoriasis, a disease or disorder associated with abnormal cell proliferation ((e.g., cancer, psoriasis etc.), cancer (e.g., prostate cancer, lung cancer, breast cancer, Non-small cell lung carcinomas (NSCLCs), leukemia etc.), inflammation, glioma, carcinoma, neuropathology, tumors, vascular diseases, cell damage, brain tumors, hypercholesterolemia, liposarcoma, coronary heart disease, coronary artery disease, glomerulonephritis, venous thrombosis and a pathological process.

[0072] In another preferred embodiment, the antisense oligonucleotides modulate the expression and/or function of NRF2 in patients suffering from or at risk of developing diseases or disorders associated with NRF2. Examples of such diseases or disorders associated with NRF2 comprise: Alzheimer's disease, multiple sclerosis, senile plaques, cerebral amyloid angiopathy, atherosclerosis, glioblastoma, amyloid deposition, neurodegenerative diseases, neurofibrillary tangles, dementia, choriocarcinoma, astrocytoma, amyloidosis, hyperlipidemia, neurodegeneration, neoplastic transformation, prostate cancer, atherosclerotic plaque, obstruction, AIDS, metastasis, myocardial infarction, pulmonary fibrosis, necrosis, shock, melanoma, colorectal carcinoma, genetic susceptibility, psoriasis, cancer, inflammation, glioma, carcinoma, breast cancer, neuropathology, tumors, prostate carcinoma, vascular diseases, cell damage, brain tumors, Non-small cell lung carcinomas (NSCLCs), hypercholesterolemia, liposarcoma, coronary heart disease, immunodeficiency, coronary artery disease, glomerulonephritis, venous thrombosis, pathological processes or leukemia.

[0073] In a preferred embodiment, the oligonucleotides are specific for polynucleotides of NRF2, which includes, without limitation noncoding regions. The NRF2 targets comprise variants of NRF2; mutants of NRF2, including SNPs; noncoding sequences of NRF2; alleles, fragments and the like. Preferably the oligonucleotide is an antisense RNA molecule.

[0074] In accordance with embodiments of the invention, the target nucleic acid molecule is not limited to NRF2 polynucleotides alone but extends to any of the isoforms, receptors, homologs, non-coding regions and the like of NRF2.

[0075] In another preferred embodiment, an oligonucleotide targets a natural antisense sequence (natural antisense to the coding and non-coding regions) of NRF2 targets, including, without limitation, variants, alleles, homologs, mutants, derivatives, fragments and complementary sequences thereto. Preferably the oligonucleotide is an antisense RNA or DNA molecule.

[0076] In another preferred embodiment, the oligomeric compounds of the present invention also include variants in which a different base is present at one or more of the nucleotide positions in the compound. For example, if the first nucleotide is an adenine, variants may be produced which contain thymidine, guanosine, cytidine or other natural or unnatural nucleotides at this position. This may be done at any of the positions of the antisense compound. These compounds are then tested using the methods described herein to determine their ability to inhibit expression of a target nucleic acid.

[0077] In some embodiments, homology, sequence identity or complementarity, between the antisense compound and target is from about 50% to about 60%. In some embodiments, homology, sequence identity or complementarity, is from about 60% to about 70%. In some embodiments, homology, sequence identity or complementarity, is from about 70% to about 80%. In some embodiments, homology, sequence identity or complementarity, is from about 80% to about 90%. In some embodiments, homology, sequence identity or complementarity, is about 90%, about 92%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100%.

[0078] An antisense compound is specifically hybridizable when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a loss of activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences under conditions in which specific binding is desired. Such conditions include, i.e., physiological conditions in the case of in vivo assays or therapeutic treatment, and conditions in which assays are performed in the case of in vitro assays.

[0079] An antisense compound, whether DNA, RNA, chimeric, substituted etc, is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

[0080] In another preferred embodiment, targeting of NRF2 including without limitation, antisense sequences which are identified and expanded, using for example, PCR, hybridization etc., one or more of the sequences set forth as SEQ ID NO: 3, and the like, modulate the expression or function of NRF2. In one embodiment, expression or function is up-regulated as compared to a control. In another preferred embodiment, expression or function is down-regulated as compared to a control.

[0081] In another preferred embodiment, oligonucleotides comprise nucleic acid sequences set forth as SEQ ID NOS: 4 to 6 including antisense sequences which are identified and expanded, using for example, PCR, hybridization etc. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds and the like.

Examples of modified bonds or internucleotide linkages comprise phosphorothioate, phosphorodithioate or the like. In another preferred embodiment, the nucleotides comprise a phosphorus derivative. The phosphorus derivative (or modified phosphate group) which may be attached to the sugar or sugar analog moiety in the modified oligonucleotides of the present invention may be a monophosphate, diphosphate, triphosphate, alkylphosphate, alkanephosphate, phosphorothioate and the like. The preparation of the above-noted phosphate analogs, and their incorporation into nucleotides, modified nucleotides and oligonucleotides, *per se*, is also known and need not be described here.

[0082] The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

[0083] In embodiments of the present invention oligomeric antisense compounds, particularly oligonucleotides, bind to target nucleic acid molecules and modulate the expression and/or function of molecules encoded by a target gene. The functions of DNA to be interfered comprise, for example, replication and transcription. The functions of RNA to be interfered comprise all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The functions may be up-regulated or inhibited depending on the functions desired.

[0084] The antisense compounds, include, antisense oligomeric compounds, antisense oligonucleotides, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other oligomeric compounds that hybridize to at least a portion of the target nucleic acid. As such, these compounds may be introduced in the form of single-stranded, double-stranded, partially single-stranded, or circular oligomeric compounds.

[0085] Targeting an antisense compound to a particular nucleic acid molecule, in the context of this invention, can be a multistep process. The process usually begins with the identification of a target nucleic acid whose function is to be modulated. This target nucleic acid may be, for example, a cellular gene (or mRNA transcribed from the gene) whose



expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target nucleic acid encodes Nuclear factor (erythroid-derived 2)-like 2 (NRF2).

[0086] The targeting process usually also includes determination of at least one target region, segment, or site within the target nucleic acid for the antisense interaction to occur such that the desired effect, e.g., modulation of expression, will result. Within the context of the present invention, the term "region" is defined as a portion of the target nucleic acid having at least one identifiable structure, function, or characteristic. Within regions of target nucleic acids are segments. "Segments" are defined as smaller or sub-portions of regions within a target nucleic acid. "Sites," as used in the present invention, are defined as positions within a target nucleic acid.

[0087] In a preferred embodiment, the antisense oligonucleotides bind to the natural antisense sequences of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) and modulate the expression and/or function of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) (SEQ ID NO: 1).

Examples of antisense sequences include SEQ ID NOS: 3 to 6.

[0088] In another preferred embodiment, the antisense oligonucleotides bind to one or more segments of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides and modulate the expression and/or function of Nuclear factor (erythroid-derived 2)-like 2 (NRF2). The segments comprise at least five consecutive nucleotides of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) sense or antisense polynucleotides.

[0089] In another preferred embodiment, the antisense oligonucleotides are specific for natural antisense sequences of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) wherein binding of the oligonucleotides to the natural antisense sequences of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) modulate expression and/or function of Nuclear factor (erythroid-derived 2)-like 2 (NRF2).

[0090] In another preferred embodiment, oligonucleotide compounds comprise sequences set forth as SEQ ID NOS: 4 to 6, antisense sequences which are identified and expanded, using for example, PCR, hybridization etc. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds and the like. Examples of modified bonds or internucleotide linkages comprise phosphorothioate, phosphorodithioate or the like. In another preferred embodiment, the nucleotides comprise a phosphorus derivative. The phosphorus derivative (or modified phosphate group) which may be attached to the sugar

or sugar analog moiety in the modified oligonucleotides of the present invention may be a monophosphate, diphosphate, triphosphate, alkylphosphate, alkanephosphate, phosphorothioate and the like. The preparation of the above-noted phosphate analogs, and their incorporation into nucleotides, modified nucleotides and oligonucleotides, per se, is also known and need not be described here.

[0091] Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes has a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG; and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). Eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA transcribed from a gene encoding Nuclear factor (erythroid-derived 2)-like 2 (NRF2), regardless of the sequence(s) of such codons. A translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively).

[0092] The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon. Consequently, the "start codon region" (or "translation initiation codon region") and the "stop codon region" (or "translation termination codon region") are all regions that may be targeted effectively with the antisense compounds of the present invention.

[0093] The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is

also a region which may be targeted effectively. Within the context of the present invention, a targeted region is the intragenic region encompassing the translation initiation or termination codon of the open reading frame (ORF) of a gene.

[0094] Another target region includes the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA (or corresponding nucleotides on the gene). Still another target region includes the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA (or corresponding nucleotides on the gene). The 5' cap site of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap site. Another target region for this invention is the 5' cap region.

[0095] Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. In one embodiment, targeting splice sites, i.e., intron-exon junctions or exon-intron junctions, is particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular splice product is implicated in disease. An aberrant fusion junction due to rearrangement or deletion is another embodiment of a target site. mRNA transcripts produced via the process of splicing of two (or more) mRNAs from different gene sources are known as "fusion transcripts". Introns can be effectively targeted using antisense compounds targeted to, for example, DNA or pre-mRNA.

[0096] In another preferred embodiment, the antisense oligonucleotides bind to coding and/or non-coding regions of a target polynucleotide and modulate the expression and/or function of the target molecule.

[0097] In another preferred embodiment, the antisense oligonucleotides bind to natural antisense polynucleotides and modulate the expression and/or function of the target molecule.

[0098] In another preferred embodiment, the antisense oligonucleotides bind to sense polynucleotides and modulate the expression and/or function of the target molecule.

[0099] Alternative RNA transcripts can be produced from the same genomic region of DNA. These alternative transcripts are generally known as "variants". More specifically, "pre-mRNA variants" are transcripts produced from the same genomic DNA that differ from other transcripts produced from the same genomic DNA in either their start or stop position and contain both intronic and exonic sequence.

[00100] Upon excision of one or more exon or intron regions, or portions thereof during splicing, pre-mRNA variants produce smaller "mRNA variants". Consequently, mRNA variants are processed pre-mRNA variants and each unique pre-mRNA variant must always produce a unique mRNA variant as a result of splicing. These mRNA variants are also known as "alternative splice variants". If no splicing of the pre-mRNA variant occurs then the pre-mRNA variant is identical to the mRNA variant.

[00101] Variants can be produced through the use of alternative signals to start or stop transcription. Pre-mRNAs and mRNAs can possess more than one start codon or stop codon. Variants that originate from a pre-mRNA or mRNA that use alternative start codons are known as "alternative start variants" of that pre-mRNA or mRNA. Those transcripts that use an alternative stop codon are known as "alternative stop variants" of that pre-mRNA or mRNA. One specific type of alternative stop variant is the "polyA variant" in which the multiple transcripts produced result from the alternative selection of one of the "polyA stop signals" by the transcription machinery, thereby producing transcripts that terminate at unique polyA sites. Within the context of the invention, the types of variants described herein are also embodiments of target nucleic acids.

[00102] The locations on the target nucleic acid to which the antisense compounds hybridize are defined as at least a 5-nucleotide long portion of a target region to which an active antisense compound is targeted.

[00103] While the specific sequences of certain exemplary target segments are set forth herein, one of skill in the art will recognize that these serve to illustrate and describe particular embodiments within the scope of the present invention. Additional target segments are readily identifiable by one having ordinary skill in the art in view of this disclosure.

[00104] Target segments 5-100 nucleotides in length comprising a stretch of at least five (5) consecutive nucleotides selected from within the illustrative preferred target segments are considered to be suitable for targeting as well.

[00105] Target segments can include DNA or RNA sequences that comprise at least the 5 consecutive nucleotides from the 5'-terminus of one of the illustrative preferred target segments (the remaining nucleotides being a consecutive stretch of the same DNA or RNA beginning immediately upstream of the 5'-terminus of the target segment and continuing until the DNA or RNA contains about 5 to about 100 nucleotides). Similarly preferred target segments are represented by DNA or RNA sequences that comprise at least the 5 consecutive nucleotides from the 3'-terminus of one of the illustrative preferred target segments (the remaining nucleotides being a consecutive stretch of the same DNA or RNA beginning immediately downstream of the 3'-terminus of the target segment and continuing until the DNA or RNA contains about 5 to about 100 nucleotides). One having skill in the art armed with the target segments illustrated herein will be able, without undue experimentation, to identify further preferred target segments.

[00106] Once one or more target regions, segments or sites have been identified, antisense compounds are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

[00107] In embodiments of the invention the oligonucleotides bind to an antisense strand of a particular target. The oligonucleotides are at least 5 nucleotides in length and can be synthesized so each oligonucleotide targets overlapping sequences such that oligonucleotides are synthesized to cover the entire length of the target polynucleotide. The targets also include coding as well as non coding regions.

[00108] In one embodiment, it is preferred to target specific nucleic acids by antisense oligonucleotides. Targeting an antisense compound to a particular nucleic acid, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a non coding polynucleotide such as for example, non coding RNA (ncRNA).

[00109] RNAs can be classified into (1) messenger RNAs (mRNAs), which are translated into proteins, and (2) non-protein-coding RNAs (ncRNAs). ncRNAs comprise microRNAs, antisense transcripts and other Transcriptional Units (TU) containing a high density of stop codons and lacking any extensive "Open Reading Frame". Many ncRNAs appear to start from initiation sites in 3' untranslated regions (3'UTRs) of protein-coding loci. ncRNAs are often rare and at least half of the ncRNAs that have been sequenced by the FANTOM consortium

seem not to be polyadenylated. Most researchers have for obvious reasons focused on polyadenylated mRNAs that are processed and exported to the cytoplasm. Recently, it was shown that the set of non-polyadenylated nuclear RNAs may be very large, and that many such transcripts arise from so-called intergenic regions (Cheng, J. *et al.* (2005) *Science* 308 (5725), 1149-1154; Kapranov, P. *et al.* (2005). *Genome Res* 15 (7), 987-997). The mechanism by which ncRNAs may regulate gene expression is by base pairing with target transcripts. The RNAs that function by base pairing can be grouped into (1) cis encoded RNAs that are encoded at the same genetic location, but on the opposite strand to the RNAs they act upon and therefore display perfect complementarity to their target, and (2) trans-encoded RNAs that are encoded at a chromosomal location distinct from the RNAs they act upon and generally do not exhibit perfect base-pairing potential with their targets.

[00110] Without wishing to be bound by theory, perturbation of an antisense polynucleotide by the antisense oligonucleotides described herein can alter the expression of the corresponding sense messenger RNAs. However, this regulation can either be discordant (antisense knockdown results in messenger RNA elevation) or concordant (antisense knockdown results in concomitant messenger RNA reduction). In these cases, antisense oligonucleotides can be targeted to overlapping or non-overlapping parts of the antisense transcript resulting in its knockdown or sequestration. Coding as well as non-coding antisense can be targeted in an identical manner and that either category is capable of regulating the corresponding sense transcripts – either in a concordant or discordant manner. The strategies that are employed in identifying new oligonucleotides for use against a target can be based on the knockdown of antisense RNA transcripts by antisense oligonucleotides or any other means of modulating the desired target.

[00111] *Strategy 1:* In the case of discordant regulation, knocking down the antisense transcript elevates the expression of the conventional (sense) gene. Should that latter gene encode for a known or putative drug target, then knockdown of its antisense counterpart could conceivably mimic the action of a receptor agonist or an enzyme stimulant.

[00112] *Strategy 2:* In the case of concordant regulation, one could concomitantly knock down both antisense and sense transcripts and thereby achieve synergistic reduction of the conventional (sense) gene expression. If, for example, an antisense oligonucleotide is used to achieve knockdown, then this strategy can be used to apply one antisense oligonucleotide targeted to the sense transcript and another antisense oligonucleotide to the corresponding

antisense transcript, or a single energetically symmetric antisense oligonucleotide that simultaneously targets overlapping sense and antisense transcripts.

[00113] According to the present invention, antisense compounds include antisense oligonucleotides, ribozymes, external guide sequence (EGS) oligonucleotides, siRNA compounds, single- or double-stranded RNA interference (RNAi) compounds such as siRNA compounds, and other oligomeric compounds which hybridize to at least a portion of the target nucleic acid and modulate its function. As such, they may be DNA, RNA, DNA-like, RNA-like, or mixtures thereof, or may be mimetics of one or more of these. These compounds may be single-stranded, doublestranded, circular or hairpin oligomeric compounds and may contain structural elements such as internal or terminal bulges, mismatches or loops. Antisense compounds are routinely prepared linearly but can be joined or otherwise prepared to be circular and/or branched. Antisense compounds can include constructs such as, for example, two strands hybridized to form a wholly or partially double-stranded compound or a single strand with sufficient self-complementarity to allow for hybridization and formation of a fully or partially double-stranded compound. The two strands can be linked internally leaving free 3' or 5' termini or can be linked to form a continuous hairpin structure or loop. The hairpin structure may contain an overhang on either the 5' or 3' terminus producing an extension of single stranded character. The double stranded compounds optionally can include overhangs on the ends. Further modifications can include conjugate groups attached to one of the termini, selected nucleotide positions, sugar positions or to one of the internucleoside linkages. Alternatively, the two strands can be linked via a non-nucleic acid moiety or linker group. When formed from only one strand, dsRNA can take the form of a self-complementary hairpin-type molecule that doubles back on itself to form a duplex. Thus, the dsRNAs can be fully or partially double stranded. Specific modulation of gene expression can be achieved by stable expression of dsRNA hairpins in transgenic cell lines, however, in some embodiments, the gene expression or function is up regulated. When formed from two strands, or a single strand that takes the form of a self-complementary hairpin-type molecule doubled back on itself to form a duplex, the two strands (or duplex-forming regions of a single strand) are complementary RNA strands that base pair in Watson-Crick fashion.

[00114] Once introduced to a system, the compounds of the invention may elicit the action of one or more enzymes or structural proteins to effect cleavage or other modification of the target nucleic acid or may work via occupancy-based mechanisms. In general, nucleic acids

(including oligonucleotides) may be described as "DNA-like" (i.e., generally having one or more 2'-deoxy sugars and, generally, T rather than U bases) or "RNA-like" (i.e., generally having one or more 2'-hydroxyl or 2'-modified sugars and, generally U rather than T bases). Nucleic acid helices can adopt more than one type of structure, most commonly the A- and B-forms. It is believed that, in general, oligonucleotides which have B-form-like structure are "DNA-like" and those which have A-formlike structure are "RNA-like." In some (chimeric) embodiments, an antisense compound may contain both A- and B-form regions.

[00115] In another preferred embodiment, the desired oligonucleotides or antisense compounds, comprise at least one of: antisense RNA, antisense DNA, chimeric antisense oligonucleotides, antisense oligonucleotides comprising modified linkages, interference RNA (RNAi), short interfering RNA (siRNA); a micro, interfering RNA (miRNA); a small, temporal RNA (stRNA); or a short, hairpin RNA (shRNA); small RNA-induced gene activation (RNAa); small activating RNAs (saRNAs), or combinations thereof.

[00116] dsRNA can also activate gene expression, a mechanism that has been termed "small RNA-induced gene activation" or RNAa. dsRNAs targeting gene promoters induce potent transcriptional activation of associated genes. RNAa was demonstrated in human cells using synthetic dsRNAs, termed "small activating RNAs" (saRNAs). It is currently not known whether RNAa is conserved in other organisms.

[00117] Small double-stranded RNA (dsRNA), such as small interfering RNA (siRNA) and microRNA (miRNA), have been found to be the trigger of an evolutionary conserved mechanism known as RNA interference (RNAi). RNAi invariably leads to gene silencing via remodeling chromatin to thereby suppress transcription, degrading complementary mRNA, or blocking protein translation. However, in instances described in detail in the examples section which follows, oligonucleotides are shown to increase the expression and/or function of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides and encoded products thereof. dsRNAs may also act as small activating RNAs (saRNA). Without wishing to be bound by theory, by targeting sequences in gene promoters, saRNAs would induce target gene expression in a phenomenon referred to as dsRNA-induced transcriptional activation (RNAa).

[00118] In a further embodiment, the "preferred target segments" identified herein may be employed in a screen for additional compounds that modulate the expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides. "Modulators" are those compounds that decrease or increase the expression of a nucleic acid molecule encoding Nuclear factor



(erythroid-derived 2)-like 2 (NRF2) and which comprise at least a 5-nucleotide portion that is complementary to a preferred target segment. The screening method comprises the steps of contacting a preferred target segment of a nucleic acid molecule encoding sense or natural antisense polynucleotides of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) with one or more candidate modulators, and selecting for one or more candidate modulators which decrease or increase the expression of a nucleic acid molecule encoding Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides, e.g. SEQ ID NOS: 4 to 6. Once it is shown that the candidate modulator or modulators are capable of modulating (e.g. either decreasing or increasing) the expression of a nucleic acid molecule encoding Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides, the modulator may then be employed in further investigative studies of the function of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides, or for use as a research, diagnostic, or therapeutic agent in accordance with the present invention.

**[00119]** Targeting the natural antisense sequence preferably modulates the function of the target gene. For example, the NRF2 gene (e.g. accession number NM\_006164, Fig. 2). In a preferred embodiment, the target is an antisense polynucleotide of the NRF2 gene. In a preferred embodiment, an antisense oligonucleotide targets sense and/or natural antisense sequences of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides (e.g. accession number NM\_006164, Fig. 2), variants, alleles, isoforms, homologs, mutants, derivatives, fragments and complementary sequences thereto. Preferably the oligonucleotide is an antisense molecule and the targets include coding and noncoding regions of antisense and/or sense NRF2 polynucleotides.

**[00120]** The preferred target segments of the present invention may be also be combined with their respective complementary antisense compounds of the present invention to form stabilized double-stranded (duplexed) oligonucleotides.

**[00121]** Such double stranded oligonucleotide moieties have been shown in the art to modulate target expression and regulate translation as well as RNA processing via an antisense mechanism. Moreover, the double-stranded moieties may be subject to chemical modifications (Fire *et al.*, (1998) *Nature*, 391, 806-811; Timmons and Fire, (1998) *Nature*, 395, 854; Timmons *et al.*, (2001) *Gene*, 263, 103-112; Tabara *et al.*, (1998) *Science*, 282, 430-431; Montgomery *et al.*, (1998) *Proc. Natl. Acad. Sci. USA*, 95, 15502-15507; Tuschl *et al.*, (1999) *Genes Dev.*, 13, 3191-3197; Elbashir *et al.*, (2001) *Nature*, 411, 494-498; Elbashir

*et al.*, (2001) *Genes Dev.* 15, 188-200). For example, such double-stranded moieties have been shown to inhibit the target by the classical hybridization of antisense strand of the duplex to the target, thereby triggering enzymatic degradation of the target (Tijsterman *et al.*, (2002) *Science*, 295, 694-697).

5 [00122] In a preferred embodiment, an antisense oligonucleotide targets Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides (e.g. accession number NM\_006164), variants, alleles, isoforms, homologs, mutants, derivatives, fragments and complementary sequences thereto. Preferably the oligonucleotide is an antisense molecule.

10 [00123] In accordance with embodiments of the invention, the target nucleic acid molecule is not limited to Nuclear factor (erythroid-derived 2)-like 2 (NRF2) alone but extends to any of the isoforms, receptors, homologs and the like of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) molecules.

15 [00124] In another preferred embodiment, an oligonucleotide targets a natural antisense sequence of NRF2 polynucleotides, for example, polynucleotides set forth as SEQ ID NO: 3, and any variants, alleles, homologs, mutants, derivatives, fragments and complementary sequences thereto. Examples of antisense oligonucleotides are set forth as SEQ ID NOS: 4 to 6.

20 [00125] In one embodiment, the oligonucleotides are complementary to or bind to nucleic acid sequences of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) antisense, including without limitation noncoding sense and/or antisense sequences associated with Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides and modulate expression and/or function of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) molecules.

25 [00126] In another preferred embodiment, the oligonucleotides are complementary to or bind to nucleic acid sequences of NRF2 natural antisense, set forth as SEQ ID NO: 3 and modulate expression and/or function of NRF2 molecules.

[00127] In a preferred embodiment, oligonucleotides comprise sequences of at least 5 consecutive nucleotides of SEQ ID NOS: 4 to 6 and modulate expression and/or function of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) molecules.

30 [00128] The polynucleotide targets comprise NRF2, including family members thereof, variants of NRF2; mutants of NRF2, including SNPs; noncoding sequences of NRF2; alleles

of NRF2; species variants, fragments and the like. Preferably the oligonucleotide is an antisense molecule.

[00129] In another preferred embodiment, the oligonucleotide targeting Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides, comprise: antisense RNA, interference RNA (RNAi), short interfering RNA (siRNA); micro interfering RNA (miRNA); a small, temporal RNA (stRNA); or a short, hairpin RNA (shRNA); small RNA-induced gene activation (RNAa); or, small activating RNA (saRNA).

[00130] In another preferred embodiment, targeting of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides, e.g. SEQ ID NO: 3, modulates the expression or function of these targets. In one embodiment, expression or function is up-regulated as compared to a control. In another preferred embodiment, expression or function is down-regulated as compared to a control.

[00131] In another preferred embodiment, antisense compounds comprise sequences set forth as SEQ ID NOS: 4 to 6. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds and the like.

[00132] In another preferred embodiment, SEQ ID NOS: 4 to 6 comprise one or more LNA nucleotides.

[00133] Table 1 shows exemplary antisense oligonucleotides useful in the methods of the invention.

Sequence ID	Oligo Name	Sequence
SEQ ID NO:4	CUR- 0030	UAGGAAAGACAUUCCUCAUUCCUCAUUUAGGCAA
SEQ ID NO:5	CUR -0032	UUCUGCUACCAUAUCCACCACUCCAC
SEQ ID NO:6	CUR - 0509	T*A*G*G*A*A*A*G*A*C*A*T*T*C*C*T*C*A*T

[00134] The modulation of a desired target nucleic acid can be carried out in several ways known in the art. For example, antisense oligonucleotides, siRNA etc. Enzymatic nucleic acid molecules (e.g., ribozymes) are nucleic acid molecules capable of catalyzing one or more of a variety of reactions, including the ability to repeatedly cleave other separate nucleic acid molecules in a nucleotide base sequence-specific manner. Such enzymatic nucleic acid molecules can be used, for example, to target virtually any RNA transcript (Zaug *et al.*, 324,

*Nature* 429 1986; Cech, 260 *JAMA* 3030, 1988; and Jefferies *et al.*, 17 *Nucleic Acids Research* 1371, 1989).

[00135] Because of their sequence-specificity, trans-cleaving enzymatic nucleic acid molecules show promise as therapeutic agents for human disease (Usman & McSwiggen, (1995) *Ann. Rep. Med. Chem.* 30, 285-294; Christoffersen and Marr, (1995) *J. Med. Chem.* 38, 2023-2037). Enzymatic nucleic acid molecules can be designed to cleave specific RNA targets within the background of cellular RNA. Such a cleavage event renders the mRNA non-functional and abrogates protein expression from that RNA. In this manner, synthesis of a protein associated with a disease state can be selectively inhibited.

[00136] In general, enzymatic nucleic acids with RNA cleaving activity act by first binding to a target RNA. Such binding occurs through the target binding portion of a enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through complementary base pairing, and once bound to the correct site, acts enzymatically to cut the target RNA. Strategic cleavage of such a target RNA will destroy its ability to direct synthesis of an encoded protein. After an enzymatic nucleic acid has bound and cleaved its RNA target, it is released from that RNA to search for another target and can repeatedly bind and cleave new targets.

[00137] Several approaches such as in vitro selection (evolution) strategies (Orgel, (1979) *Proc. R. Soc. London, B* 205, 435) have been used to evolve new nucleic acid catalysts capable of catalyzing a variety of reactions, such as cleavage and ligation of phosphodiester linkages and amide linkages, (Joyce, (1989) *Gene*, 82, 83-87; Beaudry *et al.*, (1992) *Science* 257, 635-641; Joyce, (1992) *Scientific American* 267, 90-97; Breaker *et al.*, (1994) *TIBTECH* 12, 268; Bartel *et al.*, (1993) *Science* 261:1411- 1418; Szostak, (1993) *TIBS* 17, 89-93; Kumar *et al.*, (1995) *FASEB J.*, 9, 1183; Breaker, (1996) *Curr. Op. Biotech.*, 7, 442).

[00138] The development of ribozymes that are optimal for catalytic activity would contribute significantly to any strategy that employs RNA-cleaving ribozymes for the purpose of regulating gene expression. The hammerhead ribozyme, for example, functions with a catalytic rate (kcat) of about 1 min<sup>-1</sup> in the presence of saturating (10 mM) concentrations of Mg<sup>2+</sup> cofactor. An artificial "RNA ligase" ribozyme has been shown to catalyze the corresponding self-modification reaction with a rate of about 100 min<sup>-1</sup>. In addition, it is known that certain modified hammerhead ribozymes that have substrate binding arms made

of DNA catalyze RNA cleavage with multiple turn-over rates that approach 100 min<sup>-1</sup>. Finally, replacement of a specific residue within the catalytic core of the hammerhead with certain nucleotide analogues gives modified ribozymes that show as much as a 10-fold improvement in catalytic rate. These findings demonstrate that ribozymes can promote chemical transformations with catalytic rates that are significantly greater than those displayed in vitro by most natural self-cleaving ribozymes. It is then possible that the structures of certain self-cleaving ribozymes may be optimized to give maximal catalytic activity, or that entirely new RNA motifs can be made that display significantly faster rates for RNA phosphodiester cleavage.

[00139] Intermolecular cleavage of an RNA substrate by an RNA catalyst that fits the "hammerhead" model was first shown in 1987 (Uhlenbeck, O. C. (1987) *Nature*, 328: 596-600). The RNA catalyst was recovered and reacted with multiple RNA molecules, demonstrating that it was truly catalytic.

[00140] Catalytic RNAs designed based on the "hammerhead" motif have been used to cleave specific target sequences by making appropriate base changes in the catalytic RNA to maintain necessary base pairing with the target sequences (Haseloff and Gerlach, (1988) *Nature*, 334, 585; Walbot and Bruening, (1988) *Nature*, 334, 196; Uhlenbeck, O. C. (1987) *Nature*, 328: 596-600; Koizumi, M., *et al.* (1988) *FEBS Lett.*, 228: 228-230). This has allowed use of the catalytic RNA to cleave specific target sequences and indicates that catalytic RNAs designed according to the "hammerhead" model may possibly cleave specific substrate RNAs in vivo. (see Haseloff and Gerlach, (1988) *Nature*, 334, 585; Walbot and Bruening, (1988) *Nature*, 334, 196; Uhlenbeck, O. C. (1987) *Nature*, 328: 596-600).

[00141] RNA interference (RNAi) has become a powerful tool for modulating gene expression in mammals and mammalian cells. This approach requires the delivery of small interfering RNA (siRNA) either as RNA itself or as DNA, using an expression plasmid or virus and the coding sequence for small hairpin RNAs that are processed to siRNAs. This system enables efficient transport of the pre-siRNAs to the cytoplasm where they are active and permit the use of regulated and tissue specific promoters for gene expression.

[00142] In a preferred embodiment, an oligonucleotide or antisense compound comprises an oligomer or polymer of ribonucleic acid (RNA) and/or deoxyribonucleic acid (DNA), or a mimetic, chimera, analog or homolog thereof. This term includes oligonucleotides composed of naturally occurring nucleotides, sugars and covalent internucleoside (backbone) linkages as

well as oligonucleotides having non-naturally occurring portions which function similarly. Such modified or substituted oligonucleotides are often desired over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for a target nucleic acid and increased stability in the presence of nucleases.

5 [00143] According to the present invention, the oligonucleotides or "antisense compounds" include antisense oligonucleotides (e.g. RNA, DNA, mimetic, chimera, analog or homolog thereof), ribozymes, external guide sequence (EGS) oligonucleotides, siRNA compounds, single- or double-stranded RNA interference (RNAi) compounds such as siRNA compounds, saRNA, aRNA, and other oligomeric compounds which hybridize to at least a portion of the  
10 target nucleic acid and modulate its function. As such, they may be DNA, RNA, DNA-like, RNA-like, or mixtures thereof, or may be mimetics of one or more of these. These compounds may be single-stranded, double-stranded, circular or hairpin oligomeric compounds and may contain structural elements such as internal or terminal bulges, mismatches or loops. Antisense compounds are routinely prepared linearly but can be joined  
15 or otherwise prepared to be circular and/or branched. Antisense compounds can include constructs such as, for example, two strands hybridized to form a wholly or partially double-stranded compound or a single strand with sufficient self-complementarity to allow for hybridization and formation of a fully or partially double-stranded compound. The two strands can be linked internally leaving free 3' or 5' termini or can be linked to form a  
20 continuous hairpin structure or loop. The hairpin structure may contain an overhang on either the 5' or 3' terminus producing an extension of single stranded character. The double stranded compounds optionally can include overhangs on the ends. Further modifications can include conjugate groups attached to one of the termini, selected nucleotide positions, sugar positions or to one of the internucleoside linkages. Alternatively, the two strands can be linked via a  
25 non-nucleic acid moiety or linker group. When formed from only one strand, dsRNA can take the form of a self-complementary hairpin-type molecule that doubles back on itself to form a duplex. Thus, the dsRNAs can be fully or partially double stranded. Specific modulation of gene expression can be achieved by stable expression of dsRNA hairpins in transgenic cell lines (Hammond *et al.*, (1991) *Nat. Rev. Genet.*, 2, 110-119; Matzke *et al.*, (2001) *Curr. Opin.*  
30 *Genet. Dev.*, 11, 221-227; Sharp, (2001) *Genes Dev.*, 15, 485-490). When formed from two strands, or a single strand that takes the form of a self-complementary hairpin-type molecule doubled back on itself to form a duplex, the two strands (or duplex-forming regions of a single strand) are complementary RNA strands that base pair in Watson-Crick fashion.

[00144] Once introduced to a system, the compounds of the invention may elicit the action of one or more enzymes or structural proteins to effect cleavage or other modification of the target nucleic acid or may work via occupancy-based mechanisms. In general, nucleic acids (including oligonucleotides) may be described as "DNA-like" (i.e., generally having one or more 2'-deoxy sugars and, generally, T rather than U bases) or "RNA-like" (i.e., generally having one or more 2'-hydroxyl or 2'-modified sugars and, generally U rather than T bases). Nucleic acid helices can adopt more than one type of structure, most commonly the A- and B-forms. It is believed that, in general, oligonucleotides which have B-form-like structure are "DNA-like" and those which have A-formlike structure are "RNA-like." In some (chimeric) embodiments, an antisense compound may contain both A- and B-form regions.

[00145] The antisense compounds in accordance with this invention can comprise an antisense portion from about 5 to about 80 nucleotides (i.e. from about 5 to about 80 linked nucleosides) in length. This refers to the length of the antisense strand or portion of the antisense compound. In other words, a single-stranded antisense compound of the invention comprises from 5 to about 80 nucleotides, and a double-stranded antisense compound of the invention (such as a dsRNA, for example) comprises a sense and an antisense strand or portion of 5 to about 80 nucleotides in length. One of ordinary skill in the art will appreciate that this comprehends antisense portions of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80 nucleotides in length, or any range therewithin.

[00146] In one embodiment, the antisense compounds of the invention have antisense portions of 10 to 50 nucleotides in length. One having ordinary skill in the art will appreciate that this embodies oligonucleotides having antisense portions of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides in length, or any range therewithin. In some embodiments, the oligonucleotides are 15 nucleotides in length.

[00147] In one embodiment, the antisense or oligonucleotide compounds of the invention have antisense portions of 12 or 13 to 30 nucleotides in length. One having ordinary skill in the art will appreciate that this embodies antisense compounds having antisense portions of 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 nucleotides in length, or any range therewithin.

[00148] In another preferred embodiment, the oligomeric compounds of the present invention also include variants in which a different base is present at one or more of the nucleotide positions in the compound. For example, if the first nucleotide is an adenosine, variants may be produced which contain thymidine, guanosine or cytidine at this position. This may be done at any of the positions of the antisense or dsRNA compounds. These compounds are then tested using the methods described herein to determine their ability to inhibit expression of a target nucleic acid.

[00149] In some embodiments, homology, sequence identity or complementarity, between the antisense compound and target is from about 40% to about 60%. In some embodiments, homology, sequence identity or complementarity, is from about 60% to about 70%. In some embodiments, homology, sequence identity or complementarity, is from about 70% to about 80%. In some embodiments, homology, sequence identity or complementarity, is from about 80% to about 90%. In some embodiments, homology, sequence identity or complementarity, is about 90%, about 92%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100%.

[00150] In another preferred embodiment, the antisense oligonucleotides, such as for example, nucleic acid molecules set forth in SEQ ID NOS: 3 to 6 comprise one or more substitutions or modifications. In one embodiment, the nucleotides are substituted with locked nucleic acids (LNA).

[00151] In another preferred embodiment, the oligonucleotides target one or more regions of the nucleic acid molecules sense and/or antisense of coding and/or non-coding sequences associated with NRF2 and the sequences set forth as SEQ ID NOS: 1, 3. The oligonucleotides are also targeted to overlapping regions of SEQ ID NOS: 1, 3.

[00152] Certain preferred oligonucleotides of this invention are chimeric oligonucleotides. "Chimeric oligonucleotides" or "chimeras," in the context of this invention, are oligonucleotides which contain two or more chemically distinct regions, each made up of at least one nucleotide. These oligonucleotides typically contain at least one region of modified nucleotides that confers one or more beneficial properties (such as, for example, increased nuclease resistance, increased uptake into cells, increased binding affinity for the target) and a region that is a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target,



thereby greatly enhancing the efficiency of antisense modulation of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected  
5 by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art. In one preferred embodiment, a chimeric oligonucleotide comprises at least one region modified to increase target binding affinity, and, usually, a region that acts as a substrate for RNase H. Affinity of an oligonucleotide for its target (in this case, a nucleic acid encoding ras) is routinely determined by measuring the  $T_m$  of an oligonucleotide/target pair,  
10 which is the temperature at which the oligonucleotide and target dissociate; dissociation is detected spectrophotometrically. The higher the  $T_m$ , the greater is the affinity of the oligonucleotide for the target.

[00153] Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides  
15 and/or oligonucleotides mimetics as described above. Such; compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures comprise, but are not limited to, US patent nos. 5,013,830; 5,149,797; 5, 220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, each of which is herein incorporated by  
20 reference.

[00154] In another preferred embodiment, the region of the oligonucleotide which is modified comprises at least one nucleotide modified at the 2' position of the sugar, most preferably a 2'-Oalkyl, 2'-O-alkyl-O-alkyl or 2'-fluoro-modified nucleotide. In other preferred embodiments, RNA modifications include 2'-fluoro, 2'-amino and 2' O-methyl modifications on the ribose of  
25 pyrimidines, abasic residues or an inverted base at the 3' end of the RNA. Such modifications are routinely incorporated into oligonucleotides and these oligonucleotides have been shown to have a higher  $T_m$  (i.e., higher target binding affinity) than; 2'-deoxyoligonucleotides against a given target. The effect of such increased affinity is to greatly enhance RNAi oligonucleotide inhibition of gene expression. RNase H is a cellular endonuclease that  
30 cleaves the RNA strand of RNA:DNA duplexes; activation of this enzyme therefore results in cleavage of the RNA target, and thus can greatly enhance the efficiency of RNAi inhibition. Cleavage of the RNA target can be routinely demonstrated by gel electrophoresis. In another preferred embodiment, the chimeric oligonucleotide is also modified to enhance nuclease

resistance. Cells contain a variety of exo- and endo-nucleases which can degrade nucleic acids. A number of nucleotide and nucleoside modifications have been shown to make the oligonucleotide into which they are incorporated more resistant to nuclease digestion than the native oligodeoxynucleotide. Nuclease resistance is routinely measured by incubating oligonucleotides with cellular extracts or isolated nuclease solutions and measuring the extent of intact oligonucleotide remaining over time, usually by gel electrophoresis. Oligonucleotides which have been modified to enhance their nuclease resistance survive intact for a longer time than unmodified oligonucleotides. A variety of oligonucleotide modifications have been demonstrated to enhance or confer nuclease resistance. Oligonucleotides which contain at least one phosphorothioate modification are presently more preferred. In some cases, oligonucleotide modifications which enhance target binding affinity are also, independently, able to enhance nuclease resistance. Some desirable modifications can be found in De Mesmaeker *et al.* (1995) *Acc. Chem. Res.*, 28:366-374.

**[00155]** Specific examples of some preferred oligonucleotides envisioned for this invention include those comprising modified backbones, for example, phosphorothioates, phosphotriesters, methyl phosphonates, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages. Most preferred are oligonucleotides with phosphorothioate backbones and those with heteroatom backbones, particularly  $\text{CH}_2\text{--NH--O--CH}_2$ ,  $\text{CH}_2\text{--N(CH}_3\text{)--O--CH}_2$  [known as a methylene(methylimino) or MMI backbone],  $\text{CH}_2\text{--O--N(CH}_3\text{)--CH}_2$ ,  $\text{CH}_2\text{--N(CH}_3\text{)--N(CH}_3\text{)--CH}_2$  and  $\text{O--N(CH}_3\text{)--CH}_2\text{--CH}_2$  backbones, wherein the native phosphodiester backbone is represented as  $\text{O--P--O--CH}_2$ . The amide backbones disclosed by De Mesmaeker *et al.* (1995) *Acc. Chem. Res.* 28:366-374 are also preferred. Also preferred are oligonucleotides having morpholino backbone structures (Summerton and Weller, U.S. Pat. No. 5,034,506). In other preferred embodiments, such as the peptide nucleic acid (PNA) backbone, the phosphodiester backbone of the oligonucleotide is replaced with a polyamide backbone, the nucleotides being bound directly or indirectly to the aza nitrogen atoms of the polyamide backbone (Nielsen *et al.* (1991) *Science* 254, 1497). Oligonucleotides may also comprise one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH, SH, SCH<sub>3</sub>, F, OCN, OCH<sub>3</sub>, OCH<sub>3</sub>, OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub> or O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub> where n is from 1 to about 10; C<sub>1</sub> to C<sub>10</sub> lower alkyl, alkoxyalkoxy, substituted lower alkyl, alkaryl or aralkyl; Cl; Br; CN; CF<sub>3</sub>; OCF<sub>3</sub>; O--, S--, or N-alkyl; O--, S--, or N-alkenyl; SOCH<sub>3</sub>; SO<sub>2</sub>CH<sub>3</sub>; ONO<sub>2</sub>; NO<sub>2</sub>; N<sub>3</sub>; NH<sub>2</sub>;

heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy [2'-O-CH<sub>2</sub> CH<sub>2</sub> OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl)] (Martin *et al.*, (1995) *Helv. Chim. Acta*, 78, 486). Other preferred modifications include 2'-methoxy (2'-O--CH<sub>3</sub>), 2'- propoxy (2'-OCH<sub>2</sub> CH<sub>2</sub>CH<sub>3</sub>) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyls in place of the pentofuranosyl group.

[00156] Oligonucleotides may also include, additionally or alternatively, nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleotides include adenine (A), guanine (G), thymine (T), cytosine (C) and uracil (U). Modified nucleotides include nucleotides found only infrequently or transiently in natural nucleic acids, e.g., hypoxanthine, 6-methyladenine, 5-Me pyrimidines, particularly 5-methylcytosine (also referred to as 5-methyl-2' deoxycytosine and often referred to in the art as 5-Me-C), 5- hydroxymethylcytosine (HMC), glycosyl HMC and gentobiosyl HMC, as well as synthetic nucleotides, e.g., 2-aminoadenine, 2-(methylamino)adenine, 2-(imidazolylalkyl)adenine, 2- (aminoalkylamino)adenine or other heterosubstituted alkyladenines, 2-thiouracil, 2-thiothymine, 5- bromouracil, 5-hydroxymethyluracil, 8-azaguanine, 7-deazaguanine, N<sup>6</sup> (6-aminohexyl)adenine and 2,6-diaminopurine. (Kornberg, A., DNA Replication, W. H. Freeman & Co., San Francisco, 1980, pp75-77; Gebeyehu, G., (1987) *et al. Nucl. Acids Res.* 15:4513). A "universal" base known in the art, e.g., inosine, may be included. 5-Me-C substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C. (Sanghvi, Y. S., in Crooke, S. T. and Lebleu, B., eds., Antisense Research and Applications, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions.

[00157] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety, a cholesteryl moiety (Letsinger *et al.*, (1989) *Proc. Natl. Acad. Sci. USA* 86, 6553), cholic acid (Manoharan *et al.* (1994) *Bioorg. Med. Chem. Let.*

4, 1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660, 306; Manoharan *et al.* (1993) *Bioorg. Med. Chem. Lett.* 3, 2765), a thiocholesterol (Oberhauser *et al.*, (1992) *Nucl. Acids Res.* 20, 533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras *et al.* *EMBO J.* 1991, 10, 111; Kabanov *et al.* (1990) *FEBS Lett.* 259, 327; Svinarchuk *et al.* (1993) *Biochimie* 75, 49), a phospholipid, e.g., dihexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero- 3-H-phosphonate (Manoharan *et al.* (1995) *Tetrahedron Lett.* 36, 3651; Shea *et al.* (1990) *Nucl. Acids Res.* 18, 3777), a polyamine or a polyethylene glycol chain (Manoharan *et al.* (1995) *Nucleosides & Nucleotides*, 14, 969), or adamantane acetic acid (Manoharan *et al.* (1995) *Tetrahedron Lett.* 36, 3651). Oligonucleotides comprising lipophilic moieties, and methods for preparing such oligonucleotides are known in the art, for example, U.S. Pat. Nos. 5,138,045, 5,218,105 and 5,459,255.

[00158] It is not necessary for all positions in a given oligonucleotide to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single oligonucleotide or even at within a single nucleoside within an oligonucleotide. The present invention also includes oligonucleotides which are chimeric oligonucleotides as hereinbefore defined.

[00159] In another embodiment, the nucleic acid molecule of the present invention is conjugated with another moiety including but not limited to abasic nucleotides, polyether, polyamine, polyamides, peptides, carbohydrates, lipid, or polyhydrocarbon compounds. Those skilled in the art will recognize that these molecules can be linked to one or more of any nucleotides comprising the nucleic acid molecule at several positions on the sugar, base or phosphate group.

[00160] The oligonucleotides used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including Applied Biosystems. Any other means for such synthesis may also be employed; the actual synthesis of the oligonucleotides is well within the talents of one of ordinary skill in the art. It is also well known to use similar techniques to prepare other oligonucleotides such as the phosphorothioates and alkylated derivatives. It is also well known to use similar techniques and commercially available modified amidites and controlled-pore glass (CPG) products such as biotin, fluorescein, acridine or psoralen-modified amidites and/or CPG (available from Glen Research, Sterling

VA) to synthesize fluorescently labeled, biotinylated or other modified oligonucleotides such as cholesterol-modified oligonucleotides.

[00161] In accordance with the invention, use of modifications such as the use of LNA monomers to enhance the potency, specificity and duration of action and broaden the routes of administration of oligonucleotides comprised of current chemistries such as MOE, ANA, FANA, PS etc (Uhlman, *et al.* (2000) *Current Opinions in Drug Discovery & Development* Vol. 3 No 2). This can be achieved by substituting some of the monomers in the current oligonucleotides by LNA monomers. The LNA modified oligonucleotide may have a size similar to the parent compound or may be larger or preferably smaller. It is preferred that such LNA-modified oligonucleotides contain less than about 70%, more preferably less than about 60%, most preferably less than about 50% LNA monomers and that their sizes are between about 5 and 25 nucleotides, more preferably between about 12 and 20 nucleotides.

[00162] Preferred modified oligonucleotide backbones comprise, but not limited to, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates comprising 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

[00163] Representative United States patents that teach the preparation of the above phosphorus containing linkages comprise, but are not limited to, US patent nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference.

[00164] Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These comprise those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane

backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

[00165] Representative United States patents that teach the preparation of the above oligonucleosides comprise, but are not limited to, US patent nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264, 562; 5, 264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596, 086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623, 070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, each of which is herein incorporated by reference.

[00166] In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds comprise, but are not limited to, US patent nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference . Further teaching of PNA compounds can be found in Nielsen, *et al.* (1991) *Science* 254, 1497-1500.

[00167] In another preferred embodiment of the invention the oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular- CH<sub>2</sub>-NH-O-CH<sub>2</sub>-, -CH<sub>2</sub>-N (CH<sub>3</sub>)-O-CH<sub>2</sub>-known as a methylene (methylimino) or MMI backbone, - CH<sub>2</sub>-O-N (CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>N(CH<sub>3</sub>)-N(CH<sub>3</sub>) CH<sub>2</sub>-and-O-N(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>2</sub>- wherein the native phosphodiester backbone is represented as-O-P-O-CH<sub>2</sub>- of the above referenced US patent no. 5,489,677, and the amide backbones of the above referenced US patent no. 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced US patent no. 5,034,506.

[00168] Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C to CO alkyl or C2 to CO alkenyl and alkynyl. Particularly preferred are O (CH<sub>2</sub>)<sub>n</sub> OmCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>)<sub>2</sub> where n and m can be from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C to CO, (lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification comprises 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2- methoxyethyl) or 2'-MOE) (Martin *et al.*, (1995) *Helv. Chim. Acta*, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification comprises 2'-dimethylaminoethoxyethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples herein below, and 2'- dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'- DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N (CH<sub>2</sub>)<sub>2</sub>.

[00169] Other preferred modifications comprise 2'-methoxy (2'-O CH<sub>3</sub>), 2'-aminopropoxy (2'-O CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures comprise, but are not limited to, US patent nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514, 785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646, 265; 5,658,873; 5,670,633; and 5,700,920, each of which is herein incorporated by reference.

[00170] Oligonucleotides may also comprise nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleotides comprise the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleotides comprise other synthetic and natural

nucleotides such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2- aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudo-uracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylquanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine.

[00171] Further, nucleotides comprise those disclosed in United States Patent No. 3,687,808, those disclosed in 'The Concise Encyclopedia of Polymer Science And Engineering', pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch *et al.*, 'Angewandte Chemie, International Edition', 1991, 30, page 613, and those disclosed by Sanghvi, Y.S., Chapter 15, 'Antisense Research and Applications', pages 289-302, Crooke, S.T. and Lebleu, B. ea., CRC Press, 1993. Certain of these nucleotides are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These comprise 5-substituted pyrimidines, 6- azapyrimidines and N-2, N-6 and O-6 substituted purines, comprising 2-aminopropyladenine, 5- propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds, 'Antisense Research and Applications', CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[00172] Representative United States patents that teach the preparation of the above noted modified nucleotides as well as other modified nucleotides comprise, but are not limited to, US patent nos. 3,687,808, as well as 4,845,205; 5,130,302; 5,134,066; 5,175, 273; 5, 367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,596,091; 5,614,617; 5,750,692, and 5,681,941, each of which is herein incorporated by reference.

[00173] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates, which enhance the activity, cellular distribution, or cellular uptake of the oligonucleotide.



[00174] Such moieties comprise but are not limited to, lipid moieties such as a cholesterol moiety (Letsinger *et al.*, (1989) *Proc. Natl. Acad. Sci. USA*, 86, 6553-6556), cholic acid (Manoharan *et al.*, (1994) *Bioorg. Med. Chem. Lett.*, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan *et al.*, (1992) *Ann. N. Y. Acad. Sci.*, 660, 306-309; Manoharan *et al.*, (1993) *Bioorg. Med. Chem. Lett.*, 3, 2765-2770), a thiocholesterol (Oberhauser *et al.*, (1992) *Nucl. Acids Res.*, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Kabanov *et al.*, (1990) *FEBS Lett.*, 259, 327-330; Svinarchuk *et al.*, (1993) *Biochimie* 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan *et al.*, (1995) *Tetrahedron Lett.*, 36, 3651-3654; Shea *et al.*, (1990) *Nucl. Acids Res.*, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Mancharan *et al.*, (1995) *Nucleosides & Nucleotides*, 14, 969-973), or adamantane acetic acid (Manoharan *et al.*, (1995) *Tetrahedron Lett.*, 36, 3651-3654), a palmityl moiety (Mishra *et al.*, (1995) *Biochim. Biophys. Acta*, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-t oxysterol moiety (Crooke *et al.*, (1996) *J. Pharmacol. Exp. Ther.*, 277, 923-937).

[00175] Representative United States patents that teach the preparation of such oligonucleotides conjugates comprise, but are not limited to, US patent nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552, 538; 5,578,717, 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486, 603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762, 779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082, 830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5, 245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241, 5,391, 723; 5,416,203, 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5, 565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599, 928 and 5,688,941, each of which is herein incorporated by reference.

[00176] *Drug discovery*: The compounds of the present invention can also be applied in the areas of drug discovery and target validation. The present invention comprehends the use of the compounds and preferred target segments identified herein in drug discovery efforts to elucidate relationships that exist between Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides and a disease state, phenotype, or condition. These methods include detecting or modulating Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides comprising contacting a sample, tissue, cell, or organism with the compounds of the present invention, measuring the nucleic acid or protein level of Nuclear factor (erythroid-derived 2)-like 2

(NRF2) polynucleotides and/or a related phenotypic or chemical endpoint at some time after treatment, and optionally comparing the measured value to a non-treated sample or sample treated with a further compound of the invention. These methods can also be performed in parallel or in combination with other experiments to determine the function of unknown genes for the process of target validation or to determine the validity of a particular gene product as a target for treatment or prevention of a particular disease, condition, or phenotype.

*Assessing Up-regulation or Inhibition of Gene Expression:*

[00177] Transfer of an exogenous nucleic acid into a host cell or organism can be assessed by directly detecting the presence of the nucleic acid in the cell or organism. Such detection can be achieved by several methods well known in the art. For example, the presence of the exogenous nucleic acid can be detected by Southern blot or by a polymerase chain reaction (PCR) technique using primers that specifically amplify nucleotide sequences associated with the nucleic acid. Expression of the exogenous nucleic acids can also be measured using conventional methods including gene expression analysis. For instance, mRNA produced from an exogenous nucleic acid can be detected and quantified using a Northern blot and reverse transcription PCR (RT-PCR).

[00178] Expression of RNA from the exogenous nucleic acid can also be detected by measuring an enzymatic activity or a reporter protein activity. For example, antisense modulatory activity can be measured indirectly as a decrease or increase in target nucleic acid expression as an indication that the exogenous nucleic acid is producing the effector RNA. Based on sequence conservation, primers can be designed and used to amplify coding regions of the target genes. Initially, the most highly expressed coding region from each gene can be used to build a model control gene, although any coding or non coding region can be used. Each control gene is assembled by inserting each coding region between a reporter coding region and its poly(A) signal. These plasmids would produce an mRNA with a reporter gene in the upstream portion of the gene and a potential RNAi target in the 3' non-coding region. The effectiveness of individual antisense oligonucleotides would be assayed by modulation of the reporter gene. Reporter genes useful in the methods of the present invention include acetohydroxyacid synthase (AHAS), alkaline phosphatase (AP), beta galactosidase (LacZ), beta glucuronidase (GUS), chloramphenicol acetyltransferase (CAT), green fluorescent protein (GFP), red fluorescent protein (RFP), yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), horseradish peroxidase (HRP), luciferase (Luc), nopaline synthase

(NOS), octopine synthase (OCS), and derivatives thereof. Multiple selectable markers are available that confer resistance to ampicillin, bleomycin, chloramphenicol, gentamycin, hygromycin, kanamycin, lincomycin, methotrexate, phosphinothricin, puromycin, and tetracycline. Methods to determine modulation of a reporter gene are well known in the art, and include, but are not limited to, fluorometric methods (e.g. fluorescence spectroscopy, Fluorescence Activated Cell Sorting (FACS), fluorescence microscopy), antibiotic resistance determination.

[00179] NRF2 protein and mRNA expression can be assayed using methods known to those of skill in the art and described elsewhere herein. For example, immunoassays such as the ELISA can be used to measure protein levels. NRF2 antibodies for ELISAs are available commercially, e.g., from R&D Systems (Minneapolis, MN), Abcam, Cambridge, MA.

[00180] In embodiments, NRF2 expression (e.g., mRNA or protein) in a sample (e.g., cells or tissues in vivo or in vitro) treated using an antisense oligonucleotide of the invention is evaluated by comparison with NRF2 expression in a control sample. For example, expression of the protein or nucleic acid can be compared using methods known to those of skill in the art with that in a mock-treated or untreated sample. Alternatively, comparison with a sample treated with a control antisense oligonucleotide (e.g., one having an altered or different sequence) can be made depending on the information desired. In another embodiment, a difference in the expression of the NRF2 protein or nucleic acid in a treated vs. an untreated sample can be compared with the difference in expression of a different nucleic acid (including any standard deemed appropriate by the researcher, e.g., a housekeeping gene) in a treated sample vs. an untreated sample.

[00181] Observed differences can be expressed as desired, e.g., in the form of a ratio or fraction, for use in a comparison with control. In embodiments, the level of NRF2 mRNA or protein, in a sample treated with an antisense oligonucleotide of the present invention, is increased or decreased by about 1.25-fold to about 10-fold or more relative to an untreated sample or a sample treated with a control nucleic acid. In embodiments, the level of NRF2 mRNA or protein is increased or decreased by at least about 1.25-fold, at least about 1.3-fold, at least about 1.4-fold, at least about 1.5-fold, at least about 1.6-fold, at least about 1.7-fold, at least about 1.8-fold, at least about 2-fold, at least about 2.5-fold, at least about 3-fold, at least about 3.5-fold, at least about 4-fold, at least about 4.5-fold, at least about 5-fold, at least about 5.5-fold, at least about 6-fold, at least about 6.5-fold, at least about 7-fold, at least about 7.5-

fold, at least about 8-fold, at least about 8.5-fold, at least about 9-fold, at least about 9.5-fold, or at least about 10-fold or more.

*Kits, Research Reagents, Diagnostics, and Therapeutics*

[00182] The compounds of the present invention can be utilized for diagnostics, therapeutics, and prophylaxis, and as research reagents and components of kits. Furthermore, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes or to distinguish between functions of various members of a biological pathway.

[00183] For use in kits and diagnostics and in various biological systems, the compounds of the present invention, either alone or in combination with other compounds or therapeutics, are useful as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

[00184] As used herein the term "biological system" or "system" is defined as any organism, cell, cell culture or tissue that expresses, or is made competent to express products of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) genes. These include, but are not limited to, humans, transgenic animals, cells, cell cultures, tissues, xenografts, transplants and combinations thereof.

[00185] As one non limiting example, expression patterns within cells or tissues treated with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds that affect expression patterns.

[00186] Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, (2000) *FEBS Lett.*, 480, 17-24; Celis, *et al.*, (2000) *FEBS Lett.*, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, *et al.*, (2000) *Drug Discov. Today*, 5, 415- 425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, (1999) *Methods Enzymol.*, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, *et al.*, (2000) *Proc. Natl. Acad. Sci. U.S.A.*, 97, 1976-81), protein arrays and proteomics (Celis, *et al.*, (2000) *FEBS Lett.*, 480, 2-16; Jungblut, *et al.*, *Electrophoresis*, 1999, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, *et al.*,

FEBS Lett., 2000, 480, 2-16; Larsson, et al., J. Biotechnol., 2000, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, *et al.*, (2000) Anal. Biochem. 286, 91-98; Larson, *et al.*, (2000) *Cytometry* 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, (2000) *Curr. Opin. Microbiol.* 3, 316-21), comparative genomic hybridization (Carulli, *et al.*, (1998) *J. Cell Biochem. Suppl.*, 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, (1999) *Eur. J. Cancer*, 35, 1895-904) and mass spectrometry methods (To, Comb. (2000) *Chem. High Throughput Screen*, 3, 235-41).

[00187] The compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding Nuclear factor (erythroid-derived 2)-like 2 (NRF2). For example, oligonucleotides that hybridize with such efficiency and under such conditions as disclosed herein as to be effective Nuclear factor (erythroid-derived 2)-like 2 (NRF2) modulators are effective primers or probes under conditions favoring gene amplification or detection, respectively. These primers and probes are useful in methods requiring the specific detection of nucleic acid molecules encoding Nuclear factor (erythroid-derived 2)-like 2 (NRF2) and in the amplification of said nucleic acid molecules for detection or for use in further studies of Nuclear factor (erythroid-derived 2)-like 2 (NRF2). Hybridization of the antisense oligonucleotides, particularly the primers and probes, of the invention with a nucleic acid encoding Nuclear factor (erythroid-derived 2)-like 2 (NRF2) can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabeling of the oligonucleotide, or any other suitable detection means. Kits using such detection means for detecting the level of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in a sample may also be prepared.

[00188] The specificity and sensitivity of antisense are also harnessed by those of skill in the art for therapeutic uses. Antisense compounds have been employed as therapeutic moieties in the treatment of disease states in animals, including humans. Antisense oligonucleotide drugs have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that antisense compounds can be useful therapeutic modalities that can be configured to be useful in treatment regimes for the treatment of cells, tissues and animals, especially humans.

[00189] For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides is treated by administering antisense compounds in

accordance with this invention. For example, in one non-limiting embodiment, the methods comprise the step of administering to the animal in need of treatment, a therapeutically effective amount of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) modulator. The Nuclear factor (erythroid-derived 2)-like 2 (NRF2) modulators of the present invention effectively modulate the activity of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) or modulate the expression of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) protein. In one embodiment, the activity or expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in an animal is inhibited by about 10% as compared to a control. Preferably, the activity or expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in an animal is inhibited by about 30%. More preferably, the activity or expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in an animal is inhibited by 50% or more. Thus, the oligomeric compounds modulate expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) mRNA by at least 10%, by at least 50%, by at least 25%, by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 75%, by at least 80%, by at least 85%, by at least 90%, by at least 95%, by at least 98%, by at least 99%, or by 100% as compared to a control.

[00190] In one embodiment, the activity or expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) and/or in an animal is increased by about 10% as compared to a control. Preferably, the activity or expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in an animal is increased by about 30%. More preferably, the activity or expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in an animal is increased by 50% or more. Thus, the oligomeric compounds modulate expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) mRNA by at least 10%, by at least 50%, by at least 25%, by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 75%, by at least 80%, by at least 85%, by at least 90%, by at least 95%, by at least 98%, by at least 99%, or by 100% as compared to a control.

[00191] For example, the reduction of the expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) may be measured in serum, blood, adipose tissue, liver or any other body fluid, tissue or organ of the animal. Preferably, the cells contained within said fluids, tissues or organs being analyzed contain a nucleic acid molecule encoding Nuclear factor (erythroid-derived 2)-like 2 (NRF2) peptides and/or the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) protein itself.

[00192] The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of a compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the compounds and methods of the invention may also be useful prophylactically.

5 *Conjugates*

[00193] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates that enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. These moieties or conjugates can include conjugate groups covalently bound to functional groups such as primary or  
10 secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugate groups include cholesterol, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines,  
15 coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve uptake, enhance resistance to degradation, and/or strengthen sequence-specific hybridization with the target nucleic acid. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve uptake, distribution, metabolism or excretion of the compounds of the present invention.  
20 Representative conjugate groups are disclosed in International Patent Application No. PCT/US92/09196, filed Oct. 23, 1992, and U.S. Pat. No. 6,287,860, which are incorporated herein by reference. Conjugate moieties include, but are not limited to, lipid moieties such as a cholesterol moiety, cholic acid, a thioether, e.g., hexyl-5- tritylthiol, a thiocholesterol, an aliphatic chain, e.g., dodecandiol or undecyl residues, a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-Hphosphonate, a polyamine  
25 or a polyethylene glycol chain, or adamantane acetic acid, a palmityl moiety, or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety. Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen,  
30 dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic.

[00194] Representative United States patents that teach the preparation of such oligonucleotides conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941.

#### *Formulations*

[00195] The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as foreexample, liposomes, receptor-targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption-assisting formulations include, but are not limited to, U.S. Pat. Nos. 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,165; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

[00196] Although, the antisense oligonucleotides do not need to be administered in the context of a vector in order to modulate a target expression and/or function, embodiments of the invention relates to expression vector constructs for the expression of antisense oligonucleotides, comprising promoters, hybrid promoter gene sequences and possess a strong constitutive promoter activity, or a promoter activity which can be induced in the desired case.

[00197] In an embodiment, invention practice involves administering at least one of the foregoing antisense oligonucleotides with a suitable nucleic acid delivery system. In one embodiment, that system includes a non-viral vector operably linked to the polynucleotide. Examples of such nonviral vectors include the oligonucleotide alone (e.g. any one or more of



SEQ ID NOS: 4 to 6) or in combination with a suitable protein, polysaccharide or lipid formulation.

[00198] Additionally suitable nucleic acid delivery systems include viral vector, typically sequence from at least one of an adenovirus, adenovirus-associated virus (AAV), helper-dependent adenovirus, retrovirus, or hemagglutinin virus of Japan-liposome (HVJ) complex. Preferably, the viral vector comprises a strong eukaryotic promoter operably linked to the polynucleotide e.g., a cytomegalovirus (CMV) promoter.

[00199] Additionally preferred vectors include viral vectors, fusion proteins and chemical conjugates. Retroviral vectors include Moloney murine leukemia viruses and HIV-based viruses. One preferred HIV-based viral vector comprises at least two vectors wherein the gag and pol genes are from an HIV genome and the env gene is from another virus. DNA viral vectors are preferred. These vectors include pox vectors such as orthopox or avipox vectors, herpesvirus vectors such as a herpes simplex I virus (HSV) vector [Geller, A.I. *et al.*, (1995) *J. Neurochem*, 64: 487; Lim, F., *et al.*, in DNA Cloning: Mammalian Systems, D. Glover, Ed. (Oxford Univ. Press, Oxford England) (1995); Geller, A.I. *et al.*, (1993) *Proc Natl. Acad. Sci.*: U.S.A.:90 7603; Geller, A.I., *et al.*, (1990) *Proc Natl. Acad. Sci USA*: 87:1149], Adenovirus Vectors (LeGal LaSalle *et al.*, *Science*, 259:988 (1993); Davidson, *et al.*, (1993) *Nat. Genet.* 3: 219; Yang, *et al.*, (1995) *J. Virol.* 69: 2004) and Adeno-associated Virus Vectors (Kaplitt, M.G., *et al.*, (1994) *Nat. Genet.* 8:148).

[00200] The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof.

[00201] The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto. For oligonucleotides, preferred examples of pharmaceutically acceptable salts and their uses are further described in U.S. Pat. No. 6,287,860, which is incorporated herein by reference.

[00202] The present invention also includes pharmaceutical compositions and formulations that include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local

or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration.

[00203] For treating tissues in the central nervous system, administration can be made by, e.g., injection or infusion into the cerebrospinal fluid. Administration of antisense RNA into cerebrospinal fluid is described, e.g., in U.S. Pat. App. Pub. No. 2007/0117772, "Methods for slowing familial ALS disease progression," incorporated herein by reference in its entirety.

[00204] When it is intended that the antisense oligonucleotide of the present invention be administered to cells in the central nervous system, administration can be with one or more agents capable of promoting penetration of the subject antisense oligonucleotide across the blood-brain barrier. Injection can be made, e.g., in the entorhinal cortex or hippocampus. Delivery of neurotrophic factors by administration of an adenovirus vector to motor neurons in muscle tissue is described in, e.g., U.S. Pat. No. 6,632,427, "Adenoviral-vector-mediated gene transfer into medullary motor neurons," incorporated herein by reference. Delivery of vectors directly to the brain, e.g., the striatum, the thalamus, the hippocampus, or the substantia nigra, is known in the art and described, e.g., in U.S. Pat. No. 6,756,523, "Adenovirus vectors for the transfer of foreign genes into cells of the central nervous system particularly in brain," incorporated herein by reference. Administration can be rapid as by injection or made over a period of time as by slow infusion or administration of slow release formulations.

[00205] The subject antisense oligonucleotides can also be linked or conjugated with agents that provide desirable pharmaceutical or pharmacodynamic properties. For example, the antisense oligonucleotide can be coupled to any substance, known in the art to promote penetration or transport across the blood-brain barrier, such as an antibody to the transferrin receptor, and administered by intravenous injection. The antisense compound can be linked with a viral vector, for example, that makes the antisense compound more effective and/or increases the transport of the antisense compound across the blood-brain barrier. Osmotic blood brain barrier disruption can also be accomplished by, e.g., infusion of sugars including,

but not limited to, meso erythritol, xylitol, D(+) galactose, D(+) lactose, D(+) xylose, dulcitol, myo-inositol, L(-) fructose, D(-) mannitol, D(+) glucose, D(+) arabinose, D(-) arabinose, cellobiose, D(+) maltose, D(+) raffinose, L(+) rhamnose, D(+) melibiose, D(-) ribose, adonitol, D(+) arabitol, L(-) arabitol, D(+) fucose, L(-) fucose, D(-) lyxose, L(+) lyxose, and  
5 L(-) lyxose, or amino acids including, but not limited to, glutamine, lysine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glycine, histidine, leucine, methionine, phenylalanine, proline, serine, threonine, tyrosine, valine, and taurine. Methods and materials for enhancing blood brain barrier penetration are described, e.g., in U. S. Patent No. 4,866,042, "Method for the delivery of genetic material across the blood brain barrier,"  
10 6,294,520, "Material for passage through the blood-brain barrier," and 6,936,589, "Parenteral delivery systems," all incorporated herein by reference in their entirety.

[00206] The subject antisense compounds may be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, for example, liposomes, receptor-targeted molecules, oral, rectal, topical or other formulations,  
15 for assisting in uptake, distribution and/or absorption. For example, cationic lipids may be included in the formulation to facilitate oligonucleotide uptake. One such composition shown to facilitate uptake is LIPOFECTIN (available from GIBCO-BRL, Bethesda, MD).

[00207] Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration. Pharmaceutical compositions and formulations for  
20 topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

[00208] The pharmaceutical formulations of the present invention, which may conveniently  
25 be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if  
30 necessary, shaping the product.

[00209] The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups,

soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances that increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also  
5 contain stabilizers.

[00210] Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, foams and liposome-containing formulations. The pharmaceutical compositions and formulations of the present invention may comprise one or more penetration enhancers, carriers, excipients or other active or inactive ingredients.

10 [00211] Emulsions are typically heterogeneous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. Emulsions may contain additional components in addition to the dispersed phases, and the active drug that may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Microemulsions are included as an embodiment of the present invention. Emulsions and their uses are well  
15 known in the art and are further described in U.S. Pat. No. 6,287,860.

[00212] Formulations of the present invention include liposomal formulations. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers. Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior  
20 that contains the composition to be delivered. Cationic liposomes are positively charged liposomes that are believed to interact with negatively charged DNA molecules to form a stable complex. Liposomes that are pH-sensitive or negatively-charged are believed to entrap DNA rather than complex with it. Both cationic and noncationic liposomes have been used to deliver DNA to cells.

25 [00213] Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids. When incorporated into liposomes, these specialized lipids result in liposomes with enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome  
30 comprises one or more glycolipids or is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. Liposomes and their uses are further described in U.S. Pat. No. 6,287,860.

[00214] The pharmaceutical formulations and compositions of the present invention may also include surfactants. The use of surfactants in drug products, formulations and in emulsions is well known in the art. Surfactants and their uses are further described in U.S. Pat. No. 6,287,860, which is incorporated herein by reference.

5 [00215] In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs. Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating  
10 agents, and non-chelating nonsurfactants. Penetration enhancers and their uses are further described in U.S. Pat. No. 6,287,860, which is incorporated herein by reference.

[00216] One of skill in the art will recognize that formulations are routinely designed according to their intended use, i.e. route of administration.

[00217] Preferred formulations for topical administration include those in which the  
15 oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoyl-phosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl  
20 DOTAP and dioleoyl-phosphatidyl ethanolamine DOTMA).

[00218] For topical or other administration, oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters, pharmaceutically acceptable salts thereof, and  
25 their uses are further described in U.S. Pat. No. 6,287,860.

[00219] Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitabets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those  
30 in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts and fatty

acids and their uses are further described in U.S. Pat. No. 6,287,860, which is incorporated herein by reference. Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally, in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents and their uses are further described in U.S. Pat. No. 6,287,860, which is incorporated herein by reference.

[00220] Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions that may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

[00221] Certain embodiments of the invention provide pharmaceutical compositions containing one or more oligomeric compounds and one or more other chemotherapeutic agents that function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to cancer chemotherapeutic drugs such as daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bischloroethyl- nitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5- azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclo-phosphoramidate, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUDR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. Combinations of

antisense compounds and other non-antisense drugs are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

[00222] In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. For example, the first target may be a particular antisense sequence of Nuclear factor (erythroid-derived 2)-like 2 (NRF2), and the second target may be a region from another nucleotide sequence. Alternatively, compositions of the invention may contain two or more antisense compounds targeted to different regions of the same Nuclear factor (erythroid-derived 2)-like 2 (NRF2) nucleic acid target. Numerous examples of antisense compounds are illustrated herein and others may be selected from among suitable compounds known in the art. Two or more combined compounds may be used together or sequentially.

*Dosing:*

[00223] The formulation of therapeutic compositions and their subsequent administration (dosing) is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC50s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01  $\mu$ g to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01  $\mu$ g to 100 g per kg of body weight, once or more daily, to once every 20 years.

[00224] In embodiments, a patient is treated with a dosage of drug that is at least about 1, at least about 2, at least about 3, at least about 4, at least about 5, at least about 6, at least about

7, at least about 8, at least about 9, at least about 10, at least about 15, at least about 20, at least about 25, at least about 30, at least about 35, at least about 40, at least about 45, at least about 50, at least about 60, at least about 70, at least about 80, at least about 90, or at least about 100 mg/kg body weight. Certain injected dosages of antisense oligonucleotides are described, e.g., in U.S. Pat. No. 7,563,884, "Antisense modulation of PTP1B expression," incorporated herein by reference in its entirety.

[00225] While various embodiments of the present invention have been described above, it should be understood that they have been presented by way of example only, and not limitation. Numerous changes to the disclosed embodiments can be made in accordance with the disclosure herein without departing from the spirit or scope of the invention. Thus, the breadth and scope of the present invention should not be limited by any of the above described embodiments.

[00226] All documents mentioned herein are incorporated herein by reference. All publications and patent documents cited in this application are incorporated by reference for all purposes to the same extent as if each individual publication or patent document were so individually denoted. By their citation of various references in this document, Applicants do not admit any particular reference is "prior art" to their invention. Embodiments of inventive compositions and methods are illustrated in the following examples.

### EXAMPLES

[00227] The following non-limiting Examples serve to illustrate selected embodiments of the invention. It will be appreciated that variations in proportions and alternatives in elements of the components shown will be apparent to those skilled in the art and are within the scope of embodiments of the present invention.

*Example 1: Design of antisense oligonucleotides specific for a nucleic acid molecule antisense to a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) and/or a sense strand of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide*

[00228] As indicated above the term "oligonucleotide specific for" or "oligonucleotide targets" refers to an oligonucleotide having a sequence (i) capable of forming a stable complex with a portion of the targeted gene, or (ii) capable of forming a stable duplex with a portion of an mRNA transcript of the targeted gene.



[00229] Selection of appropriate oligonucleotides is facilitated by using computer programs that automatically align nucleic acid sequences and indicate regions of identity or homology. Such programs are used to compare nucleic acid sequences obtained, for example, by searching databases such as GenBank or by sequencing PCR products. Comparison of nucleic acid sequences from a range of species allows the selection of nucleic acid sequences that display an appropriate degree of identity between species. In the case of genes that have not been sequenced, Southern blots are performed to allow a determination of the degree of identity between genes in target species and other species. By performing Southern blots at varying degrees of stringency, as is well known in the art, it is possible to obtain an approximate measure of identity. These procedures allow the selection of oligonucleotides that exhibit a high degree of complementarity to target nucleic acid sequences in a subject to be controlled and a lower degree of complementarity to corresponding nucleic acid sequences in other species. One skilled in the art will realize that there is considerable latitude in selecting appropriate regions of genes for use in the present invention.

[00230] An antisense compound is "specifically hybridizable" when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a modulation of function and/or activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and under conditions in which assays are performed in the case of in vitro assays

[00231] The hybridization properties of the oligonucleotides described herein can be determined by one or more in vitro assays as known in the art. For example, the properties of the oligonucleotides described herein can be obtained by determination of binding strength between the target natural antisense and a potential drug molecules using melting curve assay.

[00232] The binding strength between the target natural antisense and a potential drug molecule (Molecule) can be estimated using any of the established methods of measuring the strength of intermolecular interactions, for example, a melting curve assay.

[00233] Melting curve assay determines the temperature at which a rapid transition from double-stranded to single-stranded conformation occurs for the natural antisense/Molecule complex. This temperature is widely accepted as a reliable measure of the interaction strength between the two molecules.

[00234] A melting curve assay can be performed using a cDNA copy of the actual natural antisense RNA molecule or a synthetic DNA or RNA nucleotide corresponding to the binding site of the Molecule. Multiple kits containing all necessary reagents to perform this assay are available (e.g. Applied Biosystems Inc. MeltDoctor kit). These kits include a suitable buffer solution containing one of the double strand DNA (dsDNA) binding dyes (such as ABI HRM dyes, SYBR Green, SYTO, etc.). The properties of the dsDNA dyes are such that they emit almost no fluorescence in free form, but are highly fluorescent when bound to dsDNA.

[00235] To perform the assay the cDNA or a corresponding oligonucleotide are mixed with Molecule in concentrations defined by the particular manufacturer's protocols. The mixture is heated to 95 °C to dissociate all pre-formed dsDNA complexes, then slowly cooled to room temperature or other lower temperature defined by the kit manufacturer to allow the DNA molecules to anneal. The newly formed complexes are then slowly heated to 95 °C with simultaneous continuous collection of data on the amount of fluorescence that is produced by the reaction. The fluorescence intensity is inversely proportional to the amounts of dsDNA present in the reaction. The data can be collected using a real time PCR instrument compatible with the kit (e.g. ABI's StepOne Plus Real Time PCR System or LightTyper instrument, Roche Diagnostics, Lewes, UK).

[00236] Melting peaks are constructed by plotting the negative derivative of fluorescence with respect to temperature ( $-d(\text{Fluorescence})/dT$ ) on the y-axis) against temperature (x-axis) using appropriate software (for example LightTyper (Roche) or SDS Dissociation Curve, ABI). The data is analyzed to identify the temperature of the rapid transition from dsDNA complex to single strand molecules. This temperature is called  $T_m$  and is directly proportional to the strength of interaction between the two molecules. Typically,  $T_m$  will exceed 40 °C.

*Example 2: Modulation of NRF2 polynucleotides*

*Treatment of HepG2 cells with antisense oligonucleotides*

[00237] HepG2 cells from ATCC (cat# HB-8065) were grown in growth media (MEM/EBSS (Hyclone cat #SH30024, or Mediatech cat # MT-10-010-CV) +10% FBS (Mediatech cat# MT35- 011-CV)+ penicillin/streptomycin (Mediatech cat# MT30-002-CI)) at 37°C and 5% CO<sub>2</sub>. One day before the experiment the cells were replated at the density of  $1.5 \times 10^5/\text{ml}$  into 6 well plates and incubated at 37°C and 5% CO<sub>2</sub>. On the day of the experiment the media in the 6 well plates was changed to fresh growth media. All antisense oligonucleotides were diluted to the concentration of 20 µM. Two µl of this solution was incubated with 400 µl of

Opti-MEM media (Gibco cat#31985-070) and 4 µl of Lipofectamine 2000 (Invitrogen cat# 11668019) at room temperature for 20 min and applied to each well of the 6 well plates with HepG2 cells. A Similar mixture including 2 µl of water instead of the oligonucleotide solution was used for the mock-transfected controls. After 3-18 h of incubation at 37°C and 5% CO<sub>2</sub> the media was changed to fresh growth media. 48 h after addition of antisense oligonucleotides the media was removed and RNA was extracted from the cells using SV Total RNA Isolation System from Promega (cat # Z3105) or RNeasy Total RNA Isolation kit from Qiagen (cat# 74181) following the manufacturers' instructions. 600 ng of RNA was added to the reverse transcription reaction performed using Verso cDNA kit from Thermo Scientific (cat#AB1453B) or High Capacity cDNA Reverse Transcription Kit (cat# 4368813) as described in the manufacturer's protocol. The cDNA from this reverse transcription reaction was used to monitor gene expression by real time PCR using ABI Taqman Gene Expression Mix (cat#4369510) and primers/probes designed by ABI (Applied Biosystems Taqman Gene Expression Assay: Hs00232352\_m1 by Applied Biosystems Inc., Foster City CA). The following PCR cycle was used: 50°C for 2 min, 95°C for 10 min, 40 cycles of (95°C for 15 seconds, 60°C for 1 min) using StepOne Plus Real Time PCR Machine (Applied Biosystems).

[00238] Fold change in gene expression after treatment with antisense oligonucleotides was calculated based on the difference in 18S-normalized dCt values between treated and mock-transfected samples.

#### *Results:*

[00239] Real time PCR results show that the levels of NRF2 mRNA in HepG2 cells are significantly increased 48 h after treatment with two of the oligos designed to NRF2 antisense BU656954.1 (Fig 1A).

#### *Treatment of CHP212 cells with antisense oligonucleotides*

[00240] CHP212 cells from ATCC (cat# CRL-2273) were grown in growth media (MEM/EBSS (Hyclone cat #SH30024, or Mediatech cat # MT-10-010-CV) +10% FBS (Mediatech cat# MT35- 011-CV)+ penicillin/streptomycin (Mediatech cat# MT30-002-CI)) at 37°C and 5% CO<sub>2</sub>. One day before the experiment the cells were replated at the density of  $1.5 \times 10^5$ /ml into 6 well plates and incubated at 37°C and 5% CO<sub>2</sub>. On the day of the experiment the media in the 6 well plates was changed to fresh growth media. All antisense oligonucleotides were diluted to the concentration of 20 µM. Two µl of this solution was

incubated with 400 µl of Opti-MEM media (Gibco cat#31985-070) and 4 µl of Lipofectamine 2000 (Invitrogen cat# 11668019) at room temperature for 20 min and applied to each well of the 6 well plates with CHP212 cells. A Similar mixture including 2 µl of water instead of the oligonucleotide solution was used for the mock-transfected controls. After 3-18 h of incubation at 37°C and 5% CO<sub>2</sub> the media was changed to fresh growth media. 48 h after addition of antisense oligonucleotides the media was removed and RNA was extracted from the cells using SV Total RNA Isolation System from Promega (cat # Z3105) or RNeasy Total RNA Isolation kit from Qiagen (cat# 74181) following the manufacturers' instructions. 600 ng of RNA was added to the reverse transcription reaction performed using Verso cDNA kit from Thermo Scientific (cat#AB1453B) or High Capacity cDNA Reverse Transcription Kit (cat# 4368813) as described in the manufacturer's protocol. The cDNA from this reverse transcription reaction was used to monitor gene expression by real time PCR using ABI Taqman Gene Expression Mix (cat#4369510) and primers/probes designed by ABI (Applied Biosystems Taqman Gene Expression Assay: Hs00232352\_m1 by Applied Biosystems Inc., Foster City CA). The following PCR cycle was used: 50°C for 2 min, 95°C for 10 min, 40 cycles of (95°C for 15 seconds, 60°C for 1 min) using StepOne Plus Real Time PCR Machine (Applied Biosystems).

[00241] Fold change in gene expression after treatment with antisense oligonucleotides was calculated based on the difference in 18S-normalized dCt values between treated and mock-transfected samples.

#### *Results:*

[00242] Real time PCR results show that the levels of NRF2 mRNA in CHP212 cells are significantly increased 48 h after treatment with two of the oligos designed to NRF2 antisense BU656954.1 (Fig 1B).

[00243] Although the invention has been illustrated and described with respect to one or more implementations, equivalent alterations and modifications will occur to others skilled in the art upon the reading and understanding of this specification and the annexed drawings. In addition, while a particular feature of the invention may have been disclosed with respect to only one of several implementations, such feature may be combined with one or more other features of the other implementations as may be desired and advantageous for any given or particular application.

[00244] The Abstract of the disclosure will allow the reader to quickly ascertain the nature of the technical disclosure. It is submitted with the understanding that it will not be used to interpret or limit the scope or meaning of the following claims.

## CLAIMS

What is claimed is:

1. A method of modulating a function of and/or the expression of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising:

contacting said cells or tissues with at least one antisense oligonucleotide 5 to 30 nucleotides in length wherein said at least one oligonucleotide has at least 50% sequence identity to a reverse complement of a polynucleotide comprising 5 to 30 consecutive nucleotides within nucleotides 1 to 5808 of SEQ ID NO: 3 (Figure 3); thereby modulating a function of and/or the expression of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

2. A method of modulating a function of and/or the expression of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising:

contacting said cells or tissues with at least one antisense oligonucleotide 5 to 30 nucleotides in length wherein said at least one oligonucleotide has at least 50% sequence identity to a reverse complement of a natural antisense of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide; thereby modulating a function of and/or the expression of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

3. A method of modulating a function of and/or the expression of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising:

contacting said cells or tissues with at least one antisense oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to an antisense oligonucleotide to the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide; thereby modulating a function of and/or the expression of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

4. A method of modulating a function of and/or the expression of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising:

contacting said cells or tissues with at least one antisense oligonucleotide that targets a region of a natural antisense oligonucleotide of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide; thereby modulating a function of and/or the expression of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

5. The method of claim 4, wherein a function of and/or the expression of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) is increased *in vivo* or *in vitro* with respect to a control.

6. The method of claim 4, wherein the at least one antisense oligonucleotide targets a natural antisense sequence of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide.

7. The method of claim 4, wherein the at least one antisense oligonucleotide targets a nucleic acid sequence comprising coding and/or non-coding nucleic acid sequences of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide.

8. The method of claim 4, wherein the at least one antisense oligonucleotide targets overlapping and/or non-overlapping sequences of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide.

9. The method of claim 4, wherein the at least one antisense oligonucleotide comprises one or more modifications selected from: at least one modified sugar moiety, at least one modified internucleoside linkage, at least one modified nucleotide, and combinations thereof.

10. The method of claim 9, wherein the one or more modifications comprise at least one modified sugar moiety selected from: a 2'-O-methoxyethyl modified sugar moiety, a 2'-methoxy modified sugar moiety, a 2'-O-alkyl modified sugar moiety, a bicyclic sugar moiety, and combinations thereof.

11. The method of claim 9, wherein the one or more modifications comprise at least one modified internucleoside linkage selected from: a phosphorothioate, 2'-O-methoxyethyl (MOE), 2'-fluoro, alkylphosphonate, phosphorodithioate,

alkylphosphonothioate, phosphoramidate, carbamate, carbonate, phosphate triester, acetamidate, carboxymethyl ester, and combinations thereof.

12. The method of claim 9, wherein the one or more modifications comprise at least one modified nucleotide selected from: a peptide nucleic acid (PNA), a locked nucleic acid (LNA), an arabino-nucleic acid (FANA), an analogue, a derivative, and combinations thereof.

13. The method of claim 1, wherein the at least one oligonucleotide comprises at least one oligonucleotide sequences set forth as SEQ ID NOS: 4 to 6.

14. A method of modulating a function of and/or the expression of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) gene in mammalian cells or tissues *in vivo* or *in vitro* comprising:

contacting said cells or tissues with at least one short interfering RNA (siRNA) oligonucleotide 5 to 30 nucleotides in length, said at least one siRNA oligonucleotide being specific for an antisense polynucleotide of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide, wherein said at least one siRNA oligonucleotide has at least 50% sequence identity to a complementary sequence of at least about five consecutive nucleic acids of the antisense and/or sense nucleic acid molecule of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide; and, modulating a function of and/or the expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in mammalian cells or tissues *in vivo* or *in vitro*.

15. The method of claim 14, wherein said oligonucleotide has at least 80% sequence identity to a sequence of at least about five consecutive nucleic acids that is complementary to the antisense and/or sense nucleic acid molecule of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide.

16. A method of modulating a function of and/or the expression of Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in mammalian cells or tissues *in vivo* or *in vitro* comprising:

contacting said cells or tissues with at least one antisense oligonucleotide of about 5 to 30 nucleotides in length specific for noncoding and/or coding sequences of a sense and/or natural antisense strand of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide wherein said at least one antisense oligonucleotide has at least 50% sequence identity to at least one nucleic acid sequence set forth as SEQ ID NOS: 1, 3;



and, modulating the function and/or expression of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) in mammalian cells or tissues *in vivo* or *in vitro*.

17. A synthetic, modified oligonucleotide comprising at least one modification wherein the at least one modification is selected from: at least one modified sugar moiety; at least one modified internucleotide linkage; at least one modified nucleotide, and combinations thereof; wherein said oligonucleotide is an antisense compound which hybridizes to and modulates the function and/or expression of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) gene *in vivo* or *in vitro* as compared to a normal control.

18. The oligonucleotide of claim 17, wherein the at least one modification comprises an internucleotide linkage selected from the group consisting of: phosphorothioate, alkylphosphonate, phosphorodithioate, alkylphosphonothioate, phosphoramidate, carbamate, carbonate, phosphate triester, acetamidate, carboxymethyl ester, and combinations thereof.

19. The oligonucleotide of claim 17, wherein said oligonucleotide comprises at least one phosphorothioate internucleotide linkage.

20. The oligonucleotide of claim 17, wherein said oligonucleotide comprises a backbone of phosphorothioate internucleotide linkages.

21. The oligonucleotide of claim 17, wherein the oligonucleotide comprises at least one modified nucleotide, said modified nucleotide selected from: a peptide nucleic acid, a locked nucleic acid (LNA), analogue, derivative, and a combination thereof.

22. The oligonucleotide of claim 17, wherein the oligonucleotide comprises a plurality of modifications, wherein said modifications comprise modified nucleotides selected from: phosphorothioate, alkylphosphonate, phosphorodithioate, alkylphosphonothioate, phosphoramidate, carbamate, carbonate, phosphate triester, acetamidate, carboxymethyl ester, and a combination thereof.

23. The oligonucleotide of claim 17, wherein the oligonucleotide comprises a plurality of modifications, wherein said modifications comprise modified nucleotides selected from: peptide nucleic acids, locked nucleic acids (LNA), analogues, derivatives, and a combination thereof.

24. The oligonucleotide of claim 17, wherein the oligonucleotide comprises at least one modified sugar moiety selected from: a 2'-O-methoxyethyl modified sugar moiety, a 2'-methoxy modified sugar moiety, a 2'-O-alkyl modified sugar moiety, a bicyclic sugar moiety, and a combination thereof.
- 5 25. The oligonucleotide of claim 17, wherein the oligonucleotide comprises a plurality of modifications, wherein said modifications comprise modified sugar moieties selected from: a 2'-O-methoxyethyl modified sugar moiety, a 2'-methoxy modified sugar moiety, a 2'-O-alkyl modified sugar moiety, a bicyclic sugar moiety, and a combination thereof.
- 10 26. The oligonucleotide of claim 17, wherein the oligonucleotide is of at least about 5 to 30 nucleotides in length and hybridizes to an antisense and/or sense strand of a Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide wherein said oligonucleotide has at least about 20% sequence identity to a complementary sequence of at least about five consecutive nucleic acids of the antisense and/or sense coding and/or noncoding nucleic acid sequences of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide.
- 15 27. The oligonucleotide of claim 17, wherein the oligonucleotide has at least about 80% sequence identity to a complementary sequence of at least about five consecutive nucleic acids of the antisense and/or sense coding and/or noncoding nucleic acid sequence of the Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide.
- 20 28. The oligonucleotide of claim 17, wherein said oligonucleotide hybridizes to and modulates expression and/or function of at least one Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide *in vivo* or *in vitro*, as compared to a normal control.
- 25 29. The oligonucleotide of claim 17, wherein the oligonucleotide comprises the sequences set forth as SEQ ID NOS: 4 to 6.
- 30 30. A composition comprising one or more oligonucleotides specific for one or more Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotides, said polynucleotides comprising antisense sequences, complementary sequences, alleles, homologs, isoforms, variants, derivatives, mutants, fragments, or combinations thereof.

31. The composition of claim 30, wherein the oligonucleotides have at least about 40% sequence identity as compared to any one of the nucleotide sequences set forth as SEQ ID NOS: 4 to 6.

32. The composition of claim 30, wherein the oligonucleotides comprise nucleotide sequences set forth as SEQ ID NOS: 4 to 6.

33. The composition of claim 32, wherein the oligonucleotides set forth as SEQ ID NOS: 4 to 6 comprise one or more modifications or substitutions.

34. The composition of claim 33, wherein the one or more modifications are selected from: phosphorothioate, methylphosphonate, peptide nucleic acid, locked nucleic acid (LNA) molecules, and combinations thereof.

35. A method of preventing or treating a disease associated with at least one Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide and/or at least one encoded product thereof, comprising:

administering to a patient a therapeutically effective dose of at least one antisense oligonucleotide that binds to a natural antisense sequence of said at least one Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide and modulates expression of said at least one Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide; thereby preventing or treating the disease associated with the at least one Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide and/or at least one encoded product thereof.

36. The method of claim 35, wherein a disease associated with the at least one Nuclear factor (erythroid-derived 2)-like 2 (NRF2) polynucleotide is selected from: chronic obstructive pulmonary disease (COPD), multiple sclerosis, a hepatic disease or disorder, a gastrointestinal disease or disorder, diabetes, autoimmunity, an immune related disease or disorder, an immunodeficiency (e.g., AIDS), a neurological disease or disorder, a neurodegenerative disease or disorder, a disease or disorder or condition associated with oxidative stress, an eye disease (e.g., age-related macular degeneration, cataracts, light retinopathy, retinopathy of prematurity etc.), a skin disease, asthma, arteriosclerosis, a chronic inflammatory diseases or condition (e.g., vasculitis, pulmonary bronchitis, rheumatoid arthritis, osteoarthritis, hepatitis, pancreatitis, dermatitis, esophagitis, ulcerative colitis, Crohn's disease, conjunctivitis etc.), nerve repair and paralysis, neuroendocrine differentiation, an inflammatory

disease, a muscular disease or disorder, diseases or disorders associated with infectious organisms, senile plaques, cerebral amyloid angiopathy, atherosclerosis, glioblastoma, amyloid deposition, neurofibrillary tangles, dementia, choriocarcinoma, astrocytoma, amyloidosis, hyperlipidemia, neoplastic transformation, an atherosclerotic plaque, an atherosclerotic obstruction, metastasis, myocardial infarction, pulmonary fibrosis, necrosis, shock, melanoma, colorectal carcinoma, genetic susceptibility, psoriasis, a disease or disorder associated with abnormal cell proliferation ((e.g., cancer, psoriasis etc.), cancer (e.g., prostate cancer, lung cancer, breast cancer, Non-small cell lung carcinomas (NSCLCs), leukemia etc.), inflammation, glioma, carcinoma, neuropathology, tumors, vascular diseases, cell damage, brain tumors, hypercholesterolemia, liposarcoma, coronary heart disease, coronary artery disease, glomerulonephritis, venous thrombosis and a pathological process.

37. A method of identifying and selecting at least one oligonucleotide for in vivo administration comprising: selecting a target polynucleotide associated with a disease state; identifying at least one oligonucleotide comprising at least five consecutive nucleotides which are complementary to the selected target polynucleotide or to a polynucleotide that is antisense to the selected target polynucleotide; measuring the thermal melting point of a hybrid of an antisense oligonucleotide and the target polynucleotide or the polynucleotide that is antisense to the selected target polynucleotide under stringent hybridization conditions; and selecting at least one oligonucleotide for in vivo administration based on the information obtained.

FIGURE 1A





FIGURE 1B



## FIGURE 2

(SEQ ID NO: 1)

>gi| 166295208|ref| NM\_006164| Homo sapiens nuclear factor (erythroid-derived 2)-like 2 (NFE2L2), transcript variant 1, mRNA.

AAATCAGGGAGGCGCAGCTCCTACACCAACGCCTTTCCGGGGCTCCGGGTGTGTTTGTTCCTCAACTGTTTAAACTG  
TTTCAAAGCGTCCGAACCTCCAGCGACCTTCGCAAACAACTCTTTATCTCGCGGGCGAGAGCGCTGCCCTTATTTG  
CGGGGGAGGGCAAACCTGAACGCCGGCACCGGGGAGCTAACGGAGACCTCCTCTAGGTCCCCCGCTGCTGGGACC  
CCAGCTGGCAGTCCCTTTCCCGCCCCCGGACCGCGAGCTTCTTGCCTCAGCCCCGGCGCGGTGGGGGATTTTCGG  
AAGCTCAGCCCCGCGCGGCCGGCGGGGAAGGAAGGGCCCCGACTCTTGCCCCGCCCTTGTTGGGGCGGGAGGCGGA  
GCGGGGACAGGGGCCCGCGGCGTGTAGCCGATTACCGAGTGCCGGGGAGCCCCGAGGAGCCGCCGACGCAGCCGC  
CACCGCCGCCGCCGCCGCCAGAGCCGCCCTGTCCGCGCCGCGCTCGGCAGCCGGAACAGGGCCGCCGTCGG  
GGAGCCCCAACACACGGTCCACAGCTCATCATGATGGACTTGGAGCTGCCGCCCGGGACTCCCGTCCCAGCAG  
GACATGGATTTGATTGACATACTTTGGAGGCAAGATATAGATCTTGGAGTAAGTCGAGAAGTATTTGACTTCAGT  
CAGCGACGGAAAGAGTATGAGCTGGAAAAACAGAAAAAACTTGAAAAGGAAAGACAAGAACAACCTCCAAAAGGAG  
CAAGAGAAAGCCTTTTTTCGCTCAGTTACAACCTAGATGAAGAGACAGGTGAATTTCTCCCAATTCAGCCAGCCCAG  
CACATCCAGTCAGAAACCAGTGGATCTGCCAACTACTCCAGGTTGCCACATTCCCAAATCAGATGCTTTGTAC  
TTTGATGACTGCATGCAGCTTTTGGCGCAGACATTCCCGTTTGTAGATGACAATGAGGTTTCTTCGGCTACGTTT  
CAGTCACTTGTTTCTGATATTCCCGGTACATCGAGAGCCCAGTCTTCATTGCTACTAATCAGGCTCAGTCACCT  
GAACTTCTGTTGCTCAGGTAGCCCCCTGTTGATTTAGACGGTATGCAACAGGACATTGAGCAAGTTTGGGAGGAG  
CTATTATCCATTCTGAGTTACAGTGTCTTAATATTGAAAATGACAAGCTGGTTGAGACTACCATGGTTCCAAGT  
CCAGAAGCCAAACTGACAGAAGTTGACAATTATCATTTTTACTCATCTATACCCTCAATGGAAAAAGAGTAGGT  
AACTGTAGTCCACATTTTCTTAATGCTTTTGGAGATTCTTTCAGCAGCATCCTCTCCACAGAAGACCCCAACCAG  
TTGACAGTGAACCTATTAAATTCAGATGCCACAGTCAACACAGATTTTGGTGATGAATTTTATTCTGCTTTTATA  
GCTGAGCCCAGTATCAGCAACAGCATGCCCTCACCTGCTACTTTAAGCCATTCATCTCTGAACCTCTAAATGGG  
CCCATTGATGTTTCTGATCTATCACTTTGCAAAGCTTTCAACCAAAACCACCTGAAAGCACAGCAGAATTCAAT  
GATTCTGACTCCGGCATTTCCTAAACACAAAGTCCAGTGTGGCATCACCAGAACCTCAGTGGAAATCTTCCAGC  
TATGGAGACACACTACTTGGCCTCAGTGATTCTGAAGTGGAAGAGCTAGATAGTGCCCCCTGGAAAGTGTCAAACAG  
AATGGTCTTAAACACCAGTACATTCTTCTGGGGATATGGTACAACCCCTTGTCACCATCTCAGGGGCAGAGCACT  
CACGTGCATGATGCCCAATGTGAGAACACACCAGAGAAAGAATTGCCCTGTAAGTCTGGTCAATCGGAAAAACCCCA  
TTCACAAAAGACAAACATTTCAAGCCGCTTGGAGGCTCATCTCACAAGAGATGAACTTAGGGCAAAAGCTCTCCAT  
ATCCCATTCCCTGTAGAAAAAATCATTAACCTCCCTGTTGTTGACTTCAACGAAATGATGTCCAAAGAGCAGTTC  
AATGAAGCTCAACTTGCATTAATTCGGGATATACGTAGGAGGGGTAAGAATAAAGTGGCTGCTCAGAATTGCAGA  
AAAAGAAAACCTGGAAAATATAGTAGAACTAGAGCAAGATTTAGATCATTGAAAGATGAAAAAGAAAAATTGCTC  
AAAGAAAAGGAGAAAATGACAAAAGCCTTCACCTACTGAAAAACAACCTCAGCACCTTATATCTCGAAGTTTTTC  
AGCATGCTACGTGATGAAGATGGAAAACCTTATTCTCCTAGTGAATACTCCCTGCAGCAAACAAGAGATGGCAAT  
GTTTTCTTGTTCCTCCAAAAGTAAGAAGCCAGATGTTAAGAAAAACTAGATTTAGGAGGATTTGACCTTTTCTGAG  
CTAGTTTTTTTTGTACTATTATACTAAAAGCTCCTACTGTGATGTGAAATGCTCATACTTTATAAGTAATTTCTATG  
CAAAATCATAGCCAAAACCTAGTATAGAAAATAATACGAACTTTAAAAAGCATTGGAGTGTCAGTATGTTGAATC  
AGTAGTTTCACTTTAACTGTAAACAATTTCTTAGGACACCATTGTTGGGCTAGTTTCTGTGTAAGTGTAATACTAC  
AAAACTTATTTATACTGTTCTTATGTCATTTGTTATATTATAGATTTATATGATGATATGACATCTGGCTAAA  
AAGAAATTATTGCAAACTAACCCTATGTACTTTTTTATAAATACTGTATGGACAAAAAATGGCATTTTTTTATA  
TTAAATTGTTTAGCTCTGGCAAAAAAAAAAATTTAAGAGCTGGTACTAATAAAGGATTATTATGACTGTTAAA  
TTAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

(SEQ ID NO: 2)

>hg18\_knownGene\_uc010fra.1 range=chr2:177805251-177836511 5'pad=0 3'pad=0  
strand=- repeatMasking=none

AAGGGACAGGTTGGAGCTGTTGATCTGTTGCGCAATTGCTATTTTCCCCAGAGCGGCTTTGTCTTTGGATTTAGC  
GTTTCAGAATTGCAATTCCAAAATGTGTAAGACGGGATATTCTCTTCTGTGCTGTCAAGGgtaagagttgagag  
gtagattagaatttctgttgccttttagtctgttagtaatttttgccttcagctattatttctccctgagtact  
ttatatatgtttccttttcagttgagaatttgcctcaatttcttaacatgttccccctcttctgcaggggagag  
agtggaacgcttgcgtttcaaacacttgctaatttctgtgaattgttataaaagtgaagaaagtgttctgctca  
tcctttgtagaaactttaaaagtagatatttatatttctaacttcttttgtaaagtgaatttttaggaaaaaattg  
gaattcaaggaaatgtgtacttgatgtacagtaaatacgtttatgtctgttaagtgaagttttcgtttaattcc

aaagatctattgttaaagttttaagttatggacaatgtatttaattgtgcttttttccccctttaaagttgtttgtc  
ttgaacttttccccagtccttcatttgggattgtattcttctggttccaactagtgaagaattaaattgtaccttcc  
gatttatttaatatagctgctgttcaagagttatctataggaatgcttgtttgactgaaggatataatgataaaaatg  
aacttttagtttttctgatttccggaattgtctacagccctctattatttttgaatttttctttctttgtctgtgta  
atataacatttccttatacaaaaatgcgtgaaacatatgtaccgcattaagtgtgtattttccaattacgtttgat  
aaaaataactgcttttctacctttgtaatcagacatcgatttgcataatttgaaaacagaaaaagaacaagaaaat  
ttagaattgattttgtttatgattcatatttagattgttgtcatccataaagatttgaacagagtcaaacatttttc  
ttgaccttttgtcaaaaaaaactctccaggtgtgaagtggtagtaggatatagtttttcaccttctgtgcagagg  
atgtgttgataaacagggcagtaaggagagcccttagattcagtatctccattaggtggaaggaacctaccattac  
tcttttcaagggtgtcttggagagataaagcactttcctagatgagggtgaagttgatattgaaatagaccgat  
gagattatttttaagtcaaacaaaagaaaaataggacaggcctctcatctcctgaatgaatgttaagcaagaac  
aacttgacttttttagatatgtgaaaaatctcttgagaggttgtggaagcagcatgtaggaggataaatgcataac  
tcacatcagtccttttacgcattaaaaataacttgggccattttgaaatctttctttcttgccttgggttcttata  
agcaatgttggggggaaaaactgactctgtctttaggattaccaggaaaagttgaaagccaaatgcatcatttttc  
ttaatatattgtttttattgaggataaatcatgtttaacaaagctgcatgaaaaaaaggaacccaacaagtacatt  
tttttttctgggaatttctctgtcctgactgaagacatttaaaggggttttgtgtcaatatttcctccttaac  
tgttcctcaagcctgggttaacttttcagtatgcaggaagaatttggcactaagataataatagataagttactg  
ggcacgtccctgggtgccagacataattgcttcacatgcataattttacagtaacctcacgggaaatattattat  
ccctgggtttccaggttgggaaaaccaggcataatagtttagatcaggcattaggttaggaaagggttaggagagagg  
catagagaaaggccaggacctgctaagggtgaccatactattgagtggccaaggtagcattcaaacaggtttgact  
cctgggggcttctaataccacttccagactgtacaaaactccttaaaactctaaataggaagaaaagctcacac  
tttaaaaagtgggttaagttgagatttgcctattctgggggaaggagagggccggaatataaaaagattaatgtctgc  
agcttttccatttagcaagacatttaatactgtatgtgacgtggggaggtggacctgttaaatacaggggagag  
tgaatacaatccatcaatgggtaccactgtgggtccttttaaggaagacacagaggcctcggtggattcaatttag  
aaaggcaggactgttttttgttttttgttttttttttttttgagatgaagtcttgttctgttggccaggctgga  
atgcagtggcctgatcttggctcactgcagcctctacttcccgggttcaagtgattctcctgcttcagcctcccg  
agtagctgggattataggcaccactgcccggctaatttttgtatttttagtagagatgggggtttaccatgttg  
gccaggctgggctgggtctcgaactcctgacctcaggtgatccaccgcctcggcctcccaaagtgttgggattac  
aggtgtgagccactgcgccagtcctgtttcttaataataataatagtttttaaaaatttctgatgtttggcaatgtcc  
aagattctgcctcacatttttccagaaactgccttttttcttaattaaatctcttagctactggagtcctttgcctt  
tgtttgccttgctacttgggtacatttcttctgttccattggctccttttgttttgaagtgtgctcaaaaaataag  
tttggtagtaagaatatggtgaagagtggtgacagacaggggttaaacaggctgaaggaaagagagcagaatgggg  
aaaatttgagttgaaaacagctagcagaaagctgatggcgcaaattcaaccctaatactcatgttttggcacttct  
gaactatagcaagcataagggctcaactcttatctagttgaagcttacttgggtctttgagaatagtacatctca  
aaattaaacacttttcttagtccttattgttaagttgtttttggaattattgtcttatttgtctgaattaaagtgtg  
ctgcaggacttataggatctgatggatgatttgcctaaaagttgggtccagggtcaacctcagggtcatttaaatgct  
tgggtccagccactttctgaaagttgactgaaatatgttgcacagcagggttagaggaggacaagtttactagatgt  
catttttaacctgcttgtcttcatctctgactgtagaaatgtatattcattataagtttggccaaaatagagaat  
agagtttattctttgatcatctaatcttaggagctattgaaatttactgctacctgcttttaaaataagcgcca  
acaaacattttattaaaagggctttaagggtataggccaagatttattcaattgctaattgggtatgttttctgtaa  
gaaagcaaaaatatgttcaagtgtataatattaaatattaacattcagaaacttggagaaatacagatggcctttt  
ttttttgagacgcagtccttgcctctgtcgcgccagggtggagtgacgtggcatgatctccgctcactgcaagctcca  
cctcccgggttcacgccatttctcctgccccagcctcccagtagctgggtactacaggccaccgcccaccagcctg  
gctaattttttgtatttttagtagagatgggggttccactgtgttagccaggatgatctcgatctcctgacctcat  
gatccgctgctcagcctcccaaagtgtgggattacaagcttaagccaccgcgctggcctagatggatttct  
tgtaccagactataccctagacatgaaaacaagactgttaaacaaaatcctgtgctacctaaagtgttgagccttc  
tgtgtgtcagtttctgccataataaataactcttaactatgattattaagtataattaagaactttgagtagggga  
ggcacgtttaatcattcgtttttacggcatatgtactataataacctgtgtgttgcaaatcctgggtattttaaagt  
ctttttgtgtattttgagcatgtaagtttatttgggtccaaactgctgggtactagtaaagacaagtccagggcataa  
gtaggacttacagcaccaggttcgtcttttttttttgagacagagtcctcagtcggtcaccacagggtggagtgcat  
tggtagagctgtgggtcactgcacctccacctcccaggttcaagcgatttctcctgcctcagcctcccaagtagc  
tgggattacaagtgtgtgccgccacgcccgttaatttttgtatttttagtagagatggagtttactatgtttgg  
ccaggctgatcccgcctcaggtgatgtgccaccttgggtctccgaaagtgtgggattataggcggtgagccact  
gcgcctgacctccaagtcctcttcaaaggctctgttgataattgactcttgagagtcattacagtcctcaac



agcctgtattctgaatatattgtttttaattctaagacaatgctaaatagccattccaaataaggtgagaactgag  
acttttagggcctgtattatcagctgtacactgacaagctctgaactctattgctttttcaaatacagagaagaaaa  
atgagtaagaaaagaaaaatatggatcacttaggctttgatgcctcctctgagtttatagtgtttaccttctgta  
ttacttaggtgatcctctctaattggcatttttaaattctctgttgatacttgtcttacttatgtatctgttttttt  
tcctctatagtagatagaatttatagatatatatgtatttttaaagtaagatacaatttacgtaccttacaattta  
cccatttaattctaattcaggggttatgcaaccctatgaatttagtagtttattcatagagttatgcaaccatcact  
gtaatcaattttcaacaccctcaaaagaaaccccatatccattagcagtcacttcccattttcccctcaactctcc  
tggcccttggttaactactcagttattttctgtacctatggatttgccattctggacatatcatgtaaatggaat  
agcatgtaaatggaatcatacagttatttatccttctgttgctggcttctttcacagcaatatgttttcaagggtt  
aaaaatgttgcaacgtatatcaatagcgtcattcctttttatggctgcatactattccattgtatggctatacca  
cattttattttatccagttgatgatcatttgagttgtttccactttttatgtttctgtttgtttcttacagcagtg  
gttctcaaactgttgtatattagactcaccatcagttatttaaaaaactaatgcccaggccatatccctcatgaaa  
tcaaaatctctgggggtggggccaggcatcactattttttaagattctgaggggattctaacgagcagtcagg  
tttgagagccagtgccctagggcagcagtcacacaaccttttagcaccagggaactagttttgtggaagacagttt  
ttccatagaaggaggtagaggatggtttcaggatgaaactgttccacctcagatcatcaggcattagattgtcct  
aaggagctggcaacctacatcccttgcatactcagtttacaacagggttccggcttctctgataaaactaatgctg  
ctgctaattctgacaggaggtggagctcaggcggaatgctcgcctactgctcacctcctgctgtgcgccctggtt  
cctaattggggcactgaccctccgcagccagcaggtggggaccctgcccctaggggaccacccctcagtgcttgg  
atccattgaagtatatagtgttttagaaaatctgggttaggatgtcttatttaaatcatgaaagcaagtgttttg  
tctttgatcattgcccttcacagccttacagttaacacctacattcaggaactgtgtttcaaagtgcctggcgag  
tgtttgaaactcactaaatatataaacctgggctcattaaacccctccagaagatttgggttaactttgttatga  
aagggttccctgaagtgggtcagcaggattgctctgtgttttttcaggaatgtggcctgtgataacttgcaatc  
tagattatttggaaatagcactgagagaagccagtgaggacagaatgctcagaagcttgggataggtgttgaaca  
tcctggaggccaggacggaacactgtcttactctaggaagctgtgttccctgggcctcattatcttccctccgttaa  
aaacaaaagacttaaatctccacagcagctttcagcaacttcatttttgggttctctgtatctgcctgataaagt  
cccactttgtagtggctcccacttataatttacctgaatggcttttgggttgacataatttggaaactggggctaac  
ttccaaactgttggcaacttgtgtgtgggtgtgctgcccacagcacagcagtcaccttgagagacttgatgggtg  
tggtgggtgggttggggggttctgaagctggcttagccccagccctatacacccacccacagatgggtgggaacaa  
gcccagaagagagtgggttaactctgtccactgtggcctccacagccaaggttgccaggcagagctcgcaaggccc  
aattccagctcttgtctttgaccgttgcccctttgtgttgggggggtgtatttagtcacctttctggaagcatgct  
tttctaattctagtcatcagtagtttgttgccttttaagattttgaaaatggatcctgttattttacttaggagtt  
tcgtattgaatgggtgtacataatgtgattcaagtacctcaaaacagaaggacttcagtttaagatttaggctctat  
gcaacatacacttcttgcattttctcattcaatgtcctttccttttttttttttttttttttttgcagaatgtag  
ctgacattcagagtagatttagtaccttcaatgtctgtgtgaaagaaatgaccttaatatgaggacaatattgact  
gtgtatttagggggccactgttaaggcatatagaattttgctttatttcagacctgacaatctcttgtctgctc  
tgcttccgcataaaattataactgcacatggatgtaaaacccaacctattccctgcctgagggactagaatag  
aggaagaatgactatagttctttgttgccttttgtgaaggtaacaggcacagaggtatgatgcatgatggaatt  
atatacctcttcttgagggtgttggaggctgactaaggacctgtacttttttttgggttgttttgactggtg  
ggccaggagacctgtgttgtgtctagagagtggttgaagaacctgaatatctcgaaaaaagaaaaataattttt  
ttaaccattaaaaattcctggtagtgacttccctctggcaagtaaaaaactctcattttccttaaaaaatgagagtt  
ttttacttgcaataggaaaacttgccaatttttcaagttatttttattcttgatgattctcaatgagacataatt  
aaaatacacatagaaaaaacaaccgtaggccaggtgcagtggtcacaccagtaatcccaacactttgggaagtc  
aagggtgggaggatcactagagaccaggagtttgagatgagcctgggcaacataattgagacactgtctcctcaaaa  
aattaaaaaaattagctggatccagtggtgcacacctgcaattgtagctatttgggagggttaagggtggaagaata  
gcttgagcccaggagtttgaagggtgcagtgagccataatcatgccactgtactccagcccaggcaacagagtaag  
atcatgtctcaatgaaaaaaagaaaaatcaaccctagtggactggaacagggtggtttactttgtgtcagctgc  
agtccccactcgccaaccacatgttcttctcctgctgggtcccaagcagagaccagaaccaagaatgagaat  
catcctgtgggtgcagggtgtatcttctcctccagagacagcacttcacctctgttagaaacttctttaccaca  
ctgctgtttgggtgaaatccttagggcagtgacttccaaactgtgatccaggggccacccttgggtgggaggatca  
ggaaggagggggaatgtatcttaagaaaagttgggggaagctcatatctcttctcgttgatcctgagaagtaaaag  
tctttcctcctcagagagatgcgggtagacatgactcgcctagatagaagctcattcatctccctcccttctgcc  
tctgcaggactccttggaaagtctgggtccccgggagtatgcggctcttgcctgtgtgtacaggtttccaaagttca  
cttgataacagtacaattgtgctgtaaattgtgcagtaattggagactgaggaaacaattagtctttcatcttcat  
tcatgttaccagctcacttccatgggtgtgaaaagagccagactttggaggttaggagacatctgaatacagacag

aactggcaagattatggggaagaaaagggaaaagtaataatattttcaaagagggttaagccatttggccaggac  
aacactgcttgccaggttagcagaactgggattttgactgtccataatccctattctctgcttgttttactccttcc  
ctctcttctttcttttttcttattgtctaatgacctacagacattctgtcagctgctaaagaatgagggactccttt  
ataaagattgataggacagtgtcgctacttaaggagttccttccatgtggaccagtggggcgctgttgacgcat  
ggggctggacttctcttttggagcagagtccctgttccacggatcgctgtgacaagcagaaaagtgtcccccacccacc  
tccatacaaatccctaattgctcctgggtgatacaaccgccccagtggaagcacgggtggtgcatggacttctagaga  
ccatttgaagccaggacattgctgcttgttcttaggaaatggcagaatattactttcttgtttttgtaggggaag  
cctgccatctcaccaggtggcagcttgtttataccttttatcctaacctgaagcaggtgatgggacagaggatcat  
cgaatataatgaaagtgtccacaaaggtagagtctgggtatattttatgtcaggttagagtgttgtccagat  
cccttcacacagagcaacatttaatatggtaattgttactgctgaagttggccatttctcggaacctacagctgca  
aggaattggtaacaaaaggataaactaaacattgttactatttttaattttcaaggagatgaagttaaaatcatat  
atgtcatatctctctggggcttagccaccttttctgctgggcactttttgaagaagtctgaatactgagatagga  
gagtaaaggggggaaagtaagtttgccacttctcatcctttttctgaccatcagcctgaggggaagtaactagaa  
tccgctaagaaaattcacttaatcagcagttgattgattgttatattgtccacattgcaaaattctttaaggat  
atttgaagatattatcacatttgttctcttacaatctattgtattgtttatttgaaggggtgagtgtaatatgtt  
cttaagatatcgtactttatttccggtagcatatccaagaaataatttagaagattgttaattgggaatgatgtt  
aaataatttttctatgactagtagttgggacatatatgatataaaacatgggtatctttgttacttaagtaatttg  
aactctaaccatataatcttaggttagacaatgaaaatactgaattgttagtctaaataatacacgtactcat  
tccgcaaatattttattaggtatctacacaccttaggaatctttcagggcatgaagctgttttaattctccaagat  
aatgcagcacgtatcccccttgaggggacatttgtttttcaaattgtcctctgtgtctcctctgcttaggatatg  
agttcctctaagactgtatttttaattcttttgtgtaattctgatgtctagcgccatgcctgggtgcttacgtaatag  
ttgctcagtcagtttgagggtgagtcatttaactgccctctaggagcttgatatttaataaaactagtctgtga  
tacatgacaatgctgtgacagtgcagatgagaaaatgtgacttctaactggggtgctcagagaggatatcctgga  
ggaactgccattttaactagacttttcttttaagcaggtagattttgtttgggttcaggcttcagctggaagccca  
agcaatgggtgcagatatgagctggactatgacagactattgggcccagtgagggtgacaaggttaacttgggggtt  
gccatagagggccttcagtgctcatgtgcaactgtttcaagtttgtagcaggcatctggaaaccatggaagaagat  
tctggaaagggcagggcaggttaaactggatgtggataggaagttactgtggcaggtgaattggaagatggacagg  
ttggaggcagggagatgggtgagcgggctcctgccaaaggtgtaggtctgagcattgttgtgacggcttagttggtt  
tgacatcagcacacagattgaggatctgtcattgttaactctaaattgtatccttgttagacatgtaaaaataact  
tttttagagcctccgttttttaaggagggaagtggttgtgctcaagcttgcatgctccgctgttctctgtgctttaa  
aaatactcatccaccagtgtggtcactgatgaaaggggaggaaaaactagccagaagttgccattttgtctaaag  
aatggattcatttcttctcaaggtggcacagagagagtttctctctttttgccccctctcttagattgatttact  
atttttagtaaatcttaacagtctgatccttgcctccaattacaatgattagcaatatattgcccgaagacaagaaa  
aaacaatttctctcttttctctcctatgagcttttctgtgagccccaagcagacagatgtggacatctagcaa  
tgctgttaacaacagctgcagtcctgttgactctctgcaacactgggcacgggtacaaagtgtctctgacaaactct  
tctgccctttgcaatccttactacgccctgtgaggtgaggagtagttcttctctgttttgtaggtgaagaactgagg  
caaagagaggttatgtcatttagatttccaaataggggtgtgttccgtttcatcatctttagattgatgttactgt  
gacatcatcttctagaattaacaaagtgaactgggcacaggaaatagcacatggcacaccttcagcaaatgggtg  
gttgtgatcattttcatcattccatttcttcttgaaaaaaatcccttttttttttttttttttgagacagagtctc  
actctgtcgccaggctggactgggtgtgatctcggtcactgcaagctccgcctcccgggttcacgccattcttc  
tgccctcagcctcctgagtagctgggactacaggcacggggccaccacgcctgggctaatttttttgtatttttagta  
gagacggagtttaccgtgttagccaggatgggtctcgaactcctgacctcgatccgcccacctcggcctccca  
aagtgtggtgattacaggcatgagccaccacgcctggcaaaaatccctcgagttccagtgtaaacactcacgccc  
ttttttgtgtttgtctataaacctctcatttctccttgggtcttcttgtccttgtgtgaggtcctgggtgtgtgga  
gactgaggtcttctgagacagaaaaccaaccccatagcaggcctgggtgtgtgcccctagaatggcagaggccccac  
acgttgccccggctcctctggaagaccctccagcctgtctgctggattccttgtacttatgaggattgttaaacc  
atctcataggattcctttccagactcagcccttcacttgtggcagcatttcttactctgaggttctgggcctaga  
aaccagtggtccataggcaaggtctagggacagagagagttttagaaagtgggaatgcatcagtggaaggctaa  
atccaaacaggaactgagctggcctgggtggctttgtccttttgccatcttcgtaaccctttaaggctttaaga  
cacttcttttgaataactaccccaactaaatttggccagagctgtgaagggaagaaaatctaagacccttgag  
ggagatgacagccgaacgttttacatttttaaccaccttgggtgttgccatgcacagctgatacataaacaagtaa  
tggcctcatcttctattcttttatttactgggttagaccacagaaccctaaaaggtagatgcttctcatgggtcattt  
ggcatttgaacaggagctgaattttattaatagtactaataacctcatccttggacatccttgggttgtccgatat  
ttttaaggggcatttacatcctttgtttttcatatggaaaactttccgtaagggctgggtatcatcttggccttc

cattgcaggaacagcacagggaggttcggtggctagggccacagttacccacgcagctaggtagcagcagaggccc  
cacctggcacccttggtctcctaagcacatcgcaactgtgcagcctccctgaggaagcctcgagtggataggaac  
tctggttctttaagataaacctgagttggatctcactgtcaccatttatcagatggggcgatgtcttcaggcaaa  
agtacttaacctctctgtttttgtttccttactccaaaatggtaatgatcatactactacccacctgggtggatt  
ggtgtgcatatgaaggaggtcatggggacaagtgggttggaagagtgttgagcacatagaaagtacaagataaatg  
gtggccattatgttactacttagactatatcatacttgtctttatctggtgcttagaatcttgacactgagatctt  
tcattctcagatcttaatatgaaagcttaccaggtaaacactaccccaacataattatcttagtgggttcatacat  
gtgaaagcagttgaaatacaatttctgaactccgggttaatttatatccttagttggaacaggtagtagccctccc  
agcacctaaggcctctcacttctatccattcttcagatgaccttgcataatcatggtaattgaatagcatctg  
caccataacattgttgaaataacatcaaaatgatccatttagccatcatgagccttccctagtaattcagactcaca  
ccagttctttcaggttattgagtagcttttacaataattgctaacaggtacaaatactgtctcaggtattttgga  
aggggtgaattttccatttgatttcagtgatcttttagggaggttaatttgcaaaaccgaagagtttggttccgtcaa  
agtcacccttaaggagtggtggaggaccacattgtagggttatactgcactgtccctcatgcttgaaatttggtc  
atgcaaccaaggaactgaatttttaatttcgttaaattttgatttaaattttaaaacagaaatagtataaaatat  
ttttctgttaaataacattgtagtggttggaagactacatttcgtgttaccatgctgggtaagatttctttag  
tgtgtattgtgtttgtcacctcggttaataattttcttatttttggttgcatgttaaataattttggatatattg  
ggttaaataaatatattgttaaataatttaattcacttggtttatttttttggttggtgaggttaggctactaaaaat  
ttaaataattacccatgtggattgcattatatttctattagtgctgctttagctgcaagtagcctatttaaaaaaa  
tcattattttcagacttatggacttcatagaatcagaggtatttagaggaaaggatcttgggggtcacctaattcc  
agctccaacattttgcaagtgaataagaccaaggccagaggtacaataaaatgactgacctcctcccagatgggg  
actggatcccagtggttttatagcctttgccttggtcctgggtttacttggaacatagcagacatcagagctg  
cagagaacttctaccagatccggagtgagaaatgccttatttatttatttatttatttatttatttatttatttattt  
ccgggtattgtgtggccttaggctgtagttagtgccagaaacatgggtcactgcagcctcaacctcctgggtc  
caggatcctcatgcctcaaccttctgagtagccggggccacaggtacgcgccaccacgcctggctatttttttt  
ttctcttttgtagagactgcgtcttccatggttgccaggtgatataaaaactcctgggtcaagtggctcctcct  
accttggtcccccaaagtgtgggataacaggtgtgagccatcatgcctgttttggcagtttttagggggccacg  
ctaattttgagccaggtttgcgttgataatgtaatgctggaggtgggcaggggacatgatctgcctgaaaggg  
ctgagaactgggaggtccttattcctatagatttagagagcaaaggccacagtccttacagaaccacacaatattcag  
agcttgtgggttgagagaaggaaagtcctctgaaaaataaaaaaagagtacagaggccagagggcggtggctcatg  
cctgtaatgccagcactttgggagggccagggcagggcggtacacctaagggtgaggagatcgagaccagcctgacca  
acatggagaaaccctgtctctactaaaaatgcaaaattggccaggtatgggtggcgatgcctgtaatccaagcta  
cttgggaggtgagggcaagagaatcgcttgaaacccgggagggcggttgcggtgagctgagatcacgccattgc  
actccagcctgggcaacaagagtgaactctgtctcaaaaaaaaagaaacagagtacaggtcagacattgggca  
attttctcataacaatttcataggccatttaatttggttacagacttgaatagcacacaacagttctctaagggtc  
agtttccaaagtaattcctaataatttaattgaatttttaagtgtatagaaattgcagtaggaaagtctagcttga  
tctaatttgtgttaggttgaccaggtgacaaccacccacccctggattctgctcagttaggtcagacctggggcc  
agttgcccatttagcatctcaggcctcaggatcctcatttataaacctggggttgaaactgcggtctccacgatc  
actgtcagctctccttagtcagttgggtgttctaaatgtagtttagtggtccagtaaacacctgttcctgagtgat  
acatctttaaggagccggttagatgggtcaacctgggtgacttgctttttcaagtctgccttgctattagaaag  
gctgagcttactgattttgcctgtcacgtttgagtggtcctgagactttgcccagcctcggtcttatctgcgggt  
acccagcctctgcattccttgcccctacaaaatgtgctgccagttccaaaggccacaaatgaaattagcttggc  
tgagcatggcttaggcacacaccagtggttaaagaaatgctgtttgctggctgacacttctggagtggaagtta  
ttcttttttcttttttttttttttttttttgcgggggtggggcagagttttgctcttggttggccaggttgagtgcaa  
tgccacgatctcggttgagcgatcttgggtcaccataacctccgcctcctgggttcaagcgatttcttgcct  
cagcctctggagtagctgagattataggctcccgctaccacgcccggctaattttttgtatttttagtagagaca  
gggtttcaccatgttggtcaggtgtgttgaactcctgaccttacgatccacctgcctcggtcctccaaagtgc  
tggtattacaggttgagccacggcgccggcctttattcctttctttggcctcaattttattgttaagagatag  
tggtgaaggcccaattttgaaatgatcctggaattaatgtaggaggtcctgtataagcagttaatgtgtatttat  
tgagtgggtaccctgtgccaatatgatacaaggtgtgaaggacacatgagttgggaaaggcatggggaaatgcac  
gctcacactgctgggtgcgagtgtaaaaggtacagtcctctatgtagggtattttggcagtatatactctttgaccc  
agcaactccacttttaagtatttatcttagggataccctcacacatttatgaaatgatttatatacaaggatatt  
cattatagcaatatttgaatggcaaaaagaaacaaaagtcagatgggggactgattaaataattatgttattaa  
gtaatactgtttcagtagcatctgtgtaatgaaataatgtccagtcattaaaaatagtgaggcaagctctgggtgc  
agtggctcacgcctgtaatccagcaccttgggaggtgaagccggaggattgcttgagaccagcctgggcaaca

tagtgagaccctgtctgtacaaaaaaaaaaaaaaaaaagtaaaatattagccaggcatagtggcacatgcctgta  
atctcagctattcaggaggctgaggtgggaagattgcttgagcccaggaggttgaggctgcagtgcagctgtgatt  
ctgccactgcactccagcctgggcaacagaacaagaccctatcttagaaaaaagaaaaaagaaaaagggaca  
acttacaactctgtgccctgatgtagaacatctccaaaatatattaaatgagtggtgggatactatgcttatatg  
cgattgtgtgtattttaatggcttcctaagaaaacaaaaaaaaaacctgatactagtgccttgaggagggtta  
actgggaaattacttttgcacttttaaatcttataccatgtacacctgggtatctatttttaaaaagccattttc  
atgccctaaatgagtttagtcagaccatagtaagaaaatcctgtaagacatacaaagtatggggtaattactggc  
tttatgggaagactgatttcaatgcaaaactctccctatgcaagggttgctctgatttcaatgctaagctctc  
tttaggcaagttacgttggcctgatttgcctgttaagtcgtatggggcaggcaagtctgagtgccagagggcag  
acctttaactcttagtttccctgccttgaggagaaggcacagggttaaagtggtttccagaacgtgtctgttgggt  
gtgaacaccaaggagcagtgaggaagcttccagggtcaagggtcagaaggctctggactggagtcctgcctctgct  
acttgggtgaatatgtgatttggggcagggttgcttaatccccgaagcctcaacgtgttcatctagaaatgaagatt  
acaggctggagcgcagtggtcatgcctgtaatccagcactttgggaggctgaggcaggcagatcatctgaggtc  
aggagttcgagaccagcttggccaacatggcaaaaccccatccctactaaaaatacaaaaattagctgggcatgg  
tggcagggtgctgtaatcccaactactcgggagggtgaggcaggagaatcacttcaaccaggagggtggagggtg  
cagtgagctgagattgtgccactgcactccagcctaggcaacagagtgagactctgtctcaaaaaaaaaaagac  
tacaatacctggcccaactccctcatgctactgtagtgggtcaaatagagatagcaccatgaacactcctgta  
aaccatgaagtatgcaaacaccagggtgtaatagaagctgttaggtacctgtgaggccagcagacaagagcaagag  
atgctaattttaaagaattaagggtgaagcaagatcttttccctctgccaataacttggcaagagttgtaaaa  
ttagaaaagtagacccttagtagtttgataatcctttgacctctgacctgcacaaatgatctcaccctttagac  
ctgttcccttatttgcacaaacaagaggggtgtctgggtgattcctgtgggtcccttccagttgtagcgttccgtg  
acagtggtggcattaacagtaattcccatcttggctgagatggatgagtcataactgaaaagtcaaaaatacga  
ggaagaagtctctttatgtatgagaattttccttgagctagagactcagagcctcttggggaggagatgagggaa  
aacattgccaccaccaagaaggagacgaaggatgtataaagagaatggagatgtatttacttttttttttttt  
tttttttttgagacagagtctcactctgtccttctggctggagtgacgtggcagatcttgggttactgcaatc  
tccacctcccagggttcacgccattctcctgcctcagcctcccaagtagctgggacacaggcgccaccaccacgc  
ccagctaattttttgtgttttttagtagagatgggggtttcaccttagccaggatgggtctccatctcctgacctcg  
tgatccgcccgcctcggcctcccaaagtgtctgggattacaggcgtgtgccaccgcacccagccgagattatttac  
ttcttgtgtgttcttgccatctccatagcctcctttgctttcaaatagccacttgggctgggccttttgaaaaa  
gatttaacccttgtttatttgaaggatattcacacagtagggcagttatacactttgtggcagtcgttacctctt  
ctgtgaggtctttctgtgcagatttagtggaacctcctccatgctgcctgggcttctacttgccttctcatcactt  
tgtctcataactcagatcagagactgtgtcttactcatctttgtatccctggcatctgaggggggtggatctgat  
gggtagatggattagtgatccacttagggtatttacactaatcttgttgatcttgactactacctttataaccac  
caccattctacaagttttaacttttttttttttttgagacagagttttgctcttgtcaccagcgtggagtgc  
tgggcgatcttggctcactgcaacctctgcctccccgggttcaagcgattctcctgcctcagcctcccgagtag  
ctgggactacaggcaccgcaccatgcctggctaatttttttgatttttagtagagacgggggtttcacctgtg  
tagccaggatgggtctcaatctcctgatcctgtgatccacctacctcagcctcctaaagtgtctgggattacaggcg  
tgagccaccgcacccagcaaggatcttttttttttttttaaagctaagtgtttatacatataataatattgtt  
aataaaatgtatctaaaattcataaccaccagcacacaagggtctccggctagcagcagtaaatggcatttagtact  
tacttagcccccaagcctggatgactgactatactgacttattattacttttagctattttgtaggtagttacatc  
tgaaaattgaccaggcacagtggttcatgcctgtaatctcagcactttgggagaccaaggcaggaggatcagttg  
gggccaagaatttgagaccagcatgagtaacatagcaagaacctctctacaaaaataaaaaattaaccaggcat  
ggtgatgcatgctacttgggaggccaagggtgggaggatcacttgagcctgggaattagaagctgcagtgcagtag  
gaagggtgccactgcactccagcctgggcaacagagcaagactcttaatctcttttaaaaaaaaaaatctgaaaagg  
accccaaggattgaactttatctttacattttatctttacaaatccaggatcaaaagagaggggtaaaggatcactt  
ggggagtttagaagtggcagaagtatctgtagaagccttacggtgtgaatgtgctggattgtgtgtgccaccct  
catgggggacttcccttgaggatgaggggtacaccagcatgggcagcaaaagctgtttgacctcaagggtggagatct  
gagccacctggctcagtcctatgccagtttaatttctatggagggttggttaaagtaggttagggagttggatccaat  
gttccctgaagtttctgttggtaatatggcaggggtcgggtcgggtgtggggagggttggtatcattctaagggcag  
tacaagacctaatcccaaagggtgagaagttcctctctgtggaggacagcatttagaactcaatacatataggt  
tgttcaaaaatggggaagagattagaaaaattatcctaatgtttgcttgtcaaaccaagccattctctcagcaa  
agtagtgaaacccttacataagtcctttgaagcctagcagcagtatcctactcaaacacaaaaccacccccaaaa  
aacaaccccccaaaaacccctccttggggactcagatgtgaacctctcatggaaaattcagtagcatgtgctttt  
tgtagcaggtaggatcacctgctgttttgaaaattaacttttttctgacatctttttgaccgaactctgcatgt

gcctcttgagaaaataatccaaattcttgcctttcttcactttccagttttgctttatctttgcatcatgctctgc  
agtttacagacaagcataatgggtgagagttccagtaaattggctggatcatatttcttagattgagtcagtgact  
gaggaacttggttgggaacaggaagcagcccatgaaaatggctgcacatagtaacctgccagattgaagaaaccg  
gaccactttctagtggaagataaaaagggcagaaaaatggttctataactcaggggaatggtaaaatttgtaacta  
gggttttaacttggtgaataatacgtctttgtgtgattttgggctttttatgctgttaaaagtctctagttatt  
caagataagggaaaggaaaacttggattcgtggtaactgttgctacctcttaaatgtctctgctgatgtatacaa  
catgttgctaggccagttattttttttaaggcactgtagttacatagaaataaacctggcaacatttacacttgg  
cctcgataggctgtaaggccccctttgtctccagttattctgtgtttctgtatttttaacacctactttacactta  
ctgtagcaacacaatcaagatagcaaatcagttctaataaaaccaaagtactggaatgtgggaaatagaaagaat  
gacctagaattcttgctgagaaaacaattggattgtctatattcttgtaatgcagtggtgagagaagccagagaag  
taaataaggcagatacatatctatgaaatgtaattagtcacagcaagatgtattttaaaaacacatctgcttgg  
gtcattttgctttatctaataaggattacgactgcaaggcagagatttgtaaataagggagatctctgttgtttt  
tgttttgtttttgctttgtttgttttaagacagggctggagggcagtagatgatcatggctcactgcaacctt  
catctcctgggctcaagtgtcctcccacatcagtttcccaagtagctgggactacagcagtgctcaccatgtc  
tggctaatttttaattttttgtagagttgaggcctgtgttgccaggtggctctaaactcctgggctcaagt  
gattctcacacctcagcctcccaaagagctgggattacaggtgtgagccaccgtgcctggctgttattttatcaa  
ttcctcaaaattcaagcatgccctcaaataatttttgagcctcatatagctcttaatttggtttatattcatcca  
gatggtaggatatgattcatgttccacctgggttttcttcagaaatacttagacagtgctatggaaaaaacatt  
gaaccagatttagcttttaggtaaataataataacacctgggaataatatgaagaatgaaatgaaagaaagaaaa  
atataatcaagtgttaacttagactctgaatgtctggtaactaagatttggtatgaaggaaggatctcaggggaataa  
cattaaaaataagatttttttttaaatggaaatacatgattttaagaaggaataacttaagacattccaactttata  
ctacatgaaagtaaaaagatcaaggatcagttttctctttggagcaaggctcagtttctgctttctgttgagagtt  
agtagtgtgtataattaatttttttaacagtttattgatgaaattttatttttagaaggaaatctcccaccacct  
tcctgtcattctcagggcaagatagagggtaggagaggggaaggagactcgggggaatgggaggtgcaagcctta  
attagacatttgagtgtctaaaagattacttttcatttcaaatactttgtctcagtttactgaatagctttaatt  
taggacagggattgtattatctttgctttatcttttccaataattttggatcatctaagcatgatagatgctggtt  
tattcaaggcactaaaattgacttgcaaaagacataatttaatatggattctgtgttcaagtcctttcttctttgt  
ttatcctgggtacaaagatgtttgttttaaaagtgggtatgagtttaattgtccaacttcagatttccctaaaatgtgt  
aacagattaccagttggcaaatgatgatataattttttaacatgaataatttctgttaggccagatgggtgatt  
taacatgaattgtccaacatttgtgtttcttatcaaattgtcttcttttgtctaaagccaagctagcaacataatt  
tggaacacactcaggaatcactaaaggacaagagcaaaaaaaattcagacaccaagtgccagacggcatttgctc  
catatgccatgcaaaggctcctggccagtgctcgcagtgcccttggaagctttctcaccctgttgaagaaagta  
tctgtgcagtgctgtgcaacccatctacctgctttcctctctctacccaaaccctacttatctaacactttgtac  
tgtggccttccctcatgggggacttgcaacttcccaagccctggatttctgtgcagaagagcagttttagttct  
tccttaggcacttccctttcctcctgtccagtgctcatgttcttgtttgttatgggctccaccagacgattattt  
cctcttagatgatgtgtgagccttgacagagagctgggtcatttgtggaataaatcatgtgcccagtcacacggct  
ccttacaggaaatgggtgggtgcataattagcttagcacaaaatacagtagtgagcttggtggtcccaacagtgagt  
cttcagtttaacttctgctgttttttttttttttttaacataaagctttgaatgtttcatacagtatcaaatc  
tctgcaggattttgctgctgaaacagatgtgaaaatctgagtagcacagagtatgacattaaggccataacaat  
aaacctattgttgctgcattctaaatttgatcaagatgtctgtttttgggaatccaaaatagttgcttcaaattt  
agcttttttcttttatgaagcagttttgttttgttttggagacctctcatagcaagaaaaaacattacata  
aacttttttttgagacaaggtctcactttgttaccaggtggagtgcaagtgccagatcttggtcactgcagc  
ctcgacctcctgggctcaagcagtcctcccaccttagtccccacgtagctgggactacaggcacatgccaccac  
gcccggctaatttttgtatttttagtagagacaggggtttccacctgttgccaggtgggtctcgaactcctgagc  
acaagcagatctgcccgcctcgccctcccaaagtgttaagattataggcgtgagccaccatgcctggccacataaa  
tataaacttaaaaaatacacaggaaaacctttcagaacactttgaaacctattttgggtcctattacgattgcactg  
aatgtccccagcctaataagagcaattagatgtcatttgggtggcattcttgggtggagacagtaagctagggcc  
gcaggagctgcttagtctcccaggtccagcattgtttcaggtaacgtgaacaggatgctcaagagctagtctaa  
ataatgcttcatgtcttatttttctgtttgttgataaggattcataaagatgtctttttctggtcatgtcaagg  
gaaattcaaataaaaaacattgaattgggttgagctgtccattgaagggtactcacagcaggaattgagtgga  
attgaatgtgtgtgtgtgtatgtttgtaaacatttatattagagtgatgcaaatgtatatgggctgcccactg  
ccatcaccattacccccaccaacagtaacaaaaccacctcaagagcaatatccactaaaacttctaatttgctt  
attaatttgctttatttttgaataattcagaatctctgagatcaaaatccttttttattacaacatgttaat  
agtagtctaattcagacctgcctgaggaagaaccaagcaatgaagctgtccatgattttagaagttaattggct

cattctttctgaccatctaggtagtcaccaacattatcttaggctccccctagtttaggtccacaacctcattatta  
tccattggcctggacttttgggtctcgcttatacagcaagtagtagagcctatcactgtcaacatgtaaacattttgc  
atattttacatttatagagagatacttttcaagtcgctcatttgatccactaacatctgtatgcttgggtaggat  
ggaataattaaacaatgtatagctaaagccatccaaacccctcaaataattatcatctgatgggaaggaaaagg  
gggaaaggaaattaaatagttatcttcaattcttggcatgattgacactgacaattataaattccattcaatattt  
tcttcagctcatgaggggttgggtgtaaggatactttgcaagtcagatttatgagcagaagatgtttgcaaccta  
aattgcttcagttgcatgaaaaagtcacaaccttttactgatttttatcctgggatctccttaacaagaaaaaac  
ctcaggcccaggaagtctaggggtgagggcagggatgtgtcaaagtacacttttccccacgtacagggacaagtatt  
tgagtttctttgattgactttgcaaagagctttgcacactttgcacatcacttcttgagatgggtgggaacatgg  
actgcctattttacagagttcaattaagatcgagggatcatgcattcctttcatttttgggtgctagtacctagaa  
gatgggaacatattgaacacttaataccttgcttgggcagatagctccagtttgagaaagtctagacagggatc  
cctgctatgtatctagatttactgagaagaagcgcccaagctaatacagaattttaaagtaggtttctcagctgg  
gcgtgggtggctcacacctgtaatcccagcactttgggagggccgagggcagggcagatcacttgaggtcaggggttcg  
agaccagcctggccaacatggcgaaatcctgtctctactaaaaatagaaaaattagccgggcatgggtggcacgca  
cctgtaatcccagctattcaggaggtgaggtgagagaactgcttgaacctggagggcggaagttgcagtgaagct  
gagattgtgccactgcactccagcctgggagacagagcgagactccgtctcaaaaaaaaaataaaaaaaagtaggt  
ttctctatacctttgaattttctgtctcccatgataacctttgggtaaacatgtccatcagaatggaaacaaatc  
atatttttaaaatagaaattattctcatacctatttttagtgggaatccccctgagaagatagccatgtatagtcga  
ctgaatgctctgtggttgctattttgttgctcttggccaggaaactgatctaggtgacctacagacttaactctga  
cctgcagactggccgcttcatactaccaagattccaaagccaagtgtatatacagtcggcccatgcagcctctgg  
ttccatgatgcaaccaatggcagattgggtcaaaaaatatttggaaaaaggccaagcacagtgagggtggttcatgc  
ctgtaatcctagcactttgggaagctgagggcaggaaaaatcaatggagcccggaattcaagaccagcctgggcaa  
aatggcaagacgcatctctacaaaaagtaattagctgggcttggtggcatgtgcctgtagtcccagctacttgg  
gaggtgaggtgggaggatcacttgagcccaggtgaaggctgcagtgaagccatgattgcatcaccacactccagc  
ctgggtgacagagtgaagccctatctcaaaatgagtaataaataataaaatttcaaaaggaaaaagtgga  
aaaacccgaaaaaataacaatacacacataaaatacagtataacaactatttacacagcatttacactttactag  
gtattgtaagtaatccagggtgagttaaagtataaaagggtatgcataggttatttgcagcaaatgccatgc  
cattttataatagtgtaggctcatggctaataaagcatgaacccgactgaagctgcctgggtatgtgtctttgtt  
cagccactttctggtagtatcttttttcttaacctcagtatcttatactgtataacaggaatgttttctctgta  
taaggctgttttgagggaatagtttctatatgtaaagtacttagaacagtgcatagcatataggaagcccttaata  
gatactggcttaaagataattaggtatgtcctagttaggtatacctaactacaagtgtactttaactatgccta  
ctttgtgtgctttacaatatataattgttcaaggctagaataatgagcccatgagcaggggtttcatagcatttaa  
gattccaacgccaaccttcaatcatggacacaggtggcctgtgccatctagctagctgcgctgcctccaggtgg  
gtggcctgctcctctgttggataaaaaatagcctgactcctcactgaaacaaccacatgttatttgcgtccttgg  
gaagctgaaagtaatagaactctaagctctggcgttgccagctgtcagcagaatgaaatgcaaagtgtgacgtgtc  
tacactgtaccactggccttcaagtcaccaactcccggtcccccatcttgattttttccattctcaaatatt  
atctccctataggtgatttgtggaatggaaacagatttagatttctacctatttagatttttccaaagactaa  
aattacaaatcagaaaaactgctattgacacatttttgggttaggaaaatctggatgtggttccatgcctagcct  
agtcataataaattctctccccatgtatcttttgtgtgtgtgaaacagggctcttgtcacccaagctggagtgcag  
tggtacaatcacagctcaccatggccttgacttccctgggctcaagcaatcctctcacttcagtctccctaatagt  
tgggactacaggctcgtatcaccacgtgcaactaatttgttttgatttttgggtagaacaggtgtcactatggtg  
cccaggctagtaactcctgtgtcctcagcaatcctcctgccttggcctcccaactcctgggattacaggcctgag  
tcaccaagctcagcctctcctcatcaaaaccattccctctttgtaagattcttccatgtcttaaaacttaatac  
ctttaatccttccctggagccatagttttatttatttatttatttatgtagacaggggtctcactttgttgcc  
cgggctggagtgcagtgcacgatcatagctcactgctgccatgacctcctggactgaagtgatcctcctacctc  
agcctcctgagtagctgggaccacaggtgtgagccaccacaccagttgttgttgttgttgttgttgttgttgtt  
gtagaacaggggtctctccatgttgccaggctggtgcggaactcctgggctcaagcaatccacctgcctcagcc  
tcccaaagtgtgggcttgtaggcgtgagccaccgcacctgtccctgtctgcttttattttattttttttgaga  
tgaggtcttgcctgttgccaggctggagtgcagtgggtgcgatctcagctgactgcaacctctgcctcccaggt  
tcaagccgattctcctgcctcagcctcctgagtagctgggattacaggcacgtaccaccacacccggctgatttt  
tgtatttttagtagagacgtgggttcccatgttggccaggctgatctcgaactccagacctcaggtgatccacc  
caccttggcctcccaaagtgtgggattacaggcatataggttaagccacagagataagagtagcaagcagtgtg  
gggagaagataatctagtgtaggaaagatggaaagaatgggtgatatttcacaactgctagattgacactttaact  
tgagagttatccctctataaactggatatttgcgtgagcaagtttaaagaaaactccatgtattcagtaattgtt

tccagacccccacttctcttatataagccagtggttagagtgttgcctctaatttctatattccattaaacaagg  
gtgggatttcttctcattcaaaccatttgtgactttgcccttagtgacctctaccatcacccctatcattaatga  
tccaagtgattagaatggatggctatgtgtttgtaggttgtttgttgcctttaatgaatagattcattgaatgga  
attaactgtgtcttcttagaacattggatctgtgttctcatttagattgtatttgtaatctccccacttcccacc  
atcaacagtggcataatgtgaattaatttatgtgttatctgtcatttaaaaaacatgagctctctccttccctttt  
tttgtcttaaacatagGACATGGATTGTGATTGACATACTTTGGAGGCAAGATATAGATCTTGGAGTAAGTCGAGA  
AGTATTTGACTTCAGTCAGCGACGGAAAGAGTATGAGCTGGAAAAACAGAAAAAACTTGAAAAGGAAAGACAAGA  
ACAACTCCAAAAGGAGCAAGAGAAAGCCTTTTTCGCTCAGTTACAACCTAGATGAAGAGACAGGTGAATTTCTCCC  
AATTCAGCCAGCCAGCACATCCAGTCAGAAACCAGTGGATCTGCCAACTACTCCAGgtacagagtactcagtt  
cttgggaaagttatggcaggtttaaggaaacactgagcaaggaattaaaatatctggatttgagttccagctttg  
cctttcttttacttaaccttgtcaaactctactttccaacctcagcctcctgatgagttcagtaacctaacctgagt  
tcagtaaccaacctaattgatcttaacagtggttattgtgaggattgggaagacttaagttacaccaagagttttg  
taaagtatagaacatcctgtgaaggatcaagtagcagcaacagaagtagtagcaggagaaccaagtagcagggat  
tactggcgttactgtgtgtgtggcaagcactgttttaagaacatatactgactgatttaattaacacaactatatta  
attagttaccattatccctgttttatctacgatgagcaactgaagctcacaaggttaaataatttgcccagatc  
actcagaaaattggaggagctaagatctgaacccatgtggtctagttcaaattgtgcatcaaagtgtatctctgaa  
ataagataaatatttacttaacttgattataaatattttatgaacatcaattattgaatatttagcttggcaatg  
gaatatttaaccatttttgttttcccttgtgtcattcccttttatcagGTTGCCACATTCCCAAATCAGATGCT  
TTGTACTTTGATGACTGCATGCAGCTTTTGGCGCAGACATTCCTGTTGTAGATGACAATGAGGTGAGGTATAAA  
ATAACCTGGTTAATAGAAAACTCCATCATAACTATAAAATAACAATCTATTCTATGTAAGTCCCGTCAATGAAT  
CTCCATTTAAAAGAATAAAAAACATTTTATAGGAGGAAATTTTAAACCAAGGAAATACTCTTGTCAGGAAACCTT  
AGCCTATAAATAACTTTACAATTAAGAAAAAAAAAACCTTCACACAATACAAAAACCAAAACCATTGACTATTG  
CATAGCCAGAAACATGGACAGCATAACCATGGAAACAAATAACCCATTTGCTGCAAGTATCTAAGAGGTTTGGTG  
AGTAAAGAGCCAGCTGGGCAATAAACGAAGACTTGTTCAGTTAACAATTTTAATAAATCTGTTTTATCTAGTACC  
ACTGTGCTAGATATTATATAAACTAAATCCTAAAGATTGTACTTACGCATTTTAAAGTTTACTTTCAAATGCTTA  
AGCTGAAACAGACCAGCAAATTATAAATTTGAGTCAGTGGGGTAGGAAAAAAGATTTGTTATTTACAAACGGGG  
TCATGACTGGTTAGTAAGTAGAGAGACACAGAACTGCAGCTGATTCCATTTTGTGTTTGTAGTGGTGCCTTAGAGC  
TTACTCATCCCCTGTTGGTGGAGACTCATAAATCAATGCCTTATCAATTTTAGGTTTCTTCGGCTACGTTTCAG  
TCACTTGTTCCTGATATTCCCGGTCACATCGAGAGCCAGTCTTCATTGCTACTAATCAGGCTCAGTCACCTGAA  
ACTTCTGTTGCTCAGGTAGCCCCGTGTTGATTTAGACGGTATGCAACAGGACATTGAGCAAGTTTGGGAGGAGCTA  
TTATCCATTCTGAGTTACAGGTAATAAATAGAAATGTAATACTGGAGATTTTTTTTATATTCACTGCCTTTAG  
TCATTCTGATTATTTATATACCACCTATTTATAGGAAGGATTGGAGGGTGCTATTAACCTTA

### FIGURE 3

Natural antisense sequence (BU656954.1): SEQ ID NO: 3

AACAGACTGGCGGCGCGCGGAAAACGCGTCACGTGACGACTGGCCCCGCCTCTCCTCTCGGTCCCATATTGA  
ACTCGAGTTGGAAGAGGCGAGTCCGGTCTCAAAATGGAGGTAAACCGCCGCCGGTCGCCCCAGCCCGAC  
TCCGGCCGTCGCCGTCGCCCGGGGGGAGGAGGGCCATGATCCAAAGGAACAGAGCAGTTGAGAAAAC  
GTTTATTGGTGGTCTGAGCTTTGAAACTACAGATGATAGTTTACGAGAACATTTGAGAAATGGGGCACACTC  
ACAGATTGTGTGGTAATGAGAGACCCCCAAACAAAACGTTCCAGGGGCTTTGGTTTTGTGACTTATTCTGTGT  
TGAAGAGGTGGATGCAGCAATGTGTGCTCGACCACACAAGGTTGATGGGCGTGTAGTGGAAACCAAGAGAG  
CTGTTTCTAGAGAGGATTCTGTAAAGCCTGGTGCCCATCTAACAGTGAAGAAAATTTTGTGGTGGTATTAAA  
GAAGATACAGAAGAATATAATTTGAGAGACTACTTTGAAAAGTATGGCAAGATTGAAACCATAGAAGTTATGG  
AAGACAGGCAGAGTGGAAAAAGAGAGGATTGCTTTGTAACCTTTGATGATCATGATACAGTTGATAAAAT  
TGTTGTTTCAGAAATACCACACTATTAATGGGCATAATTGTGAAGTGAAGAAAAGGCCCTTTCTAAACAAGAGATGC  
AGTCTGCTGGATCACAGAGAGGTCGTGGAGGTGGATCTGGCAATTTTATGGGTCGCGGAGGGAACCTTTGGAG  
GTGGTGGAGGTAATTTGGCCGTGGTGGAACTTTGGTGGAGAGGAGGCTATGGTGGTGGAGGTGGTGGC  
AGCAGAGGTAGTTATGGAGGAGGTGATGGTGGATATAATGGATTTGGAGGTGATGGTGGCAACTATGGCGG  
TGGTCTGGTTATAGTAGTAGAGGGGGCTATGGTGGTGGTGGACCAGGATATGGAAACCAAGGTGGTGGAT  
ATGGTGGAGGTGGAGGATATGATGGTTACAATGAAGGAGGAAATTTGGCGGTGGTAACTATGGTGGTGGT

GGGAACTATAATGATTTTGGAAATTATAGTGGACAACAGCAATCAAATTATGGACCCATGAAAGGGGGCAGTT  
TTGGTGGAGAAGCTCGGGCAGTCCCTATGGTGGTGGTTATGGATCTGGTGGTGGAAAGTGGTGGATATGGTA  
GCAGAAGGTTCTAAAAACAGCAGAAAAAGGGCTACAGTTCTTAGCAGGAGAGAGAGCGAGGAGTTGTCAGGA  
AAGCTGCAGGTTACTTTGAGACAGTCGTCCCAAATGCATTAGAGGAACTGTAAAAATCTGCCACAGAAGGAAC  
GATGATCCATAGTCAGAAAAGTTACTGCAGCTTAAACAGGAAACCCCTTCTTGTTGAGGACTGTCATAGCCACAG  
TTTGCAAAAAGTGCAGCTATTGATTAATGCAATGTAGTGTCAATTAGATGTACATTCTGAGGTCTTTTATCTGT  
TGTAGCTTTGTCTTTTCTTTTCTTTTCATTACATCAGGTATATTGCCCTGTAAATTGTGGTAGTGGTACCAGGA  
ATAAAAAATTAAGGAATTTTAACTTTTCAATATTTGTGTAGTTCAGTTTTTCTACATTTTAGTACAGAACTTTA  
ACAAAATGCAGTTTCGAAGGTGTTTCCTTGTGAGTTAACAAGTAAAGAAGATCATTGTTAATTACTATTTTGTAT  
GAATTTTGCTAAAGTTAACTGTAAAGAAACACCTGCTGACTTGCAGTTTAAGGGGAATCTATTCTCCCATTTCC  
AAACCATGATATGAATGGGCGCTGACATGTGGAGAGAATAGATAATTTGTGTGTTTGCAATGTGTGTTTGA  
TAAATAGGATTGGGTATTTAAATTAGCATTTGTGAATTTAATAGCATTAAGATTACCTTCAAATGAAAAAAAT  
CTCAAAATTTCTATTTGGTTTTGTGCATTTTCTTTTAAAATGTAATCATATGATTTTAGTGTGTTAGACTTGCTG  
AGTCTAGCTGTGTTAGAACATCTCTATTCTACATTTACCTTGGTCAAATTTGAACTGCTGCCATAGGTTTTGG  
GTGTAAAGAATGTTTACTGCCCTCCATTTAAATTCTGAAAAGGGATGGTGGATGTTTTCCCTCTCCTACGTTAGA  
AACCATTCTTAAAACTTTTGAATATAGAACCATTAAGCCTGCTATATCTGAGCAAATTAGTGGGTACCTTTT  
TTTTCTATTTAAAGCACAAAGAGGCCATAAATCTTGAGTTACTTTAAATCTTTTTTTTGATACAAGTTTTGAGA  
GCAAGAGAATAAAAAATCATGTGTTATTAACCCCTAACTGGCTGGCATGCTTTCCTGTTTGTATTCTATACATTT  
TGCTGGATGAAACCAAGGATAGTTCAGGTATAATTGTCCAAAATAACCTAACTGCAGCAGAAATGTAGCACAG  
TTGCTTAGTACAGGCTTCTCACTTCTACAGACCTGAATTCAAATTTGGATAGTCTGAGTCTTAAATTTCCAAA  
GAACACACTGTTATTTCTGTGTATATTTCAACATAAATCATGTTGTTACCAATTTGTTTGAAGGCCCTGGTTG  
AGAAGAGTTTTAGTTAATAAGGTCATATATACATATATTAATATAAACCAATGTCTACTGTTTTGCTCCAGCTAG  
TGCTTACAGTTTCATTGAGCCCTGAGTATGTGCCCTGCTGTTACTCTCTTTGGTAGTTGAACGTTGAATTCAG  
TCTTTTGTTTAAGAAGTACTAAGCAAACAAGCAATAAAAAGGGGAATGGGGTGTGCTAGTGTGTTGAATATGC  
TCTCTGTTGCTCTAATTCTGTGCCTCTGTGCATTAATATTTGGATGCATGCAATGCCAGCATGGAAATTTGGTCT  
TCACACATACTGCAGTTTTCCAGAAACATTCACAAACCAATAAATGTAACAGACATTCCATTTGTTAATGGGCAT  
ATATGTGAAAAGCAGTGTAGAAAATAGGCTAATATTAGAAAATGGTTAAGTCCTAAATAACTTCAAGTGTGGT  
TATATAATGGACACTGTCAATGTTTCATAACTTAAACCTGGGTACCTGGTCAAATAATGCTTGGGAAACATTAA  
AATTGAGCTAAATTGTCTCAAGTTCTTTTATTCATATAAATAAAGTTTAAAGGAATGGGGGAGATTAAACATTC  
CTGTTTTATGTTTGTGAAATTGTTTGACACAACCTTGACAGTATCCTTTAATGGCATGAGGTTAATTGTACTGTT  
AACCAACTTTCTATGTTCTGGAAGTATTATAGTGAACCAATTCACAGTAAGTTGATGTTTACAACCTATAAG  
CAGGTGAAATCTGTGTATGTGACCTGTTTATAAGTTGTATTAGCTTAGCTCTTGTGAACAGTGTGGAAAAGTAA  
GCCATGAGGAGAGCGATTTAACCACCTTTAAGGACCTAAGATGTGCTTTTAAAGCACAGTGTGGATCACAGA  
AACTCACTAAGACAGGACTTCAGCAGCCTTTTGTGTTTGGACAAGTCAGCATAAATAAAGAATGACAAGGCAG  
CAGCAAGAGCTTCACTACAGAGAAGTGAAGGCATAAGATACTATGATGATAGTGAACACTTTCCAAAAGCT  
AGTTAAATCTGCTTATTACAACCTGAAATATCGAAGAAAGTCTAGCAGGAAGGAGCTCTTCGCCTTTTGAACAT  
CAATGAGAGATAGTTGCCACAGTCACTAGGTCTAGCATTTAGACCTGCAAGGAAGGGCAATAAGCATTAGGTA  
AGGCTTGAATTTGAATTTTTTCACTAATTAAGAGTAATTTTTTGTAAGCAAGGTAAGAGTAATCTTTTGATT  
TGCAGGTTGAATGAGAACCCTACTTGCCTAAATGAGGAATGTCTTTCCTACCATCTAAATACGAAGGTTTCTG  
GCTGGGTAAAGTTTGTAGTTGACAGTAAACCTGATGACACCATTTGTTTCCCTGCAAGTCTACATTACATATT  
CACAACCTTTGTCCCTCTCTAGTAGGCACATTGGAAAAATTTCTCAACTGAAAACCTTGGTACCATGTCCTAC  
ACGTTTTAAACCTTAGTTTTAAAAATCCCTGCGAAATAGCCATAAGTATTCATATCAAGTCAGTTGTGACTCC  
TTGTGTATACAATTCATTTTTGTGTCTTCAGGGTAACTCAATTTTTGGTAAAGTGGTTTCAGCTTTTGTGAAA  
ACCGTTTTGGTGTGTAAGCATGACACACAACAGACTCAGTAAGCTGCCCCATCCTCATACTAGGAAAACACCTTC  
AAAGGAACATTAAAAGTTACCAGGGCCAGGCACAGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCTG  
AGGCAGATGGATCCCAAGTCCAGGAATTTGAGACGAGCCTGGGCAACATAGTGAGAGCCTGTCAACAAAAAA



TAGAAAAATTAGTTGGGCTTGGTGATACACATCTGTAGTCCCAGCTATTTGGGAGGCTGCCTTGATATCAGGC  
 AGTCGAGGCTGCAGTGAGCTGACTGCCCCACTGTATTCCAGCCTGGGTGACCCCATCTCAAAGAAGAAAAGTT  
 ACCAGATGTCATGGGTAAAGGTTGGTCTTCAAGTGGCCTCATAAGTTGTCTTGCAATTTAAATTCAGGGAATTCA  
 TTGGACCAATAGGTTACATTTTCGTTCTTTTTTGTGTTTGGTTCATCTGTTAAGCAGTGGGGGCCTAATTACTGC  
 TCCTTTGTAAAAACACATTTTCCCAAAGAACACTGAATTACCGTTCAAAGTGGTTGTTGATGGGTAATAAGGGC  
 TGTTTTGCTGCCCCAAAAGGGCTTAACAATTTAGTCGGATAGTTTACTTAAAAAAAAAATCCTTTGGAGACA  
 TACTGAAAATGCAAACTAGTTTCTAAATTATCAATTCCTACATGAAGAAGCAGTTTGCCAGAGTTTAGTCTCA  
 GAAAATGACTGGTTGGCTCTATTTAAATCAGAACCCAATTTCTACGCGTGTGAATAAGGTAACAGCCTTTGAT  
 GAATTTCTTCACAACATGGTTTTAGTGAAGCAAACATTTTTTTTTTAAGGGCATTGTTCTTTCTAGTTTATTTCT  
 TTTTATGAAATAAAATTATTTATTTAAACAGTTCCATTGTCGTTTCTGAAAACACTACAGTATTCTCAGAAGTTGTA  
 GCAGCAGTAAAAAAAAAAAAAGTTGTTATATAAGTGATTGGGGCAGATTTAACTGATTTTGTTAAACCAATTTGT  
 AAGTTACTGCTTCTAATATTACACTTCTAAAAAGCTGAATTTATACTCATGTCCTAAAGGAGAATATGTGGTAAT  
 AAAGTATATTTGTTAAGTAACTAATTGAAATAGGCTTGGTTTTAAGAGTTCCAGTATATAATAATCACAAATTG  
 AAACCTGACAGTATCTTGGGAGTTCCAGTAATGTCACAAATTAGTGAATAAGCATGCCAGTGTGCAAGGGTAA  
 TGTAAGGATTGTTAGCCTATCTAAATATTCAAATTACTTTAAACTTAAGTATGTTTTCTGATTTTAAAGAATTC  
 AGAAGTGTTCTGTAATGGATTGAGATGTTTCATTTGTAGTATAATGAAATGTTTACAGAAAGATAACTTTTTTCAT  
 TAAAATATTTTGTAAATGTGTGTGTTGTTTTGTCACCTCACAAATGTTGATGTGACTTAAACACTATAGGTGAAT  
 ATTTTGACTTATTTTACCAGTAAGTAATAAAACAACAGGAACTTG

FIGURE 4

Sequence ID	Sequence
SEQ ID NO:4	UAGGAAAGACAUUCCUCAUCCUCAUUUAGGCAA
SEQ ID NO:5	UUCUGCUACCAUAUCCACCACUCCAC
SEQ ID NO:6	T*A*G*G*A*A*A*G*A*C*A*T*T*C*C*T*C*A*T

FIGURE 5

Sequence ID	Sequence
SEQ ID NO:7	ATCCUUUCUGUAAGGAGUAAAUCCG
SEQ ID NO:8	AAGACGAUGGUUAUAGGUGGUGAAGG