




(86) Date de dépôt PCT/PCT Filing Date: 2011/07/08
(87) Date publication PCT/PCT Publication Date: 2012/01/12
(45) Date de délivrance/Issue Date: 2022/01/25
(85) Entrée phase nationale/National Entry: 2013/01/09
(86) N° demande PCT/PCT Application No.: US 2011/043306
(87) N° publication PCT/PCT Publication No.: 2012/006497
(30) Priorité/Priority: 2010/07/09 (US61/363,039)

(51) Cl.Int./Int.Cl. *C07K 5/12* (2006.01),
A61K 38/12 (2006.01), *A61P 1/12* (2006.01),
C07K 7/64 (2006.01), *G01N 33/68* (2006.01)
(72) Inventeurs/Inventors:
ZADINA, JAMES E., US;
HACKLER, LASZLO, US
(73) Propriétaires/Owners:
UNITED STATES DEPARTMENT OF VETERANS
AFFAIRS, US;
THE ADMINISTRATORS OF THE TULANE
EDUCATIONAL FUND, US
(74) Agent: CASSAN MACLEAN IP AGENCY INC.

(54) Titre : ANALOGUES DE L'AGONISTE DU RECEPTEUR OPIOIDE MU DES ENDOMORPHINES
(54) Title: MU OPIOID RECEPTOR AGONIST ANALOGS OF THE ENDOMORPHINS

Tyr-c[D-Lys-Trp-Phe-Glu]-NH₂
(SEQ ID NO:1)



C₄₀H₄₈N₈O₇
MW=752.3646

(57) Abrégé/Abstract:

The invention relates to cyclic peptide agonists that bind to the mu (morphine) opioid receptor and their use in the treatment of acute and/or chronic pain. Embodiments of the invention are directed to cyclic pentapeptide and hexapeptide analogs of endomorphin that have (i) a carboxy- terminal extension with an amidated hydrophilic amino acid and (ii) a substitution in amino acid position 2. These peptide analogs exhibit increased solubility compared to similar tetrapeptide analogs while maintaining favorable or improved therapeutic ratios of analgesia to side effects.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2012/006497 A3(43) International Publication Date
12 January 2012 (12.01.2012)

(51) International Patent Classification:

C07K 5/12 (2006.01) *G01N 33/68* (2006.01)
C07K 7/64 (2006.01) *A61P 1/12* (2006.01)
A61K 38/12 (2006.01)

(21) International Application Number:

PCT/US2011/043306

(22) International Filing Date:

8 July 2011 (08.07.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/363,039 9 July 2010 (09.07.2010) US

(71) Applicants (for all designated States except US): **THE ADMINISTRATORS OF THE TULANE EDUCATIONAL FUND** [US/US]; 1430 Tulane Avenue, New Orleans, Louisiana 70112 (US). **UNITED STATES DEPARTMENT OF VETERANS AFFAIRS** [US/US]; 810 Vermont Avenue, NW, Washington, District of Columbia 20420 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ZADINA, James E.** [US/US]; 3813 Courtland Drive, Metairie, Louisiana 70002 (US). **HACKLER, Laszlo** [HU/US]; 1111 Lake Ave, #115, Metairie, Louisiana 70005 (US).

(74) Agents: **OLSON, Arne M.** et al.; Olson & Cepuritis, Ltd., 20 N. Wacker Dr., 36th Floor, Chicago, Illinois 60606 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:

24 May 2012

(54) Title: MU OPIOID RECEPTOR AGONIST ANALOGS OF THE ENDOMORPHINS

Tyr-c[D-Lys-Trp-Phe-Glu]-NH₂
 (SEQ ID NO:1)



C₄₀H₄₈N₈O₇
 MW=752.3646

FIG. 1

(57) Abstract: The invention relates to cyclic peptide agonists that bind to the mu (morphine) opioid receptor and their use in the treatment of acute and/or chronic pain. Embodiments of the invention are directed to cyclic pentapeptide and hexapeptide analogs of endomorphin that have (i) a carboxy-terminal extension with an amidated hydrophilic amino acid and (ii) a substitution in amino acid position 2. These peptide analogs exhibit increased solubility compared to similar tetrapeptide analogs while maintaining favorable or improved therapeutic ratios of analgesia to side effects.

WO 2012/006497 A3

MU OPIOID RECEPTOR AGONIST ANALOGS OF THE ENDOMORPHINS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application Serial No. 61/363,039, filed on July 9, 2010.

STATEMENT OF GOVERNMENT SUPPORT

[0002] [This paragraph intentionally left blank.]

FIELD OF THE INVENTION

[0003] The present invention relates to peptide agonists that bind to the mu (morphine) opioid receptor and their use in the treatment of acute and chronic pain.

BACKGROUND OF THE INVENTION

[0004] Activation of the mu opioid receptor is among the most effective means of alleviating a wide range of pain conditions. Of the recently cloned opioid receptors e.g., mu (3,19,20), delta (6,9), and kappa (12-14), the vast majority of clinically used opioids act at the mu receptor. As illustrated in genetically altered "knock-out" mice, the absence of the mu receptor eliminates the analgesic effects of morphine (8), illustrating its central role in opioid-induced pain relief. The unique effectiveness of mu agonists can be attributed to several factors, including their presence in numerous regions of the nervous system that regulate pain processing and activation of multiple mechanisms that limit pain transmission (e.g., inhibiting release of excitatory

transmitters from the peripheral nervous system and decreasing cellular excitability in the central nervous system).

[0005] Limitations on the use of opioids result from negative side effects, including abuse liability, respiratory depression, and cognitive and motor impairment. Major efforts to develop compounds that maintain analgesic properties while reducing the negative side effects have met with limited success. This is evident from the recent epidemic of prescription drug abuse.

Numerous attempts at targeting alternative mechanisms of pain relief to avoid these side effects have generally been met with similar problems, i.e., a profile of adverse effects that are different from opioids, but often sufficiently serious to warrant removal from the market (e.g., COX inhibitors) or lack of approval to enter the market (e.g., TRP receptor antagonists). Over 100 million patients annually in the United States experience acute or chronic pain and frequently do not achieve adequate relief from existing drugs due to limited efficacy or excessive side effects.

[0006] Elderly patients tend to show greater sensitivity to severe pain and recent guidelines of the American Geriatric Society suggest greater use of opioids and reduction of non-steroidal anti-inflammatory drugs (NSAIDs) (10). Impairment of motor and cognitive function can be more debilitating in the elderly than in younger patients, particularly due to increased risk of fractures (7). Opioids with reduced motor and cognitive impairment are therefore a growing unmet need.

[0007] Natural endogenous peptides from bovine and human brain that are highly selective for the mu opioid receptor relative to the delta or kappa receptor have been described (22 and US Patent No. 6,303,578). These peptides are potent analgesics and have shown promise of reduced abuse liability (21) and respiratory

depression (4,5), as measured in rodent studies. The limited metabolic stability of the natural peptides led to the development of cyclized, D-amino acid-containing tetrapeptide analogs of the endomorphins (U.S. Patent No. 5,885,958) of sufficient metabolic stability to produce potent analgesia in rodents after peripheral administration. A lead compound from this group reportedly was 3-fold more potent than morphine in alleviating neuropathic pain and showed reduced rewarding properties in animal models that are correlated with abuse potential. While these results are promising, the development of additional compounds showing equal or better properties is desirable. The instant invention addresses this need by providing peptide analogs having unexpectedly better solubility and side-effect profiles than the previously described materials.

SUMMARY OF THE INVENTION

[0008] An embodiment of the instant invention is directed to pentapeptide and hexapeptide analogs of endomorphins that differ from the previously described tetrapeptide analogs by having (i) a carboxy-terminal extension with an amidated hydrophilic amino acid, (ii) a substitution in amino acid position 2; or (iii) a combination of (i) and (ii). The pentapeptide and hexapeptide analogs of the present invention exhibit increased solubility relative to the tetrapeptides while maintaining favorable therapeutic ratios of analgesia-to-side effects.

[0009] The compounds of the present invention are cyclic peptides that act as mu opioid receptor agonists with high affinity. These compounds provide relief of acute pain, chronic pain, or both, and comprise or consist of compounds of Formula I:

(I) H-Tyr-c[X₁-X₂-X₃-X₄]-X₅. X₁ and X₄ each independently is an acidic amino acid (i.e., an amino acid comprising a carboxylic acid-substituted side-chain) or a basic amino acid (i.e., an

amino acid comprising an amino-substituted side-chain), with the proviso that if X_1 is an acidic amino acid (e.g., D-Asp or D-Glu), then X_4 is a basic amino acid (e.g., Lys, Orn, Dpr, or Dab), and *vice versa*. Preferably, X_1 is D-Asp, D-Glu, D-Lys, D-Orn, D-Dpr or D-Dab; while X_4 preferably is Asp, Glu, Lys, Orn, Dpr or Dab. X_2 and X_3 each independently is an aromatic amino acid (i.e., an amino acid comprising an aromatic group in the side chain thereof). For example, X_2 preferably is Trp, Phe, or N-alkyl-Phe, where the alkyl group preferably comprises 1 to about 6 carbon atoms, i.e., a (C_1 to C_6) alkyl group. X_3 preferably is Phe, D-Phe, or p-Y-Phe where Y is NO_2 , F, Cl, or Br. X_5 is selected from the group consisting of -NHR, Ala-NHR, Arg-NHR, Asn-NHR, Asp-NHR, Cys-NHR, Glu-NHR, Gln-NHR, Gly-NHR, His-NHR, Ile-NHR, Leu-NHR, Met-NHR, Orn-NHR, Phe-NHR, Pro-NHR, Ser-NHR, Thr-NHR, Trp-NHR, Tyr-NHR, and Val-NHR; where R is H or an alkyl group (e.g. a (C_1 to C_{10}) alkyl group such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, hexyl, isohexyl, heptyl, or isoheptyl). The peptide of Formula I is cyclic (shown as " $c[X_1-X_2-X_3-X_4]$ " in the formula) by virtue of an amide linkage between the carboxylic acid and amino substituents of the side chains of amino acid residues X_1 and X_4 . For example, the linkage can be an amide bond formed between the side chain amino group of the D-Lys, D-Orn, D-Dpr, D-Dab, Lys, Orn, Dpr, or Dab with the side chain carboxyl group of D-Asp, D-Glu, Asp, or Glu.

[0010] In one embodiment of the invention directed to a peptide of Formula I, X_5 is NHR, R is H, and X_5 can be $-NH_2$ (i.e., the peptide is an amidated pentapeptide), or Ala- NH_2 , Arg- NH_2 , Asn- NH_2 , Asp- NH_2 , Cys- NH_2 , Glu- NH_2 , Gln- NH_2 , Gly- NH_2 , His- NH_2 , Ile- NH_2 , Leu- NH_2 , Met- NH_2 , Orn- NH_2 , Phe- NH_2 , Pro- NH_2 , Ser- NH_2 , Thr- NH_2 , Trp- NH_2 , Tyr- NH_2 , or Val- NH_2 (i.e., the peptide is an amidated hexapeptide). In one particular embodiment, X_5 is NH_2 . In other

particular embodiments, X₅ is Ala-NH₂, Arg-NH₂, Asn-NH₂, Asp-NH₂, Cys-NH₂, Glu-NH₂, Gln-NH₂, Gly-NH₂, His-NH₂, Ile-NH₂, Leu-NH₂, Met-NH₂, Orn-NH₂, Phe-NH₂, Pro-NH₂, Ser-NH₂, Thr-NH₂, Trp-NH₂, Tyr-NH₂, or Val-NH₂.

[0011] Another embodiment of the invention is directed to a peptide of Formula I, wherein X₁ is D-Asp, D-Glu, D-Lys, or D-Orn; and X₄ is Asp, Glu, Lys, or Orn.

[0012] Another embodiment of the invention is directed to a compound of Formula I, wherein X₅ is NHR and R is a (C₁ to C₁₀) alkyl.

[0013] Another embodiment of the invention is directed to a peptide of Formula I, wherein the aromatic amino acid of X₂ is Trp, Phe, or N-alkyl-Phe, and the alkyl group of N-alkyl-Phe is a (C₁ to C₆) alkyl. In one particular embodiment, X₂ is N-methyl-Phe (N-Me-Phe).

[0014] Another embodiment of the invention is directed to a peptide of Formula I, wherein the aromatic amino acid residue of either X₂ or X₃ is Phe, D-Phe, Trp, D-Trp, D-Tyr, N-alkyl-Phe, and the alkyl group of N-alkyl-Phe is (C₁ to C₁₀) alkyl or p-Y-Phe, wherein Y is NO₂, F, Cl, or Br.

[0015] Another embodiment of the invention is directed to a peptide of Formula I, wherein the aromatic amino acid of X₃ is Phe, D-Phe, or p-Y-Phe, wherein Y is NO₂, F, Cl, or Br. In one particular embodiment, X₃ is p-Cl-Phe.

[0016] Another embodiment of the invention is directed to a peptide of Formula I selected from the group consisting of:

Tyr-c[D-Lys-Trp-Phe-Glu]-NH₂ (SEQ ID NO:1);

Tyr-c[D-Glu-Phe-Phe-Lys]-NH₂ (SEQ ID NO:2);

Tyr-c[D-Lys-Trp-Phe-Glu]-Gly-NH₂ (SEQ ID NO:3);

Tyr-c[D-Glu-Phe-Phe-Lys]-Gly-NH₂ (SEQ ID NO:4);

Tyr-c[D-Lys-Trp-Phe-Asp]-NH₂(SEQ ID NO:5);

Tyr-c[D-Glu-N-Me-Phe-Phe-Lys]-NH₂ (SEQ ID NO:6); and

Tyr-c[D-Orn-Phe-p-Cl-Phe-Asp]-Val-NH₂ (SEQ ID NO:7).

[0017] Another aspect of the invention is directed to a pharmaceutical composition comprising a peptide of Formula I and a pharmaceutically acceptable carrier (e.g., a diluent or excipient).

[0018] Yet another aspect of the invention is directed to the use of a peptide of Formula I in a method of treating a patient having a condition that responds to an opioid, or a condition for which opioid treatment is standard in the art. Such a method comprises or consists of administering to the patient an effective amount of a peptide of Formula I of the invention.

Particular embodiments of this method can be followed for the purpose of providing at least one effect selected from (i) analgesia (pain relief), (ii) relief from a gastrointestinal disorder such as diarrhea, (iii) therapy for an opioid drug dependence, and (iv) treatment of any condition for which an opioid is indicated. In some embodiments the peptides of Formula I can be used to treat acute or chronic pain. Uses for the peptides of Formula I also include, but are not be limited to, use as antimigraine agents, immunomodulatory agents, immunosuppressive agents or antiarthritic agents. Certain embodiments of the methods of the present invention, such as treatment of pain or opioid drug dependence, are directed to patients having a history of opioid substance abuse. In certain embodiments of the present methods, the peptide is administered parenterally (e.g., intravenous). This invention also relates to a peptide of Formula I for use in one of said methods of treatment.

[0019] Another aspect of the invention is directed to a method of activating or regulating a mu-opioid receptor by contacting the mu-opioid receptor with a compound of the invention, as well as the use of the peptide of Formula I in such a treatment.

[0020] Another aspect of the invention is directed to a method of measuring the quantity of a mu opioid receptor in a sample using a peptide of Formula I. This method can comprise or consist of the following steps: (i) contacting a sample suspected of containing a mu opioid receptor with a peptide of Formula I to form a compound-receptor complex, (ii) detecting the complex, and (iii) quantifying the amount of complex formed.

[0021] Another aspect of the invention is directed to the use of a peptide of Formula I to perform a competitive assay method of detecting the presence of a molecule that binds to a mu opioid receptor. This method can comprise or consist of the following steps: (i) contacting a sample suspected of containing a molecule that binds to a mu opioid receptor with a mu opioid receptor and a peptide of Formula I, wherein the compound and receptor form a compound-receptor complex; (ii) measuring the amount of the complex formed in step (i); and (iii) comparing the amount of complex measured in step (ii) with the amount of a complex formed between the mu opioid receptor and the peptide in the absence of said sample.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] FIG. 1 shows Tyr-c[D-Lys-Trp-Phe-Glu]-NH₂ (SEQ ID NO:1), which is described as "Compound 1" in the following disclosure. The structural and basic molecular formulae, as well as the molecular weight (MW), are shown for Compound 1.

[0023] FIG. 2 shows Tyr-c[D-Glu-Phe-Phe-Lys]-NH₂ (SEQ ID NO:2), which is described as "Compound 2" in the following disclosure. The structural and basic molecular formulae, as well as the molecular weight (MW), are shown for Compound 2.

[0024] FIG. 3 shows Tyr-c[D-Glu-Phe-Phe-Lys]-Gly-NH₂ (SEQ ID NO:4), which is described as "Compound 4" in the following disclosure. The structural and basic molecular formulae, as well as the molecular weight (MW), are shown for Compound 4.

[0025] FIG. 4 shows opioid receptor binding activity for Compound 1. (A) mu receptor binding of "Compound 1" (triangles) or DAMGO (squares). (B) Antagonist activity of Compound 1 against binding of SNC80 to delta receptor.

[0026] FIG. 5 shows effects of compounds on antinociception and respiration. (A) Effects of Compounds 1 and 2 on antinociception as compared with morphine. **= $p < 0.01$. (B) Effects of Compounds 1 and 2 on respiratory minute volume (MV) over a 20-minute period as compared to morphine. * $p < 0.05$. *** $p < 0.001$.

[0027] FIG. 6 shows the effects of Compound 2 on antinociception and motor impairment. (A) The effects of Compound 2 (filled triangles) and morphine sulfate (MS, filled squares) on antinociception were measured by the tail flick (TF) test. Also, the effects of Compound 2 (open triangles) and morphine sulfate (open squares) on motor behavior were measured. (* = $p < 0.05$). (B) The bar graph shows the ratio of the area under the curve (AUC) for percent motor impairment relative to the AUC for percent antinociception. This ratio is significantly greater (* $p < 0.05$) for morphine than for Compound 2, consistent with greater motor impairment relative to analgesia for morphine.

[0028] FIG. 7 shows the effects of compounds on drug abuse liability. (A) The effects of Compound 1 (filled triangles), morphine (filled squares), and vehicle (filled circles) on antinociception were measured by the tail flick (TF) test. * $p < 0.05$. (B) The cumulative doses of either morphine or Compound 1 that were shown to produce maximal antinociception as shown in (A) were tested for the ability to induce conditioned place preference (CPP).

*** $p < 0.01$.

[0029] FIG. 8 shows a graph of paw pressure required to elicit a paw withdrawal response at times before and after spared nerve injury surgery for rats treated with Compound 2, compared to rats treated with morphine or no analgesic.

DETAILED DESCRIPTION OF THE INVENTION

[0030] Peptides of Formula I, which are cyclic pentapeptide and hexapeptide analogs of endomorphin-1 (Tyr-Pro-Trp-Phe-NH₂, SEQ ID NO:8) and endomorphin-2 (Tyr-Pro-Phe-Phe-NH₂, SEQ ID NO:9) were prepared. Non-limiting examples of peptides with the composition of Formula I include Compounds 1-7 below, wherein the side chains of amino acid residues 2 (X₁) and 5 (X₄) in the sequence are linked by an amide bond between the side-chains thereof. The formulac of Compounds 1-7 are shown in Table 1.

Table 1.

Compound	H-Tyr-	X₁-	X₂-	X₃-	X₄-	X₅	SEQ ID NO:
1.	Tyr-	c[D-Lys	Trp	Phe	Glu]	NH ₂	(SEQ ID NO:1)
2.	Tyr-	c[D-Glu	Phe	Phe	Lys]	NH ₂	(SEQ ID NO:2)
3.	Tyr-	c[D-Lys	Trp	Phc	Glu]	Gly-NH ₂	(SEQ ID NO:3)
4.	Tyr-	c[D-Glu	Phe	Phe	Lys]	Gly-NH ₂	(SEQ ID NO:4)
5.	Tyr-	c[D-Lys	Trp	Phe	Asp]	NH ₂	(SEQ ID NO:5)
6.	Tyr-	c[D-Glu	N-Me-Phe	Phe	Lys]	NH ₂	(SEQ ID NO:6)
7.	Tyr-	c[D-Orn	Phe	p-Cl-Phe	Asp]	Val-NH ₂	(SEQ ID NO:7)

[0031] In some embodiments, the peptides of Formula I includes peptides with an N-alkylated phenylalanine in position 3 (X₂). Alkyl groups suitable in the peptides of the present invention include (C₁ to C₁₀) alkyl groups, preferably (C₁ to C₆) alkyl groups (e.g., methyl or ethyl).

Compound 6 illustrates a cyclic analog whose linear primary amino acid sequence contains an N-methylated phenylalanine in position 3. Other peptides of this invention include compounds wherein the amino acid at position 4 (X₃) is p-Y-phenylalanine, wherein Y is NO₂, F, Cl or Br, in order to enhance receptor binding and potency. An exemplary peptide (Compound 7), whose linear primary amino acid sequence is provided in SEQ ID NO:7, has a p-chlorophenylalanine (p-Cl-Phe) in position 4.

[0032] Compounds 1 (Figure 1), 2 (Figure 2), 5 and 6 are examples of cyclic pentapeptides, and Compounds 3, 4 (Figure 3) and 7 are examples of cyclic hexapeptides of the instant invention.

[0033] For reference, the abbreviations for amino acids described herein include alanine (Ala), arginine (Arg), asparagine (Asn), aspartic acid (Asp), cysteine (Cys), glutamine (Gln), glutamic acid (Glu), glycine (Gly), histidine (His), isoleucine (Ile), leucine (Leu), lysine (Lys), methionine

(Met), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr), tryptophan (Trp), tyrosine (Tyr), valine (Val), ornithine (Orn), naphthylalanine (Nal), 2,3-diaminopropionic acid (Dpr), and 2,4-diaminobutyric acid (Dab). The L- or D-enantiomeric forms of these and other amino acids can be included in the peptides of Formula I. Other amino acids, or derivatives or unnatural forms thereof such as those listed in the 2009/2010 Aldrich Handbook of Fine Chemicals can be used in preparing compounds of the invention.

[0034] In Formula I, X₁ can be, for example, D-Asp, D-Glu, D-Lys, D-Orn, D-Dpr or D-Dab, and X₄ can be, for example, Asp, Glu, Lys, Orn, Dpr or Dab. In general, an amino acid or derivative thereof can be used as X₁ or X₄ if it contains either an amino group or a carboxyl group in its side chain. In some embodiments, the amino acid used for X₁ can be a D-enantiomeric form of such amino acid.

[0035] X₂ and X₃ in Formula I are aromatic amino acids. Examples of such amino acids are unsubstituted or substituted aromatic amino acids selected from the group consisting of phenylalanine, heteroarylalanine, naphthylalanine (Nal), homophenylalanine, histidine, tryptophan, tyrosine, arylglycine, heteroarylglycine, thyroxine, aryl-beta-alanine, and heteroaryl-beta-alanine. Examples of substituted versions of these aromatic amino acids are disclosed in U.S. Patent No. 7,629,319. As used herein, "aromatic amino acid" refers to an α -amino acid comprising an aromatic group (including aromatic hydrocarbon and aromatic heterocyclic groups) in the side-chain thereof.

[0036] In some embodiments, X₂ in Formula I can be N-alkyl-Phe, where the alkyl group comprises 1 to about 6 carbon atoms. Alternatively, the alkyl group can comprise about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 carbons, for example. The alkyl group can be a methyl (i.e., X₂ is N-Me-Phe), ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, hexyl, isohexyl, heptyl, or isoheptyl group, or any other branched form thereof, for example. By definition, the alkyl group of N-alkyl-Phe is linked to the α -amino group of phenylalanine. This alpha amino group is involved in an amide bond with the X₁ residue in certain peptides of the invention; therefore, the alpha amino group of X₂ (when N-alkyl-Phe) as it exists in such peptides is a tertiary amide.

[0037] In some embodiments X₃ in Formula I is para-Y-Phe (p-Y-Phe), where Y is NO₂, F, Cl, or Br, for example. For example, X₃ can be p-Cl-Phe. Alternatively, the NO₂, F, Cl, or Br groups can be linked in the ortho or meta positions of the phenyl ring of Phe. Any aromatic amino acid incorporated in the compounds of the invention such as at X₂ or X₃ can have the above groups linked thereto in the ortho, meta, or para positions.

Solubility.

[0038] The solubility of the peptides of Formula I (e.g., in saline or physiologic buffer) typically is enhanced relative to the prior art tetrapeptide analogs of the endomorphins. Addition of a hydrophilic amino acid and amidated C-terminus to the relatively hydrophobic tetrapeptide sequences Tyr-c-[D-Lys-Trp-Phe] (SEQ ID NO:10) and Tyr-c-[D-Lys-Phe-Phe] (SEQ ID NO:11), resulted in an unexpectedly high improvement in solubility while maintaining or improving functionality. For example, Compound 1 was soluble in water, saline and 20% PEG/saline at about 43, 21 and 90 mg/mL, respectively, compared to less than about 2 mg/mL for the previously described compounds. While increases in solubility are associated with

improved pharmaceutical delivery properties, higher solubility is also often associated with reduced functional activity (e.g., receptor binding) that may depend on lipophilicity.

Surprisingly however, as described in examples below, the functional properties of the compounds of the invention are not diminished, and indeed are generally improved.

Methods of Preparation of the Peptides of Formula I.

[0039] The peptides of Formula I can be prepared by conventional solution phase (2) or solid phase (18) methods with the use of proper protecting groups and coupling agents; see references 2 and 20. Such methods generally utilize various protecting groups on the various amino acid residues of the peptides. A suitable deprotection method is employed to remove specified or all of the protecting groups, including splitting off the resin if solid phase synthesis is applied. The peptides can be synthesized, for example, as described below.

[0040] Peptides of Formula I were synthesized on Rink Amide™ resin via Fmoc chemistry. A t-butyl group was used for Tyr, Glu, Asp side chain protection and Boc was used for Lys, Orn and Trp side chain protection. All materials were obtained from EMD Biosciences, Inc (San Diego, CA). The peptide was assembled on Rink Amide resin by repetitive removal of the Fmoc protecting group and coupling of protected amino acid. HBTU (O-benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate; CAS # 94790-37-1) and HOBT (N-hydroxybenzotriazole; CAS # 2592-95-2) were used as coupling reagents in N,N-dimethylformamide (DMF) and diisopropylethylamine (DIPEA) was used as a base. The resin was treated with an aqueous cocktail of trifluoroacetic acid and triisopropylsilane (TFA/TIS/H₂O

cocktail) for cleavage and removal of the side chain protecting groups. Crude peptide was precipitated with diethyl ether and collected by filtration.

[0041] Cyclization of the linear Fmoc-Tyr-c[X₁-X₂-X₃-X₄]-X₅ precursors: About 1 mmol of peptide was dissolved in about 1000 mL DMF and about 2 mmol DIPEA was added to the solution, followed by a solution of HBTU (about 1.1 mmol) and HOBT (about 1.1 mmol) in about 100 mL DMF. The reaction mixture was stirred at room temperature overnight. Solvent was removed *in vacuo*. The resulting solid residue was washed with 5% citric acid, saturated NaCl, saturated NaHCO₃, and water. The final solid was washed with diethyl ether and dried under high vacuum.

[0042] Preparation of Tyr-c[X₁-X₂-X₃-X₄]-X₅ peptides. The solids obtained above were dissolved in 20% piperidine/DMF. The mixture was stirred at room temperature for about 1 hour. Solvent was removed *in vacuo*. Residues were dissolved in 10% aqueous acetonitrile (MeCN/H₂O) and lyophilized.

[0043] Purification of the crude lyophilized peptides was performed with reverse phase high performance liquid chromatography (RP-HPLC). The HPLC system Gold 32 Karat™ (Beckman) consisting of the programmable solvent module 126 and the diode array detector module 168 was used in the purification and the purity control of the peptides. Reverse phase HPLC was performed using a gradient made from two solvents: (A) 0.1 % TFA in water and (B) 0.1% TFA in acetonitrile. For preparative runs, a VYDAC® 218TP510 column (250 x 10 mm; Alltech Associates, Inc.) was used with a gradient of 5-20% solvent B in solvent A over a period of 10 min, 20-25% B over a period of 30 minutes, 25-80% B over a period of 1 minute and isocratic elution over 9 minutes at a flow rate of about 4 mL/min, absorptions being measured at both 214

and 280 nm. The same gradient was used for analytical runs on a VYDAC®218TP54 column (250 x 4.6 mm) at a flow rate of about 1 mL/min.

Pharmaceutical Preparations.

[0044] The instant invention also provides pharmaceutical preparations which contain a pharmaceutically effective amount of the peptides in a pharmaceutically acceptable carrier (e.g., a diluent, complexing agent, additive, excipient, adjuvant and the like). The peptide can be present for example in a salt form, a micro-crystal form, a nano-crystal form, a co-crystal form, a nanoparticle form, a microparticle form, or an amphiphilic form. The carrier can be an organic or inorganic carrier that is suitable for external, enteral or parenteral applications. The peptides of the present invention can be compounded, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, liposomes, suppositories, intranasal sprays, solutions, emulsions, suspensions, aerosols, targeted chemical delivery systems (15), and any other form suitable for use. Non-limiting examples of carriers that can be used include water, glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea and other carriers suitable for use in manufacturing preparations, in solid, semisolid, liquid or aerosol form. In addition auxiliary, stabilizing, thickening and coloring agents and perfumes can be used.

The present invention also provides pharmaceutical compositions useful for treating pain and related conditions, as described herein. The pharmaceutical compositions comprise at least one peptide of Formula I in combination with a pharmaceutically acceptable carrier, vehicle, or diluent, such as an aqueous buffer at a physiologically acceptable pH (e.g., pH 7 to 8.5), a polymer-based nanoparticle vehicle, a liposome, and the like. The pharmaceutical compositions

can be delivered in any suitable dosage form, such as a liquid, gel, solid, cream, or paste dosage form. In one embodiment, the compositions can be adapted to give sustained release of the peptide.

[0045] In some embodiments, the pharmaceutical compositions include, but are not limited to, those forms suitable for oral, rectal, nasal, topical, (including buccal and sublingual), transdermal, vaginal, parenteral (including intramuscular, subcutaneous, and intravenous), spinal (epidural, intrathecal), and central (intracerebroventricular) administration. The compositions can, where appropriate, be conveniently provided in discrete dosage units. The pharmaceutical compositions of the invention can be prepared by any of the methods well known in the pharmaceutical arts. Some preferred modes of administration include intravenous (iv), topical, subcutaneous, oral and spinal.

[0046] Pharmaceutical formulations suitable for oral administration include capsules, cachets, or tablets, each containing a predetermined amount of one or more of the peptides, as a powder or granules. In another embodiment, the oral composition is a solution, a suspension, or an emulsion. Alternatively, the peptides can be provided as a bolus, electuary, or paste. Tablets and capsules for oral administration can contain conventional excipients such as binding agents, fillers, lubricants, disintegrants, colorants, flavoring agents, preservatives, or wetting agents. The tablets can be coated according to methods well known in the art, if desired. Oral liquid preparations include, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs. Alternatively, the compositions can be provided as a dry product for constitution with water or another suitable vehicle before use. Such liquid preparations can contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may

include edible oils), preservatives, and the like. The additives, excipients, and the like typically will be included in the compositions for oral administration within a range of concentrations suitable for their intended use or function in the composition, and which are well known in the pharmaceutical formulation art. The peptides of the present invention will be included in the compositions within a therapeutically useful and effective concentration range, as determined by routine methods that are well known in the medical and pharmaceutical arts. For example, a typical composition can include one or more of the peptides at a concentration in the range of at least about 0.01 nanomolar to about 1 molar, preferably at least about 1 nanomolar to about 100 millimolar.

[0047] Pharmaceutical compositions for parenteral, spinal, or central administration (e.g. by bolus injection or continuous infusion) or injection into amniotic fluid can be provided in unit dose form in ampoules, pre-filled syringes, small volume infusion, or in multi-dose containers, and preferably include an added preservative. The compositions for parenteral administration can be suspensions, solutions, or emulsions, and can contain excipients such as suspending agents, stabilizing agents, and dispersing agents. Alternatively, the peptides can be provided in powder form, obtained by aseptic isolation of sterile solid or by lyophilization from solution, for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use. The additives, excipients, and the like typically will be included in the compositions for parenteral administration within a range of concentrations suitable for their intended use or function in the composition, and which are well known in the pharmaceutical formulation art. The peptides of the present invention will be included in the compositions within a therapeutically useful and effective concentration range, as determined by routine methods that are well known in the

medical and pharmaceutical arts. For example, a typical composition can include one or more of the peptides at a concentration in the range of at least about 0.01 nanomolar to about 100 millimolar, preferably at least about 1 nanomolar to about 10 millimolar.

[0048] Pharmaceutical compositions for topical administration of the peptides to the epidermis (mucosal or cutaneous surfaces) can be formulated as ointments, creams, lotions, gels, or as a transdermal patch. Such transdermal patches can contain penetration enhancers such as linalool, carvacrol, thymol, citral, menthol, t-anethole, and the like. Ointments and creams can, for example, include an aqueous or oily base with the addition of suitable thickening agents, gelling agents, colorants, and the like. Lotions and creams can include an aqueous or oily base and typically also contain one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, coloring agents, and the like. Gels preferably include an aqueous carrier base and include a gelling agent such as cross-linked polyacrylic acid polymer, a derivatized polysaccharide (e.g., carboxymethyl cellulose), and the like. The additives, excipients, and the like typically will be included in the compositions for topical administration to the epidermis within a range of concentrations suitable for their intended use or function in the composition, and which are well known in the pharmaceutical formulation art. The peptides of the present invention will be included in the compositions within a therapeutically useful and effective concentration range, as determined by routine methods that are well known in the medical and pharmaceutical arts. For example, a typical composition can include one or more of the peptides at a concentration in the range of at least about 0.01 nanomolar to about 1 molar, preferably at least about 1 nanomolar to about 100 millimolar.

[0049] Pharmaceutical compositions suitable for topical administration in the mouth (e.g., buccal or sublingual administration) include lozenges comprising the peptide in a flavored base, such as sucrose, acacia, or tragacanth; pastilles comprising the peptide in an inert base such as gelatin and glycerin or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier. The pharmaceutical compositions for topical administration in the mouth can include penetration enhancing agents, if desired. The additives, excipients, and the like typically will be included in the compositions of topical oral administration within a range of concentrations suitable for their intended use or function in the composition, and which are well known in the pharmaceutical formulation art. The peptides of the present invention will be included in the compositions within a therapeutically useful and effective concentration range, as determined by routine methods that are well known in the medical and pharmaceutical arts. For example, a typical composition can include one or more of the peptides at a concentration in the range of at least about 0.01 nanomolar to about 1 molar, preferably at least about 1 nanomolar to about 100 millimolar.

[0050] A pharmaceutical composition suitable for rectal administration comprises a peptide of the present invention in combination with a solid or semisolid (e.g., cream or paste) carrier or vehicle. For example, such rectal compositions can be provided as unit dose suppositories. Suitable carriers or vehicles include cocoa butter and other materials commonly used in the art. The additives, excipients, and the like typically will be included in the compositions of rectal administration within a range of concentrations suitable for their intended use or function in the composition, and which are well known in the pharmaceutical formulation art. The peptides of the present invention will be included in the compositions within a therapeutically useful and

effective concentration range, as determined by routine methods that are well known in the medical and pharmaceutical arts. For example, a typical composition can include one or more of the peptides at a concentration in the range of at least about 0.01 nanomolar to about 1 molar, preferably at least about 1 nanomolar to about 100 millimolar.

[0051] According to one embodiment, pharmaceutical compositions of the present invention suitable for vaginal administration are provided as pessaries, tampons, creams, gels, pastes, foams, or sprays containing a peptide of the invention in combination with carriers as are known in the art. Alternatively, compositions suitable for vaginal administration can be delivered in a liquid or solid dosage form. The additives, excipients, and the like typically will be included in the compositions of vaginal administration within a range of concentrations suitable for their intended use or function in the composition, and which are well known in the pharmaceutical formulation art. The peptides of the present invention will be included in the compositions within a therapeutically useful and effective concentration range, as determined by routine methods that are well known in the medical and pharmaceutical arts. For example, a typical composition can include one or more of the peptides at a concentration in the range of at least about 0.01 nanomolar to about 1 molar, preferably at least about 1 nanomolar to about 100 millimolar.

[0052] Pharmaceutical compositions suitable for intra-nasal administration are also encompassed by the present invention. Such intra-nasal compositions comprise a peptide of the invention in a vehicle and suitable administration device to deliver a liquid spray, dispersible powder, or drops. Drops may be formulated with an aqueous or non-aqueous base also comprising one or more dispersing agents, solubilizing agents, or suspending agents. Liquid sprays are conveniently

delivered from a pressurized pack, an insufflator, a nebulizer, or other convenient means of delivering an aerosol comprising the peptide. Pressurized packs comprise a suitable propellant such as dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide, or other suitable gas as is well known in the art. Aerosol dosages can be controlled by providing a valve to deliver a metered amount of the peptide. Alternatively, pharmaceutical compositions for administration by inhalation or insufflation can be provided in the form of a dry powder composition, for example, a powder mix of the peptide and a suitable powder base such as lactose or starch. Such powder composition can be provided in unit dosage form, for example, in capsules, cartridges, gelatin packs, or blister packs, from which the powder can be administered with the aid of an inhalator or insufflator. The additives, excipients, and the like typically will be included in the compositions of intra-nasal administration within a range of concentrations suitable for their intended use or function in the composition, and which are well known in the pharmaceutical formulation art. The peptides of the present invention will be included in the compositions within a therapeutically useful and effective concentration range, as determined by routine methods that are well known in the medical and pharmaceutical arts. For example, a typical composition can include one or more of the peptides at a concentration in the range of at least about 0.01 nanomolar to about 1 molar, preferably at least about 1 nanomolar to about 100 millimolar.

[0053] Optionally, the pharmaceutical compositions of the present invention can include one or more other therapeutic agent, e.g., as a combination therapy. The additional therapeutic agent will be included in the compositions within a therapeutically useful and effective concentration range, as determined by routine methods that are well known in the medical and pharmaceutical

arts. The concentration of any particular additional therapeutic agent may be in the same range as is typical for use of that agent as a monotherapy, or the concentration may be lower than a typical monotherapy concentration if there is a synergy when combined with a peptide of the present invention.

[0054] In another aspect, the present invention provides for the use of the peptides of Formula I for treatment of pain, treatment of discomfort associated with gastrointestinal disorders, and treatment of drug dependence. Methods for providing analgesia (alleviating or reducing pain), relief from gastrointestinal disorders such as diarrhea, and therapy for drug dependence in patients, such as mammals, including humans, comprise administering to a patient suffering from one of the aforementioned conditions an effective amount of a peptide of Formula I. Diarrhea may be caused by a number of sources, such as infectious disease, cholera, or an effect or side-effect of various drugs or therapies, including those used for cancer therapy. Preferably, the peptide is administered parenterally or enterally. The dosage of the effective amount of the peptides can vary depending upon the age and condition of each individual patient to be treated. However, suitable unit dosages typically range from about 0.01 to about 100 mg. For example, a unit dose can be in the range of about 0.2 mg to about 50 mg. Such a unit dose can be administered more than once a day, e.g., two or three times a day.

[0055] All of the embodiments of the peptides of Formula I can be in the "isolated" state. For example, an "isolated" peptide is one that has been completely or partially purified. In some instances, the isolated compound will be part of a greater composition, buffer system or reagent mix. In other circumstances, the isolated peptide may be purified to homogeneity. A composition may comprise the peptide or compound at a level of at least about 50, 80, 90, or

95% (on a molar basis or weight basis) of all the other species that are also present therein.

Mixtures of the peptides of Formula I may be used in practicing methods provided by the invention.

[0056] Additional embodiments of the current invention are directed towards methods of using the peptides of Formula I disclosed herein in medicinal formulations or as therapeutic agents, for example. These methods may involve the use of a single peptide, or multiple peptides in combination (i.e., a mixture). Accordingly, certain embodiments of the invention are drawn to medicaments comprising the peptides of Formula I, and methods of manufacturing such medicaments.

[0057] As used herein, the terms "reducing," "inhibiting," "blocking," "preventing", "alleviating," or "relieving" when referring to a compound (e.g., a peptide), mean that the compound brings down the occurrence, severity, size, volume, or associated symptoms of a condition, event, or activity by at least about 7.5%, 10%, 12.5%, 15%, 17.5%, 20%, 22.5%, 25%, 27.5%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, or 100% compared to how the condition, event, or activity would normally exist without application of the compound or a composition comprising the compound. The terms "increasing," "elevating," "enhancing," "upregulating", "improving," or "activating" when referring to a compound mean that the compound increases the occurrence or activity of a condition, event, or activity by at least about 7.5%, 10%, 12.5%, 15%, 17.5%, 20%, 22.5%, 25%, 27.5%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 100%, 150%, 200%, 250%, 300%, 400%, 500%, 750%, or 1000% compared to how the condition, event, or activity would normally exist without application of the compound or a composition comprising the compound.

[0058] The following examples are included to demonstrate certain aspects of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples, which represent techniques known to function well in practicing the invention, can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific disclosed embodiments and still obtain a like or similar result without departing from the spirit and scope of the invention. The examples are provided for illustration purposes only and are not intended to be limiting.

Example 1: Binding and activation of human opioid receptors.

[0059] The peptides of Formula I showed surprisingly high affinity (subnanomolar) for the human mu opioid receptor with selective binding relative to the delta and kappa opioid receptors. The compounds were tested in standard binding assays using ³H-DAMGO (tritiated [D-Ala², N-Me-Phe⁴, Gly-ol]-enkephalin; CAS # 78123-71-4), ³H-DPDPE (CAS# 88373-73-3), and ³H-U69593 (CAS# 96744-75-1) to label mu, delta and kappa receptors, respectively, in membranes from CHO cells expressing human cloned receptors. As shown in Table 2, endomorphin-1 (EM1, SEQ ID NO:8) and endomorphin -2 (EM2, SEQ ID NO:9) are the most selective endogenous mu agonists previously reported. Analogs based on these natural opioids show greater affinity for the mu receptor, albeit with less selectivity. Tetrapeptide endomorphin analogs described earlier (U.S. Patent No. 5,885,958; ck1, Tyr-c[D-Lys-Trp-Phe] (SEQ ID NO:10); ck2, Tyr-c[D-Lys-Phe-Phe] (SEQ ID NO:11)) showed the highest affinity of the compounds tested. Peptides of Formula I, which include a hydrophilic amino acid and amidated

carboxy-terminus (Compounds 1, 2, 5) retained high affinity binding, but increased selectivity for the mu receptor.

Table 2: Compound binding to opioid receptors.

	K _i (nM)		Kappa	Selectivity	
	Mu	Delta		Delta/Mu	Kappa/Mu
Morphine	0.92	242	56	264	61
DAMGO	0.78	589	334	754	429
EM1	2.07	1215	>10000	587	>5000
EM2	1.32	5704	>10000	4328	>5000
ck1	0.32	28	35	90	111
ck2	0.36	3	12	9	33
Compound 1	0.49	132	128	267	260
Compound 2	0.73	69	71	94	98
Compound 5	0.43	140	29	328	67

[0060] Receptor activation: GTPγS functional assay. Functional activation of the three opioid receptors was tested in standard assays in which the non-hydrolysable GTP analog, ³⁵S-GTPγS, was used to quantify activation of cloned human opioid receptors expressed in cell membranes. Figure 4A shows that Compound 1 is a full efficacy agonist with significantly greater potency than the reference compound, DAMGO. Figure 4B shows that Compound 1 exhibits unexpected full efficacy as a delta antagonist; i.e., it is able to inhibit the delta activation produced by an ED₈₀ dose of the reference delta agonist, SNC80 (CAS # 156727-74-1). Table 3 shows that all agonists tested are potent activators of the mu receptor, with EC₅₀ (median effective concentration) values at low-nanomolar to sub-nanomolar concentrations. All compounds were found to be full efficacy (>90%) agonists at the mu receptor. The endomorphins and the compounds of Formula I of the invention show remarkable selectivity for

receptor activation, with delta activation below 50% at concentrations up to 10 μ M, reflecting selectivity >100000. Compounds 1 and 3, however, showed full efficacy delta antagonism; Compound 1 exhibited this antagonism at a relatively low concentration.

Table 3: Opioid receptor activation by compounds.

	Agonist EC ₅₀ (nM)			Selectivity		Delta Antagonist	
	mu	delta	kappa	delta/mu	kappa/mu	IC 50	efficacy
MS ^a	3.90	1245	2404	319	616		
DAMGO	1.98	3641	13094	1839	6613		
ck1	0.21	138	469.51	658	2236		
ck2	0.15	7	206.11	44	1374		
EM1	1.82	>100000	>100000	>50000	>50000	4287	100
EM2	8.44	>100000	>100000	>10000	>10000	30000	88
Comp. 1	0.15	>100000	963.79	>500000	6425	105	93
Comp. 2	0.99	>100000	12114.00	>100000	12236	2750	51
Comp. 5	0.22	>100000	740.34	>400000	3365	557	100

^amorphine sulfate

[0061] Receptor activation: Beta-arrestin recruitment. Beta-arrestin is an intracellular protein that is recruited to the mu opioid receptor following activation by agonists. It has been shown to activate intracellular signaling pathways that in many cases are independent of well-known G-protein mediated pathways. It has recently been shown that beta-arrestin knockout mice exhibit altered responses to morphine, including increased analgesia and decreased side effects such as tolerance, respiratory depression, and constipation (16). These results indicate that the analgesic and side-effects of morphine are separable by manipulation of cell signaling processes. These findings also provide support for the recent concept known variously as “functional selectivity”, “biased agonism”, “agonist directed signaling” and other descriptions.

According to this concept, agonists capable of producing a different cascade of signaling at a given receptor could produce a different profile of desired and undesired effects relative to other agonists for that receptor. Three of the analogs of this invention were tested and showed patterns of beta-arrestin recruitment (ranging from high potency with low efficacy to moderate potency with significant efficacy) that were different from each other and from morphine. Together with the differential analgesic/side-effect profiles relative to morphine described in previous examples, the beta arrestin results suggest that these compounds exhibit “functional selectivity”, favoring analgesia over adverse side-effects.

[0062] Beyond the value of high mu agonist selectivity (i.e., exclusion of potential side-effects resulting from activation of multiple receptors), delta antagonism is expected to attenuate opioid-induced tolerance, dependence, and reward. As first shown in 1991 (1) and supported in numerous studies since, delta antagonists can reduce morphine-induced tolerance and dependence, while maintaining or enhancing analgesia. Recent studies (11) have also shown reduced rewarding properties of mu agonist/delta antagonists as reflected in the conditioned place preference (CPP) test described below. The activity of the peptides of Formula I (e.g., Compound 1) as mu agonists/delta antagonists as well as at mu/delta receptor dimers indicate that the peptides will produce effective analgesia with reduced tolerance, dependence, and reward (18).

Example 2: Providing analgesia of greater duration, but with reduced respiratory depression, relative to morphine after intravenous administration.

[0063] Respiratory depression is a major safety issue in the use of opioids. An opioid providing analgesia as effective as that produced by morphine, but with less respiratory depression, would be a major advance for the safe use of opioid analgesics. Effectiveness after systemic administration, such as intravenous (i.v.) injection, is unusual for peptide-based compounds, and would be critical for the clinical utility thereof. Two peptides (Compounds 1 and 2) were tested for their effects on respiration (minute ventilation) and duration of antinociception relative to morphine. Rats with indwelling jugular catheters were placed in a BUXCO™ whole body plethysmograph apparatus for determining multiple respiratory parameters. For 20 minutes following i.v. injection of vehicle (saline), baseline minute ventilation was determined. Animals were then injected with morphine or test compound and changes from baseline were determined for 20 minutes, the period of maximal inhibition of minute ventilation by all compounds. The standard tail-flick (TF) test was used to determine antinociception. A baseline test was conducted before placing the animal in the BUXCO™ chamber, at the end of the 20-minute respiratory test, and at every 20 minutes thereafter until the TF latency returned to below 2x baseline TF. Baseline latencies were 3-4 seconds and a cut-off time ("maximal antinociception") was set at 9 seconds to avoid tissue damage.

[0064] Figure 5A shows that 10 mg/kg doses of Compounds 1 and 2 produced significantly longer antinociception than all other treatments (**= $p < 0.01$) and 5.6 mg/kg doses produced antinociception similar to the 10 mg dose of morphine. Despite the greater antinociceptive effect of Compounds 1 and 2, significantly (* $p < 0.05$) less inhibition of respiration was observed in

both doses of Compound 1 and in the 5.6 mg/kg dose of Compound 2 (Figure 5B). These results indicate an unexpected and clearly safer therapeutic profile for the peptides of Formula I over the current standard opioid analgesic.

Example 3: Providing analgesia of greater duration than morphine with reduced impairment of neuromotor coordination and cognitive function.

[0065] Neuromotor and cognitive impairment are characteristics of opioids that are of particular importance in two populations, i.e., military combat troops, where escape from immediate danger can require unimpaired motor and cognitive skills, and the elderly, where these impairments can exacerbate compromised function including impaired balance, which can lead to increased risk of fractures.

Example 3a: Neuromotor coordination.

[0066] Figure 6A illustrates that Compound 2 produces significantly greater antinociception, but significantly reduced motor impairment, relative to morphine (MS). Both compounds were administered by cumulative intravenous (i.v.) doses in rats. Increasing quarter-log doses were given every 20 minutes, and a tail flick (TF) test (a test of latency to remove the tail from a hot light beam) followed by a rotorod test were conducted about 15 minutes after each injection. Escalating doses were given until each animal showed greater than 90% maximum possible effect (%MPE) on the TF test, determined as: $[(\text{latency to TF minus baseline latency}) / (9 \text{ sec maximum (cut off) time to avoid tissue damage) minus baseline}] \times 100$. The animal was then placed on a rod that rotated at speeds escalating to 13 revolutions per minute (RPM) over 3 minutes, and the latency to fall from the rod was determined. Only animals that consistently remained on the rod for the full 180 seconds during training in the drug-naïve state were tested.

%Maximum Possible Inhibition (%MPI) of motor coordination was determined as $100 - (\text{latency to fall}/180 \times 100)$.

[0067] The two compounds showed similar onset to maximal antinociception, but Compound 2 produced significantly longer antinociception, as reflected by TF latencies significantly ($* = p < 0.05$) longer than those of the morphine group at 135 and 155 minutes (Figure 6A). Despite this greater antinociception, the motor impairment was significantly less than that of morphine (Figure 6B, $* p < 0.05$). The impairment of motor behavior by morphine was significantly above that of vehicle controls ($* p < 0.05$) while that of Compound 2 was not.

Example 3b: Cognitive impairment.

[0068] A widely used standard test of cognitive function is the Morris Water Maze (MWM). During training, rats learn to find a hidden escape platform based on spatial memory. Average latency to the platform, as well as average distance from the platform (a measure unaffected by swim speed), decrease as the task is acquired and provide indices of spatial memory. After 4 days of training, an injection of morphine produced impairment of spatial memory, as reflected by a significant increase in the latency to, and average distance from, the platform. By contrast, Compound 2, at doses that provide equal or greater antinociception than morphine, did not produce significant impairment. These results indicate an unexpected and superior therapeutic profile of the peptides of Formula 1 with regard to cognitive function relative to the current standard opioid analgesic.

Example 4: Providing analgesia of greater duration, but reduced reward, relative to morphine.

[0069] Opioids remain the standard treatment for relief of severe pain, but diversion of pain medications for non-pain use has become a serious national problem (see U.S. Department of Health and Human Services Substance Abuse and Mental Health Services Administration, found at world wide website oas.samhsa.gov/2k9/painRelievers/nonmedicalTrends.pdf). Considerable efforts in academia and industry have focused on "tamper-proof" versions of opioid medications, but there has been little success in developing opioids that provide highly effective analgesia with minimal abuse potential. The conditioned place preference (CPP) paradigm is a widely accepted model for demonstrating rewarding properties of drugs, and all major classes of abused drugs produce CPP, including opioids such as morphine and heroin. Briefly, animals are first allowed, on Day 1, to freely explore a 3-compartment apparatus consisting of a small "start box" and two larger compartments that are perceptually distinct (gray vs. black and white stripes in this example). For the next three days, the animals are given an i.v. injection of drug and confined to one compartment, and vehicle is given in the other. The time at which the drug or vehicle is given (a.m. or p.m.) is counterbalanced, as is the compartment in which the drug is given (preferred or non-preferred, as determined during the baseline test). This unbiased design allows for detection of both drug preference and drug aversion. After three days of conditioning (Days 2, 3 and 4), the animal is allowed free access to all compartments on Day 5 in the drug-free state and the change in absolute time and proportion of time spent in the drug-paired compartment are determined. A significant increase in the time or proportion of time spent in the drug-paired compartment on the post-conditioning test day relative to that on the pre-

conditioning baseline test is interpreted as a conditioned place preference, reflective of rewarding properties and potential abuse liability.

[0070] When the cumulative doses of either morphine or Compound 1 that were shown to produce maximal antinociception (Figure 7A) were tested for the ability to induce CPP (Figure 7B), morphine produced a significant (***) $p < 0.01$ increase in the time spent on the drug side, while Compound 1 did not, even though significantly (* $p < 0.05$) greater antinociception (Figure 7A) was observed with Compound 1 from 140-180 minutes after its injection. These findings are consistent with less abuse liability for the novel analogs relative to morphine.

Example 5: Alleviation of Chronic Pain.

[0071] Chronic pain affects a large proportion of the population. One form of chronic pain, neuropathic pain, is particularly difficult to treat. Figure 8 shows that Compound 2 provides unexpectedly potent relief of neuropathic pain induced by the spared nerve injury (SNI) model in the rat. Prior to SNI surgery (“pre-surgery”), an average pressure of about 177 g applied to the hindpaw with a Randall-Selitto device was required to elicit a paw withdrawal response. About 7-10 days post-surgery, the animals showed hyperalgesia, indicated by a reduction in the average pressure (to about 70 g) required to elicit withdrawal. Morphine and Compound 2 were administered by intrathecal delivery in doses chosen to produce full alleviation of the hyperalgesia. Cumulative dosing showed that the ED_{50} for the return to the pre-surgical baseline response was about 1.4 μg for morphine and about 0.018 μg for Compound 2. On a molar basis, Compound 2 was 160-fold more potent than morphine against neuropathic pain. Similar results were observed after other forms of chronic pain including post-incisional (post-operative) and inflammatory pain induced by Complete Freund’s Adjuvant (CFA). The foregoing examples are

illustrative, but not exhaustive, as to the types of acute or chronic pain for which the peptides of Formula I are effective.

[0072] Chronic administration of morphine has been shown in several studies to produce an inflammatory response in the spinal cord, reflected by increased activation of glial cells (astroglia and microglia) and “opioid induced hyperalgesia”. When morphine and Compound 1, at doses producing similar antinociception after initial dosing, were administered twice/day for three days, morphine produced greater glial activation than did compound 1. These results suggest that compounds of Formula 1 will induce fewer side-effects associated with inflammatory responses compared to morphine.

[0073] [This paragraph intentionally left blank.]

[0074] Preferred embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Variations of those preferred embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations

thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

REFERENCES

[0075] The following references are referred to in this application:

- (1) Abdelhamid E. E., Sultana M., Portoghese P. S. and Takemori A. E. (1991) Selective blockage of *delta* opioid receptors prevents the development of morphine tolerance and dependence in mice. *J. Pharmacol. Exp. Ther.* **258**, 299-303;
- (2) Bodanszky M. (1993) *Peptide Chemistry: A Practical Textbook*. Springer-Verlag, New York;
- (3) Chen Y., Mestek A., Liu J., Hurley J. A. and Yu L. (1993) Molecular cloning and functional expression of a μ -opioid receptor from rat brain. *Mol. Pharmacol.* **44**, 8-12;
- (4) Czaplá M. A., Gozal D., Alea O. A., Beckerman R. C. and Zadina J. E. (2000) Differential cardiorespiratory effects of endomorphin 1, endomorphin 2, DAMGO, and morphine. *Am. J. Respir. Crit Care Med* **162**, 994-999;
- (5) Czaplá M. A. and Zadina J. E. (2005) Reduced suppression of CO₂-induced ventilatory stimulation by endomorphins relative to morphine. *Brain Res.* **1059**, 159-166;
- (6) Evans C. J., Keith D. E., Jr., Morrison H., Magendzo K. and Edwards R. H. (1992) Cloning of a delta opioid receptor by functional expression. *Science* **258**, 1952-1955;
- (7) Gianni W., Ceci M., Bustacchini S., Corsonello A., Abbatecola A. M., Brancati A. M., Assisi A., Scuteri A., Cipriani L. and Lattanzio F. (2009) Opioids for the treatment of chronic non-cancer pain in older people. *Drugs Aging* **26 Suppl 1**, 63-73;
- (8) Kieffer B. L. (1999) Opioids: first lessons from knockout mice. *Trends Pharmacol. Sci* **20**, 19-26;
- (9) Kieffer B. L., Befort K., Gaveriaux-Ruff C. and Hirth C. G. (1992) The δ -opioid receptor: isolation of a cDNA by expression cloning and pharmacological characterization. *Proc. Natl. Acad. Sci U. S A* **89**, 12048-12052;

- (10) Kuehn B. M. (2009) New pain guideline for older patients: avoid NSAIDs, consider opioids. *JAMA* **302**, 19;
- (11) Lenard N. R., Daniels D. J., Portoghese P. S. and Roerig S. C. (2007) Absence of conditioned place preference or reinstatement with bivalent ligands containing mu-opioid receptor agonist and delta-opioid receptor antagonist pharmacophores. *Eur. J. Pharmacol.* **566**, 75-82;
- (12) Meng F., Xie G. X., Thompson R. C., Mansour A., Goldstein A., Watson S. J. and Akil H. (1993) Cloning and pharmacological characterization of a rat κ opioid receptor. *Proc. Natl. Acad. Sci U. S A* **90**, 9954-9958;
- (13) Minami M., Toya T., Katao Y., Maekawa K., Nakamura S., Onogi T., Kaneko S. and Satoh M. (1993) Cloning and expression of a cDNA for the rat κ -opioid receptor. *FEBS Lett.* **329**, 291-295;
- (14) Nishi M., Takeshima H., Fukuda K., Kato S. and Mori K. (1993) cDNA cloning and pharmacological characterization of an opioid receptor with high affinities for κ -subtype-selective ligands. *FEBS Lett.* **330**, 77-80;
- (15) Prokai-Tatrai K., Prokai L. and Bodor N. (1996) Brain-targeted delivery of a leucine-enkephalin analogue by retrometabolic design. *J. Med Chem.* **39**, 4775-4782;
- (16) Rachal, KM, JKL Walker, and LM Bohn (2005) Morphine Side Effects in β -Arrestin 2 Knockout Mice. *J. Pharmacol. Exp. Ther.* **314**, 1195-1201
- (17) Rozenfeld R. and Devi L. A. (2010) Receptor heteromerization and drug discovery. *Trends Pharmacol. Sci* **31**, 124-130;
- (18) Stewart J. M. and Young J. D. (1984) *Solid Phase Peptide Synthesis*. Pierce Chemical Company;
- (19) Thompson R. C., Mansour A., Akil H. and Watson S. J. (1993) Cloning and pharmacological characterization of a rat μ opioid receptor. *Neuron* **11**, 903-913;
- (20) Wang J. B., Johnson P. S., Persico A. M., Hawkins A. L., Griffin C. A. and Uhl G. R. (1994) Human μ opiate receptor. cDNA and genomic clones, pharmacologic characterization and chromosomal assignment. *FEBS Lett.* **338**, 217-222;

(21) Wilson A. M., Soignier R. D., Zadina J. E., Kastin A. J., Nores W. L., Olson R. D. and Olson G. A. (2000) Dissociation of analgesic and rewarding effects of endomorphin-1 in rats. *Peptides* **21**, 1871-1874; and

(22) Zadina J. E., Hackler L., Ge L. J. and Kastin A. J. (1997) A potent and selective endogenous agonist for the μ -opiate receptor. *Nature* **386**, 499-502.

What is claimed is:

1. A cyclic peptide of Formula I:



wherein

X_1 is an acidic or basic D-amino acid; and X_4 is an acidic or basic amino acid;

X_2 is Trp;

X_3 is an aromatic amino acid;

X_5 is NHR, Ala-NHR, Arg-NHR, Asn-NHR, Asp-NHR, Cys-NHR, Glu-NHR, Gln-NHR, Gly-NHR, His-NHR, Ile-NHR, Leu-NHR, Met-NHR, Orn-NHR, Phe-NHR, Pro-NHR, Ser-NHR, Thr-NHR, Trp-NHR, Tyr-NHR, or Val-NHR, wherein R is H or an alkyl group; and there is an amide bond between an amino group and a carboxylic acid group on side chains of amino acids X_1 and X_4 ;

with the proviso that when X_1 is an acidic D-amino acid, then X_4 is a basic amino acid; and when X_1 is a basic D-amino acid, then X_4 is an acidic amino acid.

2. The peptide of claim 1, wherein:

(i) X_1 is selected from the group consisting of D-Lys and D-Orn; and X_4 is selected from the group consisting of D-Asp, D-Glu, Asp, and Glu; or

(ii) X_1 is selected from the group consisting of D-Asp and D-Glu; and X_4 is selected from the group consisting of D-Lys, D-Orn, Lys, and Orn.

3. The peptide of claim 1, wherein X_3 is p-Cl-Phe.

4. The peptide of claim 1, wherein R is H and X_5 is Ala-NH₂, Arg-NH₂, Asn-NH₂, Asp-NH₂, Cys-NH₂, Glu-NH₂, Gln-NH₂, Gly-NH₂, His-NH₂, Ile-NH₂, Leu-NH₂, Met-NH₂, Orn-NH₂, Phe-NH₂, Pro-NH₂, Ser-NH₂, Thr-NH₂, Trp-NH₂, Tyr-NH₂, or Val-NH₂.

5. The peptide of claim 1, wherein the alkyl group is a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, hexyl, isohexyl, heptyl, or isoheptyl group.

6. The peptide of claim 1, selected from the group consisting of:
Tyr-c[D-Lys-Trp-Phe-Glu]-NH₂ (SEQ ID NO:1),
Tyr-c[D-Lys-Trp-Phe-Glu]-Gly-NH₂ (SEQ ID NO:3), and
Tyr-c[D-Lys-Trp-Phe-Asp]-NH₂ (SEQ ID NO:5).
7. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a peptide according to any one of claims 1 to 6.
8. A peptide according to any one of claims 1 to 6 for use in a pain medication.
9. A peptide of any one of claims 1 to 6 for use in treating pain.
10. The use of a peptide of any one of claims 1 to 6 in the manufacture of a medicament for treating pain.
11. An *in vitro* method of activating a mu-opioid receptor, wherein the method comprises contacting the mu-opioid receptor with a peptide according to any one of claims 1 to 6.
12. A method for measuring the quantity of a mu opioid receptor in a sample, comprising:
(i) contacting a sample suspected of containing a mu opioid receptor with a peptide to form a compound-receptor complex, wherein the peptide is a peptide of any one of claims 1 to 6;
(ii) detecting the complex formed in step (i); and
(iii) quantifying the amount of complex detected in step (ii).
13. A competitive assay method for detecting the presence of a molecule that binds to a mu opioid receptor comprising:

(i) contacting a sample suspected of containing a molecule that binds to a mu opioid receptor with a mu opioid receptor and a peptide according to any one of claims 1 to 6, wherein the peptide and receptor form a compound-receptor complex;

(ii) measuring the amount of the complex formed in step (i); and

(iii) comparing the amount of complex measured in step (ii) with the amount of a complex formed between the mu opioid receptor and the peptide in the absence of the sample.

14. A peptide according to any one of claims 1 to 6 for use in treating pain from a gastrointestinal disorder.

15. A cyclic peptide wherein the peptide consists of the amino acid sequence of Tyr-c[D-Lys-Trp-Phe-Glu]-NH₂ (SEQ ID NO: 1) or wherein the peptide consists of the amino acid sequence of Tyr-c[D-Lys-Trp-Phe-Asp]-NH₂ (SEQ ID NO:5).

16. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and the peptide of claim 15.

17. Use of the peptide of claim 15 for treating pain.

18. A cyclic peptide of Formula I:

(I) H-Tyr-c[X₁-X₂-X₃-X₄]-X₅, wherein

X₁ is an acidic D-amino acid or a basic D-amino acid; and X₄ is an acidic or basic amino acid;

X₂ is Trp, Phe, or N-methyl-Phe;

X₃ is Phe or p-Cl-Phe;

X₅ is Ala-NHR, Gly-NHR, Ile-NHR, Leu-NHR, or Val-NHR, wherein R is H or an alkyl group; and there is an amide bond between an amino group and a carboxylic acid group on side chains of amino acids X₁ and X₄;

with the proviso that when X₁ is an acidic D-amino acid, then X₄ is a basic amino acid;
and

when X₁ is a basic D-amino acid, then X₄ is an acidic amino acid.

19. The peptide of claim 18, wherein:
 - (i) X₁ is selected from the group consisting of D-Lys and D-Orn; and X₄ is selected from the group consisting of D-Asp, D-Glu, Asp, and Glu; or
 - (ii) X₁ is selected from the group consisting of D-Asp and D-Glu; and X₄ is selected from the group consisting of D-Lys, D-Orn, Lys, and Orn.
20. The peptide of claim 18 or 19, wherein X₂ is Trp or Phe.
21. The peptide of claim 18 or 19, wherein X₂ is N-methyl-Phe.
22. The peptide of any one of claims 18 to 21, wherein X₃ is Phe.
23. The peptide of any one of claims 18 to 22, wherein R is H and X₅ is Ala-NH₂, Gly-NH₂, Ile-NHR, Leu-NH₂, or Val-NH₂.
24. The peptide of any one of claims 18 to 22, wherein the alkyl group is a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, hexyl, isohexyl, heptyl, or isoheptyl group.
25. A peptide having the formula:
Tyr-c[D-Lys-Trp-Phe-Glu]-Gly-NH₂ (SEQ ID NO:3).
26. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a peptide according to any one of claims 18 to 25.
27. A method for measuring the quantity of a mu opioid receptor in a sample, comprising:

- (i) contacting a sample suspected of containing a mu opioid receptor with a peptide to form a compound-receptor complex, wherein the peptide is a peptide of any one of claims 18 to 25;
- (ii) detecting the complex formed in step (i); and
- (iii) quantifying the amount of complex detected in step (ii).

28. A competitive assay method for detecting the presence of a molecule that binds to a mu opioid receptor comprising:

- (i) contacting a sample suspected of containing a molecule that binds to a mu opioid receptor with a mu opioid receptor and a peptide according to any one of claims 18 to 25, wherein the peptide and receptor form a compound-receptor complex;
- (ii) measuring the amount of the complex formed in step (i); and
- (iii) comparing the amount of complex measured in step (ii) with the amount of a complex formed between the mu opioid receptor and the peptide in the absence of the sample.

29. A pharmaceutical composition for treating pain comprising a pharmaceutically acceptable carrier and the peptide of any one of claims 18 to 25.

30. A pharmaceutical composition for treating pain from a gastrointestinal disorder comprising a pharmaceutically acceptable carrier and the peptide of any one of claims 18 to 25.

31. Use of a peptide of any one of claims 18 to 25 for the preparation of a medicament for treating pain.

32. Use of a peptide of any one of claims 18 to 25 for the preparation of a medicament for treating pain from a gastrointestinal disorder.

33. A cyclic peptide of Formula I:

(I) H-Tyr-cyclo[X₁-X₂-X₃-X₄]-X₅,

wherein

X₁ is a basic D-amino acid and X₄ is an acidic amino acid;
X₂ is Trp and X₃ is an aromatic amino acid; and
X₅ is NHR, wherein R is H or an alkyl group; and there is an amide bond between an amino group and a carboxylic acid group on side chains of amino acids X₁ and X₄.

34. The peptide of claim 33, wherein:
X₁ is selected from the group consisting of D-Lys and D-Orn and X₄ is selected from the group consisting of D-Asp, D-Glu, Asp, and Glu.

35. The peptide of claim 33, wherein:
X₃ is selected from the group consisting of Phe, D-Phe, and p-Y-Phe, wherein Y is NO₂, F, Cl, or Br.

36. The peptide of claim 33, wherein X₃ is p-Cl-Phe.

37. The peptide of claim 33, wherein R is H.

38. The peptide of claim 33, wherein R is an alkyl group selected from methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, hexyl, isohexyl, heptyl, and isoheptyl group.

39. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and the peptide of claim 33.

40. The peptide of claim 33, wherein X₁ is D-Lys.

41. The peptide of claim 33, wherein X₄ is Asp or Glu.

42. A cyclic peptide of Formula 1:

(I) H-Tyr-c[X₁-X₂-X₃-X₄]-X₅, wherein

X₁ and X₄ each independently is an acidic amino acid or a basic amino acid;

X₂ is Phe;

X₃ is an aromatic amino acid;

X₅ is Ala-NHR, Arg-NHR, Asn-NHR, Asp-NHR, Cys-NHR, Glu-NHR, Gln-NHR, Gly-NHR, His-NHR, Ile-NHR, Leu-NHR, Met-NHR, Orn-NHR, Phe-NHR, Pro-NHR, Ser-NHR, Thr-NHR, Trp-NHR, Tyr-NHR, or Val-NHR, wherein R is H or an alkyl group; and there is an amide bond between an amino group and a carboxylic acid group on side chains of amino acids X₁ and X₄;

with the proviso that when X₁ is an acidic amino acid, then X₄ is a basic amino acid; and when X₁ is a basic amino acid, then X₄ is an acidic amino acid.

43. The peptide of claim 42, wherein:

(i) X₁ is selected from the group consisting of D-Lys, D-Orn, Lys, and Orn; and X₄ is selected from the group consisting of D-Asp, D-Glu, Asp, and Glu; or

(ii) X₁ is selected from the group consisting of D-Asp, D-Glu, Asp, and Glu; and X₄ is selected from the group consisting of D-Lys, D-Orn, Lys, and Orn.

44. The peptide of claim 42, wherein X₃ is p-Cl-Phe.

45. The peptide of claim 42, wherein X₁ is a D-amino acid.

46. The peptide of claim 42, wherein R is H and X₅ is Ala-NH₂, Arg-NH₂, Asn-NH₂, Asp-NH₂, Cys-NH₂, Glu-NH₂, Gln-NH₂, Gly-NH₂, His-NH₂, Ile-NH₂, Leu-NH₂, Met-NH₂, Orn-NH₂, Phe-NH₂, Pro-NH₂, Ser-NH₂, Thr-NH₂, Trp-NH₂, Tyr-NH₂ or Val-NH₂.

47. The peptide of claim 42, wherein the alkyl group is a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, hexyl, isohexyl, heptyl, or isoheptyl group.

48. The peptide of claim 42, selected from the group consisting of:

Tyr-c[D-Glu-Phe-Phe-Lys]-Gly-NH₂ (SEQ ID NO:4), and

Tyr-c[D-Orn-Phe-p-Cl-Phe-Asp]-Val-NH₂ (SEQ ID NO:7).

49. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a peptide according to any one of claims 42 to 48.

50. A peptide according to any one of claims 42 to 48 for use in treating pain.

51. An *in vitro* method of activating a mu opioid receptor, wherein the method comprises contacting the mu opioid receptor with a peptide according to any one of claims 44 to 48.

52. A method for measuring the quantity of a mu opioid receptor in a sample, comprising:

(i) contacting a sample suspected of containing a mu opioid receptor with a peptide to form a compound-receptor complex, wherein the peptide is a peptide of any one of claims 42 to 48;

(ii) detecting the complex formed in step (i); and

(iii) quantifying the amount of complex detected in step (ii).

53. A competitive assay method for detecting the presence of a molecule that binds to a mu opioid receptor comprising:

(i) contacting a sample suspected of containing a molecule that binds to a mu opioid receptor with a mu opioid receptor and a peptide of any one of claims 42 to 48, wherein the peptide and receptor form a compound-receptor complex;

(ii) measuring the amount of the complex formed in step (i); and

(iii) comparing the amount of complex measured in step (ii) with the amount of a complex formed between the mu opioid receptor and the peptide in the absence of the sample.

54. A peptide of any one of claims 42 to 48 for use in treating pain from a gastrointestinal disorder.

55. A cyclic peptide of Formula I:

(I) H-Tyr-c[X₁-X₂-X₃-X₄]-X₅,

X₁ is an acidic D-amino acid or a basic D-amino acid; and X₄ is an acidic or basic amino acid;

X₂ is Trp;

X₃ is Phe or p-Cl-Phe; and

X₅ is NHR, wherein R is H or an alkyl group; and there is an amide bond between an amino group and a carboxylic acid group on side chains of amino acids X₁ and X₄;

with the proviso that when X₁ is an acidic D-amino acid, then X₄ is a basic amino acid; and when X₁ is a basic D-amino acid, then X₄ is an acidic amino acid.

56. The peptide of claim 55, wherein:

(i) X₁ is selected from the group consisting of D-Lys and D-Orn; and X₄ is selected from the group consisting of D-Asp, D-Glu, Asp, and Glu; or (ii) X₁ is selected from the group consisting of D-Asp and D-Glu; and X₄ is selected from the group consisting of D-Lys, D-Orn, Lys, and Orn.

57. The peptide of claim 55, wherein X₃ is D-Phe or p-Cl-Phe.

58. The peptide of claim 57, wherein X₃ is p-Cl-Phe.


59. The peptide of claim 55, wherein the alkyl group is a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, hexyl, isohexyl, heptyl, or isoheptyl group.

60. A pharmaceutical composition for treating pain comprising a pharmaceutically acceptable carrier and a peptide of any one of claims 55 to 59.

61. A pharmaceutical composition for treating pain from a gastrointestinal disorder comprising a pharmaceutically acceptable carrier and a peptide of any one of claims 55 to 59.


62. Use of a peptide of any one of claims 55 to 59 for the preparation of a medicament for treating pain.

63. Use of a peptide of any one of claims 55 to 59 for the preparation of a medicament for treating pain from a gastrointestinal disorder.

1/6**Tyr-c[D-Lys-Trp-Phe-Glu]-NH₂**
(SEQ ID NO:1)**C₄₀H₄₈N₈O₇****MW=752.3646****FIG. 1**

2/6

Tyr-c[D-Glu-Phe-Phe-Lys]-NH₂
(SEQ ID NO:2)




C₃₈H₄₇N₇O₇

MW=713.3535

FIG. 2

3/6

Tyr-c[D-Glu-Phe-Phe-Lys]-Gly-NH₂
(SEQ ID NO:4)



C₄₀H₅₀N₈O₈

MW=770.3752

FIG. 3

4/6




FIG. 4




FIG. 5

5/6




FIG. 6

6/6




FIG. 7

Reversal of Neuropathic Pain





FIG. 8

Tyr-c[D-Lys-Trp-Phe-Glu]-NH₂
(SEQ ID NO:1)



C₄₀H₄₉N₉O₇

MW=752.3646