



(22) Date de dépôt/Filing Date: 1993/08/20

(41) Mise à la disp. pub./Open to Public Insp.: 1995/02/21

(45) Date de délivrance/Issue Date: 2004/04/06

(51) Cl.Int.⁵/Int.Cl.⁵ C07D 409/06, A61K 31/41

(72) Inventeurs/Inventors:

SEO, AKIRA, JP;
HIRAGA, KUNIKAZU, JP;
KOGA, HIROYASU, JP;
NIWANO, YOSHIMI, JP


(73) Propriétaire/Owner:

NIHON NOHYAKU CO., LTD., JP

(74) Agent: RICHES, MCKENZIE & HERBERT LLP

(54) Titre : COMPOSITION ACCELERANT LA CICATRISATION

(54) Title: COMPOSITION FOR ACCELERATING HEALING OF WOUND



(I)


(57) Abrégé/Abstract:

A composition for accelerating healing of a wound is disclosed, comprising as an active ingredient a compound represented by formula (I): (see formula I) wherein R represents a hydrogen atom, a phenyl group or a phenyl group substituted with a halogen atom, a C₁-C₆ alkyl group, a hydroxyl group or a C₁-C₆ alkoxy group; and X represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group.



ABSTRACT OF THE DISCLOSURE

A composition for accelerating healing of a wound is disclosed, comprising as an active ingredient a compound represented by formula (I):



wherein R represents a hydrogen atom, a phenyl group or a phenyl group substituted with a halogen atom, a C₁-C₆ alkyl group, a hydroxyl group or a C₁-C₆ alkoxy group; and X represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group.

COMPOSITION FOR ACCELERATING HEALING OF WOUND

FIELD OF THE INVENTION

This invention relates to a composition for accelerating healing of a wound.

BACKGROUND OF THE INVENTION

A wound is a kind of injuries to the body and includes incisions, wounds of gastrointestinal tracts, gastric ulcers, burns, detachment, lacerations, incision, pressure ulcers called decubitus or bed sore, erosion, and lesions of surface tissues due to infection. In particular, it is desirable to accelerate healing of a surgical wound by a positive and direct treatment without relying on spontaneous cure because a patient having undergone a surgical operation generally has considerably reduced physical strength. Further, bed sore not only causes pain to a patient but entails a heavy expenditure for healing, and is expected to give rise to a great social problem with the recent increase in aged population. Healing of these wounds generally depends on synthesis of connective tissues and epithelial tissues by cell proliferation. A drug which stimulates or accelerates the differentiation and proliferation of cells at the lesion is believed valuable in healing of a wound.

Known drugs for accelerating healing of a wound mainly include extracts of naturally occurring substances, e.g., aloe, antibiotics, antiphlogistics, kallikrein, adenine, nicotinic acid, allantoin, vitamin A, zinc, c-AMP derivatives (see JP-A-


63-107935, the term "JP-A" as used herein means an "unexamined published Japanese patent application"), etc., and improved formulations for improving absorbability of the active ingredient have been proposed. With the recent various dermatohistological elucidations, an attempt at applying an epidermal growth factor (EGF) to postoperative healing of a wound has also been reported (see JP-A-3-106823).

However, the known compounds exemplified above have not been considered to have sufficient effects on healing a wound. Accordingly, it has been demanded to find a drug which directly acts on the develop to heal a wound.

SUMMARY OF THE INVENTION

Under the above-mentioned circumstances, the present inventors have extensively conducted investigations for the purpose of providing a drug for topical use effective for healing of a skin wound. As a result, they have found an excellent effect in a compound represented by formula (I) shown below and completed the present invention.

The present invention provides a composition for accelerating healing of a wound comprising as an active ingredient a compound represented by formula (I):




(I)

wherein R represents a hydrogen atom, a phenyl group or a phenyl group substituted with a halogen atom, a C₁-C₆ alkyl group, a hydroxyl group or a C₁-C₆ alkoxy group; and X represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group.

Although many of the compounds represented by formula (I) are known to exhibit antifungal (see U.S. Patent 4,636,519, U.S. Patent 4,738,976 and JP-B-4-68308) and antimicotic activities (see U.S. Patent 4,636,519 and U.S. Patent 4,738,976), it is unknown that the compounds of formula (I) possess an action of accelerating healing of a skin wound.

In another aspect, the present invention provides use of a composition for accelerating healing of a wound of the present invention for accelerating healing of a wound in a patient in need of acceleration for healing of a wound.


In another aspect, the present invention provides a composition for accelerating healing of a wound comprising a compound represented by formula (I):



wherein R represents a hydrogen atom, a phenyl group or a phenyl group substituted with a halogen atom, a C₁-C₆ alkyl group, a hydroxyl group or a C₁-C₆ alkoxy group; and X

represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group; and pharmaceutically acceptable carriers.


In another aspect, the present invention provides use of a compound represented by formula (I):




wherein R represents a hydrogen atom, a phenyl group or a phenyl group substituted with a halogen atom, a C₁-C₆ alkyl group, a hydroxyl group or a C₁-C₆ alkoxy group; and X represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group for accelerating healing of a wound.

DETAILED DESCRIPTION OF THE INVENTION

While the compounds of formula (I) each embrace two geometrical isomers as illustrated below, the compounds may be used in the present invention either as one of the isomers or as an isomeric mixture.



E-Form




Z-Form

Specific examples of the compounds which can be used in the present invention are shown in Tables 1 and 2 below for illustrative purposes only but not for limitation. Physicochemical properties of these compounds are also shown. In the Tables, "mp" is a melting point, "n_D" is a refractive

2104515

index, and the δ values (ppm) in NMR spectra are those measured in CDCl_3 using TMS as a standard.

TABLE 1












(Ia)

Compound No.	R	Physicochemical Properties
1	H	mp: 125.6°C
2		mp: 111-112°C (E-form)
3		mp: 119.4°C (Z-form)
4		mp: 141.5°C (E-form)
5		n_D^{24} : 1.6083
6		viscous oily substance (Z-form) NMR δ : 3.54-4.20 (m, 2H), 5.69 (dd, 1H), 7.00-7.75 (m, 7H)
7		viscous oily substance (E-form) NMR δ : 3.45-4.10 (m, 2H), 5.76 (dd, 1H), 7.00-7.85 (m, 7H)
8		viscous oily substance (Z-form) NMR δ : 3.6-4.0 (m, 2H), 5.22 (dd, 1H), 6.9-7.81 (m, 7H)

/To be cont'd

TABLE 1 (cont'd.)

Compound No.	R	Physicochemical Properties
9		mp: 148.8°C (E-form)
10		viscous oily substance (Z-form)
11		mp: 119°C (E-form)
12		viscous oily substance NMR δ : 3.6-3.9 (m, 2H), 5.1-5.5 (m, 1H), 6.9-7.8 (m, 7H)
13		n_D^{13} : 1.6313 (Z-form)
14		mp: 123.3°C (E-form)
15		n_D^{24} : 1.6360
16		n_D^{16} : 1.6196
17		viscous oily substance (Z-form)

/To be cont'd

TABLE 1 (cont'd.)






Compound No.	R	Physicochemical Properties
18		viscous oily substance (E-form) NMR δ : 3.71 (d, 2H), 3.82 (s, 3H), 5.68 (t, 1H), 6.87, 7.00 and 6.72 (each 1H, H of azole ring), 7.00- 7.60 (m, 4H)
19		viscous oily substance (Z-form) NMR δ : 3.75 (d, 2H), 3.77 (s, 3H), 5.17 (t, 1H), 6.74-7.60 (m, 7H)
20		viscous oily substance (E-form) NMR δ : 3.67 (d, 2H), 3.80 (s, 3H), 5.24 (t, 1H), 6.70-7.65 (m, 7H)
21		mp: 219-221°C (E-form)

TABLE 2








(Ib)

Compound No.	R	Physicochemical Properties
22	H	mp: 102.6°C
23		viscous oily substance
24		mp: 71-73°C (E-form)
25		mp: 75-77°C (Z-form)
26		n_D^{19} : 1.6558
27		viscous oily substance NMR δ : 3.82 (d, 2H), 5.31 (t, 1H), 7.18-7.51 (m, 4H), 8.11 (s, 1H), 8.39 (s, 1H)
28		viscous oily substance
29		$n_D^{20.5}$: 1.6232


/To be cont'd

TABLE 2 (cont'd.)

Compound No.	R	Physicochemical Properties
30		mp: 115.0°C
31		n_D^{24} : 1.6162
32		viscous oily substance
33		viscous oily substance
34		mp: 214-216°C (E-form)

R is preferably o-chlorophenyl, p-hydroxyphenyl or p-tolyl group.

Of the compounds represented by formula (I), those represented by formula (I'):



wherein X is as defined above,
are novel compounds.

The compound of the present invention can be converted to the salt by techniques known in the art. Examples of such salts include hydrochlorides, sulfates and nitrate.

The composition for accelerating healing of a wound according to the present invention can be prepared as various formulations such as solutions, emulsions, ointments, creams, lotions, poultices, tablets, granules, capsules, ampuls, etc. by compounding with pharmaceutically acceptable carriers. Specific examples of the pharmaceutically acceptable carriers are polyethylene glycol, 1,2-propanediol, glycerole stearate, spermaceti, isopropyl myristate, polysolvate, stearyl alcohol, cetanol, sorbitan monostearate, methyl p-hydroxybenzoate, propyl p-hydroxybenzoate, dibutylhydroxytoluene, calcium hydrogenphosphate, lactose, starch, heavy magnesium oxide, water, glucose solution and polyvinyl alcohol.

The compound of formula (I) is usually mixed with a base in an amount usually of from 0.01 to 50% by weight, preferably from 0.05 to 10% by weight, and more preferably from 0.1 to 3% by weight, based on the base weight.

The preparations of the present invention can be given orally or parenterally. When the preparations are orally given to adult patients, it is desirable that the patients are dosed with 1 to 100 mg/kg of body weight.

The present invention will now be illustrated in greater detail with reference to Formulation Examples and Test Examples, but it should be understood that the present

invention is not to be construed as being limited thereto. All the parts and percents are by weight unless otherwise indicated.

Synthesis Example for the compounds represented by formula (I) is described below. The other compounds are able to prepare according to the similar method described in U.S. Patent 4,636,519.

SYNTHESIS EXAMPLE

Synthesis of 2-(1-Imidazolyl)-2-{4-(hydroxyphenyl)-1,3-dithiolane-2-ylidene}acetonitrile (Compound No. 21)

In 40 ml of dimethyl sulfoxide were dissolved 4.6 g (0.004 mol) of 1-cyanomethylimidazole, 3.1 g (0.005 mol) of carbon disulfide and 5.2 g of potassium hydroxide powder by stirring at room temperature for 1 hour to prepare a dithiolate solution. The solution was added dropwise to a solution of 14.8 g (0.04 mol) of 1,2-dibromo-1-(4-t-butoxyphenyl)ethane in 40 ml of dimethyl sulfoxide at 30°C, followed by stirring for 2 hours. To the reaction mixture was added 200 ml of ice-water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water, dried over magnesium sulfate, and concentrated to obtain a mixture of geometric isomers as a viscous oily substance. Purification of the product by silica gel chromatography gave 2.4 g (yield: 16.8%) of a Z-isomer and 4.5 g (yield: 31.5%) of an E-isomer both as a viscous oily substance.

The resulting E-isomer was dissolved in 10 ml of trifluoroacetic acid, and the solution was stirred at room temperature for 1 hour. The reaction mixture was concentrated under reduced pressure, and the residue was dissolved in ethyl acetate and washed with an aqueous sodium hydrogencarbonate. The organic layer was dried over magnesium sulfate, and the solvent was removed by distillation under reduced pressure. Recrystallization of the residue from ethanol yielded 2.4 g (66.5%) of the titled compound (mp: 219-221°C).

FORMULATION EXAMPLE 1

Compound of the invention	1 part
Polyethylene glycol 400	99 parts

The above components were mixed to dissolve the active ingredient to prepare a solution for topical application.

FORMULATION EXAMPLE 2

Compound of the invention	2 parts
Polyethylene glycol 400	49 parts
Polyethylene glycol 4000	49 parts

The above components were mixed while heating to dissolve the active ingredient and cooled to prepare an ointment.

FORMULATION EXAMPLE 3

Compound of the invention	3 parts
1,2-Propanediol	5 parts
Glycerol stearate	5 parts
Spermaceti	5 parts

Isopropyl myristate	10 parts
---------------------	----------

Polysolvate	4 parts
-------------	---------

A mixture of the above components was warmed and cooled, and 68 parts of water were added thereto while stirring to prepare a cream.

FORMULATION EXAMPLE 4

Compound of the invention	0.1 part
---------------------------	----------

Stearyl alcohol	5.0 parts
-----------------	-----------

Cetanol	5.0 parts
---------	-----------

Middle-chain fatty acid triglyceride	10.0 parts
--------------------------------------	------------

Isopropyl myristate	5.0 parts
---------------------	-----------

Polysolvate 60	4.0 parts
----------------	-----------

Sorbitan monostearate	1.0 part
-----------------------	----------

Methyl p-hydroxybenzoate	0.14 part
--------------------------	-----------

Propyl p-hydroxybenzoate	0.06 part
--------------------------	-----------

Dibutylhydroxytoluene	0.02 part
-----------------------	-----------

Purified water	the balance
----------------	-------------

The above components were mixed in a usual manner to prepare a cream.

FORMULATION EXAMPLE 5

Compound of the invention	50 parts
---------------------------	----------

Starch	10 parts
--------	----------

Lactose	15 parts
---------	----------

Crystalline cellulose	20 parts
-----------------------	----------

Polyvinyl alcohol	5 parts
-------------------	---------

Water	30 parts
-------	----------

The above-mentioned components were homogenously kneaded, granulated, dried and sieved to obtain granules.

TEST EXAMPLE 1

Animals were shaved on their back, and two circular excisional wounds (1 cm diameter) per animal were made by cutting out the full-thickness skin with scissors. From the next day of the wounding, a cream containing 0.5% or 1% of Compound No. 4 which prepared in accordance with Formulation Example 4 was topically applied daily to the wounded site in the amount of 60 mg/site/day everyday. For comparison, a commercially available 5% Solcoseryl^{*} ointment (produced by Tobishi pharmaceutical Co., Ltd.) was applied in the same amount. The wound area was treated on the 4th, 6th, and 8th days of application, and the square measure was calculated by an Image analyzer (SPICCA-II,^{*} manufactured by Olympus Co., Ltd.). Date is expressed as % of the initial wound area which was obtained on the next day of wounding. On the 5th, 7th or 9th day of application, two rats per group were sacrificed, and the involved skin was excised for light microscopic observation.

The results of measurement of the wound area are shown in Table 3 below. Each area ratio based on the initial wound area as 100%. Because two of the animals of each group were put to death for histopathological examinations on the 5th, 7th and 9th days of application, the number of involved sites under observation reduced by two on each killing.

*Trade-mark

TABLE 3

<u>Test Group</u>	<u>Area Ratio of Wound (%)</u>		
	<u>4th Day</u>	<u>6th Day</u>	<u>8th Day</u>
Untreated group	87	74	41
Control group (treated with the base alone)	99	70	40
5.0% Solcoseryl [*] group	87	35	11
0.5% Compound 4 group	75	40	9
1.0% Compound 4 group	77	39	11

As is apparent from Table 3, the compound of the present invention shows a higher or equal activity of accelerating healing of a wound than or to Solcoseryl^{*} used as a reference drug. Histopathological examinations also revealed the activity of the compound in acceleration of angiogenesis, elogenation and differentiation of the epidermis, and formation of granulation tissue in the dermis which are important for the healing process of a wound.

TEST EXAMPLE 2

Rats were wounded on their back in the same manner as in Test Example 1. From the next day, a 1% polyethylene glycol 400 (PEG 400) solution of a test compound was applied daily in a volume of 0.1 ml/site/day, and the area ratio of the wound based on the initial wound was measured on the 5th and 7th day of application. The results obtained are shown in Tables 4 and 5 below.

*Trade-mark

TABLE 4

<u>Test Group</u>	<u>Area Ratio of Lesion (%)</u>	
	<u>5th Day</u>	<u>7th Day</u>
Untreated group	69	47
Control group (treated with only PEG 400)	62	34
Compound No. 1	67	25
Compound No. 2	63	25
Compound No. 3	52	20
Compound No. 4	51	21
Compound No. 22	62	25
Compound No. 24	65	29

TABLE 5

<u>Test Group</u>	<u>Area Ratio of Lesion (%)</u>	
	<u>5th Day</u>	<u>7th Day</u>
Untreated group	61	36
PEG 400 group	57	31
Compound No. 4	37	22
Compound No. 15	43	21
Compound No. 20	47	28
Compound No. 21	33	15
Compound No. 24	39	25

As shown in Tables 4 and 5, the compounds according to the present invention as well as Compound No. 4 possess a wound healing accelerating activity.

TEST EXAMPLE 3

Cotton pellets weighing 39-41 mg were sterilized by autoclaving, immersed with ethanol solution of Compound No. 4 (0.2 ml/pellet), and dried under negative pressure. Under light ether anesthesia, two pellets were subcutaneously implanted under bilateral Scapula of rats, respectively, through dorsal skin incision. Each animal (each group containing 10 rats) was intramuscularly injected a 1:1 mixture of penicillin G (1×10^4 units/ml) and streptomycin (8 mg/ml) in a volume of 0.2 ml to prevent it from suffering from bacterial infection. Animals were sacrificed 7 days after implantation of pellets, and each granulation tissue including the pellet was excised to measure its dry weight. The results are shown in Table 6.

TABLE 6

<u>Test Group</u>	<u>Dry weight of granulation tissue including cotton pellet (mg)</u>
Untreated group	102.1 $\pm$ 5.2
Control group (treated with the base only)	93.0 $\pm$ 4.7
0.5 mg/pellet Compound No. 4	108.7 $\pm$ 10.4
1.0 mg/pellet Compound No. 4	123.7 $\pm$ 7.1
2.0 mg/pellet Compound No. 4	136.4 $\pm$ 14.1
4.0 mg/pellet Compound No. 4	130.6 $\pm$ 6.0

As shown in Table 6, Compound No. 4 increased the dry weight of granulation tissue by the cotton pellet implantation


to show an activity of accelerating the formation of granulation tissue. The increases by the treatment with Compound No. 4 at dose more than 1 mg/pellet were significantly greater than those in controls.

As described above, a wound of the skin, etc. can be healed rapidly by applying the compound of the present invention as a composition for accelerating healing of the wound.

While the invention has been described in detail and with reference to specific examples thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.

CLAIMS

1. A composition for accelerating healing of a wound comprising as the active ingredient a compound represented by formula (I):



wherein R represents a hydrogen atom, a phenyl group or a phenyl group substituted with a halogen atom, a C₁-C₆ alkyl group, a hydroxyl group or a C₁-C₆ alkoxy group; and X represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group; and pharmaceutically acceptable carriers.


2. The composition for accelerating healing of a wound as claimed in claim 1, wherein R is o-chlorophenyl group, p-hydroxyphenyl group or p-tolyl group.

3. The composition according to claim 1 or 2, which is for use in topical application.

4. Use of a composition for accelerating healing of a wound of claim 1 or 2 for accelerating healing of a wound in a patient in need of acceleration for healing of a wound.


5. Use according to claim 4, wherein the composition is suitable for topical application.

6. A compound represented by formula (I'):



wherein X represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group.

7. A composition for accelerating healing of a wound comprising a compound represented by formula (I):




wherein R represents a hydrogen atom, a phenyl group or a phenyl group substituted with a halogen atom, a C₁-C₆ alkyl group, a hydroxyl group or a C₁-C₆ alkoxy group; and X represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group; and pharmaceutically acceptable carriers.

8. The composition for accelerating healing of a wound as claimed in claim 7, wherein R is o-chlorophenyl group, p-hydroxyphenyl group or p-tolyl group.

9. The composition according to claim 7 or 8, which is for use in topical application.


10. Use of a compound represented by formula (I):



wherein R represents a hydrogen atom, a phenyl group or a phenyl group substituted with a halogen atom, a C₁-C₆ alkyl group, a hydroxyl group or a C₁-C₆ alkoxy group; and X represents a 1-imidazolyl group or a 1,2,4-triazol-1-yl group for accelerating healing of a wound.

11. Use as claimed in claim 10, wherein R is o-chlorophenyl group, p-hydroxyphenyl group or p-tolyl group.

12. Use according to claim 10 or 11, wherein the compound is in a form suitable for topical application.



(I)