

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

09 November 2023 (09.11.2023)



(10) International Publication Number

WO 2023/215812 A1

(51) International Patent Classification:

A23F 3/14 (2006.01) A23L 2/60 (2006.01)

A23F 3/16 (2006.01) A23L 2/38 (2006.01)

A23F 3/40 (2006.01) A23L 2/58 (2006.01)

(21) International Application Number:

PCT/US2023/066583

(22) International Filing Date:

04 May 2023 (04.05.2023)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/338,235 04 May 2022 (04.05.2022) US

(71) Applicant: THE COCA-COLA COMPANY [US/US];

One Coca-cola Plaza, Nw, Atlanta, GA 30313 (US).

(72) Inventors: PRAKASH, Indra; 9750 Talisman Drive, Al-

pharetta, GA 30022 (US). HIGIRO, Juvenal; 1008 Water Oak Trace, Acworth, GA 30102 (US).

(74) Agent: SMYTHE, Nicole; King & Spalding, 1180

Peachtree Street, Atlanta, GA 30309 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CV, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, MG, MK, MN, MU, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, CV, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SC, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, ME, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: TEA BEVERAGES WITH IMPROVED TASTE

(57) Abstract: Tea beverages are provided, wherein the tea beverages comprise at least one tea product, at least one high potency sweetener, and at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. Methods of preparing the tea beverages are also provided.



WO 2023/215812 A1

TEA BEVERAGES WITH IMPROVED TASTE

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application No. 63/338,235, filed May 4, 2022, which is incorporated herein in its entirety.

FIELD OF THE INVENTION

The present invention relates generally to tea beverages sweetened with certain high potency sweeteners and modified with certain organic salts of sodium and/or potassium.

BACKGROUND OF THE INVENTION

Because of its unique aroma and taste, black tea is widely consumed in the world in various forms. Iced tea is a popular summer drink. Fresh iced tea is a cold drink prepared by brewing black tea followed by rapid cooling, thereby providing a cool and refreshing taste while still retaining the aroma of black tea. Large-scale preparation of packaged tea beverages, however, e.g., iced tea, is known to provide an astringent (chalky) and bitter taste.

To improve the taste of packaged black tea beverages, e.g., iced tea, sugar can be added (sucrose, fructose, glucose, lactic acid, etc.). However, this leads to increased caloric content and astringent and off-tastes may remain.

Typically, sugar-sweetened iced tea beverages contain from 30 grams of sugar to 60 grams of sugar per 18.5 US fluid ounces. Consumers increasingly prefer lower calorie versions of beverages generally, including tea beverages. Various approaches to reduced calorie tea beverages have been attempted. Substitution of synthetic high potency sweeteners for sugar provides the taste of sweetened iced tea beverages without the calories, e.g., Lipton's Zero Sugar Southern Sweet Iced Tea utilizes sucralose instead of sucrose and Gold Peak's Zero Sugar Sweet Tea utilizes aspartame and acesulfame K instead of sucrose.

However, consumers increasingly prefer natural high potency sweeteners. Stevia-based and monk fruit-based (*Luo Han Guo*) natural high potency sweeteners are popular choices for beverage formulations generally. However, these sweeteners are known to elicit off-tastes that are objectionable to consumers and taste different than sucrose, e.g., delayed sweetness onset, lingering sweet aftertaste, bitter taste, metallic taste, astringent taste, cooling taste, licorice-like taste, watery taste, and poor mouthfeel.

US 20080226773 describes a diet tea beverage having less than about 5 calories per 8 ounces sweetened with rebaudioside A instead of sucrose, optionally also containing erythritol and tagatose. However, rebaudioside A is known to elicit objectionable sweetness lingering and bitterness at high concentrations (e.g., 400-600 ppm).

Therefore, there remains a need for low-calorie and zero-calorie packaged tea beverages that have improved taste and flavor profiles.

SUMMARY OF THE INVENTION

The present invention relates to tea beverages comprising certain high potency sweeteners and certain organic salts of potassium or sodium. The organic salts of potassium or sodium act to modulate the sensory profile of the tea beverage to provide improved sensory properties over comparable tea beverages in the absence of said organic salts. The potassium or sodium salts disclosed herein also perform better than certain magnesium and calcium salts.

In one aspect, the present invention provides a tea beverage comprising:

- i. at least one tea product;
- ii. at least one high potency sweetener selected from the group consisting of rebaudioside A, rebaudioside M, rebaudioside AM, rebaudioside E, rebaudioside N, rebaudioside O, rebaudioside B, monk fruit juice or concentrate, monk fruit extract, siamenoside I, mogroside V, thaumatin (and variants thereof), brazzein (and variants thereof), monellin (and variants thereof), sweet truffle protein (and variants thereof), sucralose, potassium acesulfame, aspartame, neotame, saccharin, cyclamate, and combinations thereof; and
- iii. at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate.

The at least one high potency sweetener is present in a sweetening amount.

In certain embodiments, the at least one high potency sweetener comprises rebaudioside M. The rebaudioside M can be provided as a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, e.g., about 95% rebaudioside M by weight.

In other certain embodiments, the at least one high potency sweetener comprises monk fruit extract comprising about 45% or greater mogroside V by weight.

In other embodiments, the at least one high potency sweetener comprises (i) a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, e.g., about 95% rebaudioside M by weight and (ii) monk fruit extract comprising about 45% or greater mogroside V by weight on a dry basis.

In some embodiments, the tea beverages disclosed herein contain a single sodium salt selected from the group consisting of sodium gluconate, sodium citrate and sodium lactate. In other embodiments, the tea beverages disclosed herein contain a single potassium salt selected from the group consisting of potassium gluconate, potassium citrate, and potassium lactate.

The tea beverages disclosed herein are low-calorie or zero-calorie beverages. In certain embodiments, the tea beverages have a from about 0 to about 40 calories, or from about 0 to about 10 calories per 8 US fluid ounce serving.

The tea beverages can further comprise least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, or combinations thereof.

In another aspect, the present invention provides a method of preparing a tea beverage.

In one embodiment, a method of preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining.

In another embodiment, a method of preparing a tea beverage comprises combining (i) at least one tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency

sweetener; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate with water to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii) and the water.

Once all of the ingredients are combined, the tea beverage is pasteurized then cooled. The pasteurized tea beverage can be packaged, e.g., hot-filled into bottles prior to cooling. The cooled beverages can then be stored for distribution or immediate consumption.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

“Fruit extract,” as used herein, refers to extracts of fruits. Fruit extract may be produced from fruit peels, seeds and/or pulp

“Juice”, as used herein, refers to a beverage suitable for human consumption containing a liquid extracted from a fruit, vegetable, or plant. Examples of juices include, without limitation, fruit juice, vegetable juice, plant juice, plant water, and plant sap. Examples of juices also include juices that are reconstituted from-concentrate (FC) juices, not-from-concentrate (NFC) juices, purees, nectars or combinations thereof. Exemplary “juice beverages” include “fruit beverages,” “fruit juices,” “fruit mixed juices”, “fruit grains-containing fruit juices”, “fruit/vegetable mixed juices”, and “fruit juice-containing beverages” defined in the “Ordinance for Enforcement of the Fair Competition Code for Labeling of Fruit Beverages and the like” enforced in 2016.

“Reduced calorie,” as used herein, refers to a beverage comprising a mixture of caloric sweeteners (e.g., sucrose) and one or more non-sucrose sweeteners. Reduced-calorie beverages include mid-calorie beverages and low-calorie beverages.

“Mid-calorie beverage,” as used herein, refers to a beverage that has from 41 to 60 calories per 8 oz. serving.

“Low-calorie beverage,” as used herein, refers to a beverage that has from 6 to 40 calories per 8 oz. serving.

“Zero-calorie beverage,” as used herein, refers to a beverage that has less than 5 calories per 8 oz. serving.

“Natural high potency sweetener” or “NHPS” as used herein, refers to any sweetener found naturally in nature and characteristically has a sweetness potency greater than sucrose, fructose, or glucose, yet has less calories. The natural high potency sweetener can be provided as a pure compound or, alternatively, as part of an extract.

“No salty taste”, as used herein, refers to an inability to detect salty flavor in a beverage. Methods of determining whether a beverage tastes salty are known in the art, e.g., J. Giguère, et al., “Abstract 18991: Salt Taste Detection and Recognition Thresholds - Reliability of a Rapid Sensory Analysis Method”, Circulation, November 25, 2014, Vol 130, Issue suppl 2. The temporal stability of a rapid sensory analysis based on the 3-alternative forced-choice (3-AFC) method (ASTM E679) was tested with 30 adult volunteers. Detection Threshold (DT) and Recognition Threshold (RT) for salt were determined using a series of ascending concentrations.

“Synthetic high potency sweetener,” as used herein, refers to any composition which is not found naturally in nature and characteristically has a sweetness potency greater than sucrose, fructose, or glucose, yet has less calories.

“Tea product,” as used herein, refers to a product made from the leaves of the traditional tea plant (*Camellia sinensis*). The tea product can be in the form of brewed tea or a tea extract.

II. Tea Beverages

The present disclosure relates to tea beverages having low or zero calorie content. The tea beverages comprise at least one tea product, at least one high potency sweetener, and at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate.

It has been found that the described sodium and potassium salts can be used to effectively improve the taste attributes of the at least one high potency sweetener and provide tea beverages with improved sensory properties compared to a corresponding tea beverage in the absence of the sodium or potassium salt. Exemplary taste attribute modulations include decreasing or eliminating bitterness, decreasing or eliminating bitter linger, decreasing or eliminating sourness, decreasing or eliminating astringency, decreasing or eliminating saltiness, decreasing or eliminating metallic notes, improving mouthfeel, decreasing or eliminating sweetness linger, increasing sweetness onset and increasing sweetness intensity. Multiple taste attributes can be modulated simultaneously, such that the salt-containing tea beverage, overall, has more sucrose-sweetened characteristics compared to a corresponding

tea beverage without the salt(s). Methods of quantifying improvement in sucrose-sweetened characteristics are known in the art and include taste testing and histogram mapping with isosweet sucrose-sweetened beverage controls.

The tea beverages of the present invention contain at least one tea product, e.g., brewed tea or a tea extract. The tea can be any desired type of tea, for example, white, yellow, green, oolong, black, red, post-fermented, or a mixture thereof.

In certain embodiments, the tea beverage comprises brewed tea, e.g., black tea. Brewed tea is prepared by extracting the tea leaves with hot water (i.e., brewing), then separating the brewed tea from the leaves. In such embodiments, the brewed tea comprises at least about 50% by weight of the tea beverage, such as, for example, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 99%. In some embodiments, the beverage ingredients are added directly to the brewed tea to prepare the tea beverage. In other embodiments, the brewed tea is diluted with a volume of water, then the beverage ingredients are added to prepare the tea beverage.

Tea extract can be provided in liquid or powder form and is available from commercial sources or can be prepared. Tea extract is generally prepared by extracting the tea leaves with one or more solvents, then concentrating and/or drying as desired to provide a liquid concentrate or dried powder.

In embodiments where the tea beverage comprises water and a tea extract, e.g., a black tea extract, the tea extract is present in an amount from about 0.01% to about 6% by weight, such as, for example, from about 0.01% to about 1.5% by weight, from about 0.01% to about 0.1% or from about 0.05% to about 1.0%.

The tea beverages of the present invention are mainly water. The remaining ingredients are dissolved in the water. Purified water or water of a standard beverage quality can be employed in order not to adversely affect beverage taste, odor, or appearance. The water typically will be clear, colorless, free from objectionable minerals, tastes and odors, free from organic matter, low in alkalinity and of acceptable microbiological quality based on industry and government standards applicable at the time of producing the beverage. In certain embodiments, the water used in the tea beverages is "treated water," which refers to water that has been treated to reduce the total dissolved solids of the water prior to optional supplementation, e.g., with calcium. Methods of producing treated water are known to those of ordinary skill in the art and include deionization, distillation, filtration and reverse osmosis ("r-o"), among others. The terms "treated water," "purified water," "demineralized water,"

“distilled water,” and “r-o water” are understood to be generally synonymous, referring to water from which substantially all mineral content has been removed, typically containing no more than about 500 ppm total dissolved solids, e.g. 250 ppm total dissolved solids.

In exemplary embodiments, water comprises at least about 95% by weight of the tea beverage, e.g., at least about 96%, at least about 97%, at least about 98%, or at least about 99%.

The tea beverages of the present invention preferably contain from about 0 to about 40 calories per 8 US fluid ounce (240 mL) serving, such as, for example, from about 0 to about 30, from about 0 to about 20, from about 0 to about 10, from about 10 to about 40, from about 10 to about 30, from about 10 to about 20, from about 20 to about 40, from about 20 to about 30 or from about 30 to about 40.

The high potency sweetener is present in the tea beverage in a sweetening amount, i.e., in an amount above its sweetness recognition threshold such that it is perceptibly sweet. The high potency sweetener can be any known high potency sweetener, including natural and synthetic high potency sweeteners.

Non-limiting examples of natural high potency sweeteners include stevia sweeteners and steviol glycoside sweeteners, such as rebaudioside M, rebaudioside D, rebaudioside A, rebaudioside AM, rebaudioside N, rebaudioside O, rebaudioside E, steviolmonoside, steviolbioside, rubusoside, dulcoside B, dulcoside A, rebaudioside B, rebaudioside G, stevioside, rebaudioside C, rebaudioside F, rebaudioside I, rebaudioside H, rebaudioside L, rebaudioside K, rebaudioside J, rebaudioside M2, rebaudioside D2, rebaudioside S, rebaudioside T, rebaudioside U, rebaudioside V, rebaudioside W, rebaudioside Z1, rebaudioside Z2, rebaudioside IX, enzymatically glucosylated steviol glycosides, and combinations thereof.

In a particular embodiment, the high potency sweetener is a steviol glycoside sweetener selected from the group consisting of rebaudioside A, rebaudioside M, rebaudioside AM, rebaudioside E, rebaudioside N, rebaudioside O, and rebaudioside B.

Steviol glycoside sweeteners can be provided in pure form or as part of a mixture. The steviol glycoside mixture sweetener typically has a total steviol glycoside content of about 95% by weight or greater on a dry basis. The remaining 5% comprises other non-steviol glycoside compounds, e.g. by-products from extraction or purification processes. In some embodiments, the steviol glycoside blend sweetener has a total steviol glycoside content of about 96% or greater, about 97% or greater, about 98% or greater or about 99% or greater.

In certain embodiments, a steviol glycoside mixture comprises at least about 5% of a particular steviol glycoside by weight on a dry basis, such as, for example, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95% or at least about 97%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside A by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside A by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside M by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside M by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside AM by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside AM by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside E by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside E by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside D by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside D by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside E by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside E by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside N by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside N by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside O by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside O by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The steviol glycoside mixture may comprise at least about 50% rebaudioside B by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80%, from about 80% to about 99%, or from about 90% to about 99%.

The steviol glycoside mixture may comprise at least about 95% rebaudioside B by weight on a dry basis, such as, for example, from about 95% to about 99%, from about 96% to about 99%, from about 97% to about 99%, or from about 98% to about 99%.

The concentration of the steviol glycoside sweetener in the tea beverage can vary from about 25 ppm to about 600 ppm, such as, for example, from about 25 ppm to about 500 ppm, from about 25 ppm to about 400 ppm, from about 25 ppm to about 300 ppm, from about 25 ppm to about 200 ppm, from about 25 ppm to about 100 ppm, from about 100 ppm to about 600 ppm, from about 100 ppm to about 500 ppm, from about 100 ppm to about 400 ppm, from about 100 ppm to about 300 ppm, from about 100 ppm to about 200 ppm, from about 200 ppm to about 600 ppm, from about 200 ppm to about 500 ppm, from about 200 ppm to about 400 ppm, from about 200 ppm to about 300 ppm, from about 300 ppm to about 600 ppm, from about 300 ppm to about 500 ppm, from about 300 ppm to about 400 ppm, from about 400 ppm to about 600 ppm, from about 400 ppm to about 500 ppm, or from about 500 ppm to about 600 ppm. In a particular embodiment, the steviol glycoside sweetener is present in a concentration from about 100 ppm to about 600 ppm.

Exemplary natural high potency sweeteners also includes Luo Han Guo and the related mogroside compounds, such as monk fruit juice or concentrate, monk fruit extract, mogroside V, grosmogroside I, mogroside IA, mogroside IE, 11-oxomogroside IA, mogroside II, mogroside II A, mogroside II B, mogroside II E, 7-oxomogroside II E, mogroside III, Mogroside IIIe, 11-oxomogroside IIIe, 11- deoxymogroside III, mogroside IV, Mogroside IVA, 11-oxomogroside IV, 11-oxomogroside IVA, mogroside V, isomogroside V, 11-deoxymogroside V, 7-oxomogroside V, 11-oxomogroside V, isomogroside V, mogroside VI, mogrol, 11-oxomogrol, siamenoside I, isomers of siamenoside I (e.g. those disclosed in 20170119032; incorporated by reference in its entirety), 11-oxo-siamenoside I, 11-oxo-isomers of siamenoside I, (3 β ,9 β ,10 α ,11 α ,24R)-3-[(4-O- β -D-glucopyranosyl-6-O- β -D-glucopyranosyl]-25-hydroxyl-9-methyl-19-norlanost-5-en-24-yl-[2-O- β -D-glucopyranosyl-6-O- β -D-glucopyranosyl]- β -D-glucopyranoside); (3 β , 9 β , 10 α , 11 α , 24R)-[(2-O- β -D-glucopyranosyl-6-O- β -D-glucopyranosyl- β -D- glucopyranosyl)oxy]-25-hydroxy-9-methyl-19-norlanost-5-en-24-yl-[2-O- β -D-glucopyranosyl-6-O- β -D-glucopyranosyl]- β -D-glucopyranoside); and (3 β , 9 β , 10 α , 11 α , 24R)-[(2-O- β -D-glucopyranosyl-6-O- β -D-glucopyranosyl- β -D-glucopyranosyl)oxy]-25-hydroxy-9-methyl-19-norlanost-5-en-24-yl-[2-O- β -D-glucopyranosyl-6-O- β -D-glucopyranosyl]- β -D-glucopyranoside).

In particular embodiments, the mogroside sweetener is siamenoside I or mogroside V.

Mogroside sweeteners can be provided in pure form or as part of a mixture. In certain embodiments, a mogroside mixture comprises at least about 5% of a particular mogroside by weight on a dry basis, such as, for example, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95% or at least about 97%.

The mogroside mixture may comprise at least about 50% siamenoside I by weight on a dry basis, such as, for example, from about 50% to about 99%, from about 50% to about 80%, from about 50% to about 70%, from about 50% to about 60%, from about 60% to about 99%, from about 60% to about 80%, from about 60% to about 70%, from about 70% to about 99%, from about 70% to about 80% and from about 80% to about 99%.

The mogroside mixture may comprise at least about 25% mogroside V by weight on a dry basis, such as, for example, from about 25% to about 99%, from about 25% to about 80%, from about 25% to about 70%, from about 25% to about 60%, from about 25% to about 50%, from about 25% to about 40%, or from about 25% to about 30%.

In one embodiment, the mogroside sweetener is monk fruit extract. In certain embodiments, the monk fruit extract comprises about 45% or greater mogroside V by weight on a dry basis, such as, for example, about 50% or greater mogroside V, about 55% or greater mogroside V, about 60% or greater mogroside V, about 65% or greater mogroside V, about 70% or greater mogroside V, about 75% or greater mogroside V, about 80% or greater mogroside V, about 85% or greater mogroside V, about 90% or greater mogroside V or about 95% or greater mogroside V.

In another embodiment, the mogroside sweetener is monk fruit juice or monk fruit juice concentrate. The concentrate typically comprises at least 3.5 wt% mogroside V.

The concentration of the mogroside sweetener or mogroside mixture sweetener in the tea beverage can vary from about 25 ppm to about 600 ppm, such as, for example, from about 25 ppm to about 500 ppm, from about 25 ppm to about 400 ppm, from about 25 ppm to about 300 ppm, from about 25 ppm to about 200 ppm, from about 25 ppm to about 100 ppm, from about 100 ppm to about 600 ppm, from about 100 ppm to about 500 ppm, from about 100 ppm to about 400 ppm, from about 100 ppm to about 300 ppm, from about 100 ppm to about 200 ppm, from about 200 ppm to about 600 ppm, from about 200 ppm to about 500 ppm, from about 200 ppm to about 400 ppm, from about 200 ppm to about 300 ppm, from about 300 ppm to about 600 ppm, from about 300 ppm to about 500 ppm, from about 300 ppm to about 400 ppm, from about 400 ppm to about 600 ppm, from about 400 ppm to about 500 ppm or from about 500 ppm to about 600 ppm.

Other exemplary natural high potency sweeteners include Amai proteins, monatin and its salts (monatin SS, RR, RS, SR), curculin, glycyrrhizic acid and its salts, thaumatin (and variants thereof, e.g., thaumatin I, thaumatin II) monellin (and variants thereof), miraculin, mabinlin, brazzein (and variants thereof), sweet truffle protein (and variants thereof), hernandulcin, phyllodulcin, glycyphyllin, phloridzin, trilobatin, baiyunoside, osladin, polypodoside A, pterocaryoside A, pterocaryoside B, mukurozioside, phlomisioside I, periandrin I, abrusoside A, and cyclocarioside I.

Sweet truffle protein refers to the sweet proteins recently identified from fungal proteins, e.g., *M. terfezoides* gleba, also called "Myd polypeptides" according to US Patent Application No. 2021/0401013, incorporated herein by reference.

Non-limiting examples of synthetic high potency sweeteners include sucralose, potassium acesulfame, aspartame, alitame, saccharin, neohesperidin dihydrochalcone

synthetic derivatives, cyclamate, neotame, dulcin, suosan, cyclamate, saccharin, advantame, and salts thereof.

In exemplary embodiments, the high potency sweetener is selected from the group consisting of rebaudioside A, rebaudioside M, rebaudioside AM, rebaudioside E, rebaudioside N, rebaudioside O, rebaudioside B, monk fruit juice or concentrate, monk fruit extract, siamenoside I, mogroside V, thaumatin (and variants thereof), brazzein (and variants thereof), monellin (and variants thereof), sweet truffle protein (and variants thereof), sucralose, potassium acesulfame, neotame, aspartame, saccharin, cyclamate, and combinations thereof.

In certain embodiments, the at least one high potency sweetener comprises rebaudioside M. In other embodiments, the at least one high potency sweetener comprises a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis. In yet other embodiments, the at least one high potency sweetener comprises a steviol glycoside mixture comprising at least about 95% rebaudioside M by weight on a dry basis. In still further embodiments, the at least one high potency sweetener comprises rebaudioside M and monk fruit extract. In yet another embodiment, the at least one high potency sweetener comprises (i) a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis and (ii) monk fruit extract. In a still further embodiment, the at least one high potency sweetener comprises (i) a steviol glycoside mixture comprising at least about 95% rebaudioside M by weight on a dry basis and (ii) monk fruit extract.

The high potency sweetener is present in a sweetening amount, i.e., in an amount above the sweetness recognition threshold such that it is perceptibly sweet. The concentration of the high potency sweetener can vary from about 1 ppm to about 600 ppm, from about 1 ppm to about 500 ppm, from about 1 ppm to about 400 ppm, about 1 ppm to about 300 ppm, from about 1 ppm to about 200 ppm, from about 1 ppm to about 100 ppm, from about 1 ppm to about 50 ppm, from about 1 ppm to about 25 ppm, from about 1 ppm to about 15 ppm or about 1 ppm to about 10 ppm.

In certain embodiments, the tea beverage does not contain high potency synthetic sweeteners, e.g., aspartame, acesulfame K, sucralose, etc.

The sweetness of the tea beverages herein is provided by the at least one high potency sweetener. Therefore, in some embodiments, the tea beverages of the present invention do not contain added sugar or caloric sweeteners, e.g., sucrose, glucose, fructose, high fructose corn syrup, cane sugar, honey, and agave syrup and the like.

In other embodiments, the tea beverages use a mixture of caloric and non-caloric (high potency) sweeteners.

Suitable carbohydrate sweeteners include, but are not limited to, sucrose, glyceraldehyde, dihydroxyacetone, erythrose, threose, erythrulose, arabinose, lyxose, ribose, xylose, ribulose, xylulose, allose, altrose, galactose, glucose, gulose, idose, mannose, talose, fructose, psicose, sorbose, tagatose, mannoheptulose, sedoheltulose, octolose, fucose, rhamnose, arabinose, turanose, sialose, high fructose corn syrup and combinations thereof.

The amount of the carbohydrate sweetener in the tea beverage can vary from about 1 wt% to about 10 wt%, such as, for example, from about 4 wt% to about 10 wt%, about 5 wt% to about 10 wt%, about 6 wt% to about 10 wt%, about 7 wt% to about 10 wt%, about 8 wt% to about 10 wt% or about 9 wt% to about 10 wt%. In certain embodiments, the carbohydrate sweetener is present in an amount from about 1 wt% to about 3 wt%.

Other suitable sweeteners include allulose, allose, sucrose, fructose, glucose, propylene glycol, glycerol, erythritol, arabinitol, maltitol, lactitol, sorbitol, mannitol, xylitol, tagatose, trehalose, galactose, rhamnose, cyclodextrin (e.g., α -cyclodextrin, β -cyclodextrin, and γ -cyclodextrin), ribulose, threose, arabinose, xylose, lyxose, allose, altrose, mannose, idose, lactose, maltose, invert sugar, isotrehalose, neotrehalose, palatinose isomaltulose, erythrose, deoxyribose, gulose, idose, talose, erythrulose, xylulose, allulose, turanose, cellobiose, glucosamine, mannosamine, fucose, fuculose, glucuronic acid, gluconic acid, glucono-lactone, abequose, galactosamine, xylo-oligosaccharides (xylotriose, xylobiose and the like), gentio-oligosaccharides (gentiobiose, gentiotriose, gentiotetraose and the like), galacto-oligosaccharides, sorbose, ketotriose (dihydroxyacetone), aldotriose (glyceraldehyde), nigero-oligosaccharides, fructooligosaccharides (kestose, nystose and the like), maltotetraose, inaltotriol, tetrasaccharides, mannan-oligosaccharides, malto-oligosaccharides (maltotriose, maltotetraose, maltopentaose, maltohexaose, maltoheptaose and the like), dextrans, lactulose, melibiose, raffinose, rhamnose, ribose, isomerized, liquid sugars such as high fructose corn/starch syrup ("HFCS/HFSS," e.g., HFCS55, HFCS42, or HFCS90), coupling sugars, soybean oligosaccharides, glucose syrup and combinations thereof. It is understood that D- or L-configurations can be used when applicable.

Exemplary sugar alcohol sweeteners include, but are not limited to, sorbitol, mannitol, lactitol, maltitol, xylitol, erythritol and combinations thereof.

At least one sugar alcohol can be present in the tea beverage in an amount from about 0.1 wt% to about 3.5 wt%, such as, for example, from about 0.5 wt% to about 3.5 wt%, from

about 0.5 wt% to about 3.0 wt%, from about 0.5 wt% to about 2.5 wt%, from about 0.5 wt% to about 2.0 wt%, from about 0.5 wt% to about 1.5 wt%, from about 0.5 wt% to about 1.0 wt%, from about 1.0 wt% to about 3.5 wt%, from about 1.0 wt% to about 3.0 wt%, from about 1.0 wt% to about 2.5 wt%, from about 1.0 wt% to about 2.0 wt%, from about 1.0 wt% to about 1.5 wt%, from about 1.5 wt% to about 3.5 wt%, from about 1.5 wt% to about 3.0 wt%, from about 1.5 wt% to about 2.5 wt%, from about 1.5 wt% to about 2.0 wt%, from about 2.0 wt% to about 3.5 wt%, from about 2.0 wt% to about 3.0 wt%, from about 2.0 wt% to about 2.5 wt%, from about 2.5 wt% to about 3.5 wt%, from about 2.5 wt% to about 3.0 wt% or from about 3.0 wt% to about 3.5 wt%.

Other suitable calorie sweeteners include cane sugar, honey, and agave syrup.

The tea beverages of the present invention comprise at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. In certain embodiments, the at least one salt is a sodium salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, and combinations thereof. In other certain embodiments, the at least one salt is a potassium salt selected from the group consisting of potassium gluconate, potassium citrate, potassium lactate, and combinations thereof. It has been found that, in some instances, salts containing the same cation (here, sodium or potassium) provide superior taste attribute modulation than mixtures of sodium and potassium-containing salts.

The concentration of the at least one salt in the tea beverage can be from about 100 ppm to about 300 ppm, such as, for example, from about 100 ppm to about 250 ppm, from about 100 ppm to about 200 ppm, from about 100 ppm to about 150 ppm, from about 150 ppm to about 300 ppm, from about 150 ppm to about 250 ppm, from about 150 to about 200 ppm, from about 200 ppm to about 300 ppm, from about 200 ppm to about 250 ppm, from about 250 ppm to about 300 ppm. It has been found that such concentrations of the at least one salt provide superior results compared to both lower and higher concentrations. For example, higher concentrations provide undesirable flavor changes.

It has been found that, in some instances, the described potassium and sodium salts provide superior taste attribute modulation compared to magnesium or calcium salts. Therefore, in some embodiments, the tea beverages disclosed herein do not contain magnesium or calcium salts. In other embodiments, the tea beverages disclosed herein do not contain magnesium or calcium salts of gluconate, citrate or lactate.

It has also been found that, in some instances, a single sodium or potassium salt contemplated here, e.g., potassium gluconate or sodium gluconate, provides superior taste attribute modulation to mixtures of sodium-containing salts or potassium-containing salts. Therefore, in some embodiments, the tea beverages disclosed herein contain a single sodium salt selected from the group consisting of sodium gluconate, sodium citrate, and sodium lactate. In other embodiments, the tea beverages disclosed herein contain a single potassium salt selected from the group consisting of potassium gluconate, potassium citrate, and potassium lactate.

In certain embodiments, the tea beverages do not contain potassium, sodium, calcium, or magnesium salts of inorganic ions, e.g., chloride, fluoride, phosphate, monohydrogen phosphate, dihydrogen phosphate, sulfate, bisulfate, nitrate, carbonate, or bicarbonate.

The tea beverages of the present invention can be served hot, at room temperature or chilled (4 °C). In a particular embodiment, the tea beverage is an iced tea beverage.

In one embodiment, the tea beverage is a packaged tea beverage in, e.g., an aluminum bottle, plastic (e.g., PET) bottle, or glass bottle.

The tea beverages of the present invention can also contain at least one natural flavor, e.g., fruit or herb flavors. The flavor can be provided in any suitable form.

Exemplary fruit flavors include, but are not limited to, açai, apple, apricot, avocado, banana, blackberry, black currant, blueberry, breadfruit, cantaloupe, carambola, cherimoya, cherry, clementine, coconut, cranberry, custard-apple, date, dragonfruit, durian, elderberry, feijoa, fig, gooseberry, grapefruit, grape, guava, honeydew melon, jackfruit, java-plum, jujube fruit, kiwifruit, kumquat, lemon, lime, longan, loquat, lychee, mandarin, mango, mangosteen, mulberry, nectarine, orange, papaya, passion fruit, peach, pear, persimmon, pitaya (dragonfruit), pineapple, pitanga, plantain, plum, pomegranate, prickly pear, prune, pummelo, quince, raspberry, rhubarb, rose-apple, sapodilla, sapote, soursop, strawberry, sugar-apple, tamarine, tangerine, watermelon, and combinations thereof. In a particular embodiment, the tea beverage contains lemon and/or orange flavor.

Exemplary herb flavors include, but are not limited to chamomile, peppermint, hibiscus, lavender, marigold, chrysanthemum, ginger, spearmint, turmeric, and combinations thereof.

The tea beverages of the present invention can also contain at least one juice. Suitable juices include, but are not limited to, juices of citrus fruit (e.g., orange, grapefruit, lemon, lime,

tangerine, tangelo), apricot, apple, kumquat, mango, pear, peach, pineapple, papaya, passion fruit, grape, strawberry, raspberry, cranberry, currant, bean, blueberry, blackberry, acai, lychee, kiwi, pomegranate, watermelon, rhubarb, aronia, tomato, celery, cucurbits, onion, watercress, cucumber, carrot, parsley, beet, asparagus, potato, turnip, rutabaga, and combinations thereof.

The at least one juice can be provided in any form, e.g., fresh juice, juices, that are reconstituted from-concentrate (FC) juices, purees and/or nectars. Exemplary juices include lemon juice, including lemon juice from concentrate, peach puree, and mango puree.

The tea beverages of the present invention can also contain at least one fruit extract. Any suitable fruit extract can be used. Exemplary fruits that could be used to produce a fruit extract include, but are not limited to: apple, blueberry, pear, lemon, lime, mandarin, tangerine, grapefruit, cranberry, orange, strawberry, grape, kiwi, pineapple, passion fruit, mango, guava, raspberry, cherry, grapefruit, orange, lemon, lime, mandarin, mango, passion fruit, guava, and combinations thereof.

In a particular embodiment, the fruit extract is lemon extract.

The tea beverages of the present invention can also contain at least one herbal extract. Any suitable herbal extract can be used. Exemplary extracts include, but are not limited, yerba mate, chamomile, peppermint, hibiscus, marigold, chrysanthemum, ginger, turmeric.

The tea beverages of the present invention can also contain at least one herbal tea, i.e., brewed plant materials of something other than the traditional tea plant (*Camellia sinensis*). Any suitable herbal tea can be used. Suitable herbal teas include, but are not limited to, rooibos, lemon grass, tulsi, lemon peel, lemon myrtle, yerba mate, chamomile, peppermint, hibiscus, marigold, chrysanthemum, ginger, turmeric. Herbal teas also include teas made from plant material of any of the fruit listed above, i.e., apricot, apple, kumquat, mango, pear, peach, pineapple, papaya, passion fruit, grape, strawberry, raspberry, cranberry, currant, bean, blueberry, blackberry, acai, lychee, kiwi, pomegranate, watermelon, rhubarb, aronia, tomato, celery, cucurbits, onion, watercress, cucumber, carrot, parsley, beet, asparagus, potato, turnip, rutabaga, and combinations thereof.

The tea beverages can also contain at least one organic acid, which serves one or more additional functions, including, for example, lending tartness to the taste of the beverage, enhancing palatability, increasing thirst quenching effect, and acting as a mild preservative. Exemplary organic acids include citric acid, malic acid, ascorbic acid, tartaric acid, lactic acid, adipic acid, fumaric acid, gluconic acid, succinic acid, and maleic acid. The particular acid or

acids chosen and the amount used will depend, in part, on the other ingredients, the desired shelf life, as well as effects on the beverage pH, titratable acidity, and taste. The tea beverage contains at least one organic acid in an amount from about 0.1% to about 1.0% by weight, e.g., about 0.2% to about 0.7% by weight, or about 0.3% to about 0.6% by weight. In a particular embodiment, the tea beverage contains citric acid in an amount from about 0.1% to about 1.0% by weight.

A non-limiting example of the pH range of the tea beverage may be from about 2.5 to about 7. In a particular embodiment, the pH of an iced tea beverage is about 2.6 to about 4.1.

The amount of sucrose in a reference solution may be described in degrees Brix (°Bx). One degree Brix is 1 gram of sucrose in 100 grams of solution and represents the strength of the solution as percentage by weight (% w/w) (strictly speaking, by mass). In embodiments where sucrose is used in addition to the at least one high potency sweetener, the tea beverages of the present invention are from about 2 °Bx to about 7 °Bx, such as, for example, from about 4 °Bx to about 6 °Bx, or about 5 °Bx to about 6 °Bx.

A. Formulations

A tea beverage of the present invention comprises (i) at least one tea product; (ii) at least one high potency sweetener; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In one embodiment, a tea beverage comprises (i) at least one tea product selected from a brewed tea and a tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In a more particular embodiment, a tea beverage comprises (i) at least one tea product selected from a brewed tea and a tea extract, wherein the tea is selected from the group

consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener selected from the group consisting of rebaudioside A, rebaudioside M, rebaudioside AM, rebaudioside E, rebaudioside N, rebaudioside O, rebaudioside B, monk fruit juice or concentrate, monk fruit extract, siamenoside I, mogroside V, thaumatin (and variants thereof), brazzein (and variants thereof), monellin (and variants thereof), sweet truffle protein (and variants thereof), sucralose, potassium acesulfame, aspartame, neotame, saccharin, cyclamate, and combinations thereof; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. The tea beverage comprises the at least one high potency sweetener in an amount from about 1 ppm to about 600 ppm, as discussed above. The tea beverage comprises the at least one salt in an amount from about 100 ppm to about 300 ppm, as discussed above. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In a particular embodiment, a tea beverage comprises (i) black tea extract or brewed black tea; (ii) at least one high potency sweetener selected from the group consisting of rebaudioside A, rebaudioside M, rebaudioside AM, rebaudioside E, rebaudioside N, rebaudioside O, rebaudioside B, monk fruit juice or concentrate, monk fruit extract, siamenoside I, mogroside V, thaumatin (and variants thereof), brazzein (and variants thereof), monellin (and variants thereof), sweet truffle protein (and variants thereof), sucralose, potassium acesulfame, aspartame, neotame, saccharin, cyclamate, and combinations thereof; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. The tea beverage comprises the at least one high potency sweetener in an amount from about 1 ppm to about 600 ppm, as discussed above. The tea beverage comprises the at least one salt in an amount from about 100 ppm to about 300 ppm, as discussed above. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In a more particular embodiment, a tea beverage comprises (i) black tea extract or brewed black tea; (ii) at least one high potency sweetener comprising rebaudioside M; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. The tea beverage comprises rebaudioside M in an amount from about 25 to about 600 ppm, or about

100 ppm to about 600 ppm, as discussed above. The tea beverage comprises the at least one salt in an amount from about 100 pm to about 300 ppm, as discussed above. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In an even more particular embodiment, a tea beverage comprises (i) black tea extract or brewed black tea; (ii) at least one high potency sweetener comprising a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. The tea beverage comprises the steviol glycoside mixture comprising rebaudioside M in an amount from about 25 to about 600 ppm, or about 100 ppm to about 600 ppm, as discussed above. The tea beverage comprises the at least one salt in an amount from about 100 pm to about 300 ppm, as discussed above. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In an even more particular embodiment, a tea beverage comprises (i) black tea extract or brewed black tea; (ii) at least one high potency sweetener comprising a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis; and (iii) a salt selected from (a) sodium gluconate and (b) potassium gluconate. The tea beverage comprises the steviol glycoside mixture comprising rebaudioside M in an amount from about 25 to about 600 ppm, or about 100 ppm to about 600 ppm, as discussed above. The tea beverage comprises the salt in an amount from about 100 pm to about 300 ppm, as discussed above. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In another embodiment, a tea beverage comprises (i) black tea extract or brewed black tea; (ii) at least one high potency sweetener comprising rebaudioside M and monk fruit extract; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. The tea beverage comprises rebaudioside M in an amount from about 25 to about 600 ppm, or about 100 ppm to about 600 ppm, as discussed above. The tea beverage comprises monk fruit extract in an amount from about 25 ppm to about 100 ppm. The tea beverage comprises

the at least one salt in an amount from about 100 pm to about 300 ppm, as discussed above. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In a more particular embodiment, a tea beverage comprises (i) black tea extract or brewed black tea; (ii) at least one high potency sweetener comprising (a) a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis and (b) monk fruit extract; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate. The tea beverage comprises the steviol glycoside mixture comprising rebaudioside M in an amount from about 25 to about 600 ppm, or about 100 ppm to about 600 ppm, as discussed above. The tea beverage comprises the monk fruit extract in an amount from about 25 ppm to about 100 ppm, as discussed above. The tea beverage comprises the at least one salt in an amount from about 100 pm to about 300 ppm, as discussed above. As discussed above, the tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

In a still more particular embodiment, a tea beverage comprises (i) black tea extract or brewed black tea; (ii) at least one high potency sweetener comprising (a) a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis and (b) monk fruit extract; and (iii) at least one salt selected from (a) sodium gluconate and (b) a mixture of potassium citrate and potassium lactate. The tea beverage comprises the steviol glycoside mixture comprising rebaudioside M in an amount from about 25 to about 600 ppm, or about 100 ppm to about 600 ppm, as discussed above. The tea beverage comprises the monk fruit extract in an amount from about 25 to about 100 ppm, as discussed above. The tea beverage comprises the at least one salt in an amount from about 100 pm to about 300 ppm, as discussed above. The tea beverage can optionally comprise at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, and combinations thereof.

B. Functional Ingredients

The tea beverages described herein optionally include at least one functional ingredient described herein below.

Exemplary functional ingredients include, but are not limited to, saponins, antioxidants, dietary fiber sources, fatty acids, vitamins, glucosamine, minerals, preservatives, hydration agents, probiotics, prebiotics, weight management agents, osteoporosis management agents, phytoestrogens, long chain primary aliphatic saturated alcohols, phytosterols and combinations thereof.

In certain embodiments, the functional ingredient is at least one saponin. As used herein, the at least one saponin may comprise a single saponin or a plurality of saponins as a functional ingredient for the composition provided herein. Saponins are glycosidic natural plant products comprising an aglycone ring structure and one or more sugar moieties. Non-limiting examples of specific saponins for use in particular embodiments of the invention include group A acetyl saponin, group B acetyl saponin, and group E acetyl saponin. Several common sources of saponins include soybeans, which have approximately 5% saponin content by dry weight, soapwort plants (*Saponaria*), the root of which was used historically as soap, as well as alfalfa, aloe, asparagus, grapes, chickpeas, yucca, and various other beans and weeds. Saponins may be obtained from these sources by using extraction techniques well known to those of ordinary skill in the art. A description of conventional extraction techniques can be found in U.S. Pat. Appl. No. 2005/0123662.

In certain embodiments, the functional ingredient is at least one antioxidant. As used herein, "antioxidant" refers to any substance which inhibits, suppresses, or reduces oxidative damage to cells and biomolecules.

Examples of suitable antioxidants for embodiments of this invention include, but are not limited to, vitamins, vitamin cofactors, minerals, hormones, carotenoids, carotenoid terpenoids, non-carotenoid terpenoids, flavonoids, flavonoid polyphenolics (e.g., bioflavonoids), flavonols, flavones, phenols, polyphenols, esters of phenols, esters of polyphenols, nonflavonoid phenolics, isothiocyanates, and combinations thereof. In some embodiments, the antioxidant is vitamin A, vitamin C, vitamin E, ubiquinone, mineral selenium, manganese, melatonin, α -carotene, β -carotene, lycopene, lutein, zeaxanthin, cryptoxanthin, resveratrol, eugenol, quercetin, catechin, gossypol, hesperetin, curcumin, ferulic acid, thymol, hydroxytyrosol, tumeric, thyme, olive oil, lipoic acid, glutathione, glutamine, oxalic acid, tocopherol-derived compounds, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethylenediaminetetraacetic acid (EDTA), tert-butylhydroquinone, acetic acid, pectin, tocotrienol, tocopherol, coenzyme Q10, zeaxanthin, astaxanthin, canthaxanthin, saponins, limonoids, kaempferol, myricetin, isorhamnetin, proanthocyanidins, quercetin, rutin, luteolin, apigenin, tangeretin, hesperetin, naringenin, erodictyol, flavan-3-ols (e.g., anthocyanidins), gallocatechins, epicatechin and its gallate forms, epigallocatechin and its gallate forms

(EGCG) theaflavin and its gallate forms, thearubigins, isoflavone, phytoestrogens, genistein, daidzein, glycitein, anthocyanins, cyaniding, delphinidin, malvidin, pelargonidin, peonidin, petunidin, ellagic acid, gallic acid, salicylic acid, rosmarinic acid, cinnamic acid and its derivatives (e.g., ferulic acid), chlorogenic acid, chicoric acid, gallotannins, ellagitannins, anthoxanthins, betacyanins and other plant pigments, silymarin, citric acid, lignan, antinutrients, bilirubin, uric acid, R- α -lipoic acid, N-acetylcysteine, emblicanin, apple extract, apple skin extract (applephenon), rooibos extract red, rooibos extract, green, hawthorn berry extract, red raspberry extract, green coffee antioxidant (GCA), aronia extract 20%, grape seed extract (VinOseed), cocoa extract, hops extract, mangosteen extract, mangosteen hull extract, cranberry extract, pomegranate extract, pomegranate hull extract, pomegranate seed extract, hawthorn berry extract, pomella pomegranate extract, cinnamon bark extract, grape skin extract, bilberry extract, pine bark extract, pycnogenol, elderberry extract, mulberry root extract, wolfberry (gogi) extract, blackberry extract, blueberry extract, blueberry leaf extract, raspberry extract, turmeric extract, citrus bioflavonoids, black currant, ginger, acai powder, green coffee bean extract, green tea extract, and phytic acid, or combinations thereof. In alternate embodiments, the antioxidant is a synthetic antioxidant such as butylated hydroxytoluene or butylated hydroxyanisole, for example. Other sources of suitable antioxidants for embodiments of this invention include, but are not limited to, fruits, vegetables, tea, cocoa, chocolate, spices, herbs, rice, organ meats from livestock, yeast, whole grains, or cereal grains.

Particular antioxidants belong to the class of phytonutrients called polyphenols (also known as "polyphenolics"), which are a group of chemical substances found in plants, characterized by the presence of more than one phenol group per molecule. Suitable polyphenols for embodiments of this invention include catechins, proanthocyanidins, procyanidins, anthocyanins, quercetin, rutin, resveratrol, isoflavones, curcumin, punicalagin, ellagitannin, hesperidin, naringin, citrus flavonoids, chlorogenic acid, other similar materials, and combinations thereof.

In one embodiment, the antioxidant is a catechin such as, for example, epigallocatechin gallate (EGCG). In another embodiment, the antioxidant is chosen from proanthocyanidins, procyanidins or combinations thereof. In particular embodiments, the antioxidant is an anthocyanin. In still other embodiments, the antioxidant is chosen from quercetin, rutin or combinations thereof. In one embodiment, the antioxidant is resveratrol. In another embodiment, the antioxidant is an isoflavone. In still another embodiment, the antioxidant is curcumin. In a yet further embodiment, the antioxidant is chosen from

punicalagin, ellagitannin or combinations thereof. In a still further embodiment, the antioxidant is chlorogenic acid.

In certain embodiments, the functional ingredient is at least one dietary fiber. Numerous polymeric carbohydrates having significantly different structures in both composition and linkages fall within the definition of dietary fiber. Such compounds are well known to those skilled in the art, non-limiting examples of which include non-starch polysaccharides, lignin, cellulose, methylcellulose, the hemicelluloses, β -glucans, pectins, gums, mucilage, waxes, inulins, oligosaccharides, fructooligosaccharides, cyclodextrins, chitins, and combinations thereof. Although dietary fiber generally is derived from plant sources, indigestible animal products such as chitins are also classified as dietary fiber. Chitin is a polysaccharide composed of units of acetylglucosamine joined by $\beta(1-4)$ linkages, similar to the linkages of cellulose.

In certain embodiments, the functional ingredient is at least one fatty acid. As used herein, "fatty acid" refers to any straight chain monocarboxylic acid and includes saturated fatty acids, unsaturated fatty acids, long chain fatty acids, medium chain fatty acids, short chain fatty acids, fatty acid precursors (including omega-9 fatty acid precursors), and esterified fatty acids. As used herein, "long chain polyunsaturated fatty acid" refers to any polyunsaturated carboxylic acid or organic acid with a long aliphatic tail. As used herein, "omega-3 fatty acid" refers to any polyunsaturated fatty acid having a first double bond as the third carbon-carbon bond from the terminal methyl end of its carbon chain. In particular embodiments, the omega-3 fatty acid may comprise a long chain omega-3 fatty acid. As used herein, "omega-6 fatty acid" any polyunsaturated fatty acid having a first double bond as the sixth carbon-carbon bond from the terminal methyl end of its carbon chain.

Suitable omega-3 fatty acids for use in embodiments of the present invention can be derived from algae, fish, animals, plants, or combinations thereof, for example. Examples of suitable omega-3 fatty acids include, but are not limited to, linolenic acid, alpha-linolenic acid, eicosapentaenoic acid, docosahexaenoic acid, stearidonic acid, eicosatetraenoic acid and combinations thereof. In some embodiments, suitable omega-3 fatty acids can be provided in fish oils, (e.g., menhaden oil, tuna oil, salmon oil, bonito oil, and cod oil), microalgae omega-3 oils or combinations thereof. In particular embodiments, suitable omega-3 fatty acids may be derived from commercially available omega-3 fatty acid oils such as Microalgae DHA oil (from Martek, Columbia, MD), OmegaPure (from Omega Protein, Houston, TX), Marinol C-38 (from Lipid Nutrition, Channahon, IL), Bonito oil and MEG-3 (from Ocean Nutrition, Dartmouth, NS), Evogel (from Symrise, Holzminden, Germany), Marine Oil, from tuna or salmon (from

Arista Wilton, CT), OmegaSource 2000, Marine Oil, from menhaden and Marine Oil, from cod (from OmegaSource, RTP, NC).

Suitable omega-6 fatty acids include, but are not limited to, linoleic acid, gamma-linolenic acid, dihommo-gamma-linolenic acid, arachidonic acid, eicosadienoic acid, docosadienoic acid, adrenic acid, docosapentaenoic acid and combinations thereof.

Suitable esterified fatty acids for embodiments of the present invention include, but are not limited to, monoacylglycerols containing omega-3 and/or omega-6 fatty acids, diacylglycerols containing omega-3 and/or omega-6 fatty acids, or triacylglycerols containing omega-3 and/or omega-6 fatty acids and combinations thereof.

In certain embodiments, the functional ingredient is glucosamine, optionally further comprising chondroitin sulfate.

In certain embodiments, the functional ingredient is at least one preservative. In particular embodiments, the preservative is chosen from antimicrobials, antioxidants, antienzymatics or combinations thereof. Non-limiting examples of antimicrobials include sulfites, propionates, benzoates, sorbates, nitrates, nitrites, bacteriocins, salts, sugars, acetic acid, dimethyl dicarbonate (DMDC), ethanol, and ozone. In one embodiment, the preservative is a sulfite. Sulfites include, but are not limited to, sulfur dioxide, sodium bisulfite, and potassium hydrogen sulfite. In another embodiment, the preservative is a propionate. Propionates include, but are not limited to, propionic acid, calcium propionate, and sodium propionate. In yet another embodiment, the preservative is a benzoate. Benzoates include, but are not limited to, sodium benzoate and benzoic acid. In still another embodiment, the preservative is a sorbate. Sorbates include, but are not limited to, potassium sorbate, sodium sorbate, calcium sorbate, and sorbic acid. In a still further embodiment, the preservative is a nitrate and/or a nitrite. Nitrates and nitrites include, but are not limited to, sodium nitrate and sodium nitrite. In another embodiment, the at least one preservative is a bacteriocin, such as, for example, nisin. In still another embodiment, the preservative is ethanol. In yet another embodiment, the preservative is ozone. Non-limiting examples of antienzymatics suitable for use as preservatives in particular embodiments of the invention include ascorbic acid, citric acid, and metal chelating agents such as ethylenediaminetetraacetic acid (EDTA).

In certain embodiments, the functional ingredient is at least one hydration agent. In another particular embodiment, the hydration agent is a carbohydrate to supplement energy stores burned by muscles. Suitable carbohydrates for use in particular embodiments of this invention are described in U.S. Patent Numbers 4,312,856, 4,853,237, 5,681,569, and

6,989,171. Non-limiting examples of suitable carbohydrates include monosaccharides, disaccharides, oligosaccharides, complex polysaccharides or combinations thereof. Non-limiting examples of suitable types of monosaccharides for use in particular embodiments include trioses, tetroses, pentoses, hexoses, heptoses, octoses, and nonoses. Non-limiting examples of specific types of suitable monosaccharides include glyceraldehyde, dihydroxyacetone, erythrose, threose, erythrulose, arabinose, lyxose, ribose, xylose, ribulose, xylulose, allose, altrose, galactose, glucose, gulose, idose, mannose, talose, fructose, psicose, sorbose, tagatose, mannoheptulose, sedoheltulose, octolose, and sialose. Non-limiting examples of suitable disaccharides include sucrose, lactose, and maltose. Non-limiting examples of suitable oligosaccharides include saccharose, maltotriose, and maltodextrin. In other particular embodiments, the carbohydrates are provided by a corn syrup, a beet sugar, a cane sugar, a juice, or a tea.

In another particular embodiment, the hydration agent is a flavanol that provides cellular rehydration. Flavanols are a class of natural substances present in plants, and generally comprise a 2-phenylbenzopyrone molecular skeleton attached to one or more chemical moieties. Non-limiting examples of suitable flavanols for use in particular embodiments of this invention include catechin, epicatechin, galocatechin, epigallocatechin, epicatechin gallate, epigallocatechin 3-gallate, theaflavin, theaflavin 3-gallate, theaflavin 3'-gallate, theaflavin 3,3' gallate, thearubigin or combinations thereof. Several common sources of flavanols include tea plants, fruits, vegetables, and flowers. In preferred embodiments, the flavanol is extracted from green tea.

In a particular embodiment, the hydration agent is a glycerol solution to enhance exercise endurance. The ingestion of a glycerol containing solution has been shown to provide beneficial physiological effects, such as expanded blood volume, lower heart rate, and lower rectal temperature.

In certain embodiments, the functional ingredient is chosen from at least one probiotic, prebiotic and combination thereof. The probiotic is a beneficial microorganism that affects the human body's naturally-occurring gastrointestinal microflora. Examples of probiotics include, but are not limited to, bacteria of the genus *Lactobacilli*, *Bifidobacteria*, *Streptococci*, or combinations thereof, that confer beneficial effects to humans. In particular embodiments of the invention, the at least one probiotic is chosen from the genus *Lactobacilli*. According to other particular embodiments of this invention, the probiotic is chosen from the genus *Bifidobacteria*. In a particular embodiment, the probiotic is chosen from the genus *Streptococcus*.

Probiotics that may be used in accordance with this invention are well-known to those of skill in the art. Non-limiting examples of foodstuffs comprising probiotics include yogurt, sauerkraut, kefir, kimchi, fermented vegetables, and other foodstuffs containing a microbial element that beneficially affects the host animal by improving the intestinal microbalance.

Prebiotics, in accordance with the embodiments of this invention, include, without limitation, mucopolysaccharides, oligosaccharides, polysaccharides, amino acids, vitamins, nutrient precursors, proteins and combinations thereof. According to a particular embodiment of this invention, the prebiotic is chosen from dietary fibers, including, without limitation, polysaccharides and oligosaccharides. Non-limiting examples of oligosaccharides that are categorized as prebiotics in accordance with particular embodiments of this invention include fructooligosaccharides, inulins, isomalto-oligosaccharides, lactilol, lactosucrose, lactulose, pyrodextrins, soy oligosaccharides, transgalacto-oligosaccharides, and xylo-oligosaccharides. In other embodiments, the prebiotic is an amino acid. Although a number of known prebiotics break down to provide carbohydrates for probiotics, some probiotics also require amino acids for nourishment.

Prebiotics are found naturally in a variety of foods including, without limitation, bananas, berries, asparagus, garlic, wheat, oats, barley (and other whole grains), flaxseed, tomatoes, Jerusalem artichoke, onions and chicory, greens (e.g., dandelion greens, spinach, collard greens, chard, kale, mustard greens, turnip greens), and legumes (e.g., lentils, kidney beans, chickpeas, navy beans, white beans, black beans).

In certain embodiments, the functional ingredient is at least one weight management agent. As used herein, "a weight management agent" includes an appetite suppressant and/or a thermogenesis agent. As used herein, the phrases "appetite suppressant", "appetite satiation compositions", "satiety agents", and "satiety ingredients" are synonymous. The phrase "appetite suppressant" describes macronutrients, herbal extracts, exogenous hormones, anorectics, anorexigenics, pharmaceutical drugs, and combinations thereof, that when delivered in an effective amount, suppress, inhibit, reduce, or otherwise curtail a person's appetite. The phrase "thermogenesis agent" describes macronutrients, herbal extracts, exogenous hormones, anorectics, anorexigenics, pharmaceutical drugs, and combinations thereof, that when delivered in an effective amount, activate or otherwise enhance a person's thermogenesis or metabolism.

Suitable weight management agents include macronutrients selected from the group consisting of proteins, carbohydrates, dietary fats, and combinations thereof. Consumption of proteins, carbohydrates, and dietary fats stimulates the release of peptides with appetite-

suppressing effects. For example, consumption of proteins and dietary fats stimulates the release of the gut hormone cholecystokinin (CCK), while consumption of carbohydrates and dietary fats stimulates release of Glucagon-like peptide 1 (GLP-1).

Suitable macronutrient weight management agents also include carbohydrates. Carbohydrates generally comprise sugars, starches, cellulose and gums that the body converts into glucose for energy. Carbohydrates often are classified into two categories, digestible carbohydrates (e.g., monosaccharides, disaccharides, and starch) and non-digestible carbohydrates (e.g., dietary fiber). Studies have shown that non-digestible carbohydrates and complex polymeric carbohydrates having reduced absorption and digestibility in the small intestine stimulate physiologic responses that inhibit food intake. Accordingly, the carbohydrates embodied herein desirably comprise non-digestible carbohydrates or carbohydrates with reduced digestibility. Non-limiting examples of such carbohydrates include polydextrose; inulin; monosaccharide-derived polyols such as erythritol, mannitol, xylitol, and sorbitol; disaccharide-derived alcohols such as isomalt, lactitol, and maltitol; and hydrogenated starch hydrolysates. Carbohydrates are described in more detail herein below.

In another particular embodiment, the weight management agent is a dietary fat. Dietary fats are lipids comprising combinations of saturated and unsaturated fatty acids. Polyunsaturated fatty acids have been shown to have a greater satiating power than mono-unsaturated fatty acids. Accordingly, the dietary fats embodied herein desirably comprise poly-unsaturated fatty acids, non-limiting examples of which include triacylglycerols.

In another particular embodiment, the weight management agent is an herbal extract. Extracts from numerous types of plants have been identified as possessing appetite suppressant properties. Non-limiting examples of plants whose extracts have appetite suppressant properties include plants of the genus *Hoodia*, *Trichocaulon*, *Caralluma*, *Stapelia*, *Orbea*, *Asclepias*, and *Camelia*. Other embodiments include extracts derived from *Gymnema Sylvestre*, Kola Nut, Citrus Aurantium, Yerba Mate, *Griffonia Simplicifolia*, Guarana, myrrh, guggul Lipid, and black current seed oil.

The herbal extracts may be prepared from any type of plant material or plant biomass. Non-limiting examples of plant material and biomass include the stems, roots, leaves, dried powder obtained from the plant material, and sap or dried sap. The herbal extracts generally are prepared by extracting sap from the plant and then spray-drying the sap. Alternatively, solvent extraction procedures may be employed. Following the initial extraction, it may be desirable to further fractionate the initial extract (e.g., by column chromatography) in order to

obtain an herbal extract with enhanced activity. Such techniques are well known to those of ordinary skill in the art.

In one embodiment, the herbal extract is derived from a plant of the genus *Hoodia*. A sterol glycoside of *Hoodia*, known as P57, is believed to be responsible for the appetite-suppressant effect of the *Hoodia* species. In another embodiment, the herbal extract is derived from a plant of the genus *Caralluma*, non-limiting examples of which include caratuberside A, caratuberside B, bouceroside I, bouceroside II, bouceroside III, bouceroside IV, bouceroside V, bouceroside VI, bouceroside VII, bouceroside VIII, bouceroside IX, and bouceroside X. In another embodiment, the at least one herbal extract is derived from a plant of the genus *Trichocaulon*. *Trichocaulon* plants are succulents that generally are native to southern Africa, similar to *Hoodia*, and include the species *T. piliferum* and *T. officinale*. In another embodiment, the herbal extract is derived from a plant of the genus *Stapelia* or *Orbea*. Not wishing to be bound by any theory, it is believed that the compounds exhibiting appetite suppressant activity are saponins, such as pregnane glycosides, which include stavarosides A, B, C, D, E, F, G, H, I, J, and K. In another embodiment, the herbal extract is derived from a plant of the genus *Asclepias*. Not wishing to be bound by any theory, it is believed that the extracts comprise steroidal compounds, such as pregnane glycosides and pregnane aglycone, having appetite suppressant effects.

In another particular embodiment, the weight management agent is an exogenous hormone having a weight management effect. Non-limiting examples of such hormones include CCK, peptide YY, ghrelin, bombesin and gastrin-releasing peptide (GRP), enterostatin, apolipoprotein A-IV, GLP-1, amylin, somastatin, and leptin.

In another embodiment, the weight management agent is a pharmaceutical drug. Non-limiting examples include phentermine, diethylpropion, phendimetrazine, sibutramine, rimonabant, oxyntomodulin, floxetine hydrochloride, ephedrine, phenethylamine, or other stimulants.

In certain embodiments, the functional ingredient is at least one osteoporosis management agent. In certain embodiments, the osteoporosis management agent is at least one calcium source. According to a particular embodiment, the calcium source is any compound containing calcium, including salt complexes, solubilized species, and other forms of calcium. Non-limiting examples of calcium sources include amino acid chelated calcium, calcium carbonate, calcium oxide, calcium hydroxide, calcium sulfate, calcium chloride, calcium phosphate, calcium hydrogen phosphate, calcium dihydrogen phosphate, calcium

citrate, calcium malate, calcium citrate malate, calcium gluconate, calcium tartrate, calcium lactate, solubilized species thereof, and combinations thereof.

According to a particular embodiment, the osteoporosis management agent is a magnesium source. The magnesium source is any compound containing magnesium, including salt complexes, solubilized species, and other forms of magnesium. Non-limiting examples of magnesium sources include magnesium chloride, magnesium citrate, magnesium gluceptate, magnesium gluconate, magnesium lactate, magnesium hydroxide, magnesium picolate, magnesium sulfate, solubilized species thereof, and mixtures thereof. In another particular embodiment, the magnesium source comprises an amino acid chelated or creatine chelated magnesium.

In other embodiments, the osteoporosis agent is chosen from vitamins D, C, K, their precursors and/or beta-carotene and combinations thereof.

Numerous plants and plant extracts also have been identified as being effective in the prevention and treatment of osteoporosis. Non-limiting examples of suitable plants and plant extracts as osteoporosis management agents include species of the genus *Taraxacum* and *Amelanchier*, as disclosed in U.S. Patent Publication No. 2005/0106215, and species of the genus *Lindera*, *Artemisia*, *Acorus*, *Carthamus*, *Carum*, *Cnidium*, *Curcuma*, *Cyperus*, *Juniperus*, *Prunus*, *Iris*, *Cichorium*, *Dodonaea*, *Epimedium*, *Erigonoum*, *Soya*, *Mentha*, *Ocimum*, *thymus*, *Tanacetum*, *Plantago*, *Spearmint*, *Bixa*, *Vitis*, *Rosemarinus*, *Rhus*, and *Anethum*, as disclosed in U.S. Patent Publication No. 2005/0079232.

In certain embodiments, the functional ingredient is at least one phytoestrogen. Phytoestrogens are compounds found in plants which can typically be delivered into human bodies by ingestion of the plants or the plant parts having the phytoestrogens. As used herein, "phytoestrogen" refers to any substance which, when introduced into a body causes an estrogen-like effect of any degree. For example, a phytoestrogen may bind to estrogen receptors within the body and have a small estrogen-like effect.

Examples of suitable phytoestrogens for embodiments of this invention include, but are not limited to, isoflavones, stilbenes, lignans, resorcylic acid lactones, coumestans, coumestrol, equol, and combinations thereof. Sources of suitable phytoestrogens include, but are not limited to, whole grains, cereals, fibers, fruits, vegetables, black cohosh, agave root, black currant, black haw, chasteberry, cramp bark, dong quai root, devil's club root, false unicorn root, ginseng root, groundsel herb, licorice, liferoot herb, motherwort herb, peony root, raspberry leaves, rose family plants, sage leaves, sarsaparilla root, saw palmetto berry, wild

yam root, yarrow blossoms, legumes, soybeans, soy products (e.g., miso, soy flour, soymilk, soy nuts, soy protein isolate, tempen, or tofu) chick peas, nuts, lentils, seeds, clover, red clover, dandelion leaves, dandelion roots, fenugreek seeds, green tea, hops, red wine, flaxseed, garlic, onions, linseed, borage, butterfly weed, caraway, chaste tree, vitex, dates, dill, fennel seed, gotu kola, milk thistle, pennyroyal, pomegranates, southernwood, soya flour, tansy, and root of the kudzu vine (pueraria root) and the like, and combinations thereof.

Isoflavones belong to the group of phytonutrients called polyphenols. In general, polyphenols (also known as "polyphenolics"), are a group of chemical substances found in plants, characterized by the presence of more than one phenol group per molecule.

Suitable phytoestrogen isoflavones in accordance with embodiments of this invention include genistein, daidzein, glycitein, biochanin A, formononetin, their respective naturally occurring glycosides and glycoside conjugates, matairesinol, secoisolariciresinol, enterolactone, enterodiol, textured vegetable protein, and combinations thereof.

Suitable sources of isoflavones for embodiments of this invention include, but are not limited to, soy beans, soy products, legumes, alfalfa sprouts, chickpeas, peanuts, and red clover.

In certain embodiments, the functional ingredient is at least one long chain primary aliphatic saturated alcohol. Long-chain primary aliphatic saturated alcohols are a diverse group of organic compounds. The term alcohol refers to the fact these compounds feature a hydroxyl group (-OH) bound to a carbon atom. Non-limiting examples of particular long-chain primary aliphatic saturated alcohols for use in particular embodiments of the invention include the 8 carbon atom 1-octanol, the 9 carbon 1-nonanol, the 10 carbon atom 1-decanol, the 12 carbon atom 1-dodecanol, the 14 carbon atom 1-tetradecanol, the 16 carbon atom 1-hexadecanol, the 18 carbon atom 1-octadecanol, the 20 carbon atom 1-eicosanol, the 22 carbon 1-docosanol, the 24 carbon 1-tetracosanol, the 26 carbon 1-hexacosanol, the 27 carbon 1-heptacosanol, the 28 carbon 1-octacosol, the 29 carbon 1-nonacosanol, the 30 carbon 1-triacontanol, the 32 carbon 1-dotriacontanol, and the 34 carbon 1-tetracontanol.

In one embodiment, the long-chain primary aliphatic saturated alcohol is a policosanol. Policosanol is the term for a mixture of long-chain primary aliphatic saturated alcohols composed primarily of 28 carbon 1-octacosol and 30 carbon 1-triacontanol, as well as other alcohols in lower concentrations such as 22 carbon 1-docosanol, 24 carbon 1-tetracosanol, 26 carbon 1-hexacosanol, 27 carbon 1-heptacosanol, 29 carbon 1-nonacosanol, 32 carbon 1-dotriacontanol, and 34 carbon 1-tetracontanol.

In certain embodiments, the functional ingredient is at least one phytosterol, phytostanol or combination thereof. As used herein, the phrases “stanol”, “plant stanol” and “phytostanol” are synonymous. Plant sterols and stanols are present naturally in small quantities in many fruits, vegetables, nuts, seeds, cereals, legumes, vegetable oils, bark of the trees and other plant sources. Sterols are a subgroup of steroids with a hydroxyl group at C-3. Generally, phytosterols have a double bond within the steroid nucleus, like cholesterol; however, phytosterols also may comprise a substituted side chain (R) at C-24, such as an ethyl or methyl group, or an additional double bond. The structures of phytosterols are well known to those of skill in the art.

At least 44 naturally-occurring phytosterols have been discovered, and generally are derived from plants, such as corn, soy, wheat, and wood oils; however, they also may be produced synthetically to form compositions identical to those in nature or having properties similar to those of naturally-occurring phytosterols. Non-limiting suitable phytosterols include, but are not limited to, 4-desmethylsterols (e.g., β -sitosterol, campesterol, stigmasterol, brassicasterol, 22-dehydrobrassicasterol, and Δ^5 -avenasterol), 4-monomethyl sterols, and 4,4-dimethyl sterols (triterpene alcohols) (e.g., cycloartenol, 24-methylenecycloartanol, and cyclobranol).

As used herein, the phrases “stanol”, “plant stanol” and “phytostanol” are synonymous. Phytostanols are saturated sterol alcohols present in only trace amounts in nature and also may be synthetically produced, such as by hydrogenation of phytosterols. Suitable phytostanols include, but are not limited to, β -sitostanol, campestanol, cycloartanol, and saturated forms of other triterpene alcohols.

Both phytosterols and phytostanols, as used herein, include the various isomers such as the α and β isomers. The phytosterols and phytostanols of the present invention also may be in their ester form. Suitable methods for deriving the esters of phytosterols and phytostanols are well known to those of ordinary skill in the art, and are disclosed in U.S. Patent Numbers 6,589,588, 6,635,774, 6,800,317, and U.S. Patent Publication Number 2003/0045473. Non-limiting examples of suitable phytosterol and phytostanol esters include sitosterol acetate, sitosterol oleate, stigmasterol oleate, and their corresponding phytostanol esters. The phytosterols and phytostanols of the present invention also may include their derivatives.

C. Additives

The tea beverages described herein can further include at least one additive. Exemplary additives include, but not limited to, carbohydrates, polyols, amino acids and their corresponding salts, poly-amino acids and their corresponding salts, sugar acids and their

corresponding salts, nucleotides, organic acids, inorganic acids, bitter compounds, caffeine, flavorants and flavoring ingredients, astringent compounds, proteins or protein hydrolysates, surfactants, emulsifiers, plant extracts, flavonoids, alcohols, polymers and combinations thereof.

The term "polyol", as used herein, refers to a molecule that contains more than one hydroxyl group. A polyol may be a diol, triol, or a tetraol which contains 2, 3, and 4 hydroxyl groups respectively. A polyol also may contain more than 4 hydroxyl groups, such as a pentaol, hexaol, heptaol, or the like, which contain 5, 6, or 7 hydroxyl groups, respectively. Additionally, a polyol also may be a sugar alcohol, polyhydric alcohol, or polyalcohol which is a reduced form of carbohydrate, wherein the carbonyl group (aldehyde or ketone, reducing sugar) has been reduced to a primary or secondary hydroxyl group. Non-limiting examples of polyols in some embodiments include maltitol, mannitol, sorbitol, lactitol, xylitol, isomalt, propylene glycol, glycerol (glycerin), threitol, galactitol, palatinose, reduced isomalto-oligosaccharides, reduced xylo-oligosaccharides, reduced gentio-oligosaccharides, reduced maltose syrup, reduced glucose syrup, and sugar alcohols or any other carbohydrates capable of being reduced which do not adversely affect taste.

Suitable amino acid additives include, but are not limited to, aspartic acid, arginine, glycine, glutamic acid, proline, threonine, theanine, cysteine, cystine, alanine, valine, tyrosine, leucine, arabinose, trans-4-hydroxyproline, isoleucine, asparagine, serine, lysine, histidine, ornithine, methionine, carnitine, aminobutyric acid (α -, β -, and/or δ -isomers), glutamine, hydroxyproline, taurine, norvaline, sarcosine, and their salt forms such as sodium or potassium salts or acid salts. The amino acid additives also may be in the D- or L-configuration and in the mono-, di-, or tri-form of the same or different amino acids. Additionally, the amino acids may be α -, β -, γ - and/or δ -isomers if appropriate. Combinations of the foregoing amino acids and their corresponding salts (e.g., sodium, potassium, calcium, magnesium salts or other alkali or alkaline earth metal salts thereof, or acid salts) also are suitable additives in some embodiments. The amino acids may be natural or synthetic. The amino acids also may be modified. Modified amino acids refers to any amino acid wherein at least one atom has been added, removed, substituted, or combinations thereof (e.g., N-alkyl amino acid, N-acyl amino acid, or N-methyl amino acid). Non-limiting examples of modified amino acids include amino acid derivatives such as trimethyl glycine, N-methyl-glycine, and N-methyl-alanine. As used herein, modified amino acids encompass both modified and unmodified amino acids. As used herein, amino acids also encompass both peptides and polypeptides (e.g., dipeptides, tripeptides, tetrapeptides, and pentapeptides) such as glutathione and L-alanyl-L-glutamine.

Suitable polyamino acid additives include poly-L-aspartic acid, poly-L-lysine (e.g., poly-L- α -lysine or poly-L- ϵ -lysine), poly-L-ornithine (e.g., poly-L- α -ornithine or poly-L- ϵ -ornithine), poly-L-arginine, other polymeric forms of amino acids, and salt forms thereof (e.g., calcium, potassium, sodium, or magnesium salts such as L-glutamic acid mono sodium salt). The poly-amino acid additives also may be in the D- or L-configuration. Additionally, the poly-amino acids may be α -, β -, γ -, δ -, and ϵ -isomers if appropriate. Combinations of the foregoing poly-amino acids and their corresponding salts (e.g., sodium, potassium, calcium, magnesium salts or other alkali or alkaline earth metal salts thereof or acid salts) also are suitable additives in some embodiments. The poly-amino acids described herein also may comprise copolymers of different amino acids. The poly-amino acids may be natural or synthetic. The poly-amino acids also may be modified, such that at least one atom has been added, removed, substituted, or combinations thereof (e.g., N-alkyl poly-amino acid or N-acyl poly-amino acid). As used herein, poly-amino acids encompass both modified and unmodified poly-amino acids. For example, modified poly-amino acids include, but are not limited to, poly-amino acids of various molecular weights (MW), such as poly-L- α -lysine with a MW of 1,500, MW of 6,000, MW of 25,200, MW of 63,000, MW of 83,000, or MW of 300,000.

Suitable sugar acid additives include, but are not limited to, aldonic, uronic, aldaric, alginic, gluconic, glucuronic, glucaric, galactaric, galacturonic, and salts thereof (e.g., sodium, potassium, calcium, magnesium salts or other physiologically acceptable salts), and combinations thereof.

Suitable nucleotide additives include, but are not limited to, inosine monophosphate ("IMP"), guanosine monophosphate ("GMP"), adenosine monophosphate ("AMP"), cytosine monophosphate (CMP), uracil monophosphate (UMP), inosine diphosphate, guanosine diphosphate, adenosine diphosphate, cytosine diphosphate, uracil diphosphate, inosine triphosphate, guanosine triphosphate, adenosine triphosphate, cytosine triphosphate, uracil triphosphate, alkali or alkaline earth metal salts thereof, and combinations thereof. The nucleotides described herein also may comprise nucleotide-related additives, such as nucleosides or nucleic acid bases (e.g., guanine, cytosine, adenine, thymine, uracil).

Suitable organic acid additives include any compound which comprises a -COOH moiety, such as, for example, C2-C30 carboxylic acids, substituted hydroxyl C2-C30 carboxylic acids, butyric acid (ethyl esters), substituted butyric acid (ethyl esters), benzoic acid, substituted benzoic acids (e.g., 2,4-dihydroxybenzoic acid), substituted cinnamic acids, hydroxyacids, substituted hydroxybenzoic acids, anisic acid substituted cyclohexyl carboxylic acids, tannic acid, aconitic acid, lactic acid, tartaric acid, citric acid, isocitric acid, gluconic acid, glucoheptonic acids, adipic acid, hydroxycitric acid, malic acid, fruitaric acid (a blend of malic,

fumaric, and tartaric acids), fumaric acid, maleic acid, succinic acid, chlorogenic acid, salicylic acid, creatine, caffeic acid, bile acids, acetic acid, ascorbic acid, alginic acid, erythorbic acid, polyglutamic acid, glucono delta lactone, and their alkali or alkaline earth metal salt derivatives thereof. In addition, the organic acid additives also may be in either the D- or L-configuration.

Suitable bitter compound additives include, but are not limited to, caffeine, quinine, urea, bitter orange oil, naringin, quassia, and salts thereof.

Suitable flavorants and flavoring ingredient additives include, but are not limited to, vanillin, vanilla extract, mango extract, cinnamon, citrus, coconut, ginger, viridiflorol, almond, menthol (including menthol without mint), grape skin extract, and grape seed extract. "Flavorant" and "flavoring ingredient" are synonymous and can include natural or synthetic substances or combinations thereof. Flavorants also include any other substance which imparts flavor and may include natural or non-natural (synthetic) substances which are safe for human or animals when used in a generally accepted range. Non-limiting examples of proprietary flavorants include Döhler™ Natural Flavoring Sweetness Enhancer K14323 (Döhler™, Darmstadt, Germany), Symrise™ Natural Flavor Mask for Sweeteners 161453 and 164126 (Symrise™, Holzminden, Germany), Natural Advantage™ Bitterness Blockers 1, 2, 9 and 10 (Natural Advantage™, Freehold, New Jersey, U.S.A.), and Sucramask™ (Creative Research Management, Stockton, California, U.S.A.).

Suitable polymer additives include, but are not limited to, chitosan, pectin, pectic, pectinic, polyuronic, polygalacturonic acid, starch, food hydrocolloid or crude extracts thereof (e.g., gum acacia senegal (Fibergum™), gum acacia seyal, carageenan), poly-L-lysine (e.g., poly-L- α -lysine or poly-L- ϵ -lysine), poly-L-ornithine (e.g., poly-L- α -ornithine or poly-L- ϵ -ornithine), polypropylene glycol, polyethylene glycol, poly(ethylene glycol methyl ether), polyarginine, polyaspartic acid, polyglutamic acid, polyethylene imine, alginic acid, sodium alginate, propylene glycol alginate, and sodium polyethyleneglycolalginate, sodium hexametaphosphate and its salts, and other cationic polymers and anionic polymers.

Suitable protein or protein hydrolysate additives include, but are not limited to, bovine serum albumin (BSA), whey protein (including fractions or concentrates thereof such as 90% instant whey protein isolate, 34% whey protein, 50% hydrolyzed whey protein, and 80% whey protein concentrate), soluble rice protein, soy protein, protein isolates, protein hydrolysates, reaction products of protein hydrolysates, glycoproteins, and/or proteoglycans containing amino acids (e.g., glycine, alanine, serine, threonine, asparagine, glutamine, arginine, valine, isoleucine, leucine, norvaline, methionine, proline, tyrosine, hydroxyproline, and the like),

collagen (e.g., gelatin), partially hydrolyzed collagen (e.g., hydrolyzed fish collagen), and collagen hydrolysates (e.g., porcine collagen hydrolysate).

Suitable surfactant additives include, but are not limited to, polysorbates (e.g., polyoxyethylene sorbitan monooleate (polysorbate 80), polysorbate 20, polysorbate 60), sodium dodecylbenzenesulfonate, dioctyl sulfosuccinate or dioctyl sulfosuccinate sodium, sodium dodecyl sulfate, cetylpyridinium chloride (hexadecylpyridinium chloride), hexadecyltrimethylammonium bromide, sodium cholate, carbamoyl, choline chloride, sodium glycocholate, sodium taurodeoxycholate, lauric arginate, sodium stearyl lactylate, sodium taurocholate, lecithins, sucrose oleate esters, sucrose stearate esters, sucrose palmitate esters, sucrose laurate esters, and other emulsifiers, and the like.

Suitable flavonoid additives are classified as flavonols, flavones, flavanones, flavan-3-ols, isoflavones, or anthocyanidins. Non-limiting examples of flavonoid additives include, but are not limited to, catechins (e.g., green tea extracts such as Polyphenon™ 60, Polyphenon™ 30, and Polyphenon™ 25 (Mitsui Norin Co., Ltd., Japan), polyphenols, rutins (e.g., enzyme modified rutin Sanmelin™ AO (San-fi Gen F.F.I., Inc., Osaka, Japan)), neohesperidin, naringin, neohesperidin dihydrochalcone, and the like.

Suitable alcohol additives include, but are not limited to, ethanol.

Suitable astringent compound additives include, but are not limited to, tannic acid, europium chloride (EuCl_3), gadolinium chloride (GdCl_3), terbium chloride (TbCl_3), alum, tannic acid, and polyphenols (e.g., tea polyphenols).

III. Methods

Methods of preparing the tea beverages of the present invention are also provided.

In one embodiment, a method of preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining. All of the components and their respective amounts in the tea beverage are described above.

In a more particular embodiment, a method of preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener selected from the group consisting of rebaudioside A, rebaudioside M, rebaudioside AM, rebaudioside E, rebaudioside N, rebaudioside O, rebaudioside B, monk fruit juice or concentrate, monk fruit extract, siamenoside I, mogroside V, thaumatin (and variants thereof), brazzein (and variants thereof), monellin (and variants thereof), sweet truffle protein (and variants thereof), sucralose, potassium acesulfame, aspartame, neotame, saccharin, cyclamate, and combinations thereof; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining. All of the components and their respective amounts in the tea beverage are described above.

In a more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising rebaudioside M; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining. All of the components and their respective amounts in the tea beverage are described above.

In an even more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium

citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining. All of the components and their respective amounts in the tea beverage are described above.

In an even more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis; and (iii) a salt selected from (a) sodium gluconate and (b) potassium gluconate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining. All of the components and their respective amounts in the tea beverage are described above.

In another embodiment, a method for preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising rebaudioside M and monk fruit extract; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining. All of the components and their respective amounts in the tea beverage are described above.

In a more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener

comprising (a) a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis and (b) monk fruit extract; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining. All of the components and their respective amounts in the tea beverage are described above.

In a still more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one brewed tea selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising (a) a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis and (b) monk fruit extract; and (iii) at least one salt selected from (a) sodium gluconate and (b) a mixture of potassium citrate and potassium lactate to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii). The method can optionally include a step of diluting the brewed tea with a volume of water prior to combining. All of the components and their respective amounts in the tea beverage are described above.

In a more particular embodiment, a method of preparing a tea beverage comprises combining (i) at least one tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener selected from the group consisting of rebaudioside A, rebaudioside M, rebaudioside AM, rebaudioside E, rebaudioside N, rebaudioside O, rebaudioside B, monk fruit juice or concentrate, monk fruit extract, siamenoside I, mogroside V, thaumatin (and variants thereof), brazzein (and variants thereof), monellin (and variants thereof), sweet truffle protein (and variants thereof), sucralose, potassium acesulfame, aspartame, neotame, saccharin, cyclamate, and combinations thereof; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate with water to provide a tea beverage. The method

optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii) and the water. All of the components and their respective amounts in the tea beverage are described above.

In a more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising rebaudioside M; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate with water to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii) and the water. All of the components and their respective amounts in the tea beverage are described above.

In an even more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate with water to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii) and the water. All of the components and their respective amounts in the tea beverage are described above.

In an even more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis; and (iii) a salt selected from (a) sodium gluconate and (b) potassium gluconate

with water to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii) and the water. All of the components and their respective amounts in the tea beverage are described above.

In another embodiment, a method for preparing a tea beverage comprises combining (i) at least one tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising rebaudioside M and monk fruit extract; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate with water to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii) and the water. All of the components and their respective amounts in the tea beverage are described above.

In a more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising (a) a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on a dry basis and (b) monk fruit extract; and (iii) at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate with water to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii) and the water. All of the components and their respective amounts in the tea beverage are described above.

In a still more particular embodiment, a method for preparing a tea beverage comprises combining (i) at least one tea extract, wherein the tea is selected from the group consisting of white, yellow, green, oolong, black, red, and a combination thereof; (ii) at least one high potency sweetener comprising (a) a steviol glycoside mixture comprising at least about 80% rebaudioside M by weight on a dry basis, or at least about 95% rebaudioside M by weight on

a dry basis and (b) monk fruit extract; and (iii) at least one salt selected from (a) sodium gluconate and (b) a mixture of potassium citrate and potassium lactate with water to provide a tea beverage. The method optionally comprises combining (iv) at least one caloric sweetener, (v) at least one natural flavor, (vi) at least one fruit juice, (vii) at least one fruit extract, (viii) at least one herbal extract, (ix) at least one herbal tea, (x) at least one organic acid, (xi) at least one functional ingredient, and/or (xii) at least one additive with (i)-(iii) and the water. All of the components and their respective amounts in the tea beverage are described above.

Methods of preparing tea beverages are known in the art. For example, once all of the ingredients are combined, the tea beverage is pasteurized then cooled. The pasteurized tea beverage can be packaged, e.g., hot-filled into bottles prior to cooling. The cooled beverages can then be stored for distribution or immediate consumption.

EXAMPLES

Materials and Methods

Commercial natural sweeteners rebaudioside-M (95% purity), Monk fruit extract (50% Mogroside V) were used to sweeten the tea base. Food grade salts with high purity (>95%) were acquired from different suppliers.

Table 1. Compounds Used and Maximum Levels in Beverages

Ingredients	MAX USED (ppm)
Tri-calcium citrate tetrahydrate	70
Calcium lactate pentahydrate	70
Magnesium lactate dihydrate	200
Sodium gluconate	230
Tri-potassium citrate	140
Potassium lactate	140
Potassium gluconate	230

Tea Beverages

The ingredients were dissolved in filtered water to complete dissolution. The beverages were pasteurized (89-91°C), hot-filled in 300 ml PET bottles, then cooled in ice water. Beverages were stored in 4 °C and tasted cold. Beverage brix was 0.4° and titratable acidity was 0.189% w/v as citric acid. Table 2 shows the ingredients list for the control samples (Control 1 with 472 ppm of rebaudioside M; Control 2 with 425 ppm of rebaudioside M and 50 ppm of Monk fruit extract) without salts. For test samples 1-14, different salts were added on top of each control sample in the amount (ppm) as shown in Tables 3 and 4.

Table 2. Tea Beverage Composition

Ingredients	Control 1 Rebaudioside M (grams)	Control 2 Rebaudioside M + Monk fruit extract (grams)
Filtered water	99.565	99.565
Black tea powder	0.09	0.09
Citric acid	0.189	0.189
Sodium citrate	0.0525	0.0525
Rebaudioside-M 95%	0.0472	0.0425
Monk fruit extract	-	0.005
Lemon flavor	0.0563	0.0563
TOTAL (grams)	100	100

Table 3. Organic Salt Combinations – Lower Levels (Test Samples 1-7)

Ingredients	Test 1 (ppm)	Test 2 (ppm)	Test 3 (ppm)	Test 4 (ppm)	Test 5 (ppm)	Test 6 (ppm)	Test 7 (ppm)
Potassium citrate	70		35				
Potassium lactate		70	35				
Sodium gluconate				115			
Calcium citrate					35	35	
Calcium lactate					35		
Magnesium lactate						100	
Potassium gluconate							115

Table 4. Organic Salt Combinations – Higher Levels (Test Samples 8-14)

Ingredients	Test 8 (ppm)	Test 9 (ppm)	Test 10 (ppm)	Test 11 (ppm)	Test 12 (ppm)	Test 13 (ppm)	Test 14 (ppm)
Potassium citrate	140		70				
Potassium lactate		140	70				
Sodium gluconate				230			
Calcium citrate					70	70	
Calcium lactate					70		

Magnesium lactate						200	
Potassium gluconate							230

Sensory Evaluation

The beverages were evaluated blindly by six expert panelists. Samples were coded and randomly presented to the panelists. Panelists were instructed to eat an unsalted cracker and rinse the mouth with water before and in between samples. The maximum samples for each session was set at 6 samples to avoid fatigue. For each sample, panelists were instructed to take 3 sips, then write down their evaluation comments.

Sensory Results

From above samples, Test 1, 2, 3, 4 and 7 were much preferred by panelists who thought that there was significant taste improvement. Test 5 and 6 were least preferred by panelists and thought to be close to control sample.

Table 5. Tea Beverages with Rebaudioside-M – Low Level of Organic Salts

Beverages with Salts	Panelists Comments
Control 1 (no added organic salts)	More sweetness lingering, more licorice aftertaste
Test 1	Sweeter compared to control, cleaner tea flavor, no sweet lingering aftertaste
Test 2	Sweeter compared to control, cleaner tea flavor, no sweet lingering aftertaste, nice flavor release
Test 3	2 nd choice, preferred over Test 1 and Test 2 samples, sweeter than control, more rounded flavor and flavor release
Test 4	Best sample in this set, clean flavor, sweeter than control sample, nice tea flavor, no off-notes
Test 5	Some sweetness lingering at end, not different from control sample
Test 6	Some sweetness lingering at end comparable to Test sample 5, not different from control sample
Test 7	3 rd choice, sweeter than control, balanced flavor, well-liked by panelists

Table 6. Tea Beverages with Rebaudioside-M + Monk Fruit – Lower Organic Salts

Beverages with Salts	Panelists Comments
Control 1 (no added organic salts)	More sweetness lingering, more bitter and licorice aftertaste, less orange flavor
Test 1	Less lingering, more tea and lemon flavor than control, sweetness is more rounded.
Test 2	Flavor is more enhanced, and linger is reduced in comparison to control, compared to sample 1 there is slight linger but sample is still good.
Test 3 – K citrate + K lactate	Best sample in this set, flavor is enhanced with no sweetness lingering, mouthfeel isn't chalky or powdery like control, combination of these two salts works better than individual salts
Test 4 – Na gluconate	2 nd most liked sample in set, flavor is enhanced more than Test 3, linger is cut and mouthfeel is good.
Test 5	Slight sweetness lingering and slight powdery mouthfeel, taste improvement from control but not as good as other salts.
Test 6	Sample is an improvement from control, slight linger but better than control, flavor is slightly suppressed and not enhanced like other samples, has slight chalky powdery mouthfeel
Test 7	Sample is least preferred among test samples, sweetness is improved but flavor is muted, sample tastes better than control

Table 7. Tea Beverages with Rebaudioside-M – Higher Organic Salts

Beverages with Salts	Panelists Comments
Control 2 (no added organic salts)	More sweetness lingering, more bitter and licorice aftertaste, less orange flavor
Test 1	Noticeable difference from control, less powdery and astringent, mouth feel is much cleaner, lingering is suppressed, flavor is enhanced
Test 2	Much less sweetness lingering compared to control, cleaner flavor
Test 3	Preferred over Test 1 and Test 2 samples.

	Good balance between sweetness and lemon tea, linger is cut, and isn't as powdery as control
Test 4- Na gluconate	Favorite sample in this set, flavor is enhanced and gives good balance with sweetness, no astringent taste, major taste improvement from control
Test 5	Good sample, better taste compared to control, but not as much of an improvement compared to other test samples, linger is cut and flavor is enhanced.
Test 6	Sweeter than control, less aftertaste compared to control
Test 7- K gluconate	2 nd sample liked after Test 4. Flavor is enhanced and the sweetness is much more rounded with good balance, no powdery or astringent aftertaste

Table 8. Tea Beverages with Rebaudioside-M + Monk Fruit – Higher Organic Salts

Beverages with Salts	Panelists Comments
Control 2 (no added organic salts)	More sweetness lingering, more bitter and licorice aftertaste, less orange flavor
Test 1	Sample shows significant taste improvement from control, sweetness is much more rounded and sweeter than control, no linger or powdery chalky mouthfeel, flavor is enhanced more tea notes.
Test 2	Sample is improvement from control, much less linger but is slightly chalky and powdery compared to first sample, there is more tea flavor and lemon notes.
Test 3 – K citrate + K lactate	Best sample in set, combination of these salts works much better than individual salts, no linger, astringent or powdery notes and mouthfeel is much improved, tea flavor is enhanced.
Test 4 – Na gluconate	2 nd most liked sample in set, no linger and powdery or astringent notes, more desired tea flavor notes, significant improvement from control.
Test 5	Sample is improvement from control but is not as preferred as other samples, slight notes of saltiness, flavor is enhanced
Test 6	Significant sweet linger reduction compared to control, nice and enhanced tea flavor

Test 7	Significant sweet linger reduction compared to control, nice and enhanced tea notes, no chalky or astringent notes from control
--------	---

Overall, organic salts significantly improved the taste (reduced astringency, more mouthfeel, more rounded flavor) with more impact from higher salt concentrations, and panelists preferred this set over the set with lower organic salt concentrations.

CLAIMS

1. A tea beverage comprising:
 - i. at least one tea product;
 - ii. at least one high potency sweetener selected from the group consisting of rebaudioside A, rebaudioside M, rebaudioside AM, rebaudioside E, rebaudioside N, rebaudioside O, rebaudioside B, monk fruit juice or concentrate, monk fruit extract, siamenoside I, mogroside V, thaumatin (and variants thereof), brazzein (and variants thereof), monellin (and variants thereof), sweet truffle protein (and variants thereof), sucralose, potassium acesulfame, aspartame, neotame, saccharin, cyclamate, and combinations thereof; and
 - iii. at least one salt selected from the group consisting of sodium gluconate, sodium citrate, sodium lactate, potassium gluconate, potassium citrate, and potassium lactate.
2. The tea beverage of claim 1, wherein the at least one tea product is selected from a brewed tea and a tea extract, wherein the tea of the brewed tea and the tea extract is selected from the group consisting of white, yellow, green, oolong, black, red, post-fermented, or a combination thereof.
3. The tea beverage of claim 2, wherein the at least one tea product is brewed black tea or a black tea extract.
4. The tea beverage of claim 2, wherein the at least one tea product is a tea extract and the tea beverage further comprises water in an amount of at least about 95% by weight, or at least about 99% by weight.
5. The tea beverage of claim 1, wherein the concentration of the at least one high potency sweetener is from about 1 ppm to about 600 ppm.
6. The tea beverage of claim 5, wherein the at least one high potency sweetener comprises rebaudioside M.
7. The tea beverage of claim 6, wherein the concentration of the at least one high potency sweetener is from about 25 to about 600 ppm.
8. The tea beverage of claim 6, wherein the high potency sweetener further comprises monk fruit extract comprising about 45% or greater mogroside V by weight.

9. The tea beverage of claim 8, wherein the concentration of the monk fruit extract is from about 25 ppm to about 100 ppm.
10. The tea beverage of claim 1, wherein the at least one salt is present in a concentration from about 100 ppm to about 300 ppm.
11. The tea beverage of claim 6, wherein the at least one salt is potassium gluconate or sodium gluconate in a concentration from about 100 ppm to about 300 ppm
12. The tea beverage of claim 9, wherein the at least one salt is (a) sodium gluconate or (b) a mixture of potassium citrate and potassium lactate in a concentration from about 100 ppm to about 300 ppm.
13. The tea beverage of claim 10, wherein the at least one salt is (a) sodium gluconate or (b) a mixture of potassium citrate and potassium lactate in a concentration from about 100 ppm to about 300 ppm.
14. The tea beverage of claim 1, wherein the tea beverage does not contain potassium, sodium, calcium, or magnesium salts of inorganic ions and/or does not contain magnesium or calcium salts of gluconate, citrate or lactate.
15. The tea beverage of claim 1, wherein the tea beverage has from about 0 to about 40 calories.
16. The tea beverage of claim 1, wherein the tea beverage does not contain added sugar.
17. The tea beverage of claim 1, wherein the tea beverage further comprises at least one caloric sweetener.
18. The tea beverage of claim 1, further comprising at least one caloric sweetener, at least one natural flavor, at least one fruit juice, at least one fruit extract, at least one herbal extract, at least one herbal tea, at least one organic acid, or combinations thereof.
19. The tea beverage of claim 3, wherein the at least one salt is potassium gluconate or sodium gluconate.
20. The tea beverage of claim 3, wherein the at least one salt is (a) sodium gluconate or (b) a mixture of potassium citrate and potassium lactate.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2023/066583

A. CLASSIFICATION OF SUBJECT MATTER

A23F 3/14(2006.01)i; A23F 3/16(2006.01)i; A23F 3/40(2006.01)i; A23L 2/60(2006.01)i; A23L 2/38(2006.01)i;
A23L 2/58(2006.01)i

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A23F 3/14(2006.01); A23F 3/16(2006.01); A23F 3/40(2006.01); A23L 1/221(2006.01); A23L 2/00(2006.01);
A23L 2/60(2006.01); A23L 27/30(2016.01); A23L 29/30(2016.01); C07K 14/43(2006.01)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models
Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: tea beverage, high potency sweetener, salt, black tea, Camellia sinensis, rebaudioside M, monk fruit extract, sodium gluconate, potassium gluconate, potassium citrate, potassium lactate, calorie, sugar-free beverage

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2022-0015384 A1 (SUNTORY HOLDINGS LIMITED) 20 January 2022 (2022-01-20) claims 1, 8-11; paragraphs [0038]-[0057], [0064], [0075], [0084], [0092], [0094], [0184], [0203], [0289]-[0293], [0329], [0335]; tables 10, 12	1-20
A	US 2019-0183148 A1 (SUNTORY HOLDINGS LIMITED) 20 June 2019 (2019-06-20) paragraphs [0020], [0032], [0038]; table 2	1-20
A	WO 2020-205978 A1 (THE COCA-COLA COMPANY) 08 October 2020 (2020-10-08) claims 1, 2, 5, 6, 13, 15; pages 55, 56	1-20
A	US 2019-0216119 A1 (MERISANT US, INC.) 18 July 2019 (2019-07-18) paragraphs [0002], [0024], [0044], [0063]	1-20
A	US 2008-0226776 A1 (ROY, GLENN et al.) 18 September 2008 (2008-09-18) claims 1, 5, 7, 9; paragraphs [0030], [0049]	1-20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

“A” document defining the general state of the art which is not considered to be of particular relevance
“D” document cited by the applicant in the international application
“E” earlier application or patent but published on or after the international filing date
“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
“O” document referring to an oral disclosure, use, exhibition or other means
“P” document published prior to the international filing date but later than the priority date claimed

“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

“&” document member of the same patent family

Date of the actual completion of the international search

28 August 2023

Date of mailing of the international search report

28 August 2023

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
189 Cheongsa-ro, Seo-gu, Daejeon
35208, Republic of Korea

Facsimile No. +82-42-481-8578

Authorized officer

HEO, Joo Hyung

Telephone No. +82-42-481-5373

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/US2023/066583

Patent document cited in search report			Publication date (day/month/year)	Patent family member(s)			Publication date (day/month/year)
US	2022-0015384	A1	20 January 2022	AU	2019-393558	A1	11 June 2020
				AU	2019-393558	A1	01 July 2021
				CN	113163784	A	23 July 2021
				EP	3892102	A1	13 October 2021
				SG	11202105908	A	29 July 2021
				WO	2020-116628	A1	11 June 2020
US	2019-0183148	A1	20 June 2019	AU	2017-239890	A1	08 November 2018
				AU	2017-239890	A1	05 October 2017
				AU	2017-239890	B2	03 October 2019
				AU	2019-284092	A1	23 January 2020
				AU	2019-284092	B2	08 April 2021
				CN	109152392	A	04 January 2019
				EP	3437484	A1	06 February 2019
				JP	6257862	B1	10 January 2018
				MY	190325	A	14 April 2022
				NZ	747260	A	31 January 2020
				SG	11201808405	A	30 October 2018
				US	10750764	B2	25 August 2020
				US	11122824	B2	21 September 2021
				US	2020-0345041	A1	05 November 2020
				WO	2017-170990	A1	05 October 2017
WO	2020-205978	A1	08 October 2020	AU	2020-253391	A1	28 October 2021
				AU	2020-253391	A1	08 October 2020
				BR	112021019734	A2	08 March 2022
				CA	3135373	A1	08 October 2020
				CN	113939197	A	14 January 2022
				EP	3945859	A1	09 February 2022
				IL	286833	A	31 October 2021
				IL	286833	D0	31 October 2021
				JP	2022-519941	A	25 March 2022
				KR	10-2021-0134813	A	10 November 2021
				MX	2021011908	A	15 December 2021
				SG	11202110820	A	28 October 2021
				US	2022-361544	A1	17 November 2022
US	2019-0216119	A1	18 July 2019	WO	2019-143916	A1	25 July 2019
US	2008-0226776	A1	18 September 2008	AR	065755	A1	01 July 2009
				BR	PI0803108	A2	30 August 2011
				CA	2656306	A1	18 September 2008
				CN	101668435	A	10 March 2010
				EP	2120606	A1	25 November 2009
				EP	2489282	A2	22 August 2012
				EP	2489282	A3	05 December 2012
				JP	2010-521178	A	24 June 2010
				JP	2012-143255	A	02 August 2012
				MX	2009009722	A	07 October 2009
				RU	2008151094	A	27 June 2010
				RU	2424733	C2	27 July 2011
				WO	2008-112979	A1	18 September 2008