



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/12/18
(87) Date publication PCT/PCT Publication Date: 2019/06/27
(85) Entrée phase nationale/National Entry: 2020/06/18
(86) N° demande PCT/PCT Application No.: KR 2018/016166
(87) N° publication PCT/PCT Publication No.: 2019/124951
(30) Priorités/Priorities: 2017/12/19 (US62/607,876);
2018/06/01 (US62/679,598); 2018/09/18 (US62/732,946)

(51) Cl.Int./Int.Cl. *A61K 31/404* (2006.01),
A61K 31/222 (2006.01), *A61K 31/439* (2006.01),
A61K 31/506 (2006.01), *A61K 45/06* (2006.01),
A61P 35/00 (2006.01), *G01N 33/574* (2006.01),
G01N 33/68 (2006.01)

(71) Demandeur/Applicant:
GPCR THERAPEUTICS, INC., KR

(72) Inventeurs/Inventors:
SEEN, DONGSEUNG, KR;
KIM, EUNHEE, KR;
JEONG, JAE-YEON, KR;
YANG, CHANG SOO, KR;
LEE, MILIM, KR;

(54) Titre : INHIBITEURS D'HETEROMERES GPCR ET LEURS UTILISATIONS
(54) Title: GPCR HETEROMER INHIBITORS AND USES THEREOF

(57) **Abrégé/Abstract:**

This invention relates to inhibitors of CXCR4-G protein-coupled receptor (GPCR) heteromers (CXCR4-GPCR heteromers) associated with cancers, where CXCR4 forms a functional heteromer with other G protein-coupled receptors (GPCRx). More specifically, this invention relates to GPCRx that form heteromers with CXCR4, which upon co-stimulation with CXCR4 agonists and GPCRx agonists leads to enhanced signaling downstream of CXCR4. This invention also provides for the use of inhibitors of the interacting GPCR partner of the CXCR4-GPCRx heteromer or CXCR4-GPCRx heteromer-specific inhibitors including inhibitors of the formation of the CXCR4-GPCRx heteromer and CXCR4-GPCRx heteromer-specific antibodies, and in the diagnosis and/or therapy for cancer.

(72) **Inventeurs(suite)/Inventors(continued):** KIM, SOHUI, KR; HUH, WON-KI, KR; SONG, YONG BHUM, KR;
PARK, CHUL O, KR; PARK, HYERYUNG, KR; LEE, JI YEONG, KR

(74) **Agent:** TORYS LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
27 June 2019 (27.06.2019)(10) International Publication Number
WO 2019/124951 A9

(51) International Patent Classification:

A61K 31/404 (2006.01) *A61K 45/06* (2006.01)
A61K 31/439 (2006.01) *A61P 35/00* (2006.01)
A61K 31/222 (2006.01) *G01N 33/574* (2006.01)
A61K 31/506 (2006.01) *G01N 33/68* (2006.01)

(21) International Application Number:

PCT/KR2018/016166

(22) International Filing Date:

18 December 2018 (18.12.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/607,876 19 December 2017 (19.12.2017) US
 62/679,598 01 June 2018 (01.06.2018) US
 62/732,946 18 September 2018 (18.09.2018) US

(71) Applicant: **GPCR THERAPEUTICS, INC.** [KR/KR];
 Room 407, Nakseongdae R & D Center, 38, Nakseong-
 dae-ro, Gwanak-gu, Seoul 08790 (KR).

(72) Inventors: **SEEN, Dong Seung**; 107-3502, Banpo Riche
 Apt., 35, Gomurae-ro, Seocho-gu, Seoul 06597 (KR). **KIM,
 Eun Hee**; 227-2201, Park Rio Apt., 435, Olympic-ro, Song-
 pa-gu, Seoul 05507 (KR). **JEONG, Jae Yeon**; 101-1005,
 Sangdo Emcotown Central Park Apt., 346-1, Sangdo-ro,
 Dongjak-gu, Seoul 07039 (KR). **YANG, Chang Soo**;
 101, 13-38, Sillim-ro 7ga-gil, Gwanak-gu, Seoul 08815
 (KR). **LEE, Mi Lim**; 101, 67, Bogeun-gil, Gwanak-gu,
 Seoul 08835 (KR). **KIM, So Hui**; 201, 20, Janggunbong
 7-gil, Gwanak-gu, Seoul 08784 (KR). **HUH, Won Ki**;
 101-1005, Sangdo Emcotown Central Park Apt., 346-1,
 Sangdo-ro, Dongjak-gu, Seoul 07039 (KR). **SONG, Yong
 Bhum**; 1008-602, Daedeok Techno Valley 10 Apt., 61
 Baeul, 2-ro, Yuseong-gu, Daejeon 34020 (KR). **PARK,
 Chul O**; 107, Sejong Minitel, 72 Bogeun 6-gil, Gwanak-
 gu, Seoul 08840 (KR). **PARK, Hye Ryung**; 43 Gin-
 gorang-ro 4-gil, Gwangjin-gu, Seoul 04916 (KR). **LEE,
 Ji Yeong**; 23-35, Nambusunhwan-ro 226-gil, Gwanak-gu,
 Seoul 08788 (KR).

(74) Agent: **LEE & KO IP**; 25F, 63, Namdaemun-ro, Jung-gu,
 Seoul 04532 (KR).

(81) Designated States (*unless otherwise indicated, for every
 kind of national protection available*): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
 CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
 DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
 HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
 KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
 OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
 SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
 TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
 kind of regional protection available*): ARIPO (BW, GH,
 GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
 UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
 TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
 MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
 TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
 KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))
- with information concerning authorization of rectification
 of an obvious mistake under Rule 91.3 (b) (Rule 48.2(i))

(48) Date of publication of this corrected version:

07 November 2019 (07.11.2019)

(15) Information about Correction:

see Notice of 07 November 2019 (07.11.2019)

(54) Title: GPCR HETEROMER INHIBITORS AND USES THEREOF

(57) Abstract: This invention relates to inhibitors of CXCR4-G protein-coupled receptor (GPCR) heteromers (CXCR4-GPCR heteromers) associated with cancers, where CXCR4 forms a functional heteromer with other G protein-coupled receptors (GPCRx). More specifically, this invention relates to GPCR_x that form heteromers with CXCR4, which upon co-stimulation with CXCR4 agonists and GPCR_x agonists leads to enhanced signaling downstream of CXCR4. This invention also provides for the use of inhibitors of the interacting GPCR partner of the CXCR4-GPCR_x heteromer or CXCR4-GPCR_x heteromer-specific inhibitors including inhibitors of the formation of the CXCR4-GPCR_x heteromer and CXCR4-GPCR_x heteromer-specific antibodies, and in the diagnosis and/or therapy for cancer.



WO 2019/124951 A9

Description

Title of Invention: GPCR HETEROMER INHIBITORS AND USES THEREOF

Technical Field

- [1] This application claims the benefit of priority from U.S. Provisional Application No. 62/607,876, filed December 19, 2017, and further claims the benefit of priority from U.S. Provisional Application No. 62/679,598, filed June 1, 2018, and further claims the benefit of priority from U.S. Provisional Application No. 62/732,946, filed September 18, 2018. Each of the foregoing related applications, in their entirety, are incorporated herein by reference.
- [2] In addition, each of the references identified herein, in their entirety, are incorporated herein by reference.

Background Art

- [3] The invention disclosed herein relates generally to inhibitors of novel, functional GPCR heteromers, and more specifically to inhibitors of CXC receptor 4 (CXCR4)-G protein-coupled receptor (GPCR) heteromers that display enhanced signaling downstream of CXCR4 as a result of functional heteromer formation and that are associated with cancers and other diseases.
- [4] G protein-coupled receptors (GPCRs) are seven-transmembrane domain cell surface receptors that are coupled to G proteins. GPCRs mediate diverse sensory and physiological responses by perceiving stimuli including light, odorants, hormones, neurotransmitters, chemokines, small lipid molecules, and nucleotides. There are approximately 800 GPCR genes in human genome, and more than half of them are predicted to encode sensory receptors such as olfactory, visual, and taste receptors (Bjarnadottir, et al., 2006). The remaining 350 GPCRs have physiologically important roles in embryonic development, behavior, mood, cognition, regulation of blood pressure, heart rate, and digestive processes, regulation of immune system and inflammation, maintenance of homeostasis, and growth and metastasis of cancers (Filmore 2004, Overington, et al., 2006). GPCRs are associated with many diseases and are the targets of approximately 40% of all prescription drugs (Filmore 2004).
- [5] CXC receptor 4 (CXCR4) is a member of the chemokine receptor family GPCR. CXCR4 is expressed on most of the hematopoietic cell types, bone marrow stem cells, endothelial progenitor cells, vascular endothelial cells, neurons and neuronal stem cells, microglia and astrocytes (Klein and Rubin 2004, Griffith, et al., 2014). CXCR4 responds to its ligand C-X-C Motif Chemokine ligand 12 (CXCL12), also known as Stromal cell-derived factor 1 (SDF-1), and has essential roles in the embryonic de-

velopment of the hematopoietic, cardiovascular, and nervous systems (Griffith, et al., 2014). CXCR4 was discovered as a co-receptor for human immunodeficiency virus (HIV), and has important roles in the homing of hematopoietic stem cells (HSCs) to the bone marrow, inflammation, immune surveillance of tissues, and tissue re-generation in adult (Chatterjee, et al. 2014). Mutations in the C-terminus of CXCR4 cause persistent CXCR4 activation, leading to a congenital immune deficiency called WHIM syndrome (Warts, Hypogammaglobulinemia, Infections, and Myelokathexis) characterized by neutropenia and B cell lymphopenia (Hernandez, et al., 2003; Kawai, 2009). CXCR4 also has essential roles in T and B lymphocyte development within lymphoid organs and the thymus during development and in adult (Allen, et al., 2004; Ara, et al., 2003).

- [6] CXCR4 is implicated in various immune and autoimmune diseases, such as HIV infection, ischaemia, wound healing, rheumatoid arthritis, systemic lupus erythematosus (SLE), interstitial pneumonias, vascular disease, multiple sclerosis, pulmonary fibrosis, and allergic airway disease (Chu et al., 2017; Debnath, et al., 2013; Domanska, et al., 2013). The involvement of CXCR4 in rheumatoid arthritis was demonstrated by the increased accumulation of CXCR4-positive T-cells in arthritic joints, and a reduction in collagen-induced arthritis in CXCR4-deficient mice (Buckley et al., 2000; Chung et al., 2010). Moreover, CXCR4 antagonist, AMD3100, alleviated collagen-induced arthritis significantly in a mouse model (De Klerck et al., 2005). CXCR4 also regulates pulmonary fibrosis by recruiting circulatory fibroblasts and bone marrow-derived progenitor cells during lung injury, and AMD3100 demonstrated a preventive effect in bleomycin-induced mouse pulmonary fibrosis (Song et al., 2010). CXCR4/CXCL12 axis is also implicated in the pathogenesis of SLE. In mouse models of lupus and patients with SLE, inflammatory cells such as monocytes, neutrophils, and B-cells showed increased expression of CXCR4 and migrated toward skin and lung that overexpress CXCL12 predominantly (Chong and Mohan, 2009; Wang et al., 2009; Wang et al., 2010). CXCR4 antagonist, CTCE-9908, prolonged survival and greatly improved disease conditions and nephritis in a mouse model of lupus (Wang et al., 2009). Furthermore, CXCR4 is also involved in brain and cardiac diseases including brain injury, stroke, myocardial infarction, atherosclerosis and injury-induced vascular restenosis (Cheng et al., 2017; Domanska, et al., 2013; Doring, et al., 2014).

- [7] The involvement of CXCR4 in cancer was first noticed when B cells from patients with chronic lymphocytic leukemia (B-CCL) express high levels of functional CXCR4 on the surface, showing enhanced calcium mobilization and actin polymerization upon CXCL12 exposure, and migration towards bone marrow stromal cells that secrete CXCL12 (Burger, et al., 1999).

- [8] CXCR4 is subsequently characterized to be responsible for breast cancer metastasis to organs that express higher levels of CXCL12 such as lymph nodes, bone marrow, lung, and liver (Muller, et al., 2001). After initial discovery, increasing evidence indicates that CXCR4 is associated with a variety of different cancers and has multiple potential roles in malignancy. CXCR4 is overexpressed in more than 23 human cancers, including breast cancer, lung cancer, brain cancer, kidney cancer (or renal cell carcinoma), pancreatic cancer, ovarian cancer, prostate cancer, melanoma, leukemia, multiple myeloma, gastrointestinal cancers, and soft tissue sarcomas, and regarded as a poor prognosis marker (Domanska, et al., 2013; Chatterjee, et al., 2014; Furusato et al., 2010). CXCR4 is the only chemokine receptor that is expressed by the majority of cancer types (Liang et al., 2015), and stimulates tumor cell growth, survival, and invasiveness in response to CXCL12 secreted by the cancer cells and the surrounding cancer-associated cells (Burger and Kipps, 2006; Chatterjee et al., 2014; Domanska et al., 2013).
- [9] Systematic meta-analyses using databases including PubMed, EMBASE, and Cochrane library indicate significant association between CXCR4 over-expression and poorer progression-free survival and overall survival in various cancers including hematological malignancy, breast cancer, colorectal cancer, esophageal cancer, head and neck cancer, renal cancer, lung cancer, gynecologic cancer, liver cancer, prostate cancer, and gallbladder cancer (Du et al., 2015; Hu et al., 2015; Li et al., 2017; Wang et al., 2016; Zhao et al., 2015).
- [10] CXCR4/CXCL12 axis plays a central role in tumor growth, invasion, angiogenesis, vasculogenesis, metastasis, drug resistance, and cancer cell-tumor microenvironment interaction (D'Alterio et al., 2012; Domanska et al., 2013; Guo et al., 2016)
- [11] The critical role of CXCR4 in cancer cell proliferation and tumor growth was demonstrated in various experimental models *in vitro* and *in vivo* such as orthotopic, subcutaneous human xenograft, and transgenic mouse models using CXCR4 antagonists (Domanska et al., 2013). The Daoy medulloblastoma cells and U87 glioblastoma cells showed CXCR4 expression and exhibited dose-dependent increase in proliferation to a gradient of CXCL12 *in vitro*, and systemic administration of AMD3100 inhibited the growth of intracranial U87 and Daoy cell xenografts (Rubin et al., 2003).
- [12] Involvement of CXCR4 in the metastasis of cancer cells towards CXCL12 expressing organs was also demonstrated in pancreatic, thyroid, melanoma, prostate, and colon cancer xenograft models (Bartolome et al., 2009; De Falco et al., 2007; Taichman et al., 2002; Wang et al., 2008; Zeelenberg et al., 2003).
- [13] The main mechanism of action described for the small molecules or peptides antagonists of CXCR4 is centered on their ability to mobilize malignant cells from the

BM, thereby sensitizing them to chemotherapy. These agents have shown limitations regarding short half-lives, making their adequate management over long periods of time difficult (Hendrix et al., 2000). In contrast, therapeutic monoclonal antibodies have the advantage of having more prolonged half-lives, and are suitable for less frequent dosing. Additionally, human IgG antibodies have the ability to induce cell death upon binding to their target protein on cancer cells, via interaction with Fc-receptor on effector cells, including antibody-dependent cell mediated cytotoxicity/phagocytosis (ADCC/ADCP) (Jiang et al., 2011). Such cytotoxic mechanism of action are not inherent to small molecules or peptides, and have been demonstrated to play a key role in the clinical activity of several therapeutic antibodies (Wang et al., 2015).

- [14] Targeting CXCR4 using a neutralizing anti-CXCR4 antibody or CXCR4 specific antagonists inhibited primary tumor growth as well as metastasis to secondary organs in breast cancer, colon cancer, hepatocellular carcinoma, osteosarcoma, and melanoma (De Falco et al., 2007; Hassan et al., 2011; Huang et al., 2009; Kim et al., 2008; Muller et al., 2001; Schimanski et al., 2006; Smith et al., 2004; Zeelenberg et al., 2003). In a transgenic breast cancer mouse model, inhibition of CXCR4 with CTCE-9908 reduced not only the growth of primary tumor but also the expression of vascular endothelial growth factor (VEGF) and AKT phosphorylation (Hassan et al., 2011).

- [15] Cancer stem cells (CSCs) are a population of cancer cells with properties such as infinite self-renewal, potential to differentiate into multiple cancer lineages, ability to adopt a quiescent state, and intrinsic high resistance to chemo- and radio-therapy. CSCs are considered as the major cause of cancer relapse and recurrence after standard anti-proliferative therapy. Therefore, targeting cancer stem cells are expected to provide more effective therapeutic interventions to eradicate the cancer and prevent relapse (Batlle and Clevers, 2017; Reya et al., 2001; Wurth, 2016). Interestingly, CSCs also express CXCR4, and CXCR4 directs the trafficking and metastasis of these cells to the CXCL12-rich microenvironments such as bone marrow and subventricular zone in the brain that favor cancer stem cell maintenance, survival and growth. CXCR4 antagonists have been shown to mobilize CSCs from these protective microenvironments, and sensitize them to conventional chemo- and radiotherapy, and anti-angiogenic therapy (Burger and Kipps, 2006; Burger and Peled, 2009; Furusato et al., 2010; Redondo-Munoz et al., 2006; Walenkamp et al., 2017; Wurth, 2016).

- [16] Increasing evidence showed that tumor mass contains various cell types such as stromal fibroblasts, immune cells, endothelial cells, connective tissue, and extracellular matrix in addition to cancer cells that constitute tumor microenvironment (TME) or cancer cell niches. CXCR4/CXCL12 axis plays pivotal roles in tumor cell-microenvironment interaction that support tumor structure, growth, angiogenesis, and evasion of immune surveillance in various cancers of both the hematopoietic and non-

hematopoietic system (Burger and Kipps, 2006; Burger and Peled, 2009; Walenkamp et al., 2017). CXCL12 can promote tumor angiogenesis by recruiting endothelial cells to the TME directly, or indirectly by attracting CXCR4-positive inflammatory cells to the tumor mass, and making them to secrete proangiogenic factors (Owen and Mohamadzadeh, 2013; Walenkamp et al., 2017).

- [17] Growing evidence indicates that CXCR4/CXCL12 axis contributes to the lack of tumor responsiveness to angiogenesis inhibitors. Vascular endothelial growth factor (VEGF) was considered as the major pro-angiogenic factor in cancer, and was targeted for anti-angiogenic therapy in patients with rectal carcinoma using an anti-VEGF antibody bevacizumab (Genentech). Surprisingly, bevacizumab increased the expression of CXCL12 and CXCR4 in cancer cells, and increased plasma levels of CXCL12 in these patients were associated with rapid disease progression and metastasis (Owen and Mohamadzadeh, 2013; Xu et al., 2009). Therefore, the efficacy of a combination therapy using bevacizumab and plerixafor was evaluated for recurrent glioma (ClinicalTrials.gov identifier: NCT01339039). However, the study was terminated due to a low accrual rate (Walenkamp et al., 2017).
- [18] Inhibition of CXCR4/CXCL12 axis has been demonstrated to disrupt the tumor microenvironment (TME) and expose the tumor cells to immune attack by decreasing the infiltration of myeloid-derived suppressor cells, by increasing the ratio of CD8+ cytotoxic T cells to Treg cells, or eliminating tumor re-vascularization (Burger et al., 2011; Domanska et al., 2013; Walenkamp et al., 2017). Administration of CXCR4 antagonists, AMD3100 or T22, acted synergistically with immune checkpoint inhibitors such as cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4) antibodies, programmed cell death protein 1 (PD-1), and programmed cell death ligand 1 (PD-L1) antibodies and greatly increased their antitumor activity in a model of pancreatic ductal adenocarcinoma, advanced hepatocellular carcinoma (HCC), and CXCR4-transduced B16 melanoma (Chen et al., 2015; Feig et al., 2013; Lee et al., 2006; Scala, 2015; Walenkamp et al., 2017). These results indicate that targeting CXCR4/CXCL12 axis offers benefit to the conventional immune checkpoint inhibitors.
- [19] Various drugs targeting CXCR4 have been developed (Peled, et al., 2012; Debnath, et al., 2013; Walenkamp, et al., 2017). CXCR4 inhibitors can be divided into 5 categories:
- [20] (1) non-peptide small molecule antagonists, such as AMD3100 (plerixafor, Mozobil™, Genzyme (MA, USA); (Cashen et al., 2007; Donzella et al., 1998), AMD070 (AMD11070; Crawford and Alan Kaller, 2008; Stone et al., 2007), AMD3465 (Genzyme Corp., Biochem Pharmacol. 2009 Oct 15;78(8):993-1000; Bodart et al., 2009; Ling et al., 2013), GSK812397 (Jenkinson et al., 2010; Planesas et al., 2015), KRH-3955 (Murakami et al., 2009; Nakasone et al., 2013), KRH-1636

- (Ichiyama et al., 2003), D-[Lys3] GHRP-6 (Patel et al., 2012), TG-0054 (Burixafor, TaiGen Biotechnology Co., Ltd.; de Nigris et al., 2012; Hsu et al., 2015), WZ811 (Zhan et al., 2007), MSX-122 (Metastatix, Inc.; Liang et al., 2012), and 508MCI (Compound 26; Zhu et al., 2010);
- [21] (2) small-peptide antagonists, such as anti-HIV peptides T22 (Masuda et al., 1992), T134 and T140 (Tamamura et al., 1998), BKT140 (a/k/a BL-8040; TF14016; 4F-Benzoyl-TN14003, Biokine Therapeutics Ltd. (Rehovot, Israel); Fahham et al., 2012; Otani et al., 2012), ALX40-4C (Canadian company Allelix Biopharmaceuticals (ON, Canada); Doranz et al., 2001), GST-NT21MP (Yang et al., 2014), FC131 (Tanaka et al., 2009; Tanaka et al., 2008), FC122 (Inokuchi et al., 2011), POL6326 (de Nigris et al., 2012), and LY2510924 (Lilly) (Peng et al., 2015);
- [22] (3) antibodies to CXCR4, such as ulocuplumab (MDX1338/BMS-936564; Kuhne et al., 2013), PF-06747143 (Pfizer; Liu et al., 2017), 12G5 (Endres et al., 1996), i-bodies (AD-114, AD-114-6H, AD-114-Im7-FH, and AD-114-PA600-6H; AdAlta; Griffiths et al., 2016; Griffiths et al., 2018), nanobodies (238D2 and 238D4; Jahnichen et al., 2010; Proc. Natl Acad. Sci. 2010, USA 107(47), 20565-20570), and ALX-0651 (Ablynx, biparatopic nanobody, ClinicalTrials.gov Identifier: NCT01374503);
- [23] (4) ligand (CXCL12) analogs with suppressive activity, such as CTCE-9908 (Chemokine Therapeutics (BC, Canada); Wong et al., 2014); and
- [24] (5) radiolabeled CXCR4 ligands, such as [99mTc]O2-AMD3100 (Hartimath et al., 2013), [68Ga]pentixafor (Demmer et al., 2011; Gourni et al., 2011), [177Lu]pentixather, and [90Y]pentixather (Herrmann et al., 2016).
- [25] AMD3100 (JM 3100, plerixafor, marketed as Mozobil) is a CXCR4-specific small molecule antagonist that inhibits CXCL12-mediated calcium mobilization and chemotaxis in various cell types (Hatse, Princen et al. 2002), and prevents tumor growth in mouse xenograft models (Rubin, Kung et al. 2003, Cho, Yoon et al. 2013, Liao, Fu et al. 2015). AMD3100 was originally developed for HIV therapy (as an HIV entry blocker that specifically antagonizes CXCR4, one of the HIV entry co-receptors), but was approved by the U.S. Food and Drug Administration only for stem cell mobilization in patients with lymphoma and multiple myeloma in 2008 (Keating 2011). During the initial clinical trial of AMD3100 for HIV infection, this compound was noted to cause hematopoietic stem cell (HSC, CD34⁺) mobilization from bone marrow to peripheral blood (Broxmeyer et al., 2005; De Clercq, 2003; Liles et al., 2003).
- [26] Development of AMD3100 for HIV therapy was suspended due to significant side effects including thrombocytopenia, premature ventricular contractions, and leukocytosis following long-term inhibition of CXCR4/CXCL12 axis (Hendrix, et al., 2004; Peled, et al., 2012). Although investigating the possibility of the use of AMD3100 as an anticancer drug is underway, lack of oral availability and some severe

side effects accompanying with long-term use need to be overcome (Peled, et al., 2012). Instead, subcutaneous AMD3100 was approved in combination with G-CSF for autologous stem cell mobilization and transplantation for patients in the US with non-Hodgkin lymphoma (NHL) or multiple myeloma for up to 4 consecutive days, and for patients in Europe with multiple myeloma or lymphoma for 2-4 consecutive days and for up to 7 consecutive days, respectively (DiPersio et al., 2009a; DiPersio et al., 2009b; Keating, 2011). So far, AMD3100 is the only CXCR4 antagonist approved by the FDA.

- [27] AMD3100 has been shown to inhibit CXCL12-mediated calcium mobilization and chemotaxis in various cancer cell types expressing CXCR4 (Hatse et al., 2002), and has demonstrated significant anti-tumor activity with decreased metastasis and increased overall survival in various mouse xenograft models (Burger et al., 2011; Chatterjee et al., 2014; Cho et al., 2013; Debnath et al., 2013; Domanska et al., 2013; Liao et al., 2015; Rubin et al., 2003; Walenkamp et al., 2017).
- [28] However, AMD3100 exhibits partial agonism for CXCR4 *in vitro* and increases the proliferation of melanoma cells (Kim et al., 2010; Zhang et al., 2002). AMD3100 also acts as a positive allosteric modulator of CXCR7, an alternative chemokine receptor for CXCL12, and increases CXCL12 binding to CXCR7 (Kalatskaya et al., 2009). Like CXCR4, CXCR7 is also highly expressed by many types of cancers and is associated with tumor metastasis (Decaillot et al., 2011; Zabel et al., 2011). Due to the complex properties of AMD3100, the exact role of AMD3100 on cancer needs to be investigated carefully.
- [29] AMD3100 has been clinically evaluated as anticancer drug in combination with conventional cytotoxic drugs such as mitoxantrone, etoposide, cytarabine, daunorubicin, azacitidine, lenalidomide, decitabine, clofarabine, fludarabine, and/or idarubicin; receptor tyrosine kinase inhibitors AC220 and sorafenib; HSP90 inhibitor (Ganetespib); G-CSF; proteasome inhibitor (bortezomib); or monoclonal antibody (rituximab) in hematological malignancies including acute myeloid leukemia (AML), MM, Myelodysplastic Syndromes (MDS), CLL, and small lymphocytic lymphoma (SLL) (ClinicalTrials.gov Identifier: NCT00512252, NCT00694590, NCT00903968, NCT00990054, NCT01065129, NCT01373229, NCT01352650, NCT01236144, NCT01301963, NCT01160354, NCT01027923, NCT00943943, and NCT01435343). AMD3100 was examined for interrupting lymphoid tumor microenvironment communication (NCT01610999) and for chemosensitization in combination with other anti-cancer drugs in patients with relapsed or refractory AML, ALL, and MDS (ClinicalTrials.gov identifier: NCT00906945 and NCT01319864).
- [30] AMD3100 has been or is also being evaluated in solid tumors including high grade glioma, Ewing's sarcoma, neuroblastoma, pancreatic, ovarian, and colorectal cancers

alone or in combination with angiogenesis inhibitor bevacizumab (ClinicalTrials.gov identifier: NCT01339039, NCT01977677, NCT01288573, NCT02179970, and NCT03277209).

- [31] AMD3100 has been tested or is currently being evaluated for the mobilization of the hematopoietic stem/progenitor/precursor cells in patients with various hematological malignancies and diseases including AML, MDS, neutropenia, beta-thalassemia, sickle cell disease, WHIMS, Fanconi anemia, Wiskott-Aldrich Syndrome, systemic mastocytosis ((Domanska et al., 2013), NCT01058993, NCT01206075, NCT03226691, NCT00967785, NCT02678533, NCT03019809, and NCT00001756). It has been or currently being evaluated for recruiting CD34+ cells, endothelial progenitor cells (EPC), and/or CD117+ progenitor cells in patients with diabetes, wounds, critical limb ischemia, chronic obstructive pulmonary disease (COPD), cystic fibrosis, pulmonary fibrosis (ClinicalTrials.gov identifier: NCT02056210, NCT02790957, NCT01916577, NCT03182426).
- [32] Although investigating the possibility of the use of AMD3100 as an anticancer drug is underway, lack of oral availability and some severe side effects accompanying with long-term use need to be overcome (Peled et al., 2012).
- [33] The orally available CXCR4 antagonist, AMD070 (also referred to herein as AMD-070, AMD11070, AMD-11070, or X4P-001; X4 Pharmaceuticals) has demonstrated antitumor activity in various tumor models (Morimoto et al., 2016; O'Boyle et al., 2013; Parameswaran et al., 2011) and has been studied in a phase I/II clinical trial in HIV infected subjects (ClinicalTrials.gov identifier: NCT00089466, (Debnath et al., 2013). AMD070 is now in Phase II/III clinical trial in patients with WHIM syndrome, advanced melanoma, and renal cell carcinoma alone or together with pembrolizumab, nivolumab, or axitinib (ClinicalTrials.gov identifier: NCT03005327, NCT02823405, NCT02923531, and NCT02667886).
- [34] The efficacy of small peptide antagonists of CXCR4, T140 and its analogs TN14003, TC14012, and BKT140 (also referred to herein as BL-8040, TF14016, 4F-Benzoyl-TN14003, BioLineRx, Ltd.) for blocking CXCR4 has been demonstrated in numerous preclinical studies including stem cell mobilization, small-cell lung cancer (SCLC), breast cancer, melanoma, AML, chronic myeloid leukemia, MM, pancreatic cancer, and rheumatoid arthritis (Burger et al., 2011). Currently, BKT140 is under clinical development for hematopoietic stem and progenitor cell mobilization in various hematopoietic malignancies (ClinicalTrials.gov identifier: NCT02502968, NCT03154827, NCT02763384, NCT02639559, and NCT02462252). In addition, BKT140 is in Phase II clinical trials for interfering tumor microenvironment interaction in subjects with AML in combination with cytarabine (ClinicalTrials.gov identifier: NCT02502968). It is also investigated in various cancers such as AML,

gastric cancer, non-small-cell lung cancer (NSCLC), and metastatic pancreatic cancers together with immune checkpoint inhibitors such as Pembrolizumab (Keytruda, Merck) and Atezolizumab (Tecentriq, Genentech/Roche) (ClinicalTrials.gov Identifier: NCT03154827, NCT03281369, NCT03337698, NCT02907099, NCT03193190, and NCT02826486).

- [35] Similarly, CTCE-9908 (Chemokine Therapeutics Corp., Canada), a peptide inhibitor of CXCR4, reduced metastases in mouse models of osteosarcoma and melanoma (Kim et al., 2008). After phase I study in patients with advanced solid cancers, CTCE-9908 was designated as an orphan drug for the treatment of osteosarcoma by the FDA in 2005. There is no further development after phase I/II clinical trial has been completed in 2008 (Debnath, et al., 2013).
- [36] MSX-122 (Altiris Therapeutics), a small molecule CXCR4 antagonist, has been abandoned after clinical trial in patients with solid tumors (ClinicalTrials.gov identifier: NCT00591682). Two major setbacks have been reported in 2017 in the field of developing CXCR4 antagonists as anti-cancer drugs. Bristol-Myers Squibb (BMS) discontinued a phase I/II study of ulocuplumab (BMS-936564 / MDX1338, (Kuhne et al., 2013), a fully human anti-CXCR4 antibody, in patients with solid tumors for lack of efficacy (<https://seekingalphacom/article/4057548-bristol-failure-makes-small-dent-cxcr4-blocking-approach?page=2>). Also, Eli Lilly decided to abandon a few anti-cancer programs including LY2510924, which is a CXCR4 peptide antagonist (<http://www.fiercebitech.com/biotech/lilly-puts-two-thirds-mid-phase-cancer-pipeline-up-for-sale-major-shake-up-r-d-priorities>) after a series of clinical trials in solid tumors (ClinicalTrials.gov identifier: NCT02737072, NCT01391130, and NCT01439568). These events indicate that development of CXCR4 antagonists as anti-cancer drugs, especially for solid tumors, is an ongoing challenge.
- [37] Other small molecule antagonists that are currently under clinical investigation, include:
- [38] USL311 (Upsher-Smith) is under phase I and phase II in patients with solid tumors and glioblastoma multiforme (GBM), respectively (ClinicalTrials.gov identifier: NCT02765165); GMI-1359 (Glycomimetics) is in phase I trial (ClinicalTrials.gov identifier: NCT02931214).
- [39] PF-06747143 (Pfizer), a humanized anti-CXCR4 antibody, is in phase I clinical trial in patients with AML either alone or combined with chemotherapy (ClinicalTrials.gov identifier: NCT02954653, (Liu et al., 2017)).
- [40] AD-114 (i-body, AdAlta; (Griffiths et al., 2018) is a humanized shark anti-CXCR4 antibody and received orphan drug designation from United States FDA for a drug candidate used for treatment of Idiopathic Pulmonary Fibrosis in 2017

(<https://www.reuters.com/article/idUSFWN1F70Y0>). AD-114 study is mainly focused on fibrotic conditions, including wet age-related macular degeneration and non-alcoholic fatty liver disease

(<https://lungdiseasenews.com/2017/08/23/adalta-present-research-on-investigative-therapy-ad-114-at-ipf-summit/>).

- [41] POL6326 (balixafortide, Polyphor), a peptide CXCR4 antagonist, has been under clinical evaluation in subjects with breast cancer (ClinicalTrials.gov identifier: NCT01837095), hematologic malignancies (ClinicalTrials.gov identifier: NCT01413568), and stem cell mobilization and healing in patients with acute myocardial infarction (ClinicalTrials.gov identifier: NCT01905475).
- [42] Radiolabeled CXCR4 ligands such as [^{99m}Tc]O2-AMD3100 and [⁶⁸Ga]Pentixafor has been used for preclinical or clinical SPECT or PET imaging of CXCR4 expression (Demmer et al., 2011; Gourni et al., 2011; Hartimath et al., 2013; Walenkamp et al., 2017). PET imaging with [⁶⁸Ga]Pentixafor has demonstrated increased expression of CXCR4 not only in hematologic and solid tumors such as leukemia, lymphoma, MM, adrenocortical carcinoma, and SCLC, but also in other pathological conditions such as splenosis, stroke, atherosclerosis, and myocardial infarction (Walenkamp et al., 2017).
- [43] A peptide CXCR4 ligand labeled with α - or β -emitters ([¹⁷⁷Lu]pentixather and [⁹⁰Y]pentixather) has been tested in more than 30 therapies as CXCR4-directed endoradiotherapy together with standard chemotherapy in patients with hematologic malignancies (Walenkamp et al., 2017).
- [44] Although various drugs antagonizing CXCR4 or antibody targeting CXCR4 have been developed and are under investigation, be it alone, in combination with conventional anticancer therapies, or with an immune checkpoint inhibitor, success has been limited so far (Peled, et al., 2012, Debnath, et al., 2013, Walenkamp, et al., 2017). Due to the pivotal roles of CXCR4 in B and T lymphocyte development and immune surveillance, long-term or persistent inhibition of CXCR4 with CXCR4 antagonist would potentially evoke immune system and hematopoietic dysfunctions and expose cancer patients to risk of immune suppression (Burger, 2009). Hematopoietic stem cells (HSCs) are normally protected in bone marrow niches. If CXCR4 antagonists are administered with cytotoxic drugs or radiotherapy, HSCs mobilized into periphery would be exposed to the effects of cytotoxic treatment which could exacerbate cytopenias. Cardiac complications observed in patients treated with AMD3100 also raised general concern in prolonged use of CXCR4 antagonists as anticancer drugs.
- [45] To avoid potential side effects associated with conventional CXCR4 antagonists and to develop more efficient anti-cancer drugs targeting CXCR4, new paradigm for designing CXCR4 inhibitor is urgently required.
- [46] Recently, GPCR heteromers offer new possibilities for developing more specific and

disease-restricted therapeutics. Traditionally, GPCRs were considered to be monomeric since monomeric GPCRs can activate G proteins by inducing conformational changes of the heptahelical domain upon ligand binding (Pin et al., 2008; Okada et al., 2001). However, increasing evidence demonstrates that GPCRs may form oligomers, either homo- or hetero-oligomers, and GPCR heteromers can exhibit specific properties that clearly distinguish them from existing well-defined monomers, such as signaling pathways, ligand binding affinities, internalization, and recycling (De Falco et al., 2007; Ferre et al., 2010; Gomes et al., 2016). Therefore, GPCR oligomerization may provide a way to increase the diversity of GPCR entities with a limited number of genes (Park and Palczewski, 2005). Thus, identification of novel GPCR heteromers would offer new possibilities for understanding the roles of GPCR heteromers in specific tissues and in specific diseases, as well as providing new opportunities for developing more efficient therapeutics with less side effects. (Milligan 2008; Rozenfeld and Devi 2010; Gomes et al., 2016; Farran 2017).

- [47] CXCR4 also forms heteromers with different GPCRs. So far CXCR4 is known to interact with GPCRs in the chemokine receptor family (CCR2 (Rodriguez-Frade et al., 2004; Sohy et al., 2007; Sohy et al., 2009; Armando et al., 2014), CCR5 (Agrawal et al., 2004; Rodriguez-Frade et al., 2004; Sohy et al., 2007, Sohy et al., 2009; Martinez-Munoz et al., 2014), CXCR3 (Watts et al., 2013), and CXCR7 (Sierro et al., 2007; Levoye et al., 2009; Decaillot et al., 2011)), chemerin chemokine-like receptor 1 (CMKLR1) (de Poorter et al., 2013), δ -opioid receptor (OPRD) (Pello et al. 2008; Burbassi et al., 2010), and adrenergic receptor family (ADRA1A (Tripathi et al., 2015), ADRA1B (Tripathi et al., 2015), and ADRB2 (LaRocca et al. 2010; Nakai et al., 2014)).
- [48] The existence of CXCR4-CCR2 and CXCR4-CCR5 heteromers was first noticed by studying HIV infection. Rodriguez-Frade et al. showed that CCR2-01, a CCR2-specific monoclonal antibody, did not compete with CCL2 for binding to CCR2 or trigger CCR2 signaling, but blocked replication of monotropic (R5) and T-tropic (X4) HIV strains by inducing oligomerization of CCR2 with CCR5 or CXCR4 (Rodriguez-Frade et al., 2004). Agrawal et al. also showed that CCR5D32 specifically inhibited CCR5 and CXCR4 cell surface expression by heteromerization with CCR5 and CXCR4, thereby inhibiting HIV coreceptor activity of CCR5 and CXCR4 in CD4+ cells (Agrawal et al., 2004). Co-expression of CCR5 was shown to prevent HIV-1 gp120 binding to the cell surface and to reduce X4 HIV-1 infectivity by inducing CCR5-CD4-CXCR4 oligomerization and conformational changes in CD4 and CXCR4 (Martinez-Munoz et al., 2014).
- [49] Sohy et al. reported the existence of negative binding cooperativity between the subunits of CXCR4-CCR2 and CXCR4-CCR5 heteromers, i.e. the ligands of one

receptor competed for the binding of a specific tracer of the other, in recombinant cell lines and primary leukocytes (Sohy et al., 2007; Sohy et al., 2009). They also demonstrated that TAK-779, the CCR2 and CCR5 antagonist, prevented calcium mobilization and chemotaxis initiated by the CXCR4 agonist CXCL12 in cells co-expressing CCR2 and CXCR4 or CCR5 and CXCR4 (Sohy et al., 2007; Sohy et al., 2009). Armando et al. showed that CXCR4 and CCR2 can form homo- and hetero-oligomers, and co-activation of CCR2 and CXCR4 with the human monocyte chemotactic protein 1 (MCP-1) and CXCL12 caused a synergistic increase in calcium mobilization (Armando et al., 2014).

- [50] Watts et al. identified CXCR4-CXCR3 heteromers in HEK293T cells and showed negative binding cooperativity for endogenous agonists and small CXCR3 agonist VUF10661, but not for CXCR3 antagonists VUF10085 nor CXCR4 antagonist AMD3100 (Watts et al., 2013).
- [51] CXCR4 also interacts with ACKR3, also known as CXCR7, a chemokine family GPCR that binds the CXCL12 and CXCL11, but is unable to couple with G proteins upon agonist binding. Sierro et al. identified CXCR4-ACKR3 heteromers as functional heteromers that displayed enhanced CXCL12-induced calcium signaling and altered ERK1/2 signaling in HEK293 cells (Sierro et al., 2007). Lovoye et al., reported an opposite result showing that CXCR4-ACKR3 heteromers exhibited reduced G α i and calcium responses upon exposure to CXCL12 in CHO-K1 cells (Levoye et al., 2009). Decaillot et al. noticed that co-expression of CXCR4 and ACKR3 constitutively recruited β -arrestin to CXCR4/ACKR3 heteromers and potentiated CXCL12-mediated β -arrestin-dependent downstream signaling pathways, including ERK1/2, p38 MAPK, and enhanced cell migration upon exposure to CXCL12 (Decaillot et al., 2011).
- [52] CXCR4 also interacts with the chemerin chemokine-like receptor 1 (CMKLR1, also known as ChemR23) (de Poorter et al., 2013). Although CXCR4-CMKLR1 heteromers display negative agonist binding cooperativity similar to CXCR4-CCR2 and CXCR4-CCR5 heteromers reported by Sohy et al. (Sohy et al., 2007; Sohy et al., 2009), AMD3100 did not cross inhibit chemerin binding or inhibit calcium mobilization induced by CXCL12 (de Poorter et al., 2013).
- [53] CXCR4 and the δ -opioid receptor (DOR) are widely distributed in brain tissues and immune cells. Pello et al. reported that CXCR4 and DOR can form heteromers, and simultaneous stimulation with both agonists suppressed the activation of G α i signaling, and prevented cell migration toward CXCL12, although individual agonist elicit robust G α i signaling (Pello et al., 2008). Burbassi et al. also noticed increased CXCR4-DOR heteromers and reduced coupling of CXCR4 to G-proteins in brain tissue and cultured glia from μ -opioid receptor (MOR)-deficient mice (Burbassi et al., 2010). CXCR4 function was rescued with a DOR antagonist, indicating that DOR is involved in the

- suppression of CXCR4 in glia by forming CXCR4-DOR heteromers (Burbassi et al., 2010).
- [54] CXCR4 is also known to interact with α - and β -adrenergic receptors (α -AR and β -AR). LaRocca et al. reported that stimulation of CXCR4 with CXCL12 negatively regulates β -AR-induced cAMP accumulation and PKA-dependent phosphorylation of phospholamban in adult rat ventricular myocytes using non-selective β -AR agonist, isoproterenol (LaRocca et al., 2010). They showed the co-expression of CXCR4 and ADRB2 (β 2-AR) on the cardiac myocytes, and physical association of CXCR4 with ADRB2 using co-immunoprecipitation and bioluminescence resonance energy transfer, suggesting the CXCR4-ADRB2 heteromer as a novel cardiac modulator.
- [55] Nakai et al. studied the function of ADRB2 on lymphocytes. Stimulation of lymphocytes with ADRB2-selective agonists suppressed egress of lymphocytes from lymph nodes and produced lymphopenia in mice (Nakai et al., 2014). ADRB2 physically interacted with CCR7 and CXCR4, and activation of ADRB2 enhanced retention-promoting signals through CCR7-ADRB2 and CXCR4-ADRB2 heteromers, and subsequently reduced lymphocyte egress from lymph nodes.
- [56] Tripathi et al. revealed the presence of CXCR4-ADRA1A (α 1A-AR) and CXCR4-ADRA1B (α 1B-AR) heteromers on the surface of vascular smooth muscle cells (VSMC) (Tripathi et al., 2015). A peptide derived from the second trans-membrane helix of CXCR4 disrupted the interaction between ADRA1A/B and CXCR4, inhibited calcium mobilization and contraction of VSMC upon α 1-AR stimulation. Activation of CXCR4 with CXCL12 increased the potency of α 1-AR agonists on blood pressure response in rats, suggesting that CXCR4-ADRA1A/B heteromers could be a novel pharmacological target for blood pressure regulation.
- [57] CXCR4 was reported to interact with CNR2 (Cannabinoid Receptor 2, aka CB2) (Coke et al., 2016; Scarlett et al., 2018). Simultaneous activation of CXCR4 and CB2 with both agonists resulted in reduction of ERK1/2 activation, calcium mobilization and cellular chemotaxis. These results show that cannabinoid system can negatively modulate CXCR4 function and tumor progression.
- [58] As described above, GPCRs forming heteromers with CXCR4 have been studied within a limited number of GPCR families, such as chemokine, adrenergic, and opioid receptor families. Considering the major role and increased expression of CXCR4 in a variety of pathological conditions, there is a great potential for the existence of different CXCR4-GPCR_x heteromers that confer unique features to a specific disease. However, due to the large number of GPCR genes (about 800) and difficulties in establishing high-throughput proximity-based screening techniques and functional assays, the identification of new CXCR4-GPCR_x heteromers has been a major challenge.

Disclosure of Invention

Technical Problem

- [59] Thus, there exists in the art a need for identifying functional GPCR heteromers, such as CXCR4-GPCR_x, and developing their inhibitors for use as GPCR heteromer-targeting cancer therapeutics with higher efficacy and lower side effects. This invention satisfies this need and provides related advantages.

Solution to Problem

- [60] By the use of adenovirus high-throughput system (AdHTS) that allows rapid and efficient generation of multiple recombinant adenoviruses simultaneously (Choi, et al., 2012), and the adenovirus-based bimolecular fluorescence complementation (BiFC) assay (patent; Song, 2014), inventors screened large number of GPCRs for their association with CXCR4 in U-2 OS cells.
- [61] In one aspect, provided herein is a method for treatment, amelioration, prevention, or diagnosis, of a cancer in a subject having CXCR4-GPCR_x heteromer, the method comprising: administering to the subject a therapeutically effective amount of an inhibitor of a CXCR4-GPCR_x heteromer, wherein: GPCR_x heteromerizes with CXCR4 in the subject, the heteromerization of GPCR_x with CXCR4 is accompanied by enhancement of signaling downstream of CXCR4; and the enhancement of signaling downstream of CXCR4 is suppressed by the inhibitor of the CXCR4-GPCR_x heteromer.
- [62] In another aspect, provided herein, is a method for treating cancer in a patient having a cell containing a CXCR4-GPCR_x heteromer, the method comprising: administering to the patient an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein:
- [63] i) the CXCR4-GPCR_x heteromer has enhanced downstream signaling; and
- [64] ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient.
- [65] In another aspect, provided herein, is a method of suppressing enhanced downstream signaling from a CXCR4-GPCR_x heteromer in a cell of a patient suffering from cancer, the method comprising: administering to the patient an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein:
- [66] i) the CXCR4-GPCR_x heteromer has enhanced downstream signaling; and
- [67] ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient.
- [68] In another aspect, provided herein, is a pharmaceutical kit for use in treating cancer

in a patient having a cell containing a CXCR4-GPCR α heteromer, the pharmaceutical kit comprising: an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; wherein the CXCR4-GPCR α heteromer has enhanced downstream signaling.

[69] In another aspect, provided herein, is a pharmaceutical composition for use in treating cancer in a patient having a cell containing a CXCR4-GPCR α heteromer, the pharmaceutical composition comprising:

[70] i) an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and

[71] ii) a pharmaceutically acceptable carrier;

[72] wherein the CXCR4-GPCR α heteromer has enhanced downstream signaling.

[73] In another aspect, provided herein, is a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, wherein the CXCR4-GPCR α heteromer has enhanced downstream signaling, the method comprising:

[74] 1) determining whether the patient has the cancer cell containing the CXCR4-GPCR α heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if:

[75] i) the patient's cancer cell containing said CXCR4-GPCR α heteromer; or

[76] ii) a CXCR4-GPCR α heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR α heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient derived cell(s) containing the CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR α heteromer; and

[77] 2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer to the cancer patient.

[78] In another aspect, provided herein, is a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, the method comprising:

[79] 1) determining whether the patient's cancer cell contains the CXCR4-GPCR α heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR α heteromer is present in the patient's cancer cell; wherein:

[80] a) the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR,

CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and

- [81] b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay; and
- [82] 2) if the patient's cancer cell contains said CXCR4-GPCR α heteromer, then internally administering an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer to the patient.
- [83] In another aspect, provided herein, is a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer having enhanced downstream signaling, the method comprising:
- [84] 1) determining whether the patient has the cancer cell containing the CXCR4-GPCR α heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if:
 - [85] i) the patient's cancer cell contains said CXCR4-GPCR α heteromer; or
 - [86] ii) a CXCR4-GPCR α heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR α heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; and
- [87] 2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and
- [88] 3) if the patient does not have a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR α inhibitor as a single inhibitor.
- [89] In another aspect, provided herein, is a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, the method comprising:
- [90] 1) determining whether the patient's cancer cell contains the CXCR4-GPCR α heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR α heteromer is present in the patient's cancer cell; wherein:
 - [91] a) the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and

- [92] b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, enzyme-linked immunosorbent assay (ELISA), flow cytometry, RNAseq, qRT-PCR, microarray, or a fluorescent animal assay; and
- [93] 2) if the patient's cancer cell contains said CXCR4-GPCR α heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and
- [94] 3) if the patient's cancer cell does not contain said CXCR4-GPCR α heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR α inhibitor as a single inhibitor.
- [95] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the heteromerization of CXCR4 with GPCR α is evaluated by a proximity-based assay.
- [96] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the proximity-based assay is selected from the group consisting of bimolecular fluorescence complementation (BiFC), proximity ligation assay (PLA), fluorescence resonance energy transfer (FRET), bioluminescence resonance energy transfer (BRET), cysteine crosslinking, co-immunoprecipitation, and a combination of TR-FRET and SNAP-tag (Comps-Agrar et al., 2011).
- [97] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the heteromerization of CXCR4 with GPCR α is evaluated by a co-internalization assay.
- [98] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the enhancement of cell signaling downstream of CXCR4 is evaluated by an intracellular Ca²⁺ assay.
- [99] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR α is selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, APLNR, C5AR1, CALCR, CCR5, CHRM1, GALR1, EDNRB, HRH1, MLNR, NTSR1, PTGER2, PTGER3, SSTR2, and TACR3.
- [100] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR α is selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, PTGER2, and TACR3.
- [101] In certain embodiments of the method of treating, method of suppressing, pharma-

ceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.

[102] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.

[103] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.

[104] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3.

[105] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, HRH1, NTSR1, and TACR3.

[106] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3.

[107] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3.

[108] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADRB2, CHRM1, EDNRB, HRH1, and TACR3.

[109] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADRB2, CHRM1, HRH1, and TACR3.

[110] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADRB2, EDNRB, HRH1, and TACR3.

[111] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR_x is selected from the group consisting of ADRB2, EDNRB, and HRH1.

- [112] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR α is selected from the group consisting of ADRB2, CHRM1, and HRH1.
- [113] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR α is selected from the group consisting of ADRB2, HRH1, and TACR3.
- [114] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, GPCR α is selected from the group consisting of ADRB2 and HRH1.
- [115] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of the CXCR4-GPCR α heteromer is, or comprises, an inhibitor of the CXCR4.
- [116] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of CXCR4 is selected from the group consisting of: AD-114, AD-114-6H, AD-114-Im7-FH, AD-114-PA600-6H, ALX-0651, ALX40-4C, AMD070 (AMD11070, X4P-001), AMD3100 (plerixafor), AMD3465, ATI 2341, BKT140 (BL-8040; TF14016; 4F-Benzoyl-TN14003), CTCE-9908, CX549,,D-[Lys3] GHRP-6, FC122, FC131, GMI-1359, GSK812397, GST-NT21MP, isothiouraea-1a, isothiouraea-1t (IT1t), KRH-1636, KRH-3955, LY2510924, LY2624587, MSX-122, N-[¹¹C]Methyl-AMD3465, PF-06747143, POL6326, SDF-1 1-9[P2G] dimer, SDF1 P2G, T134, T140, T22, TC 14012, TG-0054 (Burixafor), USL311, ulocuplumab (MDX1338/BMS-936564), viral macrophage inflammatory protein-II (vMIP-II), WZ811, 12G5, 238D2, 238D4, [⁶⁴Cu]-AMD3100, [⁶⁴Cu]-AMD3465, [⁶⁸Ga]pentixafor, [⁹⁰Y]pentixather, [^{99m}Tc]O₂ - AMD3100, [¹⁷⁷Lu]pentixather, and 508MCl (Compound 26).
- [117] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of the CXCR4-GPCR α heteromer is, or comprises, an antagonist of the CXCR4.
- [118] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the antagonist of CXCR4 is selected from the group consisting of ALX40-4C, AMD070 (AMD11070, X4P-001), AMD3100 (plerixafor), AMD3465, ATI 2341, BKT140 (BL-8040; TF14016; 4F-Benzoyl-TN14003), CTCE-9908, CX549,,D-[Lys3] GHRP-6, FC122, FC131, GMI-1359, GSK812397, GST-NT21MP, isothiouraea-1a, isothiouraea-1t (IT1t), KRH-1636, KRH-3955, LY2510924, MSX-122, N-[¹¹C]Methyl-AMD3465, POL6326, SDF-1 1-9[P2G] dimer, SDF1 P2G, T134, T140, T22, TC 14012, TG-0054 (Burixafor), USL311, viral macrophage inflammatory protein-II (vMIP-II), WZ811, [⁶⁴Cu]-AMD3100, [⁶⁴Cu]-AMD3465, [⁶⁸Ga]pentixafor, [⁹⁰Y]pentixather,

[99mTc]O₂-AMD3100, [177Lu]pentixather, and 508MCl (Compound 26).

- [119] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of the CXCR4-GPCR_x heteromer is, or comprises, an antibody of the CXCR4.
- [120] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the antibody of CXCR4 is selected from the group consisting of AD-114, AD-114-6H, AD-114-Im7-FH, AD-114-PA600-6H, ALX-0651, LY2624587, PF-06747143, ulocuplumab (MDX1338/BMS-936564), 12G5, 238D2, and 238D4.
- [121] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of the CXCR4-GPCR_x heteromer is, or further comprises, an inhibitor of the GPCR_x.
- [122] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of the GPCR_x is concurrently or sequentially administered with the inhibitor of the CXCR4.
- [123] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of the CXCR4-GPCR_x heteromer is, or comprises, an antagonist, an inverse agonist, an allosteric modulator, an antibody or binding portion thereof, a ligand, or any combinations, of GPCR_x.
- [124] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of a molecule selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, PTGER2, and TACR3.
- [125] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of a molecule selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.
- [126] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of a molecule selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.
- [127] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of a molecule selected from the group consisting of ADCYAP1R1,

ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.

- [128] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3.
- [129] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, HRH1, NTSR1, and TACR3.
- [130] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3.
- [131] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3.
- [132] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADRB2, CHRM1, EDNRB, HRH1, and TACR3.
- [133] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADRB2, CHRM1, HRH1, and TACR3.
- [134] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADRB2, EDNRB, HRH1, and TACR3.
- [135] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADRB2, EDNRB, and HRH1.
- [136] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADRB2, CHRM1, and HRH1.

- [137] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADRB2, HRH1, and TACR3.
- [138] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of a molecule selected from the group consisting of ADRB2 and HRH1.
- [139] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of ADCYAP1R1 selected from the group consisting of: M65, Max.d.4, MK-0893, N-stearyl-[Nle¹⁷] neurotensin-(6-11)/VIP-(7-28), PACAP-(6-38), and PG 97-269.
- [140] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of ADORA2B selected from the group consisting of:
3-isobutyl-8-pyrrolidinoxanthine, alloxazine, AS16, AS70, AS74, AS94, AS95, AS96, AS99, AS100, AS101, ATL802, BW-A1433, caffeine, CGS 15943, CPX, CSC, CVT-6883, DAX, DEPX, derenofylline, DPCPX, FK-453, I-ABOPX, istradefylline, KF26777, LAS38096, LUF5981, MRE 2029F20, MRE 3008F20, MRS1191, MRS1220, MRS1523, MRS1706, MRS1754, MSX-2, OSIP339391, pentoxifylline, preladenant, PSB-10, PSB-11, PSB36, PSB603, PSB-0788, PSB1115, rolofylline, SCH 58261, SCH442416, ST-1535, theophylline, tonapofylline, vipadenant, xanthine amine congener, XCC, and ZM-241385.
- [141] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of ADORA3 selected from the group consisting of: ATL802, BW-A1433, caffeine, CGS 15943, CSC, CVT-6883, derenofylline, dextriguldipine, DPCPX, FK-453, flavanone, flavone, galangin, I-ABOPX, istradefylline, KF26777, LAS38096, LUF5981, MRE 2029F20, MRE 3008F20, MRE 3010F20, MRS1041, MRS1042, MRS1067, MRS1088, MRS1093, MRS1097, MRS1177, MRS1186, MRS1191, MRS1191, MRS1220, MRS1476, MRS1486, MRS1505, MRS1523, MRS1754, MRS928, MSX-2, nicardipine, preladenant, PSB-10, PSB-11, PSB36, PSB603, PSB1115, rolofylline, sakuranetin, SCH 58261, SCH442416, ST-1535, theophylline, tonapofylline, vipadenant, visnagin, VUF5574, VUF8504, VUF8507, xanthine amine congener, and ZM-241385.
- [142] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR α is an inhibitor of ADRB2 selected from the group consisting of: Alprenolol, atenolol,

betaxolol, bupranolol, butoxamine, carazolol, carvedilol, CGP 12177, cicloprolol, ICI 118551, ICYP, labetalol, levobetaxolol, levobunolol, LK 204-545, metoprolol, nadolol, NIHP, NIP, propafenone, propranolol, sotalol, SR59230A, and timolol.

- [143] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of ADRB2 is Carvedilol.
- [144] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of C5AR1 inhibitor is selected from the group consisting of: A8^{Δ71-73}, AcPhe-Orn-Pro-D-Cha-Trp-Arg, avacopan, C089, CHIPS, DF2593A, JPE1375, L-156,602, NDT9520492, *N*-methyl-Phe-Lys-Pro-D-Cha-Trp-D-Arg-CO₂H, PMX205, PMX53, RPR121154, and W54011.
- [145] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of CALCR selected from the group consisting of α -CGRP-(8-37) (human), AC187, CT-(8-32) (salmon), and olcegepant.
- [146] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of CALCR is AC187.
- [147] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of CHRM1 selected from the group consisting of 3-Quinuclidinyl benzilate (QNB), 4-DAMP, acridinium, AE9C90CB, AFDX384, amitriptyline, AQ-RA 741, atropine, benztropine, biperiden, darifenacin, dicyclomine, dosulepin, ethopropazine, glycopyrrolate, guanylpirenzepine, hexahydrodifenidol, hexahydrosiladifenidol, hexocyclium, himbacine, ipratropium, lithocholylcholine, methoctramine, ML381, muscarinic toxin 1, muscarinic toxin 2, muscarinic toxin 3, *N*-methyl scopolamine, otenzepad, oxybutynin, *p*-F-HHSiD, pirenzepine, propantheline, (R,R)-quinuclidinyl-4-fluoromethyl-benzilate, scopolamine, silahexocyclium, solifenacin, telenzepine, tiotropium, tolterodine, trihexyphenidyl, tripitramine, UH-AH 37, umeclidinium, and VU0255035.
- [148] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of CHRM1 is Oxybutynin, umeclidinium, and VU0255035.
- [149] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of EDNRB selected from the group consisting of A192621, ambrisentan, atrasentan, bosentan (RO 470203, Tracleer);, BQ788, IRL 2500, K-8794, macitentan,

RES7011, Ro 46-8443, SB209670, SB217242 (enrasentan), TAK 044, and tezosentan (RO610612).

- [150] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of EDNRB is bosentan.
- [151] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of HRH1 selected from the group consisting of (-)-chlorpheniramine, (+)-chlorpheniramine, (-)-*trans*-H₂-PAT, (+)-*cis*-H₂-PAT, (+)-*trans*-H₂-PAT, (±)-*cis*-H₂-PAT, (±)-*trans*-H₂-PAT, (*R*)-cetirizine, (*S*)-cetirizine, 9-OH-risperidone, A-317920, A-349821, ABT-239, alimemazine, amitriptyline, aripiprazole, arpromidine, asenapine, astemizole, AZD3778, azelastine, BU-E 47, cetirizine, chlorpheniramine, chlorpromazine, ciproxifan, clemastine, clobenpropit, clozapine, conessine, cyclizine, cyproheptadine, desloratadine, diphenhydramine, dosulepin, doxepin, epinastine, fexofenadine, fluphenazine, fluspirilene, haloperidol, hydroxyzine, impromidine, INCB-38579, JNJ-39758979, ketotifen, loratadine, loxapine, MK-0249, molindone, olanzapine, perphenazine, pimozide, pipamperone, pitolisant, promethazine, pyrilamine, quetiapine, risperidone, sertindole, terfenadine, thioridazine, thiothixene, trifluoperazine, tripeleennamine, triprolidine, ziprasidone, and zotepine.
- [152] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of HRH1 is cetirizine, pyrilamine, hydroxyzine, or loratadine.
- [153] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of MLNR selected from the group consisting of GM-109, MA-2029, and OHM-11526.
- [154] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of MLNR is MA-2029.
- [155] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x is an inhibitor of NTSR1 selected from the group consisting of Meclinertant, SR48527, SR48692, and SR142948A.
- [156] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of NTSR1 is Merclinertant.
- [157] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of GPCR_x

is an inhibitor of TACR3 selected from the group consisting of [Trp⁷, β-Ala⁸] neurokinin A-(4-10), AZD2624, FK 224, GR138676, GSK 172981, GSK 256471, N',2-diphenylquinoline-4-carbohydrazide 8m, N',2-diphenylquinoline-4-carbohydrazide, osanetant, PD 154740, PD 161182, PD157672, saredutant, SB 218795, SB 222200, SB 235375, SCH 206272, SSR 146977, and talnetant.

- [158] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of TACR3 is SSR146977.
- [159] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of the CXCR4-GPCR_x heteromer is a protein-protein interaction (PPI) inhibitor.
- [160] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the cancer is selected from the group consisting of breast cancer, lung cancer, brain cancer, kidney cancer, pancreatic cancer, ovarian cancer, prostate cancer, melanoma, multiple myeloma, gastrointestinal cancers, renal cell carcinoma, soft tissue sarcomas, hepatocellular carcinoma, stomach cancer, colorectal cancer, esophageal cancer, and leukemia.
- [161] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of CXCR4-GPCR_x heteromer is administered to the subject in a pharmaceutical composition.
- [162] Further disclosed herein is a method for assessing response, or potential response, of a subject having CXCR4-GPCR_x heteromer, to treatment, amelioration, or prevention of a cancer, the method comprising: obtaining a sample from the subject; detecting heteromerization of CXCR4 and GPCR_x in the sample; and based at least in part on detection of the heteromerization of CXCR4 and GPCR_x, assessing the subject's response, or potential response, to the treatment, amelioration, or prevention of a cancer.
- [163] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the heteromerization of CXCR4 and GPCR_x is accompanied by enhancement of signaling downstream of CXCR4.
- [164] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the enhancement of cell signaling downstream of CXCR4 is evaluated by an intracellular Ca²⁺ assay.
- [165] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the heteromerization of

CXCR4 with GPCR α is evaluated by a proximity-based assay.

- [166] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the proximity-based assay is selected from the group consisting of bimolecular fluorescence complementation (BiFC), proximity ligation assay (PLA), fluorescence resonance energy transfer (FRET), bioluminescence resonance energy transfer (BRET), cysteine crosslinking, and co-immunoprecipitation.
- [167] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the heteromerization of CXCR4 with GPCR α is evaluated by a co-internalization assay.
- [168] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4-GPCR α heteromer has, causes, or produces, the enhanced downstream signaling.
- [169] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the enhanced downstream signaling results from the CXCR4-GPCR α heteromer, such as results from agonism of the CXCR4-GPCR α heteromer, results from CXCR4 agonism of the CXCR4-GPCR α heteromer, or results from GPCR α agonism of the CXCR4-GPCR α heteromer.
- [170] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the enhanced downstream signaling is downstream of the CXCR4, the respective GPCR α , or the CXCR4-GPCR α heteromer.
- [171] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the cancer patient, such as suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the patient's cancer cells.
- [172] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the enhanced downstream signaling from the CXCR4-GPCR α heteromer is determined by an intracellular Ca²⁺ assay.
- [173] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the enhanced downstream signaling is an enhanced amount of calcium mobilization.
- [174] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the enhanced amount of

calcium mobilization is a synergistic amount of calcium mobilization.

- [175] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4-GPCR α heteromer has two or more of the following characteristics:
- [176] 1) the CXCR4-GPCR α heteromer components in a cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allosterism, as determined via one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay;
- [177] 2) an enhanced amount of calcium mobilization, such that: a) either CXCR4 or the respective GPCR α in an individual protomer context in a cell, upon co-stimulation with CXCL12 and a respective selective GPCR α agonist, results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; and b) the CXCR4-GPCR α heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; as determined via a calcium mobilization assay; or
- [178] 3) a CXCR4-GPCR α heteromer-selective reagent: i) alters heteromer-specific properties of the CXCR4-GPCR α heteromer in a patient derived cell, ii) alters heteromer-specific function of the CXCR4-GPCR α heteromer in a patient derived cell; iii) alters heteromer-specific properties of a patient derived cell containing the CXCR4-GPCR α heteromer; or iv) decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR α heteromer.
- [179] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4-GPCR α heteromer components in the cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allosterism, as determined via one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay.
- [180] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the proximity-based assay is, or comprises, resonance energy transfer (RET), bioluminescence RET (BRET), fluorescence RET (FRET), time-resolved fluorescence RET (TR-FRET), antibody-aided FRET, ligand-aided FRET, bimolecular fluorescence complementation

(BiFC), or a proximity ligation assay (PLA).

[181] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4-GPCR α heteromer exhibits the enhanced amount of calcium mobilization, such that:

[182] a) either the CXCR4 or the respective GPCR α in an individual protomer context in the cell upon co-stimulation with CXCL12 and a respective selective GPCR α agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; and

[183] b) the CXCR4-GPCR α heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist;

[184] as determined via a calcium mobilization assay.

[185] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4-GPCR α heteromer-selective reagent: i) alters heteromer-specific properties of the CXCR4-GPCR α heteromer in the patient derived cell; ii) alters heteromer-specific function of the CXCR4-GPCR α heteromer in the patient derived cell; iii) alters heteromer-specific properties of the patient derived cell containing the CXCR4-GPCR α heteromer; or iv) decreases cell proliferation of the patient derived cell containing the CXCR4-GPCR α heteromer.

[186] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4-GPCR α heteromer-selective reagent is a CXCR4 inhibitor, a GPCR α inhibitor, or a CXCR4-GPCR α heteromer inhibitor.

[187] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4-GPCR α heteromer-selective reagent is a CXCR4 antagonist, a GPCR α antagonist, or a CXCR4-GPCR α heteromer antagonist.

[188] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the method administers an inhibitor selected from the group consisting of: the CXCR4 inhibitor, the GPCR α inhibitor, or the CXCR4-GPCR α heteromer inhibitor.

[189] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor is administered as a pharmaceutical composition.

[190] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor is administered as a pharmaceutical composition.

ceutical composition, or pharmaceutical kit, disclosed herein, the administered inhibitor is the CXCR4 inhibitor.

- [191] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the administered inhibitor is the GPCR α inhibitor.
- [192] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the administered inhibitor is the inhibitor of the CXCR4-GPCR α heteromer.
- [193] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the method administers the combination of inhibitors selected from the group consisting of: the CXCR4 inhibitor, the GPCR α inhibitor, and the inhibitor of the CXCR4-GPCR α heteromer.
- [194] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the combination of inhibitors are administered sequentially, concurrently, or simultaneously.
- [195] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4 inhibitor is an antagonist of CXCR4, an inverse agonist of CXCR4, a partial antagonist of CXCR4, an allosteric modulator of CXCR4, an antibody of CXCR4, an antibody fragment of CXCR4, a ligand of CXCR4, or an antibody-drug conjugate of CXCR4.
- [196] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the GPCR α inhibitor is an antagonist of GPCR α , an inverse agonist of GPCR α , a partial antagonist of GPCR α , an allosteric modulator of GPCR α , an antibody of GPCR α , an antibody fragment of GPCR α , a ligand of GPCR α , or an antibody-drug conjugate of GPCR α .
- [197] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor of the CXCR4-GPCR α heteromer is an antagonist of the CXCR4-GPCR α heteromer, an inverse agonist of the CXCR4-GPCR α heteromer, a partial antagonist of the CXCR4-GPCR α heteromer, an allosteric modulator of the CXCR4-GPCR α heteromer, an antibody of the CXCR4-GPCR α heteromer, an antibody fragment of the CXCR4-GPCR α heteromer, a ligand of the CXCR4-GPCR α heteromer, a protein-protein interaction (PPI) inhibitor of the CXCR4-GPCR α heteromer, or an antibody-drug conjugate of the CXCR4-GPCR α heteromer.
- [198] In certain embodiments of the method of treating or method of suppressing, disclosed herein, the method further comprises detecting the CXCR4-GPCR α heteromer in the cancer patient.
- [199] In certain embodiments of the method of treating or method of suppressing, disclosed

herein, the method further comprises identifying the CXCR4-GPCR α heteromer in the cancer patient.

[200] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the method further comprises:

[201] i) obtaining or having obtained a biological sample from the cancer patient;

[202] ii) conducting or having conducted a diagnostic assay to determine presence, identity, or presence and identity, of a CXCR4-GPCR α heteromer in the obtained biological sample from the cancer patient; and

[203] iii) selecting the inhibitor or combination of inhibitors to suppress enhanced downstream signaling from the CXCR4-GPCR α heteromer.

[204] In certain embodiments of the method of treating or method of suppressing, disclosed herein, the patient's biological sample is a biological fluid sample or a biological tissue sample.

[205] In certain embodiments of the method of treating or method of suppressing, disclosed herein, a liquid biopsy is performed on the biological fluid sample or the biological tissue sample.

[206] In certain embodiments, the biological fluid sample includes circulating tumor cells (CTCs), tumor-derived cell-free DNA (cfDNA), circulating small RNAs, and extracellular vesicles including exosomes, from bodily fluids as disclosed, for example, in Campos CDM et al., "Molecular Profiling of Liquid Biopsy Samples for Precision Medicine," Cancer J. 2018 Mar/Apr;24(2):93-103, which is incorporated hereby in its entirety.

[207] In certain embodiments of the method of treating or method of suppressing, disclosed herein, the biological fluid sample is a blood sample, a plasma sample, a saliva sample, a cerebral fluid sample, an eye fluid sample, or a urine sample.

[208] In certain embodiments of the method of treating or method of suppressing, disclosed herein, the biological tissue sample is an organ tissue sample, a bone tissue sample, or a tumor tissue sample.

[209] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the cancer cell contains the CXCR4-GPCR α heteromer.

[210] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, a normal, non-cancerous cell, does not contain the CXCR4-GPCR α heteromer.

[211] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR α heteromer

is decreased in the range of 5-100% more upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient.

- [212] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor.
- [213] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the efficacy of a GPCR α inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor.
- [214] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR α heteromer is decreased in the range of 5-100% more upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient.
- [215] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor.
- [216] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the efficacy of a GPCR α inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor.
- [217] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the method administers a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer.
- [218] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the method administers a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer.

ceutical composition, or pharmaceutical kit, disclosed herein, the method administers a combination of a CXCR4 inhibitor and a GPCR $\times$ inhibitor.

[219] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the method administers a CXCR4-GPCR $\times$ heteromer inhibitor.

[220] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the administering of the combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR $\times$ heteromer in the cancer patient in the range of between 5-2000 fold, relative to single inhibitor administration.

[221] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the administering of the combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR $\times$ heteromer in the cancer patient in the range of between 5-1500 fold, relative to single inhibitor administration.

[222] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the administering of the combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR $\times$ heteromer in the cancer patient in the range of between 500-1500 fold, relative to single inhibitor administration.

[223] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the administering of the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR $\times$ heteromer in the cancer patient in the range of between 5-2000 fold, relative to suppression of downstream signaling from either a CXCR4 protomer or a GPCR $\times$ protomer in their respective individual protomer context.

[224] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the administering of the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR $\times$ heteromer in the cancer patient in the range of between 5-1500 fold, relative to suppression of downstream signaling from either a CXCR4 protomer or a GPCR $\times$ protomer in their respective individual protomer context.

[225] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the administering of the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR $\times$ heteromer in the cancer patient in the range of between 500-1500 fold, relative to suppression of downstream signaling from either a CXCR4 protomer or a GPCR $\times$ protomer in their respective individual protomer context.

- [226] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the cancer cell of the patient contains the CXCR4-GPCR α heteromer in a greater concentration than a normal, non-cancerous cell from said patient,
- [227] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the presence of the CXCR4-GPCR α heteromer in the cancer cell of the patient identifies a sub-population of CXCR4-mediated cancer patients.
- [228] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the presence of the CXCR4-GPCR α heteromer in the cancer cell of the patient is a biomarker of a sub-population of CXCR4-mediated cancer patients.
- [229] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the biomarker of the sub-population of CXCR4-mediated cancer patients allows for precision medicine, patient stratification, or patient classification.
- [230] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the biomarker of the sub-population of CXCR4-mediated cancer patients allows for GPCR-based precision cancer therapeutics.
- [231] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor is an antibody.
- [232] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the CXCR4-GPCR α heteromer-selective reagent is an antibody.
- [233] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the antibody is a bi-specific antibody of the CXCR4-GPCR α heteromer.
- [234] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the antibody is a heteromer-specific antibody of the CXCR4-GPCR α heteromer.
- [235] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the inhibitor is a bi-specific ligand(s) of the CXCR4-GPCR α heteromer.
- [236] In certain embodiments of the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, disclosed herein, the antibody is an antibody-drug conjugate (ADC), as disclosed in, for example, Beck et al., "Strategies

and challenges for the next generation of antibody-drug conjugates", *Nature Reviews Drug Discovery*, 16:315-337, (2017) and Lambert, et al., "Antibody-Drug Conjugates for Cancer Treatment", *Annual Review of Medicine*, 69:191-207 (2018), each of which are incorporated hereby in its entirety.

[237]

Brief Description of Drawings

[238]

FIGURE 1: Shows schematic drawing of bimolecular fluorescence complementation (BiFC) assay. GPCR A is fused with N-terminal fragment of yellow fluorescent protein (YFP) Venus (VN) and GPCR B is fused with C-terminal fragment of Venus (VC). When GPCR A and B form heteromer, the complementary VN and VC are close enough to form functional Venus.

[239]

FIGURES 2a-2u: Shows identification of CXCR4-interacting GPCRs using BiFC assay. Representative images that showed negative BiFC signal; CXCR4-VN and HA-VC (FIG. 2a), and CXCR4-VN and GCGR-VC (FIG. 2c). Representative images of CXCR4 and GPCR_x that showed positive BiFC signal; CXCR4-VN and CXCR4-VC (FIGS. 2b), CXCR4-VN and ADCYAP1R1-VC (FIG. 2d), CXCR4-VC and ADORA2B-VN (FIG. 2e), CXCR4-VN and ADORA3-VC (FIG. 2f), CXCR4-VN and ADRB2-VC (FIG. 2g), CXCR4-VN and APLNR-VC (FIG. 2h), CXCR4-VN and C5AR1-VC (FIG. 2i), CXCR4-VN and CALCR-VC (FIG. 2j), CXCR4-VN and CCR5-VC (FIG. 2k), CXCR4-VN and CHRM1-VC (FIG. 2l), CXCR4-VN and GALR1-VC (FIG. 2m), CXCR4-VN and EDNRB-VC (FIG. 2n), CXCR4-VN and HRH1-VC (FIG. 2o), CXCR4-VN and MLNR-VC (FIG. 2p), CXCR4-VN and NTSR1-VC (FIG. 2q), CXCR4-VN and PTGER2-VC (FIG. 2r), CXCR4-VC and PTGER3-VN (FIG. 2s), CXCR4-VN and SSTR2-VC (FIG. 2t), and CXCR4-VN and TACR3-VC (FIG. 2u).

[240]

FIGURES 3a-3b: Shows principle of GPCR co-internalization study. Cells co-expressing CXCR4-GFP and GPCR_x are treated with GPCR_x specific agonist. (FIG. 3a) CXCR4 and GPCR_x does not physically interact with each other. GPCR_x agonist induces internalization of GPCR_x, but not CXCR4-GFP. (FIG. 3b) CXCR4 and GPCR_x physically interact and forms heteromer. GPCR_x agonist induces internalization of GPCR_x, and CXCR4-GFP is co-internalized with GPCR_x.

[241]

FIGURES 4a-4q: Shows co-internalization of CXCR4-EGFP upon stimulation of GPCR_x with its corresponding agonist (control: CXCR4-GFP (FIG. 4a)). U-2 OS cells were co-transduced with adenoviruses encoding CXCR-EGFP and GPCR_x-VC, and the following GPCR_x partners were examined to form heteromer with CXCR4-EGFP and to induce co-internalization of the CXCR4-EGFP: GPCR_x represents ADCYAP1R1 (FIG. 4b), ADORA2B (FIG. 4c), ADORA3 (FIG. 4d), ADRB2 (FIG.

4e), APLNR (FIG. 4f), C5AR1 (FIG. 4g), CCR5 (FIG. 4h), CHRM1 (FIG. 4i), GALR1 (FIG. 4j), EDNRB (FIG. 4k), HRH1 (FIG. 4l), MLNR (FIG. 4m), NTSR1 (FIG. 4n), PTGER3 (FIG. 4o), SSTR2 (FIG. 4p), and TACR3 (FIG. 4q).

[242] **FIGURES 5a-5d:** Shows enhancement of the calcium response in cells co-expressing CXCR4 and ADRB2 upon co-stimulation with their respective selective agonists. MDA-MB-231 human breast cancer cells were transduced with adenoviruses encoding CXCR4 and HA-VC (FIG. 5a), ADRB2 and HA-VC (FIG. 5b), or CXCR4 and ADRB2 (FIG. 5c). Adenoviruses encoding HA-VC were used to adjust the total amount of adenoviruses transduced. Cells were allowed to express GPCR for 2 days, incubated with Cal-520 AM for 2 hr, and were treated with 15 nM of CXCL12, 100 nM of salmeterol (ADRB2-selective agonist), or CXCL12 and salmeterol together. Calcium mobilization was measured using FlexStation 3 Multi-Mode Microplate Reader. The results were normalized for base-line activity. Data represent three independent experiments (mean $\pm$ SEM). (FIG. 5d) Calcium mobilization was quantified by calculating the area-under-the-curve (AUC) of each graph in A-C. Data were normalized to CXCL12-stimulated calcium response in cells expressing CXCR4 alone. Sum represents the calculated additive effect of the responses obtained after stimulation of CXCL12 and salmeterol individually in cells co-expressing CXCR4 and ADRB2, to allow the visualization of the potentiation. ***P < 0.001; Student's t test.

[243] **FIGURES 6a-6l:** Shows enhancement of the calcium response in cells co-expressing CXCR4 and GPCR α upon co-stimulation with their respective selective agonists as shown in FIGS 5a-c. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and HA-VC, GPCR α and HA-VC, or CXCR4 and GPCR α . GPCR α represents ADCYAP1R1 (FIG. 6a), ADORA2B (FIG. 6b), ADORA3 (FIG. 6c), C5AR1 (FIG. 6d), CALCR (FIG. 6e), CHRM1 (FIG. 6f), EDNRB (FIG. 6g), HRH1 (FIG. 6h), MLNR (FIG. 6i), NTSR1 (FIG. 6j), PTGER2 (FIG. 6k), and TACR3 (FIG. 6l). Cells were incubated with Cal-520 AM, and were treated with CXCL12, GPCR α agonist, or CXCL12 and GPCR α agonist together. Calcium responses were quantified as described in FIG. 5. Data represent three independent experiments (mean $\pm$ SEM). *P < 0.05, **P < 0.01, ***P < 0.001; Student's t test.

[244] **FIGURES 7a-7e:** Shows examples of CXCR4-GPCR α heteromer that failed to show enhanced calcium signaling in cells co-expressing CXCR4 and GPCR α in the presence of agonists to both GPCRs. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and HA-VC, GPCR α and HA-VC, or CXCR4 and GPCR α . GPCR α represents APLNR (FIG. 7a), CCR5 (FIG. 7b), GALR1 (FIG. 7c), PTGER3 (FIG. 7d), and SSTR2 (FIG. 7e). Cells were incubated with Cal-520 AM, and were treated with CXCL12, GPCR α agonist, or CXCL12 and GPCR α agonist together. Calcium mobilization was quantified as described in FIG. 5. Data represent three independent ex-

periments (mean $\pm$ SEM). Student's t test. ns: not significant.

- [245] **FIGURES 8a-8l:** Shows co-treatment of both antagonists efficiently suppressed the enhanced calcium response when cells expressing CXCR4-GPCR α heteromer were stimulated with CXCL12 and GPCR α agonists simultaneously. MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and GPCR α . GPCR α represents ADRB2 (FIG. 8a), CHRM1 (FIGS. 8b-8d), EDNRB (FIG. 8e), HRH1(endo) (FIG. 8f and FIG. 8g), HRH1 (FIG. 8h and FIG. 8i), MLNR (FIG. 8j), NTSR1 (FIG. 8k), and TACR3 (FIG. 8l). In F & G, cells were transduced with adenoviruses encoding CXCR4, and endogenous HRH1 responses were examined with 50 nM of histamine. Cells were incubated with Cal-520 AM for 2 hr, GPCR antagonists or vehicle for 30 min, and stimulated with indicated amounts of CXCL12, GPCR α agonist, or CXCL12 and GPCR α agonist. Calcium mobilization was quantified as described in FIG. 5d. Data represent three independent experiments (mean $\pm$ SEM). *P < 0.05, **P < 0.01, ***P < 0.001; Student's t test.
- [246] **FIGURE 9:** Shows principle of internalization inhibition study. Cells co-expressing CXCR4-GFP and GPCR α are treated with CXCL12 (SDF-1) and /or GPCR α specific antagonist. (scenario A) CXCR4 and GPCR α forms heteromer. CXCR4 agonist, CXCL12 (SDF-1) induced internalization of CXCR4-GFP alone or CXCR4-GFP with GPCR α . (scenario B) GPCR α antagonist does not induce internalization of CXCR4-GFP. (scenario C) Internalization of CXCL12 (SDF-1) stimulated CXCR4-GFP is inhibited by GPCR α specific antagonist.
- [247] **FIGURES 10a-10c:** Shows inhibition of internalization by GPCR α antagonist for heteromers. (FIG. 10a) CXCR4-ADRB2; (FIG. 10b) CXCR4-CHRM1; and (FIG. 10c) CXCR4-HRH1. (FIG. 10a) CXCR4-GFP expressing U-2 OS cells were transduced with adenovirus encoding ADRB2. Treatment by CXCL12 (SDF-1), a CXCR4 agonist, induced CXCR4-ADRB2 internalization. But Carvedilol, an ADRB2 antagonist, did not induce internalization. Co-treatment with CXCL12 and Carvedilol partially inhibited CXCL12 induced CXCR4-ADRB2 internalization. (FIG. 10b) CXCR4-GFP expressing U-2 OS cells were transduced with adenovirus encoding CHRM1. Treatment by CXCL12 induced CXCR4-CHRM1 internalization. But Oxybutynin or Umeclidinium, CHRM1 antagonists, did not induce internalization. Co-treatment with CXCL12 and Oxybutynin or Umeclidinium inhibited CXCL12 induced CXCR4-CHRM1 internalization. (FIG. 10c) CXCR4-GFP expressing U-2 OS cells were transduced with adenovirus encoding HRH1. Treatment by CXCL12 induced CXCR4-HRH1 internalization. But Prometazine, Hydroxyzine, or Loratadine, HRH1 antagonists, did not induce internalization. Co-treatment with CXCL12 and Prometazine, Hydroxyzine, or Loratadine inhibited CXCL12 induced CXCR4-HRH1 internalization.

- [248] **FIGURES 11a-11c:** Shows the effect of GPCR α antagonist on the survival of patient derived cells (PDCs) from cancer patients. (FIG. 11a) Effect of ADRB2 antagonist (Carvedilol) on the survival of PDC. Carvedilol induced significantly decreased survival of cells. (IC₅₀=11.69 μ M) (FIG. 11b) Effect of CHRM1 antagonists (Oxybutynin, Umeclidinium) on the survival of PDC. Oxybutynin and Umeclidinium each showed significantly decreased survival of cells at IC₅₀=3.04 μ M and 4.03 μ M respectively. (FIG. 11c) Effect of HRH1 antagonists (Promethazine, Hydroxyzine, and Loratadine) on the survival of PDC. Promethazine, Hydroxyzine, and Loratadine each showed decreased survival of PDC at IC₅₀=18.39 μ M, 12.79 μ M and 5.29 μ M, respectively.
- [249] **FIGURES 12a-12c:** Shows CXCR4-ADRB2 heterodimer detection by PLA and qRT-PCR in U-2 OS cells over-expressing CXCR4 and ADRB2. CXCR4-GFP expressing U-2 OS cells were transduced with adenovirus encoding ADRB2 at different MOIs for 2 days. Then CXCR4-ADRB2 co-expressing U-2 OS cells were performed PLA. (FIG. 12a) Image of CXCR4-ADRB2 heteromer detection by PLA. (FIG. 12b) Increase of the PLA signal proportionate to the expression level of ADRB2 in a dose dependent manner. (FIG. 12c) Data from qRT-PCR showing endogenous ADRB2 expression level of U-2 OS cells.
- [250] **FIGURES 13a-13b:** Shows CXCR4-ADRB2 heteromer detection in PDC by PLA. GBM originated PDCs (sample IDs : 986T, 948T, 783T, 777T, 352T1, 352T2, 578T, 559T, 464T, 448T, 096T) were plated at chamber slide and CXCR4-ADRB2 heteromer were detected by PLA with CXCR4 and ADRB2 specific antibodies. (FIG. 13a) Image of CXCR4-ADRB2 heteromer detection. Nuclei were visualized with DAPI staining and CXCR4-ADRB2 heteromers were shown as small dots. (FIG. 13b) Ratio of CXCR4-ADRB2 heteromer in PDC.
- [251] **FIGURES 14a-14b:** Shows CXCR4-CHRM1 heteromer detection in PDC by PLA. GBM originated PDCs (sample IDs : 986T, 948T, 783T, 777T, 352T1, 352T2, 578T, 559T, 464T, 448T, 096T) were plated at chamber slide and CXCR4-CHRM1 heteromer were detected by PLA with CXCR4 and CHRM1 specific antibodies. (FIG. 14a) Image of CXCR4-CHRM1 heteromer detection. Nuclei were visualized with DAPI staining and CXCR4-ADRB2 heteromers were shown as small dots. (FIG. 14b) Ratio of CXCR4-CHRM1 heteromer in PDC.
- [252] **FIGURES 15a-15b:** Shows results from CXCR4-GPCR α heteromer detection in PDX. GBM originated PDXs (sample IDs; 777T, 783T, 948T, 559T) were detected CXCR4-ADRB2 heteromer by PLA with CXCR4 and ADRB2 specific antibodies. (FIG. 15a) Image of CXCR4-ADRB2 heteromer detection. Nuclei were visualized with DAPI staining and CXCR4-ADRB2 heteromers were shown as small dots. (FIG. 15b) Ratio of CXCR4-ADRB2 heteromer in PDX.

- [253] **FIGURES 16a-16b:** Shows the enhancement of the calcium response in cells co-expressing CXCR4 and GPCR α upon co-stimulation with their respective agonists. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and ADRB2 (FIG. 16a) or CXCR4 and HRH1 (FIG. 16b). Cells were cultured for 3 days, stained with Cal-520 AM, and were treated with either CXCL12 (30 nM) alone, increasing doses of histamine or salmeterol alone, or increasing doses of histamine or salmeterol in combination with 30 nM of CXCL12. Calcium mobilization was measured using FlexStation 3. Sum represents the calculated additive value of the responses evoked by 30 nM of CXCL12 alone (open square) and GPCR α ligand alone at indicated doses (filled circle). Sum graph was depicted as a broken line with inverted triangles. Statistically significant differences between the sum (inverted triangle) and co-treatment (filled square) at each point were determined by Student's *t* test. **P* < 0.05; ***P* < 0.01; ****P* < 0.001; Data represent mean $\pm$ SD (*n* = 3).
- [254] **FIGURES 17a-17c:** Shows no enhancement of the calcium response in cells co-expressing CXCR4 and GPCR α upon co-stimulation with their respective agonists. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and APLNR (FIG. 17a), CXCR4 and PTGER3 (FIG. 17b), or CXCR4 and SSTR2 (FIG. 17c). Cells were cultured for 3 days, stained with Cal 6 dye, and were treated with either CXCL12 alone (20 nM), increasing doses of Apelin-13, PGE2, or octreotide alone, or increasing doses of Apelin-13, PGE2 or octreotide in combination with 20 nM of CXCL12. Calcium mobilization was measured using FlexStation 3. Sum represents the calculated additive value of the responses evoked by 20 nM of CXCL12 alone (open square) and GPCR α ligand alone at indicated doses (filled circle). Sum graph was depicted as a broken line with inverted triangles. Statistically significant differences between the sum (inverted triangle) and co-treatment (filled square) at each point were determined by Student's *t* test. No statistical difference was observed at any point. Data represent mean $\pm$ SD (*n* = 3).
- [255] **FIGURES 18a-18b:** Shows the enhancement of the calcium response in wild-type MDA-MB-231 cells upon co-stimulation with CXCL12 and histamine. (FIG. 18a) MDA-MB-231 cells were stimulated with CXCL12 (100 nM), histamine (10 nM), or CXCL12 and histamine together, and Ca²⁺ responses elicited by endogenous CXCR4 and HRH1 were measured. (FIG. 18b) Calcium mobilization was quantified by calculating the area-under-the-curve (AUC) of each graph. Sum represents the calculated additive value of the responses evoked by each agonist. The data represent three independent experiments done in triplicates (mean $\pm$ SEM). Statistically significant difference between the sum and co-treatment was determined by Student's *t* test. **P* < 0.05.
- [256] **FIGURES 19a-19b:** Shows enhanced migration of MDA-MB-231 cells in response

to a co-stimulation with CXCL12 (30 nM) and histamine (50 nM). MDA-MB-231 cells were transduced with 1 MOI of lentivirus encoding CXCR4, and chemotactic migration of the cells toward CXCL12, histamine, CXCL12 and histamine together in the presence or absence of pyrilamine (1 μ M), a HRH1-selective inverse agonist, was evaluated. Chemotaxis was quantified by counting the migrated cells on the lower surface of the membrane of 10 fields per chamber. (FIG. 19a) A representative picture of each group. (FIG. 19b) Although histamine itself did not induce migration of the MDA-MB-231 cells, it significantly increased the migration of the cells upon co-stimulation with CXCL12. The enhanced migration of MDA-MB-231 cells was due to endogenous HRH1 since the addition of pyrilamine (1 μ M), a HRH1-selective inverse agonist, completely abolished the enhanced response. The data represent mean $\pm$ SEM ($n = 3$ or 5). Statistically significant differences were determined by Student's t test. * $P < 0.05$.

[257] **FIGURE 20:** Shows efficient suppression of the enhanced calcium response by co-treatment of anti-CXCR4 antibody and ADRB2 antagonist when cells expressing CXCR4-ADRB2 heteromer were simultaneously stimulated with CXCL12 and ADRB2 agonist. MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and ADRB2. Cells were treated with ADRB2 antagonist (Carvedilol), anti-CXCR4 antibody (12G5), or vehicle of indicated concentration and incubated with Cal 6 for 2 hours. Cells were subsequently stimulated with indicated amounts of CXCL12, ADRB2 agonist (Salmeterol), or CXCL12 and ADRB2 agonist.

[258] **FIGURES 21a-21b:** Shows detection of CXCR4-GPCR heteromer by PLA in U-2 OS cells over-expressing CXCR4 and GPCR α . U-2 OS cells expressing CXCR4-GFP were transduced with adenovirus encoding CHRM1 or HRH1 at different MOIs for 2 days. Subsequently, PLA was performed on U-2 OS cells co-expressing CXCR4-GPCR α using CXCR4- and GPCR α - specific antibodies. The PLA signal increased in proportion to the expression level of CHRM1 (FIG. 21a) and HRH1 (FIG. 21b) in a dose-dependent manner.

[259] **FIGURES 22a-22b:** (FIG. 22a) Shows images of three mice that were transplanted with a parental cell A549, A549-CXCR4 overexpressing CXCR4 stably and A549-CXCR4-ADRB2 stably overexpressing CXCR4-ADRB2 heteromer, respectively, at 28 days after transplantation; (FIG. 22b) Shows graph comparing tumor growth between transplanted cells from FIG. 22a; tumor growth was monitored every third or fourth day by measuring the length (L) and width (W) of the tumor and calculating tumor volume in the basis of the following formula: Volume = $0.5 LW^2$. The results are presented as the mean $\pm$ standard deviation of 3 individuals.

[260] **FIGURES 23a-23g:** Shows calcium response in MDAMB231 cells coexpressing CXCR4 and GPCR α upon co-stimulation with CXCL12 and endogenous ligand for

GPCRx. MDAMB231 cells were transduced with adenoviruses encoding CXCR4 and HA-VC, GPCR α and HA-VC, or CXCR4 and GPCR α where GPCR α represents ADCYAP1R1 (PAC1) (FIG. 23a), ADORA2B (FIG. 23b), ADORA3 (FIG. 23c), CHRM1 (FIG. 23d), EDNRB (FIG. 23e), MLNR (FIG. 23f), and TACR3 (FIG. 23g).

- [261] **FIGURES 24a-24b:** Shows calcium response in MDAMB231 cells co-expressing CXCR4 and ADCYAP1R1 upon co-stimulation with their selective endogenous ligands. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and ADCYAP1R1. (FIG. 24a) Cells were treated with PACAP38 (1 nM, ADCYAP1R1 selective endogenous ligand) alone, increasing doses of CXCL12 alone, or increasing doses of CXCL12 in combination with 1 nM of PACAP38. (FIG. 24b) Cells were treated with CXCL12 alone, increasing doses of PACAP38 alone, or increasing doses of PACAP38 in combination with 15 nM of CXCL12.
- [262] **FIGURES 25a-25b:** Shows calcium response in MDA-MB-231 cells co-expressing CXCR4 and TACR3 upon co-stimulation with their selective endogenous ligands. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and TACR3. (FIG. 25a) Cells were treated with Neurokinin B (0.4 nM, TACR3 selective endogenous ligand) alone, increasing doses of CXCL12 alone, or increasing doses of CXCL12 in combination with 0.4 nM of Neurokinin B. (FIG. 25b) Cells were treated with CXCL12 alone (30 nM), increasing doses of Neurokinin B alone, or increasing doses of Neurokinin B in combination with 30 nM of CXCL12.
- [263] **FIGURES 26a-26b:** Shows validation of CRISPR/Cas9-mediated CXCR4 and HRH1 gene editing in MDA-MB-231 cells. (FIG. 26a) MDA^{CXCR4+, HRH1-} cells were generated by disrupting HRH1 gene in MDA-MB-231 cells stably expressing CXCR4 (MDA^{CXCR4+}), by transducing lentiviruses encoding Cas9 and guide RNA targeting HRH1. The absence of functional HRH1 was confirmed by measuring calcium responses upon exposure to histamine. (FIG. 26b) CXCR4 gene was edited in MDA-MB-231 cells using CRISPR/Cas-9 system and expression of CXCR4 was detected using immunoblotting.
- [264] **FIGURES 27a-27b:** Shows enhanced calcium response in MDA-MB-231 cells upon co-treatment with CXCL12 and histamine is abrogated in the absence of HRH1. (FIG. 27a) MDAMB231 cells stably overexpressing CXCR4 (MDA^{CXCR4+}) or MDA^{CXCR4+} cells depleted with HRH1 using CRISPR/Cas9 system (MDA^{CXCR4+, HRH1-}) were treated with CXCL12 alone (50 nM), increasing concentrations of histamine, or histamine and CXCL12 together. (FIG. 27b) MDA^{CXCR4+} cells or MDA^{CXCR4+, HRH1-} cells were treated with histamine alone (15 nM), increasing concentrations of CXCL12, or CXCL12 and histamine together.
- [265] **FIGURES 28a-28b:** Shows enhanced calcium response in MDA-MB-231 cells upon co-treatment with CXCL12 and histamine is abrogated in the absence of CXCR4.

(FIG. 28a) MDA-MB-231 cells (MDA^{WT}) or MDA^{WT} cells depleted with CXCR4 using CRISPR/Cas9 system (MDA^{CXCR4-}) were treated with histamine alone (15 nM), increasing concentrations of CXCL12, or histamine and CXCL12 together. (FIG. 28b) MDA-MB-231 cells or MDA^{CXCR4-} cells were treated with CXCL12 alone (100 nM), increasing concentrations of histamine alone, or CXCL12 and histamine together, and Ca²⁺ responses were measured.

[266] **FIGURES 29a-29b:** Shows anti-tumor effect of CXCR4 inhibitor, ADRB2 inhibitor, or CXCR4 and ADRB2 inhibitor in mice transplanted with A549-CXCR4-ADRB2 cell line stably overexpressing CXCR4-ADRB2 heteromer. Figure 29a is a graph comparing tumor growth rates for the *in vivo* antitumor effect of CXCR4 inhibitor LY2510924, ADRB2 inhibitor Carvedilol, or combination of LY2510924 and Carvedilol. Figure 29b is a graph comparing tumor growth rates for the antitumor effect of CXCR4 inhibitor AMD070, ADRB2 inhibitor Carvedilol, or combination of AMD070 and Carvedilol. Tumor growth was monitored every third or fourth day by measuring the length (*L*) and width (*W*) of the tumor and calculating tumor volume in the basis of the following formula: Volume = 0.5 *LW*². The results are presented as the mean ± standard deviation of 10 individuals.

[267] Unless indicated otherwise, the following includes abbreviations for terms disclosed herein: acute myeloid leukemia (AML), Adenosine A3 Receptor (ADORA3), Adenosine Receptor A2b (ADORA2B), adenovirus high-throughput system (AdHTS), Adenylate Cyclase Activating Polypeptide 1 (Pituitary) Receptor Type I (ADCYAP1R1), Adrenoceptor Alpha 1A (ADRA1A), Adrenoceptor Beta 2 (ADRB2), Apelin Receptor (APLNR), Atypical chemokine receptor 3 (ACKR3), bimolecular fluorescence complementation (BiFC), Bioluminescence Resonance Energy Transfer (BRET), bovine serum albumin (BSA), Calcitonin Receptor (CALCR), Cancer stem cells (CSCs), C-C chemokine receptor type 2 (CCR2), chemerin chemokine-like receptor 1 (CMKLR1), Cholinergic Receptor Muscarinic 1 (CHRM1), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), chronic obstructive pulmonary disease (COPD), Complement C5a Receptor 1 (C5AR1), C-terminal fragment of Venus (VC), C-X-C Motif Chemokine ligand 12 (CXCL12), CXCR4, cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4), δ -opioid receptor (OPRD), Endothelin Receptor Type B (EDNRB), enzyme-linked immunosorbent assay (ELISA), formalin-fixed paraffin-embedded (FFPE), fluorescence resonance energy transfer (FRET), G protein-coupled receptor (GPCR), Galanin Receptor 1 (GALR1), glioblastoma multiforme (GBM), Glucagon receptor (GCGR), GPCR heteromer identification technology (GPCR-HIT), Granulocyte-colony stimulating factor (G-CSF), hematopoietic stem cells (HSCs), hepatocellular carcinoma (HCC), Histamine Receptor H1 (HRH1), human immunodeficiency virus

(HIV), International Union of Basic and Clinical Pharmacology Committee on Receptor Nomenclature and Drug Classification (NC-IUPHAR), μ -opioid receptor (MOR), Motilin Receptor (MLNR), Multiple myeloma (MM), multiplicity of infection (MOI), Myelodysplastic Syndromes (MDS), Neurotensin Receptor 1 (NTSR1), non-Hodgkin lymphoma (NHL), non-small-cell lung cancer (NSCLC), N-terminal fragments of Venus (VN), patient derived cell (PDC), Patient-Derived Xenograft (PDX), positron emission tomography (PET), Computed Tomography (CT), programmed cell death ligand 1 (PD-L1), programmed cell death protein 1 (PD-1), Prostaglandin E Receptor 2 (PTGER2), Prostaglandin E Receptor 3 (PTGER3), proximity ligation assay (PLA), reverse transcription-quantitative polymerase chain reaction (RT-qPCR), Single-photon emission computed tomography (SPECT), small lymphocytic lymphoma (SLL), small-cell lung cancer (SCLC), Somatostatin Receptor 2 (SSTR2), Stromal cell-derived factor 1 (SDF-1), systemic lupus erythematosus (SLE), Tachykinin Receptor 3 (TACR3), Threshold cycles (Ct), time-resolved FRET (TR-FRET), tumor microenvironment (TME), Vascular endothelial growth factor (VEGF), vascular smooth muscle cells (VSMC), WHIM syndrome (Warts, Hypogammaglobulinemia, Infections, and Myelokathexis), green fluorescence protein (GFP), and yellow fluorescence protein (YFP).

Mode for the Invention

- [268] CXCR4 plays important roles in tumor formation and progression, but the development of CXCR4 antagonists as anticancer drugs has not been successful possibly due to side effects and lack of efficacy within acceptable dose ranges. Recently, various CXCR4-GPCR heteromers with distinct physiological and pharmacological properties were reported, but their roles in cancer biology or possibilities for developing anti-cancer therapeutics targeting CXCR4-GPCR heteromers have not been clearly understood.
- [269] Traditionally GPCRs were believed to function as monomers that interact with hetero-trimeric G proteins upon ligand binding, and drugs were developed based on monomeric or homomeric GPCRs (Milligan 2008). This view has changed drastically by the discoveries that GPCRs can form heteromers, and heteromerization is obligatory for some GPCRs. GPCR heteromerization is known to alter GPCR maturation and cell surface delivery, ligand binding affinity, signaling intensity and pathways, as well as receptor desensitization and recycling (Terrillon and Bouvier 2004; Ferre et al., 2010; Rozenfeld and Devi 2010; Gomes et al., 2016; Farran 2017). Different GPCR heteromers display distinct functional and pharmacological properties, and GPCR heteromerization can vary depending on cell types, tissues, and diseases or pathological conditions (Terrillon and Bouvier 2004; Ferre et al., 2010; Rozenfeld and Devi 2010;

Gomes et al., 2016; Farran 2017). Now GPCR heteromerization is regarded as a general phenomenon, and deciphering GPCR heteromerization opens new avenues for understanding receptor function, physiology, roles in diseases and pathological conditions. Accordingly, identification of GPCR heteromers and their functional properties offers new opportunity for developing new pharmaceuticals or finding new use of old drugs with fewer side effects, higher efficacy, and increased tissue selectivity (Ferre et al., 2010; Rozenfeld and Devi 2010; Farran 2017).

[270] The identification of bona fide GPCR heteromer requires intensive and critical evaluation. To distinguish GPCR heteromers from simple association of GPCRs, researchers in this field and the International Union of Basic and Clinical Pharmacology Committee on Receptor Nomenclature and Drug Classification (NC-IUPHAR) have declared GPCR heteromer as "macromolecular complex composed of at least two (functional) receptor units [protomers] with biochemical properties that are demonstrably different from those of its individual components" and these heteromers exist in native tissue (Ferre et al., 2009; Gomes et al., 2016; Pin et al., 2007). They proposed three criteria to demonstrate GPCR heteromers: (1) Heteromers should exhibit appropriate co-localization and interaction to enable allosterism using co-immunoprecipitation, in situ hybridization, or proximity-based techniques including proximity ligation assays in cells/tissues that express both receptors and not in cells/tissues that lack one of the receptors; (2) Heteromers should exhibit distinct properties such as changes in signaling, ligand binding, and/or trafficking, only in cells/tissues expressing both receptors but not in cells/tissues that lack one of the receptors; and (3) Heteromer-selective reagents should alter heteromer-specific properties. Heteromer-selective reagents include heteromer-selective antibodies, membrane-permeable peptides, and bivalent/bifunctional ligands (Gomes et al., 2016; Pin et al., 2007). Although many GPCR heteromers have been identified *in vitro* using recombinant receptors expressed in heterologous cells, only a few have demonstrated novel properties, and very few have shown evidence for GPCR heteromerization in native tissue due to technical problems (Gomes et al., 2016). NC-IUPHAR announced that one should provide evidence that satisfies at least two of the above three criteria for approval of new GPCR heteromers (Pin et al., 2007).

[271] As disclosed herein, to establish whether criterion 1 of 3 is satisfied (relating to whether heteromer components colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allosterism) in determining the presence/existence of a CXCR4-GPCR α heteromer, one or more of the following methods may be utilized to including, but not limited to: co-internalization assays; co-localization assays (determining co-localization of the receptor protomers within a cellular compartment) such as *in situ* hybridization, immunohistochemistry, or immunoelectron mi-

croscopy; proximity-based assays, such as proximity-based biophysical techniques, including resonance energy transfer (RET), bioluminescence RET (BRET), fluorescence RET (FRET), time-resolved fluorescence RET (TR-FRET), antibody-aided FRET, ligand-aided FRET, bimolecular fluorescence complementation (BiFC), and proximity ligation assays (PLA); co-immunoprecipitation assays; or fluorescent animals. For example, BiFC, co-internalization assay, or PLA, were utilized to evaluate whether a CXCR4-GPCR α heteromer satisfied criterion 1 of 3.

- [272] As disclosed herein, to establish whether criterion 2 of 3 is satisfied (relating to whether a heteromer exhibits properties distinct from those of the individual protomers), such as a CXCR4-GPCR α heteromer that results in an enhanced downstream signaling, for example an enhanced calcium mobilization (such as determined by a calcium mobilization assay), a two-tiered approach was utilized on those CXCR4-GPCR α heteromers that satisfied criterion 1 of 3 discussed above: (1) determine the presence/absence of an enhanced downstream signaling, for example an enhanced calcium mobilization (synergism) in the individual protomer context - comparing calcium mobilization in cells co-expressing HA-VC and one of the protomers (either CXCR4 or GPCR α) upon (a) co-stimulation with CXCL12 and the respective selective agonist, relative to (b) stimulation with either CXCL12 alone or the respective selective agonist alone; and (2) determine the presence/absence of an enhanced downstream signaling, for example an enhanced calcium mobilization (synergism) in the CXCR4-GPCR α heteromer context - comparing calcium mobilization in cells co-expressing CXCR4 and GPCR α upon (a) co-stimulation with CXCL12 and the respective selective agonist, relative to (b) the sum of stimulation with either CXCL12 alone or the respective selective agonist alone. As disclosed herein, to satisfy criterion 2 of 3 and be considered a CXCR4-GPCR α heteromer that results in an enhanced downstream signaling, for example an enhanced calcium mobilization, there must be (1) an absence of an enhanced downstream signaling, for example an enhanced calcium mobilization in either protomer context (i.e., the CXCR4 and HA-VC context or the GPCR α and HA-VC context), and (2) a presence of an enhanced downstream signaling, for example an enhanced calcium mobilization in the CXCR4-GPCR α heteromer context. In both the protomer context and the CXCR4-GPCR α heteromer context, the concentration of CXCL12 utilized to stimulate the cells (either as a single agent or in combination with the respective selective GPCR α agonist) and the concentration of the selective GPCR α agonist (either an endogenous agonist or a known selective agonist for the respective GPCR α) utilized to stimulate the cells (either as a single agent or in combination with CXCL12) are independently at a concentration of 100x the EC₅₀ concentration or lower. For example, the concentration of CXCL12 utilized to stimulate the cells (either as a single agent or

in combination with the respective selective GPCR α agonist) was at a concentration of 15 nM (which is approximately the EC₅₀ concentration against CXCR4).

[273] As disclosed herein, to establish whether criterion 3 of 3 is satisfied (relating to whether heteromer-selective reagents should alter heteromer-specific properties) in determining the presence/existence of a CXCR4-GPCR α heteromer, patient derived cells, having satisfied criterion 1 of 3 and 2 of 3, are effected in the presence of an antagonist (a CXCR4 antagonist, a GPCR α antagonist, or a CXCR4-GPCR α heteromer antagonist), such as effecting cell proliferation of the patient derived cells containing a CXCR4-GPCR α heteromer.

[274] In some embodiments, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, as disclosed herein, include or rely upon establishing that an association of CXCR4 and a GPCR α in a cell satisfies at least two of the following criteria or characteristics to be considered a CXCR4-GPCR α heteromer, comprising: 1) the CXCR4-GPCR α heteromer components in a cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allostereism, as determined via one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay; 2) an enhanced amount of calcium mobilization, such that: a) either CXCR4 or the respective GPCR α in an individual protomer context in a cell, upon co-stimulation with CXCL12 and a respective selective GPCR α agonist, results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; and b) the CXCR4-GPCR α heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; as determined via a calcium mobilization assay; or 3) a CXCR4-GPCR α heteromer-selective reagent: i) alters heteromer-specific properties of the CXCR4-GPCR α heteromer in a patient derived cell; ii) alters heteromer-specific function of the CXCR4-GPCR α heteromer in a patient derived cell; iii) alters heteromer-specific properties of a patient derived cell containing the CXCR4-GPCR α heteromer; or iv) decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR α heteromer. In some embodiments, the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific properties of the CXCR4-GPCR α heteromer in a patient derived cell, as determined by at least one of the following methods: PLA, radioligand binding assays, [³⁵S]GTP- γ S Binding assays, Calcium assay, cAMP assay, GTPase assay, PKA activation, ERK1/2

and/or Akt/PKB Phosphorylation assays, Src and STAT3 phosphorylation assays, CRE-reporter assay, NFAT-RE-reporter assay, SRE-reporter assay, SRF-RE reporter assay, NF-kB-RE reporter assay, Secreted alkaline phosphatase Assay, Inositol 1-Phosphate Production assay, Adenylyl Cyclase Activity assay, analysis of target gene expression by RT-PCR, RT-qPCR, RNAseq, next generation sequencing (NGS), or microarray. In some embodiments, the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific function of the CXCR4-GPCR α heteromer in a patient derived cell, as determined by at least one of the following methods: PLA, radioligand binding assays, [35 S]GTP- γ S Binding assays, Calcium assay, cAMP assay, GTPase assay, PKA activation, ERK1/2 and/or Akt/PKB Phosphorylation assays, Src and STAT3 phosphorylation assays, CRE-reporter assay, NFAT-RE-reporter assay, SRE-reporter assay, SRF-RE reporter assay, NF-kB-RE reporter assay, Secreted alkaline phosphatase Assay, Inositol 1-Phosphate Production assay, Adenylyl Cyclase Activity assay, analysis of target gene expression by RT-PCR, RT-qPCR, RNAseq, or microarray. In some embodiments, the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific properties of a patient derived cell containing the CXCR4-GPCR α heteromer, as determined by at least one of the following methods: assays on proliferation, migration, invasion, and drug resistance (survival) of cancer cells, modulation of immune cell function, angiogenesis, vasculogenesis, metastasis, drug resistance, tissue microarray (TMA), and cancer cell-tumor microenvironment (TME) interaction. For example, in some embodiments, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, as disclosed herein, include or rely upon establishing that an association of CXCR4 and a GPCR α in a cell satisfies at least two of the following criteria or characteristics to be considered a CXCR4-GPCR α heteromer, comprising: 1) the CXCR4-GPCR α heteromer components in a cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allostereism, as determined via one or more of the following: a co-internalization assay, bimolecular fluorescence complementation (BiFC), or a proximity ligation assay (PLA); 2) an enhanced amount of calcium mobilization, such that: a) either CXCR4 or the respective GPCR α in an individual protomer context in a cell, upon co-stimulation with CXCL12 and a respective selective GPCR α agonist, results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; and b) the CXCR4-GPCR α heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; as determined via

a calcium mobilization assay; or 3) a CXCR4-GPCR_x heteromer-selective reagent: i) alters heteromer-specific properties of the CXCR4-GPCR_x heteromer in a patient derived cell; ii) alters heteromer-specific function of the CXCR4-GPCR_x heteromer in a patient derived cell; iii) alters heteromer-specific properties of a patient derived cell containing the CXCR4-GPCR_x heteromer; or iv) decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR_x heteromer. In some embodiments, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, as disclosed herein, include or rely upon establishing that an association of CXCR4 and a GPCR_x in a cell satisfies criteria 1 and 2 to be considered a CXCR4-GPCR_x heteromer. In some embodiments, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, as disclosed herein, include or rely upon establishing that an association of CXCR4 and a GPCR_x in a cell satisfies criteria 1 and 3 to be considered a CXCR4-GPCR_x heteromer. In some embodiments, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, as disclosed herein, include or rely upon establishing that an association of CXCR4 and a GPCR_x in a cell satisfies criteria 2 and 3 to be considered a CXCR4-GPCR_x heteromer. In some embodiments, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, as disclosed herein, include or rely upon establishing that an association of CXCR4 and a GPCR_x in a cell satisfies criteria 1, 2, and 3, to be considered a CXCR4-GPCR_x heteromer.

- [275] As disclosed herein, the CXCR4-GPCR_x heteromer satisfying at least two of the three criteria may have, cause, or produce an enhanced downstream signaling. The enhanced downstream signaling may be the result from the CXCR4-GPCR_x heteromer, such as from agonism of the CXCR4-GPCR_x heteromer, from CXCR4 agonism of the CXCR4-GPCR_x heteromer, from GPCR_x agonism of the CXCR4-GPCR_x heteromer, and/or from CXCR4 agonism and GPCR_x agonism of the CXCR4-GPCR_x heteromer. In some embodiments, the enhanced downstream signaling may be downstream of the CXCR4, the respective GPCR_x, or the CXCR4-GPCR_x heteromer. In some embodiments, the enhanced downstream signaling may be from the CXCR4-GPCR_x heteromer, relative to downstream signaling from a CXCR4 protomer or a respective GPCR_x protomer in their respective individual protomer context. In some embodiments, the enhanced downstream signaling may be from the CXCR4-GPCR_x heteromer, relative to downstream signaling from a CXCR4 protomer in an individual protomer context. In some embodiments, the enhanced downstream signaling may be from the CXCR4-GPCR_x heteromer, relative to downstream signaling from a respective GPCR_x protomer in an individual protomer context. In some embodiments, the enhanced downstream signaling may be from the CXCR4-GPCR_x heteromer,

relative to downstream signaling from a CXCR4 protomer and a respective GPCR α protomer in their respective individual protomer context. The enhanced downstream signaling from said CXCR4-GPCR α heteromer, may in some embodiments, be suppressed in the cancer patient, such as suppressed in the patient's cancer cells. In some embodiments, the enhanced downstream signaling from said CXCR4-GPCR α heteromer, may be an enhanced amount of calcium mobilization (or synergistic amount of calcium mobilization), which may be determined by an intracellular Ca²⁺ assay, such as a calcium mobilization assay.

- [276] As disclosed herein, the CXCR4-GPCR α heteromer satisfying at least two of the three criteria may have, cause, or produce an enhanced downstream signaling, wherein the enhanced downstream signaling is an enhanced amount of calcium mobilization. The enhanced amount of calcium mobilization from the CXCR4-GPCR α heteromer may be a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is at least 10% greater, than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay. In some embodiments, the enhanced amount of calcium mobilization from the CXCR4-GPCR α heteromer may be a synergistic amount of calcium mobilization that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is at least 10% greater, than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay. For example, the enhanced amount of calcium mobilization (or synergistic amount of calcium mobilization) from the CXCR4-GPCR α heteromer may be a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is at least 20% greater, at least 30% greater, at least 40% greater, at least 50% greater, at least 75% greater, at least 90% greater, at least 100% greater, at least 150% greater, or at least 200% greater, than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay. In some embodiments, the enhanced amount of calcium mobilization (or synergistic amount of calcium mobilization) from the CXCR4-GPCR α heteromer may be a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, may be between 10-100% greater than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay, for example, may be a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the re-

spective selective GPCR α agonist, may be between 25-100% greater, 50-100% greater, 75-100% greater, or 100-200% greater, than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.

[277] In some embodiments, according to the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, as disclosed herein, the CXCR4-GPCR α heteromer satisfies at least two of the three criteria, thereby having, causing, or producing, an enhanced downstream signaling, wherein the enhanced downstream signaling is an enhanced amount of calcium mobilization, such that: a) either the CXCR4 or the respective GPCR α in an individual protomer context in the cell upon co-stimulation with CXCL12 and a respective selective GPCR α agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; and b) the CXCR4-GPCR α heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; as determined via a calcium mobilization assay. For example, in some embodiments, the enhanced downstream signaling from the CXCR4-GPCR α heteromer is an enhanced amount of calcium mobilization, such that: i) the calcium mobilization from the protomer CXCR4 or GPCR α , in the individual protomer context in the cell, is non-synergistic, as determined via calcium mobilization assay; and ii) the calcium mobilization from the CXCR4-GPCR α heteromer in the cell is synergistic, as determined via a calcium mobilization assay. In some embodiments, the individual protomer context may be: a) the individual protomer CXCR4 in the cell, in the absence of the respective individual protomer GPCR α ; or b) the respective individual protomer GPCR α in the cell, in the absence of the individual protomer CXCR4; upon co-stimulation with CXCL12 and a respective selective GPCR α agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay. In some embodiments, the individual protomer context may be, independently: a) the individual protomer CXCR4 in the cell, in the absence of the respective individual protomer GPCR α ; and b) the respective individual protomer GPCR α in the cell, in the absence of the individual protomer CXCR4; upon co-stimulation with CXCL12 and a respective selective GPCR α agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the

CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay. For example, in some embodiments, the CXCR4-GPCR_x heteromer, upon co-stimulation with the CXCL12 and the respective selective GPCR_x agonist, may result in a calcium mobilization amount that is greater than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay.

- [278] The term "CXCR4" as used herein refers to C-X-C Motif Chemokine Receptor 4, also identified by unique database identifiers (IDs) and alternate names as shown in Table 1 (Chatterjee et al., 2014; Debnath et al., 2013; Domanska et al., 2013; Guo et al., 2016; Peled et al., 2012; Roccato et al., 2014; Walenkamp et al., 2017).
- [279] The terms "GPCR_x" as used herein refers to GPCRs that were used in this study to investigate if these GPCRs interact with CXCR4 and show properties distinct from those of individual protomers, including ADCYAP1R1 (ADCYAP Receptor Type I), ADORA2B (Adenosine A2b Receptor), ADORA3 (Adenosine A3 Receptor), ADRB2 (Adrenoceptor Beta 2), APLNR (Apelin Receptor), C5AR1 (Complement C5a Receptor 1), CALCR (Calcitonin Receptor), CCR5 (Chemokine (C-C Motif) Receptor 5), CHRM1 (Cholinergic Receptor Muscarinic 1), GALR1 (Galanin Receptor 1), EDNRB (Endothelin Receptor Type B), HRH1 (Histamine Receptor H1), MLNR (Motilin Receptor), NTSR1 (Neurotensin Receptor 1), PTGER2 (Prostaglandin E Receptor 2), PTGER3 (Prostaglandin E Receptor 3), SSTR2 (Somatostatin Receptor 2), and TACR3 (Tachykinin Receptor 3), also identified by unique database identifiers (IDs) and alternate names as shown in Table 1 and Table 2.
- [280] Table 1 shown below provides the nomenclature of GPCR_x that form heteromers with CXCR4 and synergistically enhance Ca²⁺ response upon co-stimulation with both agonists, and Table 2 shown below provides the nomenclature of GPCR_x that form heteromers with CXCR4, but do not synergistically enhance Ca²⁺ response upon co-stimulation with both agonists.

[281] [Table 1]

Gene name	Full name	Other names	IDs
CXCR4	C-X-C Motif Chemokine Receptor 4	Leukocyte-Derived Seven Transmembrane Domain Receptor; Lipopolysaccharide-Associated Protein 3; Stromal Cell-Derived Factor 1 Receptor; Chemokine (C-X-C Motif) Receptor 4; LPS-Associated Protein 3 Seven Transmembrane Helix Receptor; C-X-C Chemokine Receptor Type 4; Neuropeptide Y Receptor Y3; Neuropeptide Y3 Receptor; Chemokine Receptor; Seven-Transmembrane-Segment Receptor, Spleen; Chemokine (C-X-C Motif), Receptor 4 (Fusin); SDF-1 Receptor; CD184 Antigen; Fusin; LAP-3; LESTR; NPYRL; FB22; HM89; LCR1; D2S201E; HSY3RR; NPYY3R; CXC-R4; CXCR-4; CD184; NPY3R; WHIMS; LAP3; NPYR; WHIM	GCID: GC02M136114HGNC: 2561Entrez Gene: 7852Ensembl: ENSG00000121966OMIM: 162643UniProtKB: P61073
ADCYAP1R1	ADCYAP Receptor Type I	Adenylate Cyclase Activating Polypeptide 1 (Pituitary) Receptor Type I; Pituitary Adenylate Cyclase-Activating Polypeptide Type 1	GCID: GC07P031058HGNC: 242Entrez Gene: 117Ensembl: ENSG00000078549OMIM: 102981UniProtKB:

		Receptor; PACAP Type I Receptor; PACAP Receptor 1; PACAP-R1; Pituitary Adenylate Cyclase Activating Polypeptide 1 Receptor Type I Hiphop; Pituitary Adenylate Cyclase-Activating Polypeptide Type I Receptor; PACAP-R-1; PACAPRI;PACAPR; PAC1R; PAC1	P41586
ADORA2B	Adenosine A2b Receptor	Adenosine Receptor A2b; ADORA2	GCID: GC17P015927HGNC: 264Entrez Gene: 136Ensembl: ENSG00000170425OM IM: 600446UniProtKB: P29275
ADORA3	Adenosine A3 Receptor	Adenosine Receptor A3; A3AR	GCID: GC01M111499HGNC: 268Entrez Gene: 140Ensembl: ENSG00000282608OM IM: 600445UniProtKB: P0DMS8
ADRB2	Adrenoceptor Beta 2	Adrenergic, Beta-2-, Receptor, Surface; Beta-2 Adrenoreceptor; Beta-2 Adrenoceptor; ADRB2R; B2AR; Adrenoceptor Beta 2, Surface; Adrenoceptor Beta 2 Surface; Beta-2 Adrenergic Receptor; Catecholamine Receptor;BETA2AR; ADRBR; BAR	GCID: GC05P148825HGNC: 286Entrez Gene: 154Ensembl: ENSG00000169252OM IM: 109690UniProtKB: P07550

C5AR1	Complement C5a Receptor 1	Complement Component 5a Receptor 1; Complement Component 5 Receptor 1 (C5a Ligand); C5a Anaphylatoxin Chemotactic Receptor 1; C5a Anaphylatoxin Chemotactic Receptor; Complement Component 5 Receptor 1; C5a Anaphylatoxin Receptor; C5a-R; C5R1; C5AR; CD88 Antigen; C5a Ligand; CD88; C5a	GCID: GC19P047290HGNC: 1338Entrez Gene: 728Ensembl: ENSG00000197405OM IM: 113995UniProtKB: P21730
CALCR	Calcitonin Receptor	CT-R; CTR1; CRT; CTR	GCID: GC07M093424HGNC: 1440Entrez Gene: 799Ensembl: ENSG00000004948OM IM: 114131UniProtKB: P30988
CHRM1	Cholinergic Receptor Muscarinic 1	Acetylcholine Receptor, Muscarinic 1; Muscarinic Acetylcholine Receptor M1; HM1; M1R; M1	GCID: GC11M062927HGNC: 1950Entrez Gene: 1128Ensembl: ENSG00000168539OM IM: 118510UniProtKB: P11229
EDNRB	Endothelin Receptor Type B	Endothelin Receptor Non-Selective Type; ET-BR; ET-B; ETRB; Endothelin Receptor Subtype B1; ABCDS; HSCR2; ETB1; ETBR; WS4A; HSCR; ETB	GCID: GC13M077895HGNC: 3180Entrez Gene: 1910Ensembl: ENSG00000136160OM IM: 131244UniProtKB: P24530
HRH1	Histamine Receptor H1	HH1R; H1R; Histamine Receptor, Subclass H1; Histamine H1 Receptor;	GCID: GC03P011113HGNC: 5182Entrez Gene:

		HisH1; H1-R	3269Ensembl: ENSG00000196639OM IM: 600167UniProtKB: P35367
MLNR	Motilin Receptor	G Protein-Coupled Receptor 38; GPR38; MTLR1; G-Protein Coupled Receptor 38; MTLR	GCID: GC13P049220HGNC: 4495Entrez Gene: 2862Ensembl: ENSG00000102539OM IM: 602885UniProtKB: O43193
NTSR1	Neurotensin Receptor 1	High-Affinity Levo- cabastine-Insensitive Neu- rotensin Receptor;Neurotensin Receptor 1 (High Affinity); NT-R-1; NTR1; NTRH; Neurotensin Receptor Type 1;NTRR; NTR	GCID: GC20P062708HGNC: 8039Entrez Gene: 4923Ensembl: ENSG00000101188OM IM: 162651UniProtKB: P30989
TACR3	Tachykinin Receptor 3	Neurokinin Beta Receptor; Neurokinin B Receptor; NK-3 Receptor; NK-3R; NK3RNKR; Neuromedin-K Receptor;TAC3RL; TAC3 R; HH11	GCID: GC04M103586HGNC: 11528Entrez Gene: 6870Ensembl: ENSG00000169836OM IM: 162332UniProtKB: P29371

[282] *GCID: Genecards identification

[283] HGNC: HUGO Gene Nomenclature Committee

[284]

[285] [Table 2]

Gene name	Full name	Other names	IDs
<i>APLNR</i>	Apelin Receptor	G-Protein Coupled Receptor HG11; Angiotensin II Receptor-Like 1; Angiotensin Receptor-Like 1; APJ (Apelin) Receptor; G Protein-Coupled Receptor APJ; G-Protein Coupled Receptor APJ; HG11 Orphan Receptor; APJ Receptor; AGTRL1; APJ; APJR; HG11	GCID: GC11M057233HG NC: 339Entrez Gene: 187Ensembl: ENSG00000134817 OMIM: 600052UniProtKB: P35414
<i>CCR5</i>	Chemokine (C-C Motif) Receptor 5	C-C Chemokine Receptor Type 5; Chemokine Receptor CCR5; Chemokine (C-C Motif) Receptor 5 (Gene/Pseudogene); C-C Motif Chemokine Receptor 5 A159A; Chemokine Receptor CCR5 Delta32; CD195 Antigen; HIV-1 Fusion Coreceptor; CC-CKR-5; ChemR13; CMKBR5; CCR-5; C-C CKR-5; CCCKR5; IDDM22; CD195; CKR-5; CKR5	GCID: GC03P046382HGN C: 1606Entrez Gene: 1234Ensembl: ENSG00000160791 OMIM: 601373UniProtKB: P51681
<i>GALR1</i>	Galanin Receptor 1	GALNR1; GAL1-R; GALR-1; GALNR; Galanin Receptor Type 1	GCID: GC18P077250HGN C: 4132Entrez Gene: 2587Ensembl: ENSG00000166573 OMIM: 600377UniProtKB: P47211
<i>PTGER2</i>	Prostaglandin E Receptor 2	Prostaglandin E Receptor 2 (Subtype EP2), 53kDa;	GCID: GC14P052314HGN

		Prostaglandin E Receptor 2 (Subtype EP2), 53kD; PGE2 Receptor EP2 Subtype; PGE Receptor EP2 Subtype; Prostanoid EP2 Receptor; Prostaglandin E2 Receptor EP2 Subtype; EP2	C: 9594Entrez Gene: 5732Ensembl: ENSG00000125384 OMIM: 176804UniProtKB: P43116
<i>PTGER3</i>	Prostaglandin E Receptor 3	Prostaglandin E Receptor 3 (Subtype EP3); PGE2 Receptor EP3 Subtype; Prostanoid EP3 Receptor;PGE2-R; Prostaglandin E Receptor EP3 Subtype 3 Isoform; Prostaglandin E2 Receptor EP3 Subtype; Prostaglandin Receptor (PGE-2); PGE Receptor, EP3 Subtype; PGE Receptor EP3 Subtype; EP3-III; EP3-II; EP3-IV;EP3-VI; EP3-I; EP3e; EP3	GCID: GC01M070852HG NC: 9595Entrez Gene: 5733Ensembl: ENSG00000050628 OMIM: 176806UniProtKB: P43115
<i>SSTR2</i>	Somatostatin Receptor 2	SRIF-1; SS2R; Somatostatin Receptor Type 2; SS-2-R; SS2-R	GCID: GC17P073165HGN C: 11331Entrez Gene: 6752Ensembl: ENSG00000180616 OMIM: 182452UniProtKB: P30874

[286] *GCID: Genecards identification

[287] HGNC: HUGO Gene Nomenclature Committee

[288]

[289] Further information regarding the GPCRs evaluated herein as heteromers with CXCR4 are detailed below:

[290] (1) ADCYAP1R1 -- Pituitary adenylate cyclase-activating polypeptide type I receptor also known as PAC1, is a protein that in humans is encoded by the

ADCYAP1R1 gene. This receptor binds pituitary adenylate cyclase activating peptide. ADCYAP1R1 is a membrane-associated protein and shares significant homology with members of the G-protein coupled class B glucagon/secretin receptor family. This receptor mediates diverse biological actions of adenylate cyclase activating polypeptide 1 and is positively coupled to adenylate cyclase (Vaudry et al., 2000). This is a receptor for PACAP-27 and PACAP-38. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase. ADCYAP1R1 may regulate the release of adrenocorticotropin, luteinizing hormone, growth hormone, prolactin, epinephrine, and catecholamine. ADCYAP1R1 may play a role in spermatogenesis and sperm motility. ADCYAP1R1 causes smooth muscle relaxation and secretion in the gastrointestinal tract (Ogi et al., 1993).

[291] ADCYAP1R1 is expressed in the adrenal medulla, pancreatic acini, uterus, myenteric plexus and brain (Reubi, 2000; Reubi et al., 2000). It is also expressed in the trigeminal, otic and superior cervical ganglia (prejunctional) and cerebral arteries (postjunctional) (Knutsson and Edvinsson, 2002). Diseases associated with ADCYAP1R1 include post-traumatic stress disorder (Lowe et al., 2015) and anxiety disorder. Among its related pathways are G alpha(s) signalling events and RET signaling (Cooper et al., 2013).

[292] (2) ADORA2B -- The adenosine A2B receptor, also known as ADORA2B, is a G-protein coupled adenosine receptor, and also denotes the human adenosine A2b receptor gene which encodes it. Adenosine functions as a signaling molecule through the activation of four distinct adenosine receptors - ADORA1, ADORA2A, ADORA2B, and ADORA3, also called as A1, A2A, A2B, and A3, respectively. This integral membrane protein stimulates adenylate cyclase activity in the presence of adenosine. This protein also interacts with netrin-1, which is involved in axon elongation. These receptors are widely expressed and have been implicated in several biological functions, both physiological and pathological. Both ADORA2A and ADORA2B receptors are important in cancer immunology. The tumor microenvironment is hypoxic, which promotes the expression of CD39 and CD73 by immune cells. CD39 and CD73 are ectonucleotidases that convert ATP to adenosine, elevating concentrations of adenosine locally. Adenosine is a crucial mediator of altered immune function in cancer as it binds ADORA2A and ADORA2B receptors on lymphocytes silencing the antitumor immune response (Chen et al., 2013).

[293] ADORA2B has a role in regulating CXCR4 expression *in vivo* and in protecting against vascular lesion formation (Yang et al., 2008) and promotes progression of human oral cancer (Kasama et al., 2015). Identification of a pharmacologically tractable Fra-1/ADORA2B axis promoting breast cancer metastasis (Desmet et al., 2013). ADORA2B receptors display high expression levels in the cecum, colon and

- bladder, with lower levels in the lung, blood vessels and eye (Fagerberg et al., 2014).
- [294] (3) ADORA3 -- The adenosine A3 receptor, also known as ADORA3, is an adenosine receptor, but also denotes the human gene encoding it. ADORA3 receptors are G-protein coupled receptors that couple to Gi/Gq and are involved in a variety of intracellular signaling pathways and physiological functions. It mediates a sustained cardioprotective function during cardiac ischemia, it is involved in the inhibition of neutrophil degranulation in neutrophil-mediated tissue injury, it has been implicated in both neuroprotective and neurodegenerative effects, and it may also mediate both cell proliferation and cell death (Gao et al., 2008; Miwatashi et al., 2008).
- [295] Cordycepin induces apoptosis in human bladder cancer cells via activation of ADORA3 receptors (Cao et al., 2017). Jafari et al. showed that ADORA3 receptor agonist inhibited survival of breast cancer stem cells via GLI-1 and ERK1/2 pathway (Jafari et al., 2017). ADORA3 is broadly expressed in adrenal (RPKM 4.4), small intestine (RPKM 2.9) and 20 other tissues containing brain, bladder, lymph node and colon (Fagerberg et al., 2014).
- [296] (4) ADRB2 -- The beta-2 adrenergic receptor (β 2 adrenoreceptor), also known as ADRB2, is a cell membrane-spanning beta-adrenergic receptor that interacts with epinephrine, a hormone and neurotransmitter (ligand synonym, adrenaline) whose signaling, via a downstream L-type calcium channel interaction, mediates physiologic responses such as smooth muscle relaxation and bronchodilation (Gregorio et al., 2017). ADRB2 functions in muscular system such as smooth muscle relaxation, motor nerve terminals, glycogenolysis and in circulatory system such as heart muscle contraction, cardiac output increase. In the normal eye, beta-2 stimulation by salbutamol increases intraocular pressure via net. In digestive system, the ADRB2 induces glycogenolysis and gluconeogenesis in liver and insulin secretion from pancreas (Fitzpatrick, 2004).
- [297] ADRB2 signaling in the cardiac myocyte is modulated by interactions with CXCR4 (LaRocca et al., 2010). Norepinephrine attenuates CXCR4 expression and the corresponding invasion of MDA-MB-231 breast cancer cells via ADRB2 (Wang et al., 2015a). ADRB2 is expressed in several cancers such as pancreatic, prostate (Braadland et al., 2014; Xu et al., 2017), renal and breast cancer (Choy et al., 2016).
- [298] (5) C5AR1 -- The C5a receptor also known as complement component 5a receptor 1 (C5AR1) or CD88 (Cluster of Differentiation 88) is a G protein-coupled receptor for C5a. It functions as a complement receptor. C5AR1 modulates inflammatory responses, obesity, development and cancers. (Gerard and Gerard, 1994).
- [299] The C5AR1 is expressed on granulocytes, monocytes, dendritic cells, hepatoma-derived cell line HepG2, astrocytes, microglia (Klos et al., 2013). C5AR1 is related in several diseases such as inflammatory bowel disease, rheumatoid arthritis, psoriasis,

experimental allergic encephalomyelitis, multiple sclerosis, meningitis, and brain trauma (Lee et al., 2008).

- [300] (6) CALCR -- The calcitonin receptor (CT) is a G protein-coupled receptor that binds the peptide hormone calcitonin and is involved in maintenance of calcium homeostasis, particularly with respect to bone formation and metabolism (Dacquin et al., 2004; Davey et al., 2008). CT works by activating the G-proteins Gs and Gq often found on osteoclasts, on cells in the kidney, and on cells in a number of regions of the brain. It may also affect the ovaries in women and the testes in men. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase. The calcitonin receptor is thought to couple to the heterotrimeric guanosine triphosphate-binding protein that is sensitive to cholera toxin. The physiological effects of CT such as inhibition of osteoclast mediated bone resorption or increased Ca²⁺ excretion by the kidney are mediated by high affinity CTs (Albrandt et al., 1995).
- [301] Loss of function mutation in CT identifies highly aggressive glioblastoma with poor outcome (Pal et al., 2018). Expression of CTRs in mammary gland, cartilage, nose and ear has been reported. The developmental regulation of CALCR gene expression in a variety of the above tissues suggests a role for CTRs in the morphogenesis of these tissues (Pondel, 2000).
- [302] (7) CHRM1 -- The muscarinic cholinergic receptors belong to a larger family of G protein-coupled receptors. The functional diversity of these receptors is defined by the binding of acetylcholine and includes cellular responses such as adenylate cyclase inhibition, phosphoinositide degeneration, and potassium channel mediation. Muscarinic receptors influence many effects of acetylcholine in the central and peripheral nervous system (Rang, 2003). The muscarinic cholinergic receptor 1 is involved in mediation of vagally-induced bronchoconstriction and in the acid secretion of the gastrointestinal tract. It is predominantly found bound to G proteins of class Gq that use upregulation of phospholipase C and, therefore, inositol trisphosphate and intracellular calcium as a signaling pathway (Qin et al., 2011).
- [303] Muscarinic receptors are widely distributed throughout the body and control distinct functions according to location and subtype (CHRM1 - CHRM5). They are predominantly expressed in the parasympathetic nervous system where they exert both inhibitory and excitatory effects. This receptor is found mediating slow excitatory post-synaptic potential at the ganglion in the postganglionic nerve, is common in exocrine glands and in the CNS (Johnson, 2002). Diseases associated with CHRM1 include asthma and heart block, congenital and Pelizaeus-Merzbacher disease. Activators of CHRM1 muscarinic acetylcholine receptors (mAChRs) may provide novel treatments for schizophrenia and Alzheimer's disease (Marlo et al., 2009).
- [304] (8) EDNRB -- Endothelin receptor type B (ETB) receptor was cloned shortly after

the ETA receptor (EDBRA). Similarly to the EDNRA, it belongs to class A of G protein-coupled heptahelical receptors with an external amino terminus and internal carboxy terminus and binding sites intrinsic to the heptahelical portions of the receptor. Similar to the EDNRA, the endogenous agonist ET-1 has a high affinity for the EDNRB. The pharmacology of the EDNRB receptor is somewhat richer than that of the EDNRA in that multiple agonists of the EDNRB are recognized, including sarafotoxin 6c (S6c) and IRL1620. Selective antagonists of the EDNRB include BQ788, A192621, RES7011, and IRL2500. The EDNRB has been localized to cardiovascular tissues, the pulmonary system, neurons, bone, pancreas, and kidney (Watts, 2010).

[305] EDNRB is a G protein-coupled receptor which activates a phosphatidylinositol-calcium second messenger system. Its ligand, endothelin, consists of a family of three potent vasoactive peptides: ET1, ET2, and ET3. In melanocytic cells the EDNRB gene is regulated by the microphthalmia-associated transcription factor. Mutations in either gene are links to Waardenburg syndrome (Sato-Jin et al., 2008). The multigenic disorder, Hirschsprung disease type 2, is due to mutation in EDNRB gene (Tanaka et al., 1998). Diseases associated with EDNRB include Waardenburg syndrome, Type 4A and Abcd syndrome. Among its related pathways are calcium signaling pathway and prostaglandin synthesis and regulation of Waardenburg Syndrome (Sato-Jin et al., 2008).

[306] (9) HRH1 -- The H1 receptor is a histamine receptor belonging to the family of rhodopsin-like G-protein-coupled receptors. This receptor is activated by the biogenic amine histamine. The HRH1 is linked to an intracellular G-protein (Gq) that activates phospholipase C and the inositol triphosphate (IP3) signaling pathway. Antihistamines, which act on this receptor, are used as anti-allergy drugs. The crystal structure of the receptor has been determined and used to discover new HRH1 ligands in structure-based virtual screening studies (de Graaf et al., 2011). A subunit of the G-protein subsequently dissociates and affects intracellular messaging including downstream signaling accomplished through various intermediaries such as cyclic AMP, cyclic GMP, calcium, and nuclear factor kappa B. It functions in immune-cell chemotaxis, pro-inflammatory cytokine production, expression of cell adhesion molecules, and other allergic and inflammatory conditions (Canonica and Blaiss, 2011).

[307] In peripheral tissues, the HRH1 mediates the contraction of smooth muscles, increase in capillary permeability due to contraction of terminal venules, and catecholamine release from adrenal medulla, as well as mediating neurotransmission in the central nervous system. It is also known to contribute to the pathophysiology of allergic diseases such as atopic dermatitis, asthma, anaphylaxis and allergic rhinitis (Xie and He, 2005). It is expressed in smooth muscles, on vascular endothelial cells, in the

heart, and in the central nervous system (de Graaf et al., 2011).

- [308] (10) MLNR -- Motilin receptor is a G protein-coupled receptor that binds motilin. Motilin in turn is an intestinal peptide that stimulates contraction of gut smooth muscle. Also, the MLNR mediates progastric effects. MLNR is an important therapeutic target for the treatment of hypomotility disorders (Depoortere, 2001).
- [309] It is found at its highest concentrations in the nerves of the antral wall of the stomach and is also found at significant levels throughout the smooth muscle of the upper gastrointestinal tract (Kitazawa et al., 1995). Diseases associated with MLNR include gastroparesis and diabetic autonomic neuropathy (Kitazawa et al., 1997).
- [310] (11) NTSR1 -- Neurotensin receptor 1 belongs to the large superfamily of G-protein coupled receptors. Neurotensin is a 13-amino acid peptide originally isolated from hypothalamic and later from intestines of bovine. In the brain, neurotensin is exclusively found in nerve cells, fibers, and terminals, whereas the majority of peripheral neurotensin is found in the endocrine N-cells located in the intestine. Central or peripheral injections of neurotensin produce completely different pharmacological effects indicating that the peptide does not cross the blood-brain barrier. NTSR1 mediates the multiple functions of neurotensin, such as hypotension, hyperglycemia, hypothermia, antinociception, and regulation of intestinal motility and secretion (Vincent, 1995).
- [311] Signaling is effected via G proteins that activate a phosphatidylinositol-calcium second messenger system. Signaling leads to the activation of downstream MAP kinases and protects cells against apoptosis (Heakal et al., 2011). Swift et al. showed altered expression of neurotensin receptors is associated with the differentiation state of prostate cancer (Swift et al., 2010).
- [312] (12) PTGER2 -- Prostaglandin E2 receptor 2, also known as EP2, is a prostaglandin receptor for prostaglandin E2 (PGE2) encoded by the human gene PTGER2. PTGER2 is classified as a relaxant type of prostanoid receptor based on its ability, upon activation, to relax certain types of smooth muscle. When initially bound to PGE2 or any other of its agonists, it mobilizes G proteins containing the G α s-G $\beta\gamma$ complex. The G α s-G $\beta\gamma$ complexes dissociate into their G α s and G $\beta\gamma$ subunits which in turn regulate cell signaling pathways. In particular, G α s stimulates adenylyl cyclase to raise cellular levels of cAMP thereby activating PKA; PKA activates various types of signaling molecules such as the transcription factor CREB which lead to different types of functional responses depending on cell type (Markovic et al., 2017; Woodward et al., 2011). PTGER2 also activates the GSK-3 pathway which regulates cell migratory responses and innate immune responses including pro-inflammatory cytokine and interleukin production and Beta-catenin pathway which regulates not only cell-cell adhesion but also activates the Wnt signaling pathway which, in turn, stimulates the transcription of genes responsible for regulating cell migration and proliferation

- (Woodward et al., 2011).
- [313] The PTGER2 receptor can act as a tumor promoter. PTGER2 gene knockout mice have less lung, breast, skin, and colon cancers following exposure to carcinogens. Knockout of this gene in mice with the adenomatous polyposis coli mutation also causes a decrease in the size and number of pre-cancerous intestinal polyps that the animals develop. These effects are commonly ascribed to the loss of PTGER2-mediated: vascular endothelial growth factor production and thereby of tumor vascularization; regulation of endothelial cell motility and survival; interference with transforming growth factor- β 's anti-cell proliferation activity; and, more recently, regulation of host anti-tumor immune responses (O'Callaghan and Houston, 2015).
- [314] PTGER2 is widely distributed in humans. Its protein is expressed in human small intestine, lung, media of arteries and arterioles of the kidney, thymus, uterus, brain cerebral cortex, brain striatum, brain hippocampus, corneal epithelium, corneal chorio-capillaries, Myometriuml cells, eosinophiles, sclera of the eye, articular cartilage, the corpus cavernosum of the penis, and airway smooth muscle cells; its mRNA is expressed in gingival fibroblasts, monocyte-derived dendritic cells, aorta, corpus cavernosum of the penis, articular cartilage, airway smooth muscle, and airway epithelial cells. In rats, the receptor protein and/or mRNA has been found in lung, spleen, intestine, skin, kidney, liver, long bones, and rather extensively throughout the brain and other parts of the central nervous system (Yagami et al., 2016).
- [315] Pre-clinical studies indicate that PTGER2 may be a target for treating and/or preventing particular human disorders involving: allergic diseases such as asthma (particular aspirin and nonsteroidal inflammatory drug-induced asthma syndromes) and rhinitis (Machado-Carvalho et al., 2014); glaucoma (Doucette and Walter, 2017); various diseases of the nervous system (Yagami et al., 2016); fractures, osteoporosis, and other bone abnormalities (Li et al., 2007); pulmonary fibrosis; certain forms of malignant disease such as colon cancer including those that arise from Adenomatous polyposis coli mutations (O'Callaghan and Houston, 2015); and salt-sensitive forms of hypertension (Yang and Du, 2012); This receptor has also been suggested to be a target for contraception (Sugimoto et al., 2015).
- [316] (13) TACR3 -- Tachykinin receptor 3, also known as TACR3, function as receptors for tachykinins. The tachykinins are a family of peptides that comprise substance P (SP), neurokinin A (NKA), neurokinin B (NKB) and the species divergent endokinins including endokinin B (EKB) in humans. These tachykinins are encoded on three different genes, preprotachykinin 1 (TAC1) encoding SP and NKA, TAC3 encoding NKB and TAC4 encoding EKB. Three tachykinin receptors have been identified, which interact with these tachykinins: TACR1, TACR2, and TACR3, also called as NK1, NK2 and NK3, respectively, whereby SP and EKB show the greatest potency for

TACR1, NKA for TACR2 and NKB for TACR3. Receptor affinities are specified by variations in the 5'-end of the sequence. Tachykinin receptor-3 (TACR3) is the mediator of biologic actions encoded by the C-terminal sequence of tachykinins, for which neurokinin B is a more potent agonist than neurokinin A or substance P (Page et al., 2003).

- [317] It is reported that four human pedigrees with severe congenital gonadotropin deficiency and pubertal failure in which all affected individuals are homozygous for loss-of-function mutations in TAC3 (encoding Neurokinin B) or its receptor TACR3 (encoding NK3R) (Topaloglu et al., 2009). NKB, a member of the substance P-related tachykinin family, is known to be highly expressed in hypothalamic neurons that also express kisspeptin (Goodman et al., 2007), a recently identified regulator of gonadotropin-releasing hormone secretion (Gianetti and Seminara, 2008).
- [318] (14) APLNR -- The apelin receptor (also known as the APJ receptor) is a G protein-coupled receptor which binds apelin and Apela/ELABELA/Toddler (Medhurst et al., 2003). Receptor for apelin receptor early endogenous ligand (APELA) and apelin (APLN) hormones coupled to G proteins that inhibit adenylate cyclase activity. APLNR plays a key role in early development such as gastrulation and heart morphogenesis by acting as a receptor for APELA hormone. APLNR plays also a role in various processes in adults such as regulation of blood vessel formation, blood pressure, heart contractility, and heart failure by acting as a receptor for APLN hormone. A decade of investigations has shown that the apelinergic system has a broad range of biological functions, playing an important role particularly in maintaining homeostasis of the cardiovascular system and fluid metabolism. The activation of APLNR causes a broad spectrum of biochemical changes including cAMP suppression, phosphorylation of protein kinase B (Akt), ERK1/25 and p70S6K, calcium mobilization⁷ and nitric oxide synthase (NOS). However, the specificity linking these biochemical activities to a certain biological function is yet to be determined (Wang et al., 2015c).
- [319] Also, it functions in embryonic and tumor angiogenesis (Wu et al., 2017) and as a human immunodeficiency virus (HIV-1) coreceptor (Zhou et al., 2003). Both APLNR and apelin are expressed in many tissues including heart, lung, endothelium, kidney and brain (Wang et al., 2015c).
- [320] (15) CCR5 -- C-C chemokine receptor type 5, also known as CCR5 or CD195, is a protein on the surface of white blood cells that is involved in the immune system as it acts as a receptor for chemokines. This is the process by which T cells are attracted to specific tissue and organ targets. Many forms of HIV, the virus that causes AIDS, initially use CCR5 to enter and infect host cells. Certain individuals carry a mutation known as CCR5-Δ32 in the CCR5 gene, protecting them against these strains of HIV.

Certain populations have inherited the Delta 32 mutation resulting in the genetic deletion of a portion of the CCR5 gene. Homozygous carriers of this mutation are resistant to M-tropic strains of HIV-1 infection (Hutter et al., 2009). CCR5's cognate ligands include CCL3, CCL4 (also known as MIP 1 α and 1 β , respectively), and CCL3L1. CCR5 furthermore interacts with CCL5 (Struyf et al., 2001).

- [321] It is likely that CCR5 plays a role in inflammatory responses to infection, though its exact role in normal immune function is unclear. Regions of this protein are also crucial for chemokine ligand binding, functional response of the receptor, and HIV co-receptor activity (Barmania and Pepper, 2013). CCR5 is predominantly expressed on T cells, macrophages, dendritic cells, eosinophils, microglia and a subpopulation of ether breast or prostate cancer cells (Sicoli et al., 2014; Velasco-Velazquez et al., 2012).
- [322] (16) GALR1 -- Galanin receptor 1 (GAL1) is a G-protein coupled receptor encoded by the GALR1 gene. The ubiquitous neuropeptide galanin controls numerous functions such as endocrine secretions, intestinal motility, and behavioral activities. These regulatory effects of galanin are mediated through the interaction with specific membrane receptors and involve the pertussis toxin-sensitive guanine nucleotide binding proteins Gi/Go as transducing elements (Habert-Ortoli et al., 1994).
- [323] GALR1 has anti-proliferative effects in oral squamous cell carcinoma. GALR1 protein and mRNA expression and GAL secretion were detected at variable levels in immortalized human oral keratinocytes and human oropharyngeal squamous cell carcinoma cell lines. Upon competitive inhibition of GALR1, proliferation was up-regulated in immortalized and malignant keratinocytes (Henson et al., 2005)
- [324] Stevenson et al. report the stem cell marker and regulator, galanin and GALR1 as key determinants of drug resistance and potential therapeutic targets for combating drug resistance. Mechanistically, they identify a novel role for the GALR1-galanin receptor-ligand axis as an important upstream regulator of expression of the anti-apoptotic protein FLIPL. Clinically, galanin mRNA was found to be overexpressed in colorectal tumours, and notably, high galanin mRNA expression correlated with poor disease-free survival in early stage disease (Stevenson et al., 2012). GALR1 is widely expressed in the brain and spinal cord, as well as in peripheral sites such as the small intestine and heart (Fagerberg et al., 2014)
- [325] (17) PTGER3 -- Prostaglandin EP3 receptor (53kDa), also known as EP3, is a prostaglandin receptor for prostaglandin E2 (PGE2) encoded by the human gene PTGER3; it is one of four identified EP receptors, the others being PTGER1, PTGER2, and PTGER4, also called as EP1, EP2, and EP4, respectively, all of which bind with and mediate cellular responses to PGE2 and also, but generally with lesser affinity and responsiveness, certain other prostanoids. PTGER3 activation promotes duodenal secretion in mice; this function is mediated by PTGER3 activation in humans. EP

receptor functions can vary with species and most of the functional studies cited here have not translated their animal and tissue models to humans (Moreno, 2017).

- [326] Studies of the direct effects of PTGER3 activation on cancer in animal and tissue models give contradictory results suggesting that this receptor does not play an important role in carcinogenesis. However, some studies suggest an indirect pro-carcinogenic function for the PTGER3: The growth and metastasis of implanted Lewis lung carcinoma cells, a mouse lung cancer cell line, is suppressed in PTGER3-deficient mice. This effect was associated with a reduction in the levels of Vascular endothelial growth factor and matrix metalloproteinase-9 expression in the tumor's stroma; expression of the pro-lymphangiogenic growth factor, VEGF-C and its receptor, VEGFR3; and a tumor-associated angiogenesis and lymphangiogenesis (O'Callaghan and Houston, 2015).
- [327] PTGER3 is widely distributed in humans. Its protein and/or mRNA is expressed in kidney (i.e. glomeruli, Tamm-Horsfall protein negative late distal convoluted tubules, connecting segments, cortical and medullary collecting ducts, media and endothelial cells of arteries and arterioles); stomach (vascular smooth muscle and gastric fundus mucosal cells); thalamus (anterior, ventromedial, laterodorsal, paraventricular and central medial nuclei); intestinal mucosal epithelia at the apex of crypts; myometrium (stromal cells, endothelial cells, and, in pregnancy, placenta, chorion, and amnion); mouth gingival fibroblasts; and eye (corneal endothelium and keratocytes, trabecular cells, ciliary epithelium, and conjunctival and iridal stroma cells, and retinal Müller cells) (Norel, 2016).
- [328] (18) SSTR2 -- Somatostatin receptor type 2 is a protein that in humans is encoded by the SSTR2 gene. Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein-coupled receptors that are expressed in a tissue-specific manner. SSTR2 has been shown to interact with shank2 (Zitzer et al., 1999).
- [329] SSTR2 encodes somatostatin receptor that can inhibit the cell proliferation of solid tumors. SSTR2 promoter hypermethylation is associated with the risk and progression of laryngeal squamous cell carcinoma in males (Shen et al., 2016). Most pituitary adenomas express SSTR2, but other somatostatin receptors are also found (Miller et al., 1995). Somatostatin analogs (i.e. Octreotide, Lanreotide) are used to stimulate this receptors, and thus to inhibit further tumor proliferation (Zatelli et al., 2007). SSTR2 is expressed in highest levels in cerebrum and kidney (Fagerberg et al., 2014).
- [330] The term "inhibitor" as used herein refers to molecule that inhibits or suppresses the enhanced function of CXCR4-GPCR_x heteromer. Non-limiting examples of the inhibitor of the invention that can be used for treatment, amelioration, or prevention of

a cancer or related symptoms include GPCR α antagonist, GPCR α inverse agonist, GPCR α positive and negative allosteric modulator, CXCR4-GPCR α heteromer-specific antibody or its antigen binding portions including single-domain antibody-like scaffolds, bivalent ligands which have a pharmacophore selective for CXCR4 joined by a spacer arm to a pharmacophore selective for GPCR α , bispecific antibody against CXCR4 and GPCR α , radiolabeled CXCR4 ligand linked with GPCR α ligand, and small molecule ligands that inhibit heteromer-selective signaling. Certain examples of inhibitors against GPCR α that form heteromers with CXCR4 and enhance Ca²⁺ response upon co-stimulation with both agonists are listed in Table 3.

[331] The term "antagonist" as used herein refers to a type of receptor ligand or drug that blocks or dampens a biological response by binding to and blocking a receptor, also called blockers. Antagonists have affinity but no efficacy for their cognate receptors, and their binding disrupts the interaction and inhibit the function of an agonist or inverse agonist at the cognate receptors. Certain examples of antagonists against GPCR α that form heteromers with CXCR4 and enhance Ca²⁺ response upon co-stimulation with both agonists are listed in Table 3.

[332] Table 3. Examples of inhibitors against GPCR α that form heteromers with CXCR4 and enhances Ca²⁺ response upon co-stimulation with both agonists.

[333] [Table 3]

Gene name	Antagonists/Inverse agonists	Antibodies/nobodies/i-bodies/others
CXCR4	ALX40-4C, AMD070 (AMD11070, X4P-001), AMD3100 (plerixafor), AMD3465, ATI 2341, BKT140 (BL-8040; TF14016; 4F-Benzoyl-TN14003), CTCE-9908, CX549, D-[Lys3] GHRP-6, FC122, FC131, GMI-1359, GSK812397, GST-NT21MP, isothioureia-1a, isothioureia-1t (IT1t), KRH-1636, KRH-3955, LY2510924, MSX-122, N-[¹¹ C]Methyl-AMD3465, POL6326, SDF-1 1-9[P2G] dimer, SDF1 P2G, T134, T140, T22, TC 14012, TG-0054 (Burixafor), USL311, viral macrophage inflammatory protein-II (vMIP-II), WZ811, [⁶⁴ Cu]-AMD3100, [⁶⁴ Cu]-AMD3465, [⁶⁸ Ga]pentixafor, [⁹⁰ Y]pentixather, [^{99m} Tc]O ₂ -AMD3100, [¹⁷⁷ Lu]pentixather, and 508MCl (Compound 26).	AD-114, AD-114-6H, AD-114-Im7-FH, AD-114-PA60 0-6H, ALX-0651, LY2624587, PF-06747143, ulocuplumab (MDX1338/BMS-936564), 12G5, 238D2, and 238D4
ADCYAP1R1	M65, Max.d.4, MK-0893, N-stearyl-[Nle ¹⁷] neurotensin-(6-11)/VIP-(7-28), PACAP-(6-38), and PG 97-269	
ADORA2B	3-isobutyl-8-pyrrolidinoxanthine, alloxazine, AS16, AS70, AS74, AS94, AS95, AS96, AS99, AS100, AS101, ATL802, BW-A1433, caffeine, CGS 15943, CPX, CSC, CVT-6883, DAX, DEPX, derenofylline, DPCPX, FK-453, I-ABOPX, istradefylline, KF26777, LAS38096, LUF5981, MRE 2029F20, MRE 3008F20, MRS1191, MRS1220, MRS1523, MRS1706, MRS1754, MSX-2, OSIP339391, pentoxifylline, preladenant, PSB-10, PSB-11, PSB36, PSB603, PSB-0788, PSB1115, rolofylline, SCH 58261, SCH442416, ST-1535, theophylline, tonapofylline, vipadenant, xanthine amine congener, XCC, and ZM-241385.	
ADORA3	ATL802, BW-A1433, caffeine, CGS 15943, CSC, CVT-6883, derenofylline, dextriguldipine, DPCPX, FK-453, flavanone, flavone, galangin, I-ABOPX, istradefylline, KF26777, LAS38096, LUF5981, MRE 2029F20, MRE 3008F20, MRE 3010F20, MRS1041, MRS1042,	

	MRS1067, MRS1088, MRS1093, MRS1097, MRS1177, MRS1186, MRS1191, MRS1191, MRS1220, MRS1476, MRS1486, MRS1505, MRS1523, MRS1754, MRS928, MSX-2, nicardipine, preladenant, PSB-10, PSB-11, PSB36, PSB603, PSB1115, rollofylline, sakuranetin, SCH 58261, SCH442416, ST-1535, theophylline, tonapofylline, vipadenant, visnagin, VUF5574, VUF8504, VUF8507, xanthine amine congener, and ZM-241385	
ADRB2	Alprenolol, atenolol, betaxolol, bupranolol, butoxamine, carazolol, carvedilol, CGP 12177, cicloprolol, ICI 118551, ICYP, labetalol, levobetaxolol, levobunolol, LK 204-545, metoprolol, nadolol, NIHP, NIP, propafenone, propranolol, sotalol, SR59230A, and timolol.	
C5AR1	A8 ^{Δ71-73} , AcPhe-Orn-Pro-D-Cha-Trp-Arg, avacopan, C089, CHIPS, DF2593A, JPE1375, L-156,602, NDT9520492, <i>N</i> -methyl-Phe-Lys-Pro-D-Cha-Trp-D-Arg-CO ₂ H, PMX205, PMX53, RPR121154, and W54011.	
CALCR	α -CGRP-(8-37) (human), AC187, CT-(8-32) (salmon), and olcegepant.	
CHRM1	3-Quinuclidinyl benzilate (QNB), 4-DAMP, aclidinium, AE9C90CB, AFDX384, amitriptyline, AQ-RA 741, atropine, benzatropine, biperiden, darifenacin, di-cyclomine, dosulepin, ethopropazine, glycopyrrolate, guanylpirenzepine, hexahydrodifenidol, hexahydrosiladifenidol, hexocyclium, himbacine, ipratropium, lithocholylcholine, methoctramine, ML381, muscarinic toxin 1, muscarinic toxin 2, muscarinic toxin 3, N-methyl scopolamine, otenzepad, oxybutynin, <i>p</i> -F-HHSiD, pirenzepine, propantheline, (R,R)-quinuclidinyl-4-fluoromethyl-benzilate, scopolamine, silahexocyclium, solifenacin, telenzepine, tiotropium, tolterodine, trihexyphenidyl, tripitramine, UH-AH 37, umeclidinium, and VU0255035.	
EDNRB	A192621, ambrisentan, atrasentan, bosentan (RO	

	470203, Tracleer);, BQ788, IRL 2500, K-8794, macitentan, RES7011, Ro 46-8443, SB209670, SB217242 (enrasentan), TAK 044, and tezosentan (RO610612).	
HRH1	(-)-chlorpheniramine, (+)-chlorpheniramine, (-)- <i>trans</i> -H ₂ -PAT, (+)- <i>cis</i> -H ₂ -PAT, (+)- <i>trans</i> -H ₂ -PAT, (±)- <i>cis</i> -H ₂ -PAT, (±)- <i>trans</i> -H ₂ -PAT, (<i>R</i>)-cetirizine, (<i>S</i>)-cetirizine, 9-OH-risperidone, A-317920, A-349821, ABT-239, alimemazine, amitriptyline, aripiprazole, arpromidine, asenapine, astemizole, AZD3778, azelastine, BU-E 47, cetirizine, chlorpheniramine, chlorpromazine, ciproxifan, clemastine, clobenpropit, clozapine, conessine, cyclizine, cyproheptadine, desloratadine, diphenhydramine, dosulepin, doxepin, epinastine, fexofenadine, fluphenazine, fluspirilene, haloperidol, hydroxyzine, imipromidine, INCB-38579, JNJ-39758979, ketotifen, loratadine, loxapine, MK-0249, molindone, olanzapine, perphenazine, pimozide, pipamperone, pitolisant, promethazine, pyrilamine, quetiapine, risperidone, sertindole, terfenadine, thioridazine, thiothixene, trifluoperazine, tripeleennamine, triprolidine, ziprasidone, and zotepine.	
MLNR	GM-109, MA-2029, and OHM-11526.	
NTSR1	Meclinetant, SR48527, SR48692, and SR142948A	
TACR3	[Trp ⁷ , β-Ala ⁸] neurokinin A-(4-10), AZD2624, FK 224, GR138676, GSK 172981, GSK 256471, N',2-diphenylquinoline-4-carbohydrazide 8m, N',2-diphenylquinoline-4-carbohydrazide, osanetant, PD 154740, PD 161182, PD157672, saredutant, SB 218795, SB 222200, SB 235375, SCH 206272, SSR 146977, and talnetant.	

[334] In some embodiments, the method for treating cancer in a patient having a cell containing a CXCR4-GPCR_x heteromer, or the method of suppressing enhanced downstream signaling from a CXCR4-GPCR_x heteromer in a cell of a patient suffering from cancer, as disclosed herein, may comprise administering, or the pharmaceutical kit or pharmaceutical kit, comprise, administering to the patient an inhibitor

or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein: i) the CXCR4-GPCR_x heteromer has enhanced downstream signaling; and ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient. In some embodiments, the pharmaceutical kit or pharmaceutical composition for use in treating cancer in a patient having a cell containing a CXCR4-GPCR_x heteromer, as disclosed herein, may comprise an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein the CXCR4-GPCR_x heteromer has enhanced downstream signaling. For example, in some embodiments, the CXCR4 inhibitor is an antagonist of CXCR4, an inverse agonist of CXCR4, a partial antagonist of CXCR4, an allosteric modulator of CXCR4, an antibody of CXCR4, an antibody fragment of CXCR4, a ligand of CXCR4, or an antibody-drug conjugate of CXCR4. For example, in some embodiments, the GPCR_x inhibitor is an antagonist of GPCR_x, an inverse agonist of GPCR_x, a partial antagonist of GPCR_x, an allosteric modulator of GPCR_x, an antibody of GPCR_x, an antibody fragment of GPCR_x, a ligand of GPCR_x, or an antibody-drug conjugate of GPCR_x. For example, in some embodiments, the inhibitor of the CXCR4-GPCR_x heteromer is an antagonist of the CXCR4-GPCR_x heteromer, an inverse agonist of the CXCR4-GPCR_x heteromer, a partial antagonist of the CXCR4-GPCR_x heteromer, an allosteric modulator of the CXCR4-GPCR_x heteromer, an antibody of the CXCR4-GPCR_x heteromer, an antibody fragment of the CXCR4-GPCR_x heteromer, a ligand of the CXCR4-GPCR_x heteromer, a protein-protein interaction (PPI) inhibitor of the CXCR4-GPCR_x heteromer, or an antibody-drug conjugate of the CXCR4-GPCR_x heteromer.

[335] In some embodiments, the method for treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, as disclosed herein, includes a combination of inhibitors selected from the group consisting of: the CXCR4 inhibitor, the GPCR_x inhibitor, and the inhibitor of the CXCR4-GPCR_x heteromer, wherein the combination may be in a single pharmaceutical composition or a plurality of separate pharmaceutical compositions for each respective inhibitor. In some embodiments, the combination of inhibitors comprises the CXCR4 inhibitor and the GPCR_x inhibitor. In some embodiments, the combination of inhibitors comprises the CXCR4 inhibitor and the inhibitor of the CXCR4-GPCR_x heteromer. In some embodiments, the combination of inhibitors comprises the GPCR_x inhibitor and the inhibitor of the CXCR4-GPCR_x heteromer. In some embodiments, the combination of inhibitors are administered sequentially, concurrently, or simultaneously. In some embodiments, the combination of inhibitors are administered as a pharmaceutical composition further comprising a phar-

maceutically acceptable carrier. In some embodiments, the combination of inhibitors are administered as separate pharmaceutical compositions, wherein the separate pharmaceutical compositions independently further comprise a pharmaceutically acceptable carrier. In some embodiments, the method for treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, as disclosed herein, comprises a therapeutically effective amount or a sub-therapeutically effective amount of the CXCR4 inhibitor, such as administering a therapeutically effective amount of the CXCR4 inhibitor to the patient. In some embodiments, the method for treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, as disclosed herein, comprises a therapeutically effective amount or a sub-therapeutically effective amount of the GPCR α inhibitor, such as administering a therapeutically effective amount of the GPCR α inhibitor to the patient. In some embodiments, the method for treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, as disclosed herein, comprises a therapeutically effective amount or a sub-therapeutically effective amount of the inhibitor of the CXCR4-GPCR α heteromer, such as administering a therapeutically effective amount of the inhibitor of the CXCR4-GPCR α heteromer to the patient.

- [336] In some embodiments, the method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, as disclosed herein, may comprise: 1) determining whether the patient has the cancer cell containing the CXCR4-GPCR α heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if: i) the patient's cancer cell containing said CXCR4-GPCR α heteromer; or ii) a CXCR4-GPCR α heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR α heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient derived cell(s) containing the CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR α heteromer; and 2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer to the cancer patient. In some embodiments, the method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, as disclosed herein, may comprise: 1) determining whether the patient's cancer cell contains the CXCR4-GPCR α heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR α heteromer is present in the patient's cancer cell; wherein: a) the GPCR α of the CXCR4-GPCR α heteromer is selected from the group

consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay, such as via a co-internalization assay, bimolecular fluorescence complementation (BiFC), or a proximity ligation assay (PLA); and 2) if the patient's cancer cell contains said CXCR4-GPCR_x heteromer, then internally administering an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer to the patient. In some embodiments, the patient's biological sample is a biological fluid sample or is a biological tissue sample. In some embodiments, a liquid biopsy is performed on the biological fluid sample or performed on the biological tissue sample. In some embodiments, the biological fluid sample is a blood sample, a plasma sample, a saliva sample, a cerebral fluid sample, an eye fluid sample, or a urine sample. In certain embodiments, the biological fluid sample includes circulating tumor cells (CTCs), tumor-derived cell-free DNA (cfDNA), circulating small RNAs, and extracellular vesicles including exosomes, from bodily fluids as disclosed, for example, in Campos CDM et al., "Molecular Profiling of Liquid Biopsy Samples for Precision Medicine," *Cancer J.* 2018 Mar/Apr;24(2):93-103, which is incorporated hereby in its entirety. In some embodiments, the biological tissue sample is an organ tissue sample, a bone tissue sample, or a tumor tissue sample.

[337] The term "heteromer" as used herein refers to macromolecular complex composed of at least two GPCR units [protomers with biochemical properties that are demonstrably different from those of its individual components. Heteromerization can be evaluated by in situ hybridization, immunohistochemistry, RNAseq, Reverse transcription-quantitative PCR (RT-qPCR, realtime PCR), microarray, proximity ligation assay (PLA), time-resolved FRET (TR-FRET), whole-body Single-photon emission computed tomography (SPECT) or Positron Emission Tomography/Computed Tomography (PET/CT).

[338] The phrase "effective amount" as used herein refers to an amount sufficient to effect beneficial or desired results. An effective amount can be administered in one or more administrations, applications or dosages. Such delivery is dependent on a number of variables including the time period for which the individual dosage unit is to be used, the bioavailability of the agent, the route of administration, etc.

[339] The phrase "therapeutically effective amount" as used herein refers to the amount of a therapeutic agent (e.g., an inhibitor, an antagonist, or any other therapeutic agent

provided herein) which is sufficient to reduce, ameliorate, and/or prevent the severity and/or duration of a cancer and/or a symptom related thereto. A therapeutically effective amount of a therapeutic agent can be an amount necessary for the reduction, amelioration, or prevention of the advancement or progression of a cancer, reduction, amelioration, or prevention of the recurrence, development or onset of a cancer, and/or to improve or enhance the prophylactic or therapeutic effect of another therapy (e.g., a therapy other than the administration of a inhibitor, an antagonist, or any other therapeutic agent provided herein).

[340] The phrase "therapeutic agent" refers to any agent that can be used in the treatment, amelioration, prevention, or management of a cancer and/or a symptom related thereto. In certain embodiments, a therapeutic agent refers to an inhibitor of CXCR4-GPCR α heteromer of the invention. A therapeutic agent can be an agent which is well known to be useful for, or has been or is currently being used for the treatment, amelioration, prevention, or management of a cancer and/or a symptom related thereto.

[341] The phrase "intracellular Ca²⁺ assay," "calcium mobilization assay," or their variants as used herein refer to cell-based assay to measure the calcium flux associated with GPCR activation or inhibition. The method utilizes a calcium sensitive fluorescent dye that is taken up into the cytoplasm of most cells. The dye binds the calcium released from intracellular store and its fluorescence increases. The change in the fluorescence intensity is directly correlated to the amount of intracellular calcium that is released into cytoplasm in response to ligand activation of the receptor of interest.

[342] The phrase "proximity-based assay" as used herein refers to biophysical and biochemical techniques that are able to monitor proximity and/or binding of two protein molecules *in vitro* (in cell lysates) and in live cells, including bioluminescence resonance energy transfer (BRET), fluorescence resonance energy transfer (FRET), bimolecular fluorescence complementation (BiFC), Proximity ligation assay (PLA), cysteine crosslinking, and co-immunoprecipitation (Ferre et al., 2009; Gomes et al., 2016).

[343] Alternative methods for detecting heteromer formation include, but are not limited to: immunostaining (Bushlin et al., 2012; Decaillot et al., 2008); immunoelectron microscopy (Fernandez-Duenas et al., 2015); BRET (Pfleger and Eidne, 2006); Time-resolved FRET assays (Fernandez-Duenas et al., 2015); In Situ Hybridization (He et al., 2011); FRET (Lohse et al., 2012); β -arrestin recruitment assay using GPCR heteromer identification technology (GPCR-HIT, Dimerix Bioscience) (Mustafa and Pfleger, 2011) using BRET, FRET, BiFC, Bimolecular Luminescence Complementation, enzyme fragmentation assay, and Tango Tango GPCR assay system (Thermo Fisher Scientific) (Mustafa, 2010); PRESTO-Tango system (Kroeze et al., 2015); regulated secretion/aggregation technology (ARIAD Pharmaceuticals) (Hansen

et al., 2009); Receptor Selection and Amplification Technology (ACADIA Pharmaceuticals) (Hansen et al., 2009); DimerScreen (Cara Therapeutics) (Mustafa, 2010); Dimer/interacting protein translocation assay (Patobios) (Mustafa, 2010); Co-immunoprecipitation (Abd Alla et al., 2009); GPCR internalization assays using surface enzyme-linked immunosorbent assay (ELISA) (Decaillot et al., 2008) or Flow Cytometry (Law et al., 2005); Whole Cell Phosphorylation Assays (Pfeiffer et al., 2002); and Proximity-ligation assay (PLA) (Frederick et al., 2015).

[344] Alternative methods for detecting changes in pharmacological properties, signaling properties, and/or trafficking properties, in cells expressing both CXCR4 and GPCR α include, but are not limited to: Radioligand Binding Assays (Bushlin et al., 2012; Pfeiffer et al., 2002); Cell Surface Biotinylation and Immunoblotting (He et al., 2011); immunostaining (Bushlin et al., 2012; Decaillot et al., 2008); immunoelectron microscopy (Fernandez-Duenas et al., 2015); [35 S]GTP S Binding assays (Bushlin et al., 2012); Calcium imaging or assays using dyes such as Fura 2-acetomethoxy ester (Molecular Probes), Fluo-4 NW calcium dye (Thermo Fisher Scientific), or FLIPR5 dye (Molecular Devices); cAMP assays using radioimmunoassay kit (Amersham Biosciences); AlphaScreen (PerkinElmer Life Sciences); Parameter Cyclic AMP Assay (R&D Systems); femto cAMP kit (Cisbio); cAMP Direct Immunoassay Kit (Calbiochem) or GloSensor cAMP assay (Promega); GTPase assay (Pello et al., 2008); PKA activation (Stefan et al., 2007); ERK1/2 and/or Akt/PKB Phosphorylation Assays (Callen et al., 2012); Src and STAT3 phosphorylation assays (Rios et al., 2006); reporter assays such as cAMP response element (CRE); nuclear factor of activated T-cells response element (NFAT-RE); serum response element (SRE); serum response factor response element (SRF-RE); and NF- κ B-response element luciferase reporter assays; Secreted alkaline phosphatase Assay (Decaillot et al., 2011); Measurement of Inositol 1-Phosphate Production Using TR-FRET or [3 H]myo-Inositol (Mustafa et al., 2012); RT-qPCR for measuring downstream target gene expression (Mustafa et al., 2012); and Adenylyl Cyclase Activity (George et al., 2000); next generation sequencing (NGS); and any other assay that can detect a change in receptor function as a result of receptor heterodimerization.

[345] The phrase "protein-protein interaction inhibitor," "PPI inhibitor," or their variants as used herein refer to any molecules that can interfere with protein-protein interactions. Protein-protein interaction, unlike enzyme-substrate interaction involving well-defined binding pockets, is a transient interaction or association between proteins over relatively large areas and is often driven by electrostatic interactions, hydrophobic interactions, hydrogen bonds, and/or Van der Waals forces. PPI inhibitors may include, but not limited to, membrane-permeable peptides or lipid fused to a peptide sequence that disrupts the GPCR heteromeric interface, for example, transmembrane helix, intra-

cellular loop, or C-terminal tail of GPCR_x. The PPI inhibitor of the CXCR4-GPCR_x heteromer, for example, may be a membrane-permeable peptide or cell-penetrating peptide (CPP) conjugated with peptide that targets the CXCR4-GPCR_x heteromeric interface(s), or may be a cell-penetrating lipidated peptide targeting the CXCR4-GPCR_x heteromeric interface(s).

- [346] For example, the membrane-permeable peptide or cell-penetrating peptide includes: HIV-1 TAT peptides, such as TAT₄₈₋₆₀ and TAT₄₉₋₅₇; Penetratins, such as pAntp(43-58); Polyarginines (Rn such as R5 to R12); Diatos peptide vector 1047 (DPV1047, Vectocell®); MPG (HIV gp41 fused to the nuclear localization signal (NLS) of the SV40 large T antigen); Pep-1 (tryptophan-rich cluster fused to the NLS of SV40 large T antigen); pVEC peptide (vascular endothelial cadherin); p14 alternative reading frame (ARF) protein-based ARF(1-22); N-terminus of the unprocessed bovine prion protein BPrPr(1-28); Model amphipathic peptide (MAP); Transportans; Azurin-derived p28 peptide; amphipathic β -sheet peptides, such as VT5; proline-rich CPPs, such as Bac 7 (Bac1-24); hydrophobic CPPs, such as C105Y derived from α 1-Antitrypsin; PFVYLI derived from synthetic C105Y; Pep-7 peptide (CHL8 peptide phage clone); and modified hydrophobic CPPs, such as stapled peptides and prenylated peptides (Guidotti et al., 2017; Kristensen et al., 2016). The membrane-permeable peptide or cell-penetrating peptide can further include, for example, TAT-derived cell-penetrating peptides, signal sequence-based (e.g., NLS) cell-penetrating peptides, hydrophobic membrane translocating sequence (MTS) peptides, and arginine-rich molecular transporters. The cell-penetrating lipidated peptide includes, for example, pepducins, such as ICL1/2/3, C-tail- short palmitoylated peptides (Covic et al., 2002; O'Callaghan et al., 2012).
- [347] The peptide(s) that target the CXCR4-GPCR_x heteromeric interface may be, for example, a transmembrane domain of CXCR4, transmembrane domain of GPCR_x, intracellular loop of CXCR4, intracellular loop of GPCR_x, C-terminal domain of CXCR4, or C-terminal domain of GPCR_x, extracellular loop of CXCR4, extracellular loop of GPCR_x, N-terminal region of CXCR4, or N-terminal region of GPCR_x.
- [348] In some embodiments, the invention provides a pharmaceutical composition (sometimes referred to herein as "pharmaceutical formulations") having an inhibitor of a CXCR4-GPCR_x heteromer of the invention and a pharmaceutically acceptable carrier. A pharmaceutically acceptable carrier that can be used in the pharmaceutical compositions of the invention include any of the standard pharmaceutical carriers known in the art, such as physiologically acceptable carriers, excipients or stabilizers, for example, phosphate buffered saline solution, water and emulsions such as an oil and water emulsion, and various types of wetting agents. These pharmaceutical compositions can be prepared in liquid unit dose forms or any other dosing form that is

sufficient for delivery of the inhibitor of a CXCR4-GPCR α heteromer of the invention to the target area of the subject in need of treatment. For example, the pharmaceutical compositions can be prepared in any manner appropriate for the chosen mode of administration, e.g., intravascular, intramuscular, subcutaneous, intradermal, intrathecal, etc. Other optional components, e.g., pharmaceutical grade stabilizers, buffers, preservatives, excipients and the like can be readily selected by one of skill in the art. The preparation of a pharmaceutically composition, having due regard to pH, isotonicity, stability and the like, is within the level of skill in the art.

[349] Pharmaceutical formulations containing one or more inhibitors of a CXCR4-GPCR α heteromer of the invention provided herein can be prepared for storage by mixing the inhibitors having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (Remington's Pharmaceutical Sciences (1990) Mack Publishing Co., Easton, PA), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol; low molecular weight polypeptides of less than about 10 residues; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counterions such as sodium; metal complexes, e.g., Zn-protein complexes; and/or non-ionic surfactants such as TWEENTM, PLURONICSTM, or polyethylene glycol (PEG).

[350] Thus, in some embodiments, the invention provides a method for treatment, amelioration, or prevention of a disease in a subject in need thereof. The methods of the invention can include administering a therapeutically effective amount of a pharmaceutical composition provided herein to the subject. For example, the pharmaceutical composition can include one or more inhibitors of a CXCR4-GPCR α heteromer provided herein. Diseases that can be treated or prevented using the methods of the invention include cancer, tumor, metastasis, and/or angiogenesis. In particular, the methods of the invention are useful for treating cancers or related symptoms wherein the cells of cancer, tumor, and/or microenvironment expresses the CXCR4-GPCR α heteromer. Non-limiting examples of cancers or tumors that can be treated, ameliorated, or prevented using the methods of the invention include tumors of the gas-

trointestinal tract, for example, breast cancer, lung cancer, small cell carcinoma of the lung, hepatocellular carcinoma, brain cancer, kidney cancer, pancreatic cancer or pancreatic adenocarcinoma, ovarian cancer, prostate cancer, melanoma, lymphoma, leukemia, multiple myeloma, renal cell carcinoma, soft tissue sarcoma, gastrointestinal cancer, stomach cancer, colon cancer, colorectal cancer, colorectal adenocarcinoma, bladder adenocarcinoma, esophageal cancer, and adenocarcinoma of the stomach, esophagus, throat, and urogenital tract.

- [351] In some embodiments, the invention provides a method for treating cancer in a patient having a cell containing a CXCR4-GPCR α heteromer, or provides a method of suppressing enhanced downstream signaling from a CXCR4-GPCR α heteromer in a cell of a patient suffering from cancer, the method comprising: administering to the patient an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; wherein: i) the CXCR4-GPCR α heteromer has enhanced downstream signaling; and ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the cancer patient.
- [352] In some embodiments, the invention provides a pharmaceutical kit for use in treating cancer in a patient having a cell containing a CXCR4-GPCR α heteromer, the pharmaceutical kit comprising: an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; wherein the CXCR4-GPCR α heteromer has enhanced downstream signaling. In some embodiments, the invention provides a pharmaceutical composition for use in treating cancer in a patient having a cell containing a CXCR4-GPCR α heteromer, the pharmaceutical composition comprising: i) an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and ii) a pharmaceutically acceptable carrier; wherein the CXCR4-GPCR α heteromer has enhanced downstream signaling.
- [353] In some embodiments, according to the method of treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, provided herein, the progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR α heteromer is decreased in the range of 5-100% more upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient, such as decreased in the range of 5-100% more, 10-100% more, 20-100% more, 30-100% more, 40-100% more, 50-100% more, 60-100% more, 75-100% more, 5-75% more, 5-50% more, or 5-25% more, upon administration of the combination of inhibitors, relative to administering

the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient. In some embodiments, the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; for example, selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, and HRH1; selected from the group consisting of: ADRB2, CHRM1, and HRH1; or selected from the group consisting of: ADRB2, HRH1, and TACR3. In some embodiments, the GPCR α inhibitor is selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CALCR inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; for example, selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor;

selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, and HRH1 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, and HRH1 inhibitor; or selected from the group consisting of: ADRB2 inhibitor, HRH1, inhibitor and TACR3 inhibitor.

[354] In some embodiments, according to the method of treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, provided herein, the efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor, such as increased in the range of 5-1750%, 5-1500%, 5-1250%, 5-1000%, 5-900%, 5-800%, 5-700%, 5-500%, 5-400%, 5-250%, 5-200%, 5-100%, 5-75%, 5-50%, 5-40%, 5-30%, 5-25%, 100-2000%, 200-2000%, 300-2000%, 500-2000%, 750-2000%, 1000-2000%, 1250-2000%, 1500-2000%, 5-1500%, 25-1500%, 50-1500%, 75-1500%, 100-1500%, 200-1500%, 300-1500%, 500-1500%, 750-1500%, 1000-1500%, or 1250-1500%, when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor. In some embodiments, the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; for example, selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, and HRH1; selected from the group consisting of: ADRB2,

CHRM1, and HRH1; or selected from the group consisting of: ADRB2, HRH1, and TACR3. In some embodiments, the GPCR_x inhibitor is selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CALCR inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; for example, selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, and HRH1 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, and HRH1 inhibitor; or selected from the group consisting of: ADRB2 inhibitor, HRH1, inhibitor and TACR3 inhibitor.

- [355] In some embodiments, according to the method of treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, provided herein, the efficacy of a GPCR_x inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR_x heteromer, relative to efficacy of the GPCR_x inhibitor when administered as a single inhibitor, such as increased in the range of 5-1750%, 5-1500%, 5-1250%, 5-1000%, 5-900%, 5-800%, 5-700%, 5-500%, 5-400%, 5-250%, 5-200%, 5-100%, 5-75%, 5-50%, 5-40%, 5-30%, 5-25%, 100-2000%, 200-2000%, 300-2000%, 500-2000%, 750-2000%, 1000-2000%, 1250-2000%, 1500-2000%, 5-1500%,

25-1500%, 50-1500%, 75-1500%, 100-1500%, 200-1500%, 300-1500%, 500-1500%, 750-1500%, 1000-1500%, or 1250-1500%, when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor. In some embodiments, the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; for example, selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, and HRH1; selected from the group consisting of: ADRB2, CHRM1, and HRH1; or selected from the group consisting of: ADRB2, HRH1, and TACR3. In some embodiments, the GPCR α inhibitor is selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CALCR inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; for example, selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADORA2B inhibitor,

ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, and HRH1 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, and HRH1 inhibitor; or selected from the group consisting of: ADRB2 inhibitor, HRH1, inhibitor and TACR3 inhibitor.

[356] In some embodiments, according to the method of treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, provided herein, the method administers a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; or the pharmaceutical kit or pharmaceutical composition comprises a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer. In some embodiments, the combination of inhibitors is a combination of two inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer. In some embodiments, according to the method of treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, provided herein, the method administers a combination of a CXCR4 inhibitor and a GPCR_x inhibitor; or the pharmaceutical kit or pharmaceutical composition comprises a combination of a CXCR4 inhibitor and a GPCR_x inhibitor. In some embodiments, according to the method of treating, method of suppressing, pharmaceutical kit, or pharmaceutical composition, provided herein, the method administers a CXCR4-GPCR_x heteromer inhibitor; or the pharmaceutical kit or pharmaceutical composition comprises a CXCR4-GPCR_x heteromer inhibitor.

[357] In some embodiments, according to the method of treating or method of suppressing provided herein, or according to the use of the pharmaceutical kit or pharmaceutical composition provided herein, the administering of the combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient in the range of between 5-2000 fold, relative to single inhibitor administration, such as suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient in the range of between 5-1750 fold, 5-1500 fold, 5-1250 fold, 5-1000 fold, 5-900 fold, 5-800 fold, 5-700 fold, 5-500 fold,

[illegible]

TACR3 inhibitor; selected from the group consisting of: ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, and HRH1 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, and HRH1 inhibitor; or selected from the group consisting of: ADRB2 inhibitor, HRH1, inhibitor and TACR3 inhibitor.

- [358] In some embodiments, according to the method of treating or method of suppressing provided herein, or according to the use of the pharmaceutical kit or pharmaceutical composition provided herein, the administering of the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient in the range of between 5-2000 fold, relative to suppression of downstream signaling from either a CXCR4 protomer or a GPCR_x protomer in their respective individual protomer context, such as suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient in the range of between 5-1750 fold, 5-1500 fold, 5-1250 fold, 5-1000 fold, 5-900 fold, 5-800 fold, 5-700 fold, 5-500 fold, 5-400 fold, 5-250 fold, 5-200 fold, 5-100 fold, 5-75 fold, 5-50 fold, 5-40 fold, 5-30 fold, 5-25 fold, 100-2000 fold, 200-2000 fold, 300-2000 fold, 500-2000 fold, 750-2000 fold, 1000-2000 fold, 1250-2000 fold, 1500-2000 fold, 5-1500 fold, 25-1500 fold, 50-1500 fold, 75-1500 fold, 100-1500 fold, 200-1500 fold, 300-1500 fold, 500-1500 fold, 750-1500 fold, 1000-1500 fold, or 1250-1500 fold, relative to suppression of downstream signaling from either a CXCR4 protomer or a GPCR_x protomer in their respective individual protomer context. In some embodiments, the GPCR_x of the CXCR4-GPCR_x heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; for example, selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting

of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, and HRH1; selected from the group consisting of: ADRB2, CHRM1, and HRH1; or selected from the group consisting of: ADRB2, HRH1, and TACR3. In some embodiments, the GPCR_x inhibitor is selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CALCR inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; for example, selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, and HRH1 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, and HRH1 inhibitor; or selected from the group consisting of: ADRB2 inhibitor, HRH1, inhibitor and TACR3 inhibitor.

[359] Initial testing and evaluation of an inhibitor, or combination of inhibitors, regarding

whether effective, or therapeutically effective, in suppressing an enhanced downstream signaling from a CXCR4-GPCR α heteromer, according to the methods disclosed herein, and/or in determining an IC₅₀ value according to the assays disclosed herein, may utilize a concentration of the inhibitor in the range of between 1-10 μ M (or each of the inhibitors of the combination at concentrations in the range of between 1-10 μ M), such as at a concentration of 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 μ M. If suppression of the signal by the inhibitor is not appreciable enough for a determinable measurement, then a greater concentration of the inhibitor may be used to better evaluate a determinable measurement, such as an IC₅₀ value. If suppression of the signal by the inhibitor is very strong, then a lower concentration of the inhibitor may be used to better evaluate a determinable measurement, such as an IC₅₀ value.

[360] It is understood that modifications which do not substantially affect the activity of the various embodiments of this invention are also provided within the definition of the invention provided herein. Accordingly, the following examples are intended to illustrate but not limit the invention disclosed herein.

[361]

[362] **EXAMPLES**

[363] **Example 1. Evaluation of CXCR4-GPCR α heteromer formation by BiFC assay.**

[364] To identify novel CXCR4-GPCR α heteromers, we made recombinant adenoviruses encoding 143 GPCRs fused with N-terminal fragments of yellow fluorescent protein Venus (VN) and 147 GPCRs fused with C-terminal fragment of Venus (VC) as described in Song et al. (Song et al., 2014; SNU patent; Song, thesis). CXCR4-GPCR heteromers were identified using bimolecular fluorescence complementation (BiFC) assay (FIG. 1), in which two complementary VN and VC fragments of Venus re-constitute a fluorescent signal only when both fragments are close enough through interaction between two different proteins to which they are fused (Hu et al., 2002).

[365] U-2 OS cells were plated in 96-well plates, were co-transduced with 30 MOI each of adenoviruses encoding CXCR4-VN and GPCR α -VC or CXCR4-VC and GPCR α -VN, and were allowed to express GPCRs for 2 days. After staining the cells with Hoechst 33342, BiFC and nuclear images were obtained from three fields per well using IN Cell Analyzer 1000. Images of about 200 cells from each well were analyzed with Multi-target analysis software in IN Cell Developer ToolBox (GE Healthcare, Waukesha, WI). Cell boundary was marked based on Hoechst signal, and fluorescence intensity per cell was measured. Cells with fluorescence intensities higher than background level were considered as BiFC positive cells. Dead cells that showed extremely high intensities were excluded from the cell count. Positive cells were determined, and positive cell count ratio ("BiFC score") was calculated as (positive cells/total cells) x 100 (see Table 4 below).

- [366] When CXCR4-VN was co-expressed with HA-VC (FIG. 2a) or GCGR-VC, a GPCR encoding glucagon receptor (FIG. 2c), no yellow fluorescence protein (YFP) signal (BiFC signal) was observed. In contrast, when CXCR4-VN was co-expressed with CXCR4-VC (FIG. 2b), BiFC signal was observed in the plasma membrane and in the cytoplasm. Strong BiFC signal was observed in the plasma membrane and in the cytoplasm when CXCR4-VN was co-transduced with ADCYAP1R1-VC (FIG. 2d), ADORA3-VC (FIG. 2f), ADRB2-VC (FIG. 2g), APLNR-VC (FIG. 2h), C5AR1-VC (FIG. 2i), CALCR-VC (FIG. 2j), CCR5-VC (FIG. 2k), CHRM1-VC (FIG. 2l), GALR1-VC (FIG. 2m), EDNRB-VC (FIG. 2n), HRH1-VC (FIG. 2o), MLNR-VC (FIG. 2p), NTSR1-VC (FIG. 2q), PTGER2-VC (FIG. 2r), SSTR2-VC (FIG. 2t), and TACR3-VC (FIG. 2u). Robust BiFC signal was also observed when cells were co-transduced with CXCR4-VC and ADORA2B-VN (FIG. 2e) or PTGER3-VN (FIG. 2s). Cells that showed BiFC fluorescence signal higher than background level was counted as BiFC positive cells, and BiFC score was calculated.
- [367] Protein-protein interaction can be affected by fusion tags such as fluorescence protein fragments in BiFC or renilla luciferase in BRET through interfering with the expression, folding, or localization of the partner protein. Partner protein can also impair the expression or folding of the fusion tags, and affect proximity-based assay result. Thus, the absence of signal between two proteins in specific combination does not necessarily imply that the proteins do not interact, but simply that the attached donor and acceptor molecules are in particular conformation that does not allow interaction to occur (Eidne et al., 2002; Kerppola, 2006).
- [368] Therefore, CXCR4 and GPCR_x that gave BiFC signal in either CXCR4-VN and GPCR_x-VC or CXCR4-VC and GPCR_x-VN combination were considered as interacting proteins. The BiFC score of CXCR4-VN and CXCR4-VC pair, a well-known homomer, was 9.9. Thus, CXCR4-GPCR_x pairs that gave BiFC score equal or higher than 10 were selected as candidates for CXCR-GPCR_x heteromer as shown in Table 4.
- [369]
- [370] Table 4. GPCRs that exhibited BiFC score equal or higher than 10 when co-expressed with CXCR4.

[371] [Table 4]

GPCRx		BiFC score	GPCRx		BiFC score
		CXCR4-VN & GPCRx-VC			CXCR4-VN & GPCRx-VC
1	ADCYAP1R1	23	10	GALR1	60
2	ADORA2B	11*	11	EDNRB	41
3	ADORA3	43	12	HRH1	45
4	ADRB2	24	13	MLNR	10
5	APLNR	63	14	NTSR1	48
6	C5AR1	42	15	PTGER2	58
7	CALCR	53	16	PTGER3	13*
8	CCR5	44	17	SSTR2	30
9	CHRM1	24	18	TACR3	33

*CXCR4-VC and GPCR_x-VN were used.

[372]

[373] Sixteen GPCRs, ADCYAP1R1, ADORA3, ADRB2, APLNR, C5AR1, CALCR, CCR5, CHRM1, GALR1, EDNRB, HRH1, MLNR, NTSR1, PTGER2, SSTR2, and TACR3, were identified as CXCR4-interacting GPCRs in CXCR4-VN and GPCR_x-VC combination and two GPCRs, ADORA2B and PTGER3, gave BiFC score higher than 10 in CXCR4-VC and GPCR_x-VN combination as shown in Table 4. Among those, ADRB2 and CCR5 have been reported to form heteromers with CXCR4 (LaRocca 2010, Nakai 2014, (Agrawal et al., 2004; LaRocca et al., 2010; Rodriguez-Frade et al., 2004; Sohy et al., 2007; Sohy et al., 2009; Martinez-Munoz et al., 2014)), and the remaining 16 GPCRs were found to be novel CXCR4-GPCR_x heteromers that have not been reported to the best of our knowledge.

[374] Among GPCRs that have been known to interact with CXCR4, ADRA1A and CXCR3 were also included in our BiFC assay. These GPCRs were excluded from further investigation since they gave BiFC signal of less than 10 with CXCR4: ADRA1A (CXCR4-VN-ADRA1A-VC: BiFC score 7.85) and CXCR3 (BiFC score 1.16).

[375] Cannabinoid Receptor 2 (CB2, sometimes referred to as CNR2) was identified as a CXCR4-interacting GPCR in our BiFC assay (CNR2-VN-CXCR4-VC configuration: BiFC score 4.25; CNR2-VC-CXCR4-VN configuration: BiFC score 38.1), and co-internalization assay as described in FIGS. 3a-3b and 4a-4q. But CNR2 was not selected as the final candidate because it failed to exhibit an enhanced downstream signaling in the calcium mobilization assay (see Example 3 below).

[376]

[377] **Example 2. Evaluation of CXCR4-GPCR_x heteromer formation by co-internalization assay.**

- [378] Some of the GPCR heteromers are known to exhibit, as a result of the heteromerization, altered trafficking properties such as maturation of the partner GPCR (GABA(B) receptor) (White, 1998), agonist-mediated internalization of the partner GPCR from the cell surface (DOR-GRPR, A2A-D2R) (Hillion et al., 2002; Liu et al., 2011; Torvinen et al., 2005), and changes in the localization of the partner GPCR from an intracellular compartment to the cell surface (DOR-CB1) (Rozenfeld et al., 2012).
- [379] Co-internalization of pairs of co-expressed GPCRs in response to agonists selective for only one of the pair has been used to confirm GPCR heteromerization (Milligan, 2008). To examine if GPCR_x modulates the trafficking of CXCR4 when co-expressed and forms CXCR4-GPCR_x heteromer and also to confirm the physical interaction between CXCR4 and GPCR_x identified using BiFC assay, cells were co-transduced with adenoviruses encoding CXCR4-GFP and GPCR_x and GFP images were obtained before and 30 min after stimulation with GPCR_x agonist. Loss of GFP expression on the cell surface or appearance of GFP granules inside the cells were considered as CXCR4-GFP co-internalization with GPCR_x (FIGS. 3a-3b).
- [380] In FIGS. 4a-4q, cells were stimulated with GPCR_x agonists specific for CXCR4 (FIG. 4a), ADCYAP1R1 (FIG. 4b), ADORA2B (FIG. 4c), ADORA3 (FIG. 4d), ADRB2 (FIG. 4e), APLNR (FIG. 4f), C5AR1 (FIG. 4g), CCR5 (FIG. 4h), CHRM1 (FIG. 4i), GALR1 (FIG. 4j), EDNRB (FIG. 4k), HRH1 (FIG. 4l), MLNR (FIG. 4m), NTSR1 (FIG. 4n), PTGER3 (FIG. 4o), SSTR2 (FIG. 4p), and TACR3 (FIG. 4q) as follows: 10 nM of CXCL12 (FIG. 4a), 1 μ M of vasoactive intestinal peptide (VIP) (FIG. 4b), 1 μ M of BAY 60-6583 (FIG. 4c), 1 μ M of CGS21680 (FIG. 4d), 100 nM of formoterol (FIG. 4e), 1 μ M of apelin-13 (FIG. 4f), 100 nM of C5a (FIG. 4g), 100 nM of CCL2 (FIG. 4h), 1 μ M of acetylcholine (FIG. 4i), 100 nM of galanin (FIG. 4j), 1 μ M of endothelin 1 (FIG. 4k), 1 μ M of histamine (FIG. 4l), 100 nM of motilin (FIG. 4m), 1 μ M of neurotensin (FIG. 4n), 1 μ M of PGE2 (FIG. 4o), 1 μ M of SRIF-14 (FIG. 4p), and 1 μ M of senktide (FIG. 4q), respectively. Images were obtained before and 30 min after agonist stimulation, and analyzed using IN Cell Analyzer 2000. The selection of the concentration of these GPCR_x agonists was initially set at the EC₅₀ concentration (or alternatively, the K_i or K_d concentration) for the respective specific GPCR_x, and if the resulting signal was too intense, then the concentration was lowered below the EC₅₀, and if the resulting signal was too weak, then the concentration was increased to no more than 10,000x the EC₅₀ concentration.
- [381] Stimulation of cells expressing CXCR4-GFP with CXCL12 resulted in re-location of GFP from the plasma membrane to distinct intracellular granules, showing internalization of surface CXCR4-GFP into the cytoplasm (FIG. 4a). Among 18 CXCR4-GPCR_x heteromers identified by BiFC assay, 16 heteromers containing the following GPCR_x induced internalization of CXCR4 as revealed by increased intra-

cellular GFP granules and reduced GFP signal in the plasma membrane in cells co-expressing CXCR4-GFP and GPCR α , confirming heteromer co-internalization: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, APLNR, C5AR1, CCR5, CHRM1, GALR1, EDNRB, HRH1, MLNR, NTSR1, PTGER3, SSTR2, and TACR3 (FIG. 4b-q). CXCR4-GPCR α heteromers containing CALCR and PTGER2, on the other hand, did not show co-internalization of the heteromer upon GPCR α agonist treatment. These results demonstrate that certain GPCR α partners identified in the BiFC assay do form CXCR4-GPCR α heteromers in cells co-expressing both GPCRs, as evidenced by altered cellular trafficking properties of the heteromer.

[382]

[383] **Example 3. Evaluation by Ca²⁺ mobilization assay of enhanced CXCR4 downstream signaling upon CXCR4-GPCR α heteromer formation and inhibition of the enhanced signaling.**

[384] To further examine if these CXCR4-GPCR α heteromers exhibit properties distinct from those of the individual protomers (in cells lacking one of the receptors), we investigated calcium signaling in the presence of either or both agonists in cells expressing either GPCR or both GPCRs together. In this example, as noted in Example 2, the selection of the concentration of these GPCR α agonists was initially set at the EC₅₀ concentration (or alternatively, the K_i or K_d concentration) for the respective specific GPCR α , and if the resulting Ca²⁺ signal was too intense, then the concentration was lowered below the EC₅₀, and if the resulting Ca²⁺ signal was too weak, then the concentration was increased to no more than 100x the EC₅₀ concentration.

[385] When MDA-MB-231 human breast cancer cells were transduced with adenoviruses encoding CXCR4, treatment of CXCL12 evoked intracellular calcium mobilization (FIG. 5a). Stimulation of the cells with salmeterol, an ADRB2-selective agonist, did not induce calcium response, demonstrating the calcium response evoked by CXCL12 was mediated by CXCR4. Co-treatment of the cells with CXCL12 and salmeterol induced similar calcium response compared to the one induced by CXCL12 alone. In cells overexpressing ADRB2 alone, CXCL12 did not induce calcium response while salmeterol induced calcium response, showing that salmeterol induces calcium response through ADRB2 (FIG. 5b). Co-treatment of both agonists induced similar calcium response to the one stimulated by salmeterol alone.

[386] In cells overexpressing both CXCR4 and ADRB2, stimulation with each agonist induced calcium responses similar to the ones shown in cells expressing CXCR4 or ADRB2 alone (FIGS. 5c vs 5a and 5b). In contrast, co-treatment of both agonists together significantly increased the calcium response compared to the ones evoked by individual agonists (FIGS. 5c & 5d). The enhanced calcium signaling was observed only in cells expressing both CXCR4 and ADRB2, but not in cells expressing either

CXCR4 or ADRB2 alone. These results clearly demonstrate that CXCR4-ADRB2 heteromer exhibit properties distinct from those of the individual GPCRs.

[387]

[388]

To examine if the other GPCRs identified to interact with CXCR4 in BiFC and co-internalization assay also show distinct properties in calcium signaling, experiments shown in FIGS. 5a-5d were performed for other identified GPCR_x partners. In cells co-expressing CXCR4 and either ADCYAP1R1 (FIG. 6a), ADORA2B (FIG. 6b), ADORA3 (FIG. 6c), C5AR1 (FIG. 6d), CALCR (FIG. 6e), CHRM1 (FIG. 6f), EDNRB (FIG. 6g), HRH1 (FIG. 6h), MLNR (FIG. 6i), NTSR1 (FIG. 6j), PTGER2 (FIG. 6k), or TACR3 (FIG. 6l), co-treatment of CXCL12 and respective GPCR_x agonist significantly increased calcium responses compared with the sum of calcium responses that were induced by individual agonists as shown in FIGS. 5c and 5d, and as shown in FIGS. 6a-6l. The cells co-expressing the CXCR4 and the GPCR_x (either ADCYAP1R1 (FIG. 6a), ADORA2B (FIG. 6b), ADORA3 (FIG. 6c), C5AR1 (FIG. 6d), CALCR (FIG. 6e), CHRM1 (FIG. 6f), EDNRB (FIG. 6g), HRH1 (FIG. 6h), MLNR (FIG. 6i), NTSR1 (FIG. 6j), PTGER2 (FIG. 6k), or TACR3 (FIG. 6l)) exhibited an enhanced calcium mobilization upon co-stimulation (or co-treatment) with the CXCL12 and the respective selective GPCR_x agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR_x agonist, as determined by the calcium mobilization assay. Notably, in cells co-expressing CXCR4 and CNR2, CXCL12-induced calcium response was significantly reduced, rather than enhanced, upon addition of CXCL12 alone or together with JWH-133, a selective CNR2 agonist, compared with the CXCL12-mediated calcium response observed in cells expressing CXCR4 alone (data not shown).

[389]

Within the above set of co-expressed systems, it was also noted that an enhanced amount of calcium mobilization was observed in the individual protomer context in cells involving CXCR4 and PTGER2 (FIG. 6k) (an individual protomer context, *e.g.*, either CXCR4 expressed in the absence of the respective GPCR_x, or the respective GPCR_x expressed in the absence of the CXCR4), upon co-stimulation with CXCL12 and the respective selective GPCR_x agonist. Specifically, an enhanced amount of calcium mobilization was observed for PTGER2 (FIG. 6k), in the individual protomer context, in both situations where CXCR4 was expressed in the absence of the respective GPCR_x, and where the respective GPCR_x (PTGER2) was expressed in the absence of the CXCR4, upon co-stimulation with CXCL12 and the respective selective GPCR_x agonist (5 μ M of PGE₂), the calcium mobilization amount resulting from the individual protomer context systems were greater than the sum of calcium mobilization amounts resulting from single agonist stimulations with either the CXCL12 or the re-

spective selective GPCR α agonist (5 μ M of PGE₂).

- [390] Accordingly, from these observations, those systems co-expressing CXCR4 and the GPCR α selected from the group consisting of ADCYAP1R1 (FIG. 6a), ADORA2B (FIG. 6b), ADORA3 (FIG. 6c), C5AR1 (FIG. 6d), CALCR (FIG. 6e), CHRM1 (FIG. 6f), EDNRB (FIG. 6g), HRH1 (FIG. 6h), MLNR (FIG. 6i), NTSR1 (FIG. 6j), and TACR3 (FIG. 6l), are considered to have: (i) exhibited an enhanced calcium mobilization upon co-stimulation (or co-treatment) with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist, as determined by the calcium mobilization assay; (ii) exhibited a calcium mobilization amount equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist in an individual protomer context; and thus (iii) constituted CXCR4-GPCR α heteromers exhibiting properties distinct from those of the individual GPCR. The enhanced calcium responses were not observed in cells expressing individual GPCRs, thereby demonstrating that these CXCR4-GPCR α heteromers exhibit properties distinct from those of the individual GPCR.
- [391] The enhanced calcium signaling induced by small amounts of individual agonists together clearly indicated that CXCR4-GPCR α heteromers would increase the sensitivity and dynamic range of these receptors at limiting ligand concentrations *in vivo*, and may be associated with poor prognosis. The enhanced calcium signaling by the CXCR4-GPCR α heteromers also indicates that the current diagnosis based on the level of CXCR4 expression alone needs to be changed to consider the expression levels of CXCR4 and GPCR α .
- [392] In contrast, GPCRs such as APLNR (FIG. 7a), CCR5 (FIG. 7b), GALR1 (FIG. 7c), PTGER3 (FIG. 7d), and SSTR2 (FIG. 7e) did not show enhanced calcium responses when cells expressing both receptors were co-treated with both agonists, although these GPCRs were shown to interact with CXCR4 in BiFC (Table 5) and CXCR4-GFP co-internalization assay. These results clearly demonstrate that the synergistic increase in calcium response shown in FIGS. 5a-5d and 6a-6l is a unique feature of CXCR4-GPCR α heteromers, that is not shared by other CXCR4-GPCR α heteromers such as CXCR4-APLNR, CXCR4-CCR5, CXCR4-GALR1, CXCR4-PTGER3, and CXCR4-SSTR2.
- [393] Table 5. GPCRs that exhibited BiFC score equal or higher than 10 when co-expressed with CXCR4 but did not show enhanced Ca²⁺ signaling upon co-treatment of both agonists.

[394] [Table 5]

CXCR4-GPCR _x heteromers that do not show enhanced Ca ²⁺ signaling upon co-treatment of both agonists		
GPCR _x		BiFC score
		CXCR4-VN & GPCR _x -VC
1	APLNR	63
2	CCR5	44
3	GALR1	60
4	PTGER3*	13
5	SSTR2	30

*CXCR4-VC and GPCR_x-VN were used.

- [395] It was further examined if enhanced calcium responses in cells co-expressing CXCR4 and GPCR_x upon co-stimulation with CXCL12 and GPCR_x ligands are inhibited by GPCR_x antagonists. In cells co-expressing CXCR4 and either ADRB2 (FIG. 8a), CHRM1 (FIG. 8b & FIG. 8c), HRH1 (FIGS. 8f-8i), MLNR (FIG. 8j), or NTSR1 (FIG. 8k), treatment of 10 μ M of ADRB2 antagonist carvedilol (FIG. 8a), 1 μ M of CHRM1-selective antagonist VU0255035 (FIG. 8b), 10 μ M of CHRM1 antagonist oxybutynin (FIG. 8c), 1 μ M of HRH1-selective antagonist cetirizine (FIG. 8f), 1 μ M of HRH1-selective antagonist pyrilamine (FIG. 8g), 10 μ M of HRH1 antagonist hydroxyzine (FIG. 8h), 10 μ M of HRH1-selective antagonist loratadine (FIG. 8i), 1 μ M of MLNR-selective antagonist MA-2029 (FIG. 8j) and 1 μ M of NTSR1-selective antagonist meclizine (FIG. 8k), respectively, suppressed the enhanced calcium signaling significantly. And co-treatment of both antagonists together resulted in a more complete suppression (FIGS. 8a-8c, and 8f-8k). These results demonstrated that GPCR_x antagonists could be used as an efficient therapeutics against CXCR4-GPCR_x heteromers where GPCR_x represents ADRB2, CHRM1, HRH1, MLNR, and NTSR1.
- [396] In cells co-expressing CXCR4 and CHRM1, 10 μ M of muscarinic acetylcholine receptor antagonist umeclidinium reduced the enhanced calcium signaling, although not statistically significant (FIG. 8d). In these cells, co-treatment of AMD3100 and umeclidinium almost completely inhibited calcium response induced by simultaneous addition of CXCL12 and bethanechol.
- [397] In cells co-expressing CXCR4 and either EDNRB (FIG. 8e) or TACR3 (FIG. 8l), treatment of either 1 μ M of AMD3100 alone (FIG. 8e and FIG. 8l), or 1 μ M of endothelin receptor antagonist bosentan (FIG. 8e) or 1 μ M of TACR3-selective antagonist SSR 146977 alone (FIG. 8l) failed to inhibit enhanced calcium response significantly. But when the cells were co-treated with AMD3100 and bosentan (FIG. 8e) or AMD3100 and SSR 146977 together (FIG. 8l), the enhanced calcium responses were significantly inhibited.

[398] In cells co-expressing CXCR4 and either CHRM1 (FIG. 8b), EDNRB (FIG. 8e), HRH1 (FIG. 8f and 8g), or MLNR (FIG. 8j), although 1 μ M of AMD3100 alone failed to inhibit enhanced calcium signaling, co-treatment of 1 μ M of both antagonists together significantly suppressed the enhanced calcium signaling.

[399] These results clearly demonstrate that co-treatment of small doses of antagonists targeting each protomer provides novel therapeutic tools to efficiently suppress CXCR4-GPCR α heteromer response while avoiding side effects associated with high doses of individual antagonists.

[400]

[401] **Example 4. Inhibition of internalization by GPCR α antagonists.**

[402] To further study if co-internalization of CXCR4 heterodimer was blocked by partner GPCR α antagonists, internalization inhibition assay was performed. As shown in FIGS. 4b-4q, CXCR4-GFP expressing U-2 OS cells were co-internalized by partner GPCR α specific agonist when CXCR4 and GPCR α were simultaneously transduced to cells (control: CXCR4-GFP (FIG. 4a)). If CXCR4 formed heterodimer with GPCR α and co-internalized by partner GPCR α , it can be blocked by GPCR α specific antagonist.

[403] U-2 OS cells stably expressing CXCR4-GFP were transduced with Adenoviruses encoding GPCR α (ADRB2, CHRM1, HRH1). After 2 days, images were obtained before and 20 minutes after stimulating cells with CXCR4-specific agonist, CXCL12 (SDF-1) (20 nM), and/ or GPCR α -specific antagonist (10 μ M). Internalization of CXCR4-GFP was observed as GFP granules using IN Cell Analyzer 2500. Loss of GFP expression on the cell surface or appearance of GFP granules inside the cells were considered as CXCR4-GFP co-internalization. CXCR4 agonist, CXCL12 induced internalization of CXCR4-GFP with GPCR α (FIGS. 10a-c, first columns). GPCR α antagonists, treated as follows, had no effect on the internalization of the heteromer (FIGS. 10a-c, second columns): Carvedilol, an ADRB2 antagonist (FIG. 10a); Oxybutynin and Umeclidinium, CHRM1 antagonists (FIG. 10b); Prometazine, Hydroxyzine, and Loratadine, HRH1 antagonists (FIG. 10c). Internalization of CXCL12 stimulated CXCR4-GFP with GPCR α were inhibited by GPCR α specific antagonists (FIGS. 10a-c, third columns).

[404] These data demonstrate that CXCR4-GPCR α co-internalization was heteromer specific event and CXCR4 and GPCR α formed heterodimer. These data further demonstrate that, by inhibition of internalization of CXCR4 heteromer, abnormal downstream signal in CXCR4-GPCR α heteromer overexpressed cells, such as cancers, can be blocked for therapeutics purposes.

[405]

[406] **Example 5. Evaluation by cell proliferation assay of phenotypic effects of**

inhibitor of CXCR4-GPCRx heteromer signaling on tumor growth.

- [407] Several CXCR4 antagonists have been developed, but none have been approved as anticancer drugs until now. To overcome the limitation of CXCR4 inhibitors and to develop CXCR4 heteromer-based therapeutic agent, we tested the effect of GPCR α antagonist on cell proliferation.
- [408] We prepared single cell suspensions of resected and dissociated glioblastoma tissues from patients. (provided by Samsung Seoul hospital in Seoul, Korea) These cells were cultured under conditions optimal for propagation and non-differentiation of normal neuronal stem cell. Media were composed serum free Neurobasal media supplemented with basic FGF and EGF.
- [409] The effect of GPCR α antagonist on the survival of patient derived cells (PDCs) were assessed using ATPlite (PerkinElmer, Cat. No. 6016739 reagent. ATPlite is an Adenosine Triphosphate monitoring system based on firefly luciferase. This luminescence assay is the alternative to colorimetric, fluorometric and radioisotopic assays for the quantitative evaluation of proliferation and cytotoxicity of cultured mammalian cells. Cells were seeded in 384-well plate at 500 cells/well in 40 μ l culture media. After overnight growth, the cells were cultured for 7 days in the presence of several dose of GPCR α antagonist or DMSO alone. After the 7-day incubation, 15 μ l ATPlite was added into the each well and the plate were shaken for 5 minutes in an orbital shaker at 700 rpm. The luminescent signal was detected within 30 minutes at PerkinElmer TopCount detection instrument. The cell viability was calculated using the equation: Cell viability (%) = (OD of antagonist treatment/OD of DMSO only treatment) x 100%.
- [410] As shown in FIGS. 11a-c, when Carvedilol, ADRB2 specific antagonist, was treated in PDC expressing CXCR4 and ADRB2, growth of cells was inhibited significantly. (IC₅₀=11.69 μ M, FIG. 11a). Oxybutynin and Umeclidinium, CHRM1 antagonists also inhibited on the survival of PDC. Oxybutynin or Umeclidinium each showed significantly decreased survival of cells at IC₅₀=3.04 μ M or 4.03 μ M, respectively. (FIG. 11b). Promethazine, Hydroxyzine, and or Loratadine as a HRH1 antagonist each showed decreased survival of PDC at IC₅₀=18.39 μ M, 12.79 μ M or 5.29 μ M, respectively.(FIG. 11c).
- [411] These results demonstrate that CXCR4 heteromer induced abnormal cell proliferation can be blocked by partner GPCR α specific antagonist in CXCR4-GPCRx heteromer expressing cells, indicating that inhibition of cancer cell growth using partner GPCR α antagonist in CXCR4-GPCRx heteromer bearing patients can overcome the limitations of CXCR4 inhibitors alone as cancer therapeutics.

[412]

[413] **Example 6. Evaluation of CXCR4-GPCRx heteromer formation in patient**

derived cells (PDC) using Proximity ligation assay (PLA).

- [414] To investigate the existence of GPCR complexes in native tissues, various approaches such as atomic force microscopy (Fotiadis et al., 2006), co-immunoprecipitation (Gomes et al., 2004) and binding or functional assays (Wreggett and Wells, 1995) have been used. The most convenient methods to monitor interactions are based on resonance energy transfer performed with labeled proteins. The labeling can be performed by selective probes such as antibodies or fluorescent ligands (Roess et al., 2000; Patel et al., 2002).
- [415] Bazin et al. employed a time-resolved fluorescence resonance energy transfer (TR-FRET)-based approach that offers a much higher signal-to-noise ratio (Bazin et al., 2002). FRET is based on the transfer of energy between two fluorophores, a donor and an acceptor, when in close proximity. Molecular interactions between biomolecules can be assessed by coupling each partner with a fluorescent label and by detecting the level of energy transfer. Introducing a time delay of approximately 50 to 150 μ seconds between the system excitation and fluorescence measurement allows the signal to be cleared of all non-specific short-lived emissions.
- [416] Proximity ligation assay (PLA) is a technology that extends the capabilities of traditional immunoassays to include direct detection of proteins, protein interactions and modifications with high specificity and sensitivity (Gullberg et al., 2004). Two primary antibodies raised in different species recognize the target antigen on the proteins of interest. Secondary antibodies directed against the constant regions of the different primary antibodies, called PLA probes, bind to the primary antibodies. Each of the PLA probes has a unique short DNA strand attached to it. If the PLA probes are in close proximity (that is, if the two original proteins of interest are in close proximity, or part of a protein complex, as shown in the figures), the DNA strands can participate in rolling circle DNA synthesis when appropriate substrates and enzymes are added. The DNA synthesis reaction results in several-hundred fold amplification of the DNA circle. Next, fluorescent-labeled complementary oligonucleotide probes are added, and they bind to the amplified DNA. The resulting high concentration of fluorescence is easily visible as a distinct bright spot when viewed with a fluorescence microscope (Gustafsdottir et al., 2005).
- [417] CXCR4 overexpressing cell line, U2OS-CXCR4, was infected with ADRB2 expressing adenovirus, Ad-ADRB2 at the dose of 0, 2.5, 10, 40 MOIs for 2 days. PLA was performed as described previously (Brueggemann et al., 2014; Tripathi et al., 2014). To perform PLA, infected cells were fixed with 4% paraformaldehyde (PFA) on sixteen-well tissue culture slides. Slides were blocked with blocking solution provided by Duolink and incubated with mouse anti-CXCR4 (1:200, Santacruz, Sc-53534), Rabbit anti-ADRB2(1:200, Thermoscientific, PA5-33333), Rabbit anti-CHRM1(1:200,

Ls bio, Ls-C313301) at 37°C for 1 h in a humidifying chamber. Slides were then washed and incubated (1 h at 37°C) with secondary anti-rabbit and anti-mouse antibodies conjugated with plus and minus Duolink II PLA probes. Slides were washed again and then incubated with ligation-ligase solution (30 min at 37°C) followed by incubation with amplification-polymerase solution (2 h at 37°C). Slides were then mounted with minimal volume of Duolink II mounting medium with 4',6-diamidino-2-phenylindole (DAPI) for 15-30 min, and PLA signals [Duolink In Situ Detection Reagents Green (λ excitation/emission 495/527 nm) or Red (λ excitation/emission 575/623 nm) were identified as fluorescent spots under a IN Cell analyzer 2500.

[418] As shown in the FIGS. 12a-12b, the PLA signal increases in a dose dependent manner as the expression level of ADRB2. FIG. 12a: Images of PLA signal from U2OS cells expressing CXCR4-ADRB2 heteromer over a series of MOIs (multiplicity of infection). FIG. 12b: The red signal spots were counted and calculated by normalization against negative control. The PLA signal increased proportionate to the expression level of ADRB2 in a dose dependent manner. FIG. 12c: To investigate endogenous ADRB2 expression, qRT-PCR was performed with ADRB2 specific primers. As the results show, endogenous ADRB2 expression level of U2OS cell was quite high, indicating that the PLA signal is detected even in the section without virus infection (at 0 MOI for ADRB2).

[419] Traditionally Glioblastoma (GBM) is the most common and lethal primary brain tumor. Preclinical cancer biology has largely relied on the use of human cancer cell lines *in vitro* and the xenograft process of established these cell lines. However, the process of establishing conventional cell lines results in irreversible loss of important biological properties and, as a result, the xenograft tumor models do not maintain genomic and phenotypic characteristics present in the original tumor.

[420] Patient derived cell (PDC) derived directly from glioblastoma harbor extensive similarities to normal neural stem cells and recapitulate the genotype, gene expression patterns, and *in vivo* biology of human glioblastomas.

[421] To perform PLA with PDC samples, the patient derived cells were plated and fixed with 4 % PFA on sixteen-well tissue culture slides. Slides were blocked with blocking solution provided by Duolink and incubated with mouse anti-CXCR4 (1:200, Santa Cruz, Sc-53534), rabbit anti-ADRB2 (1:200, Thermo Scientific, PA5-33333), rabbit anti-CHRM1 (1:200, Lsbio, Ls-C313301) at 37 °C for 1 h in a humidifying chamber. Slides were then washed and incubated (1 h at 37 °C) with secondary anti-rabbit and anti-mouse antibodies conjugated with plus and minus Duolink II PLA probes. Slides were washed again and then incubated with ligation-ligase solution (30 min at 37 °C) followed by incubation with amplification-polymerase solution (2 h at 37 °C). Slides

were then mounted with minimal volume of Duolink II mounting medium with 4',6'-diamidino-2-phenylindole (DAPI) for 15-30 min, and PLA signals [Duolink In Situ Detection Reagents Green (λ excitation/emission 495/527 nm) or Red (λ excitation/emission 575/623 nm)] were identified as fluorescent spots under the IN Cell analyzer 2500.

[422] As shown in FIGS. 13a-13b and FIGS. 14a-14b, PLA ratio refers to the CXCR4 GPCR $\times$ heteromer, and the frequency of the heteromer formation varies depending on the patient. PLA ratio was calculated as: number of fluorescent spots in PDC sample / number of fluorescent spots in negative control. Negative control (NC) represents background fluorescence signal, indicated by the number of spots when only the secondary antibody conjugated with plus and minus Duolink II PLA probes is treated without primary antibody (mouse anti-CXCR4, rabbit anti-ADRB2, rabbit anti-CHRM1) treatment in PLA processing. These data demonstrate quantitative analysis of CXCR4-ADRB2 heterodimer in cancer patient samples.

[423] **Example 7. Evaluation of CXCR4-GPCR $\times$ heteromer formation *in vivo* using PDX model.**

[424] To perform PLA with PDX samples, the glioblastoma patient derived FFPE samples were used (provided by Samsung Seoul hospital in Seoul, Korea). After FFPE sample were de-paraffinized and performed heat induced antigen retrieval for 15 minutes at 100°C. Slides were blocked with blocking solution provided by Duolink and incubated with rabbit anti-CXCR4 (1:200, Thermoscientific, PA3305), mouse anti-ADRB2 (1:200, Santacruz, Sc-271322), at 37 °C for 1 h in a humidifying chamber. The other process was same as described above (PLA with PDC).

[425] In the FIG. 15a, nuclei were visualized with DAPI staining, and CXCR4-ADRB4 heteromers were stained with PLA as small dots. As shown in FIG. 15b, PLA ratio is different according to the patient and based on this result, indicating that it is possible to perform personalized medicine by the companion diagnostics.

[426]

[427] **Example 8. Evaluation by Ca²⁺ mobilization assay of enhanced CXCR4 downstream signaling upon CXCR4-GPCR $\times$ heteromer formation.**

[428] In cells overexpressing both CXCR4 and ADRB2, stimulation with salmeterol alone did not elicit calcium mobilization dose-dependently (FIG. 16a). But when the cells were co-stimulated with salmeterol in the presence of CXCL12, calcium signaling was greatly enhanced in a broad range of salmeterol concentrations, such as concentrations in the range of between 10 nM to 300 nM.

[429] Similarly, co-treatment with histamine and CXCL12 also significantly enhanced calcium responses in cells overexpressing both CXCR4 and HRH1 over a broad range of histamine concentrations (0.3 nM to 300 nM), even at histamine concentrations that

did not evoke any calcium responses when treated alone (0.3 nM and 1 nM) (FIG. 16b). Considering that the concentration of histamine in human plasma and glomeruli is below 10 nM and 2 μ M, respectively (Sedor and Abboud, 1984), this result indicates that enhanced calcium signaling by CXCR4-HRH1 heteromer can occur at physiological levels of histamine concentration *in vivo*.

[430] In cells expressing CXCR4-APLNR (FIG. 17a), CXCR4-PTGER3 (FIG. 17b), or CXCR4-SSTR2 (FIG. 17c), stimulation with GPCR α agonist (Apelin-13, PGE₂, or octreotide, respectively) alone increased calcium signal dose-dependently. However, unlike in cells expressing CXCR4-ADRB2 or CXCR4-HRH1, calcium signal was not enhanced any further upon co-stimulation with both agonists (CXCL12 and GPCR α agonist) in any dose ranges tested in cells expressing CXCR4-APLNR, CXCR4-PTGER3, or CXCR4-SSTR2. These results are consistent with the results shown in FIGS. 7a, 7d, 7e, respectively, and show that the lack of signal enhancement in cells expressing CXCR4-APLNR or CXCR4-SSTR2 upon co-stimulation was not due to the use of a specific dose of GPCR α agonist, but due to the intrinsic nature of these heteromers. These results also clearly demonstrate that the enhanced calcium responses shown in FIG. 6a-6l and FIGS. 16a-16b are unique features of CXCR4-GPCR α heteromers such as CXCR4-ADRB2 or CXCR4-HRH1 described in FIGS. 5c-5d and FIG. 6h, respectively.

[431]

[432] **Example 9. Evaluation of enhanced Ca²⁺ mobilization upon co-stimulation of endogenous CXCR4-HRH1 heteromer in MDA-MB-231 cells.**

[433] Using RT-qPCR, the endogenous expression levels of CXCR4 and HRH1 in MDA-MB-231 cells were measured. The threshold cycles (C_t) of CXCR4 and HRH1 were 28.5 and 27.7, respectively, when 12.5 ng of total RNA was analyzed. Stimulation of MDA-MB-231 cells with either CXCL12 (100 nM), or histamine (10 nM) alone, elicit weak calcium responses (FIG. 18a). But when the cells were co-stimulated with CXCL12 and histamine together, greatly enhanced calcium response was observed similar to the enhancement observed in cells overexpressing CXCR4 and HRH1 together (FIGS. 18a and 18b, compared with FIG. 16b). The result clearly indicates that the enhanced calcium response upon co-stimulation with both agonists could occur in native cells expressing both CXCR4 and HRH1 together.

[434]

[435] **Example 10. Evaluation of enhanced migration of cancer cells by CXCR4-HRH1 heteromer upon co-stimulation with both ligands.**

[436] Since MDA-MB-231 cells express about twice more HRH1 mRNA compared with CXCR4 mRNA when measured by RT-qPCR, the MDA-MB-231 cells were transduced with small amount of lentivirus encoding CXCR4 (1 MOI) and chemotactic

migration of the cells toward CXCL12 and HRH1 was measured (FIGS. 19a and 19b). Histamine (50 nM) alone, sufficient to produce calcium signaling as shown in FIG. 27a (see below), did not induce cell migration. On the other hand, when treated together with CXCL12, histamine significantly enhanced the migration of MDA-MB-231 cells towards CXCL12. However, in the presence of pyrilamine, a HRH1-selective inverse agonist, co-stimulation of the cells with CXCL12 and histamine failed to enhance the cell migration elicited by CXCL12. The results clearly demonstrate that the enhancement of observed cancer cell migration was specifically induced by HRH1, and not by other histamine receptor subtypes.

[437]

[438] **Example 11. Evaluation by Ca²⁺ mobilization assay of enhanced CXCR4 downstream signaling upon CXCR4-GPCR_x heteromer formation and inhibition of the enhanced signaling.**

[439] In cells overexpressing both CXCR4 and ADRB2, co-treatment of both agonists CXCL12, a CXCR4 agonist and salmeterol, an ADRB2-selective agonist significantly increased the calcium response compared to the ones evoked by individual agonists (FIG. 20). These results clearly demonstrate that CXCR4-ADRB2 heteromer exhibit properties distinct from those of the individual GPCRs.

[440] It was examined if enhanced calcium responses in cells co-expressing CXCR4 and ADRB2 upon co-stimulation with CXCL12 and ADRB2 ligands are inhibited by anti-CXCR4 antibody as a CXCR4 antagonist. In cells co-expressing CXCR4 and ADRB2, treatment of 2 µg of anti-CXCR4 antibody, 12G5, suppressed the enhanced calcium signaling as shown in FIG. 20. And co-treatment of both antagonists (Carvedilol, ADRB2 antagonist and 12G5, CXCR4 antagonist together) resulted in a more significant suppression. These results demonstrated that anti-CXCR4 antibody and ADRB2 antagonist could be used as an efficient therapeutics against CXCR4-ADRB2 heteromer.

[441] Utilizing the Calcium mobilization assay, MDA-MB-231 human breast cancer cells were seeded at 20,000 cells per well in a black clear bottom 96-well plate (Corning Costar, #3340) in 100 µL of RPMI 1640 supplemented with 10% FBS. The next day, cells were co-transduced with 10 MOI of CXCR4 and 30 MOI of GPCR_x. After 2 days, cells were treated with ADRB2 antagonist, Carvedilol (Tocris), anti-CXCR4 antibody, 12G5 (Thermo Scientific, 35-8800) with indicated amounts and incubated with Cal 6 (FLIPR® Calcium 6 Assay Kit by Molecular Devices, Cat. R8191) for 2 hr. And then, cells were stimulated with indicated amounts of CXCL12, ADRB2 agonist, or CXCL12 and ADRB2 agonist. Calcium mobilization was measured using FlexStation 3 Multi-Mode Microplate Reader. The results were normalized for baseline activity. Calcium mobilization was quantified by calculating the area-

under-the-curve (AUC) of each graph. Data were normalized to CXCL12-stimulated calcium response in cells expressing CXCR4 alone. Data represent three independent experiments (mean $\pm$ SEM). *P < 0.05; Student's t test.

[442]

[443] **Example 12. Evaluation of CXCR4-GPCR α heteromer formation in CXCR4-GPCR α expressing cell using PLA.**

[444] A process for screening a CXCR4-GPCR α heteromer is essential for the treatment of CXCR4-GPCR α targeted anticancer drug. First, for the quantitative detection of CXCR4 and GPCR α heteromer, PLA was performed in CXCR4-GPCR α expressing cells. CXCR4 overexpressing cell line, the U2OS-CXCR4 was infected with GPCR α expressing adenovirus, Ad-GPCR α at the dose of 0, 2.5, 10, 40 MOIs for 2 days. And then, the transduced cells were fixed with 4% paraformaldehyde and performed PLA was performed. The number of PLA signals means quantitatively demonstrating the formation of CXCR4-GPCR α heteromerization.

[445] As shown in the FIGS. 21a-21b, the PLA signal increased proportionate to the expression level of CHRM1 (FIG. 21a) and HRH1 (FIG. 21b) in a dose dependent manner. These results demonstrate detection of different types of CXCR-GPCR α heteromer and quantitative analysis of CXCR4-GPCR α heteromer in cancer patient samples.

[446] Utilizing the PLA in cells, CXCR4 overexpression cell lines, the U2OS-CXCR4 was infected with GPCR α expressing adenovirus, Ad-GPCR α at the dose of 0, 2.5, 10, 40 MOIs for 2 days. PLA was performed as described previously (Brueggemann et al., 2014; Tripathi et al., 2014). To perform PLA, infected cells were fixed with 4% PFA on sixteen-well tissue culture slides. Slides were blocked with blocking solution provided by Duolink and incubated with mouse anti-CXCR4 (Santacruz, Sc-53534), rabbit anti-CHRM1 (LS Bio, LS-C313301), or rabbit anti-HRH1 (Thermoscientific, PA5-27817) at 37°C for 1 h in a humidifying chamber. Slides were then washed and incubated (1 h at 37°C) with secondary anti-rabbit and anti-mouse antibodies conjugated with plus and minus Duolink II PLA probes. Slides were washed again and then incubated with ligation-ligase solution (30 min at 37°C) followed by incubation with amplification-polymerase solution (2 h at 37°C). Slides were then mounted with minimal volume of Duolink II mounting medium with 4',6-diamidino-2phenylindole (DAPI) for 15-30 min, and PLA signals [Duolink In Situ Detection Reagents Green (λ excitation/emission 495/527 nm) or Red (λ excitation/emission 575/623 nm)] were identified as fluorescent spots under a IN Cell analyzer 2500.

[447]

[448] **Example 13. Ca²⁺ Mobilization Inhibition of the Enhanced CXCR4 Downstream Signaling upon CXCR4-GPCR α heteromer Formation - Comparison**

of Single Inhibitor Treatment to Combination Inhibitor Treatment.

- [449] The data shown above (see Example 3, FIG. 8a) demonstrated that an increased Ca^{2+} signal due to CXCR4-ADRB2 heteromer formation was significantly decreased when treated simultaneously with a CXCR4 inhibitor and an ADRB2 inhibitor. The degree of inhibition (measured as IC_{50} values for Ca^{2+} response) was compared for a series of CXCR4 inhibitors, evaluated as single treatment in cells expressing CXCR4 alone (individual protomer context), as single treatment in CXCR4-ADRB2 heteromer-expressing cells, and as co-treatment with an ADRB2 inhibitor (Carvedilol; 10 μM) in CXCR4-ADRB2 heteromer-expressing cells.
- [450] MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 only, or CXCR4 and ADRB. The cells were cultured for 2 days and were treated CXCR4 inhibitor alone or co-treated with a CXCR4 inhibitor and an ADRB2 inhibitor (Carvedilol; 10 μM). The cells were then stained with Cal-6 for 2 hours and stimulated with CXCR4 agonist (CXCL12; 20 nM) and ADRB2 agonist (Salmeterol; 1 μM). Calcium mobilization was measured using FlexStation3, the results of which are shown in Table 6, showing IC_{50} of Ca^{2+} response in: (1) MDA-MB-231 cells expressing CXCR4 alone (individual protomer context), treated with CXCR4 inhibitor only (2nd column); (2) MDA-MB-231 cells expressing CXCR4-ADRB2 heteromer, treated simultaneously with CXCR4 inhibitor and ADRB2 inhibitor Carvedilol (3rd column); and (3) MDA-MB-231 cells expressing CXCR4-ADRB2 heteromer, treated only with CXCR4 inhibitor (4th column).

[451]

[452] [Table 6]

CXCR4 inhibitor	CXCR4	CXCR4+ADRB2 (+ Carvedilol)	CXCR4+ADRB2 (- Carvedilol)
	IC_{50} [nM]	IC_{50} [nM]	IC_{50} [nM]
AMD3100	57.80±20.36	0.053±0.07	28.65±13.23
AMD070	43.37±0.57	1.40±2.2	24.69±8.88
12G5	80.19±48.5	2.784±0.86	13.95±10.04
Ulocuplumab	0.42±0.09	0.0008±0.0002	1.12±0.56
BKT140	434±4.38	0.81±0.41	130.40±49.78
TZ14011	23.93±3.58	0.02±0.02	17.78±7.84

[453]

- [454] As shown in Table 6, the IC_{50} value of Ca^{2+} response was CXCR4 inhibitor dependent in the context of single treatment of MDA-MB-231 cells expressing CXCR4 alone, and in the single treatment context and the co-treatment context with an ADRB2 inhibitor of MDA-MB-231 cells expressing CXCR4-ADRB2 heteromer. Depending on the identity of the CXCR4 inhibitor, the IC_{50} value of Ca^{2+} response in

the CXCR4-ADRB2 heteromer context decreased by 1400 times or more when treated in combination with an ADRB2 inhibitor (3rd column), relative to single treatment with the CXCR4 inhibitor alone (4th column). For instance, the IC₅₀ value of Ca²⁺ response in the CXCR4-ADRB2 heteromer context decreased about 540 fold (from 28.65 nM to 0.053 nM), about 1400 fold (from 1.12 nM to 0.0008 nM), or about 890 fold (from 17.78 nM to 0.02 nM), upon co-treatment of Carvedilol with AMD3100, Ulocuplumab, or TZ14011, relative to single treatment with AMD3100, Ulocuplumab, or TZ14011, respectively. These results suggest that co-treating with a CXCR4 inhibitor and ADRB2 inhibitor more effectively inhibits increased Ca²⁺ response than single treatment with a CXCR4 inhibitor alone in CXCR4-ADRB2 heteromer expressing cells.

[455] To determine the possibility that CXCR4-ADRB2 heteromer formation induces conformation change and/or changes the binding affinity for the CXCR4 inhibitor, the IC₅₀ values of Ca²⁺ response were compared between single treatment with a CXCR4 inhibitor of cells expressing CXCR4 alone and cells expressing CXCR4-ADRB2 heteromer. The results showed that single treatment with a CXCR4 inhibitor altered the IC₅₀ value only by about 0.4-5 fold in cells expressing CXCR4-ADRB2 heteromer (4th column), relative to cells expressing CXCR4 alone (2nd column). These results suggest that co-treatment with CXCR4 inhibitor and ADRB2 inhibitor can increase, to a dramatic extent for certain CXCR4 inhibitors, the therapeutic efficacy towards CXCR4-ADRB2 heteromer expressing patients and/or patient cells/tissues, compared with single treatment with CXCR4 inhibitor.

[456]

[457] **Example 14. Effect of CXCR4 -ADRB2 heteromer on Tumor Growth.**

[458] To investigate the effect of CXCR4-ADRB2 heteromer on tumor growth, cell lines stably overexpressing CXCR4 alone or CXCR4-ADRB2 heteromer were prepared in A549 lung cancer cell, and the same amount of cells (1x10⁷ cell / head) were injected subcutaneously in nude mice to compare tumor growth rates. As shown in FIGS. 22a-22b, the tumor size at 28 days after transplantation was 351.4 ± 214.7 mm³ for A549, 726.9 ± 259.6 mm³ for A549-CXCR4, and 1012.2 ± 556.1 mm³ for A549-CXCR4-ADRB2. The tumor growth rate of mice transplanted with the A549-CXCR4 overexpressing CXCR4 was faster than that of mice bearing only parental A549, and the fastest tumor growth was observed in mice transplanted with cells overexpressing the CXCR4-ADRB2 heteromer. These results suggest that the formation of CXCR4-ADRB2 heteromer synergistically increases the Ca²⁺ response and thus promotes tumor growth.

[459] Images of three mice which were transplanted with either parental cell A549, A549-CXCR4 stably overexpressing CXCR4, or A549-CXCR4-ADRB2 stably over-

expressing CXCR4-ADRB2 heteromer, at 28 days after transplantation, are shown in FIG. 22a. These images show that the tumor size of the mouse bearing the cell expressing CXCR4-ADRB2 heteromer is the largest (accelerated the most) among them. The tumor growth rates of these three different mice over time are graphically illustrated in FIG. 22b. Tumor growth was monitored every third or fourth day by measuring the length (L) and width (W) of the tumor and calculating tumor volume on the basis of the following formula: $\text{Volume} = 0.5 \text{ LW}^2$. The mice bearing CXCR4-expressing cell show relatively fast tumor growth as compared to the mice bearing parental cell A549, and the tumor growth is the fastest in mice transplanted with cells overexpressing the CXCR4-ADRB2 heteromer.

[460]

[461] **Example 15. Evaluation by Ca^{2+} Mobilization Assay of Enhanced CXCR4 Downstream Signaling Upon Co-stimulation of CXCR4-GPCR α heteromer with Endogenous Ligands.**

[462] The enhancement of the calcium response in MDA-MB-231 cells co-expressing CXCR4 and GPCR α upon co-stimulation with CXCL12 and endogenous ligand for GPCR α , are shown in FIGS. 23a-23g. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and HA-VC, GPCR α and HA-VC, or CXCR4 and GPCR α where GPCR α represents ADCYAP1R1 (FIG. 23a), ADORA2B (FIG. 23b), ADORA3 (FIG. 23c), CHRM1 (FIG. 23d), EDNRB (FIG. 23e), MLNR (FIG. 23f), and TACR3 (FIG. 23g). Cells were treated with CXCL12 alone, GPCR α endogenous ligand alone, or CXCL12 and GPCR α ligand together. Calcium mobilization was calculated as described in FIG. 5d. Statistically significant difference between the sum (square with dots) and co-treatment (filled square) was determined by Student's *t* test. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; Mean $\pm$ SEM ($n = 3$).

[463] In FIG. 6a, enhanced Ca^{2+} signaling was observed in cells expressing CXCR4 and ADCYAP1R1 together, but not in cells expressing individual GPCR alone, upon co-treatment with CXCL12 and VIP, a non-selective endogenous ADCYAP1R1 ligand. When the cells were co-treated with CXCL12 and PACAP-38, an ADCYAP1R1-selective endogenous ligand, enhanced Ca^{2+} signaling was also observed (FIG. 23a). The result demonstrates that enhanced signaling of CXCR4-ADCYAP1R1 heteromer upon co-stimulation with CXCL12 is not only limited to VIP, but also selective ADCYAP1R1 natural ligand. It further suggests this enhancement could happen *in vivo* in cells expressing CXCR4 and ADCYAP1R1 together by their natural ligands.

[464] Enhanced Ca^{2+} signaling was observed in cells expressing CXCR4 and ADORA2B (FIG. 6b) or CXCR4 and ADORA3 together (FIG. 6c), but not in cells expressing individual GPCR alone, upon co-treatment with CXCL12 and BAY 60-6583

(ADORA2B-selective agonist), or CXCL12 and CI-IB-MECA (ADORA3-selective agonist), respectively. When these cells were co-treated with CXCL12 and adenosine, endogenous ligand for all adenosine receptors, enhanced Ca^{2+} signaling was also observed (FIGS. 23b and 23c). The result demonstrates that natural endogenous ligand, not only synthetic agonists, enhances downstream signaling of CXCR4-ADORA2B and CXCR4-ADORA3 heteromers upon co-stimulation with CXCL12. It further suggests that synergistic enhancement of downstream signaling could happen *in vivo* in cells expressing CXCR4 and ADORA2B or CXCR4 and ADORA3 together by their natural ligands.

[465] In FIG. 6f, enhanced Ca^{2+} signaling was observed in cells expressing CXCR4 and CHRM1 together, but not in cells expressing individual GPCR alone, upon co-treatment with CXCL12 and bethanechol, a synthetic CHRM1 agonist. When the cells were co-treated with CXCL12 and acetylcholine, endogenous ligand for all acetylcholine receptors, enhanced Ca^{2+} signaling was also observed (FIG. 23d). The result demonstrates that natural endogenous ligand, not only synthetic agonist, enhances signaling of CXCR4-CHRM1 heteromer upon co-stimulation with CXCL12. It further suggests that synergistic enhancement of downstream signaling could happen *in vivo* in cells expressing CXCR4 and CHRM1 together by their natural ligands.

[466] In FIG. 6g, enhanced Ca^{2+} signaling was observed in cells expressing CXCR4 and EDNRB together, but not in cells expressing individual GPCR alone, upon co-treatment with CXCL12 and BQ3020, an EDNRB-selective agonist. When the cells were co-treated with CXCL12 and endothelin-1, endogenous ligand for endothelin receptors, enhanced Ca^{2+} signaling was also observed (FIG. 23e). The result demonstrates that natural endogenous ligand, not only synthetic agonist, enhances signaling of CXCR4-EDNRB heteromer upon co-stimulation with CXCL12. It further suggests that synergistic enhancement of downstream signaling could happen *in vivo* in cells expressing CXCR4 and EDNRB together by their natural ligands.

[467] In FIG. 6i, enhanced Ca^{2+} signaling was observed in cells expressing CXCR4 and MLNR together, but not in cells expressing individual GPCR alone, upon co-treatment with CXCL12 and roxithromycin, an antibiotic as well as full MLNR agonist. When the cells were co-treated with CXCL12 and motilin, a selective endogenous ligand for MLNR, enhanced Ca^{2+} signaling was also observed (FIG. 23f). The result demonstrates that natural endogenous ligand, not only synthetic agonist, enhances signaling of CXCR4-MLNR heteromer upon co-stimulation with CXCL12. It further suggests that synergistic enhancement of downstream signaling could happen *in vivo* in cells expressing CXCR4 and MLNR together by their natural ligands.

[468] In FIG. 6l, enhanced Ca^{2+} signaling was observed in cells expressing CXCR4 and TACR3 together, but not in cells expressing individual GPCR alone, upon co-

treatment with CXCL12 and senktide, a TACR3-selective agonist. When the cells were co-treated with CXCL12 and neurokinin B, selective endogenous ligand for TACR3, enhanced Ca^{2+} signaling was also observed (FIG. 23g). The result demonstrates that natural endogenous ligand, not only synthetic agonist, enhances signaling of CXCR4-TACR3 heteromer upon co-stimulation with CXCL12. It further suggests that synergistic enhancement of downstream signaling could happen *in vivo* in cells expressing CXCR4 and EDNRB together by their natural ligands.

[469]

[470] **Example 16. Evaluation by Ca^{2+} Mobilization Assay of Enhanced CXCR4 Downstream Signaling Upon Co-stimulation of CXCR4-GPCR α heteromer.**

[471] In MDA-MB-231 cells co-expressing CXCR4 and ADCYAP1R1, stimulation with selective endogenous ligands CXCL12 alone or PACAP38 alone, respectively, produced calcium signaling dose-dependently (FIGS. 24a-24b). MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and ADCYAP1R1. As shown in FIG. 24a, addition of small amount of ADCYAP1R1 selective endogenous ligand (PACAP38; 1 nM) significantly enhanced calcium response in a broad range of CXCL12 concentrations compared with the sum values calculated by adding the responses obtained by PACAP38 alone and CXCL12 alone at indicated doses (FIG. 24a). Maximal Ca^{2+} response evoked by CXCL12 alone was taken as 100%. Sum represents the calculated additive value of the responses evoked by 1 nM of PACAP38 and CXCL12 alone at indicated doses. Similarly, as shown in FIG. 24b, addition of CXCR4 selective endogenous ligand (CXCL12; 15 nM) significantly enhanced downstream response in a broad range of PACAP38 concentrations compared with the sum values calculated by adding the responses obtained by CXCL12 alone and PACAP38 alone at indicated doses, even at concentrations that did not evoke any response when treated alone (0.03 ~ 0.3 nM) (FIG. 24b). Maximal Ca^{2+} response evoked by PACAP38 alone was taken as 100%. Sum represents the calculated additive value of the responses evoked by 15 nM of CXCL12 alone and PACAP38 alone at indicated doses. Statistically significant differences between the sum and co-treatment at each point were determined by Student's *t* test. **P* < 0.05; ***P* < 0.01; ****P* < 0.001; Mean $\pm$ SD (*n* = 3). These results suggest an enhanced response in cells co-expressing CXCR4 and ADCYAP1R1 *in vivo* in the presence of small amounts of CXCL12 and ADCYAP1R1 ligands together.

[472] The enhancement of calcium response in a broad range of ligand concentrations in cells co-expressing CXCR4 and TACR3 is shown in FIGS. 25a-25b. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 and TACR3. Stimulation of the cells with CXCL12 alone or Neurokinin B alone, respectively, produced calcium signaling dose-dependently (FIGS. 25a-25b). As shown in FIG. 25a, addition of small

amount of Neurokinin B (0.4 nM, TACR3-selective endogenous ligand) significantly enhanced calcium response in a broad range of CXCL12 concentrations compared with the sum values calculated by adding the responses obtained by Neurokinin B alone and CXCL12 alone at indicated doses. Maximal calcium response evoked by CXCL12 alone was taken as 100%. Sum represents the calculated additive value of the responses evoked by 0.4 nM of Neurokinin B alone and CXCL12 alone at indicated doses. Similarly, as shown in FIG. 25b, addition of CXCL12 significantly enhanced downstream response in a broad range of Neurokinin B concentrations compared with the sum values calculated by adding the responses obtained by CXCL12 alone and Neurokinin B alone at indicated doses. Maximal response evoked by Neurokinin B alone was taken as 100%. Sum represents the calculated additive value of the responses evoked by 30 nM of CXCL12 alone and Neurokinin B alone at indicated doses. Statistically significant differences between the sum and co-treatment at each point were determined by Student's *t* test. **P* < 0.05; ***P* < 0.01; Mean ± SD (*n* = 3). These results suggest an enhanced response in cells co-expressing CXCR4 and TACR3 *in vivo* in the presence of small amounts of CXCL12 and Neurokinin B together.

[473]

[474]

Example 17. Confirmation of a Loss of Heteromer-Specific Property Upon Deletion of One of the Protomers.

[475]

In FIG. 8f & 8g, enhanced calcium signaling was observed in MDA-MB-231 cells overexpressing CXCR4 alone upon co-treatment with CXCL12 and histamine. Using RT-qPCR, MDA-MB-231 cells were previously shown to express HRH1 as well as low level of CXCR4 mRNAs, with about twice more HRH1 mRNA being expressed compared with CXCR4 mRNA (Example 9). To confirm if the enhanced calcium signaling shown in FIG. 8f & 8g is due to the presence of endogenous HRH1 expression, and not to the presence of other histamine receptors, HRH1 gene in MDA CXCR4+ cells was deleted using CRISPR/Cas9 system. MDA-MB-231 cells stably expressing CXCR4 (MDA^{CXCR4+}) were transduced with lentiviruses encoding Cas9 and guide RNA targeting HRH1. The presence of functional HRH1 was detected by measuring calcium responses upon exposure to histamine. While MDA^{CXCR4+} cells showed dose-dependent increase in calcium signaling upon exposure to histamine, MDA^{CXCR4+, HRH1-} cells did not show any calcium response even at 1 μM histamine (FIG. 26a). The result implies almost complete absence of functional HRH1 in MDA CXCR4+, HRH1- cells.

[476]

When MDA-MB-231 cells stably overexpressing CXCR4 (MDA^{CXCR4+} cells) were treated with histamine, dose-dependent increase in calcium signaling was observed (FIG. 27a). When the cells were treated with histamine in the presence of CXCL12 (50 nM), significantly enhanced calcium response was obtained in a broad range of

histamine concentrations, resulting in increased potency and efficacy as evidenced by the changes in EC_{50} and E_{max} values. However, in MDA^{CXCR4+} cells depleted with HRH1 using CRISPR/Cas9 system (MDA^{CXCR4+, HRH1-} cells), no histamine-mediated response was obtained in the absence or presence of CXCL12 (50 nM), reconfirming the absence of functional HRH1 in these cells. Maximal calcium response evoked by histamine alone in MDA^{CXCR4+} cells was taken as 100%. Sum represents the calculated additive value of the responses evoked by 50 nM of CXCL12 alone and histamine alone at indicated doses.

[477] In MDA^{CXCR4+} cells, stimulation with CXCL12 produced calcium response dose-dependently (FIG. 27b). Addition of non-signaling concentration of histamine (15 nM) significantly enhanced calcium response in a broad range of CXCL12 concentrations, resulting in greatly increased potency and efficacy as evidenced by the changes in EC_{50} and E_{max} values. CXCL12-mediated calcium response in MDA^{CXCR4+, HRH1-} cells was similar to the response observed in MDA^{CXCR4+} cells. However, addition of histamine failed to increase CXCL12-mediated calcium response in MDA^{CXCR4+, HRH1-} cells, demonstrating the loss of heteromer-specific property upon deletion of HRH1. Maximal response evoked by CXCL12 alone in MDA^{CXCR4+} cells was taken as 100%. Sum represents the calculated additive value of the responses evoked by 15 nM of histamine alone and CXCL12 alone at indicated doses. Statistically significant differences between the sum and co-treatment at each point were determined by Student's *t* test. **P* < 0.05; ***P* < 0.01; ****P* < 0.001; Mean ± SD (n = 3). EC_{50} and E_{max} values were calculated using GraphPad Prism software.

[478] In FIGS. 18a-18b, significant enhancement in calcium signaling was also observed in wildtype MDA-MB-231 cells in the presence of CXCL12 and histamine together although individual ligand alone only produced faint signals. To confirm if the enhanced response is due to the endogenous expression of CXCR4, CXCR4 gene was targeted using CRISPR/Cas9 system, and expression of CXCR4 was detected using immunoblotting. Expression of CXCR4 was significantly decreased in MDA-MB-231 cells treated with guide RNA targeting CXCR4 compared to the expression in cells treated with control non-targeting guide RNA (FIG. 26b).

[479] Enhanced calcium response in MDA-MB-231 cells upon co-treatment with CXCL12 and histamine is abrogated in the absence of CXCR4. As shown in FIG. 28a, in MDA-MB-231 cells (MDA-MB-231 cells (MDA^{WT}), but not in MDA^{WT} cells depleted with CXCR4 using CRISPR/Cas9 system (MDA^{CXCR4-}), enhanced signaling in the presence of small amount of histamine (15 nM) was evident in a broad range of CXCL12 concentrations although CXCL12-mediated dose-dependent increase in calcium response was not observed due to the low level expression of CXCR. Maximal calcium response evoked by CXCL12 alone in MDA^{WT} cells was taken as 100%. Sum represents the

calculated additive value of the responses evoked by 15 nM of histamine alone and CXCL12 alone at indicated doses. Deletion of CXCR4 completely abrogate the enhanced signaling in the presence of histamine, demonstrating loss of heteromer-specific property upon deletion of CXCR4.

[480] Similarly, as shown in FIG. 28b, addition of CXCL12 enhanced histamine-mediated response in a broad range of histamine concentrations in MDA-MB-231 cells compared with the sum values calculated by adding the responses obtained by CXCL12 alone and histamine alone at indicated doses. The enhanced signaling shown in MDA-MB-231 cells was not observed in MDA^{CXCR4-} cells, although MDA^{CXCR4-} cells retained histamine responses. Maximal response evoked by histamine alone in MDA^{WT} cells was taken as 100%. Sum represents the calculated additive value of the responses evoked by 100 nM of CXCL12 alone and histamine alone at indicated doses. Statistically significant differences between the sum and co-treatment at each point were determined by Student's *t* test. ***P* < 0.01; ****P* < 0.001; Mean ± SD (*n* = 3). EC₅₀ and E_{max} values were calculated using GraphPad Prism software. Taken together, these results demonstrate that CXCR4-HRH1 heteromer is responsible for the enhanced signaling upon co-treatment with CXCL12 and histamine in wildtype MDA-MB-231 cells.

[481]

[482] **Example 18. Ca²⁺ Mobilization Inhibition of the Enhanced CXCR4 Downstream Signaling upon CXCR4-GPCR_x heteromer Formation - Comparison of Single Inhibitor Treatment to Combination Inhibitor Treatment.**

[483] Overview: As illustrated below (see Tables 7-14), co-treatment by various combinations of a CXCR4 inhibitor and a GPCR_x inhibitor in a series of CXCR4-GPCR_x heteromer-expressing cells results in a significantly reduced calcium response compared to single treatment by a CXCR4 inhibitor alone. The GPCR_x protomers of the series of CXCR4-GPCR_x heteromers evaluated in Tables 7-14 are: ADRB2, HRH1, ADCYAP1R1, C5AR1, CALCR, EDNRB, MLNR, and TACR3, respectively. MDA-MB-231 cells were transduced with adenoviruses encoding CXCR4 only, or CXCR4 and the specified GPCR_x. The cells were cultured for 2 days and were treated CXCR4 antagonist alone or co-treatment of CXCR4 antagonist and the specified GPCR_x antagonist. And then, cells were stained with Cal-6 for 2 hours and stimulated with CXCL12 (20 nM) and the specified GPCR_x agonist. Calcium mobilization was measured using FlexStation3. A value in the column labeled "None" is the IC₅₀ of Ca²⁺ response in the specified CXCR4-GPCR_x heteromer-expressing MDA-MB-231 treated with CXCR4 inhibitor only (in the absence of a specified GPCR_x antagonist). A value in the remaining columns is the IC₅₀ of Ca²⁺ response in the specified CXCR4-GPCR_x heteromer-expressing MDA-MB-231 from the simultaneous

treatment with CXCR4 antagonist and the specified GPCR $\times$ antagonist.

[484]

[485] The efficiency of CXCR4 inhibitors in suppressing the enhanced signaling of CXCR4-ADRB2 heteromer was measured in the absence or presence of the representative ADRB2 inhibitors Carazolol or Propranolol (**Table 7**). MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and ADRB2, and Ca²⁺ signaling was measured upon co-stimulating the cells with CXCL12 (20 nM) and Salmeterol (1 μ M). Inhibitors were treated 30 min before agonist stimulation. Ca²⁺ mobilization was measured as described in **Fig. 5**, and IC₅₀ values were calculated using GraphPad Prism software.

[486]

[487] [Table 7]

CXCR4 inhibitors	ADRB2 inhibitors		
	None	Carazolol (10 μ M)	Propranolol (10 μ M)
	IC ₅₀ [nM]	IC ₅₀ [nM]	IC ₅₀ [nM]
AMD3100	22.10 $\pm$ 8.53	2.69 $\pm$ 3.41	6.23 $\pm$ 4.90
Ulocuplumab	0.226 $\pm$ 0.144	0.009 $\pm$ 0.004	0.041 $\pm$ 0.024
BKT140	123.86 $\pm$ 76.91	26.66 $\pm$ 19.32	66.14 $\pm$ 53.77

[488]

[489] CXCR4 inhibitors AMD-3100, Ulocuplumab and BKT140 suppressed the CXCR4-ADRB2 signaling more efficiently (greater potency) in the presence of 10 μ M of Carazolol or Propranolol as shown with the reduced IC₅₀ values compared to the IC₅₀ values of CXCR4 inhibitors in the absence of ADRB2 inhibitors. Depending on the identity of the CXCR4 inhibitor, the IC₅₀ value of Ca response in the CXCR4-ADRB2 heteromer context decreased by up to about 25 times or more when treated in combination with an ADRB2 inhibitor (Carazolol or Propranolol), relative to single treatment with the CXCR4 inhibitor alone ("None" column). These results suggest that co-treating with a CXCR4 inhibitor and ADRB2 inhibitor more effectively inhibits increased Ca response than single treatment with a CXCR4 inhibitor alone in CXCR4-ADRB2 heteromer expressing cells.

[490]

The efficiency of CXCR4 inhibitors in suppressing the enhanced signaling of CXCR4-HRH1 heteromer was measured in the absence or presence of the representative HRH1 inhibitors Hydroxyzine, Prometazine, or Cyproheptadine (**Table 8**). MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and HRH1, and Ca²⁺ signaling was measured upon co-stimulating the cells with CXCL12 (20 nM) and Histamine (1 nM). Inhibitors were treated 30 min before agonist stimulation. Ca²⁺ mobilization was measured as described in **Fig. 5**, and IC₅₀ values were calculated using GraphPad Prism software.

[491]

[492] [Table 8]

CXCR4 inhibitors	HRH1 inhibitors			
	None	Hydroxyzine (10 μ M)	Promethazine (10 μ M)	Cyproheptadine (10 μ M)
	IC ₅₀ [nM]	IC ₅₀ [nM]	IC ₅₀ [nM]	IC ₅₀ [nM]
AMD3100	39.07 $\pm$ 14.5	1.58 $\pm$ 1.14	2.59 $\pm$ 2.06	0.75 $\pm$ 0.49
Ulocuplumab	0.5 $\pm$ 0.44	0.003 $\pm$ 0.003	0.000098 $\pm$ 0.0001	0.003 $\pm$ 0.001
BKT140	232.3 $\pm$ 91.95	53.75 $\pm$ 21.24	1.66 $\pm$ 1.13	9.71 $\pm$ 7.17

[493]

[494] CXCR4 inhibitors AMD-3100, Ulocuplumab and BKT140 suppressed the CXCR4-HRH1 signaling more efficiently (greater potency) in the presence of 10 μ M of Hydroxyzine, Promethazine, or Cyproheptadine as shown with the reduced IC₅₀ values compared to the IC₅₀ values of CXCR4 inhibitors in the absence of HRH1 inhibitors. Depending on the identity of the CXCR4 inhibitor, the IC₅₀ value of Ca response in the CXCR4-HRH1 heteromer context decreased by up to about 5,100 times or more when treated in combination with an HRH1 inhibitor (Hydroxyzine, Promethazine, or Cyproheptadine), relative to single treatment with the CXCR4 inhibitor alone ("None" column). These results suggest that co-treating with a CXCR4 inhibitor and HRH1 inhibitor more effectively inhibits increased Ca response than single treatment with a CXCR4 inhibitor alone in CXCR4-HRH1 heteromer expressing cells.

[495] The efficiency of CXCR4 inhibitors in suppressing the enhanced signaling of CXCR4-ADCYAP1R1 heteromer was measured in the absence or presence of the representative ADCYAP1R1 inhibitors M65 or PACAP-(6-38) (**Table 9**). MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and ADCYAP1R1, and Ca²⁺ signaling was measured upon co-stimulating the cells with CXCL12 (20 nM) and PACAP-38 (1 nM). Inhibitors were treated 30 min before agonist stimulation. Ca²⁺ mobilization was measured as described in **Fig. 5**, and IC₅₀ values were calculated using GraphPad Prism software.

[496]

[497] [Table 9]

CXCR4 inhibitors	ADCYAP1R1 inhibitors		
	None	M65 (1 μ M)	PACAP-(6-38) (1 μ M)
	IC ₅₀ [nM]	IC ₅₀ [nM]	IC ₅₀ [nM]
AMD3100	546.8 $\pm$ 337.8	20.21 $\pm$ 12.83	2.42 $\pm$ 0.09
BKT140	213.08 $\pm$ 142.72	7.46 $\pm$ 1.93	19.56 $\pm$ 3.1

[498]

[499] CXCR4 inhibitors AMD-3100 and BKT140 suppressed the CXCR4-ADCYAP1R1

signaling more efficiently (greater potency) in the presence of 1 μ M of M65 or 1 μ M of PACAP-(6-38) as shown with the reduced IC₅₀ values compared to the IC₅₀ values of CXCR4 inhibitors in the absence of ADCYAP1R1 inhibitors. Depending on the identity of the CXCR4 inhibitor, the IC₅₀ value of Ca response in the CXCR4-ADCYAP1R1 heteromer context decreased by up to about 225 times or more when treated in combination with an ADCYAP1R1 inhibitor (M65 or PACAP-(6-38)), relative to single treatment with the CXCR4 inhibitor alone ("None" column). These results suggest that co-treating with a CXCR4 inhibitor and ADCYAP1R1 inhibitor more effectively inhibits increased Ca response than single treatment with a CXCR4 inhibitor alone in CXCR4-ADCYAP1R1 heteromer expressing cells.

[500] The efficiency of CXCR4 inhibitors in suppressing the enhanced signaling of CXCR4-C5AR1 heteromer was measured in the absence or presence of the representative C5AR1 inhibitor W54011 (**Table 10**). MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and C5AR1, and Ca²⁺ signaling was measured upon co-stimulating the cells with CXCL12 (20 nM) and C5a (0.03 nM). Inhibitors were treated 30 min before agonist stimulation. Ca²⁺ mobilization was measured as described in **Fig. 5**, and IC₅₀ values were calculated using GraphPad Prism software.

[501]

[502] [Table 10]

CXCR4 inhibitors	C5AR1 inhibitors	
	None	W54011 (10 μ M)
	IC ₅₀ [nM]	IC ₅₀ [nM]
AMD3100	769.34 $\pm$ 240.77	434.00 $\pm$ 82.94
Ulocuplumab	3.80 $\pm$ 0.22	1.06 $\pm$ 0.25
BKT140	118.01 $\pm$ 56.52	9.72 $\pm$ 6.87

[503]

[504] CXCR4 inhibitors AMD-3100, BKT140, and Ulocuplumab suppressed the CXCR4-C5AR1 signaling more efficiently (greater potency) in the presence of 10 μ M of W54011, as shown with the reduced IC₅₀ values compared to the IC₅₀ values of CXCR4 inhibitors in the absence of C5AR1 inhibitors. Depending on the identity of the CXCR4 inhibitor, the IC₅₀ value of Ca response in the CXCR4-C5AR1 heteromer context decreased by up to about 12 times or more when treated in combination with an C5AR1 inhibitor (W54011), relative to single treatment with the CXCR4 inhibitor alone ("None" column). These results suggest that co-treating with a CXCR4 inhibitor and C5AR1 inhibitor more effectively inhibits increased Ca response than single treatment with a CXCR4 inhibitor alone in CXCR4-C5AR1 heteromer expressing

cells.

[505] The efficiency of CXCR4 inhibitors in suppressing the enhanced signaling of CXCR4-CALCR heteromer was measured in the absence or presence of the representative CALCR inhibitor CT-(8-32) (**Table 11**). MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and CALCR, and Ca²⁺ signaling was measured upon co-stimulating the cells with CXCL12 (20 nM) and calcitonin (100 nM). Inhibitors were treated 30 min before agonist stimulation. Ca²⁺ mobilization was measured as described in **Fig. 5**, and IC₅₀ values were calculated using GraphPad Prism software.

[506]

[507] [Table 11]

CXCR4 inhibitor	CALCR inhibitors	
	None	CT-(8-32) (salmon) (10 μ M)
	IC ₅₀ [nM]	IC ₅₀ [nM]
AMD3100	2677.67 $\pm$ 953.23	12.54 $\pm$ 14.7
BKT140	1041.6 $\pm$ 756.9	15.62 $\pm$ 3.71

[508]

[509] CXCR4 inhibitors AMD-3100 and BKT140 suppressed the CXCR4-CALCR signaling more efficiently (greater potency) in the presence of 10 μ M of CT-(8-32) as shown with the reduced IC₅₀ values compared to the IC₅₀ values of CXCR4 inhibitors in the absence of CALCR inhibitors. Depending on the identity of the CXCR4 inhibitor, the IC₅₀ value of Ca response in the CXCR4-CALCR heteromer context decreased by up to about 210 times or more when treated in combination with an CALCR inhibitor (CT-(8-32)), relative to single treatment with the CXCR4 inhibitor alone ("None" column). These results suggest that co-treating with a CXCR4 inhibitor and CALCR inhibitor more effectively inhibits increased Ca response than single treatment with a CXCR4 inhibitor alone in CXCR4-CALCR heteromer expressing cells.

[510] The efficiency of CXCR4 inhibitors in suppressing the enhanced signaling of CXCR4-EDNRB heteromer was measured in the absence or presence of the representative EDNRB inhibitors ambrisentan or bosentan (**Table 12**). MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and EDNRB, and Ca²⁺ signaling was measured upon co-stimulating the cells with CXCL12 (20 nM) and BQ3020 (0.5 nM). Inhibitors were treated 30 min before agonist stimulation. Ca²⁺ mobilization was measured as described in **Fig. 5**, and IC₅₀ values were calculated using GraphPad Prism software.

[511] [Table 12]

CXCR4 inhibitor	EDNRB inhibitors		
	None	Ambrisentan (10 μ M)	Bosentan (10 μ M)
	IC ₅₀ [nM]	IC ₅₀ [nM]	IC ₅₀ [nM]
AMD3100	926.10 $\pm$ 346.23	287.13 $\pm$ 88.75	169.12 $\pm$ 147.30
Ulocuplumab	14.11 $\pm$ 2.31	2.24 $\pm$ 0.18	4.68 $\pm$ 2.81
BKT140	2680.5 $\pm$ 3927.44	529.37 $\pm$ 122.26	8.50 $\pm$ 7.37

[512]

[513] CXCR4 inhibitors AMD-3100, BKT140, and Ulocuplumab suppressed the CXCR4-EDNRB signaling more efficiently (greater potency) in the presence of 10 μ M of ambrisentan or 10 μ M of bosentan, as shown with the reduced IC₅₀ values compared to the IC₅₀ values of CXCR4 inhibitors in the absence of EDNRB inhibitors. Depending on the identity of the CXCR4 inhibitor, the IC₅₀ value of Ca response in the CXCR4-EDNRB heteromer context decreased by up to about 315 times or more when treated in combination with an EDNRB inhibitor (ambrisentan or bosentan), relative to single treatment with the CXCR4 inhibitor alone ("None" column). These results suggest that co-treating with a CXCR4 inhibitor and EDNRB inhibitor more effectively inhibits increased Ca response than single treatment with a CXCR4 inhibitor alone in CXCR4-EDNRB heteromer expressing cells.

[514] The efficiency of CXCR4 inhibitors in suppressing the enhanced signaling of CXCR4-MLNR heteromer was measured in the absence or presence of the representative MLNR inhibitor MA-2029 (**Table 13**). MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and MLNR, and Ca²⁺ signaling was measured upon co-stimulating the cells with CXCL12 (20 nM) and motilin (0.2 nM). Inhibitors were treated 30 min before agonist stimulation. Ca²⁺ mobilization was measured as described in **Fig. 5**, and IC₅₀ values were calculated using GraphPad Prism software.

[515] [Table 13]

CXCR4 inhibitor	MLNR inhibitors	
	None	MA-2029 (10 μ M)
	IC ₅₀ [nM]	IC ₅₀ [nM]
AMD3100	946.1 $\pm$ 171.4	199.23 $\pm$ 64.34
Ulocuplumab	25.51 $\pm$ 16.48	2.25 $\pm$ 1.30
BKT140	643.6 $\pm$ 162.9	93.39 $\pm$ 155.20

[516]

[517] CXCR4 inhibitors AMD-3100, BKT140, and Ulocuplumab suppressed the CXCR4-MLNR signaling more efficiently (greater potency) in the presence of 10 μ M

of MA-2029 as shown with the reduced IC_{50} values compared to the IC_{50} values of CXCR4 inhibitors in the absence of MLNR inhibitor. Depending on the identity of the CXCR4 inhibitor, the IC_{50} value of Ca response in the CXCR4-MLNR heteromer context decreased by up to about 11 times or more when treated in combination with an MLNR inhibitor (MA-2029), relative to single treatment with the CXCR4 inhibitor alone ("None" column). These results suggest that co-treating with a CXCR4 inhibitor and MLNR inhibitor more effectively inhibits increased Ca response than single treatment with a CXCR4 inhibitor alone in CXCR4-MLNR heteromer expressing cells.

[518] The efficiency of CXCR4 inhibitors in suppressing the enhanced signaling of CXCR4-TACR3 heteromer was measured in the absence or presence of the representative TACR3 inhibitors SB 222200, osanetant, or talnetant (**Table 14**). MDA-MB-231 cells were co-transduced with adenoviruses encoding CXCR4 and TACR3, and Ca^{2+} signaling was measured upon co-stimulating the cells with CXCL12 (20 nM) and neurokinin B (0.3 nM). Inhibitors were treated 30 min before agonist stimulation. Ca^{2+} mobilization was measured as described in **Fig. 5**, and IC_{50} values were calculated using GraphPad Prism software.

[519] [Table 14]

CXCR4 inhibitor	TACR3 inhibitors			
	None	SB-222200 (10 μ M)	Osanetant (10 μ M)	Talnetant (10 μ M)
	IC_{50} [nM]	IC_{50} [nM]	IC_{50} [nM]	IC_{50} [nM]
AMD3100	2680.33 $\pm$ 210.49	343.13 $\pm$ 199.54	567.53 $\pm$ 322.61	179.63 $\pm$ 52.11
Ulocuplumab	50.65 $\pm$ 23.57	3.07 $\pm$ 1.18	4.61 $\pm$ 4.36	3.30 $\pm$ 0.89
BKT140	1497.67 $\pm$ 419.98	217.90 $\pm$ 92.86	194.33 $\pm$ 91.07	230.30 $\pm$ 33.86

[520]

[521] CXCR4 inhibitors AMD-3100, BKT140, and ulocuplumab suppressed the CXCR4-TACR3 signaling more efficiently (greater potency) in the presence of 10 μ M of SB-222200, 10 μ M of osanetant, or 10 μ M of talnetant as shown with the reduced IC_{50} values compared to the IC_{50} values of CXCR4 inhibitors in the absence of TACR3 inhibitors. Depending on the identity of the CXCR4 inhibitor, the IC_{50} value of Ca response in the CXCR4-TACR3 heteromer context decreased by up to about 16 times or more when treated in combination with an TACR3 inhibitor (SB-222200, osanetant, or talnetant), relative to single treatment with the CXCR4 inhibitor alone ("None" column). These results suggest that co-treating with a CXCR4 inhibitor and TACR3 inhibitor more effectively inhibits increased Ca response than single treatment with a CXCR4 inhibitor alone in CXCR4-TACR3 heteromer expressing cells.

[522]

[523] **Example 19. Anti-tumor effect of CXCR4-ADRB2 heteromer inhibitors on Tumor Growth.**

- [524] In Figure 22, we observed that the CXCR4 and ADRB2 overexpressing cells bearing mice showed a dramatic increase in tumor proliferation compared to the A549 parent cell bearing mice. This suggests that CXCR4-ADRB2 heteromer promotes tumor proliferation.
- [525] To investigate the anti-tumor effect of CXCR4-ADRB2 heteromer inhibitors on tumor growth, the A549-CXCR4-ADRB2, stably overexpressing CXCR4-ADRB2 heteromer (1×10^7 cell / head) was injected subcutaneously in nude mice. When tumor sizes were reached an average of 50-100mm³, CXCR4 inhibitor, ADRB2 inhibitor alone or combination of CXCR4 and ADRB2 inhibitors were treated.
- [526] Figure 29a is a graph comparing tumor growth rates for the *in vivo* antitumor effect of CXCR4 inhibitor LY2510924 (3 mg/kg), ADRB2 inhibitor Carvedilol (30 mg/kg), or combination of LY2510924 (3 mg/kg) and Carvedilol (30 mg/kg). Figure 29b is a graph comparing tumor growth rates for the antitumor effect of CXCR4 inhibitor AMD070 (10 mg/kg), ADRB2 inhibitor Carvedilol (30 mg/kg), or combination of AMD070 (10 mg/kg) and Carvedilol (30 mg/kg). The tumor growth was monitored every third or fourth day by measuring the length (*L*) and width (*W*) of the tumor and calculating tumor volume based on the following formula: Volume = $0.5 LW^2$.
- [527] As shown in **FIG. 29a**, CXCR4-ADRB2 overexpressing mice were more inhibited tumor growth by the CXCR4 inhibitor, LY2510924 or the ADRB2 inhibitor, Carvedilol than the control mice. In addition, the combination of LY2510924 and Carvedilol demonstrated that the tumor growth inhibitory effect was superior to the single administration group. More specifically CXCR4-ADRB2 overexpressing A549 bearing mice treated with vehicle as a control reached an average tumor volume of $554.2 \pm 152.7 \text{ mm}^3$ at day 21 post-treatment, as compared to LY2510924, Carvedilol or LY2510924 and Carvedilol, which reached an average tumor volume of $488.9 \pm 135.2 \text{ mm}^3$, and $432.0 \pm 206.4 \text{ mm}^3$, $356.3 \pm 125.4 \text{ mm}^3$ respectively, in the same time period.
- [528] A similar pattern was observed in the group treated with AMD070, another inhibitor of CXCR4.
- [529] **FIG. 29b** shows CXCR4-ADRB2 heteromer-overexpressing cell lines were transplanted at nude mouse and when tumor size reached 50-100 mm³, CXCR4 inhibitor, AMD070 or ADRB2 inhibitor, Carvedilol was administered orally alone or in combination for 23 days. As shown in **FIG. 29b**, the size of the tumor on the 23 days after administration of the drug was $618.5 \pm 190.9 \text{ mm}^3$ in the control group, $543.2 \pm 260.4 \text{ mm}^3$ or $510.4 \pm 139.9 \text{ mm}^3$ in the group treated with AMD070 or Carvedilol, respectively. The tumor size of drug treated group is shown to be smaller than that of control group. In addition, in the combination of AMD070 and Carvedilol, the size of the tumor was $418.0 \pm 238.4 \text{ mm}^3$, which indicates that the antitumor effect is superior to that of the single administration group.

[530] These results suggest that co-treatment of CXCR4 and ADRB2 inhibitors may provide a better therapeutic effect for CXCR4 and ADRB2 heteromer expressing patients.

[531]

[532] All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

[533] While preferred embodiments have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

[534]

[535] ***Exemplary Embodiments***

[536] In an embodiment, a method for treating cancer in a subject having a cell containing a CXCR4-GPCR_x heteromer, the method comprising: administering to the subject a therapeutically effective amount of an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer.

[537] In an embodiment, a method for treating cancer in a subject having CXCR4-GPCR_x heteromer, the method comprising: administering to the subject a therapeutically effective amount of an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein the cell containing the CXCR4-GPCR_x heteromer has enhanced downstream signaling.

[538] In an embodiment, a method for treatment, amelioration, or prevention of a cancer in a subject having CXCR4-GPCR_x heteromer, the method comprising: administering to the subject a therapeutically effective amount of an inhibitor of a CXCR4-GPCR_x heteromer, wherein: GPCR_x heteromerizes with CXCR4 in the subject, the heteromerization of GPCR_x with CXCR4 is accompanied by enhancement of signaling downstream of CXCR4; and the enhancement of signaling downstream of CXCR4 is suppressed by the inhibitor of the CXCR4-GPCR_x heteromer.

[539] In an embodiment, a method for assessing response, or potential response, of a subject having CXCR4-GPCR_x heteromer to treatment, amelioration, or prevention of a cancer, the method comprising: obtaining a sample from the subject; detecting heteromerization of CXCR4 and GPCR_x in the sample; and based at least in part on detection of the heteromerization of CXCR4 and GPCR_x, assessing the subject's response, or potential response, to the treatment, amelioration, or prevention of a

cancer.

[540] In an embodiment, a method for treating cancer in a patient having a cell containing a CXCR4-GPCR_x heteromer, the method comprising: administering to the patient an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein:

[541] i) the CXCR4-GPCR_x heteromer has enhanced downstream signaling; and

[542] ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient.

[543] In an embodiment, a method of suppressing enhanced downstream signaling from a CXCR4-GPCR_x heteromer in a cell of a patient suffering from cancer, the method comprising: administering to the patient an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein:

[544] i) the CXCR4-GPCR_x heteromer has enhanced downstream signaling; and

[545] ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient.

[546] In an embodiment, a pharmaceutical kit for use in treating cancer in a patient having a cell containing a CXCR4-GPCR_x heteromer, the pharmaceutical kit comprising: an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein the CXCR4-GPCR_x heteromer has enhanced downstream signaling.

[547] In an embodiment, a pharmaceutical composition for use in treating cancer in a patient having a cell containing a CXCR4-GPCR_x heteromer, the pharmaceutical composition comprising:

[548] i) an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; and

[549] ii) a pharmaceutically acceptable carrier;

[550] wherein the CXCR4-GPCR_x heteromer has enhanced downstream signaling.

[551] In an embodiment, a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR_x heteromer, wherein the CXCR4-GPCR_x heteromer has enhanced downstream signaling, the method comprising:

[552] 1) determining whether the patient has the cancer cell containing the CXCR4-GPCR_x heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if:

- [553] i) the patient's cancer cell contains said CXCR4-GPCR α heteromer; or
- [554] ii) a CXCR4-GPCR α heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR α heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient derived cell(s) containing the CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR α heteromer; and
- [555] 2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer to the cancer patient.
- [556] In an embodiment, a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, the method comprising:
- [557] 1) determining whether the patient's cancer cell contains the CXCR4-GPCR α heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR α heteromer is present in the patient's cancer cell; wherein:
- [558] a) the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and
- [559] b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay; and
- [560] 2) if the patient's cancer cell contains said CXCR4-GPCR α heteromer, then internally administering an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer to the patient.
- [561] In an embodiment, a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer having enhanced downstream signaling, the method comprising:
- [562] 1) determining whether the patient has the cancer cell containing the CXCR4-GPCR α heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if:
- [563] i) the patient's cancer cell contains said CXCR4-GPCR α heteromer; or
- [564] ii) a CXCR4-GPCR α heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR α heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient derived cell(s) containing said

CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; and

[565] 2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and

[566] 3) if the patient does not have a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR α inhibitor as a single inhibitor.

[567] In an embodiment, a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer having enhanced downstream signaling, the method comprising:

[568] 1) determining whether the patient has the cancer cell containing the CXCR4-GPCR α heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if:

[569] i) the patient's cancer cell contains said CXCR4-GPCR α heteromer; or

[570] ii) a CXCR4-GPCR α heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR α heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; and

[571] 2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and

[572] 3) if the patient does not have a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR α inhibitor as a single inhibitor;

[573] wherein:

[574] a) progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR α heteromer is decreased in the range of 5-100% more upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient;

[575] b) efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor; and/or

- [576] c) efficacy of a GPCR α inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor.
- [577] In an embodiment, a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, the method comprising:
- [578] 1) determining whether the patient's cancer cell contains the CXCR4-GPCR α heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR α heteromer is present in the patient's cancer cell; wherein:
- [579] a) the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and
- [580] b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, enzyme-linked immunosorbent assay (ELISA), flow cytometry, RNAseq, qRT-PCR, microarray, or a fluorescent animal assay; and
- [581] 2) if the patient's cancer cell contains said CXCR4-GPCR α heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and
- [582] 3) if the patient's cancer cell does not contain said CXCR4-GPCR α heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR α inhibitor as a single inhibitor.
- [583] In an embodiment, a method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, the method comprising:
- [584] 1) determining whether the patient's cancer cell contains the CXCR4-GPCR α heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR α heteromer is present in the patient's cancer cell; wherein:
- [585] a) the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and
- [586] b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, enzyme-linked immunosorbent assay (ELISA), flow

cytometry, RNAseq, qRT-PCR, microarray, or a fluorescent animal assay; and

[587] 2) if the patient's cancer cell contains said CXCR4-GPCR α heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and

[588] 3) if the patient's cancer cell does not contain said CXCR4-GPCR α heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR α inhibitor as a single inhibitor;

[589] wherein:

[590] a) progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR α heteromer is decreased in the range of 5-100% more upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient;

[591] b) efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor; and/or

[592] c) efficacy of a GPCR α inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor.

[593] In certain embodiments, one or more than one (including for instance all) of the following further embodiments may comprise each of the other embodiments or parts thereof.

[594] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer has, causes, or produces, the enhanced downstream signaling.

[595] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling results from the CXCR4-GPCR α heteromer.

[596] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling results from agonism of the CXCR4-GPCR α heteromer.

[597] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and

any one or more of the further embodiments herein, wherein the enhanced downstream signaling results from CXCR4 agonism of the CXCR4-GPCR_x heteromer.

[598] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling results from GPCR_x agonism of the CXCR4-GPCR_x heteromer.

[599] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling results from CXCR4 agonism and GPCR_x agonism of the CXCR4-GPCR_x heteromer.

[600] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling is downstream of the CXCR4, the respective GPCR_x, or the CXCR4-GPCR_x heteromer.

[601] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling is downstream of the CXCR4.

[602] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling is downstream of the respective GPCR_x.

[603] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling is downstream of the CXCR4-GPCR_x heteromer.

[604] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling from the CXCR4-GPCR_x heteromer is relative to downstream signaling from a CXCR4 protomer or a respective GPCR_x protomer in their respective individual protomer context.

[605] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling from the CXCR4-GPCR_x heteromer is relative to downstream signaling from

a CXCR4 protomer in an individual protomer context.

- [606] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling from the CXCR4-GPCR α heteromer is relative to downstream signaling from a respective GPCR α protomer in an individual protomer context.
- [607] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling from the CXCR4-GPCR α heteromer is relative to downstream signaling from a CXCR4 protomer and a respective GPCR α protomer in their respective individual protomer context.
- [608] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the cancer patient.
- [609] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the patient's cancer cells.
- [610] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method further comprises determining or diagnosing the presence of a CXCR4-GPCR α heteromer.
- [611] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method further comprises determining or diagnosing the presence of a CXCR4-GPCR α heteromer in a cancer patient.
- [612] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method further comprises determining or diagnosing the presence of a CXCR4-GPCR α heteromer in a cancer cell or cancer tissue.
- [613] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and

any one or more of the further embodiments herein, wherein the method further comprises determining or diagnosing the presence of a CXCR4-GPCR α heteromer in a cancer cell or cancer tissue obtained from a cancer patient.

- [614] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method further comprises determining or diagnosing the presence of a CXCR4-GPCR α heteromer; and wherein the determining or diagnosing comprises:
- [615] 1) obtaining or having obtained a biological sample (e.g., cells or tissue, such as cells or tissue from a cancer patient); and
- [616] 2) performing or having performed an assay on the biological sample, wherein the assay is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay.
- [617] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling from the CXCR4-GPCR α heteromer is determined by an intracellular Ca²⁺ assay.
- [618] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced downstream signaling is an enhanced amount of calcium mobilization.
- [619] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced amount of calcium mobilization from the CXCR4-GPCR α heteromer is a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is at least 10% greater, at least 20% greater, at least 30% greater, at least 40% greater, at least 50% greater, at least 75% greater, or at least 90% greater, than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.
- [620] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced amount of calcium mobilization from the CXCR4-GPCR α heteromer is a calcium mobilization

amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is between 10-100% greater than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.

- [621] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced amount of calcium mobilization from the CXCR4-GPCR α heteromer is a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is between 25-100% greater than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.
- [622] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced amount of calcium mobilization is determined by an intracellular Ca²⁺ assay.
- [623] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the intracellular Ca²⁺ assay is a calcium mobilization assay.
- [624] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced amount of calcium mobilization is a synergistic amount of calcium mobilization.
- [625] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the synergistic amount of calcium mobilization from the cells containing the CXCR4-GPCR α heteromer is a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is at least 10% greater, at least 20% greater, at least 30% greater, at least 40% greater, at least 50% greater, at least 75% greater, or at least 90% greater, than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.
- [626] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and

any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer has two or more of the following characteristics:

- [627] 1) the CXCR4-GPCR α heteromer components in a cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allostereism, as determined via one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay;
- [628] 2) an enhanced amount of calcium mobilization, such that:
- [629] a) either CXCR4 or the respective GPCR α in an individual protomer context in a cell, upon co-stimulation with CXCL12 and a respective selective GPCR α agonist, results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; and
- [630] b) the CXCR4-GPCR α heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist;
- [631] as determined via a calcium mobilization assay; or
- [632] 3) a CXCR4-GPCR α heteromer-selective reagent:
- [633] i) alters heteromer-specific properties of the CXCR4-GPCR α heteromer in a patient derived cell;
- [634] ii) alters heteromer-specific function of the CXCR4-GPCR α heteromer in a patient derived cell;
- [635] iii) alters heteromer-specific properties of a patient derived cell containing the CXCR4-GPCR α heteromer; or
- [636] iv) decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR α heteromer.
- [637] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific properties of the CXCR4-GPCR α heteromer in a patient derived cell, as determined by at least one of the following methods: PLA, radioligand binding assays, [35S]GTP-rS Binding assays, Calcium assay, cAMP assay, GTPase assay, PKA activation, ERK1/2 and/or Akt/PKB Phosphorylation assays, Src and STAT3 phosphorylation assays, CRE-reporter assay, NFAT-RE-reporter assay, SRE-reporter assay, SRF-RE reporter assay, NF-kB-RE reporter assay, Secreted alkaline phosphatase Assay, Inositol 1-Phosphate Production

assay, Adenylyl Cyclase Activity assay, analysis of target gene expression by RT-PCR, RT-qPCR, RNAseq, or microarray.

[638] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR_x heteromer-selective reagent alters heteromer-specific function of the CXCR4-GPCR_x heteromer in a patient derived cell, as determined by at least one of the following methods: PLA, radioligand binding assays, [35S]GTP-rS Binding assays, Calcuim assay, cAMP assay, GTPase assay, PKA activation, ERK1/2 and/or Akt/PKB Phosphorylation assays, Src and STAT3 phosphorylation assays, CRE-reporter assay, NFAT-RE-reporter assay, SRE-reporter assay, SRF-RE reporter assay, NF-κB-RE reporter assay, Secreted alkaline phosphatase Assay, Inositol 1-Phosphate Production assay, Adenylyl Cyclase Activity assay, analysis of target gene expression by RT-PCR, RT-qPCR, RNAseq, next generation sequencing (NGS), or microarray.

[639] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR_x heteromer-selective reagent alters heteromer-specific properties of a patient derived cell containing the CXCR4-GPCR_x heteromer, as determined by at least one of the following methods: assays on proliferation, migration, invasion, and drug resistance (survival) of cancer cells, modulation of immune cell function, angiogenesis, vasculogenesis, metastasis, drug resistance, tissue microarray (TMA), and cancer cell-tumor microenvironment (TME) interaction.

[640] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR_x heteromer components comprise individual protomers CXCR4 and GPCR_x.

[641] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the cell containing either the CXCR4 or the respective GPCR_x in an individual protomer context comprises, independently:

[642] i) the individual protomer CXCR4 in the absence of the respective individual protomer GPCR_x; or

[643] ii) the respective individual protomer GPCR_x in the absence of the individual protomer CXCR4;

[644] respectively.

[645] In a further embodiment, the method of treating, method of suppressing, pharma-

ceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the cell containing the CXCR4 in an individual protomer context comprises said individual protomer CXCR4 in the absence of the respective individual protomer GPCR_x.

[646] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the cell containing the respective GPCR_x in an individual protomer context comprises said respective individual protomer GPCR_x in the absence of the individual protomer CXCR4.

[647] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR_x heteromer components in the cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allostereism, as determined via one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, or a fluorescent animal assay.

[648] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the proximity-based assay is, or comprises, resonance energy transfer (RET), bioluminescence RET (BRET), fluorescence RET (FRET), time-resolved fluorescence RET (TR-FRET), antibody-aided FRET, ligand-aided FRET, bimolecular fluorescence complementation (BiFC), or a proximity ligation assay (PLA).

[649] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR_x heteromer components in the cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allostereism, as determined via one or more of the following: a co-internalization assay, bimolecular fluorescence complementation (BiFC), or a proximity ligation assay (PLA).

[650] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the patient's cancer cell contains the CXCR4-GPCR_x heteromer.

[651] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR_x

heteromer exhibits the enhanced amount of calcium mobilization, such that:

- [652] a) either the CXCR4 or the respective GPCR_x in an individual protomer context in the cell upon co-stimulation with CXCL12 and a respective selective GPCR_x agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR_x agonist; and
- [653] b) the CXCR4-GPCR_x heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR_x agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR_x agonist;
- [654] as determined via a calcium mobilization assay.
- [655] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein:
- [656] i) the calcium mobilization from the protomer CXCR4 or GPCR_x, in the individual protomer context in the cell, is non-synergistic, as determined via calcium mobilization assay; and
- [657] ii) the calcium mobilization from the CXCR4-GPCR_x heteromer in the cell is synergistic, as determined via a calcium mobilization assay.
- [658] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein in the individual protomer context:
- [659] a) the individual protomer CXCR4 in the cell, in the absence of the respective individual protomer GPCR_x; or
- [660] b) the respective individual protomer GPCR_x in the cell, in the absence of the individual protomer CXCR4;
- [661] upon co-stimulation with CXCL12 and a respective selective GPCR_x agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay.
- [662] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein in the individual protomer context, independently:
- [663] a) the individual protomer CXCR4 in the cell, in the absence of the respective individual protomer GPCR_x; and

- [664] b) the respective individual protomer GPCR_x in the cell, in the absence of the individual protomer CXCR4;
- [665] upon co-stimulation with CXCL12 and a respective selective GPCR_x agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay.
- [666] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR_x heteromer upon co-stimulation with the CXCL12 and the respective selective GPCR_x agonist results in a calcium mobilization amount that is greater than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay.
- [667] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the calcium mobilization amount resulting from the co-stimulation of the CXCR4-GPCR_x heteromer is an enhanced amount of calcium mobilization, relative to the sum of calcium mobilizations resulting from single agonist stimulation of said CXCR4-GPCR_x heteromer, as determined via a calcium mobilization assay.
- [668] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced amount of calcium mobilization is at least 10% greater, at least 20% greater, at least 30% greater, at least 40% greater, at least 50% greater, at least 75% greater, or at least 90% greater, than the sum of calcium mobilizations resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay.
- [669] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the enhanced amount of calcium mobilization is a synergistic amount of calcium mobilization.
- [670] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method administers a CXCR4-GPCR_x heteromer-selective reagent.

- [671] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent:
- [672] i) alters heteromer-specific properties of the CXCR4-GPCR α heteromer in the patient derived cell;
- [673] ii) alters heteromer-specific function of the CXCR4-GPCR α heteromer in the patient derived cell;
- [674] iii) alters heteromer-specific properties of the patient derived cell containing the CXCR4-GPCR α heteromer; or
- [675] iv) decreases cell proliferation of the patient derived cell containing the CXCR4-GPCR α heteromer.
- [676] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific properties of the CXCR4-GPCR α heteromer in the patient derived cell.
- [677] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific function of the CXCR4-GPCR α heteromer in the patient derived cell.
- [678] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific properties of the patient derived cell containing the CXCR4-GPCR α heteromer.
- [679] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent decreases cell proliferation of the patient derived cell containing the CXCR4-GPCR α heteromer.
- [680] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent is a CXCR4 inhibitor, a GPCR α inhibitor, or a CXCR4-GPCR α heteromer inhibitor.
- [681] In a further embodiment, the method of treating, method of suppressing, pharma-

ceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent is a CXCR4 antagonist, a GPCR α antagonist, or a CXCR4-GPCR α heteromer antagonist.

[682] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer-selective reagent is administered as a pharmaceutical composition.

[683] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method administers an inhibitor selected from the group consisting of: the CXCR4 inhibitor, the GPCR α inhibitor, or the CXCR4-GPCR α heteromer inhibitor.

[684] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the inhibitor is administered as a pharmaceutical composition.

[685] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the administered inhibitor is the CXCR4 inhibitor.

[686] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the administered inhibitor is the GPCR α inhibitor.

[687] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the administered inhibitor is the inhibitor of the CXCR4-GPCR α heteromer.

[688] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method administers the combination of inhibitors selected from the group consisting of: the CXCR4 inhibitor, the GPCR α inhibitor, and the inhibitor of the CXCR4-GPCR α heteromer.

[689] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the combination of inhibitors is administered as a pharmaceutical composition.

- [690] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the combination of inhibitors comprises the CXCR4 inhibitor and the GPCR $\times$ inhibitor.
- [691] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the combination of inhibitors comprises the CXCR4 inhibitor and the inhibitor of the CXCR4-GPCR $\times$ heteromer.
- [692] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the combination of inhibitors comprises the GPCR $\times$ inhibitor and the inhibitor of the CXCR4-GPCR $\times$ heteromer.
- [693] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the combination of inhibitors are administered sequentially, concurrently, or simultaneously.
- [694] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the combination of inhibitors are administered as a pharmaceutical composition further comprising a pharmaceutically acceptable carrier.
- [695] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the combination of inhibitors are administered as separate pharmaceutical compositions, wherein the separate pharmaceutical compositions independently further comprise a pharmaceutically acceptable carrier.
- [696] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4 inhibitor is an antagonist of CXCR4, an inverse agonist of CXCR4, a partial antagonist of CXCR4, an allosteric modulator of CXCR4, an antibody of CXCR4, an antibody fragment of CXCR4, or a ligand of CXCR4.
- [697] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR $\times$ inhibitor is an

antagonist of GPCR_x, an inverse agonist of GPCR_x, a partial antagonist of GPCR_x, an allosteric modulator of GPCR_x, an antibody of GPCR_x, an antibody fragment of GPCR_x, or a ligand of GPCR_x.

- [698] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the inhibitor of the CXCR4-GPCR_x heteromer is an antagonist of the CXCR4-GPCR_x heteromer, an inverse agonist of the CXCR4-GPCR_x heteromer, a partial antagonist of the CXCR4-GPCR_x heteromer, an allosteric modulator of the CXCR4-GPCR_x heteromer, an antibody of the CXCR4-GPCR_x heteromer, an antibody fragment of the CXCR4-GPCR_x heteromer, a ligand of the CXCR4-GPCR_x heteromer, or a protein-protein interaction (PPI) inhibitor of the CXCR4-GPCR_x heteromer.
- [699] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4 inhibitor, the GPCR_x inhibitor, or the inhibitor of the CXCR4-GPCR_x heteromer, is an antibody-drug conjugate.
- [700] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a therapeutically effective amount of the inhibitor of the CXCR4-GPCR_x heteromer is administered to the patient.
- [701] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a sub-therapeutically effective amount of the inhibitor of the CXCR4-GPCR_x heteromer is administered to the patient.
- [702] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a therapeutically effective amount of the CXCR4 inhibitor is administered to the patient.
- [703] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a sub-therapeutically effective amount of the CXCR4 inhibitor is administered to the patient.
- [704] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4 inhibitor is

selected from the group consisting of:

- [705] AD-114, AD-114-6H, AD-114-Im7-FH, AD-114-PA600-6H, ALX-0651, ALX40-4C, AMD070 (AMD11070, X4P-001), AMD3100 (plerixafor), AMD3465, ATI 2341, BKT140 (BL-8040; TF14016; 4F-Benzoyl-TN14003), CTCE-9908, CX549,,D-[Lys3] GHRP-6, FC122, FC131, GMI-1359, GSK812397, GST-NT21MP, isothioureia-1a, isothioureia-1t (IT1t), KRH-1636, KRH-3955, LY2510924, LY2624587, MSX-122, N-[¹⁴C]Methyl-AMD3465, PF-06747143, POL6326, SDF-1 1-9[P2Gdimer, SDF1 P2G, T134, T140, T22, TC 14012, TG-0054 (Burixafor), USL311, ulocuplumab (MDX1338/BMS-936564), viral macrophage inflammatory protein-II (vMIP-II), WZ811, 12G5, 238D2, 238D4, [⁶⁴Cu]-AMD3100, [⁶⁴Cu]-AMD3465, [⁶⁸Ga]pentixafor, [⁹⁰Y]pentixather, [^{99m}Tc]O₂-AMD3100, [¹⁷⁷Lu]pentixather, and 508MCl (Compound 26).
- [706] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a therapeutically effective amount of the GPCR_x inhibitor is administered to the patient.
- [707] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a sub-therapeutically effective amount of the GPCR_x inhibitor is administered to the patient.
- [708] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is selected from the group consisting of:
- [709] ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.
- [710] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is ADRB2 or HRH1.
- [711] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is ADCYAP1R1.
- [712] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the ADCYAP1R1 inhibitor is selected from the group consisting of:

- [713] M65, Max.d.4, MK-0893, N-stearyl-[Nle¹⁷]neurotensin-(6-11)/VIP-(7-28), PACAP-(6-38), and PG 97-269.
- [714] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is ADORA2B.
- [715] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the ADORA2B inhibitor is selected from the group consisting of:
- [716] 3-isobutyl-8-pyrrolidinoxanthine, alloxazine, AS16, AS70, AS74, AS94, AS95, AS96, AS99, AS100, AS101, ATL802, BW-A1433, caffeine, CGS 15943, CPX, CSC, CVT-6883, DAX, DEPX, derenofylline, DPCPX, FK-453, I-ABOPX, istradefylline, KF26777, LAS38096, LUF5981, MRE 2029F20, MRE 3008F20, MRS1191, MRS1220, MRS1523, MRS1706, MRS1754, MSX-2, OSIP339391, pentoxifylline, preladenant, PSB-10, PSB-11, PSB36, PSB603, PSB-0788, PSB1115, rolodylline, SCH 58261, SCH442416, ST-1535, theophylline, tonapodylline, vipadenant, xanthine amine congener, XCC, and ZM-241385.
- [717] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is ADORA3.
- [718] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the ADORA3 inhibitor is selected from the group consisting of:
- [719] ATL802, BW-A1433, caffeine, CGS 15943, CSC, CVT-6883, derenofylline, dexniguldipine, DPCPX, FK-453, flavanone, flavone, galangin, I-ABOPX, istradefylline, KF26777, LAS38096, LUF5981, MRE 2029F20, MRE 3008F20, MRE 3010F20, MRS1041, MRS1042, MRS1067, MRS1088, MRS1093, MRS1097, MRS1177, MRS1186, MRS1191, MRS1191, MRS1220, MRS1476, MRS1486, MRS1505, MRS1523, MRS1754, MRS928, MSX-2, nicardipine, preladenant, PSB-10, PSB-11, PSB36, PSB603, PSB1115, rolodylline, sakuranetin, SCH 58261, SCH442416, ST-1535, theophylline, tonapodylline, vipadenant, visnagin, VUF5574, VUF8504, VUF8507, xanthine amine congener, and ZM-241385.
- [720] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is ADRB2.
- [721] In a further embodiment, the method of treating, method of suppressing, pharma-

ceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the ADRB2 inhibitor is selected from the group consisting of:

- [722] Alprenolol, atenolol, betaxolol, bupranolol, butoxamine, carazolol, carvedilol, CGP 12177, cicloprolol, ICI 118551, ICYP, labetalol, levobetaxolol, levobunolol, LK 204-545, metoprolol, nadolol, NIHP, NIP, propafenone, propranolol, sotalol, SR59230A, and timolol.
- [723] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is C5AR1.
- [724] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the C5AR1 inhibitor is selected from the group consisting of:
- [725] A8^{Δ71-73}, AcPhe-Orn-Pro-D-Cha-Trp-Arg, avacopan, C089, CHIPS, DF2593A, JPE1375, L-156,602, NDT9520492, *N*-methyl-Phe-Lys-Pro-D-Cha-Trp-D-Arg-CO₂H, PMX205, PMX53, RPR121154, and W54011.
- [726] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is CALCR.
- [727] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CALCR inhibitor is selected from the group consisting of:
- [728] α -CGRP-(8-37) (human), AC187, CT-(8-32) (salmon), and olcegepant.
- [729] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is CHRM1.
- [730] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CHRM1 inhibitor is selected from the group consisting of:
- [731] 3-Quinuclidinyl benzilate (QNB), 4-DAMP, aclidinium, AE9C90CB, AFDX384, amitriptyline, AQ-RA 741, atropine, benztropine, biperiden, darifenacin, dicyclomine, dosulepin, ethopropazine, glycopyrrolate, guanylpirenzepine, hexahydrodifenidol, hexahydrosiladifenidol, hexocyclium, himbacine, ipratropium, lithocholylcholine, methoctramine, ML381, muscarinic toxin 1, muscarinic toxin 2, muscarinic toxin 3, *N*-methyl scopolamine, otenzepad, oxybutynin, *p*-F-HHSiD, pirenzepine, propantheline,

(R,R)-quinuclidinyl-4-fluoromethyl-benzilate, scopolamine, silahexocyclium, so-lifenacin, telenzepine, tiotropium, tolterodine, trihexyphenidyl, tripitramine, UH-AH 37, umeclidinium, and VU0255035.

- [732] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is EDNRB.
- [733] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the EDNRB inhibitor is selected from the group consisting of:
- [734] A192621, ambrisentan, atrasentan, bosentan (RO 470203, Tracleer); BQ788, IRL 2500, K-8794, macitentan, RES7011, Ro 46-8443, SB209670, SB217242 (enrasentan), TAK 044, and tezosentan (RO610612).
- [735] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is HRH1.
- [736] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the HRH1 inhibitor is selected from the group consisting of:
- [737] (-)-chlorpheniramine, (+)-chlorpheniramine, (-)-*trans*-H₂-PAT, (+)-*cis*-H₂-PAT, (+)-*trans*-H₂-PAT, (±)-*cis*-H₂-PAT, (±)-*trans*-H₂-PAT, (R)-cetirizine, (S)-cetirizine, 9-OH-risperidone, A-317920, A-349821, ABT-239, alimemazine, amitriptyline, aripiprazole, arpromidine, asenapine, astemizole, AZD3778, azelastine, BU-E 47, cetirizine, chlorpheniramine, chlorpromazine, ciproxifan, clemastine, clobenpropit, clozapine, conessine, cyclizine, cyproheptadine, desloratadine, diphenhydramine, dosulepin, doxepin, epinastine, fexofenadine, fluphenazine, fluspirilene, haloperidol, hydroxyzine, impromidine, INCB-38579, JNJ-39758979, ketotifen, loratadine, loxapine, MK-0249, molindone, olanzapine, perphenazine, pimozide, pipamperone, pitolisant, promethazine, pyrillamine, quetiapine, risperidone, sertindole, terfenadine, thioridazine, thiothixene, trifluoperazine, tripeleminamine, triprolidine, ziprasidone, and zotepine.
- [738] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is MLNR.
- [739] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the MLNR inhibitor is

selected from the group consisting of:

[740] GM-109, MA-2029, and OHM-11526.

[741] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is NTSR1.

[742] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the NTSR1 inhibitor is selected from the group consisting of:

[743] Meclinetant, SR48527, SR48692, and SR142948A.

[744] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR_x is TACR3.

[745] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the TACR3 inhibitor is selected from the group consisting of:

[746] [Trp⁷, β-Ala⁸]neurokinin A-(4-10), AZD2624, FK 224, GR138676, GSK 172981, GSK 256471, N',2-diphenylquinoline-4-carbohydrazide 8m, N',2-diphenylquinoline-4-carbohydrazide, osanetant, PD 154740, PD 161182, PD157672, saredutant, SB 218795, SB 222200, SB 235375, SCH 206272, SSR 146977, and talnetant.

[747] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the respective selective GPCR_x agonist is natural ligand of the respective GPCR_x., respectively.

[748] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method further comprises detecting the CXCR4-GPCR_x heteromer in the cancer patient.

[749] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method further comprises identifying the CXCR4-GPCR_x heteromer in the cancer patient.

[750] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method further comprises:

- [751] i) obtaining or having obtained a biological sample from the cancer patient;
- [752] ii) conducting or having conducted a diagnostic assay to determine presence, identity, or presence and identity, of a CXCR4-GPCR $\times$ heteromer in the obtained biological sample from the cancer patient; and
- [753] iii) selecting the inhibitor or combination of inhibitors to suppress enhanced downstream signaling from the CXCR4-GPCR $\times$ heteromer.
- [754] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the cancer is selected from the group consisting of: breast cancer, lung cancer, brain cancer, kidney cancer, pancreatic cancer, ovarian cancer, prostate cancer, melanoma, multiple myeloma, gastrointestinal cancers, renal cell carcinoma, soft tissue sarcomas, hepatocellular carcinoma, stomach cancer, colorectal cancer, esophageal cancer, and leukemia.
- [755] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the patient's biological sample is a biological fluid sample.
- [756] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a liquid biopsy is performed on the biological fluid sample. In some embodiments, the biological fluid sample includes circulating tumor cells (CTCs), tumor-derived cell-free DNA (cfDNA), circulating small RNAs, and extracellular vesicles including exosomes, from bodily fluids as disclosed, for example, in Campos CDM et al., "Molecular Profiling of Liquid Biopsy Samples for Precision Medicine," *Cancer J.* 2018 Mar/Apr;24(2):93-103, which is incorporated hereby in its entirety.
- [757] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the biological fluid sample is a blood sample, a plasma sample, a saliva sample, a cerebral fluid sample, an eye fluid sample, or a urine sample.
- [758] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the patient's biological sample is a biological tissue sample.
- [759] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a liquid biopsy is

performed on the biological tissue sample.

- [760] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the biological tissue sample is an organ tissue sample, a bone tissue sample, or a tumor tissue sample.
- [761] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the cancer cell contains the CXCR4-GPCR α heteromer.
- [762] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein a normal, non-cancerous cell, does not contain the CXCR4-GPCR α heteromer.
- [763] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the cancer cell of the patient contains the CXCR4-GPCR α heteromer in a greater concentration than a normal, non-cancerous cell from said patient, for example, 10% greater concentration than a normal, non-cancerous cell from said patient, such as 25% greater, 50% greater, 100% greater, 200% greater, or 300% greater, than a normal, non-cancerous cell from said patient.
- [764] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR α heteromer contained in the cancer cell of the patient is in a greater concentration than a normal, non-cancerous cell from said patient, for example, 10% greater concentration than a normal, non-cancerous cell from said patient, such as 25% greater, 50% greater, 100% greater, 200% greater, or 300% greater, than a normal, non-cancerous cell from said patient; and wherein the GPCR α of the CXCR4-GPCR α heteromer contained in the cancer cell of the patient is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.
- [765] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the presence of the CXCR4-GPCR α heteromer in the cancer cell of the patient identifies a sub-population of CXCR4-mediated cancer patients.
- [766] In a further embodiment, the method of treating, method of suppressing, pharma-

ceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the presence of the CXCR4-GPCR_x heteromer in the cancer cell of the patient is a biomarker of a sub-population of CXCR4-mediated cancer patients.

[767] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the biomarker of the sub-population of CXCR4-mediated cancer patients allows for precision medicine, patient stratification, or patient classification.

[768] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the biomarker of the sub-population of CXCR4-mediated cancer patients allows for GPCR-based precision cancer therapeutics.

[769] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the presence of the CXCR4-GPCR_x heteromer in the cancer cell of the patient identifies a sub-population of CXCR4-mediated cancer patients; and wherein the GPCR_x of the CXCR4-GPCR_x heteromer present in the cancer cell of the patient is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.

[770] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the presence of the CXCR4-GPCR_x heteromer in the cancer cell of the patient is a biomarker of a sub-population of CXCR4-mediated cancer patients; and wherein the GPCR_x of the CXCR4-GPCR_x heteromer present in the cancer cell of the patient is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.

[771] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the inhibitor is an antibody.

[772] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the CXCR4-GPCR_x heteromer-selective reagent is an antibody.

- [773] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the antibody is a bi-specific antibody of the CXCR4-GPCR α heteromer.
- [774] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the antibody is a heteromer-specific antibody of the CXCR4-GPCR α heteromer.
- [775] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the inhibitor is a bi-specific ligand(s) of the CXCR4-GPCR α heteromer.
- [776] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the antibody is an antibody-drug conjugate (ADC), as disclosed in, for example, Beck et al., "Strategies and challenges for the next generation of antibody-drug conjugates", *Nature Reviews Drug Discovery*, 16:315-337, (2017) and Lambert, et al., "Antibody-Drug Conjugates for Cancer Treatment", *Annual Review of Medicine*, 69:191-207 (2018), each of which are incorporated hereby in its entirety.
- [777] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method comprises: administering to the patient an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; wherein: i) the CXCR4-GPCR α heteromer has enhanced downstream signaling; and ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the cancer patient.
- [778] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the pharmaceutical kit or pharmaceutical composition, such as for use in treating cancer in a patient having a cell containing a CXCR4-GPCR α heteromer, comprises: an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; wherein the CXCR4-GPCR α heteromer has enhanced downstream signaling.
- [779] In a further embodiment, the method of treating, method of suppressing, pharma-

ceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR α heteromer is decreased in the range of 5-100% more, 10-100% more, 20-100% more, 30-100% more, 40-100% more, 50-100% more, 60-100% more, 75-100% more, 5-75% more, 5-50% more, or 5-25% more, upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient.

[780] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the efficacy of a CXCR4 inhibitor is increased in the range of 5-2000%, 5-1750%, 5-1500%, 5-1250%, 5-1000%, 5-900%, 5-800%, 5-700%, 5-500%, 5-400%, 5-250%, 5-200%, 5-100%, 5-75%, 5-50%, 5-40%, 5-30%, 5-25%, 100-2000%, 200-2000%, 300-2000%, 500-2000%, 750-2000%, 1000-2000%, 1250-2000%, 1500-2000%, 5-1500%, 25-1500%, 50-1500%, 75-1500%, 100-1500%, 200-1500%, 300-1500%, 500-1500%, 750-1500%, 1000-1500%, or 1250-1500%, when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor.

[781] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the efficacy of a GPCR α inhibitor is increased in the range of 5-2000%, 5-1750%, 5-1500%, 5-1250%, 5-1000%, 5-900%, 5-800%, 5-700%, 5-500%, 5-400%, 5-250%, 5-200%, 5-100%, 5-75%, 5-50%, 5-40%, 5-30%, 5-25%, 100-2000%, 200-2000%, 300-2000%, 500-2000%, 750-2000%, 1000-2000%, 1250-2000%, 1500-2000%, 5-1500%, 25-1500%, 50-1500%, 75-1500%, 100-1500%, 200-1500%, 300-1500%, 500-1500%, 750-1500%, 1000-1500%, or 1250-1500%, when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor.

[782] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method administers a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; or the pharmaceutical kit or pharmaceutical composition comprises a combination of in-

hibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer.

[783] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the combination of inhibitors is a combination of two inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer.

[784] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method administers a combination of a CXCR4 inhibitor and a GPCR_x inhibitor; or the pharmaceutical kit or pharmaceutical composition comprises a combination of a CXCR4 inhibitor and a GPCR_x inhibitor.

[785] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the method administers a CXCR4-GPCR_x heteromer inhibitor; or the pharmaceutical kit or pharmaceutical composition comprises a CXCR4-GPCR_x heteromer inhibitor.

[786] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the administering of the combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient in the range of between 5-2000 fold, 5-1750 fold, 5-1500 fold, 5-1250 fold, 5-1000 fold, 5-900 fold, 5-800 fold, 5-700 fold, 5-500 fold, 5-400 fold, 5-250 fold, 5-200 fold, 5-100 fold, 5-75 fold, 5-50 fold, 5-40 fold, 5-30 fold, 5-25 fold, 100-2000 fold, 200-2000 fold, 300-2000 fold, 500-2000 fold, 750-2000 fold, 1000-2000 fold, 1250-2000 fold, 1500-2000 fold, 5-1500 fold, 25-1500 fold, 50-1500 fold, 75-1500 fold, 100-1500 fold, 200-1500 fold, 300-1500 fold, 500-1500 fold, 750-1500 fold, 1000-1500 fold, or 1250-1500 fold, relative to single inhibitor administration.

[787] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the administering of the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient in the range of between 5-2000 fold, 5-1750 fold, 5-1500 fold, 5-1250 fold, 5-1000 fold, 5-900 fold, 5-800 fold, 5-700 fold, 5-500 fold, 5-400 fold, 5-250 fold, 5-200 fold, 5-100 fold, 5-75 fold,

5-50 fold, 5-40 fold, 5-30 fold, 5-25 fold, 100-2000 fold, 200-2000 fold, 300-2000 fold, 500-2000 fold, 750-2000 fold, 1000-2000 fold, 1250-2000 fold, 1500-2000 fold, 5-1500 fold, 25-1500 fold, 50-1500 fold, 75-1500 fold, 100-1500 fold, 200-1500 fold, 300-1500 fold, 500-1500 fold, 750-1500 fold, 1000-1500 fold, or 1250-1500 fold, relative to suppression of downstream signaling from either a CXCR4 protomer or a GPCR α protomer in their respective individual protomer context.

[788] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; for example, selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, CHRM1, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADORA2B, ADORA3, ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, NTSR1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, CHRM1, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, HRH1, and TACR3; selected from the group consisting of: ADRB2, EDNRB, and HRH1; selected from the group consisting of: ADRB2, CHRM1, and HRH1; or selected from the group consisting of: ADRB2, HRH1, and TACR3.

[789] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the GPCR α inhibitor is selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CALCR inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; for example, selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, C5AR1 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor,

CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, MLNR inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADCYAP1R1 inhibitor, ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADORA2B inhibitor, ADORA3 inhibitor, ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, NTSR1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, HRH1 inhibitor, and TACR3 inhibitor; selected from the group consisting of: ADRB2 inhibitor, EDNRB inhibitor, and HRH1 inhibitor; selected from the group consisting of: ADRB2 inhibitor, CHRM1 inhibitor, and HRH1 inhibitor; or selected from the group consisting of: ADRB2 inhibitor, HRH1, inhibitor and TACR3 inhibitor.

[790] In a further embodiment, the method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of the above embodiments and any one or more of the further embodiments herein, wherein the initial concentration of the inhibitor is in the range of between 1-10 μ M (or each of the inhibitors of the combination at concentrations in the range of between 1-10 μ M), such as at a concentration of 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 μ M, in testing and/or evaluating said inhibitor, or combination of inhibitors, regarding whether effective, or therapeutically effective, in suppressing an enhanced downstream signaling from a CXCR4-GPCR $\times$ heteromer.

[791]

[792] **Materials and Methods**

[793] **Reagents**

[794] BAY 60-6583, CGS 21680, C5a, acetylcholine, motilin, neurotensin, senktide, salmeterol, CI-IB-MECA, salmon calcitonin, BQ-3020, octreotide, VU0255035, cetirizine, pyrilamine, MA-2029, meclizine, and SSR 146977 were purchased from Tocris Bioscience (Ellisville, MO, USA). Galanin, endothelin-1, histamine, Prostaglandin E2 (PGE2), SRIF-14 (somatostatin), and bethanechol were purchased from Sigma-Aldrich (St Louis, Mo., USA). CXCL12, vasoactive intestinal peptide (VIP), and CCL2 were purchased from R&D systems (Minneapolis, Minnesota, USA). Formoterol, carvedilol, oxybutynin, bosentan, hydroxyzine, and loratadine were from

Prestwick Chemical (Illkirch, France). Apelin-13, roxithromycin, AMD3100, umeclidinium were obtained from Cayman Chemical Company (Ann Arbor, MI, USA), Selleckchem (Huston, TX, USA), Medchem express (Princeton, NJ, USA), and Santa Cruz Biotechnology (Santa Cruz, CA, USA), respectively.

[795]

[796] **Construction of adenoviral vectors containing GPCR cDNAs.**

[797] Vectors containing BiFC fragments, pCS2+VNm10 and pBiFC-VC155, were obtained from Chang-Deng Hu (Hu et al., 2002) and James Smith (Saka et al., 2007), respectively. The VNm10 is a Venus VN154 variant containing L46F and L64F mutations. To construct pAdBiFC-VN and pAdBiFC-VC vectors, DNAs containing VNm10 and VC155 were amplified by PCR and cloned into pShuttle-CMV. pAdBiFC-VN and pAdBiFC-VC vectors were obtained through homologous recombination between pAdEasy-1 and either pShuttle-CMV-VNm10 or pShuttle-CMV-VC155 using AdEasy vector system according to the manufacturer's instructions (Qbiogene, Carlsbad, CA). Human GPCR cDNA clones were obtained from the Missouri S&T cDNA Resource Center (Rolla, MO, USA). GPCR cDNAs were amplified by PCR and cloned into pDONR201 vector. Adenoviruses encoding GPCR, GPCR-VN, GPCR-VC, and GPCR-EGFP were obtained through *in vitro* LR recombination between entry clones containing GPCR and either pAdHTS, pAdBiFC-VN, pAdBiFC-VC, or pAdHTS-GFPC vectors, and subsequent transfection into 293A cells using AdHTS system as described previously (Choi et al., 2012; Song et al., 2014).

[798]

[799] **Cell culture**

[800] U-2 OS cells and MDA-MB-231 cells were purchased from the American Type Culture Collection (Manassas, VA, USA), and 293A cells were from Invitrogen (Carlsbad, CA, USA). U-2 OS cells, MDA-MB-231 cells, and 293A cells were grown in McCoy's 5A medium, RPMI 1640, and Modified Eagle's medium, respectively, in the presence of 10% fetal bovine serum (FBS), 100 units/ml of penicillin, and 100 µg/ml of streptomycin. Cells were cultured with 5% CO₂ at 37 °C.

[801]

[802] **BiFC assay**

[803] U-2 OS cells were seeded at a density of 3000 cells per well in a black 96-well clear-bottom plate in 100 µl of McCoy's 5A medium supplemented with 10% FBS. On the following day, the cells were transduced with 30 MOI each of adenoviruses encoding GPCR-VN and GPCR-VC. Three days post-transduction, cells were fixed with 2 % formaldehyde and nuclei were stained with Hoechst 33342 (Invitrogen, Carlsbad, CA). Images were acquired using IN cell Analyzer 1000 and analyzed with IN Cell Developer ToolBox (GE Healthcare, Waukesha, WI). BiFC and nuclear images were

visualized using a $\times 20$ objective and 360-nm (Hoechst) and 480-nm excitation filters, and monitored through 460- and 535-nm emission filters, respectively, with a 61002 trichroic mirror.

[804]

[805] **Co-internalization assay**

[806] U-2 OS cells were seeded at a density of 3000 cells per well in a black 96-well clear-bottom plate in 100 μ l of McCoy's 5A medium supplemented with 10% FBS. On the following day, the cells were co-transduced with adenoviruses encoding CXCR4-EGFP (10 MOI) and GPCR-VC or GPCR-VN (30 MOI). Two days post-transduction, cells were stimulated with GPCR α agonists for 30 min and fixed with 2% formaldehyde. Images were obtained using IN cell analyzer 2000 using an excitation wavelength of 480 nm and an emission wavelength of 535 nm.

[807]

[808] **Calcium mobilization assay**

[809] MDA-MB-231 human breast cancer cells were seeded at 20,000 cells per well in a black clear bottom 96-well plate (Corning Costar, #3340) in 100 μ l of RPMI 1640 supplemented with 10% FBS. The next day, cells were co-transduced with 10 MOI of CXCR4 and 30 MOI of HA-VC, 10 MOI of HA-VC and 30 MOI of GPCR α , or 10 MOI of CXCR4 and 30 MOI of GPCR α . Adenoviruses encoding HA-VC were used to adjust the total amount of adenoviruses transduced. After 2 days, cells were washed twice with assay buffer (Hank's balanced salt solution without phenol red, supplemented with 0.1% BSA and 20 mM HEPES, pH7.4), and stained with 5 μ g/ml of Cal-520 AM (AAT Bioquest, Sunnyvale, CA, USA) diluted in assay buffer for 2 hr. Cells were washed with assay buffer three times, and incubated at 37 °C for another 30 min. Plate was loaded on a FlexStation 3 Multi-Mode Microplate Reader (Molecular Devices, Sunnyvale, CA, USA) and GPCR α agonists were added. Intracellular Ca²⁺ mobilization was measured at 37 °C using an excitation wavelength of 490 nm and an emission wavelength of 525 nm. Antagonists or vehicles were added 30 min before agonist treatment.

[810]

[811] **Internalization inhibition assay**

[812] CXCR4-GFP expressed U-2 OS cells were seeded at a density of 5000 cells per well in a black 96-well clear-bottom plate in 100 μ l of McCoy's 5A medium supplemented with 10% FBS. On the following day, the cells were transduced with adenoviruses encoding GPCR α (30 MOI). Two days post-transduction, cells were stimulated with SDF-1 and/or GPCR α agonists for 20 min and fixed with 4% paraformaldehyde. Images were obtained using IN cell analyzer 2500 using an excitation wavelength of 480 nm and an emission wavelength of 535 nm.

[813]

[814] **Proliferation assay**

[815] PDCs were seeded in 384-well plate at 500 ea /well in 40 ul culture media. After overnight growth, the cells were cultured for 7 days in the presence of several dose of GPCR α antagonist or DMSO alone. After the 7-day incubation, 15 ul ATPlite (PerkinElmer, Cat. No. 6016739) was added into the each well and the plates were shaken for 5 minutes in an orbital shaker at 700 rpm. The luminescent signal was detected within 30 minutes at PerkinElmer TopCount detection instrument. The cell viability was calculated using the equation: Cell viability (%) = (OD of antagonist treatment/OD of DMSO only treatment) x 100%.

[816]

[817] **Proximity ligation assay in cell**

[818] CXCR4 overexpression cell lines, the U2OS-CXCR4 was infected with ADRB2 expressing adenovirus, Ad-ADRB2 at the dose of 0, 2.5, 10, 40 MOIs for 2 days. Proximity ligation assays (PLA) were performed as described previously (Brueggemann et al., 2014; Tripathi et al., 2014). To perform PLA, infected cells were fixed with 4% paraformaldehyde (PFA) on sixteen-well tissue culture slides. Slides were blocked with blocking solution provided by Duolink and incubated with mouse anti-CXCR4 (1:200, Santacruz, Sc-53534), Rabbit anti-ADRB2 (1:200, Thermo-scientific, PA5-33333) at 37°C for 1 h in a humidifying chamber. Slides were then washed and incubated (1 h at 37°C) with secondary anti-rabbit and anti-mouse antibodies conjugated with plus and minus Duolink II PLA probes. Slides were washed again and then incubated with ligation-ligase solution (30 min at 37°C) followed by incubation with amplification-polymerase solution (2 h at 37°C). Slides were then mounted with minimal volume of Duolink II mounting medium with 4',6-diamidino-2phenylindole (DAPI) for 15-30 min, and PLA signals [Duolink In Situ Detection Reagents Green (λ excitation/emission 495/527 nm) or Red (λ excitation/emission 575/623 nm)] were identified as fluorescent spots under a IN Cell analyzer 2500.

[819]

[820] **Proximity ligation assay in PDC**

[821] PDCs were seeded in 96-chamber slide at 100000 ea /well in 100 ul culture media. To perform PLA, PDCs were fixed with 4% paraformaldehyde (PFA) and were blocked with blocking solution provided by Duolink and incubated with mouse anti-CXCR4 (1:200, Santacruz, Sc-53534), Rabbit anti-ADRB2 (1:200, Thermo-scientific, PA5-33333), Rabbit anti-CHRM1 (1:200, Ls bio, Ls-C313301) at 37°C for 1 h in a humidifying chamber. Slides were then washed and incubated (1 h at 37°C) with secondary anti-rabbit and anti-mouse antibodies conjugated with plus and minus

Duolink II PLA probes. Slides were washed again and then incubated with ligation-ligase solution (30 min at 37°C) followed by incubation with amplification-polymerase solution (2 h at 37°C). Slides were then mounted with minimal volume of Duolink II mounting medium with 4',6-diamidino-2phenylindole (DAPI) for 15-30 min, and PLA signals [Duolink In Situ Detection Reagents Green (λ excitation/emission 495/527 nm) or Red (λ excitation/emission 575/623 nm)] were identified as fluorescent spots under a IN Cell analyzer 2500.

[822]

[823] **Proximity ligation assay in PDX**

[824] To perform PLA with PDX samples, the glioblastoma patient derived FFPE samples were used (provided by Samsung Seoul hospital in Seoul, Korea). After FFPE sample were de-paraffinized and performed heat induced antigen retrieval for 15 minutes at 100°C. Slides were blocked with blocking solution provided by Duolink and incubated with rabbit anti-CXCR4 (1:200, Thermoscientific, PA3305), mouse anti-ADRB2 (1:200, Santacruz, Sc-271322), at 37 °C for 1 h in a humidifying chamber. The other process was same as described above (PLA with PDC).

[825]

[826] **Chemotactic Migration Assay**

[827] Transwell plates (8 μ m pore size polycarbonate membrane, 6.5-mm diameter) were coated with 50 μ g/mL collagen for 2 h at 37 °C (Corning Inc.). MDA-MB-231 cells were serum-starved 24 h and then plated in serum-free medium containing 0.5% BSA in the top chamber (20,000 cells/100 μ L). Antagonists or inverse agonists were added to the cells 30 min before plating the cells in the transwell plates. Attractants were added in the bottom chamber. Antagonists or inverse agonists were also included in the bottom chamber. After 3 h at 37 °C, cells on the top transwell membrane were removed using a cotton swab, fixed, and stained with 0.1% Crystal violet. Chemotaxis was quantified by counting the migrated cells on the lower surface of the membrane of 10 fields per chamber at 10 $\times$ objective.

[828]

[829] **Reverse Transcription-quantitative Polymerase Chain Reaction (RT-qPCR)**

[830] Total RNA was extracted using TRIzol (Invitrogen) and cDNA was synthesized from 1 μ g of total RNA after treatment with DNase I (Sigma). RT-qPCR was conducted using SensiFAST SYBR kit (Bioline). Primer sequences are as follows:

[831] ADRB2-F: 5'-CTCTTCCATCGTGTCTCTCTAC-3' (SEQ ID NO:1),

[832] ADRB2-R: 5'-AATCTTCTGGAGCTGCCTTT-3' (SEQ ID NO:2);

[833] HRH1-F: 5'- CCTCTGCTGGATCCCTTATTTC-3' (SEQ ID NO:3),

[834] HRH1-R: 5'-GGTTCAGTGTGGAGTTGATGTA-3' (SEQ ID NO:4);

[835] CXCR4-F: 5'- CCACCATCTACTCCATCATCTTC-3' (SEQ ID NO:5),

- [836] CXCR4-R: 5'-ACTTGTCGTCATGCTTCTC-3' (SEQ ID NO:6);
 [837] β -actin-F: 5'-GGAAATCGTGCGTGACATTAAG-3' (SEQ ID NO:7),
 [838] β -actin-R: 5'-AGCTCGTAGCTCTTCTCCA-3' (SEQ ID NO:8); GAPDH-F:
 [839] 5'-ATGACATCAAGAAGGTGGTGAA-3' (SEQ ID NO:9), GAPDH-R:
 [840] 5'-GCTGTTGAAGTCAGAGGAGAC-3' (SEQ ID NO:10).
 [841] Threshold cycles (Ct) were calculated by QuantStudio 3 Real-Time PCR system (Thermo Fisher Scientific).

[842]

[843] **Lentivirus production**

- [844] pLenti CMV Hygro DEST (w117-1) was a gift from Eric Campeau & Paul Kaufman (Addgene plasmid # 17454) (Campeau et al., 2009). CXCR4 cDNA was inserted into the lentiviral vector using LR recombination. Lentiviruses encoding CXCR4 were produced using ViraPower Lentiviral Expression Systems (Invitrogen).

[845]

[846] **Construction of Stable Cell Lines Expressing CXCR4 alone or CXCR4 and ADRB2 heteromer**

- [847] To establish stable CXCR4 expressing cell lines, lentiviral stock (containing the packaged pLenti6-CXCR4 expression construct which inserted with CXCR4 gene in pLenti-CMV Hygro DEST (Addgene, #17454)), were produced by co-transfecting the ViraPower Lentiviral Packaging Mix (Invitrogen, K497500) and pLenti6-CXCR4 expression construct into 293FT producer cell line. Transduction of this lentiviral stock into A549 cell line was performed and followed by selection with hygromycin (100 μ g/mL) and blasticidin (5 μ g/mL). To establish stable CXCR4-ADRB2 heteromer expressing cell lines, lentiviral stock (containing the packaged pLenti6/V5-ADRB2 expression construct which inserted with ADRB2 gene in pLenti6/V5-DEST Gateway™ Vector (Invitrogen, V49610)), were produced by co-transfecting the ViraPower Packaging Mix and pLenti6/V5-ADRB2 expression construct into 293FT producer cell line. Transduction of this lentiviral stock into A549-CXCR4 cell line was performed and followed by selection with hygromycin (100 μ g/mL) and blasticidin (5 μ g/mL). Then clones resistant to antibiotics were selected and performed RT-qPCR and immunofluorescence to confirm the expression of the inserted gene, CXCR4 and ADRB2.

[848]

[849] **Mouse Xenograft Model**

- [850] Five week-old, female Balb/c-nu/nu mice were obtained from Envigo (France) and maintained in specific pathogen-free animal facility. All protocols for animal use and euthanasia were approved by the Qubest Bio (South Korea) Animal Experimental Ethics Committee based on the Animal Protection Act. The 1×10^7 cells of A549,

A549-CXCR4 or A549-CXCR4-ADRB2 were suspended in 100 μ L of phosphate-buffered saline (PBS) and were implanted by subcutaneously in axillary region between the clavicular and chest wall on the right side of the mouse. The tumor growth was monitored every third or fourth day by measuring the length (L) and width (W) of the tumor with an electronic caliper and calculating tumor volume in the basis of the following formula: Volume = $0.5 LW^2$.

[851]

[852] **Generation of CXCR4 or HRH1 Knockout Cells Using CRISPR/Cas9 System**

[853] CRISPR guide RNAs targeting CXCR4 and HRH1, and non-targeting guide RNA cloned in lentiCRISPR v2 vector were purchased from GenScript (Piscataway, NJ) (Sanjana et al., 2014). The guide RNA sequences are as follows:

[854] TGTTGGCTGCCTTACTACAT (CXCR4 gRNA #1; (SEQ ID NO:11)),

[855] CGATCAAGTCCGCCACCGAG (HRH1 gRNA #3; (SEQ ID NO:12)), and

[856] ACGGAGGCTAAGCGTCGCAA (non-targeting gRNA; (SEQ ID NO:13)).

Lentiviruses were produced using ViraPower Lentiviral Expression Systems (Invitrogen). MDA-MB-231 cells and MDA-MB-231 cells overexpressing CXCR4 (MDA^{CXCR4+}) were transduced with lentiviruses encoding non-targeting gRNA, CXCR4 gRNA, or HRH1 gRNAs. Cells were selected with puromycin (3 μ M) for 2 weeks and loss of CXCR4 and HRH1 was estimated with immunoblotting and calcium responses, respectively. For detecting CXCR4, anti-CXCR4 rabbit monoclonal antibody (Abcam, #ab124824) was used.

[857]

[858] **Source of GPCR α inhibitors**

[859] Ulocuplumab was purchased from Creative Biolabs (Shirley, NY, USA). BKT140 was purchased from Chem Scene (Monmouth Junction, NJ, USA). Carazolol was purchased from Santacruz Biotech (Dallas, TX, USA). Osanetant was purchased from Axon Medchem, (Groningen, Netherland). M65 and CT-(8-32) (salmon) were from Bachem (San Diego, CA, USA). PACAP-(6-38), W54011, PMX205, PMX53, AC187, SB 222200, and Talnetant were from Tocris Bioscience (Ellisville, MO, USA). Propranolol, Prometazine, Cyproheptadine, Hydroxyzine, Ambrisentan and macitentan were from Prestwick Chemical (Illkirch, France), and Selleckchem (Huston, TX, USA), respectively.

[860]

[861] ***In vivo* studies**[862] For the antitumor efficacy test using CXCR4 and ADRB2 inhibitor, A549-CXCR4-ADRB2 cell line overexpressing CXCR4 and ADRB2 in A549, lung cancer cell line was subcutaneously administered (1×10^7 cell in 100 μ L PBS / head) to BALB/c-nu. When Tumor size reached 50-100 mm³, groups of ten mice were admin-

istrated with vehicle (1% dimethylcellulose in PBS), CXCR4 inhibitor (3 mg/kg of LY2510924 formulated in PBS, 10 mg/kg of AMD070 formulated in DMSO), ADRB2 inhibitor (30 mg/kg of Carvedilol formulated in 1% dimethylcellulose in PBS) or combination of CXCR4 and ADRB2 inhibitor, AMD070 and Carvedilol, or LY2510924 and Carvedilol once a day for 21-23 days. LY2510924 was subcutaneously injected and AMD070 or Carvedilol was administrated orally. The tumor growth was monitored every third or fourth day by measuring the length(*L*) and width(*W*) of the tumor: Volume = $0.5 LW^2$.

[863]

[864] **References**

[865] Each of the following references, in their entirety, are incorporated herein by reference:

[866] Abd Alla, J., Reeck, K., Langer, A., Streichert, T., and Quitterer, U. (2009). Calcitriculin enhances B2 bradykinin receptor maturation and heterodimerization. *Biochem Biophys Res Commun* 387, 186-190;

[867] Agrawal, L., Lu, X., Qingwen, J., VanHorn-Ali, Z., Nicolescu, I.V., McDermott, D.H., Murphy, P.M., and Alkhatib, G. (2004). Role for CCR5Delta32 protein in resistance to R5, R5X4, and X4 human immunodeficiency virus type 1 in primary CD4+ cells. *J Virol* 78, 2277-2287;

[868] Albrandt, K., Brady, E.M., Moore, C.X., Mull, E., Sierzega, M.E., and Beaumont, K. (1995). Molecular cloning and functional expression of a third isoform of the human calcitonin receptor and partial characterization of the calcitonin receptor gene. *Endocrinology* 136, 5377-5384;

[869] Allen, C.D., Ansel, K.M., Low, C., Lesley, R., Tamamura, H., Fujii, N., and Cyster, J.G. (2004). Germinal center dark and light zone organization is mediated by CXCR4 and CXCR5. *Nat Immunol* 5, 943-952;

[870] Ara, T., Itoi, M., Kawabata, K., Egawa, T., Tokoyoda, K., Sugiyama, T., Fujii, N., Amagai, T., and Nagasawa, T. (2003). A role of CXC chemokine ligand 12/stromal cell-derived factor-1/pre-B cell growth stimulating factor and its receptor CXCR4 in fetal and adult T cell development *in vivo*. *J Immunol* 170, 4649-4655;

[871] Armando, S., Quoyer, J., Lukashova, V., Maiga, A., Percherancier, Y., Heveker, N., Pin, J.P., Prezeau, L., and Bouvier, M. (2014). The chemokine CXC4 and CC2 receptors form homo- and heterooligomers that can engage their signaling G-protein effectors and betaarrestin. *FASEB J* 28, 4509-4523;

[872] Barmania, F., and Pepper, M.S. (2013). C-C chemokine receptor type five (CCR5): An emerging target for the control of HIV infection. *Appl Transl Genom* 2, 3-16;

[873] Bartolome, R.A., Ferreira, S., Miquilena-Colina, M.E., Martinez-Prats, L., Soto-Montenegro, M.L., Garcia-Bernal, D., Vaquero, J.J., Agami, R., Delgado, R., Desco,

- M., *et al.* (2009). The chemokine receptor CXCR4 and the metalloproteinase MT1-MMP are mutually required during melanoma metastasis to lungs. *Am J Pathol* 174, 602-612;
- [874] Batlle, E., and Clevers, H. (2017). Cancer stem cells revisited. *Nat Med* 23, 1124-1134;
- [875] Bazin, H., Trinquet, E., and Mathis, G. (2002). Time resolved amplification of cryptate emission: a versatile technology to trace biomolecular interactions. *J Biotechnol* 82, 233-250;
- [876] Beck a et al., "Strategies and challenges for the next generation of antibody-drug conjugates", *Nature Reviews Drug Discovery*, 16:315-337, (2017);
- [877] Bjarnadottir, T.K., Gloriam, D.E., Hellstrand, S.H., Kristiansson, H., Fredriksson, R., and Schioth, H.B. (2006). Comprehensive repertoire and phylogenetic analysis of the G protein-coupled receptors in human and mouse. *Genomics* 88, 263-273;
- [878] Bodart, V., Anastassov, V., Darkes, M.C., Idzan, S.R., Labrecque, J., Lau, G., Mosi, R.M., Neff, K.S., Nelson, K.L., Ruzek, M.C., *et al.* (2009). Pharmacology of AMD3465: a small molecule antagonist of the chemokine receptor CXCR4. *Biochem Pharmacol* 78, 993-1000;
- [879] Braadland, P.R., Ramberg, H., Grytli, H.H., and Tasken, K.A. (2014). beta-Adrenergic Receptor Signaling in Prostate Cancer. *Front Oncol* 4, 375;
- [880] Broxmeyer, H.E., Orschell, C.M., Clapp, D.W., Hangoc, G., Cooper, S., Plett, P.A., Liles, W.C., Li, X., Graham-Evans, B., Campbell, T.B., *et al.* (2005). Rapid mobilization of murine and human hematopoietic stem and progenitor cells with AMD3100, a CXCR4 antagonist. *J Exp Med* 201, 1307-1318;
- [881] Brueggemann, L.I., Mackie, A.R., Cribbs, L.L., Freda, J., Tripathi, A., Majetschak, M., and Byron, K.L. (2014). Differential protein kinase C-dependent modulation of Kv7.4 and Kv7.5 subunits of vascular Kv7 channels. *J Biol Chem* 289, 2099-2111;
- [882] Buckley, C.D., Amft, N., Bradfield, P.F., Pilling, D., Ross, E., Arenzana-Seisdedos, F., Amara, A., Curnow, S.J., Lord, J.M., Scheel-Toellner, D., *et al.* (2000). Persistent induction of the chemokine receptor CXCR4 by TGF-beta 1 on synovial T cells contributes to their accumulation within the rheumatoid synovium. *J Immunol* 165, 3423-3429;
- [883] Burbassi, S., Sengupta, R., and Meucci, O. (2010). Alterations of CXCR4 function in mu-opioid receptor-deficient glia. *Eur J Neurosci* 32, 1278-1288;
- [884] Burger, J.A., Burger, M., and Kipps, T.J. (1999). Chronic lymphocytic leukemia B cells express functional CXCR4 chemokine receptors that mediate spontaneous migration beneath bone marrow stromal cells. *Blood* 94, 3658-3667;
- [885] Burger, J.A., and Kipps, T.J. (2006). CXCR4: a key receptor in the crosstalk between tumor cells and their microenvironment. *Blood* 107, 1761-1767;

- [886] Burger, J.A., and Peled, A. (2009). CXCR4 antagonists: targeting the microenvironment in leukemia and other cancers. *Leukemia* 23, 43-52;
- [887] Burger, J.A., Stewart, D.J., Wald, O., and Peled, A. (2011). Potential of CXCR4 antagonists for the treatment of metastatic lung cancer. *Expert Rev Anticancer Ther* 11, 621-630;
- [888] Bushlin, I., Gupta, A., Stockton, S.D., Jr., Miller, L.K., and Devi, L.A. (2012). Dimerization with cannabinoid receptors allosterically modulates delta opioid receptor activity during neuropathic pain. *PLoS One* 7, e49789;
- [889] Callen, L., Moreno, E., Barroso-Chinea, P., Moreno-Delgado, D., Cortes, A., Mallol, J., Casado, V., Lanciego, J.L., Franco, R., Lluís, C., *et al.* (2012). Cannabinoid receptors CB1 and CB2 form functional heteromers in brain. *J Biol Chem* 287, 20851-20865;
- [890] Campeau, E., Ruhl, V.E., Rodier, F., Smith, C.L., Rahmberg, B.L., Fuss, J.O., Campisi, J., Yaswen, P., Cooper, P.K., and Kaufman, P.D. (2009). A versatile viral system for expression and depletion of proteins in mammalian cells. *PLoS One* 4, e6529;
- [891] Canonica, G.W., and Blaiss, M. (2011). Antihistaminic, anti-inflammatory, and antiallergic properties of the nonsedating second-generation antihistamine desloratadine: a review of the evidence. *World Allergy Organ J* 4, 47-53;
- [892] Cao, H.L., Liu, Z.J., and Chang, Z. (2017). Cordycepin induces apoptosis in human bladder cancer cells via activation of A3 adenosine receptors. *Tumour Biol* 39, 1010428317706915;
- [893] Cashen, A.F., Nervi, B., and DiPersio, J. (2007). AMD3100: CXCR4 antagonist and rapid stem cell-mobilizing agent. *Future Oncol* 3, 19-27;
- [894] Chatterjee, S., Behnam Azad, B., and Nimmagadda, S. (2014). The intricate role of CXCR4 in cancer. *Adv Cancer Res* 124, 31-82;
- [895] Chen, J.F., Eltzschig, H.K., and Fredholm, B.B. (2013). Adenosine receptors as drug targets--what are the challenges? *Nat Rev Drug Discov* 12, 265-286;
- [896] Chen, Y., Ramjiawan, R.R., Reiberger, T., Ng, M.R., Hato, T., Huang, Y., Ochiai, H., Kitahara, S., Unan, E.C., Reddy, T.P., *et al.* (2015). CXCR4 inhibition in tumor microenvironment facilitates anti-programmed death receptor-1 immunotherapy in sorafenib-treated hepatocellular carcinoma in mice. *Hepatology* 61, 1591-1602;
- [897] Cheng, X., Wang, H., Zhang, X., Zhao, S., Zhou, Z., Mu, X., Zhao, C., and Teng, W. (2017). The Role of SDF-1/CXCR4/CXCR7 in Neuronal Regeneration after Cerebral Ischemia. *Front Neurosci* 11, 590;
- [898] Cho, K.S., Yoon, S.J., Lee, J.Y., Cho, N.H., Choi, Y.D., Song, Y.S., and Hong, S.J. (2013). Inhibition of tumor growth and histopathological changes following treatment with a chemokine receptor CXCR4 antagonist in a prostate cancer xenograft model.

- Oncol Lett 6, 933-938;
- [899] Choi, E.W., Seen, D.S., Song, Y.B., Son, H.S., Jung, N.C., Huh, W.K., Hahn, J.S., Kim, K., Jeong, J.Y., and Lee, T.G. (2012). AdHTS: a high-throughput system for generating recombinant adenoviruses. *J Biotechnol* 162, 246-252;
- [900] Chong, B.F., and Mohan, C. (2009). Targeting the CXCR4/CXCL12 axis in systemic lupus erythematosus. *Expert Opin Ther Targets* 13, 1147-1153;
- [901] Choy, C., Raytis, J.L., Smith, D.D., Duenas, M., Neman, J., Jandial, R., and Lew, M.W. (2016). Inhibition of beta2-adrenergic receptor reduces triple-negative breast cancer brain metastases: The potential benefit of perioperative beta-blockade. *Oncol Rep* 35, 3135-3142;
- [902] Chu, T., Shields, L.B.E., Zhang, Y.P., Feng, S.Q., Shields, C.B., and Cai, J. (2017). CXCL12/CXCR4/CXCR7 Chemokine Axis in the Central Nervous System: Therapeutic Targets for Remyelination in Demyelinating Diseases. *Neuroscientist* 23, 627-648;
- [903] Chung, S.H., Seki, K., Choi, B.I., Kimura, K.B., Ito, A., Fujikado, N., Saijo, S., and Iwakura, Y. (2010). CXC chemokine receptor 4 expressed in T cells plays an important role in the development of collagen-induced arthritis. *Arthritis Res Ther* 12, R188;
- [904] Comps-Agrar, L., Maurel, D., Rondard, P., Pin, J.P., Trinquet, E., and Prezeau, L. (2011). Cell-surface protein-protein interaction analysis with time-resolved FRET and snap-tag technologies: application to G protein-coupled receptor oligomerization. *Methods Mol Biol* 756, 201-214;
- [905] Cooper, A.J., Narasimhan, S., Rickels, K., and Lohoff, F.W. (2013). Genetic polymorphisms in the PACAP and PAC1 receptor genes and treatment response to venlafaxine XR in generalized anxiety disorder. *Psychiatry Res* 210, 1299-1300;
- [906] Covic, L., Gresser, A.L., Talavera, J., Swift, S., and Kuliopulos, A. (2002). Activation and inhibition of G protein-coupled receptors by cell-penetrating membrane-tethered peptides. *Proc Natl Acad Sci USA*, 99, 643-648;
- [907] Crawford, J.B., Chen, G., Gauthier, D., Wilson, T., Carpenter, B., Baird, I.R., McEachern, E., and Alan Kaller, C.H., Bem Atsma, Renato T. Skerlj, and Gary J. Bridger (2008). AMD070, a CXCR4 Chemokine Receptor Antagonist: Practical Large-Scale Laboratory Synthesis. *Organic Process Research & Development* 12, 823-830;
- [908] D'Alterio, C., Barbieri, A., Portella, L., Palma, G., Polimeno, M., Riccio, A., Ierano, C., Franco, R., Scognamiglio, G., Bryce, J., *et al.* (2012). Inhibition of stromal CXCR4 impairs development of lung metastases. *Cancer Immunol Immunother* 61, 1713-1720;
- [909] Dacquin, R., Davey, R.A., Laplace, C., Levasseur, R., Morris, H.A., Goldring, S.R., Gebre-Medhin, S., Galson, D.L., Zajac, J.D., and Karsenty, G. (2004). Amylin inhibits bone resorption while the calcitonin receptor controls bone formation *in vivo*. *J Cell*

- Biol 164, 509-514;
- [910] Davey, R.A., Turner, A.G., McManus, J.F., Chiu, W.S., Tjahyono, F., Moore, A.J., Atkins, G.J., Anderson, P.H., Ma, C., Glatt, V., *et al.* (2008). Calcitonin receptor plays a physiological role to protect against hypercalcemia in mice. *J Bone Miner Res* 23, 1182-1193;
- [911] De Clercq, E. (2003). The bicyclam AMD3100 story. *Nat Rev Drug Discov* 2, 581-587;
- [912] De Falco, V., Guarino, V., Avilla, E., Castellone, M.D., Salerno, P., Salvatore, G., Faviana, P., Basolo, F., Santoro, M., and Melillo, R.M. (2007). Biological role and potential therapeutic targeting of the chemokine receptor CXCR4 in undifferentiated thyroid cancer. *Cancer Res* 67, 11821-11829;
- [913] de Graaf, C., Kooistra, A.J., Vischer, H.F., Katritch, V., Kuijter, M., Shiroishi, M., Iwata, S., Shimamura, T., Stevens, R.C., de Esch, I.J., *et al.* (2011). Crystal structure-based virtual screening for fragment-like ligands of the human histamine H(1) receptor. *J Med Chem* 54, 8195-8206;
- [914] De Klerck, B., Geboes, L., Hatse, S., Kelchtermans, H., Meyvis, Y., Vermeire, K., Bridger, G., Billiau, A., Schols, D., and Matthys, P. (2005). Pro-inflammatory properties of stromal cell-derived factor-1 (CXCL12) in collagen-induced arthritis. *Arthritis Res Ther* 7, R1208-1220;
- [915] de Nigris, F., Schiano, C., Infante, T., and Napoli, C. (2012). CXCR4 inhibitors: tumor vasculature and therapeutic challenges. *Recent Pat Anticancer Drug Discov* 7, 251-264;
- [916] de Poorter, C., Baertsoen, K., Lannoy, V., Parmentier, M., and Springael, J.Y. (2013). Consequences of ChemR23 heteromerization with the chemokine receptors CXCR4 and CCR7. *PLoS One* 8, e58075;
- [917] Debnath, B., Xu, S., Grande, F., Garofalo, A., and Neamati, N. (2013). Small molecule inhibitors of CXCR4. *Theranostics* 3, 47-75;
- [918] Decaillot, F.M., Kazmi, M.A., Lin, Y., Ray-Saha, S., Sakmar, T.P., and Sachdev, P. (2011). CXCR7/CXCR4 heterodimer constitutively recruits beta-arrestin to enhance cell migration. *J Biol Chem* 286, 32188-32197;
- [919] Decaillot, F.M., Rozenfeld, R., Gupta, A., and Devi, L.A. (2008). Cell surface targeting of mu-delta opioid receptor heterodimers by RTP4. *Proc Natl Acad Sci U S A* 105, 16045-16050;
- [920] Demmer, O., Gourni, E., Schumacher, U., Kessler, H., and Wester, H.J. (2011). PET imaging of CXCR4 receptors in cancer by a new optimized ligand. *ChemMedChem* 6, 1789-1791;
- [921] Depoortere, I. (2001). Motilin and motilin receptors: characterization and functional significance. *Verh K Acad Geneesk Belg* 63, 511-529;

- [922] Desmet, C.J., Gallenne, T., Prieur, A., Rey, F., Visser, N.L., Wittner, B.S., Smit, M.A., Geiger, T.R., Laoukili, J., Iskit, S., *et al.* (2013). Identification of a pharmacologically tractable Fra-1/ADORA2B axis promoting breast cancer metastasis. *Proc Natl Acad Sci U S A* 110, 5139-5144;
- [923] DiPersio, J.F., Micallef, I.N., Stiff, P.J., Bolwell, B.J., Maziarz, R.T., Jacobsen, E., Nademanee, A., McCarty, J., Bridger, G., Calandra, G., *et al.* (2009a). Phase III prospective randomized double-blind placebo-controlled trial of plerixafor plus granulocyte colony-stimulating factor compared with placebo plus granulocyte colony-stimulating factor for autologous stem-cell mobilization and transplantation for patients with non-Hodgkin's lymphoma. *J Clin Oncol* 27, 4767-4773;
- [924] DiPersio, J.F., Stadtmauer, E.A., Nademanee, A., Micallef, I.N., Stiff, P.J., Kaufman, J.L., Maziarz, R.T., Hosing, C., Fruehauf, S., Horwitz, M., *et al.* (2009b). Plerixafor and G-CSF versus placebo and G-CSF to mobilize hematopoietic stem cells for autologous stem cell transplantation in patients with multiple myeloma. *Blood* 113, 5720-5726;
- [925] Domanska, U.M., Kruizinga, R.C., Nagengast, W.B., Timmer-Bosscha, H., Huls, G., de Vries, E.G., and Walenkamp, A.M. (2013). A review on CXCR4/CXCL12 axis in oncology: no place to hide. *Eur J Cancer* 49, 219-230;
- [926] Donzella, G.A., Schols, D., Lin, S.W., Este, J.A., Nagashima, K.A., Maddon, P.J., Allaway, G.P., Sakmar, T.P., Henson, G., De Clercq, E., *et al.* (1998). AMD3100, a small molecule inhibitor of HIV-1 entry via the CXCR4 co-receptor. *Nat Med* 4, 72-77;
- [927] Doranz, B.J., Fillion, L.G., Diaz-Mitoma, F., Sitar, D.S., Sahai, J., Baribaud, F., Orsini, M.J., Benovic, J.L., Cameron, W., and Doms, R.W. (2001). Safe use of the CXCR4 inhibitor ALX40-4C in humans. *AIDS Res Hum Retroviruses* 17, 475-486;
- [928] Doring, Y., Pawig, L., Weber, C., and Noels, H. (2014). The CXCL12/CXCR4 chemokine ligand/receptor axis in cardiovascular disease. *Front Physiol* 5, 212;
- [929] Doucette, L.P., and Walter, M.A. (2017). Prostaglandins in the eye: Function, expression, and roles in glaucoma. *Ophthalmic Genet* 38, 108-116;
- [930] Du, Y., Long, Q., Guan, B., and Mu, L. (2015). Prognostic Value of High CXCR4 Expression in Renal Cell Carcinoma: A System Review and Meta-Analysis. *Dis Markers* 2015, 568980;
- [931] Eidne, K.A., Kroeger, K.M., and Hanyaloglu, A.C. (2002). Applications of novel resonance energy transfer techniques to study dynamic hormone receptor interactions in living cells. *Trends Endocrinol Metab* 13, 415-421;
- [932] Endres, M.J., Clapham, P.R., Marsh, M., Ahuja, M., Turner, J.D., McKnight, A., Thomas, J.F., Stoeckenau-Haggarty, B., Choe, S., Vance, P.J., *et al.* (1996). CD4-independent infection by HIV-2 is mediated by fusin/CXCR4. *Cell* 87, 745-756;

- [933] Fagerberg, L., Hallstrom, B.M., Oksvold, P., Kampf, C., Djureinovic, D., Odeberg, J., Habuka, M., Tahmasebpoor, S., Danielsson, A., Edlund, K., *et al.* (2014). Analysis of the human tissue-specific expression by genome-wide integration of transcriptomics and antibody-based proteomics. *Mol Cell Proteomics* *13*, 397-406;
- [934] Fahham, D., Weiss, I.D., Abraham, M., Beider, K., Hanna, W., Shlomai, Z., Eizenberg, O., Zamir, G., Izhar, U., Shapira, O.M., *et al.* (2012). *In vitro* and *in vivo* therapeutic efficacy of CXCR4 antagonist BKT140 against human non-small cell lung cancer. *J Thorac Cardiovasc Surg* *144*, 1167-1175 e1161;
- [935] Farran, B. (2017). An update on the physiological and therapeutic relevance of GPCR oligomers. *Pharmacol Res* *117*, 303-327;
- [936] Feig, C., Jones, J.O., Kraman, M., Wells, R.J., Deonaraine, A., Chan, D.S., Connell, C.M., Roberts, E.W., Zhao, Q., Caballero, O.L., *et al.* (2013). Targeting CXCL12 from FAP-expressing carcinoma-associated fibroblasts synergizes with anti-PD-L1 immunotherapy in pancreatic cancer. *Proc Natl Acad Sci U S A* *110*, 20212-20217;
- [937] Fernandez-Duenas, V., Taura, J.J., Cottet, M., Gomez-Soler, M., Lopez-Cano, M., Ledent, C., Watanabe, M., Trinquet, E., Pin, J.P., Lujan, R., *et al.* (2015). Untangling dopamine-adenosine receptor-receptor assembly in experimental parkinsonism in rats. *Dis Model Mech* *8*, 57-63;
- [938] Ferre, S., Baler, R., Bouvier, M., Caron, M.G., Devi, L.A., Durroux, T., Fuxe, K., George, S.R., Javitch, J.A., Lohse, M.J., *et al.* (2009). Building a new conceptual framework for receptor heteromers. *Nat Chem Biol* *5*, 131-134;
- [939] Ferre, S., Navarro, G., Casado, V., Cortes, A., Mallol, J., Canela, E.I., Lluís, C., and Franco, R. (2010). G protein-coupled receptor heteromers as new targets for drug development. *Prog Mol Biol Transl Sci* *91*, 41-52;
- [940] Filmore, D. (2004). It's a GPCR world. *Modern Drug Discovery American Chemical Society 2004 (November)*, 24-28;
- [941] Fitzpatrick, D., Purves, D., Augustine, G. (2004). "Table 20:2" (Mass: Sunderland);
- [942] Fotiadis, D., Jastrzebska, B., Philippsen, A., Muller, D.J., Palczewski, K., and Engel, A. (2006). Structure of the rhodopsin dimer: a working model for G-protein-coupled receptors. *Curr Opin Struct Biol* *16*, 252-259;
- [943] Frederick, A.L., Yano, H., Trifilieff, P., Vishwasrao, H.D., Biezonski, D., Meszaros, J., Urizar, E., Sibley, D.R., Kellendonk, C., Sonntag, K.C., *et al.* (2015). Evidence against dopamine D1/D2 receptor heteromers. *Mol Psychiatry* *20*, 1373-1385;
- [944] Furusato, B., Mohamed, A., Uhlen, M., and Rhim, J.S. (2010). CXCR4 and cancer. *Pathol Int* *60*, 497-505;
- [945] Gao, Z.G., Ye, K., Goblyos, A., Ijzerman, A.P., and Jacobson, K.A. (2008). Flexible modulation of agonist efficacy at the human A3 adenosine receptor by the imidazoquinoline allosteric enhancer LUF6000. *BMC Pharmacol* *8*, 20;

- [946] George, S.R., Fan, T., Xie, Z., Tse, R., Tam, V., Varghese, G., and O'Dowd, B.F. (2000). Oligomerization of mu- and delta-opioid receptors. Generation of novel functional properties. *J Biol Chem* 275, 26128-26135;
- [947] Gerard, C., and Gerard, N.P. (1994). C5A anaphylatoxin and its seven trans-membrane-segment receptor. *Annu Rev Immunol* 12, 775-808;
- [948] Gianetti, E., and Seminara, S. (2008). Kisspeptin and KISS1R: a critical pathway in the reproductive system. *Reproduction* 136, 295-301;
- [949] Gomes, I., Ayoub, M.A., Fujita, W., Jaeger, W.C., Pflieger, K.D., and Devi, L.A. (2016). G Protein-Coupled Receptor Heteromers. *Annu Rev Pharmacol Toxicol* 56, 403-425;
- [950] Gomes, I., Gupta, A., Filipovska, J., Szeto, H.H., Pintar, J.E., and Devi, L.A. (2004). A role for heterodimerization of mu and delta opiate receptors in enhancing morphine analgesia. *Proc Natl Acad Sci U S A* 101, 5135-5139;
- [951] Goodman, R.L., Lehman, M.N., Smith, J.T., Coolen, L.M., de Oliveira, C.V., Jafarzadehshirazi, M.R., Pereira, A., Iqbal, J., Caraty, A., Ciofi, P., *et al.* (2007). Kisspeptin neurons in the arcuate nucleus of the ewe express both dynorphin A and neurokinin B. *Endocrinology* 148, 5752-5760;
- [952] Gourni, E., Demmer, O., Schottelius, M., D'Alessandria, C., Schulz, S., Dijkgraaf, I., Schumacher, U., Schwaiger, M., Kessler, H., and Wester, H.J. (2011). PET of CXCR4 expression by a (68)Ga-labeled highly specific targeted contrast agent. *J Nucl Med* 52, 1803-1810;
- [953] Gregorio, G.G., Masureel, M., Hilger, D., Terry, D.S., Juette, M., Zhao, H., Zhou, Z., Perez-Aguilar, J.M., Hauge, M., Mathiasen, S., *et al.* (2017). Single-molecule analysis of ligand efficacy in beta2AR-G-protein activation. *Nature* 547, 68-73;
- [954] Griffith, J.W., Sokol, C.L., and Luster, A.D. (2014). Chemokines and chemokine receptors: positioning cells for host defense and immunity. *Annu Rev Immunol* 32, 659-702;
- [955] Griffiths, K., Dolezal, O., Cao, B., Nilsson, S.K., See, H.B., Pflieger, K.D., Roche, M., Gorry, P.R., Pow, A., Viduka, K., *et al.* (2016). i-bodies, Human Single Domain Antibodies That Antagonize Chemokine Receptor CXCR4. *J Biol Chem* 291, 12641-12657;
- [956] Griffiths, K., Habel, D.M., Jaffar, J., Binder, U., Darby, W.G., Hosking, C.G., Skerra, A., Westall, G.P., Hogaboam, C.M., and Foley, M. (2018). Anti-fibrotic Effects of CXCR4-Targeting i-body AD-114 in Preclinical Models of Pulmonary Fibrosis. *Sci Rep* 8, 3212;
- [957] Guidotti, G., Brambilla, L., and Rossi, D. (2017). Cell-Penetrating Peptides: From Basic Research to Clinics. *Trends Pharmacol Sci* 38, 406-424;
- [958] Gullberg, M., Gustafsdottir, S.M., Schallmeiner, E., Jarvius, J., Bjarnegard, M.,

- Betsholtz, C., Landegren, U., and Fredriksson, S. (2004). Cytokine detection by antibody-based proximity ligation. *Proc Natl Acad Sci U S A* *101*, 8420-8424;
- [959] Guo, F., Wang, Y., Liu, J., Mok, S.C., Xue, F., and Zhang, W. (2016). CXCL12/CXCR4: a symbiotic bridge linking cancer cells and their stromal neighbors in oncogenic communication networks. *Oncogene* *35*, 816-826;
- [960] Gustafsdottir, S.M., Schallmeiner, E., Fredriksson, S., Gullberg, M., Soderberg, O., Jarvius, M., Jarvius, J., Howell, M., and Landegren, U. (2005). Proximity ligation assays for sensitive and specific protein analyses. *Anal Biochem* *345*, 2-9;
- [961] Habert-Ortoli, E., Amiranoff, B., Loquet, I., Laburthe, M., and Mayaux, J.F. (1994). Molecular cloning of a functional human galanin receptor. *Proc Natl Acad Sci U S A* *91*, 9780-9783;
- [962] Hansen, J.L., Hansen, J.T., Speerschnieder, T., Lyngso, C., Erikstrup, N., Burstein, E.S., Weiner, D.M., Walther, T., Makita, N., Iiri, T., *et al.* (2009). Lack of evidence for AT1R/B2R heterodimerization in COS-7, HEK293, and NIH3T3 cells: how common is the AT1R/B2R heterodimer? *J Biol Chem* *284*, 1831-1839;
- [963] Hartimath, S.V., Domanska, U.M., Walenkamp, A.M., Rudi, A.J.O.D., and de Vries, E.F. (2013). [(9)(9)mTc]O(2)-AMD3100 as a SPECT tracer for CXCR4 receptor imaging. *Nucl Med Biol* *40*, 507-517;
- [964] Hassan, S., Buchanan, M., Jahan, K., Aguilar-Mahecha, A., Gaboury, L., Muller, W.J., Alsawafi, Y., Mourskaia, A.A., Siegel, P.M., Salvucci, O., *et al.* (2011). CXCR4 peptide antagonist inhibits primary breast tumor growth, metastasis and enhances the efficacy of anti-VEGF treatment or docetaxel in a transgenic mouse model. *Int J Cancer* *129*, 225-232;
- [965] Hatse, S., Princen, K., Bridger, G., De Clercq, E., and Schols, D. (2002). Chemokine receptor inhibition by AMD3100 is strictly confined to CXCR4. *FEBS Lett* *527*, 255-262;
- [966] He, S.Q., Zhang, Z.N., Guan, J.S., Liu, H.R., Zhao, B., Wang, H.B., Li, Q., Yang, H., Luo, J., Li, Z.Y., *et al.* (2011). Facilitation of mu-opioid receptor activity by preventing delta-opioid receptor-mediated codegradation. *Neuron* *69*, 120-131;
- [967] Heikal, Y., Woll, M.P., Fox, T., Seaton, K., Levenson, R., and Kester, M. (2011). Neurotensin receptor-1 inducible palmitoylation is required for efficient receptor-mediated mitogenic-signaling within structured membrane microdomains. *Cancer Biol Ther* *12*, 427-435;
- [968] Hendrix, C.W., Collier, A.C., Lederman, M.M., Schols, D., Pollard, R.B., Brown, S., Jackson, J.B., Coombs, R.W., Glesby, M.J., Flexner, C.W., *et al.* (2004). Safety, pharmacokinetics, and antiviral activity of AMD3100, a selective CXCR4 receptor inhibitor, in HIV-1 infection. *J Acquir Immune Defic Syndr* *37*, 1253-1262;
- [969] Hendrix, C.W., Flexner, C., MacFarland, R.T., Giandomenico, C., Fuchs, E.J.,

- Redpath, E., Bridger, G., and Henson, G.W. (2000). Pharmacokinetics and safety of AMD-3100, a novel antagonist of the CXCR-4 chemokine receptor, in human volunteers. *Antimicrob Agents Chemother* 44, 1667-1673;
- [970] Henson, B.S., Neubig, R.R., Jang, I., Ogawa, T., Zhang, Z., Carey, T.E., and D'Silva, N.J. (2005). Galanin receptor 1 has anti-proliferative effects in oral squamous cell carcinoma. *J Biol Chem* 280, 22564-22571;
- [971] Hernandez, P.A., Gorlin, R.J., Lukens, J.N., Taniuchi, S., Bohinjec, J., Francois, F., Klotman, M.E., and Diaz, G.A. (2003). Mutations in the chemokine receptor gene CXCR4 are associated with WHIM syndrome, a combined immunodeficiency disease. *Nat Genet* 34, 70-74;
- [972] Herrmann, K., Schottelius, M., Lapa, C., Osl, T., Poschenrieder, A., Hanscheid, H., Luckerath, K., Schreder, M., Bluemel, C., Knott, M., *et al.* (2016). First-in-Human Experience of CXCR4-Directed Endoradiotherapy with ¹⁷⁷Lu- and ⁹⁰Y-Labeled Pentixather in Advanced-Stage Multiple Myeloma with Extensive Intra- and Extramedullary Disease. *J Nucl Med* 57, 248-251;
- [973] Hillion, J., Canals, M., Torvinen, M., Casado, V., Scott, R., Terasmaa, A., Hansson, A., Watson, S., Olah, M.E., Mallol, J., *et al.* (2002). Coaggregation, cointernalization, and codesensitization of adenosine A2A receptors and dopamine D2 receptors. *J Biol Chem* 277, 18091-18097;
- [974] Hsu, W.T., Jui, H.Y., Huang, Y.H., Su, M.Y., Wu, Y.W., Tseng, W.Y., Hsu, M.C., Chiang, B.L., Wu, K.K., and Lee, C.M. (2015). CXCR4 Antagonist TG-0054 Mobilizes Mesenchymal Stem Cells, Attenuates Inflammation, and Preserves Cardiac Systolic Function in a Porcine Model of Myocardial Infarction. *Cell Transplant* 24, 1313-1328;
- [975] Hu, C.D., Chinenov, Y., and Kerppola, T.K. (2002). Visualization of interactions among bZIP and Rel family proteins in living cells using bimolecular fluorescence complementation. *Mol Cell* 9, 789-798;
- [976] Hu, F., Miao, L., Zhao, Y., Xiao, Y.Y., and Xu, Q. (2015). A meta-analysis for CXCR4 chemokine receptor type 4 as a prognostic marker and potential drug target in hepatocellular carcinoma. *Drug Des Devel Ther* 9, 3625-3633;
- [977] Huang, E.H., Singh, B., Cristofanilli, M., Gelovani, J., Wei, C., Vincent, L., Cook, K.R., and Lucci, A. (2009). A CXCR4 antagonist CTCE-9908 inhibits primary tumor growth and metastasis of breast cancer. *J Surg Res* 155, 231-236;
- [978] Hutter, G., Nowak, D., Mossner, M., Ganepola, S., Mussig, A., Allers, K., Schneider, T., Hofmann, J., Kucherer, C., Blau, O., *et al.* (2009). Long-term control of HIV by CCR5 Delta32/Delta32 stem-cell transplantation. *N Engl J Med* 360, 692-698;
- [979] Ichiyama, K., Yokoyama-Kumakura, S., Tanaka, Y., Tanaka, R., Hirose, K., Bannai, K., Edamatsu, T., Yanaka, M., Niitani, Y., Miyano-Kurosaki, N., *et al.* (2003). A

- duodenally absorbable CXC chemokine receptor 4 antagonist, KRH-1636, exhibits a potent and selective anti-HIV-1 activity. *Proc Natl Acad Sci U S A* *100*, 4185-4190;
- [980] Inokuchi, E., Oishi, S., Kubo, T., Ohno, H., Shimura, K., Matsuoka, M., and Fujii, N. (2011). Potent CXCR4 antagonists containing amidine type Peptide bond isosteres. *ACS Med Chem Lett* *2*, 477-480;
- [981] Jafari, S.M., Panjehpour, M., Aghaei, M., Joshaghani, H.R., and Enderami, S.E. (2017). A3 Adenosine Receptor Agonist Inhibited Survival of Breast Cancer Stem Cells via GLI-1 and ERK1/2 Pathway. *J Cell Biochem* *118*, 2909-2920;
- [982] Jahnichen, S., Blanchetot, C., Maussang, D., Gonzalez-Pajuelo, M., Chow, K.Y., Bosch, L., De Vrieze, S., Serruys, B., Ulrichs, H., Vandeveld, W., *et al.* (2010). CXCR4 nanobodies (VHH-based single variable domains) potently inhibit chemotaxis and HIV-1 replication and mobilize stem cells. *Proc Natl Acad Sci U S A* *107*, 20565-20570;
- [983] Jenkinson, S., Thomson, M., McCoy, D., Edelstein, M., Danehower, S., Lawrence, W., Wheelan, P., Spaltenstein, A., and Gudmundsson, K. (2010). Blockade of X4-tropic HIV-1 cellular entry by GSK812397, a potent noncompetitive CXCR4 receptor antagonist. *Antimicrob Agents Chemother* *54*, 817-824;
- [984] Jiang, X.R., Song, A., Bergelson, S., Arroll, T., Parekh, B., May, K., Chung, S., Strouse, R., Mire-Sluis, A., and Schenerman, M. (2011). Advances in the assessment and control of the effector functions of therapeutic antibodies. *Nat Rev Drug Discov* *10*, 101-111;
- [985] Johnson, G. (2002). *PDQ Pharmacology* (2nd ed.) (Hamilton, Ontario: BC Decker Inc.);
- [986] Kalatskaya, I., Berchiche, Y.A., Gravel, S., Limberg, B.J., Rosenbaum, J.S., and Heveker, N. (2009). AMD3100 is a CXCR7 ligand with allosteric agonist properties. *Mol Pharmacol* *75*, 1240-1247;
- [987] Kasama, H., Sakamoto, Y., Kasamatsu, A., Okamoto, A., Koyama, T., Minakawa, Y., Ogawara, K., Yokoe, H., Shiiba, M., Tanzawa, H., *et al.* (2015). Adenosine A2b receptor promotes progression of human oral cancer. *BMC Cancer* *15*, 563;
- [988] Kawai, T., and Malech, H.L. (2009). WHIM syndrome: congenital immune deficiency disease. *Curr Opin Hematol* *16*, 20-26;
- [989] Keating, G.M. (2011). Plerixafor: a review of its use in stem-cell mobilization in patients with lymphoma or multiple myeloma. *Drugs* *71*, 1623-1647;
- [990] Kerppola, T.K. (2006). Design and implementation of bimolecular fluorescence complementation (BiFC) assays for the visualization of protein interactions in living cells. *Nat Protoc* *1*, 1278-1286;
- [991] Kim, H.Y., Hwang, J.Y., Kim, S.W., Lee, H.J., Yun, H.J., Kim, S., and Jo, D.Y. (2010). The CXCR4 Antagonist AMD3100 Has Dual Effects on Survival and Pro-

- liferation of Myeloma Cells *In vitro*. Cancer Res Treat 42, 225-234;
- [992] Kim, S.Y., Lee, C.H., Midura, B.V., Yeung, C., Mendoza, A., Hong, S.H., Ren, L., Wong, D., Korz, W., Merzouk, A., *et al.* (2008). Inhibition of the CXCR4/CXCL12 chemokine pathway reduces the development of murine pulmonary metastases. Clin Exp Metastasis 25, 201-211;
- [993] Kitazawa, T., Taneike, T., and Ohga, A. (1995). Excitatory action of [Leu13]motilin on the gastrointestinal smooth muscle isolated from the chicken. Peptides 16, 1243-1252;
- [994] Kitazawa, T., Taneike, T., and Ohga, A. (1997). Functional characterization of neural and smooth muscle motilin receptors in the chicken proventriculus and ileum. Regul Pept 71, 87-95;
- [995] Klein, R.S., and Rubin, J.B. (2004). Immune and nervous system CXCL12 and CXCR4: parallel roles in patterning and plasticity. Trends Immunol 25, 306-314;
- [996] Klos, A., Wende, E., Wareham, K.J., and Monk, P.N. (2013). International Union of Basic and Clinical Pharmacology. [corrected]. LXXXVII. Complement peptide C5a, C4a, and C3a receptors. Pharmacol Rev 65, 500-543;
- [997] Knutsson, M., and Edvinsson, L. (2002). Distribution of mRNA for VIP and PACAP receptors in human cerebral arteries and cranial ganglia. Neuroreport 13, 507-509;
- [998] Kristensen, M., Birch, D., and Morck Nielsen, H. (2016). Applications and Challenges for Use of Cell-Penetrating Peptides as Delivery Vectors for Peptide and Protein Cargos. Int J Mol Sci 17;
- [999] Kroeze, W.K., Sassano, M.F., Huang, X.P., Lansu, K., McCorvy, J.D., Giguere, P.M., Sciaky, N., and Roth, B.L. (2015). PRESTO-Tango as an open-source resource for interrogation of the druggable human GPCRome. Nat Struct Mol Biol 22, 362-369;
- [1000] Kuhne, M.R., Mulvey, T., Belanger, B., Chen, S., Pan, C., Chong, C., Cao, F., Niekro, W., Kempe, T., Henning, K.A., *et al.* (2013). BMS-936564/MDX-1338: a fully human anti-CXCR4 antibody induces apoptosis *in vitro* and shows antitumor activity *in vivo* in hematologic malignancies. Clin Cancer Res 19, 357-366;
- [1001] Lambert, et al., "Antibody-Drug Conjugates for Cancer Treatment", Annual Review of Medicine, 69:191-207 (2018);
- [1002] LaRocca, T.J., Schwarzkopf, M., Altman, P., Zhang, S., Gupta, A., Gomes, I., Alvin, Z., Champion, H.C., Haddad, G., Hajjar, R.J., *et al.* (2010). beta2-Adrenergic receptor signaling in the cardiac myocyte is modulated by interactions with CXCR4. J Cardiovasc Pharmacol 56, 548-559;
- [1003] Law, P.Y., Erickson-Herbrandson, L.J., Zha, Q.Q., Solberg, J., Chu, J., Sarre, A., and Loh, H.H. (2005). Heterodimerization of mu- and delta-opioid receptors occurs at the cell surface only and requires receptor-G protein interactions. J Biol Chem 280, 11152-11164;

- [1004] Lee, C.H., Kakinuma, T., Wang, J., Zhang, H., Palmer, D.C., Restifo, N.P., and Hwang, S.T. (2006). Sensitization of B16 tumor cells with a CXCR4 antagonist increases the efficacy of immunotherapy for established lung metastases. *Mol Cancer Ther* 5, 2592-2599;
- [1005] Lee, H., Whitfeld, P.L., and Mackay, C.R. (2008). Receptors for complement C5a. The importance of C5aR and the enigmatic role of C5L2. *Immunol Cell Biol* 86, 153-160;
- [1006] Levoe, A., Balabanian, K., Baleux, F., Bachelier, F., and Lagane, B. (2009). CXCR7 heterodimerizes with CXCR4 and regulates CXCL12-mediated G protein signaling. *Blood* 113, 6085-6093;
- [1007] Li, M., Thompson, D.D., and Paralkar, V.M. (2007). Prostaglandin E(2) receptors in bone formation. *Int Orthop* 31, 767-772;
- [1008] Li, Y.P., Pang, J., Gao, S., Bai, P.Y., Wang, W.D., Kong, P., and Cui, Y. (2017). Role of CXCR4 and SDF1 as prognostic factors for survival and the association with clinicopathology in colorectal cancer: A systematic meta-analysis. *Tumour Biol* 39, 1010428317706206;
- [1009] Liang, J.X., Gao, W., Liang, Y., and Zhou, X.M. (2015). Chemokine receptor CXCR4 expression and lung cancer prognosis: a meta-analysis. *Int J Clin Exp Med* 8, 5163-5174;
- [1010] Liang, Z., Zhan, W., Zhu, A., Yoon, Y., Lin, S., Sasaki, M., Klapproth, J.M., Yang, H., Grossniklaus, H.E., Xu, J., *et al.* (2012). Development of a unique small molecule modulator of CXCR4. *PLoS One* 7, e34038;
- [1011] Liao, Y.X., Fu, Z.Z., Zhou, C.H., Shan, L.C., Wang, Z.Y., Yin, F., Zheng, L.P., Hua, Y.Q., and Cai, Z.D. (2015). AMD3100 reduces CXCR4-mediated survival and metastasis of osteosarcoma by inhibiting JNK and Akt, but not p38 or Erk1/2, pathways in *in vitro* and mouse experiments. *Oncol Rep* 34, 33-42;
- [1012] Liles, W.C., Broxmeyer, H.E., Rodger, E., Wood, B., Hubel, K., Cooper, S., Hangoc, G., Bridger, G.J., Henson, G.W., Calandra, G., *et al.* (2003). Mobilization of hematopoietic progenitor cells in healthy volunteers by AMD3100, a CXCR4 antagonist. *Blood* 102, 2728-2730;
- [1013] Ling, X., Spaeth, E., Chen, Y., Shi, Y., Zhang, W., Schober, W., Hail, N., Jr., Konopleva, M., and Andreeff, M. (2013). The CXCR4 antagonist AMD3465 regulates oncogenic signaling and invasiveness *in vitro* and prevents breast cancer growth and metastasis *in vivo*. *PLoS One* 8, e58426;
- [1014] Liu, S.H., Gu, Y., Pascual, B., Yan, Z., Hallin, M., Zhang, C., Fan, C., Wang, W., Lam, J., Spilker, M.E., *et al.* (2017). A novel CXCR4 antagonist IgG1 antibody (PF-06747143) for the treatment of hematologic malignancies. *Blood Adv* 1, 1088-1100;

- [1015] Liu, X.Y., Liu, Z.C., Sun, Y.G., Ross, M., Kim, S., Tsai, F.F., Li, Q.F., Jeffry, J., Kim, J.Y., Loh, H.H., *et al.* (2011). Unidirectional cross-activation of GRPR by MOR1D uncouples itch and analgesia induced by opioids. *Cell* 147, 447-458;
- [1016] Lohse, M.J., Nuber, S., and Hoffmann, C. (2012). Fluorescence/bioluminescence resonance energy transfer techniques to study G-protein-coupled receptor activation and signaling. *Pharmacol Rev* 64, 299-336;
- [1017] Lowe, S.R., Pothén, J., Quinn, J.W., Rundle, A., Bradley, B., Galea, S., Ressler, K.J., and Koenen, K.C. (2015). Gene-by-social-environment interaction (GxSE) between ADCYAP1R1 genotype and neighborhood crime predicts major depression symptoms in trauma-exposed women. *J Affect Disord* 187, 147-150;
- [1018] Machado-Carvalho, L., Roca-Ferrer, J., and Picado, C. (2014). Prostaglandin E2 receptors in asthma and in chronic rhinosinusitis/nasal polyps with and without aspirin hypersensitivity. *Respir Res* 15, 100;
- [1019] Markovic, T., Jakopin, Z., Dolenc, M.S., and Mlinaric-Rascan, I. (2017). Structural features of subtype-selective EP receptor modulators. *Drug Discov Today* 22, 57-71;
- [1020] Marlo, J.E., Niswender, C.M., Days, E.L., Bridges, T.M., Xiang, Y., Rodriguez, A.L., Shirey, J.K., Brady, A.E., Nalawajko, T., Luo, Q., *et al.* (2009). Discovery and characterization of novel allosteric potentiators of M1 muscarinic receptors reveals multiple modes of activity. *Mol Pharmacol* 75, 577-588;
- [1021] Martinez-Munoz, L., Barroso, R., Dyrhaug, S.Y., Navarro, G., Lucas, P., Soriano, S.F., Vega, B., Costas, C., Munoz-Fernandez, M.A., Santiago, C., *et al.* (2014). CCR5/CD4/CXCR4 oligomerization prevents HIV-1 gp120IIIB binding to the cell surface. *Proc Natl Acad Sci U S A* 111, E1960-1969;
- [1022] Ueda, M., Nakashima, H., Ueda, T., Naba, H., Ikoma, R., Otaka, A., Terakawa, Y., Tamamura, H., Ibuka, T., Murakami, T., *et al.* (1992). A novel anti-HIV synthetic peptide, T-22 ([Tyr5,12,Lys7]-polyphemusin II). *Biochem Biophys Res Commun* 189, 845-850;
- [1023] Medhurst, A.D., Jennings, C.A., Robbins, M.J., Davis, R.P., Ellis, C., Winborn, K.Y., Lawrie, K.W., Hervieu, G., Riley, G., Bolaky, J.E., *et al.* (2003). Pharmacological and immunohistochemical characterization of the APJ receptor and its endogenous ligand apelin. *J Neurochem* 84, 1162-1172;
- [1024] Miller, G.M., Alexander, J.M., Bikkal, H.A., Katznelson, L., Zervas, N.T., and Klibanski, A. (1995). Somatostatin receptor subtype gene expression in pituitary adenomas. *J Clin Endocrinol Metab* 80, 1386-1392;
- [1025] Milligan, G. (2008). A day in the life of a G protein-coupled receptor: the contribution to function of G protein-coupled receptor dimerization. *Br J Pharmacol* 153 Suppl 1, S216-229;
- [1026] Miwatashi, S., Arikawa, Y., Matsumoto, T., Uga, K., Kanzaki, N., Imai, Y.N., and

- Ohkawa, S. (2008). Synthesis and biological activities of 4-phenyl-5-pyridyl-1,3-thiazole derivatives as selective adenosine A3 antagonists. *Chem Pharm Bull (Tokyo)* 56, 1126-1137;
- [1027] Moreno, J.J. (2017). Eicosanoid receptors: Targets for the treatment of disrupted intestinal epithelial homeostasis. *Eur J Pharmacol* 796, 7-19;
- [1028] Imoto, M., Matsuo, Y., Koide, S., Tsuboi, K., Shamoto, T., Sato, T., Saito, K., Takahashi, H., and Takeyama, H. (2016). Enhancement of the CXCL12/CXCR4 axis due to acquisition of gemcitabine resistance in pancreatic cancer: effect of CXCR4 antagonists. *BMC Cancer* 16, 305;
- [1029] Muller, A., Homey, B., Soto, H., Ge, N., Catron, D., Buchanan, M.E., McClanahan, T., Murphy, E., Yuan, W., Wagner, S.N., *et al.* (2001). Involvement of chemokine receptors in breast cancer metastasis. *Nature* 410, 50-56;
- [1030] Murakami, T., Kumakura, S., Yamazaki, T., Tanaka, R., Hamatake, M., Okuma, K., Huang, W., Toma, J., Komano, J., Yanaka, M., *et al.* (2009). The novel CXCR4 antagonist KRH-3955 is an orally bioavailable and extremely potent inhibitor of human immunodeficiency virus type 1 infection: comparative studies with AMD3100. *Antimicrob Agents Chemother* 53, 2940-2948;
- [1031] Mustafa, S., Ayoub, M.A., Pflieger, K.D.G. (2010). Uncovering GPCR heteromer-biased ligands. *Drug Discov Today Technol* 7, e1-e94;
- [1032] Mustafa, S., and Pflieger, K.D. (2011). G protein-coupled receptor heteromer identification technology: identification and profiling of GPCR heteromers. *J Lab Autom* 16, 285-291;
- [1033] Mustafa, S., See, H.B., Seeber, R.M., Armstrong, S.P., White, C.W., Ventura, S., Ayoub, M.A., and Pflieger, K.D. (2012). Identification and profiling of novel alpha1A-adrenoceptor-CXC chemokine receptor 2 heteromer. *J Biol Chem* 287, 12952-12965;
- [1034] Nakai, A., Hayano, Y., Furuta, F., Noda, M., and Suzuki, K. (2014). Control of lymphocyte egress from lymph nodes through beta2-adrenergic receptors. *J Exp Med* 211, 2583-2598;
- [1035] Nakasone, T., Kumakura, S., Yamamoto, M., Murakami, T., and Yamamoto, N. (2013). Single oral administration of the novel CXCR4 antagonist, KRH-3955, induces an efficient and long-lasting increase of white blood cell count in normal macaques, and prevents CD4 depletion in SHIV-infected macaques: a preliminary study. *Med Microbiol Immunol* 202, 175-182;
- [1036] Norel, X., Jones, R.L., Giembycz, M., Narumiya, S., Woodward, D.F., Coleman, R.A., Abramovitz, M., Breyer, R.M., Hills, R. (2016). Prostanoid receptors: EP3 receptor. *IUPHAR/BPS Guide to Pharmacology*;
- [1037] O'Boyle, G., Swidenbank, I., Marshall, H., Barker, C.E., Armstrong, J., White, S.A.,

- Fricker, S.P., Plummer, R., Wright, M., and Lovat, P.E. (2013). Inhibition of CXCR4-CXCL12 chemotaxis in melanoma by AMD11070. *Br J Cancer* 108, 1634-1640;
- [1038] O'Callaghan, K., Kuliopulos, A., and Covic, L. (2012). Turning receptors on and off with intracellular pepducins: new insights into G-protein-coupled receptor drug development. *J Biol Chem* 287, 12787-12796;
- [1039] O'Callaghan, G., and Houston, A. (2015). Prostaglandin E2 and the EP receptors in malignancy: possible therapeutic targets? *Br J Pharmacol* 172, 5239-5250;
- [1040] Ogi, K., Miyamoto, Y., Masuda, Y., Habata, Y., Hosoya, M., Ohtaki, T., Masuo, Y., Onda, H., and Fujino, M. (1993). Molecular cloning and functional expression of a cDNA encoding a human pituitary adenylate cyclase activating polypeptide receptor. *Biochem Biophys Res Commun* 196, 1511-1521;
- [1041] Okada, T., Ernst, O.P., Palczewski, K., and Hofmann, K.P. (2001). Activation of rhodopsin: new insights from structural and biochemical studies. *Trends Biochem Sci* 26, 318-324;
- [1042] Otani, Y., Kijima, T., Kohmo, S., Oishi, S., Minami, T., Nagatomo, I., Takahashi, R., Hirata, H., Suzuki, M., Inoue, K., *et al.* (2012). Suppression of metastases of small cell lung cancer cells in mice by a peptidic CXCR4 inhibitor TF14016. *FEBS Lett* 586, 3639-3644;
- [1043] Overington, J.P., Al-Lazikani, B., and Hopkins, A.L. (2006). How many drug targets are there? *Nat Rev Drug Discov* 5, 993-996;
- [1044] Owen, J.L., and Mohamadzaheh, M. (2013). Macrophages and chemokines as mediators of angiogenesis. *Front Physiol* 4, 159;
- [1045] Page, N.M., Bell, N.J., Gardiner, S.M., Manyonda, I.T., Brayley, K.J., Strange, P.G., and Lowry, P.J. (2003). Characterization of the endokinins: human tachykinins with cardiovascular activity. *Proc Natl Acad Sci U S A* 100, 6245-6250;
- [1046] Pal, J., Patil, V., Kumar, A., Kaur, K., Sarkar, C., and Somasundaram, K. (2018). Loss-of-Function Mutations in Calcitonin Receptor (CALCR) Identify Highly Aggressive Glioblastoma with Poor Outcome. *Clin Cancer Res* 24, 1448-1458;
- [1047] Parameswaran, R., Yu, M., Lim, M., Groffen, J., and Heisterkamp, N. (2011). Combination of drug therapy in acute lymphoblastic leukemia with a CXCR4 antagonist. *Leukemia* 25, 1314-1323;
- [1048] Park, P.S., and Palczewski, K. (2005). Diversifying the repertoire of G protein-coupled receptors through oligomerization. *Proc Natl Acad Sci U S A* 102, 8793-8794;
- [1049] Patel, K., Dixit, V.D., Lee, J.H., Kim, J.W., Schaffer, E.M., Nguyen, D., and Taub, D.D. (2012). Identification of ghrelin receptor blocker, D-[Lys3] GHRP-6 as a CXCR4 receptor antagonist. *Int J Biol Sci* 8, 108-117;
- [1050] Patel, R.C., Kumar, U., Lamb, D.C., Eid, J.S., Rocheville, M., Grant, M., Rani, A.,

- Hazlett, T., Patel, S.C., Gratton, E., *et al.* (2002). Ligand binding to somatostatin receptors induces receptor-specific oligomer formation in live cells. *Proc Natl Acad Sci U S A* 99, 3294-3299;
- [1051] Peled, A., Wald, O., and Burger, J. (2012). Development of novel CXCR4-based therapeutics. *Expert Opin Investig Drugs* 21, 341-353;
- [1052] Pello, O.M., Martinez-Munoz, L., Parrillas, V., Serrano, A., Rodriguez-Frade, J.M., Toro, M.J., Lucas, P., Monterrubio, M., Martinez, A.C., and Mellado, M. (2008). Ligand stabilization of CXCR4/delta-opioid receptor heterodimers reveals a mechanism for immune response regulation. *Eur J Immunol* 38, 537-549;
- [1053] Peng, S.B., Zhang, X., Paul, D., Kays, L.M., Gough, W., Stewart, J., Uhlik, M.T., Chen, Q., Hui, Y.H., Zamek-Gliszczynski, M.J., *et al.* (2015). Identification of LY2510924, a novel cyclic peptide CXCR4 antagonist that exhibits antitumor activities in solid tumor and breast cancer metastatic models. *Mol Cancer Ther* 14, 480-490;
- [1054] Pfeiffer, M., Koch, T., Schroder, H., Laugsch, M., Holtt, V., and Schulz, S. (2002). Heterodimerization of somatostatin and opioid receptors cross-modulates phosphorylation, internalization, and desensitization. *J Biol Chem* 277, 19762-19772;
- [1055] Pfleger, K.D., and Eidne, K.A. (2006). Illuminating insights into protein-protein interactions using bioluminescence resonance energy transfer (BRET). *Nat Methods* 3, 165-174;
- [1056] Pin, J.P., Neubig, R., Bouvier, M., Devi, L., Filizola, M., Javitch, J.A., Lohse, M.J., Milligan, G., Palczewski, K., Parmentier, M., *et al.* (2007). International Union of Basic and Clinical Pharmacology. LXVII. Recommendations for the recognition and nomenclature of G protein-coupled receptor heteromultimers. *Pharmacol Rev* 59, 5-13;
- [1057] Planesas, J.M., Perez-Nueno, V.I., Borrell, J.I., and Teixido, J. (2015). Studying the binding interactions of allosteric agonists and antagonists of the CXCR4 receptor. *J Mol Graph Model* 60, 1-14;
- [1058] Pondel, M. (2000). Calcitonin and calcitonin receptors: bone and beyond. *Int J Exp Pathol* 81, 405-422;
- [1059] Qin, K., Dong, C., Wu, G., and Lambert, N.A. (2011). Inactive-state preassembly of G(q)-coupled receptors and G(q) heterotrimers. *Nat Chem Biol* 7, 740-747;
- [1060] Rang, H.P., Dale, M.M., Ritter, J.M., Moore, P.K. (2003). "Ch. 10", *Pharmacology* (5th ed.). (Elsevier Churchill Livingstone);
- [1061] Redondo-Munoz, J., Escobar-Diaz, E., Samaniego, R., Terol, M.J., Garcia-Marco, J.A., and Garcia-Pardo, A. (2006). MMP-9 in B-cell chronic lymphocytic leukemia is up-regulated by alpha4beta1 integrin or CXCR4 engagement via distinct signaling pathways, localizes to podosomes, and is involved in cell invasion and migration. *Blood* 108, 3143-3151;

- [1062] Reubi, J.C. (2000). *In vitro* evaluation of VIP/PACAP receptors in healthy and diseased human tissues. Clinical implications. *Ann N Y Acad Sci* 921, 1-25;
- [1063] Reubi, J.C., Laderach, U., Waser, B., Gebbers, J.O., Robberecht, P., and Laissue, J.A. (2000). Vasoactive intestinal peptide/pituitary adenylate cyclase-activating peptide receptor subtypes in human tumors and their tissues of origin. *Cancer Res* 60, 3105-3112;
- [1064] Reya, T., Morrison, S.J., Clarke, M.F., and Weissman, I.L. (2001). Stem cells, cancer, and cancer stem cells. *Nature* 414, 105-111;
- [1065] Rios, C., Gomes, I., and Devi, L.A. (2006). mu opioid and CB1 cannabinoid receptor interactions: reciprocal inhibition of receptor signaling and neuritogenesis. *Br J Pharmacol* 148, 387-395;
- [1066] Roccaro, A.M., Sacco, A., Purschke, W.G., Moschetta, M., Buchner, K., Maasch, C., Zboralski, D., Zollner, S., Vonhoff, S., Mishima, Y., *et al.* (2014). SDF-1 inhibition targets the bone marrow niche for cancer therapy. *Cell Rep* 9, 118-128;
- [1067] Rodriguez-Frade, J.M., del Real, G., Serrano, A., Hernanz-Falcon, P., Soriano, S.F., Vila-Coro, A.J., de Ana, A.M., Lucas, P., Prieto, I., Martinez, A.C., *et al.* (2004). Blocking HIV-1 infection via CCR5 and CXCR4 receptors by acting in trans on the CCR2 chemokine receptor. *EMBO J* 23, 66-76;
- [1068] Roess, D.A., Horvat, R.D., Munnely, H., and Barisas, B.G. (2000). Luteinizing hormone receptors are self-associated in the plasma membrane. *Endocrinology* 141, 4518-4523;
- [1069] Rozenfeld, R., Bushlin, I., Gomes, I., Tzavaras, N., Gupta, A., Neves, S., Battini, L., Gusella, G.L., Lachmann, A., Ma'ayan, A., *et al.* (2012). Receptor heteromerization expands the repertoire of cannabinoid signaling in rodent neurons. *PLoS One* 7, e29239;
- [1070] Rozenfeld, R., and Devi, L.A. (2010). Receptor heteromerization and drug discovery. *Trends Pharmacol Sci* 31, 124-130;
- [1071] Rubin, J.B., Kung, A.L., Klein, R.S., Chan, J.A., Sun, Y., Schmidt, K., Kieran, M.W., Luster, A.D., and Segal, R.A. (2003). A small-molecule antagonist of CXCR4 inhibits intracranial growth of primary brain tumors. *Proc Natl Acad Sci U S A* 100, 13513-13518;
- [1072] Saka, Y., Hagemann, A.I., Piepenburg, O., and Smith, J.C. (2007). Nuclear accumulation of Smad complexes occurs only after the midblastula transition in *Xenopus*. *Development* 134, 4209-4218;
- [1073] Sanjana, N.E., Shalem, O., and Zhang, F. (2014). Improved vectors and genome-wide libraries for CRISPR screening. *Nat Methods* 11, 783-784;
- [1074] Sato-Jin, K., Nishimura, E.K., Akasaka, E., Huber, W., Nakano, H., Miller, A., Du, J., Wu, M., Hanada, K., Sawamura, D., *et al.* (2008). Epistatic connections between

- microphthalmia-associated transcription factor and endothelin signaling in Waardenburg syndrome and other pigmentary disorders. *FASEB J* 22, 1155-1168;
- [1075] Scala, S. (2015). Molecular Pathways: Targeting the CXCR4-CXCL12 Axis-Untapped Potential in the Tumor Microenvironment. *Clin Cancer Res* 21, 4278-4285;
- [1076] Schimanski, C.C., Bahre, R., Gockel, I., Muller, A., Frerichs, K., Horner, V., Teufel, A., Simiantonaki, N., Biesterfeld, S., Wehler, T., *et al.* (2006). Dissemination of hepatocellular carcinoma is mediated via chemokine receptor CXCR4. *Br J Cancer* 95, 210-217;
- [1077] Sedor, J.R., and Abboud, H.E. (1984). Actions and metabolism of histamine in glomeruli and tubules of the human kidney. *Kidney Int* 26, 144-152;
- [1078] Shen, Z., Chen, X., Li, Q., Zhou, C., Li, J., Ye, H., and Duan, S. (2016). SSTR2 promoter hypermethylation is associated with the risk and progression of laryngeal squamous cell carcinoma in males. *Diagn Pathol* 11, 10;
- [1079] Sicoli, D., Jiao, X., Ju, X., Velasco-Velazquez, M., Ertel, A., Addya, S., Li, Z., Ando, S., Fatatis, A., Paudyal, B., *et al.* (2014). CCR5 receptor antagonists block metastasis to bone of v-Src oncogene-transformed metastatic prostate cancer cell lines. *Cancer Res* 74, 7103-7114;
- [1080] Sierro, F., Biben, C., Martinez-Munoz, L., Mellado, M., Ransohoff, R.M., Li, M., Woehl, B., Leung, H., Groom, J., Batten, M., *et al.* (2007). Disrupted cardiac development but normal hematopoiesis in mice deficient in the second CXCL12/SDF-1 receptor, CXCR7. *Proc Natl Acad Sci U S A* 104, 14759-14764;
- [1081] Smith, M.C., Luker, K.E., Garbow, J.R., Prior, J.L., Jackson, E., Piwnica-Worms, D., and Luker, G.D. (2004). CXCR4 regulates growth of both primary and metastatic breast cancer. *Cancer Res* 64, 8604-8612;
- [1082] Sohy, D., Parmentier, M., and Springael, J.Y. (2007). Allosteric transinhibition by specific antagonists in CCR2/CXCR4 heterodimers. *J Biol Chem* 282, 30062-30069;
- [1083] Sohy, D., Yano, H., de Nadai, P., Urizar, E., Guillabert, A., Javitch, J.A., Parmentier, M., and Springael, J.Y. (2009). Hetero-oligomerization of CCR2, CCR5, and CXCR4 and the protean effects of "selective" antagonists. *J Biol Chem* 284, 31270-31279;
- [1084] Song, J.S., Kang, C.M., Kang, H.H., Yoon, H.K., Kim, Y.K., Kim, K.H., Moon, H.S., and Park, S.H. (2010). Inhibitory effect of CXC chemokine receptor 4 antagonist AMD3100 on bleomycin induced murine pulmonary fibrosis. *Exp Mol Med* 42, 465-472;
- [1085] Song, Y.B., Park, C.O., Jeong, J.Y., and Huh, W.K. (2014). Monitoring G protein-coupled receptor activation using an adenovirus-based beta-arrestin bimolecular fluorescence complementation assay. *Anal Biochem* 449, 32-41;
- [1086] Stefan, E., Aquin, S., Berger, N., Landry, C.R., Nyfeler, B., Bouvier, M., and Michnick, S.W. (2007). Quantification of dynamic protein complexes using Renilla lu-

- ciferase fragment complementation applied to protein kinase A activities *in vivo*. *Proc Natl Acad Sci U S A* 104, 16916-16921;
- [1087] Stevenson, L., Allen, W.L., Turkington, R., Jithesh, P.V., Proutski, I., Stewart, G., Lenz, H.J., Van Schaeybroeck, S., Longley, D.B., and Johnston, P.G. (2012). Identification of galanin and its receptor GalR1 as novel determinants of resistance to chemotherapy and potential biomarkers in colorectal cancer. *Clin Cancer Res* 18, 5412-5426;
- [1088] Stone, N.D., Dunaway, S.B., Flexner, C., Tierney, C., Calandra, G.B., Becker, S., Cao, Y.J., Wiggins, I.P., Conley, J., MacFarland, R.T., *et al.* (2007). Multiple-dose escalation study of the safety, pharmacokinetics, and biologic activity of oral AMD070, a selective CXCR4 receptor inhibitor, in human subjects. *Antimicrob Agents Chemother* 51, 2351-2358;
- [1089] Struyf, S., Menten, P., Lenaerts, J.P., Put, W., D'Haese, A., De Clercq, E., Schols, D., Proost, P., and Van Damme, J. (2001). Diverging binding capacities of natural LD78beta isoforms of macrophage inflammatory protein-1alpha to the CC chemokine receptors 1, 3 and 5 affect their anti-HIV-1 activity and chemotactic potencies for neutrophils and eosinophils. *Eur J Immunol* 31, 2170-2178;
- [1090] Sugimoto, Y., Inazumi, T., and Tsuchiya, S. (2015). Roles of prostaglandin receptors in female reproduction. *J Biochem* 157, 73-80;
- [1091] Swift, S.L., Burns, J.E., and Maitland, N.J. (2010). Altered expression of neurotensin receptors is associated with the differentiation state of prostate cancer. *Cancer Res* 70, 347-356;
- [1092] Taichman, R.S., Cooper, C., Keller, E.T., Pienta, K.J., Taichman, N.S., and McCauley, L.K. (2002). Use of the stromal cell-derived factor-1/CXCR4 pathway in prostate cancer metastasis to bone. *Cancer Res* 62, 1832-1837;
- [1093] Tamamura, H., Xu, Y., Hattori, T., Zhang, X., Arakaki, R., Kanbara, K., Omagari, A., Otaka, A., Ibuka, T., Yamamoto, N., *et al.* (1998). A low-molecular-weight inhibitor against the chemokine receptor CXCR4: a strong anti-HIV peptide T140. *Biochem Biophys Res Commun* 253, 877-882;
- [1094] Tanaka, H., Moroi, K., Iwai, J., Takahashi, H., Ohnuma, N., Hori, S., Takimoto, M., Nishiyama, M., Masaki, T., Yanagisawa, M., *et al.* (1998). Novel mutations of the endothelin B receptor gene in patients with Hirschsprung's disease and their characterization. *J Biol Chem* 273, 11378-11383;
- [1095] Tanaka, T., Nomura, W., Narumi, T., Esaka, A., Oishi, S., Ohashi, N., Itotani, K., Evans, B.J., Wang, Z.X., Peiper, S.C., *et al.* (2009). Structure-activity relationship study on artificial CXCR4 ligands possessing the cyclic pentapeptide scaffold: the exploration of amino acid residues of pentapeptides by substitutions of several aromatic amino acids. *Org Biomol Chem* 7, 3805-3809;

- [1096] Tanaka, T., Tsutsumi, H., Nomura, W., Tanabe, Y., Ohashi, N., Esaka, A., Ochiai, C., Sato, J., Itotani, K., Murakami, T., *et al.* (2008). Structure-activity relationship study of CXCR4 antagonists bearing the cyclic pentapeptide scaffold: identification of the new pharmacophore. *Org Biomol Chem* 6, 4374-4377;
- [1097] Terrillon, S., and Bouvier, M. (2004). Roles of G-protein-coupled receptor dimerization. *EMBO Rep* 5, 30-34;
- [1098] Topaloglu, A.K., Reimann, F., Guclu, M., Yalin, A.S., Kotan, L.D., Porter, K.M., Serin, A., Mungan, N.O., Cook, J.R., Imamoglu, S., *et al.* (2009). TAC3 and TACR3 mutations in familial hypogonadotropic hypogonadism reveal a key role for Neurokinin B in the central control of reproduction. *Nat Genet* 41, 354-358;
- [1099] Torvinen, M., Torri, C., Tombesi, A., Marcellino, D., Watson, S., Lluís, C., Franco, R., Fuxe, K., and Agnati, L.F. (2005). Trafficking of adenosine A2A and dopamine D2 receptors. *J Mol Neurosci* 25, 191-200;
- [1100] Tripathi, A., Davis, J.D., Staren, D.M., Volkman, B.F., and Majetschak, M. (2014). CXC chemokine receptor 4 signaling upon co-activation with stromal cell-derived factor-1 α and ubiquitin. *Cytokine* 65, 121-125;
- [1101] Tripathi, A., Vana, P.G., Chavan, T.S., Brueggemann, L.I., Byron, K.L., Tarasova, N.I., Volkman, B.F., Gaponenko, V., and Majetschak, M. (2015). Heteromerization of chemokine (C-X-C motif) receptor 4 with α 1A/B-adrenergic receptors controls α 1-adrenergic receptor function. *Proc Natl Acad Sci U S A* 112, E1659-1668;
- [1102] Vaudry, D., Gonzalez, B.J., Basille, M., Yon, L., Fournier, A., and Vaudry, H. (2000). Pituitary adenylate cyclase-activating polypeptide and its receptors: from structure to functions. *Pharmacol Rev* 52, 269-324;
- [1103] Velasco-Velazquez, M., Jiao, X., De La Fuente, M., Pestell, T.G., Ertel, A., Lisanti, M.P., and Pestell, R.G. (2012). CCR5 antagonist blocks metastasis of basal breast cancer cells. *Cancer Res* 72, 3839-3850;
- [1104] Vincent, J.P. (1995). Neurotensin receptors: binding properties, transduction pathways, and structure. *Cell Mol Neurobiol* 15, 501-512;
- [1105] Walenkamp, A.M.E., Lapa, C., Herrmann, K., and Wester, H.J. (2017). CXCR4 Ligands: The Next Big Hit? *J Nucl Med* 58, 77S-82S;
- [1106] Wang, A., Fairhurst, A.M., Tus, K., Subramanian, S., Liu, Y., Lin, F., Igarashi, P., Zhou, X.J., Batteux, F., Wong, D., *et al.* (2009). CXCR4/CXCL12 hyperexpression plays a pivotal role in the pathogenesis of lupus. *J Immunol* 182, 4448-4458;
- [1107] Wang, A., Guilpain, P., Chong, B.F., Chouzenoux, S., Guillevin, L., Du, Y., Zhou, X.J., Lin, F., Fairhurst, A.M., Boudreaux, C., *et al.* (2010). Dysregulated expression of CXCR4/CXCL12 in subsets of patients with systemic lupus erythematosus. *Arthritis Rheum* 62, 3436-3446;
- [1108] Wang, F., Li, S., Zhao, Y., Yang, K., Chen, M., Niu, H., Yang, J., Luo, Y., Tang, W.,

- and Sheng, M. (2016). Predictive role of the overexpression for CXCR4, C-Met, and VEGF-C among breast cancer patients: A meta-analysis. *Breast* 28, 45-53;
- [1109] Wang, L.P., Jin, J., Lv, F.F., Cao, J., Zhang, J., Wang, B.Y., Shao, Z.M., Hu, X.C., and Wang, Z.H. (2015a). Norepinephrine attenuates CXCR4 expression and the corresponding invasion of MDA-MB-231 breast cancer cells via beta2-adrenergic receptors. *Eur Rev Med Pharmacol Sci* 19, 1170-1181;
- [1110] Wang, W., Erbe, A.K., Hank, J.A., Morris, Z.S., and Sondel, P.M. (2015b). NK Cell-Mediated Antibody-Dependent Cellular Cytotoxicity in Cancer Immunotherapy. *Front Immunol* 6, 368;
- [1111] Wang, Z., Ma, Q., Liu, Q., Yu, H., Zhao, L., Shen, S., and Yao, J. (2008). Blockade of SDF-1/CXCR4 signalling inhibits pancreatic cancer progression *in vitro* via inactivation of canonical Wnt pathway. *Br J Cancer* 99, 1695-1703;
- [1112] Wang, Z., Yu, D., Wang, M., Wang, Q., Kouznetsova, J., Yang, R., Qian, K., Wu, W., Shuldiner, A., Sztalryd, C., *et al.* (2015c). Elabela-apelin receptor signaling pathway is functional in mammalian systems. *Sci Rep* 5, 8170;
- [1113] Watts, A.O., van Lipzig, M.M., Jaeger, W.C., Seeber, R.M., van Zwam, M., Vinet, J., van der Lee, M.M., Siderius, M., Zaman, G.J., Boddeke, H.W., *et al.* (2013). Identification and profiling of CXCR3-CXCR4 chemokine receptor heteromer complexes. *Br J Pharmacol* 168, 1662-1674;
- [1114] Watts, S.W. (2010). Endothelin receptors: what's new and what do we need to know? *Am J Physiol Regul Integr Comp Physiol* 298, R254-260;
- [1115] White, J.H., Wise, A., Main, M.J., Green, A., Fraser, N.J., Disney, G.H., Barnes, A.A., Emson, P., Foord, S.M., Marshall, F.H. (1998). Heterodimerization is required for the formation of a functional GABA(B) receptor. *Nature* 396, 679-682;
- [1116] Wong, D., Kandagatla, P., Korz, W., and Chinni, S.R. (2014). Targeting CXCR4 with CTCE-9908 inhibits prostate tumor metastasis. *BMC Urol* 14, 12;
- [1117] Woodward, D.F., Jones, R.L., and Narumiya, S. (2011). International Union of Basic and Clinical Pharmacology. LXXXIII: classification of prostanoid receptors, updating 15 years of progress. *Pharmacol Rev* 63, 471-538;
- [1118] Wreggett, K.A., and Wells, J.W. (1995). Cooperativity manifest in the binding properties of purified cardiac muscarinic receptors. *J Biol Chem* 270, 22488-22499;
- [1119] Wu, L., Chen, L., and Li, L. (2017). Apelin/APJ system: A novel promising therapy target for pathological angiogenesis. *Clin Chim Acta* 466, 78-84;
- [1120] Wurth, R., Florio, T. (2016). Subventricular zone microenvironment protects glioblastoma cells from radiotherapy cytotoxicity: role of the chemokine CXCL12. *Translational Cancer Research* 5(Suppl 6):, S1098-S1101;
- [1121] Xie, H., and He, S.H. (2005). Roles of histamine and its receptors in allergic and inflammatory bowel diseases. *World J Gastroenterol* 11, 2851-2857;

- [1122] Xu, H., Fu, S., Chen, Q., Gu, M., Zhou, J., Liu, C., Chen, Y., and Wang, Z. (2017). The function of oxytocin: a potential biomarker for prostate cancer diagnosis and promoter of prostate cancer. *Oncotarget* 8, 31215-31226;
- [1123] Xu, L., Duda, D.G., di Tomaso, E., Ancukiewicz, M., Chung, D.C., Lauwers, G.Y., Samuel, R., Shellito, P., Czito, B.G., Lin, P.C., *et al.* (2009). Direct evidence that bevacizumab, an anti-VEGF antibody, up-regulates SDF1alpha, CXCR4, CXCL6, and neuropilin 1 in tumors from patients with rectal cancer. *Cancer Res* 69, 7905-7910;
- [1124] Yagami, T., Koma, H., and Yamamoto, Y. (2016). Pathophysiological Roles of Cyclooxygenases and Prostaglandins in the Central Nervous System. *Mol Neurobiol* 53, 4754-4771;
- [1125] Yang, D., Koupenova, M., McCrann, D.J., Kopeikina, K.J., Kagan, H.M., Schreiber, B.M., and Ravid, K. (2008). The A2b adenosine receptor protects against vascular injury. *Proc Natl Acad Sci U S A* 105, 792-796;
- [1126] Yang, Q., Zhang, F., Ding, Y., Huang, J., Chen, S., Wu, Q., Wang, Z., Wang, Z., and Chen, C. (2014). Antitumour activity of the recombination polypeptide GST-NT21MP is mediated by inhibition of CXCR4 pathway in breast cancer. *Br J Cancer* 110, 1288-1297;
- [1127] Yang, T., and Du, Y. (2012). Distinct roles of central and peripheral prostaglandin E2 and EP subtypes in blood pressure regulation. *Am J Hypertens* 25, 1042-1049;
- [1128] Zabel, B.A., Lewen, S., Berahovich, R.D., Jaen, J.C., and Schall, T.J. (2011). The novel chemokine receptor CXCR7 regulates trans-endothelial migration of cancer cells. *Mol Cancer* 10, 73;
- [1129] Zatelli, M.C., Ambrosio, M.R., Bondanelli, M., and Uberti, E.C. (2007). Control of pituitary adenoma cell proliferation by somatostatin analogs, dopamine agonists and novel chimeric compounds. *Eur J Endocrinol* 156 Suppl 1, S29-35;
- [1130] Zeelenberg, I.S., Ruuls-Van Stalle, L., and Roos, E. (2003). The chemokine receptor CXCR4 is required for outgrowth of colon carcinoma micrometastases. *Cancer Res* 63, 3833-3839;
- [1131] Zhan, W., Liang, Z., Zhu, A., Kurtkaya, S., Shim, H., Snyder, J.P., and Liotta, D.C. (2007). Discovery of small molecule CXCR4 antagonists. *J Med Chem* 50, 5655-5664;
- [1132] Zhang, W.B., Navenot, J.M., Haribabu, B., Tamamura, H., Hiramatsu, K., Omagari, A., Pei, G., Manfredi, J.P., Fujii, N., Broach, J.R., *et al.* (2002). A point mutation that confers constitutive activity to CXCR4 reveals that T140 is an inverse agonist and that AMD3100 and ALX40-4C are weak partial agonists. *J Biol Chem* 277, 24515-24521;
- [1133] Zhao, H., Guo, L., Zhao, H., Zhao, J., Weng, H., and Zhao, B. (2015). CXCR4 overexpression and survival in cancer: a system review and meta-analysis. *Oncotarget* 6, 5022-5040;
- [1134] Zhou, N., Fan, X., Mukhtar, M., Fang, J., Patel, C.A., DuBois, G.C., and Pomerantz,

- R.J. (2003). Cell-cell fusion and internalization of the CNS-based, HIV-1 co-receptor, APJ. *Virology* 307, 22-36;
- [1135] Zhu, A., Zhan, W., Liang, Z., Yoon, Y., Yang, H., Grossniklaus, H.E., Xu, J., Rojas, M., Lockwood, M., Snyder, J.P., *et al.* (2010). Dipyrimidine amines: a novel class of chemokine receptor type 4 antagonists with high specificity. *J Med Chem* 53, 8556-8568; and
- [1136] Zitzer, H., Honck, H.H., Bachner, D., Richter, D., and Kreienkamp, H.J. (1999). Somatostatin receptor interacting protein defines a novel family of multidomain proteins present in human and rodent brain. *J Biol Chem* 274, 32997-33001.

Claims

- [Claim 1] A method for treatment, amelioration, or prevention of a cancer in a subject having CXCR4-GPCR_x heteromer, the method comprising: administering to the subject a therapeutically effective amount of an inhibitor of a CXCR4-GPCR_x heteromer, wherein: GPCR_x heteromerizes with CXCR4 in the subject, the heteromerization of GPCR_x with CXCR4 is accompanied by enhancement of signaling downstream of CXCR4; and the enhancement of signaling downstream of CXCR4 is suppressed by the inhibitor of the CXCR4-GPCR_x heteromer.
- [Claim 2] The method of claim 1, wherein the heteromerization of CXCR4 with GPCR_x is evaluated by a proximity-based assay.
- [Claim 3] The method of claim 2, wherein the proximity-based assay is selected from the group consisting of bimolecular fluorescence complementation (BiFC), proximity ligation assay (PLA), fluorescence resonance energy transfer (FRET), bioluminescence resonance energy transfer (BRET), cysteine crosslinking, and co-immunoprecipitation.
- [Claim 4] The method of any one of claims 1-3, wherein the heteromerization of CXCR4 with GPCR_x is evaluated by a co-internalization assay.
- [Claim 5] The method of any one of claims 1-4, wherein the enhancement of cell signaling downstream of CXCR4 is evaluated by an intracellular Ca²⁺ assay.
- [Claim 6] The method of any one of claims 1-5, wherein GPCR_x is selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, APLNR, C5AR1, CALCR, CCR5, CHRM1, GALR1, EDNRB, HRH1, MLNR, NTSR1, PTGER2, PTGER3, SSTR2, and TACR3.
- [Claim 7] The method of any one of claims 1-6, wherein the inhibitor of the CXCR4-GPCR_x heteromer is an inhibitor of the CXCR4.
- [Claim 8] The method of claim 7, wherein the inhibitor of CXCR4 is selected from the group consisting of AD-114, AD-114-6H, AD-114-Im7-FH, AD-114-PA600-6H, ALX-0651, ALX40-4C, AMD070 (AMD11070, X4P-001), AMD3100 (plerixafor), AMD3465, ATI 2341, BKT140 (BL-8040; TF14016; 4F-Benzoyl-TN14003), CTCE-9908, CX549, D-[Lys3] GHRP-6, FC122, FC131, GMI-1359, GSK812397, GST-NT21MP, isothioureia-1a, isothioureia-1t (IT1t), KRH-1636, KRH-3955, LY2510924, LY2624587, MSX-122, N-[¹¹C]Methyl-AMD3465,

PF-06747143, POL6326, SDF-1 1-9[P2G] dimer, SDF1 P2G, T134, T140, T22, TC 14012, TG-0054 (Burixafor), USL311, ulocuplumab (MDX1338/BMS-936564), viral macrophage inflammatory protein-II (vMIP-II), WZ811, 12G5, 238D2, 238D4, [⁶⁴Cu]-AMD3100, [⁶⁴Cu]-AMD3465, [⁶⁸Ga]pentixafor, [⁹⁰Y]pentixather, [^{99m}Tc]O₂ - AMD3100, [¹⁷⁷Lu]pentixather, and 508MCl (Compound 26).

- [Claim 9] The method of claim 7, wherein the inhibitor of the CXCR4-GPCR_x heteromer is an antagonist of the CXCR4.
- [Claim 10] The method of claim 9, wherein the antagonist of CXCR4 is selected from the group consisting of ALX40-4C, AMD070 (AMD11070, X4P-001), AMD3100 (plerixafor), AMD3465, ATI 2341, BKT140 (BL-8040; TF14016; 4F-Benzoyl-TN14003), CTCE-9908, CX549,,D-[Lys3] GHRP-6, FC122, FC131, GMI-1359, GSK812397, GST-NT21MP, isothioureia-1a, isothioureia-1t (IT1t), KRH-1636, KRH-3955, LY2510924, MSX-122, N-[¹¹C]Methyl-AMD3465, POL6326, SDF-1 1-9[P2G] dimer, SDF1 P2G, T134, T140, T22, TC 14012, TG-0054 (Burixafor), USL311, viral macrophage inflammatory protein-II (vMIP-II), WZ811, [⁶⁴Cu]-AMD3100, [⁶⁴Cu]-AMD3465, [⁶⁸Ga]pentixafor, [⁹⁰Y]pentixather, [^{99m}Tc]O₂-AMD3100, [¹⁷⁷Lu]pentixather, and 508MCl (Compound 26).
- [Claim 11] The method of claims 7, wherein the inhibitor of the CXCR4-GPCR_x heteromer further comprises an inhibitor of the GPCR_x.
- [Claim 12] The method of claims 11, wherein the inhibitor of GPCR_x is concurrently or sequentially administered with the inhibitor of CXCR4.
- [Claim 13] The method of any one of claims 1-12, wherein the inhibitor of the CXCR4-GPCR_x heteromer is, or comprises a combination of, an antagonist, an inverse agonist, an allosteric modulator, an antibody or binding portion thereof, or a ligand, of GPCR_x.
- [Claim 14] The method of claim 13, wherein the antagonist of GPCR_x is an antagonist of a molecule selected from the group consisting of ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, PTGER2, and TACR3.
- [Claim 15] The method of claim 14, wherein the antagonist of GPCR_x is an antagonist of ADRB2 selected from the group consisting of Alprenolol, atenolol, betaxolol, bupranolol, butoxamine, carazolol, carvedilol, CGP 12177, cicloprolol, ICI 118551, ICYP, labetalol, levobetaxolol, levobunolol, LK 204-545, metoprolol, nadolol, NIHP, NIP, propafenone, propranolol, sotalol, SR59230A, and timolol.

- [Claim 16] The method of claim 15, wherein the antagonist of ADRB2 is Carvedilol.
- [Claim 17] The method of claim 14, wherein the antagonist of GPCR_x is an antagonist of CALCR selected from the group consisting of α -CGRP-(8-37) (human), AC187, CT-(8-32) (salmon), and olcegepant.
- [Claim 18] The method of claim 17, wherein the antagonist of CALCR is AC187.
- [Claim 19] The method of claim 14, wherein the antagonist of GPCR_x is an antagonist of CHRM1 selected from the group consisting of 3-Quinuclidinyl benzilate (QNB), 4-DAMP, aclidinium, AE9C90CB, AFDX384, amitriptyline, AQ-RA 741, atropine, benztropine, biperiden, darifenacin, dicyclomine, dosulepin, ethopropazine, glycopyrrolate, guanylpirenzepine, hexahydrodifenidol, hexahydrosiladifenidol, hexocyclium, himbacine, ipratropium, lithocholylcholine, methoctramine, ML381, muscarinic toxin 1, muscarinic toxin 2, muscarinic toxin 3, N-methyl scopolamine, otenzepad, oxybutynin, *p*-F-HHSiD, pirenzepine, propantheline, (R,R)-quinuclidinyl-4-fluoromethyl-benzilate, scopolamine, silahexocyclium, solifenacin, telenzepine, tiotropium, tolterodine, trihexyphenidyl, tripitramine, UH-AH 37, umeclidinium, and VU0255035.
- [Claim 20] The method of claim 19, wherein the antagonist of CHRM1 is selected from the group consisting of Oxybutynin, umeclidinium, and VU0255035.
- [Claim 21] The method of claim 14, wherein the antagonist of GPCR_x is an antagonist of EDNRB selected from the group consisting of A192621, ambrisentan, atrasentan, bosentan (RO 470203, Tracleer);, BQ788, IRL 2500, K-8794, macitentan, RES7011, Ro 46-8443, SB209670, SB217242 (enrasentan), TAK 044, and tezosentan (RO610612).
- [Claim 22] The method of claim 21, wherein the antagonist of EDNRB is bosentan.
- [Claim 23] The method of claim 14, wherein the antagonist of GPCR_x is an antagonist of HRH1 selected from the group consisting of (-)-chlorpheniramine, (+)-chlorpheniramine, (-)-*trans*-H₂-PAT, (+)-*cis*-H₂-PAT, (+)-*trans*-H₂-PAT, ($\pm$)-*cis*-H₂-PAT, ($\pm$)-*trans*-H₂-PAT, (R)-cetirizine, (S)-cetirizine, 9-OH-risperidone, A-317920, A-349821, ABT-239, alimemazine, amitriptyline, aripiprazole, arpromidine, asenapine, astemizole, AZD3778, azelastine, BU-E 47, cetirizine, chlorpheniramine, chlorpromazine, ciproxifan, clemastine,

clobenpropit, clozapine, conessine, cyclizine, cyproheptadine, desloratadine, diphenhydramine, dosulepin, doxepin, epinastine, fexofenadine, fluphenazine, fluspirilene, haloperidol, hydroxyzine, impromidine, INCB-38579, JNJ-39758979, ketotifen, loratadine, loxapine, MK-0249, molindone, olanzapine, perphenazine, pimozide, pipamperone, pitolisant, promethazine, pyrilamine, quetiapine, risperidone, sertindole, terfenadine, thioridazine, thiothixene, trifluoperazine, tripeleennamine, triprolidine, ziprasidone, and zotepine.

- [Claim 24] The method of claim 23, wherein the antagonist of HRH1 is cetirizine, pyrilamine, hydroxyzine, or loratadine.
- [Claim 25] The method of claim 14, wherein the antagonist of GPCR_x is an antagonist of MLNR selected from the group consisting of GM-109, MA-2029, and OHM-11526.
- [Claim 26] The method of claim 25, wherein the antagonist of MLNR is MA-2029.
- [Claim 27] The method of claim 14, wherein the antagonist of GPCR_x is an antagonist of NTSR1 selected from the group consisting of Meclinertant, SR48527, SR48692, and SR142948A.
- [Claim 28] The method of claim 27, wherein the antagonist of NTSR1 is Merclinertant.
- [Claim 29] The method of claim 14, wherein the antagonist of GPCR_x is an antagonist of TACR3 selected from the group consisting of [Trp⁷, β-Ala⁸] neurokinin A-(4-10), AZD2624, FK 224, GR138676, GSK 172981, GSK 256471, N',2-diphenylquinoline-4-carbohydrazide 8m, N',2-diphenylquinoline-4-carbohydrazide, osanetant, PD 154740, PD 161182, PD157672, saredutant, SB 218795, SB 222200, SB 235375, SCH 206272, SSR 146977, and talnetant.
- [Claim 30] The method of claim 29, wherein the antagonist of TACR3 is SSR146977.
- [Claim 31] The method of any one of claims 1-30, wherein the inhibitor of the CXCR4-GPCR_x heteromer is a protein-protein interaction (PPI) inhibitor.
- [Claim 32] The method of any one of claims 1-31, wherein the cancer is selected from the group consisting of breast cancer, lung cancer, brain cancer, kidney cancer, pancreatic cancer, ovarian cancer, prostate cancer, melanoma, multiple myeloma, gastrointestinal cancers, renal cell carcinoma, soft tissue sarcomas, hepatocellular carcinoma, stomach cancer, colorectal cancer, esophageal cancer, and leukemia.
- [Claim 33] The method of any one of claims 1-32, wherein the inhibitor of

CXCR4-GPCR α heteromer is administered to the subject in a pharmaceutical composition.

- [Claim 34] A method for assessing response, or potential response, of a subject having CXCR4-GPCR α heteromer to treatment, amelioration, or prevention of a cancer, the method comprising:
obtaining a sample from the subject;
detecting heteromerization of CXCR4 and GPCR α in the sample; and
based at least in part on detection of the heteromerization of CXCR4 and GPCR α , assessing the subject's response, or potential response, to the treatment, amelioration, or prevention of a cancer.
- [Claim 35] A method of claim 34, wherein the heteromerization of CXCR4 and GPCR α is accompanied by enhancement of signaling downstream of CXCR4.
- [Claim 36] The method of claim 35, wherein the enhancement of cell signaling downstream of CXCR4 is evaluated by an intracellular Ca²⁺ assay.
- [Claim 37] The method of any one of claims 34-36, wherein the heteromerization of CXCR4 with GPCR α is evaluated by a proximity-based assay.
- [Claim 38] The method of claim 37, wherein the proximity-based assay is selected from the group consisting of bimolecular fluorescence complementation (BiFC), proximity ligation assay (PLA), fluorescence resonance energy transfer (FRET), bioluminescence resonance energy transfer (BRET), cysteine crosslinking, and co-immunoprecipitation.
- [Claim 39] The method of any one of claims 34-38, wherein the heteromerization of CXCR4 with GPCR α is evaluated by a co-internalization assay.
- [Claim 40] A method for treating cancer in a patient having a cell containing a CXCR4-GPCR α heteromer, the method comprising:
administering to the patient an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer;
wherein:
i) the CXCR4-GPCR α heteromer has enhanced downstream signaling;
and
ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the cancer patient.
- [Claim 41] A method of suppressing enhanced downstream signaling from a CXCR4-GPCR α heteromer in a cell of a patient suffering from cancer, the method comprising:

administering to the patient an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein:

- i) the CXCR4-GPCR_x heteromer has enhanced downstream signaling; and
- ii) the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR_x heteromer in the cancer patient.

[Claim 42]

A pharmaceutical kit for use in treating cancer in a patient having a cell containing a CXCR4-GPCR_x heteromer, the pharmaceutical kit comprising:

an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; wherein the CXCR4-GPCR_x heteromer has enhanced downstream signaling.

[Claim 43]

A pharmaceutical composition for use in treating cancer in a patient having a cell containing a CXCR4-GPCR_x heteromer, the pharmaceutical composition comprising:

- i) an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; and
 - ii) a pharmaceutically acceptable carrier;
- wherein the CXCR4-GPCR_x heteromer has enhanced downstream signaling.

[Claim 44]

A method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR_x heteromer, wherein the CXCR4-GPCR_x heteromer has enhanced downstream signaling, the method comprising:

- 1) determining whether the patient has the cancer cell containing the CXCR4-GPCR_x heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if:

- i) the patient's cancer cell contains said CXCR4-GPCR_x heteromer; or
- ii) a CXCR4-GPCR_x heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR_x heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient

derived cell(s) containing the CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR α heteromer; and

2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer to the cancer patient.

[Claim 45]

A method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer, the method comprising:

1) determining whether the patient's cancer cell contains the CXCR4-GPCR α heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR α heteromer is present in the patient's cancer cell; wherein:

a) the GPCR α of the CXCR4-GPCR α heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and

b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, enzyme-linked immunosorbent assay (ELISA), flow cytometry, RNAseq, qRT-PCR, microarray, or a fluorescent animal assay; and

2) if the patient's cancer cell contains said CXCR4-GPCR α heteromer, then internally administering an inhibitor or a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer to the patient.

[Claim 46]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-45, wherein the CXCR4-GPCR α heteromer has, causes, or produces, the enhanced downstream signaling.

[Claim 47]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-45, wherein the enhanced downstream signaling results from the CXCR4-GPCR α heteromer.

- [Claim 48] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-45, wherein the enhanced downstream signaling results from agonism of the CXCR4-GPCR α heteromer.
- [Claim 49] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-45, wherein the enhanced downstream signaling results from CXCR4 agonism of the CXCR4-GPCR α heteromer.
- [Claim 50] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-45, wherein the enhanced downstream signaling results from GPCR α agonism of the CXCR4-GPCR α heteromer.
- [Claim 51] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-45, wherein the enhanced downstream signaling results from CXCR4 agonism and GPCR α agonism of the CXCR4-GPCR α heteromer.
- [Claim 52] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-51, wherein the enhanced downstream signaling is downstream of the CXCR4, the respective GPCR α , or the CXCR4-GPCR α heteromer.
- [Claim 53] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-51, wherein the enhanced downstream signaling is downstream of the CXCR4.
- [Claim 54] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-51, wherein the enhanced downstream signaling is downstream of the respective GPCR α .
- [Claim 55] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-51, wherein the enhanced downstream signaling is downstream of the CXCR4-GPCR α heteromer.
- [Claim 56] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-51, wherein the enhanced downstream signaling from the CXCR4-GPCR α heteromer is relative to downstream signaling from a CXCR4 protomer or a respective GPCR α protomer in their respective individual protomer context.
- [Claim 57] The method of treating, method of suppressing, pharmaceutical com-

position, or pharmaceutical kit, of any one of claims 40-51, wherein the enhanced downstream signaling from the CXCR4-GPCR α heteromer is relative to downstream signaling from a CXCR4 protomer in an individual protomer context.

[Claim 58] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-51, wherein the enhanced downstream signaling from the CXCR4-GPCR α heteromer is relative to downstream signaling from a respective GPCR α protomer in an individual protomer context.

[Claim 59] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-51, wherein the enhanced downstream signaling from the CXCR4-GPCR α heteromer is relative to downstream signaling from a CXCR4 protomer and a respective GPCR α protomer in their respective individual protomer context.

[Claim 60] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-59, wherein the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the cancer patient.

[Claim 61] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-59, wherein the administered inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the patient's cancer cells.

[Claim 62] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-61, wherein the enhanced downstream signaling from the CXCR4-GPCR α heteromer is determined by an intracellular Ca²⁺ assay.

[Claim 63] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 62, wherein the intracellular Ca²⁺ assay is a calcium mobilization assay.

[Claim 64] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-63, wherein the enhanced downstream signaling is an enhanced amount of calcium mobilization.

[Claim 65] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 64, wherein the enhanced

amount of calcium mobilization from the CXCR4-GPCR α heteromer is a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is greater than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.

[Claim 66] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 64, wherein the enhanced amount of calcium mobilization from the CXCR4-GPCR α heteromer is a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is at least 10% greater, at least 20% greater, at least 30% greater, at least 40% greater, at least 50% greater, at least 75% greater, or at least 90% greater, than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.

[Claim 67] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 64, wherein the enhanced amount of calcium mobilization from the CXCR4-GPCR α heteromer is a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is at least 100% greater than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.

[Claim 68] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 64-67, wherein the enhanced amount of calcium mobilization is determined by an intracellular Ca²⁺ assay.

[Claim 69] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 68, wherein the intracellular Ca²⁺ assay is a calcium mobilization assay.

[Claim 70] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 64-67, wherein the enhanced amount of calcium mobilization is a synergistic amount of calcium mobilization.

[Claim 71]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 70, wherein the synergistic amount of calcium mobilization from the cells containing the CXCR4-GPCR α heteromer is a calcium mobilization amount that, upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist, is at least 10% greater, at least 20% greater, at least 30% greater, at least 40% greater, at least 50% greater, at least 75% greater, or at least 90% greater, than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.

[Claim 72]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-70, wherein the CXCR4-GPCR α heteromer has two or more of the following characteristics:

- 1) the CXCR4-GPCR α heteromer components in a cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allostery, as determined via one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, enzyme-linked immunosorbent assay (ELISA), flow cytometry, RNAseq, qRT-PCR, microarray, or a fluorescent animal assay;
- 2) an enhanced amount of calcium mobilization, such that:
 - a) either CXCR4 or the respective GPCR α in an individual protomer context in a cell, upon co-stimulation with CXCL12 and a respective selective GPCR α agonist, results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; and
 - b) the CXCR4-GPCR α heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; as determined via a calcium mobilization assay; or
- 3) a CXCR4-GPCR α heteromer-selective reagent:
 - i) alters heteromer-specific properties of the CXCR4-GPCR α heteromer

in a patient derived cell;

ii) alters heteromer-specific function of the CXCR4-GPCR_x heteromer in a patient derived cell;

iii) alters heteromer-specific properties of a patient derived cell containing the CXCR4-GPCR_x heteromer; or

iv) decreases cell proliferation of a patient derived cell(s) containing the CXCR4-GPCR_x heteromer.

[Claim 73] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-72, wherein the CXCR4-GPCR_x heteromer components comprise individual protomers CXCR4 and GPCR_x.

[Claim 74] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 72 or claim 73, wherein the cell containing either the CXCR4 or the respective GPCR_x in an individual protomer context comprises, independently:

i) the individual protomer CXCR4 in the absence of the respective individual protomer GPCR_x; or

ii) the respective individual protomer GPCR_x in the absence of the individual protomer CXCR4;

respectively.

[Claim 75] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 72-74, wherein the cell containing the CXCR4 in an individual protomer context comprises said individual protomer CXCR4 in the absence of the respective individual protomer GPCR_x.

[Claim 76] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 72-74, wherein the cell containing the respective GPCR_x in an individual protomer context comprises said respective individual protomer GPCR_x in the absence of the individual protomer CXCR4.

[Claim 77] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-76, wherein the CXCR4-GPCR_x heteromer components in the cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allosterism, as determined via one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, enzyme-linked

immunosorbent assay (ELISA), flow cytometry, RNAseq, qRT-PCR, microarray, or a fluorescent animal assay.

[Claim 78]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 77, wherein the proximity-based assay is, or comprises, resonance energy transfer (RET), bioluminescence RET (BRET), fluorescence RET (FRET), time-resolved fluorescence RET (TR-FRET), antibody-aided FRET, ligand-aided FRET, bimolecular fluorescence complementation (BiFC), or a proximity ligation assay (PLA).

[Claim 79]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-76, wherein the CXCR4-GPCR α heteromer components in the cell colocalize and physically interact, either directly or via intermediate proteins acting as conduits for allostereism, as determined via one or more of the following: a co-internalization assay, bimolecular fluorescence complementation (BiFC), or a proximity ligation assay (PLA).

[Claim 80]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-79, wherein the patient's cancer cell contains the CXCR4-GPCR α heteromer.

[Claim 81]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-80, wherein the CXCR4-GPCR α heteromer exhibits the enhanced amount of calcium mobilization, such that:

- a) either the CXCR4 or the respective GPCR α in an individual protomer context in the cell upon co-stimulation with CXCL12 and a respective selective GPCR α agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist; and
 - b) the CXCR4-GPCR α heteromer exhibits an enhanced calcium mobilization upon co-stimulation with the CXCL12 and the respective selective GPCR α agonist relative to the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist;
- as determined via a calcium mobilization assay.

[Claim 82]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-80, wherein:

- i) the calcium mobilization from the protomer CXCR4 or GPCR α , in

the individual protomer context in the cell, is non-synergistic, as determined via calcium mobilization assay; and

ii) the calcium mobilization from the CXCR4-GPCR_x heteromer in the cell is synergistic, as determined via a calcium mobilization assay.

[Claim 83]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-82, wherein in the individual protomer context:

a) the individual protomer CXCR4 in the cell, in the absence of the respective individual protomer GPCR_x; or

b) the respective individual protomer GPCR_x in the cell, in the absence of the individual protomer CXCR4;

upon co-stimulation with CXCL12 and a respective selective GPCR_x agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay.

[Claim 84]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-82, wherein in the individual protomer context, independently:

a) the individual protomer CXCR4 in the cell, in the absence of the respective individual protomer GPCR_x; and

b) the respective individual protomer GPCR_x in the cell, in the absence of the individual protomer CXCR4;

upon co-stimulation with CXCL12 and a respective selective GPCR_x agonist results in a calcium mobilization amount that is equal to or less than the sum of calcium mobilization amounts resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay.

[Claim 85]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-84, wherein the CXCR4-GPCR_x heteromer upon co-stimulation with the CXCL12 and the respective selective GPCR_x agonist results in a calcium mobilization amount that is greater than the sum of calcium mobilization amounts resulting from single agonist stimulation of said cells with either the CXCL12 or the respective selective GPCR_x agonist, as determined via a calcium mobilization assay.

[Claim 86]

The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 85, wherein the calcium mobi-

lization amount resulting from the co-stimulation of the CXCR4-GPCR α heteromer is an enhanced amount of calcium mobilization, relative to the sum of calcium mobilizations resulting from single agonist stimulation of said CXCR4-GPCR α heteromer, as determined via a calcium mobilization assay.

- [Claim 87] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 86, wherein the enhanced amount of calcium mobilization is at least 10% greater, at least 20% greater, at least 30% greater, at least 40% greater, at least 50% greater, at least 75% greater, or at least 90% greater, than the sum of calcium mobilizations resulting from single agonist stimulation with either the CXCL12 or the respective selective GPCR α agonist, as determined via a calcium mobilization assay.
- [Claim 88] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 86 or claim 87, wherein the enhanced amount of calcium mobilization is a synergistic amount of calcium mobilization.
- [Claim 89] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-71, wherein the method administers a CXCR4-GPCR α heteromer-selective reagent.
- [Claim 90] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 72-89, wherein the CXCR4-GPCR α heteromer-selective reagent:
- i) alters heteromer-specific properties of the CXCR4-GPCR α heteromer in the patient derived cell;
 - ii) alters heteromer-specific function of the CXCR4-GPCR α heteromer in the patient derived cell;
 - iii) alters heteromer-specific properties of the patient derived cell containing the CXCR4-GPCR α heteromer; or
 - iv) decreases cell proliferation of the patient derived cell containing the CXCR4-GPCR α heteromer.
- [Claim 91] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 90, wherein the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific properties of the CXCR4-GPCR α heteromer in the patient derived cell.
- [Claim 92] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 90, wherein the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific

- function of the CXCR4-GPCR α heteromer in the patient derived cell.
- [Claim 93] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 90, wherein the CXCR4-GPCR α heteromer-selective reagent alters heteromer-specific properties of the patient derived cell containing the CXCR4-GPCR α heteromer.
- [Claim 94] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 90, wherein the CXCR4-GPCR α heteromer-selective reagent decreases cell proliferation of the patient derived cell containing the CXCR4-GPCR α heteromer.
- [Claim 95] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 89-94, wherein the CXCR4-GPCR α heteromer-selective reagent is a CXCR4 inhibitor, a GPCR α inhibitor, or a CXCR4-GPCR α heteromer inhibitor.
- [Claim 96] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 89-94, wherein the CXCR4-GPCR α heteromer-selective reagent is a CXCR4 antagonist, a GPCR α antagonist, or a CXCR4-GPCR α heteromer antagonist.
- [Claim 97] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 89-96, wherein the CXCR4-GPCR α heteromer-selective reagent is administered as a pharmaceutical composition.
- [Claim 98] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-88, wherein the method administers an inhibitor selected from the group consisting of: the CXCR4 inhibitor, the GPCR α inhibitor, or the CXCR4-GPCR α heteromer inhibitor.
- [Claim 99] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 98, wherein the inhibitor is administered as a pharmaceutical composition.
- [Claim 100] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 99, wherein the administered inhibitor is the CXCR4 inhibitor.
- [Claim 101] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 99, wherein the administered inhibitor is the GPCR α inhibitor.
- [Claim 102] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 99, wherein the administered inhibitor is the CXCR4-GPCR α heteromer inhibitor.

- position, or pharmaceutical kit, of claim 99, wherein the administered inhibitor is the inhibitor of the CXCR4-GPCR α heteromer.
- [Claim 103] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-88, wherein the method administers the combination of inhibitors selected from the group consisting of: the CXCR4 inhibitor, the GPCR α inhibitor, and the inhibitor of the CXCR4-GPCR α heteromer.
- [Claim 104] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 103, wherein the combination of inhibitors is administered as a pharmaceutical composition.
- [Claim 105] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 103, wherein the combination of inhibitors comprises the CXCR4 inhibitor and the GPCR α inhibitor.
- [Claim 106] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 103, wherein the combination of inhibitors comprises the CXCR4 inhibitor and the inhibitor of the CXCR4-GPCR α heteromer.
- [Claim 107] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 103, wherein the combination of inhibitors comprises the GPCR α inhibitor and the inhibitor of the CXCR4-GPCR α heteromer.
- [Claim 108] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 103-107, wherein the combination of inhibitors are administered sequentially, concurrently, or simultaneously.
- [Claim 109] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 103-108, wherein the combination of inhibitors are administered as a pharmaceutical composition further comprising a pharmaceutically acceptable carrier.
- [Claim 110] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 103-109, wherein the combination of inhibitors are administered as separate pharmaceutical compositions, wherein the separate pharmaceutical compositions independently further comprise a pharmaceutically acceptable carrier.
- [Claim 111] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-110, wherein the CXCR4 inhibitor is an antagonist of CXCR4, an inverse agonist of

CXCR4, a partial antagonist of CXCR4, an allosteric modulator of CXCR4, an antibody of CXCR4, an antibody fragment of CXCR4, or a ligand of CXCR4.

[Claim 112] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-111, wherein the GPCR α inhibitor is an antagonist of GPCR α , an inverse agonist of GPCR α , a partial antagonist of GPCR α , an allosteric modulator of GPCR α , an antibody of GPCR α , an antibody fragment of GPCR α , or a ligand of GPCR α .

[Claim 113] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-112, wherein the inhibitor of the CXCR4-GPCR α heteromer is an antagonist of the CXCR4-GPCR α heteromer, an inverse agonist of the CXCR4-GPCR α heteromer, a partial antagonist of the CXCR4-GPCR α heteromer, an allosteric modulator of the CXCR4-GPCR α heteromer, an antibody of the CXCR4-GPCR α heteromer, an antibody fragment of the CXCR4-GPCR α heteromer, a ligand of the CXCR4-GPCR α heteromer, or a protein-protein interaction (PPI) inhibitor of the CXCR4-GPCR α heteromer.

[Claim 114] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-110, wherein the CXCR4 inhibitor, the GPCR α inhibitor, or the inhibitor of the CXCR4-GPCR α heteromer, is an antibody-drug conjugate.

[Claim 115] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-114, wherein a therapeutically effective amount of the inhibitor of the CXCR4-GPCR α heteromer is administered to the patient.

[Claim 116] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-114, wherein a sub-therapeutically effective amount of the inhibitor of the CXCR4-GPCR α heteromer is administered to the patient.

[Claim 117] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-116, wherein a therapeutically effective amount of the CXCR4 inhibitor is administered to the patient.

[Claim 118] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-116, wherein a sub-therapeutically effective amount of the CXCR4 inhibitor is ad-

- ministered to the patient.
- [Claim 119] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-118, wherein the CXCR4 inhibitor is selected from the group consisting of: AD-114, AD-114-6H, AD-114-Im7-FH, AD-114-PA600-6H, ALX-0651, ALX40-4C, AMD070 (AMD11070, X4P-001), AMD3100 (plerixafor), AMD3465, ATI 2341, BKT140 (BL-8040; TF14016; 4F-Benzoyl-TN14003), CTCE-9908, CX549,,D-[Lys3] GHRP-6, FC122, FC131, GMI-1359, GSK812397, GST-NT21MP, isothioureia-1a, isothioureia-1t (IT1t), KRH-1636, KRH-3955, LY2510924, LY2624587, MSX-122, N-[¹¹C]Methyl-AMD3465, PF-06747143, POL6326, SDF-1 1-9[P2G] dimer, SDF1 P2G, T134, T140, T22, TC 14012, TG-0054 (Burixafor), USL311, ulocuplumab (MDX1338/BMS-936564), viral macrophage inflammatory protein-II (vMIP-II), WZ811, 12G5, 238D2, 238D4, [⁶⁴Cu]-AMD3100, [⁶⁴Cu]-AMD3465, [⁶⁸Ga]pentixafor, [⁹⁰Y]pentixather, [^{99m}Tc]O₂ - AMD3100, [¹⁷⁷Lu]pentixather, and 508MCl (Compound 26).
- [Claim 120] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-119, wherein a therapeutically effective amount of the GPCR_x inhibitor is administered to the patient.
- [Claim 121] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-119, wherein a sub-therapeutically effective amount of the GPCR_x inhibitor is administered to the patient.
- [Claim 122] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-121, wherein the GPCR_x is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3.
- [Claim 123] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is ADRB2 or HRH1.
- [Claim 124] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is ADCYAP1R1.
- [Claim 125] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 124, wherein the

ADCYAP1R1 inhibitor is selected from the group consisting of: M65, Max.d.4, MK-0893, N-stearyl-[Nle¹⁷] neurotensin-(6-11)/VIP-(7-28), PACAP-(6-38), and PG 97-269.

[Claim 126] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is ADORA2B.

[Claim 127] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 126, wherein the ADORA2B inhibitor is selected from the group consisting of:
3-isobutyl-8-pyrrolidinoxanthine, alloxazine, AS16, AS70, AS74, AS94, AS95, AS96, AS99, AS100, AS101, ATL802, BW-A1433, caffeine, CGS 15943, CPX, CSC, CVT-6883, DAX, DEPX, derenofylline, DPCPX, FK-453, I-ABOPX, istradefylline, KF26777, LAS38096, LUF5981, MRE 2029F20, MRE 3008F20, MRS1191, MRS1220, MRS1523, MRS1706, MRS1754, MSX-2, OSIP339391, pentoxifylline, preladenant, PSB-10, PSB-11, PSB36, PSB603, PSB-0788, PSB1115, rolofylline, SCH 58261, SCH442416, ST-1535, theophylline, tonapofylline, vipadenant, xanthine amine congener, XCC, and ZM-241385.

[Claim 128] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is ADORA3.

[Claim 129] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 128-70, wherein the ADORA3 inhibitor is selected from the group consisting of:
ATL802, BW-A1433, caffeine, CGS 15943, CSC, CVT-6883, derenofylline, dexniguldipine, DPCPX, FK-453, flavanone, flavone, galangin, I-ABOPX, istradefylline, KF26777, LAS38096, LUF5981, MRE 2029F20, MRE 3008F20, MRE 3010F20, MRS1041, MRS1042, MRS1067, MRS1088, MRS1093, MRS1097, MRS1177, MRS1186, MRS1191, MRS1191, MRS1220, MRS1476, MRS1486, MRS1505, MRS1523, MRS1754, MRS928, MSX-2, nicardipine, preladenant, PSB-10, PSB-11, PSB36, PSB603, PSB1115, rolofylline, sakuranetin, SCH 58261, SCH442416, ST-1535, theophylline, tonapofylline, vipadenant, visnagin, VUF5574, VUF8504, VUF8507, xanthine amine congener, and ZM-241385.

[Claim 130] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein

the GPCR_x is ADRB₂.

- [Claim 131] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 130, wherein the ADRB₂ inhibitor is selected from the group consisting of:
Alprenolol, atenolol, betaxolol, bupranolol, butoxamine, carazolol, carvedilol, CGP 12177, cicloprolol, ICI 118551, ICYP, labetalol, levo-betaxolol, levobunolol, LK 204-545, metoprolol, nadolol, NIHP, NIP, propafenone, propranolol, sotalol, SR59230A, and timolol.
- [Claim 132] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is C5AR₁.
- [Claim 133] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 132, wherein the C5AR₁ inhibitor is selected from the group consisting of:
A8^{Δ71-73}, AcPhe-Orn-Pro-D-Cha-Trp-Arg, avacopan, C089, CHIPS, DF2593A, JPE1375, L-156,602, NDT9520492, *N*-methyl-Phe-Lys-Pro-D-Cha-Trp-D-Arg-CO₂H, PMX205, PMX53, RPR121154, and W54011.
- [Claim 134] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is CALCR.
- [Claim 135] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 134, wherein the CALCR inhibitor is selected from the group consisting of:
 α -CGRP-(8-37) (human), AC187, CT-(8-32) (salmon), and olcegepant.
- [Claim 136] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is CHRM₁.
- [Claim 137] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 136, wherein the CHRM₁ inhibitor is selected from the group consisting of:
3-Quinuclidinyl benzilate (QNB), 4-DAMP, acridinium, AE9C90CB, AFDX384, amitriptyline, AQ-RA 741, atropine, benztropine, biperiden, darifenacin, dicyclomine, dosulepin, ethopropazine, glycopyrrolate, guanylpirenzepine, hexahydrodifenidol, hexahydrosiladifenidol, hexocyclium, himbacine, ipratropium, lithocholylcholine, methoctramine, ML381, muscarinic toxin 1, muscarinic toxin 2, muscarinic toxin 3, *N*-methyl scopolamine, otenzepad, oxybutynin, *p*-

F-HHSiD, pirenzepine, propantheline, (R,R)-quinuclidinyl-4-fluoromethyl-benzilate, scopolamine, silahexocyclium, solifenacin, telenzepine, tiotropium, tolterodine, trihexyphenidyl, tripitramine, UH-AH 37, umeclidinium, and VU0255035.

[Claim 138] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is EDNRB.

[Claim 139] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 138, wherein the EDNRB inhibitor is selected from the group consisting of: A192621, ambrisentan, atrasentan, bosentan (RO 470203, Tracleer);, BQ788, IRL 2500, K-8794, macitentan, RES7011, Ro 46-8443, SB209670, SB217242 (enrasentan), TAK 044, and tezosentan (RO610612).

[Claim 140] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is HRH1.

[Claim 141] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 140, wherein the HRH1 inhibitor is selected from the group consisting of: (-)-chlorpheniramine, (+)-chlorpheniramine, (-)-*trans*-H₂-PAT, (+)-*cis*-H₂-PAT, (+)-*trans*-H₂-PAT, (±)-*cis*-H₂-PAT, (±)-*trans*-H₂-PAT, (*R*)-cetirizine, (*S*)-cetirizine, 9-OH-risperidone, A-317920, A-349821, ABT-239, alimemazine, amitriptyline, aripiprazole, arpromidine, asenapine, astemizole, AZD3778, azelastine, BU-E 47, cetirizine, chlorpheniramine, chlorpromazine, ciproxifan, clemastine, clobenpropit, clozapine, conessine, cyclizine, cyproheptadine, desloratadine, diphenhydramine, dosulepin, doxepin, epinastine, fexofenadine, fluphenazine, fluspirilene, haloperidol, hydroxyzine, impromidine, INCB-38579, JNJ-39758979, ketotifen, loratadine, loxapine, MK-0249, molindone, olanzapine, perphenazine, pimozide, pipamperone, pitolisant, promethazine, pyrilamine, quetiapine, risperidone, sertindole, terfenadine, thioridazine, thiothixene, trifluoperazine, tripeleminamine, triprolidine, ziprasidone, and zotepine.

[Claim 142] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is MLNR.

- [Claim 143] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 142, wherein the MLNR inhibitor is selected from the group consisting of:
GM-109, MA-2029, and OHM-11526.
- [Claim 144] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is NTSR1.
- [Claim 145] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 144, wherein the NTSR1 inhibitor is selected from the group consisting of:
Meclinetant, SR48527, SR48692, and SR142948A.
- [Claim 146] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-122, wherein the GPCR_x is TACR3.
- [Claim 147] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 146, wherein the TACR3 inhibitor is selected from the group consisting of:
[Trp⁷, β-Ala⁸] neurokinin A-(4-10), AZD2624, FK 224, GR138676, GSK 172981, GSK 256471, N',2-diphenylquinoline-4-carbohydrazide 8m, N',2-diphenylquinoline-4-carbohydrazide, osanetant, PD 154740, PD 161182, PD157672, saredutant, SB 218795, SB 222200, SB 235375, SCH 206272, SSR 146977, and talnetant.
- [Claim 148] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-147, wherein the respective selective GPCR_x agonist is natural ligand of the respective GPCR_x., respectively.
- [Claim 149] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-148, wherein the method further comprises detecting the CXCR4-GPCR_x heteromer in the cancer patient.
- [Claim 150] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-149, wherein the method further comprises identifying the CXCR4-GPCR_x heteromer in the cancer patient.
- [Claim 151] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-150, wherein the method further comprises:
i) obtaining or having obtained a biological sample from the cancer

patient;

ii) conducting or having conducted a diagnostic assay to determine presence, identity, or presence and identity, of a CXCR4-GPCR $\times$ heteromer in the obtained biological sample from the cancer patient; and

iii) selecting the inhibitor or combination of inhibitors to suppress enhanced downstream signaling from the CXCR4-GPCR $\times$ heteromer.

[Claim 152] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-151, wherein the cancer is selected from the group consisting of: breast cancer, lung cancer, brain cancer, kidney cancer, pancreatic cancer, ovarian cancer, prostate cancer, melanoma, multiple myeloma, gastrointestinal cancers, renal cell carcinoma, soft tissue sarcomas, hepatocellular carcinoma, stomach cancer, colorectal cancer, esophageal cancer, and leukemia.

[Claim 153] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-152, wherein the patient's biological sample is a biological fluid sample.

[Claim 154] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 153, wherein a liquid biopsy is performed on the biological fluid sample.

[Claim 155] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 153 or claim 154, wherein the biological fluid sample is a blood sample, a plasma sample, a saliva sample, a cerebral fluid sample, an eye fluid sample, or a urine sample.

[Claim 156] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-152, wherein the patient's biological sample is a biological tissue sample.

[Claim 157] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 156, wherein a liquid biopsy is performed on the biological tissue sample.

[Claim 158] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of claim 156 or claim 157, wherein the biological tissue sample is an organ tissue sample, a bone tissue sample, or a tumor tissue sample.

[Claim 159] A method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR $\times$ heteromer having enhanced downstream signaling, the method comprising:

1) determining whether the patient has the cancer cell containing the

CXCR4-GPCR α heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if:

- i) the patient's cancer cell contains said CXCR4-GPCR α heteromer; or
 - ii) a CXCR4-GPCR α heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR α heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; and
- 2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer; and
- 3) if the patient does not have a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR α inhibitor as a single inhibitor.

[Claim 160]

A method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR α heteromer having enhanced downstream signaling, the method comprising:

- 1) determining whether the patient has the cancer cell containing the CXCR4-GPCR α heteromer having enhanced downstream signaling by: obtaining or having obtained a biological sample from the patient; and performing or having performed an assay on the biological sample to determine if:
 - i) the patient's cancer cell contains said CXCR4-GPCR α heteromer; or
 - ii) a CXCR4-GPCR α heteromer-selective reagent: alters heteromer-specific properties or function of said CXCR4-GPCR α heteromer in a patient derived cell(s); alters heteromer-specific properties of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; or decreases cell proliferation of a patient derived cell(s) containing said CXCR4-GPCR α heteromer; and
- 2) if the patient has a cancer cell containing said CXCR4-GPCR α heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor

of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; and

3) if the patient does not have a cancer cell containing said CXCR4-GPCR_x heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR_x inhibitor as a single inhibitor;

wherein:

a) progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR_x heteromer is decreased in the range of 5-100% more upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR_x inhibitor as the single inhibitor to said patient;

b) efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR_x inhibitor to the patient having said cancer cell containing the CXCR4-GPCR_x heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor; and/or

c) efficacy of a GPCR_x inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR_x heteromer, relative to efficacy of the GPCR_x inhibitor when administered as a single inhibitor.

[Claim 161]

A method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR_x heteromer, the method comprising:

1) determining whether the patient's cancer cell contains the CXCR4-GPCR_x heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR_x heteromer is present in the patient's cancer cell; wherein:

a) the GPCR_x of the CXCR4-GPCR_x heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and

b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, enzyme-linked immunosorbent assay (ELISA), flow cytometry,

RNAseq, qRT-PCR, microarray, or a fluorescent animal assay; and
 2) if the patient's cancer cell contains said CXCR4-GPCR_x heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; and

3) if the patient's cancer cell does not contain said CXCR4-GPCR_x heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR_x inhibitor as a single inhibitor.

[Claim 162]

A method for treating cancer in a patient having a cancer cell containing a CXCR4-GPCR_x heteromer, the method comprising:

1) determining whether the patient's cancer cell contains the CXCR4-GPCR_x heteromer by: obtaining or having obtained a biological sample from the patient and performing or having performed an assay on the biological sample to determine if said CXCR4-GPCR_x heteromer is present in the patient's cancer cell; wherein:

a) the GPCR_x of the CXCR4-GPCR_x heteromer is selected from the group consisting of: ADCYAP1R1, ADORA2B, ADORA3, ADRB2, C5AR1, CALCR, CHRM1, EDNRB, HRH1, MLNR, NTSR1, and TACR3; and

b) the assay performed on the biological sample is or comprises one or more of the following: a co-internalization assay, a colocalization assay, in situ hybridization, immunohistochemistry, immunoelectron microscopy, a proximity-based assay, a co-immunoprecipitation assay, enzyme-linked immunosorbent assay (ELISA), flow cytometry, RNAseq, qRT-PCR, microarray, or a fluorescent animal assay; and

2) if the patient's cancer cell contains said CXCR4-GPCR_x heteromer, then internally administering to the cancer patient a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR_x, and an inhibitor of the CXCR4-GPCR_x heteromer; and

3) if the patient's cancer cell does not contain said CXCR4-GPCR_x heteromer, then internally administering to the cancer patient either the CXCR4 inhibitor or the GPCR_x inhibitor as a single inhibitor; wherein:

a) progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR_x heteromer is decreased in the range of 5-100% more upon administration of the combination of inhibitors,

relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient;

b) efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor; and/or

c) efficacy of a GPCR α inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor.

[Claim 163] The method of treating of any one of claims 159-162, wherein progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR α heteromer is decreased in the range of 5-100% more upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient.

[Claim 164] The method of treating of any one of claims 159-163, wherein efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor.

[Claim 165] The method of treating of any one of claims 159-164, wherein efficacy of a GPCR α inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor.

[Claim 166] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-165, wherein progression of the cancer in the patient having said cancer cell containing the CXCR4-GPCR α heteromer is decreased in the range of 5-100% more upon administration of the combination of inhibitors, relative to administering the CXCR4 inhibitor or GPCR α inhibitor as the single inhibitor to said patient.

[Claim 167] The method of treating, method of suppressing, pharmaceutical com-

position, or pharmaceutical kit, of any one of claims 40-166, wherein efficacy of a CXCR4 inhibitor is increased in the range of 5-2000% when administered in combination with the GPCR α inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the CXCR4 inhibitor when administered as a single inhibitor.

[Claim 168] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-167, wherein efficacy of a GPCR α inhibitor is increased in the range of 5-2000% when administered in combination with the CXCR4 inhibitor to the patient having said cancer cell containing the CXCR4-GPCR α heteromer, relative to efficacy of the GPCR α inhibitor when administered as a single inhibitor.

[Claim 169] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-168, wherein the method administers a combination of inhibitors selected from the group consisting of: an inhibitor of CXCR4, an inhibitor of GPCR α , and an inhibitor of the CXCR4-GPCR α heteromer.

[Claim 170] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-169, wherein the method administers a combination of a CXCR4 inhibitor and a GPCR α inhibitor.

[Claim 171] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-170, wherein the administering of the combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the cancer patient in the range of between 5-2000 fold, relative to single inhibitor administration.

[Claim 172] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-171, wherein the method administers a CXCR4-GPCR α heteromer inhibitor.

[Claim 173] The method of treating, method of suppressing, pharmaceutical composition, or pharmaceutical kit, of any one of claims 40-172, wherein the administering of the inhibitor or combination of inhibitors suppresses the enhanced downstream signaling from said CXCR4-GPCR α heteromer in the cancer patient in the range of between 5-2000 fold, relative to suppression of downstream signaling from either a CXCR4 protomer or a GPCR α protomer in their re-

spective individual protomer context.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 47

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME


THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 47


NOTE: For additional volumes, please contact the Canadian Patent Office


NOM DU FICHIER / FILE NAME :


NOTE POUR LE TOME / VOLUME NOTE:


[FIG. 1]



























[FIG. 3a]

[FIG. 3b]


[FIG. 4a]

CXCR4-GFP**Pre-stimulation****CXCL12**


[FIG. 4b]

CXCR4-GFP & ADCYAP1R1**Pre-stimulation****VIP**


[FIG. 4c]

CXCR4-GFP & ADORA2B**Pre-stimulation****BAY 60-6583**

[FIG. 4d]

CXCR4-GFP & ADORA3**Pre-stimulation****CGS21680**

[FIG. 4e]


CXCR4-GFP & ADRB2**Pre-stimulation****Formoterol**

[FIG. 4f]

CXCR4-GFP & APLNR

Pre-stimulation

Apelin-13




[FIG. 4g]

CXCR4-GFP & C5AR1

Pre-stimulation

C5a




[FIG. 4h]

CXCR4-GFP & CCR5

Pre-stimulation

CCL2




[FIG. 4i]

CXCR4-GFP & CHRM1

Pre-stimulation

Acetylcholine




[FIG. 4j]


CXCR4-GFP & GALR1

Pre-stimulation


galanin



[FIG. 4k]

CXCR4-GFP & EDNRB**Pre-stimulation****Endothelin 1**

[FIG. 4l]


CXCR4-GFP & HRH1**Pre-stimulation****Histamine**

[FIG. 4m]

CXCR4-GFP & MLNR

Pre-stimulation

Motilin




[FIG. 4n]


CXCR4-GFP & NTSR1

Pre-stimulation


Neurotensin



[FIG. 4o]

CXCR4-GFP & PTGER3**Pre-stimulation****PGE2**

[FIG. 4p]


CXCR4-GFP & SSTR2**Pre-stimulation****SRIF-14**


[FIG. 4q]

CXCR4-GFP & TACR3


Pre-stimulation

Senktide







CXCR4 + ADRB2




[FIG. 5d]

Quantification








[FIG. 6c]




[FIG. 6d]












PTGER2




TACR3




[FIG. 7a] APLNR




[FIG. 7b] CCR5




[FIG. 7c] GALR1 [FIG. 7d] PTGER3




[FIG. 7e]






[Fig. 8a]







[FIG. 8d]
CXCR4-CHRM1











[Fig. 8I]




[FIG. 9]




[FIG. 10a]


CXCR4-ADRB2




[FIG. 10b]

CXCR4-CHRM1






[FIG. 10c (continued)]




[FIG. 11a]

CXCR4-ADRB2




[FIG. 11b]



[FIG. 11c]

CXCR4-HRH1



DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 47

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 47

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE: