



(86) **Date de dépôt PCT/PCT Filing Date:** 2015/04/30
(87) **Date publication PCT/PCT Publication Date:** 2015/11/19
(45) **Date de délivrance/Issue Date:** 2018/07/17
(85) **Entrée phase nationale/National Entry:** 2016/11/09
(86) **N° demande PCT/PCT Application No.:** IB 2015/053174
(87) **N° publication PCT/PCT Publication No.:** 2015/173683
(30) **Priorité/Priority:** 2014/05/14 (US61/993,138)


(51) **Cl.Int./Int.Cl.** **C07D 471/04** (2006.01),
A61K 31/395 (2006.01), **A61K 31/437** (2006.01),
C07D 487/04 (2006.01)

(72) **Inventeurs/Inventors:**
COE, JOTHAM WADSWORTH, US;
JONES, PETER, US;
SABNIS, YOGESH ANIL, BE;
STROHBACH, JOSEPH WALTER, US;
WAKENHUT, FLORIAN MICHEL, DE;
WHITLOCK, GAVIN ALISTAIR, GB;
DEHNHARDT, CHRISTOPH MARTIN, CA

(73) **Propriétaire/Owner:**
PFIZER INC., US

(74) **Agent:** SMART & BIGGAR

(54) **Titre : PYRAZOLOPYRIDINES ET PYRAZOLOPYRIMIDINES**
(54) **Title: PYRAZOLOPYRIDINES AND PYRAZOLOPYRIMIDINES**



(57) **Abrégé/Abstract:**

A compound having the structure: A compound having the structure (I) or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein A and A' are independently C

(57) Abrégé(suite)/Abstract(continued):

or N, where C may be unsubstituted or substituted by halo or C₁-C₆ alkyl; R and R⁰ are independently selected from the group consisting of H, C₁-C₆ alkyl, hydroxy(C₁-C₆ alkyl), phenyl(C₁-C₆ alkyl), and -(CH₂)_n-W, where W is C₃-C₈ cycloalkyl, phenyl, naphthyl, 5- or 6-membered heteroaryl or heterocyclic containing 1-3 N, S and/or O atoms, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' and SR', where R' and R'' are independently C₁-C₆ alkyl or C₃-C₈ cycloalkyl, etc.; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, etc.; or, R and R₀ and the N atom to which they are bonded together form a monocyclic or bicyclic heterocyclic ring which may be unsubstituted or substituted by (a) halo, hydroxy, heteroaryl, C₁-C₆ alkyl, C₁-C₆ alkoxy, etc., or (b) -(CH₂)_n-W, where W is C₃-C₈ cycloalkyl, phenyl, etc.; R¹ is H, halo or cyano; R² and R^{2'} are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms; X is a bond, -CO-, -CONH-, -SO₂-, -SONH-, or -(CH₂)_m-; R³ is H, C₁-C₄ alkyl, phenyl, naphthyl, 5- or 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic, etc., or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent -Y-R⁴ and/or 1-4 substituents each independently selected from R⁵; with the proviso that when X is -CO- or -SO₂-, R³ is not H; Y is a bond, -(CH₂)_m- or -O-; R⁴ is (a) H, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, oxo, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, etc.; (b) phenyl or naphthyl, said phenyl and naphthyl being optionally substituted with 1-5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, cyano, -OR⁶, -NR⁷R⁸, etc.; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, etc.; R⁶ is H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl, etc.; R⁷ and R⁸ are each independently H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C₁-C₆ alkyl is optionally substituted by C₃-C₈ cycloalkyl, halo, etc., and said heterocyclic ring being optionally substituted by one or more C₁-C₆ alkyl or C₃-C₈ cycloalkyl groups; R⁹ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl; and, m and n are independently 0, 1, 2 or 3. The invention also relates to pharmaceutically acceptable salts of these compounds and to pharmaceutically acceptable solvates thereof; to compositions containing such compounds; and to the uses of such compounds in the treatment of various diseases, particularly asthma and COPD.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

WIPO | PCT

(10) International Publication Number
WO 2015/173683 A1(43) International Publication Date
19 November 2015 (19.11.2015)

(51) International Patent Classification:

C07D 471/04 (2006.01) A61K 31/4162 (2006.01)
C07D 487/04 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/IB2015/053174

(22) International Filing Date:

30 April 2015 (30.04.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/993,138 14 May 2014 (14.05.2014) US

(71) Applicant: **PFIZER INC.** [US/US]; 235 East 42nd Street, New York, New York 10017 (US).

(72) Inventors: **COE, Jotham Wadsworth**; 8 Bush Hill Drive, Niantic, Connecticut 06357 (US). **DEHNHARDT, Christoph Martin**; 4400 Buchanan Street, Apt. 1803, Burnaby, British Columbia V5C 0E3 (CA). **JONES, Peter**; 61 Henderson Street, Arlington, Massachusetts 02474 (US). **SABNIS, Yogesh Anil**; Avenue de la Colombie 12, B-1000 Brussels (BE). **STROHBACH, Joseph Walter**; 114 Crockett Creek Drive, Wentzville, Missouri 63385 (US).

WAKENHUT, Florian Michel; Am Hang 40, 34212 Melsungen (DE). **WHITLOCK, Gavin Alistair**; 165 Dover Road, Sandwich Kent CT130DD (GB).


(74) Agent: **KLEIMAN, Gabriel L.**; 235 East 42nd Street, MS 235/9/64, New York, NY 10017 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on next page]

(54) Title: PYRAZOLOPYRIDINES AND PYRAZOLOPYRIMIDINES



(57) Abstract: A compound having the structure: A compound having the structure (I) or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein A and A' are independently C or N, where C may be unsubstituted or substituted by halo or C₁-C₆ alkyl; R and R⁰ are independently selected from the group consisting of H, C₁-C₆ alkyl, hydroxy(C₁-C₆ alkyl), phenyl(C₁-C₆ alkyl), and -(CH₂)_n-W, where W is C₃-C₈ cycloalkyl, phenyl, naphthyl, 5- or 6-membered heteroaryl or heterocyclic containing 1 -3 N, S and/or O atoms, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' and SR', where R' and R'' are independently C₁-C₆ alkyl or C₃-C₈ cycloalkyl, etc.; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, etc.; or, R and R₀ and the N atom to which they are bonded together form a monocyclic or bicyclic heterocyclic ring which may be unsubstituted or substituted by (a) halo, hydroxy, heteroaryl, C₁-C₆ alkyl, C₁-C₆ alkoxy, etc., or (b) -(CH₂)_n-W, where W is C₃-C₈cycloalkyl, phenyl, etc.; R¹ is H, halo or cyano; R² and R^{2'} are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms; X is a bond, -CO-, -CONH-, -SO₂-, -SONH-, or -(CH₂)_m; R³ is H, C₁-C₄ alkyl, phenyl, naphthyl, 5- or 6-membered heteroaryl or heterocyclic containing 1 -3 N atoms, a 5-membered heteroaryl or heterocyclic, etc., or (c) 2 O or S atoms and 0-2 N atoms; wherein each of

[Continued on next page]

WO 2015/173683 A1

SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Declarations under Rule 4.17:

- *as to the identity of the inventor (Rule 4.17(i))*
- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

- *with international search report (Art. 21(3))*

said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent -Y-R⁴ and/or 1 -4 substituents each independently selected from R⁵; with the proviso that when X is -CO- or -SO₂-, R³ is not H; Y is a bond, -(CH₂)_m- or -O-; R⁴ is (a) H, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, oxo, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, etc.; (b) phenyl or naphthyl, said phenyl and naphthyl being optionally substituted with 1 -5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, cyano, -OR⁶, -NR⁷R⁸, etc.; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, etc.; R⁶ is H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl, etc.; R⁷ and R⁸ are each independently H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1 -2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C₁-C₆ alkyl is optionally substituted by C₃-C₈ cycloalkyl, halo, etc., and said heterocyclic ring being optionally substituted by one or more C₁-C₆ alkyl or C₃-C₈ cycloalkyl groups; R⁹ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl; and, *m* and *n* are independently 0, 1, 2 or 3. The invention also relates to pharmaceutically acceptable salts of these compounds and to pharmaceutically acceptable solvates thereof; to compositions containing such compounds; and to the uses of such compounds in the treatment of various diseases, particularly asthma and COPD.

PYRAZOLOPYRIDINES AND PYRAZOLOPYRIMIDINES

Field of the Invention

The present invention relates to pyrazolopyridines and pyrazolopyrimidines,
5 pharmaceutical compositions comprising such compounds and their use as
medicaments. More particularly, the present invention provides 6-phenyl-1H-
pyrazolopyridines derivatives and 6-phenyl-1H-pyrazolopyrimidines derivatives which
are Janus Kinase (JAK) inhibitors and useful for the treatment of allergic and respiratory
conditions, particularly chronic obstructive pulmonary disease.

10

Background

Chronic obstructive pulmonary disease (COPD) is the fourth leading cause of
death in the US and is characterized by airflow obstruction that is not fully reversible
with bronchodilators. The airflow limitation is usually progressive and is associated with
an abnormal inflammatory response of the lungs to noxious particles or gases, primarily
15 cigarette smoke. Symptoms are typically breathing-related (e.g. chronic cough,
exertional dyspnea, expectoration and wheeze). Patients experience periods of stable
disease interspersed with inflammatory exacerbations resulting in acute decline in lung
function and often hospitalization.

Current treatment guidelines recommend bronchodilators as the mainstay of
20 COPD drug treatment. However, anti-inflammatory inhaled corticosteroids (ICS) and
bronchodilator/inhaled corticosteroid combination products, are extensively used.
Whilst inhaled corticosteroids do provide some benefits with respect to short term lung
function improvements and exacerbation frequency, they do not address the
corticosteroid-refractory inflammation which is characteristic of this disease and thought
25 to play a key role in disease progression. There is a clear medical need for anti-
inflammatory therapies in COPD that will address the chronic inflammatory component
of the disease and ultimately provide symptomatic relief, a reduction in exacerbation
frequency and an amelioration of exacerbation severity.

The Janus kinase (JAK) family of receptor associated tyrosine kinases, JAK 1,
30 JAK 2, JAK 3 and tyrosine kinase 2 (TYK2), are involved in signal transduction
associated with a variety of inflammatory cytokines. JAK kinases can function as either
hetero or homo-dimers, phosphorylating STAT transcription factors which regulate
inflammatory gene transcription. Oral JAK 1/JAK 3 inhibitors such as CP-690550 have

shown impressive anti-inflammatory activity in inflammatory diseases such as rheumatoid arthritis and psoriasis.

Many JAK dependent cytokines are thought to play key roles in the pathology of COPD which involves the interplay of multiple inflammatory cells such as T lymphocytes, neutrophils, macrophages and lung epithelium. For example the JAK 1/JAK 3 heterodimer plays a key role in T lymphocyte survival and activation whereas JAK 2 is thought to be critical for regulation of neutrophil activation and apoptosis. JAK 1 and JAK 2 play an important role in IL-13 mediated inflammatory signaling in macrophages, which is thought to link acute inflammatory events to chronic progressive disease. Importantly JAK 1, JAK 2 and TYK 2 also play an important role in signaling mediated by $\text{IFN}\gamma$, a cytokine associated with the chronic inflammation observed in COPD, which modulates the activity of T cells, epithelium and macrophages whilst not being modulated by corticosteroids.


Macrophage phagocytosis of bacteria is impaired in the lungs of COPD patients, potentially in part due to high local $\text{IFN}\gamma$ levels. In vitro studies with isolated patient cells have shown that JAK inhibitors increase phagocytotic rate in the presence of $\text{IFN}\gamma$. Consequently, as well as exerting a direct anti-inflammatory effect, JAK inhibitors may also increase the ability of the lung to maintain a sterile environment.

JAK inhibitors are therefore likely to have utility in the treatment of a range of inflammatory diseases, including lung diseases such as COPD, asthma and pulmonary vascular disease. Compounds which have a broad inhibitory activity across the range of Janus kinases, in particular, are likely to have a potent anti-inflammatory effect. However, such a selectivity profile can also lead to undesirable side-effects in systemically circulating compounds, particularly anemia and neutropenia associated with JAK 2 inhibition. For the treatment of lung diseases, it is therefore particularly favorable to provide JAK inhibitors which can be administered by inhalation and which inhibit Janus kinases locally in the lung without having a significant systemic exposure.

There is thus a need to provide new JAK inhibitors that are potent, selective inhibitors of Janus kinases with appropriate metabolic stability and pharmacokinetic properties, particularly compounds which can be administered by inhalation and are active in lung tissue whilst having poor systemic penetration or high systemic lability.

Summary of the Invention

The present invention provides pyrazolopyridines and pyrazolopyrimidines which are potent and selective inhibitors of Janus kinases, including a compound having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein A and A' are independently C or N, where C may be unsubstituted or substituted by halo or C₁-C₆ alkyl;

R and R⁰ are independently selected from the group consisting of H, C₁-C₆ alkyl, hydroxy(C₁-C₆ alkyl), phenyl(C₁-C₆ alkyl), and -(CH₂)_n-W, where W is C₃-C₈ cycloalkyl, phenyl, naphthyl, 5- or 6-membered heteroaryl or heterocyclic containing 1-3 N, S and/or O atoms, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' and SR', where R' and R'' are independently C₁-C₆ alkyl or C₃-C₈ cycloalkyl, amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, phenyl, heteroaryl, or heterocyclic; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -CONR'R'', NR'COR'', -NR'CONR'R'', -NR'CO₂R'', -(CH₂)_n-SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, hydroxyalkylamino, heterocyclic, or -(CH₂)_n-W', where W' is hydroxy, C₃-C₈ cycloalkyl, phenyl, naphthyl, heterocyclic, or 5- or 6-membered heteroaryl containing 1-3 N, S and/or O atoms;

or, R and R⁰ and the N atom to which they are bonded together form a monocyclic or bicyclic heterocyclic ring which may be unsubstituted or substituted by (a) halo, hydroxy, heteroaryl, C₁-C₆ alkyl, C₁-C₆ alkoxy, (C₁-C₆ alkoxy)C₁-C₆ alkoxy, aryl(C₁-C₆ alkoxy), aryloxy, amino, aminoacyl, C₁-C₆ alkylaminoacyl, arylalkylaminoacyl, di(C₁-C₆ alkyl)aminoacyl, -SO₂-R', -SO₂-NR''-(CH₂)_n-W, -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' is independently amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, C₁-C₆ alkyl or C₃-C₈ cycloalkyl, or (b) -(CH₂)_n-W, where W is C₃-C₈ cycloalkyl, phenyl, naphthyl,

heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N atoms, $-\text{SO}_2\text{-R}'$, $-\text{NHSO}_2\text{-R}'$, $-\text{NR}''\text{SO}_2\text{-R}'$ or SR' , where R' and R'' is independently alkyl or cycloalkyl; wherein each of said phenyl, aryl, or heteroaryl may be unsubstituted or substituted by halo, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, cyano, or hydroxy;

5 R^1 is H, cyano or halo; R^2 and $\text{R}^{2'}$ are independently H, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ alkylthio, or $\text{C}_3\text{-C}_8$ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

X is a bond, $-\text{CO}-$, $-\text{CONH}-$, $-\text{SO}_2-$, $-\text{SONH}-$, or $-(\text{CH}_2)_m--$;

R^3 is H, $\text{C}_1\text{-C}_4$ alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic
10 containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, or an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N atoms or (b) 1 O
15 or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent $-\text{Y-R}^4$ and/or 1-4 substituents each independently selected from R^5 ; with the proviso that when X is $-\text{CO}-$ or $-\text{SO}_2-$, R^3 is not H;

Y is a bond, $-(\text{CH}_2)_m-$ or $-\text{O}-$;

20 R^4 is (a) H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, halo, oxo, $-\text{OR}^6$, $-\text{NR}^7\text{R}^8$, $-\text{SR}^6$, $-\text{SOR}^9$, $-\text{SO}_2\text{R}^9$, $-\text{COR}^6$, $-\text{OCOR}^6$, $-\text{COOR}^6$, $-\text{NR}^6\text{COR}^6$, $-\text{CONR}^7\text{R}^8$, $-\text{NR}^6\text{SO}_2\text{R}^9$, $-\text{SO}_2\text{NR}^7\text{R}^8$, $-\text{NR}^6\text{CONR}^7\text{R}^8$, $-\text{NR}^6\text{COOR}^9$ and $-\text{NR}^6\text{SO}_2\text{NR}^7\text{R}^8$; (b) phenyl or naphthyl, said phenyl and naphthyl being optionally substituted with 1-5 substituents selected from $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, halo, $-\text{CN}$, $-\text{OR}^6$, $-\text{NR}^7\text{R}^8$, $-\text{SR}^6$, $-\text{SOR}^9$, $-\text{SO}_2\text{R}^9$, $-\text{COR}^6$, $-\text{OCOR}^6$,
25 $-\text{COOR}^6$, $-\text{NR}^6\text{COR}^6$, $-\text{CONR}^7\text{R}^8$, $-\text{NR}^6\text{SO}_2\text{R}^9$, $-\text{SO}_2\text{NR}^7\text{R}^8$, $-\text{NR}^6\text{CONR}^7\text{R}^8$, $-\text{NR}^6\text{COOR}^9$ and $-\text{NR}^6\text{SO}_2\text{NR}^7\text{R}^8$; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, halo, oxo, $-\text{OR}^6$, $-\text{NR}^7\text{R}^8$, $-\text{SR}^6$, $-\text{SOR}^9$, $-\text{SO}_2\text{R}^9$, $-\text{COR}^6$, $-\text{OCOR}^6$, $-\text{COOR}^6$,
30 $-\text{NR}^6\text{COR}^6$, $-\text{CONR}^7\text{R}^8$, $-\text{NR}^6\text{SO}_2\text{R}^9$, $-\text{SO}_2\text{NR}^7\text{R}^8$, $-\text{NR}^6\text{CONR}^7\text{R}^8$, $-\text{NR}^6\text{COOR}^9$ and $-\text{NR}^6\text{SO}_2\text{NR}^7\text{R}^8$;


R^5 is $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, halo, cyano, $-\text{OR}^6$, $-\text{NR}^7\text{R}^8$, $-\text{SR}^6$, $-\text{SOR}^9$, $-\text{SO}_2\text{R}^9$, $-\text{COR}^6$, $-\text{OCOR}^6$, $-\text{COOR}^6$, $-\text{NR}^6\text{COR}^6$, $-\text{CONR}^7\text{R}^8$, $-\text{NR}^6\text{SO}_2\text{R}^9$, $-\text{SO}_2\text{NR}^7\text{R}^8$, $-\text{NR}^6\text{CONR}^7\text{R}^8$, $-\text{NR}^6\text{COOR}^9$ or $-\text{NR}^6\text{SO}_2\text{NR}^7\text{R}^8$;

R^6 is H, C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl, said C_1 - C_6 alkyl is optionally substituted by $-NR^7R^8$ or a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, cyano, hydroxy and cyano;

R^7 and R^8 are each independently H, C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C_1 - C_6 alkyl is optionally substituted by C_3 - C_8 cycloalkyl, halo, cyano, hydroxy, amino, $(C_1$ - C_6 alkyl)amino or $di(C_1$ - C_6 alkyl)amino and said heterocyclic ring being optionally substituted by one or more C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl groups;

R^9 is C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl; and, m and n are independently 0, 1, 2 or 3.

The invention also provides a compound of formula (Ia) having the structure:



(Ia)

or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

A, A'' and A''' are independently C or N, where C may be unsubstituted or substituted by halo or C_1 - C_6 alkyl;

R^1 is H, cyano or halo;

R^2 and $R^{2'}$ are independently H, C_1 - C_6 alkyl, cyano, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, or C_3 - C_8 cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

R^3 is H, C_1 - C_4 alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, or

an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N atoms or (b) 1 O or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent -Y-R⁴ and/or 1-4 substituents each independently selected from R⁵;

5 Y is a bond, -(CH₂)_m- or -O-;

R⁴ is (a) H, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, oxo, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸; (b) phenyl or naphthyl, said phenyl and naphthyl being optionally substituted with 1-5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, -CN, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, oxo, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸;

15 R⁵ is C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, cyano, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ or -NR⁶SO₂NR⁷R⁸;

20 R⁶ is H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl, said C₁-C₆ alkyl is optionally substituted by -NR⁷R⁸ or a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, hydroxy and cyano;

25 R⁷ and R⁸ are each independently H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C₁-C₆ alkyl is optionally substituted by C₃-C₈ cycloalkyl, halo, cyano, hydroxy, amino, (C₁-C₆ alkyl)amino or di(C₁-C₆ alkyl)amino and said heterocyclic ring being optionally substituted by one or more C₁-C₆ alkyl or C₃-C₈ cycloalkyl groups;


R⁹ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

30 R¹⁰ is -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, C₁-C₆ alkylamino, di(C₁-C₆

alkyl)amino, heterocyclic, $-(CH_2)_n-W'$, where W' is hydroxy, C_3-C_8 cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C_1-C_6 alkyl, C_1-C_6 alkoxy, aryloxy, $-SO_2-R'$, $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently phenyl, C_1-C_6 alkyl or C_3-C_8 cycloalkyl;

R^{11} and R^{12} are each independently H, hydroxy, halo, cyano, C_1-C_6 alkyl or C_3-C_8 cycloalkyl; and, m and n are independently 0, 1, 2, or 3.

The invention further provides a compound of formula (Ib) having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

A'' and A''' are independently C or N, where C may be unsubstituted or substituted by halo or C_1-C_6 alkyl;

R^1 is H, cyano or halo;

R^2 and R^2 are independently H, C_1-C_6 alkyl, cyano, C_1-C_6 alkoxy, C_1-C_6 alkylthio, or C_3-C_8 cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

R^3 is H, C_1-C_4 alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, or an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N atoms or (b) 1 O or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent $-Y-R^4$ and/or 1-4 substituents each independently selected from R^5 ;

Y is a bond, $-(CH_2)_m-$ or $-O-$;

R^4 is (a) H, C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$; (b) phenyl or naphthyl, said phenyl and naphthyl being optionally substituted with 1-5 substituents selected from C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, $-CN$, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$;

R^5 is C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, cyano, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ or $-NR^6SO_2NR^7R^8$;

R^6 is H, C_1-C_6 alkyl or C_3-C_8 cycloalkyl, said C_1-C_6 alkyl is optionally substituted by $-NR^7R^8$ or a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, hydroxy and cyano;

R^7 and R^8 are each independently H, C_1-C_6 alkyl or C_3-C_8 cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C_1-C_6 alkyl is optionally substituted by C_3-C_8 cycloalkyl, halo, cyano, hydroxy, amino, $(C_1-C_6 \text{ alkyl})\text{amino}$ or $\text{di}(C_1-C_6 \text{ alkyl})\text{amino}$ and said heterocyclic ring being optionally substituted by one or more C_1-C_6 alkyl or C_3-C_8 cycloalkyl groups;


R^9 is C_1-C_6 alkyl or C_3-C_8 cycloalkyl;

R^{10} is $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently hydrogen, C_1-C_6 alkyl, C_3-C_8 cycloalkyl, phenyl, amino, C_1-C_6 alkylamino, $\text{di}(C_1-C_6 \text{ alkyl})\text{amino}$, heterocyclic, $-(CH_2)_n-W'$, where W' is hydroxy, C_3-C_8 cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano,

hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently phenyl, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

R¹¹ and R¹² are each independently H, hydroxy, halo, cyano, C₁-C₆ alkyl or C₃-C₈ cycloalkyl; and, *m* and *n* are independently 0, 1, 2 or 3. In another embodiment, the invention provides the compound of formula Ib wherein R¹⁰ is -NR''SO₂-R' and R' and R'' are both C₁-C₆ alkyl.

The invention also provides a compound of formula (Ic) having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

R¹ is H, cyano or halo;

R² and R² are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

R³ is H, C₁-C₄ alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, or an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N atoms or (b) 1 O or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent -Y-R⁴ and/or 1-4 substituents each independently selected from R⁵;

Y is a bond, -(CH₂)_m- or -O-;

R⁴ is (a) H, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, oxo, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸; (b) phenyl or naphthyl, said phenyl

and naphthyl being optionally substituted with 1-5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, -CN, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸; or (c) a 3 to 8-membered saturated or partially unsaturated
 5 monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, oxo, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸;

10 R⁵ is C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, cyano, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ or -NR⁶SO₂NR⁷R⁸;

R⁶ is H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl, said C₁-C₆ alkyl is optionally substituted by -NR⁷R⁸ or a 3 to 8-membered saturated or partially unsaturated monocyclic
 15 heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, cyano, hydroxy and cyano;

R⁷ and R⁸ are each independently H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-
 20 membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C₁-C₆ alkyl is optionally substituted by C₃-C₈ cycloalkyl, halo, cyano, hydroxy, amino, (C₁-C₆ alkyl)amino or di(C₁-C₆ alkyl)amino and said heterocyclic ring being optionally substituted by one or more C₁-C₆ alkyl or C₃-C₈ cycloalkyl groups;


R⁹ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

25 R¹⁰ is -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, heterocyclic, -(CH₂)_n-W', where W' is hydroxy, C₃-C₈ cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may
 30 be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently phenyl, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

R¹¹ and R¹² are each independently H, hydroxy, halo, cyano, C₁-C₆ alkyl or C₃-C₈ cycloalkyl; and, *m* and *n* are independently 0, 1, 2 or 3. In certain embodiments, the

invention provides a compound of formula Ic wherein R^{10} is $-NR''SO_2-R'$ and R' and R'' are both C_1-C_6 alkyl.

The invention additionally provides the compound of formula (Id) having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

A and A' are independently C or N, where C may be unsubstituted or substituted
10 by C_1-C_6 alkyl;


R and R^0 are independently selected from the group consisting of H, C_1-C_6 alkyl, hydroxy(C_1-C_6 alkyl), phenyl(C_1-C_6 alkyl), and $-(CH_2)_n-W$, where W is C_3-C_8 cycloalkyl, phenyl, naphthyl, 5- or 6-membered heteroaryl or heterocyclic containing 1-3 N, S and/or O atoms, $-SO_2-R'$, $-NHSO_2-R'$, $-NR''SO_2-R'$ and SR' , where R' and R'' are
15 independently C_1-C_6 alkyl or C_3-C_8 cycloalkyl, amino, C_1-C_6 alkylamino, di(C_1-C_6 alkyl)amino, phenyl, heteroaryl, or heterocyclic; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C_1-C_6 alkyl, C_1-C_6 alkoxy, aryloxy, $-SO_2-R'$, $-CONR'R''$, $NR'COR''$, $-NR'CONR'R''$, $-NR'CO_2R''$, $-(CH_2)_n-SO_2-R'$, $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently C_1-C_6 alkyl, C_3-C_8
20 cycloalkyl, phenyl, amino, hydroxyalkylamino, heterocyclic, or $-(CH_2)_n-W'$, where W' is hydroxy, C_3-C_8 cycloalkyl, phenyl, naphthyl, heterocyclic, or 5- or 6-membered heteroaryl containing 1-3 N, S and/or O atoms;

or, R and R^0 and the N atom to which they are bonded together form a
25 monocyclic or bicyclic heterocyclic ring which may be unsubstituted or substituted by (a) halo, hydroxy, heteroaryl, C_1-C_6 alkyl, C_1-C_6 alkoxy, (C_1-C_6 alkoxy) C_1-C_6 alkoxy, aryl(C_1-C_6 alkoxy), aryloxy, amino, aminoacyl, C_1-C_6 alkylaminoacyl, arylalkylaminoacyl, di(C_1-C_6 alkyl)aminoacyl, $-SO_2-R'$, $-SO_2-NR''-(CH_2)_n-W$, $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' is independently amino, C_1-C_6 alkylamino, di(C_1-C_6 alkyl)amino, C_1-C_6 alkyl or
30 C_3-C_8 cycloalkyl, or (b) $-(CH_2)_n-W$, where W is C_3-C_8 cycloalkyl, phenyl, naphthyl,

heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N atoms, $-\text{SO}_2\text{-R}'$, $-\text{NHSO}_2\text{-R}'$, $-\text{NR}''\text{SO}_2\text{-R}'$ or SR' , where R' and R'' is independently alkyl or cycloalkyl; wherein each of said phenyl, aryl, or heteroaryl may be unsubstituted or substituted by halo, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, cyano, or hydroxy;

5 R^1 is H, cyano or halo; R^2 and R^2 are independently H, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ alkylthio, or $\text{C}_3\text{-C}_8$ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms; and, n is 0, 1, 2 or 3.

The invention additionally provides the compound of formula (Ie) having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:


15 A, A', A'' and A''' are independently C or N, where C may be unsubstituted or substituted by halo or $\text{C}_1\text{-C}_6$ alkyl;

R^1 is H, cyano or halo;

R^2 and R^2 are independently H, $\text{C}_1\text{-C}_6$ alkyl, cyano, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ alkylthio, or $\text{C}_3\text{-C}_8$ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

20 R^{10} is $-\text{NHSO}_2\text{-R}'$, $-\text{NR}''\text{SO}_2\text{-R}'$ or SR' where R' and R'' is independently $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_8$ cycloalkyl; and,

R^{11} and R^{12} are each independently H, hydroxy, halo, cyano, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_8$ cycloalkyl. In certain embodiments, the invention provides a compound having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

5 A'' and A''' are independently C or N, where C may be unsubstituted or substituted by halo or C₁-C₆ alkyl;


 R¹ is H, cyano or halo;

 R² and R² are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or
10 more fluorine atoms;

 R¹⁰ is -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, heterocyclic, -(CH₂)_n-W', where W' is hydroxy, C₃-C₈ cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms;
15 wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently phenyl, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

 R¹¹ and R¹² are each independently H, hydroxy, halo, cyano, C₁-C₆ alkyl or C₃-C₈
20 cycloalkyl; and, *m* and *n* are independently 0, 1, 2 or 3.

The also provides the compound of formula (Ig) having the structure:



(lg)


or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

R^1 is H, cyano or halo;

5 R^2 and R^2 are independently H, C_1 - C_6 alkyl, cyano, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, or C_3 - C_8 cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

R^{10} is $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently hydrogen, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, phenyl, amino, C_1 - C_6 alkylamino, di(C_1 - C_6 alkyl)amino, heterocyclic, $-(CH_2)_n-W'$, where W' is hydroxy, C_3 - C_8 cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, aryloxy, $-SO_2-R'$, $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently phenyl, C_1 - C_6 alkyl or C_3 - C_8 cyclo alkyl; and, n is 0, 1, 2 or 3. In certain embodiments, the invention provides the compound of formula lg wherein R^{10} is $-NR''SO_2-R'$ and R' and R'' are both C_1 - C_6 alkyl.

The invention also provides the compound of formula (lh) having the structure:



(lh)

or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

A and A' are independently C or N, where C may be unsubstituted or substituted by halo or C_1 - C_6 alkyl;

25 R and R^0 are independently selected from the group consisting of H, C_1 - C_6 alkyl, hydroxy(C_1 - C_6 alkyl), phenyl(C_1 - C_6 alkyl), and $-(CH_2)_n-W$, where W is C_3 - C_8 cycloalkyl, phenyl, naphthyl, 5- or 6-membered heteroaryl or heterocyclic containing 1-3 N, S and/or O atoms, $-SO_2-R'$, $-NHSO_2-R'$, $-NR''SO_2-R'$ and SR' , where R' and R'' are independently C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl, amino, C_1 - C_6 alkylamino, di(C_1 - C_6 alkyl)amino, phenyl, heteroaryl, or heterocyclic; wherein each of said alkyl, cycloalkyl,

heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -CONR'R'', NR'COR'', -NR'CONR'R'', -NR'CO₂R'', -(CH₂)_n-SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, hydroxyalkylamino, heterocyclic, or -(CH₂)_n-W', where W' is hydroxy, C₃-C₈ cycloalkyl, phenyl, naphthyl, heterocyclic, or 5- or 6-membered heteroaryl containing 1-3 N, S and/or O atoms;

or, R and R⁰ and the N atom to which they are bonded together form a monocyclic or bicyclic heterocyclic ring which may be unsubstituted or substituted by (a) halo, hydroxy, heteroaryl, C₁-C₆ alkyl, C₁-C₆ alkoxy, (C₁-C₆ alkoxy)C₁-C₆ alkoxy, aryl(C₁-C₆ alkoxy), aryloxy, amino, aminoacyl, C₁-C₆ alkylaminoacyl, arylalkylaminoacyl, di(C₁-C₆ alkyl)aminoacyl, -SO₂-R', -SO₂-NR''-(CH₂)_n-W, -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' is independently amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, C₁-C₆ alkyl or C₃-C₈ cycloalkyl, or (b) -(CH₂)_n-W, where W is C₃-C₈ cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N atoms, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR', where R' and R'' is independently alkyl or cycloalkyl; wherein each of said phenyl, aryl, or heteroaryl may be unsubstituted or substituted by halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, cyano, or hydroxy;

R¹ is H, cyano or halo;

R² and R² are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

R³ is H, C₁-C₄ alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, or an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N atoms or (b) 1 O or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent -Y-R⁴ and/or 1-4 substituents each independently selected from R⁵; with the proviso that when X is -CO- or -SO₂-, R³ is not H;

Y is a bond, -(CH₂)_m- or -O-;

R^4 is (a) H, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$; (b) phenyl or naphthyl, said phenyl and naphthyl being optionally substituted with 1-5 substituents selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, $-CN$, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$;


R^5 is C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, $-CN$, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ or $-NR^6SO_2NR^7R^8$;

R^6 is H, C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl, said C_1 - C_6 alkyl is optionally substituted by $-NR^7R^8$ or a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, hydroxy and cyano;

R^7 and R^8 are each independently H, C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C_1 - C_6 alkyl is optionally substituted by C_3 - C_8 cycloalkyl, halo, cyano, hydroxy, amino, $(C_1$ - C_6 alkyl)amino or $di(C_1$ - C_6 alkyl)amino and said heterocyclic ring being optionally substituted by one or more C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl groups;

R^9 is C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl; and, m and n are independently 0, 1, 2 or 3. In certain embodiments, the invention provides the compound of formula Ih wherein R^{10} is $-NR''SO_2-R'$ and R' and R'' are both C_1 - C_6 alkyl.

The invention also provides the compound of formula (Ii) having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

5 R^1 is H, cyano or halo;

R^2 and R^2 are independently H, C_1 - C_6 alkyl, cyano, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, or C_3 - C_8 cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

10 R^3 is H, C_1 - C_4 alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, or an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N atoms or (b) 1 O or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1
15 substituent -Y- R^4 and/or 1-4 substituents each independently selected from R^5 ;

Y is a bond, $-(CH_2)_m-$ or -O-;

20 R^4 is (a) H, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$; (b) phenyl or naphthyl, said phenyl and naphthyl being optionally substituted with 1-5 substituents selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, cyano, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, halo, cyano, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, -

25

OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸;

R⁵ is C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, -CN, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ or -NR⁶SO₂NR⁷R⁸;

R⁶ is H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl, said C₁-C₆ alkyl is optionally substituted by -NR⁷R⁸ or a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, hydroxy and cyano;

R⁷ and R⁸ are each independently H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C₁-C₆ alkyl is optionally substituted by C₃-C₈ cycloalkyl, halo, cyano, hydroxy, amino, (C₁-C₆ alkyl)amino or di(C₁-C₆ alkyl)amino and said heterocyclic ring being optionally substituted by one or more C₁-C₆ alkyl or C₃-C₈ cycloalkyl groups; R⁹ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

R¹⁰ is -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, heterocyclic, -(CH₂)_n-W', where W' is hydroxy, C₃-C₈ cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently phenyl, C₁-C₆ alkyl or C₃-C₈ cycloalkyl; and, m and n are independently 0, 1, 2 or 3.

In certain embodiments, the invention provides a compound of formula Ig wherein R¹⁰ is -NR''SO₂-R' and R' and R'' are both C₁-C₆ alkyl.

In specific embodiments, the invention provides a compound selected from the group consisting of:

4-({2-[ethyl(ethylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)-phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(ethylsulfonyl)amino]benzyl}amino)-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-
[(ethylsulfonyl)(methyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-([2-(4-hydroxyphenyl)ethyl]amino)-N-
5 methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-methylpropyl)amino]-1H-
pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-(2-ethyl-5-fluoro-4-
hydroxyphenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({5-fluoro-2-
[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-fluoro-6-
[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
15 carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-
[ethyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

4-[(cyclopentylmethyl)amino]-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-1H-
20 pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({5-methyl-2-
[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

4-({2-[ethyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-
25 trifluoro-ethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[(ethylsulfonyl)(methyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-
trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-fluoro-2-
[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
30 carboxamide;

4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-
2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-
methylpropyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(cyclopentylmethyl)amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-
N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-fluoro-6-
[methyl(methylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
5 carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-
[methyl(methylsulfonyl)amino]benzyl}-amino)-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({2-
10 [methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({5-methyl-2-
[methyl(methylsulfonyl)amino]-benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(phenylsulfonyl)-
15 amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-{4-[(phenylsulfonyl)amino]-
phenyl}ethyl)-amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[(2-
20 hydroxyethyl)(methylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-
c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-
[methyl(phenylsulfonyl)amino]benzyl}-amino)-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[(2-
25 hydroxyethyl)(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-hydroxy-2-
[methyl(methylsulfonyl)amino]benzyl}-amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
30 carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-hydroxy-2-
[methyl(methylsulfonyl)amino]benzyl}-amino)-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}-amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

5 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[(3-hydroxyphenyl)sulfonyl](methyl)-amino)benzyl]amino]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({4-hydroxy-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 4-({2-[ethyl(methylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(phenylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

15 4-({2-[ethyl(phenylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(methylsulfonyl)amino]-benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

20 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-{{(1R)-1-{2-[methyl(methylsulfonyl)amino]-phenyl}ethyl}amino}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-{{(1S)-1-{2-[methyl(methylsulfonyl)-amino]phenyl}ethyl}amino}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

25 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-[methyl(phenylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({4-hydroxy-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

30 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(phenylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(phenylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(phenylsulfonyl)-amino]pyrazin-2-yl)methyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

N-ethyl-4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(4-[methyl(methylsulfonyl)-amino]pyridin-3-yl)methyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-[methyl(methylsulfonyl)amino]-pyridin-3-yl)methyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-methoxy-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(2-[ethyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-[methyl(methylsulfonyl)-amino]pyridin-3-yl)methyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)-amino]pyridin-2-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-[(2-[methyl(methylsulfonyl)amino]benzyl)amino]-N-(6-methylpyridin-3-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-[(1,3,3-trimethylureido)benzyl]-amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-[(2-(N-methyl-1H-pyrazole-4-sulfonamido)benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(2-N,1-dimethyl-1H-imidazole-4-sulfonamido)benzyl]amido)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[(2-methoxyethyl)sulfonyl]-(methyl)amino)benzyl]amino]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(2-[ethyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-(sulfamoylmethyl)benzyl)-amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-(methyl[6-(morpholin-4-yl)pyridin-3-yl]sulfonyl)amino)benzyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-(methyl[3-(morpholin-4-yl)propyl]sulfonyl)amino)benzyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(5-methyl-2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

5 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino}benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino}benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 4-[(5-chloro-2-[ethyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(5-chloro-2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

15 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(sulfamoyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide ;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((N-(2-hydroxyethyl)sulfamoyl)(methyl)-aminobenzyl)amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

20 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-{6-[(2-hydroxyethyl)amino]pyridin-3-yl}-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

25 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-(6-methylpyridin-3-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

30 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(5-fluoro-2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(2-[ethyl(methylsulfonyl)amino]-5-fluoropyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-methyl-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-((2-(N-ethylethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-((2-(N-ethylmethylsulfonamido)-5-fluorobenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoro-ethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-((5-chloro-2-(N-ethylmethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoro-ethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylethylsulfonamido)benzyl)-amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-(((5-methyl-2-(N-methylmethylsulfonamido)pyridin-3-yl)methyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(methylsulfonyl)amino]-5-methylpyridin-3-yl)methyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-fluoro-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)-phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(methylsulfonyl)amino]-5-methylbenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-((2-(N-methylmethylsulfonamido)benzyl)-amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(N-methylmethylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

- 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-((2-(N-methylphenylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylphenylsulfonamido)-benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 5 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-hydroxy-2-(N-methylmethylsulfon-amido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylmethylsulfon-amido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 10 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-hydroxy-2-(N-methylmethylsulfon-amido)benzyl)amino)-N-(6-((2-hydroxyethyl)amino)pyridin-3-yl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-(((3-(N-methylmethyl-sulfonamido)pyrazin-2-yl)methyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 15 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-(2-hydroxyethyl)methyl-sulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-(2-hydroxyethyl)methylsulfon-amido)benzyl)amino)-N-methyl-1H-pyrazolo[3,4-
- 20 d]pyrimidine-3-carboxamide;
- 4-((2-(N-ethylmethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)-phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 4-((5-fluoro-2-(N-methylmethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 25 4-((2-(N-ethylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 4-((2-(N-methylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 30 4-((2-(N-ethylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;
- 4-((2-(N-methylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(methyl(sulfamoyl)amino)-benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(methyl(N-methylsulfamoyl)amino)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

5 4-[4-(7,8-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]-3-ethylphenol formate;

3-ethyl-4-[4-(3-phenoxyazetidin-1-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate;

3-ethyl-4-{4-[6-(4-methyl-1H-imidazol-1-yl)-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl}phenol formate;

3-ethyl-4-{4-[6-(2-methoxyethoxy)-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl}phenol formate;

1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-3-methylazetidin-3-ol formate;

15 2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-N-[2-(pyrrolidin-1-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide formate;

N-benzyl-2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-1,2,3,4-tetrahydroiso-quinoline-5-carboxamide formate;

4-{4-[7-(benzyloxy)-6-methoxy-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl}-3-ethylphenol formate;

4-[4-(5-chloro-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]-3-ethylphenol formate;

4-chloro-3-({1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]azetidin-3-yl}oxy)benzonitrile formate;

25 3-ethyl-4-[4-(6-fluoro-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate;

3-ethyl-4-[4-(8-methoxy-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate;

30 N-{2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-1,2,3,4-tetrahydroisoquinolin-5-yl}methanesulfonamide formate;

4-(4-{[2-(biphenyl-4-yl)ethyl]amino}-1H-pyrazolo[4,3-c]pyridin-6-yl)-3-ethylphenol formate;

N-[2-({[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide hydrochloride;

1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-N,N-dimethylpyrrolidine-3-sulfonamide (racemic);

N-[2-({[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-3-methylphenyl]-N-methylmethanesulfonamide diethylamine salt;

5 3-ethyl-4-[4-(4-methoxypiperidin-1-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol diethylamine salt;

N-[2-({[2-(3,4-dimethoxyphenyl)ethyl][6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide hydrochloride;

10 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](2-{4-[(methylsulfon-yl)amino]phenyl)ethyl)amino}methyl)phenyl]-N-methylmethanesulfonamide hydrochloride;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-phenyl]-N-methylmethanesulfonamide;

15 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-4-methylphenyl]-N-methylmethanesulfonamide hydrochloride;

N-[4-chloro-2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide hydrochloride;

20 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-3-fluorophenyl]-N-methylmethanesulfonamide hydrochloride;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-phenyl]-N-methylethanesulfonamide hydrochloride;

N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl]ethanesulfonamide hydrochloride;

25 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-propylmethanesulfonamide;

N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl]methanesulfonamide;

30 N-butyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl]methanesulfonamide;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl][2-(morpholin-4-yl)ethyl]amino}methyl)phenyl]-N-methylmethanesulfonamide;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl][2-(morpholin-4-yl)ethyl]amino}methyl)phenyl]-N-methylmethanesulfonamide;

N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)-amino)methyl]-4-methylphenyl]methanesulfonamide;

N-[2-({ethyl[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]-4-methylphenyl]-N-methylmethanesulfonamide;

5 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](propyl)amino)methyl]-4-methylphenyl]-N-methylmethanesulfonamide;

N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino)methyl]phenyl]methanesulfonamide hydrochloride;

10 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino)methyl]phenyl]-N-methylmethanesulfonamide hydrochloride;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl]-N-(2-hydroxyethyl)methanesulfonamide;

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl}-N-methylmethanesulfonamide;

15 N-(2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino)methyl]phenyl)-N-methylmethanesulfonamide;

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]-4-methylphenyl}-N-methylmethanesulfonamide;

20 4-{4-[(cyclopropylmethyl)amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-2-fluoro-5-(2,2,2-trifluoroethyl)phenol;

4-{4-[(2-cyclopropylethyl)amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-2-fluoro-5-(2,2,2-trifluoroethyl)phenol;

2-fluoro-4-{4-[(2-methylpropyl)amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-5-(2,2,2-trifluoroethyl)phenol;

25 4-[4-(butylamino)-1H-pyrazolo[3,4-d]pyrimidin-6-yl]-2-fluoro-5-(2,2,2-trifluoroethyl)phenol;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]amino)methyl]-phenyl]-N-methylmethanesulfonamide hydrochloride;

30 N-(2-(((6-(2-ethyl-4-hydroxy-6-methylphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-phenyl)-N-methylmethanesulfonamide;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-3-(1H-imidazol-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl]-N-methylmethanesulfonamide;

N-[2-({[3-(4,5-dimethyl-1H-imidazol-2-yl)-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl]-N-methylmethanesulfonamide;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-3-(4-methyl-1H-imidazol-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

4-[3-(5-benzyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-4-{[2-(methylsulfonyl)ethyl]-amino}-1H-pyrazolo[4,3-c]pyridin-6-yl]-2-fluoro-5-(2,2,2-trifluoroethyl)phenol;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-3-(1H-pyrazol-1-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(1H-pyrazol-1-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(5-methyl-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-3-(5-methyl-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-[5-(6-methylpyridin-3-yl)-4H-1,2,4-triazol-3-yl]-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

4-(5-{6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl}-4H-1,2,4-triazol-3-yl)piperidine-1-carboxamide;

N-(2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-{5-[1-(pyrrolidin-1-ylacetyl)piperidin-4-yl]-4H-1,2,4-triazol-3-yl}-1H-pyrazolo[3,4-d]pyrimidin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]amino)methyl}-4-hydroxyphenyl)-N-methylmethanesulfonamide;

N-{2-[(3-(5-acetyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[3,4-d]pyrimidin-4-yl]amino)methyl}-4-hydroxyphenyl)-N-methylmethanesulfonamide;

N-[2-(dimethylamino)ethyl]-2-{6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl}-1,4,6,7-tetrahydro-5H-imidazo[4,5-c]pyridine-5-carboxamide; and,

4-(3-(5-benzyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-4-((3-hydroxy-2-methylpropyl)-amino)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)-2-fluoro-5-(2,2,2-trifluoroethyl)phenol.

Preferred embodiments of the invention include:

- 5 4-((2-[(ethylsulfonyl)(methyl)amino]benzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- 4-((2-[ethyl(ethylsulfonyl)amino]benzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- 4-((2-[ethyl(methylsulfonyl)amino]-5-methylbenzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- 10 4-((2-[ethyl(methylsulfonyl)amino]-5-fluorobenzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- 4-((2-[ethyl(methylsulfonyl)amino]benzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- 15 4-((5-chloro-2-[ethyl(methylsulfonyl)amino]benzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-((2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-((5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl)amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- 20 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-((5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- and,
- 25 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-((2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;
- or, a pharmaceutically acceptable salt thereof.

More preferred embodiments of the invention include 4-((2-[(ethylsulfonyl)(methyl)amino]benzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide, 4-((2-[ethyl(ethylsulfonyl)amino]benzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide, 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-((2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide, 4-((2-[ethyl(methylsulfonyl)amino]-5-fluorobenzyl)amino)-6-[5-fluoro-4-hydroxy-

2-(2,2,2-trifluoro-ethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide, 4-({2-[ethyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide, or a pharmaceutically acceptable salt thereof.

5 In other aspects, the invention provides a pharmaceutical composition comprising any pyrazolopyridine and pyrazolopyrimidine compound set forth herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or salt, and a pharmaceutically acceptable excipient.

The present invention also provides a method of treating a disease or condition
10 for which a JAK inhibitor is indicated, in a subject in need of such treatment, comprising administering to the subject a therapeutically effective amount of any compound set forth herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or salt.

The present invention further provides a method of treating a disease or
15 condition selected from allergic rhinitis, nasal congestion, rhinorrhea, perennial rhinitis, nasal inflammation, asthma of all types, chronic obstructive pulmonary disease, chronic or acute bronchoconstriction, chronic bronchitis, small airways obstruction, emphysema, chronic eosinophilic pneumonia, adult respiratory distress syndrome, exacerbation of airways hyper-reactivity consequent to other drug therapy, pulmonary vascular disease,
20 pulmonary arterial hypertension, acute lung injury, bronchiectasis, sinusitis, allergic conjunctivitis, idiopathic pulmonary fibrosis or atopic dermatitis, comprising administering to the subject a therapeutically effective amount of any compound set forth herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or salt.

25 The present invention also provides a method of treating chronic obstructive pulmonary disease, comprising administering to the subject a therapeutically effective amount of any compound set forth herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or salt.

The present invention also provides a method of treating a disease or condition
30 selected from inflammation, neuroinflammation, arthritis, rheumatoid arthritis, spondyloarthropathies, systemic lupus erythematosus arthritis, osteoarthritis, gouty arthritis, pain, fever, pulmonary sarcoidosis, silicosis, cardiovascular disease, atherosclerosis, myocardial infarction, thrombosis, congestive heart failure and cardiac reperfusion injury, cardiomyopathy, stroke, ischaemia, reperfusion injury, brain edema,

brain trauma, neurodegeneration, liver disease, inflammatory bowel disease, Crohn's disease, ulcerative colitis, nephritis, retinitis, retinopathy, macular degeneration, glaucoma, diabetes (type 1 and type 2), diabetic neuropathy, viral and bacterial infection, myalgia, endotoxic shock, toxic shock syndrome, autoimmune disease, osteoporosis, multiple sclerosis, endometriosis, menstrual cramps, vaginitis, candidiasis, cancer, fibrosis, obesity, muscular dystrophy, polymyositis, Alzheimer's disease, skin flushing, eczema, psoriasis, atopic dermatitis and sunburn, comprising administering to the subject a therapeutically effective amount of any compound set forth herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or salt.

The disease or condition for which a JAK inhibitor is indicated is preferably an allergic or respiratory condition such as allergic rhinitis, nasal congestion, rhinorrhea, perennial rhinitis, nasal inflammation, asthma of all types, chronic obstructive pulmonary disease (COPD), chronic or acute bronchoconstriction, chronic bronchitis, small airways obstruction, emphysema, chronic eosinophilic pneumonia, adult respiratory distress syndrome, exacerbation of airways hyper-reactivity consequent to other drug therapy, pulmonary vascular disease (including pulmonary arterial hypertension), acute lung injury, bronchiectasis, sinusitis, allergic conjunctivitis, idiopathic pulmonary fibrosis or atopic dermatitis, particularly asthma or chronic obstructive pulmonary disease, most particularly chronic obstructive pulmonary disease.

Other diseases and conditions of interest are inflammation (including neuroinflammation), arthritis (including rheumatoid arthritis, spondyloarthropathies, systemic lupus erythematosus arthritis, osteoarthritis and gouty arthritis), pain, fever, pulmonary sarcoidosis, silicosis, cardiovascular disease (including atherosclerosis, myocardial infarction, thrombosis, congestive heart failure and cardiac reperfusion injury), cardiomyopathy, stroke, ischaemia, reperfusion injury, brain edema, brain trauma, neurodegeneration, liver disease, inflammatory bowel disease (including Crohn's disease and ulcerative colitis), nephritis, retinitis, retinopathy, macular degeneration, glaucoma, diabetes (including type 1 and type 2 diabetes), diabetic neuropathy, viral and bacterial infection, myalgia, endotoxic shock, toxic shock syndrome, autoimmune disease, osteoporosis, multiple sclerosis, endometriosis, menstrual cramps, vaginitis, candidiasis, cancer, fibrosis, obesity, muscular dystrophy, polymyositis, Alzheimer's disease, skin flushing, eczema, psoriasis, atopic dermatitis and sunburn.

Types of asthma include atopic asthma, non-atopic asthma, allergic asthma, atopic bronchial IgE-mediated asthma, bronchial asthma, essential asthma, true asthma, intrinsic asthma caused by pathophysiologic disturbances, extrinsic asthma caused by environmental factors, essential asthma of unknown or inapparent cause, 5 bronchitic asthma, emphysematous asthma, exercise-induced asthma, allergen induced asthma, cold air induced asthma, occupational asthma, infective asthma caused by bacterial, fungal, protozoal, or viral infection, non-allergic asthma, incipient asthma, wheezy infant syndrome and bronchiolitis.

The treatment of asthma includes palliative treatment for the symptoms and 10 conditions of asthma such as wheezing, coughing, shortness of breath, tightness in the chest, shallow or fast breathing, nasal flaring (nostril size increases with breathing), retractions (neck area and between or below the ribs moves inward with breathing), cyanosis (gray or bluish tint to skin, beginning around the mouth), runny or stuffy nose, and headache.

15 The present invention also provides any of the uses, methods or compositions as defined above wherein the compound of formula (I)-(II), or pharmaceutically acceptable salt thereof, or pharmaceutically acceptable solvate of said compound or salt, is used in combination with another pharmacologically active compound, particularly one of the functionally-defined classes or specific compounds listed below. Generally, the 20 compounds of the combination will be administered together as a formulation in association with one or more pharmaceutically acceptable excipients.

Suitable agents for use in combination therapy with a compound of formula (I)-(II), or pharmaceutically acceptable salt thereof, or pharmaceutically acceptable solvate of said compound or salt, particularly in the treatment of respiratory disease, include: a 25 5-lipoxygenase activating protein (FLAP) antagonist; a leukotriene antagonist (LTRA) such as an antagonist of LTB₄, LTC₄, LTD₄, LTE₄, CysLT₁ or CysLT₂, e.g. montelukast or zafirlukast; a histamine receptor antagonist, such as a histamine type 1 receptor antagonist or a histamine type 2 receptor antagonist, e.g. loratidine, fexofenadine, desloratidine, levocetirizine, methapyrilene or cetirizine; an α 1-adrenoceptor agonist or 30 an α 2-adrenoceptor agonist, e.g. phenylephrine, methoxamine, oxymetazoline or methylnorephrine; a muscarinic M3 receptor antagonist, e.g. tiotropium or ipratropium; a dual muscarinic M3 receptor antagonist/ β 2 agonist; a PDE inhibitor, such as a PDE3 inhibitor, a PDE4 inhibitor or a PDE5 inhibitor, e.g. theophylline, sildenafil, vardenafil, tadalafil, ibudilast, cilomilast or roflumilast; sodium cromoglycate or sodium nedocromil;

a cyclooxygenase (COX) inhibitor, such as a non-selective inhibitor (e.g. AspirinTM or ibuprofen) or a selective inhibitor (e.g. celecoxib or valdecoxib); a glucocorticosteroid, e.g. fluticasone, mometasone, dexamethasone, prednisolone, budesonide, ciclesonide or beclamethasone; an anti-inflammatory monoclonal antibody, e.g. infliximab, adalimumab, 5 tanezumab, ranibizumab, bevacizumab or mepolizumab; a β 2 agonist, e.g. salmeterol, albuterol, salbutamol, fenoterol or formoterol, particularly a long-acting β 2 agonist; an integrin antagonist, e.g. natalizumab; an adhesion molecule inhibitor, such as a VLA-4 antagonist; a kinin B₁ or B₂ receptor antagonist; an immunosuppressive agent, such as an inhibitor of the IgE pathway (e.g. omalizumab) or cyclosporine; a matrix 10 metalloprotease (MMP) inhibitor, such as an inhibitor of MMP-9 or MMP-12; a tachykinin NK₁, NK₂ or NK₃ receptor antagonist; a protease inhibitor, such as an inhibitor of elastase, chymase or cathepsin G; an adenosine A_{2a} receptor agonist; an adenosine A_{2b} receptor antagonist; a urokinase inhibitor; a dopamine receptor agonist (e.g. ropinirole), particularly a dopamine D2 receptor agonist (e.g., bromocriptine); a modulator 15 of the NF κ B pathway, such as an IKK inhibitor; a further modulator of a cytokine signalling pathway such as an inhibitor of JAK kinase, syk kinase, p38 kinase, SPHK-1 kinase, Rho kinase, EGF-R or MK-2; a mucolytic, mucokinetic or anti-tussive agent; an antibiotic; an antiviral agent; a vaccine; a chemokine; an epithelial sodium channel (ENaC) blocker or Epithelial sodium channel (ENaC) inhibitor; a nucleotide receptor 20 agonist, such as a P2Y2 agonist; a thromboxane inhibitor; niacin; a 5-lipoxygenase (5-LO) inhibitor, e.g. Zileuton; an adhesion factor, such as VLAM, ICAM or ELAM; a CRTH2 receptor (DP₂) antagonist; a prostaglandin D₂ receptor (DP₁) antagonist; a haematopoietic prostaglandin D2 synthase (HPGDS) inhibitor; interferon- β ; a soluble human TNF receptor, e.g. Etanercept; a HDAC inhibitor; a phosphoinositide 3-kinase 25 gamma (PI3K γ) inhibitor; a phosphoinositide 3-kinase delta (PI3K δ) inhibitor; a CXCR-1 or a CXCR-2 receptor antagonist; an IRAK-4 inhibitor; and, a TLR-4 or TLR-9 inhibitor, including the pharmaceutically acceptable salts of the specifically named compounds and the pharmaceutically acceptable solvates of said specifically named compounds and salts.

30 Besides being useful for human treatment, compounds of formula (I)-(li) are also useful for veterinary treatment of companion animals, exotic animals and farm animals.

Detailed Description of the Invention

Unless otherwise defined herein, scientific and technical terms used in connection with the present invention have the meanings that are commonly understood
5 by those of ordinary skill in the art.

The phrase “therapeutically effective” is intended to qualify the amount of compound or pharmaceutical composition, or the combined amount of active ingredients in the case of combination therapy. This amount or combined amount will achieve the goal of treating the relevant condition.

10 The term “treatment,” as used herein to describe the present invention and unless otherwise qualified, means administration of the compound, pharmaceutical composition or combination to effect preventative, palliative, supportive, restorative or curative treatment. The term treatment encompasses any objective or subjective improvement in a subject with respect to a relevant condition or disease.

15 The term “preventive treatment,” as used herein to describe the present invention, means that the compound, pharmaceutical composition or combination is administered to a subject to inhibit or stop the relevant condition from occurring in a subject, particularly in a subject or member of a population that is significantly predisposed to the relevant condition.

20 The term “palliative treatment,” as used herein to describe the present invention, means that the compound, pharmaceutical composition or combination is administered to a subject to remedy signs and/or symptoms of a condition, without necessarily modifying the progression of, or underlying etiology of, the relevant condition.

The term “supportive treatment,” as used herein to describe the present
25 invention, means that the compound, pharmaceutical composition or combination is administered to a subject as a part of a regimen of therapy, but that such therapy is not limited to administration of the compound, pharmaceutical composition or combination. Unless otherwise expressly stated, supportive treatment may embrace preventive, palliative, restorative or curative treatment, particularly when the compounds or
30 pharmaceutical compositions are combined with another component of supportive therapy.

The term “restorative treatment,” as used herein to describe the present invention, means that the compound, pharmaceutical composition or combination is administered to a subject to modify the underlying progression or etiology of a condition.

Non-limiting examples include an increase in forced expiratory volume in one second (FEV₁) for lung disorders, decreased rate of a decline in lung function over time, inhibition of progressive nerve destruction, reduction of biomarkers associated and correlated with diseases or disorders, a reduction in relapses, improvement in quality of life, reduced time spent in hospital during an acute exacerbation event and the like.

The term "curative treatment," as used herein to describe the present invention, means that compound, pharmaceutical composition or combination is administered to a subject for the purpose of bringing the disease or disorder into complete remission, or that the disease or disorder is undetectable after such treatment.

The term "selective", when used to describe a functionally-defined receptor ligand or enzyme inhibitor means selective for the defined receptor or enzyme subtype as compared with other receptor or enzyme subtypes in the same family. For instance, a selective PDE5 inhibitor is a compound which inhibits the PDE5 enzyme subtype more potently than any other PDE enzyme subtype. Such selectivity is preferably at least 2 fold (as measured using conventional binding assays), more preferably at least 10 fold, most preferably at least 100 fold.

The term "alkyl", alone or in combination, means an acyclic, saturated hydrocarbon group of the formula C_nH_{2n+1} which may be linear or branched. Examples of such groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl and hexyl. Unless otherwise specified, an alkyl group comprises from 1 to 6 carbon atoms.

The carbon atom content of alkyl and various other hydrocarbon-containing moieties is indicated by a prefix designating a lower and upper number of carbon atoms in the moiety, that is, the prefix C_i-C_j indicates a moiety of the integer "i" to the integer "j" carbon atoms, inclusive. Thus, for example, C₁-C₆ alkyl refers to alkyl of one to six carbon atoms, inclusive.

The term "hydroxy," as used herein, means an OH radical.

Het³ is a saturated or partially saturated (i.e. non aromatic) heterocycle and may be attached via a ring nitrogen atom (when the heterocycle is attached to a carbon atom) or a ring carbon atom (in all cases). Equally, when substituted, the substituent may be located on a ring nitrogen atom (if the substituent is joined through a carbon atom) or a ring carbon atom (in all cases). Specific examples include oxiranyl, aziridinyl, oxetanyl, azetidiny, tetrahydrofuranyl, pyrrolidinyl, tetrahydropyranyl, piperidinyl, 1,4-dioxanyl, morpholinyl, piperazinyl, azepanyl, oxepanyl, oxazepanyl and diazepinyl.

Het³ may be fully saturated or partially unsaturated, i.e. may have one or more degrees of unsaturation but may not be fully aromatic.

Het¹ is an aromatic heterocycle and may be attached via a ring carbon atom (in all cases) or a ring nitrogen atom with an appropriate valency (when the heterocycle is attached to a carbon atom). Equally, when substituted, the substituent may be located on a ring carbon atom (in all cases) or a ring nitrogen atom with an appropriate valency (if the substituent is joined through a carbon atom). Specific examples include thienyl, furanyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl and pyrazinyl.

Het² is an aromatic heterocycle and may be attached via a ring carbon atom (in all cases) or a ring nitrogen atom with an appropriate valency (when the heterocycle is attached to a carbon atom). Equally, when substituted, the substituent may be located on a ring carbon atom (in all cases) or a ring nitrogen atom with an appropriate valency (if the substituent is joined through a carbon atom). Het² is aromatic and is therefore necessarily a fused bicycle. Specific examples include imidazo[2,1-b][1,3]thiazolyl, benzofuranyl, benzothienyl, indolyl, benzimidazolyl, indazolyl, benzotriazolyl, pyrrolo[2,3-b]pyridyl, pyrrolo[2,3-c]pyridyl, pyrrolo[3,2-c]pyridyl, pyrrolo[3,2-b]pyridyl, imidazo[4,5-b]pyridyl, imidazo[4,5-c]pyridyl, pyrazolo[4,3-d]pyridyl, pyrazolo[4,3-c]pyridyl, pyrazolo[3,4-c]pyridyl, pyrazolo[3,4-b]pyridyl, isoindolyl, indazolyl, purinyl, indoliziny, imidazo[1,2-a]pyridyl, imidazo[1,5-a]pyridyl, pyrazolo[1,5-a]pyridyl, pyrrolo[1,2-b]pyridazinyl, imidazo[1,2-c]pyrimidinyl, quinoliny, isoquinoliny, cinnoliny, quinazoliny, quinoxaliny, phthalaziny, 1,6-naphthyridiny, 1,7-naphthyridiny, 1,8-naphthyridiny, 1,5-naphthyridiny, 2,6-naphthyridiny, 2,7-naphthyridiny, pyrido[3,2-d]pyrimidinyl, pyrido[4,3-d]pyrimidinyl, pyrido[3,4-d]pyrimidinyl, pyrido[2,3-d]pyrimidinyl, pyrido[2,3-d]pyrazinyl, pyrido[3,4-b]pyrazinyl, pyrimido[5,4-d]pyrimidinyl, pyrazino[2,3-b]pyrazinyl and pyrimido[4,5-d]pyrimidine.

The term "cycloalkyl" means a means a monocyclic, saturated hydrocarbon group of the formula C_nH_{2n-1}. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl. Unless otherwise specified, a cycloalkyl group comprises from 3 to 8 carbon atoms.

The term "oxo" means a doubly bonded oxygen. The term "alkoxy" means a radical comprising an alkyl radical that is bonded to an oxygen atom, such as a methoxy radical. Examples of such radicals include methoxy, ethoxy, propoxy, isopropoxy, butoxy and tert-butoxy. The term "halo" means, fluoro, chloro, bromo or iodo.

As used herein, the terms “co-administration”, “co-administered” and “in combination with”, referring to a combination of a compound of formula (I)-(II) and one or more other therapeutic agents, includes the following:

- simultaneous administration of such a combination of a compound of formula (I)-(II) and a further therapeutic agent to a patient in need of treatment, when such components are formulated together into a single dosage form which releases said components at substantially the same time to said patient;
- substantially simultaneous administration of such a combination of a compound of formula (I)-(II) and a further therapeutic agent to a patient in need of treatment, when such components are formulated apart from each other into separate dosage forms which are taken at substantially the same time by said patient, whereupon said components are released at substantially the same time to said patient;
- sequential administration of such a combination of a compound of formula (I)-(II) and a further therapeutic agent to a patient in need of treatment, when such components are formulated apart from each other into separate dosage forms which are taken at consecutive times by said patient with a significant time interval between each administration, whereupon said components are released at substantially different times to said patient; and
- sequential administration of such a combination of a compound of formula (I)-(II) and a further therapeutic agent to a patient in need of treatment, when such components are formulated together into a single dosage form which releases said components in a controlled manner.

The term ‘excipient’ is used herein to describe any ingredient other than a compound of formula (I)-(II). The choice of excipient will to a large extent depend on factors such as the particular mode of administration, the effect of the excipient on solubility and stability, and the nature of the dosage form. The term “excipient” encompasses diluent, carrier or adjuvant.

One way of carrying out the invention is to administer a compound of formula (I)-(II) in the form of a prodrug. Thus, certain derivatives of a compound of formula (I)-(II) which may have little or no pharmacological activity themselves can, when administered into or onto the body, be converted into a compound of formula (I)-(II) having the desired activity, for example by hydrolytic cleavage, particularly hydrolytic cleavage promoted by an esterase or peptidase enzyme. Such derivatives are referred to as ‘prodrugs’. Further information on the use of prodrugs may be found in ‘Pro-drugs as

Novel Delivery Systems', Vol. 14, ACS Symposium Series (T. Higuchi and W. Stella) and 'Bioreversible Carriers in Drug Design', Pergamon Press, 1987 (Ed. E. B. Roche, American Pharmaceutical Association). Reference can also be made to Nature Reviews/Drug Discovery, 2008, **7**, 355 and Current Opinion in Drug Discovery and
 5 Development, 2007, **10**, 550.

Prodrugs in accordance with the invention can, for example, be produced by replacing appropriate functionalities present in the compounds of formula (I)-(II) with certain moieties known to those skilled in the art as 'pro-moieties' as described, for example, in 'Design of Prodrugs' by H. Bundgaard (Elsevier, 1985).

10 Thus, a prodrug in accordance with the invention is (a) an ester or amide derivative of a carboxylic acid in a compound of formula (I)-(II); (b) an ester, carbonate, carbamate, phosphate or ether derivative of a hydroxyl group in a compound of formula (I)-(II); (c) an amide, imine, carbamate or amine derivative of an amino group in a compound form formula (I)-(II); (d) a thioester, thiocarbonate, thiocarbamate or sulphide
 15 derivatives of a thiol group in a compound of formula (I)-(II); or (e) an oxime or imine derivative of a carbonyl group in a compound of formula (I)-(II).

Some specific examples of prodrugs in accordance with the invention include:

(i) where a compound of formula (I)-(II) contains a carboxylic acid functionality

20 (-COOH), an ester thereof, such as a compound wherein the hydrogen of the carboxylic acid functionality of the compound of formula (I)-(II) is replaced by C₁-C₈ alkyl (e.g. ethyl) or (C₁-C₈ alkyl)C(=O)OCH₂- (e.g. ^tBuC(=O)OCH₂-);

(ii) where a compound of formula (I)-(II) contains an alcohol functionality (-OH), an ester thereof, such as a compound wherein the hydrogen of the alcohol
 25 functionality of the compound of formula (I)-(II) is replaced by -CO(C₁-C₈ alkyl) (e.g. methylcarbonyl) or the alcohol is esterified with an amino acid;

(iii) where a compound of formula (I)-(II) contains an alcohol functionality (-OH), an ether thereof, such as a compound wherein the hydrogen of the alcohol
 30 functionality of the compound of formula (I)-(II) is replaced by (C₁-C₈ alkyl)C(=O)OCH₂- or -CH₂OP(=O)(OH)₂;

(iv) where a compound of formula (I)-(II) contains an alcohol functionality (-OH), a phosphate thereof, such as a compound wherein the hydrogen of the alcohol
 functionality of the compound of formula (I)-(II) is replaced by -P(=O)(OH)₂ or -P(=O)(ONa)₂ or -P(=O)(O⁻)₂Ca²⁺;

(v) where a compound of formula (I)-(II) contains a primary or secondary amino functionality ($-\text{NH}_2$ or $-\text{NHR}$ where $\text{R} \neq \text{H}$), an amide thereof, for example, a compound wherein, as the case may be, one or both hydrogens of the amino functionality of the compound of formula (I)-(II) is/are replaced by $(\text{C}_1\text{-C}_{10})$ alkanoyl, $-\text{COCH}_2\text{NH}_2$ or the amino group is derivatised with an amino acid;

(vi) where a compound of formula (I)-(II) contains a primary or secondary amino functionality ($-\text{NH}_2$ or $-\text{NHR}$ where $\text{R} \neq \text{H}$), an amine thereof, for example, a compound wherein, as the case may be, one or both hydrogens of the amino functionality of the compound of formula (I)-(II) is/are replaced by $-\text{CH}_2\text{OP}(=\text{O})(\text{OH})_2$.

Certain compounds of formula (I)-(II) may themselves act as prodrugs of other compounds of formula (I)-(II). It is also possible for two compounds of formula (I)-(II) to be joined together in the form of a prodrug. In certain circumstances, a prodrug of a compound of formula (I)-(II) may be created by internally linking two functional groups in a compound of formula (I)-(II), for instance by forming a lactone.

References below to compounds of formula (I)-(II) are taken to include the compounds themselves and prodrugs thereof. The invention includes such compounds of formula (I)-(II) as well as pharmaceutically acceptable salts of such compounds and pharmaceutically acceptable solvates of said compounds and salts. Pharmaceutically acceptable salts of the compounds of formula (I)-(II) include acid addition and base salts.

Suitable acid addition salts are formed from acids which form non-toxic salts. Examples include the acetate, adipate, aspartate, benzoate, besylate, bicarbonate/carbonate, bisulfate/sulfate, borate, camsylate, citrate, cyclamate, edisylate, esylate, formate, fumarate, gluceptate, gluconate, glucuronate, hexafluorophosphate, hibenzate, hydrochloride/chloride, hydrobromide/bromide, hydroiodide/iodide, isethionate, lactate, malate, maleate, malonate, mesylate, methylsulfate, naphthylate, 2-napsylate, nicotinate, nitrate, orotate, oxalate, palmitate, pamoate, phosphate/hydrogen phosphate/dihydrogen phosphate, pyroglutamate, saccharate, stearate, succinate, tannate, tartrate, tosylate, trifluoroacetate, naphthalene-1,5-disulfonic acid and xinofoate salts.

Suitable base salts are formed from bases which form non-toxic salts. Examples include the aluminium, arginine, benzathine, calcium, choline, diethylamine, diolamine, glycine, lysine, magnesium, meglumine, olamine, potassium, sodium, tromethamine and zinc salts.

Hemisalts of acids and bases may also be formed, for example, hemisulfate and hemicalcium salts. For a review on suitable salts, see Handbook of Pharmaceutical Salts: Properties, Selection, and Use by Stahl and Wermuth (Wiley-VCH, 2002).

Pharmaceutically acceptable salts of compounds of formula (I)-(II) may be
5 prepared by one or more of three methods:

- (i) by reacting a compound of formula (I)-(II) with the desired acid or base;
- (ii) by removing an acid- or base-labile protecting group from a suitable precursor of a compound of formula (I)-(II) or by ring-opening a suitable cyclic precursor, for example, a lactone or lactam, using the desired acid or base; or
- 10 (iii) by converting one salt of the compound of formula (I)-(II) to another by reaction with an appropriate acid or base or by means of a suitable ion exchange column.

All three reactions are typically carried out in solution. The resulting salt may precipitate out and be collected by filtration or may be recovered by evaporation of the
15 solvent. The degree of ionisation in the resulting salt may vary from completely ionised to almost non-ionised.

The compounds of formula (I)-(II), and pharmaceutically acceptable salts thereof, may exist in unsolvated and solvated forms. The term 'solvate' is used herein to describe a molecular complex comprising the compound of formula (I)-(II), or a
20 pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable solvent molecules, for example, ethanol. The term 'hydrate' may be employed when said solvent is water.

A currently accepted classification system for organic hydrates is one that defines isolated site, channel, or metal-ion coordinated hydrates - see Polymorphism in
25 Pharmaceutical Solids by K. R. Morris (Ed. H. G. Brittain, Marcel Dekker, 1995). Isolated site hydrates are ones in which the water molecules are isolated from direct contact with each other by intervening organic molecules. In channel hydrates, the water molecules lie in lattice channels where they are next to other water molecules. In metal-ion coordinated hydrates, the water molecules are bonded to the metal ion.

30 When the solvent or water is tightly bound, the complex will have a well-defined stoichiometry independent of humidity. When, however, the solvent or water is weakly bound, as in channel solvates and hygroscopic compounds, the water/solvent content will be dependent on humidity and drying conditions. In such cases, non-stoichiometry will be the norm.

Also included within the scope of the invention are multi-component complexes (other than salts and solvates) wherein the drug and at least one other component are present in stoichiometric or non-stoichiometric amounts. Complexes of this type include clathrates (drug-host inclusion complexes) and co-crystals. The latter are typically
5 defined as crystalline complexes of neutral molecular constituents which are bound together through non-covalent interactions, but could also be a complex of a neutral molecule with a salt. Co-crystals may be prepared by melt crystallization, by recrystallization from solvents, or by physically grinding the components together. *Cf.* Chem. Commun., 17, 1889-1896, by O. Almarsson and M. J. Zaworotko (2004). For a
10 general review of multi-component complexes, see J. Pharm. Sci., 64 (8), 1269-1288, by Halebian (1975).

The compounds of the invention may exist in a continuum of solid states ranging from fully amorphous to fully crystalline. The term 'amorphous' refers to a state in which the material lacks long range order at the molecular level and, depending upon
15 temperature, may exhibit the physical properties of a solid or a liquid. Typically such materials do not give distinctive X-ray diffraction patterns and, while exhibiting the properties of a solid, are more formally described as a liquid. Upon heating, a change from solid to liquid properties occurs which is characterised by a change of state, typically second order ('glass transition'). The term 'crystalline' refers to a solid phase in
20 which the material has a regular ordered internal structure at the molecular level and gives a distinctive X-ray diffraction pattern with defined peaks. Such materials when heated sufficiently will also exhibit the properties of a liquid, but the change from solid to liquid is characterised by a phase change, typically first order ('melting point').

The compounds of formula (I)-(II) may also exist in a mesomorphic state
25 (mesophase or liquid crystal) when subjected to suitable conditions. The mesomorphic state is intermediate between the true crystalline state and the true liquid state (either melt or solution). Mesomorphism arising as the result of a change in temperature is described as 'thermotropic' and that resulting from the addition of a second component, such as water or another solvent, is described as 'lyotropic'. Compounds that have the
30 potential to form lyotropic mesophases are described as 'amphiphilic' and consist of molecules which possess an ionic (such as $-\text{COO}^-\text{Na}^+$, $-\text{COO}^-\text{K}^+$, or $-\text{SO}_3^-\text{Na}^+$) or non-ionic (such as $-\text{N}^+\text{N}(\text{CH}_3)_3$) polar head group. For more information, see Crystals and the Polarizing Microscope by N. H. Hartshorne and A. Stuart, 4th Edition (Edward Arnold, 1970).

Hereinafter all references to compounds of formula (I)-(II) include references to pharmaceutically acceptable salts, solvates, multi-component complexes and liquid crystals thereof and to solvates, multi-component complexes and liquid crystals of pharmaceutically acceptable salts thereof.

5 The compounds of formula (I)-(II) may exhibit polymorphism and/or one or more kinds of isomerism (e.g. optical, geometric or tautomeric isomerism). The compounds of formula (I)-(II) may also be isotopically labelled. Such variation is implicit to the compounds of formula (I)-(II) defined as they are by reference to their structural features and therefore within the scope of the invention.

10 Compounds of formula (I)-(II) containing one or more asymmetric carbon atoms can exist as two or more stereoisomers. Where a compound of formula (I)-(II) contains an alkenyl or alkenylene group, geometric cis/trans (or Z/E) isomers are possible. Where structural isomers are interconvertible via a low energy barrier, tautomeric isomerism ('tautomerism') can occur. This can take the form of proton tautomerism in
15 compounds of formula (I)-(II) containing, for example, an imino, keto, or oxime group, or so-called valence tautomerism in compounds which contain an aromatic moiety. It follows that a single compound may exhibit more than one type of isomerism.

 The pharmaceutically acceptable salts of compounds of formula (I)-(II) may also contain a counterion which is optically active (e.g. d-lactate or l-lysine) or racemic (e.g.
20 dl-tartrate or dl-arginine).

 Cis/trans isomers may be separated by conventional techniques well known to those skilled in the art, for example, chromatography and fractional crystallization.

 Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the
25 racemate (or the racemate of a salt or derivative) using, for example, chiral high pressure liquid chromatography (HPLC). Alternatively, the racemate (or a racemic precursor) may be reacted with a suitable optically active compound, for example, an alcohol, or, in the case where the compound of formula (I)-(II) contains an acidic or basic moiety, a base or acid such as 1-phenylethylamine or tartaric acid. The resulting
30 diastereomeric mixture may be separated by chromatography and/or fractional crystallization and one or both of the diastereoisomers converted to the corresponding pure enantiomer(s) by means well known to a skilled person. Chiral compounds of formula (I)-(II) (and chiral precursors thereof) may be obtained in enantiomerically-enriched form using chromatography, typically HPLC, on an asymmetric resin with a

mobile phase consisting of a hydrocarbon, typically heptane or hexane, containing from 0 to 50% by volume of isopropanol, typically from 2% to 20%, and from 0 to 5% by volume of an alkylamine, typically 0.1% diethylamine. Concentration of the eluate affords the enriched mixture. Chiral chromatography using sub- and supercritical fluids
5 may be employed. Methods for chiral chromatography useful in some embodiments of the present invention are known in the art (see, for example, Smith, Roger M., Loughborough University, Loughborough, UK; Chromatographic Science Series (1998), 75 (Supercritical Fluid Chromatography with Packed Columns), pp. 223-249 and references cited therein). In some relevant examples herein, columns were obtained
10 from Chiral Technologies, Inc, West Chester, Pennsylvania, USA, a subsidiary of Daicel[®] Chemical Industries, Ltd., Tokyo, Japan.

When any racemate crystallizes, crystals of two different types are possible. The first type is the racemic compound (true racemate) referred to above wherein one homogeneous form of crystal is produced containing both enantiomers in equimolar
15 amounts. The second type is the racemic mixture or conglomerate wherein two forms of crystal are produced in equimolar amounts each comprising a single enantiomer. While both of the crystal forms present in a racemic mixture have identical physical properties, they may have different physical properties compared to the true racemate. Racemic mixtures may be separated by conventional techniques known to those skilled in the art.
20 See, for example, Stereochemistry of Organic Compounds by E. L. Eliel and S. H. Wilen (Wiley, 1994).

The present invention includes all pharmaceutically acceptable isotopically-labelled compounds of formula (I)-(II) wherein one or more atoms are replaced by atoms having the same atomic number, but an atomic mass or mass number different from the
25 atomic mass or mass number which predominates in nature. Isotopically-labelled compounds of formula (I)-(II) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples and Preparations using an appropriate isotopically-labelled reagent in place of the non-labelled reagent previously employed. In particular,
30 hydrogen atoms may be replaced by deuterium atoms since such deuterated compounds are sometimes more resistant to metabolism.

Also included within the scope of the invention are active metabolites of compounds of formula (I)-(II), that is, compounds formed *in vivo* upon administration of

the drug, often by oxidation or dealkylation. Some examples of metabolites in accordance with the invention include

(i) where a compound of formula (I)-(II) contains a methyl group, an hydroxymethyl derivative thereof ($-\text{CH}_3 \rightarrow -\text{CH}_2\text{OH}$);

5 (ii) where a compound of formula (I)-(II) contains an alkoxy group, an hydroxy derivative thereof ($-\text{OR} \rightarrow -\text{OH}$);

(iii) where a compound of formula (I)-(II) contains a tertiary amino group, a secondary amino derivative thereof ($-\text{NRR}' \rightarrow -\text{NHR}$ or $-\text{NHR}'$);

(iv) where a compound of formula (I)-(II) contains a secondary amino group, a
10 primary derivative thereof ($-\text{NHR} \rightarrow -\text{NH}_2$);

(v) where a compound of formula (I)-(II) contains a phenyl moiety, a phenol derivative thereof ($-\text{Ph} \rightarrow -\text{PhOH}$); and

(vi) where a compound of formula (I)-(II) contains an amide group, a carboxylic acid derivative thereof ($-\text{CONH}_2 \rightarrow \text{COOH}$).

15 For administration to human patients, the total daily dose of a compound of formula (I)-(II) is typically in the range of 0.01mg to 500mg depending, of course, on the mode of administration. In another embodiment of the present invention, the total daily dose of a compound of formula (I)-(II) is typically in the range of 0.1 mg to 300mg. In yet another embodiment of the present invention, the total daily dose of a compound of
20 formula (I)-(II) is typically in the range of 1mg to 30mg. The total daily dose may be administered in single or divided doses and may, at the physician's discretion, fall outside of the typical range given herein. These dosages are based on an average human subject having a weight of about 65kg to 70kg. The physician will readily be able to determine doses for subjects whose weight falls outside this range, such as infants
25 and the elderly.

In the case of dry powder inhalers and aerosols, the dosage unit is determined by means of a prefilled capsule, blister or pocket or by a system that utilises a gravimetrically fed dosing chamber. Units in accordance with the invention are typically arranged to administer a metered dose or "puff" containing from 1 to 5000 μg of drug.
30 The overall daily dose will typically be in the range 1 μg to 20mg which may be administered in a single dose or, more usually, as divided doses throughout the day.

A compound of formula (I)-(II) can be administered per se, or in the form of a pharmaceutical composition, which, as active constituent contains an efficacious dose

of at least one compound of the invention, in addition to customary pharmaceutically innocuous excipients and/or additives.

Pharmaceutical compositions suitable for the delivery of compounds of the present invention and methods for their preparation will be readily apparent to those skilled in the art. Such compositions and methods for their preparation may be found, for example, in Remington's Pharmaceutical Sciences, 19th Edition (Mack Publishing Company, 1995).

Compounds of formula (I)-(II) may be administered orally. Oral administration may involve swallowing, so that the compound enters the gastrointestinal tract, or buccal or sublingual administration may be employed by which the compound enters the blood stream directly from the mouth. Formulations suitable for oral administration include solid formulations such as tablets, capsules containing particulates, liquids, or powders, lozenges (including liquid-filled), chews, multi- and nano-particulates, gels, solid solution, liposome, films, ovules, sprays and liquid formulations.

Liquid formulations include suspensions, solutions, syrups and elixirs. Such formulations may be employed as fillers in soft or hard capsules and typically comprise a carrier, for example, water, ethanol, polyethylene glycol, propylene glycol, methylcellulose, or a suitable oil, and one or more emulsifying agents and/or suspending agents. Liquid formulations may also be prepared by the reconstitution of a solid, for example, from a sachet.

Compounds of formula (I)-(II) may also be used in fast-dissolving, fast-disintegrating dosage forms such as those described in Expert Opinion in Therapeutic Patents, 11 (6), 981-986, by Liang and Chen (2001).

For tablet dosage forms, depending on dose, the drug may make up from 1 weight % to 80 weight % of the dosage form, more typically from 5 weight % to 60 weight % of the dosage form. In addition to the drug, tablets generally contain a disintegrant. Examples of disintegrants include sodium starch glycolate, sodium carboxymethyl cellulose, calcium carboxymethyl cellulose, croscarmellose sodium, crospovidone, polyvinylpyrrolidone, methyl cellulose, microcrystalline cellulose, lower alkyl-substituted hydroxypropyl cellulose, starch, pregelatinised starch and sodium alginate. Generally, the disintegrant will comprise from 1 weight % to 25 weight %. In one embodiment of the present invention, the disintegrant will comprise from 5 weight % to 20 weight % of the dosage form. Binders are generally used to impart cohesive qualities to a tablet formulation. Suitable binders include microcrystalline cellulose,

gelatin, sugars, polyethylene glycol, natural and synthetic gums, polyvinylpyrrolidone, pregelatinised starch, hydroxypropyl cellulose and hydroxypropyl methylcellulose. Tablets may also contain diluents, such as lactose (monohydrate, spray-dried monohydrate, anhydrous and the like), mannitol, xylitol, dextrose, sucrose, sorbitol, 5 microcrystalline cellulose, starch and dibasic calcium phosphate dihydrate. Tablets may also optionally comprise surface active agents, such as sodium lauryl sulfate and polysorbate 80, and glidants such as silicon dioxide and talc. When present, surface active agents may comprise from 0.2 weight % to 5 weight % of the tablet, and glidants may comprise from 0.2 weight % to 1 weight % of the tablet. Tablets also generally 10 contain lubricants such as magnesium stearate, calcium stearate, zinc stearate, sodium stearyl fumarate, and mixtures of magnesium stearate with sodium lauryl sulfate. Lubricants generally comprise from 0.25 weight % to 10 weight %. In one embodiment of the present invention, lubricants comprise from 0.5 weight % to 3 weight % of the tablet. Other possible ingredients include anti-oxidants, colorants, flavoring agents, 15 preservatives and taste-masking agents.

Exemplary tablets contain up to about 80% drug, from about 10 weight % to about 90 weight % binder, from about 0 weight % to about 85 weight % diluent, from about 2 weight % to about 10 weight % disintegrant, and from about 0.25 weight % to about 10 weight % lubricant.

20 Tablet blends may be compressed directly or by roller to form tablets. Tablet blends or portions of blends may alternatively be wet-, dry-, or melt-granulated, melt congealed, or extruded before tableting. The final formulation may comprise one or more layers and may be coated or uncoated; it may even be encapsulated. Formulations of tablets are discussed in *Pharmaceutical Dosage Forms: Tablets*, Vol. 1, 25 by H. Lieberman and L. Lachman (Marcel Dekker, New York, 1980).

Consumable oral films for human or veterinary use are typically pliable water-soluble or water-swellaable thin film dosage forms which may be rapidly dissolving or mucoadhesive and typically comprise a compound of formula (I)-(li), a film-forming polymer, a binder, a solvent, a humectant, a plasticiser, a stabiliser or emulsifier, a 30 viscosity-modifying agent and a solvent. Some components of the formulation may perform more than one function. The film-forming polymer may be selected from natural polysaccharides, proteins, or synthetic hydrocolloids and is typically present in the range 0.01 to 99 weight %, more typically in the range 30 to 80 weight %. Other possible ingredients include anti-oxidants, colorants, flavorings and flavor enhancers,

preservatives, salivary stimulating agents, cooling agents, co-solvents (including oils), emollients, bulking agents, anti-foaming agents, surfactants and taste-masking agents. Films in accordance with the invention are typically prepared by evaporative drying of thin aqueous films coated onto a peelable backing support or paper. This may be done
5 in a drying oven or tunnel, typically a combined coater dryer, or by freeze-drying or vacuuming.

Solid formulations for oral administration may be formulated to be immediate and/or modified release. Modified release includes delayed, sustained, pulsed, controlled, targeted and programmed release. Suitable modified release formulations for
10 the purposes of the invention are described in US Patent No. 6,106,864. Details of other suitable release technologies such as high energy dispersions and osmotic and coated particles are to be found in Pharmaceutical Technology On-line, 25(2), 1-14, by Verma et al (2001). The use of chewing gum to achieve controlled release is described in WO-A-00/35298.

15 Compounds of formula (I)-(II) may also be administered directly into the blood stream, into muscle, or into an internal organ. Such parenteral administration includes intravenous, intraarterial, intraperitoneal, intrathecal, intraventricular, intraurethral, intrasternal, intracranial, intramuscular, intra-articular and subcutaneous administration. Suitable devices for parenteral administration include needle (including microneedle)
20 injectors, needle-free injectors and infusion techniques.

Compounds of the invention may also be administered topically to the skin or mucosa, that is, dermally or transdermally.

The compounds of formula (I)-(II) can also be administered intranasally or by inhalation, typically in the form of a dry powder (either alone, as a mixture, for example,
25 in a dry blend with lactose, or as a mixed component particle, for example, mixed with phospholipids, such as phosphatidylcholine) from a dry powder inhaler, as an aerosol spray from a pressurised container, pump, spray, atomiser (preferably an atomiser using electrohydrodynamics to produce a fine mist), or nebuliser, with or without the use of a suitable propellant, such as 1,1,1,2-tetrafluoroethane or 1,1,1,2,3,3,3-
30 heptafluoropropane, or as nasal drops. For intranasal use, the powder may comprise a bioadhesive agent, for example, chitosan or cyclodextrin. Delivery by inhalation is the preferred route of administration for the compounds of the present invention.

The pressurised container, pump, spray, atomizer, or nebuliser contains a solution or suspension of the compound of formula (I)-(II) comprising, for example,

ethanol, aqueous ethanol, or a suitable alternative agent for dispersing, solubilising, or extending release of the compound, a propellant as solvent and an optional surfactant, such as sorbitan trioleate, oleic acid, or an oligolactic acid.

Prior to use in a dry powder or suspension formulation, the drug product is
5 micronised to a size suitable for delivery by inhalation (typically less than 5 microns). This may be achieved by any appropriate comminuting method, such as spiral jet milling, fluid bed jet milling, supercritical fluid processing to form nanoparticles, high pressure homogenisation, or spray drying.

Capsules (made, for example, from gelatin or hydroxypropylmethylcellulose),
10 blisters and cartridges for use in an inhaler or insufflator may be formulated to contain a powder mix of the compound of the invention, a suitable powder base such as lactose or starch and a performance modifier such as l-leucine, mannitol, or magnesium stearate. The lactose may be anhydrous or in the form of the monohydrate, preferably the latter. Other suitable excipients include dextran, glucose, maltose, sorbitol, xylitol,
15 fructose, sucrose and trehalose.

A suitable solution formulation for use in an atomiser using electrohydrodynamics to produce a fine mist may contain from 1µg to 20mg of the compound of the invention per actuation and the actuation volume may vary from 1µl to 100µl. A typical formulation may comprise a compound of formula (I)-(II), propylene glycol, sterile water, ethanol and
20 sodium chloride. Alternative solvents which may be used instead of propylene glycol include glycerol and polyethylene glycol.

Suitable flavors, such as menthol and levomenthol, or sweeteners, such as saccharin or saccharin sodium, may be added to those formulations of the invention intended for intranasal administration. Formulations for intranasal administration may be
25 formulated to be immediate and/or modified release using, for example, PGLA. Modified release includes delayed, sustained, pulsed, controlled, targeted and programmed release.

Compounds of formula (I)-(II) may also be administered directly to the eye or ear, typically in the form of drops of a micronised suspension or solution in isotonic, pH-
30 adjusted, sterile saline.

Compounds of formula (I)-(II) may be combined with soluble macromolecular entities, such as cyclodextrin and suitable derivatives thereof or polyethylene glycol-containing polymers, in order to improve their solubility, dissolution rate, taste, bioavailability and/or stability when using any of the aforementioned modes of

administration. Drug-cyclodextrin complexes, for example, are found to be generally useful for most dosage forms and administration routes. Both inclusion and non-inclusion complexes may be used. As an alternative to direct complexation with the drug, the cyclodextrin may be used as an auxiliary additive, i.e. as a carrier, diluent, or solubilizer. Most commonly used for these purposes are alpha-, beta- and gamma-cyclodextrins, examples of which may be found in international patent publications WO91/11172, WO94/02518 and WO98/55148.

Inasmuch as it may be desirable to administer a combination of active compounds, for example, for the purpose of treating a particular disease or condition, it is within the scope of the present invention that two or more pharmaceutical compositions, at least one of which contains a compound of formula (I)-(II), may conveniently be combined in the form of a kit suitable for coadministration of the compositions. Thus, a kit of the invention comprises two or more separate pharmaceutical compositions, at least one of which contains a compound of formula (I)-(II), and means for separately retaining said compositions, such as a container, divided bottle, or divided foil packet. An example of such a kit is the familiar blister pack used for the packaging of tablets, capsules and the like. Such a kit is particularly suitable for administering different dosage forms, for example, oral and parenteral, for administering separate compositions at different dosage intervals, or for titrating the separate compositions against one another. To assist compliance, the kit typically comprises directions for administration and may be provided with a so-called memory aid.


The compounds of the invention may be prepared by any method known in the art for the preparation of compounds of analogous structure. In particular, the compounds of the invention can be prepared by the procedures described by reference to the Schemes that follow, or by the specific methods described in the Examples, or by processes similar to either.

The skilled person will appreciate that the experimental conditions set forth in the schemes that follow are illustrative of suitable conditions for effecting the transformations shown, and that it may be necessary or desirable to vary the precise conditions employed for the preparation of compounds of formula (I)-(II). It will be further appreciated that it may be necessary or desirable to carry out the transformations in a different order from that described in the schemes, or to modify one or more of the transformations, to provide the desired compound of the invention.

In addition, the skilled person will appreciate that it may be necessary or desirable at any stage in the synthesis of compounds of the invention to protect one or more sensitive groups, so as to prevent undesirable side reactions. In particular, it may be necessary or desirable to protect amino or carboxylic acid groups. The protecting groups used in the preparation of the compounds of the invention may be used in conventional manner. See, for example, those described in 'Greene's Protective Groups in Organic Synthesis' by Theodora W Greene and Peter G M Wuts, third edition, (John Wiley and Sons, 1999), in particular chapters 7 ("Protection for the Amino Group") and 5 ("Protection for the Carboxyl Group"), which also describes methods for the removal of such groups.

All of the derivatives of the formula (I)-(II) can be prepared by the procedures described in the general methods presented below or by routine modifications thereof. The present invention also encompasses any one or more of these processes for preparing the derivatives of formula (I)-(II), in addition to any novel intermediates used therein. The person skilled in the art will appreciate that the following reactions may be heated thermally or under microwave irradiation.

According to a first process, compounds of formula (I) may be prepared from compounds of formula (IX) and (VIII), as illustrated by Scheme 1.



Scheme 1

Wherein X is –CONH–; Hal is Cl, Br or iodo; M is boronic acid or boronate ester; PG¹ is tert-butyl, 2,4-dimethoxybenzyl; PG² is silylethoxymethyl, tetrahydropyranyl; PG³ is silylethoxymethyl, benzyl, or methyl.

5 It may be necessary or desirable to interchange the protecting groups in this Scheme to provide the highest yielding transformations.

Compounds of formulae (X), (IX), (VIII) and (IV) are commercially available or may be synthesized by those skilled in the art according to the literature or preparations described herein.

10 Compounds of formula (I)-(II) may be prepared from compounds of formula (II) according to process step (vi), a deprotection step mediated by either an organic acid, a Lewis acid or hydrogenation, or a sequential combination of each required. Preferred conditions comprise TFA and/or boron tribromide in a suitable organic solvent such as DCM or neat, at room or elevated temperatures and/or hydrogenation using a suitable
15 catalyst such as 10% Pd/C in an organic solvent such as EtOH at room temperature.

Wherein compounds of formula (I)-(II) are racemic, chiral separation may be employed to afford the two enantiomers. Wherein compounds of formula (I) include an R group that contains oxooxazolidine, this may be reacted with a suitable organic base to effect an open chain R group. Preferred conditions comprise sodium hydroxide at
20 from 0 °C to room temperature for 18 hours.

Compounds of formula (II) may be prepared from compounds of formula (III) according to process step (v), a carbonylation step in the presence of a suitable amine of formula (X) or (XI), a suitable palladium catalyst, and an organic base and a suitable solvent heated either in a sealed tube or under microwave irradiation. Typical conditions
25 comprise molybdenum hexacarbonyl with DBU and palladium acetate heated to 100 °C either thermally for 45 minutes or under microwave irradiation for 10 minutes in the presence of a compound of formula (X) or (XI), such as methylamine or 88% ammonia in a suitable organic solvent such as THF. Alternatively carbon monoxide gas (typically at 1 – 100 atmospheres) can be used in place of molybdenum hexacarbonyl in the
30 carbonylation step.


Compounds of formula (III) may be prepared from compounds of formula (IV) and (V) according to process step (iv), an N-oxide rearrangement step with compounds of formula (IV) and an organic base in a suitable organic solvent at elevated

temperatures. Preferred conditions comprise triethylamine in DMF at elevated temperatures of between 80-100 °C for 18 hours.

Compounds of formula (V) may be prepared from compounds of formula (VI) according to process step (iii), an oxidation reaction. Preferred conditions comprise mCPBA in DCM at 0 °C for 18 hours. Compounds of formula (VI) may be prepared from compounds of formula (VII) according to process step (ii), an electrophilic halogenation reaction. Typically, compounds (VII) have the PG² protecting group removed by methods known to those skilled in the art prior to electrophilic halogenation. Preferred conditions comprise N-iodosuccinimide in DMF at from 0 °C to room temperature for 18 hours followed by subsequent reprotection with PG².

Compounds of formula (VII) may be prepared from compounds of formula (IX) and (VIII) according to process step (i), a Suzuki cross-coupling reaction with compounds of formula (V). Suzuki cross-coupling is conveniently effected in the presence of a suitable catalyst, e.g., palladium or nickel and a base. Typical conditions comprise a boronic acid or ester, a palladium catalyst with phosphine ligands in an organic solvent at elevated temperatures. Preferred Suzuki conditions comprise palladium acetate with phosphine ligand S-Phos, and potassium phosphate in ethanol at 80 °C for 18 hours.

According to a second process, compounds of formula (I) may be prepared from compounds of formula (VI) as illustrated by Scheme 2.



Scheme 2

Wherein X is -CONH-; Hal is Cl, Br or I, PG¹ is tert-butyl, 2,4-dimethoxybenzyl; PG² is silylethoxymethyl, tetrahydropyranyl; PG³ is silylethoxymethyl, benzyl, or methyl.


Compounds of formulae (XII) are commercially available or may be synthesized by those skilled in the art according to the literature or preparations described herein.

5 Compounds of formula (VI) are described in Scheme 1.

Compounds of formula (I) may be prepared from compounds of formula (XIII) according to process steps (vii) and (vi), a nucleophilic aromatic substitution reaction with compounds of formula (XII) followed by a deprotection step. Typical conditions comprise heating to 90 °C with compounds of formula (XII) in a suitable organic solvent
10 with a suitable organic base, followed by deprotection as described in Scheme 1. Preferred conditions comprise DIPEA in n-butanol at 90 °C for 18 hours or triethylamine in DMF at 80-100 °C for 6 hours followed by TFA in DCM followed by boron tribromide in DCM. Alternatively compounds of formula (I) may be prepared from compounds of formula (XIII) and formula (XII) using a cross coupling reaction followed by deprotection
15 if required. Typical conditions comprise a suitable metal catalyst in the presence of an inorganic base with an organic ligand. Preferred conditions comprise Pd₂(dba)₃ with BINAP and cesium carbonate in toluene at elevated temperatures of 80-140 °C either thermally or under microwave irradiation.

Compounds of formula (XIII) may be prepared from compounds of formula (XIV) according to process steps (iii) and (viii), an oxidation reaction followed by an N-oxide
20 rearrangement-halogenation reaction. Typical conditions comprise oxidation as described in Scheme 1 process step (iii) followed by stirring the N-oxide in a suitable organic solvent at temperatures of 0-10 °C with electrophilic halogenating reagents. Preferred conditions comprise mCPBA in DCM followed by either POCl₃ or oxalyl
25 chloride in DCM. Compounds of formula (XIV) may be prepared from compounds of formula (VI) and either (X) or (XI) according to process step (v) as described in Scheme 1.

According to a third process, compounds of formula (I) may be prepared from compounds of formula (III) as illustrated by Scheme 3.



Scheme 3


Wherein X is -CONH- ; Hal is Cl, Br or I; PG^1 is tert-butyl, 2,4-dimethoxybenzyl; PG^2 is silylethoxymethyl, tetrahydropyranyl; PG^3 is silylethoxymethyl, benzyl, or methyl.

Compounds of formulae (XI) and (XI) are commercially available or may be synthesized by those skilled in the art according to the literature or preparations described herein. Compounds of formula (III) are described in Scheme 1. Compounds of formula (I) may be prepared from compounds of formula (II) according to process step (vi) as described in Scheme 1. Compounds of formula (II) may be prepared from compounds of formula (XV) according to process step (ix), an amide bond formation reaction with compounds of formula (X) or (XI) with activation of the carboxylic acid via a mixed anhydride or using a suitable base such as DIPEA and a suitable coupling agent such as HATU, BOP. Preferred conditions comprise isobutyl chloroformate in THF with NMM or BOP or HATU in DMF with DIPEA as base.

Process step (vi) may be performed before process step (ix) to obtain compounds of formula (I) in Scheme 3.

Compounds of formula (XV) may be prepared from compounds of formula (III) according to process step (v) as described in Scheme 1 but in the absence of compounds of formula (X) and (XI) in a solvent such as methanol with water added if necessary.

According to a fourth process, compounds of formula (I) may be prepared from compounds of formula (XX) and (XIV) as illustrated by Scheme 4.



Scheme 4

Wherein X is $-\text{CONH}-$; Hal is Cl, Br or I; PG^1 is tert-butyl, 2,4-dimethoxybenzyl;
 PG^2 is silylethoxymethyl, tetrahydropyranyl; PG^3 is silylethoxymethyl, benzyl, or methyl;
 5 PG^4 is carboxybenzyl.


Compounds of formulae (XVII), (X) and (XI) are commercially available or may
 be synthesized by those skilled in the art according to the literature or preparations
 described herein. Compounds of formula (XIV) are described in Scheme 2.
 Compounds of formula (XXI) are described in Scheme 5. Compounds of formula (I)
 10 may be prepared from compounds of formula (XVI) according to process step (vi) as
 described in Scheme 1.

Compounds of formula (XVI) may be prepared from compounds of formula (XVII)
 and (XVIII) according to process step (ix), a sulfonamide formation step. Preferred
 conditions comprise reacting compounds of formula (XVII) with compounds of formula
 15 (XVIII) in a suitable organic solvent such as THF at from 0 °C to room temperature for

18 hours. Alternatively a base may be added to facilitate the reaction such as sodium hydride. Compounds of formula (XVIII) may be prepared from compounds of formula (XIX) according to process step (vi) a deprotection reaction as described in Scheme 1. Preferred conditions comprise palladium on carbon in ethanol at room temperature under hydrogenation at 30 psi for 1 hour.

Compounds of formula (XIX) may be prepared from compounds of formula (XIV) according to process steps (iii) and (iv), an oxidation of compounds of formula (XIV) followed by a rearrangement step with compounds of formula (XX) as described in Scheme 1.

According to a fifth process, compounds of formula (IV) may be prepared from compounds of formula (XXIV) as illustrated by Scheme 5.



Compounds of formulae (XXIV), (XVII) and (XXVI) are commercially available or may be synthesized by those skilled in the art according to the literature or preparations described herein. Compounds of formula (IV) may be prepared from compounds of formula (XXI) according to process step (xv), a reaction to form a carbamate activating group in the presence of an inorganic base. Preferred conditions comprise sodium carbonate in DCM with 4-nitrophenylchloroformate.


Compounds of formula (XXI) may be prepared from compounds of formula (XXII) according to process step (xiv), a reduction step in the presence of a metal catalyst and an inorganic hydrogen donor or under an atmosphere of hydrogen. Preferred conditions

comprise $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ with sodium borohydride and di-tert butyl dicarbonate in methanol followed by 4M HCl in dioxane or 10% palladium on carbon in acetic acid or RaneyTM Nickel in methanolic ammonia under an atmosphere of 40 psi of hydrogen at room temperature for 18 hours.

- 5 Compounds of formula (XXII) may be prepared from compounds of formula (XXV) according to process step (xiii), an alkylation reaction with compounds of formula (XXVI) in the presence of a quaternary ammonium salt. Preferred conditions comprise benzyltriethylammonium chloride and 40% aqueous sodium hydroxide solution in THF with compounds of formula (XXVI). Compounds of formula (XXII) may also be prepared
- 10 from compounds of formula (XXIII) according to process step (xii), an alkylation reaction in the presence of an inorganic base. Preferred conditions comprise potassium carbonate in acetone with compounds of formula (XXVI) or Mitsunobu conditions with compounds of formula (XVIII) using DEAD in THF.

- 15 Compounds of formulae (XXV) and (XXIII) may be prepared from compounds of formula (XXIV) and (XVII) according to process step (xi) a sulfonamide formation reaction. Preferred conditions comprise stirring in pyridine at from 0 °C to room temperature or in the presence of LiHMDS in THF. Compounds of formula (XXIII) may also be prepared from sulfonamides reacting with halo-substituted heterocycles in the presence of a base such as cesium carbonate in acetonitrile.

- 20 According to a sixth process, compounds of formula (II) may be prepared from compounds of formula (VII) as illustrated by Scheme 6.




Scheme 6

Compounds of formula (VII) may be prepared as described in Scheme 1. Compounds of formulae (XII) are commercially available or may be synthesized by those skilled in the art according to the literature or preparations described herein. Compounds of formula (I) may be prepared from compounds of formula (XXIX) according to reaction step (vi), a deprotection step as described in Scheme 1.

Compounds of formula (XXIX) may be prepared according to reaction step (xvi), an N-oxide rearrangement step effected by employment of a dehydrating agent such as PyBrop with amines of formula (XII). Preferred conditions comprise PyBrop with DIPEA in a suitable organic solvent such as DCM at room temperature. Alternatively the N-oxide rearrangement step may employ acetic anhydride to afford the hydroxy intermediate followed by interconversion to triflate. The triflate may then be converted to compounds of formula (XXIX) by heating with amines of formula (XII). Typical conditions comprise heating the N-oxide with triethylamine and acetic anhydride, followed by triflic anhydride with pyridine in DCM at room temperature, and finally heating with compounds of formula (XII) with triethylamine in DMF.

Compounds of formula (XXVIII) may be prepared from compounds of formula (VII) according to process step (iii) as described in Scheme 1.

According to a seventh process, compounds of formula (I) may be prepared from compounds of formula (VII) as illustrated by Scheme 7.




Scheme 7

Compounds of formula (III) may be prepared as described in Scheme 1. Compounds of formulae (XXX) or (XXXI) are commercially available or may be synthesized by those skilled in the art according to the literature or preparations described herein.

5 Compounds of formula (I)-(II) may be prepared from compounds of formula (III) according to process steps (xvii) and (vi), a cross coupling reaction, such as a Stille reaction or a Buchwald reaction followed by a deprotection step if required. Typical conditions for a Stille cross coupling reaction comprise a suitable tin reagent in the presence of one or two metal catalysts in a suitable organic solvent at elevated
10 temperatures with compounds of formula (XXX). Preferred conditions comprise bis(tributyltin) and copper(I)iodide with tetrakis(triphenylphosphine)palladium in toluene at 100 °C. Typical conditions for a Buchwald reaction comprise a copper catalyst and a suitable organic ligand in the presence of an inorganic base at elevated temperatures. Preferred conditions comprise cuprous oxide and 4,7-dimethoxy-1,10-phenanthroline
15 with cesium carbonate and PEG in DMSO at 110 °C.

Compounds of formula (I) may also be prepared from compounds of formula (XXXV) according to process step (xviii) a heterocyclic cyclization reaction, either directly from the nitrile or the carboxylic acid, or via an acyl hydrazone from the carboxylic acid. Preferred conditions comprise heating with the required nitrile or
20 hydrazone in butanol at elevated temperatures under microwave irradiation. Compounds of formula (XXXV) may be prepared from compounds of formula (III) according to process step (v) as described in Scheme 3 to afford the carboxylic acid, or using zinc cyanide and tetrakis(triphenylphosphine)palladium in DMF at elevated temperatures under microwave irradiation to afford the nitrile.

25 According to an eighth process, compounds of formula (I)-(II) may be prepared from compounds of formula (XXXIII) as illustrated by Scheme 8.



Scheme 8

Wherein M is boronic acid or ester.

Compounds of formula (XXXIII) are either commercially available or prepared as described herein. Compounds of formula (I)-(II) may be prepared from compounds of formula (XXXII) and (VIII) according to process steps (i) and (vi), a Suzuki cross-coupling reaction followed by deprotection as described in Scheme 1. Preferred
 5 conditions for the Suzuki step comprise PEPPSI-IPr catalyst with potassium carbonate in toluene at elevated temperatures. Compounds of formula (XXXII) may be prepared from compounds of formula (XXXIII) and (XII) according to process step (vii), a nucleophilic aromatic substitution reaction as described in Scheme 2.

The skilled person will further appreciate that compounds of formula (I)-(II) may
 10 be interconverted to other compounds of formula (I)-(II) by functional group manipulation, or suitably protected compounds of formula (I)-(II) may be interconverted to other suitably protected compounds of formula (I)-(II) followed by a deprotection step to afford compounds of formula (I)-(II)

Typical interconversions include:

15 Wherein R or R⁰ contains a ketone or aldehyde functionality, these may be reduced using a suitable reducing agent such as sodium borohydride;

Wherein R or R⁰ contain an amine, these may be interconverted to a urea, an amide, a sulfonamide or a sulfamide followed by suitable deprotection as required.

20 Wherein compounds of formula (XVI) contain an R'' group that has a leaving group such as halo, an alkylation may occur with amines such as morpholine.

In the non-limiting Examples and Preparations that are set out later in the description, and in the aforementioned Schemes, the following the abbreviations, definitions and analytical procedures may be referred to:

ACE-Cl is 1-chloroethylchloroformate;

25 BBr₃ is boron tribromide;

BINAP is 2,2'-bis(diphenylphosphino)-1,1'-binaphthalene;

BOP is (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate;

Cbz is benzyloxycarbonyl;

30 Cs₂CO₃ is cesium carbonate;

DBU is diazabicyclo[5.4.0]undec-7-ene;

DCM is dichloromethane;

DEAD is diethylazodicarboxylate;

DIPEA is N-ethyldiisopropylamine, N,N-diisopropylethylamine;

- DMAP is dimethylaminopyridine;
 DMF is dimethyl formamide;
 EDCI.HCl is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride;
 EtOAc is ethylacetate;
 5 HATU is 1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-
 oxid hexafluorophosphate;
 HBTU is *N,N,N',N'*-Tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uronium
 hexafluorophosphate;
 HCl is hydrochloric acid;
 10 HOBt is 1-hydroxybenzotriazole;
 IPA is isopropanol;
 KOAc is potassium acetate;
 LiHMDS is lithium (bistrimethylsilyl)amide
m-CPBA is meta chloroperoxy benzoic acid
 15 MeCN is acetonitrile;
 MeOH is methanol;
 NaBH₄ is sodium borohydride;
 NaHCO₃ is sodium hydrogen carbonate;
 NaH is sodium hydride;
 20 NaOH is sodium hydroxide;
 NBS is N-bromosuccinimide;
 NiCl₂·6H₂O is nickel dichloride hydrate;
 NMM is N-methylmorpholine;
 NMP is N-methyl-2-pyrrolidone;
 25 PeppiTM-IPr is [1,3-Bis(2,6-Diisopropylphenyl)imidazol-2-ylidene](3-
 chloropyridyl)palladium(II) dichloride;
 Pd/C is palladium on carbon;
 Pd₂(dba)₃ is tris(dibenzylideneacetone)dipalladium;
 Pd(dppf)2Cl₂ is 1,1'-bis(diphenylphosphino)ferrocene-palladium(II)dichloride;
 30 Pd(OAc)₂ is palladium acetate;
 Pd(PPh₃)₄ is tetrakis(triphenylphosphine)palladium (0)
 PEG is polyethylene glycol;
 POCl₃ is phosphorus oxychloride;
 PTSA is paratoluenesulfonic acid;

PyBrop is (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate;

SEM is silylethoxymethyl;

SPhos is 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl;

TBDMS is tertbutyldimethylsilyl;

5 TBME is tert-butyl dimethyl ether;

t-BuOK is potassium tert-butoxide;

TEA is triethylamine;

TES is triethylsilyl;

Tf is triflate which is trifluoromethanesulfonate;

10 TFA is trifluoroacetic acid;

THF is tetrahydrofuran;

THP is tetrahydropyran; and,

TLC is thin layer chromatography.

¹H and ¹⁹F Nuclear magnetic resonance (NMR) spectra were in all cases
 15 consistent with the proposed structures. Characteristic chemical shifts (δ) are given in
 parts-per-million downfield from tetramethylsilane (for ¹H-NMR) and upfield from
 trichloro-fluoro-methane (for ¹⁹F NMR) using conventional abbreviations for designation
 of major peaks: e.g. s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad.
 The following abbreviations have been used for common solvents: CDCl₃,
 20 deuteriochloroform; d₆-DMSO, deuterodimethylsulphoxide; and CD₃OD,
 deuteromethanol. Where appropriate, tautomers may be recorded within the NMR data;
 and some exchangeable protons may not be visible. Mass spectra, MS (m/z), were
 recorded using either electrospray ionisation (ESI) or atmospheric pressure chemical
 ionisation (APCI). Where relevant and unless otherwise stated the m/z data provided
 25 are for isotopes ¹⁹F, ³⁵Cl, ⁷⁹Br and ¹²⁷I. Wherein preparative TLC or silica gel
 chromatography has been used, one skilled in the art may choose any combination of
 solvents to purify the desired compound.

Either IUPAC or ACD Labs naming packages have been used, and are
 interchangeably employed throughout the Examples and Preparations.

30 **Preparative HPLC:**

Where singleton compounds are purified by preparative HPLC, these are two
 methods used, shown below:

Detection for both analytical and preparative QC:

Detectors: ELSD; Polymer Labs PL-ELS 2100, UV; Waters 2487 detector at 225 and 255 nm

Mass Spectrometer; Waters ZQ using electrospray ionization.

5 **Preparative Method 1 Acidic Conditions**

Column: Gemini NX C18, 5µm 21.2 x 100 mm; Temperature: Ambient;
Detection: ELSD-MS; Mobile Phase A: 0.1% formic acid in water; Mobile Phase B:
0.1% formic acid in acetonitrile; Gradient: initial 0% B, 1 min – 5% B; 7 mins –
95% B; 9 mins – 95% B; 9.1 mins – 5% B; 10 mins -5% B; Flow rate: 18 mL/min;

10 Injection volume: 1000 µL

Preparative Method 2 Basic Conditions

Column: Gemini NX C18, 5µm 21.2 x 100 mm; Temperature: Ambient; Detection:
ELSD-MS; Mobile Phase A: 0.1% diethylamine in water; Mobile Phase B: 0.1%
diethylamine in acetonitrile; Gradient: initial 0%B, 1 min – 5% B; 7 mins – 95% B; 9
15 mins – 95% B; 9.1 mins – 5% B; 10 mins -5% B; Flow rate: 18 mL/min; Injection
volume: 1000 µL

Analytical LCMS QC:

Column: Gemini C18 50 x 4.6 mm, 3 micron; 5 minutes run.
20 Gradient initial - 95% A, 5% B; 3 mins – 95% B; hold to 4 mins then back to 5% B
at 4.1-5 mins. Flow rate 1.5 mL/min
Acidic conditions: Mobile Phase A: 0.1% Formic acid in Water. Mobile Phase B:
0.1% Formic acid in acetonitrile
Basic conditions: Mobile Phase A: 0.1% ammonia in water; Mobile Phase B:
25 0.1% Ammonia in acetonitrile.

Example 1

4-({2-[Ethyl(ethylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-
trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

30 To a solution of N-ethyl-N-(2-{{(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethyl-
silyl)ethoxy}methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-
c]pyridin-4-yl)amino]methyl}phenyl)ethanesulfonamide (**Preparation 62**, 170 mg, 0.18
mmol) in 2M methylamine in THF (2.6 mL) was added molybdenum hexacarbonyl
(48.18 mg, 0.181 mmol), DBU (82.77 ml, 0.544 mmol) and palladium acetate (2.85 mg,

0.01 mmol). The reaction was heated at 100 °C under microwave irradiation for 10 minutes. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 30% EtOAc in hexanes. The residue was dissolved in TFA (3 mL) and stirred at room temperature for 30 minutes. The reaction was
5 concentrated *in vacuo*, dissolved in MeOH, cooled in ice and treated with ethylene diamine. The reaction was stirred at room temperature for 2 hours before concentrating *in vacuo*. The residue was purified using silica gel column chromatography eluting with EtOAc to afford the title compound (60 mg, 58% over 2 steps).

¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.90 (t, 3H), 1.15 (t, 3H), 2.85 (d, 3H), 3.21
10 (m, 2H), 3.44-3.63 (m, 4H), 4.72-4.74 (m, 1H), 4.86-4.88 (m, 1H), 6.69 (s, 1H), 6.92 (d, 1H), 7.19 (d, 1H), 7.31 (m, 2H), 7.40 (m, 2H), 8.85 (m, 1H), 9.77 (m, 1H), 10.07 (s, 1H), 13.71 (s, 1H).

MS m/z 609 [M+H]⁺

15 The following Examples (**Examples 2 –25**) were prepared according to the method described for **Example 1** using the appropriate pyrazolo-pyridine and Purification Method (PM) below if different from the method described:

Purification Method A: Silica gel column chromatography eluting with between 40-60% EtOAc in hexanes.

20 **Purification Method B:** Silica gel column chromatography or Preparative TLC eluting with 4% MeOH in DCM.

Purification Method C: Silica gel column chromatography followed by Preparative TLC eluting both with up to 30% MeOH in DCM.

Purification Method D: Silica gel column chromatography eluting with EtOAc.

25

Ex	Name	Data
2	4-({2-[ethyl(ethylsulfonyl)-amino]benzyl}amino)-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 555 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.85 (t, 3H), 0.93 (t, 3H), 1.20 (t, 3H), 2.84 (d, 3H), 3.15-3.31 (m, 2H), 3.50-3.70 (m, 2H), 4.69-4.71 (m, 1H), 4.95-4.98 (m, 1H), 6.62 (s, 1H), 6.74 (d, 1H), 6.97 (s, 1H), 7.31 (m, 2H), 7.41 (m, 2H), 8.79 (m, 1H), 9.64 (m, 1H). Using N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-{{2-(trimethylsilyl)ethoxy}-methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)-methyl]phenyl]ethanesulfonamide (Preparation 63) and PM A.
3	6-(2-ethyl-5-fluoro-4-hydroxy-phenyl)-4-({2-[(ethylsulfonyl)-(methyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 541 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.86 (t, 3H), 1.22 (t, 3H), 2.49 (m, 2H), 2.84 (d, 3H), 3.35 (m, 2H), 4.70-5.00 (br m, 2H), 6.63 (s, 1H), 6.78 (d, 1H), 7.03 (d, 1H), 7.32 (m, 2H), 7.41 (m, 2H), 8.80 (t, 1H), 9.66 (t, 1H), 9.76 (s, 1H), 13.63 (s, 1H). Using N-[2-({[6-(2-ethyl-5-fluoro-4-{{2-(trimethylsilyl)ethoxy}-methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl]phenyl]-N-methylethanesulfonamide (Preparation 64) and PM A.
4	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-{{2-(4-hydroxyphenyl)ethyl}amino}-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 450 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.06 (t, 3H), 2.66-2.84 (m, 7H), 3.62 (m, 2H), 6.61 (s, 1H), 6.65 (m, 2H), 6.86 (m, 1H), 7.05 (m, 2H), 7.15 (m, 1H), 8.73 (m, 1H), 9.28 (m, 1H). Using 4-(2-{{[6-(2-ethyl-5-fluoro-4-{{2-(trimethylsilyl)ethoxy}-methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino}ethyl)phenol (Preparation 65) and PM B.
5	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-methylpropyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 386 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.97 (d, 6H), 1.08 (t, 3H), 1.89 (m, 1H), 2.69 (m, 2H), 2.84 (d, 3H), 3.35 (m, 2H), 6.58 (s, 1H), 6.85 (d, 1H), 7.09 (d, 1H), 8.74 (m, 1H), 9.31 (m, 1H), 9.79 (s, 1H), 13.50 (s, 1H). N-ethyl-N-(2-{{[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl]phenyl)ethanesulfonamide (Preparation 66) and PM A.

6	4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 561 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.83 (t, 3H), 2.44 (m, 2H), 2.86 (d, 3H), 3.05 (s, 3H), 3.08 (s, 3H), 4.69-4.87 (br m, 2H), 6.66 (s, 1H), 6.78 (d, 1H), 7.03 (d, 1H), 7.36-7.39 (m, 2H), 7.54 (d, 1H), 8.83 (m, 1H), 9.69 (m, 1H), 9.77 (s, 1H), 13.67 (s, 1H). N-[4-chloro-2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide (Preparation 67) and PM A.
7	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 545 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.85 (t, 3H), 2.45 (br m, 2H), 2.85 (d, 3H), 3.05 (s, 3H), 3.08 (s, 3H), 4.69 (br m, 1H), 4.90 (br m, 1H), 6.65 (s, 1H), 6.78 (d, 1H), 7.02 (d, 1H), 7.10-7.18 (m, 2H), 7.55 (m, 1H), 8.83 (m, 1H), 9.68 (m, 1H), 9.77 (s, 1H), 13.66 (s, 1H). N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}-4-fluorophenyl)-N-methylmethanesulfonamide (Preparation 68) and PM A.
8	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-fluoro-6-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 545 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.02 (t, 3H), 2.66 (m, 2H), 3.02 (s, 6H), 4.57-5.00 (br m, 2H), 6.61 (s, 1H), 6.85 (d, 1H), 7.06 (d, 1H), 7.25 (m, 1H), 7.36-7.45 (m, 2H), 8.71 (m, 1H), 9.44 (m, 1H), 9.79 (s, 1H), 13.58 (s, 1H). N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl)-3-fluorophenyl)-N-methylmethanesulfonamide (Preparation 69) and PM A.
9	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-[ethyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 541 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.85 (t, 3H), 0.93 (t, 3H), 2.45 (m, 2H), 2.84 (d, 3H), 3.02 (s, 3H), 3.49 (m, 1H), 3.68 (m, 1H), 4.70 (m, 1H), 4.97 (m, 1H), 6.62 (s, 1H), 6.77 (d, 1H), 6.98 (d, 1H), 7.31 (m, 2H), 7.44 (m, 2H), 8.79 (m, 1H), 9.63 (m, 1H), 9.73 (br s, 1H), 13.62 (br s, 1H). N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}-phenyl]methanesulfonamide (Preparation 70) and PM A.

10	4-[(cyclopentylmethyl)amino]-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 412 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.04 (t, 3H), 1.28 (m, 2H), 1.54-1.61 (m, 4H), 1.77 (m, 2H), 2.15 (m, 1H), 2.65 (m, 2H), 2.84 (d, 3H), 3.41 (m, 2H), 6.58 (s, 1H), 6.85 (d, 1H), 7.12 (d, 1H), 8.73 (m, 1H), 9.29 (m, 1H), 9.78 (s, 1H), 13.54 (br s, 1H). N-(cyclopentylmethyl)-6-(2-ethyl-5-fluoro-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl)-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridin-4-amine (Preparation 71) and PM A.
11	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-((5-methyl-2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 595 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.21 (s, 3H), 2.85 (d, 3H), 3.01 (s, 3H), 3.07 (s, 3H), 3.62 (m, 2H), 4.64-4.83 (m, 2H), 6.70 (s, 1H), 6.93 (d, 1H), 7.13 (m, 1H), 7.19 (m, 2H), 7.37 (d, 1H), 8.82 (m, 1H), 9.71 (m, 1H), 10.09 (s, 1H), 13.70 (s, 1H). 6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-N-methyl-4-((5-methyl-2-(N-methylmethylsulfonylamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 72) and PM C.
12	4-((2-[ethyl(methylsulfonyl)amino]benzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 595 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.92 (t, 3H), 2.85 (s, 1H), 3.02 (s, 3H), 3.50 (m, 2H), 3.64 (m, 2H), 4.72 (m, 1H), 4.93 (m, 1H), 6.69 (s, 1H), 6.93 (d, 1H), 7.16 (d, 1H), 7.31-7.45 (m, 4H), 8.84 (m, 1H), 9.75 (m, 1H), 10.07 (s, 1H), 13.70 (br s, 1H). N-ethyl-N-(2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-phenyl)methane sulphonamide (Preparation 73) and PM D.
13	4-((2-[(ethylsulfonyl)methyl]amino)benzyl)amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 595 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.21 (t, 3H), 2.85 (d, 3H), 3.11 (s, 3H), 3.32 (m, 2H), 3.57 (m, 2H), 4.72-4.84 (br m, 2H), 6.70 (s, 1H), 6.94 (d, 1H), 7.18-7.47 (m, 4H), 8.84 (m, 1H), 9.78 (m, 1H), 10.08 (br s, 1H), 13.70 (br s, 1H). N-(2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)phenyl)-N-methylethane Sulphonamide (Preparation 74) and PM A.

14	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 599 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.86 (d, 3H), 3.05 (s, 3H), 3.08 (s, 3H), 3.55 (m, 2H), 4.65-4.90 (m, 2H), 6.73 (s, 1H), 7.06-7.21 (m, 4H), 8.86 (m, 1H), 9.80 (m, 1H), 10.10 (s, 1H), 13.74 (s, 1H). Using N-(4-fluoro-2-{{(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl}phenyl)-N-methylmethanesulfonamide (Preparation 75) and PM D.
15	4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 613 [M-H] ⁻ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.05 (d, 3H), 3.05 (s, 3H), 3.07 (s, 3H), 3.52 (m, 2H), 4.75-4.90 (m, 2H), 6.73 (s, 1H), 6.94 (s, 1H), 7.21 (d, 1H), 7.33 (m, 1H), 7.39 (m, 1H), 7.54 (d, 1H), 8.86 (m, 1H), 9.79 (m, 1H), 10.08 (s, 1H), 13.74 (s, 1H). Using N-(4-chloro-2-{{(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}aminomethyl}phenyl)-N-methylmethanesulfonamide (Preparation 61) and PM C.
16	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-methylpropyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 440 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.97 (s, 3H), 0.98 (s, 3H), 1.90 (m, 1H), 2.84 (d, 3H), 3.31 (m, 2H), 4.10 (m, 2H), 6.67 (s, 1H), 7.05 (d, 1H), 7.30 (d, 1H), 8.77 (m, 1H), 9.39 (m, 1H), 10.13 (s, 1H), 13.62 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-N-(2-methylpropyl)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-amine (Preparation 78) and PM A.
17	4-[(cyclopentylmethyl)amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 466 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.28 (m, 2H), 1.52-1.61 (m, 4H), 1.79 (m, 1H), 2.15 (m, 1H), 3.39 (m, 2H), 4.12 (m, 2H), 6.67 (s, 1H), 7.02 (d, 1H), 7.30 (d, 1H), 8.76 (m, 1H), 9.37 (m, 1H), 10.13 (s, 1H), 13.62 (br s, 1H). N-(cyclopentylmethyl)-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-amine (Preparation 77) and PM A.
18	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-fluoro-6-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 599 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.80 (d, 3H), 3.01 (s, 3H), 3.02 (s, 3H), 3.80-4.25 (br m, 2H), 4.55-5.10 (br m, 1H), 6.70 (s, 1H), 7.04 (d, 1H), 7.28 (m, 2H), 7.43 (m, 2H), 8.74 (m, 1H), 9.54 (m, 1H), 10.14 (s, 1H), 13.66 (s, 1H).

	yl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	N-(3-fluoro-2-{{(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{[2-(trimethylsilyl)-ethoxy]methoxy}phenyl]-3-iodo-1-{{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl}phenyl)-N-methylmethanesulfonamide (Preparation 76) and PM A.
19	6-(2-ethyl-5-fluoro-4-hydroxy-phenyl)-N-methyl-4-{{2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 527 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.88 (t, 3H), 2.84 (d, 3H), 3.04 (s, 3H), 3.10 (s, 3H), 3.31 (m, 2H), 4.69 (br m, 1H), 4.94 (br m, 1H), 6.63 (s, 1H), 6.79 (d, 1H), 7.04 (d, 1H), 7.30 (m, 2H), 7.42 (m, 1H), 7.49 (m, 1H), 8.80 (m, 1H), 9.63 (m, 1H), 9.75 (s, 1H), 13.61 (br s, 1H). N-[2-{{(6-(2-ethyl-5-fluoro-4-{{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{{[2-(trimethylsilyl)-ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl}phenyl)-N-methylmethanesulfonamide (Preparation 79) and PM A.
20	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-{{2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 581 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.64 (d, 3H), 3.04 (s, 3H), 3.10 (s, 3H), 3.62 (q, 2H), 4.72 (br s, 1H), 4.88 (br s, 1H), 6.54 (s, 1H), 6.95 (d, 1H), 7.21 (d, 1H), 7.30 (m, 2H), 7.37 (d, 1H), 7.45 (m, 1H), 8.82 (t, 1H), 9.75 (t, 1H), 10.09 (s, 1H), 13.70 (s, 1H). N-(2-{{(6-[5-fluoro-2-(2,2,2-trifluoro-ethyl)-4-{{[2-(trimethylsilyl)ethoxy]-methoxy}phenyl]-3-iodo-1-{{[2-(tri-methylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino]-methyl}phenyl)-N-methylmethanesulfonamide (Preparation 105) and PM A.
21	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-{{5-methyl-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 541 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.90 (t, 3H), 2.23 (s, 3H), 2.84 (d, 3H), 3.01 (s, 3H), 3.08 (s, 3H), 3.39 (m, 2H), 4.65 (br m, 1H), 4.90 (br m, 1H), 6.63 (s, 1H), 6.80 (d, 1H), 7.06 (d, 1H), 7.14 (m, 1H), 7.23 (m, 1H), 7.37 (m, 1H), 8.80 (m, 1H), 9.60 (m, 1H), 9.76 (s, 1H), 13.61 (br s, 1H). N-[2-{{(6-(2-ethyl-5-fluoro-4-{{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{{[2-(trimethylsilyl)ethoxy]-methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl)-4-methylphenyl)-N-methylmethanesulfonamide (Preparation 80) and PM A.

22	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(phenylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 589 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.89 (t, 3H), 3.03 (s, 3H), 3.17 (d, 3H), 3.31 (m, 2H), 4.70 (br m, 1H), 5.00 (br m, 1H), 6.56 (d, 1H), 6.64 (s, 1H), 6.79 (d, 1H), 7.05 (d, 1H), 7.16 (m, 1H), 7.28 (m, 1H), 7.44 (m, 1H), 7.59-7.67 (m, 4H), 7.73 (m, 1H), 8.80 (m, 1H), 9.66 (m, 1H), 9.75 (br s, 1H), 13.62 (br s, 1H). N-[2-({[6-(2-ethyl-5-fluoro-4-({2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-({2-(trimethylsilyl)ethoxy}methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylbenzenesulfonamide (Preparation 81) and PM B.
23	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-{4-[(phenylsulfonyl)amino]phenyl}ethyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 589 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.01 (t, 3H), 2.69 (m, 2H), 2.78-2.83 (m, 5H), 3.61 (q, 2H), 6.60 (s, 1H), 6.85 (d, 1H), 6.98 (m, 2H), 7.12 (m, 3H), 7.48-7.57 (m, 3H), 7.72 (d, 1H), 8.73 (m, 1H), 9.25 (m, 1H), 10.13 (s, 1H), 13.55 (s, 1H). N-[4-(2-({[6-(2-ethyl-5-fluoro-4-({2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-({2-(trimethylsilyl)ethoxy}methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}ethyl)phenyl]benzenesulfonamide (Preparation 82) and PM B.
24	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[(2-hydroxyethyl)-(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 611 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.85 (d, 3H), 3.08 (s, 3H), 3.32 (m, 2H), 3.55-3.64 (m, 4H), 4.76-4.93 (m, 3H), 6.70 (s, 1H), 6.93 (d, 1H), 7.21 (d, 1H), 7.30-7.36 (m, 3H), 7.46 (m, 1H), 8.84 (m, 1H), 9.73 (m, 1H), 10.08 (s, 1H), 13.71 (br s, 1H). N-(2-({[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-({2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-({2-(trimethylsilyl)ethoxy}methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino]-methyl}phenyl)-N-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]methanesulfonamide (Preparation 87) and PM B.
25 (Intermediate)	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-methoxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 557 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.95 (t, 3H), 2.55 (m, 2H), 2.82 (d, 3H), 3.05 (s, 3H), 3.10 (s, 3H), 3.76 (s, 3H), 4.50-4.90 (m, 2H), 6.62 (s, 1H), 6.81 (d, 1H), 6.92 (dd, 1H), 7.04-7.07 (m, 2H), 7.34 (d, 1H), 8.78 (m, 1H), 9.54 (m, 1H), 9.76 (s, 1H), 13.60 (s, 1H). Using N-[2-({[6-(2-ethyl-5-fluoro-4-({2-(trimethylsilyl)ethoxy}-methoxy}phenyl)-3-iodo-1-({2-(trimethylsilyl)ethoxy}methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-5-methoxyphenyl]-N-methyl-methanesulfonamide (Preparation 88) and PM B.

The following Examples (**Examples 26 – 28**) were prepared according to the method described for **Example 1** using ammonia in THF and the appropriate pyrazolopyridine and Purification Method (PM) below if different from the method described.

5 **Purification Method E:** Silica gel column chromatography or preparative TLC eluting with 4% MeOH in DCM.

Example	Name	Data
26	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(phenylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 629 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.03 (s, 3H), 3.70 (m, 2H), 4.68 (m, 1H), 4.92 (m, 1H), 6.60 (d, 1H), 6.71 (s, 1H), 6.97 (d, 1H), 7.13-7.29 (m, 3H), 7.44 (d, 1H), 7.59-7.75 (m, 5H), 7.86 (br s, 1H), 8.21 (br s, 1H), 9.75 (m, 1H), 10.10 (s, 1H), 13.69 (s, 1H). Using N-(2-{{(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)-methyl}phenyl)-N-methylbenzene-sulfonamide (Preparation 86).
27	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[(2-hydroxyethyl)(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 596 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.07 (s, 3H), 3.32 (m, 2H), 3.56-3.67 (m, 4H), 4.73-4.92 (m, 3H), 6.70 9s, 1H), 6.94 (d, 1H), 7.18 (d, 1H), 7.21-7.45 (m, 4H), 7.83 (br s, 1H), 8.19 (br s, 1H), 9.74 (m, 1H), 10.10 (br s, 1H), 13.70 (br s, 1H). Using N-(2-{{(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl}phenyl)-N-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]methanesulfonamide (Preparation 87) and PM E.
28 (intermediate)	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-methoxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 543 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.95 (t, 3H), 2.59 (m, 2H), 3.04 (s, 3H), 3.16 (s, 3H), 3.31 (s, 3H), 3.76 (s, 3H), 4.65 (br m, 2H), 6.62 (s, 1H), 6.79 (d, 1H), 6.91 (d, 1H), 7.04-7.07 (m, 2H), 7.34 (d, 1H), 7.78 (br s, 1H), 8.14 (br s, 1H), 9.53 (m, 1H), 9.76 (s, 1H), 13.57 (s, 1H). Using N-[2-{{(6-(2-ethyl-5-fluoro-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl)-5-methoxyphenyl]-N-methylmethane-sulfonamide (Preparation 88) and PM E.

Example 29

6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-hydroxy-2-[methyl
(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide

To a solution of 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-methoxy-2-[methyl
(methyl-sulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide (**Example 28**, 80 mg, 0.14 mmol) in DCM (10 mL) was added boron
tribromide (0.09 mL, 1 mmol) at 0°C. The reaction was stirred at room temperature for 2
10 hours before the addition of further boron tribromide (0.09 mL, 1 mmol) and further
stirring for 2 hours. The reaction was partitioned between DCM and saturated aqueous
sodium bicarbonate solution, the organic layer was collected, dried over sodium sulfate
and concentrated *in vacuo*. The residue was purified using Preparative HPLC to afford
the title compound (32 mg, 41%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.94 (t, 3H),
15 2.59 (m, 2H), 2.81 (d, 3H), 3.02 (s, 3H), 3.07 (s, 3H), 4.69 (br m, 2H), 6.61 (s, 1H), 6.73
(dd, 1H), 6.79-6.83 (m, 2H), 7.05 (d, 1H), 7.23 (d, 1H), 8.76 (m, 1H), 9.49 (m, 1H), 9.60
(br s, 1H). MS m/z 543 [M+H]⁺

Example 30

6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-hydroxy-2-
[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound was prepared according to the method described for
Example 29 using 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-methoxy-2-
[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide
(**Example 27**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.96 (t, 3H), 2.60 (m, 2H), 3.01 (s,
25 3H), 3.08 (s, 3H), 4.65 (br m, 2H), 6.61 (s, 1H), 6.74 (dd, 1H), 6.80-6.83 (m, 2H), 7.09
(d, 1H), 7.25 (d, 1H), 7.77 (br s, 1H), 8.13 (br s, 1H), 9.49 (m, 1H), 9.56 (s, 1H), 9.76 (s,
1H), 13.56 (s, 1H). MS m/z 529 [M+H]⁺

Example 31

6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-4-({5-hydroxy-2-[methyl
(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-
carboxamide

To a solution of N-[2-({[6-(2-ethyl-5-fluoro-4-{{2-(trimethylsilyl)-ethoxy}meth-
oxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]-pyridin-4-yl]-
amino)methyl)-4-methoxyphenyl]-N-methylmethanesulfonamide (**Preparation 83**, 250

mg, 0.28 mmol) in 2M methylamine in THF (3 mL) was added DBU (0.13 mL, 0.85 mmol), palladium acetate (4.43 mg, 0.02 mmol) and molybdenum hexacarbonyl (75 mg, 0.28 mmol) and the reaction was heated to 100 °C for 10 minutes under microwave irradiation. The reaction was cooled, concentrated *in vacuo* and purified directly using silica gel column chromatography eluting with 45% EtOAc in hexanes. The resulting oil was dissolved in DCM (15 mL) and cooled to 0°C. BBr₃ (0.10 mL, 1.07 mmol) was added and the reaction stirred at room temperature for 6 hours. The reaction was concentrated *in vacuo* and partitioned between saturated aqueous sodium bicarbonate solution and EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC eluting with 5% MeOH in DCM to afford the title compound as a white solid (43 mg, 27% over two steps). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.88 (t, 3H), 2.84 (d, 3H), 2.98 (s, 3H), 3.05 (s, 3H), 3.31 (m, 2H), 4.61 (m, 1H), 4.86 (m, 1H), 6.66 (m, 2H), 6.78 (m, 2H), 7.05 (d, 1H), 7.26 (d, 1H), 8.80 (m, 1H), 9.51 (s, 1H), 9.59 (m, 1H), 9.74 (s, 1H), 13.62 (s, 1H). MS m/z 543 [M+H]⁺

Example 32

6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[(3-hydroxyphenyl)sulfonyl](methyl)amino)benzyl)amino]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of N-(2-[(6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-[(2-(trimethylsilyl)ethoxy)methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl)amino]methyl)-phenyl)-3-methoxy-N-methylbenzenesulfonamide (**Preparation 106**, 330 mg, 0.37 mmol) in 2M methylamine in THF (2 mL) was added DBU (0.16 mL, 1.19 mmol), palladium acetate (5.86 mg, 0.03 mmol) and molybdenum hexacarbonyl (99 mg, 0.37 mmol) and the reaction was heated to 100 °C for 10 minutes under microwave irradiation. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 35% EtOAc in hexanes. The resulting oil was treated with TFA (0.5 mL) and the solution stirred at room temperature for 30 minutes before concentrating *in vacuo*. Ethylene diamine (0.5 mL) was added and the reaction stirred at room temperature for 15 minutes before pouring onto ice-water and extracting into 20% IPA in DCM. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 45% EtOAc in hexanes. The residue was dissolved in DCM (10 mL) and boron tribromide (0.18 mL, 1.89 mmol) was added dropwise at 0°C.

The reaction was stirred at room temperature for 2 hours followed by quenching with saturated aqueous sodium bicarbonate solution and extracting into 20% IPA in DCM. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography followed by
 5 preparative TLC eluting both with 57% EtOAc in hexanes to afford the title compound as a yellow solid (25 mg, 10% over 3 steps).

¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.85 (d, 3H), 3.01 (s, 3H), 3.59-3.65 (m, 2H), 4.71 (m, 1H), 4.95 (m, 1H), 6.63 (d, 1H), 6.71 (s, 1H), 6.96 (d, 1H), 7.02 (m, 1H), 7.09 (m, 2H), 7.14-7.29 (m, 3H), 7.41 (m, 2H), 8.86 (m, 1H), 9.78 (m, 1H), 10.10 (s, 1H),
 10 10.15 (s, 1H), 13.71 (s, 1H). MS m/z 659 [M+H]⁺

The following Examples (**Examples 33 – 37**) were prepared according to the method described for **Example 32** using the appropriate pyrazolo-pyridine, and Purification Method (PM) below if different from the method described:

Purification Method F: Silica gel column chromatography or preparative TLC eluting
 15 with 4% MeOH in DCM.

Ex	Name	Data
33	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({4-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 597 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.83 (d, 3H), 3.02 (s, 3H), 3.17 (s, 3H), 3.77 (m, 2H), 4.70 (br m, 2H), 6.69-6.73 (m, 2H), 6.83 (m, 1H), 6.97 (m, 1H), 7.20-7.25 (m, 2H), 8.80 (m, 1H), 9.56 (s, 1H), 9.60 (m, 1H), 10.10 (s, 1H), 13.67 (s, 1H). Using N-(2-{[(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl)amino]methyl}-5-methoxyphenyl)-N-methylmethane-sulfonamide (Preparation 89).
34	4-({2-[ethyl(methylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 611 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.90 (t, 3H), 2.88 (d, 3H), 2.96 (s, 3H), 3.39-3.51 (m, 2H), 3.54-3.68 (m, 2H), 4.60 (m, 1H), 4.80 (m, 1H), 6.66 (m, 1H), 6.69 (s, 1H), 6.78 (s, 1H), 6.93 (d, 1H), 7.20 (m, 2H), 7.95 (br s, 1H), 8.85 (m, 1H), 9.50 (s, 1H), 9.70 (m, 1H), 10.07 (br s, 1H), 13.71 (br s, 1H). Using N-ethyl-N-(2-{[(6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl)amino]methyl}-4-methoxyphenyl)methanesulfonamide (Preparation 107).

35	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(phenylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 659 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.86 (d, 3H), 2.97 (s, 3H), 3.63 (m, 2H), 4.58 (m, 1H), 4.86 (m, 1H), 6.34 (m, 1H), 6.47 (m, 1H), 6.71 (s, 1H), 6.77 (s, 1H), 6.93 (d, 1H), 7.23 (d, 1H), 7.58-7.73 (m, 5H), 8.85 (m, 1H), 9.51 (s, 1H), 9.71 (m, 1H), 10.07 (s, 1H), 13.70 (s, 1H). Using N-(2-({[6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-{2-(trimethylsilyl)ethoxy}-methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}-4-methoxy-phenyl)-N-methylbenzene-sulfonamide (Preparation 108).
36	4-({2-[ethyl(phenylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 673 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.85 (t, 3H), 2.86 (d, 3H), 3.22 (m, 1H), 3.53 (m, 1H), 3.68 (m, 2H), 4.52 (m, 1H), 4.81 (m, 1H), 6.41 (d, 1H), 6.48 (dd, 1H), 6.70 (s, 1H), 6.78 (d, 1H), 6.94 (d, 1H), 7.21 (d, 1H), 7.56-7.70 (m, 5H), 8.83 (m, 1H), 9.52 (s, 1H), 9.67 (m, 1H), 10.06 (s, 1H), 13.71 (s, 1H). Using N-ethyl-N-(2-({[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-3-iodo-1-{2-(trimethylsilyl)ethoxy}-methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}-4-methoxyphenyl)benzenesulfonamide (Preparation 90).
37	6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 538 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.45 (m, 2H), 0.61 (m, 2H), 2.09 (m, 1H), 2.84 (s, 3H), 3.06 (s, 3H), 3.14 (s, 3H), 4.70 (br m, 1H), 5.00 (br m, 1H), 6.46 (m, 1H), 6.79 (s, 1H), 7.70 (m, 1H), 7.31 (m, 2H), 7.41 (m, 1H), 7.50 (m, 1H), 8.78 (m, 1H), 9.62 (m, 1H), 9.73 (m, 1H), 13.61 (m, 1H). Using N-[2-({[6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1-{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide (Preparation 109) and PM F.

Examples 38 and 39

6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({(1R)-1-{2-[methyl(methylsulfonyl)amino]phenyl}ethyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide and 6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({(1S)-1-{2-[methyl(methylsulfonyl)amino]phenyl}ethyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compounds were prepared according to the method described for **Example 1** using racemic N-[2-(1-{{[6-(2-ethyl-5-fluoro-4-{2-(trimethylsilyl)ethoxy]-

methoxy}phenyl)-3-iodo-1-{{2-(trimethyl-silyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}ethyl)phenyl]-N-methylmethanesulfonamide (**Preparation 84**). The residue was purified using silica gel column chromatography eluting with 6% MeOH in DCM followed by chiral separation using chiral preparative HPLC to afford the separated
 5 enantiomers.

Fraction 1: 44 mg, 100% ee, registered as (R) – enantiomer **Example 38**

¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.86 (t, 3H), 1.53 (d, 3H), 2.21 (m, 2H), 2.83 (s, 3H), 2.90 (d, 3H), 3.08 (s, 3H), 5.47 (m, 1H), 6.50 (s, 1H), 6.73 (m, 2H), 7.21-7.45 (m, 4H), 8.82 (m, 1H), 9.69 (br s, 1H), 9.86 (m, 1H), 13.57 (br s, 1H). MS m/z 541
 10 [M+H]⁺

Fraction 2: 41 mg, 87.5% ee, registered as (S) – enantiomer **Example 39**

¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.86 (t, 3H), 1.53 (d, 3H), 2.21 (m, 2H), 2.83 (s, 3H), 2.90 (d, 3H), 3.08 (s, 3H), 5.47 (m, 1H), 6.50 (s, 1H), 6.73 (m, 2H), 7.21-7.45 (m, 4H), 8.82 (m, 1H), 9.69 (br s, 1H), 9.86 (m, 1H), 13.57 (br s, 1H). MS m/z 541
 15 [M+H]⁺

Example 40

6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-[methyl(phenylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

20 To a solution of N-[2-{{6-(2-ethyl-5-fluoro-4-{{2-(trimethylsilyl)ethoxy}-methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylbenzenesulfonamide (**Preparation 81**, 400 mg, 0.44 mmol) in MeOH (2 mL) was added DBU (0.20 mL, 1.31 mmol), palladium acetate (6.85 mg, 0.03 mmol) and molybdenum hexacarbonyl (115 mg, 0.44 mmol) and the reaction
 25 was heated to 125 °C under microwave irradiation for 20 minutes. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 70% EtOAc in hexanes. The residue was dissolved in THF (4 mL) and cooled to -20°C. NMM (0.021 mL, 0.19 mmol) followed by isobutylchloroformate (0.03 mL, 0.19 mmol) were added and the reaction stirred at this temperature for 2 hours.
 30 Aqueous ammonia was then added and the reaction stirred at room temperature for 1 hour. The reaction was quenched by the addition of water and extracted into EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 40% EtOAc in hexanes. The residue was treated with TFA (2 mL) and stirred for 2 hours

before concentrating *in vacuo*. Ethylene diamine (0.5 mL) was added and the reaction stirred at room temperature for 1 hour before concentrating *in vacuo*, pouring onto ice-water and extracting into EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using Preparative TLC
 5 eluting with 60% EtOAc in hexanes to afford the title compound (23 mg, 10% over 3 steps).

¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.92 (t, 3H), 3.04 (s, 3H), 3.40 (m, 2H), 4.70 (br m, 1H), 5.00 (br m, 1H), 6.55 (d, 1H), 6.64 (s, 1H), 6.80 (d, 1H), 7.06 (m, 1H), 7.14 (m, 1H), 7.29 (m, 1H), 7.47 (m, 1H), 7.59-7.67 (m, 4H), 7.73 (m, 1H), 7.83 (br s, 1H), 8.18 (br s, 1H), 9.65 (m, 1H), 9.77 (s, 1H), 13.60 (br s, 1H). MS m/z 575 [M+H]⁺
 10

Example 41

6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({4-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

A solution of 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-({2-(trimethylsilyl)ethoxy}methoxy)phenyl]-4-({4-methoxy-2-[methyl(methylsulfonyl)amino]-benzyl}-amino)-1-([2-(trimethylsilyl)ethoxy]-methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide
 15 (Preparation 21, 102 mg, 0.15 mmol) in TFA (5 mL) was stirred at room temperature for 30 minutes before concentrating *in vacuo*. Ethylene diamine (0.5 mL) was added and the reaction stirred at room temperature for 15 minutes before concentrating *in vacuo*, pouring onto ice-water and extracting into EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC to afford a white solid. The solid was dissolved in DCM (5 mL) and boron tribromide (0.108 mL, 1.14 mmol) was added dropwise at 0 °C and stirred at room temperature for 2 hours. The reaction was washed with saturated
 20 aqueous sodium bicarbonate solution, the organic layer collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC eluting with 5% MeOH in DCM to afford the title compound (37 mg, 54%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.02 (s, 3H), 3.08 (s, 3H), 3.79 (m, 2H), 4.65 (br m, 2H), 6.69-6.73 (m, 2H), 6.83 (m, 1H), 6.99 (d, 1H), 7.20-7.25 (m, 2H), 7.81 (s, 1H), 8.17 (s, 1H), 9.59 (m, 2H), 10.10 (br s, 1H), 13.70 (br s, 1H). MS m/z 583 [M+H]⁺
 30

The following Examples (Examples 42 – 54) were prepared according to the method described for Example 41 using the appropriate pyrazolo-pyridine, and Purification Method (PM) as described below if different from the method described:

Purification Method G: Silica gel column chromatography eluting with 5 – 7% MeOH in DCM followed by preparative HPLC.

Ex	Name	Data
42	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(phenylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 645 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.98 (s, 3H), 3.69 (m, 2H), 4.57 (m, 1H), 4.83 (m, 1H), 6.35 (m, 1H), 6.47 (m, 1H), 6.71 (s, 1H), 6.79 (d, 1H), 6.96 (d, 1H), 7.24 (d, 1H), 7.58-7.73 (m, 5H), 7.87 (br s, 1H), 8.22 (br s, 1H), 9.53 (s, 1H), 9.71 (m, 1H), 10.09 (s, 1H), 13.69 (s, 1H). Using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-methoxy-2-[methyl(phenylsulfonyl)amino]benzyl}amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 24).
43	4-({2-[ethyl(phenylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 657 [M-H] ⁻ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.85 (t, 3H), 3.23 (m, 1H), 3.67 (m, 1H), 3.70 (m, 2H), 4.50 (m, 1H), 4.79 (m, 1H), 6.40 (m, 1H), 6.50 (m, 1H), 6.70 (s, 1H), 6.80 (s, 1H), 6.95 (d, 1H), 7.22 (d, 1H), 7.55-7.68 (m, 5H), 7.85 (br s, 1H), 8.20 (br s, 1H), 9.52 (s, 1H), 9.68 (m, 1H), 10.05 (s, 1H), 13.67 (br s, 1H). Using N-ethyl-N-(2-{{(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)-ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl}-4-methoxyphenyl)benzenesulfonamide (Preparation 25).
44	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({3-[methyl(phenylsulfonyl)amino]pyrazin-2-yl}methyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 645 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.87 (m, 6H), 3.62 (m, 2H), 5.06 (m, 2H), 6.68 (s, 1H), 6.97 (d, 1H), 7.15 (d, 1H), 7.61 (m, 4H), 7.73 (m, 1H), 8.33 (s, 1H), 8.58 (s, 1H), 8.78 (m, 1H), 9.89 (m, 1H), 10.08 (s, 1H), 13.67 (s, 1H). Using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({3-[methyl(phenylsulfonyl)amino]pyrazin-2-yl}methyl)amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 39).

45	4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methylamino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 597 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.85 (t, 3H), 2.85 (d, 3H), 3.10 (s, 3H), 3.49 (q, 2H), 3.69 (q, 2H), 4.99 (m, 2H), 6.68 (s, 1H), 6.95 (d, 1H), 7.10 (d, 1H), 8.54 (s, 1H), 8.61 (s, 1H), 8.79 (m, 1H), 9.82 (m, 1H), 10.07 (s, 1H), 13.67 (s, 1H). Using 4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methylamino]-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 40) and PM G.
46	N-ethyl-4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methylamino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 611 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.86 (t, 3H), 1.14 (t, 3H), 3.35 (m, 2H), 3.53 (m, 2H), 3.67 (m, 2H), 4.99 (m, 2H), 6.68 (s, 1H), 6.94 (d, 1H), 7.13 (d, 1H), 8.54 (s, 1H), 8.60 (s, 1H), 8.83 (m, 1H), 9.81 (m, 1H), 10.07 (s, 1H), 13.68 (s, 1H). Using N-ethyl-4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methylamino]-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 41).
47	6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-4-[(2-[methyl(methylsulfonyl)amino]benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 525 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.46 (m, 2H), 0.65 (m, 2H), 2.09 (m, 1H), 3.05 (s, 3H), 3.14 (s, 3H), 4.70 (br m, 1H), 5.00 (br m, 1H), 6.45 (d, 1H), 6.80 (s, 1H), 7.09 (d, 1H), 7.32 (m, 2H), 7.45 (m, 2H), 7.81 (s, 1H), 8.16 (s, 1H), 9.62 (m, 1H), 9.73 (s, 1H), 13.59 (br s, 1H). Using 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-4-[(2-[methyl(methylsulfonyl)amino]benzyl)amino]-1-[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 26) and PM G.
48	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(4-[methyl(methylsulfonyl)amino]pyridin-3-yl)methylamino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 582 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.83 (s, 3H), 2.90 (s, 3H), 3.70 (s, 3H), 3.85 (m, 2H), 4.47 (m, 2H), 6.72 (s, 1H), 7.00 (d, 1H), 7.20-7.26 (m, 2H), 7.77 (s, 1H), 7.87 (m, 1H), 8.78 (m, 1H), 9.67 (m, 1H), 10.13 (s, 1H), 13.67 (s, 1H). Using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(4-[methyl(methylsulfonyl)amino]pyridin-3-yl)methylamino]-1-[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 42).

49	6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 540 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.45 (m, 2H), 0.61 (m, 2H), 2.00 (m, 1H), 2.84 (d, 3H), 3.11 (s, 6H), 4.85 (m, 2H), 6.46 (d, 1H), 6.81 (s, 1H), 7.02 (d, 1H), 7.37 (m, 1H), 7.83 (d, 1H), 8.40 (m, 1H), 8.80 (m, 1H), 9.69-9.73 (m, 2H), 13.64 (s, 1H). Using 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-N-methyl-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 110).
50	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(5-methoxy-2-[methyl(methylsulfonyl)amino]benzyl)amino]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 597 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.86 (d, 3H), 2.98 (s, 3H), 3.04 (s, 3H), 3.64 (m, 2H), 4.61 (m, 1H), 4.82 (m, 1H), 6.63 (dd, 1H), 6.70 (s, 1H), 6.76 (m, 1H), 6.95 (d, 1H), 7.19-7.27 (m, 2H), 8.84 (m, 1H), 9.50 (s, 1H), 9.70 (m, 1H), 10.09 (s, 1H), 13.70 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-4-[(5-methoxy-2-[methyl(methylsulfonyl)amino]benzyl)amino]-N-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 46).
51	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 583 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.86 (d, 3H), 3.06 (s, 3H), 3.13 (s, 3H), 3.60 (m, 2H), 4.95 (m, 2H), 6.68 (s, 1H), 6.96 (d, 1H), 7.16 (d, 1H), 8.49 (s, 1H), 8.59 (s, 1H), 8.80 (m, 1H), 9.85 (m, 1H), 10.10 (s, 1H), 13.68 (s, 1H). Using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 47).
52	4-[(2-[ethyl(methylsulfonyl)-amino]pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 582 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.86 (t, 3H), 3.07 (s, 3H), 3.55-3.67 (m, 4H), 4.84 (m, 2H), 6.72 (s, 1H), 6.95 (d, 1H), 7.19 (d, 1H), 7.38 (m, 1H), 7.82 (d, 1H), 7.87 (br s, 1H), 8.22 (br s, 1H), 8.44 (m, 1H), 9.80 (m, 1H), 10.07 (s, 1H), 13.71 (s, 1H). Using N-(2,4-dimethoxybenzyl)-4-[(2-[ethyl(methylsulfonyl)-amino]pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 52).

53	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 582 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.86 (d, 3H), 3.08 (s, 3H), 3.12 (s, 3H), 3.62 (q, 2H), 4.83 (m, 2H), 6.73 (s, 1H), 6.95 (d, 1H), 7.21 (d, 1H), 7.36 (m, 1H), 7.79 (d, 1H), 8.41 (m, 1H), 8.85 (m, 1H), 9.80 (m, 1H), 10.10 (s, 1H), 13.73 (s, 1H). Using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 53).
54	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 660 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.86 (d, 3H), 3.02 (s, 3H), 3.61 (m, 2H), 4.61 (m, 1H), 4.83 (m, 1H), 6.39 (d, 1H), 6.50 (m, 1H), 6.71 (s, 1H), 6.80 (m, 1H), 6.95 (d, 1H), 7.16-7.24 (m, 2H), 7.65 (m, 1H), 8.06 (m, 1H), 8.78 (s, 1H), 8.87 (m, 1H), 9.57 (s, 1H), 9.74 (m, 1H), 10.08 (s, 1H), 13.71 (br s, 1H). Using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-methoxy-2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-N-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 54).

Example 55

6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)amino]pyridin-2-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

5

A solution of 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[(2-(trimethylsilyl)ethoxy)methoxy]phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)amino]pyridin-2-yl)methyl]amino]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 43**, 110 mg, 0.13 mmol) in TFA (5 mL) was stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo*, dissolved in MeOH and cooled in ice-water. Ethylene diamine was added dropwise and stirred for 1 hour. The reaction was quenched by the addition of water and extracted into EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using Preparative HPLC to afford the title compound (18 mg, 26%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.84 (d, 3H), 3.07 (s, 6H), 3.61 (m, 2H), 4.70-5.20 (br m, 2H), 6.66 (s, 1H), 6.94 (d, 1H),

10

15

7.16 (d, 1H), 7.37 (m, 1H), 7.92 (m, 1H), 8.47 (m, 1H), 9.77 (m, 1H), 10.08 (br s, 1H), 13.63 (br s, 1H). MS m/z 582 [M+H]⁺

The following Examples (**Examples 56 – 73**) were prepared according to the method described for **Example 55** using the appropriate pyrazolo-pyridine and Purification Method below if different from the method described

Purification Method H: Preparative TLC.

Purification Method I: Preparative HPLC.

Ex	Name	Data
56	6-(2-ethyl-5-fluoro-4-hydroxy-phenyl)-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-(6-methylpyridin-3-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 604 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.89 (t, 3H), 2.44 (s, 3H), 3.04 (s, 3H), 3.11 (s, 3H), 3.32 (m, 2H), 4.75 (m, 1H), 5.00 (m, 1H), 6.70 (s, 1H), 6.78 (d, 1H), 7.06 (d, 1H), 7.24-7.32 (m, 3H), 7.42-7.49 (m, 2H), 8.13 (dd, 1H), 8.87 (d, 1H), 9.25 (t, 1H), 9.79 (s, 1H), 10.83 (s, 1H), 13.89 (s, 1H). Using 6-(2-ethyl-5-fluoro-4-({2-(trimethylsilyl)ethoxy}methoxy}phenyl)-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-(6-methylpyridin-3-yl)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 20) and PM H.
57	6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((1,3,3-trimethylureido)benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 610 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.83 (d, 3H), 3.00 (s, 3H), 3.64 (m, 2H), 4.84 (br m, 2H), 6.70 (s, 1H), 6.96 (d, 1H), 7.19-7.46 (m, 5H), 8.83 (m, 1H), 9.74 (m, 1H), 10.09 (s, 1H), 13.70 (s, 1H). Using 6-(5-fluoro-2-(2,2,2-trifluoro-ethyl)-4-((2-(trimethylsilyl)ethoxy)-methoxy)phenyl)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-4-((2-(1,3,3-trimethylureido)benz-yl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 13) and PM H.

58	6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(N-methyl-1H-pyrazole-4-sulfonamido)benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 633 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.85 (d, 3H), 2.99 (s, 3H), 3.66 (m, 2H), 4.74 (m, 1H), 4.97 (m, 1H), 6.71 (m, 2H), 6.96 (m, 1H), 7.17-7.28 (m, 3H), 7.40 (m, 1H), 7.73 (s, 1H), 8.30 (s, 1H), 8.83 (m, 1H), 9.76 (m, 1H), 10.10 (s, 1H), 13.71 (s, 1H), 13.75 (s, 1H). Using 6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-trimethylsilyl)ethoxy)methoxy)phenyl)-N-methyl-4-((2-(N-methyl-1H-pyrazole-4-sulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 3) and PM H.
59	4-((2-N,1-dimethyl-1H-imidazole-4-sulfonamido)benzyl)amido)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 647 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.84 (d, 3H), 3.08 (s, 3H), 3.70 (m, 5H), 4.68 (m, 1H), 4.95 (m, 1H), 6.71 (s, 1H), 6.96 (m, 2H), 7.16-7.26 (m, 3H), 7.36 (m, 1H), 7.72 (s, 1H), 7.88 (s, 1H), 8.82 (m, 1H), 9.73 (m, 1H), 10.08 (s, 1H), 13.69 (s, 1H). Using 4-((2-(N,1-dimethyl-1H-imidazole-4-sulfonamido)benzyl)amino)-6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 8) and PM H.
60	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-((2-methoxyethyl)sulfonyl)(methyl)amino)benzyl)amino]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 625 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.84 (d, 3H), 3.09 (s, 3H), 3.26 (s, 3H), 3.44-3.68 (m, 6H), 4.73-4.86 (m, 2H), 6.70 (s, 1H), 6.94 (d, 1H), 7.21 (d, 1H), 7.30 (m, 2H), 7.38 (m, 1H), 7.50 (m, 1H), 8.83 (m, 1H), 9.77 (m, 1H), 10.08 (br s, 1H), 13.70 (br s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-4-[(2-((2-methoxyethyl)sulfonyl)(methyl)amino)benzyl)amino]-N-methyl-1-([2-(trimethylsilyl)ethoxy)methyl]-1H-pyrazolo-[4,3-c]pyridine-3-carboxamide (Preparation 9) and PM H using 50% EtOAc in hexanes.

61	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 630 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.08 (s, 3H), 3.69 (m, 2H), 4.71 (m, 1H), 4.92 (m, 1H), 5.75 (d, 1H), 6.71 (s, 1H), 6.97 (m, 1H), 7.17 (m, 1H), 7.24 (d, 1H), 7.30 (t, 1H), 7.44 (d, 1H), 7.65-7.86 (br s, 1H), 8.07 (m, 1H), 8.21 (br s, 1H), 8.79 (m, 1H), 8.89 (m, 1H), 9.78 (t, 1H), 10.09 (s, 1H), 13.69 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-4-{{2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 10).
62	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 568 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.09 (s, 3H), 3.12 (s, 3H), 3.65 (m, 2H), 4.81 (m, 2H), 6.73 (s, 1H), 6.96 (d, 1H), 7.21 (d, 1H), 7.37 (m, 1H), 7.82 (m, 1H), 7.86 (br s, 1H), 8.21 (br s, 1H), 8.41 (m, 1H), 9.81 (t, 1H), 10.09 (s, 1H), 13.69 (s, 1H). Using 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 15).
63	4-[(2-[ethyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 596 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.84 (t, 3H), 2.83 (d, 3H), 3.05 (s, 3H), 3.52-3.64 (m, 4H), 4.83 (m, 2H), 6.70 (s, 1H), 6.91 (d, 1H), 7.15 (d, 1H), 7.35 (m, 1H), 7.76 (m, 1H), 8.42 (m, 1H), 8.83 (m, 1H), 9.76 (m, 1H), 10.04 (s, 1H), 13.71 (br s, 1H). Using -[(2-[ethyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-N-methyl-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 48) and PM I.
64	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-{{2-(sulfamoylmethyl)benzyl}amino}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 567 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.84 (d, 3H), 3.76 (q, 2H), 4.42 (s, 2H), 4.84 (m, 2H), 6.71 (s, 1H), 6.87 (s, 2H), 6.98 (d, 1H), 7.20-7.36 (m, 5H), 8.83 (m, 1H), 9.75 (m, 1H), 10.10 (br s, 1H), 13.70 (br s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-N-methyl-4-{{2-(sulfamoylmethyl)benzyl}amino}-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 49).

65	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-{{2-(methyl{[6-(morpholin-4-yl)pyridin-3-yl]sulfonyl}amino)benzyl}amino}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 715 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.00 (s, 3H), 3.61-3.74 (m, 10H), 4.73 (m, 1H), 4.94 (m, 1H), 6.71 (s, 1H), 6.70 (d, 1H), 6.91 (d, 1H), 6.95 (d, 1H), 7.19-7.30 (m, 3H), 7.43 (m, 1H), 7.64 (m, 1H), 7.86 (br s, 1H), 8.21 (br s, 1H), 8.28 (m, 1H), 9.75 (m, 1H), 10.10 (s, 1H), 13.68 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-4-{{2-(methyl{[6-(morpholin-4-yl)pyridin-3-yl]sulfonyl}amino)benzyl}amino}-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 5).
66	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-{{2-(methyl{[3-(morpholin-4-yl)propyl]sulfonyl}amino)benzyl}amino}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 680 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.81 (m, 2H), 2.29 (m, 6H), 3.12 (s, 3H), 3.23 (m, 2H), 3.57 (m, 4H), 3.62 (m, 2H), 4.70-5.00 (m, 2H), 6.70 (s, 1H), 6.95 (d, 1H), 7.22 (d, 1H), 7.31-7.34 (m, 2H), 7.41-7.46 (m, 2H), 7.84 (s, 1H), 8.20 (s, 1H), 9.75 (m, 1H), 10.08 (s, 1H), 13.67 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-4-{{2-(methyl{[3-(morpholin-4-yl)propyl]sulfonyl}amino)benzyl}amino}-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 2).
67	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(5-methyl-2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 596 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.21 (s, 3H), 2.86 (d, 3H), 3.05 (s, 3H), 3.09 (s, 3H), 3.54 (m, 2H), 4.77 (m, 2H), 6.72 (s, 1H), 6.95 (d, 1H), 7.19 (d, 1H), 7.59 (s, 1H), 8.23 (s, 1H), 8.83 (m, 1H), 9.78 (m, 1H), 10.09 (br s, 1H), 13.72 (br s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-N-methyl-4-[(5-methyl-2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 44).

68	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 644 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.84 (d, 3H), 3.07 (s, 3H), 3.62 (m, 2H), 4.74 (m, 1H), 4.93 (m, 1H), 6.66 (d, 1H), 6.71 (s, 1H), 6.96 (d, 1H), 7.15-7.23 (m, 2H), 7.28 (t, 1H), 7.42 (m, 1H), 7.65 (m, 1H), 8.06 (m, 1H), 8.80-8.83 (m, 2H), 8.89 (m, 1H), 9.76 (m, 1H), 10.07 (s, 1H), 13.70 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoro-ethyl)-4-{{2-(trimethylsilyl)-ethoxy}methoxy}phenyl]-N-methyl-4-({2-[methyl(pyridin-3-ylsulfonyl)-amino]benzyl}amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 45) and PM H using 4% MeOH in DCM.
69	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino}benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 658 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.58 (s, 3H), 2.85 (d, 3H), 3.05 (s, 3H), 3.65 (m, 2H), 4.74 (m, 1H), 4.93 (m, 1H), 6.68 (d, 1H), 6.71 (s, 1H), 6.96 (d, 1H), 7.16-7.23 (m, 2H), 7.29 (t, 1H), 7.43 (d, 1H), 7.50 (d, 1H), 7.93 (m, 1H), 8.65 (m, 1H), 8.82 (m, 1H), 9.75 (m, 1H), 10.08 (s, 1H), 13.70 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-N-methyl-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino}benzyl)amino]-1-{{2-(trimethylsilyl)-ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 6).
70	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino}benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 644 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.58 (s, 3H), 3.06 (s, 3H), 3.67 (m, 2H), 4.71 (m, 1H), 4.92 (m, 1H), 6.69 (d, 1H), 6.71 (s, 1H), 6.97 (d, 1H), 7.16-7.24 (m, 2H), 7.30 (t, 1H), 7.46 (d, 1H), 7.50 (d, 1H), 7.91 (br s, 1H), 7.93 (dd, 1H), 8.21 (br s, 1H), 8.65 (m, 1H), 9.77 (m, 1H), 10.09 (s, 1H), 13.68 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino}benzyl)amino]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 7) and using aqueous ammonia instead of ethylene diamine.

71	4-[(5-chloro-2-ethyl(methylsulfonyl)amino)pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 616 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.85 (t, 3H), 3.49-3.65 (m, 4H), 4.81 (m, 2H), 6.74 (s, 1H), 6.95 (d, 1H), 7.18 (d, 1H), 7.81 (m, 1H), 7.90 (br s, 1H), 8.24 (br s, 1H), 8.51 (m, 1H), 9.80 (m, 1H), 10.08 (s, 1H), 13.74 (s, 1H). Using N-tert-butyl-4-[(5-chloro-2-ethyl(methylsulfonyl)amino)pyridin-3-yl)methyl]amino]-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy} phenyl]-1-{2-(trimethylsilyl)-ethoxy}methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 50).
72	4-[(5-chloro-2-methyl(methylsulfonyl)amino)pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 602 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.07 (s, 3H), 3.16 (s, 3H), 3.58 (q, 2H), 4.79 (m, 2H), 6.75 (s, 1H), 6.96 (d, 1H), 7.18 (d, 1H), 7.80 (d, 1H), 7.89 (br s, 1H), 8.23 (br s, 1H), 8.47 (d, 1H), 9.82 (m, 1H), 10.10 (s, 1H), 13.73 (s, 1H). Using N-tert-butyl-4-[(5-chloro-2-methyl(methylsulfonyl)amino)pyridin-3-yl)methyl]amino]-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy} phenyl]-1-{2-(trimethylsilyl)-ethoxy}methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 51).
73	6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(sulfamoyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 528 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.87 (t, 3H), 2.53 (m, 2H), 2.84 (s, 3H), 3.01 (s, 3H), 3.39 (m, 2H), 4.70 (br s, 1H), 5.00 (br s, 1H), 6.54 (s, 1H), 6.79 (d, 1H), 7.03 (m, 2H), 7.24 (m, 1H), 7.37 (m, 1H), 7.43 (m, 1H), 8.78 (m, 1H), 9.60 (m, 1H), 9.75 (s, 1H), 13.60 (s, 1H). Using 6-(2-ethyl-5-fluoro-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-N-methyl-4-({2-[methyl(sulfamoyl)-amino]benzyl}amino)-1-{2-(trimethylsilyl)-ethoxy}methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 1).

Example 74

6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((N-(2-hydroxyethyl)-sulfamoyl)(methyl)aminobenzyl)amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

5

The title compound was prepared according to the method described for **Example 55** using 6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)-methoxy)phenyl)-N-methyl-4-((2-(N-methyl-2-oxooxazolidine-3-sulfonamido)benzyl)-amino)-1-((2-(trimethylsilyl)-ethoxy)-methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

10 (**Preparation 58**). The residue was treated with 6M NaOH (0.5 mL) at 0 °C and stirred

at room temperature for 18 hours. The reaction was acidified with HCl at 0°C, and the resulting precipitate was filtered, extracted into EtOAc and concentrated *in vacuo*. The residue was purified using preparative TLC. ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.07 (m, 3H), 2.84 (s, 3H), 3.02 (s, 3H), 3.39 (m, 2H), 3.69 (m, 2H), 4.70 (m, 2H), 4.90 (br m, 2H), 6.94 (s, 1H), 6.96 (m, 1H), 7.20-7.46 (m, 5H), 8.82 (m, 1H), 9.71 (m, 1H), 10.10 (br s, 1H), 13.70 (s, 1H). MS m/z 626 [M+H]⁺

Example 75

6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-{6-[(2-hydroxyethyl)amino]pyridin-3-yl}-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-4-({5-methoxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1-{{2-(trimethyl-silyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxylic acid (**Preparation 11**, 100 mg, 0.12 mmol) in DCM (3 mL) was added boron tribromide (0.08 mL, 0.82 mmol) and the reaction was stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo* and triturated with ether/pentane. The resulting solid was dissolved in DMF (2 mL) and 2-[(5-aminopyridin-2-yl)amino]ethanol (51 mg, 0.33 mmol) followed by DIPEA (0.07 mL, 0.17 mmol) were added. HATU (159 mg, 0.42 mmol) was added and the reaction stirred at room temperature for 18 hours. The reaction was partitioned between EtOAc and water, the organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC to afford the title compound (15 mg, 13%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.98 (s, 3H), 3.03 (s, 3H), 3.32 (m, 2H), 3.52 (m, 2H), 3.64 (m, 2H), 4.60 (m, 1H), 4.71 (m, 1H), 4.83 (m, 1H), 6.45 (m, 1H), 6.53 (d, 1H), 6.63 (dd, 1H), 6.75 (s, 1H), 6.78 (m, 1H), 6.96 (d, 1H), 7.21-7.26 (m, 2H), 7.78 (dd, 1H), 8.31 (s, 1H), 9.46 (m, 2H), 10.10 (s, 1H), 10.48 (s, 1H), 13.86 (s, 1H). MS m/z 719 [M+H]⁺

Example 76

6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(methylsulfonyl)amino]-benzyl}amino)-N-(6-methylpyridin-3-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}-phenyl]-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-1-{{2-(trimethyl-silyl)ethoxy}-methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxylic acid (**Preparation 12**, 100 mg, 0.12 mmol) in DMF (3 mL) was added 6-methylpyridin-3-amine (65 mg, 0.60 mmol),

DIPEA (0.13 mL, 0.73 mmol) and BOP (267 mg, 0.60 mmol) and the reaction was stirred at room temperature for 18 hours before concentrating *in vacuo*. The residue was partitioned between ice-water and EtOAc, the organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 5% MeOH in DCM. The residue was treated with TFA (3 mL) and stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo*, dissolved in MeOH and cooled in ice-water. Ethylene diamine was added until the solution became basic, with stirring for 15 minutes. The solution was concentrated *in vacuo* and purified using preparative HPLC to afford the title compound (35 mg, 29%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.45 (s, 3H), 3.04 (s, 3H), 3.11 (s, 3H), 3.62 (m, 2H), 4.80 (m, 1H), 4.95 (m, 1H), 6.77 (s, 1H), 6.96 (d, 1H), 7.20-7.33 (m, 4H), 7.41 (m, 1H), 7.49 (m, 1H), 8.13 (dd, 1H), 8.87 (d, 1H), 9.33 (t, 1H), 10.85 (br s, 1H). MS m/z 658 [M+H]⁺

Example 77

6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide 6-(4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-methylmethyl-sulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 27**, 60 mg, 0.07 mmol) was treated with TFA (8 mL) and heated to reflux for 18 hours. The reaction was cooled, concentrated *in vacuo*, quenched by the addition of saturated aqueous sodium bicarbonate solution and extracted into EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The crude residue was purified using preparative TLC to afford the title compound (21 mg, 51%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.04 (s, 3H), 3.14 (s, 3H), 3.69 (m, 2H), 4.70 (br m, 1H), 4.90 (m, 1H), 6.70 (s, 1H), 6.96 (d, 1H), 7.22 (d, 1H), 7.27-7.34 (m, 2H), 7.40 (m, 1H), 7.49 (m, 1H), 7.85 (br s, 1H), 8.20 (br s, 1H), 9.75 (m, 1H), 10.09 (s, 1H), 13.70 (s, 1H). MS m/z 567 [M+H]⁺

Example 78

6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-fluoro-2-[methyl(methyl-sulfonyl)amino]pyridin-3-yl}methyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

6-(4-Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((5-fluoro-2-(N-methylmethylsulfonamido)pyridin-3-yl)methyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo-[4,3-c]pyridine-3-carboxamide (**Preparation 55**, 80 mg, 0.10 mmol) was

treated with TFA (10 mL) and heated to reflux for 18 hours. The reaction was cooled, concentrated *in vacuo* and partitioned between saturated aqueous sodium bicarbonate solution and EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC to afford the title compound (20 mg, 35%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.07 (s, 3H), 3.12 (s, 3H), 3.57 (q, 2H), 4.80 (m, 2H), 6.75 (s, 1H), 6.95 (d, 1H), 7.18 (d, 1H), 7.62 (dd, 1H), 7.89 (br s, 1H), 8.24 (br s, 1H), 8.42 (m, 1H), 9.85 (m, 1H), 10.11 (br s, 1H), 13.73 (br s, 1H). MS m/z 586 [M+H]⁺

The following Examples (**Examples 79 – 91**) were prepared according to the method described for **Example 78** using the appropriate pyrazolo-pyridine.

Example	Name	Data
79	4-[(2-[ethyl(methylsulfonyl)amino]-5-fluoropyridin-3-yl)methylamino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 600 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.85 (t, 3H), 3.07 (s, 3H), 3.51 (q, 2H), 3.61 (q, 2H), 4.08 (m, 1H), 6.74 (s, 1H), 6.94 (d, 1H), 7.19 (d, 1H), 7.61 (m, 1H), 7.90 (br s, 1H), 8.24 (br s, 1H), 8.45 (s, 1H), 9.84 (t, 1H), 10.08 (s, 1H), 13.74 (s, 1H). Using 6-(4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((5-fluoro-2-(N-ethylmethylsulfonamido)pyridin-3-yl)methylamino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 56).
80	4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methylamino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 583 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.86 (t, 3H), 3.10 (s, 3H), 3.59 (q, 2H), 3.68 (q, 2H), 4.97 (m, 2H), 6.60 (s, 1H), 6.97 (d, 1H), 7.15 (d, 1H), 7.82 (br s, 1H), 8.16 (br s, 1H), 8.56 (s, 1H), 8.61 (s, 1H), 9.85 (br s, 1H), 10.09 (s, 1H), 13.66 (s, 1H). Using 6-(4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((3-(N-ethylmethylsulfonamido)pyrazin-2-yl)methylamino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 57).

81	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-((5-methyl-2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 581 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.20 (s, 3H), 3.01 (s, 3H), 3.07 (s, 3H), 3.64 (m, 2H), 4.64 (m, 1H), 4.88 (m, 1H), 6.94 (s, 1H), 6.96 (d, 1H), 7.14 (m, 1H), 7.21 (m, 2H), 7.37 (m, 1H), 7.84 (br s, 1H), 8.18 (br s, 1H), 9.69 (s, 1H), 10.09 (s, 1H), 13.67 (s, 1H). Using 6-(4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((5-methyl-2-(N-methylmethylsulfonylamino)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 28).
82	4-((2-(N-ethylethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 595 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.90 (t, 3H), 1.19 (t, 3H), 3.19 (m, 2H), 3.51-3.64 (m, 4H), 4.70 (m, 1H), 4.87 (m, 1H), 6.69 (s, 1H), 6.94 (d, 1H), 7.19 (d, 1H), 7.30 (m, 2H), 7.42 (m, 2H), 7.85 (br s, 1H), 8.10 (br s, 1H), 9.77 (m, 1H), 10.06 (s, 1H), 13.67 (s, 1H). Using 6-(4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-ethylethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 32).
83	4-((2-(N-ethylmethylsulfonamido)-5-fluorobenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 599 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.89 (t, 3H), 3.02 (s, 3H), 3.43-3.64 (m, 4H), 4.71 (m, 1H), 4.85 (m, 1H), 6.72 (s, 1H), 6.93 (d, 1H), 7.10-7.19 (m, 3H), 7.51 (m, 1H), 7.88 (br s, 1H), 8.22 (br s, 1H), 9.79 (m, 1H), 10.05 (s, 1H), 13.71 (s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-ethylmethylsulfonamido)-5-fluorobenzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 34).
84	4-((5-chloro-2-(N-ethylmethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 615 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.01 (t, 3H), 3.16 (s, 3H), 3.58-3.76 (m, 4H), 4.82 (m, 1H), 4.98 (m, 1H), 6.85 (s, 1H), 7.06 (d, 1H), 7.32 (m, 1H), 7.52 (m, 2H), 7.62 (m, 1H), 8.01 (br s, 1), 8.35 (br s, 1H), 9.91 (t, 1H), 10.18 (s, 1H), 13.85 (s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-ethylmethylsulfonamido)-5-chlorobenzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 35).

85	6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylethylsulfonamido)benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 581 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.19 (t, 3H), 3.11 (s, 3H), 3.23-3.31 (m, 2H), 3.58 (q, 2H), 4.60-4.90 (m, 2H), 6.70 (s, 1H), 6.95 (d, 1H), 7.21 (d, 1H), 7.26-7.34 (m, 2H), 7.40-7.47 (m, 2H), 7.86 (br s, 1H), 8.21 (br s, 1H), 9.78 (t, 1H), 10.09 (s, 1H), 13.68 (s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-methylethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 37).
86	6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-(((5-methyl-2-(N-methylmethylsulfonamido)pyridin-3-yl)methyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 582 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.22 (s, 3H), 3.05 (s, 3H), 3.09 (s, 3H), 3.61 (q, 2H), 4.75 (m, 2H), 6.72 (s, 1H), 6.96 (d, 1H), 7.19 (d, 1H), 7.61 (br s, 1H), 7.86 (br s, 1H), 8.23 (m, 2H), 9.77 (m, 1H), 10.09 (s, 1H), 13.70 (s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((5-methyl-2-(N-methylmethylsulfonamido)pyridin-3-yl)methyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 38).
87	4-[(2-[ethyl(methylsulfonyl)amino]-5-methylpyridin-3-yl)methyl]amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 596 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.86 (t, 3H), 2.22 (s, 3H), 3.04 (s, 3H), 3.55-3.63 (m, 4H), 4.78 (m, 2H), 6.72 (s, 1H), 6.94 (d, 1H), 7.20 (d, 1H), 7.60 (br s, 1H), 7.87 (br s, 1H), 8.22 (s, 1H), 8.26 (s, 1H), 9.76 (t, 1H), 10.07 (br s, 1H), 13.71 (br s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((2-(N-ethylmethylsulfonamido)-5-methylpyridin-3-yl)methyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 33).
88	6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-((5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 611 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.05 (s, 3H), 3.08 (s, 3H), 3.58 (m, 2H), 4.70 (m, 1H), 4.83 (m, 1H), 6.73 (s, 1H), 6.94 (d, 1H), 7.09-7.22 (m, 3H), 7.56 (m, 1H), 7.89 (s, 1H), 8.24 (s, 1H), 9.80 (m, 1H), 10.11 (br s, 1H), 13.72 (br s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((5-fluoro-2-(N-methylmethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 59).

89	4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 601 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.05 (s, 3H), 3.08 (s, 3H), 3.56 (m, 2H), 4.66-4.82 (m, 2H), 6.73 (s, 1H), 6.94 (d, 1H), 7.20 (d, 1H), 7.38 (m, 2H), 7.53 (m, 1H), 7.88 (s, 1H), 8.22 (s, 1H), 9.77 (m, 1H), 10.09 (s, 1H), 13.71 (br s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((5-chloro-2-(N-methylmethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 30).
90	4-({2-[ethyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 581 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.88 (t, 3H), 3.02 (s, 3H), 3.46-3.68 (m, 4H), 4.69 (m, 1H), 2.92 (m, 1H), 6.69 (s, 1H), 6.94 (d, 1H), 7.17 (d, 1H), 7.32 (m, 2H), 7.45 (m, 2H), 7.86 (br s, 1H), 8.22 (br s, 1H), 9.76 (m, 1H), 10.08 (s, 1H), 13.69 (s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-ethylmethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 36).
91	4-({2-[ethyl(methylsulfonyl)amino]-5-methylbenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 595 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.90 (t, 3H), 2.21 (s, 3H), 2.99 (s, 3H), 3.46 (m, 1H), 3.61 (m, 1H), 4.64 (m, 1H), 4.86 (m, 1H), 6.69 (s, 1H), 6.92 (d, 1H), 7.12 (m, 1H), 7.18-7.21 (m, 2H), 7.30 (d, 1H), 7.84 (s, 1H), 8.19 (s, 1H), 9.69 (m, 1H), 10.06 (s, 1H), 13.67 (s, 1H). Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-ethylmethylsulfonamido)-5-methylbenzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 31).

Example 92**6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide**

5 6-(4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((5-methoxy-2-(N-methylmethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 29**, 100 mg, 0.12 mmol) was heated to 100 °C in neat TFA (15 mL) for 18 hours. The reaction was cooled, concentrated *in vacuo* and partitioned between saturated aqueous sodium bicarbonate solution and EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column

10

chromatography eluting with 55% EtOAc in hexanes. The residue was stirred with neat boron tribromide (8 eq) at 0 °C for 4 hours. The reaction was partitioned between DCM and saturated aqueous sodium bicarbonate solution, the organic layer collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC eluting with 5% MeOH in DCM to afford the title compound (30 mg, 51%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.98 (s, 3H), 3.04 (s, 3H), 3.66 (m, 2H), 4.58 (m, 1H), 4.79 (m, 1H), 6.63 (dd, 1H), 6.70 (s, 1H), 6.78 (m, 1H), 6.93 (d, 1H), 7.19-7.26 (m, 2H), 7.85 (br s, 1H), 8.20 (br s, 1H), 9.50 (s, 1H), 9.70 (m, 1H), 10.08 (s, 1H), 13.67 (s, 1H). MS m/z 583 [M+H]⁺

Example 93

6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-((2-(N-methylmethylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

To a solution of N-(2-(((6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 272, 249 mg, 0.36 mmol) and methylamine in THF (3 mL, 2M) was added molybdenum hexacarbonyl (96 mg, 0.36 mmol) and palladium acetate (5.7 mg, 0.025 mmol) followed by DBU (165 mg, 1.09 mmol) and the reaction was heated to 100 °C under microwave irradiation for 10 minutes. The reaction was concentrated *in vacuo*, diluted with EtOAc and filtered through CeliteTM. The filtrate was concentrated *in vacuo* and purified using silica gel column chromatography followed by Preparative TLC. The residue was dissolved in ethanol (7 mL) and hydrogenated with Pd(OH)₂ (15 mg) at 40 psi for 16 hours. The reaction was filtered through celite and concentrated *in vacuo*. The residue was triturated with pentane and ether to afford the title compound as an off white solid (42 mg, 65%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.98 (t, 3H), 2.83 (m, 4H), 3.07 (s, 3H), 3.18 (s, 3H), 3.39 (m, 1H), 4.90 (br m, 1H), 5.05 (br m, 1H), 6.82 (m, 1H), 7.35-7.56 (m, 5H), 8.86 (m, 1H), 9.81 (t, 1H), 10.02 (br s, 1H), 13.97 (br s, 1H). MS m/z 528 [M+H]⁺

Example 94

6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(N-methylmethanesulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

The title compound was prepared according to the method described for
5 **Preparation 93** using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (**Preparation 276**). The residue was purified using silica gel

column chromatography eluting with 60% EtOAc in hexanes. ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.82 (d, 3H), 3.07 (s, 3H), 3.17 (s, 3H), 4.27 (m, 2H), 4.91-5.01 (br m, 2H), 6.97 (m, 1H), 7.32-7.41 (m, 3H), 7.54 (m, 1H), 7.67 (m, 1H), 8.87 (t, 1H), 9.91 (t, 1H), 10.37 (br s, 1H), 14.03 (br s, 1H). MS m/z 582 [M+H]⁺

5

Example 95

6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-4-((2-(N-methylphenylsulfonamido)benzyl)amino)-1H-pyrazolo[3.4-d]pyrimidine-3-carboxamide

To a solution of 6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-4-((2-(N-methylphenylsulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo-[3,4-d]pyrimidine-3-carboxylic acid (**Preparation 258**, 300mg, 837.14mmol) in anhydrous THF (10 mL) was added NMM (0.06 mL, 0.57 mmol) and isobutylchloroformate (0.07 mL, 0.57 mmol) at -20 °C and the reaction mixture was stirred at this temperature for 2 hours. Aqueous ammonia (0.6 mL) was added and the reaction stirred at room temperature for 1 hour. The reaction was quenched by the addition of water and EtOAc. The organic layer was separated, washed with water, brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 66% EtOAc in hexanes. The residue was dissolved in TFA and stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo* and dissolved in MeOH (5 mL), cooling to 0°C. Ethylene diamine was added drop-wise until the solution showed a basic pH. The reaction was extracted into 20% IPA in DCM, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC to afford the title compound (30 mg, 27%).

¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.96 (t, 3H), 2.89 (q, 2H), 3.10 (s, 3H), 4.90 (br m, 1H), 5.08 (br m, 1H), 6.57 (m, 1H), 6.81 (m, 1H), 7.19 (m, 1H), 7.32 (m, 1H), 7.46 (m, 1H), 7.55-7.67 (m, 5H), 7.73 (m, 1H), 7.91 (br s, 1H), 8.26 (br s, 1H), 9.86 (t, 1H), 10.05 (br s, 1H), 13.96 (br s, 1H). MS m/z 575 [M+H]⁺

25

Example 96

6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylphenylsulfonamido)benzyl)amino)-1H-pyrazolo[3.4-d]pyrimidine-3-carboxamide

The title compound was prepared according to the method described for **Example 95** using 6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-4-((2-(N-methylphenylsulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (**Preparation 261**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.10 (s, 3H), 4.32 (m, 2H), 4.92 (m, 1H), 5.06 (m,

30

1H), 6.56 (m, 1H), 6.99 (m, 1H), 7.17 (m, 1H), 7.32 (m, 1H), 7.44 (m, 1H), 7.61-7.67 (m, 6H), 7.93 (br s, 1H), 8.28 (br s, 1H), 9.93 (t, 1H), 10.41 (br s, 1H), 14.04 (br s, 1H). MS m/z 630 [M+H]⁺

Example 97

5 6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-hydroxy-2-(N-methylmethylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

To a solution of 6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-methoxy-2-(N-methylmethylsulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo-[3,4-d]pyrimidine-3-carboxylic acid
 10 (**Preparation 269**, 0.1g, 0.13 mmol), HOBT (36 mg, 0.27 mmol) and EDCI (51 mg, 0.27 mmol) in dichloromethane (6 mL) at 0 °C was added ammonium chloride (36 mg, 0.67 mmol) and DIPEA (0.12 mL, 0.67 mmol) and the reaction was stirred at room temperature for 14 hours. The reaction was concentrated *in vacuo* and the residue diluted with ethyl acetate. The organic solution was washed with brine, dried over
 15 sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 52% EtOAc in hexanes. The residue (62 mg, 0.084 mmol) was dissolved in DCM (5 mL) at 0 °C and boron tribromide (0.08 mL, 0.83 mmol) was added. The reaction was stirred at room temperature for 1 hour. The reaction was concentrated *in vacuo*, diluted with methanol (5 ml) and treated with ethylene diamine
 20 until the pH was basic, stirring for 1 hour. The solvent was removed *in vacuo* and the residue was partitioned between ethyl acetate and water, the organic extracts were dried over sodium sulfate and purified by Preparative TLC to afford the title compound as an off-white solid (25mg, 51%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.01 (s, 3H), 3.12 (s, 3H), 4.30 (m, 2H), 4.80 (m, 1H), 4.91 (m, 1H), 6.68-6.71 (m, 1H), 6.76-6.77 (m, 1H), 6.97-7.00 (m, 1H), 7.31-7.33 (m, 1H), 7.69-7.72 (m, 1H), 7.93 (br s, 1H), 8.28 (br s, 1H), 9.61 (br s, 1H), 9.87 (t, 1H), 10.40 (br s, 1H). MS m/z 584 [M+H]⁺
 25

Example 98

6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylmethylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide
 30 The title compound was prepared according to the method described by **Example 97** using 6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylmethylsulfonamido)-benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-o[3,4-d]pyrimidine-3-carboxylic acid (**Preparation 270**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.07 (s, 3H), 3.18 (s, 3H), 4.29 (m, 2H), 4.88-5.00 (br m, 2H), 6.98 (m, 1H),

7.32-7.43 (m, 3H), 7.54 (m, 1H), 7.68 (m, 1H), 7.92 (br s, 1H), 8.27 (br s, 1H), 9.92 (t, 1H), 10.39 (br s, 1H), 14.02 (br s, 1H). MS m/z 568 [M+H]⁺

Example 99

5 6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-hydroxy-2-(N-methylmethylsulfonamido)benzyl)amino)-N-(6-((2-hydroxyethyl)amino)pyridin-3-yl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

To a solution of 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-hydroxy-2-(N-methyl methylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (**Example 174**, 160 mg, 0.27 mmol) and 2-[(5-aminopyridin-2-yl)amino]ethanol (84 mg, 0.54 mmol) in DMF (5 mL) was added HATU (312 mg, 0.82 mmol) and DIPEA (0.12 mL, 0.68 mmol) and the reaction was stirred at room temperature for 18 hours. The reaction was purified directly by Preparative HPLC to afford the title compound (48 mg, 24%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.02 (s, 3H), 3.12 (s, 3H), 3.32 (m, 2H), 3.51 (m, 2H), 4.30 (m, 2H), 4.81-4.93 (m, 2H), 6.44-6.52 (m, 2H), 6.68 (m, 1H), 6.76 (m, 1H), 6.97 (m, 1H), 7.33 (m, 1H), 7.69-7.77 (m, 2H), 8.31 (d, 1H), 9.63 (t, 1H), 10.53 (br s, 1H). MS m/z 720 [M+H]⁺

Example 100

20 6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-(((3-(N-methylmethyl-sulfonamido)pyrazin-2-yl)methyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

To a solution of N-(3-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)pyrazin-2-yl)-N-methylmethanesulfonamide (**Preparation 277**, 220 mg, 0.33 mmol) in methylamine in THF (3 mL) was added molybdenum hexacarbonyl (87.77 mg, 0.33 mmol), Pd(OAc)₂ (5.18 mg, 0.07 mmol) and DBU (0.15 mL, 0.99 mmol) and the reaction was heated to 100 °C under microwave irradiation for 10 minutes. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography followed by preparative TLC. The residue was dissolved in DCM (5 mL) and treated with boron tribromide (0.11 mL, 1.17 mmol) at 0 °C and stirred at room temperature for 18 hours.

30 The reaction was concentrated *in vacuo* and partitioned between DCM and saturated aqueous sodium bicarbonate solution. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC eluting with 5% MeOH in DCM to afford the title compound as a white solid (23 mg, 47%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.84 (s, 3H), 3.15 (s,

3H), 3.19 (s, 3H), 4.21 (m, 2H), 5.13 (m, 2H), 6.97 (m, 1H), 7.58 (m, 1H), 8.53 (d, 1H), 8.63 (d, 1H), 8.86 (m, 1H), 10.06 (m, 1H), 10.37 (br s, 1H), 14.04 (br s, 1H). MS m/z 584 [M+H]⁺

The following Examples (**Examples 101 – 104**) were prepared according to the method described for **Example 1** using the appropriate pyrazolo-pyrimidine as described below:

Ex	Name	Data
101	6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-(2-hydroxyethyl)methylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide	MS m/z 598 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.10 (s, 3H), 3.30-3.39 (m, 2H), 3.64 (m, 2H), 4.26 (m, 2H), 4.81 (m, 1H), 4.94 (m, 2H), 6.99 (m, 1H), 7.39 (m, 3H), 7.51 (m, 1H), 7.68 (m, 1H), 7.90 (m, 1H), 8.26 (m, 1H), 9.89 (s, 1H), 10.32 (br s, 1H), 14.01 (br s, 1H). Using N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-(2-((tert-butyldimethylsilyl)oxy)ethyl)methanesulfonamide (Preparation 265) with ammonia in THF.
102	6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-(2-hydroxyethyl)methylsulfonamido)benzyl)amino)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide	MS m/z 612 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.83 (m, 3H), 3.10 (s, 3H), 3.33-3.40 (m, 2H), 3.64 (m, 2H), 4.21-4.30 (m, 2H), 4.85 (t, 1H), 5.01 (m, 2H), 6.96 (m, 1H), 7.36-7.38 (m, 3H), 7.51 (m, 1H), 7.67 (m, 1H), 8.88 (m, 1H), 9.87 (t, 1H), 10.40 (br s, 1H). Using N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-(2-((tert-butyldimethylsilyl)oxy)ethyl)methanesulfonamide (Preparation 265).

103	4-((2-(N-ethylmethanesulfonamido)-benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide	MS m/z 594 [M-H] ⁻ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.00 (t, 3H), 2.83 (d, 3H), 3.06 (s, 3H), 3.55 (m, 1H), 3.69 (m, 1H), 4.18 (m, 2H), 4.87 (m, 1H), 4.92 (m, 1H), 6.96 (m, 1H), 7.33-7.40 (m, 3H), 7.51 (m, 1H), 7.65 (m, 1H), 8.88 (t, 1H), 9.91 (t, 1H), 10.36 (br s, 1H), 14.05 (br s, 1H). Using N-ethyl-N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)-methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)methanesulfonamide (Preparation 263).
104	4-((5-fluoro-2-(N-methylmethanesulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide	MS m/z 600 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.85 (d, 3H), 3.08 (s, 3H), 3.16 (s, 3H), 4.21 (m, 2H), 4.91-4.99 (m, 2H), 6.95 (m, 1H), 7.10-7.22 (m, 2H), 7.60-7.66 (m, 2H), 8.89 (m, 1H), 9.92 (t, 1H). Using N-(4-fluoro-2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)-phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 268).

Example 105

4-((2-(N-Ethylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

5 The title compound was prepared according to the method described for **Example 32** using N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-methyl)-4-methoxyphenyl)-benzenesulfonamide (**Preparation 266**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.93 (t, 3H), 2.85 (d, 3H), 3.24 (m, 1H), 3.77 (m, 1H), 4.29 (m, 2H),
10 4.85-4.93 (m, 2H), 6.36 (m, 1H), 6.54 (m, 1H), 6.76 (m, 1H), 6.98 (m, 1H), 7.58-7.72 (m, 5H), 8.89 (m, 1H), 9.62 (s, 1H), 9.85 (br s, 1H), 10.10 (br s, 1H), 14.10 (br s, 1H). MS m/z 672 [M-H]⁻

Example 106

4-((2-(N-Methylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

15

The title compound was prepared according to the method described for **Example 32** using N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethyl-silyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-amino)methyl)-4-methoxyphenyl)-benzenesulfonamide (**Preparation 267**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.85 (s, 3H), 3.05 (s, 3H), 4.34 (m, 2H), 4.86 (m, 1H), 4.97 (m, 1H), 6.32 (m, 1H), 6.51 (m, 1H), 6.76 (m, 1H), 6.98 (m, 1H), 7.60-7.74 (m, 6H), 8.91 (m, 1H), 9.63 (s, 1H), 9.89 (t, 1H), 10.40 (br s, 1H), 14.10 (br s, 1H). MS m/z 660 [M+H]⁺

The following Examples (**Examples 107 – 108**) were prepared according to the method described for **Example 41** using the appropriate pyrazolo-pyrimidine as described below:

Ex	Name	Data
107	4-((2-(N-ethylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide	MS m/z 660 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.94 (t, 3H), 3.20 (m, 1H), 3.78 (m, 1H), 4.24-4.30 (m, 2H), 4.81 (m, 1H), 4.97 (m, 1H), 6.35 (m, 1H), 6.53 (m, 1H), 6.79 (m, 1H), 7.00 (m, 1H), 7.57-7.72 (m, 6H), 7.95 (br s, 1H), 8.30 (br s, 1H), 9.63 (s, 1H), 9.87 (t, 1H), 10.37 (br s, 1H), 14.05 (br s, 1H). Using 4-((2-(N-ethylphenylsulfon-amido)-5-methoxybenzyl)amino)-6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoro-ethyl)phenyl)-1-((2-(trimethylsilyl)-ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (Preparation 253).
108	4-((2-(N-methylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide	MS m/z 646 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.05 (s, 3H), 4.08 (m, 2H), 4.82 (m, 1H), 4.96 (m, 1H), 6.32 (m, 1H), 6.50 (m, 1H), 6.78 (m, 1H), 7.01 (m, 1H), 7.60-7.75 (m, 6H), 7.95 (br s, 1H), 8.30 (br s, 1H), 9.63 (s, 1H), 9.89 (t, 1H), 10.38 (s, 1H), 14.05 (br s, 1H). Using 4-((2-(N-methylphenylsulfonamido)-5-methoxybenzyl)amino)-6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)-phenyl)-1-((2-(trimethylsilyl)-ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide (Preparation 254).

Example 109

6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(methyl(sulfamoyl)amino)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

To a solution of tert-butyl 6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(methylamino)benzyl)amino)-3-(methylcarbamoyl)-1H-pyrazolo[3,4-d]pyrimidine-1-


carboxylate (**Preparation 256**, 50 mg, 0.08 mmol) in anhydrous THF (5 mL) was added sodium hydride (3 mg, 0.08 mmol) at 0°C. The reaction was stirred for 10 minutes before the addition of sulfamoyl chloride (7 mg, 0.06 mmol) and further stirring at 0 °C for 2.5 hours. The reaction was quenched by the addition of ice-water and extracted into EtOAc. The organic layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative TLC and dissolved in DCM (5 mL). The solution was treated with boron tribromide (0.08 mL, 0.8 mmol) and stirred at room temperature for 18 hours. The reaction was concentrated *in vacuo* and partitioned between DCM and saturated aqueous sodium bicarbonate solution. The organic layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative TLC to afford the title compound as a white solid (11 mg, 32%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.86 (s, 3H), 3.08 (s, 3H), 4.29 (m, 2H), 4.86-5.05 (m, 2H), 6.97 (m, 1H), 7.09 (s, 2H), 7.25-7.35 (m, 3H), 7.46 (m, 1H), 7.69-7.72 (m, 1H), 8.88 (t, 1H), 9.88 (t, 1H), 10.37 (s, 1H), 14.02 (s, 1H). MS m/z 583 [M+H]⁺

Example 110

6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(methyl(N-methylsulfamoyl)amino)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

To a solution of tert-butyl 6-(4-((tert-butoxycarbonyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(methylamino)benzyl)amino)-3-(methylcarbamoyl)-1H-pyrazolo-[3,4-d]pyrimidine-1-carboxylate (**Preparation 255**, 56 mg, 0.08 mmol) in THF (3 mL) was added sodium hydride (2 mg, 0.08 mmol) at 0°C. The reaction was stirred at room temperature for 2 minutes before the addition of methanesulfonyl chloride (10 mg, 0.08 mmol) and further stirring for 18 hours. The reaction was quenched by the addition of ice-water and extracted into EtOAc. The organic layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative TLC and treated with 4M HCl in dioxane (0.3 mL). The reaction was stirred at room temperature for 3 hours before concentrating *in vacuo* and triturating with pentane-ether to afford the title compound. ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.62 (d, 3H), 2.86 (d, 3H), 3.05 (s, 3H), 4.33 (m, 2H), 4.95 (m, 2H), 6.98 (m, 1H), 7.28-7.49 (m, 4H), 7.73 (m, 1H), 8.89 (t, 1H), 9.92 (t, 1H), 10.38 (s, 1H), 14.03 (s, 1H). MS m/z 595 [M-H]⁻

Library Protocol 1



A 0.1 M solution of 6-(2-ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 328**, 700 μ L, 70 μ mol) in toluene was added to an amine of formula (XII) (200 μ mol, 2.9 eq) and the solution degassed with nitrogen. Cesium carbonate (45 mg, 140 μ mol) was added followed by $\text{Pd}_2(\text{dba})_3$ (3.4 mg, 3.5 μ mol) and BINAP (2.2 mg, 3.5 μ mol) and the reaction further degassed with nitrogen. The reaction was shaken at 80 $^{\circ}\text{C}$ for 16 hours before concentrating *in vacuo*. Water (1 mL) was added followed by EtOAc (1 mL) and the mixture filtered. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. To the residue was added a solution of HCl in EtOH (1 mL, v:v 1:6) and the reaction shaken at 80 $^{\circ}\text{C}$ for 2 hours. The reaction was cooled, concentrated *in vacuo* and purified using one of the Preparative HPLC methods described below:

Preparative HPLC

Method A: Agella Venusil ASB C18, 150x21.2mmx5 μ m; Acetonitrile-Water (0.225% formic acid); Flow rate: 35 mL/min; Gradient time 8 mins.

Method B: Boston Symmetrix ODS-H, 150x30mmx5 μ m; Acetonitrile-Water (0.225% formic acid); Flow rate: 30 mL/min; Gradient time 10 mins.

Method C: DIKMA Diamonsil(2) C18, 200x20mmx5 μ m; Acetonitrile-Water (0.225% formic acid); Flow rate: 30 mL/min; Gradient time 10 mins.

LCMS:

Method 1

A: 0.0375% TFA in water; B: 0.01875% TFA in MeCN; Column: XBridge C18, 2.1x50 mmx5 μ m; Gradient: From 99% [A] and 1% [B] to 95% [A] and 5% [B] in 0.6 min, further to 100% [B] in 4.0 min and finally back to initial condition in 4.3 min, 0.8 mL/min flow rate.

Method 2

A: 0.0375% TFA in water; B: 0.01875% TFA in MeCN; Column: XBridge C18, 2.1x50 mmx5 μ m; Gradient: From 90% [A] and 10% [B] to 100% [B] in 4 min and finally back to initial condition in 4.3 min, 0.8 mL/min flow rate.

Method 3

A: 0.0375% TFA in water; B: 0.01875% TFA in MeCN; Column: XBridge C18, 2.1x50 mmx5µm; Gradient: From 75% [A] and 25% [B] to 100% [B] in 3.5 min and finally back to initial condition in 4.0 min, 0.8 mL/min flow rate.

5 The compounds of the Examples in the table below (**Examples 111 – 124**) were prepared and purified from 6-(2-ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 328**) and the appropriate amine according to Library Protocol 1. The compounds were isolated as their formate salts.

Ex	Name	MS Data/HPLC Organic gradient/Amine
111	4-[4-(7,8-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]-3-ethylphenol formate	MS m/z 431 [M+H] ⁺ Rt = 1.72 min HPLC organic gradient: 21-51%. 1,2,3,4-tetrahydro-7,8-dimethoxy-isoquinoline (Chem. & Pharm. Bull. (1998), 46 (6), 918-927).
112	3-ethyl-4-[4-(3-phenoxyazetidin-1-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate	MS m/z 387 [M+H] ⁺ Rt = 1.73 min HPLC organic gradient: 21-51%. 3-phenoxyazetidine.
113	3-ethyl-4-[4-[6-(4-methyl-1H-imidazol-1-yl)-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate	MS m/z 451 [M+H] ⁺ Rt = 2.16 min HPLC organic gradient: 3-33%. 6-(4-methyl-1H-imidazol-1-yl)-1,2,3,4-tetrahydroisoquinoline hydrochloride (Preparation 345).
114	3-ethyl-4-[4-[6-(2-methoxyethoxy)-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate	MS m/z 445 [M+H] ⁺ Rt = 2.64 min HPLC organic gradient: 18-48%. 1,2,3,4-tetrahydro-6-(2-methoxyethoxy)-isoquinoline.
115	1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-3-methylazetidin-3-ol formate	MS m/z 325 [M+H] ⁺ Rt = 2.11 min HPLC organic gradient: 3-33%. 3-methyl-3-azetidinol (J. Med. Chem. (2010), 53(9), 3645-3674).

116	2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-N-[2-(pyrrolidin-1-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide formate	MS m/z 547 [M+H] ⁺ Rt = 2.22 min HPLC organic gradient: 6-36%. N-(2-(pyrrolidin-1-yl)ethyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide (Preparation 346).
117	N-benzyl-2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-1,2,3,4-tetrahydroisoquinoline-5-carboxamide formate	MS m/z 504 [M+H] ⁺ Rt = 2.50 min HPLC organic gradient: 20-50%. N-benzyl-1,2,3,4-tetrahydroisoquinoline-5-carboxamide hydrochloride (Preparation 347).
118	4-{4-[7-(benzyloxy)-6-methoxy-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl}-3-ethylphenol formate	MS m/z 507 [M+H] ⁺ Rt = 2.20 min HPLC organic gradient: 26-56%. 1,2,3,4-tetrahydro-6-methoxy-7-(phenylmethoxy)-isoquinoline (Heterocycles (1989), 28(1), 295-301).
119	4-[4-(5-chloro-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]-3-ethylphenol formate	MS m/z 405 [M+H] ⁺ Rt = 2.60 min HPLC organic gradient: 22-52%. 1,2,3,4-tetrahydro-5-chloro-isoquinoline .
120	4-chloro-3-({1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]azetidin-3-yl}oxy)benzonitrile formate	MS m/z 446 [M+H] ⁺ Rt = 2.56 min HPLC organic gradient: 21-51%. 3-(azetidin-3-yloxy)-4-chlorobenzonitrile hydrochloride (Preparation 348).
121	3-ethyl-4-[4-(6-fluoro-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate	MS m/z 389 [M+H] ⁺ Rt = 2.70 min HPLC organic gradient: 19-49%. 1,2,3,4-tetrahydro-6-fluoro-isoquinoline.
122	3-ethyl-4-[4-(8-methoxy-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate	MS m/z 401 [M+H] ⁺ Rt = 1.82 min HPLC organic gradient: 14-44%. 1,2,3,4-tetrahydro-8-methoxyisoquinoline.

123	N-{2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-1,2,3,4-tetrahydroisoquinolin-5-yl}methanesulfonamide formate	MS m/z 464 [M+H] ⁺ Rt = 2.16 min HPLC organic gradient: 13-43%. N-1,2,3,4-tetrahydroisoquinolin-5-yl}methanesulfonamide hydrochloride.
124	4-(4-([2-(biphenyl-4-yl)ethyl]amino)-1H-pyrazolo[4,3-c]pyridin-6-yl)-3-ethylphenol formate	MS m/z 435 [M+H] ⁺ Rt = 2.20 min HPLC organic gradient: 25-55%. 2-([1,1'-biphenyl]-4-yl)ethanamine.

Example 125**N-[2-([6-(2-Ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl]-N-methylmethanesulfonamide hydrochloride**

5 To a solution of 6-(2-ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 328**, 14 mg, 0.02 mmol) in toluene (0.5 mL) was added N-[2-aminomethyl]phenyl]-N-methylmethanesulfonamide (**PCT Intl. Appl. 2010058846**, 9 mg, 0.03 mmol), cesium carbonate (14 mg, 0.04 mmol), Pd(OAc)₂ (0.9 mg, 0.004 mmol) and BINAP (3.7 mg, 0.006 mmol). The reaction was degassed with nitrogen followed by heating to 150 °C under microwave irradiation for 15 minutes. The reaction was filtered, washing through with DCM and the filtrate concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 1-5% MeOH in DCM. The residue was dissolved in MeOH (1 mL) and cHCl (0.2 mL) was added and the reaction heated to 80 °C for 3 hours. The reaction was cooled, concentrated *in vacuo* and triturated with DCM to afford the title compound as the hydrochloride salt (10 mg, 50%).
¹H NMR (400MHz, DMSO-d₆): δ ppm 0.75 (t, 3H), 2.40 (q, 2H), 3.00 (s, 3H), 3.03 (s, 3H), 4.55 (br m, 1H), 4.95 (br m, 1H), 6.50 (m, 3H), 7.00 (m, 1H), 7.25 (m, 2H), 7.40-7.50 (m, 2H), 7.80 (t, 1H), 8.20 (s, 1H), 9.25 (s, 1H), 12.95 (s, 1H). MS m/z 452 [M+H]⁺

20

The following compounds of the Examples below (**Examples 126 – 130**) were prepared according to the method described for **Example 125** above using 6-(2-ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 328**) or 6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 331**) or 6-(2-

25

ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-
1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 330**) and the
appropriate amine. Purification took place according to the Purification Method (PM)
described or one of the following below. Compounds were isolated as the free parent,
5 diethylamine salt or hydrochloride salt as described below:

Purification Method J: The residue was dissolved in DMSO (0.9 mL) and triethylamine (0.1 mL) and purified using Preparative HPLC.

Ex	Name	MS Data/PM/Amine
126	Racemic 1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-N,N-dimethylpyrrolidine-3-sulfonamide (free parent).	MS m/z 416 [M+H] ⁺ Rt = 2.75 min PM J. Racemic N,N-dimethyl-3-pyrrolidinesulfonamide hydrochloride.
127	N-[2-({[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-3-methylphenyl]-N-methylmethanesulfonamide diethylamine salt	MS m/z 466 [M+H] ⁺ Rt = 2.99 min PM J. N-[2-(aminomethyl)-3-methylphenyl]-N-methylmethanesulfonamide (PCT Intl. Appl. 2012045195).
128	3-ethyl-4-[4-(4-methoxypiperidin-1-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol diethylamine salt	MS m/z 353 [M+H] ⁺ Rt = 2.13 min PM J. 4-methoxypiperidine.
129	N-[2-({[2-(3,4-dimethoxyphenyl)ethyl][6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 634 [M+H] ⁺ ¹ H NMR (400MHz, DMSO-d ₆): δ ppm 0.85 (br s, 3H), 2.95 (m, 2H), 3.09 (s, 6H), 3.64 (s, 3H), 3.69 (s, 3H), 4.03 (m, 1H), 5.09 (m, 2H), 6.72-6.83 (m, 3H), 6.91 (br s, 1H), 7.16-7.32 (br s, 6H), 7.60 (br s, 1H), 7.98 (br s, 1H), 8.24 (br s, 1H), 9.80 (br s, 1H), 10.29 (br s, 1H). N-(2-(((3,4-dimethoxyphenethyl)amino)-methyl)phenyl)-N-methylmethanesulfonamide (Preparation 367).

130	N-[2-({6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl}(2-{4-[(methylsulfonyl)amino]phenyl}ethyl)amino)methyl)phenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 667 [M+H] ⁺ ¹ H NMR (400MHz, DMSO-d ₆): δ ppm 0.94-1.06 (m, 3H), 1.75-1.88 (m, 4H), 2.66 (s, 3H), 2.71-2.81 (m, 2H), 2.95-3.05 (s, 6H), 4.01 (m, 2H), 5.09 (m, 2H), 6.90-6.98 (m, 2H), 7.09-7.11 (m, 2H), 7.19-7.21 (m, 2H), 7.29-7.37 (m, 3H), 7.42-7.60 (m, 1H), 9.61 (s, 1H). N-methyl-N-(2-(((4-(methylsulfonamido)-phenethyl)amino)methyl) phenyl)methanesulfonamide (Preparation 369).
-----	--	--

Example 131**N-[2-({6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl)phenyl]-N-methylmethanesulfonamide**

5 To a stirring solution of 4-nitrophenyl {2-[methyl(methylsulfonyl)amino]benzyl}carbamate (**Preparation 166**, 2.02 g, 5.32 mmol) and triethylamine (2.12 mL, 15.33 mmol) in anhydrous DMF (20 mL) was added 6-(2-ethyl-5-fluoro-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 332**, 1.5 g, 3.07 mmol) and the reaction

10 was heated to 80 °C for 15 hours. The reaction was concentrated *in vacuo* and partitioned between water and ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 1% MeOH in DCM. The residue was dissolved in MeOH (10 mL) and cHCl (8 mL) was added. The reaction was heated at from 65-80

15 °C for 6 hours before cooling and concentrating *in vacuo*. The residue was triturated with MeCN/ether to afford the title compound as the hydrochloride salt (400 mg, 42%).
¹H NMR (400MHz, DMSO-d₆): δ ppm 0.98 (m, 3H), 2.32-2.41 (m, 2H), 3.09-3.18 (m, 6H), 5.11 (m, 2H), 6.98 (s, 2H), 7.09-7.21 (m, 2H), 7.21 (s, 2H), 7.27 (br s, 21H), 7.34-7.36 (m, 1H), 7.42-7.48 (m, 3H), 7.63 (br s, 1H), 8.70 (br s, 1H), 10.04 (br s, 1H), 10.35

20 (br s, 1H), 12.20 (br s, 1H), 14.17 (br s, 1H). MS m/z 470 [M+H]⁺

The following compounds of the Examples below (**Examples 132 – 136**) were prepared according to the method described for **Example 131** above using 6-(2-ethyl-5-fluoro-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 332**) or 6-(2-ethyl-4-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 333**) and the appropriate aminocarbamate. The compounds were

25

isolated as their hydrochloride salts. Purification was carried out according to the Purification Method (PM) described or one of the following below:

Purification Method K: Trituration with pentane-ether.

Ex	Name	MS Data/PM/Aminocarbamate
132	N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}-4-methylphenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 484 [M+H] ⁺ ¹ H NMR (400MHz, DMSO-d ₆): δ ppm 1.00 (s, 3H), 2.30 (m, 5H), 3.14 (s, 6H), 4.77 (m, 2H), 6.97 (s, 2H), 7.21 (d, 2H), 7.33 (d, 1H), 7.45 (s, 1H), 8.68 (s, 1H), 9.94 (s, 1H), 10.33 (s, 1H), 12.17 (s, 1H), 14.16 (s, 1H). PM K. 4-nitrophenyl {5-methyl-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 165).
133	N-[4-chloro-2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 504 [M+H] ⁺ ¹ H NMR (400MHz, DMSO-d ₆): δ ppm 0.87 (t, 3H), 3.01 (s, 3H), 3.24 (s, 3H), 4.85-5.07 (br s, 4H), 6.87 (s, 1H), 6.93 (d, 1H), 7.13 (d, 1H), 7.45 (d, 1H), 7.53-7.59 (m, 2H), 8.48 (s, 1H), 9.96 (br s, 1H), 10.34 (br s, 1H), 12.32 (br s, 1H), 14.15 (br s, 1H). 4-nitrophenyl{5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 156)
134	N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}-3-fluorophenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 488 [M+H] ⁺ ¹ H NMR (400MHz, DMSO-d ₆): δ ppm 1.05 (t, 3H), 2.49 (s, 2H), 3.05 (s, 3H), 3.21 (s, 3H), 4.94 (s, 2H), 6.98 (t, 2H), 7.27 (d, 1H), 7.36 (t, 1H), 7.52 (m, 2H), 8.72 (s, 1H), 10.38 (s, 1H), 12.56 (s, 1H). 4-nitrophenyl{2-[ethyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 162).
135	N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl]-N-methylethanesulfonamide hydrochloride	MS m/z 488 [M+H] ⁺ ¹ H NMR (400MHz, DMSO-d ₆): δ ppm 0.98 (m, 3H), 1.25 (m, 3H), 2.39-2.41 (m, 1H), 2.53 (s, 1H), 3.20 (m, 1H), 5.08-5.10 (m, 2H), 6.99 (m, 2H), 7.27 (s, 1H), 7.43-7.47 (m, 3H), 7.61 (m, 1H), 8.65 (br s, 1H), 10.01 (s, 1H), 10.32 (br s, 1H), 12.15 (br s, 1H), 13.01 (br s, 1H), 14.22 (br s, 1H). 4-nitrophenyl{2-[(ethylsulfonyl)(methyl)amino]benzyl}carbamate (Preparation 159).

136	N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]ethanesulfonamide hydrochloride	MS m/z 498 [M+H] ⁺ ¹ H NMR (400MHz, DMSO-d ₆): δ ppm 1.04 (m, 5H), 1.24 (m, 1H), 2.32 (m, 1H), 3.08 (m, 1H), 3.61-3.70 (m, 2H), 4.94 (m, 2H), 7.00 (m, 2H), 7.28 (m, 1H), 7.39-7.56 (m, 4H), 8.30-8.61 (m, 2H), 9.95 (br s, 1H), 10.31 (br s, 1H), 12.00 (br s, 1H), 13.00 (br s, 1H), 14.15 (br s, 1H). 4-nitrophenyl[ethyl(ethylsulfonyl)amino]benyl]-carbamate (Preparation 157).
------------	---	---

Example 137

N-[2-({[6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-propylmethanesulfonamide

5 To 6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 331**, 200 mg, 0.30 mmol) and N-(2-(aminomethyl)phenyl)-N-propylmethanesulfonamide trifluoroacetate (**Preparation 349**, 172 mg, 0.71 mmol) in DMF (5 mL) was added triethylamine (0.19 mL, 1.4 mmol) and the reaction was heated
10 to 110 °C for 2 hours. The reaction was cooled and partitioned between EtOAc and water. The organic layer was collected, washed with water, dried over magnesium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 1:1 EtOAc:Heptanes. The residue was dissolved in MeOH (5 mL) and cHCl (1.5 mL) was added and the reaction heated to 60 °C for 18 hours. The
15 reaction was cooled, concentrated *in vacuo* and purified using preparative HPLC to afford the title compound as the free parent. Rt = 2.85 minutes; MS m/z 498 [M+H]⁺

The following compounds of the Examples below (**Examples 138 – 144**) were prepared according to the method described for **Example 137** above using 6-(2-ethyl-5-
20 fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo-[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 331**) and the appropriate amine. Deprotection was carried out as described or using TFA/TES in place of cHCl. Purification was carried out according to the Purification Method (PM) described or one of the following below. The compounds were all isolated as free
25 parents.

Purification Method L: The reaction mixture was quenched by the addition of saturated aqueous NaHCO₃ solution and extracted into DCM. The organic layer was

collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative HPLC.

Ex	Name	MS Data/PM/Amine
138	N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)methanesulfonamide	MS m/z 484 [M+H] ⁺ Rt = 2.75 min N-[2-(aminomethyl)phenyl]-N-ethylmethanesulfonamide hydrochloride (Preparation 188).
139	N-butyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)methanesulfonamide	MS m/z 512 [M+H] ⁺ Rt = 2.98 min N-(2-(aminomethyl)phenyl)-N-butylmethanesulfonamide trifluoroacetate (Preparation 350).
140	N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl][2-(morpholin-4-yl)ethyl]amino}-methyl)phenyl]-N-methylmethanesulfonamide	MS m/z 583 [M+H] ⁺ Rt = 1.98 min N-methyl-N-(2-(((2-morpholinoethyl)amino)methyl)phenyl)methanesulfonamide (Preparation 353).
141	N-butyl-N-(2-(((6-(2-ethyl-5-fluoro-4-(1-oxidanyl)phenyl)-1H-pyrazolo[4,5-c]pyridin-4-yl)(methyl)amino)methyl)phenyl)methanesulfonamide	MS m/z 526 [M+H] ⁺ Rt = 3.00 min N-butyl-N-(2-((methylamino)methyl)phenyl)methanesulfonamide (Preparation 354).
142	N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino)methyl}-4-methylphenyl)methanesulfonamide	MS m/z 512 [M+H] ⁺ Rt = 2.03 min N-ethyl-N-(4-methyl-2-((methylamino)methyl)phenyl)methanesulfonamide (Preparation 355).
143	N-[2-({ethyl[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}-4-methylphenyl]-N-methylmethanesulfonamide	MS m/z 512 [M+H] ⁺ Rt = 2.89 min N-(2-((ethylamino)methyl)-4-methylphenyl)-N-methylmethanesulfonamide (Preparation 356). PM L.

144	N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](propyl)amino}methyl)-4-methylphenyl]-N-methylmethanesulfonamide	MS m/z 526 [M+H] ⁺ Rt = 3.04 min N-methyl-N-(4-methyl-2-((propylamino)methyl)phenyl)methanesulfonamide (Preparation 357). PM L.
-----	--	--

Example 145**N-Ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino}methyl)phenyl]methanesulfonamide hydrochloride**

5 To a solution of 6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 330**, 400 mg, 0.64 mmol) and N-ethyl-N-(2-((methylamino)methyl)phenyl)methanesulfonamide (**Preparation 358**, 234 mg, 0.96 mmol) in toluene (8 mL) was added cesium carbonate (420 mg, 1.29 mmol) and the mixture was

10 degassed with nitrogen for 5 minutes. Pd(OAc)₂ (16 mg, 0.064 mmol) and BINAP (60 mg, 0.096 mmol) were added and the reaction was heated to 140 °C under microwave irradiation for 30 minutes. The reaction was partitioned between water and ethyl acetate, the organic layer collected, washed with brine, dried over sodium sulfate, and concentrated *in vacuo*. The residue was purified using silica gel column

15 chromatography eluting with 36% EtOAc in hexanes. The residue (147 mg, 0.22 mmol) was dissolved in MeOH (10mL) and cHCl (10mL) was added with heating to 65 °C for 4 hours. The reaction was concentrated *in vacuo* and triturated with pentane-ether to afford the title compound as the hydrochloride salt (125 mg, 34%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 1.00-1.13 (m, 6H), 2.56 (m, 2H), 3.07 (s, 3H), 3.43 (s, 3H), 3.66 (m, 2H), 5.19 (br s, 2H), 6.82 (s, 1H), 6.88 (d, 1H), 7.09 (d, 1H), 7.23 (t, 3H), 7.33 (t, 1H), 7.39 (t, 1H), 7.53 (d, 1H), 8.16 (s, 1H), 10.31 (br s, 1H), 11.95 (br s, 1H), 14.16 (br s, 1H). MS m/z 496 [M-H]⁺

20

Example 146**N-[2-({[6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino}methyl)phenyl]-N-methylmethanesulfonamide hydrochloride**

25 The title compound was prepared according to the method described for **Example 145** using 6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate (**Preparation 330**), N-methyl-N-(2-((methylamino)methyl)phenyl)methanesulfonamide

(**Preparation 359**). The residue was triturated with pentane-ether and further purified by Preparative TLC to afford the title compound as the hydrochloride salt (60 mg, 59%).
¹H NMR (400MHz, DMSO-d₆): δ ppm 0.91 (m, 3H), 2.54 (m, 2H), 3.09 (s, 3H), 3.16 (s, 3H), 3.37 (s, 3H), 5.10 (s, 2H), 6.67 (s, 1H), 6.79 (d, 1H), 7.01 (d, 1H), 7.09 (d, 1H), 7.28 (t, 1H), 7.34 (t, 1H), 7.55 (d, 1H), 8.06 (s, 1H), 9.74 (s, 1H), 13.08 (s, 1H). MS m/z 482 [M-H]⁻

Example 147

N-[2-({6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl]phenyl]-N-(2-hydroxyethyl)methanesulfonamide

To a solution of N-(2-(benzyloxy)ethyl)-N-(2-(((6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)methanesulfonamide (**Preparation 327**, 79 mg, 0.09 mmol) in 1:1 MeOH:EtOH (10 mL) was added ammonium formate (1 mg, 0.09 mmol) followed by palladium hydroxide (4 mg). The reaction was heated to 70 °C for 18 hours before cooling and filtering through celite. The filtrate was concentrated *in vacuo* and partitioned between EtOAc and water. The organic layer was collected, washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was dissolved in DCM (3 mL) and TFA (141 μL, 1.84 mmol) followed by triethylsilane (21.5 μL, 0.18 mmol) were added. The reaction was heated to 70 °C for 72 hours before cooling and quenching with saturated aqueous NaHCO₃ solution. The reaction was extracted into EtOAc, and the organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was triturated with DCM to afford the title compound (16 mg, 43%).¹H NMR (400MHz, CDCl₃): δ ppm 0.90 (t, 3H), 2.25-2.40 (m, 2H), 3.18 (s, 3H), 3.25 (m, 2H), 3.75-3.80 (m, 1H), 3.83-3.95 (m, 1H), 4.75 (m, 1H), 5.30 (m, 1H), 6.70 (s, 1H), 6.80 (m, 1H), 6.95 (m, 1H), 7.40 (m, 2H), 7.50 (m, 1H), 7.63 (m, 1H), 8.28 (s, 1H). MS m/z 500 [M+H]⁺

Example 148

N-[2-({6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl]phenyl]-N-methylmethanesulfonamide

To a solution of 6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 334**, 100 mg, 0.21 mmol) in DCM (1 mL) was added N-[2-(aminomethyl)phenyl]-N-methylmethanesulfonamide (**PCT Intl. Appl. 2010058846**, 60 mg, 0.28 mmol) followed by PyBrop (130 mg, 0.28 mmol) and DIPEA (0.14 mL, 0.81 mmol). The reaction was

stirred at room temperature for 18 hours. The reaction was poured into saturated aqueous NaHCO_3 solution and extracted with DCM three times. The organic layers were collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 1:1 EtOAc in hexanes. The residue was dissolved in DCM (0.6 mL) and TFA (0.2 mL) followed by TES (0.05 mL) were added at 0°C. The reaction was stirred at room temperature for 18 hours before being quenched with saturated aqueous NaHCO_3 solution. The mixture was extracted into DCM, the organic layer collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 60% EtOAc in heptanes. The residue was dissolved in DCM (1 mL) and BBr_3 (0.72 mL) was added at 0°C. The reaction was stirred at room temperature for 18 hours before concentrating *in vacuo* and purifying by preparative HPLC to afford the title compound. MS m/z 524 $[\text{M}+\text{H}]^+$ R_t = 2.39 minutes.

Example 149

N-(2-({6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl}(methyl)amino)methyl)phenyl)-N-methylmethanesulfonamide


The title compound was prepared according to the method described for **Example 148** using N-methyl-N-(2-((methylamino)methyl)phenyl)methanesulfonamide (**Preparation 359**). MS m/z 538 $[\text{M}+\text{H}]^+$ R_t = 2.52 minutes.

Example 150

N-(2-({6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl}amino)methyl)-4-methylphenyl)-N-methylmethanesulfonamide

The title compound was prepared according to the method described for **Example 148** using the free base of N-[2-(aminomethyl)-4-methylphenyl]-N-methylmethanesulfonamide hydrochloride (**Preparation 189**). MS m/z 538 $[\text{M}+\text{H}]^+$ R_t = 2.47 minutes.

Library Protocol 2



To a 0.2M solution of amines of formula (XII) (1 mL, 200 μ mol) in nBuOH was added a 0.2M solution of 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (**PCT Intl. Appl. 2013014567**, 1 mL, 200 μ mol) followed by DIPEA (120 μ L, 700 μ mol). The reaction was heated to 80 °C for 16 hours before concentrating *in vacuo*. The residue was dissolved in 1:1 MeOH:toluene (1.5 mL). To the solution was added potassium carbonate (62 mg, 450 μ mol), PeppsiTM-IPr (3 mg, 4.5 μ mol) and (2-
 5 {2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-(2,2,2-trifluoroethyl)phenoxy}methoxy}ethyl)(trimethyl)silane (**Preparation 150**, 400 μ mol). The reaction was heated to 100 °C under microwave irradiation for 25 minutes before
 10 concentrating *in vacuo*. The residue was dissolved in EtOAc (5 mL) and washed with water (3 mL) and brine (3 mL). The organic extract was dried over sodium sulfate and concentrated *in vacuo*. The residue was dissolved in TFA (1 mL) and stirred at room temperature for 16 hours. The reaction was concentrated *in vacuo* and azeotroped with toluene. The residue was dissolved in MeOH and ethylenediamine (35 μ L, 500 μ mol)
 15 was added with stirring at room temperature for 18 hours. The reaction was concentrated *in vacuo*, dissolved in DMSO (1mL) and purified using preparative HPLC as described below:

LCMS:

A: 0.05% formic acid in water; B: MeCN; Column: RESTEK C18,
 20 30x2.1mmx3 μ m; Gradient: From 98% [A] and 2% [B] to 90% [A] and 10% [B] in 1 min, further to 98% [B] in 2 min and finally back to initial condition in 2.90 min, 1.5 mL/min flow rate.

Preparative HPLC:

Method A: Gemini NXC18 (100x20mmx5 μ); Acetonitrile-water (20 mM NH₄CO₃);
 25 Flow rate 20 mL/min; Gradient time 10 mins for 10-75% organic elution.

Method B: reposit Gold C18 (250x20mmx5 μ); Acetonitrile-water (20 mM NH₄CO₃); Flow rate 20 mL/min; Gradient time 18 mins for 10-70% organic elution.

The compounds of the Examples in the table below (**Examples 151 – 154**) were
 30 prepared and purified from 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (**PCT Intl. Appl. 2013014567**), (2-
 {2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-(2,2,2-trifluoroethyl)phenoxy}methoxy}ethyl)(trimethyl)silane
 (**Preparation 150**) and the appropriate amine according to Library Protocol 2.

Ex	Name	MS Data/Amine
151	4-{4-[(cyclopropylmethyl)-amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-2-fluoro-5-(2,2,2-trifluoroethyl)phenol	MS m/z 382 [M+H] ⁺ Rt = 1.53 minutes Cyclopropylmethylamine
152	4-{4-[(2-cyclopropylethyl)amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-2-fluoro-5-(2,2,2-trifluoroethyl)phenol	MS m/z 396 [M+H] ⁺ Rt = 1.57 minutes Cyclopropylethylamine
153	2-fluoro-4-{4-[(2-methylpropyl)amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-5-(2,2,2-trifluoroethyl)phenol	MS m/z 384 [M+H] ⁺ Rt = 1.56 minutes Isobutylamine
154	4-[4-(butylamino)-1H-pyrazolo[3,4-d]pyrimidin-6-yl]-2-fluoro-5-(2,2,2-trifluoroethyl)phenol	MS m/z 384 [M+H] ⁺ Rt = 1.57 minutes Butylamine

Example 155

N-[2-({[6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide hydrochloride

5 The title compound was prepared according to the method described for **Example 125** using N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (**Preparation 312**) and (2-([2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-ethylphenoxy]methoxy)ethyl)(trimethyl) silane (**PCT Intl. Appl. WO2013014567A1**). SPhos
10 was used as the ligand and the final residue was triturated with pentane/ether to afford the hydrochloride salt. ¹H NMR (400MHz, DMSO-d₆): δ ppm 0.91 (t, 3H), 2.66 (m, 2H), 3.06 (s, 3H), 3.13 (s, 3H), 4.82 (br s, 1H), 5.06 (br s, 1H), 6.86 (d, 1H), 7.33-7.46 (m, 4H), 7.55 (d, 1H), 8.55 (br s, 1H), 10.33 (br s, 1H), 14.56 (br s, 1H). MS m/z 471 [M+H]⁺

Example 156

15 N-(2-(((6-(2-Ethyl-4-hydroxy-6-methylphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

The title compound was prepared according to the method described for **Preparation 299** pyrimidines using 2-(4-(benzyloxy)-2-ethyl-6-methylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**Preparation 342**) followed by treating the residue with
20 TFA at reflux. The reaction was concentrated *in vacuo* and partitioned between EtOAc and saturated aqueous sodium bicarbonate solution. The organic extracts were washed

with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative HPLC. ¹H NMR (400MHz, DMSO-d₆): δ ppm 0.63-0.82 (m, 3H), 1.86 (m, 5H), 2.91 (s, 3H), 2.94 (s, 3H), 4.37 (br m, 1H), 4.85 (br m, 1H), 6.36 (br m, 3H), 7.20 (m, 2H), 7.33 (m, 2H), 8.17 (m, 1H), 9.03 (m, 1H), 12.88 (br s, 1H). MS m/z 466 [M+H]⁺

Example 157

N-[2-({[6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-3-(1H-imidazol-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide

A solution of N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide (**Preparation 79**, 200 mg, 0.22 mmol) and 2-bromo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-imidazole (**J. Org. Chem. (2010) 75 (15) 4911-4920**, 62.102 mg, 0.22 mmol) in toluene (2 mL) was degassed with nitrogen for 5 minutes. Bis(tributyltin) (0.27 mL, 0.54 mmol) and copper (I) iodide (8.53 mg, 0.045 mmol) were added followed by Pd(PPh₃)₄ (25.88 mg, 0.022 mmol) and the reaction was heated to 100 °C for 6.5 hours. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 15% EtOAc in hexanes. The residue (80 mg, 0.086 mmol) was treated with TFA (2 mL) and the solution stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo*, dissolved in MeOH (5 mL) and cooled in ice water. Ethylene diamine was added dropwise until the solution was basic, with stirring for 15 minutes. The solution was concentrated *in vacuo* and purified using silica gel column chromatography eluting with 60% EtOAc in hexanes to afford the title compound (25 mg, 54%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 0.91 (t, 3H), 2.43 (m, 2H), 3.06 (s, 3H), 3.10 (s, 3H), 4.70 (br m, 1H), 5.00 (br m, 1H), 6.58 (m, 1H), 6.77 (m, 1H), 7.05 (m, 1H), 7.10 (s, 1H), 7.26-7.31 (m, 3H), 7.47 (m, 2H), 9.74 (s, 1H), 10.89 (t, 1H), 12.93 (s, 1H), 13.29 (s, 1H). MS m/z 536 [M+H]⁺

Example 158

N-[2-({[3-(4,5-Dimethyl-1H-imidazol-2-yl)-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide

The title compound was prepared according to the method described by **Example 157** using N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide (**Preparation 79**) and 2-bromo-

4,5-dimethyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole (**Preparation 386**) at 115 °C under microwave irradiation for 30 minutes. ¹H NMR (400MHz, DMSO-d₆): δ ppm 0.97 (t, 3H), 2.56 (m, 2H), 3.04 (s, 3H), 3.10 (s, 3H), 4.60 (br m, 1H), 5.00 (br m, 1H), 6.55 (s, 1H), 6.79 (m, 1H), 7.08 (m, 1H), 7.35 (m, 2H), 7.49 (m, 1H), 7.51 (m, 1H), 9.74 (s, 1H), 10.95 (t, 1H), 12.40 (s, 1H), 13/15 (s, 1H). MS m/z 564 [M+H]⁺

Example 159

N-[2-({[6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-3-(4-methyl-1H-imidazol-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide

The title compound was prepared according to the method described by **Example 157** using N-[2-({[6-(2-ethyl-5-fluoro-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide (**Preparation 79**) and 2-iodo-5-methyl-1H-imidazole at 115 °C under microwave irradiation for 30 minutes. Following deprotection the residue was purified using preparative HPLC. ¹H NMR (400MHz, DMSO-d₆): δ ppm 0.95 (t, 1.5H), 0.99 (t, 1.5H), 1.99 (s, 1.5H), 2.32 (s, 1.5H), 3.05 (m, 3H), 3.10 (m, 3H), 4.66 (br m, 1H), 4.99 (br m, 1H), 6.56 (m, 1H), 6.77-6.90 (m, 2H), 7.01-7.09 (m, 1H), 7.27-7.36 (m, 2H), 7.46-7.61 (m, 2H), 9.74 (br s, 1H), 10.95 (m, 1H), 12.55 (s, 0.5H), 12.69 (s, 0.5H), 13.22 (br s, 1H). MS m/z 550 [M+H]⁺

Example 160 Intermediate

2-Fluoro-4-(4-((2-(methylthio)ethyl)amino)-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl)-5-(2,2,2-trifluoroethyl)phenol

The title compound was prepared according to the method described by **Example 157** using 6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-N-(2-(methylthio)ethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-amine (**Preparation 379**) and tert-butyl 2-iodo-6,7-dihydro-1H-imidazo[4,5-c]pyridine-5(4H)-carboxylate (**WO2013014567**). MS m/z 522 [M+H]⁺

Example 161

4-[3-(5-Benzyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-4-{{2-(methylsulfanyl)-ethyl}amino}-1H-pyrazolo[4,3-c]pyridin-6-yl]-2-fluoro-5-(2,2,2-trifluoroethyl)phenol

To a suspension of anhydrous magnesium sulfate (40 mg, 0.33 mmol) and 2-fluoro-4-(4-((2-(methylthio)ethyl)amino)-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl)-5-(2,2,2-trifluoroethyl)phenol (**Example 161**, 44 mg, 0.08 mmol) in methanol (2.5 mL) was added a solution of benzaldehyde (0.017 mL,

0.17 mmol) in methanol (2.5 mL). The reaction was stirred for 1 hour at 55 °C before cooling to room temperature and adding sodium cyanoborohydride (10.6 mg, 0.17 mmol). The reaction was stirred at room temperature for 18 hours. The reaction was filtered and concentrated *in vacuo*. The residue was partitioned between 20% IPA in DCM and water. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative TLC to afford the title compound (20 mg, 39%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 1.90 (s, 1.5H), 1.98 (s, 1.5H), 2.66-2.88 (m, 6H), 3.47 (m, 2H), 3.64-3.72 (m, 4H), 4.03-4.11 (m, 2H), 6.62 (s, 1H), 7.01 (m, 1H), 7.27-7.38 (m, 6H), 10.11 (s, 1H), 10.68 (t, 0.5H), 10.75 (t, 0.5H), 12.46 (s, 0.5H), 12.58 (s, 0.5H), 13.23 (s, 1H). MS m/z 612 [M+H]⁺

Example 162

N-[2-({[6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-3-(1H-pyrazol-1-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide

To a solution of N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide (**Preparation 79**, 250 mg, 0.29 mmol) in DMSO (0.5 mL) was added pyrazole (19.88 mg, 0.29 mmol), PEG (500 mg), cesium carbonate (133 mg, 0.41 mmol), cuprous oxide (1.25 mg, 0.01 mmol) and 4,7-dimethoxy-1,10-phenanthroline (5.61 mg, 0.023 mmol) and the reaction was heated to 110 °C for 18 hours. The reaction was cooled, diluted with EtOAc and washed with water and brine. The organic layer was dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 27% EtOAc in hexanes. The residue (77 mg, 0.097 mmol) was treated with TFA (2.5 mL) and stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo*, dissolved in MeOH (5 mL) and cooled in ice water. Ethylene diamine was added dropwise until the solution was basic, with stirring for 15 minutes. The solution was concentrated *in vacuo* and purified using silica gel column chromatography eluting with 6% MeOH in DCM to afford the title compound (32 mg, 62%).

¹H NMR (400MHz, DMSO-d₆): δ ppm 0.93 (t, 3H), 2.49 (m, 2H), 3.05 (s, 3H), 3.07 (s, 3H), 4.67 (br m, 1H), 5.00 (br m, 1H), 6.60 (s, 1H), 6.66 (m, 1H), 6.80 (m, 1H), 7.05 (m, 1H), 7.31 (m, 2H), 7.45 (m, 2H), 7.91 (s, 1H), 8.54 (s, 1H), 9.61 (t, 1H), 9.76 (s, 1H), 13.11 (s, 1H). MS m/z 536 [M+H]⁺

Example 163

N-{2-[(6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(1H-pyrazol-1-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl]phenyl}-N-methylmethanesulfonamide

The title compound was prepared according to the method described for
 5 **Example 162** using N-methyl-N-(2-[(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[(2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-[(2-(trimethylsilyl)ethoxy)methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl]phenyl)methanesulfonamide (**Preparation 105**) and pyrazole. Following deprotection the residue was purified using silica gel column chromatography eluting with 45% EtOAc in hexanes. ¹H NMR (400MHz, DMSO-d₆): δ ppm 3.06 (m, 6H), 3.40 (m, 2H), 4.64 (br m, 1H), 4.93 (br m, 1H), 6.66 (m, 2H), 6.94 (m, 1H), 7.19-7.22 (m, 1H), 7.28-7.34 (m, 2H), 7.41-7.50 (m, 2H), 7.93 (s, 1H), 8.57 (s, 1H), 9.71 (t, 1H), 10.11 (br s, 1H), 13.21 (br s, 1H). MS m/z 590 [M+H]⁺

Example 164

N-{2-[(6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(5-methyl-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl]phenyl}-N-methylmethanesulfonamide

To a solution of acetimidamide hydrochloride (33 mg, 0.35 mmol) in 2-methoxyethanol (3 mL) was added DIPEA (0.087 mL, 0.50 mmol) followed by N-(2-[(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[(2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-(hydrazinecarbonyl)-1-[(2-(tri-methylsilyl)ethoxy)methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl]phenyl)-N-methyl-methanesulfonamide (**Preparation 374**, 120 mg, 0.143 mmol). The reaction was stirred at 85 °C for 18 hours. The reaction was quenched with water and extracted with DCM. The organic extracts were collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using
 25 silica gel column chromatography eluting with 27% EtOAc in DCM. The residue (100 mg, 0.12 mmol) was treated with TFA (2 mL) and stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo*, dissolved in MeOH (5 mL) and cooled in ice water. Ethylene diamine was added dropwise until the solution was basic, with stirring for 15 minutes. The solution was concentrated *in vacuo* and purified using
 30 preparative TLC to afford the title compound (40 mg, 54%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 1.75 (s, 3H), 3.04 -3.12 (m, 6H), 3.68 (m, 2H), 4.70 (br m, 1H), 5.00 (br m, 1H), 6.67 (m, 1H), 6.97 (m, 1H), 7.20-7.22 (m, 1H), 7.25-7.29 (m, 2H), 7.34 (m, 2H), 7.85 (m, 1H), 10.08-10.25 (m, 1H), 13.36 (s, 0.5H), 13.71 (s, 0.5H), 14.02 (s, 0.5H), 14.52 (s, 0.5H). MS m/z 605 [M+H]⁺

Example 165

N-[2-({[6-(2-Ethyl-5-fluoro-4-hydroxyphenyl)-3-(5-methyl-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide

To a solution of 6-[5-fluoro-2-ethyl-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-4-
 5 ({{2-[methyl(methylsulfonyl)amino]benzyl}amino)-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxylic acid (**Preparation 373**, 180 mg, 0.23 mmol) in THF (12 mL) was added hydrazine hydrochloride (39.82 mg, 0.58 mmol), BOP (257 mg, 0.58 mmol) and DIPEA (0.122 mL, 0.70 mmol). The reaction was stirred at room temperature for 18 hours. The reaction was quenched with water, extracted into DCM, the organic
 10 extracts collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 8% MeOH in DCM. The residue was added to a solution of acetimidamide hydrochloride (30 mg, 0.32 mmol) and DIPEA (0.08 mL, 0.44 mmol) in 2-methoxyethanol (3 mL) and the reaction was heated to 85 °C for 18 hours. The reaction was cooled and quenched by the addition of
 15 water and extracted into DCM. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative TLC. The residue was treated with TFA (2 mL) and stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo*, dissolved in MeOH (5 mL) and cooled in ice water. Ethylene diamine was added dropwise until the
 20 solution was basic, with stirring for 15 minutes. The solution was concentrated *in vacuo* and purified using preparative TLC to afford the title compound (30 mg, 88%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 0.97 (t, 3H), 2.20 (s, 3H), 2.56 (m, 2H), 3.04 (s, 3H), 3.12 (s, 3H), 4.65 (br m, 1H), 5.00 (br m, 1H), 6.66 (m, 1H), 6.83 (m, 1H), 7.02 (m, 1H), 7.33 (m, 2H), 7.51-7.57 (m, 2H), 9.78 (m, 1H), 10.13 (m, 1H), 13.27 (s, 0.5H), 13.62 (s,
 25 0.5H), 13.99 (s, 0.5H), 14.49 (s, 0.5H). MS m/z 549 [M-H]⁺

Example 166

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-[5-(6-methylpyridin-3-yl)-4H-1,2,4-triazol-3-yl]-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl}-N-methylmethanesulfonamide

30 To a solution of N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(hydrazinecarbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (**Preparation 376**, 160 mg, 0.22 mmol) in n-butanol (2 mL) was added 5-cyano-2-methyl pyridine (65 mg, 0.55 mmol) and potassium carbonate (16 mg, 0.12 mmol). The reaction mixture was heated to 150

°C under microwave irradiation for 50 minutes. The reaction was cooled and concentrated *in vacuo*. The residue was partitioned between water and ethyl acetate. The organic extracts were washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 15% MeOH in DCM followed by preparative TLC. The residue was treated with boron tribromide (0.047 mL 0.47 mmol) and the reaction stirred at room temperature for 30 minutes. The reaction was concentrated *in vacuo* and partitioned between saturated aqueous NaHCO₃ and EtOAc. The organic extracts were dried over sodium sulfate, concentrated *in vacuo* and purified by preparative TLC to afford the title compound (19 mg, 41%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 1.22 (s, 3H), 3.03 (s, 3H), 3.07 (s, 3H), 3.71 (m, 2H), 4.75 (br m, 1H), 5.05 (br m, 1H), 6.78 (s, 1H), 6.96 (m, 1H), 7.22-7.38 (m, 4H), 7.40 (m, 2H), 7.75 (m, 1H), 8.93 (m, 1H), 10.01 (br m, 1H), 13.80 (br m, 1H). MS m/z 682 [M+H]⁺

Example 167

4-(5-{6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-{5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino}-1H-pyrazolo[3,4-d]pyrimidin-3-yl}-4H-1,2,4-triazol-3-yl)piperidine-1-carboxamide

To a solution of N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(5-(piperidin-4-yl)-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 371**, 70 mg, 0.097 mmol) in DCM (12 mL) was added triethylamine (0.02 mL, 0.146 mmol) and trimethylsilylisocyanate (0.013 mL, 0.097 mmol). The reaction was quenched with water, and the organic extracts were dried over sodium sulfate and concentrated *in vacuo*. The residue was dissolved in DCM (10 mL) and boron tribromide (0.041 mL, 0.41 mmol) was added at 0 °C and stirred for 3 hours. The reaction was quenched with saturated aqueous sodium bicarbonate solution followed by extraction with 20% IPA in DCM. The organic extracts were collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative TLC eluting with 10% MeOH in DCM to afford the title compound (34 mg, 78%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 1.42 (m, 2H), 1.75 (m, 2H), 2.67-2.75 (m, 2H), 3.01 (s, 3H), 3.08 (s, 3H), 3.17 (m, 1H), 3.89 (m, 2H), 4.34 (m, 2H), 4.69 (m, 1H), 4.99 (m, 1H), 5.93 (br s, 2H), 6.76 (m, 1H), 6.86 (m, 1H), 6.99 (m, 1H), 7.36 (m, 1H), 7.75 (m, 1H), 9.66 (br s, 1H), 10.38-10.50 (m, 2H), 13.72 (br s, 0.5H), 14.05 (br s, 0.5H), 14.20 (br s, 0.5H), 14.72 (br s, 0.5H). MS m/z 734 [M+H]⁺

Example 168

N-(2-(((6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-{5-[1-(pyrrolidin-1-yl)acetyl]piperidin-4-yl}-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

5 To a solution of N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(5-(piperidin-4-yl)-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-phenyl)-N-methylmethanesulfonamide (**Preparation 372**, 90mg, 0.13mmol) and 2-(pyrrolidin-1-yl)acetic acid (21 mg, 0.13 mmol) in DCM (10 mL) was added DIPEA (0.065 mL, 0.39 mmol) followed by BOP (58 mg, 0.13 mmol). The reaction was allowed
10 to stir at room temperature for 18 hours. The reaction was concentrated *in vacuo* and partitioned between 20% IPA in DCM and water. The organic extracts were washed with brine, dried and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 12% MeOH in DCM. The residue was dissolved in DCM and treated with boron tribromide (0.083 mL, 0.87 mmol) at 0°C. The reaction was
15 stirred at room temperature for 18 hours before the addition of another aliquot of boron tribromide (0.25 mL) and further stirring for 3 hours. The reaction was quenched by the addition of saturated aqueous sodium bicarbonate solution and extracted with 20% IPA in DCM. The organic extract was separated, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with
20 15% MeOH in DCM to afford the title compound (41 mg, 42%).

¹H NMR (400MHz, DMSO-d₆): δ ppm 1.33 (m, 1H), 1.48 (m, 1H), 1.71-1.90 (m, 6H), 2.59-2.70 (m, 3H), 2.97 (m, 1H), 3.06 (m, 4H), 3.13 (s, 3H), 3.31 (m, 2H), 3.90 (m, 1H), 4.08 (m, 1H), 4.19 (m, 1H), 4.31 (m, 2H), 4.79 (br m, 1H), 5.12 (br m, 1H), 6.99 (m, 1H), 7.35-7.59 (m, 4H), 7.73 (m, 1H), 10.44 (m, 2H), 13.93 (br s, 1H), 14.55 (br s, 1H).
25 MS m/z 786 [M+H]⁺

Example 169

N-(2-(((6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-hydroxyphenyl)-N-methylmethanesulfonamide

30 The title compound was prepared according to the method described for **Example 29** using N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 370**).
¹H NMR (400MHz, DMSO-d₆): δ ppm 2.32 -2.67 (m, 2H), 2.91 (m, 2H), 3.02 (s, 3H),

3.10 (s, 3H), 3.32 (m, 2H), 4.34 (m, 2H), 4.67 (m, 1H), 5.00 (m, 1H), 6.72 (m, 1H), 6.74 (m, 1H), 6.98 (m, 1H), 7.33 (m, 1H), 7.72 (m, 1H), 9.65 (m, 1H), 11.26-11.33 (m, 1H), 12.62 (m, 1H), 13.59 (m, 1H). MS m/z 662 [M+H]⁺

Example 170

5 N-{2-[(3-(5-Acetyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[3,4-d]pyrimidin-4-yl}amino)methyl]-4-hydroxyphenyl}-N-methylmethanesulfonamide

To a solution of N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 370**, 10 200 mg, 0.29 mmol) and DIPEA (0.096 ml, 0.58 mmol) in DCM (35 mL) at 0 °C was added acetyl chloride (0.021 mL, 0.29 mmol) and the reaction stirred at room temperature for 2 hours. The reaction was concentrated *in vacuo* and partitioned between EtOAc and water. The organic phase was dried, concentrated *in vacuo* and 15 purified by silica gel column chromatography eluting with 5% MeOH in DCM. The residue was dissolved in DCM (10 mL) and treated with boron tribromide (0.083 mL, 0.87 mmol) at 0°C. The reaction was stirred at room temperature for 18 hours before the addition of another aliquot of boron tribromide (0.25 mL) with further stirring for 3 hours. The reaction was quenched by the addition of saturated aqueous sodium 20 bicarbonate solution and extracted with 20% IPA in DCM. The organic extract was separated, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative TLC eluting with 10% MeOH in DCM to afford the title compound (25 mg, 38%). ¹H NMR (400MHz, DMSO-d₆): δ ppm (tautomers?) 2.50 (m, 3H), 2.62 (m, 2H), 3.02 (s, 3H), 3.15 (s, 3H), 3.70 (m, 3H), 4.35 (m, 4H), 4.54 (m, 1H), 5.00 (m, 1H), 6.71-7.00 (m, 3H), 7.35 (m, 1H), 7.70 (m, 1H), 9.60 (m, 1H), 10.37 (m, 1H), 11.54- 25 11.67 (m, 1H), 13.65 (s, 1H). MS m/z 704 [M+H]⁺

Example 171

N-[2-(Dimethylamino)ethyl]-2-{6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl}-1,4,6,7-tetrahydro-5H-imidazo[4,5-c]pyridine-5-carboxamide

30

To a solution of N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 370**, 80

mg, 0.116 mmol) in DCM (5 mL), was added boron tribromide (0.077 mL, 0.81 mmol) at 0 °C and stirred for 2 hours. Another aliquot of boron tribromide (7 eq) was added and the reaction mixture was stirred at room temperature for a further 2 hours. The reaction was quenched by the addition of saturated aqueous sodium bicarbonate solution and
 5 extracted with 20% IPA in DCM. The organic extract was separated, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by preparative TLC. The residue was dissolved in THF (1 mL) and added to a solution of N,N-dimethylamine (6 mL) in THF (1.5 mL) and bromoethylisocyanate (0.02 mL, 0.18 mmol) that had stirred at 0 °C for 10 minutes. The reaction was stirred at room temperature for 18 hours. The
 10 reaction was concentrated *in vacuo* and purified using preparative TLC to afford the title compound (15 mg, 21%). 25 minute HPLC QC (Sunfire C18 (150x4.6mmx5u), mobile phase A = MeCN, mobile phase B = 10 mM ammonium acetate in water Rt = 2.59 minutes. MS m/z 776 [M+H]⁺

Example 172 Intermediate

15 Racemic 2-Fluoro-4-(4-((3-hydroxy-2-methylpropyl)amino)-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)-5-(2,2,2-trifluoroethyl)phenol

The title compound was prepared according to the method described for **Example 157** using tert-butyl 2-iodo-6,7-dihydro-1H-imidazo[4,5-c]pyridine-5(4H)-carboxylate (**WO2013014567**) and racemic 3-((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)-methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-2-methylpropan-1-ol (**Preparation 380**). MS m/z 521 [M+H]⁺
 20

Example 173

25 4-(3-(5-Benzyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-4-((3-hydroxy-2-methylpropyl)amino)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)-2-fluoro-5-(2,2,2-trifluoroethyl)phenol

The title compound was prepared according to the method described for **Example 161** using racemic 2-fluoro-4-(4-((3-hydroxy-2-methylpropyl)amino)-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)-5-(2,2,2-trifluoroethyl)phenol (**Example 172**). The residue was purified by preparative HPLC. 10 minute HPLC QC (Gemini NX-C18 (50x4.6mmx3u), mobile phase A = 0.05% formic acid in water, mobile phase B = MeCN Rt = 4.20 minutes MS m/z 611 [M+H]⁺
 30

Example 174 Intermediate

6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-hydroxy-2-(N-methylmethylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid

The title compound was prepared according to the method described for
 5 **Preparation 11** using N-(2-(((6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-hydroxyphenyl)-N-methylmethanesulfonamide (**Preparation 271**). MS m/z 585 [M+H]⁺

Preparation 1

6-(2-Ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-N-methyl-4-{[2-(methyl(sulfamoyl)amino)benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-N-methyl-4-{[2-(methylamino)benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 14**, 500 mg, 0.705 mmol) in THF
 15 (2 mL) was added NaH (28.2 mg, 0.70 mmol) at 0 °C followed by dropwise addition of sulfamoyl chloride (97 mg, 0.84 mmol). The reaction was allowed to warm to room temperature for 1 hour. The reaction was quenched with water and extracted with ethyl acetate. The organic extracts were separated, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography
 20 eluting with 40% EtOAc in Hexanes to afford the title compound as an off-white solid (350 mg, 62%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.11 (s, 9H), -0.02 (s, 9H), 0.80 (t, 2H), 0.84-0.91 (m, 5H), 2.59 (m, 2H), 2.83 (s, 3H), 3.01 (s, 3H), 3.57 (t, 2H), 3.75 (t, 2H), 4.75 (br s, 1H), 5.00 (br s, 1H), 5.29 (s, 2H), 5.72 (s, 2H), 6.96 (s, 1H), 7.04 (s, 1H), 7.13 (m, 1H), 7.22-7.43 (m, 4H), 8.85 (m, 1H), 9.68 (m, 1H). MS m/z 788 [M+H]⁺

Preparation 2

6-[5-Fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-4-{[2-(methyl{[3-(morpholin-4-yl)propyl]sulfonyl}amino)benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 4-{[2-((3-chloropropyl)sulfonyl)(methyl)amino]benzyl}amino]-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 4**,
 30 290 mg, 0.32 mmol) in EtOH (2 mL) was added morpholine (0.5 mL) and the reaction was heated to 110 °C under microwave irradiation for 75 minutes. The reaction was cooled, concentrated *in vacuo* and partitioned between EtOAc and water. The organic

layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using preparative TLC to afford the title compound (70 mg, 80%). MS m/z 940 [M+H]⁺

Preparation 3

5 6-(5-Fluoro-2-(2,2,2-trifluoroethyl)-4-((2-trimethylsilyl)ethoxy)methoxy)phenyl)-N-methyl-4-((2-(N-methyl-1H-pyrazole-4-sulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-fluoro-4-[[2-(trimethylsilyl)ethoxy]-methoxy]phenyl)-N-methyl-4-[[2-(methylamino)benzyl]amino]-1-[[2-(trimethylsilyl)ethoxy]-methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 10
17, 150 mg, 0.19 mmol) in THF (10 mL) was added 1H-pyrazole-4-sulfonylchloride (0.03 mL, 0.19 mmol) at 0°C. The reaction mixture was allowed to stir at room temperature for 18 hours. The reaction was concentrated *in vacuo* and purified using silica gel column chromatography to afford the title compound (76 mg, 43%). ¹H NMR
15 (400 MHz, DMSO-d₆): δ ppm -0.11 (s, 9H), -0.03 (s, 9H), 0.79-0.89 (m, 4H), 2.86 (d, 3H), 2.99 (s, 3H), 3.57 (t, 2H), 3.76 (m, 4H), 4.80 (br m, 1H), 4.97 (br m, 1H), 5.30 (s, 2H), 5.73 (s, 2H), 6.70 (m, 1H), 7.09 (s, 1H), 7.18-7.71 (m, 5H), 7.71 (s, 1H), 8.29 (s, 1H), 8.90 (m, 1H), 9.84 (m, 1H), 13.75 (s, 1H). MS m/z 893 [M+H]⁺

20 The following Preparations (Preparations 4 – 10) were prepared according to the method described for Preparation 3 using either 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)-ethoxy]methoxy]phenyl]-4-[[2-(methylamino)benzyl]amino]-1-[[2-(trimethylsilyl)ethoxy]-methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 16)
25 or 6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-fluoro-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl)-N-methyl-4-[[2-(methylamino)benzyl]amino]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 17) and the appropriate sulfonyl chloride as described below:

30

Preparation Number	Name	Data
4	4-[(2-[(3-chloropropyl)sulfonyl]-(methyl)amino)benzyl)amino]-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 889 [M+H] ⁺ Using 3-chloropropanesulfonyl chloride.
5	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-4-[[2-(methyl[(6-morpholin-4-yl)pyridin-3-yl)sulfonyl]amino)benzyl]amino]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 975 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm -0.11 (s, 9H), -0.03 (s, 9H), 0.80-0.89 (m, 4H), 2.99 (s, 3H), 3.56-3.84 (m, 14H), 4.78 (m, 1H), 4.93 (m, 1H), 5.30 (s, 2H), 5.73 (s, 2H), 6.80 (d, 1H), 6.90 (d, 1H), 7.10 (s, 1H), 7.19-7.33 (m, 3H), 7.42 (m, 1H), 7.63 (m, 1H), 7.98 (s, 1H), 8.27 (m, 2H), 9.83 (m, 1H). Using 6-morpholin-4-yl-pyridine-3-sulfonyl chloride.
6	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-N-methyl-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino)benzyl)amino]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 918 [M+H] ⁺ Using 6-methylpyridine-3-sulfonyl chloride.
7	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino)benzyl)amino]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 904 [M+H] ⁺ Using 6-methylpyridine-3-sulfonyl chloride.

8	4-((2-(N,1-dimethyl-1H-imidazole-4-sulfonamido)benzyl)amino)-6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-N-methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 907 [M+H] ⁺ Using 1-methyl-1H-imidazole-4-sulfonylchloride.
9	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}-methoxy}phenyl]-4-[(2-((2-methoxyethyl)sulfonyl)(methyl)amino)benzyl]amino]-N-methyl-1-{2-(trimethylsilyl)ethoxy}methyl-1H-pyrazolo-[4,3-c]pyridine-3-carboxamide	MS m/z 885 [M+H] ⁺ Using 2-methoxyethanesulfonyl chloride.
10	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}-methoxy}phenyl]-4-[(2-[methyl-(pyridin-3-ylsulfonyl)amino]benzyl)amino]-1-{2-(trimethylsilyl)ethoxy}methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	Using pyridine-3-sulfonyl chloride. Taken directly on to the next step.

Preparation 11

6-[5-Fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-4-({5-methoxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1-{2-(trimethylsilyl)ethoxy}methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxylic acid

To a solution of N-(2-[(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}-methoxy}phenyl]-3-iodo-1-{2-(trimethylsilyl)ethoxy}methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)-amino]methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 85**, 250 mg, 0.26 mmol) in MeOH (4 mL) was added molybdenum hexacarbonyl (84.91 mg, 0.32 mmol), DBU (0.119 mL, 0.80 mmol) and Pd(OAc)₂ (4 mg, 0.02 mmol). The reaction was heated to 125 °C for 15 minutes under microwave irradiation. The reaction was cooled, diluted with EtOAc and filtered through celite. The filtrate was concentrated *in vacuo* and purified using silica gel column chromatography eluting with 10% MeOH in DCM to afford the title compound (100 mg, 51%). MS m/z 858 [M+H]⁺

Preparation 12

6-[5-Fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-4-
 ({[2-(methyl(methylsulfonyl)amino)benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl})-1H-
 pyrazolo[4,3-c]pyridine-3-carboxylic acid

5 The title compound was prepared according to the method described for **Preparation 11** using N-methyl-N-(2-{[(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl})-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl)methanesulfonamide (**Preparation 105**). MS m/z 829 [M+H]⁺

Preparation 13

6-(5-Fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-N-
 methyl-1-((2-(trimethylsilyl)ethoxy)methyl)-4-((2-(1,3,3-trimethylureido)benzyl)amino)-
 1H-pyrazolo[4,3-c]pyridine-3-carboxamide

10 To a solution of 6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-fluoro-4-{[2-(trimethylsilyl)ethoxy]-methoxy}phenyl)-N-methyl-4-{[2-(methylamino)benzyl]amino}-1-
 15 {[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 17**, 50 mg, 0.06 mmol) in THF (5 mL) was added sodium hydride (1.88 mg, 0.08 mmol) at 0°C. After stirring for 2 minutes, dimethylsulfamoyl chloride (15 mg, 0.11 mmol) was added and the reaction was stirred for 1 hour. The reaction was
 20 partitioned between EtOAc and water, the organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was dissolved in DMF (1 mL) and treated with cesium carbonate (64 mg, 0.19 mmol) followed by methyl iodide (27 mg, 0.19 mmol). The reaction was stirred at room temperature for 18 hours before quenching with ammonium chloride and extraction with EtOAc. The organic layer was
 25 collected and purified using preparative TLC to afford the title compound (45 mg, 78%). MS m/z 870 [M+H]⁺

Preparation 14

6-(2-Ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-N-methyl-4-{[2-
 (methyl-amino)benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-
 c]pyridine-3-carboxamide

30 To a solution of benzyl [2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-(methylcarbamoyl)-1-{[2-(trimethylsilyl)ethoxy]methyl})-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl)methylcarbamate (**Preparation 18**, 775 mg, 0.91 mmol) in EtOH (25 mL) was added 10% Pd/C (100 mg) and the reaction

was hydrogenated at room temperature at 30 psi for 1 hour. The reaction was filtered, the filtrate was concentrated *in vacuo* and the residue was purified by silica gel column chromatography eluting with 15% EtOAc in hexanes to afford the title compound (530 mg, 81%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.11 (s, 9H), -0.01 (s, 9H), 0.81 (t, 2H), 0.88 (t, 2H), 1.00 (t, 2H), 2.38 (s, 3H), 2.69 (m, 2H), 2.82 (s, 3H), 3.56 (t, 2H), 3.77 (t, 2H), 4.59 (m, 2H), 5.33 (s, 2H), 5.71 (s, 2H), 6.02 (m, 1H), 6.44 (m, 1H), 6.55 (m, 1H), 6.98 (s, 1H), 7.04-7.19 (m, 2H), 7.22 (m, 1H), 8.84 (m, 1H), 9.69 (m, 1H). MS m/z 709 [M+H]⁺

Preparation 15

10 6-[5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound was prepared according to the method described for **Preparation 14** using 6-[4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 22**). The residue was purified by silica gel column chromatography eluting with 25% EtOAc in DCM. ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.58 (m, 2H), 1.74 (m, 1H), 1.93-2.10 (m, 3H), 3.10 (s, 1H), 3.12 (s, 3H), 3.61 (m, 1H), 3.74 (m, 2H), 3.90 (m, 1H), 4.82 (m, 2H), 5.88 (m, 1H), 6.95 (d, 1H), 7.03 (s, 1H), 7.25 (d, 1H), 7.36 (m, 1H), 7.81 (m, 1H), 7.97 (br s, 1H), 8.19 (br s, 1H), 8.41 (m, 1H), 9.82 (t, 1H), 10.15 (s, 1H). MS m/z 652 [M+H]⁺

Preparation 16

25 6-[5-Fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-4-{[2-(methyl-amino)benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound was prepared according to the method described for **Preparation 14** using benzyl (2-{[(3-carbamoyl-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl)-amino]methyl}phenyl)methylcarbamate (**Preparation 23**). The residue was purified by silica gel column chromatography eluting with 30% EtOAc in hexanes. MS m/z 749 [M+H]⁺

Preparation 17

6-(5-Fluoro-2-(2,2,2-trifluoroethyl)-4-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-N-methyl-4-{[2-(methylamino)benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

5 The title compound was prepared according to the method described for **Preparation 14** using benzyl [2-({[6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-(methylcarbamoyl)-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl)methylcarbamate (**Preparation 19**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.11 (s, 9H), -0.01 (s, 9H), 0.81 (t, 2H), 0.89 (t, 2H), 2.50 (s, 3H), 2.84 (d, 3H), 3.58 (t, 2H), 3.76 (t, 2H), 3.95 (q, 10 2H), 4.58 (d, 2H), 5.33 (s, 2H), 5.73 (s, 2H), 5.74 (br s, 1H), 6.45-6.54 (m, 2H), 7.05-7.11 (m, 3H), 7.35-7.39 (m, 2H), 8.87 (m, 1H), 9.73 (m, 1H). MS m/z 763 [M+H]⁺

Preparation 18

Benzyl [2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-(methylcarbamoyl)-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl)methylcarbamate

To a solution of benzyl [2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl)methylcarbamate (**Preparation 91**, 1 20 g, 1 mmol) in methylamine/THF (10 mL) was added DBU (0.49 mL, 3.23 mmol), Pd(OAc)₂ (17 mg, 0.08 mmol) followed by molybdenum hexacarbonyl (0.29 mg, 1.09 mmol). The reaction was heated to 100 °C under microwave irradiation for 10 minutes. The reaction was cooled, concentrated *in vacuo* and diluted with EtOAc. The mixture was filtered through celite, the filtrate concentrated *in vacuo* and purified using silica gel 25 column chromatography eluting with 47% EtOAc in hexanes to afford the title compound (775 mg, 84%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.10 (s, 9H), -0.01 (s, 9H), 0.79 (t, 2H), 0.90 (m, 5H), 2.57 (m, 2H), 2.83 (d, 3H), 3.08 (s, 1H), 3.59 (t, 2H), 3.74 (t, 2H), 4.53 (m, 1H), 4.71 (m, 1H), 4.88 (m, 1H), 5.00 (m, 1H), 5.27 (s, 1H), 5.75 (s, 2H), 6.98 (s, 1H), 7.08-7.42 (m, 11H), 8.83 (m, 1H), 9.67 (m, 1H). MS m/z 843 [M+H]⁺

30

Preparation 19

Benzyl [2-({[6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-(methylcarbamoyl)-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl)methylcarbamate

A solution of 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide-5-oxide (**Preparation 117**, 3.2 g, 4.96 mmol) in DMF (100 mL) was treated with benzyl methyl[2-({[(4-nitrophenoxy)carbonyl]amino}-methyl)phenyl]carbamate (**Preparation 178**, 2.68 g, 6.16 mmol) and triethylamine (0.68 mL, 4.96 mmol) and heated at 80 °C for 16 hours. Further benzyl methyl[2-({[(4-nitrophenoxy)carbonyl]amino}-methyl)phenyl]carbamate (1.24 eq) and triethylamine (1 eq) were added and the reaction allowed to continue for 6 hours. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography to afford the title compound as an oil (4.2 g, 94%). MS m/z 897 [M+H]⁺

Preparation 20

6-(2-Ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-4-{[2-methyl(methylsulfonyl)amino]benzyl}amino)-N-(6-methylpyridin-3-yl)-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound was prepared according to the method described for **Preparation 18** using N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]-N-methylmethanesulfonamide (**Preparation 79**) and 6-methylpyridin-3-amine with DBU at 100 °C for 10 minutes under microwave irradiation. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 7% heptanes in EtOAc. MS m/z 864 [M+H]⁺

Preparation 21

6-[5-Fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-4-({[4-methoxy-2-[methyl(methylsulfonyl)amino]benzyl]amino)-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of N-(2-({[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]-amino}methyl)-5-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 89**, 350 mg, 0.37 mmol) in MeOH (2 mL) was added DBU (0.16 mL, 1.19 mmol), palladium acetate (5.86 mg, 0.03 mmol) and molybdenum hexacarbonyl (99 mg, 0.37 mmol) and the reaction was heated to 100 °C for 10 minutes under microwave irradiation. The reaction was cooled, concentrated *in vacuo* and purified directly using silica gel column chromatography eluting with 12% MeOH in DCM. The resulting residue was dissolved in anhydrous THF (5 mL) and NMM (0.033 mL, 0.30 mmol) was

added followed by isobutylchloroformate (0.04 mL, 0.30 mmol) at -20°C. The reaction was stirred for 2 hours at this temperature before the addition of aqueous ammonia (0.5 mL) with further stirring for 1 hour. The reaction was partitioned between EtOAc and water, the organic layer collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 42% EtOAc in hexanes to afford the title compound (102 mg, 32% over 2 steps). MS m/z 857 [M+H]⁺

The following Preparations (**Preparations 22 – 26**) were prepared according to the method described for **Preparation 21** using the appropriate iodo intermediate as described below:

Preparation Number	Name	Data
22	6-[4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 742 [M+H] ⁺ Using N-{3-[(6-[4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl]pyridin-2-yl}-N-methylmethanesulfonamide (Preparation 93).
23	Benzyl (2-[(3-carbamoyl-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-1-{2-(trimethylsilyl)ethoxy}methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl}phenyl)methylcarbamate	MS m/z 883 [M+H] ⁺ Using benzyl (2-[(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-{2-(trimethylsilyl)ethoxy}methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl}phenyl)methylcarbamate (Preparation 92).
24	6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(5-methoxy-2-[methyl(phenylsulfonyl)amino]benzyl)amino)-1-{2-(trimethylsilyl)ethoxy}methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 803 [M+H] ⁺ Using N-(2-[(6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-{2-(trimethylsilyl)ethoxy}methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl]-5-methoxyphenyl)-N-methylmethanesulfonamide (Preparation 108).

25	N-ethyl-N-(2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 933 [M+H] ⁺ Using N-ethyl-N-(2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide (Preparation 90).
26	6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-4-([2-(methyl(methylsulfonyl)amino)benzyl]amino)-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 669 [M+H] ⁺ Using N-[2-((6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl]-N-methylmethanesulfonamide (Preparation 109).

Preparation 27

6-(4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-methylmethyl-sulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

5

To a solution of N-(2-((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methyl-methanesulfonamide (**Preparation 60**, 5.8 g, 7.04 mmol) in THF (15 mL) was added molybdenum hexacarbonyl (1.872 g, 7.04 mmol), DBU (3.15 mL) and Pd(OAc)₂ (111 mg, 0.15 mmol), and t-butyl amine (6 mL). The reaction was heated in a sealed tube to 100 °C for 45 minutes. The reaction was cooled, filtered and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 29% EtOAc in hexanes to afford the title compound (4 g, 71%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.45 (s, 9H), 1.67-1.73 (m, 2H), 1.91-2.02 (m, 2H), 2.44 (m, 2H), 3.05 (s, 3H), 3.10 (s, 3H), 3.66-3.95 (m, 4H), 4.78-4.86 (m, 2H), 5.21 (s, 2H), 5.86 (m, 1H), 7.04 (s, 1H), 7.27-7.50 (m, 11H), 7.73 (s, 1H), 9.66 (t, 1H). MS m/z 797 [M+H]⁺

15

The following Preparations (**Preparations 28 – 38**) were prepared according to the method described for **Preparation 27** using the appropriate iodo intermediate as described below:

20

Preparation Number	Name	Data
28	6-(4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl-N-(tert-butyl)-4-((5-methyl-2-(N-methylmethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 811 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-4-yl)amino)methyl)-4-methylphenyl)-N-methylmethanesulfonamide (Preparation 94).
29	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((5-methoxy-2-(N-methylmethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 826 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (Preparation 103).
30	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((5-chloro-2-(N-methylmethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 831 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-4-yl)amino)methyl)-4-chlorophenyl)-N-methylmethanesulfonamide (Preparation 101).
31	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((2-(N-ethylmethylsulfonamido)-5-methylbenzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 825 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-4-yl)amino)methyl)-4-methylphenyl)-N-ethylmethanesulfonamide (Preparation 104).
32	6-(4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl-N-(tert-butyl)-4-((2-(N-ethylethylsulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 825 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-4-yl)amino)methyl)phenyl)-N-ethylethanesulfonamide (Preparation 95).

33	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((2-(N-ethylmethanesulfonamido)-5-methylpyridin-3-yl)methyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 826 [M+H] ⁺ Using N-(3-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-5-methylpyridin-2-yl)-N-ethylmethanesulfonamide (Preparation 100).
34	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((2-(N-ethylmethanesulfonamido)-5-fluorobenzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 829 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-4-fluorophenyl)-N-ethylmethanesulfonamide (Preparation 96).
35	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((2-(N-ethylmethanesulfonamido)-5-chlorobenzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 845 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-4-chlorophenyl)-N-ethylmethanesulfonamide (Preparation 97).
36	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((2-(N-ethylmethanesulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 811 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-ethylmethanesulfonamide (Preparation 102).
37	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((2-(N-methylethanesulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 811 [M+H] ⁺ Using N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methylethanesulfonamide (Preparation 98).
38	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((5-methyl-2-(N-methylmethanesulfonamido)pyridin-3-yl)methyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 812 [M+H] ⁺ Using N-(3-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-5-methylpyridin-2-yl)-N-methylmethanesulfonamide (Preparation 99).

Preparation 39

6-[5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-
[methyl(phenylsulfonyl)amino]pyrazin-2-yl)methyl]amino]-1-[2-
(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 4-chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 118**, 150 mg, 0.27 mmol) in n-Butanol (4 mL) was added N-[3-(aminomethyl)pyrazin-2-yl]-N-methylbenzenesulfonamide (**Preparation 219**, 114 mg, 0.41 mmol) and DIPEA (0.17 mL 0.96 mmol). The reaction was heated to 90 °C in a sealed tube for 18 hours. The reaction was quenched by the addition of water and extracted with EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 30-50% EtOAc in hexanes to afford the title compound as a yellow solid (110 mg, 51%). MS m/z 789 [M+H]⁺

The following Preparations (**Preparations 40 – 59**) were prepared according to the method described for **Preparation 39** using the appropriate chloropyridine and the appropriate amine as described below:

Preparation Number	Name	Data
40	4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino]-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 741 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 118) and N-[3-(aminomethyl)pyrazin-2-yl]-N-ethylmethanesulfonamide (Preparation 221).

41	N-ethyl-4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino]-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-[2-(trimethylsilyl)ethoxy]methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 755 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-ethyl-1-[2-(trimethylsilyl)ethoxy]methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 119) and N-[3-(aminomethyl)pyrazin-2-yl]-N-ethylmethanesulfonamide (Preparation 221).
42	6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(4-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1-[2-(trimethylsilyl)ethoxy]methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 726 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-[2-(trimethylsilyl)ethoxy]methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 118) and N-[3-(aminomethyl)pyridin-4-yl]methanesulfonamide (Preparation 197).
43	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)amino]pyridin-2-yl)methyl]amino]-1-[2-(trimethylsilyl)ethoxy]methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 842 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-[2-(trimethylsilyl)ethoxy]methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 122) and N-(2-(aminomethyl)pyridine-3-yl)-N-methylmethane sulphonamide (PCT Intl. Appl. 2008129380).
44	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-4-[(5-methyl-2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1-[2-(trimethylsilyl)ethoxy]methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 856 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-[2-(trimethylsilyl)ethoxy]methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 122) and N-[3-(aminomethyl)-5-methylpyridin-2-yl]-N-methylmethanesulfonamide (Preparation 198).

45	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-4-({2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 904 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 122) and N-[2-(aminomethyl)phenyl]-N-methylpyridine-3-sulfonamide hydrochloride (Preparation 220).
46	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-4-({5-methoxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 871 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 122) and N-[2-(aminomethyl)-4-methoxyphenyl]-N-(methylsulfonyl)methanesulfonamide hydrochloride (Preparation 191).
47	6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({3-[methyl(methylsulfonyl)amino]pyrazin-2-yl}methyl)amino)-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 727 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 118) and N-(3-(aminomethyl)pyrazin-2-yl)-N-methylmethanesulfonamide (PCT Intl. Appl. 2008129380).
48	4-({[2-[ethyl(methylsulfonyl)amino]pyridin-3-yl}methyl]amino)-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 856 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 122) and N-[3-(aminomethyl)pyridin-2-yl]-N-ethylmethanesulfonamide (Preparation 218).

49	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-N-methyl-4-{{2-(sulfamoylmethyl)-benzyl}amino}-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 826 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-N-methyl-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 122) and 1-[2-(aminomethyl)phenyl]methanesulfonamide hydrochloride (Preparation 199).
50	N-tert-butyl-4-{{5-chloro-2-[ethyl(methylsulfonyl)amino]pyridin-3-yl}methyl}amino]-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 932 [M+H] ⁺ Using N-tert-butyl-4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 120) and N-[3-(aminomethyl)-5-chloropyridin-2-yl]-N-ethylmethanesulfonamide hydrochloride (Preparation 201).
51	N-tert-butyl-4-{{5-chloro-2-[methyl(methylsulfonyl)amino]pyridin-3-yl}methyl}amino]-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 918 [M+H] ⁺ Using N-tert-butyl-4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 120) and N-[3-(aminomethyl)-5-chloropyridin-2-yl]-N-methylmethanesulfonamide hydrochloride (Preparation 202).
52	N-(2,4-dimethoxybenzyl)-4-{{2-[ethyl(methylsulfonyl)amino]pyridin-3-yl}methyl}amino]-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 876 [M+H] ⁺ Using 4-chloro-N-(2,4-dimethoxybenzyl)-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 121) and N-[3-(aminomethyl)pyridin-2-yl]-N-ethylmethanesulfonamide (Preparation 218).
53	6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-{{2-[methyl(methylsulfonyl)amino]pyridin-3-yl}methyl}amino]-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 726 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 118) and N-[3-(aminomethyl)pyridin-2-yl]-N-methylmethanesulfonamide (Preparation 217).

54	6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-methoxy-2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-N-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 818 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 118) and N-[2-(aminomethyl)-4-methoxyphenyl]-N-methylpyridine-3-sulfonamide (Preparation 212).
55	6-((4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((5-fluoro-2-(N-methylmethylsulfonamido)pyridin-3-yl)methyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 816 [M+H] ⁺ Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 123) and N-[3-(aminomethyl)-5-fluoropyridin-2-yl]-N-methylmethanesulfonamide hydrochloride (Preparation 203).
56	6-((4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((5-fluoro-2-(N-ethylmethylsulfonamido)pyridin-3-yl)methyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 830 [M+H] ⁺ Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 123) and N-[3-(aminomethyl)-5-fluoropyridin-2-yl]-N-ethylmethanesulfonamide hydrochloride (Preparation 204).
57	6-((4-benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-(((3-(N-ethylmethylsulfonamido)pyrazin-2-yl)methyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 813 [M+H] ⁺ Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 123) and N-[3-(aminomethyl)pyrazin-2-yl]-N-ethylmethanesulfonamide (Preparation 221).
58	6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-N-methyl-4-((2-(N-methyl-2-oxooxazolidine-3-sulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 813 [M+H] ⁺ Using 4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-N-methyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 122) and tert-butyl 2-(N-methyl-2-oxooxazolidine-3-sulfonamido)benzyl-carbamate hydrochloride (Preparation 208).

59	6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-((5-fluoro-2-(N-methylmethanesulfonamido)benzyl)amino)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 814 [M+H] ⁺ Using 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 123) and N-[2-(aminomethyl)-4-fluorophenyl]-N-methylmethanesulfonamide hydrochloride (Preparation 186).
----	---	---

Preparation 60

N-(2-(((6-(4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

To a solution of 6-[4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 112**, 110 mg, 0.18 mmol) in DMF (5 mL) was added 4-nitrophenyl {2-[methyl(methylsulfonyl)amino]benzyl}carbamate (**Preparation 166**, 82 mg, 0.22 mmol) followed by triethylamine (0.06 mL, 0.438 mmol). The reaction was heated to 100 °C for 16 hours. Further 4-nitrophenyl {2-[methyl(methylsulfonyl)amino]benzyl}carbamate (1.2 eq) and triethylamine (2.5 eq) were added and the reaction continued at 100 °C for 18 hours. The reaction was cooled, concentrated *in vacuo* and partitioned between ice-water and EtOAc. The organic layer was collected, washed with saturated aqueous potassium carbonate solution, brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 49% EtOAc in hexanes to afford the title compound (120 mg, 83%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.50-1.75 (m, 3H), 1.89 (m, 2H), 2.32 (m, 1H), 3.05 (s, 3H), 3.06 (s, 3H), 3.53 (m, 1H), 3.70-3.73 (m, 2H), 3.87 (m, 1H), 4.74 (m, 1H), 4.92 (m, 1H), 5.21 (s, 2H), 5.78 (m, 1H), 6.82 (t, 1H), 7.11 (s, 1H), 7.27-7.50 (m, 11H). MS m/z 824 [M+H]⁺

Preparation 61

N-(4-Chloro-2-[[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl)-N-methylmethanesulfonamide

A solution of 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]-methoxy]phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 114**, 650 mg, 0.91 mmol) in DMF was treated with 4-nitrophenyl {5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (**Preparation 156**, 564.76 mg,

1.36 mmol) and triethylamine (0.31 mL, 2.27 mmol) and the reaction was heated to 90 °C for 16 hours. Further 4-nitrophenyl {5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (1.5 eq) and TEA (1.5 eq) were added and the reaction was heated to 90 °C for a further 4 hours. The reaction was cooled and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 30% EtOAc in hexanes to afford the title compound (365 mg, 42%). MS m/z 944 [M³⁵Cl+H]⁺

The following Preparations (**Preparations 62 – 105**) were prepared according to the method described for **Preparation 61** using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 114**) or 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 113**) or 6-[4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 112**) and the appropriate aminocarbamate.

Preparation Number	Name	Data
62	N-ethyl-N-(2-{{6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino}methyl}phenyl)ethanesulfonamide	MS m/z 938 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(ethylsulfonyl)amino]benzyl}carbamate (Preparation 157).
63	N-ethyl-N-[2-{{6-(2-ethyl-5-fluoro-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl}amino}methyl}phenyl)ethanesulfonamide	MS m/z 884 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(ethylsulfonyl)amino]benzyl}carbamate (Preparation 157).

64	N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}meth-yl)phenyl]-N-methylethanesulfonamide	MS m/z 870 [M+H] ⁺ Using 4-nitrophenyl {2-[(ethylsulfonyl)(methyl)amino]benzyl}carbamate (Preparation 159).
65	4-(2-{[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-ethyl)phenol	MS m/z 779 [M+H] ⁺ Using 4-nitrophenyl [2-(4-hydroxyphenyl)ethyl]carbamate (Preparation 158).
66	6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-N-(2-methylpropyl)-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-amine	MS m/z 715 [M+H] ⁺ Using 4-nitrophenyl (2-methylpropyl)carbamate (Preparation 160).
67	N-[4-chloro-2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl]-N-methylmethane sulfonamide	MS m/z 890 [M+H] ⁺ Using 4-nitrophenyl {5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 156).
68	N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)-4-fluorophenyl]-N-methyl-methane sulfonamide	MS m/z 874 [M+H] ⁺ Using 4-nitrophenyl {5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 161).
69	N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-3-fluorophenyl]-N-methylmethane sulfonamide	MS m/z 874 [M+H] ⁺ Using 4-nitrophenyl {2-fluoro-6-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 162).

70	N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl]methanesulfonamide	MS m/z 870 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 163).
71	N-(cyclopentylmethyl)-6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-amine	MS m/z 743 [M ¹²⁹ I+H] ⁺ Using 4-nitrobenzyl (cyclopentylmethyl)carbamate (Preparation 164).
72	6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-N-methyl-4-((5-methyl-2-(N-methylmethylsulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 924 [M+H] ⁺ Using 4-nitrophenyl {5-methyl-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 165).
73	N-ethyl-N-(2-({[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl)methanesulfonamide	MS m/z 924 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 163).
74	N-(2-({[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl)-N-methylethane sulfonamide	MS m/z 924 [M+H] ⁺ Using 4-nitrophenyl {2-[(ethylsulfonyl)(methyl)amino]benzyl}carbamate (Preparation 159).
75	N-(4-fluoro-2-({[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 928 [M+H] ⁺ Using 4-nitrophenyl {5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 161).

76	N-(3-fluoro-2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)-phenyl)-N-methylmethanesulfonamide	MS m/z 928 [M+H] ⁺ Using 4-nitrophenyl {2-fluoro-6-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 162).
77	N-(cyclopentylmethyl)-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-amine	MS m/z 795 [M+H] ⁺ Using 4-nitrobenzyl (cyclopentylmethyl)carbamate (Preparation 164).
78	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-iodo-N-(2-methylpropyl)-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-amine	MS m/z 771 [M ¹²⁹ I+H] ⁺ Using 4-nitrophenyl (2-methylpropyl)carbamate (Preparation 160).
79	N-[2-((6-(2-ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl)-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)phenyl]-N-methylmethanesulfonamide	MS m/z 856 [M+H] ⁺ Using 4-nitrophenyl {2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 166).
80	N-[2-((6-(2-ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl)-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)-4-methylphenyl]-N-methylmethanesulfonamide	MS m/z 870 [M+H] ⁺ Using 4-nitrophenyl {5-methyl-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 165).
81	N-[2-((6-(2-ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl)-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)phenyl]-N-methylbenzene sulfonamide	MS m/z 918 [M+H] ⁺ Using 4-nitrophenyl {2-[methyl(phenylsulfonyl)amino]benzyl}carbamate (Preparation 167).

82	N-[4-(2-{[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-ethyl)phenyl]benzenesulfonamide	MS m/z 918 [M+H] ⁺ 4-nitrophenyl (2-{4-[(phenylsulfonyl)amino]phenyl}ethyl) carbamate (Preparation 168)
83	N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)-4-methoxyphenyl]-N-methylmethanesulfonamide	MS m/z 886 [M+H] ⁺ 4-nitrophenyl {2-[bis(methylsulfonyl)amino]-5-methoxybenzyl}carbamate (Preparation 169).
84	Racemic N-[2-(1-{[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]-methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}ethyl)phenyl]-N-methylmethanesulfonamide	MS m/z 870 [M+H] ⁺ Racemic 4-nitrophenyl (1-{2-[methyl(methylsulfonyl)amino]phenyl}ethyl)carbamate (Preparation 170).
85	N-(2-{[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]-methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)-4-methoxyphenyl)-N-methylmethane-sulfonamide	MS m/z 940 [M+H] ⁺ 4-nitrophenyl {2-[bis(methylsulfonyl)amino]-5-methoxybenzyl}carbamate (Preparation 169).
86	N-(2-{[6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]-methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}methyl)phenyl)-N-methylbenzenesulfonamide	MS m/z 972 [M+H] ⁺ Using 4-nitrophenyl {2-[methyl(phenylsulfonyl)amino]benzyl}carbamate (Preparation 167).

87	N-(2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)-phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]methanesulfonamide	MS m/z 1024 [M+H] ⁺ Using 4-nitrobenzyl (2-((methylsulfonyl)[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]amino)benzyl)carbamate (Preparation 171).
88	N-[2-((6-(2-ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl)-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-5-methoxyphenyl]-N-methylmethanesulfonamide	MS m/z 886 [M+H] ⁺ Using 4-nitrophenyl {4-methoxy-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 172).
89	N-(2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-5-methoxyphenyl)-N-methylmethanesulfonamide	MS m/z 940 [M+H] ⁺ Using 4-nitrophenyl {4-methoxy-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 172).
90	N-ethyl-N-(2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 1016 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(phenylsulfonyl)amino]-5-methoxybenzyl}carbamate (Preparation 176).
91	benzyl [2-((6-(2-ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl)-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl]methylcarbamate	MS m/z 912 [M+H] ⁺ Using benzyl methyl[2-((4-nitrophenoxy)carbonyl)amino)methyl]phenyl]carbamate (Preparation 178).

92	benzyl (2-((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)-phenyl)methylcarbamate	MS m/z 966 [M+H] ⁺ Using benzyl methyl[2-((4-nitrophenoxy)carbonyl)amino]methylphenylcarbamate (Preparation 178).
93	N-{3-[(6-[4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo-[4,3-c]pyridin-4-yl]amino)-methyl}pyridin-2-yl)-N-methylmethanesulfonamide	MS m/z 825 [M+H] ⁺ Using 4-nitrophenyl {2-[methyl(methylsulfonyl)amino]pyridin-3-yl}methylcarbamate (Preparation 177).
94	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo-[4,3-c]pyridine-4-yl)amino)methyl)-4-methylphenyl)-N-methylmethanesulfonamide	MS m/z 838 [M+H] ⁺ Using 4-nitrophenyl {5-methyl-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 165).
95	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo-[4,3-c]pyridine-4-yl)amino)methyl)-phenyl)-N-ethylethanesulfonamide	MS m/z 852 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(ethylsulfonyl)amino]benzyl}carbamate (Preparation 157).
96	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-4-fluorophenyl)-N-ethylmethanesulfonamide	MS m/z 856 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]-5-fluorobenzyl}carbamate (Preparation 179).
97	N-(2-(((6-(4-(benzyloxy)-5-chloro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-4-fluorophenyl)-N-ethylmethanesulfonamide	MS m/z 871 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]-5-chlorobenzyl}carbamate (Preparation 180).
98	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-phenyl)-N-methylethanesulfonamide	MS m/z 838 [M+H] ⁺ Using 4-nitrophenyl {2-[(ethylsulfonyl)(methyl)amino]benzyl}carbamate (Preparation 159).

99	N-(3-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-5-methylpyridin-2-yl)-N-methylmethanesulfonamide	MS m/z 839 [M+H] ⁺ Using 4-nitrobenzyl ((5-methyl-2-(N-methylmethanesulfonamido)pyridin-3-yl)methyl)carbamate (Preparation 181).
100	N-(3-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)-5-methylpyridin-2-yl)-N-ethylmethanesulfonamide	MS m/z 853 [M+H] ⁺ Using 4-nitrophenyl ((2-(N-ethylmethanesulfonamido)-5-methylpyridin-3-yl)methyl)carbamate (Preparation 182).
101	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-4-chlorophenyl)-N-methylmethanesulfonamide	MS m/z 858 [M+H] ⁺ Using 4-nitrophenyl {5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 156).
102	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)-phenyl)-N-ethylmethanesulfonamide	MS m/z 838 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 163).
103	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide	MS m/z 854 [M+H] ⁺ Using 4-nitrophenyl {2-[bis(methylsulfonyl)amino]-5-methoxybenzyl}carbamate (Preparation 169).
104	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)-4-methylphenyl)-N-ethylmethanesulfonamide	MS m/z 852 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]-5-methylbenzyl}carbamate (Preparation 183).
105	N-methyl-N-(2-(((6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)-methyl)-phenyl)methanesulfonamide	MS m/z 910 [M+H] ⁺ Using 4-nitrophenyl {2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 166).

The following Preparations (**Preparations 106 – 109**) were prepared according to the method described for **Preparation 61** using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-**{**[2-(trimethylsilyl)ethoxy]methyl**}**-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 111**) or 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1-**{**[2-(trimethylsilyl)ethoxy]methyl**}**-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 115**) and the appropriate aminocarbamate.

Preparation Number	Name	Data
106	N-(2- { [(6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1- { [2-(trimethylsilyl)ethoxy]methyl } -1H-pyrazolo[4,3-c]pyridin-4-yl)amino]-methyl}phenyl)-3-methoxy-N-methylbenzenesulfonamide	MS m/z 886 [M+H] ⁺ Using 4-nitrophenyl (2- { [(3-methoxyphenyl)sulfonyl](methyl)amino}benzyl)carbamate (Preparation 173).
107	N-ethyl-N-(2- { [(6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1- { [2-(trimethylsilyl)ethoxy]methyl } -1H-pyrazolo[4,3-c]pyridin-4-yl)amino]-methyl}-4-methoxyphenyl)-methanesulfonamide	MS m/z 838 [M+H] ⁺ Using 4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]-5-methoxybenzyl}carbamate (Preparation 174).
108	N-(2- { [(6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1- { [2-(trimethylsilyl)ethoxy]methyl } -1H-pyrazolo[4,3-c]pyridin-4-yl)amino]-methyl}-4-methoxyphenyl)-N-methylbenzenesulfonamide	MS m/z 886 [M+H] ⁺ Using 4-nitrophenyl {5-methoxy-2-[methyl(phenylsulfonyl)amino]benzyl}carbamate (Preparation 175).
109	N-[2- { [(6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1- { [2-(trimethylsilyl)ethoxy]methyl } -1H-pyrazolo[4,3-c]pyridin-4-yl)amino}methyl}phenyl]-N-methylmethanesulfonamide	MS m/z 752 [M+H] ⁺ Using 4-nitrophenyl{2-[methyl(methylsulfonyl)amino]benzyl}carbamate (Preparation 166).

Preparation 110

6-(2-Cyclopropyl-5-fluoro-4-methoxyphenyl)-N-methyl-4-[(2-[methyl(methylsulfonyl)-amino]pyridin-3-yl)methyl]amino]-1-[(2-(trimethylsilyl)ethoxy)methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

5 The title compound was prepared according to the method described for **Preparation 61** using 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-N-methyl-1-[(2-(trimethylsilyl)ethoxy)methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide 5-oxide (**Preparation 116**) and 4-nitrophenyl ((2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl)carbamate (**Preparation 177**). MS m/z 684 [M+H]⁺

Preparation 111

6-[5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-[(2-(trimethylsilyl)ethoxy)methyl]-1H-pyrazolo[4,3-c]pyridine 5-oxide

To a stirred solution of 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-[(2-(trimethylsilyl)ethoxy)methyl]-1H-pyrazolo[4,3-c]pyridine (**Preparation 131**, 15 5.50 g, 9.45 mmol) in dry DCM (550 mL) at 0°C, was added mCPBA (1.79 g, 10.40 mmol) portionwise followed by stirring at room temperature for 16 hours. The reaction was quenched with saturated aqueous sodium bicarbonate solution and saturated aqueous sodium bisulphite solution, the organic extracts separated, dried and purified by silica gel column chromatography eluting with EtOAc to afford the title compound 20 (3.40 g, 50%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.12 (s, 9H), 0.783 (m, 2H), 3.46-3.75 (m, 4H), 3.92 (s, 3H), 5.77 (m, 2H), 7.25 (d, 1H), 7.36 (d, 1H), 8.13 (s, 1H), 8.63 (s, 1H). MS m/z 598 [M+H]⁺

Preparation 112

6-[4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine 5-oxide

To a stirred solution of 6-[4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 132**, 17.7 g, 29 mmol) in anhydrous DCM (900 mL) at 0 °C was added mCPBA (7.51 g, 43.5 mmol) and the reaction was stirred warming to room temperature for 18 hours. The reaction 30 was quenched by the addition of saturated sodium sulphite solution (600 mL) followed by saturated aqueous sodium bicarbonate solution (600 mL). The organic layer was collected, washed with water (3 x 50 mL), brine (2 x 50 mL), dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 56-80% EtOAc in hexanes to afford the title compound (13

g, 71%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.55-1.67 (m, 3H), 1.98 (m, 2H), 2.33 (m, 1H), 3.51 (m, 1H), 3.68 (m, 2H), 3.88 (m, 1H), 5.28 (s, 2H), 5.90 (m, 1H), 7.30-7.51 (m, 7H), 8.60 (s, 1H), 8.80 (s, 1H).

- 5 The following Preparations (**Preparations 113 – 117**) were prepared according to the method described for **Preparation 111** using the appropriate pyrrolopyridine as described below:

Preparation Number	Name	Data
113	6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)-phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine 5-oxide	MS m/z 598 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm -0.147 (s, 9H), -0.02 (s, 9H), 0.78 (t, 2H), 0.92 (t, 2H), 0.97 (t, 2H), 2.31-2.49 (br m, 2H), 3.55 (br t, 2H), 3.78 (t, 2H), 5.35 (s, 1H), 5.74 (s, 2H), 7.14 (d, 1H), 7.24 (d, 2H), 8.05 (s, 1H), 8.59 (s, 1H). Using 6-(2-ethyl-5-fluoro-4-[[2-(trimethylsilyl)ethoxy]-methoxy]phenyl)-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine (Preparation 144).
114	6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine 5-oxide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm -0.11 (s, 9H), 0.00 (s, 9H), 0.78 (t, 2H), 0.89 (t, 2H), 3.52 (m, 3H), 3.60-3.75 (m, 3H), 5.37 (m, 2H), 5.71 (m, 2H), 7.31 (d, 1H), 7.42 (d, 1H), 8.14 (s, 1H), 8.64 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine (Preparation 135).
115	6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine 5-oxide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm -0.11 (s, 9H), 0.64 (m, 4H), 0.81 (t, 2H), 1.71 (m, 1H), 3.55 (t, 2H), 3.89 (s, 3H), 5.74 (s, 2H), 6.78 (d, 1H), 7.14 (d, 1H), 8.06 (s, 1H), 8.59 (s, 1H). Using 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine (Preparation 136).

116	6-(2-cyclopropyl-5-fluoro-4-methoxy-phenyl)-3-iodo-1-{{2-(trimethyl-silyl)ethoxy}-methyl}-1H-pyrazolo-[4,3-c]pyridine 5-oxide	MS m/z 487 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm -0.10 (s, 9H), 0.64 (m, 4H), 0.83 (t, 2H), 1.74 (m, 1H), 2.83 (d, 3H), 3.58 (t, 2H), 3.89 (s, 3H), 5.80 (s, 2H), 6.77 (d, 1H), 7.13 (d, 1H), 8.14 (s, 1H), 8.64 (m, 1H), 8.95 (s, 1H). Using 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-N-methyl-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 128).
117	6-[5-fluoro-2-(2,2,2-trifluoro-ethyl)-4-{{2-(trimethylsilyl)-ethoxy}-methoxy}phenyl]-N-methyl-1-{{2-(trimethylsilyl)-ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide-5-oxide	MS m/z 645 [M+H] ⁺ Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-N-methyl-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (Preparation 130).

Preparation 118

4-Chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

5 **Step 1**

To a solution of 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-{{2-(trimethylsilyl)ethoxy}methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

(**Preparation 125**, 2 g, 3.90 mmol) in anhydrous DCM (30 mL) was added mCPBA (1.2 g, 4.29 mmol) at 0 °C and the reaction was stirred at room temperature for 18 hours.

10 The reaction was quenched by the addition of saturated aqueous sodium bisulfite and sodium bicarbonate solutions and extracted into DCM. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 15% MeOH in DCM to afford the intermediate N-oxide.

15 **Step 2**

This intermediate was dissolved in DMF (20 mL) and oxalyl chloride (2.43 mL, 28.38 mmol) was added at 0 °C with stirring for 1 hour. The reaction was quenched by the addition of water and extracted into EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel

20 column chromatography eluting with 17% EtOAc in DCM to afford the title compound (400 mg, 26%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.11 (s, 9H), 0.82 (m, 2H), 2.85

(d, 3H), 3.57 (t, 2H), 3.92 (s, 3H), 4.10 (q, 2H), 5.85 (s, 2H), 7.37 (d, 1H), 7.49 (d, 1H), 8.11 (s, 1H), 8.69 (m, 1H). MS m/z 547 [M³⁵Cl+H]⁺

Preparation 119

4-Chloro-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-ethyl-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound was prepared according to the method described for **Preparation 118** using N-ethyl-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 129**).

MS m/z 561 [M+H]⁺

Preparation 120

N-tert-Butyl-4-chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound may be prepared according to the method described for **Preparation 118**, Step 1 using N-tert-butyl-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 127**). The N-oxide intermediate (1.3 g, 1.94 mmol) was dissolved in DCM (150 mL) with triethylamine (0.35 mL, 2.52 mmol) and POCl₃ (0.23 mL, 2.52 mmol) was added at 0°C. The reaction was stirred for 1 hour at 10 °C before quenching with saturated aqueous sodium bicarbonate solution and extraction with DCM. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 15% EtOAc in DCM to afford a yellow oil (530 mg, 39%).
¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.08 (s, 9H), -0.01 (s, 9H), 0.84 (t, 2H), 0.90 (t, 2H), 1.41 (s, 9H), 3.58 (t, 2H), 3.77 (t, 2H), 4.07 (m, 2H), 5.37 (s, 2H), 5.84 (s, 2H), 7.46 (m, 2H), 8.08 (s, 1H), 8.33 (s, 1H). MS m/z 705 [M+H]⁺

Preparation 121

4-Chloro-N-(2,4-dimethoxybenzyl)-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound may be prepared according to the method described for **Preparation 118** using N-(2,4-dimethoxybenzyl)-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 126**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.10 (s, 9H),

0.85 (t, 2H), 3.58 (t, 2H), 3.75 (s, 3H), 3.81 (s, 3H), 3.92 (s, 3H), 4.09 (q, 2H), 4.43 (m, 2H), 5.85 (s, 2H), 6.50 (m, 1H), 6.58 (s, 1H), 7.24 (d, 1H), 7.35 (d, 1H), 7.49 (d, 1H), 8.11 (s, 1H), 8.97 (m, 1H). MS m/z 683 [M+H]⁺

Preparation 122

5 4-Chloro-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound may be prepared according to the method described for
Preparation 118 using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide (**Preparation 130**). The N-oxide intermediate (800 mg, 1.24 mmol) was dissolved in DCM (7 mL) and a solution of POCl₃ (0.148 mL, 1.6 mmol) in DCM (3 mL) was added dropwise at 0°C. The reaction was stirred for 30 minutes before the addition of water and extraction into DCM. The organic layer was
 10 collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 14% EtOAc in DCM to afford the title compound as a yellow solid (600 mg, 73%).

¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.11 (s, 9H), -0.01 (s, 9H), 0.82 (t, 2H), 0.90 (t, 3H), 2.85 (d, 3H), 3.57 (t, 2H), 3.77 (t, 2H), 5.37 (s, 2H), 5.85 (s, 2H), 7.43-7.51
 20 (m, 2H), 8.11 (s, 1H), 8.67 (m, 1H). MS m/z 629 [M+H]⁺

Preparation 123

6-(4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide
 25 (**Preparation 124**, 3.80 g, 6.5 mmol) in anhydrous DCM (250 mL) was added mCPBA (1.68 g, 9.75 mol) at 0 °C and the reaction was stirred at room temperature for 18 hours. The reaction was quenched by the addition of saturated aqueous sodium sulphite solution and saturated aqueous sodium bicarbonate solution and extracted into EtOAc.
 30 The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with EtOAc to afford the intermediate N-oxide, that was dissolved in DCM (300 mL). To the solution was added triethylamine (1.07 mL, 7.74 mmol) followed by POCl₃ (0.62 mL, 6.71 mmol) at 0°C. The reaction was stirred at 10 °C for 1 hour before the addition of ice-water. The

reaction was extracted into DCM, the organic layer was collected, dried over sodium sulfate and concentrated *in vacuo* to afford the title compound as the desired chloro isomer confirmed by nOe irradiation of the remaining pyridyl proton.

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.41 (s, 9H), 1.56-1.80 (m, 3H), 1.95-2.04 (m, 2H), 2.37 (m, 1H), 3.76 (m, 1H), 3.91-4.18 (m, 3H), 5.27 (s, 2H), 6.01 (m, 1H), 7.34-7.55 (m, 7H), 8.03 (s, 1H), 8.38 (br s, 1H). MS m/z 619 [M+H]⁺

Preparation 124

6-(4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-N-(tert-butyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 6-[4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 132**, 8 g, 13 mmol) in THF (30 mL) and tert-butylamine (16 mL) was added molybdenum hexacarbonyl (3.48 g, 13 mmol), DBU (5.86 mL, 39.25 mmol) and Pd(OAc)₂ (180 mg, 1.3 mmol). The reaction was heated in a sealed tube at 100 °C for 1 hour. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography to afford the title compound. ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.46 (s, 9H), 1.60-1.74 (m, 4H), 1.97-2.01 (m, 2H), 3.76-3.82 (m, 1H), 3.94-4.22 (m, 3H), 5.27 (s, 2H), 6.03 (m, 1H), 7.36-7.52 (m, 7H), 7.60 (m, 1H), 8.01 (m, 1H), 9.41 (s, 1H). MS m/z 585 [M+H]⁺

Preparation 125

6-[5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

To a solution of 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 131**, 3.4 g, 5.85 mmol) and 2M methylamine in THF (30 mL) was added palladium acetate (92 mg, 0.41 mmol), DBU (2.62 mL, 17.54 mmol) and molybdenum hexacarbonyl (1.55 g, 5.85 mmol). The reaction was heated in a sealed tube at 100 °C for 60 minutes before concentrating *in vacuo*. The residue was diluted with EtOAc, filtered through celite and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 30% EtOAc in hexanes to afford the title compound as a yellow solid (2 g, 67%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.15 (s, 9H), 0.80 (t, 2H), 2.85 (d, 3H), 3.56 (t, 2H), 3.92 (s, 3H), 4.18 (q, 2H), 5.87 (s, 2H), 7.33 (d, 1H), 7.41 (d, 1H), 8.07 (s, 1H), 8.66 (m, 1H), 9.45 (s, 1H). MS m/z 513 [M+H]⁺

The following Preparations (**Preparations 126 – 129**) were prepared according to the method described for **Preparation 125** using the appropriate pyrrolopyridine and amine as described below:

Preparation Number	Name	Data
126	N-(2,4-dimethoxybenzyl)-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 649 [M+H] ⁺ Using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridine (Preparation 131) and 2,4-dimethoxybenzylamine.
127	N-tert-butyl-6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy)methoxy}phenyl]-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	MS m/z 671 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm -0.13 (s, 9H), -0.01 (s, 9H), 0.81 (t, 2H), 0.91 (t, 2H), 1.45 (s, 9H), 3.58 (t, 2H), 3.80 (t, 2H), 4.03 (m, 2H), 5.36 (s, 2H), 6.02 (s, 2H), 7.43 (m, 2H), 7.66 (s, 1H), 8.08 (s, 1H). Using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{{2-(trimethylsilyl)ethoxy)methoxy}phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridine (Preparation 135) and <i>tert</i> -butylamine.
128	6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-N-methyl-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	Taken on directly to the next step. Using 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridine (Preparation 136) and methylamine.
129	N-ethyl-6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm -0.14 (s, 9H), 0.80 (t, 2H), 1.17 (t, 3H), 3.37 (m, 2H), 3.56 (t, 2H), 3.89 (s, 3H), 4.18 (m, 2H), 5.87 (s, 2H), 7.33 (d, 1H), 7.44 (d, 1H), 8.08 (s, 1H), 8.72 (m, 1H), 9.45 (s, 1H). Using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-{{2-(trimethylsilyl)ethoxy)methyl}-1H-pyrazolo[4,3-c]pyridine (Preparation 131) with ethylamine.

Preparation 130

6-[5-Fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-N-methyl-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxamide

The title compound may be prepared according to the method described for
 5 **Preparation 18** using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]-methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine (**Preparation 135**) with methylamine. ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.15 (s, 9H), -0.01 (s, 9H), 0.80 (t, 2H), 0.91 (t, 2H), 2.85 (d, 3H), 3.58 (t, 2H), 3.78 (t, 2H), 4.10 (q, 2H), 5.36 (s, 2H), 5.87 (s, 2H), 7.43 (m, 2H), 8.08 (s, 1H), 8.64 (m, 1H), 9.46 (s, 1H).
 10 MS m/z 629 [M+H]⁺

Preparation 131

6-[5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine

To a suspension of NaH (0.59 g, 24.93 mmol) in dry DMF (100 mL) at 0 °C was
 15 added a solution of 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1H-pyrazolo[4,3-c]pyridine (**Preparation 137**, 7.50 g, 16.62 mmol) in DMF (100 mL). The reaction was stirred for 30 minutes before the addition of SEM-chloride (4.42 mL, 24.93 mmol) drop-wise. The reaction was stirred for 1 hour before being quenched with ice-water and extracted into EtOAc. The organic layer was washed with water, brine, dried
 20 over sodium sulfate and concentrated *in vacuo*. The crude residue was purified using silica gel column chromatography to afford the title compound as a yellow liquid (5.50 g, 47%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.15 (s, 9H), 0.76 (m, 2H), 3.51 (m, 2H), 3.91 (s, 3H), 4.11 (q, 2H), 5.81 (s, 2H), 7.33 (d, 1H), 7.42 (d, 1H), 7.99 (s, 1H), 8.85 (s, 1H). MS m/z 582 [M+H]⁺

Preparation 132

6-[4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine

To a solution of 2-fluoro-4-[3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]-5-(2,2,2-trifluoroethyl)phenol (**Preparation 133**, 21.8 g, 41.8 mmol) in
 30 acetone (200 mL) was treated with benzyl bromide (7.5 mL, 62.7 mmol) and potassium carbonate (14.4 g, 104 mmol) and the reaction was heated to reflux. The reaction was cooled, filtered, the filtrate collected and concentrated *in vacuo*. The residue was taken up in EtOAc, washed with water, brine, dried over sodium sulfate and concentrated *in*

vacuo. The residue was purified using silica gel column chromatography eluting with 18% EtOAc in hexane to afford the title compound (23 g, 90%).

¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.57-1.71 (m, 3H), 2.01 (m, 2H), 2.37 (m, 1H), 3.74 (m, 1H), 3.91 (m, 1H), 4.03 (m, 1H), 4.16 (m, 1H), 5.27 (s, 2H), 5.97 (m, 1H),
 5 7.34-7.51 (m, 7H), 7.95 (s, 1H), 8.83 (s, 1H). MS m/z 612 [M+H]⁺

Preparation 133

2-Fluoro-4-[3-iodo-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]-5-(2,2,2-trifluoroethyl)phenol

To a solution of 2-fluoro-4-(3-iodo-1H-pyrazolo[4,3-c]pyridin-6-yl)-5-(2,2,2-trifluoroethyl)phenol (**Preparation 134**, 40.8 g, 93 mmol) in DMF (500 mL) was added dihydropyran (17 mL, 187 mmol) and PTSA (7.10 g, 37 mmol) and the reaction was heated to 80 °C for 18 hours. Additional dihydropyran (2 eq) and PTSA (0.4 eq) were added and the reaction continued at this temperature for a further 2 hours followed by cooling to room temperature for 18 hours. The reaction was quenched by the addition of
 15 saturated aqueous sodium bicarbonate solution dropwise, and concentrated *in vacuo*. The aqueous residue was extracted into EtOAc, washed with water, brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 17-20% EtOAc in hexanes to afford the title compound as a yellow solid (22 g, 45%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.44-
 20 1.70 (m, 3H), 2.02 (m, 2H), 2.37 (m, 1H), 3.75 (m, 1H), 3.88-4.06 (m, 2H), 4.15 (m, 1H), 5.97 (m, 1H), 7.10 (d, 1H), 7.40 (d, 1H), 7.91 (s, 1H), 8.81 (s, 1H), 10.31 (s, 1H). MS m/z 522 [M+H]⁺

Preparation 134

2-Fluoro-4-(3-iodo-1H-pyrazolo[4,3-c]pyridin-6-yl)-5-(2,2,2-trifluoroethyl)phenol

A solution of 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1H-pyrazolo[4,3-c]pyridine (**Preparation 137**, 44 g, 97 mmol) in DCM (350 mL) was treated with boron tribromide (46 mL, 488 mmol) at 0°C, and the reaction was allowed to stir warming to room temperature over 5 hours. The reaction was concentrated *in vacuo* and treated with saturated aqueous sodium bicarbonate solution. The resulting
 30 precipitate was filtered and dried under vacuum to afford the title compound as a white solid (41 g, 97%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 4.00 (q, 2H), 7.09 (d, 1H), 7.37 (d, 1H), 7.58 (s, 1H), 8.81 (s, 1H), 10.26 (s, 1H), 13.95 (s, 1H). MS m/z 438 [M+H]⁺

Preparation 135

6-[5-Fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine

The title compound was prepared according to the method described for
 5 **Preparation 131** using 6-[4-{[tert-butyl(dimethyl)silyl]oxy}-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1H-pyrazolo[4,3-c]pyridine (**Preparation 138**). The reaction conditions cause deprotection of the TBDMS ether and subsequent re-protection with SEM-chloride. ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.14 (s, 9H), -0.03 (s, 9H), 0.78 (t, 2H), 0.90 (t, 2H), 3.56 (t, 2H), 3.77 (t, 2H), 4.09 (m, 2H), 5.36 (s, 2H), 5.81 (s, 2H),
 10 7.42 (m, 2H), 8.01 (s, 1H), 8.86 (s, 1H). MS m/z 698 [M+H]⁺

Preparation 136

6-(2-Cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine

15 The title compound was prepared according to the method described for **Preparation 131** using 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1H-pyrazolo[4,3-c]pyridine (**Preparation 139**). Taken on to the next step without further purification.

Preparation 137

20 6-[5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1H-pyrazolo[4,3-c]pyridine

To a solution of 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine (**Preparation 140**, 11.20 g, 34.43 mmol) in DMF (200 mL) at 0 °C was added N-iodosuccinimide (9.29 g, 41.32 mmol). The reaction was stirred at
 25 room temperature for 16 hours. The reaction was diluted with ethyl acetate, washed with saturated aqueous sodium bicarbonate solution and saturated aqueous sodium thiosulfate solution. The organic extracts were separated, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 22% EtOAc in hexanes to afford the title compound as a
 30 white solid (7.50 g, 48%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.91 (s, 3H), 4.11 (q, 2H), 7.32 (d, 1H), 7.46 (d, 1H), 7.62 (s, 1H), 8.83 (s, 1H), 13.90 (br s, 1H). MS m/z 452 [M+H]⁺

Preparation 138**6-[4-{{tert-Butyl(dimethyl)silyl}oxy}-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1H-pyrazolo[4,3-c]pyridine**

The title compound was prepared according to the method described for
 5 **Preparation 137** using 6-[4-{{tert-butyl(dimethyl)silyl}oxy}-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo-[4,3-c]pyridine (**Preparation 141**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.24 (s, 6H), 0.99 (s, 9H), 4.08 (m, 2H), 7.16 (d, 1H), 7.46 (d, 1H), 7.65 (s, 1H), 8.83 (s, 1H), 13.99 (br s, 1H). MS m/z 552 [M+H]⁺

Preparation 139**6-(2-Cyclopropyl-5-fluoro-4-methoxyphenyl)-3-iodo-1H-pyrazolo[4,3-c]pyridine**

The title compound was prepared according to the method described for
Preparation 137 using 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 143**). The residue was triturated with pentane and ether. ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.69 (m, 2H), 0.80 (m, 2H), 2.09 (m, 1H), 3.88 (s, 3H), 6.73 (d, 1H), 7.30 (d, 1H), 7.65 (s, 1H), 8.83 (s, 1H), 13.88 (s, 1H).
 15

Preparation 140**6-[5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine**

To a solution of 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 145**, 14.50 g, 35.41
 20 mmol) in dioxane (150 mL) was added 4M HCl in dioxane (60 mL). The reaction was stirred at room temperature for 16 hours before concentrating *in vacuo*. The residue was partitioned between EtOAc and saturated aqueous sodium bicarbonate solution. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo* to afford the title compound that was used directly in the next
 25 reaction (11.20 g, 97%). MS m/z 326 [M+H]⁺

Preparation 141**6-[4-{{tert-Butyl(dimethyl)silyl}oxy}-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine**

To a solution of 2-fluoro-4-(1H-pyrazolo[4,3-c]sulfate-6-yl)-5-(2,2,2-trifluoroethyl)phenol (**Preparation 142**, 13 g, 41.76 mmol) and 2,6 lutidine (7.29 mL, 62.65 mmol) in anhydrous THF (500 mL) at 0 °C was added TBDMS-triflate (11.52 mL, 50.12 mmol) and the reaction was stirred at room temperature for 18 hours. The reaction was concentrated *in vacuo* and partitioned between water and ethyl acetate. The organic layer was collected, washed with brine, dried over sodium sulfate and

concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 20% EtOAc in hexanes to afford the title compound as a white solid (11 g, 62%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.24 (s, 6H), 0.99 (s, 9H), 4.09 (m, 2H), 7.17 (d, 1H), 7.45 (d, 1H), 7.63 (s, 1H), 8.33 (s, 1H), 9.15 (s, 1H), 13.57 (br s, 1H).

5

Preparation 142

2-Fluoro-4-(1H-pyrazolo[4,3-c]pyridin-6-yl)-5-(2,2,2-trifluoroethyl)phenol

6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 146**, 24 g, 45.66 mmol) was dissolved in TFA (48 mL) at 0 °C and stirred for 1 hour. The reaction was concentrated *in vacuo* and taken up in MeOH. Ethylene diamine (2.4 mL) was added at 0 °C and the reaction stirred for 20 minutes. The reaction was concentrated *in vacuo* and partitioned between IPA:DCM (1:9) and water. The organic extract was dried with sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography to afford the title compound as an off-white solid (13 g, 91%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.98 (m, 2H), 7.08 (d, 1H), 7.33 (d, 1H), 7.57 (s, 1H), 8.32 (s, 1H), 9.14 (s, 1H), 10.21 (br s, 1H), 13.52 (br s, 1H).

15

Preparation 143

6-(2-Cyclopropyl-5-fluoro-4-methoxyphenyl)-1H-pyrazolo[4,3-c]pyridine

To a solution of 6-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 148**, 5 g, 13.61 mmol) in MeOH (25 mL) was added concentrated HCl (3.5 mL) at 0 °C and the reaction was stirred for 16 hours. The reaction was concentrated *in vacuo* and partitioned between saturated aqueous NaHCO₃ solution and 25% IPA in DCM. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo* to afford the title compound as a white solid (3.3 g, 85%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.71 (m, 2H), 0.81 (m, 2H), 2.12 (m, 1H), 3.88 (s, 3H), 6.71 (d, 1H), 7.30 (d, 1H), 7.66 (s, 1H), 8.32 (s, 1H), 9.16 (s, 1H), 13.46 (s, 1H). MS m/z 284 [M+H]⁺

25

Preparation 144

6-(2-Ethyl-5-fluoro-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl)-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine

30

The title compound was prepared according to the methods described by **Preparations 142, 141, 137 and 131** using 6-[2-ethyl-5-fluoro-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 147**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.13 (s, 9H), -0.01

(s, 9H), 0.79 (t, 2H), 0.91 (t, 2H), 0.99 (t, 2H), 2.66 (q, 2H), 3.55 (t, 2H), 3.78 (t, 2H), 5.35 (s, 2H), 5.80 (s, 2H), 7.22 (m, 2H), 7.88 (s, 1H), 8.84 (s, 1H).

Preparation 145

6-[5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine

A solution of palladium acetate (0.47 g, 2.10 mmol) and S-Phos (0.86 g, 2.10 mmol) in ethanol (75 mL) was heated at 50 °C for 45 minutes after purging with nitrogen (Solution A). Meanwhile a solution of 6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 149**, 10g, 42.07 mmol) in ethanol (75 mL) was treated with 2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**PCT Intl. Appl. 2013014567**, 21.08 g, 63.10 mmol) and an aqueous solution of potassium phosphate (17.86 g, 84.14 mmol) in water (50 mL) followed by purging with nitrogen for 10 minutes (Solution B). Solution A was added to Solution B and the reaction heated to 80 °C for 18 hours before cooling and concentrating *in vacuo*. The residue was partitioned between ethyl acetate and water, the organic extracts collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 15% EtOAc in hexanes to afford the title compound as a yellow liquid (14.10 g, 82%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.58 (m, 2H), 1.73 (m, 1H), 1.99 (m, 2H), 2.42 (m, 1H), 3.78 (m, 1H), 3.92 (m, 4H), 4.09 (m, 1H), 4.26 (m, 1H), 5.98 (d, 1H), 7.33 (d, 1H), 7.48 (d, 1H), 7.92 (s, 1H), 8.37 (s, 1H), 9.15 (s, 1H). MS m/z 410 [M+H]⁺

Preparation 146

6-[5-Fluoro-2-(2,2,2-trifluoroethyl)-4-{2-(trimethylsilyl)ethoxy}methoxy}phenyl]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine

The title compound was prepared according to the method described for **Preparation 145** using 6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 149**) and (2-{[2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-(2,2,2-trifluoroethyl)phenoxy] methoxy}ethyl)(trimethyl)silane (**Preparation 150**). The residue was purified using silica gel column chromatography eluting with 9% EtOAc in hexanes (19 g, 86%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.02 (s, 9H), 0.91 (m, 2H), 1.58 (m, 2H), 1.73 (m, 1H), 2.05 (m, 2H), 2.41 (m, 1H), 3.73 (m, 2H), 3.92-4.21 (m, 2H), 5.35 (s, 2H), 5.98 (m, 1H), 7.40 (d, 1H), 7.50 (d, 1H), 7.93 (s, 1H), 8.38 (s, 1H), 9.16 (s, 1H).

Preparation 147**6-[2-Ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine**

The title compound was prepared according the method described for
 5 **Preparation 145** using (2-{[2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-ethyl-phenoxy] methoxy}ethyl)(trimethyl)silane (**PCT Intl. Appl. WO2013014567A1**) and 6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 149**). The residue was purified using silica gel column chromatography eluting with 30% EtOAc in hexanes (34 g, 85%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.01 (s, 9H),
 10 0.92 (m, 2H), 1.05 (m, 3H), 1.58 (m, 2H), 1.73 (m, 1H), 2.03 (m, 2H), 2.41 (m, 1H), 2.65 (m, 2H), 3.72-3.92 (m, 4H), 5.34 (s, 2H), 5.93 (m, 1H), 7.21-7.28 (m, 2H), 7.80 (s, 1H), 8.36 (s, 1H), 9.14 (s, 1H). MS m/z 526 [M+H]⁺

Preparation 148**6-(2-Cyclopropyl-5-fluoro-4-methoxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine**

The title compound was prepared according the method described for
Preparation 145 using 2-(2-cyclopropyl-5-fluoro-4-methoxyphenyl)-4,4,5,5-tetramethyl-
 20 1,3,2-dioxaborolane (**Preparation 151**) and 6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 149**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.65-0.85 (m, 4H), 1.58 (m, 2H), 1.71 (m, 1H), 2.01 (m, 2H), 2.11 (m, 1H), 2.40 (m, 1H), 3.31 (s, 3H), 3.74 (m, 1H), 3.90 (m, 1H), 5.94 (m, 1H), 6.74 (d, 1H), 7.31 (d, 1H), 7.91 (s, 1H), 8.35 (s, 1H), 9.15 (s, 1H). MS m/z 368 [M+H]⁺

Preparation 149**6-Chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine**

To a solution of 6-chloro-1H-pyrazolo[4,3-c]pyridine (75 g, 488.37 mmol) in DCM (2 L) was added dihydropyran (66.98 mL, 732.56 mmol) followed by *para*-toluenesulfonic acid (18.58 g, 97.67 mmol) and the reaction was heated to reflux for 18
 30 hours. Further *para*-toluenesulfonic acid (0.1 eq) and dihydropyran (0.75 eq) were added and the reaction continued heating at reflux for 6 hours. The reaction was cooled and quenched with saturated aqueous sodium bicarbonate solution. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 17%

EtOAc in hexanes followed by trituration with ether to afford the title compound as a pale yellow solid (83 g, 72%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.59 (m, 2H), 1.71 (m, 1H), 2.02 (m, 2H), 2.29 (m, 1H), 3.74 (m, 1H), 3.89 (m, 1H), 5.91 (m, 1H), 7.93 (s, 1H), 8.38 (s, 1H), 8.94 (s, 1H).

5

Preparation 150

(2-{[2-Fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-(2,2,2-trifluoroethyl)phenoxy]methoxy}ethyl)(trimethyl)silane

To a solution of (2-{[4-bromo-2-fluoro-5-(2,2,2-trifluoroethyl)phenoxy]methoxy}ethyl)-(trimethyl)silane (**Preparation 152**, 34 g, 84.31 mmol), in dry 1,4-dioxane (1 L) was added bis(pinacolato)diboron (21.41 g, 84.31 mmol) followed by KOAc (24.82 g, 252.95 mmol). The reaction mixture was purged with nitrogen for 20 minutes before the addition of Pd(dppf)₂Cl₂ (6.886g, 8.432mmol) followed by further degassing for 20 minutes. The reaction was heated to reflux for 18 hours before cooling to room temperature and concentrating *in vacuo*. The residue was suspended in EtOAc and filtered through a bed of Celite. The filtrate was washed with water, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 10% EtOAc in hexanes to afford the title compound as an oil (31 g, 82%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.05 (s, 9H), 0.87 (m, 2H), 1.32 (s, 12H), 3.74 (m, 2H), 3.92 (m, 2H), 5.32 (s, 2H), 7.29 (d, 1H), 7.42 (d, 1H).

20

Preparation 151

2-(2-Cyclopropyl-5-fluoro-4-methoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

The title compound was prepared according to the method described for **Preparation 150** using 1-bromo-2-cyclopropyl-5-fluoro-4-methoxybenzene (**Preparation 154**). Taken on to the next step as is.

25

Preparation 152

(2-{[4-Bromo-2-fluoro-5-(2,2,2-trifluoroethyl)phenoxy]methoxy}ethyl)(trimethyl)silane

To a solution of 4-bromo-2-fluoro-5-(2,2,2-trifluoroethyl)phenol (**Preparation 153**, 25 g, 91.57 mmol) in DCM (200 mL) was added DIPEA (17.54 mL, 100.73 mmol) at room temperature followed by SEM-Cl (17.86 mL, 100 mmol) drop-wise at 0 °C and the reaction was stirred at room temperature for 4 hours. The reaction was partitioned between DCM and water, the organic layer was collected, washed with brine, dried over

30

sodium sulfate and concentrated *in vacuo* to afford the title compound as an oil (34 g, 92%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.03 (s, 9H), 0.87 (m, 2H), 3.70-3.79 (m, 4H), 5.29 (s, 2H), 7.42 (d, 1H), 7.70 (d, 1H).

Preparation 153

4-Bromo-2-fluoro-5-(2,2,2-trifluoroethyl)phenol

To a solution of 1-bromo-5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)benzene (**PCT Intl. Appl. 2013014567**, 30 g, 104.51 mmol) in DCM (800 mL) at 0 °C was added boron tribromide (130.91 g, 522 mmol) drop-wise and the reaction was stirred at room temperature for 16 hours. The reaction was quenched by the addition of saturated aqueous NaHCO₃ solution and extracted into DCM. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo* to afford the title compound as a white solid that was taken directly on to the next step (22 g, 77%).

Preparation 154

1-Bromo-2-cyclopropyl-5-fluoro-4-methoxybenzene

To a solution of 4-cyclopropyl-1-fluoro-2-methoxybenzene (**Preparation 155**, 8.7 g, 41 mmol) in DMF (250 mL) at 0 °C was added NBS (7.40 g, 41 mmol) and the reaction was stirred at room temperature for 3 hours. The reaction was partitioned between EtOAc and brine, the organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with hexanes to afford the title compound as a colorless oil (9.5 g, 92%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.73 (m, 2H), 0.96 (m, 2H), 2.01 (m, 1H), 3.82 (s, 3H), 6.72 (d, 1H), 7.50 (d, 1H).

Preparation 155

4-Cyclopropyl-1-fluoro-2-methoxybenzene

To a solution of 5-bromo-2-fluoroanisole (10 g, 48.77 mmol) in toluene (100 mL) was added water (10 mL), cyclopropyl boronic acid (5.44 g, 63 mmol), tricyclohexylphosphine (1.37 g, 4.87 mmol) and potassium phosphate (36.3 g, 170 mmol). The reaction was degassed with nitrogen before the addition of Pd(OAc)₂ (547 mg, 2.44 mmol) followed by heating to 100 °C for 3 hours. The reaction was cooled and partitioned between EtOAc and brine. The organic layer was collected, concentrated *in vacuo* and purified using silica gel column chromatography to afford the title compound as a colorless oil (9.7 g, quant).

¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.65 (m, 2H), 0.92 (m, 2H), 1.88 (m, 1H), 3.81 (s, 3H), 6.61 (m, 1H), 6.82 (m, 1H), 7.06 (m, 1H).

Preparation 156**4-Nitrophenyl {5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate**

To a solution of N-[2-(aminomethyl)-4-chlorophenyl]-N-methylmethanesulfonamide hydrochloride (**Preparation 211**, 3.20 g, 11.39 mmol) and sodium carbonate (3.62 g, 34.18 mmol) in DCM (50 mL) at 0°C, was added 4-nitrophenylchloroformate (2.52 g, 12.53 mmol) and the reaction was stirred at room temperature for 18 hours. The reaction was concentrated *in vacuo* and the residue purified by silica gel column chromatography eluting with 55-100% EtOAc in hexanes to afford the title compound as a pale yellow solid (2.20 g, 46%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.08 (s, 3H), 3.17 (s, 3H), 4.32 (br m, 1H), 4.52 (br m, 1H), 7.40-7.61 (m, 5H), 8.25 (m, 2H).

The following Preparations (**Preparations 157 – 183**) were prepared according to the method described for **Preparation 156** using the appropriate amine as described below, and taken directly on to the next step;

Preparation Number	Name	Data/SM
157	4-nitrophenyl {2-[ethyl(ethylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)phenyl]-N-ethylethanesulfonamide hydrochloride (Preparation 184).
158	4-nitrophenyl [2-(4-hydroxyphenyl)ethyl]carbamate	2-(4-hydroxyphenyl)ethanamine.
159	4-nitrophenyl {2-[(ethylsulfonyl)(methyl)amino]benzyl}carbamate	N-[2-(aminomethyl)phenyl]-N-methylethanesulfonamide hydrochloride (Preparation 185).
160	4-nitrophenyl (2-methylpropyl)carbamate	2-methylpropylamine.
161	4-nitrophenyl {5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)-4-fluorophenyl]-N-methylmethanesulfonamide hydrochloride (Preparation 186).
162	4-nitrophenyl {2-fluoro-6-[methyl(methylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)-3-fluorophenyl]-N-methylmethanesulfonamide hydrochloride (Preparation 187).
163	4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)phenyl]-N-ethylmethanesulfonamide hydrochloride (Preparation 188).
164	4-nitrobenzyl (cyclopentylmethyl)carbamate	1-cyclopentylmethanamine

165	4-nitrophenyl {5-methyl-2-[methyl(methylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)-4-methylphenyl]-N-methylmethanesulfonamide hydrochloride (Preparation 189).
166	4-nitrophenyl {2-[methyl(methylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)phenyl]-N-methylmethanesulfonamide (PCT Intl. Appl. 2010058846).
167	4-nitrophenyl {2-[methyl(phenylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)phenyl]-N-methylbenzenesulfonamide hydrochloride (Preparation 190).
168	4-nitrophenyl {2-{4-[(phenylsulfonyl)amino]phenyl}172ulph}carbamate	N-[4-(2-aminoethyl)phenyl]benzenesulfonamide (Preparation 210).
169	4-nitrophenyl {2-[bis(methylsulfonyl)amino]-5-methoxybenzyl}carbamate	N-[2-(aminomethyl)-4-methoxyphenyl]-N-(methylsulfonyl)methanesulfonamide hydrochloride (Preparation 191).
170	Racemic 4-nitrophenyl {1-{2-[methyl(methylsulfonyl)amino]phenyl}ethyl}carbamate	Racemic N-[2-(1-aminoethyl)phenyl]-N-methylmethanesulfonamide (Preparation 209).
171	4-nitrobenzyl {2-{(methylsulfonyl)[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]amino}benzyl}carbamate	N-[2-(aminomethyl)phenyl]-N-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]methanesulfonamide (Preparation 216).
172	4-nitrophenyl {4-methoxy-2-[methyl(methylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)-5-methoxyphenyl]-N-methylmethanesulfonamide hydrochloride (Preparation 192).
173	4-nitrophenyl {2-[(3-methoxyphenyl)sulfonyl](methyl)amino}benzyl}carbamate	N-[2-(aminomethyl)phenyl]-3-methoxy-N-methylbenzenesulfonamide hydrochloride (Preparation 193).
174	4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]-5-methoxybenzyl}carbamate	N-[2-(aminomethyl)-4-methoxyphenyl]-N-ethylmethanesulfonamide hydrochloride (Preparation 194).
175	4-nitrophenyl {5-methoxy-2-[methyl(phenylsulfonyl)amino]benzyl}carbamate	N-[2-(aminomethyl)-4-methoxyphenyl]-N-methylbenzenesulfonamide hydrochloride (Preparation 195).
176	4-nitrophenyl {2-[ethyl(phenylsulfonyl)amino]-5-methoxybenzyl}carbamate	N-[2-(aminomethyl)-4-methoxyphenyl]-N-ethylbenzenesulfonamide hydrochloride (Preparation 196).
177	4-nitrophenyl {2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl}carbamate	N-[3-(aminomethyl)pyridin-2-yl]-N-methylmethanesulfonamide (Preparation 217).
178	benzyl methyl{2-[(4-nitrophenoxy)carbonyl]amino}methyl}phenyl}carbamate	Benzyl [2-(aminomethyl)phenyl]methylcarbamate hydrochloride (Preparation 249).
179	4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]-5-fluorobenzyl}carbamate	N-[2-(aminomethyl)-4-fluorophenyl]-N-ethylmethanesulfonamide (Preparation 205).

180	4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]-5-chlorobenzyl}carbamate	N-[2-(aminomethyl)-4-chlorophenyl]-N-ethylmethanesulfonamide (Preparation 206).
181	4-nitrobenzyl ((5-methyl-2-(N-methylmethylsulfonamido)pyridin-3-yl)methyl)carbamate	N-[3-(aminomethyl)-5-methylpyridin-2-yl]-N-methylmethanesulfonamide hydrochloride (Preparation 198).
182	4-nitrophenyl ((2-(N-ethylmethylsulfonamido)-5-methylpyridin-3-yl)methyl)carbamate	N-[3-(aminomethyl)-5-methylpyridin-2-yl]-N-ethylmethanesulfonamide hydrochloride (Preparation 207).
183	4-nitrophenyl {2-[ethyl(methylsulfonyl)amino]-5-methylbenzyl}carbamate	N-[2-(aminomethyl)-4-methylphenyl]-N-ethylmethanesulfonamide hydrochloride (Preparation 200).

The following Preparations (**Preparations 184 – 207**) were prepared according to the methods described in the three steps below:

- 1) **Preparation 236**
- 2) **Preparation 213**
- 3) **Preparation 211**

using the appropriate alkyl halide as described below:

Preparation Number	Name	Data/SM
184	N-[2-(aminomethyl)-phenyl]-N-ethylethanesulfonamide hydrochloride	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.06 (s, 3H), 3.22 (s, 3H), 4.09 (b s, 1H), 4.21 (b s, 1H), 7.47-7.52 (m, 2H), 7.62-7.63 (m, 2H), 8.23 (b s, 1H). Using N-(2-cyanophenyl)-N-(ethylsulfonyl)ethanesulfonamide (Preparation 241) and ethyl iodide.
185	N-[2-(aminomethyl)phenyl]-N-methylethanesulfonamide hydrochloride	MS m/z 229 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.26 (t, 3H), 3.06-3.56 (br m, 4H), 4.03-4.19 (br m, 2H), 7.42-7.49 (m, 2H), 7.60 (m, 1H), 7.68 (m, 1H), 8.48 (br s, 3H). Using N-(2-cyanophenyl)-N-(ethylsulfonyl)ethanesulfonamide (Preparation 241) and methyl iodide.

186	N-[2-(aminomethyl)-4-fluorophenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 233 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.06 (s, 3H), 3.56 (s, 3H), 4.02 (br s, 1H), 4.20 (br s, 1H), 7.32-7.37 (m, 1H), 7.56-7.59 (m, 1H), 7.68-7.71 (m, 1H), 8.49 (br s, 2H). Using N-(4-fluoro-2-cyanophenyl)-N-(methylsulfonyl)-methanesulfonamide (Preparation 239) and methyl iodide.
187	N-[2-(aminomethyl)-3-fluorophenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 233 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.06 (s, 3H), 3.27 (s, 3H), 4.05 (br s, 1H), 4.20 (br s, 1H), 7.36-7.41 (t, 1H), 7.52-7.62 (m, 2H), 8.38 (br s, 2H). Using N-(3-fluoro-2-cyanophenyl)-N-(methylsulfonyl)-methanesulfonamide (Preparation 240) and methyl iodide.
188	N-[2-(aminomethyl)phenyl]-N-ethylmethanesulfonamide hydrochloride	MS m/z 229 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.26 (t, 3H), 3.13 (m, 1H), 3.24 (s, 3H), 3.39 (m, 1H), 4.03 (br s, 1H), 4.19 (br s, 1H), 7.44-7.49 (m, 2H), 7.59-7.70 (m, 2H), 8.48 (br s, 2H). Using N-(2-cyanophenyl)-N-(methylsulfonyl)-methanesulfonamide (Preparation 238) and ethyl iodide.
189	N-[2-(aminomethyl)-4-methylphenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 229 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.34 (s, 3H), 3.03 (s, 3H), 3.19 (s, 3H), 3.98 (br s, 1H), 4.17 (br s, 1H), 7.29-7.31 (d, 1H), 7.47-7.51 (m, 2H), 8.33 (br s, 2H). Using N-(4-methyl-2-cyanophenyl)-N-(methylsulfonyl)-methanesulfonamide (Preparation 242) and methyl iodide.
190	N-[2-(aminomethyl)phenyl]-N-methylbenzenesulfonamide hydrochloride	MS m/z 277 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.10 (s, 3H), 4.20 (br m, 1H), 4.40 (br m, 1H), 6.51 (m, 1H), 7.16 (m, 1H), 7.33 (m, 3H), 7.64 (m, 4H), 7.77 (m, 1H). Using N-(2-cyanophenyl)-N-(phenylsulfonyl)-benzenesulfonamide (Preparation 243) and methyl iodide.
191	N-[2-(aminomethyl)-4-methoxyphenyl]-N-(methylsulfonyl)-methanesulfonamide hydrochloride	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.05 (s, 3H), 3.18 (s, 3H), 3.80 (s, 3H), 3.99 (m, 1H), 4.18 (m, 1H), 7.02 (dd, 1H), 7.26 (d, 1H), 7.54 (d, 1H), 8.43 (br s, 3H). Using N-(2-cyano-4-methoxyphenyl)-N-(methylsulfonyl)-methanesulfonamide (Preparation 244).

192	N-[2-(aminomethyl)-5-methoxyphenyl]-N-methylmethanesulfonamide hydrochloride	MS m/z 245 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.07 (s, 3H), 3.22 (s, 3H), 3.81 (s, 3H), 3.93 (br m, 1H), 4.13 (br m, 1H), 7.07 (dd, 1H), 7.17 (d, 1H), 7.58 (d, 1H), 8.24 (br s, 3H). Using N-(2-cyano-5-methoxyphenyl)-N-methylmethanesulfonamide (Preparation 224).
193	N-[2-(aminomethyl)-phenyl]-3-methoxy-N-methylbenzenesulfonamide hydrochloride	MS m/z 307 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.37 (s, 3H), 3.78 (s, 3H), 4.11 (m, 1H), 4.31 (m, 1H), 6.61 (m, 1H), 6.99 (m, 1H), 7.17 (d, 1H), 7.28-7.35 (m, 2H), 7.42-7.46 (m, 1H), 7.56 (t, 1H), 7.73 (m, 1H), 8.53 (br s, 3H). Using N-(2-cyanophenyl)-3-methoxy-N-methylbenzenesulfonamide (Preparation 225).
194	N-[2-(aminomethyl)-4-methoxyphenyl]-N-ethylmethanesulfonamide hydrochloride	MS m/z 259 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.99 (t, 3H), 3.02 (3.46 (m, 1H), 3.71 (m, 1H), 3.81 (s, 3H), 4.09 (m, 2H), 7.02 (dd, 1H), 7.30 (d, 1H), 7.50 (d, 1H), 8.40 (br s, 3H). N-(2-cyano-4-methoxyphenyl)-N-(methylsulfonyl)-methanesulfonamide (Preparation 244) and ethyl iodide.
195	N-[2-(aminomethyl)-4-methoxyphenyl]-N-methylbenzenesulfonamide hydrochloride	MS m/z 307 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.11 (s, 3H), 3.54 (s, 3H), 4.06 (m, 1H), 4.30 (m, 1H), 6.45 (d, 1H), 6.82 (m, 1H), 7.30 (d, 1H), 7.58-7.75 (m, 4H), 7.77 (m, 1H), 8.41 (br s, 3H). N-(4-methoxy-2-cyanophenyl)-N-(phenylsulfonyl)benzenesulfonamide (Preparation 246) and methyl iodide.
196	N-[2-(aminomethyl)-4-methoxyphenyl]-N-ethylbenzenesulfonamide hydrochloride	MS m/z 321 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.96 (t, 3H), 3.20 (m, 1H), 3.78 (s, 3H), 3.86 (m, 1H), 4.15 (m, 2H), 6.47 (d, 1H), 6.85 (m, 1H), 7.31 (s, 1H), 7.61 (m, 4H), 7.76 (m, 1H), 8.40 (br s, 3H). N-(4-methoxy-2-cyanophenyl)-N-(phenylsulfonyl)benzenesulfonamide (Preparation 246) and ethyl iodide.
197	N-(3-(aminomethyl)pyridin-4-yl)-N-methylmethanesulfonamide	MS m/z 216 [M+H] ⁺ Using N-(3-cyanopyridin-4-yl)methanesulfonamide (Preparation 248) and methyl iodide.

198	N-[3-(aminomethyl)-5-methylpyridin-2-yl]-N-methylmethanesulfonamide hydrochloride	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.36 (s, 3H), 3.09 (s, 3H), 3.18 (s, 3H), 7.90 (d, 1H), 8.26 (br s, 3H), 8.39 (d, 1H). MS m/z 230 [M+H] ⁺ Using N-(3-cyano-5-methylpyridin-2-yl)-N-(methylsulfonyl)methanesulfonamide (Preparation 245) and methyl iodide.
199	1-[2-(aminomethyl)-phenyl]methanesulfonamide hydrochloride	MS m/z 201 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 4.18 (br m, 2H), 4.48 (s, 2H), 7.00 (s, 2H), 7.43 (m, 3H), 7.53 (m, 1H), 8.23 (br s, 3H). Using Steps 2 and 3 only with (2-cyanophenyl)methanesulfonamide.
200	N-[2-(aminomethyl)-4-methylphenyl]-N-ethylmethanesulfonamide hydrochloride	MS m/z 243 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.98 (t, 3H), 2.50 (s, 3H), 3.03 (s, 3H), 3.49 (m, 1H), 3.70 (m, 1H), 4.09 (m, 2H), 7.29 (m, 1H), 7.45 (d, 1H), 7.51 (s, 1H), 8.42 (br s, 3H). MS m/z 243 [M+H] ⁺ Using N-(4-methyl-2-cyanophenyl)-N-(methylsulfonyl)-methanesulfonamide (Preparation 242) and ethyl iodide.
201	N-[3-(aminomethyl)-5-chloropyridin-2-yl]-N-ethylmethanesulfonamide hydrochloride	MS m/z 264 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.99 (t, 3H), 3.07 (s, 3H), 3.70 (q, 2H), 4.17 (m, 2H), 8.31 (s, 1H), 8.44 (br s, 3H), 8.67 (s, 1H). Using steps 2 and 3 only with N-(5-chloro-3-cyanopyridin-2-yl)-N-ethylmethanesulfonamide (Preparation 232).
202	N-[3-(aminomethyl)-5-chloropyridin-2-yl]-N-methylmethanesulfonamide hydrochloride	MS m/z 250 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.98 (s, 3H), 3.29 (s, 3H), 4.39 (m, 2H), 8.46 (s, 1H), 8.57 (m, 1H), 8.70 (br s, 3H). Using steps 2 and 3 only with N-(5-chloro-3-cyanopyridin-2-yl)-N-methylmethanesulfonamide (Preparation 233).
203	N-[3-(aminomethyl)-5-fluoropyridin-2-yl]-N-methylmethanesulfonamide hydrochloride	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.07 (s, 3H), 3.19 (s, 3H), 4.18 (br s, 2H), 8.10 (dd, 1H), 8.40 (br s, 3H), 8.60 (d, 1H). Using steps 2 and 3 only with N-(5-fluoro-3-cyanopyridin-2-yl)-N-methylmethanesulfonamide (Preparation 231).
204	N-[3-(aminomethyl)-5-fluoropyridin-2-yl]-N-ethylmethanesulfonamide hydrochloride	MS m/z 250 [M+H] ⁺ Using steps 2 and 3 only with N-(5-fluoro-3-cyanopyridin-2-yl)-N-ethylmethanesulfonamide (Preparation 235). Taken on directly to the next step.

205	N-[2-(aminomethyl)-4-fluorophenyl]-N-ethylmethanesulfonamide	MS m/z 247 [M+H] ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.98 (t, 3H), 3.07 (s, 3H), 3.48 (m, 1H), 3.64 (m, 1H), 3.71-3.83 (q, 2H), 7.13 (m, 1H), 7.44 (m, 2H). Using N-(4-fluoro-2-cyanophenyl)-N-(methylsulfonyl)-methanesulfonamide (Preparation 239), ethyl iodide and washing with saturated aqueous sodium bicarbonate solution.
206	N-[2-(aminomethyl)-4-chlorophenyl]-N-ethylmethanesulfonamide hydrochloride	MS m/z 263 [M+H] ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.00 (t, 3H), 3.55 (m, 1H), 3.75 (m, 1H), 4.03-4.20 (m, 2H), 7.55 (dd, 1H), 7.62 (m, 1H), 7.83 (d, 1H), 8.54 (br s, 3H). Using N-(4-chloro-2-cyanophenyl)-N-methylmethanesulfonamide (Preparation 237).
207	N-[3-(aminomethyl)-5-methylpyridin-2-yl]-N-ethylmethanesulfonamide hydrochloride	MS m/z 244 [M+H] ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.95 (t, 3H), 2.49 (s, 3H), 3.59 (s, 3H), 3.65 (q, 2H), 4.10 (m, 2H), 8.07 (d, 1H), 8.40 (d, 1H), 8.71 (br s, 3H). Using N-(3-cyano-5-methylpyridin-2-yl)-N-(methylsulfonyl)methanesulfonamide (Preparation 245) and ethyl iodide.

Preparation 208

tert-Butyl 2-(N-methyl-2-oxooxazolidine-3-sulfonamido)benzylcarbamate
hydrochloride

5 To a suspension of tert-butyl (2-((2-oxooxazolidine)-3-sulfonamido)benzyl)carbamate (**Preparation 251**, 2.5 g, 6.73 mmol) and anhydrous potassium carbonate (2.32 g, 16.82 mmol) in acetone (300 mL) was added methyl iodide (2.39 g, 16.83 mmol) and the reaction heated to reflux for 18 hours. The reaction was cooled and concentrated in vacuo. The residue was partitioned between water and

10 DCM, the organic layer was collected, washed with brine and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 35% EtOAc in hexanes before being treated with 4M HCl in dioxane (7 mL) and stirring at room temperature for 18 hours. The reaction was concentrated *in vacuo* and triturated with ether/pentane to afford the title compound as the hydrochloride salt (1.60 g, 68%

15 over 2 steps). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.36 (s, 3H), 3.56 (m, 1H), 3.98 (m, 1H), 4.08 (m, 1H), 4.19 (m, 1H), 4.44 (m, 2H), 7.52-7.68 (m, 4H), 8.36 (br s, 3H). MS m/z 285 [M+H]⁺

Preparation 209

Racemic N-[2-(1-Aminoethyl)phenyl]-N-methylmethanesulfonamide

To a solution of N-(2-acetylphenyl)-N-methylmethanesulfonamide (**Preparation 246**, 10 g, 43.99 mmol) in EtOH (150 mL) was added triethylamine (7.93 mL, 57 mmol) and hydroxylamine hydrochloride (3.98 g, 57 mmol) and the reaction was heated to 80 °C for 18 hours. The reaction was cooled, diluted with EtOAc, washed with brine, dried
 5 over sodium sulfate and concentrated *in vacuo*. The residue was dissolved in MeOH (50 mL) and ammonium formate (2.15 g, 34 mmol) and activated zinc dust (2.25 g, 34 mmol) were added. The reaction was heated to reflux for 18 hours. The reaction was filtered through celite and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 12% MeOH in DCM to afford the title
 10 compound as a colorless oil (1.2 g, 77%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.48 (d, 3H), 3.16 (s, 3H), 3.22 (s, 3H), 4.76 (m, 1H), 7.40-7.80 (m, 4H), 8.32 (s, 1H). MS m/z 229 [M+H]⁺

Preparation 210

N-[4-(2-Aminoethyl)phenyl]benzenesulfonamide

15 To a solution of 2-(2-aminoethyl)aniline (30 g, 220 mmol) in DCM (700 mL) at 0 °C was added triethylamine (36.8 mL, 264 mmol) followed by tert-butylidicarbonate (52.9 g, 242 mmol) and the reaction was allowed to warm to room temperature stirring for 2 hours. The reaction was added to water (500 mL), the organic layer collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo* to afford a yellow oil.
 20 The oil was dissolved in DCM (400 mL) and pyridine (20 mL), and benzenesulfonyl chloride (26.1 mL, 203 mmol) was added. The reaction was stirred at room temperature for 48 hours. Further benzenesulfonyl chloride (6.51 mL, 0.3 eq) was added and the reaction continued for 24 hours. The reaction was washed with 1M aqueous HCl solution (500 mL), concentrated aqueous ammonia solution (400 mL), brine (500 mL),
 25 dried over sodium sulfate and concentrated *in vacuo*. The residue was recrystallised from EtOAc/Ether to afford a white solid. The solid was dissolved in dioxane (200 mL), 4M HCl in dioxane (282 mL) was added and the reaction was stirred at room temperature for 18 hours. The reaction was concentrated *in vacuo* and the residue suspended in hot MeOH (150 mL). 7M ammonia in MeOH (150 mL) was added and the
 30 solution cooled. The resulting precipitate was collected and purified further using silica gel column chromatography eluting with 0.4% NH₃ in 10-15% MeOH in DCM to afford the title compound as a yellow solid (16.8 g, 26% over three steps).

¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.64 (t, 2H), 2.82 (t, 2H), 6.95-7.05 (m, 4H), 7.50-7.60 (m, 3H), 7.53 (m, 2H). MS m/z 275 [M-H]⁻

Preparation 211**N-[2-(Aminomethyl)-4-chlorophenyl]-N-methylmethanesulfonamide hydrochloride**

To a solution of tert-butyl {5-chloro-2-[methyl(methylsulfonyl)-amino]benzyl}carbamate (**Preparation 213**, 8.2 g, 23 mmol) in MeOH (100 mL) was added 4M HCl in dioxane (100 mL) at 0 °C and the reaction was stirred at room temperature for 5 hours. The reaction was concentrated *in vacuo* and triturated with a 1:1 mixture of MeCN:ether to afford the title compound as the hydrochloride salt (8.00 g, 100%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.07 (s, 3H), 3.20 (s, 3H), 4.05 (br m, 1H), 4.20 (br m, 1H), 7.58 (m, 1H), 7.66 (m, 1H), 7.73 (m, 1H), 8.32 (br s, 3H). MS m/z 249 [M+H]

Preparation 212**N-[2-(Aminomethyl)-4-methoxyphenyl]-N-methylpyridine-3-sulfonamide dihydrochloride**

The title compound was prepared according to the method described for **Preparation 211** using tert-butyl {5-methoxy-2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}carbamate (**Preparation 214**). MS m/z 308 [M+H]⁺

Preparation 213**tert-Butyl {5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}carbamate**

To a solution of N-(4-chloro-2-cyanophenyl)-N-methylmethanesulfonamide (**Preparation 236**, 6.20 g, 25.40 mmol) in MeOH (150 mL) was added di-tert-butyl dicarbonate (11.71 mL, 50.82 mmol) and NiCl₂·6H₂O (1.20 g, 5.08 mmol). The reaction was cooled to 0 °C and NaBH₄ (9.61 g, 254 mmol) was added portion-wise. The reaction was stirred at room temperature for 6 hours before being quenched by the addition of diethylenetriamine with stirring for 30 minutes. The reaction was concentrated *in vacuo* and partitioned between EtOAc and saturated aqueous sodium bicarbonate. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography to afford the title compound (8.20 g, 93%). ¹H NMR (400 MHz,

DMSO-d₆): 3.06(s, 3H), 3.21(s, 3H), 4.01-4.19 (m, 2H), 7.54-7.57 (q, 1H), 7.65-7.68 (d, 1H), 7.82-7.83 (d, 1H) MS m/z 349 [M+H]⁺ and 249 [M-Boc+H]⁺

Preparation 214**tert-Butyl {5-methoxy-2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}carbamate**

To a solution of tert-butyl [5-methoxy-2-(methylamino)benzyl]carbamate (**Preparation 215**, 3.8 g, 10 mmol) in THF (20 mL) was added NaH (373 mg, 15 mmol)

at 0 °C and the reaction was stirred for 15 minutes before the addition of pyridine-3-sulfonyl chloride (1.36 mL, 11 mmol) dropwise. The reaction was stirred at room temperature for 18 hours before being quenched with water and extracted into EtOAc. The organic layer was collected, washed with water, brine, dried over sodium sulfate
 5 and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 55% EtOAc in hexanes to afford the title compound (3.3 g, 78%). MS m/z 407 [M-H]-

Preparation 215

tert-Butyl [5-methoxy-2-(methylamino)benzyl]carbamate

10 The title compound was prepared according to the method described for **Preparation 213** using 5-methoxy-2-(methylamino)benzonitrile (**Preparation 250**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.37 (s, 9H), 2.66 (d, 3H), 3.63 (s, 3H), 3.95 (m, 2H), 4.77 (br s, 1H), 6.45 (d, 1H), 6.63 (br s, 1H), 6.69 (dd, 1H), 7.23 (br t, 1H). MS m/z 267 [M+H]⁺

Preparation 216

N-[2-(Aminomethyl)phenyl]-N-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]methanesulfonamide

To a solution of N-(2-cyanophenyl)-N-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]methanesulfonamide (**Preparation 222**, 8 g, 25 mmol) in MeOH (100 mL)
 20 was added NiCl₂·6H₂O (1.17 g, 5 mmol) followed by sodium borohydride (6.53 g, 172 mmol) at 0°C. The reaction was stirred at room temperature for 4 hours before being quenched by the addition of diethylenetetramine. The reaction was concentrated *in vacuo* and purified using silica gel column chromatography eluting with 10% MeOH in DCM to afford the title compound (4.9 g, 60%). Taken on directly to the next step.

Preparation 217

N-[3-(Aminomethyl)pyridin-2-yl]-N-methylmethanesulfonamide

To a solution of N-(3-cyanopyridin-2-yl)-N-methylmethanesulfonamide (**Preparation 230**, 10 g, 47 mmol) in methanolic ammonia (100 mL) was added Raney Nickel (2 g) and the reaction was hydrogenated at 40 psi at room temperature for 18
 30 hours. The reaction was filtered through celite, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 10% MeOH in DCM to afford the title compound (7.5 g, 74%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.13 (s, 6H), 3.82 (br s, 2H), 7.44 (m, 1H), 8.03 (m, 1H), 8.37 (m, 1H).

Preparation 218**N-[3-(Aminomethyl)pyridin-2-yl]-N-ethylmethanesulfonamide**

The title compound was prepared according to the method described for **Preparation 217** using N-(3-cyanopyridin-2-yl)-N-methylmethanesulfonamide (**Preparation 234**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 0.94 (t, 3H), 3.06 (s, 3H), 3.63 (q, 2H), 3.86 (br s, 2H), 7.46 (m, 1H), 8.06 (m, 1H), 8.41 (m, 1H). MS m/z 230 [M+H]

Preparation 219**N-[3-(Aminomethyl)pyrazin-2-yl]-N-methylbenzenesulfonamide**

A solution of N-(3-cyanopyrazin-2-yl)-N-methylbenzenesulfonamide (**Preparation 228**, 7.2 g, 32 mmol) in AcOH (100 mL) was purged under nitrogen for 15 minutes followed by the addition of 10% Pd-C (1.4 g) and hydrogenated under at 40 psi hydrogen in a Parr-shaker for 18 hours. The reaction was filtered through celite, concentrated *in vacuo*, neutralized with 1N NaOH and extracted with DCM. The organic layer was collected, dried over sodium sulfate, concentrated *in vacuo* and purified by silica gel column chromatography eluting with 10% MeOH in DCM to afford the title compound as a yellow solid (4.1 g, 46%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.06 (s, 3H), 4.04 (s, 2H), 7.17 (m, 4H), 7.33 (m, 1H), 8.34 (d, 1H), 8.64 (d, 1H). MS m/z 279 [M+H]⁺

Preparation 220**N-[2-(Aminomethyl)phenyl]-N-methylpyridine-3-sulfonamide hydrochloride**

To a solution of tert-butyl {2-[(pyridin-3-ylsulfonyl)amino]benzyl}carbamate (**Preparation 227**, 4.57 g, 12 mmol) in acetone (100 mL) was added potassium carbonate (5.20 g, 38 mmol) followed by methyl iodide (1.56 mL, 25 mmol). The reaction was heated to reflux for 2 hours. The reaction was evaporated to dryness and partitioned between water & ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate, evaporated and purified by silica gel column chromatography eluting with 52% EtOAc in hexanes. The residue was dissolved in MeOH (25 mL) and 4M HCl in dioxane (25 mL) was added with stirring at room temperature for 4 hours. The reaction was concentrated *in vacuo* and triturated with MeCN-ether to afford the title compound as the hydrochloride salt (4.2 g, 100%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.19 (s, 3H), 4.14 (m, 1H), 4.32 (m, 1H), 6.65 (d, 1H), 7.31 (t, 1H), 7.47 (t, 1H), 7.67-7.72 (m, 2H), 7.99 (m, 1H), 8.36 (br s, 3H), 8.71 (m, 1H), 8.95 (m, 1H),

Preparation 221**N-[3-(Aminomethyl)pyrazin-2-yl]-N-ethylmethanesulfonamide**

The title compound was prepared according to the method described for **Preparation 219** using N-(3-cyanopyrazin-2-yl)-N-ethylmethanesulfonamide
 5 (Preparation 229). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.00 (t, 3H), 1.86 (br s, 2H), 3.13 (s, 3H), 3.68 (q, 2H), 3.95 (s, 2H). MS m/z 231 [M+H]⁺

Preparation 222**N-(2-Cyanophenyl)-N-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]methanesulfonamide**

To a suspension of N-(2-cyanophenyl)methanesulfonamide (Preparation 223, 7
 10 g, 25 mmol) and polymer bound triphenylphosphine (14 g, 53 mmol) in anhydrous THF (100 mL) was added DEAD (8.42 mL, 53 mmol) followed by 2-(tetrahydro-pyran-2-yloxy)ethanol (7.82 g, 146 mmol) drop-wise at 0°C. The reaction was stirred at room temperature for 5 hours before filtering through celite. The filtrate was concentrated *in vacuo* and purified using silica gel column chromatography eluting with 30-35% EtOAc
 15 in hexanes to afford the title compound (8 g, 69%). ¹H NMR (400 MHz, MeOD): δ ppm 1.42-1.62 (m, 6H), 3.14 (s, 3H), 3.40-3.55 (m, 2H), 3.70-4.00 (m, 5H), 7.53 (m, 1H), 7.69 (m, 1H), 7.72-7.81 (m, 2H).

Preparation 223

N-(2-Cyanophenyl)-N-methylmethanesulfonamide To a solution of N-(2-cyanophenyl)-N-(methylsulfonyl)methanesulfonamide (Preparation 238, 300 g, 1.09
 20 mol) in THF (2 L) was added 40% aqueous sodium hydroxide (2 L), benzyl triethylammonium chloride (24.91 g, 0.100 mol), and iodomethane (81.68 mL, 1.31 mol.) The reaction was stirred at room temperature for 18 hours. The reaction was diluted with EtOAc and partitioned with brine. The organic layer was collected, dried over
 25 sodium sulfate and concentrated *in vacuo*. The residue was triturated in pentane-ether to afford the title compound (208 g, 90%). ¹H NMR (400 MHz, CDCl₃): δ ppm 3.11 (s, 3H), 3.38 (s, 3H), 7.43-7.47 (t, 1H), 7.52-7.55 (m, 1H), 7.63-7.71 (m, 2H); MS m/z 211[M-H]⁻

Preparation 224**N-(2-Cyano-5-methoxyphenyl)-N-methylmethanesulfonamide**

To a stirred solution of 4-methoxy-2-(methylamino)benzonitrile (Preparation 226, 11 g, 68 mmol) in THF at -78°C, 1M LiHMDS in THF (108.5 mL) was added drop wise. The solution was stirred for 30 minutes followed by the addition of methanesulfonyl chloride (7.92 mL, 102 mmol). The reaction was stirred for 1 hour before quenching with

saturated aqueous ammonium chloride solution and extracting into EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 40% EtOAc in hexanes to afford the title compound as a white solid (12 g, 73%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.13 (s, 3H), 3.26 (s, 3H), 3.88 (s, 3H), 7.11 (dd, 1H), 7.28 (d, 1H), 7.83 (d, 1H).

Preparation 225

N-(2-Cyanophenyl)-3-methoxy-N-methylbenzenesulfonamide

The title compound was prepared according to the method described for **Preparation 224** using 2-(methylamino)benzonitrile and 3-methoxybenzenesulfonyl chloride. Taken on directly to the next step. MS m/z 303 [M+H]⁺

Preparation 226

4-Methoxy-2-(methylamino)benzonitrile

To a solution of 2-fluoro-4-methoxybenzonitrile (1 g, 6.61 mmol) in MeCN (10 mL) was added 40% aqueous methylamine (20 mL) and the reaction heated to 60 °C in a sealed tube. The reaction was cooled, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 60% EtOAc in hexanes to afford the title compound as a white solid (600 mg, 56%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.76 (d, 3H), 3.78 (s, 3H), 6.12 (m, 2H), 6.22 (m, 1H), 7.37 (m, 1H). MS m/z 163 [M+H]⁺

Preparation 227

tert-Butyl {2-[(pyridin-3-ylsulfonyl)amino]benzyl}carbamate

To a solution of (2-amino-benzyl)-carbamic acid tert butyl ester (3.2 g, 14 mmol) in pyridine (25 mL) was added pyridine-3-sulfonylchloride (1.75 mL, 14 mmol) at 0°C. The reaction was stirred at room temperature for 4 hours before concentrating *in vacuo*. The residue was partitioned between EtOAc and water, the organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 65% EtOAc in hexanes to afford the title compound (4.5 g, 87%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.38 (s, 9H), 4.07 (m, 2H), 6.80 (d, 1H), 7.13 (m, 1H), 7.21 (m, 2H), 7.32 (m, 1H), 7.62 (m, 1H), 8.05 (m, 1H), 8.78-8.83 (m, 2H), 9.92 (s, 1H). MS m/z 364 [M+H]⁺

Preparation 228

N-(3-Cyanopyrazin-2-yl)-N-methylbenzenesulfonamide

To a solution of 2-chloro-3-cyanopyrazine (5 g, 35.94 mmol) and Cs₂CO₃ (16.27 g, 50 mmol) in acetonitrile (75 mL) was added N-methylbenzenesulfonamide (7.37 g, 43 mmol) and the reaction heated to 80 °C for 3 hours. The reaction mixture was

concentrated *in vacuo* and the residue partitioned between water and EtOAc. The organic layer was collected, dried, concentrated *in vacuo* and purified by silica gel column chromatography eluting with 50% EtOAc in hexanes to afford the title compound (7.2 g, 73%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.12 (s, 3H), 7.64 (m, 4H), 7.76 (m, 1H), 8.80 (d, 1H), 8.85 (d, 1H).

Preparation 229

N-(3-Cyanopyrazin-2-yl)-N-ethylmethanesulfonamide

The title compound was prepared according to the method described for **Preparation 228** using N-ethylmethanesulfonamide. ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.09 (t, 3H), 3.20 (s, 3H), 3.83 (q, 2H), 8.89 (d, 1H), 8.97 (d, 1H).

Preparation 230

N-(3-Cyanopyridin-2-yl)-N-methylmethanesulfonamide

To a solution of 2-chloronicotinonitrile (10 g, 71.9 mmol) in MeCN (200 mL) was added cesium carbonate (32.5 g, 99 mmol) followed by N-methylmethanesulfonamide (9.42 g, 86 mmol) and the reaction was heated to 80 °C for 3 hours. The reaction was cooled, concentrated *in vacuo* and partitioned between EtOAc and water. The organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 50% EtOAc in hexanes to afford the title compound (12.9 g, 85%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.21 (s, 3H), 3.28 (s, 3H), 7.62 (m, 1H), 8.45 (d, 1H), 8.77 (d, 1H). MS m/z 212 [M+H]⁺

The following Preparations (**Preparations 231 – 235**) were prepared according to the method described for **Preparation 230** using the appropriate chloropyridine and sulfonamide as described below, and taken directly on to the next step:

Preparation Number	Name	Data/SM
231	N-(5-fluoro-3-cyanopyridin-2-yl)-N-methylmethanesulfonamide	MS m/z 230 [M+H] ⁺ Using N-methylmethanesulfonamide and 2-chloro-5-fluoronicotinonitrile.

232	N-(5-chloro-3-cyanopyridin-2-yl)-N-ethylmethanesulfonamide	MS m/z 260 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.17 (t, 3H), 3.10 (s, 3H), 3.87 (q, 2H), 8.02 (s, 1H), 8.62 (s, 1H). Using N-ethylmethanesulfonamide and 2,5-dichloronicotinonitrile.
233	N-(5-chloro-3-cyanopyridin-2-yl)-N-methylmethanesulfonamide	MS m/z 246 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.21 (s, 3H), 3.27 (s, 3H), 8.73 (d, 1H), 8.86 (d, 1H). Using N-methylmethanesulfonamide and 2,5-dichloronicotinonitrile.
234	N-(3-cyanopyridin-2-yl)-N-ethylmethanesulfonamide	MS m/z 226 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.05 (t, 3H), 3.17 (s, 3H), 3.77 (q, 2H), 7.67 (m, 1H), 8.49 (m, 1H), 8.84 (m, 1H). Using 2-chloronicotinonitrile and N-ethylmethanesulfonamide.
235	N-(5-fluoro-3-cyanopyridin-2-yl)-N-ethylmethanesulfonamide	MS m/z 244 [M+H] ⁺ Using N-ethylmethanesulfonamide and 2-chloro-5-fluoronicotinonitrile.

Preparation 236**N-(4-Chloro-2-cyanophenyl)-N-methylmethanesulfonamide**

A solution of N-(4-chloro-2-cyanophenyl)-N-(methylsulfonyl)methanesulfonamide (Preparation 237, 8.00 g, 25.91 mmol) in THF (100 mL) and 40% aqueous NaOH solution (100 mL) was cooled to 0°C. Benzyltriethylammonium chloride (0.59 g, 2.591 mmol) and MeI (5.64 mL, 90.68 mmol) were added and the reaction was stirred for 18 hours. The reaction was partitioned between EtOAc and water and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 30% EtOAc in hexanes to afford the title compound as a yellow solid (6.20 g, 97%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.14 (s, 3H), 3.26 (s, 3H), 7.78 (d, 1H), 7.88 (dd, 1H), 8.13 (d, 1H).

Preparation 237**N-(4-Chloro-2-cyanophenyl)-N-(methylsulfonyl)methanesulfonamide**

To a solution of 2-amino-5-chloro-benzonitrile (5.00 g, 32.77 mmol) in pyridine (100 mL) at 0 °C was added methanesulphonylchloride (10.21 mL, 131.07 mmol) and the reaction stirred at room temperature for 18 hours. The reaction was concentrated *in vacuo* and partitioned between 2N HCl and EtOAc. The organic layer was collected,

washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was triturated with 1:1 acetonitrile:ether to afford the title compound (8.00 g, 79%).

¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.61 (s, 6H), 7.90 (d, 1H), 7.98 (dd, 1H), 8.31 (d, 1H).

5

The following Preparations (**Preparations 238 – 246**) were prepared according to the method described for **Preparation 237** using the appropriate aniline as described below:

Preparation Number	Name	Data/SM
238	N-(2-cyanophenyl)-N-(methylsulfonyl)methanesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.10 (t, 3H), 3.12 (s, 3H), 3.67-3.73 (q, 2H), 7.59 (t, 1H), 7.73-7.75 (d, 1H), 7.82 (t, 1H), 7.93 (d, 1H). Using 2-aminobenzonitrile.
239	N-(4-fluoro-2-cyanophenyl)-N-(methylsulfonyl)methanesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.61 (s, 6H), 7.70-7.80 (m, 1H), 7.93-7.96 (m, 1H), 8.11-8.14 (m, 1H). Using 2-amino-5-fluorobenzonitrile.
240	N-(3-fluoro-2-cyanophenyl)-N-(methylsulfonyl)methanesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.10 (s, 3H), 3.20 (s, 3H), 7.54-7.58 (m, 1H), 7.63-7.65 (m, 1H), 7.83-7.89 (m, 1H). Using 2-amino-6-fluorobenzonitrile.
241	N-(2-cyanophenyl)-N-(ethylsulfonyl)ethanesulfonamide	Taken on directly to the next step as crude. Using 2-aminobenzonitrile.
242	N-(4-methyl-2-cyanophenyl)-N-(methylsulfonyl)methanesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 2.44 (s, 3H), 3.50 (s, 6H), 7.35-7.37 (d, 1H), 7.48-7.50 (d, 1H), 7.59 (s, 1H). Using 2-amino-5-methylbenzonitrile.
243	N-(2-cyanophenyl)-N-(phenylsulfonyl)benzenesulfonamide	MS m/z 399 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 7.18 (m, 1H), 7.65-7.88 (m, 12H), 8.02 (m, 1H). Using 2-aminobenzonitrile and benzenesulfonyl chloride.
244	N-(2-cyano-4-methoxyphenyl)-N-(methylsulfonyl)methanesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 3.57 (s, 6H), 3.87 (s, 3H), 7.37 (dd, 1H), 7.65 (d, 1H), 7.75 (d, 1H). Using 2-amino-4-methoxybenzonitrile and methanesulfonyl chloride.

245	N-(3-cyano-5-methylpyridin-2-yl)-N-(methylsulfonyl)methanesulfonamide	Taken directly on to the next step. Using 2-amino-5-methylpyridine-3-carbonitrile and methanesulfonyl chloride.
246	N-(4-methoxy-2-cyanophenyl)-N-(phenylsulfonyl)benzenesulfonamide	Taken directly on to the next step. Using 2-amino-5-methylpyridine-3-carbonitrile and methanesulfonyl chloride.

Preparation 247**N-(2-Acetylphenyl)-N-methylmethanesulfonamide**

To a solution of N-[2-(aminomethyl)phenyl]-N-methylmethanesulfonamide (**Preparation 252**, 10.5 g, 49 mmol) in acetone (250 mL) was added potassium carbonate (13.59 g, 98.47 mmol) and methyl iodide (6.13 mL, 98.47 mmol) at 0 °C followed by heating to 60 °C for 4 hours. The reaction was cooled and concentrated *in vacuo*. The residue was partitioned between EtOAc and water, the organic layer was collected, washed with brine, dried and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 40% EtOAc in hexanes to afford the title compound (10 g, 89%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.54 (s, 3H), 2.97 (s, 3H), 3.29 (s, 3H), 7.46 (m, 1H), 7.60-7.64 (m, 3H). MS m/z 228 [M+H]⁺

Preparation 248**N-[3-(Aminomethyl)pyridin-4-yl]methanesulfonamide**

The title compound was prepared according to the method described for **Preparation 247** using 3-cyano-4-aminopyridine. ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.92 (s, 3H), 7.31 (d, 1H), 7.99 (d, 1H), 8.66 (s, 1H), 13.14 (br s, 1H). MS m/z 198 [M+H]⁺

Preparation 249**Benzyl [2-(aminomethyl)phenyl]methylcarbamate hydrochloride**

To a solution of tert-butyl [2-(methylamino)benzyl]carbamate (**PCT Intl. Appl. 2004046107**, 1.7 g, 7.2 mmol) in THF (25 mL) at 0 °C was added NaH (0.25 g, 10.8 mmol) followed by Cbz-chloride (1.22 g, 7.2 mmol) and catalytic DMAP (9 mg, 0.72 mmol). The reaction was heated to reflux for 2 hours before cooling, quenching with water and extracting into EtOAc. The organic extracts were dried over sodium sulfate, concentrated *in vacuo* and purified by silica gel column chromatography. The residue was dissolved in MeOH (10 mL) and 20% HCl in dioxane (10 mL) was added with stirring at room temperature for 18 hours. The reaction was concentrated *in vacuo*, and

trituated with pentane and ether to afford the title compound as the hydrochloride salt (1.2 g, 91%). MS m/z 271 [M+H]⁺

Preparation 250

5-Methoxy-2-(methylamino)benzonitrile

5 To a solution of 2-amino-5-methoxybenzonitrile (10 g, 67 mmol) in DMF (100 mL) was added tBuOK (9.46 g, 84 mmol) followed by dimethyl oxalate (11.95 g, 101 mmol) at 0 °C and the reaction was heated to 120 °C for 4 hours. The reaction was cooled, quenched by the addition of ice-water and extracted into EtOAc. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The
10 residue was purified using silica gel column chromatography to afford the title compound (2.9 g, 27%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.72 (s, 3H), 3.67 (s, 3H), 5.77 (br s, 1H), 6.67 (d, 1H), 7.06 (d, 1H), 7.11 (dd, 1H).

Preparation 251

tert-Butyl 2-(2-oxooxazolidine-3-sulfonamido)benzylcarbamate

15 To a solution of chlorosulfonyl isocyanate (2 g, 8.99 mmol) in dry DCM (20 mL) at 0 °C was added bromoethanol (0.60 mL, 8.19 mmol) and the reaction was stirred at room temperature for 10 minutes. Triethylamine (2.74 mL, 19.79 mmol) in DCM was added followed by (2-amino-benzyl)-carbamic acid tert-butyl ester (1.9 g, 8.19 mmol) and the reaction stirred at room temperature for 18 hours. The reaction was quenched
20 with water, extracted into DCM, the organic layer was collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 20% EtOAc in hexanes to afford the title compound (2.5 g, 75%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.43 (s, 9H), 3.81 (t, 2H), 4.25 (m, 2H), 4.36 (t, 2H), 7.23-7.37 (m, 5H), 10.42 (br s, 1H).

25

Preparation 252

N-[2-(Aminomethyl)phenyl]-N-methylmethanesulfonamide

To a solution of 1-(2-aminophenyl)ethanone (2 g, 14.8 mmol) in pyridine (20 mL) at 0 °C was added methanesulfonyl chloride (4.6 mL, 59 mmol) and the reaction was stirred at room temperature for 18 hours. The reaction was concentrated *in vacuo* and
30 partitioned between 2N HCl and EtOAc. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo* to afford the title compound as a brown solid (1.6 g, 51%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.65 (s, 3H), 3.18 (s, 3H), 7.23 (t, 1H), 7.58-7.67 (m, 2H), 8.07 (d, 1H). MS m/z 212 [M-H]⁺

Preparation 253

4-((2-(N-Ethylphenylsulfonamido)-5-methoxybenzyl)amino)-6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

To a solution of 4-((2-(N-ethylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (**Preparation 259**, 300 mg, 0.36 mmol) in anhydrous THF (4 mL) was added NMM (0.06 mL, 0.58 mmol) and isobutylchloroformate (0.07 mL, 0.58 mmol) at -20 °C and the reaction was stirred at this temperature for 30 minutes. Aqueous ammonia (0.2 mL) was added and the reaction stirred at room temperature for 30 minutes. The reaction was diluted with water and extracted into EtOAc. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 40% EtOAc in hexanes to afford the title compound (180 mg, 60%). MS m/z 818 [M+H]⁺

Preparation 254

4-((2-(N-Methylphenylsulfonamido)-5-methoxybenzyl)amino)-6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide

The title compound was prepared according to the method described for **Preparation 253** using 4-((2-(N-methylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (**Preparation 260**). MS m/z 804 [M+H]⁺

Preparation 255

tert-Butyl 6-(4-((tert-butoxycarbonyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(methylamino)benzyl)amino)-3-(methylcarbamoyl)-1H-pyrazolo[3,4-d]pyrimidine-1-carboxylate

To a solution of (2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(methylcarbamoyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate (**Preparation 256**, 80 mg, 0.13 mmol) in anhydrous THF was added boron tribromide (0.08 mL, 0.9 mmol) at 0 °C and the reaction was stirred at this temperature for 2 hours. Another aliquot of boron tribromide was added (7eq) and the reaction continued stirring for a further 2 hours. The reaction was concentrated *in vacuo* and dissolved in DCM. The solution was washed with saturated aqueous sodium bicarbonate solution, brine, dried over sodium sulfate and

concentrated *in vacuo*. The residue was purified using preparative TLC. The residue was dissolved in anhydrous THF (5 mL). To this solution at 0 °C was added triethylamine (0.04 mL, 0.3 mmol) followed by di-tert-butyl dicarbonate (0.05 mL, 0.21 mmol) and catalytic DMAP (1 mg, 0.008 mmol). The reaction was stirred at room temperature for 1 hour before concentrating *in vacuo*. The residue was purified by preparative TLC to afford the title compound. MS m/z 704 [M+H]⁺

Preparation 256

tert-Butyl 6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(methylamino)benzyl)amino)-3-(methylcarbamoyl)-1H-pyrazolo[3,4-d]pyrimidine-1-carboxylate

10

To a solution of benzyl (2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(methylcarbamoyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate (**Preparation 257**, 150 mg, 0.23 mmol) in anhydrous THF (5 mL) at 0 °C was added triethylamine (0.08 mL, 0.575 mmol) followed by di-tert-butyl dicarbonate (60 mg, 0.27 mmol) and a catalytic amount of DMAP. The reaction was stirred at room temperature for 18 hours. The reaction was partitioned between EtOAc and brine, the organic layer collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 32% EtOAc in hexanes. The residue was dissolved in ethanol (15 mL) and was hydrogenated at 30 psi for 1 hour over 10% palladium on carbon (10 mg). The reaction was filtered through celite, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 32% EtOAc in hexanes to afford the title compound as a white solid (50 mg, 80%). MS m/z 618 [M+H]⁺

20

Preparation 257

25

Benzyl (2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(methylcarbamoyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate

The title compound was prepared according to the method described for **Preparation 18** using benzyl (2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate (**Preparation 284**). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 2.82 (s, 3H), 3.17 (s, 3H), 3.88 (s, 3H), 4.26-4.39 (m, 2H), 4.65 (m, 1H), 4.79-4.90 (m, 2H), 5.03 (m, 1H), 7.09-7.42 (m, 10H), 7.70 (m, 1H), 8.83 (t, 1H), 9.90 (t, 1H), 14.07 (s, 1H).

30

The following **Preparations (Preparations 258 – 261)** were prepared according to the method described for **Preparation 11** using the appropriate pyrazolopyrimidine as described below.

5

Preparation Number	Name	Data/SM
258	6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-4-((2-(N-methylphenylsulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo-[3,4-d]pyrimidine-3-carboxylic acid	MS m/z 835 [M-H] ⁻ N-(2-(((6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-methyl)phenyl)-N-methylbenzenesulfonamide (Preparation 262).
259	4-((2-(N-ethylphenylsulfonamido)-5-methoxybenzyl)amino)-6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid	MS m/z 819 [M-H] ⁻ N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide (Preparation 266).
260	6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-methoxy-2-(N-methylphenylsulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid	MS m/z 805 [M+H] ⁺ N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide (Preparation 267).
261	6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-4-((2-(N-methylphenylsulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid	MS m/z 889 [M-H] ⁻ N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide (Preparation 264).

Preparation 262

N-(2-(((6-(2-Ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide

- 5 To a suspension of NaH (0.163 g, 6.79 mmol) in dry DMF (50 mL) was added N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-2-ethyl-5-fluorophenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide (**Preparation 274**, 2.1 g, 2.71 mmol) at 0 °C and the reaction stirred for 15 minutes. SEMCI was then added (1.06 mL, 5.97 mmol) and the reaction allowed to warm to room temperature.
- 10 The reaction was quenched with water, partitioned between EtOAc and brine, the organic layer collected, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography to afford the title compound (760 mg, 43%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm -0.10 (s, 9H), -0.05 (s, 9H), 0.89 (m, 4H), 1.03 (t, 3H), 2.88 (m, 2H), 3.09 (s, 3H), 3.57 (t, 2H), 3.73 (t, 2H), 4.93 (m, 1H), 5.16 (m,
- 15 1H), 5.31 (s, 2H), 5.61 (s, 2H), 6.54 (m, 1H), 7.10 (d, 1H), 7.17 (t, 1H), 7.30 (t, 1H), 7.42 (m, 1H), 7.46 (m, 1H), 7.62-7.67 (m, 4H), 7.76 (m, 1H), 7.95 (s, 1H). MS m/z 919 [M+H]⁺

- The following **Preparations (Preparations 263 – 268)** were prepared according
- 20 to the method described for **Preparation 131** using either DMF or THF and the appropriate pyrazolopyrimidine as described below.

Preparation Number	Name	Data/SM
263	N-ethyl-N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-methyl)phenyl)methanesulfonamide	MS m/z 925 [M+H] ⁺ N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-ethylmethanesulfonamide (Preparation 280).
264	N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide	MS m/z 973 [M+H] ⁺ N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide (Preparation 278).
265	N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-(2-((tert-butyldimethylsilyl)oxy)ethyl)methanesulfonamide	Taken on directly to the next step. N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-(2-((tert-butyldimethylsilyl)oxy)ethyl)methanesulfonamide (Preparation 279).
266	N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 901 [M+H] ⁺ N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide (Preparation 281).

267	N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-	MS m/z 885 [M-H] ⁻ N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide
268	N-(4-fluoro-2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 813 [M+H] ⁺ N-(4-fluoro-2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 283).

Preparation 269

6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-methoxy-2-(N-methylmethylsulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid

The title compound was prepared according to the methods described for **Preparations 262 and 258** using N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 273**). MS m/z 743 [M+H]⁺

Preparation 270

6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylmethylsulfonamido)-benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid

The title compound was prepared according to the methods described for **Preparations 262 and 258** using N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (**Preparation 275**). MS m/z 713 [M+H]⁺

Preparation 271

N-(2-(((6-(5-Fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-hydroxyphenyl)-N-methylmethanesulfonamide

To a solution of N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethane sulfonamide (**Preparation 273**, 300 mg, 0.43 mmol) in DCM (10 mL) at

0 °C was added boron tribromide (0.28 mL, 3.02 mmol). The reaction was stirred at room temperature for 30 minutes before concentrating *in vacuo*. The residue was partitioned between ethyl acetate and water, the organic layer was collected, dried and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 10% MeOH in DCM to afford the title compound as a white solid (250 mg, 87%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 3.02 (s, 3H), 3.12 (s, 3H), 4.11-4.21 (m, 2H), 4.80 (m, 1H), 4.94 (m, 1H), 6.67 (m, 1H), 6.76 (m, 1H), 6.97 (m, 1H), 7.29-7.33 (m, 2H), 7.57 (m, 1H), 9.58 (s, 1H), 10.36 (s, 1H), 13.88 (s, 1H). MS m/z 667 [M+H]⁺

10

The following **Preparations (Preparations 272 – 284)** were prepared according to the method described for **Preparation 137** in an organic solvent such as DCM or DMF and using the appropriate pyrazolopyrimidine as described below.

Preparation Number	Name	Data/SM
272	N-(2-(((6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 687 [M+H] ⁺ N-(2-(((6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 285).
273	N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide	MS m/z 695 [M+H] ⁺ N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (Preparation 286).
274	N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-2-ethyl-5-fluoro-phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-methyl)phenyl)-N-methylbenzenesulfonamide	MS m/z 773 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.18 (s, 6H), 0.93 (t, 3H), 0.95 (s, 9H), 2.81 (q, 2H), 3.06 (s, 3H), 4.89 (m, 1H), 5.17 (m, 1H), 6.53 (d, 1H), 7.14 (d, 1H), 7.18 (m, 1H), 7.31 (m, 1H), 7.33-7.45 (m, 3H), 7.61-7.66 (m, 4H), 7.76 (m, 1H). N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-2-ethyl-5-fluorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide (Preparation 294).

275	N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 665 [M+H] ⁺ N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 287).
276	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 741 [M+H] ⁺ N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 288).
277	N-(3-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-methyl)pyrazin-2-yl)-N-methylmethanesulfonamide	MS m/z 665 [M-H] ⁻ N-(3-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)pyrazin-2-yl)-N-methylmethanesulfonamide (Preparation 289).
278	N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-methyl)phenyl)-N-methylbenzenesulfonamide	MS m/z 827 [M+H] ⁺ N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide (Preparation 295).
279	N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-methyl)phenyl)-N-(2-((tert-butyldimethylsilyl)oxy)ethyl)methanesulfonamide	MS m/z 909 [M+H] ⁺ N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-(2-((tert-butyldimethylsilyl)oxy)ethyl)methanesulfonamide (Preparation 296).
280	N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-ethylmethanesulfonamide	MS m/z 779 [M+H] ⁺ N-(2-(((6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-ethylmethanesulfonamide (Preparation 297).

281	N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 771 [M+H] ⁺ N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide (Preparation 290).
282	N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 757 [M+H] ⁺ N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide (Preparation 291).
283	N-(4-fluoro-2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 683 [M+H] ⁺ N-(4-fluoro-2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 292).
284	Benzyl (2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate	MS m/z 721 [M+H] ⁺ Benzyl (2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate (Preparation 293).

The following Preparations (Preparations 285 – 293) were prepared according to the method described for Preparation 140 using either 4M HCl in dioxane or cHCl in MeOH with the appropriate pyrazolopyrimidine as described below.

5

Preparation Number	Name	Data/SM
285	N-(2-(((6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 561 [M+H] ⁺ N-(2-(((6-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 299).

286	N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide	MS m/z 569 [M+H] ⁺ N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (Preparation 300).
287	N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 539 [M+H] ⁺ N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 302).
288	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 615 [M+H] ⁺ N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 303).
289	N-(3-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)pyrazin-2-yl)-N-methylmethanesulfonamide	MS m/z 541 [M+H] ⁺ N-(3-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)pyrazin-2-yl)-N-methylmethanesulfonamide (Preparation 304).
290	N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 645 [M+H] ⁺ N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide (Preparation 308).
291	N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 631 [M+H] ⁺ N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide (Preparation 309).
292	N-(4-fluoro-2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 557 [M+H] ⁺ N-(4-fluoro-2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 310).

293	Benzyl (2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate	MS m/z 595 [M+H] ⁺ Benzyl (2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate (Preparation 311).
-----	---	---

Preparation 294

N-(2-(((6-(4-((tert-Butyldimethylsilyl)oxy)-2-ethyl-5-fluorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide

5 N-(2-(((6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzene sulfonamide (**Preparation 301**, 2.6 g, 3.48 mmol) was treated with TFA (5 mL) and the reaction stirred at room temperature for 30 minutes before concentrating *in vacuo*. The residue was diluted with methanol (20 mL), cooled in ice-water and

10 treated with a drop-wise addition of ethylene diamine until the solution became basic. The solution was concentrated *in vacuo* and purified using silica gel column chromatography eluting with EtOAc. The residue (1.7 g, 3.19 mmol) was dissolved in anhydrous THF (10 mL) and 2,6 lutidine (0.55 mL, 4.78 mmol) was added followed by TBDMS-triflate (0.88 mL, 3.83 mmol) at 0°C. The reaction was stirred for 18 hours. The

15 reaction was concentrated *in vacuo* and partitioned between water and ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate, concentrated *in vacuo* and purified by silica gel column chromatography eluting with 20% EtOAc in hexanes to afford the title compound as a white solid (1.9 g, 92%). MS m/z 647 [M+H]⁺

20 The following **Preparations (Preparations 295 – 297)** were prepared according to the method described for **Preparation 294** using the appropriate pyrazolopyrimidine as described below.

25 |

Preparation Number	Name	SM
295	N-(2-(((6-(4-((tert-butyl)dimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide	MS m/z 701 [M+H] ⁺ N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide (Preparation 305).
296	N-(2-(((6-(4-((tert-butyl)dimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-(2-((tert-butyl)dimethylsilyl)oxy)ethylmethanesulfonamide	MS m/z 783 [M+H] ⁺ N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-(2-((tetrahydro-2H-pyran-2-yl)oxy)ethyl)methanesulfonamide (Preparation 316).
297	N-(2-(((6-(4-((tert-butyl)dimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-ethylmethanesulfonamide	MS m/z 653 [M+H] ⁺ N-ethyl-N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)methanesulfonamide (Preparation 307).

Preparation 298

4-Chloro-6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine

5

To a solution of 4-(benzyloxy)-6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (**Preparation 306**, 12.5 g, 19.75 mmol) in THF (100 mL) was added 10% palladium on carbon (1.5 g) and the reaction was hydrogenated at 50 psi for 18 hours. The reaction was filtered and the filtrate was concentrated *in vacuo*. The residue was purified using silica gel column chromatography to afford a white solid. 5 g (9.22 mmol) was dissolved in DMF (50 mL) and cooled to 0°C. Oxalyl chloride (7.96 mL, 92 mmol) was added and the reaction stirred at room temperature for 6 hours. The reaction was quenched with water and extracted into EtOAc. The organic layer was collected, dried over sodium

10

sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography to afford the title compound (1.2 g, 23%). MS *m/z* 561 [M+H]⁺

Preparation 299

N-(2-(((6-(4-(Benzyloxy)-2-ethyl-5-fluorophenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

A solution of palladium acetate (42 mg, 0.19 mmol) and S-Phos (77 mg, 0.19 mmol) in ethanol (10 mL) was heated at 50 °C for 45 minutes after purging with nitrogen (Solution A). Meanwhile a solution of N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

(**Preparation 312**, 1.7 g, 3.77 mmol) in ethanol (30 mL) was treated with 2-(4-(benzyloxy)-2-ethyl-5-fluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

(**Preparation 323**, 1.83 g, 5.13 mmol) and an aqueous solution of potassium phosphate (1.6 g, 7.54 mmol) in water (12 mL). This solution was purged with nitrogen for 10 minutes (Solution B). Solution A was added to Solution B and the mixture heated at 80

°C for 18 hours. The reaction was cooled, concentrated *in vacuo*. The resulting black solid was suspended in ethyl acetate filtered through celite. The filtrate was concentrated *in vacuo*, the residue was taken up in EtOAc, washed with water, brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 40% EtOAc in hexanes to afford the title compound as a fluffy white solid (1.58 g, 65%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.02 (m, 3H), 1.55 (m, 2H), 1.74 (m, 1H), 1.85 (m, 1H), 2.00 (m, 1H), 2.43 (m, 1H), 2.84 (m, 2H), 3.07 (s, 3H), 3.16 (s, 3H), 3.63 (m, 1H), 3.93 (m, 1H), 4.75 (br m, 1H), 5.00 (br m, 1H), 5.22 (s, 2H), 5.85 (m, 1H), 7.10 (m, 1H), 7.30-7.54 (m, 10H), 8.22 (s, 1H), MS *m/z* 645 [M+H]⁺

The following **Preparations (Preparations 300 – 311)** were prepared according to the method described for **Preparation 299** using the appropriate chloro-pyrazolopyrimidine and arylboronic ester as described below.

Preparation Number	Name	Data	SM
300	N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.56 (m, 2H), 1.73 (m, 1H), 1.89 (m, 1H), 1.98-2.01 (m, 1H), 2.42-2.50 (m, 1H), 3.04 (s, 3H), 3.13 (s, 3H), 3.66 (m, 4H), 3.88 (s, 3H), 3.93 (m, 1H), 4.29-4.34 (m, 2H), 4.70 (m, 1H), 5.00 (m, 1H), 5.87 (m, 1H), 6.91 (m, 2H), 7.23 (d, 1H), 7.48 (d, 1H), 7.78 (m, 1H), 8.24 (s, 1H), 8.85 (t, 1H).	2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (PCT Intl. Appl. 2013014567) and N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (Preparation 314).
301	N-(2-(((6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide	MS m/z 747 [M+H] ⁺	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide (Preparation 313) and 2-[2-ethyl-5-fluoro-4-[[2-(trimethylsilyl)ethoxy]methoxy]phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (PCT Intl. Appl. 2013014567).

302	N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	MS m/z 623 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.56 (m, 2H), 1.73 (m, 1H), 1.86 (m, 1H), 2.04 (m, 2H), 3.08 (s, 3H), 3.18 (s, 3H), 3.65 (m, 1H), 3.88 (s, 3H), 3.92-3.96 (m, 1H), 4.30-4.38 (m, 2H), 4.75 (br s, 1H), 5.00 (br s, 1H), 5.87 (m, 1H), 7.20 (m, 1H), 7.30-7.38 (m, 3H), 7.54 (m, 1H), 7.70-7.73 (m, 1H), 8.24 (s, 1H), 8.90 (t, 1H).	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 312) and 2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (PCT Intl. Appl 2013014567).
303	N-(2-(((6-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.56 (m, 2H), 1.74 (m, 1H), 1.89 (m, 1H), 2.00 (m, 2H), 3.07 (s, 3H), 3.16 (s, 3H), 3.67 (m, 1H), 3.93 (m, 1H), 4.22-4.37 (m, 2H), 4.80 (br m, 1H), 5.00 (br m, 1H), 5.23 (s, 2H), 5.86 (m, 1H), 7.29-7.55 (m, 10H), 7.73 (m, 1H), 8.24 (s, 1H), 8.89 (t, 1H).	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (Preparation 312) and 2-(4-(benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (Preparation 325).
304	N-(3-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)pyrazin-2-yl)-N-methylmethanesulfonamide	MS m/z 625 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.56 (m, 2H), 1.65 (m, 1H), 1.89 (m, 1H), 2.00 (m, 1H), 2.49 (m, 1H), 3.13 (s, 3H), 3.18 (s, 3H), 3.66 (m, 1H), 3.88 (m, 4H), 4.25-4.30 (m, 2H), 5.03 (m, 2H), 5.85 (m, 1H), 7.19 (m, 1H), 7.58 (m, 1H), 8.29 (s, 1H), 8.53 (d, 1H), 8.60 (m, 1H), 9.03 (t, 1H).	N-(3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)pyrazin-2-yl)-N-methylmethanesulfonamide (Preparation 315) and 2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (PCT Intl. Appl 2013014567).

305	N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide	MS m/z 801 [M+H] ⁺	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide (Preparation 313) and (2-([2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-(2,2,2-trifluoroethyl)-phenoxy]methoxy)ethyl)(trimethyl)silane (Preparation 150).
306	4-(benzyloxy)-6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine	MS m/z 634 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm - 0.01 (s, 9H), 0.89 (t, 2H), 1.583(m, 2H), 1.75 (m, 1H), 1.92-2.05 (m, 2H), 2.44 (m, 1H), 3.67 (m, 1H), 3.77 (t, 2H), 4.01 (m, 1H), 4.34-4.47 (m, 2H), 5.39 (s, 2H), 5.70 (s, 2H), 5.95 (m, 1H), 7.36-7.54 (m, 6H), 8.01 (m, 1H), 8.34 (s, 1H).	4-(benzyloxy)-6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (Preparation 322) and (2-([2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-(2,2,2-trifluoroethyl)-phenoxy]methoxy)ethyl)(trimethyl)silane (Preparation 150).
307	N-ethyl-N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)methane-sulfonamide	MS m/z 753 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm - 0.05 (s, 9H), 0.80 (t, 2H), 1.00 (t, 3H), 1.56 (m, 2H), 1.75 (m, 1H), 1.98 (m, 1H), 2.04 (m, 1H), 2.49 (m, 1H), 3.06 (s, 3H), 3.54 (m, 1H), 3.57-3.76 (m, 4H), 3.93 (m, 1H), 4.20 (m, 2H), 4.85 (m, 1H), 5.00 (m, 1H), 5.32 (s, 2H), 5.87 (m, 1H), 7.29-7.39 (m, 4H), 7.52 (m, 1H), 7.66 (m, 1H), 8.26 (s, 1H), 8.91 (t, 1H).	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-ethylmethane-sulfonamide (Preparation 317) and (2-([2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-(2,2,2-trifluoroethyl)-phenoxy]methoxy)ethyl)(trimethyl)silane (Preparation 150).

308	N-ethyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 729 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 0.93 (t, 3H), 1.57 (m, 2H), 1.75 (m, 1H), 1.90 (m, 1H), 2.01 (m, 1H), 2.43 (m, 1H), 3.21 (m, 1H), 3.63-3.68 (m, 4H), 3.79-3.96 (m, 4H), 4.05-4.35 (m, 2H), 4.81 (m, 1H), 4.99 (m, 1H), 5.90 (m, 1H), 6.50 (m, 1H), 6.75 (m, 1H), 6.90 (m, 1H), 7.25 (m, 1H), 7.60-7.80 (m, 6H), 8.30 (s, 1H), 8.85 (t, 1H).	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-ethylbenzenesulfonamide (Preparation 318) and 2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (PCT Intl. Appl 2013014567).
309	N-methyl-N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)benzenesulfonamide	MS m/z 715 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.56 (m, 2H), 1.74 (m, 1H), 1.98 (m, 1H), 2.00 (m, 1H), 2.50 (m, 1H), 3.33 (s, 3H), 3.63-3.68 (m, 4H), 3.88-3.96 (m, 4H), 4.25-4.40 (m, 2H), 4.73 (m, 1H), 5.03 (m, 1H), 5.86 (m, 1H), 6.45 (m, 1H), 6.71 (m, 1H), 6.92 (m, 1H), 7.22 (m, 1H), 7.61-7.79 (m, 6H), 8.26 (s, 1H), 8.90 (t, 1H).	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylbenzenesulfonamide (Preparation 319) and 2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (PCT Intl. Appl 2013014567).
310	N-(4-fluoro-2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.56 (m, 2H), 1.74 (m, 1H), 1.86 (m, 1H), 2.01 (m, 1H), 2.42 (m, 1H), 3.08 (s, 3H), 3.16 (s, 3H), 3.60 (m, 1H), 3.88 (s, 3H), 3.96 (m, 1H), 4.03-4.31 (m, 2H), 4.76 (m, 1H), 5.01 (m, 1H), 5.88 (m, 1H), 7.11-7.22 (m, 3H), 7.61-7.71 (m, 2H), 8.24 (s, 1H), 8.93 (t, 1H).	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-fluorophenyl)-N-methylmethanesulfonamide (Preparation 320) and 2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetra-methyl-1,3,2-dioxaborolane (PCT Intl. Appl 2013014567).

311	Benzyl 2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-(methyl)carbamate	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.56 (m, 2H), 1.73 (m, 1H), 1.85 (m, 1H), 2.01 (m, 1H), 2.42 (m, 1H), 3.18 (s, 3H), 3.69 (m, 1H), 3.87 (s, 3H), 3.93 (m, 1H), 4.27-4.50 (m, 2H), 4.57 (m, 1H), 4.75 (m, 1H), 4.93-5.03 (m, 2H), 5.86 (m, 1H), 7.12-7.39 (m, 10H), 7.71 (m, 1H), 8.22 (s, 1H), 8.85 (t, 1H).	benzyl 2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate (Preparation 321) and 2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (PCT Intl. Appl. 2013014567).
-----	--	---	--

Preparation 312

N-(2-(((6-Chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

5 To a stirred solution of 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (**PCT Intl. Appl. 2013014567**, 3 g, 10.99 mmol) in anhydrous n-butanol (12 mL), containing DIPEA (6.69 mL, 38.45 mmol) was added N-[2-(aminomethyl)phenyl]-N-methylmethane sulphonamide hydrochloride (**PCT Intl. Appl. 2010058846**, 2.76 g, 10.98 mmol) and the reaction was heated in a sealed tube at 90 °C for 16 hours. The

10 reaction was concentrated *in vacuo* and the residue partitioned between water & ethyl acetate. The organic extracts were washed with brine, dried over sodium sulfate, concentrated *in vacuo* and triturated with pentane-ether to afford the title compound as an off white solid (3.5 g, 71%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.17 (m, 2H), 1.70 (m, 1H), 1.81 (m, 1H), 1.98 (m, 1H), 2.36 (m, 1H), 3.09 (s, 3H), 3.25 (s, 3H), 3.66 (m,

15 1H), 3.94 (m, 1H), 4.60 (br m, 1H), 4.90 (br m, 1H), 5.72 (m, 1H), 7.33-7.42 (m, 3H), 7.56 (m, 1H), 8.22 (s, 1H), 9.26 (m, 1H).

The following **Preparations (Preparations 313 – 321)** were prepared according to the method described for **Preparation 312** using 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (**PCT Intl. Appl. 2013014567**) or other suitable pyrazolopyrimidine and benzylamine as described below.

20

Preparation Number	Name	Data	SM
313	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylbenzenesulfonamide	¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.54 (m, 2H), 1.71 (m, 1H), 1.84 (m, 1H), 1.98 (m, 1H), 2.32 (m, 1H), 3.19 (s, 3H), 3.75 (m, 1H), 3.90 (m, 1H), 0.89 (br m, 1H), 5.00 (br m, 1H), 5.75 (m, 2H), 6.58 (m, 1H), 7.19 (t, 1H), 7.35 (t, 1H), 7.45 (m, 1H), 7.63-7.78 (m, 5H), 8.23 (s, 1H), 9.30 (m, 1H).	N-[2-(amino-methyl)phenyl]-N-methylbenzenesulfonamide hydrochloride (Preparation 190).
314	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide	MS m/z 481 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.54 (m, 2H), 1.73 (m, 1H), 1.84 (m, 1H), 1.98 (m, 1H), 2.37 (m, 1H), 3.06 (s, 3H), 3.21 (s, 3H), 3.63-3.77 (m, 4H), 3.94 (m, 1H), 4.54 (m, 1H), 4.95 (m, 1H), 5.74 (m, 1H), 6.94 (m, 2H), 7.48 (m, 1H), 8.22 (s, 1H), 9.22 (t, 1H).	N-[2-(aminomethyl)-4-methoxyphenyl]-N-(methylsulfonyl)methanesulfonamide hydrochloride (Preparation 191).
315	N-(3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)pyrazin-2-yl)-N-methylmethanesulfonamide	MS m/z 453 [M+H] ⁺ ¹ H NMR (400 MHz, DMSO-d ₆): δ ppm 1.54 (m, 2H), 1.73 (m, 1H), 1.85 (m, 1H), 1.98 (m, 1H), 2.33 (m, 1H), 3.15 (s, 3H), 3.31 (s, 3H), 3.66 (m, 1H), 3.90 (m, 1H), 4.90 (m, 2H), 5.73 (m, 1H), 8.24 (s, 1H), 8.54 (m, 1H), 8.62 (m, 1H), 9.44 (t, 1H).	N-(3-(aminomethyl)pyrazin-2-yl)-N-methylmethanesulfonamide hydrochloride (PCT Intl. Appl. 2008129380).

316	N-(2-(((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-(2-((tetrahydro-2H-pyran-2-yl)oxy)ethyl)methanesulfonamide	MS m/z 854 [M+H] ⁺	4-chloro-6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (Preparation 298) and N-[2-(aminomethyl)-phenyl]-N-[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]-methanesulfonamide (Preparation 216).
317	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-ethylmethanesulfonamide	MS m/z 465 [M+H] ⁺	N-[2-(aminomethyl)phenyl]-N-ethylmethanesulfonamide hydrochloride (Preparation 188).
318	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-ethylbenzenesulfonamide	MS m/z 557 [M+H] ⁺	N-[2-(aminomethyl)-4-methoxyphenyl]-N-ethylbenzenesulfonamide hydrochloride (Preparation 196).
319	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylbenzenesulfonamide	MS m/z 543 [M+H] ⁺	N-[2-(aminomethyl)-4-methoxyphenyl]-N-methylbenzenesulfonamide
320	N-(2-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-fluorophenyl)-N-methylmethanesulfonamide	MS m/z 469 [M+H] ⁺	N-[2-(aminomethyl)-4-fluorophenyl]-N-methylmethanesulfonamide hydrochloride (Preparation 186).

321	benzyl (2-(((6-chloro-1-(tetra-hydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)(methyl)carbamate	MS m/z 507 [M+H] ⁺	Benzyl [2-(aminomethyl)phenyl]methylcarbamate hydrochloride (Preparation 249).
-----	--	-------------------------------	---

Preparation 3224-(Benzyloxy)-6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine

To a suspension of NaH (0.48 g, 20.13 mmol) in THF (50 mL) at 0 °C was added
 5 benzyl alcohol (1.98 g, 18.30 mol) slowly. The mixture was allowed to stir for 45 minutes
 at 0 °C followed by the addition of 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-
 pyrazolo[3,4-d]pyrimidine (**PCT Intl. Appl. 2013014567**, 5 g, 18.30 mmol). The reaction
 was stirred at room temperature for 2 hours before being quenched with brine. The
 solution was extracted into EtOAc, washed with brine, dried over sodium sulfate and
 10 concentrated *in vacuo*. The residue was purified by silica gel column chromatography
 to afford the title compound (3.4 g, 54%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.56
 (m, 2H), 1.74-1.78 (m, 1H), 1.87-1.91 (m, 1H), 1.98-2.02 (m, 1H), 2.33-2.43 (m, 1H),
 3.71 (m, 1H), 3.92 (m, 1H), 5.62 (s, 2H), 5.86 (m, 1H), 7.38 (m, 3H), 7.55 (m, 2H), 8.37
 (s, 1H). MS m/z 345 [M+H]⁺

Preparation 3232-(4-(Benzyloxy)-2-ethyl-5-fluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

The title compound was prepared according to the method described for
Preparation 150 using 1-(benzyloxy)-4-bromo-5-ethyl-2-fluorobenzene (**Preparation**
324). Taken on directly to the next step.

Preparation 3241-(Benzyloxy)-4-bromo-5-ethyl-2-fluorobenzene

To a solution of 4-bromo-5-ethyl-2-fluorophenol (**PCT Intl. Appl. 2013014567**, 3
 g, 13.69 mmol) in acetone (30 mL) was added benzyl bromide (2.57 g, 15.06 mmol) and
 the reaction heated to reflux with potassium carbonate (2.83 g, 20.54 mmol) for 18
 25 hours. The reaction was filtered, concentrated *in vacuo* and purified using silica gel
 column chromatography eluting with 5% EtOAc in hexanes to afford the title compound
 as a colorless oil (3.20 g, 76%). ¹H NMR (400 MHz, DMSO-d₆): δ ppm 1.13 (t, 3H), 2.62
 (q, 2H), 5.18 (s, 2H), 7.26-7.51 (m, 7H).

Preparation 325

2-(4-(Benzyloxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane

The title compound was prepared according to the methods described for
 5 **Preparations 323 and 324** using 4-bromo-2-fluoro-5-(2,2,2-trifluoroethyl)phenol
 (**Preparation 326**). Taken on directly to the next step.

Preparation 326

4-Bromo-2-fluoro-5-(2,2,2-trifluoroethyl)phenol

To a solution of 1-bromo-5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)benzene
 10 (**WO2013014567**, 88.5 g, 308.31 mmol) at 0 °C in DCM (2000 mL), was added boron
 tribromide (204.56 mL, 2158.17 mmol) and the reaction was stirred at room temperature
 for 18 hours. The reaction was quenched by the addition of cold water dropwise at 0°C.
 The organic layer was separated, the aqueous extracts washed twice with DCM, the
 organic extracts combined, washed with brine, dried, concentrated *in vacuo* and
 15 triturated with pentane to afford the title compound as a white solid (78 g, 93%). ¹H
 NMR (400 MHz, DMSO-d₆): δ ppm 3.66-3.74 (m, 2H), 7.10 (d, 1H), 7.52 (d, 1H), 10.50
 (br s, 1H).

Preparation 327

N-(2-(Benzyloxy)ethyl)-N-(2-(((6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)-
 20 methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-
 yl)amino)methyl)phenyl)methanesulfonamide

A solution of 6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-
 ((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl
 trifluoromethanesulfonate (**Preparation 331**, 100 mg, 0.15 mmol), N-(2-
 25 (aminomethyl)phenyl)-N-(2-(benzyloxy)ethyl)methanesulfonamide hydrochloride
 (**Preparation 366**, 64 mg, 0.225 mmol) and triethylamine (62 µL, 0.45 mmol) in DMF (2
 mL) was heated to between 80-90 °C for 36 hours. The reaction was cooled and
 partitioned between EtOAc (50 mL) and water (50 mL). The organic layer was collected,
 further washed with water and brine, dried over sodium sulfate and concentrated *in*
 30 *vacuo*. The residue was purified using silica gel column chromatography eluting with
 EtOAc in heptanes to afford the title compound (51 mg, 40%). ¹H NMR (400MHz,
 CDCl₃): δ ppm 0.00 (s, 9H), 0.50 (s, 9H), 0.95 (m, 3H), 1.05 (m, 3H), 1.15 (m, 3H), 2.80
 (m, 2H), 3.10 (s, 3H), 3.50 (m, 1H), 3.60 (m, 4H), 3.80 (m, 1H), 3.90 (m, 2H), 4.05-4.20

(m, 1H), 4.60 (m, 2H), 5.15 (m, 1H), 5.20 (s, 2H), 5.60 (s, 2H), 6.15 (m, 1H), 6.80 (s, 1H), 7.10-7.40 (m, 10H), 7.70 (m, 1H), 7.90 (s, 1H). MS m/z 851 [M+H]⁺

Preparation 328

5 6-(2-Ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate

Triflic anhydride (0.21 mL, 1.25 mmol) was added dropwise to a solution of 6-(2-ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-ol (**Preparation 329**, 495 mg, 0.96 mmol) and pyridine (0.34 mL, 4.2 mmol) in DCM (5 mL) at 0°C. The reaction was stirred at room
10 temperature for 4 hours. The reaction was diluted with water (45 mL), acidified to pH=3 with citric acid and extracted with EtOAc (2 x 45 mL). The organic layers were combined, washed with a dilute solution of citric acid at pH=3 (45 mL), saturated aqueous NaHCO₃ solution, brine, dried over sodium sulfate and concentrated *in vacuo* to afford the title compound that was used directly in the next reaction.

15 **Preparation 329**

6-(2-Ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-ol

To a solution of 6-(2-ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 333**, 8 g, 15.5 mmol) in THF (160 mL) was added TEA (3.13 g, 31 mmol) dropwise, followed by
20 the addition of acetic anhydride (23.7 g, 232.5 mmol) dropwise at room temperature. The reaction was heated to 65 °C for 18 hours. The reaction was cooled and quenched by the addition of saturated aqueous NaHCO₃ solution (60 mL), and stirred for 10 hours. The reaction was diluted with water and extracted into EtOAc. The organic layer was
25 collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography to afford the title compound as an oil (5 g, 63%). ¹H NMR (400MHz, CDCl₃): δ ppm -0.06 (s, 9H), 0.0 (s, 9H), 0.84-0.88 (m, 2H), 0.94-0.98 (m, 2H), 1.12-1.16 (m, 3H), 2.62-2.64 (m, 2H), 3.55-3.59 (m, 2H), 3.75-3.79 (m, 2H), 5.26 (s, 2H), 5.57 (s, 2H), 6.42 (s, 1H), 6.92-6.96 (m,
30 1H), 6.99 (s, 1H), 7.21-7.25 (m, 1H), 8.14 (s, 1H), 9.15 (s, 1H). MS m/z 516 [M+H]⁺

Preparation 330

6-(2-Ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate

The title compound was prepared according to the methods described for Preparations 328 and 329 using 6-(2-ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (Preparation 332). ¹H NMR (400MHz, DMSO-d₆): δ ppm -0.01 (s, 9H), 0.91 (t, 2H), 1.04 (t, 3H), 1.59 (m, 2H), 1.71 (m, 1H), 2.30 (m, 2H), 2.37 (m, 2H), 2.70 (m, 2H), 3.78 (t, 3H), 3.90 (d, 1H), 5.36 (s, 2H), 6.05 (d, 1H), 7.25 (d, 1H), 7.35 (d, 1H), 8.10 (s, 1H), 8.52 (s, 1H).

Preparation 331

6-(2-Ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl trifluoromethanesulfonate

The title compound was prepared according to the methods described for Preparations 328 and 329 using 6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine 5-oxide (Preparation 335). Used directly in the next reaction.

Preparation 332

6-(2-Ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine 5-oxide

To a stirred solution of 6-[2-ethyl-5-fluoro-4-([2-(trimethylsilyl)ethoxy]methoxy)phenyl]-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (Preparation 147, 24 g, 50.88 mmol) in anhydrous DCM (300 mL) was added mCPBA (33.52 g, 117 mmol) and the reaction was stirred at room temperature for 18 hours. The reaction was quenched with saturated aqueous NaHCO₃ solution and extracted into DCM. The organic layer was collected, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 10% heptanes in EtOAc to afford the title compound as a yellow solid (14.5 g, 58%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 0.01 (s, 9H), 0.91-0.95 (t, 2H), 1.00-1.01 (t, 3H), 1.56 (s, 2H), 1.66-1.69 (m, 1H), 1.95-1.98 (m, 2H), 2.28-2.36 (m, 3H), 3.69-3.80 (m, 3H), 3.71-3.80 (m, 3H), 3.86 (d, 1H), 5.34 (s, 2H), 5.94 (d, 1H), 7.16-7.23 (m, 2H), 7.94 (s, 1H), 8.20 (s, 1H), 8.91 (s, 1H). MS m/z 488 [M+H]⁺

Preparation 333

6-(2-Ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine 5-oxide

To a solution of 6-(2-ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine (Preparation 336, 7 g, 14

mmol) in DCM (100 mL) was added m-CPBA (5.6 g, 28 mmol) at room temperature, and the reaction was stirred for 5 hours. The reaction was washed with 10% aqueous NaHSO₃ solution and saturated aqueous NaHCO₃ solution. The organic layer was separated, dried over sodium sulfate and concentrated *in vacuo* to afford the title compound (7 g, 97%). ¹H NMR (400MHz, CDCl₃): δ ppm 0.01 (s, 9H), 0.09 (s, 9H), 0.92-0.96 (m, 2H), 1.03-1.07 (m, 2H), 1.17-1.21 (m, 3H), 2.41-2.56 (m, 1H), 2.68-2.81 (m, 1H), 3.62-3.66 (m, 2H), 3.82-3.87 (m, 2H), 5.33 (s, 2H), 5.75 (s, 2H), 6.99-7.08 (m, 1H), 7.12 (m, 1H), 7.24 (d, 1H), 7.57 (d, 1H), 7.57 (d, 1H), 8.12 (s, 1H). MS m/z 516 [M+H]⁺

Preparation 334

6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine 5-oxide

The title compound was prepared according to the method described for **Preparation 111** using 6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 337**). MS m/z 472 [M+H]⁺

Preparation 335

6-(2-Ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine 5-oxide

The title compound was prepared according to the method described for **Preparation 111** using 6-(2-ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 338**). MS m/z 534 [M+H]⁺

Preparation 336

6-(2-Ethyl-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine

To a solution of 6-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine

(**Preparation 340**, 6.7 g, 23.67 mmol) in DMSO (120 mL) was added (2-((3-ethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)methoxy)ethyl)trimethylsilane (**Preparation 343**, 9.8 g, 26.03 mmol), potassium phosphate (18.88 g, 71.01 mmol) and water (12 mL) at room temperature. Pd(PPh₃)₄ (2.7 g, 2.3 mmol) was added, the reaction degassed under vacuum and refilled with nitrogen, and heated to 100 °C for 18 hours. The reaction was poured into ice water (200 mL) and extracted with

EtOAc. The organic layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography to afford the title compound as a yellow oil (6 g, 51%). Taken on directly to the next step.

Preparation 337

5 6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine

The title compound was prepared according to the method described for **Preparation 336** using 2-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**PCT Intl. Appl. 2013014567**) and 6-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 340**). ¹H NMR (400MHz, CDCl₃): δ ppm 0.00 (s, 9H), 0.95 (m, 2H), 3.62 (m, 2H), 3.82 (q, 2H), 4.02 (s, 10 3H), 5.80 (s, 2H), 7.10 (m, 1H), 7.30 (m, 1H), 7.60 (s, 1H), 8.25 (s, 1H), 9.12 (s, 1H).

Preparation 338

15 6-(2-Ethyl-5-fluoro-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine

The title compound was prepared according to the method described for **Preparation 336** using (2-([2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-ethylphenoxy] methoxy)ethyl)(trimethyl)-silane (**PCT Intl. Appl. WO2013014567A1**) and 6-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 20 340**). ¹H NMR (400MHz, CDCl₃): δ ppm 0.10 (s, 9H), 0.10 (s, 9H), 0.93-1.00 (m, 2H), 1.04-1.08 (m, 2H), 1.13-1.22 (m, 3H), 2.76 (q, 2H), 3.66 (m, 2H), 3.91 (m, 2H), 5.38 (s, 2H), 5.82 (s, 2H), 7.21-7.25 (m, 2H), 7.56 (s, 1H), 8.24 (s, 1H), 9.21 (s, 1H). MS m/z 518 [M+H]⁺

Preparation 339

25 N-(2-(((6-Chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

The title compound was prepared according to the method described for **Preparation 312** using 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine (**Preparation 341**) and N-[2-(aminomethyl)phenyl]-N-methylmethanesulfonamide (**PCT Intl. Appl. 2010058846**). ¹H NMR (400MHz, DMSO-d₆): δ ppm 1.55 (m, 2H), 1.69 (m, 1H), 1.88-2.00 (m, 2H), 2.29 (m, 1H), 3.04 (s, 1H), 3.27 (s, 3H), 3.69 (m, 1H), 3.85 (m, 1H), 4.55 (br m, 1H), 4.91 (br m, 1H), 5.67 (m, 1H), 6.91 (s, 1H), 7.32- 30 7.52 (m, 4H), 8.24 (s, 1H), 8.29 (t, 1H). MS m/z 449 [M+H]⁺

Preparation 340**6-Chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridine**

To a solution of 6-chloro-1H-pyrazolo[4,3-c]pyridine (8.5 g, 55 mmol) in anhydrous THF (200 mL) was added NaH (60% dispersion in oil, 2.3 g, 58 mmol) at 0°C. After stirring at room temperature for 20 minutes, SEMCl (9.67 g, 58.06 mmol) was added dropwise at 0°C. The reaction was stirred at room temperature for 2 hours before quenching with water and extracting into EtOAc. The organic layer was separated, dried over Na₂SO₄ and concentrated *in vacuo*. The residue was purified by silica gel column chromatography to afford the title compound as a yellow oil (14 g, 90%). Taken on directly to the next step.

Preparation 341**4,6-Dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[4,3-c]pyridine**

The title compound was prepared according to the method described for **Preparation 149** using 4,6-dichloro-1H-pyrazolo[4,3-c]pyridine. ¹H NMR (400MHz, DMSO-d₆): δ ppm 1.58-1.61 (m, 3H), 1.97-2.03 (m, 2H), 2.31-2.34 (m, 1H), 3.76-3.80 (s, 1H), 3.84-3.91 (m, 1H), 5.92-5.95 (d, 1H), 8.07 (s, 1H), 8.46 (s, 1H). MS m/z 272 [M+H]⁺

Preparation 342**2-((4-Benzyloxy)-2-ethyl-6-methylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane**

To a solution of 3-ethyl-4-iodo-5-methylphenol (J. Med. Chem. (2005), 48(2), 586-592, 500 mg, 1.90 mmol) in acetone (20 mL) was added benzylbromide (1.43 mL, 2.86 mmol) and potassium carbonate (658 mg, 4.77 mol). The reaction was heated to 70 °C for 18 hours. The reaction was cooled, filtered and concentrated *in vacuo*. The residue was dissolved in EtOAc and washed with water, brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with hexanes. The residue was dissolved in anhydrous DMSO (1.6 mL) and bis(pinacolato)diboron (1032 mg, 4.06 mmol) and KOAc (543 mg, 5.54 mmol) were added. The reaction was purged under argon for 10 minutes before the addition of Pd(dppf)₂Cl₂ (135 mg, 0.18 mmol) followed by degassing for another 10 minutes and then heating to 80 °C for 18 hours. The reaction was cooled, concentrated *in vacuo* and suspended in EtOAc. The suspension was filtered through celite and the filtrate washed with water, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 5% EtOAc in hexanes to afford the title compound. ¹H NMR (400MHz, DMSO-d₆): δ ppm 1.10 (t,

3H), 1.30 (s, 12H), 2.28 (s, 3H), 2.60 (q, 2H), 5.06 (s, 2H), 6.63 (m, 2H), 7.31-7.44 (m, 5H).

Preparation 343

(2-((3-Ethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)methoxy)ethyl)trimethylsilane

To a solution of (2-((4-bromo-3-ethylphenoxy)methoxy)ethyl)trimethylsilane (**Preparation 344**, 300 mg, 0.9 mmol) in dioxane (5 mL) was added bispinacolatodiboron (276 mg, 1.09 mmol), Pd(PPh₃)₄ (105 mg, 0.09 mmol) and potassium phosphate (384 mg, 1.81 mmol) and the reaction was heated to 80 °C for 18 hours. The reaction was cooled and partitioned between water and EtOAc, eluted through a phase separation cartridge and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 0-50% DCM in heptanes to afford the title compound. Taken on directly to the next step.

Preparation 344

(2-((4-Bromo-3-ethylphenoxy)methoxy)ethyl)trimethylsilane

To a solution of 4-bromo-3-ethylphenol (9 g, 44.8 mmol) in DCM (100 mL) was added DIPEA (8.6 mL, 49.3 mmol) followed by SEMCI (8.73 mL, 49.3 mmol) and the reaction was stirred at room temperature for 18 hours. The reaction was washed with water, 1N aqueous HCl solution and saturated aqueous sodium hydrogen carbonate solution, brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 3% EtOAc in hexanes to afford the title compound. ¹H NMR (400MHz, CDCl₃): δ ppm 0.00 (s, 9H), 0.90 (m, 2H), 1.25 (m, 3H), 2.75 (m, 2H), 3.75 (m, 2H), 5.20 (s, 2H), 6.80 (m, 1H), 7.00 (d, 1H), 7.40 (d, 1H).

Preparation 345

6-(4-Methyl-1H-imidazol-1-yl)-1,2,3,4-tetrahydroisoquinoline hydrochloride

To a solution of 6-fluoro-3,4-dihydro-2H-isoquinolin-1-one (13 g, 79 mmol) in DMSO (150 mL) was added 4-methylimidazole (7.8 g, 95 mmol) followed by cesium carbonate (38 g, 118.5 mmol) and the reaction was heated to 125 °C for 18 hours. The reaction was cooled and extracted into chloroform/isopropanol (v:v 3:1, 500 mL) three times. The organic layers were combined, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. A portion of the residue (9 g, 39.6 mmol) was dissolved in THF and cooled to 0°C. LiAlH₄ (3 g, 79.2 mmol) was added portionwise, and the reaction heated to 60 °C for 18 hours. The reaction was cooled and quenched by the

addition of 10% NaOH solution (6 mL), before filtration and concentration *in vacuo*. The residue was purified by silica gel column chromatography eluting with 30-100% EtOAc in petroleum ether followed by the addition of 2N HCl in EtOAc. The resulting precipitate was filtered to afford the title compound as the hydrochloride salt (11.6 g, 42%). ¹H
 5 NMR (400MHz, DMSO-d₆): δ ppm 2.34 (s, 3H), 3.08 (m, 2H), 3.37 (m, 2H), 4.30 (m, 2H), 7.48 (m, 1H), 7.62-7.68 (m, 2H), 8.00 (s, 1H), 9.61 (s, 1H), 9.91 (br s, 2H).

Preparation 346

N-(2-(Pyrrolidin-1-yl)ethyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide

To a solution of 1,2,3,4-tetrahydro-2-(2,2,2-trifluoroacetyl)-7-isoquinoline sulfonyl
 10 chloride (400 mg, 1.2 mmol) in MeOH (5 mL) was added 2-(pyrrolidin-1-yl)ethanamine in excess and the reaction stirred at room temperature for 30 minutes. Water (1 mL) followed by potassium carbonate (150 mg, 1.4 mmol) were added and the reaction stirred at room temperature for 18 hours. The reaction was concentrated *in vacuo* and the residue dissolved in DCM. The suspension was filtered and the filtrate purified by
 15 silica gel column chromatography eluting with 10-100% (90:10:1 DCM:MeOH:NH₃) in DCM to afford the title compound (130 mg, 35%). MS m/z 310 [M+H]⁺

Preparation 347

N-Benzyl-1,2,3,4-tetrahydroisoquinoline-5-carboxamide hydrochloride

To a solution of 2-(tert-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinoline-5-
 20 carboxylic acid (200 mg, 0.721 mmol) and DIPEA (87 μL, 0.793 mmol) in DCM (10 mL) was added HBTU (301 mg, 0.793 mmol) followed by a solution of benzylamine (151 μL 0.865 mmol) in DCM (5 mL) and the reaction was stirred at room temperature for 72 hours. The reaction was washed with water (1 mL), 1N HCl (aq) (1 mL) and 1N NaOH (aq) (1 mL). The organic layer was dried over sodium sulfate and concentrated *in*
 25 *vacuo*. The residue was dissolved in MeOH (5 mL) and 4N HCl in dioxane (3 mL) was added. The reaction was stirred at room temperature for 18 hours. The reaction was concentrated *in vacuo* and triturated with diethylether to afford the title compound as the hydrochloride salt (200 mg, quant.).

¹H NMR (400MHz, DMSO-d₆): δ ppm 3.09 (t, 2H), 3.33 (m, 2H), 4.28 (t, 2H), 4.44
 30 (d, 2H), 5.19 (s, 2H), 5.95 (d, 1H), 7.23-7.42 (m, 8H), 8.93 (t, 1H), 9.49 (br s, 2H).

Preparation 348

3-(Azetidin-3-yloxy)-4-chlorobenzonitrile hydrochloride

To a solution of 1-benzhydryl-3-azetidiny methanesulfonate (44.6 g, 0.147 mol) and 2-chloro-5-cyanophenol (22.6 g, 0.147 mol) in MeCN (600 mL) was added Cs₂CO₃

(62.3 g, 0.19 mol). The reaction was stirred at 80 °C for 24 hours. The reaction was filtered, and the filtrate was concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 5-20% EtOAc in petroleum ether. The residue was dissolved in dichloroethane (550 mL) and potassium carbonate (66.4 g, 0.48 mmol) followed by ACE-Cl (20.5 g, 0.14 mmol) was added. The reaction was heated to reflux for 2 hours. The reaction was concentrated *in vacuo* and the residue was recrystallized with MeOH to afford the title compound as the hydrochloride salt (13.8 g, 59%). ¹H NMR (400MHz, MeOD): δ ppm 4.21-4.25 (m, 2H), 4.61-4.66 (m, 2H), 5.25-5.31 (m, 1H), 7.30 (s, 1H), 7.40-7.43 (d, 1H), 7.63-7.65 (d, 1H).

10

Preparation 349**N-(2-(Aminomethyl)phenyl)-N-propylmethanesulfonamide trifluoroacetate**

To a solution of tert-butyl 2-(N-propylmethanesulfonamido)benzylcarbamate (**Preparation 351**, 265 mg, 0.77 mmol) in DCM (2 mL) was added TFA (0.5 mL) and the reaction stirred at room temperature for 1 hour. The reaction was diluted with DCM and washed with a 1:1 mixture of 880 NH₃ in water (20 mL). The organic layer was collected, dried over magnesium sulfate and concentrated *in vacuo* to afford the title compound as the trifluoroacetate salt (172 mg, 92%). ¹H NMR (400MHz, CDCl₃): δ ppm 0.89 (t, 3H), 1.41-1.55 (m, 2H), 2.96 (s, 3H), 3.39-3.46 (m, 1H), 3.62-3.69 (m, 1H), 3.83-3.87 (m, 1H), 4.10-4.20 (m, 1H), 7.19-7.22 (m, 1H), 7.28-7.32 (m, 1H), 7.36-7.40 (m, 1H), 7.57 (d, 1H).

20

Preparation 350**N-(2-(Aminomethyl)phenyl)-N-butylmethanesulfonamide trifluoroacetate**

The title compound was prepared according to the method described by **Preparation 349** using tert-butyl 2-(N-butylmethanesulfonamido)benzylcarbamate (**Preparation 354**) and isolated as the trifluoroacetate salt. ¹H NMR (400MHz, CDCl₃): δ ppm 0.87 (t, 3H), 1.26-1.52 (m, 4H), 2.95 (s, 3H), 3.41-3.48 (m, 1H), 3.67-3.75 (m, 1H), 3.83-3.86 (m, 1H), 4.10-4.14 (m, 1H), 7.19-7.21 (m, 1H), 7.28-7.32 (m, 1H), 7.37-7.41 (m, 1H), 7.58 (d, 1H).

25

Preparation 351**tert-Butyl 2-(N-propylmethanesulfonamido)benzylcarbamate**

30

The title compound was prepared according to the method described by **Preparation 213** using N-(2-cyanophenyl)-N-propylmethanesulfonamide (**Preparation 352**). Taken on directly to the next step.

Preparation 352**N-(2-Cyanophenyl)-N-propylmethanesulfonamide**

To a solution of N-(2-cyanophenyl)methanesulfonamide (**Preparation 223**, 500 mg, 2.55 mmol) in NMP (10 mL) was added sodium hydride (148 mg, 3.83 mmol) and the reaction stirred for 30 minutes at room temperature. Propyl iodide (1.74 mL, 3.83 mmol) was added and the reaction was stirred at room temperature for 18 hours. The reaction was quenched by the addition of water and extracted into EtOAc. The organic layer was collected, washed with water, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 30% EtOAc in heptanes to afford the title compound (505 mg, 83%). ¹H NMR (400MHz, CDCl₃): δ ppm 0.94 (t, 3H), 1.48-1.60 (m, 2H), 3.11 (s, 3H), 3.71 (t, 2H), 7.46-7.54 (m, 2H), 7.65-7.69 (m, 1H), 7.72-7.74 (m, 1H).

Preparation 353**N-Methyl-N-(2-(((2-morpholinoethyl)amino)methyl)phenyl)methanesulfonamide**

Sodium hydride (76 mg, 1.92 mmol) was added to a solution of tert-butyl 2-(N-methylmethylsulfonamido)benzylcarbamate (**WO 2010058846**, 200 mg, 0.64 mmol) in NMP and the reaction was stirred at 0 °C for 30 minutes. 2-morpholinoethanamine (226 mg, 0.96 mL) was added and the reaction stirred at room temperature for 18 hours. The reaction was quenched by the addition of water and extracted into EtOAc. The organic layer was collected, dried and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 100:10:1 DCM:MeOH:TEA. The residue was dissolved in DCM (2 mL) and TFA (1 mL) was added. The reaction was stirred at room temperature for 1 hour. The reaction was concentrated *in vacuo* to afford the title compound as the trifluoroacetate salt. MS m/z 328 [M+H]⁺

25

The following Preparations (**Preparations 354 – 359**) were prepared according to the method described by **Preparation 353** using the appropriate sulphonamide and alkyl halide as described below. The compounds were isolated according to the described experimental or by dissolving in DCM (20 mL) and washing with a 1:1 mixture of ammonium hydroxide:water. The organic layer was collected, dried over magnesium sulfate and concentrated *in vacuo* to afford the title compound that was used in the next reaction directly.

30

<u>Preparation Number</u> <u>Preparation No.</u>	Name	SM	Data
354	N-butyl-N-(2-((methylamino)methyl)phenyl)methanesulfonamide	tert-butyl 2-(N-butylmethylsulfonamido)benzylcarbamate (Preparation 364) and methyl iodide.	¹ H NMR (400MHz, CDCl ₃): δ ppm 0.88 (t, 3H), 1.27-1.58 (m, 4H), 2.50 (s, 3H), 2.95 (s, 3H), 3.44-3.50 (m, 1H), 3.67-3.76 (m, 2H), 3.95-4.00 (m, 1H), 7.20-7.23 (m, 1H), 7.28-7.32 (m, 1H), 7.34-7.38 (m, 1H), 7.61 (d, 1H).
355	N-ethyl-N-(4-methyl-2-((methylamino)methyl)phenyl)methanesulfonamide	tert-butyl 2-(N-ethylmethylsulfonamido)-5-methylbenzylcarbamate (Preparation 361) and methyl iodide.	Taken on directly to the next step.
356	N-(2-((ethylamino)methyl)-4-methylphenyl)N-methylmethanesulfonamide	tert-butyl 5-methyl-2-(N-methylmethylsulfonamido)benzylcarbamate (Preparation 360) and ethyl iodide.	¹ H NMR (400MHz, CDCl ₃): δ ppm 1.21 (t, 3H), 2.32 (s, 3H), 2.82 (m, 2H), 2.99 (s, 3H), 3.21 (s, 3H), 3.95 (s, 2H), 7.07 (m, 2H), 7.40 (s, 1H).
357	N-methyl-N-(4-methyl-2-((propylamino)methyl)phenyl)methanesulfonamide	tert-butyl 5-methyl-2-(N-methylmethylsulfonamido)benzylcarbamate (Preparation 360) and propyl iodide.	¹ H NMR (400MHz, CDCl ₃): δ ppm 0.95 (t, 3H), 1.60 (q, 2H), 2.83 (d, 3H), 2.31 (s, 3H), 2.95 (s, 3H), 3.20 (s, 3H), 3.95 (s, 2H), 7.07 (m, 2H), 7.40 (s, 1H).
358	N-ethyl-N-(2-((methylamino)methyl)phenyl)methanesulfonamide	tert-butyl 2-(N-ethylmethylsulfonamido)benzylcarbamate (Preparation 362) and methyl iodide.	Taken on directly to the next step.
359	N-methyl-N-(2-((methylamino)methyl)phenyl)methanesulfonamide	tert-butyl 2-(N-methylmethylsulfonamido)benzylcarbamate (Preparation 363) and methyl iodide.	Taken on directly to the next step.

The following Preparations (**Preparations 360 – 364**) were prepared according to the methods described by **Preparations 351 and 352** using the appropriate sulphonamide and alkyl halide as described below:

<u>Preparation Number</u> Prep-No	Name	SM	Data
360	tert-butyl 5-methyl-2-(N-methylmethanesulfonamido)benzylcarbamate	N-(2-cyano-4-methylphenyl)-methanesulfonamide (Preparation 365) and methyl iodide.	Taken on directly to the next step.
361	tert-butyl 2-(N-ethyl-methylsulfonamido)-5-methylbenzylcarbamate	N-(2-cyano-4-methylphenyl)-methanesulfonamide (Preparation 365) and ethyl iodide.	MS m/z 343 [M+H] ⁺ Taken on directly to the next step.
362	tert-butyl 2-(N-ethyl-methylsulfonamido)benzylcarbamate	N-(2-cyanophenyl)methanesulfonamide (Preparation 223) and ethyl iodide.	MS m/z 329 [M+H] ⁺ Taken on directly to the next step.
363	tert-butyl 2-(N-methylmethanesulfonamido)benzylcarbamate	N-(2-cyanophenyl)methanesulfonamide (Preparation 223) and methyl iodide	MS m/z 315 [M+H] ⁺ Taken on directly to the next step.
364	tert-butyl 2-(N-butylmethanesulfonamido)benzylcarbamate	N-(2-cyanophenyl)methanesulfonamide (Preparation 223) and butyl iodide	¹ H NMR (400MHz, CDCl ₃): δ ppm 0.87 (t, 3H), 1.26-1.55 (m, 13H), 2.94 (s, 3H), 3.39-3.46 (m, 1H), 3.71-3.78 (m, 1H), 4.30-4.35 (m, 1H), 4.56-4.61 (m, 1H), 5.35 (br s, 1H), 7.19-7.21 (m, 1H), 7.30-7.39 (m, 2H), 7.60 (d, 1H).

Preparation 365**N-(2-Cyano-4-methylphenyl)methanesulfonamide**

The title compound was prepared according to the method described for

5 **Preparation 223** using methyl iodide and N-(4-methyl-2-cyanophenyl)-N-

(methylsulfonyl)methanesulfonamide (**Preparation 242**). Taken on directly to the next step.

Preparation 366

N-(2-(Aminomethyl)phenyl)-N-(2-(benzyloxy)ethyl)methanesulfonamide hydrochloride

The title compound was prepared according to the methods described for **Preparations 222, 213 and 211** using 2-(benzyloxy)ethanol and N-(2-cyanophenyl)methanesulfonamide. MS m/z 335 [M+H]⁺

Preparation 367

N-(2-(((3,4-Dimethoxyphenethyl)amino)methyl)phenyl)-N-methylmethanesulfonamide

To a solution of 2-(3,4-dimethoxyphenyl)-N-(2-(N-methylmethylsulfonamido)benzyl)acetamide (**Preparation 368**, 800 mg, 2.03 mmol) in THF (15 mL) was added borane-dimethylsulfide (2M in THF, 2.55 mL, 5.10 mmol) and the reaction was heated to reflux for 2.5 hours. The reaction was cooled, concentrated *in vacuo* and the residue dissolved in methanol (12 mL). The solution was treated with 6N HCl (8 mL) and heated to reflux for 2 hours. The reaction was concentrated *in vacuo* and the residue basified with 3N NaOH solution. The aqueous layer was extracted into 10% MeOH/DCM, the organic extracts were washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified using silica gel column chromatography eluting with 7% MeOH in DCM to afford the title compound (390 mg, 51%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 2.63-2.71 (m, 4H), 3.04 (s, 3H), 3.17 (s, 3H), 3.71 (s, 6H), 3.80 (br s, 2H), 6.68-6.83 (m, 3H), 7.28-7.51 (m, 4H). MS m/z 379 [M+H]⁺

Preparation 368

2-(3,4-Dimethoxyphenyl)-N-(2-(N-methylmethylsulfonamido)benzyl)acetamide

To a solution of N-[2-(aminomethyl)phenyl]-N-methylmethanesulfonamide (PCT Intl. Appl. 2010058846, 1 g, 3.64 mmol) and 2-(3,4-dimethoxyphenyl)acetic acid (786 mg, 4.00 mmol) in THF (20 mL) was added propylphosphonic anhydride (2.9 g, 9.11 mmol) followed by DIPEA (2.21 mL, 12.68 mmol) and the reaction was stirred at room temperature for 14 hours. The reaction was concentrated *in vacuo* and the residue was partitioned between EtOAc and saturated aqueous sodium bicarbonate solution. The organic layer was collected, washed with brine, dried, concentrated *in vacuo* and purified using silica gel column chromatography eluting with 4% MeOH in DCM to afford

the title compound as a white solid (540 mg, 38%). ¹H NMR (400MHz, DMSO-d₆): δ ppm 3.05 (s, 3H), 3.14 (s, 3H), 3.40 (s, 2H), 3.71 (s, 6H), 4.26 (br s, 1H), 4.46 (br s, 1H), 6.77-6.88 (m, 3H), 7.26-7.34 (m, 3H), 7.45-7.47 (m, 1H), 8.34 (t, 1H). MS m/z 393 [M+H]⁺

5

Preparation 369

N-Methyl-N-(2-(((4-(methylsulfonamido)phenethyl)amino)methyl)phenyl)methanesulfonamide

The title compound was prepared according to the methods described for **Preparations 368 and 367** using 2-(4-(methylsulfonamido)phenyl)acetic acid. ¹H NMR (400MHz, DMSO-d₆): δ ppm 2.66-2.73 (m, 4H), 2.92 (s, 3H), 3.04 (s, 3H), 3.13 (s, 3H), 3.80 (br s, 2H), 7.09-7.17 (m, 4H), 7.29-7.34 (m, 2H), 7.42-7.49 (m, 2H), 9.52 (br s, 1H). MS m/z 412 [M+H]⁺

10

Preparation 370

N-(2-(((6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide

15

The title compound was prepared according to the method described for **Example 157** using N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 273**) and tert-butyl 2-iodo-6,7-dihydro-1H-imidazo[4,5-c]pyridine-5(4H)-carboxylate (**WO2013014567**) using HCl in dioxane for the deprotection step. MS m/z 690 [M+H]⁺

20

Preparation 371

N-(2-(((6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(5-(piperidin-4-yl)-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide

25

To a solution of N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(hydrazinecarbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide (**Preparation 375**, 450 mg, 0.59 mol) in butanol (2 mL) was added tert-butyl 4-cyanopiperidine-1-carboxylate (624 mg, 2.97 mmol) and the reaction was heated to 150 °C under microwave irradiation for 50 minutes. The reaction was cooled, filtered and concentrated *in vacuo*. The residue was purified using preparative HPLC. The residue was treated with TFA (2 mL) and stirred at room temperature for 30 minutes. The

30

reaction was concentrated *in vacuo*, dissolved in MeOH (5 mL) and cooled in ice water. Ethylene diamine was added dropwise until the solution was basic, with stirring for 1 hour. The solution was concentrated *in vacuo* and extracted into 20% IPA in DCM. The organic layer was washed with water, dried over sodium sulfate and concentrated *in vacuo* to afford the title compound that was used directly in the next reaction. MS m/z 719 [M+H]⁺

Preparation 372

N-(2-(((6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(5-(piperidin-4-yl)-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

10

The title compound was prepared according to the method described by **Preparation 371** using N-(2-(((3-cyano-6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide (**Preparation 377**) and tert-butyl 4-(hydrazinecarbonyl)piperidine-1-carboxylate in the presence of potassium carbonate. MS m/z 689 [M+H]⁺

15

Preparation 373

6-[5-Fluoro-2-ethyl-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-4-{[2-[methyl(methylsulfonyl)amino]benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxylic acid

20

The title compound may be prepared according to the method described for **Preparation 11** using N-[2-({[6-(2-ethyl-5-fluoro-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl)-3-iodo-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide (**Preparation 79**). Taken on directly to the next step.

25

Preparation 374

N-(2-(((6-(5-Fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-(hydrazinecarbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

30

To a solution of 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-{[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-4-{[2-[methyl(methylsulfonyl)amino]benzyl]amino}-1-{[2-(trimethylsilyl)ethoxy]methyl}-1H-pyrazolo[4,3-c]pyridine-3-carboxylic acid (**Preparation 12**, 0.55 g, 0.66 mmol) in

MeOH/Toluene (15 mL) was added 2M trimethylsilyldiazomethane in THF (0.997 mL, 1.99 mmol) dropwise at 0°C. The reaction was stirred for 2 hours at room temperature. The reaction was concentrated *in vacuo* and the residue was purified using silica gel column chromatography eluting with 8% MeOH in DCM. The residue was dissolved in
 5 MeOH (5 mL) and hydrazine monohydrate (40.12 mg, 0.80 mmol) was added. The reaction was heated to reflux for 18 hours. The reaction was concentrated *in vacuo* and the residue was purified using neutral alumina column chromatography eluting with 50% EtOAc in hexanes to afford the title compound (297 mg, 66%). MS m/z 842 [M+H]⁺

Preparation 375

10 N-(2-(((6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(hydrazinecarbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)-4-methoxyphenyl)-N-methylmethanesulfonamide

The title compound was prepared according to the method described by **Preparation 374** using 6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-methoxy-2-(N-methylmethyl-sulfonamido)benzyl)amino)-1-((2-(trimethylsilyl)ethoxy)-
 15 methyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxylic acid (**Preparation 269**). MS m/z 757 [M+H]⁺

Preparation 376

20 N-(2-(((6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-(hydrazinecarbonyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

The - was prepared according to the methods described for **Preparations 11 and 374** using N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-tri-fluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methyl-methanesulfonamide (**Preparation 378**). ¹H NMR (400MHz, DMSO-d₆): δ ppm -
 25 0.11 (s, 9H), 0.83 (m, 2H), 1.23 (br s, 2H), 3.05 (s, 3H), 3.11 (s, 3H), 3.57 (m, 2H), 3.77 (m, 2H), 3.86 (s, 3H), 4.68 (m, 2H), 4.80 (br m, 1H), 4.90 (br m, 1H), 5.72 (s, 2H), 7.08 (s, 1H), 7.20 -7.51 (m, 6H), 9.68 (t, 1H), 10.17 (m, 1H). MS m/z 726 [M+H]⁺

Preparation 377

30 N-(2-(((3-Cyano-6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

To a solution of N-(2-(((6-(5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-

phenyl)-N-methylmethanesulfonamide (**Preparation 378**, 1.2 g, 1.51 mmol) in DMF (10 mL), was added zinc cyanide (0.19 g, 1.66 mmol) and Pd(PPh₃)₄ (0.05 mg, 0.04 mmol). The reaction was degassed with nitrogen and heated to 120 °C under microwave irradiation for 20 minutes. The reaction was quenched with water and extracted into ethyl acetate. The organic extracts were dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 48% EtOAc in hexanes to afford the title compound (610 mg, 58%). ¹H NMR (400MHz, DMSO-d₆): δ ppm -0.07 (s, 9H), 0.86 (t, 2H), 3.08 (s, 3H), 3.17 (s, 3H), 3.65 (t, 2H), 3.88 (s, 3H), 4.22 (m, 2H), 4.89 (br m, 1H), 5.00 (br m, 1H), 5.75 (s, 2H), 7.24-7.41 (m, 4H), 7.55-7.59 (m, 2H), 8.29 (t, 1H). MS m/z 694 [M+H]⁺

Preparation 378

N-(2-(((6-(5-Fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)phenyl)-N-methylmethanesulfonamide

The title compound was prepared according to the method described for **Preparation 61** using 6-[5-fluoro-4-methoxy-2-(2,2,2-trifluoroethyl)phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]-methyl]-1H-pyrazolo[4,3-c]pyridine (**Preparation 111**) and 4-nitrophenyl {2-[methyl(methylsulfonyl)amino]benzyl}carbamate (**Preparation 166**). MS m/z 794 [M+H]⁺

Preparation 379

6-(5-Fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-N-(2-(methylthio)ethyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[4,3-c]pyridin-4-amine

The title compound was prepared according to the method described for **Preparation 61** using 6-[5-fluoro-2-(2,2,2-trifluoroethyl)-4-[[2-(trimethylsilyl)ethoxy]methoxy}phenyl]-3-iodo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1H-pyrazolo[4,3-c]pyridine 5-oxide (**Preparation 114**) and 4-nitrophenyl (2-(methylthio)ethyl)carbamate (**Preparation 385**). MS m/z 787 [M+H]⁺

Preparation 380

Racemic 3-(((6-(5-Fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-3-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-2-methylpropan-1-ol

The title compound was prepared according to the method described for **Preparation 131** using racemic N-(3-((tert-butyldimethylsilyl)oxy)-2-methylpropyl)-6-(4-

((tert-butyldimethylsilyl)-oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine (**Preparation 381**). MS m/z 786 [M+H]⁺

Preparation 381

Racemic N-(3-((tert-Butyldimethylsilyl)oxy)-2-methylpropyl)-6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-amine

The title compound was prepared according to the method described for **Preparation 137** using racemic N-(3-((tert-butyldimethylsilyl)oxy)-2-methylpropyl)-6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine (**Preparation 382**). ¹H NMR (400MHz, DMSO-d₆): δ ppm -0.01 (s, 6H), 0.22 (s, 6H), 0.79 (m, 9H), 0.98 (m, 12H), 2.04 (m, 1H), 3.48 (m, 1H), 3.56 (m, 2H), 3.68 (m, 1H), 4.45 (m, 2H), 6.71 (m, 1H), 7.13 (m, 1H), 7.81 (m, 1H), 13.88 (s, 1H). MS m/z 754 [M+H]⁺

Preparation 382

Racemic N-(3-((tert-Butyldimethylsilyl)oxy)-2-methylpropyl)-6-(4-((tert-butyldimethylsilyl)oxy)-5-fluoro-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine

The title compound was prepared according to the methods described by **Preparations 142 and 141** using racemic 3-((6-(5-fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-2-methylpropan-1-ol (**Preparation 383**). MS m/z 626 [M-H]⁻

Preparation 383

Racemic 3-((6-(5-Fluoro-2-(2,2,2-trifluoroethyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-2-methylpropan-1-ol

The title compound was prepared according to the method described for **Preparation 299** using racemic 3-((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-2-methylpropan-1-ol (**Preparation 384**) and (2-([2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5-(2,2,2-trifluoroethyl)phenoxy]-methoxy)ethyl) (trimethyl)silane (**Preparation 150**). MS m/z 614 [M+H]⁺

Preparation 384

Racemic 3-((6-Chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)amino)-2-methylpropan-1-ol

The title compound was prepared according to the method described by **Preparation 299** using racemic 3-amino-2-methylpropan-1-ol. MS m/z 326 [M+H]⁺

Preparation 385

4-Nitrophenyl (2-(methylthio)ethyl)carbamate

5 The title compound was prepared according to the method described for **Preparation 156** using 2-(methylthio)ethanamine. Taken on directly to the next step.

Preparation 386

2-Bromo-4,5-dimethyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole

To solution of 4,5-dimethyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole
10 (**Preparation 387**, 270 mg, 1.19 mmol) in anhydrous THF (3 mL) at -78°C, was added butyllithium (0.54 mL, 1.31 mmol) dropwise. The reaction was kept at -78 °C for 15 minutes before the addition of carbon tetrabromide (474 mg, 1.43 mmol) in THF (2 mL). The reaction was warmed to room temperature before quenching with ammonium chloride and extracting into EtOAc. The organic layer was collected, washed with water,
15 brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 3% MeOH in DCM to afford the title compound as a colorless oil (220 mg, 60%). ¹H NMR (400MHz, DMSO-d₆): δ ppm - 0.01 (s, 9H), 0.87 (t, 2H), 2.01 (s, 3H), 2.14 (s, 3H), 3.51 (t, 2H), 5.18 (s, 2H).

Preparation 387

4,5-Dimethyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-imidazole

A suspension of NaH (124 mg, 3.12 mmol) in DMF (3 mL) was added a solution of 4,5-dimethyl-1H-imidazole (200 mg, 2.08 mmol) in DMF (2 mL) at 0°C. The suspension was stirred for 15 minutes before the dropwise addition of SEM chloride (0.44 mL, 2.49 mmol). The reaction was stirred at room temperature for 1 hour, then
25 partitioned between ethyl acetate and water. The combined organic extracts were washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel column chromatography eluting with 5% MeOH in DCM to afford the title compound as a colorless oil (270 mg, 57%). ¹H NMR (400MHz, DMSO-d₆): δ ppm -0.01 (s, 9H), 0.82 (t, 2H), 2.01 (s, 3H), 2.09 (s, 3H), 3.43 (t, 2H), 5.18 (s,
30 2H), 7.52 (s, 1H).

Biological Evaluation

JAK Caliper Enzyme Assay at 1mM ATP

Test article was solubilized in dimethyl sulfoxide (DMSO) to a stock concentration of 30 mM. An 11-point half log dilution series was created in DMSO with a top concentration of 600 μ M. The test compound plate also contained positive control wells containing a known inhibitor to define 100% inhibition and negative control wells containing DMSO to define no inhibition. The compound plates were diluted 1 to 60 resulting in a top final assay compound concentration of 10 μ M and a 2% DMSO concentration.

Test article and assay controls were added to a 384-well plate. Reaction mixtures contained 20 mM HEPES, pH 7.4, 10 mM magnesium chloride, 0.01% bovine serum albumin (BSA), 0.0005% TweenTM 20, 1 mM ATP and 1 μ M peptide substrate. The JAK1 and TYK2 assays contained 1 μ M of the IRStide peptide (5FAM-KKSRGDYMTMQID) and the JAK2 and JAK3 assays contained 1 μ M of the JAKtide peptide (FITC-KGGEEEEYFELVKK). The assays were initiated by the addition of 20 nM JAK1, 1 nM JAK2, 1 nM JAK3 or 1 nM TYK2 enzyme and were incubated at room temperature for three hours for JAK1, 60 minutes for JAK2, 75 minutes for JAK3 or 135 minutes for TYK2. Enzyme concentrations and incubation times were optimized for each new enzyme preps and were modified slightly over time to ensure 20%-30% phosphorylation. The assays were stopped with a final concentration of 10 mM EDTA, 0.1% Coating Reagent and 100 mM HEPES, pH=7.4. The assay plates were placed on a Caliper Life Science Lab Chip 3000 (LC3000) instrument, and each well was sampled using appropriate separation conditions to measure the unphosphorylated and phosphorylated peptide.

A549 Cell Assay: Inhibition of pSTAT3

An assay measuring the efficacy of JAK inhibitors on the functional response of recombinant human interferon γ (rhIFN γ) stimulated STAT-3 phosphorylation in the A549 human epithelial cell line.

81797008

Method

5 A549 cells (ATCC #CCL-185), were plated at 30,000 cells/well in 96 well flat bottomed tissue culture plates (BD#353072) in 200 μ L of growth medium (DMEM, Pfizer media prep, with 10% Fetal Bovine Serum, Sigma #F4135, 2 mM L-Glutamine, Pfizer media prep, 100 U/mL penicillin, Pfizer media prep, and 200 μ g/ml streptomycin, Pfizer media prep), and cultured at 37°C, 5% CO₂ incubator for 18 hours. Growth medium was removed by vacuum aspiration (V&P Scientific #vp187bp-60), and 90 μ L of pre-warmed assay medium (DMEM with 0.2% BSA, Miltenyi # 130-091-376) was added to each well

and incubated for 15 minutes at 37°C. 10 µL of vehicle control or test compound (final concentration range of 0.3 nM to 10 µM with 0.1% DMSO) was added to the cells. Plates were incubated at 37°C for 1 hour. After compound incubation, 10 µL of 220 ng/mL recombinant human IFN γ (R&D Systems #285-IF, final rhIFN γ concentration of 20 ng/mL) was added to the cells and plates were incubated for 30 minutes at 37°C. Wells containing A549 cells, medium with 0.1% DMSO and no rhIFN γ were used as background controls. After rhIFN γ stimulation, media was aspirated from each well and 35 µL /well of iced-cold MSD lysis buffer containing protease and phosphatase inhibitors from Phospho-STAT3 Tyr705 assay kit (Meso-Scale Discovery #K150DID) was added to each well. Plates were incubated at 4°C with shaking for 30 minutes. Cell lysates were assayed following the MSD Phospho-STAT3 Tyr705 assay kit protocol to detect pSTAT3.

Data were collected and transformed into percent inhibition and calculated using the following formula:

$$\% \text{ Inhibition} = \left(1 - \left(\frac{\text{Compound pSTAT3} - \text{Basal pSTAT3}}{\text{No Compound Control pSTAT3} - \text{Basal pSTAT3}} \right) \right) * 100$$

Data were graphically displayed as percent inhibition using GraphPad Prism 4.0, and IC₅₀ curves were fitted using a point to point analysis.

Human T cell assay: Inhibition of pSTAT5

An assay measuring the efficacy of JAK inhibitors on the functional response of recombinant human interleukin-2 (rhIL-2) stimulated STAT5 phosphorylation in isolated human T cells.

Method:

Human whole blood from individual donors was collected from the phlebotomy unit on-site. Peripheral venous blood (30-60 mL) from healthy volunteers of either sex was used as the source of T cells. T cell isolation from venous whole blood is routinely performed in a class II microbiological safety cabinet. Each sample was collected into between 3 and 6 10 mL Sodium Heparin Vacutainer tubes (BD #367874). The blood was poured into sterile 50 mL conicals (Corning #430828) and incubated with the T cell Rosette Sep Cocktail (Stemcell Technologies #15061) at 50 µL /mL antibody/blood ratio for 20 min. with shaking at room temperature. The blood/antibody mixture was then diluted 1:2 with PBS (Pfizer







media prep)/2%FBS (Sigma #F4135) and 30 mL of the mixture was layered onto 15 mL Ficoll-Hypaque (GE Healthcare #17-1440-03) in 50 mL conical tubes. The tubes were then centrifuged at 1200 xg for 20 min. at room temperature with no brake. Following centrifugation the T cells formed a buffy coat between the Ficoll-Hypaque and plasma layers. The plasma above the buffy coats was removed to within 5 mm of the buffy coat using a sterile Pasteur pipette. The buffy coats were then collected into fresh sterile 50 mL conical tubes containing 25 mL PBS/2%FBS (2 buffy coats per 50 mL conical). PBS/2%FBS was added to the buffy coat cells such that the final volume in the tube was 50 mL. The tubes were then centrifuged at 200 xg for 15 min. at room temperature. The supernatant was discarded and the pellet re-suspended in 10-20 mL of DMEM (Pfizer media prep) Assay media/0.2% BSA (Miltlenyi #130-091-376). A differential cell count was performed using a haemocytometer and cells were diluted to 1.1×10^6 T cells/ml in DMEM/0.2% BSA media. Compounds (10 mM-0.3 μ M) were diluted with Hanks Balanced Salt Solution (HBSS) (Sigma #H6648) at 1:100 dilution. Immediately following cell isolation and compound dilution, 90 μ L/well of T cells ($\sim 1 \times 10^6$ / ml) in assay medium (DMEM+0.2% BSA) was added to the VWR Deep well V bottom plate (#3906-520-300). 10 μ L/well of compound (final concentration range of 10 μ M - 0.3 nM with 0.1% DMSO) or 0.1% DMSO in HBSS as controls was added to the appropriate wells. Plates were incubated for 1 hour at 37°C, 5% CO₂ incubator. 10 μ L of 3.3 μ g/mL rhIL-2 (R&D Systems #202-IL) was added to the cells (300 ng/ml final assay concentration). Wells containing T cells, medium with 0.1% DMSO and no rhIL-2 were used as background controls. Plates were incubated for 15 min at 37°C. After rhIL-2 stimulation, 800 μ L of cold PBS/0.1%BSA was added and the plates were centrifuged at 1400 rpm for 5 min at 4°C. Supernatant was aspirated and 100 μ L of ice-cold MSD lysis buffer containing protease and phosphatase inhibitors from Phospho-STAT5a/b Tyr694 assay kit (Meso-Scale Discovery #K150IGD) was added to the cell pellet. Plates were shaken for 30 min. at 4°C and then frozen overnight. The following day, cell lysates were assayed using the MSD ELISA kit protocol for detection of pSTAT5.








Data were collected and transformed into percent inhibition and calculated using the following formula:








$$\% \text{ Inhibition} = \left(1 - \left(\frac{\text{Compound pSTAT5} - \text{Basal pSTAT5}}{\text{No Compound Control pSTAT5} - \text{Basal pSTAT5}} \right) \right) * 100$$








Data were graphically displayed as percent inhibition using GraphPad Prism 4.0, and IC_{50} curves were fitted using a point to point analysis.








Table 1. Data for JAK Caliper™ Kinase assays at 1mM ATP and Cell Based assays








		Caliper™ Assays at 1mM ATP				Cell Based Assays	
Ex.	Structure	JAK 1 IC ₅₀ (nM)	JAK2 IC ₅₀ (nM)	JAK3 IC ₅₀ (nM)	TYK2 IC ₅₀ (nM)	A549 cell assay: Inhibition of pSTAT3 IC ₅₀ (nM)	Human T cell assay: Inhibition of pSTAT5 IC ₅₀ (nM)
125		0.2	1.0	4.4	5.6		
126		8.1	23.5	35.1	668.3		
127		6.0	14.9	53.3	624.9		
131		<0.5	2.1	10.2	4.7	7.4	24.3
124		87.6	182.6	156.7	4234.5		
123		3.3	4.5	9.5	82.1		








128		3.6	12.6	30.5	514.3		
132		19.0	31.4	18.4	1293.1	256.5	206.4
111		13.8	52.4	193.6	1540.2		
112		12.1	82.9	67.5	5170.8		
113		10.7	26.3	42.4	723.1		
114		10.1	25.9	56.2	289.2		
115		4.3	10.1	16.8	229.0		








116		3.8	10.1	23.3	162.7		
117		7.2	5.0	41.6	425.0		
118		67.4	118.9	339.6	2730.0		
119		39.4	46.5	111.0	561.1		
120		10.8	127.8	89.2	3320.1		
121		9.7	16.9	40.8	302.1		
122		14.9	25.1	78.1	478.4		








137		<0.1	0.3	3.3	45.4		
138		0.6	2.6	14.2	4.8	5.4	
139		8.5	16.7	27.5	1992.6		
133		1.2	5.9	30.5	249.3	10.1	34.7
135		0.7	3.4	17.3	8.4	3.3	
145		1.0	4.5	16.2	15.8	15.7	157.3
141		19.5	43.2	61.6	7069.4		








140		18.1	53.5	131.1	1410.3		
147		1.1	4.7	18.5	49.4		
142		3.4	7.9	27.9	885.1		
143		40.4	43.6	121.9	4663.2		
144		55.3	70.7	185.9	8734.3		
146		0.9	5.1	19.1	49.9	7.6	62.3
148		1.8	6.3	22.8	145.6	14.9	








149		2.7	14.5	22.7	763.5		
150		21.6	48.9	77.8	4156.1		
19		0.9	5.9	30.9	4.8	11.9	20.2
134		<0.5	1.7	7.6	20.3		258.6
136		0.8	4.3	22.4	8.7	13.3	
155		1.2	5.0	15.5	61.0	30.1	
129		11.6	43.7	180.1	56.6	62.4	








130		43.8	154.9	1358.2	194.1	728.6	
5		2.1	7.5	31.9	7.0	1.7	
20		3.6	16.6	95.3	20.9	18.8	31.9
3		1.9	9.6	47.5	6.8	16.2	
6		3.1	15.8	84.9	15.3	23.4	
8		1.7	7.5	33.8	5.7	6.4	
7		1.7	9.2	49.0	6.5	5.7	








21		2.7	13.2	67.9	11.4	35.6	
9		1.7	10.8	60.0	7.2	14.9	
2		3.4	17.0	84.5	12.1	46.8	
10		4.7	19.0	93.1	17.9	26	
11		8.7	36.9	206.6	229.4	72.4	
14		4.2	20.5	99.7	43.2	20.7	
1		8.0	38.1	176.8	36.5	67.4	








15		13.6	55.0	342.4	618.3	92.4	
93		2.0	8.4	27.4	13.9	31.9	40.4
16		6.3	23.0	130.8	66.0	65.1	
18		4.0	17.4	73.0	31.0	97	
17		18.5	52.9	255.7	142.4	173.9	
13		5.3	26.6	118.0	35.7	96.7	
12		6.4	31.5	120.6	24.8	44.9	








4		1.7	6.2	26.6	15.3	29.7	
56		6.6	26.1	134.8	19.9	276.9	53.4
31		1.3	7.0	33.0	4.6	74	48.2
22		17.5	81.3	359.0	59.1	52.8	
23		5.8	25.7	110.7	83.2	51.4	
165		3.5	16.9	78.0	14.6	566.7	91.3
38		4.0	25.0	121.8	17.8	30.8	








39		32.3	48.2	278.5	320.3	60.9	
162		4.1	23.6	90.6	15.4	20	
157		3.4	15.0	76.5	11.7	26.2	
94		5.4	27.7	71.2	503.4	125.5	
158		5.3	27.8	114.0	33.3	31.3	
159		4.3	18.5	83.3	16.1	18.7	
163		32.9	150.2	491.0	355.9	98.6	








76		29.6	112.9	475.0	315.5	324.8	120.7
171		27.5	128.9	428.9	150.2	1575.2	1090.4
92		2.1	13.5	45.7	14.5	398.2	343.4
77		3.3	15.1	63.6	27.1	32.9	59.9
166		20.9	97.4	322.5	126.5	697.3	590.8
164		4.5	22.2	91.0	24.2	3022.5	3897.0
98		4.9	18.3	39.6	600.3	723.6	324.5








170		3.2	12.7	38.0	373.4	1356.3	767.2
169		10.5	22.9	59.8	305.7	>10000	
109		6.4	31.2	107.3	674.2		
73		5.4	18.8	80.9	33.9	35.6	
167		4.1	16.0	47.4	310.0	>10000	
168		15.7	59.5	121.7	1156.7	1327.4	
161		16.8	54.3	157.6	2260.9	>10000	








102		5.6	35.3	110.6	68.6	3357.9	
105		3.2	14.4	51.2	14.0	7023.2	
110		5.4	26.9	120.4	35.0	>10000	
74		14.7	68.7	208.9	215.7		
50		1.7	10.7	47.4	10.5	320.8	
106		41.2	283.2	210.6	6111.8	2517.2	
51		3.5	20.8	66.9	226.4	182.5	








40		4.6	23.1	130.3	18.1	50	
173		9.3	50.3	133.9	180.1	220.3	
26		19.9	91.1	293.3	217.5	124.5	
107		50.0	196.3	589.0	172.5	898.7	
95		6.7	27.0	82.4	89.3	59.9	
108		50.5	123.5	197.7	5692.8	1504.2	
100		8.7	32.3	61.9	1149.1	6880.8	








152		41.6	228.0	156.7	3673.3	133.2	
151		33.9	168.9	124.7	2976.2	133.4	
154		77.6	301.6	228.0	5136.3		
153		74.0	241.2	207.0	7458.4		
24		2.9	16.5	69.6	29.9	226.4	
27		2.9	14.7	53.5	48.8	873.6	
101		10.5	41.3	51.1	1956.2	>10000	








29		1.5	8.5	33.2	19.0	86.9	
30		1.0	5.8	23.3	29.0	163.5	
99		6.8	26.9	62.4	425.6	2465.9	
33		5.7	25.0	75.2	281.9	490.1	
103		5.0	31.0	92.6	349.0	235	
32		60.4	276.8	839.2	373.4	627.5	
90		2.7	18.1	77.1	23.2	30.4	








34		5.1	27.8	98.4	24.6	300.6	
41		27.4	142.0	383.2	1188.3	1185.0	
48		67.6	188.4	148.5	6329.8	>10000	
75		61.1	320.9	755.3	1935.5	1345.1	
35		39.9	249.0	816.9	256.4	437.4	
96		30.5	122.6	247.5	1734.4	2871.2	
97		55.1	204.6	414.6	2193.5	2073.3	








36		75.0	401.0	1226.6	364.4	433.1	
68		14.2	72.7	252.8	72.3	43.9	
42		25.3	158.4	534.0	233.3	645	
43		55.1	302.0	1068.7	438.7	763.5	
45		4.0	18.9	68.7	148.5	46.5	
57		21.1	93.4	408.7	148.7	60.4	
46		32.9	101.0	179.0	1032.8	103.4	



44		25.8	85.2	292.6	999.0	59.7	
58		10.3	54.4	236.5	67.5	173.7	
59		5.3	23.5	112.4	22.6	290.5	
53		9.7	33.1	102.3	242.9	21.3	
88		6.6	41.1	139.7	127.3	31.5	
52		2.9	15.0	46.3	109.2	74.3	
62		2.9	14.4	39.5	278.0	71.8	

63		5.0	22.5	82.1	107.0	14.6	
104		10.4	66.1	112.7	1377.4	292.2	
54		6.1	32.9	122.7	25.9	134.6	
64		3.3	11.1	36.9	153.1	212.7	
55		2.1	13.7	54.0	45.9	155.6	
37		5.0	19.7	52.0	169.0	47.4	
60		6.6	29.2	115.2	56.5	34.4	

47		10.6	28.4	54.3	320.2	281.9	
65		23.3	83.8	283.3	125.8	160.9	
69		16.2	63.7	256.8	66.8	91.6	
61		7.3	30.2	110.5	52.3	71.2	
70		12.0	52.4	207.9	79.4	46.2	
89		14.6	57.1	211.6	916.9	46.3	
67		16.3	48.2	72.8	2028.3	104.1	


66		3.5	14.8	60.0	81.3	221.2	
86		38.4	110.3	81.7	4819.5	425	
49		78.0	181.1	248.0	3603.3	385.3	
87		30.6	87.8	133.4	2486.5	213.2	
84		27.6	142.3	537.3	880.1	32.9	
83		12.1	63.3	201.2	139.1	30	
91		28.7	127.7	462.5	865.2	26.3	

71		44.1	156.5	359.0	2753.5	110.1	
72		44.8	134.9	197.9	3445.7	94.1	
156		25.5	120.4	176.8	9995.0	72.4	
85		7.8	47.2	155.9	81.4	12.9	
78		5.9	39.4	89.9	930.1	27.2	
79		10.7	59.1	175.0	698.4	30.7	
80		2.9	16.2	49.2	167.4	197	

82		13.7	83.9	313.1	121.2	20.9	
81		4.7	27.1	102.8	297.5		

CLAIMS:

1. A compound having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

10 A, A'' and A''' are independently C or N, where C may be unsubstituted or substituted by halo or C₁-C₆ alkyl;

R¹ is H, cyano or halo;

15 R² and R^{2'} are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

20 R³ is H, C₁-C₄ alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, or an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N

atoms or (b) 1 O or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent $-Y-R^4$ and/or 1-4 substituents each independently selected from R^5 ;

5

Y is a bond, $-(CH_2)_m-$ or $-O-$;

R^4 is (a) H, C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ or $-NR^6SO_2NR^7R^8$; (b) phenyl or naphthyl, said phenyl or naphthyl being optionally substituted with 1-5 substituents selected from the group consisting of C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, $-CN$, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from the group consisting of O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from the group consisting of C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$;

20

R^5 is C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, $-CN$, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ or $-NR^6SO_2NR^7R^8$;

25

R^6 is H, C_1-C_6 alkyl or C_3-C_8 cycloalkyl, said C_1-C_6 alkyl is optionally substituted by $-NR^7R^8$ or a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from the group consisting of O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from the group consisting of C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, hydroxy and cyano;

30

R^7 and R^8 are each independently H, C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C_1 - C_6 alkyl is optionally substituted by C_3 - C_8 cycloalkyl, halo, hydroxy, amino, (C_1 - C_6 alkyl)amino or di(C_1 - C_6 alkyl)amino and said heterocyclic ring being optionally substituted by one or more C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl groups;


R^9 is C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl;

10

R^{10} is $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently hydrogen, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, phenyl, amino, C_1 - C_6 alkylamino, di(C_1 - C_6 alkyl)amino, heterocyclic, $-(CH_2)_n-W'$, where W' is hydroxy, C_3 - C_8 cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, aryloxy, $-SO_2-R'$, $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently phenyl, C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl;

R^{11} and R^{12} are each independently H, hydroxy, halo, cyano, C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl; and, m and n are independently 0, 1, 2, or 3.

2. A compound having the structure:



(Ib)

or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

5

A'' and A''' are independently C or N, where C may be unsubstituted or substituted by halo or C₁-C₆ alkyl;

R¹ is H, cyano or halo;

10

R² and R^{2'} are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

15

R³ is H, C₁-C₄ alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, or an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N atoms or (b) 1 O or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent -Y-R⁴ and/or 1-4 substituents each independently selected from R⁵;

20

Y is a bond, $-(CH_2)_m-$ or $-O-$;

R^4 is (a) H, C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ or $-NR^6SO_2NR^7R^8$; (b) phenyl or naphthyl, said phenyl or naphthyl being optionally substituted with 1-5 substituents selected from the group consisting of C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, $-CN$, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$; or (c) a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from the group consisting of O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from the group consisting of C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, oxo, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ and $-NR^6SO_2NR^7R^8$;

R^5 is C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, $-CN$, $-OR^6$, $-NR^7R^8$, $-SR^6$, $-SOR^9$, $-SO_2R^9$, $-COR^6$, $-OCOR^6$, $-COOR^6$, $-NR^6COR^6$, $-CONR^7R^8$, $-NR^6SO_2R^9$, $-SO_2NR^7R^8$, $-NR^6CONR^7R^8$, $-NR^6COOR^9$ or $-NR^6SO_2NR^7R^8$;

R^6 is H, C_1-C_6 alkyl or C_3-C_8 cycloalkyl, said C_1-C_6 alkyl is optionally substituted by $-NR^7R^8$ or a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from the group consisting of O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from the group consisting of C_1-C_6 alkyl, C_3-C_8 cycloalkyl, halo, hydroxy and cyano;

R^7 and R^8 are each independently H, C_1-C_6 alkyl or C_3-C_8 cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C_1-C_6 alkyl is optionally substituted by C_3-C_8 cycloalkyl,

halo, hydroxy, amino, (C₁-C₆ alkyl)amino or di(C₁-C₆ alkyl)amino and said heterocyclic ring being optionally substituted by one or more C₁-C₆ alkyl or C₃-C₈ cycloalkyl groups;

5 R⁹ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl;


R¹⁰ is -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, heterocyclic, -(CH₂)_n-W', where W' is hydroxy, C₃-C₈ cycloalkyl, phenyl, naphthyl,
 10 heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently phenyl, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

15

R¹¹ and R¹² are each independently H, hydroxy, halo, cyano, C₁-C₆ alkyl or C₃-C₈ cycloalkyl; and, *m* and *n* are independently 0, 1, 2 or 3.

3. The compound of claim 2 wherein R¹⁰ is -NR''SO₂-R' and R' and R'' are both C₁-C₆
 20 alkyl.

4. A compound having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

R¹ is H, cyano or halo;

5

R² and R^{2'} are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

10 R³ is H, C₁-C₄ alkyl, phenyl, naphthyl, 6-membered heteroaryl or heterocyclic containing 1-3 N atoms, a 5-membered heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N atoms, a 10-membered bicyclic heteroaryl or heterocyclic containing 1-4 N atoms, a 9-membered bicyclic heteroaryl or heterocyclic containing either (a) 1-4 N atoms or (b) 1 O or S atom and 0-3 N
15 atoms, or an 8-membered bicyclic heteroaryl or heterocyclic containing (a) 1-4 N atoms or (b) 1 O or S atom and 1-3 N atoms or (c) 2 O or S atoms and 0-2 N atoms; wherein each of said phenyl, naphthyl, heteroaryl or heterocyclic is optionally substituted by alkyl, 1 substituent -Y-R⁴ and/or 1-4 substituents each independently selected from R⁵;

20

Y is a bond, -(CH₂)_m- or -O-;

R⁴ is (a) H, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, oxo, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸,
25 -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ or -NR⁶SO₂NR⁷R⁸; (b) phenyl or naphthyl, said phenyl or naphthyl being optionally substituted with 1-5 substituents selected from the group consisting of C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, -CN, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸; or (c) a 3 to 8-membered saturated
30 or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from the group consisting of O and N, said heteroaryl being optionally

substituted by 1-5 substituents selected from the group consisting of C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, oxo, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ and -NR⁶SO₂NR⁷R⁸;

5

R⁵ is C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, -CN, -OR⁶, -NR⁷R⁸, -SR⁶, -SOR⁹, -SO₂R⁹, -COR⁶, -OCOR⁶, -COOR⁶, -NR⁶COR⁶, -CONR⁷R⁸, -NR⁶SO₂R⁹, -SO₂NR⁷R⁸, -NR⁶CONR⁷R⁸, -NR⁶COOR⁹ or -NR⁶SO₂NR⁷R⁸;

10 R⁶ is H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl, said C₁-C₆ alkyl is optionally substituted by -NR⁷R⁸ or a 3 to 8-membered saturated or partially unsaturated monocyclic heteroaryl, containing 1 or 2 heteroatoms selected from the group consisting of O and N, said heteroaryl being optionally substituted by 1-5 substituents selected from the group consisting of C₁-C₆ alkyl, C₃-C₈ cycloalkyl, halo, hydroxy and cyano;

15

R⁷ and R⁸ are each independently H, C₁-C₆ alkyl or C₃-C₈ cycloalkyl or are taken together with the nitrogen atom to which they are attached to form a 4-, 5- or 6-membered saturated heterocyclic ring containing 1-2 nitrogen atoms or 1 nitrogen and 1 oxygen atom, said C₁-C₆ alkyl is optionally substituted by C₃-C₈ cycloalkyl, halo, hydroxy, amino, (C₁-C₆ alkyl)amino or di(C₁-C₆ alkyl)amino and said heterocyclic ring being optionally substituted by one or more C₁-C₆ alkyl or C₃-C₈ cycloalkyl groups;

20

R⁹ is C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

25

R¹⁰ is -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, heterocyclic, -(CH₂)_n-W', where W' is hydroxy, C₃-C₈ cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy,

30


81797008

C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently phenyl, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

5 R¹¹ and R¹² are each independently H, hydroxy, halo, cyano, C₁-C₆ alkyl or C₃-C₈ cycloalkyl; and, *m* and *n* are independently 0, 1, 2 or 3.

5. The compound of claim 4 wherein R¹⁰ is -NR''SO₂-R' and R' and R'' are both C₁-C₆ alkyl.

10 6. A compound having the structure:



15 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

A, A', A'' and A''' are independently C or N, where C may be unsubstituted or substituted by halo or C₁-C₆ alkyl;


20 R¹ is H, cyano or halo;

R² and R^{2'} are independently H, C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, C₁-C₆ alkylthio, or C₃-C₈ cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

R^{10} is $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' is independently C_1-C_6 alkyl or C_3-C_8 cycloalkyl; and,

- 5 R^{11} and R^{12} are each independently H, hydroxy, halo, cyano, C_1-C_6 alkyl or C_3-C_8 cycloalkyl.

7. A compound having the structure:



or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:


- 15 A'' and A''' are independently C or N, where C may be unsubstituted or substituted by halo or C_1-C_6 alkyl;

R^1 is H, cyano or halo;

- 20 R^2 and $R^{2'}$ are independently H, C_1-C_6 alkyl, cyano, C_1-C_6 alkoxy, C_1-C_6 alkylthio, or C_3-C_8 cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

- R^{10} is $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently hydrogen, C_1-C_6 alkyl, C_3-C_8 cycloalkyl, phenyl, amino, C_1-C_6 alkylamino, di(C_1-C_6 alkyl)amino, heterocyclic, $-(CH_2)_n-W'$, where W' is hydroxy, C_3-C_8 cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein
- 5 each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyanohydroxy, C_1-C_6 alkyl, C_1-C_6 alkoxy, aryloxy, $-SO_2-R'$, $-NHSO_2-R'$, $-NR''SO_2-R'$ or SR' where R' and R'' are independently phenyl, C_1-C_6 alkyl or C_3-C_8 cycloalkyl;
- 10 R^{11} and R^{12} are each independently H, hydroxy, halo, cyano, C_1-C_6 alkyl or C_3-C_8 cycloalkyl; and, m and n are independently 0, 1, 2 or 3.

8. A compound having the structure:



15

or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or pharmaceutically acceptable salt, wherein:

R^1 is H, cyano or halo;

20

R^2 and $R^{2'}$ are independently H, C_1-C_6 alkyl, cyano, C_1-C_6 alkoxy, C_1-C_6 alkylthio, or C_3-C_8 cycloalkyl where alkyl, alkoxy, or cycloalkyl is optionally substituted by one or more fluorine atoms;

R¹⁰ is -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, phenyl, amino, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, heterocyclic, -(CH₂)_n-W', where W' is hydroxy, C₃-C₈ cycloalkyl, phenyl, naphthyl, heterocyclic, 5- or 6-membered heteroaryl containing 1-3 N and/or O atoms; wherein
 5 each of said alkyl, cycloalkyl, heterocyclic, phenyl, naphthyl or heteroaryl may be unsubstituted or substituted by phenyl, heteroaryl, heterocyclic, halo, cyano, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, aryloxy, -SO₂-R', -NHSO₂-R', -NR''SO₂-R' or SR' where R' and R'' are independently phenyl, C₁-C₆ alkyl or C₃-C₈ cyclo alkyl; and, *n* is 0, 1, 2 or 3.

10

9. The compound of claim 8 wherein R¹⁰ is -NR''SO₂-R' and R' and R'' are both C₁-C₆ alkyl.

10. The compound selected from the group consisting of:

15

4-({2-[ethyl(ethylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)-phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(ethylsulfonyl)amino]benzyl}amino)-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

20

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-[(ethylsulfonyl)(methyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-{{2-(4-hydroxyphenyl)ethyl}amino}-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

25

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-methylpropyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

30

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-fluoro-6-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

5 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-[ethyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(cyclopentylmethyl)amino]-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({5-methyl-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

15 4-({2-[(ethylsulfonyl)(methyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-fluoro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

20 4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-methylpropyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

25 4-[(cyclopentylmethyl)amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-fluoro-6-[methyl(methylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

30 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-({2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

5 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({5-methyl-2-[methyl(methylsulfonyl)amino]-benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(phenylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-{4-[(phenylsulfonyl)amino]phenyl}ethyl)-amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[(2-hydroxyethyl)(methylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

15 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(phenylsulfonyl)amino]benzyl}-amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[(2-hydroxyethyl)(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

20 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}-amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

25 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({4-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}-amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}-amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

30 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[(3-hydroxyphenyl)sulfonyl](methyl)-amino)benzyl]amino]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({4-hydroxy-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

5 4-({2-[ethyl(methylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(phenylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 4-({2-[ethyl(phenylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

15 6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-({2-[methyl(methylsulfonyl)amino]-benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-([(1R)-1-{2-[methyl(methylsulfonyl)amino]-phenyl}ethyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

20 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-([(1S)-1-{2-[methyl(methylsulfonyl)-amino]phenyl}ethyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-[methyl(phenylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

25 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({4-hydroxy-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

30 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(phenylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(phenylsulfonyl)amino]-5-hydroxybenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(phenylsulfonyl)-amino]pyrazin-2-yl)methyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methyl)amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

N-ethyl-4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methyl)amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(4-[methyl(methylsulfonyl)-amino]pyridin-3-yl)methyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-cyclopropyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-[(2-[methyl(methylsulfonyl)amino]-pyridin-3-yl)methyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-methoxy-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)amino]pyrazin-2-yl)methyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(2-[ethyl(methylsulfonyl)amino]pyridin-3-yl)methyl)amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-[methyl(methylsulfonyl)-amino]pyridin-3-yl)methyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

5 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(3-[methyl(methylsulfonyl)-amino]pyridin-2-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-(6-methylpyridin-3-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((1,3,3-trimethylureido)benzyl)-amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(N-methyl-1H-pyrazole-4-sulfonamido)benzyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

15 4-((2-N,1-dimethyl-1H-imidazole-4-sulfonamido)benzyl)amido)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[(2-methoxyethyl)sulfonyl]-(methyl)amino)benzyl]amino]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

20 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

25 4-[(2-[ethyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[[2-(sulfamoylmethyl)benzyl]-amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[[2-(methyl{[6-(morpholin-4-yl)pyridin-3-yl]sulfonyl}amino)benzyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

5 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[[2-(methyl{[3-(morpholin-4-yl)propyl]sulfonyl}amino)benzyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(5-methyl-2-[methyl(methyl-sulfonyl)amino]pyridin-3-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-([2-[methyl(pyridin-3-ylsulfonyl)amino]benzyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

15 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-methyl-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino}benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(2-{methyl[(6-methylpyridin-3-yl)sulfonyl]amino}benzyl)amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

20 4-[(5-chloro-2-[ethyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(5-chloro-2-[methyl(methylsulfonyl)amino]pyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

25 6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-([2-[methyl(sulfamoyl)amino]benzyl]amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide ;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((N-(2-hydroxyethyl)sulfamoyl)(methyl)-aminobenzyl)amino)-N-methyl-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

30 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-N-{6-[(2-hydroxyethyl)amino]pyridin-3-yl}-4-([5-hydroxy-2-

[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(methylsulfonyl)amino]benzyl}amino)-N-(6-methylpyridin-3-yl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-[(5-fluoro-2-[methyl(methylsulfonyl)-amino]pyridin-3-yl)methyl]amino]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(2-[ethyl(methylsulfonyl)amino]-5-fluoropyridin-3-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(3-[ethyl(methylsulfonyl)amino]pyrazin-2-yl)methyl]amino]-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-methyl-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-((2-(N-ethylethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-((2-(N-ethylmethylsulfonamido)-5-fluorobenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoro-ethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-((5-chloro-2-(N-ethylmethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoro-ethyl)phenyl)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylethylsulfonamido)benzyl)-amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-(((5-methyl-2-(N-methylmethylsulfonamido)pyridin-3-yl)methyl)amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-[(2-[ethyl(methylsulfonyl)amino]-5-methylpyridin-3-yl)methyl]amino-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

5 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-fluoro-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({5-chloro-2-[methyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

10 4-({2-[ethyl(methylsulfonyl)amino]benzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)-phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

4-({2-[ethyl(methylsulfonyl)amino]-5-methylbenzyl}amino)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

15 6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(methylsulfonyl)-amino]benzyl}amino)-1H-pyrazolo[4,3-c]pyridine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-N-methyl-4-((2-(N-methylmethylsulfonamido)benzyl)-amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

20 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(N-methylmethylsulfon-amido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-4-((2-(N-methylphenylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

25 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylphenylsulfonamido)-benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

30 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-hydroxy-2-(N-methylmethylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-methylmethylsulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

5 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((5-hydroxy-2-(N-methylmethylsulfonamido)benzyl)amino)-N-(6-((2-hydroxyethyl)amino)pyridin-3-yl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-(((3-(N-methylmethyl-sulfonamido)pyrazin-2-yl)methyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

10 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-(2-hydroxyethyl)methyl-sulfonamido)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

15 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-4-((2-(N-(2-hydroxyethyl)methylsulfon-amido)benzyl)amino)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

4-((2-(N-ethylmethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)-phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

20 4-((5-fluoro-2-(N-methylmethylsulfonamido)benzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

4-((2-(N-ethylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

25 4-((2-(N-methylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

4-((2-(N-ethylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

30 4-((2-(N-methylphenylsulfonamido)-5-hydroxybenzyl)amino)-6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(methyl(sulfamoyl)amino)-benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

5 6-(5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl)-N-methyl-4-((2-(methyl(N-methylsulfamoyl)amino)benzyl)amino)-1H-pyrazolo[3,4-d]pyrimidine-3-carboxamide;

4-[4-(7,8-dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]-3-ethylphenol formate;

3-ethyl-4-[4-(3-phenoxyazetidin-1-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate;

10 3-ethyl-4-{4-[6-(4-methyl-1H-imidazol-1-yl)-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl}phenol formate;

3-ethyl-4-{4-[6-(2-methoxyethoxy)-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl}phenol formate;

15 1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-3-methylazetidin-3-ol formate;

2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-N-[2-(pyrrolidin-1-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide formate;

N-benzyl-2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-1,2,3,4-tetrahydroisoquinoline-5-carboxamide formate;

20 4-{4-[7-(benzyloxy)-6-methoxy-3,4-dihydroisoquinolin-2(1H)-yl]-1H-pyrazolo[4,3-c]pyridin-6-yl}-3-ethylphenol formate;

4-[4-(5-chloro-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]-3-ethylphenol formate;

25 4-chloro-3-({1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]azetidin-3-yl}oxy)benzonitrile formate;

3-ethyl-4-[4-(6-fluoro-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate;

3-ethyl-4-[4-(8-methoxy-3,4-dihydroisoquinolin-2(1H)-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol formate;

30 N-{2-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-1,2,3,4-tetrahydroisoquinolin-5-yl}methanesulfonamide formate;

4-(4-([2-(biphenyl-4-yl)ethyl]amino)-1H-pyrazolo[4,3-c]pyridin-6-yl)-3-ethylphenol formate;

N-[2-([6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl]-N-methylmethanesulfonamide hydrochloride;

5 1-[6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]-N,N-dimethylpyrrolidine-3-sulfonamide (racemic);

N-[2-([6-(2-ethyl-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl)-3-methylphenyl]-N-methylmethanesulfonamide diethylamine salt;

10 3-ethyl-4-[4-(4-methoxypiperidin-1-yl)-1H-pyrazolo[4,3-c]pyridin-6-yl]phenol diethylamine salt;

N-[2-([2-(3,4-dimethoxyphenyl)ethyl][6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl]-N-methylmethanesulfonamide hydrochloride;

15 N-[2-([6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](2-{4-[(methylsulfon-yl)amino]phenyl}ethyl)amino)methyl]phenyl]-N-methylmethanesulfonamide hydrochloride;

N-[2-([6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl)-phenyl]-N-methylmethanesulfonamide;

20 N-[2-([6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl)-4-methylphenyl]-N-methylmethanesulfonamide hydrochloride;

N-[4-chloro-2-([6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl]-N-methylmethanesulfonamide hydrochloride;

N-[2-([6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl)-3-fluorophenyl]-N-methylmethanesulfonamide hydrochloride;

25 N-[2-([6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl)-phenyl]-N-methylethanesulfonamide hydrochloride;

N-ethyl-N-[2-([6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)-methyl]phenyl]ethanesulfonamide hydrochloride;

30 N-[2-([6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl]-N-propylmethanesulfonamide;

N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl]methanesulfonamide;

N-butyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl]methanesulfonamide;

5 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl][2-(morpholin-4-yl)ethyl]amino}-methyl)phenyl]-N-methylmethanesulfonamide;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl][2-(morpholin-4-yl)ethyl]amino}-methyl)phenyl]-N-methylmethanesulfonamide;

10 N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)-amino}-methyl)-4-methylphenyl]methanesulfonamide;

N-[2-({ethyl[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)-4-methylphenyl]-N-methylmethanesulfonamide;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](propyl)amino}-methyl)-4-methylphenyl]-N-methylmethanesulfonamide;

15 N-ethyl-N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino}-methyl)phenyl]methanesulfonamide hydrochloride;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino}-methyl)phenyl]-N-methylmethanesulfonamide hydrochloride;

20 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino}-methyl)phenyl]-N-(2-hydroxyethyl)methanesulfonamide;

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]phenyl}-N-methylmethanesulfonamide;

N-(2-{[6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl](methyl)amino}methyl)phenyl)-N-methylmethanesulfonamide;

25 N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl]-4-methylphenyl}-N-methylmethanesulfonamide;

4-{4-[(cyclopropylmethyl)amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-2-fluoro-5-(2,2,2-trifluoroethyl)phenol;

30 4-{4-[(2-cyclopropylethyl)amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-2-fluoro-5-(2,2,2-trifluoroethyl)phenol;

2-fluoro-4-{4-[(2-methylpropyl)amino]-1H-pyrazolo[3,4-d]pyrimidin-6-yl}-5-(2,2,2-trifluoroethyl)phenol;

4-[4-(butylamino)-1H-pyrazolo[3,4-d]pyrimidin-6-yl]-2-fluoro-5-(2,2,2-trifluoroethyl)phenol;

5 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]amino)methyl}-phenyl]-N-methylmethanesulfonamide hydrochloride;

N-(2-(((6-(2-ethyl-4-hydroxy-6-methylphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl)amino)methyl)-phenyl)-N-methylmethanesulfonamide;

10 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-3-(1H-imidazol-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

N-[2-({[3-(4,5-dimethyl-1H-imidazol-2-yl)-6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-3-(4-methyl-1H-imidazol-2-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

15 4-[3-(5-benzyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-4-{[2-(methylsulfanyl)ethyl]-amino}-1H-pyrazolo[4,3-c]pyridin-6-yl]-2-fluoro-5-(2,2,2-trifluoroethyl)phenol;

N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-3-(1H-pyrazol-1-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

20 N-{2-[[6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(1H-pyrazol-1-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl}-N-methylmethanesulfonamide;

N-{2-[[6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(5-methyl-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl}-N-methylmethanesulfonamide;

25 N-[2-({[6-(2-ethyl-5-fluoro-4-hydroxyphenyl)-3-(5-methyl-4H-1,2,4-triazol-3-yl)-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl)-N-methylmethanesulfonamide;

N-{2-[[6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-[5-(6-methylpyridin-3-yl)-4H-1,2,4-triazol-3-yl]-1H-pyrazolo[4,3-c]pyridin-4-yl]amino)methyl}phenyl}-N-methylmethanesulfonamide;

4-(5-{6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl}-4H-1,2,4-triazol-3-yl)piperidine-1-carboxamide;

5 N-(2-{{6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-{5-[1-(pyrrolidin-1-yl)acetyl]piperidin-4-yl}-4H-1,2,4-triazol-3-yl}-1H-pyrazolo[3,4-d]pyrimidin-4-yl}amino)methyl}phenyl)-N-methyl-methanesulfonamide;

N-{2-[(6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-3-(4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl}amino)methyl]-4-hydroxyphenyl}-N-methylmethanesulfonamide;

10 N-{2-[(3-(5-acetyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-1H-pyrazolo[3,4-d]pyrimidin-4-yl}amino)methyl]-4-hydroxyphenyl}-N-methylmethanesulfonamide;

15 N-[2-(dimethylamino)ethyl]-2-{6-[5-fluoro-4-hydroxy-2-(2,2,2-trifluoroethyl)phenyl]-4-({5-hydroxy-2-[methyl(methylsulfonyl)amino]benzyl}amino)-1H-pyrazolo[3,4-d]pyrimidin-3-yl}-1,4,6,7-tetrahydro-5H-imidazo[4,5-c]pyridine-5-carboxamide; and,

4-(3-(5-benzyl-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridin-2-yl)-4-((3-hydroxy-2-methylpropyl)-amino)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)-2-fluoro-5-(2,2,2-trifluoroethyl)phenol.

20

11. A pharmaceutical composition comprising a compound as defined in any one of claims 1-10, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or salt, and a pharmaceutically acceptable excipient.

25


12. Use of a therapeutically effective amount of a compound as defined in any one of claims 1-10, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or salt, for treating allergic rhinitis, nasal congestion, rhinorrhea, perennial rhinitis, nasal inflammation, asthma of all types, chronic obstructive pulmonary disease, chronic or acute bronchoconstriction, chronic bronchitis, small airways obstruction, emphysema, chronic eosinophilic pneumonia,

30

adult respiratory distress syndrome, exacerbation of airways hyper-reactivity consequent to other drug therapy, pulmonary vascular disease, pulmonary arterial hypertension, acute lung injury, bronchiectasis, sinusitis, allergic conjunctivitis, idiopathic pulmonary fibrosis or atopic dermatitis in a subject.

5

13. Use of a therapeutically effective amount of a compound as defined in any one of claims 1-10, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of said compound or salt, for treating inflammation, neuroinflammation, arthritis, rheumatoid arthritis, spondyloarthropathies, systemic lupus erythematosus arthritis, osteoarthritis, gouty arthritis, pain, fever, pulmonary sarcoidosis, silicosis, cardiovascular disease, atherosclerosis, myocardial infarction, thrombosis, congestive heart failure and cardiac reperfusion injury, cardiomyopathy, stroke, ischaemia, reperfusion injury, brain edema, brain trauma, neurodegeneration, liver disease, inflammatory bowel disease, Crohn's disease, ulcerative colitis, nephritis, retinitis, retinopathy, macular degeneration, glaucoma, diabetes type 1 or type 2, diabetic neuropathy, viral and bacterial infection, myalgia, endotoxic shock, toxic shock syndrome, autoimmune disease, osteoporosis, multiple sclerosis, endometriosis, menstrual cramps, vaginitis, candidiasis, cancer, fibrosis, obesity, muscular dystrophy, polymyositis, Alzheimer's disease, skin flushing, eczema, psoriasis, atopic dermatitis or sunburn, in a subject.



(Ia)