



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2274967 C 2009/10/13

(11)(21) **2 274 967**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1997/12/12
(87) Date publication PCT/PCT Publication Date: 1998/06/25
(45) Date de délivrance/Issue Date: 2009/10/13
(85) Entrée phase nationale/National Entry: 1999/06/15
(86) N° demande PCT/PCT Application No.: US 1997/023015
(87) N° publication PCT/PCT Publication No.: 1998/026796
(30) Priorité/Priority: 1996/12/16 (US08/767,169)

(51) Cl.Int./Int.Cl. *A61K 38/22* (2006.01)
(72) Inventeur/Inventor:
YOUNG, ANDREW A., NZ
(73) Propriétaire/Owner:
AMYLIN PHARMACEUTICALS, INC., US
(74) Agent: SMART & BIGGAR

(54) Titre : METHODES ET COMPOSITIONS POUR LE TRAITEMENT DE LA DOULEUR
(54) Title: METHODS AND COMPOSITIONS FOR TREATING PAIN

(57) **Abrégé/Abstract:**

Methods for treating pain are disclosed which comprise administration of a therapeutically effective amount of an amylin or an amylin agonist alone or in conjunction with a narcotic analgesic, for example, morphine or pentazocine.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/22	A1	(11) International Publication Number: WO 98/26796 (43) International Publication Date: 25 June 1998 (25.06.98)
(21) International Application Number: PCT/US97/23015 (22) International Filing Date: 12 December 1997 (12.12.97) (30) Priority Data: 08/767,169 16 December 1996 (16.12.96) US (71) Applicant: AMYLIN PHARMACEUTICALS, INC. [US/US]; 9373 Towne Centre Drive, San Diego, CA 92121 (US). (72) Inventor: YOUNG, Andrew, A.; 9514 Easter Way, San Diego, CA 92121 (US). (74) Agents: DUFT, Bradford, J. et al.; Lyon & Lyon LLP, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHODS AND COMPOSITIONS FOR TREATING PAIN (57) Abstract Methods for treating pain are disclosed which comprise administration of a therapeutically effective amount of an amylin or an amylin agonist alone or in conjunction with a narcotic analgesic, for example, morphine or pentazocine.		

WO 98/26796

PCT/US97/23015

METHODS AND COMPOSITIONS FOR TREATING PAIN**Field of the Invention**

The present invention relates to methods for treating pain. More particularly, the invention relates to the use of an amylin or agonist of amylin in the treatment of pain, either alone or in combination with a narcotic analgesic.

Background**Amylin**

The structure and biology of amylin have previously been reviewed. See, for example, Rink et al., *Trends in Pharmaceutical Sciences*, 14:113-118 (1993); Gaeta and Rink, *Med. Chem. Res.*, 3:483-490 (1994); and, Pittner et al., *J. Cell. Biochem.*, 55S:19-28 (1994). Amylin is a 37 amino acid protein hormone. It was isolated, purified and chemically characterized as the major component of amyloid deposits in the islets of pancreases of deceased human Type 2 diabetics (Cooper et al., *Proc. Natl. Acad. Sci. USA*, 84:8628-8632 (1987)). The amylin molecule has two important post-translational modifications: the C-terminus is amidated, and the cysteines in positions 2 and 7 are cross-linked to form an N-terminal loop. The sequence of the open reading frame of the human amylin gene shows the presence of the Lys-Arg dibasic amino acid proteolytic cleavage signal, prior to the N-terminal codon for Lys, and the Gly prior to the Lys-Arg proteolytic signal at the C-terminal position, a typical sequence for amidation by protein amidating enzyme, PAM (Cooper et al.,

Biochem. Biophys. Acta, 1014:247-258 (1989)). Amylin is the subject of United States Patent No. 5,367,052, issued November 22, 1995.

In Type 1 diabetes, amylin has been shown to be deficient and combined replacement with insulin has been proposed as a preferred treatment over insulin alone in all forms of diabetes. The use of amylin and other amylin agonists for the treatment of diabetes mellitus is the subject of United States Patent No. 5,175,145, issued December 29, 1992. Pharmaceutical compositions containing amylin and amylin plus insulin are described in United States Patent No. 5,124,314, issued June 23, 1992.

Excess amylin action has been said to mimic key features of Type 2 diabetes and amylin blockade has been proposed as a novel therapeutic strategy. It has been disclosed in United States Patent No. 5,266,561, issued November 30, 1993, that amylin causes reduction in both basal and insulin-stimulated incorporation of labeled glucose into glycogen in skeletal muscle. The latter effect was also disclosed to be shared by calcitonin gene related peptide (CGRP) (see also Leighton and Cooper, *Nature*, 335:632-635 (1988)). Amylin and CGRP were approximately equipotent, showing marked activity at 1 to 10 nM. Amylin is also reported to reduce insulin-stimulated uptake of glucose into skeletal muscle and reduce glycogen content (Young et al., *Amer. J. Physiol.*, 259:45746-1 (1990)). The treatment of Type 2 diabetes and insulin resistance with amylin antagonists is disclosed.

Both the chemical structure and the gene sequence of amylin have been said to support the determination that it is a biologically active or "messenger" molecule. The chemical structure is nearly 50% identical to the CGRPs, also

WO 98/26796

PCT/US97/23015

37 amino acid proteins which are widespread neurotransmitters with many potent-biological actions, including vasodilation. Amylin and CGRP share the ²Cys-⁷Cys disulphide bridge and the C-terminal amide, both of which are essential for full biologic activity (Cooper *et al.*, *Proc. Natl. Acad. Sci. USA*, 85:7763-7766 (1988)).

Amylin reportedly may be one member of a family of related peptides which includes CGRP, insulin, insulin-like growth factors and the relaxins and which share common genetic heritage (Cooper *et al.*, *Prog. Growth Factor Research*, 1:99-105 (1989)). The two peptides calcitonin and CGRP-1 share common parentage in the calcitonin gene where alternative processing of the primary mRNA transcript leads to the generation of the two distinct peptides, which share only limited sequence homology (about 30%) (Amara *et al.*, *Science*, 229:1094-1097 (1985)). The amylin gene sequence is typical for a secreted messenger protein, with the mRNA coding a prepropeptide with processing sites for production of the secreted protein within the Golgi or secretory granules. Amylin is mainly co-localized with insulin in beta cell granules and may share the proteolytic processing enzymes that generate insulin from pro-insulin.

Amylin is primarily synthesized in pancreatic beta cells and is secreted in response to nutrient stimuli such as glucose and arginine. Studies with cloned beta-cell tumor lines (Moore *et al.*, *Biochem. Biophys. Res. Commun.*, 179(1) (1991)), isolated islets (Kanatsuka *et al.*, *FEBS Letts.*, 259(1), 199-201 (1989)) and perfused rat pancreases (Ogawa *et al.*, *J. Clin. Invest.*, 85:973-976 (1990)) have shown that short pulses, 10 to 20 minutes, of nutrient secretagogues such as glucose and arginine, stimulate release of amylin as well as insulin. The molar amylin:insulin ratio of the

WO 98/26796

PCT/US97/23015

secreted proteins varies between preparations from about 0.01 to 0.4, but appears not to vary much with acute stimuli in any one preparation. However, during prolonged stimulation by elevated glucose, the amylin:insulin ratio can progressively increase (Gedulin et al., *Biochem. Biophys. Res. Commun.*, 180(1):782-789 (1991)). Thus, perhaps because gene expression and rate of translation are independently controlled, amylin and insulin are not always secreted in a constant ratio.

It has been discovered that certain actions of amylin are similar to known non-metabolic actions of CGRP and calcitonin; however, the metabolic actions of amylin discovered during investigations of this newly identified protein appear to reflect its primary biologic role. At least some of these metabolic actions are mimicked by CGRP, albeit at doses which are markedly vasodilatory (see, e.g., Leighton et al., *Nature*, 335:632-635 (1988)); Molina et al., *Diabetes*, 39:260-265 (1990)).

The first discovered action of amylin was the reduction of insulin-stimulated incorporation of glucose into glycogen in rat skeletal muscle (Leighton et al., *Nature*, 335:632-635 (1988)); the muscle was made "insulin-resistant." Subsequent work with rat soleus muscle *ex-vivo* and *in vitro* has indicated that amylin reduces glycogen synthase activity, promotes conversion of glycogen phosphorylase from the inactive b form to the active a form, promotes net loss of glycogen (in the presence or absence of insulin), increases glucose-6-phosphate levels, and can increase lactate output (see, e.g., Deems et al., *Biochem. Biophys. Res. Commun.*, 181(1):116-120 (1991)); Young et al., *FEBS Letts*, 281(1,2):149-151 (1991)). Amylin, like epinephrine, appears not to affect glucose transport per se (e.g., Pittner et al.,

FEBS Letts., 365(1):98-100 (1995)). Studies of amylin and insulin dose-response relations show that amylin acts as a noncompetitive or functional antagonist of insulin in skeletal muscle (Young *et al.*, *Am. J. Physiol.*, 263(2):E274-E281 (1992)). There is no evidence that amylin interferes with insulin binding to its receptors, or the subsequent activation of insulin receptor tyrosine kinase (Follett *et al.*, *Clinical Research*, 39(1):39A (1991)); Koopmans *et al.*, *Diabetologia*, 34:218-224 (1991)). The actions of amylin on skeletal muscle resemble those of adrenaline (epinephrine). However, while adrenaline's actions are believed to be mediated largely by cAMP, some workers have concluded that amylin's actions are not mediated by cAMP (see Deems *et al.*, *Biochem. Biophys. Res. Commun.*, 181(1):116-120 (1991)). Others report that amylin does activate adenyl cyclase and increases cAMP in skeletal muscle (Pittner *et al.*, *Biochim Biophys. Acta*, 1267:75-82 (1995); Moore and Rink, *Diabetes*, 42 (Suppl 1):257A (abstract 821) (1993)), consistent with transduction of its effect on glycogen metabolism via cAMP-dependent protein kinase phosphorylation of synthase and phosphorylase.

It is believed that amylin acts through receptors present in plasma membranes. It has been reported that amylin works in skeletal muscle via a receptor-mediated mechanism that promotes glycogenolysis, by activating the rate-limiting enzyme for glycogen breakdown, phosphorylase a (Young *et al.*, *FEBS Letts.*, 281:149-151 (1991)). Studies of amylin and CGRP, and the effect of selective antagonists, suggest that amylin acts via its own receptor (Beaumont *et al.*, *Br. J. Pharmacol.*, 115(5):713-715 (1995); Wang *et al.*, *FEBS Letts.*, 219:195-198 (1991 b)), counter to the conclusion of other workers that amylin may act primarily at CGRP

receptors (e.g., Chantry et al., *Biochem. J.*, 277:139-143 (1991)); Galeazza et al., *Peptides*, 12:585-591 (1991)); Zhu et al., *Biochem. Biophys. Res. Commun.*, 177(2):771-776 (1991)). Recently, amylin receptors and their use in various methods for screening and assaying for amylin agonist and antagonist compounds were described in United States Patent No. 5,264,372, issued November 23, 1993.

While amylin has marked effects on hepatic fuel metabolism in vivo, there is no general agreement as to what amylin actions are seen in isolated hepatocytes or perfused liver. The available data do not support the idea that amylin promotes hepatic glycogenolysis, i.e., it does not act like glucagon (e.g., Stephens et al., *Diabetes*, 40:395-400 (1991); Gomez-Foix et al., *Biochem J.*, 276:607-610 (1991)). It has been suggested that amylin may act on the liver to promote conversion of lactate to glycogen and to enhance the amount of glucose able to be liberated by glucagon (see Roden et al., *Diabetologia*, 35:116-120 (1992)). Thus, amylin could act as an anabolic partner to insulin in liver, in contrast to its catabolic action in muscle.

In fat cells, contrary to its adrenaline-like action in muscle, amylin has no detectable actions on insulin-stimulated glucose uptake, incorporation of glucose into triglyceride, CO₂ production (Cooper et al., *Proc. Natl. Acad. Sci.*, 85:7763-7766 (1988)) epinephrine-stimulated lipolysis, or insulin-inhibition of lipolysis (Lupien and Young, "Diabetes Nutrition and Metabolism - Clinical and Experimental," Vol. 6(1), pages 1318 (February 1993)). Amylin thus exerts tissue-specific effects, with direct action on skeletal muscle, marked indirect (via supply of substrate) and perhaps direct effects on liver, while

WO 98/26796

PCT/US97/23015

adipocytes appear "blind" to the presence or absence of amylin.

It has also been reported that amylin can have marked effects on secretion of insulin. In isolated islets (Ohsawa *et al.*, *Biochem. Biophys. Res. Commun.*, 160(2):961-967 (1989)), in the perfused pancreas (Silvestre *et al.*, *Reg. Pept.*, 31:23-31 (1991)), and in the intact rat (Young *et al.*, *Mol. Cell. Endocrinol.*, 84:R1-R5 (1992)), some experiments indicate that amylin inhibits insulin secretion. Other workers, however, have been unable to detect effects of amylin on isolated β -cells, on isolated islets, or in the whole animal (see Broderick *et al.*, *Biochem. Biophys. Res. Commun.*, 177:932-938 (1991) and references therein).

In preclinical experiments, a striking effect of amylin *in vivo* is stimulation of a rapid rise in plasma lactate, followed by a rise in plasma glucose (Young *et al.*, *FEBS Letts.*, 281(1,2):149-151 (1991)). Evidence indicates that the increased lactate provides substrate for glucose production and that amylin actions can occur independent of changes in insulin or glucagon. In "glucose clamp" experiments, amylin infusions cause "insulin resistance", both by reducing peripheral glucose disposal, and by limiting insulin-mediated suppression of hepatic glucose output (e.g., Frontoni *et al.*, *Diabetes*, 40:568-573 (1991)); Koopmans *et al.*, *Diabetologia*, 34:218-224 (1991)).

In lightly anesthetized rats which were fasted for 18 hours to deplete their stores of hepatic glycogen, amylin injections stimulated rises in plasma lactate from about 0.5 to 1.5 mM followed by a prolonged increase in plasma glucose levels from about 6 to 11 mM. These effects were observed for both intravenous and subcutaneous injections (Young *et al.*, *FEBS Letts.*, 281(1,2):149-151 (1991)). The effects of

amylin in fed animals differ quantitatively from its effects in fasted animals. In fed rats, with presumably normal liver glycogen stores, amylin causes a more pronounced and prolonged rise in plasma lactate; however, there is only a modest rise in plasma glucose. It has been suggested that amylin promotes the "return limb" of the Cori cycle, *i.e.*, muscle glycogen via breakdown to lactate provides substrate for hepatic gluconeogenesis and glycogen production and probably triglyceride synthesis. Insulin drives the forward limb, *i.e.*, uptake of glucose into muscle and production of muscle glycogen. Insulin and amylin can thus be seen as partners in regulating the "indirect" pathway of post-prandial hepatic glycogen repletion. "Insulin resistance" in muscle and liver may be under normal, physiologic regulation by amylin.

Amylin or amylin agonists potently inhibit gastric emptying in rats (Young *et al.*, *Diabetologia* 38(6):642-648 (1995)), dogs (Brown *et al.*, *Diabetes* 43(Suppl 1):172A (1994)) and humans (Macdonald *et al.*, *Diabetologia* 38(Suppl 1):A32 (abstract 118) (1995)). Because gastric emptying is accelerated in amylin-deficient type 1 diabetic BB rats (Young *et al.*, *Diabetologia*, *supra*; Nowak *et al.*, *J. Lab. Clin. Med.*, 123(1):110-6 (1994)) and in rats treated with the selective amylin antagonist, AC187 (Gedulin *et al.*, *Diabetologia*, 38(Suppl 1):A244 (1995)), the effect of amylin on gastric emptying appears to be physiological (operative at concentrations that normally circulate).

Control of nutrient release from the stomach is becoming recognized as an important component of overall fuel homeostasis. In human volunteers, over a range of carbohydrate concentrations in a liquid meal, energy release from the stomach was remarkably constant at ~2 kcal/min

(Brener et al., *Gastroenterology*, 85(1):76-82 (1983)), equivalent to ~500 mg glucose/min. This release rate is about the same as the rate of glucose disposal that insulin-sensitive individuals can attain at their peak plasma insulin concentrations (1nM) (Young et al., *Am. J. Physiol.* 254 (2 Pt 1):E231-236 (1988)). Thus, the rate at which carbohydrate is released from the stomach and absorbed is normally matched to the rate at which it can be metabolized.

Several feedback loops may control nutrient efflux from the stomach. Peptides that are candidates as participants in such control loops should (1) change in response to meals, and (2) potentially modulate gastric emptying. Peptides known to inhibit gastric emptying are limited to amylin, cholecystokinin (CCK), glucagon-like peptide-1 (GLP-1), secretin and gastrin releasing peptide/bombesin (GRP). GRP does not change with meals. Secretin is secreted in response to acid (but not nutrients) entering the duodenum. Of the peptides that are secreted in response to nutrients (amylin, CCK and GLP-1), only amylin and GLP-1 are secreted in response to glucose ingestion. CCK is secreted in response to fat ingestion. Of the hormones that might therefore mediate feedback control of the gastric release and subsequent absorption of glucose (amylin and GLP-1), amylin is the more potent (Young et al., *Metabolism Clinical and Experimental* 45(1):1-3 (1996)), and it appears that amylin may be a major regulator of carbohydrate absorption, at least in rodents (Young et al., *Biochemical Society Transactions* 23(2):325-331 (1995)).

Non-metabolic actions of amylin include vasodilator effects which may be mediated by interaction with CGRP vascular receptors. Reported *in vivo* tests suggest that amylin is at least about 100 to 1000 times less potent than

WO 98/26796

PCT/US97/23015

CGRP as a vasodilator (Brain et al., *Eur. J. Pharmacol.*, 183:2221 (1990); Wang et al., *FEBS Letts.*, 291:195-198 (1991)). The effect of amylin on regional hemodynamic actions, including renal blood flow, in conscious rats was recently reported (Gardiner et al., *Diabetes*, 40:948-951 (1991)). The authors noted that infusion of rat amylin was associated with greater renal vasodilation and less mesenteric vasoconstriction than is seen with infusion of human α -CGRP. They concluded that, by promoting renal hyperemia to a greater extent than did α -CGRP, rat amylin could cause less marked stimulation of the renin-angiotensin system, and thus, less secondary angiotensin II-mediated vasoconstriction. It was also noted, however, that during coinfusion of human α -⁸⁻³⁷CGRP and rat amylin renal and mesenteric vasoconstrictions were unmasked, presumably due to unopposed vasoconstrictor effects of angiotensin II, and that this finding is similar to that seen during coinfusion of human A-CGRP and human α -⁸⁻³⁷CGRP (*id.* at 951).

Injected into the brain, or administered peripherally, amylin has been reported to suppress food intake, e.g., Chance et al., *Brain Res.*, 539:352-354 (1991)), an action shared with CGRP and calcitonin. The effective concentrations at the cells that mediate this action are not known. Amylin has also been reported to have effects both on isolated osteoclasts where it caused cell quiescence, and *in vivo* where it was reported to lower plasma calcium by up to 20% in rats, in rabbits, and in humans with Paget's disease (see, e.g., Zaidi et al., *Trends in Endocrinal. and Metab.*, 4:255-259 (1993)). From the available data, amylin seems to be 10 to 30 times less potent than human calcitonin for these actions. Interestingly, it was reported that amylin appeared to increase osteoclast cAMP production but not to increase

WO 98/26796

PCT/US97/23015

cytosolic Ca^{2+} , while calcitonin does both (Alam et al., *Biochem. Biophys. Res. Commun.*, 179(1):134-139 (1991)). It was suggested, though not established, that calcitonin may act via two receptor types and that amylin may interact with one of these.

Infusing amylin receptor antagonists may be used to alter glucoregulation. $^{8-37}\text{CGRP}$ is a demonstrated amylin blocker *in vitro* and *in vivo* (Wang et al., *Biochem. Biophys. Res. Commun.*, 181(3):1288-1293 (1991)), and was found to alter glucoregulation following an arginine infusion in fed rats (Young et al., *Mol. Cell. Endocrinol.*, 84:R1-R5 (1992)). The initial increase in glucose concentration is attributed to arginine-stimulated glucagon secretion from islet alpha cells; the subsequent restoration of basal glucose is attributed to insulin action along with changes in other glucoregulatory hormones. When the action of amylin is blocked by preinfusion of $^{8-37}\text{hCGRP}$, the initial glucose increase is not significantly different, but there is a subsequent fall in glucose concentration to well below the basal level, which is restored only after some 80 minutes. Thus, glucoregulation following this challenge with an islet secretagogue was reportedly altered by infusion of an amylin receptor antagonist. Additionally, insulin concentrations were measured at half-hour intervals and it was found that insulin concentration 30 minutes following the arginine infusion was almost twice as high in animals infused with an amylin receptor antagonist as in the normal controls. $^{8-37}\text{CGRP}$ is also an effective CGRP antagonist. However, very similar results were seen with another amylin antagonist, AC66, which is selective for amylin receptors compared with CGRP receptors (Young et al., *Mol. Cell. Endocrinol.*, 84:R1-R5 (1992)). These results are said to support the conclusion

that blockade of amylin action can exert important therapeutic benefits in Type 2 diabetes.

It has also been discovered that, surprisingly in view of its previously described renal vasodilator and other properties, amylin markedly increases plasma renin activity in intact rats when given subcutaneously in a manner that avoids any disturbance of blood pressure. This latter point is important because lowered blood pressure is a strong stimulus to renin release. Amylin antagonists, such as amylin receptor antagonists, including those selective for amylin receptors compared to CGRP and/or calcitonin receptors, can be used to block the amylin-evoked rise of plasma renin activity. The use of amylin antagonists to treat renin-related disorders is described in United States Patent No. 5,376,638, issued December 27, 1994.

In normal humans, fasting amylin levels from 1 to 10pM and post-prandial or post-glucose levels of 5 to 20pM have been reported (e.g., Hartter et al., *Diabetologia*, 34:52-54 (1991); Sanke et al., *Diabetologia*, 34:129-132 (1991); Koda et al., *The Lancet*, 339:1179-1180 (1992)). In obese, insulin-resistant individuals, post-food amylin levels can go higher, reaching up to about 50pM. For comparison, the values for fasting and post-prandial insulin are 20 to 50pM, and 100 to 300 pM respectively in healthy people, with perhaps 3-to 4-fold higher levels in insulin-resistant people. In Type 1 diabetes, where beta cells are destroyed, amylin levels are at or below the levels of detection and do not rise in response to glucose (Koda et al., *The Lancet*, 339:1179-1180 (1992)). In normal mice and rats, basal amylin levels have been reported from 30 to 100 pM, while values up to 600 pM have been measured in certain insulin-resistant, diabetic strains of rodents (e.g., Huang et al.,

WO 98/26796

PCT/US97/23015

Hypertension, 19:I-101-I-109 (1991); Gill et al., *Life Sciences*, 48:703-710 (1991)).

Patients with Type 1 diabetes, in addition to a lack of insulin, are reported to have marked amylin deficiency. As noted above, data show that amylin expression and secretion by pancreatic beta cells is absent or well below normal in Type 1 diabetes. In several animal models of Type 1 diabetes, amylin secretion and gene expression are depressed (Cooper et al., *Diabetes*, 497-500 (1991); Ogawa et al., *J. Clin. Invest.*, 85:973-976 (1990)). Measurements of plasma amylin in Type 1 diabetic patients show that amylin is deficient in these patients after an overnight fast, and that a glucose load does not elicit any increase in amylin levels (Koda et al., *The Lancet*, 339:1179-1180 (1992)).

Pain

Pain is defined by the International Association for the Study of Pain as "an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage." Merskey, *Pain* 6:249 (1979). Many diseases and medical procedures are characterized by the sense of pain in different ways for a patient. For example, during and after surgical operations, the manifestation of pain is high. The same is true for many disorders involving trauma, such as burns. Pain is also present in most inflammatory conditions and in association with tumor-related diseases or treatment of those. Different therapies are used to achieve an analgesic effect. Drugs such as narcotic analgesics, for example, morphine, are used for reducing severe pain, for example that associated with surgical operations. Another type of pharmaceutical used to reduce pain is the sedative agent, which include barbiturates

WO 98/26796

PCT/US97/23015

and benzodiazepines. Many of these drugs have side effects such as depressant action on respiration and circulation. Several of these drugs also may induce hypnotic effects. Non-steroidal drugs are also used to treat pain and inflammation, and appear to act by preventing the synthesis of prostaglandins.

Chronic or intractable pain, such as may occur in conditions such as degenerative bone diseases and cancer, is a debilitating condition which is treated with a variety of analgesic agents, for example, narcotic analgesics compounds such as morphine. Chronic pain can also outlast the onset of any known or suspected physical cause. It can occur long after a known injury or disease or it can occur without any known physical cause whatsoever. It can also be accompanied by known tissue pathology, such as chronic inflammation which occurs in some types of arthritis.

In general, brain pathways governing the perception of pain are incompletely understood. Sensory afferent synaptic connections to the spinal cord, termed "nociceptive pathways" have been documented in some detail. In the first leg of such pathways, C- and A-fibers which project from peripheral sites to the spinal cord carry nociceptive signals. Polysynaptic junctions in the dorsal horn of the spinal cord are involved in the relay and modulation of sensations of pain to various regions of the brain, including the periaqueductal grey region (McGreer). Analgesia, or the absence or reduction of pain in response to a stimulation that would normally be painful, can be effected directly by decreasing transmission along such nociceptive pathways. Analgesic opiates are thought to act by mimicking the effects of endorphin or enkephalin peptide-containing neurons, which terminate presynaptically at the C- or A-fiber terminals and

which, when they fire, inhibit release of substance P. Descending pathways from the brain are also inhibitory on C- and A-fiber firing.

Narcotic analgesics, for example, opiates such as morphine, while effective in producing analgesia, may induce tolerance in a subject, so that increasing doses are required to achieve a satisfactory effect. At high doses, such compounds produce side effects, including respiratory depression, which can be life threatening, and nausea and vomiting, which are unpleasant and can be life threatening in a sedated patient. Such compounds may also produce physical dependence in a subject. Dependence appears to be related to the dose of narcotic analgesic taken and the period of time over which it is taken. Therefore, compounds which serve as either a replacement for, or as an adjunct to, narcotic analgesic treatment have utility in the treatment of pain, particularly, pain of the chronic type.

SUMMARY OF THE INVENTION

We have now discovered, surprisingly, in view of contrary reports from others (see, e.g., Boual, et al., *supra*), that amylin and amylin agonists, for example, the amylin agonist analogue ^{25,28,29}Pro-h-amylin (also referred to as "pramlintide" and previously referred to as "AC-0137"), can be used for analgesia in mammals.

The present invention is directed to novel methods for treating pain comprising the administration of an amylin or an amylin agonist, for example, the amylin agonist analogue ^{25,28,29}Pro-h-amylin, wherein the agonist is not a calcitonin. By "calcitonin" is meant the human peptide hormone calcitonin and species variations of it, such as rat calcitonin, salmon calcitonin, eel calcitonin (including

WO 98/26796

PCT/US97/23015

aminosuberic eel calcitonin). In one aspect, the invention is directed to a method of treating pain in a mammalian subject comprising administering to said subject an effective analgesic amount of an amylin or such an amylin agonist. By "effective analgesic amount" is meant an amount effective to reduce or eliminate pain in response to a stimulation that would normally be painful, and thus includes a complete analgesic effect (absence of pain) as well as what is sometimes referred to as "hypalgesic" effect or reduction in pain.

The term "amylin" is understood to include compounds such as those defined in U.S. Patent No. 5,234,906, issued August 10, 1993, for "Hyperglycemic Compositions," the contents of which are hereby incorporated by reference. For example, it includes the human peptide hormone referred to as amylin and secreted from the beta cells of the pancreas, and species variations of it. "Amylin agonist" is also a term known in the art, and refers to a compound which mimics effects of amylin. An amylin agonist may be a peptide or a non-peptide compound, and include amylin agonist analogues.

The term "amylin agonist analogue" is understood to refer to derivatives of an amylin which act as amylin agonists, normally, it is presently believed, by virtue of binding to or otherwise directly or indirectly interacting with an amylin receptor or other receptor or receptors with which amylin itself may interact to elicit a biological response. Useful amylin agonist analogues include those identified in an International Application, WPI Acc. No. 93-182488/22, entitled "New Amylin Agonist Peptides Used for Treatment and Prevention of Hypoglycemia and Diabetes Mellitus," the contents of which is also hereby incorporated by reference.

In preferred embodiment, the amylin agonist is an amylin agonist analogue, preferably, ^{25,28,29}Pro-h-amylin. In yet other preferred embodiments, the method further comprises administering a narcotic analgesic. By "narcotic analgesic" is meant the naturally occurring opium alkaloids, semi-synthetic and synthetic derivatives. Examples of such compounds include morphine, pentazocine, hydromorphone, oxymorphone, levorphanol, methadone, meperidine, anileridine, alphaprodine, fentanyl, codeine, oxycodone and hydrocodone. Preferred narcotic analgesics include morphine and pentazocine.

In another aspect, the invention is directed to a method of enhancing the analgesic activity of a narcotic analgesic (such as morphine or pentazocine) comprising administering said narcotic analgesic along with an amylin or an amylin agonist described herein. This synergism enables the use of lower doses of either or both drugs with a concomitant reduction in the risk of possible side effects. In a preferred embodiment, the amylin agonist is an amylin agonist analogue, preferably, ^{25,28,29}Pro-h-amylin.

In a related aspect, the invention features pharmaceutical compositions comprising a therapeutically effective amount of a mixture of (1) an amylin and/or an amylin agonist as described herein, preferably, the amylin agonist analogue, ^{25,28,29}Pro-h-amylin, and (2) a narcotic analgesic, preferably morphine or pentazocine.

Such compositions may include pharmaceutically acceptable salts of amylin or an amylin agonist, and/or pharmaceutically acceptable salts of a narcotic analgesic. Such compositions may further comprise a pharmaceutically acceptable carrier.

75331-42

Due to amylin's reported vasodilator action (Brain et al, Am. J. Pathl., 136:487-490 (1990)), amylin or an amylin agonist, either alone or in combination with an opiate or opiate agonist, may be especially useful in the treatment of vascular headache, such as migraine.

Thus, in one aspect, the present invention provides use of an amylin, or of an amylin analogue that acts as an agonist, in the manufacture of a medicament for the treatment or prevention of pain in a mammalian subject.

10 In another aspect, the present invention provides use of an amylin, or of an amylin analogue that acts as an agonist, for the treatment or prevention of pain in a mammalian subject.

15 In another aspect, the present invention provides use of an amylin, or of an amylin analogue that acts as an agonist, in the manufacture of a medicament for enhancing the analgesic activity of a narcotic analgesic, wherein the medicament is for use in combination with the narcotic analgesic.

20 In another aspect, the present invention provides use of an amylin, or of an amylin analogue that acts as an agonist, for enhancing the analgesic activity of a narcotic analgesic, wherein the amylin or amylin analogue is for use in combination with the narcotic analgesic.

25 In another aspect, the present invention provides A pharmaceutical composition comprising: (1) an amylin or an amylin analogue that acts as an agonist, or a pharmaceutically acceptable salt thereof; and (2) a narcotic analgesic, or a pharmaceutically acceptable salt thereof.

30 In another aspect, the present invention provides a commercial package comprising a pharmaceutical composition

75331-42

described herein, together with instructions for use for treating or preventing pain in a mammalian subject.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 shows the results of a time course study of the effect of amylin administration on number of writhes per 10 minutes, 5 minutes post acetic acid administration when 0.1 mg/kg amylin (closed circles) or saline (open circles) was injected intraperitoneally (ip) 5, 15, 30 and 60 minutes prior to acetic acid injection in mice. The error bars indicate mean $\pm$ standard error for each timepoint.

FIGURE 2 shows the results of a time course study of the effect of amylin administration on number of writhes per 10 minutes, 5 minutes post acetic acid administration when 0.1 mg/kg amylin (closed circles) or saline (open circles) was injected subcutaneously (sc) 5, 15, 30 and 60 minutes prior to acetic acid injection in mice. The error bars indicate mean $\pm$ standard error for each timepoint.

FIGURE 3 shows the results of a dose response study of the effect of administration of various doses (0.001 mg/kg, 0.003 mg/kg, 0.01 mg/kg, 1.0 mg/kg and 10.0 mg/kg) of amylin on the number of writhes per 10 minutes, 5 minutes post acetic acid administration when injected intraperitoneally (ip) 30 minutes prior to acetic acid injection in mice. The error bars indicate mean $\pm$ standard error for each timepoint.

FIGURE 4 shows the results of a dose response study of the effect of administration of various doses (0.001 mg/kg, 0.003 mg/kg, 0.01 mg/kg, 1.0 mg/kg and 10.0 mg/kg) of amylin on the number of writhes per 10 minutes, 5 minutes

WO 98/26796

PCT/US97/23015

post acetic acid administration when injected subcutaneously (sc) 30 minutes prior to acetic acid injection in mice. The error bars indicate mean $\pm$ standard error for each timepoint.

FIGURE 5 shows the percent inhibition (mean $\pm$ standard error) of number of writhes per 10 minutes, 5 minutes post acetic acid administration, in mice injected subcutaneously (sc) with .003 mg/kg amylin (closed squares) or various doses (0.01 mg/kg, 0.1 mg/kg or 3.0 mg/kg) of morphine (open squares), compared to saline controls.

FIGURE 6 shows the percent inhibition (mean $\pm$ standard error) of number of writhes per 10 minutes, 5 minutes post acetic acid administration, in mice injected subcutaneously (sc) with 0.01 mg/kg amylin (closed squares) or various doses (0.01 mg/kg, 0.1 mg/kg or 3.0 mg/kg) morphine (open squares), compared to saline controls.

FIGURE 7 is an isobologram plot characterizing 30% inhibition of writhing response.

FIGURE 8 is an isobologram plot characterizing 35% inhibition of writhing response.

FIGURE 9 is an isobologram plot characterizing 40% inhibition of writhing response.

FIGURE 10 is an isobologram plot characterizing 45% inhibition of writhing response.

DETAILED DESCRIPTION OF THE INVENTION

The rate of writhing activity in response to intraperitoneal injection of dilute acetic acid in mice is a commonly used surrogate of pain in animal studies of analgesia. As described in Example 1, rat amylin, when injected 30 minutes prior to acetic acid injection (both intraperitoneally and subcutaneously) in mice, significantly reduced the number of writhes in the 10-minute period

beginning 5 minutes after acetic acid injection compared to control ($P < 0.001$ for ip injection; $P < 0.01$ for sc injection). As indicated in Figures 1 and 2, the peak analgesic effect of amylin occurred at 30 minutes prior to acetic acid injection. Based on these studies, a 30-minute timepoint was selected for use in dose response studies.

As described in Example 2, various doses of amylin were evaluated for analgesic effect. The results shown Figures 3 and 4 indicate that the lowest effective dose of amylin in these experiments was 0.01 mg/kg ($P < 0.05$ sc; $P < 0.01$ ip). A dose of 0.1 mg/kg of amylin was also effective in inducing analgesia ($P < 0.01$ sc; $P < 0.05$ ip). At a higher doses administered (1 mg/kg and 10 mg/kg), amylin no longer exhibited significant analgesic activity ($P < 0.05$). These examples demonstrate the effectiveness of amylin and amylin agonists including amylin agonist analogues in averting nociceptive behavior (writhing) in mice injected with acetic acid. These results also indicate that the analgesic effectiveness of amylin or amylin agonists may be dose-dependent.

Example 3 describes experiments in which the analgesic activity of various dosages of amylin and morphine was examined. These experiments used the same mouse writhing assay procedures as in the experiments described in Examples 1 and 2. In one experiment, a dose of rat amylin which had been shown to be ineffective (0.003 mg/kg) in inducing analgesia under the experimental conditions in Example 2 (or saline control) was combined with 3 doses of morphine: 0.01 mg/kg, 0.1 mg/kg and 3.0 mg/kg. In another experiment, a dose of rat amylin which had been shown to be effective (0.01 mg/kg) in inducing analgesia under the experimental

conditions in Example 2 (or saline control) was combined with the same 3 doses of morphine.

The results shown in Figures 5 and 6 indicate that administration of morphine alone at doses of 0.01 mg/kg and 0.1 mg/kg did not significantly inhibit the writhing response ($P > 0.05$), but morphine alone was effective in significantly inhibiting the writhing response at a dose of 3.0 mg/kg ($P < 0.001$). Amylin plus morphine showed an increased efficacy in reducing analgesia compared to morphine alone at combinations of: (1) 0.003 mg/kg amylin plus 0.1 mg/kg morphine ($P < 0.05$); (2) 0.01 mg/kg amylin plus 0.01 mg/kg morphine ($P < 0.05$); and (3) 0.01 mg/kg amylin plus 0.1 mg/kg morphine ($P < 0.05$). Thus, a combination of a non-analgesic dose of amylin and a non-analgesic dose of morphine can provide an analgesic effect.

Example 4 describes analysis of the results of the writhing studies of Example 3 to further study the interaction between amylin and morphine. The results were graphed in isobolograms according to the method of Berebaum, J. Theor. Biol. 144:413 (1985). The isobologram is a quantitative method for measuring interactions between equieffective dosages of drugs to indicate synergy, additive effect or antagonism.

As shown in Figures 7-10, the interaction of amylin and morphine on acetic acid-induced writhing in mice is synergistic. That is, the resulting activity of amylin and morphine together is greater than the sum of the activities of the individual components.


Example 5 describes the analysis of additional narcotic analogesics following the methods of Examples 1-4.

Activity as amylin agonists can be ascertained by performing various screening assays, including the nucleus

accumbens receptor binding assay described below in Example R, the soleus muscle assay described below in Example S, a gastric emptying assay described below in Example T or U, or by the ability to induce hypocalcemia or reduce postprandial hyperglycemia in mammals, as described herein.

The nomenclature of various amylin agonist analogue compounds useful in the present invention and referred to herein can be used to indicate both the peptide that the sequence is based on and the modifications made to any basic peptide amylin sequence, such as human amylin. An amino acid preceded by a superscript number indicates that the named amino acid replaces the amino acid normally present at the amino acid position of the superscript in the basic amino acid sequence. For example, "¹⁸Arg^{25,28}Pro-h-amylin" refers to a peptide based on the sequence of "h-amylin" or "human-amylin" having the following substitutions: Arg replacing His at residue 18, Pro replacing Ala at residue 25, and Pro replacing Ser at residue 28. The term "des-¹Lys-h-amylin" refers to a peptide based on the sequence of human amylin, with the first, or N-terminal, amino acid deleted.


Amylin agonist analogues useful in this invention include amylin agonist analogues having the following amino acid sequence:



wherein A₁ is hydrogen, Lys, Ser, Ala, des-α-amino Lys, or acetylated Lys; B₁ is Ala, Ser or Thr; C₁ is Val, Leu or Ile; D₁ is His or Arg; E₁ is Ser or Thr; F₁ is Ser, Thr, Gln or Asn; G₁ is Asn, Gln or His; H₁ is Phe, Leu or Tyr; I₁ is Ala or Pro; J₁ is Ile, Val, Ala or Leu; K₁ is Ser, Pro, Leu, Ile

or Thr; L₁ is Ser, Pro or Thr; M₁ is Asn, Asp or Gln; X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage; and Z is hydroxy, amino, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, aryloxy or aralkyloxy; provided that (a) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Phe, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Ser, and M₁ is Asn; (b) when A₁ is Lys, B₁ is Ala, C₁ is Ile, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Pro, and M₁ is Asn; (c) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Thr, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Ala, J₁ is Ile, K₁ is Ser, L₁ is Pro, and M₁ is Asn; (d) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val, K₁ is Pro, L₁ is Pro, and M₁ is Asn; (e) when A₁ is Lys, B₁ is Ala, C₁ is Val, D₁ is His, E₁ is Ser, F₁ is Asn, G₁ is Asn, H₁ is Leu, I₁ is Pro, J₁ is Val, K₁ is Ser, L₁ is Pro and M₁ is Asn; or (f) when A₁ is Lys, B₁ is Thr, C₁ is Val, D₁ is Arg, E₁ is Ser, F₁ is Ser, G₁ is His, H₁ is Leu, I₁ is Ala, J₁ is Ala, K₁ is Leu, L₁ is Pro and M₁ is Asp; then one or more of any of A₁ to M₁ is not an L-amino acid and Z is not amino.

Preferred such amylin agonist analogues include amylin agonist analogues having the following amino acid sequence:



wherein

A₁ is Lys, Ala, Ser or hydrogen;

B₁ is Ala;

WO 98/26796

PCT/US97/23015

C₁ is Val;

D₁ is His or Arg;

E₁ is Ser;

F₁ is Ser;

G₁ is Asn;

H₁ is Phe;

I₁ is Ala or Pro;

J₁ is Ile, Val, Ala or Leu;

K₁ is Ser, Pro, Leu, Ile or Thr;

L₁ is Ser, Pro or Thr;

M₁ is Asn, Asp or Gln;

X and Y are independently selected residues having side chains which are chemically bonded to each other to form an intramolecular linkage; and Z is amino, hydroxy, alkylamino, dialkylamino, cycloalkylamino, arylamino, aralkylamino, alkyloxy, arlyoxy or aralkyloxy; provided that when A₁ is Lys, D₁ is His, I₁ is Ala, J₁ is Ile, K₁ is Ser and L₁ is Ser, then one or more of any of A₁ to M₁ is not an L-amino and Z is not amino.

Suitable side chains for X and Y include groups derived from alkyl sulfhydryls which may form disulfide bonds; alkyl acids and alkyl amines which may form cyclic lactams; alkyl aldehydes or alkyl halides and alkylamines which may condense and be reduced to form an alkyl amine bridge; or side chains which may be connected to form an alkyl, alkenyl, alkynyl, ether or thioether bond. Preferred alkyl chains include lower alkyl groups having from about 1 to about 6 carbon atoms.

As used herein, the following terms have the following meanings unless expressly stated to the contrary:

The term "alkyl" refers to both straight- and branched-chain alkyl groups. The term "lower alkyl" refers

to both straight- and branched-chain alkyl groups having a total of from 1 to 6 carbon atoms and includes primary, secondary and tertiary alkyl groups. Typical lower alkyls include, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl, n-hexyl, and the like.

The term "aryl" refers to carbocyclic aromatic groups of 6 to 14 carbon atoms such as phenyl and naphthyl, as well as heterocyclic aromatic groups containing 1 to 3 heteroatoms (nitrogen, oxygen, sulfur, etc.) such as pyridyl, triazolopyrazine, pyrimidine and the like.

The term "aralkyl" refers to an "aryl" group of 6 to 10 carbon atoms directly attached to an "alkyl" group of 1 to 4 carbon atoms and includes for example benzyl, p-chlorobenzyl, p-methylbenzyl, and 2-phenylethyl.

The term "cycloalkyl" refers to cyclic alkyl groups of 5 to 8 carbon atoms.

Biologically active derivatives of the above agonist analogues are also included within the scope of amylin agonist analogues useful in the present invention in which the stereochemistry of individual amino acids may be inverted from (L)/S to (D)/R at one or more specific sites. Also included within the scope of amylin agonist analogues useful in the present invention are the agonist analogues modified by glycosylation of Asn, Ser and/or Thr residues.

Biologically active agonist analogues of amylin which contain less peptide character are also included in the scope of amylin agonist analogues useful in the present invention. Such peptide mimetics may include, for example, one or more of the following substitutions for -CO-NH- amide bonds: depsipeptides (-CO-O-), iminomethylenes (-CH₂-NH-), trans-alkenes (-CH=CH-), β -enaminonitriles (-C(=CH-CN)-NH-),

thioamides (-CS-NH-), thiomethylenes (-S-CH₂- or -CH₂-S-), methylenes (-CH₂-CH₂-) and retro-amides (-NH-CO-).

The above-described amylin agonist analogues form salts with various inorganic and organic acids and bases. Such salts include salts prepared with organic and inorganic acids, for example, HCl, HBr, H₂SO₄, H₃PO₄, trifluoroacetic acid, acetic acid, formic acid, methanesulfonic acid, toluenesulfonic acid, maleic acid, fumaric acid and camphorsulfonic acid. Salts prepared with bases include, for example, ammonium salts, alkali metal salts (such as sodium and potassium salts) and alkali earth salts (such as calcium and magnesium salts). Acetate, hydrochloride, and trifluoroacetate salts are preferred.

The salts may be formed by conventional means, as by reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion on a suitable ion exchange resin.

The above-described amylin agonist analogues include various stereoisomers. In the preferred amylin agonist analogues, the chiral centers on the peptide backbone are all S.

Preferred amylin agonist analogue compounds include des-¹Lys-h-amylin, ²⁸Pro-h-amylin, ^{25,28,29}Pro-h-amylin, ¹⁸Arg^{25,28}Pro-h-amylin, and des-¹Lys¹⁸Arg^{25,28}Pro-h-amylin, all of which show amylin agonist activity in vivo in treated test animals, provoking marked hyperlactemia followed by hyperglycemia. In addition to having activities characteristic of amylin, certain preferred compounds have also been found to possess more desirable solubility and

stability characteristics when compared to human amylin. These preferred compounds include $^{25}\text{Pro}^{26}\text{Val}^{28,29}\text{Pro-h-amylin}$, $^{25,28,29}\text{Pro-h-amylin}$, and $^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$. Preferred human analogues of human amylin have substitutions of Arg at position 18, Leu at position 23, Val at position 26, Pro at position 25, Pro at position 28 and/or Pro at position 29, which may be missing the first amino acid (i.e., des-Lys compounds).

The methods of the present invention can employ an amylin, or an amylin agonist as described herein, including amylin agonist analogues, for example, amylin agonist analogues such as $^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$, des- $^1\text{Lys}^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$, $^{18}\text{Arg}^{25,28,29}\text{Pro-h-amylin}$, des- $^1\text{Lys}^{18}\text{Arg}^{25,28,29}\text{Pro-h-amylin}$, $^{25,28,29}\text{Pro-h-amylin}$, des- $^1\text{Lys}^{25,28,29}\text{Pro-h-amylin}$, and $^{25}\text{Pro}^{26}\text{Val}^{25,28}\text{Pro-h-amylin}$. Examples of other suitable amylin agonist analogues include: $^{23}\text{Leu}^{25}\text{Pro}^{26}\text{Val}^{28,29}\text{Pro-h-amylin}$; $^{23}\text{Leu}^{25}\text{Pro}^{26}\text{Val}^{28}\text{Pro-h-amylin}$; des- $^1\text{Lys}^{23}\text{Leu}^{25}\text{Pro}^{26}\text{Val}^{28}\text{Pro-h-amylin}$; $^{18}\text{Arg}^{23}\text{Leu}^{25}\text{Pro}^{26}\text{Val}^{28}\text{Pro-h-amylin}$; $^{18}\text{Arg}^{23}\text{Leu}^{25,28,29}\text{Pro-h-amylin}$; $^{18}\text{Arg}^{23}\text{Leu}^{25,28}\text{Pro-h-amylin}$; $^{17}\text{Ile}^{23}\text{Leu}^{25,28,29}\text{Pro-h-amylin}$; $^{17}\text{Ile}^{25,28,29}\text{Pro-h-amylin}$; des- $^1\text{Lys}^{17}\text{Ile}^{23}\text{Leu}^{25,28,29}\text{Pro-h-amylin}$; $^{17}\text{Ile}^{18}\text{Arg}^{23}\text{Leu-h-amylin}$; $^{17}\text{Ile}^{18}\text{Arg}^{23}\text{Leu}^{26}\text{Val}^{29}\text{Pro-h-amylin}$; $^{17}\text{Ile}^{18}\text{Arg}^{23}\text{Leu}^{25}\text{Pro}^{26}\text{Val}^{28,29}\text{Pro-h-amylin}$; $^{13}\text{Thr}^{21}\text{His}^{23}\text{Leu}^{26}\text{Ala}^{28}\text{Leu}^{29}\text{Pro}^{31}\text{Asp-h-amylin}$; $^{13}\text{Thr}^{21}\text{His}^{23}\text{Leu}^{26}\text{Ala}^{29}\text{Pro}^{31}\text{Asp-h-amylin}$; des- $^1\text{Lys}^{13}\text{Thr}^{21}\text{His}^{23}\text{Leu}^{26}\text{Ala}^{28}\text{Pro}^{31}\text{Asp-h-amylin}$; $^{13}\text{Thr}^{18}\text{Arg}^{21}\text{His}^{23}\text{Leu}^{26}\text{Ala}^{29}\text{Pro}^{31}\text{Asp-h-amylin}$; $^{13}\text{Thr}^{18}\text{Arg}^{21}\text{His}^{23}\text{Leu}^{28,29}\text{Pro}^{31}\text{Asp-h-amylin}$; and, $^{13}\text{Thr}^{18}\text{Arg}^{21}\text{His}^{23}\text{Leu}^{25}\text{Pro}^{26}\text{Ala}^{28,29}\text{Pro}^{31}\text{Asp-h-amylin}$.

Still further amylin agonists including amylin agonist analogues are disclosed in the above-noted WPI Acc. No. 93-182488/22, "New Amylin Agonist Peptides Used for

WO 98/26796

PCT/US97/23015

Treatment and Prevention of Hypoglycemia and Diabetes Mellitus."

The activity of amylin agonists may be evaluated using certain biological assays described herein. The receptor binding assay can identify putative amylin agonists and antagonists and can be used to evaluate binding, while the soleus muscle assay distinguishes between amylin agonist and antagonist action in glucose metabolism in muscle. Gastric emptying assays such as those discussed below can also be used to screen for amylin agonists. Effects of amylin or amylin agonists on pain can be identified, evaluated, or screened for using the methods described in Examples 1-5 below, or other art-known or equivalent methods for determining analgesic effect. See, e.g., Tjolsen, et al., Handbook of Lab Animal Science, Vol. II, Chapter 12, Animal Models in Pain Research (Svendsen, Ed., CRC Press, 1994). Preferred, amylin agonist compounds exhibit activity in the receptor binding assay on the order of less than about 1 to 5 nM, preferably less than about 1 nM and more preferably less than about 50 pM. In the soleus muscle assay, preferred amylin agonist compounds show EC₅₀ values on the order of less than about 1 to 10 micromolar.

The receptor binding assay, a competition assay which measures the ability of compounds to bind specifically to membrane-bound amylin receptors, is described in United States Patent No. 5,264,372, issued November 23, 1993, the disclosure of which is incorporated herein by reference. A preferred source of the membrane preparations used in the assay is the basal forebrain which comprises membranes from the nucleus accumbens and surrounding regions. Compounds being assayed compete for binding to these receptor preparations with ¹²⁵I Bolton Hunter rat amylin. Competition

curves, wherein the amount bound (B) is plotted as a function of the log of the concentration of ligand are analyzed by computer, using analyses by nonlinear regression to a 4-parameter logistic equation (Inplot program; GraphPAD Software, San Diego, California) or the ALLFIT program of DeLean *et al.* (ALLFIT, Version 2.7 (NIH, Bethesda, MD 20892)). Munson and Rodbard, *Anal. Biochem.* 107:220-239 (1980).

Assays of biological activity of amylin agonists, in the soleus muscle may be performed using previously described methods (Leighton, B. and Cooper, *Nature*, 335:632-635 (1988); Cooper, *et al.*, *Proc. Natl. Acad. Sci. USA* 85:7763-7766 (1988)). In summary, amylin agonist activity may be assessed by measuring the inhibition of insulin-stimulated glycogen synthesis in soleus muscle.

Methods of measuring the rate of gastric emptying are disclosed in, for example, Young *et al.*, *Diabetologia*, 38(6):642-648 (1995). Briefly, in a phenol red method, conscious rats receive by gavage an acoloric gel containing methyl cellulose and a phenol red indicator. Twenty minutes after gavage, animals are anesthetized using halothane, the stomach exposed and clamped at the pyloric and lower esophageal sphincters, removed and opened into an alkaline solution. Stomach content may be derived from the intensity of the phenol red in the alkaline solution, measured by absorbance at a wavelength of 560 nm. Briefly, in a tritiated glucose method, conscious rats are gavaged with tritiated glucose in water. The rats are gently restrained by the tail, the tip of which is anesthetized using lidocaine. Tritium in the plasma separated from tail blood is collected at various timepoints and detected in a beta

WO 98/26796

PCT/US97/23015

counter. Test compounds are administered 1 minute before gavage.

Amylins and peptide amylin agonists may be prepared using standard solid-phase peptide synthesis techniques and preferably an automated or semiautomated peptide synthesizer. Typically, using such techniques, an α -N-carbamoyl protected amino acid and an amino acid attached to the growing peptide chain on a resin are coupled at room temperature in an inert solvent such as dimethylformamide, N-methylpyrrolidinone or methylene chloride in the presence of coupling agents such as dicyclohexylcarbodiimide and 1-hydroxybenzotriazole in the presence of a base such as diisopropylethylamine. The α -N-carbamoyl protecting group is removed from the resulting peptide-resin using a reagent such as trifluoroacetic acid or piperidine, and the coupling reaction repeated with the next desired N-protected amino acid to be added to the peptide chain. Suitable N-protecting groups are well known in the art, with t-butyloxycarbonyl (tBoc) and fluorenylmethoxycarbonyl (Fmoc) being preferred herein.

The solvents, amino acid derivatives and 4-methylbenzhydryl-amine resin used in the peptide synthesizer may be purchased from Applied Biosystems Inc. (Foster City, CA). The following side-chain protected amino acids may be purchased from Applied Biosystems, Inc.: Boc-Arg(Mts), Fmoc-Arg(Pmc), Boc-Thr(Bzl), Fmoc-Thr(t-Bu), Boc-Ser(Bzl), Fmoc-Ser(t-Bu), Boc-Tyr(BrZ), Fmoc-Tyr(t-Bu), Boc-Lys(Cl-Z), Fmoc-Lys(Boc), Boc-Glu(Bzl), Fmoc-Glu(t-Bu), Fmoc-His(Trt), Fmoc-Asn(Trt), and Fmoc-Gln(Trt). Boc-His(BOM) may be purchased from Applied Biosystems, Inc. or Bachem Inc. (Torrance, CA). Anisole, methylsulfide, phenol, ethanedithiol, and thioanisole may be obtained from Aldrich Chemical Company (Milwaukee, WI). Air Products and Chemicals

WO 98/26796

PCT/US97/23015

(Allentown, PA) supplies HF. Ethyl ether, acetic acid and methanol may be purchased from Fisher Scientific (Pittsburgh, PA).

Solid phase peptide synthesis may be carried out with an automatic peptide synthesizer (Model 430A, Applied Biosystems Inc., Foster City, CA) using the NMP/HOBt (Option 1) system and TboC or Fmoc chemistry (see, Applied Biosystems User's Manual for the ABI 430A Peptide Synthesizer, Version 1.3B July 1, 1988, section 6, pp. 49-70, Applied Biosystems, Inc., Foster City, CA) with capping. Boc-peptide-resins may be cleaved with HF (-5°C to 0°C, 1 hour). The peptide may be extracted from the resin with alternating water and acetic acid, and the filtrates lyophilized. The Fmoc-peptide resins may be cleaved according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc., 1990, pp. 6-12). Peptides may be also be assembled using an Advanced Chem Tech Synthesizer (Model MPS 350, Louisville, Kentucky). Peptides may be purified by RP-HPLC (preparative and analytical) using a Waters Delta Prep 3000 system. A C4, C8 or C18 preparative column (10 μ , 2.2 x 25 cm; Vydac, Hesperia, CA) may be used to isolate peptides, and purity may be determined using a C4, C8 or C18 analytical column (5 μ , 0.46 x 25 cm; Vydac). Solvents (A=0.1% TFA/water and B=0.1% TFA/CH₃CN) may be delivered to the analytical column at a flowrate of 1.0 ml/min and to the preparative column at 15 ml/min. Amino acid analyses may be performed on the Waters Pico Tag system and processed using the Maxima program. Peptides may be hydrolyzed by vapor-phase acid hydrolysis (115°C, 20-24 h). Hydrolysates may be derivatized and analyzed by standard methods (Cohen, et al., The Pico Tag Method: A Manual of Advanced Techniques for Amino Acid Analysis, pp. 11-52, Millipore Corporation, Milford, MA

(1989)). Fast atom bombardment analysis may be carried out by M-Scan, Incorporated (West Chester, PA). Mass calibration may be performed using cesium iodide or cesium iodide/glycerol. Plasma desorption ionization analysis using time of flight detection may be carried out on an Applied Biosystems Bio-Ion 20 mass spectrometer.

Peptide compounds useful in the invention may also be prepared using recombinant DNA techniques, using methods now known in the art. See, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d Ed., Cold Spring Harbor (1989). Non-peptide compounds useful in the present invention may be prepared by art-known methods.

The compounds referenced above may form salts with various inorganic and organic acids and bases. Such salts include salts prepared with organic and inorganic acids, for example, HCl, HBr, H₂SO₄, H₃PO₄, trifluoroacetic acid, acetic acid, formic acid, methanesulfonic acid, toluenesulfonic acid, maleic acid, fumaric acid and camphorsulfonic acid. Salts prepared with bases include ammonium salts, alkali metal salts, e.g., sodium and potassium salts, and alkali earth salts, e.g., calcium and magnesium salts. Acetate, hydrochloride, and trifluoroacetate salts are preferred. The salts may be formed by conventional means, as by reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion on a suitable ion exchange resin.

Compositions useful in the invention may conveniently be provided in the form of formulations suitable for parenteral (including intravenous, intramuscular and

WO 98/26796

PCT/US97/23015

subcutaneous) or nasal or oral administration. In some cases, it will be convenient to provide an amylin agonist and another analgesic agent, for example, an opiate, such as morphine, in a single composition or solution for administration together. In other cases, it may be more advantageous to administer another analgesic agent separately from said amylin or amylin agonist. A suitable administration format may best be determined by a medical practitioner for each patient individually. Suitable pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, e.g., Remington's Pharmaceutical Sciences by E.W. Martin. See also Wang, Y.J. and Hanson, M.A. "Parenteral Formulations of Proteins and Peptides: Stability and Stabilizers," Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2S (1988).

Compounds useful in the invention can be provided as parenteral compositions for injection or infusion. They can, for example, be suspended in an inert oil, suitably a vegetable oil such as sesame, peanut, olive oil, or other acceptable carrier. Preferably, they are suspended in an aqueous carrier, for example, in an isotonic buffer solution at a pH of about 5.6 to 7.4. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH buffering agents. Useful buffers include for example, sodium acetate/acetic acid buffers. A form of repository or "depot" slow release preparation may be used so that therapeutically effective amounts of the preparation are delivered into the bloodstream over many hours or days following transdermal injection or delivery.

WO 98/26796

PCT/US97/23015

The desired isotonicity may be accomplished using sodium chloride or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol, polyols (such as mannitol and sorbitol), or other inorganic or organic solutes. Sodium chloride is preferred particularly for buffers containing sodium ions.

If desired, solutions of the above compositions may be thickened with a thickening agent such as methyl cellulose. They may be prepared in emulsified form, either water in oil or oil in water. Any of a wide variety of pharmaceutically acceptable emulsifying agents may be employed including, for example, acacia powder, a non-ionic surfactant (such as a Tween), or an ionic surfactant (such as alkali polyether alcohol sulfates or sulfonates, e.g., a Triton).

Compositions useful in the invention are prepared by mixing the ingredients following generally accepted procedures. For example, the selected components may be simply mixed in a blender or other standard device to produce a concentrated mixture which may then be adjusted to the final concentration and viscosity by the addition of water or thickening agent and possibly a buffer to control pH or an additional solute to control tonicity.

For use by the physician, the compositions will be provided in dosage unit form containing an amount of an amylin or amylin agonist, for example, an amylin agonist analogue compound with or without another analgesic agent which will be effective in one or multiple doses to control pain at the selected level. Therapeutically effective amounts of an amylin or amylin agonist, such as an amylin agonist analogue, for use in the control of pain are those that decrease pain. As will be recognized by those in the

field, an effective amount of therapeutic agent will vary with many factors including the age and weight of the patient, the patient's physical condition, the action to be obtained and other factors.

The effective single, divided or continuous analgesic doses of the compounds, for example, including $^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$, $\text{des-}^{1}\text{Lys}^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$, $^{18}\text{Arg}^{25,28,29}\text{Pro-h-amylin}$, $\text{des-}^{1}\text{Lys}^{18}\text{Arg-}^{25,28,29}\text{Pro-h-amylin}$, $^{25,28,29}\text{Pro-h-amylin}$, $\text{des-}^{1}\text{Lys}^{25,28,29}\text{Pro-h-amylin}$, and $^{25}\text{Pro}^{26}\text{Val}^{25,28}\text{Pro-h-amylin}$, will typically be in the range of 0.01 or 0.03 to about 5 mg/day, preferably about 0.01 or 0.5 to 2 mg/day and more preferably about 0.01 or 0.1 to 1 mg/day, for a 70 kg patient, administered in a single, divided or continuous doses. The exact dose to be administered is determined by the attending clinician and is dependent upon where the particular compound lies within the above quoted range, as well as upon the age, weight and condition of the individual. Administration should begin at the first sign of pain. Administration may be by injection or infusion, preferably intravenous, subcutaneous or intramuscular. Orally active compounds may be taken orally, however dosages should be increased 5-10 fold.

Generally, in treating or preventing pain, the compounds of this invention may be administered to patients in need of such treatment in a dosage ranges similar to those given above, however, the compounds may be administered more frequently, for example, one, two, or three times a day or continuously.

To assist in understanding the present invention, the following Examples are included which describe the results of a series of experiments. The experiments relating to this invention should not, of course, be construed as

specifically limiting the invention and such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are considered to fall within the scope of the invention as described herein and hereinafter claimed.

EXAMPLE 1

Time Course of Amylin Action

Male Swiss Webster mice (NIH/Sw) obtained from Harlan (Madison, Wisconsin) and weighing 20-35 g were group housed with free access to food and water and maintained in a stable environment (12:12 light:dark cycle; $23 \pm 1^{\circ}\text{C}$). All animals were habituated to the test room for at least one day prior to any experimentation, and were tested once between 07:30 and 14:00.

Rat amylin was synthesized by standard solid phase peptide synthesis methods. Morphine was obtained from Steris Laboratories (Phoenix, Arizona). All drugs were dissolved in physiological saline, and given in a dose volume 10 ml/kg body weight.

The mouse writhing assay procedure used was a modification of a procedure disclosed in Hendershot and Forsaith, *J. Pharmacol. Expt. Therap.*, 125:237-240 (1959). Each mouse was allowed to habituate to the observation box for at least 15 minutes prior to testing. Each mouse was given an intraperitoneal injection of a 2% acetic acid solution to produce a writhing reaction, characterized by a wave on contraction of the abdominal musculature followed by the extension of the hind limbs. The number of writhes per animal was counted during a 10-minute interval starting 5 minutes after acetic acid injection.

WO 98/26796

PCT/US97/23015

0.1 mg/kg of rat amylin was administered subcutaneously (sc) or intraperitoneally (ip) at 5, 15, 30 and 60 minutes prior to acetic acid injection in mice. Saline injections were used as a negative control.

The results are shown in Figures 1 and 2. Amylin, when injected 30 minutes prior to acetic acid injection (both ip and sc), significantly reduced the number of writhes per 10-minute period 5 minutes post acetic acid injection compared to the saline-treated animals ($P < 0.001$ for ip injection; $P < 0.01$ for sc injection). The peak effect of amylin occurred at 30 minutes prior to acetic acid injection. The 30-minute timepoint was selected for use in the dose response studies described in Example 2 below.

EXAMPLE 2

Dose Response of Amylin Action

The same experimental procedures used in the experiments described in Example 1 were used in the following dose response studies. Subcutaneous and intraperitoneal injections of rat amylin (0.001, 0.003, 0.01, 0.1, 1.0 and 10.0 mg/kg) were given 30 minutes prior to acetic acid injection. Saline was used as a negative control. Morphine (1.0 mg/kg) was used as a positive control. The results are shown in Figures 3 and 4. The lowest effective analgesic dose of amylin in these experiments was 0.01 mg/kg ($P < 0.05$ sc; $P < 0.01$ ip). A dose of 0.1 mg/kg was also effective ($P < 0.01$ sc; $P < 0.05$ ip). At higher doses administered (1 mg/kg and 10 mg/kg), amylin no longer exhibited significant analgesic activity ($P > 0.05$).

EXAMPLE 3

Analgesic Actions of Amylin and Morphine

The ability of amylin to enhance the analgesic effect of morphine is shown in these experiments, using the same experimental procedures described in Example 1. In one experiment, a dose of rat amylin which had been shown to be ineffective (0.003 mg/kg) in inducing analgesia under the experimental conditions in Example 2 was combined with 3 doses of morphine: 0.01 mg/kg, 0.1 mg/kg and 3.0 mg/kg. In another experiment, a dose of rat amylin which had been shown to be effective (0.01 mg/kg) in inducing analgesia under the experimental conditions in Example 2 (or saline control) was combined with the same 3 doses of morphine.

The results are shown in Figures 5 and 6. Administration of morphine alone at doses of 0.01 mg/kg and 0.1 mg/kg did not significantly inhibit the writhing response ($P > 0.05$). Morphine alone was effective in significantly inhibiting the writhing response at a dose of 3.0 mg/kg ($P < 0.001$). Amylin plus morphine showed an increased efficacy in reducing analgesia compared to morphine alone at combinations of: (1) 0.003 mg/kg amylin plus 0.1 mg/kg morphine ($P < 0.05$); (2) 0.01 mg.kg amylin plus 0.01 mg/kg morphine ($P < 0.05$); and (3) 0.01 mg.kg amylin plus 0.1 mg/kg morphine ($P < 0.05$). Thus, a combination of a non-analgesic dose of amylin and a non-analgesic dose of morphine can provide an analgesic effect.

EXAMPLE 4

Isobologram Analysis of Interaction of Amylin and Morphine

To further study the interaction between amylin and morphine, the results of the writhing studies were graphed in isobolograms according to the method of Berebaum, "The

WO 98/26796

PCT/US97/23015

unexpected effect of a combination of agents: the general solution," J. Theor. Biol. 114:413 (1985). The isobologram is a quantitative method for measuring interactions between dosages of drugs that are equieffective in relationship to a common pharmacological endpoint. In this instance, the writhing test was used to estimate a common level of analgesic dose-ratio combination. In the isobolographic figures, areas of dose additional, synergism and/or antagonism are clearly defined by reference to a theoretical straight (addition) line connecting the points on each axis. According to the isobologram theory, any points falling under the curve represents unexpectedly enhanced analgesic activity and any points located above the line would represent unexpectedly diminished analgesic activity.

The synergistic interaction of amylin and morphine on acetic acid-induced writhing in mice is demonstrated by the data in Figures 7-10. In the isobolograms the analgesic effect of amylin alone is presented on the ordinate and that of morphine alone is presented on the abscissa. The percent of inhibition for each compound and the combination of amylin and morphine [ANDREW/SUNIL: WHERE DID DATA FOR DIFFERENT DOSAGES OF MORPHINE ALONE COME FROM?] were derived from sigmoidal dose-response curves from the data depicted in Figure 4 (dose-response for amylin given sc) and Figure 5 (combination of amylin (3 μ g/kg, sc) and morphine (0.01, 0.1 and 3 μ g/kg, 1P). Figures 7-10 depict the synergistic interaction of amylin and morphine at 30%, 35%, 40% and 45%, respectively, indicating the existence of unexpectedly enhanced analgesic activity of combinations of amylin and morphine. That is, the resulting activity of amylin and

WO 98/26796

PCT/US97/23015

morphine together is greater than the activity expected from the sum of the activities of the individual components.

EXAMPLE 5

Analysis of Additional Narcotic Analgesics

Following the procedures of Examples 1-4, inclusive, experiments are performed substituting an equi-analgesic amount of each of: hydromorphone, oxymorphone, levorphanol, methadone, meperidine, alphapradine, fentanyl, codeine, oxycodone or hydrocodone for the morphine of Examples 1-4.

[ANDREW: OTHER COMPOUNDS?]

EXAMPLE A

Preparation of ²⁸Pro-human-Amylin

Solid phase synthesis of this analogue of human ("h-") amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Ac^m-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid hydrofluoric acid ("HF") in the presence of dimethylsulfide and anisole. The ²⁸Pro-h-amylin was purified by preparative HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: (M+1)/e=3914.

EXAMPLE B**Preparation of $^{25}\text{Pro}^{26}\text{Val}^{28,29}\text{Pro-h-Amylin}$**

Solid phase synthesis of this amylin analogue using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7} -[disulfide]amylin-MBHA-resin was obtained by treatment with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The $^{25}\text{Pro}^{26}\text{Val}^{28,29}\text{Pro-h-amylin}$ was purified by preparative HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: (M+1)/e=3936.

EXAMPLE C**Preparation of $^{2,7}\text{Cyclo-}[^2\text{Asp}, ^7\text{Lys}]\text{-h-Amylin}$**

Solid phase synthesis of this amylin analogue using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. ^2Asp and ^7Lys were introduced with Boc- $^2\text{Asp}(\text{Fmoc})$ -OH and Boc- $^7\text{Lys}(\text{Fmoc})$ -OH. Following selective side-chain deprotection with piperidine the side-chain to side-chain ($^2\text{Asp-}^7\text{Lys}$) cyclization was carried out using benzotriazol-1-yl-oxy-tris(dimethylamino)-phosphonium hexafluorophosphate (BOP reagent). Cyclization was as described in Di Maio, et al., *J. Med. Chem.*, 33:661-667 (1990); Felix et al., *Int. J. Pept. Prot. Res.*, 32:441 (1988). The $^{2,7}\text{cyclo-}[^2\text{Asp}, ^7\text{Lys}]\text{amylin-MBHA-resin}$ obtained after cyclization was cleaved with liquid HF in the presence

WO 98/26796

PCT/US97/23015

of dimethylsulfide and anisole. The ^{2,7}cyclo-[²Asp,⁷Lys]-h-amylin was purified by preparative HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. FAB mass spec:
(M+1)/e=3925.

EXAMPLE D

Preparation of des-¹Lys-h-Amylin

Solid phase synthesis of des-¹Lys-h-amylin (also represented as ²⁻³⁷h-amylin) using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Ac^m-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The des-¹Lys-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec:
(M+H)⁺=3,775.

EXAMPLE E

Preparation of ¹Ala-h-Amylin

Solid phase synthesis of ¹Ala-h-amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide

synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Acm-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The ¹Ala-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: (M+H)⁺=3,847.

EXAMPLE F

Preparation of ¹Ser-h-Amylin

Solid phase synthesis of ¹Ser-h-amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Acm-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The ¹Ser-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: (M+H)⁺=3,863.

EXAMPLE G**Preparation of ²⁹Pro-h-Amylin**

Solid phase synthesis of this analogue of human amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Acm-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The ²⁹Pro-h-amylin was purified by preparative HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: (M+H)⁺=3916.

EXAMPLE H**Preparation of ^{25,28}Pro-h-Amylin**

Solid phase synthesis of ^{25,28}Pro-h-amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Acm-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The ^{25,28}Pro-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid

WO 98/26796

PCT/US97/23015

analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: $(M+H)^+ = 3,939$.

EXAMPLE I

Preparation of des-¹Lys^{25,28}Pro-h-Amylin

Solid phase synthesis of des-¹Lys^{25,28}Pro-h-amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Ac^m-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The des-¹Lys^{25,28}Pro-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: $(M+H)^+ = 3,811$.

EXAMPLE J

Preparation of ¹⁸Arg^{25,28}Pro-h-Amylin

Solid phase synthesis of ¹⁸Arg^{25,28}Pro-h-amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Ac^m-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence

of dimethylsulfide and anisole. The $^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$ was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: $(\text{M}+\text{H})^+=3,959$.

EXAMPLE K

Preparation of des- $^1\text{Lys}^{18}\text{Arg}^{25,28}\text{Pro-h-Amylin}$

Solid phase synthesis of des- $^1\text{Lys}^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$ using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7} -[disulfide]amylin-MBHA-resin was obtained by treatment of Ac^m-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The des- $^1\text{Lys}^{18}\text{Arg}^{25,28}\text{Pro-h-amylin}$ was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: $(\text{M}+\text{H})^+=3,832$.

EXAMPLE L

Preparation of $^{18}\text{Arg}^{25,28,29}\text{Pro-h-Amylin}$

Solid phase synthesis of $^{18}\text{Arg}^{25,28,29}\text{Pro-h-amylin}$ using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The

^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Acm-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The ¹⁸Arg^{25,28,29}Pro-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: (M+H)⁺=3,971.

EXAMPLE M

Preparation of des-¹Lys¹⁸Arg^{25,28,29}Pro-h-Amylin

Solid phase synthesis of des-¹Lys¹⁸Arg^{25,28,29}Pro-h-amylin using methylbenzhydrylamine anchor-bond resin and N^α-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Acm-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The des-¹Lys¹⁸Arg^{25,28,29}Pro-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: (M+H)⁺=3,843.

EXAMPLE N**Preparation of ^{25,28,29}Pro-h-Amylin**

Solid phase synthesis of ^{25,28,29}Pro-h-amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Ac^m-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The ^{25,28,29}Pro-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: (M+H)⁺=3,949.

EXAMPLE O**Preparation of des-¹Lys^{25,28,29}Pro-h-Amylin**

Solid phase synthesis of des-¹Lys^{25,28,29}Pro-h-amylin using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection was carried out by standard peptide synthesis methods. The ^{2,7}-[disulfide]amylin-MBHA-resin was obtained by treatment of Ac^m-protected cysteines with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization was achieved the resin and side chain protecting groups were cleaved with liquid HF in the presence of dimethylsulfide and anisole. The des-¹Lys^{25,28,29}Pro-h-amylin was purified by preparative reversed-phase HPLC. The peptide was found to be homogeneous by analytical HPLC and capillary electrophoresis and the

WO 98/26796

PCT/US97/23015

structure confirmed by amino acid analysis and sequence analysis. The product gave the desired mass ion. FAB mass spec: $(M+H)^+=3,823$.

EXAMPLE P

Preparation of des-¹Lys²⁵Pro²⁶Val^{28,29}Pro-h-Amylin

Solid phase synthesis of this h-amylin analogue using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection is carried out by standard peptide synthesis methods, and the ^{2,7}-[disulfide]amylin-MBHA-resin obtained by treatment with thallium (III) trifluoroacetate in trifluoroacetic acid. After cyclization is achieved the resin and side chain protecting groups are cleaved with liquid HF in the presence of dimethylsulfide and anisole. The des-¹Lys²⁵Pro²⁶Val^{28,29}Pro-h-amylin is then purified by preparative HPLC.

EXAMPLE Q

Preparation of [(D)-¹¹Arg]-Amylin

Solid phase synthesis of this amylin analogue using methylbenzhydrylamine anchor-bond resin and N^a-Boc/benzyl-side chain protection is carried out by standard peptide synthesis methods. (D)-¹¹Arg is introduced with Boc-(D)-¹¹Arg(Mtr)-OH. The ^{2,7}-[disulfide]amylin-MBHA-resin, obtained by treatment with thallium (III) trifluoroacetate in trifluoroacetic acid, is cyclized and the resin and side chain protecting groups are cleaved with liquid HF in the presence of dimethylsulfide and anisole. The [(D)-¹¹Arg]-amylin is then purified by preparative HPLC.

EXAMPLE R

WO 98/26796

PCT/US97/23015

Receptor Binding Assay

Evaluation of the binding of compounds to amylin receptors was carried out as follows. ^{125}I -rat amylin (Bolton-Hunter labeled at the N-terminal lysine) was purchased from Amersham Corporation (Arlington Heights, IL). Specific activities at time of use ranged from 1950 to 2000 Ci/mmol. Unlabeled peptides were obtained from BACHEM Inc. (Torrance, CA) and Peninsula Laboratories (Belmont, CA).

Male Sprague-Dawley rats (200-250) grams were sacrificed by decapitation. Brains were removed to cold phosphate-buffered saline (PBS). From the ventral surface, cuts were made rostral to the hypothalamus, bounded laterally by the olfactory tracts and extending at a 45° angle medially from these tracts. This basal forebrain tissue, containing the nucleus accumbens and surrounding regions, was weighed and homogenized in ice-cold 20 mM HEPES buffer (20 mM HEPES acid, pH adjusted to 7.4 with NaOH at 23°C). Membranes were washed three times in fresh buffer by centrifugation for 15 minutes at $48,000 \times g$. The final membrane pellet was resuspended in 20 mM HEPES buffer containing 0.2 mM phenylmethylsulfonyl fluoride (PMSF).

To measure ^{125}I -amylin binding, membranes from 4 mg original wet weight of tissue were incubated with ^{125}I -amylin at 12-16 pM in 20 mM HEPES buffer containing 0.5 mg/ml bacitracin, 0.5 mg/ml bovine serum albumin, and 0.2 mM PMSF. Solutions were incubated for 60 minutes at 23°C . Incubations were terminated by filtration through GF/B glass fiber filters (Whatman Inc., Clifton, NJ) which had been presoaked for 4 hours in 0.3% polyethyleneimine in order to reduce nonspecific binding of radiolabeled peptides. Filters were washed immediately before filtration with 5 ml cold PBS, and immediately after filtration with 15 ml cold PBS. Filters

WO 98/26796

PCT/US97/23015

were removed and radioactivity assessed in a gamma-counter at a counting efficiency of 77%. Competition curves were generated by measuring binding in the presence of 10^{-12} to 10^{-6} M unlabeled test compound and were analyzed by nonlinear regression using a 4-parameter logistic equation (Inplot program; GraphPAD Software, San Diego).

In this assay, purified human amylin binds to its receptor at a measured IC_{50} of about 50 pM. Results for test compounds are set forth in Table I, showing that each of the compounds has significant receptor binding activity.

EXAMPLE S

Soleus Muscle Assay

Determination of amylin agonist activity of compounds was carried out using the soleus muscle assay as follows. Male Harlan Sprague-Dawley rats of approximately 200g mass were used in order to maintain mass of the split soleus muscle less than 40mg. The animals were fasted for 4 hours prior to sacrifice by decapitation. The skin was stripped from the lower limb which was then pinned out on corkboard. The tendo achilles was cut just above os calcis and *m. gastrocnemius* reflected out from the posterior aspect of the tibia. *M. soleus*, a small 15-20mm long, 0.5mm thick flat muscle on the bone surface of *m. gastrocnemius* was then stripped clear and the perimysium cleaned off using fine scissors and forceps. *M. soleus* was then split into equal parts using a blade passed antero-posteriorly through the belly of the muscle to obtain a total of 4 muscle strips from each animal. After dissecting the muscle from the animal, it was kept for a short period in physiological saline. It was not necessary that the muscle be held under tension as this

had no demonstrable effects on radioglucose incorporation into glycogen.

Muscles were added to 50mL Erlenmeyer flasks containing 10mL of a pregassed Krebs-Ringer bicarbonate buffer containing (each liter) NaCl 118.5 mmol (6.93g), KCl 5.94 mmol (443mg), CaCl₂ 2.54 mmol (282mg), MgSO₄ 1.19 mmol (143mg), KH₂PO₄ 1.19 mmol (162mg), NaHCO₃ 25 mmol (2.1g), 5.5mmol glucose (1g) and recombinant human insulin (Humulin-R, Eli Lilly, IN) and the test compound, as detailed below. pH at 37°C was verified as being between 7.1 and 7.4. Muscles were assigned to different flasks so that the 4 muscle pieces from each animal were evenly distributed among the different assay conditions. The incubation media were gassed by gently blowing carbogen (95% O₂, 5% CO₂) over the surface while being continuously agitated at 37°C in an oscillating water bath. After a half-hour "preincubation" period, 0.5μCi of U-¹⁴C-glucose was added to each flask which was incubated for a further 60 minutes. Each muscle piece was then rapidly removed, blotted and frozen in liquid N₂, weighed and stored for subsequent determination of ¹⁴C-glycogen.

¹⁴C-glycogen determination was performed in a 7mL scintillation vial. Each frozen muscle specimen was placed in a vial and digested in 1mL 60% potassium hydroxide at 70°C for 45 minutes under continuous agitation. Dissolved glycogen was precipitated out onto the vial by the addition of 3mL absolute ethanol and overnight cooling at -20°C. The supernatant was gently aspirated, the glycogen washed again with ethanol, aspirated and the precipitate dried under vacuum. All ethanol is evaporated to avoid quenching during scintillation counting. The remaining glycogen was

WO 98/26796

PCT/US97/23015

redissolved in 1mL water and 4mL scintillation fluid and counted for ^{14}C .

The rate of glucose incorporation into glycogen (expressed in $\mu\text{mol/g/hr}$) was obtained from the specific activity of ^{14}C -glucose in the 5.5mM glucose of the incubation medium, and the total ^{14}C counts remaining in the glycogen extracted from each muscle. Dose/response curves were fitted to a 4-parameter logistic model using a least-squares iterative routine (ALLFIT, v2.7, NIH, MD) to derive EC_{50} 's. Since EC_{50} is log-normally distributed, it is expressed $\pm$ standard error of the logarithm. Pairwise comparisons were performed using t-test based routines of SYSTAT (Wilkinson, "SYSTAT: the system for statistics," SYSTAT Inc., Evanston IL (1989)).

Dose response curves were generated with muscles added to media containing 7.1nM (1000 $\mu\text{U/mL}$) insulin and each test compound added at final (nominal) concentrations of 0, 1, 3, 10, 30, 100, 300 and 1000nM. Each assay also contained internal positive controls consisting of a single batch of archived rat amylin, lyophilized and stored at -70°C .

Human amylin is a known hyperglycemic peptide, and EC_{50} measurements of amylin preparations in the soleus muscle assay range typically from about 1-10 nM, although some commercial preparations which are less than 90% pure have higher EC_{50} 's due to the presence of contaminants that result in a lower measured activity. Results for test compounds are set forth in Table I, showing that each of the compounds has amylin activity.

TABLE I

	Receptor Binding	Soleus
Muscle		

WO 98/26796

PCT/US97/23015

<u>EC₅₀ (nM)</u>	<u>Assay IC₅₀ (pM)</u>	<u>Assay</u>
1) ²⁸ Pro-h-Amylin	15.0	2.64
2) ²⁵ Pro ²⁶ Val ^{28,29} Pro-h-Amylin	18.0	4.68
3) ^{2,7} Cyclo- [² Asp, ⁷ Lys] -h-Amylin	310.0	6.62
4) ²⁻³⁷ h-Amylin	236.0	1.63
5) ¹ Ala-h-Amylin	148.0	12.78
6) ¹ Ser-h-Amylin	33.0	8.70
7) ²⁹ Pro-h-Amylin	64.0	3.75
8) ^{25,28} Pro-h-Amylin	26.0	13.20
9) des- ¹ Lys ^{25,28} Pro-h-Amylin	85.0	7.70
10) ¹⁸ Arg ^{25,28} Pro-h-Amylin	32.0	2.83
11) des- ¹ Lys ¹⁸ Arg ^{25,28} Pro-h-Amylin	82.0	3.77
12) ¹⁸ Arg ^{25,28,29} Pro-h-Amylin	21.0	1.25
13) des- ¹ Lys ¹⁸ Arg ^{25,28,29} Pro-h-Amylin	21.0	1.86
14) ^{25,28,29} Pro-h-Amylin	10.0	3.71
15) des- ¹ Lys ^{25,28,29} Pro-h-Amylin	14.0	4.15

EXAMPLE T**PHENOL RED GASTRIC EMPTYING ASSAY**

Gastric emptying was measured using a modification (Plourde et al., Life Sci. 53:857-862 (1993)) of the original method of Scarpignato et al. (Arch. Int. Pharmacodyn. Ther. 246:286-295 (1980)). Briefly, conscious rats received by gavage. 1.5 mL of an acoloric gel containing 1.5% methyl cellulose (M-0262, Sigma Chemical Co., St. Louis, MO) and 0.05% phenol red indicator. Twenty minutes after gavage, rats were anesthetized using 5% halothane, the stomach exposed and clamped at the pyloric and lower esophageal

WO 98/26796

PCT/US97/23015

sphincters using artery forceps, removed and opened into an alkaline solution which was made up to a fixed volume. Stomach content was derived from the intensity of the phenol red in the alkaline solution, measured by absorbance at a wavelength of 560 nm. In most experiments, the stomach was clear. In other experiments, particulate gastric contents were centrifuged to clear the solution for absorbance measurements. Where the diluted gastric contents remained turbid, the spectroscopic absorbance due to phenol red was derived as the difference between that present in alkaline vs acetified diluent. In separate experiments on 7 rats, the stomach and small intestine were both excised and opened into an alkaline solution. The quantity of phenol red that could be recovered from the upper gastrointestinal tract within 29 minutes of gavage was $89 \pm 4\%$; dye which appeared to bind irrecoverably to the gut luminal surface may have accounted for the balance. To compensate for this small loss, percent of stomach contents remaining after 20 minutes were expressed as a fraction of the gastric contents recovered from control rats sacrificed immediately after gavage in the same experiment. Percent gastric emptying contents remaining = $(\text{absorbance at 20 min}) / (\text{absorbance at 0 min})$. Dose response curves for gastric emptying were fitted to a 4-parameter logistic model using a least-squares iterative routine (ALLFIT, v2.7, NIH, Bethesda, MD) to derive ED_{50} s. Since ED_{50} is log-normally distributed, it is expressed $\pm$ standard error of the logarithm. Pairwise comparisons were performed using one-way analysis of variance and the Student-Newman-Keuls multiple comparisons test (Instat v2.0, GraphPad Software, San Diego, CA) using $P < 0.05$ as the level of significance.

In dose response studies, rat amylin (Bachem, Torrance, CA) dissolved in 0.15M saline, was administered as

WO 98/26796

PCT/US97/23015

a 0.1 mL subcutaneous bolus in doses of 0, 0.01, 0.1, 1, 10 or 100 μ g 5 minutes before gavage in Harlan Sprague Dawley (non-diabetic) rats fasted 20 hours and diabetic BB rats fasted 6 hours. When subcutaneous amylin injections were given 5 minutes before gavage with phenol red indicator, there was a dose-dependent suppression of gastric emptying (data not shown). Suppression of gastric emptying was complete in normal HSD rats administered 1 μ g of amylin, and in diabetic rats administered 10 μ g ($P = 0.22, 0.14$). The ED_{50} for inhibition of gastric emptying in normal rats was 0.43 μ g (0.60 nmol/kg) $\pm$ 0.19 log units, and was 2.2 μ (2.3 nmol/kg) $\pm$ 0.18 log units in diabetic rats.

EXAMPLE U

TRITIATED GLUCOSE GASTRIC EMPTYING ASSAY

Conscious, non-fasted, Harlan Sprague Dawley rats were restrained by the tail, the tip of which was anesthetized using 2% lidocaine. Tritium in plasma separated from tail blood collected 0, 15, 30, 60, 90 and 120 minutes after gavage was detected in a beta counter. Rats were injected subcutaneously with 0.1 mL saline containing 0, 0.1, 0.3, 1, 10 or 100 μ g of rat amylin 1 minute before gavage ($n=8,7,5,5,5$, respectively). After gavage of saline pre-injected rats with tritiated glucose, plasma tritium increased rapidly ($t_{1/2}$ of about 8 minutes) to an asymptote that slowly declined. Subcutaneous injection with amylin dose-dependently slowed and/or delayed the absorption of the label. Plasma tritium activity was integrated over 30 minutes to obtain the areas under the curve plotted as a function of amylin dose. The ED_{50} derived from the logistic fit was 0.35 μ g of amylin.

75331-42

CLAIMS:

1. Use of an amylin, or of an amylin analogue that acts as an agonist, in the manufacture of a medicament for the treatment or prevention of pain in a mammalian subject.
- 5 2. Use of an amylin, or of an amylin analogue that acts as an agonist, for the treatment or prevention of pain in a mammalian subject.
3. The use according to claim 1 or 2, wherein said amylin analogue is based on human amylin.
- 10 4. The use according to claim 3, wherein said amylin analogue is ^{25,28,29}Pro-h-amylin.
5. The use according to any one of claims 1 to 4, wherein said medicament is for use in combination with a narcotic analgesic.
- 15 6. The use according to claim 5, wherein said narcotic analgesic is morphine, pentazocine, hydromorphone, oxymorphone, levorphanol, methadone, meperidine, anileridine, alphaprodine, fentanyl, codeine, oxycodone or hydrocodone.
- 20 7. The use according to claim 6, wherein said narcotic analgesic is morphine.
8. The use according to claim 6, wherein said narcotic analgesic is pentazocine.
9. Use of an amylin, or of an amylin analogue that
25 acts as an agonist, in the manufacture of a medicament for enhancing the analgesic activity of a narcotic analgesic, wherein the medicament is for use in combination with the narcotic analgesic.

75331-42

10. Use of an amylin, or of an amylin analogue that acts as an agonist, for enhancing the analgesic activity of a narcotic analgesic, wherein the amylin or amylin analogue is for use in combination with the narcotic analgesic.

5 11. The use of claim 9 or 10, wherein said amylin analogue is based on human amylin.

12. The use according to claim 11, wherein said amylin analogue is ^{25,28,29}Pro-h-amylin.

13. The use according to any one of claims 9 to 12,
10 wherein said narcotic analgesic is morphine, pentazocine, hydromorphone, oxymorphone, levorphanol, methadone, meperidine, anileridine, alphaprodine, fentanyl, codeine, oxycodone or hydrocodone.

14. The use according to claim 13, wherein said
15 narcotic analgesic is morphine.

15. The use according to claim 13, wherein said narcotic analgesic is pentazocine.

16. A pharmaceutical composition comprising: (1) an amylin or an amylin analogue that acts as an agonist, or a
20 pharmaceutically acceptable salt thereof; and (2) a narcotic analgesic, or a pharmaceutically acceptable salt thereof.

17. The pharmaceutical composition according to claim 16 further comprising a pharmaceutically acceptable carrier.

25 18. The pharmaceutical composition according to claim 16 or 17 wherein said amylin analogue is based on human amylin.

19. The pharmaceutical composition according to claim 18 wherein said amylin analogue is ^{25,28,29}Pro-h-amylin.

75331-42

20. The pharmaceutical composition according to any one of claims 16 to 19, wherein said narcotic analgesic is morphine, pentazocine, hydromorphone, oxymorphone, levorphanol, methadone, meperidine, anileridine,
5 alphaprodine, fentanyl, codeine, oxycodone or hydrocodone.

21. The pharmaceutical composition according to claim 20, wherein said narcotic analgesic is morphine.

22. The pharmaceutical composition according to claim 20, wherein said narcotic analgesic is pentazocine.

10 23. A commercial package comprising a pharmaceutical composition according to any one of claims 16 to 22, together with instructions for use for treating or preventing pain in a mammalian subject.


SMART & BIGGAR

OTTAWA, CANADA

PATENT AGENTS

1/10


FIG. 1.



ns = not significant, ** = P < 0.01, *** = P < 0.001

2/10

FIG. 2.



ns = not significant, ** = P < 0.01

FIG. 3.




FIG. 4






FIG. 5.



ns= NOT SIGNIFICANT, * = P<0.05, ** = P<0.01, *** = P<0.001

6/10

FIG. 6.




Mean $\pm$ SE % inhibition of the writhing response after s.c administration of saline (open squares) or rat-amylin, 3 μ g/kg (closed squares) in animals treated with 0.01, 0.1 and 3 mg/kg doses of morphine (i.p); N $\geq$ 6

* = P < 0.05

7/10

FIG. 7.



Mean $\pm$ SE % inhibition of the writhing response after s.c administration of saline (open squares) or rat-amylin, 10 μ g/kg (closed squares) in animals treated with 0.01, 0.1 and 3 mg/kg doses of morphine (i.p); N $\geq$ 8

* = P < 0.05

8/10

FIG. 8.






FIG. 9.



ns=NOT SIGNIFICANT * =P<0.05, ** =P<0.01, ***=P<0.001

FIG. 10.



MORPHINE AMYLIN OR SALINE GIVEN sc, NALOXONE GIVEN ip, 30 MINUTES PRIOR TO ACETIC ACID INJECTION

ns=NOT SIGNIFICANT, **=P<0.01 ***=P<0.0001