

(19)



(11)

EP 1 523 528 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
22.02.2012 Bulletin 2012/08

(51) Int Cl.:
C09B 69/10 (2006.01) **C09B 57/00** (2006.01)
G03F 7/00 (2006.01) **C07D 487/04** (2006.01)
C07D 209/00 (2006.01) **C07D 209/00** (2006.01)

(21) Application number: **03764989.4**

(86) International application number:
PCT/EP2003/007638

(22) Date of filing: **15.07.2003**

(87) International publication number:
WO 2004/009710 (29.01.2004 Gazette 2004/05)

(54) **POLYMERISABLE DIKETOPYRROLOPYRROLES, USE OF SUCH COMPOUNDS IN COLOUR FILTERS AND POLYMERS PREPARED FROM THESE COMPOUNDS**

POLYREAKTIONSFÄHIGE DIKETOPYRROLOPYRROLE, DEREN VERWENDUNG IN FARBFILTERN UND DARAUS HERGESTELLTE POLYMERE

DICETOPYRROLOPYRROLES POLYMERISABLES, UTILISATION DE CES COMPOSES DANS DES FILTRES COULEURS ET POLYMERES PREPARES A BASE DE CES COMPOSES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**

(30) Priority: **22.07.2002 EP 02405640**

(43) Date of publication of application:
20.04.2005 Bulletin 2005/16

(73) Proprietor: **BASF SE
67056 Ludwigshafen (DE)**

(72) Inventors:
• **ADAM, Jean-Marie
F-68128 Rosenau (FR)**
• **DE KEYZER, Gerardus
CH-4125 Riehen (CH)**

(56) References cited:
EP-A- 0 337 951 EP-A- 0 654 711
EP-A- 0 787 730 EP-A- 0 787 731
EP-A- 0 811 625 EP-A- 0 894 798
WO-A-99/54332

EP 1 523 528 B1


Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description


[0001] The present invention relates to the use of polymerisable diketopyrrolopyrroles in colour filters, which can themselves be used for example in electro-optical systems such as TV screens, liquid crystal displays, charge coupled devices, plasma displays or electroluminescent displays and the like. These may be, for example, active (twisted nematic) or passive (supertwisted nematic) ferroelectric displays or light-emitting diodes.

[0002] EP-A-811625 discloses rheology improving agents based on N-substituted diaryldiketopyrrolopyrroles and a process for preparing 2,5-bis(hydroxymethyl)-3,6-diarylpyrrolo-pyrrole-1,4-dione. EP-A-894798 discloses diketopyrrolopyrroles N-substituted via a spacer with a polymerizable group and polymers thereof. EP-A-654711 discloses compositions comprising diketopyrrolopyrroles and their use is the production of color Filters


[0003] Accordingly, in one aspect the present invention relates to colour filters comprising a transparent substrate and a layer comprising from 1 to 75% by weight, preferably from 5 to 50% by weight, with particular preference from 25 to 40% by weight, based on the overall weight of the layer, of a diketopyrrolopyrrole of the general formula (I) dispersed in a high molecular mass organic material:



wherein A and B independently of one another are a group of the formula




or



wherein

R₃ and R₄ independently of one another are hydrogen, halogen, C₁-C₁₈alkyl, C₁-C₁₈alkoxy, -NR₁₈R₁₇, -CONHR₁₈, -COOR₁₉, -SO₂NH-R₂₀, C₁-C₁₈alkoxycarbonyl, C₁-C₁₈alkylaminocarbonyl, -CN, -NO₂, trifluoromethyl, C₅-C₇cycloalkyl, -C=N-(C₁-C₁₈alkyl),



5

imidazolyl, pyrazolyl, triazolyl, piperazinyl, pyrrolyl, oxazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, morpholinyl, piperidinyl or pyrrolidinyl,

G is $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{CH}=\text{N}-$, $-\text{N}=\text{N}-$, $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{CONH}-$ or $-\text{NR}_9-$,

10 R_5 and R_6 independently of one another are hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_{18} alkoxy or $-\text{CN}$,

R_7 and R_8 independently of one another are hydrogen, halogen or C_1 - C_6 alkyl and R_9 is hydrogen or C_1 - C_6 alkyl,

R_1 and R_2 are independently of each other C_1 - C_{18} alkyl, C_1 - C_{18} alkyl, which is interrupted one or more times by O or S, C_6 - C_{12} aryl, C_7 - C_{12} aralkyl, or a group of the formula $-\text{C}(\text{O})\text{OR}_{10}$, wherein R_{10} is C_1 - C_{18} alkyl, C_5 - C_{10} cycloalkyl, C_6 - C_{12} aryl, or C_7 - C_{12} aralkyl, or

15 a group of the formula




wherein

20 X_2 is an alkylene, arylene, aralkylene or cycloalkylene spacer containing optionally one or more groups $-\text{O}-$, $-\text{S}-$, $-\text{NR}_{14}-$, $-\text{CO}-$, $-\text{CONH}-$, $-\text{CONR}_{15}-$, or $-\text{COO}-$ as linking bridge,

X_3 is $-\text{OH}$, $-\text{NH}_2$, $-\text{C}(\text{R}_{11})=\text{CH}_2$, $-\text{OC}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$, $-\text{C}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$, C_5 - C_7 cycloalkenyl,

25



30

or

$-\text{OC}(\text{O})-\text{N}-\text{X}_4-\text{N}-\text{C}(\text{O})-\text{O}-\text{X}_5-\text{O}-\text{C}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$ wherein

R_{11} is hydrogen, or C_1 - C_4 alkyl, or halogen,

R_{12} is hydrogen, C_1 - C_4 alkyl, or halogen,

35 R_{13} is hydrogen, C_1 - C_4 alkyl, or C_6 - C_{12} aryl,

R_{14} and R_{15} are independently of each other hydrogen, C_1 - C_8 alkyl, or C_6 - C_{12} aryl,

R_{16} , R_{17} , R_{18} and R_{20} are independently of each other hydrogen, C_1 - C_{18} alkyl, C_6 - C_{12} aryl, or C_1 - C_{12} aralkyl,

R_{19} is C_1 - C_{18} alkyl, C_6 - C_{12} aryl, or C_7 - C_{12} aryl, and

X_4 and X_5 are independently of each other an alkylene, arylene, aralkylene or cycloalkylene spacer,

40 R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 can also be a group of formula



wherein

45 X_1 is a single bond, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{CONH}-$, $-\text{COO}-$, $-\text{SO}_2-\text{NH}-$, or $-\text{SO}_2-\text{O}-$, and

X_2 and X_3 are as defined above,


with the proviso that at least one, preferably two, of the groups of the formula (II) and/or (III) is present per molecule.

[0004] X_1 is preferably a single bond, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{CONH}-$, or $-\text{SO}_2-\text{NH}-$.

50 **[0005]** X_2 is preferably an alkylene spacer containing optionally one or more, in particular one or two groups, $-\text{O}-$, $-\text{S}-$, $-\text{NR}_{14}-$, $-\text{CO}-$, $-\text{CONH}-$, $-\text{CONR}_{15}-$, or $-\text{COO}-$ as linking bridge. The number of carbon atoms of the alkylene spacer is preferably greater than 4, more preferred greater than 6 and most preferred in the range of 8 to 16.

[0006] X_3 is preferably $-\text{OH}$, $-\text{NH}_2$, $-\text{OC}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$, $-\text{C}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$, or $-\text{OC}(\text{O})-\text{N}-\text{X}_4-\text{N}-\text{C}(\text{O})-\text{O}-\text{X}_5-\text{O}-\text{C}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$; wherein R_{12} is hydrogen, or methyl, and X_4 and X_5 are as defined above.

55 **[0007]** Preferably the diketopyrrolopyrroles have the general formula



15 wherein R₁ and R₂ are independently of each other a group of the formula



wherein

20 X₂ is an alkylene, arylene, aralkylene or cycloalkylene spacer containing optionally one or more groups -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO- as linking bridge,

X₃ is OH, NH₂, -C(R₁₁)=CH₂, -OC(O)-C(R₁₂)=CH₂, -C(O)-C(R₁₂)=CH₂, or -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂; wherein

R₁₁ is hydrogen, or methyl,

25 R₁₂ is hydrogen, or methyl,

R₁₄ and R₁₅ are independently of each other hydrogen, C₁-C₈alkyl, or C₆-C₁₂aryl, and X₄ and X₅ are as defined above;

R₃ and R₄ independently of one another are C₁-C₁₈alkyl, C₁-C₁₈alkoxy, -NR₁₆R₁₇, -CONHR₁₆, -COOR₁₉, -SO₂NH-R₂₀, C₁-C₁₈alkoxycarbonyl, C₁-C₁₈alkylaminocarbonyl, wherein R₁₆, R₁₇, R₁₈, R₁₉ and R₂₀ are C₁-C₁₈alkyl, wherein diketopyrrolopyrroles are more preferred, wherein R₁ and R₂ are independently of each other a radical of the formula




wherein

X₂ is C₁-C₁₈alkylene and


35 X₃ is -NH₂, -OH, -CH=CH₂, -C(CH₃)=CH₂, -OCO-CH=CH₂ or -OCO-C(CH₃)=CH₂, -CO-CH=CH₂ or -CO-C(CH₃)=CH₂ and/or

wherein R₃ and R₄ are independently of each other C₁-C₁₈alkylmercapto, C₁-C₁₈alkoxy, or -NR₁₆R₁₇, wherein one of the groups R₁₆ and R₁₇ is hydrogen and the other is C₁-C₁₈alkyl or both groups R₁₆ and R₁₇ are independently of each other C₁-C₁₈alkyl; or -


40 of the general formula



55 in which R₂₁ and R₂₂ are independently of one another hydrogen, C₁-C₁₈alkyl, C₁-C₁₀alkyl which is interrupted one or more times by O or S, C₇-C₁₂aralkyl or a group of the formula



in which R₅ is C₁-C₁₈alkyl,
R₂₃ and R₂₄ independently of one another are a group of formula



wherein

X₁ is a single bond, -O-, -S-, -NH-, -CONH-, -COO-, -SO₂-NH-, or -SO₂-O-,

X₂ is an alkylene, arylene, aralkylene or cycloalkylene spacer containing optionally one or more groups -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO- as linking bridge,


X₃ is -OH, -NH₂, -C(R₁₁)=CH₂, -OC(O)-C(R₁₂)=CH₂, -C(O)-C(R₁₂)=CH₂, or -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂; wherein

R₁₁ is hydrogen, or methyl,

R₁₂ is hydrogen, or methyl,

R₁₄ and R₁₅ are independently of each other hydrogen, C₁-C₈alkyl, or C₆-C₁₂aryl, and

X₄ and X₅ are independently of each other an alkylene, arylene, aralkylene or cycloalkylene spacer, wherein a diketopyrrolopyrrole is more preferred, wherein R₂₃ and R₂₄ independently of one another are a group of formula



wherein

X₁ is -S-, -SO₂NH- or -NH-,

X₂ is a C₁-C₁₈alkylene group, and

X₃ is -OH, -NH₂, -CH=CH₂, -C(CH₃)=CH₂, -OCO-CH=CH₂ or -OCO-C(CH₃)=CH₂, -CO-CH=CH₂, or -CO-C(CH₃)=CH₂, and/or

diketopyrrolopyrroles, wherein R₂₁ and R₂₂ independently of one another are hydrogen, or C₁-C₁₈alkyl.

[0008] In contrast to conventional pigments the diketopyrrolopyrroles of formula I do not tend to aggregate and, hence, show very good dispersibility. Color filters prepared by using the diketopyrrolopyrroles of formula I have high transparence and pure hue. In addition, they facilitate adjustment of color points and enable a large choice of shades.

[0009] The pigments will generally be used in the manufacture of colour filters as a dispersion in an organic solvent or water. There are several ways to manufacture these colour filters, which follow two mainstreams:

- Direct patterning during applying;
- Patterning after applying the pigment.

Direct patterning can be obtained by several printing techniques, such as impact (off-set, flexography, stamping, letter-press etc.) as well as non-impact (ink jet techniques).

Other direct patterning techniques are based on lamination processes, electronic discharging processes like electro-deposition and some special colour proofing methods, like the so-called Chromalin™ process (DuPont).

[0010] For impact printing techniques, the pigment may be dispersed in water or organic solvents by standard de-agglomeration methods (Skandex, Dynamill, Dispermat and the like) in the presence of a dispersant and a polymeric binder to produce an ink. Any dispersion technique known in the field, including the choice of solvent, dispersant and binder, can be used. The type of ink and its viscosity depend on the application technique and are well-known to the skilled artisan. Most usual binders, to which the invention is of course not limited, are (meth)acrylates, epoxies, PVA, polyimids, Novolak systems and the like as well as combinations of these polymers.

The ink dispersion then can be printed on all kind of standard printing machines. Curing of the binder system is preferably achieved by a heating process. The three colours can be applied at once or in different printing steps with intermediate drying and/or curing steps, for example one colour at the time in three printing steps.

Inks for use in ink jet, for example piezo or bubble jet, can be prepared likewise. They generally contain a pigment dispersed in water and/or one or a mixture of many hydrophilic organic solvents in combination with a dispersant and a binder.

For ink jet printing a standard ink jet printer can be used or a dedicated printer can be built in order to optimize for example the printing speed etc.

[0011] For lamination techniques, like thermal transfer and the like, a web system has to be made: The pigment is

dispersed in a solvent or water with dispersant and binder and coated on a foil and dried. The pigment/binder system can be patternwise or uniformly transferred to a colour filter substrate with the help of energy (UV, IR, heat, pressure etc.). Depending on the technique used, the colourant for example may be transferred alone (dye diffusion or sublimation transfer), or the colourant dispersion may be entirely transferred including the binder (wax transfer).

For electrodeposition, the pigment has to be dispersed in water together with an ionized polymer. By means of an electrical current, the ionized polymer is deionized at the anode or the cathode and, being insoluble then, deposited together with the pigments. This can be done on patterned or patternwise shielded, by a photoresist, (transparent) photoconductors like ITO etc.

The Chromalin™ process makes use of a photosensitive material, deposited on a colour filter substrate. The material becomes tacky upon UV exposure. The so called 'toner', comprising a mixture or compound of pigment and polymer, is distributed on the substrate and sticks on the tacky parts. This process has to be done three to four times for R,G,B and eventually black.

Patterning after applying is a method based mostly on the known photoresist technology, wherein the pigment is dispersed in the photoresist composition. Other methods are indirect patterning with the help of a separate photoresist or lamination techniques.

The pigment may be dispersed into photoresists by any standard method such as described above for the printing processes. The binder systems may also be identical. Further suitable compositions are described for example in EP 654711, WO 98/45756 or WO 98/45757.

Photoresists comprise a photoinitiator and a poly-crosslinkable monomer (negative radical polymerization), a material to crosslink the polymers itself (for example a photoacid generator or the like) or a material to chemically change the solubility of the polymer in certain developing media. This process, however, can also be done with heat (for example using thermal arrays or an NIR beam) instead of UV, in the case of some polymers which undergo chemical changes during heating processes, resulting in changes of solubility in the mentioned developing media. A photoinitiator is then not needed.

The photosensitive or heat sensible material is coated on a colour filter substrate, dried and UV(or heat) irradiated, sometimes again baked (photoacid generators) and developed with a developing medium (mostly a base). In this last step only the non-exposed (negative systems) or only the exposed (positive systems) parts are washed away, giving the wanted pattern. This operation has to be repeated for all the colours used.

Photosensitive lamination techniques are using the same principle, the only difference being the coating technique. A photosensitive system is applied as described above, however on a web instead of a colour filter substrate. The foil is placed on the colour filter substrate and the photosensitive layer is transferred with the help of heat and/or pressure.

Indirect processes, with the above mentioned polymeric binders without a photosensitive component, make use of an extra photoresist, coated on top of the pigmented resist. During the patterning of the photoresist, the pigmented resist is patterned as well. The photoresist has to be removed afterwards.

More details about the manufacture of colour filters can be found in text books, reviews and other scientific articles. The skilled artisan will associate the instant invention with the use of any such known technique as well.

For example, which is of course in no way limitative, substantially colourless methacrylic resins are commonly used in colour filters, examples thereof which are known to the skilled artisan being copolymers of aromatic methacrylates with methacrylic acid of M_w from 30'000 to 60'000. Such resins are highly appropriated to make films by spin-coating.

The colour filters of the invention contain the pigment compositions of the invention judiciously in a concentration of from 1 to 75% by weight, preferably from 5 to 50% by weight, with particular preference from 25 to 40% by weight, based on the overall weight of the pigmented layer.

The invention therefore likewise provides a colour filter comprising a transparent substrate and a layer comprising from 1 to 75% by weight, preferably from 5 to 50% by weight, with particular preference from 25 to 40% by weight, based on the overall weight of the layer, of a pigment composition of the invention or the individual components of said composition dispersed in a high molecular mass organic material. The substrate is preferably essentially colourless ($T \geq 95\%$ all over the visible range from 400 to 700 nm).

The instant printing inks or photoresists for making colour filters contain the pigment compositions of the invention judiciously in a concentration of from 0.01 to 40% by weight, preferably from 1 to 25% by weight, with particular preference from 5 to 10% by weight, based on the overall weight of the printing ink or photoresist.

The invention therefore likewise provides a composition for making colour filters comprising from 0.01 to 40% by weight, preferably from 1 to 25% by weight, with particular preference from 5 to 10% by weight, based on the overall weight of the composition, of a pigment composition of the invention dispersed therein.

This pigment composition also may additionally contain other colorants of different structure. The additional components will shift the mixture's spectrum hypsochromically or batho-chromically depending on their own hue. The skilled artisan will appreciate by himself which colorants can additionally be used, and in which amounts, depending on the desired colour. In certain cases, it is advantageous to use the inventive compositions in mixture or in combination with other additives such as wetting agents, surfactants, defoamers, antioxidants, UV absorbers, light stabilizers, plastisizers, or

general texture improving agents and so forth. Generally such additives can be used in a concentration from about 0.1 to 25 percent, preferably from about 0.2 to 15 % and most preferably from about 0.5 to 8 %, by weight based on the total weight of (a), (b) and (c).

Suitable surfactants include anionic surfactants such as alkylbenzene- or alkylnaphthalene-sulfonates, alkylsulfosuccinates or naphthalene formaldehyde sulfonates; cationic surfactants including, for example, quaternary salts such as benzyl tributyl ammonium chloride; or nonionic or amphoteric surfactants such as polyoxyethylene surfactants and alkyl- or amidopropyl betaines, respectively.

Suitable texture improving agents are, for example, fatty acids such as stearic acid or behenic acid, and fatty amines such as laurylamine and stearylamine. In addition, fatty alcohols or ethoxylated fatty alcohols, polyols such as aliphatic 1,2-diols or epoxidized soy bean oil, waxes, resin acids and resin acid salts may be used for this purpose.

Suitable UV stabilizers are, for example, the known benzotriazole derivatives (see, for example, US-B1-6,184,375, column 20, line 48 to column 21, line 15) and 2-(2-hydroxyphenyl)-1,3,5-triazines (see, for example, US-B1-6,184,375, column 22, line 41 to 65) known under the trade name TINUVIN® or CIBA® Fast H Liquid an aryl sulfonated benzotriazole, both being products of CIBA Specialty Chemicals Corporation.

[0012] The term "halogen" is generally iodine, fluorine, bromine or chlorine, preferably bromine or chlorine.

[0013] C_1 - C_4 alkyl group is methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl or tert-butyl; C_1 - C_8 alkyl is, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-amyl, tert-amyl or hexyl; C_1 - C_{18} alkyl is, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-amyl, tert-amyl, hexyl, heptyl, octyl, 2-ethylhexyl, nonyl, decyl, dodecyl, tetradecyl, hexadecyl or octadecyl.

[0014] The term "alkylene" means in general linear or branched C_1 - C_{18} alkylene, wherein examples of preferred linear representatives are for example $-(CH_2)_4-$, $-(CH_2)_5-$, $-(CH_2)_6-$, $-(CH_2)_7-$, $-(CH_2)_8-$, $-(CH_2)_9-$, $-(CH_2)_{10}-$, $-(CH_2)_{11}-$, $-(CH_2)_{12}-$, $-(CH_2)_{13}-$, $-(CH_2)_{14}-$, $-(CH_2)_{15}-$, $-(CH_2)_{16}-$, $-(CH_2)_{17}-$, $-(CH_2)_{18}-$, preferably C_4 - C_{16} alkylene such as $-(CH_2)_4-$, $-(CH_2)_5-$, $-(CH_2)_6-$, $-(CH_2)_7-$, $-(CH_2)_8-$, $-(CH_2)_9-$, $-(CH_2)_{10}-$, $-(CH_2)_{11}-$, or $-(CH_2)_{12}-$.

[0015] The "alkoxy group" in C_1 - C_{18} alkoxy or in C_1 - C_{18} alkoxycarbonyl can be linear or branched and is for example methoxy, ethoxy, n-propoxy, isopropoxy, butyloxy, hexyloxy, decyloxy, dodecyloxy, hexadecyloxy or octadecyloxy, preferably C_1 - C_8 alkoxy such as methoxy, ethoxy, n-propoxy, isopropoxy, butyloxy, hexyloxy, or octyloxy.

[0016] C_1 - C_{18} alkylmercapto is, for example, methylmercapto, ethylmercapto, propylmercapto, butylmercapto, octylmercapto, decylmercapto, hexadecylmercapto or octadecylmercapto.

[0017] C_1 - C_{18} alkylamino is, both alone and in C_1 - C_{18} alkylaminocarbonyl, for example methylamino, ethylamino, propylamino, hexylamino, decylamino, hexadecylamino or octadecylamino, preferably C_1 - C_6 alkylamino such as methylamino, ethylamino, propylamino or hexylamino.

[0018] C_5 - C_{10} cycloalkyl is preferably C_5 - C_7 cycloalkyl, such as, for example, cyclopentyl or cyclohexyl, especially cyclohexyl.


[0019] C_5 - C_7 cycloalkenyl is mono- or bicyclic cycloalkenyl, for example cyclopentenyl, cyclohexenyl or norbornenyl.

[0020] " C_6 - C_{12} aryl" is typically phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, which may be unsubstituted or substituted by, for example, one or more C_1 - C_4 alkyl groups, C_1 - C_4 alkoxy groups, or halogen atoms.

[0021] " C_7 - C_{12} aralkyl" is typically benzyl, 2-benzyl-2-propyl, β -phenyl-ethyl, α,α -dimethylbenzyl, ω -phenyl-butyl, or ω,ω -dimethyl- ω -phenyl-butyl, in which both the aliphatic hydrocarbon group and aromatic hydrocarbon group may be unsubstituted or substituted by, for example, one or more C_1 - C_4 alkyl groups, C_1 - C_4 alkoxy groups, or halogen atoms.


[0022] The term "alkylene (spacer)" is typically C_1 - C_{30} alkylene, preferably C_1 - C_{18} alkylene, and embraces the linear as well as the branched representatives and can be, for example, $-CH_2$ - and C_2 - C_{30} alkylene, such as $-(CH_2)_2-$, $-CH(Me)-$, $-(CH_2)_3-$, $-CH_2-CH(Me)-$, $-C(Me)_2-$, $-(CH_2)_4-$, $-(CH_2)_5-$, $-(CH_2)_6-$, $-(CH_2)_7-$, $-(CH_2)_8-$, $-(CH_2)_9-$, $-(CH_2)_{10}-$, $-(CH_2)_{11}-$, $-(CH_2)_{12}-$, $-(CH_2)_{13}-$, $-(CH_2)_{14}-$, $-(CH_2)_{15}-$, $-(CH_2)_{16}-$, $-(CH_2)_{17}-$, $-(CH_2)_{18}-$, $-(CH_2)_{19}-$, $-(CH_2)_{20}-$, $-(CH_2)_{21}-$, $-(CH_2)_{22}-$, $-(CH_2)_{23}-$, $-(CH_2)_{24}-$, $-(CH_2)_{25}-$, $-(CH_2)_{26}-$, $-(CH_2)_{27}-$, $-(CH_2)_{28}-$, $-(CH_2)_{29}-$, $-(CH_2)_{30}-$, preferably $-CH_2-$, $-(CH_2)_2-$, $-(CH_2)_3-$, $-(CH_2)_4-$, $-(CH_2)_5-$, $-(CH_2)_6-$, $-(CH_2)_7-$, $-(CH_2)_8-$, $-(CH_2)_9-$, $-(CH_2)_{10}-$, $-(CH_2)_{11}-$, $-(CH_2)_{12}-$, $-(CH_2)_{13}-$, $-(CH_2)_{14}-$, $-(CH_2)_{15}-$, $-(CH_2)_{16}-$, $-(CH_2)_{17}-$, $-(CH_2)_{18}-$, and also $-CH(C_2-C_{30}alkylene)-$. The "alkylene spacer" can optionally comprise one or more, in particular one or two groups selected from $-O-$, $-S-$, $-NR_{14}-$, $-CO-$, $-CONH-$, $-CONR_{15}-$, or $-COO-$ as linking group. C_1 - C_{30} alkylene can, for example, be interrupted several times by $-O-$, $-S-$, $-NH-$ or $-C(O)NH-$, such as $-(CH_2)_2-O-(CH_2)_2-$, $-(CH_2)_2-O-(CH_2)_2-$, $-(CH_2)_2-S-(CH_2)_2-$, $-CH_2-CH-CH_2-O-(CH_2)_p-CH_3$, wherein p is an integer from 1 to 10; or $-CHX_{13}CH_2-(X_{14})_n-OH$, wherein X_{13} is C_1 - C_8 alkyl, X_{14} is an alkylene oxide monomer, preferably ethylene oxide or propylene oxide, or alkylene amino monomer, preferably amino ethylene or amino propylene, and n is an integer from 1 to 10, preferably 1 to 5; or $-(CH_2)_2-NH-(CH_2)_2-$ or $-(CH_2)_2-C(O)NH-(CH_2)_2-$.

[0023] "Arylene (spacer)" is an unsubstituted or substituted carbocyclic or heterocyclic arylene group, preferably containing 6 to 14 carbon atoms, typically phenylene, naphthylene, anthracenylene, anthraquinonylene, pyridinylene, quinolinyne, preferably a group




wherein X_{11} is a single bond in ortho-, meta- or para-position, or -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO- in ortho-, meta- or para-position; para-phenylene and paraphenylenoxy are preferred.

[0024] "Aralkylene (spacer)" is an unsubstituted or substituted carbocyclic or heterocyclic aralkylene group, preferably containing 6 to 14 carbon atoms, preferably a group wherein X_1 is a single bond in ortho-, meta- or para-position, or -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO- in ortho-, meta- or para-position, and X_{12} is alkylene, or a group




wherein X_{12} is alkylene in ortho-, meta- or para-position and X_{11} is a single bond, -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO-.

[0025] "Cycloalkylene (spacer)" is an unsubstituted or substituted carbocyclic or heterocyclic cycloalkylene group, preferably containing 6 to 14 carbon atoms, typically cyclohexylene, preferably a group



wherein X_{11} is a single bond in 2-, 3- or 4-position, or -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO- in 2-, 3- or 4-position; 4-cyclohexylene and 4-cyclohexylenoxy are preferred.

[0026] Another aspect of the present invention is directed to new diketopyrrolopyrroles. These are at the one hand diketopyrrolopyrroles of the general formula



(IV),

wherein

R_1 and R_2 are independently of each other a group of the formula



wherein

X_2 is an alkylene, arylene, aralkylene or cycloalkylene spacer containing optionally one or more groups -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO- as linking bridge,

X_3 is OH, NH₂, -C(R₁₁)=CH₂, -OC(O)-C(R₁₂)=CH₂, -C(O)-C(R₁₂)=CH₂, or -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂; wherein

R_{11} is hydrogen, or methyl,

R_{12} is hydrogen, or methyl,


R_{14} and R_{15} are independently of each other hydrogen, C_1 - C_8 alkyl, or C_6 - C_{12} aryl, and X_4 and X_5 are as defined above, R_3 and R_4 independently of one another are $-NR_{16}R_{17}$, wherein R_{16} and R_{17} are C_1 - C_{18} alkyl, preferably diketopyrrolopyrroles, wherein R_1 and R_2 are independently of each other a radical of the formula




wherein

X_2 is C_1 - C_{18} alkylene and

X_3 is $-NH_2$, $-OH$, $-CH=CH_2$, $-C(CH_3)=CH_2$, $-OCO-CH=CH_2$ or $-OCO-C(CH_3)=CH_2$, $-CO-CH=CH_2$ or $-CO-C(CH_3)=CH_2$; and on the other hand diketopyrrolopyrroles of the general formula



in which R_{21} and R_{22} are independently of one another C_1 - C_{18} alkyl R_{23} and R_{24} independently of one another are a group of formula



wherein


X_1 is $-S-$, $-SO_2NH-$ or $-NH-$,

X_2 is a C_1 - C_{18} alkylene group, and

X_3 is $-OH$, $-NH_2$, $-CH=CH_2$, $-C(CH_3)=CH_2$, $-OCO-CH=CH_2$, $-OCO-C(CH_3)=CH_2$, $-CO-CH=CH_2$, or $-CO-C(CH_3)=CH_2$.

[0027] Preferably R_1 is R_2 , R_3 is R_4 in formula IV and R_{21} is R_{22} , R_{23} is R_{24} in formula V. In addition, the sum of the number of carbon atoms of R_1 and R_3 (as well as R_2 and R_4) and R_{21} and R_{23} (as well as R_{22} and R_{24}) is preferably 12 ± 6 , most preferred 12 ± 4 .

[0028] Particularly preferred are diketopyrrolopyrroles of the Formula




wherein R_3 and R_4 are preferably identical and are C_1 - C_{16} alkyl, C_1 - C_{16} alkoxy, $-NR_{16}R_{17}$, $-CONHR_{18}$, or $-SO_2NH-R_{20}$, wherein R_{16} and R_{17} are independently of each other both C_1 - C_{16} alkyl or one of the groups R_{16} and R_{17} is hydrogen and the other is C_1 - C_{16} alkyl, R_{18} , R_{19} and R_{20} are C_1 - C_{16} alkyl, and R_{18} is C_1 - C_{15} alkyl;

X_2 is a group $-(X_{22})_{n2}-X_{23}-$, wherein $n2$ is 0 or 1, X_{22} is $-\text{CONH}-$, $-\text{CONR}_{15}-$, or $-\text{COO}-$, X_{23} is an alkylene having 2 to 16 carbon atoms, preferably $-(\text{CH}_2)_{n3}-$, wherein $n3$ is 2 to 16, and X_3 is $-\text{OH}$, $-\text{NH}_2$, $-\text{C}(\text{R}_{11})=\text{CH}_2$, $-\text{C}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$, $-\text{OC}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$ or $-\text{OC}(\text{O})-\text{N}-\text{X}_4-\text{N}-\text{C}(\text{O})-\text{O}-\text{X}_5-\text{O}-\text{C}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$; wherein


R_{11} is hydrogen, or methyl,

R_{12} is hydrogen, or methyl,

and X_4 and X_5 are as defined above; preferably X_4 is p-phenylene, 4,4'-diphenylene, 1,4-cyclohexylene, or $-(\text{CH}_2)_{n4}-$, wherein $n4$ is 3 to 6, and diketopyrrolopyrroles of the formula




in which R_{21} and R_{22} are independently of one another hydrogen, $\text{C}_1\text{-C}_{16}$ alkyl, $\text{C}_1\text{-C}_{16}$ alkyl which is interrupted one or more times by O or S, $\text{C}_7\text{-C}_{12}$ aralkyl or a group of the formula



in which R_5 is $\text{C}_1\text{-C}_{18}$ alkyl,

R_{23} and R_{24} independently of one another are a group of formula



wherein

X_1 is $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, or $-\text{SO}_2-\text{NH}-$,

X_2 is an alkylene having 2 to 16 carbon atoms, preferably $-(\text{CH}_2)_{n3}-$, wherein $n3$ is 2 to 16,


X_3 is $-\text{OH}$, $-\text{NH}_2$, $-\text{C}(\text{R}_{11})=\text{CH}_2$, $-\text{C}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$, $-\text{OC}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$ or $\text{C}(\text{O})-\text{N}-\text{X}_4-\text{N}-\text{C}(\text{O})-\text{O}-\text{X}_5-\text{O}-\text{C}(\text{O})-\text{C}(\text{R}_{12})=\text{CH}_2$; wherein

R_{11} is hydrogen, or methyl,


R_{12} is hydrogen, or methyl,

and X_4 and X_5 are as defined above; preferably X_4 is p-phenylene, 4,4'-diphenylene, 1,4-cyclohexylene, or $-(\text{CH}_2)_{n4}-$, wherein $n4$ is 3 to 6.


[0029] Particularly preferred are the following compounds:




$n = 4 \text{ to } 6 \text{ or } 8 \text{ to } 16$, in particular 12;



$n = 8 \text{ to } 16$, in particular 12;



$n = 8 \text{ to } 16$, in particular 12; or



$n = 8 \text{ to } 16$, in particular 12.

[0030] The diketopyrrolopyrroles are known or can be prepared according to known procedures, see for example: EP-A-61426, EP-A-94911, WO99/54332, WO00/14126, EP-A-787 730 and EP-A-787731.

It has been observed, that some of the diketopyrrolopyrroles of the general formula I, especially IV and V, show fluorescence (photoluminescence). Said diketopyrrolopyrroles can be used in certain full-color OLED displays with one or few luminescent materials as color changing materials (instead of color filters). Color changing materials work by absorbing light of shorter wavelength (e. g., blue light) and then emitting light of longer wavelength by fluorescence (photoluminescence) (e. g., red or green light). The use of color changing materials is known in the art (see e. g., US-B-5,126,214 (Idemitsu Kosan Co., Ltd.) and US-B- 5,294,870 (Eastman Kodak Co.)).

Besides the use of the diketopyrrolopyrroles for preparing color filters the diketopyrrolopyrroles (DPPs) IV or V can also be used for the preparation of polymers. The polymerisation of the inventive DPPs IV or V is usually carried out in a manner known per se, if desired in the presence of a suitable, preferably customary, comonomer carrying e.g. at least one carbon-carbon double bond or of a polymer carrying polyreactive groups. In a preferred embodiment of this invention, coloured (co-)polymers can be prepared by polyreacting a mixture consisting of novel DPP monomers and other customary and suitable copolymerisable monomers in liquid phase such as in a melt, solution, suspension and emulsion. Suitable copolymerisable monomers to be mentioned are, for example, the group of the acrylates, methacrylates and other customary vinylic monomers such as styrene and its customary monomer derivatives or 2-N-vinylpyrrolidone. Particularly preferred acrylates are monofunctional acrylates such as butanediolmonoacrylate, 2-hydroxyethylacrylate, butylacrylate, 2-ethylhexylacrylate, phenoxyethylacrylate, tetrahydrofurfurylacrylate, polypropylene glycol monoacrylate, bifunctional acrylates such as 1,6-hexanedioldiacrylate, tripropylene glycol diacrylate, polyethylene glycol(200) diacrylate and polyethylene glycol(400) diacrylate, ethoxylated and propoxylated neopentyl glycol diacrylate, polyfunctional acrylates such as trimethylolpropanetriacrylate, pentaerythritoltriacylate, ethoxylated or propoxylated trimethylolpropane-triacrylate, propoxylated glycerol triacrylate, tris(2-hydroxyethyl) isocyanurate triacrylate as well as their mixtures with

one another. Such compounds are sufficiently known from, inter alia, "Strahlendhardtung", Curt R. Vincentzu Verlag, Hannover, p. 83 to 92 (1996).

These novel DPP polymers are usually prepared by commonly known methods, e.g. either by a polyreaction, i.e. by polymerisation (thermal or photochemical), polycondensation or polyaddition, or by a polymer-analogous reaction, i.e. by reacting the novel DPP compounds containing suitable reactive groups with polymers already obtained which in turn contain reactive groups (grafting). Photochemical polymerisation is preferred, especially if Q is an acrylic or methacrylic radical.

[0031] It is, thus, possible to prepare e.g. vinyl polymers, allyl polymers, vinyl ester polymers, vinylamide polymers, vinyl acetate polymers or vinyl ketone polymers from DPP monomers; the reactive groups of which contain C=C bonds; polyaldehydes, polyisocyanates, polyepoxides, polyethers, polyacetones or polylactams from monofunctional DPP monomers, the reactive groups of which contain hetero atoms; and from bifunctional DPP monomers, the reactive groups of which contain hetero atoms, via polycondensation, polyesters, polyamides, polyimides or polycarbonates and, via polyaddition, polyepoxides, polyurethanes or polyimides, the polymerisation being, for example, a radical, cationic or anionic polymerisation, coordination polymerisation or group transfer polymerisation.

[0032] DPP monomers which contain amino and hydroxy groups can also be reacted with an polyisocyanate to prepare a monomeric or polymeric isocyanate, which can be reacted with one or more hydroxy-functional acrylates or (meth)acrylates to obtain DPP compounds having groups $-OC(O)-N-X_4-N-C(O)-O-X_5-O-C(O)-C(R_{12})=CH_2$.

[0033] Examples of polyisocyanates are aliphatic diisocyanates, such as trimethylene diisocyanate, tetramethylene diisocyanate, hexamethylene diisocyanate, pentamethylene diisocyanate, 1,2-dipropylene diisocyanate, 2,4,4'- or 2,2,4'-trimethylhexamethylene diisocyanate, and aromatic polyisocyanates, such as m-phenylene diisocyanate, p-phenylene diisocyanate, 4,4'-diphenyl diisocyanate, 1,5-naphthalene diisocyanate, 4,4'-diphenylmethane diisocyanate, 2,4- or 2,6-tolylene diisocyanate, 4,4'-toluidine diisocyanate and 4,4'-diphenyl ether diisocyanate, and alicyclic diisocyanates, such as 1,3- or 1,4-cyclohexane diisocyanate, isophorone diisocyanate and 4,4'-methylenebis(cyclohexyl isocyanate).

[0034] Examples of hydroxy-functional (meth)acrylates are 2-hydroxyethyl(meth)acrylate, 2-hydroxypropyl(meth)acrylate, 2- or 4-hydroxybutyl(meth)acrylate, pentandiol mono(meth)acrylate, trimethylolpropane di(meth)acrylate, and pentaerythritol tri(meth)acrylate.

[0035] Typical examples of the preparation of DPP polymers, starting from the novel DPP monomers IV and V, are (a) the polymerisation for the preparation of DPP polyacrylates or polymethacrylates by radical thermal polymerisation of DPP acrylates or DPP methacrylates, i.e. DPPs of formula IV or V containing an acrylic or methacrylic group, or the radical photopolymerisation of DPP acrylates or DPP methacrylates, (b) the polycondensation for the preparation of DPP-containing polyesters from DPPs of formula IV or V containing a hydroxy group, or the preparation of DPP polycarbonates from DPP diols and phosgene, (c) the polyaddition for the preparation of DPP polyurethanes from DPP diols and diisocyanates, and also (d) the polymer-analogous reaction, e.g. the reaction of a DPP alcohol with a polymer prepared from styrene and maleic acid anhydride, which thus contains anhydride groups, to a polymer containing DPP mono- or diester groups.

[0036] Where required, the novel DPP polymers also contain additives, such as light stabilisers, antioxidants and UV absorbers, which may be added during or after the actual polymerisation, e.g. also during the processing of the polymers (extrusion). These additives can themselves also contain polyreactive groups and can in this case be copolymerised together with the DPP monomers IV or V.

[0037] The actual preparation of polymers is known from the state of the art (described, inter alia, in Houben-Weyl "Methoden der Organischen Chemie", "Makromolekulare Stoffe", Vol. E20, parts 1-3 (1986, 1987)).

[0038] If the DPPs of formula IV or V contain $-CH=CH_2$, acrylate or methacrylate groups, the polymerisation can be carried out e.g. photochemically, one of the customary photoinitiators (see e.g. "Chemistry & Technology of UV & EB Formulations for Coatings, Inks and Paints, Vol. 3: Photoinitiators for Free Radical and Cationic Polymerization" 1991, p. 1115-325) usually being added to the reaction mixture in an amount in the range from typically 0.5 to 5 % by weight, based on the sum of all monomers used.

[0039] Another preferred embodiment of this invention thus relates to polymers based on the diketopyrrolopyrroles IV or V, which are obtainable by polyreacting a mixture consisting of

(A) from 0.5 to 20, preferably from 1 to 10 % by weight, based on the sum of the components (A) and (B), of a diketopyrrolopyrrole IV or V, and

(B) from 99.5 to 80, preferably from 99 to 90 % by weight, based on the sum of the components (A) and (B), of a monomer which is copolymerisable with the diketopyrrolopyrroles IV and V,

the sums of (A) and (B) making up 100 % by weight.

[0040] The DPP polymers prepared according to this invention are particularly suitable for colouring high molecular weight organic materials, such as biopolymers, plastic materials, including fibres, glasses, ceramic products, for formulations of decorative cosmetics, for the preparation of inks, printing inks, paint systems, in particular automotive lacquers

and photosensitive coatings, photo- and electroconducting polymers, fluorescent brighteners, photocell aggregates, coloured photoresists and dispersion paints, and the diketopyrrolopyrroles of this invention can also be used in the biomedical field, for example for the preparation of diagnostic agents, and in the fields of impact and non impact printing and photo/repro in general.

[0041] Illustrative examples of suitable high molecular weight organic materials which can be coloured with the novel DPP polymers, are vinyl polymers, such as polystyrene, poly- α -methylstyrene, poly-p-methylstyrene, poly-p-hydroxystyrene, poly-p-hydroxyphenylstyrene, polymethyl methacrylate and polyacrylamide as well as the corresponding methacrylic compounds, polymethylmaleate, polyacrylonitrile, polymethacrylonitrile, polyvinyl chloride, polyvinyl fluoride, polyvinylidene chloride, polyvinylidene fluoride, polyvinyl acetate, polymethyl vinyl ether and polybutyl vinyl ether; polymers derived from maleinimide and/or maleic anhydride, such as copolymers of maleic anhydride with styrene; polyvinyl pyrrolidone; ABS; ASA; polyamides; polyimides; polyamidimides; polysulfones; polyether sulfones; polyphenylene oxides; polyurethane; polyureae; polycarbonates; polyarylenes; polyarylenesulfides; polyepoxides; polyolefins, such as polyethylene and polypropylene; polyalkadienes; biopolymers and their derivatives, such as cellulose, cellulose ethers and cellulose esters, such as ethylcellulose, nitrocellulose, cellulose acetate and cellulose butyrate, starch, chitin, chitosan, gelatin, zein; natural resins; synthetic resins, such as alkyd resins, acrylic resins, phenolic resins, epoxy resins, aminoformaldehyde resins, such as urea/formaldehyde resins and melamine/formaldehyde resins; rubber; casein; silicone and silicone resins; caoutchouc, chlorinated rubber; and also polymers which are used, for example, as binders in paint systems, such as novolaks derived from C₁-C₆aldehydes, such as formaldehyde and acetaldehyde and a binuclear or mononuclear, preferably mononuclear, phenol which, if desired, is substituted by one or two C₁-C₉alkyl groups, one or two halogen atoms or a phenyl ring, such as o-, m- or p-cresol, xylene, p-tert-butylphenol, o-, m- or p-nonylphenol, p-chlorophenol or p-phenylphenol, or from a compound containing more than one phenolic group, typically resorcinol, bis(4-hydroxyphenyl)methane or 2,2-bis(4-hydroxyphenyl)propane; as well as suitable mixtures of the cited materials.

[0042] Particularly preferred high molecular weight organic materials, in particular for the preparation of a paint system, printing ink or ink, are e.g. cellulose ethers and cellulose esters, such as ethylcellulose, nitrocellulose, cellulose acetate and cellulose butyrate, natural resins or synthetic resins (polymerisation or condensation resins), such as aminoplasts, in particular urea/ formaldehyde resins and melamine/formaldehyde resins, alkyd resins, phenolic plastics, polycarbonates, polyolefins, polystyrene, polyvinyl chloride, polyamides, polyurethanes, polyesters, ABS, ASA, polyphenylene oxides, rubber, casein, silicone and silicone resins as well as their possible mixtures with one another.

[0043] It is also possible to use high molecular weight organic materials in dissolved form as film formers, such as boiled linseed oil, nitrocellulose, alkyd resins, phenolic resins, melamine/formaldehyde resins and urea/formaldehyde resins and also acrylic resins.

[0044] The cited high molecular weight organic compounds can be used singly or in mixtures, e.g. as granulates, plastic compounds, melts or in the form of solutions, in particular for the preparation of spinning solutions, paint systems, coating compositions, inks or printing inks.

[0045] The novel DPP polymers are used for mass colouring polyvinyl chloride, polyamides and, in particular, polyolefins such as polyethylene and polypropylene, and for the preparation of paint systems, including powder coatings, inks, printing inks and paints.

Examples of preferred binders for paint systems to be mentioned are alkyd/melamine surface coating resins, acryl/melamine surface coating resins, cellulose acetate/ cellulose butyrate paints and two-component paints based on acrylic resins crosslinkable with polyisocyanate.

[0046] The novel DPP polymers can be added to the material to be coloured in any desired amount, depending on the end use requirements. In the case of high molecular weight organic materials, for example, the novel DPP polymers can be used in amounts in the range from 0.01 to 40, preferably from 0.1 to 20 % by weight, based on the total weight of the high molecular weight organic material.

[0047] The high molecular weight organic materials are normally coloured with the novel DPP polymers such that said polymers, if desired in the form of masterbatches, are admixed to the high molecular weight organic materials using customary suitable appliances, for example extruders, roll mills, mixing or milling apparatus. The material thus treated is then usually brought into the desired final form by methods known per se, such as calendering, moulding, extruding, coating, casting or injection moulding.

[0048] It is possible to dissolve DDPs of formula IV or V containing photocurable groups such as acrylate or methacrylate groups in other photocurable monomers, either with or without solvent, in the latter case e.g. by melting and dissolving, and then to mix them with corresponding photoinitiators and to coat suitable substrates therewith, the coatings being cured, i.e. polymerised, by means of actinic radiation, preferably UV radiation.

[0049] The novel DPP monomers can be polyreacted in an extruder together with other monomers, in particular those customarily used for the preparation of the above-mentioned polymers (reactive extrusion, in analogy to the process described, inter alia, in EP-A 337 951). Copolymers prepared in this manner usually have the same spectrum of use as the blends of novel DPP polymers and high molecular weight organic materials mentioned so far.

[0050] To obtain different shades, the novel DPP polymers can be advantageously added in admixture with fillers, transparent and opaque white, coloured and/or black pigments and conventional luster pigments in the desired amount.

[0051] To prepare paint systems, coating compositions, inks and printing inks, the corresponding high molecular weight organic substances, such as binders, synthetic resin dispersions and the like, and the novel DPP compounds or polymers, are usually dispersed or dissolved, if desired together with customary additives, such as fillers, paint auxiliaries, siccatives, plasticisers and/or additional pigments, in a shared solvent or solvent mixture. This may be effected by dispersing or dissolving the individual components by themselves or also several together and only then bringing all the components together, or by adding all of them in one go. For printing applications, all conventional industrial printing methods may be used, such as screen printing, rotogravure, bronze printing, flexographic printing and offset printing.


[0052] The polymers and copolymers prepared according to this invention and based on the novel DPP monomers IV or V show excellent migration and fastness characteristics and have high transparence and pure hue as compared to the corresponding polymers of the state of the art.

[0053] The following examples further describe some preferred embodiments of the invention, but do not limit the scope of the invention. In the examples, all parts are by weight unless otherwise indicated.


Examples

Example 1:

[0054] 2.4 parts of NaH were added to the stirred mixture of 9.4 parts of compound:




and 0.6 parts of tetraethylammonium bromide in 400 parts N-methylpyrrolidone. The suspension was stirred for two hours and 13.7 parts of 6-chlorohexanol dissolved in 50 parts N-methylpyrrolidone were added dropwise over 30 minutes. The suspension was heated to 80-85°C and stirred at that temperature over 15 hours. The reaction mixture was cooled to room temperature and diluted with 400 parts of water. The homogeneous suspension was stirred during 1 hour and then filtered. The filter cake was washed with water and dried at 60° in vacuum. 10.8 parts of compound:




were obtained.

Example 2:

[0055] 2.4 parts of NaH were added to the stirred mixture of 9.6 parts of compound:



and 0.6 parts of tetraethylammonium bromide in 400 parts N-methylpyrrolidone. The suspension was stirred for two hours and 16 parts of 1-bromododecane dissolved in 50 parts N-methylpyrrolidone were added dropwise over 30 minutes. The suspension was heated to 80-85°C and stirred at that temperature over 15 hours. The reaction mixture was cooled to room temperature and diluted with 400 parts of water. The homogeneous suspension was stirred during 1 hour and then filtered. The filter cake was washed with water and dried at 60° in vacuum. 11.6 parts of compound:




were obtained.

[0056] Using a similar procedure the following products were obtained:

Example 3:


[0057]



(The starting material was obtained according to example 2 of WO99/54332)


Example 4:

[0058]



Example 5:

[0059] 8.7 parts of compound obtained in example 1, 6.3 parts of methacrylic chloride, 0.5 parts pyridine, and 0.2 parts thiodiphenylamine were dissolved in 300 parts dichloromethane and stirred at room temperature during 6 hours. The reaction mixture was filtered and the filter cake was washed with 500 parts of dichloromethane and dried. 8.5 parts of compound:




were obtained.

Example 6 (Preparation of a Color Filter device (LCD):

[0060] 2.78 g of the composition of example 5, 0.12 g of Solspers® 22000 (Avecia), 0.58 g of Solspers® 24000 (Avecia), 13.33 g of methoxy propyl acetate (RER 600™, Olin Corp.) and 4.77 g of a 40% solution of an acrylate polymer in methoxy propyl acetate are added, diluted with 16.80 g of methoxy propyl acetate, then spin-coated on borosilicate glass plates (Corning 1737) at 1000 rpm. The coated glass plates are dried for 2 minutes at 100°C, then for 5 minutes at 200°C on a hot plate to obtain a uniform red film of thickness 0.4-0.5 μm and excellent transparency and pure hue. Substantially, no sizeable crystals are observed under the optical microscope.

[0061] In a manner similar to example 1 the following compounds can be obtained:




Example	n
7	8
8	9
9	10
10	11
11	12
12	13
13	14

EP 1 523 528 B1

(continued)


Example	n
14	15
15	16

In a manner similar to example 2 the following compounds can be obtained:




Example	n
16	8
17	9
18	10
19	11
20	12
21	13
22	14
23	15
24	16

[0062] In a manner similar to example 3 the following compounds can be obtained:



Example	n
25	8
26	9
27	10
28	11
29	12
30	13
31	14
32	15
33	16

[0063] In a manner similar to example 4 the following compounds can be obtained:




Example	n
34	8

(continued)


Example	n
35	9
36	10
37	11
38	12
39	13
40	14
41	15
42	16

Claims


1. A colour filter comprising a transparent substrate and a layer comprising from 1 to 75% by weight, preferably from 5 to 50% by weight, with particular preference from 25 to 40% by weight, based on the overall weight of the layer, of a diketopyrrolopyrrole of the general formula (I) dispersed in a high molecular mass organic material:



wherein A and B independently of one another are a group of the formula




or



wherein

R_3 and R_4 independently of one another are hydrogen, halogen, C_1 - C_{18} alkyl, C_1 - C_{18} alkoxy, $-NR_{16}R_{17}$, $-CONHR_{18}$, $-COOR_{19}$, $-SO_2NH-R_{20}$, C_1 - C_{18} alkoxycarbonyl, C_1 - C_{11} alkylaminocarbonyl, $-CN$, $-NO_2$, trifluoromethyl, C_5 - C_7 cycloalkyl, $-C=N-(C_1-C_{18}alkyl)$,



imidazolyl, pyrazolyl, triazolyl, piperazinyl, pyrrolyl, oxazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, morpholinyl, piperidinyl or pyrrolidinyl,

G is $-CH_2-$, $-CH(CH_3)-$, $-C(CH_3)_2-$, $-CH=N-$, $-N=N-$, $-O-$, $-S-$, $-SO-$, $-SO_2-$, $-CONH-$ or $-NR_9-$,

R_5 and R_6 independently of one another are hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_{18} alkoxy or $-CN$;

R_7 and R_8 independently of one another are hydrogen, halogen or C_1 - C_6 alkyl and R_9 is hydrogen or C_1 - C_6 alkyl,


R_1 and R_2 are independently of each other C_1 - C_{18} alkyl, C_1 - C_{18} alkyl, which is interrupted one or more times by O or S, C_8 - C_{12} aryl, C_7 - C_{12} aralkyl, or a group of the formula $-C(O)OR_{10}$, wherein R_{10} is C_1 - C_{18} alkyl, C_5 - C_{10} cycloalkyl, C_8 - C_{12} aryl, or C_7 - C_{12} aralkyl, or a group of the formula



wherein

X_2 is an alkylene, arylene, aralkylene or cycloalkylene spacer containing optionally one or more groups $-O-$, $-S-$, $-NR_{14}-$, $-CO-$, $-CONH-$, $-CONR_{15}-$, or $-COO-$ as linking bridge,

X_3 is OH, NH_2 , $-C(R_{11})=CH_2$, $-OC(O)-C(R_{12})=CH_2$, $-C(O)-C(R_{12})=CH_2$, C_5 - C_7 cycloalkenyl,



or $-OC(O)-N-X_4-N-C(O)-O-X_5-O-C(O)-C(R_{12})=CH_2$ wherein

R_{11} is hydrogen, or C_1 - C_4 alkyl, or halogen,

R_{12} is hydrogen, C_1 - C_4 alkyl, or halogen,

R_{13} is hydrogen, C_1 - C_4 alkyl, or C_6 - C_{12} aryl,

R_{14} and R_{15} are independently of each other hydrogen, C_1 - C_8 alkyl, or C_6 - C_{12} aryl,

R_{16} , R_{17} , R_{18} and R_{20} are independently of each other hydrogen, C_1 - C_{18} alkyl, C_6 - C_{12} aryl, or C_7 - C_{12} aralkyl,

R_{19} is C_1 - C_{18} alkyl, C_6 - C_{12} aryl, or C_7 - C_{12} aryl, and

X_4 and X_5 are independently of each other an alkylene, arylene, aralkylene or cycloalkylene spacer,

R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 can also be a group of formula




wherein

X_1 is $-O-$, $-S-$, $-NH-$, $-CONH-$, $-COO-$, $-SO_2NH-$, or $-SO_2O-$, and

X_2 and X_3 are as defined above,

with the proviso that at least one, preferably two, of the groups of the formula (II) and/or (III) is present per molecule.

2. A colour filter according to claim 1, wherein the pigment has the general formula



wherein R_1 and R_2 are independently of each other a group of the formula



wherein

X_2 is an alkylene, allylene, aralkylene or cycloalkylene spacer containing optionally a group $-O-$, $-S-$, $-NR_4-$, $-CO-$, $-CONH-$, $-CONR_{15}-$, or $-COO-$ as linking bridge,

X_3 is $-OH$, $-NH_2$, $-C(R_{11})=CH_2$, $-OC(O)-C(R_{12})=CH_2$, $-C(O)-C(R_{12})=CH_2$, or $-OC(O)-N-X_4-N-C(O)-O-X_5-O-C(O)-C(R_{12})=CH_2$; wherein


R_{11} is hydrogen, or methyl,

R_{12} is hydrogen, or methyl,


R_{14} and R_{15} are independently of each other hydrogen, C_1-C_8 alkyl, or C_6-C_{12} aryl, and X_4 and X_5 are as defined in claim 1,

R_3 and R_4 independently of one another are hydrogen, halogen, C_1-C_{18} alkyl, C_1-C_{18} alkoxy, $-NR_{16}R_{17}$, $-CONHR_{18}$, $-COOR_{19}$, $-SO_2NH-R_{20}$, C_1-C_{18} alkoxycarbonyl, C_1-C_{18} alkylaminocarbonyl, $-CN$, $-NO_2$, trifluoromethyl, C_5-C_7 cycloalkyl, wherein R_{16} , R_{17} , R_{18} , R_{19} and R_{20} are as defined in claim 1.

3. A colour filter according to claim 1, wherein the pigment has the general formula




in which R_{21} and R_{22} are independently of one another hydrogen, C_1-C_{18} alkyl, C_1-C_{18} alkyl which is interrupted one or more times by O or S, C_7-C_{12} aralkyl or a group of the formula



in which R_5 is C_1 - C_{18} alkyl,

R_{23} and R_{24} independently of one another are a group of formula



wherein

X_1 is -O-, -S-, -NH-, -CONH-, -COO-, -SO₂-NH-, or -SO₂-O-,

X_2 is an alkylene, ary lene, aralkylene or cycloalkylene spacer containing optionally one or more groups -O-, S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO- as linking bridge,


X_3 is -OH, -NH₂, -C(R₁₁)=CH₂, -OC(O)-C(R₁₂)=CH₂, -C(O)-C(R₁₂)=CH₂, or -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂; wherein

R_{11} is hydrogen, or methyl,

R_{12} is hydrogen, or methyl,

R_{14} and R_{15} are independently of each other hydrogen, C_1 - C_8 alkyl, or C_6 - C_{12} aryl, and X_4 and X_5 are as defined in claim 1.

4. A diketopyrrolopyrrole of the general formula



wherein R_1 and R_2 are independently of each other a group of the formula



wherein

X_2 is an alkylene, arylene, aralkylene or cycloalkylene spacer containing optionally one or more groups -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, or -COO- as linking bridge,

X_3 is OH, NH₂, -C(R₁₁)=CH₂, -OC(O)-C(R₁₂)=CH₂, -C(O)-C(R₁₂)=CH₂, or -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂; wherein

R_{11} is hydrogen, or methyl,

R_{12} is hydrogen, or methyl,

R_{14} and R_{15} are independently of each other hydrogen, C_1 - C_8 alkyl, or C_6 - C_{12} aryl, and X_4 and X_5 are as defined in claim 1,

R_3 and R_4 independently of one another are -NR₁₆R₁₇, wherein R_{16} and R_{17} are C_1 - C_{18} alkyl.

5. A diketopyrrolopyrrole according to claim 4, wherein R_1 and R_2 are independently of each other a radical of the formula




wherein


X_2 is C_1 - C_{18} alkylene and

X_3 is -NH₂, -OH, -CH=CH₂, -C(CH₃)=CH₂, -OCO-CH=CH₂, -OCO-C(CH₃)=CH₂, -CO-CH=CH₂ or -CO-C(CH₃)=CH₂.

6. A diketopyrrolopyrrole of the general formula



15 in which R_{21} and R_{22} are independently of one another C_1 - C_{18} alkyl,
 R_{23} and R_{24} independently of one another are a group of formula




wherein

X_1 is -S-, -SO₂NH- or -NH-,

X_2 is a C_1 - C_{18} alkylene group, and

X_3 is -OH, -NH₂, -CH=CH₂, -C(CH₃)=CH₂, -OCO-CH=CH₂, -OCO-C(CH₃)=CH₂, -CO-CH=CH₂, or -CO-C(CH₃)=CH₂.

25 7. A diketopyrrolopyrrole according to claim 6, which is a compound of formulae



45 wherein n is 8 to 16.

8. A polymer, obtainable by polyreacting a mixture consisting of

50 (A) from 0.5 to 20, preferably from 1 to 10 % by weight, based on the sum of the components (A) and (B), of a
diketopyrrolopyrrole IV or V, and


(B) from 99.5 to 80, preferably from 99 to 90 % by weight, based on the sum of the components (A) and (B), of
a monomer which is copolymerisable with the diketopyrrolopyrroles IV and V,

the sums of (A) and (B) making up 100 % by weight.


55 9. Use of a diketopyrrolopyrrole of the general formula (I) for manufacturing a colour filter according to claim 1.

Patentansprüche


1. Farbfilter mit einem transparenten Substrat und einer Schicht, die 1 bis 75 Gew.-%, vorzugsweise 5 bis 50 Gew.-%, besonders bevorzugt 25 bis 40 Gew.-%, bezogen auf das Gesamtgewicht der Schicht, eines Diketopyrrolopyrrols der allgemeinen Formel (I) in einem hochmolekularen organischen Material dispergiert umfasst:



worin A und B unabhängig voneinander für eine Gruppe der Formel




oder



stehen, worin

R₃ und R₄ unabhängig voneinander für Wasserstoff, Halogen, C₁-C₁₈-Alkyl, C₁-C₁₈-Alkoxy, -NR₁₆R₁₇, -CONHR₁₈, -COOR₁₉, -SO₂NH-R₂₀, C₁-C₁₈-Alkoxycarbonyl, C₁-C₁₈-Alkylaminocarbonyl, -CN, -NO₂, Trifluormethyl, C₅-C₇-Cycloalkyl, -C=N- (C₁-C₁₈-Alkyl),



Imidazolyl, Pyrazolyl, Triazolyl, Piperazinyl, Pyrrolyl, Oxazolyl, Benzoxazolyl, Benzothiazolyl, Benzimidazolyl, Morpholinyl, Piperidinyl oder Pyrrolidinyl stehen,

G für -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH=N-, -N=N-, -O-, -S-, -SO-, -SO₂-, -CONH- oder -NR₉- steht;


R₅ und R₆ unabhängig voneinander für Wasserstoff, Halogen, C₁-C₆-Alkyl, C₁-C₁₈-Alkoxy oder -CN stehen, R₇ und R₈ unabhängig voneinander für Wasserstoff, Halogen oder C₁-C₆-Alkyl stehen und R₉ für Wasserstoff oder C₁-C₆-Alkyl steht,

R_1 und R_2 unabhängig voneinander für C_1 - C_{18} -Alkyl, C_1 - C_{18} -Alkyl, das ein- oder mehrfach durch O oder S unterbrochen ist, C_6 - C_{12} -Aryl, C_7 - C_{12} -Aralkyl oder eine Gruppe der Formel $-C(O)OR_{10}$, worin R_{10} für C_1 - C_{18} -Alkyl, C_5 - C_{10} -Cycloalkyl, C_6 - C_{12} -Aryl oder C_7 - C_{12} -Aralkyl steht, oder eine Gruppe der Formel



stehen, worin

X_2 für einen Alkyl-, Arylen-, Aralkyl- oder Cycloalkyl-Spacer, der gegebenenfalls eine oder mehrere Gruppen $-O-$, $-S-$, $-NR_{14}-$, $-CO-$, $-CONH-$, $-CONR_{15}-$ oder $-COO-$ als Verknüpfungsbrücke enthält, steht, X_3 für OH , NH_2 , $-C(R_{11})=CH_2$, $-OC(O)-C(R_{12})=CH_2$, $-C(O)-C(R_{12})=CH_2$, C_5 - C_7 -Cycloalkenyl,



oder $-OC(O)-N-X_4-N-C(O)-O-X_5-O-C(O)-C(R_{12})=CH_2$ steht; worin

R_{11} für Wasserstoff oder C_1 - C_4 -Alkyl oder Halogen steht,

R_{12} für Wasserstoff, C_1 - C_4 -Alkyl oder Halogen steht, R_{13} für Wasserstoff, C_1 - C_4 -Alkyl oder C_6 - C_{12} -Aryl steht,

R_{14} und R_{15} unabhängig voneinander für Wasserstoff, C_1 - C_8 -Alkyl oder C_6 - C_{12} -Aryl stehen,

R_{16} , R_{17} , R_{18} und R_{20} unabhängig voneinander für Wasserstoff, C_1 - C_{18} -Alkyl, C_6 - C_{12} -Aryl oder C_7 - C_{12} -Aralkyl stehen,

R_{19} für C_1 - C_{18} -Alkyl, C_6 - C_{12} -Aryl oder C_7 - C_{12} -Aralkyl steht und

X_4 und X_5 unabhängig voneinander für einen Alkyl-, Arylen-, Aralkyl- oder Cycloalkyl-Spacer stehen,

R_3 , R_4 , R_5 , R_6 , R_7 und R_8 auch für eine Gruppe der Formel




stehen können, worin

X_1 für $-O-$, $-S-$, $-NH-$, $-CONH-$, $-COO-$, $-SO_2-NH-$ oder $-SO_2-O-$ steht und

X_2 und X_3 die oben angegebene Bedeutung besitzen, mit der Maßgabe, dass pro Molekül mindestens eine, vorzugsweise zwei, der Gruppen der Formel (II) und/oder Formel (III) vorliegen.

2. FarbfILTER nach Anspruch 1, wobei das Pigment die allgemeine Formel



aufweist, worin R_1 und R_2 unabhängig voneinander für eine Gruppe der Formel



stehen, worin

X_2 für einen Alkyl-, Arylen-, Aralkyl- oder Cycloalkyl-Spacer, der gegebenenfalls eine Gruppe $-O-$, $-S-$, $-NR_{14}-$, $-CO-$, $-CONH-$, $-CONR_{15}-$ oder $-COO-$ als Verknüpfungsbrücke enthält, steht,

X_3 für $-OH$, $-NH_2$, $-C(R_{11})=CH_2$, $-OC(O)-C(R_{12})=CH_2$, $-C(O)-C(R_{12})=CH_2$ oder $-OC(O)-N-X_4-N-C(O)-O-X_5-O-C(O)-C(R_{12})=CH_2$ steht; worin

R₁₁ für Wasserstoff oder Methyl steht,


R₁₂ für Wasserstoff oder Methyl steht,

R₁₄ und R₁₅ unabhängig voneinander für Wasserstoff, C₁-C₈-Alkyl oder C₆-C₁₂-Aryl stehen und


X₄ und X₅ die in Anspruch 1 angegebene Bedeutung besitzen,

R₃ und R₄ unabhängig voneinander für Wasserstoff, Halogen, C₁-C₁₈-Alkyl, C₁-C₁₈-Alkoxy, -NR₁₆R₁₇, -CONHR₁₈, -COOR₁₉, -SO₂NH-R₂₀, C₁-C₁₈-Alkoxy-carbonyl, C₁-C₁₈-Alkylaminocarbonyl, -CN, -NO₂, Trifluormethyl, C₅-C₇-Cycloalkyl stehen, worin R₁₆, R₁₇, R₁₈, R₁₉ und R₂₀ die in Anspruch 1 angegebene Bedeutung besitzen.

3. Farbfilter nach Anspruch 1, wobei das Pigment die allgemeine Formel




aufweist, worin R₂₁ und R₂₂ unabhängig voneinander für Wasserstoff, C₁-C₁₈-Alkyl, C₁-C₁₈-Alkyl, das ein- oder mehrfach durch O oder S unterbrochen ist, C₇-C₁₂-Aralkyl oder eine Gruppe der Formel



worin R₅ für C₁-C₁₈-Alkyl steht, stehen,

R₂₃ und R₂₄ unabhängig voneinander für eine Gruppe der Formel



stehen, worin

X₁ für -O-, -S-, -NH-, -CONH-, -COO-, -SO₂-NH- oder -SO₂-O- steht,

X₂ für einen Alkylen-, Arylen-, Aralkylen- oder Cycloalkylen-Spacer, der gegebenenfalls eine oder mehrere Gruppen -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅- oder -COO- als Verknüpfungsbrücke enthält, steht,

X₃ für -OH, -NH₂, -C(R₁₁)=CH₂, -OC(O)-C(R₁₂)=CH₂, -C(O)-C(R₁₂)=CH₂ oder -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂ steht; worin


R₁₁ für Wasserstoff oder Methyl steht,

R₁₂ für Wasserstoff oder Methyl steht,

R₁₄ und R₁₅ unabhängig voneinander für Wasserstoff, C₁-C₈-Alkyl oder C₆-C₁₂-Aryl stehen und

X₄ und X₅ die in Anspruch 1 angegebene Bedeutung besitzen.

4. Diketopyrrolopyrrol der allgemeinen Formel



worin R_1 und R_2 unabhängig voneinander eine Gruppe der Formel



stehen, worin

X_2 für einen Alkylen-, Arylen-, Aralkylen- oder Cycloalkylen-Spacer, der gegebenenfalls eine oder mehrere Gruppen -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅- oder -COO- als Verknüpfungsbrücke enthält, steht,

20 X_3 für OH, NH₂, -C(R₁₁)=CH₂, -OC(O)-C(R₁₂)=CH₂, -C(O)-C(R₁₂)=CH₂ oder -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂ steht; worin

R_{11} für Wasserstoff oder Methyl steht,

R_{12} für Wasserstoff oder Methyl steht,

R_{19} und R_{15} unabhängig voneinander für Wasserstoff, C₁-C₈-Alkyl oder C₆-C₁₂-Aryl stehen und

25 X_4 und X_5 die in Anspruch 1 angegebene Bedeutung besitzen,

R_3 und R_4 unabhängig voneinander für -NR₁₆R₁₇ stehen, worin R_{16} und R_{17} für C₁-C₁₈-Alkyl stehen.

5. Diketopyrrolopyrrol nach Anspruch 4, worin R_1 und R_2 unabhängig voneinander für einen Rest der Formel




stehen, worin

X_2 für C₁-C₁₈-Alkylen steht und


35 X_3 für -NH₂, -OH, -CH=CH₂, -C(CH₃)=CH₂, -OCO-CH=CH₂, -OCO-C(CH₃)=CH₂, -CO-CH=CH₂ oder -CO-C(CH₃)=CH₂ steht.

6. Diketopyrrolopyrrol der allgemeinen Formel



worin R_{21} und R_{22} unabhängig voneinander für C₁-C₁₈-Alkyl stehen,

R_{23} und R_{24} unabhängig voneinander für eine Gruppe der Formel




stehen, worin

X_1 für -S-, -SO₂-NH- oder -NH- steht,

55 X_2 für eine C₁-C₁₈-Alkylengruppe steht und

X₃ für -OH, -NH₂, -CH=CH₂, -C(CH₃)=CH₂, -OCO-CH=CH₂, -OCO-C(CH₃)=CH₂, -CO-CH=CH₂, oder -CO-C(CH₃)=CH₂ steht.

7. Diketopyrrolopyrrol nach Anspruch 6, bei dem es sich um eine Verbindung der Formeln



worin n für 8 bis 16 steht, handelt.


8. Polymer, erhältlich durch Polyreaktion einer Mischung, die aus

(A) 0,5 bis 20 Gew.-%, vorzugsweise 1 bis 10 Gew.-%, bezogen auf die Summe der Komponenten (A) und (B), eines Diketopyrrolopyrrols IV oder V und
(B) 99,5 bis 80 Gew.-%, vorzugsweise 99 bis 90 Gew.-%, bezogen auf die Summe der Komponenten (A) und (B), eines Monomers, das mit den Diketopyrrolopyrrolen IV und V copolymerisierbar ist, besteht, wobei sich die Summen von (A) und (B) auf 100 Gew.-% belaufen.



9. Verwendung eines Diketopyrrolopyrrols der allgemeinen Formel (I) zur Herstellung eines Farbfilters gemäß Anspruch 1.

Revendications


1. Filtre de couleur comprenant un substrat transparent et une couche comprenant de 1 à 75% en poids, préférablement de 5 à 50% en poids, particulièrement préférablement de 25 à 40% en poids, sur la base du poids global de la couche, de dicétopyrrolopyrrole de formule générale (I) dispersé dans un matériau organique de masse moléculaire élevée :



dans laquelle A et B sont, indépendamment l'un de l'autre, un groupement de formule




ou



dans lesquelles

R_3 et R_4 sont, indépendamment l'un de l'autre, hydrogène, halogène, C_1 - C_{18} alkyle, C_1 - C_{18} alcoxy, $-NR_{16}R_{17}$, $-CONHR_{18}$, $-COOR_{19}$, $-SO_2NH-R_{20}$, C_1 - C_{18} alcoxycarbonyl, C_1 - C_{18} alkylaminocarbonyl, $-CN$, $-NO_2$, trifluorométhyle, C_5 - C_7 cycloalkyle, $-C=N-$ (C_1 - C_{18} alkyl),



imidazole, pyrazole, triazole, pipérazine, pyrrole, oxazole, benzoxazole, benzothiazole, benzimidazole, morpholine, pipéridine ou pyrrolidine,

G est $-CH_2-$, $-CH(CH_3)-$, $-C(CH_3)_2-$, $-CH=N-$, $-N=N-$, $-O-$, $-S-$, $-SO-$, $-SO_2-$, $-CONH-$ ou $-NR_9-$,

R_5 et R_6 sont, indépendamment l'un de l'autre, hydrogène, halogène, C_1 - C_6 alkyle, C_1 - C_{18} alcoxy ou $-CN$,

R_7 et R_8 sont, indépendamment l'un de l'autre, hydrogène, halogène ou C_1 - C_6 alkyle et R_9 est hydrogène ou C_1 - C_6 alkyle,


R_1 et R_2 sont, indépendamment l'un de l'autre, C_1 - C_{18} alkyle, C_1 - C_{18} alkyle qui est interrompu une ou plusieurs fois par O ou S , C_6 - C_{12} aryle, C_7 - C_{12} aralkyle, ou un groupement de formule $-C(O)OR_{10}$, où R_{10} est C_1 - C_{18} alkyle, C_5 - C_{10} cycloalkyle, C_6 - C_{12} aryle, ou C_7 - C_{12} aralkyle, ou


un groupement de formule



dans laquelle X_2 est un espaceur alkylène, arylène, aralkylène ou cycloalkylène contenant éventuellement un ou plusieurs groupements $-O-$, $-S-$, $-NR_{14}-$, $-CO-$, $-CONH-$, $-CONR_{15}-$, ou $-COO-$ comme pont de liaison,

X_3 est OH , NH_2 , $-C(R_{11})=CH_2$, $-OC(O)-C(R_{12})=CH_2$, $-C(O)-C(R_{12})=CH_2$, C_5 - C_7 cycloalcénylo,





ou $-OC(O)-N-X_4-N-C(O)-O-X_5-O-C(O)-C(R_{12})=CH_2$; où

R_{11} est hydrogène, ou C_1-C_4 alkyle, ou halogène,

R_{12} est hydrogène, C_1-C_4 alkyle, ou halogène,

R_{13} est hydrogène, C_1-C_4 alkyle, ou C_6-C_{12} aryle,

R_{14} et R_{15} sont, indépendamment l'un de l'autre, hydrogène, C_1-C_8 alkyle, ou C_6-C_{12} aryle,

R_{16} , R_{17} , R_{18} et R_{20} sont, indépendamment l'un de l'autre, hydrogène, C_1-C_{18} alkyle, C_6-C_{12} aryle, ou C_7-C_{12} aralkyle,

R_{19} est C_1-C_{18} alkyle, C_6-C_{12} aryle, ou C_7-C_{12} aryle, et

X_4 et X_5 sont, indépendamment l'un de l'autre, un espaceur alkylène, arylène, aralkylène ou cycloalkylène,

R_3 , R_4 , R_5 , R_6 , R_7 , et R_8 peuvent aussi être un groupement de formule




dans laquelle X_1 est $-O-$, $-S-$, $-NH-$, $-CONH-$, $-COO-$, $-SO_2-NH-$, ou $-SO_2-O-$, et

X_2 et X_3 sont tels que définis ci-dessus,

à condition qu'au moins un, préférablement deux, parmi les groupements de formule (II) et/ou (III) soient présents par molécule.

2. Filtre de couleur selon la revendication 1, dans lequel le pigment répond à la formule générale



dans laquelle

R_1 et R_2 sont, indépendamment l'un de l'autre, un groupement de formule



dans laquelle

X_2 est un espaceur alkylène, arylène, aralkylène ou cycloalkylène contenant éventuellement un groupement $-O-$, $-S-$, $-NR_{14}-$, $-CO-$, $-CONH-$, $-CONR_{15}-$, ou $-COO-$ comme pont de liaison,

X_3 est $-OH$, $-NH_2$, $-C(R_{11})=CH_2$, $-OC(O)-C(R_{12})=CH_2$, $-C(O)-C(R_{22})=CH_2$, ou $-OC(O)-N-X_4-N-C(O)-O-X_5-O-C(O)-C(R_{12})=CH_2$; où


R_{11} est hydrogène, ou méthyle,

R_{12} est hydrogène, ou méthyle,

R_{14} et R_{15} sont, indépendamment l'un de l'autre, hydrogène, C_1-C_8 alkyle, ou C_6-C_{12} aryle, et


X_4 et X_5 sont tels que définis selon la revendication 1, R_3 et R_4 sont, indépendamment l'un de l'autre, hydrogène, halogène, C_1-C_{18} alkyle, C_1-C_{18} alcoxy, $-NR_{16}R_{17}$, $-CONHR_{18}$, $-COOR_{19}$, $-SO_2NH-R_{20}$, C_1-C_{18} alcoxycarbonyl, C_1-C_{18} alkylaminocarbonyl, $-CN$, $-NO_2$, trifluorométhyle, C_5-C_7 cycloalkyle, où R_{16} , R_{17} , R_{18} , R_{19} et R_{20} sont tels que définis selon la revendication 1.

3. Filtre de couleur selon la revendication 1, dans lequel le pigment répond à la formule générale




dans laquelle

15 R_{21} et R_{22} sont, indépendamment l'un de l'autre, hydrogène, C_1 - C_{18} alkyle, C_1 - C_{18} alkyle qui est interrompu une ou plusieurs fois par O ou S, C_7 - C_{12} aralkyle ou un groupement de formule



dans laquelle R_5 est C_1 - C_{18} alkyle,

R_{23} et R_{24} sont, indépendamment l'un de l'autre, un groupement de formule



dans laquelle

X_1 est -O-, -S-, -NH-, -CONH-, -COO-, -SO₂-NH-, ou -SO₂-O-,

30 X_2 est un espaceur alkylène, arylène, aralkylène ou cycloalkylène contenant éventuellement un ou plusieurs groupements -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, ou -COO- comme pont de liaison,

X_3 est -OH-, -NH₂-, -C(R₁₁)=CH₂-, -OC(O)-C(R₁₂)=CH₂-, -C(O)-C(R₁₂)=CH₂-, ou -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂; où


R_{11} est hydrogène, ou méthyle,

R_{12} est hydrogène, ou méthyle,

35 R_{14} et R_{15} sont, indépendamment l'un de l'autre, hydrogène, C_1 - C_8 alkyle, ou C_6 - C_{12} aryle, et

X_4 et X_5 sont tels que définis selon la revendication 1.

4. Dicétopyrrolopyrrole de formule générale



dans laquelle

R_1 et R_2 sont, indépendamment l'un de l'autre, un groupement de formule



dans laquelle

X_2 est un espaceur alkylène, arylène, aralkylène ou cycloalkylène contenant éventuellement un ou plusieurs grou-


pements -O-, -S-, -NR₁₄-, -CO-, -CONH-, -CONR₁₅-, ou -COO- comme pont de liaison,
X₃ est OH, NH₂, -C(R₁₁)=CH₂, -OC(O)-C(R₁₂)=CH₂, -C(O)-C(R₁₂)=CH₂, ou -OC(O)-N-X₄-N-C(O)-O-X₅-O-C(O)-C(R₁₂)=CH₂ ; où
R₁₁ est hydrogène, ou méthyle,
R₁₂ est hydrogène, ou méthyle,
R₁₄ et R₁₅ sont, indépendamment l'un de l'autre, hydrogène, C₁-C₈alkyle, ou C₆-C₁₂aryle, et X₄ et X₅ sont tels que
définis selon la revendication 1,
R₃ et R₄ sont, indépendamment l'un de l'autre, -NR₁₆R₁₇, où R₁₆ et R₁₇ sont C₁-C₁₈alkyle.

5. Dicétopyrrolopyrrole selon à revendication 4, dans lequel R₁ et R₂ sont, indépendamment l'un de l'autre, un radical de formule




dans laquelle
X₂ est C₁-C₁₈alkylène et
X₃ est -NH₂, -OH, -CH=CH₂, -C(CH₃)=CH₂, -OCO-CH=CH₂, -OCO-C(CH₃)=CH₂, -CO-CH=CH₂ ou -CO-C(CH₃)=CH₂.

6. Dicétopyrrolopyrrole de formule générale




dans laquelle
R₂₁ et R₂₂ sont, indépendamment l'un de l'autre, C₁-C₁₈alkyle,
R₂₃ et R₂₄ sont, indépendamment l'un de l'autre, un groupement de formule



dans laquelle
X₁ est -S-, -SO₂-NH-, ou -NH-,
X₂ est un groupement C₁-C₁₈alkylène, et
X₃ est -OH, -NH₂, -CH=CH₂, -C(CH₃)=CH₂, -OCO-CH=CH₂, -OCO-C(CH₃)=CH₂, -CO-CH=CH₂, ou -CO-C(CH₃)=CH₂.

7. Dicétopyrrolopyrrole selon la revendication 6, qui est un composé de formules



dans lesquelles n va de 8 à 16.

8. Polymère, pouvant être obtenu par la poly-réaction d'un mélange constitué par

(A) de 0,5 à 20, préférablement de 1 à 10% en poids, sur la base de la somme des composants (A) et (B), d'un dicetopyrrolopyrrole IV ou V, et

(B) de 99,5 à 80, préférablement de 99 à 90% en poids, sur la base de la somme des composants (A) et (B), d'un monomère qui est copolymérisable avec les dicetopyrrolopyrroles IV et V,

les sommes de (A) et (B) constituant 100% en poids.

9. Utilisation d'un dicetopyrrolopyrrole de formule générale (I), pour la fabrication d'un filtre de couleur selon la revendication 1.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 811625 A [0002]
- EP 894798 A [0002]
- EP 654711 A [0002] [0011]
- WO 9845756 A [0011]
- WO 9845757 A [0011]
- US 6184375 B1 [0011]
- EP 61426 A [0030]
- EP 94911 A [0030]
- WO 9954332 A [0030] [0057]
- WO 0014126 A [0030]
- EP 787730 A [0030]
- EP 787731 A [0030]
- US 5126214 B [0030]
- US 5294870 B [0030]
- EP 337951 A [0049]

Non-patent literature cited in the description

- Strahlenhdtung. Curt R.Vincentzu Verlag, 1996, 83-92 [0030]
- HOUBEN-WEYL. Methoden der Organischen Chemie. *Makromolekulare Stoffe*, 1986, vol. E20, 1-3 [0037]
- Chemistry & Technology of UV & EB Formulations for Coatings, Inks and Paints, Vol. 3: Photoinitiators for Free Radical and Cationic Polymerization. 1991, vol. 3, 1115-325 [0038]