

[54] Title: METHOD OF INHIBITING INTERLEUKIN -1-RELEASE

[76] Inventor (s): GEORGE KU, of Cincinnati, and NIALL DOHERTY, of West Chester, both of Ohio, U.S.A.

[73] Assignee (s): MERRELL DOW PHARMACEUTICALS INC., of Cincinnati, Ohio, U.S.A.

[22] Filed: March 16, 1988

[21] Application Serial No: 36645

FOREIGN APPLICATION PRIORITY DATA

[31] Number (s) : 026,587; 151,521

[32] Date (s) : 3-17-87; 2-18-88

[33] Country (ies) : U.S.A.

[52] PH Class 514/476; 514/653, 514/721

[51] Int. Class A61K 31/27, A61K 31/075, A61K 31/135

[58] Field of Search 514/476, 514/653, 514/721
A61K 31/27, A61K 31/075, A61K 31/135

[56] Reference (s) Cited and/or Considered: None

[67]

ABSTRACT

Methods useful for inhibiting the release of interleukin-1 and for alleviating interleukin-1 mediated conditions, such as IL-1-mediated inflammation, comprising administration of an effective amount of a pharmaceutically acceptable antioxidant compound such as diulfirn, tetrakis β -(2,6-di-tert-butyl-4-hydroxyphenyl)propionyloxy methyl methane or 2,4-dicobutyl-6-(N,N-dimethylaminomethyl)-phenol.



36644

91 JUN -3 P1:30

26427

METHOD OF INHIBITING INTERLEUKIN-1

RELEASE

ABSTRACT

Methods useful for inhibiting the release of interleukin-1 and for alleviating interleukin-1 mediated conditions, such as IL-1-mediated inflammation, comprising administration of an effective amount of a pharmaceutically acceptable antioxidant compound such as disulfiram, tetrakis β -(2,6-di-tert-butyl-4-hydroxyphenyl)propionyloxy methyl/methane or 2,4-di-isobutyl-6-(N,N-dimethylaminomethyl)-phenol.

26427

BACKGROUND OF THE INVENTION

1. Field of the Invention

Interleukin-1 (IL-1) is the name for a family of molecules which have multiple biological effects. The name interleukin-1 was proposed in 1979; and earlier literature reports refer to it by some other name. Murphy, British Journal of Rheumatology, 1985; 24 (suppl. 1): 6-9, and Oppenheim et al., Immunology Today, vol. 2, 45-55 (1986), IL-1 is secreted by stimulated macrophages, and has several significant biological effects, such as mediation of T-lymphocyte proliferation and pyrogenic and proinflammatory effects.

IL-1 activities are summarized in the two above papers. IL-1 has been described to mediate the acute phase response in inflammation, and to have pyrogenic and proinflammatory effects. IL-1 induces connective tissue changes, and has been demonstrated to induce the release of degradative enzymes from mesenchymal cells that are present at sites of bony erosion in inflammatory disease states, such as rheumatoid arthritis. Billingham, Brit. J. Rheumatology, 1985; 24 (suppl 1): 25-28. Dayer, Brit. J. Rheumatology, 1985; 24 (suppl 1): 15-20. The production of acute phase proteins in the hepatocytes during the acute phase of inflammation is mediated by IL-1.

26427

Whicher, Brit. J. Rheumatology, 1985: 24 (suppl 1):
21-24.

IL-1 is also involved as a mediator in the
inflammatory skin disease, psoriasis, Camp et al.,
5 J. Immunology 1986: 137: 3469-3474, and Ristow, Proc.
Natl. Acad. Sci. USA 1987: 84: 1940-1944. It is cy-
totoxic for insulin producing beta cells in the pan-
creas, and is thus a causative factor in the develop-
ment of diabetes mellitus Bendtsen et al., Science 1986:
10 232: 1545-1547 and Marx, Science 1988: 239: 257-258.
IL-1 also appears to be involved in the development
of atherosclerotic lesions or atherosclerotic plaque.
Marx. Science 1988: 239: 257-258. In the absence or
suppression of endogenous prostaglandins, IL-1 stimu-
15 lates growth and proliferation of vascular smooth
muscle cells, which could lead to thickening of vas-
cular walls, such as occurs in atherogenesis, Libby
et al., Fed. Proc. Mar. 1, 1987: Vol. 46, No. 3: 975.
Abstract 3837.

20 It would be advantageous to control the re-
lease of IL-1, and to be able to treat IL-1-mediated
effects. It would also be advantageous to control
or treat IL-1 mediated inflammations, without pro-
duction of concomitant side effects known to accom-
25 pany the use of antiinflammatory steroids and non-

26427

steroidal antiinflammatory agents.

SUMMARY OF THE INVENTION

5 It has now been found that certain pharmaceu-
tically-acceptable compounds can be used to inhibit
the release of IL-1, and thus to control or treat IL-
1 mediated conditions. Compounds useful in practice-
ing the method of the invention include disulfiram,
tetrakis [3-(2,6-di-tert-butyl-4-hydroxyphenyl)pro-
pionyloxy methyl]methane and 2,4-di-isobutyl-6-(N,N-
10 dimethylaminomethyl)-phenol. Although the compounds
have diverse structures, the compounds all have anti-
oxidant activity. Tetrakis [3-(2,6-di-tert-butyl-4-
hydroxyphenyl)propionyloxy methyl]methane (also named as
Irganox 1010 or as benzenepropanoic acid: 3,5-bis(1,1-
15 dimethylethyl)-4-hydroxy-, tetraester with 2,2-bis-
(hydroxymethyl)-1,3-propanediol) is commercially used
as an antioxidant. The casual mechanism of any rela-
tionship between antioxidant activity of the compounds
and their ability to inhibit IL-1 release is not
20 known, and the invention is not limited to any parti-
cular theoretical mechanism.

Such compounds can be administered to animals
to inhibit secretion of IL-1; to inhibit or treat IL-1-
mediated effects; and to inhibit or treat IL-1-mediated
25 inflammation.

26427

5 In the method of the invention, one or more compound is administered to an animal, typically to a mammal in need of inhibition of IL-1 secretion, inhibition of IL-1-mediated effects, or inhibition of IL-1-mediated inflammation, in an amount effective to produce such inhibition. The compounds can be administered to inhibit or treat IL-1-mediated effects in conditions such as inflammation, psoriasis, atherosclerosis, and diabetes.

10 The compounds can be administered by conventional routes, oral administration being preferred. The dosage to be employed will vary according to factors such as the species, age, weight and condition of the particular animal being treated, and the particular compound employed. Optimum dosages in
15 particular situations can be determined by conventional dose range finding techniques.

20 In general, dosage levels for the use of the compounds to inhibit IL-1 release can be ascertained by conventional range finding studies. The compounds are preferably administered orally at daily dosages from about one to about 300 milligrams of active ingredient per kilogram of animal body weight. Useful
25 results in inhibition of IL-1 release have been obtained with daily oral dosages of 100 milligrams per

26427

kilogram of animal body weight (mg/kg).

5 Although some of the compounds of the method of the invention, such as disulfiram, are known to produce pharmacologic effects unrelated to IL-1 release, they can be usefully employed in the method of the invention with animals which are not in need of such other pharmacologic action. In certain cases, the other pharmacologic action may be regarded as an undesirable side effect, when the desired result is inhibition of IL-1 release. Thus, with disulfiram, 10 for example, concomitant administration of ethanol should be avoided, to avoid the well known reaction to alcohol.

15 The compounds used in the invention can be formulated in conventional pharmaceutically-acceptable carriers to provide unit dosage forms convenient for administration. In general, known dosage forms and carrier materials can be used.

20 The invention is further illustrated in the following Example.

EXAMPLE

Peritoneal macrophages obtained from CD-1 mice were collected and washed once with RPMI-1640 medium (GIBCO, Grand Island, N.Y.) containing 100

26427

Units/ml penicillin, 100 µg/ml streptomycin and 25 µg/ml fungizone (GIBCO, Grand Island, N.Y.). Cells were suspended at 6×10^6 cells per ml, and one ml aliquots were plated into 6-well flat-bottom plates. After one hour incubation at 37°C. in a humidified air chamber containing 5% CO₂, non-adherent cells were removed and 1 ml. RPMI medium (with or without lipopolysaccharide (LPS)-100 µg/well) was added to each well; LPS stimulates macrophages to release IL-1. Incubation was continued for 6 hours, after which the culture supernatant was collected and filtered through 0.22 micrometer Acrodisc filters (Gelman, Ann Arbor, MI). The fluid was stored at a temperature of -70°C. until assayed activity for IL-1 activity.

IL-1 activity was determined by the C3H/HeJ thymocyte proliferation assay of Mizel et al., J. Immunol. 120: 1497 (1978). In this procedure, thymocytes of C3H/HeJ mice are incubated with the macrophage culture supernatant in the presence of phytohemagglutinin, and pulsed by incubation with ³H-thymidine. The cells are then harvested and radioactivity is determined by liquid scintillation counting. IL-1 activity was expressed as units defined according to Mizel et al., J. Immunol. 120: 1497 (1978).

Compounds were tested in this procedure by oral

26427

administration to CD-1 mice 40, 24 and 16 hours prior to collection of peritoneal macrophages. The dosage used as 100 mg/kg. The compounds and results obtained are set out in the following table.

5

Compound	Percent Reduction of IL-1 Secretion
Disulfiram	79.0%
10 Tetrakis [3-(2,6-tert.-butyl-4-hydroxyphenyl)propionyloxy methyl]-methane	66.8%
2,4-Di-isobutyl-6-(N,N-dimethylamino-methyl)-phenol	92.5%

15

The above results indicate significant inhibition of IL-1 release was obtained with the test compounds.

26427

WHAT IS CLAIMED IS:

1. A method of inhibiting the release of interleukin-1 in animals which comprises administering to an animal in need thereof an amount of disulfiram effective to inhibit the release of interleukin-1.

5 2. Method of Claim 1 wherein the animal is suffering from inflammation and the compound is administered in an amount sufficient to alleviate the inflammation.

10 3. Method of Claim 1 wherein the animal is suffering from psoriasis and the compound is administered in an amount sufficient to alleviate the psoriasis.

4. Method of Claim 1 wherein the animal is suffering from diabetes and the compound is administered in an amount sufficient to alleviate the diabetes.

15 5. Method of Claim 1 wherein the animal is suffering from atherosclerosis and the compound is administered in an amount sufficient to alleviate the atherosclerosis.

GEORGE KU
NIALL DOHERTY
Inventors