



(51) International Patent Classification:

A61K 39/00 (2006.01) C12N 5/07 (2010.01)
A61K 39/35 (2006.01)

(21) International Application Number:

PCT/US2015/048225

(22) International Filing Date:

3 September 2015 (03.09.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/045,131 3 September 2014 (03.09.2014) US

(71) Applicant: MODERNA THERAPEUTICS, INC.
[US/US]; 320 Bent Street, Cambridge, Massachusetts
02141 (US).

(72) Inventors: HOGE, Stephen G.; 66 Summit Avenue,
Brookline, Massachusetts 02446 (US). HUANG, Eric Yi-
Chun; 90 Wareham Street #302, Boston, Massachusetts
02118 (US). O'DEA, Louis Saint Laurence; 566 Main
St., Hingham, Massachusetts 02043 (US).

(74) Agents: WARD, Donna T. et al.; 142A Main Street, Gro-
ton, Massachusetts 01450 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: TOLEROGENIC COMPOSITIONS AND METHODS

FIG. 1A





FIG. 1B



(57) Abstract: The invention relates to tolerogenic compositions and methods for the preparation, manufacture and therapeutic use thereof.

**Published:**

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

TOLEROGENIC COMPOSITIONS AND METHODS

REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Patent Application No. 62/045,131, filed September 3, 2014, entitled Compositions and Methods for Tolerizing Cellular Systems, the contents of which is herein incorporated by reference in its entirety.

REFERENCE TO SEQUENCE LISTING

[0002] The present application is being filed along with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled M147PCTSEQLIST.txt, created on September 3, 2015 which is 931,522 bytes in size. The information in the electronic format of the sequence listing is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0003] The present invention is directed to compositions and methods for tolerizing cellular systems. Specifically, in one aspect, the invention relates to tolerogenic polynucleotides, e.g., tolerogenic modified RNA in combination with one or more antigens (which may be delivered as a tolerogenic polynucleotide) useful in tolerizing cell systems. The tolerogenic polynucleotides of the invention may encode peptides, polypeptides or multiple proteins. The tolerogenic polynucleotides or tolerogenic polypeptides, collectively tolerogenic molecules, of interest may be used in therapeutic, clinical and/or research settings.

BACKGROUND OF THE INVENTION

[0004] Advances in the general understanding of immunomodulation have identified several drivers of tolerance and/or suppressors of adaptive immune responses. T-regulatory cells (Tregs) are critical suppressors of T effector cells and Tregs are now understood to be the critical balancers against autoimmunity and for modulating immune responses to avoid damage to self (Wing & Sagakuchi, Nat Immunol. 2010 Jan;11(1):7-13).

[0005] Dendritic cells (DCs) and other antigen presenting cells (APCs) also play a critical role in presenting antigens to T cells, including Tregs. The context of these antigen presentations, including APC to T cell signaling through both cell contact (e.g.,

TCR/MHC2/co receptor) and non-contact (e.g., cytokines) mechanisms, can either promote tolerance or induce potent adaptive responses and high affinity T effector cell responses. These in turn can potentiate and lock in B-cell antibody responses (Watanabe et al, *Autoimmunity*, 1999, Vol. 31, No. 4, Pages 273-282; Luo, et al., *PNAS*, 2007; vol. 104; no. 8; 2821–2826).

[0006] Several mechanisms for natural modulation of the T reg vs. T effector balance towards a tolerant phenotype for self-and non-self-antigens have also been described, particularly the use of immunosuppressive cytokines, antigen presenting modulators, co-inhibitory factors and T regulatory cell epitopes (Tregitopes).

[0007] For example, several cytokines with potent anti-inflammatory profiles have been identified. These can drive polarity of the adaptive immune response towards tolerance, often through Tregs. In addition to classical examples like TGF-beta (Luo et al, *PNAS*, 2007, Vol. 104, No. 8, Pages 2821-2826), and IL-10 several novel families have been described including IL-12 and IL-37 (see reviews in Banchereau et al, *Nature Immunology*, 2012, Vol. 13, No. 10, Pages 925-931; Cheng et al, *Journal of Biological Chemistry*, 2011, Vol. 286, No. 20, Pages 18013-18025; Wing & Sakaguchi, *Nature Immunology*, 2010, Vol. 11, No. 1, Pages 7-13). In most, if not all, cases these cytokines act locally as non-contact signaling molecules between APCs and T cells or between T regs and T effector cells (T effectors).

[0008] Antigen presenting modifiers have also been identified. For example, SOCS1 mRNA can attenuate antigen presentation (Evel-Kabler et al, *Journal of Clinical Investigation*, 2006, Vol. 116, No. 1, Pages 90-100). Similarly several molecules have been identified that decrease MHC2 antigen presentation.

[0009] Several co-inhibitory molecules also exist which signal through contact as part of the T-cell receptor (TCR) synapse. These include ILT3, GITR, and CTLA4. ILT3 expression on monocytes and DCs has been associated with transplant acceptance in allografts (Chang et al, *Nature Immunology*, 2002, Vol. 3, No. 3, Pages 237-243) and delays onset of T1D in NOD-scid mice (Vlad et al, *Diabetes*, 2008, Vol. 57, Pages 1878-1886). See also Suci-Foca et al, *Journal of Immunology*, 2007, Vol. 178, Pages 7432-7441.

[00010] T regulatory cell epitopes or Tregitopes, first identified bio-informatically as conserved epitopes on Fc regions of circulating IgG (Cousens et al, *J Clin Immunol*, 2012, Vol 33 Suppl 1, Pages S43-S49), have been the focus of a recent study showing some impact of co-administration of antigens and Tregs in models of autoimmune type 1 diabetes.

[00011] The importance of Tregs in tolerance cannot be understated, with several specific cytokines being identified to play a role in the process to date (e.g., IL-10, TGF-beta) see Hoffman et al, *Molecular Therapy*, 2011, Vol. 19, No. 7, Pages 1263-1272, for example. It is crucial to note that context matters in the sense of adaptive immune function and microenvironments. For example, hepatic gene transfer can be used to induce tolerance even to non-native (e.g., human in mouse/non-human primate/canine) proteins (LoDuca et al, *Curr Gene Ther*, 2009, vol. 9(2), Pages 104-114; Finn et al, *Blood*, 2010, Vol. 116, Pages 5842-5848). Similar findings have been observed in human trials of gene therapy where pharmacological immunomodulation, including cyclosporine A, rituximab, and high dose systemic steroids, can modulate or prevent development of neutralizing antibodies (NAbs). Mingozi et al, *Molecular Therapy*, 2012, Vol. 20, No. 7, Pages 1410-1416.

[00012] The gene therapy field has also made significant advances over the last decade in addressing the problem of adaptive immune responses to transgenes, both self (tolerance breaking) and non-self (no native peptide). see High, *Blood*, 2012, Vol 120, Pages 4482-4487.

[00013] The specific administration locale or effective microenvironment also plays a strong role in positive outcomes and is the predominant reason that recombinant approaches are not as successful since expression is more widespread for such tolerogenic compositions. For example, systemically administered nanoparticles, including ionizable and cationic lipid nanoparticles, are usually taken up preferentially in the liver and by APCs as well as myeloid cells (DCs, monocytes) in the blood. It is envisioned that adjusting nanoparticle size and the addition of targeting epitopes can drive the preference towards APCs/DCs and away from hepatocytes. Furthermore, it is also envisioned that cell, tissue or organ specific targeting may enhance or improve tolerance. To this end, engineering microRNA binding sites into the 3'UTR of the

transcript that destabilize the transcript in hepatocytes can also be used to further tune translation of the tolerogenic target towards DCs and the cells of the immune system (e.g., mir122 “knock down” in hepatocytes) and the specific antigen / transgene towards the target cells (e.g. mir142.3p “knock down” in lymphocytes).

[00014] The present invention addresses the present and long-felt need for tolerogenic compounds and/or compositions, including methods of using such compound and compositions, for the selective tuning of the adaptive immune system as interventions which expand Treg cells may offer novel treatment options in a variety of clinical settings. To this end, the present invention provides tolerogenic compositions including polynucleotides which may be used alone or in conjunction with other therapeutic modalities, including antigens, adjuvants and/or other polynucleotides (whether tolerizing or not), to alter or modulate tolerance in cellular systems. In this manner, the profile or signature of an immune system microenvironment may be fine tuned using the embodiments of the invention like a rheostat or regulator to accept or reject the presentation of one or more antigens, adjuvants or therapeutic modalities.

SUMMARY OF THE INVENTION

[00015] Described herein are compositions, methods, processes, kits and devices for the design, preparation, manufacture and/or formulation of polynucleotides (whether protein coding or not) which function to alter the adaptive immune response of a cell or cell systems to one or more antigens, adjuvants or therapeutic modalities or to change the innate immune profile or signature of a cell or cell system in response to contact with such antigen adjuvant or therapeutic modality. The compositions or formulations of the polynucleotides may comprise phosphatidylserine (PS).

[00016] Provided herein are methods of inducing tolerance in a cell system to an antigen comprising contacting a cellular system with a tolerogenic composition comprising phosphatidylserine (PS), the antigen and one or more polynucleotides which may encode a tolerogenic polypeptide of interest.

[00017] Also provided herein are methods of inducing Treg cell activity in a cellular system comprising contacting the cellular system with a tolerogenic composition comprising phosphatidylserine (PS), an antigen and one or more polynucleotides which may encode a tolerogenic polypeptide of interest.

[00018] In one embodiment, the one or more tolerogenic polynucleotides may comprise a chemically modified mRNA and/or may encode an immunomodulatory polypeptide. The immunomodulatory polypeptide may encode an inhibitor of mTOR, IL-2, an anti-IL-2 complex, IL-10, TGF- β , IL-35, galectin-1, IL-23, IL-27, IL-35, IL-37 or antibody such as, but not limited to, an antibody reactive to CD3, CD40, CD40 ligand, CD4, and CTLA-4. The antibody may be reactive with a member

[00019] The chemically modified mRNA may comprise at least one region which is codon optimized.

[00020] The tolerogenic compositions described herein may be formulated in any formulation described herein such as, but not limited to a lipid nanoparticle (LNP). The tolerogenic composition may comprise phosphatidylserine (PS).

[00021] The LNP formulation may be administered by the methods described herein including, but not limited to, systemically. The tolerogenic compositions described herein may include phosphatidylserine (PS).

[00022] The tolerogenic compositions described herein may be co-administered with phosphatidylserine (PS).

[00023] Provided herein is a method of treating an autoimmune disease (e.g., lupus), inflammatory disease (e.g., colitis, Crohn's disease, allergic encephalitis), allograft transplant/graft vs. host disease (GVHD), diabetes or multiple sclerosis, comprising contacting a cell, tissue or organism with a tolerogenic composition comprising an antigen and one or more polynucleotides encoding a tolerogenic polypeptide of interest.

[00024] Provided herein is a method of treating a subject with an environmental hypersensitivity or allergic reaction to a protein or antigen comprising contacting the subject with a tolerogenic composition described herein. The environmental hypersensitivity or allergic reaction may be to a protein, antigen or other trigger such as, but not limited to, plants, grass, tree and other plant pollens, mammal, animal hair or dander, insect, insect parts, excreta or venom, helminthes, non-mammals, birds and reptiles, foods (e.g., legumes, nuts and fish), bacteria, fungi and molds, drugs, metals, wool and latex.

[00025] The subject may be contacted using a micro-dosing regimen that occurs over a period of time such as, but not limited to, at least 1 hour, at least 2 hours, at least 3 hours,

at least 4 hours, at least 5 hours, at least 6 hours, at least 7 hours, at least 8 hours, at least 9 hours, at least 10 hours, at least 11 hours, at least 12 hours, at least 13 hours, at least 14 hours, at least 15 hours, at least 16 hours, at least 17 hours, at least 18 hours, at least 19 hours, at least 20 hours, at least 21 hours, at least 22 hours, at least 23 hours, at least 1 day, at least 36 hours, at least 2 days, at least 3 days, at least 4 days, at least 5 days, at least 6 days, at least 1 week, at least 2 weeks, at least 3 weeks, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months or more than 6 months. The amount of antigen and one or more polynucleotides in the tolerogenic composition may be increased during the micro-dosing regimen.

[00026] The details of various embodiments of the invention are set forth in the description below. Other features, objects, and advantages of the invention will be apparent from the description and the drawings, and from the appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[00027] The foregoing and other objects, features and advantages will be apparent from the following description of particular embodiments of the invention, as illustrated in the accompanying drawings in which like reference characters refer to the same parts throughout the different views. The drawings are not necessarily to scale, emphasis instead being placed upon illustrating the principles of various embodiments of the invention.

[00028] FIG. 1 comprises FIG 1A and FIG 1B showing a schematic of an IVT polynucleotide construct. FIG 1A is a schematic of an IVT polynucleotide construct taught in commonly owned US Patent No. 8,999,380, the contents of which are incorporated herein by reference. FIG 1B is a schematic of an IVT polynucleotide construct.

[00029] FIG. 2 is a schematic of a series of chimeric polynucleotides of the present invention. Such chimeric polynucleotides may function alone or in combination with another molecule as a polynucleotide encoding a tolerogenic polypeptide of interest.

[00030] FIG. 3 is a schematic of a series of chimeric polynucleotides illustrating various patterns of positional modifications and showing regions analogous to those regions of an mRNA polynucleotide. Such chimeric polynucleotides may function alone

or in combination with another molecule as a polynucleotide encoding a tolerogenic polypeptide of interest.

[00031] FIG. 4 is a schematic of a series of chimeric polynucleotides illustrating various patterns of positional modifications based on Formula I. Such chimeric polynucleotides may function alone or in combination with another molecule as a polynucleotide encoding a tolerogenic polypeptide of interest.

[00032] FIG. 5 is a is a schematic of a series of chimeric polynucleotides illustrating various patterns of positional modifications based on Formula I and further illustrating a blocked or structured 3' terminus. Such polynucleotides may function alone or in combination with another molecule as a polynucleotide encoding a tolerogenic polypeptide of interest.

DETAILED DESCRIPTION OF THE INVENTION

[00033] The present invention provides compositions and methods for tolerizing cellular systems, in particular for the modulation of adaptive immunity using tolerogenic molecules (e.g., tolerogenic polynucleotides which may comprise modified RNA and/or mRNA constructs), including promoting tolerance for gene therapy, improving the ability to “repeat dose” one or more polynucleotides as a therapy, and treating autoimmune diseases such as lupus, inflammatory diseases such as colitis, Chron’s disease, allergic encephalitis, and/or allograft transplant/graft vs. host disease (GVHD), diabetes or multiple sclerosis.

[00034] The combination of polynucleotides encoding tolerogenic signaling polypeptides and a delivery technology (e.g., lipid nanoparticles or LNPs) with a propensity for uptake in antigen presenting cells (APCs) provides a platform for driving tolerance to specific antigens or epitopes contained therein that may be co-administered to those APCs/dendritic cells (DCs). Some practical applications of the present invention include: (1) treatment of autoimmune disease, (2) creating or improving tolerance to gene therapy transgenes, (3) creating or improving tolerance to repeat modified mRNA therapy that is immunogenic, and (4) creating or improving tolerance to allografts/transplants of solid organs.

[00035] In one embodiment, a tolerogenic composition comprising at least one polynucleotide encoding a tolerogenic polypeptide may be used as a tolerance inducing

therapeutic in transplant. The composition may comprise or may be co-administered with phosphatidylserine (PS). As a non-limiting example, harvested organs may be contacted with a tolerogenic composition through a single perfusion. As another non-limiting example, the transplant donor may be administered a tolerogenic composition comprising polynucleotides encoding tolerogenic polypeptides of interest such as tolerogenic signals based on the donor HLA haplotype to the transplant recipient.

[00036] In one embodiment, a tolerogenic composition comprising at least one polynucleotide encoding a tolerogenic polypeptide may be used to treat and/or prevent autoimmunity. “Autoimmunity” is the failure of an organism in recognizing its own constituent parts as self, thus leading to an immune response against its own cells and tissues. Autoimmunity is an immune response to foreign antigens (alloantigens) from members of the same species. Diseases that result from the aberrant immune response are referred to as “autoimmune diseases” where an organism loses the ability to ignore “self” and react to “non-self.”

[00037] In one embodiment, co-administration of a tolerogenic composition comprising at least one polynucleotide encoding a tolerogenic polypeptide may be used to induce tolerance and to create recipient Tregs for key donor antigens.

[00038] In another embodiment, a tolerogenic composition comprising at least one polynucleotide encoding a tolerogenic polypeptide may be used in allogenic bone marrow transplantation (BMT) to instruct tolerance for the allografted immune system in order to reduce or eliminate graft versus host disease (GVHD).

[00039] In another embodiment, a tolerogenic composition comprising at least one polynucleotide encoding a tolerogenic polypeptide may be used in a formulation with phosphatidylserine in allogenic bone marrow transplantation (BMT) to further enhance tolerance for the allografted immune system in order to reduce or eliminate GVHD, by inhibiting maturation and antigen presentation of dendritic cells.

[00040] Described herein are compositions (including pharmaceutical compositions) and methods for the design, preparation, manufacture and/or formulation of polynucleotides, specifically IVT polynucleotides, chimeric polynucleotides and/or circular polynucleotides encoding at least one tolerogenic molecule and/or fragment thereof.

[00041] In some embodiments, the polynucleotides are administered in combination with one or more antigens, adjuvants or other molecule (including other tolerogenic polynucleotides) in tolerogenic compositions.

[00042] Also provided are systems, processes, devices and kits for the selection, design and/or utilization of the polynucleotides described herein.

[00043] According to the present invention, the polynucleotides are preferably modified in a manner as to avoid the deficiencies of other molecules of the art. Consequently, provided herein, therefore, are polynucleotides (also referred to as polynucleotides, irrespective of whether they are synthesized via IVT or chimeric means or whether they are linear or circular) which improve one or more of the stability and/or clearance in tissues, receptor uptake and/or kinetics, cellular access, engagement with translational machinery, mRNA half-life, translation efficiency, immune evasion, protein production capacity, secretion efficiency (when applicable), accessibility to circulation, protein half-life and/or modulation of a cell's status, function and/or activity while also functioning to modulate or alter at least one adaptive immune response of a cell or cell system.

[00044] Polynucleotides of the present invention may be administered alone or in combination with other polynucleotides encoding tolerogenic polypeptides of interest (of any type) or with other molecules to alter self or non-self responsiveness of cells or cellular systems. To this end, a first polynucleotide encoding a tolerogenic polypeptide of interest may alter the adaptive immune response to itself, its encoded polypeptide or to another distinct polynucleotide encoding a tolerogenic polypeptide of interest or the polypeptide encoded therein.

[00045] In one embodiment, provided herein are tolerogenic compositions comprising at least one antigen, adjuvant or other molecule and at least one polynucleotide encoding a tolerogenic polypeptide of interest.

[00046] In one embodiment, at least one polynucleotide encoding a tolerogenic polypeptide of interest may be formulated in a tolerogenic composition with at least one antigen, adjuvant or other molecule. The tolerogenic composition may be delivered to a cell, tissue or subject alone or in combination with other polynucleotides, adjuvants, antigens and/or other molecules.

[00047] In one embodiment, at least one polynucleotide encoding a tolerogenic polypeptide of interest may be co-administered in a tolerogenic composition with at least one antigen, adjuvant or other molecule. The polynucleotide and the antigen, adjuvant or other molecule may be formulated in the same tolerogenic composition or in distinct tolerogenic compositions. As a non-limiting example, the polynucleotide encoding a tolerogenic polypeptide of interest may be formulated in a tolerogenic composition which may also comprise at least one antigen, adjuvant or other molecule. As another non-limiting example, the polynucleotide encoding a tolerogenic polypeptide of interest may be formulated in a first tolerogenic composition and the antigen, adjuvant or other molecule may be formulated in a second tolerogenic composition. The first and second composition may be co-administered to a cell, tissue or organism.

[00048] As used herein, “co-administered” means the administration of two or more components. These components for co-administration include, but are not limited to adjuvants, antigens, active ingredients, polynucleotides, amino acids, inactive ingredients and excipients. Co-administration refers to the administration of two or more components simultaneously or with a time lapse between administration such as 1 second, 5 seconds, 10 seconds, 15 seconds, 30 seconds, 45 seconds, 1 minute, 2 minutes, 3 minutes, 4 minutes, 5 minutes, 6 minutes, 7 minutes, 8 minutes, 9 minutes, 10 minutes, 11 minutes, 12 minutes, 13 minutes, 14 minutes, 15 minutes, 16 minutes, 17 minutes, 18 minutes, 19 minutes, 20 minutes, 21 minutes, 22 minutes, 23 minutes, 24 minutes, 25 minutes, 26 minutes, 27 minutes, 28 minutes, 29 minutes, 30 minutes, 31 minutes, 32 minutes, 33 minutes, 34 minutes, 35 minutes, 36 minutes, 37 minutes, 38 minutes, 39 minutes, 40 minutes, 41 minutes, 42 minutes, 43 minutes, 44 minutes, 45 minutes, 46 minutes, 47 minutes, 48 minutes, 49 minutes, 50 minutes, 51 minutes, 52 minutes, 53 minutes, 54 minutes, 55 minutes, 56 minutes, 57 minutes, 58 minutes, 59 minutes, 1 hour, 1.5 hours, 2 hours, 2.5 hours, 3 hours, 4 hours, 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, 10 hours, 11 hours, 12 hours, 13 hours, 14 hours, 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23 hours, 1 day, 1.5 days, 2 days, or more than 3 days.

[00049] It is further appreciated that certain features of the present disclosure, which are, for clarity, described in the context of separate embodiments, can also be provided in

combination in a single embodiment. Conversely, various features of the present disclosure which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

I. Compositions of the Invention

Polynucleotides

[00050] The present invention provides nucleic acid molecules, specifically polynucleotides which, in some embodiments, encode one or more peptides or polypeptides of interest such as, but not limited to, the tolerogenic polypeptides taught herein. The term “nucleic acid,” in its broadest sense, includes any compound and/or substance that comprise a polymer of nucleotides. These polymers are often referred to as polynucleotides. When referring to polynucleotides of the invention, it should be understood that the term “polynucleotides” broadly embraces the polynucleotides encoding tolerogenic polypeptides of interest regardless of their method of synthesis (e.g., IVT or chemically synthesized or combinations thereof); structure, (e.g., linear or circular or combinations thereof); or coding capacity (e.g., protein coding or non-coding).

[00051] The polynucleotides of the present invention function to promote tolerance in cells or cellular systems, whether to self or non-self antigens. As used herein, the term “tolerance” when referring to cells or cellular systems means an antigen-specific nonresponsiveness to a challenge, where the “antigen” comprises a polynucleotide of the present invention. By non-responsiveness” it is meant that administration or contact with a specific antigen produces at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99% or 100% reduction in responsiveness as compared to the cell or cellular response in the absence of the polynucleotide encoding a tolerogenic polypeptide of interest or the tolerogenic polypeptide. Therefore, the tolerogenic molecules of the present invention may be referred to, or considered, antigens.

[00052] Such antigen challenges may be delivered with or without an adjuvant. Adjuvants may be strong adjuvants or weak adjuvants. As used herein, the term “tolerogenic” in reference to a polynucleotide or polypeptide refers to a molecule which functions to induce, promote, or improve tolerance. Such induction, promotion or improvement may be via the modulation of the balance between Tregs and T effector

cells. A polynucleotide of the invention may encode a tolerogenic polypeptide or it may encode a noncoding tolerogenic polynucleotide.

[00053] Exemplary nucleic acids or polynucleotides of the invention include, but are not limited to, ribonucleic acids (RNAs), deoxyribonucleic acids (DNAs), threose nucleic acids (TNAs), glycol nucleic acids (GNAs), peptide nucleic acids (PNAs), locked nucleic acids (LNAs, including LNA having a β - D-ribo configuration, α -LNA having an α -L-ribo configuration (a diastereomer of LNA), 2'-amino-LNA having a 2'-amino functionalization, and 2'-amino- α -LNA having a 2'-amino functionalization), ethylene nucleic acids (ENA), cyclohexenyl nucleic acids (CeNA) or hybrids or combinations thereof.

[00054] In one embodiment, linear polynucleotides of the present invention which are made using only *in vitro* transcription (IVT) enzymatic synthesis methods are referred to as "IVT polynucleotides." Methods of making IVT polynucleotides are known in the art and are described in co-pending International Publication Nos. WO2013151666, WO2013151667, WO2013151668, WO2013151663, WO2013151669, WO2013151670, WO2013151664, WO2013151665, WO2013151736, WO2013151671 and WO2013151672, and in some embodiments, the antigen-specific response that is to be altered arises from a challenge to the cell or cellular system by any one of the polynucleotides or polypeptides taught therein, the contents of each of which are herein incorporated by reference in their entireties.

[00055] In another embodiment, the polynucleotides of the present invention which have portions or regions which differ in size and/or chemical modification pattern, chemical modification position, chemical modification percent or chemical modification population and combinations of the foregoing are known as "chimeric polynucleotides." A "chimera" according to the present invention is an entity having two or more incongruous or heterogeneous parts or regions. As used herein a "part" or "region" of a polynucleotide is defined as any portion of the polynucleotide which is less than the entire length of the polynucleotide.

[00056] In yet another embodiment, the polynucleotides of the present invention that are circular are known as "circular polynucleotides" or "circP." As used herein, "circular polynucleotides" or "circP" means a single stranded circular polynucleotide which acts

substantially like, and has the properties of, an RNA. The term “circular” is also meant to encompass any secondary or tertiary configuration of the circP.

[00057] In some embodiments, the polynucleotide includes from about 30 to about 100,000 nucleotides (e.g., from 30 to 50, from 30 to 100, from 30 to 250, from 30 to 500, from 30 to 1,000, from 30 to 1,500, from 30 to 3,000, from 30 to 5,000, from 30 to 7,000, from 30 to 10,000, from 30 to 25,000, from 30 to 50,000, from 30 to 70,000, from 100 to 250, from 100 to 500, from 100 to 1,000, from 100 to 1,500, from 100 to 3,000, from 100 to 5,000, from 100 to 7,000, from 100 to 10,000, from 100 to 25,000, from 100 to 50,000, from 100 to 70,000, from 100 to 100,000, from 500 to 1,000, from 500 to 1,500, from 500 to 2,000, from 500 to 3,000, from 500 to 5,000, from 500 to 7,000, from 500 to 10,000, from 500 to 25,000, from 500 to 50,000, from 500 to 70,000, from 500 to 100,000, from 1,000 to 1,500, from 1,000 to 2,000, from 1,000 to 3,000, from 1,000 to 5,000, from 1,000 to 7,000, from 1,000 to 10,000, from 1,000 to 25,000, from 1,000 to 50,000, from 1,000 to 70,000, from 1,000 to 100,000, from 1,500 to 3,000, from 1,500 to 5,000, from 1,500 to 7,000, from 1,500 to 10,000, from 1,500 to 25,000, from 1,500 to 50,000, from 1,500 to 70,000, from 1,500 to 100,000, from 2,000 to 3,000, from 2,000 to 5,000, from 2,000 to 7,000, from 2,000 to 10,000, from 2,000 to 25,000, from 2,000 to 50,000, from 2,000 to 70,000, and from 2,000 to 100,000).

[00058] In one embodiment, the polynucleotides of the present invention may encode at least one peptide or polypeptide of interest. In another embodiment, the polynucleotides of the present invention may be non-coding.

[00059] In one embodiment, the length of a region encoding at least one peptide polypeptide of interest of the polynucleotides present invention is greater than about 30 nucleotides in length (e.g., at least or greater than about 35, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,900, 2,000, 2,500, and 3,000, 4,000, 5,000, 6,000, 7,000, 8,000, 9,000, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 or up to and including 100,000 nucleotides). As used herein, such a region may be referred to as a “coding region” or “region encoding.”

[00060] In one embodiment, the polynucleotides of the present invention is or functions as a messenger RNA (mRNA). As used herein, the term “messenger RNA”

(mRNA) refers to any polynucleotide which encodes at least one peptide or polypeptide of interest and which is capable of being translated to produce the encoded peptide polypeptide of interest *in vitro*, *in vivo*, *in situ* or *ex vivo*.

[00061] In one embodiment, the polynucleotides of the present invention may be structurally modified or chemically modified. As used herein, a “structural” modification is one in which two or more linked nucleosides are inserted, deleted, duplicated, inverted or randomized in a polynucleotide without significant chemical modification to the nucleotides themselves. Because chemical bonds will necessarily be broken and reformed to effect a structural modification, structural modifications are of a chemical nature and hence are chemical modifications. However, structural modifications will result in a different sequence of nucleotides. For example, the polynucleotide “ATCG” may be chemically modified to “AT-5meC-G”. The same polynucleotide may be structurally modified from “ATCG” to “ATCCCG”. Here, the dinucleotide “CC” has been inserted, resulting in a structural modification to the polynucleotide.

[00062] In one embodiment, the polynucleotides of the present invention, such as IVT polynucleotides or circular polynucleotides, may have a uniform chemical modification of all or any of the same nucleoside type or a population of modifications produced by mere downward titration of the same starting modification in all or any of the same nucleoside type, or a measured percent of a chemical modification of all any of the same nucleoside type but with random incorporation, such as where all uridines are replaced by a uridine analog, e.g., pseudouridine. In another embodiment, the polynucleotides may have a uniform chemical modification of two, three, or four of the same nucleoside type throughout the entire polynucleotide (such as all uridines and all cytosines, etc. are modified in the same way).

[00063] When the polynucleotides of the present invention are chemically and/or structurally modified the polynucleotides may be referred to as “modified polynucleotides.”

[00064] In one embodiment, the polynucleotides of the present invention may include a sequence encoding a self-cleaving peptide. The self-cleaving peptide may be, but is not limited to, a 2A peptide. As a non-limiting example, the 2A peptide may have the protein sequence: GSGATNFSLLKQAGDVEENPGP (SEQ ID NO: 1), fragments or variants

thereof. In one embodiment, the 2A peptide cleaves between the last glycine and last proline. As another non-limiting example, the polynucleotides of the present invention may include a polynucleotide sequence encoding the 2A peptide having the protein sequence GSGATNFSLLKQAGDVEENPGP (SEQ ID NO: 1) fragments or variants thereof.

[00065] One such polynucleotide sequence encoding the 2A peptide is GGAAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAGGCTGGAGACGTGGAG GAGAACCCTGGACCT (SEQ ID NO: 2). The polynucleotide sequence of the 2A peptide may be modified or codon optimized by the methods described herein and/or are known in the art.

[00066] In one embodiment, this sequence may be used to separate the coding region of two or more polypeptides of interest. As a non-limiting example, the sequence encoding the 2A peptide may be between a first coding region A and a second coding region B (A-2Apep-B). The presence of the 2A peptide would result in the cleavage of one long protein into protein A, protein B and the 2A peptide. Protein A and protein B may be the same or different peptides or polypeptides of interest. In another embodiment, the 2A peptide may be used in the polynucleotides of the present invention to produce two, three, four, five, six, seven, eight, nine, ten or more proteins.

IVT Polynucleotide Architecture

[00067] Traditionally, the basic components of an mRNA molecule include at least a coding region, a 5'UTR, a 3'UTR, a 5' cap and a poly-A tail. The IVT polynucleotides of the present invention may function as mRNA but are distinguished from wild-type mRNA in their functional and/or structural design features which serve to overcome existing problems of effective polypeptide production using nucleic-acid based therapeutics.

[00068] Figure 1 shows a primary construct **100** of an IVT polynucleotide of the present invention. As used herein, "primary construct" refers to a polynucleotide of the present invention which encodes one or more polypeptides of interest and which retains sufficient structural and/or chemical features to allow the polypeptide of interest encoded therein to be translated.

[00069] According to FIG. 1A and 1B, the primary construct **100** of an IVT polynucleotide here contains a first region of linked nucleotides **102** that is flanked by a first flanking region **104** and a second flanking region **106**. The first flanking region **104** may include a sequence of linked nucleosides which function as a 5' untranslated region (UTR) such as the 5' UTR of any of the nucleic acids encoding the native 5'UTR of the polypeptide or a non-native 5'UTR such as, but not limited to, a heterologous 5'UTR or a synthetic 5'UTR. The polypeptide of interest may comprise at its 5' terminus one or more signal sequences encoded by a signal sequence region **103**. The flanking region **104** may comprise a region of linked nucleotides comprising one or more complete or incomplete 5' UTRs sequences which may be completely codon optimized or partially codon optimized. The flanking region **104** may include at least one nucleic acid sequence including, but not limited to, miR sequences, TERZAKTM sequences and translation control sequences. The flanking region **104** may also comprise a 5' terminal cap **108**. The 5' terminal capping region **108** may include a naturally occurring cap, a synthetic cap or an optimized cap. Non-limiting examples of optimized caps include the caps taught by Rhoads in US Patent No. US7074596 and International Patent Publication No. WO2008157668, WO2009149253 and WO2013103659, the contents of each of which are herein incorporated by reference in its entirety. The second flanking region **106** may comprise a region of linked nucleotides comprising one or more complete or incomplete 3' UTRs which may encode the native 3' UTR of the polypeptide or a non-native 3'UTR such as, but not limited to, a heterologous 3'UTR or a synthetic 3' UTR. The second flanking region **106** may be completely codon optimized or partially codon optimized. The flanking region **106** may include at least one nucleic acid sequence including, but not limited to, miR sequences and translation control sequences. The flanking region **106** may also comprise a 3' tailing sequence **110**. The 3' tailing sequence **110** may include a synthetic tailing region **112** and/or a chain terminating nucleoside **114**. Non-limiting examples of a synthetic tailing region include a polyA tail, a polyC tail, a polyA-G quartet and/or a stem loop sequence. Non-limiting examples of chain terminating nucleosides include 2'-O methyl, F and locked nucleic acids (LNA).

[00070] Bridging the 5' terminus of the first region **102** and the first flanking region **104** is a first operational region **105**. Traditionally this operational region comprises a

Start codon. The operational region may alternatively comprise any translation initiation sequence or signal including a Start codon.

[00071] Bridging the 3' terminus of the first region **102** and the second flanking region **106** is a second operational region **107**. Traditionally this operational region comprises a Stop codon. The operational region may alternatively comprise any translation initiation sequence or signal including a Stop codon. Multiple serial stop codons may also be used in the IVT polynucleotide. In one embodiment, the operation region of the present invention may comprise two stop codons. The first stop codon may be "TGA" or "UGA" and the second stop codon may be selected from the group consisting of "TAA," "TGA," "TAG," "UAA," "UGA" or "UAG."

[00072] The shortest length of the first region of the primary construct of the IVT polynucleotide of the present invention can be the length of a nucleic acid sequence that is sufficient to encode for a dipeptide, a tripeptide, a tetrapeptide, a pentapeptide, a hexapeptide, a heptapeptide, an octapeptide, a nonapeptide, or a decapeptide. In another embodiment, the length may be sufficient to encode a peptide of 2-30 amino acids, e.g. 5-30, 10-30, 2-25, 5-25, 10-25, or 10-20 amino acids. The length may be sufficient to encode for a peptide of at least 11, 12, 13, 14, 15, 17, 20, 25 or 30 amino acids, or a peptide that is no longer than 40 amino acids, e.g. no longer than 35, 30, 25, 20, 17, 15, 14, 13, 12, 11 or 10 amino acids. Examples of dipeptides that the polynucleotide sequences can encode or include, but are not limited to, carnosine and anserine.

[00073] The length of the first region of the primary construct of the IVT polynucleotide encoding the polypeptide of interest of the present invention is greater than about 30 nucleotides in length (e.g., at least or greater than about 35, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,900, 2,000, 2,500, and 3,000, 4,000, 5,000, 6,000, 7,000, 8,000, 9,000, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 or up to and including 100,000 nucleotides).

[00074] In some embodiments, the IVT polynucleotide includes from about 30 to about 100,000 nucleotides (e.g., from 30 to 50, from 30 to 100, from 30 to 250, from 30 to 500, from 30 to 1,000, from 30 to 1,500, from 30 to 3,000, from 30 to 5,000, from 30 to 7,000, from 30 to 10,000, from 30 to 25,000, from 30 to 50,000, from 30 to 70,000,

from 100 to 250, from 100 to 500, from 100 to 1,000, from 100 to 1,500, from 100 to 3,000, from 100 to 5,000, from 100 to 7,000, from 100 to 10,000, from 100 to 25,000, from 100 to 50,000, from 100 to 70,000, from 100 to 100,000, from 500 to 1,000, from 500 to 1,500, from 500 to 2,000, from 500 to 3,000, from 500 to 5,000, from 500 to 7,000, from 500 to 10,000, from 500 to 25,000, from 500 to 50,000, from 500 to 70,000, from 500 to 100,000, from 1,000 to 1,500, from 1,000 to 2,000, from 1,000 to 3,000, from 1,000 to 5,000, from 1,000 to 7,000, from 1,000 to 10,000, from 1,000 to 25,000, from 1,000 to 50,000, from 1,000 to 70,000, from 1,000 to 100,000, from 1,500 to 3,000, from 1,500 to 5,000, from 1,500 to 7,000, from 1,500 to 10,000, from 1,500 to 25,000, from 1,500 to 50,000, from 1,500 to 70,000, from 1,500 to 100,000, from 2,000 to 3,000, from 2,000 to 5,000, from 2,000 to 7,000, from 2,000 to 10,000, from 2,000 to 25,000, from 2,000 to 50,000, from 2,000 to 70,000, and from 2,000 to 100,000).

[00075] According to the present invention, the first and second flanking regions of the IVT polynucleotide may range independently from 15-1,000 nucleotides in length (e.g., greater than 30, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, and 900 nucleotides or at least 30, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, and 1,000 nucleotides).

[00076] According to the present invention, the tailing sequence of the IVT polynucleotide may range from absent to 500 nucleotides in length (e.g., at least 60, 70, 80, 90, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, or 500 nucleotides). Where the tailing region is a polyA tail, the length may be determined in units of or as a function of polyA Binding Protein binding. In this embodiment, the polyA tail is long enough to bind at least 4 monomers of PolyA Binding Protein. PolyA Binding Protein monomers bind to stretches of approximately 38 nucleotides. As such, it has been observed that polyA tails of about 80 nucleotides and 160 nucleotides are functional.

[00077] According to the present invention, the capping region of the IVT polynucleotide may comprise a single cap or a series of nucleotides forming the cap. In this embodiment the capping region may be from 1 to 10, e.g. 2-9, 3-8, 4-7, 1-5, 5-10, or at least 2, or 10 or fewer nucleotides in length. In some embodiments, the cap is absent.

[00078] According to the present invention, the first and second operational regions of the IVT polynucleotide may range from 3 to 40, e.g., 5-30, 10-20, 15, or at least 4, or 30 or fewer nucleotides in length and may comprise, in addition to a Start and/or Stop codon, one or more signal and/or restriction sequences.

[00079] In one embodiment, the IVT polynucleotides of the present invention may be structurally modified or chemically modified. When the IVT polynucleotides of the present invention are chemically and/or structurally modified the polynucleotides may be referred to as “modified IVT polynucleotides.”

[00080] In one embodiment, if the IVT polynucleotides of the present invention are chemically modified they may have a uniform chemical modification of all or any of the same nucleoside type or a population of modifications produced by mere downward titration of the same starting modification in all or any of the same nucleoside type, or a measured percent of a chemical modification of all any of the same nucleoside type but with random incorporation, such as where all uridines are replaced by a uridine analog, e.g., pseudouridine. In another embodiment, the IVT polynucleotides may have a uniform chemical modification of two, three, or four of the same nucleoside type throughout the entire polynucleotide (such as all uridines and all cytosines, etc. are modified in the same way).

[00081] In one embodiment, the IVT polynucleotides of the present invention may include a sequence encoding a self-cleaving peptide, described herein, such as but not limited to the 2A peptide. The polynucleotide sequence of the 2A peptide in the IVT polynucleotide may be modified or codon optimized by the methods described herein and/or are known in the art.

[00082] In one embodiment, this sequence may be used to separate the coding region of two or more polypeptides of interest in the IVT polynucleotide.

[00083] In one embodiment, the IVT polynucleotide of the present invention may be structurally and/or chemically modified. When chemically modified and/or structurally modified the IVT polynucleotide may be referred to as a “modified IVT polynucleotide.”

[00084] In one embodiment, the IVT polynucleotide may encode at least one peptide or polypeptide of interest. In another embodiment, the IVT polynucleotide may encode two or more peptides or polypeptides of interest. Non-limiting examples of peptides or

polypeptides of interest include heavy and light chains of antibodies, an enzyme and its substrate, a label and its binding molecule, a second messenger and its enzyme or the components of multimeric proteins or complexes.

[00085] IVT polynucleotides (such as, but not limited to, primary constructs), formulations and compositions comprising IVT polynucleotides, and methods of making, using and administering IVT polynucleotides are described in co-pending International Publication Nos. WO2013151666, WO2013151667, WO2013151668, WO2013151663, WO2013151669, WO2013151670, WO2013151664, WO2013151665, WO2013151736, WO2013151671 and WO2013151672; the contents of each of which are herein incorporated by reference in their entirety.

Chimeric Polynucleotide Architecture

[00086] The chimeric polynucleotides of the present invention maintain a modular organization similar to IVT polynucleotides, but the chimeric polynucleotides comprise one or more structural and/or chemical modifications or alterations which impart useful properties to the polynucleotide. As such, the chimeric polynucleotides which are modified mRNA molecules of the present invention are termed “chimeric modified mRNA” or “chimeric mRNA.”

[00087] Chimeric polynucleotides have portions or regions which differ in size and/or chemical modification pattern, chemical modification position, chemical modification percent or chemical modification population and combinations of the foregoing.

[00088] Examples of parts or regions, where the chimeric polynucleotide functions as an mRNA and encodes a polypeptide of interest include, but are not limited to, untranslated regions (UTRs, such as the 5' UTR or 3' UTR), coding regions, cap regions, polyA tail regions, start regions, stop regions, signal sequence regions, and combinations thereof. Figure 2 illustrates certain embodiments of the chimeric polynucleotides of the invention which may be used as mRNA. Figure 3 illustrates a schematic of a series of chimeric polynucleotides identifying various patterns of positional modifications and showing regions analogous to those regions of an mRNA polynucleotide. Regions or parts that join or lie between other regions may also be designed to have subregions. These are shown in the figure.

[00089] In some embodiments, the chimeric polynucleotides of the invention have a structure comprising Formula I as described in paragraphs [00090] – [00099] in International Patent Publication No. WO2015058069, the contents of which are herein incorporated by reference in its entirety.

[00090] In some embodiments the chimeric polynucleotide of Formula I encodes one or more peptides or polypeptides of interest. Such encoded molecules may be encoded across two or more regions.

[00091] In one embodiment, at least one of the regions of linked nucleosides of A may comprise a sequence of linked nucleosides which can function as a 5' untranslated region (UTR). The sequence of linked nucleosides may be a natural or synthetic 5' UTR. As a non-limiting example, the chimeric polynucleotide may encode a polypeptide of interest and the sequence of linked nucleosides of A may encode the native 5' UTR of a polypeptide encoded by the chimeric polynucleotide or the sequence of linked nucleosides may be a non-heterologous 5' UTR such as, but not limited to a synthetic UTR.

[00092] In another embodiment, at least one of the regions of linked nucleosides of A may be a cap region. The cap region may be located 5' to a region of linked nucleosides of A functioning as a 5'UTR. The cap region may comprise at least one cap such as, but not limited to, Cap0, Cap1, ARCA, inosine, N1-methyl-guanosine, 2'fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, 2-azido-guanosine, Cap2 and Cap4.

[00093] In one embodiment, at least one of the regions of linked nucleosides of B may comprise at least one open reading frame of a nucleic acid sequence. The nucleic acid sequence may be codon optimized and/or comprise at least one modification.

[00094] In one embodiment, at least one of the regions of linked nucleosides of C may comprise a sequence of linked nucleosides which can function as a 3' UTR. The sequence of linked nucleosides may be a natural or synthetic 3' UTR. As a non-limiting example, the chimeric polynucleotide may encode a polypeptide of interest and the sequence of linked nucleosides of C may encode the native 3' UTR of a polypeptide encoded by the chimeric polynucleotide or the sequence of linked nucleosides may be a non-heterologous 3' UTR such as, but not limited to a synthetic UTR.

[00095] In one embodiment, at least one of the regions of linked nucleosides of A comprises a sequence of linked nucleosides which functions as a 5' UTR and at least one of the regions of linked nucleosides of C comprises a sequence of linked nucleosides which functions as a 3' UTR. In one embodiment, the 5' UTR and the 3' UTR may be from the same or different species. In another embodiment, the 5' UTR and the 3' UTR may encode the native untranslated regions from different proteins from the same or different species.

[00096] Figures 4 and 5 provide schematics of a series of chimeric polynucleotides illustrating various patterns of positional modifications based on Formula I as well as those having a blocked or structured 3' terminus.

[00097] Chimeric polynucleotides, including the parts or regions thereof, of the present invention may be classified as hemimers, gapmers, wingmers, or blockmers.

[00098] As used herein, a "hemimer" is chimeric polynucleotide comprising a region or part which comprises half of one pattern, percent, position or population of a chemical modification(s) and half of a second pattern, percent, position or population of a chemical modification(s). Chimeric polynucleotides of the present invention may also comprise hemimer subregions. In one embodiment, a part or region is 50% of one and 50% of another.

[00099] In one embodiment the entire chimeric polynucleotide can be 50% of one and 50% of the other. Any region or part of any chimeric polynucleotide of the invention may be a hemimer. Types of hemimers include pattern hemimers, population hemimers or position hemimers. By definition, hemimers are 50:50 percent hemimers.

[000100] As used herein, a "gapmer" is a chimeric polynucleotide having at least three parts or regions with a gap between the parts or regions. The "gap" can comprise a region of linked nucleosides or a single nucleoside which differs from the chimeric nature of the two parts or regions flanking it. The two parts or regions of a gapmer may be the same or different from each other.

[000101] As used herein, a "wingmer" is a chimeric polynucleotide having at least three parts or regions with a gap between the parts or regions. Unlike a gapmer, the two flanking parts or regions surrounding the gap in a wingmer are the same in degree or kind. Such similarity may be in the length of number of units of different modifications

or in the number of modifications. The wings of a wingmer may be longer or shorter than the gap. The wing parts or regions may be 20, 30, 40, 50, 60 70, 80, 90 or 95% greater or shorter in length than the region which comprises the gap.

[000102] As used herein, a “blockmer” is a patterned polynucleotide where parts or regions are of equivalent size or number and type of modifications. Regions or subregions in a blockmer may be 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490 or 500, nucleosides long.

[000103] Chimeric polynucleotides, including the parts or regions thereof, of the present invention having a chemical modification pattern are referred to as “pattern chimeras.” Pattern chimeras may also be referred to as blockmers. Pattern chimeras are those polynucleotides having a pattern of modifications within, across or among regions or parts.

[000104] Patterns of modifications within a part or region are those which start and stop within a defined region. Patterns of modifications across a part or region are those patterns which start in on part or region and end in another adjacent part or region. Patterns of modifications among parts or regions are those which begin and end in one part or region and are repeated in a different part or region, which is not necessarily adjacent to the first region or part.

[000105] The regions or subregions of pattern chimeras or blockmers may have simple alternating patterns such as ABAB[AB]_n where each “A” and each “B” represent different chemical modifications (at least one of the base, sugar or backbone linker), different types of chemical modifications (e.g., naturally occurring and non-naturally occurring), different percentages of modifications or different populations of modifications. The pattern may repeat n number of times where n=3-300. Further, each A or B can represent from 1-2500 units (e.g., nucleosides) in the pattern. Patterns may also be alternating multiples such as AABBAABB[AABB]_n (an alternating double multiple) or AAABBBAAABBB[AAABBB]_n (an alternating triple multiple) pattern. The pattern may repeat n number of times where n=3-300.

[000106] Different patterns may also be mixed together to form a second order pattern. For example, a single alternating pattern may be combined with a triple alternating pattern to form a second order alternating pattern A'B'. One example would be [ABABAB][AAABBBAAABBB] [ABABAB][AAABBBAAABBB] [ABABAB][AAABBBAAABBB], where [ABABAB] is A' and [AAABBBAAABBB] is B'. In like fashion, these patterns may be repeated n number of times, where n=3-300.

[000107] Patterns may include three or more different modifications to form an ABCABC[ABC]_n pattern. These three component patterns may also be multiples, such as AABBBCCAABBBCC[AABBBCC]_n and may be designed as combinations with other patterns such as ABCABCAABBBCCABCABCAABBBCC, and may be higher order patterns.

[000108] Regions or subregions of position, percent, and population modifications need not reflect an equal contribution from each modification type. They may form series such as “1-2-3-4”, “1-2-4-8”, where each integer represents the number of units of a particular modification type. Alternatively, they may be odd only, such as ‘1-3-3-1-3-1-5’ or even only “2-4-2-4-6-4-8” or a mixture of both odd and even number of units such as “1-3-4-2-5-7-3-3-4”.

[000109] Pattern chimeras may vary in their chemical modification by degree (such as those described above) or by kind (e.g., different modifications).

[000110] Chimeric polynucleotides, including the parts or regions thereof, of the present invention having at least one region with two or more different chemical

modifications of two or more nucleoside members of the same nucleoside type (A, C, G, T, or U) are referred to as “positionally modified” chimeras. Positionally modified chimeras are also referred to herein as “selective placement” chimeras or “selective placement polynucleotides”. As the name implies, selective placement refers to the design of polynucleotides which, unlike polynucleotides in the art where the modification to any A, C, G, T or U is the same by virtue of the method of synthesis, can have different modifications to the individual As, Cs, Gs, Ts or Us in a polynucleotide or region thereof. For example, in a positionally modified chimeric polynucleotide, there may be two or more different chemical modifications to any of the nucleoside types of As, Cs, Gs, Ts, or Us. There may also be combinations of two or more to any two or more of the same nucleoside type. For example, a positionally modified or selective placement chimeric polynucleotide may comprise 3 different modifications to the population of adenines in the molecule and also have 3 different modifications to the population of cytosines in the construct—all of which may have a unique, non-random, placement.

[000111] Chimeric polynucleotides, including the parts or regions thereof, of the present invention having a chemical modification percent are referred to as “percent chimeras.” Percent chimeras may have regions or parts which comprise at least 1%, at least 2%, at least 5%, at least 8%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% positional, pattern or population of modifications. Alternatively, the percent chimera may be completely modified as to modification position, pattern, or population. The percent of modification of a percent chimera may be split between naturally occurring and non-naturally occurring modifications.

[000112] Chimeric polynucleotides, including the parts or regions thereof, of the present invention having a chemical modification population are referred to as “population chimeras.” A population chimera may comprise a region or part where nucleosides (their base, sugar or backbone linkage, or combination thereof) have a select population of modifications. Such modifications may be selected from functional populations such as modifications which induce, alter or modulate a phenotypic outcome. For example, a functional population may be a population or selection of chemical

modifications which increase the level of a cytokine. Other functional populations may individually or collectively function to decrease the level of one or more cytokines. Use of a selection of these like-function modifications in a chimeric polynucleotide would therefore constitute a “functional population chimera.” As used herein, a “functional population chimera” may be one whose unique functional feature is defined by the population of modifications as described above or the term may apply to the overall function of the chimeric polynucleotide itself. For example, as a whole the chimeric polynucleotide may function in a different or superior way as compared to an unmodified or non-chimeric polynucleotide.

[000113] It should be noted that polynucleotides which have a uniform chemical modification of all of any of the same nucleoside type or a population of modifications produced by mere downward titration of the same starting modification in all of any of the same nucleoside type, or a measured percent of a chemical modification of all any of the same nucleoside type but with random incorporation, such as where all uridines are replaced by a uridine analog, e.g., pseudouridine, are not considered chimeric. Likewise, polynucleotides having a uniform chemical modification of two, three, or four of the same nucleoside type throughout the entire polynucleotide (such as all uridines and all cytosines, etc. are modified in the same way) are not considered chimeric polynucleotides. One example of a polynucleotide which is not chimeric is the canonical pseudouridine/5-methyl cytosine modified polynucleotide of the prior art. These uniform polynucleotides are arrived at entirely via *in vitro* transcription (IVT) enzymatic synthesis; and due to the limitations of the synthesizing enzymes, they contain only one kind of modification at the occurrence of each of the same nucleoside type, i.e., adenosine (A), thymidine (T), guanosine (G), cytidine (C) or uridine (U), found in the polynucleotide. Such polynucleotides may be characterized as IVT polynucleotides.

[000114] The chimeric polynucleotides of the present invention may be structurally modified or chemically modified. When the chimeric polynucleotides of the present invention are chemically and/or structurally modified the polynucleotides may be referred to as “modified chimeric polynucleotides.”

[000115] In some embodiments of the invention, the chimeric polynucleotides may encode two or more peptides or polypeptides of interest. Such peptides or polypeptides of

interest include the heavy and light chains of antibodies, an enzyme and its substrate, a label and its binding molecule, a second messenger and its enzyme or the components of multimeric proteins or complexes.

[000116] The regions or parts of the chimeric polynucleotides of the present invention may be separated by a linker or spacer moiety. Such linkers or spaces may be nucleic acid based or non-nucleosidic.

[000117] In one embodiment, the chimeric polynucleotides of the present invention may include a sequence encoding a self-cleaving peptide described herein, such as, but not limited to, a 2A peptide. The polynucleotide sequence of the 2A peptide in the chimeric polynucleotide may be modified or codon optimized by the methods described herein and/or are known in the art.

[000118] Notwithstanding the foregoing, the chimeric polynucleotides of the present invention may comprise a region or part which is not positionally modified or not chimeric as defined herein.

[000119] For example, a region or part of a chimeric polynucleotide may be uniformly modified at one or more A, T, C, G, or U but according to the invention, the polynucleotides will not be uniformly modified throughout the entire region or part.

[000120] Regions or parts of chimeric polynucleotides may be from 15-1000 nucleosides in length and a polynucleotide may have from 2-100 different regions or patterns of regions as described herein.

[000121] In one embodiment, chimeric polynucleotides encode one or more polypeptides of interest. In another embodiment, the chimeric polynucleotides are substantially non-coding. In another embodiment, the chimeric polynucleotides have both coding and non-coding regions and parts.

[000122] Figure 2 illustrates the design of certain chimeric polynucleotides of the present invention when based on the scaffold of the polynucleotide of Figure 1. Shown in the figure are the regions or parts of the chimeric polynucleotides where patterned regions represent those regions which are positionally modified and open regions illustrate regions which may or may not be modified but which are, when modified, uniformly modified. Chimeric polynucleotides of the present invention may be completely positionally modified or partially positionally modified. They may also have

subregions which may be of any pattern or design. Shown in Figure 2 are a chimeric subregion and a hemimer subregion.

[000123] In one embodiment, the shortest length of a region of the chimeric polynucleotide of the present invention encoding a peptide can be the length that is sufficient to encode for a dipeptide, a tripeptide, a tetrapeptide, a pentapeptide, a hexapeptide, a heptapeptide, an octapeptide, a nonapeptide, or a decapeptide. In another embodiment, the length may be sufficient to encode a peptide of 2-30 amino acids, e.g. 5-30, 10-30, 2-25, 5-25, 10-25, or 10-20 amino acids. The length may be sufficient to encode for a peptide of at least 11, 12, 13, 14, 15, 17, 20, 25 or 30 amino acids, or a peptide that is no longer than 40 amino acids, e.g. no longer than 35, 30, 25, 20, 17, 15, 14, 13, 12, 11 or 10 amino acids. Examples of dipeptides that the polynucleotide sequences can encode or include, but are not limited to, carnosine and anserine.

[000124] In one embodiment, the length of a region of the chimeric polynucleotide of the present invention encoding the peptide or polypeptide of interest is greater than about 30 nucleotides in length (e.g., at least or greater than about 35, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,900, 2,000, 2,500, and 3,000, 4,000, 5,000, 6,000, 7,000, 8,000, 9,000, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 or up to and including 100,000 nucleotides). As used herein, such a region may be referred to as a “coding region” or “region encoding.”

[000125] In some embodiments, the chimeric polynucleotide includes from about 30 to about 100,000 nucleotides (e.g., from 30 to 50, from 30 to 100, from 30 to 250, from 30 to 500, from 30 to 1,000, from 30 to 1,500, from 30 to 3,000, from 30 to 5,000, from 30 to 7,000, from 30 to 10,000, from 30 to 25,000, from 30 to 50,000, from 30 to 70,000, from 100 to 250, from 100 to 500, from 100 to 1,000, from 100 to 1,500, from 100 to 3,000, from 100 to 5,000, from 100 to 7,000, from 100 to 10,000, from 100 to 25,000, from 100 to 50,000, from 100 to 70,000, from 100 to 100,000, from 500 to 1,000, from 500 to 1,500, from 500 to 2,000, from 500 to 3,000, from 500 to 5,000, from 500 to 7,000, from 500 to 10,000, from 500 to 25,000, from 500 to 50,000, from 500 to 70,000, from 500 to 100,000, from 1,000 to 1,500, from 1,000 to 2,000, from 1,000 to 3,000, from 1,000 to 5,000, from 1,000 to 7,000, from 1,000 to 10,000, from 1,000 to 25,000,

from 1,000 to 50,000, from 1,000 to 70,000, from 1,000 to 100,000, from 1,500 to 3,000, from 1,500 to 5,000, from 1,500 to 7,000, from 1,500 to 10,000, from 1,500 to 25,000, from 1,500 to 50,000, from 1,500 to 70,000, from 1,500 to 100,000, from 2,000 to 3,000, from 2,000 to 5,000, from 2,000 to 7,000, from 2,000 to 10,000, from 2,000 to 25,000, from 2,000 to 50,000, from 2,000 to 70,000, and from 2,000 to 100,000).

[000126] According to the present invention, regions or subregions of the chimeric polynucleotides may also range independently from 15-1,000 nucleotides in length (e.g., greater than 30, 40, 45, 50, 55, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 550, 600, 650, 700, 750, 800, 850, 900 and 950 nucleotides or at least 30, 40, 45, 50, 55, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950 and 1,000 nucleotides).

[000127] According to the present invention, regions or subregions of chimeric polynucleotides may range from absent to 500 nucleotides in length (e.g., at least 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, or 500 nucleotides). Where the region is a polyA tail, the length may be determined in units of or as a function of polyA Binding Protein binding. In this embodiment, the polyA tail is long enough to bind at least 4 monomers of PolyA Binding Protein. PolyA Binding Protein monomers bind to stretches of approximately 38 nucleotides. As such, it has been observed that polyA tails of about 80 nucleotides to about 160 nucleotides are functional. The chimeric polynucleotides of the present invention which function as an mRNA need not comprise a polyA tail.

[000128] According to the present invention, chimeric polynucleotides which function as an mRNA may have a capping region. The capping region may comprise a single cap or a series of nucleotides forming the cap. In this embodiment the capping region may be from 1 to 10, e.g. 2-9, 3-8, 4-7, 1-5, 5-10, or at least 2, or 10 or fewer nucleotides in length. In some embodiments, the cap is absent.

[000129] The present invention contemplates chimeric polynucleotides which are circular or cyclic. As the name implies circular polynucleotides are circular in nature meaning that the termini are joined in some fashion, whether by ligation, covalent bond,

common association with the same protein or other molecule or complex or by hybridization. Any of the circular polynucleotides as taught in, for example, International Patent Publication No. WO2015034925 (Attorney Docket number M51.20), the contents of which are incorporated herein by reference in their entirety, may be made chimeric according to the present invention.

[000130] Chimeric polynucleotides, formulations and compositions comprising chimeric polynucleotides, and methods of making, using and administering chimeric polynucleotides are also described in International Patent Publication No. WO2015034928 (Attorney Docket No. M057.20), the contents of which is incorporated by reference in its entirety.

Circular Polynucleotide Architecture

[000131] The present invention contemplates polynucleotides which are circular or cyclic. As the name implies circular polynucleotides are circular in nature meaning that the termini are joined in some fashion, whether by ligation, covalent bond, common association with the same protein or other molecule or complex or by hybridization. Any of the circular polynucleotides as taught in, for example, International Patent Publication No. WO2015034925 (Attorney Docket number M51.20), the contents of which are incorporated herein by reference in their entirety.

[000132] Circular polynucleotides of the present invention may be designed according to the circular RNA construct scaffolds shown in Figures 6-12 of International Patent Publication No. WO2015058069, the contents of which are herein incorporated by reference in its entirety. Figures 6-12 are described in paragraphs [000145] – [000154] of International Patent Publication No. WO2015058069, the contents of which are herein incorporated by reference in its entirety. Such polynucleotides are circular polynucleotides or circular constructs.

[000133] The circular polynucleotides or circPs of the present invention which encode at least one peptide or polypeptide of interest are known as circular RNAs or circRNA. As used herein, “circular RNA” or “circRNA” means a circular polynucleotide that can encode at least one peptide or polypeptide of interest. The circPs of the present invention which comprise at least one sensor sequence and do not encode a peptide or polypeptide of interest are known as circular sponges or circSP. As used herein, “circular sponges,”

“circular polynucleotide sponges” or “circSP” means a circular polynucleotide which comprises at least one sensor sequence and does not encode a polypeptide of interest. As used herein, “sensor sequence” means a receptor or pseudo-receptor for endogenous nucleic acid binding molecules. Non-limiting examples of sensor sequences include, microRNA binding sites, microRNA seed sequences, microRNA binding sites without the seed sequence, transcription factor binding sites and artificial binding sites engineered to act as pseudo-receptors and portions and fragments thereof.

[000134] The circPs of the present invention which comprise at least one sensor sequence and encode at least one peptide or polypeptide of interest are known as circular RNA sponges or circRNA-SP. As used herein, “circular RNA sponges” or “circRNA-SP” means a circular polynucleotide which comprises at least one sensor sequence and at least one region encoding at least one peptide or polypeptide of interest.

[000135] Circular polynucleotides, formulations and compositions comprising circular polynucleotides, and methods of making, using and administering circular polynucleotides are also described in International Patent Publication No. WO2015034925, the contents of which is incorporated by reference in its entirety.

Multimers of Polynucleotides

[000136] According to the present invention, multiple distinct chimeric polynucleotides and/or IVT polynucleotides may be linked together through the 3'-end using nucleotides which are modified at the 3'-terminus. Chemical conjugation may be used to control the stoichiometry of delivery into cells. For example, the glyoxylate cycle enzymes, isocitrate lyase and malate synthase, may be supplied into cells at a 1:1 ratio to alter cellular fatty acid metabolism. This ratio may be controlled by chemically linking chimeric polynucleotides and/or IVT polynucleotides using a 3'-azido terminated nucleotide on one polynucleotides species and a C5-ethynyl or alkynyl-containing nucleotide on the opposite polynucleotide species. The modified nucleotide is added post-transcriptionally using terminal transferase (New England Biolabs, Ipswich, MA) according to the manufacturer's protocol. After the addition of the 3'-modified nucleotide, the two polynucleotides species may be combined in an aqueous solution, in the presence or absence of copper, to form a new covalent linkage via a click chemistry mechanism as described in the literature.

[000137] In another example, more than two chimeric polynucleotides and/or IVT polynucleotides may be linked together using a functionalized linker molecule. For example, a functionalized saccharide molecule may be chemically modified to contain multiple chemical reactive groups (SH-, NH₂-, N₃, etc...) to react with the cognate moiety on a 3'-functionalized mRNA molecule (i.e., a 3'-maleimide ester, 3'-NHS-ester, alkynyl). The number of reactive groups on the modified saccharide can be controlled in a stoichiometric fashion to directly control the stoichiometric ratio of conjugated chimeric polynucleotides and/or IVT polynucleotides.

[000138] In one embodiment, the chimeric polynucleotides and/or IVT polynucleotides may be linked together in a pattern. The pattern may be a simple alternating pattern such as CD[CD]_x where each "C" and each "D" represent a chimeric polynucleotide, IVT polynucleotide, different chimeric polynucleotides or different IVT polynucleotides. The pattern may repeat x number of times, where x= 1-300. Patterns may also be alternating multiples such as CCDD[CCDD]_x (an alternating double multiple) or CCCDDD[CCDDDD]_x (an alternating triple multiple) pattern. The alternating double multiple or alternating triple multiple may repeat x number of times, where x= 1-300.

Conjugates and Combinations of Polynucleotides

[000139] In order to further enhance protein production, polynucleotides of the present invention can be designed to be conjugated to other polynucleotides, dyes, intercalating agents (*e.g.* acridines), cross-linkers (*e.g.* psoralene, mitomycin C), porphyrins (TPPC4, texaphyrin, Sapphyrin), polycyclic aromatic hydrocarbons (*e.g.*, phenazine, dihydrophenazine), artificial endonucleases (*e.g.* EDTA), alkylating agents, phosphate, amino, mercapto, PEG (*e.g.*, PEG-40K), MPEG, [MPEG]₂, polyamino, alkyl, substituted alkyl, radiolabeled markers, enzymes, haptens (*e.g.* biotin), transport/absorption facilitators (*e.g.*, aspirin, vitamin E, folic acid), synthetic ribonucleases, proteins, *e.g.*, glycoproteins, or peptides, *e.g.*, molecules having a specific affinity for a co-ligand, or antibodies *e.g.*, an antibody, that binds to a specified cell type such as a cancer cell, endothelial cell, or bone cell, hormones and hormone receptors, non-peptidic species, such as lipids, lectins, carbohydrates, vitamins, cofactors, or a drug.

[000140] Conjugation may result in increased stability and/or half-life and may be particularly useful in targeting the polynucleotides to specific sites in the cell, tissue or organism.

[000141] According to the present invention, the polynucleotides may be administered with, conjugated to or further encode one or more of RNAi agents, siRNAs, shRNAs, miRNAs, miRNA binding sites, antisense RNAs, ribozymes, catalytic DNA, tRNA, RNAs that induce triple helix formation, aptamers or vectors, and the like.

Bifunctional Polynucleotides

[000142] In one embodiment of the invention are bifunctional polynucleotides (e.g., bifunctional IVT polynucleotides, bifunctional chimeric polynucleotides or bifunctional circular polynucleotides). As the name implies, bifunctional polynucleotides are those having or capable of at least two functions. These molecules may also by convention be referred to as multi-functional.

[000143] The multiple functionalities of bifunctional polynucleotides may be encoded by the RNA (the function may not manifest until the encoded product is translated) or may be a property of the polynucleotide itself. It may be structural or chemical.

Bifunctional modified polynucleotides may comprise a function that is covalently or electrostatically associated with the polynucleotides. Further, the two functions may be provided in the context of a complex of a chimeric polynucleotide and another molecule.

[000144] Bifunctional polynucleotides may encode peptides which are anti-proliferative. These peptides may be linear, cyclic, constrained or random coil. They may function as aptamers, signaling molecules, ligands or mimics or mimetics thereof. Anti-proliferative peptides may, as translated, be from 3 to 50 amino acids in length. They may be 5-40, 10-30, or approximately 15 amino acids long. They may be single chain, multichain or branched and may form complexes, aggregates or any multi-unit structure once translated.

Noncoding Polynucleotides

[000145] As described herein, provided are polynucleotides having sequences that are partially or substantially not translatable, e.g., having a noncoding region. As one non-limiting example, the noncoding region may be the first region of the IVT polynucleotide or the circular polynucleotide. Alternatively, the noncoding region may be a region other

than the first region. As another non-limiting example, the noncoding region may be the A, B and/or C region of the chimeric polynucleotide.

[000146] Such molecules are generally not translated, but can exert an effect on protein production by one or more of binding to and sequestering one or more translational machinery components such as a ribosomal protein or a transfer RNA (tRNA), thereby effectively reducing protein expression in the cell or modulating one or more pathways or cascades in a cell which in turn alters protein levels. The polynucleotide may contain or encode one or more long noncoding RNA (lncRNA, or lincRNA) or portion thereof, a small nucleolar RNA (sno-RNA), micro RNA (miRNA), small interfering RNA (siRNA) or Piwi-interacting RNA (piRNA). Examples of such lncRNA molecules and RNAi constructs designed to target such lncRNA any of which may be encoded in the polynucleotides are taught in International Publication, WO2012/018881 A2, the contents of which are incorporated herein by reference in their entirety.

Tolerogenic Polypeptides of Interest

[000147] Polynucleotides of the present invention may encode one or more tolerogenic peptides or polypeptides of interest. They may also affect the levels, signaling or function of one or more peptides or polypeptides. Tolerogenic polypeptides of interest, according to the present invention include any of those taught in, for example, those listed in Table 6 of International Publication Nos. WO2013151666, WO2013151667, WO2013151668, WO2013151663, WO2013151669, WO2013151670, WO2013151664, WO2013151665, WO2013151736, WO2013151671 and WO2013151672 and Table 178 of International Publication No. WO2013151671; the contents of each of which are herein incorporated by reference in their entireties.

[000148] According to the present invention, the polynucleotide may be designed to encode one or more polypeptides of interest or fragments thereof. Such polypeptide of interest may include, but is not limited to, whole polypeptides, a plurality of polypeptides or fragments of polypeptides, which independently may be encoded by one or more regions or parts or the whole of a polynucleotide. As used herein, the term “polypeptides of interest” refer to any polypeptide which is selected to be encoded within, or whose function is affected by, the polynucleotides of the present invention.

[000149] As used herein, “polypeptide” means a polymer of amino acid residues (natural or unnatural) linked together most often by peptide bonds. The term, as used herein, refers to proteins, polypeptides, and peptides of any size, structure, or function. In some instances the polypeptide encoded is smaller than about 50 amino acids and the polypeptide is then termed a peptide. If the polypeptide is a peptide, it will be at least about 2, 3, 4, or at least 5 amino acid residues long. Thus, polypeptides include gene products, naturally occurring polypeptides, synthetic polypeptides, homologs, orthologs, paralogs, fragments and other equivalents, variants, and analogs of the foregoing. A polypeptide may be a single molecule or may be a multi-molecular complex such as a dimer, trimer or tetramer. They may also comprise single chain or multichain polypeptides such as antibodies or insulin and may be associated or linked. Most commonly disulfide linkages are found in multichain polypeptides. The term polypeptide may also apply to amino acid polymers in which one or more amino acid residues are an artificial chemical analogue of a corresponding naturally occurring amino acid.

[000150] The term “polypeptide variant” refers to molecules which differ in their amino acid sequence from a native or reference sequence. The amino acid sequence variants may possess substitutions, deletions, and/or insertions at certain positions within the amino acid sequence, as compared to a native or reference sequence. Ordinarily, variants will possess at least about 50% identity (homology) to a native or reference sequence, and preferably, they will be at least about 80%, more preferably at least about 90% identical (homologous) to a native or reference sequence.

[000151] In some embodiments “variant mimics” are provided. As used herein, the term “variant mimic” is one which contains one or more amino acids which would mimic an activated sequence. For example, glutamate may serve as a mimic for phospho-threonine and/or phospho-serine. Alternatively, variant mimics may result in deactivation or in an inactivated product containing the mimic, e.g., phenylalanine may act as an inactivating substitution for tyrosine; or alanine may act as an inactivating substitution for serine.

[000152] “Homology” as it applies to amino acid sequences is defined as the percentage of residues in the candidate amino acid sequence that are identical with the residues in the amino acid sequence of a second sequence after aligning the sequences and introducing

gaps, if necessary, to achieve the maximum percent homology. Methods and computer programs for the alignment are well known in the art. It is understood that homology depends on a calculation of percent identity but may differ in value due to gaps and penalties introduced in the calculation.

[000153] By “homologs” as it applies to polypeptide sequences means the corresponding sequence of other species having substantial identity to a second sequence of a second species.

[000154] “Analog” is meant to include polypeptide variants which differ by one or more amino acid alterations, e.g., substitutions, additions or deletions of amino acid residues that still maintain one or more of the properties of the parent or starting polypeptide.

[000155] The present invention contemplates several types of compositions which are polypeptide based including variants and derivatives. These include substitutional, insertional, deletion and covalent variants and derivatives. The term “derivative” is used synonymously with the term “variant” but generally refers to a molecule that has been modified and/or changed in any way relative to a reference molecule or starting molecule.

[000156] As such, polynucleotides encoding peptides or polypeptides containing substitutions, insertions and/or additions, deletions and covalent modifications with respect to reference sequences, in particular the polypeptide sequences disclosed herein, are included within the scope of this invention. For example, sequence tags or amino acids, such as one or more lysines, can be added to the peptide sequences of the invention (e.g., at the N-terminal or C-terminal ends). Sequence tags can be used for peptide purification or localization. Lysines can be used to increase peptide solubility or to allow for biotinylation. Alternatively, amino acid residues located at the carboxy and amino terminal regions of the amino acid sequence of a peptide or protein may optionally be deleted providing for truncated sequences. Certain amino acids (e.g., C-terminal or N-terminal residues) may alternatively be deleted depending on the use of the sequence, as for example, expression of the sequence as part of a larger sequence which is soluble, or linked to a solid support.

[000157] “Substitutional variants” when referring to polypeptides are those that have at least one amino acid residue in a native or starting sequence removed and a different

amino acid inserted in its place at the same position. The substitutions may be single, where only one amino acid in the molecule has been substituted, or they may be multiple, where two or more amino acids have been substituted in the same molecule.

[000158] As used herein the term “conservative amino acid substitution” refers to the substitution of an amino acid that is normally present in the sequence with a different amino acid of similar size, charge, or polarity. Examples of conservative substitutions include the substitution of a non-polar (hydrophobic) residue such as isoleucine, valine and leucine for another non-polar residue. Likewise, examples of conservative substitutions include the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, and between glycine and serine. Additionally, the substitution of a basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue such as aspartic acid or glutamic acid for another acidic residue are additional examples of conservative substitutions. Examples of non-conservative substitutions include the substitution of a non-polar (hydrophobic) amino acid residue such as isoleucine, valine, leucine, alanine, methionine for a polar (hydrophilic) residue such as cysteine, glutamine, glutamic acid or lysine and/or a polar residue for a non-polar residue.

[000159] “Insertional variants” when referring to polypeptides are those with one or more amino acids inserted immediately adjacent to an amino acid at a particular position in a native or starting sequence. “Immediately adjacent” to an amino acid means connected to either the alpha-carboxy or alpha-amino functional group of the amino acid.

[000160] “Deletional variants” when referring to polypeptides are those with one or more amino acids in the native or starting amino acid sequence removed. Ordinarily, deletional variants will have one or more amino acids deleted in a particular region of the molecule.

[000161] “Covalent derivatives” when referring to polypeptides include modifications of a native or starting protein with an organic proteinaceous or non-proteinaceous derivatizing agent, and/or post-translational modifications. Covalent modifications are traditionally introduced by reacting targeted amino acid residues of the protein with an organic derivatizing agent that is capable of reacting with selected side-chains or terminal residues, or by harnessing mechanisms of post-translational modifications that function in

selected recombinant host cells. The resultant covalent derivatives are useful in programs directed at identifying residues important for biological activity, for immunoassays, or for the preparation of anti-protein antibodies for immunoaffinity purification of the recombinant glycoprotein. Such modifications are within the ordinary skill in the art and are performed without undue experimentation.

[000162] Certain post-translational modifications are the result of the action of recombinant host cells on the expressed polypeptide. Glutaminyl and asparaginyl residues are frequently post-translationally deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues may be present in the polypeptides produced in accordance with the present invention.

[000163] Other post-translational modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the alpha-amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)).

[000164] “Features” when referring to polypeptides are defined as distinct amino acid sequence-based components of a molecule. Features of the polypeptides encoded by the polynucleotides of the present invention include surface manifestations, local conformational shape, folds, loops, half-loops, domains, half-domains, sites, termini or any combination thereof.

[000165] As used herein when referring to polypeptides the term “surface manifestation” refers to a polypeptide based component of a protein appearing on an outermost surface.

[000166] As used herein when referring to polypeptides the term “local conformational shape” means a polypeptide based structural manifestation of a protein which is located within a definable space of the protein.

[000167] As used herein when referring to polypeptides the term “fold” refers to the resultant conformation of an amino acid sequence upon energy minimization. A fold may occur at the secondary or tertiary level of the folding process. Examples of secondary level folds include beta sheets and alpha helices. Examples of tertiary folds include

domains and regions formed due to aggregation or separation of energetic forces.

Regions formed in this way include hydrophobic and hydrophilic pockets, and the like.

[000168] As used herein the term “turn” as it relates to protein conformation means a bend which alters the direction of the backbone of a peptide or polypeptide and may involve one, two, three or more amino acid residues.

[000169] As used herein when referring to polypeptides the term “loop” refers to a structural feature of a polypeptide which may serve to reverse the direction of the backbone of a peptide or polypeptide. Where the loop is found in a polypeptide and only alters the direction of the backbone, it may comprise four or more amino acid residues. Oliva et al. have identified at least 5 classes of protein loops (J. Mol Biol 266 (4): 814-830; 1997). Loops may be open or closed. Closed loops or “cyclic” loops may comprise 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acids between the bridging moieties. Such bridging moieties may comprise a cysteine-cysteine bridge (Cys-Cys) typical in polypeptides having disulfide bridges or alternatively bridging moieties may be non-protein based such as the dibromozyl agents used herein.

[000170] As used herein when referring to polypeptides the term “half-loop” refers to a portion of an identified loop having at least half the number of amino acid residues as the loop from which it is derived. It is understood that loops may not always contain an even number of amino acid residues. Therefore, in those cases where a loop contains or is identified to comprise an odd number of amino acids, a half-loop of the odd-numbered loop will comprise the whole number portion or next whole number portion of the loop (number of amino acids of the loop/2+/-0.5 amino acids). For example, a loop identified as a 7 amino acid loop could produce half-loops of 3 amino acids or 4 amino acids ($7/2=3.5\pm0.5$ being 3 or 4).

[000171] As used herein when referring to polypeptides the term “domain” refers to a motif of a polypeptide having one or more identifiable structural or functional characteristics or properties (e.g., binding capacity, serving as a site for protein-protein interactions).

[000172] As used herein when referring to polypeptides the term “half-domain” means a portion of an identified domain having at least half the number of amino acid residues as the domain from which it is derived. It is understood that domains may not always

contain an even number of amino acid residues. Therefore, in those cases where a domain contains or is identified to comprise an odd number of amino acids, a half-domain of the odd-numbered domain will comprise the whole number portion or next whole number portion of the domain (number of amino acids of the domain/2+/-0.5 amino acids). For example, a domain identified as a 7 amino acid domain could produce half-domains of 3 amino acids or 4 amino acids ($7/2=3.5\pm0.5$ being 3 or 4). It is also understood that sub-domains may be identified within domains or half-domains, these subdomains possessing less than all of the structural or functional properties identified in the domains or half domains from which they were derived. It is also understood that the amino acids that comprise any of the domain types herein need not be contiguous along the backbone of the polypeptide (i.e., nonadjacent amino acids may fold structurally to produce a domain, half-domain or subdomain).

[000173] As used herein when referring to polypeptides the terms “site” as it pertains to amino acid based embodiments is used synonymously with “amino acid residue” and “amino acid side chain.” A site represents a position within a peptide or polypeptide that may be modified, manipulated, altered, derivatized or varied within the polypeptide based molecules of the present invention.

[000174] As used herein the terms “termini” or “terminus” when referring to polypeptides refers to an extremity of a peptide or polypeptide. Such extremity is not limited only to the first or final site of the peptide or polypeptide but may include additional amino acids in the terminal regions. The polypeptide based molecules of the present invention may be characterized as having both an N-terminus (terminated by an amino acid with a free amino group (NH₂)) and a C-terminus (terminated by an amino acid with a free carboxyl group (COOH)). Proteins of the invention are in some cases made up of multiple polypeptide chains brought together by disulfide bonds or by non-covalent forces (multimers, oligomers). These sorts of proteins will have multiple N- and C-termini. Alternatively, the termini of the polypeptides may be modified such that they begin or end, as the case may be, with a non-polypeptide based moiety such as an organic conjugate.

[000175] Once any of the features have been identified or defined as a desired component of a polypeptide to be encoded by the polynucleotide of the invention, any of

several manipulations and/or modifications of these features may be performed by moving, swapping, inverting, deleting, randomizing or duplicating. Furthermore, it is understood that manipulation of features may result in the same outcome as a modification to the molecules of the invention. For example, a manipulation which involved deleting a domain would result in the alteration of the length of a molecule just as modification of a nucleic acid to encode less than a full length molecule would.

[000176] Modifications and manipulations can be accomplished by methods known in the art such as, but not limited to, site directed mutagenesis or *a priori* incorporation during chemical synthesis. The resulting modified molecules may then be tested for activity using *in vitro* or *in vivo* assays such as those described herein or any other suitable screening assay known in the art.

[000177] According to the present invention, the polypeptides may comprise a consensus sequence which is discovered through rounds of experimentation. As used herein a “consensus” sequence is a single sequence which represents a collective population of sequences allowing for variability at one or more sites.

[000178] As recognized by those skilled in the art, protein fragments, functional protein domains, and homologous proteins are also considered to be within the scope of polypeptides of interest of this invention. For example, provided herein is any protein fragment (meaning a polypeptide sequence at least one amino acid residue shorter than a reference polypeptide sequence but otherwise identical) of a reference protein 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or greater than 100 amino acids in length. In another example, any protein that includes a stretch of about 20, about 30, about 40, about 50, or about 100 amino acids which are about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 95%, or about 100% identical to any of the sequences described herein can be utilized in accordance with the invention. In certain embodiments, a polypeptide to be utilized in accordance with the invention includes 2, 3, 4, 5, 6, 7, 8, 9, 10, or more mutations as shown in any of the sequences provided or referenced herein.

Types of Polypeptides of Interest

[000179] The polynucleotides of the present invention may be designed to encode tolerogenic polypeptides of interest.

[000180] In one embodiment, polynucleotides may encode variant polypeptides which have a certain identity with a reference polypeptide sequence. As used herein, a “reference polypeptide sequence” refers to a starting polypeptide sequence. Reference sequences may be wild type sequences or any sequence to which reference is made in the design of another sequence. A “reference polypeptide sequence” may, e.g., be any one of those polypeptides disclosed in Table 6 of U International Publication Nos.

WO2013151666, WO2013151667, WO2013151668, WO2013151663, WO2013151669, WO2013151670, WO2013151664, WO2013151665, WO2013151736, WO2013151671 and WO2013151672 and Table 178 of International Publication No. WO2013151671; the contents of each of which are herein incorporated by reference in their entireties.

[000181] Reference molecules (polypeptides or polynucleotides) may share a certain identity with the designed molecules (polypeptides or polynucleotides). The term “identity” as known in the art, refers to a relationship between the sequences of two or more peptides, polypeptides or polynucleotides, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between them as determined by the number of matches between strings of two or more amino acid residues or nucleosides. Identity measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., “algorithms”). Identity of related peptides can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; and Carillo et al., SIAM J. Applied Math. 48, 1073 (1988).

[000182] In some embodiments, the encoded polypeptide variant may have the same or a similar activity as the reference polypeptide. Alternatively, the variant may have an altered activity (e.g., increased or decreased) relative to a reference polypeptide. Generally, variants of a particular polynucleotide or polypeptide of the invention will

have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% but less than 100% sequence identity to that particular reference polynucleotide or polypeptide as determined by sequence alignment programs and parameters described herein and known to those skilled in the art. Such tools for alignment include those of the BLAST suite (Stephen F. Altschul, Thomas L. Madden, Alejandro A. Sch  ffer, Jinghui Zhang, Zheng Zhang, Webb Miller, and David J. Lipman (1997), "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", Nucleic Acids Res. 25:3389-3402.) Other tools are described herein, specifically in the definition of "Identity."

[000183] Default parameters in the BLAST algorithm include, for example, an expect threshold of 10, Word size of 28, Match/Mismatch Scores 1, -2, Gap costs Linear. Any filter can be applied as well as a selection for species specific repeats, e.g., *Homo sapiens*. *Polynucleotides having Untranslated Regions (UTRs)*

[000184] The polynucleotides of the present invention may comprise one or more regions or parts which act or function as an untranslated region. Where polynucleotides are designed to encode at least one polypeptide of interest, the polynucleotides may comprise one or more of these untranslated regions.

[000185] By definition, wild type untranslated regions (UTRs) of a gene are transcribed but not translated. In mRNA, the 5'UTR starts at the transcription start site and continues to the start codon but does not include the start codon; whereas, the 3'UTR starts immediately following the stop codon and continues until the transcriptional termination signal. There is growing body of evidence about the regulatory roles played by the UTRs in terms of stability of the nucleic acid molecule and translation. The regulatory features of a UTR can be incorporated into the polynucleotides of the present invention to, among other things, enhance the stability of the molecule. The specific features can also be incorporated to ensure controlled down-regulation of the transcript in case they are misdirected to undesired organs sites.

[000186] Tables 1 and 2 provide a listing of exemplary UTRs which may be utilized in the polynucleotides of the present invention. Shown in Table 1 is a listing of a 5'-untranslated region of the invention. Variants of 5' UTRs may be utilized wherein one or more nucleotides are added or removed to the termini, including A, T, C or G.

Table 1. 5'-Untranslated Regions

5' UTR Identifier	Name/ Description	SEQ ID NO.
5UTR-001	Upstream UTR	3
5UTR-002	Upstream UTR	4
5UTR-003	Upstream UTR	5
5UTR-004	Upstream UTR	6
5UTR-005	Upstream UTR	7
5UTR-006	Upstream UTR	8
5UTR-007	Upstream UTR	9
5UTR-008	Upstream UTR	10
5UTR-009	Upstream UTR	11
5UTR-010	Upstream UTR	12
5UTR-011	Upstream UTR	13
5UTR-012	Upstream UTR	14
5UTR-013	Upstream UTR	15
5UTR-014	Upstream UTR	16
5UTR-015	Upstream UTR	17
5UTR-016	Upstream UTR	18
5UTR-017	Upstream UTR	19

[000187] Shown in Table 2 is a listing of 3'-untranslated regions of the invention. Variants of 3' UTRs may be utilized wherein one or more nucleotides are added or removed to the termini, including A, T, C or G.

Table 2. 3'-Untranslated Regions

3' UTR Identifier	Name/ Description	SEQ ID NO.
3UTR-001	Creatine Kinase	20
3UTR-002	Myoglobin	21
3UTR-003	α -actin	22
3UTR-004	Albumin	23
3UTR-005	α -globin	24
3UTR-006	G-CSF	25
3UTR-007	Colla2; collagen, type I, alpha 2	26
3UTR-008	Col6a2; collagen, type VI, alpha 2	27
3UTR-009	RPN1; ribophorin I	28
3UTR-010	LRP1; low density lipoprotein receptor-related protein 1	29
3UTR-011	Nnt1; cardiotrophin-like cytokine factor 1	30
3UTR-012	Col6a1; collagen, type VI, alpha 1	31
3UTR-013	Calr; calreticulin	32
3UTR-014	Colla1; collagen, type I, alpha 1	33
3UTR-015	Plod1; procollagen-lysine, 2-oxoglutarate 5-dioxygenase 1	34
3UTR-016	Nucb1; nucleobindin 1	35
3UTR-017	α -globin	36

5' UTR and Translation Initiation

[000188] Natural 5'UTRs bear features which play roles in translation initiation. They harbor signatures like Kozak sequences which are commonly known to be involved in the process by which the ribosome initiates translation of many genes. Kozak sequences have the consensus CCR(A/G)CCAUGG, where R is a purine (adenine or guanine) three bases upstream of the start codon (AUG), which is followed by another 'G'. 5'UTR also have been known to form secondary structures which are involved in elongation factor binding.

[000189] By engineering the features typically found in abundantly expressed genes of specific target organs, one can enhance the stability and protein production of the polynucleotides of the invention. For example, introduction of 5' UTR of liver-expressed mRNA, such as albumin, serum amyloid A, Apolipoprotein A/B/E, transferrin, alpha fetoprotein, erythropoietin, or Factor VIII, could be used to enhance expression of a nucleic acid molecule, such as a polynucleotides, in hepatic cell lines or liver. Likewise, use of 5' UTR from other tissue-specific mRNA to improve expression in that tissue is possible for muscle (MyoD, Myosin, Myoglobin, Myogenin, Herculin), for endothelial cells (Tie-1, CD36), for myeloid cells (C/EBP, AML1, G-CSF, GM-CSF, CD11b, MSR, Fr-1, i-NOS), for leukocytes (CD45, CD18), for adipose tissue (CD36, GLUT4, ACRP30, adiponectin) and for lung epithelial cells (SP-A/B/C/D). Untranslated regions useful in the design and manufacture of polynucleotides include, but are not limited, to those disclosed in co-pending, co-owned International Patent Publication No. WO2014164253 (Attorney Docket Number M42.20), the contents of which are incorporated herein by reference in its entirety.

[000190] Other non-UTR sequences may also be used as regions or subregions within the polynucleotides. For example, introns or portions of introns sequences may be incorporated into regions of the polynucleotides of the invention. Incorporation of intronic sequences may increase protein production as well as polynucleotide levels.

[000191] Combinations of features may be included in flanking regions and may be contained within other features. For example, the ORF may be flanked by a 5' UTR which may contain a strong Kozak translational initiation signal and/or a 3' UTR which

may include an oligo(dT) sequence for templated addition of a poly-A tail. 5'UTR may comprise a first polynucleotide fragment and a second polynucleotide fragment from the same and/or different genes such as the 5'UTRs described in US Patent Application Publication No. 20100293625, herein incorporated by reference in its entirety.

[000192] Co-pending, co-owned International Patent Publication No. WO2014164253 (Attorney Docket Number M42.20) provides a listing of exemplary UTRs which may be utilized in the polynucleotide of the present invention as flanking regions. Variants of 5' or 3' UTRs may be utilized wherein one or more nucleotides are added or removed to the termini, including A, T, C or G.

[000193] It should be understood that any UTR from any gene may be incorporated into the regions of the polynucleotide. Furthermore, multiple wild-type UTRs of any known gene may be utilized. It is also within the scope of the present invention to provide artificial UTRs which are not variants of wild type regions. These UTRs or portions thereof may be placed in the same orientation as in the transcript from which they were selected or may be altered in orientation or location. Hence a 5' or 3' UTR may be inverted, shortened, lengthened, made with one or more other 5' UTRs or 3' UTRs. As used herein, the term "altered" as it relates to a UTR sequence, means that the UTR has been changed in some way in relation to a reference sequence. For example, a 3' or 5' UTR may be altered relative to a wild type or native UTR by the change in orientation or location as taught above or may be altered by the inclusion of additional nucleotides, deletion of nucleotides, swapping or transposition of nucleotides. Any of these changes producing an "altered" UTR (whether 3' or 5') comprise a variant UTR.

[000194] In one embodiment, a double, triple or quadruple UTR such as a 5' or 3' UTR may be used. As used herein, a "double" UTR is one in which two copies of the same UTR are encoded either in series or substantially in series. For example, a double beta-globin 3' UTR may be used as described in US Patent publication 20100129877, the contents of which are incorporated herein by reference in its entirety.

[000195] It is also within the scope of the present invention to have patterned UTRs. As used herein "patterned UTRs" are those UTRs which reflect a repeating or alternating pattern, such as ABABAB or AABBAABBAABB or ABCABCABC or variants thereof

repeated once, twice, or more than 3 times. In these patterns, each letter, A, B, or C represent a different UTR at the nucleotide level.

[000196] In one embodiment, flanking regions are selected from a family of transcripts whose proteins share a common function, structure, feature of property. For example, polypeptides of interest may belong to a family of proteins which are expressed in a particular cell, tissue or at some time during development. The UTRs from any of these genes may be swapped for any other UTR of the same or different family of proteins to create a new polynucleotide. As used herein, a “family of proteins” is used in the broadest sense to refer to a group of two or more polypeptides of interest which share at least one function, structure, feature, localization, origin, or expression pattern.

[000197] The untranslated region may also include translation enhancer elements (TEE). As a non-limiting example, the TEE may include those described in US Application No. 20090226470, herein incorporated by reference in its entirety, and those known in the art.

3' UTR and the AU Rich Elements

[000198] Natural or wild type 3' UTRs are known to have stretches of Adenosines and Uridines embedded in them. These AU rich signatures are particularly prevalent in genes with high rates of turnover. Based on their sequence features and functional properties, the AU rich elements (AREs) can be separated into three classes (Chen et al, 1995): Class I AREs contain several dispersed copies of an AUUUA motif within U-rich regions. C-Myc and MyoD contain class I AREs. Class II AREs possess two or more overlapping UUAUUUA(U/A)(U/A) nonamers. Molecules containing this type of AREs include GM-CSF and TNF- α . Class III AREs are less well defined. These U rich regions do not contain an AUUUA motif. c-Jun and Myogenin are two well-studied examples of this class. Most proteins binding to the AREs are known to destabilize the messenger, whereas members of the ELAV family, most notably HuR, have been documented to increase the stability of mRNA. HuR binds to AREs of all the three classes. Engineering the HuR specific binding sites into the 3' UTR of nucleic acid molecules will lead to HuR binding and thus, stabilization of the message *in vivo*.

[000199] Introduction, removal or modification of 3' UTR AU rich elements (AREs) can be used to modulate the stability of polynucleotides of the invention. When

engineering specific polynucleotides, one or more copies of an ARE can be introduced to make polynucleotides of the invention less stable and thereby curtail translation and decrease production of the resultant protein. Likewise, AREs can be identified and removed or mutated to increase the intracellular stability and thus increase translation and production of the resultant protein. Transfection experiments can be conducted in relevant cell lines, using polynucleotides of the invention and protein production can be assayed at various time points post-transfection. For example, cells can be transfected with different ARE-engineering molecules and by using an ELISA kit to the relevant protein and assaying protein produced at 6 hour, 12 hour, 24 hour, 48 hour, and 7 days post-transfection.

microRNA Binding Sites

[000200] microRNAs (or miRNA) are 19-25 nucleotide long noncoding RNAs that bind to the 3'UTR of nucleic acid molecules and down-regulate gene expression either by reducing nucleic acid molecule stability or by inhibiting translation. The polynucleotides of the invention may comprise one or more microRNA target sequences, microRNA sequences, or microRNA seeds. Such sequences may correspond to any known microRNA such as those taught in US Publication US2005/0261218, US Publication US2005/0059005, US Patent Publication No. 20140200261 (Attorney Docket No. M037.10), US Patent Publication No. 20140147454 (Attorney Docket No. M039.11), International Patent Publication No. WO2014113089 (Attorney Docket No. M037.20), and International Patent Publication No. WO2014081507 (Attorney Docket No. M039.21), the contents of each of which are incorporated herein by reference in their entirety.

[000201] A microRNA sequence comprises a “seed” region, i.e., a sequence in the region of positions 2-8 of the mature microRNA, which sequence has perfect Watson-Crick complementarity to the miRNA target sequence. A microRNA seed may comprise positions 2-8 or 2-7 of the mature microRNA. In some embodiments, a microRNA seed may comprise 7 nucleotides (e.g., nucleotides 2-8 of the mature microRNA), wherein the seed-complementary site in the corresponding miRNA target is flanked by an adenine (A) opposed to microRNA position 1. In some embodiments, a microRNA seed may comprise 6 nucleotides (e.g., nucleotides 2-7 of the mature microRNA), wherein the

seed-complementary site in the corresponding miRNA target is flanked by an adenine (A) opposed to microRNA position 1. See for example, Grimson A, Farh KK, Johnston WK, Garrett-Engle P, Lim LP, Bartel DP; Mol Cell. 2007 Jul 6; 27(1):91-105; each of which is herein incorporated by reference in their entirety. The bases of the microRNA seed have complete complementarity with the target sequence. By engineering microRNA target sequences into the polynucleotides (e.g., in a 3'UTR like region or other region) of the invention one can target the molecule for degradation or reduced translation, provided the microRNA in question is available. This process will reduce the hazard of off target effects upon nucleic acid molecule delivery. Identification of microRNA, microRNA target regions, and their expression patterns and role in biology have been reported (Bonauer et al., Curr Drug Targets 2010 11:943-949; Anand and Cheresch Curr Opin Hematol 2011 18:171-176; Contreras and Rao Leukemia 2012 26:404-413 (2011 Dec 20. doi: 10.1038/leu.2011.356); Bartel Cell 2009 136:215-233; Landgraf et al, Cell, 2007 129:1401-1414; each of which is herein incorporated by reference in its entirety).

[000202] For example, if the nucleic acid molecule is an mRNA and is not intended to be delivered to the liver but ends up there, then miR-122, a microRNA abundant in liver, can inhibit the expression of the gene of interest if one or multiple target sites of miR-122 are engineered into the 3' UTR region of the polynucleotides. Introduction of one or multiple binding sites for different microRNA can be engineered to further decrease the longevity, stability, and protein translation of polynucleotides.

[000203] As used herein, the term "microRNA site" refers to a microRNA target site or a microRNA recognition site, or any nucleotide sequence to which a microRNA binds or associates. It should be understood that "binding" may follow traditional Watson-Crick hybridization rules or may reflect any stable association of the microRNA with the target sequence at or adjacent to the microRNA site.

[000204] Conversely, for the purposes of the polynucleotides of the present invention, microRNA binding sites can be engineered out of (i.e. removed from) sequences in which they occur, e.g., in order to increase protein expression in specific tissues. For example, miR-122 binding sites may be removed to improve protein expression in the liver. Regulation of expression in multiple tissues can be accomplished through introduction or removal of one or several microRNA binding sites.

[000205] Examples of tissues where microRNA are known to regulate mRNA, and thereby protein expression, include, but are not limited to, liver (miR-122), muscle (miR-133, miR-206, miR-208), endothelial cells (miR-17-92, miR-126), myeloid cells (miR-142-3p, miR-142-5p, miR-16, miR-21, miR-223, miR-24, miR-27), adipose tissue (let-7, miR-30c), heart (miR-1d, miR-149), kidney (miR-192, miR-194, miR-204), and lung epithelial cells (let-7, miR-133, miR-126). MicroRNA can also regulate complex biological processes such as angiogenesis (miR-132) (Anand and Cheresh Curr Opin Hematol 2011 18:171-176; herein incorporated by reference in its entirety).

[000206] Expression profiles, microRNA and cell lines useful in the present invention include those taught in, for example, International Patent Publication Nos. WO2014081507 (Attorney Docket Number M39) and WO2014113089 (Attorney Docket Number M37) each filed July 23, 2013, the contents of which are incorporated by reference in their entirety.

[000207] In the polynucleotides of the present invention, binding sites for microRNAs that are involved in such processes may be removed or introduced, in order to tailor the expression of the polynucleotides expression to biologically relevant cell types or to the context of relevant biological processes. A listing of microRNA, miR sequences and miR binding sites is listed in Table 9 of U.S. Provisional Application No. 61/753,661 filed January 17, 2013, in Table 9 of U.S. Provisional Application No. 61/754,159 filed January 18, 2013, and in Table 7 of U.S. Provisional Application No. 61/758,921 filed January 31, 2013, each of which are herein incorporated by reference in their entireties.

[000208] Examples of use of microRNA to drive tissue or disease-specific gene expression are listed (Getner and Naldini, Tissue Antigens. 2012, 80:393-403; herein incorporated by reference in its entirety). In addition, microRNA seed sites can be incorporated into mRNA to decrease expression in certain cells which results in a biological improvement. An example of this is incorporation of miR-142 sites into a UGT1A1-expressing lentiviral vector. The presence of miR-142 seed sites reduced expression in hematopoietic cells, and as a consequence reduced expression in antigen-presenting cells, leading to the absence of an immune response against the virally expressed UGT1A1 (Schmitt et al., Gastroenterology 2010; 139:999-1007; Gonzalez-Asequinolaza et al. Gastroenterology 2010, 139:726-729; both herein incorporated by

reference in its entirety) . Incorporation of miR-142 sites into modified mRNA could not only reduce expression of the encoded protein in hematopoietic cells, but could also reduce or abolish immune responses to the mRNA-encoded protein. Incorporation of miR-142 seed sites (one or multiple) into mRNA would be important in the case of treatment of patients with complete protein deficiencies (UGT1A1 type I, LDLR-deficient patients, CRIM-negative Pompe patients, etc.) .

[000209] Lastly, through an understanding of the expression patterns of microRNA in different cell types, polynucleotides can be engineered for more targeted expression in specific cell types or only under specific biological conditions. Through introduction of tissue-specific microRNA binding sites, polynucleotides could be designed that would be optimal for protein expression in a tissue or in the context of a biological condition.

[000210] Transfection experiments can be conducted in relevant cell lines, using engineered polynucleotides and protein production can be assayed at various time points post-transfection. For example, cells can be transfected with different microRNA binding site-engineering polynucleotides and by using an ELISA kit to the relevant protein and assaying protein produced at 6 hour, 12 hour, 24 hour, 48 hour, 72 hour and 7 days post-transfection. *In vivo* experiments can also be conducted using microRNA-binding site-engineered molecules to examine changes in tissue-specific expression of formulated polynucleotides.

[000211] In one embodiment, a polynucleotide encoding a tolerogenic polypeptide of interest may comprise at least one terminal modification such as, but not limited to, the terminal modifications described in US Patent Publication No. 20140147454 (Attorney Docket No. M039.11) and International Patent Publication No. WO2014081507 (Attorney Docket No. M039.21), the contents of each of which are incorporated herein by reference in their entirety. As a non-limiting example, the terminal modification may be at least one miR-122 sequence or fragment such as at least one miR-122 binding site in the 3'UTR of the polynucleotide. The polynucleotide may be formulated in any of the formulations described herein such as, but not limited to, a lipid nanoparticle to target translation of the polynucleotide in myeloid cells. The size of a lipid nanoparticle comprising the polynucleotide may be optimized to specifically target APCs or DCs

and/or the nanoparticles may comprise at least one targeting epitope to target APCs or DCs.

Regions having a 5' Cap

[000212] The 5' cap structure of a natural mRNA is involved in nuclear export, increasing mRNA stability and binds the mRNA Cap Binding Protein (CBP), which is responsible for mRNA stability in the cell and translation competency through the association of CBP with poly(A) binding protein to form the mature cyclic mRNA species. The cap further assists the removal of 5' proximal introns removal during mRNA splicing.

[000213] Endogenous mRNA molecules may be 5'-end capped generating a 5'-ppp-5'-triphosphate linkage between a terminal guanosine cap residue and the 5'-terminal transcribed sense nucleotide of the mRNA molecule. This 5'-guanylate cap may then be methylated to generate an N7-methyl-guanylate residue. The ribose sugars of the terminal and/or antiterminal transcribed nucleotides of the 5' end of the mRNA may optionally also be 2'-O-methylated. 5'-decapping through hydrolysis and cleavage of the guanylate cap structure may target a nucleic acid molecule, such as an mRNA molecule, for degradation.

[000214] In some embodiments, polynucleotides may be designed to incorporate a cap moiety. Modifications to the polynucleotides of the present invention may generate a non-hydrolyzable cap structure preventing decapping and thus increasing mRNA half-life. Because cap structure hydrolysis requires cleavage of 5'-ppp-5' phosphodiester linkages, modified nucleotides may be used during the capping reaction. For example, a Vaccinia Capping Enzyme from New England Biolabs (Ipswich, MA) may be used with α -thio-guanosine nucleotides according to the manufacturer's instructions to create a phosphorothioate linkage in the 5'-ppp-5' cap. Additional modified guanosine nucleotides may be used such as α -methyl-phosphonate and seleno-phosphate nucleotides.

[000215] Additional modifications include, but are not limited to, 2'-O-methylation of the ribose sugars of 5'-terminal and/or 5'-antiterminal nucleotides of the polynucleotide (as mentioned above) on the 2'-hydroxyl group of the sugar ring. Multiple distinct 5'-cap

structures can be used to generate the 5'-cap of a nucleic acid molecule, such as a polynucleotide which functions as an mRNA molecule.

[000216] Cap analogs, which herein are also referred to as synthetic cap analogs, chemical caps, chemical cap analogs, or structural or functional cap analogs, differ from natural (i.e. endogenous, wild-type or physiological) 5'-caps in their chemical structure, while retaining cap function. Cap analogs may be chemically (i.e. non-enzymatically) or enzymatically synthesized and/or linked to the polynucleotides of the invention.

[000217] For example, the Anti-Reverse Cap Analog (ARCA) cap contains two guanines linked by a 5'-5'-triphosphate group, wherein one guanine contains an N7 methyl group as well as a 3'-O-methyl group (i.e., N7,3'-O-dimethyl-guanosine-5'-triphosphate-5'-guanosine (m^7G -3'mppp-G; which may equivalently be designated 3' O-Me- $m^7G(5')$ ppp(5')G). The 3'-O atom of the other, unmodified, guanine becomes linked to the 5'-terminal nucleotide of the capped polynucleotide. The N7- and 3'-O-methylated guanine provides the terminal moiety of the capped polynucleotide.

[000218] Another exemplary cap is mCAP, which is similar to ARCA but has a 2'-O-methyl group on guanosine (i.e., N7,2'-O-dimethyl-guanosine-5'-triphosphate-5'-guanosine, m^7Gm -ppp-G).

[000219] In one embodiment, the cap is a dinucleotide cap analog. As a non-limiting example, the dinucleotide cap analog may be modified at different phosphate positions with a boranophosphate group or a phosphoroselenoate group such as the dinucleotide cap analogs described in US Patent No. US 8,519,110, the contents of which are herein incorporated by reference in its entirety.

[000220] In another embodiment, the cap is a cap analog is a N7-(4-chlorophenoxyethyl) substituted dinucleotide form of a cap analog known in the art and/or described herein. Non-limiting examples of a N7-(4-chlorophenoxyethyl) substituted dinucleotide form of a cap analog include a N7-(4-chlorophenoxyethyl)-G(5')ppp(5')G and a N7-(4-chlorophenoxyethyl)- $m^{3'-O}G(5')$ ppp(5')G cap analog (See e.g., the various cap analogs and the methods of synthesizing cap analogs described in Kore et al. Bioorganic & Medicinal Chemistry 2013 21:4570-4574; the contents of which are herein incorporated by reference in its entirety). In another embodiment, a cap analog of the present invention is a 4-chloro/bromophenoxyethyl analog.

[000221] While cap analogs allow for the concomitant capping of a polynucleotide or a region thereof, in an in vitro transcription reaction, up to 20% of transcripts can remain uncapped. This, as well as the structural differences of a cap analog from an endogenous 5'-cap structures of nucleic acids produced by the endogenous, cellular transcription machinery, may lead to reduced translational competency and reduced cellular stability.

[000222] Polynucleotides of the invention may also be capped post-manufacture (whether IVT or chemical synthesis), using enzymes, in order to generate more authentic 5'-cap structures. As used herein, the phrase “more authentic” refers to a feature that closely mirrors or mimics, either structurally or functionally, an endogenous or wild type feature. That is, a “more authentic” feature is better representative of an endogenous, wild-type, natural or physiological cellular function and/or structure as compared to synthetic features or analogs, etc., of the prior art, or which outperforms the corresponding endogenous, wild-type, natural or physiological feature in one or more respects. Non-limiting examples of more authentic 5'cap structures of the present invention are those which, among other things, have enhanced binding of cap binding proteins, increased half-life, reduced susceptibility to 5' endonucleases and/or reduced 5'decapping, as compared to synthetic 5'cap structures known in the art (or to a wild-type, natural or physiological 5'cap structure). For example, recombinant Vaccinia Virus Capping Enzyme and recombinant 2'-O-methyltransferase enzyme can create a canonical 5'-5'-triphosphate linkage between the 5'-terminal nucleotide of a polynucleotide and a guanine cap nucleotide wherein the cap guanine contains an N7 methylation and the 5'-terminal nucleotide of the mRNA contains a 2'-O-methyl. Such a structure is termed the Cap1 structure. This cap results in a higher translational-competency and cellular stability and a reduced activation of cellular pro-inflammatory cytokines, as compared, e.g., to other 5'cap analog structures known in the art. Cap structures include, but are not limited to, 7mG(5')ppp(5')N,pN2p (cap 0), 7mG(5')ppp(5')NlmpNp (cap 1), and 7mG(5')-ppp(5')NlmpN2mp (cap 2).

[000223] As a non-limiting example, capping chimeric polynucleotides post-manufacture may be more efficient as nearly 100% of the chimeric polynucleotides may be capped. This is in contrast to ~80% when a cap analog is linked to a chimeric polynucleotide in the course of an in vitro transcription reaction.

[000224] According to the present invention, 5' terminal caps may include endogenous caps or cap analogs. According to the present invention, a 5' terminal cap may comprise a guanine analog. Useful guanine analogs include, but are not limited to, inosine, N1-methyl-guanosine, 2'fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, and 2-azido-guanosine.

Viral Sequences

[000225] Additional viral sequences such as, but not limited to, the translation enhancer sequence of the barley yellow dwarf virus (BYDV-PAV), the Jaagsiekte sheep retrovirus (JSRV) and/or the Enzootic nasal tumor virus (See e.g., International Pub. No. WO2012129648; herein incorporated by reference in its entirety) can be engineered and inserted in the polynucleotides of the invention and can stimulate the translation of the construct *in vitro* and *in vivo*. Transfection experiments can be conducted in relevant cell lines at and protein production can be assayed by ELISA at 12hr, 24hr, 48hr, 72 hr and day 7 post-transfection.

IRES Sequences

[000226] Further, provided are polynucleotides which may contain an internal ribosome entry site (IRES). First identified as a feature Picorna virus RNA, IRES plays an important role in initiating protein synthesis in absence of the 5' cap structure. An IRES may act as the sole ribosome binding site, or may serve as one of multiple ribosome binding sites of an mRNA. Polynucleotides containing more than one functional ribosome binding site may encode several peptides or polypeptides that are translated independently by the ribosomes ("multicistronic nucleic acid molecules"). When polynucleotides are provided with an IRES, further optionally provided is a second translatable region. Examples of IRES sequences that can be used according to the invention include without limitation, those from picornaviruses (e.g. FMDV), pest viruses (CFFV), polio viruses (PV), encephalomyocarditis viruses (ECMV), foot-and-mouth disease viruses (FMDV), hepatitis C viruses (HCV), classical swine fever viruses (CSFV), murine leukemia virus (MLV), simian immune deficiency viruses (SIV) or cricket paralysis viruses (CrPV).

Poly-A tails

[000227] During RNA processing, a long chain of adenine nucleotides (poly-A tail) may be added to a polynucleotide such as an mRNA molecule in order to increase stability. Immediately after transcription, the 3' end of the transcript may be cleaved to free a 3' hydroxyl. Then poly-A polymerase adds a chain of adenine nucleotides to the RNA. The process, called polyadenylation, adds a poly-A tail that can be between, for example, approximately 80 to approximately 250 residues long, including approximately 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240 or 250 residues long.

[000228] PolyA tails may also be added after the construct is exported from the nucleus.

[000229] According to the present invention, terminal groups on the poly A tail may be incorporated for stabilization. Polynucleotides of the present invention may include des-3' hydroxyl tails. They may also include structural moieties or 2'-O-methyl modifications as taught by Junjie Li, et al. (Current Biology, Vol. 15, 1501–1507, August 23, 2005, the contents of which are incorporated herein by reference in its entirety).

[000230] The polynucleotides of the present invention may be designed to encode transcripts with alternative polyA tail structures including histone mRNA. According to Norbury, "Terminal uridylation has also been detected on human replication-dependent histone mRNAs. The turnover of these mRNAs is thought to be important for the prevention of potentially toxic histone accumulation following the completion or inhibition of chromosomal DNA replication. These mRNAs are distinguished by their lack of a 3' poly(A) tail, the function of which is instead assumed by a stable stem-loop structure and its cognate stem-loop binding protein (SLBP); the latter carries out the same functions as those of PABP on polyadenylated mRNAs" (Norbury, "Cytoplasmic RNA: a case of the tail wagging the dog," Nature Reviews Molecular Cell Biology; AOP, published online 29 August 2013; doi:10.1038/nrm3645) the contents of which are incorporated herein by reference in its entirety.

[000231] Unique poly-A tail lengths provide certain advantages to the polynucleotides of the present invention.

[000232] Generally, the length of a poly-A tail, when present, is greater than 30 nucleotides in length. In another embodiment, the poly-A tail is greater than 35 nucleotides in length (e.g., at least or greater than about 35, 40, 45, 50, 55, 60, 70, 80, 90,

100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,900, 2,000, 2,500, and 3,000 nucleotides). In some embodiments, the polynucleotide or region thereof includes from about 30 to about 3,000 nucleotides (e.g., from 30 to 50, from 30 to 100, from 30 to 250, from 30 to 500, from 30 to 750, from 30 to 1,000, from 30 to 1,500, from 30 to 2,000, from 30 to 2,500, from 50 to 100, from 50 to 250, from 50 to 500, from 50 to 750, from 50 to 1,000, from 50 to 1,500, from 50 to 2,000, from 50 to 2,500, from 50 to 3,000, from 100 to 500, from 100 to 750, from 100 to 1,000, from 100 to 1,500, from 100 to 2,000, from 100 to 2,500, from 100 to 3,000, from 500 to 750, from 500 to 1,000, from 500 to 1,500, from 500 to 2,000, from 500 to 2,500, from 500 to 3,000, from 1,000 to 1,500, from 1,000 to 2,000, from 1,000 to 2,500, from 1,000 to 3,000, from 1,500 to 2,000, from 1,500 to 2,500, from 1,500 to 3,000, from 2,000 to 3,000, from 2,000 to 2,500, and from 2,500 to 3,000).

[000233] In one embodiment, the poly-A tail is designed relative to the length of the overall polynucleotide or the length of a particular region of the polynucleotide. This design may be based on the length of a coding region, the length of a particular feature or region or based on the length of the ultimate product expressed from the polynucleotides.

[000234] In this context the poly-A tail may be 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100% greater in length than the polynucleotide or feature thereof. The poly-A tail may also be designed as a fraction of the polynucleotides to which it belongs. In this context, the poly-A tail may be 10, 20, 30, 40, 50, 60, 70, 80, or 90% or more of the total length of the construct, a construct region or the total length of the construct minus the poly-A tail. Further, engineered binding sites and conjugation of polynucleotides for Poly-A binding protein may enhance expression.

[000235] Additionally, multiple distinct polynucleotides may be linked together via the PABP (Poly-A binding protein) through the 3'-end using modified nucleotides at the 3'-terminus of the poly-A tail. Transfection experiments can be conducted in relevant cell lines at and protein production can be assayed by ELISA at 12hr, 24hr, 48hr, 72 hr and day 7 post-transfection.

[000236] In one embodiment, the polynucleotides of the present invention are designed to include a polyA-G Quartet region. The G-quartet is a cyclic hydrogen bonded array of

four guanine nucleotides that can be formed by G-rich sequences in both DNA and RNA. In this embodiment, the G-quartet is incorporated at the end of the poly-A tail. The resultant polynucleotide is assayed for stability, protein production and other parameters including half-life at various time points. It has been discovered that the polyA-G quartet results in protein production from an mRNA equivalent to at least 75% of that seen using a poly-A tail of 120 nucleotides alone.

Start codon region

[000237] In some embodiments, the polynucleotides of the present invention may have regions that are analogous to or function like a start codon region.

[000238] In one embodiment, the translation of a polynucleotide may initiate on a codon which is not the start codon AUG. Translation of the polynucleotide may initiate on an alternative start codon such as, but not limited to, ACG, AGG, AAG, CTG/CUG, GTG/GUG, ATA/AUA, ATT/AUU, TTG/UUG (see Touriol et al. *Biology of the Cell* 95 (2003) 169-178 and Matsuda and Mauro *PLoS ONE*, 2010 5:11; the contents of each of which are herein incorporated by reference in its entirety). As a non-limiting example, the translation of a polynucleotide begins on the alternative start codon ACG. As another non-limiting example, polynucleotide translation begins on the alternative start codon CTG or CUG. As yet another non-limiting example, the translation of a polynucleotide begins on the alternative start codon GTG or GUG.

[000239] Nucleotides flanking a codon that initiates translation such as, but not limited to, a start codon or an alternative start codon, are known to affect the translation efficiency, the length and/or the structure of the polynucleotide. (See e.g., Matsuda and Mauro *PLoS ONE*, 2010 5:11; the contents of which are herein incorporated by reference in its entirety). Masking any of the nucleotides flanking a codon that initiates translation may be used to alter the position of translation initiation, translation efficiency, length and/or structure of a polynucleotide.

[000240] In one embodiment, a masking agent may be used near the start codon or alternative start codon in order to mask or hide the codon to reduce the probability of translation initiation at the masked start codon or alternative start codon. Non-limiting examples of masking agents include antisense locked nucleic acids (LNA) polynucleotides and exon-junction complexes (EJCs) (See e.g., Matsuda and Mauro

describing masking agents LNA polynucleotides and EJC (PLoS ONE, 2010 5:11); the contents of which are herein incorporated by reference in its entirety).

[000241] In another embodiment, a masking agent may be used to mask a start codon of a polynucleotide in order to increase the likelihood that translation will initiate on an alternative start codon.

[000242] In one embodiment, a masking agent may be used to mask a first start codon or alternative start codon in order to increase the chance that translation will initiate on a start codon or alternative start codon downstream to the masked start codon or alternative start codon.

[000243] In one embodiment, a start codon or alternative start codon may be located within a perfect complement for a miR binding site. The perfect complement of a miR binding site may help control the translation, length and/or structure of the polynucleotide similar to a masking agent. As a non-limiting example, the start codon or alternative start codon may be located in the middle of a perfect complement for a miR-122 binding site. The start codon or alternative start codon may be located after the first nucleotide, second nucleotide, third nucleotide, fourth nucleotide, fifth nucleotide, sixth nucleotide, seventh nucleotide, eighth nucleotide, ninth nucleotide, tenth nucleotide, eleventh nucleotide, twelfth nucleotide, thirteenth nucleotide, fourteenth nucleotide, fifteenth nucleotide, sixteenth nucleotide, seventeenth nucleotide, eighteenth nucleotide, nineteenth nucleotide, twentieth nucleotide or twenty-first nucleotide.

[000244] In another embodiment, the start codon of a polynucleotide may be removed from the polynucleotide sequence in order to have the translation of the polynucleotide begin on a codon which is not the start codon. Translation of the polynucleotide may begin on the codon following the removed start codon or on a downstream start codon or an alternative start codon. In a non-limiting example, the start codon ATG or AUG is removed as the first 3 nucleotides of the polynucleotide sequence in order to have translation initiate on a downstream start codon or alternative start codon. The polynucleotide sequence where the start codon was removed may further comprise at least one masking agent for the downstream start codon and/or alternative start codons in order to control or attempt to control the initiation of translation, the length of the polynucleotide and/or the structure of the polynucleotide.

Stop Codon Region

[000245] In one embodiment, the polynucleotides of the present invention may include at least two stop codons before the 3' untranslated region (UTR). The stop codon may be selected from TGA, TAA and TAG. In one embodiment, the polynucleotides of the present invention include the stop codon TGA and one additional stop codon. In a further embodiment the additional stop codon may be TAA. In another embodiment, the polynucleotides of the present invention include three stop codons.

Signal Sequences

[000246] The polynucleotides may also encode additional features which facilitate trafficking of the polypeptides to therapeutically relevant sites. One such feature which aids in protein trafficking is the signal sequence. As used herein, a "signal sequence" or "signal peptide" is a polynucleotide or polypeptide, respectively, which is from about 9 to 200 nucleotides (3-60 amino acids) in length which is incorporated at the 5' (or N-terminus) of the coding region or polypeptide encoded, respectively. Addition of these sequences result in trafficking of the encoded polypeptide to the endoplasmic reticulum through one or more secretory pathways. Some signal peptides are cleaved from the protein by signal peptidase after the proteins are transported.

[000247] Additional signal sequences which may be utilized in the present invention include those taught in, for example, databases such as those found at <http://www.signalpeptide.de/> or <http://proline.bic.nus.edu.sg/spdb/>. Those described in US Patents 8,124,379; 7,413,875 and 7,385,034 are also within the scope of the invention and the contents of each are incorporated herein by reference in their entirety.

Tolerogenic Molecules encoded by polynucleotides

[000248] According to the present invention, the polynucleotides may encode at least one polypeptide of interest. Certain tolerogenic molecules of the present invention are listed in Table 3. Shown in Table 3, in addition to the name and description of the gene encoding the polypeptide of interest, where applicable, (Tolerogenic Molecule Description) are the ENSEMBL Transcript ID (ENST), the ENSEMBL Protein ID (ENSP) and the optimized open reading frame sequence ID (Optimized ORF SEQ ID).

[000249] For any particular gene there may exist one or more variants or isoforms. Non-limiting examples of these variants or isoforms, if known, are shown in the table as

well. It will be appreciated by those of skill in the art that disclosed in the Table are potential flanking regions. These are encoded in each ENST transcript either to the 5' (upstream) or 3' (downstream) of the ORF or coding region. The coding region is definitively and specifically disclosed by teaching the ENSP and/or the Protein Sequence. Consequently, the sequences taught flanking that encode the protein are considered regions that flank the ORF or coding region. It is also possible to further characterize the 5' and 3' regions that flank the ORF or coding region by utilizing one or more available databases or algorithms. Databases have annotated the features contained in the regions that flank the ORF or coding region of the ENST transcripts and these are available in the art. Also described in Table 3, if known are optimized transcript and/or ORF sequences. In Table 3, "N/A" means not applicable.

Table 3. Tolerogenic Molecules

No.	Tolerogenic Molecule Description	ENSP	Protein SEQ ID NO	ENST	Trans SEQ ID NO	Optimized Trans SEQ ID NO	Optimized ORF SEQ ID NO
1	CD274 molecule	370985	37	381573	172		220, 259, 298, 337, 376
2	CD274 molecule	370989	38	381577	173		221, 260, 299, 338, 377
3	CD40 molecule, TNF receptor superfamily member 5	361350	39	372276	174		222, 261, 300, 339, 378
4	CD40 molecule, TNF receptor superfamily member 5	361352	40	372278	175		223, 262, 301, 340, 379
5	CD40 molecule, TNF receptor superfamily member 5	361359	41	372285	176		224, 263, 302, 341, 380
6	CD40 ligand	359662	42	370628	177		225, 264, 303, 342, 381
7	CD40 ligand	359663	43	370629	178		226, 265, 304, 343, 382
8	Anti-CD40L domain antibody (dAb)	N/A	44				
9	Mutated IgG-4 chain C region	N/A	45				
10	Membrane bound IgM CH4 of IgM	N/A	46				
11	Membrane bound IgG-4 chain with a VK signal	N/A	47				

	peptide, a FLAG tag, an anti-CD40L dAb and an immunoglobulin mu chain						
12	Secreted IgG-4 chain with a VK signal peptide, a FLAG tag and an anti-CD40L dAb	N/A	48				
13	lymphocyte antigen 75	263636	49	263636	179		227, 266, 305, 344, 383
14	lymphocyte antigen 75	451446	50	553424	180		228, 267, 306, 345, 384
15	lymphocyte antigen 75	451511	51	554112	181		229, 268, 307, 346, 385
16	CD52 molecule	363330	52	374213	182		230, 269, 308, 347, 386
17	fms-related tyrosine kinase 3 ligand	204637	53	204637	183		231, 270, 309, 348, 387, 415, 420
18	fms-related tyrosine kinase 3 ligand	341305	54	344019	184		232, 271, 310, 349, 388, 416, 421
19	fms-related tyrosine kinase 3 ligand	469613	55	594009	185		233, 272, 311, 350, 389
20	lectin, galactoside-binding, soluble, 1	215909	56	215909	186		234, 273, 312, 351, 390
21	leucine rich repeat containing 32	260061	57	260061	187		235, 274, 313, 352, 391
22	leucine rich repeat containing 32	385766	58	404995	188		236, 275, 314, 353, 392
23	leucine rich repeat containing 32	384126	59	407242	189		237, 276, 315, 354, 393
24	leucine rich repeat containing 32	413331	60	421973	190		238, 277, 316, 355, 394
25	indoleamine 2,3-dioxygenase 1	430950	61	518237	191		239, 278, 317, 356, 395
26	indoleamine 2,3-dioxygenase 1	429297	62	518804	192		240, 279, 318, 357, 396
27	indoleamine 2,3-dioxygenase 1	428716	63	519154	193		241, 280, 319, 358, 397

28	indoleamine 2,3-dioxygenase 1	430505	64	522495	194		242, 281, 320, 359, 398
29	indoleamine 2,3-dioxygenase 1	429933	65	522840	195		243, 282, 321, 360, 399
30	indoleamine 2,3-dioxygenase 2	426447	66	389060	196		244, 283, 322, 361, 400
31	indoleamine 2,3-dioxygenase 2	443432	67	502986	197		245, 284, 323, 362, 401
32	interleukin 10	412237	68	423557	198		213, 246, 285, 324, 363, 402, 425-511
33	interleukin 10					216	
34	interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35)	303231	69	305579	199		247, 286, 325, 364, 403, 417, 422
35	interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35)	419046	70	466512	200		248, 287, 326, 365, 404, 418, 423
36	interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35)	420184	71	480787	201		249, 288, 327, 366, 405, 419, 424
37	interleukin 12A	N/A	72				
38	Epstein-Barr virus induced 3	221847	73	221847	202		250, 289, 328, 367, 406
39	Epstein-Barr virus induced 3	N/A	74				
40	IL12-alpha, 2A peptide, EBI3 construct with a sequence of 6 histidines	N/A	75				
41	IL12-alpha, GS linker, EBI3 construct with a FLAG tag	N/A	76				
42	EBI3, GS linker, IL12-alpha and an Fc region	N/A	77				
43	interleukin 37	263326	78	263326	203		251, 290, 329, 368, 407
44	interleukin 37	309883	79	311328	204		252, 291, 330, 369, 408

45	interleukin 37	263328	80	349806	205		253, 292, 331, 370, 409
46	interleukin 37	263327	81	352179	206		254, 293, 332, 371, 410
47	interleukin 37	309208	82	353225	207		255, 294, 333, 372, 411
48	leukocyte immunoglobulin-like receptor subfamily B (with TM and ITIM domains) member 4	408995	83	430952	208		256, 295, 334, 373, 412
49	leukocyte immunoglobulin-like receptor subfamily B (with TM and ITIM domains) member 4	N/A	84	N/A	209		
50	Extracellular ILT3	N/A	85				
51	Extracellular ILT3-IgG1 Hinge - IgG1-Fc	N/A	86				
52	Extracellular ILT3-GS region-IgG4 CH2 CH3	N/A	87				
53	suppressor of cytokine signaling 1	329418	88	332029	210		257, 296, 335, 374, 413
54	suppressor of cytokine signaling 1			N/A	211		
55	suppressor of cytokine signaling 1						214
56	suppressor of cytokine signaling 1 cDNA					217	
57	suppressor of cytokine signaling 1						215
58	suppressor of cytokine signaling 1 with mir-122 in 3'UTR cDNA					218	
59	TCR inhibitory peptide	N/A	89				
60	TCR inhibitory peptide	N/A	90				
61	TCR inhibitory peptide	N/A	91				
62	TCR inhibitory peptide	N/A	92				
63	TCR inhibitory peptide	N/A	93				
64	TCR inhibitory peptide	N/A	94				
65	TCR inhibitory peptide	N/A	95				
66	TCR inhibitory peptide	N/A	96				
67	TCR inhibitory peptide	N/A	97				
68	TCR inhibitory peptide	N/A	98				
69	TCR inhibitory peptide	N/A	99				
70	TCR inhibitory peptide	N/A	100				

71	TCR inhibitory peptide	N/A	101				
72	TCR inhibitory peptide	N/A	102				
73	TCR inhibitory peptide	N/A	103				
74	TCR inhibitory peptide	N/A	104				
75	TCR inhibitory peptide	N/A	105				
76	TCR inhibitory peptide	N/A	106				
77	TCR inhibitory peptide	N/A	107				
78	TCR inhibitory peptide	N/A	108				
79	TCR inhibitory peptide	N/A	109				
80	TCR inhibitory peptide	N/A	110				
81	TCR inhibitory peptide	N/A	111				
82	TCR inhibitory peptide	N/A	112				
83	TCR inhibitory peptide	N/A	113				
84	TCR inhibitory peptide	N/A	114				
85	TCR inhibitory peptide	N/A	115				
86	TCR inhibitory peptide	N/A	116				
87	TCR inhibitory peptide	N/A	117				
88	TCR inhibitory peptide	N/A	118				
89	TCR inhibitory peptide	N/A	119				
90	TCR inhibitory peptide	N/A	120				
91	TCR inhibitory peptide	N/A	121				
92	TCR inhibitory peptide	N/A	122				
93	TCR inhibitory peptide	N/A	123				
94	TCR inhibitory peptide	N/A	124				
95	TCR inhibitory peptide	N/A	125				
96	TCR inhibitory peptide	N/A	126				
97	TCR inhibitory peptide	N/A	127				
98	TCR inhibitory peptide	N/A	128				
99	TCR inhibitory peptide	N/A	129				
100	TCR inhibitory peptide	N/A	130				
101	TCR inhibitory peptide	N/A	131				
102	TCR inhibitory peptide	N/A	132				
103	TCR inhibitory peptide	N/A	133				
104	TCR inhibitory peptide	N/A	134				
105	TCR inhibitory peptide	N/A	135				
106	TCR inhibitory peptide	N/A	136				
107	TCR inhibitory peptide	N/A	137				
108	TCR inhibitory peptide	N/A	138				
109	TCR inhibitory peptide	N/A	139				
110	TCR inhibitory peptide	N/A	140				
111	TCR inhibitory peptide	N/A	141				

112	TCR inhibitory peptide	N/A	142				
113	TCR inhibitory peptide	N/A	143				
114	TCR inhibitory peptide	N/A	144				
115	TCR inhibitory peptide	N/A	145				
116	TCR inhibitory peptide	N/A	146				
117	TCR inhibitory peptide	N/A	147				
118	TCR inhibitory peptide	N/A	148				
119	TCR inhibitory peptide	N/A	149				
120	TCR inhibitory peptide	N/A	150				
121	TCR inhibitory peptide	N/A	151				
122	TCR inhibitory peptide	N/A	152				
123	TCR inhibitory peptide	N/A	153				
124	TCR inhibitory peptide	N/A	154				
125	transforming growth factor, beta 1	221930	155	221930	212	219	258, 297, 336, 375, 414, 512-598

[000250] In one embodiment, polynucleotides of the present invention encode PD-L1. Wu *et al.* (2013, *Cellular & Molecular Immunity*; 10; 393-402) have suggested that IL-10 and TGF-beta are needed to maintain dendritic cells in tolerogenic state and that PD-L1 is needed for direct cell contact between the cell types and hence it plays important role in Treg expansion. The study showed that tolerogenic dendritic cells promote expansion of Tregs via PD-L1 on their surface and reciprocally Tregs facilitate Tol-DCs to maintain transplantation tolerance of pancreatic islets induced by apoptotic cells via secreting IL-10 and TGF-beta. Consequently, the polynucleotides taught herein provide alternative means of tolerizing cells and cellular systems.

[000251] In some embodiments, the tolerogenic polynucleotides or compositions thereof encoding PD-L1 and/or anti-CD40 ligand antibody may be used before, during or after allogenic bone marrow (BM) transplantation to prevent or ameliorate destruction of tissue allografts. Such treatments may improve or aid in the establishment of hematopoietic chimerism where both recipient and donor bone marrow cells coexist. In one embodiment, the PD-L1 encoding polynucleotides effect tolerization of both CD4 and CD8 T cells. In some embodiments, CD8 T-cells are tolerized and in this embodiment, tolerogenic polynucleotides may encode LAG-3, a homolog of CD4 that binds to MHC class II. Materials and methods for this embodiment can be found at, for

example, Lucas et al, Blood, 2011, 117; 5532-5540, the contents of which are incorporated herein by reference in their entirety.

[000252] In some embodiments, tolerogenic polynucleotides may encode anti-CD40 ligand antibodies for the treatment of inflammation or obesity. The polynucleotides of the present invention may be evaluated for such outcomes using the methods as described in Poggi, et al, Arteriosclerosis Thromb. Vasc. Biol, 2011; 31; 2251-2260, the contents of which are incorporated herein by reference in their entirety.

[000253] In some embodiments, anti-CD205 antibodies or CD-205-IgG fusion proteins may be used as antigen carriers in order to tolerize cellular systems. Such antibodies or fusion proteins alone or in combination with an antigen may be delivered with or encoded by any of the tolerogenic polynucleotides of the present invention. Such CD205 fusion proteins and/or CD205 antibodies and/or antigens which may be encoded are taught in for example, Shrimpton, et al, Molecular Immunology, 2009; 46; 1229-1239, the contents of which are incorporated herein by reference in their entirety. Validation may be performed using the methods taught in Shrimpton.

[000254] In some embodiments, the tolerogenic polynucleotides of the present invention may encode soluble CD52. Such a polynucleotide may be used to suppress certain classes of T cells such as such as CD52hiCD4+ cells. In some embodiments, tolerogenic polynucleotides encoding soluble CD52 may be administered to a subject having or suspected of having diabetes. In some embodiments, the tolerogenic polynucleotide encoding soluble CD52 alters the T cell activation as compared to the response to the auto-antigen GAD65. The tolerogenic polynucleotides of the present invention may be evaluated for therapeutic outcomes by the methods described by, for example, Bandala-Sanchez et al, Nature Immunology, 2013, Vol. 14, Pages 741-748; Danke, et al., 2005, J. Autoimmun. Vol. 25, No. 4, Pages 303-311; and Yang et al., 2008, J. Autoimmun. Vol. 31, No. 1, Pages 30-41, the contents of each of which are incorporated herein by reference in their entirety.

[000255] The polynucleotides of the present invention may encode any of the monovalent anti-CD40L antibody polypeptides or fragments thereof disclosed in US Patent 7,563,443, the contents of which are incorporated herein by reference in its

entirety. Such polynucleotides may be used for the treatment of immune related disorders or any disease or condition associated with CD40L function.

[000256] Polynucleotides of the present invention may also encode anti-CD40 antibodies, anti CD154 antibodies and/or Fc-silent anti-CD154 domain antibodies such as those described by Pinelli et al (Am. J. Transplantation, 2013, 1-10, DOI: 10.1111/sjt.12417), the contents of which are incorporated herein by reference in their entirety. Such polynucleotides are useful in prolonging graft survival, inhibition of alloreactive T cell expansion, attenuation of cytokine production of antigen-specific T cells and promotion of the conversion of Foxp⁺ induced Tregs.

[000257] Polynucleotides of the present invention may encode the human anti-CD40 monoclonal antibody, ASKP1240, as disclosed in Oura et al, Am. J. Transplantation, 2012, 12, 1740-54, the contents of which are incorporated herein by reference in its entirety. Such polynucleotides are useful in effecting long term hepatic allograft acceptance. In some embodiments, the polynucleotides of the present invention encode one or more polypeptides or fragments which act to block CD40 (e.g., by blocking binding to CD154) and function as immunosuppressants in the treatment of liver transplantation.

[000258] Polynucleotides of the present invention may be administered in combination with donor specific transfusion (DST). Such co-administration may be used in the treatment of or support of prolongation of islet, cardiac, skin and/or kidney allograft survival. Such methods are described in, for example, Ferrer et al., 2012, PLoS ONE, volume 7, issue 7, e40559, the contents of which are incorporated herein by reference in their entirety.

[000259] Polynucleotides of the present invention may be administered alone or in combination with other polynucleotides or molecules to diminish the number of interferon-gamma producing alloreactive CD8⁺ T cells and/or to reduce the intra-graft expression of inflammatory chemokines.

[000260] In one embodiment, anti-CD70 antibodies may be encoded by the polynucleotides of the present invention and administered either alone or in combination with anti-CD154. Such polynucleotides may be administered along with polynucleotides encoding anti-LFA1 antibodies such as those of Dai et al., to prolong the survival of heart

allografts (Transplant Immunology, 2011, 195-202), the contents of which are incorporated herein by reference in its entirety.

[000261] In some embodiments, the tolerogenic polynucleotides may encode one or more cytokines which permit tolerance to self or non-self antigens. Such cytokines include TGF-beta (including the inactive latent form and the processed form), IL-27, IL-35 and/or IL37, IL-2, IL-10, IL-19, IL-20, IL-22, IL-24, IL-26, including any of the extended IL-10 superfamily as taught by Commins et al, 2008, J. Allergy Clin. Immunol., 121, 1108-1111, the contents of which are incorporated herein by reference in their entirety. These polynucleotides may be co-administered with either a self or non-self antigen.

[000262] In some embodiments, the tolerogenic polynucleotides may encode the Ig-Like Transcript 3 (ILT3) for use in modulating cytokine signaling, allogenic tumor transplantation, GVHD, and/or to induce effective anti-tumor responses in cellular systems. Such polynucleotides or compositions thereof may be evaluated using the methods and/or materials taught in, for example, Suciu-Foca, et al., J. Immunology, 2007, 178, 7432-7441; Vlad and Sucio-Foca, Exp. and Mol. Pathology, 2012, 93, 294-301; Torres-Aguilar, J. Immunol., 2010, 184, 1765-1775; Rutella, et al., Blood, 2006, 108, 1435-1440; Chang, et al., J. Immunol., 2010, 185, 5714-5722; Janikashvili, et al., Clin. and Dev. Immunol., 2011, Article ID 430394; Cheng, et al., J.B.C., 2011, 286, 18013-18025; Chang, et al., J. Immunol., 2012, 188, 000; Chang, et al., Nature Immunology, 2002, 3(3), 237; Steinman, et al., Annu. Rev. Immunol., 2003, 21, 685-711; Vlad, et al., Diabetes, 2008, 57, 1878; Ge, et al., Transplant Immunology, 2012, 26, 19-26, the contents of each of which are incorporated herein by reference in their entirety.

[000263] In some embodiments, the tolerogenic polynucleotides may encode the SOCS1 for use in inhibiting cytokine signaling and/or to induce effective anti-tumor responses in cellular systems. Such polynucleotides or compositions thereof may be evaluated using the methods and/or materials taught in, for example, Evel-Kabler et al., J. Clin. Invest. 2006, 116(1); Yoshimura et al., Arthritis Research & Therapy, 2005, 7(3), 100; the contents of each of which are incorporated herein by reference in their entirety.

[000264] In some embodiments, the tolerogenic polynucleotides may encode the IL-35 for use in treating arthritis, limiting chronic inflammatory disease such as asthma and

inflammatory bowel disease or colitis or in particular microenvironments such as intestinal infection with *Trichuris muris* or tumor microenvironments. Such polynucleotides or compositions thereof may be evaluated using the methods and/or materials taught in, for example, Collison et al., *Nature*, 2007, 450(22), 566-571; Wirtz, et al., *Gastroenterology*, 2011, 141(5), 1875-1886; Kochetkova et al., *J. Immunol.* 2010, 184, 7144-7153; Neidbala, et al., *Eur. J. Immunol.*, 2007, 37, 3021-3029; Collison, *Nature Immunity*, 2010, 11(12), 1093, the contents of each of which are incorporated herein by reference in their entirety.

[000265] In some embodiments, the tolerogenic polynucleotides may encode the IL-37 (formerly known as IL-1H4 and IL-1F7) for use in suppressing inflammatory and/or immune responses. Such polynucleotides or compositions thereof are taught in, and may be evaluated using the methods and/or materials of, for example, Bufler et al., *PNAS*, 2002, 99(21), 13723-13728; Nold et al., *Nature Immunol.* 2010, 11(11), 1014; Kumar et al., *Cytokine*, 2002, 18(2), 61-71; Boraschi, et al., *Eur. Cytokine Netw.*, 2011, 22(3), 127-47; and Sharma et al., *J. Immunol.*, 2008, 180, 5477-5482, the contents of each of which is incorporated herein by reference in their entirety.

[000266] Tolerogenic polynucleotides encoding IL-2 or compositions thereof may be administered to subjects for the amelioration and/or treatment of autoimmune anemia, diabetes, general autoimmunity, humoral immune responses, systemic autoimmunity, dermatomyositis, influenza, or inflammatory bowel disease.

[000267] Tolerogenic polynucleotides encoding IL-10 or compositions thereof may be administered to subjects for the amelioration and/or treatment of infections, intestinal homeostasis including colitis, inflammatory bowel disease, autoimmune diseases including lupus, cancers such as melanoma and infectious diseases including leishmaniasis and tuberculosis, Chron's disease, and psoriasis.

[000268] Tolerogenic polynucleotides encoding IL-27, IL-35 and/or IL37 or compositions thereof may be administered to subjects for the amelioration and/or treatment of pathogen-associated conditions or diseases.

[000269] In some embodiments, the tolerogenic polynucleotides may encode the hematopoietic growth factor, Flt3L for use in promoting immune tolerance. Such polynucleotides or compositions thereof may be evaluated using the encoded Flt3 ligand,

methods and/or materials taught in, for example, Klein et al, Eur. J. Immunology, 2013, 43, 533-539, the contents of which are incorporated herein by reference in their entirety.

[000270] In some embodiments, the tolerogenic polynucleotides may encode the growth factor, TGFbeta 1. Such polynucleotides or compositions thereof may be evaluated using encoded TGFbeta1, in methods and/or materials taught in, for example, Li, et al., BMB Reports, 201245(9), 509-514, the contents of which are incorporated herein by reference in their entirety.

[000271] In some embodiments, the tolerogenic polynucleotides may encode the galectin-1 for use in suppressing autoimmune neuroinflammation. Such polynucleotides may be evaluated using the encoded galectin-1, an endogenous glycan-binding protein, the methods and/or materials taught in, for example, Ilarregui, et al, Nature Immunology, 2009, vol 10, no. 9, 981, the contents of which are incorporated herein by reference in their entirety.

[000272] In some embodiments, the tolerogenic polynucleotides may encode a soluble GARP (glycoprotein A repetitions predominant) protein for use as an immunomodulator of inflammatory diseases including transplant rejection, autoimmunity and allergy. Such polynucleotides may be evaluated by encoding the soluble GARP (sGARP) or using the methods and/or materials taught in, for example, Hahn, et al, Blood, 2013, 122(7); 1182-1191 or the methods and/or materials taught in, for example, Wang, et al., PNAS, 2009, 106(32), 13439-13444, the contents of both of which are incorporated herein by reference in their entirety.

[000273] In some embodiments, the tolerogenic polynucleotides of the present invention may encode one or more tregitopes, tregitope conjugates or tregitope cores, fragments or cores having flanks. Included in the tregitope family are any of those disclosed in, for example, US Patent 7,884,184, van der Marel, et al., World J. Gastroent. 2012, 18(32) 4288-4299; Hui, et al., Mol. Therapy, 2013, 21(9), 1727-1737; Elyaman et al., Neurol. Res. International, 2011 Article ID256460; De Groot, et al., Blood 2008, 112, 3303-3311; Cousens et al., J. Diab. Research, 2013, Article ID 621693; Cousens, et al., Autoimmunity Reviews, 2012, doi: 10.1016, the contents of which are incorporated herein by reference in its entirety.

[000274] Evaluation of such encoded tregitopes may be performed using the materials and/or methods taught in, US Patent 7,884,184, van der Marel, et al., World J. Gastroenterol. 2012, 18(32) 4288-4299; Hui, et al., Mol. Therapy, 2013, 21(9), 1727-1737; Elyaman et al., Neurol. Res. International, 2011 Article ID256460; De Groot, et al., Blood 2008, 112, 3303-3311; Cousens et al., J. Diab. Research, 2013, Article ID 621693; Cousens, et al., Autoimmunity Reviews, 2012, doi: 10.1016, the contents of which are incorporated herein by reference in its entirety.

[000275] In some embodiments, the tolerogenic polynucleotides may encode one or more proteins from a parasite such as a helminth, including but not limited to the Sj16 protein from *Schistosoma Japonicum* or a total extract from *Fasciola hepatica*, or high molecular weight components of *Ascaris suum*, or soluble worm extract (SWA) or eggs (SEA) of *Schistosoma mansoni*. Such tolerogenic polynucleotides or compositions thereof are useful in the treatment of wounds to improve wound healing, diabetes, inflammation or protection against parasitic infections. Encoded proteins or peptides or extracts include those taught in, for example Hu et al, International J. Parasitol. 2009; 38, 191-200; Hu et al, J. Parasitol. 2012; 98(2), 274-283; Carranza et al, 2012, PLoS ONE, vol. 7, Issue 7, 1-9; Sun et al, 2012, Parasite Immunology, 34, 430-439; Zacccone et al., 2010, J. Biomed. And Biotech., ArticleID 795210, doi:10.1155/2010/79521; Silva et al., 2006, Eur. J. Immunol., 36, 3227-3237; Gravano and Vignali, 2012, Cell Mol. Life Sci, 69(12), 1997-2008; and Gause et al, 2013, Nature Reviews: Immunology, 13, 607-614, the contents of each of which is incorporated herein by reference in their entireties.

Evaluation of such encoded proteins or peptides may be performed using the materials and/or methods taught in, for example, Hu et al, International J. Parasitol. 2009; 38, 191-200; Hu et al, J. Parasitol. 2012; 98(2), 274-283; Carranza et al, 2012, PLoS ONE, vol. 7, Issue 7, 1-9; Sun et al, 2012, Parasite Immunology, 34, 430-439; Zacccone et al., 2010, J. Biomed. And Biotech., ArticleID 795210, doi:10.1155/2010/79521; Silva et al., 2006, Eur. J. Immunol., 36, 3227-3237, the contents of each of which is incorporated herein by reference in their entirety.

[000276] In some embodiments, the tolerogenic polynucleotides may encode one or more peptides for use in the inhibition of Tcell receptor (TCR) signaling. Such tolerogenic polynucleotides or compositions thereof are useful in the treatment of a

disease or condition where T-cells are involved and are taught in for example US Publication 2013/0039948 published February 14, 2013, the contents of which are incorporated herein by reference in its entirety.

[000277] Evaluation of such encoded proteins or peptides may be performed using the materials and/or methods taught in, for example, example US Publication 2013/0039948 published February 14, 2013, the contents of which are incorporated herein by reference in its entirety.

[000278] In some embodiments, the tolerogenic polynucleotides may encode the tryptophan catabolizing enzyme indoleamine 2,3,-dioxygenase (IDO). Such tolerogenic polynucleotides or compositions thereof are useful in the treatment of systemic autoimmune disease. Evaluation of such encoded proteins or peptides may be performed using the materials and/or methods taught in, for example, Ravishankar et al, 2012, PNAS, Early Edition, doi/10.1073/pnas.1117736109, 1-6; Munn and Mellor, 2013, Cell, 34(3), 137-143; Curti, et al, 2009, Blood, 113, 2394-2401, the contents of each of which are incorporated herein by reference in their entirety.

[000279] In some embodiments, the tolerogenic polynucleotides may encode the foxp3 transcription factor. Such tolerogenic polynucleotides or compositions thereof may also comprise a cell penetrating peptide such as the N-terminal fragment of long PTEN isoform. These may be delivered to target APCs using any formulation taught herein, such as lipid nanoparticles. These may also incorporate or comprise one or more microRNA, microRNA binding site or precursor to minimize non-APC, non-DC and non lymphocytic translation. Evaluation of such encoded proteins or peptides may be performed using the materials and/or methods taught in, for example Hopkins et al, 2013, Science, Vol. 341, No. 6144, Pages 399-402.

Protein Cleavage Signals and Sites

[000280] In one embodiment, the polypeptides of the present invention may include at least one protein cleavage signal containing at least one protein cleavage site. The protein cleavage site may be located at the N-terminus, the C-terminus, at any space between the N- and the C- termini such as, but not limited to, half-way between the N- and C-termini, between the N-terminus and the half way point, between the half way point and the C-terminus, and combinations thereof.

[000281] In one embodiment, the polynucleotides of the present invention may be engineered such that the polynucleotide contains at least one encoded protein cleavage signal. The encoded protein cleavage signal may be located in any region including but not limited to before the start codon, after the start codon, before the coding region, within the coding region such as, but not limited to, half way in the coding region, between the start codon and the half way point, between the half way point and the stop codon, after the coding region, before the stop codon, between two stop codons, after the stop codon and combinations thereof.

[000282] The encoded protein cleavage signal may include, but is not limited to, a proprotein convertase (or prohormone convertase), thrombin and/or Factor Xa protein cleavage signal.

[000283] As a non-limiting example, U.S. Pat. No. 7,374,930 and U.S. Pub. No. 20090227660, herein incorporated by reference in their entireties, use a furin cleavage site to cleave the N-terminal methionine of GLP-1 in the expression product from the Golgi apparatus of the cells. In one embodiment, the polypeptides of the present invention include at least one protein cleavage signal and/or site with the proviso that the polypeptide is not GLP-1.

Insertions and Substitutions

[000284] In one embodiment, the 5'UTR of the polynucleotide may be replaced by the insertion of at least one region and/or string of nucleosides of the same base. The region and/or string of nucleotides may include, but is not limited to, at least 3, at least 4, at least 5, at least 6, at least 7 or at least 8 nucleotides and the nucleotides may be natural and/or unnatural. As a non-limiting example, the group of nucleotides may include 5-8 adenine, cytosine, thymine, a string of any of the other nucleotides disclosed herein and/or combinations thereof.

[000285] In one embodiment, the 5'UTR of the polynucleotide may be replaced by the insertion of at least two regions and/or strings of nucleotides of two different bases such as, but not limited to, adenine, cytosine, thymine, any of the other nucleotides disclosed herein and/or combinations thereof. For example, the 5'UTR may be replaced by inserting 5-8 adenine bases followed by the insertion of 5-8 cytosine bases. In another

example, the 5'UTR may be replaced by inserting 5-8 cytosine bases followed by the insertion of 5-8 adenine bases.

[000286] In one embodiment, the polynucleotide may include at least one substitution and/or insertion downstream of the transcription start site which may be recognized by an RNA polymerase. As a non-limiting example, at least one substitution and/or insertion may occur downstream the transcription start site by substituting at least one nucleic acid in the region just downstream of the transcription start site (such as, but not limited to, +1 to +6). Changes to region of nucleotides just downstream of the transcription start site may affect initiation rates, increase apparent nucleotide triphosphate (NTP) reaction constant values, and increase the dissociation of short transcripts from the transcription complex curing initial transcription (Briebe et al, Biochemistry (2002) 41: 5144-5149; herein incorporated by reference in its entirety). The modification, substitution and/or insertion of at least one nucleoside may cause a silent mutation of the sequence or may cause a mutation in the amino acid sequence.

[000287] In one embodiment, the polynucleotide may include the substitution of at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12 or at least 13 guanine bases downstream of the transcription start site.

[000288] In one embodiment, the polynucleotide may include the substitution of at least 1, at least 2, at least 3, at least 4, at least 5 or at least 6 guanine bases in the region just downstream of the transcription start site. As a non-limiting example, if the nucleotides in the region are GGGAGA the guanine bases may be substituted by at least 1, at least 2, at least 3 or at least 4 adenine nucleotides. In another non-limiting example, if the nucleotides in the region are GGGAGA the guanine bases may be substituted by at least 1, at least 2, at least 3 or at least 4 cytosine bases. In another non-limiting example, if the nucleotides in the region are GGGAGA the guanine bases may be substituted by at least 1, at least 2, at least 3 or at least 4 thymine, and/or any of the nucleotides described herein.

[000289] In one embodiment, the polynucleotide may include at least one substitution and/or insertion upstream of the start codon. For the purpose of clarity, one of skill in the art would appreciate that the start codon is the first codon of the protein coding region

whereas the transcription start site is the site where transcription begins. The polynucleotide may include, but is not limited to, at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7 or at least 8 substitutions and/or insertions of nucleotide bases. The nucleotide bases may be inserted or substituted at 1, at least 1, at least 2, at least 3, at least 4 or at least 5 locations upstream of the start codon. The nucleotides inserted and/or substituted may be the same base (e.g., all A or all C or all T or all G), two different bases (e.g., A and C, A and T, or C and T), three different bases (e.g., A, C and T or A, C and T) or at least four different bases. As a non-limiting example, the guanine base upstream of the coding region in the polynucleotide may be substituted with adenine, cytosine, thymine, or any of the nucleotides described herein. In another non-limiting example the substitution of guanine bases in the polynucleotide may be designed so as to leave one guanine base in the region downstream of the transcription start site and before the start codon (see Esvelt *et al.* Nature (2011) 472(7344):499-503; the contents of which is herein incorporated by reference in its entirety). As a non-limiting example, at least 5 nucleotides may be inserted at 1 location downstream of the transcription start site but upstream of the start codon and the at least 5 nucleotides may be the same base type.

Incorporating Post Transcriptional Control Modulators

[000290] In one embodiment, the polynucleotides of the present invention may include at least one post transcriptional control modulator. These post transcriptional control modulators may be, but are not limited to, small molecules, compounds and regulatory sequences. As a non-limiting example, post transcriptional control may be achieved using small molecules identified by PTC Therapeutics Inc. (South Plainfield, NJ) using their GEMSTM (Gene Expression Modulation by Small-Molecules) screening technology.

[000291] The post transcriptional control modulator may be a gene expression modulator which is screened by the method detailed in or a gene expression modulator described in International Publication No. WO2006022712, herein incorporated by reference in its entirety. Methods identifying RNA regulatory sequences involved in translational control are described in International Publication No. WO2004067728, herein incorporated by reference in its entirety; methods identifying compounds that modulate untranslated region dependent expression of a gene are described in

International Publication No. WO2004065561, herein incorporated by reference in its entirety.

[000292] In one embodiment, the polynucleotides of the present invention may include at least one post transcriptional control modulator is located in the 5' and/or the 3' untranslated region of the polynucleotides of the present invention.

[000293] In another embodiment, the polynucleotides of the present invention may include at least one post transcription control modulator to modulate premature translation termination. The post transcription control modulators may be compounds described in or a compound found by methods outlined in International Publication No. WO2004010106, WO2006044456, WO2006044682, WO2006044503 and WO2006044505, each of which is herein incorporated by reference in its entirety. As a non-limiting example, the compound may bind to a region of the 28S ribosomal RNA in order to modulate premature translation termination (See e.g., WO2004010106, herein incorporated by reference in its entirety).

[000294] In one embodiment, polynucleotides of the present invention may include at least one post transcription control modulator to alter protein expression. As a non-limiting example, the expression of VEGF may be regulated using the compounds described in or a compound found by the methods described in International Publication Nos. WO2005118857, WO2006065480, WO2006065479 and WO2006058088, each of which is herein incorporated by reference in its entirety.

[000295] The polynucleotides of the present invention may include at least one post transcription control modulator to control translation. In one embodiment, the post transcription control modulator may be a RNA regulatory sequence. As a non-limiting example, the RNA regulatory sequence may be identified by the methods described in International Publication No. WO2006071903, herein incorporated by reference in its entirety.

II. Design, Synthesis and Quantitation of Polynucleotides

Codon Optimization

[000296] The polynucleotides, their regions or parts or subregions may be codon optimized. Codon optimization methods are known in the art and may be useful in efforts to achieve one or more of several goals. These goals include to match codon frequencies

in target and host organisms to ensure proper folding, bias GC content to increase mRNA stability or reduce secondary structures, minimize tandem repeat codons or base runs that may impair gene construction or expression, customize transcriptional and translational control regions, insert or remove protein trafficking sequences, remove/add post translation modification sites in encoded protein (e.g. glycosylation sites), add, remove or shuffle protein domains, insert or delete restriction sites, modify ribosome binding sites and mRNA degradation sites, to adjust translational rates to allow the various domains of the protein to fold properly, or to reduce or eliminate problem secondary structures within the polynucleotide. Codon optimization tools, algorithms and services are known in the art, non-limiting examples include services from GeneArt (Life Technologies), DNA2.0 (Menlo Park CA) and/or proprietary methods. In one embodiment, the ORF sequence is optimized using optimization algorithms. Codon options for each amino acid are given in Table 4.

Table 4. Codon Options

Amino Acid	Single Letter Code	Codon Options
Isoleucine	I	ATT, ATC, ATA
Leucine	L	CTT, CTC, CTA, CTG, TTA, TTG
Valine	V	GTT, GTC, GTA, GTG
Phenylalanine	F	TTT, TTC
Methionine	M	ATG
Cysteine	C	TGT, TGC
Alanine	A	GCT, GCC, GCA, GCG
Glycine	G	GGT, GGC, GGA, GGG
Proline	P	CCT, CCC, CCA, CCG
Threonine	T	ACT, ACC, ACA, ACG
Serine	S	TCT, TCC, TCA, TCG, AGT, AGC
Tyrosine	Y	TAT, TAC
Tryptophan	W	TGG
Glutamine	Q	CAA, CAG
Asparagine	N	AAT, AAC
Histidine	H	CAT, CAC
Glutamic acid	E	GAA, GAG
Aspartic acid	D	GAT, GAC
Lysine	K	AAA, AAG
Arginine	R	CGT, CGC, CGA, CGG, AGA, AGG
Selenocysteine	Sec	UGA in mRNA in presence of Selenocysteine insertion element (SECIS)
Stop codons	Stop	TAA, TAG, TGA

[000297] Features, which may be considered beneficial in some embodiments of the present invention, may be encoded by regions of the polynucleotide and such regions may be upstream (5') or downstream (3') to a region which encodes a polypeptide. These regions may be incorporated into the polynucleotide before and/or after codon optimization of the protein encoding region or open reading frame (ORF). It is not required that a polynucleotide contain both a 5' and 3' flanking region. Examples of such features include, but are not limited to, untranslated regions (UTRs), Kozak sequences, an oligo(dT) sequence, and detectable tags and may include multiple cloning sites which may have XbaI recognition.

[000298] In some embodiments, a 5' UTR and/or a 3' UTR region may be provided as flanking regions. Multiple 5' or 3' UTRs may be included in the flanking regions and may be the same or of different sequences. Any portion of the flanking regions, including none, may be codon optimized and any may independently contain one or more different structural or chemical modifications, before and/or after codon optimization.

[000299] After optimization (if desired), the polynucleotides components are reconstituted and transformed into a vector such as, but not limited to, plasmids, viruses, cosmids, and artificial chromosomes. For example, the optimized polynucleotide may be reconstituted and transformed into chemically competent *E. coli*, yeast, neurospora, maize, drosophila, etc. where high copy plasmid-like or chromosome structures occur by methods described herein.

[000300] Synthetic polynucleotides and their nucleic acid analogs play an important role in the research and studies of biochemical processes. Various enzyme-assisted and chemical-based methods have been developed to synthesize polynucleotides and nucleic acids.

Enzymatic Methods

[000301] The polynucleotides of the present invention may be synthesized using enzymatic methods known in the art. As a non-limiting example, enzymatic methods, *in vitro* transcription-enzymatic synthesis and RNA polymerases useful for synthesis are described in co-pending International Publication No. WO2015034928, the contents of which are herein incorporated by reference, such as in paragraphs [000276] – [000297].

Solid-phase chemical synthesis

The polynucleotides of the present invention (e.g., chimeric polynucleotides or circular polynucleotides) may be manufactured in whole or in part using solid phase techniques. As a non-limiting example, solid phase techniques useful for synthesis are described in co-pending International Publication No. WO2015034928, the contents of which are herein incorporated by reference, such as in paragraphs [000298] – [000307]. Liquid Phase Chemical Synthesis

[000302] The synthesis of chimeric polynucleotides or circular polynucleotides of the present invention by the sequential addition of monomer building blocks may be carried out in a liquid phase. A covalent bond is formed between the monomers or between a terminal functional group of the growing chain and an incoming monomer. Functional groups not involved in the reaction must be temporarily protected. After the addition of each monomer building block, the reaction mixture has to be purified before adding the next monomer building block. The functional group at one terminal of the chain has to be deprotected to be able to react with the next monomer building blocks. A liquid phase synthesis is labor- and time-consuming and cannot not be automated. Despite the limitations, liquid phase synthesis is still useful in preparing short polynucleotides in a large scale. Because the system is homogenous, it does not require a large excess of reagents and is cost-effective in this respect.

Combination of Different Synthetic Methods

[000303] The synthetic methods discussed above each has its own advantages and limitations. Attempts have been conducted to combine these methods to overcome the limitations. Such combinations of methods are within the scope of the present invention. As a non-limiting example, combinations of the different synthetic methods useful for the present invention are described in co-pending International Publication No. WO2015034928, the contents of which are herein incorporated by reference, such as in paragraphs [000309] – [000312].

Small Region Synthesis

[000304] Regions or subregions of the polynucleotides of the present invention may comprise small RNA molecules such as siRNA, and therefore may be synthesized in the same manner. There are several methods for preparing siRNA, such as chemical synthesis using appropriately protected ribonucleoside phosphoramidites, *in vitro*

transcription, siRNA expression vectors, and PCR expression cassettes. Sigma-Aldrich® is one of the siRNA suppliers and synthesizes their siRNA using ribonucleoside phosphoramidite monomers protected at the 2' position with a t-butylmethylsilyl (TBDMS) group. The solid-phase chemical synthesis is carried out with SIGMA-ALDRICH®'s Ultra Fast Parallel Synthesis (UFPS) and Ultra Fast Parallel Deprotection (UFPD) to achieve high coupling efficiency and fast deprotection. The final siRNA products may be purified with HPLC or PAGE. Such methods may be used to synthesize regions or subregions of chimeric polynucleotides.

[000305] *In vitro* transcription and expression from a vector or a PCR-generated siRNA cassette require appropriate templates to produce siRNAs. The commercially available AMBION® SILENCER® siRNA construction kit produces siRNA by *in vitro* transcription of DNA templates and contains the enzymes, buffers, primers needed. Such methods may be used to synthesize regions or subregions of chimeric polynucleotides.

Ligation of Polynucleotide Regions or Subregions

[000306] Polynucleotides such as chimeric polynucleotides and/or circular polynucleotides may be prepared by ligation of one or more regions or subregions. As a non-limiting example, methods for the ligation of polynucleotide regions or subregions useful in the present invention are described in co-pending International Publication No. WO2015034928, the contents of which are herein incorporated by reference, such as in paragraphs [000315] – [000322].

Modification and Conjugation of Polynucleotides

[000307] Non-natural modified nucleotides may be introduced to polynucleotides or nucleic acids during synthesis or post-synthesis of the chains to achieve desired functions or properties. The modifications may be on internucleotide lineage, the purine or pyrimidine bases, or sugar. The modification may be introduced at the terminal of a chain or anywhere else in the chain; with chemical synthesis or with a polymerase enzyme. For example, hexitol nucleic acids (HNAs) are nuclease resistant and provide strong hybridization to RNA. Short messenger RNAs (mRNAs) with hexitol residues in two codons have been constructed (Lavrik et al., *Biochemistry*, 40, 11777-11784 (2001), the contents of which are incorporated herein by reference in their entirety). The

antisense effects of a chimeric HNA gapmer oligonucleotide comprising a phosphorothioate central sequence flanked by 5' and 3' HNA sequences have also been studied (See e.g., Kang et al., *Nucleic Acids Research*, vol. 32(4), 4411-4419 (2004), the contents of which are incorporated herein by reference in their entirety). The preparation and uses of modified nucleotides comprising 6-member rings in RNA interference, antisense therapy or other applications are disclosed in US Pat. Application No. 2008/0261905, US Pat. Application No. 2010/0009865, and PCT Application No. WO97/30064 to Herdewijn et al.; the contents of each of which are herein incorporated by reference in their entirety). Modified nucleic acids and their synthesis are disclosed in copending International Patent Publication No. WO2013052523 (Attorney Docket Number M09), the contents of which are incorporated herein by reference in their entirety. The synthesis and strategy of modified polynucleotides is reviewed by Verma and Eckstein in *Annual Review of Biochemistry*, vol. 76, 99-134 (1998), the contents of which are incorporated herein by reference in their entirety.

[000308] Either enzymatic or chemical ligation methods can be used to conjugate polynucleotides or their regions with different functional blocks, such as fluorescent labels, liquids, nanoparticles, delivery agents, etc. The conjugates of polynucleotides and modified polynucleotides are reviewed by Goodchild in *Bioconjugate Chemistry*, vol. 1(3), 165-187 (1990), the contents of which are incorporated herein by reference in their entirety. US Pat. No. 6,835,827 and US Pat. No. 6,525,183 to Vinayak et al. (the contents of each of which are herein incorporated by reference in their entirety) teach synthesis of labeled oligonucleotides using a labeled solid support.

Quantification

[000309] In one embodiment, the polynucleotides of the present invention may be quantified in exosomes or when derived from one or more bodily fluid. As used herein "bodily fluids" include peripheral blood, serum, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, bronchoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, sweat, fecal matter, hair, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice,

lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, and umbilical cord blood. Alternatively, exosomes may be retrieved from an organ selected from the group consisting of lung, heart, pancreas, stomach, intestine, bladder, kidney, ovary, testis, skin, colon, breast, prostate, brain, esophagus, liver, and placenta.

[000310] In the exosome quantification method, a sample of not more than 2mL is obtained from the subject and the exosomes isolated by size exclusion chromatography, density gradient centrifugation, differential centrifugation, nanomembrane ultrafiltration, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof. In the analysis, the level or concentration of a polynucleotide may be an expression level, presence, absence, truncation or alteration of the administered construct. It is advantageous to correlate the level with one or more clinical phenotypes or with an assay for a human disease biomarker. The assay may be performed using construct specific probes, cytometry, qRT-PCR, real-time PCR, PCR, flow cytometry, electrophoresis, mass spectrometry, or combinations thereof while the exosomes may be isolated using immunohistochemical methods such as enzyme linked immunosorbent assay (ELISA) methods. Exosomes may also be isolated by size exclusion chromatography, density gradient centrifugation, differential centrifugation, nanomembrane ultrafiltration, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof.

[000311] These methods afford the investigator the ability to monitor, in real time, the level of polynucleotides remaining or delivered. This is possible because the polynucleotides of the present invention differ from the endogenous forms due to the structural or chemical modifications.

[000312] In one embodiment, the polynucleotide may be quantified using methods such as, but not limited to, ultraviolet visible spectroscopy (UV/Vis). A non-limiting example of a UV/Vis spectrometer is a NANODROP® spectrometer (ThermoFisher, Waltham, MA). The quantified polynucleotide may be analyzed in order to determine if the polynucleotide may be of proper size, check that no degradation of the polynucleotide has occurred. Degradation of the polynucleotide may be checked by methods such as, but not limited to, agarose gel electrophoresis, HPLC based purification methods such as, but not limited to, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase

HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC), liquid chromatography-mass spectrometry (LCMS), capillary electrophoresis (CE) and capillary gel electrophoresis (CGE).

Purification

[000313] Purification of the polynucleotides described herein may include, but is not limited to, polynucleotide clean-up, quality assurance and quality control. Clean-up may be performed by methods known in the arts such as, but not limited to, AGENCOURT® beads (Beckman Coulter Genomics, Danvers, MA), poly-T beads, LNA™ oligo-T capture probes (EXIQON® Inc, Vedbaek, Denmark) or HPLC based purification methods such as, but not limited to, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC). The term “purified” when used in relation to a polynucleotide such as a “purified polynucleotide” refers to one that is separated from at least one contaminant. As used herein, a “contaminant” is any substance which makes another unfit, impure or inferior. Thus, a purified polynucleotide (e.g., DNA and RNA) is present in a form or setting different from that in which it is found in nature, or a form or setting different from that which existed prior to subjecting it to a treatment or purification method.

[000314] A quality assurance and/or quality control check may be conducted using methods such as, but not limited to, gel electrophoresis, UV absorbance, or analytical HPLC.

[000315] In another embodiment, the polynucleotides may be sequenced by methods including, but not limited to reverse-transcriptase-PCR.

III. Modifications

[000316] As used herein in a polynucleotide (such as a chimeric polynucleotide, IVT polynucleotide or a circular polynucleotide), the terms “chemical modification” or, as appropriate, “chemically modified” refer to modification with respect to adenosine (A), guanosine (G), uridine (U), thymidine (T) or cytidine (C) ribo- or deoxyribonucleotides in one or more of their position, pattern, percent or population. Generally, herein, these terms are not intended to refer to the ribonucleotide modifications in naturally occurring 5'-terminal mRNA cap moieties.

[000317] In a polypeptide, the term “modification” refers to a modification as compared to the canonical set of 20 amino acids.

[000318] The modifications may be various distinct modifications. In some embodiments, the regions may contain one, two, or more (optionally different) nucleoside or nucleotide modifications. In some embodiments, a modified polynucleotide, introduced to a cell may exhibit reduced degradation in the cell, as compared to an unmodified polynucleotide.

[000319] Modifications which are useful in the present invention include, but are not limited to those in Table 5. Noted in the table are the symbol of the modification, the nucleobase type and whether the modification is naturally occurring or not.

Table 5. Modifications

Name	Symbol	Base	Naturally Occurring
2-methylthio-N6-(cis-hydroxyisopentenyl)adenosine	ms2i6A	A	YES
2-methylthio-N6-methyladenosine	ms2m6A	A	YES
2-methylthio-N6-threonyl carbamoyladenosine	ms2t6A	A	YES
N6-glycinylicarbamoyladenosine	g6A	A	YES
N6-isopentenyladenosine	i6A	A	YES
N6-methyladenosine	m6A	A	YES
N6-threonylicarbamoyladenosine	t6A	A	YES
1,2'-O-dimethyladenosine	m1Am	A	YES
1-methyladenosine	m1A	A	YES
2'-O-methyladenosine	Am	A	YES
2'-O-ribosyladenosine (phosphate)	Ar(p)	A	YES
2-methyladenosine	m2A	A	YES
2-methylthio-N6 isopentenyladenosine	ms2i6A	A	YES
2-methylthio-N6-hydroxynorvalyl carbamoyladenosine	ms2hn6A	A	YES
2'-O-methyladenosine	m6A	A	YES
2'-O-ribosyladenosine (phosphate)	Ar(p)	A	YES
isopentenyladenosine	Iga	A	YES
N6-(cis-hydroxyisopentenyl)adenosine	io6A	A	YES
N6,2'-O-dimethyladenosine	m6Am	A	YES
N ⁶ ,2'-O-dimethyladenosine	m ⁶ Am	A	YES
N6,N6,2'-O-trimethyladenosine	m62Am	A	YES
N6,N6-dimethyladenosine	m62A	A	YES
N6-acetyladenosine	ac6A	A	YES
N6-hydroxynorvalylcarbamoyladenosine	hn6A	A	YES
N6-methyl-N6-threonylicarbamoyladenosine	m6t6A	A	YES

2-methyladenosine	m ² A	A	YES
2-methylthio-N ⁶ -isopentenyladenosine	ms ² i ⁶ A	A	YES
7-deaza-adenosine	--	A	NO
N1-methyl-adenosine	--	A	NO
N6, N6 (dimethyl)adenine	--	A	NO
N6-cis-hydroxy-isopentenyl-adenosine	--	A	NO
α-thio-adenosine	--	A	NO
2 (amino)adenine	--	A	NO
2 (aminopropyl)adenine	--	A	NO
2 (methylthio) N6 (isopentenyl)adenine	--	A	NO
2-(alkyl)adenine	--	A	NO
2-(aminoalkyl)adenine	--	A	NO
2-(aminopropyl)adenine	--	A	NO
2-(halo)adenine	--	A	NO
2-(halo)adenine	--	A	NO
2-(propyl)adenine	--	A	NO
2'-Amino-2'-deoxy-ATP	--	A	NO
2'-Azido-2'-deoxy-ATP	--	A	NO
2'-Deoxy-2'-a-aminoadenosine TP	--	A	NO
2'-Deoxy-2'-a-azidoadenosine TP	--	A	NO
6 (alkyl)adenine	--	A	NO
6 (methyl)adenine	--	A	NO
6-(alkyl)adenine	--	A	NO
6-(methyl)adenine	--	A	NO
7 (deaza)adenine	--	A	NO
8 (alkenyl)adenine	--	A	NO
8 (alkynyl)adenine	--	A	NO
8 (amino)adenine	--	A	NO
8 (thioalkyl)adenine	--	A	NO
8-(alkenyl)adenine	--	A	NO
8-(alkyl)adenine	--	A	NO
8-(alkynyl)adenine	--	A	NO
8-(amino)adenine	--	A	NO
8-(halo)adenine	--	A	NO
8-(hydroxyl)adenine	--	A	NO
8-(thioalkyl)adenine	--	A	NO
8-(thiol)adenine	--	A	NO
8-azido-adenosine	--	A	NO
aza adenine	--	A	NO
deaza adenine	--	A	NO
N6 (methyl)adenine	--	A	NO
N6-(isopentyl)adenine	--	A	NO
7-deaza-8-aza-adenosine	--	A	NO

7-methyladenine	--	A	NO
1-Deazaadenosine TP	--	A	NO
2'Fluoro-N6-Bz-deoxyadenosine TP	--	A	NO
2'-OMe-2-Amino-ATP	--	A	NO
2'O-methyl-N6-Bz-deoxyadenosine TP	--	A	NO
2'-a-Ethynyladenosine TP	--	A	NO
2-aminoadenine	--	A	NO
2-Aminoadenosine TP	--	A	NO
2-Amino-ATP	--	A	NO
2'-a-Trifluoromethyladenosine TP	--	A	NO
2-Azidoadenosine TP	--	A	NO
2'-b-Ethynyladenosine TP	--	A	NO
2-Bromoadenosine TP	--	A	NO
2'-b-Trifluoromethyladenosine TP	--	A	NO
2-Chloroadenosine TP	--	A	NO
2'-Deoxy-2',2'-difluoroadenosine TP	--	A	NO
2'-Deoxy-2'-a-mercaptopadenosine TP	--	A	NO
2'-Deoxy-2'-a-thiomethoxyadenosine TP	--	A	NO
2'-Deoxy-2'-b-aminoadenosine TP	--	A	NO
2'-Deoxy-2'-b-azidoadenosine TP	--	A	NO
2'-Deoxy-2'-b-bromoadenosine TP	--	A	NO
2'-Deoxy-2'-b-chloroadenosine TP	--	A	NO
2'-Deoxy-2'-b-fluoroadenosine TP	--	A	NO
2'-Deoxy-2'-b-iodoadenosine TP	--	A	NO
2'-Deoxy-2'-b-mercaptopadenosine TP	--	A	NO
2'-Deoxy-2'-b-thiomethoxyadenosine TP	--	A	NO
2-Fluoroadenosine TP	--	A	NO
2-Iodoadenosine TP	--	A	NO
2-Mercaptopadenosine TP	--	A	NO
2-methoxy-adenine	--	A	NO
2-methylthio-adenine	--	A	NO
2-Trifluoromethyladenosine TP	--	A	NO
3-Deaza-3-bromoadenosine TP	--	A	NO
3-Deaza-3-chloroadenosine TP	--	A	NO
3-Deaza-3-fluoroadenosine TP	--	A	NO
3-Deaza-3-iodoadenosine TP	--	A	NO
3-Deazaadenosine TP	--	A	NO
4'-Azidoadenosine TP	--	A	NO
4'-Carbocyclic adenosine TP	--	A	NO
4'-Ethynyladenosine TP	--	A	NO
5'-Homo-adenosine TP	--	A	NO
8-Aza-ATP	--	A	NO
8-bromo-adenosine TP	--	A	NO

8-Trifluoromethyladenosine TP	--	A	NO
9-Deazaadenosine TP	--	A	NO
2-aminopurine	--	A/G	NO
7-deaza-2,6-diaminopurine	--	A/G	NO
7-deaza-8-aza-2,6-diaminopurine	--	A/G	NO
7-deaza-8-aza-2-aminopurine	--	A/G	NO
2,6-diaminopurine	--	A/G	NO
7-deaza-8-aza-adenine, 7-deaza-2-aminopurine	--	A/G	NO
2-thiocytidine	s2C	C	YES
3-methylcytidine	m3C	C	YES
5-formylcytidine	f5C	C	YES
5-hydroxymethylcytidine	hm5C	C	YES
5-methylcytidine	m5C	C	YES
N4-acetylcytidine	ac4C	C	YES
2'-O-methylcytidine	Cm	C	YES
2'-O-methylcytidine	Cm	C	YES
5,2'-O-dimethylcytidine	m5 Cm	C	YES
5-formyl-2'-O-methylcytidine	f5Cm	C	YES
lysidine	k2C	C	YES
N4,2'-O-dimethylcytidine	m4Cm	C	YES
N4-acetyl-2'-O-methylcytidine	ac4Cm	C	YES
N4-methylcytidine	m4C	C	YES
N4,N4-Dimethyl-2'-OMe-Cytidine TP	--	C	YES
4-methylcytidine	--	C	NO
5-aza-cytidine	--	C	NO
Pseudo-iso-cytidine	--	C	NO
pyrrolo-cytidine	--	C	NO
α -thio-cytidine	--	C	NO
2-(thio)cytosine	--	C	NO
2'-Amino-2'-deoxy-CTP	--	C	NO
2'-Azido-2'-deoxy-CTP	--	C	NO
2'-Deoxy-2'-a-aminocytidine TP	--	C	NO
2'-Deoxy-2'-a-azidocytidine TP	--	C	NO
3 (deaza) 5 (aza)cytosine	--	C	NO
3 (methyl)cytosine	--	C	NO
3-(alkyl)cytosine	--	C	NO
3-(deaza) 5 (aza)cytosine	--	C	NO
3-(methyl)cytidine	--	C	NO
4,2'-O-dimethylcytidine	--	C	NO
5 (halo)cytosine	--	C	NO
5 (methyl)cytosine	--	C	NO
5 (propynyl)cytosine	--	C	NO
5 (trifluoromethyl)cytosine	--	C	NO

5-(alkyl)cytosine	--	C	NO
5-(alkynyl)cytosine	--	C	NO
5-(halo)cytosine	--	C	NO
5-(propynyl)cytosine	--	C	NO
5-(trifluoromethyl)cytosine	--	C	NO
5-bromo-cytidine	--	C	NO
5-iodo-cytidine	--	C	NO
5-propynyl cytosine	--	C	NO
6-(azo)cytosine	--	C	NO
6-aza-cytidine	--	C	NO
aza cytosine	--	C	NO
deaza cytosine	--	C	NO
N4 (acetyl)cytosine	--	C	NO
1-methyl-1-deaza-pseudoisocytidine	--	C	NO
1-methyl-pseudoisocytidine	--	C	NO
2-methoxy-5-methyl-cytidine	--	C	NO
2-methoxy-cytidine	--	C	NO
2-thio-5-methyl-cytidine	--	C	NO
4-methoxy-1-methyl-pseudoisocytidine	--	C	NO
4-methoxy-pseudoisocytidine	--	C	NO
4-thio-1-methyl-1-deaza-pseudoisocytidine	--	C	NO
4-thio-1-methyl-pseudoisocytidine	--	C	NO
4-thio-pseudoisocytidine	--	C	NO
5-aza-zebularine	--	C	NO
5-methyl-zebularine	--	C	NO
pyrrolo-pseudoisocytidine	--	C	NO
zebularine	--	C	NO
(E)-5-(2-Bromo-vinyl)cytidine TP	--	C	NO
2,2'-anhydro-cytidine TP hydrochloride	--	C	NO
2'Fluor-N4-Bz-cytidine TP	--	C	NO
2'Fluoro-N4-Acetyl-cytidine TP	--	C	NO
2'-O-Methyl-N4-Acetyl-cytidine TP	--	C	NO
2'O-methyl-N4-Bz-cytidine TP	--	C	NO
2'-a-Ethynylcytidine TP	--	C	NO
2'-a-Trifluoromethylcytidine TP	--	C	NO
2'-b-Ethynylcytidine TP	--	C	NO
2'-b-Trifluoromethylcytidine TP	--	C	NO
2'-Deoxy-2',2'-difluorocytidine TP	--	C	NO
2'-Deoxy-2'-a-mercaptopcytidine TP	--	C	NO
2'-Deoxy-2'-a-thiomethoxycytidine TP	--	C	NO
2'-Deoxy-2'-b-aminocytidine TP	--	C	NO
2'-Deoxy-2'-b-azidocytidine TP	--	C	NO
2'-Deoxy-2'-b-bromocytidine TP	--	C	NO

2'-Deoxy-2'-b-chlorocytidine TP	--	C	NO
2'-Deoxy-2'-b-fluorocytidine TP	--	C	NO
2'-Deoxy-2'-b-iodocytidine TP	--	C	NO
2'-Deoxy-2'-b-mercaptopcytidine TP	--	C	NO
2'-Deoxy-2'-b-thiomethoxycytidine TP	--	C	NO
2'-O-Methyl-5-(1-propynyl)cytidine TP	--	C	NO
3'-Ethynylcytidine TP	--	C	NO
4'-Azidocytidine TP	--	C	NO
4'-Carbocyclic cytidine TP	--	C	NO
4'-Ethynylcytidine TP	--	C	NO
5-(1-Propynyl)ara-cytidine TP	--	C	NO
5-(2-Chloro-phenyl)-2-thiocytidine TP	--	C	NO
5-(4-Amino-phenyl)-2-thiocytidine TP	--	C	NO
5-Aminoallyl-CTP	--	C	NO
5-Cyanocytidine TP	--	C	NO
5-Ethynylara-cytidine TP	--	C	NO
5-Ethynylcytidine TP	--	C	NO
5'-Homo-cytidine TP	--	C	NO
5-Methoxycytidine TP	--	C	NO
5-Trifluoromethyl-Cytidine TP	--	C	NO
N4-Amino-cytidine TP	--	C	NO
N4-Benzoyl-cytidine TP	--	C	NO
pseudoisocytidine	--	C	NO
7-methylguanosine	m7G	G	YES
N2,2'-O-dimethylguanosine	m2Gm	G	YES
N2-methylguanosine	m2G	G	YES
wyosine	imG	G	YES
1,2'-O-dimethylguanosine	m1Gm	G	YES
1-methylguanosine	m1G	G	YES
2'-O-methylguanosine	Gm	G	YES
2'-O-ribosylguanosine (phosphate)	Gr(p)	G	YES
2'-O-methylguanosine	Gm	G	YES
2'-O-ribosylguanosine (phosphate)	Gr(p)	G	YES
7-aminomethyl-7-deazaguanosine	preQ1	G	YES
7-cyano-7-deazaguanosine	preQ0	G	YES
archaeosine	G+	G	YES
methylwyosine	mimG	G	YES
N2,7-dimethylguanosine	m2,7G	G	YES
N2,N2,2'-O-trimethylguanosine	m22Gm	G	YES
N2,N2,7-trimethylguanosine	m2,2,7G	G	YES
N2,N2-dimethylguanosine	m22G	G	YES
N ² ,7,2'-O-trimethylguanosine	m ^{2,7} Gm	G	YES
6-thio-guanosine	--	G	NO

7-deaza-guanosine	--	G	NO
8-oxo-guanosine	--	G	NO
N1-methyl-guanosine	--	G	NO
α -thio-guanosine	--	G	NO
2 (propyl)guanine	--	G	NO
2-(alkyl)guanine	--	G	NO
2'-Amino-2'-deoxy-GTP	--	G	NO
2'-Azido-2'-deoxy-GTP	--	G	NO
2'-Deoxy-2'-a-aminoguanosine TP	--	G	NO
2'-Deoxy-2'-a-azidoguanosine TP	--	G	NO
6 (methyl)guanine	--	G	NO
6-(alkyl)guanine	--	G	NO
6-(methyl)guanine	--	G	NO
6-methyl-guanosine	--	G	NO
7 (alkyl)guanine	--	G	NO
7 (deaza)guanine	--	G	NO
7 (methyl)guanine	--	G	NO
7-(alkyl)guanine	--	G	NO
7-(deaza)guanine	--	G	NO
7-(methyl)guanine	--	G	NO
8 (alkyl)guanine	--	G	NO
8 (alkynyl)guanine	--	G	NO
8 (halo)guanine	--	G	NO
8 (thioalkyl)guanine	--	G	NO
8-(alkenyl)guanine	--	G	NO
8-(alkyl)guanine	--	G	NO
8-(alkynyl)guanine	--	G	NO
8-(amino)guanine	--	G	NO
8-(halo)guanine	--	G	NO
8-(hydroxyl)guanine	--	G	NO
8-(thioalkyl)guanine	--	G	NO
8-(thiol)guanine	--	G	NO
aza guanine	--	G	NO
deaza guanine	--	G	NO
N (methyl)guanine	--	G	NO
N-(methyl)guanine	--	G	NO
1-methyl-6-thio-guanosine	--	G	NO
6-methoxy-guanosine	--	G	NO
6-thio-7-deaza-8-aza-guanosine	--	G	NO
6-thio-7-deaza-guanosine	--	G	NO
6-thio-7-methyl-guanosine	--	G	NO
7-deaza-8-aza-guanosine	--	G	NO
7-methyl-8-oxo-guanosine	--	G	NO

N2,N2-dimethyl-6-thio-guanosine	--	G	NO
N2-methyl-6-thio-guanosine	--	G	NO
1-Me-GTP	--	G	NO
2'Fluoro-N2-isobutyl-guanosine TP	--	G	NO
2'O-methyl-N2-isobutyl-guanosine TP	--	G	NO
2'-a-Ethynylguanosine TP	--	G	NO
2'-a-Trifluoromethylguanosine TP	--	G	NO
2'-b-Ethynylguanosine TP	--	G	NO
2'-b-Trifluoromethylguanosine TP	--	G	NO
2'-Deoxy-2',2'-difluoroguanosine TP	--	G	NO
2'-Deoxy-2'-a-mercaptopguanosine TP	--	G	NO
2'-Deoxy-2'-a-thiomethoxyguanosine TP	--	G	NO
2'-Deoxy-2'-b-aminoguanosine TP	--	G	NO
2'-Deoxy-2'-b-azidoguanosine TP	--	G	NO
2'-Deoxy-2'-b-bromoguanosine TP	--	G	NO
2'-Deoxy-2'-b-chloroguanosine TP	--	G	NO
2'-Deoxy-2'-b-fluoroguanosine TP	--	G	NO
2'-Deoxy-2'-b-iodoguanosine TP	--	G	NO
2'-Deoxy-2'-b-mercaptopguanosine TP	--	G	NO
2'-Deoxy-2'-b-thiomethoxyguanosine TP	--	G	NO
4'-Azidoguanosine TP	--	G	NO
4'-Carbocyclic guanosine TP	--	G	NO
4'-Ethynylguanosine TP	--	G	NO
5'-Homo-guanosine TP	--	G	NO
8-bromo-guanosine TP	--	G	NO
9-Deazaguanosine TP	--	G	NO
N2-isobutyl-guanosine TP	--	G	NO
1-methylinosine	m1I	I	YES
inosine	I	I	YES
1,2'-O-dimethylinosine	m1Im	I	YES
2'-O-methylinosine	Im	I	YES
7-methylinosine		I	NO
2'-O-methylinosine	Im	I	YES
epoxyqueuosine	oQ	Q	YES
galactosyl-queuosine	galQ	Q	YES
mannosylqueuosine	manQ	Q	YES
queuosine	Q	Q	YES
allylamino-thymidine	--	T	NO
aza thymidine	--	T	NO
deaza thymidine	--	T	NO
deoxy-thymidine	--	T	NO
2'-O-methyluridine	--	U	YES
2-thiouridine	s2U	U	YES

3-methyluridine	m3U	U	YES
5-carboxymethyluridine	cm5U	U	YES
5-hydroxyuridine	ho5U	U	YES
5-methyluridine	m5U	U	YES
5-taurinomethyl-2-thiouridine	tm5s2U	U	YES
5-taurinomethyluridine	tm5U	U	YES
dihydrouridine	D	U	YES
pseudouridine	Ψ	U	YES
(3-(3-amino-3-carboxypropyl)uridine	acp3U	U	YES
1-methyl-3-(3-amino-5-carboxypropyl)pseudouridine	m1acp3Ψ	U	YES
1-methylpseduouridine	m1Ψ	U	YES
1-methyl-pseudouridine	--	U	YES
2'-O-methyluridine	Um	U	YES
2'-O-methylpseudouridine	Ψm	U	YES
2'-O-methyluridine	Um	U	YES
2-thio-2'-O-methyluridine	s2Um	U	YES
3-(3-amino-3-carboxypropyl)uridine	acp3U	U	YES
3,2'-O-dimethyluridine	m3Um	U	YES
3-Methyl-pseudo-Uridine TP	--	U	YES
4-thiouridine	s4U	U	YES
5-(carboxyhydroxymethyl)uridine	chm5U	U	YES
5-(carboxyhydroxymethyl)uridine methyl ester	mchm5U	U	YES
5,2'-O-dimethyluridine	m5Um	U	YES
5,6-dihydro-uridine	--	U	YES
5-aminomethyl-2-thiouridine	nm5s2U	U	YES
5-carbamoylmethyl-2'-O-methyluridine	ncm5Um	U	YES
5-carbamoylmethyluridine	ncm5U	U	YES
5-carboxyhydroxymethyluridine	--	U	YES
5-carboxyhydroxymethyluridine methyl ester	--	U	YES
5-carboxymethylaminomethyl-2'-O-methyluridine	cmnm5Um	U	YES
5-carboxymethylaminomethyl-2-thiouridine	cmnm5s2U	U	YES
5-carboxymethylaminomethyl-2-thiouridine	--	U	YES
5-carboxymethylaminomethyluridine	cmnm5U	U	YES
5-carboxymethylaminomethyluridine	--	U	YES
5-Carbamoylmethyluridine TP	--	U	YES
5-methoxycarbonylmethyl-2'-O-methyluridine	mcm5Um	U	YES
5-methoxycarbonylmethyl-2-thiouridine	mcm5s2U	U	YES
5-methoxycarbonylmethyluridine	mcm5U	U	YES
5-methoxyuridine	mo5U	U	YES
5-methyl-2-thiouridine	m5s2U	U	YES
5-methylaminomethyl-2-selenouridine	mnm5se2U	U	YES
5-methylaminomethyl-2-thiouridine	mnm5s2U	U	YES

5-methylaminomethyluridine	mm5U	U	YES
5-Methyldihydrouridine	--	U	YES
5-Oxyacetic acid- Uridine TP	--	U	YES
5-Oxyacetic acid-methyl ester-Uridine TP	--	U	YES
N1-methyl-pseudo-uridine	--	U	YES
uridine 5-oxyacetic acid	cm5U	U	YES
uridine 5-oxyacetic acid methyl ester	mcm5U	U	YES
3-(3-Amino-3-carboxypropyl)-Uridine TP	--	U	YES
5-(iso-Pentenylaminomethyl)- 2-thiouridine TP	--	U	YES
5-(iso-Pentenylaminomethyl)-2'-O-methyluridine TP	--	U	YES
5-(iso-Pentenylaminomethyl)uridine TP	--	U	YES
5-propynyl uracil	--	U	NO
α -thio-uridine	--	U	NO
1 (aminoalkylamino-carbonylethylenyl)-2(thio)-pseudouracil	--	U	NO
1 (aminoalkylaminocarbonylethylenyl)-2,4-(dithio)pseudouracil	--	U	NO
1 (aminoalkylaminocarbonylethylenyl)-4(thio)pseudouracil	--	U	NO
1 (aminoalkylaminocarbonylethylenyl)-pseudouracil	--	U	NO
1 (aminocarbonylethylenyl)-2(thio)-pseudouracil	--	U	NO
1 (aminocarbonylethylenyl)-2,4-(dithio)pseudouracil	--	U	NO
1 (aminocarbonylethylenyl)-4(thio)pseudouracil	--	U	NO
1 (aminocarbonylethylenyl)-pseudouracil	--	U	NO
1 substituted 2(thio)-pseudouracil	--	U	NO
1 substituted 2,4-(dithio)pseudouracil	--	U	NO
1 substituted 4 (thio)pseudouracil	--	U	NO
1 substituted pseudouracil	--	U	NO
1-(aminoalkylamino-carbonylethylenyl)-2-(thio)-pseudouracil	--	U	NO
1-Methyl-3-(3-amino-3-carboxypropyl)pseudouridine TP	--	U	NO
1-Methyl-3-(3-amino-3-carboxypropyl)pseudo-UTP	--	U	NO
1-Methyl-pseudo-UTP	--	U	NO
2 (thio)pseudouracil	--	U	NO
2' deoxy uridine	--	U	NO
2' fluorouridine	--	U	NO
2-(thio)uracil	--	U	NO
2,4-(dithio)pseudouracil	--	U	NO
2' methyl, 2' amino, 2' azido, 2' fluoro-guanosine	--	U	NO
2'-Amino-2'-deoxy-UTP	--	U	NO

2'-Azido-2'-deoxy-UTP	--	U	NO
2'-Azido-deoxyuridine TP	--	U	NO
2'-O-methylpseudouridine	--	U	NO
2' deoxy uridine	2' dU	U	NO
2' fluorouridine	--	U	NO
2'-Deoxy-2'-a-aminouridine TP	--	U	NO
2'-Deoxy-2'-a-azidouridine TP	--	U	NO
2-methylpseudouridine	m3Ψ	U	NO
3 (3 amino-3 carboxypropyl)uracil	--	U	NO
4 (thio)pseudouracil	--	U	NO
4-(thio)pseudouracil	--	U	NO
4-(thio)uracil	--	U	NO
4-thiouracil	--	U	NO
5 (1,3-diazole-1-alkyl)uracil	--	U	NO
5 (2-aminopropyl)uracil	--	U	NO
5 (aminoalkyl)uracil	--	U	NO
5 (dimethylaminoalkyl)uracil	--	U	NO
5 (guanidiniumalkyl)uracil	--	U	NO
5 (methoxycarbonylmethyl)-2-(thio)uracil	--	U	NO
5 (methoxycarbonyl-methyl)uracil	--	U	NO
5 (methyl) 2 (thio)uracil	--	U	NO
5 (methyl) 2,4 (dithio)uracil	--	U	NO
5 (methyl) 4 (thio)uracil	--	U	NO
5 (methylaminomethyl)-2 (thio)uracil	--	U	NO
5 (methylaminomethyl)-2,4 (dithio)uracil	--	U	NO
5 (methylaminomethyl)-4 (thio)uracil	--	U	NO
5 (propynyl)uracil	--	U	NO
5 (trifluoromethyl)uracil	--	U	NO
5-(2-aminopropyl)uracil	--	U	NO
5-(alkyl)-2-(thio)pseudouracil	--	U	NO
5-(alkyl)-2,4 (dithio)pseudouracil	--	U	NO
5-(alkyl)-4 (thio)pseudouracil	--	U	NO
5-(alkyl)pseudouracil	--	U	NO
5-(alkyl)uracil	--	U	NO
5-(alkynyl)uracil	--	U	NO
5-(allylamino)uracil	--	U	NO
5-(cyanoalkyl)uracil	--	U	NO
5-(dialkylaminoalkyl)uracil	--	U	NO
5-(dimethylaminoalkyl)uracil	--	U	NO
5-(guanidiniumalkyl)uracil	--	U	NO
5-(halo)uracil	--	U	NO
5-(1,3-diazole-1-alkyl)uracil	--	U	NO
5-(methoxy)uracil	--	U	NO

5-(methoxycarbonylmethyl)-2-(thio)uracil	--	U	NO
5-(methoxycarbonyl-methyl)uracil	--	U	NO
5-(methyl) 2(thio)uracil	--	U	NO
5-(methyl) 2,4 (dithio)uracil	--	U	NO
5-(methyl) 4 (thio)uracil	--	U	NO
5-(methyl)-2-(thio)pseudouracil	--	U	NO
5-(methyl)-2,4 (dithio)pseudouracil	--	U	NO
5-(methyl)-4 (thio)pseudouracil	--	U	NO
5-(methyl)pseudouracil	--	U	NO
5-(methylaminomethyl)-2 (thio)uracil	--	U	NO
5-(methylaminomethyl)-2,4(dithio)uracil	--	U	NO
5-(methylaminomethyl)-4-(thio)uracil	--	U	NO
5-(propynyl)uracil	--	U	NO
5-(trifluoromethyl)uracil	--	U	NO
5-aminoallyl-uridine	--	U	NO
5-bromo-uridine	--	U	NO
5-iodo-uridine	--	U	NO
5-uracil	--	U	NO
6 (azo)uracil	--	U	NO
6-(azo)uracil	--	U	NO
6-aza-uridine	--	U	NO
allyamino-uracil	--	U	NO
aza uracil	--	U	NO
deaza uracil	--	U	NO
N3 (methyl)uracil	--	U	NO
P pseudo-UTP-1-2-ethanoic acid	--	U	NO
pseudouracil	--	U	NO
4-Thio-pseudo-UTP	--	U	NO
1-carboxymethyl-pseudouridine	--	U	NO
1-methyl-1-deaza-pseudouridine	--	U	NO
1-propynyl-uridine	--	U	NO
1-taurinomethyl-1-methyl-uridine	--	U	NO
1-taurinomethyl-4-thio-uridine	--	U	NO
1-taurinomethyl-pseudouridine	--	U	NO
2-methoxy-4-thio-pseudouridine	--	U	NO
2-thio-1-methyl-1-deaza-pseudouridine	--	U	NO
2-thio-1-methyl-pseudouridine	--	U	NO
2-thio-5-aza-uridine	--	U	NO
2-thio-dihydropseudouridine	--	U	NO
2-thio-dihydrouridine	--	U	NO
2-thio-pseudouridine	--	U	NO
4-methoxy-2-thio-pseudouridine	--	U	NO
4-methoxy-pseudouridine	--	U	NO

4-thio-1-methyl-pseudouridine	--	U	NO
4-thio-pseudouridine	--	U	NO
5-aza-uridine	--	U	NO
dihydropseudouridine	--	U	NO
(±)1-(2-Hydroxypropyl)pseudouridine TP	--	U	NO
(2R)-1-(2-Hydroxypropyl)pseudouridine TP	--	U	NO
(2S)-1-(2-Hydroxypropyl)pseudouridine TP	--	U	NO
(E)-5-(2-Bromo-vinyl)ara-uridine TP	--	U	NO
(E)-5-(2-Bromo-vinyl)uridine TP	--	U	NO
(Z)-5-(2-Bromo-vinyl)ara-uridine TP	--	U	NO
(Z)-5-(2-Bromo-vinyl)uridine TP	--	U	NO
1-(2,2,2-Trifluoroethyl)-pseudo-UTP	--	U	NO
1-(2,2,3,3,3-Pentafluoropropyl)pseudouridine TP	--	U	NO
1-(2,2-Diethoxyethyl)pseudouridine TP	--	U	NO
1-(2,4,6-Trimethylbenzyl)pseudouridine TP	--	U	NO
1-(2,4,6-Trimethyl-benzyl)pseudo-UTP	--	U	NO
1-(2,4,6-Trimethyl-phenyl)pseudo-UTP	--	U	NO
1-(2-Amino-2-carboxyethyl)pseudo-UTP	--	U	NO
1-(2-Amino-ethyl)pseudo-UTP	--	U	NO
1-(2-Hydroxyethyl)pseudouridine TP	--	U	NO
1-(2-Methoxyethyl)pseudouridine TP	--	U	NO
1-(3,4-Bis-trifluoromethoxybenzyl)pseudouridine TP	--	U	NO
1-(3,4-Dimethoxybenzyl)pseudouridine TP	--	U	NO
1-(3-Amino-3-carboxypropyl)pseudo-UTP	--	U	NO
1-(3-Amino-propyl)pseudo-UTP	--	U	NO
1-(3-Cyclopropyl-prop-2-ynyl)pseudouridine TP	--	U	NO
1-(4-Amino-4-carboxybutyl)pseudo-UTP	--	U	NO
1-(4-Amino-benzyl)pseudo-UTP	--	U	NO
1-(4-Amino-butyl)pseudo-UTP	--	U	NO
1-(4-Amino-phenyl)pseudo-UTP	--	U	NO
1-(4-Azidobenzyl)pseudouridine TP	--	U	NO
1-(4-Bromobenzyl)pseudouridine TP	--	U	NO
1-(4-Chlorobenzyl)pseudouridine TP	--	U	NO
1-(4-Fluorobenzyl)pseudouridine TP	--	U	NO
1-(4-Iodobenzyl)pseudouridine TP	--	U	NO
1-(4-Methanesulfonylbenzyl)pseudouridine TP	--	U	NO
1-(4-Methoxybenzyl)pseudouridine TP	--	U	NO
1-(4-Methoxy-benzyl)pseudo-UTP	--	U	NO
1-(4-Methoxy-phenyl)pseudo-UTP	--	U	NO
1-(4-Methylbenzyl)pseudouridine TP	--	U	NO
1-(4-Methyl-benzyl)pseudo-UTP	--	U	NO

1-(4-Nitrobenzyl)pseudouridine TP	--	U	NO
1-(4-Nitro-benzyl)pseudo-UTP	--	U	NO
1(4-Nitro-phenyl)pseudo-UTP	--	U	NO
1-(4-Thiomethoxybenzyl)pseudouridine TP	--	U	NO
1-(4-Trifluoromethoxybenzyl)pseudouridine TP	--	U	NO
1-(4-Trifluoromethylbenzyl)pseudouridine TP	--	U	NO
1-(5-Amino-pentyl)pseudo-UTP	--	U	NO
1-(6-Amino-hexyl)pseudo-UTP	--	U	NO
1,6-Dimethyl-pseudo-UTP	--	U	NO
1-[3-(2-{2-[2-(2-Aminoethoxy)-ethoxy]-ethoxy}-ethoxy)-propionyl]pseudouridine TP	--	U	NO
1-{3-[2-(2-Aminoethoxy)-ethoxy]-propionyl } pseudouridine TP	--	U	NO
1-Acetylpsseudouridine TP	--	U	NO
1-Alkyl-6-(1-propynyl)-pseudo-UTP	--	U	NO
1-Alkyl-6-(2-propynyl)-pseudo-UTP	--	U	NO
1-Alkyl-6-allyl-pseudo-UTP	--	U	NO
1-Alkyl-6-ethynyl-pseudo-UTP	--	U	NO
1-Alkyl-6-homoallyl-pseudo-UTP	--	U	NO
1-Alkyl-6-vinyl-pseudo-UTP	--	U	NO
1-Allylpsseudouridine TP	--	U	NO
1-Aminomethyl-pseudo-UTP	--	U	NO
1-Benzoylpsseudouridine TP	--	U	NO
1-Benzyloxymethylpsseudouridine TP	--	U	NO
1-Benzyl-pseudo-UTP	--	U	NO
1-Biotinyl-PEG2-pseudouridine TP	--	U	NO
1-Biotinylpsseudouridine TP	--	U	NO
1-Butyl-pseudo-UTP	--	U	NO
1-Cyanomethylpsseudouridine TP	--	U	NO
1-Cyclobutylmethyl-pseudo-UTP	--	U	NO
1-Cyclobutyl-pseudo-UTP	--	U	NO
1-Cycloheptylmethyl-pseudo-UTP	--	U	NO
1-Cycloheptyl-pseudo-UTP	--	U	NO
1-Cyclohexylmethyl-pseudo-UTP	--	U	NO
1-Cyclohexyl-pseudo-UTP	--	U	NO
1-Cyclooctylmethyl-pseudo-UTP	--	U	NO
1-Cyclooctyl-pseudo-UTP	--	U	NO
1-Cyclopentylmethyl-pseudo-UTP	--	U	NO
1-Cyclopentyl-pseudo-UTP	--	U	NO
1-Cyclopropylmethyl-pseudo-UTP	--	U	NO
1-Cyclopropyl-pseudo-UTP	--	U	NO
1-Ethyl-pseudo-UTP	--	U	NO
1-Hexyl-pseudo-UTP	--	U	NO

1-Homoallylpseudouridine TP	--	U	NO
1-Hydroxymethylpseudouridine TP	--	U	NO
1-iso-propyl-pseudo-UTP	--	U	NO
1-Me-2-thio-pseudo-UTP	--	U	NO
1-Me-4-thio-pseudo-UTP	--	U	NO
1-Me-alpha-thio-pseudo-UTP	--	U	NO
1-Methanesulfonylmethylpseudouridine TP	--	U	NO
1-Methoxymethylpseudouridine TP	--	U	NO
1-Methyl-6-(2,2,2-Trifluoroethyl)pseudo-UTP	--	U	NO
1-Methyl-6-(4-morpholino)-pseudo-UTP	--	U	NO
1-Methyl-6-(4-thiomorpholino)-pseudo-UTP	--	U	NO
1-Methyl-6-(substituted phenyl)pseudo-UTP	--	U	NO
1-Methyl-6-amino-pseudo-UTP	--	U	NO
1-Methyl-6-azido-pseudo-UTP	--	U	NO
1-Methyl-6-bromo-pseudo-UTP	--	U	NO
1-Methyl-6-butyl-pseudo-UTP	--	U	NO
1-Methyl-6-chloro-pseudo-UTP	--	U	NO
1-Methyl-6-cyano-pseudo-UTP	--	U	NO
1-Methyl-6-dimethylamino-pseudo-UTP	--	U	NO
1-Methyl-6-ethoxy-pseudo-UTP	--	U	NO
1-Methyl-6-ethylcarboxylate-pseudo-UTP	--	U	NO
1-Methyl-6-ethyl-pseudo-UTP	--	U	NO
1-Methyl-6-fluoro-pseudo-UTP	--	U	NO
1-Methyl-6-formyl-pseudo-UTP	--	U	NO
1-Methyl-6-hydroxyamino-pseudo-UTP	--	U	NO
1-Methyl-6-hydroxy-pseudo-UTP	--	U	NO
1-Methyl-6-iodo-pseudo-UTP	--	U	NO
1-Methyl-6-iso-propyl-pseudo-UTP	--	U	NO
1-Methyl-6-methoxy-pseudo-UTP	--	U	NO
1-Methyl-6-methylamino-pseudo-UTP	--	U	NO
1-Methyl-6-phenyl-pseudo-UTP	--	U	NO
1-Methyl-6-propyl-pseudo-UTP	--	U	NO
1-Methyl-6-tert-butyl-pseudo-UTP	--	U	NO
1-Methyl-6-trifluoromethoxy-pseudo-UTP	--	U	NO
1-Methyl-6-trifluoromethyl-pseudo-UTP	--	U	NO
1-Morpholinomethylpseudouridine TP	--	U	NO
1-Pentyl-pseudo-UTP	--	U	NO
1-Phenyl-pseudo-UTP	--	U	NO
1-Pivaloylpseudouridine TP	--	U	NO
1-Propargylpseudouridine TP	--	U	NO
1-Propyl-pseudo-UTP	--	U	NO
1-propynyl-pseudouridine	--	U	NO
1-p-tolyl-pseudo-UTP	--	U	NO

1-tert-Butyl-pseudo-UTP	--	U	NO
1-Thiomethoxymethylpseudouridine TP	--	U	NO
1-Thiomorpholinomethylpseudouridine TP	--	U	NO
1-Trifluoroacetylpsudouridine TP	--	U	NO
1-Trifluoromethyl-pseudo-UTP	--	U	NO
1-Vinylpseudouridine TP	--	U	NO
2,2'-anhydro-uridine TP	--	U	NO
2'-bromo-deoxyuridine TP	--	U	NO
2'-F-5-Methyl-2'-deoxy-UTP	--	U	NO
2'-OMe-5-Me-UTP	--	U	NO
2'-OMe-pseudo-UTP	--	U	NO
2'-a-Ethynyluridine TP	--	U	NO
2'-a-Trifluoromethyluridine TP	--	U	NO
2'-b-Ethynyluridine TP	--	U	NO
2'-b-Trifluoromethyluridine TP	--	U	NO
2'-Deoxy-2',2'-difluorouridine TP	--	U	NO
2'-Deoxy-2'-a-mercaptopuridine TP	--	U	NO
2'-Deoxy-2'-a-thiomethoxyuridine TP	--	U	NO
2'-Deoxy-2'-b-aminouridine TP	--	U	NO
2'-Deoxy-2'-b-azidouridine TP	--	U	NO
2'-Deoxy-2'-b-bromouridine TP	--	U	NO
2'-Deoxy-2'-b-chlorouridine TP	--	U	NO
2'-Deoxy-2'-b-fluorouridine TP	--	U	NO
2'-Deoxy-2'-b-iodouridine TP	--	U	NO
2'-Deoxy-2'-b-mercaptopuridine TP	--	U	NO
2'-Deoxy-2'-b-thiomethoxyuridine TP	--	U	NO
2-methoxy-4-thio-uridine	--	U	NO
2-methoxyuridine	--	U	NO
2'-O-Methyl-5-(1-propynyl)uridine TP	--	U	NO
3-Alkyl-pseudo-UTP	--	U	NO
4'-Azidouridine TP	--	U	NO
4'-Carbocyclic uridine TP	--	U	NO
4'-Ethynyluridine TP	--	U	NO
5-(1-Propynyl)ara-uridine TP	--	U	NO
5-(2-Furanyl)uridine TP	--	U	NO
5-Cyanouridine TP	--	U	NO
5-Dimethylaminouridine TP	--	U	NO
5'-Homo-uridine TP	--	U	NO
5-iodo-2'-fluoro-deoxyuridine TP	--	U	NO
5-Phenylethynyluridine TP	--	U	NO
5-Trideuteromethyl-6-deuterouridine TP	--	U	NO
5-Trifluoromethyl-Uridine TP	--	U	NO
5-Vinylarauridine TP	--	U	NO

6-(2,2,2-Trifluoroethyl)-pseudo-UTP	--	U	NO
6-(4-Morpholino)-pseudo-UTP	--	U	NO
6-(4-Thiomorpholino)-pseudo-UTP	--	U	NO
6-(Substituted-Phenyl)-pseudo-UTP	--	U	NO
6-Amino-pseudo-UTP	--	U	NO
6-Azido-pseudo-UTP	--	U	NO
6-Bromo-pseudo-UTP	--	U	NO
6-Butyl-pseudo-UTP	--	U	NO
6-Chloro-pseudo-UTP	--	U	NO
6-Cyano-pseudo-UTP	--	U	NO
6-Dimethylamino-pseudo-UTP	--	U	NO
6-Ethoxy-pseudo-UTP	--	U	NO
6-Ethylcarboxylate-pseudo-UTP	--	U	NO
6-Ethyl-pseudo-UTP	--	U	NO
6-Fluoro-pseudo-UTP	--	U	NO
6-Formyl-pseudo-UTP	--	U	NO
6-Hydroxyamino-pseudo-UTP	--	U	NO
6-Hydroxy-pseudo-UTP	--	U	NO
6-Iodo-pseudo-UTP	--	U	NO
6-iso-Propyl-pseudo-UTP	--	U	NO
6-Methoxy-pseudo-UTP	--	U	NO
6-Methylamino-pseudo-UTP	--	U	NO
6-Methyl-pseudo-UTP	--	U	NO
6-Phenyl-pseudo-UTP	--	U	NO
6-Phenyl-pseudo-UTP	--	U	NO
6-Propyl-pseudo-UTP	--	U	NO
6-tert-Butyl-pseudo-UTP	--	U	NO
6-Trifluoromethoxy-pseudo-UTP	--	U	NO
6-Trifluoromethyl-pseudo-UTP	--	U	NO
Alpha-thio-pseudo-UTP	--	U	NO
Pseudouridine 1-(4-methylbenzenesulfonic acid) TP	--	U	NO
Pseudouridine 1-(4-methylbenzoic acid) TP	--	U	NO
Pseudouridine TP 1-[3-(2-ethoxy)]propionic acid	--	U	NO
Pseudouridine TP 1-[3-{2-(2-[2-(2-ethoxy)-ethoxy]-ethoxy)-ethoxy}]propionic acid	--	U	NO
Pseudouridine TP 1-[3-{2-(2-[2-{2(2-ethoxy)-ethoxy}-ethoxy]-ethoxy)-ethoxy}]propionic acid	--	U	NO
Pseudouridine TP 1-[3-{2-(2-[2-ethoxy]-ethoxy)-ethoxy}]propionic acid	--	U	NO
Pseudouridine TP 1-[3-{2-(2-ethoxy)-ethoxy}]propionic acid	--	U	NO
Pseudouridine TP 1-methylphosphonic acid	--	U	NO

Pseudouridine TP 1-methylphosphonic acid diethyl ester	--	U	NO
Pseudo-UTP-N1-3-propionic acid	--	U	NO
Pseudo-UTP-N1-4-butanoic acid	--	U	NO
Pseudo-UTP-N1-5-pentanoic acid	--	U	NO
Pseudo-UTP-N1-6-hexanoic acid	--	U	NO
Pseudo-UTP-N1-7-heptanoic acid	--	U	NO
Pseudo-UTP-N1-methyl-p-benzoic acid	--	U	NO
Pseudo-UTP-N1-p-benzoic acid	--	U	NO
wybutosine	yW	W	YES
hydroxywybutosine	OHyW	W	YES
isowyosine	imG2	W	YES
peroxywybutosine	o2yW	W	YES
undermodified hydroxywybutosine	OHyW*	W	YES
4-demethylwyosine	imG-14	W	YES

[000320] Other modifications which may be useful in the polynucleotides of the present invention are listed in Table 6.

Table 6. Additional Modification types

Name	Type
2,6-(diamino)purine	Other
1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl	Other
1,3-(diaza)-2-(oxo)-phenthiazin-1-yl	Other
1,3-(diaz)-2-(oxo)-phenoxazin-1-yl	Other
1,3,5-(triaz)-2,6-(diox)-naphthalene	Other
2 (amino)purine	Other
2,4,5-(trimethyl)phenyl	Other
2' methyl, 2' amino, 2' azido, 2' fluoro-cytidine	Other
2' methyl, 2' amino, 2' azido, 2' fluoro-adenine	Other
2' methyl, 2' amino, 2' azido, 2' fluoro-uridine	Other
2'-amino-2'-deoxyribose	Other
2-amino-6-Chloro-purine	Other
2-aza-inosinyl	Other
2'-azido-2'-deoxyribose	Other
2'fluoro-2'-deoxyribose	Other
2'-fluoro-modified bases	Other
2'-O-methyl-ribose	Other
2-oxo-7-aminopyridopyrimidin-3-yl	Other
2-oxo-pyridopyrimidine-3-yl	Other
2-pyridinone	Other
3 nitropyrrole	Other
3-(methyl)-7-(propynyl)isocarbostyryl	Other

3-(methyl)isocarbostyryl	Other
4-(fluoro)-6-(methyl)benzimidazole	Other
4-(methyl)benzimidazole	Other
4-(methyl)indolyl	Other
4,6-(dimethyl)indolyl	Other
5 nitroindole	Other
5 substituted pyrimidines	Other
5-(methyl)isocarbostyryl	Other
5-nitroindole	Other
6-(aza)pyrimidine	Other
6-(azo)thymine	Other
6-(methyl)-7-(aza)indolyl	Other
6-chloro-purine	Other
6-phenyl-pyrrolo-pyrimidin-2-on-3-yl	Other
7-(aminoalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenthiazin-1-yl	Other
7-(aminoalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl	Other
7-(aminoalkylhydroxy)-1,3-(diaz)-2-(oxo)-phenoxazin-1-yl	Other
7-(aminoalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenthiazin-1-yl	Other
7-(aminoalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenoxazin-1-yl	Other
7-(aza)indolyl	Other
7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl	Other
7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenthiazin-1-yl	Other
7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl	Other
7-(guanidiniumalkylhydroxy)-1,3-(diaz)-2-(oxo)-phenoxazin-1-yl	Other
7-(guanidiniumalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenthiazin-1-yl	Other
7-(guanidiniumalkylhydroxy)-1,3-(diaz)-2-(oxo)-phenoxazin-1-yl	Other
7-(propynyl)isocarbostyryl	Other
7-(propynyl)isocarbostyryl, propynyl-7-(aza)indolyl	Other
7-deaza-inosinyl	Other
7-substituted 1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl	Other
7-substituted 1,3-(diaz)-2-(oxo)-phenoxazin-1-yl	Other
9-(methyl)-imidizopyridinyl	Other
aminoindolyl	Other
anthracenyl	Other
bis-ortho-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl	Other
bis-ortho-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl	Other
difluorotolyl	Other
hypoxanthine	Other
imidizopyridinyl	Other
inosinyl	Other
isocarbostyryl	Other
isoguanisine	Other
N2-substituted purines	Other

N6-methyl-2-amino-purine	Other
N6-substituted purines	Other
N-alkylated derivative	Other
naphthalenyl	Other
nitrobenzimidazolyl	Other
nitroimidazolyl	Other
nitroindazolyl	Other
nitropyrazolyl	Other
nubularine	Other
O6-substituted purines	Other
O-alkylated derivative	Other
ortho-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl	Other
ortho-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl	Other
Oxoformycin TP	Other
para-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl	Other
para-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl	Other
pentacenyl	Other
phenanthracenyl	Other
phenyl	Other
propynyl-7-(aza)indolyl	Other
pyrenyl	Other
pyridopyrimidin-3-yl	Other
pyridopyrimidin-3-yl, 2-oxo-7-amino-pyridopyrimidin-3-yl	Other
pyrrolo-pyrimidin-2-on-3-yl	Other
pyrrolopyrimidinyl	Other
pyrrolopyrizinyl	Other
stilbenzyl	Other
substituted 1,2,4-triazoles	Other
tetracenyl	Other
tubercidine	Other
xanthine	Other
Xanthosine-5'-TP	Other
2-thio-zebularine	Other
5-aza-2-thio-zebularine	Other
7-deaza-2-amino-purine	Other
pyridin-4-one ribonucleoside	Other
2-Amino-riboside-TP	Other
Formycin A TP	Other
Formycin B TP	Other
Pyrrolosine TP	Other
2'-OH-ara-adenosine TP	Other
2'-OH-ara-cytidine TP	Other
2'-OH-ara-uridine TP	Other

2'-OH-ara-guanosine TP	Other
5-(2-carbomethoxyvinyl)uridine TP	Other
N6-(19-Amino-pentaoxanonadecyl)adenosine TP	Other

[000321] The polynucleotides can include any useful linker between the nucleosides. Such linkers, including backbone modifications are given in Table 7.

Table 7. Linker modifications

Name	TYPE
3'-alkylene phosphonates	Linker
3'-amino phosphoramidate	Linker
alkene containing backbones	Linker
aminoalkylphosphoramidates	Linker
aminoalkylphosphotriesters	Linker
boranophosphates	Linker
-CH ₂ -O-N(CH ₃)-CH ₂ -	Linker
-CH ₂ -N(CH ₃)-N(CH ₃)-CH ₂ -	Linker
-CH ₂ -NH-CH ₂ -	Linker
chiral phosphonates	Linker
chiral phosphorothioates	Linker
formacetyl and thioformacetyl backbones	Linker
methylene (methylimino)	Linker
methylene formacetyl and thioformacetyl backbones	Linker
methyleneimino and methylenehydrazino backbones	Linker
morpholino linkages	Linker
-N(CH ₃)-CH ₂ -CH ₂ -	Linker
oligonucleosides with heteroatom internucleoside linkage	Linker
phosphinates	Linker
phosphoramidates	Linker
phosphorodithioates	Linker
phosphorothioate internucleoside linkages	Linker
phosphorothioates	Linker
phosphotriesters	Linker
PNA	Linker
siloxane backbones	Linker
sulfamate backbones	Linker
sulfide sulfoxide and sulfone backbones	Linker
sulfonate and sulfonamide backbones	Linker
thionoalkylphosphonates	Linker
thionoalkylphosphotriesters	Linker

thionophosphoramidates	Linker
------------------------	--------

[000322] The polynucleotides can include any useful modification, such as to the sugar, the nucleobase, or the internucleoside linkage (*e.g.* to a linking phosphate / to a phosphodiester linkage / to the phosphodiester backbone). One or more atoms of a pyrimidine nucleobase may be replaced or substituted with optionally substituted amino, optionally substituted thiol, optionally substituted alkyl (*e.g.*, methyl or ethyl), or halo (*e.g.*, chloro or fluoro). In certain embodiments, modifications (*e.g.*, one or more modifications) are present in each of the sugar and the internucleoside linkage. Modifications according to the present invention may be modifications of ribonucleic acids (RNAs) to deoxyribonucleic acids (DNAs), threose nucleic acids (TNAs), glycol nucleic acids (GNAs), peptide nucleic acids (PNAs), locked nucleic acids (LNAs) or hybrids thereof). Additional modifications are described herein.

[000323] In some embodiments, the polynucleotides of the invention do not substantially induce an innate immune response of a cell into which the mRNA is introduced. Features of an induced innate immune response include 1) increased expression of pro-inflammatory cytokines, 2) activation of intracellular PRRs (RIG-I, MDA5, etc., and/or 3) termination or reduction in protein translation.

[000324] In certain embodiments, it may be desirable to intracellularly degrade a polynucleotide introduced into the cell. For example, degradation of a polynucleotide may be preferable if precise timing of protein production is desired. Thus, in some embodiments, the invention provides a polynucleotide containing a degradation domain, which is capable of being acted on in a directed manner within a cell.

[000325] Any of the regions of the polynucleotides may be chemically modified as taught herein or as taught in International Publication Number WO2013052523 (Attorney Docket Number M9) and International Publication Number WO2014093924 (Attorney Docket Number M36) the contents of each of which are incorporated herein by reference in its entirety.

Modified Polynucleotide Molecules

[000326] The present invention also includes building blocks, *e.g.*, modified ribonucleotides, and modified ribonucleotides, of polynucleotide molecules. For

example, these building blocks can be useful for preparing the polynucleotides of the invention. Such building blocks are taught in International Publication Number WO2013052523 (Attorney Docket Number M9) and International Publication Number WO2014093924 (Attorney Docket Number M36) the contents of each of which are incorporated herein by reference in its entirety.

Modifications on the Sugar

[000327] The modified nucleosides and nucleotides (e.g., building block molecules), which may be incorporated into a polynucleotide (e.g., RNA or mRNA, as described herein), can be modified on the sugar of the ribonucleic acid. For example, the 2' hydroxyl group (OH) can be modified or replaced with a number of different substituents. Exemplary substitutions at the 2'-position include, but are not limited to, H, halo, optionally substituted C₁₋₆ alkyl; optionally substituted C₁₋₆ alkoxy; optionally substituted C₆₋₁₀ aryloxy; optionally substituted C₃₋₈ cycloalkyl; optionally substituted C₃₋₈ cycloalkoxy; optionally substituted C₆₋₁₀ aryloxy; optionally substituted C₆₋₁₀ aryl-C₁₋₆ alkoxy, optionally substituted C₁₋₁₂ (heterocyclyl)oxy; a sugar (e.g., ribose, pentose, or any described herein); a polyethyleneglycol (PEG), -O(CH₂CH₂O)_nCH₂CH₂OR, where R is H or optionally substituted alkyl, and n is an integer from 0 to 20 (e.g., from 0 to 4, from 0 to 8, from 0 to 10, from 0 to 16, from 1 to 4, from 1 to 8, from 1 to 10, from 1 to 16, from 1 to 20, from 2 to 4, from 2 to 8, from 2 to 10, from 2 to 16, from 2 to 20, from 4 to 8, from 4 to 10, from 4 to 16, and from 4 to 20); "locked" nucleic acids (LNA) in which the 2'-hydroxyl is connected by a C₁₋₆ alkylene or C₁₋₆ heteroalkylene bridge to the 4'-carbon of the same ribose sugar, where exemplary bridges included methylene, propylene, ether, or amino bridges; aminoalkyl, as defined herein; aminoalkoxy, as defined herein; amino as defined herein; and amino acid, as defined herein

[000328] Generally, RNA includes the sugar group ribose, which is a 5-membered ring having an oxygen. Exemplary, non-limiting modified nucleotides include replacement of the oxygen in ribose (e.g., with S, Se, or alkylene, such as methylene or ethylene); addition of a double bond (e.g., to replace ribose with cyclopentenyl or cyclohexenyl); ring contraction of ribose (e.g., to form a 4-membered ring of cyclobutane or oxetane); ring expansion of ribose (e.g., to form a 6- or 7-membered ring having an additional carbon or heteroatom, such as for anhydrohexitol, altritol, mannitol, cyclohexanyl,

cyclohexenyl, and morpholino that also has a phosphoramidate backbone); multicyclic forms (e.g., tricyclo; and “unlocked” forms, such as glycol nucleic acid (GNA) (e.g., R-GNA or S-GNA, where ribose is replaced by glycol units attached to phosphodiester bonds), threose nucleic acid (TNA, where ribose is replaced with α -L-threofuranosyl-(3'→2')) , and peptide nucleic acid (PNA, where 2-amino-ethyl-glycine linkages replace the ribose and phosphodiester backbone). The sugar group can also contain one or more carbons that possess the opposite stereochemical configuration than that of the corresponding carbon in ribose. Thus, a polynucleotide molecule can include nucleotides containing, e.g., arabinose, as the sugar. Such sugar modifications are taught International Publication No. WO2013052523 (Attorney Docket Number M9) and International Publication No. WO2014093924 (Attorney Docket Number M36) the contents of each of which are incorporated herein by reference in its entirety.

Modifications on the Nucleobase

[000329] The present disclosure provides for modified nucleosides and nucleotides. As described herein “nucleoside” is defined as a compound containing a sugar molecule (e.g., a pentose or ribose) or a derivative thereof in combination with an organic base (e.g., a purine or pyrimidine) or a derivative thereof (also referred to herein as “nucleobase”). As described herein, “nucleotide” is defined as a nucleoside including a phosphate group. The modified nucleotides may be synthesized by any useful method, as described herein (e.g., chemically, enzymatically, or recombinantly to include one or more modified or non-natural nucleosides). The polynucleotides may comprise a region or regions of linked nucleosides. Such regions may have variable backbone linkages. The linkages may be standard phosphoester linkages, in which case the polynucleotides would comprise regions of nucleotides.

[000330] The modified nucleotide base pairing encompasses not only the standard adenosine-thymine, adenosine-uracil, or guanosine-cytosine base pairs, but also base pairs formed between nucleotides and/or modified nucleotides comprising non-standard or modified bases, wherein the arrangement of hydrogen bond donors and hydrogen bond acceptors permits hydrogen bonding between a non-standard base and a standard base or between two complementary non-standard base structures. One example of such non-

standard base pairing is the base pairing between the modified nucleotide inosine and adenine, cytosine or uracil.

[000331] The modified nucleosides and nucleotides can include a modified nucleobase. Examples of nucleobases found in RNA include, but are not limited to, adenine, guanine, cytosine, and uracil. Examples of nucleobase found in DNA include, but are not limited to, adenine, guanine, cytosine, and thymine. Such modified nucleobases (including the distinctions between naturally occurring and non-naturally occurring) are taught in International Publication No. WO2013052523 (Attorney Docket Number M9) and International Publication No. WO2014093924 (Attorney Docket Number M36) the contents of each of which are incorporated herein by reference in its entirety.

Combinations of Modified Sugars, Nucleobases, and Internucleoside Linkages

[000332] The polynucleotides of the invention can include a combination of modifications to the sugar, the nucleobase, and/or the internucleoside linkage. These combinations can include any one or more modifications described herein.

[000333] Examples of modified nucleotides and modified nucleotide combinations are provided below in Tables 8 and 9. These combinations of modified nucleotides can be used to form the polynucleotides of the invention. Unless otherwise noted, the modified nucleotides may be completely substituted for the natural nucleotides of the polynucleotides of the invention. As a non-limiting example, the natural nucleotide uridine may be substituted with a modified nucleoside described herein. In another non-limiting example, the natural nucleotide uridine may be partially substituted (e.g., about 0.1%, 1%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 99.9%) with at least one of the modified nucleoside disclosed herein. Any combination of base/sugar or linker may be incorporated into the polynucleotides of the invention and such modifications are taught in International Publication Number WO2013052523 (Attorney Docket Number M9) and in International Publication Number WO2014093924 (Attorney Docket Number M36) the contents of each of which are incorporated herein by reference in its entirety.

Table 8. Combinations

<u>Modified Nucleotide</u>	<u>Modified Nucleotide Combination</u>
α -thio-cytidine	α -thio-cytidine/5-iodo-uridine
	α -thio-cytidine/N1-methyl-pseudouridine

	α -thio-cytidine/ α -thio-uridine
	α -thio-cytidine/5-methyl-uridine
	α -thio-cytidine/pseudo-uridine
	about 50% of the cytosines are α -thio-cytidine
pseudoisocytidine	pseudoisocytidine/5-iodo-uridine
	pseudoisocytidine/N1-methyl-pseudouridine
	pseudoisocytidine/ α -thio-uridine
	pseudoisocytidine/5-methyl-uridine
	pseudoisocytidine/pseudouridine
	about 25% of cytosines are pseudoisocytidine
	pseudoisocytidine/about 50% of uridines are N1-methyl-pseudouridine and about 50% of uridines are pseudouridine
pyrrolo-cytidine	pseudoisocytidine/about 25% of uridines are N1-methyl-pseudouridine and about 25% of uridines are pseudouridine
	pyrrolo-cytidine/5-iodo-uridine
	pyrrolo-cytidine/N1-methyl-pseudouridine
	pyrrolo-cytidine/ α -thio-uridine
	pyrrolo-cytidine/5-methyl-uridine
	pyrrolo-cytidine/pseudouridine
5-methyl-cytidine	about 50% of the cytosines are pyrrolo-cytidine
	5-methyl-cytidine/5-iodo-uridine
	5-methyl-cytidine/N1-methyl-pseudouridine
	5-methyl-cytidine/ α -thio-uridine
	5-methyl-cytidine/5-methyl-uridine
	5-methyl-cytidine/pseudouridine
	about 25% of cytosines are 5-methyl-cytidine
	about 50% of cytosines are 5-methyl-cytidine
	5-methyl-cytidine/5-methoxy-uridine
	5-methyl-cytidine/5-bromo-uridine
	5-methyl-cytidine/2-thio-uridine
	5-methyl-cytidine/about 50% of uridines are 2-thio-uridine
	about 50% of uridines are 5-methyl-cytidine/ about 50% of uridines are 2-thio-uridine
N4-acetyl-cytidine	N4-acetyl-cytidine /5-iodo-uridine
	N4-acetyl-cytidine /N1-methyl-pseudouridine
	N4-acetyl-cytidine / α -thio-uridine
	N4-acetyl-cytidine /5-methyl-uridine
	N4-acetyl-cytidine /pseudouridine
	about 50% of cytosines are N4-acetyl-cytidine
	about 25% of cytosines are N4-acetyl-cytidine
	N4-acetyl-cytidine /5-methoxy-uridine
	N4-acetyl-cytidine /5-bromo-uridine
	N4-acetyl-cytidine /2-thio-uridine
	about 50% of cytosines are N4-acetyl-cytidine/ about 50% of uridines are 2-thio-uridine

Table 9. Combinations

1-(2,2,2-Trifluoroethyl)pseudo-UTP
1-Ethyl-pseudo-UTP
1-Methyl-pseudo-U-alpha-thio-TP
1-methyl-pseudouridine TP, ATP, GTP, CTP
1-methyl-pseudo-UTP/5-methyl-CTP/ATP/GTP
1-methyl-pseudo-UTP/CTP/ATP/GTP
1-Propyl-pseudo-UTP
25 % 5-Aminoallyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Aminoallyl-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Bromo-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Bromo-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Bromo-CTP + 75 % CTP/1-Methyl-pseudo-UTP
25 % 5-Carboxy-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Carboxy-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Ethyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Ethyl-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Ethynyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Ethynyl-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Fluoro-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Fluoro-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Formyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Formyl-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Hydroxymethyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Hydroxymethyl-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Iodo-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Iodo-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Methoxy-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Methoxy-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Methyl-CTP + 75 % CTP/25 % 5-Methoxy-UTP + 75 % 1-Methyl-pseudo-UTP
25 % 5-Methyl-CTP + 75 % CTP/25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Methyl-CTP + 75 % CTP/50 % 5-Methoxy-UTP + 50 % 1-Methyl-pseudo-UTP
25 % 5-Methyl-CTP + 75 % CTP/50 % 5-Methoxy-UTP + 50 % UTP
25 % 5-Methyl-CTP + 75 % CTP/5-Methoxy-UTP

25 % 5-Methyl-CTP + 75 % CTP/75 % 5-Methoxy-UTP + 25 % 1-Methyl-pseudo-UTP
25 % 5-Methyl-CTP + 75 % CTP/75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Phenyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Phenyl-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Trifluoromethyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % 5-Trifluoromethyl-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % 5-Trifluoromethyl-CTP + 75 % CTP/1-Methyl-pseudo-UTP
25 % N4-Ac-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % N4-Ac-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % N4-Bz-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % N4-Bz-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % N4-Methyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % N4-Methyl-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25 % Pseudo-iso-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
25 % Pseudo-iso-CTP + 75 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
25% 5-Bromo-CTP/75% CTP/ Pseudo-UTP
25% 5-methoxy-UTP/25% 5-methyl-CTP/ATP/GTP
25% 5-methoxy-UTP/5-methyl-CTP/ATP/GTP
25% 5-methoxy-UTP/75% 5-methyl-CTP/ATP/GTP
25% 5-methoxy-UTP/CTP/ATP/GTP
25% 5-methoxy-UTP/50% 5-methyl-CTP/ATP/GTP
2-Amino-ATP
2-Thio-CTP
2-thio-pseudouridine TP, ATP, GTP, CTP
2-Thio-pseudo-UTP
2-Thio-UTP
3-Methyl-CTP
3-Methyl-pseudo-UTP
4-Thio-UTP
50 % 5-Bromo-CTP + 50 % CTP/1-Methyl-pseudo-UTP
50 % 5-Hydroxymethyl-CTP + 50 % CTP/1-Methyl-pseudo-UTP
50 % 5-methoxy-UTP/5-methyl-CTP/ATP/GTP
50 % 5-Methyl-CTP + 50 % CTP/25 % 5-Methoxy-UTP + 75 % 1-Methyl-pseudo-UTP
50 % 5-Methyl-CTP + 50 % CTP/25 % 5-Methoxy-UTP + 75 % UTP

50 % 5-Methyl-CTP + 50 % CTP/50 % 5-Methoxy-UTP + 50 % 1-Methyl-pseudo-UTP
50 % 5-Methyl-CTP + 50 % CTP/50 % 5-Methoxy-UTP + 50 % UTP
50 % 5-Methyl-CTP + 50 % CTP/5-Methoxy-UTP
50 % 5-Methyl-CTP + 50 % CTP/75 % 5-Methoxy-UTP + 25 % 1-Methyl-pseudo-UTP
50 % 5-Methyl-CTP + 50 % CTP/75 % 5-Methoxy-UTP + 25 % UTP
50 % 5-Trifluoromethyl-CTP + 50 % CTP/1-Methyl-pseudo-UTP
50% 5-Bromo-CTP/ 50% CTP/Pseudo-UTP
50% 5-methoxy-UTP/25% 5-methyl-CTP/ATP/GTP
50% 5-methoxy-UTP/50% 5-methyl-CTP/ATP/GTP
50% 5-methoxy-UTP/75% 5-methyl-CTP/ATP/GTP
50% 5-methoxy-UTP/CTP/ATP/GTP
5-Aminoallyl-CTP
5-Aminoallyl-CTP/ 5-Methoxy-UTP
5-Aminoallyl-UTP
5-Bromo-CTP
5-Bromo-CTP/ 5-Methoxy-UTP
5-Bromo-CTP/1-Methyl-pseudo-UTP
5-Bromo-CTP/Pseudo-UTP
5-bromocytidine TP, ATP, GTP, UTP
5-Bromo-UTP
5-Carboxy-CTP/ 5-Methoxy-UTP
5-Ethyl-CTP/5-Methoxy-UTP
5-Ethynyl-CTP/5-Methoxy-UTP
5-Fluoro-CTP/ 5-Methoxy-UTP
5-Formyl-CTP/ 5-Methoxy-UTP
5-Hydroxy- methyl-CTP/ 5-Methoxy-UTP
5-Hydroxymethyl-CTP
5-Hydroxymethyl-CTP/1-Methyl-pseudo-UTP
5-Hydroxymethyl-CTP/5-Methoxy-UTP
5-hydroxymethyl-cytidine TP, ATP, GTP, UTP
5-Iodo-CTP/ 5-Methoxy-UTP
5-Me-CTP/5-Methoxy-UTP
5-Methoxy carbonyl methyl-UTP
5-Methoxy-CTP/5-Methoxy-UTP

5-methoxy-uridine TP, ATP, GTP, UTP
5-methoxy-UTP
5-Methoxy-UTP
5-Methoxy-UTP/ N6-Isopentenyl-ATP
5-methoxy-UTP/25% 5-methyl-CTP/ATP/GTP
5-methoxy-UTP/5-methyl-CTP/ATP/GTP
5-methoxy-UTP/75% 5-methyl-CTP/ATP/GTP
5-methoxy-UTP/CTP/ATP/GTP
5-Methyl-2-thio-UTP
5-Methylaminomethyl-UTP
5-Methyl-CTP/ 5-Methoxy-UTP
5-Methyl-CTP/ 5-Methoxy-UTP(cap 0)
5-Methyl-CTP/ 5-Methoxy-UTP(No cap)
5-Methyl-CTP/25 % 5-Methoxy-UTP + 75 % 1-Methyl-pseudo-UTP
5-Methyl-CTP/25 % 5-Methoxy-UTP + 75 % UTP
5-Methyl-CTP/50 % 5-Methoxy-UTP + 50 % 1-Methyl-pseudo-UTP
5-Methyl-CTP/50 % 5-Methoxy-UTP + 50 % UTP
5-Methyl-CTP/5-Methoxy-UTP/N6-Me-ATP
5-Methyl-CTP/75 % 5-Methoxy-UTP + 25 % 1-Methyl-pseudo-UTP
5-Methyl-CTP/75 % 5-Methoxy-UTP + 25 % UTP
5-Phenyl-CTP/ 5-Methoxy-UTP
5-Trifluoro- methyl-CTP/ 5-Methoxy-UTP
5-Trifluoromethyl-CTP
5-Trifluoromethyl-CTP/ 5-Methoxy-UTP
5-Trifluoromethyl-CTP/1-Methyl-pseudo-UTP
5-Trifluoromethyl-CTP/Pseudo-UTP
5-Trifluoromethyl-UTP
5-trifluoromethylcytidine TP, ATP, GTP, UTP
75 % 5-Aminoallyl-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Aminoallyl-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Bromo-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Bromo-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Carboxy-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Carboxy-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP

75 % 5-Ethyl-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Ethyl-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Ethynyl-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Ethynyl-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Fluoro-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Fluoro-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Formyl-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Formyl-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Hydroxymethyl-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Hydroxymethyl-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Iodo-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Iodo-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Methoxy-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Methoxy-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-methoxy-UTP/5-methyl-CTP/ATP/GTP
75 % 5-Methyl-CTP + 25 % CTP/25 % 5-Methoxy-UTP + 75 % 1-Methyl-pseudo-UTP
75 % 5-Methyl-CTP + 25 % CTP/25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Methyl-CTP + 25 % CTP/50 % 5-Methoxy-UTP + 50 % 1-Methyl-pseudo-UTP
75 % 5-Methyl-CTP + 25 % CTP/50 % 5-Methoxy-UTP + 50 % UTP
75 % 5-Methyl-CTP + 25 % CTP/5-Methoxy-UTP
75 % 5-Methyl-CTP + 25 % CTP/75 % 5-Methoxy-UTP + 25 % 1-Methyl-pseudo-UTP
75 % 5-Methyl-CTP + 25 % CTP/75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Phenyl-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Phenyl-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Trifluoromethyl-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % 5-Trifluoromethyl-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % 5-Trifluoromethyl-CTP + 25 % CTP/1-Methyl-pseudo-UTP
75 % N4-Ac-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % N4-Ac-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % N4-Bz-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % N4-Bz-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % N4-Methyl-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP
75 % N4-Methyl-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75 % Pseudo-iso-CTP + 25 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP

75 % Pseudo-iso-CTP + 25 % CTP/ 75 % 5-Methoxy-UTP + 25 % UTP
75% 5-Bromo-CTP/25% CTP/ 1-Methyl-pseudo-UTP
75% 5-Bromo-CTP/25% CTP/ Pseudo-UTP
75% 5-methoxy-UTP/25% 5-methyl-CTP/ATP/GTP
75% 5-methoxy-UTP/50% 5-methyl-CTP/ATP/GTP
75% 5-methoxy-UTP/75% 5-methyl-CTP/ATP/GTP
75% 5-methoxy-UTP/CTP/ATP/GTP
8-Aza-ATP
Alpha-thio-CTP
CTP/25 % 5-Methoxy-UTP + 75 % 1-Methyl-pseudo-UTP
CTP/25 % 5-Methoxy-UTP + 75 % UTP
CTP/50 % 5-Methoxy-UTP + 50 % 1-Methyl-pseudo-UTP
CTP/50 % 5-Methoxy-UTP + 50 % UTP
CTP/5-Methoxy-UTP
CTP/5-Methoxy-UTP (cap 0)
CTP/5-Methoxy-UTP(No cap)
CTP/75 % 5-Methoxy-UTP + 25 % 1-Methyl-pseudo-UTP
CTP/75 % 5-Methoxy-UTP + 25 % UTP
CTP/UTP(No cap)
N1-Me-GTP
N4-Ac-CTP
N4Ac-CTP/1-Methyl-pseudo-UTP
N4Ac-CTP/5-Methoxy-UTP
N4-acetyl-cytidine TP, ATP, GTP, UTP
N4-Bz-CTP/ 5-Methoxy-UTP
N4-methyl CTP
N4-Methyl-CTP/ 5-Methoxy-UTP
Pseudo-iso-CTP/ 5-Methoxy-UTP
PseudoU-alpha-thio-TP
pseudouridine TP, ATP, GTP, CTP
pseudo-UTP/5-methyl-CTP/ATP/GTP
UTP-5-oxyacetic acid Me ester
Xanthosine

[000334] According to the invention, polynucleotides of the invention may be synthesized to comprise the combinations or single modifications of Tables 8 or 9.

[000335] Where a single modification is listed, the listed nucleoside or nucleotide represents 100 percent of that A, U, G or C nucleotide or nucleoside having been modified. Where percentages are listed, these represent the percentage of that particular A, U, G or C nucleobase triphosphate of the total amount of A, U, G, or C triphosphate present. For example, the combination: 25 % 5-Aminoallyl-CTP + 75 % CTP/ 25 % 5-Methoxy-UTP + 75 % UTP refers to a polynucleotide where 25% of the cytosine triphosphates are 5-Aminoallyl-CTP while 75% of the cytosines are CTP; whereas 25% of the uracils are 5-methoxy UTP while 75% of the uracils are UTP. Where no modified UTP is listed then the naturally occurring ATP, UTP, GTP and/or CTP is used at 100% of the sites of those nucleotides found in the polynucleotide. In this example all of the GTP and ATP nucleotides are left unmodified.

IV. Pharmaceutical Compositions

Formulation, Administration, Delivery and Dosing

[000336] The present invention provides polynucleotides compositions and complexes in combination with one or more pharmaceutically acceptable excipients. Pharmaceutical compositions may optionally comprise one or more additional active substances, e.g. therapeutically and/or prophylactically active substances. Pharmaceutical compositions of the present invention may be sterile and/or pyrogen-free. General considerations in the formulation and/or manufacture of pharmaceutical agents may be found, for example, in *Remington: The Science and Practice of Pharmacy* 21st ed., Lippincott Williams & Wilkins, 2005 (incorporated herein by reference in its entirety).

[000337] In some embodiments, compositions are administered to humans, human patients or subjects. For the purposes of the present disclosure, the phrase “active ingredient” generally refers to polynucleotides to be delivered as described herein.

[000338] Although the descriptions of pharmaceutical compositions provided herein are principally directed to pharmaceutical compositions which are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to any other animal, e.g., to non-human animals, e.g.

non-human mammals. Modification of pharmaceutical compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist can design and/or perform such modification with merely ordinary, if any, experimentation. Subjects to which administration of the pharmaceutical compositions is contemplated include, but are not limited to, humans and/or other primates; mammals, including commercially relevant mammals such as cattle, pigs, horses, sheep, cats, dogs, mice, and/or rats; and/or birds, including commercially relevant birds such as poultry, chickens, ducks, geese, and/or turkeys.

[000339] Formulations of the pharmaceutical compositions described herein may be prepared by any method known or hereafter developed in the art of pharmacology. In general, such preparatory methods include the step of bringing the active ingredient into association with an excipient and/or one or more other accessory ingredients, and then, if necessary and/or desirable, dividing, shaping and/or packaging the product into a desired single- or multi-dose unit.

[000340] Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition in accordance with the invention will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. By way of example, the composition may comprise between 0.1% and 100%, e.g., between .5 and 50%, between 1-30%, between 5-80%, at least 80% (w/w) active ingredient.

Formulations

[000341] The polynucleotides of the invention can be formulated using one or more excipients to: (1) increase stability; (2) increase cell transfection; (3) permit the sustained or delayed release (e.g., from a depot formulation of the polynucleotide); (4) alter the biodistribution (e.g., target the polynucleotide to specific tissues or cell types); (5) increase the translation of encoded protein *in vivo*; and/or (6) alter the release profile of encoded protein *in vivo*. In addition to traditional excipients such as any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, excipients

of the present invention can include, without limitation, lipidoids, liposomes, lipid nanoparticles, polymers, lipoplexes, core-shell nanoparticles, peptides, proteins, cells transfected with polynucleotides (e.g., for transplantation into a subject), hyaluronidase, nanoparticle mimics and combinations thereof. Accordingly, the formulations of the invention can include one or more excipients, each in an amount that together increases the stability of the polynucleotide, increases cell transfection by the polynucleotide, increases the expression of polynucleotides encoded protein, and/or alters the release profile of polynucleotide encoded proteins. Further, the polynucleotides of the present invention may be formulated using self-assembled nucleic acid nanoparticles.

[000342] Formulations of the pharmaceutical compositions described herein may be prepared by any method known or hereafter developed in the art of pharmacology. In general, such preparatory methods include the step of associating the active ingredient with an excipient and/or one or more other accessory ingredients.

[000343] A pharmaceutical composition in accordance with the present disclosure may be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. As used herein, a “unit dose” refers to a discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject and/or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.

[000344] Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition in accordance with the present disclosure may vary, depending upon the identity, size, and/or condition of the subject being treated and further depending upon the route by which the composition is to be administered. For example, the composition may comprise between 0.1% and 99% (w/w) of the active ingredient. By way of example, the composition may comprise between 0.1% and 100%, e.g., between .5 and 50%, between 1-30%, between 5-80%, at least 80% (w/w) active ingredient.

[000345] In some embodiments, the formulations described herein may contain at least one polynucleotide. As a non-limiting example, the formulations may contain 1, 2, 3, 4 or 5 polynucleotide.

[000346] In one embodiment, the formulations described herein may comprise more than one type of polynucleotide. In one embodiment, the formulation may comprise a chimeric polynucleotide in linear and circular form. In another embodiment, the formulation may comprise a circular polynucleotide and an IVT polynucleotide. In yet another embodiment, the formulation may comprise an IVT polynucleotide, a chimeric polynucleotide and a circular polynucleotide.

[000347] In one embodiment the formulation may contain polynucleotide encoding proteins selected from categories such as, but not limited to, human proteins, veterinary proteins, bacterial proteins, biological proteins, antibodies, immunogenic proteins, therapeutic peptides and proteins, secreted proteins, plasma membrane proteins, cytoplasmic and cytoskeletal proteins, intracellular membrane bound proteins, nuclear proteins, proteins associated with human disease and/or proteins associated with non-human diseases. In one embodiment, the formulation contains at least three polynucleotides encoding proteins. In one embodiment, the formulation contains at least five polynucleotide encoding proteins.

[000348] Pharmaceutical formulations may additionally comprise a pharmaceutically acceptable excipient, which, as used herein, includes, but is not limited to, any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, and the like, as suited to the particular dosage form desired. Various excipients for formulating pharmaceutical compositions and techniques for preparing the composition are known in the art (see Remington: The Science and Practice of Pharmacy, 21st Edition, A. R. Gennaro, Lippincott, Williams & Wilkins, Baltimore, MD, 2006; incorporated herein by reference in its entirety). The use of a conventional excipient medium may be contemplated within the scope of the present disclosure, except insofar as any conventional excipient medium may be incompatible with a substance or its derivatives, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition.

[000349] In some embodiments, the particle size of the lipid nanoparticle may be increased and/or decreased. The change in particle size may be able to help counter

biological reaction such as, but not limited to, inflammation or may increase the biological effect of the modified mRNA delivered to mammals.

[000350] Pharmaceutically acceptable excipients used in the manufacture of pharmaceutical compositions include, but are not limited to, inert diluents, surface active agents and/or emulsifiers, preservatives, buffering agents, lubricating agents, and/or oils. Such excipients may optionally be included in the pharmaceutical formulations of the invention.

Lipidoids

[000351] The synthesis of lipidoids has been extensively described and formulations containing these compounds are particularly suited for delivery of polynucleotides (see Mahon et al., *Bioconjug Chem.* 2010 21:1448-1454; Schroeder et al., *J Intern Med.* 2010 267:9-21; Akinc et al., *Nat Biotechnol.* 2008 26:561-569; Love et al., *Proc Natl Acad Sci U S A.* 2010 107:1864-1869; Siegwart et al., *Proc Natl Acad Sci U S A.* 2011 108:12996-3001; all of which are incorporated herein in their entireties).

[000352] While these lipidoids have been used to effectively deliver double stranded small interfering RNA molecules in rodents and non-human primates (see Akinc et al., *Nat Biotechnol.* 2008 26:561-569; Frank-Kamenetsky et al., *Proc Natl Acad Sci U S A.* 2008 105:11915-11920; Akinc et al., *Mol Ther.* 2009 17:872-879; Love et al., *Proc Natl Acad Sci U S A.* 2010 107:1864-1869; Leuschner et al., *Nat Biotechnol.* 2011 29:1005-1010; all of which is incorporated herein in their entirety), the present disclosure describes their formulation and use in delivering polynucleotides.

[000353] Complexes, micelles, liposomes or particles can be prepared containing these lipidoids and therefore, can result in an effective delivery of the polynucleotide, as judged by the production of an encoded protein, following the injection of a lipidoid formulation via localized and/or systemic routes of administration. Lipidoid complexes of polynucleotides can be administered by various means including, but not limited to, intravenous, intramuscular, or subcutaneous routes.

[000354] *In vivo* delivery of nucleic acids may be affected by many parameters, including, but not limited to, the formulation composition, nature of particle PEGylation, degree of loading, polynucleotide to lipid ratio, and biophysical parameters such as, but not limited to, particle size (Akinc et al., *Mol Ther.* 2009 17:872-879; herein incorporated

by reference in its entirety). As an example, small changes in the anchor chain length of poly(ethylene glycol) (PEG) lipids may result in significant effects on *in vivo* efficacy. Formulations with the different lipidoids, including, but not limited to penta[3-(1-laurylamino)propionyl]-triethylenetetramine hydrochloride (TETA-5LAP; aka 98N12-5, see Murugaiah et al., *Analytical Biochemistry*, 401:61 (2010); herein incorporated by reference in its entirety), C12-200 (including derivatives and variants), and MD1, can be tested for *in vivo* activity.

[000355] The lipidoid referred to herein as “98N12-5” is disclosed by Akinc et al., *Mol Ther.* 2009 17:872-879 and is incorporated by reference in its entirety.

[000356] The lipidoid referred to herein as “C12-200” is disclosed by Love et al., *Proc Natl Acad Sci U S A.* 2010 107:1864-1869 and Liu and Huang, *Molecular Therapy*. 2010 669-670; both of which are herein incorporated by reference in their entirety. The lipidoid formulations can include particles comprising either 3 or 4 or more components in addition to polynucleotides. As an example, formulations with certain lipidoids, include, but are not limited to, 98N12-5 and may contain 42% lipidoid, 48% cholesterol and 10% PEG (C14 alkyl chain length). As another example, formulations with certain lipidoids, include, but are not limited to, C12-200 and may contain 50% lipidoid, 10% distearylphosphatidyl choline, 38.5% cholesterol, and 1.5% PEG-DMG.

[000357] In one embodiment, a polynucleotide formulated with a lipidoid for systemic intravenous administration can target the liver. For example, a final optimized intravenous formulation using polynucleotides, and comprising a lipid molar composition of 42% 98N12-5, 48% cholesterol, and 10% PEG-lipid with a final weight ratio of about 7.5 to 1 total lipid to polynucleotides, and a C14 alkyl chain length on the PEG lipid, with a mean particle size of roughly 50–60 nm, can result in the distribution of the formulation to be greater than 90% to the liver.(see, Akinc et al., *Mol Ther.* 2009 17:872-879; herein incorporated by reference in its entirety). In another example, an intravenous formulation using a C12-200 (see US provisional application 61/175,770 and published international application WO2010129709, each of which is herein incorporated by reference in their entirety) lipidoid may have a molar ratio of 50/10/38.5/1.5 of C12-200/distearylphosphatidyl choline/cholesterol/PEG-DMG, with a weight ratio of 7 to 1 total lipid to polynucleotides, and a mean particle size of 80 nm may be effective to

deliver polynucleotides to hepatocytes (see, Love et al., Proc Natl Acad Sci U S A. 2010 107:1864-1869 herein incorporated by reference in its entirety). In another embodiment, an MD1 lipidoid-containing formulation may be used to effectively deliver polynucleotides to hepatocytes *in vivo*.

[000358] The characteristics of optimized lipidoid formulations for intramuscular or subcutaneous routes may vary significantly depending on the target cell type and the ability of formulations to diffuse through the extracellular matrix into the blood stream. While a particle size of less than 150 nm may be desired for effective hepatocyte delivery due to the size of the endothelial fenestrae (see, Akinc et al., Mol Ther. 2009 17:872-879 herein incorporated by reference in its entirety), use of a lipidoid-formulated polynucleotides to deliver the formulation to other cells types including, but not limited to, endothelial cells, myeloid cells, and muscle cells may not be similarly size-limited.

[000359] Use of lipidoid formulations to deliver siRNA *in vivo* to other non-hepatocyte cells such as myeloid cells and endothelium has been reported (see Akinc et al., Nat Biotechnol. 2008 26:561-569; Leuschner et al., Nat Biotechnol. 2011 29:1005-1010; Cho et al. Adv. Funct. Mater. 2009 19:3112-3118; 8th International Judah Folkman Conference, Cambridge, MA October 8-9, 2010; each of which is herein incorporated by reference in its entirety). Effective delivery to myeloid cells, such as monocytes, lipidoid formulations may have a similar component molar ratio. Different ratios of lipidoids and other components including, but not limited to, distearylphosphatidyl choline, cholesterol and PEG-DMG, may be used to optimize the formulation of the polynucleotide for delivery to different cell types including, but not limited to, hepatocytes, myeloid cells, muscle cells, etc. For example, the component molar ratio may include, but is not limited to, 50% C12-200, 10% distearylphosphatidyl choline, 38.5% cholesterol, and %1.5 PEG-DMG (see Leuschner et al., Nat Biotechnol 2011 29:1005-1010; herein incorporated by reference in its entirety). The use of lipidoid formulations for the localized delivery of nucleic acids to cells (such as, but not limited to, adipose cells and muscle cells) via either subcutaneous or intramuscular delivery, may not require all of the formulation components desired for systemic delivery, and as such may comprise only the lipidoid and the polynucleotide.

[000360] Combinations of different lipidoids may be used to improve the efficacy of polynucleotides directed protein production as the lipidoids may be able to increase cell transfection by the polynucleotide; and/or increase the translation of encoded protein (see Whitehead et al., Mol. Ther. 2011, 19:1688-1694, herein incorporated by reference in its entirety).

Liposomes, Lipoplexes, and Lipid Nanoparticles

[000361] The polynucleotides of the invention can be formulated using one or more liposomes, lipoplexes, or lipid nanoparticles. In one embodiment, pharmaceutical compositions of polynucleotides include liposomes. Liposomes are artificially-prepared vesicles which may primarily be composed of a lipid bilayer and may be used as a delivery vehicle for the administration of nutrients and pharmaceutical formulations. Liposomes can be of different sizes such as, but not limited to, a multilamellar vesicle (MLV) which may be hundreds of nanometers in diameter and may contain a series of concentric bilayers separated by narrow aqueous compartments, a small unilamellar vesicle (SUV) which may be smaller than 50 nm in diameter, and a large unilamellar vesicle (LUV) which may be between 50 and 500 nm in diameter. Liposome design may include, but is not limited to, opsonins or ligands in order to improve the attachment of liposomes to unhealthy tissue or to activate events such as, but not limited to, endocytosis. Liposomes may contain a low or a high pH in order to improve the delivery of the pharmaceutical formulations.

[000362] The formation of liposomes may depend on the physicochemical characteristics such as, but not limited to, the pharmaceutical formulation entrapped and the liposomal ingredients, the nature of the medium in which the lipid vesicles are dispersed, the effective concentration of the entrapped substance and its potential toxicity, any additional processes involved during the application and/or delivery of the vesicles, the optimization size, polydispersity and the shelf-life of the vesicles for the intended application, and the batch-to-batch reproducibility and possibility of large-scale production of safe and efficient liposomal products.

[000363] As a non-limiting example, liposomes such as synthetic membrane vesicles may be prepared by the methods, apparatus and devices described in US Patent Publication No. US20130177638, US20130177637, US20130177636, US20130177635,

US20130177634, US20130177633, US20130183375, US20130183373 and US20130183372, the contents of each of which are herein incorporated by reference in its entirety.

[000364] In one embodiment, pharmaceutical compositions described herein may include, without limitation, liposomes such as those formed from 1,2-dioleoyloxy-*N,N*-dimethylaminopropane (DODMA) liposomes, DiLa2 liposomes from Marina Biotech (Bothell, WA), 1,2-dilinoleoyloxy-3-dimethylaminopropane (DLin-DMA), 2,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA), and MC3 (US20100324120; herein incorporated by reference in its entirety) and liposomes which may deliver small molecule drugs such as, but not limited to, DOXIL® from Janssen Biotech, Inc. (Horsham, PA).

[000365] In one embodiment, pharmaceutical compositions described herein may include, without limitation, liposomes such as those formed from the synthesis of stabilized plasmid-lipid particles (SPLP) or stabilized nucleic acid lipid particle (SNALP) that have been previously described and shown to be suitable for oligonucleotide delivery *in vitro* and *in vivo* (see Wheeler et al. Gene Therapy. 1999 6:271-281; Zhang et al. Gene Therapy. 1999 6:1438-1447; Jeffs et al. Pharm Res. 2005 22:362-372; Morrissey et al., Nat Biotechnol. 2005 2:1002-1007; Zimmermann et al., Nature. 2006 441:111-114; Heyes et al. J Contr Rel. 2005 107:276-287; Semple et al. Nature Biotech. 2010 28:172-176; Judge et al. J Clin Invest. 2009 119:661-673; deFougerolles *Hum Gene Ther.* 2008 19:125-132; U.S. Patent Publication No US20130122104; all of which are incorporated herein in their entireties). The original manufacture method by Wheeler et al. was a detergent dialysis method, which was later improved by Jeffs et al. and is referred to as the spontaneous vesicle formation method. The liposome formulations are composed of 3 to 4 lipid components in addition to the polynucleotide. As an example a liposome can contain, but is not limited to, 55% cholesterol, 20% distearylphosphatidyl choline (DSPC), 10% PEG-S-DSG, and 15% 1,2-dioleoyloxy-*N,N*-dimethylaminopropane (DODMA), as described by Jeffs et al. As another example, certain liposome formulations may contain, but are not limited to, 48% cholesterol, 20% DSPC, 2% PEG-c-DMA, and 30% cationic lipid, where the cationic lipid can be 1,2-distearloxy-*N,N*-

dimethylaminopropane (DSDMA), DODMA, DLin-DMA, or 1,2-dilinolenyloxy-3-dimethylaminopropane (DLenDMA), as described by Heyes et al.

[000366] In some embodiments, liposome formulations may comprise from about 25.0% cholesterol to about 40.0% cholesterol, from about 30.0% cholesterol to about 45.0% cholesterol, from about 35.0% cholesterol to about 50.0% cholesterol and/or from about 48.5% cholesterol to about 60% cholesterol. In a preferred embodiment, formulations may comprise a percentage of cholesterol selected from the group consisting of 28.5%, 31.5%, 33.5%, 36.5%, 37.0%, 38.5%, 39.0% and 43.5%. In some embodiments, formulations may comprise from about 5.0% to about 10.0% DSPC and/or from about 7.0% to about 15.0% DSPC.

[000367] In one embodiment, pharmaceutical compositions may include liposomes which may be formed to deliver polynucleotides which may encode at least one immunogen or any other polypeptide of interest. The polynucleotide may be encapsulated by the liposome and/or it may be contained in an aqueous core which may then be encapsulated by the liposome (see International Pub. Nos. WO2012031046, WO2012031043, WO2012030901 and WO2012006378 and US Patent Publication No. US20130189351, US20130195969 and US20130202684; the contents of each of which are herein incorporated by reference in their entirety).

[000368] In another embodiment, liposomes may be formulated for targeted delivery. As a non-limiting example, the liposome may be formulated for targeted delivery to the liver. The liposome used for targeted delivery may include, but is not limited to, the liposomes described in and methods of making liposomes described in US Patent Publication No. US20130195967, the contents of which are herein incorporated by reference in its entirety.

[000369] In another embodiment, the polynucleotide which may encode an immunogen may be formulated in a cationic oil-in-water emulsion where the emulsion particle comprises an oil core and a cationic lipid which can interact with the polynucleotide anchoring the molecule to the emulsion particle (see International Pub. No. WO2012006380; herein incorporated by reference in its entirety).

[000370] In one embodiment, the polynucleotides may be formulated in a water-in-oil emulsion comprising a continuous hydrophobic phase in which the hydrophilic phase is

dispersed. As a non-limiting example, the emulsion may be made by the methods described in International Publication No. WO201087791, herein incorporated by reference in its entirety.

[000371] In another embodiment, the lipid formulation may include at least cationic lipid, a lipid which may enhance transfection and a least one lipid which contains a hydrophilic head group linked to a lipid moiety (International Pub. No. WO2011076807 and U.S. Pub. No. 20110200582; the contents of each of which is herein incorporated by reference in their entirety). In another embodiment, the polynucleotides encoding an immunogen may be formulated in a lipid vesicle which may have crosslinks between functionalized lipid bilayers (see U.S. Pub. No. 20120177724, the contents of which is herein incorporated by reference in its entirety).

[000372] In one embodiment, the polynucleotides may be formulated in a liposome as described in International Patent Publication No. WO2013086526, herein incorporated by reference in its entirety. The polynucleotides may be encapsulated in a liposome using reverse pH gradients and/or optimized internal buffer compositions as described in International Patent Publication No. WO2013086526, herein incorporated by reference in its entirety.

[000373] In one embodiment, the pharmaceutical compositions may be formulated in liposomes such as, but not limited to, DiLa2 liposomes (Marina Biotech, Bothell, WA), SMARTICLES® (Marina Biotech, Bothell, WA), neutral DOPC (1,2-dioleoyl-sn-glycero-3-phosphocholine) based liposomes (e.g., siRNA delivery for ovarian cancer (Landen et al. Cancer Biology & Therapy 2006 5(12)1708-1713); herein incorporated by reference in its entirety) and hyaluronan-coated liposomes (Quiet Therapeutics, Israel).

[000374] In one embodiment, the cationic lipid may be a low molecular weight cationic lipid such as those described in US Patent Application No. 20130090372, the contents of which are herein incorporated by reference in its entirety.

[000375] In one embodiment, the polynucleotides may be formulated in a lipid vesicle which may have crosslinks between functionalized lipid bilayers.

[000376] In one embodiment, the polynucleotides may be formulated in a liposome comprising a cationic lipid. The liposome may have a molar ratio of nitrogen atoms in the cationic lipid to the phosphates in the RNA (N:P ratio) of between 1:1 and 20:1 as

described in International Publication No. WO2013006825, herein incorporated by reference in its entirety. In another embodiment, the liposome may have a N:P ratio of greater than 20:1 or less than 1:1.

[000377] In one embodiment, the polynucleotides may be formulated in a lipid-polycation complex. The formation of the lipid-polycation complex may be accomplished by methods known in the art and/or as described in U.S. Pub. No. 20120178702, herein incorporated by reference in its entirety. As a non-limiting example, the polycation may include a cationic peptide or a polypeptide such as, but not limited to, polylysine, polyornithine and/or polyarginine and the cationic peptides described in International Pub. No. WO2012013326 or US Patent Pub. No. US20130142818; each of which is herein incorporated by reference in its entirety. In another embodiment, the polynucleotides may be formulated in a lipid-polycation complex which may further include a neutral lipid such as, but not limited to, cholesterol or dioleoyl phosphatidylethanolamine (DOPE).

[000378] In one embodiment, the polynucleotide may be formulated in an aminoalcohol lipidoid. Aminoalcohol lipidoids which may be used in the present invention may be prepared by the methods described in U.S. Patent No. 8,450,298, herein incorporated by reference in its entirety.

[000379] The liposome formulation may be influenced by, but not limited to, the selection of the cationic lipid component, the degree of cationic lipid saturation, the nature of the PEGylation, ratio of all components and biophysical parameters such as size. In one example by Semple et al. (Semple et al. Nature Biotech. 2010 28:172-176; herein incorporated by reference in its entirety), the liposome formulation was composed of 57.1 % cationic lipid, 7.1% dipalmitoylphosphatidylcholine, 34.3 % cholesterol, and 1.4% PEG-c-DMA. As another example, changing the composition of the cationic lipid could more effectively deliver siRNA to various antigen presenting cells (Basha et al. Mol Ther. 2011 19:2186-2200; herein incorporated by reference in its entirety). In some embodiments, liposome formulations may comprise from about 35 to about 45% cationic lipid, from about 40% to about 50% cationic lipid, from about 50% to about 60% cationic lipid and/or from about 55% to about 65% cationic lipid. In some embodiments, the ratio

of lipid to mRNA in liposomes may be from about 5:1 to about 20:1, from about 10:1 to about 25:1, from about 15:1 to about 30:1 and/or at least 30:1.

[000380] In some embodiments, the ratio of PEG in the lipid nanoparticle (LNP) formulations may be increased or decreased and/or the carbon chain length of the PEG lipid may be modified from C14 to C18 to alter the pharmacokinetics and/or biodistribution of the LNP formulations. As a non-limiting example, LNP formulations may contain from about 0.5% to about 3.0%, from about 1.0% to about 3.5%, from about 1.5% to about 4.0%, from about 2.0% to about 4.5%, from about 2.5% to about 5.0% and/or from about 3.0% to about 6.0% of the lipid molar ratio of PEG-c-DOMG as compared to the cationic lipid, DSPC and cholesterol. In another embodiment the PEG-c-DOMG may be replaced with a PEG lipid such as, but not limited to, PEG-DSG (1,2-Distearoyl-sn-glycerol, methoxypolyethylene glycol), PEG-DMG (1,2-Dimyristoyl-sn-glycerol) and/or PEG-DPG (1,2-Dipalmitoyl-sn-glycerol, methoxypolyethylene glycol). The cationic lipid may be selected from any lipid known in the art such as, but not limited to, DLin-MC3-DMA, DLin-DMA, C12-200 and DLin-KC2-DMA.

[000381] In one embodiment, the polynucleotides may be formulated in a lipid nanoparticle such as those described in International Publication No. WO2012170930, herein incorporated by reference in its entirety.

[000382] In one embodiment, the formulation comprising the polynucleotide is a nanoparticle which may comprise at least one lipid. The lipid may be selected from, but is not limited to, DLin-DMA, DLin-K-DMA, 98N12-5, C12-200, DLin-MC3-DMA, DLin-KC2-DMA, DODMA, PLGA, PEG, PEG-DMG, PEGylated lipids and amino alcohol lipids. In another aspect, the lipid may be a cationic lipid such as, but not limited to, DLin-DMA, DLin-D-DMA, DLin-MC3-DMA, DLin-KC2-DMA, DODMA and amino alcohol lipids. The amino alcohol cationic lipid may be the lipids described in and/or made by the methods described in US Patent Publication No. US20130150625, herein incorporated by reference in its entirety. As a non-limiting example, the cationic lipid may be 2-amino-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-[[[(9Z,2Z)-octadeca-9,12-dien-1-yloxy]methyl]propan-1-ol (Compound 1 in US20130150625); 2-amino-3-[(9Z)-octadec-9-en-1-yloxy]-2-[[[(9Z)-octadec-9-en-1-yloxy]methyl]propan-1-ol (Compound 2 in US20130150625); 2-amino-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-

[(octyloxy)methyl]propan-1-ol (Compound 3 in US20130150625); and 2-(dimethylamino)-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-[[9Z,12Z)-octadeca-9,12-dien-1-yloxy]methyl}propan-1-ol (Compound 4 in US20130150625); or any pharmaceutically acceptable salt or stereoisomer thereof.

[000383] In one embodiment, the cationic lipid may be selected from, but not limited to, a cationic lipid described in International Publication Nos. WO2012040184, WO2011153120, WO2011149733, WO2011090965, WO2011043913, WO2011022460, WO2012061259, WO2012054365, WO2012044638, WO2010080724, WO201021865, WO2008103276, WO2013086373 and WO2013086354, US Patent Nos. 7,893,302, 7,404,969, 8,283,333, and 8,466,122 and US Patent Publication No. US20100036115, US20120202871, US20130064894, US20130129785, US20130150625, US20130178541 and US20130225836; the contents of each of which are herein incorporated by reference in their entirety. In another embodiment, the cationic lipid may be selected from, but not limited to, formula A described in International Publication Nos. WO2012040184, WO2011153120, WO2011149733, WO2011090965, WO2011043913, WO2011022460, WO2012061259, WO2012054365, WO2012044638 and WO2013116126 or US Patent Publication No. US20130178541 and US20130225836; the contents of each of which is herein incorporated by reference in their entirety. In yet another embodiment, the cationic lipid may be selected from, but not limited to, formula CLI-CLXXIX of International Publication No. WO2008103276, formula CLI-CLXXIX of US Patent No. 7,893,302, formula CLI-CLXXXII of US Patent No. 7,404,969 and formula I-VI of US Patent Publication No. US20100036115, formula I of US Patent Publication No. US20130123338; each of which is herein incorporated by reference in their entirety. As a non-limiting example, the cationic lipid may be selected from (20Z,23Z)-N,N-dimethylnonacos-20,23-dien-10-amine, (17Z,20Z)-N,N-dimethylhexacos-17,20-dien-9-amine, (1Z,19Z)-N,N-dimethylpentacos-16,19-dien-8-amine, (13Z,16Z)-N,N-dimethyldocos-13,16-dien-5-amine, (12Z,15Z)-N,N-dimethylhenicos-12,15-dien-4-amine, (14Z,17Z)-N,N-dimethyltricos-14,17-dien-6-amine, (15Z,18Z)-N,N-dimethyltetracos-15,18-dien-7-amine, (18Z,21Z)-N,N-dimethylheptacos-18,21-dien-10-amine, (15Z,18Z)-N,N-dimethyltetracos-15,18-dien-5-amine, (14Z,17Z)-N,N-dimethyltricos-14,17-dien-4-amine, (19Z,22Z)-N,N-dimethyloctacos-19,22-dien-9-

amine, (18Z,21 Z)-N,N-dimethylheptacos-18,21-dien-8-amine, (17Z,20Z)-N,N-dimethylhexacos-17,20-dien-7-amine, (16Z,19Z)-N,N-dimethylpentacos-16,19-dien-6-amine, (22Z,25Z)-N,N-dimethylhentriaconta-22,25-dien-10-amine, (21 Z,24Z)-N,N-dimethyltriaconta-21,24-dien-9-amine, (18Z)-N,N-dimethylheptacos-18-en-10-amine, (17Z)-N,N-dimethylhexacos-17-en-9-amine, (19Z,22Z)-N,N-dimethyloctacos-19,22-dien-7-amine, N,N-dimethylheptacosan-10-amine, (20Z,23Z)-N-ethyl-N-methylnonacos-20,23-dien-10-amine, 1-[(11Z,14Z)-1-nonylicos-11,14-dien-1-yl]pyrrolidine, (20Z)-N,N-dimethylheptacos-20-en-10-amine, (15Z)-N,N-dimethylheptacos-15-en-10-amine, (14Z)-N,N-dimethylnonacos-14-en-10-amine, (17Z)-N,N-dimethylnonacos-17-en-10-amine, (24Z)-N,N-dimethyltritriacont-24-en-10-amine, (20Z)-N,N-dimethylnonacos-20-en-10-amine, (22Z)-N,N-dimethylhentriacont-22-en-10-amine, (16Z)-N,N-dimethylpentacos-16-en-8-amine, (12Z,15Z)-N,N-dimethyl-2-nonylhenicos-12,15-dien-1-amine, (13Z,16Z)-N,N-dimethyl-3-nonyldocos-13,16-dien-1-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]heptadecan-8-amine, 1-[(1S,2R)-2-hexylcyclopropyl]-N,N-dimethylnonadecan-10-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]nonadecan-10-amine, N,N-dimethyl-21-[(1S,2R)-2-octylcyclopropyl]henicosan-10-amine, N,N-dimethyl-1-[(1S,2S)-2-[(1R,2R)-2-pentylcyclopropyl]methyl]cyclopropyl]nonadecan-10-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]hexadecan-8-amine, N,N-dimethyl-[(1R,2S)-2-undecylcyclopropyl]tetradecan-5-amine, N,N-dimethyl-3-{7-[(1S,2R)-2-octylcyclopropyl]heptyl}dodecan-1-amine, 1-[(1R,2S)-2-heptylcyclopropyl]-N,N-dimethyloctadecan-9-amine, 1-[(1S,2R)-2-decylcyclopropyl]-N,N-dimethylpentadecan-6-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]pentadecan-8-amine, R-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-(octyloxy)propan-2-amine, S-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-(octyloxy)propan-2-amine, 1-{2-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-1-[(octyloxy)methyl]ethyl}pyrrolidine, (2S)-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-[(5Z)-oct-5-en-1-yloxy]propan-2-amine, 1-{2-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-1-[(octyloxy)methyl]ethyl}azetidine, (2S)-1-(hexyloxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, (2S)-1-(heptyloxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-(nonyloxy)-3-

[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-[(9Z)-octadec-9-en-1-yloxy]-3-(octyloxy)propan-2-amine; (2S)-N,N-dimethyl-1-[(6Z,9Z,12Z)-octadeca-6,9,12-trien-1-yloxy]-3-(octyloxy)propan-2-amine, (2S)-1-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethyl-3-(pentyloxy)propan-2-amine, (2S)-1-(hexyloxy)-3-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethylpropan-2-amine, 1-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, 1-[(13Z,16Z)-docosa-13,16-dien-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, (2S)-1-[(13Z,16Z)-docosa-13,16-dien-1-yloxy]-3-(hexyloxy)-N,N-dimethylpropan-2-amine, (2S)-1-[(13Z)-docos-13-en-1-yloxy]-3-(hexyloxy)-N,N-dimethylpropan-2-amine, 1-[(13Z)-docos-13-en-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, 1-[(9Z)-hexadec-9-en-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, (2R)-N,N-dimethyl-H(1-metoyloctyl)oxy]-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, (2R)-1-[(3,7-dimethyloctyl)oxy]-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-(octyloxy)-3-({8-[(1S,2S)-2-[(1R,2R)-2-pentylcyclopropyl]methyl}cyclopropyl]octyl}oxy)propan-2-amine, N,N-dimethyl-1-{{8-(2-ocylcyclopropyl)octyl}oxy}-3-(octyloxy)propan-2-amine and (11E,20Z,23Z)-N,N-dimethylnonacos-11,20,2-trien-10-amine or a pharmaceutically acceptable salt or stereoisomer thereof.

[000384] In one embodiment, the lipid may be a cleavable lipid such as those described in International Publication No. WO2012170889, herein incorporated by reference in its entirety.

[000385] In another embodiment, the lipid may be a cationic lipid such as, but not limited to, Formula (I) of U.S. Patent Application No. US20130064894, the contents of which are herein incorporated by reference in its entirety.

[000386] In one embodiment, the cationic lipid may be synthesized by methods known in the art and/or as described in International Publication Nos. WO2012040184, WO2011153120, WO2011149733, WO2011090965, WO2011043913, WO2011022460, WO2012061259, WO2012054365, WO2012044638, WO2010080724, WO201021865, WO2013086373 and WO2013086354; the contents of each of which are herein incorporated by reference in their entirety.

[000387] In another embodiment, the cationic lipid may be a trialkyl cationic lipid. Non-limiting examples of trialkyl cationic lipids and methods of making and using the trialkyl cationic lipids are described in International Patent Publication No. WO2013126803, the contents of which are herein incorporated by reference in its entirety.

[000388] In one embodiment, the LNP formulations of the polynucleotides may contain PEG-c-DOMG at 3% lipid molar ratio. In another embodiment, the LNP formulations polynucleotides may contain PEG-c-DOMG at 1.5% lipid molar ratio.

[000389] In one embodiment, the pharmaceutical compositions of the polynucleotides may include at least one of the PEGylated lipids described in International Publication No. WO2012099755, herein incorporated by reference.

[000390] In one embodiment, the LNP formulation may contain PEG-DMG 2000 (1,2-dimyristoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000]. In one embodiment, the LNP formulation may contain PEG-DMG 2000, a cationic lipid known in the art and at least one other component. In another embodiment, the LNP formulation may contain PEG-DMG 2000, a cationic lipid known in the art, DSPC and cholesterol. As a non-limiting example, the LNP formulation may contain PEG-DMG 2000, DLin-DMA, DSPC and cholesterol. As another non-limiting example the LNP formulation may contain PEG-DMG 2000, DLin-DMA, DSPC and cholesterol in a molar ratio of 2:40:10:48 (see e.g., Geall et al., Nonviral delivery of self-amplifying RNA vaccines, PNAS 2012; PMID: 22908294; herein incorporated by reference in its entirety).

[000391] In one embodiment, the LNP formulation may be formulated by the methods described in International Publication Nos. WO2011127255 or WO2008103276, the contents of each of which is herein incorporated by reference in their entirety. As a non-limiting example, the polynucleotides described herein may be encapsulated in LNP formulations as described in WO2011127255 and/or WO2008103276; each of which is herein incorporated by reference in their entirety.

[000392] In one embodiment, the polynucleotides described herein may be formulated in a nanoparticle to be delivered by a parenteral route as described in U.S. Pub. No. US20120207845; the contents of which are herein incorporated by reference in its entirety.

[000393] In one embodiment, the polynucleotides may be formulated in a lipid nanoparticle made by the methods described in US Patent Publication No US20130156845 or International Publication No WO2013093648 or WO2012024526, each of which is herein incorporated by reference in its entirety.

[000394] The lipid nanoparticles described herein may be made in a sterile environment by the system and/or methods described in US Patent Publication No. US20130164400, herein incorporated by reference in its entirety.

[000395] In one embodiment, the LNP formulation may be formulated in a nanoparticle such as a nucleic acid-lipid particle described in US Patent No. 8,492,359, the contents of which are herein incorporated by reference in its entirety. As a non-limiting example, the lipid particle may comprise one or more active agents or therapeutic agents; one or more cationic lipids comprising from about 50 mol % to about 85 mol % of the total lipid present in the particle; one or more non-cationic lipids comprising from about 13 mol % to about 49.5 mol % of the total lipid present in the particle; and one or more conjugated lipids that inhibit aggregation of particles comprising from about 0.5 mol % to about 2 mol % of the total lipid present in the particle. The nucleic acid in the nanoparticle may be the polynucleotides described herein and/or are known in the art.

[000396] In one embodiment, the LNP formulation may be formulated by the methods described in International Publication Nos. WO2011127255 or WO2008103276, the contents of each of which are herein incorporated by reference in their entirety. As a non-limiting example, modified RNA described herein may be encapsulated in LNP formulations as described in WO2011127255 and/or WO2008103276; the contents of each of which are herein incorporated by reference in their entirety.

[000397] In one embodiment, LNP formulations described herein may comprise a polycationic composition. As a non-limiting example, the polycationic composition may be selected from formula 1-60 of US Patent Publication No. US20050222064; the content of which is herein incorporated by reference in its entirety. In another embodiment, the LNP formulations comprising a polycationic composition may be used for the delivery of the modified RNA described herein *in vivo* and/or *in vitro*.

[000398] In one embodiment, the LNP formulations described herein may additionally comprise a permeability enhancer molecule. Non-limiting permeability enhancer

molecules are described in US Patent Publication No. US20050222064; the content of which is herein incorporated by reference in its entirety.

[000399] In one embodiment, the pharmaceutical compositions may be formulated in liposomes such as, but not limited to, DiLa2 liposomes (Marina Biotech, Bothell, WA), SMARTICLES® (Marina Biotech, Bothell, WA), neutral DOPC (1,2-dioleoyl-sn-glycero-3-phosphocholine) based liposomes (e.g., siRNA delivery for ovarian cancer (Landen et al. Cancer Biology & Therapy 2006 5(12)1708-1713); herein incorporated by reference in its entirety) and hyaluronan-coated liposomes (Quiet Therapeutics, Israel).

[000400] In one embodiment, the polynucleotides may be formulated in a lyophilized gel-phase liposomal composition as described in US Publication No. US2012060293, herein incorporated by reference in its entirety.

[000401] The nanoparticle formulations may comprise a phosphate conjugate. The phosphate conjugate may increase *in vivo* circulation times and/or increase the targeted delivery of the nanoparticle. Phosphate conjugates for use with the present invention may be made by the methods described in International Application No. WO2013033438 or US Patent Publication No. US20130196948, the contents of each of which are herein incorporated by reference in its entirety. As a non-limiting example, the phosphate conjugates may include a compound of any one of the formulas described in International Application No. WO2013033438, herein incorporated by reference in its entirety.

[000402] The nanoparticle formulation may comprise a polymer conjugate. The polymer conjugate may be a water soluble conjugate. The polymer conjugate may have a structure as described in U.S. Patent Application No. 20130059360, the contents of which are herein incorporated by reference in its entirety. In one aspect, polymer conjugates with the polynucleotides of the present invention may be made using the methods and/or segmented polymeric reagents described in U.S. Patent Application No. 20130072709, herein incorporated by reference in its entirety. In another aspect, the polymer conjugate may have pendant side groups comprising ring moieties such as, but not limited to, the polymer conjugates described in US Patent Publication No. US20130196948, the contents of which is herein incorporated by reference in its entirety.

[000403] The nanoparticle formulations may comprise a conjugate to enhance the delivery of nanoparticles of the present invention in a subject. Further, the conjugate may

inhibit phagocytic clearance of the nanoparticles in a subject. In one aspect, the conjugate may be a “self” peptide designed from the human membrane protein CD47 (e.g., the “self” particles described by Rodriguez et al (Science 2013 339, 971-975), herein incorporated by reference in its entirety). As shown by Rodriguez et al. the self-peptides delayed macrophage-mediated clearance of nanoparticles which enhanced delivery of the nanoparticles. In another aspect, the conjugate may be the membrane protein CD47 (e.g., see Rodriguez et al. Science 2013 339, 971-975, herein incorporated by reference in its entirety). Rodriguez et al. showed that, similarly to “self” peptides, CD47 can increase the circulating particle ratio in a subject as compared to scrambled peptides and PEG coated nanoparticles.

[000404] In one embodiment, the polynucleotides of the present invention are formulated in nanoparticles which comprise a conjugate to enhance the delivery of the nanoparticles of the present invention in a subject. The conjugate may be the CD47 membrane or the conjugate may be derived from the CD47 membrane protein, such as the “self” peptide described previously. In another aspect the nanoparticle may comprise PEG and a conjugate of CD47 or a derivative thereof. In yet another aspect, the nanoparticle may comprise both the “self” peptide described above and the membrane protein CD47.

[000405] In another aspect, a “self” peptide and/or CD47 protein may be conjugated to a virus-like particle or pseudovirion, as described herein for delivery of the polynucleotides of the present invention.

[000406] In another embodiment, pharmaceutical compositions comprising the polynucleotides of the present invention and a conjugate which may have a degradable linkage. Non-limiting examples of conjugates include an aromatic moiety comprising an ionizable hydrogen atom, a spacer moiety, and a water-soluble polymer. As a non-limiting example, pharmaceutical compositions comprising a conjugate with a degradable linkage and methods for delivering such pharmaceutical compositions are described in US Patent Publication No. US20130184443, the contents of which are herein incorporated by reference in its entirety.

[000407] The nanoparticle formulations may be a carbohydrate nanoparticle comprising a carbohydrate carrier and a polynucleotide. As a non-limiting example, the carbohydrate

carrier may include, but is not limited to, an anhydride-modified phytoglycogen or glycogen-type material, phytoglycogen octenyl succinate, phytoglycogen beta-dextrin, anhydride-modified phytoglycogen beta-dextrin. (See e.g., International Publication No. WO2012109121; the contents of which are herein incorporated by reference in its entirety).

[000408] Nanoparticle formulations of the present invention may be coated with a surfactant or polymer in order to improve the delivery of the particle. In one embodiment, the nanoparticle may be coated with a hydrophilic coating such as, but not limited to, PEG coatings and/or coatings that have a neutral surface charge. The hydrophilic coatings may help to deliver nanoparticles with larger payloads such as, but not limited to, polynucleotides within the central nervous system. As a non-limiting example nanoparticles comprising a hydrophilic coating and methods of making such nanoparticles are described in US Patent Publication No. US20130183244, the contents of which are herein incorporated by reference in its entirety.

[000409] In one embodiment, the lipid nanoparticles of the present invention may be hydrophilic polymer particles. Non-limiting examples of hydrophilic polymer particles and methods of making hydrophilic polymer particles are described in US Patent Publication No. US20130210991, the contents of which are herein incorporated by reference in its entirety.

[000410] In another embodiment, the lipid nanoparticles of the present invention may be hydrophobic polymer particles.

[000411] Lipid nanoparticle formulations may be improved by replacing the cationic lipid with a biodegradable cationic lipid which is known as a rapidly eliminated lipid nanoparticle (reLNP). Ionizable cationic lipids, such as, but not limited to, DLinDMA, DLin-KC2-DMA, and DLin-MC3-DMA, have been shown to accumulate in plasma and tissues over time and may be a potential source of toxicity. The rapid metabolism of the rapidly eliminated lipids can improve the tolerability and therapeutic index of the lipid nanoparticles by an order of magnitude from a 1 mg/kg dose to a 10 mg/kg dose in rat. Inclusion of an enzymatically degraded ester linkage can improve the degradation and metabolism profile of the cationic component, while still maintaining the activity of the reLNP formulation. The ester linkage can be internally located within the lipid chain or it

may be terminally located at the terminal end of the lipid chain. The internal ester linkage may replace any carbon in the lipid chain.

[000412] In one embodiment, the internal ester linkage may be located on either side of the saturated carbon.

[000413] In one embodiment, an immune response may be elicited by delivering a lipid nanoparticle which may include a nanospecies, a polymer and an immunogen. (U.S. Publication No. 20120189700 and International Publication No. WO2012099805; each of which is herein incorporated by reference in their entirety). The polymer may encapsulate the nanospecies or partially encapsulate the nanospecies. The immunogen may be a recombinant protein, a modified RNA and/or a polynucleotide described herein. In one embodiment, the lipid nanoparticle may be formulated for use in a vaccine such as, but not limited to, against a pathogen.

[000414] Lipid nanoparticles may be engineered to alter the surface properties of particles so the lipid nanoparticles may penetrate the mucosal barrier. Mucus is located on mucosal tissue such as, but not limited to, oral (e.g., the buccal and esophageal membranes and tonsil tissue), ophthalmic, gastrointestinal (e.g., stomach, small intestine, large intestine, colon, rectum), nasal, respiratory (e.g., nasal, pharyngeal, tracheal and bronchial membranes), genital (e.g., vaginal, cervical and urethral membranes). Nanoparticles larger than 10-200 nm which are preferred for higher drug encapsulation efficiency and the ability to provide the sustained delivery of a wide array of drugs have been thought to be too large to rapidly diffuse through mucosal barriers. Mucus is continuously secreted, shed, discarded or digested and recycled so most of the trapped particles may be removed from the mucosa tissue within seconds or within a few hours. Large polymeric nanoparticles (200nm -500nm in diameter) which have been coated densely with a low molecular weight polyethylene glycol (PEG) diffused through mucus only 4 to 6-fold lower than the same particles diffusing in water (Lai et al. PNAS 2007 104(5):1482-487; Lai et al. Adv Drug Deliv Rev. 2009 61(2): 158-171; each of which is herein incorporated by reference in their entirety). The transport of nanoparticles may be determined using rates of permeation and/or fluorescent microscopy techniques including, but not limited to, fluorescence recovery after photobleaching (FRAP) and high resolution multiple particle tracking (MPT). As a non-limiting example,

compositions which can penetrate a mucosal barrier may be made as described in U.S. Pat. No. 8,241,670 or International Patent Publication No. WO2013110028, the contents of each of which are herein incorporated by reference in its entirety.

[000415] The lipid nanoparticle engineered to penetrate mucus may comprise a polymeric material (i.e. a polymeric core) and/or a polymer-vitamin conjugate and/or a tri-block co-polymer. The polymeric material may include, but is not limited to, polyamines, polyethers, polyamides, polyesters, polycarbamates, polyureas, polycarbonates, poly(styrenes), polyimides, polysulfones, polyurethanes, polyacetylenes, polyethylenes, polyethyleneimines, polyisocyanates, polyacrylates, polymethacrylates, polyacrylonitriles, and polyarylates. The polymeric material may be biodegradable and/or biocompatible. Non-limiting examples of biocompatible polymers are described in International Patent Publication No. WO2013116804, the contents of which are herein incorporated by reference in its entirety. The polymeric material may additionally be irradiated. As a non-limiting example, the polymeric material may be gamma irradiated (See e.g., International App. No. WO201282165, herein incorporated by reference in its entirety). Non-limiting examples of specific polymers include poly(caprolactone) (PCL), ethylene vinyl acetate polymer (EVA), poly(lactic acid) (PLA), poly(L-lactic acid) (PLLA), poly(glycolic acid) (PGA), poly(lactic acid-co-glycolic acid) (PLGA), poly(L-lactic acid-co-glycolic acid) (PLLGA), poly(D,L-lactide) (PDLA), poly(L-lactide) (PLLA), poly(D,L-lactide-co-caprolactone), poly(D,L-lactide-co-caprolactone-co-glycolide), poly(D,L-lactide-co-PEO-co-D,L-lactide), poly(D,L-lactide-co-PPO-co-D,L-lactide), polyalkyl cyanoacrylate, polyurethane, poly-L-lysine (PLL), hydroxypropyl methacrylate (HPMA), polyethyleneglycol, poly-L-glutamic acid, poly(hydroxy acids), polyanhydrides, polyorthoesters, poly(ester amides), polyamides, poly(ester ethers), polycarbonates, polyalkylenes such as polyethylene and polypropylene, polyalkylene glycols such as poly(ethylene glycol) (PEG), polyalkylene oxides (PEO), polyalkylene terephthalates such as poly(ethylene terephthalate), polyvinyl alcohols (PVA), polyvinyl ethers, polyvinyl esters such as poly(vinyl acetate), polyvinyl halides such as poly(vinyl chloride) (PVC), polyvinylpyrrolidone, polysiloxanes, polystyrene (PS), polyurethanes, derivatized celluloses such as alkyl celluloses, hydroxyalkyl celluloses, cellulose ethers, cellulose esters, nitro celluloses, hydroxypropylcellulose, carboxymethylcellulose,

polymers of acrylic acids, such as poly(methyl(meth)acrylate) (PMMA), poly(ethyl(meth)acrylate), poly(butyl(meth)acrylate), poly(isobutyl(meth)acrylate), poly(hexyl(meth)acrylate), poly(isodecyl(meth)acrylate), poly(lauryl(meth)acrylate), poly(phenyl(meth)acrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate) and copolymers and mixtures thereof, polydioxanone and its copolymers, polyhydroxyalkanoates, polypropylene fumarate, polyoxymethylene, poloxamers, poly(ortho)esters, poly(butyric acid), poly(valeric acid), poly(lactide-co-caprolactone), PEG-PLGA-PEG and trimethylene carbonate, polyvinylpyrrolidone. The lipid nanoparticle may be coated or associated with a co-polymer such as, but not limited to, a block co-polymer (such as a branched polyether-polyamide block copolymer described in International Publication No. WO2013012476, herein incorporated by reference in its entirety), and (poly(ethylene glycol))- (poly(propylene oxide))-(poly(ethylene glycol)) triblock copolymer (see e.g., US Publication 20120121718 and US Publication 20100003337 and U.S. Pat. No. 8,263,665; each of which is herein incorporated by reference in their entirety). The co-polymer may be a polymer that is generally regarded as safe (GRAS) and the formation of the lipid nanoparticle may be in such a way that no new chemical entities are created. For example, the lipid nanoparticle may comprise poloxamers coating PLGA nanoparticles without forming new chemical entities which are still able to rapidly penetrate human mucus (Yang et al. Angew. Chem. Int. Ed. 2011 50:2597-2600; the contents of which are herein incorporated by reference in its entirety). A non-limiting scalable method to produce nanoparticles which can penetrate human mucus is described by Xu et al. (See e.g., J Control Release 2013, 170(2):279-86; the contents of which are herein incorporated by reference in its entirety).

[000416] The vitamin of the polymer-vitamin conjugate may be vitamin E. The vitamin portion of the conjugate may be substituted with other suitable components such as, but not limited to, vitamin A, vitamin E, other vitamins, cholesterol, a hydrophobic moiety, or a hydrophobic component of other surfactants (e.g., sterol chains, fatty acids, hydrocarbon chains and alkylene oxide chains).

[000417] The lipid nanoparticle engineered to penetrate mucus may include surface altering agents such as, but not limited to, polynucleotides, anionic proteins (e.g., bovine

serum albumin), surfactants (e.g., cationic surfactants such as for example dimethyldioctadecyl-ammonium bromide), sugars or sugar derivatives (e.g., cyclodextrin), nucleic acids, polymers (e.g., heparin, polyethylene glycol and poloxamer), mucolytic agents (e.g., N-acetylcysteine, mugwort, bromelain, papain, clerodendrum, acetylcysteine, bromhexine, carbocysteine, eprazinone, mesna, ambroxol, sobrerol, domiodol, letosteine, stepronin, tiopronin, gelsolin, thymosin β 4 dornase alfa, neltenexine, erdosteine) and various DNases including rhDNase.. The surface altering agent may be embedded or enmeshed in the particle's surface or disposed (e.g., by coating, adsorption, covalent linkage, or other process) on the surface of the lipid nanoparticle. (see e.g., US Publication 20100215580 and US Publication 20080166414 and US20130164343; each of which is herein incorporated by reference in their entirety).

[000418] In one embodiment, the mucus penetrating lipid nanoparticles may comprise at least one polynucleotide described herein. The polynucleotide may be encapsulated in the lipid nanoparticle and/or disposed on the surface of the particle. The polynucleotide may be covalently coupled to the lipid nanoparticle. Formulations of mucus penetrating lipid nanoparticles may comprise a plurality of nanoparticles. Further, the formulations may contain particles which may interact with the mucus and alter the structural and/or adhesive properties of the surrounding mucus to decrease mucoadhesion which may increase the delivery of the mucus penetrating lipid nanoparticles to the mucosal tissue.

[000419] In another embodiment, the mucus penetrating lipid nanoparticles may be a hypotonic formulation comprising a mucosal penetration enhancing coating. The formulation may be hypotonic for the epithelium to which it is being delivered. Non-limiting examples of hypotonic formulations may be found in International Patent Publication No. WO2013110028, the contents of which are herein incorporated by reference in its entirety.

[000420] In one embodiment, in order to enhance the delivery through the mucosal barrier the formulation may comprise or be a hypotonic solution. Hypotonic solutions were found to increase the rate at which muco-inert particles such as, but not limited to, mucus-penetrating particles, were able to reach the vaginal epithelial surface (See e.g., Ensign et al. Biomaterials 2013 34(28):6922-9; the contents of which is herein incorporated by reference in its entirety).

[000421] In one embodiment, the polynucleotide is formulated as a lipoplex, such as, without limitation, the ATUPLEX™ system, the DACC system, the DBTC system and other siRNA-lipoplex technology from Silence Therapeutics (London, United Kingdom), STEMFECT™ from STEMGENT® (Cambridge, MA), and polyethylenimine (PEI) or protamine-based targeted and non-targeted delivery of nucleic acids (Aleku et al. *Cancer Res.* 2008 68:9788-9798; Strumberg et al. *Int J Clin Pharmacol Ther* 2012 50:76-78; Santel et al., *Gene Ther* 2006 13:1222-1234; Santel et al., *Gene Ther* 2006 13:1360-1370; Gutbier et al., *Pulm Pharmacol. Ther.* 2010 23:334-344; Kaufmann et al. *Microvasc Res* 2010 80:286-293; Weide et al. *J Immunother.* 2009 32:498-507; Weide et al. *J Immunother.* 2008 31:180-188; Pascolo *Expert Opin. Biol. Ther.* 4:1285-1294; Fotin-Mleczek et al., 2011 *J. Immunother.* 34:1-15; Song et al., *Nature Biotechnol.* 2005, 23:709-717; Peer et al., *Proc Natl Acad Sci U S A.* 2007 6;104:4095-4100; deFougerolles *Hum Gene Ther.* 2008 19:125-132; all of which are incorporated herein by reference in its entirety).

[000422] In one embodiment such formulations may also be constructed or compositions altered such that they passively or actively are directed to different cell types *in vivo*, including but not limited to hepatocytes, immune cells, tumor cells, endothelial cells, antigen presenting cells, and leukocytes (Akinc et al. *Mol Ther.* 2010 18:1357-1364; Song et al., *Nat Biotechnol.* 2005 23:709-717; Judge et al., *J Clin Invest.* 2009 119:661-673; Kaufmann et al., *Microvasc Res* 2010 80:286-293; Santel et al., *Gene Ther* 2006 13:1222-1234; Santel et al., *Gene Ther* 2006 13:1360-1370; Gutbier et al., *Pulm Pharmacol. Ther.* 2010 23:334-344; Basha et al., *Mol. Ther.* 2011 19:2186-2200; Fenske and Cullis, *Expert Opin Drug Deliv.* 2008 5:25-44; Peer et al., *Science.* 2008 319:627-630; Peer and Lieberman, *Gene Ther.* 2011 18:1127-1133; all of which are incorporated herein by reference in its entirety). One example of passive targeting of formulations to liver cells includes the DLin-DMA, DLin-KC2-DMA and DLin-MC3-DMA-based lipid nanoparticle formulations which have been shown to bind to apolipoprotein E and promote binding and uptake of these formulations into hepatocytes *in vivo* (Akinc et al. *Mol Ther.* 2010 18:1357-1364; herein incorporated by reference in its entirety). Formulations can also be selectively targeted through expression of different ligands on their surface as exemplified by, but not limited by, folate, transferrin,

N-acetylgalactosamine (GalNAc), and antibody targeted approaches (Kolhatkar et al., *Curr Drug Discov Technol.* 2011 8:197-206; Musacchio and Torchilin, *Front Biosci.* 2011 16:1388-1412; Yu et al., *Mol Membr Biol.* 2010 27:286-298; Patil et al., *Crit Rev Ther Drug Carrier Syst.* 2008 25:1-61; Benoit et al., *Biomacromolecules.* 2011 12:2708-2714; Zhao et al., *Expert Opin Drug Deliv.* 2008 5:309-319; Akinc et al., *Mol Ther.* 2010 18:1357-1364; Srinivasan et al., *Methods Mol Biol.* 2012 820:105-116; Ben-Arie et al., *Methods Mol Biol.* 2012 757:497-507; Peer 2010 *J Control Release.* 20:63-68; Peer et al., *Proc Natl Acad Sci U S A.* 2007 104:4095-4100; Kim et al., *Methods Mol Biol.* 2011 721:339-353; Subramanya et al., *Mol Ther.* 2010 18:2028-2037; Song et al., *Nat Biotechnol.* 2005 23:709-717; Peer et al., *Science.* 2008 319:627-630; Peer and Lieberman, *Gene Ther.* 2011 18:1127-1133; all of which are incorporated herein by reference in its entirety).

[000423] In one embodiment, the polynucleotide is formulated as a solid lipid nanoparticle. A solid lipid nanoparticle (SLN) may be spherical with an average diameter between 10 to 1000 nm. SLN possess a solid lipid core matrix that can solubilize lipophilic molecules and may be stabilized with surfactants and/or emulsifiers. In a further embodiment, the lipid nanoparticle may be a self-assembly lipid-polymer nanoparticle (see Zhang et al., *ACS Nano*, 2008, 2 (8), pp 1696–1702; the contents of which are herein incorporated by reference in its entirety). As a non-limiting example, the SLN may be the SLN described in International Patent Publication No. WO2013105101, the contents of which are herein incorporated by reference in its entirety. As another non-limiting example, the SLN may be made by the methods or processes described in International Patent Publication No. WO2013105101, the contents of which are herein incorporated by reference in its entirety.

[000424] Liposomes, lipoplexes, or lipid nanoparticles may be used to improve the efficacy of polynucleotides directed protein production as these formulations may be able to increase cell transfection by the polynucleotide; and/or increase the translation of encoded protein. One such example involves the use of lipid encapsulation to enable the effective systemic delivery of polyplex plasmid DNA (Heyes et al., *Mol Ther.* 2007 15:713-720; herein incorporated by reference in its entirety). The liposomes, lipoplexes, or lipid nanoparticles may also be used to increase the stability of the polynucleotide.

[000425] In one embodiment, the polynucleotides of the present invention can be formulated for controlled release and/or targeted delivery. As used herein, “controlled release” refers to a pharmaceutical composition or compound release profile that conforms to a particular pattern of release to effect a therapeutic outcome. In one embodiment, the polynucleotides may be encapsulated into a delivery agent described herein and/or known in the art for controlled release and/or targeted delivery. As used herein, the term “encapsulate” means to enclose, surround or encase. As it relates to the formulation of the compounds of the invention, encapsulation may be substantial, complete or partial. The term “substantially encapsulated” means that at least greater than 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.9 or greater than 99.999% of the pharmaceutical composition or compound of the invention may be enclosed, surrounded or encased within the delivery agent. “Partially encapsulation” means that less than 10, 10, 20, 30, 40 50 or less of the pharmaceutical composition or compound of the invention may be enclosed, surrounded or encased within the delivery agent. Advantageously, encapsulation may be determined by measuring the escape or the activity of the pharmaceutical composition or compound of the invention using fluorescence and/or electron micrograph. For example, at least 1, 5, 10, 20, 30, 40, 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.99 or greater than 99.99% of the pharmaceutical composition or compound of the invention are encapsulated in the delivery agent.

[000426] In one embodiment, the controlled release formulation may include, but is not limited to, tri-block co-polymers. As a non-limiting example, the formulation may include two different types of tri-block co-polymers (International Pub. No. WO2012131104 and WO2012131106; each of which is herein incorporated by reference in its entirety).

[000427] In another embodiment, the polynucleotides may be encapsulated into a lipid nanoparticle or a rapidly eliminated lipid nanoparticle and the lipid nanoparticles or a rapidly eliminated lipid nanoparticle may then be encapsulated into a polymer, hydrogel and/or surgical sealant described herein and/or known in the art. As a non-limiting example, the polymer, hydrogel or surgical sealant may be PLGA, ethylene vinyl acetate (EVAc), poloxamer, GELSITE® (Nanotherapeutics, Inc. Alachua, FL), HYLENEX® (Halozyme Therapeutics, San Diego CA), surgical sealants such as fibrinogen polymers

(Ethicon Inc. Cornelia, GA), TISSELL® (Baxter International, Inc Deerfield, IL), PEG-based sealants, and COSEAL® (Baxter International, Inc Deerfield, IL).

[000428] In another embodiment, the lipid nanoparticle may be encapsulated into any polymer known in the art which may form a gel when injected into a subject. As another non-limiting example, the lipid nanoparticle may be encapsulated into a polymer matrix which may be biodegradable.

[000429] In one embodiment, the polynucleotide formulation for controlled release and/or targeted delivery may also include at least one controlled release coating. Controlled release coatings include, but are not limited to, OPADRY®, polyvinylpyrrolidone/vinyl acetate copolymer, polyvinylpyrrolidone, hydroxypropyl methylcellulose, hydroxypropyl cellulose, hydroxyethyl cellulose, EUDRAGIT RL®, EUDRAGIT RS® and cellulose derivatives such as ethylcellulose aqueous dispersions (AQUACOAT® and SURELEASE®).

[000430] In one embodiment, the controlled release and/or targeted delivery formulation may comprise at least one degradable polyester which may contain polycationic side chains. Degradable polyesters include, but are not limited to, poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), and combinations thereof. In another embodiment, the degradable polyesters may include a PEG conjugation to form a PEGylated polymer.

[000431] In one embodiment, the controlled release and/or targeted delivery formulation comprising at least one polynucleotide may comprise at least one PEG and/or PEG related polymer derivatives as described in US Patent No. 8,404,222, herein incorporated by reference in its entirety.

[000432] In another embodiment, the controlled release delivery formulation comprising at least one polynucleotide may be the controlled release polymer system described in US20130130348, herein incorporated by reference in its entirety.

[000433] In one embodiment, the polynucleotides of the present invention may be encapsulated in a therapeutic nanoparticle. Therapeutic nanoparticles may be formulated by methods described herein and known in the art such as, but not limited to, International Pub Nos. WO2010005740, WO2010030763, WO2010005721, WO2010005723, WO2012054923, US Pub. Nos. US20110262491, US20100104645,

US20100087337, US20100068285, US20110274759, US20100068286, US20120288541, US20130123351 and US20130230567 and US Pat No. 8,206,747, 8,293,276, 8,318,208 and 8,318,211; the contents of each of which are herein incorporated by reference in their entirety. In another embodiment, therapeutic polymer nanoparticles may be identified by the methods described in US Pub No. US20120140790, herein incorporated by reference in its entirety.

[000434] In one embodiment, the therapeutic nanoparticle may be formulated for sustained release. As used herein, “sustained release” refers to a pharmaceutical composition or compound that conforms to a release rate over a specific period of time. The period of time may include, but is not limited to, hours, days, weeks, months and years. As a non-limiting example, the sustained release nanoparticle may comprise a polymer and a therapeutic agent such as, but not limited to, the polynucleotides of the present invention (see International Pub No. 2010075072 and US Pub No. US20100216804, US20110217377 and US20120201859, each of which is herein incorporated by reference in their entirety). In another non-limiting example, the sustained release formulation may comprise agents which permit persistent bioavailability such as, but not limited to, crystals, macromolecular gels and/or particulate suspensions (see US Patent Publication No US20130150295, the contents of which is herein incorporated by reference in its entirety).

[000435] In one embodiment, the therapeutic nanoparticles may be formulated to be target specific. As a non-limiting example, the therapeutic nanoparticles may include a corticosteroid (see International Pub. No. WO2011084518; herein incorporated by reference in its entirety). In one embodiment, the therapeutic nanoparticles may be formulated to be cancer specific. As a non-limiting example, the therapeutic nanoparticles may be formulated in nanoparticles described in International Pub No. WO2008121949, WO2010005726, WO2010005725, WO2011084521 and US Pub No. US20100069426, US20120004293 and US20100104655, each of which is herein incorporated by reference in their entirety.

[000436] In one embodiment, the nanoparticles of the present invention may comprise a polymeric matrix. As a non-limiting example, the nanoparticle may comprise two or more polymers such as, but not limited to, polyethylenes, polycarbonates,

polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polylysine, poly(ethylene imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester) or combinations thereof.

[000437] In one embodiment, the therapeutic nanoparticle comprises a diblock copolymer. In one embodiment, the diblock copolymer may include PEG in combination with a polymer such as, but not limited to, polyethylenes, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polylysine, poly(ethylene imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester) or combinations thereof. In another embodiment, the diblock copolymer may comprise the diblock copolymers described in European Patent Publication No. the contents of which are herein incorporated by reference in its entirety. In yet another embodiment, the diblock copolymer may be a high-X diblock copolymer such as those described in International Patent Publication No. WO2013120052, the contents of which are herein incorporated by reference in its entirety.

[000438] As a non-limiting example the therapeutic nanoparticle comprises a PLGA-PEG block copolymer (see US Pub. No. US20120004293 and US Pat No. 8,236,330, each of which is herein incorporated by reference in their entirety). In another non-limiting example, the therapeutic nanoparticle is a stealth nanoparticle comprising a diblock copolymer of PEG and PLA or PEG and PLGA (see US Pat No 8,246,968 and International Publication No. WO2012166923, the contents of each of which are herein incorporated by reference in its entirety). In yet another non-limiting example, the therapeutic nanoparticle is a stealth nanoparticle or a target-specific stealth nanoparticle as described in US Patent Publication No. US20130172406, the contents of which are herein incorporated by reference in its entirety.

[000439] In one embodiment, the therapeutic nanoparticle may comprise a multiblock copolymer (See e.g., U.S. Pat. No. 8,263,665 and 8,287,910 and US Patent Pub. No. US20130195987; the contents of each of which are herein incorporated by reference in its entirety).

[000440] In yet another non-limiting example, the lipid nanoparticle comprises the block copolymer PEG-PLGA-PEG (see e.g., the thermosensitive hydrogel (PEG-PLGA-PEG) was used as a TGF-beta1 gene delivery vehicle in Lee et al. Thermosensitive Hydrogel as a Tgf- β 1 Gene Delivery Vehicle Enhances Diabetic Wound Healing. Pharmaceutical Research, 2003 20(12): 1995-2000; as a controlled gene delivery system in Li et al. Controlled Gene Delivery System Based on Thermosensitive Biodegradable Hydrogel. Pharmaceutical Research 2003 20(6):884-888; and Chang et al., Non-ionic amphiphilic biodegradable PEG-PLGA-PEG copolymer enhances gene delivery efficiency in rat skeletal muscle. J Controlled Release. 2007 118:245-253; each of which is herein incorporated by reference in its entirety). The polynucleotides of the present invention may be formulated in lipid nanoparticles comprising the PEG-PLGA-PEG block copolymer.

[000441] In one embodiment, the therapeutic nanoparticle may comprise a multiblock copolymer (See e.g., U.S. Pat. No. 8,263,665 and 8,287,910 and US Patent Pub. No. US20130195987; the contents of each of which are herein incorporated by reference in its entirety).

[000442] In one embodiment, the block copolymers described herein may be included in a polyion complex comprising a non-polymeric micelle and the block copolymer. (See e.g., U.S. Pub. No. 20120076836; herein incorporated by reference in its entirety).

[000443] In one embodiment, the therapeutic nanoparticle may comprise at least one acrylic polymer. Acrylic polymers include but are not limited to, acrylic acid, methacrylic acid, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, amino alkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), polycyanoacrylates and combinations thereof.

[000444] In one embodiment, the therapeutic nanoparticles may comprise at least one poly(vinyl ester) polymer. The poly(vinyl ester) polymer may be a copolymer such as a

random copolymer. As a non-limiting example, the random copolymer may have a structure such as those described in International Application No. WO2013032829 or US Patent Publication No US20130121954, the contents of which are herein incorporated by reference in its entirety. In one aspect, the poly(vinyl ester) polymers may be conjugated to the polynucleotides described herein. In another aspect, the poly(vinyl ester) polymer which may be used in the present invention may be those described in, herein incorporated by reference in its entirety.

[000445] In one embodiment, the therapeutic nanoparticle may comprise at least one diblock copolymer. The diblock copolymer may be, but it not limited to, a poly(lactic acid-poly(ethylene)glycol copolymer (see e.g., International Patent Publication No. WO2013044219; herein incorporated by reference in its entirety). As a non-limiting example, the therapeutic nanoparticle may be used to treat cancer (see International publication No. WO2013044219; herein incorporated by reference in its entirety).

[000446] In one embodiment, the therapeutic nanoparticles may comprise at least one cationic polymer described herein and/or known in the art.

[000447] In one embodiment, the therapeutic nanoparticles may comprise at least one amine-containing polymer such as, but not limited to polylysine, polyethylene imine, poly(amidoamine) dendrimers, poly(beta-amino esters) (See e.g., U.S. Pat. No. 8,287,849; herein incorporated by reference in its entirety) and combinations thereof.

[000448] In another embodiment, the nanoparticles described herein may comprise an amine cationic lipid such as those described in International Patent Application No. WO2013059496, the contents of which are herein incorporated by reference in its entirety. In one aspect the cationic lipids may have an amino-amine or an amino-amide moiety.

[000449] In one embodiment, the therapeutic nanoparticles may comprise at least one degradable polyester which may contain polycationic side chains. Degradable polyesters include, but are not limited to, poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), and combinations thereof. In another embodiment, the degradable polyesters may include a PEG conjugation to form a PEGylated polymer.

[000450] In another embodiment, the therapeutic nanoparticle may include a conjugation of at least one targeting ligand. The targeting ligand may be any ligand

known in the art such as, but not limited to, a monoclonal antibody. (Kirpotin et al, Cancer Res. 2006 66:6732-6740; herein incorporated by reference in its entirety).

[000451] In one embodiment, the therapeutic nanoparticle may be formulated in an aqueous solution which may be used to target cancer (see International Pub No. WO2011084513 and US Pub No. US20110294717, each of which is herein incorporated by reference in their entirety).

[000452] In one embodiment, the therapeutic nanoparticle comprising at least one polynucleotide may be formulated using the methods described by Podobinski et al in US Patent No. 8,404,799, the contents of which are herein incorporated by reference in its entirety.

[000453] In one embodiment, the polynucleotides may be encapsulated in, linked to and/or associated with synthetic nanocarriers. Synthetic nanocarriers include, but are not limited to, those described in International Pub. Nos. WO2010005740, WO2010030763, WO201213501, WO2012149252, WO2012149255, WO2012149259, WO2012149265, WO2012149268, WO2012149282, WO2012149301, WO2012149393, WO2012149405, WO2012149411, WO2012149454 and WO2013019669, and US Pub. Nos. US20110262491, US20100104645, US20100087337 and US20120244222, each of which is herein incorporated by reference in their entirety. The synthetic nanocarriers may be formulated using methods known in the art and/or described herein. As a non-limiting example, the synthetic nanocarriers may be formulated by the methods described in International Pub Nos. WO2010005740, WO2010030763 and WO201213501 and US Pub. Nos. US20110262491, US20100104645, US20100087337 and US2012024422, each of which is herein incorporated by reference in their entirety. In another embodiment, the synthetic nanocarrier formulations may be lyophilized by methods described in International Pub. No. WO2011072218 and US Pat No. 8,211,473; the content of each of which is herein incorporated by reference in their entirety. In yet another embodiment, formulations of the present invention, including, but not limited to, synthetic nanocarriers, may be lyophilized or reconstituted by the methods described in US Patent Publication No. US20130230568, the contents of which are herein incorporated by reference in its entirety.

[000454] In one embodiment, the synthetic nanocarriers may contain reactive groups to release the polynucleotides described herein (see International Pub. No. WO20120952552 and US Pub No. US20120171229, each of which is herein incorporated by reference in their entirety).

[000455] In one embodiment, the synthetic nanocarriers may contain an immunostimulatory agent to enhance the immune response from delivery of the synthetic nanocarrier. As a non-limiting example, the synthetic nanocarrier may comprise a Th1 immunostimulatory agent which may enhance a Th1-based response of the immune system (see International Pub No. WO2010123569 and US Pub. No. US20110223201, each of which is herein incorporated by reference in its entirety).

[000456] In one embodiment, the synthetic nanocarriers may be formulated for targeted release. In one embodiment, the synthetic nanocarrier is formulated to release the polynucleotides at a specified pH and/or after a desired time interval. As a non-limiting example, the synthetic nanoparticle may be formulated to release the polynucleotides after 24 hours and/or at a pH of 4.5 (see International Pub. Nos. WO2010138193 and WO2010138194 and US Pub Nos. US20110020388 and US20110027217, each of which is herein incorporated by reference in their entirety).

[000457] In one embodiment, the synthetic nanocarriers may be formulated for controlled and/or sustained release of the polynucleotides described herein. As a non-limiting example, the synthetic nanocarriers for sustained release may be formulated by methods known in the art, described herein and/or as described in International Pub No. WO2010138192 and US Pub No. 20100303850, each of which is herein incorporated by reference in their entirety.

[000458] In one embodiment, the polynucleotides may be formulated for controlled and/or sustained release wherein the formulation comprises at least one polymer that is a crystalline side chain (CYSC) polymer. CYSC polymers are described in U.S. Patent No. 8,399,007, herein incorporated by reference in its entirety.

[000459] In one embodiment, the synthetic nanocarrier may be formulated for use as a vaccine. In one embodiment, the synthetic nanocarrier may encapsulate at least one polynucleotide which encode at least one antigen. As a non-limiting example, the synthetic nanocarrier may include at least one antigen and an excipient for a vaccine

dosage form (see International Pub No. WO2011150264 and US Pub No. US20110293723, each of which is herein incorporated by reference in their entirety). As another non-limiting example, a vaccine dosage form may include at least two synthetic nanocarriers with the same or different antigens and an excipient (see International Pub No. WO2011150249 and US Pub No. US20110293701, each of which is herein incorporated by reference in their entirety). The vaccine dosage form may be selected by methods described herein, known in the art and/or described in International Pub No. WO2011150258 and US Pub No. US20120027806, each of which is herein incorporated by reference in their entirety).

[000460] In one embodiment, the synthetic nanocarrier may comprise at least one polynucleotide which encodes at least one adjuvant. As non-limiting example, the adjuvant may comprise dimethyldioctadecylammonium-bromide, dimethyldioctadecylammonium-chloride, dimethyldioctadecylammonium-phosphate or dimethyldioctadecylammonium-acetate (DDA) and an apolar fraction or part of said apolar fraction of a total lipid extract of a mycobacterium (See e.g., U.S. Pat. No. 8,241,610; herein incorporated by reference in its entirety). In another embodiment, the synthetic nanocarrier may comprise at least one polynucleotide and an adjuvant. As a non-limiting example, the synthetic nanocarrier comprising and adjuvant may be formulated by the methods described in International Pub No. WO2011150240 and US Pub No. US20110293700, each of which is herein incorporated by reference in its entirety.

[000461] In one embodiment, the synthetic nanocarrier may encapsulate at least one polynucleotide which encodes a peptide, fragment or region from a virus. As a non-limiting example, the synthetic nanocarrier may include, but is not limited to, the nanocarriers described in International Pub No. WO2012024621, WO201202629, WO2012024632 and US Pub No. US20120064110, US20120058153 and US20120058154, each of which is herein incorporated by reference in their entirety.

[000462] In one embodiment, the synthetic nanocarrier may be coupled to a polynucleotide which may be able to trigger a humoral and/or cytotoxic T lymphocyte (CTL) response (See e.g., International Publication No. WO2013019669, herein incorporated by reference in its entirety).

[000463] In one embodiment, the polynucleotides may be encapsulated in, linked to and/or associated with zwitterionic lipids. Non-limiting examples of zwitterionic lipids and methods of using zwitterionic lipids are described in US Patent Publication No. US20130216607, the contents of which are herein incorporated by reference in its entirety. In one aspect, the zwitterionic lipids may be used in the liposomes and lipid nanoparticles described herein.

[000464] In one embodiment, the polynucleotides may be formulated in colloid nanocarriers as described in US Patent Publication No. US20130197100, the contents of which are herein incorporated by reference in its entirety.

[000465] In one embodiment, the nanoparticle may be optimized for oral administration. The nanoparticle may comprise at least one cationic biopolymer such as, but not limited to, chitosan or a derivative thereof. As a non-limiting example, the nanoparticle may be formulated by the methods described in U.S. Pub. No. 20120282343; herein incorporated by reference in its entirety.

[000466] In some embodiments, LNPs comprise the lipid KL52 (an amino-lipid disclosed in U.S. Application Publication No. 2012/0295832 expressly incorporated herein by reference in its entirety). Activity and/or safety (as measured by examining one or more of ALT/AST, white blood cell count and cytokine induction) of LNP administration may be improved by incorporation of such lipids. LNPs comprising KL52 may be administered intravenously and/or in one or more doses. In some embodiments, administration of LNPs comprising KL52 results in equal or improved mRNA and/or protein expression as compared to LNPs comprising MC3.

[000467] In some embodiments, polynucleotides may be delivered using smaller LNPs. Such particles may comprise a diameter from below 0.1 μm up to 100 nm such as, but not limited to, less than 0.1 μm , less than 1.0 μm , less than 5 μm , less than 10 μm , less than 15 μm , less than 20 μm , less than 25 μm , less than 30 μm , less than 35 μm , less than 40 μm , less than 50 μm , less than 55 μm , less than 60 μm , less than 65 μm , less than 70 μm , less than 75 μm , less than 80 μm , less than 85 μm , less than 90 μm , less than 95 μm , less than 100 μm , less than 125 μm , less than 150 μm , less than 175 μm , less than 200 μm , less than 225 μm , less than 250 μm , less than 275 μm , less than 300 μm , less than 325 μm , less than 350 μm , less than 375 μm , less than 400 μm , less than 425 μm , less than 450 μm , less

than 475 um, less than 500 um, less than 525 um, less than 550 um, less than 575 um, less than 600 um, less than 625 um, less than 650 um, less than 675 um, less than 700 um, less than 725 um, less than 750 um, less than 775 um, less than 800 um, less than 825 um, less than 850 um, less than 875 um, less than 900 um, less than 925 um, less than 950 um, less than 975 um,

[000468] In another embodiment, polynucleotides may be delivered using smaller LNPs which may comprise a diameter from about 1 nm to about 100 nm, from about 1 nm to about 10 nm, about 1 nm to about 20 nm, from about 1 nm to about 30 nm, from about 1 nm to about 40 nm, from about 1 nm to about 50 nm, from about 1 nm to about 60 nm, from about 1 nm to about 70 nm, from about 1 nm to about 80 nm, from about 1 nm to about 90 nm, from about 5 nm to about 100 nm, from about 5 nm to about 10 nm, about 5 nm to about 20 nm, from about 5 nm to about 30 nm, from about 5 nm to about 40 nm, from about 5 nm to about 50 nm, from about 5 nm to about 60 nm, from about 5 nm to about 70 nm, from about 5 nm to about 80 nm, from about 5 nm to about 90 nm, about 10 to about 50 nm, from about 20 to about 50 nm, from about 30 to about 50 nm, from about 40 to about 50 nm, from about 20 to about 60 nm, from about 30 to about 60 nm, from about 40 to about 60 nm, from about 20 to about 70 nm, from about 30 to about 70 nm, from about 40 to about 70 nm, from about 50 to about 70 nm, from about 60 to about 70 nm, from about 20 to about 80 nm, from about 30 to about 80 nm, from about 40 to about 80 nm, from about 50 to about 80 nm, from about 60 to about 80 nm, from about 20 to about 90 nm, from about 30 to about 90 nm, from about 40 to about 90 nm, from about 50 to about 90 nm, from about 60 to about 90 nm and/or from about 70 to about 90 nm.

[000469] In some embodiments, such LNPs are synthesized using methods comprising microfluidic mixers. Exemplary microfluidic mixers may include, but are not limited to a slit interdigital micromixer including, but not limited to those manufactured by Microinnova (Allerheiligen bei Wildon, Austria) and/or a staggered herringbone micromixer (SHM) (Zhigaltsev, I.V. et al., Bottom-up design and synthesis of limit size lipid nanoparticle systems with aqueous and triglyceride cores using millisecond microfluidic mixing have been published (Langmuir. 2012. 28:3633-40; Belliveau, N.M. et al., Microfluidic synthesis of highly potent limit-size lipid nanoparticles for in vivo

delivery of siRNA. *Molecular Therapy-Nucleic Acids*. 2012. 1:e37; Chen, D. et al., Rapid discovery of potent siRNA-containing lipid nanoparticles enabled by controlled microfluidic formulation. *J Am Chem Soc*. 2012. 134(16):6948-51; each of which is herein incorporated by reference in its entirety). In some embodiments, methods of LNP generation comprising SHM, further comprise the mixing of at least two input streams wherein mixing occurs by microstructure-induced chaotic advection (MICA). According to this method, fluid streams flow through channels present in a herringbone pattern causing rotational flow and folding the fluids around each other. This method may also comprise a surface for fluid mixing wherein the surface changes orientations during fluid cycling. Methods of generating LNPs using SHM include those disclosed in U.S. Application Publication Nos. 2004/0262223 and 2012/0276209, each of which is expressly incorporated herein by reference in their entirety.

[000470] In one embodiment, the polynucleotides of the present invention may be formulated in lipid nanoparticles created using a micromixer such as, but not limited to, a Slit Interdigital Microstructured Mixer (SIMM-V2) or a Standard Slit Interdigital Micro Mixer (SSIMM) or Caterpillar (CPMM) or Impinging-jet (IJMM) from the Institut für Mikrotechnik Mainz GmbH, Mainz Germany).

[000471] In one embodiment, the polynucleotides of the present invention may be formulated in lipid nanoparticles created using microfluidic technology (see Whitesides, George M. The Origins and the Future of Microfluidics. *Nature*, 2006 442: 368-373; and Abraham et al. Chaotic Mixer for Microchannels. *Science*, 2002 295: 647-651; each of which is herein incorporated by reference in its entirety). As a non-limiting example, controlled microfluidic formulation includes a passive method for mixing streams of steady pressure-driven flows in micro channels at a low Reynolds number (See e.g., Abraham et al. Chaotic Mixer for Microchannels. *Science*, 2002 295: 647-651; which is herein incorporated by reference in its entirety).

[000472] In one embodiment, the polynucleotides of the present invention may be formulated in lipid nanoparticles created using a micromixer chip such as, but not limited to, those from Harvard Apparatus (Holliston, MA) or Dolomite Microfluidics (Royston, UK). A micromixer chip can be used for rapid mixing of two or more fluid streams with a split and recombine mechanism.

[000473] In one embodiment, the polynucleotides of the invention may be formulated for delivery using the drug encapsulating microspheres described in International Patent Publication No. WO2013063468 or U.S. Patent No. 8,440,614, each of which is herein incorporated by reference in its entirety. The microspheres may comprise a compound of the formula (I), (II), (III), (IV), (V) or (VI) as described in International patent application WO2013063468, the contents of which are herein incorporated by reference in its entirety. In another aspect, the amino acid, peptide, polypeptide, lipids (APPL) are useful in delivering the polynucleotides of the invention to cells (see International Patent Publication No. WO2013063468, herein incorporated by reference in its entirety).

[000474] In one embodiment, the polynucleotides of the invention may be formulated in lipid nanoparticles having a diameter from about 10 to about 100 nm such as, but not limited to, about 10 to about 20 nm, about 10 to about 30 nm, about 10 to about 40 nm, about 10 to about 50 nm, about 10 to about 60 nm, about 10 to about 70 nm, about 10 to about 80 nm, about 10 to about 90 nm, about 20 to about 30 nm, about 20 to about 40 nm, about 20 to about 50 nm, about 20 to about 60 nm, about 20 to about 70 nm, about 20 to about 80 nm, about 20 to about 90 nm, about 20 to about 100 nm, about 30 to about 40 nm, about 30 to about 50 nm, about 30 to about 60 nm, about 30 to about 70 nm, about 30 to about 80 nm, about 30 to about 90 nm, about 30 to about 100 nm, about 40 to about 50 nm, about 40 to about 60 nm, about 40 to about 70 nm, about 40 to about 80 nm, about 40 to about 90 nm, about 40 to about 100 nm, about 50 to about 60 nm, about 50 to about 70 nm, about 50 to about 80 nm, about 50 to about 90 nm, about 50 to about 100 nm, about 60 to about 70 nm, about 60 to about 80 nm, about 60 to about 90 nm, about 60 to about 100 nm, about 70 to about 80 nm, about 70 to about 90 nm, about 70 to about 100 nm, about 80 to about 90 nm, about 80 to about 100 nm and/or about 90 to about 100 nm.

[000475] In one embodiment, the lipid nanoparticles may have a diameter from about 10 to 500 nm.

[000476] In one embodiment, the lipid nanoparticle may have a diameter greater than 100 nm, greater than 150 nm, greater than 200 nm, greater than 250 nm, greater than 300 nm, greater than 350 nm, greater than 400 nm, greater than 450 nm, greater than 500 nm, greater than 550 nm, greater than 600 nm, greater than 650 nm, greater than 700 nm,

greater than 750 nm, greater than 800 nm, greater than 850 nm, greater than 900 nm, greater than 950 nm or greater than 1000 nm.

[000477] In one aspect, the lipid nanoparticle may be a limit size lipid nanoparticle described in International Patent Publication No. WO2013059922, the contents of which are herein incorporated by reference in its entirety. The limit size lipid nanoparticle may comprise a lipid bilayer surrounding an aqueous core or a hydrophobic core; where the lipid bilayer may comprise a phospholipid such as, but not limited to, diacylphosphatidylcholine, a diacylphosphatidylethanolamine, a ceramide, a sphingomyelin, a dihydrosphingomyelin, a cephalin, a cerebroside, a C8-C20 fatty acid diacylphosphatidylcholine, and 1-palmitoyl-2-oleoyl phosphatidylcholine (POPC). In another aspect the limit size lipid nanoparticle may comprise a polyethylene glycol-lipid such as, but not limited to, DLPE-PEG, DMPE-PEG, DPPC-PEG and DSPE-PEG.

[000478] In one embodiment, the polynucleotides may be delivered, localized and/or concentrated in a specific location using the delivery methods described in International Patent Publication No. WO2013063530, the contents of which are herein incorporated by reference in its entirety. As a non-limiting example, a subject may be administered an empty polymeric particle prior to, simultaneously with or after delivering the polynucleotides to the subject. The empty polymeric particle undergoes a change in volume once in contact with the subject and becomes lodged, embedded, immobilized or entrapped at a specific location in the subject.

[000479] In one embodiment, the polynucleotides may be formulated in an active substance release system (See e.g., US Patent Publication No. US20130102545, herein incorporated by reference in its entirety). The active substance release system may comprise 1) at least one nanoparticle bonded to an oligonucleotide inhibitor strand which is hybridized with a catalytically active nucleic acid and 2) a compound bonded to at least one substrate molecule bonded to a therapeutically active substance (e.g., polynucleotides described herein), where the therapeutically active substance is released by the cleavage of the substrate molecule by the catalytically active nucleic acid.

[000480] In one embodiment, the polynucleotides may be formulated in a nanoparticle comprising an inner core comprising a non-cellular material and an outer surface comprising a cellular membrane. The cellular membrane may be derived from a cell or a

membrane derived from a virus. As a non-limiting example, the nanoparticle may be made by the methods described in International Patent Publication No. WO2013052167, herein incorporated by reference in its entirety. As another non-limiting example, the nanoparticle described in International Patent Publication No. WO2013052167, herein incorporated by reference in its entirety, may be used to deliver the polynucleotides described herein.

[000481] In one embodiment, the polynucleotides may be formulated in porous nanoparticle-supported lipid bilayers (protocells). Protocells are described in International Patent Publication No. WO2013056132, the contents of which are herein incorporated by reference in its entirety.

[000482] In one embodiment, the polynucleotides described herein may be formulated in polymeric nanoparticles as described in or made by the methods described in US Patent No. 8,420,123 and 8,518,963 and European Patent No. EP2073848B1, the contents of each of which are herein incorporated by reference in their entirety. As a non-limiting example, the polymeric nanoparticle may have a high glass transition temperature such as the nanoparticles described in or nanoparticles made by the methods described in US Patent No. 8,518,963, the contents of which are herein incorporated by reference in its entirety. As another non-limiting example, the polymer nanoparticle for oral, parenteral and topical formulations may be made by the methods described in European Patent No. EP2073848B1, the contents of which are herein incorporated by reference in its entirety.

[000483] In another embodiment, the polynucleotides described herein may be formulated in nanoparticles used in imaging. The nanoparticles may be liposome nanoparticles such as those described in US Patent Publication No US20130129636, herein incorporated by reference in its entirety. As a non-limiting example, the liposome may comprise gadolinium(III)2-{4,7-bis-carboxymethyl-10-[(N,N-distearylamidomethyl-N'-amido-methyl]-1,4,7,10-tetra-azacyclododec-1-yl}-acetic acid and a neutral, fully saturated phospholipid component (see e.g., US Patent Publication No US20130129636, the contents of which is herein incorporated by reference in its entirety).

[000484] In one embodiment, the nanoparticles which may be used in the present invention are formed by the methods described in U.S. Patent Application No. US20130130348, the contents of which is herein incorporated by reference in its entirety.

[000485] The nanoparticles of the present invention may further include nutrients such as, but not limited to, those which deficiencies can lead to health hazards from anemia to neural tube defects (see e.g., the nanoparticles described in International Patent Publication No WO2013072929, the contents of which is herein incorporated by reference in its entirety). As a non-limiting example, the nutrient may be iron in the form of ferrous, ferric salts or elemental iron, iodine, folic acid, vitamins or micronutrients.

[000486] In one embodiment, the polynucleotides of the present invention may be formulated in a swellable nanoparticle. The swellable nanoparticle may be, but is not limited to, those described in U.S. Patent No. 8,440,231, the contents of which is herein incorporated by reference in its entirety. As a non-limiting embodiment, the swellable nanoparticle may be used for delivery of the polynucleotides of the present invention to the pulmonary system (see e.g., U.S. Patent No. 8,440,231, the contents of which is herein incorporated by reference in its entirety).

[000487] The polynucleotides of the present invention may be formulated in polyanhydride nanoparticles such as, but not limited to, those described in U.S. Patent No. 8,449,916, the contents of which is herein incorporated by reference in its entirety.

[000488] The nanoparticles and microparticles of the present invention may be geometrically engineered to modulate macrophage and/or the immune response. In one aspect, the geometrically engineered particles may have varied shapes, sizes and/or surface charges in order to incorporated the polynucleotides of the present invention for targeted delivery such as, but not limited to, pulmonary delivery (see e.g., International Publication No WO2013082111, the contents of which is herein incorporated by reference in its entirety). Other physical features the geometrically engineering particles may have include, but are not limited to, fenestrations, angled arms, asymmetry and surface roughness, charge which can alter the interactions with cells and tissues. As a non-limiting example, nanoparticles of the present invention may be made by the methods described in International Publication No WO2013082111, the contents of which is herein incorporated by reference in its entirety.

[000489] In one embodiment, the nanoparticles of the present invention may be water soluble nanoparticles such as, but not limited to, those described in International Publication No. WO2013090601, the contents of which is herein incorporated by reference in its entirety. The nanoparticles may be inorganic nanoparticles which have a compact and zwitterionic ligand in order to exhibit good water solubility. The nanoparticles may also have small hydrodynamic diameters (HD), stability with respect to time, pH, and salinity and a low level of non-specific protein binding.

[000490] In one embodiment the nanoparticles of the present invention may be developed by the methods described in US Patent Publication No. US20130172406, the contents of which are herein incorporated by reference in its entirety.

[000491] In one embodiment, the nanoparticles of the present invention are stealth nanoparticles or target-specific stealth nanoparticles such as, but not limited to, those described in US Patent Publication No. US20130172406; the contents of which is herein incorporated by reference in its entirety. The nanoparticles of the present invention may be made by the methods described in US Patent Publication No. US20130172406, the contents of which are herein incorporated by reference in its entirety.

[000492] In another embodiment, the stealth or target-specific stealth nanoparticles may comprise a polymeric matrix. The polymeric matrix may comprise two or more polymers such as, but not limited to, polyethylenes, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polyesters, polyanhydrides, polyethers, polyurethanes, polymethacrylates, polyacrylates, polycyanoacrylates or combinations thereof.

[000493] In one embodiment, the nanoparticle may be a nanoparticle-nucleic acid hybrid structure having a high density nucleic acid layer. As a non-limiting example, the nanoparticle-nucleic acid hybrid structure may be made by the methods described in US Patent Publication No. US20130171646, the contents of which are herein incorporated by reference in its entirety. The nanoparticle may comprise a nucleic acid such as, but not limited to, polynucleotides described herein and/or known in the art.

[000494] At least one of the nanoparticles of the present invention may be embedded in the core a nanostructure or coated with a low density porous 3-D structure or coating which is capable of carrying or associating with at least one payload within or on the surface of the nanostructure. Non-limiting examples of the nanostructures comprising at least one nanoparticle are described in International Patent Publication No.

WO2013123523, the contents of which are herein incorporated by reference in its entirety.

Polymers, Biodegradable Nanoparticles, and Core-Shell Nanoparticles

[000495] The polynucleotides of the invention can be formulated using natural and/or synthetic polymers. Non-limiting examples of polymers which may be used for delivery include, but are not limited to, DYNAMIC POLYCONJUGATE® (Arrowhead Research Corp., Pasadena, CA) formulations from MIRUS® Bio (Madison, WI) and Roche Madison (Madison, WI), PHASERX™ polymer formulations such as, without limitation, SMARTT POLYMER TECHNOLOGY™ (PHASERX®, Seattle, WA), DMRI/DOPE, poloxamer, VAXFECTIN® adjuvant from Vical (San Diego, CA), chitosan, cyclodextrin from Calando Pharmaceuticals (Pasadena, CA), dendrimers and poly(lactic-co-glycolic acid) (PLGA) polymers. RONDEL™ (RNAi/Oligonucleotide Nanoparticle Delivery) polymers (Arrowhead Research Corporation, Pasadena, CA) and pH responsive co-block polymers such as, but not limited to, PHASERX® (Seattle, WA).

[000496] A non-limiting example of chitosan formulation includes a core of positively charged chitosan and an outer portion of negatively charged substrate (U.S. Pub. No. 20120258176; herein incorporated by reference in its entirety). Chitosan includes, but is not limited to N-trimethyl chitosan, mono-N-carboxymethyl chitosan (MCC), N-palmitoyl chitosan (NPCS), EDTA-chitosan, low molecular weight chitosan, chitosan derivatives, or combinations thereof.

[000497] In one embodiment, the polymers used in the present invention have undergone processing to reduce and/or inhibit the attachment of unwanted substances such as, but not limited to, bacteria, to the surface of the polymer. The polymer may be processed by methods known and/or described in the art and/or described in International Pub. No. WO2012150467, herein incorporated by reference in its entirety.

[000498] A non-limiting example of PLGA formulations include, but are not limited to, PLGA injectable depots (e.g., ELIGARD® which is formed by dissolving PLGA in 66% N-methyl-2-pyrrolidone (NMP) and the remainder being aqueous solvent and leuprolide. Once injected, the PLGA and leuprolide peptide precipitates into the subcutaneous space).

[000499] Many of these polymer approaches have demonstrated efficacy in delivering oligonucleotides *in vivo* into the cell cytoplasm (reviewed in deFougerolles *Hum Gene Ther.* 2008 19:125-132; herein incorporated by reference in its entirety). Two polymer approaches that have yielded robust *in vivo* delivery of nucleic acids, in this case with small interfering RNA (siRNA), are dynamic polyconjugates and cyclodextrin-based nanoparticles (see e.g., US Patent Publication No. US20130156721, herein incorporated by reference in its entirety). The first of these delivery approaches uses dynamic polyconjugates and has been shown *in vivo* in mice to effectively deliver siRNA and silence endogenous target mRNA in hepatocytes (Rozema et al., *Proc Natl Acad Sci U S A.* 2007 104:12982-12887; herein incorporated by reference in its entirety). This particular approach is a multicomponent polymer system whose key features include a membrane-active polymer to which nucleic acid, in this case siRNA, is covalently coupled via a disulfide bond and where both PEG (for charge masking) and *N*-acetylgalactosamine (for hepatocyte targeting) groups are linked via pH-sensitive bonds (Rozema et al., *Proc Natl Acad Sci U S A.* 2007 104:12982-12887; herein incorporated by reference in its entirety). On binding to the hepatocyte and entry into the endosome, the polymer complex disassembles in the low-pH environment, with the polymer exposing its positive charge, leading to endosomal escape and cytoplasmic release of the siRNA from the polymer. Through replacement of the *N*-acetylgalactosamine group with a mannose group, it was shown one could alter targeting from asialoglycoprotein receptor-expressing hepatocytes to sinusoidal endothelium and Kupffer cells. Another polymer approach involves using transferrin-targeted cyclodextrin-containing polycation nanoparticles. These nanoparticles have demonstrated targeted silencing of the *EWS-FLII* gene product in transferrin receptor-expressing Ewing's sarcoma tumor cells (Hu-Lieskovan *et al.*, *Cancer Res.* 2005 65: 8984-8982; herein incorporated by reference in its entirety) and siRNA formulated in these nanoparticles was well tolerated in non-human

primates (Heidel *et al.*, Proc Natl Acad Sci USA 2007 104:5715-21; herein incorporated by reference in its entirety). Both of these delivery strategies incorporate rational approaches using both targeted delivery and endosomal escape mechanisms.

[000500] The polymer formulation can permit the sustained or delayed release of polynucleotides (e.g., following intramuscular or subcutaneous injection). The altered release profile for the polynucleotide can result in, for example, translation of an encoded protein over an extended period of time. The polymer formulation may also be used to increase the stability of the polynucleotide. Biodegradable polymers have been previously used to protect nucleic acids other than polynucleotide from degradation and been shown to result in sustained release of payloads in vivo (Rozema *et al.*, Proc Natl Acad Sci U S A. 2007 104:12982-12887; Sullivan *et al.*, Expert Opin Drug Deliv. 2010 7:1433-1446; Convertine *et al.*, Biomacromolecules. 2010 Oct 1; Chu *et al.*, Acc Chem Res. 2012 Jan 13; Manganiello *et al.*, Biomaterials. 2012 33:2301-2309; Benoit *et al.*, Biomacromolecules. 2011 12:2708-2714; Singha *et al.*, Nucleic Acid Ther. 2011 2:133-147; deFougerolles Hum Gene Ther. 2008 19:125-132; Schaffert and Wagner, Gene Ther. 2008 16:1131-1138; Chaturvedi *et al.*, Expert Opin Drug Deliv. 2011 8:1455-1468; Davis, Mol Pharm. 2009 6:659-668; Davis, Nature 2010 464:1067-1070; each of which is herein incorporated by reference in its entirety).

[000501] In one embodiment, the pharmaceutical compositions may be sustained release formulations. In a further embodiment, the sustained release formulations may be for subcutaneous delivery. Sustained release formulations may include, but are not limited to, PLGA microspheres, ethylene vinyl acetate (EVAc), poloxamer, GELSITE® (Nanotherapeutics, Inc. Alachua, FL), HYLENEX® (Halozyne Therapeutics, San Diego CA), surgical sealants such as fibrinogen polymers (Ethicon Inc. Cornelia, GA), TISSELL® (Baxter International, Inc Deerfield, IL), PEG-based sealants, and COSEAL® (Baxter International, Inc Deerfield, IL).

[000502] As a non-limiting example modified mRNA may be formulated in PLGA microspheres by preparing the PLGA microspheres with tunable release rates (e.g., days and weeks) and encapsulating the modified mRNA in the PLGA microspheres while maintaining the integrity of the modified mRNA during the encapsulation process. EVAc are non-biodegradable, biocompatible polymers which are used extensively in

pre-clinical sustained release implant applications (e.g., extended release products Ocusert a pilocarpine ophthalmic insert for glaucoma or progestasert a sustained release progesterone intrauterine device; transdermal delivery systems Testoderm, Duragesic and Selegiline; catheters). Poloxamer F-407 NF is a hydrophilic, non-ionic surfactant triblock copolymer of polyoxyethylene-polyoxypropylene-polyoxyethylene having a low viscosity at temperatures less than 5°C and forms a solid gel at temperatures greater than 15°C. PEG-based surgical sealants comprise two synthetic PEG components mixed in a delivery device which can be prepared in one minute, seals in 3 minutes and is reabsorbed within 30 days. GELSITE® and natural polymers are capable of in-situ gelation at the site of administration. They have been shown to interact with protein and peptide therapeutic candidates through ionic interaction to provide a stabilizing effect.

[000503] Polymer formulations can also be selectively targeted through expression of different ligands as exemplified by, but not limited by, folate, transferrin, and N-acetylgalactosamine (GalNAc) (Benoit et al., *Biomacromolecules*. 2011 12:2708-2714; Rozema et al., *Proc Natl Acad Sci U S A*. 2007 104:12982-12887; Davis, *Mol Pharm*. 2009 6:659-668; Davis, *Nature* 2010 464:1067-1070; each of which is herein incorporated by reference in its entirety).

[000504] The polynucleotides of the invention may be formulated with or in a polymeric compound. The polymer may include at least one polymer such as, but not limited to, polyethenes, polyethylene glycol (PEG), poly(l-lysine)(PLL), PEG grafted to PLL, cationic lipopolymer, biodegradable cationic lipopolymer, polyethylenimine (PEI), cross-linked branched poly(alkylene imines), a polyamine derivative, a modified poloxamer, a biodegradable polymer, elastic biodegradable polymer, biodegradable block copolymer, biodegradable random copolymer, biodegradable polyester copolymer, biodegradable polyester block copolymer, biodegradable polyester block random copolymer, multiblock copolymers, linear biodegradable copolymer, poly[α -(4-aminobutyl)-L-glycolic acid) (PAGA), biodegradable cross-linked cationic multi-block copolymers, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polylysine,

poly(ethylene imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), acrylic polymers, amine-containing polymers, dextran polymers, dextran polymer derivatives or combinations thereof .

[000505] As a non-limiting example, the polynucleotides of the invention may be formulated with the polymeric compound of PEG grafted with PLL as described in U.S. Pat. No. 6,177,274; herein incorporated by reference in its entirety. The formulation may be used for transfecting cells *in vitro* or for *in vivo* delivery of polynucleotide. In another example, the polynucleotide may be suspended in a solution or medium with a cationic polymer, in a dry pharmaceutical composition or in a solution that is capable of being dried as described in U.S. Pub. Nos. 20090042829 and 20090042825; each of which are herein incorporated by reference in their entireties.

[000506] As another non-limiting example the polynucleotides of the invention may be formulated with a PLGA-PEG block copolymer (see US Pub. No. US20120004293 and US Pat No. 8,236,330, herein incorporated by reference in their entireties) or PLGA-PEG-PLGA block copolymers (See U.S. Pat. No. 6,004,573, herein incorporated by reference in its entirety). As a non-limiting example, the polynucleotides of the invention may be formulated with a diblock copolymer of PEG and PLA or PEG and PLGA (see US Pat No 8,246,968, herein incorporated by reference in its entirety).

[000507] A polyamine derivative may be used to deliver nucleic acids or to treat and/or prevent a disease or to be included in an implantable or injectable device (U.S. Pub. No. 20100260817 (now U.S. Patent No. 8,460,696) the contents of each of which is herein incorporated by reference in its entirety). As a non-limiting example, a pharmaceutical composition may include the polynucleotide and the polyamine derivative described in U.S. Pub. No. 20100260817 (now U.S. Patent No. 8,460,696; the contents of which are incorporated herein by reference in its entirety. As a non-limiting example the polynucleotides of the present invention may be delivered using a polyamine polymer such as, but not limited to, a polymer comprising a 1,3-dipolar addition polymer prepared by combining a carbohydrate diazide monomer with a diyne unit comprising oligoamines (U.S. Pat. No. 8,236,280; herein incorporated by reference in its entirety).

[000508] The polynucleotides of the invention may be formulated with at least one acrylic polymer. Acrylic polymers include but are not limited to, acrylic acid,

methacrylic acid, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, amino alkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), polycyanoacrylates and combinations thereof.

[000509] In one embodiment, the polynucleotides of the present invention may be formulated with at least one polymer and/or derivatives thereof described in International Publication Nos. WO2011115862, WO2012082574 and WO2012068187 and U.S. Pub. No. 20120283427, each of which are herein incorporated by reference in their entireties. In another embodiment, the polynucleotides of the present invention may be formulated with a polymer of formula Z as described in WO2011115862, herein incorporated by reference in its entirety. In yet another embodiment, the polynucleotides may be formulated with a polymer of formula Z, Z' or Z'' as described in International Pub. Nos. WO2012082574 or WO2012068187 and U.S. Pub. No. 2012028342, each of which are herein incorporated by reference in their entireties. The polymers formulated with the modified RNA of the present invention may be synthesized by the methods described in International Pub. Nos. WO2012082574 or WO2012068187, each of which are herein incorporated by reference in their entireties.

[000510] The polynucleotides of the invention may be formulated with at least one acrylic polymer. Acrylic polymers include but are not limited to, acrylic acid, methacrylic acid, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, amino alkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), polycyanoacrylates and combinations thereof.

[000511] Formulations of polynucleotides of the invention may include at least one amine-containing polymer such as, but not limited to polylysine, polyethylene imine, poly(amidoamine) dendrimers, poly(amine-co-esters) or combinations thereof. As a non-limiting example, the poly(amine-co-esters) may be the polymers described in and/or made by the methods described in International Publication No WO2013082529, the contents of which are herein incorporated by reference in its entirety.

[000512] For example, the polynucleotides of the invention may be formulated in a pharmaceutical compound including a poly(alkylene imine), a biodegradable cationic

lipopolymer, a biodegradable block copolymer, a biodegradable polymer, or a biodegradable random copolymer, a biodegradable polyester block copolymer, a biodegradable polyester polymer, a biodegradable polyester random copolymer, a linear biodegradable copolymer, PAGA, a biodegradable cross-linked cationic multi-block copolymer or combinations thereof. The biodegradable cationic lipopolymer may be made by methods known in the art and/or described in U.S. Pat. No. 6,696,038, U.S. App. Nos. 20030073619 and 20040142474 each of which is herein incorporated by reference in their entireties. The poly(alkylene imine) may be made using methods known in the art and/or as described in U.S. Pub. No. 20100004315, herein incorporated by reference in its entirety. The biodegradable polymer, biodegradable block copolymer, the biodegradable random copolymer, biodegradable polyester block copolymer, biodegradable polyester polymer, or biodegradable polyester random copolymer may be made using methods known in the art and/or as described in U.S. Pat. Nos. 6,517,869 and 6,267,987, the contents of which are each incorporated herein by reference in their entirety. The linear biodegradable copolymer may be made using methods known in the art and/or as described in U.S. Pat. No. 6,652,886. The PAGA polymer may be made using methods known in the art and/or as described in U.S. Pat. No. 6,217,912 herein incorporated by reference in its entirety. The PAGA polymer may be copolymerized to form a copolymer or block copolymer with polymers such as but not limited to, poly-L-lysine, polyarginine, polyornithine, histones, avidin, protamines, polylactides and poly(lactide-co-glycolides). The biodegradable cross-linked cationic multi-block copolymers may be made by methods known in the art and/or as described in U.S. Pat. No. 8,057,821, 8,444,992 or U.S. Pub. No. 2012009145 each of which are herein incorporated by reference in their entireties. For example, the multi-block copolymers may be synthesized using linear polyethyleneimine (LPEI) blocks which have distinct patterns as compared to branched polyethyleneimines. Further, the composition or pharmaceutical composition may be made by the methods known in the art, described herein, or as described in U.S. Pub. No. 20100004315 or U.S. Pat. Nos. 6,267,987 and 6,217,912 each of which are herein incorporated by reference in their entireties.

[000513] The polynucleotides of the invention may be formulated with at least one degradable polyester which may contain polycationic side chains. Degradable polyesters

include, but are not limited to, poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), and combinations thereof. In another embodiment, the degradable polyesters may include a PEG conjugation to form a PEGylated polymer.

[000514] The polynucleotides of the invention may be formulated with at least one crosslinkable polyester. Crosslinkable polyesters include those known in the art and described in US Pub. No. 20120269761, the contents of which is herein incorporated by reference in its entirety.

[000515] The polynucleotides of the invention may be formulated in or with at least one cyclodextrin polymer. Cyclodextrin polymers and methods of making cyclodextrin polymers include those known in the art and described in US Pub. No. 20130184453, the contents of which are herein incorporated by reference in its entirety.

[000516] In one embodiment, the polynucleotides of the invention may be formulated in or with at least one crosslinked cation-binding polymers. Crosslinked cation-binding polymers and methods of making crosslinked cation-binding polymers include those known in the art and described in International Patent Publication No. WO2013106072, WO2013106073 and WO2013106086, the contents of each of which are herein incorporated by reference in its entirety.

[000517] In one embodiment, the polynucleotides of the invention may be formulated in or with at least one branched polymer. Branched polymers and methods of making branched polymers include those known in the art and described in International Patent Publication No. WO2013113071, the contents of each of which are herein incorporated by reference in its entirety.

[000518] In one embodiment, the polynucleotides of the invention may be formulated in or with at least PEGylated albumin polymer. PEGylated albumin polymer and methods of making PEGylated albumin polymer include those known in the art and described in US Patent Publication No. US20130231287, the contents of each of which are herein incorporated by reference in its entirety.

[000519] In one embodiment, the polymers described herein may be conjugated to a lipid-terminating PEG. As a non-limiting example, PLGA may be conjugated to a lipid-terminating PEG forming PLGA-DSPE-PEG. As another non-limiting example, PEG conjugates for use with the present invention are described in International Publication

No. WO2008103276, herein incorporated by reference in its entirety. The polymers may be conjugated using a ligand conjugate such as, but not limited to, the conjugates described in U.S. Pat. No. 8,273,363, herein incorporated by reference in its entirety.

[000520] In one embodiment, the polynucleotides disclosed herein may be mixed with the PEGs or the sodium phosphate/sodium carbonate solution prior to administration. In another embodiment, a polynucleotides encoding a protein of interest may be mixed with the PEGs and also mixed with the sodium phosphate/sodium carbonate solution. In yet another embodiment, polynucleotides encoding a protein of interest may be mixed with the PEGs and a polynucleotides encoding a second protein of interest may be mixed with the sodium phosphate/sodium carbonate solution.

[000521] In one embodiment, the polynucleotides described herein may be conjugated with another compound. Non-limiting examples of conjugates are described in US Patent Nos. 7,964,578 and 7,833,992, each of which are herein incorporated by reference in their entireties. In another embodiment, modified RNA of the present invention may be conjugated with conjugates of formula 1-122 as described in US Patent Nos. 7,964,578 and 7,833,992, each of which are herein incorporated by reference in their entireties. The polynucleotides described herein may be conjugated with a metal such as, but not limited to, gold. (See e.g., Giljohann et al. Journ. Amer. Chem. Soc. 2009 131(6): 2072-2073; herein incorporated by reference in its entirety). In another embodiment, the polynucleotides described herein may be conjugated and/or encapsulated in gold-nanoparticles. (International Pub. No. WO201216269 and U.S. Pub. No. 20120302940 and US20130177523; the contents of each of which is herein incorporated by reference in its entirety).

[000522] As described in U.S. Pub. No. 20100004313, herein incorporated by reference in its entirety, a gene delivery composition may include a nucleotide sequence and a poloxamer. For example, the polynucleotides of the present invention may be used in a gene delivery composition with the poloxamer described in U.S. Pub. No. 20100004313.

[000523] In one embodiment, the polymer formulation of the present invention may be stabilized by contacting the polymer formulation, which may include a cationic carrier, with a cationic lipopolymer which may be covalently linked to cholesterol and polyethylene glycol groups. The polymer formulation may be contacted with a cationic

lipopolymer using the methods described in U.S. Pub. No. 20090042829 herein incorporated by reference in its entirety. The cationic carrier may include, but is not limited to, polyethylenimine, poly(trimethylenimine), poly(tetramethylenimine), polypropylenimine, aminoglycoside-polyamine, dideoxy-diamino-b-cyclodextrin, spermine, spermidine, poly(2-dimethylamino)ethyl methacrylate, poly(lysine), poly(histidine), poly(arginine), cationized gelatin, dendrimers, chitosan, 1,2-Dioleoyl-3-Trimethylammonium-Propane(DOTAP), N-[1-(2,3-dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride (DOTMA), 1-[2-(oleoyloxy)ethyl]-2-oleyl-3-(2-hydroxyethyl)imidazolinium chloride (DOTIM), 2,3-dioleoyloxy-N-[2(sperminecarboxamido)ethyl]-N,N-dimethyl-1-propanaminium trifluoroacetate (DOSPA), 3B-[N—(N',N'-Dimethylaminoethane)-carbamoyl]Cholesterol Hydrochloride (DC-Cholesterol HCl) diheptadecylamidoglycyl spermidine (DOGS), N,N-distearyl-N,N-dimethylammonium bromide (DDAB), N-(1,2-dimyristyloxyprop-3-yl)-N,N-dimethyl-N-hydroxyethyl ammonium bromide (DMRIE), N,N-dioleoyl-N,N-dimethylammonium chloride (DODAC) and combinations thereof. As a non-limiting example, the polynucleotides may be formulated with a cationic lipopolymer such as those described in U.S. Patent Application No. 20130065942, herein incorporated by reference in its entirety.

[000524] The polynucleotides of the invention may be formulated in a polyplex of one or more polymers (See e.g., U.S. Pat. No. 8,501,478, U.S. Pub. No. 20120237565 and 20120270927 and 20130149783 and International Patent Pub. No. WO2013090861; the contents of each of which is herein incorporated by reference in its entirety). As a non-limiting example, the polyplex may be formed using the novel alpha-aminoamidine polymers described in International Publication No. WO2013090861, the contents of which are herein incorporated by reference in its entirety. As another non-limiting example, the polyplex may be formed using the click polymers described in US Patent No. 8,501,478, the contents of which is herein incorporated by reference in its entirety.

[000525] In one embodiment, the polyplex comprises two or more cationic polymers. The cationic polymer may comprise a poly(ethylene imine) (PEI) such as linear PEI. In another embodiment, the polyplex comprises p(TETA/CBA) its PEGylated analog p(TETA/CBA)-g-PEG2k and mixtures thereof (see e.g., US Patent Publication No.

US20130149783, the contents of which are herein incorporated by reference in its entirety.

[000526] The polynucleotides of the invention can also be formulated as a nanoparticle using a combination of polymers, lipids, and/or other biodegradable agents, such as, but not limited to, calcium phosphate. Components may be combined in a core-shell, hybrid, and/or layer-by-layer architecture, to allow for fine-tuning of the nanoparticle so to delivery of the polynucleotide, polynucleotides may be enhanced (Wang et al., *Nat Mater.* 2006 5:791-796; Fuller et al., *Biomaterials.* 2008 29:1526-1532; DeKoker et al., *Adv Drug Deliv Rev.* 2011 63:748-761; Endres et al., *Biomaterials.* 2011 32:7721-7731; Su et al., *Mol Pharm.* 2011 Jun 6;8(3):774-87; herein incorporated by reference in its entirety). As a non-limiting example, the nanoparticle may comprise a plurality of polymers such as, but not limited to hydrophilic-hydrophobic polymers (e.g., PEG-PLGA), hydrophobic polymers (e.g., PEG) and/or hydrophilic polymers (International Pub. No. WO20120225129; the contents of which is herein incorporated by reference in its entirety).

[000527] As another non-limiting example the nanoparticle comprising hydrophilic polymers for the polynucleotides may be those described in or made by the methods described in International Patent Publication No. WO2013119936, the contents of which are herein incorporated by reference in its entirety.

[000528] In one embodiment, the biodegradable polymers which may be used in the present invention are poly(ether-anhydride) block copolymers. As a non-limiting example, the biodegradable polymers used herein may be a block copolymer as described in International Patent Publication No WO2006063249, herein incorporated by reference in its entirety, or made by the methods described in International Patent Publication No WO2006063249, herein incorporated by reference in its entirety.

[000529] In another embodiment, the biodegradable polymers which may be used in the present invention are alkyl and cycloalkyl terminated biodegradable lipids. As a non-limiting example, the alkyl and cycloalkyl terminated biodegradable lipids may be those described in International Publication No. WO2013086322 and/or made by the methods described in International Publication No. WO2013086322; the contents of which are herein incorporated by reference in its entirety.

[000530] In yet another embodiment, the biodegradable polymers which may be used in the present invention are cationic lipids having one or more biodegradable group located in a lipid moiety. As a non-limiting example, the biodegradable lipids may be those described in US Patent Publication No. US20130195920, the contents of which are herein incorporated by reference in its entirety.

[000531] Biodegradable calcium phosphate nanoparticles in combination with lipids and/or polymers have been shown to deliver polynucleotides *in vivo*. In one embodiment, a lipid coated calcium phosphate nanoparticle, which may also contain a targeting ligand such as anisamide, may be used to deliver the polynucleotide, polynucleotides of the present invention. For example, to effectively deliver siRNA in a mouse metastatic lung model a lipid coated calcium phosphate nanoparticle was used (Li et al., J Contr Rel. 2010 142: 416-421; Li et al., J Contr Rel. 2012 158:108-114; Yang et al., Mol Ther. 2012 20:609-615; herein incorporated by reference in its entirety). This delivery system combines both a targeted nanoparticle and a component to enhance the endosomal escape, calcium phosphate, in order to improve delivery of the siRNA.

[000532] In one embodiment, calcium phosphate with a PEG-polyanion block copolymer may be used to delivery polynucleotides (Kazikawa et al., J Contr Rel. 2004 97:345-356; Kazikawa et al., J Contr Rel. 2006 111:368-370; the contents of each of which are herein incorporated by reference in its entirety).

[000533] In one embodiment, a PEG-charge-conversional polymer (Pitella et al., Biomaterials. 2011 32:3106-3114; the contents of which are herein incorporated by reference in its entirety) may be used to form a nanoparticle to deliver the polynucleotides of the present invention. The PEG-charge-conversional polymer may improve upon the PEG-polyanion block copolymers by being cleaved into a polycation at acidic pH, thus enhancing endosomal escape.

[000534] In one embodiment, a polymer used in the present invention may be a pentablock polymer such as, but not limited to, the pentablock polymers described in International Patent Publication No. WO2013055331, herein incorporated by reference in its entirety. As a non-limiting example, the pentablock polymer comprises PGA-PCL-PEG-PCL-PGA, wherein PEG is polyethylene glycol, PCL is poly(E-caprolactone), PGA is poly(glycolic acid), and PLA is poly(lactic acid). As another non-limiting example, the

pentablock polymer comprises PEG-PCL- PLA-PCL-PEG, wherein PEG is polyethylene glycol, PCL is poly(E-caprolactone), PGA is poly(glycolic acid), and PLA is poly(lactic acid).

[000535] In one embodiment, a polymer which may be used in the present invention comprises at least one diepoxide and at least one aminoglycoside (See e.g., International Patent Publication No. WO2013055971, the contents of which are herein incorporated by reference in its entirety). The diepoxide may be selected from, but is not limited to, 1,4 butanediol diglycidyl ether (1,4 B), 1,4-cyclohexanedimethanol diglycidyl ether (1,4 C), 4-vinylcyclohexene diepoxide (4VCD), ethyleneglycol diglycidyl ether (EDGE), glycerol diglycidyl ether (GDE), neopentylglycol diglycidyl ether (NPDGE), poly(ethyleneglycol) diglycidyl ether (PEGDE), poly(propyleneglycol) diglycidyl ether (PPGDE) and resorcinol diglycidyl ether (RDE). The aminoglycoside may be selected from, but is not limited to, streptomycin, neomycin, framycetin, paromomycin, ribostamycin, kanamycin, amikacin, arbekacin, bekanamycin, dibekacin, tobramycin, spectinomycin, hygromycin, gentamicin, netilmicin, sisomicin, isepamicin, verdamicin, astromicin, and apramycin. As a non-limiting example, the polymers may be made by the methods described in International Patent Publication No. WO2013055971, the contents of which are herein incorporated by reference in its entirety. As another non-limiting example, compositions comprising any of the polymers comprising at least one diepoxide and at least one aminoglycoside may be made by the methods described in International Patent Publication No. WO2013055971, the contents of which are herein incorporated by reference in its entirety.

[000536] In one embodiment, a polymer which may be used in the present invention may be a cross-linked polymer. As a non-limiting example, the cross-linked polymers may be used to form a particle as described in US Patent No. 8,414,927, the contents of which are herein incorporated by reference in its entirety. As another non-limiting example, the cross-linked polymer may be obtained by the methods described in US Patent Publication No. US20130172600, the contents of which are herein incorporated by reference in its entirety.

[000537] In another embodiment, a polymer which may be used in the present invention may be a cross-linked polymer such as those described in US Patent No. 8,461,132, the

contents of which are herein incorporated by reference in its entirety. As a non-limiting example, the cross-linked polymer may be used in a therapeutic composition for the treatment of a body tissue. The therapeutic composition may be administered to damaged tissue using various methods known in the art and/or described herein such as injection or catheterization.

[000538] In one embodiment, a polymer which may be used in the present invention may be a di-aliphatic substituted pegylated lipid such as, but not limited to, those described in International Patent Publication No. WO2013049328, the contents of which are herein incorporated by reference in its entirety.

[000539] In one embodiment, a block copolymer is PEG-PLGA-PEG (see e.g., the thermosensitive hydrogel (PEG-PLGA-PEG) was used as a TGF-beta1 gene delivery vehicle in Lee et al. Thermosensitive Hydrogel as a Tgf- β 1 Gene Delivery Vehicle Enhances Diabetic Wound Healing. *Pharmaceutical Research*, 2003 20(12): 1995-2000; as a controlled gene delivery system in Li et al. Controlled Gene Delivery System Based on Thermosensitive Biodegradable Hydrogel. *Pharmaceutical Research* 2003 20(6):884-888; and Chang et al., Non-ionic amphiphilic biodegradable PEG-PLGA-PEG copolymer enhances gene delivery efficiency in rat skeletal muscle. *J Controlled Release*. 2007 118:245-253; each of which is herein incorporated by reference in its entirety) may be used in the present invention. The present invention may be formulated with PEG-PLGA-PEG for administration such as, but not limited to, intramuscular and subcutaneous administration.

[000540] In another embodiment, the PEG-PLGA-PEG block copolymer is used in the present invention to develop a biodegradable sustained release system. In one aspect, the polynucleotides of the present invention are mixed with the block copolymer prior to administration. In another aspect, the polynucleotides acids of the present invention are co-administered with the block copolymer.

[000541] In one embodiment, the polymer used in the present invention may be a multi-functional polymer derivative such as, but not limited to, a multi-functional N-maleimidyl polymer derivatives as described in US Patent No US8454946, the contents of which are herein incorporated by reference in its entirety.

[000542] The use of core-shell nanoparticles has additionally focused on a high-throughput approach to synthesize cationic cross-linked nanogel cores and various shells (Siegwart et al., Proc Natl Acad Sci U S A. 2011 108:12996-13001; the contents of which are herein incorporated by reference in its entirety). The complexation, delivery, and internalization of the polymeric nanoparticles can be precisely controlled by altering the chemical composition in both the core and shell components of the nanoparticle. For example, the core-shell nanoparticles may efficiently deliver siRNA to mouse hepatocytes after they covalently attach cholesterol to the nanoparticle.

[000543] In one embodiment, a hollow lipid core comprising a middle PLGA layer and an outer neutral lipid layer containing PEG may be used to delivery of the polynucleotide, polynucleotides of the present invention. As a non-limiting example, in mice bearing a luciferase-expressing tumor, it was determined that the lipid-polymer-lipid hybrid nanoparticle significantly suppressed luciferase expression, as compared to a conventional lipoplex (Shi et al, Angew Chem Int Ed. 2011 50:7027-7031; herein incorporated by reference in its entirety).

[000544] In one embodiment, the lipid nanoparticles may comprise a core of the polynucleotides disclosed herein and a polymer shell. The polymer shell may be any of the polymers described herein and are known in the art. In an additional embodiment, the polymer shell may be used to protect the polynucleotides in the core.

[000545] Core-shell nanoparticles for use with the polynucleotides of the present invention are described and may be formed by the methods described in U.S. Pat. No. 8,313,777 or International Patent Publication No. WO2013124867, the contents of each of which are herein incorporated by reference in their entirety.

[000546] In one embodiment, the core-shell nanoparticles may comprise a core of the polynucleotides disclosed herein and a polymer shell. The polymer shell may be any of the polymers described herein and are known in the art. In an additional embodiment, the polymer shell may be used to protect the polynucleotides in the core.

[000547] In one embodiment, the polymer used with the formulations described herein may be a modified polymer (such as, but not limited to, a modified polyacetal) as described in International Publication No. WO2011120053, the contents of which are herein incorporated by reference in its entirety.

[000548] In one embodiment, the formulation may be a polymeric carrier cargo complex comprising a polymeric carrier and at least one nucleic acid molecule. Non-limiting examples of polymeric carrier cargo complexes are described in International Patent Publications Nos. WO2013113326, WO2013113501, WO2013113325, WO2013113502 and WO2013113736 and European Patent Publication No. EP2623121, the contents of each of which are herein incorporated by reference in their entireties. In one aspect the polymeric carrier cargo complexes may comprise a negatively charged nucleic acid molecule such as, but not limited to, those described in International Patent Publication Nos. WO2013113325 and WO2013113502, the contents of each of which are herein incorporated by reference in its entirety.

[000549] In one embodiment, a pharmaceutical composition may comprise polynucleotides of the invention and a polymeric carrier cargo complex. The polynucleotides may encode a protein of interest such as, but not limited to, an antigen from a pathogen associated with infectious disease, an antigen associated with allergy or allergic disease, an antigen associated with autoimmune disease or an antigen associated with cancer or tumor disease (See e.g., the antigens described in International Patent Publications Nos. WO2013113326, WO2013113501, WO2013113325, WO2013113502 and WO2013113736 and European Patent Publication No. EP2623121, the contents of each of which are herein incorporated by reference in their entireties).

[000550] As a non-limiting example, the core-shell nanoparticle may be used to treat an eye disease or disorder (See e.g. US Publication No. 20120321719, the contents of which are herein incorporated by reference in its entirety).

[000551] In one embodiment, the polymer used with the formulations described herein may be a modified polymer (such as, but not limited to, a modified polyacetal) as described in International Publication No. WO2011120053, the contents of which are herein incorporated by reference in its entirety.

Peptides and Proteins

[000552] The polynucleotides of the invention can be formulated with peptides and/or proteins in order to increase transfection of cells by the polynucleotide. In one embodiment, peptides such as, but not limited to, cell penetrating peptides and proteins and peptides that enable intracellular delivery may be used to deliver pharmaceutical

formulations. A non-limiting example of a cell penetrating peptide which may be used with the pharmaceutical formulations of the present invention includes a cell-penetrating peptide sequence attached to polycations that facilitates delivery to the intracellular space, e.g., HIV-derived TAT peptide, penetratins, transportans, or hCT derived cell-penetrating peptides (see, e.g., Caron et al., *Mol. Ther.* 3(3):310-8 (2001); Langel, *Cell-Penetrating Peptides: Processes and Applications* (CRC Press, Boca Raton FL, 2002); El-Andaloussi et al., *Curr. Pharm. Des.* 11(28):3597-611 (2003); and Deshayes et al., *Cell. Mol. Life Sci.* 62(16):1839-49 (2005), all of which are incorporated herein by reference in their entirety). The compositions can also be formulated to include a cell penetrating agent, e.g., liposomes, which enhance delivery of the compositions to the intracellular space. Polynucleotides of the invention may be complexed to peptides and/or proteins such as, but not limited to, peptides and/or proteins from Aileron Therapeutics (Cambridge, MA) and Permeon Biologics (Cambridge, MA) in order to enable intracellular delivery (Cronican et al., *ACS Chem. Biol.* 2010 5:747-752; McNaughton et al., *Proc. Natl. Acad. Sci. USA* 2009 106:6111-6116; Sawyer, *Chem Biol Drug Des.* 2009 73:3-6; Verdine and Hilinski, *Methods Enzymol.* 2012;503:3-33; all of which are herein incorporated by reference in its entirety).

[000553] In one embodiment, the cell-penetrating polypeptide may comprise a first domain and a second domain. The first domain may comprise a supercharged polypeptide. The second domain may comprise a protein-binding partner. As used herein, “protein-binding partner” includes, but are not limited to, antibodies and functional fragments thereof, scaffold proteins, or peptides. The cell-penetrating polypeptide may further comprise an intracellular binding partner for the protein-binding partner. The cell-penetrating polypeptide may be capable of being secreted from a cell where the polynucleotide may be introduced.

[000554] Formulations of the including peptides or proteins may be used to increase cell transfection by the polynucleotide, alter the biodistribution of the polynucleotide (e.g., by targeting specific tissues or cell types), and/or increase the translation of encoded protein. (See e.g., International Pub. No. WO2012110636 and WO2013123298; the contents of which are herein incorporated by reference in its entirety).

[000555] In one embodiment, the cell penetrating peptide may be, but is not limited to, those described in US Patent Publication No US20130129726, US20130137644 and US20130164219, each of which is herein incorporated by reference in its entirety.

Cells

[000556] The polynucleotides of the invention can be transfected *ex vivo* into cells, which are subsequently transplanted into a subject. As non-limiting examples, the pharmaceutical compositions may include red blood cells to deliver modified RNA to liver and myeloid cells, virosomes to deliver modified RNA in virus-like particles (VLPs), and electroporated cells such as, but not limited to, from MAXCYTE® (Gaithersburg, MD) and from ERYTECH® (Lyon, France) to deliver modified RNA. Examples of use of red blood cells, viral particles and electroporated cells to deliver payloads other than polynucleotides have been documented (Godfrin et al., Expert Opin Biol Ther. 2012 12:127-133; Fang et al., Expert Opin Biol Ther. 2012 12:385-389; Hu et al., Proc Natl Acad Sci U S A. 2011 108:10980-10985; Lund et al., Pharm Res. 2010 27:400-420; Huckriede et al., J Liposome Res. 2007;17:39-47; Cusi, Hum Vaccin. 2006 2:1-7; de Jonge et al., Gene Ther. 2006 13:400-411; all of which are herein incorporated by reference in its entirety).

[000557] The polynucleotides may be delivered in synthetic VLPs synthesized by the methods described in International Pub No. WO2011085231 and WO2013116656 and US Pub No. 20110171248, the contents of each of which are herein incorporated by reference in their entireties.

[000558] Cell-based formulations of the polynucleotides of the invention may be used to ensure cell transfection (e.g., in the cellular carrier), alter the biodistribution of the polynucleotide (e.g., by targeting the cell carrier to specific tissues or cell types), and/or increase the translation of encoded protein.

Introduction Into Cells

[000559] A variety of methods are known in the art and suitable for introduction of nucleic acid into a cell, including viral and non-viral mediated techniques. Examples of typical non-viral mediated techniques include, but are not limited to, electroporation, calcium phosphate mediated transfer, nucleofection, sonoporation, heat shock, magnetofection, liposome mediated transfer, microinjection, microprojectile mediated

transfer (nanoparticles), cationic polymer mediated transfer (DEAE-dextran, polyethylenimine, polyethylene glycol (PEG) and the like) or cell fusion.

[000560] The technique of sonoporation, or cellular sonication, is the use of sound (e.g., ultrasonic frequencies) for modifying the permeability of the cell plasma membrane. Sonoporation methods are known to those in the art and are used to deliver nucleic acids *in vivo* (Yoon and Park, Expert Opin Drug Deliv. 2010 7:321-330; Postema and Gilja, Curr Pharm Biotechnol. 2007 8:355-361; Newman and Bettinger, Gene Ther. 2007 14:465-475; all herein incorporated by reference in their entirety). Sonoporation methods are known in the art and are also taught for example as it relates to bacteria in US Patent Publication 20100196983 and as it relates to other cell types in, for example, US Patent Publication 20100009424, each of which are incorporated herein by reference in their entirety.

[000561] Electroporation techniques are also well known in the art and are used to deliver nucleic acids *in vivo* and clinically (Andre et al., Curr Gene Ther. 2010 10:267-280; Chiarella et al., Curr Gene Ther. 2010 10:281-286; Hojman, Curr Gene Ther. 2010 10:128-138; all herein incorporated by reference in their entirety). Electroporation devices are sold by many companies worldwide including, but not limited to BTX® Instruments (Holliston, MA) (e.g., the AgilePulse In Vivo System) and Inovio (Blue Bell, PA) (e.g., Inovio SP-5P intramuscular delivery device or the CELLECTRA® 3000 intradermal delivery device). In one embodiment, polynucleotides may be delivered by electroporation as described in Example 9.

Micro-Organ

[000562] The polynucleotides may be contained in a micro-organ which can then express an encoded polypeptide of interest in a long-lasting therapeutic formulation. In one aspect, the micro-organ may comprise a vector comprising a nucleic acid sequence (e.g., the polynucleotides of the present invention) encoding a polypeptide of interest, operably linked to one or more regulatory sequences. Micro-organs are described in co-pending International Patent Publication No. WO2015038892, the contents of which is incorporated by reference in its entirety, such as, but not limited to, in paragraphs [000672] – [000677].

Hyaluronidase

[000563] The intramuscular or subcutaneous localized injection of polynucleotides of the invention can include hyaluronidase, which catalyzes the hydrolysis of hyaluronan. By catalyzing the hydrolysis of hyaluronan, a constituent of the interstitial barrier, hyaluronidase lowers the viscosity of hyaluronan, thereby increasing tissue permeability (Frost, Expert Opin. Drug Deliv. (2007) 4:427-440; herein incorporated by reference in its entirety). It is useful to speed their dispersion and systemic distribution of encoded proteins produced by transfected cells. Alternatively, the hyaluronidase can be used to increase the number of cells exposed to a polynucleotide of the invention administered intramuscularly or subcutaneously.

Nanoparticle Mimics

[000564] The polynucleotides of the invention may be encapsulated within and/or absorbed to a nanoparticle mimic. A nanoparticle mimic can mimic the delivery function organisms or particles such as, but not limited to, pathogens, viruses, bacteria, fungus, parasites, prions and cells. As a non-limiting example the polynucleotides of the invention may be encapsulated in a non-viron particle which can mimic the delivery function of a virus (see International Pub. No. WO2012006376 and US Patent Publication No. US20130171241 and US20130195968, the contents of each of which are herein incorporated by reference in its entirety).

Nanotubes

[000565] The polynucleotides of the invention can be attached or otherwise bound to at least one nanotube such as, but not limited to, rosette nanotubes, rosette nanotubes having twin bases with a linker, carbon nanotubes and/or single-walled carbon nanotubes, The polynucleotides may be bound to the nanotubes through forces such as, but not limited to, steric, ionic, covalent and/or other forces. Nanotubes are described in co-pending International Patent Publication No. WO2015038892, the contents of which is incorporated by reference in its entirety, such as, but not limited to, in paragraphs [000679] – [000684].

Conjugates

[000566] The polynucleotides of the invention include conjugates, such as a polynucleotide covalently linked to a carrier or targeting group, or including two

encoding regions that together produce a fusion protein (e.g., bearing a targeting group and therapeutic protein or peptide).

[000567] The conjugates of the invention include a naturally occurring substance, such as a protein (e.g., human serum albumin (HSA), low-density lipoprotein (LDL), high-density lipoprotein (HDL), or globulin); an carbohydrate (e.g., a dextran, pullulan, chitin, chitosan, inulin, cyclodextrin or hyaluronic acid); or a lipid. The ligand may also be a recombinant or synthetic molecule, such as a synthetic polymer, e.g., a synthetic polyamino acid, an oligonucleotide (e.g. an aptamer). Examples of polyamino acids include polyamino acid is a polylysine (PLL), poly L-aspartic acid, poly L-glutamic acid, styrene-maleic acid anhydride copolymer, poly(L-lactide-co-glycolid) copolymer, divinyl ether-maleic anhydride copolymer, N-(2-hydroxypropyl)methacrylamide copolymer (HMPA), polyethylene glycol (PEG), polyvinyl alcohol (PVA), polyurethane, poly(2-ethylacrylic acid), N-isopropylacrylamide polymers, or polyphosphazene. Example of polyamines include: polyethylenimine, polylysine (PLL), spermine, spermidine, polyamine, pseudopeptide-polyamine, peptidomimetic polyamine, dendrimer polyamine, arginine, amidine, protamine, cationic lipid, cationic porphyrin, quaternary salt of a polyamine, or an alpha helical peptide.

[000568] Representative U.S. patents that teach the preparation of polynucleotide conjugates, particularly to RNA, include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941; 6,294,664; 6,320,017; 6,576,752; 6,783,931; 6,900,297; 7,037,646; each of which is herein incorporated by reference in their entireties.

[000569] In one embodiment, the conjugate of the present invention may function as a carrier for the polynucleotides of the present invention. The conjugate may comprise a cationic polymer such as, but not limited to, polyamine, polylysine, polyalkylenimine,

and polyethylenimine which may be grafted to with poly(ethylene glycol). As a non-limiting example, the conjugate may be similar to the polymeric conjugate and the method of synthesizing the polymeric conjugate described in U.S. Pat. No. 6,586,524 herein incorporated by reference in its entirety.

[000570] A non-limiting example of a method for conjugation to a substrate is described in US Patent Publication No. US20130211249, the contents of which are herein incorporated by reference in its entirety. The method may be used to make a conjugated polymeric particle comprising a polynucleotide.

[000571] The conjugates can also include targeting groups, e.g., a cell or tissue targeting agent, e.g., a lectin, glycoprotein, lipid or protein, e.g., an antibody, that binds to a specified cell type such as a kidney cell. A targeting group can be a thyrotropin, melanotropin, lectin, glycoprotein, surfactant protein A, Mucin carbohydrate, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-gulucosamine multivalent mannose, multivalent fucose, glycosylated polyaminoacids, multivalent galactose, transferrin, bisphosphonate, polyglutamate, polyaspartate, a lipid, cholesterol, a steroid, bile acid, folate, vitamin B12, biotin, an RGD peptide, an RGD peptide mimetic or an aptamer.

[000572] Targeting groups can be proteins, e.g., glycoproteins, or peptides, e.g., molecules having a specific affinity for a co-ligand, or antibodies e.g., an antibody, that binds to a specified cell type such as a cancer cell, endothelial cell, or bone cell. Targeting groups may also include hormones and hormone receptors. They can also include non-peptidic species, such as lipids, lectins, carbohydrates, vitamins, cofactors, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-gulucosamine multivalent mannose, multivalent fucose, or aptamers. The ligand can be, for example, a lipopolysaccharide, or an activator of p38 MAP kinase.

[000573] The targeting group can be any ligand that is capable of targeting a specific receptor. Examples include, without limitation, folate, GalNAc, galactose, mannose, mannose-6P, aptamers, integrin receptor ligands, chemokine receptor ligands, transferrin, biotin, serotonin receptor ligands, PSMA, endothelin, GCPII, somatostatin, LDL, and HDL ligands. In particular embodiments, the targeting group is an aptamer. The aptamer can be unmodified or have any combination of modifications disclosed herein.

[000574] As a non-limiting example, the targeting group may be a glutathione receptor (GR)-binding conjugate for targeted delivery across the blood-central nervous system barrier (See e.g., US Patent Publication No. US2013021661012, the contents of which are herein incorporated by reference in its entirety).

[000575] In one embodiment, the conjugate of the present invention may be a synergistic biomolecule-polymer conjugate. The synergistic biomolecule-polymer conjugate may be long-acting continuous-release system to provide a greater therapeutic efficacy. The synergistic biomolecule-polymer conjugate may be those described in US Patent Publication No. US20130195799, the contents of which are herein incorporated by reference in its entirety.

[000576] In another embodiment, the conjugate which may be used in the present invention may be an aptamer conjugate. Non-limiting examples of aptamers conjugates are described in International Patent Publication No. WO2012040524, the contents of which are herein incorporated by reference in its entirety. The aptamer conjugates may be used to provide targeted delivery of formulations comprising polynucleotides.

[000577] In one embodiment, the conjugate which may be used in the present invention may be an amine containing polymer conjugate. Non-limiting examples of amine containing polymer conjugate are described in US Patent No. US 8,507,653, the contents of which are herein incorporated by reference in its entirety. The factor IX moiety polymer conjugate may be comprise releasable linkages to release the polynucleotides upon and/or after delivery to a subject.

[000578] In one embodiment, pharmaceutical compositions of the present invention may include chemical modifications such as, but not limited to, modifications similar to locked nucleic acids.

[000579] Representative U.S. Patents that teach the preparation of locked nucleic acid (LNA) such as those from Santaris, include, but are not limited to, the following: U.S. Pat. Nos. 6,268,490; 6,670,461; 6,794,499; 6,998,484; 7,053,207; 7,084,125; and 7,399,845, each of which is herein incorporated by reference in its entirety.

[000580] Representative U.S. patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each

of which is herein incorporated by reference. Further teaching of PNA compounds can be found, for example, in Nielsen *et al.*, Science, 1991, 254, 1497-1500.

[000581] Some embodiments featured in the invention include polynucleotides with phosphorothioate backbones and oligonucleotides with other modified backbones, and in particular --CH₂--NH--CH₂--, --CH₂--N(CH₃)--O--CH₂--[known as a methylene (methylinino) or MMI backbone], --CH₂--O--N(CH₃)--CH₂--, --CH₂--N(CH₃)--N(CH₃)--CH₂-- and --N(CH₃)--CH₂--CH₂--[wherein the native phosphodiester backbone is represented as --O—P(O)₂--O--CH₂--] of the above-referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above-referenced U.S. Pat. No. 5,602,240. In some embodiments, the polynucleotides featured herein have morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

[000582] Modifications at the 2' position may also aid in delivery. Preferably, modifications at the 2' position are not located in a polypeptide-coding sequence, i.e., not in a translatable region. Modifications at the 2' position may be located in a 5'UTR, a 3'UTR and/or a tailing region. Modifications at the 2' position can include one of the following at the 2' position: H (i.e., 2'-deoxy); F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Exemplary suitable modifications include O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. In other embodiments, the polynucleotides include one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties, or a group for improving the pharmacodynamic properties, and other substituents having similar properties. In some embodiments, the modification includes a 2'-methoxyethoxy (2'-O--CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin *et al.*, *Helv. Chim. Acta*, 1995, 78:486-504) i.e., an alkoxy-alkoxy group. Another exemplary modification is 2'-dimethylaminooxyethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples herein

below, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), *i.e.*, 2'-O--CH₂--O--CH₂--N(CH₂)₂, also described in examples herein below. Other modifications include 2'-methoxy (2'-OCH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked dsRNAs and the 5' position of 5' terminal nucleotide.

Polynucleotides of the invention may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative U.S. patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920; the contents of each of which is herein incorporated by reference in their entirety.

[000583] In still other embodiments, the polynucleotide is covalently conjugated to a cell penetrating polypeptide. The cell-penetrating peptide may also include a signal sequence. The conjugates of the invention can be designed to have increased stability; increased cell transfection; and/or altered the biodistribution (e.g., targeted to specific tissues or cell types).

[000584] In one embodiment, the polynucleotides may be conjugated to an agent to enhance delivery. As a non-limiting example, the agent may be a monomer or polymer such as a targeting monomer or a polymer having targeting blocks as described in International Publication No. WO2011062965, herein incorporated by reference in its entirety. In another non-limiting example, the agent may be a transport agent covalently coupled to the polynucleotides of the present invention (See e.g., U.S. Pat. Nos. 6,835,393 and 7,374,778, each of which is herein incorporated by reference in its entirety). In yet another non-limiting example, the agent may be a membrane barrier transport enhancing agent such as those described in U.S. Pat. Nos. 7,737,108 and 8,003,129, each of which is herein incorporated by reference in its entirety.

[000585] In another embodiment, polynucleotides may be conjugated to SMARTT POLYMER TECHNOLOGY® (PHASERX®, Inc. Seattle, WA).

[000586] In another aspect, the conjugate may be a peptide that selectively directs the nanoparticle to neurons in a tissue or organism. As a non-limiting example, the peptide used may be, but is not limited to, the peptides described in US Patent Publication No. US20130129627, herein incorporated by reference in its entirety.

[000587] In yet another aspect, the conjugate may be a peptide that can assist in crossing the blood-brain barrier.

Self-Assembled Nanoparticles

[000588] Self-assembled nanoparticles have a well-defined size which may be precisely controlled as the nucleic acid strands may be easily reprogrammable. The polynucleotides described herein may be formulated in self-assembled nanoparticles. Nucleic acid self-assembled nanoparticles are described in International Patent Publication No. WO2014152211, the contents of which are herein incorporated by reference in its entirety, such as in paragraphs [000740] - [000743]. Polymer-based self-assembled nanoparticles are described in International Patent Publication No. WO2014152211, the contents of which are herein incorporated by reference in its entirety, such as in paragraphs [000744] - [000749].

Self-Assembled Macromolecules

[000589] The polynucleotides may be formulated in amphiphilic macromolecules (AMs) for delivery. AMs comprise biocompatible amphiphilic polymers which have an alkylated sugar backbone covalently linked to poly(ethylene glycol). In aqueous solution, the AMs self-assemble to form micelles. Non-limiting examples of methods of forming AMs and AMs are described in US Patent Publication No. US20130217753, the contents of which are herein incorporated by reference in its entirety.

Inorganic Nanoparticles

[000590] The polynucleotides of the present invention may be formulated in inorganic nanoparticles (U.S. Pat. No. 8,257,745, herein incorporated by reference in its entirety). The inorganic nanoparticles may include, but are not limited to, clay substances that are water swellable. As a non-limiting example, the inorganic nanoparticle may include synthetic smectite clays which are made from simple silicates (See e.g., U.S. Pat. No. 5,585,108 and 8,257,745 each of which are herein incorporated by reference in their entirety).

[000591] In one embodiment, the inorganic nanoparticles may comprise a core of the polynucleotides disclosed herein and a polymer shell. The polymer shell may be any of the polymers described herein and are known in the art. In an additional embodiment, the polymer shell may be used to protect the polynucleotides in the core.

Semi-conductive and Metallic Nanoparticles

[000592] The polynucleotides of the present invention may be formulated in water-dispersible nanoparticle comprising a semiconductive or metallic material (U.S. Pub. No. 20120228565; herein incorporated by reference in its entirety) or formed in a magnetic nanoparticle (U.S. Pub. No. 20120265001 and 20120283503; each of which is herein incorporated by reference in its entirety). The water-dispersible nanoparticles may be hydrophobic nanoparticles or hydrophilic nanoparticles.

[000593] In one embodiment, the semi-conductive and/or metallic nanoparticles may comprise a core of the polynucleotides disclosed herein and a polymer shell. The polymer shell may be any of the polymers described herein and are known in the art. In an additional embodiment, the polymer shell may be used to protect the polynucleotides in the core.

Surgical Sealants: Gels and Hydrogels

[000594] In one embodiment, the polynucleotides disclosed herein may be encapsulated into any hydrogel known in the art which may form a gel when injected into a subject. Surgical sealants such as gels and hydrogels are described in International Patent Publication No. WO2014152211, the contents of which are herein incorporated by reference in its entirety, such as in paragraphs [000762] - [000809].

Suspension formulations

[000595] In some embodiments, suspension formulations are provided comprising polynucleotides, water immiscible oil depots, surfactants and/or co-surfactants and/or co-solvents. Combinations of oils and surfactants may enable suspension formulation with polynucleotides. Delivery of polynucleotides in a water immiscible depot may be used to improve bioavailability through sustained release of mRNA from the depot to the surrounding physiologic environment and prevent polynucleotides degradation by nucleases.

[000596] In some embodiments, suspension formulations of mRNA may be prepared using combinations of polynucleotides, oil-based solutions and surfactants. Such formulations may be prepared as a two-part system comprising an aqueous phase comprising polynucleotides and an oil-based phase comprising oil and surfactants. Exemplary oils for suspension formulations may include, but are not limited to sesame oil and Miglyol (comprising esters of saturated coconut and palmkernel oil-derived caprylic and capric fatty acids and glycerin or propylene glycol), corn oil, soybean oil, peanut oil, beeswax and/or palm seed oil. Exemplary surfactants may include, but are not limited to Cremophor, polysorbate 20, polysorbate 80, polyethylene glycol, transcutool, Capmul®, labrasol, isopropyl myristate, and/or Span 80. In some embodiments, suspensions may comprise co-solvents including, but not limited to ethanol, glycerol and/or propylene glycol.

[000597] Suspensions may be formed by first preparing polynucleotides formulation comprising an aqueous solution of polynucleotide and an oil-based phase comprising one or more surfactants. Suspension formation occurs as a result of mixing the two phases (aqueous and oil-based). In some embodiments, such a suspension may be delivered to an aqueous phase to form an oil-in-water emulsion. In some embodiments, delivery of a suspension to an aqueous phase results in the formation of an oil-in-water emulsion in which the oil-based phase comprising polynucleotides forms droplets that may range in size from nanometer-sized droplets to micrometer-sized droplets. In some embodiments, specific combinations of oils, surfactants, cosurfactants and/or co-solvents may be utilized to suspend polynucleotides in the oil phase and/or to form oil-in-water emulsions upon delivery into an aqueous environment.

[000598] In some embodiments, suspensions may provide modulation of the release of polynucleotides into the surrounding environment. In such embodiments, polynucleotides release may be modulated by diffusion from a water immiscible depot followed by resolubilization into a surrounding environment (e.g. an aqueous environment).

[000599] In some embodiments, polynucleotides within a water immiscible depot (e.g. suspended within an oil phase) may result in altered polynucleotides stability (e.g. altered degradation by nucleases).

[000600] In some embodiments, polynucleotides may be formulated such that upon injection, an emulsion forms spontaneously (e.g. when delivered to an aqueous phase). Such particle formation may provide a high surface area to volume ratio for release of polynucleotides from an oil phase to an aqueous phase.

[000601] In one embodiment, the polynucleotides may be formulated in a nanoemulsion such as, but not limited to, the nanoemulsions described in US Patent No. 8,496,945, the contents of which are herein incorporated by reference in its entirety. The nanoemulsions may comprise nanoparticles described herein. As a non-limiting example, the nanoparticles may comprise a liquid hydrophobic core which may be surrounded or coated with a lipid or surfactant layer. The lipid or surfactant layer may comprise at least one membrane-integrating peptide and may also comprise a targeting ligand (see e.g., US Patent No. 8,496,945, the contents of which are herein incorporated by reference in its entirety).

Cations and Anions

[000602] Formulations of polynucleotides disclosed herein may include cations or anions. In one embodiment, the formulations include metal cations such as, but not limited to, Zn^{2+} , Ca^{2+} , Cu^{2+} , Mg^{+} and combinations thereof. As a non-limiting example, formulations may include polymers and a polynucleotides complexed with a metal cation (See e.g., U.S. Pat. Nos. 6,265,389 and 6,555,525, each of which is herein incorporated by reference in its entirety).

[000603] In some embodiments, cationic nanoparticles comprising combinations of divalent and monovalent cations may be formulated with polynucleotides. Such nanoparticles may form spontaneously in solution over a given period (e.g. hours, days, etc.). Such nanoparticles do not form in the presence of divalent cations alone or in the presence of monovalent cations alone. The delivery of polynucleotides in cationic nanoparticles or in one or more depot comprising cationic nanoparticles may improve polynucleotide bioavailability by acting as a long-acting depot and/or reducing the rate of degradation by nucleases.

Molded Nanoparticles and Microparticles

[000604] The polynucleotides disclosed herein may be formulated in nanoparticles and/or microparticles. These nanoparticles and/or microparticles may be molded into any

size shape and chemistry. As an example, the nanoparticles and/or microparticles may be made using the PRINT® technology by LIQUIDA TECHNOLOGIES® (Morrisville, NC) (See e.g., International Pub. No. WO2007024323; the contents of which are herein incorporated by reference in its entirety).

[000605] In one embodiment, the molded nanoparticles may comprise a core of the polynucleotides disclosed herein and a polymer shell. The polymer shell may be any of the polymers described herein and are known in the art. In an additional embodiment, the polymer shell may be used to protect the polynucleotides in the core.

[000606] In one embodiment, the polynucleotides of the present invention may be formulated in microparticles. The microparticles may contain a core of the polynucleotides and a cortex of a biocompatible and/or biodegradable polymer. As a non-limiting example, the microparticles which may be used with the present invention may be those described in U.S. Patent No. 8,460,709, U.S. Patent Publication No. US20130129830 and International Patent Publication No WO2013075068, each of which is herein incorporated by reference in its entirety. As another non-limiting example, the microparticles may be designed to extend the release of the polynucleotides of the present invention over a desired period of time (see e.g., extended release of a therapeutic protein in U.S. Patent Publication No. US20130129830, herein incorporated by reference in its entirety).

[000607] The microparticle for use with the present invention may have a diameter of at least 1 micron to at least 100 microns (e.g., at least 1 micron, at least 5 micron, at least 10 micron, at least 15 micron, at least 20 micron, at least 25 micron, at least 30 micron, at least 35 micron, at least 40 micron, at least 45 micron, at least 50 micron, at least 55 micron, at least 60 micron, at least 65 micron, at least 70 micron, at least 75 micron, at least 80 micron, at least 85 micron, at least 90 micron, at least 95 micron, at least 97 micron, at least 99 micron, and at least 100 micron).

NanoJackets and NanoLiposomes

[000608] The polynucleotides disclosed herein may be formulated in NanoJackets and NanoLiposomes by Keystone Nano (State College, PA). NanoJackets are made of compounds that are naturally found in the body including calcium, phosphate and may also include a small amount of silicates. Nanojackets may range in size from 5 to 50 nm

and may be used to deliver hydrophilic and hydrophobic compounds such as, but not limited to, polynucleotides.

[000609] NanoLiposomes are made of lipids such as, but not limited to, lipids which naturally occur in the body. NanoLiposomes may range in size from 60-80 nm and may be used to deliver hydrophilic and hydrophobic compounds such as, but not limited to, polynucleotides. In one aspect, the polynucleotides disclosed herein are formulated in a NanoLiposome such as, but not limited to, Ceramide NanoLiposomes.

Pseudovirions

[000610] In one embodiment, the polynucleotides disclosed herein may be formulated in Pseudovirions (e.g., pseudo-virions). Pseudovirions are described in paragraphs [000593] – [000598] of International Patent Publication No. WO2015058069, the contents of which are herein incorporated by reference in its entirety.

Minicells

[000611] In one aspect, the polynucleotides may be formulated in bacterial minicells. As a non-limiting example, bacterial minicells may be those described in International Publication No. WO2013088250 or US Patent Publication No. US20130177499, the contents of each of which are herein incorporated by reference in its entirety. The bacterial minicells comprising therapeutic agents such as polynucleotides described herein may be used to deliver the therapeutic agents to brain tumors.

Semi-solid Compositions

[000612] In one embodiment, the polynucleotides may be formulated with a hydrophobic matrix to form a semi-solid composition. As a non-limiting example, the semi-solid composition or paste-like composition may be made by the methods described in International Patent Publication No WO201307604, herein incorporated by reference in its entirety. The semi-solid composition may be a sustained release formulation as described in International Patent Publication No WO201307604, herein incorporated by reference in its entirety.

[000613] In another embodiment, the semi-solid composition may further have a micro-porous membrane or a biodegradable polymer formed around the composition (see e.g., International Patent Publication No WO201307604, herein incorporated by reference in its entirety).

[000614] The semi-solid composition using the polynucleotides of the present invention may have the characteristics of the semi-solid mixture as described in International Patent Publication No WO201307604, herein incorporated by reference in its entirety (e.g., a modulus of elasticity of at least 10^{-4} N·mm⁻², and/or a viscosity of at least 100mPa·s).

Exosomes

[000615] In one embodiment, the polynucleotides may be formulated in exosomes. The exosomes may be loaded with at least one polynucleotide and delivered to cells, tissues and/or organisms. As a non-limiting example, the polynucleotides may be loaded in the exosomes described in International Publication No. WO2013084000, herein incorporated by reference in its entirety.

Silk-Based Delivery

[000616] In one embodiment, the polynucleotides may be formulated in a sustained release silk-based delivery system. The silk-based delivery system may be formed by contacting a silk fibroin solution with a therapeutic agent such as, but not limited to, the polynucleotides described herein and/or known in the art. As a non-limiting example, the sustained release silk-based delivery system which may be used in the present invention and methods of making such system are described in US Patent Publication No. US20130177611, the contents of which are herein incorporated by reference in its entirety.

Microparticles

[000617] In one embodiment, formulations comprising polynucleotides may comprise microparticles. The microparticles may comprise a polymer described herein and/or known in the art such as, but not limited to, poly(α -hydroxy acid), a polyhydroxy butyric acid, a polycaprolactone, a polyorthoester and a polyanhydride. The microparticle may have adsorbent surfaces to adsorb biologically active molecules such as polynucleotides. As a non-limiting example microparticles for use with the present invention and methods of making microparticles are described in US Patent Publication No. US2013195923 and US20130195898 and US Patent No. 8,309,139 and 8,206,749, the contents of each of which are herein incorporated by reference in its entirety.

[000618] In another embodiment, the formulation may be a microemulsion comprising microparticles and polynucleotides. As a non-limiting example, microemulsions

comprising microparticles are described in US Patent Publication No. US2013195923 and US20130195898 and US Patent No. 8,309,139 and 8,206,749, the contents of each of which are herein incorporated by reference in its entirety.

Amino Acid Lipids

[000619] In one embodiment, the polynucleotides may be formulated in amino acid lipids. Amino acid lipids are lipophilic compounds comprising an amino acid residue and one or more lipophilic tails. Non-limiting examples of amino acid lipids and methods of making amino acid lipids are described in US Patent No. 8,501,824, the contents of which are herein incorporated by reference in its entirety.

[000620] In one embodiment, the amino acid lipids have a hydrophilic portion and a lipophilic portion. The hydrophilic portion may be an amino acid residue and a lipophilic portion may comprise at least one lipophilic tail.

[000621] In one embodiment, the amino acid lipid formulations may be used to deliver the polynucleotides to a subject.

[000622] In another embodiment, the amino acid lipid formulations may deliver a polynucleotide in releasable form which comprises an amino acid lipid that binds and releases the polynucleotides. As a non-limiting example, the release of the polynucleotides may be provided by an acid-labile linker such as, but not limited to, those described in U.S. Patent Nos. 7,098,032, 6,897,196, 6,426,086, 7,138,382, 5,563,250, and 5,505,931, the contents of each of which are herein incorporated by reference in its entirety.

Microvesicles

[000623] In one embodiment, polynucleotides may be formulated in microvesicles. Non-limiting examples of microvesicles include those described in US Patent Publication No. US20130209544, the contents of which are herein incorporated by reference in its entirety.

[000624] In one embodiment, the microvesicle is an ARRDC1-mediated microvesicles (ARMMs). Non-limiting examples of ARMMs and methods of making ARMMs are described in International Patent Publication No. WO2013119602, the contents of which are herein incorporated by reference in its entirety.

Interpolyelectrolyte Complexes

[000625] In one embodiment, the polynucleotides may be formulated in an interpolyelectrolyte complex. Interpolyelectrolyte complexes are formed when charge-dynamic polymers are complexed with one or more anionic molecules. Non-limiting examples of charge-dynamic polymers and interpolyelectrolyte complexes and methods of making interpolyelectrolyte complexes are described in US Patent No. 8,524,368, the contents of which is herein incorporated by reference in its entirety.

Crystalline Polymeric Systems

[000626] In one embodiment, the polynucleotides may be formulated in crystalline polymeric systems. Crystalline polymeric systems are polymers with crystalline moieties and/or terminal units comprising crystalline moieties. Non-limiting examples of polymers with crystalline moieties and/or terminal units comprising crystalline moieties termed “CYC polymers,” crystalline polymer systems and methods of making such polymers and systems are described in US Patent No. US 8,524,259, the contents of which are herein incorporated by reference in its entirety.

Phosphatidylserine (PS)

[000627] In one embodiment, the tolerogenic compositions and/or polynucleotides of the invention may comprise phosphatidylserine (PS). PS is an anionic phospholipid which is restricted to the inner surface of the plasma membrane. While not wishing to be bound by theory, PS translocates to the cell surface during early apoptosis where it serves as a marker for rapid uptake by phagocytes. PS is an inner surface plasma membrane phospholipid that translocates to the cell surface during early apoptosis where it serves as a marker for rapid uptake by phagocytes and immature dendritic cells. Several studies have shown that apoptotic cells inhibit dendritic cell maturation, thereby preventing autoimmunity (Williams et al., Immunology, 2007, Vol. 124, Pages 89-101, Stuart et al, Journal of Immunology, 2002, Vol. 168, Pages 1627-1635; the contents of each of which are herein incorporated by reference in their entirety), and PS may play a critical role in this process by inhibiting maturation and self-antigen presentation of dendritic cells (Wallet et al, Clinical Medicine & Research, 2005, Vol. 3, Pages 166-175; the contents of which are herein incorporated by reference in its entirety). PS inhibits the expression of MHC, and the expression of molecules associated with maturation of dendritic cells and co-stimulatory molecules, the secretion of IL-12p70, and the ability to activate CD4 and

CD8 T cells by dendritic cells (Kim et al. Immunity, 2004 Vol. 21, Pages 643-653, Chen et al, Journal of Immunology, 2004, Vol. 173, Pages 2985–2994, Doffek et al, Molecular Immunology, 2011, Vol. 48, Pages 1771-1777, Wallet et al, Clinical Medicine & Research, 2005, Vol. 3, Pages 166-175; the contents of each of which are herein incorporated by reference in their entirety). Moreover, as described by Gaitonde et al., FVIII complexed with PS significantly reduced the development of antibody response against FVIII in Hemophilia A mice, concurrent with an increase in number of Treg, suggesting the involvement of Tregs in PS-mediated hypo-responsiveness. (Journal of Biological Chemistry, 2013, Vol 288, Pages 17051-17056; the contents of which are herein incorporated by reference in its entirety).

[000628] In one embodiment, the tolerogenic composition and/or polynucleotides of the invention may comprise a phosphatidylserine (PS) and may inhibit the activation of dendritic cells (DCs) and may reduce adaptive immunity.

[000629] In another embodiment, the tolerogenic compositions and/or polynucleotides of the invention can be formulated in a liposome, lipid nanoparticle and/or polymer which comprises a phosphatidylserine (PS) to modulate adaptive immune responses.

[000630] In one embodiment, any of the formulations described herein may comprise phosphatidyleserine (PS) and at least one polynucleotide described herein.

[000631] In one embodiment, any of the tolerogenic compositions described herein may be co-administered with phosphatidyleserine (PS).

[000632] In one embodiment, any of the tolerogenic compositions described herein may be co-formulated with phosphatidyleserine (PS).

Excipients

[000633] Pharmaceutical formulations may additionally comprise a pharmaceutically acceptable excipient, which, as used herein, includes, but are not limited to, any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, flavoring agents, stabilizers, antioxidants, osmolality adjusting agents, pH adjusting agents and the like, as suited to the particular dosage form desired. Various excipients for formulating pharmaceutical compositions and techniques for preparing the composition are known in the art (see *Remington: The*

Science and Practice of Pharmacy, 21st Edition, A. R. Gennaro (Lippincott, Williams & Wilkins, Baltimore, MD, 2006; incorporated herein by reference in its entirety). The use of a conventional excipient medium may be contemplated within the scope of the present disclosure, except insofar as any conventional excipient medium is incompatible with a substance or its derivatives, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated to be within the scope of this invention.

[000634] In some embodiments, a pharmaceutically acceptable excipient may be at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% pure. In some embodiments, an excipient is approved for use for humans and for veterinary use. In some embodiments, an excipient may be approved by United States Food and Drug Administration. In some embodiments, an excipient may be of pharmaceutical grade. In some embodiments, an excipient may meet the standards of the United States Pharmacopoeia (USP), the European Pharmacopoeia (EP), the British Pharmacopoeia, and/or the International Pharmacopoeia.

[000635] Pharmaceutically acceptable excipients used in the manufacture of pharmaceutical compositions include, but are not limited to, inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering agents, lubricating agents, and/or oils. Such excipients may optionally be included in pharmaceutical compositions. The composition may also include excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and/or perfuming agents.

[000636] Exemplary diluents include, but are not limited to, calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, *etc.*, and/or combinations thereof.

[000637] Exemplary granulating and/or dispersing agents include, but are not limited to, potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose and wood products, natural sponge,

cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinyl-pyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (VEEGUM[®]), sodium lauryl sulfate, quaternary ammonium compounds, *etc.*, and/or combinations thereof.

[000638] Exemplary surface active agents and/or emulsifiers include, but are not limited to, natural emulsifiers (e.g. acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (e.g. bentonite [aluminum silicate] and VEEGUM[®] [magnesium aluminum silicate]), long chain amino acid derivatives, high molecular weight alcohols (e.g. stearyl alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (e.g. carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (e.g. carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan fatty acid esters (e.g. polyoxyethylene sorbitan monolaurate [TWEEN[®]20], polyoxyethylene sorbitan [TWEEN[®]60], polyoxyethylene sorbitan monooleate [TWEEN[®]80], sorbitan monopalmitate [SPAN[®]40], sorbitan monostearate [SPAN[®]60], sorbitan tristearate [SPAN[®]65], glyceryl monooleate, sorbitan monooleate [SPAN[®]80]), polyoxyethylene esters (e.g. polyoxyethylene monostearate [MYRJ[®]45], polyoxyethylene hydrogenated castor oil, polyethoxylated castor oil, polyoxymethylene stearate, and SOLUTOL[®]), sucrose fatty acid esters, polyethylene glycol fatty acid esters (e.g. CREMOPHOR[®]), polyoxyethylene ethers, (e.g. polyoxyethylene lauryl ether [BRIJ[®]30]), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid, ethyl laurate, sodium lauryl sulfate, PLUORINC[®]F 68, POLOXAMER[®]188, cetrimonium bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, *etc.* and/or combinations thereof.

[000639] Exemplary binding agents include, but are not limited to, starch (*e.g.* cornstarch and starch paste); gelatin; sugars (*e.g.* sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol); amino acids (*e.g.*, glycine); natural and synthetic gums (*e.g.* acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum silicate (VEEGUM[®]), and larch arabogalactan); alginates; polyethylene oxide; polyethylene glycol; inorganic calcium salts; silicic acid; polymethacrylates; waxes; water; alcohol; *etc.*; and combinations thereof.

[000640] Exemplary preservatives may include, but are not limited to, antioxidants, chelating agents, antimicrobial preservatives, antifungal preservatives, alcohol preservatives, acidic preservatives, and/or other preservatives. Oxidation is a potential degradation pathway for mRNA, especially for liquid mRNA formulations. In order to prevent oxidation, antioxidants can be added to the formulation. Exemplary antioxidants include, but are not limited to, alpha tocopherol, ascorbic acid, acorbyl palmitate, benzyl alcohol, butylated hydroxyanisole, EDTA, m-cresol, methionine, butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, thioglycerol and/or sodium sulfite. Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA), citric acid monohydrate, disodium edetate, dipotassium edetate, edetic acid, fumaric acid, malic acid, phosphoric acid, sodium edetate, tartaric acid, and/or trisodium edetate. Exemplary antimicrobial preservatives include, but are not limited to, benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and/or thimerosal. Exemplary antifungal preservatives include, but are not limited to, butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and/or sorbic acid. Exemplary alcohol preservatives include, but are not limited to, ethanol, polyethylene glycol, phenol,

phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and/or phenylethyl alcohol. Exemplary acidic preservatives include, but are not limited to, vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and/or phytic acid. Other preservatives include, but are not limited to, tocopherol, tocopherol acetate, deteroxime mesylate, cetrimide, butylated hydroxyanisol (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium metabisulfite, GLYDANT PLUS[®], PHENONIP[®], methylparaben, GERMALL[®] 115, GERMABEN[®] II, NEOLONE[™], KATHON[™], and/or EUXYL[®].

[000641] In some embodiments, the pH of polynucleotide solutions are maintained between pH 5 and pH 8 to improve stability. Exemplary buffers to control pH may include, but are not limited to sodium phosphate, sodium citrate, sodium succinate, histidine (or histidine-HCl), sodium carbonate, and/or sodium malate. In another embodiment, the exemplary buffers listed above may be used with additional monovalent counterions (including, but not limited to potassium). Divalent cations may also be used as buffer counterions; however, these are not preferred due to complex formation and/or mRNA degradation.

[000642] Exemplary buffering agents may also include, but are not limited to, citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, D-gluconic acid, calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate, potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, *etc.*, and/or combinations thereof.

[000643] Exemplary lubricating agents include, but are not limited to, magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, glyceryl behenate, hydrogenated

vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, *etc.*, and combinations thereof.

[000644] Exemplary oils include, but are not limited to, almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macademia nut, mallow, mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughy, palm, palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and/or combinations thereof.

[000645] Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and/or perfuming agents can be present in the composition, according to the judgment of the formulator.

[000646] Exemplary additives include physiologically biocompatible buffers (e.g., trimethylamine hydrochloride), addition of chelants (such as, for example, DTPA or DTPA-bisamide) or calcium chelate complexes (as for example calcium DTPA, CaNaDTPA-bisamide), or, optionally, additions of calcium or sodium salts (for example, calcium chloride, calcium ascorbate, calcium gluconate or calcium lactate). In addition, antioxidants and suspending agents can be used.

Cryoprotectants for mRNA

[000647] In some embodiments, polynucleotide formulations may comprise cyroprotectants. As used herein, the term “cryoprotectant” refers to one or more agent that when combined with a given substance, helps to reduce or eliminate damage to that substance that occurs upon freezing. In some embodiments, cryoprotectants are combined with polynucleotides in order to stabilize them during freezing. Frozen storage of mRNA between -20°C and -80°C may be advantageous for long term (e.g. 36 months) stability

of polynucleotide. In some embodiments, cryoprotectants are included in polynucleotide formulations to stabilize polynucleotide through freeze/thaw cycles and under frozen storage conditions. Cryoprotectants of the present invention may include, but are not limited to sucrose, trehalose, lactose, glycerol, dextrose, raffinose and/or mannitol. Trehalose is listed by the Food and Drug Administration as being generally regarded as safe (GRAS) and is commonly used in commercial pharmaceutical formulations.

Bulking agents

[000648] In some embodiments, polynucleotide formulations may comprise bulking agents. As used herein, the term “bulking agent” refers to one or more agents included in formulations to impart a desired consistency to the formulation and/or stabilization of formulation components. In some embodiments, bulking agents are included in lyophilized polynucleotide formulations to yield a “pharmaceutically elegant” cake, stabilizing the lyophilized polynucleotides during long term (e.g. 36 month) storage. Bulking agents of the present invention may include, but are not limited to sucrose, trehalose, mannitol, glycine, lactose and/or raffinose. In some embodiments, combinations of cryoprotectants and bulking agents (for example, sucrose/glycine or trehalose/mannitol) may be included to both stabilize polynucleotides during freezing and provide a bulking agent for lyophilization.

[000649] Non-limiting examples of formulations and methods for formulating the polynucleotides of the present invention are also provided in International Publication No WO2013090648 filed December 14, 2012, the contents of which are incorporated herein by reference in their entirety.

Inactive Ingredients

[000650] In some embodiments, polynucleotide formulations may comprise at least one excipient which is an inactive ingredient. As used herein, the term “inactive ingredient” refers to one or more inactive agents included in formulations. In some embodiments, all, none or some of the inactive ingredients which may be used in the formulations of the present invention may be approved by the US Food and Drug Administration (FDA). A non-exhaustive list of inactive ingredients and the routes of administration the inactive ingredients may be formulated in are described in 4 of co-pending International Publication No. WO2014152211 (Attorney Docket No. M030).

Delivery

[000651] The present disclosure encompasses the delivery of polynucleotides for any of therapeutic, pharmaceutical, diagnostic or imaging by any appropriate route taking into consideration likely advances in the sciences of drug delivery. Delivery may be naked or formulated.

Naked Delivery

[000652] The polynucleotides of the present invention may be delivered to a cell naked. As used herein in, “naked” refers to delivering polynucleotides free from agents which promote transfection. For example, the polynucleotides delivered to the cell may contain no modifications. The naked polynucleotides may be delivered to the cell using routes of administration known in the art and described herein.

Formulated Delivery

[000653] The polynucleotides of the present invention may be formulated, using the methods described herein. The formulations may contain polynucleotides which may be modified and/or unmodified. The formulations may further include, but are not limited to, cell penetration agents, a pharmaceutically acceptable carrier, a delivery agent, a bioerodible or biocompatible polymer, a solvent, and a sustained-release delivery depot. The formulated polynucleotides may be delivered to the cell using routes of administration known in the art and described herein.

[000654] The compositions may also be formulated for direct delivery to an organ or tissue in any of several ways in the art including, but not limited to, direct soaking or bathing, via a catheter, by gels, powder, ointments, creams, gels, lotions, and/or drops, by using substrates such as fabric or biodegradable materials coated or impregnated with the compositions, and the like.

Administration

[000655] The polynucleotides of the present invention may be administered by any route which results in a therapeutically effective outcome. These include, but are not limited to enteral (into the intestine), gastroenteral, epidural (into the dura matter), oral (by way of the mouth), transdermal, peridural, intracerebral (into the cerebrum), intracerebroventricular (into the cerebral ventricles), epicutaneous (application onto the skin), intradermal, (into the skin itself), subcutaneous (under the skin), nasal

administration (through the nose), intravenous (into a vein), intravenous bolus, intravenous drip, intraarterial (into an artery), intramuscular (into a muscle), intracardiac (into the heart), intraosseous infusion (into the bone marrow), intrathecal (into the spinal canal), intraperitoneal, (infusion or injection into the peritoneum), intravesical infusion, intravitreal, (through the eye), intracavernous injection (into a pathologic cavity) intracavitary (into the base of the penis), intravaginal administration, intrauterine, extra-amniotic administration, transdermal (diffusion through the intact skin for systemic distribution), transmucosal (diffusion through a mucous membrane), transvaginal, insufflation (snorting), sublingual, sublabial, enema, eye drops (onto the conjunctiva), in ear drops, auricular (in or by way of the ear), buccal (directed toward the cheek), conjunctival, cutaneous, dental (to a tooth or teeth), electro-osmosis, endocervical, endosinusal, endotracheal, extracorporeal, hemodialysis, infiltration, interstitial, intra-abdominal, intra-amniotic, intra-articular, intrabiliary, intrabronchial, intrabursal, intracartilaginous (within a cartilage), intracaudal (within the cauda equine), intracisternal (within the cisterna magna cerebellomedularis), intracorneal (within the cornea), dental intracornal, intracoronary (within the coronary arteries), intracorporus cavernosum (within the dilatable spaces of the corporus cavernosa of the penis), intradiscal (within a disc), intraductal (within a duct of a gland), intraduodenal (within the duodenum), intradural (within or beneath the dura), intraepidermal (to the epidermis), intraesophageal (to the esophagus), intragastric (within the stomach), intragingival (within the gingivae), intraileal (within the distal portion of the small intestine), intralesional (within or introduced directly to a localized lesion), intraluminal (within a lumen of a tube), intralymphatic (within the lymph), intramedullary (within the marrow cavity of a bone), intrameningeal (within the meninges), intraocular (within the eye), intraovarian (within the ovary), intrapericardial (within the pericardium), intrapleural (within the pleura), intraprostatic (within the prostate gland), intrapulmonary (within the lungs or its bronchi), intrasinal (within the nasal or periorbital sinuses), intraspinal (within the vertebral column), intrasynovial (within the synovial cavity of a joint), intratendinous (within a tendon), intratesticular (within the testicle), intrathecal (within the cerebrospinal fluid at any level of the cerebrospinal axis), intrathoracic (within the thorax), intratubular (within the tubules of an organ), intratumor (within a tumor), intratympanic (within the aurus

media), intravascular (within a vessel or vessels), intraventricular (within a ventricle), iontophoresis (by means of electric current where ions of soluble salts migrate into the tissues of the body), irrigation (to bathe or flush open wounds or body cavities), laryngeal (directly upon the larynx), nasogastric (through the nose and into the stomach), occlusive dressing technique (topical route administration which is then covered by a dressing which occludes the area), ophthalmic (to the external eye), oropharyngeal (directly to the mouth and pharynx), parenteral, percutaneous, periarticular, peridural, perineural, periodontal, rectal, respiratory (within the respiratory tract by inhaling orally or nasally for local or systemic effect), retrobulbar (behind the pons or behind the eyeball), intramyocardial (entering the myocardium), soft tissue, subarachnoid, subconjunctival, submucosal, topical, transplacental (through or across the placenta), transtracheal (through the wall of the trachea), transtympanic (across or through the tympanic cavity), ureteral (to the ureter), urethral (to the urethra), vaginal, caudal block, diagnostic, nerve block, biliary perfusion, cardiac perfusion, photopheresis or spinal. In specific embodiments, compositions may be administered in a way which allows them cross the blood-brain barrier, vascular barrier, or other epithelial barrier. In one embodiment, a formulation for a route of administration may include at least one inactive ingredient. Non-limiting examples of routes of administration and inactive ingredients which may be included in formulations for the specific route of administration is shown in Table 9 of International Patent Publication No. WO2015058069, the contents of which are herein incorporated by reference in its entirety. In Table 9 of WO2015058069, “AN” means anesthetic, “CNBLK” means cervical nerve block, “NBLK” means nerve block, “IV” means intravenous, “IM” means intramuscular and “SC” means subcutaneous.

[000656] Non-limiting routes of administration for the polynucleotides of the present invention are described below.

Parenteral and Injectable Administration

[000657] Liquid dosage forms for parenteral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and/or elixirs. In addition to active ingredients, liquid dosage forms may comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol,

ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and/or perfuming agents. In certain embodiments for parenteral administration, compositions are mixed with solubilizing agents such as CREMOPHOR[®], alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and/or combinations thereof.

[000658] A pharmaceutical composition for parenteral administration may comprise at least one inactive ingredient. Any or none of the inactive ingredients used may have been approved by the US Food and Drug Administration (FDA). A non-exhaustive list of inactive ingredients for use in pharmaceutical compositions for parenteral administration includes hydrochloric acid, mannitol, nitrogen, sodium acetate, sodium chloride and sodium hydroxide.

[000659] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing agents, wetting agents, and/or suspending agents. Sterile injectable preparations may be sterile injectable solutions, suspensions, and/or emulsions in nontoxic parenterally acceptable diluents and/or solvents, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid can be used in the preparation of injectables. The sterile formulation may also comprise adjuvants such as local anesthetics, preservatives and buffering agents.

[000660] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, and/or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[000661] Injectable formulations may be for direct injection into a region of a tissue, organ and/or subject. As a non-limiting example, a tissue, organ and/or subject may be directly injected a formulation by intramyocardial injection into the ischemic region. (See e.g., Zangi et al. Nature Biotechnology 2013; the contents of which are herein incorporated by reference in its entirety).

[000662] In order to prolong the effect of an active ingredient, it is often desirable to slow the absorption of the active ingredient from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microcapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

Rectal and Vaginal Administration

[000663] In one embodiment, the polynucleotides described here may be formulated for rectal and vaginal administration by the methods or compositions described in International Patent Publication No. WO2014152211, the contents of which are incorporated by reference in its entirety, such as in paragraphs [000910] - [000913].

Oral Administration

[000664] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and/or elixirs. In addition to active ingredients, liquid dosage forms may comprise inert diluents and/or excipients commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn,

germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and/or perfuming agents. In certain embodiments for parenteral administration, compositions are mixed with solubilizing agents such as CREMOPHOR[®], alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and/or combinations thereof.

[000665] Syrups and elixirs can be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol, glucose or sucrose. Such formulations can also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions can be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension can be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation can also be a sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

[000666] Suspensions for oral dosage may contain the active materials in a mixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients may be suspending agents, as a non-limiting example the suspending agents may be sodium carboxymethylcellulose, methylcellulose, hydropropyl-methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents can be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate; or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol

monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

[000667] Oily suspensions for oral dosage can be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions can contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents and flavoring agents can be added to provide palatable oral preparations. These compositions can be preserved by the addition of an anti-oxidant such as ascorbic acid

[000668] The oral dosage may also be in the form of oil-in-water emulsions. The oily phase can be a vegetable oil or a mineral oil or mixtures of these. Suitable emulsifying agents can be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

[000669] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, an active ingredient is mixed with at least one inert, pharmaceutically acceptable excipient such as sodium citrate or dicalcium phosphate and/or fillers or extenders (*e.g.* starches, lactose, sucrose, glucose, mannitol, and silicic acid), binders (*e.g.* carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia), humectants (*e.g.* glycerol), disintegrating agents (*e.g.* agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate), solution retarding agents (*e.g.* paraffin), absorption accelerators (*e.g.* quaternary ammonium compounds), wetting agents (*e.g.* cetyl alcohol and glycerol monostearate), absorbents (*e.g.* kaolin and bentonite clay), and lubricants (*e.g.* talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate), and mixtures thereof. In the case of capsules, tablets and pills, the dosage form

may comprise buffering agents. The solid dosage forms may also dissolve once they come in contact with liquid such as, but not limited to, saliva and bile.

[000670] Compositions intended for oral use can be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions can contain one or more such sweetening agents, flavoring agents, coloring agents or preservative agents in order to provide pharmaceutically elegant and palatable preparations.

[000671] Solid dosage forms may be uncoated or they can be coated by known techniques. In some cases such coatings can be prepared by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed.

[000672] Formulations for oral use can also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

[000673] Dosage forms for oral delivery may also be chewable or may be suckable (e.g., lozenge form). The chewable dosage forms may be sustained release formulations such as, but not limited to, the sustained release compositions described in International Publication No WO2013082470 and US Publication No US20130142876, each of which is herein incorporated by reference in its entirety. The chewable dosage forms may comprise amphipathic lipids such as, but not limited to, those described in International Publication No WO2013082470 and US Publication No US20130142876, each of which is herein incorporated by reference in its entirety.

Topical or Transdermal Administration

[000674] In one embodiment, the polynucleotides described here may be formulated for topical or transdermal administration by the methods or compositions described in International Patent Publication No. WO2014152211, the contents of which are incorporated by reference in its entirety, such as in paragraphs [000925] - [000941].

Depot Administration

[000675] In one embodiment, the polynucleotides described here may be formulated for depot administration by the methods or compositions described in International Patent Publication No. WO2014152211, the contents of which are incorporated by reference in its entirety, such as in paragraphs [000942] - [000948].

Pulmonary Administration

[000676] In one embodiment, the polynucleotides described here may be formulated for pulmonary administration by the methods or compositions described in International Patent Publication No. WO2014152211, the contents of which are incorporated by reference in its entirety, such as in paragraphs [000949] - [000954].

Intranasal, nasal and buccal Administration

[000677] In one embodiment, the polynucleotides described here may be formulated for intranasal, nasal or buccal administration by the methods or compositions described in International Patent Publication No. WO2014152211, the contents of which are incorporated by reference in its entirety, such as in paragraphs [000955] - [000958].

Ophthalmic and Auricular (Otic) Administration

[000678] In one embodiment, the polynucleotides described here may be formulated for ophthalmic or auricular (otic) administration by the methods or compositions described in International Patent Publication No. WO2014152211, the contents of which are incorporated by reference in its entirety, such as in paragraphs [000959] - [000965].

Payload Administration: Detectable Agents and Therapeutic Agents

[000679] The polynucleotides described herein can be used in a number of different scenarios in which delivery of a substance (the “payload”) to a biological target is desired, for example delivery of detectable substances for detection of the target, or delivery of a therapeutic agent. Detection methods can include, but are not limited to, both imaging *in vitro* and *in vivo* imaging methods, *e.g.*, immunohistochemistry, bioluminescence imaging (BLI), Magnetic Resonance Imaging (MRI), positron emission tomography (PET), electron microscopy, X-ray computed tomography, Raman imaging, optical coherence tomography, absorption imaging, thermal imaging, fluorescence reflectance imaging, fluorescence microscopy, fluorescence molecular tomographic imaging, nuclear magnetic resonance imaging, X-ray imaging, ultrasound imaging,

photoacoustic imaging, lab assays, or in any situation where tagging/staining/imaging is required.

[000680] The polynucleotides can be designed to include both a linker and a payload in any useful orientation. For example, a linker having two ends is used to attach one end to the payload and the other end to the nucleobase, such as at the C-7 or C-8 positions of the deaza-adenosine or deaza-guanosine or to the N-3 or C-5 positions of cytosine or uracil. The polynucleotide of the invention can include more than one payload (e.g., a label and a transcription inhibitor), as well as a cleavable linker. In one embodiment, the modified nucleotide is a modified 7-deaza-adenosine triphosphate, where one end of a cleavable linker is attached to the C7 position of 7-deaza-adenine, the other end of the linker is attached to an inhibitor (e.g., to the C5 position of the nucleobase on a cytidine), and a label (e.g., Cy5) is attached to the center of the linker (see, e.g., compound 1 of A*pCp C5 Parg Capless in Fig. 5 and columns 9 and 10 of U.S. Pat. No. 7,994,304, incorporated herein by reference). Upon incorporation of the modified 7-deaza-adenosine triphosphate to an encoding region, the resulting polynucleotide having a cleavable linker attached to a label and an inhibitor (e.g., a polymerase inhibitor). Upon cleavage of the linker (e.g., with reductive conditions to reduce a linker having a cleavable disulfide moiety), the label and inhibitor are released. Additional linkers and payloads (e.g., therapeutic agents, detectable labels, and cell penetrating payloads) are described herein and in International Publication No. WO2013151666 (Attorney Docket Number M300), the contents of which are incorporated herein by reference in their entirety.

[000681] For example, the polynucleotides described herein can be used in reprogramming induced pluripotent stem cells (iPS cells), which can directly track cells that are transfected compared to total cells in the cluster. In another example, a drug that may be attached to the polynucleotides via a linker and may be fluorescently labeled can be used to track the drug *in vivo*, e.g. intracellularly. Other examples include, but are not limited to, the use of a polynucleotides in reversible drug delivery into cells.

[000682] The polynucleotides described herein can be used in intracellular targeting of a payload, e.g., detectable or therapeutic agent, to specific organelle. Exemplary intracellular targets can include, but are not limited to, the nuclear localization for

advanced mRNA processing, or a nuclear localization sequence (NLS) linked to the mRNA containing an inhibitor.

[000683] In addition, the polynucleotides described herein can be used to deliver therapeutic agents to cells or tissues, *e.g.*, in living animals. For example, the polynucleotides described herein can be used to deliver highly polar chemotherapeutics agents to kill cancer cells. The polynucleotides attached to the therapeutic agent through a linker can facilitate member permeation allowing the therapeutic agent to travel into a cell to reach an intracellular target.

[000684] In one example, the linker is attached at the 2'-position of the ribose ring and/or at the 3' and/or 5' position of the polynucleotides (See *e.g.*, International Pub. No. WO2012030683, herein incorporated by reference in its entirety). The linker may be any linker disclosed herein, known in the art and/or disclosed in International Pub. No. WO2012030683, herein incorporated by reference in its entirety.

[000685] In another example, the polynucleotides can be attached to the polynucleotides a viral inhibitory peptide (VIP) through a cleavable linker. The cleavable linker can release the VIP and dye into the cell. In another example, the polynucleotides can be attached through the linker to an ADP-ribosylate, which is responsible for the actions of some bacterial toxins, such as cholera toxin, diphtheria toxin, and pertussis toxin. These toxin proteins are ADP-ribosyltransferases that modify target proteins in human cells. For example, cholera toxin ADP-ribosylates G proteins modifies human cells by causing massive fluid secretion from the lining of the small intestine, which results in life-threatening diarrhea.

[000686] In some embodiments, the payload may be a therapeutic agent such as a cytotoxin, radioactive ion, chemotherapeutic, or other therapeutic agent. A cytotoxin or cytotoxic agent includes any agent that may be detrimental to cells. Examples include, but are not limited to, taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxyanthracenedione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, puromycin, maytansinoids, *e.g.*, maytansinol (see U.S. Pat. No. 5,208,020 incorporated herein in its entirety), rachelmycin (CC-1065, see U.S. Pat. Nos. 5,475,092, 5,585,499,

and 5,846,545, all of which are incorporated herein by reference), and analogs or homologs thereof. Radioactive ions include, but are not limited to iodine (e.g., iodine 125 or iodine 131), strontium 89, phosphorous, palladium, cesium, iridium, phosphate, cobalt, yttrium 90, samarium 153, and praseodymium. Other therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thiotepa chlorambucil, rachelmycin (CC-1065), melphalan, carmustine (BSNU), lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine, vinblastine, taxol and maytansinoids).

[000687] In some embodiments, the payload may be a detectable agent, such as various organic small molecules, inorganic compounds, nanoparticles, enzymes or enzyme substrates, fluorescent materials, luminescent materials (e.g., luminol), bioluminescent materials (e.g., luciferase, luciferin, and aequorin), chemiluminescent materials, radioactive materials (e.g., ^{18}F , ^{67}Ga , $^{81\text{m}}\text{Kr}$, ^{82}Rb , ^{111}In , ^{123}I , ^{133}Xe , ^{201}Tl , ^{125}I , ^{35}S , ^{14}C , ^3H , or $^{99\text{m}}\text{Tc}$ (e.g., as pertechnetate (technetate(VII), TcO_4^-)), and contrast agents (e.g., gold (e.g., gold nanoparticles), gadolinium (e.g., chelated Gd), iron oxides (e.g., superparamagnetic iron oxide (SPIO), monocrystalline iron oxide nanoparticles (MIONs), and ultrasmall superparamagnetic iron oxide (USPIO)), manganese chelates (e.g., Mn-DPDP), barium sulfate, iodinated contrast media (iohexol), microbubbles, or perfluorocarbons). Such optically-detectable labels include for example, without limitation, 4-acetamido-4'-isothiocyanatostilbene-2,2'-disulfonic acid; acridine and derivatives (e.g., acridine and acridine isothiocyanate); 5-(2'-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS); 4-amino-N-[3-vinylsulfonyl]phenyl]naphthalimide-3,5 disulfonate; N-(4-anilino-1-naphthyl)maleimide; anthranilamide; BODIPY; Brilliant Yellow; coumarin and derivatives (e.g., coumarin, 7-amino-4-methylcoumarin (AMC, Coumarin 120), and 7-amino-4-trifluoromethylcoumarin (Coumarin 151)); cyanine dyes; cyanosine; 4',6-diaminidino-2-phenylindole (DAPI); 5' 5"-dibromopyrogallol-sulfonaphthalein (Bromopyrogallol Red);

7-diethylamino-3-(4'-isothiocyanatophenyl)-4-methylcoumarin; diethylenetriamine pentaacetate; 4,4'-diisothiocyanatodihydro-stilbene-2,2'-disulfonic acid; 4,4'-diisothiocyanatostilbene-2,2'-disulfonic acid; 5-[dimethylamino]-naphthalene-1-sulfonyl chloride (DNS, dansylchloride); 4-dimethylaminophenylazophenyl-4'-isothiocyanate (DABITC); eosin and derivatives (e.g., eosin and eosin isothiocyanate); erythrosin and derivatives (e.g., erythrosin B and erythrosin isothiocyanate); ethidium; fluorescein and derivatives (e.g., 5-carboxyfluorescein (FAM), 5-(4,6-dichlorotriazin-2-yl)aminofluorescein (DTAF), 2',7'-dimethoxy-4'5'-dichloro-6-carboxyfluorescein, fluorescein, fluorescein isothiocyanate, X-rhodamine-5-(and-6)-isothiocyanate (QFITC or XRITC), and fluorescamine); 2-[2-[3-[[1,3-dihydro-1,1-dimethyl-3-(3-sulfopropyl)-2H-benz[e]indol-2-ylidene]ethylidene]-2-[4-(ethoxycarbonyl)-1-piperazinyl]-1-cyclopenten-1-yl]ethenyl]-1,1-dimethyl-3-(3-sulfopropyl)-1H-benz[e]indolium hydroxide, inner salt, compound with n,n-diethylethanamine(1:1) (IR144); 5-chloro-2-[2-[3-[(5-chloro-3-ethyl-2(3H)-benzothiazol-ylidene)ethylidene]-2-(diphenylamino)-1-cyclopenten-1-yl]ethenyl]-3-ethyl benzothiazolium perchlorate (IR140); Malachite Green isothiocyanate; 4-methylumbelliferone orthocresolphthalein; nitrotyrosine; pararosaniline; Phenol Red; B-phycoerythrin; o-phthaldialdehyde; pyrene and derivatives(e.g., pyrene, pyrene butyrate, and succinimidyl 1-pyrene); butyrate quantum dots; Reactive Red 4 (CIBACRONTM Brilliant Red 3B-A); rhodamine and derivatives (e.g., 6-carboxy-X-rhodamine (ROX), 6-carboxyrhodamine (R6G), lissamine rhodamine B sulfonyl chloride rhodamine (Rhod), rhodamine B, rhodamine 123, rhodamine X isothiocyanate, sulforhodamine B, sulforhodamine 101, sulfonyl chloride derivative of sulforhodamine 101 (Texas Red), N,N,N',N'tetramethyl-6-carboxyrhodamine (TAMRA) tetramethyl rhodamine, and tetramethyl rhodamine isothiocyanate (TRITC)); riboflavin; rosolic acid; terbium chelate derivatives; Cyanine-3 (Cy3); Cyanine-5 (Cy5); cyanine-5.5 (Cy5.5), Cyanine-7 (Cy7); IRD 700; IRD 800; Alexa 647; La Jolla Blue; phthalocyanine; and naphthalocyanine.

[000688] In some embodiments, the detectable agent may be a non-detectable precursor that becomes detectable upon activation (e.g., fluorogenic tetrazine-fluorophore constructs (e.g., tetrazine-BODIPY FL, tetrazine-Oregon Green 488, or tetrazine-BODIPY TMR-X) or enzyme activatable fluorogenic agents (e.g., PROSENSE® (VisEn

Medical))). In vitro assays in which the enzyme labeled compositions can be used include, but are not limited to, enzyme linked immunosorbent assays (ELISAs), immunoprecipitation assays, immunofluorescence, enzyme immunoassays (EIA), radioimmunoassays (RIA), and Western blot analysis.

Combinations

[000689] The polynucleotides may be used in combination with one or more other therapeutic, prophylactic, diagnostic, or imaging agents. By “in combination with,” it is not intended to imply that the agents must be administered at the same time and/or formulated for delivery together, although these methods of delivery are within the scope of the present disclosure. Compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. In some embodiments, the present disclosure encompasses the delivery of pharmaceutical, prophylactic, diagnostic, or imaging compositions in combination with agents that may improve their bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body. As a non-limiting example, the polynucleotides may be used in combination with a pharmaceutical agent for the treatment of cancer or to control hyperproliferative cells. In U.S. Pat. No. 7,964,571, herein incorporated by reference in its entirety, a combination therapy for the treatment of solid primary or metastasized tumor is described using a pharmaceutical composition including a DNA plasmid encoding for interleukin-12 with a lipopolymer and also administering at least one anticancer agent or chemotherapeutic. Further, the polynucleotides of the present invention that encodes anti-proliferative molecules may be in a pharmaceutical composition with a lipopolymer (see e.g., U.S. Pub. No. 20110218231, herein incorporated by reference in its entirety, claiming a pharmaceutical composition comprising a DNA plasmid encoding an anti-proliferative molecule and a lipopolymer) which may be administered with at least one chemotherapeutic or anticancer agent (See e.g., the “Combination” Section in US Patent No. 8,518,907 and International Patent Publication No. WO201218754; the contents of each of which are herein incorporated by reference in its entirety).

[000690] The polynucleotides and pharmaceutical formulations thereof may be administered to a subject alone or used in combination with or include one or more other therapeutic agents, for example, anticancer agents. Thus, combinations of polynucleotides with other anti-cancer or chemotherapeutic agents are within the scope of the invention. Examples of such agents can be found in *Cancer Principles and Practice of Oncology* by V. T. Devita and S. Hellman (editors), 6th edition (Feb. 15, 2001), Lippincott Williams & Wilkins Publishers. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the cancer involved. Such anti-cancer agents include, but are not limited to, the following: estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic/cytostatic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors and other angiogenesis inhibitors, inhibitors of cell proliferation and survival signaling, apoptosis inducing agents and agents that interfere with cell cycle checkpoints. The polynucleotides may also be useful in combination with any therapeutic agent used in the treatment of HCC, for example, but not limitation sorafenib. Polynucleotides may be particularly useful when co-administered with radiation therapy.

[000691] In certain embodiments, the polynucleotides may be useful in combination with known anti-cancer agents including the following: estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors, and other angiogenesis inhibitors.

[000692] Examples of estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic agents, a hypoxia activatable, proteasome inhibitors, microtubule inhibitors/microtubule-stabilising agents, topoisomerase inhibitors, inhibitors of mitotic kinesins, histone deacetylase inhibitors, inhibitors of kinases involved in mitotic progression, antiproliferative agents, monoclonal antibody targeted therapeutic agents, HMG-CoA reductase inhibitors, prenyl-protein transferase inhibitors, angiogenesis inhibitors, therapeutic agents that modulate or inhibit angiogenesis, agents that interfere with cell cycle checkpoints, agents that interfere with

receptor tyrosine kinases (RTKs), inhibitors of cell proliferation and survival signaling pathway, apoptosis inducing agents, NSAIDs that are selective COX-2 inhibitors, inhibitors of COX-2, compounds that have been described as specific inhibitors of COX-2, angiogenesis inhibitors, tyrosine kinase inhibitors, compounds other than anti-cancer compounds, inhibitor of inherent multidrug resistance (MDR), anti-emetic agents to treat nausea or emesis, and neurokinin-1 receptor antagonists, are described in co-pending International Patent Publication No. WO2015038892, the contents of which is incorporated by reference in its entirety, such as, but not limited to, in paragraphs [000925] – [000953] and [000954] – [000957].

[000693] Another embodiment of the instant invention is the use of the polynucleotides in combination with gene therapy for the treatment of cancer. For an overview of genetic strategies to treating cancer see Hall et al. (*Am J Hum Genet* 61:785-789 (1997)) and Kufe et al. (*Cancer Medicine*, 5th Ed, pp 876-889, BC Decker, Hamilton, 2000). Gene therapy can be used to deliver any tumor suppressing gene. Examples of such genes include, but are not limited to, p53, which can be delivered via recombinant virus-mediated gene transfer (see U.S. Pat. No. 6,069,134, for example), a uPA/uPAR antagonist (“Adenovirus-Mediated Delivery of a uPA/uPAR Antagonist Suppresses Angiogenesis-Dependent Tumor Growth and Dissemination in Mice,” *Gene Therapy*, August 5(8):1105-13 (1998)), and interferon gamma (*J Immunol* 164:217-222 (2000)).

[000694] Polynucleotides may also be useful for treating or preventing cancer, including bone cancer, in combination with bisphosphonates (understood to include bisphosphonates, diphosphonates, bisphosphonic acids and diphosphonic acids). Examples of bisphosphonates include but are not limited to: etidronate (Didronel), pamidronate (Aredia), alendronate (Fosamax), risedronate (Actonel), zoledronate (Zometa), ibandronate (Boniva), incadronate or cimadronate, clodronate, EB-1053, minodronate, neridronate, piridronate and tiludronate including any and all pharmaceutically acceptable salts, derivatives, hydrates and mixtures thereof.

[000695] Polynucleotides may also be useful for treating cancer in combination with the following therapeutic agents: abarelix (Plenaxis Depot®); aldesleukin (Prokine®); Aldesleukin (Proleukin®); Alemtuzumab (Campath®); alitretinoin (Panretin®); allopurinol (Zyloprim®); altretamine (Hexylen®); amifostine (Ethyol®); anastrozole

(Arimidex®); arsenic trioxide (Trisenox®); asparaginase (Elspar®); azacitidine (Vidaza®); bendamustine hydrochloride (Treanda®); bevacuzimab (Avastin®); bexarotene capsules (Targretin®); bexarotene gel (Targretin®); bleomycin (Blenoxane®); bortezomib (Velcade®); brefeldin A; busulfan intravenous (Busulfex®); busulfan oral (Myleran®); calusterone (Methosarb®); capecitabine (Xeloda®); carboplatin (Paraplatin®); carmustine (BCNU®, BiCNU®); carmustine (Gliadel®); carmustine with Polifeprosan 20 Implant (Gliadel Wafer®); celecoxib (Celebrex); cetuximab (Erbix®); chlorambucil (Leukeran®); cisplatin (Platinol®); cladribine (Leustatin 2-CdA®); clofarabine (Clolar®); cyclophosphamide (Cytosan®, Neosar®); cyclophosphamide (Cytosan Injection®); cyclophosphamide (Cytosan Tablet®); cytarabine (Cytosar-U®); cytarabine liposomal (DepoCyt); dacarbazine (DTIC-Dome®); dactinomycin, actinomycin D (Cosmegen®); dalteparin sodium injection (Fragmin®); Darbepoetin alfa (Aranesp®); dasatinib (Sprycel®); daunorubicin liposomal (DanuoXome®); daunorubicin, daunomycin (Daunorubicin®); daunorubicin, daunomycin (Cerubidine®); degarelix (Firmagon®); Denileukin diftitox (Ontak®); dexrazoxane (Zinecard®); dexrazoxane hydrochloride (Totect®); didemnin B; 17-DMAG; docetaxel (Taxotere®); doxorubicin (Adriamycin PFS®); doxorubicin (Adriamycin®, Rubex®); doxorubicin (Adriamycin PFS Injection®); doxorubicin liposomal (Doxil®); dromostanolone propionate (Dromostanolone®); dromostanolone propionate (Masterone Injection®); eculizumab injection (Soliris®); Elliott's B Solution (Elliott's B Solution®); eltrombopag (Promacta®); epirubicin (Ellence®); Epoetin alfa (Epogen®); erlotinib (Tarceva®); estramustine (Emcyt®); ethinyl estradiol; etoposide phosphate (Etopophos®); etoposide, VP-16 (Vepesid®); everolimus tablets (Afinitor®); exemestane (Aromasin®); ferumoxitol (Feraheme Injection®); Filgrastim (Neupogen®); floxuridine (intraarterial) (FUDR®); fludarabine (Fludara®); fluorouracil, 5-FU (Adrucil®); fulvestrant (Faslodex®); gefitinib (Iressa®); geldanamycin; gemcitabine (Gemzar®); gemtuzumab ozogamicin (Mylotarg®); goserelin acetate (Zoladex Implant®); goserelin acetate (Zoladex®); histrelin acetate (Histrelin Implant®); hydroxyurea (Hydrea®); Ibritumomab Tiuxetan (Zevalin®); idarubicin (Idamycin®); ifosfamide (IFEX®); imatinib mesylate (Gleevec®); interferon alfa 2a (Roferon A®); Interferon alfa-2b (Intron A®); iobenguane 1123 injection (AdreView®); irinotecan

(Camptosar®); ixabepilone (Ixempra®); lapatinib tablets (Tykerb®); lenalidomide (Revlimid®); letrozole (Femara®); leucovorin (Wellcovorin®, Leucovorin®); Leuprolide Acetate (Eligard®); levamisole (Ergamisol®); lomustine, CCNU (CeeBU); meclorethamine, nitrogen mustard (Mustargen®); megestrol acetate (Megace®); melphalan, L-PAM (Alkeran®); mercaptopurine, 6-MP (Purinethol®); mesna (Mesnex®); mesna (Mesnex Tabs®); methotrexate (Methotrexate®); methoxsalen (Uvadex®); 8-methoxypsoralen; mitomycin C (Mutamycin®); mitotane (Lysodren®); mitoxantrone (Novantrone®); mitramycin; nandrolone phenpropionate (Durabolin-50); nelarabine (Arranon®); nilotinib (Tasigna®); Nofetumomab (Verluma®); ofatumumab (Arzerra®); Oprelvekin (Neumega®); oxaliplatin (Eloxatin®); paclitaxel (Paxene®); paclitaxel (Taxol®); paclitaxel protein-bound particles (Abraxane®); palifermin (Kepivance®); pamidronate (Aredia®); panitumumab (Vectibix®); pazopanib tablets (Votrient™); pegademase (Adagen (Pegademase Bovine)®); pegaspargase (Oncaspar®); Pegfilgrastim (Neulasta®); pemetrexed disodium (Alimta®); pentostatin (Nipent®); pipobroman (Vercyte®); plerixafor (Mozobil®); plicamycin, mithramycin (Mithracin®); porfimer sodium (Photofrin®); pralatrexate injection (Folotyn®); procarbazine (Matulane®); quinacrine (Atabrine®); rapamycin; Rasburicase (Elitek®); raloxifene hydrochloride (Evista®); Rituximab (Rituxan®); romidepsin (Istodax®); romiplostim (Nplate®); sargramostim (Leukine®); Sargramostim (Prokine); sorafenib (Nexavar); streptozocin (Zanosar®); sunitinib maleate (Sutent); talc (Sclerosol); tamoxifen (Nolvadex); temozolomide (Temodar); temsirolimus (Torisel); teniposide, VM-26 (Vumon®); testolactone (Teslac®); thioguanine, 6-TG (Thioguanine®); thiopurine; thiotepa (Thioplex®); topotecan (Hycamtin®); toremifene (Fareston); Tositumomab (Bexxar); Tositumomab/I-131 tositumomab (Bexxar®); trans-retinoic acid; Trastuzumab (Herceptin®); tretinoin, ATRA (Vesanoid®); triethylenemelamine; Uracil Mustard (Uracil Mustard Capsules®); valrubicin (Valstar®); vinblastine (Velban®); vincristine (Oncovin®); vinorelbine (Navelbine®); vorinostat (Zolinza®); wortmannin; and zoledronate (Zometa®).

[000696] The combinations referred to above can conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical compositions

comprising a combination as defined above together with a pharmaceutically acceptable diluent or carrier represent a further aspect of the invention.

[000697] The individual compounds of such combinations can be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations. In one embodiment, the individual compounds will be administered simultaneously in a combined pharmaceutical formulation.

[000698] It will further be appreciated that therapeutically, prophylactically, diagnostically, or imaging active agents utilized in combination may be administered together in a single composition or administered separately in different compositions. In general, it is expected that agents utilized in combination will be utilized at levels that do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination will be lower than those utilized individually. In one embodiment, the combinations, each or together may be administered according to the split dosing regimens described herein.

Dosing

[000699] The present invention provides methods comprising administering modified mRNAs and their encoded proteins or complexes in accordance with the invention to a subject in need thereof. Nucleic acids, proteins or complexes, or pharmaceutical, imaging, diagnostic, or prophylactic compositions thereof, may be administered to a subject using any amount and any route of administration effective for preventing, treating, diagnosing, or imaging a disease, disorder, and/or condition (*e.g.*, a disease, disorder, and/or condition relating to working memory deficits). The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the disease, the particular composition, its mode of administration, its mode of activity, and the like. Compositions in accordance with the invention are typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions of the present invention may be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective, prophylactically effective, or appropriate imaging dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity

of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

[000700] In certain embodiments, compositions in accordance with the present invention may be administered at dosage levels sufficient to deliver from about 0.0001 mg/kg to about 100 mg/kg, from about 0.001 mg/kg to about 0.05 mg/kg, from about 0.005 mg/kg to about 0.05 mg/kg, from about 0.001 mg/kg to about 0.005 mg/kg, from about 0.05 mg/kg to about 0.5 mg/kg, from about 0.01 mg/kg to about 50 mg/kg, from about 0.1 mg/kg to about 40 mg/kg, from about 0.5 mg/kg to about 30 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.1 mg/kg to about 10 mg/kg, or from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic, diagnostic, prophylactic, or imaging effect (see e.g., the range of unit doses described in International Publication No WO2013078199, herein incorporated by reference in its entirety). The desired dosage may be delivered three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three weeks, or every four weeks. In certain embodiments, the desired dosage may be delivered using multiple administrations (*e.g.*, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations). When multiple administrations are employed, split dosing regimens such as those described herein may be used.

[000701] In one embodiment, administration of polynucleotides in split-dose regimens may be used to produce higher levels of proteins in mammalian subjects. As used herein, a “split dose” is the division of single unit dose or total daily dose into two or more doses, *e.g.*, two or more administrations of the single unit dose. As used herein, a “single unit dose” is a dose of any therapeutic administered in one dose/at one time/single route/single point of contact, *i.e.*, single administration event. As used herein, a “total daily dose” is an amount given or prescribed in 24 hr period. It may be administered as a single unit dose. In one embodiment, the polynucleotides of the present invention are

administered to a subject in split doses. The polynucleotides may be formulated in buffer only or in a formulation described herein.

[000702] In one embodiment, the polynucleotides described herein may be administered using a micro-dosing regimen in order to induce tolerance to the polynucleotide, its excipients or the protein encoded by the polynucleotide. As a non-limiting example, the micro-dosing regimen may be administered in at least two doses (e.g., at least 2 doses, at least 3 doses, at least 4 doses, at least 5 doses, at least 6 doses, at least 7 doses, at least 8 doses, at least 9 doses, at least 10 doses, at least 11 doses, at least 12 doses, at least 13 doses, at least 14 doses, at least 15 doses, at least 16 doses, at least 17 doses, at least 18 doses, at least 19 doses, at least 20 doses, at least 21 doses, at least 22 doses, at least 23 doses, at least 24 doses, at least 25 doses, at least 26 doses, at least 27 doses, at least 28 doses, at least 29 doses, at least 30 doses, at least 31 doses, at least 32 doses, at least 33 doses, at least 34 doses, at least 35 doses, at least 36 doses, at least 37 doses, at least 38 doses, at least 39 doses, at least 40 doses, at least 41 doses, at least 42 doses, at least 43 doses, at least 44 doses, at least 45 doses, at least 46 doses, at least 47 doses, at least 48 doses, at least 49 doses, at least 50 doses, at least 51 doses, at least 52 doses, at least 53 doses, at least 54 doses, at least 55 doses, at least 56 doses, at least 57 doses, at least 58 doses, at least 59 doses, at least 60 doses, at least 61 doses, at least 62 doses, at least 63 doses, at least 64 doses, at least 65 doses, at least 66 doses, at least 67 doses, at least 68 doses, at least 69 doses, at least 70 doses, at least 71 doses, at least 72 doses, at least 73 doses, at least 74 doses, at least 75 doses, at least 76 doses, at least 77 doses, at least 78 doses, at least 79 doses, at least 80 doses, at least 81 doses, at least 82 doses, at least 83 doses, at least 84 doses, at least 85 doses, at least 86 doses, at least 87 doses, at least 88 doses, at least 89 doses, at least 90 doses, at least 91 doses, at least 92 doses, at least 93 doses, at least 94 doses, at least 95 doses, at least 96 doses, at least 97 doses, at least 98 doses, at least 99 doses, at least 100 doses or more than 100 doses), each of which are at levels as low as one millionth the desired therapeutic dose. As another non-limiting example, the micro-dosing regimen may be staged so the amount of polynucleotide administered may be increased over a pre-determined period of time (e.g., at least 1 hour, at least 2 hours, at least 3 hours, at least 4 hours, at least 5 hours, at least 6 hours, at least 7 hours, at least 8 hours, at least 9 hours, at least 10 hours, at least 11 hours, at least 12

hours, at least 13 hours, at least 14 hours, at least 15 hours, at least 16 hours, at least 17 hours, at least 18 hours, at least 19 hours, at least 20 hours, at least 21 hours, at least 22 hours, at least 23 hours, at least 1 day, at least 36 hours, at least 2 days, at least 3 days, at least 4 days, at least 5 days, at least 6 days, at least 1 week, at least 2 weeks, at least 3 weeks, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months or more than 6 months) and may rise progressively over a period of time until desired effect is achieved (e.g., an efficacy dose) or safety considerations (e.g., toxicity) are rectified.

Dosage Forms

[000703] A pharmaceutical composition described herein can be formulated into a dosage form described herein, such as a topical, intranasal, intratracheal, or injectable (e.g., intravenous, intraocular, intravitreal, intramuscular, intracardiac, intraperitoneal, and subcutaneous).

Liquid dosage forms

[000704] Liquid dosage forms for parenteral administration are described in co-pending International Patent Publication No. WO2015038892, the contents of which is incorporated by reference in its entirety, such as, but not limited to, in paragraph [0001037].

Injectable

[000705] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art and may include suitable dispersing agents, wetting agents, and/or suspending agents. Sterile injectable preparations may be sterile injectable solutions, suspensions, and/or emulsions in nontoxic parenterally acceptable diluents and/or solvents, for example, a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed include, but are not limited to, water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid can be used in the preparation of injectables.

Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, and/or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

In order to prolong the effect of an active ingredient, it may be desirable to slow the absorption of the active ingredient from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the polynucleotides then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered polynucleotides may be accomplished by dissolving or suspending the polynucleotides in an oil vehicle. Injectable depot forms are made by forming microcapsule matrices of the polynucleotides in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of polynucleotides to polymer and the nature of the particular polymer employed, the rate of polynucleotides release can be controlled. Examples of other biodegradable polymers include, but are not limited to, poly(orthoesters) and poly(anhydrides). Depot injectable formulations may be prepared by entrapping the polynucleotides in liposomes or microemulsions which are compatible with body tissues.

Pulmonary

[000706] Pulmonary and intranasal formulations for delivery and administration are described in co-pending International Patent Publication No. WO2013151666, the contents of which is incorporated by reference in its entirety, such as, but not limited to, in paragraphs [000766] – [000781].

Coatings or Shells

[000707] Solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type may be employed as fillers in soft and

hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

Multi-dose and repeat-dose administration

[000708] In some embodiments, compounds and/or compositions of the present invention may be administered in two or more doses (referred to herein as “multi-dose administration”). Such doses may comprise the same components or may comprise components not included in a previous dose. Such doses may comprise the same mass and/or volume of components or an altered mass and/or volume of components in comparison to a previous dose. In some embodiments, multi-dose administration may comprise repeat-dose administration. As used herein, the term “repeat-dose administration” refers to two or more doses administered consecutively or within a regimen of repeat doses comprising substantially the same components provided at substantially the same mass and/or volume. In some embodiments, subjects may display a repeat-dose response. As used herein, the term “repeat-dose response” refers to a response in a subject to a repeat-dose that differs from that of another dose administered within a repeat-dose administration regimen. In some embodiments, such a response may be the expression of a protein in response to a repeat-dose comprising mRNA. In such embodiments, protein expression may be elevated in comparison to another dose administered within a repeat-dose administration regimen or protein expression may be reduced in comparison to another dose administered within a repeat-dose administration regimen. Alteration of protein expression may be from about 1% to about 20%, from about 5% to about 50% from about 10% to about 60%, from about 25% to about 75%, from about 40% to about 100% and/or at least 100%. A reduction in expression of mRNA administered as part of a repeat-dose regimen, wherein the level of protein translated from the administered RNA is reduced by more than 40% in comparison to another dose within the repeat-dose regimen is referred to herein as “repeat-dose resistance.”

Properties of the Pharmaceutical Compositions

[000709] The pharmaceutical compositions described herein can be characterized by one or more of the following properties:

Bioavailability

[000710] The polynucleotides, when formulated into a composition with a delivery agent as described herein, can exhibit an increase in bioavailability as compared to a composition lacking a delivery agent as described herein. As used herein, the term “bioavailability” refers to the systemic availability of a given amount of polynucleotides administered to a mammal. Bioavailability can be assessed by measuring the area under the curve (AUC) or the maximum serum or plasma concentration ($C_{\max}$) of the unchanged form of a compound following administration of the compound to a mammal. AUC is a determination of the area under the curve plotting the serum or plasma concentration of a compound along the ordinate (Y-axis) against time along the abscissa (X-axis).

Generally, the AUC for a particular compound can be calculated using methods known to those of ordinary skill in the art and as described in G. S. Banker, *Modern Pharmaceutics, Drugs and the Pharmaceutical Sciences*, v. 72, Marcel Dekker, New York, Inc., 1996, herein incorporated by reference in its entirety.

[000711] The $C_{\max}$ value is the maximum concentration of the compound achieved in the serum or plasma of a mammal following administration of the compound to the mammal. The $C_{\max}$ value of a particular compound can be measured using methods known to those of ordinary skill in the art. The phrases “increasing bioavailability” or “improving the pharmacokinetics,” as used herein mean that the systemic availability of a first polynucleotides, measured as AUC, $C_{\max}$, or $C_{\min}$ in a mammal is greater, when co-administered with a delivery agent as described herein, than when such co-administration does not take place. In some embodiments, the bioavailability of the polynucleotides can increase by at least about 2%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%.

[000712] In some embodiments, liquid formulations of polynucleotides may have varying in vivo half-life, requiring modulation of doses to yield a therapeutic effect. To address this, in some embodiments of the present invention, polynucleotides formulations may be designed to improve bioavailability and/or therapeutic effect during repeat administrations. Such formulations may enable sustained release of polynucleotides

and/or reduce polynucleotide degradation rates by nucleases. In some embodiments, suspension formulations are provided comprising polynucleotides, water immiscible oil depots, surfactants and/or co-surfactants and/or co-solvents. Combinations of oils and surfactants may enable suspension formulation with polynucleotides. Delivery of polynucleotides in a water immiscible depot may be used to improve bioavailability through sustained release of polynucleotides from the depot to the surrounding physiologic environment and/or prevent polynucleotide degradation by nucleases.

[000713] In some embodiments, cationic nanoparticles comprising combinations of divalent and monovalent cations may be formulated with polynucleotides. Such nanoparticles may form spontaneously in solution over a given period (e.g. hours, days, etc.). Such nanoparticles do not form in the presence of divalent cations alone or in the presence of monovalent cations alone. The delivery of polynucleotides in cationic nanoparticles or in one or more depot comprising cationic nanoparticles may improve polynucleotide bioavailability by acting as a long-acting depot and/or reducing the rate of degradation by nucleases.

Therapeutic Window

[000714] The polynucleotides, when formulated into a composition with a delivery agent as described herein, can exhibit an increase in the therapeutic window of the administered polynucleotides composition as compared to the therapeutic window of the administered polynucleotides composition lacking a delivery agent as described herein. As used herein “therapeutic window” refers to the range of plasma concentrations, or the range of levels of therapeutically active substance at the site of action, with a high probability of eliciting a therapeutic effect. In some embodiments, the therapeutic window of the polynucleotides when co-administered with a delivery agent as described herein can increase by at least about 2%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%.

Volume of Distribution

[000715] The polynucleotides, when formulated into a composition with a delivery agent as described herein, can exhibit an improved volume of distribution (V_{dist}), e.g., reduced or targeted, relative to a composition lacking a delivery agent as described herein. The volume of distribution (V_{dist}) relates the amount of the drug in the body to the concentration of the drug in the blood or plasma. As used herein, the term “volume of distribution” refers to the fluid volume that would be required to contain the total amount of the drug in the body at the same concentration as in the blood or plasma: V_{dist} equals the amount of drug in the body/concentration of drug in blood or plasma. For example, for a 10 mg dose and a plasma concentration of 10 mg/L, the volume of distribution would be 1 liter. The volume of distribution reflects the extent to which the drug is present in the extravascular tissue. A large volume of distribution reflects the tendency of a compound to bind to the tissue components compared with plasma protein binding. In a clinical setting, V_{dist} can be used to determine a loading dose to achieve a steady state concentration. In some embodiments, the volume of distribution of the polynucleotides when co-administered with a delivery agent as described herein can decrease at least about 2%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%.

Biological Effect

[000716] In one embodiment, the biological effect of the modified mRNA delivered to the animals may be categorized by analyzing the protein expression in the animals. The protein expression may be determined from analyzing a biological sample collected from a mammal administered the modified mRNA of the present invention. In one embodiment, the expression protein encoded by the modified mRNA administered to the mammal of at least 50 pg/ml may be preferred. For example, a protein expression of 50-200 pg/ml for the protein encoded by the modified mRNA delivered to the mammal may be seen as a therapeutically effective amount of protein in the mammal.

Detection of Polynucleotides Acids by Mass Spectrometry

[000717] Mass spectrometry (MS) is an analytical technique that can provide structural and molecular mass/concentration information on molecules after their conversion to

ions. The molecules are first ionized to acquire positive or negative charges and then they travel through the mass analyzer to arrive at different areas of the detector according to their mass/charge (m/z) ratio. Methods of detecting polynucleotides are described in co-pending International Patent Publication No. WO2015038892, the contents of which is incorporated by reference in its entirety, such as, but not limited to, in paragraphs [0001055] – [0001067].

V. Uses of polynucleotides of the Invention

[000718] The polynucleotides of the present invention are designed, in preferred embodiments, to provide for avoidance or evasion of deleterious bio-responses such as the immune response and/or degradation pathways, overcoming the threshold of expression and/or improving protein production capacity, improved expression rates or translation efficiency, improved drug or protein half life and/or protein concentrations, optimized protein localization, to improve one or more of the stability and/or clearance in tissues, receptor uptake and/or kinetics, cellular access by the compositions, engagement with translational machinery, secretion efficiency (when applicable), accessibility to circulation, and/or modulation of a cell's status, function and/or activity.

Therapeutics

Therapeutic Agents

[000719] The polynucleotides of the present invention, such as modified nucleic acids and modified RNAs, and the proteins translated from them described herein can be used as therapeutic or prophylactic agents. They are provided for use in medicine. For example, a polynucleotide described herein can be administered to a subject, wherein the polynucleotides is translated *in vivo* to produce a therapeutic or prophylactic polypeptide in the subject. Provided are compositions, methods, kits, and reagents for diagnosis, treatment or prevention of a disease or condition in humans and other mammals. The active therapeutic agents of the invention include polynucleotides, cells containing polynucleotides or polypeptides translated from the polynucleotides.

[000720] In certain embodiments, provided herein are combination therapeutics containing one or more polynucleotides containing translatable regions that encode for a protein or proteins that boost a mammalian subject's immunity along with a protein that induces antibody-dependent cellular toxicity. For example, provided herein are

therapeutics containing one or more nucleic acids that encode trastuzumab and granulocyte-colony stimulating factor (G-CSF). In particular, such combination therapeutics are useful in Her2+ breast cancer patients who develop induced resistance to trastuzumab. (See, e.g., Albrecht, Immunotherapy. 2(6):795-8 (2010)).

[000721] Provided herein are methods of inducing translation of a recombinant polypeptide in a cell population using the polynucleotides described herein. Such translation can be *in vivo*, *ex vivo*, *in culture*, or *in vitro*. The cell population is contacted with an effective amount of a composition containing a nucleic acid that has at least one nucleoside modification, and a translatable region encoding the recombinant polypeptide. The population is contacted under conditions such that the nucleic acid is localized into one or more cells of the cell population and the recombinant polypeptide is translated in the cell from the nucleic acid.

[000722] An “effective amount” of the composition is provided based, at least in part, on the target tissue, target cell type, means of administration, physical characteristics of the nucleic acid (e.g., size, and extent of modified nucleosides), and other determinants. In general, an effective amount of the composition provides efficient protein production in the cell, preferably more efficient than a composition containing a corresponding unmodified nucleic acid. Increased efficiency may be demonstrated by increased cell transfection (i.e., the percentage of cells transfected with the nucleic acid), increased protein translation from the nucleic acid, decreased nucleic acid degradation (as demonstrated, e.g., by increased duration of protein translation from a modified nucleic acid), or reduced innate immune response of the host cell.

[000723] Aspects of the invention are directed to methods of inducing *in vivo* translation of a recombinant polypeptide in a mammalian subject in need thereof. Therein, an effective amount of a composition containing a nucleic acid that has at least one structural or chemical modification and a translatable region encoding the recombinant polypeptide is administered to the subject using the delivery methods described herein. The nucleic acid is provided in an amount and under other conditions such that the nucleic acid is localized into a cell of the subject and the recombinant polypeptide is translated in the cell from the nucleic acid. The cell in which the nucleic

acid is localized, or the tissue in which the cell is present, may be targeted with one or more than one rounds of nucleic acid administration.

[000724] In certain embodiments, the administered polynucleotides directs production of one or more recombinant polypeptides that provide a functional activity which is substantially absent in the cell, tissue or organism in which the recombinant polypeptide is translated. For example, the missing functional activity may be enzymatic, structural, or gene regulatory in nature. In related embodiments, the administered polynucleotides directs production of one or more recombinant polypeptides that increases (e.g., synergistically) a functional activity which is present but substantially deficient in the cell in which the recombinant polypeptide is translated.

[000725] In other embodiments, the administered polynucleotides directs production of one or more recombinant polypeptides that replace a polypeptide (or multiple polypeptides) that is substantially absent in the cell in which the recombinant polypeptide is translated. Such absence may be due to genetic mutation of the encoding gene or regulatory pathway thereof. In some embodiments, the recombinant polypeptide increases the level of an endogenous protein in the cell to a desirable level; such an increase may bring the level of the endogenous protein from a subnormal level to a normal level or from a normal level to a super-normal level.

[000726] Alternatively, the recombinant polypeptide functions to antagonize the activity of an endogenous protein present in, on the surface of, or secreted from the cell. Usually, the activity of the endogenous protein is deleterious to the subject; for example, due to mutation of the endogenous protein resulting in altered activity or localization. Additionally, the recombinant polypeptide antagonizes, directly or indirectly, the activity of a biological moiety present in, on the surface of, or secreted from the cell. Examples of antagonized biological moieties include lipids (e.g., cholesterol), a lipoprotein (e.g., low density lipoprotein), a nucleic acid, a carbohydrate, a protein toxin such as shiga and tetanus toxins, or a small molecule toxin such as botulinum, cholera, and diphtheria toxins. Additionally, the antagonized biological molecule may be an endogenous protein that exhibits an undesirable activity, such as a cytotoxic or cytostatic activity.

[000727] The recombinant proteins described herein may be engineered for localization within the cell, potentially within a specific compartment such as the nucleus, or are

engineered for secretion from the cell or translocation to the plasma membrane of the cell.

[000728] In some embodiments, modified mRNAs and their encoded polypeptides in accordance with the present invention may be used for treatment of any of a variety of diseases, disorders, and/or conditions, including but not limited to one or more of the following: autoimmune disorders (*e.g.* diabetes, lupus, multiple sclerosis, psoriasis, rheumatoid arthritis); inflammatory disorders (*e.g.* arthritis, pelvic inflammatory disease); infectious diseases (*e.g.* viral infections (*e.g.*, HIV, HCV, RSV), bacterial infections, fungal infections, sepsis); neurological disorders (*e.g.* Alzheimer's disease, Huntington's disease; autism; Duchenne muscular dystrophy); cardiovascular disorders (*e.g.* atherosclerosis, hypercholesterolemia, thrombosis, clotting disorders, angiogenic disorders such as macular degeneration); proliferative disorders (*e.g.* cancer, benign neoplasms); respiratory disorders (*e.g.* chronic obstructive pulmonary disease); digestive disorders (*e.g.* inflammatory bowel disease, ulcers); musculoskeletal disorders (*e.g.* fibromyalgia, arthritis); endocrine, metabolic, and nutritional disorders (*e.g.* diabetes, osteoporosis); urological disorders (*e.g.* renal disease); psychological disorders (*e.g.* depression, schizophrenia); skin disorders (*e.g.* wounds, eczema); blood and lymphatic disorders (*e.g.* anemia, hemophilia); *etc.*

[000729] Diseases characterized by dysfunctional or aberrant protein activity include cystic fibrosis, sickle cell anemia, epidermolysis bullosa, amyotrophic lateral sclerosis, and glucose-6-phosphate dehydrogenase deficiency. The present invention provides a method for treating such conditions or diseases in a subject by introducing nucleic acid or cell-based therapeutics containing the polynucleotides provided herein, wherein the polynucleotides encode for a protein that antagonizes or otherwise overcomes the aberrant protein activity present in the cell of the subject. Specific examples of a dysfunctional protein are the missense mutation variants of the cystic fibrosis transmembrane conductance regulator (CFTR) gene, which produce a dysfunctional protein variant of CFTR protein, which causes cystic fibrosis.

[000730] Diseases characterized by missing (or substantially diminished such that proper (normal or physiological protein function does not occur) protein activity include cystic fibrosis, Niemann-Pick type C, β thalassemia major, Duchenne muscular

dystrophy, Hurler Syndrome, Hunter Syndrome, and Hemophilia A. Such proteins may not be present, or are essentially non-functional. The present invention provides a method for treating such conditions or diseases in a subject by introducing nucleic acid or cell-based therapeutics containing the polynucleotides provided herein, wherein the polynucleotides encode for a protein that replaces the protein activity missing from the target cells of the subject. Specific examples of a dysfunctional protein are the nonsense mutation variants of the cystic fibrosis transmembrane conductance regulator (CFTR) gene, which produce a nonfunctional protein variant of CFTR protein, which causes cystic fibrosis.

[000731] Thus, provided are methods of treating cystic fibrosis in a mammalian subject by contacting a cell of the subject with a polynucleotide having a translatable region that encodes a functional CFTR polypeptide, under conditions such that an effective amount of the CFTR polypeptide is present in the cell. Preferred target cells are epithelial, endothelial and mesothelial cells, such as the lung, and methods of administration are determined in view of the target tissue; i.e., for lung delivery, the RNA molecules are formulated for administration by inhalation.

[000732] In another embodiment, the present invention provides a method for treating hyperlipidemia in a subject, by introducing into a cell population of the subject with a modified mRNA molecule encoding Sortilin, a protein recently characterized by genomic studies, thereby ameliorating the hyperlipidemia in a subject. The *SORT1* gene encodes a trans-Golgi network (TGN) transmembrane protein called Sortilin. Genetic studies have shown that one of five individuals has a single nucleotide polymorphism, rs12740374, in the 1p13 locus of the *SORT1* gene that predisposes them to having low levels of low-density lipoprotein (LDL) and very-low-density lipoprotein (VLDL). Each copy of the minor allele, present in about 30% of people, alters LDL cholesterol by 8 mg/dL, while two copies of the minor allele, present in about 5% of the population, lowers LDL cholesterol 16 mg/dL. Carriers of the minor allele have also been shown to have a 40% decreased risk of myocardial infarction. Functional *in vivo* studies in mice describes that overexpression of *SORT1* in mouse liver tissue led to significantly lower LDL-cholesterol levels, as much as 80% lower, and that silencing *SORT1* increased LDL cholesterol

approximately 200% (Musunuru K et al. From noncoding variant to phenotype via *SORT1* at the 1p13 cholesterol locus. *Nature* 2010; 466: 714-721).

[000733] In another embodiment, the present invention provides a method for treating hematopoietic disorders, cardiovascular disease, oncology, diabetes, cystic fibrosis, neurological diseases, inborn errors of metabolism, skin and systemic disorders, and blindness. The identity of molecular targets to treat these specific diseases has been described (Templeton ed., *Gene and Cell Therapy: Therapeutic Mechanisms and Strategies*, 3rd Edition, Boca Raton, FL: CRC Press; herein incorporated by reference in its entirety).

[000734] Provided herein, are methods to prevent infection and/or sepsis in a subject at risk of developing infection and/or sepsis, the method comprising administering to a subject in need of such prevention a composition comprising a polynucleotide precursor encoding an anti-microbial polypeptide (*e.g.*, an anti-bacterial polypeptide), or a partially or fully processed form thereof in an amount sufficient to prevent infection and/or sepsis. In certain embodiments, the subject at risk of developing infection and/or sepsis may be a cancer patient. In certain embodiments, the cancer patient may have undergone a conditioning regimen. In some embodiments, the conditioning regimen may include, but is not limited to, chemotherapy, radiation therapy, or both. As a non-limiting example, a polynucleotide can encode Protein C, its zymogen or prepro-protein, the activated form of Protein C (APC) or variants of Protein C which are known in the art. The polynucleotides may be chemically modified and delivered to cells. Non-limiting examples of polypeptides which may be encoded within the chemically modified mRNAs of the present invention include those taught in US Patents 7,226,999; 7,498,305; 6,630,138 each of which is incorporated herein by reference in its entirety. These patents teach Protein C like molecules, variants and derivatives, any of which may be encoded within the chemically modified molecules of the present invention.

[000735] Further provided herein, are methods to treat infection and/or sepsis in a subject, the method comprising administering to a subject in need of such treatment a composition comprising a polynucleotide precursor encoding an anti-microbial polypeptide (*e.g.*, an anti-bacterial polypeptide), *e.g.*, an anti-microbial polypeptide described herein, or a partially or fully processed form thereof in an amount sufficient to

treat an infection and/or sepsis. In certain embodiments, the subject in need of treatment is a cancer patient. In certain embodiments, the cancer patient has undergone a conditioning regimen. In some embodiments, the conditioning regimen may include, but is not limited to, chemotherapy, radiation therapy, or both.

[000736] In certain embodiments, the subject may exhibit acute or chronic microbial infections (*e.g.*, bacterial infections). In certain embodiments, the subject may have received or may be receiving a therapy. In certain embodiments, the therapy may include, but is not limited to, radiotherapy, chemotherapy, steroids, ultraviolet radiation, or a combination thereof. In certain embodiments, the patient may suffer from a microvascular disorder. In some embodiments, the microvascular disorder may be diabetes. In certain embodiments, the patient may have a wound. In some embodiments, the wound may be an ulcer. In a specific embodiment, the wound may be a diabetic foot ulcer. In certain embodiments, the subject may have one or more burn wounds. In certain embodiments, the administration may be local or systemic. In certain embodiments, the administration may be subcutaneous. In certain embodiments, the administration may be intravenous. In certain embodiments, the administration may be oral. In certain embodiments, the administration may be topical. In certain embodiments, the administration may be by inhalation. In certain embodiments, the administration may be rectal. In certain embodiments, the administration may be vaginal.

Allergens or Environment Hypersensitivity

[000737] In certain embodiments, the polynucleotides of the present invention may be used to treat a subject with an environmental hypersensitivity or allergic reactions. As a non-limiting example, the polynucleotides of the present invention may be used to generate non-human proteins that may play a role in the induction of hypersensitivity or allergic reactions in humans or other mammals. Allergen Specific Immunotherapy (SIT) for respiratory allergic diseases is able to significantly improve symptoms as reduce the need for symptomatic medication, and may also show long-term clinical effects (*Jacobsen et al. Clinical and Translational Allergy 2012, 2:8*; the contents of which are herein incorporated by reference in its entirety). In one embodiment, the polynucleotides of the present invention may encode proteins or antigens or other triggers for an allergic reaction such as, but are not limited to, plants, grass, tree and other plant pollens,

mammal, animal hair or dander, insect, insect parts, excreta or venom, helminthes, non-mammals, birds and reptiles, foods (e.g., legumes, nuts and fish), bacteria, fungi and molds, drugs, metals, wool and latex..

[000738] In one embodiment, the polynucleotides described herein may be administered using a micro-dosing regimen in order to induce tolerance to an allergen or hypersensitivity to the environment. As a non-limiting example, the micro-dosing regimen may be administered in at least two doses (e.g., at least 2 doses, at least 3 doses, at least 4 doses, at least 5 doses, at least 6 doses, at least 7 doses, at least 8 doses, at least 9 doses, at least 10 doses, at least 11 doses, at least 12 doses, at least 13 doses, at least 14 doses, at least 15 doses, at least 16 doses, at least 17 doses, at least 18 doses, at least 19 doses, at least 20 doses, at least 21 doses, at least 22 doses, at least 23 doses, at least 24 doses, at least 25 doses, at least 26 doses, at least 27 doses, at least 28 doses, at least 29 doses, at least 30 doses, at least 31 doses, at least 32 doses, at least 33 doses, at least 34 doses, at least 35 doses, at least 36 doses, at least 37 doses, at least 38 doses, at least 39 doses, at least 40 doses, at least 41 doses, at least 42 doses, at least 43 doses, at least 44 doses, at least 45 doses, at least 46 doses, at least 47 doses, at least 48 doses, at least 49 doses, at least 50 doses, at least 51 doses, at least 52 doses, at least 53 doses, at least 54 doses, at least 55 doses, at least 56 doses, at least 57 doses, at least 58 doses, at least 59 doses, at least 60 doses, at least 61 doses, at least 62 doses, at least 63 doses, at least 64 doses, at least 65 doses, at least 66 doses, at least 67 doses, at least 68 doses, at least 69 doses, at least 70 doses, at least 71 doses, at least 72 doses, at least 73 doses, at least 74 doses, at least 75 doses, at least 76 doses, at least 77 doses, at least 78 doses, at least 79 doses, at least 80 doses, at least 81 doses, at least 82 doses, at least 83 doses, at least 84 doses, at least 85 doses, at least 86 doses, at least 87 doses, at least 88 doses, at least 89 doses, at least 90 doses, at least 91 doses, at least 92 doses, at least 93 doses, at least 94 doses, at least 95 doses, at least 96 doses, at least 97 doses, at least 98 doses, at least 99 doses, at least 100 doses or more than 100 doses), each of which are at levels as low as one millionth the desired therapeutic dose. As another non-limiting example, the micro-dosing regimen may be staged so the amount of polynucleotide administered may be increased over a pre-determined period of time (e.g., at least 1 hour, at least 2 hours, at least 3 hours, at least 4 hours, at least 5 hours, at least 6 hours, at least 7 hours, at least 8

hours, at least 9 hours, at least 10 hours, at least 11 hours, at least 12 hours, at least 13 hours, at least 14 hours, at least 15 hours, at least 16 hours, at least 17 hours, at least 18 hours, at least 19 hours, at least 20 hours, at least 21 hours, at least 22 hours, at least 23 hours, at least 1 day, at least 36 hours, at least 2 days, at least 3 days, at least 4 days, at least 5 days, at least 6 days, at least 1 week, at least 2 weeks, at least 3 weeks, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months or more than 6 months) and may rise progressively over a period of time until desired effect is achieved (e.g., an efficacy dose) or safety considerations (e.g., toxicity) are rectified.

[000739] In certain embodiments, the polynucleotides may be used to treat a subject with a medical condition that is considered to be wholly or partially related to an exogenous antigen such as, but not limited to, Asthma, Rhino-conjunctivitis, Lupus Erythematosus, Crohn's disease, Ulcerative Colitis, Celiac disease or Type 1 Diabetes Mellitus, multiple sclerosis, cancers. Antigen-specific immunotherapy has been shown useful to induce immune tolerance, in a number of autoimmune disorders, including in diabetes (reviewed in *Roep and Peakman, Cold Spring Harb Perspect Med* 2012, 2:a007781, *Han et al, Am J Transl Res*, 2013 Vol 5(4):379-392, and *Peakman, F1000 Biology Reports* 2012, 4:19;the content of each of which is herein incorporated by reference in its entirety), multiple sclerosis (reviewed in *Badawi et al, Clin Immunol.*, 2012, Vol 144(2), 127–138;the content of which is herein incorporated by reference in its entirety).), and in cancer where antigen from the patient's own tumor can be used as adjuvant therapy to elicit an long-term anti-tumor immune response immune response (*Baars et al, British Journal of Cancer*, 2002, 86, 1230 – 1234;the content of which is herein incorporated by reference in its entirety). Dosing may be performed as micro-dosing regimen described herein.

[000740] In certain embodiments, the polynucleotides may be used to treat a subject to induce tolerance to a potential antigen/allergen that is administered in a therapeutic indication. Dosing may be performed as micro-dosing regimen described herein.

[000741] In certain embodiments, to induce tolerance the admin polynucleotides described herein may be administered by a route of administration described herein such

as, but not limited to, intradermal, sublingual, intranasal, oral, intramuscular, subcutaneous, intravenous, vaginal and rectal.

[000742] Other aspects of the present disclosure relate to transplantation of cells containing polynucleotides to a mammalian subject. Administration of cells to mammalian subjects is known to those of ordinary skill in the art, and include, but is not limited to, local implantation (e.g., topical or subcutaneous administration), organ delivery or systemic injection (e.g., intravenous injection or inhalation), and the formulation of cells in pharmaceutically acceptable carrier. Such compositions containing polynucleotides can be formulated for administration intramuscularly, transarterially, intraperitoneally, intravenously, intranasally, subcutaneously, endoscopically, transdermally, or intrathecally. In some embodiments, the composition may be formulated for extended release.

[000743] The subject to whom the therapeutic agent may be administered suffers from or may be at risk of developing a disease, disorder, or deleterious condition. Provided are methods of identifying, diagnosing, and classifying subjects on these bases, which may include clinical diagnosis, biomarker levels, genome-wide association studies (GWAS), and other methods known in the art.

[000744] In another embodiment, it may be useful to optimize the expression of a specific polypeptide in a cell line or collection of cell lines of potential interest, particularly a polypeptide of interest such as a protein variant of a reference protein having a known activity. In one embodiment, provided is a method of optimizing expression of a polypeptide of interest in a target cell, by providing a plurality of target cell types, and independently contacting with each of the plurality of target cell types a modified mRNA encoding a polypeptide. Additionally, culture conditions may be altered to increase protein production efficiency. Subsequently, the presence and/or level of the polypeptide of interest in the plurality of target cell types is detected and/or quantitated, allowing for the optimization of a polypeptide of interest's expression by selection of an efficient target cell and cell culture conditions relating thereto. Such methods may be useful when the polypeptide of interest contains one or more post-translational modifications or has substantial tertiary structure, which often complicate efficient protein production.

Protein recovery

[000745] The protein of interest may be preferably recovered from the culture medium as a secreted polypeptide, or it can be recovered from host cell lysates if expressed without a secretory signal. It may be necessary to purify the protein of interest from other recombinant proteins and host cell proteins in a way that substantially homogenous preparations of the protein of interest are obtained. The cells and/or particulate cell debris may be removed from the culture medium or lysate. The product of interest may then be purified from contaminant soluble proteins, polypeptides and nucleic acids by, for example, fractionation on immunoaffinity or ion-exchange columns, ethanol precipitation, reverse phase HPLC (RP-HPLC), SEPHADEX® chromatography, chromatography on silica or on a cation exchange resin such as DEAE. Methods of purifying a protein heterologous expressed by a host cell are well known in the art.

[000746] Methods and compositions described herein may be used to produce proteins which are capable of attenuating or blocking the endogenous agonist biological response and/or antagonizing a receptor or signaling molecule in a mammalian subject. For example, IL-12 and IL-23 receptor signaling may be enhanced in chronic autoimmune disorders such as multiple sclerosis and inflammatory diseases such as rheumatoid arthritis, psoriasis, lupus erythematosus, ankylosing spondylitis and Chron's disease (Kikly K, Liu L, Na S, Sedgwich JD (2006) Cur. Opin. Immunol. 18(6): 670-5). In another embodiment, a nucleic acid encodes an antagonist for chemokine receptors. Chemokine receptors CXCR-4 and CCR-5 are required for HIV entry into host cells (Arenzana-Seisdedos F et al, (1996) Nature. Oct 3; 383 (6599):400).

Gene Silencing

[000747] The polynucleotides described herein are useful to silence (i.e., prevent or substantially reduce) expression of one or more target genes in a cell population. A polynucleotide encoding a polypeptide of interest capable of directing sequence-specific histone H3 methylation is introduced into the cells in the population under conditions such that the polypeptide is translated and reduces gene transcription of a target gene via histone H3 methylation and subsequent heterochromatin formation. In some embodiments, the silencing mechanism is performed on a cell population present in a mammalian subject. By way of non-limiting example, a useful target gene is a mutated

Janus Kinase-2 family member, wherein the mammalian subject expresses the mutant target gene suffers from a myeloproliferative disease resulting from aberrant kinase activity.

[000748] Co-administration of polynucleotides and RNAi agents are also provided herein.

Modulation of Biological Pathways

[000749] The rapid translation polynucleotides introduced into cells provides a desirable mechanism of modulating target biological pathways. Such modulation includes antagonism or agonism of a given pathway. In one embodiment, a method is provided for antagonizing a biological pathway in a cell by contacting the cell with an effective amount of a composition comprising a polynucleotide encoding a polypeptide of interest, under conditions such that the polynucleotides is localized into the cell and the polypeptide is capable of being translated in the cell from the polynucleotides, wherein the polypeptide inhibits the activity of a polypeptide functional in the biological pathway. Exemplary biological pathways are those defective in an autoimmune or inflammatory disorder such as multiple sclerosis, rheumatoid arthritis, psoriasis, lupus erythematosus, ankylosing spondylitis colitis, or Crohn's disease; in particular, antagonism of the IL-12 and IL-23 signaling pathways are of particular utility. (See Kikly K, Liu L, Na S, Sedgwick JD (2006) Curr. Opin. Immunol. 18 (6): 670–5).

[000750] Further, provided are polynucleotides encoding an antagonist for chemokine receptors; chemokine receptors CXCR-4 and CCR-5 are required for, e.g., HIV entry into host cells (Arenzana-Seisdedos F et al, (1996) Nature. Oct 3;383(6599):400).

[000751] Alternatively, provided are methods of agonizing a biological pathway in a cell by contacting the cell with an effective amount of a polynucleotide encoding a recombinant polypeptide under conditions such that the nucleic acid is localized into the cell and the recombinant polypeptide is capable of being translated in the cell from the nucleic acid, and the recombinant polypeptide induces the activity of a polypeptide functional in the biological pathway. Exemplary agonized biological pathways include pathways that modulate cell fate determination. Such agonization is reversible or, alternatively, irreversible.

Expression of Ligand or Receptor on Cell Surface

[000752] In some aspects and embodiments of the aspects described herein, the polynucleotides described herein can be used to express a ligand or ligand receptor on the surface of a cell (*e.g.*, a homing moiety). A ligand or ligand receptor moiety attached to a cell surface can permit the cell to have a desired biological interaction with a tissue or an agent *in vivo*. A ligand can be an antibody, an antibody fragment, an aptamer, a peptide, a vitamin, a carbohydrate, a protein or polypeptide, a receptor, *e.g.*, cell-surface receptor, an adhesion molecule, a glycoprotein, a sugar residue, a therapeutic agent, a drug, a glycosaminoglycan, or any combination thereof. For example, a ligand can be an antibody that recognizes a cancer-cell specific antigen, rendering the cell capable of preferentially interacting with tumor cells to permit tumor-specific localization of a modified cell. A ligand can confer the ability of a cell composition to accumulate in a tissue to be treated, since a preferred ligand may be capable of interacting with a target molecule on the external face of a tissue to be treated. Ligands having limited cross-reactivity to other tissues are generally preferred.

[000753] In some cases, a ligand can act as a homing moiety which permits the cell to target to a specific tissue or interact with a specific ligand. Such homing moieties can include, but are not limited to, any member of a specific binding pair, antibodies, monoclonal antibodies, or derivatives or analogs thereof, including without limitation: Fv fragments, single chain Fv (scFv) fragments, Fab' fragments, F(ab')₂ fragments, single domain antibodies, camelized antibodies and antibody fragments, humanized antibodies and antibody fragments, and multivalent versions of the foregoing; multivalent binding reagents including without limitation: monospecific or bispecific antibodies, such as disulfide stabilized Fv fragments, scFv tandems ((SCFV)₂ fragments), diabodies, tribodies or tetrabodies, which typically are covalently linked or otherwise stabilized (*i.e.*, leucine zipper or helix stabilized) scFv fragments; and other homing moieties include for example, aptamers, receptors, and fusion proteins.

[000754] In some embodiments, the homing moiety may be a surface-bound antibody, which can permit tuning of cell targeting specificity. This is especially useful since highly specific antibodies can be raised against an epitope of interest for the desired targeting site. In one embodiment, multiple antibodies are expressed on the surface of a

cell, and each antibody can have a different specificity for a desired target. Such approaches can increase the avidity and specificity of homing interactions.

[000755] A skilled artisan can select any homing moiety based on the desired localization or function of the cell, for example an estrogen receptor ligand, such as tamoxifen, can target cells to estrogen-dependent breast cancer cells that have an increased number of estrogen receptors on the cell surface. Other non-limiting examples of ligand/receptor interactions include CCRI (*e.g.*, for treatment of inflamed joint tissues or brain in rheumatoid arthritis, and/or multiple sclerosis), CCR7, CCR8 (*e.g.*, targeting to lymph node tissue), CCR6, CCR9, CCR10 (*e.g.*, to target to intestinal tissue), CCR4, CCR10 (*e.g.*, for targeting to skin), CXCR4 (*e.g.*, for general enhanced transmigration), HCELL (*e.g.*, for treatment of inflammation and inflammatory disorders, bone marrow), Alpha4beta7 (*e.g.*, for intestinal mucosa targeting), VLA-4/VCAM-1 (*e.g.*, targeting to endothelium). In general, any receptor involved in targeting (*e.g.*, cancer metastasis) can be harnessed for use in the methods and compositions described herein.

VI. Kits and Devices

Kits

[000756] The invention provides a variety of kits for conveniently and/or effectively carrying out methods of the present invention. Typically kits will comprise sufficient amounts and/or numbers of components to allow a user to perform multiple treatments of a subject(s) and/or to perform multiple experiments.

[000757] In one aspect, the present invention provides kits comprising the molecules (polynucleotides) of the invention. In one embodiment, the kit comprises one or more functional antibodies or function fragments thereof.

[000758] Said kits can be for protein production, comprising a first polynucleotides comprising a translatable region. The kit may further comprise packaging and instructions and/or a delivery agent to form a formulation composition. The delivery agent may comprise a saline, a buffered solution, a lipidoid or any delivery agent disclosed herein.

[000759] In one embodiment, the buffer solution may include sodium chloride, calcium chloride, phosphate and/or EDTA. In another embodiment, the buffer solution may include, but is not limited to, saline, saline with 2mM calcium, 5% sucrose, 5% sucrose

with 2mM calcium, 5% Mannitol, 5% Mannitol with 2mM calcium, Ringer's lactate, sodium chloride, sodium chloride with 2mM calcium and mannose (See e.g., U.S. Pub. No. 20120258046; herein incorporated by reference in its entirety). In a further embodiment, the buffer solutions may be precipitated or it may be lyophilized. The amount of each component may be varied to enable consistent, reproducible higher concentration saline or simple buffer formulations. The components may also be varied in order to increase the stability of modified RNA in the buffer solution over a period of time and/or under a variety of conditions. In one aspect, the present invention provides kits for protein production, comprising: a polynucleotide comprising a translatable region, provided in an amount effective to produce a desired amount of a protein encoded by the translatable region when introduced into a target cell; a second polynucleotide comprising an inhibitory nucleic acid, provided in an amount effective to substantially inhibit the innate immune response of the cell; and packaging and instructions.

[000760] In one aspect, the present invention provides kits for protein production, comprising a polynucleotide comprising a translatable region, wherein the polynucleotide exhibits reduced degradation by a cellular nuclease, and packaging and instructions.

[000761] In one aspect, the present invention provides kits for protein production, comprising a polynucleotide comprising a translatable region, wherein the polynucleotide exhibits reduced degradation by a cellular nuclease, and a mammalian cell suitable for translation of the translatable region of the first nucleic acid.

Devices

[000762] The present invention provides for devices which may incorporate polynucleotides that encode polypeptides of interest. These devices contain in a stable formulation the reagents to synthesize a polynucleotide in a formulation available to be immediately delivered to a subject in need thereof, such as a human patient

[000763] Devices for administration may be employed to deliver the polynucleotides of the present invention according to single, multi- or split-dosing regimens taught herein. Such devices are taught in, for example, International Publication No. WO2013151666 (Attorney Docket Number M300), the contents of which are incorporated herein by reference in their entirety.

[000764] Method and devices known in the art for multi-administration to cells, organs and tissues are contemplated for use in conjunction with the methods and compositions disclosed herein as embodiments of the present invention. These include, for example, those methods and devices having multiple needles, hybrid devices employing for example lumens or catheters as well as devices utilizing heat, electric current or radiation driven mechanisms.

[000765] According to the present invention, these multi-administration devices may be utilized to deliver the single, multi- or split doses contemplated herein. Such devices are taught for example in, International Publication No. WO2013151666 (Attorney Docket Number M300), the contents of which are incorporated herein by reference in their entirety.

[000766] In one embodiment, the polynucleotide is administered subcutaneously or intramuscularly via at least 3 needles to three different, optionally adjacent, sites simultaneously, or within a 60 minutes period (e.g., administration to 4, 5, 6, 7, 8, 9, or 10 sites simultaneously or within a 60 minute period).

Methods and Devices utilizing catheters and/or lumens

[000767] Methods and devices using catheters and lumens may be employed to administer the polynucleotides of the present invention on a single, multi- or split dosing schedule. Such methods and devices are described in International Publication No. WO2013151666 (Attorney Docket Number M300), the contents of which are incorporated herein by reference in their entirety.

Methods and Devices utilizing electrical current

[000768] Methods and devices utilizing electric current may be employed to deliver the polynucleotides of the present invention according to the single, multi- or split dosing regimens taught herein. Such methods and devices are described in International Publication No. WO2013151666 (Attorney Docket Number M300), the contents of which are incorporated herein by reference in their entirety.

VII. Definitions

[000769] At various places in the present specification, substituents of compounds of the present disclosure are disclosed in groups or in ranges. It is specifically intended that

the present disclosure include each and every individual subcombination of the members of such groups and ranges

[000770] *About*: As used herein, the term “about” means +/- 10% of the recited value.

[000771] *Administered in combination*: As used herein, the term “administered in combination” or “combined administration” means that two or more agents are administered to a subject at the same time or within an interval such that there may be an overlap of an effect of each agent on the patient. In some embodiments, they are administered within about 60, 30, 15, 10, 5, or 1 minute of one another. In some embodiments, the administrations of the agents are spaced sufficiently closely together such that a combinatorial (*e.g.*, a synergistic) effect is achieved.

[000772] *Adjuvant*: As used herein, the term “adjuvant” means a substance that enhances a subject’s immune response to an antigen.

[000773] *Animal*: As used herein, the term “animal” refers to any member of the animal kingdom. In some embodiments, “animal” refers to humans at any stage of development. In some embodiments, “animal” refers to non-human animals at any stage of development. In certain embodiments, the non-human animal is a mammal (*e.g.*, a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, or a pig). In some embodiments, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, and worms. In some embodiments, the animal is a transgenic animal, genetically-engineered animal, or a clone.

[000774] *Antigen*: As used herein, the term “antigen” refers to the substance that binds specifically to the respective antibody. An antigen may originate either from the body, such as cancer antigen used herein, or from the external environment, for instance, from infectious agents.

[000775] *Antigens of interest or desired antigens*: As used herein, the terms “antigens of interest” or “desired antigens” include those proteins and other biomolecules provided herein that are immunospecifically bound by the antibodies and fragments, mutants, variants, and alterations thereof described herein. Examples of antigens of interest include, but are not limited to, insulin, insulin-like growth factor, hGH, tPA, cytokines, such as interleukins (IL), *e.g.*, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, interferon (IFN) alpha, IFN beta,

IFN gamma, IFN omega or IFN tau, tumor necrosis factor (TNF), such as TNF alpha and TNF beta, TNF gamma, TRAIL; G-CSF, GM-CSF, M-CSF, MCP-1 and VEGF.

[000776] *Approximately:* As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[000777] *Associated with:* As used herein, the terms “associated with,” “conjugated,” “linked,” “attached,” and “tethered,” when used with respect to two or more moieties, means that the moieties are physically associated or connected with one another, either directly or via one or more additional moieties that serves as a linking agent, to form a structure that is sufficiently stable so that the moieties remain physically associated under the conditions in which the structure is used, *e.g.*, physiological conditions. An “association” need not be strictly through direct covalent chemical bonding. It may also suggest ionic or hydrogen bonding or a hybridization based connectivity sufficiently stable such that the “associated” entities remain physically associated.

[000778] *Bifunctional:* As used herein, the term “bifunctional” refers to any substance, molecule or moiety which is capable of or maintains at least two functions. The functions may affect the same outcome or a different outcome. The structure that produces the function may be the same or different. For example, bifunctional modified RNAs of the present invention may encode a cytotoxic peptide (a first function) while those nucleosides which comprise the encoding RNA are, in and of themselves, cytotoxic (second function). In this example, delivery of the bifunctional modified RNA to a cancer cell would produce not only a peptide or protein molecule which may ameliorate or treat the cancer but would also deliver a cytotoxic payload of nucleosides to the cell should degradation, instead of translation of the modified RNA, occur.

[000779] *Biocompatible*: As used herein, the term “biocompatible” means compatible with living cells, tissues, organs or systems posing little to no risk of injury, toxicity or rejection by the immune system.

[000780] *Biodegradable*: As used herein, the term “biodegradable” means capable of being broken down into innocuous products by the action of living things.

[000781] *Biologically active*: As used herein, the phrase “biologically active” refers to a characteristic of any substance that has activity in a biological system and/or organism. For instance, a substance that, when administered to an organism, has a biological effect on that organism, is considered to be biologically active. In particular embodiments, a polynucleotide of the present invention may be considered biologically active if even a portion of the polynucleotides is biologically active or mimics an activity considered biologically relevant.

[000782] *Cancer stem cells*: As used herein, “cancer stem cells” are cells that can undergo self-renewal and/or abnormal proliferation and differentiation to form a tumor.

[000783] *Chimera*: As used herein, “chimera” is an entity having two or more incongruous or heterogeneous parts or regions.

[000784] *Chimeric polynucleotide*: As used herein, “chimeric polynucleotides” are those nucleic acid polymers having portions or regions which differ in size and/or chemical modification pattern, chemical modification position, chemical modification percent or chemical modification population and combinations of the foregoing.

[000785] *Compound*: As used herein, the term “compound,” is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted.

[000786] The compounds described herein can be asymmetric (*e.g.*, having one or more stereocenters). All stereoisomers, such as enantiomers and diastereomers, are intended unless otherwise indicated. Compounds of the present disclosure that contain asymmetrically substituted carbon atoms can be isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically active starting materials are known in the art, such as by resolution of racemic mixtures or by stereoselective synthesis. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present disclosure. Cis and trans geometric isomers of

the compounds of the present disclosure are described and may be isolated as a mixture of isomers or as separated isomeric forms.

[000787] Compounds of the present disclosure also include tautomeric forms.

Tautomeric forms result from the swapping of a single bond with an adjacent double bond and the concomitant migration of a proton. Tautomeric forms include prototrophic tautomers which are isomeric protonation states having the same empirical formula and total charge. Examples prototrophic tautomers include ketone – enol pairs, amide – imidic acid pairs, lactam – lactim pairs, amide – imidic acid pairs, enamine – imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, such as, 1H- and 3H-imidazole, 1H-, 2H- and 4H- 1,2,4-triazole, 1H- and 2H-isoindole, and 1H- and 2H-pyrazole. Tautomeric forms can be in equilibrium or sterically locked into one form by appropriate substitution.

[000788] Compounds of the present disclosure also include all of the isotopes of the atoms occurring in the intermediate or final compounds. “Isotopes” refers to atoms having the same atomic number but different mass numbers resulting from a different number of neutrons in the nuclei. For example, isotopes of hydrogen include tritium and deuterium.

[000789] The compounds and salts of the present disclosure can be prepared in combination with solvent or water molecules to form solvates and hydrates by routine methods.

[000790] *Committed:* As used herein, the term “committed” means, when referring to a cell, when the cell is far enough into the differentiation pathway where, under normal circumstances, it will continue to differentiate into a specific cell type or subset of cell type instead of into a different cell type or reverting to a lesser differentiated cell type.

[000791] *Conserved:* As used herein, the term “conserved” refers to nucleotides or amino acid residues of a polynucleotide sequence or polypeptide sequence, respectively, that are those that occur unaltered in the same position of two or more sequences being compared. Nucleotides or amino acids that are relatively conserved are those that are conserved amongst more related sequences than nucleotides or amino acids appearing elsewhere in the sequences.

[000792] In some embodiments, two or more sequences are said to be “completely conserved” if they are 100% identical to one another. In some embodiments, two or more sequences are said to be “highly conserved” if they are at least 70% identical, at least 80% identical, at least 90% identical, or at least 95% identical to one another. In some embodiments, two or more sequences are said to be “highly conserved” if they are about 70% identical, about 80% identical, about 90% identical, about 95%, about 98%, or about 99% identical to one another. In some embodiments, two or more sequences are said to be “conserved” if they are at least 30% identical, at least 40% identical, at least 50% identical, at least 60% identical, at least 70% identical, at least 80% identical, at least 90% identical, or at least 95% identical to one another. In some embodiments, two or more sequences are said to be “conserved” if they are about 30% identical, about 40% identical, about 50% identical, about 60% identical, about 70% identical, about 80% identical, about 90% identical, about 95% identical, about 98% identical, or about 99% identical to one another. Conservation of sequence may apply to the entire length of a polynucleotide or polypeptide or may apply to a portion, region or feature thereof.

[000793] *Controlled Release:* As used herein, the term “controlled release” refers to a pharmaceutical composition or compound release profile that conforms to a particular pattern of release to effect a therapeutic outcome.

[000794] *Cyclic or Cyclized:* As used herein, the term “cyclic” refers to the presence of a continuous loop. Cyclic molecules need not be circular, only joined to form an unbroken chain of subunits. Cyclic molecules such as the engineered RNA or mRNA of the present invention may be single units or multimers or comprise one or more components of a complex or higher order structure.

[000795] *Cytostatic:* As used herein, “cytostatic” refers to inhibiting, reducing, suppressing the growth, division, or multiplication of a cell (*e.g.*, a mammalian cell (*e.g.*, a human cell)), bacterium, virus, fungus, protozoan, parasite, prion, or a combination thereof.

[000796] *Cytotoxic:* As used herein, “cytotoxic” refers to killing or causing injurious, toxic, or deadly effect on a cell (*e.g.*, a mammalian cell (*e.g.*, a human cell)), bacterium, virus, fungus, protozoan, parasite, prion, or a combination thereof.

[000797] *Delivery*: As used herein, “delivery” refers to the act or manner of delivering a compound, substance, entity, moiety, cargo or payload.

[000798] *Delivery Agent*: As used herein, “delivery agent” refers to any substance which facilitates, at least in part, the *in vivo* delivery of a polynucleotide to targeted cells.

[000799] *Destabilized*: As used herein, the term “destable,” “destabilize,” or “destabilizing region” means a region or molecule that is less stable than a starting, wild-type or native form of the same region or molecule.

[000800] *Detectable label*: As used herein, “detectable label” refers to one or more markers, signals, or moieties which are attached, incorporated or associated with another entity that is readily detected by methods known in the art including radiography, fluorescence, chemiluminescence, enzymatic activity, absorbance and the like. Detectable labels include radioisotopes, fluorophores, chromophores, enzymes, dyes, metal ions, ligands such as biotin, avidin, streptavidin and haptens, quantum dots, and the like. Detectable labels may be located at any position in the peptides or proteins disclosed herein. They may be within the amino acids, the peptides, or proteins, or located at the N- or C- termini.

[000801] *Diastereomer*: As used herein, the term “diastereomer,” means stereoisomers that are not mirror images of one another and are non-superimposable on one another.

[000802] *Digest*: As used herein, the term “digest” means to break apart into smaller pieces or components. When referring to polypeptides or proteins, digestion results in the production of peptides.

[000803] *Differentiated cell*: As used herein, the term “differentiated cell” refers to any somatic cell that is not, in its native form, pluripotent. Differentiated cell also encompasses cells that are partially differentiated.

[000804] *Differentiation*: As used herein, the term “differentiation factor” refers to a developmental potential altering factor such as a protein, RNA or small molecule that can induce a cell to differentiate to a desired cell-type.

[000805] *Differentiate*: As used herein, “differentiate” refers to the process where an uncommitted or less committed cell acquires the features of a committed cell.

[000806] *Distal*: As used herein, the term “distal” means situated away from the center or away from a point or region of interest.

[000807] *Dosing regimen*: As used herein, a “dosing regimen” is a schedule of administration or physician determined regimen of treatment, prophylaxis, or palliative care.

[000808] *Dose splitting factor (DSF)*-ratio of PUD of dose split treatment divided by PUD of total daily dose or single unit dose. The value is derived from comparison of dosing regimens groups.

[000809] *Enantiomer*: As used herein, the term “enantiomer” means each individual optically active form of a compound of the invention, having an optical purity or enantiomeric excess (as determined by methods standard in the art) of at least 80% (i.e., at least 90% of one enantiomer and at most 10% of the other enantiomer), preferably at least 90% and more preferably at least 98%.

[000810] *Encapsulate*: As used herein, the term “encapsulate” means to enclose, surround or encase.

[000811] *Encoded protein cleavage signal*: As used herein, “encoded protein cleavage signal” refers to the nucleotide sequence which encodes a protein cleavage signal.

[000812] *Engineered*: As used herein, embodiments of the invention are “engineered” when they are designed to have a feature or property, whether structural or chemical, that varies from a starting point, wild type or native molecule.

[000813] *Effective Amount*: As used herein, the term “effective amount” of an agent is that amount sufficient to effect beneficial or desired results, for example, clinical results, and, as such, an “effective amount” depends upon the context in which it is being applied. For example, in the context of administering an agent that treats cancer, an effective amount of an agent is, for example, an amount sufficient to achieve treatment, as defined herein, of cancer, as compared to the response obtained without administration of the agent.

[000814] *Exosome*: As used herein, “exosome” is a vesicle secreted by mammalian cells or a complex involved in RNA degradation.

[000815] *Expression*: As used herein, “expression” of a nucleic acid sequence refers to one or more of the following events: (1) production of an RNA template from a DNA sequence (e.g., by transcription); (2) processing of an RNA transcript (e.g., by splicing,

editing, 5' cap formation, and/or 3' end processing); (3) translation of an RNA into a polypeptide or protein; and (4) post-translational modification of a polypeptide or protein.

[000816] *Feature*: As used herein, a “feature” refers to a characteristic, a property, or a distinctive element.

[000817] *Formulation*: As used herein, a “formulation” includes at least a polynucleotide and a delivery agent.

[000818] *Fragment*: A “fragment,” as used herein, refers to a portion. For example, fragments of proteins may comprise polypeptides obtained by digesting full-length protein isolated from cultured cells.

[000819] *Functional*: As used herein, a “functional” biological molecule is a biological molecule in a form in which it exhibits a property and/or activity by which it is characterized.

[000820] *Homology*: As used herein, the term “homology” refers to the overall relatedness between polymeric molecules, *e.g.* between nucleic acid molecules (*e.g.* DNA molecules and/or RNA molecules) and/or between polypeptide molecules. In some embodiments, polymeric molecules are considered to be “homologous” to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% identical or similar. The term “homologous” necessarily refers to a comparison between at least two sequences (polynucleotide or polypeptide sequences). In accordance with the invention, two polynucleotide sequences are considered to be homologous if the polypeptides they encode are at least about 50%, 60%, 70%, 80%, 90%, 95%, or even 99% for at least one stretch of at least about 20 amino acids. In some embodiments, homologous polynucleotide sequences are characterized by the ability to encode a stretch of at least 4–5 uniquely specified amino acids. For polynucleotide sequences less than 60 nucleotides in length, homology is determined by the ability to encode a stretch of at least 4–5 uniquely specified amino acids. In accordance with the invention, two protein sequences are considered to be homologous if the proteins are at least about 50%, 60%, 70%, 80%, or 90% identical for at least one stretch of at least about 20 amino acids.

[000821] *Identity*: As used herein, the term “identity” refers to the overall relatedness between polymeric molecules, *e.g.*, between polynucleotide molecules (*e.g.* DNA

molecules and/or RNA molecules) and/or between polypeptide molecules. Calculation of the percent identity of two polynucleotide sequences, for example, can be performed by aligning the two sequences for optimal comparison purposes (*e.g.*, gaps can be introduced in one or both of a first and a second nucleic acid sequences for optimal alignment and non-identical sequences can be disregarded for comparison purposes). In certain embodiments, the length of a sequence aligned for comparison purposes is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or 100% of the length of the reference sequence. The nucleotides at corresponding nucleotide positions are then compared. When a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which needs to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. For example, the percent identity between two nucleotide sequences can be determined using methods such as those described in *Computational Molecular Biology*, Lesk, A. M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D. W., ed., Academic Press, New York, 1993; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; *Computer Analysis of Sequence Data, Part I*, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; each of which is incorporated herein by reference. For example, the percent identity between two nucleotide sequences can be determined using the algorithm of Meyers and Miller (CABIOS, 1989, 4:11-17), which has been incorporated into the ALIGN program (version 2.0) using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. The percent identity between two nucleotide sequences can, alternatively, be determined using the GAP program in the GCG software package using an NWSgapdna.CMP matrix. Methods commonly employed to determine percent identity between sequences include, but are not limited to those disclosed in Carillo, H., and Lipman, D., SIAM J Applied

Math., 48:1073 (1988); incorporated herein by reference. Techniques for determining identity are codified in publicly available computer programs. Exemplary computer software to determine homology between two sequences include, but are not limited to, GCG program package, Devereux, J., *et al.*, *Nucleic Acids Research*, 12(1), 387 (1984)), BLASTP, BLASTN, and FASTA Altschul, S. F. *et al.*, *J. Molec. Biol.*, 215, 403 (1990)).

[000822] *Infectious Agent*: As used herein, the phrase “infectious agent” means an agent capable of producing an infection.

[000823] *Inhibit expression of a gene*: As used herein, the phrase “inhibit expression of a gene” means to cause a reduction in the amount of an expression product of the gene. The expression product can be an RNA transcribed from the gene (*e.g.*, an mRNA) or a polypeptide translated from an mRNA transcribed from the gene. Typically a reduction in the level of an mRNA results in a reduction in the level of a polypeptide translated therefrom. The level of expression may be determined using standard techniques for measuring mRNA or protein.

[000824] *Infectious agent*: As used herein, an “infectious agent” refers to any microorganism, virus, infectious substance, or biological product that may be engineered as a result of biotechnology, or any naturally occurring or bioengineered component of any such microorganism, virus, infectious substance, or biological product, can cause emerging and contagious disease, death or other biological malfunction in a human, an animal, a plant or another living organism.

[000825] *Influenza*: As used herein, “influenza” or “flu” is an infectious disease of birds and mammals caused by RNA viruses of the family Orthomyxoviridae, the influenza viruses.

[000826] *Isomer*: As used herein, the term “isomer” means any tautomer, stereoisomer, enantiomer, or diastereomer of any compound of the invention. It is recognized that the compounds of the invention can have one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as double-bond isomers (*i.e.*, geometric E/Z isomers) or diastereomers (*e.g.*, enantiomers (*i.e.*, (+) or (-)) or cis/trans isomers). According to the invention, the chemical structures depicted herein, and therefore the compounds of the invention, encompass all of the corresponding stereoisomers, that is, both the stereomerically pure form (*e.g.*, geometrically pure, enantiomerically pure, or

diastereomerically pure) and enantiomeric and stereoisomeric mixtures, e.g., racemates. Enantiomeric and stereoisomeric mixtures of compounds of the invention can typically be resolved into their component enantiomers or stereoisomers by well-known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and stereoisomers can also be obtained from stereomerically or enantiomerically pure intermediates, reagents, and catalysts by well-known asymmetric synthetic methods.

[000827] *In vitro*: As used herein, the term “*in vitro*” refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, in a Petri dish, etc., rather than within an organism (e.g., animal, plant, or microbe).

[000828] *In vivo*: As used herein, the term “*in vivo*” refers to events that occur within an organism (e.g., animal, plant, or microbe or cell or tissue thereof).

[000829] *Isolated*: As used herein, the term “isolated” refers to a substance or entity that has been separated from at least some of the components with which it was associated (whether in nature or in an experimental setting). Isolated substances may have varying levels of purity in reference to the substances from which they have been associated. Isolated substances and/or entities may be separated from at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or more of the other components with which they were initially associated. In some embodiments, isolated agents are more than about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a substance is “pure” if it is substantially free of other components. *Substantially isolated*: By “substantially isolated” is meant that the compound is substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the compound of the present disclosure. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compound of the present disclosure, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

[000830] *IVT Polynucleotide*: As used herein, an “IVT polynucleotide” is a linear polynucleotide which may be made using only *in vitro* transcription (IVT) enzymatic synthesis methods.

[000831] *Linker*: As used herein, a “linker” refers to a group of atoms, e.g., 10-1,000 atoms, and can be comprised of the atoms or groups such as, but not limited to, carbon, amino, alkylamino, oxygen, sulfur, sulfoxide, sulfonyl, carbonyl, and imine. The linker can be attached to a modified nucleoside or nucleotide on the nucleobase or sugar moiety at a first end, and to a payload, e.g., a detectable or therapeutic agent, at a second end. The linker may be of sufficient length as to not interfere with incorporation into a nucleic acid sequence. The linker can be used for any useful purpose, such as to form polynucleotide multimers (e.g., through linkage of two or more chimeric polynucleotides molecules or IVT polynucleotides) or polynucleotides conjugates, as well as to administer a payload, as described herein. Examples of chemical groups that can be incorporated into the linker include, but are not limited to, alkyl, alkenyl, alkynyl, amido, amino, ether, thioether, ester, alkylene, heteroalkylene, aryl, or heterocyclyl, each of which can be optionally substituted, as described herein. Examples of linkers include, but are not limited to, unsaturated alkanes, polyethylene glycols (e.g., ethylene or propylene glycol monomeric units, e.g., diethylene glycol, dipropylene glycol, triethylene glycol, tripropylene glycol, tetraethylene glycol, or tetraethylene glycol), and dextran polymers and derivatives thereof. Other examples include, but are not limited to, cleavable moieties within the linker, such as, for example, a disulfide bond (-S-S-) or an azo bond (-N=N-), which can be cleaved using a reducing agent or photolysis. Non-limiting examples of a selectively cleavable bond include an amido bond can be cleaved for example by the use of tris(2-carboxyethyl)phosphine (TCEP), or other reducing agents, and/or photolysis, as well as an ester bond can be cleaved for example by acidic or basic hydrolysis.

[000832] *MicroRNA (miRNA) binding site*: As used herein, a microRNA (miRNA) binding site represents a nucleotide location or region of a nucleic acid transcript to which at least the “seed” region of a miRNA binds.

[000833] *Modified*: As used herein “modified” refers to a changed state or structure of a molecule of the invention. Molecules may be modified in many ways including

chemically, structurally, and functionally. In one embodiment, the mRNA molecules of the present invention are modified by the introduction of non-natural nucleosides and/or nucleotides, e.g., as it relates to the natural ribonucleotides A, U, G, and C. Noncanonical nucleotides such as the cap structures are not considered “modified” although they differ from the chemical structure of the A, C, G, U ribonucleotides.

[000834] *Mucus*: As used herein, “mucus” refers to the natural substance that is viscous and comprises mucin glycoproteins.

[000835] *Naturally occurring*: As used herein, “naturally occurring” means existing in nature without artificial aid.

[000836] *Neutralizing antibody*: As used herein, a “neutralizing antibody” refers to an antibody which binds to its antigen and defends a cell from an antigen or infectious agent by neutralizing or abolishing any biological activity it has.

[000837] *Non-human vertebrate*: As used herein, a “non-human vertebrate” includes all vertebrates except *Homo sapiens*, including wild and domesticated species. Examples of non-human vertebrates include, but are not limited to, mammals, such as alpaca, banteng, bison, camel, cat, cattle, deer, dog, donkey, gayal, goat, guinea pig, horse, llama, mule, pig, rabbit, reindeer, sheep water buffalo, and yak.

[000838] *Off-target*: As used herein, “off target” refers to any unintended effect on any one or more target, gene, or cellular transcript.

[000839] *Open reading frame*: As used herein, “open reading frame” or “ORF” refers to a sequence which does not contain a stop codon in a given reading frame.

[000840] *Operably linked*: As used herein, the phrase “operably linked” refers to a functional connection between two or more molecules, constructs, transcripts, entities, moieties or the like.

[000841] *Optionally substituted*: Herein a phrase of the form “optionally substituted X” (e.g., optionally substituted alkyl) is intended to be equivalent to “X, wherein X is optionally substituted” (e.g., “alkyl, wherein said alkyl is optionally substituted”). It is not intended to mean that the feature “X” (e.g. alkyl) *per se* is optional.

[000842] *Part*: As used herein, a “part” or “region” of a polynucleotide is defined as any portion of the polynucleotide which is less than the entire length of the polynucleotide.

[000843] *Peptide*: As used herein, “peptide” is less than or equal to 50 amino acids long, e.g., about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 amino acids long.

[000844] *Paratope*: As used herein, a “paratope” refers to the antigen-binding site of an antibody.

[000845] *Patient*: As used herein, “patient” refers to a subject who may seek or be in need of treatment, requires treatment, is receiving treatment, will receive treatment, or a subject who is under care by a trained professional for a particular disease or condition.

[000846] *Pharmaceutically acceptable*: The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[000847] *Pharmaceutically acceptable excipients*: The phrase “pharmaceutically acceptable excipient,” as used herein, refers any ingredient other than the compounds described herein (for example, a vehicle capable of suspending or dissolving the active compound) and having the properties of being substantially nontoxic and non-inflammatory in a patient. Excipients may include, for example: antiadherents, antioxidants, binders, coatings, compression aids, disintegrants, dyes (colors), emollients, emulsifiers, fillers (diluent), film formers or coatings, flavors, fragrances, glidants (flow enhancers), lubricants, preservatives, printing inks, sorbents, suspending or dispersing agents, sweeteners, and waters of hydration. Exemplary excipients include, but are not limited to: butylated hydroxytoluene (BHT), calcium carbonate, calcium phosphate (dibasic), calcium stearate, croscarmellose, crosslinked polyvinyl pyrrolidone, citric acid, crospovidone, cysteine, ethylcellulose, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, maltitol, mannitol, methionine, methylcellulose, methyl paraben, microcrystalline cellulose, polyethylene glycol, polyvinyl pyrrolidone, povidone, pregelatinized starch, propyl paraben, retinyl palmitate, shellac, silicon dioxide, sodium carboxymethyl cellulose, sodium citrate, sodium starch glycolate, sorbitol, starch (corn), stearic acid, sucrose, talc, titanium dioxide, vitamin A, vitamin E, vitamin C, and xylitol.

[000848] *Pharmaceutically acceptable salts*: The present disclosure also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, “pharmaceutically acceptable salts” refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form (e.g., by reacting the free base group with a suitable organic acid). Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. Representative acid addition salts include acetate, acetic acid, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzenesulfonic acid, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. The pharmaceutically acceptable salts of the present disclosure include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present disclosure can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company,

Easton, Pa., 1985, p. 1418, *Pharmaceutical Salts: Properties, Selection, and Use*, P.H. Stahl and C.G. Wermuth (eds.), Wiley-VCH, 2008, and Berge et al., *Journal of Pharmaceutical Science*, 66, 1-19 (1977), each of which is incorporated herein by reference in its entirety.

[000849] *Pharmaceutically acceptable solvate*: The term “pharmaceutically acceptable solvate,” as used herein, means a compound of the invention wherein molecules of a suitable solvent are incorporated in the crystal lattice. A suitable solvent is physiologically tolerable at the dosage administered. For example, solvates may be prepared by crystallization, recrystallization, or precipitation from a solution that includes organic solvents, water, or a mixture thereof. Examples of suitable solvents are ethanol, water (for example, mono-, di-, and tri-hydrates), *N*-methylpyrrolidinone (NMP), dimethyl sulfoxide (DMSO), *N,N'*-dimethylformamide (DMF), *N,N'*-dimethylacetamide (DMAC), 1,3-dimethyl-2-imidazolidinone (DMEU), 1,3-dimethyl-3,4,5,6-tetrahydro-2-(1H)-pyrimidinone (DMPU), acetonitrile (ACN), propylene glycol, ethyl acetate, benzyl alcohol, 2-pyrrolidone, benzyl benzoate, and the like. When water is the solvent, the solvate is referred to as a “hydrate.”

[000850] *Pharmacokinetic*: As used herein, “pharmacokinetic” refers to any one or more properties of a molecule or compound as it relates to the determination of the fate of substances administered to a living organism. Pharmacokinetics is divided into several areas including the extent and rate of absorption, distribution, metabolism and excretion. This is commonly referred to as ADME where: (A) Absorption is the process of a substance entering the blood circulation; (D) Distribution is the dispersion or dissemination of substances throughout the fluids and tissues of the body; (M) Metabolism (or Biotransformation) is the irreversible transformation of parent compounds into daughter metabolites; and (E) Excretion (or Elimination) refers to the elimination of the substances from the body. In rare cases, some drugs irreversibly accumulate in body tissue.

[000851] *Physicochemical*: As used herein, “physicochemical” means of or relating to a physical and/or chemical property.

[000852] *Polypeptide per unit drug (PUD)*: As used herein, a PUD or product per unit drug, is defined as a subdivided portion of total daily dose, usually 1 mg, pg, kg, etc., of a

product (such as a polypeptide) as measured in body fluid or tissue, usually defined in concentration such as pmol/mL, mmol/mL, etc. divided by the measure in the body fluid.

[000853] *Preventing*: As used herein, the term “preventing” refers to partially or completely delaying onset of an infection, disease, disorder and/or condition; partially or completely delaying onset of one or more symptoms, features, or clinical manifestations of a particular infection, disease, disorder, and/or condition; partially or completely delaying onset of one or more symptoms, features, or manifestations of a particular infection, disease, disorder, and/or condition; partially or completely delaying progression from an infection, a particular disease, disorder and/or condition; and/or decreasing the risk of developing pathology associated with the infection, the disease, disorder, and/or condition.

[000854] *Prodrug*: The present disclosure also includes prodrugs of the compounds described herein. As used herein, “prodrugs” refer to any substance, molecule or entity which is in a form predicate for that substance, molecule or entity to act as a therapeutic upon chemical or physical alteration. Prodrugs may be covalently bonded or sequestered in some way and which release or are converted into the active drug moiety prior to, upon or after administered to a mammalian subject. Prodrugs can be prepared by modifying functional groups present in the compounds in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compounds. Prodrugs include compounds wherein hydroxyl, amino, sulfhydryl, or carboxyl groups are bonded to any group that, when administered to a mammalian subject, cleaves to form a free hydroxyl, amino, sulfhydryl, or carboxyl group respectively. Preparation and use of prodrugs is discussed in T. Higuchi and V. Stella, “Pro-drugs as Novel Delivery Systems,” Vol. 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are hereby incorporated by reference in their entirety.

[000855] *Proliferate*: As used herein, the term “proliferate” means to grow, expand or increase or cause to grow, expand or increase rapidly. “Proliferative” means having the ability to proliferate. “Anti-proliferative” means having properties counter to or inapposite to proliferative properties.

[000856] *Progenitor cell*: As used herein, the term “progenitor cell” refers to cells that have greater developmental potential relative to a cell which it can give rise to by differentiation.

[000857] *Prophylactic*: As used herein, “prophylactic” refers to a therapeutic or course of action used to prevent the spread of disease.

[000858] *Prophylaxis*: As used herein, a “prophylaxis” refers to a measure taken to maintain health and prevent the spread of disease. An “immune prophylaxis” refers to a measure to produce active or passive immunity to prevent the spread of disease.

[000859] *Protein cleavage site*: As used herein, “protein cleavage site” refers to a site where controlled cleavage of the amino acid chain can be accomplished by chemical, enzymatic or photochemical means.

[000860] *Protein cleavage signal*: As used herein “protein cleavage signal” refers to at least one amino acid that flags or marks a polypeptide for cleavage.

[000861] *Protein of interest*: As used herein, the terms “proteins of interest” or “desired proteins” include those provided herein and fragments, mutants, variants, and alterations thereof.

[000862] *Proximal*: As used herein, the term “proximal” means situated nearer to the center or to a point or region of interest.

[000863] *Pseudouridine*: As used herein, pseudouridine refers to the C-glycoside isomer of the nucleoside uridine. A “pseudouridine analog” is any modification, variant, isoform or derivative of pseudouridine. For example, pseudouridine analogs include but are not limited to 1-carboxymethyl-pseudouridine, 1-propynyl-pseudouridine, 1-taurinomethyl-pseudouridine, 1-taurinomethyl-4-thio-pseudouridine, 1-methylpseudouridine ($m^1\psi$), 1-methyl-4-thio-pseudouridine ($m^1s^4\psi$), 4-thio-1-methyl-pseudouridine, 3-methyl-pseudouridine ($m^3\psi$), 2-thio-1-methyl-pseudouridine, 1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-1-deaza-pseudouridine, dihydropseudouridine, 2-thio-dihydropseudouridine, 2-methoxyuridine, 2-methoxy-4-thio-uridine, 4-methoxy-pseudouridine, 4-methoxy-2-thio-pseudouridine, N1-methyl-pseudouridine, 1-methyl-3-(3-amino-3-carboxypropyl)pseudouridine ($acp^3\psi$), and 2'-O-methyl-pseudouridine (ψm).

[000864] *Purified*: As used herein, “purify,” “purified,” “purification” means to make substantially pure or clear from unwanted components, material defilement, admixture or imperfection.

[000865] *Repeated transfection*: As used herein, the term “repeated transfection” refers to transfection of the same cell culture with a polynucleotide a plurality of times. The cell culture can be transfected at least twice, at least 3 times, at least 4 times, at least 5 times, at least 6 times, at least 7 times, at least 8 times, at least 9 times, at least 10 times, at least 11 times, at least 12 times, at least 13 times, at least 14 times, at least 15 times, at least 16 times, at least 17 times at least 18 times, at least 19 times, at least 20 times, at least 25 times, at least 30 times, at least 35 times, at least 40 times, at least 45 times, at least 50 times or more.

[000866] *Sample*: As used herein, the term “sample” or “biological sample” refers to a subset of its tissues, cells or component parts (e.g. body fluids, including but not limited to blood, mucus, lymphatic fluid, synovial fluid, cerebrospinal fluid, saliva, amniotic fluid, amniotic cord blood, urine, vaginal fluid and semen). A sample further may include a homogenate, lysate or extract prepared from a whole organism or a subset of its tissues, cells or component parts, or a fraction or portion thereof, including but not limited to, for example, plasma, serum, spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, blood cells, tumors, organs. A sample further refers to a medium, such as a nutrient broth or gel, which may contain cellular components, such as proteins or nucleic acid molecule.

[000867] *Signal Sequences*: As used herein, the phrase “signal sequences” refers to a sequence which can direct the transport or localization of a protein.

[000868] *Single unit dose*: As used herein, a “single unit dose” is a dose of any therapeutic administered in one dose/at one time/single route/single point of contact, i.e., single administration event.

[000869] *Similarity*: As used herein, the term “similarity” refers to the overall relatedness between polymeric molecules, e.g. between polynucleotide molecules (e.g. DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Calculation of percent similarity of polymeric molecules to one another can be performed

in the same manner as a calculation of percent identity, except that calculation of percent similarity takes into account conservative substitutions as is understood in the art.

[000870] *Split dose*: As used herein, a “split dose” is the division of single unit dose or total daily dose into two or more doses.

[000871] *Stable*: As used herein “stable” refers to a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and preferably capable of formulation into an efficacious therapeutic agent.

[000872] *Stabilized*: As used herein, the term “stabilize”, “stabilized,” “stabilized region” means to make or become stable.

[000873] *Stereoisomer*: As used herein, the term “stereoisomer” refers to all possible different isomeric as well as conformational forms which a compound may possess (e.g., a compound of any formula described herein), in particular all possible stereochemically and conformationally isomeric forms, all diastereomers, enantiomers and/or conformers of the basic molecular structure. Some compounds of the present invention may exist in different tautomeric forms, all of the latter being included within the scope of the present invention.

[000874] *Subject*: As used herein, the term “subject” or “patient” refers to any organism to which a composition in accordance with the invention may be administered, e.g., for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include animals (e.g., mammals such as mice, rats, rabbits, non-human primates, and humans) and/or plants.

[000875] *Substantially*: As used herein, the term “substantially” refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of interest. One of ordinary skill in the biological arts will understand that biological and chemical phenomena rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term “substantially” is therefore used herein to capture the potential lack of completeness inherent in many biological and chemical phenomena.

[000876] *Substantially equal*: As used herein as it relates to time differences between doses, the term means plus/minus 2%.

[000877] *Substantially simultaneously*: As used herein and as it relates to plurality of doses, the term means within 2 seconds.

[000878] *Suffering from*: An individual who is “suffering from” a disease, disorder, and/or condition has been diagnosed with or displays one or more symptoms of a disease, disorder, and/or condition.

[000879] *Susceptible to*: An individual who is “susceptible to” a disease, disorder, and/or condition has not been diagnosed with and/or may not exhibit symptoms of the disease, disorder, and/or condition but harbors a propensity to develop a disease or its symptoms. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition (for example, cancer) may be characterized by one or more of the following: (1) a genetic mutation associated with development of the disease, disorder, and/or condition; (2) a genetic polymorphism associated with development of the disease, disorder, and/or condition; (3) increased and/or decreased expression and/or activity of a protein and/or nucleic acid associated with the disease, disorder, and/or condition; (4) habits and/or lifestyles associated with development of the disease, disorder, and/or condition; (5) a family history of the disease, disorder, and/or condition; and (6) exposure to and/or infection with a microbe associated with development of the disease, disorder, and/or condition. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition will develop the disease, disorder, and/or condition. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition will not develop the disease, disorder, and/or condition.

[000880] *Sustained release*: As used herein, the term “sustained release” refers to a pharmaceutical composition or compound release profile that conforms to a release rate over a specific period of time.

[000881] *Synthetic*: The term “synthetic” means produced, prepared, and/or manufactured by the hand of man. Synthesis of polynucleotides or polypeptides or other molecules of the present invention may be chemical or enzymatic.

[000882] *Targeted Cells*: As used herein, “targeted cells” refers to any one or more cells of interest. The cells may be found *in vitro*, *in vivo*, *in situ* or in the tissue or organ of an organism. The organism may be an animal, preferably a mammal, more preferably a human and most preferably a patient.

[000883] *Therapeutic Agent*: The term “therapeutic agent” refers to any agent that, when administered to a subject, has a therapeutic, diagnostic, and/or prophylactic effect and/or elicits a desired biological and/or pharmacological effect.

[000884] *Therapeutically effective amount*: As used herein, the term “therapeutically effective amount” means an amount of an agent to be delivered (*e.g.*, nucleic acid, drug, therapeutic agent, diagnostic agent, prophylactic agent, *etc.*) that is sufficient, when administered to a subject suffering from or susceptible to an infection, disease, disorder, and/or condition, to treat, improve symptoms of, diagnose, prevent, and/or delay the onset of the infection, disease, disorder, and/or condition.

[000885] *Therapeutically effective outcome*: As used herein, the term “therapeutically effective outcome” means an outcome that is sufficient in a subject suffering from or susceptible to an infection, disease, disorder, and/or condition, to treat, improve symptoms of, diagnose, prevent, and/or delay the onset of the infection, disease, disorder, and/or condition.

[000886] *Total daily dose*: As used herein, a “total daily dose” is an amount given or prescribed in 24 hr period. It may be administered as a single unit dose.

[000887] *Totipotency*: As used herein, “totipotency” refers to a cell with a developmental potential to make all of the cells found in the adult body as well as the extra-embryonic tissues, including the placenta.

[000888] *Transcription factor*: As used herein, the term “transcription factor” refers to a DNA-binding protein that regulates transcription of DNA into RNA, for example, by activation or repression of transcription. Some transcription factors effect regulation of transcription alone, while others act in concert with other proteins. Some transcription factor can both activate and repress transcription under certain conditions. In general, transcription factors bind a specific target sequence or sequences highly similar to a specific consensus sequence in a regulatory region of a target gene. Transcription factors may regulate transcription of a target gene alone or in a complex with other molecules.

[000889] *Transcription*: As used herein, the term “transcription” refers to methods to introduce exogenous nucleic acids into a cell. Methods of transfection include, but are not limited to, chemical methods, physical treatments and cationic lipids or mixtures.

[000890] *Transdifferentiation*: As used herein, “transdifferentiation” refers to the capacity of differentiated cells of one type to lose identifying characteristics and to change their phenotype to that of other fully differentiated cells.

[000891] *Treating*: As used herein, the term “treating” refers to partially or completely alleviating, ameliorating, improving, relieving, delaying onset of, inhibiting progression of, reducing severity of, and/or reducing incidence of one or more symptoms or features of a particular infection, disease, disorder, and/or condition. For example, “treating” cancer may refer to inhibiting survival, growth, and/or spread of a tumor. Treatment may be administered to a subject who does not exhibit signs of a disease, disorder, and/or condition and/or to a subject who exhibits only early signs of a disease, disorder, and/or condition for the purpose of decreasing the risk of developing pathology associated with the disease, disorder, and/or condition.

[000892] *Unmodified*: As used herein, “unmodified” refers to any substance, compound or molecule prior to being changed in any way. Unmodified may, but does not always, refer to the wild type or native form of a biomolecule. Molecules may undergo a series of modifications whereby each modified molecule may serve as the “unmodified” starting molecule for a subsequent modification.

[000893] *Unipotent*: As used herein, “unipotent” when referring to a cell means to give rise to a single cell lineage.

[000894] *Vaccine*: As used herein, the phrase “vaccine” refers to a biological preparation that improves immunity to a particular disease.

[000895] *Viral protein*: As used herein, the phrase “viral protein” means any protein originating from a virus.

Equivalents and Scope

[000896] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments in accordance with the invention described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the appended claims.

[000897] In the claims, articles such as “a,” “an,” and “the” may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or

descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[000898] It is also noted that the term “comprising” is intended to be open and permits but does not require the inclusion of additional elements or steps. When the term “comprising” is used herein, the term “consisting of” is thus also encompassed and disclosed.

[000899] Where ranges are given, endpoints are included. Furthermore, it is to be understood that unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or subrange within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

[000900] In addition, it is to be understood that any particular embodiment of the present invention that falls within the prior art may be explicitly excluded from any one or more of the claims. Since such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the compositions of the invention (*e.g.*, any nucleic acid or protein encoded thereby; any method of production; any method of use; *etc.*) can be excluded from any one or more claims, for any reason, whether or not related to the existence of prior art.

[000901] All cited sources, for example, references, publications, databases, database entries, and art cited herein, are incorporated into this application by reference, even if not expressly stated in the citation. In case of conflicting statements of a cited source and the instant application, the statement in the instant application shall control.

[000902] Section and table headings are not intended to be limiting.

EXAMPLES**Example 1. Manufacture of Polynucleotides**

[000903] According to the present invention, the manufacture of polynucleotides and or parts or regions thereof may be accomplished utilizing the methods taught in International Patent Publication No. WO2014152027 (Attorney Docket number M500), the contents of which is incorporated herein by reference in its entirety.

[000904] Purification methods may include those taught in International Patent Publication No. WO2014152030 (Attorney Docket number M501); International Patent Publication No. WO2014152031 (Attorney Docket number M502); each of which is incorporated herein by reference in its entirety.

[000905] Detection and characterization methods of the polynucleotides may be performed as taught in International Patent Publication No. WO2014144039 (Attorney Docket number M505), each of which is incorporated herein by reference in its entirety.

[000906] Characterization of the polynucleotides of the invention may be accomplished using a procedure selected from the group consisting of polynucleotide mapping, reverse transcriptase sequencing, charge distribution analysis, and detection of RNA impurities, wherein characterizing comprises determining the RNA transcript sequence, determining the purity of the RNA transcript, or determining the charge heterogeneity of the RNA transcript. Such methods are taught in, for example, International Patent Publication No. WO2014144711 (Attorney Docket number M506) and International Patent Publication No. WO2014144767 (Attorney Docket number M507) the contents of each of which is incorporated herein by reference in its entirety.

Example 2. Chimeric polynucleotide synthesis: triphosphate route**Introduction**

[000907] According to the present invention, two regions or parts of a chimeric polynucleotide may be joined or ligated using triphosphate chemistry.

[000908] According to this method, a first region or part of 100 nucleotides or less is chemically synthesized with a 5' monophosphate and terminal 3'desOH or blocked OH. If the region is longer than 80 nucleotides, it may be synthesized as two strands for ligation.

[000909] If the first region or part is synthesized as a non-positionally modified region or part using in vitro transcription (IVT), conversion the 5' monophosphate with subsequent capping of the 3' terminus may follow.

[000910] Monophosphate protecting groups may be selected from any of those known in the art.

[000911] The second region or part of the chimeric polynucleotide may be synthesized using either chemical synthesis or IVT methods. IVT methods may include an RNA polymerase that can utilize a primer with a modified cap. Alternatively, a cap of up to 80 nucleotides may be chemically synthesized and coupled to the IVT region or part.

[000912] It is noted that for ligation methods, ligation with DNA T4 ligase, followed by treatment with DNase should readily avoid concatenation.

[000913] The entire chimeric polynucleotide need not be manufactured with a phosphate-sugar backbone. If one of the regions or parts encodes a polypeptide, then it is preferable that such region or part comprise a phosphate-sugar backbone.

[000914] Ligation is then performed using any known click chemistry, orthoclick chemistry, solulink, or other bioconjugate chemistries known to those in the art.

Synthetic route

[000915] The chimeric polynucleotide is made using a series of starting segments. Such segments include:

[000916] (a) Capped and protected 5' segment comprising a normal 3'OH (SEG. 1)

[000917] (b) 5' triphosphate segment which may include the coding region of a polypeptide and comprising a normal 3'OH (SEG. 2)

[000918] (c) 5' monophosphate segment for the 3' end of the chimeric polynucleotide (e.g., the tail) comprising cordycepin or no 3'OH (SEG. 3)

[000919] After synthesis (chemical or IVT), segment 3 (SEG. 3) is treated with cordycepin and then with pyrophosphatase to create the 5' monophosphate.

[000920] Segment 2 (SEG. 2) is then ligated to SEG. 3 using RNA ligase. The ligated polynucleotide is then purified and treated with pyrophosphatase to cleave the diphosphate. The treated SEG.2-SEG. 3 construct is then purified and SEG. 1 is ligated to the 5' terminus. A further purification step of the chimeric polynucleotide may be performed.

[000921] Where the chimeric polynucleotide encodes a polypeptide, the ligated or joined segments may be represented as: 5'UTR (SEG. 1), open reading frame or ORF (SEG. 2) and 3'UTR+PolyA (SEG. 3).

[000922] The yields of each step may be as much as 90-95%.

Example 3: PCR for cDNA Production

[000923] PCR procedures for the preparation of cDNA are performed using 2x KAPA HIFI™ HotStart ReadyMix by Kapa Biosystems (Woburn, MA). This system includes 2x KAPA ReadyMix 12.5 µl; Forward Primer (10 uM) 0.75 µl; Reverse Primer (10 uM) 0.75 µl; Template cDNA -100 ng; and dH₂O diluted to 25.0 µl. The reaction conditions are at 95° C for 5 min. and 25 cycles of 98° C for 20 sec, then 58° C for 15 sec, then 72° C for 45 sec, then 72° C for 5 min. then 4° C to termination.

[000924] The reverse primer of the instant invention incorporates a poly-T₁₂₀ for a poly-A₁₂₀ in the mRNA. Other reverse primers with longer or shorter poly(T) tracts can be used to adjust the length of the poly(A) tail in the polynucleotide mRNA.

[000925] The reaction is cleaned up using Invitrogen's PURELINK™ PCR Micro Kit (Carlsbad, CA) per manufacturer's instructions (up to 5 µg). Larger reactions will require a cleanup using a product with a larger capacity. Following the cleanup, the cDNA is quantified using the NANODROP™ and analyzed by agarose gel electrophoresis to confirm the cDNA is the expected size. The cDNA is then submitted for sequencing analysis before proceeding to the *in vitro* transcription reaction.

Example 4. *In vitro* Transcription (IVT)

[000926] The *in vitro* transcription reaction generates polynucleotides containing uniformly modified polynucleotides. Such uniformly modified polynucleotides may comprise a region or part of the polynucleotides of the invention. The input nucleotide triphosphate (NTP) mix is made in-house using natural and un-natural NTPs.

[000927] A typical *in vitro* transcription reaction includes the following:

- | | | |
|---|---|--------|
| 1 | Template cDNA | 1.0 µg |
| 2 | 10x transcription buffer (400 mM Tris-HCl pH 8.0, 190 mM MgCl ₂ , 50 mM DTT, 10 mM Spermidine) | 2.0 µl |
| 3 | Custom NTPs (25mM each) | 7.2 µl |
| 4 | RNase Inhibitor | 20 U |

- 5 T7 RNA polymerase 3000 U
- 6 dH₂O Up to 20.0 µl. and
- 7 Incubation at 37° C for 3 hr-5 hrs.

[000928] The crude IVT mix may be stored at 4° C overnight for cleanup the next day. 1 U of RNase-free DNase is then used to digest the original template. After 15 minutes of incubation at 37° C, the mRNA is purified using Ambion's MEGACLEAR™ Kit (Austin, TX) following the manufacturer's instructions. This kit can purify up to 500 µg of RNA. Following the cleanup, the RNA is quantified using the NanoDrop and analyzed by agarose gel electrophoresis to confirm the RNA is the proper size and that no degradation of the RNA has occurred.

5. Enzymatic Capping

[000929] Capping of a polynucleotide is performed as follows where the mixture includes: IVT RNA 60 µg-180µg and dH₂O up to 72 µl. The mixture is incubated at 65° C for 5 minutes to denature RNA, and then is transferred immediately to ice.

[000930] The protocol then involves the mixing of 10x Capping Buffer (0.5 M Tris-HCl (pH 8.0), 60 mM KCl, 12.5 mM MgCl₂) (10.0 µl); 20 mM GTP (5.0 µl); 20 mM S-Adenosyl Methionine (2.5 µl); RNase Inhibitor (100 U); 2'-O-Methyltransferase (400U); Vaccinia capping enzyme (Guanylyl transferase) (40 U); dH₂O (Up to 28 µl); and incubation at 37° C for 30 minutes for 60 µg RNA or up to 2 hours for 180 µg of RNA.

[000931] The polynucleotide is then purified using Ambion's MEGACLEAR™ Kit (Austin, TX) following the manufacturer's instructions. Following the cleanup, the RNA is quantified using the NANODROP™ (ThermoFisher, Waltham, MA) and analyzed by agarose gel electrophoresis to confirm the RNA is the proper size and that no degradation of the RNA has occurred. The RNA product may also be sequenced by running a reverse-transcription-PCR to generate the cDNA for sequencing.

Example 6. PolyA Tailing Reaction

[000932] Without a poly-T in the cDNA, a poly-A tailing reaction must be performed before cleaning the final product. This is done by mixing Capped IVT RNA (100 µl); RNase Inhibitor (20 U); 10x Tailing Buffer (0.5 M Tris-HCl (pH 8.0), 2.5 M NaCl, 100 mM MgCl₂)(12.0 µl); 20 mM ATP (6.0 µl); Poly-A Polymerase (20 U); dH₂O up to 123.5 µl and incubation at 37° C for 30 min. If the poly-A tail is already in the transcript, then

the tailing reaction may be skipped and proceed directly to cleanup with Ambion's MEGACLEAR™ kit (Austin, TX) (up to 500 µg). Poly-A Polymerase is preferably a recombinant enzyme expressed in yeast.

[000933] It should be understood that the processivity or integrity of the polyA tailing reaction may not always result in an exact size polyA tail. Hence polyA tails of approximately between 40-200 nucleotides, e.g., about 40, 50, 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 150-165, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164 or 165 are within the scope of the invention.

Example 7. Natural 5' Caps and 5' Cap Analogues

[000934] 5'-capping of polynucleotides may be completed concomitantly during the *in vitro*-transcription reaction using the following chemical RNA cap analogs to generate the 5'-guanosine cap structure according to manufacturer protocols: 3'-O-Me-m7G(5')ppp(5') G [the ARCA cap]; G(5')ppp(5')A; G(5')ppp(5')G; m7G(5')ppp(5')A; m7G(5')ppp(5')G (New England BioLabs, Ipswich, MA). 5'-capping of modified RNA may be completed post-transcriptionally using a Vaccinia Virus Capping Enzyme to generate the "Cap 0" structure: m7G(5')ppp(5')G (New England BioLabs, Ipswich, MA). Cap 1 structure may be generated using both Vaccinia Virus Capping Enzyme and a 2'-O methyl-transferase to generate: m7G(5')ppp(5')G-2'-O-methyl. Cap 2 structure may be generated from the Cap 1 structure followed by the 2'-O-methylation of the 5'-antepenultimate nucleotide using a 2'-O methyl-transferase. Cap 3 structure may be generated from the Cap 2 structure followed by the 2'-O-methylation of the 5'-preantepenultimate nucleotide using a 2'-O methyl-transferase. Enzymes are preferably derived from a recombinant source.

[000935] When transfected into mammalian cells, the modified mRNAs have a stability of between 12-18 hours or more than 18 hours, e.g., 24, 36, 48, 60, 72 or greater than 72 hours.

Example 8. Capping Assays

A. Protein Expression Assay

[000936] Polynucleotides encoding a polypeptide, containing any of the caps taught herein can be transfected into cells at equal concentrations. 6, 12, 24 and 36 hours post-

transfection the amount of protein secreted into the culture medium can be assayed by ELISA. Synthetic polynucleotides that secrete higher levels of protein into the medium would correspond to a synthetic polynucleotide with a higher translationally-competent Cap structure.

B. Purity Analysis Synthesis

[000937] Polynucleotides encoding a polypeptide, containing any of the caps taught herein can be compared for purity using denaturing Agarose-Urea gel electrophoresis or HPLC analysis. Polynucleotides with a single, consolidated band by electrophoresis correspond to the higher purity product compared to polynucleotides with multiple bands or streaking bands. Synthetic polynucleotides with a single HPLC peak would also correspond to a higher purity product. The capping reaction with a higher efficiency would provide a more pure polynucleotide population.

C. Cytokine Analysis

[000938] Polynucleotides encoding a polypeptide, containing any of the caps taught herein can be transfected into cells at multiple concentrations. 6, 12, 24 and 36 hours post-transfection the amount of pro-inflammatory cytokines such as TNF-alpha and IFN-beta secreted into the culture medium can be assayed by ELISA. Polynucleotides resulting in the secretion of higher levels of pro-inflammatory cytokines into the medium would correspond to a polynucleotides containing an immune-activating cap structure.

D. Capping Reaction Efficiency

[000939] Polynucleotides encoding a polypeptide, containing any of the caps taught herein can be analyzed for capping reaction efficiency by LC-MS after nuclease treatment. Nuclease treatment of capped polynucleotides would yield a mixture of free nucleotides and the capped 5'-5-triphosphate cap structure detectable by LC-MS. The amount of capped product on the LC-MS spectra can be expressed as a percent of total polynucleotide from the reaction and would correspond to capping reaction efficiency. The cap structure with higher capping reaction efficiency would have a higher amount of capped product by LC-MS.

Example 9. Agarose Gel Electrophoresis of Modified RNA or RT PCR Products

[000940] Individual polynucleotides (200-400 ng in a 20 µl volume) or reverse transcribed PCR products (200-400 ng) are loaded into a well on a non-denaturing 1.2%

Agarose E-Gel (Invitrogen, Carlsbad, CA) and run for 12-15 minutes according to the manufacturer protocol.

Example 10. Nanodrop Modified RNA Quantification and UV Spectral Data

[000941] Modified polynucleotides in TE buffer (1 μ l) are used for Nanodrop UV absorbance readings to quantitate the yield of each polynucleotide from a chemical synthesis or *in vitro* transcription reaction.

Example 11. Formulation of Modified mRNA Using Lipidoids

[000942] Polynucleotides are formulated for *in vitro* experiments by mixing the polynucleotides with the lipidoid at a set ratio prior to addition to cells. *In vivo* formulation may require the addition of extra ingredients to facilitate circulation throughout the body. To test the ability of these lipidoids to form particles suitable for *in vivo* work, a standard formulation process used for siRNA-lipidoid formulations may be used as a starting point. After formation of the particle, polynucleotide is added and allowed to integrate with the complex. The encapsulation efficiency is determined using a standard dye exclusion assays.

Example 12. Method of Screening for Protein Expression

A. Electrospray Ionization

[000943] A biological sample which may contain proteins encoded by a polynucleotide administered to the subject is prepared and analyzed according to the manufacturer protocol for electrospray ionization (ESI) using 1, 2, 3 or 4 mass analyzers. A biologic sample may also be analyzed using a tandem ESI mass spectrometry system.

Patterns of protein fragments, or whole proteins, are compared to known controls for a given protein and identity is determined by comparison.

B. Matrix-Assisted Laser Desorption/Ionization

[000944] A biological sample which may contain proteins encoded by one or more polynucleotides administered to the subject is prepared and analyzed according to the manufacturer protocol for matrix-assisted laser desorption/ionization (MALDI).

[000945] Patterns of protein fragments, or whole proteins, are compared to known controls for a given protein and identity is determined by comparison.

C. Liquid Chromatography-Mass spectrometry-Mass spectrometry

[000946] A biological sample, which may contain proteins encoded by one or more polynucleotides, may be treated with a trypsin enzyme to digest the proteins contained

within. The resulting peptides are analyzed by liquid chromatography-mass spectrometry-mass spectrometry (LC/MS/MS). The peptides are fragmented in the mass spectrometer to yield diagnostic patterns that can be matched to protein sequence databases via computer algorithms. The digested sample may be diluted to achieve 1 ng or less starting material for a given protein. Biological samples containing a simple buffer background (e.g. water or volatile salts) are amenable to direct in-solution digest; more complex backgrounds (e.g. detergent, non-volatile salts, glycerol) require an additional clean-up step to facilitate the sample analysis.

[000947] Patterns of protein fragments, or whole proteins, are compared to known controls for a given protein and identity is determined by comparison.

Example 13. Cyclization and/or concatemerization

[000948] According to the present invention, a polynucleotide may be cyclized, or concatemerized, to generate a translation competent molecule to assist interactions between poly-A binding proteins and 5'-end binding proteins. The mechanism of cyclization or concatemerization may occur through at least 3 different routes: 1) chemical, 2) enzymatic, and 3) ribozyme catalyzed. The newly formed 5'-/3'-linkage may be intramolecular or intermolecular.

[000949] In the first route, the 5'-end and the 3'-end of the nucleic acid contain chemically reactive groups that, when close together, form a new covalent linkage between the 5'-end and the 3'-end of the molecule. The 5'-end may contain an NHS-ester reactive group and the 3'-end may contain a 3'-amino-terminated nucleotide such that in an organic solvent the 3'-amino-terminated nucleotide on the 3'-end of a synthetic mRNA molecule will undergo a nucleophilic attack on the 5'-NHS-ester moiety forming a new 5'-/3'-amide bond.

[000950] In the second route, T4 RNA ligase may be used to enzymatically link a 5'-phosphorylated nucleic acid molecule to the 3'-hydroxyl group of a nucleic acid forming a new phosphodiester linkage. In an example reaction, 1 µg of a nucleic acid molecule is incubated at 37°C for 1 hour with 1-10 units of T4 RNA ligase (New England Biolabs, Ipswich, MA) according to the manufacturer's protocol. The ligation reaction may occur in the presence of a split polynucleotide capable of base-pairing with both the 5'- and 3'-region in juxtaposition to assist the enzymatic ligation reaction.

[000951] In the third route, either the 5'-or 3'-end of the cDNA template encodes a ligase ribozyme sequence such that during *in vitro* transcription, the resultant nucleic acid molecule can contain an active ribozyme sequence capable of ligating the 5'-end of a nucleic acid molecule to the 3'-end of a nucleic acid molecule. The ligase ribozyme may be derived from the Group I Intron, Group I Intron, Hepatitis Delta Virus, Hairpin ribozyme or may be selected by SELEX (systematic evolution of ligands by exponential enrichment). The ribozyme ligase reaction may take 1 to 24 hours at temperatures between 0 and 37°C. It is to be understood that the words which have been used are words of description rather than limitation, and that changes may be made within the purview of the appended claims without departing from the true scope and spirit of the invention in its broader aspects.

[000952] While the present invention has been described at some length and with some particularity with respect to the several described embodiments, it is not intended that it should be limited to any such particulars or embodiments or any particular embodiment, but it is to be construed with references to the appended claims so as to provide the broadest possible interpretation of such claims in view of the prior art and, therefore, to effectively encompass the intended scope of the invention.

[000953] All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, section headings, the materials, methods, and examples are illustrative only and not intended to be limiting.

Claims

1. A method of inducing tolerance in a cell system to an antigen comprising contacting said cellular system with phosphatidylserine (PS) and a tolerogenic composition comprising the antigen, one or more polynucleotides.
2. A method of inducing Treg cell activity in a cellular system comprising contacting said system with phosphatidylserine (PS) and a tolerogenic composition comprising an antigen, one or more polynucleotides and phosphatidylserine (PS).
3. The method of claim 1 or 2, wherein the phosphatidylserine is co-administered with the tolerogenic composition.
4. The method of any of claims 1-3, wherein said one or more polynucleotides comprises a chemically modified mRNA, said chemically modified mRNA encoding an immunomodulatory polypeptide.
5. The method of claim 4, wherein said immunomodulatory polypeptide is an antibody.
6. The method of claim 4, wherein said immunomodulatory polypeptide is selected from the group consisting of an inhibitor of mTOR, IL-2, an anti-IL-2 complex, IL-10, TGF- β , IL-35, galectin-1, IL-23, IL-27, IL-35 and IL-37.
7. The method of claim 5, wherein said antibody is specifically reactive with a member selected from the group consisting of CD3, CD40, CD40 ligand, CD4, and CTLA-4.
8. The method of any of claims 3-7, wherein said mRNA comprises at least one region which is codon optimized.
9. The method of any of claims 1-8, wherein the tolerogenic composition is formulated in a lipid nanoparticle (LNP).
10. The method of claim 8, wherein the LNP comprises phosphatidylserine (PS).

11. The method of claim 9, wherein the formulated LNP composition is administered systemically.
12. A method of treating an autoimmune disease, inflammatory diseases, allograft transplant/graft vs. host disease (GVHD), diabetes or multiple sclerosis, comprising contacting a cell, tissue or organism with phosphatidylserine (PS) and a tolerogenic composition comprising an antigen and one or more polynucleotides encoding a tolerogenic polypeptide of interest.
13. The method of claim 12, wherein the autoimmune disease is lupus.
14. The method of claim 12, wherein the inflammatory disease is selected from the group consisting of colitis, Crohn's disease, allergic encephalitis.
15. The method of any of claims 12-14, wherein the tolerogenic composition is formulated in a lipid nanoparticle (LNP).
16. The method of claim 15, wherein the LNP comprises phosphatidylserine (PS).
17. The method of claim 15, wherein the formulated LNP composition is administered systemically.
18. The method of any of claims 12-14, wherein the phosphatidylserine is co-administered with the tolerogenic composition.
19. A method of treating a subject with an environmental hypersensitivity or allergic reaction to a protein or antigen comprising contacting said subject with a tolerogenic composition comprising the antigen and one or more polynucleotides.
20. The method of claim 19, wherein the subject is contacted using a micro-dosing regimen.
21. The method of claim 20, wherein the micro-dosing regimen occurs over a period of time selected from the group consisting of at least 1 hour, at least 2 hours, at least 3 hours, at least 4 hours, at least 5 hours, at least 6 hours, at least 7 hours, at least 8 hours, at least 9 hours, at least 10 hours, at least 11

hours, at least 12 hours, at least 13 hours, at least 14 hours, at least 15 hours, at least 16 hours, at least 17 hours, at least 18 hours, at least 19 hours, at least 20 hours, at least 21 hours, at least 22 hours, at least 23 hours, at least 1 day, at least 36 hours, at least 2 days, at least 3 days, at least 4 days, at least 5 days, at least 6 days, at least 1 week, at least 2 weeks, at least 3 weeks, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months or more than 6 months.

22. The method of claim 21, wherein the dose of the amount of antigen and one or more polynucleotides in the tolerogenic composition increases during the micro-dosing regimen.
23. The method of claim 19, wherein the environmental hypersensitivity or allergic reaction is to a protein or antigen or other trigger selected from the group consisting of plants, grass, tree and other plant pollens, mammal, animal hair or dander, insect, insect parts, excreta or venom, helminthes, non-mammals, birds and reptiles, foods (e.g., legumes, nuts and fish), bacteria, fungi and molds, drugs, metals, wool and latex.

FIG. 1A




FIG. 1B




FIG. 2




FIG. 3




FIG. 4


Patterns of positional modification

A	(L)	B	(L)	C	(L)
5' [NNNNN _n] _x L1 [NNNNNNNNNNNNNN _o] _y L2 [XXXXXXXXXXXXX _p] _z L3 3'					
5' [NNNNN _n] _x L1 [XXXXXXXXXXXXXXXXX _o] _y L2 [XXXXXXXXXXXXX _p] _z L3 3'					
5' [XXXXXX _n] _x L1 [XXXXXXXXXXXXXXXXX _o] _y L2 [XXXXXXXXXXXXX _p] _z L3 3'					
5' [NNNNN _n] _x L1 [NNNNNNNNNNNNNN _o] _y L2 [XXXXXXXXXXXXX _p] _z L3 3'					

A, B, C- Region or Part of chimeric polynucleotide
 N-nucleoside
 n, o, p-number of nucleosides
 x, y, z-number of regions
 X-selective placement nucleoside
 L-linker (optional)

FIG. 5

Blocked or structured 3' termini



A, B, C- Region or Part of chimeric polynucleotide
 N-nucleoside
 n, o, p-number of nucleosides
 x, y, z-number of regions
 X-selective placement nucleoside
 K-non-nucleosidic moiety or conjugate
 L-linker (optional)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US15/48225

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 39/00, 39/35; C12N 5/07 (2015.01)

CPC - A61K 39/00, 31/685; C07K 16/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8): A61K 39/00, 39/35, 31/685; C12N 5/07, 5/16; C07K 16/28, 16/18 (2015.01)

CPC: A61K 38/00, 39/00, 31/685; C07K 16/28, 16/18

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatSeer (US, EP, WO, JP, DE, GB, CN, FR, KR, ES, AU, IN, CA, INPADOC Data); Google; Google Scholar; PubMed; ScienceDirect; induc*, mak*, generat*, toleranc*, cell*, system*, antigen*, phosphatidylserine, 'PS,' tolerogenic*, composition*, formula*, polynucleotid*, 'Moderna Therapeutics,' 'Stephen Hoge,' 'Eric Huang,' 'Louis O'Dea,' 'Treg,' 'regulatory T cells,' co-administer*, concurrent*, administer*

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/184976 A2 (NORTHWESTERN UNIVERSITY, et al.) 12 December 2013; abstract; paragraphs [0006], [0016]-[0018], [0020], [0022], [0029], [0058], [00112], [00127], [00157].	1-3
A	FREUND'S ADJUVANT, COMPLETE AND INCOMPLETE: Product Information Sheet: Product Number F 5881 and F 5506. Sigma-Aldrich. 08 April 1998.	1-3

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

06 January 2016 (06.01.2016)

Date of mailing of the international search report

14 JAN 2016

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US15/48225

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 4-11
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

.-***-Please See Supplemental Page-***-

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Group I: Claims 1-3

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

.-***-Continued from Box No. III: Observations Where Unity of Invention Is Lacking:

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I: Claims 1-3 are directed toward methods of inducing tolerance in a cell system and inducing Treg cell activity in a cellular system.

Group II: Claims 12-18 are directed toward a method of treating an autoimmune disease, inflammatory diseases, allograft transplant/graft vs. host disease (GVHD), diabetes or multiple sclerosis.

Group III: Claims 19-23 are directed toward a method of treating a subject with an environmental hypersensitivity or allergic reaction.

The inventions listed as Groups I-III do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the special technical features of Group I include inducing Treg cell activity, which is not present in either of Groups II or III, the special technical features of Group II including treating multiple sclerosis, which is not present in Groups I or III, the special technical features of Group III including treating a subject with an environmental hypersensitivity, which is not present in either of Groups I or II.

Groups I-III share the technical features including: contacting a subject or cellular system with a tolerogenic composition comprising an antigen and one or more polynucleotides. Groups I and II share the technical features including contacting a cellular system with phosphatidylserine (PS) and a tolerogenic composition comprising an antigen, and one or more polynucleotides. Groups II and III share the technical features including methods of treating a disease in a subject.

However, these shared technical features are previously disclosed by WO 2013/184976 A2 to Northwestern University, et al. (hereinafter 'Northwestern').

Northwestern discloses contacting a subject (administering a composition to a subject; paragraph [0017]) with a tolerogenic composition (with a composition that induces tolerance; paragraph [0017]) comprising an antigen (comprising one or more antigens; paragraph [0017]) and one or more polynucleotides (and complete Freund's Adjuvant (and one or more polynucleotides), paragraphs [0044], [0284]; wherein complete Freund's Adjuvant comprises killed mycobacteria comprising genomic DNA polynucleotides (see Sigma Specifications for CFA)); contacting a cellular system (administering a composition to a subject having an immune system; paragraphs [0006], [0017], [0127]) with phosphatidylserine (PS) (comprising liposomes comprising phosphatidylserine; paragraphs [0031], [0112]) and a tolerogenic composition (a composition that induces tolerance; paragraph [0017]) comprising an antigen (comprising one or more antigens; paragraph [0017]), and one or more polynucleotides (and complete Freund's Adjuvant (and one or more polynucleotides), paragraphs [0044], [0284]; wherein complete Freund's Adjuvant comprises killed mycobacteria comprising genomic DNA polynucleotides (see Sigma Specifications for CFA)); and treating a disease in a subject (paragraph [0056]).

Since none of the special technical features of the Groups I-III inventions is found in more than one of the inventions, and since all of the shared technical features are previously disclosed by the Northwestern reference, unity of invention is lacking.