

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 June 2006 (22.06.2006)

PCT

(10) International Publication Number
WO 2006/063639 A1

(51) International Patent Classification:

A61K 31/205 (2006.01) *A61P 25/00* (2006.01)
A61P 43/00 (2006.01) *A61P 21/00* (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:

PCT/EP2005/012054

(22) International Filing Date:

10 November 2005 (10.11.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

RM2004A000606

13 December 2004 (13.12.2004) IT

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicant (for all designated States except US):
SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE S.P.A. [IT/IT]; Viale Shakesepare, 47, I-00144 Rome (IT).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **KOVERECH, Aleardo** [IT/IT]; 200, Via Aurelia Antica, I-00165 Rome (IT).

(74) Agent: **CAMPANELLI, Domenico**; c/o Sigma-Tau Industrie Farmaceutiche Riunite S.p.A., Via Pontina, Km 30,400, I-00040 Pomezia (IT).

WO 2006/063639 A1

(54) Title: USE OF ACETYL L-CARNITINE FOR THE TREATMENT OF FIBROMYALGIC SYNDROME

(57) Abstract: The use of acetyl L-carnitine or one of its pharmaceutically acceptable salts is described for the preparation of a medicament and/or dietetic product for the treatment of fibromyalgic syndrome.

Use of acetyl L-carnitine for the treatment of fibromyalgic syndrome

The present invention relates to the use of acetyl L-carnitine for the preparation of a medicament for the treatment of fibromyalgic syndrome.

Fibromyalgic syndrome or fibromyalgia is a chronic pain syndrome characterised by diffuse arthromyalgias accompanied to varying extents by sleep disorders, muscle-tension headache, irritable bowel syndrome, and asthenia.

The symptoms sometimes present in isolation and sometimes in association with other rheumatic diseases or with endocrine and metabolic diseases, particularly dysthyroidism (Arthritis Rheum 1983; 26: 817-824; Seminar Arthritis Rheum 1981;11:151-71).

The condition is distinctly more prevalent in subjects of female gender and is more frequent in patients aged from 25 to 55 years.

The cause of fibromyalgia is still unknown and various pathogenetic mechanisms are currently postulated, such as psychic disturbances, alteration of normal sleep activity and of the pain perception threshold, and functional disorders affecting skeletal muscle with the presence of morphological alterations of the muscle fibre cells (Scand. J. Rheumatol. 1986; 15: 1-6) or various metabolic deficits (Bull Rheum Dis 1978; 26: 928-931; J. Rheumatol 1992; 19:90-94).

The diagnosis of fibromyalgia is obtained on a purely clinical basis (medical history and physical examination) since laboratory tests (haematology and blood chemistry) and instrumental investigations (radiology, electrophysiology) yield no diagnostically useful indications in this clinical picture.

Medical history and physical examination are therefore decisive for

the purposes of identifying this syndrome. The location of the pain, described as experienced "all over", the abundance and colourfulness of the terms the patient uses to describe it, the plentiful array of associated symptoms, including headache, irritable bowel syndrome, and sleep disorders, and the history of numerous examinations by specialists and instrumental investigations may suggest the diagnosis of a clinical picture indicative of a "fibrositic situation". The symptoms are generally experienced mainly in the morning, deteriorate with fatigue, physical activity, stress and changes in the weather, but improve with massage and periods of rest and recreation. At the physical examination an excessive muscular tension is noted, even to the extent of an actual contracture affecting typical muscular districts such as the sternocleidomastoid, trapezius and paravertebral muscles where tender points need to be sought (Br Med J 1994; 309: 696-9).

Essential for the diagnosis of fibromyalgia is the demonstration of deep tender points with areas of increased soft tissue consistency (Arthritis Rheum 1990; 33(2): 160-172). Tender points differ from trigger points, in that they do not cause referred and irradiated pain, but prove tender only in the site stimulated. The simplest technique for detecting tender points is simple digital pressure (approximately 4 kg with the digital pulp) or by pinching the skin at points corresponding to definite joints and tendon insertions. Pressure on these points is accompanied by an exaggerated reaction of withdrawal from the examiner, jump sign, and neurovegetative phenomena such as dermographia, horripilation, and cold sweats. The tender points most commonly used for diagnostic purposes are nine in

number (bilaterally), and to exemplify the diagnostic and classificatory approach criteria have been defined for locating and evaluating the tender points present in the patient (Arthritis Rheum 1990; 33(2): 160-172).

Previous therapeutic uses of acetyl L-carnitine are already known..

US 4,751,242 describes the use of acetyl L-carnitine for the treatment of peripheral neuropathies.

US 4,362.719 describes the use of L-carnitine and acyl L-carnitines for the treatment of juvenile onset diabetes mellitus.

US 5192805 relates to the use of acetyl L-carnitine in the therapeutic treatment of coma.

US 6,037,3721128 relates to the use of acetyl L-carnitine, isovaleryl L-carnitine and propionyl L-carnitine to increase IGF-1 levels, for the treatment of lateral amyotrophic sclerosis, neuropathies of the optic and olfactory nerves, trigeminal nerve neuralgia and other pathologies.

US 6,037,372 relates to the use of an L-carnitine alkanoyl, including acetyl L-carnitine, for the treatment of pathologies mediated by glutamate such as epilepsy, schizophrenia, chronic fatigue syndrome, lateral amyotrophic sclerosis and others.

Acetyl L-carnitine, a commercially available product, can be prepared with the process described by R. Krinmberg, and W. Wittandt, in Biochem. Z. 251, 229 (1932).

Fibromyalgia, despite being a clinical condition with a benign course, is difficult to treat (Rheum Dis Clin N Am 1989; 15:61-71).

The patients often use anti-inflammatory drugs which, in addition to having no pharmacological rationale in this condition, are ineffective.

Muscle relaxants and antidepressant drugs have also been used for the treatment of this disease, but only improve the quality of sleep (Arthritis Rheum 1986; 29: 1371-7; Arthritis Rheum 1994, 37: 3240).

Good results, albeit short-lasting, have also been obtained by means of local infiltration of the tender points with anaesthetics.

There is therefore a strongly perceived need for the availability of new drugs useful for the treatment of fibromyalgia which are more effective and do not present the drawbacks of the above-mentioned known drugs.

It has now been found that acetyl L-carnitine or one of its pharmaceutically acceptable salts is a useful compound for the treatment of fibromyalgia.

What is meant by pharmaceutically acceptable salt of acetyl L-carnitine is any salt prepared by addition of an acid to acetyl L-carnitine inner salt, and which does not give rise to unwanted toxic or side effects. The formation of salts by addition of an acid is well known in pharmaceutical technology.

Non-limitative examples of such salts are chloride, bromide, orotate, aspartate, acid aspartate, citrate, acid citrate, magnesium citrate, phosphate, acid phosphate, fumarate, acid fumarate, magnesium fumarate, glycerophosphate, lactate, maleate and acid maleate, mucate, oxalate, acid oxalate, pamoate, acid pamoate, sulphate, acid sulphate, glucose phosphate, tartrate, acid tartrate, magnesium tartrate, 2-amino ethanesulphonate, magnesium 2-amino ethanesulphonate, methanesulphonate, choline tartrate, trichloroacetate, and trifluoroacetate.

The subject of the present invention is therefore the use of acetyl L-carnitine or of one of its pharmaceutically acceptable salts for the preparation of a medicament and/or dietetic product for the treatment of fibromyalgia.

The acetyl L-carnitine can be in any form suitable for oral or parenteral administration. It can be administered in the form of a unit dose or in divided doses, in amounts ranging from 40 mg to 4 g/day.

The preferred dose according to the present invention is 1.5 g per day, administered at the start of treatment and for the first 2 weeks at the dose of 500 mg intramuscularly (i.m.) and 500 mg (twice daily) orally (os), the treatment thereafter being continued with oral doses of 500 mg three times daily.

The daily dose will depend, according to the judgement of the primary care physician, on the patient's weight, age and condition. Larger doses of acetyl L-carnitine can be administered thanks to the extremely low toxicity of said active ingredient.

The following examples illustrate the invention.

A multicentre, randomized, placebo-controlled, double-blind clinical trial was conducted, the aim of which was to evaluate the efficacy of acetyl L-carnitine in patients with fibromyalgic syndrome.

For the evaluation of the efficacy of the compound according to the present invention evoked pain was monitored (as measured with the "Pressure Threshold Meter" algometer) in the osteoarticular regions indicated by "The American College of Rheumatology" (ACR) as being "tender points" characteristic of fibromyalgia syndrome. [WOLFE F,

SMYTHE HA, YUNUS MB, *et al.* The American College of Rheumatology Criteria for the classification of fibromyalgia: report of the multicenter criteria committee; *Arthritis Rheum* 1990; 33(2): 160-172].

Also monitored were additional secondary parameters which will be discussed later.

The evaluation of the tolerability of the treatment was done by means of routine haematological and blood-chemistry tests and by recording adverse events.

The patients included in the trial were patients of either gender aged from 25 to 65 years, who matched up to the above-mentioned American College of Rheumatology diagnostic criteria for fibromyalgia syndrome.

The diagnostic criteria for fibromyalgic syndrome had to be confirmed by the presence of all four of the following conditions:

- bilateral diffuse pain in the axial skeleton;
- bilateral diffuse pain in the upper limbs;
- bilateral diffuse pain in the lower limbs;
- pain located in 11 or more of the 18 anatomical regions listed in the map of tender points (ACR tender points) evoked with a standardised pressure manoeuvre.

Patients with any one of the following conditions were not recruited into the trial:

- presence of infections or inflammatory processes affecting the osteoarticular or tegumental apparatus;

- history or presence of clinically important osteoarticular disease capable of confounding the evaluation of the study treatments (e.g. history of trauma);
- previous orthopaedic surgery, spondyloarthritis, connectivitis etc.;
- clinical or laboratory evidence of dysthyroidism;
- presence of systemic diseases (diabetes, kidney, cardiovascular and respiratory disease, psychiatric illness) or major pathological conditions capable of interfering with the evaluation of the study treatments;
- continuous use of anti-inflammatory steroids during the last 3 months prior to the trial;
- recent initiation of oestrogen-progestogen hormone replacement therapy (< 1 year);
- therapy with antidepressant drugs during the last 6 months prior to the trial;
- therapy with non-steroidal anti-inflammatory drugs during the last 3 days prior to the trial;
- therapy with analgesic drugs during the last 7 days prior to the trial;
- participation in a clinical trial during the previous 6 months;
- pregnant women, breastfeeding women, or women of childbearing age not adopting an adequate contraceptive method.

Acetyl L-carnitine was administered for a period of 10 consecutive weeks at a dose of 1,500 mg/day.

During the first two weeks it was administered at the dose of 500 mg by the intramuscular route (i.m.) and 500 mg (twice daily) by the oral route (os).

Over the following eight weeks acetyl L-carnitine was administered orally at doses of 500 mg three times daily.

The results obtained are given in Table 1 here below.

Table 1

Visit	Mean variation compared to baseline		P (Student's t-test)
	Placebo (N=47)	Acetyl L-carnitine (N=42)	
2 weeks	0.16 (0.39)	0.24 (0.64)	-
6 weeks	0.45 (0.57)	0.55 (0.64)	-
10 weeks	0.34 (0.66)	0.71 (0.71)	<0.05

The results obtained, presented in Table 1, show that at the end of therapy the patients treated with the compound according to the present invention had a statistically significant improvement in pain evoked in the osteoarticular regions (the above-mentioned ACR regions) indicated as tender points characterised by fibromyalgic syndrome as compared to the control group.

In addition, at the end of the treatment period the patients treated with the compound according to the invention presented statistically significant improvements (assessed with visual analogue scales) in secondary clinical parameters relating to general well-being, mental health, and physical health in general ($P<0.05$).

Pomezia,

CLAIMS

1. Use of acetyl L-carnitine or one of its pharmaceutically acceptable salts for the preparation of a medicament and/or dietetic product for the treatment of fibromyalgia.
2. Use according to claim 1, in which the pharmaceutically acceptable salt of acetyl L-carnitine is selected from the group consisting of chloride, bromide, orotate, aspartate, acid aspartate, citrate, acid citrate, magnesium citrate, phosphate, acid phosphate, fumarate, acid fumarate, magnesium fumarate, glycerophosphate, lactate, maleate and acid maleate, mucate, oxalate, acid oxalate, pamoate, acid pamoate, sulphate, acid sulphate, glucose phosphate, tartrate, acid tartrate, magnesium tartrate, 2-amino ethanesulphonate, magnesium 2-amino ethanesulphonate, methanesulphonate, choline tartrate, trichloroacetate, and trifluoroacetate.

Pomezia,

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2005/012054

A. CLASSIFICATION OF SUBJECT MATTER A61K31/205 A61P43/00 A61P25/00 A61P21/00			
According to International Patent Classification (IPC) or to both national classification and IPC			
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A61K A61P			
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched			
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, EMBASE, BIOSIS, CHEM ABS Data			
C. DOCUMENTS CONSIDERED TO BE RELEVANT			
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	
Y	ONOFRJ M ET AL: "L-ACETYL CARNITINE AS A NEW THERAPEUTIC APPROACH FOR PERIPHERAL NEUROPATHIES WITH PAIN" INTERNATIONAL JOURNAL OF CLINICAL PHARMACOLOGY RESEARCH, BIOSCIENCE EDIPRINT, GENEVA,, CH, vol. 15, no. 1, 1995, pages 9-15, XP001105959 ISSN: 0251-1649 abstract	1,2	
Y	WO 02/096409 A (SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE S.P.A; CALVANI, MENOTTI; MOS) 5 December 2002 (2002-12-05) abstract; claims	1,2	
Y	US 2004/198837 A1 (MENDEL CARL M ET AL) 7 October 2004 (2004-10-07) abstract	1,2	
<input type="checkbox"/> Further documents are listed in the continuation of Box C.		<input checked="" type="checkbox"/> See patent family annex.	
* Special categories of cited documents :			
A document defining the general state of the art which is not considered to be of particular relevance			
E earlier document but published on or after the international filing date			
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)			
O document referring to an oral disclosure, use, exhibition or other means			
P document published prior to the international filing date but later than the priority date claimed			
T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention			
X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone			
Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.			
& document member of the same patent family			
Date of the actual completion of the international search		Date of mailing of the international search report	
8 February 2006		16/02/2006	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, TX. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Herrera, S	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2005/012054

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 02096409	A 05-12-2002	CA	2448242 A1	05-12-2002
		CZ	20033220 A3	16-06-2004
		EP	1399141 A1	24-03-2004
		HU	0401294 A2	28-12-2004
		IT	RM20010293 A1	29-11-2002
		JP	2004532865 T	28-10-2004
		MX	PA03010919 A	27-02-2004
		SK	15842003 A3	08-06-2004
		US	2004152781 A1	05-08-2004
US 2004198837	A1 07-10-2004	US	6803387 B1	12-10-2004