

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)


PCT

(10) International Publication Number
WO 01/85129 A2

- (51) International Patent Classification⁷: A61K 7/48
 - (21) International Application Number: PCT/US01/14533
 - (22) International Filing Date: 4 May 2001 (04.05.2001)
 - (25) Filing Language: English
 - (26) Publication Language: English
 - (30) Priority Data:
09/565,716 5 May 2000 (05.05.2000) US
 - (71) Applicant: JOHNSON & JOHNSON CONSUMER COMPANIES, INC. [US/US]; 199 Grandview Road, Skillman, NJ 08558 (US).
 - (72) Inventors: COLE, Curtis, A.; 9 Orchard Road, Ringoes, NJ 08551 (US). FLACK, Laura, E.; 1054 Knollwood Drive, Birmingham, AL 35242 (US). KAMINSKI, Claudia; 490 Shire Road, Milford, NJ 08848 (US). VAN LEEUWEN, Claudia; 5, rue Arthur Papavoine, F-27100 Le Vaudreuil (FR). LUKENBACH, Elvin, R.; 160 Klinesville Road, Flemington, NJ 08822 (US).
 - (74) Agents: JOHNSON, Philip, S. et al.; Johnson & Johnson, One Johnson and Johnson Plaza, New Brunswick, NJ 08933 (US).
 - (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
 - (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: SKIN CARE COMPOSITION



(57) Abstract: There are provided compositions which include a retinoid and preferably retinol; a dermatologically active acid; and a volatile base, such as ammonium hydroxide. Another embodiment of the invention includes compositions comprising a retinoid and preferably retinol; a dermatologically active acid; a volatile base; and a second neutralizing agent. There are also provided compositions which include a retinoid, a neutralized ammonium salt of a dermatologically active acid, and optionally a neutralized salt, other than ammonium salt, of an acid. Further provided are methods for reducing fine lines, wrinkles, skin roughness, and pore size and for increasing the clarity of a skin surface, cellular turnover, skin radiance, skin smoothness, skin permeation or collagen synthesis in a mammal in need thereof. Compositions as described above are administered topically to the skin of the animal.

WO 01/85129 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

SKIN CARE COMPOSITION

5 This application is a continuation-in-part of
U.S. Serial No. 09/436,867, filed November 9, 1999, which
claims priority from U.S. Serial No. 09/325,452, filed
June 3, 1999, and U.S. Serial No. 60/017,956, filed
November 12, 1998, all of which are herein incorporated
10 by reference.

FIELD OF THE INVENTION

 This invention relates to skin care
compositions which include, in a single formulation, the
15 beneficial ingredients for aging or photodamaged skin,
retinol and an acid.

BACKGROUND OF THE INVENTION

 Retinol or vitamin A alcohol is useful in the
20 reduction of fine lines, wrinkles, and mottled
hyperpigmentation in skin. Hydroxy acids, and
particularly alpha-hydroxy acids, are useful in
increasing the clarity of the skin surface, increasing
cellular turnover, and increasing skin radiance and

smoothness. Ascorbic acid has skin permeation and collagen synthesis activity.

Retinol is physically unstable and rapidly degrades when stored at a pH below about 5. Acids such as hydroxy acids, and particularly alpha-hydroxy acids and ascorbic acid, on the other hand, are not active in increasing skin cell turnover, exfoliation, skin permeation, and/or collagen synthesis at pHs above about 5.

Consequently, retinol and hydroxy acids and/or ascorbic acid have generally been packaged separately. Retinol typically is packaged in a vehicle at a pH above about 5, while alpha-hydroxy acids and ascorbic acid are packaged at a pH of about 4 or below. Therefore, one must apply two separate products in order to achieve the benefit of both of these ingredients.

The present inventors have discovered a single composition which include both of these ingredients, in which both of these ingredients are stable, and in which both of these ingredients are active upon application to the skin.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graphic illustration of skin pH over time after treatment.

Figure 2 is a graphic illustration of cell proliferation measured as slope of fluorescence after treatment.

Figure 3 is a graphic illustration comparing the activity of ammonium hydroxide and sodium hydroxide neutralized alpha-hydroxy acids in combination with retinol.

Figure 4 is a graphic illustration of skin pH over time before and after treatment.

SUMMARY OF THE INVENTION

According to one embodiment of the present invention there are provided compositions which include:

- (A) a retinoid and preferably retinol;
- (B) a dermatologically active acid; and
- (C) a volatile base, such as, for example, a volatile compound comprising an amine such as ammonium hydroxide or 2-dimethylaminoethanol (N,N-dimethylethanolamine or DMAE). Volatile bases have a vapor pressure typically below atmospheric pressure, preferably below about 700 mm Hg, and more preferably below about 600 mm Hg. The volatile base preferably

evaporates upon contact with skin. The compositions preferably contain an acid neutralizing effective amount of ammonium hydroxide or DMAE.

Another embodiment of the present invention provides compositions which include:

- (A) a retinoid and preferably retinol;
- (B) a dermatologically active acid;
- (C) a volatile base; and
- (D) at least one second neutralizing agent.

According to yet another embodiment of the present invention, there are provided compositions which include:

- (A) retinol; and
- (B) a neutralized ammonium salt of a

dermatologically active acid (e.g. the ammonium salt formed by a volatile compound comprising an amine such as ammonium hydroxide or a volatile alkanolamine such as DMEA). Examples of such salts include ammonium glycolate and N₁N-dimethylethanolammonium glycolate. Optionally, a second neutralized salt of a dermatologically active acid is included in the compositions.

Further provided are methods for reducing fine lines, wrinkles, skin roughness, and pore size and for increasing the clarity of a skin surface, cellular

turnover, skin radiance and skin smoothness in an animal, for example, a mammal, such as a human, in need thereof.

Compositions as described above are administered topically to the skin of the animal.

5 Methods for preparing the compositions above are also provided.

Other features and advantages of the invention will be apparent from the detailed description of the invention, the drawings, and the claims.

10

DETAILED DESCRIPTION OF THE INVENTION

In one embodiment, the present formulations provide compositions which have a storage pH of about 5 or higher. This provides storage stability for the retinoid compound (e.g. retinol). However, the pH of these compositions drops to below 5 when applied to the skin. This allows the hydroxy acid(s) and/or other skin beneficial acids(s) therein to become active upon application of the composition to the skin.

20 Retinoids suitable for use in the present invention preferably are unstable or pH sensitive in that they are chemically and physically unstable at relatively low pH such as, for example a pH below about 5, such as retinol and derivatives thereof. Suitable retinoids

include, but are not limited to retinol and derivatives thereof, such as retinyl palmitate and retinyl acetate; retinaldehyde; and like compounds that bind to retinoid receptors.

5 Retinol is also known as vitamin A alcohol. Retinol is chemically and physically unstable at a pH below about 5. It is useful in reducing fine lines at wrinkles in skin. It is also useful in reducing mottled hyperpigmentation of skin. Other retinoids having pH
10 dependent stability may also be used in combination with or in place of retinol in the present invention.

 The dermatologically active acid may be a cosmetically active acid or a pharmaceutically active acid, such as, for example, a hydroxy acid, ascorbic acid
15 or a derivative thereof, lipoic acid, dihydrolipoic acid, or a combination thereof.

 Hydroxy acids useful in the present invention are either alpha- or beta-hydroxy acids, poly-hydroxy acids, or any combinations of any of the foregoing.
20 Preferably, the hydroxy acid is an alpha-hydroxy acid. Examples of alpha hydroxy acids include, but are not limited to, glycolic acid, malic acid, tartaric acid, pyuric acid, citric acid, or any combination of any of the foregoing. Special mention is made of glycolic acid.

Beta-hydroxy acids include, but are not limited to, salicylic acid.

Other suitable hydroxy acids are disclosed in U.S. Patent No. 5, 889,054, which is hereby incorporated by reference.

Other acids suitable for use in the present invention include, but are not limited to, ascorbic acid and derivatives thereof, lipoic acid, and dihydrolipoic acid. Suitable ascorbic acid derivatives include, but are not limited to, magnesium ascorbyl phosphate; sodium ascorbyl phosphate; sodium ascorbate; and ascorbyl glucosides.

Suitable second neutralizing agents which may be included in the composition include, but are not limited to, alkali hydroxides, such as sodium hydroxide and potassium hydroxide; and organic bases, such as alkanolamines, including, but not limited to, diethanolamine, triethanolamine, DMAE and aminobutanol; ammonium hydroxide, and amino acids, including, but not limited to, arginine and lysine; and any combination of any of the foregoing. A preferred second neutralizing agent is sodium hydroxide.

When utilized, ammonium hydroxide is typically added as a solution containing from about 27 to about 31

percent by weight of ammonium hydroxide based upon 100 percent by weight of total ammonium hydroxide solution.

The compositions of the present invention may also include other adjuvants, such as, for example, vehicles including, but not limited to, water or alcohol; humectants, including, but not limited to, glycerin; buffering agents including, but not limited to, citric acid and sodium citrate; viscosity adjusters, including, but not limited to, carbomer gelling agents, gum derivatives, and other viscosity controlling, decreasing, and increasing agents; preservatives including, but not limited to, parabens, such as methylparaben and propylparaben, and phenoxyethanol; emulsifiers including, but not limited to, polysorbate 80, glyceryl distearate, POE 10 stearyl ether, steareth 10, ceateareth 20 and stearyl alcohol, and cetareth 20 and cetearyl alcohol; conditioning agents including, but not limited to, octyl hydroxystearate, stearyl alcohol, lactose, and dimethicone; emollients including, but not limited to, cholesterol NF, petrolatum, mineral oils and esters including, but not limited to, isopropyl myristate, isopropyl palmitate, 1-decene polymer (hydrogenated), and C₁₂-C₁₅ alcohol benzoates; thickeners, including, but not limited to, binders, polyacrylamide, C₁₃-C₁₄ isoparaffin,

and laureth-7; antioxidants, including, but not limited to ascorbic acid, butylated hydroxytoluene (BHT), tocopheryl acetate, and the like; UV stabilizers; UV radiation absorbers (sunscreen filters); fragrances; colorants; chelating agents, including, but not limited to, disodium ethylenediaminetetraacetate (EDTA); or any combinations of any of the foregoing. Examples of these adjuvants are disclosed in the International Cosmetic Ingredient Dictionary and Handbook, 7th Ed. (1997)

These compositions can be formulated as creams, gels, or liquids, and preferably are prepared as lotions.

These compositions can be prepared as liposomes, including, but not limited to, unilamellar, multilamellar, or paucilamellar vesicles; nanospheres; microsponges; emulsions, such as a multiple emulsion and a cleansing emulsion; or any combination of any of the foregoing by methods known to those skilled in the art. In one embodiment, the composition is prepared as a paucilamellar vesicle (e.g. containing the retinoid and/or the dermatologically active acid or salt thereof) having, for example, between 2 and 10 lipid bilayers and a lipophilic core which may contain an apolar oil or wax.

The compositions are typically neutralized to a pH above about 4.5, preferably ranging from about 4.5 to

about 8 and most preferably from about 5 to about 6. The amount of the volatile base (e.g. ammonium hydroxide or DMAE) and optionally second neutralizing agent useful herein is that amount sufficient to adjust the pH of the compositions to the above pH ranges. The amount of volatile base in the compositions of the present invention is preferably that amount sufficient to adjust the pH of the acid from about 4.0 or less to at least about 5.

A preferred method of preparation includes neutralizing the composition to a pH of about 4.0 or less with the aforementioned second neutralizing agent and then further neutralizing the composition to a pH of at least about 5 with the volatile base.

The amount of retinoid in these compositions is typically a fine line-, wrinkle-, or mottled pigmentation-reducing effective amount. The amount of retinoid (e.g., retinol) is at least about 0.001 percent by weight, (e.g., about 0.01 to about 10 percent, such as about 0.01 to about 1 percent), based upon 100 percent by weight of total composition.

The amount of acid, ammonium salt of acid, or other salt of the acid is typically a skin surface clarity, cellular turnover-, skin radiance-, skin

smoothness-, skin permeation-, or collagen synthesis-
increasing effective amount. Preferably, this amount
ranges from about 0.1 to about 20 percent by weight based
upon 100 percent by weight of total composition. More
5 preferably this amount ranges from about 1 to about 12
percent by weight, and most preferably, this amount is
from about 4 to about 8 percent by weight, based upon 100
percent by weight of total composition.

The composition preferably contains from about
10 1 to about 99 percent, and more preferably from about 60
to about 95 percent by weight of water, based upon 100
percent by weight of total composition.

Generally, the composition contains sufficient
thickener to impart body to the composition without
15 causing it to become so viscous as to hinder
spreadability of the composition. The composition also
preferably contains up to about 5 percent by weight of a
viscosity adjuster, up to about 20 percent by weight of
an emollient, from about 0.1 to about 10 percent by
20 weight of an emulsifier, up to about 5 percent by weight
of a spreading agent, up to about 10 percent by weight of
a thickener, a preservative, a chelating agent, and a
humectant, based upon 100 percent weight of total
composition. More preferably, the composition contains

from about 0.1 to about 2 percent by weight of a
viscosity adjuster, from about 3 to about 5 percent by
weight of an emulsifier, from about 1 to about 2 percent
by weight of a spreading agent, an antimicrobially
5 effective amount of a preservative, and from about 3 to
about 5 percent by weight of a thickener, based upon 100
percent weight of total composition.

Without being bound by any theory, applicants
believe that by using a salt of the acid and a volatile
10 base, the storage pH of the present composition can
remain above 5, thereby providing a stable atmosphere for
the retinol or any other pH sensitive ingredient.
However, when applied to the skin, the pH of the salt of
the acid changes by volatilization of the volatile base
15 (e.g., the ammonium). The pH then drops to a range in
which the acid can cause beneficial changes.

The compositions can be applied topically to a
mammal, and preferably a human, in need of a retinoid,
acids, or a combination thereof. Typically, the amount
20 applied will be that amount effective to accomplish the
purpose of application.

The following examples illustrate the invention
without limitation. All amounts are given as weight

percentages based upon 100 percent by weight of total composition unless noted otherwise.

Example 1

5 A retinol/alpha-hydroxy acid composition having the formulation of Table 1 and a pH of about 6 and containing paucilamellar vesicles was prepared by a shear mixing method. The apparatus used to prepare the liposomes by the shear mixing method is described in U.S. Patent No. 4,895,452, which is hereby incorporated by
10 reference. A mixture containing the appropriate amounts of the ingredients for the lipid phase was heated in a container at about 75° C until all of the lipids melted. The lipid melt was then cooled to about 65° C. The
15 ingredients for the aqueous phase were mixed together, heated to about 75° C to dissolve them, and then cooled to about 60° C. The lipid melt and aqueous phase mixture were then poured into separate holding reservoirs of the shear mixing apparatus. The positive displacement pump
20 for the lipid melt and aqueous phase mixture feed lines were turned on. The feed rate was adjusted to 1 part lipid to 4 parts aqueous phase. The aqueous phase mixture and lipid melt were fed through injection jets

into a cylindrical mixing chamber tangentially with respect to the cylinder wall. In the mixing chamber, the two streams of flowing liquid intersect in such a manner as to cause shear mixing that leads to the formation of liposomes. The liposomes formed were then withdrawn through an exit tube and transferred to a Cafero mixing vesicle. The liposomes were cooled to 40° C, under mixing at 200 rpm. After cooling, the single addition components listed in Table 1, were added sequentially. The resultant mixture was then mixed at 200 rpm for about 30 minutes. The formulation was allowed to cool to room temperature under ambient conditions.

Table 1			
Retinol/Alpha-Hydroxy Acid Liposome Formulation-pH6			
<u>TRADE NAME</u>	<u>CHEMICAL NAME</u>	<u>FUNCTION</u>	<u>%WT/W</u> <u>T</u>
<u>AQUEOUS PHASE</u>			
Deionized Water	D.I. Water	Vehicle	60.93
Glycerin 916	Glycerin	Humectant	4
Citric Acid	Citric Acid	Buffering Agent	0.13
Sodium Citrate	Sodium Citrate	Buffering Agent	0.5
Sodium Chloride	Sodium Chloride	Viscosity Adjuster	0.1
Methyl Parasept	Methylparaben	Preservative	0.25
Propyl Parasept	Propylparaben	Preservative	0.15
Tween 80	Polysorbate 80	Emulsifier	0.7

Glypure (70%)	Glycolic Acid	Skin Conditioner	5.71
NH ₄ OH [^]	Ammonium Hydroxide (27 to 31% Solution)	pH Adjuster (pH=6)	3.2
<u>LIPID PHASE</u>			
Wickenol 171	Octyl Hydroxystearate	Conditioning Agent	5.8
Kessco GDS	Glyceryl Distearate	Emulsifier	2.8
Cholesterol, NH	Cholesterol NF	Emulsifer	1
BRIJ 76	POE 10 Stearyl Ether	Emulsifer	1.4
Protocol ST 20G	Ceteareth 20 and Stearyl Alcohol	Emulsifier	3
Protocol CS 20D	Ceteareth 20 and Stearyl Alcohol	Emulsifier	3
Stearyl Alcohol	Stearyl Alcohol	Skin Conditioner	0.5
Retinol 50CJ**	Retinol in Polysorbate- 20	Skin Conditioner	0.4
BHT	BHT	Antioxidant	0.1
Vitamin E Acetate	Tocopheryl Acetate	Antioxidant	0.1
<u>SINGLE ADDITION COMPONENTS</u>			
Emeressence 1160	Phenoxyethanol	Preservative	0.73
Dimethicone 47V	100 Centistoke Dimethicone	Skin Conditioner	2.5
Sepigel 305	Polyacrylamide, C13-24 Isoparrifin and Laureth- 7	Thickener	3

**Retinol 50CJ is available from BASF of Mount Olive, NJ, and contains 50% by weight of retinol.

[^]Amount of NH₄OH required to reach pH of 6 is estimated; each batch will be titrated to pH=6.

5

The formulation was applied to the skin, and the pH of the skin was measured over time. Results are illustrated in Figure 1. The pH of the preparation

10

dropped to about 4.1 within 15 minutes of application.

This reduced the skin pH to about 4.

Comparative Example 1A

5 A retinol/alpha-hydroxy acid containing
composition having the formulation of Table 2 and a pH of
about 4 was prepared as described in Example 1. The
amount of ammonium hydroxide in this composition was
approximately half the amount incorporated in the
10 composition of Example 1.

<u>Table 2</u>			
Retinol/Alpha-Hydroxy Acid Liposome Formulation - pH4			
<u>TRADE NAME</u>	<u>CHEMICAL NAME</u>	<u>FUNCTION</u>	<u>%WT/WT</u>
AQUEOUS PHASE	(qs with DI water)		
Deionized Water	D.I. Water	Vehicle	62.43
Glycerin 916	Glycerin	Humectant	4
Citric Acid	Citric Acid	Buffering Agent	0.13
Sodium Citrate	Sodium Citrate	Buffering Agent	0.5
Sodium Chloride	Sodium Chloride	Viscosity Adjuster	0.1
Methyl Parasept	Methylparaben	Preservative	0.25
Propyl Parasept	Propylparaben	Preservative	0.15
Tween 80	Polysorbate 80	Emulsifier	0.7
Glypure (70%)	Glycolic Acid	Skin Conditioner	5.71
NH ₄ OH [^]	Ammonium	pH Adjuster	1.7

	Hydroxide 27 to 31% Solution	(pH=4)	
<u>LIPID PHASE</u>			
Wickenol 171	Octyl Hydroxystearate	Conditioning Agent	5.8
Kessco GDS	Glyceryl Distearate	Emulsifier	2.8
Cholesterol, NH	Cholesterol NF	Emollient	1
BRIJ 76	POE 10 Stearyl Ether	Emulsifier	1.4
Protocol ST 20G	Ceteareth 20 and Stearyl Alcohol	Emulsifier	3
Protocol CS 20D	Ceteareth 20 and Stearyl Alcohol	Emulsifier	3
Stearyl Alcohol	Stearyl Alcohol	Skin Conditioner	0.5
Retinol 50CJ**	Retinol in Polysorbate-20	Skin Conditioner	0.4
BHT	BHT	Antioxidant	0.1
Vitamin E Acetate	Tocopheryl Acetate	Antioxidant	0.1
<u>SINGLE ADDITION COMPONENTS</u>			
Emeressence 1160	Phenoxyethanol	Preservative	0.73
Dimethicone 47V	100 Centistoke Dimethicone	Skin Conditioner	2.5
Sepigel 305	Polyacrylamide, C ₁₃₋₂₄ Isoparrifin and Laureth-7	Thickener	3

**Retinol 50CJ is available from BASF of Mount Olive, NJ, and contains 50% by weight of retinol.

^Amount of NH₄OH required to reach pH of 4 is estimated.

5

The formulation was applied to skin, and the pH of the skin was measured over time. Results are illustrated in Figure 1.

10

Comparative Example 1B

A retinol/alpha-hydroxy acid containing composition was prepared as described in Example 1 above, except sodium hydroxide was substituted for the ammonium hydroxide.

The formulation was applied to skin, and the pH of the skin was measured over time. Results are illustrated in Figure 1.

Comparative Example 1C

An alpha-hydroxy acid containing composition having 8 percent by weight sodium glycolate at a pH of about 3.5 and no retinol was prepared as described in Example 1 above.

The formulation was applied to skin, and the pH of the skin was measured over time. Results are illustrated in Figure 1.

Example 2

A composition containing 0.15 percent by weight of retinol and 4 percent by weight of glycolic acid, neutralized with ammonium hydroxide to a pH of about 6, was prepared as described in Example 1 above.

An *in vivo* study of proliferative activity on skin was conducted. The marker of proliferative activity is an increase in fluorescent signal in the ultraviolet portion of the light spectrum. Over the course of 11
5 days of application, the fluorescence of the epidermis (exciting with 296 nm radiation, monitoring fluorescence at 340 nm) increases with increased proliferation activity. This fluorescence marker also increases after another proliferation inducing treatment such as tape-
10 stripping, and has been shown to correlate with increased cell turnover-rate as measured by increased loss of epidermal stain, dansyl chloride.

The slope of the increased fluorescence is illustrated in Figure 2.

15

Comparative Example 2A

An *in vivo* study as described in Example 2 was conducted using a preparation containing no glycolic acid or retinol at pH 6 (placebo).

20

The slope of the increased fluorescence is illustrated in Figure 2.

Comparative Example 2B

An *in vivo* study as described in Example 2 was conducted using a preparation containing 4 percent by weight of partially neutralized glycolic acid at pH 4 without retinol (Avon ANEW®).

The slope of the increased fluorescence is illustrated in Figure 2.

Comparative Example 2C

An *in vivo* study as described in Example 2 was conducted using a preparation containing 8 percent by weight of glycolic acid partially neutralized at pH 3.8 without retinol (Neutrogena HEALTHY SKIN®).

The slope of the increased fluorescence is illustrated in Figure 2.

Comparative Example 2D

An *in vivo* study as described in Example 2 was conducted on untreated skin.

The slope of the increased fluorescence is illustrated in Figure 2.

Figure 2 illustrates a significant increase in fluorescence activity and, therefore, cell proliferation in the retinol/glycolic acid preparation of Example 2 in

comparison with both a placebo (Example 2A) and untreated skin (Example 2D).

Figure 2 also illustrates a significant increase in fluorescence activity and, therefore, cell proliferation in the retinol/glycolic acid preparation of Example 2 which is similar to that of glycolic acid containing products having pH=s of about 4 (Comparative Examples 2B-D).

Example 3

A composition containing 0.15 percent by weight of retinol and 4 percent by weight of glycolic acid neutralized to pH 5.5 with ammonium hydroxide as in Example 1 was prepared.

Fluorescence was measured as in Example 2. Results are illustrated in Figure 3.

Comparative Example 3A

A composition containing 0.15 percent by weight of retinol and 4 percent by weight of glycolic acid neutralized to pH 5.5 with sodium hydroxide as in Example 1 was prepared.

Fluorescence was measured as in Example 2. Results are illustrated in Figure 3.

Comparative Example 3B

The fluorescence of untreated skin was measured as in Example 2. Results are illustrated in Figure 3.

5

Figure 3 illustrates that while ammonium glycolate (Example 3) dissociates when applied to the skin, sodium glycolate apparently does not (Comparative Example 3A). The latter results in little change in proliferative activity of the skin, and thus no apparent skin benefit.

10

Example 4

A composition prepared as in Example 1 was stored for 13 weeks at 40° C (simulating 2 years of ambient aging). This preparation retained 87% of the original retinol content after storage.

15

Comparative Example 4A

A composition prepared in Comparative Example 1A was stored for 13 weeks at 40° C (simulating 2 years of ambient aging). This preparation retained only 52% of the original retinol content after storage.

20

Example 5

A retinol/alpha-hydroxy acid containing composition having the formulation of Table 3 and containing paucilamellar vesicles was prepared as in Example 1 above. After the single addition components were added, a slurry of water and Cabopol ETD 2020 was added to the composition. Mirasil DM 100 and Phenoxetol were added thereto sequentially under mixing at 200 rpm for about 30 minutes. The formulation was allowed to cool to about 25E C under ambient conditions. The composition did not contain ammonium hydroxide.

Table 3

TRADE NAME	CHEMICAL NAME	FUNCTION	% WT/WT
<u>LIPID PHASE</u>			
Brij 76	Steareth-10	Emulsifier	1.4
Kessco GDS	Glyceryl Distearate	Emulsifier	2.8
Cholesterol NF	Cholesterol	Emulsifier	1
Procol ST 20G	Ceteareth-20 & Stearyl Alcohol	Emulsifier	3
Procol CS 20D	Cereareth-20 & Cetearyl Alcohol	Emulsifier	3
Lanol S	Stearyl Alcohol	Skin Conditioner	0.5
Wickenol 171	Octyl Hydroxystearate	Conditioning Agent	5.8014
BHT	BHT	Antioxidant	0.1
Tween 80	Polysorbate 80	Emulsifier	0.7
Retinol 50CJ**	Retinol in Polysorbate-20	Skin Conditioner	0.25

<u>AQUEOUS PHASE</u>			
Eau purifiee	Aqua	Vehicle	41.0843
Pricerin 9099	Glycerin	Humectant	4
Methylparaben	Methylparaben	Preservative	0.25
Propylparaben	Propylparaben	Preservative	0.15
Disodium EDTA	Disodium EDTA	Chelator	0.1
Lactose Rectapur	Lactose	Humectant	5
Glypure 70%	Glycolic acid (70%)	Skin Conditioner	5.7143
Sodium Hydroxide	Sodium Hydroxide	pH Adjuster	1.32
Eau purifiee	Aqua	Vehicle	20
Carbopol ETD 2020	Acrylates/ C10-30 Alkyl Acrylate crosspolymer	Thickener	0.6
<u>SINGLE ADDITION COMPONENTS</u>			
Mirasil DM 100	Dimethicone	Skin Conditioner	2.5
Phenoxytol	Phenoxyethanol	Preservative	0.73

**Retinol 50CJ is available from BASF of Mount Olive, NJ, and contains 50% by weight of retinol.

5

A control having the formulation of Table 3 was prepared excluding ammonium hydroxide and sodium hydroxide (Example 5A). The composition and control were applied to skin, and the pH of the skin was measured over time. Results are illustrated in Figure 4.

10

Example 6

A retinol/alpha-hydroxy acid containing composition having the formulation of Table 4 and a pH of about 5.8 was prepared as described in Example 5, except 3% by weight of ammonium hydroxide was substituted for the sodium hydroxide in Example 5.

Table 4

TRADE NAME	CHEMICAL NAME	FUNCTION	% WT/WT
Brij 76	Steareth-10	Emulsifier	1.4
<u>LIPID PHASE</u>			
Kessco GDS	Glyceryl Distearate	Emulsifier	2.8
Cholesterol NF	Cholesterol	Emulsifier	1
Procol ST 20G	Ceteareth-20 & Stearyl Alcohol	Emulsifier	3
Procol CS 20D	Cereareth-20 & Cetearyl Alcohol	Emulsifier	3
Lanol S	Stearyl Alcohol	Emulsifier	0.5
Wickenol 171	Octyl Hydroxystearate	Emulsifier	5.8014
BHT	BHT	Antioxidant	0.1
Tween 80	Polysorbate 80	Emulsifier	0.7
Retinol 50CJ**	Retinol in Polysorbate-20	Skin Conditioner	0.25
Eau purifiee	Aqua	Vehicle	39.4043
<u>AQUEOUS PHASE</u>			
Pricerin 9099	Glycerin	Humectant	4
Methylparaben	Methylparaben	Preservative	0.25
Propylparaben	Propylparaben	Preservative	0.15
Disodium EDTA	Disodium EDTA	Chelator	0.1

Lactose Rectapur	Lactose	Humectant	5
Glypure 70%	Glycolic acid (70%)	Skin Conditioner	5.7143
Ammonium Hydroxide	Ammonium Hydroxide (30%)	pH Adjuster	3
Eau purifiee	Aqua	Vehicle	20
Carbopol ETD 2020	Acrylates/ C10-30 Alkyl Acrylate crosspolymer	Thickener	0.6
<u>SINGLE ADDITION COMPONENTS</u>			
Mirasil DM 100	Dimethicone	Skin Conditioner	2.5
Phenoxetol	Phenoxyethanol	Preservative	0.73

**Retinol 50CJ is available from BASF of Mount Olive, NJ, and contains 50% by weight of retinol.

5 A control having the formulation of Table 4 was prepared excluding ammonium hydroxide (Example 6A). The composition and control were applied to skin, and the pH of the skin was measured over time. Results are illustrated in Figure 4.

10

Examples 7 and 8

Two retinol/alpha-hydroxy acid containing liposomal compositions having the formulations of Table 5 below are prepared as follows.

15

Table 5

TRADE NAME	CHEMICAL NAME	Function	Example 7 (%W/W)	Example 8 (% W/W)	Ranges
<u>LIPID PHASE</u>					
Glyceryl Dilaurate	Glyceryl Dilaurate	Nonionic Surfactant	2.8	2.8	1.4-8.4
Cholesterol	Cholesterol	Nonionic Surfactant	0.9	0.9	0.45-2.7
POE 10 Stearyl Alcohol	POE 10 Stearyl Alcohol	Nonionic Surfactant	2.5	2.5	1.25-7.5
Laureth-9	Laureth-9	Nonionic Surfactant	1.24	1.24	0.62-3.72
Butylated Hydroxytoluene (BHT)	BHT	Anti-oxidant	0.05	0.05	0-3
Retinol 50C™	Retinol in Polysorbate-20	Skin Conditioner	0.2	0.4	0.01-2
<u>AQUEOUS PHASE</u>					
Citric Acid	Citric Acid	Anti-oxidant	0.4	0.4	0.1-0.8
Trisodium Citrate dihydrate	Trisodium Citrate dihydrate	Buffer	0.6	0.6	0.1-0.8
Ascorbic Acid	Ascorbic Acid	Anti-oxidant	0.01	0.01	0.01-0.1
Glycerin	Glycerin	Humectant	0	4.0	0-20
Disodium EDTA	Disodium EDTA	Chelating Agent Preservative	0.2	0.2	0.01-0.2
Phenoxyethanol	Phenoxyethanol	Preservative	0.5	0.5	0.01-0.5
Methylparaben	Methylparaben	Preservative	0.25	0.25	0.01-0.2
Propylparaben	Propylparaben	Preservative	0.15	0.15	0.01-0.2
Glypure (70%)		Skin Conditioner	5.71	5.71	0.01-15
Ammonium Hydroxide (27 to 31%)	Ammonium Hydroxide (27 to 31%)	pH adjuster (pH=6)	3.2	3.2	0.01-10
Water	Water	Carrier	81.29	77.06	40-90

These compositions may be prepared by the following two methods.

5

1. Shear Mixing Method: Appropriate amounts of the lipid phase ingredients are mixed in a container heated to about 75° C until all the lipids have melted. The lipid melt is then cooled to about 65° C. The aqueous phase ingredients are mixed and heated to about 75° C to dissolve them and then cooled to about 60° C. The lipid melt and aqueous phase mixture are poured into separate holding reservoirs of a shear mixing apparatus for preparing liposomes as described in U.S. Patent No. 4,895,452. The positive displacement pump for the lipid and aqueous feed lines is turned on. The feed rate will depend on the desired viscosity of the composition. For a thinner consistency, a feed rate of 1 part lipid to 9 parts aqueous phase may be utilized. For thicker consistencies, a feed rate of 1 part lipid phase to 4 parts aqueous phase may be utilized. After the feed rate is adjusted, valves to the feed lines are opened and the aqueous phase mixture and lipid melt are fed through injection jets into a cylindrical mixing chamber tangentially with respect to the cylinder wall. In the

10

15

20

25

mixing chamber, the two streams of liquid intersect in such a manner as to cause shear mixing that causes the formation of liposomes. The liposomes are then withdrawn through an exit tube.

5

2. Syringe Method: Appropriate amounts of the lipid phase ingredients are mixed in a beaker at 75° C until the lipids melt. The lipid melt is drawn into a syringe, which was preheated in a water bath to about 75° C. A
10 second syringe containing appropriate amounts of the aqueous phase ingredients is preheated in a water bath to about 70° C. The two syringes were then connected via a 3-way metal stopcock. The ratio of aqueous phase mixture to lipid phase mixture was about 4:1 or 4 ml of aqueous
15 phase mixture to 1 ml of lipid phase mixture. The ratio of aqueous phase mixture to lipid phase mixture can be adjusted to obtain the desired viscosity. After injecting the aqueous phase mixture into the lipid phase mixture, the resulting mixture is rapidly mixed back and
20 forth between the two syringes several times until the contents cool to about 25-30° C.

Examples 9 and 10

Two oil-in-water emulsions of the present invention are shown in Table 6.

Table 6

TRADE NAME	CHEMICAL NAME	Function	Example 9 (%W/W)	Example 10 (%W/W)	Ranges
<u>OIL PHASE</u>					
Cetearyl Glucoside	Cetearyl Glucoside	Surfactant	1.4	1.4	0.1-2.8
C12-15 Alkyl Benzoate	C12-15 Alkyl Benzoate	Surfactant	4.0	4.0	1-6
Octyl Hydroxystearate	Octyl Hydroxystearate	Emollient	1.0	1.0	0-5
Dimethicone	Dimethicone	Spreading Agent	1.0	1.0	0-5
Cyclomethicone	Cyclomethicone	Spreading Agent	1.0	1.0	0-5
Cetyl Alcohol	Cetyl Alcohol	Emollient	2.5	2.5	0-4
Butylated Hydroxytoluene	BHT	Anti-oxidant	0.05	0.05	0-3
Octyl Methoxycinnamate	Octyl Methoxycinnamate	Sunscreen	6.0	6.0	0-10
Propylparaben	Propylparaben	Preservative	0.5	0.1	0-0.5
Vitamin E acetate	Vitamin E acetate	Anti-oxidant	0.5	0.5	0-0.5
Retinol	Retinol	Anti-Wrinkle	0.25	0.4	0.01-5
Tocopherol Acetate	Tocopherol Acetate	Anti-oxidant	0.5	0.5	0-0.5
<u>AQUEOUS PHASE</u>					
Glycerin	Glycerin	Humectant	3.0	3.0	0-20
D-Pathenol	D-Pathenol	Pro-Vitamin	0.5	0.5	0-5
Disodium EDTA	Disodium EDTA	Chelator, whitening agent	0.1	0.1	0.01-1
Methyl Paraben	Methyl Paraben	Preservative	0.2	0.2	0-0.3
Carbomer		Thickener	0.35	0.35	0-3

Glycolic acid (70%)	Glycolic acid (70%)	Skin Conditioner	5.71	5.71	0-15
Ammonium Hydroxide	Ammonium Hydroxide	pH adjuster	3.2	3.2	0-1
Deionized Water	Deionized Water	Carrier	68.19	68.04	50-80

Each emulsion is prepared by mixing the oil phase ingredients and heating the mixture to about 85° C.

The oil phase mixture is then cooled to about 60° C.

5 In a separate vessel, the carbomer is slowly added to the water. After mixing for about 10 minutes the remaining aqueous phase ingredients are added and the mix is heated to about 60° C.

10 The two phases are then combined, mixed for about 10 minutes, and cooled to room temperature. One or more depigmentation agents may be added to the formulations in these examples.

Examples 11 and 12

15 Two water-in-oil emulsions of the present invention are shown in Table 7.

Table 7

TRADE NAME	CHEMICAL NAME	Function	Example 11 (%W/W)	Example 12 (%W/W)	Preferred Ranges
<u>OIL PHASE</u>					
Mineral Oil	Mineral Oil	Emollient	25.0	25.0	40-80
Sorbitan Monooleate	Sorbitan Monooleate	Surfactant	5.0	5.0	1-6

Stearyl Alcohol	Stearyl Alcohol	Emollient	25.0	25.0	20-60
Dimethicone	Dimethicone	Spreading Agent	1.0	1.0	1-5
Cetyl Alcohol	Cetyl Alcohol	Emollient	2.0	2.0	0.1-10
Hydrogenated Lecithin	Hydrogenated Lecithin	Anti-oxidant	3.0	3.0	0-10
Parsol MCX		Sunscreen	3.0	3.0	0-10
Butylated Hydroxytoluene	BHT	Anti-oxidant	0.05	0.05	0-3
Retinol	Retinol	Anti-Wrinkle	0.25	0.4	0.01-5
Propylparaben	Propylparaben	Preservative	0.5	0.5	0.01-0.5
Vitamin E acetate	Vitamin E acetate	Anti-oxidant	0.5	0.5	0.01-0.5
<u>AQUEOUS PHASE</u>					
Glycerin	Glycerin	Humectant	3.0	3.0	0-20
Methyl Paraben	Methyl Paraben	Preservative	0.2	0.2	0.01-0.3
Water	Water	Carrier	22.59	22.44	20-45
Glycolic acid (70%)	Glycolic acid (70%)	Skin Conditioner	5.71	5.71	0-15
Ammonium Hydroxide	Ammonium Hydroxide	pH adjuster	3.2	3.2	0-1

Each emulsion is prepared by melting stearyl alcohol and mineral oil at about 70° C. The other oil phase ingredients are added and the mixture is heated to about 75° C. The aqueous phase ingredients are dissolved in water and warmed to about 70° C. The aqueous phase mixture is added to the oil phase mixture. The resulting mixture is stirred until it congeals.

Example 13

An oil-in-water emulsion of the present invention is shown in Table 8.

The emulsion is prepared by mixing the oil phase ingredients and heating the mixture to about 85° C. The oil phase mixture is then cooled to about 60° C. In a separate vessel, the water phase ingredients are added, mixed and heated to about 60° C.

The two phases are then combined, mixed for about 10 minutes, and cooled to about 35° C, at which time the post additions are added and mixed, followed by the addition and mixing of the glycolic acid/malic acid/deionized water buffer pre-mix. The retinol 50C is then added and mixed last.

Table 8

CTFA Name	Trade Name	FUNCTION	Wt. %
<u>OIL PHASE</u>			
C12-15 Alkyl Benzoate	Finsolv TN	Solubilizing Agent	4.00
Octyl Hydroxystearate	Wickenol 171	emolient	1.00
Dimethicone, 100 centistoke	Dimethicone 47V-100	emolient	1.00
Steareth 2	Brij 72	emulsifier	0.60
Cetyl Alcohol	Cetal	emolient, emulsifier	2.50
Steareth 20	Brij 721	emulsifier	0.90

BHT	BHT	antioxidant for cosmetics	0.10
Pemulen TR1	Acrylates/C10-30 Alkyl Acrylate Crosspolymer	thickener/emulsifier	0.50
<u>WATER PHASE</u>			
Deionized Water	Water	Solvent	62.59
Disodium EDTA	EDTA	Chelating Agent	0.10
Glycerin	Glycerin 916 99.7% USP	humectant, emollient	3.00
Panthenol	D-Panthenol U.S.P. FCC	moisturizing agent	0.50
Phenoxyethanol	Emeressence 1160	perservative	0.73
Methylparaben	Methylparaben	preservative	0.35
Propylparaben	Propylparaben	preservative	0.17
<u>POST ADDITIONS</u>			
<i>DMAE/Tyrosine Pre-mix</i> L-Tyrosine	L-Tyrosine	active	0.50
Deionized Water DMAE	Deionized Water 2-(dimethylamino)-ethanol	solvent active	15.00 3.00
<u>BUFFER PRE-MIX</u>			
Glycolic Acid Malic Acid Deionized Water	Glypure 70 Malic Acid Deionized Water	buffer buffer Solvent	1.20 0.84 1.32
Retinol 50C	Vitamin A alcohol in Polysorbate 80	vitamin A	0.10

All patents, publications, applications, and test methods mentioned herein are hereby incorporated by reference.

Many variations of the present invention will suggest themselves to those skilled in the art in light of the above, detailed description. All such obvious variations are within the full intended scope of the appended claims.

5

Claims:

1. A composition comprising:

(A) a retinoid;

5 (B) a dermatologically active acid; and

(C) 2-dimethylaminoethanol.

2. A composition as defined in claim 1, wherein
said retinoid is retinol.

10

3. A composition as defined in claim 2, wherein
said dermatologically active acid is selected from the
group consisting of a hydroxy acid, ascorbic acid and
derivatives thereof, lipoic acid, dihydrolipoic acid, or
15 a combination thereof.

4. A composition as defined in claim 3,
wherein said hydroxy acid is an alpha-hydroxy acid.

20

5. A composition as defined in claim 4,
wherein said alpha-hydroxy acid is selected from the
group consisting of malic acid, tartaric acid, lactic
acid, pyruvic acid, citric acid, or any combination of
any of the foregoing.

6. A composition as defined in claim 5,
wherein said alpha-hydroxy acid is glycolic acid.

5 7. A composition as defined in claim 2,
wherein said retinol comprises from about 0.01 to about
10 percent by weight, based upon 100 percent by weight of
total composition.

10 8. A composition as defined in claim 7,
wherein the amount of said acid ranges from about 0.1 to
about 20 percent by weight, based upon 100 percent by
weight of total composition.

15 9. A composition as defined in claim 7, said
composition comprises from about 0.01 to about 10 percent
by weight of retinol and from about 0.1 to about 20
percent by weight of said glycolic acid, based upon 100
percent of total composition.

20 10. A composition as defined in claim 2,
wherein the amount of 2-dimethylaminoethanol is effective
to neutralize said composition to a pH ranging from about
4.5 to about 8.

11. A composition as defined in claim 10,
wherein the amount of 2-dimethylaminoethanol is
sufficient to neutralize said composition to a pH ranging
5 from about 5 to about 6.

12. A composition as defined in claim 2,
wherein said retinol is within a paucilamellar vesicle.

10 13. A composition as defined in claim 2,
further comprising a second neutralizing agent.

14. A composition as defined in claim 13,
wherein said second neutralizing agent comprises an
15 alkali hydroxide, alkanolamine, amino acid, or any
combination of any of the foregoing.

15. A composition as defined in claim 14,
wherein said second neutralizing agent comprises sodium
20 hydroxide, potassium hydroxide, diethanolamine,
triethanolamine, aminobutanol, arginine, lysine, or any
combination of any of the foregoing.

16. A composition comprising:

(A) retinoid; and

(B) an N,N-dimethylethanolammonium salt of a dermatologically active acid.

5 17. A composition as defined in claim 16,
wherein said retinoid is retinol.

10 18. A composition as defined in claim 17,
wherein said dermatologically active acid is selected
from the group consisting of a hydroxy acid, ascorbic
acid and derivatives thereof, lipoic acid, dihydrolipoic
acid, or a combination thereof.

15 19. A composition as defined in claim 18,
wherein said hydroxy acid is an alpha-hydroxy acid.

20 20. A composition as defined in claim 19,
wherein said alpha-hydroxy acid is selected from the
group consisting of malic acid, tartaric acid, lactic
acid, pyruvic acid, citric acid, or any combination of
any of the foregoing.

21. A composition as defined in claim 16,
wherein said N,N-dimethylethanolammonium salt is N,N-
dimethylethanolammonium glycolate.

5 22. A method for reducing fine lines,
wrinkles, skin roughness, and pore size and for
increasing the clarity of a skin surface, cellular
turnover, skin radiance, skin smoothness, skin
permeation, or collagen synthesis in a mammal in need
10 thereof, said method comprising topically administering a
composition as defined in claim 1 to said mammal.

15 23. A method for reducing fine lines,
wrinkles, skin roughness, and pore size and for
increasing the clarity of a skin surface, cellular
turnover, skin radiance, skin smoothness, skin
permeation, or collagen synthesis in a mammal in need
thereof, said method comprising topically administering a
composition as defined in claim 16 to said mammal.

Figure 1




Figure 2




Figure 3




Figure 4

