



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/08/09
(87) Date publication PCT/PCT Publication Date: 2019/02/14
(85) Entrée phase nationale/National Entry: 2020/02/07
(86) N° demande PCT/PCT Application No.: US 2018/046061
(87) N° publication PCT/PCT Publication No.: 2019/032863
(30) Priorité/Priority: 2017/08/11 (US62/544,083)

(51) Cl.Int./Int.Cl. *C07D 405/14* (2006.01),
A61K 31/4365 (2006.01), *A61K 31/496* (2006.01),
A61K 31/4995 (2006.01), *A61K 31/5386* (2006.01),
A61K 31/551 (2006.01), *A61P 35/00* (2006.01),
C07D 471/04 (2006.01), *C07D 487/04* (2006.01),
C07D 495/04 (2006.01), *C07D 519/00* (2006.01)

(71) Demandeur/Applicant:
FORMA THERAPEUTICS, INC., US

(72) Inventeurs/Inventors:
ZABLOCKI, MARY-MARGARET, US;
GUERIN, DAVID J., US;
NG, PUI YEE, US;
WANG, ZHONGGUO, US;

(54) Titre : CARBOXAMIDES UTILISEES EN TANT QU'INHIBITEURS DE PROTEASE SPECIFIQUE DE L'UBIQUITINE
(54) Title: CARBOXAMIDES AS UBIQUITIN-SPECIFIC PROTEASE INHIBITORS

(57) **Abrégé/Abstract:**

The present disclosure relates to modulators, such as inhibitors, of at least one pathway chosen from USP28 and USP25, pharmaceutical compositions comprising the inhibitors, and methods of using the inhibitors. The modulators, such as inhibitors, of at least one pathway chosen from USP28 and USP25 can be useful in the treatment of cancers, among other ailments.

(72) **Inventeurs(suite)/Inventors(continued):** SHELEKHIN, TATIANA, US; CARAVELLA, JUSTIN, US; LI, HONGBIN, US; IOANNIDIS, STEPHANOS, US

(74) **Agent:** BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
14 February 2019 (14.02.2019)



(10) International Publication Number
WO 2019/032863 A1

(51) International Patent Classification:

C07D 405/14 (2006.01) A61K 31/496 (2006.01)
C07D 471/04 (2006.01) A61K 31/551 (2006.01)
C07D 487/04 (2006.01) A61K 31/4995 (2006.01)
C07D 495/04 (2006.01) A61K 31/5386 (2006.01)
C07D 519/00 (2006.01) A61K 31/4365 (2006.01)
A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2018/046061

(22) International Filing Date:

09 August 2018 (09.08.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/544,083 11 August 2017 (11.08.2017) US

(71) Applicant: **FORMA THERAPEUTICS, INC.** [US/US];
500 Arsenal Street, Suite 100, Watertown, Massachusetts
02472 (US).

(72) Inventor; and

(71) Applicant: **ZABLOCKI, Mary-Margaret** [US/US]; 500
Arsenal Street, Suite 100, Watertown, Massachusetts 02472
(US).

(72) Inventors: **GUERIN, David J.**; 500 Arsenal Street, Suite
100, Watertown, Massachusetts 02472 (US). **NG, Pui
Yee**; 500 Arsenal Street, Suite 100, Watertown, Massa-
chusetts 02472 (US). **WANG, Zhongguo**; 500 Arsenal
Street, Suite 100, Watertown, Massachusetts 02472 (US).
SHELEKHIN, Tatiana; 500 Arsenal Street, Suite 100,
Watertown, Massachusetts 02472 (US). **CARAVELLA,
Justin**; 500 Arsenal Street, Suite 100, Watertown, Mass-
achusetts 02472 (US). **LI, Hongbin**; 500 Arsenal Street,
Suite 100, Watertown, Massachusetts 02472 (US). **IOAN-
NIDIS, Stephanos**; 500 Arsenal Street, Suite 100, Water-
town, Massachusetts 02472 (US).

(74) Agent: **HERZFELD, Deborah M.** et al.; 125 Cambridge
Park Drive, Suite 301, Cambridge, Massachusetts 02140
(US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,

OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with amended claims (Art. 19(1))

(54) Title: CARBOXAMIDES AS UBIQUITIN-SPECIFIC PROTEASE INHIBITORS

(57) Abstract: The present disclosure relates to modulators, such as inhibitors, of at least one pathway chosen from USP28 and USP25, pharmaceutical compositions comprising the inhibitors, and methods of using the inhibitors. The modulators, such as inhibitors, of at least one pathway chosen from USP28 and USP25 can be useful in the treatment of cancers, among other ailments.

WO 2019/032863 A1

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 326

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 326

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

CARBOXAMIDES AS UBIQUITIN-SPECIFIC PROTEASE INHIBITORS

TECHNICAL FIELD

[0001] The present disclosure is directed to modulators of at least one pathway chosen from ubiquitin-specific protease 28 (USP28) and/or ubiquitin-specific protease 25 (USP25) useful in the treatment of diseases or disorders associated with at least one pathway chosen from USP28 and USP25 enzymes. Specifically, the disclosure is concerned with chemical entities and compositions inhibiting at least one pathway chosen from USP28 and USP25, methods of treating diseases or disorders associated with at least one pathway chosen from USP28 and USP25, and methods of synthesis of these compounds.

BACKGROUND

[0002] USP28 and USP25 are cysteine isopeptidases of the USP sub-family of DUBs containing three distinct domains: an N-terminal UBA-like domain; a pair of ubiquitin-interacting motifs (UIM) and a USP domain that is predicted to have the conserved fold of the USP sub-family (Nijman et al., *Cell* **2005**, 123, 773-786; Komander et al., *Mol. Cell Bio.* **2009**, 10, 550-563). USP28 and USP25 exert their function through regulating the stability of a plethora of cellular proteins. USP28 has been characterized as a tumor-promoting factor and has been found to stabilize many oncoproteins. USP25 has been characterized as a tumor-promoting factor and as a regulator of cellular responses related to autoimmune disease, inflammation, and infectious diseases (such as viruses and bacteria).

[0003] Amplification, deletions and mutations of USP28 have been identified in multiple cancer types, including breast cancer, AML, ovarian cancer, and colorectal cancer. (cbioportal; <http://www.cbioportal.org>; Diefenbacher et al., *J. of Clin. Investi.* **2014**, 124, 3407-3418; Popov et al., *Nat. Cell. Biol.* **2007**, 9, 729-731). Furthermore, USP28 overexpression has been correlated with poor prognosis in patients with glioblastoma, non-small cell lung carcinoma and bladder cancers suggesting that USP28 plays an important role in tumorigenesis of these tumor types. (Wang et al. *Exp. Biol. Med.* **2016**, 255-264; Zhang et al. *J. Cell. Mol. Med.* **2015**, 19, 799-805; Guo et al., *Tumor Bio.* **2014**, 35, 4017-4022).

[0004] A large-scale shRNA screen has also identified a role of USP28 in the control of the stability of MYC protein. (Popov, *Nat. Cell. Biol.*, 765-774). MYC is a master regulator of the transcription of genes involved in cell growth, proliferation and apoptosis and is essential for tumor initiation and maintenance in many tumor types. (Meyer et al., *Nat. Rev. Cancer* **2008**, 8, 976-990; Conacci-Sorrell et al., *Cold Spring Harb. Perspect. Med.* **2014**, 4, 1-24; Huang et al., *Cold Spring Harb. Perspect. Med.* **2013**; Roussel et al., *Cold Spring Harb. Perspect. Med.* **2013**; Gabay et al., *Cold Spring Harb. Perspect. Med.* **2014**; Schmitz et al., *Cold Spring Harb. Perspect. Med.* **2014**). In addition, MYC is the most frequently amplified oncogene in human cancer, with alterations in many tumor types including breast, lung and prostate. (Beroukhi et al., *Nature* **2010**, 463, 899-905). Knockdown of the USP28 gene has been shown to lead to a decrease of MYC protein and an associated inhibition of growth in a panel of human cancer cell lines in vitro. (Popov, *Nat. Cell Biol.*, 765-774).

[0005] USP28 has also been reported to be required to impart stability on the LSD1 (lysine-specific demethylase 1) protein. (Wu et al., *Cell Rep.* **2013**, 5, 224-236). LSD1 is a histone demethylase that complexes with many partner proteins to control cellular pluripotency and differentiation. (Metzger et al. *Nature* **2005**, 437, 436-439; Toffolo et al, *J. Neurochem.* **2014** 128, 603-616, 2014; Periz et al., *PloS Biology* **2015**). Knockdown of USP28 in tumor cells has been shown to lead to the destabilization of LSD1 protein, the suppression of cancer stem cell (CSC)-like characteristics in vitro, and the inhibition of tumor growth in vivo. (Wu, *Cell Rep.*, 224-236). Small molecule inhibitors of LSD1 have shown antitumor activity in models of AML and Ewing sarcoma. (Sankar et al., “Reversible LSD1 inhibition interferes with global EWS/ETS transcriptional activity and impedes Ewing sarcoma tumor growth” *Clin Cancer Res.* **2014** 4584-4597; Schenk et al., *Nat. Med.* **2012**, 18, 605-611). Thus, USP28 inhibition represents an alternate approach to targeting LSD1 in these tumor types.

[0006] USP28 inhibition has also been shown to reduce NICD1-Levels and to lead to inhibition of the NOTCH pathway activity. (Diefenbacher et al.). NOTCH signaling controls diverse cellular differentiation decisions and drives tumorigenesis in certain tumor types. NOTCH1 is a potent T-cell oncogene, with >50% of T-cell acute lymphoblastic leukemia (T-ALL) cases carrying activating mutations in NOTCH1. (Weng et al. *Science* **2004**, 306, 269-271). Increased NOTCH1 protein levels have also been associated with disease progression in colon

cancer. (Meng et al., *Cancer Res.* **2009**, *69*, 573-582). NOTCH1 rearrangements lead to constitutive pathway activation and drive tumorigenesis in many cancer types, including triple-negative breast cancer. (Stoeck et al., *Cancer Discov.* **2014**, *4*, 1154-1167).


[0007] Other reported substrates of USP28 include c-Jun, Cyclin E, HIF-1 α , Claspin, 53BP1, and Mdc1, many of which play important roles in tumorigenesis in humans. (Diefenbacher et al.; Flügel et al. *Blood* **2012**, *119*, 1292-1301; Zhang et al., “A role for the deubiquitinating enzyme USP28 in control of the DNA-damage response” *Cell* **2006**, *126*, 529-542). Interestingly, many USP28 substrates are recognized by FBW7, the substrate recognition subunit of SCF (FBW7) E3 ubiquitin ligase. (Diefenbacher et al.). FBW7 recognizes USP28 substrates in a phosphorylation-dependent manner and targets them for ubiquitination ultimately leading to their proteasomal degradation. The antagonizing roles of USP28 and FBW7 on their shared oncoprotein substrates indicate the intricate nature of protein stability control and may provide additional therapeutic opportunities for cancer treatment.

[0008] Mice with a germline knockout of USP28 have been shown to be viable and fertile, confirming that USP28 activity is not required for normal development and reproductive function. (Knobel et al., *Molecular and Cellular Biology* **2014**, *34*, 2062-2074). Conditional knockout of USP28 in mouse intestine led to the reduction of oncoproteins including c-Myc, active NOTCH (NICD1) and c-JUN which was associated with decreased intestinal cell proliferation and enhanced differentiation. More importantly, intestinal tumorigenesis induced by APC mutation was effectively blocked with acute USP28 depletion suggesting that USP28 could be an appealing target to reduce tumor burden and improve survival for intestinal cancers. (Diefenbacher et al.).

[0009] In summary, USP28 and USP25 play important roles in promoting tumorigenesis in cells and modulating immune responses. Its major role being in the deubiquitination and stabilization of diverse oncoproteins and epigenetic drivers and immunomodulatory proteins among other cellular factors, which are necessary for immune responses and tumor initiation and growth in humans. Inhibition of USP28 and/or USP25 with small molecule inhibitors therefore can be developed for medical use, such as for the treatment for cancer such as lung cancer. For this reason, there remains a considerable need for novel and potent small molecule inhibitors of USP28 and/or USP25.

SUMMARY

[0010] The present disclosure relates to chemical entities chosen from compounds of Formula (I)



and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, wherein:

X is chosen from C(R)₂ and O;

Y is chosen from C and N;

R is chosen from H, halogens, C₁-C₆ alkyl, -OH, and -CN, wherein any R group containing hydrogen can have one or more hydrogen replaced with deuterium;

R₁ is chosen from 6-12 membered heteroaryls optionally substituted with one or more substituent chosen from R₅ and R₆, wherein a 6-membered nonfused heteroaryl is substituted with one or more R₆, and further wherein any R₁ group containing hydrogen can have one or more hydrogen replaced with deuterium;

R₂ is chosen from N-linked and C-linked 4-12 membered heterocyclyls, and an O linked to a heterocyclyl, wherein the heterocyclyls are optionally substituted with one or more R₅, a sulfur member of the heterocyclyls can be S(O) or S(O)₂, and further wherein any R₂ group containing hydrogen can have one or more hydrogen replaced with deuterium;

R₃ is independently chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH, and wherein the cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₇;

R₄ is chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein alkyls are optionally substituted with one or more substituent independently

chosen from (C₁-C₆) alkoxy and -OH, and wherein the cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₅, and further wherein any R₄ group containing hydrogen can have one or more hydrogen replaced with deuterium;

R₅ is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH, and wherein any R₅ group containing hydrogen can have one or more hydrogen replaced with deuterium;

R₆ is chosen from -N(C₁-C₆)alkyl-aryls, -N(C₁-C₆)alkyl-heteroaryls, and -N(C₁-C₆)alkyl-heterocyclyl groups, wherein the groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups, and further wherein any R₆ group containing hydrogen can have one or more hydrogen replaced with deuterium;

R₇ is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH;
and

n is 0, 1, 2, or 3.

[0011] The compound of Formula (I) can be a compound, or pharmaceutically acceptable salt thereof, wherein:

X is chosen from C(R)₂ (e.g., C(R)(R) where each R is the same or different) and O;

Y is chosen from C (e.g., CH or C(R) where R is the same or different from R in moiety X) and N;

R is independently chosen from H, halogens, C₁-C₆ alkyl, -OH, and -CN;

R₁ is chosen from 6-12 membered heteroaryls optionally substituted with one or more substituent chosen from R₅ and R₆, wherein a 6-membered nonfused heteroaryl is substituted with one or more R₆;

R₂ is chosen from N-linked 4-12 membered heterocyclyls, C-linked 4-12 membered heterocyclyls, and O-linker to a heterocyclyl, wherein the N-linked, C-linked or O-linker heterocyclyls are optionally substituted with one or more R₅;

R₃ is independently chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH, and wherein the cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₇;

R₄ is chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH, and wherein the cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₅;

R₅ is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH;

R₆ is chosen from -N-(C₁-C₆)alkyl-aryls, -N-(C₁-C₆)alkyl-heteroaryls, and -N-(C₁-C₆)alkyl-heterocyclyl groups, wherein the groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups;

R₇ is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH; and

n is 0, 1, 2, or 3.

[0012] In other examples, the compound of Formula (I) can be a compound, or pharmaceutically acceptable salt thereof, wherein:

X is chosen from C(R)₂ (e.g., C(R)(R) where each R is the same or different) and O;

Y is chosen from C (e.g., CH or C(R) where R is the same or different from R in moiety X) and N;

each R in X and Y is independently chosen from H, halogens, C₁-C₆ alkyl, -OH, and -CN;

R₁ is chosen from heteroaryls optionally substituted with one or more substituent chosen from R₅ and R₆, wherein the heteroaryl is optionally substituted with one or more R₆;

R₂ is chosen from N-linked-heterocyclyls, C-linked-heterocyclyls, and O-linker-heterocyclyls, wherein the N-linked, C-linked or O-linker heterocyclyls are optionally substituted with one or more R₅;

each R₃ is independently chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein the alkyls are each optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH, and wherein the cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₇;

each R₄ is independently chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein alkyls are each optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH, and wherein the cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₅;

each R₅ is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein the alkyls are each optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH;

each R₆ is independently chosen from amino-alkyl-aryls (e.g., -N-(C₁-C₆)alkyl-aryls), amino-alkyl-heteroaryls (e.g., -N-(C₁-C₆)alkyl-heteroaryls), amino-alkyl-cyclyl and amino-alkyl-heterocyclyl (e.g., -N-(C₁-C₆)alkyl-heterocyclyls) groups, wherein these groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups;

each R₇ is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein the alkyls are each optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH; and

n is 0, 1, 2, or 3.

[0013] Another aspect of the present disclosure relates to a method of treating a disease or disorder associated with inhibition of USP28. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with inhibition of USP28 an effective amount of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0014] Another aspect of the disclosure relates to a method of treating a disease or disorder associated with inhibition of USP25. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with inhibition of USP25 an effective amount of a at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0015] Another aspect of the disclosure relates to a method of treating a disease or disorder associated with inhibition of at least one pathway chosen from USP28 and USP25. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with inhibition of at least one pathway chosen from USP28 and USP25 effective amount of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0016] Another aspect of the disclosure is directed to a method of inhibiting USP28. The method involves administering to a patient in need thereof an effective amount of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, hydrates, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0017] Another aspect of the disclosure is directed to a method of inhibiting USP25. The method involves administering to a patient in need thereof an effective amount of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, hydrates, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0018] Another aspect of the disclosure is directed to a method of inhibiting at least one pathway chosen from USP28 and USP25. The method involves administering to a patient in need thereof an effective amount of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0019] Another aspect of the disclosure relates to a method of treating cancer. The method comprises administering to a patient in need thereof an effective amount of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0020] Another aspect of the disclosure is directed to pharmaceutical compositions comprising at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, and a pharmaceutically acceptable carrier. The pharmaceutical acceptable carrier may further include an excipient, diluent, or surfactant.

[0021] Another aspect of the present disclosure relates to at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, for use in the manufacture of a medicament for treating a disease associated with inhibiting USP28.

[0022] Another aspect of the present disclosure relates to the use of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, in the treatment of a disease associated with inhibiting USP28.

[0023] Another aspect of the present disclosure relates to the use of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, in the treatment of a disease associated with inhibiting USP25.

[0024] Another aspect of the present disclosure relates to the use of at least one chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, in the treatment of a disease associated with inhibiting at least one pathway chosen from USP28 and USP25.

DESCRIPTION OF THE EMBODIMENTS

[0025] Compounds useful for inhibiting USP 28 and/or USP25 are disclosed herein, including USP25 Inhibitor compounds, USP28 Inhibitor compounds and USP28/25 Inhibitor compounds as defined herein. The USP28/25 Inhibitor, USP28 Inhibitor and/or USP25 Inhibitor

compounds can be a compound disclosed herein, including a compound of Formula (I), a compound of Formula (II), a compound of Formula (III), a compound of Formula (IV), a compound of Formula (V), a compound of Formula (VI), and/or a compound of Formula (VII).

[0026] The term “USP28 Inhibitor” compound as used herein refers to a compound disclosed herein (e.g., a compound of Formula (I), a compound of Formula (II), a compound of Formula (III), a compound of Formula (IV), a compound of Formula (V), a compound of Formula (VI), and/or a compound of Formula (VII)) having an IC_{50} of 2 micromolar or less in the Ubiquitin-Rhodamine 110 Assay for USP28 as described in Example A-1(a) and/or the Ubiquitin-Rhodamine 110 Assay for USP28 as described in Example A-1(b) herein. For example, the USP28 Inhibitor can be a compound of a formula disclosed herein having an IC_{50} value of up to 2 micromolar using the Ubiquitin-Rhodamine 110 Assay for USP28 as described in Example A-1(a), including IC_{50} values ranging from 0.001 – 2 micromolar, preferably 0.001-0.2 micromolar, and more preferably 0.001-0.05 micromolar. The USP28 Inhibitor can be a compound of a formula disclosed herein having an IC_{50} value of up to 2 micromolar using the Ubiquitin-Rhodamine 110 Assay for USP28 as described in Example A-1(b), including IC_{50} values ranging from 0.001 – 2 micromolar, preferably 0.001-0.2 micromolar, more preferably from 0.001-0.05 micromolar. The USP28 Inhibitor can be a compound of a formula disclosed herein having an IC_{50} values of up to 2 micromolar using both the Ubiquitin-Rhodamine 110 Assay for USP28 as described in Example A-1(a) and the Ubiquitin-Rhodamine 110 Assay for USP28 as described in Example A-1(b), including IC_{50} values of 0.001 – 2 micromolar, preferably 0.001-0.2 micromolar, more preferably 0.001-0.05 micromolar for both assays.


[0027] The term “USP25 Inhibitor” as used herein refers to a compound disclosed herein (e.g., a compound of Formula (I), a compound of Formula (II), a compound of Formula (III), a compound of Formula (IV), a compound of Formula (V), a compound of Formula (VI), and/or a compound of Formula (VII)) having an IC_{50} of 2 micromolar or less in the Ubiquitin-Rhodamine 110 Assay for USP25 as described in Example A-2 herein. For example, the USP25 Inhibitor can be a compound of a formula disclosed herein having an IC_{50} value of up to 2 micromolar using the Ubiquitin-Rhodamine 110 Assay for USP25 as described in Example A-2, including IC_{50} values ranging from 0.001 – 2 micromolar, preferably 0.001-0.2 micromolar, more preferably 0.001-0.05 micromolar. The USP25 Inhibitor can be a compound of a formula disclosed herein having an

IC₅₀ value of up to 2 micromolar using the Ubiquitin-Rhodamine 110 Assay for USP25 as described in Example A-2, including IC₅₀ values ranging from 0.001 – 2 micromolar, preferably 0.001-0.2 micromolar, more preferably 0.001-0.05 micromolar. The USP25 Inhibitor can be a compound of a formula disclosed herein having an IC₅₀ values of up to 2 micromolar using both the Ubiquitin-Rhodamine 110 Assay for USP25 as described in Example A-2 and the Ubiquitin-Rhodamine 110 Assay for USP25 as described in Example A-2, including IC₅₀ values ranging from 0.001 – 2 micromolar, preferably 0.001-0.2 micromolar, more preferably 0.001-0.05 micromolar for both assays.

[0028] The term “USP28/25 Inhibitor” as used herein refers to a compound disclosed herein (e.g., a compound of Formula (I), a compound of Formula (II), a compound of Formula (III), a compound of Formula (IV), a compound of Formula (V), a compound of Formula (VI), and/or a compound of Formula (VII)) that is a USP28 Inhibitor or a USP25 Inhibitor or both a USP28 Inhibitor and USP25 Inhibitor, as defined herein.

[0029] Optionally, any one or more hydrogen atoms in a compound of Formula (I), Formula (II), a compound of Formula (III), a compound of Formula (IV), a compound of Formula (V), a compound of Formula (VI), and/or a compound of Formula (VII)) can independently be replaced with deuterium or other hydrogen isotope.

[0030] In a first aspect of the disclosure, the chemical entities are chosen from compounds of Formula (I):



and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, are described wherein X, Y, R₁, R₂, R₃, R₄, and n are as described herein above.

[0031] The details of the disclosure are set forth in the accompanying description below. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, illustrative methods and materials are now described.

Other features, objects, and advantages of the disclosure will be apparent from the description and from the claims. In the specification and the appended claims, the singular forms also include the plural unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. All patents and publications cited in this specification are incorporated herein by reference in their entireties.

[0032] The articles "a" and "an" are used in this disclosure to refer to one or more than one (*e.g.*, to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

[0033] The term "and/or" is used in this disclosure to mean either "and" or "or" unless indicated otherwise.

[0034] The term "optionally substituted" is understood to mean that a given chemical moiety (*e.g.*, an alkyl group) can (but is not required to) be bonded other substituents (*e.g.*, heteroatoms). For instance, an alkyl group that is optionally substituted can be a fully saturated alkyl chain (*e.g.*, a pure hydrocarbon). Alternatively, the same optionally substituted alkyl group can have substituents different from hydrogen. For instance, it can, at any point along the chain be bounded to a halogen atom, a hydroxyl group, or any other substituent described herein. Thus the term "optionally substituted" means that a given chemical moiety has the potential to contain other functional groups, but does not necessarily have any further functional groups. Suitable substituents used in the optional substitution of the described groups include, without limitation, halogen, oxo, -OH, -CN, -COOH, -CH₂CN, -O-(C₁-C₆) alkyl, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, -O-(C₂-C₆) alkenyl, -O-(C₂-C₆) alkynyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -OH, -OP(O)(OH)₂, -OC(O)(C₁-C₆) alkyl, -C(O)(C₁-C₆) alkyl, -OC(O)O(C₁-C₆) alkyl, -NH₂, -NH((C₁-C₆) alkyl), -N((C₁-C₆) alkyl)₂, -NHC(O)(C₁-C₆) alkyl, -C(O)NH(C₁-C₆) alkyl, -S(O)₂(C₁-C₆) alkyl, -S(O)NH(C₁-C₆) alkyl, and S(O)N((C₁-C₆) alkyl)₂. The substituents can themselves be optionally substituted. "Optionally substituted" as used herein also refers to substituted or unsubstituted whose meaning is described below.

[0035] As used herein, the term "substituted" means that the specified group or moiety bears one or more suitable substituents wherein the substituents may connect to the specified group or moiety at one or more positions. For example, an aryl substituted with a cycloalkyl may indicate

that the cycloalkyl connects to one atom of the aryl with a bond or by fusing with the aryl and sharing two or more common atoms.

[0036] As used herein, the term “unsubstituted” means that the specified group bears no substituents.

[0037] Unless otherwise specifically defined, the term "aryl" refers to cyclic, aromatic hydrocarbon groups that have 1 to 3 aromatic rings, including monocyclic or bicyclic groups such as phenyl, biphenyl or naphthyl. Where containing two aromatic rings (bicyclic, etc.), the aromatic rings of the aryl group may be joined at a single point (*e.g.*, biphenyl), or fused (*e.g.*, naphthyl). The aryl group may be optionally substituted by one or more substituents, *e.g.*, 1 to 5 substituents, at any point of attachment. Exemplary substituents include, but are not limited to, -H, -halogen, -O-(C₁-C₆) alkyl, (C₁-C₆) alkyl, -O-(C₂-C₆) alkenyl, -O-(C₂-C₆) alkynyl, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, -OH, -OP(O)(OH)₂, -OC(O)(C₁-C₆) alkyl, -C(O)(C₁-C₆) alkyl, -OC(O)O(C₁-C₆) alkyl, -NH₂, NH((C₁-C₆) alkyl), N((C₁-C₆) alkyl)₂, -S(O)₂-(C₁-C₆) alkyl, -S(O)NH(C₁-C₆) alkyl, and -S(O)N((C₁-C₆) alkyl)₂. The substituents can themselves be optionally substituted. Furthermore when containing two fused rings the aryl groups herein defined may have an unsaturated or partially saturated ring fused with a fully saturated ring. Exemplary ring systems of these aryl groups include, but are not limited to, phenyl, biphenyl, naphthyl, anthracenyl, phenalenyl, phenanthrenyl, indanyl, indenyl, tetrahydronaphthalenyl, tetrahydrobenzoannulenyl, and the like.

[0038] Unless otherwise specifically defined, "heteroaryl" means a monovalent monocyclic aromatic radical of 5 to 24 ring atoms or a polycyclic aromatic radical, containing one or more ring heteroatoms selected from N, O, and S, the remaining ring atoms being C. Heteroaryl as herein defined also means a bicyclic heteroaromatic group wherein the heteroatom is selected from N, O, and S. The aromatic radical is optionally substituted independently with one or more substituents described herein. Examples include, but are not limited to, furyl, thienyl, pyrrolyl, pyridyl, pyrazolyl, pyrimidinyl, imidazolyl, isoxazolyl, oxazolyl, oxadiazolyl, pyrazinyl, indolyl, thiophen-2-yl, quinolyl, benzopyranyl, isothiazolyl, thiazolyl, thiadiazole, indazole, benzimidazolyl, thieno[3,2-b]thiophene, triazolyl, triazinyl, imidazo[1,2-b]pyrazolyl, furo[2,3-c]pyridinyl, imidazo[1,2-a]pyridinyl, indazolyl, pyrrolo[2,3-c]pyridinyl, pyrrolo[3,2-c]pyridinyl, pyrazolo[3,4-c]pyridinyl, thieno[3,2-c]pyridinyl, thieno[2,3-c]pyridinyl, thieno[2,3-b]pyridinyl,

benzothiazolyl, indolyl, indolinyl, indolinonyl, dihydrobenzothiophenyl, dihydrobenzofuranyl, benzofuran, chromanyl, thiochromanyl, tetrahydroquinoliny, dihydrobenzothiazine, dihydrobenzoxanyl, quinoliny, isoquinoliny, 1,6-naphthyridiny, benzo[de]isoquinoliny, pyrido[4,3-b][1,6]naphthyridiny, thieno[2,3-b]pyraziny, quinazolinyl, tetrazolo[1,5-a]pyridiny, [1,2,4]triazolo[4,3-a]pyridiny, isoindolyl, pyrrolo[2,3-b]pyridiny, pyrrolo[3,4-b]pyridiny, pyrrolo[3,2-b]pyridiny, imidazo[5,4-b]pyridiny, pyrrolo[1,2-a]pyrimidiny, tetrahydro pyrrolo[1,2-a]pyrimidiny, dibenzo[b,d] thiophene, pyridin-2-one, furo[3,2-c]pyridiny, furo[2,3-c]pyridiny, 1H-pyrido[3,4-b][1,4] thiaziny, benzooxazolyl, benzoisoxazolyl, furo[2,3-b]pyridiny, benzothiophenyl, 1,5-naphthyridiny, furo[3,2-b]pyridine, [1,2,4]triazolo[1,5-a]pyridiny, benzo [1,2,3]triazolyl, imidazo[1,2-a]pyrimidiny, [1,2,4]triazolo[4,3-b]pyridaziny, benzo[c][1,2,5]thiadiazolyl, benzo[c][1,2,5]oxadiazole, 1,3-dihydro-2H-benzo[d]imidazol-2-one, 3,4-dihydro-2H-pyrazolo [1,5-b][1,2]oxaziny, 4,5,6,7-tetrahydropyrazolo[1,5-a]pyridiny, thiazolo[5,4-d]thiazolyl, imidazo[2,1-b][1,3,4]thiadiazolyl, thieno[2,3-b]pyrrolyl, thieno[2,3-d]thiazole, 1a,2,3,7b-tetrahydro-1*H*-cyclopropa[c][1,8]naphthyridine, 3H-indolyl, and derivatives thereof. Furthermore the terms "heteroaryl" and "heteroar-", as used herein, also include groups in which a heteroaromatic ring is fused to one or more aryl, cycloaliphatic, or heterocyclyl rings, where the radical or point of attachment is on the heteroaromatic ring. Nonlimiting examples include indolinyl, indolinonyl, dihydrobenzothiophenyl, dihydrobenzofuran, chromanyl, thiochromanyl, tetrahydroquinoliny, dihydrobenzothiazine, 3,4-dihydro-1H-isoquinoliny, 2,3-dihydrobenzofuran, indolinyl, indolyl, isoindolyl and dihydrobenzoxanyl.

[0039] Halogen or “halo” refers to fluorine, chlorine, bromine, or iodine.

[0040] Alkyl refers to a straight or branched chain saturated hydrocarbon containing 1-12 carbon atoms. Examples of a (C₁-C₆) alkyl group include, but are not limited to, methyl, ethyl, propyl, butyl, pentyl, hexyl, isopropyl, isobutyl, sec-butyl, *tert*-butyl, isopentyl, neopentyl, and isohexyl.

[0041] “Alkoxy” refers to a straight or branched chain saturated hydrocarbon containing 1-12 carbon atoms containing a terminal “O” in the chain, *e.g.*, -O(alkyl). Examples of alkoxy groups include without limitation, methoxy, ethoxy, propoxy, butoxy, *t*-butoxy, or pentoxy groups.

[0042] The term “alkylene” or “alkylenyl” refers to a divalent alkyl radical. Any of the above mentioned monovalent alkyl groups may be an alkylene by abstraction of a second hydrogen atom

from the alkyl. As herein defined, alkylene may also be a C₀-C₆ alkylene. An alkylene may further be a C₀-C₄ alkylene. Typical alkylene groups include, but are not limited to, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH₂CH₂-, -CH₂CH(CH₃)-, -CH₂C(CH₃)₂-, -CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, and the like.

[0043] “Cycloalkyl” or “carbocyclyl” means monocyclic or polycyclic saturated carbon rings containing 3-18 carbon atoms. Examples of cycloalkyl groups include, without limitations, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, norboranyl, norbornyl, bicyclo[2.2.2]octanyl, or bicyclo[2.2.2]octenyl and derivatives thereof. A C₃-C₈ cycloalkyl is a cycloalkyl group containing between 3 and 8 carbon atoms. A cycloalkyl group can be fused (*e.g.*, decalin) or bridged (*e.g.*, norbornane).

[0044] “Heterocyclyl” or “heterocycloalkyl” monocyclic or polycyclic rings containing carbon and heteroatoms taken from oxygen, nitrogen, or sulfur and wherein there is not delocalized π electrons (aromaticity) shared among the entire ring carbon or heteroatoms. The heterocycloalkyl ring structure may be substituted by one or more substituents. The substituents can themselves be optionally substituted. Examples of heterocyclyl rings include, but are not limited to, oxetanyl, azetadanyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrolidinyl, oxazolinyl, oxazolidinyl, thiazolinyl, thiazolidinyl, pyranal, thiopyranal, tetrahydropyranyl, dioxalinyl, piperidinyl, morpholinyl, thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S-dioxide, piperazinyl, azepinyl, oxepinyl, diazepinyl, tropanyl, oxazolidinonyl, and homotropanyl. As used herein, “heterocyclyl” and “heterocycloalkyl” also includes bridged and spirocyclic ring systems where at least one atom is a heteroatom. A heterocyclic ring as a substituent may attach via a ring heteroatom (*e.g.* “N-linked”) or via a ring carbon (*e.g.* “C-linked”).

[0045] The term “hydroxyalkyl” means an alkyl group as defined above, where the alkyl group is substituted with one or more OH groups. Examples of hydroxyalkyl groups include HO-CH₂-, HO-CH₂-CH₂- and CH₃-CH(OH)-.

[0046] The term “haloalkyl” as used herein refers to an alkyl group, as defined herein, which is substituted one or more halogen. Examples of haloalkyl groups include, but are not limited to, trifluoromethyl, difluoromethyl, pentafluoroethyl, trichloromethyl, etc.

[0047] The term “haloalkoxy” as used herein refers to an alkoxy group, as defined herein, which is substituted one or more halogen. Examples of haloalkyl groups include, but are not limited to, trifluoromethoxy, difluoromethoxy, pentafluoroethoxy, trichloromethoxy, etc.

[0048] The term “cyano” as used herein means a substituent having a carbon atom joined to a nitrogen atom by a triple bond, *e.g.*, $C\equiv N$.

[0049] The term "solvate" refers to a complex of variable stoichiometry formed by a solute and solvent. Such solvents for the purpose of the disclosure may not interfere with the biological activity of the solute. Examples of suitable solvents include, but are not limited to, water, MeOH, EtOH, and AcOH. Solvates wherein water is the solvent molecule are typically referred to as hydrates. Hydrates include compositions containing stoichiometric amounts of water, as well as compositions containing variable amounts of water.

[0050] The term "isomer" refers to compounds that have the same composition and molecular weight but differ in physical and/or chemical properties. The structural difference may be in constitution (geometric isomers) or in the ability to rotate the plane of polarized light (stereoisomers). With regard to stereoisomers, the compounds of Formula (I) may have one or more asymmetric carbon atom and may occur as racemates, racemic mixtures and as individual enantiomers or diastereomers.

[0051] The disclosure also includes pharmaceutical compositions comprising an effective amount of a disclosed compound and a pharmaceutically acceptable carrier.

[0052] The term "treating" with regard to a subject, refers to improving at least one symptom of the subject's disorder. Treating includes curing, improving, or at least partially ameliorating the disorder.

[0053] The term "disorder" is used in this disclosure to mean, and is used interchangeably with, the terms disease, condition, or illness, unless otherwise indicated.

[0054] The term "administer", "administering", or "administration" as used in this disclosure refers to either directly administering a disclosed compound or pharmaceutically acceptable salt of the disclosed compound or a composition to a subject, or administering a prodrug derivative or analog of the compound or pharmaceutically acceptable salt of the compound or composition to the subject, which can form an equivalent amount of active compound within the subject's body.

[0055] The term "prodrug," as used in this disclosure, means a compound which is convertible in vivo by metabolic means (*e.g.*, by hydrolysis) to a disclosed compound.

[0056] The term "cancer" includes, but is not limited to, the following cancers: bladder cancer, breast cancer (*e.g.*, ductal carcinoma), cervical cancer (*e.g.*: squamous cell carcinoma), colorectal cancer (*e.g.*, adenocarcinoma), esophageal cancer (*e.g.*, squamous cell carcinoma), gastric cancer (*e.g.*: adenocarcinoma, medulloblastoma, colon cancer, choriocarcinoma, squamous cell carcinoma), head and neck cancer, hematologic cancer (*e.g.*, acute lymphocytic anemia, acute myeloid leukemia, acute lymphoblastic B cell leukemia, anaplastic large cell lymphoma, B-cell lymphoma, Burkitt's lymphoma, chronic lymphocytic leukemia, chronic eosinophilic leukemia/hypereosinophilic syndrome, chronic myeloid leukemia, Hodgkin's lymphoma, mantle cell lymphoma, multiple myeloma, T-cell acute lymphoblastic leukemia), lung cancer (*e.g.*, bronchioloalveolar adenocarcinoma, mesothelioma, mucoepidermoid carcinoma, small-cell lung cancer, non-small cell lung cancer, adenocarcinoma, squamous cell carcinoma), liver cancer (*e.g.*, hepatocellular carcinoma), lymphoma, neurological cancer (*e.g.*, glioblastoma, neuroblastoma, neuroglioma), ovarian (*e.g.*, adenocarcinoma), pancreatic cancer (*e.g.*, ductal carcinoma), prostate cancer (*e.g.*, adenocarcinoma), renal cancer (*e.g.*, renal cell carcinoma, clear cell renal carcinoma), sarcoma (*e.g.*, chondrosarcoma, Ewings sarcoma, fibrosarcoma, multipotential sarcoma, osteosarcoma, rhabdomyosarcoma, synovial sarcoma), skin cancer (*e.g.*, melanoma, epidermoid carcinoma, squamous cell carcinoma), thyroid cancer (*e.g.*, medullary carcinoma), and uterine cancer.


[0057] The present disclosure relates to chemical entities chosen from compounds and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, capable of inhibiting at least one pathway chosen from USP28 and USP25, which are useful for the treatment of diseases and disorders associated with modulation of at least one pathway chosen from USP28 and USP25. The disclosure further relates to chemical entities chosen from compounds and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, which are useful for inhibiting at least one pathway chosen from USP28 and USP25.

[0058] In any of the embodiments as disclosed herein, the cancer can be any cancer in any organ, for example, a cancer is selected from the group consisting of glioma, thyroid carcinoma,

breast carcinoma, small-cell lung carcinoma, non-small-cell carcinoma, gastric carcinoma, colon carcinoma, gastrointestinal stromal carcinoma, pancreatic carcinoma, bile duct carcinoma, CNS carcinoma, ovarian carcinoma, endometrial carcinoma, prostate carcinoma, renal carcinoma, anaplastic large-cell lymphoma, leukemia, multiple myeloma, mesothelioma, and melanoma, and combinations thereof.


[0059] The present disclosure relates to chemical entities chosen from compounds and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, capable of inhibiting at least one pathway chosen from USP28 and USP25, which are useful for the treatment of diseases and disorders associated with modulation of at least one pathway chosen from USP28 and/or USP25 enzyme. The present disclosure further relates to chemical entities chosen from compounds and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, which are useful for inhibiting at least one pathway chosen from USP28 and USP25.

[0060] In one embodiment, the chemical entities are chosen from compounds of Formula (II):




wherein X, Y, R₁, R₂, R₃, R₄, and n are as described herein above, and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0061] In another embodiment, the chemical entities are chosen from compounds of Formula (III):




wherein X, Y, R₁, R₂, R₃, R₄, and n are as described herein above, and pharmaceutically acceptable salts, hydrates, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0062] In another embodiment, the chemical entities are chosen from compounds of Formula (IV):




wherein X, Y, R₁, R₂, R₃, R₄, and n are as described herein above and pharmaceutically acceptable salts, hydrates, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0063] In another embodiment, the chemical entities are chosen from compounds of Formula (V):



wherein X, Y, R₁, R₂, R₃, R₄, and n are as described herein above and pharmaceutically acceptable salts, hydrates, solvates, prodrugs, stereoisomers, and tautomers thereof.

[0064] In some embodiments, the chemical entities are chosen from compounds of Formula (VI):



or a pharmaceutically acceptable salt thereof,

wherein:

X is chosen from C(R)(R'') and O;

each of Y₁, Y₂, and Y₃ is independently chosen from C(R₃) and N;

R' is chosen from H, deuterium, and CH₃;

each of R and R'' is independently chosen from H, halogens, -OH, -CN, C₁-C₆ alkyl optionally substituted with one or more R_i,

or R and R'' together with the carbon they are attached form a spirocyclic cyclopropyl optionally substituted with one or more R_i, wherein any R, R'', or R_i group being or containing hydrogen can independently have one or more hydrogen replaced with deuterium;

each R_i is independently chosen from halogen, -OH, and CH₃;

R₁ is chosen from 6-12 membered fused and nonfused heteroaryls optionally substituted with one or more substituent chosen from R₅ and/or R₆, and further wherein any R₁ group containing hydrogen can have one or more hydrogen replaced with deuterium;

R₂ is chosen from N-linked 4-12 membered heterocyclyls, C-linked 4-12 membered heterocyclyls, and an O linker attached to a 4-12 membered heterocyclyl, wherein the 4-12 membered heterocyclyls are optionally substituted with one or more R₅ (which can be the same or different from the one or more R₅ of R₁), and further wherein any hydrogen in a R₂ group can have one or more hydrogen replaced with deuterium;

each R₃ is independently chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein each of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₇;

R₄ is chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein each of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl are optionally substituted with one or more R₅, and further wherein any R₄ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₅ is independently chosen from –OH, –NH₂, amido-(C₁-C₆) alkyl, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of –NH₂, amido-(C₁-C₆) alkyl, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, –NH₂, and –OH, and wherein any R₅ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₆ is independently chosen from -amino alkyl-aryls, -amino alkyl-heteroaryls, -amino alkyl-cyclyl, and -amino alkyl-heterocyclyl groups, wherein each of the R₆ groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups, and further wherein any R₆ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₇ is independently chosen from –OH, –NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of –NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and –OH;

and

n is 0, 1, 2, or 3.

[0065] The compound of Formula (VI) or a pharmaceutically acceptable salt thereof, wherein:

X is chosen from C(R)(R'') and O;

each of Y₁, Y₂, and Y₃ is independently chosen from C(R₃) and N;

R' is chosen from H, and CH₃;

each of R and R'' is independently chosen from H, halogens, -OH, -CN, C₁-C₆ alkyl optionally substituted with one or more R_i,

or R and R'' together with the carbon they are attached form a spirocyclic cycloalkyl (e.g., spirocyclic cyclopropyl) optionally substituted with one or more R_i, wherein any R, R'';

each R_i is independently chosen from halogen, -OH, and CH₃;

R₁ is chosen from a fused or nonfused heteroaryls (e.g., 6-12 membered fused or nonfused heteroaryls) optionally substituted with one or more substituent chosen from R₅ and/or R₆ (e.g., a 6-membered nonfused heteroaryl optionally substituted with one or more R₆);

R₂ is chosen from N-linked-heterocyclyls (e.g., 4-12 membered heterocyclyls), C-linked-heterocyclyls (e.g., 4-12 membered heterocyclyls), and O-linker-heterocyclyls (e.g., 4-12



membered heterocyclyls such as), wherein any of the -N-linked-, -C-linked or -O-linked (e.g., 4-12 membered) heterocyclyls is optionally substituted with one or more R₅ (which can be the same or different from the one or more R₅ of R₁);

each R₃ is independently chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein each of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₇;

R₄ is chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups, wherein each of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, and heteroaryl are optionally substituted with one or more R₅;

each R₅ is independently chosen from -OH, -NH₂, amido-(C₁-C₆) alkyl, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, amido-(C₁-C₆) alkyl, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, -NH₂, and -OH;

each R₆ is independently chosen from -amino alkyl-aryls, -amino alkyl-heteroaryls, -amino alkyl-cyclyl, and -amino alkyl-heterocyclyl groups, wherein each of the R₆ groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups;

each R₇ is independently chosen from –OH, –NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of –NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and –OH; and

n is 0, 1, 2, or 3.

[0066] In some embodiments, a compound of Formula (I), Formula (II), Formula (III), Formula (IV), Formula (V) and/or Formula (VI) above includes X (if present) as CH₂ (e.g., R and R'' are both hydrogen in Formula (VI)). In other embodiments, a compound of Formula (I), Formula (II), Formula (III), Formula (IV), Formula (V) and/or Formula (VI) above includes X (if present) as O.

[0067] In another embodiment, the chemical entities are chosen from compounds of Formula (VI) and pharmaceutically acceptable salts thereof, wherein:

X is chosen from C(R)(R'') and O;

each of Y₁, Y₂, and Y₃ is independently chosen from C(R₃) and N;

R' is chosen from H and CH₃;

each of R and R'' is independently chosen from H, halogens, -OH, -CN, C₁-C₆ alkyl optionally substituted with one or more R_i;

each R_i is independently chosen from halogen, -OH, and CH₃;

R₁ is chosen from 6-12 membered fused and nonfused heteroaryls optionally substituted with one or more R₅, wherein a 6-membered nonfused heteroaryl is substituted with one or more R₆;

R₂ is chosen from N-linked and C-linked 4-12 membered heterocyclyls, wherein the 4-12 membered heterocyclyls are optionally substituted with one or more R₅ (which can be the same or different from the one or more R₅ of R₁);

each R₃ is independently chosen from H, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, and -CN, wherein each of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, and (C₁-C₆) haloalkoxy groups are optionally substituted with one or more R₇;

R_4 is chosen from H, (C₁-C₆) alkyl, halogen, -OH, and -CN, wherein the (C₁-C₆) alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH;

each R_5 is independently chosen from -OH, -NH₂, -NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein alkyls are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and -OH;


each R_6 is independently chosen from -NH(C₁-C₆)alkyl-aryls, -NH(C₁-C₆)alkyl-heteroaryls, -NH(C₁-C₆)alkyl-cyclyl and -NH(C₁-C₆)alkyl-heterocyclyl groups, wherein each of the R_6 groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups;

each R_7 is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein the (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy and -OH;

and

n is 0, 1, 2, or 3.

[0068] In another embodiment, the chemical entities are chosen from compounds of Formula (VII):



or a pharmaceutically acceptable salt thereof, wherein

Y is chosen from C(R₃) and N;

R' is chosen from H, deuterium, and CH₃;

R₁ is chosen from 6-11 membered heteroaryls optionally substituted with one or more substituent chosen from R₅ and/or R₆;

R₂ is chosen from N-linked 4-12 membered heterocyclyls and C-linked 4-12 membered heterocyclyls, wherein the heterocyclyls are optionally substituted with one or more R₅, and further wherein any R₂ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₃ (if present) is independently chosen from H, deuterium, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, wherein each of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₇;

each R₄ is chosen from H, deuterium, (C₁-C₆) alkyl, halogen, -OH, -CN, and further wherein any R₄ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₅ (if present) is independently chosen from -OH, -NH₂, NHC(O)CH₃, -C(O)NHCH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, -NHC(O)CH₃, -C(O)NHCH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and -OH, and wherein any R₅ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₆ (if present) is chosen from -NH(C₁-C₆)alkyl-aryls, -NH(C₁-C₆)alkyl-heteroaryls, -NH(C₁-C₆)alkyl-heterocyclyl groups, and -NH(C₁-C₆)alkyl-heterocyclyl groups, wherein each of the R₆ groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups, and further wherein any R₆ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₇ (if present) is independently chosen from –OH, –NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of –NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and –OH; and

n is 0, 1, 2, or 3.

[0069] In some embodiments of the Formulae above, X is CH₂. In another embodiments, X is O.

[0070] In some embodiments of the Formulae above, R₁ is chosen from 6-12 membered heteroaryls optionally substituted with one or more R₅. In some embodiments of the Formulae above, R₁ is chosen from 6 membered heteroaryls substituted with one or more R₆. In some embodiments of the Formulae above, any R₁ group or optional substituent containing hydrogen can have one or more hydrogen replaced with deuterium.

[0071] In some embodiments of the Formulae above, R₂ is chosen from N-linked 4-10 membered heterocyclyls optionally substituted with one or more R₅, and wherein a sulfur member of the heterocyclyls can be S(O) or S(O)₂. In some embodiments of the Formulae above R₂ is chosen from C-linked 4-10 membered heterocyclyls optionally substituted with one or more R₅, and wherein a sulfur member of the heterocyclyls can be S(O) or S(O)₂. In some embodiments of the Formulae above R₂ is chosen from O linked to a heterocyclic entity that is optionally substituted with one or more R₅, and wherein a sulfur member of the heterocyclyls can be S(O) or S(O)₂. In some embodiments of the Formulae above, any R₂ group or optional substituent containing hydrogen can have one or more hydrogen replaced with deuterium.





















[0072] In some embodiments of the Formulae above, R₃ is independently chosen from H, (C₁-C₆) alkyl, (C₁-C₆) haloalkoxy, halogen, and –CN.


















[0073] In some embodiments of the Formulae above, R₄ is chosen from H and (C₁-C₆) alkyls. In some embodiments of the Formulae above, one or more hydrogen of R₄ can be replaced with deuterium.

[0074] In some embodiments of the Formulae above, n is 0, 1, or 2. In another embodiment, n is 0 or 1. In yet another embodiment, n is 1, 2, or 3. In another embodiment, n is 1 or 2. In another embodiment, n is 2 or 3. In another embodiment, n is 0. In another embodiment, n is 1. In another embodiment, n is 2. In another embodiment, n is 3.

[0075] In some embodiments of the Formulae above, R₁, optionally substituted with R₅ and/or R₆, is chosen from the groups of Table A. Preferably, a compound of Formula (I), Formula (II), Formula (III), Formula (IV), Formula (V), Formula (VI) and/or Formula (VII) comprises R₁ (alone or as substituted with one of more R₅ and/or R₆) that is selected from the groups in Table A below.

























Table A:

























			
			
			
			
			























			
			
			
			
			



















[0076] In some embodiments of the Formulae above, R₂, optionally substituted with R₅, is chosen from the groups of Table B below. Preferably, a compound of Formula (I), Formula (II), Formula (III), Formula (IV), Formula (V), Formula (VI) and/or Formula (VII) can comprises R₂ (alone or as substituted with one of more R₅ and/or R₆, which can be the same or different) that is selected from the groups in Table B below.

Table B


			
			
			
			
			
			


[0077] Preferably, a compound of Formula (I), Formula (II), Formula (III), Formula (IV), Formula (V), Formula (VI) and/or Formula (VII) can comprises both R₁ (alone or as substituted with one of more R₅ and/or R₆, which can be the same or different) that is selected from the groups in Table A above, and R₂ (alone or as substituted with one of more R₅ and/or R₆, which can be the same or different) that is selected from the groups in Table B above.

[0078] In some embodiments, the chemical entities are chosen from compounds of Formula (IIa):



(IIa),


or a pharmaceutically acceptable salt thereof, wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and n are as defined in Formula (I), and/or from compounds of Formula (IIaa):



(IIaa),

or a pharmaceutically acceptable salt thereof, wherein R_1 , R_2 , R_3 , R_5 , R_6 , R_7 and n are as defined in Formula (I).


In some embodiments, the chemical entities are chosen from compounds of Formula (IIb):



(IIb'),

or a pharmaceutically acceptable salt thereof, wherein R' , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and n are each independently as defined Formula (VI).

[0079] In some embodiments, the chemical entities are chosen from compounds of Formula (IIaa):



(IIaa),


or a pharmaceutically acceptable salt thereof,
wherein

R₁ is chosen from 8-9-membered heteroaryls substituted with one or more substituent chosen from R₅ and R₆;


R₂ is chosen from N-linked 6-12 membered heterocyclyls or C-linked 6-12 membered heterocyclyls, wherein the 6-12 membered heterocyclyls are optionally substituted with one or more R₅;

R₃ is independently chosen from H, (C₁-C₆) alkyl, halogen, and -CN; and wherein R₅, R₆, R₇ and n are each independently as defined in Formula (I).


[0080] In preferred embodiments of Formula (IIaa), R₁, optionally substituted with R₅ and/or R₆, is chosen from



R₂, optionally substituted with R₅, is chosen from



[0081] In some embodiments, the chemical entities are chosen from compounds of Formula (IIb'):



wherein R' is chosen from H and CH₃;


R₁ is chosen from 8-9 membered heteroaryls substituted with one or more substituent chosen from R₅ and R₆;

R₂ is chosen from N-linked 6-12 membered heterocyclyls or C-linked 6-12 membered heterocyclyls, wherein the 6-12 membered heterocyclyls are optionally substituted with one or more R₅;


R₃ is independently chosen from H and halogen; and

R₅, R₆, R₇ and n are each independently as defined in Formula (VI).


[0082] In preferred embodiments of Formula (IIb'), R₁, optionally substituted with R₅ and/or R₆, is chosen from



R₂, optionally substituted with R₅, is chosen from




[0083] In some embodiments, the chemical entities are chosen from compounds of Formula (IIIa):




or a pharmaceutically acceptable salt thereof,

wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and n are each independently as defined Formula (I); and/or from compounds Formula (IIIb):



or a pharmaceutically acceptable salt thereof, wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and n are each independently as defined Formula (I).

[0084] In some embodiments, the chemical entities are chosen from compounds of Formula (IIIaa):



or a pharmaceutically acceptable salt thereof, wherein


R_1 is chosen from 8-11 membered heteroaryls optionally substituted with one or more R_5 ;

R_2 is chosen from N-linked 4-12 membered heterocyclyls and C-linked 4-12 membered heterocyclyls, optionally substituted with one or more R_5 ;

each R_5 (if present) is independently chosen from $-OH$, $-NH_2$, $NHC(O)CH_3$, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_1-C_6) haloalkyl, (C_1-C_6) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and $-C(O)-$ heterocycloalkyl groups, wherein each of $-NH_2$, $-NHC(O)CH_3$, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_1-C_6) haloalkyl, (C_1-C_6) haloalkoxy, cycloalkyl, heterocycloalkyl, and $-C(O)-$ heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C_1-C_6) alkoxy, $-NH_2$, and $-OH$; and

n is 0, 1, 2, or 3.

[0085] In preferred embodiments of Formula (IIIaa), R_1 is chosen from



wherein B is chosen from a bond or C;

Z is chosen from N, S, C(Rii);


Rii is chosen from H, CH₃ and R₅;

R₂ is chosen from N-linked 5-8 membered heterocyclyls substituted with one to three R₅;


each R₅ (if present) is independently chosen from -OH, -NH₂, NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, -NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and -OH; and

n is 0, 1, 2, or 3.


[0086] In further preferred embodiments of Formula (IIIaa), R₁, optionally substituted with R₅, is chosen from



R₂, optionally substituted with R₅, is chosen from




[0087] In some embodiments, the chemical entities are chosen from compounds of Formula (IVa):




or a pharmaceutically acceptable salt thereof,
wherein R_1 , R_2 , R_4 , R_5 , R_6 , R_7 and n are each independently as defined Formula (I).

[0088] In some embodiments, the chemical entities are chosen from compounds of Formula (Va):



or a pharmaceutically acceptable salt thereof,
wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 and n are each independently as defined Formula (I).


[0089] In some embodiments, the chemical entity is chosen from compounds of Formula (Vaa):




or a pharmaceutically acceptable salt thereof,

wherein R_1 is chosen from 8-membered heteroaryls substituted with one or more substituent chosen from R_5 and R_6 ; R_2 is chosen from N-linked 5-membered heterocyclyls; and R_5 , R_6 , R_7 and n are each independently as defined Formula (I).

In further preferred embodiments of Formula (Vaa), R_1 substituted with R_5 is chosen from




[0090] In some embodiments, the chemical entities are chosen from compounds of Formula (VIIa'):



or a pharmaceutically acceptable salt thereof,

wherein R', R₁, R₂, R₃, R₄, R₅, R₆, R₇ and n are each independently as defined Formula (VI).

[0091] In at least one embodiment, the chemical entities are chosen from compounds of Formula (VIIaa):



or a pharmaceutically acceptable salt thereof, wherein


R₁ is chosen from 8-9 membered heteroaryls optionally substituted with one or more R₅;

R₂ is chosen from N-linked 4-12 membered heterocyclyls optionally substituted with one or more R₅;

each R₅ (if present) is independently chosen from -OH, -NH₂, NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, -NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and -OH; and

n is 0, 1, 2, or 3.

[0092] In preferred embodiments of Formula (VIIaa), R₁ is chosen from



wherein B is chosen from a bond or C;

Z is chosen from N, S, C(Rii);


Rii is chosen from H, CH₃ and R₅;

R₂ is chosen from N-linked 5-8 membered heterocyclyls substituted with one to three R₅;


each R₅ (if present) is independently chosen from -OH, -NH₂, NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, -NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and -OH; and

n is 0, 1, 2, or 3.

[0093] In further preferred embodiments of Formula (VIIaa), R₁, optionally substituted with R₅, is chosen from



R₂, optionally substituted with R₅, is chosen from





[00100] The compounds of the disclosure may contain asymmetric or chiral centers, and, therefore, exist in different stereoisomeric forms. It is intended that all stereoisomeric forms of the compounds of the disclosure as well as mixtures thereof, including racemic mixtures, form part of the present disclosure. In addition, the present disclosure embraces all geometric and positional isomers. For example, if a compound of the disclosure incorporates a double bond or a fused ring, both the cis- and trans-forms, as well as mixtures, are embraced within the scope of the disclosure. Each compound herein disclosed includes all the enantiomers that conform to the general structure of the compound. The compounds may be in a racemic or enantiomerically pure form, or any other form in terms of stereochemistry. The assay results may reflect the data collected for the racemic form, the enantiomerically pure form, or any other form in terms of stereochemistry.

[00101] Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (*e.g.*, chiral auxiliary such as a chiral alcohol or Mosher's acid chloride), separating the diastereomers and converting (*e.g.*, hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some of the compounds of the disclosure may be atropisomers (*e.g.*, substituted biaryls) and are considered as part of this disclosure. Enantiomers can also be separated by use of a chiral HPLC column.

[00102] It is also possible that the compounds of the disclosure may exist in different tautomeric forms, and all such forms are embraced within the scope of the disclosure. Also, for example, all keto-enol and imine-enamine forms of the compounds are included in the disclosure.

[00103] All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds (including those of the salts, solvates, esters and prodrugs of the compounds as well as the salts, solvates and esters of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms, are contemplated within the scope of this disclosure, as are positional isomers (such as, for example, 4-pyridyl and 3-pyridyl). (For example, if a compound of Formula (I) incorporates a double bond or a fused ring, both the cis- and trans-forms, as well as mixtures, are embraced within

the scope of the disclosure. Also, for example, all keto-enol and imine-enamine forms of the compounds are included in the disclosure.) Individual stereoisomers of the compounds of the disclosure may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present disclosure can have the S or R configuration as defined by the IUPAC 1974 Recommendations. The use of the terms “salt”, “solvate”, “ester,” “prodrug” and the like, is intended to equally apply to the salt, solvate, ester and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or prodrugs of the compounds disclosed herein.

[00104] The compounds disclosed herein may form salts which are also within the scope of this disclosure.

[00105] The present disclosure relates to compounds which are modulators of at least one pathway chosen from USP28 and USP25. In one embodiment, the compounds of the present disclosure are inhibitors of at least one pathway chosen from USP28 and USP25.

[00106] The present disclosure is directed to chemical entities chosen from compounds as described herein and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, and pharmaceutical compositions comprising at least one chemical entity chosen from compounds as described herein, and pharmaceutically acceptable salts, hydrates, solvates, prodrugs, stereoisomers, and tautomers thereof.

[00107] Another aspect of the disclosure relates to a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with modulation of USP28. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 an effective amount the compositions and chemical entities of Formula (I). In one embodiment, the disease or disorder is cancer.

[00108] In another aspect, the present disclosure is directed to a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with inhibition of USP28. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 an effective amount the compositions and chemical entities of Formula (I). In one embodiment, the disease or disorder is cancer.

[00109] In another aspect, the present disclosure is directed to a method of inhibiting USP28. The method involves administering to a patient in need thereof an effective amount of a chemical entity of Formula (I).

[00110] Another aspect of the disclosure relates to a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with modulation of USP25. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP25 an effective amount the compositions and chemical entities of Formula (I). In one embodiment, the disease or disorder is cancer.

[00111] In another aspect, the present disclosure is directed to a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with inhibition of USP25. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP25 an effective amount the compositions and chemical entities of Formula (I). In one embodiment, the disease or disorder is cancer.

[00112] In another aspect, the present disclosure is directed to a method of inhibiting USP25. The method involves administering to a patient in need thereof an effective amount of a chemical entity of Formula (I).

[00113] Another aspect of the disclosure relates to a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with modulation of USP28. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 an effective amount the compositions and chemical entities of Formula (I). In one embodiment, the disease or disorder is cancer. In another aspect, the present disclosure is directed to a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with inhibition of USP28. The method comprises administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 an effective amount the compositions and chemical entities of Formula (I). In one embodiment, the disease or disorder is cancer. The method can comprise administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 and/or USP25 an effective amount of a pharmaceutical composition comprising a USP28 Inhibitor, USP25 Inhibitor, and/or USP28/25 Inhibitor as disclosed herein. In another aspect, the present disclosure is directed to a method of inhibiting at least one pathway chosen from USP28 and USP25. The method involves

administering to a patient in need thereof an effective amount of a chemical entity of Formula (I). The method can comprise administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 and/or USP25 an effective amount of a pharmaceutical composition comprising a USP28 Inhibitor, USP25 Inhibitor, and/or USP28/25 Inhibitor as disclosed herein.

[00114] Another aspect of the present disclosure relates to a method of treating, preventing, inhibiting, or eliminating a disease or disorder in a patient associated with the inhibition of USP28, the method comprising administering to a patient in need thereof an effective amount of a chemical entity of Formula (I). In one embodiment, the disease or disorder is cancer. The method can comprise administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 and/or USP25 an effective amount of a pharmaceutical composition comprising a USP28 Inhibitor, USP25 Inhibitor, and/or USP28/25 Inhibitor as disclosed herein.

[00115] Another aspect of the present disclosure relates to a method of treating, preventing, inhibiting, or eliminating a disease or disorder in a patient associated with the inhibition of USP25, the method comprising administering to a patient in need thereof an effective amount of a chemical entity of Formula (I). In one embodiment, the disease or disorder is cancer. The method can comprise administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 and/or USP25 an effective amount of a pharmaceutical composition comprising a USP28 Inhibitor, USP25 Inhibitor, and/or USP28/25 Inhibitor as disclosed herein.

[00116] Another aspect of the present disclosure relates to a method of treating, preventing, inhibiting, or eliminating a disease or disorder in a patient associated with the inhibition of at least one pathway chosen from USP28 and USP25, the method comprising administering to a patient in need thereof an effective amount of a chemical entity of Formula (I). In one embodiment, the disease or disorder is cancer. The method can comprise administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 and/or USP25 an effective amount of a pharmaceutical composition comprising a USP28 Inhibitor, USP25 Inhibitor, and/or USP28/25 Inhibitor as disclosed herein.

[00117] In another aspect, the present disclosure relates to a method of treating, preventing, inhibiting, or eliminating cancer. The method comprises administering to a patient in need of a treatment for cancer an effective amount of a chemical entity chosen from compounds of Formula

(I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof. The method can comprise administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 and/or USP25 an effective amount of a pharmaceutical composition comprising a USP28 Inhibitor, USP25 Inhibitor, and/or USP28/25 Inhibitor as disclosed herein.

[00118] Another aspect of the present disclosure relates to a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, for use in a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with inhibiting USP28. In one embodiment, the disease or disorder is cancer. The method can comprise administering to a patient in need of a treatment for diseases or disorders associated with modulation of USP28 and/or USP25 an effective amount of a pharmaceutical composition comprising a USP28 Inhibitor, USP25 Inhibitor, and/or USP28/25 Inhibitor as disclosed herein.

[00119] In another aspect, the present disclosure relates to a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, for use in a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with inhibiting USP25. In one embodiment, the disease or disorder is cancer.

[00120] Another aspect of the present disclosure relates to a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, for use in a method of treating, preventing, inhibiting, or eliminating a disease or disorder associated with inhibiting at least one pathway chosen from USP28 and USP25. In one embodiment, the disease or disorder is cancer.

[00121] In another aspect, the present disclosure relates to a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, for use in a method for treating, preventing, inhibiting, or eliminating cancer.

[00122] Another aspect of the present disclosure relates to the use of a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs,

stereoisomers, and tautomers thereof, in the manufacture of a medicament for treating, preventing, inhibiting, or eliminating a disease or disorder associated with inhibiting USP28. In one embodiment, the disease or disorder is cancer.

[00123]

[00124] Another aspect of the present disclosure relates to the use of a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, in the manufacture of a medicament for treating, preventing, inhibiting, or eliminating a disease or disorder associated with inhibiting at least one pathway chosen from USP28 and USP25. In one embodiment, the disease or disorder is cancer.

[00125] In another aspect, the present disclosure relates to the use of a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof, in the manufacture of a medicament for treating, preventing, inhibiting, or eliminating cancer.

[00126] In other embodiments, the present disclosure relates to the use of an inhibitor of USP28 for the preparation of a medicament used in the treatment, prevention, inhibition or elimination of a disease or disorder associated with cancer.

[00127] The present disclosure also relates to the use of an inhibitor of USP28 for the preparation of a medicament used in the treatment, prevention, inhibition, or elimination of a disease or condition mediated by USP28, wherein the medicament comprises a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof. The present disclosure also relates to the use of an inhibitor of USP28 for the preparation of a medicament used in the treatment, prevention, inhibition, or elimination of a disease or condition mediated by USP28, wherein the medicament comprises a chemical entity chosen from compounds of Formulae (II), (III), (IV), (V), (VI), (VII), (IIa), (IIaa), (IIb'), (IIIa), (IIIb), (IIIaa), (IVa), (Va), (Vaa), (VIIa') or Formula (VIIaa) that is USP28 Inhibitor, a USP25 Inhibitor and/or a USP 28/25 Inhibitor as defined herein.

[00128] The present disclosure also relates to the use of an inhibitor of USP25 for the preparation of a medicament used in the treatment, prevention, inhibition, or elimination of a disease or condition mediated by USP25, wherein the medicament comprises a chemical entity

chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof. The present disclosure also relates to the use of an inhibitor of USP25 for the preparation of a medicament used in the treatment, prevention, inhibition, or elimination of a disease or condition mediated by USP25, wherein the medicament comprises a chemical entity chosen from compounds of Formulae (II), (III), (IV), (V), (VI), (VII), (IIa), (IIaa), (IIb'), (IIIa), (IIIb), (IIIaa), (IVa), (Va), (Vaa), (VIIa') or Formula (VIIaa) that is USP28 Inhibitor, a USP25 Inhibitor and/or a USP 28/25 Inhibitor as defined herein.

[00129] In another aspect, the present disclosure relates to a method for the manufacture of a medicament for treating, preventing, inhibiting, or eliminating a disease or condition mediated by at least one pathway chosen from USP28 and USP25, wherein the medicament comprises a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof. These mechanisms have been shown to be useful in treating cancer such as lung cancer and brain cancer.

[00130] In some embodiments, the patient is selected for treatment based on gene amplification and/or elevated tumor expression of USP28, MYC, LSD1, NICD1, and/or reduced expression of FBXW7 relative to tissue-matched expression.

[00131] In some embodiments, the patient is selected for treatment based on gene amplification and/or elevated tumor expression of USP28, USP25, MYC, LSD1, NICD1, and/or reduced expression of FBXW7 relative to tissue-matched expression.

[00132] In some embodiments, administration of a compound of Formula (I) or a pharmaceutical composition comprising a compound of the present disclosure and a pharmaceutically acceptable carrier induces a change in the cell cycle, cell viability, cell apoptosis, or differentiation.

[00133] For example, the change in the cell cycle or cell viability or differentiation may be indicated by decreased tumor levels of MYC, LSD1, NICD1, PIM1, CDK1, POLA2, HEY1, and/or CCND1, and/or increased levels of CD86, p21, LGALS4, and/or DLL1.

[00134] Another aspect of the disclosure is directed to pharmaceutical compositions comprising a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof and a pharmaceutically acceptable

carrier. The pharmaceutical acceptable carrier may further include an excipient, diluent, or surfactant. Another aspect of the disclosure is directed to pharmaceutical compositions comprising a chemical entity chosen from compounds of Formulae (II), (III), (IV), (V), (VI), (VII), (IIa), (IIaa), (IIb'), (IIIa), (IIIb), (IIIaa), (IVa), (Va), (Vaa), (VIIa') or Formula (VIIaa) that is USP28 Inhibitor, a USP25 Inhibitor and/or a USP 28/25 Inhibitor as defined herein.

[00135] In one embodiment, are provided methods of treating a disease or disorder associated with modulation of USP28 including cancer comprising administering to a patient suffering from at least one of said diseases or disorder a chemical entity of Formula (I). The methods of treating a disease or disorder associated with modulation of USP28 including cancer can comprise administering to a patient suffering from at least one of said diseases or disorder a chemical entity of Formulae (II), (III), (IV), (V), (VI), (VII), (IIa), (IIaa), (IIb'), (IIIa), (IIIb), (IIIaa), (IVa), (Va), (Vaa), (VIIa') or Formula (VIIaa) that is USP28 Inhibitor, a USP25 Inhibitor and/or a USP 28/25 Inhibitor as defined herein.

[00136] In another embodiment, are provided methods of treating a disease or disorder associated with modulation of USP25 including cancer, comprising administering to a patient suffering from at least one of said diseases or disorder a chemical entity of Formula (I). The methods of treating a disease or disorder associated with modulation of USP25 including cancer, can comprise administering to a patient suffering from at least one of said diseases or disorder a chemical entity of Formulae (II), (III), (IV), (V), (VI), (VII), (IIa), (IIaa), (IIb'), (IIIa), (IIIb), (IIIaa), (IVa), (Va), (Vaa), (VIIa') or Formula (VIIaa) that is USP28 Inhibitor, a USP25 Inhibitor and/or a USP 28/25 Inhibitor as defined herein.

[00137] In another embodiment, are provided methods of treating a disease or disorder associated with modulation of at least one pathway chosen from USP28 and USP25 including cancer, comprising administering to a patient suffering from at least one of said diseases or disorder a chemical entity of Formula (I). The methods of treating a disease or disorder associated with modulation of at least one pathway chosen from USP28 and USP25 including cancer, can also comprise administering to a patient suffering from at least one of said diseases or disorder a chemical entity of Formulae (II), (III), (IV), (V), (VI), (VII), (IIa), (IIaa), (IIb'), (IIIa), (IIIb), (IIIaa), (IVa), (Va), (Vaa), (VIIa') or Formula (VIIaa) that is USP28 Inhibitor, a USP25 Inhibitor and/or a USP 28/25 Inhibitor as defined herein.

[00138] One therapeutic use of the compounds or compositions of the present disclosure which inhibit USP28 is to provide treatment to patients or subjects suffering from cancer.

[00139] Another therapeutic use of the compounds or compositions of the present disclosure which inhibit USP25 is to provide treatment to patients or subjects suffering from cancer.

[00140] Another therapeutic use of the compounds or compositions of the present disclosure which inhibit at least one pathway chosen from USP28 and USP25 is to provide treatment to patients or subjects suffering from cancer.

[00141] The compounds of the disclosure can be administered in effective amounts to treat or prevent a disorder and/or prevent the development thereof in subjects.

[00142] Administration of the disclosed compounds can be accomplished via any mode of administration for therapeutic agents.

[00143] Another aspect of the disclosure is directed to pharmaceutical compositions comprising a chemical entity chosen from compounds of Formula (I), and pharmaceutically acceptable salts, solvates, prodrugs, stereoisomers, and tautomers thereof and a pharmaceutically acceptable carrier. The pharmaceutical acceptable carrier may further include an excipient, diluent, or surfactant.

[00144] The dosage regimen utilizing the disclosed compound is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal or hepatic function of the patient; and the particular disclosed compound employed. A physician or veterinarian of ordinary skill in the art can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

[00145] Non-limiting examples of compounds according to Formulae (I)-(VII) of the disclosure include those of Tables 9-25 below.

Method of Synthesizing the Compounds

[00146] The compounds of the present disclosure can be prepared in a number of ways known to those skilled in the art of organic synthesis. The compounds of the present disclosure may be made by a variety of methods, including standard chemistry. Suitable synthetic routes are depicted in the Schemes provided herein. The compounds disclosed herein may be prepared by methods


known in the art of organic synthesis as set forth in part by the following synthetic schemes. In the schemes described herein, it is well understood that protecting groups for sensitive or reactive groups are employed where necessary in accordance with general principles of chemistry. Protecting groups are manipulated according to standard methods of organic synthesis (T. W. Greene and P. G. M. Wuts, "Protective Groups in Organic Synthesis", Third edition, Wiley, New York 1999). These groups are removed at a convenient stage of the compound synthesis using methods that are readily apparent to those skilled in the art. The selection processes, as well as the reaction conditions and order of their execution, shall be consistent with the preparation of compounds disclosed herein (including, e.g., compounds of Formula (I)).

[00147] The compounds described herein may be made from commercially available starting materials or synthesized using known organic, inorganic, and/or enzymatic processes. Those skilled in the art will recognize if a stereocenter exists in the compounds disclosed herein. Accordingly, the present disclosure includes both possible stereoisomers (unless specified in the synthesis) and includes not only racemic compounds but the individual enantiomers and/or diastereomers as well. When a compound is desired as a single enantiomer or diastereomer, it may be obtained by stereospecific synthesis or by resolution of the final product or any convenient intermediate. Resolution of the final product, an intermediate, or a starting material may be affected by any suitable method known in the art. See, for example, "Stereochemistry of Organic Compounds" by E. L. Eliel, S. H. Wilen, and L. N. Mander (Wiley-Interscience, 1994).

[00148] By way of example, compounds of the present disclosure can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereof as appreciated by those skilled in the art. Preferred methods include but are not limited to those methods described below. General procedures to prepare compounds of the instant invention are described in General Scheme 1. An appropriately substituted and protected bicyclic intermediate **1** can be reacted with an appropriately substituted protected amine intermediate **2** under palladium-catalyzed carbon-nitrogen coupling protocols using an appropriate palladium complex, ligand, and base (such as but not limited to: RuPhos 3rd generation palladium precatalyst and cesium carbonate) in a suitable solvent such as toluene at an appropriate temperature (such as 100 °C) to afford intermediate **3**. The protecting group 1 (PG₁; typically a Cbz group) can be removed under suitable deprotection conditions (such as but not

limited to: hydrogen (gas), with palladium on carbon in an appropriate solvent such as methanol, ethanol, or ethyl acetate) to afford amine intermediate **4**. The suitably substituted amine intermediate **4** can be reacted with a suitably substituted carboxylic acid under amide coupling conditions (such as but not limited to: the coupling reagents EDC and HOBt with an appropriate base such as Et₃N or DIEA in a solvent such as DMF or DMA) to afford the penultimate amide intermediate **5**. The protecting group 2 (PG₂; typically a boc group) can be removed under appropriate conditions such as TFA in a solvent such as DCM or HCl in a solvent such as MeOH or dioxane to afford the final compounds **6**. The final compounds can be typically purified by preparative HPLC and isolated as the free base. In the case where mixtures of enantiomers and/or diastereomers are formed, the individual stereoisomers can be purified at an appropriate stage, in many cases by chiral HPLC.


General Scheme 1




General Methods for the Preparation of Selected Intermediates

[00149] General procedures to prepare intermediates of the instant invention are described in Intermediate General Scheme 1. An appropriately substituted leaving group containing starting material **1** (LG₁; typically a bromide) can be reacted with the lithiated chiral auxiliary **2** (formed by reacting **2** with a strong base such as nBuLi) in an appropriate solvent such as THF at low temperature (typically -78 °C) to afford intermediate **3**. Hydrolysis to remove the auxiliary under conditions such as aqueous HCl in a solvent such as acetonitrile, followed by reduction (typically using NaBH₄ in a solvent such as MeOH) can afford amino alcohol **4**. Amino alcohol **4** can be cyclized using a strong base such as NaH in a solvent such as DMSO at low temperature (typically -70 °C) to afford Intermediate **4**. Addition of an appropriate protecting group (PG₁; typically a CBz group) to the corresponding intermediate can result in an appropriately substituted bicyclic intermediate **5**.


Intermediate General Scheme 1.



[00150] General procedures to prepare intermediates of the instant invention are described in Intermediate General Scheme 2. An appropriately substituted ketone **1** can be reacted under reductive amination conditions (typically with ammonium acetate as the amine source, NaBH₃CN as the reductant, and a solvent such as MeOH) to afford the amine intermediate **2**. Amine **2** is then protected (PG₁; typically a CBz group) to afford the appropriately substituted intermediate **3**.

Intermediate General Scheme 2.


[00151] General procedures to prepare intermediates of the instant invention are described in Intermediate General Scheme 3. Acid chloride **1** can be reacted under a lewis acid promoted ethylene addition (typically using $AlCl_3$ and ethylene gas in a solvent such as DCM) to afford ketone **2**. Ketone **2** can then either be converted to an oxime then reduced (using *O*-Methylhydroxylamine hydrochloride, pyridine and EtOH to form the oxime; reduction using hydrogen gas, Raney Ni in EtOH solvent) or reacted under reductive amination conditions (typically with ammonium acetate as the amine source, $NaBH_3CN$ as the reductant, and a solvent such as MeOH) to afford amine **3**. Amine **3** can be then protected (PG_1 ; typically a CBz group) to afford the appropriately substituted intermediate **4**.

Intermediate General Scheme 3.


[00152] General procedures to prepare intermediates of the instant invention are described in Intermediate General Scheme 4. 2-Pyridone **2** can be obtained by the Michael addition of enamino ketone (typically generated in situ by treatment of a mono-protected cyclohexane-1,4-dione such as **1** with ammonia in methanol) with propynoic ester. Pyridone **2** can be converted to incorporate

a leaving group (LG; typically a triflate group which can be obtained by treatment with triflic anhydride in the presence of a base, such as triethylamine). Following removal of the ketone protecting group, ketone **4** can then be reacted under reductive amination conditions (typically with ammonium acetate as the amine source, NaBH₃CN as the reductant, and a solvent such as MeOH) to afford amine **5**. Amine **5** can be then protected (PG₁; typically a Cbz group) to afford the appropriately substituted intermediate **6**.


Intermediate General Scheme 4.




[00153] General procedures to prepare intermediates of the instant invention are described in Intermediate General Scheme 5. Ketone **1** can be condensed with dimethyl-formamide dimethyl acetal in a solvent (such as toluene) to provide enamino ketone **2**. Ketone **2** can be treated with quinidine and a base (such as sodium ethoxide) to provide pyrimidin-2-amine **3**. Pyrimidin-2-amine **3** can be converted to incorporate a leaving group (LG; typically a chloride group which can be obtained by treatment with CuCl₂ and tert-butyl nitrite) to afford the appropriately substituted intermediate **4**.

Intermediate General Scheme 5.

[00154] General procedures to prepare intermediates of the instant invention are described in Intermediate General Scheme 6. An appropriately substituted intermediate **1** can be reacted with an appropriately substituted boronic ester **2** (or boronic acid) under palladium catalyzed carbon-carbon bond forming conditions (using an appropriate palladium catalyst such as Pd(dppf)Cl_2 dichloromethane complex) in a solvent such as 1,4-dioxane/water mixture, and a base such as potassium carbonate at a temperature such as 100 °C) to afford the coupled intermediate **3**. The olefin double bond present in intermediate **3** can then be functionalized using standard methods such as but not limited to: olefin reduction and epoxidation followed by reduction of the resultant epoxide to afford an appropriately substituted intermediate **4**.

Intermediate General Scheme 6.

[00155] General procedures to prepare intermediates of the instant invention are described in Intermediate General Scheme 7. An appropriately substituted intermediate **1** can be reacted under palladium-catalyzed boronic ester forming conditions (using a palladium catalyst such as Pd(dppf)Cl₂-CH₂Cl₂ complex and a boronic ester source such as Bis(pinacolato)diboron in a solvent such as dioxane at a temperature such as 80 °C) to afford the boronic ester **2**. Boronic ester **2** can be reacted in the presence of a suitable oxidant (such as urea-hydrogen peroxide complex in a solvent such as MeOH) to afford the phenol intermediate **3**. The phenol intermediate can then be alkylated with a suitable electrophile (for example by using Mitsunobu-type conditions: such as DIAD with PPh₃) to afford the appropriately substituted intermediate **4**.

Intermediate General Scheme 7.**EXAMPLES**

[00156] The disclosure is further illustrated by the following examples and synthesis schemes, which are not to be construed as limiting this disclosure in scope or spirit to the specific procedures herein described. It is to be understood that the examples are provided to illustrate certain embodiments and that no limitation to the scope of the disclosure is intended thereby. It is to be further understood that resort may be had to various other embodiments, modifications, and equivalents thereof which may suggest themselves to those skilled in the art without departing from the spirit of the present disclosure and/or scope of the appended claims.

Analytical Methods, Materials, and Instrumentation

[00157] Unless otherwise noted, reagents and solvents were used as received from commercial suppliers. Unless otherwise noted, reactions were conducted under an inert atmosphere of nitrogen. Proton nuclear magnetic resonance (NMR) spectra were obtained on either Bruker or Varian spectrometers at 300 or 400 MHz. Spectra are given in ppm (δ) and coupling constants, J, are reported in Hertz. Tetramethylsilane (TMS) was used as an internal standard. Purity and mass spectral data were measured using one of the two following methods. Method 1: Waters Acquity i-class ultra-performance liquid chromatography (UPLC) system with Acquity Photo Diode Array Detector, Acquity Evaporative Light Scattering Detector (ELSD) and Waters ZQ Mass

Spectrometer. Data was acquired using Waters MassLynx 4.1 software and purity characterized by UV wavelength 220 nm, evaporative light scattering detection (ELSD) and electrospray positive ion (ESI). (Column: Acquity UPLC BEH C18 1.7 μ m 2.1 x 50 mm; Flow Rate 0.6 mL/min; Solvent A (95/5/0.1%: 10 mM Ammonium Formate/Acetonitrile/Formic Acid), Solvent B (95/5/0.09%: Acetonitrile/Water/Formic Acid); gradient: 5-100% B from 0 to 2 mins, hold 100% B to 2.2 min and 5%B at 2.21 min). Method 2: SHIMADZU LCMS consisting of an UFLC 20-AD and LCMS 2020 MS detector. (Column: Shim-pack XR-ODS, 2.2 μ m, 3.0 x 50 mm; Solvent: (acetonitrile / water, containing 0.05% NH_4HCO_3)). Preparatory HPLC purifications were conducted as designated below with a Flow Rate of 20 mL/min and detection by UV wavelength 220 nm and 254 nm, unless otherwise noted. The absolute configuration of the separated enantiomers of the compounds in the examples described herein was occasionally determined. In all other cases the absolute configuration of the separated enantiomers was not determined and in those instances the configuration of the resolved materials were arbitrarily assigned as R or S in each case.

Abbreviations used in the following examples and elsewhere herein are:


Abbr	Name
ACN	acetonitrile
atm	atmospheres
Boc	t-butoxycarbonyl
BOP	(benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate
CbzCl	benzyl chloroformate
CDCl_3	deuterated chloroform
CH_2Cl_2	methylene chloride, dichloromethane
CO (g)	carbon monoxide gas
Cs_2CO_3	cesium carbonate
DAST	diethylaminosulfur trifluoride
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
DCM	methylene chloride, dichloromethane
DIEA	diisopropylethylamine
DMA	N,N-dimethylacetamide
DMF	N,N-dimethylformamide
DMSO	dimethylsulfoxide
DPPA	diphenylphosphoryl azide
EDCI	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
ESI	electrospray ionization
Et_2O	diethyl ether

Et ₃ N	triethylamine
EtOAc	ethyl acetate
EtOH	ethanol
h	hours
H ₂ O	water
HATU	1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate
HBTU	N,N,N',N'-tetramethyl-O-(1H-benzotriazol-1-yl)uronium hexafluorophosphate
HCl	hydrochloric acid
HMTA	hexamethylenetetramine
HOBt	hydroxybenzotriazole
IPA	isopropanol
K ₂ CO ₃	potassium carbonate
LDA	lithium diisopropylamide
m-CPBA	3-chloroperbenzoic acid
MeOH	methanol
MgSO ₄	magnesium sulfate
min	minutes
MS	mass spectrometry
MsCl	methanesulfonyl chloride
MTBE	methyl tert-butyl ether
Na ₂ CO ₃	sodium carbonate
Na ₂ SO ₄	sodium sulfate
NaH	sodium hydride
NaHCO ₃	sodium bicarbonate
NaOH	sodium hydroxide
NBS	N-bromosuccinimide
NH ₂ OH-HCl	hydroxylamine hydrochloride
NH ₄ Cl	ammonium chloride
NH ₄ HCO ₃	ammonium bicarbonate
NH ₄ OH	ammonium hydroxide
NMP	N-methyl-2-pyrrolidone
NMR	nuclear magnetic resonance
Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium(0)
Pd(dppf)Cl ₂	[1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II)
Pd(dppf)Cl ₂ -CH ₂ Cl ₂	[1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II)-DCM adduct
Pd/C	palladium on carbon
Pd(OAc) ₂	palladium(II) acetate
pet. ether	petroleum ether
prep-HPLC	preparatory high pressure liquid chromatography
prep-TLC	preparatory thin layer chromatography

PTSA	p-Toluenesulfonic acid
RT	retention time
RuPhos	2-dicyclohexylphosphino-2',6'-diisopropoxybiphenyl
RuPhos Pd G3	(2-Dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate
SFC	supercritical fluid chromatography
TBS	tert-butyl(dimethyl)silyl
t-BuOH	tert-butanol
t-BuOK	potassium tert-butoxide
t-BuXPhos Pd G4	methanesulfonato(2-di-t-butylphosphino-2',4',6'-tri-i-propyl-1,1'-biphenyl)(2'-methylamino-1,1'-biphenyl-2-yl)palladium(II) DCM adduct
TEDA	triethylenediamine
TFA	trifluoroacetic acid
TFAA	trifluoroacetic anhydride
THF	THF
XPhos Pd G3	(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate

PREPARATION OF INTERMEDIATES

Intermediates 1, 2, and 3. Benzyl (7-bromochroman-3-yl)carbamate, Benzyl (R)-(7-bromochroman-3-yl)carbamate, and Benzyl (S)-(7-bromochroman-3-yl)carbamate



Step 1. 7-Bromo-3-nitro-2H-chromene

[00158] A mixture of 4-bromo-2-hydroxybenzaldehyde (21.6 g, 108 mmol), 1,3-dihydro-2-benzofuran-1,3-dione (32 g, 216 mmol) and dibutylamine (7.0 g, 54 mmol) in toluene (800 mL) was heated to reflux under an atmosphere of N₂. 2-Nitroethan-1-amine (50 g, 550 mmol) was added in portions over 2 h. The mixture was stirred at reflux overnight using a Dean-Stark apparatus. After cooling to room temperature, the solids were filtered out. Eight batches were thus run in parallel and the filtrate from the eight batches were combined and concentrated under vacuum. The residue was diluted with EtOAc (2 L) and washed with 1N NaOH (2 L). The organic layer was dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:6 EtOAc/pet. ether) afforded 7-bromo-3-nitro-2H-chromene as a yellow solid. MS: (ESI, m/z): 256, 258 [M+H]⁺.

Step 2. 7-Bromochroman-3-amine

[00159] To a solution of 7-bromo-3-nitro-2H-chromene (27 g, 106 mmol) in THF (300 mL) were added BH₃ (1M in THF, 600 mL, 600 mmol) and NaBH₄ (201 mg, 5.3 mmol). The mixture was stirred overnight at 65 °C. After cooling to room temperature, the reaction was then quenched by the addition of 600 mL of MeOH and stirred for 8 h at 80 °C. After cooling to room temperature, the mixture was concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 0% to 50% B over 40 min) to afford 7-bromochroman-3-amine as a white solid. MS: (ESI, m/z): 228, 230 [M+H]⁺.

Step 3. Benzyl (7-bromochroman-3-yl)carbamate (Intermediate 1)

[00160] A solution of K₂CO₃ (19.3 g, 140 mmol) in water (150 mL) was added to a solution of 7-bromochroman-3-amine (16.0 g, 70.1 mmol) in EtOAc (300 mL). Benzyl chloroformate (17.8 g, 104 mmol) was added at -10 °C and the reaction mixture was stirred for 30 min at room temperature. The mixture was diluted with EtOAc (200 mL). The organic layer was collected, dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. The residue was washed with 1:1 EtOAc/pet. ether (200 mL) to give benzyl-(7-bromochroman-3-yl)carbamate as a white solid. MS: (ESI, m/z): 362, 364 [M+H]⁺.


Step 4. Benzyl (R)-(7-bromochroman-3-yl)carbamate (Intermediate 2) and Benzyl (S)-(7-bromochroman-3-yl)carbamate (Intermediate 3)

[00161] The racemate benzyl (7-bromochroman-3-yl)carbamate (12.5 g, 34.6 mmol) was separated by SFC (Column: ChiralArt Amylose-SA, 2x25 cm, 5 μ m; Mobile phase A: CO₂, 80%, B: EtOH, 20%; Flow rate: 40 mL/min) to afford the title compounds as follows: benzyl (R)-(7-bromochroman-3-yl)carbamate (first eluting isomer, RT = 7.98 min) as a white solid and benzyl (S)-(7-bromochroman-3-yl)carbamate (second eluting isomer, RT = 9.21 min) as a white solid.

First eluting isomer: ¹H NMR (CDCl₃, 400 MHz) δ (ppm): 7.35-7.32 (m, 5H), 7.03-7.01 (m, 2H), 6.90 (d, J = 8.8 Hz, 1H), 5.22-5.10 (m, 3H), 4.25 (s, 1 H), 4.17-4.09 (m, 2H), 3.04 (dd, J = 16.8 Hz, 4.8 Hz, 1H), 2.73 (d, J = 16.8 Hz, 1H). MS: (ESI, m/z): 362, 364 [M+H]⁺.

Second eluting isomer: ¹H NMR (CDCl₃, 400 MHz) δ (ppm): 7.35-7.32 (m, 5H), 7.03-7.01 (m, 2H), 6.90 (d, J = 8.8 Hz, 1H), 5.22-5.07 (m, 3H), 4.25 (s, 1 H), 4.18-4.09 (m, 2H), 3.04 (dd, J = 16.8 Hz, 4.8 Hz, 1H), 2.73 (d, J = 16.80 Hz, 1H). MS: (ESI, m/z): 362, 364 [M+H]⁺.

Intermediate 4-1. tert-Butyl 4-(3-amino-8-fluorochroman-7-yl)piperazine-1-carboxylate



Step 1. 3,4-Difluoro-2-hydroxybenzaldehyde

[00162] Into a 500-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen was added a 2,3-difluorophenol (10.0 g, 75.33 mmol), ACN (200 mL), HCHO (23.06 g, 738 mmol), Et₃N (21.0 mL, 146 mmol), and MgCl₂ (14.6 g, 150.28 mmol). The resulting solution was stirred for 16 h at 60 °C. The reaction mixture was cooled to 26 °C, then was diluted with 200 mL of water. The resulting solution was extracted with ethyl acetate (3 x 100 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and concentrated under

vacuum to afford a residue that was purified by silica gel chromatography with ethyl acetate/pet. ether (1:4) to afford 3,4-difluoro-2-hydroxybenzaldehyde as light yellow oil.

Step 2. tert-Butyl 4-(2-fluoro-4-formyl-3-hydroxyphenyl)piperazine-1-carboxylate

[00163] Into a 250-mL round-bottom flask was added 3,4-difluoro-2-hydroxybenzaldehyde (5 g, 31.63 mmol), tert-butyl piperazine-1-carboxylate (5.9 g, 31.68 mmol), DMSO (100 mL), and DIEA (6.1 g, 47.20 mmol). The resulting solution was stirred for 6 h at 120 °C. After cooling to room temperature, the reaction was then quenched by the addition of 100 mL of water. The resulting solution was extracted with ethyl acetate (3x100 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum to afford a residue that was purified by silica gel chromatography with ethyl acetate/pet. ether (0-30%) to afford tert-butyl 4-(2-fluoro-4-formyl-3-hydroxyphenyl)piperazine-1-carboxylate as yellow oil.

Step 3. tert-Butyl 4-(8-fluoro-3-nitro-2H-chromen-7-yl)piperazine-1-carboxylate

[00164] Into a 100-mL round-bottom flask was added tert-butyl 4-(2-fluoro-4-formyl-3-hydroxyphenyl)piperazine-1-carboxylate (185 mg, 0.48 mmol), 1,3-dihydro-2-benzofuran-1,3-dione (166 mg, 1.06 mmol, 95%), 2-nitroethanol (104 mg, 1.08 mmol), dibutylamine (37 mg, 0.27 mmol), and toluene (10 mL). The resulting solution was stirred for 8 h at 100 °C in an oil bath. The reaction mixture was cooled to room temperature, then was quenched by addition of 5 mL water, and extracted with DCM (3x10 mL). The organic layers were combined, washed with brine (3x10 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum to afford a residue that was purified by silica gel chromatography with ethyl acetate/pet. ether (1:3) to afford tert-butyl 4-(8-fluoro-3-nitro-2H-chromen-7-yl)piperazine-1-carboxylate as a red solid.


Step 4. tert-Butyl 4-(3-amino-8-fluorochroman-7-yl)piperazine-1-carboxylate

[00165] Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was added tert-butyl 4-(8-fluoro-3-nitro-2H-chromen-7-yl)piperazine-1-carboxylate (150 mg, 0.39 mmol), and THF (20 mL). Borane-THF complex (4 mL, 4 mmol) was added dropwise with stirring at 0 °C. To this reaction mixture was added NaBH₄ (73 mg, 1.93 mmol). The resulting solution was stirred for 12 h at 65°C, then was quenched by the addition of methanol (20 mL), and concentrated under vacuum to afford a residue that was purified by silica

gel chromatography with ethyl acetate/pet. ether (0-100%) to afford tert-butyl 4-(3-amino-8-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl)piperazine-1-carboxylate as yellow oil.

[00166] The following intermediate in Table 1 was prepared using standard chemical manipulations and procedures similar to those used for the preparation of Intermediate 4-1.


Table 1:

Intermediate Number	Structure and Compound Name	LRMS m/z [M+H] ⁺
4-2 ¹	 tert-Butyl 4-(3-amino-6-fluorochroman-7-yl)piperazine-1-carboxylate	352

¹Notes on procedures: Step 1 was not necessary.

Intermediate 5-1. Benzyl (R)-(7-bromo-5-fluorochroman-3-yl)carbamate

Method 1.



Step 1. 5-Bromo-2-(bromomethyl)-1,3-difluorobenzene

[00167] A mixture of 48% HBr (130 mL, 1149 mmol) and (4-bromo-2,6-difluorophenyl)methanol (40 g, 179 mmol) was heated to reflux overnight. After cooling to room temperature, the reaction mixture was poured into 80 mL of water and was extracted with hexanes (2 x 300 mL). The combined organic layers were washed with sodium bicarbonate solution, dried over MgSO₄, filtered, and concentrated to afford 5-bromo-2-(bromomethyl)-1,3-difluorobenzene. ¹H NMR (CDCl₃, 300 MHz) δ(ppm): 6.94-7.22 (m, 2H), 4.46 (s, 2H).

Step 2. (2R,5S)-2-(4-bromo-2,6-difluorobenzyl)-5-isopropyl-3,6-dimethoxy-2,5-dihydropyrazine

[00168] A solution of nBuLi (6.78 mL, 10.86 mmol, 1.6 M) in hexanes was added dropwise to a solution of (S)-2-isopropyl-3,6-dimethoxy-2,5-dihydropyrazine (2.0 g, 10.86 mmol) in 20 mL of THF at -78 °C. After stirring for 30 min at -78 °C, a solution of 5-bromo-2-(bromomethyl)-1,3-difluorobenzene (3.10 g, 10.86 mmol) in 10 mL of THF was added and the reaction mixture was stirred at -78 °C for 3 h. Then 20 mL of saturated NH₄Cl solution was added. After the reaction mixture was warmed to room temperature, 150 mL of water was added and the mixture was extracted with EtOAc three times. The combined organic layers were dried over MgSO₄, filtered, and concentrated. Purification by silica gel chromatography (eluting with 0 to 15% EtOAc/Hexanes) afforded (2R,5S)-2-(4-bromo-2,6-difluorobenzyl)-5-isopropyl-3,6-dimethoxy-2,5-dihydropyrazine. ¹H NMR (CDCl₃, 300 MHz) δ(ppm): 7.03 (d, *J* = 7.04 Hz, 2H), 4.14-4.32 (m, 1H), 3.71 (s, 3H), 3.58 (s, 3H), 3.13-3.33 (m, 1H), 2.79-2.93 (m, 1H), 2.13-2.33 (m, 1H), 1.00 (d, *J* = 7.04 Hz, 3H), 0.64 (d, *J* = 7.04 Hz, 3H). MS: (ESI, *m/z*): 389, 391 [M+H]⁺.

Step 3. Methyl (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propanoate

[00169] A solution of (2R,5S)-2-(4-bromo-2,6-difluorobenzyl)-5-isopropyl-3,6-dimethoxy-2,5-dihydropyrazine (3.1 g, 8.02 mmol) in acetonitrile (60 mL) was treated with HCl (53.5 mL, 16.04 mmol, 0.3 N). The reaction mixture was stirred at room temperature for 60 min. The reaction was made basic with sat. aq. NaHCO₃ solution and the mixture was extracted with CH₂Cl₂ three times. The combined organic layers were dried over MgSO₄, filtered, and concentrated. Purification by silica gel chromatography (eluting with 0 to 100% EtOAc/Hexanes) afforded methyl (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propanoate. ¹H NMR (CDCl₃, 300 MHz) δ(ppm): 7.07 (d, *J* = 6.74 Hz, 2H), 3.54-3.80 (m, 4H), 3.01-3.24 (m, 1H), 2.93 (s, 1H), 1.62 (br s, 2H). MS: (ESI, *m/z*): 294, 296 [M+H]⁺.

Step 4. (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propan-1-ol

[00170] To a solution of methyl (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propanoate (2.3 g, 7.85 mmol) in MeOH (70 mL) at room temperature was added NaBH₄ (1.043 g, 27.57 mmol) in portions. The mixture was stirred at room temperature overnight. Water was added and the MeOH was removed under reduced pressure. The aqueous mixture was extracted with chloroform three times. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated to afford (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propan-1-ol. ¹H NMR (CDCl₃, 300 MHz) δ(ppm): 7.08 (d, *J* = 6.74 Hz, 2H), 3.54-3.72 (m, 1H), 3.36 (dd, *J* = 10.55, 7.62 Hz, 1H), 3.09 (br s, 1H), 2.51-2.86 (m, 2H), 1.74 (br s, 3H). MS: (ESI, *m/z*): 266, 268 [M+H]⁺.

Step 5. (R)-7-bromo-5-fluorochroman-3-amine

[00171] To a solution of (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propan-1-ol (500 mg, 1.88 mmol) in DMSO (3 mL) at room temperature was added NaH (113 mg, 2.82 mmol). The mixture was stirred at room temperature for 30 min, then at 70 °C for 30 min, and kept stirring at 50 °C overnight. After cooling to room temperature, 30 mL of water, and the mixture was extracted with EtOAc three times. The combined organic layers were washed with water, dried over Na₂SO₄, filtered, and concentrated to afford crude (R)-7-bromo-5-fluorochroman-3-amine. ¹H NMR (CDCl₃, 300 MHz) δ(ppm): 6.61-6.92 (m, 2H), 4.01-4.32 (m, 1H), 3.67-3.88 (m, 1H), 3.35 (ddt, *J* = 7.00, 3.19, 1.80, 1.80 Hz, 1H), 2.79-3.03 (m, 1H), 2.46 (br d, *J* = 6.74 Hz, 1H), 1.40-1.89 (m, 2H). MS: (ESI, *m/z*): 246, 248 [M+H]⁺.


Step 6. Benzyl (R)-(7-bromo-5-fluorochroman-3-yl)carbamate

[00172] To a solution of (R)-7-bromo-5-fluorochroman-3-amine (467 mg, 1.90 mmol) and saturated solution of sodium hydrogen carbonate (10 mL) in MeOH (20 mL) at 0 °C was added benzyl chloroformate (0.405 mL, 2.85 mmol) dropwise. The mixture was stirred overnight, allowing the temperature to warm to room temperature. 20 mL of water was added and the mixture was extracted with EtOAc three times. The combined organic layers were washed with water, dried over Na₂SO₄, filtered, and concentrated. Purification by silica gel chromatography (eluting with 7% to 60% EtOAc/Hexanes) afforded benzyl (R)-(7-bromo-5-fluorochroman-3-yl)carbamate. ¹H NMR (CDCl₃, 300 MHz) δ(ppm): 7.34 (s, 5 H), 6.65-6.97 (m, 2H), 5.10 (s, 2H),

4.90-5.08 (m, 1H), 4.26 (br s, 1H), 3.97-4.21 (m, 3H), 2.89 (m, 1H), 2.78 (m, 1H). MS: (ESI, m/z): 380, 382 [M+H]⁺.

Intermediate 5-1. Benzyl (R)-(7-bromo-5-fluorochroman-3-yl)carbamate

Method 2.



Step 1. Diethyl 2-acetamido-2-(4-bromo-2,6-difluorobenzyl)malonate

[00173] To a stirred solution of diethyl 2-acetamidomalonate (59.2 g, 0.273 mol) in DMF (500 mL) was added t-BuOK (33.1 g, 0.295 mol) in portions at room temperature. The mixture was stirred at room temperature for 1 h and 5-bromo-2-(bromomethyl)-1,3-difluorobenzene (65.0 g, 0.227 mol) was added. The reaction mixture was stirred at room temperature for 3 h. Water (2000 mL) was added slowly and the mixture was stirred for 1 h. The resulting precipitate was collected by filtration, washed with water (3 x 250 mL) and dried under vacuum to give diethyl 2-acetamido-2-(4-bromo-2,6-difluorobenzyl)malonate as an off-white solid.

Step 2. 2-Acetamido-3-(4-bromo-2,6-difluorophenyl)propanoic acid

[00174] To a stirred solution of diethyl 2-acetamido-2-(4-bromo-2,6-difluorobenzyl)malonate (80 g, 0.189 mol) in ethanol (500 mL) was added a solution of NaOH (30 g, 0.758 mol) in water (500 mL). The reaction mixture was heated at reflux for 5 h and then cooled to room temperature. The mixture was adjusted to pH=5-6 with 2N aqueous HCl and heated at reflux overnight. The mixture was cooled to room temperature and adjusted to pH=8-9 with 10%

aqueous NaOH. The resulting mixture was washed with MTBE (300 mL) and the aqueous phase was adjusted to pH 2-3 with 2N aqueous HCl and then extracted with EtOAc (500 mL×2). The combined extracts were washed with brine (300 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated to dryness. The residue was treated with a mixture of EtOAc (100 mL) and pet. ether (150 mL) under stirring for 1 h. The resulting precipitate was collected by filtration, washed with pet. ether and dried under vacuum to give 2-acetamido-3-(4-bromo-2,6-difluorophenyl)propanoic acid as a white solid. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 12.86 (br s, 1H), 8.31 (br s, 1H), 7.43 (d, *J* = 6.8 Hz, 2H), 4.43 (m, 1H), 3.08-2.87 (m, 2H), 1.77 (s, 3H). MS: (ESI, *m/z*): 322, 324 [M+H]⁺.

Step 3. (R)-2-Acetamido-3-(4-bromo-2,6-difluorophenyl)propanoic acid

[00175] To a suspension of 2-acetamido-3-(4-bromo-2,6-difluorophenyl)propanoic acid (55 g, 0.171 mol) in distilled water (1.1 L) was added 10% aqueous NaOH dropwise to adjust pH to 8.5. The mixture was heated to 35-38 °C and *L*-acylase (11.0 g) was added. The reaction mixture was stirred at this temperature for 48 h while keeping the pH at 8.5 with 10% aqueous NaOH. The mixture was adjusted to pH 4-5 with 2N aqueous HCl and activated carbon (2 g) was added. The mixture was heated at 60 °C for 2 h and then cooled to room temperature. The mixture was adjusted to pH 9.5-10 with 10% aqueous NaOH and filtered. The filtrate was adjusted to pH 2-3 with 2N aqueous HCl and then extracted with EtOAc (400 mL x 2). The combined extracts were washed with 0.5N aqueous HCl (200 mL x 2) and brine (200 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated to dryness. The residue was treated with a mixture of EtOAc (60 mL) and pet. ether (80 mL) under stirring for 1 h. The resulting precipitate was collected by filtration, washed with pet. ether and dried under vacuum to give (R)-2-acetamido-3-(4-bromo-2,6-difluorophenyl)propanoic acid as a white solid.

Step 4. (R)-2-Amino-3-(4-bromo-2,6-difluorophenyl)propanoic acid hydrochloride

[00176] A mixture of (R)-2-acetamido-3-(4-bromo-2,6-difluorophenyl)propanoic acid (26 g, 80.7 mmol) in 6N aqueous HCl (260 mL) was heated under reflux for 5 h. The mixture was concentrated and the residue was dried under vacuum at 50 °C to provide crude (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propanoic acid hydrochloride as a white solid.

Step 5. (R)-Methyl 2-amino-3-(4-bromo-2,6-difluorophenyl)propanoate

[00177] To a solution of (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propanoic acid hydrochloride (25.4 g, 80.7 mmol) in MeOH (125 mL) was added MeOH/HCl (8M, 125 mL). The reaction mixture was stirred at room temperature overnight and concentrated to dryness. The residue was suspended in 5% aqueous Na₂CO₃ (250 mL) and then extracted with EtOAc (250 mL x 2). The combined extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated to give (R)-methyl 2-amino-3-(4-bromo-2,6-difluorophenyl)propanoate as an oil. ¹H NMR (CDCl₃, 400 MHz) δ(ppm): 7.08 (m, 2H), 3.73 (s, 3H), 3.70 (m, 1H), 3.09-2.87 (m, 2H), 1.55 (br s, 2H). MS: (ESI, m/z): 294, 296 [M+H]⁺.

Step 6. (R)-2-Amino-3-(4-bromo-2,6-difluorophenyl)propan-1-ol

[00178] To a stirred solution of (R)-methyl 2-amino-3-(4-bromo-2,6-difluorophenyl)propanoate (23.5 g, 79.9 mmol) in MeOH (500 mL) was added NaBH₄ (6.08 g, 159.9 mmol) portionwise. The reaction mixture was stirred at room temperature for 3 h and NaBH₄ (1.52 g, 39.9 mmol) was added. The reaction mixture was stirred at room temperature overnight. Water (500 mL) was added and MeOH was removed by evaporation under vacuum. The resulting mixture was extracted with CH₂Cl₂ (250 mL x 3) and the combined extracts were dried over anhydrous Na₂SO₄, filtered, and concentrated to afford crude (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propan-1-ol as a white solid. MS: (ESI, m/z): 266, 268 [M+H]⁺.

Step 7. (R)-7-Bromo-5-fluorochroman-3-amine

[00179] To a stirred solution of (R)-2-amino-3-(4-bromo-2,6-difluorophenyl)propan-1-ol (18.5 g, 69.5 mmol) in DMSO (100 mL) was added NaH (60% in mineral oil, 4.17 g) at room temperature. The reaction mixture was stirred at 35 °C for 3 h and ice-water (500 mL) was added carefully to quench the reaction. The resulting mixture was extracted with EtOAc (300 mL x 3) and the combined extracts were washed with brine (200 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated to give crude (R)-7-bromo-5-fluorochroman-3-amine. MS: (ESI, m/z): 246, 248 [M+H]⁺.


Step 8. Benzyl (R)-(7-bromo-5-fluorochroman-3-yl)carbamate

[00180] To a stirred mixture of crude (R)-7-bromo-5-fluorochroman-3-amine (17 g, 69.5 mmol) and saturated aqueous NaHCO₃ (200 mL) in MeOH (400 mL) was added benzyl chloroformate (17.7 g, 104.2 mmol) dropwise. The reaction mixture was stirred at room temperature overnight and

then diluted with water (500 mL). The resulting mixture was extracted with EtOAc (300 mL x 2) and the combined extracts were washed with water (200 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated. The residue was purified by silica gel chromatography to give benzyl (R)-(7-bromo-5-fluorochroman-3-yl)carbamate. MS: (ESI, m/z): 380, 382 [M+H]⁺.

[00181] The following intermediate in Table 2 may be prepared using standard chemical manipulations and procedures similar to Method 2 of Intermediate 5-1.

Table 2:


Intermediate Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺
5-2 ¹	 7-bromo-5-fluorochroman-4,4-d ₂ -3-amine	248, 250

¹Notes on procedures: Step 3 and Step 8 were not performed. In Step 1, 5-bromo-2-(bromomethyl-d₂)-1,3-difluorobenzene was prepared from methyl 4-bromo-2,6-difluorobenzoate in two steps: Methanol (0.806 ml, 19.92 mmol) was carefully added dropwise to a stirring solution of methyl 4-bromo-2,6-difluorobenzoate (5 g, 19.92 mmol), sodium tetrahydroborate-d₄ (0.834 g, 19.92 mmol) in THF (15 mL). The reaction was then allowed to stir at 70 °C for 2 h. The reaction was cooled to room temperature and 10 mL sat. aq. NH₄Cl was added. The reaction was allowed to stir at room temperature for 2 h. The organic layer was separated. The aqueous layer was extracted with 2 x 15 mL DCM. The organic layers were combined, dried over Na₂SO₄, filtered, and concentrated under vacuum to afford (4-bromo-2,6-difluorophenyl)methan-d₂-ol.

A solution of (4-bromo-2,6-difluorophenyl)methan-d₂-ol (3.00 g, 13.33 mmol) and PBr₃ (14.21 mL, 14.21 mmol) in DCM (30 mL) was stirred at 40 °C for 30 min. After cooling to room temperature, the reaction was quenched by the addition of water (7.5 mL). The resulting mixture was extracted with DCM (3 x 10 mL) and the organic layers were combined, dried over Na₂SO₄, filtered, and concentrated to afford a crude pale yellow oil. The oil was purified by normal phase chromatography using Biotage (KP-SIL 50g, 2% EtOAc/hexanes up to 25% EtOAc/hexanes. Desired fractions were combined and concentrated to afford 5-bromo-2-(bromomethyl-d₂)-1,3-difluorobenzene. ¹H NMR (CDCl₃, 300 MHz) δ(ppm): 1.54 (s, 1H) 4.46 (br d, *J* = 3.8 Hz, 1H) 7.05-7.21 (m, 2H).

Intermediate 6. tert-Butyl (R)-4-(3-amino-8-bromo-5-fluorochroman-7-yl)piperazine-1-carboxylate and

Intermediate 7-1. tert-Butyl (R)-4-(3-amino-6-cyano-5-fluorochroman-7-yl)piperazine-1-carboxylate



Step 1: tert-Butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-8-bromo-5-fluorochroman-7-yl)piperazine-1-carboxylate and tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-6-bromo-5-fluorochroman-7-yl)piperazine-1-carboxylate

[00182] Into a 20-mL vial, was added tert-butyl 4-[(3R)-3-[[[(benzyloxy)carbonyl]amino]-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate (570 mg, 1.17 mmol), THF (10 mL), and NBS (314 mg, 1.76 mmol). The resulting solution was stirred for 2 h at 25 °C, then was quenched by the addition of water (30 mL). The resulting mixture was extracted with ethyl acetate (3 x 30 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by prep-HPLC (Column: XSelect CSH Prep C18 OBD, 5 µm, 19x150 mm; Mobile phase A: water (0.05% TFA), B: ACN; Gradient: 45% B increasing to 70% B within 15 min). The collected fraction was concentrated under vacuum to afford tert-butyl 4-[(3R)-3-[[[(benzyloxy)carbonyl]amino]-8-bromo-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate (peak 1) (**Intermediate 6**) as an off-white solid and tert-butyl 4-[(3R)-3-[[[(benzyloxy)carbonyl]amino]-6-bromo-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate (peak 2) as off-white solid. MS (ESI, m/z): 564, 566 [M+H]⁺. These two compounds were carried on independently into subsequent synthetic steps.

Step 2: tert-Butyl 4-[(3R)-3-[[[(benzyloxy)carbonyl]amino]-6-cyano-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate

[00183] Into a 10-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was added tert-butyl 4-[(3R)-3-[[[(benzyloxy)carbonyl]amino]-6-bromo-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate (80 mg, 0.14 mmol), Zn(CN)₂ (13 mg, 0.11 mmol), Pd(PPh₃)₄ (8 mg, 0.01 mmol), PPh₃ (7 mg, 0.03 mmol), and NMP (5 mL). The resulting solution was stirred for 1 h at 120 °C. After cooling to 25 °C, the reaction was quenched by the addition of 10 mL of water. The resulting mixture was extracted with 3 x 20 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to afford a residue that was purified via reverse phase chromatography (Column: C18 silica gel; Mobile phase A: 0.1% TFA in H₂O, B: ACN; Flow rate: 50 mL/min; Gradient: 0% B increasing to 80% B within 30 min). The collected fractions were concentrated under vacuum to afford tert-butyl 4-[(3R)-3-[[[(benzyloxy)carbonyl]amino]-6-cyano-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate as an off-white solid. MS (ESI, m/z): 511 [M+H]⁺.


Step 3: tert-Butyl 4-[(3R)-3-amino-6-cyano-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate

[00184] Into a 25-mL round-bottom flask purged and maintained with nitrogen, was placed tert-butyl 4-[(3R)-3-[[[(benzyloxy)carbonyl]amino]-6-cyano-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate (40 mg, 0.08 mmol), ethyl acetate (4 mL), and 10% Palladium on carbon (40 mg,). The resulting mixture was stirred for 2 h at 25 °C under hydrogen atmosphere. The solids were removed by filtration through Celite and the filtrate was concentrated under vacuum to afford tert-butyl 4-[(3R)-3-amino-6-cyano-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate (**Intermediate 7-1**) as yellow oil. MS (ESI, m/z): 377 [M+H]⁺.

The following intermediate in Table 3 may be prepared using standard chemical manipulations and procedures similar to those used for the preparation of Intermediate 7-1.

Table 3:


Intermediate Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺

7-2	 tert-butyl (R)-4-(3-amino-8-cyano-5-fluorochroman-7-yl)piperazine-1-carboxylate	377

Intermediate 8. Benzyl (6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate;

Intermediate 9. Benzyl (S)-(6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate and

Intermediate 10. Benzyl (R)-(6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate



Step 1. 6-Bromo-1,2,3,4-tetrahydronaphthalen-2-amine

[00185] A solution of 6-bromo-3,4-dihydronaphthalen-2(1H)-one (5 g, 22.21 mmol), NH_4OAc (13.8 g, 179 mmol), and NaBH_3CN (1.68 g, 26.67 mmol) in MeOH (250 mL) was stirred for 1 h at room temperature. The reaction mixture was acidified with 2N HCl solution to pH 4-5 and was concentrated under vacuum. The residual solution was washed with CH_2Cl_2 (200 mL x 2). The aqueous layer was basified with 1N NaOH solution to pH 10, then extracted with CH_2Cl_2 (200 mL x 2). The combined organic layer was dried over Na_2SO_4 , filtered, and concentrated under vacuum to afford 6-bromo-1,2,3,4-tetrahydronaphthalen-2-amine as a yellow oil. MS: (ESI, m/z): 226, 228 $[\text{M}+\text{H}]^+$.

Step 2. Benzyl (6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate (Intermediate 8)

[00186] A solution of 6-bromo-1,2,3,4-tetrahydronaphthalen-2-amine (1.8 g, 7.96 mmol), benzyl chloroformate (1.6 g, 9.55 mmol), and Cs_2CO_3 (3 g, 21.71 mmol) in THF (20 mL) and

water (20 mL) was stirred at 60 °C overnight. After cooling to room temperature, the reaction mixture was extracted with EtOAc (30 mL x 3). The combined organic layer was dried over Na₂SO₄, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:100 to 1:3 EtOAc/pet. ether) afforded benzyl (6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate (**Intermediate 8**) as a solid. MS: (ESI, m/z): 360, 362 [M+H]⁺.


Step 3. Benzyl (S)-(6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate (Intermediate 9) and Benzyl (R)-(6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate (Intermediate 10)

[00187] The racemate benzyl (6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate was separated by SFC (Column: Chiralpak IA-SFC-03, 5x25 cm, 5 μm; Mobile phase A: CO₂, B: MeOH; Flow rate: 170 mL/min) to afford the title compounds as follows: benzyl (S)-(6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate (first eluting isomer, RT = 6.54 min) as a white solid and benzyl (R)-(6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate (second eluting isomer, RT = 9.06 min) as a white solid.

First eluting isomer: ¹H NMR (CDCl₃, 400 MHz) δ(ppm): 7.30-7.21 (m, 5H), 7.17-7.13 (m, 2H), 6.85-6.83 (d, *J* = 8.00 Hz, 1H), 5.02 (s, 2H), 4.70 (br, 1H), 3.95 (m, 1H), 3.01-2.96 (dd, *J* = 4.00 Hz, 16.00 Hz, 2H), 2.79-2.75 (m, 2H), 2.53-2.47 (m, 1H), 2.00-1.97 (m, 1H), 1.68-1.66 (m, 1H). MS: (ESI, m/z): 360, 362 [M+H]⁺.

Second eluting isomer: ¹H NMR (CDCl₃, 400 MHz) δ(ppm): 7.30-7.23 (m, 5H), 7.17-7.13 (m, 2H), 6.84-6.82 (d, *J* = 8.00 Hz, 1H), 5.02 (s, 2H), 4.70 (br, 1H), 4.06-3.95 (m, 1H), 3.01-2.96 (dd, *J* = 4.00 Hz, 16.0 Hz, 2H), 2.79-2.75 (m, 2H), 2.51-2.47 (m, 1H), 2.00-1.96 (m, 1H), 1.68-1.66 (m, 1H). MS: (ESI, m/z): 360, 362 [M+H]⁺.

Intermediate 11-1. Benzyl (6-bromo-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate



Step 1. 2-(4-Bromo-2-fluorophenyl)acetyl chloride

A 250-mL round-bottom flask was charged with 2-(4-bromo-2-fluorophenyl)acetic acid (10 g, 42.05 mmol), DCM (50 mL), and thionyl chloride (6.3 mL, 85.11 mmol). The resulting solution was stirred for 16 h at 40 °C. After cooling to 25 °C, the reaction mixture was concentrated under vacuum to afford 2-(4-bromo-2-fluorophenyl)acetyl chloride as brown oil.

Step 2. 6-Bromo-8-fluoro-3,4-dihydronaphthalen-2(1H)-one

A 500-mL round-bottom flask was charged with 2-(4-bromo-2-fluorophenyl)acetyl chloride (5.0 g, 18.49 mmol) and DCM (100 mL). AlCl₃ (7.15 g, 53.09 mmol) was then added in portions at 0 °C. The resulting mixture was stirred for 10 min at 0 °C, then a gentle stream of ethylene gas was bubble into the reaction mixture for 5 h at 0 °C. The reaction mixture was poured into ice and concentrated hydrochloric acid (5 mL) was added. The resulting solution was extracted with DCM (3 x 50 mL), the organic layers combined, dried over anhydrous sodium sulfate and concentrated under vacuum to afford a crude residue that was purified by column chromatography eluting with ethyl acetate/pet. ether (1:2) to afford 6-bromo-8-fluoro-3,4-dihydronaphthalen-2(1H)-one as a brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ(ppm): 2.49 (t, *J*=8.0Hz, 2H), 3.08 (t, *J*=8.0Hz, 2H), 3.49 (s, 2H), 7.40-7.42 (m, 2H).

Step 3. (2E)-6-Bromo-8-fluoro-N-methoxy-1,2,3,4-tetrahydronaphthalen-2-imine

A 250-mL round-bottom flask was charged with 6-bromo-8-fluoro-3,4-dihydronaphthalen-2(1H)-one (4.2 g, 16.07 mmol), the HCl salt of O-methylhydroxylamine (2.16 g, 25.60 mmol), ethanol (50 mL) and pyridine (5 mL, 61.50 mmol). The resulting mixture was stirred for 16 h at 80 °C in an oil bath, then was cooled to 25 °C. After cooling, the resulting mixture was concentrated under vacuum to afford a residue that was purified by column chromatography eluting with ethyl acetate/pet. ether (1:2) to afford (2E)-6-bromo-8-fluoro-N-methoxy-1,2,3,4-tetrahydronaphthalen-2-imine as a brown solid. MS: (ESI, *m/z*) 272, 274 [M+H]⁺.

Step 4. 6-Bromo-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-amine

A 250-mL round-bottom flask that was purged with nitrogen was charged with (2E)-6-bromo-8-fluoro-N-methoxy-1,2,3,4-tetrahydronaphthalen-2-imine (3.5 g, 11.58 mmol, 90%), ethanol (50 mL) and Raney Ni (2.0 g, 23.11 mmol). To this hydrogen (g) was introduced in. The resulting mixture was stirred for 48 h at 25 °C. The solids were removed by filtration over Celite. The filtrate was concentrated under vacuum to afford a residue that was purified by column chromatography




eluting with DCM/methanol (10:1) to afford 6-bromo-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-amine as a brown solid. MS: (ESI, m/z): 244, 246 $[M+H]^+$.


Step 5. Benzyl (6-bromo-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate

A 100-mL round-bottom flask was charged with 6-bromo-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-amine (930 mg, 3.43 mmol), ethyl acetate (15 mL), water (15 mL), potassium carbonate (1.58 g, 11.32 mmol), and benzyl chloroformate (780 mg, 4.53 mmol). The resulting mixture was stirred for 16 h at 60 °C in an oil bath. After cooling to 25 °C, the reaction was diluted with water (30 mL). The resulting solution was extracted with ethyl acetate (3 x 40 mL), the organic layers combined, dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum to afford a residue that was purified by column chromatography eluting with ethyl acetate/pet. ether (1:5) to afford benzyl (6-bromo-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate as an off-white solid. MS: (ESI, m/z): 378, 380 $[M+H]^+$.

[00188] The following intermediates in Table 4 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Intermediate 11-1.


Table 4:

Intermediate Number	Structure and Compound Name	LRMS m/z $[M+H]^+$
11-2 ¹	 benzyl (S)-(6-bromo-7-methyl-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate	374, 376
11-3 ¹	 benzyl (R)-(6-bromo-7-methyl-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate	374, 376
11-4 ¹	 benzyl (S)-(6-bromo-7-methyl-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate	374, 376

Intermediate Number	Structure and Compound Name	LRMS m/z [M+H] ⁺
	benzyl (S)-(6-bromo-5-methyl-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate	
11-5 ¹	 benzyl (R)-(6-bromo-5-methyl-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate	374, 376

¹**Notes on procedures:** In Step 2, a mixture of regioisomers 6-bromo-7-methyl-3,4-dihydronaphthalen-2(1H)-one and 6-bromo-5-methyl-3,4-dihydronaphthalen-2(1H)-one (3:1, respectively) were generated. The mixture was carried through Step 5. The regioisomers and enantiomers were separated by SFC using the chiral column Phenomenex Lux 5 μ m Cellulose-3 and mobile phase 50% CO₂/IPA (2 mM NH₃-MeOH) to provide **Intermediate 11-4** as the first eluting isomer, **Intermediate 11-2** as the second eluting isomer, **Intermediate 11-5** as the third eluting isomer, and **Intermediate 11-3** as the fourth eluting isomer. Stereochemistry of the separated enantiomers were arbitrarily assigned.

Intermediate 12. tert-Butyl 4-(3-aminochroman-7-yl)piperazine-1-carboxylate



Step 1. 7-Bromo-2H-chromene-3-carbonitrile

[00189] A solution of 4-bromo-2-hydroxybenzaldehyde (10 g, 47.26 mmol) and triethylenediamine (1.12 g, 9.49 mmol) in acrylonitrile (16 mL) was stirred for 24 h at 80 °C. After cooling to room temperature, the reaction was quenched with 1N NaOH (400 mL). The resulting solution was extracted with EtOAc (300 mL x 3). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 20:1-9:1 pet. ether/EtOAc) to give 7-bromo-2H-chromene-3-carbonitrile as a yellow solid. MS: (ESI, m/z): 236, 238 [M+H]⁺.

Step 2. tert-Butyl 4-(3-cyano-2H-chromen-7-yl)piperazine-1-carboxylate

[00190] A mixture of 7-bromo-2H-chromene-3-carbonitrile (1 g, 3.81 mmol), tert-butyl piperazine-1-carboxylate (950 mg, 4.85 mmol), Pd(dppf)Cl₂ (327 mg, 0.42 mmol), XPhos (191 mg, 0.38 mmol) and Cs₂CO₃ (3.9 g, 11.37 mmol) in toluene (20 mL) was stirred for 18 h at 100 °C. After cooling to room temperature, the reaction was quenched by the addition of 30 mL of water. The resulting mixture was extracted with EtOAc (30 mL x 3). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 9:1-4:1 pet. ether/EtOAc) to give 0.7 g of tert-butyl 4-(3-cyano-2H-chromen-7-yl)piperazine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 342 [M+H]⁺.

Step 3. tert-Butyl 4-(3-carbamoyl-2H-chromen-7-yl)piperazine-1-carboxylate


[00191] A solution of tert-butyl 4-(3-cyano-2H-chromen-7-yl)piperazine-1-carboxylate (500 mg, 1.32 mmol), 30% hydrogen peroxide (0.2 mL, 2.58 mmol), and potassium carbonate (304 mg, 2.09 mmol) in a mixture of DMSO (2 mL) and ethanol (10 mL) was stirred for 3 h at room temperature. The reaction was quenched by the addition of 50 mL of water. The resulting mixture was extracted with EtOAc (50 mL x 3). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 10:1-3:1 pet. ether/EtOAc) to give tert-butyl 4-(3-carbamoyl-2H-chromen-7-yl)piperazine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 360 [M+H]⁺.

Step 4. tert-Butyl 4-(3-carbamoylchroman-7-yl)piperazine-1-carboxylate

[00192] A mixture of tert-butyl 4-(3-carbamoyl-2H-chromen-7-yl)piperazine-1-carboxylate (700 mg, 1.95 mmol) and Pd/C (100 mg, 10%) in THF (50 mL) was stirred for 18 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with EtOAc) to give tert-butyl 4-(3-carbamoylchroman-7-yl)piperazine-1-carboxylate as a light yellow solid. MS: (ESI, m/z): 362 [M+H]⁺.

Step 5. tert-Butyl 4-(3-aminochroman-7-yl)piperazine-1-carboxylate

[00193] To a stirring solution of (diacetoxyiodo)benzene (468 mg, 1.45 mmol) in a mixture of DMF (3.88 mL) and water (3.88 mL) was added tert-butyl 4-(3-carbamoylchroman-7-yl)piperazine-1-carboxylate (350 mg, 0.97 mmol). The resulting solution was stirred for 18 h at room temperature. Additional (diacetoxyiodo)benzene (936 mg, 2.90 mmol) was added in two portions over 24 h. The resulting solution was diluted with 20 mL of water and was extracted with EtOAc (50 mL). The aqueous layer was concentrated under vacuum. The residue was purified by prep-HPLC (Column: SunFire Prep C18, 19x150 mm; Mobile phase A: water (0.05% NH₄HCO₃), B: ACN; Gradient: 15% B to 70% B in 15 min) to afford tert-butyl 4-(3-aminochroman-7-yl)piperazine-1-carboxylate as a light yellow solid. MS: (ESI, m/z): 334 [M+H]⁺.

Intermediate 13. tert-Butyl 3-(3-aminochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate**Step 1. 7-Chlorochroman-3-amine**

[00194] A solution of 7-chlorochroman-3-one (700 mg, 3.83 mmol) and NH₄OAc (2.37 g, 30.75 mmol) in MeOH (35 mL) was stirred for 4 h at room temperature. To this was added NaBH₃CN

(364 mg, 5.79 mmol). The resulting mixture was stirred for 14 h at room temperature and then concentrated under vacuum. The residue was diluted with 50 mL of water. The pH value of the mixture was adjusted to 5 with 1N HCl. The resulting mixture was extracted with CH₂Cl₂ (20 mL x 2). The pH value of the aqueous layer was adjusted to 10 with 1M NaOH solution. The resulting solution was extracted with CH₂Cl₂ (30 mL x 3). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum to afford 7-chlorochroman-3-amine as a light yellow oil. MS: (ESI, m/z): 184 [M+H]⁺.

Step 2. Benzyl (7-chlorochroman-3-yl)carbamate


[00195] A mixture of 7-chlorochroman-3-amine (350 mg, 1.91 mmol), potassium carbonate (786.6 mg, 5.69 mmol), and benzyl chloroformate (390 mg, 2.29 mmol) in a mixture of EtOAc (15 mL) and water (15 mL) was stirred for 3 h at 60 °C. After cooling to room temperature, the reaction mixture was poured into 10 mL of water and was extracted with EtOAc (10 mL x 3). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 7:1 pet. ether/EtOAc) to afford benzyl (7-chlorochroman-3-yl)carbamate as a white solid. MS: (ESI, m/z): 318 [M+H]⁺.

Step 3. tert-Butyl 3-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00196] Into a 20-mL sealed tube purged and maintained with an inert atmosphere of nitrogen was placed benzyl (7-chlorochroman-3-yl)carbamate (95 mg, 0.26 mmol), tert-butyl 3,8-diazabicyclo[3.2.1]octane-8-carboxylate (70 mg, 0.33 mmol), Cs₂CO₃ (294 mg, 0.90 mmol), toluene (5 mL) and RuPhos Pd G3 (25.1 mg, 0.03 mmol). The reaction mixture was treated with microwave radiation for 5 h at 120 °C. After cooling to room temperature, the reaction mixture was then poured into 10 mL of water. The resulting mixture was extracted with CH₂Cl₂ (10 mL x 3). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 2:1 pet. ether/EtOAc) to afford tert-butyl 3-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as light yellow oil. MS: (ESI, m/z): 494 [M+H]⁺.

Step 4. tert-Butyl 3-(3-(3-aminochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00197] A mixture of tert-butyl 3-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (300 mg, 0.60 mmol) and Pd/C (60 mg, 10%) in MeOH (25 mL) was stirred for 2 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated to give tert-butyl 3-(3-aminochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow oil. MS: (ESI, m/z): 360 [M+H]⁺.

Intermediate 14-1. tert-Butyl 3-(3-(3-aminochroman-7-yl)-3,9-diazabicyclo[3.3.1]nonane-9-carboxylate**Step 1. tert-Butyl 3-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-3,9-diazabicyclo[3.3.1]nonane-9-carboxylate**

[00198] A mixture of benzyl-(7-bromochroman-3-yl)carbamate, **Intermediate 1**, (400 mg, 1.07 mmol), tert-butyl 3,9-diazabicyclo[3.3.1]nonane-9-carboxylate (300 mg, 1.31 mmol), Pd(dppf)Cl₂-CH₂Cl₂ (90 mg, 0.11 mmol), Xphos (52 mg, 0.11 mmol), and Cs₂CO₃ (722 mg, 2.22 mmol) in toluene (6 mL) was stirred at 100 °C for 14 h. After cooling to room temperature, water was added and the reaction mixture was extracted with EtOAc (70 mL x 2). The combined organic layers were dried over sodium sulfate, filtered, and concentrated. Purification by prep-TLC (eluting with 1:3 EtOAc/pet. ether) afforded tert-butyl 3-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-3,9-diazabicyclo[3.3.1]nonane-9-carboxylate as a pale yellow solid. MS: (ESI, m/z): 508 [M+H]⁺.


Step 2. tert-Butyl 3-(3-(3-aminochroman-7-yl)-3,9-diazabicyclo[3.3.1]nonane-9-carboxylate

[00199] A mixture of tert-butyl 3-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-3,9-diazabicyclo[3.3.1]nonane-9-carboxylate (120 mg, 0.22 mmol) and Pd/C (60 mg, 10%) in MeOH (5 mL) was stirred for 1 h at room temperature under an atmosphere of hydrogen. The solids were

filtered away and the filtrate was concentrated under vacuum to afford tert-butyl 3-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-3,9-diazabicyclo[3.3.1]nonane-9-carboxylate as colorless oil. MS: (ESI, m/z): 374 $[M+H]^+$.


[00200] The following intermediate in Table 5 was prepared using standard chemical manipulations and procedures similar to those used for the preparation of Intermediate 14-1.

Table 5:

Intermediate Number	Structure and Name	LRMS m/z $[M+H]^+$
14-2	 tert-butyl 4-(3-aminochroman-7-yl)-2,3-dimethylpiperazine-1-carboxylate	362

¹Notes on Procedures: In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system.

Intermediate 15-1. tert-Butyl 3-(3-amino-5-fluorochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate



Step 1. 2,4-Difluoro-6-hydroxybenzaldehyde

[00201] A solution of 3,5-difluorophenol (10 g, 73 mmol), paraformaldehyde (23 g, 728 mmol), Et₃N (21 mL), and MgCl₂ (14.6 g, 153.34 mmol) in ACN (200 mL) was stirred for 14 h at 60 °C. After cooling to room temperature, the reaction was quenched by the addition of water (100 mL). The resulting mixture was extracted with DCM (2 x 200 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:10 to 1:1 EtOAc/pet. ether) afforded 2,4-difluoro-6-hydroxybenzaldehyde as a pale yellow solid. MS: (ESI, m/z): 159 [M+H]⁺.

Step 2. tert-Butyl 3-(3-fluoro-4-formyl-5-hydroxyphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00202] A solution of 2,4-difluoro-6-hydroxybenzaldehyde (600 mg, 3.80 mmol), tert-butyl 3,8-diazabicyclo[3.2.1]octane-8-carboxylate (800 mg, 3.75 mmol), and DIEA (700 mg, 5.42 mmol) in DMSO (10 mL) was stirred for 2 h at 100 °C. After cooling to room temperature the reaction was quenched by the addition of water (10 mL). The resulting mixture was extracted with DCM (2 x 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:4 EtOAc/pet. ether) afforded tert-butyl 3-(3-fluoro-4-formyl-5-hydroxyphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 351 [M+H]⁺.

Step 3. tert-Butyl 3-(3-(allyloxy)-5-fluoro-4-formylphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00203] A solution of tert-butyl 3-(3-fluoro-4-formyl-5-hydroxyphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (1.4 g, 3.60 mmol), potassium carbonate (3 g, 21.71 mmol), and allylbromide (500 mg, 4.13 mmol) in DMF (20 mL) was stirred for 2 h at 100 °C. After cooling to room temperature the reaction was quenched by the addition of water (20 mL). The resulting mixture was extracted with DCM (2 x 50 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:5 EtOAc/pet. ether) afforded tert-butyl 3-(3-(allyloxy)-5-fluoro-4-formylphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 392 [M+H]⁺.

Step 4. tert-Butyl 3-(3-(allyloxy)-5-fluoro-4-vinylphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00204] To a solution of methyltriphenylphosphonium bromide (3.3 g, 8.86 mmol) in THF (25 mL) was added sodium hydride (178 mg, 4.45 mmol, 60% dispersion in oil) in portions at 0°C. The reaction mixture was stirred for 4 h at room temperature. This was followed by the dropwise addition of a solution of tert-Butyl 3-(3-(allyloxy)-5-fluoro-4-formylphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (1.2 g, 2.77 mmol) in THF (20 mL) and stirring continued for 3 h at 30°C. The reaction was quenched by the addition of water (60 mL). The resulting mixture was extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:6 to 1:1 EtOAc/pet. ether) afforded tert-butyl 3-(3-(allyloxy)-5-fluoro-4-vinylphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a colorless oil. MS: (ESI, m/z): 389 [M+H]⁺.

Step 5. tert-Butyl 3-(5-fluoro-2H-chromen-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00205] A solution of tert-butyl 3-(3-(allyloxy)-5-fluoro-4-vinylphenyl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (800 mg, 2.06 mmol) and Grubbs Catalyst™ 2nd Gen (48 mg, 0.05 mmol) in DCM (10 mL) was stirred for 1 h at room temperature. The mixture was concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded tert-butyl 3-(5-fluoro-2H-chromen-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a colorless oil. MS: (ESI, m/z): 361 [M+H]⁺.

Step 6. tert-Butyl 3-(5-fluoro-3-nitro-2H-chromen-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00206] A solution of KNO₂ (945 mg, 11.10 mmol) and 18-crown-6 (2.2 g, 8.32 mmol) in THF (20 mL) was stirred for 1 h at room temperature. Then I₂ (2.3 g, 9.06 mmol) was added and stirring was continued for 1 h. Finally, a solution of tert-butyl 3-(5-fluoro-2H-chromen-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (1 g, 2.77 mmol) and pyridine (110 mg, 1.39 mmol) in THF (10 mL) was added to the solution and stirring was continued for 14 h. The reaction was quenched by the addition of water (20 mL). The resulting mixture was extracted with DCM (2 x 50 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and



concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:6 EtOAc/pet. ether) afforded tert-butyl 3-(5-fluoro-3-nitro-2H-chromen-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 406 [M+H]⁺.


Step 7. tert-Butyl 3-(3-amino-5-fluorochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00207] A solution of tert-butyl 3-(5-fluoro-3-nitro-2H-chromen-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (120 mg, 0.30 mmol), BH₃-THF (1 M, 20 mL, 20.0 mmol), and NaBH₄ (116 mg, 3.07 mmol) in THF (20 mL) was stirred for 14 h at 65 °C. Then methanol (20 mL) was added and stirring was continued for 4 h at 85 °C. The mixture was concentrated under vacuum. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water with 10 mM NH₄HCO₃, B: ACN; Flow rate: 50 mL/min; Gradient: 0% to 50% B over 40 min) gave tert-butyl 3-(3-amino-5-fluorochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a colorless oil. MS: (ESI, m/z): 378 [M+H]⁺.

[00208] The following intermediates in Table 6 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Intermediate 15-1.


Table 6:

Intermediate Number	Structure and Name	LRMS m/z [M+H] ⁺
15-2 ¹	 tert-butyl 4-(3-amino-6-(trifluoromethoxy)chroman-7-yl)piperazine-1-carboxylate	418
15-3	 tert-butyl 4-(3-amino-5-fluorochroman-7-yl)piperazine-1-carboxylate	352

Intermediate Number	Structure and Name	LRMS m/z [M+H] ⁺
15-4	 tert-butyl 3-(3-amino-5-fluorochroman-7-yl)-3,6-diazabicyclo[3.1.1]heptane-6-carboxylate	364

¹ **Notes on procedures:** Step 6: nitration was conducted by sonicating a solution of tert-butyl 4-[6-(trifluoromethoxy)-2H-chromen-7-yl] piperazine-1-carboxylate (800 mg, 2.00 mmol), ACN (2.2 g, 4.00 mmol), NaNO₂ (1.4 g, 20.29 mmol) and acetic acid (1.44 g, 23.98 mmol) in chloroform (40 mL) for 6 h at 50 °C. The reaction was quenched with sat. aq. NaHCO₃ solution (40 mL) and an extractive work up was performed with EtOAc.

Intermediate 16. 3-Amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxylic acid



Step 1. 2-Chloro-5-fluoro-6-methylnicotinonitrile

[00209] A mixture of 2,6-dichloro-5-fluoropyridine-3-carbonitrile (5 g, 26.18 mmol), methylboronic acid (1.58 g, 26.36 mmol), Na₂CO₃ (8.33 g, 78.54 mmol), and Pd(dppf)Cl₂-CH₂Cl₂ (958mg, 1.31 mmol) in DMF (40 mL) and water (20 mL) was stirred for 3 h at 80 °C. After cooling to room temperature, the reaction mixture was diluted with 50 mL of water. The resulting mixture was extracted with EtOAc (3 x 100 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography

(eluting with 1:5 EtOAc/pet. ether) afforded 2-chloro-5-fluoro-6-methylnicotinonitrile as a pink solid. MS: (ESI, m/z): 171 $[M+H]^+$.


Step 2. Methyl 3-amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxylate

[00210] A solution of 2-chloro-5-fluoro-6-methylnicotinonitrile (1.40 g, 8.21 mmol), KOH (1.38 g, 24.63 mmol), and methyl 2-mercaptoacetate (1.74 g, 16.42 mmol) in DMF (20 mL) and water (20 mL) was stirred for 3 h at room temperature. The pH value of the solution was adjusted to 5 with 1N HCl solution. The solids were collected by filtration to afford methyl 3-amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxylate as a yellow solid. MS: (ESI, m/z): 241 $[M+H]^+$.

Step 3. 3-Amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxylic acid

[00211] A solution of methyl 3-amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxylate (200 mg, 0.83 mmol) and NaOH (66 mg, 1.66 mmol) in MeOH (2 mL) and water (1 mL) was stirred for 1 h at 60 °C. After cooling to room temperature, the resulting mixture was concentrated under vacuum. The residue was diluted with 20 mL of water. The pH value of the mixture was adjusted to 5 with 1N HCl solution. The solids were collected by filtration to give 3-amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxylic acid as a yellow solid. MS: (ESI, m/z): 227 $[M+H]^+$.

Intermediate 17. 3-amino-5-fluorothieno[2,3-b]pyridine-2-carboxylic acid



Step 1. Methyl 3-amino-5-fluorothieno[2,3-b]pyridine-2-carboxylate


[00212] Sodium (230 mg, 10.00 mmol) was added to MeOH (20 mL) at 0 °C and the resulting mixture was stirred for 30 min at 0 °C until the sodium was consumed. To the reaction mixture was added 2-chloro-5-fluoronicotinonitrile (900 mg, 5.75 mmol) and methyl 2-mercaptoacetate (1.8 mL, 14.98 mmol). The resulting solution was stirred for 16 h at room temperature. The reaction was quenched by the addition of 50 mL of water and was extracted with DCM (3 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and

concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:1 EtOAc/pet. ether) afforded methyl 3-amino-5-fluorothieno[2,3-b]pyridine-2-carboxylate as a yellow solid. MS: (ESI, m/z): 227 [M+H]⁺.

Step 2. 3-Amino-5-fluorothieno[2,3-b]pyridine-2-carboxylic acid

[00213] A solution of methyl 3-amino-5-fluorothieno[2,3-b]pyridine-2-carboxylate (1 g, 3.76 mmol) and LiOH (100 mg, 3.97 mmol) in water (10 mL) and dioxane (10 mL) was stirred for 1 h at room temperature. The reaction was diluted with 20 mL of water and was extracted with EtOAc (3 x 30 mL). The pH value of the aqueous layer was adjusted to 6 with 6N HCl solution. The solids were collected by filtration to afford 3-amino-5-fluorothieno[2,3-b]pyridine-2-carboxylic acid as a yellow solid. MS: (ESI, m/z): 213 [M+H]⁺.

Intermediate 18. 3-Amino-6-fluorothieno[2,3-b]pyridine-2-carboxylic acid



Step 1. 2,6-Difluoronicotinonitrile

[00214] A mixture of 2,6-dichloronicotinonitrile (6.92 g, 40.00 mmol) and KF (6.98 g, 120.14 mmol) in DMF (30 mL) was stirred overnight at 90 °C. After cooling to room temperature, the reaction was quenched by the addition of 100 mL of water. The resulting mixture was extracted with EtOAc (3 x 100 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded 2,6-difluoronicotinonitrile as a white solid. MS: (ESI, m/z): 141 [M+H]⁺.

Step 2. Benzyl 2-((3-cyano-6-fluoropyridin-2-yl)thio)acetate

[00215] To a mixture of 2,6-difluoronicotinonitrile (1 g, 6.42 mmol) and NaOAc (878 mg, 10.70 mmol) in THF (20 mL) was added benzyl 2-mercaptoacetate (1.17 g, 6.42 mmol) at -70°C. The resulting solution was warmed to room temperature slowly and then stirred for 30 min. The reaction was quenched by the addition of 20 mL of water. The resulting mixture was extracted with DCM (3 x 30 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 2:25 EtOAc/pet. ether) afforded benzyl 2-((3-cyano-6-fluoropyridin-2-yl)thio)acetate as a white solid. MS: (ESI, m/z): 303[M+H]⁺.


Step 3. Benzyl 3-amino-6-fluorothieno[2,3-b]pyridine-2-carboxylate

[00216] A solution of benzyl 2-((3-cyano-6-fluoropyridin-2-yl)thio)acetate (120 mg, 0.40 mmol) in THF (2 mL) was added dropwise at -50°C to a solution of DBU (120 mg, 0.79 mmol) in THF (3 mL). The mixture was then warmed to room temperature and stirred overnight. The resulting solution was concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:1 EtOAc/pet. ether) afforded benzyl 3-amino-6-fluorothieno[2,3-b]pyridine-2-carboxylate as an off-white solid. MS: (ESI, m/z): 303 [M+H]⁺.

Step 4. 3-Amino-6-fluorothieno[2,3-b]pyridine-2-carboxylic acid

[00217] A mixture of benzyl 3-amino-6-fluorothieno[2,3-b]pyridine-2-carboxylate (89 mg, 0.29 mmol) and Pd/C (20 mg, 10%) in EtOAc (15 mL) was stirred for 1 h at room temperature. The solids were filtered out. The filtrate was concentrated under vacuum to give 3-amino-6-fluorothieno[2,3-b]pyridine-2-carboxylic acid as a light yellow solid. MS: (ESI, m/z): 213 [M+H]⁺.

Intermediate 19. 3-Amino-6-methoxythieno[2,3-b]pyridine-2-carboxylic acid



Step 1. 2-Chloro-6-methoxynicotinonitrile

[00218] Sodium (1.5 g, 65.22 mmol) was added to MeOH (25 mL) at 0 °C and the resulting mixture was stirred for 30 min at room temperature until the sodium was consumed. To the reaction mixture was added 2,6-dichloronicotinonitrile (5 g, 28.90 mmol) over 5 min, maintaining reaction temperature below 10 °C. The resulting solution was stirred overnight at room temperature. The solids were filtered away and the filtrate was concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/hexanes) afforded 2-chloro-6-methoxynicotinonitrile as a white solid. MS: (ESI, m/z): 169 [M+H]⁺.


Step 2. Methyl 3-amino-6-methoxythieno[2,3-b]pyridine-2-carboxylate

[00219] To a solution of 2-chloro-6-methoxynicotinonitrile (3.9 g, 23.13 mmol) in DMF (10 mL) was added KOH (5.2 g) at 0 °C over 5 min, followed by the addition of methyl 2-mercaptoacetate (2.46 g, 23.18 mmol). The resulting solution was stirred for 1 h at 0 °C. The reaction was quenched by the addition of 20 mL of water. The resulting mixture was extracted with EtOAc (3x30 mL). The combined organic layers were washed with water (50 mL) and brine (50 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:10 to 1:1 EtOAc/hexanes) afforded methyl 3-amino-6-methoxythieno[2,3-b]pyridine-2-carboxylate as a light yellow solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ(ppm): 8.40 (d, *J* = 8.7 Hz, 1H), 7.24 (br, 2H), 6.89 (d, *J* = 9.0 Hz, 1H), 3.93 (s, 3H), 3.77 (s, 3H). MS: (ESI, m/z): 239 [M+H]⁺.

Step 3. 3-Amino-6-methoxythieno[2,3-b]pyridine-2-carboxylic acid

[00220] A solution of methyl 3-amino-6-methoxythieno[2,3-b]pyridine-2-carboxylate (110 mg, 0.46 mmol) and LiOH (100 mg, 4.18 mmol) in THF (4 mL) and water (1.5 mL) was stirred for 2 h at 60 °C. After cooling to room temperature, the resulting mixture was concentrated under vacuum. The residue was diluted with 2 mL of water. The pH value of the solution was adjusted to 7 with 1N HCl. The solids were collected by filtration to afford 3-amino-6-methoxythieno[2,3-b]pyridine-2-carboxylic acid as a yellow solid. MS: (ESI, m/z): 225 [M+H]⁺.

Intermediate 20. 3-Amino-5-fluoro-6-methoxythieno[2,3-b]pyridine-2-carboxylic acid



Step 1. 2-Chloro-5-fluoro-6-methoxynicotinonitrile

[00221] A mixture of 2,6-dichloro-5-fluoropyridine-3-carbonitrile (3.0 g, 15.71 mmol) and MeONa (1.28 g, 23.70 mmol) in MeOH (30 mL) was stirred for 5 h at room temperature. The resulting mixture was concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded 2-chloro-5-fluoro-6-methoxynicotinonitrile as a yellow solid. MS: (ESI, m/z): 187, 189 $[M+H]^+$.


Step 2. Methyl 3-amino-5-fluoro-6-methoxythieno[2,3-b]pyridine-2-carboxylate

[00222] A solution of 2-chloro-5-fluoro-6-methoxynicotinonitrile (1.90 g, 10.22 mmol) methyl 2-mercaptoacetate (1.3 g, 12.26 mmol), and DBU (7.2 g, 47.29 mmol) in THF (30 mL) was stirred overnight at room temperature. The reaction was quenched by the addition of 50 mL of water and was extracted with EtOAc (3 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded methyl 3-amino-5-fluoro-6-methoxythieno[2,3-b]pyridine-2-carboxylate as a yellow solid. MS: (ESI, m/z): 257 $[M+H]^+$.

Step 3. 3-Amino-5-fluoro-6-methoxythieno[2,3-b]pyridine-2-carboxylic acid

[00223] A mixture of methyl 3-amino-5-fluoro-6-methoxythieno[2,3-b]pyridine-2-carboxylate (500 mg, 1.95 mmol) and LiOH (236 mg, 9.85 mmol) in THF (8 mL) and water (8 mL) was stirred overnight at 60 °C. After cooling to room temperature, the solvent was removed under vacuum. The pH value of the residue was adjusted to 7 with 3N HCl. The solids were collected by filtration to afford 3-amino-5-fluoro-6-methoxythieno[2,3-b]pyridine-2-carboxylic acid as a yellow solid. MS: (ESI, m/z): 243 $[M+H]^+$.

Intermediate 21. 3-Amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylic




Step 1. Methyl 3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylate

[00224] To a solution of 2-chloro-4,6-dimethylnicotinonitrile (2.000 g, 12.00 mmol) and DBU (5.00 g, 32.84 mmol) in DMF (20 mL) was added methyl 2-sulfanylacetate (1.019 g, 9.60 mmol) dropwise with stirring at -50°C. The resulting solution was stirred overnight at room temperature. The mixture was poured into water (50 mL) and the solids were collected by filtration to give methyl 3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylate as a light yellow solid. MS: (ESI, m/z): 237 $[M+H]^+$.

Step 2. 3-Amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylic acid

[00225] To a solution of methyl 3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylate (300 mg, 1.27 mmol) in MeOH (5 mL) was added a solution of NaOH (254 mg, 6.35 mmol) in water (5 mL). The resulting solution was stirred for 3 h at 70°C. After cooling to room temperature, the solvent was removed under vacuum. The pH value of the residue was adjusted to 6 with 3N HCl. The solids were collected by filtration to afford 3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylic acid as a yellow solid. MS: (ESI, m/z): 223 $[M+H]^+$.

Intermediate 22. 3-Amino-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxylic acid



Step 1. 6-Methyl-2-thioxo-4-(trifluoromethyl)-1,2-dihydropyridine-3-carbonitrile

[00226] A solution of 2-cyanoethanethioamide (2 g, 19.97 mmol), 1,1,1-trifluoropentane-2,4-dione (3 g, 19.47 mmol), and triethylamine (0.1 mL) in ethanol (20 mL) was stirred for 1 h at 90 °C. After cooling to room temperature, the solids were collected by filtration and dried in an oven under reduced pressure to afford 6-methyl-2-thioxo-4-(trifluoromethyl)-1,2-dihydropyridine-3-carbonitrile as a yellow solid. MS: (ESI, m/z): 219 $[M+H]^+$.

Step 2. Ethyl 3-amino-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxylate


[00227] A solution of 6-methyl-2-thioxo-4-(trifluoromethyl)-1,2-dihydropyridine-3-carbonitrile (800 mg, 3.67 mmol), ethyl 2-bromoacetate (609 mg, 3.65 mmol), and NaOEt (297 mg, 4.37 mmol) in ethanol (20 mL) was stirred overnight at 80 °C. The solvent was removed under vacuum. The residue was diluted with 30 mL of water and was extracted with ethyl acetate (3 x 30 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:10 to 1:1 EtOAc/pet. ether) afforded ethyl 3-amino-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxylate as a yellow solid. MS: (ESI, m/z): 305 $[M+H]^+$.

Step 3. 3-Amino-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxylic acid

[00228] A solution of ethyl 3-amino-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxylate (1.0 g, 3.29 mmol) and sodium hydroxide (470 mg, 11.75 mmol) in water (2 mL) and methanol (10 mL) was stirred for 2 h at 60 °C. The resulting mixture was concentrated under vacuum. The residue was diluted with 10 mL of water. The pH value of the solution was adjusted to 3 with 3N HCl. The solids were collected by filtration to give 3-amino-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxylic acid as a yellow solid. MS: (ESI, m/z): 277 $[M+H]^+$.

Intermediate 23. Methyl 6-amino-2-methylthieno[2,3-d]thiazole-5-carboxylate and

Intermediate 24. 6-Amino-2-methylthieno[2,3-d]thiazole-5-carboxylic acid



Step 1. 4-Chloro-2-methylthiazole-5-carbonitrile

[00229] A mixture of 2,4-dichlorothiazole-5-carbonitrile (1.00 g, 5.59 mmol), dimethylzinc (1M in Et₂O) (8.8 mL, 8.80 mmol), and Pd(dppf)Cl₂·CH₂Cl₂ (911 mg, 1.12 mmol) in toluene (30 mL) was stirred for 4 h at 40 °C. The reaction was quenched by the addition of 20 mL of water. The resulting mixture was extracted with EtOAc (3 x 20 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:5 EtOAc/pet. ether) afforded 4-chloro-2-methylthiazole-5-carbonitrile as a light yellow solid. MS: (ESI, m/z): 159 [M+H]⁺.

Step 2. Methyl 6-amino-2-methylthieno[2,3-d]thiazole-5-carboxylate


[00230] To a solution of 4-chloro-2-methylthiazole-5-carbonitrile (550 mg, 3.47 mmol) and DBU (1.06 g, 6.94 mmol) in THF (20 mL) was added a solution of methyl 2-mercaptoacetate (443 mg, 4.17 mmol) in THF (2 mL) dropwise with stirring at -40 °C. The resulting solution allowed to warm to room temperature while stirring overnight. The reaction was quenched with 20 mL of water. The resulting mixture was extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:7 EtOAc/pet. ether) afforded methyl 6-amino-2-methylthieno[2,3-d]thiazole-5-carboxylate as a light yellow solid. MS: (ESI, m/z): 229 [M+H]⁺.

Step 3. 6-Amino-2-methylthieno[2,3-d]thiazole-5-carboxylic acid

[00231] To a solution of methyl 6-amino-2-methylthieno[2,3-d]thiazole-5-carboxylate (174 mg, 0.76 mmol) in ACN (11 mL) was added a solution of LiOH (100 mg, 4.18 mmol) in water (5 mL). The resulting solution was stirred overnight at 30 °C, then was concentrated under vacuum. The residue was diluted with 1 mL of water. The pH value of the residue was adjusted to 7 with


1N HCl solution. The solids were collected by filtration to give 6-amino-2-methylthieno[2,3-d]thiazole-5-carboxylic acid as an off-white solid. MS: (ESI, m/z): 214 $[M+H]^+$.

Intermediate 25. 1-Ethyl-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid



[00232] To a solution of methyl 1H-pyrrolo[2,3-b]pyridine-5-carboxylate (3 g, 17.03 mmol) in DMF (80 mL) was added sodium hydride (2.04 g, 51.09 mmol, 60% dispersion in oil) in portions at 0 °C. The reaction mixture was stirred for 1 h at 0 °C. Then iodoethane (5.32 g, 34.06 mmol) was added at 0 °C. The resulting solution was stirred for 10 h at room temperature. The reaction was quenched with 10 mL of water. After stirred for 30 min, the pH value of the solution was adjusted to 7~8 with 3N HCl. The resulting mixture was extracted with EtOAc (6 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum to give of 1-ethyl-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid as a yellow solid (crude, 90% purity). MS: (ESI, m/z): 191 $[M+H]^+$.

Intermediate 26. 7-Ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxylic acid



Step 1. 3-Chloro-7-ethyl-7H-pyrrolo[2,3-c]pyridazine

[00233] To a solution of 3-chloro-7H-pyrrolo[2,3-c]pyridazine (700 mg, 4.56 mmol) in DMF (17 mL) was added NaH (365 mg, 9.12 mmol, 60% dispersion in oil) in portions at 0 °C. The reaction mixture was stirred for 30 min at 0 °C. Then iodoethane (856 mg, 5.49 mmol) was added at 0 °C and the reaction mixture was stirred for 2 h at room temperature. The reaction was quenched by the addition of 40 mL of water. The resulting mixture was extracted with EtOAc (3 x 20 mL).

The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by prep-TLC (eluting with 1:1 EtOAc/pet. ether) afforded 3-chloro-7-ethyl-7H-pyrrolo[2,3-c]pyridazine as yellow oil. MS: (ESI, m/z): 182, 183 [M+H]⁺.


Step 2. Methyl 7-ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxylate

[00234] In a 30-mL pressure tank reactor, a solution of 3-chloro-7-ethyl-7H-pyrrolo[2,3-c]pyridazine (300 mg, 1.65 mmol), Pd(dppf)Cl₂.CH₂Cl₂ (121 mg, 0.15 mmol), and Et₃N (0.69 mL, 4.96 mmol) in MeOH (15 mL) was stirred for 48 h at 120 °C under 50 atm of CO (g). After cooling to room temperature, the resulting mixture was concentrated under vacuum. The residue was diluted with 20 mL of water. The resulting mixture was extracted with EtOAc (3 x 20 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by prep-TLC (eluting with 1:1 EtOAc/pet. ether) afforded methyl 7-ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxylate as a yellow solid. MS: (ESI, m/z): 206 [M+H]⁺.

Step 3. 7-Ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxylic acid

[00235] A mixture of methyl 7-ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxylate (238 mg, 1.04 mmol), and sodium hydroxide (206 mg, 5.20 mmol) in THF (10 mL) and water (10 mL) was stirred for 18 h at room temperature. The resulting mixture was concentrated under vacuum. The residue was diluted with 10 mL of water. The pH value of the mixture was adjusted to 5 with 2N HCl. The resulting mixture was extracted with EtOAc (3 x 30 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give 7-ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxylic acid as a light yellow solid (crude). MS: (ESI, m/z): 192 [M+H]⁺.

Intermediate 27. 3-Amino-6-methylfuro[2,3-b]pyridine-2-carboxylic acid




Step 1. Ethyl 3-amino-6-methylfuro[2,3-b]pyridine-2-carboxylate

[00236] A solution of 2-chloro-6-methylnicotinonitrile (5 g, 32.77 mmol), ethyl 2-hydroxyacetate (3.36 g, 32.28 mmol), and Cs₂CO₃ (32.2 g, 98.83 mmol) in NMP (80 mL) was stirred overnight at 75 °C. After cooling to room temperature, the mixture was poured into 100 mL of water. The resulting mixture was extracted with ethyl acetate (3 x 100 mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) afforded ethyl 3-amino-6-methylfuro[2,3-b]pyridine-2-carboxylate as a pink solid. MS: (ESI, m/z): 221 [M+H]⁺.

Step 2. 3-Amino-6-methylfuro[2,3-b]pyridine-2-carboxylic acid

[00237] To a solution of ethyl 3-amino-6-methylfuro[2,3-b]pyridine-2-carboxylate (110 mg, 0.53 mmol) in methanol (1 mL) and THF (1 mL) was added a solution of LiOH (24 mg, 1.00 mmol) in water (0.5 mL) dropwise with stirring. The resulting solution was stirred overnight at room temperature. The pH value of the solution was adjusted to 8 with 1N HCl. The resulting mixture was extracted with ethyl acetate (3 x 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.1% formic acid), B: ACN; Flow rate: 50 mL/min; Gradient: 0% B to 100% B in 30 min) afforded 3-amino-6-methylfuro[2,3-b]pyridine-2-carboxylic acid as a light yellow solid. MS: (ESI, m/z): 193 [M+H]⁺.

Intermediate 28. 8-(tert-Butoxycarbonyl)-5,6,7,8-tetrahydro-1,8-naphthyridine-3-carboxylic acid



Step 1. Methyl 5,6,7,8-tetrahydro-1,8-naphthyridine-3-carboxylate

[00238] In a 30-mL pressure tank reactor, a mixture of tert-butyl 6-bromo-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate (300 mg, 0.86 mmol), Pd(dppf)Cl₂·CH₂Cl₂ (90 mg, 0.11 mmol), and Et₃N (1 mL) in MeOH (5 mL) was stirred for 48 h at 120°C under 5 atm of CO (g). After cooling to room temperature, the solvent was removed under vacuum. The residue was diluted with water (10 mL) and was extracted with EtOAc (3 x 10 mL). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:5 EtOAc/pet. ether) to afford 0.15 g of methyl 5,6,7,8-tetrahydro-1,8-naphthyridine-3-carboxylate as a white solid. MS: (ESI, m/z): 193 [M+H]⁺.


Step 2. 1-(tert-Butyl) 6-methyl 3,4-dihydro-1,8-naphthyridine-1,6(2H)-dicarboxylate

[00239] A solution of methyl 5,6,7,8-tetrahydro-1,8-naphthyridine-3-carboxylate (110 mg, 0.52 mmol), DMAP (126 mg, 1.03 mmol), and (Boc)₂O (227 mg, 1.04 mmol) in DMF (5 mL) was stirred for 1 h at room temperature. The reaction was quenched by the addition of 5 mL of water. The resulting mixture was extracted with ethyl acetate (3 x 10 mL). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:8 EtOAc/pet. ether) to afford 1-(tert-butyl) 6-methyl 3,4-dihydro-1,8-naphthyridine-1,6(2H)-dicarboxylate as a white solid. MS: (ESI, m/z): 293 [M+H]⁺.

Step 3. 8-(tert-Butoxycarbonyl)-5,6,7,8-tetrahydro-1,8-naphthyridine-3-carboxylic acid

[00240] A solution of 1-(tert-butyl) 6-methyl 3,4-dihydro-1,8-naphthyridine-1,6(2H)-dicarboxylate (280 mg, 0.86 mmol) and LiOH (81 mg, 3.38 mmol) in THF (5 mL) and water (5 mL) was stirred for 3 h at room temperature. The resulting mixture was concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.05% formic acid), B: ACN; Gradient: 0% B to 60% B in 40 min) to afford 8-(tert-butoxycarbonyl)-5,6,7,8-tetrahydro-1,8-naphthyridine-3-carboxylic acid as a white solid. MS: (ESI, m/z): 279 [M+H]⁺.

Intermediate 29. 2',3'-Dihydro-1'H-spiro[cyclopropane-1,4'-[1,8]naphthyridine]-6'-carboxylic acid



Step 1. 6-Bromo-1,2,3,4-tetrahydro-1,8-naphthyridine

[00241] To a solution of 6-bromo-3,4-dihydro-1,8-naphthyridin-2(1H)-one (5.0 g, 21.80 mmol) and NaBH₄ (4.18 g, 110.49 mmol) in THF (140 mL) was added BF₃-Et₂O (20 mL, 157.83 mmol) dropwise at 0 °C. The reaction mixture was stirred for 16 h at room temperature. 1N HCl solution (100 mL) was added and the reaction mixture was stirred for an additional 16 h at room temperature. The pH value of the mixture was then adjusted to 8 with aq. sat. NaHCO₃ solution. The resulting mixture was extracted with ethyl acetate (3 x 150 mL) and the combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give 6-bromo-1,2,3,4-tetrahydro-1,8-naphthyridine as a white solid. MS: (ESI, m/z): 213, 215 [M+H]⁺.

Step 2. tert-Butyl 6-bromo-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate

[00242] To a mixture of sodium hydride (1.41 g, 58.76 mmol, 60% dispersion in oil) in THF (100 mL) was added a solution of 6-bromo-1,2,3,4-tetrahydro-1,8-naphthyridine (5.0 g, 22.53 mmol) in THF (100 mL) at 0 °C. The reaction mixture was stirred for 30 min at 0 °C. To the mixture was then added a solution of (Boc)₂O (10.15 g, 46.51 mmol) in THF (50 mL). The resulting mixture was heated at reflux for 16 h. After cooling to room temperature, the reaction was quenched by the addition of water (150 mL). The resulting mixture was extracted with ethyl acetate (3 x 150 mL) and the combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with

1:3 EtOAc/pet. ether) afforded tert-butyl 6-bromo-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate as a light yellow solid. MS: (ESI, m/z): 313, 315 [M+H]⁺.

Step 3. tert-Butyl 6-bromo-4-oxo-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate

[00243] To a solution of tert-butyl 6-bromo-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate (2.0 g, 6.13 mmol) and NaH₂PO₄ (1.92 g, 16.00 mmol) in tert-butanol (20 mL) and water (15 mL) was added a solution of NaMnO₄·H₂O (6.13 g, 38.31 mmol) in water (5 mL) dropwise at 50 °C. The reaction mixture was stirred for 3 h at 50 °C. After cooling to room temperature, Na₂SO₃ was added and the mixture was stirred for 30 min at room temperature. The solids were filtered away and the filtrate was diluted with 50 mL of water. The resulting mixture was extracted with ethyl acetate (3 x 50 mL) and the combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded tert-butyl 6-bromo-4-oxo-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate as a white solid. MS: (ESI, m/z): 327, 329 [M+H]⁺.

Step 4. tert-Butyl 6-bromo-4-methylene-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate

[00244] A solution of methyltriphenylphosphonium bromide (3.29 g, 9.21 mmol) and t-BuOK (1M in THF) (9.2 mL, 9.02 mmol) in toluene (30 mL) was stirred for 1 h at 100 °C. To the reaction mixture was added a solution of tert-butyl 6-bromo-4-oxo-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate (1.5 g, 4.40 mmol) in toluene (5 mL). The resulting solution was stirred for 1 h at 100 °C. After cooling to room temperature, the reaction was quenched by the addition of water (50 mL). The resulting mixture was extracted with ethyl acetate (3x50 mL) and the combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:100 to 1:3 EtOAc/pet. ether) afforded tert-butyl 6-bromo-4-methylene-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate as a white solid. MS: (ESI, m/z): 325, 327 [M+H]⁺.

Step 5. tert-Butyl 6'-bromo-2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-[1,8]naphthyridine]-1'-carboxylate


[00245] To a solution of potassium hydroxide (4.8 g, 85.55 mmol) in water (7.2 mL) was added a solution of 2-nitrosopropanamide (3.76 g, 36.83 mmol) in Et₂O (30 mL) was added. The resulting solution was stirred for 1 h at 0 °C. The organic phase was separated to obtain the solution of

diazomethane in Et₂O. To a solution of tert-butyl 6-bromo-4-methylene-3,4-dihydro-1,8-naphthyridine-1(2H)-carboxylate (400 mg, 1.18 mmol) in THF (30 mL) was added the solution of diazomethane at 0 °C, followed by the addition of a mixture of Pd(OAc)₂ (28 mg, 0.12 mmol) in THF (3 mL). The reaction mixture was stirred for an additional 3 h at room temperature. The solids were filtered away and the filtrate was concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded tert-butyl 6'-bromo-2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-[1,8]naphthyridine]-1'-carboxylate as a light yellow solid. MS: (ESI, m/z): 339, 341 [M+H]⁺.

Step 6. 2',3'-Dihydro-1'H-spiro[cyclopropane-1,4'-[1,8]naphthyridine]-6'-carboxylic acid

[00246] Into a 30-mL pressure tank reactor fitted with a magnetic stir bar, was placed a mixture of tert-butyl 6'-bromo-2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-[1,8]naphthyridine]-1'-carboxylate (140 mg, 0.40 mmol), sodium acetate trihydrate (167 mg, 1.23 mmol), and Pd(dppf)Cl₂-CH₂Cl₂ (65 mg, 0.08 mmol) in DMF (6 mL) and water (2 mL). The reaction mixture was stirred for 16 h at 120 °C under an atmosphere of carbon monoxide at 50 atm. After cooling to room temperature, the reaction was quenched by the addition of 20 mL of water. The resulting solution was extracted with ethyl acetate (3 x 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water, B: ACN; Gradient: 0% B to 10% B in 10 min) afforded 2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-[1,8]naphthyridine]-6'-carboxylic acid as an off-white solid. MS: (ESI, m/z): 205 [M+H]⁺.

Intermediate 30. 3-(tert-Butoxycarbonyl)-1a,2,3,7b-tetrahydro-1H-cyclopropa[c][1,8]naphthyridine-6-carboxylic acid



Step 1. 2-(Allylamino)-5-bromonicotinaldehyde

[00247] Into two parallel 30 ml sealed tubes, each was placed 5-bromo-2-fluoronicotinaldehyde (1.83 g, 9.0 mmol), allylamine (1.03 g, 18.0 mmol) and ethanol (15 mL). The resulting solution was stirred for 3 h at 80 °C. After cooling room temperature, the resulting solution was poured into 30 mL of hydrochloric acid (1N) and the resulting mixture was stirred for 10 min and then extracted with ethyl acetate (3 x 30 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:10 EtOAc/pet. ether) afforded 2-(allylamino)-5-bromonicotinaldehyde as a light yellow solid. MS: (ESI, m/z): 241, 243 [M+H]⁺.

Step 2. N-Allyl-5-bromo-3-vinylpyridin-2-amine

[00248] A solution of methyltriphenylphosphonium bromide (6.22 g, 17.42 mmol) and potassium tert butoxide (1.96 g, 17.42 mmol) in THF (30 mL) was stirred for 1 h at room temperature. Then a solution of 2-(allylamino)-5-bromonicotinaldehyde (2.10 g, 8.71 mmol) in THF (5 mL) was added dropwise at room temperature and the reaction mixture was stirred overnight at room temperature. The reaction was quenched by the addition of 20 mL of water and was extracted with DCM (3 x 40). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting

with 1:20 EtOAc/pet. ether) afforded N-allyl-5-bromo-3-vinylpyridin-2-amine as a light yellow liquid. MS: (ESI, m/z): 239, 241 [M+H]⁺.

Step 3. tert-Butyl allyl(5-bromo-3-vinylpyridin-2-yl)carbamate

[00249] A solution of N-allyl-5-bromo-3-vinylpyridin-2-amine (590 mg, 2.47 mmol), di-tert-butyl dicarbonate (1.62 g, 7.40 mmol), triethylamine (749 mg, 7.40 mmol) and 4-dimethylaminopyridine (90 mg, 0.74 mmol) in THF (15 mL) was stirred overnight at 70 °C. After cooling to room temperature, the reaction was quenched by the addition of 50 mL of water. The resulting mixture was extracted with ethyl acetate (3 x 40 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:30 EtOAc/pet. ether) afforded tert-butyl allyl(5-bromo-3-vinylpyridin-2-yl)carbamate as an off-white solid. MS: (ESI, m/z): 339, 341 [M+H]⁺.

Step 4. tert-Butyl 6-bromo-1,8-naphthyridine-1(2H)-carboxylate

[00250] A solution of tert-butyl allyl(5-bromo-3-vinylpyridin-2-yl)carbamate (600 mg, 1.77 mmol) and dichloro[1,3-bis(2,4,6-trimethylphenyl)-2-imidazolidinylidene](benzylidene)(tricyclohexylphosphine)ruthenium(II) (Grubbs Catalyst™ 2nd Gen) (75 mg, 0.09 mmol) in DCM (10 mL) was stirred overnight at 50 °C. After cooling to room temperature, the resulting mixture was concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:20 EtOAc/pet. ether) afforded tert-butyl 6-bromo-1,8-naphthyridine-1(2H)-carboxylate as an off-white solid. MS: (ESI, m/z): 311, 313 [M+H]⁺.

Step 5. tert-Butyl 6-bromo-1,1a,2,7b-tetrahydro-3H-cyclopropa[c][1,8]naphthyridine-3-carboxylate


[00251] To a solution of potassium hydroxide (5.68 g, 101.24 mmol) in water (8 mL) was added a solution of 1-methyl-1-nitrosourea (2.98 g, 28.92 mmol) in Et₂O (40 mL) dropwise at 0 °C. The reaction mixture was stirred for 1 h at 0 °C. Then the organic phase was separated to obtain the solution of diazomethane in Et₂O. To a solution of tert-butyl 6-bromo-1,8-naphthyridine-1(2H)-carboxylate (450 mg, 1.45 mmol) in THF (15 mL) was added the solution of diazomethane in Et₂O (40 mL), followed by the addition of a solution of Pd(OAc)₂ (32 mg, 0.14 mmol) in THF (7 mL). The resulting solution was stirred overnight at room temperature. The solids were filtered away and the filtrate was concentrated under vacuum. Purification by silica gel chromatography (eluting

with gradient 1:50 to 1:20 EtOAc/pet. ether) afforded tert-butyl 6-bromo-1,1a,2,7b-tetrahydro-3H-cyclopropa[c][1,8]naphthyridine-3-carboxylate as an off-white solid. MS: (ESI, m/z): 325, 327 [M+H]⁺.

Step 6. 3-(tert-Butoxycarbonyl)-1a,2,3,7b-tetrahydro-1H-cyclopropa[c][1,8]naphthyridine-6-carboxylic acid

[00252] Into a 30-mL pressure tank reactor, was placed tert-butyl 6-bromo-1,1a,2,7b-tetrahydro-3H-cyclopropa[c][1,8]naphthyridine-3-carboxylate (200 mg, 0.62 mmol), Pd(dppf)Cl₂ (90 mg, 0.12 mmol), NaOAc (151 mg, 1.85 mmol), DMF (4.5 mL) and water (1.5 mL). The reaction mixture was stirred for 18 h at 120 °C under 50 atm of CO (g). After cooling to room temperature, the reaction mixture was diluted with 10 mL of water. The resulting mixture was extracted with ethyl acetate (3 x 10 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.1% formic acid), B: ACN; Flow rate: 50 mL/min; Gradient: 0% B to 100% B in 30 min) afforded 3-(tert-butoxycarbonyl)-1a,2,3,7b-tetrahydro-1H-cyclopropa[c][1,8]naphthyridine-6-carboxylic acid as a light yellow solid. MS: (ESI, m/z): 291 [M+H]⁺.

Intermediate 31. 6-(Benzylamino)nicotinic acid



Step 1. Methyl 6-(benzylamino)nicotinate


[00253] A mixture of methyl 6-fluoropyridine-3-carboxylate (1 g, 6.12 mmol), phenylmethanamine (1.38 g, 12.88 mmol), and potassium carbonate (2.67 g, 19.32 mmol) in DMF (10 mL) was stirred for 2 h at 80 °C. After cooling to room temperature, the reaction was quenched by the addition of 20 mL of water. The resulting mixture was extracted with ethyl acetate (3 x 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 3:10

EtOAc:pet. ether) afforded methyl 6-(benzylamino)nicotinate as a white solid. MS: (ESI, m/z): 243 [M+H]⁺.

Step 2. 6-(Benzylamino)nicotinic acid

[00254] A mixture of methyl 6-(benzylamino)nicotinate (500 mg, 1.86 mmol), methanol (20 mL), water (2 mL), and NaOH (165 mg, 4.13 mmol) was stirred for 3 h at 80 °C. After cooling to room temperature, the resulting mixture was concentrated under vacuum. The residue was dissolved in 10 mL of water. The pH value of the solution was adjusted to 6 with hydrochloric acid (6 N). The solids were collected by filtration to afford 6-(benzylamino)nicotinic acid as a white solid. MS: (ESI, m/z): 229 [M+H]⁺.

Intermediate 32. tert-Butyl 9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate



Step1: tert-Butyl 7-benzyl-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate


[00255] A solution of tert-butyl 7-benzyl-9-oxo-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate (5 g, 14.94 mmol) and DAST (12.2 g, 75.69 mmol) in CH₂Cl₂ (80 mL) was stirred for 16 h at room temperature. The reaction was then quenched by the addition of 50 mL of saturated aqueous NaHCO₃ solution. The resulting mixture was extracted with CH₂Cl₂ (100 mL x 3). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 5:1 pet. ether/EtOAc) afforded 700 mg of tert-butyl 7-benzyl-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate as a yellow solid. MS: (ESI, m/z): 353 [M+H]⁺.

Step2: tert-Butyl 9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate

[00256] A mixture of tert-butyl 7-benzyl-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate (460 mg, 1.24 mmol) and Pd/C (50 mg, 10%) in EtOAc (20 mL) was stirred at 50 °C for 2 h under an atmosphere of hydrogen. After cooling to room temperature, the solids were filtered away and the filtrate was concentrated under vacuum. The residue was purified by pre-HPLC (Column: XBridge Prep C18 OBD, 19x250 mm; Mobile phase A: water (10 mM

NH₄HCO₃), B: ACN; Gradient: 20% B to 45% B in 7 min) to afford tert-butyl 9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate as a light yellow solid. MS: (ESI, m/z): 263 [M+H]⁺.

Intermediate 33. 5-Benzyl-2-oxa-5,8-diazaspiro[3.5]nonane



Step 1. Methyl 2-(3-(nitromethyl)oxetan-3-ylamino)acetate

[00257] To a solution of oxetan-3-one (5 g, 69.38 mmol), nitromethane (5.93 g, 97.15 mmol), and Et₃N (2.1 g, 13.28 mmol) in DCM (70 mL) was added a solution of MsCl (10 g, 87.30 mmol) in DCM (70 mL) at -80 °C. Stirring continued at -80 °C for an additional 90 min. Separately, a solution of glycine ethyl ester hydrochloride (19.4 g, 139 mmol) and Et₃N (21 g, 139 mmol) in DCM (300 mL) was allowed to react, with stirring, for 10 min at room temperature. The resulting solution was added into the first reaction mixture at -80 °C in portions. After addition, the reaction mixture was stirred for 16 h at room temperature. The reaction was quenched by the addition of 100 mL of water and was extracted with 2 x 300 mL of DCM. The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by prep-HPLC (Column: XBridge Prep C18 OBD, 19x150 mm, 5 μm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 10% B to 50% B in 30 min) afforded methyl 2-(3-(nitromethyl)oxetan-3-ylamino)acetate as a yellow oil. MS: (ESI, m/z): 205 [M+H]⁺.

Step 2. 2-Oxa-5,8-diazaspiro[3.5]nonan-7-one

[00258] A suspension of methyl 2-(3-(nitromethyl)oxetan-3-ylamino)acetate (8.5 g, 41.63 mmol) and Raney Ni (2 g) in MeOH (50 mL) was stirred for 16 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum. Purification by prep-HPLC (Column: XBridge Prep C18 OBD, 19x150 mm, 5 μm;

Mobile phase A: water (10 mM NH_4HCO_3), B: ACN; Gradient: 10% B to 80% B in 30 min) afforded 2-oxa-5,8-diazaspiro[3.5]nonan-7-one as a red solid. MS: (ESI, m/z): 143 $[\text{M}+\text{H}]^+$.

Step 3. 5-Benzyl-2-oxa-5,8-diazaspiro[3.5]nonan-7-one

[00259] A mixture of 2-oxa-5,8-diazaspiro[3.5]nonan-7-one (4 g, 28.14 mmol), (bromomethyl)benzene (12 g, 70.16 mmol), Na_2CO_3 (20.9 g, 197.18 mmol), and NaI (10.5 g, 70.05 mmol) in acetonitrile (200 mL) was stirred for 3 h at 80 °C. After cooling to room temperature, the solvent was removed under vacuum. The residue was diluted with water (200 mL). The resulting mixture was extracted with DCM (2 x 300 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:10 MeOH/ CH_2Cl_2) afforded 5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonan-7-one as a yellow solid. MS: (ESI, m/z): 233 $[\text{M}+\text{H}]^+$.

Step 4. 5-Benzyl-2-oxa-5,8-diazaspiro[3.5]nonane-7-thione


[00260] A solution of 5-benzyl-2-oxa-5,8-diazaspiro [3.5]nonan-7-one (2.9 g, 11.86 mmol) and 2,4-bis(4-methoxyphenyl)-2,4-dithioxo-1,3,2,4-dithiadiphosphetane (Lawesson reagent) (2.44 g, 6.03 mmol) in THF (150 mL) was stirred for 16 h at room temperature and then stirred for an additional 3 h at 65 °C. The reaction mixture was then poured into water (100 mL) and was extracted with DCM (2 x 150 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded 5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonane-7-thione as a white solid. MS: (ESI, m/z): 249 $[\text{M}+\text{H}]^+$.

Step 5. 5-Benzyl-2-oxa-5,8-diazaspiro[3.5]nonane

[00261] A solution of 5-benzyl-2-oxa-5,8-diazaspiro [3.5]nonane-7-thione (300 mg, 1.15 mmol) and sodium borohydride (412 mg, 11.19 mmol) in THF (15 mL) and MeOH (30 mL) was stirred for 2 h at room temperature. The reaction was quenched with water (15 mL) and was stirred for an additional 14 h at room temperature. The resulting mixture was extracted with DCM (2 x 50 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by prep-HPLC (Column: XBridge Prep C18 OBD, 19x150 mm, 5 μm ; Mobile phase A: water (10 mM NH_4HCO_3), B: ACN; Gradient: 10% B to 75%

B in 30 min) afforded 5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonane as a white solid. MS: (ESI, m/z): 219 $[M+H]^+$.

Intermediate 34. tert-Butyl 2-oxa-5,8-diazaspiro[3.5]nonane-8-carboxylate




Step 1. tert-Butyl 5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonane-8-carboxylate

[00262] A solution of 5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonane, **Intermediate 31**, (200 mg, 0.82 mmol), $(Boc)_2O$ (200 mg, 0.92 mmol), Et_3N (185 mg, 1.83 mmol), and 4-dimethylaminopyridine (11 mg, 0.09 mmol) in DCM (4 mL) was stirred for 2.5 h at room temperature. The reaction was quenched by the addition of water (20 mL) and was extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:4 EtOAc/pet. ether) afforded tert-butyl 5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonane-8-carboxylate as a yellow oil. MS: (ESI, m/z): 319 $[M+H]^+$.

Step 2. tert-Butyl 2-oxa-5,8-diazaspiro[3.5]nonane-8-carboxylate

[00263] A mixture of tert-butyl 5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonane-8-carboxylate (190 mg, 0.60 mmol) and Pd/C (20 mg, 10%) in EtOAc (10 mL) was stirred for 1 h at room temperature. The solids were filtered away and the filtrate was concentrated under vacuum to afford tert-butyl 2-oxa-5,8-diazaspiro[3.5]nonane-8-carboxylate as a yellow oil. MS: (ESI, m/z): 229 $[M+H]^+$.

Intermediate 35. tert-Butyl N-(4-methoxypyrrolidin-3-yl)carbamate



Step 1. Benzyl 3-azido-4-hydroxypyrrolidine-1-carboxylate

[00264] A solution of benzyl 6-oxa-3-azabicyclo[3.1.0]hexane-3-carboxylate (5 g, 22.81 mmol), NaN₃ (3 g, 46.15 mmol), NH₄Cl (1.23 g, 22.99 mmol) in MeOH (60 mL) and water (10 mL) was stirred for 16 h at 65 °C. After cooling to room temperature, the pH value was adjusted to 7-8 with aq. 0.5N NaOH. The resulting mixture was extracted with DCM (2 x 150 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum to give benzyl 3-azido-4-hydroxypyrrolidine-1-carboxylate as light-yellow solid. MS: (ESI, m/z): 263 [M+H]⁺.

Step 2. Benzyl 3-azido-4-methoxypyrrolidine-1-carboxylate

[00265] To a solution of benzyl 3-azido-4-hydroxypyrrolidine-1-carboxylate (4 g, 15.25 mmol) in DMF (40 mL) was added NaH (1.2 g, 60% dispersion in oil) at < 10 °C. The resulting solution was stirred for 1 h at room temperature. Iodomethane (2.8 mL, 44.98 mmol) was added and stirring was continued for another 1 h. The reaction was quenched by the addition of water (50 mL). The resulting mixture was extracted with Et₂O (2 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:1 EtOAc/pet. ether) afforded benzyl 3-azido-4-methoxypyrrolidine-1-carboxylate as a colorless oil. MS: (ESI, m/z): 277 [M+H]⁺.

Step 3. Benzyl 3-amino-4-methoxypyrrolidine-1-carboxylate

[00266] A solution of benzyl 3-azido-4-methoxypyrrolidine-1-carboxylate (2 g, 7.24 mmol) and PPh₃ (2.1 g, 8.01 mmol) in THF (50 mL) and water (5 mL) was stirred for 1 h at room temperature, then for an additional 5 h at 50°C. After cooling to room temperature, the reaction was concentrated under vacuum. Purification by prep-HPLC (Column: XBridge Prep C18 OBD, 19x50 mm, 5 µm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 10% B to 80% B in 30 min) afforded benzyl 3-amino-4-methoxypyrrolidine-1-carboxylate as a colorless oil. MS: (ESI, m/z): 251 [M+H]⁺.

Step 4. Benzyl 3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidine-1-carboxylate


[00267] A solution of benzyl 3-amino-4-methoxypyrrolidine-1-carboxylate (600 mg, 2.40 mmol), Et₃N (457 mg, 4.52 mmol), and (Boc)₂O (786 mg, 3.60 mmol) in THF (12 mL) and water (12 mL) was stirred for 30 min at room temperature. The resulting mixture was extracted with

DCM (2 x 30 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 EtOAc/pet. ether) to give benzyl 3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidine-1-carboxylate as a white solid. MS: (ESI, m/z): 351 [M+H]⁺.

Step 5. tert-Butyl N-(4-methoxypyrrolidin-3-yl)carbamate

[00268] A mixture of benzyl 3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidine-1-carboxylate (200 mg, 0.57 mmol) and Pd/C (200 mg, 10%) in EtOAc (10 mL) was stirred for 3 h at room temperature under an atmosphere of hydrogen. The solids were filtered out and washed with EtOAc (3x10mL). The filtrate was concentrated under vacuum to give tert-butyl N-(4-methoxypyrrolidin-3-yl)carbamate as a white solid. MS: (ESI, m/z): 217 [M+H]⁺.

Intermediate 36. tert-butyl ((3S,4S)-4-methoxypyrrolidin-3-yl)carbamate



Step 1. Benzyl (3S,4S)-3-azido-4-hydroxypyrrolidine-1-carboxylate

[00269] To a mixture of benzyl 6-oxa-3-azabicyclo[3.1.0]hexane-3-carboxylate (10 g, 44.70 mmol) and (1R,2R)-(-)-[1,2-cyclohexanediamino-N N'-bis(3,5-di-t-butylsalicylidene)]chromium (III) chloride (Jacobsen's Chromium catalyst) (682 mg, 1.08 mmol) was added azidotrimethylsilane (6.52 g, 56.59 mmol) at 30 °C. The reaction mixture was stirred for 16 h at 30 °C. Methanol (30 mL) and trifluoroacetic acid (0.5 mL) were added. The resulting solution was stirred at 30 °C for 3 h, then concentrated under vacuum. The residue was diluted with 100 mL of water and was extracted with ethyl acetate (3x100 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:2 EtOAc/pet.ether) afforded benzyl (3S,4S)-3-azido-4-hydroxypyrrolidine-1-carboxylate as a light yellow oil. MS: (ESI, m/z): 263 [M+H]⁺.

Step 2. Benzyl (3S,4S)-3-azido-4-methoxypyrrolidine-1-carboxylate

[00270] The title compound was prepared according to the procedures of Intermediate 35, Step 2, starting from benzyl (3S,4S)-3-azido-4-hydroxypyrrolidine-1-carboxylate. MS: (ESI, m/z): 277 [M+H]⁺.

Step 3. Benzyl (3S,4S)-3-amino-4-methoxypyrrolidine-1-carboxylate


[00271] The title compound was prepared according to the procedures of Intermediate 35, Step 3, starting from benzyl (3S,4S)-3-azido-4-methoxypyrrolidine-1-carboxylate. MS: (ESI, m/z): 251 [M+H]⁺.

Step 4. Benzyl (3S,4S)-3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidine-1-carboxylate

[00272] The title compound was prepared according to the procedures of Intermediate 35, Step 4, starting from benzyl (3S,4S)-3-amino-4-methoxypyrrolidine-1-carboxylate. MS: (ESI, m/z): 351 [M+H]⁺.

Step 5. tert-butyl ((3S,4S)-4-methoxypyrrolidin-3-yl)carbamate

[00273] The title compound was prepared according to the procedures of Intermediate 35, Step 5, starting from benzyl (3S,4S)-3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidine-1-carboxylate. MS: (ESI, m/z): 217 [M+H]⁺.

Intermediate 37. tert-Butyl 3-(trifluoromethyl)piperidin-3-ylcarbamate**Step 1. 1-Benzyl 3-methyl 3-(trifluoromethyl)piperidine-1,3-dicarboxylate**

[00274] To a solution of 1-benzyl 3-methyl piperidine-1,3-dicarboxylate (2.0 g, 7.14 mmol) in THF (60 mL) was added LDA (2.0 M in THF) (10.8 mL, 21.66 mmol) dropwise at -78 °C. After the solution was stirred for 30 min at -78 °C, S-(trifluoromethyl)dibenzothiophenium trifluoromethanesulfonate (4.35 g, 10.81 mmol) was added. After addition, the resulting solution was allowed to react, with stirring, for an additional 2 h at -40 °C. The reaction was then quenched

by the addition of 30 mL of a saturated aqueous solution of NH_4Cl . The resulting mixture was extracted with EtOAc (3 x 70 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:4 EtOAc/pet. ether) afforded 1-benzyl 3-methyl 3-(trifluoromethyl)piperidine-1,3-dicarboxylate as a yellow oil. MS: (ESI, m/z): 346 $[\text{M}+\text{H}]^+$.

Step 2. 1-((Benzyloxy)carbonyl)-3-(trifluoromethyl)piperidine-3-carboxylic acid

[00275] A solution of 1-benzyl 3-methyl 3-(trifluoromethyl)piperidine-1,3-dicarboxylate (310 mg, 0.83 mmol) in MeOH (10 mL) and aq. 1N NaOH (2.7 mL, 2.70 mmol) was stirred for 2 h at 50 °C and then concentrated under vacuum. The residue was diluted with 5 mL of water. The pH value of the solution was adjusted to 3-4 with 6N HCl. The solids were collected by filtration to give 1-((benzyloxy)carbonyl)-3-(trifluoromethyl)piperidine-3-carboxylic acid as a yellow solid. MS: (ESI, m/z): 332 $[\text{M}+\text{H}]^+$.


Step 3. Benzyl 3-((tert-butoxycarbonyl)amino)-3-(trifluoromethyl)piperidine-1-carboxylate

[00276] A solution of 1-((benzyloxy)carbonyl)-3-(trifluoromethyl)piperidine-3-carboxylic acid (250 mg, 0.76 mmol), Et_3N (229 mg, 2.26 mmol), and DPPA (415 mg, 1.51 mmol) in toluene (8 mL) was stirred for 2 h at 90 °C. *tert*-Butanol (279 mg, 3.76 mmol) was added and the reaction mixture was stirred for an additional 16 h at 90 °C. After cooling to room temperature, the reaction was quenched by the addition of water (20 mL) and was extracted with EtOAc (20 mL x 3). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:10 to 1:3 EtOAc/pet. ether) afforded benzyl 3-((tert-butoxycarbonyl)amino)-3-(trifluoromethyl)piperidine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 403 $[\text{M}+\text{H}]^+$.

Step 4. *tert*-Butyl (3-(trifluoromethyl)piperidin-3-yl)carbamate

[00277] A mixture of benzyl 3-((tert-butoxycarbonyl)amino)-3-(trifluoromethyl)piperidine-1-carboxylate (90 mg, 0.21 mmol) and Pd/C (10 mg, 10%) in EtOAc (5 mL) was stirred for 1 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum to give *tert*-butyl (3-(trifluoromethyl)piperidin-3-yl)carbamate as yellow oil. MS: (ESI, m/z): 269 $[\text{M}+\text{H}]^+$.

Intermediate 38. *tert*-Butyl (4-(trifluoromethoxy)pyrrolidin-3-yl)carbamate



Step 1. Benzyl 3-azido-4-hydroxypyrrolidine-1-carboxylate

[00278] A solution of benzyl 6-oxa-3-azabicyclo[3.1.0]hexane-3-carboxylate (5 g, 22.81 mmol), NaN_3 (3 g, 46.15 mmol), NH_4Cl (1.23 g, 22.99 mmol) in MeOH (60 mL) and water (10 mL) was stirred for 16 h at 65°C . After cooling to room temperature, the pH value was adjusted to 7-8 with aq. 0.5N NaOH . The resulting mixture was extracted with DCM (2 x 150 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum to give benzyl 3-azido-4-hydroxypyrrolidine-1-carboxylate as light-yellow solid. MS: (ESI, m/z): 263 $[\text{M}+\text{H}]^+$.

Step 2. Benzyl 3-((tert-butoxycarbonyl)amino)-4-hydroxypyrrolidine-1-carboxylate

[00279] A solution of benzyl 3-azido-4-hydroxypyrrolidine-1-carboxylate (6 g, 22.88 mmol), aq. 1 N NaOH (46 mL, 46 mmol), PMe_3 (1M in THF) (70 mL, 70 mmol), and $(\text{Boc})_2\text{O}$ (15.0 g, 68.73 mmol) in THF (228 mL) was stirred for 3 h at room temperature. The reaction mixture was diluted with 50 mL of water and was extracted with CH_2Cl_2 (3 x 200 mL). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered, and concentrated under vacuum. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (10 mM NH_4HCO_3), B: ACN ; Flow rate: 50 mL/min; Gradient: 0% B to 70% B in 40 min) afforded benzyl 3-((tert-butoxycarbonyl)amino)-4-hydroxypyrrolidine-1-carboxylate as a white solid. MS: (ESI, m/z): 337 $[\text{M}+\text{H}]^+$.

Step 3. Benzyl 3-((tert-butoxycarbonyl)amino)-4-(trifluoromethoxy)pyrrolidine-1-carboxylate


[00280] A solution of benzyl 3-((tert-butoxycarbonyl)amino)-4-hydroxypyrrolidine-1-carboxylate (1 g, 2.97 mmol), Selectfluor® (5.26 g, 14.85 mmol), AgOTf (7.6 g, 29.7 mmol), KF (1.72 g, 29.61 mmol), 2-fluoropyridine (2.88 g, 29.66 mmol) and TMS-CF_3 (4.22 g, 29.68 mmol)

in EtOAc (14 mL) was stirred overnight at room temperature. The solids were filtered out and washed with CH₂Cl₂ (3 x 50 mL). The combined filtrate was concentrated under vacuum and purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Flow rate: 50 mL/min; Gradient: 0% B to 80% B in 40 min) to give benzyl 3-((tert-butoxycarbonyl)amino)-4-(trifluoromethoxy)pyrrolidine-1-carboxylate as yellow oil. MS: (ESI, m/z): 405 [M+H]⁺.

Step 4. tert-Butyl (4-(trifluoromethoxy)pyrrolidin-3-yl)carbamate

[00281] A mixture of benzyl 3-((tert-butoxycarbonyl)amino)-4-(trifluoromethoxy)pyrrolidine-1-carboxylate (190 mg, 0.47 mmol) and Pd/C (20 mg, 10%) in EtOAc (10 mL) was stirred for 2 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum to give tert-butyl (4-(trifluoromethoxy)pyrrolidin-3-yl)carbamate as a yellow oil. MS: (ESI, m/z): 271 [M+H]⁺.

Intermediate 39. 4-(tert-butyldimethylsilyloxy)piperidine



Step 1. 1-Benzyl-4-(tert-butyldimethylsilyloxy)piperidine


[00282] A solution of 1-benzylpiperidin-4-ol (1 g, 5.23 mmol), imidazole (712 mg, 10.46 mmol), and tert-butyldimethylsilyl chloride (867 mg, 5.75 mmol) in DCM (20 mL) was stirred for 3 h at room temperature and then diluted with 20 mL of water. The resulting mixture was extracted with DCM (3 x 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded 1-benzyl-4-(tert-butyldimethylsilyloxy)piperidine as colorless oil. MS: (ESI, m/z): 306 [M+H]⁺.

Step 2. 4-(tert-Butyldimethylsilyloxy)piperidine

[00283] A mixture of 1-benzyl-4-(tert-butyldimethylsilyloxy)piperidine (500 mg, 1.64 mmol) and Pd/C (50 mg, 10%) in EtOAc (20 mL) was stirred for 2 h at room temperature under an atmosphere of hydrogen. The solids were filtered out and the filtrate was concentrated under


vacuum to give 4-(tert-butyldimethylsilyloxy)piperidine as colorless oil. MS: (ESI, m/z): 216 [M+H]⁺.

Intermediate 40. 3-((tert-butyldimethylsilyl)oxy)-3-methylpiperidine



[00284] A solution of 3-methylpiperidin-3-ol (500 mg, 3.91 mmol), tert-butyldimethylsilyl chloride (497 mg, 3.30 mmol), and Et₃N (837 mg, 8.27 mmol) in DCM (20 mL) was stirred for 24 h at room temperature. The reaction was quenched by the addition of 50 mL of water. The resulting mixture was extracted with DCM (3 x 40 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:20 to 1:5 EtOAc/pet.ether) afforded 3-((tert-butyldimethylsilyl)oxy)-3-methylpiperidine as colorless oil. MS: (ESI, m/z): 230 [M+H]⁺.

Intermediate 41. tert-Butyl 7-hydroxy-5-oxa-2-azaspiro[3.4]octane-2-carboxylate



Step 1. tert-Butyl 3-(2-ethoxy-2-oxoethylidene)azetidine-1-carboxylate

[00285] To a solution of ethyl 2-(diethoxyphosphoryl)acetate (26.2 g, 116.86 mmol) in THF (60 mL) was added NaH (4.68 g, 117.01 mmol, 60% dispersion in oil) in portions at 0 °C. The resulting solution was stirred for 30 min at room temperature. To the reaction mixture was added tert-butyl 3-oxoazetidine-1-carboxylate (10 g, 58.41 mmol) with stirring at 0 °C. The resulting solution was stirred for an additional 30 min at room temperature. The reaction was then quenched by the addition of 30 mL of a water/ice mixture. The resulting solution was extracted with DCM (3 x 40 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated

under vacuum. Purification by silica gel chromatography (eluting with 1:5 EtOAc/pet. ether) afforded tert-butyl 3-(2-ethoxy-2-oxoethylidene)azetidine-1-carboxylate as yellow oil. MS: (ESI, m/z): 242 [M+H]⁺.

Step 2: 2-(tert-Butyl) 8-ethyl 7-oxo-5-oxa-2-azaspiro[3.4]octane-2,8-dicarboxylate

[00286] To a solution of NaH (1.2 g, 30.00 mmol, 60% dispersion in oil) in Et₂O (20 mL) was added methyl 2-hydroxyacetate (2.69 g, 29.86 mmol) at 0 °C. The resulting solution was stirred for 30 min at room temperature and then concentrated under vacuum. The residue was diluted with 20 mL of DMSO and tert-butyl 3-(2-ethoxy-2-oxoethylidene)azetidine-1-carboxylate (6 g, 24.87 mmol) was added. The resulting solution was stirred overnight at room temperature. The pH value of the solution was adjusted to 4-5 with hydrochloric acid (1N). The resulting mixture was extracted with Et₂O (4 x 20 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:10 to 1:5 EtOAc/pet. ether) afforded 2-(tert-butyl) 8-ethyl 7-oxo-5-oxa-2-azaspiro[3.4]octane-2,8-dicarboxylate as a light yellow solid. MS: (ESI, m/z): 300 [M+H]⁺.

Step 3. tert-Butyl 7-oxo-5-oxa-2-azaspiro[3.4]octane-2-carboxylate


[00287] A solution of 2-(tert-butyl) 8-ethyl 7-oxo-5-oxa-2-azaspiro[3.4]octane-2,8-dicarboxylate (4 g, 13.4 mmol) and NaCl (1.33 g, 22.76 mmol) in DMSO/water (10:1, 20 mL) was stirred for 2 h at 120 °C. After cooling to room temperature, the reaction was quenched with 20 mL of water. The resulting mixture was extracted with Et₂O (4 x 20 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded tert-butyl 7-oxo-5-oxa-2-azaspiro[3.4]octane-2-carboxylate as a white solid. MS: (ESI, m/z): 228 [M+H]⁺.

Step 4. tert-Butyl 7-hydroxy-5-oxa-2-azaspiro[3.4]octane-2-carboxylate

[00288] To a solution of tert-butyl 7-oxo-5-oxa-2-azaspiro[3.4]octane-2-carboxylate (1.1 g, 4.84 mmol) in THF (8 mL) was added NaH (276 mg, 7.49 mmol) in portions at 0 °C. The resulting mixture was stirred for 30 min at room temperature. The reaction was quenched with 50 mL of a water/ice mixture. The resulting mixture was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A:

water (0.5% TFA), B: ACN; Gradient: 0% to 50% B over 35 min) afforded tert-butyl 7-hydroxy-5-oxa-2-azaspiro[3.4]octane-2-carboxylate as yellow oil. MS: (ESI, m/z): 230 [M+H]⁺.

Intermediate 42. Benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate



Step 1. (4,6-Dichloropyridin-3-yl)methanol

[00289] Into a 500-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 4,6-dichloronicotinaldehyde (8 g, 43.18 mmol) and ethanol (200 mL). NaBH₄ (5.2 g, 141.21 mmol) was added in portions at 0 °C. Then the resulting solution was stirred for 2 h at 25 °C. The reaction was then quenched by the addition of water (200 mL). The resulting mixture was extracted with DCM (3 x 300 mL) and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 ethyl acetate/pet. ether) to give of ((4,6-dichloropyridin-3-yl)methanol as a colorless oil. MS: (ESI, m/z): 178,180 [M+H]⁺.

Step 2. 5-(Bromomethyl)-2,4-dichloropyridine

[00290] Into a 250-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed (4,6-dichloropyridin-3-yl)methanol (5 g, 26.68 mmol), DCM (100 mL) and PBr₃ (7.7 g, 28.45 mmol). The resulting solution was stirred for 30 min at 40 °C in an oil bath. After cooling to 25 °C, the reaction was then quenched by the addition of water (120 mL). The resulting mixture was extracted with DCM (2 x 150 mL) and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:30 ethyl acetate/pet. ether) to

give 5-(bromomethyl)-2,4-dichloropyridine as a colorless oil. MS: (ESI, m/z): 240, 242, 244 [M+H]⁺.

Step 3. (2R,5S)-2-((4,6-Dichloropyridin-3-yl)methyl)-5-isopropyl-3,6-dimethoxy-2,5-dihydropyrazine

[00291] Into a 500-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed (2S)-3,6-dimethoxy-2-(propan-2-yl)-2,5-dihydropyrazine (3 g, 15.47 mmol) and THF (200 mL). A solution of n-BuLi in n-hexane (2.5 M) (9.8 mL, 24.5 mmol) was added at -80 °C in a liquid nitrogen bath. The resulting solution was stirred for 30 min at -80 °C in a liquid nitrogen bath. Then a solution of 5-(bromomethyl)-2,4-dichloropyridine (4.7 g, 18.53 mmol) in THF (10 mL) was added. The resulting solution was allowed to react, with stirring, for an additional 1 h while the temperature was maintained at -80 °C in a liquid nitrogen bath. The reaction was then quenched by the addition of 120 mL of ammonium chloride (sat. aq.). The resulting mixture was extracted with DCM (2 x 150 mL) and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 ethyl acetate/pet. ether) to afford (2R,5S)-2-[(4,6-dichloropyridin-3-yl)methyl]-3,6-dimethoxy-5-(propan-2-yl)-2,5-dihydropyrazine as a white solid. MS: (ESI, m/z): 344, 346 [M+H]⁺.

Step 4. Methyl (2R)-2-amino-3-(4,6-dichloropyridin-3-yl)propanoate

[00292] Into a 250-mL round-bottom flask, was placed (2R,5S)-2-[(4,6-dichloropyridin-3-yl)methyl]-3,6-dimethoxy-5-(propan-2-yl)-2,5-dihydropyrazine (3.5 g, 9.66 mmol), hydrochloride acid (0.3M) (70 mL) and ACN (80 mL). The resulting solution was stirred for 2 h at 25 °C. The reaction was then quenched by the addition of 100 mL of sodium bicarbonate (sat. aq.). The resulting mixture was extracted with 2 x 150 mL of DCM and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give methyl (2R)-2-amino-3-(4,6-dichloropyridin-3-yl)propanoate as colorless oil. MS: (ESI, m/z): 249, 251 [M+H]⁺.

Step 5. (2R)-2-Amino-3-(4,6-dichloropyridin-3-yl)propan-1-ol

[00293] Into a 250-mL round-bottom flask, was placed methyl (2R)-2-amino-3-(4,6-dichloropyridin-3-yl)propanoate (1.5 g, 5.72 mmol) and methanol (50 mL). NaBH₄ (690 mg, 18.24 mmol) was added in portion at 0 °C. The resulting solution was stirred for 3 h at 25 °C. The reaction

was then quenched by the addition of water (20 mL). The resulting mixture was extracted with DCM (2 x 50 mL) and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 methanol/DCM) to afford (2R)-2-amino-3-(4,6-dichloropyridin-3-yl)propan-1-ol as a white solid. MS: (ESI, m/z): 221, 223 [M+H]⁺.


Step 6. (R)-7-Chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-amine

[00294] Into a 50-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed (2R)-2-amino-3-(4,6-dichloropyridin-3-yl)propan-1-ol (1 g, 4.30 mmol), sodium hydride (271 mg, 6.78 mmol, 60%) and DMSO (10 mL). The resulting solution was stirred for 16 h at 50 °C in an oil bath. After cooling to 25 °C, the reaction was then quenched by the addition of water (20 mL). The resulting mixture was extracted with DCM (2 x 30 mL) and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.1% formic acid), B: ACN; Flow rate: 50 mL/min; Gradient: 0% B to 30% B in 30 min). The collected fraction was concentrated under vacuum to give (R)-7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-amine as a white solid. MS: (ESI, m/z): 185, 187 [M+H]⁺.

Step 7. Benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate

[00295] Into a 50-mL round-bottom flask, was placed (R)-7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-amine (500 mg, 2.57 mmol), ethyl acetate (15 mL, 153.23 mmol), water (15 mL), benzyl chloroformate (558 mg, 3.27 mmol) and potassium carbonate (751 mg, 5.43 mmol). The resulting mixture was stirred for 30 min at 25 °C. The resulting mixture was extracted with 3 x 20 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate as a white solid. MS: (ESI, m/z): 319, 321 [M+H]⁺.

Intermediate 43. Benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)carbamate



Step 1. 3-(Bromomethyl)-2,6-dichloropyridine

[00296] Into a 500-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed (2,6-dichloropyridin-3-yl)methanol (10 g, 56.17 mmol), DCM (200 mL) and PBr_3 (15.3 g, 56.52 mmol). The resulting solution was stirred for 30 min at 40 °C in an oil bath. The pH value of the solution was adjusted to 7 with NH_4HCO_3 (sat. aq.). The resulting mixture was extracted with 3 x 200 mL of DCM. The organic layers combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 to 1:5 ethyl acetate/pet. ether) to give 3-(bromomethyl)-2,6-dichloropyridine as an off-white solid. MS: (ESI, m/z): 240, 242, 244 $[\text{M}+\text{H}]^+$.

Step 2. (2R,5S)-2-((2,6-Dichloropyridin-3-yl)methyl)-5-isopropyl-3,6-dimethoxy-2,5-dihydropyrazine

[00297] Into two 250-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed (2S)-3,6-dimethoxy-2-(propan-2-yl)-2,5-dihydropyrazine (2 g, 10.86 mmol) and THF (15 mL). This was followed by the addition of butyllithium (2.5 M) (6.5 mL, 16.25 mmol) at -78 °C. The resulting solution was stirred for 30 min. To this was added 3-(bromomethyl)-2,6-dichloropyridine (2.62 g, 10.88 mmol). The resulting solution was stirred for an additional 1 h at -78°C. The reaction was then quenched by the addition of 5 mL of NH_4HCO_3 (saturated) and the mixture was diluted with 10 mL of H_2O . The resulting mixture of two batches was combined and extracted with 3 x 50 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1/5 ethyl acetate/pet. ether) to afford (2R,5S)-

2-((2,6-dichloropyridin-3-yl)methyl)-5-isopropyl-3,6-dimethoxy-2,5-dihydropyrazine as a colorless oil. MS: (ESI, m/z): 344, 346 [M+H]⁺.

Step 3. Methyl (R)-2-amino-3-(2,6-dichloropyridin-3-yl)propanoate

[00298] Into a 500-mL round-bottom flask, was placed (2R,5S)-2-((2,6-dichloropyridin-3-yl)methyl)-5-isopropyl-3,6-dimethoxy-2,5-dihydropyrazine (5.4 g, 16.35 mmol), acetonitrile (100 mL) and hydrochloric acid (0.3 N) (157 mL). The resulting solution was stirred for 1 hour at 24 °C. The solvent was removed under vacuum and the residue was extracted with 3x100 mL of DCM. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:5 to 1:1 ethyl acetate/pet. ether) to afford to give methyl (R)-2-amino-3-(2,6-dichloropyridin-3-yl)propanoate as an off-white solid. MS: (ESI, m/z): 249, 251 [M+H]⁺.

Step 4. (R)-2-Amino-3-(2,6-dichloropyridin-3-yl)propan-1-ol

[00299] Into a 250-mL round-bottom flask, was placed methyl (R)-2-amino-3-(2,6-dichloropyridin-3-yl)propanoate (3.2 g, 12.85 mmol), methanol (15 mL), THF (60 mL) and NaBH₄ (1.46 g, 39.65 mmol). The resulting solution was stirred for 2 h at 24 °C. The reaction was then quenched by the addition of 5 mL of water. The solvent was removed under vacuum. The residue was diluted with 50 mL of water. The resulting mixture was extracted with 3 x 50 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:20 methanol/DCM) to afford (R)-2-amino-3-(2,6-dichloropyridin-3-yl)propan-1-ol as an off-white solid. MS: (ESI, m/z): 221, 223 [M+H]⁺.

Step 5. (R)-7-Chloro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-amine


[00300] Into a 250-mL round-bottom flask fitted with a hydrogen balloon, was placed (R)-2-amino-3-(2,6-dichloropyridin-3-yl)propan-1-ol (2.6 g, 11.76 mmol) and DMSO (30 mL). This was followed by the addition of sodium hydride (706 mg, 29.42 mmol, 60%) at 0 °C. The resulting solution was stirred at 24 °C for 30 min and then stirred overnight at 50 °C. After cooling to 25 °C, the reaction was then quenched by the addition of 60 mL of water. The resulting mixture was extracted with 3 x 60 mL of ethyl acetate. The organic layers combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN;

Flow rate: 50 mL/min; Gradient: 0% B to 30% B in 30 min). The collected fraction was concentrated under vacuum to give (R)-7-chloro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-amine as an off-white solid. MS: (ESI, m/z): 185, 187 [M+H]⁺.

Step 6. Benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)carbamate

[00301] Into a 100-mL round-bottom flask, was placed (R)-7-chloro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-amine (1.3 g, 6.34 mmol), ethyl acetate (15 mL), water (10 mL), potassium carbonate (1.95 g, 14.11 mmol) and benzyl chloroformate (1.44 g, 8.44 mmol). The resulting solution was stirred for 2 h at 25 °C. The reaction was then quenched by the addition of 30 mL of water. The resulting mixture was extracted with 3x30 mL of ethyl acetate and the organic layers combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by re-crystallization from pet. ether. The solids were collected by filtration to give benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)carbamate as a white solid. MS: (ESI, m/z): 319, 321 [M+H]⁺.

Intermediate 44. tert-Butyl 4-[(3R)-3-[(benzyloxy)carbonyl]amino]-3,4-dihydro-2H-1-benzopyran-7-yl]-4-fluoropiperidine-1-carboxylate



Step 1. 7-Bromo-4-methyl-3-nitro-3,4-dihydro-2H-1-benzopyran


To a solution of 7-bromo-3-nitro-2H-chromene (2.2 g, 8.59 mmol) in THF (30 mL) was added CH₃MgBr (1 M in THF) (13 mL, 13.00 mmol) at -78 °C. The resulting solution was stirred for 20 min at -78 °C. The reaction was then quenched by the addition of 20 mL of saturated aqueous NH₄Cl solution. The resulting mixture was extracted with 3 x 50 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 3:1 pet. ether/ethyl acetate) to afford 7-bromo-4-methyl-3-nitro-3,4-dihydro-2H-1-benzopyran as yellow oil. GCMS: (ESI, m/z): 271, 273 [M+H]⁺.

Step 2. 7-Bromo-4-methyl-3,4-dihydro-2H-1-benzopyran-3-amine

To a solution of $\text{BH}_3\text{-THF}$ (1M) (40 mL, 40.00 mmol) and NaBH_4 (1.26 g, 33.30 mmol) in THF (40 mL) was added 7-bromo-4-methyl-3-nitro-3,4-dihydro-2H-1-benzopyran (900 mg, 3.31 mmol). The resulting mixture was stirred in a sealed tube for 16 h at 75 °C. Then methanol (40 mL) was added and the reaction mixture was stirred for 6 h at 80 °C. After cooling to 25 °C, the solvent was removed under vacuum. The residue was diluted with 50 mL of ice/water. The resulting mixture was extracted with 3 x 50 of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.05% TFA), B: ACN; Gradient: 0% B to 40% over 45 min) to afford 7-bromo-4-methyl-3,4-dihydro-2H-1-benzopyran-3-amine as a white solid. MS: (ESI, m/z): 242, 244 $[\text{M}+\text{H}]^+$.

Step 3. Benzyl N-(7-bromo-4-methyl-3,4-dihydro-2H-1-benzopyran-3-yl) carbamate

A solution of 7-bromo-4-methyl-3,4-dihydro-2H-1-benzopyran-3-amine (500 mg, 2.07 mmol) and CbzCl (423 mg, 2.48 mmol) in ethyl acetate (8 mL) and a solution of potassium carbonate (567 mg, 4.10 mmol) in water (8 mL) was stirred for 1 h at 25 °C. The resulting solution was diluted with 20 mL of water. The resulting mixture was extracted with 3 x 20 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 3:1 pet. ether/ethyl acetate) to afford benzyl N-(7-bromo-4-methyl-3,4-dihydro-2H-1-benzopyran-3-yl)carbamate as a white solid. MS: (ESI, m/z): 376, 378 $[\text{M}+\text{H}]^+$.

Intermediate 45. tert-Butyl 6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane-1-carboxylate**Step 1. 1,4-Dibenzyl-1,4-diazepan-6-ol**

A solution of 1,3-dibromopropan-2-ol (12.37 g, 56.77 mmol), benzyl(2-(benzylamino)ethyl)amine (13.56 g, 56.42 mmol), and Et₃N (17.17 g, 169.68 mmol) in toluene (50 mL) was stirred for 48 h at 120 °C. After cooling to 25 °C, the reaction was then quenched by the addition of 100 mL of water. The resulting mixture was extracted with 2 x 100 mL of ethyl acetate and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:2 ethyl acetate/pet. ether) to afford 1,4-dibenzyl-1,4-diazepan-6-ol as yellow oil. MS: (ESI, m/z): 297 [M+H]⁺.

Step 2. 1,4-Dibenzyl-6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane

A solution of 1,4-dibenzyl-1,4-diazepan-6-ol (2.0 g, 6.61 mmol), imidazole (900 mg, 13.22 mmol) and TBS-Cl (1.2 g, 7.96 mmol) in DMF (10 mL) was stirred for 16 h at 25 °C. The reaction was then quenched by the addition of 50 mL of water. The resulting mixture was extracted with 2 x 50 mL of ethyl acetate and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:5 ethyl acetate/pet. ether) to afford 1,4-dibenzyl-6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane as yellow oil. MS: (ESI, m/z): 411 [M+H]⁺.

Step 3. 6-((tert-Butyldimethylsilyl)oxy)-1,4-diazepane


A mixture of 1,4-dibenzyl-6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane (1.5 g, 3.47 mmol) and Pd(OH)₂ on carbon (20 mg, 10%) in ethyl acetate (10 mL) was stirred under an atmosphere of hydrogen for 28 h at 20 °C. The solids were filtered out. The filtrate was concentrated under vacuum to afford 6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane as colorless oil. MS: (ESI, m/z): 231 [M+H]⁺.

Step 4. tert-Butyl 6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane-1-carboxylate

A solution of 6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane (1.2 g, 4.95 mmol), Boc₂O (700 mg, 3.21 mmol), and triethylamine (500 mg, 4.94 mmol) in DCM (20 mL) was stirred for 2 h at -40 °C in a dry ice/ethanol bath and stirred for an additional 1 h at 20 °C. The reaction was then quenched by the addition of 40 mL of water. The resulting mixture was extracted with 2 x 40 mL of ethyl acetate and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with

1:10 methanol/DCM) to afford tert-butyl 6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane-1-carboxylate as colorless oil. MS: (ESI, m/z): 331 [M+H]⁺.

Intermediate 46. tert-Butyl 6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane-1-carboxylate




To a solution of tert-butyl 2-(hydroxymethyl)piperazine-1-carboxylate (1 g, 4.39 mmol) and triethylamine (1.53 mL, 10.98 mmol) in DCM (20 mL) was added tert-butyl(chloro)dimethylsilane (693 mg, 4.60 mmol). The resulting solution was stirred overnight at 20 °C. The reaction was then quenched by the addition of 20 mL of water. The resulting mixture was extracted with 3 x 20 mL of DCM. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 20:1 ethyl DCM/methanol) to afford tert-butyl 6-((tert-butyldimethylsilyl)oxy)-1,4-diazepane-1-carboxylate as an off-colorless oil. MS: (ESI, m/z): 331 [M+H]⁺.

Intermediate 47-1. Benzyl N-[2-[(trifluoromethane)sulfonyloxy]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

Intermediate 47-2. Benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

Intermediate 47-3. Benzyl N-[(6R)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

Method 1. Chiral separation



Step 1. 7',8'-Dihydro-5'H-spiro[1,3-dioxolane-2,6'-quinoline]-2'-ol

A solution of 1,4-dioxaspiro[4.5]decan-8-one (50.0 g, 320 mmol), ethyl prop-2-ynoate (68 mL, 672 mmol) and a solution of NH_3 in MeOH (230 mL, 7M) in isopropanol (1.3 L) was stirred for 14 h at 130 °C. The reaction mixture was cooled to 20 °C. The solvent was removed under vacuum. The solids were collected by filtration. The cake was washed with 2 x 30 mL of pet. ether and dried under vacuum to give 7',8'-dihydro-5'H-spiro[1,3-dioxolane-2,6'-quinoline]-2'-ol as an off-white solid. MS (ESI, m/z): 208 $[\text{M}+\text{H}]^+$.

Step 2. 7',8'-Dihydro-5'H-spiro[1,3-dioxolane-2,6'-quinoline]-2'-yl trifluoromethanesulfonate

To a solution of 7',8'-dihydro-5'H-spiro[1,3-dioxolane-2,6'-quinoline]-2'-ol (24.0 g, 116 mmol) and triethylamine (64.4 mL, 464 mmol) in DCM (500 mL) was added a solution of trifluoromethanesulfonic anhydride (58.4 mL, 347 mmol) in DCM (100 mL) at 10 °C. The resulting solution was stirred for 1 h at 20 °C. The mixture was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 3:1 ethyl acetate/pet. ether) to afford 7',8'-dihydro-5'H-spiro[1,3-dioxolane-2,6'-quinoline]-2'-yl trifluoromethanesulfonate as a yellow oil. MS (ESI, m/z): 340 $[\text{M}+\text{H}]^+$.

Step 3. 6-Oxo-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate

A solution of 7',8'-dihydro-5'H-spiro[1,3-dioxolane-2,6'-quinoline]-2'-yl trifluoromethanesulfonate (35.0 g, 103 mmol) and hydrochloric acid (206 mL, 1N) in acetone (300

mL) was stirred for 1 h at 75 °C. The mixture was cooled to 20 °C. The acetone was concentrated under vacuum. The pH value of the residue was adjusted to 7-8 with NaHCO₃ (sat., aq.). The resulting mixture was extracted with 3 x 100 mL of DCM. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by recrystallization with 100 mL of the mixture solvent of ethyl acetate and pet. ether (10:1) to give 6-oxo-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate as an off-white solid. MS (ESI, m/z): 296 [M+H]⁺.

Step 4. 6-Amino-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate

A mixture of 6-oxo-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate (16.0 g, 54.2 mmol) and NH₄OAc (50.1 g, 650 mmol) in methanol (250 mL) was stirred for 1 h at 20 °C. NaBH₃CN (4.10 g, 65.2 mmol) was added in portions at 10 °C. The resulting mixture was allowed to react for 13 h at 20 °C. The reaction was then quenched by the addition of 50 mL of water/ice. The solvent was concentrated under vacuum. The residue was diluted with 100 mL of hydrochloric acid (1N). The resulting mixture was extracted with 2 x 100 mL of DCM. The pH value of aqueous phase was adjusted to 10 with Na₂CO₃ (sat., aq.). The aqueous phase was extracted with 4 x 100 mL of DCM. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give 6-amino-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate as a yellow oil. MS (ESI, m/z): 297 [M+H]⁺.

Step 5. Benzyl N-[2-[(trifluoromethane)sulfonyloxy]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate


A mixture of 6-amino-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate (12.0 g, 40.5 mmol), potassium carbonate (14.0 g, 101 mmol) and CbzCl (7.4 mL, 52.8 mmol) in ethyl acetate (100 mL) was stirred for 2 h at 20 °C. The reaction was then quenched by the addition of 100 mL of water/ice. The mixture was extracted with 3 x 100 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by recrystallization with 80 mL of the mixture solvent of ethyl acetate and pet. ether (10:1) to give benzyl N-[2-[(trifluoromethane)sulfonyloxy]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as an off-white solid. MS (ESI, m/z): 431 [M+H]⁺.

Step 6. Benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate and benzyl N-[(6R)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

The racemate benzyl N-[2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-1**) (7.00 g, 16.3 mmol,) was separated by SFC (Column: Chiralpak AD-H SFC, 5 x 25 cm, 5 μ m; Mobile Phase, A: CO₂: 55% and B: MeOH: 45%; Flow rate: 150 mL/min). The first eluting isomer (RT = 4.23 min) was collected and concentrated under vacuum to give benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) as an off-white solid. And the second eluting isomer (RT = 5.16 min) was collected and concentrated under vacuum to give benzyl N-[(6R)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-3**) as an off-white solid. MS (ESI, m/z) for both isomers: 431 [M+H]⁺.

Intermediate 47-2. Benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

Method 2. Chiral resolution



Step 1. (S)-6-Amino-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate

A solution of 6-amino-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate (183 g, 525.65 mmol) and L-(-)-mandelic acid (39.99 g, 262.83 mmol) in EtOH (1830 mL) was stirred at 70 °C for 2 h. The mixture was allowed to cool down to 25 °C slowly and stirred for overnight. The precipitate that was formed was collected by filtration and dried under vacuum to afford the chiral salt as a white solid. The salt was dissolved in H₂O (350 mL) and the pH value of aqueous phase was adjusted to 12 - 14 with NaOH (1 N, aq). The aqueous phase was extracted with EtOAc (500 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated under vacuum to afford the free base of the chiral amine (S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl


trifluoromethanesulfonate as a yellow oil. MS (ESI, m/z): 297 $[M+H]^+$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm): 7.79 (d, $J = 8.0$ Hz, 1H), 7.25 (d, $J = 8.4$ Hz, 1H), 3.14-3.11 (m, 1H), 2.97-2.81 (m, 3H), 2.54-2.52 (m, 1H), 2.61-2.58 (m, 2H), 1.64-1.59 (m, 1H).

Step 2. Benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

To a solution of (S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl trifluoromethanesulfonate (48.0 g, 162.02 mmol) and K_2CO_3 (55.98 g, 405.04 mmol) in EtOAc (480 mL) was added drop-wise CbzCl (35.93 g, 210.62 mmol, 29.94 mL) at 0 °C. The mixture was stirred at 20 °C for 30 min. The reaction was then quenched by water/ice (150 mL). The mixture was extracted with ethyl acetate (350 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The crude was triturated with pet. ether (200 mL) at 20 °C. The resulting precipitate was collected by filtration and dried under vacuum to afford benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a pale pink solid. MS (ESI, m/z): 431 $[M+H]^+$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm): 7.82 (d, $J = 8.0$ Hz, 1H), 7.49 (m, 1H), 7.37-7.28 (m, 6H), 5.04 (s, 2H), 3.82 (m, 1H), 3.08 (dd, $J = 16.8$ Hz, 4.8 Hz, 1H), 2.91-2.88 (m, 2H), 2.76-2.72 (m, 1H), 2.02-2.00 (m, 1H), 1.83-1.75 (m, 1H).

The following intermediate in Table 7 was prepared using standard chemical manipulations and procedures similar to those used for the preparation of Intermediate 47-1.

Table 7:

Intermediate Number	Structure and Name	LRMS m/z $[M+H]^+$
47-4 ¹	 benzyl N-[2'-(trifluoromethanesulfonyloxy)-6',7'-dihydro-5'H-spiro[cyclopropane-1,8'-quinolin]-6'-yl]carbamate	457


¹ **Notes on procedures:** In Step 1, the ketone used was 6,9-dioxadispiro[2.1.4⁵.3³]dodecan-12-one, which was prepared by the following method: (2-chloroethyl)dimethylsulfonium iodide (88.0 g, 347 mmol) was slowly added to a solution of 1,4-dioxaspiro[4.5]decan-8-one (60.0 g, 384 mmol) and t-BuOK (88.0 g, 784

mmol) in t-BuOH (1.50 L) over 4 h. The resulting solution was stirred for 1 h at 25 °C. The solids were filtered. The filtrate was concentrated under vacuum. The residue was diluted with water (400 mL). The resulting mixture was extracted with DCM (3 x 400 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified via reverse phase chromatography to afford 6,9-dioxadispiro[2.1.4⁵.3³]dodecan-12-one as a pale yellow oil. MS (ESI, m/z): 183 [M+H]⁺.

Intermediate 48-1. cis-tert-Butyl N-[4-(methoxymethyl)pyrrolidin-3-yl]carbamate;

Intermediate 48-2. tert-Butyl N-[(3R,4R)-4-(methoxymethyl)pyrrolidin-3-yl]carbamate and

Intermediate 48-3. Benzyl (3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate



Step 1. cis-Benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(hydroxymethyl)pyrrolidine-1-carboxylate

A solution of cis-benzyl 3-amino-4-(hydroxymethyl)pyrrolidine-1-carboxylate (2.00 g, 7.99 mmol), Et₃N (2.13 mL, 23.7 mmol) and (Boc)₂O (2.20 g, 10.1 mmol) in THF (40 mL) and water (20 mL) was stirred for 2 h at 20 °C. The solvent was removed under vacuum. The residue was diluted with water (20 mL). The resulting mixture was extracted with ethyl acetate (3 x 20 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by a silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to give cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(hydroxymethyl)pyrrolidine-1-carboxylate as an off-white solid. MS (ESI, m/z): 351 [M+H]⁺.

Step 2. cis-Benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate

A solution of trimethyloxoniumtetrafluoroborate (1.50 g, 10.4 mmol) in DCM (20 mL) was added to a stirring solution of 1,8-bis(dimethylamino)naphthalene (2.20 g, 10.3 mmol) and cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(hydroxymethyl)pyrrolidine-1-carboxylate (2.20 g, 6.09 mmol) in DCM (100 mL). The resulting mixture was stirred for 3 h at 40 °C. The pH of the mixture was adjusted to 4-6 with HCl (3N, aq.). The resulting mixture was extracted with ethyl acetate (3 x 100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate as white solid. MS (ESI, m/z): 365 [M+H]⁺.

Step 3. cis-tert-Butyl N-[4-(methoxymethyl)pyrrolidin-3-yl]carbamate

A mixture of cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate (1.00 g, 2.74 mmol) and palladium on carbon (1.00 g, 10%) in ethyl acetate (50 mL) was stirred for 1 h at 20 °C under a hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give cis-tert-butyl N-[4-(methoxymethyl)pyrrolidin-3-yl]carbamate (**Intermediate 48-1**) as yellow oil (crude). MS (ESI, m/z): 231 [M+H]⁺.


Step 4. tert-Butyl N-[(3R,4R)-4-(methoxymethyl)pyrrolidin-3-yl]carbamate

The racemate cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate was separated into its enantiomers by SFC (Column: Chiralpak IA-SFC, 5 x 25 cm, 5 µm; Mobile Phase, A: CO₂: 70% and B: MeOH (containing 2 mM NH₃ in MeOH): 30%; Flow rate: 150 mL/min). The first eluting isomer (RT = 3.91 min) was collected and concentrated under vacuum to give benzyl (3S,4S)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate as a yellow solid. The second eluting isomer (RT = 5.04 min) was collected and concentrated under vacuum to give benzyl (3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate (**Intermediate 48-3**) as a yellow solid. MS (ESI, m/z): 365 [M+H]⁺.

A mixture of benzyl (3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate (**Intermediate 48-3**) (0.800 g, 2.20 mmol) and palladium on carbon (0.800 g, 10%) in ethyl acetate (30 mL) was stirred for 1 h at 25 °C under a hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give tert-butyl N-[(3R,4R)-4-(methoxymethyl)pyrrolidin-3-yl]carbamate (**Intermediate 48-2**) as a yellow oil (crude). MS (ESI, m/z): 231 [M+H]⁺.

Intermediate 48-3. Benzyl (3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate

Method 2.



Step 1. Benzyl N-(2,2-dimethoxyethyl)carbamate

A solution of 2,2-dimethoxyethan-1-amine (1600 g, 15.22 mol) in toluene (8 L), was added a solution of NaOH (858 g, 21.45 mol) in water (4.42 L). This was followed by the addition of CbzCl (2598 g, 15.23 mol) dropwise with stirring at <20 °C. The resulting solution was stirred for 4 h at room temperature. The organic layer was separated and washed with 3x5 L of brine. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum to afford benzyl N-(2,2-dimethoxyethyl)carbamate as a white solid.

Step 2. Benzyl N-(2,2-dimethoxyethyl)-N-(prop-2-en-1-yl)carbamate

To a solution of benzyl N-(2,2-dimethoxyethyl)carbamate (1700 g, 7.10 mol), KOH (1755 g, 31.28 mol) and benzyltriethylammonium chloride (32.37g, 142.1 mmol) in toluene (7.82 L) was added

3-bromoprop-1-ene (1117.4 g, 9.24 mol) dropwise with stirring at room temperature. The resulting solution was stirred for 24 h at room temperature. Two batches were thus run in parallel. The reaction was then quenched by the addition of 10 L of water. The resulting solution was extracted with 2x7 L of toluene and the organic layers combined. The organic phase was washed with 2x10 L of brine. The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum to afford benzyl N-(2,2-dimethoxyethyl)-N-(prop-2-en-1-yl)carbamate as pale yellow oil.

Step 3. Benzyl N-(2-oxoethyl)-N-(prop-2-en-1-yl)carbamate

A solution of benzyl N-(2,2-dimethoxyethyl)-N-(prop-2-en-1-yl)carbamate (3900 g, 13.96 mol) in 88% formic acid (5460 mL) was stirred overnight at room temperature. The resulting mixture was concentrated under vacuum. The residue was diluted with 20 L of ethyl acetate, washed with 4x10 L of brine. The organic phase was dried over anhydrous sodium sulfate and concentrated under vacuum to afford benzyl N-(2-oxoethyl)-N-(prop-2-en-1-yl)carbamate as brown oil.

Step 4. Benzyl N-[2-(hydroxyimino)ethyl]-N-(prop-2-en-1-yl)carbamate

To a solution of benzyl N-(2-oxoethyl)-N-(prop-2-en-1-yl)carbamate (1700 g, 7.29 mol) and $\text{NH}_2\text{OH}\cdot\text{HCl}$ (638 g, 9.18 mol) in ACN (9.52 L) was added a solution of NaOAc (669 g, 8.16 mol) in H_2O (4.96 L). The resulting solution was stirred for 20 h at room temperature. Two batches were thus run in parallel. The resulting mixture was concentrated under vacuum and extracted with 3x10 L of ethyl acetate and the organic layers combined. The organic phase was washed with 3x5 L of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The crude product was purified by silica gel chromatography (eluting with 1:9 EtOAc/pet. ether) to afford benzyl N-[2-(hydroxyimino)ethyl]-N-(prop-2-en-1-yl)carbamate as colorless oil.

Step 5. cis-Benzyl 3-amino-4-(hydroxymethyl)pyrrolidine-1-carboxylate

A solution of benzyl N-[2-(hydroxyimino)ethyl]-N-(prop-2-en-1-yl)carbamate (1200 g, 4.83 mol) and xylene (6 L) was stirred overnight at 130 °C. The mixture was cooled to room temperature. HOAc (6 L) was added to the mixture. Then Zn (1200 g, 18.35 mol) was added into the mixture at 15 °C. The resulting solution was stirred overnight at room temperature in a water bath. Two batches were thus run in parallel. The batches were combined and filtered. The filtered cake was washed with xylene. The combined filtrate was diluted with 20 L of water. The mixture was extracted with xylene (4x5 L). The aqueous phase was concentrated under vacuum. The pH value

of the residue was adjusted to 9-10 with saturated sodium carbonate. The solids were filtered out. The filtrate was concentrated under vacuum. The residue was extracted with 5 L of THF and concentrated under vacuum to afford cis-benzyl 3-amino-4-(hydroxymethyl)pyrrolidine-1-carboxylate as brown oil.

Step 6. cis-Benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(hydroxymethyl)pyrrolidine-1-carboxylate

To a solution of cis-benzyl 3-amino-4-(hydroxymethyl)pyrrolidine-1-carboxylate (900 g, 3.60 mol) and triethylamine (728 g, 7.19 mol) in methanol (9 L) was added di-tert-butyl dicarbonate (863 g, 3.95 mol) dropwise with stirring at room temperature. The resulting solution was stirred overnight at room temperature. The resulting mixture was concentrated under vacuum. The crude product was purified by silica gel chromatography (eluting with 2:3 EtOAc/pet. ether) to afford cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(hydroxymethyl)pyrrolidine-1-carboxylate as a white solid.

Step 7. cis-Benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate

To a mixture of cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(hydroxymethyl)pyrrolidine-1-carboxylate (600 g, 1.71 mol) and 2,6-di-tert-butyl-4-methylpyridine (1406 g, 6.85 mol) in DCM (12 L) was added trimethyloxonium tetrafluoroborate (506.5 g, 3.42 mol). The reaction was stirred overnight at room temperature. The resulting solution was concentrated under vacuum. The crude product was purified by silica gel chromatography (eluting with 3:7 EtOAc/pet. ether) to afford cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate as a white solid.


Step 8. Benzyl (3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate

The racemate cis-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate was separated into its enantiomers by SFC (Column: Lux 5 μ m Amylose-1, 5x25 cm, 5 μ m; Mobile Phase A: CO₂: 50%, and B: MeOH: 50%; Flow rate: 160 mL/min). The first eluting isomer (RT = 4.2 min) was collected and concentrated under vacuum to give benzyl (3S,4S)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate as a white solid. MS

(ESI, m/z): 387 $[M+Na]^+$. ^1H NMR ($CDCl_3$, 300 MHz) δ (ppm): 7.31-7.44 (m, 5H), 5.14-5.36 (m, 3H), 4.28 (s, 1H), 3.46-3.70 (m, 4H), 3.32-3.46 (m, 5H), 2.54 (s, 1H), 1.47 (s, 9H).

The second eluting isomer ($RT = 5.7$ min) was collected and concentrated under vacuum to give benzyl (3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate (**Intermediate 48-3**) as a white solid. MS (ESI, m/z): 387 $[M+Na]^+$. ^1H NMR ($CDCl_3$, 300 MHz) δ (ppm): 7.33-7.40 (m, 5H), 5.09-5.40 (m, 3H), 4.28 (s, 1H), 3.46-3.73 (m, 4H), 3.32-3.46 (m, 5H), 2.53-5.55 (m, 1H), 1.46 (s, 9H).

Intermediate 49. tert-Butyl ((3S,4S)-4-(methoxy-d3)pyrrolidin-3-yl)carbamate



Step 1. Benzyl (3S,4S)-3-azido-4-(methoxy-d3)pyrrolidine-1-carboxylate

To a solution of benzyl (3S,4S)-3-azido-4-hydroxypyrrolidine-1-carboxylate (202.7 mg, 0.773 mmol) in anhydrous DMF (5 mL) was added NaH (60% dispersion in mineral oil, 37.1 mg, 0.927 mmol). The reaction was allowed to stir at room temperature for 1 h. Then, iodomethane-d3 (0.058 mL, 0.927 mmol) was added and the reaction was stirred at 22 °C for 6 h. The reaction was diluted with 5 mL of EtOAc and washed with 2 x 3 mL of H₂O. The organic layer was separated, dried over anhydrous Na₂SO₄, filtered, and concentrated to afford crude benzyl (3S,4S)-3-azido-4-(methoxy-d3)pyrrolidine-1-carboxylate as a yellow oil. MS: (ESI, m/z): 280 $[M+H]^+$.

Step 2. Benzyl (3S,4S)-3-amino-4-(methoxy-d3)pyrrolidine-1-carboxylate

A solution of benzyl (3S,4S)-3-azido-4-(methoxy-d3)pyrrolidine-1-carboxylate (215.8 mg, 0.773 mmol) and triphenylphosphine (223 mg, 0.850 mmol) in THF (1.405 mL) and water (0.14 mL) was stirred for 1 h at room temperature. The reaction was then heated at 50 °C for 5 h. After cooling to room temperature, the reaction was concentrated under vacuum. The residue was purified by

prep-HPLC (Column: Waters XBridge Prep C18 OBD 5 μ m, 19x50 mm; Mobile Phase gradient 0% to 35% ACN, 0.1% formic acid over 8 min; Flow rate: 23 mL/min) to afford benzyl (3S,4S)-3-amino-4-(methoxy-d3)pyrrolidine-1-carboxylate as a colorless oil. MS: (ESI, m/z): 254 [M+H]⁺.


Step 3. Benzyl (3S,4S)-3-((tert-butoxycarbonyl)amino)-4-(methoxy-d3)pyrrolidine-1-carboxylate

A solution of benzyl (3S,4S)-3-amino-4-(methoxy-d3)pyrrolidine-1-carboxylate (217 mg, 0.858 mmol), Et₃N (225 μ l, 1.61 mmol), and Boc₂O (0.299 mL, 1.29 mmol) in THF (1.43 mL) and water (1.43 mL) was stirred for 30 min at room temperature. The resulting mixture was extracted with 2 x 30 mL of DCM. The combined organic layers were dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum to afford benzyl (3S,4S)-3-((tert-butoxycarbonyl)amino)-4-(methoxy-d3)pyrrolidine-1-carboxylate as a colorless oil. MS: (ESI, m/z): 354 [M+H]⁺.

Step 4. tert-Butyl ((3S,4S)-4-(methoxy-d3)pyrrolidin-3-yl)carbamate

A mixture of benzyl 3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidine-1-carboxylate (200 mg, 0.57 mmol) and Pd/C (200 mg, 10%) in EtOAc (10 mL) was stirred for 3 h at room temperature under an atmosphere of hydrogen. The solids were filtered out and washed with 3 x 10 mL of EtOAc. The filtrate was concentrated under vacuum to afford tert-butyl N-(4-methoxypyrrolidin-3-yl)carbamate as a colorless oil. MS: (ESI, m/z): 220 [M+H]⁺. ¹H NMR (CDCl₃, 300 MHz) δ (ppm): 8.04-8.26 (m, 1H), 4.41-4.86 (m, 1H), 3.93-4.22 (m, 1H), 3.09-3.92 (m, 3H), 1.46 (s, 9H).

Intermediate 50. Benzyl N-(7-chloro-6,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-yl)carbamate



Step 1. 4-Chloro-3,5-difluoro-2-hydroxybenzaldehyde

A solution of 3-chloro-2,4-difluorophenol (4.00 g, 23.6 mmol) and HMTA (6.60 g, 47.2 mmol) in TFA (60 mL) was stirred for 3h at 100 °C. After cooling to 0 °C, conc. H₂SO₄ (10 mL) and H₂O (50 mL) were added into the mixture at 0 °C. The mixture was stirred for 2h at 20 °C and then diluted with H₂O (40 mL). The pH value of the mixture was adjusted to 7-8 with dibutylamine. The resulting mixture was extracted with ethyl acetate (3 x 100 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1: 5 ethyl acetate/pet. ether) to afford 4-chloro-3,5-difluoro-2-hydroxybenzaldehyde as an off-white solid. GCMS (ESI, m/z): 192,194 [M+H]⁺.

Step 2. 7-Chloro-6,8-difluoro-3-nitro-2H-chromene

2-Nitroethan-1-ol (3.72 mL, 51.9 mmol) was added to a stirring solution of 4-chloro-3,5-difluoro-2-hydroxybenzaldehyde (1.50 g, 7.77 mmol), dibutylamine (0.659 mL, 3.89 mmol) and phthalic anhydride (2.20 g, 14.8 mmol) in toluene (80 mL) via an injection pump with the flowrate of 0.5 mL/h at 125 °C. The resulting solution was refluxed for 16h at 125 °C. After cooling to 20 °C, the resulting mixture was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 ethyl acetate/pet. ether) to give 7-chloro-6,8-difluoro-3-nitro-2H-chromene as a yellow solid. MS (ESI, m/z): 248, 250 [M+H]⁺.

Step 3. 7-Chloro-8-fluoro-3,4-dihydro-2H-1-benzopyran-3-amine

A mixture of 7-chloro-6,8-difluoro-3-nitro-2H-chromene (1.30 g, 5.25 mmol), BH₃ (40 mL, 1M in THF), and NaBH₄ (399 mg, 10.5 mmol) was stirred for 16h at 65 °C. MeOH (80 mL) was added into the mixture slowly at <10 °C. The resulting solution was stirred for 8h at 80 °C. After cooling to 20 °C, the resulting mixture was concentrated under vacuum. The residue was purified via reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (10 mM NH₄HCO₃) and B: ACN (5% to 75% in 20 min)) to afford 7-chloro-8-fluoro-3,4-dihydro-2H-1-benzopyran-3-amine as an off-white solid. MS (ESI, m/z): 220, 222 [M+H]⁺.


Step 4. Benzyl N-(7-chloro-6,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-yl)carbamate

CbzCl (0.500 mL, 3.55 mmol) was added to a mixture of 7-chloro-6,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-amine (650 mg, 2.96 mmol) and potassium carbonate (822 mg, 5.92 mmol) in

ethyl acetate (20 mL) and water (20 mL) at $<10^{\circ}\text{C}$. The resulting solution was then stirred for 1h at 25°C and diluted with water (30 mL). The resulting mixture was extracted with ethyl acetate (3 x 30 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:5 ethyl acetate/pet. ether) to afford benzyl N-(7-chloro-6,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-yl)carbamate as an off-white solid. MS (ESI, m/z): 354, 356 $[\text{M}+\text{H}]^{+}$.

Intermediate 51-1. tert-Butyl N-[(3R,4S)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate

Intermediate 51-2. tert-Butyl N-[(3S,4R)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate



Step 1. Ethyl 4,4-difluoro-3-hydroxybutanoate

NaBH_4 (34.2 g, 902 mmol) was added into a stirring solution of ethyl 4,4-difluoro-3-oxobutanoate (100 g, 602 mmol) in toluene (1 L) at 0°C . The resulting mixture was stirred for 2h at 0°C , and then warmed up to 25°C and stirred for 16h. The reaction was quenched with water (400 mL). The resulting mixture was extracted with ethyl acetate (2 x 1 L). The organic layers were

combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to afford ethyl 4,4-difluoro-3-hydroxybutanoate as a light yellow oil. GCMS (ESI, m/z): 168 [M+H]⁺.

Step 2. Ethyl (2E)-4,4-difluorobut-2-enoate

To a solution of ethyl 4,4-difluoro-3-hydroxybutanoate (40.0 g, 238 mmol) and Et₃N (99.0 mL, 712 mmol) in DCM (300 mL) was added MsCl (28.0 mL, 244 mmol) dropwise at 0 °C. The resulting mixture was warmed to 25 °C and stirred for 16h. The reaction was quenched with water (500 mL). The resulting mixture was extracted with DCM (2 x 500 mL). The organic layers were combined, washed with brine (500 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 ethyl acetate/pet. ether) to afford ethyl (2E)-4,4-difluorobut-2-enoate as a yellow green oil. GCMS (ESI, m/z): 150 [M+H]⁺.

Step 3. trans-Ethyl 1-benzyl-4-(difluoromethyl)pyrrolidine-3-carboxylate

To a stirring solution of ethyl (2E)-4,4-difluorobut-2-enoate (12.0 g, 79.9 mmol) and benzyl(methoxymethyl)[(trimethylsilyl)methyl]amine (30.7 mL, 120 mmol) in DCM (250 mL) was added a solution of TFA (0.59 mL, 5.21 mmol) in DCM (20 mL) dropwise over 2 min at 0 °C. The resulting mixture was warm to 25 °C and was stirred for 14h. The reaction was quenched by the addition of water (500 mL). The resulting mixture was extracted with DCM (2 x 300 mL). The organic layers were combined, washed with sodium bicarbonate solution (sat., 300 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 to 1:5 ethyl acetate/pet. ether). The collected fraction was concentrated under vacuum to afford trans-ethyl 1-benzyl-4-(difluoromethyl)pyrrolidine-3-carboxylate as a yellow oil. MS (ESI, m/z): 284 [M+H]⁺.

Step 4. trans-Ethyl 4-(difluoromethyl)pyrrolidine-3-carboxylate

A mixture of trans-ethyl-1-benzyl-4-(difluoromethyl)pyrrolidine-3-carboxylate (5.50 g, 19.4 mmol) and Pd(OH)₂/C (2.00 g, 10%) in MeOH (30 mL) was stirred for 2h at 25 °C under hydrogen atmosphere (balloon). The solids were filtered out and the filter cake was washed with MeOH (3 x 10 mL). The filtrate was concentrated under reduced pressure to afford trans-ethyl 4-(difluoromethyl)pyrrolidine-3-carboxylate as a white solid. MS (ESI, m/z): 194 [M+H]⁺.

Step 5. trans-Benzyl 3-(difluoromethyl)-4-[ethoxy(hydroxy)methyl]pyrrolidine-1-carboxylate

CbzCl (4.05 g, 23.741 mmol) was added into a stirring solution of trans-ethyl 4-(difluoromethyl)pyrrolidine-3-carboxylate (3.90 g, 19.8 mmol) and K₂CO₃ (8.20 g, 59.3 mmol) in H₂O (20 mL) and ethyl acetate (40 mL) at 0 °C. The resulting mixture was stirred for 1 h at 25 °C. The reaction was quenched by the addition of water (100 mL). The resulting mixture was extracted with DCM (2 x 200 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reverse phase chromatography (column, C18 silica gel; Mobile phase, A: water (containing 0.1% TFA) and B: ACN (0% to 60% in 30 min)) to afford trans-benzyl 3-(difluoromethyl)-4-[ethoxy(hydroxy)methyl]pyrrolidine-1-carboxylate as a yellow oil. MS (ESI, m/z): 328 [M+H]⁺.

Step 6. trans-1-[(Benzyloxy)carbonyl]-4-(difluoromethyl)pyrrolidine-3-carboxylic acid

To a stirring solution of trans-benzyl 3-(difluoromethyl)-4-[ethoxy(hydroxy)methyl]pyrrolidine-1-carboxylate (6.00 g, 18.0 mmol) in THF (80 mL) and MeOH (20 mL) was added a solution of LiOH (4.30 g, 180 mmol) in water (10 mL). The resulting mixture was stirred for 1 h at 25 °C. The solvent was removed under vacuum. pH value of the residue was adjusted to 5-6 with NH₄Cl (sat. aq.). The resulting mixture was extracted with DCM (3 x 50 mL). The organic layers were combined, washed with sodium bicarbonate solution (sat., 50 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reverse phase chromatography (column: C18 silica gel; Mobile phase, A: water (containing 0.1% TFA) and B: ACN (0% to 60% in 30 min)) to afford trans-1-[(benzyloxy)carbonyl]-4-(difluoromethyl)pyrrolidine-3-carboxylic acid as a colorless oil. MS (ESI, m/z): 300 [M+H]⁺.

Step 7. trans-Benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(difluoromethyl)pyrrolidine-1-carboxylate

To a stirring solution of trans-1-[(benzyloxy)carbonyl]-4-(difluoromethyl)pyrrolidine-3-carboxylic acid (4.00 g, 13.1 mmol) in t-BuOH (100 mL) were added Et₃N (2.73 mL, 19.6 mmol) and DPPA (4.33 g, 15.7 mmol) dropwise at 0 °C. After addition, the resulting mixture was stirred for 2 h at 80 °C. The mixture was allowed to cool down to 25 °C and then concentrated under vacuum. The residue was purified by reverse flash chromatography (column, C18 silica gel;

Mobile phase, A: water (containing 0.1% TFA) and B: ACN (0% to 100% in 40 min)) to afford desired product. It was further purified by silica gel chromatography (eluting with 2:5 THF/pet. ether) to afford trans-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(difluoromethyl)pyrrolidine-1-carboxylate as a colorless oil. MS (ESI, m/z): 371 [M+H]⁺.

Step 8. tert-Butyl N-[(3S,4R)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate and tert-butyl N-[(3R,4S)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate


trans-Benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(difluoromethyl)pyrrolidine-1-carboxylate (1.70 g, 4.58 mmol) was separated by SFC (Column: Chiralpak AD-H SFC, 5x25 cm, 5 μm; Mobile Phase, A: CO₂ :70% and B: MeOH (containing 2 mM NH₃-MeOH): 30%; Flow rate: 180 mL/min) to afford the first eluted isomer (RT = 4.23 min) as a white solid, the stereochemistry arbitrarily assigned as tert-butyl N-[(3R,4S)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate; and the second eluted isomer (RT = 6.14 min) as a white solid, the stereochemistry arbitrarily assigned as tert-butyl N-[(3S,4R)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate. MS (ESI, m/z): 371 [M+H]⁺.

Step 9. tert-Butyl N-[(3S,4R)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate and tert-Butyl N-[(3R,4S)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate

A mixture of benzyl (3S,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(difluoromethyl)pyrrolidine-1-carboxylate (650 mg, 1.76 mmol) and Pd/C (450 mg, 10%) in ethyl acetate (14 mL) was stirred for 3 h at 25 °C under an atmosphere of hydrogen (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give tert-butyl N-[(3S,4R)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate as a colorless oil. MS (ESI, m/z): 237 [M+H]⁺.

tert-Butyl N-[(3R,4S)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate was similarly obtained from benzyl (3R,4S)-3-[[[(tert-butoxy)carbonyl]amino]-4-(difluoromethyl)pyrrolidine-1-carboxylate. MS (ESI, m/z): 237 [M+H]⁺.

Intermediate 52-1. tert-Butyl 4-[(6S)-6-amino-4-fluoro-5,6,7,8-tetrahydroquinolin-2-yl]piperazine-1-carboxylate



Step 1. Benzyl N-benzyl-N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

NaH (260 mg, 6.50 mmol, 60%) was added into a stirring solution of benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) (1.00 g, 2.23 mmol) in DMF (5 mL) in portions at 0 °C. The resulting mixture was stirred for 30 min at 0 °C. To the above mixture was added benzyl bromide (795 mg, 4.56 mmol) dropwise over 30 min at 0 °C. The resulting mixture was stirred at 26 °C for 1 h. The reaction was quenched by the addition of water (30 mL) at 0 °C. The resulting mixture was extracted with ethyl acetate (3 x 30 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reversed phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH_4HCO_3) and B: ACN (5% to 80% in 30 min)). The collected fraction was concentrated to give benzyl N-benzyl-N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a brown solid. MS (ESI, m/z): 521 $[\text{M}+\text{H}]^+$.

Step 2. Benzyl N-benzyl-N-[(6S)-2-chloro-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

A solution of benzyl N-benzyl-N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (925 mg, 1.69 mmol) and a solution of HCl (1 mL, 4M in dioxane) in dioxane (3 mL) was stirred for 15 min at 100 °C. The mixture was allowed to cool to 25 °C and concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH_4HCO_3) and B: ACN (5% to 80% in 30 min)). The collected fraction was concentrated to give benzyl N-benzyl-N-[(6S)-

2-chloro-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a brown oil. MS (ESI, m/z): 407, 409 [M+H]⁺.

Step 3. (6S)-6-[Benzyl[(benzyloxy)carbonyl]amino]-2-chloro-5,6,7,8-tetrahydroquinolin-1-ium-1-olate

A solution of benzyl N-benzyl-N-[(6S)-2-chloro-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (632 mg, 1.49 mmol) and m-CPBA (1.31 g, 7.46 mmol) in DCM (20 mL) was stirred for 16 h at 25 °C. The reaction was quenched by the addition of Na₂SO₃ (sat. aq., 30 mL) and NaHCO₃ (sat. aq., 30 mL). The resulting mixture was extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to afford (6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-2-chloro-5,6,7,8-tetrahydroquinolin-1-ium-1-olate as a white solid. MS (ESI, m/z): 423, 425 [M+H]⁺.

Step 4. (6S)-6-[Benzyl[(benzyloxy)carbonyl]amino]-2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6,7,8-tetrahydroquinolin-1-ium-1-olate

A solution of (6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-2-chloro-5,6,7,8-tetrahydroquinolin-1-ium-1-olate (470 mg, 1.06 mmol), tert-butyl piperazine-1-carboxylate (1.04 g, 5.58 mmol) and DIEA (0.990 mL, 5.57 mmol) in t-BuOH (5 mL) was stirred for 10 h at 90 °C. The resulting mixture was concentrated under vacuum. The residue was purified by reverse phase chromatography (C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (5% to 80% in 30 min)). The collected fraction was concentrated to give (6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6,7,8-tetrahydroquinolin-1-ium-1-olate as a yellow solid. MS (ESI, m/z): 573 [M+H]⁺.

Step 5. TFA salt of (6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-N,N,N-trimethyl-5,6,7,8-tetrahydroquinolin-4-aminium

A solution of (6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6,7,8-tetrahydroquinolin-1-ium-1-olate (460 mg, 0.760 mmol) in DCM (9 mL) was treated with a solution of trimethylamine (3.63 mL, 1M in THF) at 0 °C, followed by the addition of Tf₂O (0.365 mL, 2.13 mmol) dropwise at 0 °C. The resulting mixture was stirred for 1 h at 25 °C and then concentrated under vacuum. The residue was purified by reverse phase chromatography

(Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH_4HCO_3) and B: ACN (5% to 80% in 30 min)). The collected fraction was concentrated to afford the TFA salt of (6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-N,N,N-trimethyl-5,6,7,8-tetrahydroquinolin-4-aminium as a light brown solid. MS (ESI, m/z): 615 $[\text{M}+\text{H}]^+$.

Step 6. tert-Butyl 4-[(6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-4-fluoro-5,6,7,8-tetrahydroquinolin-2-yl]piperazine-1-carboxylate


A solution of the TFA salt of (6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-N,N,N-trimethyl-5,6,7,8-tetrahydroquinolin-4-aminium (230 mg, 0.300 mmol) and a solution of TBAF (3.30 mL, 1M in THF) in DMF (5 mL) was stirred for 2 h at 90 °C. The mixture was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH_4HCO_3) and B: ACN (5% to 80% in 30 min)). The collected fraction was concentrated to give tert-butyl 4-[(6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-4-fluoro-5,6,7,8-tetrahydroquinolin-2-yl]piperazine-1-carboxylate as a white solid. MS (ESI, m/z): 575 $[\text{M}+\text{H}]^+$.

Step 7. tert-Butyl 4-[(6S)-6-amino-4-fluoro-5,6,7,8-tetrahydroquinolin-2-yl]piperazine-1-carboxylate


A mixture of tert-butyl 4-[(6S)-6-[benzyl[(benzyloxy)carbonyl]amino]-4-fluoro-5,6,7,8-tetrahydroquinolin-2-yl]piperazine-1-carboxylate (100 mg, 0.170 mmol) and palladium on carbon (100 mg, 10%) in ethyl acetate (5 mL) was stirred for 3 h at 25 °C under hydrogen atmosphere. The solids were filtered out and the filter cake was washed with ethyl acetate (3 x 10 mL). The filtrate was concentrated under reduced pressure to afford tert-butyl 4-[(6S)-6-amino-4-fluoro-5,6,7,8-tetrahydroquinolin-2-yl]piperazine-1-carboxylate as a light brown solid. MS (ESI, m/z): 351 $[\text{M}+\text{H}]^+$.

[00302] The following intermediate in Table 8 was prepared using standard chemical manipulations and procedures similar to those used for the preparation of Intermediate 52-1.

Table 8

Intermediate Number	Structure and Compound Name	LRMS m/z [M+H] ⁺
52-2	 tert-butyl 3-[(6S)-6-amino-4-fluoro-5,6,7,8-tetrahydroquinolin-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate	377

Intermediate 53. trans-tert-Butyl N-[4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate



Step 1. Ethyl (2E)-4-methoxybut-2-enoate

[00303] A mixture of ethyl (2E)-4-bromobut-2-enoate (33.0 g, 171 mmol), Ag₂O (39.6 g, 171 mmol) and MeOH (100 mL) was stirred for 12 h at 25 °C without light. The solids were filtered out. The filtrate was concentrated under vacuum. The residue was purified via reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B:

ACN (5% to 75% in 20 min)). The collected fraction was concentrated under vacuum to give ethyl (2E)-4-methoxybut-2-enoate as yellow oil. MS (ESI, m/z): 145 $[M+H]^+$.

Step 2. trans-Ethyl 1-benzyl-4-(methoxymethyl)pyrrolidine-3-carboxylate

[00304] A solution of TFA (0.480 mL, 4.20 mmol) in DCM (2 mL) was added to a stirring solution of ethyl (2E)-4-methoxybut-2-enoate (9.30 g, 64.5 mmol) and benzyl(methoxymethyl)[(trimethylsilyl)methyl]amine (22.9 g, 96.7 mmol) in DCM (400 mL) at $<10\text{ }^{\circ}\text{C}$. The resulting mixture was stirred for 2 h at $25\text{ }^{\circ}\text{C}$. The reaction was quenched by the addition of NH_4HCO_3 (sat. aq., 150 mL). The resulting mixture was extracted with DCM (3 x 100 mL). The organic layers were combined, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with 1:5 ethyl acetate/pet. ether) to afford trans-ethyl 1-benzyl-4-(methoxymethyl)pyrrolidine-3-carboxylate as a yellow oil. MS (ESI, m/z): 278 $[M+H]^+$.

Step 3. trans-Ethyl 4-(methoxymethyl)pyrrolidine-3-carboxylate

[00305] A mixture of trans-ethyl 1-benzyl-4-(methoxymethyl)pyrrolidine-3-carboxylate (14.0 g, 50.4 mmol), $\text{Pd}(\text{OH})_2/\text{C}$ (8.43 g, 10%) and MeOH (200 mL) was stirred for 1 h at $25\text{ }^{\circ}\text{C}$ under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give trans-ethyl 4-(methoxymethyl)pyrrolidine-3-carboxylate as a yellow oil. MS (ESI, m/z): 188 $[M+H]^+$.

Step 4. trans-1-Benzyl 3-ethyl 4-(methoxymethyl)pyrrolidine-1,3-dicarboxylate

[00306] CbzCl (9.01 mL, 64.0 mmol) was added into a mixture of trans-ethyl 4-(methoxymethyl)pyrrolidine-3-carboxylate (8.00 g, 42.7 mmol) and K_2CO_3 (14.7 g, 106 mmol) in ethyl acetate (100 mL) and H_2O (100 mL) at $0\text{ }^{\circ}\text{C}$. The mixture was stirred for 1 h at $25\text{ }^{\circ}\text{C}$. The mixture was extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to afford trans-1-benzyl 3-ethyl 4-(methoxymethyl)pyrrolidine-1,3-dicarboxylate as a yellow oil. MS (ESI, m/z): 322 $[M+H]^+$.

Step 5. trans-1-[(Benzyloxy)carbonyl]-4-(methoxymethyl)pyrrolidine-3-carboxylic acid

[00307] A solution of trans-1-benzyl 3-ethyl 4-(methoxymethyl)pyrrolidine-1,3-dicarboxylate (5.00 g, 15.6 mmol) and NaOH (3.11 g, 77.7 mmol) in H₂O (40 mL), THF (40 mL) and MeOH (40 mL) was stirred for 2 h at 25 °C. The solvent was removed under vacuum. The pH value of the residue was adjusted to pH 6 with hydrochloric acid (3N). The resulting mixture was extracted with ethyl acetate (3 x 25 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give trans-1-[(benzyloxy)carbonyl]-4-(methoxymethyl)pyrrolidine-3-carboxylic acid as yellow oil. MS (ESI, *m/z*): 294 [M+H]⁺.

Step 6. trans-Benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate

[00308] A solution of trans-1-[(benzyloxy)carbonyl]-4-(methoxymethyl)pyrrolidine-3-carboxylic acid (3.00 g, 10.2 mmol), DPPA (3.38 g, 12.3 mmol), and Et₃N (4.25 mL, 30.6 mmol) in t-BuOH (60 mL) was stirred for 3h at 80 °C. The mixture was cooled to 25 °C and concentrated under vacuum. The residue was purified via reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (5% to 75% in 20 min)). The collected fraction was concentrated under vacuum to give a crude product. The crude product was purified by silica gel chromatography (eluting with 1:3 THF/ pet. ether) to afford trans-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidine-1-carboxylate as a colorless oil. MS (ESI, *m/z*): 365 [M+H]⁺.


Step 7. trans-tert-Butyl N-[1-benzyl-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate

[00309] NaH (59.0 mg, 1.48 mmol, 60%) was added to a solution of trans-tert-butyl N-[1-benzyl-4-(methoxymethyl)pyrrolidin-3-yl]carbamate (350 mg, 0.983 mmol) in DMF (4 mL) at 0 °C. The resulting solution was stirred for 30 min at 0 °C. Then CH₃I (0.018 mL, 0.281 mmol) was added at 0 °C. The resulting mixture was stirred for additional 2 h at 25 °C. The reaction was quenched with water (50 mL). The resulting mixture was extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to afford trans-tert-butyl N-[1-benzyl-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate as a white solid. MS (ESI, *m/z*): 379 [M+H]⁺.

Step 8. trans-tert-Butyl N-[4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate

[00310] A mixture of trans-tert-butyl N-[1-benzyl-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate (330 mg, 0.828 mmol) and palladium on carbon (300 mg, 10%) in ethyl acetate (10 mL) was stirred for 3 h at 25 °C under hydrogen atmosphere (balloon). The resulting mixture was filtered. The filtrate was concentrated under reduced pressure to afford trans-tert-butyl N-[4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate as a colorless oil. MS (ESI, m/z): 245 $[M+H]^+$.

Intermediate 54. cis-tert-Butyl N-[4-(fluoromethyl)pyrrolidin-3-yl]carbamate



Step 1. 1-Benzyl-4-(hydroxymethyl)pyrrolidin-2-one

[00311] NaBH₄ (24.0 g, 634 mmol) was added to a solution of methyl 1-benzyl-5-oxopyrrolidine-3-carboxylate (50.0 g, 214 mmol) in ethanol (800 mL) at 0 °C. The resulting solution was stirred for 4 h at 25 °C. The reaction was quenched with water (50 mL). The mixture was concentrated under vacuum. The residue was diluted with water (200 mL). The resulting mixture was extracted with DCM (3 x 500 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 10:1 dichloromethane/methanol) to give 1-benzyl-4-(hydroxymethyl)pyrrolidin-2-one as colorless oil. MS (ESI, m/z): 206 $[M+H]^+$.

Step 2. 1-Benzyl-4-(fluoromethyl)pyrrolidin-2-one

[00312] DAST (32.2 mL, 244 mmol) was added to a solution of 1-benzyl-4-(hydroxymethyl)pyrrolidin-2-one (20.0 g, 97.5 mmol) in DCM (300 mL) at -78 °C. The resulting solution was stirred for 16 h at 25 °C. The reaction was poured into ice-water (200 mL). The resulting mixture was extracted with DCM (3 x 200 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified via reverse phase chromatography (Column: C18 column; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (0% to 50% in 45 min)) to give 1-benzyl-4-(fluoromethyl)pyrrolidin-2-one as yellow oil. MS (ESI, m/z): 208 [M+H]⁺.

Step 3. trans-1-Benzyl-4-(fluoromethyl)-3-hydroxypyrrolidin-2-one

[00313] A solution of LDA (24.0 mL, 2M in THF) was added to a solution of 1-benzyl-4-(fluoromethyl)pyrrolidin-2-one (5.00 g, 23.6 mmol) in THF (100 mL) at -78 °C. The resulting solution was stirred for 1 h at -78 °C. Then O₂ was introduced in. The resulting solution was stirred for 3 h at -78 °C. The reaction was quenched by the addition of water (50 mL). The solvent was removed under vacuum. The residue was extracted with DCM (3 x 100 mL). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.1% TFA) and B: ACN (0% to 60% in 30 min)) to afford trans-1-benzyl-4-(fluoromethyl)-3-hydroxypyrrolidin-2-one as a brown oil. MS (ESI, m/z): 224 [M+H]⁺.

Step 4. trans-1-Benzyl-4-(fluoromethyl)-2-oxopyrrolidin-3-yl methanesulfonate

[00314] MsCl (2.75 mL, 35.5 mmol) was added to a stirring solution of trans-1-benzyl-4-(fluoromethyl)-3-hydroxypyrrolidin-2-one (5.40 g, 23.7 mmol) and Et₃N (6.59 mL, 47.4 mmol) in DCM (70 mL) at 0 °C. The resulting mixture was stirred for 2 h at 25 °C. The reaction was quenched by the addition of water (50 mL) at 0 °C. The resulting mixture was extracted with DCM (3 x 50 mL). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered concentrated under reduced pressure to afford trans-1-benzyl-4-(fluoromethyl)-2-oxopyrrolidin-3-yl methanesulfonate as a brown oil. MS (ESI, m/z): 302 [M+H]⁺.

Step 5. cis-3-Azido-1-benzyl-4-(fluoromethyl)pyrrolidin-2-one

[00315] A mixture of trans-1-benzyl-4-(fluoromethyl)-2-oxopyrrolidin-3-yl methanesulfonate (6.00 g, 19.5 mmol) and NaN₃ (3.81 g, 58.6 mmol) in DMF (150 mL) was stirred for 1 h at 100

°C. The mixture was allowed to cool down to 25 °C. The reaction was quenched by the addition of water (500 mL) at 25 °C. The resulting mixture was extracted with ethyl acetate (3 x 200 mL). The combined organic layers were washed with brine (300 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford cis-3-azido-1-benzyl-4-(fluoromethyl)pyrrolidin-2-one as a yellow oil. MS (ESI, m/z): 249 [M+H]⁺.

Step 6. cis-tert-Butyl N-[1-benzyl-4-(fluoromethyl)-2-oxopyrrolidin-3-yl]carbamate


[00316] A mixture of 3-azido-1-benzyl-4-(fluoromethyl)pyrrolidin-2-one (4.00 g, 15.8 mmol), Palladium on carbon (4.00 g, 10%) and di-tert-butyl dicarbonate (6.89 g, 31.6 mmol) in EtOH (100 mL) was stirred for 3 h at 25 °C under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford cis-tert-butyl N-[1-benzyl-4-(fluoromethyl)-2-oxopyrrolidin-3-yl]carbamate as a white solid. MS (ESI, m/z): 323 [M+H]⁺.

Step 7. cis-tert-Butyl N-[1-benzyl-4-(fluoromethyl)pyrrolidin-3-yl]carbamate

[00317] A solution of BH₃ (30 mL, 1M in THF) was added to a stirring solution of cis-tert-butyl N-[1-benzyl-4-(fluoromethyl)-2-oxopyrrolidin-3-yl]carbamate (2.40 g, 7.30 mmol) in THF (60 mL) at 0 °C. The resulting mixture was stirred for 16 h at 25 °C. The resulting mixture was concentrated under vacuum. Then EtOH (60 mL), H₂O (15 mL) and Et₃N (15 mL) were added. The resulting mixture was stirred for 2 h at 80 °C. The resulting mixture was cooled to 25 °C and concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 Column; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (0% to 100% in 30 min)) to afford cis-tert-butyl N-[1-benzyl-4-(fluoromethyl)pyrrolidin-3-yl]carbamate as a white solid. MS (ESI, m/z): 309 [M+H]⁺.

Step 8. cis-tert-Butyl N-[4-(fluoromethyl)pyrrolidin-3-yl]carbamate

[00318] A mixture of tert-butyl N-[1-benzyl-4-(fluoromethyl)pyrrolidin-3-yl]carbamate (450 mg, 1.43 mmol) and Pd(OH)₂/C (450 mg, 10%) in ethyl acetate (10 mL) was stirred for 2 h at 25 °C under hydrogen atmosphere (balloon). The resulting mixture filtered. The filtrate was concentrated under reduced pressure to afford cis-tert-butyl N-[4-(fluoromethyl)pyrrolidin-3-yl]carbamate as a brown oil. MS (ESI, m/z): 219 [M+H]⁺.

Intermediate 55. trans-tert-Butyl N-[4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate**Step 1. trans-Benzyl 3-azido-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate**

[00319] NaH (1.91 g, 47.7 mmol, 60%) was added into a stirring solution of trans-benzyl 3-azido-4-hydroxypyrrolidine-1-carboxylate (5.00 g, 19.1 mmol) in DMF (70 mL) at $<10^{\circ}\text{C}$. The resulting solution was stirred for 30 min at $<10^{\circ}\text{C}$. Then 1-bromo-2-methoxyethane (7.95 g, 57.2 mmol) was added to the mixture at $<10^{\circ}\text{C}$. The resulting mixture was stirred for 3 h at 26°C . The reaction was then quenched by the addition of water/ice (70 mL). The resulting mixture was extracted with ethyl acetate (3 x 100 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and the filtrate concentrated under vacuum. The residue was purified via reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (10 % to 75% in 25 min)). The collected fraction was concentrated under vacuum to give trans-benzyl 3-azido-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate as yellow oil. MS (ESI, m/z): 321 $[\text{M}+\text{H}]^{+}$.

Step 2. trans-Benzyl 3-amino-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate

[00320] A mixture of trans-benzyl 3-azido-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate (5.60 g, 17.4 mmol) and PtO_2 (2.00 g, 8.80 mmol) in ethyl acetate (100 mL) was stirred for 1 h at 26°C under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give trans-benzyl 3-amino-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate as a black oil. MS (ESI, m/z): 295 $[\text{M}+\text{H}]^{+}$.

Step 3. trans-Benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate


[00321] $(\text{Boc})_2\text{O}$ (5.45 g, 25.0 mmol) was added into a solution of trans-benzyl 3-amino-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate (5.00 g, 17.0 mmol) and Et_3N (7.08 mL, 70.0 mmol) in THF (70 mL) and H_2O (70 mL). The resulting solution was stirred for 3 h at 26°C . THF was removed under vacuum. The residue was extracted with ethyl acetate (3 x 100 mL). The organic

layers were combined, dried over anhydrous sodium sulfate, filtered and the filtrate concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to afford trans-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate as yellow oil. MS (ESI, m/z): 395 [M+H]⁺.

Step 4. trans-tert-Butyl N-[4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate

[00322] A mixture of trans-benzyl 3-[[[(tert-butoxy)carbonyl]amino]-4-(2-methoxyethoxy)pyrrolidine-1-carboxylate (1.60 g, 4.06 mmol) and palladium on carbon (1.60 g, 10%) in ethyl acetate (30 mL) was stirred for 2 h at 26 °C under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give trans-tert-butyl N-[4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate as light yellow oil. MS (ESI, m/z): 261 [M+H]⁺.

Intermediate 56. tert-Butyl N-[(3S,4S)-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate



Step 1. Benzyl (3S,4S)-3-azido-4-(propan-2-yloxy)pyrrolidine-1-carboxylate

[00323] Sodium hydride (688 mg, 17.2 mmol, 60%) was added to a stirring solution of benzyl (3S,4S)-3-azido-4-hydroxypyrrolidine-1-carboxylate (3.00 g, 11.44 mmol), 2-bromopropane (10.7 mL, 113 mmol) and KI (3.80 g, 2.89 mmol), in DMF (100 mL) at 0 °C. The resulting solution was stirred for 3 h at 50 °C. After cooling to 25 °C, the reaction was quenched by the addition of H₂O/ice (100 mL). The resulting mixture was extracted with ethyl acetate (3 x 200 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and then concentrated under vacuum. The residue was purified by reversed phase chromatography (Column, C18 silica gel; Mobile phase, A: water (containing 0.05% formic acid) and B: ACN (0% to 80% within 40 min)). The collected fraction was concentrated under vacuum to give benzyl (3S,4S)-3-azido-4-(propan-2-yloxy)pyrrolidine-1-carboxylate as colorless oil. MS (ESI, m/z): 305 [M+H]⁺.

Step 2. Benzyl (3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidine-1-carboxylate

[00324] A mixture of benzyl (3S,4S)-3-azido-4-(propan-2-yloxy)pyrrolidine-1-carboxylate (220 mg, 0.72 mmol) and PtO₂ (33.0 mg, 0.150 mmol) in ethyl acetate (10 mL) was stirred for 2 h

at 25 °C under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give benzyl (3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidine-1-carboxylate as brown oil. MS (ESI, m/z): 279 [M+H]⁺.


Step 3. Benzyl (3S,4S)-3-[[[(tert-butoxy)carbonyl]amino]-4-(propan-2-yloxy)pyrrolidine-1-carboxylate

[00325] A solution of benzyl (3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidine-1-carboxylate (200 mg, 0.720 mmol), Et₃N (0.190 mL, 1.35 mmol) and (Boc)₂O (236 mg, 1.08 mmol) in THF (5 mL) and water (5 mL) was stirred for 30 min at 25 °C. The solvent was removed under vacuum. The residue was diluted with water (10 mL). The resulting mixture was extracted with ethyl acetate (3 x 20 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by a silica gel chromatography (eluting with 1:5 ethyl acetate/pet. ether) to give benzyl (3S, 4S)-3-[[[(tert-butoxy)carbonyl]amino]-4-(propan-2-yloxy)pyrrolidine-1-carboxylate as colorless oil. MS (ESI, m/z): 379 [M+H]⁺.

Step 4. tert-Butyl N-[(3S,4S)-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate

[00326] A mixture of benzyl (3S,4S)-3-[[[(tert-butoxy)carbonyl]amino]-4-(propan-2-yloxy)pyrrolidine-1-carboxylate (250 mg, 0.660 mmol) and Palladium on carbon (250 mg, 10%) in ethyl acetate (5 mL) was stirred for 2 h at 25°C under hydrogen atmosphere (balloon). The solids were filtered out. The filtered was concentrated under vacuum to give tert-butyl N-[(3S,4S)-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate as a brown solid. MS (ESI, m/z): 245 [M+H]⁺.

Intermediate 57. Benzyl N-(2-chloro-5,6,7,8-tetrahydroquinazolin-6-yl)carbamate



Step 1. Benzyl N-[(3Z)-3-[(dimethylamino)methylidene]-4-oxocyclohexyl]carbamate

[00327] A solution of benzyl N-(4-oxocyclohexyl)carbamate (10.0 g, 40.4 mmol) and (dimethoxymethyl)dimethylamine (4.80 g, 40.4 mmol) in toluene (20 mL) was stirred for 16 h at 100 °C. The resulting mixture was concentrated under vacuum to give benzyl N-[(3Z)-3-[(dimethylamino)methylidene]-4-oxocyclohexyl]carbamate as yellow oil. MS (ESI, m/z): 303 [M+H]⁺.


Step 2. Benzyl N-(2-amino-5,6,7,8-tetrahydroquinazolin-6-yl)carbamate

[00328] A solution of guanidine hydrochloride (3.45 g, 36.1 mmol) and sodium ethoxide (2.47 g, 36.3 mmol) in ethanol (150 mL) was stirred for 30 min at 23°C. Benzyl N-[(3Z)-3-[(dimethylamino)methylidene]-4-oxocyclohexyl]carbamate (11 g, 21.8 mmol, purity: 60%) was then added. The resulting solution was stirred for overnight at 80°C. The resulting mixture was concentrated under vacuum. The residue was purified by a silica gel chromatography (eluting with 1:10 MeOH/DCM) to give benzyl N-(2-amino-5,6,7,8-tetrahydroquinazolin-6-yl)carbamate as a white solid. S (ESI, m/z): 299 [M+H]⁺.

Step 3. Benzyl N-(2-chloro-5,6,7,8-tetrahydroquinazolin-6-yl)carbamate

[00329] A mixture of benzyl N-(2-amino-5,6,7,8-tetrahydroquinazolin-6-yl)carbamate (1.85 g, 6.20 mmol), t-BuNO₂ (3.72 mL, 31.0 mmol) and CuCl₂ (4.14 g, 30.8 mmol) in ACN (40 mL) was stirred for 20 min at 60 °C. The reaction mixture was cooled to 25 °C and diethyl ether (20 mL) was added. The solids were filtered out. The filtrate was concentrated under vacuum. The residue was purified by reversed phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% NH₄HCO₃) and B: ACN (5% to 80% in 30 min)) to afford benzyl N-(2-chloro-5,6,7,8-tetrahydroquinazolin-6-yl)carbamate as a white solid. MS (ESI, m/z): 318, 320 [M+H]⁺.

Intermediate 58. tert-Butyl 4-(7-amino-5,6,7,8-tetrahydroisoquinolin-3-yl)piperazine-1-carboxylate



Step 1. RuCl-(S,S)-Ts-DPEN(hexamethylbenzene)

[00330] A solution of N-[(1S,2S)-2-amino-1,2-diphenylethyl]-4-methylbenzene-1-sulfonamide (S,S-Ts-DPEN) (1.09 g, 3.00 mmol), Et₃N (0.90 mL, 6.47 mmol), and [RuCl₂(hexamethylbenzene)]₂ (1.00 g, 1.50 mmol) in DCM (40 mL) was stirred for 5 h at 25 °C. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 MeOH/DCM) to afford RuCl-(S,S)-Ts-DPEN(hexamethylbenzene) as a red solid.

Step 2. RuCl-(S,S)-Ts-DPEN-OTf(hexamethylbenzene)

[00331] A solution of RuCl-(S,S)-Ts-DPEN(hexamethylbenzene) (1.95 g, 2.99 mmol) and silver trifluoromethanesulfonate (0.800 g, 3.17 mmol) in DCM (50 mL) was stirred for 6 h at 25 °C. The solids were filtered out. The filtrate was concentrated under vacuum to afford RuCl-(S,S)-Ts-DPEN-OTf(hexamethylbenzene) as a brown solid.

Step 3. Ethyl 6-chloro-4-methylpyridine-3-carboxylate

[00332] Sulfuric acid (15.6 mL, 306 mmol) was added into a solution of 6-chloro-4-methylpyridine-3-carboxylic acid (10.0 g, 58.3 mmol) in EtOH (250 mL). The resulting solution was stirred for 48 h at 25 °C. A saturated solution of NaHCO₃ in water (500 mL) was added to the

reaction. Then EtOH was evaporated out. The residue was extracted with DCM (2 x 500 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:15 ethyl acetate/pet. ether) to give ethyl 6-chloro-4-methylpyridine-3-carboxylate as a white solid. MS (ESI, m/z): 200, 202 [M+H]⁺.

Step 4. Methyl 3-chloro-8-hydroxy-5,6-dihydroisoquinoline-7-carboxylate

[00333] A solution of LDA (12.5 mL, 2M in THF) was added to a solution of ethyl 6-chloro-4-methylpyridine-3-carboxylate (2.00 g, 10.0 mmol) in THF (100 mL) at -78 °C. The resulting solution was stirred for 1 h at -78 °C. Then methyl prop-2-enoate (2.25 mL, 24.9 mmol) was added and the reaction was stirred for 1 h at -78 °C. The reaction was then quenched with water (50 mL). THF was removed in vacuo. The aqueous layer was extracted with DCM (3 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10 ethyl acetate/pet. ether) to give methyl 3-chloro-8-hydroxy-5,6-dihydroisoquinoline-7-carboxylate as a white solid. MS (ESI, m/z): 240, 242 [M+H]⁺.

Step 5. Methyl 3-chloro-8-hydroxy-5,6,7,8-tetrahydroisoquinoline-7-carboxylate

[00334] Et₃N (7.62 mL, 54.8 mmol) was added to a solution of formic acid (1.89 mL, 50.1 mmol) in MeOH (90 mL) at 0 °C. Then methyl 3-chloro-8-hydroxy-5,6-dihydroisoquinoline-7-carboxylate (3.00 g, 12.5 mmol) and RuCl-(S,S)-Ts-DPEN-OTf(hexamethylbenzene) (82 mg, 0.099 mmol) was added. The resulting solution was stirred for 16 h at 25 °C. The reaction was then poured into water (200 mL). The solvent was removed in vacuo. The aqueous layer was extracted with DCM (3 x 150 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:5 ethyl acetate/pet. ether) to give methyl 3-chloro-8-hydroxy-5,6,7,8-tetrahydroisoquinoline-7-carboxylate as a white solid. MS (ESI, m/z): 242, 244 [M+H]⁺.

Step 6. Methyl 3-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6-dihydroisoquinoline-7-carboxylate

[00335] A mixture of methyl 3-chloro-8-hydroxy-5,6,7,8-tetrahydroisoquinoline-7-carboxylate (0.500 g, 2.07 mmol), tert-butyl piperazine-1-carboxylate (0.750 g, 4.03 mmol), RuPhos Pd G3

(100 mg, 0.120 mmol), RuPhos (62.0 mg, 0.140 mmol) and Cs_2CO_3 (2.00 g, 6.16 mmol) in toluene (10 mL) was stirred for 16 h at 95 °C. Four batches of this reaction was set up in parallel. After cooling to room temperature, the four batches of reaction mixture were poured into water (100 mL). The resulting mixture was extracted with acyl acetate (3 x 100 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:4 ethyl acetate/pet. ether) to give methyl 3-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6-dihydroisoquinoline-7-carboxylate as a white solid). MS (ESI, m/z): 374 $[\text{M}+\text{H}]^+$.

Step 7. Methyl 3-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6-dihydroisoquinoline-7-carboxylate

[00336] A mixture of methyl 3-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6-dihydroisoquinoline-7-carboxylate (550 mg, 1.47 mmol) and Palladium on carbon (500 mg, 10%) in ethyl acetate (20 mL) was stirred for 3 h at 25 °C under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give methyl 3-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6,7,8-tetrahydroisoquinoline-7-carboxylate as a white solid. MS (ESI, m/z): 376 $[\text{M}+\text{H}]^+$.

Step 8. 3-[4-[(tert-Butoxy)carbonyl]piperazin-1-yl]-5,6,7,8-tetrahydroisoquinoline-7-carboxylic acid

[00337] A solution of LiOH (200 mg, 8.35 mmol) in water (20 mL) was added to a solution of methyl 3-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6,7,8-tetrahydroisoquinoline-7-carboxylate (500 mg, 1.33 mmol) in THF (20 mL). The resulting solution was stirred for 2 h at 25 °C. The resulting solution was concentrated under vacuum. The residue was purified via reverse phase chromatography (Column: X Bridge C18, 19 x 150 mm, 5 μm ; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (0% to 30% in 25 min)). The collected fraction was concentrated under vacuum to give 3-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6,7,8-tetrahydroisoquinoline-7-carboxylic acid as a white solid. MS (ESI, m/z): 362 $[\text{M}+\text{H}]^+$.


Step 9. tert-Butyl 4-(7-[[[(benzyloxy)carbonyl]amino]-5,6,7,8-tetrahydroisoquinolin-3-yl]piperazine-1-carboxylate

[00338] A solution of 3-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-5,6,7,8-tetrahydroisoquinoline-7-carboxylic acid (400 mg, 1.11 mmol), DPPA (476 mL, 2.20 mmol) and Et₃N (0.460 mL, 3.72 mmol) in toluene (6 mL) was stirred for 2 h at 25 °C and then 2 h at 70 °C. Then benzyl alcohol (1.14 mL, 11.0 mmol) was added. The resulting solution was stirred for 2 h at 70 °C. After cooling to room temperature, the reaction was then poured into water (30 mL). The resulting mixture was extracted with acyl acetate (3 x 20 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to give tert-butyl 4-(7-[[[(benzyloxy)carbonyl]amino]-5,6,7,8-tetrahydroisoquinolin-3-yl]piperazine-1-carboxylate as colorless oil. MS (ESI, m/z): 467 [M+H]⁺.

Step 10. tert-Butyl 4-(7-amino-5,6,7,8-tetrahydroisoquinolin-3-yl)piperazine-1-carboxylate

[00339] A mixture of tert-butyl 4-(7-[[[(benzyloxy)carbonyl]amino]-5,6,7,8-tetrahydroisoquinolin-3-yl]piperazine-1-carboxylate (200 mg, 0.430 mmol) and K₃PO₄ (228 mg, 1.07 mmol) in DMA (5 mL) was stirred for 1 h at 90 °C. After cooling to room temperature, the residue was purified via reverse phase chromatography (Column: X Bridge C18, 19 x 150 mm, 5 um; Mobile Phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (10% to 50% in 30 min)). The collected fraction was concentrated under vacuum to give tert-butyl 4-(7-amino-5,6,7,8-tetrahydroisoquinolin-3-yl)piperazine-1-carboxylate as a white solid. MS (ESI, m/z): 333 [M+H]⁺.

Intermediate 59. tert-Butyl N-[1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate



Step 1. 7-Benzyl 9-(2-hydroxyethyl) 1,4-dioxo-7-azaspiro[4.4]nonane-7,9-dicarboxylate

[00340] A mixture of 1-benzyl 3-ethyl 4-oxopyrrolidine-1,3-dicarboxylate (5.00 g, 17.2 mmol), ethane-1,2-diol (3.20 g, 51.5 mmol), and PTSA (296 mg, 1.72 mmol) in toluene (100 mL) was stirred for 3 h at 110 °C with a Dean-Stark tube. The mixture was cooled to 27 °C and concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (5% to 55% in 20 min)) to afford 7-benzyl 9-(2-hydroxyethyl) 1,4-dioxo-7-azaspiro[4.4]nonane-7,9-dicarboxylate as yellow oil. MS (ESI, m/z): 352 $[M+H]^+$.

Step 2. 7-[(Benzyloxy)carbonyl]-1,4-dioxo-7-azaspiro[4.4]nonane-9-carboxylic acid

[00341] A mixture of 7-benzyl 9-(2-hydroxyethyl) 1,4-dioxo-7-azaspiro[4.4]nonane-7,9-dicarboxylate (2.80 g, 7.96 mmol) and sodium hydroxide (199 mg, 4.98 mmol) in MeOH (40 mL), THF (40 mL) and H₂O (40 mL) was stirred for 2 h at 28 °C. THF and MeOH were removed under vacuum. The resulting mixture was acidified to pH 7-8 with HCl (1M), and then concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (5% to 30% in 20 min)) to afford 7-[(benzyloxy)carbonyl]-1,4-dioxo-7-azaspiro[4.4]nonane-9-carboxylic acid as yellow oil. MS (ESI, m/z): 308 $[M+H]^+$.

Step 3. Benzyl 9-[[tert-butoxy]carbonyl]amino]-1,4-dioxo-7-azaspiro[4.4]nonane-7-carboxylate

[00342] DPPA (632 mg, 2.30 mmol) was added to a stirring solution of 7-[(benzyloxy)carbonyl]-1,4-dioxo-7-azaspiro[4.4]nonane-9-carboxylic acid (600 mg, 1.91 mmol) and Et₃N (0.398 mL, 2.87 mmol) in t-BuOH (10 mL) at 0 °C. The resulting mixture was stirred for 2 h at 90 °C. The mixture was allowed to cool. The resulting mixture was concentrated under vacuum. The mixture was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: H₂O (containing 10 mmol/L NH₄HCO₃) and B: ACN (0% to 60% over 30 min)) to afford benzyl 9-[[tert-butoxy]carbonyl]amino]-1,4-dioxo-7-azaspiro[4.4]nonane-7-carboxylate as a yellow oil. MS (ESI, m/z): 379 $[M+H]^+$.


Step 4. tert-Butyl N-[1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate

[00343] A mixture of benzyl 9-[[[(tert-butoxy)carbonyl]amino]-1,4-dioxo-7-azaspiro[4.4]nonane-7-carboxylate (200 mg, 0.518 mmol) and Pd/C (200 mg, 10%) in EtOAc (5 mL) was stirred for 2 h at 25 °C under hydrogen atmosphere. The resulting mixture was filtered, and the filter cake was washed with ethyl acetate (3 x 5 mL). The filtrate was concentrated under reduced pressure to afford tert-butyl N-[1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate as a brown. MS (ESI, m/z): 245 $[M+H]^+$.

SYNTHETIC EXAMPLES OF COMPOUNDS OF FORMULA (I):

Example 1-1. (R)-3-Amino-6-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide and

Example 1-2. (S)-3-Amino-6-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl 4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate

[00344] A solution of tert-butyl 4-(3-aminochroman-7-yl)piperazine-1-carboxylate, **Intermediate 12**, (130 mg, 0.27 mmol), EDCI (81 mg, 0.42 mmol), HOBt (57 mg, 0.42 mmol),

DIEA (108 mg, 0.84 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (58 mg, 0.28 mmol) in DMF (5 mL) was stirred for 18 h at room temperature. The reaction was quenched with 20 mL of water and was extracted with EtOAc (50 mL x 3). The combined organic layers were washed with brine (30 mL x 2), dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. Purification by prep-HPLC (Column: XBridge Prep C18 OBD, 19x50 mm, 5 µm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 40% B to 60% B in 13 min) afforded tert-butyl 4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate.

Step 2. tert-Butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate and tert-Butyl (S)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate

[00345] The racemic mixture of tert-butyl 4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate was separated by chiral HPLC (Column: Lux cellulose-4, 0.46x5 cm, 3 µm; Mobile phase: hexanes (0.1% Et₂NH):EtOH = 60:40 in 6 min) to afford the title compounds as follows: tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate as a yellow solid (first eluting isomer, RT = 3.87 min, stereochemistry assumed) and tert-butyl (S)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate as a yellow solid (second eluting isomer, RT = 4.74 min, stereochemistry assumed).

Step 3. (R)-3-Amino-6-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide

[00346] A solution of (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate (6 mg, 0.01 mmol) and TFA (0.5 mL) in CH₂Cl₂ (2 mL) was stirred for 1 h at room temperature. The resulting mixture was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep C18 OBD, 19x50 mm, 5 µm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 5% B to 30% B in 13 min) to give (R)-3-amino-6-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide as a yellow solid. ¹H NMR (CD₃OD, 400 MHz) δ(ppm): 8.18 (d, *J* = 8.0 Hz, 1H), 7.28 (d, *J* = 8.4 Hz, 1H), 6.95 (d, *J* = 8.4 Hz, 1H), 6.54-6.56 (m, 1H), 6.41 (d, *J* = 2.4 Hz, 1H), 4.40-



4.47 (m, 1H), 4.21-4.24 (m, 1H), 3.88-3.93 (m, 1H), 3.07-3.12 (m, 4H), 2.95-3.02 (m, 5H), 2.82-2.88 (m, 1H), 2.62 (s, 3H). MS: (ESI, m/z): 424 [M+H]⁺.






Step 4. (S)-3-Amino-6-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide





[00347] A procedure similar to Step 3 was applied to (S)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperazine-1-carboxylate (8 mg, 0.02 mmol) to afford (S)-3-amino-6-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide as a white solid. ¹H NMR (CD₃OD, 300 MHz) δ(ppm): 8.18 (d, *J* = 8.4 Hz, 1H), 7.28 (d, *J* = 8.7 Hz, 1H), 6.96 (d, *J* = 8.7 Hz, 1H), 6.54-6.58 (m, 1H), 6.41 (d, *J* = 2.1 Hz, 1H), 4.39-4.86 (m, 1H), 4.22-4.25 (m, 1H), 3.88-3.96 (m, 1H), 3.04-3.10 (m, 5H), 2.89-3.02 (m, 4H), 2.81-2.87 (m, 1H), 2.63 (s, 3H). MS: (ESI, m/z): 424 [M+H]⁺.






[00348] The following examples in Table 9 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Examples 1-1 and 1-2.





Table 9:






Example Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺	¹ H NMR
1-3 ¹	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	450	(CD ₃ OD, 400 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.0 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 6.91 (d, <i>J</i> = 8.4 Hz, 1H), 6.44 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.27 (s, 1H), 4.45-4.40 (m, 1H), 4.23-4.20 (m, 1H), 3.92-3.78 (m, 1H), 3.58-3.56 (m, 2H), 3.41-3.38 (m, 2H), 2.99-2.94 (m, 1H), 2.87-2.80 (m, 3H), 2.63 (s, 3H), 1.89-1.87 (m, 4H).
1-4 ¹	 N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	450	(CD ₃ OD, 400 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.4 Hz, 1H), 7.30 (d, <i>J</i> = 8.0 Hz, 1H), 6.91 (d, <i>J</i> = 8.4 Hz, 1H), 6.44 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.28 (s, 1H), 4.45-4.40 (m, 1H), 4.23-4.20 (m, 1H), 3.92-3.78 (m, 1H), 3.58-3.56 (m, 2H), 3.41-3.38 (m, 2H), 2.99-2.94 (m, 1H), 2.87-2.80 (m, 3H), 2.63 (s, 3H), 1.89-1.87 (m, 4H).





Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
1-5 ²	 N-((3R)-7-(3,9-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	464	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.0 Hz, 1H), 7.51 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.21 (s, 2H), 6.91 (d, <i>J</i> = 8.4 Hz, 1H), 6.46 (d, <i>J</i> = 11.2 Hz, 1H), 6.24 (d, <i>J</i> = 2.4 Hz, 1H), 4.26-4.31 (m, 1H), 4.13-4.16 (m, 1H), 3.77-3.82 (m, 1H), 3.52-3.56 (m, 2H), 3.07 (s, 2H), 2.81-2.90 (m, 4H), 2.58 (s, 3H), 2.50-2.51 (m, 1H), 2.18-2.24 (m, 1H), 1.70-1.82 (m, 4H), 1.46-1.49 (m, 1H).
1-6 ²	 N-((3S)-7-(3,9-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	464	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.51 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.21 (br s, 2H), 6.91 (d, <i>J</i> = 8.4 Hz, 1H), 6.46 (dd, <i>J</i> = 2.0, 8.4 Hz, 1H), 6.24 (s, 1H), 4.31-4.26 (m, 1H), 4.16-4.13 (m, 1H), 3.80 (t, <i>J</i> = 10.0 Hz, 3H), 3.56-3.52 (m, 2H), 3.08-3.01 (m, 2H), 2.92-2.81 (m, 4H), 2.58 (s, 3H), 2.24-2.18 (m, 1H), 1.82-1.72 (m, 4H), 1.49-1.46 (m, 1H).
1-7 ³	 3-amino-N-((R)-7-((2R,3S)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.1 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 7.03 (d, <i>J</i> = 8.4 Hz, 1H), 6.72 (dd, <i>J</i> = 2.1, 8.4 Hz, 1H), 6.57 (s, 1H), 4.47-4.44 (m, 1H), 4.28-4.23 (m, 1H), 3.95-3.89 (m, 1H), 3.07-2.81 (m, 8H), 2.63 (s, 3H), 1.27 (d, <i>J</i> = 6.6 Hz, 3H), 0.97 (d, <i>J</i> = 6.3 Hz, 3H).
1-8 ³	 3-amino-N-((S)-7-((2R,3S)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.1 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 7.03 (d, <i>J</i> = 8.4 Hz, 1H), 6.72 (dd, <i>J</i> = 2.1, 8.4 Hz, 1H), 6.57 (s, 1H), 4.47-4.44 (m, 1H), 4.28-4.23 (m, 1H), 3.95-3.89 (m, 1H), 3.07-2.81 (m, 8H), 2.63 (s, 3H), 1.27 (d, <i>J</i> = 6.6 Hz, 3H), 0.97 (d, <i>J</i> = 6.3 Hz, 3H).
1-9 ³	 3-amino-N-((R)-7-((2R,3S)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ(ppm): 8.18 (d, <i>J</i> = 8.1 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 7.03 (d, <i>J</i> = 8.4 Hz, 1H), 6.67 (dd, <i>J</i> = 2.1, 8.1 Hz, 1H), 6.57 (s, 1H), 4.49-4.41 (m, 1H), 4.27-4.23 (m, 1H), 3.97-3.90 (m, 1H), 3.01-2.63 (m, 8H),





Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	3-amino-N-((R)-7-((2S,3R)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide		2.60 (s, 3H), 1.25 (d, $J = 6.6$ Hz, 3H), 0.96 (d, $J = 6.3$ Hz, 3H).
1-10 ³	 3-amino-N-((S)-7-((2S,3R)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ (ppm): 8.20 (d, $J = 8.4$ Hz, 1H), 7.30 (d, $J = 8.4$ Hz, 1H), 7.03 (d, $J = 8.1$ Hz, 1H), 6.67 (dd, $J = 2.4, 8.4$ Hz, 1H), 6.57 (s, 1H), 4.49-4.42 (m, 1H), 4.27-4.23 (m, 1H), 3.95-3.89 (m, 1H), 3.07-2.73 (m, 7H), 3.95-3.89 (m, 1H), 3.09-2.67 (m, 8H), 2.63 (s, 3H), 1.24 (d, $J = 6.6$ Hz, 3H), 0.97 (d, $J = 6.3$ Hz, 3H).
1-11 ³	 3-amino-N-((R)-7-((2R,3R)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ (ppm): 8.19 (d, $J = 8.1$ Hz, 1H), 7.30 (d, $J = 8.4$ Hz, 1H), 6.97 (d, $J = 8.4$ Hz, 1H), 6.54 (dd, $J = 2.4, 8.4$ Hz, 1H), 6.38 (s, 1H), 4.46-4.43 (m, 1H), 4.26-4.22 (m, 1H), 3.94-3.87 (m, 2H), 3.23-2.81 (m, 7H), 2.63 (s, 3H), 1.16 (d, $J = 6.9$ Hz, 3H), 0.93 (d, $J = 6.6$ Hz, 3H).
1-12 ³	 3-amino-N-((S)-7-((2R,3R)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ (ppm): 8.20 (d, $J = 8.4$ Hz, 1H), 7.30 (d, $J = 8.1$ Hz, 1H), 6.97 (d, $J = 8.4$ Hz, 1H), 6.53 (dd, $J = 2.4, 8.4$ Hz, 1H), 6.39 (s, 1H), 4.46-4.43 (m, 1H), 4.26-4.21 (m, 1H), 3.96-3.89 (m, 2H), 3.22-2.82 (m, 7H), 2.63 (s, 3H), 1.21 (d, $J = 6.3$ Hz, 3H), 0.96 (d, $J = 6.6$ Hz, 3H).
1-13 ³	 3-amino-N-((R)-7-((2S,3S)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ (ppm): 8.19 (d, $J = 8.1$ Hz, 1H), 7.30 (d, $J = 8.4$ Hz, 1H), 6.97 (d, $J = 8.4$ Hz, 1H), 6.53 (dd, $J = 2.4, 8.4$ Hz, 1H), 6.38 (s, 1H), 4.45-4.43 (m, 1H), 4.25-4.21 (m, 1H), 3.96-3.85 (m, 2H), 3.20-2.87 (m, 7H), 2.63 (s, 3H), 1.15 (d, $J = 6.9$ Hz, 3H), 0.93 (d, $J = 6.9$ Hz, 3H).





Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
1-14 ³	 3-amino-N-((S)-7-((2S,3S)-2,3-dimethylpiperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ(ppm): 8.19 (d, J = 8.4Hz, 1H), 7.30 (d, J = 8.4Hz, 1H), 6.97 (d, J = 8.4Hz, 1H), 6.53 (dd, J = 2.4, 8.4Hz, 1H), 6.38 (s, 1H), 4.45-4.42 (m, 1H), 4.26-4.22 (m, 1H), 3.94-3.85 (m, 2H), 3.20-2.81 (m, 7H), 2.63 (s, 3H), 1.15 (d, J = 6.9Hz, 3H), 0.92 (d, J = 6.9Hz, 3H).
1-15 ⁴	 (R)-3-amino-5-fluoro-6-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.29 (d, J = 10.5Hz, 1H), 7.60 (br s, 1H), 7.17 (br s, 2H), 6.90 (d, J = 8.7Hz, 1H), 6.48 (dd, J = 2.1, 8.7Hz, 1H), 6.27 (s, 1H), 4.30-4.21 (m, 1H), 4.16-4.11 (m, 1H), 3.78 (t, J = 9.9Hz, 1H), 2.97-2.94 (m, 4H), 2.88-2.78 (m, 6H), 2.55 (s, 3H), 2.47 (br s, 1H).
1-16 ⁴	 (S)-3-amino-5-fluoro-6-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.29 (d, J = 10.5Hz, 1H), 7.60 (br s, 1H), 7.17 (br s, 2H), 6.90 (d, J = 8.7Hz, 1H), 6.48 (dd, J = 2.4, 8.4Hz, 1H), 6.27 (s, 1H), 4.30-4.21 (m, 1H), 4.16-4.11 (m, 1H), 3.78 (t, J = 10.2Hz, 1H), 2.97-2.94 (m, 4H), 2.88-2.78 (m, 6H), 2.55 (s, 3H), 2.46 (br s, 1H).
1-17 ⁵	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-3-amino-5-fluorothieno[2,3-b]pyridine-2-carboxamide	454	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.70 (d, J = 2.1Hz, 1H), 8.42 (dd, J = 3.0, 9.6Hz, 1H), 7.68 (d, J = 7.5Hz, 1H), 7.22 (br s, 2H), 6.89 (d, J = 8.7Hz, 1H), 6.39 (d, J = 6.0Hz, 1H), 6.18 (s, 1H), 4.32-4.22 (m, 1H), 4.16-4.13 (m, 1H), 3.79 (t, J = 9.6Hz, 1H), 3.53-3.51 (m, 2H), 2.86-2.83 (m, 2H), 2.73-2.67 (m, 2H), 2.50-2.49 (m, 3H), 1.68-1.66 (m, 4H).
1-18 ⁵		454	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.70 (d, J = 2.1Hz, 1H), 8.42 (dd, J = 3.0, 9.6Hz, 1H), 7.68 (d, J = 7.2Hz, 1H), 7.22 (br s, 2H), 6.88 (d, J = 8.4Hz, 1H), 6.38 (d, J = 8.4Hz, 1H), 6.17 (s, 1H), 4.32-4.22 (m, 1H), 4.16-4.13 (m,




Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-3-amino-5-fluorothieno[2,3-b]pyridine-2-carboxamide		1H), 3.79 (t, J = 9.6Hz, 1H), 3.53-3.51 (m, 2H), 2.86-2.83 (m, 2H), 2.73-2.67 (m, 2H), 2.50-2.49 (m, 3H), 1.68-1.66 (m, 4H).
1-19 ⁶	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	451	(CD ₃ OD, 300 MHz) δ (ppm): 8.58 (s, 1H), 6.95 (d, J = 8.4 Hz, 1H), 6.49-6.45 (m, 1H), 6.31 (s, 1H), 4.50-4.41 (m, 1H), 4.25-4.22 (m, 1H), 3.96-3.89 (m, 1H), 3.74-3.72 (m, 2H), 3.54-3.45 (m, 2H), 3.08-2.82 (m, 4H), 2.73 (s, 3H), 1.96-1.86 (m, 4H).
1-20 ⁶	 N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	451	(CD ₃ OD, 300 MHz) δ (ppm): 8.58 (s, 1H), 6.94 (d, J = 8.4 Hz, 1H), 6.48-6.45 (m, 1H), 6.31 (s, 1H), 4.50-4.41 (m, 1H), 4.25-4.22 (m, 1H), 3.96-3.89 (m, 1H), 3.74-3.72 (m, 2H), 3.54-3.45 (m, 2H), 3.08-2.82 (m, 4H), 2.73 (s, 3H), 1.96-1.86 (m, 4H).
1-21 ⁷	 (R)-7-amino-3-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyrazine-6-carboxamide	425	(DMSO- d_6 , 400 MHz) δ (ppm): 8.64 (s, 1H), 7.79 (br s, 1H), 6.98 (br s, 2H), 6.92 (d, J = 8.4Hz, 1H), 6.50 (dd, J = 2.4, 8.8Hz, 1H), 6.29 (s, 1H), 4.33-4.26 (m, 1H), 4.18-4.14 (m, 1H), 3.84-3.79 (m, 1H), 2.98-2.92 (m, 4H), 2.88-2.83 (m, 2H), 2.81-2.79 (m, 4H), 2.65 (s, 3H), 2.39-2.32 (m, 1H).
1-22 ⁷	 (S)-7-amino-3-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyrazine-6-carboxamide	425	(DMSO- d_6 , 400 MHz) δ (ppm): 8.64 (s, 1H), 7.79 (br s, 1H), 6.98 (br s, 2H), 6.92 (d, J = 8.8Hz, 1H), 6.51 (dd, J = 2.4, 8.4Hz, 1H), 6.29 (s, 1H), 4.34-4.25 (m, 1H), 4.18-4.15 (m, 1H), 3.84-3.79 (m, 1H), 2.98-2.92 (m, 4H), 2.88-2.83 (m, 2H), 2.81-2.79 (m, 4H), 2.65 (s, 3H), 2.39-2.32 (m, 1H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
1-23 ⁸	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide	456	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 7.43 (d, <i>J</i> = 7.6Hz, 1H), 7.14 (br s, 2H), 6.87 (d, <i>J</i> = 8.8Hz, 1H), 6.37 (dd, <i>J</i> = 2.4, 8.8Hz, 1H), 6.16 (s, 1H), 4.28-4.20 (m, 1H), 4.19-4.10 (m, 1H), 3.78-3.73 (m, 1H), 3.50-3.48 (m, 2H), 3.33-3.30 (m, 3H), 2.86-2.68 (m, 5H), 2.51-2.50 (m, 2H), 1.69-1.67 (m, 4H).
1-24 ⁸	 N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide	456	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 7.43 (d, <i>J</i> = 7.6Hz, 1H), 7.14 (br s, 2H), 6.88 (d, <i>J</i> = 8.4Hz, 1H), 6.37 (dd, <i>J</i> = 2.0, 8.4Hz, 1H), 6.17 (s, 1H), 4.28-4.21 (m, 1H), 4.19-4.10 (m, 1H), 3.78-3.73 (m, 1H), 3.50-3.48 (m, 2H), 3.33-3.30 (m, 3H), 2.86-2.68 (m, 5H), 2.51-2.50 (m, 2H), 1.69-1.67 (m, 4H).
1-25 ⁹	 (R)-6-amino-2-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-d]thiazole-5-carboxamide	430	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 7.45 (br s, 1H), 7.15 (br s, 2H), 6.92 (d, <i>J</i> = 8.7Hz, 1H), 6.50 (dd, <i>J</i> = 2.4, 8.4Hz, 1H), 6.30 (s, 1H), 4.28-4.22 (m, 1H), 4.16-4.12 (m, 1H), 3.78 (t, <i>J</i> = 9.9Hz, 1H), 3.52-3.49 (m, 1H), 3.02-2.97 (m, 4H), 2.84-2.80 (m, 9H).
1-26 ⁹	 (S)-6-amino-2-methyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-d]thiazole-5-carboxamide	430	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 7.45 (br s, 1H), 7.15 (br s, 2H), 6.92 (d, <i>J</i> = 8.7Hz, 1H), 6.50 (dd, <i>J</i> = 2.4, 8.7Hz, 1H), 6.30 (s, 1H), 4.28-4.22 (m, 1H), 4.16-4.12 (m, 1H), 3.78 (t, <i>J</i> = 9.9Hz, 1H), 3.52-3.49 (m, 1H), 3.02-2.97 (m, 4H), 2.85-2.80 (m, 9H).
1-27 ¹⁰		432	(CD ₃ OD, 300 MHz) δ(ppm): 8.69 (s, 1H), 8.39 (s, 1H), 7.49 (d, <i>J</i> = 3.6Hz, 1H), 6.91 (d, <i>J</i> = 8.4Hz, 1H), 6.56 (d, <i>J</i> = 3.6Hz, 1H), 6.42 (dd, <i>J</i> = 2.4, 8.4Hz, 1H), 6.27 (s, 1H), 4.48-4.21 (m, 4H), 3.97-3.93 (m, 1H), 3.57-3.42

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-1-ethyl-1H-pyrrolo[2,3-b]pyridine-5-carboxamide		(m, 2H), 3.36-3.31 (m, 2H), 3.05-2.98 (m, 1H), 2.82-2.78 (m, 3H), 1.84-1.82 (m, 4H), 1.42 (t, $J = 7.2$ Hz, 3H).
1-28 ¹⁰	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-1-ethyl-1H-pyrrolo[2,3-b]pyridine-5-carboxamide	432	(CD ₃ OD, 300 MHz) δ (ppm): 8.69 (s, 1H), 8.39 (s, 1H), 7.49 (d, $J = 3.6$ Hz, 1H), 6.91 (d, $J = 8.4$ Hz, 1H), 6.56 (d, $J = 3.3$ Hz, 1H), 6.43 (dd, $J = 2.4$, 8.4Hz, 1H), 6.28 (s, 1H), 4.48-4.21 (m, 4H), 3.97-3.93 (m, 1H), 3.57-3.42 (m, 2H), 3.36-3.31 (m, 2H), 3.05-2.98 (m, 1H), 2.82-2.78 (m, 3H), 1.84-1.82 (m, 4H), 1.42 (t, $J = 7.2$ Hz, 3H).
1-29 ¹¹	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	468	(CD ₃ OD, 300 MHz) δ (ppm): 8.20 (d, $J = 8.4$ Hz, 1H), 7.30 (d, $J = 8.1$ Hz, 1H), 6.24-6.19 (m, 1H), 6.11 (s, 1H), 4.43-4.40 (m, 1H), 4.22-4.19 (m, 1H), 3.94-3.87 (m, 1H), 3.58-3.53 (m, 2H), 3.32-3.29 (m, 2H), 3.11-2.94 (m, 1H), 2.85-2.81 (m, 2H), 2.76-2.70 (m, 1H), 2.63 (s, 3H), 1.89-1.81 (m, 4H).
1-30 ¹¹	 N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	468	(CD ₃ OD, 300 MHz) δ (ppm): 8.20 (d, $J = 8.4$ Hz, 1H), 7.30 (d, $J = 8.4$ Hz, 1H), 6.24-6.19 (m, 1H), 6.12 (s, 1H), 4.43-4.40 (m, 1H), 4.22-4.19 (m, 1H), 3.94-3.87 (m, 1H), 3.58-3.53 (m, 2H), 3.32-3.29 (m, 2H), 3.11-2.94 (m, 1H), 2.85-2.81 (m, 2H), 2.76-2.71 (m, 1H), 2.63 (s, 3H), 1.91-1.83 (m, 4H).
1-31 ¹²	 (R)-3-amino-6-methyl-N-(7-(piperazin-1-yl)-6-(trifluoromethoxy)chroman-3-yl)-thieno[2,3-b]pyridine-2-carboxamide	508	(DMSO- <i>d</i> ₆ , 400 MHz) δ (ppm): 8.32 (d, $J = 8.4$ Hz, 1H), 7.60 (br s, 1H), 7.49 (d, $J = 8.4$ Hz, 1H), 7.23 (br s, 2H), 7.03 (s, 1H), 6.49 (s, 1H), 4.40-4.20 (m, 1H), 4.19-4.17 (m, 1H), 3.89-3.84 (m, 1H), 2.92-2.79 (m, 10H), 2.67 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	yl)thieno[2,3-b]pyridine-2-carboxamide		
1-32 ¹²	 (S)-3-amino-6-methyl-N-(7-(piperazin-1-yl)-6-(trifluoromethoxy)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	508	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.1 Hz, 1H), 7.59 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.23 (br s, 2H), 7.04 (s, 1H), 6.50 (s, 1H), 4.30-4.20 (m, 1H), 4.19-4.16 (m, 1H), 3.90-3.83 (m, 1H), 2.92-2.79 (m, 10H), 2.58 (s, 3H).
1-33 ¹³	 (R)-3-amino-N-(8-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.21 (d, <i>J</i> = 9 MHz, 1H), 7.31 (d, <i>J</i> = 9 MHz, 1H), 6.82 (d, <i>J</i> = 9 MHz, 1H), 6.62-6.56 (m, 1H), 4.51-4.42 (m, 1H), 4.36-4.30 (m, 1H), 4.31-3.93 (m, 1H), 3.04 (s, 11H), 2.63 (s, 3H).
1-34 ¹³	 (S)-3-amino-N-(8-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.21 (d, <i>J</i> = 9 MHz, 1H), 7.31 (d, <i>J</i> = 9 MHz, 1H), 6.82 (d, <i>J</i> = 9 MHz, 1H), 6.62-6.56 (m, 1H), 4.51-4.42 (m, 1H), 4.36-4.30 (m, 1H), 4.31-3.93 (m, 1H), 3.04 (s, 11H), 2.63 (s, 3H).
1-35 ¹⁴	 (R)-3-amino-N-(6-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(CD ₃ OD, 300 MHz) δ(ppm): 8.19 (d, <i>J</i> = 8.4 Hz, 1H), 7.30 (d, <i>J</i> = 8.1 Hz, 1H), 6.79 (d, <i>J</i> = 12.6 Hz, 1H), 6.47 (d, <i>J</i> = 7.5 Hz, 1H), 4.45-4.42 (m, 1H), 4.25-4.20 (m, 1H), 3.95-3.88 (m, 1H), 3.06-2.81 (m, 11H), 2.63 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
1-36 ¹⁴	 (S)-3-amino-N-(6-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.1 Hz, 1H), 7.30 (d, <i>J</i> = 8.1 Hz, 1H), 6.83 (d, <i>J</i> = 12.6 Hz, 1H), 6.51 (d, <i>J</i> = 7.5 Hz, 1H), 4.45-4.43 (m, 1H), 4.25-4.20 (m, 1H), 3.96-3.90 (m, 1H), 3.25-2.92 (m, 11H), 2.63 (s, 3H).
1-37 ¹⁵	 (R)-6-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-2-methylthieno[2,3-d]thiazole-5-carboxamide	448	(CD ₃ OD, 300 MHz) δ(ppm): 6.33 (dd, <i>J</i> = 2.4, 12.6 Hz, 1H), 6.24 (s, 1H), 4.40-4.20 (m, 1H), 4.43-4.36 (m, 1H), 4.25-4.21 (m, 1H), 3.93-3.86 (m, 1H), 3.12-3.10 (m, 4H), 3.03-2.96 (m, 5H), 2.81 (s, 3H), 2.75-2.66 (m, 1H).
1-38 ¹⁵	 (S)-6-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-2-methylthieno[2,3-d]thiazole-5-carboxamide	448	(CD ₃ OD, 300 MHz) δ(ppm): 6.34 (dd, <i>J</i> = 2.4, 12.6 Hz, 1H), 6.24 (s, 1H), 4.40-4.20 (m, 1H), 4.43-4.36 (m, 1H), 4.25-4.21 (m, 1H), 3.93-3.86 (m, 1H), 3.12-3.10 (m, 4H), 3.03-2.96 (m, 5H), 2.81 (s, 3H), 2.75-2.66 (m, 1H).
1-39 ¹⁶	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluorochroman-3-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide	474	(CD ₃ OD, 300 MHz) δ(ppm): 6.22 (dd, <i>J</i> = 2.4, 12.9 Hz, 1H), 6.12 (s, 1H), 4.40-4.37 (m, 1H), 4.23-4.18 (m, 1H), 3.91-3.85 (m, 1H), 3.64-3.62 (m, 2H), 3.42-3.31 (m, 2H), 3.11-2.65 (m, 7H), 1.87-1.85 (m, 4H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
1-40 ¹⁶	 N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluorochroman-3-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide	474	(CD ₃ OD, 300 MHz) δ(ppm): 6.21 (dd, <i>J</i> = 2.4, 12.9 Hz, 1H), 6.11 (s, 1H), 4.40-4.37 (m, 1H), 4.23-4.18 (m, 1H), 3.91-3.85 (m, 1H), 3.64-3.62 (m, 2H), 3.32-3.31 (m, 2H), 3.11-2.65 (m, 7H), 1.87-1.85 (m, 4H).
1-41 ¹⁷	 N-((3S)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)-5-fluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	454	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.7 Hz, 1H), 7.56 (br s, 1H), 7.31 (d, <i>J</i> = 8.1 Hz, 1H), 7.23 (br s, 2H), 6.13 (dd, <i>J</i> = 2.1, 12.9 Hz, 1H), 5.96 (s, 1H), 4.34-4.25 (m, 1H), 4.19-4.15 (m, 1H), 3.87-3.81 (m, 1H), 3.67-3.64 (m, 2H), 3.41-3.32 (m, 4H), 2.90-2.73 (m, 2H), 2.70 (s, 3H), 1.69 (br s, 1H), 1.50-1.45 (m, 1H).
1-42 ¹⁷	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)-5-fluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	454	(CD ₃ OD, 400 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.0 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 6.13 (dd, <i>J</i> = 2.4, 12.8 Hz, 1H), 6.06 (s, 1H), 4.45-4.40 (m, 1H), 4.25-4.21 (m, 1H), 3.95-3.87 (m, 3H), 3.55-3.49 (m, 4H), 3.01-2.96 (m, 1H), 2.77-2.70 (m, 2H), 2.65 (s, 3H), 1.68-1.67 (m, 1H).

¹ **Notes on procedures:** In Step 1, the amine **Intermediate 13** was used, and purification was by silica gel chromatography (eluting with 2:3 EtOAc/pet. ether). Step 2 chiral HPLC (Column: Chiralpak ID-2, 2x25 cm, 5 μm; Mobile phase: 40% EtOH/hexanes for 41 min) afforded Example 1-4 precursor as the first eluting isomer (RT=3.32 min) and Example 1-3 precursor as the second eluting isomer (RT=4.42 min).

² **Notes on procedures:** In Step 1, the amine **Intermediate 14-1** was used, and purification was by silica gel chromatography (eluting with 1:1 EtOAc/pet. ether). Step 2 chiral HPLC (Column: Chiralpak IB, 4.6x250 mm, 5 μm; Mobile phase A: hexanes (0.1% Et₂NH), B: EtOH; Gradient: 50% B to 50% B in 30 min) afforded Example 1-5 precursor as the first eluting isomer (RT=3.7 min) and Example 1-6 precursor as the second eluting isomer (RT=7.1 min). Step 3 purification by prep-HPLC (Column: XBridge Prep Phenyl OBD, 19x150 mm, 5 μm; Mobile phase A: Water (0.05% NH₄OH), B: ACN; Gradient: 25% B to 60% B in 7 min).

³ **Notes on procedures:** In Step 1, the amine **Intermediate 14-2** was used. Step 2 chiral separation was not performed. Step 3 prep-HPLC isolated Diastereomer Mixture 1 (containing Examples 1-7 thru 1-10,

stereochemistry assumed) and Diastereomer Mixture 2 (containing Examples 1-11 thru 1-14, stereochemistry assumed). Further purification of Mixture 1 by chiral HPLC (Column: Chiralpak ID-2, 2x25 cm, 5 μ m; Mobile phase: 20% IPA/MTBE (0.1% Et₂NH) for 23 min) provided Mixture 1a (RT=8.9 min, containing Examples 1-7 and 1-8, stereochemistry assumed) and Mixture 1b (RT=9.5 min, containing Examples 1-9 and 1-10, stereochemistry assumed). Separation of Mixture 1a (Column: Chiralpak IE, 2x25 cm, 5 μ m; Mobile phase: 50% EtOH/hexanes for 25 min; Flow rate: 15 mL/min) provided Example 1-7 (RT=19.1 min, stereochemistry assumed) and Example 1-8 (RT=23.0 min, stereochemistry assumed). Separation of Mixture 1b (Column: Chiralpak ID-2, 2x25 cm, 5 μ m; Mobile phase: 20% IPA/MTBE (0.1% Et₂NH) for 23 min) provided Example 1-9 (RT=14.6 min, stereochemistry assumed) and Example 1-10 (RT=20.1 min, stereochemistry assumed). Further purification of Mixture 2 by chiral HPLC (Column: Chiralpak ID-2, 2x25 cm, 5 μ m; Mobile phase: 30% EtOH/MTBE for 26 min; Flow rate: 16 mL/min) provided Mixture 2a (RT=8.4 min, containing Examples 1-11 and 1-12, stereochemistry assumed), Example 1-13 (RT=9.5 min, stereochemistry assumed), and Example 1-14 (RT=10.4 min, stereochemistry assumed). Separation of Mixture 2a (Column: Phenomenex Lux Cellulose-4, AXIA Packed, 2.12x25 cm, 5 μ m; Mobile Phase : 50% EtOH/Hex (0.1% Et₂NH) for 22 min) provided Example 1-11 (RT=16.0 min, stereochemistry assumed) and Example 1-12 (RT=19.9 min, stereochemistry assumed).

⁴ **Notes on procedures:** In Step 1, the carboxylic acid **Intermediate 16** was used. Step 2 chiral HPLC (Column: Chiralpak IA, 2x25 cm, 5 μ m; Mobile phase A: hexanes, B: EtOH; Flow rate: 17 mL/min; Gradient: 40% B to 60% B in 23 min) afforded Example 1-15 precursor as the first eluting isomer (RT=15.9 min) and Example 1-16 precursor as the second eluting isomer (RT=19.7 min).

⁵ **Notes on procedures:** In Step 1, the carboxylic acid **Intermediate 17** and the amine **Intermediate 13** were used. Step 2 chiral HPLC (Column: Phenomenex Lux Cellulose-4, AXIA Packed, 2.12x25 cm, 5 μ m; Mobile Phase : 50% EtOH/hexanes for 15 min) afforded Example 1-17 precursor as the first eluting isomer (RT=9.3 min) and Example 1-18 precursor as the second eluting isomer (RT=11.8 min).

⁶ **Notes on procedures:** In Step 1, the amine **Intermediate 13** was used. Step 2 chiral HPLC (Column: Chiralpak ID-2, 2x25 cm, 5 μ m; Mobile phase: 80% MeOH/CH₂Cl₂ for 15 min; Flow rate: 17 mL/min) afforded Example 1-19 precursor as the first eluting isomer (RT=10.5 min) and Example 1-20 precursor as the second eluting isomer (RT=12.2 min).

⁷ **Notes on procedures:** Step 2 chiral HPLC (Column: Chiral Art Cellulose-SB, 2x25 cm, 5 μ m; Mobile phase: 30% EtOH/Hexanes for 30 min) afforded Example 1-21 precursor as the first eluting isomer (RT=22.2 min) and Example 1-22 precursor as the second eluting isomer (RT=26.5 min).

⁸ **Notes on procedures:** In Step 1, the carboxylic acid **Intermediate 24** and the amine **Intermediate 13** were used. Step 2 chiral HPLC (Column: Chiralpak ID-2, 2x25 cm, 5 μ m; Mobile phase: 50% EtOH/hexanes for 35 min; Flow rate: 14 mL/min) afforded Example 1-23 precursor as the first eluting isomer (RT=21.6 min) and Example 1-24 precursor as the second eluting isomer (RT=29.0 min).

⁹ **Notes on procedures:** In Step 1, the carboxylic acid **Intermediate 24** was used. Step 2 chiral HPLC (Column: Chiralpak IC, 2x25 cm, 5 μ m; Mobile phase: 50% EtOH/hexanes for 25 min) afforded Example 1-25 precursor as the first eluting isomer (RT=15.8 min) and Example 1-26 precursor as the second eluting isomer (RT=20.4 min).

¹⁰ **Notes on procedures:** In Step 1, the carboxylic acid **Intermediate 25** and the amine **Intermediate 13** were used. Step 2 chiral HPLC (Column: Chiralpak ID-3, 2x25 cm, 5 μ m; Mobile phase: 50% iPrOH/hexanes for 10 min) afforded Example 1-27 precursor as the first eluting isomer (RT=4.5 min) and Example 1-28 precursor as the second eluting isomer (RT=7.2 min).

¹¹ **Notes on procedures:** In Step 1, the amine **Intermediate 15-1** was used. Step 2 chiral HPLC (Column: Chiralpak IA-2, 2x25 cm, 5 μ m; Mobile phase: 50% EtOH/hexanes for 25 min; Flow rate: 16 mL/min) afforded Example 1-29 precursor as the first eluting isomer (RT=16.6 min) and Example 1-30 precursor as the second eluting isomer (RT=21.1 min).

¹² **Notes on procedures:** In Step 1, the amine **Intermediate 15-2** was used; the amide coupling condition of HBTU and Et₃N in DMA was used. Step 2 chiral HPLC (Column: Chiralpak IC, 2x25 cm, 5 μ m; Mobile phase: 20% EtOH/MTBE) afforded Example 1-31 precursor as the first eluting isomer (RT=6.9 min) and

Example 1-32 precursor as the second eluting isomer (RT=20.0 min). In Steps 3 and 4, the crude product was treated with NH₃ in MeOH to afford the free base before purification by prep-HPLC.

¹³ **Notes on procedures:** In step 1, the amine **Intermediate 4-1** was used. Step 2 chiral HPLC (Column: Chiralpak IC, 2x25 cm, 5 μm; Mobile phase: 50% EtOH/hexanes for 22 min; Flow rate: 15 mL/min). Step 3 prep-HPLC (Column: XBridge Shield RP18 OBD, 5 μm, 19x150mm; Mobile phase: Water (with 10 mM NH₄HCO₃) and ACN, gradient (25% ACN up to 65% in 7 min)).


¹⁴ **Notes on procedures:** In step 1, the amine **Intermediate 4-2** was used. Step 2 chiral HPLC (Column: Chiralpak IA, 2.12 x 15cm, 5 μm; Mobile Phase A: hexanes, B: EtOH; Gradient: 0% to 50% B in 17 min). Step 3 prep-HPLC (Column: XBridge C18, 19 x 150 mm, 5 μm; Mobile Phase A: Water (10 mM NH₄HCO₃), B: ACN; Gradient: (10% B to 80% B in 10 min)).

¹⁵ **Notes on procedures:** In step 1 the amine **Intermediate 15-3** and the carboxylic acid **Intermediate 24** were used. The enantiomers were separated by chiral HPLC using chiral column Chiral Art Cellulose-SB and mobile phase 10% IPA/MTBE to provide the precursor to Example 1-37 as the first eluting isomer and the precursor to Example 1-38 as the second eluting isomer.

¹⁶ **Notes on procedures:** In step 1, the carboxylic acid **Intermediate 24** was used. The enantiomers were separated by chiral HPLC using chiral column Chiralpak ID-2 and mobile phase 40% EtOH/hexanes to provide the precursor to Example 1-39 as the first eluting isomer and the precursor to Example 1-40 as the second eluting isomer.

¹⁷ **Notes on procedures:** In step 1, the amine **Intermediate 15-3** was used. The enantiomers were separated by chiral HPLC using chiral column Chiralpak IG and mobile phase 50% EtOH/hexanes to provide the precursor to Example 1-41 as the first eluting isomer and the precursor to Example 1-42 as the second eluting isomer.

Example 2-1. 3-Amino-N-((3R)-7-(9,9-difluoro-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl 7-((R)-3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate

[00349] A mixture of benzyl (R)-(7-bromochroman-3-yl)carbamate, **Intermediate 2**, (400 mg, 1.08 mmol), tert-butyl 9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate, **Intermediate 32**, (288 mg, 1.06 mmol), Pd(dppf)Cl₂-CH₂Cl₂ (90 mg, 0.11 mmol), Xphos (105 mg, 0.22 mmol),

and Cs₂CO₃ (1.076 g, 3.30 mmol) in toluene (40 mL) was stirred for 16 h at 100 °C. After cooling to room temperature, the reaction was quenched with water (40 mL) and extracted with EtOAc (40 mL x 3). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:4 EtOAc/pet. ether) to give tert-butyl 7-((R)-3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate as a yellow oil. MS: (ESI, m/z): 544 [M+H]⁺.

Step 2. tert-Butyl 7-((R)-3-aminochroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate

[00350] A mixture of tert-butyl 7-((R)-3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate (110 mg, 0.19 mmol) and Pd/C (20 mg, 10%) in EtOAc (10 mL) was stirred for 2 h at 30 °C under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum to afford tert-butyl 7-((R)-3-aminochroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate as a yellow solid. MS: (ESI, m/z): 410 [M+H]⁺.

Step 3. tert-Butyl 7-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate



[00351] A solution of tert-butyl 7-((R)-3-aminochroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate (70 mg, 0.16 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (43 mg, 0.20 mmol), HOBt (45 mg, 0.33 mmol), EDCI (65 mg, 0.34 mmol), and DIEA (66 mg, 0.51 mmol) in DMF (15 mL) was stirred for 16 h at room temperature. The reaction was quenched by the addition of 25 mL of water. The resulting mixture was extracted with EtOAc (30 mL x 3). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:1 EtOAc/pet. ether) afforded tert-butyl 7-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate as a yellow solid. MS: (ESI, m/z): 600 [M+H]⁺.






Step 4. 3-Amino-N-((3R)-7-(9,9-difluoro-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide





[00352] A solution of tert-butyl 7-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-9,9-difluoro-3,7-diazabicyclo[3.3.1]nonane-3-carboxylate (50 mg, 0.08 mmol) and TFA (1 mL) in CH₂Cl₂ (3 mL) was stirred for 1 h at room temperature. The resulting mixture was concentrated under vacuum. The residue was dissolved in NH₃ (7M in MeOH) (5 mL) and stirred for 1 h. The mixture was concentrated and the crude product was purified by prep-HPLC (Column: XBridge Prep C18 OBD, 19x250 mm, 5 μm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 30% B to 50% B in 7 min) to afford 3-amino-N-((3R)-7-(9,9-difluoro-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a yellow solid. ¹H NMR (CD₃OD, 300 MHz) δ(ppm): 8.19 (d, *J* = 8.1 Hz, 1H), 7.29 (d, *J* = 8.4 Hz, 1H), 6.99 (d, *J* = 8.4 Hz, 1H), 6.64 (dd, *J* = 2.4, 8.4 Hz, 1H), 6.48 (s, 1H), 4.41-4.48 (m, 1H), 4.21-4.25 (m, 1H), 3.82-3.96 (m, 3H), 3.35-3.37 (m, 2H), 3.13-3.19 (m, 4H), 2.80-2.98 (m, 2H), 2.63 (s, 3H), 2.15-2.18 (m, 2H). MS: (ESI, *m/z*): 500 [M+H]⁺.





[00353] The following examples in Table 10 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Example 2-1.





Table 10:




Example Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺	¹ H NMR
2-2 ¹	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	436	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.49 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.21 (br s, 1H), 6.92 (d, <i>J</i> = 8.8 Hz, 1H), 6.29 (dd, <i>J</i> = 2.0, 8.4 Hz, 1H), 6.09 (s, 1H), 4.34-4.24 (m, 1H), 4.15-4.11 (m, 1H), 3.81 (t, <i>J</i> = 10.0 Hz, 1H), 3.71-3.69 (m, 2H), 3.43-3.32 (m, 6H), 2.87-2.83 (m, 2H), 2.58 (s, 3H), 1.49-1.46 (m, 1H).
2-3 ¹	 N-((3S)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	436	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.49 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.21 (br s, 1H), 6.92 (d, <i>J</i> = 8.4 Hz, 1H), 6.29 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.09 (s, 1H), 4.34-4.24 (m, 1H), 4.15-4.11 (m, 1H), 3.80 (t, <i>J</i> = 10.0 Hz, 1H), 3.71-3.69 (m, 2H), 3.43-3.32 (m, 6H), 2.87-2.83 (m, 2H), 2.58 (s, 3H), 1.47-1.45 (m, 1H).





Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
2-4 ²	 3-amino-6-methyl-N-((R)-7-((S)-2-methylpiperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(CD ₃ OD, 300 MHz) δ(ppm): 8.20 (d, J = 8.4Hz, 1H). 7.30 (d, J = 8.4Hz, 1H), 6.99 (d, J = 8.4Hz, 1H), 6.61 (d, J = 8.7Hz, 1H), 6.46 (s, 1H), 4.48-4.42 (m, 1H), 4.26-4.22 (m, 1H), 3.94-3.87 (m, 1H), 3.63-3.56 (m, 1H), 3.10-2.75 (m, 8H), 2.63 (s, 3H), 0.98 (d, J = 6.6Hz, 1H).
2-5 ³	 3-amino-6-methyl-N-((R)-7-((R)-2-methylpiperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(CD ₃ OD, 300 MHz) δ(ppm): 8.20 (d, J = 8.4Hz, 1H). 7.30 (d, J = 8.4Hz, 1H), 6.99 (d, J = 8.4Hz, 1H), 6.61 (d, J = 8.7Hz, 1H), 6.46 (s, 1H), 4.48-4.42 (m, 1H), 4.26-4.22 (m, 1H), 3.94-3.87 (m, 1H), 3.63-3.56 (m, 1H), 3.05-2.74 (m, 8H), 2.64 (s, 3H), 0.98 (d, J = 6.6Hz, 1H).
2-6 ⁴	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-8-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	450	(CD ₃ OD, 400 MHz) δ(ppm): 8.21 (d, J = 8.4Hz, 1H). 7.32 (d, J = 8.4Hz, 1H), 6.97 (d, J = 8.4Hz, 1H), 6.47 (dd, J = 2.0, 8.4Hz, 1H), 6.35 (s, 1H), 4.48-4.44 (m, 1H), 4.27-4.23 (m, 1H), 4.11-4.08 (m, 2H), 3.96-3.91 (m, 1H), 3.18-3.15 (m, 2H), 3.02-2.97 (m, 1H), 2.90-2.84 (m, 1H), 2.65 (s, 3H), 2.60-2.57 (m, 2H), 2.08-1.96 (m, 1H).
2-7 ⁵	 (R)-N-(7-(2,5-diazaspiro[3.4]octan-5-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	450	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32(d, J = 8.0 Hz, 1H), 7.50 (br s, 1H), 7.31 (d, J = 8.0 Hz, 1H), 7.20 (br s, 2H), 6.91 (d, J = 8.8 Hz, 1H), 6.66 (d, J = 5.6 Hz, 1H), 6.51 (s, 1H), 4.17-4.14 (m, 3H), 3.81 (t, J = 10.0Hz, 1H), 3.50-3.38 (m, 2H), 3.30-3.17 (m, 3H), 2.87-2.85 (m, 1H), 2.58 (s, 3H), 2.27-2.20 (m, 3H), 1.78-1.75(m, 2H).
2-8 ⁶	 3-amino-6-methyl-N-((R)-7-((S)-3-methylpiperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, J = 8.4Hz, 1H), 7.52 (br s, 1H), 7.31 (d, J = 8.4Hz, 1H), 7.22 (br s, 2H), 6.91 (d, J = 8.4Hz, 1H), 6.50 (dd, J = 2.4, 8.4Hz, 1H), 6.29 (s, 1H), 4.31-4.25 (m, 1H), 4.17-4.12 (m, 1H), 3.83-3.76 (m, 1H), 3.44-3.40 (m, 2H), 2.93-2.72 (m, 5H), 2.58 (s, 3H), 2.50-






Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	yl)thieno[2,3-b]pyridine-2-carboxamide		2.39 (m, 2H), 2.15-2.08 (m, 1H), 1.00 (d, J = 6.3 Hz, 3H).
2-9 ⁷	 3-amino-6-methyl-N-((R)-7-((R)-3-methylpiperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(DMSO- <i>d</i> ₆ , 300 MHz) δ (ppm): 8.32 (d, J = 8.4Hz, 1H), 7.52 (br s, 1H), 7.31 (d, J = 8.4Hz, 1H), 7.22 (br s, 2H), 6.91 (d, J = 8.4Hz, 1H), 6.50 (dd, J = 2.1, 8.4Hz, 1H), 6.29 (s, 1H), 4.31-4.25 (m, 1H), 4.17-4.12 (m, 1H), 3.83-3.76 (m, 1H), 3.44-3.40 (m, 2H), 2.93-2.72 (m, 5H), 2.58 (s, 3H), 2.51-2.43 (m, 1H), 2.25 (br s, 1H), 2.15-2.08 (m, 1H), 1.00 (d, J = 6.3 Hz, 3H).
2-10 ⁸	 (R)-N-(7-(2-oxa-5,8-diazaspiro[3.5]nonan-5-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	466	(DMSO- <i>d</i> ₆ , 300 MHz) δ (ppm): 8.32 (d, J = 8.1 Hz, 1H), 7.56 (br s, 1H), 7.31(d, J = 8.1 Hz, 1H), 7.22 (br s, 2H), 6.94 (d, J = 8.4 Hz, 1H), 6.27-6.23 (m, 1H), 6.05(s, 1H), 4.48-4.43 (m, 4H), 4.33-4.14 (m, 2H), 3.85-3.79 (m, 1H), 3.32-3.31 (m, 2H), 3.13-3.08 (m, 4H), 2.90-2.87 (m, 2H), 2.58 (s, 3H).
2-11 ⁹	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)chroman-3-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	437	(DMSO- <i>d</i> ₆ , 400 MHz) δ (ppm): 8.66 (s, 1H), 7.75(d, J =7.6 Hz, 1H), 6.98 (br s, 2H), 6.92(d, J =8.8Hz, 1H), 6.29 (d, J =10.8Hz, 1H), 6.09 (s, 1H), 4.33-4.28 (m, 1H), 4.16 (d, J =13.2Hz, 1H), 3.84-3.79 (m, 1H), 3.68 (d, J =6.0Hz, 2H), 3.42-3.32 (m, 4H), 2.88-2.86 (m, 2H), 2.68-2.65 (s, 3H), 2.50-2.48 (m, 1H), 1.48 (m, 1H).
2-12 ⁸	 N-((3R)-7-(3-oxa-7,9-diazabicyclo[3.3.1]nonan-7-yl)chroman-3-yl)-7-amino-3-methylthieno[2,3-b]pyridine-2-carboxamide	467	(DMSO- <i>d</i> ₆ , 400 MHz) δ (ppm): 8.66 (s, 1H), 7.77 (d, J = 7.6Hz, 1H), 6.98 (br s, 2H), 6.92 (d, J = 8.4Hz, 1H), 6.44-6.41 (m, 1H), 6.22 (s, 1H), 4.34-4.22 (m, 1H), 4.19-4.11 (m, 1H), 3.85-3.72 (m, 5H), 3.63-3.59 (m, 2H), 2.95-2.86 (m, 6H), 2.68 (s, 3H).






Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	methylthieno[2,3-b]pyrazine-6-carboxamide		
2-13 ⁸	 N-((3R)-7-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	467	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.77 (d, <i>J</i> = 7.2Hz, 1H), 6.98-6.96 (m, 3H), 6.56 (d, <i>J</i> = 8.0Hz, 1H), 4.40-4.20 (m, 1H), 4.20-4.10 (m, 1H), 3.86-3.81 (m, 1H), 3.72-3.63 (m, 4H), 3.09-3.05 (m, 2H), 3.01-2.94 (m, 4H), 2.90-2.82 (m, 2H), 2.65 (s, 3H).
2-14 ¹⁰	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)chroman-3-yl)-3-amino-6-fluorothieno[2,3-b]pyridine-2-carboxamide	440	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.64 (t, <i>J</i> = 8.0Hz, 1H), 7.61 (br s, 1H), 7.32 (br s, 2H), 7.27 (d, <i>J</i> = 8.0Hz, 1H), 6.93 (d, <i>J</i> = 8.4Hz, 1H), 6.29 (dd, <i>J</i> = 8.4, 2.0 Hz, 1H), 6.09 (br s, 1H), 4.31-4.27 (m, 1H), 4.17-4.14 (m, 1H), 3.80 (t, <i>J</i> = 10.0Hz, 1H), 3.72-3.68 (m, 2H), 3.43-3.33 (m, 4H), 2.51-2.50 (m, 2H), 2.91-2.82 (m, 2H), 1.48-1.46 (m, 1H).
2-15 ¹¹	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)chroman-3-yl)-3-amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxamide	454	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.31 (d, <i>J</i> = 10.5Hz, 1H), 7.58 (br s, 1H), 7.18 (br s, 2H), 6.92 (d, <i>J</i> = 8.7Hz, 1H), 6.28 (dd, <i>J</i> = 2.4, 8.7Hz, 1H), 6.08 (s, 1H), 4.32-4.26 (m, 1H), 4.17-4.13 (m, 1H), 3.80 (t, <i>J</i> = 9.9Hz, 1H), 3.67-3.65 (m, 2H), 3.42-3.33 (m, 4H), 2.92-2.81 (m, 2H), 2.56 (s, 3H), 2.51-2.44 (m, 1H), 1.57 (br s, 1H), 1.47-1.44 (m, 1H).
2-16 ¹²	 (R)-3-amino-5-fluoro-6-methoxy-N-(7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	458	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.33 (d, <i>J</i> = 11.1 Hz, 1H), 7.48 (br s, 1H), 7.14 (br s, 1H), 6.92 (d, <i>J</i> = 8.7Hz, 1H), 6.49 (dd, <i>J</i> = 2.4, 8.7 Hz, 1H), 6.29 (s, 1H), 4.27-4.18 (m, 1H), 4.27-4.12 (m, 1H), 4.03 (s, 1H), 3.79 (t, <i>J</i> = 9.9Hz, 1H), 2.97 - 2.94 (m, 4H), 2.86 - 2.76 (m, 6H).


Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	yl)thieno[2,3-b]pyridine-2-carboxamide		
2-17 ¹³	 (R)-3-amino-4,6-dimethyl-N-(7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 7.55(br s, 1H), 7.04 (s, 1H), 6.91 (d, <i>J</i> = 8.4Hz, 1H), 6.86 (br s, 2H), 6.50 (dd, <i>J</i> = 2.4, 8.4Hz, 1H), 6.29 (s, 1H), 4.31-4.26 (m, 1H), 4.16-4.12 (m, 1H), 3.80 (t, <i>J</i> = 9.9Hz, 1H), 2.98-2.95 (m, 4H), 2.87-2.78 (m, 6H), 2.72 (s, 3H), 2.50 (s, 3H), 2.25 (br s, 1H).
2-18 ¹⁴	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)chroman-3-yl)-3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	450	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 7.53 (br s, 1H), 7.04 (s, 1H), 6.92 (d, <i>J</i> = 8.4Hz, 1H), 6.86 (br s, 2H), 6.28 (dd, <i>J</i> = 2.1, 8.4Hz, 1H), 6.09 (s, 1H), 4.32-4.13 (m, 2H), 3.80 (t, <i>J</i> = 9.9Hz, 1H), 3.67-3.65 (m, 2H), 3.42-3.33 (m, 4H), 2.88-2.85 (m, 2H), 2.72 (s, 3H), 2.50 (s, 3H), 2.46-2.44 (m, 1H), 1.55 (br s, 1H), 1.47-1.44 (m, 1H).
2-19 ¹³	 3-amino-N-((3R)-7-(9,9-difluoro-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	514	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 7.57 (br s, 1H), 7.04 (s, 1H), 6.96 (d, <i>J</i> = 8.4 Hz, 1H), 6.87 (br s, 2H), 6.57 (dd, <i>J</i> = 2.0, 8.0Hz, 1H), 6.38 (s, 1H), 4.31-4.14 (m, 2H), 3.83-3.78 (m, 3H), 3.26-3.22 (m, 2H), 3.10-3.07 (m, 2H), 3.02-2.83 (m, 4H), 2.72 (s, 3H), 2.50 (s, 3H), 2.16-2.12 (m, 2H).
2-20 ¹⁵	 N-((3R)-7-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	480	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 7.56 (br s, 1H), 7.05 (s, 1H), 6.97 (d, <i>J</i> = 8.7Hz, 1H), 6.87 (br s, 2H), 6.57 (dd, <i>J</i> = 2.4, 8.7Hz, 1H), 6.37 (s, 1H), 4.33-4.27 (m, 1H), 4.19-4.14 (m, 1H), 3.82 (t, <i>J</i> = 9.9Hz, 1H), 3.72-3.64 (m, 4H), 3.11-2.94 (m, 6H), 2.90-2.86 (m, 2H), 2.73 (s, 3H), 2.50 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	yl)chroman-3-yl)-3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide		
2-21 ¹⁶	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)chroman-3-yl)-3-amino-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxamide	504	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 7.90 (s, 1H), 7.60 (br s, 1H), 6.92 (d, <i>J</i> = 6.3 Hz, 1H), 6.68 (br s, 2H), 6.32-6.30 (m, 1H), 6.10 (s, 1H), 4.32-4.15 (m, 2H), 3.85-3.67 (m, 3H), 3.32-3.30 (m, 3H), 2.87-2.70 (m, 3H), 2.50 (s, 3H), 1.65-1.45 (m, 2H).
2-22 ¹⁷	 3-amino-N-((3R)-7-(9,9-difluoro-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxamide	568	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 7.92 (d, <i>J</i> = 7.2Hz, 1H), 7.76 (s, 1H), 6.96 (d, <i>J</i> = 8.1Hz, 1H), 6.68-6.57 (m, 3H), 6.38 (s, 1H), 4.33-4.27 (m, 1H), 4.19-4.16 (m, 1H), 3.86-3.83 (m, 3H), 3.26-3.21 (m, 2H), 3.11-3.07 (m, 2H), 2.96-2.87 (m, 4H), 2.70 (s, 3H), 2.11-2.04 (m, 1H).
2-23 ¹⁶	 N-((3R)-7-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-3-amino-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxamide	534	(CD ₃ OD, 400 MHz) δ(ppm): 7.66 (s, 1H), 7.01 (d, <i>J</i> = 8.4Hz, 1H), 6.66-6.63 (m, 1H), 6.50 (s, 1H), 4.50-4.43 (m, 1H), 4.27-4.23 (m, 1H), 3.97-3.89 (m, 3H), 3.72-3.69 (m, 2H), 3.29-3.27 (m, 2H), 3.19-3.12 (m, 4H), 3.04-2.98 (m, 1H), 2.94-2.83 (m, 1H), 2.72 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
2-24 ¹⁸	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-[1,8]naphthyridine]-6'-carboxamide	446	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.29 (s, 1H), 7.95 (d, <i>J</i> = 7.2 Hz, 1H), 7.31 (d, <i>J</i> = 1.8 Hz, 1H), 7.21 (s, 1H), 6.87 (d, <i>J</i> = 8.7 Hz, 1H), 6.37 (dd, <i>J</i> = 2.1, 8.7 Hz, 1H), 6.16 (s, 1H), 4.24-4.11 (m, 2H), 3.75-3.68 (m, 1H), 3.47-3.44 (m, 2H), 3.37-3.34 (m, 2H), 3.30-3.28 (m, 2H), 2.89-2.72 (m, 2H), 2.67-2.64 (m, 2H), 2.27-2.21 (m, 1H), 1.64-1.60 (m, 6H), 1.02-0.98 (m, 2H), 0.76-0.73 (m, 2H).
2-25 ¹⁹	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-6-(benzylamino)nicotinamide	470	(CD ₃ OD, 300 MHz) δ(ppm): 8.47 (s, 1H), 7.83 (br s, 1H), 7.32-7.24 (m, 5H), 6.91 (d, <i>J</i> = 8.4 Hz, 1H), 6.52 (d, <i>J</i> = 8.4 Hz, 1H), 6.45 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.27 (s, 1H), 4.55 (s, 2H), 4.42-4.34 (m, 1H), 4.24-4.18 (m, 1H), 3.93-3.89 (m, 1H), 3.63-3.61 (m, 2H), 3.43-3.39 (m, 2H), 3.02-2.94 (m, 1H), 2.85-2.81 (m, 3H), 1.87-1.86 (m, 4H).
2-26 ²⁰	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)chroman-3-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide	442	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 7.41 (br s, 1H), 7.14 (br s, 2H), 6.91 (d, <i>J</i> = 8.4 Hz, 1H), 6.28 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.09 (s, 1H), 4.26-4.21 (m, 1H), 4.18-4.05 (m, 2H), 3.77 (t, <i>J</i> = 9.9 Hz, 1H), 3.68-3.67 (m, 2H), 3.43-3.56 (m, 4H), 3.18-3.16 (m, 2H), 2.84-2.81 (m, 2H), 2.79 (s, 3H), 1.47 (d, <i>J</i> = 8.1 Hz, 1H).
2-27 ²⁰	 N-((3R)-7-(3,9-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide	470	(CD ₃ OD, 300 MHz) δ(ppm): 6.96 (d, <i>J</i> = 8.1 Hz, 1H), 6.54 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.38 (s, 1H), 4.42-4.38 (m, 1H), 4.25-4.21 (m, 1H), 3.92-3.85 (m, 1H), 3.78-3.74 (m, 2H), 3.46-3.44 (m, 2H), 3.15-3.06 (m, 2H), 3.01-2.84 (m, 2H), 2.79 (s, 3H), 2.50-2.40 (m, 1H), 2.07-1.96 (m, 4H), 1.58-1.48 (m, 1H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
2-28 ²¹	 6-amino-N-((3R)-7-(9,9-difluoro-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-2-methylthieno[2,3-d]thiazole-5-carboxamide	506	(CD ₃ OD, 300 MHz) δ (ppm): 6.98 (d, J = 8.1 Hz, 1H), 6.65 (dd, J = 2.4, 8.4 Hz, 1H), 6.49 (s, 1H), 4.48-4.41 (m, 1H), 4.28-4.22 (m, 1H), 3.95-3.85 (m, 3H), 3.34-3.14 (m, 6H), 3.10-2.83 (m, 5H), 2.20-2.10 (m, 2H).
2-29 ²²	 N-((3R)-7-(3,9-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-7-ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxamide	447	(DMSO- d_6 , 300 MHz) δ (ppm): 8.89 (d, J = 7.8 Hz, 1H), 8.46(s, 1H), 8.10 (br s, 1H), 6.94 (d, J = 8.7 Hz, 1H), 6.75 (d, J = 3.3 Hz, 1H), 6.50 (d, J = 6.3 Hz, 1H), 6.32 (s, 1H), 4.52-4.49 (m, 4H), 4.20-4.04 (m, 2H), 3.60-3.57 (m, 2H), 3.16-3.14 (m, 1H), 2.97-2.89 (m, 5H), 2.54-2.51(m, 2H), 1.85-1.78 (m, 4H), 1.47 (t, J = 7.2 Hz, 3H).
2-30 ²³	 N-((3R)-7-(9,9-difluoro-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-1-ethyl-1H-pyrrolo[2,3-b]pyridine-5-carboxamide	482	(DMSO- d_6 , 400 MHz) δ (ppm): 8.76 (s, 1H), 8.65-8.47 (m, 2H), 7.81 (br s, 1H), 7.09-6.97 (m, 1H), 6.62-6.59 (m, 2H), 6.39 (s, 1H), 4.67-4.07 (m, 4H), 3.89-3.61 (m, 3H), 3.33-2.83 (m, 8H), 2.15-2.06 (m, 2H), 1.39 (t, J = 6.4Hz, 3H).
2-31 ²⁴	 (R)-3-amino-N-(8-cyano-5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	481	(DMSO- d_6 , 300 MHz) δ (ppm): 7.74(d, J = 6.6 Hz, 1H), 7.04 (s, 1H), 6.89 (s, 2H), 6.55 (d, J = 11.7 Hz, 1H), 4.36-4.32 (m, 2H), 4.12-4.08 (m, 1H), 3.06-3.03 (m, 4H), 2.91-2.89(m, 6H), 2.84-2.72 (s, 3H), 2.51-2.50 (m, 3H).
2-32 ²⁵	 (R)-3-amino-N-(8-cyano-5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	481	(DMSO- d_6 , 300 MHz) δ (ppm): 7.69(d, J = 6.6 Hz, 1H), 7.04 (s, 1H), 6.88 (s, 2H), 6.36 (s, 1H), 4.36-4.32 (m, 2H), 4.12-4.08 (m, 1H), 3.06-3.03 (m, 5H), 2.91-2.89(m, 5H), 2.84-2.72 (s, 3H), 2.51-2.50 (m, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	(R)-3-amino-N-(6-cyano-5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide		
2-33 ²⁶	 3-amino-N-((3R)-6-cyano-7-(9,9-difluoro-3,7-diazabicyclo[3.3.1]nonan-3-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	525	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.33 (d, J = 8.4Hz, 1H), 7.61 (br s, 1H), 7.57 (s, 1H), 7.31 (d, J = 8.4 Hz, 1H), 7.23 (br s, 2H), 6.62 (s, 1H), 4.30-4.25 (m, 2H), 3.96 (t, J = 10.2Hz, 1H), 3.65-3.57 (m, 2H), 3.38-3.25 (m, 5H), 3.08-2.94 (m, 4H), 2.58 (s, 3H), 2.14-2.07 (m, 2H).
2-34 ²⁷	 (S)-7-amino-3-methyl-N-(7-methyl-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	437	(CDCl ₃ , 300 MHz) δ(ppm): 8.46 (s, 1H), 6.93 (s, 1H), 6.80 (s, 1H), 4.43-4.40 (m, 1H), 3.41-3.32 (m, 4H), 3.18-3.10 (m, 5H), 2.97-2.84 (m, 3H), 2.73-2.63 (m, 5H), 2.31-2.10 (m, 5H), 1.95-1.85 (m, 1H).
2-35 ²⁷	 (R)-7-amino-3-methyl-N-(7-methyl-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	437	(CDCl ₃ , 300 MHz) δ(ppm): 8.47 (s, 1H), 6.94 (s, 1H), 6.81 (s, 1H), 4.42-4.40 (m, 1H), 3.40-3.32 (m, 4H), 3.18-3.10 (m, 5H), 2.98-2.84 (m, 3H), 2.74-2.63 (m, 5H), 2.29-2.10 (m, 5H), 1.94-1.85 (m, 1H)
2-36 ²⁷	 (S)-7-amino-3-methyl-N-(5-methyl-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	437	(CD ₃ OD, 300 MHz) δ(ppm): 8.58 (s, 1H), 6.92 (s, 2H), 4.28-4.18 (m, 1H), 3.09-3.00 (m, 5H), 2.95-2.75 (m, 7H), 2.69 (s, 3H), 2.25-2.19 (m, 4H), 1.92-1.79 (m, 1H).
2-37 ²⁷	 (R)-7-amino-3-methyl-N-(5-methyl-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	437	(CD ₃ OD, 300 MHz) δ(ppm): 8.59 (s, 1H), 6.93 (s, 2H), 4.28-4.18 (m, 1H), 3.12-3.00 (m, 5H), 2.99-2.79 (m, 7H), 2.76 (s, 3H), 2.27-2.19 (m, 4H), 1.92-1.79 (m, 1H).

Example Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺	¹ H NMR
	tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide		
2-38 ²⁸	 3-amino-N-((R)-7-((3S,4S)-3-amino-4-(methoxy-d3)pyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	457	not determined

¹ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system; the bromide **Intermediate 3** was used for Example 2-3. In Step 4, NH₃ treatment was not performed; prep-HPLC conditions (Column: XBridge Prep C18 OBD, 19x150 mm, 5 μm; Mobile phase A: water (0.05% NH₄OH), B: ACN; Gradient: 25% B to 55% B in 7 min) were used.

² **Notes on procedures:** In Step 4, NH₃ treatment was not performed; prep-HPLC conditions (Column: XBridge Prep C18 OBD, 30x150 mm, 5 μm; Mobile phase A: water (10 mM NH₄HCO₃ + 0.1% NH₄OH), B: ACN; Gradient: 15% B to 50% B in 7 min) were used.

³ **Notes on procedures:** In Step 1, the reaction was heated by microwave irradiation for 1 h at 140 °C. In Step 4, prep-HPLC conditions (Column: XBridge Shield RP18 OBD, 19x150 mm, 5 μm; Mobile phase A: Water (10 mM NH₄HCO₃), B: ACN; Gradient: 18% B to 58% B in 7 min) were used.

⁴ **Notes on procedures:** In Step 4, NH₃ treatment was not performed; prep-HPLC conditions (Column: XBridge Prep Phenyl OBD, 19x150 mm, 5 μm; Mobile phase A: Water (0.05% NH₄OH), B: MeOH; Gradient: 45% B to 75% B in 12 min) were used.

⁵ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system.

⁶ **Notes on procedures:** In Step 3, purification was by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: Water (10 mM NH₄HCO₃), B: ACN; Gradient: 0% B to 100% B in 35 min). In Step 4, NH₃ treatment was not performed; prep-HPLC conditions (Column: XBridge Prep C18 OBD, 19x150 mm, 5 μm; Mobile phase A: Water (0.05% NH₄OH), B: ACN; Gradient: 15% B to 65% B in 7 min) were used.

⁷ **Notes on procedures:** In Step 4, prep-HPLC conditions (Column: XBridge Prep C18 OBD, 19x250 mm, 10 μm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 30% B to 50% B in 7 min) were used.

⁸ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the amide coupling condition of HBTU and Et₃N in DMA was used.

⁹ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the amide coupling condition of HBTU and Et₃N in DMA was used.

¹⁰ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the carboxylic acid **Intermediate 18** and the amide coupling condition of HBTU and Et₃N in DMA were used. In Step 4, NH₃ treatment was not performed.

¹¹ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the carboxylic acid **Intermediate 16** and the amide coupling condition of HBTU and Et₃N in DMA were used.

¹² **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 20** and the amide coupling condition of HBTU and Et₃N in DMA were used.

¹³ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 21** was used.

¹⁴ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the carboxylic acid **Intermediate 21** and the amide coupling condition of HBTU and Et₃N in DMA were used.

¹⁵ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the carboxylic acid **Intermediate 21** and the amide coupling condition of HBTU and Et₃N in DMA were used.

¹⁶ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the carboxylic acid **Intermediate 22** and the amide coupling condition of HBTU and Et₃N in DMA were used.

¹⁷ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 22** and the amide coupling condition of HBTU and Et₃N in DMA were used.

¹⁸ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 29** and the amide coupling condition of HBTU and Et₃N in DMA were used. In Step 4, NH₃ treatment was not performed.

¹⁹ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 31** was used. In Step 4, prep-HPLC conditions (Column: XBridge Shield RP18 OBD, 19x150 mm, 5μm; Mobile phase A: Water (10 mM NH₄HCO₃), B: ACN; Gradient: 32% B to 62% B in 7 min) were used.

²⁰ **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the carboxylic acid **Intermediate 24** was used. In Step 4, NH₃ treatment was not performed; prep-HPLC conditions (Column: XBridge Shield RP18 OBD, 19x150 mm, 5μm; Mobile phase A: Water (0.05% NH₄OH), B: ACN; Gradient: 25% B to 50% B in 7 min) were used.

²¹ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 24** and the amide coupling condition of HBTU and Et₃N in DMA were used; purification was by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: Water (0.1% formic acid), B: ACN; Gradient: 0% B to 100% B in 40 min).

²² **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. In Step 3, the carboxylic acid **Intermediate 26** was used; purification was by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: Water (0.05% TFA), B: ACN; Gradient: 0% B to 70% B in 40 min). In Step 4, NH₃ treatment was not performed; prep-HPLC conditions (Column: XBridge Shield RP18 OBD, 19x150 mm, 5μm; Mobile phase A: Water (0.05% NH₄OH), B: ACN; Gradient: 35% B to 50% B in 8 min) were used.

²³ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 25** and the amide coupling condition of HBTU and Et₃N in DMA were used; purification was by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: Water (0.1% formic acid), B: ACN; Gradient: 0% B to 100% B in 40 min).

²⁴ **Notes on procedures:** The amine **Intermediate 7-2** and the carboxylic acid **Intermediate 21** were used. In the amide coupling step, HBTU was used as the coupling reagent with Et₃N as the base in DMA solvent. The resulting amide was purified by prep-HPLC: Column: C18 silica gel; Mobile phase, A: 10 mM NH₄HCO₃ in H₂O, B: ACN (0% to 60% B in 30 min). After the boc-deprotection step, the resulting amine was purified by prep-HPLC: Column: XBridge Prep C18 OBD Column, 19 x150 mm 5μm; mobile phase: water (0.05% NH₄OH) and ACN; gradient: 25% to 55% ACN over 8 min.

²⁵ **Notes on procedures:** The amine **Intermediate 7-1** and the carboxylic acid **Intermediate 21** were used. In the amide coupling step, HBTU was used as the coupling reagent with Et₃N as the base in DMA solvent. The resulting amide was purified by prep-HPLC: Column: C18 silica gel; Mobile phase A: 10 mM NH₄HCO₃ in H₂O, B: ACN; gradient: 0 to 60% B in 30 min. After the boc-deprotection step, the resulting amine was purified by prep-HPLC: Column: XBridge Shield RP18 OBD, 5μm, 19 x 150 mm; Mobile phase: water (0.05% NH₄OH) and ACN; gradient: 20% to 50% ACN over 11 min.

²⁶ **Notes on procedures:** In the amide coupling step, HBTU was used as the coupling reagent with Et₃N as the base in DMA solvent. The resulting amide was purified by silica gel chromatography (ethyl acetate/pet. ether, 1:1). After the boc-deprotection the crude TFA salt was treated with 7M NH₃ in methanol,

concentrated and purified by prep-HPLC: Column: X Bridge C18, 19 x150 mm, 5 μ m; Mobile Phase A: water (10 mM NH_4HCO_3), B: ACN; gradient: 10%B to 80%B in 10 min.


²⁷ **Notes on procedures:** In Step 1, the bromides **Intermediates 11-2** thru **11-5** were used. $\text{Pd}_2(\text{dba})_3/\text{Xphos}$ was used as the catalyst/ligand system.

²⁸ **Notes on procedures:** In Step 1, the amine **Intermediate 49** was used. In the coupling reaction, Xphos Pd G3 was used as the catalyst system, Cs_2CO_3 as the base, and 1,4-dioxane as the solvent. In Step 3, the amide coupling was accomplished with HATU and Et_3N in DMA at 50 °C. In Step 4, the boc-deprotection was accomplished with HCl in dioxane/EtOAc.

Example 3-1. 3-Amino-N-((R)-7-((3S,4S)-3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 3-2. 3-Amino-N-((R)-7-((3R,4R)-3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide

Method 1.



Step 1. Benzyl ((3R)-7-(3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidin-1-yl)chroman-3-yl)carbamate

[00354] A mixture of benzyl (R)-(7-bromochroman-3-yl)carbamate, **Intermediate 2**, (170 mg, 0.47 mmol), tert-butyl N-(4-methoxypyrrolidin-3-yl)carbamate, **Intermediate 35**, (100 mg, 0.46 mmol), RuPhos Pd G3 (38 mg, 0.05 mmol), RuPhos (23 mg, 0.05 mmol), and Cs_2CO_3 (300 mg,

0.92 mmol) in toluene (10 mL) was stirred for 2 h at 95 °C. After cooling to room temperature, the reaction was poured into water (20 mL). The resulting mixture was extracted with DCM (2 x 50 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel column (eluting with 1:1 EtOAc/pet. ether) afforded benzyl ((3R)-7-(3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidin-1-yl)chroman-3-yl)carbamate as a white solid. MS: (ESI, m/z): 498 [M+H]⁺.

Step 2. tert-Butyl (1-((R)-3-aminochroman-7-yl)-4-methoxypyrrolidin-3-yl)carbamate

[00355] A mixture of benzyl ((3R)-7-(3-((tert-butoxycarbonyl)amino)-4-methoxypyrrolidin-1-yl)chroman-3-yl)carbamate (80 mg, 0.16 mmol) and Pd/C (80 mg, 10%) in EtOAc (5 mL) was stirred for 3 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and washed with EtOAc (10 mL x 3). The filtrate was concentrated under vacuum to afford tert-butyl (1-((R)-3-aminochroman-7-yl)-4-methoxypyrrolidin-3-yl)carbamate as a colorless oil. MS: (ESI, m/z): 364 [M+H]⁺.

Step 3. tert-Butyl (1-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-4-methoxypyrrolidin-3-yl)carbamate

[00356] A solution of tert-butyl (1-((R)-3-aminochroman-7-yl)-4-methoxypyrrolidin-3-yl)carbamate (40 mg, 0.11 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (23 mg, 0.11 mmol), Et₃N (39 mg, 0.39 mmol), and HBTU (50 mg, 0.13 mmol) in DMA (2 mL) was stirred for 30 min at room temperature. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 10% to 80% B over 10 min) afforded tert-butyl (1-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-4-methoxypyrrolidin-3-yl)carbamate as a white solid. MS: (ESI, m/z): 554 [M+H]⁺.

Step 4. 3-Amino-N-((3R)-7-(3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00357] A solution of tert-butyl (1-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-4-methoxypyrrolidin-3-yl)carbamate (45 mg, 0.08 mmol) and TFA (1 mL) in CH₂Cl₂ (3 mL) was stirred for 30 min at room temperature. The resulting mixture was concentrated under vacuum. The crude product was purified by reverse phase chromatography

(Column: C18 silica gel; Mobile phase A: water (10 mM NH_4HCO_3), B: ACN; Gradient: 10% to 80% B over 10 min) to afford 3-amino-N-((3R)-7-(3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a colorless oil. MS: (ESI, m/z): 454 $[\text{M}+\text{H}]^+$.

Step 5. Amino-N-((R)-7-((3S,4S)-3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide and 3-Amino-N-((R)-7-((3R,4R)-3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00358] The racemic mixture of 3-amino-N-((3R)-7-(3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide (35 mg, 0.077 mmol) was separated by chiral HPLC (Column: Chiralpak IE, 2x25 cm, 5 μm ; Mobile phase: 50% EtOH/MTBE (0.1% Et_2NH) for 28 min; Flow rate: 16 mL/min) to afford the title compounds as follows:

Amino-N-((R)-7-((3S,4S)-3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a yellow solid (first eluting isomer, RT = 13.5 min). ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ (ppm): 8.32 (d, $J = 8.4$ Hz, 1H), 7.49 (br s, 1H), 7.31 (d, $J = 8.1$ Hz, 1H), 7.21 (br s, 2H), 6.87 (d, $J = 8.4$ Hz, 1H), 6.12-6.08 (br s, 1H), 5.91 (s, 1H), 4.31-4.21 (m, 1H), 4.16-4.11 (m, 1H), 3.79 (t, $J = 9.9$ Hz, 1H), 3.63-3.62 (m, 1H), 3.51-3.46 (m, 1H), 3.42-3.32 (m, 2H), 2.93 (s, 3H), 3.13-3.09 (m, 1H), 2.91-2.82 (m, 3H), 2.58 (s, 3H), 2.12-1.96 (m, 2H). MS: (ESI, m/z): 454 $[\text{M}+\text{H}]^+$;

and amino-N-((R)-7-((3R,4R)-3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a yellow solid (second eluting isomer, RT = 20.2 min). ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ (ppm): 8.32 (d, $J = 8.1$ Hz, 1H), 7.49 (br s, 1H), 7.31 (d, $J = 8.1$ Hz, 1H), 7.21 (br s, 2H), 6.86 (d, $J = 8.4$ Hz, 1H), 6.11-6.08 (br s, 1H), 5.91 (s, 1H), 4.31-4.21 (m, 1H), 4.16-4.11 (m, 1H), 3.79 (t, $J = 9.9$ Hz, 1H), 3.63-3.62 (m, 1H), 3.51-3.46 (m, 1H), 3.42-3.32 (m, 2H), 2.93 (s, 3H), 3.13-3.09 (m, 1H), 2.91-2.81 (m, 3H), 2.58 (s, 3H), 2.12-1.96 (m, 2H). MS: (ESI, m/z): 454 $[\text{M}+\text{H}]^+$.




Example 3-1. 3-Amino-N-((R)-7-((3S,4S)-3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide





Method 2.





[00359] Amino-N-((R)-7-((3S,4S)-3-amino-4-methoxypyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide can also be prepared from tert-butyl ((3S,4S)-4-methoxypyrrolidin-3-yl)carbamate, **Intermediate 36**, using procedures similar to Method 1, without need for the Step 5 chiral separation.





[00360] The following examples in Table 11 were prepared using standard chemical manipulations and procedures similar to Method 1 for the preparation of Examples 3-1 and 3-2.


Table 11:

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
3-3 ¹	 3-amino-N-((R)-7-((S)-3-amino-3-(trifluoromethyl)piperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	506	(CD ₃ OD, 400 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.4Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 6.99 (d, <i>J</i> = 8.4 Hz, 1H), 6.60 (dd, <i>J</i> = 2.4, 8.0Hz, 1H), 6.47 (s, 1H), 4.47-4.45 (m, 1H), 4.27-4.24 (m, 1H), 3.96-3.91 (m, 1H), 3.55-3.52 (m, 1H), 3.44-3.41 (m, 1H), 2.99-2.98 (m, 1H), 2.91-2.83 (m, 2H), 2.67-2.61 (m, 4H), 2.08-1.94 (m, 1H), 1.81-1.77 (m, 3H).
3-4 ¹	 3-amino-N-((R)-7-((R)-3-amino-3-(trifluoromethyl)piperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	506	(CD ₃ OD, 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4Hz, 1H), 7.52 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.22 (br s, 2H), 6.94 (d, <i>J</i> = 8.4 Hz, 1H), 6.55 (dd, <i>J</i> = 2.4, 8.4Hz, 1H), 6.37 (s, 1H), 4.31-4.27 (m, 1H), 4.18-4.14 (m, 1H), 3.81 (t, <i>J</i> = 9.6Hz, 1H), 3.47-3.44 (m, 1H), 3.34-3.31 (m, 1H), 2.88-2.78 (m, 3H), 2.64-2.59 (m, 4H), 2.01 (br s, 2H), 1.94-1.90 (m, 1H), 1.66-1.64 (m, 3H).
3-5 ²	 3-amino-N-((R)-7-((3R,4S)-3-amino-4-fluoropyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.0Hz, 1H), 7.48 (br s, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.20 (br s, 2H), 6.90 (d, <i>J</i> = 8.4 Hz, 1H), 6.11 (dd, <i>J</i> = 2.0, 8.4Hz, 1H), 5.93 (s, 1H), 4.96 (d, <i>J</i> = 54.1Hz, 1H), 4.33-4.26 (m, 1H), 4.16-4.08 (m, 1H), 3.79 (t, <i>J</i> = 10.0Hz, 1H), 3.65-3.37 (m, 6H), 3.02-2.90 (m, 1H), 3.86-3.79 (m, 2H), 2.58 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
3-6 ²	 3-amino-N-((R)-7-((3S,4R)-3-amino-4-fluoropyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.0Hz, 1H), 7.48 (br s, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.20 (br s, 2H), 6.90 (d, <i>J</i> = 8.4 Hz, 1H), 6.12 (dd, <i>J</i> = 2.0, 8.4Hz, 1H), 5.94 (s, 1H), 5.17-5.01 (m, 1H), 4.33-4.26 (m, 1H), 4.16-4.08 (m, 1H), 3.80 (t, <i>J</i> = 10.0Hz, 1H), 3.68-3.31 (m, 6H), 3.02-2.90 (m, 1H), 3.86-3.79 (m, 2H), 2.58 (s, 3H).
3-7 ³	 3-amino-N-((R)-7-((3S,4S)-3-amino-4-fluoropyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.0Hz, 1H), 7.48 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.20 (br s, 2H), 6.89 (d, <i>J</i> = 8.4 Hz, 1H), 6.13 (dd, <i>J</i> = 2.0, 8.4Hz, 1H), 5.95 (s, 1H), 4.96 (d, <i>J</i> = 52.4Hz, 1H), 4.33-4.26 (m, 1H), 4.16-4.08 (m, 1H), 3.80 (t, <i>J</i> = 10.0Hz, 1H), 3.68-3.55 (m, 2H), 3.50-3.42 (m, 1H), 3.41-3.35 (m, 1H), 3.17 (br s, 2H), 3.02-2.90 (m, 1H), 3.86-3.79 (m, 2H), 2.58 (s, 3H).
3-8 ³	 3-amino-N-((R)-7-((3R,4R)-3-amino-4-fluoropyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.0Hz, 1H), 7.48 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.20 (br s, 2H), 6.89 (d, <i>J</i> = 8.4 Hz, 1H), 6.13 (dd, <i>J</i> = 2.0, 8.4Hz, 1H), 5.96 (s, 1H), 4.96 (d, <i>J</i> = 52.8Hz, 1H), 4.33-4.26 (m, 1H), 4.16-4.08 (m, 1H), 3.80 (t, <i>J</i> = 10.0Hz, 1H), 3.68-3.55 (m, 2H), 3.50-3.42 (m, 1H), 3.41-3.35 (m, 1H), 3.17 (br s, 2H), 3.02-2.90 (m, 1H), 3.86-3.79 (m, 2H), 2.58 (s, 3H).
3-9 ⁴	 3-amino-N-((R)-7-((3R,4R)-3-amino-4-(trifluoromethoxy)pyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	508	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.51 (br s, 1H), 7.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.22 (br s, 2H), 6.89 (d, <i>J</i> = 8.4 Hz, 1H), 6.14 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 5.96 (s, 1H), 4.69-4.68 (m, 1H), 4.17-4.12 (m, 2H), 3.83-3.43 (m, 6H), 2.98-2.83 (m, 4H), 2.58 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
3-10 ⁴	 3-amino-N-((R)-7-((3S,4S)-3-amino-4-(trifluoromethoxy)pyrrolidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	508	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.51 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.22 (br s, 2H), 6.89 (d, <i>J</i> = 8.4 Hz, 1H), 6.14 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 5.96 (s, 1H), 4.69-4.68 (m, 1H), 4.17-4.12 (m, 2H), 3.83-3.42 (m, 6H), 2.98-2.83 (m, 4H), 2.58 (s, 3H).
3-11 ⁵	 3-amino-6-methyl-N-((R)-7-((R)-3-(trifluoromethyl)piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	492	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4Hz, 1H), 7.50 (br s, 1H), 7.31 (d, <i>J</i> = 8.0Hz, 1H), 7.20 (br s, 2H), 6.95 (d, <i>J</i> = 8.4Hz, 1H), 6.54 (dd, <i>J</i> = 2.4, 8.4Hz, 1H), 6.35 (s, 1H), 4.35-4.24 (m, 1H), 4.17-4.14 (m, 1H), 3.82 (t, <i>J</i> = 10.0Hz, 1H), 3.55-3.31 (m, 3H), 3.00-2.97 (m, 1H), 2.88-2.75 (m, 4H), 2.69-2.62 (m, 2H), 2.58 (s, 3H).
3-12 ⁵	 3-amino-6-methyl-N-((R)-7-((S)-3-(trifluoromethyl)piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	492	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4Hz, 1H), 7.50 (br s, 1H), 7.31 (d, <i>J</i> = 8.4Hz, 1H), 7.20 (br s, 2H), 6.95 (d, <i>J</i> = 8.4Hz, 1H), 6.54 (dd, <i>J</i> = 2.4, 8.4Hz, 1H), 6.35 (s, 1H), 4.35-4.24 (m, 1H), 4.17-4.14 (m, 1H), 3.82 (t, <i>J</i> = 10.0Hz, 1H), 3.55-3.31 (m, 3H), 3.00-2.97 (m, 1H), 2.88-2.75 (m, 4H), 2.69-2.62 (m, 2H), 2.58 (s, 3H).
3-13 ⁶	 N-((R)-7-((S)-2,7-diazaspiro[4.5]decan-2-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	478	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.1Hz, 1H), 7.47 (br s, 1H), 7.31(d, <i>J</i> = 8.1Hz, 1H), 7.20 (br s, 2H), 6.86 (d, <i>J</i> = 8.4Hz, 1H), 6.11 (d, <i>J</i> = 2.1, 8.4Hz, 1H), 5.91 (s, 1H), 4.27-4.15 (m, 1H), 4.14-4.11 (m, 1H), 3.78 (t, <i>J</i> = 9.9Hz, 1H), 3.22-3.17 (m, 3H), 2.92-2.83 (m, 3H), 2.82-2.56 (m, 7H), 1.92-1.88 (m, 1H), 1.70-1.66 (m, 1H), 1.65-1.44 (m, 4H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
3-14 ⁶	 N-((R)-7-((R)-2,7-diazaspiro[4.5]decan-2-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	478	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4Hz, 1H), 7.47 (br s, 1H), 7.31(d, <i>J</i> = 8.4Hz, 1H), 7.20 (br s, 2H), 6.86 (d, <i>J</i> = 8.4Hz, 1H), 6.12 (d, <i>J</i> = 2.4, 8.4Hz, 1H), 5.92 (s, 1H), 4.27-4.15 (m, 1H), 4.14-4.11 (m, 1H), 3.78 (t, <i>J</i> = 9.9Hz, 1H), 3.22-3.17 (m, 3H), 2.92-2.83 (m, 3H), 2.82-2.56 (m, 7H), 1.92-1.88 (m, 1H), 1.70-1.66 (m, 1H), 1.65-1.44 (m, 4H).
3-15 ⁷	 (1aS,7bR)-N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-1a,2,3,7b-tetrahydro-1H-cyclopropa[c][1,8]naphthyridine-6-carboxamide	432	(CD ₃ OD, 300 MHz) δ(ppm): 8.21 (s, 1H), 7.85 (s, 1H), 6.92 (d, <i>J</i> = 8.4 Hz, 1H), 6.45 (d, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.30 (s, 1H), 4.43-4.38 (m, 1H), 4.23-4.19 (m, 1H), 3.94-3.88 (m, 1H), 3.72-3.68 (m, 2H), 3.51-3.42 (m, 4H), 3.10-2.88 (m, 1H), 2.85-2.82 (m, 3H), 1.99-1.90 (m, 5H), 1.79-1.75 (m, 1H), 1.08-0.97 (m, 2H).
3-16 ⁷	 (1aR,7bS)-N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)chroman-3-yl)-1a,2,3,7b-tetrahydro-1H-cyclopropa[c][1,8]naphthyridine-6-carboxamide	432	(CD ₃ OD, 300 MHz) δ(ppm): 8.21 (s, 1H), 7.85 (s, 1H), 6.92 (d, <i>J</i> = 8.4 Hz, 1H), 6.46 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.30 (s, 1H), 4.41-4.38 (m, 1H), 4.23-4.19 (m, 1H), 3.94-3.88 (m, 1H), 3.72-3.69 (m, 2H), 3.51-3.42 (m, 4H), 3.10-2.88 (m, 1H), 2.85-2.82 (m, 3H), 1.99-1.90 (m, 5H), 1.79-1.75 (m, 1H), 1.08-0.97 (m, 2H).
3-17 ⁸	 3-amino-N-[(3R)-6,8-difluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	460	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.4 Hz, 1H), 7.63(d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.24 (br s, 2H), 6.80 (d, <i>J</i> = 11.2 Hz, 1H), 4.43-4.29 (m, 1H), 4.25-4.22 (m, 1H), 3.90-3.85 (m, 1H), 2.99-2.92 (m, 6H), 2.78-2.76 (m, 4H), 2.58 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
3-18 ⁸	 3-amino-N-[(3S)-6,8-difluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	460	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.0 Hz, 1H), 7.62 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.23 (br s, 2H), 6.81 (d, <i>J</i> = 10.8 Hz, 1H), 4.36-4.28 (m, 1H), 4.26-4.22 (m, 1H), 3.91-3.86 (m, 1H), 3.09-3.03 (m, 4H), 2.95-2.88 (m, 2H), 2.84-2.82 (m, 4H), 2.58 (s, 3H).

¹ **Notes on procedures:** In Step 1, the amine **Intermediate 37** was used; Pd(dppf)Cl₂-CH₂Cl₂/Xphos was used as the catalyst/ligand system. In Step 3, purification was by prep-HPLC (Column: XBridge Prep C18 OBD, 19x150 mm, 5 μm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 20% B to 50% B in 8 min). In Step 5, chiral HPLC (Column: Chiral Art Cellulose-SB, 2x25 cm, 5 μm; Mobile phase: 20% EtOH/MTBE for 10 min) provided Example 3-3 (RT=6.2 min, stereochemistry assumed) and Example 3-4 (RT=7.9 min, stereochemistry assumed).

² **Notes on procedures:** In Step 4, the crude product was treated with NH₃ (7M in MeOH) to provide the free base. In Step 5, chiral HPLC (Column: Chiralpak IA, 2x25 cm, 5 μm; Mobile phase A: hexanes/CH₂Cl₂=3:1, B: EtOH (0.1% Et₂NH); Flow rate: 14 mL/min; Gradient: hold 50% B in 25 min) provided Example 3-5 (RT=16.5 min, stereochemistry assumed) and Example 3-6 (RT=21.5 min, stereochemistry assumed).

³ **Notes on procedures:** In Step 4, the crude product was treated with NH₃ (7M in MeOH) to provide the free base. In Step 5, chiral HPLC (Column: Chiralpak IG, 2x25 cm, 5 μm; Mobile phase: 80% MeOH (0.1% IPA)/CH₂Cl₂ for 13.5 min; Flow rate: 18 mL/min) provided Example 3-7 (RT=8.6 min, stereochemistry assumed) and Example 3-8 (RT=10.6 min, stereochemistry assumed).

⁴ **Notes on procedures:** In Step 1, the amine **Intermediate 38** was used. In Step 4, purification was by prep-HPLC (Column: XBridge Prep C18 OBD, 30x150 mm, 5 μm; Mobile phase A: water (0.05% NH₄OH), B: ACN; Gradient: 15% B to 60% B in 7 min). In Step 5, the chiral column Chiralpak IC, 2x25 cm, 5 μm was used to provide Example 3-9 (RT=7.1 min, stereochemistry assumed) and Example 3-10 (RT=8.9 min, stereochemistry assumed).

⁵ **Notes on procedures:** In Step 3, the amide coupling condition of HOBt, EDCI, and DIEA in DMF was used. In Step 5, purification by chiral HPLC (Column: Chiralpak IA, 2x25 cm, 5 μm; Mobile phase: 50% EtOH/Hexanes (0.1% Et₂NH) for 23 min; Flow rate: 15 mL/min) provided Example 13-11 (RT=14.6 min, stereochemistry assumed) and Example 13-12 (RT=17.4 min, stereochemistry assumed).


⁶ **Notes on procedures:** In Step 3, the amide coupling condition of HOBt, EDCI, and DIEA in DMA was used. Step 5 chiral separation was not performed. Instead, chiral separation was performed on the products of Step 3: chiral HPLC (Column: Chiralpak IG, 20x250 mm, 5 μm; Mobile phase: 30% EtOH/MTBE for 12 min) provided the precursor to Example 3-13 (RT=6.8 min, stereochemistry assumed) and the precursor to Example 3-14 (RT=9.2 min, stereochemistry assumed).

⁷ **Notes on procedures:** In Step 1, Pd(dppf)Cl₂-CH₂Cl₂/Xphos was used as the catalyst/ligand system. In Step 3, the carboxylic acid **Intermediate 30** was used; the reverse phase chromatography was performed with 0% to 100% water (0.1% formic acid)/ACN over 30 min. In Step 4, HCl (4M in dioxane) was used. In Step 5, purification by chiral HPLC (Column: Chiralpak IG, 2x25 cm, 5 μm; Mobile phase: 30% MeOH (0.1% Et₂NH)/CH₂Cl₂ for 15 min) provided Example 13-15 (RT=9.5 min, stereochemistry assumed) and Example 13-16 (RT=11.0 min, stereochemistry assumed).

⁸ **Notes on procedures:** In Step 1, the bromide **Intermediate 50** was used. The coupling reaction was accomplished using the precatalyst tBuXphos Pd G3 and the phosphazene base P₂-Et in DMSO at 25 °C.

In Step 5, chiral separation was performed with the chiral column Chiralpak IC and mobile phase 50% EtOH/hexanes (0.1% Et₂NH).

Example 4-1. (R)-N-(7-(2-oxa-5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl (R)-(7-(5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-yl)carbamate

[00361] A mixture of benzyl (R)-(7-bromochroman-3-yl)carbamate, **Intermediate 2**, (166 mg, 0.44 mmol), 5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonane, **Intermediate 33**, (100 mg, 0.44 mmol), RuPhos Pd G3 (38 mg, 0.05 mmol), RuPhos (21 mg, 0.05 mmol), and Cs₂CO₃ (300 mg, 0.92 mmol) in toluene (5 mL) was stirred for 16 h at 95 °C. After cooling to room temperature, the reaction was poured into 10 mL of water and was extracted with DCM (2 x 20 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with 2:1 EtOAc/pet. ether) afforded benzyl (R)-(7-(5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-yl)carbamate as a white solid. MS (ESI, m/z): 500 [M+H]⁺.

Step 2. (R)-7-(2-Oxa-5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-amine


[00362] A mixture of benzyl (R)-(7-(5-benzyl-2-oxa-5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-yl)carbamate (80 mg, 0.15 mmol) and Pd/C (100 mg, 10%) in EtOAc (5 mL) was stirred for 2 h at room temperature under an atmosphere of hydrogen. The solid was filtered out and washed with MeOH (3x10 mL). The combined filtrate was concentrated under vacuum to give (R)-7-(2-oxa-

5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-amine as a light yellow oil. MS (ESI, m/z): 276 $[M+H]^+$.

Step 3. (R)-N-(7-(2-Oxa-5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00363] A solution of (R)-7-(2-oxa-5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-amine (20 mg, 0.07 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (18 mg, 0.08 mmol), Et_3N (33 mg, 0.33 mmol), and HBTU (42 mg, 0.11 mmol) in DMA (1 mL) was stirred for 30 min at room temperature. Purification by prep-HPLC (Column: X Bridge C18, 19x150 mm, 5 μm ; Mobile phase A: water (10 mM NH_4HCO_3), B: ACN; Gradient: 10%B to 80%B in 30 min) afforded (R)-N-(7-(2-oxa-5,8-diazaspiro[3.5]nonan-8-yl)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. ^1H -NMR (CD_3OD , 300 MHz) $\delta(ppm)$: 8.20 (d, $J = 8.4$ Hz, 1H), 7.30 (d, $J = 8.4$ Hz, 1H), 6.98 (d, $J = 8.4$ Hz, 1H), 6.62-6.59 (m, 1H), 6.46 (s, 1H), 4.53-4.44 (m, 5H), 4.26-4.21 (m, 1H), 3.95-3.91 (m, 1H), 3.31-3.29 (m, 2H), 3.03-2.91 (m, 3H), 2.89-2.88 (m, 3H), 2.63 (s, 3H). MS (ESI, m/z): 466 $[M+H]^+$.

Example 5-1. (R)-3-amino-N-(7-(4,4-difluoropiperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl (R)-7-(4,4-difluoropiperidin-1-yl)chroman-3-ylcarbamate

[00364] A mixture of benzyl (R)-7-bromochroman-3-ylcarbamate, **Intermediate 2**, (100 mg, 0.28 mmol), 4,4-difluoropiperidine (42 mg, 0.35 mmol), RuPhos Pd G3 (23 mg, 0.03 mmol), RuPhos (15 mg, 0.03 mmol), and Cs_2CO_3 (225 mg, 0.69 mmol) in toluene (30 mL) was stirred for

1 h at 100 °C. After cooling to room temperature, the reaction was poured into water (10 mL). The resulting mixture was extracted with EtOAc (3 x 20 mL). The combined organic layers were washed with brine (2 x 20 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel column (eluting with 1:3 EtOAc/pet. ether) afforded benzyl (R)-(7-(4,4-difluoropiperidin-1-yl)chroman-3-yl)carbamate as an off-white solid. MS: (ESI, m/z): 403 [M+H]⁺.

Step 2. (R)-7-(4,4-Difluoropiperidin-1-yl)chroman-3-amine




[00365] A mixture of benzyl (R)-(7-(4,4-difluoropiperidin-1-yl)chroman-3-yl)carbamate (98 mg, 0.24 mmol) and Pd/C (50 mg, 10%) in EtOAc (10 mL) was stirred for 2 h at 30 °C under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum to afford 60 mg of (R)-7-(4,4-difluoropiperidin-1-yl)chroman-3-amine as a pale yellow oil. MS: (ESI, m/z): 269 [M+H]⁺.

Step 3. (R)-3-Amino-N-(7-(4,4-difluoropiperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00366] A solution of (R)-7-(4,4-difluoropiperidin-1-yl)chroman-3-amine (60 mg, 0.22 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (58 mg, 0.28 mmol), Et₃N (90 mg, 0.89 mmol), and HBTU (106 mg, 0.28 mmol) in DMA (2 mL) was stirred for 30 min at room temperature. Purification by prep-HPLC (Column: XBridge Prep C₁₈ OBD, 30x150 mm, 5 µm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 40% B to 60% B in 7 min) afforded (R)-3-amino-N-(7-(4,4-difluoropiperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a yellow solid. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.32 (d, *J* = 8.0Hz, 1H), 7.52 (br s, 1H), 7.31 (d, *J* = 8.0Hz, 1H), 7.21 (br s, 2H), 6.95 (d, *J* = 8.4Hz, 1H), 6.57 (d, *J* = 8.4Hz, 1H), 6.41 (s, 1H), 4.31-4.27 (m, 1H), 4.18-4.14 (m, 1H), 3.82 (t, *J* = 10.0Hz, 1H), 3.28-3.25 (m, 4H), 2.92-2.83 (m, 2H), 2.58 (s, 3H), 2.09-1.97 (m, 4H). MS: (ESI, m/z): 459 [M+H]⁺.


[00367] The following examples in Table 12 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Example 5-1.

Table 12:

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
5-2 ¹	 (R)-3-amino-N-(7-(1,1-dioxidothiomorpholino)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	473	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.4 Hz, 1H), 7.55 (br s, 1H), 7.31 (d, <i>J</i> = 8.1 Hz, 1H), 7.22 (br s, 2H), 6.98 (d, <i>J</i> = 8.7 Hz, 1H), 6.59 (dd, <i>J</i> = 2.4, 8.4 Hz, 1H), 6.45 (s, 1H), 4.32-4.27 (m, 1H), 4.19-4.15 (m, 1H), 3.82 (t, <i>J</i> = 9.9 Hz, 1H), 3.77-3.72 (m, 4H), 3.11-3.04 (m, 4H), 2.94-2.83 (m, 2H), 2.58 (s, 3H).
5-3 ¹	 (R)-3-amino-N-(7-(3-hydroxy-3-(trifluoromethyl)azetidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	479	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4Hz, 1H), 7.52 (br s, 1H), 7.31 (d, <i>J</i> = 8.4Hz, 1H), 7.24 (s, 1H), 7.21 (br s, 2H), 6.92 (d, <i>J</i> = 8.0Hz, 1H), 6.09 (dd, <i>J</i> = 2.4, 8.4Hz, 1H), 5.95 (s, 1H), 4.33-4.26 (m, 1H), 4.16-4.12 (m, 1H), 4.08-4.05 (m, 2H), 3.80 (t, <i>J</i> = 10.0Hz, 1H), 3.75-3.73 (m, 2H), 2.87-2.84 (m, 2H), 2.58 (s, 3H).
5-4	 3-amino-N-((3R)-7-(9,9-difluoro-3-oxa-7-azabicyclo[3.3.1]nonan-7-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	501	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.4Hz, 1H), 7.51 (br s, 1H), 7.32 (d, <i>J</i> = 8.4Hz, 1H), 7.22 (br s, 2H), 6.93 (d, <i>J</i> = 8.4Hz, 1H), 6.45 (dd, <i>J</i> = 2.4, 11.6 Hz, 1H), 6.24 (s, 1H), 4.30-4.23 (m, 1H), 4.18-4.15 (m, 1H), 4.07-4.01 (m, 2H), 3.94-3.77 (m, 5H), 3.15-3.12 (m, 2H), 2.88-2.85 (m, 2H), 2.59 (s, 3H), 2.35-2.28 (m, 2H).

¹ Notes on procedures: In Step 3, the Xbridge Shield RP18 OBD column was used for the prep-HPLC purification.

Example 6-1. (R)-1-(3-(3-Amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperidine-4-carboxylic acid



Step 1. Ethyl (R)-1-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)piperidine-4-carboxylate

[00368] A mixture of benzyl (R)-(7-bromochroman-3-yl)carbamate, **Intermediate 2**, (200 mg, 0.52 mmol), ethyl piperidine-4-carboxylate (130 mg, 0.79 mmol), RuPhos Pd G3 (44 mg, 0.05 mmol), RuPhos (24 mg, 0.05 mmol), and Cs₂CO₃ (513 mg, 1.57 mmol) in toluene (30 mL) was stirred for 2 h at 90 °C. After cooling to room temperature, the solids were filtered out and the filtrate was concentrated under vacuum. Purification by silica gel column (eluting with 1:1 EtOAc/pet. ether) afforded ethyl (R)-1-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)piperidine-4-carboxylate as a yellow solid. MS: (ESI, m/z): 439 [M+H]⁺.

Step 2. Ethyl (R)-1-(3-aminochroman-7-yl)piperidine-4-carboxylate

[00369] A mixture of ethyl (R)-1-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)piperidine-4-carboxylate (180 mg, 0.32 mmol) and Pd/C (50 mg, 10%) in EtOAc (5 mL) was stirred for 2 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum to afford ethyl (R)-1-(3-aminochroman-7-yl)piperidine-4-carboxylate as a yellow oil. MS: (ESI, m/z): 305 [M+H]⁺.

Step 3. Ethyl (R)-1-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperidine-4-carboxylate

[00370] A solution of ethyl (R)-1-(3-aminochroman-7-yl)piperidine-4-carboxylate (110 mg, 0.33 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (107 mg, 0.49 mmol),




Et₃N (99 mg, 0.98 mmol), and HBTU (185 mg, 0.49 mmol) in DMA (5 mL) was stirred for 2 h at room temperature. The reaction was then quenched by the addition of 5 mL of water. The resulting mixture was extracted EtOAc (3 x 10 mL). The combined organic layers were washed with brine (2 x 30 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel column (eluting with 1:1 EtOAc/pet. ether) afforded ethyl (R)-1-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperidine-4-carboxylate as a yellow solid. MS: (ESI, m/z): 495 [M+H]⁺.

Step 4. (R)-1-(3-(3-Amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperidine-4-carboxylic acid

[00371] A mixture of ethyl (R)-1-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperidine-4-carboxylate (40 mg, 0.07 mmol) and LiOH (1.6 mg, 0.07 mmol) in THF (2 mL) and water (0.5 mL) was stirred for 16 h at room temperature. The resulting mixture was concentrated under vacuum. The residue was diluted with water (5 mL) and the pH was adjusted to 4 with aq. 3N HCl. The resulting mixture was washed with DCM (3 x 10 mL). Purification by prep-HPLC (Column: XBridge Shield RP18 OBD, 19x150 mm, 5μm; Mobile phase A: Water (10 mM NH₄HCO₃), B: ACN; Gradient: 5% B to 65% B in 7 min) afforded (R)-1-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)piperidine-4-carboxylic acid as a yellow solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ(ppm): 8.33 (d, *J* = 8.4 Hz, 1H), 7.53 (br s, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.22 (br s, 2H), 6.91 (d, *J* = 8.7 Hz, 1H), 6.52 (dd, *J* = 2.4, 8.4 Hz, 1H), 6.32 (s, 1H), 4.32-4.24 (m, 1H), 4.16-4.12 (m, 1H), 3.80 (t, *J* = 9.9 Hz, 1H), 3.57-3.52 (m, 2H), 2.86-2.82 (m, 2H), 2.73-2.65 (m, 2H), 2.58 (s, 3H), 2.50-2.28 (m, 1H), 1.91-1.84 (m, 2H), 1.69-1.57 (m, 2H). MS: (ESI, m/z): 467 [M+H]⁺.

[00372] The following examples in Table 13 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Example 6-1.


Table 13:

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
6-2 ¹	 (R)-1-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)pyrrolidine-3-carboxylic acid	453	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.50 (br s, 1H), 7.31 (d, <i>J</i> = 8.1 Hz, 1H), 7.21 (br s, 2H), 6.87 (d, <i>J</i> = 8.4 Hz, 1H), 6.13 (d, <i>J</i> = 7.2 Hz, 1H), 5.94 (s, 1H), 4.32-4.23 (m, 1H), 4.16-4.13 (m, 1H), 3.79 (t, <i>J</i> = 9.9 Hz, 1H), 3.40-3.30 (m, 2H), 3.28-3.03 (m, 3H), 2.91-2.73 (m, 2H), 2.58 (s, 3H), 2.14-2.01 (m, 2H).
6-3 ¹	 (S)-1-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)pyrrolidine-3-carboxylic acid	453	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.50 (br s, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.21 (br s, 2H), 6.87 (d, <i>J</i> = 8.4 Hz, 1H), 6.12 (dd, <i>J</i> = 2.0, 8.4 Hz, 1H), 5.94 (s, 1H), 4.34-4.21 (m, 1H), 4.16-4.09 (m, 1H), 7.79 (t, <i>J</i> = 10.0 Hz, 1H), 3.37-3.27 (m, 2H), 3.23-3.02 (m, 3H), 2.89-2.73 (m, 2H), 2.56 (s, 3H), 2.17-2.08 (m, 2H).
6-4 ²	 (S)-1-(6-(7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)azetidine-3-carboxylic acid	438	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.64 (s, 1H), 7.81 (d, <i>J</i> = 7.5 Hz, 1H), 6.91-6.85 (m, 3H), 6.21 (d, <i>J</i> = 7.8 Hz, 1H), 6.14 (s, 1H), 4.27-4.18 (m, 1H), 3.89-3.81 (m, 2H), 3.76-3.72 (m, 2H), 3.29-3.24 (m, 1H), 2.89-2.68 (m, 4H), 2.65 (s, 3H), 2.04-1.95 (m, 1H), 1.81-1.70 (m, 3H).

¹ Notes on procedures: In Step 1, methyl (3R)-pyrrolidine-3-carboxylate was used for Example 6-2 and methyl (3S)-pyrrolidine-3-carboxylate was used for Example 6-3.

² Notes on procedures: In Step 1, the bromide **Intermediate 9** was used.

Example 7-1. (R)-3-amino-N-(7-(4-hydroxypiperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl (R)-(7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-yl)carbamate

[00373] A mixture of benzyl (R)-(7-bromochroman-3-yl)carbamate, **Intermediate 2**, (200 mg, 0.55 mmol), 4-(tert-butyldimethylsilyloxy)piperidine, **Intermediate 39**, (142 mg, 0.66 mmol), RuPhos Pd G3 (46 mg, 0.05 mmol), RuPhos (26 mg, 0.06 mmol), and Cs₂CO₃ (539 mg, 1.65 mmol) in toluene (10 mL) was stirred for 3 h at 100 °C. After cooling to room temperature, the solids were filtered out and the filtrate was concentrated under vacuum. Purification by silica gel column (eluting with 1:3 EtOAc/pet. ether) afforded benzyl (R)-(7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-yl)carbamate as a yellow solid. MS: (ESI, m/z): 497 [M+H]⁺.

Step 2. (R)-7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-amine

[00374] A mixture of benzyl (R)-(7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-yl)carbamate (170 mg, 0.34 mmol) and Pd/C (170 mg, 10%) in EtOAc (6 mL) was stirred for 2 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum to afford (R)-7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-amine as a yellow solid. MS: (ESI, m/z): 363 [M+H]⁺.

Step 3. (R)-3-amino-N-(7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide





[00375] A solution of (R)-7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-amine (100 mg, 0.28 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (63 mg, 0.30 mmol), Et₃N (84 mg, 0.83 mmol), and HBTU (157 mg, 0.41 mmol) in DMA (5 mL) was stirred for 30 min at room temperature. The reaction was then quenched by the addition of 10 mL of water. The resulting mixture was extracted CH₂Cl₂ (3 x 15 mL). The combined organic layers were washed with brine (3 x 15 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel column (eluting with 1:1 EtOAc/pet. ether) afforded (R)-3-amino-N-(7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a yellow solid. MS: (ESI, m/z): 553 [M+H]⁺.






Step 4. (R)-3-amino-N-(7-(4-hydroxypiperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide


[00376] A solution of (R)-3-amino-N-(7-(4-(tert-butyldimethylsilyloxy)piperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide (30 mg, 0.05 mmol) and HCl (4N in dioxane) (1 mL) in CH₂Cl₂ (3 mL) was stirred for 10 min at room temperature. The pH was adjusted to 8 with sat. aq. NaHCO₃. Purification by prep-HPLC (Column: XBridge Prep C18 OBD, 30x150 mm, 5μm; Mobile phase A: Water (10 mM NH₄HCO₃), B: ACN; Flow rate: 60 mL/min; Gradient: 15% B to 50% B in 7 min) afforded (R)-3-amino-N-(7-(4-hydroxypiperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ(ppm): 8.32 (d, *J* = 8.1Hz, 1H), 7.52 (br s, 1H), 7.32 (d, *J* = 8.1Hz, 1H), 7.22 (br s, 2H), 6.91 (d, *J* = 8.4Hz, 1H), 6.51 (dd, *J* = 2.4, 8.4Hz, 1H), 6.31 (s, 1H), 4.65 (br s, 1H), 4.29-4.26 (m, 1H), 4.17-4.13 (m, 1H), 3.80 (t, *J* = 9.9Hz, 1H), 3.64-3.58 (m, 1H), 3.48-3.44 (m, 2H), 2.87-2.73 (m, 4H), 2.59 (s, 3H), 1.80-1.76 (m, 2H), 1.48-1.43 (m, 2H). MS: (ESI, m/z): 439 [M+H]⁺.

[00377] The following examples in Table 14 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Example 7-1.

Table 14:

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
7-2 ¹	 3-amino-N-((R)-7-((S)-3-hydroxy-3-methylpiperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	453	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.4 Hz, 1H), 7.51 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.22 (br s, 2H), 6.90 (d, <i>J</i> = 8.4 Hz, 1H), 6.48 (dd, <i>J</i> = 8.4, 2.0 Hz, 1H), 6.29 (br s, 1H), 4.39 (br s, 1H), 4.31-4.26 (m, 1H), 4.15 (dd, <i>J</i> = 10.0, 2.8 Hz, 1H), 3.80 (t, <i>J</i> = 10.0 Hz, 1H), 3.01-2.96 (m, 2H), 2.90-2.83 (m, 4H), 2.59 (s, 3H), 1.79-1.76 (m, 1H), 1.55-1.45 (m, 3H), 1.16 (s, 3H).
7-3 ¹	 3-amino-N-((R)-7-((R)-3-hydroxy-3-methylpiperidin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	453	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.4 Hz, 1H), 7.51 (br s, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.22 (br s, 2H), 6.90 (d, <i>J</i> = 8.4 Hz, 1H), 6.48 (dd, <i>J</i> = 8.4, 2.0 Hz, 1H), 6.29 (br s, 1H), 4.39 (br s, 1H), 4.34-4.25 (m, 1H), 4.16-4.13 (m, 1H), 3.80 (t, <i>J</i> = 10.0 Hz, 1H), 3.04-2.93 (m, 2H), 2.91-2.83 (m, 4H), 2.59 (s, 3H), 1.82-1.77 (m, 1H), 1.50-1.47 (m, 3H), 1.16 (s, 3H).
7-4 ²	 7-amino-N-((S)-6-((R)-6-hydroxy-1,4-diazepan-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	453	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.63 (s, 1H), 7.77-7.74 (m, 1H), 6.94-6.84 (m, 3H), 6.71-6.49 (m, 2H), 5.14-5.07 (m, 1H), 4.14-3.95 (m, 2H), 3.66-3.58 (m, 2H), 3.42-3.16 (m, 3H), 3.00-2.70 (m, 8H), 2.63 (s, 3H), 1.98-1.89 (m, 1H), 1.80-1.67 (m, 1H).
7-5 ²	 7-amino-N-((R)-6-((R)-6-hydroxy-1,4-diazepan-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	453	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.63 (s, 1H), 7.77-7.74 (m, 1H), 6.88-6.84 (m, 3H), 6.58-6.48 (m, 2H), 5.12-5.05 (m, 1H), 4.14-3.92 (m, 2H), 3.69-3.42 (m, 4H), 2.98-2.70 (m, 9H), 2.63 (s, 3H), 1.95-1.89 (m, 1H), 1.79-1.65 (m, 1H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
7-6 ²	 7-amino-N-((S)-6-((S)-6-hydroxy-1,4-diazepan-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	453	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.65 (s, 1H), 7.78 (d, <i>J</i> = 7.8 Hz, 1H), 6.91-6.86 (m, 3H), 6.59-6.50 (m, 2H), 5.19-5.12 (m, 1H), 4.17-3.98 (m, 2H), 3.69-3.23 (m, 5H), 3.02-2.72 (m, 8H), 2.65 (s, 3H), 1.96-1.75 (m, 1H), 1.24-1.16 (m, 1H).
7-7 ²	 7-amino-N-((R)-6-((S)-6-hydroxy-1,4-diazepan-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	453	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.65 (s, 1H), 7.78 (d, <i>J</i> = 7.5 Hz, 1H), 6.91-6.87 (m, 3H), 6.61-6.52 (m, 2H), 5.33-5.28 (m, 1H), 4.12-4.03 (m, 2H), 3.70-3.29 (m, 5H), 3.09-2.72 (m, 8H), 2.65 (s, 3H), 1.96-1.74 (m, 1H), 1.23-1.16 (m, 1H).
7-8 ³	 3-amino-N-((S)-6-((S)-3-(hydroxymethyl)piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ(ppm): 8.19 (d, <i>J</i> = 7.8 Hz, 1H), 7.29 (d, <i>J</i> = 7.8 Hz, 1H), 6.98 (d, <i>J</i> = 7.8 Hz, 1H), 6.80-6.73 (m, 2H), 4.28-4.20 (m, 1H), 3.59-3.46 (m, 5H), 3.14-2.91 (m, 5H), 2.80-2.63 (m, 5H), 2.48-2.40 (m, 1H), 2.18-2.10 (m, 1H), 1.89-1.78 (m, 1H).
7-9 ³	 3-amino-N-((S)-6-((R)-3-(hydroxymethyl)piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ(ppm): 8.19 (d, <i>J</i> = 7.8 Hz, 1H), 7.29 (d, <i>J</i> = 7.8 Hz, 1H), 6.98 (d, <i>J</i> = 7.8 Hz, 1H), 6.80-6.73 (m, 2H), 4.28-4.17 (m, 1H), 3.62-3.47 (m, 4H), 3.18-3.12 (m, 1H), 3.08-2.86 (m, 5H), 2.82-2.69 (m, 2H), 2.64 (s, 3H), 2.49-2.39 (m, 1H), 2.17-2.08 (m, 1H), 1.91-1.73 (m, 1H).
7-10 ³	 3-amino-N-((R)-6-((S)-3-(hydroxymethyl)piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ(ppm): 8.19 (d, <i>J</i> = 8.1 Hz, 1H), 7.29 (d, <i>J</i> = 8.1 Hz, 1H), 6.98 (d, <i>J</i> = 8.4 Hz, 1H), 6.80-6.73 (m, 2H), 4.27-4.21 (m, 1H), 3.60-3.47 (m, 4H), 3.10-2.91 (m, 6H), 2.80-2.64 (m, 5H), 2.47-2.41 (m, 1H), 2.18-2.09 (m, 1H), 1.91-1.73 (m, 1H).


Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	methylthieno[2,3-b]pyridine-2-carboxamide		
7-11 ³	 3-amino-N-((R)-6-((R)-3-(hydroxymethyl)piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	452	(CD ₃ OD, 300 MHz) δ(ppm): 8.20 (d, <i>J</i> = 7.5 Hz, 1H), 7.29 (d, <i>J</i> = 8.4 Hz, 1H), 6.98 (d, <i>J</i> = 6.6 Hz, 1H), 6.80-6.73 (m, 2H), 4.28-4.18 (m, 1H), 3.59-3.46 (m, 4H), 3.10-2.92 (m, 6H), 2.80-2.63 (m, 5H), 2.46-2.43 (m, 1H), 2.17-2.09 (m, 1H), 1.87-1.76 (m, 1H).

¹ **Notes on procedures:** In Step 1, the amine **Intermediate 40** was used. The Step 4 product was separated by chiral HPLC (Column: Chiralpak IC, 2x25 cm, 5 μm; Mobile phase A: hexanes/CH₂Cl₂=3:1, B: EtOH; Flow rate: 17 mL/min; Gradient: hold 50% B in 13 min) to provide Example 7-2 (RT=9.5 min, stereochemistry assumed) and Example 7-3 (RT=11.3 min, stereochemistry assumed).

² **Notes on procedures:** In Step 1, racemic bromide **Intermediate 8**, the amine **Intermediate 45**, and Pd(dppf)Cl₂-CH₂Cl₂/Xphos as the catalyst/ligand system were used. Amidation was performed with EDCI, HOBt, and DIEA in DMF. Following Step 4, the racemate was separated by chiral HPLC using the chiral column Chiralpak IA and mobile phase 90% MeOH/DCM (0.1% Et₂NH) to provide a first eluted sample which was a mixture of 2 isomers; Example 7-5 as the second eluted sample (stereochemistry assumed), and Example 7-7 as the third eluted sample (stereochemistry assumed). The mixture of 2 isomers was further separated by chiral HPLC using the chiral column Chiralpak IC and mobile phase 30% EtOH/MTBE (0.1% Et₂NH) to provide Example 7-4 as the first eluted isomer and Example 7-6 as the second eluted isomer (stereochemistry assumed).

³ **Notes on procedures:** In Step 1, racemic bromide **Intermediate 8**, the amine **Intermediate 46**, and Pd(dppf)Cl₂-CH₂Cl₂/Xphos as the catalyst/ligand system were used. Amidation was performed with EDCI, HOBt, and DIEA in DMF. Following Step 4, the racemate was separated by chiral HPLC using the chiral column EnantioCel-C1 and mobile phase 25% EtOH/hexanes (0.1% Et₂NH) to provide a first eluted sample 1 which was a mixture of two isomers and a second eluted sample 2 which was a mixture of two isomers. Each of the two samples were further separated by SFC using the chiral column EnantioPak A1-5 and mobile phase 50% CO₂/MeOH (0.1% iPrNH₂) to provide from sample 1, Example 7-8 as the first eluted isomer and Example 7-9 as the second eluted isomer; and from sample 2, Example 7-10 as the first eluted isomer and Example 7-11 as the second eluted isomer (stereochemistry assumed).

Example 8. (R)-3-Amino-N-(7-(4-fluoropiperidin-4-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-4-hydroxypiperidine-1-carboxylate

[00378] To a solution of benzyl (R)-(7-bromochroman-3-yl)carbamate, **Intermediate 2**, (500 mg, 1.35 mmol) in THF (13 mL) was added n-BuLi (2.5 M in hexane) (2.7 mL, 6.74 mmol) at -78 °C and stirring was continued for 1 h -78 °C. To the reaction mixture was added a solution of tert-butyl 4-oxopiperidine-1-carboxylate (1.37 g, 6.88 mmol) in THF (2 mL) at -78 °C. The resulting solution was stirred for an additional 2 h at room temperature. The reaction was quenched with sat. aq. NH₄Cl (10 mL). The resulting mixture was extracted with EtOAc (3 x 30 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-4-hydroxypiperidine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 483 [M+H]⁺.

Step 2. tert-Butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-4-fluoropiperidine-1-carboxylate

[00379] To a solution of tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-4-hydroxypiperidine-1-carboxylate (180 mg, 0.36 mmol) in DCM (10 mL) was added DAST (179 mg, 1.11 mmol) at -78 °C and stirring was continued for 1 h at -78 °C. The reaction was quenched with sat. aq. Na HCO₃ (15 mL). The resulting mixture was extracted with DCM (3 x 20 mL). The

combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:4 EtOAc/pet. ether) afforded tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-4-fluoropiperidine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 485 [M+H]⁺.

Step 3. tert-Butyl (R)-4-(3-aminochroman-7-yl)-4-fluoropiperidine-1-carboxylate

[00380] A mixture of tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)chroman-7-yl)-4-fluoropiperidine-1-carboxylate (90 mg, 0.18 mmol) and Pd/C (15 mg, 5%) in EtOAc (10 mL) was stirred for 1 h at 30 °C under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum to afford tert-butyl (R)-4-(3-aminochroman-7-yl)-4-fluoropiperidine-1-carboxylate as a yellow oil. MS: (ESI, m/z): 351 [M+H]⁺.

Step 4. tert-Butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-4-fluoropiperidine-1-carboxylate

[00381] A solution of tert-butyl (R)-4-(3-aminochroman-7-yl)-4-fluoropiperidine-1-carboxylate (65 mg, 0.15 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (54 mg, 0.26 mmol), EDCI (54 mg, 0.28 mmol), HOBt (38 mg, 0.28 mmol), and DIEA (72 mg, 0.56 mmol) in DMF (5 mL) was stirred for 16 h at room temperature. The reaction was quenched with water (15 mL) and was extracted with ethyl acetate (3x10 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:20 MeOH/CH₂Cl₂) afforded tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-4-fluoropiperidine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 541 [M+H]⁺.


Step 5. (R)-3-Amino-N-(7-(4-fluoropiperidin-4-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00382] A solution of tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)-4-fluoropiperidine-1-carboxylate (35 mg, 0.06 mmol) and HCl (4 M in dioxane) (0.2 mL) in DCM (3 mL) was stirred for 1 h at room temperature. The resulting mixture was concentrated under vacuum. The residue was purified by chiral HPLC (Column: Chiralpak IG, 20x250 mm, 5 µm; Mobile phase: 30% EtOH (8 mM NH₃-MeOH)/MTBE for 15 min) to afford (R)-3-amino-N-(7-(4-fluoropiperidin-4-yl)chroman-3-yl)-6-methylthieno[2,3-

b]pyridine-2-carboxamide as a light yellow solid. ^1H NMR ($\text{DMSO}-d_6$, 300 MHz) δ (ppm): 8.33 (d, $J = 8.1\text{Hz}$, 1H). 7.60 (br s, 1H), 7.32 (d, $J = 8.1\text{ Hz}$, 1H), 7.23 (br s, 2H), 7.12 (d, $J = 7.8\text{ Hz}$, 1H), 6.90 (dd, $J = 1.5, 8.1\text{Hz}$, 1H), 6.79 (s, 1H), 4.35-4.29 (m, 1H), 4.22-4.18 (m, 1H), 3.87 (t, $J = 9.9\text{Hz}$, 1H), 2.97-2.95 (m, 2H), 2.89-2.76 (m, 4H), 2.58 (s, 3H), 2.54-2.51 (m, 1H), 1.97-1.71 (m, 4H). MS: (ESI, m/z): 441 $[\text{M}+\text{H}]^+$.

Example 9-1. N-((R)-7-(((S)-5-oxa-2-azaspiro[3.4]octan-7-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 9-2. N-((R)-7-(((R)-5-oxa-2-azaspiro[3.4]octan-7-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl (R)-(7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)chroman-3-yl)carbamate

[00383] A mixture of benzyl (R)-(7-bromochroman-3-yl)carbamate, **Intermediate 2**, (200 mg, 0.55 mmol), 4,4,5,5-tetramethyl-2-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (169 mg, 0.67 mmol), Pd(dppf)Cl₂-CH₂Cl₂ (45 mg, 0.06 mmol), and KOAc (163 mg, 1.66 mmol) in dioxane (5 mL) was stirred for 3 h at 80 °C. After cooling to room temperature, the resulting mixture was poured into 15 mL of water. The resulting mixture was extracted with ethyl acetate 3 x 15mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:50 to 1:5 EtOAc/pet. ether) gave benzyl (R)-(7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)chroman-3-yl)carbamate as a white solid. MS: (ESI, m/z): 410 [M+H]⁺.

Step 2. Benzyl (R)-(7-hydroxychroman-3-yl)carbamate

[00384] A solution of benzyl (R)-(7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)chroman-3-yl)carbamate (75 mg, 0.18 mmol) and hydrogen peroxide-urea adduct (43 mg, 0.46 mmol) in methanol (6 mL) was stirred for 30 min at room temperature. The solvent was removed under vacuum. The residue was diluted with 5 mL of water. The solids were collected by filtration to give benzyl (R)-(7-hydroxychroman-3-yl)carbamate as a white solid. MS: (ESI, m/z): 300 [M+H]⁺.

Step 3. tert-Butyl 6-(((R)-3-(((benzyloxy)carbonyl)amino)chroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate

[00385] A solution of benzyl (R)-(7-hydroxychroman-3-yl)carbamate (250 mg, 0.84 mmol), tert-butyl 7-hydroxy-5-oxa-2-azaspiro[3.4]octane-2-carboxylate (287 mg, 1.25 mmol), PPh₃ (328 mg, 1.25 mmol) and DIAD (253 mg, 1.25 mmol) in THF (8 mL) was stirred overnight at room temperature. The resulting mixture was concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:20 to 1:2 EtOAc/pet. ether) afforded tert-butyl 6-(((R)-3-(((benzyloxy)carbonyl)amino)chroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate as a white solid. MS: (ESI, m/z): 511 [M+H]⁺.

Step 4. tert-Butyl 6-(((R)-3-aminochroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate

[00386] A mixture of tert-butyl 6-(((R)-3-(((benzyloxy)carbonyl)amino)chroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate (170 mg, 0.33 mmol) and Pd/C (80 mg, 10%) in ethyl acetate (5 mL) was stirred for 3 h at room temperature under an atmosphere of hydrogen, The solids were

filtered away and the filtrate was concentrated under vacuum to give tert-butyl 6-(((R)-3-aminochroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate as light yellow oil. MS: (ESI, m/z): 377 [M+H]⁺.

Step 5. tert-Butyl 6-(((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate

[00387] A solution of tert-butyl 6-(((R)-3-aminochroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate (110 mg, 0.29 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (73 mg, 0.35 mmol), Et₃N (89 mg, 0.88 mmol), and HBTU (133 mg, 0.35 mmol) in DMA (3 mL) was stirred for 1 h at room temperature. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.1% formic acid), B: ACN; Gradient: 0% to 70% B over 20 min) afforded tert-butyl 6-(((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate as a yellow solid. MS: (ESI, m/z): 567 [M+H]⁺.

Step 6. N-((3R)-7-((2-azaspiro[3.4]octan-6-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00388] A solution of tert-butyl 6-(((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)chroman-7-yl)oxy)-2-azaspiro[3.4]octane-2-carboxylate (75 mg, 0.13 mmol), and TFA (2 mL) in DCM (4 mL) was stirred for 1 h at room temperature. The resulting mixture was concentrated under vacuum to give N-((3R)-7-((2-azaspiro[3.4]octan-6-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as the trifluoroacetate salt as a white solid. MS: (ESI, m/z): 467 [M+H]⁺.


Step 7. N-((R)-7-(((S)-2-azaspiro[3.4]octan-6-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide and N-((R)-7-(((R)-2-azaspiro[3.4]octan-6-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00389] The racemic mixture of N-((3R)-7-((2-azaspiro[3.4]octan-6-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide TFA salt (60 mg) was separated by chiral HPLC (Column: Chiralpak IG, 20x250 mm, 5 μm; Mobile phase: 20% EtOH/MTBE (0.2% IPA) for 21 min) to afford the title compounds as follows:

First eluting isomer: N-((R)-7-(((S)-2-azaspiro[3.4]octan-6-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid (RT = 14.5 min, stereochemistry assumed) ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm): 8.32 (d, J = 8.4 Hz, 1H), 7.54 (d, J = 7.6 Hz, 1H), 7.31 (d, J = 8.4 Hz, 1H), 7.21 (s, 2H), 7.00 (d, J = 8.4 Hz, 1H), 6.46-6.43 (m, 1H), 6.35 (d, J = 2.0 Hz, 1H), 4.99 (s, 1H), 4.31-4.16 (m, 2H), 3.98-3.95 (m, 1H), 3.86-3.81 (m, 2H), 3.74-3.42 (m, 4H), 2.94-2.84 (m, 2H), 2.58 (s, 3H), 2.42-2.17 (m, 2H). MS: (ESI, m/z): 467 $[\text{M}+\text{H}]^+$.

Second eluting isomer: N-((R)-7-(((R)-2-azaspiro[3.4]octan-6-yl)oxy)chroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid (second eluting isomer, RT = 17.4 min). ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm): 8.32 (d, J = 8.4 Hz, 1H), 7.54 (d, J = 7.6 Hz, 1H), 7.31 (d, J = 8.4 Hz, 1H), 7.21 (s, 2H), 7.00 (d, J = 8.8 Hz, 1H), 6.46-6.43 (m, 1H), 6.34 (s, 1H), 4.99 (s, 1H), 4.29-4.16 (m, 2H), 3.99-3.95 (s, 1H), 3.88-3.81 (m, 2H), 3.68-3.42 (m, 4H), 2.94-2.88 (m, 2H), 2.67 (s, 3H), 2.43-2.31 (m, 2H). MS: (ESI, m/z): 467 $[\text{M}+\text{H}]^+$.

Example 10-1. (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-5-fluorochroman-7-yl)piperazine-1-carboxylate

[00390] A mixture of benzyl (R)-(7-bromo-5-fluorochroman-3-yl)carbamate, **Intermediate 5-1**, (500 mg, 1.32 mmol), tert-butyl piperazine-1-carboxylate (294 mg, 1.58 mmol), Pd(dppf)Cl $_2$ -CH $_2$ Cl $_2$ (106 mg, 0.13 mmol), Xphos (63 mg, 0.13 mmol), and Cs $_2$ CO $_3$ (1.3 g, 3.99 mmol) in toluene (15 mL) was stirred in a sealed tube at 95 °C overnight. After cooling to room temperature,

the solids were filtered out and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) to give tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-5-fluorochroman-7-yl)piperazine-1-carboxylate as an off-white solid. MS: (ESI, m/z): 486 [M+H]⁺.

Step 2. tert-Butyl (R)-4-(3-amino-5-fluorochroman-7-yl)piperazine-1-carboxylate

[00391] A mixture of tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-5-fluorochroman-7-yl)piperazine-1-carboxylate (4.9 g, 10.10 mmol) and Pd/C (0.5 g, 10%) in EtOAc (100 mL) was stirred for 1 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum to afford tert-butyl (R)-4-(3-amino-5-fluorochroman-7-yl)piperazine-1-carboxylate as yellow oil. MS: (ESI, m/z): 352 [M+H]⁺.

Step 3. tert-Butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluorochroman-7-yl)piperazine-1-carboxylate

[00392] A solution of tert-butyl (R)-4-(3-amino-5-fluorochroman-7-yl)piperazine-1-carboxylate (3.5 g, 9.96 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (2.5 g, 12.01 mmol), Et₃N (3.0 g, 29.65 mmol), and HBTU (4.5 g, 11.87 mmol) in DMA (30 mL) was stirred for 30 min at room temperature. The reaction was quenched by the addition of 60 mL of water. The resulting mixture was extracted with CH₂Cl₂ (50 mL x 3). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and concentrated under vacuum. Purification by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 0% to 100% B over 30 min; Flow rate: 90 mL/min) afforded tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluorochroman-7-yl)piperazine-1-carboxylate as an off-white solid. MS: (ESI, m/z): 542 [M+H]⁺.




Step 4. (R)-3-Amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide





[00393] A solution of tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluorochroman-7-yl)piperazine-1-carboxylate (30 mg, 0.05 mmol) and TFA (1 mL) in CH₂Cl₂ (2 mL) was stirred for 30 min at room temperature. The resulting mixture was concentrated under vacuum. The crude product was purified by prep-HPLC (Column: XBridge Prep C18 OBD, 19x50 mm, 5 μm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient:





25% B to 75% B in 7 min) to afford (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a yellow solid. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm): 8.33 (d, $J = 9$ MHz, 1H), 7.58 (d, $J = 9$ MHz, 1H), 7.31 (d, $J = 6$ MHz, 1H), 7.22 (s, 2H), 6.43-6.40 (m, 1H), 6.17 (s, 1H), 4.40-4.20 (m, 1H), 4.19-4.12 (m, 1H), 3.89-3.82 (m, 1H), 3.32 (s, 1H), 3.01-2.97 (m, 4H), 2.80-2.74 (m, 6H), 2.58 (s, 3H). MS: (ESI, m/z): 442 $[\text{M}+\text{H}]^+$.





[00394] The following examples in Table 15 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Example 10-1.

Table 15:

Example Number	Structure and Compound Name	LRMS m/z $[\text{M}+\text{H}]^+$	^1H NMR
10-2 ¹	 3-amino-N-((R)-7-((3R,4R)-3-amino-4-methoxypyrrolidin-1-yl)-5-fluorochroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	472	(DMSO- d_6 , 400 MHz) δ (ppm): 8.33 (d, $J = 8.4$ Hz, 1H), 7.56 (br s, 1H), 7.31(d, $J = 8.4$ Hz, 1H), 7.23 (br s, 2H), 5.95 (d, $J = 12.4$ Hz, 1H), 5.77 (s, 1H), 4.27 (s, 1H), 4.16-4.14 (m, 1H), 3.84-3.79 (m, 1H), 3.63-3.621 (m, 1H), 3.53-3.40 (m, 2H), 3.38-3.33 (m, 1H), 3.29 (s, 3H), 3.13-3.11 (m, 1H), 2.92-2.80 (m, 2H), 2.74-2.72 (m, 1H), 2.58 (s, 3H), 1.95 (br s, 2H).
10-3 ¹	 3-amino-N-((R)-7-((3R,4R)-3-amino-4-methoxypyrrolidin-1-yl)-5-fluorochroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	472	(DMSO- d_6 , 400 MHz) δ (ppm): 8.33 (d, $J = 8.4$ Hz, 1H), 7.56 (br s, 1H), 7.31(d, $J = 8.4$ Hz, 1H), 7.23 (br s, 2H), 5.96 (d, $J = 12.4$ Hz, 1H), 5.78 (s, 1H), 4.27 (s, 1H), 4.16-4.14 (m, 1H), 3.84-3.79 (m, 1H), 3.63-3.621 (m, 1H), 3.53-3.40 (m, 2H), 3.38-3.33 (m, 1H), 3.29 (s, 3H), 3.13-3.11 (m, 1H), 2.92-2.80 (m, 2H), 2.74-2.72 (m, 1H), 2.58 (s, 3H), 1.95 (br s, 2H).
10-4 ²	 (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)-2,2,3,3,5,5,6,6-d8)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	450	(DMSO- d_6 , 400 MHz) δ (ppm): 8.32 (d, $J = 8.0$ Hz, 1H), 7.58 (br s, 1H), 7.31 (d, $J = 8.4$ Hz, 1H), 7.22 (br s, 2H), 6.38 (d, $J = 12.4$ Hz, 1H), 6.16 (s, 1H), 4.32-4.15 (m, 2H), 3.87-3.78 (m, 1H), 2.89-2.67 (m, 2H), 2.58 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	methylthieno[2,3-b]pyridine-2-carboxamide		
10-5 ^{2,3}	 (S)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	442	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.33 (d, <i>J</i> = 9 MHz, 1H), 7.58 (d, <i>J</i> = 9 MHz, 1H), 7.31 (d, <i>J</i> = 6 MHz, 1H), 7.22 (s, 2H), 6.43-6.40 (m, 1H), 6.17 (s, 1H), 4.40-4.20 (m, 1h), 4.19-4.12 (m, 1H), 3.89-3.82 (m, 1H), 3.32 (s, 1H), 3.01-2.97 (m, 4H), 2.80-2.74 (m, 6H), 2.58 (s, 3H).
10-6 ²	 N-((3R)-7-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)-5-fluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	484	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.4 Hz, 1H), 7.58 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.23 (br s, 2H), 6.46 (dd, <i>J</i> = 2.0, 12.8 Hz, 1H), 6.24 (s, 1H), 4.32-4.27 (m, 1H), 4.20-4.15 (m, 1H), 3.86 (t, <i>J</i> = 10.0 Hz, 1H), 3.79-3.77 (m, 2H), 3.68-3.65 (m, 2H), 3.12-3.00 (m, 6H), 2.91-2.67 (m, 2 H), 2.58 (s, 3H), 2.01 (br s, 1H).
10-7 ^{2,4,5}	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)-5-fluorochroman-3-yl)-3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	468	(CD ₃ OD, 400 MHz) δ(ppm): 7.05 (s, 1H), 6.15 (d, <i>J</i> = 12.6Hz, 1H), 6.06 (s, 1H), 4.48-4.37 (m, 1H), 4.28-4.23 (m, 1H), 3.96-3.93 (m, 1H), 3.85-3.83 (m, 2H), 3.52-3.48 (m, 4H), 3.12-2.93 (m, 1H), 2.79 (s, 3H), 2.77-2.55 (m, 2H), 1.67-1.64 (m, 1H).
10-8 ^{2,5}	 N-((3R)-7-(3,6-diazabicyclo[3.1.1]heptan-3-yl)-5-fluorochroman-3-yl)-7-amino-3-methylthieno[2,3-b]pyridine-2-carboxamide	455	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.80 (d, <i>J</i> = 7.2 Hz, 1H), 6.97 (br s, 2H), 6.14 (dd, <i>J</i> = 2.0, 12.8 Hz, 1H), 5.97 (s, 1H), 4.35-3.26 (m, 1H), 4.20-4.17 (m, 1H), 3.87 (t, <i>J</i> = 9.6 Hz, 1H), 3.67-3.66 (m, 2H), 3.41-3.31 (m, 4H), 2.92-2.86 (m, 1H), 2.79-2.68 (m, 1H), 2.66 (s, 3H), 2.50-2.49 (m, 1H), 1.46-1.45 (m, 1H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	methylthieno[2,3-b]pyrazine-6-carboxamide		
10-9 ^{4,5}	 (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	456	(CD ₃ OD, 400 MHz) δ (ppm): 7.04 (s, 1H), 6.34 (dd, J = 2.4, 12.4 Hz, 1H), 6.25 (s, 1H), 4.45-4.440 (m, 1H), 4.32-4.10 (m, 1H), 3.94-3.90 (m, 1H), 3.14-3.11 (m, 4H), 3.01-2.95 (m, 5H), 2.78 (s, 3H), 2.76-2.74 (m, 1H), 2.50 (s, 3H).
10-10 ⁵	 (R)-7-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	443	(DMSO- d_6 , 400 MHz) δ (ppm): 8.66 (s, 1H), 7.85 (d, J = 7.2 Hz, 1H), 6.70 (br s, 2H), 6.38 (dd, J = 2.4, 12.8 Hz, 1H), 6.17 (s, 1H), 4.34-4.25 (m, 2H), 4.19-4.16 (m, 1H), 3.89-3.84 (m, 1H), 3.00-2.98 (m, 4H), 2.91-2.85 (m, 6H), 2.79 (s, 3H).
10-11 ⁶	 (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methoxythieno[2,3-b]pyridine-2-carboxamide	458	(DMSO- d_6 , 300 MHz) δ (ppm): 8.32 (d, J = 8.7 Hz, 1H), 7.46 (br s, 1H), 7.21 (br s, 2H), 6.68 (d, J = 9.0 Hz, 1H), 6.37 (dd, J = 2.1, 12.9 Hz, 1H), 6.17 (s, 1H), 4.30-4.21 (m, 1H), 4.19-4.16 (m, 1H), 3.93 (s, 3H), 3.83 (t, J = 9.9 Hz, 1H), 3.02-2.98 (m, 4H), 2.90-2.68 (m, 6H).
10-12 ²	 N-((3R)-7-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)-5-fluorochroman-3-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	485	(DMSO- d_6 , 300 MHz) δ (ppm): 8.67 (s, 1H), 7.85 (br s, 1H), 7.00 (br s, 2H), 6.46 (dd, J = 2.1, 12.9 Hz, 1H), 6.23 (s, 1H), 4.34-4.28 (m, 1H), 4.28-4.15 (m, 1H), 3.91-3.85 (m, 1H), 3.74-3.65 (m, 4H), 3.10-2.70 (m, 9H), 2.70-2.55 (s, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
10-13 ^{2,4}	 N-((3R)-7-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)-5-fluorochroman-3-yl)-3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	498	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 7.62 (br s, 1H), 7.04 (s, 1H), 6.87 (br s, 2H), 6.45 (dd, <i>J</i> = 2.0, 12.8 Hz, 1H), 6.23 (s, 1H), 4.30-4.25 (m, 1H), 4.19-4.16 (m, 1H), 3.86 (t, <i>J</i> = 10.0 Hz, 1H), 3.74-3.72 (m, 2H), 3.68-3.65 (m, 2H), 3.13-2.95 (m, 6H), 2.90-2.85 (m, 1H), 2.79-2.66 (m, 1H), 2.62 (s, 3H), 2.53 (s, 3H).
10-14 ^{5,7}	 (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methyl-4-(trifluoromethyl)thieno[2,3-b]pyridine-2-carboxamide	510	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 7.97(d, <i>J</i> = 6.9 Hz, 1H), 7.54 (br s, 1H), 6.68 (br s, 2H), 6.38 (d, <i>J</i> = 11.1 Hz, 1H), 6.21 (s, 1H), 4.32-4.16 (m, 2H), 3.90-3.84 (m, 1H), 3.09-2.70 (m, 10H), 2.50 (s, 3H), 2.30-2.27 (m, 1H).
10-15 ^{2,5,8}	 3-amino-N-[(3R)-5-fluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-N,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	456	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 6.60 (br s, 2H), 6.40-6.37 (m, 1H), 6.17 (s, 1H), 4.57-4.53 (m, 1H), 4.29-4.27 (m, 1H), 4.18-4.13 (m, 1H), 3.05 (s, 3H), 3.00-2.98 (m, 4H), 2.95-2.88 (m, 2H), 2.85-2.84 (m, 4H), 2.57 (s, 3H).
10-16 ^{2,3,5,8}	 3-amino-N-[(3S)-5-fluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-N,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	456	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 6.60 (br s, 2H), 6.41-6.37 (m, 1H), 6.17 (s, 1H), 4.57-4.55 (m, 1H), 4.30-4.27 (m, 1H), 4.18-4.13 (m, 1H), 3.05 (s, 3H), 3.00-2.84 (m, 7H), 2.79-2.77 (m, 4H), 2.57 (s, 3H).

¹ **Notes on procedures:** In Step 1, the amine **Intermediate 35** was used; RuPhos Pd G3/RuPhos was used as the catalyst/ligand system. After Step 4, the racemic mixture of 3-amino-N-((3R)-7-(3-amino-4-methoxypyrrolidin-1-yl)-5-fluorochroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide was

separated by chiral HPLC (Column: Chiralpak IC, 2x25 cm, 5 μ m; Mobile phase A: MTBE (0.1% Et₂NH), B: EtOH; Gradient: hold 20% B in 18 min) to provide Example 10-2 (RT=8.2 min, stereochemistry assumed) and Example 10-3 (RT=10.3 min, stereochemistry assumed).

² **Notes on procedures:** In Step 1, RuPhos Pd G3/RuPhos was used as the catalyst/ligand system.

³ **Notes on procedures:** In Step 1, benzyl (S)-(7-bromo-5-fluorochroman-3-yl)carbamate was used.

⁴ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 21** was used.

⁵ **Notes on procedures:** In Step 4, the crude product was treated with NH₃ in MeOH to afford the free base before purification by prep-HPLC.

⁶ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 19** was used.


⁷ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 22** was used.

⁸ **Notes on procedures:** Before the Cbz-deprotection (Step 2), the carbamate was methylated by treatment with NaH (2 eq.) and MeI (1.5 eq.) in DMF at 25 °C. The reaction was quenched with water and an extractive workup with EtOAc was performed, followed by purification by silica gel chromatography, to afford the corresponding tert-butyl 4-(3-[[[(benzyloxy)carbonyl](methyl)amino]-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate. MS (ESI, m/z): 500 [M+H]⁺.

Example 11-1. N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 11-2. N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

Method 1.



Step 1. tert-Butyl 3-(6-(((benzyloxy)carbonyl)amino)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00395] A mixture of benzyl (6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate, **Intermediate 8**, (2 g, 5.55 mmol), tert-butyl 3,8-diazabicyclo[3.2.1]octane-8-carboxylate (1.171 g, 5.52 mmol), Pd(dppf)Cl₂ (404.96 mg, 0.55 mmol), Xphos (527 mg, 1.11 mmol), Cs₂CO₃ (5.3 g, 16.28 mmol) in toluene (40 mL) was stirred overnight at 100 °C. After cooling to room temperature, the reaction mixture was concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:100 to 2:5 EtOAc/pet. ether) afforded tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 492 [M+H]⁺.

Step 2. tert-Butyl 3-(6-amino-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00396] A mixture of tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (1.4 g, 2.85 mmol) and Pd/C (1.4 g, 13.16 mmol) in ethanol (30 mL) was stirred for 1 h at room temperature under an atmosphere of hydrogen. The solids were filtered away and the filtrate was concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:10 MeOH/CH₂Cl₂) afforded tert-butyl 3-(6-amino-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 358 [M+H]⁺.

Step 3. tert-Butyl 3-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00397] A mixture of 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (175 mg, 0.84 mmol), tert-butyl 3-(6-amino-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (300 mg, 0.84 mmol), EDCI (193 mg, 1.01 mmol), HOBt (136 mg, 1.01 mmol), DIEA (325 mg, 2.52 mmol) in DMF (20 mL) was stirred for 18 h at 20 °C. The reaction mixture was diluted with water (50 mL) and was extracted with ethyl acetate (3x50 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. Purification by silica gel chromatography (eluting with 1:3 EtOAc/pet. ether) afforded tert-butyl

3-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate. MS: (ESI, m/z): 548 [M+H]⁺.

Step 4. tert-Butyl 3-((S)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-Butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00398] The racemic mixture of tert-butyl 3-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate was separated by chiral HPLC (Column: (R,R)-WHELK-O1-Kromasil, 0.5x25 cm, 5 µm; Mobile phase: MeOH; Detector: UV 190 to 500 nm) to afford the title compounds as follows: tert-butyl 3-((S)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid (first eluting isomer, RT = 15.5 min) and tert-butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid (second eluting isomer, RT = 20.4 min).

Step 5. N-((2S)-6-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00399] A solution of tert-butyl 3-((S)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (70 mg, 0.13 mmol) and TFA (7 mL) in DCM (28 mL) was stirred for 1 h at room temperature. The reaction mixture was concentrated under vacuum. The resulting solution was diluted with 5 mL of DCM. The pH value of the solution was adjusted to 8 with NH₃ (solution in MeOH). The resulting mixture was concentrated under vacuum. Purification by prep-HPLC (Column: XBridge BEH C18 OBD, 130 Å, 19x150 mm, 5 µm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 5% B to 52% B over 8 min) afforded N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid. ¹H NMR (CD₃OD, 400 MHz) δ(ppm): 8.19 (d, *J* = 8.0 Hz, 1H), 7.30 (d, *J* = 8.0 Hz, 1H), 6.93 (d, *J* = 8.4 Hz, 1H), 6.65 (d, *J* = 8.4 Hz, 1H), 6.58 (s, 1H), 4.24-4.23 (m, 1H), 3.63-3.60 (m, 2H), 3.45-3.41 (m, 2H), 3.02-2.68 (m, 6H), 2.64 (s, 3H), 2.14-2.08 (m, 1H), 1.92-1.75 (m, 5H). MS: (ESI, m/z): 449 [M+H]⁺.

Step 6. N-((2R)-6-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00400] A solution of tert-butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (70 mg, 0.13 mmol) and TFA (7 mL) in DCM (28 mL) was stirred for 1 h at room temperature. The reaction mixture was concentrated under vacuum. The resulting solution was diluted with 5 mL of DCM. The pH value of the solution was adjusted to 8 with NH₃ (7M in MeOH). The resulting mixture was concentrated under vacuum. Purification by prep-HPLC (Column: XBridge BEH C18 OBD, 130 Å, 19x150 mm, 5 µm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 5% B to 52% B over 8 min) afforded N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid. ¹H NMR (CD₃OD, 400 MHz) δ(ppm): 8.20 (d, *J* = 8.4 Hz, 1H), 7.30 (d, *J* = 8.4 Hz, 1H), 6.93 (d, *J* = 8.4 Hz, 1H), 6.66 (d, *J* = 8.4 Hz, 1H), 6.59 (s, 1H), 4.26-4.17 (m, 1H), 3.65-3.61 (m, 2H), 3.45-3.39 (m, 3H), 3.00-2.71 (m, 6H), 2.64 (s, 3H), 2.12-2.09 (m, 1H), 1.91-1.75 (m, 5H). MS: (ESI, *m/z*): 449 [M+H]⁺.

Example 11-1. N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

Method 2.

[00401] N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide can also be prepared according to Method 1, starting from benzyl (S)-(6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate, **Intermediate 9**, and skipping the Step 4 chiral separation.




Example 11-2. N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide





Method 2.





[00402] N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide can also be prepared according to Method 1, starting from benzyl (R)-(6-bromo-1,2,3,4-tetrahydronaphthalen-2-yl)carbamate, **Intermediate 10**, and skipping the Step 4 chiral separation.






The following examples in Table 16 were prepared using standard chemical manipulations and procedures similar to Method 1 (or Method 2 where indicated) for the preparation of Examples 11-1 and 11-2.







Table 16:





Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
11-3 ¹	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxamide	466	(CD ₃ OD, 300 MHz) δ(ppm): 8.04 (d, <i>J</i> = 9.9 Hz, 1H), 6.95 (d, <i>J</i> = 8.4 Hz, 1H), 6.73 -6.57 (m, 2H), 4.23 (m, 1H), 3.70 (s, 2H), 3.47 (m, 2H), 3.06 - 2.67 (m, 6H), 2.61 (m, 3H), 2.17 - 2.01 (m, 1H), 1.88 (m, 5H).
11-4 ¹	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxamide	466	(CD ₃ OD, 300 MHz) δ(ppm): 8.04 (d, <i>J</i> = 9.9 Hz, 1H), 6.94 (d, <i>J</i> = 8.5 Hz, 1H), 6.73-6.56 (m, 1H), 4.32-4.15 (m, 1H), 3.68 (m, 1H), 3.52 -3.40 (m, 2H), 3.05- 2.67 (m, 6H), 2.61 (m, 3H), 2.11 (m, 1H), 1.98 -.74 (m, 5H).
11-5 ²	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methoxythieno[2,3-b]pyridine-2-carboxamide	464	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.30 (d, <i>J</i> =8.7Hz, 1H), 7.40 (d, <i>J</i> =7.5Hz, 1H), 7.12 (s, 1H), 6.89-6.85 (m, 2H), 6.61-6.57 (m, 1H), 6.50 (s, 1H), 4.12-4.02 (m, 1H), 3.93 (s, 3H), 3.50 (s, 2H), 3.31 (s, 2H), 2.80-2.66 (m, 6H), 2.51-2.49 m, 1H), 1.99-1.92 (m, 1H), 1.77-1.65 (m, 5H).






Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
11-6 ²	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methoxythieno[2,3-b]pyridine-2-carboxamide	464	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 8.30 (d, <i>J</i> =8.7Hz, 1H), 7.40 (d, <i>J</i> =7.8Hz, 1H), 7.12 (s, 1H), 6.88-6.85 (m, 2H), 6.61-6.57 (m, 1H), 6.51 (s, 1H), 4.11-4.02 (m, 1H), 3.93 (s, 3H), 3.51 (s, 2H), 3.31 (s, 2H), 2.80-2.67 (m, 6H), 2.51-2.49 (m, 1H), 1.97-1.94 (m, 1H), 1.77-1.68 (m, 5H).
11-7 ³	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	449	(CD ₃ OD, 400 MHz) δ(ppm): 8.57 (s, 1H), 6.94 (d, <i>J</i> = 8.4 Hz, 1H), 6.65-6.68 (m, 1H), 6.61 (s, 1H), 4.22-4.24 (m, 1H), 3.71 (s, 2H), 3.47 (d, <i>J</i> = 15.1 Hz, 2H), 2.83 – 3.04 (m, 5H), 2.68 (s, 4H), 2.10 (s, 1H), 1.85-1.88 (m, 5H).
11-8 ³	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	449	(CD ₃ OD, 400 MHz) δ(ppm): 8.57 (s, 1H), 6.94 (d, <i>J</i> = 8.4 Hz, 1H), 6.65-6.68 (m, 1H), 6.60 (d, <i>J</i> = 2.5 Hz, 1H), 4.25 (s, 1H), 3.72 (s, 2H), 3.46 (d, <i>J</i> = 4.9 Hz, 2H), 2.83 – 3.04 (m, 5H), 2.68 (s, 4H), 2.10 (s, 1H), 1.76 – 1.94 (m, 5H).
11-9 ⁴	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-1-ethyl-1H-pyrrolo[2,3-b]pyridine-5-carboxamide	430	(CD ₃ OD, 400 MHz) δ(ppm): 8.75 (d, <i>J</i> = 2.0 Hz, 1H), 8.57-8.42 (m, 1H), 7.53 (d, <i>J</i> = 3.6 Hz, 1H), 7.06-6.87 (m, 1H), 6.72 – 6.56 (m, 3H), 4.45 – 4.21 (m, 3H), 3.61 (s, 2H), 3.56-3.37 (m, 2H), 3.13 – 2.69 (m, 6H), 2.23 – 2.12 (m, 1H), 1.94 – 1.76 (m, 5H), 1.56-1.27 (m, 3H).






Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
11-10 ⁴	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-1-ethyl-1H-pyrrolo[2,3-b]pyridine-5-carboxamide	430	(CD ₃ OD, 400 MHz) δ(ppm): 8.74 (d, <i>J</i> = 2.1 Hz, 1H), 8.51-8.42 (m, 1H), 7.52 (d, <i>J</i> = 3.6 Hz, 1H), 7.08-6.79 (m, 1H), 6.67 (m, 1H), 6.65-6.47 (m, 2H), 4.42 – 4.22 (m, 3H), 3.67 (s, 2H), 3.50 – 3.41 (m, 2H), 3.10 – 2.71 (m, 6H), 2.21 – 2.12 (m, 1H), 1.95 – 1.76 (m, 5H), 1.52-1.37 (m, 3H).
11-11 ⁵	 (S)-3-amino-N-(8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	440	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.30 (d, <i>J</i> = 8.0 Hz, 1H), 7.59(d, <i>J</i> = 7.6 Hz, 1H), 7.30 (d, <i>J</i> = 8.0 Hz, 1H), 7.15(s, 2H), 6.57-6.49 (m, 2H), 4.13-4.07 (m, 1H), 3.01-2.99 (m, 4H), 2.92-2.78 (m, 7H), 2.59-2.56 (m, 4H), 1.97-1.94 (m, 1H), 1.78-1.68 (m, 1H).
11-12 ⁵	 (R)-3-amino-N-(8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	440	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.1 Hz, 1H), 7.61(d, <i>J</i> = 7.8 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 7.17(s, 2H), 6.57-6.49 (m, 2H), 4.15-4.06 (m, 1H), 3.01-2.98 (m, 4H), 2.93-2.80 (m, 7H), 2.58-2.54(m, 4H), 1.97-1.94 (m, 1H), 1.78-1.68 (m, 1H).
11-13 ⁶	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	467	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.65 (s, 1H), 7.85 (d, <i>J</i> =7.8Hz, 1H), 6.93 (s, 2H), 6.37-6.45 (m, 2H), 4.10-4.15 (m, 1H), 3.47-3.51 (m, 2H), 3.35-3.38 (m, 2H), 2.65-2.95 (m, 6H), 2.64 (s, 3H), 2.54-2.60 (m, 1H), 1.93-2.00 (m, 1H), 1.71-1.80 (m, 1H), 1.62-1.70 (m, 4H).



Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
11-14 ⁶	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	467	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.65 (s, 1H), 7.85 (d, <i>J</i> =7.8Hz, 1H), 6.93 (s, 2H), 6.37-6.45 (m, 2H), 4.10-4.15 (m, 1H), 3.47-3.51 (m, 2H), 3.35-3.38 (m, 2H), 2.65-2.95 (m, 6H), 2.64 (s, 3H), 2.54-2.60 (m, 1H), 1.93-2.00 (m, 1H), 1.71-1.80 (m, 1H), 1.62-1.70 (m, 4H).
11-15 ⁷	 3-amino-6-methyl-N-((3R,4R)-4-methyl-7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(CD ₃ OD, 300 MHz) δ(ppm): 8.22 (d, <i>J</i> =12.0Hz, 1H), 7.32(d, <i>J</i> =8.4Hz, 1H), 7.09(d, <i>J</i> =9Hz, 1H), 6.63(dd, <i>J</i> =2.4, 8.4Hz, 1H), 6.44 (s, 1H), 4.58-4.53 (m, 1H), 4.26-4.13 (m, 2H), 3.31-3.23 (m, 1H), 3.16-3.08 (m, 4H), 3.02-2.65 (m, 4H), 2.65 (s, 3H), 1.37 (d, <i>J</i> =6.9 Hz, 3H).
11-16 ⁷	 3-amino-6-methyl-N-((3R,4S)-4-methyl-7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(CD ₃ OD, 300 MHz) δ(ppm): 8.22 (d, <i>J</i> =8.4Hz, 1H), 7.32 (d, <i>J</i> =8.4Hz, 1H), 7.13 (d, <i>J</i> =8.7Hz, 1H), 6.62 (dd, <i>J</i> =2.4, 8.4Hz, 1H), 6.42 (s, 1H), 4.24-4.20 (m, 1H), 4.16-4.10 (m, 1H), 4.00-3.94 (m, 1H), 3.13-3.08 (m, 4H), 3.02-2.94 (m, 5H), 2.65 (s, 3H), 1.37 (d, <i>J</i> =6.9 Hz, 3H)
11-17 ⁷	 3-amino-6-methyl-N-((3S,4R)-4-methyl-7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(CD ₃ OD, 300 MHz) δ(ppm): 8.22(d, <i>J</i> =8.4 Hz, 1H), 7.32 (d, <i>J</i> =8.4 Hz, 1H), 7.13 (d, <i>J</i> =9.0 Hz, 1H), 6.63 (dd, <i>J</i> =2.4, 8.4Hz, 1H), 6.42 (s, 1H), 4.24-4.20 (m, 1H), 4.16-4.10 (m, 1H), 4.00-3.94 (m, 1H), 3.13-3.08 (m, 4H), 3.02-2.94 (m, 4H), 2.65 (s, 3H), 1.37 (d, <i>J</i> =6.9 Hz, 3H).
11-18 ⁷	 3-amino-6-methyl-N-((3S,4S)-4-methyl-7-(piperazin-1-yl)chroman-3-yl)thieno[2,3-b]pyridine-2-carboxamide	438	(CD ₃ OD, 300 MHz) δ(ppm): 8.22 (d, <i>J</i> =12.0Hz, 1H), 7.32(d, <i>J</i> =8.4Hz, 1H), 7.09(d, <i>J</i> =9Hz, 1H), 6.63(dd, <i>J</i> =2.4, 8.4Hz, 1H), 6.44 (s, 1H), 4.58-4.53 (m, 1H), 4.26-4.13 (m, 2H), 3.31-3.23 (m, 1H), 3.16-3.08 (m, 4H), 3.02-2.65 (m, 4H), 2.65 (s, 3H), 1.37 (d, <i>J</i> =6.9 Hz, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
11-19 ⁸	 (S)-7-amino-3-methyl-N-(6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	423	(CD ₃ OD, 400 MHz) δ(ppm): 8.56 (s, 1H), 6.98 (d, <i>J</i> = 8.4 Hz, 1H), 6.79-6.72 (m, 2H), 4.27-4.23 (m, 1H), 3.13-2.74 (m, 12H), 2.67 (s, 3H), 2.14-2.10 (m, 1H), 1.86-1.76 (m, 1H).
11-20 ⁸	 (R)-7-amino-3-methyl-N-(6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	423	(CD ₃ OD, 400 MHz) δ(ppm): 8.57 (s, 1H), 6.98 (d, <i>J</i> = 8.4 Hz, 1H), 6.80-6.73 (m, 2H), 4.28-4.23 (m, 1H), 3.13-2.74 (m, 12H), 2.68 (s, 3H), 2.14-2.11 (m, 1H), 1.87-1.75 (m, 1H).
11-21 ⁹	 (S)-3-amino-6-methyl-N-(6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyridine-2-carboxamide	422	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.29 (s, 1H), 7.54 (d, <i>J</i> = 8.0 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 7.15 (s, 2H), 6.91 (d, <i>J</i> = 8.4 Hz, 1H), 6.72-6.70 (m, 1H), 6.62 (s, 1H), 4.13-4.09 (m, 1H), 2.98-2.96 (m, 4H), 2.87-2.67 (m, 8H), 2.58 (s, 3H), 1.98-1.95 (m, 1H), 1.79-1.71 (m, 1H).
11-22 ⁹	 (R)-3-amino-6-methyl-N-(6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyridine-2-carboxamide	422	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.30 (d, <i>J</i> = 8.4 Hz, 1H), 7.54 (d, <i>J</i> = 8.0 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.15 (s, 2H), 6.91 (d, <i>J</i> = 8.4 Hz, 1H), 6.72-6.69 (m, 1H), 6.62 (s, 1H), 4.13-4.07 (m, 1H), 2.98-2.96 (m, 4H), 2.83-2.74 (m, 8H), 2.58 (s, 3H), 1.98-1.95 (m, 1H), 1.79-1.71 (m, 1H).
11-23 ¹⁰	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-7-ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxamide	431	(CD ₃ OD, 400 MHz) δ(ppm): 8.51 (s, 1H), 8.51 (s, 1H), 7.90 (d, <i>J</i> = 3.6 Hz, 1H), 6.96 (d, <i>J</i> = 8.8 Hz, 1H), 6.75-6.67 (m, 2H), 6.61 (s, 1H), 4.59-4.54 (m, 2H), 4.41-4.36 (m, 1H), 3.62 (s, 2H), 3.46-3.43 (m, 2H), 3.13-3.07 (m, 1H), 2.95-2.80 (m, 5H), 2.19-2.16 (m, 1H), 2.00-1.84 (m, 5H), 1.56-1.53 (m, 3H).
11-24 ¹⁰	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-7-ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxamide	431	(CD ₃ OD, 400 MHz) δ(ppm): 8.51 (s, 1H), 7.90 (d, <i>J</i> = 3.2 Hz, 1H), 6.96 (d, <i>J</i> = 8.4 Hz, 1H), 6.75-6.61 (m, 3H), 4.59-4.54 (m, 2H), 4.41-4.39 (m, 1H), 3.62 (s, 2H), 3.46-3.43 (m, 2H), 3.12-3.07 (m, 1H), 2.96-2.80 (m, 5H),

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	1,2,3,4-tetrahydronaphthalen-2-yl)-7-ethyl-7H-pyrrolo[2,3-c]pyridazine-3-carboxamide		2.19-2.17 (m, 1H), 1.99-1.84 (m, 5H), 1.56-1.53 (m, 3H).
11-25 ¹¹	 7-amino-3-methyl-N-((S)-6-((S)-3-(methylamino)pyrrolidin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	437	(CD ₃ OD, 300 MHz) δ(ppm): 8.57 (s, 1H), 6.95 (d, <i>J</i> =8.4Hz, 1H), 6.48 (d, <i>J</i> =8.4Hz, 1H), 6.34 (s, 1H), 4.26-4.21 (m, 1H), 3.56-3.41 (m, 3H), 3.13-3.08 (m, 2H), 3.08-2.68 (m, 4H), 2.69 (s, 3H), 2.53 (s, 3H), 2.36-2.30 (m, 1H), 2.18-1.48 (m, 3H).
11-26 ¹¹	 7-amino-3-methyl-N-((R)-6-((S)-3-(methylamino)pyrrolidin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	437	(CD ₃ OD, 300 MHz) δ(ppm): 8.57 (s, 1H), 6.95 (d, <i>J</i> =8.4Hz, 1H), 6.48 (d, <i>J</i> =8.4Hz, 1H), 6.41 (s, 1H), 4.26-4.21 (m, 1H), 3.76-3.74 (m, 1H), 3.54-3.41 (m, 2H), 3.37-3.34 (m, 1H), 3.29-3.27 (m, 1H), 3.03-2.90 (m, 3H), 2.81-2.67 (m, 7H), 2.44-2.41 (m, 1H), 2.15-2.11 (m, 2H), 1.87-1.83 (m, 1H).
11-27 ¹²	 7-amino-3-methyl-N-((S)-6-((R)-3-(methylamino)pyrrolidin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	437	(CD ₃ OD, 400 MHz) δ(ppm): 8.58 (s, 1H), 6.92 (d, <i>J</i> =8.0Hz, 1H), 6.42 (d, <i>J</i> =8.0Hz, 1H), 6.34 (s, 1H), 4.28-4.23 (m, 1H), 3.51-3.43 (m, 1H), 3.41-3.33 (m, 2H), 3.28-3.24 (m, 1H), 3.15-3.08 (m, 1H), 3.01-2.86 (m, 3H), 2.79-2.70 (m, 1H), 2.69 (s, 3H), 2.44 (s, 3H), 2.31-2.25 (m, 1H), 2.17-2.11 (m, 1H), 1.95-1.81 (m, 2H).
11-28 ¹²	 7-amino-3-methyl-N-((R)-6-((R)-3-(methylamino)pyrrolidin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-b]pyrazine-6-carboxamide	437	(CD ₃ OD, 400 MHz) δ(ppm): 8.58 (s, 1H), 6.92 (d, <i>J</i> =8.0Hz, 1H), 6.42 (d, <i>J</i> =8.0Hz, 1H), 6.34 (s, 1H), 4.32-4.26 (m, 1H), 3.51-3.45 (m, 1H), 3.41-3.33 (m, 2H), 3.28-3.24 (m, 1H), 3.11-3.08 (m, 1H), 3.01-2.86 (m, 3H), 2.79-2.73 (m, 1H), 2.69 (s, 3H), 2.45 (s, 3H), 2.29-2.25 (m, 1H), 2.14-2.11 (m, 1H), 1.95-1.81 (m, 2H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
11-29 ¹³	 N-((S)-6-((1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[3,2-b]pyrazine-6-carboxamide	435	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.80 (d, <i>J</i> =8.0Hz, 1H), 6.92 (br s, 2H), 6.85 (d, <i>J</i> =8.40 Hz, 1H), 6.37-6.34 (br s, 1H), 6.26 (s, 1H), 4.25-4.23 (m, 1H), 4.19-4.06 (m, 1H), 3.56-3.54 (m, 1H), 3.46-3.44 (m, 1H), 2.97-2.71 (m, 7H), 2.68 (s, 3H), 2.20 (br s, 1H), 2.02-1.92 (m, 1H), 1.82-1.71 (m, 2H), 1.65-1.52 (m, 1H).
11-30 ¹³	 N-((R)-6-((1S,4S)-2,5-diazabicyclo[2.2.1]heptan-2-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	435	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.80 (d, <i>J</i> =8.0Hz, 1H), 6.92 (br s, 2H), 6.85 (d, <i>J</i> =8.4 Hz, 1H), 6.37-6.34 (br s, 1H), 6.26 (s, 1H), 4.25-4.23 (m, 1H), 4.19-4.05 (m, 1H), 3.56-3.54 (m, 1H), 3.49-3.42 (m, 1H), 2.99-2.71 (m, 7H), 2.65 (s, 3H), 2.20 (br s, 1H), 2.01-1.92 (m, 1H), 1.81-1.69 (m, 2H), 1.64-1.57 (m, 1H).
11-31 ¹⁴	 N-((S)-6-((1R,4R)-2,5-diazabicyclo[2.2.1]heptan-2-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[3,2-b]pyrazine-6-carboxamide	435	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.81 (d, <i>J</i> =7.6 Hz, 1H), 6.93 (s, 2H), 6.85 (d, <i>J</i> =8.40 Hz, 1H), 6.37-6.34 (m, 1H), 6.26 (d, <i>J</i> =2.0 Hz, 1H), 4.24 (s, 1H), 4.13-4.10 (m, 1H), 3.56 (s, 1H), 3.47-3.45 (m, 1H), 2.86-2.71 (m, 7H), 2.68-2.65 (m, 4H), 1.98-1.95 (m, 1H), 1.75-1.73 (m, 2H), 1.62-1.58 (m, 1H).
11-32 ¹⁴	 N-((R)-6-((1R,4R)-2,5-diazabicyclo[2.2.1]heptan-2-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	435	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.81 (d, <i>J</i> =7.6 Hz, 1H), 6.93 (s, 2H), 6.85 (d, <i>J</i> =8.40 Hz, 1H), 6.37-6.35 (m, 1H), 6.26 (d, <i>J</i> =1.6 Hz, 1H), 4.24 (s, 1H), 4.13-4.10 (m, 1H), 3.57 (s, 1H), 3.47-3.45 (m, 1H), 2.86-2.71 (m, 7H), 2.68-2.65 (m, 4H), 1.98-1.95 (m, 1H), 1.76-1.72 (m, 2H), 1.62-1.60 (m, 1H).
11-33 ¹⁵	 (S)-6-amino-2-methyl-N-(6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-d]thiazole-5-carboxamide	428	(DMSO-d ₆ , 300 MHz) δ(ppm): 7.45 (d, <i>J</i> =7.8 Hz, 1H), 7.06 (s, 2H), 6.88 (d, <i>J</i> =8.7 Hz, 1H), 6.68 (d, <i>J</i> =8.1 Hz, 1H), 6.60 (s, 1H), 4.10-4.00 (m, 1H), 2.95-2.93 (m, 4H), 2.80-2.63 (m, 12H), 1.94-1.91 (m, 1H), 1.74-1.69 (m, 1H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
11-34 ¹⁵	 (R)-6-amino-2-methyl-N-(6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)thieno[2,3-d]thiazole-5-carboxamide	428	(DMSO-d ₆ , 300 MHz) δ(ppm): 7.45 (d, <i>J</i> = 7.8 Hz, 1H), 7.06 (s, 2H), 6.88 (d, <i>J</i> = 8.7 Hz, 1H), 6.68 (d, <i>J</i> = 8.7 Hz, 1H), 6.60 (s, 1H), 4.10-4.00 (m, 1H), 2.95-2.94 (m, 4H), 2.85-2.63 (m, 12H), 1.95-1.91 (m, 1H), 1.76-1.67 (m, 1H).
11-35 ¹⁶	 N-((2S)-6-(3,6-diazabicyclo[3.1.1]heptan-6-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	434	(CD ₃ OD, 300 MHz) δ(ppm): 8.65-8.43 (m, 1H), 8.19 (d, <i>J</i> = 8.4Hz, 1H), 7.31 (d, <i>J</i> = 8.1Hz, 1H), 7.04 (d, <i>J</i> = 8.1Hz, 1H), 6.42-6.40 (m, 2H), 4.34 (d, <i>J</i> = 6.3Hz, 1H), 4.35-4.15 (m, 1H), 3.75 (d, <i>J</i> = 13.2Hz, 2H), 3.19 (d, <i>J</i> = 12.9Hz, 2H), 3.10-2.86 (m, 4H), 2.85-2.68 (m, 1H), 2.64 (s, 3H), 2.19-2.05 (m, 1H), 1.94-1.75 (m, 2H).
11-36 ¹⁶	 N-((2R)-6-(3,6-diazabicyclo[3.1.1]heptan-6-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	434	(CD ₃ OD, 300 MHz) δ(ppm): 8.51(s, 1H), 8.20 (d, <i>J</i> = 8.4Hz, 1H), 7.31 (d, <i>J</i> = 8.4Hz, 1H), 7.04 (d, <i>J</i> = 8.1Hz, 1H), 6.48-6.40 (m, 2H), 4.34 (d, <i>J</i> = 6.0Hz, 2H), 4.28-4.22 (m, 1H), 3.75 (d, <i>J</i> = 13.2Hz, 2H), 3.19 (d, <i>J</i> = 12.9Hz, 2H), 3.17-2.76 (m, 5H), 2.64 (s, 3H), 2.19-2.08 (m, 1H), 1.94-1.79 (m, 2H).
11-37 ¹⁷	 (R)-7-amino-N-(8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	441	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.65 (s, 1H), 7.86 (d, <i>J</i> = 7.5Hz, 1H), 6.93 (s, 2H), 6.58-6.50 (m, 2H), 4.19-4.03 (m, 1H), 3.03-2.73 (m, 11H), 2.65 (s, 3H), 2.61-2.55 (m, 1H), 2.04-1.96 (m, 1H), 1.81-1.69 (m, 1H).
11-38 ¹⁷	 (S)-7-amino-N-(8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	441	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.95 (s, 1H), 7.86 (d, <i>J</i> = 7.5Hz, 1H), 6.93 (s, 2H), 6.58-6.50 (m, 2H), 4.12-4.11 (m, 1H), 3.02-2.72 (m, 11H), 2.65 (s, 3H), 2.57-2.56 (m, 1H), 2.00-1.96 (m, 1H), 1.81-1.69 (m, 1H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
11-39 ¹⁸	 N-((2S)-6-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	482	(CD ₃ OD, 300 MHz) δ(ppm): 8.23 (d, <i>J</i> = 8.4 Hz, 1H), 7.33 (d, <i>J</i> = 8.4 Hz, 1H), 6.70-6.64 (m, 2H), 4.30-4.28 (m, 1H), 3.91 (s, 2H), 3.75 (d, <i>J</i> = 11.7 Hz, 2H), 3.21-2.97 (m, 9H), 2.67-2.56 (m, 4H), 2.15-2.06 (m, 1H), 1.93-1.83 (m, 1H).
11-40 ¹⁸	 N-((2R)-6-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	482	(CD ₃ OD, 300 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.1 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 6.66-6.62 (m, 2H), 4.30-4.28 (m, 1H), 3.88 (s, 2H), 3.72 (d, <i>J</i> = 12.3 Hz, 2H), 3.30-2.95 (m, 9H), 2.64-2.56 (m, 4H), 2.15-2.06 (m, 1H), 1.93-1.83 (m, 1H).

¹ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 16** was used. In Step 4, chiral separation was achieved by SFC (Column: Chiralpak IA-SFC-02, 5x25 mm, 5 μm; Mobile phase: CO₂ (60%)/IPA(40%); Flow rate: 40 mL/min; Detector: UV 190-500 nm) to provide the precursor to Example 11-3 (RT=9.8 min, stereochemistry assumed) and the precursor to Example 11-4 (RT=12.0 min, stereochemistry assumed).

² **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 19** was used; purification was by prep-HPLC (Column: XBridge Prep C18 OBD, 19x150 mm, 5 μm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 25% B to 55% B in 7 min). In Step 4, chiral HPLC (Column: Chiralpak IC, 2x25 cm, 5 μm; Mobile phase: 50% EtOH/hexanes for 20 min; Flow rate: 16 mL/min) provided the precursor to Example 11-5 (RT=11.4 min, stereochemistry assumed) and the precursor to Example 11-6 (RT=17.5 min, stereochemistry assumed).

³ **Notes on procedures:** In Step 4, chiral HPLC (Column: Chiralpak ID, 5x25 mm, 5 μm; Mobile phase: 50% EtOH/hexanes; Detector: UV 190-500 nm) provided the precursor to Example 11-7 (RT=11.6 min, stereochemistry assumed) and the precursor to Example 11-8 (RT=13.6 min, stereochemistry assumed).

⁴ **Notes on procedures:** Following the amidation, chiral HPLC conditions: Column: Chiralpak AD-H-SL002, 20x250 mm, 5 μm; 30% IPA/hexanes; Detector: 190 nm to 500 nm. The stereochemistry of the separated enantiomers were arbitrarily assigned. Following the boc-deprotection step, prep-HPLC conditions: Column: XBridge Shield RP18 OBD, 130 Å, 5 μm, 19 x 150 mm; Mobile phase: water (10 mM NH₄HCO₃), ACN (25% ACN to 55% over 7 min).

⁵ **Notes on procedures:** In Step 1, the bromide **Intermediate 11-1** was used. In Step 3, the crude racemate was purified by reverse-phase chromatography: Column: C18 silica gel; Mobile Phase, A: 10 mM NH₄HCO₃ in H₂O, B: ACN with 0% to 90% B in 30 min. The enantiomers were separated by chiral HPLC: Column: Phenomenex Lux 5μm Cellulose-4 AXIA Packed, 2.12 x 25cm, 5μm; 50% EtOH/hexanes for 24 min. The stereochemistry of the separated enantiomers were arbitrarily assigned. Following the boc-deprotection each enantiomer was purified by prep-HPLC (Column: XBridge Prep OBD C18, 30 x150 mm 5 μm; Mobile phase: Water (10 mM NH₄HCO₃ + 0.1% NH₃-H₂O) and ACN (18% ACN to 52% in 7 min)).

⁶ **Notes on procedures:** In Step 1, the bromide **Intermediate 11-1** was used. In Step 3, the amidation utilized HOBt, EDCI as the coupling agents with Et₃N as the base and DMF solvent. The crude racemate was purified by prep-HPLC: Column: SunFire Prep C18, 19x150 mm; Mobile phase A: water (0.05% NH₄HCO₃), B: ACN (55% ACN to 90% in 7 min). The enantiomers were separated by chiral HPLC using: Column: Phenomenex Lux 5u Cellulose-4, AXIA Packed, 250x21.2 mm, 5μm; Mobile Phase: 50% EtOH/hexanes for 19 min. The stereochemistry of the separated enantiomers were arbitrarily assigned. After the boc-deprotection step, each enantiomer was purified by prep-HPLC: Column: XBridge Prep C18 OBD, 5 μm, 19x150 mm; Mobile phase: Water (10 mM NH₄HCO₃) and ACN (35% ACN up to 60% in 5 min).

⁷ **Notes on procedures:** In Step 1, the bromide **Intermediate 44** and Ruphos Pd G3 as the catalyst were used. In Step 4, the 4 enantiomers were separated by chiral HPLC using the chiral column Cellulose SB and an eluent gradient of 0% to 5% EtOH/MTBE (0.1% Et₂NH). Stereochemistry of the separated enantiomers were arbitrarily assigned.

⁸ **Notes on procedures:** In Step 4, the enantiomers were separated by chiral HPLC using the chiral column Chiralpak ID and mobile phase MeOH. Stereochemistry of the separated enantiomers were arbitrarily assigned.

⁹ **Notes on procedures:** The enantiomers were separated after the Boc deprotection rather than in Step 4, by chiral HPLC using the chiral column Phenomenex Lux 5μm Cellulose-4 and mobile phase 50% EtOH/hexanes. Stereochemistry of the separated enantiomers were arbitrarily assigned.

¹⁰ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 26** was used. In Step 4, the enantiomers were separated by chiral HPLC using the chiral column Phenomenex Lux 5μm Cellulose-4 and mobile phase 50% EtOH/hexanes. Stereochemistry of the separated enantiomers were arbitrarily assigned.

¹¹ **Notes on procedures:** In Step 1, tert-butyl N-methyl-N-[(3S)-pyrrolidin-3-yl]carbamate was used. In Step 4, the enantiomers were separated by chiral HPLC using the chiral column Chiralpak IA and mobile phase 35% EtOH/hexanes. Stereochemistry at the tetrahydronaphthalene amide was arbitrarily assigned.

¹² **Notes on procedures:** In Step 1, tert-butyl N-methyl-N-[(3R)-pyrrolidin-3-yl]carbamate was used. In Step 4, the enantiomers were separated by chiral HPLC using the chiral column (R,R)-Whelk-O1-Kromasil and mobile phase 30% DCM/EtOH. Stereochemistry at the tetrahydronaphthalene amide was arbitrarily assigned.

¹³ **Notes on procedures:** In Step 1, tert-butyl (1S,4S)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate and Ruphos Pd G3 as the catalyst were used. In Step 4, the enantiomers were separated by chiral HPLC using the chiral column Chiralpak ID and an eluent gradient of 60% to 65% MeOH/DCM. Stereochemistry at the tetrahydronaphthalene amide was arbitrarily assigned.

¹⁴ **Notes on procedures:** In Step 1, tert-butyl (1R,4R)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate and Ruphos Pd G3 as the catalyst were used. In Step 4, the enantiomers were separated by chiral HPLC using the chiral column Chiralpak AD-H-SL001 and mobile phase 40% IPA/hexanes (0.1% Et₂NH). Stereochemistry at the tetrahydronaphthalene amide was arbitrarily assigned.

¹⁵ **Notes on procedures:** In Step 3, the carboxylic acid **Intermediate 24** was used. In Step 4, the enantiomers were separated by SFC using the chiral column Phenomenex Lux 5μm Cellulose-4 and mobile phase 60% EtOH/hexanes. Stereochemistry of the separated enantiomers were arbitrarily assigned.


¹⁶ **Notes on procedures:** Example 11 Method 2 was used. In Step 1, Ruphos Pd G3 was used as the catalyst.

¹⁷ **Notes on procedures:** In Step 1, the bromide **Intermediate 11-1** was used. In Step 4, the enantiomers were separated by SFC using the chiral column Phenomenex Lux 5μm Cellulose-4 and mobile phase 50% CO₂/EtOH:ACN (1:1). Stereochemistry of the separated enantiomers were arbitrarily assigned.

¹⁸ **Notes on procedures:** In Step 1, the bromide **Intermediate 11-1** and Ruphos Pd G3 as the catalyst were used. In Step 4, the enantiomers were separated by chiral HPLC using the chiral column Phenomenex Lux 5μm Cellulose-4 and mobile phase 50% EtOH/hexanes. Stereochemistry of the separated enantiomers were arbitrarily assigned.

Example 12-1. N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide and

Example 12-2. N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide



Step 1. tert-Butyl 3-(6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00403] A solution of tert-butyl 3-(6-amino-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (150 mg, 0.42 mmol), **Intermediate 23** methyl 6-amino-2-methylthieno[2,3-d]thiazole-5-carboxylate (98 mg, 0.43 mmol), and AlMe_3 (1M in toluene, 1.7 mL, 1.7 mmol) in toluene (6 mL) was stirred overnight at 100°C . After cooling to room temperature, the reaction was quenched by the addition of 10 mL of methanol. The solids were filtered away and the filtrate was concentrated under vacuum. Purification by silica gel chromatography (eluting with gradient 1:3-1:1 EtOAc/pet. ether) afforded tert-butyl 3-(6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid. MS: (ESI, m/z): 554 $[\text{M}+\text{H}]^+$.

Step 2. tert-Butyl 3-((S)-6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-Butyl 3-

((R)-6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00404] The racemic mixture of tert-butyl 3-(6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate was separated by chiral HPLC (Column: Phenomenex Lux cellulose-4, AXIA Packed, 250 x 21.2 mm, 5 μ m; Mobile phase: 50% Hexanes:EtOH for 21 min) to afford the title compounds as follows: tert-butyl 3-((S)-6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as white solid (first eluting isomer, RT = 12.4 min, stereochemistry assumed) and tert-butyl 3-((R)-6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as white solid (second eluting isomer, RT = 16.3 min, stereochemistry assumed). MS: (ESI, m/z): 554 [M+H]⁺.

Step 3. N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide


[00405] A solution of tert-butyl 3-((S)-6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (40 mg, 0.07 mmol) and TFA (1 mL) in DCM (5 mL) was stirred for 30 min at room temperature. The resulting solution was concentrated under vacuum. The residue was diluted with 2 mL of DCM. The pH value of the solution was adjusted to 10 with a solution of ammonia in methanol (7 M). The resulting mixture was concentrated under vacuum. Purification by prep-HPLC (Column: XBridge BEH C18 OBD, 130 Å, 19x150 mm, 5 μ m; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 25% B to 45% B over 8 min) afforded N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide as a white solid. ¹H NMR (CD₃OD, 400 MHz) δ (ppm): 6.96 (d, *J* = 8.5 Hz, 1H), 6.72-6.65 (m, 1H), 6.62 (s, 1H), 4.38-4.19 (m, 1H), 3.72 (s, 2H), 3.50 (s, 2H), 3.04 – 2.81 (m, 8H), 2.85-2.67 (m, 1H), 2.11 (d, *J* = 12.0 Hz, 1H), 2.03-1.91 (m, 4H), 1.82-1.65 (m, 1H). MS: (ESI, m/z): 454 [M+H]⁺.

Step 4. N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide

[00406] A solution of tert-butyl 3-((R)-6-(6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamido)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (50 mg, 0.09 mmol) and TFA (1 mL) in DCM (5 mL) was stirred for 30 min at room temperature. The resulting solution was concentrated under vacuum. The residue was diluted with 2 mL of DCM. The pH value of the solution was adjusted to 10 with a solution of ammonia in methanol (7 M). The resulting mixture was concentrated under vacuum. Purification by prep-HPLC (Column: XBridge BEH C18 OBD, 130 Å, 19x150 mm, 5 µm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 30% B to 55% B over 7 min) afforded N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-amino-2-methylthieno[2,3-d]thiazole-5-carboxamide as a white solid. ¹H NMR (CD₃OD, 400 MHz) δ(ppm): 6.95 (d, *J* = 8.4 Hz, 1H), 6.72 – 6.65 (m, 1H), 6.61 (d, *J* = 2.3 Hz, 1H), 4.38-4.21 (m, 1H), 3.70 (s, 2H), 3.53-3.31 (m, 2H), 3.04 – 2.81 (m, 8H), 2.81-2.71 (m, 1H), 2.15 – 2.03 (m, 1H), 2.02-1.85 (m, 4H), 1.84-1.67 (m, 1H). MS: (ESI, *m/z*): 454 [M+H]⁺.

Example 13-1. N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5,8-difluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 13-2. N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5,8-difluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide.



Step 1. tert-Butyl 3-(3-(((benzyloxy)carbonyl)amino)-5-fluorochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00407] A mixture of **Intermediate 15-1** tert-butyl 3-(3-amino-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (210 mg, 0.53 mmol), benzyl chloroformate (143 mg, 0.120 mL, 0.80 mmol), potassium carbonate (200 mg, 1.45 mmol), in ethyl acetate (2 mL) and water (2 mL) was stirred for 30 min at 20 °C. The mixture was concentrated under vacuum to a crude product that was purified by silica gel chromatography (pet. ether / EtOAc = 5:1) to afford tert-butyl 3-(3-[[[(benzyloxy) carbonyl] amino]-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid.

MS: (ESI, m/z): 512 [M+H]⁺.

Step 2. tert-Butyl 3-(3-[[[(benzyloxy)carbonyl]amino]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(3-

(((benzyloxy)carbonyl)amino)-5,6-difluorochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00408] Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed tert-butyl 3-(3-[[[(benzyloxy)carbonyl]amino]-5-fluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (350 mg, 0.65 mmol), Selectfluor (267 mg, 0.75 mmol), DCM (5 mL) and methanol (5 mL). The resulting solution was stirred for 24 h at 25 °C, then the resulting mixture was concentrated under vacuum to afford a crude residue that was purified by column chromatography eluting with ethyl acetate/pet. ether (4:1) to afford tert-butyl 3-(3-[[[(benzyloxy)carbonyl]amino]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(3-(((benzyloxy)carbonyl)amino)-5,6-difluorochroman-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate that were carried on separately to the desired final examples. Both intermediates have MS: (ESI, m/z): 530 [M+H]⁺.

Step 3. tert-Butyl 3-(3-amino-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00409] Into a flask purged and maintained with nitrogen was added tert-butyl 3-(3-[[[(benzyloxy)carbonyl]amino]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (80 mg, 0.15 mmol) and methanol (6 mL). Palladium on carbon (80 mg) was added, then the resulting suspension was stirred for 1 h at 18 °C. The solids were removed by filtration over Celite, and the resulting filtrate was concentrated under vacuum to afford tert-butyl 3-(3-amino-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as yellow oil. MS: (ESI, m/z): 396 [M+H]⁺.

Step 4. 3-(3-[3-Amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00410] Into a 50-mL round-bottom flask was added tert-butyl 3-(3-amino-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (55 mg, 0.14 mmol), DMA (2 mL), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (29 mg, 0.14 mmol), DIEA (54 mg, 0.074 mL, 0.42 mmol), EDCI (53 mg, 0.28 mmol), and HOBt (38 mg, 0.28 mmol). The resulting solution was stirred for overnight at 18°C. The resulting mixture was purified

by reversed chromatography (Column: C18 silica gel; Mobile phase: 10 mM NH_4HCO_3 in $\text{H}_2\text{O}/\text{ACN} = 100/0$ increasing to 30/70 within 30 min. The combined fractions were concentrated under vacuum to afford tert-butyl 3-(3-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 586 $[\text{M}+\text{H}]^+$.

Step 5. N-((3R)-7-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-5,8-difluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide and N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5,8-difluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00411] Into a 25-mL round-bottom flask was added tert-butyl 3-(3-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (30 mg, 0.05 mmol), DCM (1.5 mL), and trifluoroacetic acid (0.5 mL). The resulting solution was stirred for 1 h at 18 °C. The resulting mixture was concentrated under vacuum to afford racemic 3-amino-N-(7-[3,8-diazabicyclo[3.2.1]octan-3-yl]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide; trifluoroacetic acid salt as a yellow solid. MS: (ESI, m/z): 486 $[\text{M}+\text{H}]^+$.

Chiral Separation:




[00412] 3-amino-N-(7-[3,8-diazabicyclo[3.2.1]octan-3-yl]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide; trifluoroacetic acid (28 mg, 0.05 mmol) was separated by chiral HPLC (Column: ChiralArt Cellulose-SB, 2 x 25 cm, 5 μm ; Mobile Phase A: hexanes (0.1% Et_2NH), B: EtOH ; Gradient: 40% B for 19 min). The first eluting isomer ($\text{RT} = 14.40$ min) was concentrated then lyophilized to afford 3-amino-N-[(3R)-7-[3,8-diazabicyclo[3.2.1]octan-3-yl]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide (stereochemistry arbitrarily assigned) as a yellow solid. MS: (ESI, m/z): 486 $[\text{M}+\text{H}]^+$. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ (ppm): 8.32 (d, $J = 8.4$ Hz, 1H), 7.65 (d, $J = 6.8$ Hz, 1H), 7.31 (d, $J = 8.0$ Hz, 1H), 7.24 (s, 2H), 6.37-6.33 (m, 1H), 4.35-4.24 (m, 2H), 3.96-3.91 (m, 1H), 3.44 (s, 2H), 3.09-2.80 (m, 6H), 2.58 (s, 3H), 1.82-1.66 (m, 4H).




The second eluting isomer ($\text{RT} = 16.71$ min) was concentrated then lyophilized to afford 3-amino-N-[(3S)-7-[3,8-diazabicyclo[3.2.1]octan-3-yl]-5,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide (stereochemistry arbitrarily assigned) as a yellow

solid. MS: (ESI, m/z): 486 $[M+H]^+$. ^1H NMR (DMSO- d_6 , 400 MHz) δ (ppm): 8.33 (d, $J = 8.4$ Hz, 1H), 7.66 (d, $J = 7.2$ Hz, 1H), 7.31 (d, $J = 8.4$ Hz, 1H), 7.24 (s, 2H), 6.39-6.35 (m, 1H), 4.33-4.24 (m, 2H), 3.96-3.92 (m, 1H), 3.58 (s, 2H), 3.11-2.73 (m, 6H), 2.58 (s, 3H), 1.90-1.69 (m, 4H), 1.23 (s, 1H).

[00413] The following examples in Table 17 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Examples 13-1 and 13-2.

Table 17:

Example Number	Structure and Compound Name	LRMS m/z $[M+H]^+$	^1H NMR
13-3 ¹	 N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5,6-difluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	486	(DMSO- d_6 , 400 MHz) δ (ppm): 8.32(d, $J = 8.4$ Hz, 1H), 7.60(d, $J = 7.2$ Hz, 1H), 7.31(d, $J = 8.4$ Hz, 1H), 7.23(s, 2H), 6.16(d, $J = 7.2$ Hz, 1H), 4.30-4.26(m, 1H), 4.16-4.13(m, 1H), 3.89-3.84(m, 1H), 3.45(s, 2H), 3.10-2.77(m, 6H), 2.58(s, 3H), 1.82-1.77(m, 4H), 1.25(s, 1H).
13-4 ¹	 N-((3S)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5,6-difluorochroman-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	486	(DMSO- d_6 , 400 MHz) δ (ppm): 8.32(d, $J = 8.4$ Hz, 1H), 7.60(d, $J = 6.8$ Hz, 1H), 7.31(d, $J = 8.4$ Hz, 1H), 7.23(s, 2H), 6.17(d, $J = 6.4$ Hz, 1H), 4.30-4.24(m, 1H), 4.16-4.13(m, 1H), 3.89-3.84(m, 1H), 3.49(s, 2H), 3.11-2.77(m, 6H), 2.58(s, 3H), 1.84-1.68(m, 4H), 1.23(s, 1H).
13-5 ²	 (R)-3-amino-N-(6-chloro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	458,460	(DMSO- d_6 , 300 MHz) δ (ppm): 8.36 (s, 1H), 8.31 (d, $J = 8.3$ Hz, 1H), 7.72 (t, $J = 5.6$ Hz, 1H), 7.31 (d, $J = 8.3$ Hz, 1H), 7.17-7.12 (m, 4H), 6.94-6.70 (m, 2H), 5.02 (t, $J = 4.5$ Hz, 1H), 3.98-4.03 (m, 1H), 3.72-3.91 (m, 5H), 3.42-3.35 (m, 2H), 2.78-2.73 (m, 2H), 2.58 (s, 3H), 2.45-2.27 (m, 2H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
13-6 ²	 (S)-3-amino-N-(6-chloro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	458,460	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.36 (s, 1H), 8.31 (d, J = 8.3 Hz, 1H), 7.72 (t, J = 5.6 Hz, 1H), 7.31 (d, J = 8.3 Hz, 1H), 7.17-7.12 (m, 4H), 6.94-6.70 (m, 2H), 5.02 (t, J = 4.5 Hz, 1H), 3.98-4.03 (m, 1H), 3.72-3.91 (m, 5H), 3.42-3.35 (m, 2H), 2.78-2.73 (m, 2H), 2.58 (s, 3H), 2.45-2.27 (m, 2H).
13-7 ²	 (R)-3-amino-N-(8-chloro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	458,460	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.36 (s, 1H), 8.31 (d, J = 8.3 Hz, 1H), 7.72 (t, J = 5.6 Hz, 1H), 7.31 (d, J = 8.3 Hz, 1H), 7.17-7.12 (m, 4H), 6.94-6.70 (m, 2H), 5.02 (t, J = 4.5 Hz, 1H), 3.98-4.03 (m, 1H), 3.72-3.91 (m, 5H), 3.42-3.35 (m, 2H), 2.78-2.73 (m, 2H), 2.58 (s, 3H), 2.45-2.27 (m, 2H).
13-8 ²	 (S)-3-amino-N-(8-chloro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	458,460	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.36 (s, 1H), 8.31 (d, J = 8.3 Hz, 1H), 7.72 (t, J = 5.6 Hz, 1H), 7.31 (d, J = 8.3 Hz, 1H), 7.17-7.12 (m, 4H), 6.94-6.70 (m, 2H), 5.02 (t, J = 4.5 Hz, 1H), 3.98-4.03 (m, 1H), 3.72-3.91 (m, 5H), 3.42-3.35 (m, 2H), 2.78-2.73 (m, 2H), 2.58 (s, 3H), 2.45-2.27 (m, 2H).

¹ Notes on procedures: Chiral separation conditions: Column: ChiralArt Cellulose-SB, 2 x 25 cm, 5 μm; mobile phase: 30% EtOH (8 mM NH₃ in MeOH)/MTBE for 15 min.


² Notes on procedures: The halogenation reaction (step 2) was conducted using N-chlorosuccinimide in DCM solvent. The Cbz removal (step 3) was conducted using KOH/MeOH at 80 °C for 3 h. The mixture of chloro regioisomers and enantiomers (4 compounds) was separated after Step 4 using chiral HPLC: Column: XBridge BEH C18 OBD Prep Column, 130 Å, 5 μm, 19 × 150 mm; Mobile phase: water (10 mmol NH₄HCO₃) and ACN (31% ACN up to 53% in 7 min).

Example 14-1. N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide;

Example 14-2. N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide;

Example 14-3. N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 14-4. N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Mixture of racemic tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-1-bromo-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and racemic tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-3-bromo-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00414] Into a 250-mL round-bottom flask was added tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (1.2 g, 2.44 mmol) and DCM (96 mL). This was followed by the addition of a solution of NBS (478 mg, 2.69 mmol) in DCM (24 mL) dropwise with stirring at 0 °C. The resulting solution was stirred for 2 h at 20 °C, then was quenched by the addition of water (20 mL).

The resulting solution was extracted with DCM (3 x 30 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to afford a residue that was purified by silica gel chromatography eluting with ethyl acetate/pet. ether (PE/EA=100:1 to 5:1) to afford a mixture of tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-3-bromo-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-1-bromo-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 572 [M+H]⁺.

Step 2. Mixture of racemic tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-3-cyano-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00415] Into a 20-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was added a mixture of tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-3-bromo-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-1-bromo-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (460 mg, 0.81 mmol), Zn(CN)₂ (93 mg, 0.80 mmol), Pd(PPh₃)₄ (93 mg, 0.08 mmol), and N,N-dimethylformamide (9 mL). The resulting mixture was irradiated with microwave for 1 h at 120 °C. The reaction was cooled to room temperature and quenched by the addition of 20 mL of water. The resulting solution was extracted with ethyl acetate (3x15 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to afford a residue that was purified by silica gel chromatography and eluted with 100:1 to 2:1 ethyl acetate/pet. ether to afford a mixture of tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-3-cyano-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as yellow oil. MS: (ESI, m/z): 517 [M+H]⁺.

Step 3. Mixture of racemic tert-butyl 3-(6-amino-3-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and racemic tert-butyl 3-(6-amino-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00416] Into a 50-mL round-bottom flask purged and maintained with nitrogen, was added a mixture of tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-3-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (300 mg, 0.58 mmol), ethanol (20 mL), and Palladium carbon (300 mg). The resulting suspension was stirred for 1 h at 20 °C under hydrogen atmosphere. The solids were removed by filtration over celite and the filtrate was concentrated under vacuum to afford a crude residue that was purified by silica gel chromatography and eluted with DCM/methanol (DCM/MeOH=10:1) to afford a mixture of tert-butyl 3-(6-amino-3-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-amino-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as yellow oil. MS: (ESI, m/z): 383 [M+H]⁺.

Step 4. tert-Butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (Peak A, enantiomer 1);

tert-butyl 3-((S)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (Peak A, enantiomer 2);

tert-butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (Peak B, enantiomer 1);

tert-butyl 3-((S)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (Peak B, enantiomer 2).

[00417] Into a 100-mL round-bottom flask was placed the mixture of tert-butyl 3-(6-amino-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-amino-3-cyano-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (150 mg, 0.39 mmol), EDCI (90 mg, 0.47 mmol), HOBt (63 mg, 0.47 mmol), DIEA (152 mg, 1.18 mmol), N,N-dimethylformamide (10 mL), and 3-amino-6-methylthieno[2,3-

b]pyridine-2-carboxylic acid (98 mg, 0.47 mmol). The resulting solution was stirred for 2 h at 20 °C, then was quenched by the addition of 20 mL of water. The solids were collected by filtration, and the solids were purified by prep-HPLC (Column: XBridge BEH C18 OBD, 5 µm, 19 × 150 mm; Mobile phase: water (10 mM NH₄HCO₃), ACN (55% ACN up to 70% over 7 min)).

[00418] This mixture was then further purified by chiral HPLC (Column: Chiralpak OD-H, 5 µm, 20 × 250 mm; Mobile phase A: hexanes, B: EtOH, 50% B). This resulted in a single regioisomer peak A: (RT=8.45 min) and single regioisomer peak B: (RT=12.91 min).

[00419] The peak A was further purified by chiral HPLC (Column: Chiralpak IC, 5 µm, 20 × 250 mm; Mobile phase A: hexanes, B: EtOH, 50% B). This resulted in peak A, enantiomer 1: (RT=15.55 min) as a yellow solid and peak A, enantiomer 2: (RT=21.10 min) as a yellow solid. Stereochemistry of the separated peak A enantiomers were arbitrarily assigned. MS (for both enantiomers): (ESI, m/z): 573 [M+H]⁺.

[00420] The peak B was further purified by chiral HPLC (Column: Chiralpak IB, 5 µm, 20 × 250 mm; Mobile phase A: hexanes, B: EtOH, 50% B). This resulted in peak B, enantiomer 1: (RT=7.65 min) as a yellow solid and peak B, enantiomer 2: (RT=9.22 min) as a yellow solid. Stereochemistry of the separated peak B enantiomers were arbitrarily assigned. MS (for both enantiomers): (ESI, m/z): 573 [M+H]⁺.

Step 5. N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide;

N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide;

N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide



Representative procedure and spectra for the first isomer:






[00421] Into a 50-mL round-bottom flask, was added tert-butyl 3-[(6R)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-1-cyano-5,6,7,8-tetrahydronaphthalen-2-yl]-3,8-






diazabicyclo[3.2.1]octane-8-carboxylate (30 mg, 0.05 mmol), DCM (5 mL), and TFA (1 mL). The resulting solution was stirred for 1 h at 20 °C, then was concentrated under vacuum to afford a crude residue that was diluted with 5 mL of DCM. The pH of the solution was adjusted to 8 with NH₃ in MeOH (7 M), then the resulting mixture was concentrated under vacuum to a residue that was purified by prep-HPLC (Column: XBridge BEH C18 OBD Prep Column, 130 Å, 5 µm, 19 × 150 mm; Mobile phase: water (10 mmol NH₄HCO₃), ACN (35% ACN up to 56% over 7 min)) to afford N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as a light yellow solid. ¹H NMR (400 MHz, MeOH-d₄) δ(ppm): 8.20 (d, *J* = 8.3 Hz, 1H), 7.29-7.30 (m, 2H), 6.94 (d, *J* = 8.5 Hz, 1H), 4.21-4.24 (m, 1H), 3.55 (s, 2H), 3.20-3.27 (m, 2H), 2.91-3.19 (m, 5H), 2.79-2.80 (m, 1H), 2.64 (s, 3H), 2.18-2.20 (m, 3H), 1.79-1.94 (m, 3H). MS: (ESI, *m/z*): 473 [M+H]⁺.






[00422] The following examples in Table 18 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Example 14-1.






Table 18


Example Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺	¹ H NMR
14-2	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	473	(CD ₃ OD, 400 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.4 Hz, 1H), 7.31-7.28 (m, 2H), 6.94 (d, <i>J</i> = 8.8 Hz, 1H), 4.27-4.20 (m, 1H), 3.56 (s, 2H), 3.31-3.27 (m, 2H), 3.16-2.94 (m, 5H), 2.84-2.77 (m, 1H), 2.64 (s, 3H), 2.21-2.19 (m, 3H), 1.92-1.82 (m, 3H).
14-3	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	473	(CD ₃ OD, 400 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.4 Hz, 1H), 7.36-7.29 (m, 2H), 6.88 (s, 1H), 4.27-4.21 (m, 1H), 3.57 (s, 2H), 3.33-3.30 (m, 2H), 3.02-2.95 (m, 5H), 2.81-2.75 (m, 1H), 2.64 (s, 3H), 2.18-2.10 (m, 3H), 1.86-1.79 (m, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
14-4	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	473	(CD ₃ OD, 400 MHz) δ (ppm): 8.20 (d, J = 8.4 Hz, 1H), 7.37-7.30 (m, 2H), 6.88 (s, 1H), 4.27-4.21 (m, 1H), 3.60 (s, 2H), 3.34-3.30 (m, 2H), 3.03-2.97 (m, 5H), 2.81-2.75 (m, 1H), 2.64 (s, 3H), 2.20-2.10 (m, 3H), 1.87-1.80 (m, 3H).
14-5	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	474	(CD ₃ OD, 300 MHz) δ (ppm): 8.54 (s, 1H), 7.33 (s, 1H), 6.85 (s, 1H), 4.28-4.18 (m, 1H), 3.61-3.51 (m, 2H), 3.35-3.31 (m, 2H), 3.05-2.92 (m, 5H), 2.83-2.69 (m, 1H), 2.65 (s, 3H), 2.17-2.09 (m, 3H), 1.98-1.70 (m, 3H).
14-6	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	474	(CD ₃ OD, 300 MHz) δ (ppm): 8.55 (s, 1H), 7.28 (d, J = 8.4 Hz, 1H), 6.94 (d, J = 8.4 Hz, 1H), 4.27-4.19 (m, 1H), 3.61-3.59 (m, 2H), 3.33-3.31 (m, 2H), 3.18-2.88 (m, 5H), 2.83-2.74 (m, 1H), 2.65 (s, 3H), 2.25-2.12 (m, 3H), 1.92-1.75 (m, 3H).
14-7	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	474	(CD ₃ OD, 400 MHz) δ (ppm): 8.58 (s, 1H), 7.31 (d, J = 8.4 Hz, 1H), 6.97 (d, J = 8.4 Hz, 1H), 4.35-4.18 (m, 1H), 3.62-3.59 (m, 2H), 3.35-3.32 (m, 2H), 3.15-2.96 (m, 5H), 2.86-2.77 (m, 1H), 2.68 (s, 3H), 2.28-2.18 (m, 3H), 1.96-1.82 (m, 3H).
14-8	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-cyano-1,2,3,4-tetrahydronaphthalen-2-yl)-7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxamide	474	(CD ₃ OD, 400 MHz) δ (ppm): 8.54 (s, 1H), 7.32 (s, 1H), 6.84 (s, 1H), 4.28-4.17 (m, 1H), 3.53-3.49 (m, 2H), 3.28-3.23 (m, 2H), 3.06-2.92 (m, 5H), 2.81-2.70 (m, 1H), 2.64 (s, 3H), 2.18-2.08 (m, 3H), 1.89-1.75 (m, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
14-9 ^{1,2}	 (R)-3-amino-N-(5-cyano-8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	465	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.67 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.18 (s, 2H), 6.86 (d, <i>J</i> = 11.6 Hz, 1H), 4.15 (s, 1H), 3.06-2.87(m, 11H), 2.67-2.58 (m, 4H), 2.08-1.81 (m, 2H), 1.23 (s, 1H), 1.16-1.12 (m, 1H), 0.86 (s, 1H).
14-10 ^{1,2}	 (S)-3-amino-N-(5-cyano-8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	465	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.67 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.23 (s, 2H), 6.87 (d, <i>J</i> = 11.2 Hz, 1H), 4.15 (s, 1H), 3.07-2.82 (m, 11H), 2.78-2.58 (m, 4H), 2.11-1.80 (m, 2H), 1.23 (s, 1H), 1.17-1.13 (m, 1H), 0.86 (s, 1H).
14-11 ^{1,3}	 (R)-7-amino-N-(5-cyano-8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	466	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.92 (br s, 1H), 6.95 (br s, 2H), 6.88 (d, <i>J</i> = 11.6 Hz, 1H), 4.30-4.10 (m, 1H), 3.10-2.80 (m, 11H), 2.62-2.55 (m, 4H), 2.15-2.00 (m, 1H), 1.90-1.75 (m, 1H).
14-12 ^{1,3}	 (S)-7-amino-N-(5-cyano-8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	466	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.91 (br s, 1H), 6.94 (br s, 2H), 6.88 (d, <i>J</i> = 11.6 Hz, 1H), 4.30-4.10 (m, 1H), 3.10-2.80 (m, 11H), 2.62-2.55 (m, 4H), 2.15-2.00 (m, 1H), 1.90-1.75 (m, 1H).
14-13 ^{1,4}	 (R)-6-amino-N-(5-cyano-8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-2-methylthieno[2,3-d]thiazole-5-carboxamide	471	(DMSO-d ₆ , 400 MHz) δ(ppm): 7.58 (br s, 1H), 7.10 (br s, 2H), 6.85 (d, <i>J</i> = 11.6 Hz, 1H), 4.20-4.05 (m, 1H), 3.04-3.01 (m, 11H), 2.84 (s, 3H), 2.67-2.55 (m, 1H), 2.04-2.03 (m, 1H), 1.81-1.77 (m, 1H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
14-14 ^{1,4}	 (S)-6-amino-N-(5-cyano-8-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-2-methylthieno[2,3-d]thiazole-5-carboxamide	471	(DMSO-d ₆ , 400 MHz) δ(ppm): 7.58 (br s, 1H), 7.10 (br s, 2H), 6.84 (d, <i>J</i> = 11.6 Hz, 1H), 4.20-4.05 (m, 1H), 3.04-3.01 (m, 11H), 2.84 (s, 3H), 2.67-2.55 (m, 1H), 2.04-2.03 (m, 1H), 1.81-1.77 (m, 1H).
14-15 ^{1,5}	 3-amino-N-[(2S)-5-cyano-8-fluoro-6-{9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl}-1,2,3,4-tetrahydronaphthalen-2-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	507	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.69 (d, <i>J</i> = 7.2 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.19 (br s, 2H), 6.95 (d, <i>J</i> = 11.6 Hz, 1H), 4.16-4.14 (m, 1H), 3.71-3.69 (m, 2H), 3.56-3.53 (m, 2H), 3.33-3.30 (m, 2H), 3.15-2.95 (m, 7H), 2.67-2.60 (m, 1H), 2.58 (s, 3H), 2.09-2.05 (m, 1H), 1.88-1.82 (m, 1H).
14-16 ^{1,5}	 3-amino-N-[(2R)-5-cyano-8-fluoro-6-{9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl}-1,2,3,4-tetrahydronaphthalen-2-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	507	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.69 (d, <i>J</i> = 7.8 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.19 (br s, 2H), 6.95 (d, <i>J</i> = 11.6 Hz, 1H), 4.16-4.14 (m, 1H), 3.71-3.69 (m, 2H), 3.53-3.50 (m, 2H), 3.31-3.27 (m, 2H), 3.16-2.95 (m, 7H), 2.67-2.60 (m, 1H), 2.58 (s, 3H), 2.08-2.05 (m, 1H), 1.88-1.82 (m, 1H).
14-17 ^{1,6}	 3-amino-N-[(2S)-5-cyano-6-{3,8-diazabicyclo[3.2.1]octan-3-yl}-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	491	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.30 (d, <i>J</i> = 8.1 Hz, 1H), 7.66 (d, <i>J</i> = 7.5 Hz, 1H), 7.31 (d, <i>J</i> = 8.1 Hz, 1H), 7.18 (br s, 2H), 6.78 (d, <i>J</i> = 12.0 Hz, 1H), 4.15-4.14 (m, 1H), 3.44-3.42 (m, 2H), 3.33-3.25 (m, 2H), 3.05-2.82 (m, 5H), 2.61-2.55 (m, 4H), 2.07-2.03 (m, 1H), 1.94-1.75 (m, 3H), 1.66-1.63 (m, 2H).
14-18 ^{1,6}	 3-amino-N-[(2R)-5-cyano-6-{3,8-diazabicyclo[3.2.1]octan-3-yl}-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	491	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.30 (d, <i>J</i> = 8.1 Hz, 1H), 7.66 (d, <i>J</i> = 7.5 Hz, 1H), 7.31 (d, <i>J</i> = 8.1 Hz, 1H), 7.18 (br s, 2H), 6.78 (d, <i>J</i> = 11.7 Hz, 1H), 4.15-4.14 (m, 1H), 3.45-3.43 (m, 2H), 3.33-3.25 (m, 2H), 3.05-2.82 (m, 5H), 2.61-2.55 (m, 4H), 2.37 (br s, 1H), 2.07-2.03 (m, 1H), 1.94-1.75 (m, 3H), 1.66-1.63 (m, 2H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
14-19 ^{1,7}	 7-amino-N-[(2S)-5-cyano-6-{3,8-diazabicyclo[3.2.1]octan-3-yl}-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide	492	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.65 (s, 1H), 7.90 (br s, 1H), 6.94 (br s, 2H), 6.79 (d, <i>J</i> = 11.7 Hz, 1H), 4.18-4.14 (m, 1H), 3.46-3.44 (m, 2H), 3.32-3.23 (m, 2H), 3.05-2.82 (m, 6H), 2.65 (s, 3H), 2.64-2.56 (m, 1H), 2.08-1.82 (m, 4H), 1.66-1.64 (m, 2H).
14-20 ^{1,7}	 7-amino-N-[(2R)-5-cyano-6-{3,8-diazabicyclo[3.2.1]octan-3-yl}-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide	492	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.65 (s, 1H), 7.90 (br s, 1H), 6.94 (br s, 2H), 6.79 (d, <i>J</i> = 12.0 Hz, 1H), 4.18-4.15 (m, 1H), 3.44-3.42 (m, 2H), 3.32-3.22 (m, 2H), 3.05-2.82 (m, 6H), 2.65 (s, 3H), 2.64-2.56 (m, 1H), 2.08-1.82 (m, 4H), 1.68-1.64 (m, 2H).
14-21 ^{1,8}	 6-amino-N-[(2S)-5-cyano-6-{3,8-diazabicyclo[3.2.1]octan-3-yl}-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-2-methylthieno[2,3-d][1,3]thiazole-5-carboxamide	497	(DMSO-d ₆ , 400 MHz) δ(ppm): 7.57 (br s, 1H), 7.11 (br s, 2H), 6.79 (d, <i>J</i> = 12.0 Hz, 1H), 4.11 (br s, 1H), 3.45-3.43 (m, 2H), 3.27-3.22 (m, 2H), 3.04-2.82 (m, 5H), 2.79 (s, 3H), 2.61-2.52 (m, 1H), 2.05-1.96 (m, 1H), 1.94-1.86 (m, 2H), 1.84-1.76 (m, 1H), 1.66-1.63 (m, 2H).
14-22 ^{1,8}	 6-amino-N-[(2R)-5-cyano-6-{3,8-diazabicyclo[3.2.1]octan-3-yl}-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-2-methylthieno[2,3-d][1,3]thiazole-5-carboxamide	497	(DMSO-d ₆ , 400 MHz) δ(ppm): 7.59 (br s, 1H), 7.11 (br s, 2H), 6.79 (d, <i>J</i> = 12.0 Hz, 1H), 4.13-4.09 (m, 1H), 3.45-3.44 (m, 2H), 3.27-3.22 (m, 2H), 3.03-2.84 (m, 5H), 2.79 (s, 3H), 2.60-2.51 (m, 1H), 2.04-1.95 (m, 1H), 1.93-1.90 (m, 2H), 1.81-1.70 (m, 1H), 1.65-1.60 (m, 2H).
12-23 ^{1,9,10}	 3-amino-N-[(2S)-6-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5-cyano-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	495	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.63 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.17 (br s, 2H), 6.48 (d, <i>J</i> = 12.4 Hz, 1H), 4.14-4.08 (m, 1H), 3.84-3.81 (m, 1H), 3.73-3.65 (m, 2H), 3.47-3.41 (m, 1H), 3.38-3.32 (m, 1H), 3.30 (s, 3H), 3.27-3.23 (m, 1H), 3.05-3.00 (m, 1H), 2.92-2.82 (m, 2H), 2.58 (s, 3H), 2.55-2.51 (m, 1H), 2.07-2.02 (m, 1H), 1.85-1.73 (m, 2H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
12-24 ^{1,9,10}	 3-amino-N-[(2R)-6-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5-cyano-8-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	495	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.64 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.17 (br s, 2H), 6.48 (d, <i>J</i> = 12.8 Hz, 1H), 4.14-4.08 (m, 1H), 3.86-3.83 (m, 1H), 3.73-3.66 (m, 2H), 3.45-3.42 (m, 1H), 3.38-3.35 (m, 1H), 3.30 (s, 3H), 3.27-3.23 (m, 1H), 3.05-2.75 (m, 3H), 2.58 (s, 3H), 2.07-2.01 (m, 1H), 1.85-1.73 (m, 2H).

¹ **Notes on procedures:** In Step 1, only one bromination regioisomer was observed, tert-butyl 4-(6-(((benzyloxy)carbonyl)amino)-1-bromo-4-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate.

² **Notes on procedures:** Chiral separation of the enantiomers was performed after Step 5 rather than in Step 4, using the chiral column Chiralpak IA and Mobile Phase 50% EtOH/hexanes (0.1% Et₂NH). Stereochemistry of the separated enantiomers were arbitrarily assigned.

³ **Notes on procedures:** In Step 4, the amide coupling reaction was performed with HBTU and Et₃N in DMA. Chiral separation of the enantiomers was performed after Step 5 rather than in Step 4, using the chiral column Chiralpak IE and Mobile Phase 20% IPA/MTBE (0.1% Et₂NH). Stereochemistry of the separated enantiomers were arbitrarily assigned.

⁴ **Notes on procedures:** In Step 4, the carboxylic acid **Intermediate 24** was used. The amide coupling reaction was performed with HBTU and Et₃N in DMA. Chiral separation of the enantiomers was performed after Step 5 rather than in Step 4, using the chiral column Chiralpak IA and Mobile Phase 40% EtOH/hexanes-DCM (5:1) (0.1% Et₂NH). Stereochemistry of the separated enantiomers were arbitrarily assigned.

⁵ **Notes on procedures:** In Step 4, the amide coupling reaction was performed with HBTU and Et₃N in DMA. Chiral separation of the enantiomers was performed after Step 5 rather than in Step 4, using the chiral column Chiralpak IG and Mobile Phase 30% EtOH/MTBE (containing 8 mM NH₃ in MeOH). Stereochemistry of the separated enantiomers were arbitrarily assigned.

⁶ **Notes on procedures:** In Step 4, the amide coupling reaction was performed with HBTU and Et₃N in DMA. The enantiomers were separated by chiral HPLC using the chiral column Chiralpak IC and Mobile Phase 15% EtOH/MTBE. Stereochemistry of the separated enantiomers were arbitrarily assigned.

⁷ **Notes on procedures:** In Step 4, the amide coupling reaction was performed with HBTU and Et₃N in DMA. The enantiomers were separated by chiral HPLC using the chiral column Chiralpak IG and Mobile Phase 30% EtOH/MTBE. Stereochemistry of the separated enantiomers were arbitrarily assigned.


⁸ **Notes on procedures:** In Step 4, the carboxylic acid **Intermediate 24** was used. The amide coupling reaction was performed with HBTU and Et₃N in DMA. The enantiomers were separated by chiral HPLC using the chiral column Chiralpak IG and Mobile Phase 30% EtOH/MTBE. Stereochemistry of the separated enantiomers were arbitrarily assigned.

⁹ **Notes on procedures:** In Step 2, cyanation was achieved in the presence of Zn (2 eq), Zn(CN)₂ (10 eq), Pd(P(t-Bu)₃)₂ (0.5 eq), and 4,4'-di-tert-butyl-2,2'-bipyridine (1 eq) in DMA, with heating at 97 °C for 2 h.

¹⁰ **Notes on procedures:** . In Step 4, the amide coupling reaction was performed with HBTU and Et₃N in DMA. The enantiomers were separated by chiral HPLC using the chiral column Chiralpak ID-2 and Mobile Phase 15% EtOH/MTBE (0.1% Et₂NH). Stereochemistry of the separated enantiomers were arbitrarily assigned.

Example 15-1. (R)-3-Amino-6-methyl-N-(3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl)thieno[2,3-b]pyridine-2-carboxamide and

Example 15-2. (S)-3-Amino-6-methyl-N-(3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl)thieno[2,3-b]pyridine-2-carboxamide (stereoconfiguration assumed)



Step 1. Benzyl (3-nitro-5,6,7,8-tetrahydroquinolin-7-yl)carbamate

[00423] A mixture of benzyl (3-oxocyclohexyl)carbamate (247 mg, 0.999 mmol) and 1-methyl-3,5-dinitropyridin-2(1H)-one (300 mg, 1.507 mmol) in ammonia/MeOH (1M, 6 mL) was microwaved at 90°C for 30 min to provide a blackish-red solution. Eleven more samples of the same composition were run under the same conditions. The twelve samples were combined and evaporated under reduced pressure. The residue was partitioned between DCM (300 mL) and saturated aqueous NaHCO_3 (100 mL). The organic layer was washed with brine (100 mL), dried (MgSO_4), filtered, treated with silica gel, and evaporated under reduced pressure. Purification by silica gel chromatography (0 to 50% EtOAc in hexanes) provided a mixture of benzyl (3-nitro-5,6,7,8-tetrahydroquinolin-7-yl)carbamate and undesired regioisomer benzyl (3-nitro-5,6,7,8-

tetrahydroquinolin-5-yl)carbamate as a pale yellow gum. The mixture was used without further purification. MS (ESI, m/z): 213 $[M+H]^+$.

Step 2. Benzyl (3-amino-5,6,7,8-tetrahydroquinolin-7-yl)carbamate

[00424] A solution of benzyl (3-nitro-5,6,7,8-tetrahydroquinolin-7-yl)carbamate and regioisomer (2.64 g, 8.07 mmol) in ethanol (110 mL) was treated with tin(II) chloride dihydrate (9.10 g, 40.3 mmol) and conc. HCl (1 mL), then stirred at 80 °C for 75 min. The solution was allowed to cool, then was concentrated under reduced pressure to about 15 mL. The sample was made basic (pH ~9) by addition of saturated aqueous NaHCO_3 ; about midway through, the sample was diluted with DCM (100 mL). The mixture was triturated, then filtered, and the filter cake was washed with DCM (50 mL). The filter cake was triturated and sonicated in MeOH (200 mL), then the mixture was filtered. The slightly cloudy filtrate was filtered through Celite 545, then concentrated to provide a yellow solid. Purification by silica gel chromatography (0 to 15% MeOH in DCM) provided a mixture of benzyl (3-amino-5,6,7,8-tetrahydroquinolin-7-yl)carbamate and regioisomer as a yellow solid. The mixture was used without further purification. MS (ESI, m/z): 298 $[M+H]^+$.

Step 3. Benzyl (3-iodo-5,6,7,8-tetrahydroquinolin-7-yl)carbamate

[00425] A sample of benzyl (3-amino-5,6,7,8-tetrahydroquinolin-7-yl)carbamate and regioisomer (1.25 g, 4.22 mmol) was treated with ACN (17 mL). The material was cooled on a dry ice / ethylene glycol bath (approximately -10 to -15 °C) and PTSA monohydrate (1.61 g, 8.45 mmol) was added. After stirring 10 min, a solution of potassium iodide (1.05 g, 6.33 mmol) and sodium nitrite (0.436 g, 6.32 mmol) in water (3.4 mL) was added dropwise over 20 min. The dry ice bath was replaced with an ice bath, and the reddish-brown mixture was stirred at 0 °C for 3 h 40 min. The solution was diluted with EtOAc (50 mL), cooled on an ice bath and treated with saturated aqueous NaHCO_3 until basic (pH ~8-9). The organic layer was removed and the aqueous layer was extracted once more with EtOAc (50 mL). The combined organics were washed sequentially with water and brine (50 mL each), dried (Na_2SO_4), filtered, treated with silica gel, and evaporated under reduced pressure. Purification by silica gel chromatography (0 to 23% EtOAc in hexanes) provided a mixture of benzyl (3-iodo-5,6,7,8-tetrahydroquinolin-7-yl)carbamate and regioisomer as a yellow solid. The mixture was used without further purification. MS (ESI, m/z): 409 $[M+H]^+$.

Step 4. tert-Butyl 4-(7-(((benzyloxy)carbonyl)amino)-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate

[00426] A mixture of t-BuXPhos Pd G4 (76.3 mg, 94.4 μ mole), benzyl (3-iodo-5,6,7,8-tetrahydroquinolin-7-yl)carbamate and regioisomer (605.0 mg, 1.482 mmol), tert-butyl piperazine-1-carboxylate (553.5 mg, 2.97 mmol), and sodium tert-butoxide (257.1 mg, 2.68 mmol) was sealed in a 40-mL vial. The atmosphere was evacuated and replaced with nitrogen, three times. Dioxane (15 mL) was added and the solution was stirred at ambient temperature for three days. Material at the same stage from a previous run (from 40.7 mg, 0.100 mmol iodo starting material) was added. The mixture was diluted with EtOAc (~100 mL), treated with silica gel, and evaporated under reduced pressure. Purification by silica gel chromatography (0 to 87% EtOAc in hexanes) provided a mixture of tert-butyl 4-(7-(((benzyloxy)carbonyl)amino)-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate and regioisomer as a yellow foam. The mixture was used without further purification. MS (ESI, m/z): 467 [M+H]⁺.

Step 5. tert-Butyl 4-(7-amino-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate

[00427] A suspension of Pd/C (10%) (57.1 mg, 0.054 mmol) and the mixture of tert-butyl 4-(7-(((benzyloxy)carbonyl)amino)-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate and regioisomer (485 mg, 1.039 mmol) in MeOH (5 mL) was stirred at ambient temperature under an atmosphere of hydrogen overnight. The sample was filtered through Celite 545 under vacuum, and the Celite was rinsed with MeOH. The filtrate was evaporated to provide a mixture of tert-butyl 4-(7-amino-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate and regioisomer as an orange gum. The mixture was used without further purification. MS (ESI, m/z): 333 [M+H]⁺.

Step 6. tert-Butyl 4-(7-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate

[00428] A solution of tert-butyl 4-(7-amino-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate and regioisomer (130 mg, 0.391 mmol) in DCM (6 mL) was treated with 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (74 mg, 0.355 mmol), BOP (188.4 mg, 0.426 mmol), and DIEA (186.4 μ L, 1.067 mmol). The solution was stirred at ambient temperature for 4.5 h. The mixture was diluted with EtOAc (40 mL) and washed sequentially with 2M aqueous Na₂CO₃ (20 mL) and brine (2x40 mL). Heptane (~15 mL) was added, then the mixture was evaporated under reduced pressure to dryness. Purification by silica gel chromatography (0 to

100% EtOAc in hexanes) provided a mixture of tert-butyl 4-(7-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate and regioisomer as a yellow solid. MS (ESI, m/z): 523 [M+H]⁺.

Step 7. (R)-3-Amino-6-methyl-N-(3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl)thieno[2,3-b]pyridine-2-carboxamide and (S)-3-Amino-6-methyl-N-(3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl)thieno[2,3-b]pyridine-2-carboxamide

Regioisomer purification:


[00429] A solution of the mixture of tert-butyl 4-(7-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5,6,7,8-tetrahydroquinolin-3-yl)piperazine-1-carboxylate and regioisomer (97.3 mg, 0.186 mmol) in DCM (5 mL) and MeOH (1 mL) was treated with HCl (4 M in dioxane; 1.5 mL, 6.00 mmol), and allowed to stand at ambient temperature for 30 min; within 5 min a yellow precipitate began to form. The mixture was concentrated under reduced pressure. The residue was triturated in Et₂O (10 mL), then the solids were collected and washed with Et₂O (20 mL) to provide a mixture of 3-amino-6-methyl-N-(3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl)thieno[2,3-b]pyridine-2-carboxamide and regioisomer as a yellow solid. The material was purified by prep-HPLC (Column: Chiralpak IB, 250 x 4.6 mm, 5 μm; Mobile phase A: hexanes, B: Ethanol/Et₂NH; Gradient 60% B; Total Flow: 5 mL/min; Detector: 190 nm to 500 nm) to provided 3-amino-6-methyl-N-(3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl)thieno[2,3-b]pyridine-2-carboxamide as the major regioisomer peak which was then carried on to chiral separation.

Chiral Purification:

The crude product of 3-amino-6-methyl-N-[3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide (20 mg, 0.04 mmol, 90%) was separated into its enantiomers by chiral HPLC (Column: Chiralpak IC, 2 x 25cm, 5μm; Mobile phase: hexanes, 0.1% Et₂NH and ethanol (hold 50% ethanol for 30 min)). The first eluting isomer (RT = 23.0 min) was isolated as a pale yellow solid, stereochemistry arbitrarily assigned as (R)-3-amino-6-methyl-N-(3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl)thieno[2,3-b]pyridine-2-carboxamide. ¹H NMR (300 MHz, DMSO-d₆) δ(ppm): 8.30 (d, J = 8.1Hz, 1H), 8.06-8.05 (br s, 1H), 7.61 (d, J = 7.5Hz, 1H), 7.31 (d, J = 8.4Hz, 1H), 7.16-7.15 (br s, 2H), 7.02 (d, J = 2.4Hz, 1H), 4.25-4.10 (m, 1H), 3.05-3.02 (m, 5H), 2.93-2.71 (m, 8H), 2.58 (s, 3H), 2.05-1.96 (m, 1H), 1.79-1.65 (m, 1H). MS: (ESI, m/z): 423 [M+H]⁺.

The second eluting isomer (RT = 27.5 min) was isolated as a pale yellow solid, stereochemistry arbitrarily assigned as (S)-3-amino-6-methyl-N-(3-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-7-yl)thieno[2,3-b]pyridine-2-carboxamide. ¹H NMR (300 MHz, DMSO-d₆) δ(ppm): 8.30 (d, J = 8.1Hz, 1H), 8.06-8.05 (br s, 1H), 7.61 (d, J = 7.5Hz, 1H), 7.32 (d, J = 8.4Hz, 1H), 7.16-7.15 (br s, 2H), 7.02 (d, J = 2.4Hz, 1H), 4.25-4.10 (m, 1H), 3.05-3.02 (m, 5H), 2.93-2.71 (m, 8H), 2.58 (s, 3H), 2.05-1.96 (m, 1H), 1.79-1.65 (m, 1H). MS: (ESI, m/z): 423 [M+H]⁺.

Example 16. (R)-3-Amino-6-methyl-N-(7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)thieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)piperazine-1-carboxylate

[00430] A solution of benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate (**Intermediate 42**) (200 mg, 0.60 mmol), RuPhos (28 mg, 0.06 mmol), tert-butyl piperazine-1-carboxylate (175 mg, 0.94 mmol), RuPhos Pd G3 (50 mg, 0.06 mmol) and Cs₂CO₃ (410 mg, 1.26 mmol) in toluene (5 mL) was stirred for 16 h at 95 °C. After cooling to 25 °C, the reaction was then quenched by the addition of water (10 mL). The resulting mixture was extracted with DCM (3 x 10 mL) and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:2 ethyl acetate/pet. ether) to afford tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)piperazine-1-carboxylate as a white solid. MS: (ESI, m/z): 469 [M+H]⁺.

Step 2. tert-Butyl (R)-4-(3-amino-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)piperazine-1-carboxylate

[00431] A mixture of tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)piperazine-1-carboxylate (100 mg, 0.20 mmol) and Palladium on

carbon (20 mg, 10%) in ethyl acetate (10 mL) was stirred for 2 h at 25 °C under an atm of hydrogen. The solid was filtered out and the cake was washed with methanol (3x10 mL). The filtrate was concentrated under vacuum to give tert-butyl (R)-4-(3-amino-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)piperazine-1-carboxylate as a white solid. MS: (ESI, m/z): 335 [M+H]⁺.


Step 3. tert-Butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)piperazine-1-carboxylate

[00432] A solution of tert-butyl 4-[(3R)-3-amino-2H,3H,4H-pyrano[3,2-c]pyridin-7-yl]piperazine-1-carboxylate (65 mg, 0.18 mmol), HBTU (108 mg, 0.28 mmol), Et₃N (85 mg, 0.84 mmol) and 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (49 mg, 0.22 mmol) in DMA (3 mL) was stirred for 30 min at 25 °C. The reaction was then quenched by the addition of water (10 mL). The resulting mixture was extracted with DCM (2 x 10 mL) and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)piperazine-1-carboxylate as a colorless oil. MS: (ESI, m/z): 525 [M+H]⁺.

Step 4. (R)-3-Amino-6-methyl-N-(7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)thieno[2,3-b]pyridine-2-carboxamide

[00433] A solution of tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)piperazine-1-carboxylate (45 mg, 0.08 mmol) and TFA (1 mL) in DCM (3 mL) was stirred for 30 min at 25 °C and concentrated under vacuum. The residue was purified by prep-HPLC (Column: X Bridge C18, 19x150 mm, 5 μm; Mobile Phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 10% B to 80% B over 10 min) to afford (R)-3-amino-6-methyl-N-(7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)thieno[2,3-b]pyridine-2-carboxamide as a white solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ(ppm): 8.33 (d, *J* = 8.4 Hz, 1H), 7.83 (s, 1H), 7.59 (br s, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.23 (br s, 2H), 6.14 (s, 1H), 4.27- 4.20 (m, 2H), 3.92-3.86 (m, 1H), 3.48-3.35 (m, 1H), 3.33-3.22 (m, 4H), 2.84-2.81 (m, 2H), 2.76-2.72 (m, 4H), 2.58 (s, 3H). MS: (ESI, m/z): 425 [M+H]⁺.

Example 17. N-((3R)-7-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl (R)-benzyl(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate

[00434] A solution of benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate (**Intermediate 42**) (250 mg, 0.75 mmol), sodium hydride (63 mg, 1.58 mmol, 60%) and benzyl bromide (268 mg, 1.57 mmol) in DMF (5 mL) was stirred for 2 h at 25 °C. The reaction was then quenched by the addition of 15 mL of water. The resulting mixture was extracted with 3 x 15 mL of ethyl acetate and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to give benzyl (R)-benzyl(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate as a light yellow solid. MS: (ESI, m/z): 409, 411 $[M+H]^+$.

Step 2. (R)-3-(Benzyl((benzyloxy)carbonyl)amino)-7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide

[00435] A solution of benzyl (R)-benzyl(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate (300 mg, 0.70 mmol) and m-CPBA (633 mg, 3.67 mmol) in DCM (10 mL) was stirred for 16 h at 25 °C. The reaction was then quenched by the addition of 10 mL of $\text{Na}_2\text{S}_2\text{O}_3$ (sat.) and 10 mL of NaHCO_3 (sat.). The resulting mixture was extracted with 3 x 20 mL of DCM and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and

concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 10:1 DCM/methanol) to give (R)-3-(benzyl((benzyloxy)carbonyl)amino)-7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide as a light yellow solid. MS: (ESI, m/z): 425, 427 [M+H]⁺.

Step 3. (3R)-3-(Benzyl((benzyloxy)carbonyl)amino)-7-(8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide

[00436] A solution of (R)-3-(benzyl((benzyloxy)carbonyl)amino)-7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide (498 mg, 2.35 mmol) and diisopropylethylamine (303 mg, 2.34 mmol) in tert-butanol (8 mL) was stirred for 16 h at 90 °C. After cooling to 25 °C, the resulting mixture was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 10:1 DCM/methanol) to give (3R)-3-(benzyl((benzyloxy)carbonyl)amino)-7-(8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide as a yellow solid. MS: (ESI, m/z): 601 [M+H]⁺.

Step 4. (3R)-3-(Benzyl((benzyloxy)carbonyl)amino)-7-(8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide

[00437] To a solution of (3R)-3-(benzyl((benzyloxy)carbonyl)amino)-7-(8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide (100 mg, 0.16 mmol) in DCM (10 mL) and a solution of trimethylamine in THF (1M, 0.83 mL, 0.83 mmol) was added TFAA (105 mg, 0.50 mmol) at 0 °C. The resulting solution was stirred for 1 h at 25 °C and concentrated under vacuum. The residue was purified by silica gel column (eluting with 5:1 DCM/methanol) to give the intermediate (crude): ((3R)-3-(benzyl((benzyloxy)carbonyl)amino)-7-(8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)-N,N,N-trimethyl-3,4-dihydro-2H-pyrano[3,2-c]pyridin-5-aminium). The intermediate was dissolved in DMF (4 mL). TBAF (83 mg, 0.32 mmol) was added. The resulting solution was stirred for 30 min at 90 °C. After cooling to 25 °C, the reaction was then quenched by the addition of 10 mL of water. The resulting mixture was extracted with 3 x 10 mL of ethyl acetate and the organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:2 ethyl acetate/pet. ether) to give (3R)-3-(benzyl((benzyloxy)carbonyl)amino)-7-(8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide as a yellow solid. MS: (ESI, m/z): 619 [M+H]⁺.

Step 5. tert-Butyl 3-((R)-3-amino-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00438] A mixture of (3R)-3-(benzyl((benzyloxy)carbonyl)amino)-7-(8-(tert-butoxycarbonyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridine 6-oxide (58 mg, 0.09 mmol) and palladium on carbon (50 mg, 5%) in ethyl acetate (5 mL) was stirred for 2 h at 25 °C under hydrogen atmosphere. The solids were filtered out. The filtrate was concentrated under vacuum to give tert-butyl 3-((R)-3-amino-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a light yellow solid. MS: (ESI, m/z): 379 [M+H]⁺.

Step 6. tert-Butyl 3-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate


[00439] A solution of tert-butyl 3-((R)-3-amino-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (35 mg, 0.09 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (21 mg, 0.10 mmol), HBTU (57 mg, 0.15 mmol) and Et₃N (30 mg, 0.30 mmol) in DMA (3 mL) was stirred for 1 h at 25 °C. The reaction was then quenched by the addition of 10 mL of water. The resulting mixture was extracted with 3 x 15 mL of ethyl acetate and the organic layers were combined, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:2 ethyl acetate/pet. ether) to give tert-butyl 3-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 569 [M+H]⁺.

Step 7. N-((3R)-7-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

A solution of tert-butyl 3-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (25 mg, 0.04 mmol) and TFA (1 mL) in DCM (3 mL) was stirred for 1 h at 25 °C. The resulting mixture was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Shield RP18 OBD, 30 x 150 mm, 5 μm; Mobile phase A: Water (10 mM NH₄HCO₃), B: ACN; Gradient: 20% B to 47% B within 7 min). The collected fraction was concentrated under

vacuum to give N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ(ppm): 8.33 (d, *J* = 8.4 Hz, 1H), 7.62 (br s, 1H), 7.31 (d, *J* = 8.1 Hz, 1H), 7.23 (s, 2H), 5.92 (s, 1H), 4.28-4.19 (m, 2H), 3.99-3.93 (m, 1H), 3.63-3.59 (m, 2H), 3.47-3.45 (m, 2H), 2.79-2.65 (m, 5H), 2.58 (s, 3H), 1.65-1.53 (m, 4H). MS: (ESI, *m/z*): 469 [M+H]⁺.

Example 18. N-((3R)-7-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl 3-((R)-3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00440] A solution of benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)carbamate (**Intermediate 42**) (500 mg, 1.57 mmol), tert-butyl 3,8-diazabicyclo[3.2.1]octane-8-carboxylate (401 mg, 1.89 mol), RuPhos (88 mg, 0.19 mmol), RuPhos Pd G3 (129 mg, 0.15 mmol) and Cs₂CO₃ (1.03 g, 3.16 mmol) in toluene (20 mL) was stirred overnight at 95 °C. After cooling to 24 °C, the solids were filtered out and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to give tert-butyl 3-((R)-3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid. MS: (ESI, *m/z*): 495 [M+H]⁺.

Step 2. tert-Butyl 3-((R)-3-(((benzyloxy)carbonyl)amino)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00441] To a solution of tert-butyl 3-((R)-3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (500 mg, 1.01 mmol) in

DCM (25 mL) and methanol (25 mL) was added a solution of Selectfluor (358 mg, 1.01 mmol) in ACN (25 mL) at -10 °C. The resulting solution was stirred 2 h at -10 °C. The resulting mixture was concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.1% TFA), B: ACN; Flow rate: 50 mL/min; Gradient: 0% B to 80% B within 30 min). The collected fractions were concentrated under vacuum to give tert-butyl 3-((R)-3-(((benzyloxy)carbonyl)amino)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow oil. MS: (ESI, m/z): 513 [M+H]⁺.

Step 3. tert-Butyl 3-((R)-3-amino-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00442] A suspension of tert-butyl 3-((R)-3-(((benzyloxy)carbonyl)amino)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (100 mg, 0.20 mmol) and Palladium on carbon (20 mg, 10%) in ethyl acetate (5 mL) was stirred for 1.5 h at 24 °C under hydrogen atmosphere. The solids were filtered out. The filtrate was concentrated under vacuum to afford tert-butyl 3-((R)-3-amino-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow oil. MS: (ESI, m/z): 379 [M+H]⁺.


Step 4. tert-Butyl 3-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00443] A solution of tert-butyl 3-((R)-3-amino-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (80 mg, 0.13 mmol, purity 60%), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (47 mg, 0.23 mmol), Et₃N (64 mg, 0.63 mmol) and HBTU (96 mg, 0.25 mmol) in DMA (3 mL) was stirred for 30 min at 24 °C. The resulting solution was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Flow rate: 50 mL/min; Gradient: 0% B increasing to 75% B within 30 min). The collected fraction was concentrated under vacuum and to give tert-butyl 3-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow solid. MS: (ESI, m/z): 569 [M+H]⁺.

Step 5. N-((3R)-7-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

A solution of tert-butyl 3-((R)-3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-7-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (40 mg, 0.07 mmol) and TFA (1 mL) in DCM (3 mL) was stirred for 1 h at 24 °C. The resulting mixture was concentrated under vacuum. The mixture was purified by prep-HPLC (Column: XBridge Shield RP18 OBD, 5 μ m, 19x150 mm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN; Gradient: 10% B to 40% B within 2 min, increasing to 60% in 7 min)). The collected fraction was lyophilized to give N-((3R)-7-(3,8-diazabicyclo[3.2.1]octan-3-yl)-8-fluoro-3,4-dihydro-2H-pyrano[3,2-c]pyridin-3-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as a light yellow solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ (ppm): 8.33 (d, *J* = 8.1 Hz, 1H), 7.69 (s, 1H), 7.65 (br s, 1H), 7.31 (d, *J* = 8.1 Hz, 1H), 7.23 (br s, 2H), 4.32-7.29 (m, 2H), 4.05-3.95 (m, 1H), 3.60-3.48 (m, 2H), 3.48-3.40 (m, 2H), 2.97-2.91 (m, 4H), 2.58 (s, 3H), 1.72-1.68 (m, 4H). MS: (ESI, *m/z*): 469 [M+H]⁺.

Example 19. (R)-3-Amino-6-methyl-N-(7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)thieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate

[00444] A solution of benzyl (R)-4-(7-chloro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)carbamate (**Intermediate 43**) (200 mg, 0.56 mmol), tert-butyl piperazine-1-carboxylate (140 mg, 0.75 mmol), RuPhos Pd G3 (52 mg, 0.06 mmol), RuPhos (59 mg, 0.13 mmol) and Cs₂CO₃ (618 mg, 1.90 mmol) in toluene (5 mL) was stirred for 3 h at 100 °C. After cooling to 25 °C, the reaction was then quenched by the addition of 10 mL of water. The resulting mixture was extracted with 3 x 10 mL of ethyl acetate and the organic layers combined, dried over anhydrous sodium

sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to give tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate as an off-white solid. MS: (ESI, m/z): 469 [M+H]⁺.

Step 2. tert-Butyl (R)-4-(3-amino-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate

[00445] A suspension of tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate (270 mg, 0.52 mmol) and Palladium on carbon (30 mg, 10%) in ethyl acetate (20 mL) was stirred for 4 h at 25 °C. The solids were filtered out. The filtrate was concentrated under vacuum to tert-butyl (R)-4-(3-amino-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate as a brown oil. MS: (ESI, m/z): 335 [M+H]⁺.

Step 3. tert-Butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate


[00446] A solution of tert-butyl (R)-4-(3-amino-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate (50 mg, 0.13 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (34 mg, 0.16 mmol), HBTU (62 mg, 0.16 mmol) and Et₃N (45 mg, 0.44 mmol) in DMA (3 mL) was stirred for 30 min at 25 °C. The reaction was then quenched by the addition of 10 mL of water. The resulting mixture was extracted with 3 x 10 mL of ethyl acetate. The organic layers combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.1% formic acid), B: ACN; Flow rate: 50 mL/min; Gradient: 0% increasing to 100% B within 40 min). The collected fraction was concentrated under vacuum. The collected fraction was concentrated under vacuum to give tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 525 [M+H]⁺.

Step 4. (R)-3-Amino-6-methyl-N-(7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)thieno[2,3-b]pyridine-2-carboxamide

[00447] A solution of tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate (30 mg, 0.05 mmol) and TFA (1 mL) in DCM (3 mL) was stirred for 30 min at 25 °C. The resulting mixture

was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Shield RP18 OBD, 19 x 250 mm, 10 μ m; Mobile phase A: Water (10 mM NH_4HCO_3), B: ACN; Gradient: 20% B to 38% B in 8 min). The collected fraction was lyophilized to give (R)-3-amino-6-methyl-N-(7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)thieno[2,3-b]pyridine-2-carboxamide as an off-white solid. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ (ppm): 8.33 (d, $J = 8.4$ Hz, 1H), 7.55 (br s, 1H), 7.32-7.27 (m, 2H), 7.22 (br s, 2H), 6.37 (d, $J = 8.4$ Hz, 1H), 4.33-4.20 (m, 2H), 3.99-3.94 (m, 1H), 3.49-3.31 (m, 5H), 2.89-2.74 (m, 6H), 2.59 (s, 3H). MS: (ESI, m/z): 425 $[\text{M}+\text{H}]^+$.

Example 20. (R)-3-Amino-N-(6-fluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate

[00448] A solution of benzyl (R)-(7-chloro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)carbamate (**Intermediate 43**) (500 mg, 1.57 mmol), tert-butyl piperazine-1-carboxylate (352 mg, 1.19 mol), RuPhos Pd G3 (131 mg, 0.16 mmol), Ruphos (146 mg, 0.31 mmol) and Cs_2CO_3 (1.54 g, 4.72 mmol) in toluene (10 mL) was stirred for 3 h at 100 $^\circ\text{C}$. After cooling to 25 $^\circ\text{C}$, the reaction was quenched by the addition of 50 mL of water. Three batches were thus run in parallel and combined for extraction work-up with ethyl acetate (3 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to

afford tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 469 [M+H]⁺.

Step 2. tert-Butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate

[00449] To a solution of tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate (580 mg, 1.24 mmol) in DCM (25 mL) and MeOH (25 mL) was added Selectfluor (441 mg, 1.24 mmol) at -20 °C. The resulting mixture was stirred for 2 h at 25 °C. The solvent was removed under vacuum. The residue was diluted with water (30 mL). The resulting mixture was extracted with DCM (3 x 30 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:10-1:3 ethyl acetate/pet. ether) to afford tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate as an off-white solid. MS: (ESI, m/z): 487 [M+H]⁺.

Step 3. tert-Butyl (R)-4-(3-amino-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate

[00450] A suspension of tert-butyl (R)-4-(3-(((benzyloxy)carbonyl)amino)-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate (150 mg, 0.31 mmol) and Palladium on carbon (20 mg, 10%) in ethyl acetate (5 mL) was stirred for 2 h at 25 °C under hydrogen atmosphere. The solids were filtered out. The filtrate was concentrated under vacuum to give tert-butyl (R)-4-(3-amino-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate as a yellow oil. MS: (ESI, m/z): 353 [M+H]⁺.

Step 4. tert-Butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate


[00451] A solution of tert-butyl (R)-4-(3-amino-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate (90 mg, 190 μmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (56.0 mg, 270 μmol), HBTU (107 mg, 0.28 mmol) and Et₃N (77 mg, 0.76 mmol) in DMA (4 mL) was stirred for 1 h at 25 °C. The reaction was quenched with water (10 mL). The resulting mixture was extracted with ethyl acetate (3 x 10 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase

A: water (0.1% TFA), B: ACN; Flow rate: 50 mL/min; Gradient: 0% B to 70% B within 40 min). The collected fractions were concentrated under vacuum to afford tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 543 [M+H]⁺.

Step 5. (R)-3-Amino-N-(6-fluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00452] A solution of tert-butyl (R)-4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-6-fluoro-3,4-dihydro-2H-pyrano[2,3-b]pyridin-7-yl)piperazine-1-carboxylate (100 mg, 0.18 mmol) and TFA (1 mL) in DCM (3 mL) was stirred for 30 min at 25 °C. The resulting mixture was concentrated under vacuum. A solution of NH₃ in methanol (7M) (5 mL) was added. The resulting mixture was stirred for 30 min at 25 °C. The resulting mixture was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep OBD C18, 30 x 150 mm 5 µm; Mobile Phase A: Water (10 mM NH₄HCO₃), B: ACN; Flow rate: 60 mL/min; Gradient: 20% B to 45% B in 7 min) to afford (R)-3-amino-N-(6-fluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-pyrano[2,3-b]pyridin-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ(ppm): 8.32 (d, *J* = 8.1 Hz, 1H), 7.59 (br s, 1H), 7.36-7.30 (m, 2H), 7.21 (br s, 2H), 4.33-4.19 (m, 2H), 4.01-3.95 (m, 1H), 3.23-3.20 (m, 4H), 2.89-2.73 (m, 6H), 2.58 (s, 3H). MS: (ESI, m/z): 443 [M+H]⁺.

Example 21-1. N-((2R)-6-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Mixture of tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00453] A solution of tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (800 mg, 1.63 mmol) and Selectfluor (634 mg, 1.79 mmol) in ACN (16 mL) was stirred for 18 h at 20 °C. The reaction was quenched by the addition of 30 mL of water. The resulting mixture was extracted with 3x10 mL of ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1%-50% EtOAc/pet. ether) to afford a mixture of the regioisomers tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-(((benzyloxy)carbonyl)amino)-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a yellow oil. MS: (ESI, m/z): 510 [M+H]⁺.

Step 2. Mixture of tert-butyl 3-(6-amino-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-amino-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00454] A slurry of the mixture of tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-[[[(benzyloxy)carbonyl]amino]-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (508 mg, 1.00 mmol), and palladium on carbon (100 mg, 10%) in ethyl acetate (100 mL) was stirred for 1 h at 20 °C under a hydrogen atmosphere. The solids were filtered out. The resulting mixture was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1%-50% MeOH/DCM) to afford a mixture of the regioisomers tert-butyl 3-(6-amino-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-amino-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid. MS: (ESI, m/z): 376 [M+H]⁺.

Step 3. Mixture of tert-butyl 3-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00455] A solution of the mixture of tert-butyl 3-(6-amino-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-(6-amino-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (150 mg, 0.40 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (99 mg, 479.45 μmol), EDCI (92 mg, 0.48 mmol), HOBt (65 mg, 0.48 mmol), and DIEA (155 mg, 1.20 mmol) in DMF (5 mL) was stirred for 2 h at 20 °C. The reaction was quenched by the addition of 20 mL of water. The solids were collected by filtration and the crude product was purified by prep-HPLC (Column: XBridge BEH C18 OBD, 130 Å, 5 μm, 19 × 150 mm; Mobile phase A: water (10 mM NH₄HCO₃), B: ACN (60% ACN to 80% over 7 min)) to afford a mixture of the titled compounds.

[00456] Chiral separation. The mixture of regioisomers and enantiomers were partially separated by chiral HPLC (Column: Chiralpak IA, 250 x 20 mm, 5 μm; Flow: 15 mL/min; Mobile Phase 10% MeOH/MTBE; Wavelength: from 190 nm to 500 nm) to afford a mixture of tert-butyl

3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate and tert-butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (**Peak 1**: RT = 7.37 min; stereochemistry arbitrarily assigned); tert-butyl 3-((S)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid (**Peak 2**: RT = 7.69 min; stereochemistry arbitrarily assigned); and tert-butyl 3-((S)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid (**Peak 3**: RT = 13.70 min; stereochemistry arbitrarily assigned).

[00457] **Peak 1** was further purified by chiral HPLC (Column: Phenomenex Lux 5u Cellulose-4, AXIA Packed, 250 x 21.5 mm, 5 μ m; Flow: 18 mL/min, Mobile Phase: MeOH; Wavelength: from 190 nm to 500 nm) to afford tert-butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid (RT = 13.56 min; stereochemistry arbitrarily assigned), and tert-butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate as a white solid (RT = 17.31 min; stereochemistry arbitrarily assigned). MS for all 4 isomers: (ESI, m/z): 566 [M+H]⁺.




Step 4. N-((2R)-6-(3,8-Diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide






[00458] A solution of tert-butyl 3-((R)-6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-3-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (25 mg, 0.04 mmol) and TFA (1 mL) in DCM (5 mL) was stirred for 1 h at 20 °C. The resulting mixture was concentrated under vacuum. The residue was diluted with 5 mL of DCM. The pH of the solution was adjusted to 8 with NH₃ in MeOH (7 M). The resulting mixture was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep C18 OBD, 130 Å, 5 μ m, 19 x 150 mm; Mobile phase: water (10 mM NH₄HCO₃), ACN (25% ACN up to 55% over 7 min)) to afford N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid. ¹H NMR (Methanol-d₄, 400 MHz) δ (ppm): 8.20 (d, *J* = 8.0 Hz, 1H), 7.30 (d, *J* = 8.0




Hz, 1H), 6.80-6.76 (m, 2H), 4.25-4.17 (m, 1H), 3.53-3.51 (m, 2H), 3.16-3.14 (m, 2H), 3.10-2.99 (m, 2H), 2.92-2.90 (m, 2H), 2.82-2.74 (m, 2H), 2.64 (s, 3H), 2.18-2.12 (m, 1H), 2.04-2.02 (m, 2H), 1.87-1.82 (m, 3H). MS: (ESI, m/z): 466 [M+H]⁺.

[00459] The first three examples in Table 19 were prepared from the remaining three isomers isolated in Step 3 using standard chemical manipulations and procedures similar to those used for the preparation of Example 21-1. The rest of the examples in Table 19 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Examples 21-1 thru 21-4.

Table 19:

Example Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺	¹ H NMR
21-2	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-5-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	466	(CD ₃ OD, 400 MHz) δ(ppm): 8.20 (d, <i>J</i> = 8.4 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 6.84-6.75 (m, 2H), 4.25-4.16 (m, 1H), 3.66-3.62 (s, 2H), 3.21-3.18 (m, 2H), 3.05-2.95 (m, 4H), 2.82-2.75 (m, 2H), 2.64 (s, 3H), 2.18-2.07 (m, 3H), 1.91-1.81 (m, 2H), 1.81-1.76 (m, 1H).
21-3	 N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	466	(CD ₃ OD, 400 MHz) δ(ppm): 8.19 (d, <i>J</i> = 8.4 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 6.75-6.65 (m, 2H), 4.22-4.20 (m, 1H), 3.52-3.49 (m, 2H), 3.16-3.13 (m, 2H), 3.03-2.70 (m, 6H), 2.64 (s, 3H), 2.14-1.96 (m, 3H), 1.88-1.73 (m, 3H).
21-4	 N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-7-fluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	466	(CD ₃ OD, 400 MHz) δ(ppm): 8.19 (d, <i>J</i> = 8.4 Hz, 1H), 7.30 (d, <i>J</i> = 8.0 Hz, 1H), 6.78-6.67 (m, 2H), 4.25-4.17 (m, 1H), 3.67-3.63 (m, 2H), 3.21-3.18 (m, 2H), 3.03-2.72 (m, 6H), 2.64 (s, 3H), 2.11-2.05 (m, 3H), 1.90-1.79 (m, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
21-5 ¹	 (R)-3-amino-N-(5-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	440	(CD ₃ OD, 400 MHz) δ(ppm): 8.19 (d, J = 8.3 Hz, 1H), 7.38-7.29 (m, 1H), 6.94-6.78 (m, 2H), 4.32-4.19 (m, 1H), 3.09-2.94 (m, 10H), 2.85-2.70 (m, 2H), 2.64 (s, 3H), 2.25-2.13 (m, 1H), 1.82-1.75 (m, 1H).
21-6 ¹	 (S)-3-amino-N-(5-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	440	(CD ₃ OD, 400 MHz) δ(ppm): 8.19 (d, J = 8.3 Hz, 1H), 7.35-7.28 (m, 1H), 6.90-6.80 (m, 2H), 4.33-4.19 (m, 1H), 3.00 (s, 10H), 2.85-2.70 (m, 2H), 2.64 (s, 3H), 2.19-2.08 (m, 1H), 1.83-1.75 (m, 1H).
21-7 ¹	 (R)-3-amino-N-(7-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	440	(CD ₃ OD, 400 MHz) δ(ppm): 8.19 (d, J = 8.3 Hz, 1H), 7.35-7.28 (m, 1H), 6.82-6.73 (m, 2H), 4.29-4.18 (m, 1H), 3.04-2.93 (m, 9H), 2.94-2.85 (m, 2H), 2.85-2.67 (m, 1H), 2.64 (s, 3H), 2.18-2.09 (m, 1H), 1.88-1.74 (m, 1H).
21-8 ¹	 (S)-3-amino-N-(7-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide	440	(CD ₃ OD, 400 MHz) δ(ppm): 8.19 (d, J = 8.3 Hz, 1H), 7.35-7.28 (m, 1H), 6.90-6.80 (m, 2H), 4.33-4.19 (m, 1H), 3.00 (s, 10H), 2.85-2.70 (m, 2H), 2.64 (s, 3H), 2.19-2.08 (m, 1H), 1.83-1.75 (m, 1H).
21-9 ²	 (R)-7-amino-N-(5-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	441	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.64 (s, 1H), 7.81 (d, J = 7.7 Hz, 1H), 6.78-6.94 (m, 3H), 6.71 (d, J = 9.0 Hz, 1H), 4.12 (s, 1H), 2.6-2.96 (m, 16H), 1.98 (d, J = 12.1 Hz, 1H), 1.67-1.82 (m, 1H).


Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
21-10 ²	 (S)-7-amino-N-(5-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	441	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.63 (s, 1H), 7.82 (d, 1H), 6.90 (s, 2H), 6.83 (d, 1H), 6.71 (d, 1H), 4.11 (s, 1H), 2.75-2.88 (m, 13H), 2.63 (s, 3H), 1.69-2.01 (m, 2H).
21-11 ²	 (R)-7-amino-N-(7-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	441	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.63 (s, 1H), 7.82 (d, <i>J</i> = 7.7 Hz, 1H), 6.91 (s, 2H), 6.82 (d, <i>J</i> = 8.2 Hz, 2H), 4.11 (s, 1H), 2.88 (s, 10H), 2.66-2.83 (m, 1H), 2.63 (s, 5H), 2.01 (s, 1H), 1.70-1.72 (m, 1H).
21-12 ²	 (S)-7-amino-N-(7-fluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-methylthieno[2,3-b]pyrazine-6-carboxamide	441	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.64 (s, 1H), 7.82 (d, <i>J</i> = 7.7 Hz, 1H), 6.77-6.91 (m, 4H), 4.10 (br s, 1H), 2.64-2.95 (m, 16H), 2.63 (s, 5H), 2.01 (br s, 1H), 1.65-1.81 (m, 1H).

¹ **Notes on procedures:** The mixture of regioisomers and enantiomers were partially separated by chiral HPLC using the chiral column Chiralpak-AD-H-SL002 and mobile phase 50% EtOH/hexanes to provide the precursor to Example 21-7 as the first eluted sample, the precursor to Example 21-8 as the second eluted sample, and a mixture of two peaks as the third eluted sample. The mixture was further separated by chiral HPLC using the chiral column Chiralpak-IC and mobile phase 30% IPA/MTBE to provide the precursor to Example 21-5 as the first eluted sample and the precursor to Example 21-6 as the second eluted sample. Stereochemistry of the separated enantiomers were arbitrarily assigned.

² **Notes on procedures:** The mixture of regioisomers and enantiomers were partially separated by chiral HPLC using the chiral column Chiralpak-IA and mobile phase EtOH to provide the precursor to Example 21-9 as the first eluted sample, a mixture of two peaks as the second eluted sample, and the precursor to Example 21-12 as the third eluted sample. The mixture was further separated by chiral HPLC using the chiral column (R,R)-Whelk-O1-Kromasil and mobile phase 10% EtOH/MTBE to provide the precursor to Example 21-10 as the first eluted sample and the precursor to Example 21-11 as the second eluted sample. Stereochemistry of the separated enantiomers were arbitrarily assigned.

Example 22-1. (S)-3-Amino-N-(5,8-difluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 22-2. (R)-3-amino-N-(5,8-difluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl 4-(6-(benzyloxycarbonylamino)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate

[00460] To a solution of tert-butyl 4-(6-(benzyloxycarbonylamino)-4-fluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate (3.00 g, 6.20 mmol) in 1:1 DCM/MeOH (100 mL) was added Selectfluor (2.20 g, 6.20 mmol) at 0 °C. The resulting solution was stirred overnight at 20 °C. Two additional portions of Selectfluor (1.1 g, 3.1 mmol) were added at 0 °C and stirred for 4 h at 20 °C. The reaction mixture was then poured into 80 mL of water. The resulting mixture was extracted with 3 x 100 mL of DCM. The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 3:1 pet. ether/ethyl acetate) to afford tert-butyl 4-(6-(benzyloxycarbonylamino)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate as a light yellow solid. MS: (ESI, m/z): 502 [M+H]⁺.

Step 2. tert-Butyl 4-(6-amino-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate

[00461] A solution of tert-butyl 4-(6-(benzyloxycarbonylamino)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate (530 mg, 1.06 mmol) and palladium on carbon (100 mg, 10%) in ethyl acetate (10 mL) was stirred for 2 h at 20 °C under a hydrogen atmosphere. The solids were filtered out. The filtrate was concentrated under vacuum to afford tert-butyl 4-(6-amino-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 368 [M+H]⁺.

Step 3. tert-Butyl 4-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate

[00462] A solution of tert-butyl 4-(6-amino-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate (360 mg, 0.74 mmol, 75% purity), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (188 mg, 0.90 mmol), HBTU (343 mg, 0.90 mmol), and Et₃N (0.31 mL, 2.26 mmol) in DMA (5 mL) was stirred for 30 min at 20 °C. The crude product was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase A: water (0.05% formic acid), B: ACN (0% ACN up to 80% in 30 min)). The collected fraction was concentrated under vacuum to afford tert-butyl 4-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate as a yellow solid. MS: (ESI, m/z): 558 [M+H]⁺.

Step 4. Chiral separation.

[00463] The racemate tert-butyl 4-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate (250 mg) was separated by chiral HPLC (Column: ChiralArt Cellulose-SB, 2x25 cm, 5 µm; Mobile Phase: 30% EtOH/hexanes for 13 min). The first eluting isomer (RT=9.05 min) was concentrated under vacuum to afford (S)-tert-butyl 4-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate as a yellow solid (stereochemistry arbitrarily assigned). The second eluting isomer (RT=10.15 min) was concentrated under vacuum to afford (R)-tert-butyl 4-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate as a yellow solid (stereochemistry arbitrarily assigned). MS for both isomers: (ESI, m/z): 558 [M+H]⁺.

Step 5. (S)-3-Amino-N-(5,8-difluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide




[00464] A solution of (S)-tert-butyl 4-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate (60 mg, 0.11 mmol) and trifluoroacetic acid (1 mL) in DCM (3 mL) was stirred for 1 h at 20 °C. The resulting mixture was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep OBD C18, 30 x 150 mm, 5 µm; Mobile phase A: water (0.05% NH₄OH), B: ACN (15% ACN up to 50% in 7 min)). The collected fraction was lyophilized to afford (S)-3-amino-N-(5,8-difluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid. ¹H NMR (CD₃OD, 300 MHz) δ(ppm): 8.17 (d, *J* = 8.1 Hz, 1H), 7.30 (d, *J* = 8.4 Hz, 1H), 6.67-6.65 (m, 1H), 4.25-4.17 (m, 1H), 3.14-2.97 (m, 10H), 2.82-2.70 (m, 1H), 2.64-2.54 (m, 4H), 1.89-1.74 (m, 1H). MS: (ESI, *m/z*): 458 [M+H]⁺.


Step 6. (R)-3-amino-N-(5,8-difluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00465] A solution of (R)-tert-butyl 4-(6-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-1,4-difluoro-5,6,7,8-tetrahydronaphthalen-2-yl)piperazine-1-carboxylate (90 mg, 0.16 mmol) and trifluoroacetic acid (1.5 mL) in DCM (4 mL) was stirred for 1 h at 20 °C. The resulting mixture was concentrated under vacuum. The residue was purified by prep-HPLC (Column, XBridge Prep OBD C18, 30x150 mm, 5 µm; mobile phase A: water (0.05% NH₄OH), B: ACN (15% ACN up to 50% in 7 min)). The collected fraction was lyophilized to afford (R)-3-amino-N-(5,8-difluoro-6-(piperazin-1-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as an off-white solid. ¹H NMR (DMSO-*d*₆, 300 MHz) δ(ppm): 8.31 (d, *J* = 8.4Hz, 1H), 7.64 (d, *J* = 7.5Hz, 1H), 7.31 (d, *J* = 8.1Hz, 1H), 7.17 (s, 2H), 6.70-6.68 (m, 1H), 4.13-4.04 (m, 1H), 2.92-2.81 (m, 10H), 2.73-2.55 (m, 5H), 2.06-1.97 (m, 1H), 1.78-1.68 (m, 1H). MS: (ESI, *m/z*): 458 [M+H]⁺.

[00466] The following examples in Table 20 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Examples 22-1 and 22-2.

Table 20:

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
22-3 ¹	 N-((2S)-6-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)-5,8-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	500	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.66 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.18 (br s, 2H), 6.78-6.75 (m, 1H), 4.20-4.00 (m, 1H), 3.75-3.65 (m, 2H), 3.48-3.38 (m, 2H), 3.27-3.08 (m, 4H), 3.01-2.82 (m, 4H), 2.80-2.60 (m, 2H), 2.58 (s, 3H), 2.10-1.97 (m, 1H), 1.88-1.62 (m, 1H).
22-4 ¹	 N-((2R)-6-(9-oxa-3,7-diazabicyclo[3.3.1]nonan-3-yl)-5,8-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	500	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.1 Hz, 1H), 7.66 (br s, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.18 (br s, 2H), 6.78-6.75 (m, 1H), 4.20-4.00 (m, 1H), 3.75-3.65 (m, 2H), 3.48-3.38 (m, 2H), 3.27-3.08 (m, 4H), 3.01-2.82 (m, 4H), 2.80-2.60 (m, 2H), 2.58 (s, 3H), 2.10-1.98 (m, 1H), 1.88-1.62 (m, 1H).
22-5 ²	 3-amino-N-[(2S)-6-{3,8-diazabicyclo[3.2.1]octan-3-yl}-5,8-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	484	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.30 (d, <i>J</i> = 8.4 Hz, 1H), 7.63 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.17 (br s, 2H), 6.61-6.57 (m, 1H), 4.09-4.07 (m, 1H), 3.40-3.98 (m, 2H), 3.08-3.03 (m, 2H), 2.95-2.62 (m, 5H), 2.59-2.57 (m, 4H), 2.37-2.30 (m, 1H), 2.02-1.99 (m, 1H), 1.80-1.61 (m, 5H).


Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
22-6 ²	 3-amino-N-[(2R)-6-{3,8-diazabicyclo[3.2.1]octan-3-yl}-5,8-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	484	(DMSO- <i>d</i> ₆ , 300 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.64 (d, <i>J</i> = 8.0 Hz, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.17 (br s, 2H), 6.61-6.57 (m, 1H), 4.14-4.05 (m, 1H), 3.40-3.98 (m, 2H), 3.09-3.02 (m, 2H), 2.96-2.87 (m, 2H), 2.81-2.75 (m, 2H), 2.70-2.61 (m, 1H), 2.59-2.57 (m, 4H), 2.34-2.32 (m, 1H), 2.04-1.97 (m, 1H), 1.81-1.77 (m, 2H), 1.76-1.68 (m, 1H), 1.67-1.61 (m, 2H).

¹ **Notes on procedures:** In Step 4, the stereoisomers were separated by chiral HPLC using the chiral column Chiralpak IG and mobile phase 15% EtOH/MTBE. Stereochemistry of the separated enantiomers were arbitrarily assigned.

² **Notes on procedures:** In Step 4, the stereoisomers were separated by chiral HPLC using the chiral column Chiral Art Cellulose-SB and mobile phase 10% EtOH/MTBE. Stereochemistry of the separated enantiomers were arbitrarily assigned.

Example 23-1. 7-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide

Method 1.



Step 1. Benzyl N-[(6S)-2-[(3S,4S)-3-[(tert-butoxy)carbonyl]amino]-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

[00467] A mixture of benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) (500 mg, 1.16 mmol), tert-butyl N-[(3S,4S)-4-methoxypyrrolidin-3-yl]carbamate (300 mg, 1.39 mmol), RuPhos Pd G3 (195 mg, 0.230 mmol), RuPhos (108 mg, 0.230 mmol), and Cs₂CO₃ (1.14 g, 3.49 mmol) in toluene (10 mL) was stirred for 3 h at 95 °C. After cooling to 20 °C, the solids were filtered out and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 5:2 pet. ether/ethyl acetate) to give benzyl N-[(6S)-2-[(3S,4S)-3-[[tert-butoxy]carbonyl]amino]-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a brown solid. MS (ESI, m/z): 497 [M+H]⁺.

Step 2. tert-butyl N-[(3S,4S)-1-[(6S)-6-Amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-methoxypyrrolidin-3-yl]carbamate

[00468] A mixture of benzyl N-[(6S)-2-[(3S,4S)-3-[[tert-butoxy]carbonyl]amino]-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (450 mg, 0.910 mmol) and Palladium on carbon (450 mg, 10%) in ethyl acetate (10 mL) was stirred for 3 h at 20 °C under a hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give tert-butyl N-[(3S,4S)-1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-methoxypyrrolidin-3-yl]carbamate as a white solid. MS (ESI, m/z): 362 [M+H]⁺.

Step 3. tert-butyl N-[(3S,4S)-1-(6-[7-Amino-3-methylthieno[2,3-b]pyrazine-6-amido]-5,6,7,8-tetrahydroquinolin-2-yl)-4-methoxypyrrolidin-3-yl]carbamate

[00469] Triethylamine (0.400 mL, 2.92 mmol) and HBTU (377 mg, 0.990 mmol) were added to a stirring solution of tert-butyl N-[(3S,4S)-1-(6-amino-5,6,7,8-tetrahydroquinolin-2-yl)-4-methoxypyrrolidin-3-yl]carbamate (300 mg, 0.830 mmol) and 7-amino-3-methylthieno[2,3-b]pyrazine-6-carboxylic acid (174 mg, 0.830 mmol) in DMA (5 mL). The resulting solution was stirred for 2 h at 20 °C. The mixture was purified by reversed phase chromatography (Column: C18 silica gel; Mobile phase, A: water (10 mM NH₄HCO₃) and B: ACN (0% to 70% B over 30 min)). The collected fraction was concentrated under vacuum to give tert-butyl N-[(3S,4S)-1-(6-[7-amino-3-methylthieno[2,3-b]pyrazine-6-amido]-5,6,7,8-tetrahydroquinolin-2-yl)-4-methoxypyrrolidin-3-yl]carbamate as a green solid. MS (ESI, m/z): 554 [M+H]⁺.


Step 4. 7-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide

[00470] A solution of tert-butyl N-[(3S,4S)-1-[(6S)-6-[7-amino-3-methylthieno[2,3-b]pyrazine-6-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-methoxypyrrolidin-3-yl]carbamate (300 mg, 0.470 mmol) and TFA (3 mL) in DCM (10 mL) was stirred for 30 min at 20 °C and then concentrated under vacuum. The residue was treated with a solution of NH₃ in MeOH (15 mL, 7M) for 1 h at 20 °C. The resulting solution was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep C18 OBD, 19 x 150 mm 5 µm; Mobile Phase, A: water (containing 10 mM NH₄HCO₃) and B: ACN (20% to 42% B over 7 min)) to afford 7-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide as a yellow solid. MS (ESI, m/z): 454 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.65 (s, 1H), 7.81 (d, *J* = 7.6 Hz, 1H), 7.20 (d, *J* = 8.8 Hz, 1H), 6.92 (br s, 2H), 6.23 (d, *J* = 8.4 Hz, 1H), 4.28-4.12 (m, 1H), 3.62-3.59 (m, 2H), 3.48-3.41 (m, 2H), 3.36-3.33 (m, 4H), 3.14-3.12 (m, 1H), 2.84-2.73 (m, 4H), 2.66 (s, 3H), 2.05-2.00 (m, 1H), 1.90-1.83 (m, 1H), 1.79 (br s, 2H).

Example 23-1. 7-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide and

Example 23-2. 7-Amino-N-[(6R)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide


Method 2.












[00471] 7-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide was also obtained starting from the racemate benzyl N-[2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-1**) and following procedures similar to Method 1. The enantiomers of the racemate 7-amino-N-[2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide were separated by chiral HPLC using the chiral column Chiralpak IE and the eluent system 50% EtOH/MTBE (containing 2 mM NH₃ in MeOH) to provide 7-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide as a yellow solid as the first eluted isomer and 7-amino-N-[(6R)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide as a yellow solid as the second eluted isomer. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.65 (s, 1H), 7.83 (d, *J* = 7.6 Hz, 1H), 7.20 (d, *J* = 8.4 Hz, 1H), 6.93 (br, 2H), 6.24 (d, *J* = 8.8 Hz, 1H), 4.20-4.11(m, 1H), 3.63-3.56 (m, 2H), 3.47-3.44 (m, 2H), 3.33 (s, 3H), 3.15-3.13 (m, 1H), 2.84-2.73 (m, 4H), 2.65 (s, 3H), 2.05-1.98 (m, 1H), 1.91-1.83 (m, 2H). MS (ESI, *m/z*): 454 [M+H]⁺.





[00472] The following examples in Table 21 were prepared using standard chemical manipulations and procedures similar to Method 1 (or Method 2 where indicated) for the preparation of Example 23-1.



Table 21:

Example Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺	¹ H NMR
23-3 ¹	 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide	467	(DMSO- <i>d</i> ₆ , 400 MHz) δ(ppm): 7.59 (d, <i>J</i> = 7.6 Hz, 1H), 7.19 (d, <i>J</i> = 8.4 Hz, 1H), 7.04 (s, 1H), 6.81 (br s, 2H), 6.23 (d, <i>J</i> = 8.4 Hz, 1H), 4.08-4.15 (m, 1H), 3.61-3.63 (m, 2H), 3.43-3.33 (m, 3H), 3.30 (s, 3H), 3.13-3.11 (m, 1H), 2.81-2.69 (m, 7H), 2.58 (s, 3H), 1.99-1.72 (m, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
23-4 ²	 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	453	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.58 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.20-7.17 (m, 3H), 6.23 (d, <i>J</i> = 8.4 Hz, 1H), 4.16-4.07 (m, 1H), 3.62-3.58 (m, 2H), 3.47-3.41 (m, 2H), 3.29 (s, 3H), 3.12 (d, <i>J</i> = 8.4 Hz, 1H), 2.82-2.68 (m, 4H), 2.59 (s, 3H), 2.01-1.98 (m, 1H), 1.82-1.74 (m, 2H).
23-5 ²	 3-amino-N-[(6R)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	453	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.30 (d, <i>J</i> = 8.0 Hz, 1H), 7.60 (d, <i>J</i> = 8.0 Hz, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.20-7.18 (m, 3H), 6.22 (d, <i>J</i> = 8.4 Hz, 1H), 4.14-4.07 (m, 1H), 3.61-3.57 (m, 2H), 3.44-3.42 (m, 2H), 3.29 (s, 3H), 3.13 (d, <i>J</i> = 8.0 Hz, 1H), 2.81-2.68 (m, 4H), 2.58 (s, 3H), 2.01-1.98 (m, 1H), 1.82-1.74 (m, 2H).
23-6	 7-amino-3-methyl-N-[(6S)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyrazine-6-carboxamide	424	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.83 (d, <i>J</i> = 8.0 Hz, 1H), 7.24 (d, <i>J</i> = 8.4 Hz, 1H), 6.92 (br s, 2H), 6.63-6.57 (m, 1H), 4.18-4.14 (m, 1H), 3.41-3.33 (m, 3H), 2.86-2.70 (m, 9H), 2.56 (s, 3H), 2.04-2.01 (m, 1H), 1.92-1.82 (m, 1H).
23-7 ³	 3-amino-6-methyl-N-[(6S)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide	423	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.59 (d, <i>J</i> = 8.0 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.23 (d, <i>J</i> = 8.4 Hz, 1H), 7.17 (br s, 2H), 6.58 (d, <i>J</i> = 8.4 Hz, 1H), 4.18-4.16 (m, 1H), 3.34-3.32 (m, 4H), 2.94-2.68 (m, 8H), 2.58 (s, 3H), 2.08-1.90 (m, 1H), 1.88-1.80 (m, 1H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
23-8 ³	 3-amino-6-methyl-N-[(6R)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide	423	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.0 Hz, 1H) 7.59 (d, <i>J</i> = 8.0 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.23 (d, <i>J</i> = 8.4 Hz, 1H), 7.17 (br s, 2H), 6.58 (d, <i>J</i> = 8.4 Hz, 1H), 4.18-4.16 (m, 1H), 3.34-3.32 (m, 4H), 2.94-2.68 (m, 8H), 2.58 (s, 3H), 2.08-1.90 (m, 1H), 1.88-1.80 (m, 1H).
23-9 ⁴	 3-amino-6-methyl-N-[(6R)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide	429	(DMSO-d ₆ , 400 MHz) δ(ppm): 7.59 (d, <i>J</i> = 7.6 Hz, 1H), 7.23 (d, <i>J</i> = 8.4 Hz, 1H), 7.08 (br s, 2H), 6.58 (d, <i>J</i> = 8.4 Hz, 1H), 4.13-4.07 (m, 1H), 3.41-3.32 (m, 4H), 2.79-2.68 (m, 1H), 2.00-1.97 (m, 1H), 1.86-1.80 (m, 1H).
23-10	 3-amino-N-[(6S)-2-{3,6-diazabicyclo[3.1.1]heptan-3-yl}-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	435	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.58 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.25 (d, <i>J</i> = 8.4 Hz, 1H), 7.16 (br s, 2H), 6.40 (d, <i>J</i> = 8.4 Hz, 1H), 4.18-4.12 (m, 1H), 3.67-3.66 (m, 2H), 3.57-3.50 (m, 4H), 3.40-3.33 (m, 2H), 2.83-2.75 (m, 4H), 2.59 (s, 3H), 2.03-2.00 (m, 1H), 1.89-1.84 (m, 1H), 1.44 (d, <i>J</i> = 8.0 Hz, 1H).
23-11	 7-amino-N-[(6S)-2-{3,6-diazabicyclo[3.1.1]heptan-3-yl}-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide	436	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.65 (s, 1H), 7.83 (d, <i>J</i> = 8.0 Hz, 1H), 7.26 (d, <i>J</i> = 8.4 Hz, 1H), 6.92 (br s, 2H), 6.40 (d, <i>J</i> = 9.2 Hz, 1H), 4.19-4.16 (m, 1H), 3.68-3.66 (m, 2H), 3.60-3.51 (m, 4H), 3.33-3.20 (m, 2H), 2.83-2.76 (m, 4H), 2.66 (s, 3H), 2.05-2.02 (m, 1H), 1.90-1.87 (m, 1H), 1.44 (d, <i>J</i> = 8.8 Hz, 1H).
23-12	 3-amino-N-[(6S)-2-{3,6-diazabicyclo[3.1.1]heptan-3-yl}-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	449	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.8 Hz, 1H), 7.56 (d, <i>J</i> = 7.2 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.22-7.15 (m, 3H), 6.45 (d, <i>J</i> = 8.8 Hz, 1H), 4.14 (br s, 1H), 3.77-3.75

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
	3-amino-N-[(6S)-2-{3,8-diazabicyclo[3.2.1]octan-3-yl}-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide		(m, 2H), 3.47 (s, 2H), 2.82-2.68 (m, 6H), 2.59 (s, 3H), 2.02-1.83 (m, 2H), 1.63-1.54 (m, 4H).
23-13	 7-amino-N-[(6S)-2-{3,8-diazabicyclo[3.2.1]octan-3-yl}-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide	450	(DMSO-d ₆ , 300 MHz) δ(ppm): 8.65 (s, 1H), 7.82 (d, <i>J</i> = 7.6 Hz, 1H), 7.42 (d, <i>J</i> = 8.4 Hz, 1H), 6.93 (br s, 2H), 6.47-6.42 (m, 1H), 4.16 (br s, 1H), 3.77-3.73 (m, 2H), 2.85-2.66 (m, 6H), 2.58 (s, 3H), 2.41 (s, 1H), 2.03-1.81 (m, 2H), 1.67-1.54 (m, 4H).
23-14 ¹	 3-amino-4,6-dimethyl-N-[(6S)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide	437	(DMSO-d ₆ , 400 MHz) δ(ppm): 7.61 (d, <i>J</i> = 8.0 Hz, 1H), 7.24 (d, <i>J</i> = 8.8 Hz, 1H), 7.14 (s, 1H), 6.82 (br, 2H), 6.58 (d, <i>J</i> = 8.8 Hz, 1H), 4.16-4.14 (m, 1H), 3.41-3.33 (m, 6H), 2.82-2.73 (m, 12H), 2.00-1.99 (m, 1H), 1.88-1.83 (m, 1H).
23-15 ⁵	 3-amino-6-methyl-N-[(6S)-2-[(3S)-3-methylpiperazin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide	437	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.58 (d, <i>J</i> = 7.6 Hz, 1H), 7.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.23 (d, <i>J</i> = 8.8 Hz, 1H), 7.16 (br s, 2H), 6.59 (d, <i>J</i> = 8.4 Hz, 1H), 4.14-4.00 (m, 3H), 2.94-2.91 (m, 1H), 2.84-2.65 (m, 6H), 2.63-2.60 (m, 4H), 2.26-2.20 (m, 1H), 1.99-2.03 (m, 2H), 1.91-1.83 (m, 1H), 1.03-1.01 (d, <i>J</i> = 8.0 Hz, 1H).
23-16 ⁶	 3-amino-6-methyl-N-[(6S)-2-[(3R)-3-methylpiperazin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide	437	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.57 (d, <i>J</i> = 8.0 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.23 (d, <i>J</i> = 8.4 Hz, 1H), 7.16 (br s, 2H), 6.59 (d, <i>J</i> = 8.0 Hz, 1H), 4.13-4.00 (m, 3H), 2.94-2.61 (m, 7H), 2.59 (s, 3H), 2.26-2.20 (m, 2H), 1.99-1.83 (m, 2H), 1.02 (d, <i>J</i> = 6.4 Hz, 3H).

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
23-17 ⁷	 3-amino-N-[(6S)-4-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	441	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.63 (d, <i>J</i> = 7.2 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.17 (br s, 2H), 6.48 (d, <i>J</i> = 12.8 Hz, 1H), 4.22-4.12 (m, 1H), 3.36-3.20 (m, 4H), 2.90-2.84 (m, 1H), 2.82-2.75 (m, 6H), 2.59-2.54 (m, 4H), 2.04-1.98 (m, 1H), 1.92-1.83 (m, 1H).
23-18 ⁸	 N-((6S)-2-(3,8-diazabicyclo[3.2.1]octan-3-yl)-4-fluoro-5,6,7,8-tetrahydroquinolin-6-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide	467	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.63 (d, <i>J</i> = 7.9 Hz, 1H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 7.17 (br s, 2H), 6.33 (d, <i>J</i> = 12.6 Hz, 1H), 4.16-4.14 (m, 1H), 3.76-3.73 (m, 2H), 3.48-3.46 (m, 2H), 2.89-2.74 (m, 5H), 2.59-2.53 (m, 4H), 1.98-1.83 (m, 2H), 1.75-1.63 (m, 4 H).

¹ Notes on procedures: In Step 3, the carboxylic acid **Intermediate 21** was used.

² Notes on procedures: Following Method 2, the stereoisomers were separated by chiral HPLC using the chiral column Chiralpak IE and the eluent gradient 30% to 35% EtOH/MTBE (containing 2 mM NH₃ in MeOH).

³ Notes on procedures: Following Method 2, the stereoisomers were separated by chiral HPLC using the chiral column Chiralpak ID-3 and mobile phase 30% EtOH/(hexanes:DCM=3:1, containing 0.1% DIEA).

⁴ Notes on procedures: In Step 3, the carboxylic acid **Intermediate 24** was used.

⁵ Notes on procedures: In Step 1, tert-butyl (2S)-2-methylpiperazine-1-carboxylate was used.


⁶ Notes on procedures: In Step 1, tert-butyl (2R)-2-methylpiperazine-1-carboxylate was used.

⁷ Notes on procedures: Skipping Step 1 and Step 2, the amine **Intermediate 52-1** was used in Step 3.

⁸ Notes on procedures: Skipping Step 1 and Step 2, the amine **Intermediate 52-2** was used in Step 3.

Example 24-1. 3-Amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

Method 1.



Step 1. Benzyl N-[(6S)-2-[(3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

[00473] A mixture of benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) (800 mg, 1.85 mmol), tert-butyl N-[(3R,4R)-4-(methoxymethyl)pyrrolidin-3-yl]carbamate (**Intermediate 48-2**) (513 mg, 2.23 mmol), RuPhos Pd G3 (155 mg, 0.186 mmol), RuPhos (86.7 mg, 0.186 mmol) and Cs₂CO₃ (1.21 g, 3.71 mmol) in toluene (10 mL) was stirred for 3 h at 90 °C. After cooling to 25 °C, the solids were filtered out. The filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:2 ethyl acetate/pet. ether) to afford benzyl N-[(6S)-2-[(3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a white solid. MS (ESI, m/z): 511 [M+H]⁺.

Step 2. tert-Butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate

[00474] A mixture of benzyl N-[(6S)-2-[(3R,4R)-3-[[[(tert-butoxy)carbonyl]amino]-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (550 mg, 1.07 mmol) and palladium on carbon (80 mg, 10%) in ethyl acetate (5 mL) was stirred for 1 h at 25 °C under a hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give tert-butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate as a white solid (crude). MS (ESI, m/z): 377 [M+H]⁺.

Step 3. tert-Butyl N-[(3R,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate

[00475] Et₃N (0.50 mL, 3.58 mmol) and HBTU (453 mg, 1.19 mmol) were added to a stirring solution of tert-butyl N-[(3R,4R)-1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate (450 mg, 1.19 mmol) and 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (248 mg, 1.19 mmol) in DMA (4 mL). The resulting solution was stirred for 2 h at 25 °C. The mixture was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (15% to 90% over 25 min)). The collected fraction was concentrated under vacuum to give tert-butyl N-[(3R,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate as a yellow solid. MS (ESI, m/z): 567 [M+H]⁺.


Step 4. 3-Amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00476] A solution of tert-butyl N-[(3R,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate (450 mg, 0.794 mmol) and TFA (3 mL) in DCM (12 mL) was stirred for 1 h at 25 °C. The mixture was concentrated under vacuum. The residue was treated with a solution of NH₃ in MeOH (2 mL, 7M). The resulting mixture was stirred for 10 min and concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Shield RP18 OBD, 30 x150 mm, 5 µm; Mobile Phase A: water (10 mM NH₄HCO₃) and B: ACN (27% to 37% over 7 min); Flow rate: 60 mL/min). The collected fraction was lyophilized to give 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 467 [M+H]⁺. ¹H NMR (DMSO-d₆, 400 MHz) δ(ppm): 8.31 (d, *J* = 8.0 Hz, 1H), 7.59 (d, *J* = 8.0 Hz, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.19-7.17 (m, 3H), 6.20 (d, *J* = 8.4 Hz, 1H), 4.15-4.08 (m, 1H), 3.61-3.51 (m, 2H), 3.52-3.35 (m, 3H), 3.29 (s, 3H), 3.25-3.15 (m, 2H), 2.81-2.64 (m, 4H), 2.60 (s, 3H), 2.49-2.38 (m, 1H), 2.09-1.98 (m, 1H), 1.91-1.80 (m, 1H), 1.61 (br s, 2H).

Example 24-1. 3-Amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 24-2. 3-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

Method 2.




[00477] 3-Amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide was also obtained starting from the cis racemate tert-butyl N-[4-(methoxymethyl)pyrrolidin-3-yl]carbamate (**Intermediate 48-1**) and following procedures similar to Method 1. The enantiomers of the cis racemate tert-butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate (Step 3 product) were separated by chiral HPLC using the chiral column Chiralpak IA and Mobile Phase, 50% A: hexanes:DCM=3:1 and 50% B: EtOH (containing 0.1% Et₂NH) to provide tert-butyl N-[(3R,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate as a white solid as the first eluted isomer (RT=9.3 min) and tert-butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate as a white solid as the second eluted isomer (RT=11.3 min).

[00478] 3-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide was obtained from conversion of tert-butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]carbamate following procedures similar to Method 1, Step 4. ¹H NMR (DMSO-d₆, 400 MHz) δ(ppm): 8.30 (d, *J* = 8.0

Hz, 1H), 7.56 (br s, 1H), 7.31 (d, $J = 8.4$ Hz, 1H), 7.18-7.16 (m, 3H), 6.20 (d, $J = 8.4$ Hz, 1H), 4.18-4.09 (m, 1H), 3.61-3.51 (m, 2H), 3.50-3.39 (m, 3H), 3.28 (s, 3H), 3.21-3.17 (m, 2H), 2.81-2.64 (m, 4H), 2.67 (s, 3H), 2.46-2.40 (m, 1H), 2.09-1.91 (m, 1H), 1.91-1.80 (m, 1H), 1.55 (br s, 2H). MS (ESI, m/z): 467 $[M+H]^+$.

Example 25. 3-Amino-N-[(6S)-2-[(3S,4R)-3-amino-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl N-[(6S)-2-[(3S,4R)-3-[(tert-butoxy)carbonyl]amino]-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

[00479] A mixture of benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) (480 mg, 1.09 mmol), tert-butyl N-[(3S,4R)-4-(difluoromethyl)pyrrolidin-3-yl]carbamate (**Intermediate 51-2**) (284 mg, 1.20 mmol), RuPhos (51.0 mg, 0.109 mmol), RuPhos Pd G3 (91.0 mg, 0.109 mmol) and Cs_2CO_3 (890 mg, 2.73 mmol) in toluene (10 mL) was stirred for 4 h at 95 °C. After cooled to 25 °C, the solids were filtered out. The filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford benzyl N-[(6S)-2-[(3S,4R)-3-[(tert-butoxy)carbonyl]amino]-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a white solid. MS (ESI, m/z): 517 $[M+H]^+$.

Step 2. tert-butyl N-[(3S,4R)-1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(difluoromethyl)pyrrolidin-3-yl]carbamate

[00480] A mixture of benzyl N-[(6S)-2-[(3S,4R)-3-[[tert-butoxy]carbonyl]amino]-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (350 mg, 0.664 mmol) and Pd/C (350 mg, 10%) in ethyl acetate (10 mL) was stirred 2 h at 25 °C under hydrogen atmosphere. The solids were filtered out and the filter cake was washed with ethyl acetate (3 x 10 mL). The filtrate was concentrated under reduced pressure to afford tert-butyl N-[(3S,4R)-1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(difluoromethyl)pyrrolidin-3-yl]carbamate as a brown solid. MS (ESI, m/z): 383 [M+H]⁺.

Step 3. tert-Butyl N-[(3S,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(difluoromethyl)pyrrolidin-3-yl]carbamate

[00481] A solution of tert-butyl N-[(3S,4R)-1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(difluoromethyl)pyrrolidin-3-yl]carbamate (240 mg, 0.615 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (141 mg, 0.677 mmol), Et₃N (0.260 mL, 1.85 mmol) and HBTU (280 mg, 0.738 mmol) in DMA (5 mL) was stirred for 6 h at 25 °C. The mixture was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.1% TFA) and B: ACN (0% to 60% in 30 min)) to afford tert-butyl N-[(3S,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(difluoromethyl)pyrrolidin-3-yl]carbamate as a yellow solid. MS (ESI, m/z): 573 [M+H]⁺.


Step 4. 3-Amino-N-[(6S)-2-[(3S,4R)-3-amino-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00482] To a stirring solution of tert-butyl N-[(3S,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(difluoromethyl)pyrrolidin-3-yl]carbamate (300 mg, 0.513 mmol) in DCM (15 mL) was added TFA (5 mL) dropwise at 0 °C. The resulting mixture was stirred for 40 min at 25 °C. The resulting mixture was concentrated under vacuum. The resulting mixture was concentrated under vacuum. The residue was treated with NH₃ (15 mL, 7 M in MeOH). The solution was stirred for 30 min at 25 °C and concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep Phenyl OBD, 19 × 150 mm, 5 μm; Mobile Phase, A: water (10 mM NH₄HCO₃) and B: ACN (20% to 37% in 8 min); Flow rate: 60 mL/min). The product fraction was lyophilized to afford 3-amino-N-[(6S)-2-[(3S,4R)-3-amino-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 473 [M+H]⁺. ¹H

NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 8.31 (d, *J* = 8.0 Hz, 1H), 7.59 (d, *J* = 7.6 Hz, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.21 (d, *J* = 8.4 Hz, 1H), 7.17 (br s, 2H), 6.34-6.05 (m, 2H), 3.68-3.57 (m, 2H), 3.51-3.48 (m, 1H), 3.39-3.77 (m, 1H), 3.04-3.00 (m, 1H), 2.81-2.75 (m, 4H), 2.72 (s, 3H), 2.48-2.42 (m, 1H), 2.08-1.91 (m, 1H), 1.90-1.78 (m, 3H).

Example 26-1. 3-Amino-N-[(6S)-2-[(3S,4R)-3-(methoxymethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 26-2. 3-Amino-N-[(6S)-2-[(3R,4S)-3-(methoxymethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. trans-Benzyl N-[(6S)-2-(3-[[[(tert-butoxy)carbonyl](methyl)amino]-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

[00483] A mixture of benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) (200 mg, 0.446 mmol), trans-tert-butyl N-[4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate (**Intermediate 53**) (121 mg, 0.446 mmol), RuPhos (42.0 mg, 0.090 mmol), RuPhos Pd G3 (38.0 mg, 0.045 mmol) and Cs₂CO₃ (291 mg, 0.893 mmol) in toluene (4 mL) was stirred for 3 h at 90 °C. The solids were filtered out. The filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford trans-benzyl N-[(6S)-2-(3-

[[[(tert-butoxy)carbonyl](methyl)amino]-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a yellow solid. MS (ESI, m/z): 525 $[M+H]^+$.

Step 2. trans-tert-Butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate

[00484] A mixture of trans-benzyl N-[(6S)-2-(3-[[[(tert-butoxy)carbonyl](methyl)amino]-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (180 mg, 0.326 mmol) and palladium on carbon (180 mg, 10%) in ethyl acetate (10 mL) was stirred for 3 h at 25 °C under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under reduced pressure to afford trans-tert-butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate as a white solid. MS (ESI, m/z): 391 $[M+H]^+$.

Step 3. trans-tert-Butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate

[00485] HBTU (122 mg, 0.322 mmol) was added to a stirring solution of trans-tert-butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate (70.0 mg, 0.179 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (39.0 mg, 0.189 mmol) and Et₃N (0.075 mL, 0.537 mmol) in DMA (2 mL). The resulting solution was stirred for 1 h at 25 °C. The resulting mixture was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (5% to 80% in 30 min)). The collected fraction was concentrated under vacuum to give trans-tert-butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate as a yellow solid. MS (ESI, m/z): 581 $[M+H]^+$.

Step 4. tert-Butyl N-[(3R,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate and tert-butyl N-[(3S,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate

[00486] trans-tert-Butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate (40.0

mg, 0.0690 mmol) was separated by chiral HPLC (Column: Chiralpak IA, 2 x 25 cm, 5 μ m; Mobile phase, A: hexanes:DCM=3:1 (containing 0.1% Et₂NH) and B: EtOH (hold 30% in 30 min); Flow rate: 12 mL/min). The first eluting isomer (RT=11.8 min) was collected and concentrated under vacuum to give a yellow solid, stereochemistry on the pyrrolidine arbitrarily assigned as tert-butyl N-[(3S,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate. The second eluting isomer (RT=21.4 min) was collected and concentrated under vacuum to give a yellow solid, stereochemistry on the pyrrolidine arbitrarily assigned as tert-butyl N-[(3R,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate. MS (ESI, *m/z*): 581 [M+H]⁺.

Step 5. 3-Amino-N-[(6S)-2-[(3S,4R)-3-(methoxymethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00487] A solution of tert-butyl N-[(3S,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-methylcarbamate (11.0 mg, 0.0190 mmol) and TFA (0.500 mL) in DCM (1.50 mL) was stirred for 30 min at 25 °C. The resulting mixture was concentrated under vacuum. The residue was treated with a solution of NH₃ (2.00 mL, 7M in MeOH). The resulting solution was stirred for 30 min at 25 °C and concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Shield RP18 OBD, 5 μ m, 19 x 150 mm; Mobile Phase A: water (10 mM NH₄HCO₃) and B: ACN (0% to 55% in 7 min); Flow rate: 25 mL/min). The collected fraction was lyophilized to give 3-amino-N-[(6S)-2-[(3S,4R)-3-(methoxymethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, *m/z*): 481 [M+H]⁺. ¹H-NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 8.31 (d, *J* = 8.4 Hz, 1H), 7.59 (br s, 1H), 7.31 (d, *J* = 8.0 Hz, 1H), 7.20-7.17(m, 3H), 6.24 (d, *J* = 8.0 Hz, 1H), 4.14-4.10 (m, 1H), 3.64-3.59 (m, 1H), 3.54-3.50 (m, 1H), 3.46-3.27 (m, 8H), 3.17-3.10 (m, 5H), 2.55 (s, 3H), 2.35-2.30 (m, 4H), 2.02-1.97 (m, 1H), 1.88-1.82 (m, 1H).


Step 6. 3-Amino-N-[(6S)-2-[(3R,4S)-3-(methoxymethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00488] A solution of tert-butyl N-[(3R,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(methoxymethyl)pyrrolidin-3-yl]-N-

methylcarbamate (13.0 mg, 0.022 mmol) and TFA (0.500 mL) in DCM (1.50 mL) was stirred for 30 min at 25 °C. The resulting mixture was concentrated under vacuum. The residue was treated with a solution of NH₃ (2.00 mL, 7M in MeOH). The resulting solution was stirred for 30 min at 25 °C and concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Shield RP18 OBD, 5 μm, 19 x 150 mm; Mobile phase, A: water (10 mM NH₄HCO₃) and B: ACN (20% to 45% in 7 min); Flow rate: 25 mL/min). The collected fraction was lyophilized to give 3-amino-N-[(6S)-2-[(3R,4S)-3-(methoxymethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, *m/z*): 481 [M+H]⁺. ¹H-NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.31 (d, *J* = 8.0 Hz, 1H), 7.59 (br s, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.20-7.17 (m, 3H), 6.24 (d, *J* = 8.4 Hz, 1H), 4.14-4.10 (m, 1H), 3.65-3.60 (m, 1H), 3.55-3.51 (m, 1H), 3.46-3.27 (m, 5H), 3.16-3.10 (m, 2H), 3.02-3.00 (m, 1H), 2.76-2.72 (m, 4H), 2.59 (s, 3H), 2.34-2.32 (m, 4H), 2.02-1.97 (m, 1H), 1.88-1.82 (m, 1H).

Example 27-1. 3-Amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 27-2. 3-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. cis-Benzyl N-[(6S)-2-(3-[[tert-butoxy]carbonyl]amino)-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

[00489] A mixture of cis-tert-butyl N-[4-(fluoromethyl)pyrrolidin-3-yl]carbamate (**Intermediate 54**) (150 mg, 0.687 mmol), benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) (150 mg, 0.342 mmol), RuPhos (16.0 mg, 0.034 mmol), RuPhos Pd G3 (29.0mg, 0.035 mmol) and Cs₂CO₃ (278 mg, 0.853 mmol) in toluene (5 mL) was stirred for 2 h at 95 °C. After cooled to 25 °C, the solids were filtered out. The filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford cis-benzyl N-[(6S)-2-(3-[[tert-butoxy]carbonyl]amino]-4-(fluoromethyl)pyrrolidin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a white solid. MS (ESI, m/z): 499 [M+H]⁺.

Step 2. cis-tert-Butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate

[00490] A mixture of cis-benzyl N-[(6S)-2-(3-[[tert-butoxy]carbonyl]amino)-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (130 mg, 0.256 mmol) and Palladium on carbon (130 mg, 10%) in ethyl acetate (5 mL) was stirred for 2 h at 25 °C under hydrogen atmosphere (balloon). The resulting mixture was filtered. The filtrate was concentrated under reduced pressure to afford cis-tert-butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate as a brown solid. MS (ESI, m/z): 365 [M+H]⁺.

Step 3. cis-tert-Butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate

[00491] HBTU (92.0 mg, 0.243 mmol) was added to a solution of cis-tert-butyl N-[(3R,4R)-1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate (75.0 mg, 0.202 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (45.0 mg, 0.216 mmol) and Et₃N (0.084 mL, 0.603 mmol) in DMA (3 mL). The resulting solution was stirred for 1 h at 25 °C. The mixture was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (0% to 100% in 30 min)) to afford cis-tert-butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate as a yellow solid. MS (ESI, m/z): 555 [M+H]⁺.

Step 4. tert-Butyl N-[(3R,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate and tert-Butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate

[00492] cis-tert-Butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate (70.0 mg, 0.124 mmol) was separated by chiral HPLC (Column: Chiralpak IG, 20 x 250 mm, 5 μ m; Mobile phase, A: hexanes:DCM=3:1(containing 0.1% Et₂NH) and B: IPA (hold 30% in 16 min)). The first eluting isomer (RT=10.09 min) was collected and concentrated under vacuum to afford a white solid, stereochemistry on the pyrrolidine arbitrarily assigned as tert-butyl N-[(3R,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate. The second eluting isomer (RT=13.22 min) was collected and concentrated under vacuum to afford a white solid, stereochemistry on the pyrrolidine arbitrarily assigned as tert-butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate. MS (ESI, m/z): 555 [M+H]⁺.

Step 5. 3-Amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00493] A solution of tert-butyl N-[(3R,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate (15.0 mg, 0.027 mmol) and TFA (0.5 mL) in DCM (1.5 mL) was stirred for 25 min at 25 °C. The resulting mixture was concentrated under vacuum. The residue was treated with a solution of NH₃ (1.0 mL, 7M in MeOH). The resulting solution was stirred for 30 min and then concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep OBD C18 Column, 30 x 150 mm 5 μ m; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (20% to 45% in 7 min)). The collected fraction was lyophilized to afford 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 455 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 8.31 (d, *J* = 8.0 Hz, 1H), 7.58 (d, *J* = 8.4 Hz, 1H), 7.31 (d, *J* = 8.0 Hz, 1H), 7.20 (d, *J* = 8.4 Hz, 1H), 7.16 (br s, 2H), 6.23 (d, *J* = 8.4 Hz, 1H), 4.81-4.47 (m, 2H), 4.11-4.09 (m,


1H), 3.61-3.60 (m, 1H), 3.54-3.46 (m, 2H), 3.33-3.28 (m, 1H), 3.20-3.18 (m, 1H), 2.81-2.68 (m, 4H), 2.59 (s, 3H), 2.02-1.96 (m, 1H), 1.90-1.66 (m, 2H).

Step 6. 3-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00494] A solution of tert-butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(fluoromethyl)pyrrolidin-3-yl]carbamate (15.0 mg, 0.027 mmol) and TFA (0.5 mL) in DCM (1.5 mL) was stirred for 25 min at 25 °C. The resulting mixture was concentrated under vacuum. NH₃ (1.0 mL, 7M in MeOH) was added to the residue. The resulting solution was stirred for 30 min and concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep OBD C18 Column, 30x150mm 5um; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (20% to 45% in 7 min)). The collected fraction was lyophilized to afford 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 455 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.31 (d, *J* = 8.0 Hz, 1H), 7.58 (d, *J* = 7.6 Hz, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.20 (d, *J* = 8.8 Hz, 1H), 7.16 (br s, 2H), 6.23 (d, *J* = 8.4 Hz, 1H), 4.81-4.47 (m, 2H), 4.12-4.09 (m, 1H), 3.61-3.60 (m, 1H), 3.53-3.47 (m, 2H), 3.33-3.27 (m, 1H), 3.21-3.18 (m, 1H), 2.81-2.68 (m, 4H), 2.59 (s, 3H), 2.02-1.96 (m, 1H), 1.90-1.66 (m, 2H).

Example 28-1. 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 28-2. 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. trans-Benzyl N-[(6S)-2-(3-[[[(tert-butoxy)carbonyl]amino]-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

[00495] A mixture of benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) (200 mg, 0.465 mmol), trans-tert-butyl N-[4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate (**Intermediate 55**) (157 mg, 0.603 mmol), RuPhos Pd G3 (38.9 mg, 0.046 mmol), RuPhos (22.0 mg, 0.047 mmol) and Cs₂CO₃ (454 mg, 1.39 mmol) in toluene (4 mL) was stirred for 3 h at 100 °C. After cooling to 26 °C, the solids were filtered out. The filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to give trans-benzyl N-[(6S)-2-(3-[[[(tert-butoxy)carbonyl]amino]-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a yellow solid. MS (ESI, m/z): 541 [M+H]⁺.

Step 2. trans-tert-Butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate

[00496] A mixture of trans-benzyl N-[(6S)-2-(3-[[[(tert-butoxy)carbonyl]amino]-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (140 mg, 0.259 mmol) and Palladium on carbon (70.0 mg, 10%) in ethyl acetate (10 mL) was stirred for 3 h at 26 °C under an atmosphere of hydrogen (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give trans-tert-butyl N-[1-[(6S)-6-amino-5,6,7,8-

tetrahydroquinolin-2-yl]-4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate as a black solid. MS (ESI, m/z): 407 [M+H]⁺.

Step 3. trans-tert-Butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate

[00497] HBTU (112 mg, 0.295 mmol) was added to a solution of trans-tert-butyl N-[1-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate (100 mg, 0.246 mmol), Et₃N (0.103 mL, 0.741 mmol) and 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (61.0 mg, 0.293 mmol) in DMA (2 mL). The resulting solution was stirred for 1 h at 26 °C. The mixture was purified reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (20% to 85% in 25 min)). The collected fraction was concentrated under vacuum to give trans-tert-butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate as a yellow solid. MS (ESI, m/z): 597 [M+H]⁺.

Step 4. 3-Amino-N-[(6S)-2-[(3S,S-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide and 3-Amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00498] trans-tert-Butyl N-[1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate (120 mg, 0.201 mmol) was separated by chiral HPLC (Column: Chiralpak IG, 2 x 25 cm, 5 μm; Mobile phase, A: hexanes : DCM=3:1 (0.1% Et₂NH) and B: EtOH (hold 40% in 20 min)). The first eluting isomer (RT=12.26 min) was collected and concentrated under vacuum to give a yellow solid, stereochemistry on the pyrrolidine arbitrarily assigned as 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide. The second eluting isomer (RT=17.55 min) was collected and concentrated under vacuum to give a yellow solid, stereochemistry on the pyrrolidine arbitrarily assigned as 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide. MS (ESI, m/z): 597 [M+H]⁺.

Step 5. 3-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00499] A solution of tert-butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate (27.0 mg, 0.047 mmol) and TFA (1.00 mL) in DCM (3 mL) was stirred for 30 min at 26 °C. The resulting mixture was concentrated under vacuum. The residue was treated with a solution of NH₃ (2.00 mL, 7M in MeOH). The resulting solution was stirred for 30 min, and then concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep OBD C18, 19 x 150 mm, 5 μm, 13 nm; Mobile phase, A: water (10 mM NH₄HCO₃) and B: ACN (18% to 40% in 8 min), Flow rate: 60 mL/min). The collected fraction was lyophilized to give 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 497 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.31 (d, *J* = 8.4 Hz, 1H), 7.59 (d, *J* = 8.0 Hz, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.19 (d, *J* = 8.8 Hz, 1H), 7.17 (br s, 2H), 6.22 (d, *J* = 8.4 Hz, 1H), 4.14-4.10 (m, 1H), 3.75-3.74 (m, 1H), 3.64-3.55 (m, 3H), 3.49-3.31 (m, 5H), 3.24 (s, 3H), 3.14-3.12 (m, 1H), 2.82-2.72 (m, 4H), 2.59 (s, 3H), 2.03-1.82 (m, 4H).


Step 6. 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00500] A solution of tert-butyl N-[(3R,4R)-1-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-4-(2-methoxyethoxy)pyrrolidin-3-yl]carbamate (26.0 mg, 0.045 mmol) and TFA (1.00 mL) in DCM (3 mL) was stirred for 30 min at 25 °C. The resulting mixture was concentrated under vacuum. NH₃ (2.00 mL, 7M in MeOH) was added to the residue. The resulting solution was stirred for 30 min, and then concentrated under vacuum. The residue was purified by prep-HPLC (Column: X Bridge Prep OBD C18, 30 x 150 mm, 5 μm; Mobile phase, A: water (10 mM NH₄HCO₃) and B: ACN (18% to 40% in 8 min), Flow rate: 60 mL/min). The collected fraction was lyophilized to give 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 497 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.31 (d, *J* = 8.4 Hz, 1H), 7.59 (d, *J* = 8.0 Hz, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.19 (d, *J* = 8.8 Hz, 1H), 7.17 (br s, 2H), 6.22 (d, *J* = 8.4 Hz, 1H), 4.15-4.11 (m,

1H), 3.75-3.74 (m 1H), 3.63-3.59 (m, 3H), 3.48-3.33 (m, 5H), 3.24 (s, 3H), 3.14-3.11 (m, 1H), 2.80-2.72 (m, 4H), 2.59 (s, 3H), 2.01-1.82 (m, 4H).

Example 29-1. 3-Amino-N-[(6S)-2-[(9S)-9-amino-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 29-2. 3-Amino-N-[(6S)-2-[(9R)-9-amino-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl N-[(6S)-2-(9-[[[(tert-butoxy)carbonyl]amino]-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate

[00501] A mixture of benzyl N-[(6S)-2-(trifluoromethanesulfonyloxy)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-2**) (180 mg, 0.410 mmol), tert-butyl N-[1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate (**Intermediate 59**) (150 mg, 0.614 mmol), RuPhos (19.0 mg, 0.041 mmol), RuPhos Pd G3 (34.0 mg, 0.041 mmol) and Cs₂CO₃ (334 mg, 1.025 mmol) in toluene (10 mL) was stirred for 3 h at 95 °C. The mixture was allowed to cool down to 25 °C, filtered, and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to afford benzyl N-[(6S)-2-(9-[[[(tert-butoxy)carbonyl]amino]-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate as a white solid. MS (ESI, *m/z*): 525 [M+H]⁺.

Step 2. tert-Butyl N-[7-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate

[00502] A mixture of benzyl N-[(6S)-2-(9-[[[(tert-butoxy)carbonyl]amino]-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl)-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (180 mg, 0.336 mmol) and Pd/C (180 mg, 10%) in ethyl acetate (5 mL) was stirred for 2 h at 25 °C. The resulting mixture was filtered, and the filter cake was washed with ethyl acetate (3 x 5 mL). The filtrate was concentrated under reduced pressure to afford tert-butyl N-[7-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate as a brown solid. MS (ESI, m/z): 391 [M+H]⁺.

Step 3. tert-Butyl N-[7-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate

[00503] HBTU (114 mg, 0.301 mmol) was added to a solution of tert-butyl N-[7-[(6S)-6-amino-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate (100 mg, 0.251 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (57.0 mg, 0.274 mmol) and Et₃N (0.104 mL, 0.751 mmol) in DMA (3 mL). The reaction was stirred for 3 h at 25 °C. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.1% TFA) and B: ACN (0% to 60% in 30 min)) to afford tert-butyl N-[7-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate as a yellow solid. MS (ESI, m/z): 581 [M+H]⁺.

Step 4. tert-Butyl N-[(9S)-7-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate and tert-Butyl N-[(9R)-7-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate

[00504] tert-Butyl N-[7-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate (100 mg, 0.169 mmol) was separated by chiral HPLC (Column: Chiralpak IF, 2 x 25 cm, 5 μm; Mobile phase, A: MTBE (0.1% Et₂NH) and B: EtOH (hold 30% in 24 min); Flow rate: 12 mL/min). The first eluting isomer (RT=15.97 min) was collected and concentrated under vacuum to afford a yellow solid, stereochemistry on the pyrrolidine arbitrarily assigned as tert-butyl N-[(9S)-7-[(6S)-6-[3-amino-

6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate. MS (ESI, m/z): 581 $[M+H]^+$.

[00505] The second eluting isomer (RT=20.62 min) was collected and concentrated under vacuum to afford a yellow solid, stereochemistry on the pyrrolidine arbitrarily assigned as tert-butyl N-[(9R)-7-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate. MS (ESI, m/z): 581 $[M+H]^+$.

Step 5. 3-Amino-N-[(6S)-2-[(9S)-9-amino-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00506] A solution of tert-butyl N-[(9S)-7-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate (40 mg, 0.068 mmol) and TFA (0.50 mL) in DCM (1.5 mL) was stirred for 30 min at 25 °C. The resulting solution was concentrated under vacuum. NH_3 (2.0 mL, 7M in MeOH) was added to the residue. The resulting solution was stirred for 30 min at 25 °C, and then was concentrated under vacuum. The residue was purified by prep-HPLC (Column: X Bridge Prep C18 OBD, 19 x 150 mm, 5 μ m; Mobile phase, A: water (0.05% NH_4HCO_3) and B: ACN (20% to 35% in 7 min)). The collected fraction was lyophilized to afford 3-amino-N-[(6S)-2-[(9S)-9-amino-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 481 $[M+H]^+$. ^1H -NMR ($CDCl_3$, 400 MHz) δ (ppm): 7.82 (d, J = 8.0 Hz, 1H), 7.22-7.20 (m, 1H), 6.20 (d, J = 8.0 Hz, 1H), 6.06-6.02 (m, 2H), 5.52 (d, J = 7.2 Hz, 1H), 4.48-4.42 (m, 1H), 4.14-4.09 (m, 4H), 3.82-3.69 (m, 2H), 3.53-3.50 (m, 2H), 3.24-3.20 (m, 1H), 3.09-3.08 (m, 1H), 2.94-2.86 (m, 2H), 2.70-2.64 (m, 4H), 2.26-2.23 (m, 1H), 1.98-1.91 (m, 1H).


Step 6. 3-Amino-N-[(6S)-2-[(9R)-9-amino-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00507] A solution of tert-butyl N-[(9R)-7-[(6S)-6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinolin-2-yl]-1,4-dioxo-7-azaspiro[4.4]nonan-9-yl]carbamate (40.0 mg, 0.068 mmol) and TFA (0.50 mL) in DCM (1.5 mL) was stirred for 30 min at 25 °C. The resulting solution was concentrated under vacuum. NH_3 (2.0 mL, 7M in MeOH) was added to the

residue. The resulting solution was stirred for 30 min at 25 °C, and then was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep C18 OBD, 19 x 150 mm, 5 µm; Mobile phase, A: water (0.05% NH₄HCO₃) and B: ACN (20% to 35% in 7 min)). The collected fraction was lyophilized to afford 3-amino-N-[(6S)-2-[(9R)-9-amino-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, *m/z*): 481 [M+H]⁺. ¹H-NMR (CDCl₃, 400 MHz) δ(ppm): 7.82 (d, *J* = 8.4 Hz, 1H), 7.22-7.20 (m, 1H), 6.20 (d, *J* = 8.4 Hz, 1H), 6.06-6.00 (m, 2H), 5.53 (d, *J* = 6.8 Hz, 1H), 4.48-4.42 (m, 1H), 4.14-4.09 (m, 4H), 3.82-3.67 (m, 2H), 3.53-3.501 (m, 2H), 3.27-3.22 (m, 1H), 3.10-3.08 (m, 1H), 2.95-2.90 (m, 2H), 2.70-2.65 (m, 4H), 2.24-2.20 (m, 1H), 1.93-1.87 (m, 1H).

Example 30-1. 3-Amino-N-[(6S)-3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide and

Example 30-2. 3-Amino-N-[(6R)-3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl 4-(6-[[[(benzyloxy)carbonyl]amino]-5,6,7,8-tetrahydroquinolin-2-yl])piperazine-1-carboxylate

[00508] A mixture of benzyl N-[2-[(trifluoromethane)sulfonyloxy]-5,6,7,8-tetrahydroquinolin-6-yl]carbamate (**Intermediate 47-1**) (400 mg, 0.930 mmol), tert-butyl piperazine-1-carboxylate (208 mg, 1.12 mol), RuPhos (87.0 mg, 0.190 mmol), RuPhos Pd G3 (76.0 mg, 0.090 mmol), and Cs₂CO₃ (758 mg, 2.33 mmol) in toluene (15 mL) was stirred for 3 h at 95 °C. After cooling to

25 °C, the solid was filtered out. The filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to afford tert-butyl 4-(6-[[[(benzyloxy)carbonyl]amino]-5,6,7,8-tetrahydroquinolin-2-yl])piperazine-1-carboxylate as a white solid. MS (ESI, m/z): 467 [M+H]⁺.

Step 2. tert-Butyl 4-(6-[[[(benzyloxy)carbonyl]amino]-3-fluoro-5,6,7,8-tetrahydroquinolin-2-yl])piperazine-1-carboxylate

[00509] A solution of SelectFluor (683 mg, 1.93 mmol) in ACN (10 mL) was added to a stirring solution of tert-butyl 4-(6-[[[(benzyloxy)carbonyl]amino]-5,6,7,8-tetrahydroquinolin-2-yl])piperazine-1-carboxylate (600 mg, 1.29 mmol) in DCM (30 mL) and MeOH (30 mL). The resulting solution was stirred for 1 h at -20 °C. The resulting mixture was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Prep C18 OBD, 5 µm, 19 x 150 mm; Mobile Phase, A: water (0.05% TFA) and B: ACN (55% to 85% in 7 min)). The collected fraction was concentrated under vacuum to give tert-butyl 4-(6-[[[(benzyloxy)carbonyl]amino]-3-fluoro-5,6,7,8-tetrahydroquinolin-2-yl])piperazine-1-carboxylate as an off-white solid. MS (ESI, m/z): 485 [M+H]⁺.

Step 3. tert-Butyl 4-(6-amino-3-fluoro-5,6,7,8-tetrahydroquinolin-2-yl)piperazine-1-carboxylate

[00510] A mixture of tert-butyl 4-(6-[[[(benzyloxy)carbonyl]amino]-3-fluoro-5,6,7,8-tetrahydroquinolin-2-yl])piperazine-1-carboxylate (70.0 mg, 0.140 mmol) and palladium on carbon (70.0 mg, 10%) in ethyl acetate (5 mL) was stirred for 1 h at 25 °C under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to give tert-butyl 4-(6-amino-3-fluoro-5,6,7,8-tetrahydroquinolin-2-yl)piperazine-1-carboxylate as an off-white solid. MS (ESI, m/z): 351 [M+H]⁺.

Step 4. tert-Butyl 4-(6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-3-fluoro-5,6,7,8-tetrahydroquinolin-2-yl)piperazine-1-carboxylate

[00511] HBTU (58.0 mg, 0.150 mmol) was added to a solution of 4-(6-amino-3-fluoro-5,6,7,8-tetrahydroquinolin-2-yl)piperazine-1-carboxylate (45.0 mg, 0.130 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (32.0 mg, 0.150 mmol) and Et₃N (54.0 µL, 0.390 mmol) in DMA (2 mL). The resulting solution was stirred for 30 min at 25 °C. The mixture was

purified via reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH_4HCO_3) and B: ACN (0% to 85% within 25 min)). The collected fraction was concentrated under vacuum to give tert-butyl 4-(6-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-3-fluoro-5,6,7,8-tetrahydroquinolin-2-yl)piperazine-1-carboxylate as an off-white solid. MS (ESI, m/z): 541 $[\text{M}+\text{H}]^+$.


Step 5. 3-Amino-N-[(6S)-3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide and 3-Amino-N-[(6R)-3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00512] A solution of tert-butyl 4-[3-amino-6-(3-amino-6-methyl-1-benzothiophene-2-amido)-5,6,7,8-tetrahydroquinolin-2-yl]piperazine-1-carboxylate (40.0 mg, 0.070 mmol) and TFA (1.00 mL) in DCM (3 mL) was stirred for 1 h at 25 °C. The resulting mixture was concentrated under vacuum to give TFA salt of 3-amino-N-[3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a yellow solid. The enantiomers were separated by chiral HPLC (Column: Chiralpak IC, 2 x 25 cm, 5 μm ; Mobile Phase, A: Hex (0.2% Et_2NH) and B: EtOH (hold 50% in 20 min); Flow rate: 15 mL/min). The first eluting isomer (RT =11.3 min) was concentrated under vacuum and then lyophilized to afford an off-white solid, stereochemistry arbitrarily assigned as 3-amino-N-[(6S)-3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide. MS (ESI, m/z): 441 $[\text{M}+\text{H}]^+$. ^1H -NMR ($\text{DMSO}-d_6$, 300 MHz) δ (ppm): 8.30 (d, J = 8.4 Hz, 1H), 7.60 (d, J = 7.5 Hz, 1H), 7.32-7.22 (m, 2H), 7.16 (br s, 2H), 4.14 (br s, 1H), 3.25-3.21 (m, 4H), 2.83-2.72 (m, 8H), 2.58 (s, 3H), 1.98-1.86 (m, 2H).

[00513] The second eluting isomer (RT =16.9 min) was concentrated under vacuum and then lyophilized with ACN and water to afford an off-white solid, stereochemistry arbitrarily assigned as 3-amino-N-[(6R)-3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide. MS (ESI, m/z): 441 $[\text{M}+\text{H}]^+$. ^1H -NMR ($\text{DMSO}-d_6$, 300 MHz) δ (ppm): 8.30 (d, J = 8.1 Hz, 1H), 7.60 (d, J = 7.8 Hz, 1H), 7.32-7.21 (m, 2H), 7.16 (br s, 2H), 4.15 (br s, 1H), 3.22-3.20 (m, 4H), 2.83-2.72 (m, 8H), 2.58 (s, 3H), 2.07-1.83 (m, 2H).

Example 31-1. 3-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide and

Example 31-2. 3-Amino-N-[(6R)-2-[(3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl N-[2-[(3S,4S)-3-[[tert-butoxy]carbonyl]amino]-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]carbamate

[00514] A solution of benzyl N-(2-chloro-5,6,7,8-tetrahydroquinazolin-6-yl)carbamate (**Intermediate 57**) (300 mg, 0.944 mmol), tert-butyl N-[(3S,4S)-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate (**Intermediate 56**) (276 mg, 1.133 mmol) and DIEA (0.312 mL, 1.89 mmol) in DMA (10 mL) was microwaved for 1 h at 150 °C. After cooling to 25 °C, the residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (0% to 70% in 20 min)). The collected fraction was concentrated under vacuum to afford benzyl N-[2-[(3S,4S)-3-[[tert-butoxy]carbonyl]amino]-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]carbamate as a brown solid. MS (ESI, m/z): 526 [M+H]⁺.

Step 2. tert-Butyl N-[(3S,4S)-1-(6-amino-5,6,7,8-tetrahydroquinazolin-2-yl)-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate

[00515] A mixture of benzyl N-[2-[(3S,4S)-3-[[tert-butoxy]carbonyl]amino]-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]carbamate (300 mg, 0.542 mmol) and Palladium on carbon (300 mg, 10%) in ethyl acetate (20 mL) was stirred for 2 h at 20 °C under

hydrogen atmosphere (balloon). The resulting mixture was filtered and the filter cake was washed with ethyl acetate (3x10 mL). The filtrate was concentrated under vacuum to afford tert-butyl N-[(3S,4S)-1-(6-amino-5,6,7,8-tetrahydroquinazolin-2-yl)-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate as a brown solid. MS (ESI, m/z): 392 [M+H]⁺.

Step 3. tert-Butyl N-[(3S,4S)-1-[6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate

[00516] HBTU (140 mg, 0.370 mmol) was added to a solution of tert-butyl N-[(3S,4S)-1-(6-amino-5,6,7,8-tetrahydroquinazolin-2-yl)-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate (100 mg, 0.230 mmol), Et₃N (0.096 mL, 0.690 mmol) and 3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-carboxylic acid (**Intermediate 21**) (70.0 mg, 0.300 mmol) in DMA (3 mL). The resulting solution was stirred for 3 h at 25 °C. The residue was purified by reverse phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (10% to 60% in 30 min)). The collected fraction was concentrated to give tert-butyl N-[(3S,4S)-1-[6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate as a white solid. MS (ESI, m/z): 596 [M+H]⁺.

Step 4. tert-Butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate and tert-Butyl N-[(3S,4S)-1-[(6R)-6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate

[00517] tert-Butyl N-[(3S,4S)-1-[6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate (65.0 mg, 0.109 mmol) was separated by chiral HPLC (Column: Chiralpak IA, 2 x 25 cm, 5 μm; Mobile phase, A: hexanes:DCM=3:1 and B: EtOH (hold 50% in 19 min); Flow rate: 16 mL/min). The first eluting isomer (RT=11.42 min) was collected and concentrated to give a white solid, stereochemistry of the amide arbitrarily assigned as tert-butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate. The second eluting isomer (RT=15.96 min) was collected and concentrated under vacuum to give a white solid, stereochemistry of the amide arbitrarily assigned as tert-butyl N-[(3S,4S)-1-[(6R)-6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-

tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate. MS (ESI, m/z): 596 [M+H]⁺.

Step 5. 3-Amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide

[00518] A solution of tert-butyl N-[(3S,4S)-1-[(6S)-6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate (28.0 mg, 0.05 mmol) and TFA (0.5 mL) in DCM (1.5 mL) was stirred for 30 min at 25 °C. The resulting mixture was concentrated under vacuum. The residue was treated with a solution of NH₃ (1.0 mL, 7M in MeOH). The resulting solution was stirred for 30 min and concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Shield RP18 OBD, 30 x150 mm, 5 μm; Mobile phase, A: water (10 mM NH₄HCO₃) and B: ACN (25% to 45% in 7 min); Flow rate: 60 mL/min). The collected fraction was lyophilized to give 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 496 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400 MHz) δ (ppm): 8.06 (s, 1H), 7.64 (br s, 1H), 7.04 (s, 1H), 6.82 (br s, 2H), 4.14-4.11 (m, 1H), 3.78-3.73 (m, 1H), 3.71-3.70 (m, 2H), 3.57-3.53 (m, 1H), 3.37-3.32 (m, 5H), 3.25-3.22 (m, 1H), 2.82-2.62 (m, 7H), 2.01-1.98 (m, 1H), 1.91-1.83 (m, 1H), 1.74 (br s, 2H), 1.12-1.08 (m, 6H).



Step 6. 3-Amino-N-[(6R)-2-[(3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide

[00519] A solution of tert-butyl N-[(3S,4S)-1-[(6R)-6-[3-amino-4,6-dimethylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroquinazolin-2-yl]-4-(propan-2-yloxy)pyrrolidin-3-yl]carbamate (27.0 mg, 0.05 mmol) and TFA (0.5 mL) in DCM (1.5 mL) was stirred for 30 min at 25 °C. The resulting mixture was concentrated under vacuum. NH₃ (1.0 mL, 7M in MeOH) was added to the residue. The resulting solution was stirred for 30 min and concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Shield RP18 OBD, 30 x150 mm, 5 μm; Mobile phase, A: water (10 mM NH₄HCO₃) and B: ACN (25% to 50% in 7 min); Flow rate: 60 mL/min). The collected fraction was lyophilized to give 3-amino-N-[(6R)-2-[(3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinazolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 496 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400

MHz) δ (ppm): 8.07 (s, 1H), 7.65 (br s, 1H), 7.05 (s, 1H), 6.83 (br s, 2H), 4.16-4.13 (m, 1H), 3.79-3.74 (m, 1H), 3.72-3.71 (m, 2 H), 3.57-3.53 (m, 1H), 3.38-3.33 (m, 5H), 3.26-3.23 (m, 1H), 2.83-2.62 (m, 7H), 2.01-1.99 (m, 1 H), 1.92-1.83 (m, 1H), 1.74 (br s, 2H), 1.12-1.08 (m, 6H).

[00520] The following examples in Table 22 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Examples 31-1 and 31-2.


Table 22:

Example Number	Structure and Compound Name	LRMS m/z [M+H] ⁺	¹ H NMR
31-3 ¹	 3-amino-6-methyl-N-[(6'S)-2'-(piperazin-1-yl)-6',7'-dihydro-5'H-spiro[cyclopropane-1,8'-quinoline]-6'-yl]thieno[2,3-b]pyridine-2-carboxamide	449	(DMSO-d ₆ , 400 MHz) δ (ppm): 8.31 (d, J = 8.4 Hz, 1H), 7.52 (d, J = 7.6 Hz, 1H), 7.31 (d, J = 8.4 Hz, 1H), 7.20 (d, J = 8.4 Hz, 1H), 7.16 (br s, 2H), 6.58-6.51 (m, 1H), 4.33-4.27 (m, 1H), 3.44-3.27 (m, 4H), 2.88-2.82 (m, 2H), 2.75-2.68 (m, 4H), 2.59 (s, 3H), 2.39 (br s, 1H), 2.29-2.22 (m, 1H), 1.53-1.45 (m, 2H), 0.92-0.86 (m, 1H), 0.71-0.69 (m, 2H).
31-4 ¹	 3-amino-6-methyl-N-[(6'R)-2'-(piperazin-1-yl)-6',7'-dihydro-5'H-spiro[cyclopropane-1,8'-quinoline]-6'-yl]thieno[2,3-b]pyridine-2-carboxamide	449	(DMSO-d ₆ , 400 MHz) δ (ppm): 8.31 (d, J = 8.0 Hz, 1H), 7.52 (d, J = 7.6 Hz, 1H), 7.31 (d, J = 8.4 Hz, 1H), 7.20 (d, J = 8.4 Hz, 1H), 7.16 (br s, 2H), 6.52 (d, J = 8.4 Hz, 1H), 4.32-4.28 (m, 1H), 3.43-3.28 (m, 4H), 2.88-2.82 (m, 2H), 2.75-2.64 (m, 4H), 2.59 (s, 3H), 2.39-2.34 (m, 1H), 2.29-2.22 (m, 1H), 1.53-1.40 (m, 2H), 0.92-0.86 (m, 1H), 0.71-0.69 (m, 2H).

¹ **Notes on procedures:** In Step 1, the triflate **Intermediate 47-4** was used. In Step 4, the enantiomers were separated by chiral HPLC using the column Chiralpak IC and Mobile Phase 10% EtOH/MTBE (containing 10 mmol/L NH₃ in MeOH) to afford the precursor to Example 31-3 as the first eluted isomer (stereochemistry arbitrarily assigned); and the precursor to Example 31-4 as the second eluted isomer (stereochemistry arbitrarily assigned).

Example 32-1. 3-Amino-6-methyl-N-[(7S)-3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide and

Example 32-2. 3-Amino-6-methyl-N-[(7R)-3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide



Step 1. tert-Butyl 4-(7-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroisoquinolin-3-yl)piperazine-1-carboxylate

[00521] HBTU (70.0 mg, 0.180 mmol) was added to a solution of tert-butyl 4-(7-amino-5,6,7,8-tetrahydroisoquinolin-3-yl)piperazine-1-carboxylate (**Intermediate 58**) (30 mg, 0.090 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (20.0 mg, 0.100 mmol) and Et³N (0.038 mL, 0.270 mmol) in DMA (2 mL). The resulting solution was stirred for 30 min at 25 °C. The mixture was purified via reverse phase chromatography (Column: X Bridge C18, 19 x 150 mm, 5 µm; Mobile Phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (10% to 80% in 30 min)). The collected fraction was concentrated under vacuum to give tert-butyl 4-(7-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroisoquinolin-3-yl)piperazine-1-carboxylate as a white solid. MS (ESI, m/z): 523 [M+H]⁺.

Step 2. 3-Amino-6-methyl-N-[3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide


[00522] A solution of tert-butyl 4-(7-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,7,8-tetrahydroisoquinolin-3-yl)piperazine-1-carboxylate (35.0 mg, 0.070 mmol) and TFA (0.50 mL) in DCM (1.50 mL) was stirred for 30 min at 25 °C. The reaction mixture was concentrated under vacuum. Then a solution of NH₃ (2 mL, 7M in methanol) was added. The resulting solution was stirred for 30 min at 25 °C. The resulting solution was concentrated under vacuum to give 3-amino-6-methyl-N-[3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, m/z): 423 [M+H]⁺.

Step 3. 3-Amino-6-methyl-N-[(7S)-3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide and 3-Amino-6-methyl-N-[(7R)-3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide

[00523] The racemate 3-amino-6-methyl-N-[3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide (25.0 mg, 0.059 mmol) was separated into its enantiomers by chiral HPLC (Column: Chiralpak IG, 2 x 25 cm, 5 μ m; Mobile phase, A: MTBE (containing 0.1% Et₂NH) and B: ethanol (hold 50% in 20 min)). The first eluting isomer (RT= 12.37 min) was concentrated under vacuum and then lyophilized to give a white solid, stereochemistry arbitrarily assigned as 3-amino-6-methyl-N-[(7S)-3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide. MS (ESI, m/z): 423 [M+H]⁺. ¹H NMR (CD₃OD, 400 MHz) δ (ppm): 8.22 (d, *J* = 8.0 Hz, 1H), 7.89 (s, 1H), 7.33 (d, *J* = 8.0 Hz, 1H), 6.63 (s, 1H), 4.25-4.23 (m, 1 H), 3.49-3.43 (m, 4 H), 3.06-2.98 (m, 1 H), 2.97-2.92 (m, 6 H), 2.82-2.69 (m, 1 H), 2.66 (s, 3 H), 2.15-2.13 (m, 1 H), 1.89-1.81 (m, 1H). The second eluting isomer (RT= 17.03 min) was concentrated under vacuum and then lyophilized to give a white solid, stereochemistry arbitrarily assigned as 3-amino-6-methyl-N-[(7R)-3-(piperazin-1-yl)-5,6,7,8-tetrahydroisoquinolin-7-yl]thieno[2,3-b]pyridine-2-carboxamide. MS (ESI, m/z): 423 [M+H]⁺. ¹H NMR (CD₃OD, 400 MHz) δ (ppm): 8.22 (d, *J* = 8.0 Hz, 1H), 7.89 (s, 1H), 7.33 (d, *J* = 8.0 Hz, 1H), 6.63 (s, 1H), 4.25-4.23 (m, 1 H), 3.49-3.43 (m, 4 H), 3.15-2.98 (m, 1 H), 2.97-2.93 (m, 6 H), 2.82-2.72 (m, 1 H), 2.76 (s, 3 H), 2.15-2.13 (m, 1 H), 1.89-1.81 (m, 1H).

Example 33-1. (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl-4,4-d2)-6-methylthieno[2,3-b]pyridine-2-carboxamide

Example 33-2. (S)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl-4,4-d2)-6-methylthieno[2,3-b]pyridine-2-carboxamide



Step 1. 3-Amino-N-(7-bromo-5-fluorochroman-3-yl-4,4-d2)-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00524] A solution of 7-bromo-5-fluorochroman-4,4-d2-3-amine (**Intermediate 5-2**) (340 mg, 1.370 mmol), 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (342 mg, 1.645 mmol), Et₃N (0.287 ml, 2.056 mmol), and HBTU (624 mg, 1.645 mmol) in DMA (5 mL) was stirred at room temperature for 4 h. The reaction was reduced in volume by half under reduced pressure. EtOAc (5 mL) and H₂O (3 mL) were added. The organic layer was separated, dried over Na₂SO₄, filtered, and concentrated. The residue was purified by silica gel chromatography (eluting with 2% to 100% EtOAc/hexanes). Desired fractions were combined and concentrated to afford 3-amino-N-(7-bromo-5-fluorochroman-3-yl-4,4-d2)-6-methylthieno[2,3-b]pyridine-2-carboxamide as a yellow solid. MS: (ESI, m/z): 438 [M+H]⁺. ¹H NMR (CDCl₃, 300 MHz) δ(ppm): 7.61-7.83 (m, 1H), 7.03-7.13 (m, 1H), 6.69-6.90 (m, 2H), 6.00 (br s, 2H), 5.59 (br d, *J* = 6.7 Hz, 1H), 4.47-4.82 (m, 1H), 3.95-4.29 (m, 3H), 2.85-2.86 (m, 1H), 2.67-2.79 (m, 3H), 2.59 (d, *J* = 3.2 Hz, 3H).

Step 2. tert-Butyl 4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluorochroman-7-yl-4,4-d2)piperazine-1-carboxylate



[00525] A solution of 3-amino-N-(7-bromo-5-fluorochroman-3-yl-4,4-d2)-6-methylthieno[2,3-b]pyridine-2-carboxamide (150 mg, 0.342 mmol) was combined with tert-butyl piperazine-1-carboxylate (127 mg, 0.684 mmol) in anhydrous DMA (3.42 mL). A 1M in THF solution of LiHMDS (3.42 mL, 3.42 mmol) was added and the reaction was flushed with N₂ for 20 min. Pd-PEPPSI-IPent (34.2 mg, 0.034 mmol) was added and the reaction was stirred at 60 °C for 16 h. The reaction was diluted with 10 mL of EtOAc and washed with 3 mL of H₂O. The organic layer was separated, dried over anhydrous Na₂SO₄, filtered, and concentrated. The residue was purified by silica gel chromatography (eluting with 5% to 80% EtOAc/hexanes). Desired fractions were combined and concentrated to afford tert-butyl 4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluorochroman-7-yl-4,4-d2)piperazine-1-carboxylate as an off-white solid. MS: (ESI, m/z): 544 [M+H]⁺.

Step 3. 3-Amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl-4,4-d2)-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00526] HCl (4M in 1,4-dioxane, 0.276 mL, 1.104 mmol) was added to a suspension of tert-butyl 4-(3-(3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamido)-5-fluorochroman-7-yl-4,4-d₂)piperazine-1-carboxylate in EtOAc (1 mL). The reaction was stirred at 22 °C for 16 h. The reaction was concentrated to dryness under vacuum. The residue was purified by chiral HPLC using chiral column Chiralpak IA (Mobile Phase A: hexanes (0.1% Et₂NH), and B: EtOH (0.1% Et₂NH); 50% B) to afford Peak 1 whose stereochemistry was arbitrarily assigned as (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl-4,4-d₂)-6-methylthieno[2,3-b]pyridine-2-carboxamide and Peak 2 whose stereochemistry was arbitrarily assigned as (S)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl-4,4-d₂)-6-methylthieno[2,3-b]pyridine-2-carboxamide. MS: (ESI, m/z): 444 [M+H]⁺.

[00527] The following examples in Table 23 were prepared using standard chemical manipulations and procedures similar to those used for the preparation of Examples 33-1 and 33-2.


Table 23:

Example Number	Structure and Compound Name	LRMS <i>m/z</i> [M+H] ⁺	¹ H NMR
33-3 ¹	 3-amino-N-[(3R)-7-[(1R,6S)-3,8-diazabicyclo[4.2.0]octan-8-yl]-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	450	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.31 (d, <i>J</i> = 8.0 Hz, 1H), 7.50 (d, <i>J</i> = 7.6 Hz, 1H), 7.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.21 (br s, 2H), 6.87 (d, <i>J</i> = 8.4 Hz, 1H), 6.09-6.06 (m, 1H), 5.90 (s, 1H), 4.29-4.26 (m, 1H), 4.16-4.12 (m, 1H), 3.82-3.75 (m, 2H), 3.65-3.45 (m, 2H), 3.32-3.18 (m, 1H), 3.08-3.05 (m, 1H), 2.89-2.82 (m, 4H), 2.59 (s, 3H), 2.44-2.38 (m, 1H), 1.92-1.88 (m, 1H), 1.80-1.63 (m, 1H).
33-4 ¹	 3-amino-N-[(3R)-7-[(1S,6R)-3,8-diazabicyclo[4.2.0]octan-8-yl]-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide	450	(DMSO-d ₆ , 400 MHz) δ(ppm): 8.33 (d, <i>J</i> = 8.4 Hz, 1H), 7.50 (d, <i>J</i> = 7.6 Hz, 1H), 7.32 (d, <i>J</i> = 8.4 Hz, 1H), 7.21 (br s, 2H), 6.87 (d, <i>J</i> = 8.0 Hz, 1H), 6.10-6.07 (m, 1H), 5.91 (s, 1H), 4.27-4.25 (m, 1H), 4.16-4.13 (m, 1H), 3.82-3.77 (m, 2H), 3.65-3.47 (m, 2H), 3.32-3.30 (m, 1H), 3.09-3.06 (m, 1H), 2.91-2.84 (m, 4H), 2.59 (s, 3H), 2.45-2.40 (m, 1H), 1.93-1.89 (m, 1H), 1.70-1.61 (m, 1H).

	methylthieno[2,3-b]pyridine-2-carboxamide		

¹ **Notes on procedures:** In Step 1, 7-bromochroman-3-amine was used as the amine coupling partner to afford the racemate 3-amino-N-(7-bromochroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide. The enantiomers were separated by SFC using the chiral column ChiralArt Amylose-SA and Mobile Phase, A: CO₂, 65% and B: EtOH:DCM=1:1 to afford (R)-3-amino-N-(7-bromochroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as the first eluted isomer and (S)-3-amino-N-(7-bromochroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide as the second eluted isomer. In Step 2, cis-tert-butyl 3,8-diazabicyclo[4.2.0]octane-3-carboxylate was used as the amine coupling partner to afford cis-tert-butyl 8-[(3R)-3-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-3,4-dihydro-2H-1-benzopyran-7-yl]-3,8-diazabicyclo[4.2.0]octane-3-carboxylate. The cis isomers were separated by chiral HPLC using the chiral column Chiralpak IE and mobile phase 10% EtOH/MTBE (0.1% Et₂NH) to afford the precursor to Example 33-3 as the first eluted isomer (stereochemistry arbitrarily assigned) and the precursor to Example 33-4 as the second eluted isomer (stereochemistry arbitrarily assigned). In Step 3, boc-deprotection was accomplished by treatment with TFA in DCM.

Example 34. 3-Amino-N-[(3R)-7-(4-cyano-1,1-dioxo-1λ6-thian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide



[00528] n-BuLi (4.40 mL, 2.5 M in n-hexane) was added to a solution of benzyl (R)-(7-bromochroman-3-yl)carbamate (**Intermediate 2**) (800 mg, 2.16 mmol) in THF (20 mL) at -78 °C. The resulting solution was stirred for 30 min at -78 °C. Then a solution of thian-4-one (1.28 g, 11.0 mmol) in THF (3 mL) was added at -78 °C. The resulting solution was stirred for 1 h at -78 °C. The reaction was then quenched with water (30 mL). The resulting mixture was extracted with ethyl acetate (3 x 30 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:2 ethyl acetate/pet. ether) to give benzyl N-[(3R)-7-(4-hydroxythian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]carbamate as a light yellow solid. MS: (ESI, m/z): 400 [M+H]⁺.

Step 2. Benzyl N-[(3R)-7-(4-cyanothian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]carbamate

[00529] A solution of TMSCN (120 mg, 1.21 mmol) in DCM (1 mL) was added to a mixture of InBr₃ (11.0 mg, 0.030 mmol) and DCM (1 mL). Then a solution of benzyl N-[(3R)-7-(4-hydroxythian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]carbamate (120 mg, 0.300 mmol) in DCM (1 mL) was added over 30 min. The resulting mixture was stirred for 30 min at 29 °C. Three batches were thus run in parallel and combined for quenching with water (10 mL). The resulting mixture was extracted with DCM (3 x 15 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by a silica gel chromatography (eluting with 1:3 ethyl acetate/pet. ether) to give benzyl N-[(3R)-7-(4-cyanothian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]carbamate as an off-white solid. MS: (ESI, m/z): 409 [M+H]⁺.

Step 3. Benzyl N-[(3R)-7-(4-cyano-1,1-dioxo-1λ6-thian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]carbamate

[00530] A mixture of benzyl N-[(3R)-7-(4-cyanothian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]carbamate (160 mg, 0.390 mmol) and oxone (650 mg, 3.87 mmol) in 2-butanone (10 mL) was stirred for 12 h at 40 °C. The solids were filtered out. The filtrate was concentrated under vacuum. The residue was purified via a silica gel chromatography (eluting with 1:1 ethyl acetate/pet. ether) to give benzyl N-[(3R)-7-(4-cyano-1,1-dioxo-1λ6-thian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]carbamate as a white solid. MS: (ESI, m/z): 441 [M+H]⁺.


Step 4. 4-[(3R)-3-Amino-3,4-dihydro-2H-1-benzopyran-7-yl]-1,1-dioxo-1λ6-thiane-4-carbonitrile

[00531] A mixture of benzyl N-[(3R)-7-(4-cyano-1,1-dioxo-1λ6-thian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]carbamate (80.0 mg, 0.180 mmol) and palladium on carbon (80.0 mg, 10%) in ethyl acetate (10 mL) was stirred for 1 h at 29 °C under hydrogen atmosphere (balloon). The solids were filtered out. The filtrate was concentrated under vacuum to afford 4-[(3R)-3-amino-3,4-dihydro-2H-1-benzopyran-7-yl]-1,1-dioxo-1λ6-thiane-4-carbonitrile as an off-white solid. MS: (ESI, m/z): 307 $[M+H]^+$.

Step 5. 3-Amino-N-[(3R)-7-(4-cyano-1,1-dioxo-1λ6-thian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

[00532] HBTU (68.0 mg, 0.180 mmol) was added to a solution of 4-[(3R)-3-amino-3,4-dihydro-2H-1-benzopyran-7-yl]-1,1-dioxo-1λ6-thiane-4-carbonitrile (50.0 mg, 0.160 mmol), Et₃N (0.067 mL, 0.480 mmol) and 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (34.0 mg, 0.160 mmol) in DMA (2 mL). The resulting solution was stirred for 30 min at 29 °C. The mixture was purified by prep-HPLC (Column: XBridge Prep C18 OBD, 19 x 150 mm 5 μm; Mobile phase, A: water (10 mM NH₄HCO₃) and B: ACN (30% to 60% in 7 min)). The collected fraction was lyophilized to give 3-amino-N-[(3R)-7-(4-cyano-1,1-dioxo-1λ6-thian-4-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide as a light yellow solid. MS (ESI, m/z): 497 $[M+H]^+$. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.33 (d, J = 8.4 Hz, 1H), 7.63 (d, J = 7.6 Hz, 1H), 7.32 (d, J = 8.4 Hz, 2H), 7.24-7.22 (m, 3H), 7.10-7.08 (m, 1H), 6.99 (d, J = 2.0 Hz, 1H), 4.37-4.32 (m, 1H), 4.26-4.22 (m, 1H), 3.93-3.88 (m, 1H), 3.42-3.27 (m, 4H), 3.01-2.99 (m, 2H), 2.59-2.50 (m, 7H).

Example 35. 3-Amino-6-methyl-N-[(3R)-5,6,8-trifluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]thieno[2,3-b]pyridine-2-carboxamide



Step 1. Benzyl 4-(2,3,5,6-tetrafluoro-4-formylphenyl)piperazine-1-carboxylate

[00533] A solution of 2,3,4,5,6-pentafluorobenzaldehyde (10.0 g, 51.0 mmol), benzyl piperazine-1-carboxylate (11.2 g, 50.8 mmol) and DIEA (15.9 mL, 95.9 mmol) in DMF (50 mL) was stirred for 12 h at 60 °C. The reaction was quenched by the addition of water (50 mL), and extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, washed with water (6 x 100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by silica gel chromatography (eluting with 1:5 ethyl acetate/pet. ether) to give benzyl 4-(2,3,5,6-tetrafluoro-4-formylphenyl)piperazine-1-carboxylate as a white solid. MS (ESI, m/z): 397 $[M+H]^+$.

Step 2. Benzyl 4-[2,3,5,6-tetrafluoro-4-(hydroxymethyl)phenyl]piperazine-1-carboxylate

[00534] To a solution of benzyl 4-(2,3,5,6-tetrafluoro-4-formylphenyl)piperazine-1-carboxylate (4.00 g, 10.1 mmol) in methanol (40 mL) was added NaBH₄ (950 mg, 25.1 mmol) at < 10 °C. The resulting solution was stirred for 2h at 25 °C. The solvent was removed under vacuum. The residue was diluted with water (100 mL). The resulting mixture was extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give benzyl 4-[2,3,5,6-tetrafluoro-4-(hydroxymethyl)phenyl]piperazine-1-carboxylate as a white solid. MS: (ESI, m/z): 399 $[M+H]^+$.

Step 3. Benzyl 4-[4-(bromomethyl)-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate

[00535] A solution of benzyl 4-[2,3,5,6-tetrafluoro-4-(hydroxymethyl)phenyl]piperazine-1-carboxylate (5.70 g, 14.3 mmol) and PBr₃ (3.85 g, 14.2 mmol) in DCM (40 mL) was stirred for 2 h at 25 °C. The reaction was then quenched by the addition of water (40 mL). The resulting mixture was extracted with DCM (3 x 30 mL). The organic layers were combined, washed with brine (30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give benzyl 4-[4-(bromomethyl)-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate as a white solid. MS (ESI, *m/z*): 461, 463 [M+H]⁺.

Step 4. Benzyl 4-(4-[[[(2R,5S)-3,6-dimethoxy-5-(propan-2-yl)-2,5-dihydropyrazin-2-yl]methyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate

[00536] To a solution of (2S)-3,6-dimethoxy-2-(propan-2-yl)-2,5-dihydropyrazine (1.84 g, 9.99 mmol) in THF (50 mL) was added n-BuLi (6.0 mL, 2.5M in n-hexane) at -78 °C. The resulting solution was stirred for 30 min at -78 °C. To this was added benzyl 4-[4-(bromomethyl)-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate (4.60 g, 9.97 mmol) at -78 °C. The resulting solution was stirred for 2h at -78 °C. The reaction was then quenched by the addition of NH₄Cl (sat., aq.) (40 mL). The resulting mixture was extracted with ethyl acetate (3 x 30 mL). The organic layers were combined, washed with brine (3 x 30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reversed phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (5% to 75% in 30 min) to give benzyl 4-(4-[[[(2R,5S)-3,6-dimethoxy-5-(propan-2-yl)-2,5-dihydropyrazin-2-yl]methyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate as yellow oil. MS (ESI, *m/z*): 565 [M+H]⁺.

Step 5. Benzyl 4-[4-[(2R)-2-amino-3-methoxy-3-oxopropyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate

[00537] Hydrochloric acid (100 mL, 0.3N) was added into a stirring solution of benzyl 4-(4-[[[(2R,5S)-3,6-dimethoxy-5-(propan-2-yl)-2,5-dihydropyrazin-2-yl]methyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate (5.86 g, 10.3 mmol) in ACN (100 mL) at <10 °C. The resulting solution was stirred for 1h at 25 °C. The solvent was removed under vacuum. The pH value of the residue was adjusted to 9-10 with NaHCO₃ (sat. aq.). The resulting mixture was extracted with ethyl acetate (3 x 30 mL). The organic layers were combined, washed with brine (50 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give

benzyl 4-[4-[(2R)-2-amino-3-methoxy-3-oxopropyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate as a yellow solid. MS (ESI, m/z): 470 $[M+H]^+$.

Step 6. Benzyl 4-[4-[(2R)-2-amino-3-hydroxypropyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate

[00538] Into a solution of benzyl 4-[4-[(2R)-2-amino-3-methoxy-3-oxopropyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate (3.83 g, 8.16 mmol) in methanol (10 mL) and THF (40 mL) was added NaBH_4 (900 mg, 23.7 mmol) at $<10^\circ\text{C}$. The resulting solution was stirred for 3h at 25°C . The solvent was concentrated under vacuum. The residue was diluted with water (50 mL). The resulting mixture was extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, washed with brine (3 x 30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reversed phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (5% to 80% in 30 min)) to afford benzyl 4-[4-[(2R)-2-amino-3-hydroxypropyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate as a yellow oil. MS (ESI, m/z): 442 $[M+H]^+$.

Step 7. Benzyl 4-[(3R)-3-amino-5,6,8-trifluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate

[00539] Sodium hydride (34.0 mg, 1.42 mmol, 60%) was added into a stirring solution of benzyl 4-[4-[(2R)-2-amino-3-hydroxypropyl]-2,3,5,6-tetrafluorophenyl]piperazine-1-carboxylate (400 mg, 0.910 mmol) in DMSO (10 mL). The resulting solution was stirred for 30 min at 25°C and then 2h at 70°C . After cooling to 25°C , the reaction was quenched by the addition of water (40 mL). The resulting mixture was extracted with ethyl acetate (3 x 20 mL). The organic layers were combined, washed with brine (20 mL), dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified by reversed phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (20% to 70% in 30 min)) to give benzyl 4-[(3R)-3-amino-5,6,8-trifluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate as a yellow oil. MS (ESI, m/z): 422 $[M+H]^+$.

Step 8. Benzyl 4-[(3R)-3-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,8-trifluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate

[00540] To a solution of benzyl 4-[(3R)-3-amino-5,6,8-trifluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate (70.0 mg, 0.170 mmol) and 3-amino-6-methylthieno[2,3-b]pyridine-2-carboxylic acid (41.5 mg, 0.200 mmol) in DMA (2 mL) was added Et₃N (0.068 mL, 0.49 mmol) and HBTU (75.0 mg, 0.200 mmol). The resulting solution was stirred for 1 h at 25 °C. The mixture was purified by reversed phase chromatography (Column: C18 silica gel; Mobile phase, A: water (containing 0.05% TFA) and B: ACN (5% to 80% in 30 min)) to afford benzyl 4-[(3R)-3-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,8-trifluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate as a yellow oil. MS (ESI, *m/z*): 612 [M+H]⁺.

Step 9. 3-Amino-6-methyl-N-[(3R)-5,6,8-trifluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]thieno[2,3-b]pyridine-2-carboxamide

[00541] To a solution of benzyl 4-[(3R)-3-[3-amino-6-methylthieno[2,3-b]pyridine-2-amido]-5,6,8-trifluoro-3,4-dihydro-2H-1-benzopyran-7-yl]piperazine-1-carboxylate (50.0 mg, 0.080 mmol) in CH₃COOH (1.5 mL) was added hydrobromic acid (1.5 mL, 48%). The resulting solution was stirred for 1 h at 50 °C. The resulting mixture was concentrated under vacuum. The residue was purified by prep-HPLC (Column: XBridge Shield RP18 OBD Column; Mobile phase, A: water (containing 10 mmol/L NH₄HCO₃) and B: ACN (25% to 65% B in 8 min)). The collected fraction was lyophilized to give 3-amino-6-methyl-N-[(3R)-5,6,8-trifluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]thieno[2,3-b]pyridine-2-carboxamide as a white solid. MS (ESI, *m/z*): 478 [M+H]⁺. ¹H NMR (DMSO-*d*₆, 400 MHz) δ(ppm): 8.33 (d, *J* = 8.4 Hz, 1H), 7.70 (d, *J* = 6.8 Hz, 1H), 7.31 (d, *J* = 8.4 Hz, 1H), 7.25 (br s, 2H), 4.36-4.30 (m, 1H), 4.26-4.23 (m, 1H), 3.99-3.95 (m, 1H), 3.04-2.94 (m, 5H), 2.90-2.84 (m, 1H), 2.79-2.77 (m, 4H), 2.58 (s, 3H).

Example A-1(a): Biochemical Assay: Ubiquitin-Rhodamine 110 Assay for USP28 Activity.

[00542] The assay was performed in a final volume of 9 μL in assay buffer containing 20 mM Tris-HCl (pH 8.0, (1M Tris-HCl, pH 8.0 solution; Corning 46-031-CM)), 3 mM BME (2-Mercaptoethanol; Sigma 63689-25ML-F), 0.03% BGG (0.22 μM filtered, Sigma, G7516-25G), and 0.01% Triton X-100 (Sigma, T9284-10L). Nanoliter quantities of 10-point, 3-fold serial dilution in DMSO was pre-dispensed into 1536 assay plates (Corning, #3724BC) for a final test

concentration of 25 μ M to 1.3 nM, top to lowest dose, respectively. Enzyme USP28, construct His-tagged USP28-FL-mammalian, (protein expression and purification procedure described below). Concentration and incubation times were optimized for the maximal signal-to-background while maintaining initial velocity conditions at a fixed substrate concentration. The final concentration of the enzyme in the assay was 75 pM. Final substrate (Ub-Rh110; Ubiquitin-Rhodamine 110, UbiQ-126) concentration was 25 nM with [Ub-Rh110] $\ll$ K_m. 3 μ L of 2x enzyme was added to assay plates (pre-stamped with compound) preincubated with USP25 for 30 minutes and then 3 μ L of 2 x Ub-Rh110 was added to assay plates. Plates were incubated for 45 minutes at room temperature before addition of 3 μ L of stop solution (final concentration of 10 mM citric acid (Sigma, 251275-500G)). Fluorescence was read on the Envision (Excitation at 485 nm and Emission at 535 nm; Perkin Elmer) or on the PheraSTAR (Excitation at 485 nm and Emission at 535 nm; BMG Labtech).

Procedure for the Protein expression and purification for construct His-tagged USP28-FL-mammalian

[00543] Expression of USP28 (1-1077)-TEV-6*His (pTT5 vector) was carried out in Expi293f cells (sequence derived from uniprot ID: Q96RU2-1). Cells were re-suspended in lysis buffer B (50mM Bicine, pH 8.0, 20mM NaCl, 5% glycerol, 0.1% CHAPS, 5mM β -ME, 1mM PMSF, 1 μ g/ml Leupeptin, 1 μ g/ml Pepstatin) and lysed by sonication. Insoluble material was removed by centrifugation and the supernatant was loaded onto a Ni-NTA column (GE Healthcare) equilibrated with Ni Buffer A (50mM Bicine, pH 8.0, 20mM NaCl, 5% glycerol, 0.1% CHAPS, 5mM β -ME.) and washed with Ni Buffer A + 20 mM imidazole until A₂₈₀ reached baseline. The protein was eluted with Ni Buffer B (50mM Bicine, pH 8.0, 20mM NaCl, 5% glycerol, 0.1% CHAPS, 5mM β -ME, 300mM imidazole.). The protein was further purified using a SuperdexTM 200 10/300 GL column (GE Healthcare) equilibrated with 50mM Bicine, pH 8.0, 20mM NaCl, 5% glycerol, 0.1% CHAPS, 5mM β -ME. The protein was concentrated to 2.5 mg ml⁻¹, flash-frozen in liquid N₂ and stored at -80 °C.

Example A-1(b): Biochemical Assay: Ubiquitin-Rhodamine 110 Assay for USP28 Activity.

[00544] Each assay was performed in a final volume of 20 μ L in assay buffer containing 20 mM Tris-HCl (pH 8.0, (1M Tris-HCl, pH 8.0 solution; Corning 46-031-CM)), 2 mM CaCl₂ (1M Calcium Chloride solution; Sigma #21114) 2 mM BME (2-Mercaptoethanol; Sigma 63689-25ML-

F), 0.01% Prionex (0.22 μ M filtered, Sigma #G-0411), and 0.01% Triton X-100. Stock compound solutions were stored at -20 °C as 10 mM in DMSO. Up to 1 month prior to the assay, 2 mM test compounds were pre-dispensed into assay plates (Black, low volume; Corning #3820) and frozen at -20 °C. Prestamped assay plates were allowed to come to room temperature on the day of the assay. For the screen, 100 nL of 2 mM was pre-dispensed for a final screening concentration of 10 μ M (DMSO_(fc) = 0.5%). Enzyme (USP28, construct USP28 (USP28-5(1-1077)-TEV-6*His; LifeSensors) concentration and incubation times were optimized for the maximal signal-to-background while maintaining initial velocity conditions at a fixed substrate concentration. The final concentration of the enzyme in the assay was 400 pM. Final substrate (Ub-Rh110; Ubiquitin-Rhodamine 110, R&D Systems #U-555) concentration was 25 nM with [Ub-Rh110] << K_m. 10 μ L of 2x enzyme was added to assay plates (pre-stamped with compound) either simultaneously with 2 x Ub-Rh110 or preincubated with USP28 40 minutes prior to the addition of 10 μ L of 2 x Ub-Rh110 to compound plates. Plates were incubated stacked for 90 minutes at room temperature before fluorescence was read on the Envision (Excitation at 485 nm and Emission at 535 nm; Perkin Elmer) or on the PheraSTAR (Excitation at 485 nm and Emission at 535 nm; BMG Labtech).

[00545] For follow-up studies, each assay was performed in a final volume of 15 μ L in assay buffer containing 20 mM Tris-HCl (pH 8.0, (1M Tris-HCl, pH 8.0 solution; Corning 46-031-CM)), 3mM BME (2-Mercaptoethanol; Sigma 63689-25ML-F), 0.03% BGG (0.22 μ M filtered, Sigma, G7516-25G), and 0.01% Triton X-100 (Sigma, T9284-10L). Nanoliter quantities of either an 8-point or 10-point, 3-fold serial dilution in DMSO was pre-dispensed into assay plates (Perkin Elmer, ProxiPlate-384 F Plus, #) for a final test concentration of either 25 μ M to 11nM or 25 μ M to 1.3 nM, respectively. Enzyme USP28, construct USP28 (USP28-5(1-1077)-TEV-6*His; LifeSensors) concentration and incubation times were optimized for the maximal signal-to-background while maintaining initial velocity conditions at a fixed substrate concentration. The final concentration of the enzyme in the assay was 75 pM. Final substrate (Ub-Rh110; Ubiquitin-Rhodamine 110, R&D Systems #U-555) concentration was 25nM with [Ub-Rh110] << K_m. 5 μ L of 2x enzyme was added to assay plates (pre-stamped with compound) preincubated with USP28 for 30 minutes and then 5 μ L of 2 x Ub-Rh110 was added to assay plates. Plates were incubated stacked for 20 minutes at room temperature before 5 μ L of stop solution was added (final concentration of 10mM citric acid (Sigma, 251275-500G)). Fluorescence was read on the Envision

(Excitation at 485 nm and Emission at 535 nm; Perkin Elmer) or on the PheraSTAR (Excitation at 485 nm and Emission at 535 nm; BMG Labtech).

Example A-2: Biochemical Assay: Ubiquitin-Rhodamine 110 Assay for USP25 Activity.

[00546] The assay was performed in a final volume of 9 μ L in assay buffer containing 20 mM Tris-HCl (pH 8.0, (1M Tris-HCl, pH 8.0 solution; Corning 46-031-CM)), 3 mM BME (2-Mercaptoethanol; Sigma 63689-25ML-F), 0.03% BGG (0.22 μ M filtered, Sigma, G7516-25G), and 0.01% Triton X-100 (Sigma, T9284-10L). Nanoliter quantities of 10-point, 3-fold serial dilution in DMSO was pre-dispensed into 1536 assay plates (Corning, #3724BC) for a final test concentration of 25 μ M to 1.3 nM, top to lowest dose, respectively. Enzyme USP25, construct USP25-His6, (Boston Biochem E-546). Concentration and incubation times were optimized for the maximal signal-to-background while maintaining initial velocity conditions at a fixed substrate concentration. The final concentration of the enzyme in the assay was 75 pM. Final substrate (Ub-Rh110; Ubiquitin-Rhodamine 110, R&D Systems #U-555) concentration was 25 nM with [Ub-Rh110] $\ll$ K_m. 3 μ L of 2x enzyme was added to assay plates (pre-stamped with compound) preincubated with USP25 for 30 minutes and then 3 μ L of 2 x Ub-Rh110 was added to assay plates. Plates were incubated for 45 minutes at room temperature before addition of 3 μ L of stop solution (final concentration of 10 mM citric acid (Sigma, 251275-500G)). Fluorescence was read on the Envision (Excitation at 485 nm and Emission at 535 nm; Perkin Elmer) or on the PheraSTAR (Excitation at 485 nm and Emission at 535 nm; BMG Labtech).

[00547] For assay formats Example A-1(a), A-1(b), and A-2, data were reported as percent inhibition compared with control wells based on the following equation: %inh = $1 - ((\text{FLU} - \text{Ave}_{\text{Low}}) / (\text{Ave}_{\text{High}} - \text{Ave}_{\text{Low}}))$ where FLU = measured Fluorescence, Ave_{Low} = average Fluorescence of no enzyme control (n=16), and Ave_{High} = average Fluorescence of DMSO control (n=16). IC₅₀ values were determined by curve fitting of the standard 4 parameter logistic fitting algorithm included in the Activity Base software package: IDBS XE Designer Model205. Data is fitted using the Levenburg Marquardt algorithm.

[00548] The activity of compounds in the biochemical IC₅₀ assays (IC₅₀ ranges) according to the present disclosure are reported in Table 24 below according to the following:

“+”: > 2 μ M; “++”: 0.2-2 μ M; “+++”: 0.05-0.2 μ M; “++++”: 0.001-0.05 μ M.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 326

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 326

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :


NOTE POUR LE TOME / VOLUME NOTE:

AMENDED CLAIMS

received by the International Bureau on 19 December 2018 (19.12.2018)

WHAT IS CLAIMED IS:

1. A compound of Formula (IIaa):



or a pharmaceutically acceptable salt thereof, wherein

R_1 is chosen from 8-9-membered heteroaryls substituted with one or more substituent chosen from R_5 and R_6 ;

R_2 is chosen from N-linked 6-12 membered heterocyclyls or C-linked 6-12 membered heterocyclyls optionally substituted with one or more R_5 ;

R_3 is independently chosen from H, (C₁-C₆) alkyl, halogen, and -CN, wherein the (C₁-C₆) alkyl groups are optionally substituted with R_7 ;


each R_5 is independently chosen from -OH, -NH₂, -NHC(O)CH₃, -C(O)NHCH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, -NHC(O)CH₃, -C(O)NHCH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, -NH₂, and -OH;

each R_6 is independently chosen from -NH(C₁-C₆)alkyl-aryls, -NH(C₁-C₆)alkyl-heteroaryls, -NH(C₁-C₆)alkyl-cyclyl, and -NH(C₁-C₆)alkyl-heterocyclyl groups, wherein each of the R_6 groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups;

each R_7 is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and -OH; and

n is 0, 1, 2, or 3.

2. A compound of Formula (IIb'):



or a pharmaceutically acceptable salt thereof, wherein

R' is chosen from H and CH₃;

R₁ is chosen from 8-9 membered heteroaryls substituted with one or more substituent chosen from R₅ and R₆;

R₂ is chosen from N-linked 6-12 membered heterocyclyls or C-linked 6-12 membered heterocyclyls optionally substituted with one or more R₅; and

R₃ is independently chosen from H, (C₁-C₆) alkyl, halogen, and -CN, wherein the (C₁-C₆) alkyl groups are optionally substituted with R₇;

each R₅ is independently chosen from -OH, -NH₂, amido-(C₁-C₆) alkyl, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, amido-(C₁-C₆) alkyl, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, -NH₂, and -OH, and wherein any R₅ group containing hydrogen can have one or more hydrogen replaced with deuterium;


each R₆ is independently chosen from -amino alkyl-aryls, -amino alkyl-heteroaryls, -amino alkyl-cyclyl, and -amino alkyl-heterocyclyl groups, wherein each of the R₆ groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups, and further wherein any R₆ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₇ is independently chosen from –OH, –NH₂, (C₁–C₆) alkyl, (C₁–C₆) alkoxy, (C₁–C₆) haloalkyl, (C₁–C₆) haloalkoxy, halogen, cycloalkyl, –C(O)–cycloalkyl, heterocycloalkyl, and –C(O)–heterocycloalkyl groups, wherein each of –NH₂, (C₁–C₆) alkyl, (C₁–C₆) alkoxy, (C₁–C₆) haloalkyl, (C₁–C₆) haloalkoxy, halogen, cycloalkyl, –C(O)–cycloalkyl, heterocycloalkyl, and –C(O)–heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁–C₆) alkyl, (C₁–C₆) alkoxy, and –OH;


and

n is 0, 1, 2, or 3.


3. The compound of any one of claims 1-2, wherein R₁, optionally substituted with R₅ and/or R₆, is chosen from Table A.
4. The compound of any one of claims 1-3, wherein R₂, optionally substituted with R₅, is chosen from Table B.
5. The compound of any one of claims 1-3, wherein R₁, optionally substituted with R₅ and/or R₆, is chosen from




R₂, optionally substituted with R₅, is chosen from




6. The compound of any one of claims 1-2, wherein R₁, optionally substituted with R₅ and/or R₆, is chosen from



R₂, optionally substituted with R₅, is chosen from



7. A compound of Formula (VII):



or a pharmaceutically acceptable thereof, wherein

Y is chosen from C(R₃) and N;

R' is chosen from H, deuterium, and CH₃;

R₁ is chosen from 6-11 membered heteroaryls optionally substituted with one or more substituent chosen from R₅ and/or R₆;

R₂ is chosen from N-linked 4-12 membered heterocyclyls and C-linked 4-12 membered heterocyclyls, wherein the heterocyclyls are optionally substituted with one or more R₅, and further wherein any R₂ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₃ is independently chosen from H, deuterium, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, -OH, -CN, wherein each of (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups are optionally substituted with one or more R₇;

R₄ is chosen from H, deuterium, (C₁-C₆) alkyl, halogen, -OH, -CN, and further wherein any R₄ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₅ (if present) is independently chosen from -OH, -NH₂, NHC(O)CH₃, -C(O)NHCH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, -NHC(O)CH₃, -C(O)NHCH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and -OH, and wherein any R₅ group containing hydrogen can have one or more hydrogen replaced with deuterium;


each R₆ (if present) is chosen from -NH(C₁-C₆)alkyl-aryls, -NH(C₁-C₆)alkyl-heteroaryls, -NH(C₁-C₆)alkyl-heterocyclyl groups, and -NH(C₁-C₆)alkyl-heterocyclyl groups, wherein each of the R₆ groups are optionally substituted with one or more substituent chosen from -OH, -NH, halogens, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and (C₁-C₆) haloalkyl groups, and further wherein any R₆ group containing hydrogen can have one or more hydrogen replaced with deuterium;

each R₇ is independently chosen from -OH, -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, -C(O)-cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkyl, (C₁-C₆) alkoxy, and -OH;

and

n is 0, 1, 2, or 3.

8. The compound of claim 7, of Formula (IIIaa):



or a pharmaceutically acceptable thereof, wherein


R₁ is chosen from 8-11 membered heteroaryls optionally substituted with one or more R₅;

R₂ is chosen from N-linked 4-12 membered heterocyclyls and C-linked 4-12 membered heterocyclyls, optionally substituted with one or more R₅;

each R₅ (if present) is independently chosen from –OH, –NH₂, NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of –NH₂, -NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and –OH; and

n is 0, 1, 2, or 3.

9. The compound of claim 7, of Formula (VIIaa):



or a pharmaceutically acceptable thereof, wherein


R₁ is chosen from 8-9 membered heteroaryls optionally substituted with one or more R₅;

R₂ is chosen from N-linked 4-12 membered heterocyclyls optionally substituted with one or more R₅;

each R₅ (if present) is independently chosen from –OH, –NH₂, NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of –NH₂, -NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and –OH; and

n is 0, 1, 2, or 3.

10. The compound of any one of claims 8 or 9, wherein R₁ is chosen from



wherein B is chosen from a bond or C;

Z is chosen from N, S, C(Rii);


Rii is chosen from H, CH₃ and R₅;

R₂ is chosen from N-linked 5-8 membered heterocyclyls substituted with one to three R₅;


each R₅ (if present) is independently chosen from -OH, -NH₂, NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, halogen, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl groups, wherein each of -NH₂, -NHC(O)CH₃, (C₁-C₆) alkyl, (C₁-C₆) alkoxy, (C₁-C₆) haloalkyl, (C₁-C₆) haloalkoxy, cycloalkyl, heterocycloalkyl, and -C(O)-heterocycloalkyl are optionally substituted with one or more substituent independently chosen from (C₁-C₆) alkoxy, -NH₂, and -OH; and


n is 0, 1, 2, or 3.

11. The compound of any one of claims 7-10, wherein R₁, optionally substituted with R₅, is chosen from



and R₂, optionally substituted with R₅, is chosen from





12. The compound of any one of claims 1-11, wherein the compound is
- a USP28 Inhibitor compound having an IC_{50} of 0.001-2 micromolar in the Ubiquitin-Rhodamine 110 Assay for USP28 as described in Example A-1(a) herein; and/or
 - a USP28 Inhibitor compound having an IC_{50} of 0.001-2 micromolar in the Ubiquitin-Rhodamine 110 Assay for USP28 as described in Example A-1(b) herein; and/or
 - a USP25 Inhibitor compound having an IC_{50} of 0.001-2 micromolar in the Ubiquitin-Rhodamine 110 Assay for USP25 as described in Example A-2 herein.
13. The compound of any one of claims 1, 2, or 7, chosen from the compounds of any one of Tables 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 or a pharmaceutically acceptable salt thereof.
14. The compound of claim 13, chosen from the following compounds:
- 10-1:** (R)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 10-5:** (S)-3-amino-N-(5-fluoro-7-(piperazin-1-yl)chroman-3-yl)-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 10-15:** 3-amino-N-[(3R)-5-fluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-N,6-dimethylthieno[2,3-b]pyridine-2-carboxamide
- 10-16:** 3-amino-N-[(3S)-5-fluoro-7-(piperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-N,6-dimethylthieno[2,3-b]pyridine-2-carboxamide
- 11-1:** N-((2S)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide

- 11-2:** N-((2R)-6-(3,8-diazabicyclo[3.2.1]octan-3-yl)-1,2,3,4-tetrahydronaphthalen-2-yl)-3-amino-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 23-1:** 7-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide
- 23-2:** 7-amino-N-[(6R)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide
- 23-3:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide
- 23-4:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 23-5:** 3-amino-N-[(6R)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 23-6:** 7-amino-3-methyl-N-[(6S)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyrazine-6-carboxamide
- 23-7:** 3-amino-6-methyl-N-[(6S)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide
- 23-8:** 3-amino-6-methyl-N-[(6R)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide
- 23-9:** 6-amino-2-methyl-N-[(6S)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-d][1,3]thiazole-5-carboxamide
- 23-12:** 3-amino-N-[(6S)-2-{3,8-diazabicyclo[3.2.1]octan-3-yl}-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 23-13:** 7-amino-N-[(6S)-2-{3,8-diazabicyclo[3.2.1]octan-3-yl}-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide
- 23-14:** 3-amino-4,6-dimethyl-N-[(6S)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide
- 23-17:** 3-amino-N-[(6S)-4-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 23-18:** 3-amino-N-[(6S)-2-{3,8-diazabicyclo[3.2.1]octan-3-yl}-4-fluoro-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

- 24-1:** 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 24-2:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 25:** 3-amino-N-[(6S)-2-[(3S,4R)-3-amino-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 26-1:** 3-amino-N-[(6S)-2-[(3S,4R)-3-(methoxymethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 26-2:** 3-amino-N-[(6S)-2-[(3R,4S)-3-(methoxymethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 27-1:** 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 27-2:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 28-1:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 28-2:** 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(2-methoxyethoxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 29-1:** 3-amino-N-[(6S)-2-[(9S)-9-amino-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 29-2:** 3-amino-N-[(6S)-2-[(9R)-9-amino-1,4-dioxo-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 30-1:** 3-amino-N-[(6S)-3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 30-2:** 3-amino-N-[(6R)-3-fluoro-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 31-3:** 3-amino-6-methyl-N-[(6'S)-2'-(piperazin-1-yl)-6',7'-dihydro-5'H-spiro[cyclopropane-1,8'-quinoline]-6'-yl]thieno[2,3-b]pyridine-2-carboxamide
- 31-4:** 3-amino-6-methyl-N-[(6'R)-2'-(piperazin-1-yl)-6',7'-dihydro-5'H-spiro[cyclopropane-1,8'-quinoline]-6'-yl]thieno[2,3-b]pyridine-2-carboxamide

- 142:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-ethoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 203:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-5-fluoro-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 204:** 3-amino-5-fluoro-6-methyl-N-[(6S)-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide
- 205:** 3-amino-6-methyl-N-[(6S,8S)-8-methyl-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide
- 206:** 3-amino-6-methyl-N-[(6R,8S)-8-methyl-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide
- 207:** 3-amino-6-methyl-N-[(6S,8R)-8-methyl-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide
- 208:** 3-amino-6-methyl-N-[(6R,8R)-8-methyl-2-(piperazin-1-yl)-5,6,7,8-tetrahydroquinolin-6-yl]thieno[2,3-b]pyridine-2-carboxamide
- 210:** 3-amino-N-[(6S)-2-[(3S,4S)-4-amino-3-methoxy-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 211:** 3-amino-N-[(6S)-2-[(3R,4R)-4-amino-3-methoxy-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 239:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-methoxy-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 240:** 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-methoxy-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 254:** 3-amino-N-[(6S)-2-[(3S,4R)-3-amino-4-ethoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 255:** 3-amino-N-[(6S)-2-[(3R,4S)-3-amino-4-ethoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 269:** 3-amino-N-[(6S)-2-[(3S,4R)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 270:** 3-amino-N-[(6S)-2-[(3R,4S)-3-amino-4-methoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

- 279:** 7-amino-N-[(6S)-2-[(3S,4S)-4-amino-3-methoxy-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide
- 280:** 7-amino-N-[(6S)-2-[(3R,4R)-4-amino-3-methoxy-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide
- 285:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-cyclobutoxypyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 296:** 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide
- 297:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide
- 298:** 7-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide
- 299:** 7-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-3-methylthieno[2,3-b]pyrazine-6-carboxamide
- 300:** 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-3-fluoro-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 301:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-3-fluoro-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 302:** 3-amino-N-[(6R)-2-[(3R,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-3-fluoro-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 303:** 3-amino-N-[(6R)-2-[(3S,4S)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-3-fluoro-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 311:** 3-amino-N-[(6S)-2-[(3S,4S)-3-methoxy-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 312:** 3-amino-N-[(6S)-2-[(3S,4S)-3-methoxy-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide
- 320:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 321:** 3-amino-N-[(6S)-2-[(3S,4R)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

- 322:** 3-amino-N-[(6S)-2-[(3R,4S)-3-amino-4-(propan-2-yloxy)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 327:** 3-amino-N-[(6S)-2-[(3R,4S)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 328:** 3-amino-N-[(6S)-2-[(3S,4R)-3-amino-4-(methoxymethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 331:** 3-amino-N-[(6S)-2-[(3R,4S)-3-hydroxy-3-methyl-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 332:** 3-amino-N-[(6S)-2-[(3S,4R)-3-hydroxy-3-methyl-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 341:** 3-amino-N-[(6S)-2-[(3S,4R)-4-amino-3-(methoxymethyl)-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 342:** 3-amino-N-[(6S)-2-[(3R,4R)-4-amino-3-(methoxymethyl)-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 343:** 3-amino-N-[(6S)-2-[(3S,4S)-4-amino-3-(methoxymethyl)-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 344:** 3-amino-N-[(6S)-2-[(3R,4S)-4-amino-3-(methoxymethyl)-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 345:** 3-amino-N-[(6S)-2-[(3R,4R)-3-amino-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 346:** 3-amino-N-[(6S)-2-[(3S,4S)-3-amino-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 347:** 3-amino-N-[(6S)-2-[(3R,4S)-3-amino-4-(difluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 354:** 3-amino-N-[(6S)-2-{3,8-diazabicyclo[3.2.1]octan-3-yl}-5,6,7,8-tetrahydroquinolin-6-yl]-4,6-dimethylthieno[2,3-b]pyridine-2-carboxamide
- 363:** 3-amino-N-[(6S)-2-[(3R,4S)-3-amino-4-(trifluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 364:** 3-amino-N-[(6S)-2-[(3S,4R)-3-amino-4-(trifluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide

- 376:** 3-amino-N-[(6S)-2-[(3S,4R)-3-(difluoromethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 377:** 3-amino-N-[(6S)-2-[(3R,4S)-3-(difluoromethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 384:** 3-amino-N-[(6S)-2-[(3R,4S)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 385:** 3-amino-N-[(6S)-2-[(3S,4R)-3-amino-4-(fluoromethyl)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 427:** 3-amino-N-[(6S)-2-[(3R,4S)-4-amino-3-methoxy-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 428:** 3-amino-N-[(6S)-2-[(3S,4R)-4-amino-3-methoxy-3-methylpyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 430:** 3-amino-N-[(6S)-2-[(5S,9R)-9-amino-2-oxa-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 431:** 3-amino-N-[(6S)-2-[(5R,9S)-9-amino-2-oxa-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 437:** 3-amino-N-[(6S)-2-[(5R,9R)-9-amino-2-oxa-7-azaspiro[4.4]nonan-7-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 441:** 3-amino-N-[(6S)-2-[(3R,4R)-3-(fluoromethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide
- 442:** 3-amino-N-[(6S)-2-[(3S,4S)-3-(fluoromethyl)-4-(methylamino)pyrrolidin-1-yl]-5,6,7,8-tetrahydroquinolin-6-yl]-6-methylthieno[2,3-b]pyridine-2-carboxamide.