



Office de la Propriété

Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2212043 C 2005/08/30

(11)(21) 2 212 043

(12) BREVET CANADIEN  
CANADIAN PATENT

(13) C

(86) Date de dépôt PCT/PCT Filing Date: 1996/02/02  
(87) Date publication PCT/PCT Publication Date: 1996/08/08  
(45) Date de délivrance/Issue Date: 2005/08/30  
(85) Entrée phase nationale/National Entry: 1997/07/31  
(86) N° demande PCT/PCT Application No.: JP 1996/000218  
(87) N° publication PCT/PCT Publication No.: 1996/024044  
(30) Priorité/Priority: 1995/02/02 (51668/1995) JP

(51) Cl.Int.<sup>6</sup>/Int.Cl.<sup>6</sup> G01N 21/76, C12Q 1/68, G01N 33/58,  
G01N 33/52  
(72) Inventeurs/Inventors:  
TSUJI, YASUHIRO, JP;  
KAMISANGO, KEIICHI, JP;  
OTSUKA, MITSUO, JP  
(73) Propriétaire/Owner:  
CHUGAI SEIYAKU KABUSHIKI KAISHA, JP  
(74) Agent: OGILVY RENAULT LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : PROCEDE POUR DOSAGE D'UN ANALYTE EN AJUSTANT LA CHIMIOLUMINESCENCE

(54) Title: METHOD FOR ASSAY OF ANALYTE BY ADJUSTMENT OF CHEMILUMINESCENCE

(57) Abrégé/Abstract:

A method for assay of an analyte by use of a material labeled with a chemiluminescent substance, which comprises adding a quencher and/or decreasing the specific activity of a chemiluminescent substance labeled probe, thereby decreasing the quantity of chemiluminescence.

**ABSTRACT**

A method for assay of an analyte by use of a material labeled with a chemiluminescent substance, which comprises adding a quencher and/or decreasing the specific activity of a chemiluminescent substance labeled probe, thereby decreasing the quantity of chemiluminescence.

METHOD OF ASSAYING SPECIMEN SUBSTANCE BY CONTROLLING DOSE  
OF CHEMILUMINESCENCE

## TECHNICAL FIELD

5 This invention relates to a method for assay of an analyte by use of a material labeled with a chemiluminescent substance. More specifically, the invention relates to a method for assay in a broader range by decreasing the quantity of chemiluminescence.

## 10 BACKGROUND ART

Among methods for assay of an analyte in a test solution is one using a probe labeled with a chemiluminescent substance. This method is widely used as being capable of determining the amount of the analyte 15 highly sensitively by measuring the quantity of chemiluminescence. This method is useful for a suitable amount of the analyte. In the presence of a large amount of the analyte (e.g. when many copies are produced by gene amplification such as polymerase chain reaction), however, 20 the quantity of chemiluminescence by the method exceeds the determination limit of a measuring device, making accurate assay impossible. Such samples with results in excess of the upper assay limit have been determined again after dilution of the test solution. This procedure is 25 very laborious. Samples amplified by gene amplification, in particular, can cause contamination of the amplified product as a result of dilution. Utmost care has been taken to avoid the contamination, further increasing

labor. To broaden the range of assay without diluting the sample, there has been no choice but to wait for an improvement in the measuring device.

We, the inventors, have found that the foregoing 5 problems with the determination of a sample beyond the assay limits can be solved by decreasing the quantity of chemiluminescence, without requiring a laborious operation such as the dilution of the sample, or an improvement in the measuring device. This finding has led us to 10 accomplish this invention.

#### DISCLOSURE OF THE INVENTION

The present invention is a method for assay of an analyte by use of a material labeled with a chemiluminescent substance, which comprises decreasing the 15 quantity of chemiluminescence. The invention further provides a method for assay which comprises adding a quencher and/or decreasing the specific activity of a chemiluminescent substance labeled probe.

The assay of the analyte by use of a material 20 labeled with a chemiluminescent substance, indicated above, refers, for example, to reacting a sample containing an analyte with a material labeled with a chemiluminescent substance, and measuring the quantity of chemiluminescence of the conjugate to detect or determine 25 the analyte.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the effect of Phenol Red addition on the quantitative amplification and detection of Hepatitis B virus (HBV) template in the serum; and

Fig. 2 shows the effect of unlabeled probe addition 5 on the quantitative amplification and detection of HBV template in the serum.

#### BEST MODE FOR CARRYING OUT THE INVENTION

(1) Decreasing the quantity of chemiluminescence by addition of a quencher

10 The quencher used in the present invention may be any substance which can quench chemiluminescence. For example, it includes color matters and india ink (drops of india ink, supernatant of india ink). Examples of the color matters are Crystal Violet, Bromophenol Blue, 15 carminic acid, Chlorophenol Red, hematoxylin, Bromophenol Purple, Bromophenol Red, rosolic acid, Phenol Red, Cresol Red, and Methacresol Red.

When the color matter is used as the quencher, the concentration of the color matter at measurement of 20 chemiluminescence may be in the range of 0.01 to 10%, preferably 0.01 to 1%, although it differs depending on the color matter used. When india ink is used as the quencher, on the other hand, its amount at measurement of chemiluminescence may be in the range of 0.01 to 50%, 25 preferably 1 to 20%, based on the amount of the test solution. The quencher may be added at any time before measurement of chemiluminescence. For example, it may be

added either before or after the reaction between the analyte and the labeled probe.

(2) Decreasing the quantity of chemiluminescence by decreasing the specific activity of a 5 chemiluminescent substance labeled probe

To decrease the specific activity of a labeled probe in the present invention, an unlabeled probe is added to the labeled probe. The unlabeled probe may be added in an amount in the range of 0.1 to  $10^5$ , preferably 10 10 to  $10^3$ , with respect to 1 of the labeled probe.

(3) Combining the method (1) and the method (2)

In the present invention, the addition of the quencher and the reduction of the specific activity of the chemiluminescent substance labeled probe may be combined. 15 The conditions for use of both methods in this combination follow the above-described ranges.

The use of the assay method according to the present invention enables the quantity of chemiluminescence to be measured accurately, even when the 20 analyte in the sample solution is present in so large an amount as to exceed the assay limit. For instance, the analyte can be easily detected or determined quantitatively from a product produced by gene amplification of genetic information (DNA or RNA) on a 25 microorganism or cell. This action can be confirmed by Examples 1 to 4 to be offered later on.

In the method involving the addition of the quencher, not only positive signals, but also background

(noise) levels are decreased. This decrease in the quantity of chemiluminescence permits the quantitative measurement of strongly positive signals, while the decrease in the background (noise) levels makes

5 discrimination of weakly positive signals possible. In short, the addition of the quencher enables the assay of a highly positive sample without affecting the determination of a weakly positive sample. This action can be confirmed by Examples 1 to 3 to be offered later on.

10 By using the present invention to decrease the quantity of chemiluminescence, it becomes possible to detect or quantitatively determine a highly positive sample or a sample beyond the assay limits. The method of adding the quencher, in particular, reduces the background

15 (noise) level as well. Thus, a highly positive sample can be measured without influence on the assay of a weakly positive sample. Hence, the method of the present invention proves to be an excellent method which can broaden the range of assay without diluting the sample or

20 improving the measuring device.

#### EXAMPLES

The present invention will now be described in more detail with reference to Examples, whose descriptions do not limit the invention.

25 [Example 1]

##### Method

Five  $\mu$ l of human serum containing HBV-DNA sequence (Galibert, F., Mandart, E., Fitoussi, F., Tiollais, P.

and Charnay, P., Nature 281, 646-650 (1979)) (50 to 5,000 copies per amplification) was mixed with 20  $\mu$ l of an alkaline solution (pH 13), followed by heating for 5 minutes at 97°C. At room temperature, the mixture was 5 allowed to cool for 10 minutes, and then neutralized with a buffer. Two kinds of primers were added, and annealing was performed at room temperature. After DNA and RNA polymerases were added, gene amplification (using the method described in Published WO 91/01384 10 was carried out at 37°C. The amplification product and an acridinium ester labeled probe were hybridized at 60°C, whereafter the amplification product was detected by the HPA method (Arnold JR, L.J., Hammond, P.W., Wiese, W.A. and Nelson, 15 N.C., Clinical Chemistry 35, 1588-1594 (1988)). In detecting the amplification product by the measurement of chemiluminescence, the effect of addition of Phenol Red was investigated. Phenol Red was added in an amount of 0.05% to a test solution for chemiluminescence measurement 20 to measure the quantity of chemiluminescence. The results of measurement were compared with the results obtained from the testing solution containing no Phenol Red. The results are shown in Fig. 1.

#### Discussion

25 As shown in Fig. 1, the Phenol Red-free sample nearly reached the assay limit (saturation value) at about 500 GE (genome equivalents)/AMP, above which the quantity of chemiluminescence became no more linear. With the

Phenol Red-containing sample, the quantity of chemiluminescence remained linear even at 5,000 GE/AMP. This means that the addition of Phenol Red made the assay of 500 GE/AMP or more possible. Compared with the Phenol Red-free sample, the Phenol Red-containing sample markedly reduced the background (noise) level (DNA content = 0), thus permitting the assay of a weakly positive sample (about 50 GE/AMP). In other words, the addition of Phenol Red enabled the assay of a highly positive sample without 10 affecting the determination of a weakly positive sample.

[Example 2]

Method

In detecting the amplification product obtained by gene amplification as in Example 1, various amounts of 15 Phenol Red were added to study the effect of Phenol Red addition. Phenol Red was added in an amount of 0.025 to 0.2% to a test solution for chemiluminescence measurement to measure the quantity of chemiluminescence. The results of measurement were compared with the results obtained 20 using the testing solution containing no Phenol Red. The results are shown in Table 1.

Table 1 Effect of Phenol Red at determination of quantity of chemiluminescence

Phenol Red concentration (%)	Positive sample (50 GE/AMP)	Negative sample (0 GE/AMP)	Positive/negative ratio
0	237340	967	245
0.025	73120	239	306
0.05	41706	172	242
0.1	19332	114	170
0.2	7571	89	85

### Discussion

As shown in Table 1, the quantity of chemiluminescence from the positive sample and the quantity of chemiluminescence (background) from the negative sample decreased in a manner dependent on the amount of Phenol Red added. These findings demonstrate that the inventive method is available in a wide range of phenol Red concentrations.

### [Example 3]

#### Method

Tests were conducted in the same way as in Example 2, except that commercially available india ink was used instead of Phenol Red.

To a test solution for chemiluminescence measurement, 1.25 to 10% by volume of india ink was added, and the quantity of chemiluminescence was measured. The results are shown in Table 2 in comparison with the

results of measurement of the test solution free from india ink.

5 Table 2 Effect of india ink at determination of quantity of chemiluminescence

Amount of india ink added (% by volume)	Positive sample (50 GE/AMP)	Negative sample (0 GE/AMP)	Positive/negative ratio
0	291638	1689	173
1.25	30902	218	142
2.5	21292	119	179
5	6609	63	105
10	5172	43	120

#### Discussion

As shown in Table 2, the quantity of 10 chemiluminescence from the positive sample and the quantity of chemiluminescence (background) from the negative sample decreased in a manner dependent on the amount of india ink added. These findings demonstrate that the inventive method can be used in a wide range of 15 india ink concentrations.

[Example 4]

#### Method

Gene amplification was performed in the same manner as in Example 1. When the amplification product and an 20 acridinium ester labeled probe were hybridized at 60°C, an

unlabeled probe in various amounts was added. The amounts of the unlabeled probe were 10 to 1,000 relative to 1 of the labeled probe. After hybridization, the quantities of chemiluminescence in the samples were measured by the HPA 5 method. The results are shown in Fig. 2 in comparison with the results of measurement of the test solution free from the unlabeled probe.

#### Discussion

As shown in Fig. 2, the unlabeled probe-free sample 10 nearly reached the assay limit (saturation value) at about 500 to 5,000 GE/AMP, above which the quantity of chemiluminescence became no more linear. The unlabeled probe-containing samples, on the other hand, showed decreases in the quantity of chemiluminescence in a manner 15 dependent on the amount of the unlabeled probe added. When the amount of the unlabeled probe added relative to the labeled probe was 100(unlabeled probe):1(labeled probe) or more, assay was possible even at about 500,000 GE/AMP. Thus, the addition of the unlabeled probe 20 permitted assay of 500 to 500,000 GE/AMP or more.

**CLAIMS:**

1. A method for assay of an analyte which is an amplified gene product by use of a chemiluminescent substance labeled probe, which comprises decreasing a quantity of chemiluminescence by adding an unlabeled probe to the chemiluminescent substance labeled probe in order to decrease a specific activity of the chemiluminescent substance labeled probe.
2. A method for assay of an analyte which is an amplified gene product which comprises:
  - (1) providing a first probe that specifically binds to said analyte wherein said first probe is labeled with a chemiluminescent substance;
  - (2) adding an unlabeled, second probe that specifically binds to said analyte;
  - (3) contacting said first probe and said second probe with said analyte;
  - (4) quantitatively detecting the amount of chemiluminescence, wherein said second unlabeled probe decreases chemiluminescence; and
  - (5) correlating chemiluminescence to the presence of said analyte.
3. A method according to claims 1 or 2, wherein said unlabeled probe is added in amount of from 10 to  $10^3$  parts per one part of the labeled probe.
4. A method according to claim 2 which further comprises the

step of adding a quencher to said analyte before said quantitatively detecting step (4) to decrease said chemiluminescence and decrease background noise levels of said chemiluminescence.

5. A method according to claim 4 wherein the quencher is at least one member selected from the group consisting of color matter and india ink.

6. A method according to claim 2 which further comprises the step of adding a quencher to said analyte between the steps (1) and (2) or between the steps (2) and (3) to decrease said chemiluminescence and decrease background noise levels of said chemiluminescence.

7. A method according to claim 6 wherein the quencher is at least one member selected from the group consisting of color matter and india ink.

Fig. 1




Fig. 2

