



(86) Date de dépôt PCT/PCT Filing Date: 2009/05/28
(87) Date publication PCT/PCT Publication Date: 2009/12/10
(45) Date de délivrance/Issue Date: 2018/05/01
(85) Entrée phase nationale/National Entry: 2010/11/25
(86) N° demande PCT/PCT Application No.: US 2009/045519
(87) N° publication PCT/PCT Publication No.: 2009/148928
(30) Priorités/Priorities: 2008/05/29 (US61/057,183);
2009/04/17 (US61/170,561)

(51) Cl.Int./Int.Cl. *C07K 16/22* (2006.01),
A61K 39/395 (2006.01), *C07K 16/46* (2006.01),
C12N 15/00 (2006.01), *C12N 15/13* (2006.01),
C12P 21/08 (2006.01)

(72) Inventeurs/Inventors:
KIM, KYUNG JIN, US;
WANG, LIHONG, US;
PARK, HANGIL, US;
VASQUEZ, MAXIMILIANO, US

(73) Propriétaire/Owner:
GALAXY BIOTECH, LLC, US

(74) Agent: SMART & BIGGAR

(54) Titre : ANTICORPS MONOCLONAUX DIRIGES CONTRE UN FACTEUR DE CROISSANCE FIBROBLASTIQUE
BASIQUE

(54) Title: MONOCLONAL ANTIBODIES TO BASIC FIBROBLAST GROWTH FACTOR

(57) **Abrégé/Abstract:**

The present invention is directed toward a neutralizing monoclonal antibody to basic fibroblast growth factor, a pharmaceutical composition comprising same, and methods of treatment comprising administering such a pharmaceutical composition to a patient.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 December 2009 (10.12.2009)

(10) International Publication Number
WO 2009/148928 A1

(51) International Patent Classification:

C12P 21/08 (2006.01) *A61K 39/395* (2006.01)
C07K 16/00 (2006.01)

(21) International Application Number:

PCT/US2009/045519

(22) International Filing Date:

28 May 2009 (28.05.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/057,183 29 May 2008 (29.05.2008) US
61/170,561 17 April 2009 (17.04.2009) US

(71) Applicant (for all designated States except US):

GALAXY BIOTECH, LLC [US/US]; 22830 San Juan Road, Cupertino, CA 95014 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KIM, Kyung, Jin** [US/US]; 22830 San Juan Road, Cupertino, CA 95014 (US). **WANG, Lihong** [CN/US]; 275 Ventura Ave., #24, Palo Alto, CA 94306 (US). **PARK, Hangil** [KR/US]; 788 Harrison St., #520, San Francisco, CA 94107 (US). **VASQUEZ, Maximiliano** [US/US]; 3813 Louis Road, Palo Alto, CA 94303 (US).

(74) Agents: **LIEBESCHUETZ, Joe** et al.; Townsend And Townsend And Crew LLP, Two Embarcadero Center, 8th Floor, San Francisco, CA 94111-3834 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: MONOCLONAL ANTIBODIES TO BASIC FIBROBLAST GROWTH FACTOR

(57) Abstract: The present invention is directed toward a neutralizing monoclonal antibody to basic fibroblast growth factor, a pharmaceutical composition comprising same, and methods of treatment comprising administering such a pharmaceutical composition to a patient.



WO 2009/148928 A1

62451-1087

MONOCLONAL ANTIBODIES TO BASIC FIBROBLAST GROWTH FACTOR**CROSS-REFERENCES TO RELATED APPLICATIONS**

[0001] This applications claims the benefit under 35 U.S.C. § 119(e) of US Patent Application No. 61/057,183 filed May 29, 2008 and US Patent Application No. 61/170,561
5 filed April 17, 2009.

**REFERENCES TO A "SEQUENCE LISTING," A TABLE, OR A COMPUTER PROGRAM
LISTING SUBMITTED IN COMPUTER READABLE FORMAT**

[0002] This specification contains a Sequence Listing in electronic form.

10 **STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY
SPONSORED RESEARCH OR DEVELOPMENT**

[0003] The work described in this application was funded in part by Grant 5R44 CA101283-03 from the National Institutes of Health. The US Government has certain rights in this invention.

FIELD OF THE INVENTION

15 [0004] The present invention relates generally to the combination of monoclonal antibody (mAb) and recombinant DNA technologies for developing novel biologics, and more particularly, for example, to the production of monoclonal antibodies that bind to and neutralize basic Fibroblast Growth Factor.

BACKGROUND OF THE INVENTION

20 [0005] The prototypic Fibroblast Growth Factors (FGFs), acidic FGF (also called FGF1) and basic FGF (also called FGF2) were first isolated in the 1970s (FGF2 by Gospodarowicz et al., J. Biol. Chem. 250:2515, 1975). There are currently 22 known FGF family members, which can be grouped into 7 subfamilies based upon their similarity in activities and

sequences (Ornitz et al., *Genome Biol.* 2: 3005.1, 2001). However, the FGF family members bind to only four tyrosine kinase receptors (FGFR1-4) and their isoforms, which are expressed in a tissue-specific manner. The FGF1 subgroup consists of FGF1 and FGF2, which bind all four FGFRs, with FGF2 binding especially strongly to FGFR1c (Ornitz et al., *J. Biol. Chem.* 271:15292, 1996).

[0006] Human FGF2 is an 18 kDa non-glycosylated polypeptide consisting of 146 amino acids in the mature form derived from a 155 aa precursor (Ornitz et al., *Genome Biol.* 2:3005.1, 2001; Okada-Ban et al., *Int. J. Biochem. Cell. Biol.* 32:263, 2000). This 18 kDa form of FGF2 does not encode a signal sequence, but can be secreted by an unconventional energy-dependent pathway independent of the ER-Golgi complex (Mignatti et al., *J. Cell Physiol.* 151:81, 1992; Florkiewicz et al., *J. Cell Physiol.* 162:388, 1995). The single copy of the FGF2 gene also encodes four High Molecular Weight (HMW) forms of the protein, in addition to the 18 kDa form, by utilizing four alternate CUG initiation sites that provide N-terminal extensions of various sizes, resulting in proteins of 22 kDa (196 aa), 22.5 kDa (201 aa), 24 kDa (210 aa) and 34 kDa (288 aa) (Florkiewicz et al., *Proc. Natl. Acad. Sci. USA* 86:3978, 1989; Prats et al., *Proc. Natl. Acad. Sci. USA* 86:1836, 1989). The HMW forms are not secreted but are transported to the cell nucleus where they can regulate cell growth or behavior in an intracrine fashion (Delrieu, *FEBS Lett.* 468:6, 2000).

[0007] In addition to binding FGFR1-4 with high affinity, FGF2 binds to heparin sulfate proteoglycans (HSPG) with lower affinity. Although FGF2 is secreted as a monomer, cell surface HSPG dimerizes FGF2 in a non-covalent side-to-side configuration that is subsequently capable of dimerizing and activating FGF receptors (Mohammadi et al., *Cytokine Growth Factor Rev* 16:107, 2005). The binding of FGF and HSPG to the extracellular domain of FGFR induces receptor dimerization, activation and autophosphorylation.

[0008] The FGFs, and in particular FGF2, have a broad spectrum of activities on various cell types (Ornitz et al., *Genome Biol.* 2:3005.1, 2001). FGF2 stimulates proliferation of (i.e., is mitogenic for) certain cells including fibroblasts and endothelial cells and is a survival factor (anti-apoptotic) for certain cells such as neural cells (Okada-Ban, *op. cit.*). It also stimulates differentiation (morphogenesis) and migration (motility) of endothelial cells (Dow et al., *Urology* 55:800, 2000). FGF2 is involved in development, especially of the nervous system. Importantly, FGF2 is a powerful angiogenic factor (Presta et al., *Cytokine and Growth Factor Rev.* 16:159, 2005).

[0009] FGF2 and other FGFs are believed to play a role in cancer, both by stimulating angiogenesis and tumor cells directly (Presta et al., op cit.) During tumor progression, cancer cells may respond to the extracellular FGF2 secreted from the stromal cells (paracrine), and then the tumor cells themselves may secrete FGF2 and respond to it in an autocrine manner. FGF2 or its receptor FGFR1 has been shown to be expressed or overexpressed in most gliomas (Takahashi et al., Proc. Natl. Acad. Sci. USA 87:5710, 1990; Morrison et al. Cancer Res. 54:2794, 1994). FGF2 is involved in progression of prostate tumors (Dow et al., Urology 55:800, 2000) and is a key mediator of the proliferation of malignant melanomas (Wang et al., Nature Med. 3:887, 1997). Over-expression and/or involvement of FGF2 or FGFR1 in tumor progression has also been reported for salivary gland tumors (Myoken et al., J. Path. 178:429, 1996), esophageal cancer (Barclay et al., Clin. Cancer Res. 11:7683, 2005), and thyroid carcinomas (Boelaert et al., J. Clin. Endocrin. Metabol. 88:2341, 2003).

[0010] Polyclonal antibodies (antiserum) to FGF2 have been reported to inhibit tumor growth of a transplantable chondrosarcoma in mice (Baird et al., J. Cell Biochem. 30:79, 1986), neutralize various activities of FGF2 in vitro (Kurokawa et al., J. Biol. Chem. 264:7686, 1989), and to inhibit growth of U87 glioma cell intracranial xenografts when applied locally (Stan et al., J. Neurosurg. 82:1044, 1995). Among monoclonal antibodies, the DG2 mAb has been reported to neutralize activities of FGF2 in vitro and in vivo (Reilly et al., Biochem. Biophys. Res. Com. 164:736, 1989; WO 91/06668), modestly inhibit growth of rat C6 glioma xenografts in nude mice (Gross et al., J. Nat. Cancer Inst. 85:121, 1993), and modestly inhibit growth of rat chondrosarcomas when delivered intralesionally but not i.p. or i.v. (Coppola et al., Anticancer Res. 17:2033, 1997). Similarly, anti-FGF2 mAb DE6 has been reported to inhibit growth of glioma cells in vitro (Morrison et al., J. Neuroscience Res. 34:502, 1993). On the other hand, the anti-FGF2 mAbs bFM-1 and bFM-2 was reported to inhibit growth of endothelial cells in vitro but not to block tumor angiogenesis in vivo (Matsuzaki et al., Proc. Natl. Acad. Sci. USA 86:9911, 1989). The neutralizing anti-FGF2 mAb 1E6 has been reported to inhibit growth of RPMI4788 colon tumor xenografts (Aonuma et al., 19:4039, 1999). The anti-FGF2 mAb 254F1 has been reported to inhibit proliferation of human umbilical vein endothelial cells (HUVEC), while mAb FB-8 has been reported to inhibit proliferation of non-small cell lung cancer cell lines in vitro (Kuhn et al., Lung Cancer 44:167, 2004). The anti-FGF2 mAb 3H3 was reported to suppress growth of U87MG and T98G glioma and HeLa cell xenografts (Takahashi et al., FEBS Let. 288:65, 1991) and growth of the K1000 FGF2-transfected 3T3 cell line in mice (Hori et al., Cancer Res. 51:6180, 1991).

[0011] The GAL-F2 anti-FGF2 mAb described herein was discussed at the AACR 2009 Annual Meeting in Poster #1236.

BRIEF SUMMARY OF THE INVENTION

[0012] In one embodiment, the invention provides a neutralizing mAb to human basic
5 fibroblast growth factor (FGF2). The mAb inhibits at least one, and preferably several or all
biological activities of FGF2, including binding to one or more of the four FGF receptors
FGFR1-4; inducing proliferation of fibroblasts, endothelial cells, Mv 1 Lu mink lung
epithelial cells, and/or various human tumor cells; and inducing angiogenesis. The anti-FGF2
mAb can inhibit such an activity when used as a single agent. A preferred anti-FGF2 mAb
10 inhibits growth of a human tumor xenograft in a mouse. Preferably, the mAb of the invention
is genetically engineered, e.g., chimeric, humanized or human. Exemplary antibodies are
GAL-F2 and its chimeric and humanized forms, and mAbs which have the same epitope or
compete for binding with GAL-F2. Cell lines producing such antibodies are also provided.
In another embodiment, a pharmaceutical composition comprising a neutralizing anti-FGF2
15 antibody, e.g., chimeric or humanized GAL-F2, is provided. In a third embodiment, the
pharmaceutical composition is administered to a patient to treat cancer or other disease.

[0012A] The present invention as claimed relates to:

- a monoclonal antibody (mAb) that binds and neutralizes human basic
fibroblast growth factor (FGF2), and competes for binding to FGF2 with an antibody
20 produced by hybridoma PTA-8864 wherein the mAb is chimeric, humanized or human and
inhibits growth of a human tumor xenograft that expresses or responds to FGF2 in a mouse;
- an antibody which is a chimeric or humanized form of a monoclonal antibody
produced by the PTA-8864 hybridoma;
- a humanized antibody that binds human basic fibroblast growth factor (FGF2)
25 comprising a humanized light chain comprising three CDRs having the sequences
KASQSVSSDVG (SEQ ID NO:9), SGSNRYS (SEQ ID NO:10), and QQDYYSPTWT
(SEQ ID NO:11) and a humanized heavy chain comprising three CDRs having the sequences

62451-1087

NYVIN (SEQ ID NO:12), NDPYNDVSKYNEKFKG (SEQ ID NO:13) and
EGGGKYVYAMDS (SEQ ID NO:14);

- a monoclonal antibody that binds human basic fibroblast growth factor
(FGF2) comprising the three light chain CDRs having the sequences KASQSVSSDVG
5 (SEQ ID NO:9), SGSNRYS (SEQ ID NO:10), and QQDYYSPT (SEQ ID NO:11) and
three heavy chain CDRs having the sequences NYVIN (SEQ ID NO:12),
NDPYNDVSKYNEKFKG (SEQ ID NO:13) and EGGGKYVYAMDS (SEQ ID NO:14); and

- a monoclonal antibody produced by the hybridoma designated by ATCC
deposit number PTA-8864.

10 BRIEF DESCRIPTION OF THE DRAWINGS

[0013] Figure 1. Binding ELISA of GAL-F2 and control mouse mAb (mIgG) to
human FGF2 (hFGF2) and mouse FGF29 (mFGF2).

[0014] Figure 2. FGFR-Fc/FGF2-Flag Binding ELISA showing inhibition by
GAL-F2 but not control mouse mAb 5G8 for each of the four FGF receptors FGFR1-4.

15 **[0015]** Figure 3. Competitive binding assay of biotinylated GAL-F2 to human FGF2
with various anti-FGF2 mAbs.

[0016] Figure 4. Inhibition of FGF2-induced proliferation of BaF3 cells stably
transfected with FGFR1 or FGFR2, by GAL-F2 or control mAb 5G8. None means that no
FGF2 (or mAb) was applied to the cells.

20 **[0017]** Figure 5. Inhibition of FGF2-induced proliferation of Mv 1 Lu mink lung
epithelial cells by GAL-F2 or control mAb 5G8. None means that no FGF2 (or mAb) was
applied to the cells.

[0018] Figure 6. Photomicrographs of SMCC-7721 cells cloned in soft agar in the presence of control mouse mAb (mIgG), GAL-F2 or bFM-1 anti-FGF2 mAb.

[0019] Figure 7. Growth of RPMI 4788 human colon tumor xenografts in mice treated with GAL-F2 (100 μ g twice per week) or PBS alone starting from 5 days after tumor inoculation.

[0020] Figure 8. Growth of RPMI 4788 human colon tumor xenografts in mice treated with GAL-F2 or bFM-1 anti-FGF2 mAb (100 μ g twice per week) or PBS alone, starting from 6 days after tumor inoculation.

[0021] Figure 9. Growth of SMMC-7721 human hepatoma tumor xenografts in mice treated with PBS alone, GAL-F2 (100 μ g twice per week), cisplatin (100 μ g once per week), or both GAL-F2 and cisplatin. 5 mice per group.

[0022] Figure 10. Growth of HepG2 human hepatoma tumor xenografts in mice treated with PBS alone, GAL-F2 or M225 anti-EGFR mAb (100 μ g twice per week), or both GAL-F2 and M225, starting from 6 days after inoculation. 6 mice per group.

[0023] Figures 11A and 11B. Amino acid sequences of the HuGAL-F2 light chain (A) (SEQ ID NO:2) and heavy chain (B) mature variable regions (SEQ ID NO:5) are shown aligned with mouse GAL-F2 (SEQ ID NOS:1 and 4) and human acceptor V regions (SEQ ID NOS:3 and 6). The CDRs are underlined in the GAL-F2 sequences, and the amino acids substituted with mouse L2G7 amino acids are double underlined in the HuGAL-F2 sequences. The 1-letter amino acid code and Kabat numbering system are used for both the light and heavy chain.

[0024] Figure 12. Competitive binding of humanized (HuGAL-F2) and chimeric (ChGAL-F2) mAbs and control human antibody hIgG, conducted as described in the specification.

[0025] Figure 13. Amino acid sequences of the entire mature HuGAL-F2 antibody light chain (A) (SEQ ID NO:7) and heavy chain (B) (SEQ ID NO:8). The first amino acid on each line is numbered; the numbering is sequential. In the light chain, the first amino acid of the C κ region is underlined, and in the heavy chain, the first amino acids of the CH1, hinge, CH2 and CH3 regions are underlined.

62451-1087

DETAILED DESCRIPTION OF THE INVENTION

[0026] The invention provides neutralizing anti-FGF2 monoclonal antibodies, pharmaceutical compositions comprising them, and methods of using them for the treatment of disease.

1. Antibodies

5 **[0027]** Antibodies are very large, complex molecules (molecular weight of ~150,000 or about 1320 amino acids) with intricate internal structure. A natural antibody molecule contains two identical pairs of polypeptide chains, each pair having one light chain and one heavy chain. Each light chain and heavy chain in turn consists of two regions: a variable (“V”) region involved in binding the target antigen, and a constant (“C”) region that interacts with other components of the immune system.

10 The light and heavy chain variable regions come together in 3-dimensional space to form a variable region that binds the antigen (for example, a receptor on the surface of a cell). Within each light or heavy chain variable region, there are three short segments (averaging 10 amino acids in length) called the complementarity determining regions (“CDRs”). The six CDRs in an antibody variable domain (three from the light chain and three from the heavy chain) fold up together in 3-D space to form the

15 actual antibody binding site which locks onto the target antigen. The position and length of the CDRs have been precisely defined by Kabat, E. et al., Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human Services, 1983, 1987. The part of a variable region not contained in the CDRs is called the framework, which forms the environment for the CDRs. Chothia et al., J. Mol. Biol. 196:901, 1987, have defined the related concept of hypervariable regions

20 or loops determined by structure.

[0028] A humanized antibody is a genetically engineered antibody in which the CDRs from a mouse antibody (“donor antibody”, which can also be rat, hamster or other non-human species) are grafted onto a human antibody (“acceptor antibody”). The sequence of the acceptor antibody can be, for example, a mature human antibody sequence, a consensus sequence of human antibody sequences

25 or a germline sequence. Thus, a humanized antibody is an antibody having CDRs from a donor antibody and variable region framework and constant regions from a human antibody. In addition, in order to retain high binding affinity, at least one of two additional structural elements can be employed. See, US Patent No. 5,530,101 and 5,585,089, which provide detailed instructions for construction of humanized antibodies. A humanized antibody typically has a humanized heavy chain

30 and a humanized light chain. In general neither the heavy chain

variable region framework of a humanized heavy chain nor the light chain variable region framework of a humanized light chain includes more than ten or twelve substitutions resulting in residues not present in the acceptor human heavy or light chain variable region framework (including human consensus variable region frameworks and composite human variable region frameworks).

[0029] Although humanized antibodies often incorporate all six intact CDRs (as defined by Kabat) from a mouse antibody, they can also be made with less than the complete CDRs from a mouse antibody (e.g., Pascalis et al., *J. Immunol.* 169:3076, 2002). Numerous antibodies have been described in the scientific literature in which one or two CDRs can be dispensed with for binding. Padlan et al., *FASEB Journal* 9: 133-139 (1995); Vajdos et al., *Journal of Molecular Biology*, 320: 415-428 (2002); Iwahashi et al., *Mol. Immunol.* 36:1079-1091, (1999); Tamura et al., *Journal of Immunology*, 164:1432-1441 (2000). Similarly, it may be necessary to incorporate only part of the CDRs, namely the subset of CDR residues required for binding, termed the SDRs, into the humanized antibody.

[0030] CDR residues not contacting antigen and not in the SDRs can be identified based on previous studies (for example residues H60-H65 in CDRH2 are often not required), from regions of Kabat CDRs lying outside Chothia hypervariable loops, by molecular modeling and/or empirically, or as described in Gonzales et al., *Mol. Immunol.* 41: 863, 2004. If a CDR or residue(s) thereof is omitted, it is usually substituted with an amino acid occupying the corresponding position in the human acceptor sequence supplying the variable region framework sequences. The number of such substitutions to include reflects a balance of competing considerations. Such substitutions are potentially advantageous in decreasing the number of mouse amino acids in a humanized antibody and consequently decreasing potential immunogenicity. However, substitutions can also cause changes of affinity, and significant reductions in affinity are preferably avoided. Positions for substitution within CDRs and amino acids to substitute can also be selected empirically.

[0031] In the first structural element referred to above for retaining high binding affinity in a humanized antibody, the framework of the heavy chain variable region of the humanized antibody is chosen to have maximal sequence identity (between 65% and 95%) with the framework of the heavy chain variable region of the donor antibody, by suitably selecting the acceptor antibody from among the many known human antibodies. In the second structural element, in constructing the humanized antibody, selected amino acids in the framework of the human acceptor antibody (outside the CDRs) are replaced with corresponding amino acids from the donor antibody, in accordance with specified rules. Specifically, the amino

62451-1087

acids to be replaced in the framework are chosen on the basis of their ability to interact with the CDRs. For example, the replaced amino acids can be adjacent to a CDR in the donor antibody sequence or within 4-6 angstroms of a CDR in the humanized antibody as measured in 3-dimensional space.

[0032] A chimeric antibody is an antibody in which the variable region of a mouse (or other rodent) antibody is combined with the constant region of a human antibody; their construction by means of genetic engineering is well-known. Such antibodies retain the binding specificity of the mouse antibody, while being about two-thirds human. The proportion of nonhuman sequence present in mouse, chimeric and humanized antibodies suggests that the immunogenicity of chimeric antibodies is intermediate between mouse and humanized antibodies. Other types of genetically engineered antibodies that may have reduced immunogenicity relative to mouse antibodies include human antibodies made using phage display methods (Dower et al., WO91/17271; McCafferty et al., WO92/001047; Winter, WO92/20791; and Winter, FEBS Lett. 23:92, 1998) or using transgenic animals (Lonberg et al., WO93/12227; Kucherlapati WO91/10741).

[0033] As used herein, the term “human-like” antibody refers to a mAb in which a substantial portion of the amino acid sequence of one or both chains (e.g., about 50% or more) originates from human immunoglobulin genes. Hence, human-like antibodies encompass but are not limited to chimeric, humanized and human antibodies. As used herein, a mAb with “reduced immunogenicity” is one expected to have significantly less immunogenicity than a mouse antibody when administered to human patients. Such antibodies encompass chimeric, humanized and human mAbs as well as mAbs made by replacing specific amino acids in mouse antibodies that may contribute to B- or T-cell epitopes, for example exposed residues (Padlan, Mol. Immunol. 28:489, 1991). As used herein, a “genetically engineered” mAb is one for which the genes have been constructed or put in an unnatural environment (e.g., human genes in a mouse or on a bacteriophage) with the help of recombinant DNA techniques, and would therefore, e.g., not encompass a mouse mAb made with conventional hybridoma technology.

[0034] The epitope of a mAb is the region of its antigen to which the mAb binds. Two antibodies bind to the same or overlapping epitope if each competitively inhibits (blocks) binding of the other to the antigen. That is, a 1x, 5x, 10x, 20x or 100x excess of one antibody inhibits binding of the other by at least 50% but preferably 75%, 90% or even 99% as measured in a competitive binding assay (see, e.g., Junghans et al., Cancer Res. 50:1495,

1990). Alternatively, two antibodies have the same epitope if they bind the same region of the antigen or if essentially all amino acid mutations in the antigen that reduce or eliminate binding of one antibody reduce or eliminate binding of the other. Two antibodies have overlapping epitopes if some amino acid mutations that reduce or eliminate binding of one antibody reduce or eliminate binding of the other.

2. Neutralizing anti-FGF2 Antibodies

[0035] A monoclonal antibody (mAb) that binds FGF2 (i.e., an anti-FGF2 mAb) is said to neutralize FGF2, or be neutralizing, if the binding partially or completely inhibits one or more biological activities of FGF2 (i.e., when the mAb is used as a single agent). Among the biological properties of FGF2 that a neutralizing antibody may inhibit are the ability of FGF2 to bind to one or more of the four FGF receptors, to stimulate proliferation of (i.e., be mitogenic for) certain cells including fibroblasts, endothelial cells, Mv 1 Lu mink lung epithelial cells, and various human tumor cells; to stimulate differentiation and migration of cells such as endothelial cells, or to stimulate angiogenesis, for example as measured by stimulation of human vascular endothelial cell (HUVEC) proliferation or tube formation or by induction of blood vessels when applied to the chick embryo chorioallantoic membrane (CAM).

[0036] A neutralizing mAb of the invention at a concentration of, e.g., 0.01, 0.1, 0.5, 1, 2, 5, 10, 20 or 50 $\mu\text{g/ml}$ inhibits a biological function of FGF2 by about at least 50% but preferably 75%, more preferably by 90% or 95% or even 99%, and most preferably approximately 100% (essentially completely) as assayed by methods described under Examples or known in the art. Typically, the extent of inhibition is measured when the amount of FGF2 used is just sufficient to fully stimulate the biological activity, or is 1, 2, or 5 ng/ml or 0.01, 0.02, 0.05, 0.1, 0.5, 1, 3 or 10 $\mu\text{g/ml}$. Preferably, the mAb is neutralizing, i.e., inhibits the biological activity, when used as a single agent, but optionally 2 mAbs can be used together to give inhibition. Most preferably, the mAb neutralizes not just one but two, three or several of the biological activities listed above; for purposes herein, an anti-FGF2 mAb that used as a single agent neutralizes all the biological activities of FGF2 is called “fully neutralizing”, and such mAbs are most preferable. MAb of the invention are preferably specific for FGF2, that is they do not bind, or only bind to a much lesser extent (e.g., at least 10-fold less), proteins that are related to FGF2 such as the other FGFs, e.g., FGF1, and vascular endothelial growth factor (VEGF). Some mAbs of the invention bind both human FGF2 and mouse FGF2, or bind human FGF2 and one, two or more or all of mouse, rat, rabbit, chicken, dog and/or monkey (e.g., cynomolgus monkey) FGF2. Other

62451-1087

mAbs are specific for human FGF2. MAbs of the invention typically have a binding affinity (K_a) for FGF2 of at least 10^7 M^{-1} but preferably 10^8 M^{-1} or higher, and most preferably 10^9 M^{-1} or higher or even 10^{10} M^{-1} or higher.

[0037] MAbs of the invention include anti-FGF2 antibodies in their natural tetrameric form (2 light chains and 2 heavy chains) and may be of any of the known isotypes IgG, IgA, IgM, IgD and IgE and their subtypes, i.e., human IgG1, IgG2, IgG3, IgG4 and mouse IgG1, IgG2a, IgG2b, and IgG3. The mAbs of the invention are also meant to include fragments of antibodies such as Fv, Fab and F(ab')₂; bifunctional hybrid antibodies (e.g., Lanzavecchia et al., *Eur. J. Immunol.* 17:105, 1987), single-chain antibodies (Huston et al., *Proc. Natl. Acad. Sci. USA* 85:5879, 1988; Bird et al., *Science* 242:423, 1988); single-arm antibodies (Nguyen et al., *Cancer Gene Ther.* 10:840, 2003); and antibodies with altered constant regions (e.g., U.S. Patent No. 5,624,821). The mAbs may be of animal (e.g., mouse, rat, hamster or chicken) origin, or they may be genetically engineered. Rodent mAbs are made by standard methods well-known in the art, comprising multiple immunization with FGF2 in appropriate adjuvant i.p., i.v., or into the footpad, followed by extraction of spleen or lymph node cells and fusion with a suitable immortalized cell line, and then selection for hybridomas that produce antibody binding to FGF2, e.g., see under Examples. Chimeric and humanized mAbs, made by art-known methods mentioned supra, are preferred embodiments of the invention. Human antibodies made, e.g., by phage display or transgenic mice methods are also preferred (see e.g., Dower et al., McCafferty et al., Winter, Lonberg et al., Kucherlapati, supra).

[0038] The neutralizing anti-FGF2 mAb GAL-F2 described infra is an example of the invention. Once a single, archetypal anti-human-FGF2 mAb, for example GAL-F2, has been isolated that has the desired properties described herein of neutralizing FGF2, it is straightforward to generate other mAbs with similar properties, by using art-known methods. For example, mice may be immunized with FGF2 as described above, hybridomas produced, and the resulting mAbs screened for the ability to compete with the archetypal mAb for binding to FGF2. Mice can also be immunized with a smaller fragment of FGF2 containing the epitope to which GAL-F2 binds. The epitope can be localized by, e.g., screening for binding to a series of overlapping peptides spanning FGF2. Alternatively, the method of Jespers et al., *Biotechnology* 12:899, 1994, may be used to guide the selection of mAbs having the same epitope and therefore similar properties to the archetypal mAb, e.g., GAL-F2. Using phage display, first the heavy chain of the archetypal antibody is paired with a repertoire of (preferably human) light chains to select an

FGF2-binding mAb, and then the new light chain is paired with a repertoire of (preferably human) heavy chains to select a (preferably human) FGF2-binding mAb having the same epitope as the archetypal mAb. Alternatively variants of GAL-F2 can be obtained by mutagenesis of cDNA encoding the heavy and light chains of GAL-F2 obtained from the hybridoma.

[0039] Neutralizing mAbs with the same or overlapping epitope as GAL-F2, e.g., that compete for binding to FGF2 with GAL-F2, provide other examples. A chimeric or humanized form of GAL-F2 is an especially preferred embodiment. MAbs that are 90%, 95% or 99% identical to GAL-F2 in amino acid sequence of the heavy and/or light chain variable regions and maintain its functional properties, and/ or which differ from it by a small number of functionally inconsequential amino acid substitutions (e.g., conservative substitutions), deletions, or insertions are also included in the invention. MAbs having at least one and preferably all six CDR(s) that are 90%, 95% or 99% or 100% identical to corresponding CDRs of GAL-F2 are also included. Here, as elsewhere in this application, percentage sequence identities are determined with antibody sequences maximally aligned by the Kabat numbering convention. After alignment, if a subject antibody region (e.g., the entire mature variable region of a heavy or light chain) is being compared with the same region of a reference antibody, the percentage sequence identity between the subject and reference antibody regions is the number of positions occupied by the same amino acid in both the subject and reference antibody region divided by the total number of aligned positions of the two regions, with gaps not counted, multiplied by 100 to convert to percentage.

[0040] Native mAbs of the invention may be produced from their hybridomas. Genetically engineered mAbs, e.g., chimeric or humanized mAbs, may be expressed by a variety of art-known methods. For example, genes encoding their light and heavy chain V regions may be synthesized from overlapping oligonucleotides and inserted together with available C regions into expression vectors (e.g., commercially available from Invitrogen) that provide the necessary regulatory regions, e.g., promoters, enhancers, poly A sites, etc. Use of the CMV promoter-enhancer is preferred. The expression vectors may then be transfected using various well-known methods such as lipofection or electroporation into a variety of mammalian cell lines such as CHO or non-producing myelomas including Sp2/0 and NS0, and cells expressing the antibodies selected by appropriate antibiotic selection. See, e.g., US Patent No. 5,530,101. Larger amounts of antibody may be produced by growing the cells in commercially available bioreactors.

[0041] Once expressed, the mAbs or other antibodies of the invention may be purified according to standard procedures of the art such as microfiltration, ultrafiltration, protein A or G affinity chromatography, size exclusion chromatography, anion exchange chromatography, cation exchange chromatography and/or other forms of affinity chromatography based on organic dyes or the like. Substantially pure antibodies of at least about 90 or 95% homogeneity are preferred, and 98% or 99% or more homogeneity most preferred, for pharmaceutical uses.

3. Treatment Methods

[0042] The invention provides methods of treatment in which the mAb of the invention (e.g., anti-FGF2) is administered to patients having a disease (therapeutic treatment) or at risk of occurrence or recurrence of a disease (prophylactic treatment). The term "patient" includes human patients; veterinary patients, such as cats, dogs and horses; farm animals, such as cattle, sheep, and pigs; and laboratory animals used for testing purposes, such as mice and rats. The methods are particularly amenable to treatment of human patients. The mAb used in methods of treating human patients binds to the human FGF2 protein, the sequence of which is provided by, e.g., Ornitz et al., *Genome Biol.* 2: 3005.1, 2001 or Okada-Ban et al., *Int. J. Biochem. Cell. Biol.* 32:263, 2000, also Locus P09038 of Swiss-Prot database. A mAb to a human protein can also be used in other species in which the species homolog has antigenic crossreactivity with the human protein. In species lacking such crossreactivity, an antibody is used with appropriate specificity for the species homolog present in that species. However, in xenograft experiments in laboratory animals, a mAb with specificity for the human protein expressed by the xenograft is generally used.

[0043] In a preferred embodiment, the present invention provides a pharmaceutical formulation comprising the antibodies described herein. Pharmaceutical formulations of the antibodies contain the mAb in a physiologically acceptable carrier, optionally with excipients or stabilizers, in the form of lyophilized or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or acetate at a pH typically of 5.0 to 8.0, most often 6.0 to 7.0; salts such as sodium chloride, potassium chloride, etc. to make isotonic; antioxidants, preservatives, low molecular weight polypeptides, proteins, hydrophilic polymers such as polysorbate 80, amino acids, carbohydrates, chelating agents, sugars, and other standard ingredients known to those skilled in the art (Remington's Pharmaceutical Science 16th edition, Osol, A. Ed. 1980). The mAb is typically present at a concentration of 1 - 100 mg/ml, e.g., 10 mg/ml.

[0044] In another preferred embodiment, the invention provides a method of treating a patient with a disease using an anti-FGF2 mAb in a pharmaceutical formulation. The mAb prepared in a pharmaceutical formulation can be administered to a patient by any suitable route, especially parentally by intravenous infusion or bolus injection, intramuscularly or subcutaneously. Intravenous infusion can be given over as little as 15 minutes, but more often for 30 minutes, or over 1, 2 or even 3 hours. The mAb can also be injected directly into the site of disease (e.g., a tumor), or encapsulated into carrying agents such as liposomes. The dose given is sufficient at least partially to alleviate the condition being treated ("therapeutically effective dose") and is optionally 0.1 to 5 mg/kg body weight, for example 1, 2, 3 or 4 mg/kg, but may be as high as 10 mg/kg or even 15, 20 or 30 mg/kg. A fixed unit dose may also be given, for example, 100, 200, 500, 1000 or 2000 mg, or the dose may be based on the patient's surface area, e.g., 1000 mg/m². Usually between 1 and 8 doses, (e.g., 1, 2, 3, 4, 5, 6, 7 or 8) are administered to treat cancer, but 10, 20 or more doses may be given. The mAb can be administered daily, biweekly, weekly, every other week, monthly or at some other interval, depending, e.g. on the half-life of the mAb, for 1 week, 2 weeks, 4 weeks, 8 weeks, 3-6 months or longer or until the disease progresses. Repeated courses of treatment are also possible, as is chronic administration.

[0045] A combination of a dose, frequency of administration and route of administration effective to at least partially alleviate a disease present in a patient being treated is referred to as therapeutically effective regime. A combination of a dose, frequency of administration and route of administration effective to inhibit or delay onset of a disease in a patient is referred to as a prophylactically effective regime.

[0046] Diseases especially susceptible to treatment with the anti-FGF2 mAbs of this invention include solid tumors believed to require angiogenesis, or to be associated with elevated levels of FGF2, or to be associated with expression of FGF2. Such tumors, for which treatment with the anti-FGF2 mAb is appropriate, include for example ovarian cancer, breast cancer, lung cancer (small cell or non-small cell), colon cancer, prostate cancer, cervical cancer, endometrial cancer, pancreatic cancer, gastric cancer, hepatocellular carcinoma or hepatoma (liver cancer), head-and-neck tumors, melanoma, sarcomas, carcinomas, and brain tumors (e.g., gliomas such as glioblastomas). Hematologic malignancies such as leukemias and lymphomas and multiple myeloma can also be susceptible to such treatment. Other diseases associated with angiogenesis for which treatment with the anti-FGF mAbs of the invention are suitable include age-related macular

62451-1087

degeneration (AMD), diabetic retinopathy, neovascular glaucoma and other diseases of the eye; psoriasis and other diseases of the skin; and rheumatoid arthritis.

- [0047]** In a preferred embodiment, the anti-FGF2 mAb is administered in combination with (i.e., together with, that is, before, during or after) other therapy. For example, to treat cancer, the anti-FGF2 mAb may be administered together with any one or more of the known chemotherapeutic drugs, for example alkylating agents such as carmustine, chlorambucil, cisplatin, carboplatin, oxaliplatin, procarbazine, and cyclophosphamide; antimetabolites such as fluorouracil, floxuridine, fludarabine, gemcitabine, methotrexate and hydroxyurea; natural products including plant alkaloids and antibiotics such as bleomycin, doxorubicin, daunorubicin, idarubicin, etoposide, mitomycin, mitoxantrone, vinblastine, vincristine, and Taxol (paclitaxel) or related compounds such as Taxotere®; the topoisomerase 1 inhibitor irinotecan; agents specifically approved for brain tumors including temozolomide and Gliadel® wafer containing carmustine; and inhibitors of tyrosine kinases such as Gleevec® (imatinib mesylate), Sutent® (sunitinib malate), Nexavar® (sorafenib), Tarceva® (erlotinib) and Iressa® (gefitinib); inhibitors of angiogenesis; and all approved and experimental anti-cancer agents listed in WO 2005/017107 A2. The anti-FGF2 mAb may be used in combination with 1, 2, 3 or more of these other agents used in a standard chemotherapeutic regimen. Normally, the other agents are those already known to be effective for the particular type of cancer being treated. The anti-FGF2 mAb is especially useful in overcoming resistance to chemotherapeutic drugs and thereby increasing their effectiveness (see Song et al. Proc. Natl. Acad. Sci USA 97:8658, 2000).
- [0048]** Other agents with which the anti-FGF2 mAb can be administered to treat cancer include biologics such as monoclonal antibodies, including Herceptin™ against the HER2 antigen; Avastin® against VEGF; or antibodies to the Epidermal Growth Factor (EGF) receptor such as Erbitux® (cetuximab) and Vectibix® (panitumumab). Antibodies against Hepatocyte Growth Factor (HGF) are especially preferred for use with the anti-FGF2 mAb, including mAb L2G7 (Kim et al., Clin Cancer Res 12:1292, 2006 and US Patent No. 7,220,410) and particularly its chimeric and humanized forms such as HuL2G7 (WO 07115049 A2); the human anti-HGF mAbs described in WO 2005/017107 A2, particularly 2.12.1; and the HGF binding proteins described in WO 07143090 A2 or WO 07143098 A2; and other neutralizing anti-HGF mAbs that compete for binding with any of the aforementioned mAbs. A mAb that binds the cMet receptor of HGF is also preferred, for example the anti-cMet mAb OA-5D5 (Martens et al., Clin. Cancer Res. 12:6144, 2006) that has been genetically engineered to have only one “arm”, i.e. binding domain. Moreover, the

anti FGF2 mAb can be used together with any form of surgery and/or radiation therapy including external beam radiation, intensity modulated radiation therapy (IMRT) and any form of radiosurgery such as, e.g. Gamma Knife.

[0049] Treatment (e.g., standard chemotherapy) including the anti-FGF2 mAb antibody may alleviate a disease by increasing the median progression-free survival or overall survival time of patients with cancer by at least 30% or 40% but preferably 50%, 60% to 70% or even 100% or longer, compared to the same treatment (e.g., chemotherapy) but without the anti-FGF2 mAb. In addition or alternatively, treatment (e.g., standard chemotherapy) including the anti-FGF2 mAb may increase the complete response rate, partial response rate, or objective response rate (complete + partial) of patients with these tumors (e.g., ovarian, breast, lung, colon and glioblastomas especially when relapsed or refractory) by at least 30% or 40% but preferably 50%, 60% to 70% or even 100% compared to the same treatment (e.g., chemotherapy) but without the anti-FGF2 mAb.

[0050] Typically, in a clinical trial (e.g., a phase II, phase II/III or phase III trial), the aforementioned increases in median progression-free survival and/or response rate of the patients treated with chemotherapy plus the anti-FGF2 mAb, relative to the control group of patients receiving chemotherapy alone (or plus placebo), are statistically significant, for example at the $p = 0.05$ or 0.01 or even 0.001 level. The complete and partial response rates are determined by objective criteria commonly used in clinical trials for cancer, e.g., as listed or accepted by the National Cancer Institute and/or Food and Drug Administration.

4. Other Methods

[0051] The anti-FGF2 mAbs of the invention also find use in diagnostic, prognostic and laboratory methods. They may be used to measure the level of FGF2 in a tumor or in the circulation of a patient with a tumor, to determine if the level is measurable or even elevated, and therefore to follow and guide treatment of the tumor, since tumors associated with measurable or elevated levels of FGF2 will be most susceptible to treatment with the anti-FGF2 mAb. For example, a tumor associated with high levels of FGF2 would be especially susceptible to treatment with an anti-FGF2 mAb. In particular embodiments, the mAbs can be used in an ELISA or radioimmunoassay to measure the level of FGF2, e.g., in a tumor biopsy specimen or in serum or in media supernatant of FGF2-secreting cells in cell culture. The use of two anti-FGF2 mAbs binding to different epitopes (i.e., not competing for binding) will be especially useful in developing a sensitive “sandwich” ELISA to detect FGF2. For various assays, the mAb may be labeled with fluorescent molecules, spin-labeled

molecules, enzymes or radioisotopes, and may be provided in the form of kit with all the necessary reagents to perform the assay for FGF2. In other uses, the anti-FGF2 mAbs will be used to purify FGF2, e.g., by affinity chromatography.

EXAMPLES

Example 1: Reagents and Assays

[0052] Preparation of GST-FGF2, FGF2-Fc and FGF2-Flag. The cDNA sequence for human FGF2 (the precursor form with 155 amino acids; Sommer et al., 1987) was synthesized (GenScript, Inc), PCR amplified and cloned into a derivative of the pDisplay vector (Invitrogen). These plasmids were transformed into E. coli BL21(DE3) cells and FGF2 expression was induced using 1 mM IPTG. The level of FGF2 expression was determined using an FGF2 specific ELISA Kit (R&D Systems), and FGF2 was purified using heparin-Sepharose CL-6B beads (Amersham Biosciences) as described (Wiedlocha et al., Mol. Cell Biol. 16:270, 1996). Fusion proteins of FGF2 with respectively glutamine synthetase (GST-FGF2), Flag peptide (FGF2-Flag), and human IgG1 Fc domain (amino acids 216 to 446; FGF2-Fc), which were used in the FGF2/FGFR binding assays as well as for immunization, were produced similarly from the appropriate genetic constructs using standard molecular biology techniques. GST-FGF2 was purified using an anti-GST column, FGF2-Fc was purified using a protein A/G column, and FGF-Flag was used in culture supernatant after determination of the FGF2-Flag concentration. In addition, purified human FGF2 (QED Bioscience Inc.) and murine FGF2 (ProSpec-Tany Technogene Ltd.) were purchased.

[0053] Preparation of FGFR-Fc proteins. The extracellular domain (ECD) of human FGFR1 alpha (IIIc) (designated as FGFR1c) and human FGFR2C alpha (IIIc) (designated as FGFR2c) were expressed as immunoadhesin molecules. DNA fragments encoding the entire ECD of FGFR1c and FGFR2c were fused to human Fc via a polypeptide linker. These FGFR-Fc molecules were expressed by transfecting human 293F cells and selecting stable 293 transfectants in the presence of G418 (1 mg/ml) in 293 expression medium (Invitrogen). The FGFR-Fc secreted from 293F transfected cells was purified using a protein A/G column. In addition, human FGFR3c-Fc and human FGFR4-Fc fusion were obtained from R & D Systems.

[0054] Synthesis of FGF2 fragments. Peptides consisting of amino acid residues 29-44 (Peptide #1), 101-118 (Peptide #3) and 137-155 (peptide #4) of FGF2 were synthesized by SynBioSci. Extra cysteine residues were added to the c-terminus of Peptide #1 and #3 and

the N-terminus of peptide #4. These peptide fragments were then conjugated to keyhole limpet hemocyanin (KLH).

[0055] FGF2 binding ELISA. ELISA wells were coated with 50 µg/ml of heparin (Sigma) overnight at 4°C and then incubated with 0.3 -1 µg/ml of either human or mouse FGF2 for 1 hr at room temperature (RT) so that the heparin could capture the FGF2. After blocking with 2% BSA for 1 hr at RT, hybridoma culture supernatants or purified mAbs were added for 1 hr at RT. The bound anti-FGF2 antibodies were detected by the addition of HRP-goat anti-mouse IgG Fc for 1 hr at RT, followed by washing, addition of TMB substrate (Sigma) and reading at 450 nm.

[0056] FGFR-Fc/FGF2-Flag Binding ELISA. The blocking activity of anti-FGF2 mAbs was determined in the FGFR-Fc/FGF2-Flag binding ELISA. ELISA wells were coated with 2 µg/ml of goat antibody specific for human IgG-Fc overnight at 4°C. After blocking with 2% BSA for 1 hr at RT, the wells were incubated with 0.5 µg/ml of FGFR1c-Fc, FGFR2c-Fc, FGFR3c-Fc or FGFR4-Fc for 1 hr. After washing, wells were incubated with FGF2-Flag (0.2 µg/ml) in the presence of various concentrations of mAbs for 1 hr. The bound FGF2-Flag was detected by the addition of HRP-anti-Flag M2 antibody (Sigma).

Example 2: Generation of monoclonal antibodies

[0057] Balb/c mice (5-6 week old female) were immunized by injection in their rear footpads 14 times at 1 week intervals with GST-FGF2, FGF2-Fc and/or KLH conjugated FGF2 synthetic peptides, resuspended in MPL/TDM (Sigma-Aldrich), as described in the table below. Three days after the final injection, popliteal lymphoid cells were fused with P3/X63-Ag8U1 mouse myeloma cells using standard fusion methods with 35% polyethylene glycol as described (Chuntharapai et al., Methods Enzymol 288:15, 1997). Ten days after the fusion, hybridoma culture supernatants were screened for their ability to bind to FGF2 using the FGF2 binding ELISA described above. Selected mAbs were then screened for their blocking activities in the FGFR1-Fc/ FGF-Flag binding ELISA. Selected hybridomas were then cloned twice using the limiting dilution technique as described (Harlow et al., 1988).

Table: Immunization protocol

Injection No.	Antigens (in MPL/TDM) per footpad
1	GST-FGF2 10 μ g
2	GST-FGF2 5 μ g
3	GST-FGF2 2.5 μ g
4 – 7	GST-FGF2 5 μ g
8 – 9	GST-FGF2 10 μ g
10	GST-FGF2 (5 μ g) + FGF2-Fc (5 μ g)
11	FGF2-Fc (2 μ g) + Peptide #1 & # 3 (2 μ g)
12	FGF2-Fc (2 μ g) + Peptide #1, #3 & #4 (3 μ g)
13	FGF2-Fc (2 μ g) + Peptide #1, #3 & #4 (3 μ g)
14	FGF2-Fc 2 μ g

[0058] Figure 1 shows that the mAb GAL-F2 generated in this way binds to both human and mouse FGF2, approximately equally well, whereas a negative control mouse IgG mAb does not bind to FGF2. Figure 2 shows that GAL-F2 but not the control mAb 5G8 is able to inhibit binding of human FGF2 to each of the four FGF receptors FGFR1-4. In another experiment using the FGFR-Fc/FGF2-Flag Binding ELISA described above, GAL-F2 (at a concentration of 10 μ g/ml) completely inhibited the binding of FGF2 to each of the four FGFRs – FGFR1, FGFR2, FGFR3 and FGFR4 – that is, reduced the signal to background level within experimental error, while the anti-FGF2 mAbs bFM-1 and 3H3 reduced binding substantially but not completely.

Example 3: Epitope of GAL-F2

[0059] Several commercially available anti-FGF2 mAbs that have been reported to have certain neutralizing activities -- 3H3 (Calbiochem), mAb bFM-1 (Millipore) and mAb FB-8 (Abcam) -- were purchased. A competitive binding ELISA was performed to determine if GAL-F2 has the same epitope as any of these mAbs. ELISA plates were coated with 50 μ l/well of heparin (50 μ g/ml) overnight, decanted and incubated with 50 μ l/well of FGF2 for 1 hr. After blocking with 2% BSA, wells were incubated with 0.5 μ g/ml of biotinylated GAL-F2 mAb in the presence of various concentrations of each anti-FGF2 mAb to be tested. The level of biotinylated GAL-F2 binding to the FGF2 in each well was detected by the addition of HRP-streptavidin followed by TMB substrate. Figure 3 shows that none of the mAbs 3H3, bFM-1 and FB-8 nor a negative control mouse mAb (mIgG) competed for binding to FGF2 with GAL-F2, while of course GAL F2 competed with itself. Hence no previous anti-FGF2 mAb tested has the same or overlapping epitope on FGF2 as GAL-F2.

Example 4: Inhibition of FGF2-induced proliferation by GAL-F2

[0060] BaF3 cells (DSMZ, Germany) are maintained in RPMI 1640 medium (GIBCO) supplemented with 10% FCS, 10% WEHI-3 conditioned medium and antibiotics. Stable FGFR1c or FGFR2c expressing Ba/F3 cells were generated by electroporation of linearized FGFR1c and FGFR2c expression vector DNAs as described (Ornitz et al., 1996). Stable transfected cells are selected using G418 (600 μ g/ml) for 10 days. The stable Ba/F3 transfectants expressing FGFR1c or FGFR2c were shown to proliferate in response to FGF2 in the absence of WEHI-3 medium. To determine the blocking (neutralizing) activity of GAL-F2 mAb, the FGFR1c- or FGFR2c- expressing BaF3 cells (10,000 cells/ well) were washed and resuspended in RPMI with 10% FCS plus 2 μ g/ml of heparin and 20 ng/ml of FGF-2 in the presence of various concentrations of GAL-F2. After incubation for 36-48 hrs at 37°C, 5% CO₂, the level of proliferation was determined by the addition of WST-1 (Roche Applied Science) for 2 hrs. Figure 4 shows that GAL F2 inhibited FGF2-induced proliferation, almost completely at 10 μ g/ml mAb.

[0061] The blocking activities GAL-F2 on normal epithelial cells were determined using Mv 1 Lu mink lung epithelial cells (CCL-64 from ATCC). Mv 1 Lu cells (2 x 10³ cells/100 μ l/well) grown in DMEM containing 10% FCS were resuspended in serum-free DMEM and stimulated with 1-2 ng/ml of FGF2, plus 1 ng/ml of TGF- β to inhibit background proliferation, in the presence of various concentrations of mAb for 24 hrs. The level of cell proliferation was determined by the addition of WST-1 (Roche Applied Science) for 24-48 hr. Figure 5 shows that GAL-F2 inhibited FGF2-induced proliferation of the Mv 1 Lu cells, almost completely at 1 μ g/ml mAb. In a similar experiment, GAL-F2 inhibited proliferation of HUVEC induced by 10 ng/ml FGF2, by about 75% at a concentration of 0.1 μ g/ml (i.e., at an approximately equimolar ratio to FGF2) and completely at a concentration of 1.0 μ g/ml. This was somewhat better than the extent of inhibition by the bFM-1 anti-FGF2 mAb and substantially better than by the 3H3 anti-FGF2 mAb, and suggests that GAL-F2 inhibits angiogenesis, since proliferation of endothelial cells is an essential step in angiogenesis.

Example 5: Clonogenic Assay

[0062] The anti-tumor activity of GAL-F2 was investigated in vitro by its effect on the colony formation of human tumor cells in soft agar. The assay was performed as follows: 6-well plates was coated with 1.5 ml/ well of 0.6% agar in DMEM containing 10% FCS and then layered with 1.5 ml of 0.3% agar in DMEM with 10% FCS. The top agar layer was mixed with 2 x 10⁴ SMMC-7721 human hepatoma tumor cells plus 10 μ g/ml of mAb. Plates were incubated at 37°C in a humidified incubator for 10 to 14 days and then stained with

0.005% crystal violet for 1 hr and examined by photomicroscopy. Figure 6 shows that compared to cells in the presence of irrelevant control mouse IgG mAb, cells in the presence of GAL-F2 formed many fewer colonies, while the bFM-1 anti-FGF2 mAb did not reduce the number of colonies. Hence, GAL-F2 inhibited colony formation in soft agar of human tumor cells.

Example 6: Angiogenesis Assay

[0063] The anti-angiogenic activity of GAL-F2 was determined in BALB/c mice injected in the back with 0.4 ml of matrigel (BD Biosciences) with or without 30 ng of FGF2 and/or 3 μ g GAL-F2. Matrigel plugs were harvested for photography on day 6. FGF2 stimulated the formation of blood vessels in this assay, while GAL-F2 at least partially inhibited this stimulation.

Example 7: Xenograft Models

[0064] Xenograft experiments are carried out as described previously (Kim et al., Nature 362:841, 1993). Human tumor cells typically grown in complete DMEM medium are harvested in HBSS. Female athymic nude mice or NIH-III Xid/Beige/nude mice (4-6 wks old) are injected subcutaneously with $2-10 \times 10^6$ cells in 0.1 ml of HBSS in the dorsal areas. When the tumor size reaches 50-100 mm³, the mice are grouped randomly and 5 mg/kg (100 μ g total) of mAbs are administered i.p. twice per week in a volume of 0.1 ml. Tumor sizes are determined twice a week by measuring in two dimensions [length (a) and width (b)]. Tumor volume is calculated according to $V = ab^2/2$ and expressed as mean tumor volume \pm SEM. The number of mice in each treatment group is typically 5-7 mice. Statistical analysis can be performed, e.g., using Student's t test.

[0065] Figure 7 shows that GAL-F2 strongly inhibited the growth of RPMI 4788 colon tumor xenografts, and Figure 8 shows that the bFM-1 anti-FGF2 mAb inhibited xenograft growth to a lesser extent than GAL-F2. Figure 9 shows that GAL-F2 inhibited SMMC-7721 hepatoma tumor xenografts in Xid/Beige/nude mice, while the chemotherapeutic drug cisplatin at 5 mg/kg once per week inhibited xenograft growth to a lesser extent. The combination of GAL-F2 and cisplatin inhibited more strongly than either agent alone, showing an additive or synergistic effect of these agents. Figure 10 shows that GAL-F2 inhibited HepG2 hepatoma tumor xenografts in nude mice more strongly than the anti-EGF receptor mAb M225 from which Erbitux® was derived, but the combination of GAL-F2 and M225 inhibited somewhat more strongly than either agent, again showing an additive or synergistic effect of these agents. However, GAL F2 was not able to inhibit the growth of all xenografts tested, possibly because they are not dependent on FGF2 for their growth.

62451-1087

Example 8: Humanization of GAL-F2

[0066] Cloning of the light and heavy chain variable regions of the GAL-F2 mAb, construction and expression of a chimeric mAb, and design, construction, expression and purification of a humanized GAL-F2 mAb were all performed using standard methods of molecular biology, e.g. as described in US Patent Application 11/731,774 for the L2G7 mAb. The amino acid sequences of the (mature) light and heavy chain variable (V) regions of GAL-F2 are shown respectively in Fig. 11A and 11B, top lines labeled GAL-F2. More specifically, to design a humanized GAL-F2 mAb, the methods of Queen et al., US Patent Nos. 5,530,101 and 5,585,089 were generally followed. The human V_κ sequence ABA70776 and V_H sequence AAL04519, as shown respectively in Fig. 11A and 11B, bottom lines, were respectively chosen to serve as acceptor sequences for the GAL-F2 VL and VH sequences because they have particularly high framework homology (i.e., sequence identity) to them. A computer-generated molecular model of the GAL-F2 variable domain was used to locate the amino acids in the GAL-F2 framework that are close enough to the CDRs to potentially interact with them. To design the humanized GAL-F2 light and heavy chain variable regions, the CDRs from the mouse GAL-F2 mAb were first conceptually grafted into the acceptor framework regions. At framework positions where the computer model suggested significant contact with the CDRs, which may be needed to maintain the CDR conformation, the amino acids from the mouse antibody were substituted for the human framework amino acids. For the humanized GAL-F2 mAb designated HuGAL-F2, this was done at residues 1, 27 and 30 (residues 27 and 30 being within Chothia hypervariable loop H1), 48, 67, 71 and 94 of the heavy chain and at no residues in the light chain, using Kabat numbering. The light and heavy chain V region sequences of HuGAL-F2 are shown in Fig. 11A and 11B respectively, middle lines labeled HuGAL-F2, where they are aligned against the respective GAL-F2 donor and human acceptor V regions – the CDRs (as defined by Kabat) are underlined and the substituted amino acids listed above are double-underlined.

[0067] The invention provides not only a humanized GAL-F2 mAb HuGAL-F2 including the light and heavy chain V regions shown in Fig. 11, but also variant humanized GAL-F2 mAbs whose light and heavy chain variable regions differ from the sequences of HuGAL-F2 by a small number (e.g., typically no more than 1, 2, 3, 5 or 10) of replacements, deletions or insertions, usually in the framework but possibly in the CDRs. In particular, only a subset of the substitutions described above can be made in the acceptor frameworks, or additional substitution(s) can be made, e.g., the mouse GAL-F2 VH amino acid 69L may replace the

acceptor amino acid 69I, or the mouse amino acids may replace the respective amino acids in the humanized light chain at any or all of the positions 1, 3, 60 and/or 67 by Kabat numbering. On the other hand, the VH amino acid 1E (Glu) may instead be Q (Gln). Indeed, many of the framework residues not in contact with the CDRs in HuGAL-F2 can accommodate substitutions of amino acids from the corresponding positions of GAL-F2 or other mouse or human antibodies, and even many potential CDR-contact residues are also amenable to substitution or even amino acids within the CDRs may be altered. One example of a CDR substitution is to substitute a residue in a CDR with the residue occupying the corresponding position of the human acceptor sequence used to supply variable region frameworks.

[0068] Most often the replacements made in the variant humanized GAL-F2 sequences are conservative with respect to the replaced HuGAL-F2 amino acids. Amino acids can be grouped as follows for determining conservative substitutions, i.e., substitutions within a group: Group I (hydrophobic sidechains): met, ala, val, leu, ile; Group II (neutral hydrophilic side chains): cys, ser, thr; Group III (acidic side chains): asp, glu; Group IV (basic side chains): asn, gln, his, lys, arg; Group V (residues influencing chain orientation): gly, pro; and Group VI (aromatic side chains): trp, tyr, phe.

[0069] Preferably, replacements in HuGAL-F2 (whether or not conservative) have no substantial effect on the binding affinity or potency of the humanized mAb, that is, its ability to neutralize the biological activities of FGF2 (e.g., the potency in some or all of the assays described herein of the variant humanized GAL-F2 mAb is essentially the same, i.e., within experimental error, as that of HuGAL-F2). Preferably the mature variant light and heavy chain V region sequences are at least 90%, more preferably at least 95%, and most preferably at least 98% identical to the respective HuGAL-F2 mature light and heavy chain V regions. Alternatively, other human antibody variable regions with high sequence identity to those of GAL-F2 are also suitable to provide the humanized antibody framework, especially kappa V regions from human subgroup I and heavy chain V regions from human subgroup I, or consensus sequences of these subgroups.

[0070] In other humanized antibodies, at least 1, 2, 3, 4, 5, 6 or all seven of the positions of acceptor to donor substitutions mentioned in connection with the exemplified antibody (i.e., H1, H27, H30, H48, H67, H71 and 94) are preferably occupied by the residue occupying the corresponding position of the mouse donor antibody heavy chain. If the heavy chain acceptor sequence is other than AAL04519 an acceptor to donor substitution may or may not be required for the specified occupancy of a particular variable framework region position

depending whether the residue occupying the specified position is already the same between the acceptor and donor.

[0071] The exemplary mAb HuGAL-F2 discussed here has human κ and $\gamma 1$ constant regions, e.g., as presented in US Patent Application 11/731,774, and is therefore an IgG1. The complete sequences of the (mature) light and heavy chains of HuGAL-F2 are shown in Fig. 13. Thus, an exemplary humanized antibody comprises a light chain of SEQ ID NO:7 and a heavy chain of SEQ ID NO:8. While these sequences are respectively of the Km(3) and G1m(3) allotypes, it is understood that IgG1 mAbs of any (IgG1, κ) allotype are encompassed by the designation HuGAL-F2. It will also be understood that when HuGAL-F2 is manufactured by conventional procedures, one to several amino acids at the amino or carboxy terminus of the light and/or heavy chain, such as the C-terminal lysine of the heavy chain, may be missing or derivatized in a proportion or all of the molecules, and such a composition will still be encompassed by the designation HuGAL-F2 and considered a humanized GAL-F2 mAb. Humanized mAbs of other isotypes (e.g., IgG2, IgG3 and IgG4) can be made by combining the HuGAL-F2 variable regions with the appropriate human constant regions. Replacements can be made in the HuGAL-F2 constant regions to reduce or increase effector function such as complement-mediated cytotoxicity or ADCC (see, e.g., Winter et al., US Patent No. 5,624,821; Tso et al., US Patent No. 5,834,597; and Lazar et al., Proc. Natl. Acad. Sci. USA 103:4005, 2006), or to prolong half-life in humans (see, e.g., Hinton et al., J. Biol. Chem. 279:6213, 2004). Specifically but without limitation, HuGAL-F2 having mutations in the IgG constant region to a Gln at position 250 and/or a Leu at position 428 are embodiments of the present invention.

[0072] To compare the binding affinity of HuGAL-F2 with that of the mouse-human chimeric mAb ChGAL-F2, a competitive binding experiment was performed using standard ELISA technology. Specifically, human FGF2 was immobilized on a heparin-coated ELISA plate. The wells were incubated with biotinylated GAL-F2 mAb (0.5 $\mu\text{g/ml}$) in the presence of increasing concentrations of unlabeled ChGAL-F2, HuGAL-F2 or control human antibody hIgG. The level of biotinylated GAL-F2 bound was determined by the addition of HRP-streptavidin and substrate. As shown in Fig. 12, HuGAL-F2 and ChGAL-F2 competed approximately equally well, with HuGAL-F2 possibly slightly better, indicating that the binding affinity for FGF2 of HuGAL-F2 is at least as high as ChGAL-F2 and therefore as the original mouse GAL-F2 mAb. From the concentration of HuGAL-F2 required to inhibit binding of the labeled mAb by 50%, one may estimate that the binding affinity K_a of HuGAL-F2 for FGF2 is at least approximately 10^9 M^{-1} . HuGAL-F2 may also be tested in

62451-1087

any of the biological assays for FGF2 activity described herein, and will inhibit FGF2 activity comparably to the GAL-F2 mAb.

5 [0073] Although the invention has been described with reference to the presently preferred embodiments, it should be understood that various modifications can be made without departing from the invention. Unless otherwise apparent from the context any step, element, embodiment, feature or aspect of the invention can be used with any other. If more than one sequence is associated with an accession number at different times, the sequence associated with the accession number as of May 29, 2008 is intended. In the event of any discrepancy between corresponding sequences in the figures and sequences listing, the figures control.

10 [0074] The hybridoma producing the monoclonal antibody GAL-F2, ATCC Number PTA-8864, has been deposited at the American Type Culture Collection, P.O. Box 1549 Manassas, VA 20108, on January 8, 2008 under the Budapest Treaty. This deposit will be maintained at an authorized depository and replaced in the event of mutation, nonviability or destruction for a period of at least five years after the most recent request for release of a sample was received by the depository,
15 for a period of at least thirty years after the date of the deposit, or during the enforceable life of the related patent, whichever period is longest. All restrictions on the availability to the public of these cell lines will be irrevocably removed upon the issuance of a patent from the application.

SEQUENCE LISTING IN ELECTRONIC FORM

20 In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 62451-1087 Seq 18-10-2016 v3.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110> Kim, Kyung Jin
 Wang, Lihong
 Park, Hangil
 Vasquez, Maximiliano
 Galaxy Biotech, LLC

<120> MONOCLONAL ANTIBODIES TO BASIC
 FIBROBLAST GROWTH FACTOR

<130> 022382-000820PC

<140> CA 2725873

<141> 2009-05-28

<150> US 61/057,183

<151> 2008-05-29

<150> US 61/170561

<151> 2009-04-17

<160> 14

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 107

<212> PRT

<213> Mus musculus

<220>

<223> GAL-F2

<400> 1

Ser	Ile	Val	Met	Thr	Gln	Thr	Pro	Lys	Phe	Leu	Leu	Val	Ser	Ala	Gly
1				5					10					15	
Asp	Arg	Val	Thr	Met	Thr	Cys	Lys	Ala	Ser	Gln	Ser	Val	Ser	Ser	Asp
			20					25					30		
Val	Gly	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Ile
		35				40						45			
Tyr	Ser	Gly	Ser	Asn	Arg	Tyr	Ser	Gly	Val	Pro	Asp	Arg	Phe	Thr	Gly
	50					55					60				
Ser	Gly	Tyr	Gly	Thr	Asp	Phe	Thr	Phe	Thr	Ile	Ser	Thr	Val	Gln	Ala
65					70					75				80	
Glu	Asp	Leu	Ala	Val	Tyr	Phe	Cys	Gln	Gln	Asp	Tyr	Tyr	Ser	Pro	Trp
				85					90					95	
Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys					
			100					105							

<210> 2

<211> 107

<212> PRT

<213> Artificial sequence

<220>

<223> HuGAL-F2

<400> 2

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5				10						15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Ala	Ser	Gln	Ser	Val	Ser	Ser	Asp
		20						25					30		
Val	Gly	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
	35						40					45			
Tyr	Ser	Gly	Ser	Asn	Arg	Tyr	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
	50					55					60				
Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
65					70					75					80
Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Asp	Tyr	Tyr	Ser	Pro	Trp
				85					90					95	
Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys					
			100					105							

<210> 3

<211> 107

<212> PRT

<213> Homo sapiens

<220>

<223> immunoglobulin kappa light chain variable region

<400> 3

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5				10						15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Ser	Ile	Ser	Ser	Tyr
		20						25					30		
Leu	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
	35						40					45			
Tyr	Ala	Ala	Ser	Ser	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
	50					55					60				
Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
65					70					75					80
Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Ser	Tyr	Ser	Thr	Pro	Trp
				85					90					95	
Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys					
			100					105							

<210> 4

<211> 121

<212> PRT

<213> Mus musculus

<220>

<223> GAL-F2

<400> 4

Glu	Val	His	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala
1				5				10						15	


```

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
      20      25      30
Val Ile Asn Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
      35      40      45
Gly Tyr Asn Asp Pro Tyr Asn Asp Val Ser Lys Tyr Asn Glu Lys Phe
      50      55      60
Lys Gly Lys Ala Thr Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr
      65      70      75      80
Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
      85      90      95
Ala Lys Glu Gly Gly Gly Lys Tyr Val Tyr Ala Met Asp Ser Trp Gly
      100     105     110
Gln Gly Thr Ser Val Thr Val Ser Ser
      115     120

```

<210> 5
 <211> 121
 <212> PRT
 <213> Artificial sequence

<220>
 <223> HuGAL-F2

```

<400> 5
Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
  1      5      10      15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
      20      25      30
Val Ile Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Ile
      35      40      45
Gly Tyr Asn Asp Pro Tyr Asn Asp Val Ser Lys Tyr Asn Glu Lys Phe
      50      55      60
Lys Gly Arg Ala Thr Ile Thr Ser Asp Lys Ser Thr Ser Thr Ala Tyr
      65      70      75      80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
      85      90      95
Ala Lys Glu Gly Gly Gly Lys Tyr Val Tyr Ala Met Asp Ser Trp Gly
      100     105     110
Gln Gly Thr Thr Val Thr Val Ser Ser
      115     120

```

<210> 6
 <211> 122
 <212> PRT
 <213> Homo sapiens

<220>
 <223> immunoglobulin heavy chain variable region

```

<400> 6
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
  1      5      10      15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe Ser Ser Tyr
      20      25      30

```

Ala Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Gly Ile Ile Pro Ile Phe Gly Thr Ala Asn Tyr Ala Gln Lys Phe
 50 55 60
 Gln Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Val Gly Gln Leu Gly Tyr Tyr Tyr Tyr Gly Met Asp Val Trp
 100 105 110
 Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 115 120

<210> 7

<211> 214

<212> PRT

<213> Artificial sequence

<220>

<223> HuGAL-F2 antibody light chain

<400> 7

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Ser Val Ser Ser Asp
 20 25 30
 Val Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ser Gly Ser Asn Arg Tyr Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Asp Tyr Tyr Ser Pro Trp
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
 100 105 110
 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 115 120 125
 Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140
 Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160
 Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175
 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190
 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205
 Phe Asn Arg Gly Glu Cys
 210

<210> 8

<211> 451

<212> PRT

<213> Artificial sequence

<220>

<223> HuGAL-F2 antibody heavy chain

<400> 8

Glu	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ser
1				5					10					15	
Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asn	Tyr
			20				25						30		
Val	Ile	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile
		35					40					45			
Gly	Tyr	Asn	Asp	Pro	Tyr	Asn	Asp	Val	Ser	Lys	Tyr	Asn	Glu	Lys	Phe
	50					55					60				
Lys	Gly	Arg	Ala	Thr	Ile	Thr	Ser	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr
65					70					75					80
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
				85					90					95	
Ala	Lys	Glu	Gly	Gly	Gly	Lys	Tyr	Val	Tyr	Ala	Met	Asp	Ser	Trp	Gly
			100					105					110		
Gln	Gly	Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser
		115					120						125		
Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala
	130					135					140				
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val
145					150					155					160
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala
				165					170					175	
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val
			180					185					190		
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His
		195					200					205			
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys
	210					215					220				
Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly
225					230					235					240
Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met
				245					250					255	
Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His
			260					265					270		
Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val
		275					280					285			
His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr
	290					295					300				
Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly
305					310					315					320
Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile
				325					330					335	
Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val
			340					345					350		
Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser
		355					360					365			
Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu
	370					375					380				
Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro
385					390					395					400
Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val
				405					410					415	

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
 420 425 430
 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 435 440 445
 Pro Gly Lys
 450

<210> 9
 <211> 11
 <212> PRT
 <213> Artificial sequence

<220>
 <223> GAL-F2 light chain CDR1

<400> 9
 Lys Ala Ser Gln Ser Val Ser Ser Asp Val Gly
 1 5 10

<210> 10
 <211> 7
 <212> PRT
 <213> Artificial sequence

<220>
 <223> GAL-F2 light chain CDR2

<400> 10
 Ser Gly Ser Asn Arg Tyr Ser
 1 5

<210> 11
 <211> 9
 <212> PRT
 <213> Artificial sequence

<220>
 <223> GAL-F2 light chain CDR2

<400> 11
 Gln Gln Asp Tyr Tyr Ser Pro Trp Thr
 1 5

<210> 12
 <211> 5
 <212> PRT
 <213> Artificial sequence

<220>
 <223> GAL-F2 heavy chain CDR1

<400> 12
 Asn Tyr Val Ile Asn
 1 5

<210> 13
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> GAL-F2 heavy chain CDR2

<400> 13
 Asn Asp Pro Tyr Asn Asp Val Ser Lys Tyr Asn Glu Lys Phe Lys Gly
 1 5 10 15

<210> 14
 <211> 12
 <212> PRT
 <213> Artificial sequence

<220>
 <223> GAL-F2 heavy chain CDR3

<400> 14
 Glu Gly Gly Gly Lys Tyr Val Tyr Ala Met Asp Ser
 1 5 10

CLAIMS:

1. A monoclonal antibody (mAb) that binds and neutralizes human basic fibroblast growth factor (FGF2), and competes for binding to FGF2 with an antibody produced by hybridoma PTA-8864 wherein the mAb is chimeric, humanized or human and
5 inhibits growth of a human tumor xenograft that expresses or responds to FGF2 in a mouse.

2. The mAb of claim 1 which is chimeric.

3. The mAb of claim 1 which is humanized.

4. The mAb of claim 1 which is human.

5. The mAb of claim 1 which is a Fab or F(ab')₂ fragment, or single-
10 chain antibody or hybrid bifunctional antibody.

6. An antibody which is a chimeric or humanized form of a monoclonal antibody produced by the PTA-8864 hybridoma.

7. A humanized antibody that binds human basic fibroblast growth factor (FGF2) comprising a humanized light chain comprising three CDRs having the sequences
15 KASQSVSSDVG (SEQ ID NO:9), SGSNRYS (SEQ ID NO:10), and QQDYYSPTW (SEQ ID NO:11) and a humanized heavy chain comprising three CDRs having the sequences NYVIN (SEQ ID NO:12), NDPYNDVSKYNEKFKG (SEQ ID NO:13) and EGGGKYVYAMDS (SEQ ID NO:14).

8. The humanized antibody of claim 7, wherein residues H1, H27, H30, H48, H67, H71 and H94 by Kabat numbering are occupied by E, Y, T, I, A, S and K
20 respectively.

9. The humanized antibody of claim 7 wherein the light chain variable region has at least 95% sequence identity to the sequence shown in SEQ ID NO:2 and the heavy chain has at least 95% sequence identity to the sequence shown in SEQ ID NO:5,
25 wherein any variation from SEQ ID NO:2 or SEQ ID NO:5 occurs in variable region frameworks of the light chain variable region and heavy chain variable region respectively.

62451-1087

10. A monoclonal antibody that binds human basic fibroblast growth factor (FGF2) comprising the three light chain CDRs having the sequences KASQSVSSDVG (SEQ ID NO:9), SGSNRYS (SEQ ID NO:10), and QQDYYSPT (SEQ ID NO:11) and three heavy chain CDRs having the sequences NYVIN (SEQ ID NO:12),
5 NDPYNDVSKYNEKFKG (SEQ ID NO:13) and EGGGKYVYAMDS (SEQ ID NO:14).
11. A monoclonal antibody produced by the hybridoma designated by ATCC deposit number PTA-8864.
12. Hybridoma designated ATCC deposit number PTA-8864.
13. A pharmaceutical composition comprising the antibody of any one of
10 claims 1 to 11 and a physiologically acceptable carrier.
14. Use of the antibody of any one of claims 1 to 11 for treating a cancer that expresses or responds to FGF2.
15. The use of claim 14, wherein the cancer is hepatocellular carcinoma.

1/13




FIGURE 1

2/13




FIGURE 2

3/13




FIGURE 3

4/13





FIGURE 4

5/13






FIGURE 5


6/13



mIgG



GAL-F2



bFM-1

FIGURE 6

7/13




FIGURE 7

8/13




FIGURE 8

9/13




FIGURE 9

10/13




FIGURE 10

11/13

A

	1	2	3	4
	1234567890	1234567890	1234567890	1234567890
GAL-F2	SIVMTQTPKF	LLVSAGDRV	MTCKASQSVS	SDVGWYQQKP
HuGAL-F2	DIQMTQSPSS	LSASVGDRV	ITCKASQSVS	SDVGWYQQKP
ABA70776	DIQMTQSPSS	LSASVGDRV	ITCrasqsis	sylnWYQQKP

	5	6	7	8
	1234567890	1234567890	1234567890	1234567890
GAL-F2	GQSPKLLIYS	GSNRYSGVPD	RFTGSGYGTD	FTFTISTVQA
HuGAL-F2	GKAPKLLIYS	GSNRYSGVPS	RFSGSGSGTD	FTLTISLQ
ABA70776	GKAPKLLIYa	asslqsGVPS	RFSGSGSGTD	FTLTISLQ

	9	10
	1234567890	1234567890
GAL-F2	EDLAVYFCQQ	DYYSPWTFGG
HuGAL-F2	EDFATYYCQQ	DYYSPWTFGQ
ABA70776	EDFATYYCqq	systpwtFGQ

B

	1	2	3	4
	1234567890	1234567890	1234567890	1234567890
GAL-F2	EVHLQQSGPE	LVKPGASVKM	SCKASGYTFT	NYVINWVKQK
HuGAL-F2	EVQLVQSGAE	VKKPGSSVKV	SCKASGYTFT	NYVINWVRQA
AAL04519	QVQLVQSGAE	VKKPGSSVKV	SCKASGGTFS	syaisWVRQA

	5	a	6	7	8
	1234567890	12234567890	1234567890	1234567890	1234567890
GAL-F2	PGQGLEWIGY	NDPYNDVSKYN	EKFKGKATLT	SDKSSSTAYM	
HuGAL-F2	PGQGLEWIGY	NDPYNDVSKYN	EKFKGRATIT	SDKSTSTAYM	
AAL04519	PGQGLEWMGg	iipifgtanya	qkfqqRVTTIT	ADKSTSTAYM	

	abc	9	10ab	cd	11
	1222234567890	123456789000	00	1234567890123	
GAL-F2	ELSSLTSEDSAVY	YCAKEGGGKYVY	AM	DSWGQGTSTVTVSS	
HuGAL-F2	ELSSLRSED TAVY	YCAKEGGGKYVY	AM	DSWGQGT TTVTVSS	
AAL04519	ELSSLRSED TAVY	YCARvgqlgyyyygm		dvWGQGT TTVTVSS	

FIGURE 11

12/13




FIGURE 12

13/13

A

```

1                               D I Q M T Q S P S S
11  L S A S V G D R V T I T C K A S Q S V S S D V G W Y Q Q K P
41  G K A P K L L I Y S G S N R Y S G V P S R F S G S G S G T D
71  F T L T I S S L Q P E D F A T Y Y C Q Q D Y Y S P W T F G Q
101 G T K V E I K R T V A A P S V F I F P P S D E Q L K S G T A
131 S V V C L L N N F Y P R E A K V Q W K V D N A L Q S G N S Q
161 E S V T E Q D S K D S T Y S L S S T L T L S K A D Y E K H K
191 V Y A C E V T H Q G L S S P V T K S F N R G E C

```

B

```

1                               E V Q L V Q S G A E V K
13  K P G S S V K V S C K A S G Y T F T N Y V I N W V R Q A P G
43  Q G L E W I G Y N D P Y N D V S K Y N E K F K G R A T I T S
73  D K S T S T A Y M E L S S L R S E D T A V Y Y C A K E G G G
103 K Y V Y A M D S W G Q G T T V T V S S A S T K G P S V F P L
133 A P S S K S T S G G T A A L G C L V K D Y F P E P V T V S W
163 N S G A L T S G V H T F P A V L Q S S G L Y S L S S V V T V
193 P S S S L G T Q T Y I C N V N H K P S N T K V D K R V E P K
223 S C D K T H T C P P C P A P E L L G G P S V F L F P P K P K
253 D T L M I S R T P E V T C V V V D V S H E D P E V K F N W Y
283 V D G V E V H N A K T K P R E E Q Y N S T Y R V V S V L T V
313 L H Q D W L N G K E Y K C K V S N K A L P A P I E K T I S K
343 A K G Q P R E P Q V Y T L P P S R E E M T K N Q V S L T C L
373 V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P V L
403 D S D G S F F L Y S K L T V D K S R W Q Q G N V F S C S V M
433 H E A L H N H Y T Q K S L S L S P G K

```

FIGURE 13