



US 20080153771A1

(19) **United States**

(12) **Patent Application Publication**

Liu et al.

(10) **Pub. No.: US 2008/0153771 A1**

(43) **Pub. Date: Jun. 26, 2008**

(54) **COMPOSITION AND METHODS OF RNAI THERAPEUTICS FOR TREATMENT OF CANCER AND OTHER NEOVASCULARIZATION DISEASES**

(60) Provisional application No. 60/670,717, filed on Apr. 12, 2005.

Publication Classification

(76) Inventors: **Yijia Liu**, Gaithersburg, MD (US); **Patrick Y. Lu**, Rockville, MD (US); **Martin C. Woodle**, Bethesda, MD (US); **Frank Y. Xie**, Germantown, MD (US)

(51) **Int. Cl.**
A61K 31/70 (2006.01)

(52) **U.S. Cl.** 514/44

(57) **ABSTRACT**

Compositions and methods are provided for treatment of diseases involving unwanted neovascularization (NV). The invention provides treatments that control NV through selective inhibition of pro-angiogenic biochemical pathways, including inhibition of the VEGF pathway gene expression and inhibition localized at pathological NV tissues. Tissue targeted nanoparticle compositions comprising polymer conjugates and nucleic acid molecules that induce RNA interference (RNAi) are provided. The nanoparticle compositions of the invention can be used alone or in combination with other therapeutic agents such as VEGF pathway antagonists. The compositions and methods can be used for the treatment of NV diseases such as cancer, ocular disease, arthritis, and inflammatory diseases.

Correspondence Address:


ROPES & GRAY LLP
PATENT DOCKETING 39/361, 1211 AVENUE OF THE AMERICAS
NEW YORK, NY 10036-8704


(21) Appl. No.: **11/894,567**

(22) Filed: **Aug. 20, 2007**

Related U.S. Application Data

(63) Continuation of application No. PCT/US06/13645, filed on Apr. 12, 2006.





Days After Tumor Cell Inoculation

FIGURE 1




FIGURE 2




FIGURE 3






FIGURE 4



In Vitro



In Vivo




FIGURE 5

HNSCC, Head & Neck




FIGURE 6

Effects of anti-hVEGF siRNAs on VEGF secretion in MCF7/165 cells
(24 hours post-transfection)




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10





FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14




FIGURE 15

VEGF siRNA mix, 4 μ g in 1.6 μ l, Mouse flatmount model




FIGURE 16




FIGURE 17




FIGURE 18





FIGURE 19




FIGURE 20




FIGURE 21




FIGURE 22




FIGURE 23




FIGURE 24




FIGURE 25




FIGURE 26

**COMPOSITION AND METHODS OF RNAI
THERAPEUTICS FOR TREATMENT OF
CANCER AND OTHER
NEOVASCULARIZATION DISEASES**

[0001] This application is a continuation application of International Application No. PCT/US06/13645, filed Apr. 12, 2006, which designates the United States and which claims priority under 35 U.S.C. §119(e) from U.S. Provisional Application No. 60/670,717, filed Apr. 12, 2005, the contents of which are hereby incorporated by reference in their entirety. The application also relates to the following applications, the contents of which are also incorporated by reference in their entireties: International Application No. PCT/US05/03858, filed Feb. 7, 2005; International Application No. PCT/US05/03857, filed Feb. 7, 2005; and International Application No. PCT/US03/24587, filed Aug. 6, 2003.

FIELD OF THE INVENTION

[0002] The invention provides compositions and methods for treatments of diseases with unwanted neovascularization (NV), often an abnormal or excessive proliferation and growth of blood vessels. The development of NV itself often times has adverse consequences or it can be an early pathological step in disease. Despite introduction of new therapeutic antagonists of angiogenesis, including antagonists of the VEGF pathway, treatment options for controlling NV are inadequate and a large and growing unmet clinical need remains for effective treatments of NV, either to inhibit disease progression or to reverse unwanted angiogenesis. Since NV also can be a normal biological process, inhibition of unwanted NV is preferably accomplished with selectivity for a pathological tissue, which preferably requires selective delivery of therapeutic molecules to the pathological tissue.

[0003] The present invention overcomes this hurdle by providing treatments to control NV through selective inhibition of pro-angiogenic biochemical pathways, including inhibition of VEGF pathway gene expression and inhibition localized at pathological NV tissues. The present invention provides compositions and methods for using a tissue targeted nanoparticle composition comprising polymer conjugates and further comprising nucleic acid molecules that induce RNA interference (RNAi). The present invention provides compositions and methods for inhibition of individual genes or combinations of genes active in NV and more preferably in the VEGF pathway. The dsRNA nanoparticle compositions of the invention can be used alone or in combination with other therapeutic agents, including targeted therapeutics, including VEGF pathway antagonists, such as monoclonal antibodies and small molecule inhibitors, and targeted therapeutics inhibiting EGF and its receptor or PDGF and its receptors or MEK or Bcr-Abl, and immunotherapy and chemotherapy. The present invention also provides compositions and methods for the treatment of NV disease in a subject, including cancer, ocular disease, arthritis, and inflammatory diseases.

BACKGROUND OF THE INVENTION

[0004] Recent US FDA approved therapeutic agents, including Avastin, and Macugen, provide some benefit for NV diseases. Some of these agents act by binding to and inhibiting the action of Vascular Endothelial Growth Factor

(VEGF), but these agents are not effective for many patients. Other agents being evaluated in clinical studies show signs that they may provide some benefit by binding to and inhibiting the action of the receptors for VEGF, or “down stream” proteins used by these receptors for signal transduction. The picture that has emerged is that means to control this VEGF “pathway” can provide a level of control of NV that provides benefit for some patients. In addition, studies of a series of small molecule kinase inhibitors found that an inhibitor called sunitinib that has activity against multiple kinase proteins, VEGF receptor, PDGF receptor, FLT3, and Kit, offers better clinical benefit for NV diseases. However, these benefits are still inadequate for most patients and better therapeutic means to control the VEGF pathway still are needed. The agents developed to date are mostly antagonists of VEGF or its receptors, VEGF R1 and VEGF R2. One problem that has emerged with use of antagonists appears to be a response by the pathological tissues to increase production of VEGF. Thus an attractive means to improve therapeutic control of NV is to inhibit production of the VEGF pathway proteins, i.e., down regulate their gene expression, and doing so by inducing RNA interference through in vivo delivery of small interfering dsRNA oligonucleotides (siRNA).

[0005] RNA interference (RNAi) is a post-transcriptional process where a double stranded RNA inhibits gene expression in a sequence specific fashion. The RNAi process occurs in at least two steps: During one step, a long dsRNA is cleaved by an endogenous ribonuclease into shorter, 21- or 23-nucleotide-long dsRNAs. In another second step, the smaller dsRNA mediates the degradation of an mRNA molecule with a matching sequence and as a result selectively down regulating expression of that gene. This RNAi effect can be achieved by introduction of either longer double-stranded RNA (dsRNA) or shorter small interfering RNA (siRNA) to the target sequence within cells. Recently, it was demonstrated that RNAi can also be achieved by introducing of plasmid that generate dsRNA complementary to target gene.

[0006] RNAi methods have been successfully used in gene function determination experiments in *Drosophila*^(20,22,23,25), *C. elegans*^(14,15,16), and *Zebrafish*⁽²⁰⁾. In those model organisms, it has been reported that both the chemically synthesized shorter siRNA or in vitro transcribed longer dsRNA can effectively inhibit target gene expression. Methods have been reported that successfully achieved RNAi effects in non human mammalian and human cell cultures.⁽³⁹⁻⁵⁶⁾ However, RNAi effects have been difficult to observe in adult animal models⁽⁵⁷⁾. This is for several reasons including: introduction of a long double-stranded RNA into mammalian cells can trigger an antiviral immune response including up-regulation of interferon, resulting in apoptosis and death of the cells that can be either detrimental or beneficial to the desired therapeutic effect: the efficiency of dsRNA entry into the target cell is low, especially in animals; short dsRNA molecules are rapidly excreted from the blood into the urine; and RNA molecules can be degraded by RNase nuclease activity. Although RNAi has potential applications in both gene target validation and nucleic acid therapeutics, progress of the technology has been hindered due to the poor delivery of RNAi molecules into animal disease models.

[0007] It is apparent, therefore, that improved methods for delivering RNAi molecules in vivo are of great importance. It is also apparent that tissue targeted delivery of nucleic acid molecules inducing RNAi are of great importance. It is also apparent that methods for delivering nucleic acid molecules

inducing RNAi selective for VEGF pathway genes will be of great benefit for the treatment of NV diseases. These needs are addressed by the compositions and methods of the invention.

SUMMARY OF THE INVENTION

[0008] It is therefore an object of the present invention to utilize RNAi to modulate angiogenesis process in order to reverse the disease process by down regulating gene expression involved in NV pathogenesis, more specifically genes in the VEGF pathway.

[0009] It is therefore an object of the invention to provide compositions and methods for inhibiting expression of one or more VEGF pathway genes in a mammal. It is a further object of the invention to provide compositions and methods for treating NV disease by inhibiting expression of one or more VEGF pathway genes alone or in combination with other agents including antagonists of the same VEGF pathway.

[0010] In achieving these objects there has been provided a compositions and method for down regulating endogenous VEGF pathway genes, comprising administering to a tissue of the mammal a composition comprising a double-stranded RNA molecule where the RNA molecule specifically reduces or inhibits expression of the endogenous VEGF pathway gene. This down regulation of an endogenous gene may be used for treating a disease that is caused or exacerbated by activity of the VEGF pathway. The disease may be in a human.

[0011] There also has been provided a method for treating a disease in a mammal associated with undesirable expression of a VEGF pathway gene, comprising applying a nucleic acid composition comprising a dsRNA oligonucleotide, as the active pharmaceutical ingredient (API), associated with a formulation, wherein the formulation can comprise a polymer, where the nucleic acid composition is capable of reducing expression of the VEGF pathway genes and inhibiting NV in the disease. The disease may be cancer or a precancerous growth and the tissue may be, for example, a kidney tissue, breast tissue, colon tissue, a prostate tissue, a lung tissue or an ovarian tissue.

[0012] As used herein, "oligonucleotides" and similar terms based on this relate to short oligos composed of naturally occurring nucleotides as well as to oligos composed of synthetic or modified nucleotides. Oligonucleotides may be 10 or more nucleotides in length, or 15, or 16, or 17, or 18, or 19, or 20 or more nucleotides in length, or 21, or 22, or 23, or 24 or more nucleotides in length, or 25, or 26, or 27, or 28 or 29, or 30 or more nucleotides in length, 35 or more, 40 or more, 45 or more, up to about 50, nucleotides in length.

[0013] An oligonucleotide that is an siRNA may have any number of nucleotides between 15 and 30 nucleotides. In many embodiments an siRNA may have any number of nucleotides between 19 and 27 nucleotides.

[0014] In many embodiments, an siRNA may have two blunt ends, or two sticky ends, or one blunt end with one sticky end. The over hang nucleotides of a sticky end can range from one to four nucleotides or more.

[0015] In a preferred embodiment, the invention provides siRNA of 25 base pairs with blunt ends.

[0016] The terms "polynucleotide" and "oligonucleotide" are used synonymously herein.

[0017] The composition may further comprise a polymeric carrier. The polymeric carrier may comprise a cationic polymer that binds to the RNA molecule and forms nanoparticles.

The cationic polymer may be an amino acid copolymer, containing, for example, histidine and lysine residues. The polymer may comprise a branched polymer. The composition may comprise a targeted synthetic vector. The synthetic vector may comprise a cationic polymer as a nucleic acid carrier, a hydrophilic polymer as a steric protective material, and a targeting ligand as a target cell selective agent. The polymer may comprise a polyethyleneimine or a polyhistidine-lysine copolymer or a polylysine modified chemically or other effective polycationic carriers that can be used as the nucleic acid carrier module, the hydrophilic polymer may comprise a polyethylene glycol or a polyacetal or a polyoxazoline, and the targeting ligand may comprise a peptide comprising an RGD sequence or a sugar or a sugar analogue or an mAb or a fragment of an mAb, or any other effective targeting moieties.

[0018] In any of these methods, an electric field may be applied to a tissue substantially contemporaneously with the composition or subsequent to application of the composition. The composition and method of the invention comprises dsRNA oligonucleotides with a sequence matching an endogenous VEGF pathway gene or a mutated endogenous gene, and at least one mutation in the mutated gene may be in a coding or regulatory region of the gene. In any of these methods, the endogenous gene may be selected from the group consisting of VEGF pathway genes including growth factor genes, protein serine/threonine kinase genes, protein tyrosine kinase genes, protein serine/threonine phosphatase genes, protein tyrosine phosphatase genes, receptor genes, and transcription factor genes. The selected gene may include one or more genes from the group consisting of VEGF, VEGF-R1, VEGF-R2, VEGF-R3, VEGF121, VEGF165, VEGF 189, VEGF206, RAF-a, RAF-c, AKT, Ras, NF-Kb. The selected gene may include one or more genes from other biochemical pathways associated with NV including HIF, EGF, EGFr, bFGF, bFGFr, PDGF, and PDGFr. The selected gene may include one or more genes from other biochemical pathways operative in concert with NV including Her-2, c-Met, c-Myc and HGF.

[0019] The present invention also provides compositions and methods for nucleic acid agents inducing RNAi to inhibit multiple genes including cocktails of siRNA (siRNA-OC). The compositions and methods of the invention may inhibit multiple genes substantially contemporaneously or they may inhibit multiple genes sequentially. In a preferred embodiment siRNA-OC agents inhibit three VEGF pathway genes: VEGF, VEGF receptor 1, and VEGF receptor 2. In another preferred embodiment siRNA-OC are administered substantially contemporaneously. The present invention provides agents with gene inhibition selectivity derived from matching the sequence of the siRNA largely to a sequence in the targeted gene mRNA. It also provides siRNA agents with substantially similar physiochemical properties that inhibit different genes in the VEGF pathway. It also provides nanoparticle compositions largely independent of the siRNA sequence. It also provides methods for treatment of human diseases, especially NV related diseases, which can be treated with inhibitors of multiple endogenous genes. It also provides methods for treatment of human diseases by combinations of therapeutic agents administered substantially contemporaneously in some cases and, in other cases, sequentially.

[0020] One aspect of the present invention provides compositions and methods for treatment of cancer, arthritis, blindness, infectious diseases and inflammatory diseases. In another aspect of the present invention nucleic acid agents

inducing RNAi are used in concert with other therapeutic agents, such as but not limited to small molecules and monoclonal antibodies (mAb), in the same therapeutic regimen.

[0021] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] FIG. 1 shows the inhibition of DLD-1 colon xenograft tumor growth by VEGF pathway siRNA nanoparticle compositions targeted by an RGD peptide ligand alone or in combination with Avastin treatment. Stronger anti-tumor efficacy has been observed using a combination cancer therapeutics with VEGFR2-siRNA inhibitor and Avastin (Bevacizumab) in the same treatment regimen, in a human colon carcinoma (DLD-1) xenograft tumor model.

[0023] FIG. 2. VEGFR2-siRNA nanoparticle targeting neovasculature through systemic delivery demonstrated potent efficacy in colon carcinoma xenograft mouse model (DLD-1 tumor) after four repeated administrations every three days. RT-PCR results illustrate the knockdown of VEGFR2 gene expression at mRNA level.

[0024] FIG. 3. Human VEGFR2-siRNA nanoparticle targeting neovasculature through systemic delivery demonstrated potent efficacy in colon carcinoma xenograft mouse model (DLD-1 tumor) after four repeated administrations every three days. Immunohistochemistry images show down regulation of VEGFR2 and CD31 expression in the tumor tissue.

[0025] FIG. 4. While human VEGFR2-siRNA nanoparticle targeting neovasculature through systemic delivery demonstrated potent efficacy in colon carcinoma xenograft mouse model (DLD-1 tumor), no significant IFN- α induction was observed in the mouse blood stream 24 hours after the 2nd delivery.

[0026] FIG. 5. Human VEGF-siRNA (19 base pair with 3'end overhangs) is effective for silencing hVEGF expression in vitro and in vivo, at both mRNA level and protein level, evaluated by RT-PCR and ELISA. Two upper figures demonstrated significant hVEGF knockdown at both mRNA and protein levels. The lower panel shows the anti-tumor activity of siRNA-mediated hVEGF knockdown in a MCF-7/VEGF165 cell induced xenograft model, where the hVEGF expression was significantly down regulated in the tumor tissue.

[0027] FIG. 6. Human VEGF-siRNA (19 base pair with 3'end overhangs) is effective for silencing hVEGF expression in vivo resulting in tumor growth inhibition (HNSCC 1483 cell). Five times repeated intratumoral delivery with electroporation enhancement with 7 day interval.

[0028] FIG. 7. Human VEGF-siRNA (25 basepair blunt end) is more effective for silencing hVEGF expression in vitro and stronger than the 19 base pair siRNA.

[0029] FIG. 8. Human VEGF-siRNA (25 basepair blunt end) is effective for silencing hVEGF expression in vitro and has longer duration than that of the 19 basepair siRNA.

[0030] FIG. 9. Human VEGFR1-siRNA (25 base pair blunt end) is effective for silencing hVEGFR1 expression in vitro after 48 hr.

[0031] FIG. 10. Human VEGFR1-siRNA (25 base pair blunt end) is effective for silencing hVEGFR1 expression in vitro after 72 hr.

[0032] FIG. 11. Human VEGFR2-siRNA (25 base pair blunt end) is effective for silencing hVEGFR2 expression in vitro at two different time points.

[0033] FIG. 12. Human VEGF-siRNA (25 base pair blunt end) is effective for silencing hVEGF expression in vivo resulting in tumor growth inhibition, with a MDA-MB-435 breast carcinoma cell line.

[0034] FIG. 13. Concept framework for using siRNA cocktail targeting VEGF, VEGFR1 and VEGFR2 in the same treatment to block the angiogenesis pathway.

[0035] FIG. 14. siRNA cocktail targeting VEGF, VEGFR1 and VEGFR2 is more effective than any single siRNA targeting individual gene at the same dosage tested in the HSV induced ocular angiogenesis mouse model, with either local (left) or systemic (right) deliveries.

[0036] FIG. 15. Inhibition of angiogenesis by administration of VEGF pathway siRNA cocktail on day 17 or day 14 after induction of disease in ocular tissues of a murine model of retinal angiogenesis in a perfusion/flatmount measurement.

[0037] FIG. 16. siRNA cocktail targeting VEGF, VEGFR1 and VEGFR2 demonstrates very effective anti-angiogenesis activity in the retinopathy of prematurity (ROP) mouse model induced by Hypoxia.

[0038] FIG. 17. siRNA cocktail targeting VEGF₅, VEGFR1 and VEGFR2 demonstrates very effective anti-angiogenesis activity in the retinopathy of prematurity (ROP) mouse model induced by Hypoxia.

[0039] FIG. 18. siRNA cocktail targeting VEGF, VEGFR1 and VEGFR2 inhibits ocular angiogenesis significantly with different routes of administration with different carriers. IP delivery was mediated by a ligand-directed nanoparticle. siRNA Cocktail Mediated Knockdown of VEGF Pathway Factors RT-PCR Detection of mRNA levels with ROP Model.

[0040] FIG. 19. Self-assembled nanoparticle for siRNA delivery. When the pre-made RGD-PEG-PEI conjugate aqueous solution is mixed with the siRNA aqueous solution, the nanoparticles will be self-assembled as described. The nanoparticles are relatively homogeneous in size, from 50 nm to 100 nm.

[0041] FIG. 20. siRNA nanoparticle targeting neovasculature demonstrated tumor targeting property using labeled siRNA payload (left) and plasmid payload expressing Luciferase reporter gene (right).

[0042] FIG. 21. VEGFR2-siRNA nanoparticle targeting neovasculature demonstrated potent efficacy in neuroblastoma syngenic mouse model (N2A tumor) after four repeated administrations every three days.

[0043] FIG. 22. VEGFR2-siRNA nanoparticle targeting neovasculature demonstrated potent efficacy in renal carcinoma xenograft mouse model (786-O tumor) after five repeated administrations every three days.

[0044] FIG. 23. VEGFR2-siRNA nanoparticle targeting neovasculature demonstrated potent efficacy in colon carcinoma xenograft mouse model (DLD-1 tumor) after four repeated administrations every three days. Three different dosages were used in the study.

[0045] FIG. 24. VEGFR2-siRNA nanoparticle targeting neovasculature demonstrated potent efficacy in colon carcinoma xenograft mouse model (DLD-1 tumor) after four repeated administrations every three days. Four mg per kilo of VEGFR2 dosing resulted in the same anti-tumor efficacy as 5 mg per kilogram of Avastin.

[0046] FIG. 25. Manufacturing flow chart for production of an RGD-polymer conjugate and a nanoparticle-siRNA preparation starting from raw materials of a homobifunctional activated PEG, an RGD-2C peptide, a polyamine containing agent, and a mixture of dsRNA.

[0047] FIG. 26. The use of a static mixer to manufacture a nanoparticle-siRNA preparation starting from two solutions, a polymer solution comprising polymer conjugates and an siRNA solution comprising three dsRNA agents.

DETAILED DESCRIPTION

[0048] The present invention provides compositions and methods for treatment of NV diseases, which typically are characterized by attributes of multiple proteins and abnormally over-expressed disease-causing genes and multiple malfunctions of disease-causing proteins. The present invention provides nucleic acid agents, such as siRNA oligonucleotides, that activate RNA interference (RNAi) and are highly selective inhibitors of gene expression with a sequence specific manner. The present invention provides inhibition of NV by modulation of protein activity including reduction of protein expression levels and post transcriptional modification of proteins.

[0049] In cancer, the tumorigenesis process is thought to be the result of abnormal over-expression of oncogenes, angiogenesis factors, growth factors, and mutant tumor suppressors, even though under-expression of other proteins also plays a critical role. Increasing evidences supports the notion that siRNA molecules are able to “knockdown” tumorigenic genes both in vitro and in vivo, resulting in significant anti-tumor effects. The compositions and methods of the present invention demonstrate substantial knockdown of human VEGF in MCF-7 cells, MDA-MB-435 cells and 1483 cell-induced xenograft tumor models, achieving tumor growth inhibition of 40-80%, using intratumoral delivery of siRNA specifically targeting human VEGF pathway gene sequences. It is appreciated that inhibition of VEGF pathway gene expression induces anti-angiogenesis effects that alter the microvasculature in tumors and that activate tumor cell apoptosis and can enhance efficacy of cytotoxic chemotherapeutic drugs. However, to achieve significantly improved anti-tumor efficacy of anti-angiogenesis agents and chemotherapeutic drugs, a highly effective delivery method is necessary so that elevated concentrations of the drugs accumulate in the local tumor tissue, and in many instances through a systemic administration.

[0050] The present inventors have described a method of validating drug targets that determines which targets control disease pathways and therefore justify drug development efforts (see PCT/US02/31554). The present inventors also have described technologies suitable for delivery of nucleic acids into animal tissues. See WO01/47496, the contents of which are hereby incorporated by reference in their entirety. These methods enable administration of nucleic acid and achieve a significant (for example, seven-fold) increase in efficiency compared to “gold standard” nucleotide delivery reagents. Accordingly, the methods provide strong activity of

nucleic acids in tissues including activity of candidate target proteins. This platform is a powerful tool for validation of candidate genes in a tissue.

[0051] In addition, the present inventors have used these methods to achieve gene silencing in animal tissues, which is highly desired for validation of candidate target genes and as a therapeutic modality. Recently, double stranded RNA has been demonstrated to induce gene-specific silencing by a phenomenon called RNA interference (RNAi). Although the mechanism of RNAi still is not completely understood, early results suggested that the RNAi effect may be achieved in vitro in various cell types including mammalian cells. A double stranded RNA targeted against a target mRNA results in the degradation of the target, thereby causing the silencing of the corresponding gene. Large double stranded RNA is cleaved into smaller fragments of 21-23 nucleotides long by an RNase III like activity involving an enzyme Dicer. These shorter fragments, known as siRNA (small interfering RNA), are believed to mediate the cleavage of mRNA.

[0052] Although gene down-regulation by the RNAi mechanism has been studied in *C. elegans* and other lower organisms in recent past, its effectiveness in mammalian cells in culture has only recently been demonstrated. An RNAi effect recently was demonstrated in mouse using the firefly luciferase gene reporter system. To develop an RNAi technology platform for in vivo gene inhibition for research and for clinical application of nucleic acid therapeutics to treat human diseases, the present inventors performed several in vivo studies in mouse models of disease. In those experiments, either siRNA or dsRNA targeting a tumor related ligand (human VEGF) or receptor (mouse VEGFR2) was delivered to nude mice bearing xenografted human MCF-7 derived tumor or human MDA-MB-435 tumors. For the first time we were able to demonstrate that RNAi can effectively silencing target gene in tumor cells in vivo and that, as a result, tumor growth was inhibited.

[0053] The present inventors have achieved for the first time therapeutic compositions and methods for treatment of a wide variety of NV diseases using dsRNA oligonucleotides inhibiting VEGF pathway genes. The invention is described here in detail, but one skilled in the art will appreciate the full extent of the invention.

[0054] A. Potent siRNA for VEGF Pathway Gene Inhibition

[0055] The present invention provides nucleic acid agents targeting and inhibiting VEGF pathway gene sequences with a variety of physicochemical structures. One preferred embodiment of the present invention uses nucleic acid agents called siRNA including 19 base pair dsRNA with 3' overhangs and 25 base pair dsRNA with blunt ends. The siRNA of the invention with 25 base pair dsRNA with blunt ends were found to be some of the most potent inhibitors with some of the greatest duration of inhibition. Additionally, incorporation of non-naturally occurring chemical analogues are useful in the invention, including 2'-O-Methyl ribose analogues of RNA, DNA and RNA chimeric oligonucleotides, and other chemical analogues of nucleic acid oligonucleotides. The 25 basepair siRNA with and without 2'-O-Methyl ribose in the sense strand of the siRNA and targeting human VEGF sequence provide strong and durable inhibitory effect, up to 70-80% in MCF-7/VEGF165 cells and in tumors growing in animals. This inhibitory effect in MCF-7/VEGF165 cell culture lasted for 5 days. One aspect provided for by the invention is siRNA targeting human genes and the encoded

sequence also targets other mammalian species such as other primates, mice, and rats but not limited to these species. Methods well known to one skilled in the art can be used to identify and select siRNA provided by the invention. Many other forms of siRNA targeting VEGF pathway genes are provided for by the invention and others will be understood by one skilled in the art.

a. Human VEGF Specific siRNA:

25 base pair blunt ends:

hVEGF-25-siRNA-a:

Sense strand:

5'-r(CCUGAUGAGAUCGAGUACAUCUUA)-3' (SEQ ID NO: 1)

Antisense strand:

5'-r(UGAAGAUGUACUCGAUCUCAUCAGG)-3' (SEQ ID NO: 2)

hVEGF-25-siRNA-b:

Sense strand:

5'-r(GAGAGAUGAGCUUCCUACAGCACC)-3' (SEQ ID NO: 3)

Antisense strand:

5'-r(UUGUGCUGUAGGAAGCUCAUCUCUC)-3' (SEQ ID NO: 4)

hVEGF-25-siRNA-c:

Sense strand:

5'-r(CACAACAAUGUGAAUGCAGACCAA)-3' (SEQ ID NO: 5)

Antisense strand:

5'-r(UUGGUCUGCAUUCACAAUUGUUGUG)-3' (SEQ ID NO: 6)

19 base pairs with two nucleotide

(TT) overhangs

at 3':

hVEGF165

5'-r(UCGAGACCCUGGUGGACAUTT)-3' (SEQ ID NO: 7)

b. Human VEGF receptor 1 specific siRNA:

25 base pair blunt ends:

hVEGFR1-25-siRNA-a,

Sense strand:

5'-r(GCCAACAUAUUCUACAGUGUUCUUA)-3' (SEQ ID NO: 8)

Antisense strand:

5'-r(UAAGAACACUGUAGAUAUUGUUGGC)-3' (SEQ ID NO: 9)

hVEGFR1-25-siRNA-b,

Sense strand:

5'-r(CCCUCGCGGGAAGUUGUAUGGUUAA)-3' (SEQ ID NO: 10)

Antisense strand:

5'-r(UUAACCAUACAACUCCGCGGAGGG)-3' (SEQ ID NO: 11)

19 basepairs with 2 3' (TT) nucleotide overhangs:

VEGF RI (FLT)

5'-GGAGAGGACCUGAAACUGUTT (SEQ ID NO: 12)

c. Human VEGF receptor 2 specific siRNA:

25 basepair blunt ends:

hVEGFR2-25-siRNA-a,

Sense strand:

5'-r(CCUCUCUGUAAGACACUCACAAUU)-3' (SEQ ID NO: 13)

-continued

Antisense strand:

5'-r(AAUUGUGAGUGUCUUACAGAGG)-3' (SEQ ID NO: 14)

hVEGFR2-25-siRNA-b,

Sense strand:

5'-r(CCCUUGAGUCCAAUCACACAAUUA)-3' (SEQ ID NO: 15)

Antisense strand:

5'-r(UUAAUUGUGUAUUGGACUCAGGG)-3' (SEQ ID NO: 16)

hVEGFR2-25-siRNA-c,

Sense strand:

5'-r(CCAAGUAUUGAAGCAGAUGCCUUU)-3' (SEQ ID NO: 17)

Antisense strand:

5'-r(AAAGGCAUCUGCUUCAUUCACUUGG)-3' (SEQ ID NO: 18)

19 basepairs with 2 3' (TT)

nucleotide overhangs:

hVEGF R2 (KDR)

5'-CAGUAAGCGAAAGAGCCGGTT-3' (SEQ ID NO: 19)

[0056] In another embodiment, the siRNA targeting human VEGF receptor 1 has the following sequence:

Sense strand:

5'-r(CCUCAAGAGCAAACGUGACUUUUU)-3' (SEQ ID NO: 20)

Antisense strand:

5'-r(AAAUAAGUCAGUUUGUCUUGAGG)-3' (SEQ ID NO: 21)

25 Base Pair siRNA Oligos can Target the Corresponding Genes from Both Human and Mouse.

[0057] In another embodiment, the 25 base pair siRNA sequences were selected to against the corresponding mRNA sequences, including human VEGF, VEGFR1, VEGFR2, PDGFR-alpha, PDGFR-beta and EGFR genes. The sequences selected are not only specific to the human genes but also specific to the corresponding mouse genes, such as the sequences against human VEGF mRNAs are also against mouse VEGF mRNAs.

[0058] The 25 base pair siRNA oligos are very useful for mouse xenograft tumor study since the anti-tumor efficacy relies on knockdown both human and mouse genes in the tumor tissue. For example, knockdown VEGF expressions from both tumor cells (human origin) and endothelium cells (mouse origin) with the siRNA oligos targeting both sequences will provide better anti-tumor efficacy.

[0059] Meanwhile, the toxicity study with the siRNA oligos targeting both genes from testing animals (e.g. mouse or monkey) and human genes will be very useful since the actual drugs were tested for both drug agent toxicology and exaggerated pharmacology.

[0060] The following sequences are the 25 base pair siRNA oligo sequences targeting VEGF, VEGFR2, VEGFR1, PDGFR-alpha, PDGFR-beta and EGFR mRNAs from both human and mouse genes, some of them can also target to their monkey and dog counterparts.

25 Base Pair VEGF siRNA Targeting Human, Mouse, Rat, Macaque, Dog VEGF mRNA Sequences:

mhVEGF25-1: (SEQ ID NO. 22)
sense, 5'-CAAGAUCGCGACGUGUAAAUGUU-3';
antisense, 5'-AAC AUUACACGUCUGCGGAUCUUG-3'

mhVEGF25-2: (SEQ ID NO. 24)
sense, 5'-GCAGCUUGAGUUAACGAACGUACU-3';
antisense, 5'-AGUACGUUCGUUUAACUCAAGCUGC-3'

mhVEGF25-3: (SEQ ID NO. 26)
sense, 5'-CAGCUUGAGUUAACGAACGUACUU-3';
antisense, 5'-AAGUACGUUCGUUUAACUCAAGCUG-3'

mhVEGF25-4: (SEQ ID NO. 28)
sense, 5'-CCAUGCACAAGUGGUCCAGGCUGCA-3';
antisense, 5'-TGCAGCCTGGGACCACTTGGCATGG-3'

mhVEGF25-4: (SEQ ID NO. 30)
sense, 5'-CACAUAGGAGAGAGAGCUUCCUCA-3';
antisense, 5'-UGAGGAAGCUAUCUCUCCUAUGUG-3'

25 Base Pair VEGF R2 siRNA Sequences Targeting Both Human and Mouse VEGFR2 mRNA Sequences:

mhVEGFR225-1: (SEQ ID NO. 32)
sense, 5'-CCUACGGACCGUUAAGCGGGCCAUA-3';
antisense, 5'-AUUGCCCGCUUAACGGUCCGUAGG-3'

mhVEGFR225-2: (SEQ ID NO. 34)
sense, 5'-CUCAUGUCUGUUCUCAAGAUCUCA-3';
antisense, 5'-UGAGGAUCUUGAGAACAGACAUGAG-3'

mhVEGFR225-3: (SEQ ID NO. 36)
sense, 5'-CUCAUGGUGAUUGUGGAAUUCUGCA-3';
antisense, 5'-UGCAGAAUCCACAAUCACCAUGAG-3'

mhVEGFR225-4: (SEQ ID NO. 38)
sense, 5'-GAGCAUGGAAGAGGAUUCUGGACUC-3';
antisense, 5'-GAGUCCAGAAUCCUUCUCCAUAGTC-3'

mhVEGFR225-5: (SEQ ID NO. 40)
sense, 5'-CAGAACAGUAAGCGAAAGAGCCGGC-3';
antisense, 5'-GCCGCGUCUUUCGUUACUGUUCUG-3'

-continued

mhVEGFR225-6: (SEQ ID NO. 42)
sense, 5'-GACUUCUGACCUUGGAGCAUCUCA-3';
antisense, 5'-UGAGAUGCUCCAAGGUCAGGAAGUC-3'

mhVEGFR225-7: (SEQ ID NO. 44)
sense, 5'-CCUGACCUUGGAGCAUCUCAUCUGU-3';
antisense, 5'-ACAGAUGAGAUGCUCCAAGGUCAGG-3'

mhVEGFR225-8: (SEQ ID NO. 46)
sense, 5'-GCUAAGGGCAUGGAGUUCUUGGCAU-3';
antisense, 5'-AUGCCAAGAACUCCAUUGCCUUAGC-3'

25 Base Pairs VEGFR1 siRNA Sequences Targeting Both Human and Mouse VEGFR1 mRNA Sequences:

mhVEGFR125-1: (SEQ ID NO. 48)
sense, 5'-CACGCUGUUUAUUGAAAGAGUCACA-3';
antisense, 5'-UGUGACUCUUUCAUUAACAGCGUG-3'

mhVEGFR125-2: (SEQ ID NO. 50)
sense, 5'-CGCUGUUUAUUGAAAGAGUCACAGA-3';
antisense, 5'-UCUGUGACUCUUUCAUUAACAGCG-3'

mhVEGFR125-3: (SEQ ID NO. 52)
sense, 5'-CAAGGAGGGCCUCUGAUGGUGAUGU-3';
antisense, 5'-ACAUCACCAUCAGAGGCCUCCUUG-3'

mhVEGFR125-4: (SEQ ID NO. 54)
sense, 5'-CCAACUACCUCAAGAGCAAACGUGA-3';
antisense, 5'-UCACGUUUGCUCUUGAGGUAGUUGG-3'

mhVEGFR125-5: (SEQ ID NO. 56)
sense, 5'-CUACCUCAAGAGCAAACGUGACUUA-3';
antisense, 5'-UAAGUCACGUUUGCUCUUGAGGUAG-3'

mhVEGFR125-6: (SEQ ID NO. 58)
sense, 5'-CCAGAAAGUGCAUUCUACGGGACCU-3';
antisense, 5'-AGGUCCCGAUGAAUGCACUUCUGG-3'

mhVEGFR125-7: (SEQ ID NO. 60)
sense, 5'-CAUUCUACGGGACCUUGGAGCGAGA-3';
antisense, 5'-UCUCGUGCCAGGUCCGAUGAAUG-3'

mhVEGFR125-8:

-continued

(SEQ ID NO. 62)
sense, 5'-CAUCGGGACCUGGCAGCGAGAAACA-3';

(SEQ ID NO. 63)
antisense: 5'-UGUUUCUCGCGCCAGGUCCGAUG-3'

mhVEGFR125-9:
(SEQ ID NO. 64)
sense, 5'-GAGCCUGGAAAGAAUAAAACCUUU-3';

(SEQ ID NO. 65)
antisense: 5'-AAAGGUUUUGAUUCUUUCCAGGCUC-3'

mhVEGFR125-10:
(SEQ ID NO. 66)
sense, 5'-GCCUGGAAAGAAUAAAACCUUUGA-3';

(SEQ ID NO. 67)
antisense: 5'-UCAAGGUUUUGAUUCUUUCCAGGC-3'

mhVEGFR125-11:
(SEQ ID NO. 68)
sense, 5'-GCCUGGAAAGAAUAAAACCUUUGA-3';

(SEQ ID NO. 69)
antisense: 5'-UCAAGGUUUUGAUUCUUUCCAGGC-3'

mhVEGFR125-12:
(SEQ ID NO. 70)
sense, 5'-CUGAACUGAGUUUAAAAGGCACCCA-3';

(SEQ ID NO. 71)
antisense: 5'-UGGGUGCCUUUAAAACUGAGUUUCAG-3'

mhVEGFR125-13:
(SEQ ID NO. 72)
sense, 5'-GAACUGAGUUUAAAAGGCACCCAGC-3';

(SEQ ID NO. 73)
antisense: 5'-GCUGGGUGCCUUUAAAACUCAGUUG-3'

25 Base Pairs siRNA Sequences Targeting Both Human and Mouse PDGFR-Alpha:

mhPDGFRa25-1:
(SEQ ID NO. 74)
sense, 5'-GAAGAUAAUGACUCACCCUGGGGCCA-3';

(SEQ ID NO. 75)
antisense: 5'-UGGCCCCAGGUGAGUCAUUUUCUUC-3'

mhPDGFRa25-2:
(SEQ ID NO. 76)
sense, 5'-GAUAAUGACUCACCCUGGGGCCACAU-3';

(SEQ ID NO. 77)
antisense: 5'-AUGUGGGCCCCAGGUGAGUCAUUUUC-3'

mhPDGFRa25-3:
(SEQ ID NO. 78)
sense, 5'-CUCACCCUGGGGCCACAUUUGAACAU-3';

(SEQ ID NO. 79)
antisense: 5'-AUGUUCAAAUGUGGGCCCCAGGUGAG-3'

mhPDGFRa25-4:
(SEQ ID NO. 80)
sense, 5'-CUUGCUGGGAGCCUGCACCAAGUCA-3';

(SEQ ID NO. 81)
antisense: 5'-UGACUUGGUGCAGGCUCACCAAGCAAG-3'

mhPDGFRa25-5:
(SEQ ID NO. 82)
sense, 5'-GAUUCUACUUUCUACAUAAGAUA-3';

-continued

(SEQ ID NO. 83)
antisense: 5'-UGAUCUUUUGUAGAAAGUAGAAUC-3'

mhPDGFRa25-6:
(SEQ ID NO. 84)
sense, 5'-CAGAGACUGAGCGCUGACAGUGGCU-3';

(SEQ ID NO. 85)
antisense: 5'-AGCCACUGUCUGCGCUCAGUCUCUG-3'

mhPDGFRa25-7:
(SEQ ID NO. 86)
sense, 5'-GACCUGGGCAAGAGGAACAGACACA-3';

(SEQ ID NO. 87)
antisense: 5'-UGUGUCUGUUCUUCUUGCCAGGUC-3'

mhPDGFRa25-8:
(SEQ ID NO. 88)
sense, 5'-CCACCUUCAUCAAGAGAGAGGACGA-3';

(SEQ ID NO. 89)
antisense: 5'-UCGUCCUCUCUCUUGAUGAAGGUGG-3'

mhPDGFRa25-9:
(SEQ ID NO. 90)
sense, 5'-UAUGGAUUAAGCCGGUCCCAACCUGU-3';

(SEQ ID NO. 91)
antisense: 5'-ACAGGUUGGGACCCGCUUAAUCCAUA-3'

25 Base Pairs siRNA Sequences Targeting Both Human and Mouse PDGFRbeta:

mhPDGFRB25-1:
(SEQ ID NO. 92)
sense, 5'-CUGCAGAGACCUCAAAAGGUGUCCA-3';

(SEQ ID NO. 93)
antisense: 5'-UGGACACCUUUUGAGGUCUCUGCAG-3'

mhPDGFRB25-2:
(SEQ ID NO. 94)
sense, 5'-CAGAGACCUCAAAAGGUGUCCACGU-3';

(SEQ ID NO. 95)
antisense: 5'-ACGUGGACCCUUUGAGGUCUCUG-3'

mhPDGFRB25-3:
(SEQ ID NO. 96)
sense, 5'-GUGGUGGUGAUCUCAGCCAUCUUGG-3';

(SEQ ID NO. 97)
antisense: 5'-CCAGGAUGGUCUGAUAACCACCA-3'

mhPDGFRB25-4:
(SEQ ID NO. 98)
sense, 5'-GGUGGUGAUCUCAGCCAUCUUGGCC-3';

(SEQ ID NO. 99)
antisense: 5'-GGCCAGGAUGGUCUGAUAACCACCA-3'

mhPDGFRB25-5:
(SEQ ID NO. 100)
sense, 5'-GGCAAGCUGGUCAAGAUCUGUGACU-3';

(SEQ ID NO. 101)
antisense: 5'-AGUCACAGAUCUUGACCAGCUUGCC-3'

mhPDGFRB25-6:
(SEQ ID NO. 102)
sense, 5'-GCAAGCUGGUCAAGAUCUGUGACU-3';

(SEQ ID NO. 103)

-continued

antisense: 5'-AAGUCACAGAUCCUUGACCAGCUUGC-3'

mhPDGFRB25-7: (SEQ ID NO. 104)
sense, 5'-GGGUGGCACCCCUUACCCAGAGCUG-3';

(SEQ ID NO. 105)
antisense: 5'-CAGCUCUGGGUAAGGGGUGCCACCC-3'

mhPDGFRB25-8: (SEQ ID NO. 106)
sense, 5'-GCACCCCUUACCCAGAGCUGCCCAU-3';

(SEQ ID NO. 107)
antisense: 5'-AUGGGCAGCUCUGGGUAAGGGGUGC-3'

mhPDGFRB25-9: (SEQ ID NO. 108)
sense, 5'-CUUACCCAGAGCUGCCCAUGAACGA-3';

(SEQ ID NO. 109)
antisense: 5'-UCGUUCAUGGGCAGCUCUGGGUAAG-3'

mhPDGFRB25-10: (SEQ ID NO. 110)
sense, 5'-CAUGCCUCGACGAGAUCAUGAGA-3';

(SEQ ID NO. 111)
antisense: 5'-UCUCAUGAUUCGUCGAGGCAUG-3'

mhPDGFRB25-11: (SEQ ID NO. 112)
sense, 5'-GCCUCCGACGAGAUCAUGAGAUCA-3';

(SEQ ID NO. 113)
antisense: 5'-UGAUUCAUGAUUCGUCGAGGC-3'

mhPDGFRB25-12: (SEQ ID NO. 114)
sense, 5'-CCGACGAGAUCAUGAGAUCAUGCA-3';

(SEQ ID NO. 115)
antisense: 5'-UGCAUGAUUCAUAGAUUCGUCGG-3'

mhPDGFRB25-13: (SEQ ID NO. 116)
sense, 5'-GACGAGAUCAUGAGAUCAUGCAGA-3';

(SEQ ID NO. 117)
antisense: 5'-UCUGCAUGAUUCAUAGAUUCGUC-3'

25 Base Pairs siRNA Sequences Targeting Both Human and Mouse EGFR:

mhEGFR-1: (SEQ ID NO. 118)
sense, 5'-CUCUGGAUCCAGAAAGGUGAGAAAG-3';

(SEQ ID NO. 119)
antisense: 5'-CUUUCACCCUUCUGGGAUCCAGAG-3'

mhEGFR-2: (SEQ ID NO. 120)
sense, 5'-CAGCAUGUCAAGAUACAGAUUUUG-3';

(SEQ ID NO. 121)
antisense: 5'-CAAAUCUGUGAUUCUUGACAUGCUG-3'

mhEGFR-3: (SEQ ID NO. 122)
sense, 5'-GAUCACAGAUUUUGGGCUGGCCAAA-3';

(SEQ ID NO. 123)
antisense: 5'-UUUGCCAGCCCAAAUCUGUGAUC-3'

-continued

mhEGFR-4: (SEQ ID NO. 124)
sense, 5'-CAGAUUUUGGGCUGGCCAAACUGCU-3'

(SEQ ID NO. 125)
antisense: 5'-AGCAGUUUGGCCAGCCCAAAUUCUG-3'

mhEGFR-5: (SEQ ID NO. 126)
sense, 5'-C AAAGUGCCUAUCAAGUGGAGUGCA-3'

(SEQ ID NO. 127)
antisense: 5'-UGCCAUCCACUUGAUAGGCACUUUG-3'

mhEGFR-6: (SEQ ID NO. 128)
sense, 5'-CCAUCGAUGUCUACAUGAUCAUGGU-3';

(SEQ ID NO. 129)
antisense: 5'-ACCAUGAUCAUGUAGACAUCGAUGG-3'

mhEGFR-7: (SEQ ID NO. 130)
sense, 5'-CGAUGUCUACAUGAUCAUGGUCAGU-3';

(SEQ ID NO. 131)
antisense: 5'-ACUUGACCAUGAUCAUGUAGACAUCG-3'

mhEGFR-8: (SEQ ID NO. 132)
sense, 5'-CUACAUGAUCAUGGUCAAGUCUGG-3';

(SEQ ID NO. 133)
antisense: 5'-CCAGCACUUGACCAUGAUCAUGUAG-3'

mhEGFR-9: (SEQ ID NO. 134)
sense, 5'-CAUGAUCAUGGUCAAGUCUGGAGUA-3';

(SEQ ID NO. 135)
antisense: 5'-UCAUCCAGCACUUGACCAUGAUCAUG-3'

mhEGFR-10: (SEQ ID NO. 136)
sense, 5'-GGAUGAAAGAAUGCAUUUGCCAAGU-3';

(SEQ ID NO. 137)
antisense: 5'-ACUUGGCAAUGCAUUCUUUAUCC-3'

mhEGFR-11: (SEQ ID NO. 138)
sense, 5'-GACAACCCUGACUACCAGCAGGACU-3'

(SEQ ID NO. 139)
antisense: 5'-AGUCCUGCUGGUAGUCAGGGUUGUC-3'

mhEGFR-12: (SEQ ID NO. 140)
sense, 5'-CCUUCUAAAGACCAUCCAGGAGGU-3';

(SEQ ID NO. 141)
antisense: 5'-ACCUCUGGAUGGUCUUUAGAAGG-3'

[0061] Another embodiment is use of a combination of these oligos in the same drug payload. When three or more of those 25 base pair siRNAs targeting three or more genes involving in the disease process were packaged in the same nanoparticles and delivered through the same administration routs (or different routes), the treatment is e more effective since the knockdown of multiple gene expressions can effectively block the particular tumorigenic pathway. The following combinations can be further expended using the same general principles applied here.

Combinations of 25 Base Pair siRNA Oligos, Each of Them can Target the Corresponding Genes from Both Human and Mouse, can Achieve More Potent Therapeutic (Multi-Targeted) Efficacy:

1. Targeting VEGF Pathways:

VEGF-VEGFR1-VEGFR2 Cocktail A:

[0062]

mhVEGF25-1: (SEQ ID NO. 142)
sense, 5'-CAAGAUCGCGACGUGUAAAUGUU-3';
antisense, 5'-AACAUUUACACGUCUGCGGAUCUUG-3'
mhVEGFR125-3: (SEQ ID NO. 144)
sense, 5'-CAAGGAGGCCUCUGAUGGUGAUGU-3';
antisense, 5'-ACAUCACCAUCAGAGGCCUCCUUG-3'
mhVEGFR225-1: (SEQ ID NO. 146)
sense, 5'-CCUACGACCGUUAAAGCGGGCCAAU-3';
antisense, 5'-AUUGGCCCGCUUAAACGGUCCGUAGG-3'

VEGF-VEGFR1-VEGFR2 Cocktail B:

[0063]

mhVEGF25-1: (SEQ ID NO. 148)
sense, 5'-CAAGAUCGCGACGUGUAAAUGUU-3';
antisense, 5'-AACAUUUACACGUCUGCGGAUCUUG-3'
mhVEGFR125-6: (SEQ ID NO. 150)
sense, 5'-CCAGAAAGUGCAUUCACGGGACCU-3';
antisense, 5'-AGGUCCCGAUGAAUGCACUUUCUGG-3'
mhVEGFR225-2: (SEQ ID NO. 152)
sense, 5'-CUCAUGUCUGUUCUCAAGAUCUCA-3';
antisense, 5'-UGAGGAUCUUGAGAACAGACAUGAG-3'

VEGF-VEGFR1-VEGFR2 Cocktail C:

[0064]

mhVEGF25-2: (SEQ ID NO. 154)
sense, 5'-GCAGCUUGAGUUAACGAACGUACU-3';
antisense, 5'-AGUACGUUCGUUUAACUCAAGCUGC-3'
mhVEGFR125-6: (SEQ ID NO. 156)
sense, 5'-CCAGAAAGUGCAUUCACGGGACCU-3';

-continued

(SEQ ID NO. 157)
antisense: 5'-AGGUCCCGAUGAAUGCACUUUCUGG-3'
mhVEGFR225-2: (SEQ ID NO. 158)
sense, 5'-CUCAUGUCUGUUCUCAAGAUCUCA-3';
antisense: 5'-UGAGGAUCUUGAGAACAGACAUGAG-3'

[0065] Many other combinations with the 25 base pair siRNA oligos targeting VEGF, VEGFR1 and VEGFR2 respectively can be composed for the best anti-angiogenesis efficacy, by either equal or different ratios of each components.

2. Targeting both VEGF and EGF Pathways:

VEGF-VEGFR2-EGFR Cocktail A:

[0066]

mhVEGF25-1: (SEQ ID NO. 160)
sense, 5'-CAAGAUCCGACGUGUAAAUGUU-3';
antisense, 5'-AACAUUUACACGUCUGCGGAUCUUG-3'
mhVEGFR225-1: (SEQ ID NO. 162)
sense, 5'-CCUACGACCGUUAAAGCGGGCCAAU-3';
antisense, 5'-AUUGGCCCGCUUAAACGGUCCGUAGG-3'
mhEGFR-1: (SEQ ID NO. 164)
sense, 5'-CUCUGGAUCCAGAAGGUGAGAAAG-3';
antisense, 5'-CUUUCUCCACUUCUGGAUCCAGAG-3'

VEGF-VEGFR2-EGFR Cocktail B:

[0067]

mhVEGF25-1: (SEQ ID NO. 166)
sense, 5'-CAAGAUCCGACGUGUAAAUGUU-3';
antisense, 5'-AACAUUUACACGUCUGCGGAUCUUG-3'
mhVEGFR225-2: (SEQ ID NO. 168)
sense, 5'-CUCAUGUCUGUUCUCAAGAUCUCA-3';
antisense: 5'-UGAGGAUCUUGAGAACAGACAUGAG-3'
mhEGFR-5: (SEQ ID NO. 170)
sense, 5'-CAAAGUGCCUAUCAAGUGGAUGGCA-3';
antisense: 5'-UGCAUCCACUUGAUAGGCACUUUG-3'

VEGF-VEGFR2-EGFR Cocktail C:

[0068]

mhVEGF25-2: (SEQ ID NO. 172)
sense, 5'-GCAGCUUGAGUUAACGAAACGUACU-3';

(SEQ ID NO. 173)
antisense, 5'-AGUACGUUCGUUUAACUCAAGCUGC-3'

mhVEGFR225-2: (SEQ ID NO. 174)
sense, 5'-CUCAUGUCUGUUCUCAAGAUCUCA-3';

(SEQ ID NO. 175)
antisense: 5'-UGAGGAUCUUGAGAACAGACAUGAG-3'

mhEGFR-9: (SEQ ID NO. 176)
sense, 5'-CAUGAUCAUGGUCAAGUGCUGGAUGA-3';

(SEQ ID NO. 177)
antisense: 5'-UCAUCCAGCACUUGACCAUGAUCAUG-3'

[0069] Many other combinations can be made with other sequences targeting those three genes with different composition and equal or different ratios of the siRNA oligos.

3. Targeting VEGF, PDGF and EGF Pathways:

VEGFR2-PDGFRa-EGFR Cocktail A:

[0070]

mhVEGFR225-1: (SEQ ID NO. 178)
sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3'

(SEQ ID NO. 179)
antisense: 5'-AUUGCCCGCUUAACGGUCGGUAGG-3'

mhPDGFRa25-1: (SEQ ID NO. 180)
sense, 5'-GAAGAUAAUGACUCACCUGGGGCCA-3';

(SEQ ID NO. 181)
antisense: 5'-UGGCCCCAGGUGAGUCAUUUCUUC-3'

mhEGFR-1: (SEQ ID NO. 182)
sense, 5'-CUCUGGAUCCAGAAAGGUGAGAAAG-3';

(SEQ ID NO. 183)
antisense: 5'-CUUUCACCCUUCUGGGAUCCAGAG-3'

VEGFR1-VEGFR2-PDGFRa-EGFR Cocktail B:

[0071]

mhVEGFR125-6: (SEQ ID NO. 184)
sense, 5'-CCAGAAAGUGCAUUCACGGGACCU-3';

(SEQ ID NO. 185)
antisense: 5'-AGGUCCGAUGAAUGCACUUUCUGG-3'

mhVEGFR225-1: (SEQ ID NO. 186)
sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3';

(SEQ ID NO. 187)
antisense: 5'-AUUGCCCGCUUAACGGUCGGUAGG-3'

-continued

mhPDGFRa25-2: (SEQ ID NO. 188)
sense, 5'-GAUAAUGACUCACCUGGGGCCACAU-3';

(SEQ ID NO. 189)
antisense: 5'-AUGUGCCCCAGGUGAGUCAUUUAUC-3'

mhEGFR-1: (SEQ ID NO. 190)
sense, 5'-CUCUGGAUCCAGAAAGGUGAGAAAG-3';

(SEQ ID NO. 191)
antisense: 5'-CUUUCACCCUUCUGGGAUCCAGAG-3'

VEGFR2-PDGFRa-EGFR Cocktail C:

[0072]

mhVEGFR225-1: (SEQ ID NO. 192)
sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3';

(SEQ ID NO. 193)
antisense: 5'-AUUGCCCGCUUAACGGUCGGUAGG-3'

mhPDGFRa25-2: (SEQ ID NO. 194)
sense, 5'-GAUAAUGACUCACCUGGGGCCACAU-3';

(SEQ ID NO. 195)
antisense: 5'-AUGUGCCCCAGGUGAGUCAUUUAUC-3'

mhEGFR-12: (SEQ ID NO. 196)
sense, 5'-CCUUCUUAAGACCAUCCAGGAGGU-3';

(SEQ ID NO. 197)
antisense: 5'-ACCUCUGGAUGGUCUUUAAGAAGG-3'

VEGFR2-PDGFRb-EGFR Cocktail A:

[0073]

mhVEGFR225-1: (SEQ ID NO. 198)
sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3';

(SEQ ID NO. 199)
antisense: 5'-AUUGCCCGCUUAACGGUCGGUAGG-3'

mhPDGFRb25-5: (SEQ ID NO. 200)
sense, 5'-GGCAAGCUGGUCAAGAUCUGUGACU-3';

(SEQ ID NO. 201)
antisense: 5'-AGUCACAGAUCUUGACCAGCUUGCC-3'

mhEGFR-1: (SEQ ID NO. 202)
sense, 5'-CUCUGGAUCCAGAAAGGUGAGAAAG-3';

(SEQ ID NO. 203)
antisense: 5'-CUUUCACCCUUCUGGGAUCCAGAG-3'

VEGFR2-PDGFRb-EGFR Cocktail B:

[0074]

mhVEGFR225-1: (SEQ ID NO. 204)

-continued

sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3';
 (SEQ ID NO. 205)
 antisense: 5'-AUUGCCCGCUUAACGGUCCGUAGG-3'
 mhPDGFRB25-10:
 (SEQ ID NO. 206)
 sense, 5'-CAUGCCUCCGACGAGAUCUAUGAGA-3';
 (SEQ ID NO. 207)
 antisense: 5'-UCUCAUAGAUCUCGUCGGAGGCAUG-3'
 mhEGFR-1:
 (SEQ ID NO. 208)
 sense, 5'-CUCUGGAUCCAGAAAGGUGAGAAAG-3';
 (SEQ ID NO. 209)
 antisense: 5'-CUUUCACCCUUCUGGGAUCCAGAG-3'

VEGFR2-PDGFRb-EGFR Cocktail C:

[0075]

mhVEGFR225-1:
 (SEQ ID NO. 210)
 sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3';
 (SEQ ID NO. 211)
 antisense: 5'-AUUGCCCGCUUAACGGUCCGUAGG-3'
 mhPDGFRB25-5:
 (SEQ ID NO. 212)
 sense, 5'-GGCAAGCUGGUCAAGAUCUGUGACU-3';
 (SEQ ID NO. 213)
 antisense: 5'-AGUCACAGAUUCUAGCCAGCUUGCC-3'
 mhEGFR-12:
 (SEQ ID NO. 214)
 sense, 5'-CCUUCUUAAGACCAUCCAGGAGGU-3';
 (SEQ ID NO. 215)
 antisense: 5'-ACCUCUGGAUGGUCUUUAAGAAGG-3'

[0076] Many other combinations can be made with other sequences targeting those three genes with different composition and equal or different ratios of the siRNA oligos.

[0077] The combinations can also be made with more than three sequences:

VEGFR2-PDGFRab-EGFR Cocktail A:

[0078]

mhVEGFR225-1:
 (SEQ ID NO. 216)
 sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3';
 (SEQ ID NO. 217)
 antisense: 5'-AUUGCCCGCUUAACGGUCCGUAGG-3'
 mhPDGFRa25-2:
 (SEQ ID NO. 218)
 sense, 5'-GAUAAUGACUCACCGGGGCCACAU-3'
 (SEQ ID NO. 219)
 antisense: 5'-AUGUGGCCCCAGGUGAGUCAUUUUC-3'
 mhPDGFRb25-5:
 (SEQ ID NO. 220)
 sense, 5'-GGCAAGCUGGUCAAGAUCUGUGACU-3';

-continued

(SEQ ID NO. 221)
 antisense: 5'-AGUCACAGAUUCUUGACCAGCUUGCC-3'
 mhEGFR-1:
 (SEQ ID NO. 222)
 sense, 5'-CUCUGGAUCCAGAAAGGUGAGAAAG-3';
 (SEQ ID NO. 223)
 antisense: 5'-CUUUCACCCUUCUGGGAUCCAGAG-3'

VEGFR1-VEGFR2-PDGFRb-EGFR-VEGF Cocktail A:

[0079]

mhVEGFR125-6:
 (SEQ ID NO. 224)
 sense, 5'-CCAGAAAGUGCAUUCUACGGGACCU-3';
 (SEQ ID NO. 225)
 antisense: 5'-AGGUCCCGAUGAAUGCACUUUCUGG-3'
 mhVEGFR225-1:
 (SEQ ID NO. 226)
 sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3';
 (SEQ ID NO. 227)
 antisense: 5'-AUUGCCCGCUUAACGGUCCGUAGG-3'
 mhPDGFRB25-10:
 (SEQ ID NO. 228)
 sense, 5'-CAUGCCUCCGACGAGAUCUAUGAGA-3';
 (SEQ ID NO. 229)
 antisense: 5'-UCUCAUAGAUCUCGUCGGAGGCAUG-3'
 mhEGFR-1:
 (SEQ ID NO. 230)
 sense, 5'-CUCUGGAUCCAGAAAGGUGAGAAAG-3';
 (SEQ ID NO. 231)
 antisense: 5'-CUUUCACCCUUCUGGGAUCCAGAG-3'
 mhVEGF25-2:
 (SEQ ID NO. 232)
 sense, 5'-GCAGCUUGAGUUAACGAACGUACU-3';
 (SEQ ID NO. 233)
 antisense: 5'-AGUACGUUCGUUUAACUCAAGCUGC-3'

VEGFR1-VEGFR2-PDGFRb-EGFR-VEGF Cocktail C:

[0080]

mhVEGFR125-6:
 (SEQ ID NO. 234)
 sense, 5'-CCAGAAAGUGCAUUCUACGGGACCU-3';
 (SEQ ID NO. 235)
 antisense: 5'-AGGUCCCGAUGAAUGCACUUUCUGG-3'
 mhVEGFR225-1:
 (SEQ ID NO. 236)
 sense, 5'-CCUACGGACCGUUAAGCGGGCCAAU-3'
 (SEQ ID NO. 237)
 antisense: 5'-AUUGCCCGCUUAACGGUCCGUAGG-3'
 mhPDGFRa25-2:
 (SEQ ID NO. 238)
 sense, 5'-GAUAAUGACUCACCGGGGCCACAU-3';

-continued

(SEQ ID NO. 239)
antisense: 5'-AUGUGGCCCCAGGUGAGUCAUUAUC-3'

mhPDGFRB25-5:
(SEQ ID NO. 240)
sense, 5'-GGCAAGCUGGUCAAGAUCUGUGACU-3';

(SEQ ID NO. 241)
antisense: 5'-AGUCACAGAUCUUGACCAGCUUGCC-3'

mhEGFR-12:
(SEQ ID NO. 242)
sense, 5'-CCUUCUUAAGACCAUCCAGGAGGU-3';

(SEQ ID NO. 243)
antisense: 5'-ACCUCUGGAUGGUCUUUAAGAAG-3'

mhVEGF25-2:
(SEQ ID NO. 244)
sense, 5'-GCAGCUUGAGUUAACGAACGUACU-3'

(SEQ ID NO. 245)
antisense, 5'-AGUACGUUCGUUUAACUCAAGCUGC-3'

[0081] B. Combined VEGF Pathway Gene Inhibition

[0082] A number of factors were considered during development of the compositions and methods of the present invention for inhibition of NV. First, NV diseases are complex and the result of multiple proteins and abnormally over-expressed disease-causing genes and multiple malfunctions of disease-causing proteins. Second, nucleic acid agents that activate RNA interference (RNAi) are highly selective inhibitors of gene expression in a sequence specific manner. Thirdly, inhibition of NV by modulation of protein activity can be achieved by a number of methods, including an inhibition of protein function (antagonists), stimulation of protein function (agonists), reduction of protein expression levels, and post transcriptional modification of proteins. Treatment of NV disease ideally requires sophisticated therapeutic actions to effectively shut down a particular biological pathway that is critical for disease progression, including simultaneously blocking functions of both ligands and their receptor, simultaneously blocking receptor activity and down stream signaling proteins, and simultaneously blocking redundant elements of a pathway. This is the case for the NV disease and the VEGF pathway.

[0083] However, identification of a single agent selective for two or three or more proteins is difficult and oftentimes impractical, if not impossible. To overcome this difficulty, use of a combination of drugs has been something of a trend in modern medicine. In oncology applications, combined chemotherapies have achieved remarkable anti-cancer efficacy. One example is the use of docetaxel, ifosfamide and cisplatin combination therapy for treatment of oropharyngeal cancer with multiple bone metastases from prostate cancer (2). In other therapeutic areas, another example is the treatment of ulcerative colitis with combination of Corticosteroids, Metronidazole and Vancomycin (3). For treatment of insufficiently controlled type 2 diabetes, the efficacy and safety of adding rosiglitazone to a combination of glimepiride and metformin therapy were evaluated (4).

[0084] Although those clinical studies have demonstrated remarkable therapeutic efficacies, the toxicities of higher dosage and long time safeties are always major concerns, due to their different sources of origins, different manufacturing processes and different chemistry properties. To overcome these problems, an aspect of the present invention are using s

siRNA oligonucleotide gene inhibitors to provide a unique advantage to achieve combination effects with a combination of siRNA to target multiple disease causing genes in the same treatment. One advantage provided by the present invention is a result that all siRNA oligonucleotides are very similar chemically and pharmacologically, and can be from the same source of origin and same manufacturing process. Another advantage provided by the present invention is that multiple siRNA oligonucleotides can be formulated in a single preparation such as a nanoparticle preparation.

[0085] Accordingly, an aspect of the present invention is to combine siRNA agents so as to achieve specific and selective inhibition of multiple VEGF pathway genes and as a result achieve a inhibition of NV disease and a better clinical benefit. The present invention provides for many combinations of siRNA targeting including combinations that target VEGF itself together with its receptors including VEGF R1 (Flt1) and VEGF R2 (KDR), parallel growth factors including PDGF and EGF and their receptors, down stream signaling factors including RAF and AKT, and transcription factors including NFkB, and their combination. A preferred embodiment is a combination of siRNA inhibiting VEGF and two of its receptors VEGF R1 (Flt1) and VEGF R2 (KDR). Another preferred embodiment is a combination of siRNA inhibiting VEGF and its receptors, PDGF and its receptors, and EGF and its receptors. Yet another preferred embodiment is a combination of siRNA inhibiting VEGF and its receptors and down stream signaling.

[0086] The dsRNA oligonucleotides can be combined for a therapeutic for the treatment of NV disease. In one embodiment of the present invention they can be mixed together as a cocktail and in another embodiment they can be administered sequentially by the same route or by different routes and formulations and in yet another embodiment some can be administered as a cocktail and some administered sequentially. Other combinations of siRNA and methods for their combination will be understood by one skilled in the art to achieve treatment of NV diseases.

[0087] C. Combined VEGF Pathway Antagonist and Gene Inhibition

[0088] Disease is complicated and often involves multiple pathological processes as well as variations in severity of disease symptoms and, often, variations from one patient to another. Many diseases are caused by abnormal overexpression of disease-causing or disease control genes, or from foreign infectious organisms, or both. Disease progression, the development of reduced response to treatments over time and drug resistance also limit clinical benefit of a single treatment or modality. One means to overcome such limitations is through use of combinations of treatments and drugs.

[0089] Therefore, an aspect of the present invention is to combine siRNA agents with other agents so as to achieve a strong, durable, and robust inhibition of NV disease and a superior clinical benefit. The present invention provides multiple combinations of siRNA agents together with other agents, including combinations of therapeutic siRNA for VEGF itself and its receptors with antagonists of VEGF itself and its receptors (such as Avastin). The present invention also provides combinations of therapeutic siRNA agents with orally available kinase inhibitors (such as SU1 1248). The present invention also provides combinations of therapeutic siRNA agents with immunotherapy. Yet another embodiment of the present invention is to combine siRNA with antiprolif-

erative agents. Other combinations of siRNA agents and other agents will be understood by one skilled in the art to achieve treatment of NV diseases.

[0090] D. Formulation and Administration

[0091] Recent efforts towards developing tissue targetable nucleic acid delivery systems based on synthetic reagents have produced promising results. To be robust, effective delivery systems should have multiple levels of selectivity, i.e. selective localization at the disease tissue and selective inhibition of biochemical pathways driving the pathology. Moreover, the most effective current therapies require “multi-targeted” therapeutics, i.e. designer “dirty” drugs with multiple mechanisms of activity, blocking redundant pathological pathways. A superior approach would be use “smart” nanoparticles that simultaneously target disease and deliver nucleic acid agents into the target cells and into the correct subcellular compartment.

[0092] In one embodiment, the present invention provides for formulations for siRNA dsRNA oligonucleotides that comprise tissue-targetable delivery with three additional properties. These properties are nucleic acid binding into a core that can release the siRNA into the cytoplasm, protection from non-specific interactions, and tissue targeting that provides cell uptake. No one material has all of these required properties in one molecule. The invention provides for compositions and methods that use modular conjugates of three materials to combine and assemble the multiple properties required. They can be designed and synthesized to incorporate various properties and then mixed with the siRNA payload to form the nanoparticles. Based on these embodiments, a preferred embodiment comprises a modular polymer conjugate targeting neovasculature by coupling a peptide ligand specific for those cells to one end of a protective polymer, coupled at its other end to a cationic carrier for nucleic acids. This polymer conjugate has three functional domains, sometimes referred to as a tri-functional polymer (TFP). The modular design of this conjugate allows replacement and optimization of each component separately. An alternative approach has been to attach surface coatings onto preformed nanoparticles. Adsorption of a steric polymer coating onto polymers is self-limiting; once a steric layer begins to form it will impede further addition of polymer. The compositions and methods of the invention permit an efficient method for optimization of each of the three functions, largely independent of the other two functions.

Formation of Nucleic Acid Core Particle

[0093] Delivery agents for nucleic acids must assist their gaining access to the interior of cells and in a manner such that they can exert their biological activity. Efforts to address this challenge for nucleic acid therapeutics with synthetic materials include use of simple cationic lipid and polymer complexes developed as in vitro DNA transfection reagents. It was quickly found that both getting nucleic acids into cells and obtaining biological activity is extremely difficult. For example, achieving a stable nucleic acid package for transport is possible but not always easy to reconcile with the need to release the nucleic acid into the nucleus, or in the case of siRNA into the cytoplasm. Also, a large number of cationic lipids and polymers that are effective in vitro do not retain activity when administered in vivo, unfortunately for reasons still largely unknown, providing little predictive value for developing complexes active in vivo. Interest in studies with RNA have lagged far behind, until now with the recent erup-

tion of interest in siRNA. Nonetheless, lung tissue can often be transfected from an intravenous administration of DNA complexes. Similar biological activity has been observed when siRNA has been used as the payload.

[0094] Recent studies have identified one class of cationic polymers composed of defined polypeptide structures that appear to have broad capabilities. A large number of members of this class can be synthesized with defined structures covering linear and branched forms and have been found to offer biological activity both in vitro and in vivo. They have been shown to have activity with several types of nucleic acids, including plasmids and DNA or RNA oligonucleotides. The success with this class of cationic polymer appears to result from a design incorporating specific structures, including branching, and using a mixture of hard and weak bases to form a polymer with mixed cationic properties. Another advantage of this particular class is its biodegradable nature, being constructed entirely from natural amino acids, albeit non-natural branching. Having a polymer with at least two properties to use for optimization, continuing development of materials for forming effective core complexes with siRNA can be refined and focused on for further improvements to balance stability and cytoplasmic release. Optimized core forming polymers can be used as the basis for the other functions.

Protective Steric Coating

[0095] Even liposomes with an external lipid bilayer resembling the outer cellular membrane are rapidly recognized and cleared from blood. Unlike viral particles, though, nanotechnology has access to a broad range of synthetic polymer chemistry. Hydrophilic polymers, such as PEG and polyacetals and polyoxazolines, have proven effective to form a “steric” protective layer on the surface of colloidal drug delivery systems whether liposomes, polymer or electrostatic nanoparticles, reducing immune clearance from blood. The use of this steric PEG layer was first developed and most extensively studied with sterically stabilized liposomes. The present invention provides for alternative approaches, such as chemical reduction of surface charge, in addition to a steric polymer coating.

[0096] The evidence is strong that the steric barrier, and biological consequences, derive from physical, not chemical, properties. Several other hydrophilic polymers have been reported as alternatives to PEG. Physical studies on sterically stabilized liposomes have provided a strong mechanistic underpinning for physical behavior of the polymer layer and can be used to achieve similar coatings on other types of particles. However, while physical studies have shown formation of a similar polymer layer on the surface of polymer complexes with nucleic acids, and achievement of similar biological properties, we lack sufficient information today to use of the physical properties to accurately predict the desired biological properties, protection from immune clearance from blood.

[0097] Advanced studies are constructing nanoparticle complexes with controlled variations in their physical properties followed by determination of their biological properties. From the many liposome studies, it is clear that physical properties with the greatest impact on biological activity can be obtained by synthesis of a matrix of conjugates varying size of the two polymers and the grafting density. Note that while the surface steric layer function is due to physical

properties, the optimal conjugation chemistry still depends on the specific chemical nature of the steric polymer and the carrier to which it is coupled.

[0098] Methods for formation of the nanoparticles with the surface steric polymer layer are also a relevant parameter. One embodiment of the present invention provides for methods where the steric polymer is coupled to the carrier polymer to give a conjugate that self-assembles with the nucleic acid forming a nanoparticle with the steric polymer surface layer. Another embodiment is to attach surface coatings onto pre-formed nanoparticles. In self-assembly formation of the surface steric layer depends on interactions of the carrier polymer with the payload, not on penetration through a forming steric layer to react with particle surface. In this case, effects of the steric polymer on ability of the carrier polymer to bind the nucleic acid payload may have adverse effects on particle formation, and thus the surface steric layer. If this occurs, the grafting density of the steric polymer on the carrier will have exceeded its maximum, or the structural nature of the grafting is not adequate.

Surface Exposed Ligands Targeting Specific Tissues

[0099] Ability of the nanoparticle to reach the interior of the target cells, selectively, resides in its ability to induce a specific receptor mediated uptake. This is provided in the present invention by exposed ligands with the binding specificity. While many types of ligands have been studied extensively for targeting colloidal delivery systems, many have relied on antibodies. Over twenty years ago, this idea was developed for coupling antibodies to the surface of liposomes, usually referred to as immunoliposomes. One important parameter that has emerged in the impact of ligand density. The extensive studies achieved good success *in vitro* but only recently have begun to produce positive results *in vivo*, now reaching the stage of clinical development for delivery of small molecule drugs.

[0100] Antibodies tend to meet many requirements for use as the ligand, including good binding selectivity and nearly routine preparation for nearly any receptor and broad applicability of protein coupling methods regardless of nanoparticle type. Monoclonal antibodies even show signs of utility to cross the blood-brain-barrier. On the other hand, they are not ideal. As targeting ligands, antibodies are very large proteins, about 150 Kd. This makes conjugation to yield a specific orientation difficult unless an extra Cys amino acid residue can be introduced, or other unique site identified, for the chemical coupling. Truncated forms can be used to reduce size but loss of affinity can occur, such as with "scFv" constructions.

[0101] Another troubling property of antibodies is their multitude of biological activities, many conveyed in domains not associated with antigen binding, which are part of their role in the mammalian immune system. These immune activities can be exploited but they also can interfere in the nanoparticle steric layer role of avoiding immune clearance. Other proteins that are natural ligands also have been considered for targeting nanoparticles, such as transferrin for transferrin receptor and Factor VII and Factor VIIa proteins and their fragments. These agents can be good candidates but their successful use requires considerable effort to identify specific coupling chemistry, coupling without loss of binding activity, and exposure on the particle without reintroducing immune clearance.

[0102] A preferred class of ligands are small molecular weight compounds with strong selective binding affinity for internalizing receptors. Studies have been evaluated natural metabolites such as vitamins like folate and thiamine, polysaccharides like wheat germ agglutinin or sialyl Lewis^x for e-selectin, and peptide binding domains like RGD for integrins. Peptides offer a versatile class of ligand, since phage display libraries can be used to screen for natural or unnatural sequences, even with *in vivo* panning methods. Such phage display methods can permit retention of an unpaired Cys residue at one end for ease of coupling regardless of sequence. The recent *in vivo* success using an RGD peptide for targeted delivery of nanoparticles to neovasculature indicates this approach can be very effective to meet the major requirements for effective ligands: specific chemistry that doesn't interfere in ligand binding or induce immune clearance yet enables selective receptor mediated uptake at the target cells.

[0103] A preferred embodiment of the invention comprises an RGD-2C peptide ligand conjugate. The RGD-2C ligand provides localization at neovasculature as well as tumor cells nearby tumor neovasculature, intracellular internalization of the nucleic acid containing nanoparticles, and release of the nucleic acid giving effective biological activity of the nucleic acid.

[0104] Another preferred embodiment of the invention comprises mixtures of ligand-conjugates of two or more ligands, or conjugates of two or more ligands. A preferred embodiment of the invention comprises mixtures of RGD peptide ligands with Factor VII proteins or their peptide fragments or chemical analogues. Another preferred embodiment of the invention comprises mixtures of RGD peptide ligands with ligands binding e-selectins. Another preferred embodiment of the invention comprises mixtures of RGD peptide ligands with apoE proteins or their peptide fragments or chemical analogues. Another preferred embodiment of the invention comprises mixtures of RGD peptide ligands with tumor cell binding ligands such as folate, transferrin, porphyrin, bFGF, and EGF. Another preferred embodiment of the invention comprises mixtures of three or more of these ligands. A preferred embodiment of the invention comprises a conjugate of two or more ligands at one domain of a carrier and a polyamine agent at a second domain of said carrier.

[0105] Another preferred embodiment of the invention comprises mixtures of ligand conjugates. The mixture of ligand conjugates can comprise the ligands conjugated to the same carrier agent and optionally to the same domain of the carrier or the mixture of ligand conjugates can comprise the ligands conjugated to different carrier agents or optionally to different domains of carrier agents. The mixture of ligand conjugates can be further comprised of protective, fusogenic, or carrier agents, such as PEG or polyacetals or pH-sensitive polymers or polyamines or histidine-lysine copolymers.

[0106] The compositions and methods of the present invention provide for administration of RNAi comprising polymers, polymer conjugates, lipids, micelles, self-assembly colloids, nanoparticles, sterically stabilized nanoparticles, or ligand-directed nanoparticles. Targeted synthetic vectors of the type described in WO01/49324, which is hereby incorporated by reference in its entirety, may be used for systemic delivery of nucleic acids of the present invention inducing RNAi. In one embodiment, a PEI-PEG-RGD (polyethyleneimine-polyethylene glycol-argine-glycine-aspartic acid) synthetic vector can be prepared and used, for example as in

Examples 53 and 56 of WO01/49324. This vector was used to deliver RNAi systemically via intravenous injection. The skilled artisan will recognize that other targeted synthetic vector molecules known in the art may be used. For example, the vector may have an inner shell made up of a core complex comprising the RNAi and at least one complex forming reagent.

[0107] The vector also may contain a fusogenic moiety, which may comprise a shell that is anchored to the core complex, or may be incorporated directly into the core complex. The vector may further have an outer shell moiety that stabilizes the vector and reduces nonspecific binding to proteins and cells. The outer shell moiety may comprise a hydrophilic polymer, and/or may be anchored to the fusogenic moiety. The outer shell moiety may be anchored to the core complex. The vector may contain a targeting moiety that enhances binding of the vector to a target tissue and cell population. Suitable targeting moieties are known in the art and are described in detail in WO01/49324.

a. Preparation of RGD-Targeted Nanoparticles Containing VEGF Pathway siRNA:

[0108] One embodiment of the present invention provides compositions and methods for RGD-mediated ligand-directed nanoparticle preparations of anti-VEGF pathway siRNA short dsRNA molecules. FIGS. 25 and 26 show a method for the manufacture of RGD-2C-mediated tissue targeted nanoparticle containing siRNA. The targeting ligand, RGD containing peptide (ACRGDMFGCA) (SEQ ID NO: 246) is conjugated to a steric polymer such as polyethylene glycol, or other polymers with similar properties. This ligand-steric polymer conjugate is further conjugated to a polycation such as polyethyleneimine or other effective material such as a histidine-lysine copolymer. The conjugates can be covalent or non-covalent bonds and the covalent bonds can be non-cleavable or they can be cleavable such as by hydrolysis or by reducing agents. The conjugates can be prepared with hetero-bifunctional polymers, such as an NHS-PEG-VS commercially available reagent (Nektar) or hetero-bifunctional agents prepared for manufacturing, or with homo-bifunctional polymers, such as NHS-PEG-NHS commercially available reagent (Rapp Polymere) or homo-bifunctional agents prepared for manufacturing. One preferred embodiment of the present invention comprises use of a homo-bifunctional NHS-PEG-NHS agent, such as a reagent commercially available from RAPP Polymere, first to couple with an RGD-2C peptide prepared by solid phase synthesis and send to couple with a polyamine containing polymer such as PEI commercially available or a histidine-lysine copolymer prepared by solid phase synthesis.

[0109] Another preferred embodiment of the present invention comprises a first combination of an RGD-PEG-polyamine conjugate with a polyamine polymer, such as PEI or a histidine-lysine copolymer, and a second combination of the said polymer mixture with a second mixture of dsRNA oligonucleotides. A solution comprising the polymer conjugate, or comprising a mixture of a polymer conjugate with other polymer, lipid, or micelle such as materials comprising a ligand or a steric polymer or fusogen, is mixed with a solution comprising the nucleic acid, in one embodiment an siRNA targeted against specific genes of interest, in desirable ratios to obtain nanoparticles that contain siRNA.

[0110] In this embodiment, nanoparticles are formed by layered nanoparticle self-assembly comprising mixing the polymer conjugate and the nucleic acid. Non-covalent elec-

trostatic interactions between the negatively charged nucleic acid and the positively charged segment of the polymer conjugate drive the self-assembly process that leads to formation of nanoparticles. This process involves simple mixing of two solutions where one of the solutions containing the nucleic acid is added to another solution containing the polymer conjugate followed by or concurrent stirring. In one embodiment, the ratio between the positively charged components and the negatively charged components in the mixture is determined by appropriately adjusting the concentrations of each solution or by adjusting the volume of solution added. In another embodiment, the two solutions are mixed under continuous flow conditions using mixing apparatus such as static mixer (FIG. 26). In this embodiment, two or more solutions are introduced into a static mixer at rates and pressures giving a ratio of the solutions, where the streams of solutions get mixed within the static mixer. Arrangements are possible for mixers to be arranged in parallel or in series.

[0111] E. Combined Formulation and Electric Field

[0112] For certain applications, RNAi may be administered with or without application of an electric field. This can be used, for example, to deliver the RNAi molecules of the invention via direct injections into, for example, tumor tissue and directly into or nearby an angiogenic tissue or a tissue with neovasculature. The siRNA may be in a suitable pharmaceutical carrier such as, for example, a saline solution or a buffered saline solution. The present invention, thus generally described, will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.

EXAMPLES

Example 1

Tumor Growth Inhibition with Combination Use of siRNA Duplexes Targeting VEGFR2 and Bevacizumab (Avastin) against VEGF

Material and Methods.

[0113] Reagents: Avastin, monoclonal antibody against VEGF (25 mg/ml, Genetech); siRNA against mouse VEGFR2, sequence ((a) AAGCTCAGCACACAGAAAGAC (SEQ ID NO: 247); (b) AATGCGGCGGTGGTGACAGTA) (SEQ ID NO: 248); siRNA against luciferase (Qiagen); Avertin made of 1.5 gram 2,2,2-Tribromoethanol and 1.5 ml t-amyl alcohol (Cat# T4840-2, Cat#24048-6, Aldrich) in 100 ml distill water, St. Louis, Mo.]. Mice: athymus female nude mice, 5 to 6 weeks old, were purchased from TACONIC and housed conventionally. All investigations followed guidelines of the Committee on the Care of Laboratory Animals Resources, Commission of Life Sciences, National Research Council. The animal facility of Biomedical Research Institute in Rockville Maryland is fully accredited by the American Association of Laboratory Animal Care. Cells: Colon carcinoma cell line, DLD-1 (CCL-221, ATCC) was grown in RPMI 1640 medium with 2 mM L-glutamine, 1.5 g/L sodium bicarbonate, 4.5 g/L glucose, 10 mM HEPES, and 1.0 mM sodium pyruvate, 10% fetal bovine serum.

[0114] Procedure: 1) DLD-1 cells near confluence were harvested and resuspended in serum-free RPMI medium. 2) Mice were anaesthetized with Avertin, 0.4 ml/mouse i.p. 3) 100 million cells in 0.1 ml serum-free RPMI medium were injected into mice back s.c. on the left flank for establishment

of xenograft tumor model. 4) 5 days after inoculation of tumor cells, sizes of growing tumors were measured with a caliper. Mice were then randomly grouped into 6 groups, 8 mice per group. 5) Experiment design and dosing scheme were as following. 6) SiRNA was delivered through tail-vein injection after been formed a complex with RPP.

the inhibition of NV and inhibition of tumor growth was not due to a non-specific interferon-mediated effect.

Summary

[0122] This example illustrates the present invention as it provides compositions and methods for treatment of cancer

TABLE I

Regimen of anti-angiogenesis siRNA/mAb treatment of DLD-1 tumor model								
Group	Day-5	Day-6	Day-8	Day-9	Day-11	Day-12	Day-14	Day
G1 - Luc-siRNA		siRNA		siRNA		siRNA	siRJ	
G2 - Avastin	Avastin		Avastin		Avastin		Avastin	
G3 - VEGFR2-siRNA		siRNA		siRNA		siRNA		siRJ
G4 - Mouse-IgG + VEGFR2	Mouse-IgG	siRNA	Mouse-IgG	siRNA	Mouse-IgG	siRNA	Mouse-IgG	siRJ
G5 - Avastin + Luc-siRNA	Avastin	siRNA	Avastin	siRNA	Avastin	siRNA	Avastin	siRJ
G6 - Avastin + VEGFR2-siRNA	Avastin	siRNA	Avastin	siRNA	Avastin	siRNA	Avastin	siRJ

Results and Discussion

[0115] 1. RGD ligand-directed nanoparticle carrying VEGF R2 specific siRNA dsRNA itself was able to achieve tumor growth inhibition after repeated systemic intravenous administrations (FIG. 1) with only 2 mg/kg siRNA.

[0116] 2. This siRNA-based antiangiogenesis activity (Targeting VEGF R2) was able to enhance the VEGF monoclonal antibody (mAb)-mediated antiangiogenesis, resulting further tumor growth inhibition (FIG. 1).

[0117] 3. Although many biochemical pathways may be critical independently or collectively for malignant neoplasia, it is widely acknowledged that neo-angiogenesis is critical for growth of many types of cancers. Therefore neo-angiogenesis process has naturally become one of the most targeted areas for tumor drug development. Among many factors involved in neo-angiogenesis VEGF is believed an important controlling factor and one that can enable a therapeutic effect giving clinical benefit as achieved by Avastin, a monoclonal antibody against VEGF and demonstrated to inhibit growth of colon cancer. The combination of NV targeted VEGF siRNA gene inhibition shows evidence of being additive with an antagonist therapeutic when administrated in a combination protocol.

[0118] 4. The example shows the great benefit of using gene inhibitors and with antagonists such as mAb inhibitors in the same treatment regimen to improve the therapeutic outcome.

[0119] 5. RT-PCR results directly demonstrated that the VEGFR2 mRNA level in the tumor samples treated with the nanoparticle delivered VEGFR2 specific siRNA has dramatically down regulated, comparing to the expression levels in the untreated tumors (FIG. 2).

[0120] 6. Immunohistologicalchemistry (IHC analysis) indicated that the protein level knockdown has occurred for VEGFR2 and CD31 which is a marker for neo vasculature (FIG. 3).

[0121] 7. The ELISA measurement for IFN-alpha activity in the blood samples from nanoparticle/VEGFR2 siRNA treatment alone, and in combination with Avastin, did not show any significant (detectable) upregulation or induction of IFN activity (FIG. 4), indicating that

with small interfering dsRNA oligonucleotide gene inhibitors in combination with monoclonal antibodies (mAb) in the same therapeutic regimen, where the agents inhibit both expression of disease-causing genes and biological functions of disease-causing proteins, resulting in enhanced therapeutic efficacy. In addition, the example provides illustration of simultaneously blocking both the ligand (e.g. VEGF) and its receptor (e.g. VEGF R2) to provide effective inhibition of a particular biological pathway (e.g. angiogenesis pathway) thereby providing a treatment inhibiting disease progression. Therefore, combined applications of inhibitory activities contributed by two potent and targeted biological inhibitors, RNAi and mAb, is a means for effective treatment of NV diseases including cancer.

Example 2

siRNA-Mediated VEGF Silencing In Vitro and In Vivo

[0123] Human VEGF 165 specific siRNA was transfected into the MCF-7/VEGF 165 cell (hVEGF over-expression) by Lipofectamine resulting in knockdown of hVEGF mRNA in the cell (FIG. 2 upper panel). Using an electroporation procedure, MCF-7/VEGF 165 cells were transfected by hVEGF 165 specific siRNA, resulting in down regulation of hVEGF expression as determined with an ELISA analysis.

[0124] When the hVEGF165 specific siRNA were delivered through intratumoral administration repeatedly, the growth of MCF-7/VEGF 165 cell induced xenograft tumors were significantly inhibited. This inhibition was further validated when the tumor tissues were collected for testing hVEGF mRNA expression which resulted in significant down regulation based on RT-PCR analysis. Using the same hVEGF165 specific siRNA, we are also able to demonstrate tumor growth inhibition with head and neck squamous cell carcinoma (HNSCC, 1483 cell) xenograft model, through 5 repeated administrations intratumorally with seven day interval. Each injection requires 10 µg of the specific siRNA in 15 to 30 µL of RNase free aqueous solution.

Example 3

25 Basepair Blunted Double-Stranded siRNA is More Potent than Regular 19 Basepair siRNA with 3'-Overhangs in Silencing Target Gene Expression

[0125] In one of in vitro siRNA transfection studies, the MCF/165 breast cancer cells overexpressing human VEGF

(hVEGF) were transfected with either 25 basepair blunted double-stranded siRNA (hVEGF-25-siRNA-a, hVEGF-25-siRNA-b, hVEGF-25-siRNA-c, Luc-25-siRNA) or 19 basepair siRNA with 3' overhangs (hVEGF-siRNA-a, GFP-siRNA-a) using an electroporation mediated transfection method. 4×10^6 MCF7/165 cells were resuspended in 200 μ l of siPORT siRNA Electroporation Buffer (Ambion) mixed with 5 μ g siRNA and then subjected to electroporation treatment using an Electro Square Porator ECM830 (BTX, Fisher Scientific). The parameters for the electroporation are: voltage 500 v; duration of pulse 60 us; pulse number 2; pulse interval 1 second. The transfected cells were seeded to 24-well plate at a density of 5×10^4 cell/well and cultured at 37° C. incubator with 5% CO₂. At 24 hours post transfection, the culture media from each well were harvested and the concentration of human VEGF protein in the media was measured using a commercial hVEGF EKISA kit (R&D Systems).

[0126] We observed a significant stronger hVEGF target gene inhibition in 25 basepair hVEGF-siRNA treated cells than that in regular 19 basepair hVEGF siRNA treated cells. There was a more than 75% reduction of the secreted hVEGF protein in the culture media of cells transfected with 25 basepair blunted double-stranded hVEGF-25-siRNA molecules at 24 hours post siRNA transfection, compared to an about 60% hVEGF protein reduction observed in cells transfected with regular 19 basepair hVEGF-siRNA-a. Both non-specific sequence control 25 basepair blunted double-stranded siRNA (Luc-25-siRNA) or regular GFP-siRNA with 3'-overhangs did not affect VEGF expression under the same siRNA transfection condition (FIG. 7).

Example 4

25 basepair Blunted Double-Stranded siRNA Mediated an Elongated Target Gene Silencing

[0127] In another one of in vitro siRNA transfection studies, the MCF/165 breast cancer cells overexpressing human VEGF (hVEGF) were transfected with either 25 basepair blunted double-stranded siRNA (hVEGF-25-siRNA-a, hVEGF-25-siRNA-b, hVEGF-25-siRNA-c, Luc-25-siRNA) or 19 basepair siRNA with 3' overhangs (hVEGF-siRNA-a, GFP-siRNA-a) using an electroporation mediated transfection method. 4×10^6 MCF7/165 cells were resuspended in 200 μ l of siPORT siRNA Electroporation Buffer (Ambion) mixed with 2 μ g or 5 μ g siRNA and then subjected to electroporation treatment using an Electro Square Porator ECM830 (BTX, Fisher Scientific). The parameters for the electroporation are: voltage 500 v; duration of pulse 60 us; pulse number 2; pulse interval 1 second. The transfected cells were seeded to 24-well plate at a density of 5×10^4 cell/well and cultured at 37° C. incubator with 5% CO₂. At 24, 48, 72, 96, and 120 hours post transfection, the culture media from each well were harvested and replaced with fresh culture media. The concentration of human VEGF protein in the media harvested at various time points was measured using a commercial hVEGF EKISA kit (R&D). The siRNA mediated hVEGF knockdown was normalized with the hVEGF protein levels measured in cells treated with respective non-specific control siRNA.

[0128] We observed a significant stronger and elongated hVEGF target gene inhibition in 25 basepair hVEGF-siRNA treated cells than that in regular 19 basepair hVEGF siRNA treated cells at every time points tested. At 120 hours post siRNA treatment, there was still a more than 60% reduction of

the secreted hVEGF protein in the culture media of cells transfected with 5 μ g of 25 basepair blunted double-stranded hVEGF-25-siRNA molecules, compared to an less than 20% hVEGF protein reduction observed in cells transfected with 5 μ g of regular 19 basepair hVEGF-siRNA-a (FIG. 8). We also observed a dose-dependent siRNA mediated hVEGF gene inhibition for the 25 basepair blunted double-stranded hVEGF-25-siRNA molecules.

[0129] Base on our observation, the 25 basepair blunted double-stranded hVEGF-25-siRNA molecules not only gives a stronger target VEGF gene inhibition, but also results in a significant longer duration of effective target VEGF gene inhibition. For example, compared to only 48 hours of more than 60% hVEGF protein reduction achieved using 5 μ g of regular 19 basepair hVEGF siRNA, at least 120 hours of more 60% reduction of protein was achieved using 25 basepair blunted double-stranded hVEGF siRNA. Therefore, the 25 basepair blunted double-stranded siRNA are more potent target gene inhibitor that can lead to more significant therapeutic efficacy.

[0130] Additional information include (1) Luc-25-siRNA has the sequence of (sense strand 5'-rGGAACCGCUG-GAGAGCAACUGCAUA-3' (SEQ ID NO: 249) and antisense strand 5'-rCCUUGGCGACCUCUCGUUGACGUAU-3') (SEQ ID NO: 250); (2) hVEGF-siRNA-a has the sequence of (sense strand, 5'-rUCGAGACCCUG-GUGGACAuTT-3' (SEQ ID NO: 251) and antisense strand, 5'-rAUGUCCACCAGGGUCUCGAdTT-3') (SEQ ID NO: 252); (3) GFP-siRNA has the sequence of (sense strand, 5'-rGCUGACCCUGAAGUUAUCdTT-3' (SEQ ID NO: 253) and (antisense strand, 5'-rGAUGAACUUCAGGGUCAGCdTT-3') (SEQ ID NO: 254); (4) hVEGFR2-25-siRNA-c: 5'-r(CCAAGUGAUUGAAGCAGAUGCCUUU)-3' (SEQ ID NO: 255).

Example 5

25 Base Pair Blunted Double-Stranded siRNA Efficiently Knockdown Human VEGFR1 and VEGFR2 Expression In Vitro

[0131] Three 25 base pair blunted double-stranded siRNA specifically targeting human VEGFR1 were transfected in HUVEC cells through electroporation. The membrane bound VEGFR1 and free extra cellular fragment of VEGFR1 were measured 96 hours post transfection, through ELISA assay. The VEGFR1-siRNA duplex a demonstrated the strongest VEGFR1 silencing activity, compared to two other siRNA, for both cell lysate protein and the free fragments in the cell culture supernatant solution (FIGS. 9 and 10). When using the same approach to evaluate three 25 base pair blunted double-stranded siRNA targeting human VEGFR2 gene, we found all three duplexes exhibiting potent silencing activity at both 24 hours and 72 hours post transfection time points (FIG. 11).

Example 6

25 Basepair Blunted Double-Stranded siRNA Mediated Strong Anti-Tumor Efficacy in MDA-MB-435 Xenograft Tumor

[0132] 25 basepair long siRNA duplex targeting human VEGF 165 gene sequence demonstrated strong activity of tumor growth inhibition, after three repeated intratumoral administrations of 10 μ g every 5 days (FIG. 12), using a

MDA-MB-435 breast carcinoma cell line known having high expression of VEGF and bFGF proteins.

Example 7

siRNA Cocktail Targeting VEGF, VEGFR1 and VEGFR1 Genes is More Potent than Any of the Individual siRNA Targeting Only on of Those Genes in HSV Induced Ocular Neovascularization Model

[0133] Recently, we demonstrated that the siRNA cocktail containing siRNA targeting VEGF, VEGFR1 and VEGFR2 can achieve more potent antiangiogenesis efficacy than only targeting one of those genes, at the same dosage, in the animal disease model. Using this approach, we are able to efficiently block the VEGF pathway which plays the key role for pathological angiogenesis (FIG. 13).

[0134] HSV DNA that contains abundant potentially bioactive CpG-containing motifs can induce the potent angiogenesis factor vascular endothelial growth factor (VEGF) and that neutralization of VEGF with antibody minimized HSV-induced angiogenesis. A convenient model was also established in which bioactive CpG-containing oligodeoxynucleotides (ODNs) were also shown to induce neovascularization via the induction of VEGF. This model is used in the present study to evaluate the therapeutic potential of RNA interference (RNAi) to suppress VEGF expression and responsiveness.

Material and Methods

Reagents

[0135] Phosphorothioate ODNs were kindly provided by Dennis M. Klinman (Biologics Evaluation and Research, Food and Drug Administration, Washington, DC). The sequences of stimulatory ODNs used in this study were: 1466, TCAACGTTGA (SEQ ID NO: 256), and 1555, GCTAGACGTTAGCGT (SEQ ID NO: 257). Subsequent studies were performed using an equimolar mixture of ODNs 1466 and 1555. Molecular Design of Gene Targets and siRNA Three mVEGF pathway factors, mVEGF A and two mVEGF receptors (mVEGFR1 and mVEGFR2), were targeted by RNAi. For each gene target, two target sequences were assigned at different locations on the same mRNA. siRNA were designed correspondent to the above target sequences. These siRNA were designed according to the guideline proposed by Tuschl. 14,15 The designed siRNA (duplexes of sense and anti-sense strands) were synthesized by Qiagen (Valencia, Calif.). All siRNA were 21-nucleotides long doublestranded RNA oligos with a two nucleotide (TT) overhang at the 3'-end. The targeted sequences of mVEGFA were (a) AAGCCGTCCTGTGTGCCGCTG (SEQ ID NO: 258) and (b) AACGATGAAGCCCTGGAGTGC (SEQ ID NO: 259).

[0136] The targeted sequences of mVEGFR1 were:

[0137] (a) AAGTTAAAAGTGCCTGAACTG (SEQ ID NO: 260) and

[0138] (b) AAGCAGGCCAGACTCTCTTTC (SEQ ID NO: 261). The targeted sequences of mVEGFR2 were (a) AAGCTCAGCACACAGAAAGAC (SEQ ID NO: 262) and (b) ATGCGGCGGTGGTGACAGTA (SEQ ID NO: 263). The synthesis of unrelated siRNA controls, two target sequences each for LacZ and firefly luciferase were used. They were LacZ (a) AACAGTTGCGCAGCCTGAATG (SEQ ID NO: 264) and (b) AACTTAATCGCCTTGCAGCAC (SEQ ID NO: 265), Luc (a) AAGCTATGAAAC-

GATATGGGC (SEQ ID NO: 266) and (b) AACCGCTG-GAGAGCAACTGCA (SEQ ID NO: 267). Subsequent studies were conducted using an equimolar mixture of a and b for individual siRNA. Mice Female BALB/c mice (H-2d), 5 to 6 weeks old, were purchased from Harlan Sprague-Dawley (Indianapolis, Ind.) and housed conventionally. All investigations followed guidelines of the Committee on the Care of Laboratory Animals Resources, Commission of Life Sciences, National Research Council. The animal facilities of the University of Tennessee (Knoxville, Tenn.) are fully accredited by the American Association of Laboratory Animal Care.

[0139] Virus HSV-I strain RE (kindly provided by Dr. Robert Lausch, University of Alabama, Mobile, Ala.) was used in all of the procedures. Virus was grown in Vero cell monolayers (catalog no. CCL81; American Type Culture Collection, Manassas, Va.), titrated, and stored in aliquots at -80°C . until used.

In Vitro Efficacy of siRNA

[0140] To test the efficacy of RNAi in vitro, the following cell lines were used. RAW264.7 gamma NO cells (CRL-2278, ATCC) were used to test the efficiency of siVEGFA specific knockdown of VEGFA gene that is spontaneously expressed in these cells. The cells were plated in a six-well plate in RPMI with 10% fetal bovine serum overnight at 37°C . in 5% CO_2 . One day after cell plating, the cells were transfected with different concentrations of siVEGFA or siLuc (at 0, 0.1, 0.5, 1.0, or 2.0 $\mu\text{g}/2\text{ ml/well}$, respectively) using Lipofectamine 2000 (Invitrogen, Carlsbad, Calif.). Twenty-four hours later RNA from these cells was extracted for reverse transcriptase-polymerase RNA Extraction and RT-PCR). SVR cells (CRL-2280, ATCC) were used to test the efficiency of siVEGFR1-specific knockdown of VEGFR1 gene that is constitutively expressed on these cells. The cells were plated in a six-well plate in Dulbecco's modified Eagle's medium with 5% fetal bovine serum overnight at 37°C . in 5% CO_2 . One day after cell plating, the cells were transfected with different concentrations of siVEGFR1 or siLuc (at 0, 0.1, 0.5, or 1.0 $\mu\text{g}/2\text{ ml/well}$, respectively) using Lipofectamine 2000. Forty-eight hours later RNA from these cells was extracted for RSPCR to detect VEGFR1 (see RNA Extraction and NATemplate-Specific PCR) (RS-PCR). The 293 cells (CRL-1573, ATCC) were used to transfect with mVEGFR2-expressing plasmid for the detection of knockdown of exogenous mVEGFR2. The cells were plated in a six-well plate in Dulbecco's modified Eagle's medium with 5% fetal bovine serum overnight at 37°C . in 5% CO_2 . One day after cell plating, the cells were cotransfected with plasmid pCI-VEGFR2 (0.2 $\mu\text{g}/2\text{ ml/well}$) and siVEGFR2 (a, b, a & b), or siLuc (0, 0.1, 0.5, or 1.0 $\mu\text{g}/\text{well}$, respectively) using Lipofectamine 2000. Forty eight hours later RNA from these cells was extracted for RS-PCR to detect VEGFR2.

Corneal Micropocket Assay

[0141] The corneal micropocket assay used in this study observed the general protocol of Kenyon and colleagues. Pellets for insertion into the cornea were made by combining known amounts of CpG ODNs, sucralfate (10 mg, Bulch Meditec, Vaerlose, Denmark), and hydon polymer in ethanol (120 mg/1 ml ethanol; Interferon Sciences, New Brunswick, N.J.), and applying the mixture to a 15 mm^2 piece of synthetic mesh (Sefar America, Inc., Kansas City, Mo.). The mixture was allowed to air dry and fibers of the mesh were pulled apart, yielding pellets containing CpG ODNs. The micro-

pocket was made under a stereomicroscope (Leica Microsystems, Wetzlar, Germany) (four eyes per group) and pellets containing CpG ODNs were inserted into the micropocket. Angiogenesis was evaluated at days 4 and 7 after pellet implantation by using calipers (Biomedical Research Instruments, Rockville, Md.) with a stereomicroscope. The length of the neovessels originated from the limbal vessel ring toward the center of the cornea and the width of the neovessels presented in clock hours were measured. Each clock hour is equal to 30° at the circumstance. The angiogenic area was calculated according to the formula for an ellipse.

In Vivo Delivery of siRNA

[0142] Limbus was monitored at both day 4 and 7 after pellet implantation. Significant inhibition of corneal neovascularization resulted with all three test siRNA compared to those given control siLacZ at day 4 after pellet implantation ($p < 0.05$). The combination of the three tested siRNA was the most effective inhibitor, providing an approximately 60% reduction in neovascularization ($p < 0.01$).

Inhibition of CpG-Induced Neovascularization by Systemic Delivery of siRNA Targeting VEGF-Pathway Genes.

[0143] To test the anti-angiogenic effect of targeted individual siRNA and the efficiency of systemic siRNA delivery, mice with CpG ODN-containing micropockets were given a single dose i.v. of 40 µg siRNA containing either siVEGFA, siVEGFR1, siVEGFR2, a mix of the three, or control siLacZ 6 and 24 h post pellet implantation. In these experiments a polymer ("Targetran") was used that was shown in previous studies on tumor angiogenesis to facilitate extravascular delivery of siRNA. At day 4 and 7 after pellet implantation, the extent of angiogenesis was measured. All reagents used individually induced significant inhibition of neovascularization compared to the siLacZ treated group at day 4 after pellet implantation ($p < 0.05$). As observed with local administration, the mix of the three test reagents provided the most effective inhibition (average 40% inhibition, $p < 0.01$). In additional experiments, the function of the polymer vehicle was evaluated by comparing the anti-neovascularization activity of the test mix suspended in polymer or given in PBS. These experiments revealed that the use of the polymer vehicle resulted in more effective anti 16 neovascularization than was evident when the PBS vehicle was used although result was only significant at the early test period ($p < 0.05$). The results demonstrate that ocular neovascularization can be controlled by the i.v. administration of siRNA that target the VEGF system genes and that the use of the RGD-mediated dsRNA nanoparticle delivery enhanced the efficacy of the therapeutic effect.

[0144] To determine the efficient anti-angiogenic dose of siRNA in systemic delivery, mice with CpG ODN-containing micropockets were given a single dose i.v. of 10, 20, 40, 80 µg siRNA containing a mix of the siVEGFA, siVEGFR1 and siVEGFR2, or control siLuc with TargeTran vehicle at 6 and 24 h post pellet implantation. A administration of siRNA inhibited CpG induced angiogenesis in a dose dependent manner.

Therapeutic Application of siRNA Against VEGF-Pathway Genes in the HSK Model.

[0145] Previous studies have shown that VEGF is the critical angiogenic factor for induction of HSV specific angiogenesis in the HSK model. To evaluate whether administration of siRNA targeting VEGF-pathway genes inhibits the development of HSK, the corneas of mice were scarified and infected with 1-105 HSV-1 RE. Then mice were given a single dose of

10 µg (subconjunctival injection for local delivery) or 40 µg (tail vein injection for systemic delivery) mix of siRNA (an equimolar mixture of siVEGFA, siVEGFR1, and siVEGFR2) with polymer vehicle at day 1 and 3 after virus infection. As shown in FIG. 4, the angiogenesis and severity of HSK was significantly reduced in mice treated with siRNA targeting VEGF-pathway genes either locally or systemically compared to animals treated with siLuc control ($p < 0.05$). Whilst 80% of siLuc control treated eyes developed clinically evident lesions (score 2 or greater at day 10 p.i.), only 42% (local delivery) or 50% (systemic delivery) of eyes treated with siRNA targeting VEGF-pathway genes developed such lesions. In addition by day 10 p.i., the angiogenesis score was greater than 6 in 9 of 12 control eyes, but only in 5 of 12 eyes of mice treated with siRNA against VEGF-pathway genes by either local or systemic delivery. Taken together these results show that administration of siRNA against VEGF-pathway genes reduced development of HSK via inhibition of angiogenesis.

Example 8

siRNA Cocktail Targeting VEGF, VEGFR1 and VEGFR1 Genes Very Potent Anti-Angiogenesis Agent in ROP Ocular Neovascularization Model

[0146] Neovascularization in the eye is associated with various disorders, often causing severe loss of vision and eventually blindness. Among these disorders, diabetic retinopathy (DR), age-related macular degeneration (AMD), retinal vein occlusion (RVO) and retinopathy of prematurity (ROP) are prevalent. The imbalance of stimulatory factors and inhibitory factors result in NV (NV) and vascular endothelial growth factor (VEGF) is the most important factor of stimulatory factors which cause vascular permeability, dilation and endothelial cell migration, proliferation^[1].

[0147] RNA interference (RNAi) is the process of sequence-specific post-transcriptional gene silencing in a wide range of organisms, initiated by double stranded RNA (dsRNA) that is homologous in sequence to the targeting gene. In this example, small interference dsRNA oligonucleotides (siRNA) targeting VEGF path ways were used to inhibition NV induced by oxygen-induced retinopathy.

Materials and Methods

[0148] The Design and Synthesis of siRNA

[0149] siRNA was kindly provided by Intradigm Corporation. Three mVEGF pathway factors, mVEGF A and two mVEGF receptors (mVEGFR1 and mVEGFR2) were targeted by RNAi and called siMix. The siRNA targeting luciferase was called siLuc as control.

Mouse Model of Oxygen Induced Retinal Neovascularization

[0150] The model we used (Smith et al.^[2]) imitates retinopathy of prematurity. On postnatal day seven (P7) the mice and their nursing mother were placed in an airtight incubator (own production) ventilated by a mixture of oxygen and air to a final oxygen fraction of 75%±2%. Oxygen levels were

checked at least 3 times a day. On P12 the mice were returned to room air. On P17 the animals were sacrificed.

Mice

[0151] The C57BL/6 mice were purchased from center of experimental animal of Guangzhou Medical college and Guangzhou University of traditional Chinese Medicine. All investigations followed guidelines of the Committee on the Care of Laboratory Animals Resources, Commission of Life Science, National Research Council.

In Vivo Delivery of siRNA

[0152] For local delivery, siMix (4 μ g/2 μ l per eye) was delivered subconjunctivally or intravitreally in the left eye and siLuc (4 μ g/2 μ l per eye) in the right eye by a 32-gauge Hamilton syringe (Hamilton Co, Revo, Nev.) on P12 and P13 under deep anesthesia. For systemic administration, siRNA (15 μ g/50 μ l per mouse) was mixed with the RGD-PEG-PEI polymer conjugate preparation (TargeTran) and delivered intraperitoneally on P12 and P13.

Retinal Angiography^[4]

[0153] On P17 the animals were sacrificed by cardiac perfusion with a solution of 50 mg/ml fluorescein-labeled dextran in sodium chloride as described previously. Both eyes were enucleated and fixed for 0.5-1 h in 10% buffered formaldehyde at room temperature. The anterior segment was cut off and the neurosensory retina carefully removed. The retina was cut radially and flat mounted in glycerin, photoreceptors facing downward. A cover slip was placed over the retina and sealed with nail polish. Retinal whole mounts were examined by fluorescence microscopy. The areas of retinal NV were measured by soft Image-Pro Plus (Media Cybernetics, USA).

Cryosection^[5]

[0154] Eyes were removed and frozen in optimal cutting temperature embedding compound (Miles Diagnostics). Ocular frozen sections (10 μ m) were histochemically stained with biotinylated GSA. Slides were incubated in methanol/H₂O₂ for 10 min at 4° C., washed with 0.05 M Tris-buffered saline (TBS), pH 7.6, and incubated for 30 min in 10% normal bovine serum. Slides were incubated 1 h at 37° C. with biotinylated GSA and after rinsing with 0.05 M TBS, they were incubated with avidin coupled to alkaline phosphatase (Vector Laboratories) for 45 min at room temperature. After being washed for 10 min with 0.05 M TBS, slides were incubated with diaminobenzidine to give a brown reaction product and were counterstained with eosin, mounted with Cytoseal. To perform quantitative assessments, 10 μ m serial sections were cut through entire eyes, and starting from the 1st section that contained iris and extending to the last section on the other side of the eye that contained iris, every tenth section was stained with GSA and total 15 sections were stained. GSA-stained sections were examined with a microscope, and images were digitized using a digital color video camera. Image-Pro Plus software (Media Cybernetics, Silver Spring, Md., USA) was used to delineate GSA-stained cells on the surface of the retina and their area was measured. For local injection the total measurements from each eye was used as a

single experimental value. For systemic injection the mean of both eyes of a mouse was used as a single experimental value.

RT-PCR

[0155] On P14 and P17 the mouse were sacrificed and total RNA was extracted from retinae with TRIzol reagent (Invitrogen, USA). Total RNA was quantified at a ratio of 260 to 280 nm and 2 μ g RNA was converted to cDNA via cDNA synthesis Kit (Fermentas, USA). RT-generated cDNAs encoding VEGF, VEGFR1, VEGFR2 and b-actin (acting as a control of RNA integrity and as an internal standard) were amplified with RT-PCR. Amplification of 1.5 μ l of cDNA was performed with 1 μ l sense and antisense and 2.5 U of Taq polymerase. The oligonucleotide primer sequences are: mVEGF (430 bp) \square forward 5'-GAT GTC TAC CAG CGA AGC TAC TGC CGT CCG-3' (SEQ ID NO: 268) \square reverse 5'-GAA CAT CGA TGA CAA GCT TAG GTA TCG ATA caa get gcc teg cct tg-3' (SEQ ID NO: 269) \square mVEGF1 (404 bp) \square forward 5'-GTCAGC TGC TGG GAC ACC GCG GTC TTG CCT-3' (SEQ ID NO: 270) \square reverse 5'-GAA CAT CGA TGA CAA GCT TAG GTA TCG ATA tag art gaa gat tec gc-3' (SEQ ID NO: 271) \square mVEGFR2 (485 bp) \square forward 5'-TGG CTG GTC AAA CAG CTC ATC-3' (SEQ ID NO: 272) \square reverse 5'-CTC ATC CAA GGG CAA TTC ATC-3' (SEQ ID NO: 273) \square \square -actin (232 bp) \square forward 5'-CAT TGT GAT GGA CTC CGG AGA CGG-3' (SEQ ID NO: 274) \square reverse 5'-CAT CTC CTG CTG AAG TCT AGA GC-3' (SEQ ID NO: 275).

ELISA for VEGF and VEGF1, VEGFR2

[0156] On P14 and P17 the mouse were sacrificed and retinae were extracted quickly on ice. Samples were homogenized in a cell lysis buffer (Mammalian cell lysis Kit, Biotechnology Department Bio Basid Inc, Canada) and were subsequently centrifuged at 12,000 rpm for 30 min. The supernatants were analyzed for protein concentrations via the BCA protein quantitative analysis Kit (Shenery Biocolor Bioscience & Technology Company, China). Samples were diluted to a final concentration of 1 mg/ml. Levels of VEGF, VEGFR1, and VEGFR2 were determined using the Quantikine M Murine VEGF, sVEGFR1, and sVEGFR2 Immunoassay Kits, respectively (R&D Systems Inc., Minneapolis, Minn.). Six to 12 tissue samples were analyzed for each group and time point.

Results and Discussion

[0157] Fluorescein Angiographic Assessment of the Effect of siRNA on Retinal NV

[0158] The retinae of the normal P17 mouse had both superficial and deep vascular layers (joined by connecting vessels) that extended from the optic nerve to the periphery. Retinae from P17 control mouse exposed to hyperoxia contained many neo vascular tufts (high fluorescein) extending from the surface of the retina at the junction between the perfused and nonperfused retina. After subconjunctival injection of siLuc or siMix the retinae also had many NV, the areas had no obvious differences. Whereas the areas of NV after intravitreal and intraperitoneal injection of siMix were significant less than the control.

Assessment of Effects of siRNA by Histologic Quantitation of Retinal NV

[0159] Retinal NV was also assessed histologically by measured the GSA-positive cells (neovascularization) ante-

rior to the internal limiting membrane (ILM). Retinas of P17 normoxic mice had superficial, middle, deep vessels and contained no endothelial cells anterior to the ILM. Retinas of P17 mice subjected to hyperoxia contained multiple neovascular tufts on the surface with some extending into the vitreous. Retinas of mice treated with subconjunctival injection of siLuc or siMix had no difference with the areas of NV. The areas of NV after intravitreal and intraperitoneal injection of siMix were obviously reduced compared to siLuc injection. Decreased Level of VEGF, VEGFR1 VEGFR2 mRNA After Intravitreal and Intraperitoneal Injection of siRNA

[0160] To address whether treatment of siRNA against VEGF pathway genes reduces the levels the VEGF, VEGFR1, VEGFR2 mRNA, retinas were collected on P14 and P17 after intravitreal and intraperitoneal injection of siRNA. The mRNA levels were measured by RT-PCR. The expression of VEGF, VEGFR1, VEGFR2 mRNA were reduced in the retinas treated with siMix against VEGF pathway genes compared to the control retinas treated with siLuc ($p < 0.05$).

Decreased Expression of VEGF, VEGFR1, VEGFR2 Protein Levels After Intravitreal and Intraperitoneal Application of siRNA

[0161] To evaluate whether treatment of siRNA targeting VEGF pathway genes diminishes the production of VEGF, VEGFR1, VEGFR2 protein, ELISA was used to measure VEGF, VEGFR1, VEGFR2 protein in siRNA-treated retinas on P14 and P17. As shown in table 1 and 2, VEGF, VEGFR1, VEGFR2 protein levels were lower in those that received siMix compared controls given with siLuc.

[0162] Imbalance in the demand and supply of oxygen and nutrients is the initiating factor of neovascularization which induces upregulation of VEGF. VEGF is not only the potent mitogenic factor for endothelial cells; it also induces vascular permeability and dilation. These biological activities are mediated by binding VEGF to high-affinity transmembrane autophosphorylating tyrosine kinase receptors^[5]. Three distinct VEGF receptors have been identified, namely VEGFR1 (fms-line tyrosine kinase-1 or Flt-1), VEGFR2 (kinase insert domain-containing receptor or KDR) and VEGFR3 (fms-line tyrosine kinase-4 or Flt-4). VEGFR1 and VEGFR2 are predominantly expressed on vascular, VEGFR3 on lymphatic endothelium^[6]. Increased intravitreal and intraretinal levels of VEGF are associated with retinal neovascularization not only in animal model but also in patients with ischemic retinopathy. These data suggest that VEGF signaling is a good target for treatment of retinal neovascularization^[5].

[0163] Oxygen-induced retinopathy in mice is widely recognized in the world. Eric AP et al^[7] had reported that the mRNA levels of VEGF increased dramatically between 6 to 12 h after hypoxia and remained elevated for several days and then decreased toward baseline with regression of retinopathy. So we delivered siRNA on P12 and P13 in order to interfere the synthesis of VEGF before upregulation. The results of whole mount and cryosection had proved that the NV areas of retinas treated with intravitreal and intraperitoneal injection of siMix were less than injection of siLuc. That is siMix can obviously inhibit retinal NV. To explore the mechanism the retinal VEGF, VEGFR1 and VEGFR2 expressions on P14 and P17 were examined and the mRNA and protein levels were lower than the siLuc controls. These suggested it is through inhibiting VEGF pathway genes that siMix inhibit retinal NV.

[0164] However there is no distinct difference after subconjunctival injection, so it is concluded that it hadn't achieved the

effective concentration in local. After intravitreal injection there was high concentration of siRNA in vitreous and siRNA was transfected into retinal neovascular endothelial cells. But this injection can result in intraocular hemorrhage and endophthalmitis et al. Intraperitoneal injection is relative safe and siRNA can be absorbed into blood through abundant celiac capillary. But siRNA are easily trapped in the non-specific organs including liver, lungs, and spleen. So how to enhance the transfecting efficiency of siRNA is very important. The vehicle TargeTran is composed of polyethylene imine (PEI), polyethylene glycol (PEG) and arginine-glycine-aspartate peptide sequences (RGD). PEI binds to negative charges in phosphates of the siRNA. RGD motif has been identified as an integrin ligand of activated endothelial cells. Endothelial cells express a number of different integrins and integrin $\alpha v \beta 3$ and $\alpha 5 \beta 1$ have been shown to be important during angiogenesis. Both integrins are the receptor for matrix proteins with an exposed RGD tripeptide moiety and are most prominent on activated endothelial cells during angiogenesis. Thus the application of Targetran can improve the transfecting efficiency of siRNA. Kim B et al^[8] had reported the subconjunctival and intravenous injection of siMix can inhibit the CPG induced the corneal neovascularization in mice. It may also apply to other neovascular diseases.

Example 9

Ligand Directed Nanoparticle siRNA can be Specifically Delivery into Tumor with Systemic Administration

[0165] This self-assembled siRNA nanoparticle has demonstrated the neovasculature targeting property when they were systemically administrated through IV injection. The tumor specific targeting capability of this siRNA nanoparticle system has been revealed with a Luciferase report gene system.

Example 10

Ligand Directed Nanoparticle siRNA is Potent Anti-Tumor Agent Validated in Mouse Syngenic Model (Neuroglioblastoma)

[0166] Tumor targeted delivery of siRNA using the targeted nanoparticle system from iv administration is demonstrated using fluorescently labeled siRNA packaged in RGD-PEG-PEI nanoparticle.

Mouse Tumor Model

[0167] Female nude mice (6-8 weeks of age) were obtained from Taconic (Germantown, N.Y.), kept in filter-topped cages with standard rodent chow and water available ad libitum, and a 12 h light/dark cycle. Experiments were performed according to national regulations and approved by the local animal experiments ethical committee. Subcutaneous N2A tumors were induced by inoculation of 1×10^6 N2A-cells in the flank of the mice. At a tumor volume of approximately 0.5-1 cm³, mice received nanoplexes or free siRNA by i.v. injection of a solution of 0.2 ml via the tail vein. 40 μ g fluorescently-labeled siRNA was injected in the free form or as PEI- or RGD-PEG-PEI-nanoparticle. One hr after injection, tissues were dissected and examined with a dissection microscope fitted for fluorescence. Microscopic examination of tissues was performed with an Olympus SZX12 fluorescence microscope

equipped with digital camera and connected to a PC running MagnaFire 2.0 camera software (Optronics, Goleta, Calif.). Pictures were taken at equal exposure times for each tissue.

[0168] Result: Intravenous injection by tail vein of free fluorescent labeled siRNA did not show any significant accumulation in lung, liver or tumor tissue. SiRNA delivered using PEI-nanoparticle showed highest accumulation in the lung tissue followed by liver and tumor. SiRNA delivered using RGD-PEG-PEI nanoparticle showed highest level of accumulation in tumor tissue. There was a considerable reduction in lung accumulation compared to PEI-nanoparticles and very little accumulation in liver. This experiment demonstrates tumor targeted delivery of siRNA using RGD-PEG-PEI nanoparticle.

[0169] RNAi mediated inhibition of tumor angiogenesis and tumor growth were studied using tumor targeted nanoparticle formulation containing siRNA targeted against VEGFR2. Subcutaneous tumor bearing mice were treated by intravenous injection of the nanoparticle formulation at a dose of 40 μg siRNA per injection. Injections were repeated every third day and growth of the tumor was evaluated and compared with animals treated with control formulations. Inhibition of the angiogenesis was evaluated at the end of the experiment.

Mouse Tumor Model

[0170] Female nude mice (6-8 weeks of age) were obtained from Taconic (Germantown, N.Y.), kept in filter-topped cages with standard rodent chow and water available ad libitum, and a 12 h light/dark cycle. Experiments were performed according to national regulations and approved by the local animal experiments ethical committee. Subcutaneous N2A tumors were induced by inoculation of 1×10^6 N2A-cells in the flank of the mice. At a tumor volume of approximately 0.5-1 cm^3 , mice received nanoplexes or free siRNA by i.v. injection of a solution of 0.2 ml via the tail vein. The nanoparticle formulations containing siRNA were prepared by simple mixing of siRNA solutions with polymer solution at given N/P ratio.

[0171] For the tumor growth inhibition studies, the experiment was started when the tumors became palpable, at 7 days after inoculation of the tumor cells. Treatment consisted of 40 μg siRNA per mouse in RPP-nanoplexes every 3 days intravenously via the tail vein. Tumor growth was measured at regular intervals using a digital caliper by an observer blinded to treatment allocation. Each measurement consisted of tumor diameter in two directions approximately 90 degrees apart. Tumor volume was calculated as, $0.52 \times \text{longest diameter} \times \text{shortest diameter}^2$. At the end of the experiment, the animals were sacrificed and tumor tissue and surrounding skin was excised and put on a microscopy glass slide. Tissue examination for vascularization and angiogenesis was performed by microscopy using the Olympus microscope and camera equipment described above for fluorescent tissue measurements. Tissue was trans-illuminated to visualize blood vessels in the skin and a digital image was taken and stored as described above. Tissue was snap frozen immediately thereafter for Western blotting.

[0172] Result: Significant tumor growth inhibition was observed for animals treated with nanoparticle formulations containing VEGFR2 siRNA. Animals treated with non-specific siRNA did not show any substantial tumor growth inhibition compared to untreated animals. Significant inhibition of blood vessel growth was observed around the tumor tissue indicating the inhibition of angiogenesis in the VEGFR2-

siRNA treated mice. Mice treated with control siRNA showed similar blood vessel growth as the untreated animals. Western blot analysis of the tumor lysate collected from animals of different treatment groups showed substantial reduction of VEGFR2 in VEGFR2-siRNA treated animals whereas no reduction in VEGFR2 was observed in control siRNA treated animals. These experiments clearly demonstrate the delivery of siRNA into tumor tissue from intravenous administration. Effectiveness of VEGFR2 siRNA packaged in the RGD-PEG-PEI nanoparticle to inhibit tumor growth in renal cell carcinoma model was studied using a 786-O xenograft tumor model. An experimental procedure similar to example 7 was used in this study. Briefly, female nude mice (6-8 weeks of age) were obtained from Taconic (Germantown, N.Y.). Subcutaneous 786-O tumors were induced by inoculation of 5×10^6 786-O cells in the flank of the mice. At a tumor volume reached approximately 100 mm^3 , treatment was started by i.v. injection of a solution of VEGFR2-siRNA in RGD-PEG-PEI nanoparticle via the tail vein. Control treatment groups received nanoparticles containing non-specific siRNA or saline. Treatment was repeated for several days with injections every three days. Tumor volume was measured once every three days as described in example 7. Result: Significant tumor growth inhibition was observed for animals treated with VEGFR2-siRNA nanoparticle formulation. No significant tumor growth inhibition was observed for animals treated with control siRNA nanoparticles. This experiment demonstrates that the VEGFR2 siRNA delivered by tumor targeted nanoparticle formulation can achieve tumor growth inhibition.

[0173] The ligand directed siRNA nanoparticles were systemically administrated into a C57 mouse model with subcutaneous inoculation of N2A glioblastoma cells for evaluation of the impact of VEGF knockdown on the tumor angiogenesis activity. Female nude mice (6-8 weeks of age) were obtained from Taconic (Germantown, N.Y.), kept in filter-topped cages with standard rodent chow and water available ad libitum, and a 12 h light/dark cycle. The experiments were performed according to national regulations and approved by the local animal experiments ethical committee. Subcutaneous N2A tumors were induced by inoculation of 1×10^6 N2A cells in the flank of the mice. At a tumor volume of 0.5-1 cm^3 , mice received nanoplexes or free siRNA by i.v. injection of a solution of 0.2 ml via the tail vein. The nanoplex solutions were prepared as above, at N/P ratio of 2. For tissue distribution experiments, 40 mg fluorescently labeled siRNA was injected in the free form or as P- or RPP-nanoplexes. One hour after injection, the tissues were dissected and examined with a dissection microscope fitted for fluorescence. Microscopic examination of tissues was performed with an Olympus SZX12 fluorescence microscope equipped with digital camera and connected to a PC running MagnaFire 2.0 camera software (Optronics, Goleta, Calif.). Pictures were taken at equal exposure times for each tissue.

[0174] In the co-delivery experiments, plasmid and siRNA were mixed in a 1:100 molar ratio, respectively (40 mg pLuc with 13 mg siRNA), and in the sequential delivery experiments 40 mg plasmid was delivered first, followed by 40 mg siRNA 2 h later (1:300 molar ratio). The tissues were dissected, weighed and put in ice-cold reporter lysis buffer (Promega) in magnetic beads containing 2 ml tubes (Q-Biogene, Carlsbad, Calif.), 24 h after injection of the nanoplexes. Tissues were homogenized with a Fastprep FP120 magnetic homogenizer (Q-Biogene) and samples were assayed for

reporter enzyme activity using the luciferase assay system (Promega) on a Monolight 2010 luminometer (Analytical Luminescence Laboratory). In the tumor growth inhibition studies, the experiment was started when the tumors became palpable at 7 days after inoculation of the tumor cells. Treatment consisted of 40 mg siRNA per mouse in RPP-nanoplexes every 3 days intravenously via the tail vein. Tumor growth was measured at regular intervals using a digital caliper by an observer blinded to treatment allocation. Each measurement consisted of tumor diameter in two directions of 90 apart. Tumor volume was calculated as: $0.52 \times \text{longest diameter} \times \text{shortest diameter}^2$.

[0175] At the end of the experiment, the animals were sacrificed and tumor tissue and surrounding skin was excised and put on a microscopy glass slide. Tissue examination for vascularization and angiogenesis was performed by microscopy using the Olympus microscope and camera equipment described above for fluorescent tissue measurements. Tissue was trans-illuminated to visualize blood vessels in the skin and a digital image was taken and stored as described above. Tissue was snap frozen immediately thereafter for western blotting.

[0176] Tumor growth inhibition by siRNA RPP-nanoplexes. Mice were inoculated with N2A tumor cells and left untreated or treated every 3 days by tail vein injection with RPP-nanoplexes with siLacZ or siVEGF R2 at a dose of 40 mg per mouse. Treatment was started at the time-point that the tumors became palpable (20 mm³). Only VEGF R2-sequence-specific siRNA inhibited tumor growth, whereas treatment with LacZ siRNA did not affect tumor growth rate as compared with untreated controls (n=5).

Example 11

Ligand Directed Nanoparticle siRNA is Potent Anti-Tumor Agent Validated in Xenograft Tumor Model (Renal Carcinoma)

[0177] The ligand directed siRNA nanoparticles were also tested in a mouse xenograft model with a renal carcinoma cell lines. The siRNA nanoparticles were manufactured with the same materials and same procedure, using the same route of delivery as that of Example 10. Five times of repeated deliveries with three day intervals using 2 mg/kg dosage were carried out with six animals per cohort. The significant tumor growth inhibition was observed.

Example 12

Ligand Directed Nanoparticle siRNA is Potent Anti-Tumor Agent Validated in Xenograft Tumor Model (Colorectal Carcinoma)

[0178] Using siRNA nanoparticle targeting VEGFR2 gene, we further tested the antitumor efficacy in mouse xenograft model with a human colorectal carcinoma cell line, DLD-1.

[0179] Material and Methods. Reagents: Avastin, monoclonal antibody against VEGF (25 mg/ml, Genetech); siRNA against VEGFR2, sequence (Appendix 1); siRNA against luciferase (Qiagen); Avertin made of 1.5 gram 2,2,2-Tribromoethanol and 1.5 ml t-amyl alcohol (Cat# T4840-2, Cat#24048-6, Aldrich) in 100ml distill water, St. Louis, Mo.]. Mice: athymus female nude mice, 5 to 6 weeks old, were purchased from TACONIC() and housed conventionally. All investigations followed guidelines of the Committee on the Care of Laboratory Animals Resources, Commission of Life

Sciences, National Research Council. The animal facility of Biomedical Research Institute in Rockville Md. is fully accredited by the American Association of Laboratory Animal Care. Cells: Colon carcinoma cell line, DLD-1 (CCL-221, ATCC) was grown in RPMI 1640 medium with 2 mM L-glutamine, 1.5 g/L sodium bicarbonate, 4.5 g/L glucose, 10 mM HEPES, and 1.0 mM sodium pyruvate, 10% fetal bovine serum.

Reagents

[0180] Avastin, monoclonal antibody against VEGF (25 mg/ml, Genetech); siRNA against VEGFR2, sequence (Appendix 1); siRNA against luciferase (Qiagen); Avertin made of 1.5 gram 2,2,2-Tribromoethanol and 1.5 ml t-amyl alcohol (Cat# T4840-2, Cat#24048-6, Aldrich) in 100 ml distill water, St. Louis, Mo.]. Mice: athymus female nude mice, 5 to 6 weeks old, were purchased from TACONIC() and housed conventionally. All investigations followed guidelines of the Committee on the Care of Laboratory Animals Resources, Commission of Life Sciences, National Research Council. The animal facility of Biomedical Research Institute in Rockville Md. is fully accredited by the American Association of Laboratory Animal Care. Cells: Colon carcinoma cell line, DLD-1 (CCL-221, ATCC) was grown in RPMI 1640 medium with 2 mM L-glutamine, 1.5 g/L sodium bicarbonate, 4.5 g/L glucose, 10 mM HEPES, and 1.0 mM sodium pyruvate, 10% fetal bovine serum. Procedure: 1) DLD-1 cells near confluence were harvested and resuspended in serum-free RPMI medium. 2) Mice were anaesthetized with Avertin, 0.4 ml/mouse i.p. 3) 100 million cells in 0.1 ml serum-free RPMI medium were injected into mice back s.c. on the left flank for establishment of xenograft tumor model. 4) 5 days after inoculation of tumor cells, sizes of growing tumors were measured with a caliper. Mice were then randomly grouped with 7 mice per group. Three different dosing regimen were applied: 1 mg/kg, 2 mg/kg and 4 mg/kg respectively. Although the high dose at 4 mg/kg represented the strongest anti-tumor activity, there is no significant difference amount three treatment groups. Using a different comparison, it has been found that the high dose of the siRNA nanoparticle at 4 mg/kg exhibited a stronger anti-tumor efficacy than that of 5 mg/kg Avastin treatment.

REFERENCES

- [0181]** 1. Peter A, Campochiaro. Retinal and choroidal neovascularization. *J Cell Physiol*, 2000, 184:301-10.
- [0182]** 2. Smith L E, Wesolowski E, Mclellan A, et al. Oxygen-induced retinopathy in the mouse. *Invest Ophthalmol Vis Sci*, 1994, 35:101-11.
- [0183]** 3. 3 D'Amato R, Wesolowski E, Smith L E. Microscopic visualization of the retina by angiography with high-molecular-weight fluorescein-labeled dextrans in the mouse. *Microvasc Res* 1993; 46:135-42.
- [0184]** 4. Jikui shen, Rebecca samul, Joelle Zimmer, et al. Deficiency of Neuropilin 2 Suppresses VEGF-Induced Retinal NV. *Molecular medicine*, 2004, 10:12-18.
- [0185]** 5. Unsoeld A S, Junker B, Mazitschek R, et al. Local injection of receptor tyrosine kinase inhibitor MAE 87 reduces retinal neovascularization in mice. *Mol Vis*, 2004, 10:468-75.
- [0186]** 6. Clauss M. Molecular biology of the VEGF and VEGF receptor family. *Semin Thromb Hemost*, 2000, 26:561-9.

- [0187] 7. Eric A P, Robert L A, Eliot D F, et al. Vascular endothelial growth factor/vascular permeability factor expression in a mouse model of retinal neovascularization. *Proc. Natl. Acad. Sci.*, 1995, 92:905-9.
- [0188] 8. Lu P. Y. et al., Keystone Symposia, Molecular Targets for Cancer Therapy, (2003, p 219. Xu J. et al., Gene Suppression. (2003).
- [0189] 9. Lu, Patrick et al., (2002), *Cancer Gene Therapy*, Vol. 10, Supplement 1, 011.
- [0190] 10. Lu, Patrick Y et al., (2003), *Current Opinion in Molecular Therapeutics*, 5(3):225-234.
- [0191] 11. Cogoni C. et al., (2000), *Genes Dev* 10:638-643.
- [0192] 12. Guru T., (2000), *Nature* 404:804-808.
- [0193] 13. Hammond S M et al., (2001), *Nature Rev Gen* 2:110-119.
- [0194] 14. Napoli C et al., (1990), *Plant Cell* 2:279-289.
- [0195] 15. Jorgensen R A et al., (1996), *Plant Mol Biol*, 31:957-973.
- [0196] 16. Ingelbrecht I et al., (1994), *Proc Natl Acad Sci USA*, 91:10502-10506.
- [0197] 17. Cogoni C et al., *EMBO J*, 15:3153-3163.
- [0198] 18. Palauqui J C et al., (1998), *EMBO J*, 16:4738-4745.
- [0199] 19. Guo S et al., (1995), *Cell*, 81:611-620.
- [0200] 20. Fire A et al., (1998), *Nature* 391:806-811.
- [0201] 21. Timmons L. et al., (1998), *Nature* 395:854.
- [0202] 22. Timmons L. et al., (2001), *Gene* 263:103-112.
- [0203] 23. Hunter C P, (2000), *Current Biology* 10:R137-R140.
- [0204] 24. Tabara H et al., (1998), *Science* 282:430-431.
- [0205] 25. Kamath R S et al., (2000), *Genome Biology* 2:2.1-2.10.
- [0206] 26. Grishok A et al., (2000), *Science* 287:2494-2497.
- [0207] 27. Sharp P A et al., (2000), *Science* 287:2431-2433.
- [0208] 28. Sharp P A, (2001), *Genes Dev* 15:485-490.
- [0209] 29. Kennerdell J R et al., (1998), *Cell* 95:1017-1026.
- [0210] 30. Kennerdell J R et al., (2000), *Nature Biotech* 18:896-898.
- [0211] 31. Dzitoyeva S et al., (2001), *Mol Psychiatry* 6(6): 665-670.
- [0212] 32. Worby C A et al., (2001), *Sci STKE Aug.* 14, 2001 (95):PL1.
- [0213] 33. Schmid A et al., (2002), *Trends Neurosci* 25(2): 71-74.)
- [0214] 34. Hamilton A. J. et al., (1999), *Science* 286:950-952.
- [0215] 35. Hammond S et al., (2000), *Nature*, 404:293-298.
- [0216] 36. Zamore P D et al., (2000), *Cell* 101:25-33.
- [0217] 37. Hutvagner G et al., (2002), *Curr Opin Genetics & Development* 12:225-232.
- [0218] 38. Bernstein E et al., (2001), *Nature* 409:363-366.
- [0219] 39. Nykanen A et al., (2001), *Cell* 107:309-321.
- [0220] 40. Lipardi C et al., (2001), *Cell* 107:297-307.
- [0221] 41. Ketting R F et al., (1999), *Cell* 99:133-141.
- [0222] 42. Grishok A et al., (2001), *Cell* 106:23-34.
- [0223] 43. Hutvagner G et al., (2001), *Science* 293(5531): 834-838.
- [0224] 44. Ketting R F et al., (2001), *Genes Dev* 15(20): 2654-2659.
- [0225] 45. Lagos-Quintana M et al., (2001), *Science* 294: 853-858.
- [0226] 46. Lau N C et al., (2001), *Science* 294:858-862.
- [0227] 47. Lee R C et al., (2001), *Science* 294:862-864.
- [0228] 48. Ruvkun G., (2001), *Science* 294:797-799.
- [0229] 49. Manche L et al., (1992), *Mol Cell. Biol.* 12:5238-5248.
- [0230] 50. Minks M A et al., (1979), *J. Biol. Chem.* 254: 10180-10183.
- [0231] 51. Yang S et al., (2001), *Mol Cell. Biol.* 21(22): 7807-7816.
- [0232] 52. Paddison P J et al., (2002), *Proc. Natl. Acad. Sci. USA* 99(3):1443-1448. Elbashir S M et al., (2001), *Genes Dev* 15(2):188-200.
- [0233] 53. Elbashir S M et al., (2001), *Nature* 411:494-498.
- [0234] 54. Caplen N J et al., (2001), *Proc. Natl. Acad. Sci USA* 98:9746-9747.
- [0235] 55. Holen T et al., (2002), *Nucleic Acids Research* 30(8):1757-1766.
- [0236] 56. Elbashir S M et al., (2001), *EMBO J* 20:6877-6888.
- [0237] 57. Jarvis R A et al., (2001), *TechNotes* 8(5):3-5.
- [0238] 58. Brown D et al., (2002), *TechNotes* 9(1):3-5.
- [0239] 59. Brummelkamp T R et al., (2002), *Science* 296: 550-553.
- [0240] 60. Lee N S et al., (2002), *Nature Biotechnol.* 20:500-505.
- [0241] 61. Miyagishi M et al., (2002), *Nature Biotechnol.* 20:497-500.
- [0242] 62. Paddison P J et al., (2002), *Genes & Dev.* 16:948-958.
- [0243] 63. Paul C P et al., (2002), *Nature Biotechnol.* 20:505-508.
- [0244] 64. Sui G et al., (2002), *Proc. Natl. Acad. Sci USA* 99(6):5515-5520.
- [0245] 65. Yu J-Y et al., (2002), *Proc. Natl. Acad. Sci USA* 99(9):6047-6052.
- [0246] 66. McCaffrey, A. P. et al., (2002), *Nature*, Vol. 418, July, 2002.
- [0247] 67. Watanabe A, Taniguchi M. (2005). A case of oropharyngeal cancer with multiple bone metastases from prostate cancer that responded to docetaxel, ifosfamide and cisplatin combination therapy. *Gan To Kagaku Ryoho*, 32(1):65-7.
- [0248] 68. Miner J, Gillan M M, Alex P, Centola M. (2005). Steroid-Refractory Ulcerative Colitis Treated with Corticosteroids, Metronidazole and Vancomycin: a Case Report. *BMC Gastroenterol.* 5(1):3.
- [0249] 69. Orbay E, Sargin M, Sargin H, Gozu H, Bayramcli O U, Yayla A. (2004). Addition of rosiglitazone to glimepirid and metformin combination therapy in type 2 diabetes. *Endocr J.* 51(6):521-7.
- [0250] 70. Smith L E, Wesolowski E, Mclellan A, et al. Oxygen-induced retinopathy in the mouse. *Invest Ophthalmol Vis Sci*, 1994, 35:101-11.

TABLE B1-1

	human VEGF specific siRNA sequences (25 basepairs with blunt ends):
VEGF-1,	CCUGAUGAGAUCGAGUACAUCUUA (SEQ ID NO: 276)
VEGF-2,	GAGUCCAACAUCACCAUGCAGAUUA (SEQ ID NO: 277)
VEGF-3,	AGUCCAACAUCACCAUGCAGAUUAU (SEQ ID NO: 278)
VEGF-4,	CCAACAUCACCAUGCAGAUUAUGCG (SEQ ID NO: 279)

TABLE B1-1-continued

human VEGF specific siRNA sequences (25 basepairs with blunt ends):

VEGF-5,	CACCAUGCAGAUUAUGCGGAUCAA	(SEQ ID NO: 280)
VEGF-6,	GCACAUAGGAGAGAUGAGCUUCCUA	(SEQ ID NO: 281)
VEGF-7,	GAGAGAUGAGCUUCCUACAGCACAA	(SEQ ID NO: 282)

TABLE B1-3

human VEGFR1 specific siRNA sequences (25 basepairs with blunt ends):

VEGFR1-1,	CAAAGGACUUUAUCUUGUC	(SEQ ID NO: 283)
	GUGUA	
VEGFR1-2,	CCCUCGCCGGAAGUUGUAUG	(SEQ ID NO: 284)
	GUUAA	
VEGFR1-3,	CAUCACUCAGCGCAUGGCAA	(SEQ ID NO: 285)
	UAAUA	
VEGFR1-4,	CCACCACUUUAGACUGUCAU	(SEQ ID NO: 286)
	GCUAA	
VEGFR1-5,	CGGACAAGUCUAAUCUGGAG	(SEQ ID NO: 287)
	CUGAU	
VEGFR1-6,	UGACCACAUUGGCCACCAU	(SEQ ID NO: 288)
	CUGAA	
VEGFR1-7,	GAGGGCCUCUGAUGGUGAUU	(SEQ ID NO: 289)
	GUUGA	
VEGFR1-8,	CGAGCUCGGCUUUCAGGAA	(SEQ ID NO: 290)
	GAUAA	
VEGFR1-9,	CAAUCAUGCCAUCUGACA	(SEQ ID NO: 291)
	GGAAA	

TABLE B1-3-continued

human VEGFR1 specific siRNA sequences (25 basepairs with blunt ends):

VEGFR1-10,	GAAAGUAUUUCAGCUCCGAA	(SEQ ID NO: 292)
	GUUUA	

TABLE B1-5

human VEGFR2 specific siRNA sequences (25 basepairs with blunt ends):

VEGFR2-1,	CCUCGGUCAUUUAUGUCUAU	(SEQ ID NO: 293)
	GUUCA	
VEGFR2-2,	CAGAUCUCCAUUUAUUGCUU	(SEQ ID NO: 294)
	CUGUU	
VEGFR2-3,	GACCAACAUGGAGUCGUGUA	(SEQ ID NO: 295)
	CAUUA	
VEGFR2-4,	CCCUUGAGUCCAAUCACACA	(SEQ ID NO: 296)
	AUUA	
VEGFR2-5,	CCAUGUUCUUCUGGCUACUU	(SEQ ID NO: 297)
	CUUGU	
VEGFR2-6,	UCAUUCUAUUGGUCACCAU	(SEQ ID NO: 298)
	CUCAA	
VEGFR2-7,	GAGUUCUUGGCAUCGCGAAA	(SEQ ID NO: 299)
	GUGUA	
VEGFR2-8,	CAGCAGGAAUCAGUCAGUAU	(SEQ ID NO: 300)
	CUGCA	
VEGFR2-9,	CAGUGGUAUGGUUCUUGCCU	(SEQ ID NO: 301)
	CAGAA	
VEGFR2-10,	CCACACUGAGCUCUCCUCCU	(SEQ ID NO: 302)
	GUUUA	

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 302

<210> SEQ ID NO 1

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 1

ccugaugaga ucgaguacau cuuca

25

<210> SEQ ID NO 2

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

-continued

<400> SEQUENCE: 2

ugaagaugua cucgaucuca ucagg 25

<210> SEQ ID NO 3

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 3

gagagaugag cuuccuacag cacaa 25

<210> SEQ ID NO 4

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 4

uugugcugua ggaagcucau cucuc 25

<210> SEQ ID NO 5

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 5

cacaacaaau gugaugcag accaa 25

<210> SEQ ID NO 6

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 6

uuggucugca uucacauuug uugug 25

<210> SEQ ID NO 7

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 7

ucgagaccu gguggacaut t 21

<210> SEQ ID NO 8

<211> LENGTH: 25

<212> TYPE: RNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 8

gcccaacauu ucuacagugu ucuua 25

<210> SEQ ID NO 9
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 9

uaagaacacu guagaauaug uggc 25

<210> SEQ ID NO 10
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 10

cccucgccgg aaguuguauug guuaa 25

<210> SEQ ID NO 11
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 11

uuaaccauac aacuuccggc gaggg 25

<210> SEQ ID NO 12
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 12

ggagaggacc ugaaacugut t 21

<210> SEQ ID NO 13
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 13

ccucuucugu aagacacuca caauu 25

-continued

<210> SEQ ID NO 14
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 14

aaauugugagu gucuacaga agagg 25

<210> SEQ ID NO 15
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 15

cccuugaguc caaucacaca auuaa 25

<210> SEQ ID NO 16
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 16

uaaaugugu gauuggacuc aaggg 25

<210> SEQ ID NO 17
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 17

ccaagugauu gaagcagaug ccuuu 25

<210> SEQ ID NO 18
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 18

aaaggcaucu gcucaauca cuugg 25

<210> SEQ ID NO 19
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

oligonucleotide

<400> SEQUENCE: 19

caguaagcga aagagccggt t 21

<210> SEQ ID NO 20
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 20

ccucaagagc aaacgugacu uauuu 25

<210> SEQ ID NO 21
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 21

aaauaaguca cguuugcucu ugagg 25

<210> SEQ ID NO 22
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 22

caagauccgc agacguguaa auguu 25

<210> SEQ ID NO 23
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 23

aacauuuaca cgucugcgga ucuug 25

<210> SEQ ID NO 24
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 24

gcagcuugag uaaaacgaac guacu 25

<210> SEQ ID NO 25
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 25

aguacguucg uuuaacuca gcuca 25

<210> SEQ ID NO 26
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 26

cagcuugagu uaaacgaacg uacuu 25

<210> SEQ ID NO 27
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 27

aaguacguuc guuaacuca agcug 25

<210> SEQ ID NO 28
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 28

ccaugccaag uggucccagg cugca 25

<210> SEQ ID NO 29
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 29

tgcagcctgg gaccacttgg catgg 25

<210> SEQ ID NO 30
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 30

cacauaggag agaugagcuu ccuca 25

<210> SEQ ID NO 31
<211> LENGTH: 25

-continued

<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 31

ugaggaagcu caucucuccu augug 25

<210> SEQ ID NO 32
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 32

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 33
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 33

auuggcctgc uuaacggucc guagg 25

<210> SEQ ID NO 34
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 34

cucaugucug uucucaagau ccuca 25

<210> SEQ ID NO 35
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 35

ugaggauuu gagaacagac augag 25

<210> SEQ ID NO 36
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 36

cucaugguga uuguggaauu cugca 25

-continued

<210> SEQ ID NO 37
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 37

ugcagaauc cacaaucacc augag 25

<210> SEQ ID NO 38
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 38

gagcauggaa gaggauucug gacuc 25

<210> SEQ ID NO 39
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 39

gaguccagaa ucctcuucca ugctc 25

<210> SEQ ID NO 40
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 40

cagaacagua agcgaaagag cgggc 25

<210> SEQ ID NO 41
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 41

gccggcucu ucgcuuacug uucug 25

<210> SEQ ID NO 42
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 42

gacuuccuga ccuuggagca ucuca 25

<210> SEQ ID NO 43

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 43

ugagaugcuc caaggucagg aaguc 25

<210> SEQ ID NO 44

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 44

ccugaccuug gagcauca ucugu 25

<210> SEQ ID NO 45

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 45

acagaugaga ugcuccaagg ucagg 25

<210> SEQ ID NO 46

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 46

gcuaaggga uggaguucuu ggcau 25

<210> SEQ ID NO 47

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 47

augccaagaa cuccaugccc uuagc 25

<210> SEQ ID NO 48

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

oligonucleotide

<400> SEQUENCE: 48

cacgcuguuu auugaaagag ucaca 25

<210> SEQ ID NO 49
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 49

ugugacucuu ucaauaaaca gcgug 25

<210> SEQ ID NO 50
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 50

cguguuuuu ugaagaguc acaga 25

<210> SEQ ID NO 51
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 51

ucugugacuc uuucaaaaa cagcg 25

<210> SEQ ID NO 52
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 52

caaggagggc cucugauggu gaugu 25

<210> SEQ ID NO 53
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 53

acaucaccau cagaggccu ccuug 25

<210> SEQ ID NO 54
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 54

ccaacuaccu caagagcaaa cguga 25

<210> SEQ ID NO 55
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 55

ucacguuugc ucuugaggua guugg 25

<210> SEQ ID NO 56
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 56

cuaccucaag agcaaacgug acuua 25

<210> SEQ ID NO 57
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 57

uaagucacgu uugcucuuga gguag 25

<210> SEQ ID NO 58
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 58

ccagaaagug caucaucgg gaccu 25

<210> SEQ ID NO 59
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 59

aggucccgau gaaugcacuu ucugg 25

<210> SEQ ID NO 60
<211> LENGTH: 25

-continued

<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 60

cauaucaucgg gaccuggcag cgaga 25

<210> SEQ ID NO 61
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 61

ucucgcugcc aggucccgau gaaug 25

<210> SEQ ID NO 62
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 62

caucgggacc uggcagcgag aaaca 25

<210> SEQ ID NO 63
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 63

uguuucucgc ugccaggucc cgaug 25

<210> SEQ ID NO 64
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 64

gagccuggaa agaaucaaaa ccuuu 25

<210> SEQ ID NO 65
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 65

aaagguuuug auucuucca ggcuc 25

-continued

<210> SEQ ID NO 66
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 66

gccuggaaag aaucaaaacc uuuga 25

<210> SEQ ID NO 67
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 67

ucaaagguuu ugauucuuc caggc 25

<210> SEQ ID NO 68
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 68

gccuggaaag aaucaaaacc uuuga 25

<210> SEQ ID NO 69
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 69

ucaaagguuu ugauucuuc caggc 25

<210> SEQ ID NO 70
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 70

cugaacugag uuuaaaaggc accca 25

<210> SEQ ID NO 71
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 71

ugggugccuu uuaaacugag uucag 25

-continued

<210> SEQ ID NO 72
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 72

gaacugaguu uaaaaggcac ccagc 25

<210> SEQ ID NO 73
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 73

gcugggugcc uuuaaacuc aguug 25

<210> SEQ ID NO 74
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 74

gaagauaaug acucaccugg ggcca 25

<210> SEQ ID NO 75
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 75

uggccccagg ugagucauua ucuuc 25

<210> SEQ ID NO 76
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 76

gauaaugacu caccuggggc cacau 25

<210> SEQ ID NO 77
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 77

-continued

auguggcccc aggugaguca uuauc 25

<210> SEQ ID NO 78
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 78

cucaccuggg gccacauuug aacau 25

<210> SEQ ID NO 79
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 79

auguucaaaug guggccccag gugag 25

<210> SEQ ID NO 80
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 80

cuugcuggga gccugcacca aguca 25

<210> SEQ ID NO 81
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 81

ugacuuggug caggcuccca gcaag 25

<210> SEQ ID NO 82
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 82

gauucuacuu ucuacaauaa gauca 25

<210> SEQ ID NO 83
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 83

ugaucuuuuu guagaaagua gaauc 25

<210> SEQ ID NO 84

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 84

cagagacuga ggcgucagacag uggcu 25

<210> SEQ ID NO 85

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 85

agccacuguc ugcgucacagu cucug 25

<210> SEQ ID NO 86

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 86

gaccugggca agaggaacag acaca 25

<210> SEQ ID NO 87

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 87

ugugucuguu ccucugccc agguc 25

<210> SEQ ID NO 88

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 88

ccaccuucau caagagagag gacga 25

<210> SEQ ID NO 89

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 89

ucguccucuc ucuugaugaa ggugg 25

<210> SEQ ID NO 90

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 90

uauuggauuaa gccgguccca accugu 26

<210> SEQ ID NO 91

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 91

acagguuggg accggcuuaa uccaau 26

<210> SEQ ID NO 92

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 92

cugcagagac cucaaaaggu gucca 25

<210> SEQ ID NO 93

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 93

uggacaccuu uugaggucuc ugcag 25

<210> SEQ ID NO 94

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 94

cagagaccuc aaaagguguc cacgu 25

<210> SEQ ID NO 95

<211> LENGTH: 25

<212> TYPE: RNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 95

acguggacac cuuuugaggu cucug 25

<210> SEQ ID NO 96
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 96

guggugguga ucucagccau ccugg 25

<210> SEQ ID NO 97
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 97

ccaggauggc ugagaucacc accac 25

<210> SEQ ID NO 98
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 98

gguggugauc ucagccaucc uggcc 25

<210> SEQ ID NO 99
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 99

ggccaggaug gcugagauca ccacc 25

<210> SEQ ID NO 100
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 100

ggcaagcugg ucaagaucug ugacu 25

<210> SEQ ID NO 101

-continued

<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 101

agucacagau cuugaccagc uugcc 25

<210> SEQ ID NO 102
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 102

gcaagcuggu caagaucugu gacuu 25

<210> SEQ ID NO 103
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 103

aagucacaga ucuugaccag cuugc 25

<210> SEQ ID NO 104
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 104

ggguggcacc ccuaccag agcug 25

<210> SEQ ID NO 105
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 105

cagcucuggg uaaggggugc caccc 25

<210> SEQ ID NO 106
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 106

gcacccuua cccagagcug cccau 25

-continued

<210> SEQ ID NO 107
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 107

augggcagcu cuggguaagg ggugc 25

<210> SEQ ID NO 108
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 108

cuuaccaga gcugcccaug aacga 25

<210> SEQ ID NO 109
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 109

ucguucaugg gcagcucugg gaaag 25

<210> SEQ ID NO 110
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 110

caugccuccg acgagauca ugaga 25

<210> SEQ ID NO 111
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 111

ucucuagau cucgucggag gcaug 25

<210> SEQ ID NO 112
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 112

-continued

gccuccgacg agaucuauga gauca 25

<210> SEQ ID NO 113
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 113

ugaucucaua gaucucgucg gaggc 25

<210> SEQ ID NO 114
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 114

ccgacgagau cuaugagauc augca 25

<210> SEQ ID NO 115
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 115

ugcaugaucu cauagaucuc gucgg 25

<210> SEQ ID NO 116
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 116

gacgagaucu augagaucau gcaga 25

<210> SEQ ID NO 117
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 117

ucugcaugau cucauagauc ucguc 25

<210> SEQ ID NO 118
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 118

cucuggaucc cagaagguga gaaag 25

<210> SEQ ID NO 119

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 119

cuuucucacc uucugggauc cagag 25

<210> SEQ ID NO 120

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 120

cagcauguca agaucacaga uuuug 25

<210> SEQ ID NO 121

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 121

caaaaucugu gaucuugaca ugcug 25

<210> SEQ ID NO 122

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 122

gaucacagau uuugggcugg ccaa 25

<210> SEQ ID NO 123

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 123

uuuggccagc caaaaaucug ugauc 25

<210> SEQ ID NO 124

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

oligonucleotide

<400> SEQUENCE: 124

cagauuuugg gcuggccaaa cugcu 25

<210> SEQ ID NO 125
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 125

agcaguuugg ccagcccaaa aucug 25

<210> SEQ ID NO 126
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 126

caaagugccu aucaagugga uggca 25

<210> SEQ ID NO 127
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 127

ugccauccac uugauaggca cuuug 25

<210> SEQ ID NO 128
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 128

ccaucgaugu cuacaugauc auggu 25

<210> SEQ ID NO 129
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 129

accaugauca uguagacauc gaugg 25

<210> SEQ ID NO 130
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 130

cgaugucuac augaucaugg ucaagu 26

<210> SEQ ID NO 131
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 131

acuugaccuau gaucauguag acaucg 26

<210> SEQ ID NO 132
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 132

cuacaugauc auggucaagu gcugg 25

<210> SEQ ID NO 133
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 133

ccagcacuug accaugauca uguag 25

<210> SEQ ID NO 134
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 134

caugaucaug gucaagugcu ggauga 26

<210> SEQ ID NO 135
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 135

ucauccagca cuugaccaug aucaug 26

<210> SEQ ID NO 136
<211> LENGTH: 25

-continued

<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 136

ggaugaaaga augcauuugc caagu 25

<210> SEQ ID NO 137
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 137

acuuggcaaa ugcauucuuu caucc 25

<210> SEQ ID NO 138
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 138

gacaaccug acuaccagca ggacu 25

<210> SEQ ID NO 139
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 139

aguccugcug guagucaggg uuguc 25

<210> SEQ ID NO 140
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 140

ccuucuuaaa gaccuaccag gaggu 25

<210> SEQ ID NO 141
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 141

accuccugga uggucuuuaa gaagg 25

-continued

<210> SEQ ID NO 142
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 142

caagauccgc agacguguaa auguu 25

<210> SEQ ID NO 143
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 143

aacauuuaca cgucugcgga ucuug 25

<210> SEQ ID NO 144
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 144

caaggagggc cucugauggu gauu 25

<210> SEQ ID NO 145
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 145

acaucaccau cagaggccu ccuug 25

<210> SEQ ID NO 146
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 146

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 147
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 147

auuggccgc uuaacgguc guagg 25

-continued

<210> SEQ ID NO 148
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 148

caagaucgc agacguguaa auguu 25

<210> SEQ ID NO 149
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 149

aacauuaca cgucugcgga ucuug 25

<210> SEQ ID NO 150
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 150

ccagaaagug gauucaucgg gaccu 25

<210> SEQ ID NO 151
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 151

aggucccgau gaaugcacu ucugg 25

<210> SEQ ID NO 152
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 152

cucaugucug uucucaagau ccuca 25

<210> SEQ ID NO 153
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 153

-continued

ugaggauccu gagaacagac augag 25

<210> SEQ ID NO 154
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 154

gcagcuugag uaaaacgaac guacu 25

<210> SEQ ID NO 155
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 155

aguacguucg uuuaacucua gcugc 25

<210> SEQ ID NO 156
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 156

ccagaaagug caucaucgg gaccu 25

<210> SEQ ID NO 157
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 157

aggucccgau gaaugcacuu ucugg 25

<210> SEQ ID NO 158
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 158

cucaugucug uucuaagau ccuca 25

<210> SEQ ID NO 159
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 159

ugaggauccuu gagaacagac augag 25

<210> SEQ ID NO 160

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 160

caagauccgc agacguguaa auguu 25

<210> SEQ ID NO 161

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 161

aacauuuaca cgucugcgga ucuug 25

<210> SEQ ID NO 162

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 162

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 163

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 163

auuggcccgc uuaacggucc guagg 25

<210> SEQ ID NO 164

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 164

cucuggaucc cagaagguga gaaag 25

<210> SEQ ID NO 165

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 165

cuuucucacc uucugggauc cagag 25

<210> SEQ ID NO 166

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 166

caagaucgc agacguguaa auguu 25

<210> SEQ ID NO 167

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 167

aacaauuaca cgucugcgga ucuug 25

<210> SEQ ID NO 168

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 168

cucaugucug uucucaagau ccuca 25

<210> SEQ ID NO 169

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 169

ugaggauuu gagaacagac augag 25

<210> SEQ ID NO 170

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 170

caaagugccu aucaagugga uggca 25

<210> SEQ ID NO 171

<211> LENGTH: 25

<212> TYPE: RNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 171

ugccauccac uugauagga cuuug 25

<210> SEQ ID NO 172
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 172

gcagcuugag uaaaacgaac guacu 25

<210> SEQ ID NO 173
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 173

aguacguucg uuuaacuaa gcugc 25

<210> SEQ ID NO 174
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 174

cucaugucug uucucaagau ccuca 25

<210> SEQ ID NO 175
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 175

ugaggauuu gagaacagac augag 25

<210> SEQ ID NO 176
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 176

caugaucaug gucaagugcu ggauga 26

<210> SEQ ID NO 177

-continued

<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 177

ucauccagca cuugaccaug aucaug 26

<210> SEQ ID NO 178
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 178

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 179
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 179

auuggccccg uuaacggucc guagg 25

<210> SEQ ID NO 180
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 180

gaagauaaug acucaccugg ggcca 25

<210> SEQ ID NO 181
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 181

uggccccagg ugagucuuu ucuuc 25

<210> SEQ ID NO 182
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 182

cucuggaucc cagaagguga gaaag 25

-continued

<210> SEQ ID NO 183
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 183

cuuucucacc uucugggauc cagag 25

<210> SEQ ID NO 184
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 184

ccagaaagug caucaucgg gaccu 25

<210> SEQ ID NO 185
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 185

aggucccgau gaaugcacuu ucugg 25

<210> SEQ ID NO 186
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 186

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 187
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 187

auuggccgc uuaacggucc guagg 25

<210> SEQ ID NO 188
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 188

-continued

gauaaugacu caccuggggc cacau 25

<210> SEQ ID NO 189
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 189

auguggcccc aggugaguca uauac 25

<210> SEQ ID NO 190
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 190

cucuggaucc cagaagguga gaaag 25

<210> SEQ ID NO 191
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 191

cuuucucacc uucugggauc cagag 25

<210> SEQ ID NO 192
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 192

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 193
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 193

auuggcccg c uuaacggucc guagg 25

<210> SEQ ID NO 194
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 194

gauaaugacu caccuggggc cacau 25

<210> SEQ ID NO 195

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 195

auguggcccc aggugaguca uuauc 25

<210> SEQ ID NO 196

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 196

ccuucuuaaa gaccauccag gaggu 25

<210> SEQ ID NO 197

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 197

accuccugga uggucuuaa gaagg 25

<210> SEQ ID NO 198

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 198

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 199

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 199

auuggcccgc uuaacggucc guagg 25

<210> SEQ ID NO 200

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

oligonucleotide

<400> SEQUENCE: 200

ggcaagcugg ucaagaucug ugacu 25

<210> SEQ ID NO 201
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 201

agucacagau cuugaccagc uugcc 25

<210> SEQ ID NO 202
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 202

cucuggaucc cagaagguga gaaag 25

<210> SEQ ID NO 203
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 203

cuuucucacc uucugggauc cagag 25

<210> SEQ ID NO 204
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 204

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 205
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 205

auuggcccgc uuaacggucc guagg 25

<210> SEQ ID NO 206
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 206

caugccuccg acgagauca ugaga 25

<210> SEQ ID NO 207
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 207

ucucauagau cucgucggag gcaug 25

<210> SEQ ID NO 208
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 208

cucuggaucc cagaagguga gaaag 25

<210> SEQ ID NO 209
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 209

cuuucucacc uucugggauc cagag 25

<210> SEQ ID NO 210
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 210

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 211
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 211

auuggcccg uuaacggucc guagg 25

<210> SEQ ID NO 212
<211> LENGTH: 25

-continued

<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 212

ggcaagcugg ucaagaucug ugacu 25

<210> SEQ ID NO 213
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 213

agucacagau cuugaccagc uugcc 25

<210> SEQ ID NO 214
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 214

ccuucuuaaa gaccauccag gaggu 25

<210> SEQ ID NO 215
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 215

accuccugga uggucuuaa gaagg 25

<210> SEQ ID NO 216
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 216

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 217
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 217

auuggcccgc uuaacggucc guagg 25

-continued

<210> SEQ ID NO 218
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 218

gauaaugacu caccuggggc cacau 25

<210> SEQ ID NO 219
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 219

auguggcccc aggugaguca uuauc 25

<210> SEQ ID NO 220
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 220

ggcaagcugg ucaagaucug ugacu 25

<210> SEQ ID NO 221
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 221

agucacagau cuugaccagc uugcc 25

<210> SEQ ID NO 222
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 222

cucuggaucc cagaagguga gaaag 25

<210> SEQ ID NO 223
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 223

cuuucucacc uucugggauc cagag 25

-continued

<210> SEQ ID NO 224
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 224

ccagaaagug caucaucgg gaccu 25

<210> SEQ ID NO 225
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 225

aggucccgau gaaugcacuu ucugg 25

<210> SEQ ID NO 226
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 226

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 227
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 227

auuggcccg uuaacggucc guagg 25

<210> SEQ ID NO 228
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 228

caugccuccg acgagauca ugaga 25

<210> SEQ ID NO 229
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 229

-continued

ucucauagau cucgucggag gcaug 25

<210> SEQ ID NO 230
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 230

cucuggaucc cagaagguga gaaag 25

<210> SEQ ID NO 231
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 231

cuuucucacc uucugggauc cagag 25

<210> SEQ ID NO 232
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 232

gcagcuugag uuaaacgaac guacu 25

<210> SEQ ID NO 233
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 233

aguacguucg uuuaacucuaa gcugc 25

<210> SEQ ID NO 234
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 234

ccagaaagug caucaucgg gaccu 25

<210> SEQ ID NO 235
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 235

agguccccgau gaaugcacuu ucugg 25

<210> SEQ ID NO 236

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 236

ccuacggacc guuaagcggg ccaau 25

<210> SEQ ID NO 237

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 237

auuggcccg c uaaacggucc guagg 25

<210> SEQ ID NO 238

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 238

gauaaugacu caccuggggc cacau 25

<210> SEQ ID NO 239

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 239

auguggcccc aggugaguca uauac 25

<210> SEQ ID NO 240

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 240

ggcaagcugg ucaagaucug ugacu 25

<210> SEQ ID NO 241

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 241

agucacagau cuugaccagc uugcc 25

<210> SEQ ID NO 242

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 242

ccuucuuaaa gaccauccag gaggu 25

<210> SEQ ID NO 243

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 243

accuccugga uggucuuaa gaagg 25

<210> SEQ ID NO 244

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 244

gcagcuugag uaaaacgaac guacu 25

<210> SEQ ID NO 245

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 245

aguacguucg uuuaacuaa gcugc 25

<210> SEQ ID NO 246

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 246

Ala Cys Arg Gly Asp Met Phe Gly Cys Ala
1 5 10

<210> SEQ ID NO 247

<211> LENGTH: 21

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 247

aagctcagca cacagaaaga c 21

<210> SEQ ID NO 248
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 248

aatgcgggcg tggtgacagt a 21

<210> SEQ ID NO 249
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 249

ggaaccgcug gagagcaacu gcaua 25

<210> SEQ ID NO 250
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 250

ccuugcgac cucucguuga cguau 25

<210> SEQ ID NO 251
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 251

ucgagaccu ggugacaut t 21

<210> SEQ ID NO 252
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 252

auguccacca gggucucgat t 21

<210> SEQ ID NO 253

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 253

gcugaccug aaguucauct t 21

<210> SEQ ID NO 254

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Combined DNA/RNA Molecule:
Synthetic oligonucleotide

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 254

gaugaacuuc agggucagct t 21

<210> SEQ ID NO 255

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 255

ccaagugauu gaagcagaug ccuuu 25

<210> SEQ ID NO 256

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 256

tcaacgttga 10

<210> SEQ ID NO 257

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 257

gctagacggt agcgt 15

-continued

<210> SEQ ID NO 258
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 258

aagccgtcct gtgtgccgct g 21

<210> SEQ ID NO 259
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 259

aacgatgaag ccttgagtg c 21

<210> SEQ ID NO 260
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 260

aagttaaaag tgctgaact g 21

<210> SEQ ID NO 261
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 261

aagcaggcca gactctcttt c 21

<210> SEQ ID NO 262
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 262

aagctcagca cacagaaaga c 21

<210> SEQ ID NO 263
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 263

-continued

atgcgcggt ggtgacagta 20

<210> SEQ ID NO 264
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 264

aacagttgcg cagcctgaat g 21

<210> SEQ ID NO 265
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 265

aacttaatcg cttgcagca c 21

<210> SEQ ID NO 266
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 266

aagctatgaa acgatatggg c 21

<210> SEQ ID NO 267
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 267

aaccgctgga gagcaactgc a 21

<210> SEQ ID NO 268
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 268

gatgtctacc agcgaagcta ctgccgtccg 30

<210> SEQ ID NO 269
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

-continued

<400> SEQUENCE: 269

gaacatcgat gacaagctta ggtatcgata caagctgcct cgccttg 47

<210> SEQ ID NO 270

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 270

gtcagctgct gggacaccgc ggtcttgctt 30

<210> SEQ ID NO 271

<211> LENGTH: 47

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 271

gaacatcgat gacaagctta ggtatcgata tagattgaag attccgc 47

<210> SEQ ID NO 272

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 272

tggctggtca aacagctcat c 21

<210> SEQ ID NO 273

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 273

ctcatccaag ggcaattcat c 21

<210> SEQ ID NO 274

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 274

cattgtgatg gactccggag acgg 24

<210> SEQ ID NO 275

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 275

catctcctgc tgaagtctag agc 23

<210> SEQ ID NO 276
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 276

ccugaugaga ucgaguacau cuuca 25

<210> SEQ ID NO 277
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 277

gaguccaaca ucaccaugca gauua 25

<210> SEQ ID NO 278
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 278

aguccaacau caccaugcag auuau 25

<210> SEQ ID NO 279
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 279

ccaacaucac caugcagauu augcg 25

<210> SEQ ID NO 280
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 280

caccaugcag auuauugcga ucaaa 25

<210> SEQ ID NO 281
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 281

gcacauagga gagaugagcu uccua 25

<210> SEQ ID NO 282
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 282

gagaugag cuuccuacag cacaa 25

<210> SEQ ID NO 283
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 283

caaagacuu uauacuuguc gugua 25

<210> SEQ ID NO 284
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 284

cccucgccg aaguuguaug guuaa 25

<210> SEQ ID NO 285
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 285

caucacucag cgcauggcaa uaaua 25

<210> SEQ ID NO 286
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 286

ccaccacuuu agacugucau gcuaa 25

<210> SEQ ID NO 287
<211> LENGTH: 25

-continued

<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 287

cggacaaguc uaaucuggag cugau 25

<210> SEQ ID NO 288
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 288

ugaccacau uggccacau cugaa 25

<210> SEQ ID NO 289
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 289

gagggcucu gauggugauu guuga 25

<210> SEQ ID NO 290
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 290

cgagcuccg cuucaggaa gauaa 25

<210> SEQ ID NO 291
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 291

caaucaaugc cauacugaca ggaaa 25

<210> SEQ ID NO 292
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 292

gaaaguauuu cagcuccgaa guuuu 25

-continued

<210> SEQ ID NO 293
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 293

ccucggucau uuaugucuau guuca 25

<210> SEQ ID NO 294
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 294

cagaucucca uuuauugcuu cuguu 25

<210> SEQ ID NO 295
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 295

gaccaacaug gagucgugua cauaa 25

<210> SEQ ID NO 296
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 296

cccuugaguc caaucacaca auuaa 25

<210> SEQ ID NO 297
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 297

ccauguucuu cuggcuacuu cuugu 25

<210> SEQ ID NO 298
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 298

ucauucuuau uggucaccau cucaa 25

-continued

```

<210> SEQ ID NO 299
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

<400> SEQUENCE: 299

gaguucugg caucgcgaaa gugua                25

<210> SEQ ID NO 300
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

<400> SEQUENCE: 300

cagcaggaau cagucaguau cugca                25

<210> SEQ ID NO 301
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

<400> SEQUENCE: 301

cagugguauug guucugccu cagaa                25

<210> SEQ ID NO 302
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

<400> SEQUENCE: 302

ccacacugag cucuccuccu guuua                25

```

1-38. (canceled)

39. A composition selected from the group consisting of:
 a composition comprising a siRNA identified by SEQ ID NO. 2, a siRNA identified by SEQ ID NO. 9 a siRNA identified by SEQ ID NO. 14 and a pharmaceutically acceptable carrier;
 a composition comprising a siRNA identified by SEQ ID NO. 2, a siRNA identified by SEQ ID NO. 9, a siRNA identified by SEQ ID NO. 16 and a pharmaceutically acceptable carrier;
 a composition comprising a siRNA identified by SEQ ID NO. 2, a siRNA identified by SEQ ID NO. 9, a siRNA identified by SEQ ID NO. 18 and a pharmaceutically acceptable carrier;
 a composition comprising a siRNA identified by SEQ ID NO. 2, a siRNA identified by SEQ ID NO. 11, a siRNA identified by SEQ ID NO. 14 and a pharmaceutically acceptable carrier;

a composition comprising a siRNA identified by SEQ ID NO. 2, a siRNA identified by SEQ ID NO. 11, a siRNA identified by SEQ ID NO. 16 and a pharmaceutically acceptable carrier;
 a composition comprising a siRNA identified by SEQ ID NO. 2, a siRNA identified by SEQ ID NO. 11, a siRNA identified by SEQ ID NO. 18 and a pharmaceutically acceptable carrier;
 a composition comprising a siRNA identified by SEQ ID NO. 2, a siRNA with the sequence of SEQ ID NO. 21, a siRNA identified by SEQ ID NO. 14 and a pharmaceutically acceptable carrier;
 a composition comprising a siRNA identified by SEQ ID NO. 2, a siRNA with the sequence of SEQ ID NO. 21, a siRNA identified by SEQ ID NO. 16 and a pharmaceutically acceptable carrier;

42. The composition of claim 41, wherein at least one of the one or more additional nucleic acid molecules decreases the expression of a gene that promotes angiogenesis.

43. The composition of claim 41, wherein at least one of the one or more additional nucleic acid molecules decreases the expression of a gene selected from the group consisting of; VEGFR1, VEGFR2, VEGFR3, PDGF, PDGFR- α , PDGFR- β , EGF, EGFR, RAF-a, RAF-c, AKT, RAS, NFkB, HIF, bFGF, bFGFR, Her-2, c-Met, c-Myc and HGF.

44. The composition of claim 39 or 40, wherein the carrier is a nucleic acid delivery vehicle.

45. The composition of claim 44, wherein the nucleic acid delivery vehicle is synthetic.

46. The composition of claim 45 wherein the synthetic nucleic acid delivery vehicle comprises a core region comprising a cationic polymer and the nucleic acid.

47. The composition of claim 46, wherein the cationic polymer is polyethyleneimine.

48. The composition of claim 46, wherein the cationic polymer is a histidine-lysine co-polymer.

49. The composition of claim 46, wherein the synthetic nucleic acid delivery vehicle further comprises a protective hydrophilic layer.

50. The composition of claim 49, wherein the protective hydrophilic layer comprises a component selected from the group consisting of: polyethylene glycol, a polyacetal and a polyoxazoline.

51. The composition of claim 46, wherein the synthetic nucleic acid vehicle further comprises a targeting moiety.

52. The composition of claim 49, wherein the synthetic nucleic acid vehicle further comprises a targeting moiety.

53. The composition of claim 51, wherein the targeting moiety binds a tumor specific molecule or an angiogenesis-specific molecule.

54. The composition of claim 51, wherein the targeting moiety binds endothelial cells.

55. The composition of claim 54, wherein the targeting moiety binds an integrin on vascular endothelial cells.

56. The composition of claim 55, wherein the targeting moiety is a peptide comprising the amino acid sequence RGD.

57. The composition of claim 56, wherein the peptide comprising RGD is a cyclic peptide.

58. The composition of claim 55, wherein the synthetic nucleic acid delivery vehicle comprises:

- (a) a core region comprising PEI;
- (b) a protective hydrophilic layer comprising PEG; and
- (c) a cyclic RGD-containing peptide targeting moiety.

59. The composition of claim 39 or 40 comprising an additional therapeutic agent selected from the group consisting of: an anti-cancer agent, an anti-inflammatory agent and an anti-infective agent.

60. The composition of claim 39 or 40 comprising an anti-angiogenic agent.

61. The composition of claim 60, wherein the anti-angiogenic agent is an inhibitor selected from the group consisting of: an inhibitor of human VEGF, an inhibitor of human VEGFR1 and an inhibitor of human VEGFR2.

62. A method for reducing neovascularization in a subject in need thereof comprising the step of administering to the subject the composition of claim 39 or 40.

63. A method for reducing neovascularization in a subject in need thereof comprising the step of administering to the subject the composition of claim 58.

64. A method of reducing tumor growth in a subject in need thereof, comprising the step of administering to the subject the composition of claim 39 or 40.

65. A method of reducing tumor growth in a subject in need thereof, comprising the step of administering to the subject the composition of claim 58.

66. A method for decreasing one or more of VEGF, VEGFR1 or VEGFR2 protein levels in a cell comprising introducing into the cell the nucleic acid molecules of claim 39.

67. A method for decreasing the protein level of VEGF or VEGFR1, or of both VEGF and VEGFR1, in a cell comprising introducing into the cell the nucleic acid molecules of claim 40.

68. A method for decreasing the protein level of VEGF or VEGFR2, or of both VEGF and VEGFR2, in a cell comprising introducing into the cell the nucleic acid molecules of claim 40.

* * * * *