



(86) Date de dépôt PCT/PCT Filing Date: 2009/07/22
(87) Date publication PCT/PCT Publication Date: 2010/02/18
(85) Entrée phase nationale/National Entry: 2011/02/01
(86) N° demande PCT/PCT Application No.: EP 2009/059403
(87) N° publication PCT/PCT Publication No.: 2010/018049
(30) Priorité/Priority: 2008/08/11 (EP08162137.7)

(51) Cl.Int./Int.Cl. *A23D 9/02* (2006.01),
A23L 1/29 (2006.01), *C11B 1/02* (2006.01),
C11B 1/06 (2006.01)

(71) Demandeur/Applicant:
NESTEC S.A., CH

(72) Inventeurs/Inventors:
BEZELGUES, JEAN-BAPTISTE, CH;
WANG, JUNKUAN, CH;
BERTHOLET, RAYMOND, CH;
DUCRET, PIERRE, CH

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : HUILE CONTENANT UN OU PLUSIEURS PHOSPHOLIPIDES D'ACIDES GRAS POLYINSATURES A
CHAINE LONGUE DERIVES DE BIOMASSE

(54) Title: OIL CONTAINING ONE OR MORE LONG-CHAIN POLYUNSATURATED FATTY ACIDS PHOSPHOLIPIDS
DERIVED FROM BIOMASS

(57) **Abrégé/Abstract:**

The present invention relates to a method to extract long-chain polyunsaturated fatty acids from sources using emulsifiers dispersed in oil as carrier and to the oils obtainable from such a method. It also relates to the use of emulsifier in oil for improving the extraction of long- chain polyunsaturated fatty acids in the form of phospholipids and, in a further aspect, to an oil composition comprising long-chain polyunsaturated fatty acids in an amount up to 4% and phospholipids in an amount up to 15% and at least one emulsifier.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 February 2010 (18.02.2010)

(10) International Publication Number
WO 2010/018049 A1

(51) International Patent Classification:

A23D 9/02 (2006.01) *C11B 1/06* (2006.01)
C11B 1/02 (2006.01) *A23L 1/29* (2006.01)

(21) International Application Number:

PCT/EP2009/059403

(22) International Filing Date:

22 July 2009 (22.07.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

08162137.7 11 August 2008 (11.08.2008) EP

(71) Applicant (for all designated States except US):
NESTEC S.A. [CH/CH]; Avenue Nestlé 55, CH-1800
Vevey (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BEZELGUES, Jean-Baptiste** [FR/CH]; Route de Moudon 21, CH-1522 Lucens (CH). **WANG, Junkuan** [CH/CH]; Chemin des Abbesses 2B, CH-1027 Lonay (CH). **BERTHOLET, Raymond** [CH/CH]; Ch. Vers-chez-Cochard 11, CH-1807 Blonay (CH). **DUCRET, Pierre** [CH/CH]; Chemin de Cornu, CH-1113 St-Saphorin-sur-Morges (CH).

(74) Agent: **MARQUARDT, Ulf**; Avenue Nestlé 55, CH-1800 Vevey (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) Title: OIL CONTAINING ONE OR MORE LONG-CHAIN POLYUNSATURATED FATTY ACIDS PHOSPHOLIPIDS DERIVED FROM BIOMASS

(57) Abstract: The present invention relates to a method to extract long-chain polyunsaturated fatty acids from sources using emulsifiers dispersed in oil as carrier and to the oils obtainable from such a method. It also relates to the use of emulsifier in oil for improving the extraction of long-chain polyunsaturated fatty acids in the form of phospholipids and, in a further aspect, to an oil composition comprising long-chain polyunsaturated fatty acids in an amount up to 4% and phospholipids in an amount up to 15% and at least one emulsifier.



WO 2010/018049 A1

Oil containing one or more long-chain polyunsaturated
fatty acids phospholipids derived from biomass

5 **Field of the invention**

The present invention relates to a method to extract long-chain polyunsaturated fatty acids (LC-PUFAs) in the form of phospholipids from biomass using emulsifiers dispersed
10 in an oil as carrier and to the oils obtainable from such a method as a source of LC-PUFAs for food stuff, nutritional supplement, cosmetic or pharmaceutical composition. It also relates to the use of low HLB emulsifier in oil for improving the extraction of long-
15 chain polyunsaturated fatty acids in the form of phospholipids and, in a further aspect, to an oil composition comprising LC-PUFAs in an amount up to 4% and phospholipids in an amount up to 10%

20 **Background of the invention**

Long chain polyunsaturated fatty acids (LC-PUFAs) such as e.g. arachidonic acid (ARA), eicosapentaenoic (EPA) and docosahexaenoic (DHA) acids, dihomogammalinolenic (DHGLA)
25 are of interest in general public health. For several years, these fatty acids have been investigated for their roles in metabolic pathways and regulation processes. LC-PUFAs are associated with numerous health benefits such as alleviated symptoms of arthritis and a reduced risk of
30 cardiovascular and inflammatory diseases and growth development of infants.

A large number of food applications have been developed with marine oils, which are rich in LC-PUFA in the form of triacylglycerols.

5 LC-PUFAs in the form of triacylglycerols can be obtained by a process of mixing, milling and pressing the dried biomass with a vegetable oil as described for example in EP 1 239 022 and US 2006/0068076. . Although this method is efficient in obtaining LC-PUFAs when present as
10 triacylglycerol in the biomass, it leads to very poor results when the LC-PUFAs from the biomass are mainly in form of phospholipids.

Recently, there has been a growing interest in developing
15 LC-PUFAs in the form of phospholipids to enhance their bioavailability, bioefficacy and oxidative stability. Indeed, some papers assume that the nature of the glyceride structures which act as dietary carrier (e.g. triacylglycerol, phospholipids, etc.) of DHA for instance
20 could play an important role in the absorption rate of DHA at the enterocyte level and, therefore, in its accretion rate in the targeted organs such as brain or eyes.

Marine biomasses such as krill or crustaceans represent an
25 interesting source of LC-PUFAs in the form of phospholipids. The lipid composition of krill may contain for example 40-60% phospholipids, 20-30% triacylglycerols, free fatty acids and sterols. The fatty acids EPA and DHA are esterified almost exclusively in the form of
30 phospholipids, mainly in phosphatidylcholine.

Krill is commercially available as dried powder and krill oil is obtained by solvent extraction or supercritical carbon dioxide extraction. EP 1 417 211 describes a process by which extraction of phospholipids is carried
5 out by treatment with organic solvents such as ethanol, acetone, hexane. This process however can lead to the degradation of the phospholipids and requires a number of purification steps in order to get rid of solvent traces. Furthermore, the oil obtained by such a process is not
10 only expensive, but is also very bad in smell and taste.

Object of the present invention

There is therefore a need to develop a method for the extraction of LC-PUFAs, in particular LC-PUFAs in the form
15 of phospholipids, which is high yielding and does not require the use of organic solvents.

Summary of the present invention

This object is solved by means of the independent claims.
20 The dependent claims further develop the central idea of the invention.

Thus, in a first aspect the present invention relates to a method to extract long-chain polyunsaturated fatty acids
25 from a source comprising the steps of

- a. mixing the source with an oil comprising at least one emulsifier
- b. wet milling the mixture to form a slurry and
- c. pressing the resulting slurry to obtain an oil
30 comprising said long-chain polyunsaturated fatty acids

and to an oil obtainable by such a method.

For example, the oil used in step a. may be enriched with at least one emulsifier.

Additionally or alternatively step a. may involve mixing
5 the source with an oil and at least one emulsifier.

The emulsifier may be added to the oil in a weight-ratio ranging from 1:6 to 1:1, for example from 1:2 to 1:5.

10 Another aspect of the invention relates to the use of at least one emulsifier in oil for improving the extraction of long-chain polyunsaturated fatty acids in the form of phospholipids from a biomass source.

15 An oil composition comprising long-chain polyunsaturated fatty acids in an amount up to 4% and phospholipids in an amount up to 10% also forms part of the present invention.

A third aspect of the invention pertains to an oil
20 composition according to any of claims 11 to 13 for use in cosmetic and/or therapeutic treatment.

Finally, the use of an oil composition according to any of claims 11 to 13 in food also forms part of the present
25 invention.

Figures

The present invention is further described below with
30 reference to a preferred embodiment as illustrated in figure 1.

Figure 1 is a flow-chart diagram showing the different stages of a process of the present invention.

5 Figure 2 shows a volatile profile (blue curve commercial shrimp oil, black curve soft extracted shrimp oil)

10 Detailed description of the invention

The present invention relates to the extraction of long-chain polyunsaturated fatty acids. By long-chain polyunsaturated fatty acids are meant fatty acids having an aliphatic chain of at least 18 carbons. These include
15 linoleic acid, alpha-linolenic acid, dihomogammalinolenic acid, arachidonic acid, eicosapentaenoic acid, docosahexaenoic acid, etc. Preferably, the extracted long-chain polyunsaturated fatty acids are in the form of
20 phospholipids. Thus, the fatty acids are esterified to a phosphoglycerol backbone.

Long-chain polyunsaturated fatty acids are present in a number of natural sources such as vegetable fat, animal
25 fat but also in fish, zooplankton, crustaceans, microorganisms such as microalgae, fungus.

In the present invention, the preferred source is a dried biomass. More preferably, it is a marine dried biomass.
30 This biomass may be selected from zooplankton (e.g. Krill) crustacean (e.g. *Pandalus Borealis*), microalgae (e.g. *Isochrysis Galbana*) or by-products from fisheries such as

fish heads, viscerae etc. These biomasses are particularly useful when fatty acids in the form of phospholipids are to be extracted.

- 5 The method of the present invention entails a first step of mixing the source with an oil comprising at least one emulsifier.

10 The oil may be selected from any of medium-chain triglyceride (MCT) oils, vegetable oil, essential oil or any mixtures thereof. The essential oil may be for example limonene. Preferably, the oil used is a vegetable oil selected from sunflower oil, rapeseed oil, soybean oil, palm oil, coconut oil or mixtures thereof.

15

The emulsifier may be selected from any emulsifiers having a hydrophilic-lipophilic balance of less than 7. The hydrophilic-lipophilic balance (HLB) of a surfactant is a measure of the degree to which it is hydrophilic or lipophilic, determined by calculating values for the different regions of the molecule, as described by Griffin in 1949[Journal of the Society of Cosmetic Chemists 1 (1949): 311] and 1954[Journal of the Society of Cosmetic Chemists 5 (1954): 259]. Briefly, the HLB may be
25 calculated as follows:

$$\text{HLB} = 20 * M_h / M$$

30 where M_h is the molecular mass of the hydrophilic portion of the Molecule, and M is the molecular mass of the whole molecule.

The resulting HLB will be a number on an arbitrary scale of 0 to 20. An HLB value of 0 corresponds to a completely hydrophobic molecule, and a value of 20 would correspond to a molecule made up completely of hydrophilic
5 components.

Preferably, the emulsifiers used are food-grade. More preferably, the emulsifier is selected from monoglycerides, diglycerides, sorbitan monoesters such as
10 sorbitan monolaurate, sorbitan monopalmitate, sorbitan monooleate, sorbitan monostearate, polypropylene glycol monoester, polyglycerol esters or any mixtures thereof. Most preferably, the emulsifier is a monoglyceride.

15 The oil to emulsifier weight-ratio may be ranging from 1:1 to 6:1, for example from 2:1 to 5:1. Therefore it may be 2:1, 3:1, 4:1, 5:1 or any intermediate value. The concentration of emulsifier in the oil may range from 5 to 50%, for example from 10% to 40%.

20

The biomass is preferably mixed into the oil in a weight-ratio of 5:95 to 50:50.

The source of LC-PUFA phospholipids and the oil comprising
25 at least one emulsifier are then milled in a second step of the invention to form a slurry.

This grinding step may be carried out with any type of mill, preferably a colloidal mill. This step is preferably
30 carried out at less than 60°C, more preferably at about 50°C.

The milling is carried out for a period of time ranging from 10 seconds to 2 minutes. Preferably, the milling is carried out for 1 minute.

5 The slurry obtained by milling the source with the oil/emulsifier mixture comprises particles which predominantly originate from the source. Preferably, at least 90% of the particles in the slurry have a particle size of less than 500 microns, preferably less than 300
10 microns. Particle size can easily be measured by the skilled person using a particle sizer for instance laser scattering.

In a third step of the present method, the slurry is
15 pressed to obtain an oil comprising long-chain polyunsaturated fatty acids. By "pressing" is meant subjecting the slurry resulting from the previous step to pressure. Thus, pressing can be carried out under pressure of 10bars up to 500bars. For this purpose a membrane
20 filter press may be used. This typically exerts a pressure of up to 30 bars. Alternatively, a conventional hydraulic press may be used. Pressures of up to 500bars can be achieved with such hydraulic press.

25 After pressing, oil comprising long-chain polyunsaturated fatty acids can be collected.

Preferably the steps of the present method are carried out at a temperature of 30°C to 80°C.

30

Optionally, the method may comprise additional steps of removing the fines from the collected oil and deodorising

said oil. Typically, the fines may be removed using deep filter cellulose sheet.

Deodorising the resulting oil may be done by steam stripping, for instance using a 1% per hour steam injection rate under 1mbar of pressure and a temperature of between 80 and 140°C. Nitrogen stripping may equally be carried out, for example at a temperature of 80-140°C under 10mbar of pressure. Alternatively, short path molecular distillation can be carried out to deodorise the extracted oil.

The method of the invention thus allows to obtain an oil enriched with long-chain polyunsaturated fatty acids, in particular long-chain PUFAs in the form of phospholipids, without resorting to the use of organic solvents which have been traditionally used to extract polar lipids. By using neutral oil with an emulsifier, the process is natural and soft and allows the extraction of polar lipids in high yields. Indeed at least 80%, preferably at least 90% extraction yield can be achieved using the present method.

The purity of the resulting oil and its organoleptic properties are also improved compared to oils extracted with conventional organic solvents or using supercritical methods.

The resulting oil is therefore particularly suited for use not only in food, but also for use in cosmetic, pharmaceutical, nutraceutical, medical, nutritional compositions, where more natural processing methods are

required (e.g. for infant formulas). It can therefore also be used in cosmetic and/or therapeutic treatments.

One embodiment of the present invention is an oil
5 composition comprising per 100g oil composition about 2-4
g, preferably about 2.5-3.5 g, palmitic acid; about 0.5 -
4 g, preferably about 1-2 g, stearic acid; about 8-12 g,
preferably about 9-11 g, oleic acid; about 14-22 g,
preferably about 16-20 g, linoleic acid; about 0-0,5 g,
10 preferably about 0-0.2 g, linolenic acid; about 0,3 - 4 g,
preferably about 0.5-2 g eicosaenoic acid (EPA); and/or
about 0.3 - 4 g, preferably about 0.5-2 g, docosahexaenoic
acid (DHA). This oil composition is obtainable the method
described herein.

15

The present method is particularly suited for the
extraction of long-chain polyunsaturated fatty acids in
the form of phospholipids. Indeed, previous methods have
either suffered from poor yields regarding these polar
20 lipids or have made use of processes which are not cost
effective and not natural, resorting to the use of polar
solvents for example.

Referring to figure 1, MCT oil is mixed with a
25 monoglyceride (Dimodan) in a ratio of 1:1. The
oil/emulsifier mixture is then mixed with krill powder in
an amount of 55% and 45% respectively. The mixture is then
wet milled under nitrogen at 50°C for 15 minutes to give a
slurry. The resulting slurry is then pressed under 200bars
30 of pressure to yield oil enriched with long-chain
polyunsaturated fatty acids in the form of phospholipids
and carotenoids.

Therefore, the use of at least one emulsifier in an oil for improving the extraction of long-chain polyunsaturated fatty acids in the form of phospholipids from a source is
5 also part of the present invention.

Indeed, it has been shown that the yield of extraction of the long-chain polyunsaturated fatty acids is greatly improved when at least one emulsifier is used. This is all
10 the more advantageous over the prior art since extraction of long-chain polyunsaturated fatty acids in the form of phospholipids have hitherto been poorly extracted.

Using the extraction process of the invention, an oil
15 containing a carotenoid can be obtained depending on the source used. For example, when using the preferred sources of the invention, the oil obtained by the process contains astaxanthin. This carotenoid is co-extracted into the oil when carrying out the process of the invention and
20 provides the oil with additional advantages since carotenoids generally are potent natural antioxidants.

Oils comprising long-chain polyunsaturated fatty acids and an antioxidant can thus be obtained by the present method.
25

According to an aspect of the invention, an oil composition comprising long-chain polyunsaturated fatty acids in an amount up to 4% and phospholipids in an amount of up to 15% is provided. Other components include
30 carotenoids.

The amounts of polyunsaturated fatty acids and phospholipids in the present oil composition correspond to at least 80%, preferably at least 90% of the theoretically extractable amount of long-chain polyunsaturated fatty acids and phospholipids from a source.

The oil composition of the present invention may be for use in cosmetic and/or therapeutic treatment. Such oil can be used as such in food stuffs (e.g. infant formulas, baby foods, dairy products, culinary products, cocoa beverages, chocolate, ice cream, cereal bars) or in the form of capsules, gelules, or encapsulated in glassy matrices.

Examples

15

Example

Two comparative examples were carried out. The conditions used were the same in both examples, i.e. wet milling of the source and oil under a nitrogen atmosphere at 50°C for 15 seconds, followed by pressing using a conventional hydraulic press to 200bars, and collecting the extracted oil.

In the first case, the extraction process was carried out with MCT oil alone. In the second case, the same process steps were carried out, except that the MCT oil was mixed with a monoglyceride emulsifier. The resulting extraction yields are shown in the table.

30

	Theoretical %	Measured %	Yield
--	---------------	------------	-------

Source/oil mixture ratio	docosahexaenoic acid (DHA) if 100% extraction	DHA in extracted oil	(%)
45:55 Krill:MCT	1.4	0.43	30.7
45:55 Krill:MCT+Dimodan	1.4	1.15	82.15

Table 1: DHA content in oil extracted with or without emulsifier.

5 The results in the table above clearly show the improvement in the extraction yield of long-chain fatty acids when using an oil/emulsifier mixture in the extraction of long-chain polyunsaturated fatty acid (DHA). The following table shows the difference of fatty acid profile of our extract (45% Shrimp powder/ 55% mix MCT/Dimodon U) and a shrimp oil extracted with organic solvent (Chloroforme/Methanol).

10

Fatty acids	g/100g oil (solvent extract)	g/100g oil (Soft extract 45/55)
Palmitic acid	7.80	3
Stearic acid	1.85	1.60
Oleic acid	11.65	10.2
Linoleic	0.60	18.44
Linolenic	0.25	0.10
Eicosaenoic acid (EPA)	9.60	1.25
Docosahexaenoic	8.45	1.15

(DHA)		
Others	59.8	64.26

Table 2: Fatty acid profile of solvent extracted shrimp oil and soft (vegetable oil + emulsifier) extracted shrimp oil.

5

Note that the volatile profile of soft extract and commercial solvent extract are also different. As depicted below in the soft extract no acetone nor ethanol residues can be detected by SPME-GCMS.

10

Claims

- 5
1. Method to extract long-chain polyunsaturated fatty acids from a source comprising the steps of
- a. mixing the source with an oil comprising at least one emulsifier
- 10 b. milling the mixture to form a slurry and
- c. pressing the resulting slurry to obtain an oil comprising said long-chain polyunsaturated fatty acids.
- 15 2. Method according to claim 1, wherein the long-chain polyunsaturated fatty acids are linoleic acid, alpha-linolenic acid, dihomogammalinolenic acid, arachidonic acid, eicosapentaenoic acid, docosahexaenoic acid and mixtures thereof.
- 20 3. Method according to claims 1 or 2, wherein the long-chain polyunsaturated fatty acids are in the form of phospholipids.
- 25 4. Method according to any of the preceding claims, wherein the source is a dried biomass, for example selected from zooplanktons, crustaceans, microalgae, fungi or by-products from fisheries.
- 30 5. Method according to any of the preceding claims, wherein the oil is selected from MCT oil, edible vegetable oil, essential oil or mixtures thereof.

- 5 6. Method according to any of the preceding claims,
wherein the emulsifiers are selected from
emulsifiers having a hydrophilic-lipophilic balance
of less than 7.
- 10 7. Method according to any of the preceding claims,
wherein the emulsifiers are selected from
monoglycerides, diglycerides, sorbitan monoesters
such as sorbitan monolaurate, sorbitan
monopalmitate, sorbitan monooleate, sorbitan
monostearate, polypropylene glycol monoesters ,
polyglycerol esters or any mixtures thereof.
- 15 8. Method according to any of the preceding claims,
wherein the pressing is carried out under from 10 to
500 bars.
- 20 9. Use of at least one emulsifier in oil for improving
the extraction yield of long-chain polyunsaturated
fatty acids in the form of phospholipids from a
biomass.
- 25 10. Oil composition obtainable by a method according
to any of claims 1 to 8.
- 30 11. Oil composition comprising per 100g oil
composition about 2-4 g palmitic acid, about 0,5 - 4
g stearic acid, about 8-12 g oleic acid, about 14-
22 g linoleic acid, about 0-0,5 g linolenic acid,

about 0,3 - 4 g eicosaenoic acid (EPA), and about 0,3 - 4 g docosahexaenoic acid (DHA).

- 5 12. Oil composition comprising long-chain polyunsaturated fatty acids in an amount up to 4%, phospholipids in an amount up to 10%, and at least one emulsifier in an amount up to 50%.
- 10 13. Oil composition according to one of claims 10 to 12, which comprises at least one carotenoid.
- 15 14. Oil composition according to any of claims 10 to 13, for use in cosmetic and/or therapeutic treatment.
- 15 15. Use of an oil composition according to any of claims 10 to 14 in food.

1 / 2




FIG. 1




FIG. 2