



- (72) BAYER, Herbert, DE
(72) SAUTER, Hubert, DE
(72) AMMERMANN, Eberhard, DE
(72) LORENZ, Gisela, DE
(72) STRATHMANN, Siegfried, DE
(72) KÖHLE, Harald, DE
(72) RETZLAFF, Günter, DE
(71) BASF AKTIENGESELLSCHAFT, DE
(51) Int.Cl.⁶ A01N 43/54, A01N 43/72, A01N 43/64, A01N 43/56,
A01N 37/50, A01N 37/36, A01N 47/20
(30) 1995/09/25 (195 35 516.4) DE
(54) **LUTTE CONTRE DES CHAMPIGNONS NUISIBLES GRACE A
L'ASSOCIATION DE LA FENAZAQUINE AVEC UN PRINCIPE
ACTIF INHIBANT LA RESPIRATION AU NIVEAU DU
COMPLEXE CYTOCHROME III**
(54) **COMBATTING PARASITIC FUNGI WITH A COMBINATION
OF AN ACTIVE AGENT INHIBITING RESPIRATION IN THE
CYTOCHROME COMPLEX III AND OF FENAZAQUINE**

(57) L'invention concerne des agents permettant de lutter contre des champignons nuisibles, qui contiennent pour principes actifs au moins un composé inhibant la respiration au niveau du complexe cytochrome III, et de la fénazaquine. Ces agents s'utilisent notamment pour lutter contre le botrytis.

(57) The present invention relates to means for combatting parasitic fungi, containing as the active agents at least one compound which prevents respiration in the cytochrome complex III and fenazaquine. The means of the invention may be used in particular in combatting botrytis.



PCT
WELTORGANISATION FÜR GEISTIGES EIGENTUM
Internationales Büro
INTERNATIONALE ANMELDUNG VERÖFFENTLICHT NACH DEM VERTRAG ÜBER DIE
INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT)

<p>(51) Internationale Patentklassifikation ⁶ : A01N 43/54 // (A01N 43/54, 61:00, 43:56, 43:54, 37:50)</p>	A1	<p>(11) Internationale Veröffentlichungsnummer: WO 97/11606</p> <p>(43) Internationales Veröffentlichungsdatum: 3. April 1997 (03.04.97)</p>
<p>(21) Internationales Aktenzeichen: PCT/EP96/04013</p> <p>(22) Internationales Anmeldedatum: 12. September 1996 (12.09.96)</p> <p>(30) Prioritätsdaten: 195 35 516.4 25. September 1995 (25.09.95) DE</p> <p>(71) Anmelder (für alle Bestimmungsstaaten ausser US): BASF AKTIENGESELLSCHAFT [DE/DE]; D-67056 Ludwigshafen (DE).</p> <p>(72) Erfinder; und (75) Erfinder/Anmelder (nur für US): BAYER, Herbert [DE/DE]; D 3.4, D-68159 Mannheim (DE). SAUTER, Hubert [DE/DE]; Neckarpromenade 20, D-68167 Mannheim (DE). AMMERMAN, Eberhard [DE/DE]; Von-Gagern-Strasse 2, D-64646 Heppenheim (DE). LORENZ, Gisela [DE/DE]; Erlenweg 13, D-67434 Hambach (DE). STRATHMANN, Siegfried [DE/DE]; Donnersbergstrasse 9, D-67117 Limburgerhof (DE). KÖHLE, Harald [DE/DE]; Am Büschel 13, D-67273 Bobenheim (DE). RETZLAFF, Günter [DE/DE]; Schillerstrasse 34, D-67354 Römerberg (DE).</p> <p>(74) Anwalt: REITSTÖTTER, KINZEBACH & PARTNER; Postfach 86 06 49, D-81633 München (DE).</p>	<p>(81) Bestimmungsstaaten: AU, BR, CA, CN, CZ, HU, IL, JP, KR, MX, NZ, PL, RU, SK, UA, US, europäisches Patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Veröffentlicht <i>Mit internationalem Recherchenbericht. Vor Ablauf der für Änderungen der Ansprüche zugelassenen Frist. Veröffentlichung wird wiederholt falls Änderungen eintreffen.</i></p>	
<p>(54) Title: COMBATTING PARASITIC FUNGI WITH A COMBINATION OF AN ACTIVE AGENT INHIBITING RESPIRATION IN THE CYTOCHROME COMPLEX III AND OF FENAZAQUINE</p> <p>(54) Bezeichnung: SCHADPILZBEKÄMPFUNG MIT DER KOMBINATION EINES DIE ATMUNG AM CYTOCHROM-KOMPLEX III HEMMENDEN WIRKSTOFFES MIT FENAZAQUIN</p> <p>(57) Abstract The present invention relates to means for combatting parasitic fungi, containing as the active agents at least one compound which prevents respiration in the cytochrome complex III and fenazaquine. The means of the invention may be used in particular in combatting botrytis.</p> <p>(57) Zusammenfassung Die vorliegende Erfindung betrifft Mittel zur Bekämpfung von Schadpilzen, die als Wirkstoffe wenigstens eine Verbindung, welche die Atmung am Cytochrom-Komplex III hemmt, und Fenazaquin enthalten. Die erfindungsgemäßen Mittel sind insbesondere zur Bekämpfung von Botrytis brauchbar.</p>		

5 COMBATTING PARASITIC FUNGI WITH A COMBINATION OF AN ACTIVE AGENT
 INHIBITING RESPIRATION IN THE CYTOCHROME COMPLEX III AND OF
 FENAZAQUINE

10 The present invention relates to compositions for controlling
 harmful fungi and to methods of controlling harmful fungi using
 such compositions.

The literature discloses that active ingredients which inhibit
 15 the cytochrome bc₁ complex (cytochrome complex III) can be
 employed as fungicides [cf. U. Brandt, U. Haase, H. Schagger, G.
 von Jagow: "Spezifität und Wirkmechanismus der Strobilurine"
 (Specificity and Mechanism of Action of the Strobilurins), Deche-
 ma Monograph Vol. 129, 27-38, VCH Verlagsgesellschaft Weinheim,
 20 1993; J.M. Clough: Natural Product Reports, 1993, 565-574; F.
 Röhl, H. Sauter: Biochem. Soc. Trans. 22, 635 (1993)].

However, when using these active ingredients as fungicides, it
 has emerged that their activity is only transient, i.e. that re-
 25 newed growth of the fungi was observed after a short time.


It is therefore an object of the present invention to provide an
 improved possibility of controlling harmful fungi, in particular
 botrytis.

30 Surprisingly, we have found that this object is achieved by a
 composition which comprises an active ingredient which inhibits
 respiration on the cytochrome complex III in combination with fe-
 nazaquin, which is known as being acaricidally active (cf. The
 35 Pesticide Manual, 10th Edition, 1994; CAS reg. No. 120928-09-8).

The present invention therefore relates to compositions for con-
 trolling harmful fungi which comprise, in a solid or liquid
 carrier,

- 40 a) at least one active ingredient I, which inhibits respiration
 on the cytochrome complex III, and
 b) fenazaquin, of the formula


45



5


The active ingredient I is preferably a compound of the formula
IA or IB:

10



15

IA



IB

where is a double or single bond;

20 R' is -C[CO₂CH₃]=CHOCH₃, -C[CO₂CH₃]=NOCH₃,
-C[CONHCH₃]=NOCH₃, -C[CO₂CH₃]=CHCH₃, -C[CO₂CH₃]=CHCH₂CH₃,
-C[COCH₃]=NOCH₃, -C[COCH₂CH₃]=NOCH₃, -N(OCH₃)-CO₂CH₃,
-N(CH₃)-CO₂CH₃ or -N(CH₂CH₃)-CO₂CH₃;

25 R'' is an organic radical which is bonded directly or via an
oxy, mercapto, amino, or alkylamino group, or

30 together with a group X and the ring Q or T to which they
are bonded can contain an unsubstituted or substituted
bicyclic, partially or fully unsaturated system which,
besides carbon ring members, can have 1, 2 or 3 hetero
atoms independently selected from amongst oxygen, sulfur
and nitrogen;

35 R^x is -OC[CO₂CH₃]=CHOCH₃, -OC[CO₂CH₃]=CHCH₃,
-OC[CO₂CH₃]=CHCH₂CH₃, -SC[CO₂CH₃]=CHOCH₃,
-SC[CO₂CH₃]=CHCH₃, -SC[CO₂CH₃]=CHCH₂CH₃,
-N(CH₃)C[CO₂CH₃]=CHOCH₃, -N(CH₃)C[CO₂CH₃]=NOCH₃,
-CH₂C[CO₂CH₃]=CHOCH₃, -CH₂C[CO₂CH₃]=NOCH₃ or
-CH₂C[CONHCH₃]=NOCH₃;

40 R^y is oxygen, sulfur, =CH- or =N-;
n is 0, 1, 2 or 3, it being possible for the radicals X to
be identical or different if n > 1;

X is cyano, nitro, halogen, alkyl, haloalkyl, alkoxy, ha-
loalkoxy, alkylthio, or,

45 if n > 1, a C₃-C₅-alkylene, C₃-C₅-alkenylene, oxy-
C₂-C₄-alkylene, oxy-C₁-C₃-alkyleneoxy, oxy-C₂-C₄-alkeny-
lene, oxy-C₂-C₄-alkenyleneoxy or butadienediyl group

3

which is bonded to two adjacent C atoms of the phenyl ring, it being possible for these chains, in turn, to have attached to them one to three radicals which, independently of one another, are selected from amongst halogen, alkyl, haloalkyl, alkoxy, haloalkoxy and alkylthio;

Y is =C- or -N-;

Q is phenyl, pyrrolyl, thienyl, furyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, triazolyl, pyridinyl, 2-pyridinyl, pyrimidinyl or triazinyl;

and

T is phenyl, oxazolyl, thiazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl or triazinyl.

15 The substituent R" is, in particular, an alkyl, alkenyl, alkynyl, aryl, hetaryl, arylalkyl, hetarylalkyl, arylalkenyl, hetarylalkenyl, arylalkynyl or hetarylalkynyl radical which may be interrupted by one or more groups selected from amongst O, S, SO, SO₂, NR (R = H or alkyl), CO, COO, OCO, CONH, NHCO and NHCONH, or

20 a radical of the formulae which are as defined below $\text{CH}_2\text{ON} = \text{CR}^\alpha\text{CR}^\beta$ or $\text{CH}_2\text{ON} = \text{CR}^\delta\text{CR}^\epsilon = \text{NOR}^\epsilon$. These radicals may also have one or more (preferably 1, 2 or 3) substituents which, independently of one another, are selected from amongst alkyl, alkoxy, halogen, haloalkyl (in particular CF₃ and CHF₂) and aryl. The latter, in

25 turn, can have, 1, 2 or 3 substituents which, independently of one another, are selected from amongst halogen, haloalkyl (in particular CF₃ and CHF₂), phenyl, CN and phenoxy.

Such compounds and their preparation are described in the literature given in Tables I.1 to I.8 below. Compounds which are not described in these tables can be prepared by similar methods.

Halogen within the scope of the present invention is fluorine, chlorine, bromine and iodine, in particular fluorine, chlorine and bromine.

The term "alkyl" encompasses straight-chain and branched alkyl groups. They are preferably straight-chain or branched C₁-C₁₂-alkyl and in particular C₁-C₆-alkyl groups. Examples of

40 alkyl groups are alkyl such as, in particular, methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1-methylpentyl, 2-methylpentyl,

45 3-methylpentyl, 4-methylpentyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethyl-

propyl, 1-ethylbutyl, 2-ethylbutyl, 1-ethyl-2-methylpropyl, n-heptyl, 1-methylhexyl, 1-ethylpentyl, 2-ethylpentyl, 1-propylbutyl, octyl, decyl, dodecyl.

5 Haloalkyl is an alkyl group as defined above which is partially or fully halogenated by one or more halogen atoms, in particular fluorine and chlorine. There are preferably 1 to 3 halogen atoms present, the difluoromethyl or trifluoromethyl group being specially preferred.

10

What has been said above for the alkyl group and haloalkyl group applies analogously to the alkyl and haloalkyl group in alkoxy, haloalkoxy, alkylthio and haloalkylthio or similar groups.

15 The alkenyl group encompasses straight-chain and branched alkenyl groups. They are preferably straight-chain or branched C₃-C₁₂-alkenyl groups, in particular C₃-C₆-alkenyl groups. Examples of alkenyl groups are 2-propenyl, 2-butenyl, 3-butenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 2-pentenyl, 3-pentenyl, 20 4-pentenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-2-propenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 25 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 30 2,2-dimethyl-3-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl and 1-ethyl-2-methyl-2-propenyl, in 35 particular 2-propenyl, 2-butenyl, 3-methyl-2-butenyl and 3-methyl-2-pentenyl.

The alkenyl group can be partially or fully halogenated by one or more halogen atoms, in particular fluorine and chlorine. It has 40 preferably 1 to 3 halogen atoms.

The alkynyl group encompasses straight-chain and branched alkynyl groups. They are preferably straight-chain or branched C₃-C₁₂-alkynyl groups, in particular C₃-C₆-alkynyl groups.

45 Examples of alkynyl groups are 2-propynyl, 2-butylnyl, 3-butylnyl, 1-methyl-2-propynyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-methyl-3-butylnyl, 2-methyl-3-butylnyl, 1-methyl-2-butylnyl,

5

1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 2-hexynyl, 3-hexynyl, 4-alkynyl, 5-hexynyl, 1-methyl-2-pentynyl, 1-methyl-3-pentynyl, 1-methyl-4-pentynyl, 2-methyl-3-pentynyl, 2-methyl-4-pentynyl, 3-methyl-4-pentynyl, 4-methyl-2-pentynyl, 1,2-dimethyl-2-butynyl, 1,1-dimethyl-3-butynyl, 1,2-dimethyl-3-butynyl, 2,2-dimethyl-3-butynyl, 1-ethyl-2-butynyl, 1-ethyl-3-butynyl, 2-ethyl-3-butynyl and 1-ethyl-1-methyl-2-propynyl.

What has been said above about the alkenyl group and its halogen substituents, and about the alkynyl group, applies analogously to alkenyloxy and alkynyloxy.

The cycloalkyl group is preferably a C₃-C₆-cycloalkyl group, such as cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl. If the cycloalkyl group is substituted, it has preferably 1 to 3 C₁-C₄-alkyl radicals as substituents.

Cycloalkenyl is preferably a C₄-C₆-cycloalkenyl group, such as cyclobutenyl, cyclopentenyl or cyclohexenyl. If the cycloalkenyl group is substituted, it has preferably 1 to 3 C₁-C₄-alkyl radicals as substituents.

A cycloalkoxy group is preferably a C₅-C₆-cycloalkoxy group, such as cyclopentyloxy or cyclohexyloxy. If the cycloalkoxy group is substituted, it has preferably 1 to 3 C₁-C₄-alkyl radicals as substituents.

The cycloalkenyloxy group is preferably a C₅-C₆-cycloalkenyloxy group, such as cyclopentyloxy or cyclohexyloxy. If the cycloalkenyloxy group is substituted, it has preferably 1 to 3 C₁-C₄-alkyl radicals as substituents.

Aryl is preferably phenyl.



Hetaryl is preferably a 5- or 6-membered aromatic heterocycle which has 1, 2 or 3 hetero atoms which, independently of one another, are selected from amongst N, O and S. It is, in particular, pyridynyl, pyrimidynyl, thiazolyl or pyrazolyl.

Heterocyclyl is preferably a 5- or 6-membered saturated or unsaturated heterocycle which has 1, 2 or 3 hetero atoms which, independently of one another, are selected from amongst N, O and S. It is, in particular, one of the dihydro, tetrahydro and hexahydro derivatives of the radical mentioned under "hetaryl". Preferred are pyrrolidynyl, tetrahydrofuranyl, imidazolidynyl, pyrazolidynyl, oxazolidynyl, isoxazolidynyl, thiazolidynyl, isothiazolidynyl, piperidynyl or morpholynyl.

6

In a preferred embodiment, the compositions according to the invention comprise a compound of the formula IA or IB where R" is aryloxy, hetaryloxy, aryloxymethylene, hetaryloxymethylene, arylethenylene or hetarylethenylene, it being possible for these radicals to have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, halogen, CF₃, CHF₂, CN, alkoxy and phenyl which, in turn, can have 1, 2 or 3 substituents which, independently of one another, are selected from amongst halogen, CF₃, CHF₂, phenyl, CN, phenoxy, alkyl, alkoxy and haloalkoxy;

or R" is



15

where R^α is alkyl;

R^β is phenyl, pyridyl or pyrimidyl which may have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, alkoxy, halogen, haloalkoxy, CF₃ and CHF₂;

R^γ is alkyl, alkoxy, halogen, haloalkyl or hydrogen;

R^δ is hydrogen, cyano, halogen, alkyl, alkoxy, alkylthio, alkylamino, dialkylamino, alkenyl, alkenyloxy, alkenylthio, alkenylamino, N-alkenyl-N-alkylamino, alkynyl, alkynyloxy, alkynylthio, alkynylamino, N-alkynyl-N-alkylamino, it being possible for the hydrocarbon radicals of these groups to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 radicals which, independently of one another, are selected from amongst cyano, nitro, hydroxyl, alkoxy, haloalkoxy, alkoxycarbonyl, alkylthio, alkylamino, di-alkylamino, alkenyloxy, cycloalkyl, cycloalkyloxy, heterocyclyl, heterocyclyloxy, aryl, aryloxy, arylalkoxy, hetaryl, hetaryloxy and hetarylalkoxy, it being possible for the cyclic radicals, in turn, to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 groups which, independently of one another, are selected from amongst cyano, nitro, hydroxyl, alkyl, haloalkyl, cycloalkyl, alkoxy, haloalkoxy, alkoxycarbonyl, alkylthio, alkylamino, dialkylamino, alkenyl and alkenyloxy;

or
 45

7

is cycloalkyl, cycloalkyloxy, cycloalkylthio, cycloalkylamino, N-cycloalkyl-N-alkylamino, heterocyclyl, heterocyclyoxy, heterocyclylthio, heterocyclylamino, N-heterocyclyl-N-alkylamino, aryl, aryloxy, arylthio, arylamino, N-aryl-N-alkylamino, hetaryl, hetaryloxy, hetarylthio, hetarylamino or N-hetaryl-N-alkylamino, it being possible for the cyclic radicals to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 groups which, independently of one another, are selected from amongst cyano, nitro, hydroxyl, alkyl, haloalkyl, cycloalkyl, alkoxy, haloalkoxy, alkoxycarbonyl, alkylthio, alkylamino, di-alkylamino, alkenyl, alkenyloxy, benzyl, benzyloxy, aryl, aryloxy, hetaryl and hetaryloxy, it being possible for the aromatic radicals, in turn, to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 of the following groups: cyano, alkyl, haloalkyl, alkoxy, nitro;

R^E is alkyl, alkenyl or alkynyl, it being possible for these groups to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 of the following radicals: cyano, alkoxy, cycloalkyl.

Especially preferred compounds of the formula IA or IB are those where


25 R'' has one of the following meanings:

- a) phenyloxymethylene, pyridinyloxymethylene, pyrimidinyloxymethylene or pyrazolyloxymethylene, it being possible for the aromatic radical to have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, halogen, CF_3 , CHF_2 , $-C(CH_3)=NOCH_3$ and phenyl which is unsubstituted or substituted by 1, 2 or 3 halogen atoms and/or alkyl groups;
- 35 b) phenoxy or pyrimidinyloxy, unsubstituted or substituted by 1, 2 or 3 halogen atoms or one phenoxy radical which may have a halogen or cyano substituent;
- 40 c) phenylethenylene or pyrazolyethenylene, it being possible for the phenyl or pyrazolyl radical to have 1, 2 or 3 substituents which, independently of one another, are selected from amongst halogen, CF_3 , CHF_2 and phenyl.
- d) $CH_2ON=CR^aR^b$,
45 where R^a is alkyl; and

8

R^{β} is phenyl which may have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, halogen, CF_3 and CHF_2 , or is pyrimidynyl which is unsubstituted or substituted by 1 or 2 alkoxy radicals;

5



where R^{γ} is alkyl, alkoxy or halogen;

R^{δ} is alkyl, cyano, halogen, alkoxy, alkenyl or phenyl which is unsubstituted or substituted by 1, 2 or 3 halogen atoms; and

R^{ϵ} is alkyl.

15 Particularly preferred compounds are those of the formula IA, where Q is phenyl and n is 0.

Especially suitable active ingredients I are compiled in the tables which follow.

20 Table I.1A

Compounds of the formula IA, where Q is phenyl, R' is $-C(CO_2CH_3)=CHOCH_3$, n has the value 0, R'' is unsubstituted or substituted (het)aryloxymethylene, the unsubstituted or substituted (het)aryl group having the meanings given below

25

No.	Unsubst. or subst. (het)aryl	Reference
30 I.1A-1	2- CH_3 - C_6H_4	EP-A 226 917
I.1A-2	2,5-(CH_3) $_2$ - C_6H_3	EP-A 226 917
I.1A-3	2- CH_3 , 4-C[CH_3]=NO CH_3 - C_6H_3	EP-A 386 561
35 I.1A-4	2- $CH_2CH_2CH_3$, 6- CF_3 -pyrimidin-4-yl	EP-A 407 873
I.1A-5	2,4-(CH_3) $_2$ - C_6H_3	EP-A 226 917

40 Table I.1B:

Compounds of the formula IA, where R' is $-C(CO_2CH_3)=CHOCH_3$, Q is phenyl, n has the value 0, R'' is unsubstituted or substituted (het)aryloxy the unsubstituted or substituted (het)aryl group having the meanings given below

45

No.	Unsubst. or subst. (het)aryl	Reference
I.1B-1	C ₆ H ₅	EP-A 178 826
I.1B-2	6-[2-CN-C ₆ H ₄ -O]-pyrimidin-4-yl	EP-A 382 375

5

Table I.1C:

Compounds of the formula IA, where R' is -C(CO₂CH₃)=CHOCH₃, Q is phenyl, n has the value 0, R'' is unsubstituted or substituted (het)arylethylene, the unsubstituted or substituted (het)aryl group having the meanings given below

10

15	No.	Unsubst. or subst. (het)aryl	Reference
	I.1C-1	1-(2,4-Cl ₂ -C ₆ H ₃), 5-CF ₃ -pyrazol-4-yl	EP-A 528 245
	I.1C-2	1-(4-Cl-C ₆ H ₄)-pyrazol-4-yl	EP-A 378 755
20	I.1C-3	3-CF ₃ -C ₆ H ₄	EP-A 203 606
	I.1C-4	3-Cl-C ₆ H ₄	EP-A 203 606
	I.1C-5	4-C ₆ H ₅ -C ₆ H ₄	EP-A 203 606

25

Table I.1D:

Compounds of the formula IA, where Q is phenyl, R' is -C(CO₂CH₃)=CHOCH₃, n has the value 0, R'' is CH₂ON=CR^αR^β, R^α and R^β having the meanings given below

30

	No.	R ^α	R ^β	Reference
	I.1D-1	CH ₃	4-Cl-C ₆ H ₄	EP-A 370 629
35	I.1D-2	CH ₃	3-CF ₃ -C ₆ H ₄	EP-A 370 629
	I.1D-3	CH ₃	4-OCH ₂ CH ₃ -pyrimidin-2-yl	WO-A 92/18,487

40 Table I.1E:

Compounds of the formula IA, where Q is phenyl, R' is -C(CO₂CH₃)=CHOCH₃, n has the value 0, R'' is CH₂ON=CR^γCR^δ=NOR^ε, R^γ, R^δ and R^ε having the meanings given below

45

10

No.	R ^γ	R ^δ	R ^ε	Reference
I.1E-1	CH ₃	CH ₃	CH ₃	WO-A 95/21153
I.1E-2	CH ₃	CH ₃	CH ₂ CH ₃	WO-A 95/21153
I.1E-3	CH ₃	C ₆ H ₅	CH ₃	WO-A 95/21153
I.1E-4	CH ₃	C ₆ H ₅	CH ₂ CH ₃	WO-A 95/21153
I.1E-5	CH ₃	4-Cl-C ₆ H ₄	CH ₃	WO-A 95/21153
I.1E-6	CH ₃	4-Cl-C ₆ H ₄	CH ₂ CH ₃	WO-A 95/21153

Table I.2A:

- Compounds of the formula IA, where Q is phenyl, R' is
 -C(CO₂CH₃)=NOCH₃, n has the value 0, R'' is unsubstituted or sub-
 stituted (het)aryloxymethylene, the unsubstituted or substituted
 (het)aryl group having the meanings given below

No.	Unsubst. or subst. (het)aryl	Reference
I.2A-1	2-CH ₃ -C ₆ H ₄	EP-A 253 213
I.2A-2	2,5-(CH ₃) ₂ -C ₆ H ₃	EP-A 400 417
I.2A-3	2,4-(CH ₃) ₂ -C ₆ H ₃	EP-A 400 417
I.2A-4	2,3,5-(CH ₃) ₃ -C ₆ H ₂	EP-A 400 417
I.2A-5	2-Cl, 5-CH ₃ -C ₆ H ₃	EP-A 400 417
I.1A-6	2-CH ₃ , 4-C[CH ₃]=NOCH ₃ -C ₆ H ₃	EP-A 386 561

Table I.2B:

- Compounds of the formula IA, where Q is phenyl, R' is
 -C(CO₂CH₃)=NOCH₃, n has the value 0, R'' is unsubstituted or sub-
 stituted (het)aryloxy, the unsubstituted or substituted (het)aryl
 group having the meanings given below

No.	Unsubst. or subst. (het)aryl	Reference
I.2B-1	C ₆ H ₅	EP-A 253 213
I.2B-2	6-[2-CN-C ₆ H ₄ -O]-pyrimidin-4-yl	EP-A 468 684

Table I.2C:

Compounds of the formula IA, where Q is phenyl, R' is
 5 -C(CO₂CH₃)=NOCH₃, n has the value 0, R'' is CH₂ON=CR^αR^β, R^α and R^β
 having the meanings given below

No.	R ^α	R ^β	Reference
10 I.2C-1	CH ₃	4-Cl-C ₆ H ₄	EP-A 463 488
I.2C-2	CH ₃	3-Cl-C ₆ H ₄	EP-A 463 488
I.2C-3	CH ₃	4-CF ₃ -C ₆ H ₄	EP-A 463 488
I.2C-4	CH ₃	3-CF ₃ -C ₆ H ₄	EP-A 463 488
15 I.2C-5	CH ₃	4-CH ₃ -C ₆ H ₄	EP-A 463 488
I.2C-6	CH ₃	4-OCH ₂ CH ₃ -pyrimi- din-2-yl	EP-A 472 300
20 I.2C-7	CH ₃	3,5-Cl ₂ -C ₆ H ₃	EP-A 463 488

Table I.2D:

Compounds of the formula IA, where Q is phenyl, R' is
 25 -C(CO₂CH₃)=NOCH₃, n has the value 0, R'' is CH₂ON=CR^γCR^δ=NOR^ε, R^γ,
 R^δ and R^ε having the meanings given below

No.	R ^γ	R ^δ	R ^ε	Reference
30 I.2D-1	CH ₃	CH ₃	CH ₃	WO-A 95/21153
I.2D-2	CH ₃	CH ₃	CH ₂ CH ₃	WO-A 95/21153
I.2D-3	CH ₃	C ₆ H ₅	CH ₃	WO-A 95/21153
I.2D-4	CH ₃	C ₆ H ₅	CH ₂ CH ₃	WO-A 95/21153
35 I.2D-5	CH ₃	4-Cl-C ₆ H ₄	CH ₃	WO-A 95/21153
I.2D-6	CH ₃	4-Cl-C ₆ H ₄	CH ₂ CH ₃	WO-A 95/21153

Table I.3A:

40

Compounds of the formula IA, where Q is phenyl, R' is
 -C(CONHCH₃)=NOCH₃, n has the value 0, R'' is unsubstituted or sub-
 45 stituted (het)aryloxymethylene, the unsubstituted or substituted
 (het)aryl group having the meanings given below

12

No.	Unsubst. or subst. (het)aryl	Reference
I.3A-1	2-CH ₃ -C ₆ H ₄	EP-A 477 631
I.3A-2	2,5-(CH ₃) ₂ -C ₆ H ₃	EP-A 477 631
I.3A-3	2,4-(CH ₃) ₂ -C ₆ H ₃	EP-A 477 631
I.3A-4	2,3,5-(CH ₃) ₃ -C ₆ H ₂	EP-A 477 631
I.3A-5	2-CH ₃ , 4-C[CH ₃]=NOCH ₃ -C ₆ H ₃	EP-A 579 124
I.3A-6	1-[4-Cl-C ₆ H ₄]-pyrazol-3-yl	WO-A 94/19331
I.3A-7	1-[2,4-Cl ₂ -C ₆ H ₃]-pyrazol-3-yl	WO-A 94/19331

Table I.3B:

Compounds of the formula IA, where Q is phenyl, R' is -C(CONHCH₃)=NOCH₃, n has the value 0, R'' is unsubstituted or substituted (het)aryloxy, the unsubstituted or substituted (het)aryl group having the meanings given below

No.	Unsubst. or subst. (het)aryl	Reference
I.3B-1	C ₆ H ₅	EP-A 398 692
I.3B-2	6-[2-CN-C ₆ H ₄ -O]-pyrimidin-4-yl	GB-A 2 253 624

Table I.3C:

Compounds of the formula IA, where Q is phenyl, R' is -C(CONHCH₃)=NOCH₃, n has the value 0, R'' is unsubstituted or substituted (het)arylethenylene, the unsubstituted or substituted (het)aryl group having the meanings given below

No.	Unsubst. or subst. (het)aryl	Reference
I.3C-1	1-[2,4-Cl ₂ -C ₆ H ₃], 5-CF ₃ -pyrazol-4-yl	DE-A 44 23 615.8

Table I.3D:

Compounds of the formula IA, where Q is phenyl, R' is -C(CONHCH₃)=NOCH₃, n has the value 0, R'' is CH₂ON=CR^αR^β, R^α and R^β having the meanings given below

13

No.	R ^a	R ^B	Reference
I.3D-1	CH ₃	4-Cl-C ₆ H ₄	EP-A 463 488
I.3D-2	CH ₃	3-Cl ₃ -C ₆ H ₄	EP-A 463 488
I.3D-3	CH ₃	4-CF ₃ -C ₆ H ₄	EP-A 585 751
I.3D-4	CH ₃	3-CF ₃ -C ₆ H ₄	EP-A 585 751
I.3D-5	CH ₃	4-CH ₃ -C ₆ H ₄	EP-A 463 488
I.3D-6	CH ₃	3,5-Cl ₂ -C ₆ H ₃	EP-A 463 488
I.3D-7	CH ₃	2-OCH ₂ CH ₃ -pyrimi- din-2-yl	WO-A 92/13,830

15 Table I.3E:

Compounds of the formula IA, where Q is phenyl, R' is
 $-C(CONHCH_3)=NOCH_3$, n has the value 0, R'' is $CH_2ON=CR^YCR^\delta=NOR^E$, R^Y,
 R^δ and R^E having the meanings given below

20

No.	R ^Y	R ^δ	R ^E	Reference
I.3E-1	CH ₃	CH ₃	CH ₃	WO-A 95/21154
I.3E-2	CH ₃	CH ₃	CH ₂ CH ₃	WO-A 95/21154
I.3E-3	CH ₃	C ₆ H ₅	CH ₃	WO-A 95/21154
I.3E-4	CH ₃	C ₆ H ₅	CH ₂ CH ₃	WO-A 95/21154
I.3E-5	CH ₃	4-Cl-C ₆ H ₄	CH ₃	WO-A 95/21154
I.3E-6	CH ₃	4-Cl-C ₆ H ₄	CH ₂ CH ₃	WO-A 95/21154
I.3E-7	CH ₃	4-F-C ₆ H ₄	CH ₃	WO-A 95/21154

35 Table I.4A:

Compounds of the formula IA, where Q is phenyl, R' is
 $-C(CO_2CH_3)=CHCH_3$, n has the value 0, R'' is unsubstituted or sub-
 stituted (het)aryloxymethylene, the unsubstituted or substituted
 (het)aryl group having the meanings given below

40

No.	Unsubst. or subst. (het)aryl	Reference
I.4A-1	2-CH ₃ -C ₆ H ₄	EP-A 280 185
I.4A-2	2,5-(CH ₃) ₂ -C ₆ H ₃	EP-A 513 580
I.4A-3	2,4-(CH ₃) ₂ -C ₆ H ₃	EP-A 513 580
I.4A-4	2,3,5-(CH ₃) ₃ -C ₆ H ₂	EP-A 513 580

45

14

I.4A-5	2-Cl, 5-CH ₃ -C ₆ H ₃	EP-A 513 580
I.4A-6	2-CH ₃ , 4-C[CH ₃]=NOCH ₃ -C ₆ H ₃	EP-A 513 580
I.4A-7	1-[4-Cl-C ₆ H ₄]-pyrazol-3-yl	DE-A 44 15 483.6

5

Table I.4B:

Compounds of the formula IA, where Q is phenyl, R' is
 -C(CO₂CH₃)=CHCH₃, n has the value 0, R'' is unsubstituted or sub-
 10 stituted (het)aryloxy, the unsubstituted or substituted (het)aryl
 group havings the meanings given below

No.	Unsubst. or subst. (het)aryl	Reference
I.4B-1	C ₆ H ₅	EP-A 513 580

15

Table I.4C:

Compounds of the formula IA, where Q is phenyl, R' is
 -C(CO₂CH₃)=CHCH₃, n has the value 0, R'' is CH₂ON=CR^YCR^δ=NOR^ε, R^Y,
 20 R^δ and R^ε having the meanings given below

No.	R ^Y	R ^δ	R ^ε	Reference
I.4C-1	CH ₃	CH ₃	CH ₃	WO-A 95/21153
I.4C-2	CH ₃	CH ₃	CH ₂ CH ₃	WO-A 95/21153
I.4C-3	CH ₃	C ₆ H ₅	CH ₃	WO-A 95/21153
I.4C-4	CH ₃	C ₆ H ₅	CH ₂ CH ₃	WO-A 95/21153
I.4C-5	CH ₃	4-Cl-C ₆ H ₄	CH ₃	WO-A 95/21153
I.4C-6	CH ₃	4-Cl-C ₆ H ₄	CH ₂ CH ₃	WO-A 95/21153

25

30

35 Table I.5A:

Compounds of the formula IA, where Q is phenyl, R' is
 -C(CO₂CH₃)=CHCH₂CH₃, n has the value 0, R'' is unsubstituted or
 substituted (het)aryloxymethylene, the unsubstituted or substi-
 40 tuted (het)aryl group having the meanings given below

No.	Unsubst. or subst. (het)aryl	Reference
I.5A-1	2-CH ₃ -C ₆ H ₄	EP-A 513 580
I.5A-2	2,5-(CH ₃) ₂ -C ₆ H ₃	EP-A 513 580

45

15

5	I.5A-3	2,4-(CH ₃) ₂ -C ₆ H ₃	EP-A 513 580
	I.5A-4	2,3,5-(CH ₃) ₃ -C ₆ H ₂	EP-A 513 580
	I.5A-5	2-Cl, 5-CH ₃ -C ₆ H ₃	EP-A 513 580
	I.5A-6	2-CH ₃ , 4-C[CH ₃]=NOCH ₃ -C ₆ H ₃	EP-A 513 580

Table I.5B:

- 10 Compounds of the formula IA, where Q is phenyl, R' is -C(CO₂CH₃)=CHCH₂CH₃, n has the value 0, R'' is unsubstituted or substituted (het)aryloxy, the unsubstituted or substituted (het)aryl group having the meanings given below

15	No.	Unsubst. or subst. (het)aryl	Reference
	I.5B-1	C ₆ H ₅	EP-A 513 580

Table I.5C:

- 20 Compounds of the formula IA, where Q is phenyl, R' is -C(CO₂CH₃)=CHCH₂CH₃, n has the value 0, R'' is CH₂ON=CR^YCR^δ=NOR^ε, R^Y, R^δ and R^ε having the meanings given below

25	No.	R ^Y	R ^δ	R ^ε	Reference
	I.5C-1	CH ₃	CH ₃	CH ₃	WO-A 957 21153
	I.5C-2	CH ₃	CH ₃	CH ₂ CH ₃	WO-A 95/21153
	I.5C-3	CH ₃	C ₆ H ₅	CH ₃	WO-A 95/21153
	I.5C-4	CH ₃	C ₆ H ₅	CH ₂ CH ₃	WO-A 95/21153
	I.5C-5	CH ₃	4-Cl-C ₆ H ₄	CH ₃	WO-A 95/21153
30	I.5C-6	CH ₃	4-Cl-C ₆ H ₄	CH ₂ CH ₃	WO-A 95/21153
	I.5C-6	CH ₃	4-Cl-C ₆ H ₄	CH ₂ CH ₃	WO-A 95/21153

Table I.6A

- 40 Compounds of the formula IA, where Q is phenyl, R' is -C(COCH₃)=NOCH₃, n has the value 0, R'' is unsubstituted or substituted (het)aryloxymethylene, the unsubstituted or substituted (het)aryl group having the meanings given below

45	No.	Unsubst. or subst. (het)aryl	Reference
	I.6A-1	2-CH ₃ -C ₆ H ₄	EP-A 498 188
	I.6A-2	2,5-(CH ₃) ₂ -C ₆ H ₃	EP-A 498 188

16

5	I.6A-3	2,4-(CH ₃) ₂ -C ₆ H ₃	EP-A 498 188
	I.6A-4	2,3,5-(CH ₃) ₃ -C ₆ H ₂	EP-A 498 188
	I.6A-5	2-CH ₃ , 4-C[CH ₃]=NOCH ₃ -C ₆ H ₃	EP-A 498 188

Table I.6B:

Compounds of the formula IA, where Q is phenyl, R' is
 10 -C(COCH₃)=NOCH₃, n has the value 0, R'' is unsubstituted or substituted (het)aryloxy, the unsubstituted or substituted (het)aryl group having the meanings given below

15	No.	Unsubst. or subst. (het)aryl	Reference
	I.6B-1	C ₆ H ₅	EP-A 498 188
	I.6B-2	6-[2-CN-C ₆ H ₄ -O]-pyrimidin-4-yl	EP-A 498 188

Table I.7A:

20 Compounds of the formula IA, where Q is phenyl, R' is
 -C(COCH₂CH₃)=NOCH₃, n is the value 0, R'' is unsubstituted or substituted (het)aryloxymethylene, the unsubstituted or substituted
 25 (het)aryl group having the meanings given below

30	No.	Unsubst. or subst. (het)aryl	Reference
	I.7A-1	2-CH ₃ -C ₆ H ₄	EP-A 498 188
	I.7A-2	2,5-(CH ₃) ₂ -C ₆ H ₃	EP-A 498 188
	I.7A-3	2,4-(CH ₃) ₂ -C ₆ H ₃	EP-A 498 188
	I.7A-4	2,3,5-(CH ₃) ₃ -C ₆ H ₂	EP-A 498 188
35	I.7A-5	2-CH ₃ , 4-C[CH ₃]=NOCH ₃ -C ₆ H ₃	EP-A 498 188

Table I.7B:

Compounds of the formula IA, where Q is phenyl, R' is
 40 -C(COCH₂CH₃)=NOCH₃, n has the value 0, R'' is unsubstituted or substituted (het)aryloxy, the unsubstituted or substituted
 (het)aryl group having the meanings given below

45

No.	Unsubst. or subst. (het)aryl	Reference
I.7B-1	C ₆ H ₅	EP-A 498 188
I.7B-2	6-[2-CN-C ₆ H ₄ -O]-pyrimidin-4-yl	EP-A 498 188

Table I.8A:

Compounds of the formula IA, where Q is phenyl, R' is
-N(OCH₃)-CO₂CH₃, n has the value 0, R'' is unsubstituted or sub-
stituted (het)aryloxymethylene, the unsubstituted or substituted
(het)aryl group having the meanings given below

No.	Unsubst. or subst. (het)aryl	Reference
I.8A-1	2-CH ₃ -C ₆ H ₄	WO-A 93/15,046
I.8A-2	2,5-(CH ₃) ₂ -C ₆ H ₃	WO-A 93/15,046
I.8A-3	2,4-(CH ₃) ₂ -C ₆ H ₃	WO-A 93/15,046
I.8A-4	2,3,5-(CH ₃) ₃ -C ₆ H ₂	WO-A 93/15,046
I.8A-5	2-Cl, 5-CH ₃ -C ₆ H ₃	WO-A 93/15,046
I.8A-6	2-CH ₃ , 4-C[CH ₃]=NOCH ₃ -C ₆ H ₃	WO-A 93/15,046
I.8A-7	2-CH ₃ , 4-C[CH ₃]=NOCH ₂ CH ₃ -C ₆ H ₃	WO-A 93/15,046
I.8A-8	2-CH ₃ , 4-C[CH ₂ CH ₃]=NOCH ₃ -C ₆ H ₃	WO-A 93/15,046
I.8A-9	2-CH ₃ , 4-C[CH ₂ CH ₃]=NOCH ₂ CH ₃ -C ₆ H ₃	WO-A 93/15,046
I.8A-10	1-[4-Cl-C ₆ H ₄]-pyrazol-3-yl	DE-A 44 23 612.3

Table I.8B:

Compounds of the formula IA, where Q is phenyl, R' is
-N(OCH₃)=CO₂CH₃, n has the value 0, R'' is CH₂ON=CR^αR^β, R^α and R^β
have the meanings given below

No.	R ^α	R ^β	Reference
I.8B-1	CH ₃	3,5-Cl ₂ -C ₆ H ₃	WO-A 93/15,046

To produce a synergistic effect, fenazaquin and the active ingredient I are employed in a weight ratio in a range of from 20:1 to 1:20, in particular 10:1 to 1:10.

The invention also relates to a method of controlling harmful fungi, which comprises treating the fungi, their environment, or the materials, plants, seeds, soils, areas or spaces to be pro-

tected against fungal infection, with a composition as defined above, it being possible for the application of the active ingredients to be effected simultaneously, i.e. jointly or separately, or in succession.

5

The compositions according to the invention can be applied, for example, in the form of directly sprayable solutions, powders, suspensions, also highly concentrated aqueous, oily or other suspensions or dispersions, emulsions, oil dispersions, pastes, 10 dusts, materials for spreading or granules by means of spraying, atomizing, dusting, spreading or pouring. The use forms depend on the intended purposes; in any case, they should guarantee the finest possible distribution of the active ingredients according to the invention.

15

Normally, the plants are sprayed or dusted with the active ingredients or the seeds of the plants are treated with the active ingredients.

20

The formulations are prepared in a known manner, e.g. by extending the active ingredient with solvents and/or carriers, if desired using emulsifiers and dispersants, it also being possible for other organic solvents to be used as auxiliary solvents if 25 water is used as the diluent. Suitable auxiliaries are essentially: solvents, such as aromatics (e.g. xylene), chlorinated aromatics (e.g. chlorobenzenes), paraffins (e.g. mineral oil fractions), alcohols (e.g. methanol, butanol), ketones (e.g. cyclohexanone), amines (e.g. ethanolamine, dimethylformamide) and 30 water; carriers, such as ground natural minerals (e.g. kaolins, clays, talc, chalk) and ground synthetic minerals (e.g. highly-disperse silica, silicates); emulsifiers, such as non-ionic and anionic emulsifiers (e.g. polyoxyethylene fatty alcohol ethers, alkylsulfonates and arylsulfonates) and dispersants, such as lig- 35 nin-sulfite waste liquors and methylcellulose.

Suitable surfactants are the alkali metal, alkaline earth metal and ammonium salts of aromatic sulfonic acids, e.g. ligno-, phenol-, naphthalene- and dibutylnaphthalenesulfonic acid, and of 40 fatty acids, alkyl- and alkylarylsulfonates, alkyl sulfates, lauryl ether sulfates and fatty alcohol sulfates; and salts of sulfated hexa-, hepta- and octadecanols, and also fatty alcohol glycol ethers, condensates of sulfonated naphthalene and its derivatives with formaldehyde, condensates of naphthalene, or of 45 the naphthalenesulfonic acids, with phenol and formaldehyde, polyoxyethylene octylphenyl ether, ethoxylated iso-octyl-, octyl- or nonylphenol, alkylphenyl polyglycol ethers and tributylphenyl

19

polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers or polyoxypropylene alkyl ethers, lauryl alcohol polyglycol ether acetate, sorbitol esters, 5 lignin-sulfite waste liquors or methylcellulose.

Powders, materials for spreading and dusts can be prepared by mixing or concomitantly grinding the active substances with a solid carrier.

10

Granules, e.g. coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active ingredients onto solid carriers. Solid carriers are mineral earths such as silica gel, silicas, silica gels, silicates, talc, kaolin, 15 limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate and magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, such as ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal 20 and nutshell meal, cellulose powders, or other solid carriers. Examples of such preparations which comprise the active ingredients in a weight ratio of 1:1 are:

25 I. a solution of 90 parts by weight of the active ingredients and 10 parts by weight of N-methylpyrrolidone which is suitable for use in the form of microdrops;

II. a mixture of 20 parts by weight of the active ingredients, 80 parts of xylene, 10 parts by weight of the adduct of 8 to 10 30 mol of ethylene oxide to 1 mol of oleic acid N-monoethanolamide, 5 parts by weight of calcium dodecylbenzenesulfonate, 5 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil; a dispersion is obtained by finely dividing the solution in water;

35 III. an aqueous dispersion of 20 parts by weight of the active ingredients, 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil;

IV. an aqueous dispersion of 20 parts by weight of the active 40 ingredients, 25 parts by weight of cyclohexanol, 65 parts by weight of a mineral oil fraction of boiling point 210 to 280°C and 10 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil;

V. a mixture, ground in a hammer mill, of 80 parts by weight 45 of the active ingredients, 3 parts by weight of sodium diisobutyl-naphthalene-1-sulfonate, 10 parts by weight of the sodium salt

20

of a lignosulfonic acid from a sulfide waste liquor and 7 parts by weight of pulverulent silica gel; a spray mixture is obtained by finely dividing the mixture in water;

VI. an intimate mixture of 3 parts by weight of the active ingredients and 97 parts by weight of finely divided kaolin; this dust comprises 3% by weight of active ingredient;

VII. an intimate mixture of 30 parts by weight of the active ingredients, 92 parts by weight of pulverulent silica gel and 8 parts by weight of paraffin oil which had been sprayed onto the surface of this silica gel; this formulation imparts good adhesion properties to the active ingredient;

VIII. a stable aqueous dispersion of 40 parts by weight of the active ingredients, 10 parts by weight of the sodium salt of a phenylsulfonic acid/urea/formaldehyde condensate, 2 parts by weight of silica gel and 48 parts by weight of water, it being possible for this dispersion to be diluted further;

IX. a stable or oily dispersion of 20 parts by weight of the active ingredients, 2 parts by weight of calcium dodecylbenzenesulfonate, 8 parts by weight of fatty alcohol polyglycol ether, 20 parts by weight of the sodium salt of a phenolsulfonic acid/urea/formaldehyde condensate and 88 parts by weight of a paraffinic mineral oil.

The compositions according to the invention are distinguished by an outstanding activity against a broad spectrum of phytopathogenic fungi, in particular against botrytis. Some of them act systemically (i.e. they can be taken up from the treated plant without any loss in activity and may be translocated within the plant) and can be employed as foliar- and soil-acting fungicides.

They are especially important for controlling a large number of fungi which infect a plurality of crop plants, such as wheat, rye, barley, oats, rice, maize, grass, cotton, soya, coffee, sugar cane, grapevine, fruit species, ornamentals and vegetable species such as cucumbers, beans and cucurbits, and the seeds of these plants.

The compositions are applied by treating the fungi, or the seeds, plants, materials or the soil to be protected against fungal infection, with a fungicidally active amount of the active ingredients.

Application is effected before or after infection of the materials, plants or seeds by the fungi.

Specifically, the compositions are suitable for controlling the following plant diseases:

21

- Erysiphe graminis (powdery mildew) in cereals,
 Erysiphe cichoracearum and Sphaerotheca fuliginea in cucurbits,
 Podosphaera leucotricha in apples,
 Uncinula necator in grapevines,
 5 Venturia inaequalis (scab) in apples,
 Helminthosporium species in cereals,
 Septoria nodorum in wheat,
 Botrytis cinerea (gray mold) in strawberries, grapevines,
 Cercospora arachidicola in ground nuts,
 10 Pseudocercospora herpotrichoides in wheat, barley,
 Pyricularia oryzae in rice,
 Fusarium and Verticillium species in a variety of plants, Alternaria species in vegetables and fruit,
 Monilinia species in fruit, and
 15 Sclerotinia species in oilseed rape and vegetables.

The use against botrytis is preferred.

The compositions can also be employed in the protection of materials (protection of wood), e.g. against Paecilomyces variotii.

In general, the fungicidal compositions comprise from 0.1 to 95, preferably from 0.5 to 90% by weight of active ingredient.

25 Depending on the nature of the desired effect, the rates of application are from 0.02 to 3 kg of active ingredient per ha.

In the treatment of seed, amounts of from 0.001 to 50 g, preferably 0.01 to 10 g, of active ingredient are generally required per 30 kilogram of seed.

In the use form as fungicides, the compositions according to the invention can also comprise other active ingredients, e.g. herbicides, insecticides, growth regulators, fungicides, or else 35 fertilizers.

A mixture with fungicides frequently results in a widening of the fungicidal spectrum of action.

40 The following list of fungicides together with which the compounds according to the invention can be used is intended to illustrate the possible combinations, but not to impose any limitations:

45 sulfur,
 dithiocarbonate and their derivatives, such as
 iron(III) dimethyldithiocarbamate,

- zinc dimethyldithiocarbamate,
zinc ethylenebisdithiocarbamate,
manganese ethylenebisdithiocarbamate,
manganese zinc ethylenediaminebisdithiocarbamate,
5 tetramethylthiuram disulfide,
ammonia complex of zinc (N,N-ethylenebisdithiocarbamate),
ammonia complex of zinc (N,N'-propylenebisdithiocarbamate),
zinc (N,N'-propylenebisdithiocarbamate),
N,N'-polypropylenebis(thiocarbamoyl)disulfide,
10 nitro derivatives, such as
dinitro(1-methylheptyl)phenylcrotonate,
2-sec-butyl-4,6-dinitrophenyl-3,3-dimethylacrylate,
2-sec-butyl-4,6-dinitrophenylisopropyl carbonate,
di-isopropyl 5-nitroisophthalate;
15 heterocyclic substances, such as
2-heptadecyl-2-imidazoline acetate,
2,4-dichloro-6-(o-chloroanilino)-s-triazine,
O,O-diethyl phthalimidophosphonothioate,
5-amino-1-[bis(dimethylamino)phosphinyl]-3-phenyl-
20 1,2,4-triazole,
2,3-dicyano-1,4-dithioanthraquinone,
2-thio-1,3-dithiolo[4,5-b]quinoxaline,
methyl 1-(butylcarbamoyl)2-benzimidazolecarbamate,
2-methoxycarbonylaminobenzimidazole,
25 2-(furyl-(2))benzimidazole,
2-(thiazolyl-(4))benzimidazole,
N-(1,1,2,2-tetrachloroethylthio)tetrahydrophthalimide,
N-trichloromethylthiotetrahydrophthalimide,
N-trichloromethylthiophthalimide,
30 N-dichlorofluoromethylthio-N',N'-dimethyl-N-phenylsulfuric dia-
mide,
5-ethoxy-3-trichloromethyl-1,2,3-thiadiazole,
2-thiocyanatomethylthiobenzothiazole,
1,4-dichloro-2,5-dimethoxybenzene,
35 4-(2-chlorophenylhydrazono)-3-methyl-5-isoxazolone,
pyridine-2-thiol 1-oxide,
8-hydroxylquinoline or its copper salt,
2,3-dihydro-5-carboxanilido-6-methyl-1,4-oxathiine,
2,3-dihydro-5-carboxanilido-6-methyl-1,4-oxathiine 4,4-dioxide,
40 2-methyl-5,6-dihydro-4H-pyran-3-carboxanilide,
2-methylfuran-3-carboxanilide,
2,5-dimethylfuran-3-carboxanilide,
2,4,5-trimethylfuran-3-carboxanilide,
N-cyclohexyl-2,5-dimethylfuran-3-carboxamide,
45 N-cyclohexyl-N-methoxy-2,5-dimethylfuran-3-carboxamide,
2-methylbenzanilide,
2-iodobenzanilide,

23

- N-formyl-N-morpholine-2,2,2-trichloroethyl acetate,
 piperazine-1,4-diylbis(1-(2,2,2-trichloroethyl)formamide,
 1-(3,4-dichloroanilino)-1-formylamino-2,2,2-trichloroethane,
 2,6-dimethyl-N-tridecyl-morpholine or its salts,
 5 2,6-dimethyl-N-cyclododecyl-morpholine or its salts,
 N-[3-(p-tert-butylphenyl)-2-methylpropyl]-cis-2,6-dimethylmorpho-
 line,
 N-[3-(p-tert-butylphenyl)-2-methylpropyl]piperidine,
 1-[2-(2,4-dichlorophenyl)-4-ethyl-1,3-dioxolan-2-yl-
 10 ethyl]-1H-1,2,4-triazole,
 1-[2-(2,4-dichlorophenyl)-4-n-propyl]-1,3-dioxolan-2-yl-
 ethyl]-1H-1,2,4-triazole,
 N-(n-propyl)-N-(2,4,6-trichlorophenoxyethyl)-N'-imidazolyl-urea,
 1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-
 15 2-butanone,
 1-(4-chlorophenyl)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-buta-
 nol,
 α-(2-chlorophenyl)-α-(4-chlorophenyl)-5-pyrimidinemethanol,
 5-butyl-2-dimethylamino-4-hydroxyl-6-methylpyrimidine,
 20 bis-(p-chlorophenyl)-3-pyridinemethanol,
 1,2-bis-(3-ethoxycarbonyl-2-thioureido)benzene,
 1,2-bis-(3-methoxycarbonyl-2-thioureido)benzene,
 and a variety of fungicides, such as
 dodecylguanidine acetate,
 25 3-[3-(3,5-dimethyl)-2-oxycyclohexyl)-2-hydroxyethyl]]glutari-
 mide, hexachlorobenzene,
 methyl N-(2,6-dimethylphenyl)-N-(2-furoyl)-DL-alaninate,
 DL-N-(2,6-dimethylphenyl)-N-(2'-methoxyacetyl)alanine methyl
 ester,
 30 N-(2,6-dimethylphenyl)-N-chloroacetyl-D,L-2-aminobutyrolactone,
 DL-N-(2,6-dimethylphenyl)-N-(phenylacetyl)alanine methyl ester,
 5-methyl-5-vinyl-3-(3,5-dichlorophenyl)-2,4-dioxo-1,3-oxazoli-
 dine,
 3-[3,5-dichlorophenyl-(5-methyl-5-methoxymethyl)-1,3-oxazoli-
 35 dine-2,4-dione,
 3-(3,5-dichlorophenyl)-1-isopropylcarbamoylethylhydantoin,
 N-(3,5-dichlorophenyl)-1,2-dimethylcyclopropane-1,2-di-
 carboximide,
 2-cyano-[N-(ethylaminocarbonyl)-2-methoximino]acetamide,
 40 1-[2-(2,4-dichlorophenyl)pentyl]-1H-1,2,4-triazole,
 2,4-difluoro-α-(1H-1,2,4-triazolyl-1-methyl)benzhydryl alcohol,
 N-(3-chloro-2,6-dinitro-4-trifluoromethylphenyl)-5-trifluorome-
 thyl-3-chloro-2-aminopyridine,
 1-((bis(4-fluorophenyl)methylsilyl)methyl-1H-1,2,4-triazole.
 45

24



The synergistic action of the compositions according to the invention is illustrated with the aid of the use examples which follow, where the compounds of the formula I.1 to I.7 were used as active ingredients I:

5

I.1

I.2

10





15

I.3

I.4

20




25

I.5


I.6

E,E isomer (comprises 5% of Z,E isomer)

30




35



I.7

E,E side chain

40



Use Example 1

Activity against Botrytis cinerea in bell peppers

Green bell pepper disks were sprayed to drip point with aqueous preparation of active ingredient, comprising 80% of active ingredient and 20% emulsifier in the dry matter. 2 hours after the spray coating had dried on, the fruit disks were inoculated with a spore suspension of Botrytis cinerea containing 1.7×10^6 spores per ml of a 2% strength Biomalz solution. The inoculated fruit disks were subsequently incubated for 4 days in humid chambers at 18°C. The Botrytis development on the diseased fruit disks was then evaluated visually (disease level 99%).

The visually determined data for the percentage of diseased fruit area were converted into efficacies as a percentage of the untreated control. An efficacy of 0 is the same disease level as the untreated control, an efficacy of 100 is a disease level of 0 percent. The efficacies to be expected for combinations of active ingredients were calculated using Colby's formula (Colby, S. R. (Calculating synergistic and antagonistic responses of herbicide Combinations", Weeds, 15, pp. 20 - 22, 1967) and compared with the observed efficacies. The results are shown in Table 1 below.

Table 1

Active Compound	Concentration of active compound in ppm		Degree of activity in % of control	
	Fenazaquin	Active compound I.1	observed	calculated*)
Control (untreated)	---	---	0	
Fenazaquin	250	---	19	
	125	---	0	
I.1	---	63	79	
	---	31	74	
	---	16	19	
Fenazaquin + I.1	250	63	99	83
	250	31	85	79

*) calculated according to the Colby formula

The test results show that the observed efficacy in all mixing ratios exceeds the additive efficacy calculated earlier using Colby's formula, i.e. a synergistic effect is present.

5

Use Example 2

Activity against *Botrytis cinerea*

10 Pepper seedlings cv. "Neusiedler Ideal Elite" were grown until 4
- 5 leaves had developed properly and then sprayed to drip point
with aqueous suspensions comprising 80% of active ingredient and
20% of emulsifier in the dry matter. After the spray coating had
dried on, the plants were sprayed with a conidia suspension of
15 the fungus *Botrytis cinerea* and placed into a chamber with high
atmospheric humidity at 22 - 24°C. After 5 days, the disease on
the untreated control plants had developed to such an extent that
the leaf necroses formed covered a major portion of the leaves
(disease level 83%).

20

The visually determined data for the percentage of diseased fruit
area were converted into efficacies as a percentage of the un-
treated control. An efficacy of 0 is the same disease level as
the untreated control, an efficacy of 100 is a disease level of 0
25 percent. The efficacies to be expected for combinations of active
ingredients were calculated using Colby's formula (Colby, S. R.
(Calculating synergistic and antagonistic responses of herbicide
Combinations", Weeds, 15, pp. 20 - 22, 1967) and compared with
the observed efficacies. The results are shown in Table 2 below.

30

35

40

45

Table 2

Active Compound	Concentration of active compound in ppm		Degree of activity in % of control	
	Fenazaquin	Active compound I.1	observed	calculated*)
Control (untreated)	---	---	0	
Fenazaquin	500	---	4	
	250	---	4	
I.1	---	63	28	
	---	31	16	
Fenazaquin + I.1	500	63	88	33
	250	31	70	19

*) calculated according to the Colby formula

29

The test results show that the observed efficacy in all mixing ratios exceeds the additive efficacy calculated earlier using Colby's formula, i.e. a synergistic effect is present.

5

Use Example 3

Activity against Botrytis cinerea on bell peppers

10 Following the method described in Use Example 1, the following results were obtained using the compounds listed in Table 3 below:

15

20

25

30

35

40

45

Table 3

5	Active Compound	Concentration of active compound in ppm		Degree of activity in % of control	
		I.1-I.7	Fenazaquin II	observed	calculated*)
10	control (untreated)	--	--	0	
15	Fenazaquin II	--	500 250 50 25	0 0 0 0	--
	I.2	50 25	--	95 89	--
20	I.3	50 25	--	79 28	--
	I.4	50 25	--	15 34	--
25	I.5	50	--	60	--
	I.6	50 25	--	47 47	--
	I.7	50 25	--	34 42	--
30	I.2 + II	25 25	250 25	100 95	89 89
	I.3 + II	25 25	250 25	100 79	28 28
35	I.4 + II	25 25	250 25	79 66	34 34
	I.5 + II	25 50	250 50	97 79	73 60
40	I.6 + II	25 25	250 25	97 76	47 47
	I.7 + II	25 25	250 25	97 84	42 42

45 * calculated according to the Colby formula

31

The test results show that the observed efficacy in all mixing ratios exceeds the additive efficacy calculated earlier using Colby's formula, i.e. a synergistic effect is present.

5

Use Example 4

Activity against *Botrytis cinerea*

10 Following the method described in Use Example 1, the following results were obtained using the compounds listed in Table 4 below:

15

20

25

30

35

40

45

Table 4

Active compound	Concentration of active compound in ppm		Degree of activity in % of control	
	I.1-I.7	Fenazaquin II	observed	calculated*)
Control (untreated)			0	
Fenazaquin II		500 250 50 25	48 27 34 17	--
I.2	25	--	0	--
I.3	25	--	34	--
I.4	50 25	--	0 0	--
I.5	50 25	--	0 0	--
I.6	50 25	--	52 72	--
I.7	25	--	62	
I.2 + II	25	25	59	17
I.3 + II	25	25	65	46
I.4 + II	50 25 50 25	500 250 50 25	90 79 52 55	48 27 34 17
I.5 + II	50 25	50 25	52 52	34 17
I.6 + II	50 25 50 25	500 250 50 25	97 90 86 86	75 80 68 77
I.7 + II	25	25	90	68

*) calculated according to the Colby formula

The test results show that the observed efficacy in all mixing ratios exceeds the additive efficacy calculated earlier using Colby's formula, i.e. a synergistic effect is present.

5

Similar results are obtained when another individual compound from amongst those mentioned in Table I.1 to I.8 above are used in the above-described experiments.

10

15

20

25

30


35

40


45

We claim:


1. A composition for controlling harmful fungi which comprises, in a solid or liquid carrier, at least one active ingredient I, which inhibits respiration on the cytochrome complex III, and fenazaquin, of the formula.



2. A composition as claimed in claim 1 which comprises, as active ingredient I, a compound of the formula IA or IB



IA



IB

where is a double or single bond;

R' is $-\text{C}[\text{CO}_2\text{CH}_3]=\text{CHOCH}_3$, $-\text{C}[\text{CO}_2\text{CH}_3]=\text{NOCH}_3$,
 $-\text{C}[\text{CONHCH}_3]=\text{NOCH}_3$, $-\text{C}[\text{CO}_2\text{CH}_3]=\text{CHCH}_3$, $-\text{C}[\text{CO}_2\text{CH}_3]=\text{CHCH}_2\text{CH}_3$,
 $-\text{C}[\text{COCH}_3]=\text{NOCH}_3$, $-\text{C}[\text{COCH}_2\text{CH}_3]=\text{NOCH}_3$, $-\text{N}(\text{OCH}_3)-\text{CO}_2\text{CH}_3$,
 $-\text{N}(\text{CH}_3)-\text{CO}_2\text{CH}_3$ or $-\text{N}(\text{CH}_2\text{CH}_3)-\text{CO}_2\text{CH}_3$;

R'' is an organic radical which is bonded directly or via an oxy, mercapto, amino, or alkylamino group,

together with a group X and the ring Q or T to which they are bonded form an unsubstituted or substituted bicyclic, partially or fully unsaturated system which, besides carbon ring members, can have 1, 2 or 3 hetero atoms independently selected from amongst oxygen, sulfur and nitrogen;

4

R^x is -OC[CO₂CH₃]=CHOCH₃, -OC[CO₂CH₃]=CHCH₃,
-OC[CO₂CH₃]=CHCH₂CH₃, -SC[CO₂CH₃]=CHOCH₃,
-SC[CO₂CH₃]=CHCH₃, -SC[CO₂CH₃]=CHCH₂CH₃,
-N(CH₃)C[CO₂CH₃]=CHOCH₃, -N(CH₃)C[CO₂CH₃]=NOCH₃,
-CH₂C[CO₂CH₃]=CHOCH₃, -CH₂C[CO₂CH₃]=NOCH₃ or
-CH₂C[CONHCH₃]=NOCH₃;

R^y is oxygen, sulfur, =CH- or =N-;

n is 0, 1, 2 or 3, it being possible for the radicals X to be identical or different if n > 1;

X is cyano, nitro, halogen, alkyl, haloalkyl, alkoxy, haloalkoxy, alkylthio, or,

if n > 1, a C₃-C₅-alkylene, C₃-C₅-alkenylene, oxy-C₂-C₄-alkylene, oxy-C₁-C₃-alkyleneoxy, oxy-C₂-C₄-alkenylene, oxy-C₂-C₄-alkenyleneoxy or butadienediyl group which is bonded to two adjacent C atoms of the phenyl ring, it being possible for these chains, in turn, to have attached to them one to three radicals which, independently of one another, are selected from amongst halogen, alkyl, haloalkyl, alkoxy, haloalkoxy and alkylthio;

Y is =C- or -N-;


Q is phenyl, pyrrolyl, thienyl, furyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, triazolyl, pyridinyl, 2-pyridinyl, pyrimidinyl or triazinyl; and

T is phenyl, oxazolyl, thiazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl or triazinyl.

X

3. A composition as claimed in claim 2 which comprises, as active ingredient I, a compound of the formula IA or IB where R" is aryloxy, hetaryloxy, aryloxymethylene, hetaryloxymethylene, aryloethenylene or hetaryloethenylene, it being possible for these radicals to have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, halogen, CF₃, CHF₂, CN, alkoxy and phenyl which, in turn, may have 1, 2 or 3 substituents which, independently of one another, are selected from amongst halogen, CF₃, CHF₂, phenyl, CN, phenoxy, alkyl, alkoxy and haloalkoxy;

or R" is



where R^α is alkyl;

R^β is phenyl, pyridyl or pyrimidyl which may have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, alkoxy, halogen, haloalkoxy, CF₃ and CHF₂;

R^γ is alkyl, alkoxy, halogen, haloalkyl or hydrogen;


A

- R^d is hydrogen, cyano, halogen, alkyl, alkoxy, alkylthio, alkylamino, di-alkylamino, alkenyl, alkenyloxy, alkenylthio, alkenylamino, N-alkenyl-N-alkylamino, alkynyl, alkynyloxy, alkynylthio, alkynylamino, N-alkynyl-N-alkylamino, it being possible for the hydrocarbon radicals of these groups to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 radicals which, independently of one another, are selected from amongst cyano, nitro, hydroxyl, alkoxy, haloalkoxy, alkoxycarbonyl, alkylthio, alkylamino, di-alkylamino, alkenyloxy, cycloalkyl, cycloalkyloxy, heterocyclyl, heterocyclyloxy, aryl, aryloxy, arylalkoxy, hetaryl, hetaryloxy and hetarylalkoxy, it being possible for the cyclic radicals, in turn, to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 groups which, independently of one another, are selected from amongst cyano, nitro, hydroxyl, alkyl, haloalkyl, cycloalkyl, alkoxy, haloalkoxy, alkoxycarbonyl, alkylthio, alkylamino, di-alkylamino, alkenyl and alkenyloxy;
- or
- is cycloalkyl, cycloalkyloxy, cycloalkylthio, cycloalkylamino, N-cycloalkyl-N-alkylamino, heterocyclyl, heterocyclyloxy, heterocyclylthio, heterocyclylamino, N-heterocyclyl-N-alkylamino, aryl, aryloxy, arylthio, arylamino, N-aryl-N-alkylamino, hetaryl, hetaryloxy, hetarylthio, hetarylamino or N-hetaryl-N-alkylamino it being possible for the cyclic radicals to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 groups which, independently of one another, are selected from amongst cyano, nitro, hydroxyl, alkyl, haloalkyl, cycloalkyl, alkoxy, haloalkoxy, alkoxycarbonyl, alkylthio, alkylamino, di-alkylamino, alkenyl, alkenyloxy, benzyl, benzyloxy, aryl, aryloxy, hetaryl and hetaryloxy, it being possible for the aromatic radicals, in turn, to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 of the following groups: cyano, alkyl, haloalkyl, alkoxy, nitro;
- R^e is alkyl, alkenyl or alkynyl, it being possible for these groups to be partially or fully halogenated and/or to have attached to them 1, 2 or 3 of the following radicals: cyano, alkoxy, cycloalkyl.
4. A composition as claimed in claim 2 which comprises, as active ingredient I, a compound of the formula IA or IB where R'' is aryloxy, hetaryloxy, aryloxymethylene, hetaryloxymethylene, arylenylene or hetarylenylene, it being possible

x

for these radicals to have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, halogen, CF₃, CHF₂, alkoxy and phenyl which, in turn, may have 1, 2 or 3 substituents which, independently of one another are selected from amongst halogen, CF₃, CHF₂, phenyl, CN and phenoxy;

or R" is



where R^α is alkyl;

15 R^β is phenyl or pyrimidinyl which may have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, alkoxy, halogen, CF₃ and CHF₂;

20 R^γ is alkyl;

R^δ is alkyl or phenyl which, if appropriate, has 1, 2 or 3 substituents which, independently of one another are selected from amongst halogen, CF₃ and CHF₂; and

25 R^ε is alkyl.

5. A composition as claimed in claim 3 which comprises, as active ingredient I, a compound of the formula IA where R" has one of the following meanings:

30 a) phenyloxymethylene, pyridinyloxymethylene, pyrimidinyl-oxymethylene or pyrazolyloxymethylene, it being possible for the aromatic radical to have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, halogen, CF₃, CHF₂, -C(CH₃)=NOCH₃, and phenyl which is unsubstituted or substituted by 1, 2 or 3 halogen atoms and/or alkyl groups;

40 b) phenoxy or pyrimidinyl-oxo, unsubstituted or substituted by 1, 2 or 3 halogen atoms or one phenoxy radical which may have a halogen or cyano substituent;

45

-39-

X

- c) phenylethylenylene or pyrazolylenethylene, it being possible for the phenyl or pyrazolyl radical to have 1, 2 or 3 substituents which, independently of one another, are selected from amongst halogen, CF_3 , CHF_2 and phenyl.

5

- d) $\text{CH}_2\text{ON}=\text{CR}^{\alpha}\text{R}^{\beta}$,
where R^{α} is alkyl; and

10

R^{β} is phenyl which may have 1, 2 or 3 substituents which, independently of one another, are selected from amongst alkyl, halogen, CF_3 and CHF_2 , or is pyrimidynyl which is unsubstituted or substituted by 1 or 2 alkoxy radicals;

15

- e) $\text{CH}_2\text{ON}=\text{CR}^{\gamma}\text{CR}^{\delta}=\text{NOR}^{\epsilon}$,
where R^{γ} is alkyl, alkoxy or halogen;

20

R^{δ} is alkyl, cyano, halogen, alkoxy, alkenyl or phenyl which is unsubstituted or substituted by 1, 2 or 3 halogen atoms; and

R^{ϵ} is alkyl.


6. A composition as claimed in any of claims 2 to 5, where R' is
25 $-\text{C}[\text{CO}_2\text{CH}_3]=\text{NOCH}_3$, $-\text{C}[\text{CONHCN}_3]=\text{NOCH}_3$ or $-\text{C}[\text{CO}_2\text{CH}_3]=\text{CHOCH}_3$.

7. A composition as claimed in any of claims 2 to 6 which comprises, as active ingredient I, a compound of the formula IA where Q is phenyl and n is 0.

30


8. A composition as claimed in claim 1 which comprises, as active ingredient I, a compound of the following formulae:

35



40

I.1




I.2


45

-H₂O-


X




I.3



I.4




I.5



I.6

E, E-Isomer



I.7

E, E- side chain

9. A composition as claimed in any of the preceding claims which is conditioned in two parts, one part comprising the active ingredient I in a solid or liquid carrier, while the other part comprises fenazaquin in a solid or liquid carrier.
10. A method of controlling harmful fungi, which comprises treating the fungi, their environment, or the materials, plants, seeds, soils, areas or spaces to be protected against fungal infection, with a composition as claimed in any of claims 1 to 9, it being possible for the application of the active ingredients to be effected simultaneously, i.e. jointly or separately, or in succession.