



(11) **EP 3 036 993 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**29.06.2016 Bulletin 2016/26**

(51) Int Cl.:  
**A01N 43/40** (2006.01) **A01N 43/50** (2006.01)  
**A01N 43/90** (2006.01) **A01N 47/36** (2006.01)  
**A01P 13/00** (2006.01)

(21) Application number: **16152976.3**

(22) Date of filing: **27.10.2010**

(84) Designated Contracting States:  
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**  
Designated Extension States:  
**BA ME**

- **WEIMER, Monte**  
**Pittsboro, IN 46167 (US)**
- **MCVEIGH-NELSON, Andrea**  
**Indianapolis, IN 46260 (US)**
- **ELLIS, Andrew**  
**Greenville, MS 38701 (US)**

(30) Priority: **28.10.2009 US 255689 P**

(74) Representative: **f & e patent**  
**Fleischer, Engels & Partner mbB, Patentanwälte**  
**Braunsberger Feld 29**  
**51429 Bergisch Gladbach (DE)**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:  
**10773505.2 / 2 493 311**

(71) Applicant: **Dow AgroSciences LLC**  
**Indianapolis, IN 46268 (US)**

Remarks:

This application was filed on 27-01-2016 as a divisional application to the application mentioned under INID code 62.

(72) Inventors:

- **MANN, Richard**  
**Franklin, IN 46131 (US)**

(54) **SYNERGISTIC HERBICIDAL COMPOSITION CONTAINING FLUROXYPYR AND HALOSULFURON-METHYL, IMAZAMOX OR IMAZETHAPYR**

(57) An herbicidal synergistic composition containing (a) fluroxypyr and (b) an ALS inhibitor herbicide, in which the ALS inhibitor herbicide is halosulfuron-methyl, imazamox or imazethapyr, provides improved post-emergence weed control in rice, cereal and grain crops, pastures, rangelands, IVM and turf.

**EP 3 036 993 A2**

## Description

**[0001]** This invention concerns a synergistic herbicidal composition containing (a) fluroxypyr and (b) at least one herbicide selected from the group consisting of penoxsulam, halosulfuron-methyl, imazamox and imazethapyr for controlling weeds in crops, especially rice, cereal and grain crops, pastures, rangelands, industrial vegetation management (IVM) and turf. These compositions provide improved post-emergence herbicidal weed control.

**[0002]** The protection of crops from weeds and other vegetation which inhibit crop growth is a constantly recurring problem in agriculture. To help combat this problem, researchers in the field of synthetic chemistry have produced an extensive variety of chemicals and chemical formulations effective in the control of such unwanted growth. Chemical herbicides of many types have been disclosed in the literature and a large number are in commercial use.

**[0003]** In some cases, herbicidal active ingredients have been shown to be more effective in combination than when applied individually and this is referred to as "synergism." As described in the Herbicide Handbook of the Weed Science Society of America, Eighth Edition, 2002, p. 462 "'synergism' [is] an interaction of two or more factors such that the effect when combined is greater than the predicted effect based on the response to each factor applied separately." The present invention is based on the discovery that fluroxypyr, penoxsulam, halosulfuron-methyl and imazamox, already known individually for their herbicidal efficacy, display a synergistic effect when applied in combination.

**[0004]** The herbicidal compounds forming the synergistic composition of this invention are independently known in the art for their effects on plant growth.

**[0005]** The present invention concerns a synergistic herbicidal mixture comprising an herbicidally effective amount of (a) fluroxypyr and (b) an acetolactate synthase (ALS) inhibitor herbicide. ALS inhibitor herbicides include compounds from the classes of sulfonamides, sulfonylureas and imidazolinones. Particularly useful ALS inhibitor herbicides include, but are not limited to, penoxsulam, halosulfuron-methyl, imazamox and imazethapyr. The compositions may also contain an agriculturally acceptable adjuvant and/or carrier.

**[0006]** The present invention also concerns herbicidal compositions for and methods of controlling the growth of undesirable vegetation, particularly in monocot crops including rice, wheat, barley, oats, rye, sorghum, corn, maize, pastures, grasslands, rangelands, fallowland, turf, IVM and aquatics and the use of these synergistic compositions.

**[0007]** The species spectra of ALS inhibitors like penoxsulam, halosulfuron-methyl, imazamox and imazethapyr, i.e., the weed species which the respective compounds control, are broad and highly complementary with that of fluroxypyr. For example, it has been surprisingly found that a combination of penoxsulam and fluroxypyr exhibits a synergistic action in the control of Hemp sesbania (*Sesbania exaltata*; SEBEX), Texasweed (*Caperonia palustris*; CNPPA), Japanese bulrush (*Scirpus juncooides*; SCPJU), narrow-leaved plantain (*Plantago lanceolata* L.; PLALA), and barnyardgrass (*Echinochloa crus-galli*; ECHCG) at application rates equal to or lower than the rates of the individual compounds. Similarly, it has been surprisingly found that a combination of halosulfuron-methyl and fluroxypyr exhibits a synergistic action in the control of barnyardgrass (*Echinochloa crus-galli*; ECHCG), Chinese sprangletop (*Leptochloa chinensis*; LEFCH), and yellow nutsedge (*Cyperus esculentus*; CYPES) at application rates equal to or lower than the rates of the individual compounds. In addition, it has been surprisingly found that a combination of imazamox and fluroxypyr exhibits a synergistic action in the control of barnyardgrass (*Echinochloa crus-galli*; ECHCG), broadleaf signalgrass (*Brachiaria platyphylla*; BRAPP), and yellow nutsedge (*Cyperus esculentus*; CYPES) at application rates equal to or lower than the rates of the individual compounds. In addition, it has been surprisingly found that a combination of imazethapyr and fluroxypyr exhibits a synergistic action in the control of sprangletop grass (*Leptochloa* spp, LEFSS) and fall panicum (*Panicum dichotomiflorum*, PANDI) at application rates equal to or lower than the rates of the individual compounds.

**[0008]** Fluroxypyr is the common name for [(4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic acid. Its herbicidal activity is described in The Pesticide Manual, Fourteenth Edition, 2006. Fluroxypyr controls a wide range of economically important broadleaf weeds. It can be used as the acid itself or as an agriculturally acceptable salt or ester. Use as an ester is preferred, with the meptyl ester being most preferred.

**[0009]** Penoxsulam is the common name for 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy-[1,2,4]triazolo[1,5-c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide. Its herbicidal activity is described in The Pesticide Manual, Fourteenth Edition, 2006. Penoxsulam controls *Echinochloa* spp., as well as many broadleaf, sedge and aquatic weeds in rice, and *Apera* spp. in grass, as well as many broadleaf weeds in cereals.

**[0010]** Halosulfuron-methyl is the common name for methyl 3-chloro-5-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-1-methyl-1H-pyrazole-4-carboxylate. Its herbicidal activity is described in The Pesticide Manual, Fourteenth Edition, 2006. Halosulfuron-methyl controls many broadleaf and nutsedge weeds in rice, corn, sorghum, sugar cane, nuts and turf.

**[0011]** Imazamox is the common name for 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid. Its herbicidal activity is described in The Pesticide Manual, Fourteenth Edition, 2006. Imazamox controls many broadleaf weeds in corn, rape, alfalfa, peas and beans.

**[0012]** Imazethapyr is the common name for 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid. Its herbicidal activity is described in The Pesticide Manual, Fourteenth Edition, 2006.

Imazethapyr controls many grass and broadleaf weeds in alfalfa, peas, beans, soybeans and imidazolinone tolerant rice and corn.

**[0013]** The term herbicide is used herein to mean an active ingredient that kills, controls or otherwise adversely modifies the growth of plants. An herbicidally effective or vegetation controlling amount is an amount of active ingredient which causes an adversely modifying effect and includes deviations from natural development, killing, regulation, desiccation, retardation, and the like. The terms plants and vegetation include germinant seeds, emerging seedlings, plants emerging from vegetative propagules, and established vegetation.

**[0014]** Herbicidal activity is exhibited by the compounds of the synergistic mixture when they are applied directly to the plant or to the locus of the plant at any stage of growth or before planting or emergence. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of chemical applied. These and other factors can be adjusted as is known in the art to promote non-selective or selective herbicidal action. Generally, it is preferred to apply the composition of the present invention postemergence to relatively immature undesirable vegetation to achieve the maximum control of weeds.

**[0015]** In the composition of this invention, the weight ratio of fluroxypyr (acid equivalent) to ALS inhibitor herbicide (active ingredient) at which the herbicidal effect is synergistic lies within the range of between about 1:2 and about 140:1.

**[0016]** The rate at which the synergistic composition is applied will depend upon the particular type of weed to be controlled, the degree of control required, and the timing and method of application. The ALS inhibitor herbicide is applied at a rate between about 4 g ai/ha and about 100 g ai/ha and fluroxypyr is applied at a rate between about 50 g ae/ha and about 560 g ae/ha.

**[0017]** The components of the synergistic mixture of the present invention can be applied either separately or as part of a multipart herbicidal system, which can be provided as a premix or a tank mix.

**[0018]** The synergistic mixture of the present invention can be applied in conjunction with one or more other herbicides to control a wider variety of undesirable vegetation. When used in conjunction with other herbicides, the composition can be formulated with the other herbicide or herbicides, tank mixed with the other herbicide or herbicides or applied sequentially with the other herbicide or herbicides. Some of the herbicides that can be employed in conjunction with the synergistic composition of the present invention include: 2,4-D, acetochlor, acifluorfen, aclonifen, AE0172747, alachlor, amidosulfuron, aminotriazole, ammonium thiocyanate, anilifos, atrazine, AVH 301, azimsulfuron, benfuresate, bensulfuron-methyl, bentazone, benthocarb, benzobicyclon, bifenox, bispyribac-sodium, bromacil, bromoxynil, butachlor, butafenacil, butralin, cafenstrole, carbetamide, carfentrazone-ethyl, chlorflurenol, chlorimuron, chlorpropham, cinosulfuron, clethodim, clomazone, clopyralid, cloransulam-methyl, cyclosulfamuron, cycloxydim, cyhalofop-butyl, dicamba, dichlobenil, dichlorprop-P, diclosulam, diflufenican, diflufenzopyr, dimethenamid, dimethenamid-p, diquat, dithiopyr, diuron, EK2612, EPTC, esprocarb, ET-751, ethoxysulfuron, ethbenzanid, F7967, fenoxaprop, fenoxaprop-ethyl, fenoxaprop-ethyl + isoxadifen-ethyl, fentrazamide, flazasulfuron, florasulam, fluazifop, fluazifop-P-butyl, flucetosulfuron (LGC-42153), flufenacet, flufenpyr-ethyl, flumetsulam, flumiclorac-pentyl, flumioxazin, fluometuron, flupyrsulfuron, fomesafen, foramsulfuron, fumiclorac, glufosinate, glufosinate-ammonium, glyphosate, haloxyfop-methyl, haloxyfop-R, imazame-thabenz, imazapic, imazapyr, imazaquin, imazosulfuron, indanofan, indaziflam, iodiosulfuron, ioxynil, ipfencarbazone (HOK-201), IR 5790, isoproturon, isoxaben, isoxaflutole, KUH-071, lactofen, linuron, MCPA, MCPA ester & amine, mecoprop-P, mefenacet, mesosulfuron, mesotrione, metamifop, metazosulfuron (NC-620), metolachlor, metosulam, metribuzin, metsulfuron, molinate, MSMA, napropamide, nicosulfuron, norflurazon, OK-9701, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxazichlomefone, oxyfluorfen, paraquat, pendimethalin, pentoxazone, pethoxamid, picloram, picolinafen, piperophos, pretilachlor, primisulfuron, profoxydim, propachlor, propanil, propyrisulfuron (TH-547), propyza-mide, prosulfocarb, prosulfuron, pyraclonil, pyrazogyl, pyrazosulfuron, pyribenzoxim (LGC-40863), pyriftalid, pyrimino-bac-methyl, pyrimisulfan (KUH-021), pyroxsulam, pyroxasulfone (KIH-485), quinclorac, quizalofop-ethyl-D, S-3252, sethoxydim, simazine, SL-0401, SL-0402, S-metolachlor, sulcotrione, sulfentrazone, sulfosate, tebuthiuron, tefuryltrione (AVH-301), terbacil, thiazopyr, thiobencarb, triclopyr, trifluralin and tritosulfuron.

**[0019]** The synergistic composition of the present invention can, further, be used in conjunction with glyphosate, glufosinate, dicamba, imidazolinones, sulfonylureas, or 2,4-D on glyphosate-tolerant, glufosinate-tolerant, dicamba-tolerant, imidazolinone-tolerant, sulfonylurea-tolerant and 2,4-D-tolerant crops. It is generally preferred to use the synergistic composition of the present invention in combination with herbicides that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate employed. It is further generally preferred to apply the synergistic composition of the present invention and other complementary herbicides at the same time, either as a combination formulation or as a tank mix.

**[0020]** The synergistic composition of the present invention can generally be employed in combination with known herbicide safeners, such as benoxacor, benthocarb, brassinolide, cloquintocet (mexyl), cyometrinil, daimuron, dichlor-mid, dicyclonon, dimepiperate, disulfoton, fenclorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, harpin proteins, isoxadifen-ethyl, mefenpyr-diethyl, MG 191, MON 4660, naphthalic anhydride (NA), oxabetrinil, R29148 and N-phenyl-

sulfonylbenzoic acid amides, to enhance their selectivity. Cloquintocet (mexyl) is a particularly preferred safener for the synergistic compositions of the present invention, specifically antagonizing any harmful effect of the synergistic compositions on rice and cereals.

**[0021]** The synergistic mixture of fluroxypyr and penoxsulam of the present invention also provides a safening effect when applied to sunflower (*Helianthus annuus*; HELAN) and centipedegrass (*Eremochloa ophiuroides*; ERLOP). Another aspect of the present invention is a method for protecting centipedegrass and sunflower from the individual harmful effects of penoxsulam and fluroxypyr which comprises contacting centipedegrass or sunflower, or applying to the area under cultivation, a safening amount of a synergistic mixture of penoxsulam and fluroxypyr.

**[0022]** In practice, it is preferable to use the synergistic composition of the present invention in mixtures containing an herbicidally effective amount of the herbicidal components along with at least one agriculturally acceptable adjuvant or carrier. Suitable adjuvants or carriers should not be phytotoxic to valuable crops, particularly at the concentrations employed in applying the compositions for selective weed control in the presence of crops, and should not react chemically with herbicidal components or other composition ingredients. Such mixtures can be designed for application directly to weeds or their locus or can be concentrates or formulations that are normally diluted with additional carriers and adjuvants before application. They can be solids, such as, for example, dusts, granules, water dispersible granules, or wettable powders, or liquids, such as, for example, emulsifiable concentrates, solutions, emulsions or suspensions.

**[0023]** Suitable agricultural adjuvants and carriers that are useful in preparing the herbicidal mixtures of the invention are well known to those skilled in the art. Some of these adjuvants include, but are not limited to, crop oil concentrate (mineral oil (85%) + emulsifiers (15%)); nonylphenol ethoxylate; benzylcocoalkyldimethyl quaternary ammonium salt; blend of petroleum hydrocarbon, alkyl esters, organic acid, and anionic surfactant; C<sub>9</sub>-C<sub>11</sub> alkylpolyglycoside; phosphated alcohol ethoxylate; natural primary alcohol (C<sub>12</sub>-C<sub>16</sub>) ethoxylate; di-sec-butylphenol EO-PO block copolymer; polysiloxane-methyl cap; nonylphenol ethoxylate + urea ammonium nitrate; emulsified methylated seed oil; tridecyl alcohol (synthetic) ethoxylate (8EO); tallow amine ethoxylate (15 EO); PEG(400) dioleate-99.

**[0024]** Liquid carriers that can be employed include water, toluene, xylene, petroleum naphtha, crop oil, acetone, methyl ethyl ketone, cyclohexanone, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol monomethyl ether and diethylene glycol monomethyl ether, methyl alcohol, ethyl alcohol, isopropyl alcohol, amyl alcohol, ethylene glycol, propylene glycol, glycerine, *N*-methyl-2-pyrrolidinone, *N,N*-dimethyl alkylamides, dimethyl sulfoxide, liquid fertilizers and the like. Water is generally the carrier of choice for the dilution of concentrates.

**[0025]** Suitable solid carriers include talc, pyrophyllite clay, silica, attapulgius clay, kaolin clay, kieselguhr, chalk, diatomaceous earth, lime, calcium carbonate, bentonite clay, Fuller's earth, cottonseed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell flour, lignin, and the like.

**[0026]** It is usually desirable to incorporate one or more surface-active agents into the compositions of the present invention. Such surface-active agents are advantageously employed in both solid and liquid compositions, especially those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic or nonionic in character and can be employed as emulsifying agents, wetting agents, suspending agents, or for other purposes. Surfactants conventionally used in the art of formulation and which may also be used in the present formulations are described, *inter alia*, in "McCutcheon's Detergents and Emulsifiers Annual," MC Publishing Corp., Ridgewood, New Jersey, 1998 and in "Encyclopedia of Surfactants," Vol. I-III, Chemical Publishing Co., New York, 1980-81. Typical surface-active agents include salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C<sub>18</sub> ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C<sub>16</sub> ethoxylate; soaps, such as sodium stearate; alkylnaphthalene-sulfonate salts, such as sodium dibutyl-naphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; salts of mono and dialkyl phosphate esters; vegetable oils such as soybean oil, rapeseed oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; and esters of the above vegetable oils.

**[0027]** Other additives commonly used in agricultural compositions include compatibilizing agents, antifoam agents, sequestering agents, neutralizing agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids, sticking agents, dispersing agents, thickening agents, freezing point depressants, antimicrobial agents, and the like. The compositions may also contain other compatible components, for example, other herbicides, plant growth regulants, fungicides, insecticides, and the like and can be formulated with liquid fertilizers or solid, particulate fertilizer carriers such as ammonium nitrate, urea and the like.

**[0028]** The concentration of the active ingredients in the synergistic composition of the present invention is generally from 0.001 to 98 percent by weight. Concentrations from 0.01 to 90 percent by weight are often employed. In compositions designed to be employed as concentrates, the active ingredients are generally present in a concentration from 1 to 98 weight percent, preferably 5 to 90 weight percent. Such compositions are typically diluted with an inert carrier, such as water, before application, or applied as a dry or liquid formulation directly into flooded rice fields. The diluted compositions

usually applied to weeds or the locus of weeds generally contain 0.0001 to 10 weight percent active ingredient and preferably contain 0.001 to 5.0 weight percent.

**[0029]** The present compositions can be applied to weeds or their locus by the use of conventional ground or aerial dusters, sprayers, and granule applicators, by addition to irrigation or paddy water, and by other conventional means known to those skilled in the art.

**[0030]** The following examples illustrate the present invention.

#### Evaluation of Postemergence Herbicidal Activity of Mixtures in the Greenhouse

**[0031]** Seeds of the desired test plant species were planted in 80% mineral soil/20% grit planting mixture, which typically has a pH of 7.2 and an organic matter content of about 2.9 percent, in plastic pots with a surface area of 128 square centimeters (cm<sup>2</sup>). The growing medium was steam sterilized. The plants were grown for 7-19 days in a greenhouse with an approximate 14-hour (h) photoperiod which was maintained at about 29 °C during the day and 26 °C during the night. Nutrients and water were added on a regular basis and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were treated with postemergence foliar applications when they reached the third to fourth true leaf stage. All treatments were applied using a randomized complete block trial design, with 4 replications per treatment.

#### Evaluation of Postemergence Herbicidal Activity of Mixtures in the Greenhouse

**[0032]** Treatments consisted of the compounds as listed in Tables 1, 3, 5 and 6, each compound applied alone and in combination. Formulated amounts of penoxsulam, halosulfuron-methyl, imazamox and fluroxypyr-meptyl ester, were placed in 60 milliliter (mL) glass vials and dissolved in a volume of 60 mL of a water solution containing Agri-dex crop oil concentrate in a 1% volume per volume (v/v) ratio. Compound requirements are based upon a 12 mL application volume at a rate of 187 liters per hectare (L/ha). Spray solutions of the mixtures were prepared by adding the stock solutions to the appropriate amount of dilution solution to form 12 mL spray solution with active ingredients in single and two way combinations. Formulated compounds were applied to the plant material with an overhead Mandel track sprayer equipped with 8002E nozzles calibrated to deliver 187 L/ha at a spray height of 18 inches (43 centimeters (cm)) above average plant canopy.

**[0033]** The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. Treatments were rated at 7 to 21 days after application (DAA) as compared to the untreated control plants. Visual weed control was scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

#### Evaluation of Postemergence Herbicidal Activity of Mixtures in the Field

**[0034]** Field trials were conducted in rice and turf using standard herbicide small plot research methodology. Plots varied from 3 x 3 meter (m) to 3 x 10 m (width x length) with 4 replicates per treatment. The rice crop was grown using normal cultural practices for fertilization, seeding, watering, flooding and maintenance to ensure good growth of the crop and the weeds. The turf crop was a permanent established crop of centipedegrass that was grown and maintained under normal cultural practices for fertilization, watering, mowing and disease maintenance to ensure good growth of the crop and the weeds.

**[0035]** All treatments in the field trials were applied using a carbon dioxide (CO<sub>2</sub>) backpack sprayer calibrated to apply 187 L/ha spray volume. Commercially available products of penoxsulam and fluroxypyr-meptyl were mixed in water at appropriately formulated product rates to achieve the desired rates based on a unit area of application (hectare) to achieve the desired rates as shown. Treatments were rated at 7 to 33 days after application as compared to the untreated control plants. Visual weed control was scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

**[0036]** Table 2 demonstrates the herbicidal synergistic efficacy of penoxsulam + fluroxypyr-meptyl tank mixes on weed control. Table 4 demonstrates the herbicidal synergistic safening of two crops to mixtures of penoxsulam+ fluroxypyr-meptyl. Table 7 demonstrates the herbicidal synergistic efficacy of imazethapyr + fluroxypyr-meptyl tank mixes on weed control. All treatment results, both for the single product and mixtures, are an average of 3 to 4 replicates and the tank mix interactions are significant at the P>0.05 level.

**[0037]** Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 1967, 15, 20-22.).

**[0038]** The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

**[0039]** Some of the compounds tested, application rates employed, plant species tested, and results are given in Tables 1-7. All comparisons are an average of 3-4 replicates and are significant at the P>0.05 level. Rates of penoxsulam, halosulfuron-methyl, imazamox and imazethapyr are expressed in grams active ingredient/hectare (g ai/ha) and rates of fluroxypyr are expressed in grams acid equivalent (g ae)/hectare in Tables 1-7.

Table 1. Synergistic Activity of Herbicidal Compositions of Penoxsulam + Fluroxypyrmeptyl on grass weeds (*Echinochloa crus-galli* (ECHCG)) in the greenhouse.

Application Rates		% Control	
Application Rates		ECHCG	
Penoxsulam (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex
15	0	62	-
0	340	0	-
15	340	80	62
15	0	84	-
0	340	0	-
15	340	90	84
15	0	78	-
0	340	0	-
15	340	93	78

Table 2. Synergistic Activity of Herbicidal Compositions of Penoxsulam + Fluroxypyrmeptyl on broadleaf weeds (*Plantago lanceolata*, PLALA; *Caperonia palustris*, CNPPA; and *Sebex exaltata*, SEBEX) in the field.

Application Rate		% Control					
Application Rate		PLALA		CNPPA		SEBEX	
Penoxsulam (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex	Ob	Ex	Ob	Ex
15	0	13	-	-	-	-	-
0	140	0	-	-	-	-	-
15	140	83	13	-	-	-	-
35	0	-	-	67	-	-	-
0	97	-	-	45	-	-	-
35	97	-	-	93	82	-	-
35	0	-	-	67	-	-	-
0	290	-	-	60	-	-	-
35	290	-	-	97	87	-	-
22	0	-	-	-	-	76	-
0	97	-	-	-	-	57	-
22	97	-	-	-	-	100	90

**EP 3 036 993 A2**

Table 3. Synergistic Activity of Herbicidal Compositions of Penoxsulam + Fluroxypyrmeptyl on perennial rice weed *Scirpus juncooides* (SCPJU) in the greenhouse.

Application Rate		% Control	
Application Rate		SCPJU	
Penoxsulam (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex
10	0	2	-
0	70	60	-
10	70	66	60

Table 4. Synergistic Activity of Herbicidal Compositions of Penoxsulam + Fluroxypyrmeptyl on safening of injury in Sunflower (HELAN) and Centipedegrass (ERLOP) in the field.

Application Rate		% Injury			
Application Rate		HELAN		ERLOP	
Penoxsulam (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex	Ob	Ex
7.5	0	36	-	-	-
0	100	31	-	-	-
7.5	100	36	56	-	-
15	0	40	-	-	-
0	100	34	-	-	-
15	100	50	60	-	-
30	0	41	-	-	-
0	100	34	-	-	-
30	100	46	61	-	-
70	0	-	-	15	-
0	210	-	-	6	-
70	210	-	-	13	24

Table 5. Synergistic Activity of Herbicidal Compositions of Halosulfuron-methyl + Penoxsulam on rice weeds in the greenhouse (rated 21 days after application (DAA)).

Application Rate		% Control					
Application Rate		ECHCG		CYPES		LEFCH	
Halosulfuron-methyl (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex	Ob	Ex	Ob	Ex
4.4	0	-	-	68	-	-	-
0	50	-	-	1	-	-	-
4.4	50	-	-	78	68	-	-
4.4	0	-	-	68	-	-	-
0	100	-	-	3	-	-	-
4.4	100	-	-	77	69	-	-
4.4	0	-	-	68	-	2	-
0	200	-	-	8	-	15	-
4.4	200	-	-	84	71	40	17
8.8	0	3	-	73	-	-	-
0	50	6	-	1	-	-	-

EP 3 036 993 A2

(continued)

Application Rate		% Control					
Application Rate		ECHCG		CYPES		LEFCH	
Halosulfuron-methyl (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex	Ob	Ex	Ob	Ex
8.8	50	30	9	84	73	-	-
8.8	0	3	-	-	-	-	-
0	100	2	-	-	-	-	-
8.8	100	29	4	-	-	-	-
8.8	0	3	-	73	-	-	-
0	200	11	-	8	-	-	-
8.8	200	24	14	92	75	-	-
18	0	6	-	78	-	-	-
0	50	6	-	1	-	-	-
18	50	33	11	92	78	-	-
18	0	6	-	-	-	-	-
0	100	2	-	-	-	-	-
18	100	34	7	-	-	-	-
18	0	6	-	78	-	8	-
0	200	11	-	8	-	15	-
18	200	33	16	98	79	50	22

Table 6. Synergistic Activity of Herbicidal Compositions of Imazamox + Penoxsulam on rice weeds in the greenhouse (rated 21 days after application (DAA)).

Application Rate		% Control					
Application Rate		ECHCG		BRAPP		CYPES	
Imazamox (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex	Ob	Ex	Ob	Ex
4.4	0	1.3	-	3	-	-	-
0	50	1.3	-	10	-	-	-
4.4	50	32	2.4	55	12	-	-
4.4	0	1.3	-	3	-	-	-
0	100	4	-	23	-	-	-
4.4	100	66	5	50	25	-	-
4.4	0	1.3	-	3	-	-	-
0	200	6	-	20	-	-	-
4.4	200	84	7	60	21	-	-
8.8	0	29	-	46	-	-	-
0	50	1.3	-	10	-	-	-
8.8	50	100	30	76	52	-	-
8.8	0	29	-	46	-	8	-
0	100	4	-	23	-	10	-
8.8	100	100	31	86	58	39	17
8.8	0	29	-	46	-	8	-
0	200	6	-	20	-	9	-
8.8	200	100	33	92	57	33	16

EP 3 036 993 A2

(continued)

Application Rate		% Control					
		ECHCG		BRAPP		CYPES	
Imazamox (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex	Ob	Ex	Ob	Ex
18	0	85	-	-	-	-	-
0	50	1.3	-	-	-	-	-
18	50	100	85	-	-	-	-
18	0	-	-	-	-	42	-
0	200	-	-	-	-	9	-
18	200	-	-	-	-	57	47

Table 7. Synergistic Activity of Herbicidal Compositions of Imazethapyr + Fluroxypyr- meptyl on grass weeds (*Leptochloa* spp, LEFSS and *Panicum dichotomiflorum*, PANDI) in the field.

Application Rate		% Control			
		LEFSS		PANDI	
Imazethapyr (g ai/ha)	Fluroxypyr-meptyl (g ae/ha)	Ob	Ex	Ob	Ex
70	0	28	-	40	-
0	290	0	-	0	-
70	290	76	28	95	40

BRAPP = *Brachiaria platyphylla*; broadleaf signalgrass  
 CNPPA = *Caperonia palustris*; Texasweed  
 CYPES = *Cyperus esculentus*; yellow nutsedge  
 ECHCG = *Echinochloa crus-galli*; barnyardgrass  
 ERLOP, = *Eremochloa ophiuroide*; centipedegrass  
 HELAN = *Helianthus annuus*; sunflower  
 LEFCH = *Leptochlon chinensis*; Chinese sprangletop  
 LEFSS = *Leptochloa* spp, sprangletop  
 PANDI = *Panicum dichotomiflorum*  
 PLALA = *Plantago lanceolata* L.; narrow-leaved plantain  
 SCPJU = *Scirpus juncoide*; Japanese bulrush  
 SEBEX = *Sesbania exaltata*; Hemp sesbania  
 Ob = observed value (% control)  
 Ex = expected, calculated value using Colby Analysis (% control)  
 DAA = days after application  
 g ai/ha = grams active ingredient per hectare  
 g ae/ha = grams acid equivalent per hectare

ASPECTS OF THE INVENTION:

[0040]

1. A synergistic herbicidal mixture comprising an herbicidally effective amount of (a) fluroxypyr and (b) an ALS inhibitor herbicide.
2. The mixture of Aspect 1 in which fluroxypyr is the acid itself or an agriculturally acceptable salt or ester.
3. The mixture of Aspect 2 in which fluroxypyr is the meptyl ester.

4. The mixture of Aspect 1 in which the ALS inhibitor herbicide is from the sulfonamide class of herbicides.
5. The mixture of Aspect 4 in which the ALS inhibitor herbicide is penoxsulam.
- 5 6. The mixture of Aspect 1 in which the ALS inhibitor herbicide is from the sulfonurea class of herbicides.
7. The mixture of Aspect 6 in which the ALS inhibitor herbicide is halosulfuron-methyl.
8. The mixture of Aspect 1 in which the ALS inhibitor herbicide is from the imidazolinone class of herbicides.
- 10 9. The mixture of Aspect 8 in which the ALS inhibitor herbicides are imazamox and imazethapyr.
10. The mixture of Aspect 1 in which the weight ratio of fluroxypyr (acid equivalent) to the ALS inhibitor herbicide (active ingredient) is between about 1:2 and about 140:1.
- 15 11. A method for protecting centipedegrass and sunflower from the individual harmful effects of penoxsulam and fluroxypyr which comprises contacting centipedegrass or sunflower with, or applying to the area under cultivation, a safening amount of a synergistic mixture of penoxsulam and fluroxypyr.
- 20 12. An herbicidal composition comprising an herbicidally effective amount of the herbicidal mixture of Aspect 1 and an agriculturally acceptable adjuvant and/or carrier.
13. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation an herbicidally effective amount the herbicidal mixture of Aspect 1.
- 25 14. A method of controlling undesirable vegetation in rice, cereal and grain crops, pastures, rangelands, IVM and turf which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation an herbicidally effective amount the herbicidal mixture of Aspect 1.
- 30

### Claims

- 35 1. A synergistic herbicidal mixture comprising an herbicidally effective amount of (a) fluroxypyr, or an agriculturally acceptable salt or ester, and (b) an ALS inhibitor herbicide, wherein the ALS inhibitor herbicide is halosulfuron-methyl or is from the imidazolinone class of herbicides.
- 40 2. The mixture of claim 1, wherein the fluroxypyr, or an agriculturally acceptable salt or ester thereof, is the meptyl ester of fluroxypyr.
3. The mixture of claim 1 or claim 2, wherein the weight ratio of fluroxypyr (acid equivalent) to the ALS inhibitor herbicide (active ingredient) is between 1:2 and 140:1.
- 45 4. The mixture of any of claims 1-3, wherein the ALS inhibitor herbicide is halo sulfuron-methyl.
5. The mixture of any of claims 1-3, wherein the ALS inhibitor herbicide is from the imidazolinone class of herbicides and is selected from the group consisting of imazamox and imazethapyr.
- 50 6. An herbicidal composition comprising an herbicidally effective amount of the herbicidal mixture of any of claims 1-5 and an agriculturally acceptable adjuvant and/or carrier.
7. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation an herbicidally effective amount of the herbicidal mixture of any of claims 1-5.
- 55 8. The method of claim 7, wherein the undesirable vegetation is controlled in rice, cereal crops, grain crops, pastures, rangelands, industrial vegetation management, turf, wheat, barley, oats, rye, sorghum, corn, maize, grasslands, fallowland, or aquatics.

**EP 3 036 993 A2**

9. The method of claim 7 or claim 8, wherein the undesirable vegetation is controlled in rice.
10. The method of any of claims 7-9, wherein the ALS inhibitor herbicide is applied at a rate between 4 g ai/ha and 100 g ai/ha and the fluroxypyr is applied at a rate between 50 g ae/ha and 560 g ae/ha.
11. The method of any of claims 7-10, wherein the ALS inhibitor herbicide is halosulfuron-methyl and wherein the undesirable vegetation is barnyard grass, Chinese sprangletop, or yellow nutsedge.
12. The method of any of claims 7-10, wherein the ALS inhibitor herbicide is imazamox and wherein the undesirable vegetation is barnyardgrass, broadleaf signalgrass, or yellow nutsedge.
13. The method of any of claims 7-10, wherein the ALS inhibitor herbicide is imazethapyr and wherein the undesirable vegetation is sprangletop grass or fall panicum.
14. The method of any of claims 7-13, wherein the herbicidal mixture or composition is applied postemergence.
15. The method of any of claims 7-14, wherein the components of the synergistic mixture are applied either separately or as part of a multipart herbicidal system.

5

10

15

20

25

30

35

40

45

50

55

**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Non-patent literature cited in the description**

- Herbicide Handbook of the Weed Science Society of America. 2002, 462 [0003]
- The Pesticide Manual. 2006 [0008] [0009] [0010] [0011] [0012]
- McCutcheon's Detergents and Emulsifiers Annual. MC Publishing Corp, 1998 [0026]
- Encyclopedia of Surfactants. Chemical Publishing Co, 1980, vol. I-III [0026]
- COLBY, S.R. Calculation of the synergistic and antagonistic response of herbicide combinations. *Weeds*, 1967, vol. 15, 20-22 [0037]

## 摘要

包含(a)氟草烟和(b)ALS 抑制剂除草剂的除草协同组合物，其对水稻、谷类和谷物作物、牧地、牧场、IVM 和草皮提供改善的芽后杂草控制，其中 ALS 抑制剂除草剂是氯吡啶磺隆、甲氧咪草烟或咪唑乙烟酸。