



US008187839B2

(12) **United States Patent**
Li et al.(10) **Patent No.:** **US 8,187,839 B2**
(45) **Date of Patent:** **May 29, 2012**(54) **ANTI-NOTCH3 AGONIST ANTIBODIES AND
THEIR USE IN THE TREATMENT OF
NOTCH3-RELATED DISEASES**2008/0131908 A1 6/2008 Li et al.
2008/0226621 A1 9/2008 Fung et al.
2011/0223155 A1 9/2011 Fung et al.

FOREIGN PATENT DOCUMENTS

(75) Inventors: **Kang Li**, San Diego, CA (US); **Bin-Bing
Stephen Zhou**, Ho Hokus, NJ (US);
Wenjuan Wu, Houston, TX (US); **Sek
Chung Fung**, Gaithersburg, MD (US);
Sanjaya Singh, Sandy Hook, CT (US)FR 2 777 285 A1 10/1999
WO 95/15982 6/1995
WO 00/20576 4/2000
WO WO 02/24221 A2 3/2002
WO 2006/017173 A1 2/2006
WO WO 2006/053063 A2 5/2006
WO WO 2006/068822 A1 6/2006
WO WO 2006/015375 A2 9/2006
WO WO 2008/057144 A2 5/2008
WO WO 2008/150525 A1 12/2008(73) Assignee: **Genentech, Inc.**, South San Francisco,
CA (US)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

OTHER PUBLICATIONS

(21) Appl. No.: **13/033,500**(22) Filed: **Feb. 23, 2011**(65) **Prior Publication Data**

US 2011/0206675 A1 Aug. 25, 2011

Related U.S. Application Data(62) Division of application No. 11/874,682, filed on Oct.
18, 2007, now Pat. No. 7,915,390.(60) Provisional application No. 60/879,218, filed on Jan.
6, 2007, provisional application No. 60/875,597, filed
on Dec. 18, 2006, provisional application No.
60/852,861, filed on Oct. 19, 2006.(51) **Int. Cl.**
C12P 21/06 (2006.01)(52) **U.S. Cl.** **435/69.1**; 435/320.1; 435/325;
536/23.5; 530/350; 530/387.9; 530/388.22(58) **Field of Classification Search** 435/69.1,
435/320.1, 325; 536/23.5; 530/350, 387.9,
530/388.22

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,648,464 A 7/1997 Artavanis-Tsakonas et al.
5,780,300 A 7/1998 Artavanis-Tsakonas et al.
5,786,158 A 7/1998 Artavanis-Tsakonas et al.
5,789,195 A 8/1998 Artavanis-Tsakonas et al.
6,083,904 A 7/2000 Artavanis-Tsakonas
6,090,922 A 7/2000 Artavanis-Tsakonas et al.
6,149,902 A 11/2000 Artavanis-Tsakonas et al.
6,436,650 B1 8/2002 Artavanis-Tsakonas et al.
6,692,919 B1 2/2004 Artavanis-Tsakonas et al.
7,935,791 B2 5/2011 Fung et al.
7,994,285 B2 8/2011 Li et al.
2002/0151487 A1 10/2002 Nickoloff et al.
2003/0186290 A1 10/2003 Tournier-Lasserre et al.
2004/0058443 A1 3/2004 Artavanis-Tsakonas et al.
2004/0242482 A1 12/2004 Gehring et al.
2005/0112121 A1 5/2005 Artavanis-Tsakonas et al.
2005/0158859 A1 7/2005 Artavanis-Tsakonas et al.
2005/0208027 A1 9/2005 Conboy et al.
2006/0002924 A1 1/2006 Bodmer et al.
2007/0003983 A1 1/2007 Artavanis-Tsakonas et al.
2008/0107648 A1 5/2008 Noguera et al.Allenspach et al., "Notch signaling in cancer" *Cancer Biol Ther.*
1(5):466-76 (2002).Anastasi et al., "Expression of activated Notch3 in transgenic mice
enhances generation of T regulatory cells and protects against experi-
mental autoimmune diabetes" *J Immunol.* 171(9):4504-11 (Nov.
2003).Androutsellis-Theotokis et al., "Notch signalling regulates stem cell
numbers in vitro and in vivo" *Nature* 442:823-6 (Aug. 2006)Artavanis-Tsakonas et al., "Notch Signaling" *Science* 268:225-232
(Apr. 14, 1995).Artavanis-Tsakonas et al., "Notch Signaling: Cell Fate Control and
Signal Integration in Development" *Science* 284:770-776 (1999).Aster et al., "The folding and structural integrity of the first LIN-12
module of human Notch1 are calcium-dependent" *Biochemistry*
38(15):4736-42 (Apr. 1999).Bellavia et al., "Combined expression of pTalpha and Notch3 in T
cell leukemia identifies the requirement of preTCR for
leukemogenesis" *Proc Natl Acad Sci U S A.* 99(5):3788-93 (Mar.
2002).Bellavia et al., "Notch3: from subtle structural differences to func-
tional diversity" *Oncogene* 27:5092-5098 (2008).Bocchetta et al., "Notch-1 induction, a novel activity of SV40
required for growth of SV40-transformed human mesothelial cells"
Oncogene 22(1):81-9 (Jan. 2003).Bolos et al., "Notch signaling in development and cancer" *Endocr
Rev.* 28(3):339-63 (May 2007).Bray, "Notch signalling: a simple pathway becomes complex" *Nat
Rev Mol Cell Biol.* 7(9):678-89 (Sep. 2006).Buchler et al., "The Notch signaling pathway is related to
neurovascular progression of pancreatic cancer" *Ann Surg.*
242(6):791-800 (Dec. 2005).Caldas, Cristina. et al., "Humanization of the anti-CD18 antibody
6.7: an unexpected effect of a framework residue in binding to anti-
gen" *Molecular Immunology* 39:941-952 (2003).Casset, F. et al., "A peptide mimetic of an anti-CD4 monoclonal
antibody by rational design" *Biochem. & Biophys. Res. Comm.*
307:198-205 (2003).

(Continued)

Primary Examiner — Stephen Rawlings

(74) Attorney, Agent, or Firm — Danielle Pasqualone; Julia
vom Wege(57) **ABSTRACT**The present invention relates to agonist antibodies that spe-
cifically bind to Notch 3 and activate signaling. The present
invention includes antibodies binding to an epitope compris-
ing the first Lin12 domain. The present invention also
includes uses of these antibodies to treat or prevent Notch 3
related diseases or disorders.**15 Claims, 18 Drawing Sheets**

OTHER PUBLICATIONS

- Chiba, "Notch signaling in stem cell systems" *Stem Cells* 24(11):2437-47 (Nov. 2006).
- Chien, Nadine C. et al., "Significant structural and functional change of an antigen-binding site by a distant amino acid substitution: Proposal of a structural mechanism" *Proc. Natl. Acad. Sci. USA* 86:5532-5536 (Jul. 1989).
- Coffman et al., "Expression of an extracellular deletion of Notch diverts cell fate in *Xenopus* embryos" *Cell* 73(4):659-71 (May 1993).
- Davies et al., "Affinity improvement of single antibody VH domains: residues in all three hypervariable regions affect antigen binding" *Immunotechnology* 2:169-179 (1996).
- De Pascalis et al., "Grafting of "abbreviated" complementarity-determining regions containing specificity-determining residues essential for ligand contact to engineer a less immunogenic humanized monoclonal antibody" *J. Immunol.* 169:3076-3084 (2002).
- Domenga et al., "Notch3 is required for arterial identity and maturation of vascular smooth muscle cells" *Genes Dev.* 18(22):2730-5 (Nov. 2004).
- Ellisen et al., "TAN-1, the human homolog of the *Drosophila* notch gene, is broken by chromosomal translocations in T lymphoblastic neoplasms" *Cell* 66(4):649-61 (Aug. 1991).
- Fan et al., "Notch pathway inhibition depletes stem-like cells and blocks engraftment in embryonal brain tumors" *Cancer Research* 66(15):7445-52 (Aug. 2006).
- Flynn et al., "The role of Notch receptor expression in bile duct development and disease" *J. Pathol.* 204(1):55-64 (Sep. 2004).
- Fre et al., "Notch signals control the fate of immature progenitor cells in the intestine" *Nature* 435:964-8 (Jun. 2005).
- George et al. *Circulation* 97:900-906 (1998).
- Giusti et al., "Somatic Diversification of S107 from an Antiphosphocholine to an anti-DNA Autoantibody is Due to a Single Base Change in its Heavy Chain Variable Region" *Proc. Natl. Acad. Sci. USA* 84:2926-2930 (May 1987).
- Gordon et al., "Structural basis for autoinhibition of Notch" *Nat Struct Mol Biol.* 14(4):295-300 (Apr. 2007).
- Greenspan et al., "Defining epitopes: It's not as easy as it seems" *Nature Biotechnology* 7:936-937 (Oct. 1999).
- Gussow & Seemann, "Humanization of Monoclonal Antibodies" *Meth. Enzymology*, Academic Press, Inc. vol. 203:99-121 (1991).
- Haruki et al., "Dominant-negative Notch3 receptor inhibits mitogen-activated protein kinase pathway and the growth of human lung cancers" *Cancer Research* 65(9):3555-61 (May 2005).
- Hedvat et al., "Insights into extramedullary tumour cell growth revealed by expression profiling of human plasmacytomas and multiple myeloma" *Br J Haematol.* 122(5):728-44 (Sep. 2003).
- Heller et al., "Amino Acids at the Site of V-J Recombination Not Encoded by Germline Sequences" *Journal of Experimental Medicine* 166:637-646 (1987).
- Holm, P. et al., "Functional mapping and single chain construction of the anti-cytokeratin 8 monoclonal antibody TS1" *Molecular Immunology* 44:1075-1084 (2007).
- Holt et al., "Domain antibodies: proteins for therapy" *Trends Biotechnol.* 21(11):484-490 (Nov. 2003).
- Houde et al., "Overexpression of the NOTCH ligand JAG2 in malignant plasma cells from multiple myeloma patients and cell lines" *Blood* 104(12):3697-704 (Dec. 2004).
- Hu et al., "Overexpression of activated murine Notch1 and Notch3 in transgenic mice blocks mammary gland development and induces mammary tumors" *Am J Pathol.* 168(3):973-90 (Mar. 2006).
- Jang et al., "Notch signaling as a target in multimodality cancer therapy" *Curr Opin Mol Ther.* 2(1):55-65 (Feb. 2000).
- Jiang et al., "A Novel Peptide Isolated from a Phage Display Peptide Library with Trastuzumab Can Mimic Antigen Epitope of HER-2" *Journal of Biological Chemistry* 280(6):4656-4662 (2005).
- Joutel et al., "Notch signalling pathway and human diseases" *Semin Cell Dev Biol.* 9(6):619-25 (Dec. 1998).
- Joutel et al., "Notch3 mutations in CADASIL, a hereditary adult-onset condition causing stroke and dementia" *Nature* 383:707-10 (Oct. 1996).
- Joutel et al., "Skin biopsy immunostaining with a Notch3 monoclonal antibody for CADASIL diagnosis" *Lancet* 358:2049-2051 (2001).
- Jundt et al., "Jagged1-induced Notch signaling drives proliferation of multiple myeloma cells" *Blood* 103(9):3511-5 (May 2004).
- Juryneczyk et al., "Notch3 Inhibition in Myelin-Reactive T Cells Down-Regulates Protein Kinase C θ and Attenuates Experimental Autoimmune Encephalomyelitis" *Journal of Immunology* 180(4):2634-2640 (2008).
- Kadesch, "Notch signaling: a dance of proteins changing partners" *Exp Cell Res.* 260(1):1-8 (Oct. 2000).
- Kidd et al., "Sequence of the notch locus of *Drosophila melanogaster*: relationship of the encoded protein to mammalian clotting and growth factors" *Mol Cell Biol.* 6(9):3094-108 (Sep. 1986).
- Kopczynski et al., "Delta, a *Drosophila* neurogenic gene, is transcriptionally complex and encodes a protein related to blood coagulation factors and epidermal growth factor of vertebrates" *Genes Dev.* 2:1723-35 (1988).
- Leong et al., "Recent insights into the role of Notch signaling in tumorigenesis" *Blood* 107(6):2223-33 (Mar. 2006).
- Li et al., "Modulation of Notch Signaling by Antibodies Specific for the Extracellular Negative Regulatory Region of NOTCH3" *Journal of Biological Chemistry* 283(12):8046-8054 (Mar. 21, 2008).
- Lu et al., "Selection of potential markers for epithelial ovarian cancer with gene expression arrays and recursive descent partition analysis" *Clin Cancer Res.* 10(10):3291-300. (May 2004).
- MacCallum et al., "Antibody-antigen Interactions: Contact Analysis and Binding Site Topography" *J. Mol. Biol.* 262:732-745 (1996).
- Mailhos et al., "Delta4, an Endothelial Specific Notch Ligand Expressed at Sites of Physiological and Tumor Angiogenesis" *Differentiation* 69:135-144 (2001).
- Malecki et al., "Leukemia-associated mutations within the NOTCH1 heterodimerization domain fall into at least two distinct mechanistic classes" *Mol Cell Biol.* 26(12):4642-51 (Jun. 2006).
- Mariuzza et al., "The Structure Basis of Antigen-Antibody Recognition" *Ann. Rev. Biophys. Biophys. Chem.* 16:139-159 (1987).
- Nam et al., "Notch signaling as a therapeutic target" *Curr Opin Chem Biol.* 6(4):501-9 (Aug. 2002).
- Park et al., "Notch3 gene amplification in ovarian cancer" *Cancer Research* 66(12):6312-8 (Jun. 2006).
- PCT International Search Report mailed Jun. 10, 2008, in counterpart PCT Application No. PCT/US2007/081797, filed Oct. 18, 2007.
- Rao et al., "K Chain Variable Regions from Three Galactan Binding Myeloma Proteins" *Biochemistry* 17(25):5555-5559 (1978).
- Reya et al., "Stem cells, cancer, and cancer stem cells" *Nature* 414:105-11 (2001).
- Riemer et al., "Matching of trastuzumab (Herceptin) epitope mimics onto the surface of Her-2/neu—a new method of epitope definition" *Molecular Immunology* 42:1121-1124 (2005).
- Rudikoff et al., "K chain joining segments and structural diversity of antibody combining sites" *Proc. Natl. Acad. Sci. USA* 77(7):4270-4274 (1980).
- Rudikoff et al., "Single amino acid substitution altering antigen-binding specificity" *Proc. Natl. Acad. Sci. USA* 79:1979-1983 (Mar. 1982).
- Sanchez-Irizarry et al., "Notch subunit heterodimerization and prevention of ligand-independent proteolytic activation depend, respectively, on a novel domain and the LNR repeats" *Mol Cell Biol.* 24(21):9265-73 (2004).
- Screpanti et al., "Notch, a unifying target in T-cell acute lymphoblastic leukemia?" *Trends Mol Med.* 9(1):30-5 (Jan. 2003).
- Shimizu et al., "Physical interaction of Delta1, Jagged1, and Jagged2 with Notch1 and Notch3 receptors" *Biochem Biophys Res Commun.* 276(1):385-9 (Sep. 2000).
- Stancovski et al., "Mechanistic Aspects of the Opposing Effects of Monoclonal Antibodies to the ERBB2 Receptor on Tumor Growth" *Proc. Natl. Acad. Sci. USA* 88:8691-8695 (1991).
- Sullivan and Bicknell, "New molecular pathways in angiogenesis" *British Journal of Cancer* 89(2):228-231 (Jul. 21, 2003).
- Sweeney et al., "Notch 1 and 3 receptor signaling modulates vascular smooth muscle cell growth, apoptosis, and migration via a CBF-1/RBP-Jk dependent pathway" *FASEB J.* 18(12):1421-3 (Sep. 2004).
- Swiatek et al., "Notch1 is essential for postimplantation development in mice" *Genes Dev.* 8(6):707-19 (Mar. 1994).

- Taichman et al., "Notch1 and Jagged1 expression by the developing pulmonary vasculature" *Dev Dyn.* 225(2):166-75 (Oct. 2002).
- Thelu et al., "Notch signalling is linked to epidermal cell differentiation level in basal cell carcinoma, psoriasis and wound healing" *BMC Dermatol.* 2:7 (Apr. 2002).
- Vacca et al., "Notch3 and pre-TCR interaction unveils distinct NF-kappaB pathways in T-cell development and leukemia" *EMBO Journal* 25(5):1000-8 (Mar. 2006).
- Vajdos et al., "Comprehensive functional maps of the antigen-binding site of an anti-ErbB2 antibody obtained with shotgun scanning mutagenesis" *J Mol Biol.* 320:415-428 (2002).
- van Es et al., "Notch/gamma-secretase inhibition turns proliferative cells in intestinal crypts and adenomas into goblet cells" *Nature* 435:959-63 (Jun. 2005).
- van Limpt et al., "Phox2B mutations and the Delta-Notch pathway in neuroblastoma" *Cancer Lett.* 228:59-63 (Oct. 2005).
- Vardar et al., "Nuclear magnetic resonance structure of a prototype Lin12-Notch repeat module from human Notch1" *Biochemistry* 42(23):7061-7 (Jun. 2003).
- Von Boehmer, "Notch in lymphopoiesis and T cell polarization" *Nat Immunol.* 6(7):641-2 (Jul. 2005).
- Weijzen et al., "Activation of Notch-1 signaling maintains the neoplastic phenotype in human Ras-transformed cells" *Nat Med.* 8(9):979-986 (Sep. 2002).
- Weng et al., "Activating mutations of NOTCH1 in human T cell acute lymphoblastic leukemia" *Science* 306(5694):269-271 (Oct. 8, 2004).
- Wharton et al., "Nucleotide sequence from the neurogenic locus notch implies a gene product that shares homology with proteins containing EGF-like repeats" *Cell* 43:567-81 (Dec. 1985).
- Winkler et al. *Journal of Immunology* 165(8):4505-4514 (2000).
- Wu et al., "Humanization of a Murine Monoclonal Antibody by Simultaneous Optimization of Framework and CDR Residues" *J. Mol. Biol.* 294:151-162 (1999).
- Xu et al., "Antibody-Induced Growth Inhibition is Mediated Through Immunochemically and Functionally Distinct Epitopes on the Extracellular Domain of the c-erbB-2 (HER-2/neu) Gene Product p185" *International Journal of Cancer* 53(3):401-408 (Feb. 1, 1993).
- Xu et al., "Regions of *Drosophila* Notch that contribute to ligand binding and the modulatory influence of Fringe" *J Biol Chem.* 280(34):30158-65 (Aug. 2005).
- Yabe et al., "Immunohistological localization of Notch receptors and their ligands Delta and Jagged in synovial tissues of rheumatoid arthritis" *J Orthop Sci.* 10(6):589-94 (Nov. 2005).
- Yochem et al., "The *Caenorhabditis elegans* lin-12 gene encodes a transmembrane protein with overall similarity to *Drosophila* Notch" *Nature* 335:547-50 (Oct. 1988).
- Zeng et al., "Crosstalk between tumor and endothelial cells promotes tumor angiogenesis by MAPK activation of Notch signaling" *Cancer Cell* 8(1):13-23 (Jul. 2005).
- Zweidler-McKay et al., "Notch signaling is a potent inducer of growth arrest and apoptosis in a wide range of B-cell malignancies" *Blood* 106(12):3898-906 (Dec. 2005).
- U.S. Appl. No. 13/174,285, Kang Li et al.
- Maynard et al., "Antibody Engineering" *Annu. Rev. Biomed. Eng.* (2):339-379 (2000).
- Pini et al., "Design and Use of a Phage Display Library" *The Journal of Biological Chemistry* 273:21769-21776 (1998).

FIGURE 1

Amino Acid Sequence of Human Notch 3 (NP_000426)

```

1  MGPARGRRR RRRPMSPPPP PPFVRALPLL LLLAGPGAAA PPCLDGSPCA NGGRCTQLPS
61  REAACLCPPG WGERCQLED PCHSGPCAGR GVCQSSVVAG TARFSCRCPR GFRGPDCLSP
121 DPCLSPPCAH GARCSVGPDG RFLCSCPPGY QGRSCRSDVD ECRVGEPCRH GGTCLNTPGS
181 FRCQCPAGYT GPLCENPAVP CAPSPCRNGG TCRQSGDLTY DCACLPGFEG QNCEVNVDDC
241 PGHRCLNGGT CVDGVNTYNC QCPPEWTGQF CTEDVDECOQ QPNACHNGGT CFNTLGGHSC
301 VCVNGWTGES CSQNIDDCAT AVCFHGATCH DRVASFYCAC PMGKTGLLCH LDDACVSNPC
361 HEDAICDTNF VNGRAICTCP PGFTGGACDQ DVDECSIGAN PCEHLGRCVN TQGSFLCQCG
421 RGYTGPRCET DVNECLSGPC RNQATCLDRI GQFTCICMAG FTGTYCEVDI DECQSSPCVN
481 GGVCKDRVNG FSCTCPSGFS GSTCQLDVDE CASTPCRNGA KCVDQPDGYE CRCAEGFEGT
541 LCDRNVDCCS FDPCHHGRCV DGIASFSCAC APGYTGTRCE SQVDECRSQP CRHGGKCLDL
601 VDKYLCRCPS GTTGVNCEVN IDDCASNPCT FGVCRDGINR YDCVCQPGFT GPLCNVEINE
661 CASSPCGEGG SCVDGENGFR CLCPGSLPP LCLPPSHPCA HEPCSHGICY DAPGGFRCVC
721 EPGWSGPRCS QSLARDACES QPCRAGGTCS SDGMGFHCTC PPGVQGRQCE LLSPECTPNPC
781 EHGGRCESAP GQLPVCSCPQ GWQGPQCQD VDECAGPAPC GPHGICTNLA GSFSCICHGG
841 YTGPSCDQDI NDCDPNPCLN GGSCQDGVGS FSCSCLPGFA GPRCARDVDE ESLNPCGPGT
901 CTDHVASFTC TCPPGYGGFH CEQDLDFCSP SSCFNNGTCV DGVNSFSCLC RPYGTGAHCQ
961 HEADPCLSRP CLHGGVCSAA HPGFRCTCLE SFTGPQCQTL VDWCSPQPCQ NGGRCVQTGA
1021 YCLCPPGWSG RLCDIRSLPC REAAAIQIGVR LEQLCQAGGQ CVDEDSSHYC VCFEGRTGSH
1081 CEQEVDPCLA QPCQHGGTCR GYMGGYMCEC LPGYNGDNCE DDVDECASQP CQHGGSCIDL
1141 VARYLCSCFP GTLGLVCEIN EDDCGPGPFL DSGPRCLHNG TCVDLVGGFR CTCFPGYTGL
1201 ECEADINECR SGACHAAHTR DCLQDPGGGF RCLCHAGFSG PRCQTVLSPC ESQPCQHGGQ
1261 CRPSPGPGGG LTFTCHCAQP FWGPRCERVA RSCRELQCPV GVPCQQTTPRG PRCACPPGLS
1321 GPSCRSFPGS PPGASNASCA AAPCLHGGSC RPAPLAPFFR CACAQGTGP RCEAPAAPE
1381 VSEEPRCPR ACQAKRGDQR CDRECNSPGC GWGGGDCSL VGDPWRQCEA LQCWRLFNN
1441 RCDPACSSPA CLYDNFDCHA GGRERTCNPV YEKYCADHFA DGRCDQGCNT EECGWDGLDC
1501 ASEVPALLAR GVLVLTVLLP PEELLRSSAD FLQRLSAILR TSLRFRILDAH GQAMVFPYHR
1561 PSPGSEPRAR RELAPEVIGS VVMLEIDNRL CLQSPENDHC FPDQAQSAADY LGALSAVERL
1621 DFPYPLRDVR GEPLPPEPS VPLLPLLVAG AVLLLVILVL GVMVARRKRE HSTLWFPEGF
1681 SLHKDVASGH KGRREPVGQD ALGMKNMAKG ESLMGEVATD WMDTECPKAK RLKVEEPPMG
1741 AEEAVDCRQW TQHHLVAADI RVAPAMALTP PQGDADADGM DVNVRGPDGT TPLMLASFCG
1801 GALEPMPTEE DEADDTSASI ISDLICQGAQ LGARTDRTGE TALHLAARYA RADAARKLLD
1861 AGADTNAQDH SGRTPPLHTAV TADAQGVFI LIRNRSTDLD ARMADGSTAL ILAARLAVEG
1921 MVEELIASHA DVNAVDELGK SALHWAAAVN NVEATLALLK NGANKDMQDS KEETPLFLAA
1981 REGSYEAAKL LLDHFANREI TDHLDRLPD VAQERLHQDI VRLLDQPSGP RSFFPGPHGLG
2041 PLLCPPGAFL PGLKAAQSGS KKSRRPPGKA GLGPQGPRGR GKKLTACPG PLADSSVTLS
2101 PVDSLDSPRP FGGPPASPGG FPLEGPYAAA TATAVSLAQL GPGGRAGLGR QPPGGCVLSL
2161 GLLNPVAVPL DWARLPPAP PGPSFLLPLA PGPQLLNPGT PVSPQERPPP YLAVFGHGEE
2221 YPVAGAHSSP FKARFLRVPS EHPYLTPSPE SPEHWASPS PSLSDWSEST PSFATATGAM
2281 ATTTGALPAQ PLPLSVPSL AQAQTQLGPQ PEVTPKRQVL A (SEQ ID NO 1)

```

FIGURE 2A Amino acid sequence Comparison of Notch1, 2, 3 and 4.

1	- - - - - - - - - - - - - - - - - M	P P L L A P L L C L	Notch1.pro
1	- - - - - - - - - - - - - - - - - M	P A L R P A L L W A L L A L	Notch2.pro
1	K G P G A R G R R R R R R P M S	P P P V R A L P L L	Notch3.pro
1	- - - - - - - - - - - - - - - - - M Q	P P S L L L L L L L	Notch4.pro
12	A L L P A L A A R - - G P R C S Q F G E T C L N G G K C E A		Notch1.pro
16	W L C C A A P A H - - A L Q C R D G Y E P C V N E G M C V T		Notch2.pro
31	L L L A G P G A A - - A P P C L D G - S P C A N G G R C T Q		Notch3.pro
13	L L L C V S V V R P R G L L C G S F P E P C A N G G T C L S		Notch4.pro
40	A N - G T E A C V C G G A F V G P R C Q C F N P C L - S T P		Notch1.pro
44	Y H N G T G Y C K C P E G F L G E Y C Q H R D P C E - K N R		Notch2.pro
58	L P S R E A A C L C P P G W V G E R C Q L E D P C H - S G P		Notch3.pro
43	L S L G Q G T C Q C A P G F L G E T C Q F P D P C Q N A Q L		Notch4.pro
68	C K N A G T C H V V D R R - - G V A D Y A - - - - - - - C		Notch1.pro
73	C Q N G G T C V A Q A M L - - G K A T - - - - - - - C		Notch2.pro
87	C A G R G V C Q S S V V A - - G T A R - - - - - - -		Notch3.pro
73	C Q N G G S C Q A L L P A P L G L F S S P S P L T P S F L C		Notch4.pro
88	S C A L G F S G P L C L T P L D N A C L T N - P C R N G G T		Notch1.pro
91	R C A S G F T G E D C Q Y S T S H F C F V S R P C L N G S T		Notch2.pro
104	- -		Notch3.pro
103	T C L P G F T G E R C Q A K L E D P C P P S - F C S K R G R		Notch4.pro
117	C D L L T L T E Y K C R C P P G W S G K S C Q Q A D P C A S		Notch1.pro
121	C H M L S R D T Y E C T C Q V G F T G K E C Q W T D A C L S		Notch2.pro
104	- - - - - - - F S C R C P R G F R G P C S L P D P C L S		Notch3.pro
132	C H I Q A S G R P Q C S C M P G W T G E Q C Q L R D F C S A		Notch4.pro
147	N P C A N G G Q C L P - F E A S Y I C H C P P S F H G P T C		Notch1.pro
151	H P C A N G S T C T T - V A N Q F S C K C L T G F T G Q K C		Notch2.pro
126	S P C A E G A R C S V G P D G R F - C S C P P G Y Q G R S C		Notch3.pro
162	N P C V N G G V C L A - T Y P Q I Q C H C P P G F E G H A C		Notch4.pro
176	R Q D V N E C G Q K P G L C R H G G T C H N E V G S Y R C V		Notch1.pro
180	E T D V N E C D - I P G H C Q H G G T C L N L P G S Y Q C Q		Notch2.pro
156	R S D V D E C R - V G E P C R H G G T C L N T P G S F R C Q		Notch3.pro
191	E R D V N E C F Q D P G P C P K G T S C H N T L G S F Q C L		Notch4.pro
206	C R A F T G P N C E R P Y V P C S P S P C Q N G G T C R P		Notch1.pro
209	C P Q G F T G Q Y C D S L Y V P C A P S P C V N G G T C R Q		Notch2.pro
185	C P A G V T G P L C E N P A V P C A P S P C R N G G T C R Q		Notch3.pro
221	C P V G Q E G P R C E L R A G P C P P R G C S N G G T C Q T		Notch4.pro
236	T G D - - - V T H E C A C L P G F T G Q N C E E N I D D C P		Notch1.pro
239	T G D - - - F T F E C N C L P G F E G S T C E R N I D D C P		Notch2.pro
215	S G D - - - L T Y D C A C L P G F E G Q N C E V N V D D C P		Notch3.pro
251	M P E K D S T F H L C L C P P G F I G P C C E V N P D N C V		Notch4.pro
263	G N K C K N G G A C V D C V N T Y N C R C P P E W T C Q Y C		Notch1.pro
266	N H R C Q N G G V C V D C V N T Y N C R C P P Q W T C Q F C		Notch2.pro
242	C H R C L N G G T C V D C V N T Y N C Q C P P E W T C Q F C		Notch3.pro
281	S H Q C Q N G G T C Q D G L D T Y T C L C E E T W T G W D C		Notch4.pro
293	T E D V D E C Q L M - P N A C Q N G G T C H N T H G G Y N C		Notch1.pro
296	T E D V D E C L L Q - P N A C Q N G G T C A N R N G G Y G C		Notch2.pro
272	T E D V D E C Q L Q - P N A C H N G G T C F N T L G G H S C		Notch3.pro
311	S E D V D E C E A Q S P P H C R N G G T C Q N S A G S F H C		Notch4.pro

FIGURE 2B

[illegible]

FIGURE 2C

679	I D E C A G N P C E N G G T C E D G I N G F T C R C P E G Y	Notch1.pro
683	I D E C A S N P C R K G A T C I N G V N G F R C I C P E G P	Notch2.pro
658	I N E C A S S P C G E G G S C V D G E N G F R C C P P G S	Notch3.pro
627	L - - C A P N L C Q P K Q I C K D Q X D K A N C L C P D G -	Notch4.pro
709	H D P T C L S E V N E C N S N F C V H G A C R D S L N G Y K	Notch1.pro
713	H H P S C Y S Q V N E C L S N P C I H G N C T G G L S G Y K	Notch2.pro
688	L P P L C L P P S E P C A H E F C S H G I C Y D A P G G F R	Notch3.pro
654	- S P G C A P P F D N C T - - - C H H G H C Q R S S - - -	Notch4.pro
739	C D C D E G W S G T N C - - D I N N N E C E S N P C V N G G	Notch1.pro
743	C L C D A G W V G I N C - - E V D K N E C L S N P C Q N G G	Notch2.pro
718	C V C E P G W S G P R C S Q S I A R D A C E S Q P C R A G G	Notch3.pro
676	C V C D V G W T G P E C - - E A E L G G C I S A P C A H G G	Notch4.pro
767	T C K D M T S G Y V C T C R F G F S G P N C Q T N T N E C A	Notch1.pro
771	T C D N L V N G Y R C T C K K G F K G Y N C Q V N I D E C A	Notch2.pro
748	T C S S D G M G F H C T C P P G V Q G R Q C E L - - - - -	Notch3.pro
704	T C Y P Q P S G Y N C T C P T - - - - - - - - - - -	Notch4.pro
797	S N P C L N Q G T C I D D V A G Y X C N C L L P Y T G A T C	Notch1.pro
801	S N P C L N Q G T C F D D I S G Y T C H C V L P Y T G K N C	Notch2.pro
772	- -	Notch3.pro
719	- -	Notch4.pro
827	E V V L A P C A P S P C R N G G E C R Q S E D Y E S F S C V	Notch1.pro
831	Q T V L A P C S P N P C E N A A V C K E S P N F E S Y T C L	Notch2.pro
772	- - - L S P C T P N P C E H G G R C E S A P G - Q L P V C S	Notch3.pro
719	- -	Notch4.pro
857	C P T G W Q A G Q T C E V D I N E C V L - S P C R H G A S C	Notch1.pro
861	C A P G W Q G - Q R C T I D I D E C I S - K P C M N H G L C	Notch2.pro
798	C P Q G W Q G - P R C Q Q D V D E C A G P A P C G P H G I C	Notch3.pro
719	- -	Notch4.pro
886	Q N T H G G Y R C E C Q A G Y S G R N C E T D I D D C R P K	Notch1.pro
889	H N T Q G S Y M C E C P P G F S G M D C E E D I D D C L A K	Notch2.pro
827	T N L A G S F S C T C H G G Y T G P S C D Q D I N D C D P K	Notch3.pro
719	- -	Notch4.pro
916	P C H N G G S C T D G I N T A F C D C L P G F R G T F C E E	Notch1.pro
919	P C Q N G G S C M D G V N T F S C I C L P G F T G D K C Q T	Notch2.pro
857	P C L N G G S C Q D G V G S F S C S C L P G F A G P R C A R	Notch3.pro
736	P C L N G G S C N P S P G G Y Y C T C P P S H T G P Q C Q T	Notch4.pro
946	D I N E C A S D P C R N G A N C T E C V D S Y T C T C P A G	Notch1.pro
949	D M N E C L S E P C K N G G T C S D Y V N S Y T C K C Q A G	Notch2.pro
887	D V D E C L S N P C G P G - T C T E H V A S F T C T C P P G	Notch3.pro
766	S T D V C V S A P C - - - - - - - - - - - - - - -	Notch4.pro
976	F S G I H C E N N T P D C T F S S C F N G G T C V D G I N S	Notch1.pro
979	F D G V H C E N N I N E C T E S S C F N G G T C V D G I N S	Notch2.pro
916	Y G G F H C E Q D L P D C S P S S C F N G G T C V D G V N S	Notch3.pro
776	- -	Notch4.pro
1006	F T C L C P P G F T G S Y C Q H D V N - E C D S Q P C L E G	Notch1.pro
1009	F S C L C P V G F T G S F C L H E I N - E C S S H P C L N E	Notch2.pro
946	F S C L C R P G Y T G A H C Q H E A D - P C L S R P C L E G	Notch3.pro
788	F S C L C A M G F Q G P R C E G X L R P S C A D S P C R N R	Notch4.pro

FIGURE 2D

1035	G T C Q D G C G S Y R C T C P Q G Y T G P N C Q N L V H W C	Notch1.prc
1038	G T C V D G L G T Y R C S C P L G Y T G K N C Q T L V N L C	Notch2.prc
975	G V C S A A H P G F R C T C L E S F T G P Q C Q T L V D W C	Notch3.prc
818	A T C Q D S F Q G P R C L C P T G Y T G G S C Q T L M D L C	Notch4.prc
1065	D S S P C K N G G K C W Q T H T Q Y R C E C P S G W T G L Y	Notch1.prc
1068	S R S P C K N K G T C V Q K K A E S Q C L C P S G W A G A Y	Notch2.prc
1005	S R Q P C Q N G G R C V Q T G - - A Y C L C P P G W S C R L	Notch3.prc
848	A Q K P C P R N S H C L Q T G P S F H C L C L Q G W T G P L	Notch4.prc
1095	C D V P S V S C E V A A Q R Q G V D V A R L C Q H G G L C V	Notch1.prc
1096	C D V P N V S C D I A A S R R G V L V E H L C Q H S G V C -	Notch2.prc
1033	C D I R S L P C R E A A A Q I G V R L E Q L C Q A G G Q C V	Notch3.prc
878	C N L P L S S C Q K A A L S Q G I D V S S L C H N G G L C V	Notch4.prc
1125	D A G N T H H C R C Q A G Y T G S Y C E D L V D E C S P S P	Notch1.prc
1128	N A G N T H Y C Q C P L G Y T G S Y C E E Q L D E C A S N P	Notch2.prc
1063	D E D S S H Y C V C P E G R T G S H C E Q E V D P C L A Q P	Notch3.prc
908	D S G P S Y F C H C P P G F Q G S L C Q D H V N P C E S R P	Notch4.prc
1155	C Q N G A T C T D Y L G G Y S C K C V A G Y H G V N C S E E	Notch1.prc
1158	C Q H G A T C S D F I G G Y R C E C V P G Y Q G V N C E Y E	Notch2.prc
1093	C Q H G G T C R G Y M G G Y M C E C L P G Y N G D N C E D C	Notch3.prc
938	C Q N G A T C M A Q P S G Y L C Q C A P G Y D G Q N C S X E	Notch4.prc
1185	I D E C L S H P C Q N G G T C L D L P N T Y K C S C P R G T	Notch1.prc
1188	V D E C Q N Q P C Q N G G T C I D L V N H F K C S C P F G T	Notch2.prc
1123	V D E C A S Q P C Q H G G S C I D L V A R Y L C S C P F G T	Notch3.prc
968	L D A C Q S Q P C H N - - - - - - - - - - - - - - - -	Notch4.prc
1215	Q G V H C E I N V D D C N P P V D P V S R S P K C F N N G T	Notch1.prc
1218	R G L L C E E N - D D C - - - - - A R G P H C L N G G Q	Notch2.prc
1153	L G V I C E T N F D D C G P G - P P L D S G P R C L H N G T	Notch3.prc
979	- H G T	Notch4.prc
1245	C V D Q V G G Y S C T C P F G F V G E R C E G D V N E C L S	Notch1.prc
1241	C M D R I G G Y S C R C L P G F A G E R C E G D I N E C L S	Notch2.prc
1182	C V D L V G C F R C T C P P G Y T G L R C E A D I N E C R S	Notch3.prc
982	C T P K P G G F H C A C P F G F V G L R C E G D V D E C L D	Notch4.prc
1275	N P C D A R G T Q N C V Q R V N - D F H C E C R A G H T G R	Notch1.prc
1271	N P C S S E G S L D C I Q L T N - D Y L C V C R S A F T G R	Notch2.prc
1212	G A C H A A H T R D C L Q D P G G G F R C L C H A G F S G P	Notch3.prc
1012	Q P C H P T G T A A C H S I A N - A F Y C Q C L P G H T G Q	Notch4.prc
1304	R C E S V I N G C K G K P C K N G G T C A V A S N T A R G -	Notch1.prc
1300	H C E T F V D V C P Q M P C L N G G T C A V A S N M P D G -	Notch2.prc
1242	R C Q T V L S P C E S Q P C Q H G G Q C R P S P G P G G G L	Notch3.prc
1041	W C E V E I D P C H S Q P C F H G G T C E A T A G S P L G -	Notch4.prc
1333	- F I C K C P A G F E G A T C E N D A R T C G S L R C L N G	Notch1.prc
1329	- F I C R C P P G F S G A R C Q S - - - S C G Q V K C R X G	Notch2.prc
1272	T F T C H C A Q P F W G P R C E R V A R S C R E L Q C P V G	Notch3.prc
1070	- F I C H C F K G F E G P T C S H R A P S C G F H H C H H G	Notch4.prc
1362	G T C I S G F R S - - - P T C L C L G P F T G P E C Q F P A	Notch1.prc
1355	E Q C V H T A S G - - - P R C F C P S P R D - - - C E S G C	Notch2.prc
1302	V P C Q Q T F R G - - - P R C A C P P G L S G P S C R S F P	Notch3.prc
1099	G L C L P S F K P G F P P R C A C L S G Y G G P D C L T P P	Notch4.prc

FIGURE 2E

1389 S S P - - - - C L G G N P C Y N Q G T C E P T S - - E S Notch1.pro
1379 A S - - - - - S P C Q H G G S C H P Q R - - Q P Notch2.pro
1329 G S P P G A S N A S C A A A P C L H G G S C R P A P - - I A Notch3.pro
1129 A P K G - - - - C G P P S P C L Y N G S C S E T T G L G G Notch4.pro

1411 P F Y R C L C P A K F N C L L C H I L D Y S F G C G A G R D Notch1.pro
1396 P Y Y S C Q C A P P F S G S R C E L Y T - - - - - Notch2.pro
1357 P F F R C A C A Q G W T G P R C E A P A - - - - - A Notch3.pro
1154 P G F R C S C P H S S P G P R C Q K P G - - - - - Notch4.pro

1441 I P P P L I E E A C E L P E C Q E D A G N K V C S L Q C N N Notch1.pro
1416 A P P S T P P A T C L S Q Y C A D K A R D G V C D E A C N S Notch2.pro
1378 A P E V S E E P R C P R A A C Q A K R G D Q R C D R E C N S Notch3.pro
1174 - - - A K G - - - - C E G R S G D G A C D A G C S G Notch4.pro

1471 H A C G W D G G D C S L N F N D P W K N C T Q S I Q C W K Y Notch1.pro
1446 H A C Q W D G G D C S L T M E N P W A N C S S P L P C W D Y Notch2.pro
1408 P G C G W D G G D C S L S V G D P W R Q C - E A L Q C W R L Notch3.pro
1193 P G G N W D G G D C S L G V P D P W K G C P S H S R C W L L Notch4.pro

1501 F S D G H C D S Q C N S A G C L F D G F D C Q R A - - E G Q Notch1.pro
1476 I N N - Q C D E L C N T V E C L F D N F E C Q G N - - S K T Notch2.pro
1437 F N N S R C D P A C S S P A C L Y D N F D C H A G G R E R T Notch3.pro
1223 F R D G Q C E P Q C D S E E C L F D G Y D C E T P - - - P A Notch4.pro

1529 C N P L Y D Q Y C K D H F S D G H C D Q G C N S A E C E W D Notch1.pro
1503 C K - - Y D K Y C A D H F K D N H C D Q G C N S E E C G W D Notch2.pro
1467 C N P V Y E K Y C A D H F A D G R C D Q G C N T E E C G W D Notch3.pro
1250 C T P A Y D Q Y C H D H F F N G H C E K G C N T A E C G W D Notch4.pro

1559 G L D C A E L V P E R L A A G T L V V V V L M P P E Q L R N Notch1.pro
1531 G L D C A A D O P E N L A E G T L V I V V L M P P E Q L L Q Notch2.pro
1497 G L D C A S E V P A L L A R G V L V L T V L L P P E E L L R Notch3.pro
1280 G G D C R P E D G D P E W G P S L A L L V V L S P P A L D Q Notch4.pro

1589 S S F H F L R E L S R V I E T N V V F K R D A H G Q Q M I F Notch1.pro
1561 D A R S F L R A L G T L I E T N L R I K R D S Q G E L M V Y Notch2.pro
1527 S S A D F L Q R L S A T I R T S L R F R L D A H G Q A M V F Notch3.pro
1310 Q L F A L A R V L S L T I R V G L W V R K D R D G R D M V Y Notch4.pro

1619 P Y Y G R E E E L R K H P I K R A A E G W A A P D A L L G Q Notch1.pro
1591 P Y Y C E K S A A M K K - - - - - Q - - - - - R Notch2.pro
1557 P Y H R P S P G S E P R - - - - - - - - - - - Notch3.pro
1340 P Y P G A R A E K L G - - - - - - - G T R D P T Y Q E R Notch4.pro

1649 V K A S L L P G G S E G G R R R R E L D P M D V R G S I V Y Notch1.pro
1605 M T R R S L P G S - - - - - - - Q E - - - - - Notch2.pro
1569 A R R E L A P - - - - - - - - - - - E V I G S V V M Notch3.pro
1362 A A P Q T Q P L G - - - - - - - K E T D S L S A G F V V V Notch4.pro

1679 L E I D N R Q C V Q - - A S S Q C F Q S A T D V A A F L G A Notch1.pro
1625 L E I D N R Q C V Q - - D S D H C F K N T D A A A A L I A S Notch2.pro
1584 L E I D N R L C L Q S P E N D H C F P D A Q S A A D Y L G A Notch3.pro
1384 M G V D L S R C G P D H P A S R C P W D P G L L L R E L A A Notch4.pro

1707 L A S L Q S L N - - I P Y K I E A V Q S E T V E P P P P - - Notch1.pro
1653 H A I Q G T L S - - Y P - - - L V S V V S E S L T P - - - Notch2.pro
1614 L S A V E R L D - - F P Y P L S D V R G E P L E P P E - - Notch3.pro
1414 M A A V G A L E P L L P G P L L A V H P H A G T A P P A N Q Notch4.pro

FIGURE 2F

1733	- A Q L H F M Y V A A A A F V L L F F V G C G V L L S R K R	Netch1.prc
1674	- R R T Q L L Y L L A V A V V I T L F I L L G V I M A K R	Netch2.prc
1639	- P S V P L L P L L V A G A V L L L V L L V L G V M V A R R	Netch3.prc
1444	L P W P V L C S P V A G V I L L A L G A L L V L Q L I R R R	Netch4.prc
1762	R R Q H C Q L W F P E G F K V S - E A S S K - - K K R R E P I	Netch1.prc
1703	K R K H G S L W L P E G F T L R R D A S N - - H K R R E P V	Netch2.prc
1668	K R E H S T L W F P E G F S L H K D V A S G H K G R R E P V	Netch3.prc
1474	R R E H G A L W L P P C F T R R P R T Q S A P H R R R P P I	Netch4.prc
1789	G E D S V G L K P L K N - A S D G A L M D D N Q N E - - W G	Netch1.prc
1731	G Q D A V G L K N L S V Q V S E A N L G T G T S E H W V D	Netch2.prc
1698	G Q D A L G M K N M A K - - G E S L M G E V A T - W M D	Netch3.prc
1504	G E D S I G L K A L K P - - - - - K A E V D E D G - V V M	Netch4.prc
1816	D E D L E T K K F R F E E P V V L P D L D D Q T D H R Q W T	Netch1.prc
1761	D E G P Q P K K V K A E D E A L L S E E D D P I D R R P W T	Netch2.prc
1724	T E C P E A K R L K V E E P G M G - - A E E A V D C R Q W T	Netch3.prc
1527	C S G P E - - - - E G E E V G Q A E E G P P S T C Q L W S	Netch4.prc
1846	Q Q H L D A A D L R - M S A M A P T P P Q G E V D A D C M D	Netch1.prc
1791	Q Q H L E A A D I R R T P S L A L T P P Q A E Q E V D V L D	Netch2.prc
1752	Q H H L V A A D I R V A P A M A L T P P Q G D A D A D G M D	Netch3.prc
1553	L S G G C G A L P Q - - - A A M L T P P Q - E S E M E A P D	Netch4.prc
1875	V N V R G P D G F T P L M I A S C S G G G L E T G N S E E -	Netch1.prc
1821	V N V R G P D G C T P L M L A S L R G G S S D L S D E D E D	Netch2.prc
1782	V N V R G P D G F T P L M I A S F C G G A L E P M P T E E D	Netch3.prc
1579	L T T R G P D G V T P L M S A V C C G - E V Q S G T F Q G -	Netch4.prc
1904	- - R E D A P A V I S D F I Y Q G A S L H N Q T D R T G E T	Netch1.prc
1851	- A E D S S A N I I T D L V Y Q G A S L Q A Q T D R T G E M	Netch2.prc
1812	E A L D T S A S I I S D L I C Q G A Q L G A R T D R T G E T	Netch3.prc
1607	- A W L G C P E P W E P L L D G C A C P Q A E T V G T G E T	Netch4.prc
1932	A L H L A A R Y S R S D A A K R L L E A S A D A N I Q D K M	Netch1.prc
1880	A L H L A A R Y S R A L A A K R L L D A G A D A N A Q D K M	Netch2.prc
1842	A L H L A A R Y A R A C A A K R L L D A C A D T N A Q D H S	Netch3.prc
1636	P L H L A A R F S R P T A A R R L L E A G A N P N Q P D R A	Netch4.prc
1962	G R T P L H A A V S A D A Q G V F Q I L I R N R A T D L D A	Netch1.prc
1910	G R C P L H A A V A A D A Q G V F Q I L I R N R V T D L D A	Netch2.prc
1872	G R T P L H T A V T A D A Q G V F Q I L I R N R S T D L D A	Netch3.prc
1666	G R T P L H A A V A A D A R E V C Q L L I R S R Q T A V D A	Netch4.prc
1992	R M H D G T T P L I L A A R L A V E G M I E D L I N S H A D	Netch1.prc
1940	R M N D G T T P L I L A A R L A V E G M V A E L I N C Q A D	Netch2.prc
1902	R M A D G S T A L I L A A R L A V E G M V E E L I A S H A D	Netch3.prc
1696	R T E D G T T P L M L A A R L A V E D L V E E L I A A Q A D	Netch4.prc
2022	V N A V D D L G K S A L H W A A A V N N V D A A V V L L K N	Netch1.prc
1970	V N A V D D H G K S A L H W A A A V N N V E A T L L L L K N	Netch2.prc
1932	V N A V D E L G K S A L H W A A A V N N V E A T L A L L K N	Netch3.prc
1726	V G A R D K W G K T A L H W A A A V N N A R A A R S L L Q A	Netch4.prc
2052	G A N K D M Q N N R F E E T P I F L A A R E G S Y E T A K V I	Netch1.prc
2000	G A N R D M Q N K E E T P I F L A A R E G S Y E A A K I	Netch2.prc
1962	G A N K D M Q S K E E T P L F L A A R E G S Y E A A K L I	Netch3.prc
1756	G A E K D A Q N R F E Q T P I F L A A R E G A V E V A Q I I	Netch4.prc

FIGURE 2G

2082	L D H F A N R D I T D H M D R L P R D I A Q E R M H H D I V	Notch1.pro
2030	L D H F A N R D I T D H M D R L P R D V A R D R M H H D I V	Notch2.pro
1992	L D H F A N R E I T D H L D R L P R D V A Q E R L H Q D I V	Notch3.pro
1786	L G L G A A R E L R D C A G L A P A D V A H Q R N H W D L L	Notch4.pro
2112	R L L D E Y N I V R S P Q L H G A P L G G T P T L S P P L C	Notch1.pro
2060	R L L D E Y N V T P S P - - P G T V L - - T S A L S P V I C	Notch2.pro
2022	R L L D Q P S G P R S P - - P G P H G - - - - L G P L C	Notch3.pro
1816	T L L E G A G P D E A R - - - - - - - - - - - - - - -	Notch4.pro
2142	S P N G Y L G S L K P G V Q G - K K V R K P S S K - - - -	Notch1.pro
2086	G P N R S F L S L K H T P M G - K K S R R P S A K S T M P T	Notch2.pro
2045	P P G A F L P G L K A A Q S G S K K S R R P F G K - - - -	Notch3.pro
1828	- - - - - - - - H K A T P G R E A G P F P R A R - - - -	Notch4.pro
2166	G L A C G S K E A K D L K - A R R K K S Q D G K G C L L D S	Notch1.pro
2115	S L P N L A K E A K D A K G S R R K K S L S E K V Q L S E S	Notch2.pro
2070	- - - - A G L G P Q G P R G R G K K L T L A C P G P L A D S	Notch3.pro
1844	- - - - - - - - - - - - - - - T - - - - - - - - V	Notch4.pro
2195	S G M L S P V L S L E S P H G Y L S D V A S P P L L P S P -	Notch1.pro
2145	S V T L S P V L S L E S P H T Y V S D T T S S P M I T S P G	Notch2.pro
2096	S V T L S P V L S L D S P R P F G G P P A S P G G F P - - -	Notch3.pro
1846	S V S V P P H C G G A I P R C R T L S A G A C P R G G - - -	Notch4.pro
2224	F Q Q S P S V F L N H I P G M P D T H L G I S H L N V A A K	Notch1.pro
2175	I L Q A S P N P M L A T A A P P A P V H A Q H A L S F S N L	Notch2.pro
2123	- - - - - - - - - - - - - - - L E G P Y A A A T A T A	Notch3.pro
1873	- - - - - - - - - - - - - - - G A C L Q A R T W S	Notch4.pro
2254	P E M A A L G G G G R L A F E T G P P R L S H L P V A S G T	Notch1.pro
2205	H E M Q P L A H G A S T V L P S V S Q L L S H H H I V S - -	Notch2.pro
2135	V S L A Q L G G P G R A G L G R Q P P - - - - - - - - -	Notch3.pro
1883	V D L A A R G G G A Y S H C R S L S G - - - - - - - - -	Notch4.pro
2284	S T V L G S S S G G A L N F T V G G S T S L N G Q C E W L S	Notch1.pro
2233	- - - P G S G S A G S L S R - - - - L H P V P V P A D W M N	Notch2.pro
2154	- - - - - G G C V L S L G L - - - - L N P V A V P L D W A R	Notch3.pro
1902	- - - - - V G A C G - - - - - - - - - - - - - - -	Notch4.pro
2314	R L Q S G M V P N Q Y N P L R G S V A P G P L S T Q A P S L	Notch1.pro
2256	R M E V N E T - - Q Y N E M F G M V L A P A E G T H P G - -	Notch2.pro
2175	- - - - - - - - - - - - - - - L P P A P P G P S - - -	Notch3.pro
1908	- - - - - - - - - - - - - - - P T P R G R R - - - -	Notch4.pro
2344	Q H G M V G P L H S S L A A S A L S Q M M S Y Q G L P S T R	Notch1.pro
2282	- - - - - - - - I A - - P - - - - - - - - - Q S R	Notch2.pro
2185	- F L L P	Notch3.pro
1915	- -	Notch4.pro
2374	L A T Q P H L V Q T Q Q V Q F Q N L Q M Q Q Q N L Q P A N I	Notch1.pro
2288	P P E G K H I T T P R E P L P P - I V T F Q - - L I P X - -	Notch2.pro
2189	L A F G P Q L L N F G T P V S P - - - - - - - - -	Notch3.pro
1915	- - - - - - - F S A G M R G P R - - - - - - - - -	Notch4.pro
2404	Q Q Q Q S T Q P P P P P P Q P H I G V S S A A S G H L C R S	Notch1.pro
2313	- - - G S I A Q P A G A P Q P Q S T C P P A V A G P L P T M	Notch2.pro
2205	- - - - Q E R P P P Y I A V P G H G E F Y P V A C - - - -	Notch3.pro
1924	- - - - - - - F N P A I M R G R Y G V A A G R G G - - - -	Notch4.pro

FIGURE 2H

FIGURE 3 Statistics of amino acid alignment for Notch1-4

Percent of identity

Divergence

	1	2	3	4	
1		56.1	52.7	42.6	1
2	64.9		52.7	42.5	2
3	73.0	72.9		43.4	3
4	102.0	102.4	99.3		4
	1	2	3	4	

FIGURE 4AmAb 256A-13 heavy chain variable region sequence

SQVQLQQSGAELAKPGTSVKMACKASGYTFTHWMNWVKQRPGGLEWIGTINPSNDFDCN

CDR-H1

CDR-H2

QKFKDKAILTADKSSSTAYMQLSSLTSEDSAIYYCASGLTARAWFAYWGQGLVTVSAA

CDR-H3

(SEQ ID NO: 2)

FIGURE 4BmAb 256A-13 light chain (kappa) variable region sequence

RATISCRASQSVTTSNYSYMHWFQQKPGQPPKLLIKYASNLD SGV PARFSGSGGTDFTLNI

CDR-L1

CDR-L2

HPVEEEDTATFYCQHSWEIPYTFGGGTNLEIKRADAAPTV (SEQ ID NO: 3)

CDR-L3


FIGURE 5 LIGAND INDEPENDENT NOTCH3 SIGNALING


FIGURE 6 METALLOPROTEASE CLEAVAGE OF NOTCH3

FIGURE 7 NOTCH3/Fc PROTEIN CONSTRUCTS FOR EPITOPE MAPPING




FIGURE 8A

1	A	T	G	G	G	T	C	C	A	G	G	T	G	C	A	A	G	A	G	G	N3-leader Tanox.
1	A	T	G	G	G	G	C	C	G	G	G	G	G	C	C	C	G	T	G	G	N3-leader NCBI.s
21	T	A	G	A	A	G	G	C	G	T	A	G	A	A	G	G	A	G	A	C	N3-leader Tanox.
21	C	C	G	C	C	G	C	C	G	C	C	G	C	C	G	T	C	G	C	C	N3-leader NCBI.s
41	C	A	A	T	G	A	G	C	C	C	A	C	C	T	C	C	T	C	C	G	N3-leader Tanox.
41	C	G	A	T	G	T	C	G	C	C	G	C	C	A	C	C	G	C	C	A	N3-leader NCBI.s
61	C	C	A	C	C	T	C	C	A	G	T	G	A	G	A	G	C	A	C	T	N3-leader Tanox.
61	C	C	G	C	C	A	C	C	C	G	T	G	C	G	G	G	C	G	C	T	N3-leader NCBI.s
81	G	C	C	T	T	G	C	T	G	T	T	G	C	T	G	C	T	G	G		N3-leader Tanox.
81	G	C	C	C	C	T	G	C	T	G	C	T	G	C	T	G	C	T	A	G	N3-leader NCBI.s
101	C	T	G	G	A	C	C	T	G	G	T	G	C	A	G	C	A	G	C	T	N3-leader Tanox.
101	C	G	C	C	G	C	C	G	C	C	G	G	C	T	G	C	A	G	C	C	N3-leader NCBI.s
121	C	C	T	C	C	T	T	G	C	C	T	G	G	A	C						N3-leader Tanox.
121	C	C	C	C	C	T	T	G	C	C	T	G	G	A	C						N3-leader NCBI.s

FIGURE 8B

M	G	P	G	A	R	G	R	R	R	R	R	R	P	M
S	P	P	P	P	P	P	P	V	R	A	L	P	L	L
L	L	L	A	G	P	G	A	A	A	P	P	C	L	D

FIGURE 8C

EPRCPRAACQ AKRGDQRCDR ECNSPGCGWD GGDCSLSVG (SEQ ID NO 10)

FIGURE 8D

AKRGDQRCDR ECNSPGCGWD GGDCSLSVG (SEQ ID NO 11)

FIGURE 9




FIGURE 10 Summary of subdomain swap and amino acid (aa) cluster swap sequence in first LIN12 domain and Mab binding strength in ELISA assays

Expression constructs	Sequence ID	Wild type and swapped sequences of Notch3 1st LIN12 (L1) domain	256A-13	G3	Hu-Fc	% of difference
Notch3-L1	10	EPRCPRAACQAKRGDQRCDCRECNTPGCGWDGGDCSLSVG	+++	-	+++	0
L1-sub1	14	-EACELPE-----	+++	-	+++	15.34
L1-sub2	15	-----EDAGNKVCS-----	+	-	+++	17.95
L1-aa swap1	17	-----EDA-----	+++	-	++	20.51
L1-aa swap2	18	-----NKV-----	+++	-	++	7.69
L1-sub3	16	-----LQCNNHACGWDGGDCSLNFN	+	-	+++	7.69
L1-aa swap3	19	-----SLQ-----	-	-	+++	7.69
L1-aa swap4	20	-----NHA-----	+++	-	+++	7.69
L1-aa swap5	21	-----NFN	+++	-	+++	7.69

In these ELISA experiments, binding of Mab 256A-13 to Notch3-LD/Fc (L1) was used as positive control, i.e. 100% binding. Binding read-out to swap recombinant proteins were compared to that of positive standard. +++ : >50% of standard binding; ++: 10-40% of standard binding; +: 10% to minimum positive signal. -: no binding, i.e. Mab G3 mean binding read-out +/- 3 X standard error. Mab G3 is a human IgG1 control Mab used as negative control.

FIGURE 11 Alanine scanning peptides for epitope mapping of 256A-13

Expression constructs	Sequence ID	Wild type and swapped sequences of Notch3 1st LIN12 (L1) domain
Notch3-L1	10	EPRCPRAACQAKRGDQRCDRECNSPGCGWDGGDCSLSVG
AA-scan 1	22	AACQAAAGDQRC
AA-scan 2	23	ACQAKAADQRC
AA-scan 3	24	CQAKRAAQRCR
AA-scan 4	25	QAKRGAARCDRE
AA-scan 5	26	AKRGDAACDREC
AA-scan 6	27	KRGDAQAADREC
AA-scan 7	28	RGDQRAARECNS
AA-scan 8	29	GDQRCAAECNSP
AA-scan 9	30	DQRCDAACNSPG
AA-scan 10	31	QRCDRAA NSPGC
AA-scan 11	32	RCDREAA SPGCG
AA-scan 12	33	CDREC AAPGCGW
AA-scan 13	34	DRECNAACGCGWD
AA-scan 14	35	RECNSAACGWDG
AA-scan 15	36	ECNSPAA GWDGG
AA-scan 16	37	CNSPGA AWDGGD

1

ANTI-NOTCH3 AGONIST ANTIBODIES AND THEIR USE IN THE TREATMENT OF NOTCH3-RELATED DISEASES

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a divisional application of U.S. application Ser. No. 11/874,682, filed Oct. 18, 2007, now U.S. Pat. No. 7,915,390, which claims the benefit of U.S. Provisional Application No. 60/852,861, filed Oct. 19, 2006, U.S. Provisional Application No. 60/875,597, filed Dec. 18, 2006, and U.S. Provisional Application No. 60/879,218, filed Jan. 6, 2007. The disclosures of U.S. application Ser. No. 11/874,682, U.S. Provisional Application No. 60/852,861, and U.S. Provisional Application No. 60/879,218 are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

The present invention relates to anti-Notch3 agonist antibodies and their use in the amelioration, treatment, or prevention of a Notch3-related disease or disorder.

BACKGROUND OF THE INVENTION

The Notch gene was first described in 1917 when a strain of the fruit fly *Drosophila melanogaster* was found to have notched wing blades (Morgan, *Am Nat* 51:513 (1917)). The gene was cloned almost seventy years later and was determined to be a cell surface receptor playing a key role in the development of many different cell types and tissues in *Drosophila* (Wharton et al., *Cell* 43:567 (1985)). The Notch signaling pathway was soon found to be a signaling mechanism mediated by cell-cell contact and has been evolutionarily conserved from *Drosophila* to human. Notch receptors have been found to be involved in many cellular processes, such as differentiation, cell fate decisions, maintenance of stem cells, cell motility, proliferation, and apoptosis in various cell types during development and tissue homeostasis (See review Artavanis-Tsakonas, et al., *Science* 268:225 (1995)).

Mammals possess four Notch receptor proteins (designated Notch1 to Notch4) and five corresponding ligands (designated Delta Like-1 (DLL-1), Delta Like-3 (DLL-3), Delta Like-4 (DLL-4), Jagged-1 and Jagged-2). The mammalian Notch receptor genes encode ~300 kD proteins that are cleaved during their transport to the cell surface and exist as heterodimers. The extracellular portion of the Notch receptor has thirty-four epidermal growth factor (EGF)-like repeats and three cysteine-rich Notch/LIN12 repeats. The association of two cleaved subunits is mediated by sequences lying immediately N-terminal and C-terminal of the cleavage site, and these two subunits constitute the Notch heterodimerization (HD) domains (Wharton, et al., *Cell* 43:567 (1985); Kidd, et al., *Mol Cell Biol* 6:3431 (1986); Kopczynski, et al., *Genes Dev* 2:1723 (1988); Yochem, et al., *Nature* 335:547 (1988)).

At present, it is still not clear how Notch signaling is regulated by different receptors or how the five ligands differ in their signaling or regulation. The differences in signaling and/or regulation may be controlled by their expression patterns in different tissues or by different environmental cues. It has been documented that Notch ligand proteins, including Jagged/Serrate and Delta/Delta-like, specifically bind to the EGF repeat region and induce receptor-mediated Notch signaling (reviewed by Bray, *Nature Rev Mol Cell Biol*. 7:678

2

(2006), and by Kadesch, *Exp Cell Res*. 260:1 (2000)). Among the EGF repeats, the 10th to 12th repeats are required for ligand binding to the Notch receptor, and the other EGF repeats may enhance receptor-ligand interaction (Xu, et al., *J Biol. Chem.* 280:30158 (2005); Shimizu, et al., *Biochem Biophys Res Comm.* 276:385 (2000)). Although the LIN12 repeats and the dimerization domain are not directly involved in ligand binding, they play important roles in maintaining the heterodimeric protein complex, preventing ligand-independent protease cleavage and receptor activation (Sanchez-Ilizarry, et al., *Mol Cell Biol.* 24:9265 (2004); Vardar et al., *Biochem.* 42:7061 (2003)).

Normal stem cells from many tissues including intestinal and neuronal stem cells depend on Notch signaling for self-renewal and fate determination (Fre, et al., *Nature*, 435: 964 (2005); van Es, et al., *Nature*, 435: 959 (2005); Androutsellis-Theotokis, et al., *Nature*, 442: 823 (2006)). Therefore, the Notch3 agonistic antibody could have application in degenerative diseases. CADASIL (cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy) causes a type of stroke and dementia whose key features include recurrent subcortical ischaemic events and vascular dementia. CADASIL has been found to be associated with a mutant gene localized to chromosome 19 (Joutel, et al., *Nature* 383:707 (1996)). Joutel et al. identified mutations in CADASIL patients that cause serious disruption of the Notch 3 gene, indicating that Notch3 could be the defective protein in CADASIL patients. Unfortunately, this highly incapacitating and often lethal disease has remained largely undiagnosed or misdiagnosed as multiple sclerosis and Alzheimer's disease. Current studies would tend to demonstrate that it is a condition that is much more widespread than first thought.

An additional example of a Notch 3 related disease is familial hemiplegic migraine (FHM), the dominant autosomal form of migraine with aura, located in the same region of chromosome 19 as the Notch3 gene. It should be noted that more than 30% of patients suffering from CADASIL also suffer from migraine with aura. However, the latter is observed in only about 5% of the population and this observation led to the discovery of Notch3 gene involvement in the mechanism of this condition. Similarly, familial paroxysmic ataxia has been linked to a gene located in the same region of chromosome 19 and Notch3 has been implicated in this condition. Other conditions and diseases that have been linked to Notch3 include Alagille syndrome (Flynn, et al., *J Pathol* 204:55 (2004)).

Ongoing research studies are currently being pursued to identify other diseases and conditions linked to Notch3 expression and/or signaling deficiencies. In view of the large number of human diseases associated with the Notch 3 signaling pathway, it is important that new ways of preventing and treating these diseases be identified. The current invention provides novel anti-Notch 3 agonist antibodies useful for this unmet medical need.

SUMMARY OF THE INVENTION

The present invention provides novel agonist antibodies and fragments thereof that specifically bind to an epitope of the human Notch3 receptor in the LIN12 domain. Another aspect of the invention includes the epitope binding site and antibodies that bind this same epitope as the antibodies of the present invention. The antibodies of the present invention activate Notch3-mediated signaling through the Notch3 receptor independent of ligand binding.

The invention includes the amino acid sequences of the variable heavy and light chain of the antibodies and their

corresponding nucleic acid sequences. Another embodiment of the invention includes the CDR sequences of these antibodies.

Another embodiment of the present invention includes the cell lines and vectors harboring the antibody sequences of the present invention.

The present invention also includes the epitope recognized by the agonist antibodies of the invention. The present invention also includes antibodies that bind this epitope. The embodiments include a Notch 3 epitope comprising the Lin 12 domain having at least 80%, 85%, 90%, or 95% sequence identity with SEQ ID NO. 10. More particularly, the Notch 3 epitope comprises SEQ ID NO 11. The present invention includes agonist antibodies that bind this epitope.

Another embodiment of the present invention is the use of these antibodies for the preparation of a medicament or composition for the treatment of Notch 3 related diseases and disorders associated with e.g., receptor inactivation.

Another embodiment of the present invention is the use of these antibodies in the treatment of Notch 3 related diseases or disorders associated with e.g. receptor inactivation comprising the activation of said defects by, e.g., activating Notch 3 signaling independent of ligand binding. Notch 3 related disorders may include, but not limited to, CADASIL, familial hemiplegic migraine (FHM), familial paroxysmic ataxia, Alagille syndrome and other degenerative diseases.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 depicts the amino acid sequence of Notch3. The EGF repeat region extends from amino acid residue 43 to 1383; the LIN12 domain extends from amino acid residue 1384 to 1503; and the dimerization domain extends from amino acid residue 1504 to 1640.

FIG. 2 (A-H) depicts the amino acid sequence comparison between human Notch 1 (SEQ ID NO:39), Notch 2 (SEQ ID NO:40), Notch 3 (SEQ ID NO:1), and Notch 4 (SEQ ID NO:41).

FIG. 3 depicts the percent identity of Notch 1, Notch 2, Notch 3, and Notch 4.

FIGS. 4A and 4B depict the heavy and light chain variable region sequences of anti-Notch3 monoclonal antibody MAb 256A-13 (SEQ ID NO: 2 and SEQ ID NO:3, respectively), with the following CDR regions underlined: CDR-H1 (SEQ ID NO:4), CDR-H2 (SEQ ID NO:5), CDR-H3 (SEQ ID NO:6), CDR-L1 (SEQ ID NO:7), CDR-L2 (SEQ ID NO:8) and CDR-L3 (SEQ ID NO:9).

FIG. 5 depicts a luciferase reporter assay of Example 5 showing activating effects by anti-Notch3 MAbs on the Notch3 receptor.

FIG. 6 depicts the impact of Notch3 agonistic antibodies on metalloprotease cleavage of Notch3.

FIG. 7 depicts Notch3-Fc fusion protein constructs for epitope mapping of the binding site of 256A-13.

FIGS. 8A and 8B depict the comparison of the engineered Notch3 leader peptide coding sequence (SEQ ID NO:42, upper row in FIG. 8A) to the native Notch3 leader peptide coding sequence (NCBI GenBank Accession No. NM_000435) (SEQ ID NO:43, lower row in FIG. 8A) showing the changes of nucleotides (FIG. 8A) and the translated amino acid sequence of the engineered Notch leader peptide sequence (SEQ ID NO:44, FIG. 8B). FIG. 8C depicts the LIN 12 domain (SEQ ID NO:10) and FIG. 8D depicts a subdomain epitope of LIN12 (SEQ ID NO:11).

FIG. 9 depicts the generation of domain swap construct by PCR-SOE method. Arrow bars represent PCR primers. Open bar, Notch3 sequence. Filled bar, Notch1 sequence.

FIG. 10 depicts the amino acid sequences used in the Notch3 LIN12 domain epitope mapping of the MAb 256A-13.

FIG. 11 depicts the Alanine scanning peptides for linear epitope mapping of 256A-13.

DETAILED DESCRIPTION

This invention is not limited to the particular methodology, protocols, cell lines, vectors, or reagents described herein because they may vary. Further, the terminology used herein is for the purpose of describing particular embodiments only and is not intended to limit the scope of the present invention. As used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise, e.g., reference to “a host cell” includes a plurality of such host cells. Unless defined otherwise, all technical and scientific terms and any acronyms used herein have the same meanings as commonly understood by one of ordinary skill in the art in the field of the invention. Although any methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the exemplary methods, devices, and materials are described herein.

All patents and publications mentioned herein are incorporated herein by reference to the extent allowed by law for the purpose of describing and disclosing the proteins, enzymes, vectors, host cells, and methodologies reported therein that might be used with the present invention. However, nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

Terms used throughout this application are to be construed with ordinary and typical meaning to those of ordinary skill in the art. However, Applicants desire that the following terms be given the particular definition as defined below.

The phrase “substantially identical” with respect to an antibody chain polypeptide sequence may be construed as an antibody chain exhibiting at least 70%, or 80%, or 90%, or 95% sequence identity to the reference polypeptide sequence. The term with respect to a nucleic acid sequence may be construed as a sequence of nucleotides exhibiting at least about 85%, or 90%, or 95%, or 97% sequence identity to the reference nucleic acid sequence.

The term “identity” or “homology” shall be construed to mean the percentage of amino acid residues in the candidate sequence that are identical with the residue of a corresponding sequence to which it is compared, after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent identity for the entire sequence, and not considering any conservative substitutions as part of the sequence identity. Neither N- or C-terminal extensions nor insertions shall be construed as reducing identity or homology. Methods and computer programs for the alignment are well known in the art. Sequence identity may be measured using sequence analysis software.

The term “antibody” is used in the broadest sense, and specifically covers monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, and multispecific antibodies (e.g., bispecific antibodies), and antibody fragments so long as they exhibit the desired biological activity. Antibodies (Abs) and immunoglobulins (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific tar-

get, immunoglobulins include both antibodies and other antibody-like molecules which lack target specificity. The antibodies of the invention can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass. Native antibodies and immunoglobulins are usually heterotetrameric glycoproteins of about 150,000 Daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end.

As used herein, "anti-Notch3 antibody" means an antibody which binds specifically to human Notch3 in such a manner so as to activate Notch 3 signaling independent of ligand.

The term "variable" in the context of variable domain of antibodies, refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular target. However, the variability is not evenly distributed through the variable domains of antibodies. It is concentrated in three segments called complementarity determining regions (CDRs; i.e., CDR1, CDR2, and CDR3) also known as hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the target binding site of antibodies (see Kabat, et al. Sequences of Proteins of Immunological Interest, National Institute of Health, Bethesda, Md. (1987)). As used herein, numbering of immunoglobulin amino acid residues is done according to the immunoglobulin amino acid residue numbering system of Kabat, et al., unless otherwise indicated.

The term "antibody fragment" refers to a portion of a full-length antibody, generally the target binding or variable region. Examples of antibody fragments include F(ab), F(ab'), F(ab')₂ and Fv fragments. The phrase "functional fragment or analog" of an antibody is a compound having qualitative biological activity in common with a full-length antibody. For example, a functional fragment or analog of an anti-Notch3 antibody is one which can bind to a Notch3 receptor in such a manner so as to prevent or substantially reduce the ability of the receptor to bind to its ligands or initiate signaling. As used herein, "functional fragment" with respect to antibodies, refers to Fv, F(ab) and F(ab')₂ fragments. An "Fv" fragment consists of a dimer of one heavy and one light chain variable domain in a tight, non-covalent association (V_H - V_L dimer). It is in this configuration that the three CDRs of each variable domain interact to define a target binding site on the surface of the V_H - V_L dimer. Collectively, the six CDRs confer target binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for a target) has the ability to recognize and bind target, although at a lower affinity than the entire binding site.

"Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of an antibody, wherein these domains are present in a single polypeptide chain. Generally, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for target binding.

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy chain variable domain (V_H) connected to a light chain variable domain (V_L) in the same polypeptide chain. By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another changing and create two antigen-binding sites.

The F(ab) fragment contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. F(ab') fragments differ from F(ab) fragments by the addition of a few residues at the carboxyl terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. F(ab') fragments are produced by cleavage of the disulfide bond at the hinge cysteines of the F(ab')₂ pepsin digestion product. Additional chemical couplings of antibody fragments are known to those of ordinary skill in the art.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Pat. No. 4,816,567; and Morrison, et al., *Proc Natl Acad Sci USA* 81:6851 (1984)). Monoclonal antibodies are highly specific, being directed against a single target site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the target. In addition to their specificity, monoclonal antibodies are advantageous in that they may be synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies for use with the present invention may be isolated from phage antibody libraries using the well known techniques. The parent monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler, et al., *Nature* 256:495 (1975), or may be made by recombinant methods.

"Humanized" forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other target-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin template sequence. The humanized anti-

body may also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin template chosen.

The terms "cell," "cell line," and "cell culture" include progeny. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Variant progeny that have the same function or biological property, as screened for in the originally transformed cell, are included. The "host cells" used in the present invention generally are prokaryotic or eukaryotic hosts.

"Transformation" of a cellular organism, cell, or cell line with DNA means introducing DNA into the target cell so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integration. "Transfection" of a cell or organism with DNA refers to the taking up of DNA, e.g., an expression vector, by the cell or organism whether or not any coding sequences are in fact expressed. The terms "transfected host cell" and "transformed" refer to a cell in which DNA was introduced. The cell is termed "host cell" and it may be either prokaryotic or eukaryotic. Typical prokaryotic host cells include various strains of *E. coli*. Typical eukaryotic host cells are mammalian, such as Chinese hamster ovary or cells of human origin. The introduced DNA sequence may be from the same species as the host cell of a different species from the host cell, or it may be a hybrid DNA sequence, containing some foreign and some homologous DNA.

The term "vector" means a DNA construct containing a DNA sequence which is operably linked to a suitable control sequence capable of effecting the expression of the DNA in a suitable host. Such control sequences include a promoter to effect transcription, an optional operator sequence to control such transcription, a sequence encoding suitable mRNA ribosome binding sites, and sequences which control the termination of transcription and translation. The vector may be a plasmid, a phage particle, or simply a potential genomic insert. Once transformed into a suitable host, the vector may replicate and function independently of the host genome, or may in some instances, integrate into the genome itself. In the present specification, "plasmid" and "vector" are sometimes used interchangeably, as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of vectors which serve equivalent function as and which are, or become, known in the art.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including human, domestic and farm animals, nonhuman primates, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, etc.

The word "label" when used herein refers to a detectable compound or composition which can be conjugated directly or indirectly to a molecule or protein, e.g., an antibody. The label may itself be detectable (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

As used herein, "solid phase" means a non-aqueous matrix to which the antibody of the present invention can adhere. Example of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol, and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column).

As used herein, the term "Notch3-mediated disorder" means a condition or disease which is characterized by the defective or underexpressed Notch3 receptor. Specifically it

would be construed to include conditions associated with degenerative diseases such as CADASIL, FHM, familial paroxysmic ataxia, Alagille syndrome, and other degenerative diseases.

Notch 3 Receptor Immunogen for Generating Antibodies

Soluble targets or fragments thereof can be used as immunogens for generating antibodies. The antibody is directed against the target of interest. Preferably, the target is a biologically important polypeptide and administration of the antibody to a mammal suffering from a disease or disorder can result in a therapeutic benefit in that mammal. Whole cells may be used as the immunogen for making antibodies. The immunogen may be produced recombinantly or made using synthetic methods. The immunogen may also be isolated from a natural source.

For transmembrane molecules, such as receptors, fragments of these (e.g., the extracellular domain of a receptor) can be used as the immunogen. Alternatively, cells expressing the transmembrane molecule can be used as the immunogen. Such cells can be derived from a natural source (e.g., cancer cell lines) or may be cells which have been transformed by recombinant techniques to over-express the transmembrane molecule. Other forms of the immunogen useful for preparing antibodies will be apparent to those in the art.

Alternatively, a gene or a cDNA encoding human Notch3 receptor may be cloned into a plasmid or other expression vector and expressed in any of a number of expression systems according to methods well known to those of skill in the art. Methods of cloning and expressing Notch3 receptor and the nucleic acid sequence for human Notch3 receptor are known (see, for example, U.S. Pat. Nos. 5,821,332 and 5,759,546). Because of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding Notch3 receptor protein or polypeptides may be used. One may vary the nucleotide sequence by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence that codes for naturally occurring Notch3 receptor and all such variations may be considered. Any one of these polypeptides may be used in the immunization of an animal to generate antibodies that bind to human Notch3 receptor.

Recombinant Notch3 proteins from other species may also be used as immunogen to generate antibodies because of the high degree of conservation of the amino acid sequence of Notch3. A comparison between human and mouse Notch3 showed that over 90% amino acid sequences are identical between the two species.

The immunogen Notch3 receptor may, when beneficial, be expressed as a fusion protein that has the Notch3 receptor attached to a fusion segment. The fusion segment often aids in protein purification, e.g., by permitting the fusion protein to be isolated and purified by affinity chromatography, but can also be used to increase immunogenicity. Fusion proteins can be produced by culturing a recombinant cell transformed with a fusion nucleic acid sequence that encodes a protein including the fusion segment attached to either the carboxyl and/or amino terminal end of the protein. Fusion segments may include, but are not limited to, immunoglobulin Fc regions, glutathione-S-transferase, β -galactosidase, a poly-histidine segment capable of binding to a divalent metal ion, and maltose binding protein.

Recombinant Notch3 receptor protein as described in Example 1 was used to immunize mice to generate the hybridomas that produce the monoclonal antibodies of the present invention. Exemplary polypeptides comprise all or a portion of SEQ ID NO. 1 or variants thereof.

Antibody Generation

The antibodies of the present invention may be generated by any suitable method known in the art. The antibodies of the present invention may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan (Harlow, et al., *Antibodies: a Laboratory Manual*, Cold spring Harbor Laboratory Press, 2nd ed. (1988)), which is hereby incorporated herein by reference in its entirety).

For example, an immunogen as described in Example 1 may be administered to various host animals including, but not limited to, rabbits, mice, rats, etc., to induce the production of sera containing polyclonal antibodies specific for the antigen. The administration of the immunogen may entail one or more injections of an immunizing agent and, if desired, an adjuvant. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lyssolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Additional examples of adjuvants which may be employed include the MPL-TDM adjuvant (monophosphoryl lipid A, synthetic trehalose dicorynomycolate). Immunization protocols are well known in the art and may be performed by any method that elicits an immune response in the animal host chosen. Adjuvants are also well known in the art.

Typically, the immunogen (with or without adjuvant) is injected into the mammal by multiple subcutaneous or intraperitoneal injections, or intramuscularly or through IV. The immunogen may include a Notch3 polypeptide, a fusion protein, or variants thereof. Depending upon the nature of the polypeptides (i.e., percent hydrophobicity, percent hydrophilicity, stability, net charge, isoelectric point etc.), it may be useful to conjugate the immunogen to a protein known to be immunogenic in the mammal being immunized. Such conjugation includes either chemical conjugation by derivatizing active chemical functional groups to both the immunogen and the immunogenic protein to be conjugated such that a covalent bond is formed, or through fusion-protein based methodology, or other methods known to the skilled artisan. Examples of such immunogenic proteins include, but are not limited to, keyhole limpet hemocyanin, ovalbumin, serum albumin, bovine thyroglobulin, soybean trypsin inhibitor, and promiscuous T helper peptides. Various adjuvants may be used to increase the immunological response as described above.

The antibodies of the present invention comprise monoclonal antibodies. Monoclonal antibodies are antibodies which recognize a single antigenic site. Their uniform specificity makes monoclonal antibodies much more useful than polyclonal antibodies, which usually contain antibodies that recognize a variety of different antigenic sites. Monoclonal antibodies may be prepared using hybridoma technology, such as those described by Kohler, et al., *Nature* 256:495 (1975); U.S. Pat. No. 4,376,110; Harlow, et al., *Antibodies: A Laboratory Manual*, Cold spring Harbor Laboratory Press, 2nd ed. (1988) and Hammerling, et al., *Monoclonal Antibodies and T-Cell Hybridomas*, Elsevier (1981), recombinant DNA methods, or other methods known to the artisan. Other examples of methods which may be employed for producing monoclonal antibodies include, but are not limited to, the human B-cell hybridoma technique (Kosbor, et al., *Immunology Today* 4:72 (1983); Cole, et al., *Proc Natl Acad Sci USA*

80:2026 (1983)), and the EBV-hybridoma technique (Cole, et al., *Monoclonal Antibodies and Cancer Therapy*, pp. 77-96, Alan R. Liss (1985)). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the MAb of this invention may be cultivated in vitro or in vivo.

In the hybridoma model, a host such as a mouse, a humanized mouse, a mouse with a human immune system, hamster, rabbit, camel, or any other appropriate host animal, is immunized to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized in vitro. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, pp. 59-103 (1986)).

Generally, in making antibody-producing hybridomas, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, pp. 59-103 (1986)). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine or human origin. Typically, a rat or mouse myeloma cell line is employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), substances that prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these myeloma cell lines are murine myeloma lines, such as those derived from the MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, Calif. U.S. Application No., and SP2/0 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Md. USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J Immunol* 133:3001 (1984); Brodeur, et al., *Monoclonal Antibody Production Techniques and Applications*, Marcel Dekker, Inc, pp. 51-63 (1987)). The mouse myeloma cell line NSO may also be used (European Collection of Cell Cultures, Salisbury, Wilshire, UK).

The culture medium in which hybridoma cells are grown is assayed for production of monoclonal antibodies directed against Notch3. The binding specificity of monoclonal antibodies produced by hybridoma cells may be determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques are known in the art and within the skill of the artisan. The binding affinity of the monoclonal antibody to Notch3 can, for example, be determined by a Scatchard analysis (Munson, et al., *Anal Biochem* 107:220 (1980)).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the

clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, pp. 59-103 (1986)). Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium (D-MEM) or RPMI-1640 medium. In addition, the hybridoma cells may be grown in vivo as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated or isolated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-SEPHAROSE® affinity media, hydroxylapatite chromatography, gel exclusion chromatography, gel electrophoresis, dialysis, or affinity chromatography.

A variety of methods exist in the art for the production of monoclonal antibodies and thus, the invention is not limited to their sole production in hybridomas. For example, the monoclonal antibodies may be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567. In this context, the term "monoclonal antibody" refers to an antibody derived from a single eukaryotic, phage, or prokaryotic clone. DNA encoding the monoclonal antibodies of the invention is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies, or such chains from human, humanized, or other sources) (Innis, et al. In PCR Protocols. A Guide to Methods and Applications, Academic (1990), Sanger, et al., *Proc Natl Acad Sci* 74:5463 (1977)). The hybridoma cells serve as a source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, NS0 cells, Simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Pat. No. 4,816,567; Morrison, et al., *Proc Natl Acad Sci USA* 81:6851 (1984)) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain cross-linking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent cross-linking.

Antibody fragments which recognize specific epitopes may be generated by known techniques. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto, et al., *J Biochem Biophys Methods* 24:107 (1992); Brennan, et al., *Science* 229:81 (1985)). For example, Fab and F(ab')₂ fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain.

However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from an antibody phage library. Alternatively, F(ab')₂-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter, et al., *Bio/Technology* 10:163 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (Fv) (PCT patent application WO 93/16185).

For some uses, including in vivo use of antibodies in humans and in vitro detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, *Science* 229:1202 (1985); Oi, et al., *BioTechniques* 4:214 (1986); Gillies, et al., *J Immunol Methods* 125:191 (1989); U.S. Pat. Nos. 5,807, 715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entirety.

A humanized antibody is designed to have greater homology to a human immunoglobulin than animal-derived monoclonal antibodies. Humanization is a technique for making a chimeric antibody wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. Humanized antibodies are antibody molecules generated in a non-human species that bind the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and framework (FR) regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. See, e.g., U.S. Pat. No. 5,585,089; Riechmann, et al., *Nature* 332:323 (1988), which are incorporated herein by reference in their entirety. Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Pat. Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592, 106; EP 519,596; Padlan, *Molecular Immunology* 28:489 (1991); Studnicka, et al., *Protein Engineering* 7:805 (1994); Roguska, et al., *Proc Natl Acad Sci USA* 91:969 (1994)), and chain shuffling (U.S. Pat. No. 5,565,332).

Generally, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the methods of Winter and co-workers (Jones, et al., *Nature* 321:522 (1986); Riechmann, et al., *Nature* 332:323 (1988); Verhoeven, et al., *Science* 239:1534 (1988)), by substituting non-human CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been

substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possible some FR residues are substituted from analogous sites in rodent antibodies.

It is further important that humanized antibodies retain higher affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of certain residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin sequences, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is maximized, although it is the CDR residues that directly and most substantially influence antigen binding.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a non-human antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of that of the non-human parent antibody is then accepted as the human FR for the humanized antibody (Sims, et al., *J Immunol* 151:2296 (1993); Chothia, et al., *J Mol Biol* 196:901 (1987)). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter, et al., *Proc Natl Acad Sci USA* 89:4285 (1992); Presta, et al., *J Immunol* 151:2623 (1993)).

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also, U.S. Pat. Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety. The techniques of Cole, et al. and Boerder, et al. are also available for the preparation of human monoclonal antibodies (Cole, et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Riss (1985); and Boerner, et al., *J Immunol* 147:86 (1991)).

Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and diversity region may be introduced into mouse embryonic stem cells in addition to the

human heavy and light chain genes. The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce chimeric mice. The chimeric mice are then bred to produce homozygous offspring which express human antibodies. See, e.g., Jakobovitis, et al., *Proc Natl Acad Sci USA* 90:2551 (1993); Jakobovitis, et al., *Nature* 362:255 (1993); Bruggermann, et al., *Year in Immunol* 7:33 (1993); Duchosal, et al., *Nature* 355:258 (1992)). The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg, et al., *Int Rev Immunol* 13:65-93 (1995). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Pat. Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, Calif.), Genpharm (San Jose, Calif.), and Medarex, Inc. (Princeton, N.J.) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

Also human MAbs could be made by immunizing mice transplanted with human peripheral blood leukocytes, splenocytes or bone marrows (e.g., Trioma techniques of XTL). Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope (Jespers, et al., *Bio/technology* 12:899 (1988)).

Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using techniques well known to those skilled in the art (See, e.g., Greenspan, et al., *FASEB J* 7:437 (1989); Nissinoff, *J Immunol* 147:2429 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand. For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligands/receptors, and thereby block its biological activity.

The antibodies of the present invention may be bispecific antibodies. Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present inven-

tion, one of the binding specificities may be directed towards Notch3, the other may be for any other antigen, and preferably for a cell-surface protein, receptor, receptor subunit, tissue-specific antigen, virally derived protein, virally encoded envelope protein, bacterially derived protein, or bacterial surface protein, etc.

Methods for making bispecific antibodies are well known. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein, et al., *Nature* 305:537 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829 and in Traunecker, et al., *EMBO J.* 10:3655 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It may have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transformed into a suitable host organism. For further details of generating bispecific antibodies see, for example Suresh, et al., *Meth In Enzym* 121:210 (1986).

Heteroconjugate antibodies are also contemplated by the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Pat. No. 4,676,980). It is contemplated that the antibodies may be prepared in vitro using known methods in synthetic protein chemistry, including those involving cross-linking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioester bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimide and those disclosed, for example, in U.S. Pat. No. 4,676,980.

In addition, one can generate single-domain antibodies to Notch3. Examples of this technology have been described in WO9425591 for antibodies derived from Camelidae heavy chain Ig, as well in US20030130496 describing the isolation of single domain fully human antibodies from phage libraries.

One can also create a single peptide chain binding molecules in which the heavy and light chain Fv regions are connected. Single chain antibodies ("scFv") and the method of their construction are described in U.S. Pat. No. 4,946,778. Alternatively, Fab can be constructed and expressed by similar means. All of the wholly and partially human antibodies are less immunogenic than wholly murine MAbs, and the fragments and single chain antibodies are also less immunogenic.

Antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty, et al., *Nature* 348:552 (1990). Clarkson, et al., *Nature* 352:624 (1991) and Marks, et al., *J Mol Biol* 222:581 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity

(nM range) human antibodies by chain shuffling (Marks, et al., *Bio/Technology* 10:779 (1992)), as well as combinatorial infection and in vivo recombination as a strategy for constructing very large phage libraries (Waterhouse, et al., *Nuc Acids Res* 21:2265 (1993)). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences (U.S. Pat. No. 4,816,567; Morrison, et al., *Proc Natl Acad Sci USA* 81:6851 (1984)).

Another alternative is to use electrical fusion rather than chemical fusion to form hybridomas. This technique is well established. Instead of fusion, one can also transform a B cell to make it immortal using, for example, an Epstein Barr Virus, or a transforming gene. See, e.g., "Continuously Proliferating Human Cell Lines Synthesizing Antibody of Predetermined Specificity," Zurawaki, et al., in *Monoclonal Antibodies*, ed. by Kennett, et al., Plenum Press, pp. 19-33. (1980)). Anti-Notch3 MAbs can be raised by immunizing rodents (e.g., mice, rats, hamsters, and guinea pigs) with Notch3 protein, fusion protein, or its fragments expressed by either eukaryotic or prokaryotic systems. Other animals can be used for immunization, e.g., non-human primates, transgenic mice expression immunoglobulins, and severe combined immunodeficient (SCID) mice transplanted with human B lymphocytes. Hybridomas can be generated by conventional procedures by fusing B lymphocytes from the immunized animals with myeloma cells (e.g., Sp2/0 and NSO), as described earlier (Köhler, et al., *Nature* 256:495 (1975)). In addition, anti-Notch3 antibodies can be generated by screening of recombinant single-chain Fv or Fab libraries from human B lymphocytes in phage-display systems. The specificity of the MAbs to Notch3 can be tested by ELISA, Western immunoblotting, or other immunochemical techniques. The inhibitory activity of the antibodies on complement activation can be assessed by hemolytic assays, using sensitized chicken or sheep RBCs for the classical complement pathway. The hybridomas in the positive wells are cloned by limiting dilution. The antibodies are purified for characterization for specificity to human Notch3 by the assays described above. Identification of Anti-Notch-3 Antibodies

The present invention provides agonist monoclonal antibodies that activate Notch3-mediated signaling independent of ligand. In particular, the antibodies of the present invention bind to and activate Notch3. The antibodies of the present invention include the antibody designated 256A-13. The present invention also includes antibodies that bind to the same epitope as 256A-13.

Candidate anti-Notch3 antibodies were tested by enzyme linked immunosorbent assay (ELISA), Western immunoblotting, or other immunochemical techniques. Assays performed to characterize the individual antibodies are described in the Examples.

Antibodies of the invention include, but are not limited to, polyclonal, monoclonal, monovalent, bispecific, heteroconjugate, multispecific, human, humanized or chimeric antibodies, single chain antibodies, single-domain antibodies, Fab fragments, F(ab') fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above.

The antibodies may be human antigen-binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and

single-domain antibodies comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding fragments comprising any combination of variable region(s) with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be from any animal origin including birds and mammals. Preferably, the antibodies are from human, non-human primates, rodents (e.g., mouse and rat), donkey, sheep, rabbit, goat, guinea pig, camel, horse, or chicken.

As used herein, "human" antibodies" include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described infra and, for example in, U.S. Pat. No. 5,939,598 by Kuchlerlapati, et al.

The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of Notch3 or may be specific for both Notch3 as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., *J Immunol* 147:60 (1991); U.S. Pat. Nos. 4,474, 893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny, et al., *J Immunol* 148:1547 (1992).

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of Notch3 which they recognize or specifically bind. The epitope(s) or polypeptide portion(s) may be specified as described herein, e.g., by N-terminal and C-terminal positions, by size in contiguous amino acid residues, or listed in the Tables and Figures.

Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that bind Notch3 polypeptides, which have at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described herein) to Notch3 are also included in the present invention. Anti-Notch3 antibodies may also bind with a K_D of less than about 10^{-7} M, less than about 10^{-6} M, or less than about 10^{-5} M to other proteins, such as anti-Notch3 antibodies from species other than that against which the anti-Notch3 antibody is directed.

In specific embodiments, antibodies of the present invention cross-react with monkey homologues of human Notch3 and the corresponding epitopes thereof. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of the specific antigenic and/or immunogenic polypeptides disclosed herein.

Further included in the present invention are antibodies which bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide encoding Notch3 under stringent hybridization conditions. Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide of the invention. Preferred binding affinities include those with an equilibrium dissociation constant or K_D from 10^{-8} to 10^{-15} M, 10^{-8} to 10^{-12} M, 10^{-8} to 10^{-10} M, or 10^{-10} to 10^{-12} M. The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for determining competitive binding,

for example, the immunoassays described herein. In preferred embodiments, the antibody competitively inhibits binding to the epitope by at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50%.

Vectors and Host Cells

In another aspect, the present invention provides isolated nucleic acid sequences encoding an antibody variant as disclosed herein, vector constructs comprising a nucleotide sequence encoding the antibodies of the present invention, host cells comprising such a vector, and recombinant techniques for the production of the antibody.

For recombinant production of the antibody variant, the nucleic acid encoding it is isolated and inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. DNA encoding the antibody variant is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody variant). Standard techniques for cloning and transformation may be used in the preparation of cell lines expressing the antibodies of the present invention.

Vectors

Many vectors are available. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Recombinant expression vectors containing a nucleotide sequence encoding the antibodies of the present invention can be prepared using well known techniques. The expression vectors include a nucleotide sequence operably linked to suitable transcriptional or translational regulatory nucleotide sequences such as those derived from mammalian, microbial, viral, or insect genes. Examples of regulatory sequences include transcriptional promoters, operators, enhancers, mRNA ribosomal binding sites, and/or other appropriate sequences which control transcription and translation initiation and termination. Nucleotide sequences are "operably linked" when the regulatory sequence functionally relates to the nucleotide sequence for the appropriate polypeptide. Thus, a promoter nucleotide sequence is operably linked to, e.g., the antibody heavy chain sequence if the promoter nucleotide sequence controls the transcription of the appropriate nucleotide sequence.

In addition, sequences encoding appropriate signal peptides that are not naturally associated with antibody heavy and/or light chain sequences can be incorporated into expression vectors. For example, a nucleotide sequence for a signal peptide (secretory leader) may be fused in-frame to the polypeptide sequence so that the antibody is secreted to the periplasmic space or into the medium. A signal peptide that is functional in the intended host cells enhances extracellular secretion of the appropriate antibody. The signal peptide may be cleaved from the polypeptide upon secretion of antibody from the cell. Examples of such secretory signals are well known and include, e.g., those described in U.S. Pat. Nos. 5,698,435; 5,698,417; and 6,204,023.

The vector may be a plasmid vector, a single or double-stranded phage vector, or a single or double-stranded RNA or DNA viral vector. Such vectors may be introduced into cells as polynucleotides by well known techniques for introducing DNA and RNA into cells. The vectors, in the case of phage and viral vectors also may be introduced into cells as packaged or encapsulated virus by well known techniques for infection and transduction. Viral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host

cells. Cell-free translation systems may also be employed to produce the protein using RNAs derived from the present DNA constructs. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publications WO 86/05807 and WO 89/01036; and U.S. Pat. No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain.

Host Cells

The antibodies of the present invention can be expressed from any suitable host cell. Examples of host cells useful in the present invention include prokaryotic, yeast, or higher eukaryotic cells and include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., Baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

Prokaryotes useful as host cells in the present invention include gram negative or gram positive organisms such as *E. coli*, *B. subtilis*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, *Serratia*, and *Shigella*, as well as *Bacilli*, *Pseudomonas*, and *Streptomyces*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting.

Expression vectors for use in prokaryotic host cells generally comprise one or more phenotypic selectable marker genes. A phenotypic selectable marker gene is, for example, a gene encoding a protein that confers antibiotic resistance or that supplies an autotrophic requirement. Examples of useful expression vectors for prokaryotic host cells include those derived from commercially available plasmids such as the pKK223-3 vector (Pharmacia Fine Chemicals, Uppsala, Sweden), PGEM®1 vector (Promega Biotec, Madison, Wis., USA), and the pET (Novagen, Madison, Wis., USA) and pRSET (Invitrogen, Carlsbad, Calif.) series of vectors (Studier, *J Mol Biol* 219:37 (1991); Schoepfer, *Gene* 124:83 (1993)). Promoter sequences commonly used for recombinant prokaryotic host cell expression vectors include T7, (Rosenberg, et al., *Gene* 56:125 (1987)), β -lactamase (penicillinase), lactose promoter system (Chang, et al., *Nature* 275:615 (1978); Goeddel, et al., *Nature* 281:544 (1979)), tryptophan (trp) promoter system (Goeddel, et al., *Nucl Acids Res* 8:4057 (1980)), and tac promoter (Sambrook, et al., *Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory (1990)).

Yeasts or filamentous fungi useful in the present invention include those from the genus *Saccharomyces*, *Pichia*, *Actinomyces*, *Kluyveromyces*, *Schizosaccharomyces*, *Candida*, *Trichoderma*, *Neurospora*, and filamentous fungi such as *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus*. Yeast vectors will often contain an origin of replication

sequence from a 2 μ yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Suitable promoter sequences for yeast vectors include, among others, promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman, et al., *J Biol Chem* 255:2073 (1980)) or other glycolytic enzymes (Holland, et al., *Biochem* 17:4900 (1978)) such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Other suitable vectors and promoters for use in yeast expression are further described in Fleer, et al., *Gene* 107:285 (1991). Other suitable promoters and vectors for yeast and yeast transformation protocols are well known in the art. Yeast transformation protocols are well known. One such protocol is described by Hinnen, et al., *Proc Natl Acad Sci* 75:1929 (1978). The Hinnen protocol selects for Trp⁺ transformants in a selective medium.

Mammalian or insect host cell culture systems may also be employed to express recombinant antibodies. In principle, any higher eukaryotic cell culture is workable, whether from vertebrate or invertebrate culture. Examples of invertebrate cells include plant and insect cells (Luckow, et al., *Bio/Technology* 6:47 (1988); Miller, et al., *Genetics Engineering, Setlow, et al., eds. Vol. 8, pp. 277-9, Plenum Publishing* (1986); Mseda, et al., *Nature* 315:592 (1985)). For example, Baculovirus systems may be used for production of heterologous proteins. In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) may be used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Other hosts that have been identified include *Aedes*, *Drosophila melanogaster*, and *Bombyx mori*. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of AcNPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells. Moreover, plant cells cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco and also be utilized as hosts.

Vertebrate cells, and propagation of vertebrate cells, in culture (tissue culture) has become a routine procedure. See Tissue Culture, Kruse, et al., eds., Academic Press (1973). Examples of useful mammalian host cell lines are monkey kidney; human embryonic kidney line; baby hamster kidney cells; Chinese hamster ovary cells/-DHFR (CHO, Urlaub, et al., *Proc Natl Acad Sci USA* 77:4216 (1980)); mouse sertoli cells; human cervical carcinoma cells (HELA); canine kidney cells; human lung cells; human liver cells; mouse mammary tumor; and NSO cells.

Host cells are transformed with the above-described vectors for antibody production and cultured in conventional nutrient media modified as appropriate for inducing promoters, transcriptional and translational control sequences, selecting transformants, or amplifying the genes encoding the desired sequences. Commonly used promoter sequences and enhancer sequences are derived from polyoma virus, Adenovirus 2, Simian virus 40 (SV40), and human cytomegalovirus (CMV). DNA sequences derived from the SV40 viral genome may be used to provide other genetic elements for expression of a structural gene sequence in a mammalian host cell, e.g., SV40 origin, early and late promoter, enhancer,

splice, and polyadenylation sites. Viral early and late promoters are particularly useful because both are easily obtained from a viral genome as a fragment which may also contain a viral origin of replication. Exemplary expression vectors for use in mammalian host cells are commercially available.

The host cells used to produce the antibody variant of this invention may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma, St Louis, Mo.), Minimal Essential Medium (MEM, Sigma, St Louis, Mo.), RPMI-1640 (Sigma, St Louis, Mo.), and Dulbecco's Modified Eagle's Medium (DMEM, Sigma, St Louis, Mo.) are suitable for culturing host cells. In addition, any of the media described in Ham, et al., *Meth Enzymol* 58:44 (1979), Barnes, et al., *Anal Biochem* 102:255 (1980), and U.S. Pat. Nos. 4,767,704; 4,657,866; 4,560,655; 5,122,469; 5,712,163; or 6,048,728 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as X-chlorides, where X is sodium, calcium, magnesium; and phosphates), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as gentamicin drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

Polynucleotides Encoding Antibodies

The invention further provides polynucleotides or nucleic acids, e.g., DNA, comprising a nucleotide sequence encoding an antibody of the invention and fragments thereof. Exemplary polynucleotides include those encoding antibody chains comprising one or more of the amino acid sequences described herein. The invention also encompasses polynucleotides that hybridize under stringent or lower stringency hybridization conditions to polynucleotides that encode an antibody of the present invention.

The polynucleotides may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier, et al., *Bio/Techniques* 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.

Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A⁺ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated

by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

Once the nucleotide sequence and corresponding amino acid sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook, et al., *Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory (1990); Ausubel, et al., eds., *Current Protocols in Molecular Biology*, John Wiley & Sons (1998), which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

In a specific embodiment, the amino acid sequence of the heavy and/or light chain variable domains may be inspected to identify the sequences of the CDRs by well known methods, e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the regions of sequence hypervariability. Using routine recombinant DNA techniques, one or more of the CDRs may be inserted within framework regions, e.g., into human framework regions to humanize a non-human antibody, as described supra. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework regions (see, e.g., Chothia, et al., *J Mol Biol* 278: 457 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a polypeptide of the invention. Preferably, as discussed supra, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison, et al., *Proc Natl Acad Sci* 81:851 (1984); Neuberger, et al., *Nature* 312:604 (1984); Takeda, et al., *Nature* 314:452 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described supra, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine MAb and a human immunoglobulin constant region, e.g., humanized antibodies.

Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778; Bird, *Science* 242:423 (1988); Huston, et al., *Proc Natl Acad Sci USA* 85:5879 (1988); and Ward, et al., *Nature* 334:544 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in *E. coli* may also be used (Skerra, et al., *Science* 242:1038 (1988)).

Methods of Producing Anti-Notch3 Antibodies

The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques.

Recombinant expression of an antibody of the invention, or fragment, derivative, or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody or a fragment of the antibody. Once a polynucleotide encoding an antibody molecule has been obtained, the vector for the production of the antibody may be produced by recombinant DNA technology. An expression vector is constructed containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. In one aspect of the invention, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention as described above. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. Bacterial cells such as *E. coli*, and eukaryotic cells are commonly used for the expression of a recombinant antibody molecule, especially for the expression of whole recombinant antibody molecule. For example, mammalian cells such as CHO, in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus, are an effective expression system for antibodies (Foecking, et al., *Gene* 45:101 (1986); Cockett, et al., *Bio/Technology* 8:2 (1990)).

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include, but are not limited to, CHO, COS, 293, 3T3, or myeloma cells.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for one to two days in an enriched media, and then are switched to a selective media.

The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., *Cell* 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska, et al., *Proc Natl Acad Sci USA* 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy, et al., *Cell* 22:817 (1980)) genes can be employed in tk, hgprt or aprt-cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., *Proc Natl Acad Sci USA* 77:357 (1980); O'Hare, et al., *Proc Natl Acad Sci USA* 78:1527 (1981)); gpt, which confers resistance to mycophenolic acid (Mulligan, et al., *Proc Natl Acad Sci USA* 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 (Wu, et al., *Biotherapy* 3:87 (1991)); and hygromycin (Santerre, et al., *Gene* 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel, et al., eds., *Current Protocols in Molecular Biology*, John Wiley & Sons (1993); Kriegl, Gene Transfer and Expression, A Laboratory Manual, Stockton Press (1990); and in Chapters 12 and 13, Dracopoli, et al., eds., *Current Protocols in Human Genetics*, John Wiley & Sons (1994); Colberre-Garapin, et al., *J Mol Biol* 150:1 (1981), which are incorporated by reference herein in their entirety.

The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington, et al., "The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells," *DNA Cloning*, Vol. 3. Academic Press (1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse, et al., *Mol Cell Biol* 3:257 (1983)).

The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, *Nature* 322:52 (1986); Kohler, *Proc Natl Acad Sci USA* 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once an antibody molecule of the invention has been produced by an animal, chemically synthesized, or recombinantly expressed, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and size-exclusion chromatography), centrifugation, differ-

ential solubility, or by any other standard technique for the purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art, to facilitate purification.

The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide. Fused or conjugated antibodies of the present invention may be used for ease in purification. See e.g., PCT publication WO 93/21232; EP 439,095; Naramura, et al., *Immunol Lett* 39:91 (1994); U.S. Pat. No. 5,474,981; Gillies, et al., *Proc Natl Acad Sci USA* 89:1428 (1992); Fell, et al., *J Immunol* 146:2446 (1991), which are incorporated by reference in their entireties.

Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide (SEQ ID NO:38), such as the tag provided in a pQE vector (QIAGEN, Inc., Chatsworth, Calif.), among others, many of which are commercially available. As described in Gentz, et al., *Proc Natl Acad Sci USA* 86:821 (1989), for instance, hexa-histidine (SEQ ID NO:38) provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, et al., *Cell* 37:767 (1984)) and the "flag" tag.

Antibody Purification

When using recombinant techniques, the antibody variant can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the antibody variant is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, may be removed, for example, by centrifugation or ultrafiltration. Carter, et al., *Bio/Technology* 10:163 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 minutes. Cell debris can be removed by centrifugation. Where the antibody variant is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody variant. Protein A can be used to purify antibodies that are based on human IgG1, IgG2 or IgG4 heavy chains (Lindmark, et al., *J Immunol Meth* 62:1 (1983)). Protein G is recommended for all mouse isotypes and for human IgG3 (Guss, et al., *EMBO J.* 5:1567 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody variant comprises a CH3 domain, the Bakerbond ABX™ resin (J. T. Baker; Phillips-

burg, N.J.) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE® chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody variant to be recovered.

Following any preliminary purification step(s), the mixture comprising the antibody variant of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (e.g., from about 0-0.25 M salt).

Pharmaceutical Formulation

Therapeutic formulations of the polypeptide or antibody may be prepared for storage as lyophilized formulations or aqueous solutions by mixing the polypeptide having the desired degree of purity with optional "pharmaceutically-acceptable" carriers, excipients or stabilizers typically employed in the art (all of which are termed "excipients"), i.e., buffering agents, stabilizing agents, preservatives, isotonicifiers, non-ionic detergents, antioxidants, and other miscellaneous additives. See Remington's Pharmaceutical Sciences, 16th edition, Osol, Ed. (1980). Such additives must be nontoxic to the recipients at the dosages and concentrations employed.

Buffering agents help to maintain the pH in the range which approximates physiological conditions. They are preferably present at concentration ranging from about 2 mM to about 50 mM. Suitable buffering agents for use with the present invention include both organic and inorganic acids and salts thereof such as citrate buffers (e.g., monosodium citrate-disodium citrate mixture, citric acid-trisodium citrate mixture, citric acid-monosodium citrate mixture, etc.), succinate buffers (e.g., succinic acid-monosodium succinate mixture, succinic acid-sodium hydroxide mixture, succinic acid-disodium succinate mixture, etc.), tartrate buffers (e.g., tartaric acid-sodium tartrate mixture, tartaric acid-potassium tartrate mixture, tartaric acid-sodium hydroxide mixture, etc.), fumarate buffers (e.g., fumaric acid-monosodium fumarate mixture, etc.), fumarate buffers (e.g., fumaric acid-monosodium fumarate mixture, fumaric acid-disodium fumarate mixture, monosodium fumarate-disodium fumarate mixture, etc.), gluconate buffers (e.g., gluconic acid-sodium glyconate mixture, gluconic acid-sodium hydroxide mixture, gluconic acid-potassium glyconate mixture, etc.), oxalate buffer (e.g., oxalic acid-sodium oxalate mixture, oxalic acid-sodium hydroxide mixture, oxalic acid-potassium oxalate mixture, etc.), lactate buffers (e.g., lactic acid-sodium lactate mixture, lactic acid-sodium hydroxide mixture, lactic acid-potassium lactate mixture, etc.) and acetate buffers (e.g., acetic acid-sodium acetate mixture, acetic acid-sodium hydroxide mixture, etc.). Additionally, there may be mentioned phosphate buffers, histidine buffers and trimethylamine salts such as Tris.

Preservatives may be added to retard microbial growth, and may be added in amounts ranging from 0.2%-1% (w/v). Suitable preservatives for use with the present invention include phenol, benzyl alcohol, meta-cresol, methyl paraben, propyl paraben, octadecyldimethylbenzyl ammonium chloride, benzalconium halides (e.g., chloride, bromide, iodide), hexamethonium chloride, and alkyl parabens such as methyl or propyl paraben, catechol, resorcinol, cyclohexanol, and 3-pentanol.

Isotonicifiers sometimes known as "stabilizers" may be added to ensure isotonicity of liquid compositions of the present invention and include polyhydric sugar alcohols, preferably trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol and mannitol.

Stabilizers refer to a broad category of excipients which can range in function from a bulking agent to an additive which solubilizes the therapeutic agent or helps to prevent denaturation or adherence to the container wall. Typical stabilizers can be polyhydric sugar alcohols (enumerated above); amino acids such as arginine, lysine, glycine, glutamine, asparagine, histidine, alanine, ornithine, L-leucine, 2-phenylalanine, glutamic acid, threonine, etc., organic sugars or sugar alcohols, such as lactose, trehalose, stachyose, mannitol, sorbitol, xylitol, ribitol, myoinositol, galactitol, glycerol and the like, including cyclitols such as inositol; polyethylene glycol; amino acid polymers; sulfur containing reducing agents, such as urea, glutathione, thiocetic acid, sodium thioglycolate, thioglycerol, alpha.-monothioglycerol and sodium thio sulfate; low molecular weight polypeptides (i.e. <10 residues); proteins such as human serum albumin, bovine serum albumin, gelatin or immunoglobulins; hydrophylic polymers, such as polyvinylpyrrolidone monosaccharides, such as xylose, mannose, fructose, glucose; disaccharides such as lactose, maltose, sucrose and trisaccharides such as raffinose; and polysaccharides such as dextran. Stabilizers may be present in the range from 0.1 to 10,000 weights per part of weight active protein.

Non-ionic surfactants or detergents (also known as "wetting agents") may be added to help solubilize the therapeutic agent as well as to protect the therapeutic protein against agitation-induced aggregation, which also permits the formulation to be exposed to shear surface stressed without causing denaturation of the protein. Suitable non-ionic surfactants include polysorbates (20, 80, etc.), polyoxamers (184, 188 etc.), PLURONIC® polyols, polyoxyethylene sorbitan monoethers (TWEEN-20®, TWEEN-80®, etc.). Non-ionic surfactants may be present in a range of about 0.05 mg/ml to about 1.0 mg/ml, preferably about 0.07 mg/ml to about 0.2 mg/ml.

Additional miscellaneous excipients include bulking agents, (e.g., starch), chelating agents (e.g., EDTA), antioxidants (e.g., ascorbic acid, methionine, vitamin E), and cosolvents. The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide an immunosuppressive agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended. The active ingredients may also be entrapped in microcapsule prepared, for example, by coascervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsule and poly-(methylmethacrylate) microcapsule, respectively, in colloidal drug delivery systems (for example, liposomes, albumin micropheeres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Osal, Ed. (1980).

The formulations to be used for in vivo administration must be sterile. This is readily accomplished, for example, by filtration through sterile filtration membranes. Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the antibody variant, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-

release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C. resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S—S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

The amount of therapeutic polypeptide, antibody, or fragment thereof which will be effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. Where possible, it is desirable to determine the dose-response curve and the pharmaceutical compositions of the invention first in vitro, and then in useful animal model systems prior to testing in humans.

In a preferred embodiment, an aqueous solution of therapeutic polypeptide, antibody or fragment thereof is administered by subcutaneous injection. Each dose may range from about 0.5 µg to about 50 µg per kilogram of body weight, or more preferably, from about 3 µg to about 30 µg per kilogram body weight.

The dosing schedule for subcutaneous administration may vary from once a month to daily depending on a number of clinical factors, including the type of disease, severity of disease, and the subject's sensitivity to the therapeutic agent. Therapeutic Uses of Anti-Notch-3 Antibodies

It is contemplated that the antibodies of the present invention may be used to treat a mammal. In one embodiment, the antibody is administered to a nonhuman mammal for the purposes of obtaining preclinical data, for example. Exemplary nonhuman mammals to be treated include nonhuman primates, dogs, cats, rodents and other mammals in which preclinical studies are performed. Such mammals may be established animal models for a disease to be treated with the antibody or may be used to study toxicity of the antibody of interest. In each of these embodiments, dose escalation studies may be performed on the mammal.

An antibody administered alone or in combination with factor(s) can be used as a therapeutic. The present invention is directed to antibody-based therapies which involve administering antibodies of the invention to an animal, a mammal, or a human, for treating a Notch3-mediated disease, disorder, or condition. The animal or subject may be a mammal in need of a particular treatment, such as a mammal having been diagnosed with a particular disorder, e.g., one relating to Notch3. Antibodies directed against Notch3 are useful against degenerative diseases and other Notch3-associated diseases including CADASIL, FHM, Alagille syndrome, neurological and degenerative disorders in mammals, including but not limited to cows, pigs, horses, chickens, cats, dogs, non-human primates etc., as well as humans. For example, by administering

a therapeutically acceptable dose of an anti-Notch3 antibody, or antibodies, of the present invention, or a cocktail of the present antibodies, or in combination with other antibodies of varying sources, disease symptoms may be ameliorated or prevented in the treated mammal, particularly humans.

Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention as described below (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit, or prevent diseases, disorders, or conditions associated with aberrant expression and/or activity of Notch3, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of Notch3 includes, but is not limited to, alleviating at least one symptom associated with those diseases, disorders, or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

Anti-Notch3 antibodies of the present invention may be used therapeutically in a variety of diseases. The present invention provides a method for preventing or treating Notch3-mediated diseases in a mammal. The method comprises administering a disease preventing or treating amount of anti-Notch3 antibody to the mammal. The anti-Notch3 antibody binds to Notch3 and agonizes its function. Notch3 signaling has been linked to various diseases such as CADASIL, FHM, familial paroxysmic ataxia, Alagille syndrome, and other degenerative diseases and neurological disorders (Joutel, et al., *Nature* 383:707 (1996); Flynn, et al., *J Pathol* 204:55 (2004)). It is speculated that anti-Notch3 antibodies will also be effective to prevent the above mentioned diseases.

The amount of the antibody which will be effective in the treatment, inhibition, and prevention of a disease or disorder associated with aberrant expression and/or activity of Notch3 can be determined by standard clinical techniques. The dosage will depend on the type of disease to be treated, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody can be administered in treatment regimes consistent with the disease, e.g., a single or a few doses over one to several days to ameliorate a disease state or periodic doses over an extended time to inhibit disease progression and prevent disease recurrence. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 150 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent

administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the antibodies by modifications such as, for example, lipidation. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

The antibody variant composition will be formulated, dosed and administered in a manner consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The "therapeutically effective amount" of the antibody variant to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat a disease or disorder. The antibody variant need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore employed dosages.

The antibodies of the invention may be administered alone or in combination with other types of treatments.

In a preferred aspect, the antibody is substantially purified (e.g., substantially free from substances that limit its effect or produce undesired side-effects).

Various delivery systems are known and can be used to administer an antibody of the present invention, including injection, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu, et al., *J Biol Chem* 262:4429 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc.

The anti-Notch3 antibody can be administered to the mammal in any acceptable manner. Methods of introduction include but are not limited to parenteral, subcutaneous, intraperitoneal, intrapulmonary, intranasal, epidural, inhalation, and oral routes, and if desired for immunosuppressive treatment, intralesional administration. Parenteral infusions include intramuscular, intradermal, intravenous, intraarterial, or intraperitoneal administration. The antibodies or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the therapeutic antibodies or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. In addition, the antibody is suitably administered by pulse infusion, particularly with declining doses of the antibody. Preferably the dosing is given by injections, most preferably intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic.

Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent. The antibody may also be administered into the lungs of a patient in the form of a dry powder composition (See e.g., U.S. Pat. No. 6,514,496).

In a specific embodiment, it may be desirable to administer the therapeutic antibodies or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion, topical application, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering an antibody of the invention, care must be taken to use materials to which the protein does not absorb.

In another embodiment, the antibody can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249: 1527 (1990); Treat, et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein, et al., eds., pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-27; see generally *ibid.*).

In yet another embodiment, the antibody can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *Science* 249:1527 (1990); Sefton, *CRC Crit. Ref Biomed Eng* 14:201 (1987); Buchwald, et al., *Surgery* 88:507 (1980); Saudek, et al., *N Engl J Med* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer, et al., eds., CRC Press (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen, et al., eds., Wiley (1984); Ranger, et al., *J Macromol Sci Rev Macromol Chem* 23:61 (1983); see also Levy, et al., *Science* 228:190 (1985); During, et al., *Ann Neurol* 25:351 (1989); Howard, et al., *J Neurosurg* 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target.

The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of the antibody and a physiologically acceptable carrier. In a specific embodiment, the term "physiologically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such physiological carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable, or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium

stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable carriers are described in "Remington's Pharmaceutical Sciences" by E. W. Martin. Such compositions will contain an effective amount of the antibody, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In one embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration. The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

Articles of Manufacture

In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the disorders described above is provided. The article of manufacture comprises a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for preventing or treating the condition and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agent in the composition is the antibody. The label on, or associated with, the container indicates that the composition is used for treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically acceptable buffer, such as phosphate-buffered saline, Ringer's solution, and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

Antibody-Based Gene Therapy

In another aspect of the invention, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or prevent a disease or disorder associated with aberrant expression and/or activity of Notch3, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a therapeutic effect. Any of the methods for

gene therapy available can be used according to the present invention. Exemplary methods are described below.

For general reviews of the methods of gene therapy, see Goldspiel, et al., *Clinical Pharmacy* 12:488 (1993); Wu, et al., *Biotherapy* 3:87 (1991); Tolstoshev, *Ann Rev Pharmacol Toxicol* 32:573 (1993); Mulligan, *Science* 260:926 (1993); Morgan, et al., *Ann Rev Biochem* 62:191 (1993); May, *TIBTECH* 11:155 (1993).

In a one aspect, the compound comprises nucleic acid sequences encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific.

In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller, et al., *Proc Natl Acad Sci USA* 86:8932 (1989); Zijlstra, et al., *Nature* 342:435 (1989)). In specific embodiments, the expressed antibody molecule is a single chain antibody; alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids in vitro, then transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

In a specific embodiment, the nucleic acid sequences are directly administered in vivo, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated retrovirals or other viral vectors (see U.S. Pat. No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu, et al., *J Biol Chem* 262:4429 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller, et al., *Proc Natl Acad Sci USA* 86:8932 (1989); Zijlstra, et al., *Nature* 342:435 (1989)).

In a specific embodiment, viral vectors that contain nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller, et al., *Meth Enzymol* 217:581 (1993)). These retroviral vec-

tors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitate the delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen, et al., *Biotherapy* 6:291 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes, et al., *J Clin Invest* 93:644 (1994); Kiem, et al., *Blood* 83:1467 (1994); Salmons, et al., *Human Gene Therapy* 4:129 (1993); and Grossman, et al., *Curr Opin Gen and Dev* 3:110 (1993).

Adenoviruses may also be used in the present invention. Adenoviruses are especially attractive vehicles in the present invention for delivering antibodies to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky, et al., *Curr Opin Gen Dev* 3:499 (1993) present a review of adenovirus-based gene therapy. Bout, et al., *Human Gene Therapy* 5:3 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld, et al., *Science* 252:431 (1991); Rosenfeld, et al., *Cell* 68:143 (1992); Mastrangeli, et al., *J Clin Invest* 91:225 (1993); PCT Publication WO94/12649; Wang, et al., *Gene Therapy* 2:775 (1995). Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh, et al., *Proc Soc Exp Biol Med* 204:289 (1993); U.S. Pat. Nos. 5,436,146; 6,632,670; and 6,642,051).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler, et al., *Meth Enzymol* 217:599 (1993); Cohen, et al., *Meth Enzymol* 217:618 (1993); Cline, *Pharmac Ther* 29:69 (1985)) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

In a one embodiment, the cell used for gene therapy is autologous to the patient. Nucleic acid sequences encoding an antibody of the present invention are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained in vitro can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO 94/08598; Stemple, et al., *Cell* 71:973 (1992); Rheinwald, *Meth Cell Bio* 21A:229 (1980); Pittelkow, et al., *Mayo Clinic Proc* 61:771 (1986)).

EXAMPLES

Example 1

Generation of Immunogen: Notch3 Extracellular Domain-Fc Fusion Protein

Anti-Notch3 monoclonal antibodies that specifically bind to the LIN12/dimerization domain (herein after "LD") of human Notch3 were generated using a recombinant Notch3-Fc fusion protein as immunogen comprising Notch3 LD whose carboxy terminal end was fused to a gamma 1 Fc region. Specifically, the immunogen comprised amino acid residues 1378 to 1640 of Notch3 LD (See FIG. 1) and human γ 1Fc fusion protein (Notch3 LD/Fc). A control antibody was generated using the Notch3 EGF repeat region from amino acid residue 43 to 1377 as immunogen.

Notch3 protein sequence was analyzed using an internet-based research software and service (Motif Search). Human liver and pancreatic RNAs (Ambion, Inc. Austin, Tex.) were used as templates to synthesize the first strand of cDNA using a standard commercially available cDNA synthesis kit. The cDNAs encoding the Notch3 LD and the EGF repeat region were PCR-amplified in the presence of Betaine (1-2M) and DMSO (5%). The PCR-synthesized Notch3-LD DNA fragment (~0.8 kb) and Notch3-EGF repeat DNA fragment (~4 kb) were cloned into expression vectors comprising a His- γ 1Fc in the commercially available vector pSec or in the commercially available vector pCD3.1, each bearing a different antibiotic marker. This cloning resulted in two expression plasmids, one expressing a Notch3-LD/Fc fusion protein and the other expressing a Notch3-EGF/Fc fusion protein.

To facilitate the plasmid construction and to enhance the expression of the various Notch3 recombinant proteins, oligonucleotides corresponding to the leader peptide sequence comprising the first 135 base pairs of the Notch3 nucleic acid coding sequence were generated. These oligonucleotides contained some changes in the wobble coding positions to lower the GC content. All nucleotide sequence changes were silent, i.e., no amino acid sequence changes (FIGS. 8A and 8B). After annealing the oligonucleotides together, the engineered leader peptide coding sequence was linked to the rest of the coding sequence by PCR-SOE (Ho, et al., *Gene* 77:51 (1989); Horton, et al., *Bio Techniques* 8:528 (1990)) (See

FIG. 9). This leader peptide coding sequence was used in Notch3-LD/Fc and Notch3 expression constructs. Therefore, both of the Fc fusion proteins comprise a signal peptide linked to the N-terminus, and a human γ 1Fc sequence fused to the C-terminus. The amino acid sequence of Notch3-LD, including the leader peptide, is shown in FIG. 8B and SEQ ID NO:6.

Expression of Notch3-EGF/Fc and Notch3-LD/Fc fusion proteins were verified by transient transfection of the Notch3 expression plasmids into 293T (ATCC Number CRL-11268, Manassas, Va.) and CHO cells (Invitrogen, Carlsbad, Calif.), respectively. Prior to transfection, cells were cultured in DMEM (Invitrogen, Carlsbad, Calif.) growth medium containing 10% fetal calf serum (FCS), 2 mM of glutamine, and 1x essential amino acid solution followed by seeding about 3-5x10⁵ cells per well in 6-well plate and growing for approximately 24 hours. Three micrograms each of the Notch3 fusion protein expression plasmids were transfected into cells in each well using a LIPOFECTAMINE™ 2000 transfection system (Invitrogen, Carlsbad, Calif.) following the manufacturer's protocol. After transfection, the cells were cultured in fresh growth medium and cultured in a CO₂ incubator for approximately 40-48 hours before subjecting to Notch3 fusion protein expression analysis. Alternatively, after transfection, the cells were cultured in growth medium for 3-4 hours, then switched to DMEM medium containing 2% FCS and cultured for approximately 60-66 hours before drawing conditioned medium for secreted protein analysis.

Stable cell lines were generated for both Notch3-LD/Fc (His-Fc/pSec vector) and Notch3-EGF/Fc (His-Fc/pSec vector). Each plasmid was transfected into CHO cells. After transfection, the cells were cultured in DMEM growth medium overnight, then switched to growth medium with 800 μ g/ml hygromycin and cultured at least two weeks until the cells not carrying Notch3 expression plasmid were eliminated by the antibiotics. Conditioned media from the stable cell lines were subjected to Western blot analysis.

Stable or transient transfected cells were assayed for expression and secretion of Notch3-LD/Fc or Notch3-EGF/Fc fusion protein. Transfected cells harvested from culture dishes were washed once with phosphate buffered saline (PBS) and resuspended in deionized water, mixed with an equal volume of 2x protein sample loading buffer (BioRad, Hercules, Calif.) and then heated at about 100° C. for 10 minutes. Secreted protein was analyzed using conditioned medium mixed with an equal volume of 2x protein sample loading buffer and heated at 100° C. for 10 minutes. The samples were separated using 4-15% gradient SDS-PAGE. The proteins were transferred from the gel to a PVDF membrane (BioRad, Hercules, Calif.), which was blocked in 5% non-fat dry milk in PBST (PBS with 0.05% TWEEN-20®) for at least one hour prior to transfer of protein.

Notch3-EGF/Fc and Notch3-LD/Fc fusion proteins were detected by incubating with γ Fc-specific, HRP-conjugated antibody (Sigma, St Louis, Mo.) in blocking buffer for one hour at room temperature. The membrane was washed three times in PBST and developed with a chemiluminescent substrate.

For Notch3 domain/Fc fusion protein purification, CHO stable cell lines as described above were cultured in DMEM with 2% FCS for up to 5 days. One liter of conditioned medium collected, and subjected to protein-A bead-packed column for affinity binding. The column was washed with PBS, and the bound proteins were eluted in 50 mM citrate buffer (pH 2.8), and the pH was brought to neutral by adding 1 M Tris-HCl buffer (pH 8). Purity of the protein was assessed by protein gel analysis using 4-15% gradient SDS-PAGE. Protein concentration was assayed using Coomassie blue

37

reagent following the manufacturer's protocol (Pierce, Rockford, Ill.). Through this procedure, milligram quantities of Notch3-LD/Fc and Notch3-EGF/Fc protein were purified for immunization and ELISA binding assays.

Example 2

Generation of Anti-Notch3 MAbs

Male A/J mice (Harlan, Houston, Tex.), 8-12 week old, were injected subcutaneously with 25 µg of Notch3-EGF/Fc or Notch3-LD/Fc in complete Freund's adjuvant (Difco Laboratories, Detroit, Mich.) in 200 µl of PBS. Two weeks after the injections and three days prior to sacrifice, the mice were again injected intraperitoneally with 25 µg of the same antigen in PBS. For each fusion, single cell suspensions were prepared from spleen of an immunized mouse and used for fusion with Sp2/0 myeloma cells; 5×10^8 of Sp2/0 and 5×10^8 of spleen cells were fused in a medium containing 50% polyethylene glycol (M.W. 1450) (Kodak, Rochester, N.Y.) and 5% dimethylsulfoxide (Sigma, St. Louis, Mo.). The cells were then adjusted to a concentration of 1.5×10^5 spleen cells per 200 µl of the suspension in Iscove medium (Invitrogen, Carlsbad, Calif.), supplemented with 10% fetal bovine serum, 100 units/ml of penicillin, 100 µg/ml of streptomycin, 0.1 µM hypoxanthine, 0.4 µM aminopterin, and 16 µM thymidine. Two hundred microliters of the cell suspension were added to each well of about sixty 96-well plates. After around ten days, culture supernatants were withdrawn for screening their antibody-binding activity using ELISA.

The 96-well flat bottom IMMULON® II microtest plates (Dynatech, Laboratories, Chantilly, Va.) were coated using 100 µl of Notch3-EGF/Fc or Notch3-LD/Fc (0.1 µg/ml) in (PBS) containing 1× Phenol Red and 3-4 drops pHix/liter (Pierce, Rockford, Ill.) and incubated overnight at room temperature. After the coating solution was removed by flicking of the plate, 200 µl of blocking buffer containing 2% BSA in PBST containing 0.1% merthiolate was added to each well for one hour to block non-specific binding. The wells were then washed with PBST. Fifty microliters of culture supernatant from each fusion well was collected and mixed with 50 µl of blocking buffer and then added to the individual wells of the microtiter plates. After one hour of incubation, the wells were washed with PBST. The bound murine antibodies were then detected by reaction with horseradish peroxidase (HRP)-conjugated, Fc-specific goat anti-mouse IgG (Jackson ImmunoResearch Laboratories, West Grove, Pa.). HRP substrate solution containing 0.1% 3,3',5,5'-tetramethyl benzidine and 0.0003% hydrogen peroxide was added to the wells for color development for 30 minutes. The reaction was terminated by the addition of 50 ml of 2 M H₂SO₄/well. The OD at 450 nm was read with an ELISA plate reader (Molecular Devices, Sunnyvale, Calif.).

Among 185 hybridomas isolated and analyzed, one hybridoma clone from mice immunized with Notch3-LD/Fc generated a Notch3 agonist antibody 256A-13 and this antibody was further characterized. An ELISA was performed using supernatant from the hybridoma clone producing MAbs 256A-13. The results showed strong binding activity to the purified Notch3 LD/FC fusion protein to which it was generated and did not bind to human Notch1-LD/Fc (LIN/dimerization domain fused to Fc region at the carboxyl terminus) or a control human Fc protein (data not shown) (Table 1).

38

TABLE 1

ELISA OD readings of 256A-13 using hybridoma supernatant			
Target protein Hybridoma	Notch3-LD/Fc		
	supernatant	Control IgG1 MAb	256A-13
Mean		0.019	2.828
S.D.		0.002	0.047

The positive hybridoma clone from this primary ELISA screening was further isolated by single colony-picking and a second ELISA assay as described above was done to verify specific binding to the chosen immunogen. The confirmed hybridoma clone was expanded in larger scale cultures. The monoclonal antibodies (MAbs) were purified from the medium of these large scale cultures using a protein A affinity column. The anti-Notch3 agonist MAbs were then characterized using cell-based binding assays, microscopy, Western blot, and FACS analysis.

Example 3

Cell-Based Binding Assays for Anti-Notch3 MAbs

The cell-based binding assays used to characterize the anti-Notch3 MAbs required cloning a full-length of human Notch3 open reading frame into a vector, in this case PcDNATM3.1/Hygro (Invitrogen, Carlsbad, Calif.). The Notch3-coding region was synthesized by RT-PCR using human liver tumor RNA (Ambion, Inc., Austin, Tex.) as a template. The final plasmid construct, Notch3/Hygro, expressed a full-length Notch3 protein as depicted in FIG. 1. A stable cell line expressing Notch3 was generated by transfection of Notch3/Hygro plasmid construct into 293T cells (ATCC No. CRL-11268) using a LIPOFECTAMINETM 2000 kit following the same procedure as described in Example 1. After transfection, the cells were cultured in DMEM growth medium overnight, then reseeded in growth medium with 200 µg/ml hygromycin and cultured for 12-14 days. Well-isolated single colonies were picked and grown in separate wells until enough clonal cells were amplified. Stable 293T clones that were resistant to hygromycin selection and expressed high levels of Notch3 protein were identified by Western blot analysis, and by fluorescent electromicroscopy using polyclonal anti-Notch3 antibodies (R&D Systems, Minneapolis, Minn.).

A partial Notch3 expression plasmid containing only the Notch LIN12/dimerization (LD) domain and the transmembrane (TM) domain was also constructed by PCR and subcloned into PcDNATM3.1 vector.

Human Sup-T1 cell line (ATCC No. CRL-1942) naturally expressing Notch3 was also confirmed by Western blot. Sup-T1 cells were grown in RPMI1640 media containing 10% fetal calf serum, 2 mM of glutamine and 1× essential amino acid solution.

Cell-based antibody-binding was assessed using FMATM (fluorescence macro-confocal high-throughput screening) 8100 HTS System (Applied Biosystems, Foster City, Calif.) following the protocol provided by the manufacturer. Cell lines naturally expressing Notch3 or stably transfected with Notch3 expression constructs were seeded in 96-well plates. Alternatively, transiently transfected 293T or CHO cells were seeded in the 96-well plate. The cells were seeded at a density of 30,000-50,000 cells per well. After 20-24 hours, anti-Notch3 MAbs and 1×PBS reaction buffer were added to the

wells and incubated for one hour at 37° C. Cy-5-conjugated anti-mouse IgG antibody was added in the wells after removal of primary antibodies.

Cell-based antibody-binding was also assessed by fluorescence-activated cell sorter (FACS) using internally generated 293T/Notch3-stable cell line and two cancer lines, human Sup-T1 and A2780 cell lines (UK ECACC No. Cat. No. 93112519), both naturally express Notch3 (data not shown). Cells were first incubated with anti-Notch3 MABs in 1×PBS. After three washes, the cells were incubated with fluorescent molecule-conjugated secondary antibody. The cells were resuspended, fixed in 1×PBS with 0.1% paraformaldehyde, and analyzed by FACS (BD Sciences, Palo Alto, Calif.). The results indicated that 256A-13 binds to Notch3 receptor expressed either from recombinant plasmid constructs or as native protein in cultured cells (Table 2). Transiently transfected 293T cells containing a Notch3/Hygro plasmid were also stained with immunofluorescence as described above and observed by fluorescent microscopy.

TABLE 2

Binding activity of 256A-13 in cell-based FACS analysis shown as mean fluorescent intensity		
	Control IgG1	256A-13
Notch3/Hygro	24.16	32.2
Sup-T1	24.51	55.44

The cell-based FMAT and FACS analyses confirmed that MABs 256A-13 indeed binds to the Notch3 receptor expressed either from recombinant plasmid constructs or as native protein in cultured cells (Table 2 and Table 3).

TABLE 3

Summary of anti-Notch3 MABs binding activity in cell-based FMAT		
Antibody	Control IgG1	256A-13
Notch3 (full-length)	no binding	weak binding
Notch3-LDTM	no binding	strong binding

A positive binding signal was determined based on the FMAT™ system signal read-out that was significantly higher than that of the IgG1 control and other negative hybridoma clones ($p > 0.01$). The IgG1 control binding read-out was considered background. 293T cells transiently transfected with Notch3/Hygro plasmid were also stained with immunofluorescence as described above and observed by fluorescent microscopy.

The binding affinity of MAB 256A-13 was analyzed by Biacore System (Biacore Inc., Piscataway, N.J.). The antibody was directly immobilized on a chip through amine coupling (immobilization level: 200 RU), and the Notch3-LD/Fc protein (antigen) was injected at 5 different concentrations (ranging from 37.5 to 120 nM with association time between 5-8 minutes, and dissociation time between 1 and 2 hours). The running buffer and the sample buffer are PBS contained 5 mM Ca^{2+} . The chip surface was regenerated with 10 mM glycine, pH2. The antibody was characterized in duplicate. Table 4 discloses the statistical mean, standard errors and Kinetic dissociation constant (KD) calculated. The antibody has a high affinity with a KD of 280 μM , and a slow off-rate. Both the standard errors and chi square are low with a good fit (dynamic curve not shown).

TABLE 4

Characterization of MAB 256A-13 binding affinity by Biacore						
Sample	KD [pM]	ka [M-1s-1]	SE (ka)	kd [s-1]	SE (kd)	χ^2
256A-13	280	4.20e4	0.98	1.18e-5	1.02e-7	0.392

KD: 256A-13 and Notch3-LD/Fc dissociation constant. Ka: Rate of 256A-13 binding to Notch3-LD/Fc (or On-rate). Kd: Rate of 256A-13 dissociate from Notch3-LD/Fc (or Off-rate). SE: standard error.

Example 4

Western Blot Analysis of 256A-13 Binding Activity

Western blot was performed to assess the binding activity of 256A-13 to Notch3 receptor under denaturing conditions, as well as expression levels of Notch3 and other Notch-related proteins in human cell lines. Purified Notch3-LD/Fc fusion protein was combined with protein loading buffer. Protein samples were also prepared from the transiently or stably transfected cells described in Example 1, which were harvested from culture dishes, washed once with PBS, resuspended in total cellular protein extract buffer (Pierce, Rockford, Ill.), and heated at 100° C. for 10 minutes after adding equal volume of 2× protein sample loading buffer. All samples were separated by electrophoresis in a 4-15% gradient SDS-PAGE. The proteins were transferred from gel to PVDF membrane and 256A-13 was applied to the Western blot membrane as the primary detection antibody. An HRP-conjugated secondary antibody was used for detection and the signal generated using a chemiluminescent substrate as described above. Positive control antibodies against human Fc, V5 tag, Notch3 and Notch1 were purchased from (Invitrogen, R&D Systems, Santa Cruz Biotechnologies, and Orbigen).

Western blot analysis showed that MAB 256A-13 binds to Notch3-LD/Fc under denaturing condition, as well as native molecular conformation as observed in ELISA and FACS analysis.

Example 5

Assessing Functionality of 256A-13 by Luciferase Reporter Assay

A. Plasmid Constructs

The full length Notch3 expression construct described in Example 3 above was confirmed by sequencing, and is identical to the published sequence depicted in FIG. 1. The expression of Notch3 was verified by transient transfection and Western blot as described in Example 4.

To generate a luciferase reporter plasmid for Notch signaling, two complementary oligonucleotide primers containing tandem repeats of CBF1 binding motif were synthesized having the following sequences:

(SEQ ID NO 12)
5' GCTCGAGCTCGTGGGAAAATACCGTGGGAAAATGAACCGTGGGAAAATCTCTGTG

(SEQ ID NO 13)
5' GCTCGAGATTTTCCACGAGATTTTCCACGTTTC

These two oligoprimers were annealed at 65° C. in 100 mM of NaCl with each oligo at a concentration of 4 mM. After annealing to each other, the primers were extended by PCR. The PCR product was cloned into a commercially available

vector. The insert was verified by sequencing, which contains four tandem repeats of CBF1 binding motif and two flanking Xho I sites. The insert was excised using Xho I and ligated downstream of the firefly luciferase reporter coding sequence.

After luciferase reporter assay and sequencing analysis, plasmid clones with eight repeats of CBF1 binding motifs were selected and designated CBF1-Luc.

B. Stable Cell Line Generation

Two stable cell lines were generated for functional assays using human embryonic kidney cell lines (HEK293). One cell line contained the Notch3-expressing plasmid and CBF1-Luc reporter plasmid integrated into the nuclear genome. This cell line was generated by cotransfecting Notch3/hygromycin and CBF1-Luc plasmids into 293T cells using LIPO-FECTAMINE™ 2000 according to the manufacturer's protocol. Stable transfection cell clones were selected against 200 µg/ml hygromycin in DMEM growth medium, and screened by luciferase reporter assay and Western blot. A cell line with a relatively high level of Notch3 receptor expression (based on Western blot) and luciferase activity was selected for use in functional assays, and designated NC85.

C. Luciferase Reporter Assay with Notch3 Overexpressing Cells Alone

NC85 cells were cultured in the presence of MAb 256-A13 for 24 to 48 hours. The media was then removed by aspiration, cells were lysed in 1× Passive Lysis Buffer (E1501, Promega, Madison, Wis.) and luciferase activities were assayed using the Luciferase Assay System following manufacturer's protocol (E1501, Promega, Madison, Wis.) in TD-20/20 luminometer (Turner Designs Instrument, Sunnyvale, Calif.). As illustrated in FIG. 5, NC85 cells cultured in the presence of MAb 256-A13, the luciferase activity was increased almost 4 fold as compared to that with control antibody G3. The luciferase reporter assay demonstrated that MAb 256-A13 induced a dramatic increase in luciferase activity without ligand binding, while antagonist anti-Notch3 antibodies MAbs 256A-4 and 256A-8 did not (FIG. 5).

Example 8

Mapping the Binding Epitope of 256A-13

A. Epitope-Mapping Strategy and Rationale Using Notch3 Single Domain and Fc Fusion Protein Constructs

Notch3 LIN12/heterodimerization domains, also called Notch3 LIN12-dimerization domain (Notch3-LD) consisted of three LIN12 domains, 1st LIN12 (L1.), 2nd LIN12 (L2) and 3rd LIN12 (L3) (See FIG. 10). Five Notch3 single domain/Fc fusion protein expression constructs (FIG. 7) were generated, and a western blot was performed to assess which domain was sufficient for MAb 256A-13 binding. After transient transfection, the supernatants with secreted Notch3 single domain/Fc fusion proteins were analyzed by SDS-PAGE. The results showed that MAb 256A-13 only binds to Notch3-L1, and not to any other domains. ELISA experiments also showed that MAb 256A-13 has very strong binding to Notch3-L1 and weak binding to Notch3-L3, and not to other domains (Table 5).

TABLE 5

Summary of Western blot results and ELISA Readings using MAb 256A-13 against Notch3-domain/Fc fusion protein constructs				
MAb	Western blot result		ELISA OD reading	
	256A-13	Anti-human Fc	256A-13	Anti-human Fc
Notch3-LD/Fc	positive band	positive band	1.882	1.557
Notch3-L1/Fc	positive band	positive band	1.797	1.364
Notch3-L2/Fc	no band	positive band	0.015	1.337
Notch3-L3/Fc	no band	positive band	1.054	1.425
Notch3-D1/Fc	no band	positive band	0.015	1.608
Notch3-D2/Fc	no band	positive band	0.015	1.628

A. Identification of Binding Epitope(s) by Subdomain Swap

First, the agonist Notch3 MAb, 256A-13, binds to Notch3 LIN12/dimerization domain (LD), but not to the homologous human Notch1 LIN12/dimerization domain (Table 5). Second, the anti-Notch3 MAb binds to denatured Notch3 protein in Western blot as discussed in Example 4 and 8, indicating that 256A-13 binds to a single epitope or to discrete epitopes independent of each other. Third, Notch3 and Notch1 share approximately 55% amino acid sequence homology in LIN12/dimerization domain, therefore it was concluded that a subdomain swap between Notch3 and Notch1 within this region would not disrupt the protein conformation. Notch1-LD cDNA was PCR-amplified using standard PCR methods. The first strand cDNA template was synthesized from PA-1 cell total RNA (ATCC No. CRL-1572). The human IgG kappa chain leader peptide coding sequence was PCR-amplified, used as leader peptide to link to the 5' of Notch1-LD by PCR-SOE and subcloned in His-γ1Fc/pSec.

TABLE 6

ELISA OD readings of MAbs 256A-13 and control IgG1 binding to Notch3-LD/Fc or Notch1-LD/Fc				
	Notch1-LD/Fc		Notch3-LD/Fc	
	Mean	S.D.	Mean	S.D.
256A-13	0.094	0.007	4.000	0
IgG1 control	0.066	0.006	0.063	0.006

B. Generation of Subdomain Swap Fusion Protein Constructs

Based on the ELISA analysis results presented in Section A above, the target domain of the 1st LIN12 domain, or L1 was further divided into three subdomains and individually swapped with the corresponding subdomain of Notch1-L1. The subdomain swap constructs were generated using PCR-SOE (Ho, et al., *Gene* 77:51 (1989); Horton, et al., *BioTechniques* 8:528 (1990)) as illustrated in FIGS. 9 and 10. PCR and PCR-SOE reactions were performed using PCR with 1M Betaine and 5% DMSO added to the reaction. The final PCR-SOE product was subcloned and verified by sequencing. The plasmid clone with the correct insert sequence was cleaved with Nhe I and Xho I to excise the insert, which was gel-purified and subcloned. The five Notch3/Notch1 subdomain swap constructs are illustrated in FIG. 7. To facilitate the epitope mapping, the human IgG kappa chain signaling peptide was used as leader peptide in the domain swap constructs. The amino acid sequences of the subdomain constructs are shown in FIG. 10.

C. Expression of Notch3/Notch1 Subdomain Swap Fusion Protein

Notch3/Notch1-LD domain swap plasmids were transiently transfected in CHO cells using LipoFectamine 2000. CHO cells were seeded in DMEM growth medium with 10% FCS at 0.8×10^6 cells per well in 6-well plate, maintained in CO₂ incubator overnight before transfection. The cells were recovered after transfection in the growth medium for about 3 hours, then switched to DMEM with 2% FCS, and cultured for three days. The conditioned media were harvested and centrifuged at 3500 rpm for 10 minutes. The supernatant containing Notch3-LD domain swap protein secreted from CHO was collected and prepared for Western blot and ELISA binding analyses. ELISA showed that all the domain-swap fusion proteins were expressed and secreted in conditioned medium (Table 4), which was further confirmed by Western blot analysis (data not shown).

The ELISA readings used anti-human Fc antibody as detection antibody showing all the proteins were expressed in conditioned medium. Human IgG/Fc was used as a control. The starting point of human IgG/Fc coated in each well is 100 ng.

D. Epitope Binding Analysis using ELISA

The 96-well flat bottom IMMULON® II microtest plates (Dynatech, Laboratories, Chantilly, Va.) were coated with anti-human Fc antibody (Jackson ImmunoResearch) by adding 100 µl of the antibody (0.1 µg/ml) in phosphate buffered saline (PBS) containing 1× Phenol Red and 3-4 drops pHix/liter (Pierce, Rockford, Ill.), and incubated overnight at room temperature. After the coating solution was removed by flicking of the plate, 200 µl of blocking buffer containing 2% BSA in PBST and 0.1% merthiolate was added to each well for one hour to block non-specific binding. The wells were then washed with PBST. Fifty microliters of the above conditioned medium from each transfection of Notch3/Notch1 domain swap construct were collected, mixed with 50 µl of blocking buffer, and added to the individual wells of the microtiter plates. After one hour of incubation, the Notch3/Notch1-LD domain swap protein was captured by the coated anti-Fc antibody, and the wells were washed with PBST. Anti-Notch3 MAbs and isotype-matched control MAbs were serially diluted in blocking buffer as above, and 50 µl of the diluted MAbs were added in each well to assess binding to the bound Notch3/Notch1 domain swap protein. Horseradish peroxidase (HRP)-conjugated, Fc-specific goat anti-mouse IgG was used for detection. HRP substrate solution containing 0.1% 3,3',5,5'-tetramethyl benzidine and 0.0003% hydrogen peroxide was added to the wells for color development for 30 minutes. The reaction was terminated by addition of 50 ml of 2 M H₂SO₄/well. The OD at 450 nm was read with an ELISA reader. Subdomain swap constructs and clusters of mutations were similarly examined by ELISA analysis above.

ELISA binding experiments using MAb 256A-13 against the subdomain-swap proteins showed that the swap of the 1st subdomain in Notch3-L1 domain (L1) did not affect the binding, indicating that 256A-13 does not bind to this region. On the other hand, the swaps of the 2nd and 3rd subdomains in Notch3-L1 significantly reduced the binding. Therefore, those two subdomains contain the binding epitope(s) for MAb 256A-13. (FIG. 10). In contrast, isotype-matched negative control antibody, G3, does not bind to any of the domain swap fusion proteins in the ELISA assay (FIG. 10). It was concluded from the above experiments that the 1st LIN12 domain was required for MAb 256A-13 binding, and specifically within the 2nd and 3rd subdomain region.

To further map the specific epitope that MAb 256A-13 binds, the 2nd and 3rd subdomains of Notch3-L1 domain were

further divided into five amino acids clusters, and swapped with the corresponding amino acid residues in Notch1 (FIG. 10). ELISA binding assay showed that the swap from DRE (Notch3 sequence) to SQL (Notch1 sequence) completely abolished the ELISA binding activity, indicating that only this epitope is required for MAb 256A-13 binding within Notch3-L1 domain.

Pinpoint analysis of amino acid residues required for MAb 256A-13 binding is done by using di-Alanine peptide scanning. The Alanine peptides cover the DRE epitope mapped by amino acid swap analysis. The peptide is synthesized as a spot cross-linked to nylon support membrane. Antibody blot binding is assessed by dot blot. MAb G3 is used as a control IgG1. The peptide sequences are presented in FIG. 11.

Example 9

Sequencing of Anti-Notch3 MAbs

Because antibody binding properties are fully-dependent on the variable regions of both heavy chain and light chain, the variable sequences of 256A-13 were subtyped and sequenced. The antibody IgG subtype was determined using an ISOSTRIP™ mouse monoclonal antibody isotyping kit (Roche Diagnostics, Indianapolis, Ind.). The results showed that 256A-13 has an IgG₁ heavy chain and a kappa light chain.

The variable region sequences of heavy chain and light chain were decoded through RT-PCR and cDNA cloning. Total RNAs from hybridoma clones 256A-13 were isolated using an RNeasy Mini kit following manufacturer's protocol (Qiagen Sciences, Valencia, Calif.). The first strand cDNA was synthesized using the RNA template and SUPER-SCRIPTASE® III reverse transcriptase kit. The variable region of light chain and heavy chain cDNAs were PCR-amplified from the first strand cDNA using degenerative forward primers covering the 5'-end of mouse kappa chain coding region and a reverse primer matching the constant region at the juncture to the 3'-end of the variable region, or using degenerative forward primers covering the 5'-end of mouse heavy chain coding region and a constant region reverse primer in mouse heavy chain. The PCR product was cloned into a commercially available vector and sequenced by Lone Star Lab (Houston, Tex.). The nucleotide sequences were analyzed utilizing the DNASTAR® computer software program (DNASTAR, Inc., Madison, Wis.). Each anti-Notch3 MAb sequence was determined by sequences from multiple PCR clones derived from the same hybridoma clone.

The variable heavy chain region of MAb 256A-13 contains 121 amino acid residues and the light chain variable region contains 102 amino acid residues (FIGS. 4A and 4B).

Example 10

Impact of Notch3 Agonistic Antibodies on Metalloprotease Cleavage of Notch3

Notch receptor activation involves ligand induced metalloprotease cleavage at juxtamembrane site (S2) generating an extracellular subunit. This cleavage is an essential prerequisite to S3 cleavage to release the activated Notch intracellular region. To test whether the agonizing antibodies can induce ligand-independent sequential Notch activation events, including two proteolytic cleavages, 293T cells stably expressing a recombinant Notch3 receptor (NC85 cells) were treated with either G3 or 256-A13. The soluble extracellular subunits generated by proteolytic cleavage in the culture medium were detected by an ELISA assay using an antibody

45

bound to a solid surface that recognizes the Notch3 cleavage product. As shown in FIG. 6, Notch3 agonistic MAb significantly increased the generation of soluble Notch3 extracellular subunits in the conditioned medium, whereas control antibody G3 did not.

Example 12

Assay for Notch3 Related Diseases

To identify other Notch3 related diseases, one can 10 sequence the Notch3 gene from patient samples, or perform

46

immunohistochemistry to check for the under-expression of Notch3 receptor using patient tissue. In addition, one can isolate and culture cells from a patient suspected of having a Notch3 associated disease and study the impact of an agonistic antibody of the present invention on Notch3 signaling. 5

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 44

<210> SEQ ID NO 1

<211> LENGTH: 2321

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```
Met Gly Pro Gly Ala Arg Gly Arg Arg Arg Arg Arg Arg Pro Met Ser
1          5          10          15

Pro Pro Pro Pro Pro Pro Pro Val Arg Ala Leu Pro Leu Leu Leu
20          25          30

Leu Ala Gly Pro Gly Ala Ala Ala Pro Pro Cys Leu Asp Gly Ser Pro
35          40          45

Cys Ala Asn Gly Gly Arg Cys Thr Gln Leu Pro Ser Arg Glu Ala Ala
50          55          60

Cys Leu Cys Pro Pro Gly Trp Val Gly Glu Arg Cys Gln Leu Glu Asp
65          70          75          80

Pro Cys His Ser Gly Pro Cys Ala Gly Arg Gly Val Cys Gln Ser Ser
85          90          95

Val Val Ala Gly Thr Ala Arg Phe Ser Cys Arg Cys Pro Arg Gly Phe
100         105         110

Arg Gly Pro Asp Cys Ser Leu Pro Asp Pro Cys Leu Ser Ser Pro Cys
115         120         125

Ala His Gly Ala Arg Cys Ser Val Gly Pro Asp Gly Arg Phe Leu Cys
130         135         140

Ser Cys Pro Pro Gly Tyr Gln Gly Arg Ser Cys Arg Ser Asp Val Asp
145         150         155         160

Glu Cys Arg Val Gly Glu Pro Cys Arg His Gly Gly Thr Cys Leu Asn
165         170         175

Thr Pro Gly Ser Phe Arg Cys Gln Cys Pro Ala Gly Tyr Thr Gly Pro
180         185         190

Leu Cys Glu Asn Pro Ala Val Pro Cys Ala Pro Ser Pro Cys Arg Asn
195         200         205

Gly Gly Thr Cys Arg Gln Ser Gly Asp Leu Thr Tyr Asp Cys Ala Cys
210         215         220

Leu Pro Gly Phe Glu Gly Gln Asn Cys Glu Val Asn Val Asp Asp Cys
225         230         235         240

Pro Gly His Arg Cys Leu Asn Gly Gly Thr Cys Val Asp Gly Val Asn
245         250         255

Thr Tyr Asn Cys Gln Cys Pro Pro Glu Trp Thr Gly Gln Phe Cys Thr
260         265         270

Glu Asp Val Asp Glu Cys Gln Leu Gln Pro Asn Ala Cys His Asn Gly
275         280         285
```

-continued

Gly Thr Cys Phe Asn Thr Leu Gly Gly His Ser Cys Val Cys Val Asn
 290 295 300
 Gly Trp Thr Gly Glu Ser Cys Ser Gln Asn Ile Asp Asp Cys Ala Thr
 305 310 315 320
 Ala Val Cys Phe His Gly Ala Thr Cys His Asp Arg Val Ala Ser Phe
 325 330 335
 Tyr Cys Ala Cys Pro Met Gly Lys Thr Gly Leu Leu Cys His Leu Asp
 340 345 350
 Asp Ala Cys Val Ser Asn Pro Cys His Glu Asp Ala Ile Cys Asp Thr
 355 360 365
 Asn Pro Val Asn Gly Arg Ala Ile Cys Thr Cys Pro Pro Gly Phe Thr
 370 375 380
 Gly Gly Ala Cys Asp Gln Asp Val Asp Glu Cys Ser Ile Gly Ala Asn
 385 390 395 400
 Pro Cys Glu His Leu Gly Arg Cys Val Asn Thr Gln Gly Ser Phe Leu
 405 410 415
 Cys Gln Cys Gly Arg Gly Tyr Thr Gly Pro Arg Cys Glu Thr Asp Val
 420 425 430
 Asn Glu Cys Leu Ser Gly Pro Cys Arg Asn Gln Ala Thr Cys Leu Asp
 435 440 445
 Arg Ile Gly Gln Phe Thr Cys Ile Cys Met Ala Gly Phe Thr Gly Thr
 450 455 460
 Tyr Cys Glu Val Asp Ile Asp Glu Cys Gln Ser Ser Pro Cys Val Asn
 465 470 475 480
 Gly Gly Val Cys Lys Asp Arg Val Asn Gly Phe Ser Cys Thr Cys Pro
 485 490 495
 Ser Gly Phe Ser Gly Ser Thr Cys Gln Leu Asp Val Asp Glu Cys Ala
 500 505 510
 Ser Thr Pro Cys Arg Asn Gly Ala Lys Cys Val Asp Gln Pro Asp Gly
 515 520 525
 Tyr Glu Cys Arg Cys Ala Glu Gly Phe Glu Gly Thr Leu Cys Asp Arg
 530 535 540
 Asn Val Asp Asp Cys Ser Pro Asp Pro Cys His His Gly Arg Cys Val
 545 550 555 560
 Asp Gly Ile Ala Ser Phe Ser Cys Ala Cys Ala Pro Gly Tyr Thr Gly
 565 570 575
 Thr Arg Cys Glu Ser Gln Val Asp Glu Cys Arg Ser Gln Pro Cys Arg
 580 585 590
 His Gly Gly Lys Cys Leu Asp Leu Val Asp Lys Tyr Leu Cys Arg Cys
 595 600 605
 Pro Ser Gly Thr Thr Gly Val Asn Cys Glu Val Asn Ile Asp Asp Cys
 610 615 620
 Ala Ser Asn Pro Cys Thr Phe Gly Val Cys Arg Asp Gly Ile Asn Arg
 625 630 635 640
 Tyr Asp Cys Val Cys Gln Pro Gly Phe Thr Gly Pro Leu Cys Asn Val
 645 650 655
 Glu Ile Asn Glu Cys Ala Ser Ser Pro Cys Gly Glu Gly Gly Ser Cys
 660 665 670
 Val Asp Gly Glu Asn Gly Phe Arg Cys Leu Cys Pro Pro Gly Ser Leu
 675 680 685
 Pro Pro Leu Cys Leu Pro Pro Ser His Pro Cys Ala His Glu Pro Cys
 690 695 700
 Ser His Gly Ile Cys Tyr Asp Ala Pro Gly Gly Phe Arg Cys Val Cys

-continued

705	710	715	720
Glu Pro Gly Trp Ser Gly Pro Arg Cys Ser Gln Ser Leu Ala Arg Asp	725	730	735
Ala Cys Glu Ser Gln Pro Cys Arg Ala Gly Gly Thr Cys Ser Ser Asp	740	745	750
Gly Met Gly Phe His Cys Thr Cys Pro Pro Gly Val Gln Gly Arg Gln	755	760	765
Cys Glu Leu Leu Ser Pro Cys Thr Pro Asn Pro Cys Glu His Gly Gly	770	775	780
Arg Cys Glu Ser Ala Pro Gly Gln Leu Pro Val Cys Ser Cys Pro Gln	785	790	795
Gly Trp Gln Gly Pro Arg Cys Gln Gln Asp Val Asp Glu Cys Ala Gly	805	810	815
Pro Ala Pro Cys Gly Pro His Gly Ile Cys Thr Asn Leu Ala Gly Ser	820	825	830
Phe Ser Cys Thr Cys His Gly Gly Tyr Thr Gly Pro Ser Cys Asp Gln	835	840	845
Asp Ile Asn Asp Cys Asp Pro Asn Pro Cys Leu Asn Gly Gly Ser Cys	850	855	860
Gln Asp Gly Val Gly Ser Phe Ser Cys Ser Cys Leu Pro Gly Phe Ala	865	870	875
Gly Pro Arg Cys Ala Arg Asp Val Asp Glu Cys Leu Ser Asn Pro Cys	885	890	895
Gly Pro Gly Thr Cys Thr Asp His Val Ala Ser Phe Thr Cys Thr Cys	900	905	910
Pro Pro Gly Tyr Gly Gly Phe His Cys Glu Gln Asp Leu Pro Asp Cys	915	920	925
Ser Pro Ser Ser Cys Phe Asn Gly Gly Thr Cys Val Asp Gly Val Asn	930	935	940
Ser Phe Ser Cys Leu Cys Arg Pro Gly Tyr Thr Gly Ala His Cys Gln	945	950	955
His Glu Ala Asp Pro Cys Leu Ser Arg Pro Cys Leu His Gly Gly Val	965	970	975
Cys Ser Ala Ala His Pro Gly Phe Arg Cys Thr Cys Leu Glu Ser Phe	980	985	990
Thr Gly Pro Gln Cys Gln Thr Leu Val Asp Trp Cys Ser Arg Gln Pro	995	1000	1005
Cys Gln Asn Gly Gly Arg Cys Val Gln Thr Gly Ala Tyr Cys Leu	1010	1015	1020
Cys Pro Pro Gly Trp Ser Gly Arg Leu Cys Asp Ile Arg Ser Leu	1025	1030	1035
Pro Cys Arg Glu Ala Ala Ala Gln Ile Gly Val Arg Leu Glu Gln	1040	1045	1050
Leu Cys Gln Ala Gly Gly Gln Cys Val Asp Glu Asp Ser Ser His	1055	1060	1065
Tyr Cys Val Cys Pro Glu Gly Arg Thr Gly Ser His Cys Glu Gln	1070	1075	1080
Glu Val Asp Pro Cys Leu Ala Gln Pro Cys Gln His Gly Gly Thr	1085	1090	1095
Cys Arg Gly Tyr Met Gly Gly Tyr Met Cys Glu Cys Leu Pro Gly	1100	1105	1110
Tyr Asn Gly Asp Asn Cys Glu Asp Asp Val Asp Glu Cys Ala Ser	1115	1120	1125

-continued

Gln Pro	Cys Gln His Gly	Gly	Ser Cys Ile Asp	Leu Val Ala Arg
1130		1135		1140
Tyr Leu	Cys Ser Cys Pro	Pro Gly Thr Leu Gly	Val	Leu Cys Glu
1145		1150		1155
Ile Asn	Glu Asp Asp Cys	Gly	Pro Gly Pro Pro	Leu Asp Ser Gly
1160		1165		1170
Pro Arg	Cys Leu His Asn	Gly	Thr Cys Val Asp	Leu Val Gly Gly
1175		1180		1185
Phe Arg	Cys Thr Cys Pro	Pro Gly Tyr Thr Gly	Leu	Arg Cys Glu
1190		1195		1200
Ala Asp	Ile Asn Glu Cys	Arg	Ser Gly Ala Cys His	Ala Ala His
1205		1210		1215
Thr Arg	Asp Cys Leu Gln	Asp	Pro Gly Gly Gly	Phe Arg Cys Leu
1220		1225		1230
Cys His	Ala Gly Phe Ser	Gly	Pro Arg Cys Gln Thr	Val Leu Ser
1235		1240		1245
Pro Cys	Glu Ser Gln Pro	Cys	Gln His Gly Gly	Gln Cys Arg Pro
1250		1255		1260
Ser Pro	Gly Pro Gly Gly	Gly	Leu Thr Phe Thr	Cys His Cys Ala
1265		1270		1275
Gln Pro	Phe Trp Gly Pro	Arg	Cys Glu Arg Val	Ala Arg Ser Cys
1280		1285		1290
Arg Glu	Leu Gln Cys Pro	Val	Gly Val Pro Cys	Gln Gln Thr Pro
1295		1300		1305
Arg Gly	Pro Arg Cys Ala	Cys	Pro Pro Gly Leu Ser	Gly Pro Ser
1310		1315		1320
Cys Arg	Ser Phe Pro Gly	Ser	Pro Pro Gly Ala	Ser Asn Ala Ser
1325		1330		1335
Cys Ala	Ala Ala Pro Cys	Leu	His Gly Gly Ser	Cys Arg Pro Ala
1340		1345		1350
Pro Leu	Ala Pro Phe Phe	Arg	Cys Ala Cys Ala	Gln Gly Trp Thr
1355		1360		1365
Gly Pro	Arg Cys Glu Ala	Pro	Ala Ala Ala Pro	Glu Val Ser Glu
1370		1375		1380
Glu Pro	Arg Cys Pro Arg	Ala	Ala Cys Gln Ala	Lys Arg Gly Asp
1385		1390		1395
Gln Arg	Cys Asp Arg Glu	Cys	Asn Ser Pro Gly	Cys Gly Trp Asp
1400		1405		1410
Gly Gly	Asp Cys Ser Leu	Ser	Val Gly Asp Pro	Trp Arg Gln Cys
1415		1420		1425
Glu Ala	Leu Gln Cys Trp	Arg	Leu Phe Asn Asn	Ser Arg Cys Asp
1430		1435		1440
Pro Ala	Cys Ser Ser Pro	Ala	Cys Leu Tyr Asp	Asn Phe Asp Cys
1445		1450		1455
His Ala	Gly Gly Arg Glu	Arg	Thr Cys Asn Pro	Val Tyr Glu Lys
1460		1465		1470
Tyr Cys	Ala Asp His Phe	Ala	Asp Gly Arg Cys	Asp Gln Gly Cys
1475		1480		1485
Asn Thr	Glu Glu Cys Gly	Trp	Asp Gly Leu Asp	Cys Ala Ser Glu
1490		1495		1500
Val Pro	Ala Leu Leu Ala	Arg	Gly Val Leu Val	Leu Thr Val Leu
1505		1510		1515
Leu Pro	Pro Glu Glu Leu	Leu	Arg Ser Ser Ala	Asp Phe Leu Gln
1520		1525		1530

-continued

Arg Leu 1535	Ser Ala Ile Leu Arg 1540	Thr Ser Leu Arg Phe 1545	Arg Leu Asp
Ala His 1550	Gly Gln Ala Met Val 1555	Phe Pro Tyr His Arg 1560	Pro Ser Pro
Gly Ser 1565	Glu Pro Arg Ala Arg 1570	Arg Glu Leu Ala Pro 1575	Glu Val Ile
Gly Ser 1580	Val Val Met Leu Glu 1585	Ile Asp Asn Arg Leu 1590	Cys Leu Gln
Ser Pro 1595	Glu Asn Asp His Cys 1600	Phe Pro Asp Ala Gln 1605	Ser Ala Ala
Asp Tyr 1610	Leu Gly Ala Leu Ser 1615	Ala Val Glu Arg Leu 1620	Asp Phe Pro
Tyr Pro 1625	Leu Arg Asp Val Arg 1630	Gly Glu Pro Leu Glu 1635	Pro Pro Glu
Pro Ser 1640	Val Pro Leu Leu Pro 1645	Leu Leu Val Ala Gly 1650	Ala Val Leu
Leu Leu 1655	Val Ile Leu Val Leu 1660	Gly Val Met Val Ala 1665	Arg Arg Lys
Arg Glu 1670	His Ser Thr Leu Trp 1675	Phe Pro Glu Gly Phe 1680	Ser Leu His
Lys Asp 1685	Val Ala Ser Gly His 1690	Lys Gly Arg Arg Glu 1695	Pro Val Gly
Gln Asp 1700	Ala Leu Gly Met Lys 1705	Asn Met Ala Lys Gly 1710	Glu Ser Leu
Met Gly 1715	Glu Val Ala Thr Asp 1720	Trp Met Asp Thr Glu 1725	Cys Pro Glu
Ala Lys 1730	Arg Leu Lys Val Glu 1735	Glu Pro Gly Met Gly 1740	Ala Glu Glu
Ala Val 1745	Asp Cys Arg Gln Trp 1750	Thr Gln His His Leu 1755	Val Ala Ala
Asp Ile 1760	Arg Val Ala Pro Ala 1765	Met Ala Leu Thr Pro 1770	Pro Gln Gly
Asp Ala 1775	Asp Ala Asp Gly Met 1780	Asp Val Asn Val Arg 1785	Gly Pro Asp
Gly Phe 1790	Thr Pro Leu Met Leu 1795	Ala Ser Phe Cys Gly 1800	Gly Ala Leu
Glu Pro 1805	Met Pro Thr Glu Glu 1810	Asp Glu Ala Asp Asp 1815	Thr Ser Ala
Ser Ile 1820	Ile Ser Asp Leu Ile 1825	Cys Gln Gly Ala Gln 1830	Leu Gly Ala
Arg Thr 1835	Asp Arg Thr Gly Glu 1840	Thr Ala Leu His Leu 1845	Ala Ala Arg
Tyr Ala 1850	Arg Ala Asp Ala Ala 1855	Lys Arg Leu Leu Asp 1860	Ala Gly Ala
Asp Thr 1865	Asn Ala Gln Asp His 1870	Ser Gly Arg Thr Pro 1875	Leu His Thr
Ala Val 1880	Thr Ala Asp Ala Gln 1885	Gly Val Phe Gln Ile 1890	Leu Ile Arg
Asn Arg 1895	Ser Thr Asp Leu Asp 1900	Ala Arg Met Ala Asp 1905	Gly Ser Thr
Ala Leu 1910	Ile Leu Ala Ala Arg 1915	Leu Ala Val Glu Gly 1920	Met Val Glu
Glu Leu	Ile Ala Ser His Ala	Asp Val Asn Ala Val	Asp Glu Leu

-continued

1925	1930	1935
Gly Lys Ser Ala Leu His	Trp Ala Ala Ala Val	Asn Asn Val Glu
1940	1945	1950
Ala Thr Leu Ala Leu Leu	Lys Asn Gly Ala Asn	Lys Asp Met Gln
1955	1960	1965
Asp Ser Lys Glu Glu Thr	Pro Leu Phe Leu Ala	Ala Arg Glu Gly
1970	1975	1980
Ser Tyr Glu Ala Ala Lys	Leu Leu Asp His Phe	Ala Asn Arg
1985	1990	1995
Glu Ile Thr Asp His Leu	Asp Arg Leu Pro Arg	Asp Val Ala Gln
2000	2005	2010
Glu Arg Leu His Gln Asp	Ile Val Arg Leu Leu	Asp Gln Pro Ser
2015	2020	2025
Gly Pro Arg Ser Pro Pro	Gly Pro His Gly Leu	Gly Pro Leu Leu
2030	2035	2040
Cys Pro Pro Gly Ala Phe	Leu Pro Gly Leu Lys	Ala Ala Gln Ser
2045	2050	2055
Gly Ser Lys Lys Ser Arg	Arg Pro Pro Gly Lys	Ala Gly Leu Gly
2060	2065	2070
Pro Gln Gly Pro Arg Gly	Arg Gly Lys Lys Leu	Thr Leu Ala Cys
2075	2080	2085
Pro Gly Pro Leu Ala Asp	Ser Ser Val Thr Leu	Ser Pro Val Asp
2090	2095	2100
Ser Leu Asp Ser Pro Arg	Pro Phe Gly Gly Pro	Pro Ala Ser Pro
2105	2110	2115
Gly Gly Phe Pro Leu Glu	Gly Pro Tyr Ala Ala	Ala Thr Ala Thr
2120	2125	2130
Ala Val Ser Leu Ala Gln	Leu Gly Gly Pro Gly	Arg Ala Gly Leu
2135	2140	2145
Gly Arg Gln Pro Pro Gly	Gly Cys Val Leu Ser	Leu Gly Leu Leu
2150	2155	2160
Asn Pro Val Ala Val Pro	Leu Asp Trp Ala Arg	Leu Pro Pro Pro
2165	2170	2175
Ala Pro Pro Gly Pro Ser	Phe Leu Leu Pro Leu	Ala Pro Gly Pro
2180	2185	2190
Gln Leu Leu Asn Pro Gly	Thr Pro Val Ser Pro	Gln Glu Arg Pro
2195	2200	2205
Pro Pro Tyr Leu Ala Val	Pro Gly His Gly Glu	Glu Tyr Pro Val
2210	2215	2220
Ala Gly Ala His Ser Ser	Pro Pro Lys Ala Arg	Phe Leu Arg Val
2225	2230	2235
Pro Ser Glu His Pro Tyr	Leu Thr Pro Ser Pro	Glu Ser Pro Glu
2240	2245	2250
His Trp Ala Ser Pro Ser	Pro Pro Ser Leu Ser	Asp Trp Ser Glu
2255	2260	2265
Ser Thr Pro Ser Pro Ala	Thr Ala Thr Gly Ala	Met Ala Thr Thr
2270	2275	2280
Thr Gly Ala Leu Pro Ala	Gln Pro Leu Pro Leu	Ser Val Pro Ser
2285	2290	2295
Ser Leu Ala Gln Ala Gln	Thr Gln Leu Gly Pro	Gln Pro Glu Val
2300	2305	2310
Thr Pro Lys Arg Gln Val	Leu Ala	
2315	2320	

-continued

<210> SEQ ID NO 2
 <211> LENGTH: 121
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 2

```

Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Lys Pro Gly
1           5           10          15
Thr Ser Val Lys Met Ala Cys Lys Ala Ser Gly Tyr Thr Phe Thr Thr
          20          25          30
His Trp Met Asn Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp
          35          40          45
Ile Gly Thr Ile Asn Pro Ser Asn Asp Phe Thr Asp Cys Asn Gln Lys
          50          55          60
Phe Lys Asp Lys Ala Ile Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala
65          70          75          80
Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Ile Tyr Tyr
          85          90          95
Cys Ala Ser Gly Leu Thr Ala Arg Ala Trp Phe Ala Tyr Trp Gly Gln
          100         105         110
Gly Thr Leu Val Thr Val Ser Ala Ala
          115         120

```

<210> SEQ ID NO 3
 <211> LENGTH: 102
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 3

```

Arg Ala Thr Ile Ser Cys Arg Ala Ser Gln Ser Val Thr Thr Ser Asn
1           5           10          15
Tyr Ser Tyr Met His Trp Phe Gln Gln Lys Pro Gly Gln Pro Pro Lys
          20          25          30
Leu Leu Ile Lys Tyr Ala Ser Asn Leu Asp Ser Gly Val Pro Ala Arg
          35          40          45
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His Pro
          50          55          60
Val Glu Glu Glu Asp Thr Ala Thr Phe Tyr Cys Gln His Ser Trp Glu
65          70          75          80
Ile Pro Tyr Thr Phe Gly Gly Gly Thr Asn Leu Glu Ile Lys Arg Ala
          85          90          95
Asp Ala Ala Pro Thr Val
          100

```

<210> SEQ ID NO 4
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 4

```

Gly Tyr Thr Phe Thr Thr His Trp Met Asn Trp
1           5           10

```

-continued

<210> SEQ ID NO 5
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 5

Ile Asn Pro Ser Asn Asp Phe Thr Asp Cys Asn
1 5 10

<210> SEQ ID NO 6
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 6

Thr Ala Arg Ala Trp Phe Ala Tyr
1 5

<210> SEQ ID NO 7
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 7

Arg Ala Ser Gln Ser Val Thr Thr Ser Asn Tyr Ser Tyr Met His
1 5 10 15

<210> SEQ ID NO 8
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 8

Tyr Ala Ser Asn Leu Asp Ser Gly
1 5

<210> SEQ ID NO 9
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 9

Gln His Ser Trp Glu Ile Pro Tyr Thr
1 5

<210> SEQ ID NO 10
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Glu Pro Arg Cys Pro Arg Ala Ala Cys Gln Ala Lys Arg Gly Asp Gln

-continued

```

1           5           10           15
Arg Cys Asp Arg Glu Cys Asn Ser Pro Gly Cys Gly Trp Asp Gly Gly
      20           25           30

Asp Cys Ser Leu Ser Val Gly
      35

```

```

<210> SEQ ID NO 11
<211> LENGTH: 29
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 11

```

```

Ala Lys Arg Gly Asp Gln Arg Cys Asp Arg Glu Cys Asn Ser Pro Gly
1           5           10           15

Cys Gly Trp Asp Gly Gly Asp Cys Ser Leu Ser Val Gly
      20           25

```

```

<210> SEQ ID NO 12
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

```

```

<400> SEQUENCE: 12

```

```

gctcgagctc gtgggaaaat accgtgggaa aatgaaccgt gggaaaatct cgtgg      55

```

```

<210> SEQ ID NO 13
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

```

```

<400> SEQUENCE: 13

```

```

gctcgagatt ttcccacgag attttccac gggttc      35

```

```

<210> SEQ ID NO 14
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 14

```

```

Glu Glu Ala Cys Glu Leu Pro Glu Cys Gln Ala Lys Arg Gly Asp Gln
1           5           10           15

Arg Cys Asp Arg Glu Cys Asn Ser Pro Gly Cys Gly Trp Asp Gly Gly
      20           25           30

Asp Cys Ser Leu Ser Val Gly
      35

```

```

<210> SEQ ID NO 15
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 15

```

```

Glu Pro Arg Cys Pro Arg Ala Ala Cys Gln Glu Asp Ala Gly Asn Lys

```

-continued

1	5	10	15
Val Cys Ser Arg Glu Cys Asn Ser Pro Gly Cys Gly Trp Asp Gly Gly			
	20	25	30

Asp Cys Ser Leu Ser Val Gly
35

<210> SEQ ID NO 16
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 16

Glu Pro Arg Cys Pro Arg Ala Ala Cys Gln Ala Lys Arg Gly Asp Gln
1 5 10 15

Arg Cys Asp Leu Gln Cys Asn Asn His Ala Cys Gly Trp Asp Gly Gly
20 25 30

Asp Cys Ser Leu Asn Phe Asn
35

<210> SEQ ID NO 17
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 17

Glu Pro Arg Cys Pro Arg Ala Ala Cys Gln Glu Asp Ala Gly Asp Gln
1 5 10 15

Arg Cys Asp Arg Glu Cys Asn Ser Pro Gly Cys Gly Trp Asp Gly Gly
20 25 30

Asp Cys Ser Leu Ser Val Gly
35

<210> SEQ ID NO 18
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 18

Glu Pro Arg Cys Pro Arg Ala Ala Cys Gln Ala Lys Arg Gly Asn Lys
1 5 10 15

Val Cys Asp Arg Glu Cys Asn Ser Pro Gly Cys Gly Trp Asp Gly Gly
20 25 30

Asp Cys Ser Leu Ser Val Gly
35

<210> SEQ ID NO 19
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 19

Glu Pro Arg Cys Pro Arg Ala Ala Cys Gln Ala Lys Arg Gly Asp Gln

-continued

1	5	10	15
Arg Cys Ser Leu Gln Cys Asn Ser Pro Gly Cys Gly Trp Asp Gly Gly			
	20	25	30

Asp Cys Ser Leu Ser Val Gly
35

<210> SEQ ID NO 20
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 20

Glu Pro Arg Cys Pro Arg Ala Ala Cys Gln Ala Lys Arg Gly Asp Gln
1 5 10 15

Arg Cys Asp Arg Glu Cys Asn Asn His Ala Cys Gly Trp Asp Gly Gly
20 25 30

Asp Cys Ser Leu Ser Val Gly
35

<210> SEQ ID NO 21
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 21

Glu Pro Arg Cys Pro Arg Ala Ala Cys Gln Ala Lys Arg Gly Asp Gln
1 5 10 15

Arg Cys Asp Arg Glu Cys Asn Ser Pro Gly Cys Gly Trp Asp Gly Gly
20 25 30

Asp Cys Ser Leu Asn Phe Asn
35

<210> SEQ ID NO 22
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 22

Ala Ala Cys Gln Ala Ala Ala Gly Asp Gln Arg Cys
1 5 10

<210> SEQ ID NO 23
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 23

Ala Cys Gln Ala Lys Ala Ala Asp Gln Arg Cys Asp
1 5 10

<210> SEQ ID NO 24
 <211> LENGTH: 12
 <212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 24

Cys Gln Ala Lys Arg Ala Ala Gln Arg Cys Asp Arg
1 5 10

<210> SEQ ID NO 25
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 25

Gln Ala Lys Arg Gly Ala Ala Arg Cys Asp Arg Glu
1 5 10

<210> SEQ ID NO 26
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 26

Ala Lys Arg Gly Asp Ala Ala Cys Asp Arg Glu Cys
1 5 10

<210> SEQ ID NO 27
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 27

Lys Arg Gly Asp Gln Ala Ala Asp Arg Glu Cys Asn
1 5 10

<210> SEQ ID NO 28
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 28

Arg Gly Asp Gln Arg Ala Ala Arg Glu Cys Asn Ser
1 5 10

<210> SEQ ID NO 29
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 29

Gly Asp Gln Arg Cys Ala Ala Glu Cys Asn Ser Pro
1 5 10

-continued

<210> SEQ ID NO 30
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 30

Asp Gln Arg Cys Asp Ala Ala Cys Asn Ser Pro Gly
1 5 10

<210> SEQ ID NO 31
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 31

Gln Arg Cys Asp Arg Ala Ala Asn Ser Pro Gly Cys
1 5 10

<210> SEQ ID NO 32
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 32

Arg Cys Asp Arg Glu Ala Ala Ser Pro Gly Cys Gly
1 5 10

<210> SEQ ID NO 33
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 33

Cys Asp Arg Glu Cys Ala Ala Pro Gly Cys Gly Trp
1 5 10

<210> SEQ ID NO 34
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 34

Asp Arg Glu Cys Asn Ala Ala Gly Cys Gly Trp Asp
1 5 10

<210> SEQ ID NO 35
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 35

-continued

Arg Glu Cys Asn Ser Ala Ala Cys Gly Trp Asp Gly
1 5 10

<210> SEQ ID NO 36
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 36

Glu Cys Asn Ser Pro Ala Ala Gly Trp Asp Gly Gly
1 5 10

<210> SEQ ID NO 37
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
peptide

<400> SEQUENCE: 37

Cys Asn Ser Pro Gly Ala Ala Trp Asp Gly Gly Asp
1 5 10

<210> SEQ ID NO 38
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
6xHis tag

<400> SEQUENCE: 38

His His His His His His
1 5

<210> SEQ ID NO 39
<211> LENGTH: 2556
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

Met Pro Pro Leu Leu Ala Pro Leu Leu Cys Leu Ala Leu Leu Pro Ala
1 5 10 15

Leu Ala Ala Arg Gly Pro Arg Cys Ser Gln Pro Gly Glu Thr Cys Leu
20 25 30

Asn Gly Gly Lys Cys Glu Ala Ala Asn Gly Thr Glu Ala Cys Val Cys
35 40 45

Gly Gly Ala Phe Val Gly Pro Arg Cys Gln Asp Pro Asn Pro Cys Leu
50 55 60

Ser Thr Pro Cys Lys Asn Ala Gly Thr Cys His Val Val Asp Arg Arg
65 70 75 80

Gly Val Ala Asp Tyr Ala Cys Ser Cys Ala Leu Gly Phe Ser Gly Pro
85 90 95

Leu Cys Leu Thr Pro Leu Asp Asn Ala Cys Leu Thr Asn Pro Cys Arg
100 105 110

Asn Gly Gly Thr Cys Asp Leu Leu Thr Leu Thr Glu Tyr Lys Cys Arg
115 120 125

Cys Pro Pro Gly Trp Ser Gly Lys Ser Cys Gln Gln Ala Asp Pro Cys
130 135 140

-continued

Ala 145	Ser	Asn	Pro	Cys	Ala 150	Asn	Gly	Gly	Gln	Cys 155	Leu	Pro	Phe	Glu	Ala 160
Ser	Tyr	Ile	Cys	His 165	Cys	Pro	Pro	Ser	Phe 170	His	Gly	Pro	Thr	Cys	Arg 175
Gln	Asp	Val	Asn	Glu	Cys	Gly	Gln	Lys 185	Pro	Gly	Leu	Cys	Arg	His	Gly 190
Gly	Thr	Cys	His	Asn	Glu	Val	Gly	Ser	Tyr	Arg	Cys	Val	Cys	Arg	Ala 205
Thr	His	Thr	Gly	Pro	Asn	Cys 215	Glu	Arg	Pro	Tyr	Val	Pro	Cys	Ser	Pro 220
Ser	Pro	Cys	Gln	Asn	Gly 230	Gly	Thr	Cys	Arg	Pro	Thr	Gly	Asp	Val	Thr 240
His	Glu	Cys	Ala	Cys 245	Leu	Pro	Gly	Phe	Thr 250	Gly	Gln	Asn	Cys	Glu	Glu 255
Asn	Ile	Asp	Asp	Cys 260	Pro	Gly	Asn	Asn	Cys 265	Lys	Asn	Gly	Gly	Ala	Cys 270
Val	Asp	Gly	Val	Asn	Thr	Tyr	Asn	Cys	Arg	Cys	Pro	Pro	Glu	Trp	Thr 285
Gly	Gln	Tyr	Cys	Thr	Glu	Asp 295	Val	Asp	Glu	Cys	Gln	Leu	Met	Pro	Asn 300
Ala	Cys	Gln	Asn	Gly 310	Gly	Thr	Cys	His	Asn	Thr 315	His	Gly	Gly	Tyr	Asn 320
Cys	Val	Cys	Val	Asn 325	Gly	Trp	Thr	Gly	Glu	Asp 330	Cys	Ser	Glu	Asn	Ile 335
Asp	Asp	Cys	Ala	Ser 340	Ala	Ala	Cys	Phe	His 345	Gly	Ala	Thr	Cys	His	Asp 350
Arg	Val	Ala	Ser	Phe 355	Tyr	Cys	Glu	Cys	Pro	His 360	Gly	Arg	Thr	Gly	Leu 365
Leu	Cys	His	Leu	Asn 370	Asp	Ala	Cys 375	Ile	Ser	Asn 380	Pro	Cys	Asn	Glu	Gly 385
Ser	Asn	Cys	Asp	Thr 390	Asn	Pro	Val	Asn	Gly	Lys 395	Ala	Ile	Cys	Thr	Cys 400
Pro	Ser	Gly	Tyr	Thr 405	Gly	Pro	Ala	Cys	Ser 410	Gln	Asp	Val	Asp	Glu	Cys 415
Ser	Leu	Gly	Ala	Asn 420	Pro	Cys	Glu	His	Ala 425	Gly	Lys	Cys	Ile	Asn	Thr 430
Leu	Gly	Ser	Phe	Glu 435	Cys	Gln	Cys 440	Leu	Gln	Gly	Tyr	Thr	Gly	Pro	Arg 445
Cys	Glu	Ile	Asp	Val 450	Asn	Glu 455	Cys	Val	Ser	Asn 460	Pro	Cys	Gln	Asn	Asp 465
Ala	Thr	Cys	Leu	Asp 470	Gln	Ile	Gly	Glu	Phe 475	Gln	Cys	Ile	Cys	Met	Pro 480
Gly	Tyr	Glu	Gly	Val 485	His	Cys	Glu	Val	Asn 490	Thr	Asp	Glu	Cys	Ala	Ser 495
Ser	Pro	Cys	Leu	His 500	Asn	Gly	Arg	Cys 505	Leu	Asp	Lys	Ile	Asn	Glu	Phe 510
Gln	Cys	Glu	Cys	Pro 515	Thr	Gly	Phe 520	Thr	Gly	His	Leu	Cys	Gln	Tyr	Asp 525
Val	Asp	Glu	Cys	Ala 530	Ser	Thr 535	Pro	Cys	Lys	Asn 540	Gly	Ala	Lys	Cys	Leu 545
Asp	Gly	Pro	Asn	Thr 550	Tyr	Thr	Cys	Val	Cys 555	Thr	Glu	Gly	Tyr	Thr	Gly 560
Thr	His	Cys	Glu	Val	Asp	Ile	Asp	Glu	Cys	Asp	Pro	Asp	Pro	Cys	His

-continued

565					570					575					
Tyr	Gly	Ser	Cys	Lys	Asp	Gly	Val	Ala	Thr	Phe	Thr	Cys	Leu	Cys	Arg
			580					585					590		
Pro	Gly	Tyr	Thr	Gly	His	His	Cys	Glu	Thr	Asn	Ile	Asn	Glu	Cys	Ser
		595					600					605			
Ser	Gln	Pro	Cys	Arg	His	Gly	Gly	Thr	Cys	Gln	Asp	Arg	Asp	Asn	Ala
	610					615					620				
Tyr	Leu	Cys	Phe	Cys	Leu	Lys	Gly	Thr	Thr	Gly	Pro	Asn	Cys	Glu	Ile
	625				630					635					640
Asn	Leu	Asp	Asp	Cys	Ala	Ser	Ser	Pro	Cys	Asp	Ser	Gly	Thr	Cys	Leu
				645					650					655	
Asp	Lys	Ile	Asp	Gly	Tyr	Glu	Cys	Ala	Cys	Glu	Pro	Gly	Tyr	Thr	Gly
			660					665					670		
Ser	Met	Cys	Asn	Ile	Asn	Ile	Asp	Glu	Cys	Ala	Gly	Asn	Pro	Cys	His
		675					680					685			
Asn	Gly	Gly	Thr	Cys	Glu	Asp	Gly	Ile	Asn	Gly	Phe	Thr	Cys	Arg	Cys
	690					695					700				
Pro	Glu	Gly	Tyr	His	Asp	Pro	Thr	Cys	Leu	Ser	Glu	Val	Asn	Glu	Cys
	705				710					715					720
Asn	Ser	Asn	Pro	Cys	Val	His	Gly	Ala	Cys	Arg	Asp	Ser	Leu	Asn	Gly
				725					730					735	
Tyr	Lys	Cys	Asp	Cys	Asp	Pro	Gly	Trp	Ser	Gly	Thr	Asn	Cys	Asp	Ile
			740					745					750		
Asn	Asn	Asn	Glu	Cys	Glu	Ser	Asn	Pro	Cys	Val	Asn	Gly	Gly	Thr	Cys
		755					760					765			
Lys	Asp	Met	Thr	Ser	Gly	Tyr	Val	Cys	Thr	Cys	Arg	Glu	Gly	Phe	Ser
	770					775					780				
Gly	Pro	Asn	Cys	Gln	Thr	Asn	Ile	Asn	Glu	Cys	Ala	Ser	Asn	Pro	Cys
	785				790					795					800
Leu	Asn	Gln	Gly	Thr	Cys	Ile	Asp	Asp	Val	Ala	Gly	Tyr	Lys	Cys	Asn
				805					810					815	
Cys	Leu	Leu	Pro	Tyr	Thr	Gly	Ala	Thr	Cys	Glu	Val	Val	Leu	Ala	Pro
			820					825					830		
Cys	Ala	Pro	Ser	Pro	Cys	Arg	Asn	Gly	Gly	Glu	Cys	Arg	Gln	Ser	Glu
		835					840					845			
Asp	Tyr	Glu	Ser	Phe	Ser	Cys	Val	Cys	Pro	Thr	Gly	Trp	Gln	Ala	Gly
	850					855					860				
Gln	Thr	Cys	Glu	Val	Asp	Ile	Asn	Glu	Cys	Val	Leu	Ser	Pro	Cys	Arg
	865				870					875					880
His	Gly	Ala	Ser	Cys	Gln	Asn	Thr	His	Gly	Gly	Tyr	Arg	Cys	His	Cys
			885						890					895	
Gln	Ala	Gly	Tyr	Ser	Gly	Arg	Asn	Cys	Glu	Thr	Asp	Ile	Asp	Asp	Cys
			900					905					910		
Arg	Pro	Asn	Pro	Cys	His	Asn	Gly	Gly	Ser	Cys	Thr	Asp	Gly	Ile	Asn
		915					920					925			
Thr	Ala	Phe	Cys	Asp	Cys	Leu	Pro	Gly	Phe	Arg	Gly	Thr	Phe	Cys	Glu
	930					935					940				
Glu	Asp	Ile	Asn	Glu	Cys	Ala	Ser	Asp	Pro	Cys	Arg	Asn	Gly	Ala	Asn
	945				950					955					960
Cys	Thr	Asp	Cys	Val	Asp	Ser	Tyr	Thr	Cys	Thr	Cys	Pro	Ala	Gly	Phe
			965						970					975	
Ser	Gly	Ile	His	Cys	Glu	Asn	Asn	Thr	Pro	Asp	Cys	Thr	Glu	Ser	Ser
			980					985					990		

-continued

Cys	Phe	Asn	Gly	Gly	Thr	Cys	Val	Asp	Gly	Ile	Asn	Ser	Phe	Thr	Cys
	995						1000					1005			
Leu	Cys	Pro	Pro	Gly	Phe	Thr	Gly	Ser	Tyr	Cys	Gln	His	Asp	Val	
1010						1015					1020				
Asn	Glu	Cys	Asp	Ser	Gln	Pro	Cys	Leu	His	Gly	Gly	Thr	Cys	Gln	
1025						1030					1035				
Asp	Gly	Cys	Gly	Ser	Tyr	Arg	Cys	Thr	Cys	Pro	Gln	Gly	Tyr	Thr	
1040						1045					1050				
Gly	Pro	Asn	Cys	Gln	Asn	Leu	Val	His	Trp	Cys	Asp	Ser	Ser	Pro	
1055						1060					1065				
Cys	Lys	Asn	Gly	Gly	Lys	Cys	Trp	Gln	Thr	His	Thr	Gln	Tyr	Arg	
1070						1075					1080				
Cys	Glu	Cys	Pro	Ser	Gly	Trp	Thr	Gly	Leu	Tyr	Cys	Asp	Val	Pro	
1085						1090					1095				
Ser	Val	Ser	Cys	Glu	Val	Ala	Ala	Gln	Arg	Gln	Gly	Val	Asp	Val	
1100						1105					1110				
Ala	Arg	Leu	Cys	Gln	His	Gly	Gly	Leu	Cys	Val	Asp	Ala	Gly	Asn	
1115						1120					1125				
Thr	His	His	Cys	Arg	Cys	Gln	Ala	Gly	Tyr	Thr	Gly	Ser	Tyr	Cys	
1130						1135					1140				
Glu	Asp	Leu	Val	Asp	Glu	Cys	Ser	Pro	Ser	Pro	Cys	Gln	Asn	Gly	
1145						1150					1155				
Ala	Thr	Cys	Thr	Asp	Tyr	Leu	Gly	Gly	Tyr	Ser	Cys	Lys	Cys	Val	
1160						1165					1170				
Ala	Gly	Tyr	His	Gly	Val	Asn	Cys	Ser	Glu	Glu	Ile	Asp	Glu	Cys	
1175						1180					1185				
Leu	Ser	His	Pro	Cys	Gln	Asn	Gly	Gly	Thr	Cys	Leu	Asp	Leu	Pro	
1190						1195					1200				
Asn	Thr	Tyr	Lys	Cys	Ser	Cys	Pro	Arg	Gly	Thr	Gln	Gly	Val	His	
1205						1210					1215				
Cys	Glu	Ile	Asn	Val	Asp	Asp	Cys	Asn	Pro	Pro	Val	Asp	Pro	Val	
1220						1225					1230				
Ser	Arg	Ser	Pro	Lys	Cys	Phe	Asn	Asn	Gly	Thr	Cys	Val	Asp	Gln	
1235						1240					1245				
Val	Gly	Gly	Tyr	Ser	Cys	Thr	Cys	Pro	Pro	Gly	Phe	Val	Gly	Glu	
1250						1255					1260				
Arg	Cys	Glu	Gly	Asp	Val	Asn	Glu	Cys	Leu	Ser	Asn	Pro	Cys	Asp	
1265						1270					1275				
Ala	Arg	Gly	Thr	Gln	Asn	Cys	Val	Gln	Arg	Val	Asn	Asp	Phe	His	
1280						1285					1290				
Cys	Glu	Cys	Arg	Ala	Gly	His	Thr	Gly	Arg	Arg	Cys	Glu	Ser	Val	
1295						1300					1305				
Ile	Asn	Gly	Cys	Lys	Gly	Lys	Pro	Cys	Lys	Asn	Gly	Gly	Thr	Cys	
1310						1315					1320				
Ala	Val	Ala	Ser	Asn	Thr	Ala	Arg	Gly	Phe	Ile	Cys	Lys	Cys	Pro	
1325						1330					1335				
Ala	Gly	Phe	Glu	Gly	Ala	Thr	Cys	Glu	Asn	Asp	Ala	Arg	Thr	Cys	
1340						1345					1350				
Gly	Ser	Leu	Arg	Cys	Leu	Asn	Gly	Gly	Thr	Cys	Ile	Ser	Gly	Pro	
1355						1360					1365				
Arg	Ser	Pro	Thr	Cys	Leu	Cys	Leu	Gly	Pro	Phe	Thr	Gly	Pro	Glu	
1370						1375					1380				
Cys	Gln	Phe	Pro	Ala	Ser	Ser	Pro	Cys	Leu	Gly	Gly	Asn	Pro	Cys	
1385						1390					1395				

-continued

Tyr	Asn	Gln	Gly	Thr	Cys	Glu	Pro	Thr	Ser	Glu	Ser	Pro	Phe	Tyr
1400						1405					1410			
Arg	Cys	Leu	Cys	Pro	Ala	Lys	Phe	Asn	Gly	Leu	Leu	Cys	His	Ile
1415						1420					1425			
Leu	Asp	Tyr	Ser	Phe	Gly	Gly	Gly	Ala	Gly	Arg	Asp	Ile	Pro	Pro
1430						1435					1440			
Pro	Leu	Ile	Glu	Glu	Ala	Cys	Glu	Leu	Pro	Glu	Cys	Gln	Glu	Asp
1445						1450					1455			
Ala	Gly	Asn	Lys	Val	Cys	Ser	Leu	Gln	Cys	Asn	Asn	His	Ala	Cys
1460						1465					1470			
Gly	Trp	Asp	Gly	Gly	Asp	Cys	Ser	Leu	Asn	Phe	Asn	Asp	Pro	Trp
1475						1480					1485			
Lys	Asn	Cys	Thr	Gln	Ser	Leu	Gln	Cys	Trp	Lys	Tyr	Phe	Ser	Asp
1490						1495					1500			
Gly	His	Cys	Asp	Ser	Gln	Cys	Asn	Ser	Ala	Gly	Cys	Leu	Phe	Asp
1505						1510					1515			
Gly	Phe	Asp	Cys	Gln	Arg	Ala	Glu	Gly	Gln	Cys	Asn	Pro	Leu	Tyr
1520						1525					1530			
Asp	Gln	Tyr	Cys	Lys	Asp	His	Phe	Ser	Asp	Gly	His	Cys	Asp	Gln
1535						1540					1545			
Gly	Cys	Asn	Ser	Ala	Glu	Cys	Glu	Trp	Asp	Gly	Leu	Asp	Cys	Ala
1550						1555					1560			
Glu	His	Val	Pro	Glu	Arg	Leu	Ala	Ala	Gly	Thr	Leu	Val	Val	Val
1565						1570					1575			
Val	Leu	Met	Pro	Pro	Glu	Gln	Leu	Arg	Asn	Ser	Ser	Phe	His	Phe
1580						1585					1590			
Leu	Arg	Glu	Leu	Ser	Arg	Val	Leu	His	Thr	Asn	Val	Val	Phe	Lys
1595						1600					1605			
Arg	Asp	Ala	His	Gly	Gln	Gln	Met	Ile	Phe	Pro	Tyr	Tyr	Gly	Arg
1610						1615					1620			
Glu	Glu	Glu	Leu	Arg	Lys	His	Pro	Ile	Lys	Arg	Ala	Ala	Glu	Gly
1625						1630					1635			
Trp	Ala	Ala	Pro	Asp	Ala	Leu	Leu	Gly	Gln	Val	Lys	Ala	Ser	Leu
1640						1645					1650			
Leu	Pro	Gly	Gly	Ser	Glu	Gly	Gly	Arg	Arg	Arg	Arg	Glu	Leu	Asp
1655						1660					1665			
Pro	Met	Asp	Val	Arg	Gly	Ser	Ile	Val	Tyr	Leu	Glu	Ile	Asp	Asn
1670						1675					1680			
Arg	Gln	Cys	Val	Gln	Ala	Ser	Ser	Gln	Cys	Phe	Gln	Ser	Ala	Thr
1685						1690					1695			
Asp	Val	Ala	Ala	Phe	Leu	Gly	Ala	Leu	Ala	Ser	Leu	Gly	Ser	Leu
1700						1705					1710			
Asn	Ile	Pro	Tyr	Lys	Ile	Glu	Ala	Val	Gln	Ser	Glu	Thr	Val	Glu
1715						1720					1725			
Pro	Pro	Pro	Pro	Ala	Gln	Leu	His	Phe	Met	Tyr	Val	Ala	Ala	Ala
1730						1735					1740			
Ala	Phe	Val	Leu	Leu	Phe	Phe	Val	Gly	Cys	Gly	Val	Leu	Leu	Ser
1745						1750					1755			
Arg	Lys	Arg	Arg	Arg	Gln	His	Gly	Gln	Leu	Trp	Phe	Pro	Glu	Gly
1760						1765					1770			
Phe	Lys	Val	Ser	Glu	Ala	Ser	Lys	Lys	Lys	Arg	Arg	Glu	Pro	Leu
1775						1780					1785			
Gly	Glu	Asp	Ser	Val	Gly	Leu	Lys	Pro	Leu	Lys	Asn	Ala	Ser	Asp

-continued

1790	1795	1800
Gly Ala Leu Met Asp Asp Asn Gln Asn Glu Trp Gly Asp Glu Asp 1805 1810 1815		
Leu Glu Thr Lys Lys Phe Arg Phe Glu Glu Pro Val Val Leu Pro 1820 1825 1830		
Asp Leu Asp Asp Gln Thr Asp His Arg Gln Trp Thr Gln Gln His 1835 1840 1845		
Leu Asp Ala Ala Asp Leu Arg Met Ser Ala Met Ala Pro Thr Pro 1850 1855 1860		
Pro Gln Gly Glu Val Asp Ala Asp Cys Met Asp Val Asn Val Arg 1865 1870 1875		
Gly Pro Asp Gly Phe Thr Pro Leu Met Ile Ala Ser Cys Ser Gly 1880 1885 1890		
Gly Gly Leu Glu Thr Gly Asn Ser Glu Glu Glu Glu Asp Ala Pro 1895 1900 1905		
Ala Val Ile Ser Asp Phe Ile Tyr Gln Gly Ala Ser Leu His Asn 1910 1915 1920		
Gln Thr Asp Arg Thr Gly Glu Thr Ala Leu His Leu Ala Ala Arg 1925 1930 1935		
Tyr Ser Arg Ser Asp Ala Ala Lys Arg Leu Leu Glu Ala Ser Ala 1940 1945 1950		
Asp Ala Asn Ile Gln Asp Asn Met Gly Arg Thr Pro Leu His Ala 1955 1960 1965		
Ala Val Ser Ala Asp Ala Gln Gly Val Phe Gln Ile Leu Ile Arg 1970 1975 1980		
Asn Arg Ala Thr Asp Leu Asp Ala Arg Met His Asp Gly Thr Thr 1985 1990 1995		
Pro Leu Ile Leu Ala Ala Arg Leu Ala Val Glu Gly Met Leu Glu 2000 2005 2010		
Asp Leu Ile Asn Ser His Ala Asp Val Asn Ala Val Asp Asp Leu 2015 2020 2025		
Gly Lys Ser Ala Leu His Trp Ala Ala Ala Val Asn Asn Val Asp 2030 2035 2040		
Ala Ala Val Val Leu Leu Lys Asn Gly Ala Asn Lys Asp Met Gln 2045 2050 2055		
Asn Asn Arg Glu Glu Thr Pro Leu Phe Leu Ala Ala Arg Glu Gly 2060 2065 2070		
Ser Tyr Glu Thr Ala Lys Val Leu Leu Asp His Phe Ala Asn Arg 2075 2080 2085		
Asp Ile Thr Asp His Met Asp Arg Leu Pro Arg Asp Ile Ala Gln 2090 2095 2100		
Glu Arg Met His His Asp Ile Val Arg Leu Leu Asp Glu Tyr Asn 2105 2110 2115		
Leu Val Arg Ser Pro Gln Leu His Gly Ala Pro Leu Gly Gly Thr 2120 2125 2130		
Pro Thr Leu Ser Pro Pro Leu Cys Ser Pro Asn Gly Tyr Leu Gly 2135 2140 2145		
Ser Leu Lys Pro Gly Val Gln Gly Lys Lys Val Arg Lys Pro Ser 2150 2155 2160		
Ser Lys Gly Leu Ala Cys Gly Ser Lys Glu Ala Lys Asp Leu Lys 2165 2170 2175		
Ala Arg Arg Lys Lys Ser Gln Asp Gly Lys Gly Cys Leu Leu Asp 2180 2185 2190		

-continued

Ser	Ser	Gly	Met	Leu	Ser	Pro	Val	Asp	Ser	Leu	Glu	Ser	Pro	His
2195						2200					2205			
Gly	Tyr	Leu	Ser	Asp	Val	Ala	Ser	Pro	Pro	Leu	Leu	Pro	Ser	Pro
2210						2215					2220			
Phe	Gln	Gln	Ser	Pro	Ser	Val	Pro	Leu	Asn	His	Leu	Pro	Gly	Met
2225						2230					2235			
Pro	Asp	Thr	His	Leu	Gly	Ile	Gly	His	Leu	Asn	Val	Ala	Ala	Lys
2240						2245					2250			
Pro	Glu	Met	Ala	Ala	Leu	Gly	Gly	Gly	Gly	Arg	Leu	Ala	Phe	Glu
2255						2260					2265			
Thr	Gly	Pro	Pro	Arg	Leu	Ser	His	Leu	Pro	Val	Ala	Ser	Gly	Thr
2270						2275					2280			
Ser	Thr	Val	Leu	Gly	Ser	Ser	Ser	Gly	Gly	Ala	Leu	Asn	Phe	Thr
2285						2290					2295			
Val	Gly	Gly	Ser	Thr	Ser	Leu	Asn	Gly	Gln	Cys	Glu	Trp	Leu	Ser
2300						2305					2310			
Arg	Leu	Gln	Ser	Gly	Met	Val	Pro	Asn	Gln	Tyr	Asn	Pro	Leu	Arg
2315						2320					2325			
Gly	Ser	Val	Ala	Pro	Gly	Pro	Leu	Ser	Thr	Gln	Ala	Pro	Ser	Leu
2330						2335					2340			
Gln	His	Gly	Met	Val	Gly	Pro	Leu	His	Ser	Ser	Leu	Ala	Ala	Ser
2345						2350					2355			
Ala	Leu	Ser	Gln	Met	Met	Ser	Tyr	Gln	Gly	Leu	Pro	Ser	Thr	Arg
2360						2365					2370			
Leu	Ala	Thr	Gln	Pro	His	Leu	Val	Gln	Thr	Gln	Gln	Val	Gln	Pro
2375						2380					2385			
Gln	Asn	Leu	Gln	Met	Gln	Gln	Gln	Asn	Leu	Gln	Pro	Ala	Asn	Ile
2390						2395					2400			
Gln	Gln	Gln	Gln	Ser	Leu	Gln	Pro	Pro	Pro	Pro	Pro	Pro	Gln	Pro
2405						2410					2415			
His	Leu	Gly	Val	Ser	Ser	Ala	Ala	Ser	Gly	His	Leu	Gly	Arg	Ser
2420						2425					2430			
Phe	Leu	Ser	Gly	Glu	Pro	Ser	Gln	Ala	Asp	Val	Gln	Pro	Leu	Gly
2435						2440					2445			
Pro	Ser	Ser	Leu	Ala	Val	His	Thr	Ile	Leu	Pro	Gln	Glu	Ser	Pro
2450						2455					2460			
Ala	Leu	Pro	Thr	Ser	Leu	Pro	Ser	Ser	Leu	Val	Pro	Pro	Val	Thr
2465						2470					2475			
Ala	Ala	Gln	Phe	Leu	Thr	Pro	Pro	Ser	Gln	His	Ser	Tyr	Ser	Ser
2480						2485					2490			
Pro	Val	Asp	Asn	Thr	Pro	Ser	His	Gln	Leu	Gln	Val	Pro	Glu	His
2495						2500					2505			
Pro	Phe	Leu	Thr	Pro	Ser	Pro	Glu	Ser	Pro	Asp	Gln	Trp	Ser	Ser
2510						2515					2520			
Ser	Ser	Pro	His	Ser	Asn	Val	Ser	Asp	Trp	Ser	Glu	Gly	Val	Ser
2525						2530					2535			
Ser	Pro	Pro	Thr	Ser	Met	Gln	Ser	Gln	Ile	Ala	Arg	Ile	Pro	Glu
2540						2545					2550			
Ala	Phe	Lys												
2555														

<210> SEQ ID NO 40

<211> LENGTH: 2471

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 40

Met Pro Ala Leu Arg Pro Ala Leu Leu Trp Ala Leu Leu Ala Leu Trp
 1 5 10 15
 Leu Cys Cys Ala Ala Pro Ala His Ala Leu Gln Cys Arg Asp Gly Tyr
 20 25 30
 Glu Pro Cys Val Asn Glu Gly Met Cys Val Thr Tyr His Asn Gly Thr
 35 40 45
 Gly Tyr Cys Lys Cys Pro Glu Gly Phe Leu Gly Glu Tyr Cys Gln His
 50 55 60
 Arg Asp Pro Cys Glu Lys Asn Arg Cys Gln Asn Gly Gly Thr Cys Val
 65 70 75 80
 Ala Gln Ala Met Leu Gly Lys Ala Thr Cys Arg Cys Ala Ser Gly Phe
 85 90 95
 Thr Gly Glu Asp Cys Gln Tyr Ser Thr Ser His Pro Cys Phe Val Ser
 100 105 110
 Arg Pro Cys Leu Asn Gly Gly Thr Cys His Met Leu Ser Arg Asp Thr
 115 120 125
 Tyr Glu Cys Thr Cys Gln Val Gly Phe Thr Gly Lys Glu Cys Gln Trp
 130 135 140
 Thr Asp Ala Cys Leu Ser His Pro Cys Ala Asn Gly Ser Thr Cys Thr
 145 150 155 160
 Thr Val Ala Asn Gln Phe Ser Cys Lys Cys Leu Thr Gly Phe Thr Gly
 165 170 175
 Gln Lys Cys Glu Thr Asp Val Asn Glu Cys Asp Ile Pro Gly His Cys
 180 185 190
 Gln His Gly Gly Thr Cys Leu Asn Leu Pro Gly Ser Tyr Gln Cys Gln
 195 200 205
 Cys Pro Gln Gly Phe Thr Gly Gln Tyr Cys Asp Ser Leu Tyr Val Pro
 210 215 220
 Cys Ala Pro Ser Pro Cys Val Asn Gly Gly Thr Cys Arg Gln Thr Gly
 225 230 235 240
 Asp Phe Thr Phe Glu Cys Asn Cys Leu Pro Gly Phe Glu Gly Ser Thr
 245 250 255
 Cys Glu Arg Asn Ile Asp Asp Cys Pro Asn His Arg Cys Gln Asn Gly
 260 265 270
 Gly Val Cys Val Asp Gly Val Asn Thr Tyr Asn Cys Arg Cys Pro Pro
 275 280 285
 Gln Trp Thr Gly Gln Phe Cys Thr Glu Asp Val Asp Glu Cys Leu Leu
 290 295 300
 Gln Pro Asn Ala Cys Gln Asn Gly Gly Thr Cys Ala Asn Arg Asn Gly
 305 310 315 320
 Gly Tyr Gly Cys Val Cys Val Asn Gly Trp Ser Gly Asp Asp Cys Ser
 325 330 335
 Glu Asn Ile Asp Asp Cys Ala Phe Ala Ser Cys Thr Pro Gly Ser Thr
 340 345 350
 Cys Ile Asp Arg Val Ala Ser Phe Ser Cys Met Cys Pro Glu Gly Lys
 355 360 365
 Ala Gly Leu Leu Cys His Leu Asp Asp Ala Cys Ile Ser Asn Pro Cys
 370 375 380
 His Lys Gly Ala Leu Cys Asp Thr Asn Pro Leu Asn Gly Gln Tyr Ile
 385 390 395 400
 Cys Thr Cys Pro Gln Gly Tyr Lys Gly Ala Asp Cys Thr Glu Asp Val
 405 410 415

-continued

Asp Glu Cys Ala Met Ala Asn Ser Asn Pro Cys Glu His Ala Gly Lys
 420 425 430
 Cys Val Asn Thr Asp Gly Ala Phe His Cys Glu Cys Leu Lys Gly Tyr
 435 440 445
 Ala Gly Pro Arg Cys Glu Met Asp Ile Asn Glu Cys His Ser Asp Pro
 450 455 460
 Cys Gln Asn Asp Ala Thr Cys Leu Asp Lys Ile Gly Gly Phe Thr Cys
 465 470 475 480
 Leu Cys Met Pro Gly Phe Lys Gly Val His Cys Glu Leu Glu Ile Asn
 485 490 495
 Glu Cys Gln Ser Asn Pro Cys Val Asn Asn Gly Gln Cys Val Asp Lys
 500 505 510
 Val Asn Arg Phe Gln Cys Leu Cys Pro Pro Gly Phe Thr Gly Pro Val
 515 520 525
 Cys Gln Ile Asp Ile Asp Asp Cys Ser Ser Thr Pro Cys Leu Asn Gly
 530 535 540
 Ala Lys Cys Ile Asp His Pro Asn Gly Tyr Glu Cys Gln Cys Ala Thr
 545 550 555 560
 Gly Phe Thr Gly Val Leu Cys Glu Glu Asn Ile Asp Asn Cys Asp Pro
 565 570 575
 Asp Pro Cys His His Gly Gln Cys Gln Asp Gly Ile Asp Ser Tyr Thr
 580 585 590
 Cys Ile Cys Asn Pro Gly Tyr Met Gly Ala Ile Cys Ser Asp Gln Ile
 595 600 605
 Asp Glu Cys Tyr Ser Ser Pro Cys Leu Asn Asp Gly Arg Cys Ile Asp
 610 615 620
 Leu Val Asn Gly Tyr Gln Cys Asn Cys Gln Pro Gly Thr Ser Gly Val
 625 630 635 640
 Asn Cys Glu Ile Asn Phe Asp Asp Cys Ala Ser Asn Pro Cys Ile His
 645 650 655
 Gly Ile Cys Met Asp Gly Ile Asn Arg Tyr Ser Cys Val Cys Ser Pro
 660 665 670
 Gly Phe Thr Gly Gln Arg Cys Asn Ile Asp Ile Asp Glu Cys Ala Ser
 675 680 685
 Asn Pro Cys Arg Lys Gly Ala Thr Cys Ile Asn Gly Val Asn Gly Phe
 690 695 700
 Arg Cys Ile Cys Pro Glu Gly Pro His His Pro Ser Cys Tyr Ser Gln
 705 710 715 720
 Val Asn Glu Cys Leu Ser Asn Pro Cys Ile His Gly Asn Cys Thr Gly
 725 730 735
 Gly Leu Ser Gly Tyr Lys Cys Leu Cys Asp Ala Gly Trp Val Gly Ile
 740 745 750
 Asn Cys Glu Val Asp Lys Asn Glu Cys Leu Ser Asn Pro Cys Gln Asn
 755 760 765
 Gly Gly Thr Cys Asp Asn Leu Val Asn Gly Tyr Arg Cys Thr Cys Lys
 770 775 780
 Lys Gly Phe Lys Gly Tyr Asn Cys Gln Val Asn Ile Asp Glu Cys Ala
 785 790 795 800
 Ser Asn Pro Cys Leu Asn Gln Gly Thr Cys Phe Asp Asp Ile Ser Gly
 805 810 815
 Tyr Thr Cys His Cys Val Leu Pro Tyr Thr Gly Lys Asn Cys Gln Thr
 820 825 830
 Val Leu Ala Pro Cys Ser Pro Asn Pro Cys Glu Asn Ala Ala Val Cys

-continued

835					840					845					
Lys	Glu	Ser	Pro	Asn	Phe	Glu	Ser	Tyr	Thr	Cys	Leu	Cys	Ala	Pro	Gly
850						855					860				
Trp	Gln	Gly	Gln	Arg	Cys	Thr	Ile	Asp	Ile	Asp	Glu	Cys	Ile	Ser	Lys
865					870					875					880
Pro	Cys	Met	Asn	His	Gly	Leu	Cys	His	Asn	Thr	Gln	Gly	Ser	Tyr	Met
				885						890				895	
Cys	Glu	Cys	Pro	Pro	Gly	Phe	Ser	Gly	Met	Asp	Cys	Glu	Glu	Asp	Ile
			900					905					910		
Asp	Asp	Cys	Leu	Ala	Asn	Pro	Cys	Gln	Asn	Gly	Gly	Ser	Cys	Met	Asp
	915					920						925			
Gly	Val	Asn	Thr	Phe	Ser	Cys	Leu	Cys	Leu	Pro	Gly	Phe	Thr	Gly	Asp
	930					935					940				
Lys	Cys	Gln	Thr	Asp	Met	Asn	Glu	Cys	Leu	Ser	Glu	Pro	Cys	Lys	Asn
945					950					955					960
Gly	Gly	Thr	Cys	Ser	Asp	Tyr	Val	Asn	Ser	Tyr	Thr	Cys	Lys	Cys	Gln
				965					970					975	
Ala	Gly	Phe	Asp	Gly	Val	His	Cys	Glu	Asn	Asn	Ile	Asn	Glu	Cys	Thr
		980						985					990		
Glu	Ser	Ser	Cys	Phe	Asn	Gly	Gly	Thr	Cys	Val	Asp	Gly	Ile	Asn	Ser
	995					1000						1005			
Phe	Ser	Cys	Leu	Cys	Pro	Val	Gly	Phe	Thr	Gly	Ser	Phe	Cys	Leu	
	1010					1015						1020			
His	Glu	Ile	Asn	Glu	Cys	Ser	Ser	His	Pro	Cys	Leu	Asn	Glu	Gly	
	1025					1030						1035			
Thr	Cys	Val	Asp	Gly	Leu	Gly	Thr	Tyr	Arg	Cys	Ser	Cys	Pro	Leu	
	1040					1045						1050			
Gly	Tyr	Thr	Gly	Lys	Asn	Cys	Gln	Thr	Leu	Val	Asn	Leu	Cys	Ser	
	1055					1060						1065			
Arg	Ser	Pro	Cys	Lys	Asn	Lys	Gly	Thr	Cys	Val	Gln	Lys	Lys	Ala	
	1070					1075						1080			
Glu	Ser	Gln	Cys	Leu	Cys	Pro	Ser	Gly	Trp	Ala	Gly	Ala	Tyr	Cys	
	1085					1090						1095			
Asp	Val	Pro	Asn	Val	Ser	Cys	Asp	Ile	Ala	Ala	Ser	Arg	Arg	Gly	
	1100					1105						1110			
Val	Leu	Val	Glu	His	Leu	Cys	Gln	His	Ser	Gly	Val	Cys	Ile	Asn	
	1115					1120						1125			
Ala	Gly	Asn	Thr	His	Tyr	Cys	Gln	Cys	Pro	Leu	Gly	Tyr	Thr	Gly	
	1130					1135						1140			
Ser	Tyr	Cys	Glu	Glu	Gln	Leu	Asp	Glu	Cys	Ala	Ser	Asn	Pro	Cys	
	1145					1150						1155			
Gln	His	Gly	Ala	Thr	Cys	Ser	Asp	Phe	Ile	Gly	Gly	Tyr	Arg	Cys	
	1160					1165						1170			
Glu	Cys	Val	Pro	Gly	Tyr	Gln	Gly	Val	Asn	Cys	Glu	Tyr	Glu	Val	
	1175					1180						1185			
Asp	Glu	Cys	Gln	Asn	Gln	Pro	Cys	Gln	Asn	Gly	Gly	Thr	Cys	Ile	
	1190					1195						1200			
Asp	Leu	Val	Asn	His	Phe	Lys	Cys	Ser	Cys	Pro	Pro	Gly	Thr	Arg	
	1205					1210						1215			
Gly	Leu	Leu	Cys	Glu	Glu	Asn	Ile	Asp	Asp	Cys	Ala	Arg	Gly	Pro	
	1220					1225						1230			
His	Cys	Leu	Asn	Gly	Gly	Gln	Cys	Met	Asp	Arg	Ile	Gly	Gly	Tyr	
	1235					1240						1245			

-continued

Ser Cys	Arg Cys	Leu Pro	Gly	Phe Ala	Gly Glu	Arg Cys	Glu Gly
1250			1255			1260	
Asp Ile	Asn Glu	Cys Leu	Ser	Asn Pro	Cys Ser	Ser	Glu Gly Ser
1265			1270			1275	
Leu Asp	Cys Ile	Gln Leu	Thr	Asn Asp	Tyr Leu	Cys	Val Cys Arg
1280			1285			1290	
Ser Ala	Phe Thr	Gly Arg	His	Cys Glu	Thr Phe	Val	Asp Val Cys
1295			1300			1305	
Pro Gln	Met Pro	Cys Leu	Asn	Gly Gly	Thr Cys	Ala	Val Ala Ser
1310			1315			1320	
Asn Met	Pro Asp	Gly Phe	Ile	Cys Arg	Cys Pro	Pro	Gly Phe Ser
1325			1330			1335	
Gly Ala	Arg Cys	Gln Ser	Ser	Cys Gly	Gln Val	Lys	Cys Arg Lys
1340			1345			1350	
Gly Glu	Gln Cys	Val His	Thr	Ala Ser	Gly Pro	Arg	Cys Phe Cys
1355			1360			1365	
Pro Ser	Pro Arg	Asp Cys	Glu	Ser Gly	Cys Ala	Ser	Ser Pro Cys
1370			1375			1380	
Gln His	Gly Gly	Ser Cys	His	Pro Gln	Arg Gln	Pro	Pro Tyr Tyr
1385			1390			1395	
Ser Cys	Gln Cys	Ala Pro	Pro	Phe Ser	Gly Ser	Arg	Cys Glu Leu
1400			1405			1410	
Tyr Thr	Ala Pro	Pro Ser	Thr	Pro Pro	Ala Thr	Cys	Leu Ser Gln
1415			1420			1425	
Tyr Cys	Ala Asp	Lys Ala	Arg	Asp Gly	Val Cys	Asp	Glu Ala Cys
1430			1435			1440	
Asn Ser	His Ala	Cys Gln	Trp	Asp Gly	Gly Asp	Cys	Ser Leu Thr
1445			1450			1455	
Met Glu	Asn Pro	Trp Ala	Asn	Cys Ser	Ser Pro	Leu	Pro Cys Trp
1460			1465			1470	
Asp Tyr	Ile Asn	Asn Gln	Cys	Asp Glu	Leu Cys	Asn	Thr Val Glu
1475			1480			1485	
Cys Leu	Phe Asp	Asn Phe	Glu	Cys Gln	Gly Asn	Ser	Lys Thr Cys
1490			1495			1500	
Lys Tyr	Asp Lys	Tyr Cys	Ala	Asp His	Phe Lys	Asp	Asn His Cys
1505			1510			1515	
Asp Gln	Gly Cys	Asn Ser	Glu	Glu Cys	Gly Trp	Asp	Gly Leu Asp
1520			1525			1530	
Cys Ala	Ala Asp	Gln Pro	Glu	Asn Leu	Ala Glu	Gly	Thr Leu Val
1535			1540			1545	
Ile Val	Val Leu	Met Pro	Pro	Glu Gln	Leu Leu	Gln	Asp Ala Arg
1550			1555			1560	
Ser Phe	Leu Arg	Ala Leu	Gly	Thr Leu	Leu His	Thr	Asn Leu Arg
1565			1570			1575	
Ile Lys	Arg Asp	Ser Gln	Gly	Glu Leu	Met Val	Tyr	Pro Tyr Tyr
1580			1585			1590	
Gly Glu	Lys Ser	Ala Ala	Met	Lys Lys	Gln Arg	Met	Thr Arg Arg
1595			1600			1605	
Ser Leu	Pro Gly	Glu Gln	Glu	Gln Glu	Val Ala	Gly	Ser Lys Val
1610			1615			1620	
Phe Leu	Glu Ile	Asp Asn	Arg	Gln Cys	Val Gln	Asp	Ser Asp His
1625			1630			1635	
Cys Phe	Lys Asn	Thr Asp	Ala	Ala Ala	Ala Leu	Leu	Ala Ser His
1640			1645			1650	

-continued

Ala Ile	Gln Gly	Thr Leu	Ser Tyr	Pro Leu	Val Ser	Val Val	Ser
1655			1660		1665		
Glu Ser	Leu Thr	Pro Glu	Arg Thr	Gln Leu	Leu Tyr	Leu Leu	Ala
1670			1675		1680		
Val Ala	Val Val	Ile Ile	Leu Phe	Ile Ile	Leu Leu	Gly Val	Ile
1685			1690		1695		
Met Ala	Lys Arg	Lys Arg	Lys His	Gly Ser	Leu Trp	Leu Pro	Glu
1700			1705		1710		
Gly Phe	Thr Leu	Arg Arg	Asp Ala	Ser Asn	His Lys	Arg Arg	Glu
1715			1720		1725		
Pro Val	Gly Gln	Asp Ala	Val Gly	Leu Lys	Asn Leu	Ser Val	Gln
1730			1735		1740		
Val Ser	Glu Ala	Asn Leu	Ile Gly	Thr Gly	Thr Ser	Glu His	Trp
1745			1750		1755		
Val Asp	Asp Glu	Gly Pro	Gln Pro	Lys Lys	Val Lys	Ala Glu	Asp
1760			1765		1770		
Glu Ala	Leu Leu	Ser Glu	Glu Asp	Asp Pro	Ile Asp	Arg Arg	Pro
1775			1780		1785		
Trp Thr	Gln Gln	His Leu	Glu Ala	Ala Asp	Ile Arg	Arg Thr	Pro
1790			1795		1800		
Ser Leu	Ala Leu	Thr Pro	Pro Gln	Ala Glu	Gln Glu	Val Asp	Val
1805			1810		1815		
Leu Asp	Val Asn	Val Arg	Gly Pro	Asp Gly	Cys Thr	Pro Leu	Met
1820			1825		1830		
Leu Ala	Ser Leu	Arg Gly	Gly Ser	Ser Asp	Leu Ser	Asp Glu	Asp
1835			1840		1845		
Glu Asp	Ala Glu	Asp Ser	Ser Ala	Asn Ile	Ile Thr	Asp Leu	Val
1850			1855		1860		
Tyr Gln	Gly Ala	Ser Leu	Gln Ala	Gln Thr	Asp Arg	Thr Gly	Glu
1865			1870		1875		
Met Ala	Leu His	Leu Ala	Ala Arg	Tyr Ser	Arg Ala	Asp Ala	Ala
1880			1885		1890		
Lys Arg	Leu Leu	Asp Ala	Gly Ala	Asp Ala	Asn Ala	Gln Asp	Asn
1895			1900		1905		
Met Gly	Arg Cys	Pro Leu	His Ala	Ala Val	Ala Ala	Asp Ala	Gln
1910			1915		1920		
Gly Val	Phe Gln	Ile Leu	Ile Arg	Asn Arg	Val Thr	Asp Leu	Asp
1925			1930		1935		
Ala Arg	Met Asn	Asp Gly	Thr Thr	Pro Leu	Ile Leu	Ala Ala	Arg
1940			1945		1950		
Leu Ala	Val Glu	Gly Met	Val Ala	Glu Leu	Ile Asn	Cys Gln	Ala
1955			1960		1965		
Asp Val	Asn Ala	Val Asp	Asp His	Gly Lys	Ser Ala	Leu His	Trp
1970			1975		1980		
Ala Ala	Ala Val	Asn Asn	Val Glu	Ala Thr	Leu Leu	Leu Leu	Lys
1985			1990		1995		
Asn Gly	Ala Asn	Arg Asp	Met Gln	Asp Asn	Lys Glu	Glu Thr	Pro
2000			2005		2010		
Leu Phe	Leu Ala	Ala Arg	Glu Gly	Ser Tyr	Glu Ala	Ala Lys	Ile
2015			2020		2025		
Leu Leu	Asp His	Phe Ala	Asn Arg	Asp Ile	Thr Asp	His Met	Asp
2030			2035		2040		
Arg Leu	Pro Arg	Asp Val	Ala Arg	Asp Arg	Met His	His Asp	Ile

-continued

2045	2050	2055
Val Arg Leu Leu Asp Glu Tyr Asn Val Thr Pro Ser Pro Pro Gly 2060 2065 2070		
Thr Val Leu Thr Ser Ala Leu Ser Pro Val Ile Cys Gly Pro Asn 2075 2080 2085		
Arg Ser Phe Leu Ser Leu Lys His Thr Pro Met Gly Lys Lys Ser 2090 2095 2100		
Arg Arg Pro Ser Ala Lys Ser Thr Met Pro Thr Ser Leu Pro Asn 2105 2110 2115		
Leu Ala Lys Glu Ala Lys Asp Ala Lys Gly Ser Arg Arg Lys Lys 2120 2125 2130		
Ser Leu Ser Glu Lys Val Gln Leu Ser Glu Ser Ser Val Thr Leu 2135 2140 2145		
Ser Pro Val Asp Ser Leu Glu Ser Pro His Thr Tyr Val Ser Asp 2150 2155 2160		
Thr Thr Ser Ser Pro Met Ile Thr Ser Pro Gly Ile Leu Gln Ala 2165 2170 2175		
Ser Pro Asn Pro Met Leu Ala Thr Ala Ala Pro Pro Ala Pro Val 2180 2185 2190		
His Ala Gln His Ala Leu Ser Phe Ser Asn Leu His Glu Met Gln 2195 2200 2205		
Pro Leu Ala His Gly Ala Ser Thr Val Leu Pro Ser Val Ser Gln 2210 2215 2220		
Leu Leu Ser His His His Ile Val Ser Pro Gly Ser Gly Ser Ala 2225 2230 2235		
Gly Ser Leu Ser Arg Leu His Pro Val Pro Val Pro Ala Asp Trp 2240 2245 2250		
Met Asn Arg Met Glu Val Asn Glu Thr Gln Tyr Asn Glu Met Phe 2255 2260 2265		
Gly Met Val Leu Ala Pro Ala Glu Gly Thr His Pro Gly Ile Ala 2270 2275 2280		
Pro Gln Ser Arg Pro Pro Glu Gly Lys His Ile Thr Thr Pro Arg 2285 2290 2295		
Glu Pro Leu Pro Pro Ile Val Thr Phe Gln Leu Ile Pro Lys Gly 2300 2305 2310		
Ser Ile Ala Gln Pro Ala Gly Ala Pro Gln Pro Gln Ser Thr Cys 2315 2320 2325		
Pro Pro Ala Val Ala Gly Pro Leu Pro Thr Met Tyr Gln Ile Pro 2330 2335 2340		
Glu Met Ala Arg Leu Pro Ser Val Ala Phe Pro Thr Ala Met Met 2345 2350 2355		
Pro Gln Gln Asp Gly Gln Val Ala Gln Thr Ile Leu Pro Ala Tyr 2360 2365 2370		
His Pro Phe Pro Ala Ser Val Gly Lys Tyr Pro Thr Pro Pro Ser 2375 2380 2385		
Gln His Ser Tyr Ala Ser Ser Asn Ala Ala Glu Arg Thr Pro Ser 2390 2395 2400		
His Ser Gly His Leu Gln Gly Glu His Pro Tyr Leu Thr Pro Ser 2405 2410 2415		
Pro Glu Ser Pro Asp Gln Trp Ser Ser Ser Ser Pro His Ser Ala 2420 2425 2430		
Ser Asp Trp Ser Asp Val Thr Thr Ser Pro Thr Pro Gly Gly Ala 2435 2440 2445		

-continued

Gly	Gly	Gly	Gln	Arg	Gly	Pro	Gly	Thr	His	Met	Ser	Glu	Pro	Pro
	2450					2455					2460			

His	Asn	Asn	Met	Gln	Val	Tyr	Ala
2465						2470	

<210> SEQ ID NO 41

<211> LENGTH: 2002

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

Met	Gln	Pro	Pro	Ser	Leu	Leu	Leu	Leu	Leu	Leu	Leu	Leu	Leu	Cys
1				5					10				15	

Val	Ser	Val	Val	Arg	Pro	Arg	Gly	Leu	Leu	Cys	Gly	Ser	Phe	Pro	Glu
			20					25					30		

Pro	Cys	Ala	Asn	Gly	Gly	Thr	Cys	Leu	Ser	Leu	Ser	Leu	Gly	Gln	Gly
		35					40					45			

Thr	Cys	Gln	Cys	Ala	Pro	Gly	Phe	Leu	Gly	Glu	Thr	Cys	Gln	Phe	Pro
	50					55					60				

Asp	Pro	Cys	Gln	Asn	Ala	Gln	Leu	Cys	Gln	Asn	Gly	Gly	Ser	Cys	Gln
65				70					75					80	

Ala	Leu	Leu	Pro	Ala	Pro	Leu	Gly	Leu	Pro	Ser	Ser	Pro	Ser	Pro	Leu
			85						90					95	

Thr	Pro	Ser	Phe	Leu	Cys	Thr	Cys	Leu	Pro	Gly	Phe	Thr	Gly	Glu	Arg
			100					105					110		

Cys	Gln	Ala	Lys	Leu	Glu	Asp	Pro	Cys	Pro	Pro	Ser	Phe	Cys	Ser	Lys
	115						120					125			

Arg	Gly	Arg	Cys	His	Ile	Gln	Ala	Ser	Gly	Arg	Pro	Gln	Cys	Ser	Cys
	130					135					140				

Met	Pro	Gly	Trp	Thr	Gly	Glu	Gln	Cys	Gln	Leu	Arg	Asp	Phe	Cys	Ser
145					150					155					160

Ala	Asn	Pro	Cys	Val	Asn	Gly	Gly	Val	Cys	Leu	Ala	Thr	Tyr	Pro	Gln
				165					170					175	

Ile	Gln	Cys	His	Cys	Pro	Pro	Gly	Phe	Glu	Gly	His	Ala	Cys	Glu	Arg
		180						185					190		

Asp	Val	Asn	Glu	Cys	Phe	Gln	Asp	Pro	Gly	Pro	Cys	Pro	Lys	Gly	Thr
	195						200					205			

Ser	Cys	His	Asn	Thr	Leu	Gly	Ser	Phe	Gln	Cys	Leu	Cys	Pro	Val	Gly
	210					215					220				

Gln	Glu	Gly	Pro	Arg	Cys	Glu	Leu	Arg	Ala	Gly	Pro	Cys	Pro	Pro	Arg
225					230					235					240

Gly	Cys	Ser	Asn	Gly	Gly	Thr	Cys	Gln	Leu	Met	Pro	Glu	Lys	Asp	Ser
			245						250					255	

Thr	Phe	His	Leu	Cys	Leu	Cys	Pro	Pro	Gly	Phe	Ile	Gly	Pro	Gly	Cys
			260						265				270		

Glu	Val	Asn	Pro	Asp	Asn	Cys	Val	Ser	His	Gln	Cys	Gln	Asn	Gly	Gly
		275					280					285			

Thr	Cys	Gln	Asp	Gly	Leu	Asp	Thr	Tyr	Thr	Cys	Leu	Cys	Pro	Glu	Thr
	290					295					300				

Trp	Thr	Gly	Trp	Asp	Cys	Ser	Glu	Asp	Val	Asp	Glu	Cys	Glu	Ala	Gln
305					310					315					320

Gly	Pro	Pro	His	Cys	Arg	Asn	Gly	Gly	Thr	Cys	Gln	Asn	Ser	Ala	Gly
			325						330					335	

Ser	Phe	His	Cys	Val	Cys	Val	Ser	Gly	Trp	Gly	Gly	Thr	Ser	Cys	Glu
			340						345					350	

Glu 355	Asn	Leu	Asp	Asp	Cys	Ile	Ala	Ala	Thr	Cys	Ala	Pro	Gly	Ser	Thr
Cys 370	Ile	Asp	Arg	Val	Gly	Ser	Phe	Ser	Cys	Leu	Cys	Pro	Pro	Gly	Arg
Thr 385	Gly	Leu	Leu	Cys	His	Leu	Glu	Asp	Met	Cys	Leu	Ser	Gln	Pro	Cys
His	Gly	Asp	Ala	Gln	Cys	Ser	Thr	Asn	Pro	Leu	Thr	Gly	Ser	Thr	Leu
Cys	Leu	Cys	Gln	Pro	Gly	Tyr	Ser	Gly	Pro	Thr	Cys	His	Gln	Asp	Leu
Asp	Glu	Cys	Leu	Met	Ala	Gln	Gln	Gly	Pro	Ser	Pro	Cys	Glu	His	Gly
Gly	Ser	Cys	Leu	Asn	Thr	Pro	Gly	Ser	Phe	Asn	Cys	Leu	Cys	Pro	Pro
Gly 465	Tyr	Thr	Gly	Ser	Arg	Cys	Glu	Ala	Asp	His	Asn	Glu	Cys	Leu	Ser
Gln	Pro	Cys	His	Pro	Gly	Ser	Thr	Cys	Leu	Asp	Leu	Leu	Ala	Thr	Phe
His	Cys	Leu	Cys	Pro	Pro	Gly	Leu	Glu	Gly	Gln	Leu	Cys	Glu	Val	Glu
Thr	Asn	Glu	Cys	Ala	Ser	Ala	Pro	Cys	Leu	Asn	His	Ala	Asp	Cys	His
Asp	Leu	Leu	Asn	Gly	Phe	Gln	Cys	Ile	Cys	Leu	Pro	Gly	Phe	Ser	Gly
Thr 545	Arg	Cys	Glu	Glu	Asp	Ile	Asp	Glu	Cys	Arg	Ser	Ser	Pro	Cys	Ala
Asn	Gly	Gly	Gln	Cys	Gln	Asp	Gln	Pro	Gly	Ala	Phe	His	Cys	Lys	Cys
Leu	Pro	Gly	Phe	Glu	Gly	Pro	Arg	Cys	Gln	Thr	Glu	Val	Asp	Glu	Cys
Leu	Ser	Asp	Pro	Cys	Pro	Val	Gly	Ala	Ser	Cys	Leu	Asp	Leu	Pro	Gly
Ala	Phe	Phe	Cys	Leu	Cys	Pro	Ser	Gly	Phe	Thr	Gly	Gln	Leu	Cys	Glu
Val 625	Pro	Leu	Cys	Ala	Pro	Asn	Leu	Cys	Gln	Pro	Lys	Gln	Ile	Cys	Lys
Asp	Gln	Lys	Asp	Lys	Ala	Asn	Cys	Leu	Cys	Pro	Asp	Gly	Ser	Pro	Gly
Cys	Ala	Pro	Pro	Glu	Asp	Asn	Cys	Thr	Cys	His	His	Gly	His	Cys	Gln
Arg	Ser	Ser	Cys	Val	Cys	Asp	Val	Gly	Trp	Thr	Gly	Pro	Glu	Cys	Glu
Ala	Glu	Leu	Gly	Gly	Cys	Ile	Ser	Ala	Pro	Cys	Ala	His	Gly	Gly	Thr
Cys 705	Tyr	Pro	Gln	Pro	Ser	Gly	Tyr	Asn	Cys	Thr	Cys	Pro	Thr	Gly	Tyr
Thr	Gly	Pro	Thr	Cys	Ser	Glu	Glu	Met	Thr	Ala	Cys	His	Ser	Gly	Pro
Cys	Leu	Asn	Gly	Gly	Ser	Cys	Asn	Pro	Ser	Pro	Gly	Gly	Tyr	Tyr	Cys
Thr	Cys	Pro	Pro	Ser	His	Thr	Gly	Pro	Gln	Cys	Gln	Thr	Ser	Thr	Asp
Tyr	Cys	Val	Ser	Ala	Pro	Cys	Phe	Asn	Gly	Gly	Thr	Cys	Val	Asn	Arg

-continued

Pro Gly Thr Phe Ser Cys Leu Cys Ala Met Gly Phe Gln Gly Pro Arg
 785 790 795 800
 Cys Glu Gly Lys Leu Arg Pro Ser Cys Ala Asp Ser Pro Cys Arg Asn
 805 810 815
 Arg Ala Thr Cys Gln Asp Ser Pro Gln Gly Pro Arg Cys Leu Cys Pro
 820 825 830
 Thr Gly Tyr Thr Gly Gly Ser Cys Gln Thr Leu Met Asp Leu Cys Ala
 835 840 845
 Gln Lys Pro Cys Pro Arg Asn Ser His Cys Leu Gln Thr Gly Pro Ser
 850 855 860
 Phe His Cys Leu Cys Leu Gln Gly Trp Thr Gly Pro Leu Cys Asn Leu
 865 870 875 880
 Pro Leu Ser Ser Cys Gln Lys Ala Ala Leu Ser Gln Gly Ile Asp Val
 885 890 895
 Ser Ser Leu Cys His Asn Gly Gly Leu Cys Val Asp Ser Gly Pro Ser
 900 905 910
 Tyr Phe Cys His Cys Pro Pro Gly Phe Gln Gly Ser Leu Cys Gln Asp
 915 920 925
 His Val Asn Pro Cys Glu Ser Arg Pro Cys Gln Asn Gly Ala Thr Cys
 930 935 940
 Met Ala Gln Pro Ser Gly Tyr Leu Cys Gln Cys Ala Pro Gly Tyr Asp
 945 950 955 960
 Gly Gln Asn Cys Ser Lys Glu Leu Asp Ala Cys Gln Ser Gln Pro Cys
 965 970 975
 His Asn His Gly Thr Cys Thr Pro Lys Pro Gly Gly Phe His Cys Ala
 980 985 990
 Cys Pro Pro Gly Phe Val Gly Leu Arg Cys Glu Gly Asp Val Asp Glu
 995 1000 1005
 Cys Leu Asp Gln Pro Cys His Pro Thr Gly Thr Ala Ala Cys His
 1010 1015 1020
 Ser Leu Ala Asn Ala Phe Tyr Cys Gln Cys Leu Pro Gly His Thr
 1025 1030 1035
 Gly Gln Trp Cys Glu Val Glu Ile Asp Pro Cys His Ser Gln Pro
 1040 1045 1050
 Cys Phe His Gly Gly Thr Cys Glu Ala Thr Ala Gly Ser Pro Leu
 1055 1060 1065
 Gly Phe Ile Cys His Cys Pro Lys Gly Phe Glu Gly Pro Thr Cys
 1070 1075 1080
 Ser His Arg Ala Pro Ser Cys Gly Phe His His Cys His His Gly
 1085 1090 1095
 Gly Leu Cys Leu Pro Ser Pro Lys Pro Gly Phe Pro Pro Arg Cys
 1100 1105 1110
 Ala Cys Leu Ser Gly Tyr Gly Gly Pro Asp Cys Leu Thr Pro Pro
 1115 1120 1125
 Ala Pro Lys Gly Cys Gly Pro Pro Ser Pro Cys Leu Tyr Asn Gly
 1130 1135 1140
 Ser Cys Ser Glu Thr Thr Gly Leu Gly Gly Pro Gly Phe Arg Cys
 1145 1150 1155
 Ser Cys Pro His Ser Ser Pro Gly Pro Arg Cys Gln Lys Pro Gly
 1160 1165 1170
 Ala Lys Gly Cys Glu Gly Arg Ser Gly Asp Gly Ala Cys Asp Ala
 1175 1180 1185
 Gly Cys Ser Gly Pro Gly Gly Asn Trp Asp Gly Gly Asp Cys Ser

-continued

1190	1195	1200
Leu Gly Val Pro Asp Pro Trp Lys Gly Cys Pro Ser His Ser Arg		
1205	1210	1215
Cys Trp Leu Leu Phe Arg Asp Gly Gln Cys His Pro Gln Cys Asp		
1220	1225	1230
Ser Glu Glu Cys Leu Phe Asp Gly Tyr Asp Cys Glu Thr Pro Pro		
1235	1240	1245
Ala Cys Thr Pro Ala Tyr Asp Gln Tyr Cys His Asp His Phe His		
1250	1255	1260
Asn Gly His Cys Glu Lys Gly Cys Asn Thr Ala Glu Cys Gly Trp		
1265	1270	1275
Asp Gly Gly Asp Cys Arg Pro Glu Asp Gly Asp Pro Glu Trp Gly		
1280	1285	1290
Pro Ser Leu Ala Leu Leu Val Val Leu Ser Pro Pro Ala Leu Asp		
1295	1300	1305
Gln Gln Leu Phe Ala Leu Ala Arg Val Leu Ser Leu Thr Leu Arg		
1310	1315	1320
Val Gly Leu Trp Val Arg Lys Asp Arg Asp Gly Arg Asp Met Val		
1325	1330	1335
Tyr Pro Tyr Pro Gly Ala Arg Ala Glu Glu Lys Leu Gly Gly Thr		
1340	1345	1350
Arg Asp Pro Thr Tyr Gln Glu Arg Ala Ala Pro Gln Thr Gln Pro		
1355	1360	1365
Leu Gly Lys Glu Thr Asp Ser Leu Ser Ala Gly Phe Val Val Val		
1370	1375	1380
Met Gly Val Asp Leu Ser Arg Cys Gly Pro Asp His Pro Ala Ser		
1385	1390	1395
Arg Cys Pro Trp Asp Pro Gly Leu Leu Leu Arg Phe Leu Ala Ala		
1400	1405	1410
Met Ala Ala Val Gly Ala Leu Glu Pro Leu Leu Pro Gly Pro Leu		
1415	1420	1425
Leu Ala Val His Pro His Ala Gly Thr Ala Pro Pro Ala Asn Gln		
1430	1435	1440
Leu Pro Trp Pro Val Leu Cys Ser Pro Val Ala Gly Val Ile Leu		
1445	1450	1455
Leu Ala Leu Gly Ala Leu Leu Val Leu Gln Leu Ile Arg Arg Arg		
1460	1465	1470
Arg Arg Glu His Gly Ala Leu Trp Leu Pro Pro Gly Phe Thr Arg		
1475	1480	1485
Arg Pro Arg Thr Gln Ser Ala Pro His Arg Arg Arg Pro Pro Leu		
1490	1495	1500
Gly Glu Asp Ser Ile Gly Leu Lys Ala Leu Lys Pro Lys Ala Glu		
1505	1510	1515
Val Asp Glu Asp Gly Val Val Met Cys Ser Gly Pro Glu Glu Gly		
1520	1525	1530
Glu Glu Val Gly Gln Ala Glu Glu Thr Gly Pro Pro Ser Thr Cys		
1535	1540	1545
Gln Leu Trp Ser Leu Ser Gly Gly Cys Gly Ala Leu Pro Gln Ala		
1550	1555	1560
Ala Met Leu Thr Pro Pro Gln Glu Ser Glu Met Glu Ala Pro Asp		
1565	1570	1575
Leu Asp Thr Arg Gly Pro Asp Gly Val Thr Pro Leu Met Ser Ala		
1580	1585	1590

-continued

Val	Cys	Cys	Gly	Glu	Val	Gln	Ser	Gly	Thr	Phe	Gln	Gly	Ala	Trp
1595						1600					1605			
Leu	Gly	Cys	Pro	Glu	Pro	Trp	Glu	Pro	Leu	Leu	Asp	Gly	Gly	Ala
1610						1615					1620			
Cys	Pro	Gln	Ala	His	Thr	Val	Gly	Thr	Gly	Glu	Thr	Pro	Leu	His
1625						1630					1635			
Leu	Ala	Ala	Arg	Phe	Ser	Arg	Pro	Thr	Ala	Ala	Arg	Arg	Leu	Leu
1640						1645					1650			
Glu	Ala	Gly	Ala	Asn	Pro	Asn	Gln	Pro	Asp	Arg	Ala	Gly	Arg	Thr
1655						1660					1665			
Pro	Leu	His	Ala	Ala	Val	Ala	Ala	Asp	Ala	Arg	Glu	Val	Cys	Gln
1670						1675					1680			
Leu	Leu	Leu	Arg	Ser	Arg	Gln	Thr	Ala	Val	Asp	Ala	Arg	Thr	Glu
1685						1690					1695			
Asp	Gly	Thr	Thr	Pro	Leu	Met	Leu	Ala	Ala	Arg	Leu	Ala	Val	Glu
1700						1705					1710			
Asp	Leu	Val	Glu	Glu	Leu	Ile	Ala	Ala	Gln	Ala	Asp	Val	Gly	Ala
1715						1720					1725			
Arg	Asp	Lys	Trp	Gly	Lys	Thr	Ala	Leu	His	Trp	Ala	Ala	Ala	Val
1730						1735					1740			
Asn	Asn	Ala	Arg	Ala	Ala	Arg	Ser	Leu	Leu	Gln	Ala	Gly	Ala	Asp
1745						1750					1755			
Lys	Asp	Ala	Gln	Asp	Asn	Arg	Glu	Gln	Thr	Pro	Leu	Phe	Leu	Ala
1760						1765					1770			
Ala	Arg	Glu	Gly	Ala	Val	Glu	Val	Ala	Gln	Leu	Leu	Leu	Gly	Leu
1775						1780					1785			
Gly	Ala	Ala	Arg	Glu	Leu	Arg	Asp	Gln	Ala	Gly	Leu	Ala	Pro	Ala
1790						1795					1800			
Asp	Val	Ala	His	Gln	Arg	Asn	His	Trp	Asp	Leu	Leu	Thr	Leu	Leu
1805						1810					1815			
Glu	Gly	Ala	Gly	Pro	Pro	Glu	Ala	Arg	His	Lys	Ala	Thr	Pro	Gly
1820						1825					1830			
Arg	Glu	Ala	Gly	Pro	Phe	Pro	Arg	Ala	Arg	Thr	Val	Ser	Val	Ser
1835						1840					1845			
Val	Pro	Pro	His	Gly	Gly	Gly	Ala	Leu	Pro	Arg	Cys	Arg	Thr	Leu
1850						1855					1860			
Ser	Ala	Gly	Ala	Gly	Pro	Arg	Gly	Gly	Gly	Ala	Cys	Leu	Gln	Ala
1865						1870					1875			
Arg	Thr	Trp	Ser	Val	Asp	Leu	Ala	Ala	Arg	Gly	Gly	Gly	Ala	Tyr
1880						1885					1890			
Ser	His	Cys	Arg	Ser	Leu	Ser	Gly	Val	Gly	Ala	Gly	Gly	Gly	Pro
1895						1900					1905			
Thr	Pro	Arg	Gly	Arg	Arg	Phe	Ser	Ala	Gly	Met	Arg	Gly	Pro	Arg
1910						1915					1920			
Pro	Asn	Pro	Ala	Ile	Met	Arg	Gly	Arg	Tyr	Gly	Val	Ala	Ala	Gly
1925						1930					1935			
Arg	Gly	Gly	Arg	Val	Ser	Thr	Asp	Asp	Trp	Pro	Cys	Asp	Trp	Val
1940						1945					1950			
Ala	Leu	Gly	Ala	Cys	Gly	Ser	Ala	Ser	Asn	Ile	Pro	Ile	Pro	Pro
1955						1960					1965			
Pro	Cys	Leu	Thr	Pro	Ser	Pro	Glu	Arg	Gly	Ser	Pro	Gln	Leu	Asp
1970						1975					1980			
Cys	Gly	Pro	Pro	Ala	Leu	Gln	Glu	Met	Pro	Ile	Asn	Gln	Gly	Gly
1985						1990					1995			

-continued

Glu Gly Lys Lys
2000

<210> SEQ ID NO 42
<211> LENGTH: 135
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 42

atgggtccag gtgcaagagg tagaaggcgt agaaggagac caatgagccc acctcctccg 60
ccacctccag tgagagcact gcctttgctg ttgctgctgg ctggacctgg tgcagcagct 120
cctccttgcc tggac 135

<210> SEQ ID NO 43
<211> LENGTH: 135
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

atggggccgg gggcccgtag cgcgcgcgc gcgcgtgcc cgatgtgcc gccaccgcca 60
ccgccacccg tgcgggcgct gccctgctg ctgctgctag cggggccggg ggctgcagcc 120
cccccttgcc tggac 135

<210> SEQ ID NO 44
<211> LENGTH: 45
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 44

Met Gly Pro Gly Ala Arg Gly Arg Arg Arg Arg Arg Pro Met Ser
1 5 10 15
Pro Pro Pro Pro Pro Pro Pro Val Arg Ala Leu Pro Leu Leu Leu
20 25 30
Leu Ala Gly Pro Gly Ala Ala Ala Pro Pro Cys Leu Asp
35 40 45

What is claimed:

1. An isolated nucleic acid encoding a polypeptide comprising the amino acid sequence of the VH chain region of a monoclonal antibody that specifically binds to Notch3 receptor, wherein the antibody specifically binds to an epitope in SEQ ID NO:10 and activates Notch3 receptor-mediated signaling through the Notch3 receptor independent of ligand binding to the Notch3 receptor.

2. The nucleic acid of claim 1, wherein the VH chain region comprises CDR-H1 of SEQ ID NO:4, CDR-H2 of SEQ ID NO:5, and CDR-H3 of SEQ ID NO:6.

3. The nucleic acid of claim 2, wherein the VH chain region comprises SEQ ID NO:2.

4. An isolated cell comprising the nucleic acid of claim 1.

5. A method for producing a polypeptide comprising the amino acid sequence of the VH chain region of a monoclonal antibody that specifically binds to Notch3 comprising culturing the cell of claim 4 under conditions appropriate for production of the polypeptide and isolating the polypeptide so produced.

6. An isolated nucleic acid encoding a polypeptide comprising the amino acid sequence of the VL chain region of a monoclonal antibody that specifically binds to Notch3 receptor, wherein the antibody specifically binds to an epitope in SEQ ID NO:10 and activates Notch3 receptor-mediated signaling through the Notch3 receptor independent of ligand binding to the Notch3 receptor.

7. The nucleic acid of claim 6, wherein the VL chain region comprises CDR-L1 of SEQ ID NO:7, CDR-L2 of SEQ ID NO:8, and CDR-L3 of SEQ ID NO:9.

8. The nucleic acid of claim 7, wherein the VL chain region comprises SEQ ID NO:3.

9. An isolated cell comprising the nucleic acid of claim 6.

10. A method for producing a polypeptide comprising the amino acid sequence of the VL chain region of a monoclonal antibody that specifically binds to Notch3 comprising culturing the cell of claim 9 under conditions appropriate for production of the polypeptide and isolating the polypeptide so produced.

109

11. An isolated nucleic acid encoding a monoclonal anti-
body that specifically binds to Notch3 receptor, wherein the
antibody specifically binds to an epitope in SEQ ID NO:10,
and wherein the antibody activates Notch3 receptor-mediated
signaling through the Notch3 receptor independent of ligand
binding to the Notch3 receptor.

12. The nucleic acid of claim 11, wherein the antibody
comprises a VH chain region comprising CDR-H1 of SEQ ID
NO:4, CDR-H2 of SEQ ID NO:5, and CDR-H3 of SEQ ID
NO:6, and a VL chain region comprising CDR-L1 of SEQ ID
NO:7, CDR-L2 of SEQ ID NO:8, and CDR-L3 of SEQ ID
NO:9.

13. The nucleic acid of claim 12, wherein the VH chain
region comprises SEQ ID NO:2 and the VL chain region
comprises SEQ ID NO:3.

110

14. The nucleic acid of claim 11, wherein the antibody is a
humanized form of a monoclonal antibody comprising the
VH chain region of SEQ ID NO:2 and the VL chain region of
SEQ ID NO:3.

15. An isolated nucleic acid encoding a monoclonal anti-
body that specifically binds to Notch3 receptor, wherein the
antibody specifically binds to an epitope in SEQ ID NO:11,
and wherein the antibody activates Notch3 receptor-mediated
signaling through the Notch3 receptor independent of ligand
binding to the Notch3 receptor.

* * * * *