



(86) **Date de dépôt PCT/PCT Filing Date:** 2014/01/08
(87) **Date publication PCT/PCT Publication Date:** 2014/07/17
(85) **Entrée phase nationale/National Entry:** 2015/07/06
(86) **N° demande PCT/PCT Application No.:** US 2014/010607
(87) **N° publication PCT/PCT Publication No.:** 2014/110086
(30) **Priorité/Priority:** 2013/01/08 (US61/750,221)

(51) **Cl.Int./Int.Cl.** **C07D 401/14** (2006.01),
A61K 31/4184 (2006.01), **C07D 405/14** (2006.01),
C07D 409/14 (2006.01)

(71) **Demandeur/Applicant:**
SAMUMED, LLC, US

(72) **Inventeurs/Inventors:**
KUMAR KC, SUNIL, US;
HOOD, JOHN, US

(74) **Agent:** SMART & BIGGAR

(54) **Titre : INHIBITEURS DE 3-(BENZOIMIDAZOL-2-YL)-INDAZOLE DE LA VOIE DE SIGNALISATION PAR WNT ET LEURS
UTILISATIONS THERAPEUTIQUES**

(54) **Title: 3-(BENZOIMIDAZOL-2-YL)-INDAZOLE INHIBITORS OF THE WNT SIGNALING PATHWAY AND THERAPEUTIC USES
THEREOF**

(57) **Abrégé/Abstract:**

Indazole compounds for treating various diseases and pathologies are disclosed. More particularly, the present disclosure concerns the use of an indazole compound or analogs thereof, in the treatment of disorders characterized by the activation of Wnt pathway signaling (e.g., cancer, abnormal cellular proliferation, angiogenesis, Alzheimer's disease, lung disease and osteoarthritis), the modulation of cellular events mediated by Wnt pathway signaling, as well as genetic diseases and neurological conditions/disorders/diseases due to mutations or dysregulation of the Wnt pathway and/or of one or more of Wnt signaling components. Also provided are methods for treating Wnt-related disease states.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
17 July 2014 (17.07.2014)

(10) International Publication Number
WO 2014/110086 A3

- (51) International Patent Classification:
A61K 31/416 (2006.01) *A61K 31/4184* (2006.01)

(21) International Application Number:
PCT/US2014/010607

(22) International Filing Date:
8 January 2014 (08.01.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/750,221 8 January 2013 (08.01.2013) US

(71) Applicant: **SAMUMED, LLC** [US/US]; 9381 Judicial Drive, Suite 160, San Diego, California 92121 (US).

(72) Inventors; and

(71) Applicants : **KUMAR KC, Sunil** [US/US]; 10504 Clasico Court, San Diego, California 92127 (US). **HOOD, John** [US/US]; 10831 Vereda Sol del Dios, San Diego, California 92130 (US).

(74) Agents: **FOLLETT, Angela D.** et al.; Fish & Richardson P.C., P.O. Box 1022, Minneapolis, Minnesota 55440 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

- DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:
- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- (88) Date of publication of the international search report:
9 October 2014

(54) Title: 3-(BENZOIMIDAZOL-2-YL)-INDAZOLE INHIBITORS OF THE WNT SIGNALING PATHWAY AND THERAPEUTIC USES THEREOF

(57) Abstract: Indazole compounds for treating various diseases and pathologies are disclosed. More particularly, the present disclosure concerns the use of an indazole compound or analogs thereof, in the treatment of disorders characterized by the activation of Wnt pathway signaling (e.g., cancer, abnormal cellular proliferation, angiogenesis, Alzheimer's disease, lung disease and osteoarthritis), the modulation of cellular events mediated by Wnt pathway signaling, as well as genetic diseases and neurological conditions/disorders/diseases due to mutations or dysregulation of the Wnt pathway and/or of one or more of Wnt signaling components. Also provided are methods for treating Wnt-related disease states.

3-(BENZOIMIDAZOL-2-YL)-INDAZOLE INHIBITORS OF THE WNT SIGNALING PATHWAY AND THERAPEUTIC USES THEREOF

RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 61/750,221, filed January 8, 2013, which is incorporated herein by reference in its entirety.

BACKGROUND

Technical Field

[0002] This disclosure relates to inhibitors of one or more proteins in the Wnt pathway, including inhibitors of one or more Wnt proteins, and compositions comprising the same. More particularly, it concerns the use of an indazole compound or salts or analogs thereof, in the treatment of disorders characterized by the activation of Wnt pathway signaling (e.g., cancer, abnormal cellular proliferation, angiogenesis, Alzheimer's disease, lung disease, and osteoarthritis), the modulation of cellular events mediated by Wnt pathway signaling, as well as genetic diseases and neurological conditions/disorders/diseases due to mutations or dysregulation of the Wnt pathway and/or of one or more of Wnt signaling components. Also provided are methods for treating Wnt-related disease states.

Background

[0003] The Wnt growth factor family includes more than 10 genes identified in the mouse and at least 19 genes identified in the human. Members of the Wnt family of signaling molecules mediate many short-and long-range patterning processes during invertebrate and vertebrate development. The Wnt signaling pathway is known for its role in the inductive interactions that regulate growth and differentiation, and it also plays roles in the homeostatic maintenance of post-embryonic tissue integrity. Wnt stabilizes cytoplasmic β -catenin, which stimulates the expression of genes including c-myc, c jun, fra-1, and cyclin D1. In addition, misregulation of Wnt signaling can cause developmental defects and is implicated in the genesis of several human cancers. More recently, the Wnt


pathway has been implicated in the maintenance of stem or progenitor cells in a growing list of adult tissues that now includes skin, blood, gut, prostate, muscle and the nervous system.

SUMMARY

[0004] The present disclosure provides methods and reagents, involving contacting a cell with an agent, such as an indazole compound, in a sufficient amount to antagonize a Wnt activity, e.g., to reverse or control an aberrant growth state or correct a genetic disorder due to mutations in Wnt signaling components.

[0005] Some embodiments disclosed herein include Wnt inhibitors containing an indazole core. Other embodiments disclosed herein include pharmaceutical compositions and methods of treatment using these compounds.

[0006] One embodiment disclosed herein includes compounds of Formula I:



[0007] In some embodiments of Formula (I):

R^1 is -heteroaryl R^3R^4 ;

R^2 is selected from the group consisting of H, halide, C_{1-3} alkyl, -CN, -OR⁸, -OH, -(C_{1-3} alkyl)OR⁸, -NR⁹R¹⁰, -(C_{1-3} alkyl)NR⁹R¹⁰, -heteroaryl R^5 , -heterocyclyl R^6 and -aryl R^7 ;

R^3 is 1 substituent attached to the heteroaryl ring and is selected from the group consisting of H, C_{1-3} alkyl, -CF₃, -NR⁹R¹⁰, -NHC(=O)R⁸, -(C_{1-3} alkyl)heterocyclyl R^6 and -(C_{1-3} alkyl)NR⁹R¹⁰;

R^4 is 1 substituent attached to the heteroaryl ring and is selected from the group consisting of H, C_{1-3} alkyl, -CF₃, halide, -CN, -OR⁸, -OH, -(C_{1-3} alkyl)OR⁸, -NR⁹R¹⁰, -(C_{1-3} alkyl)NR⁹R¹⁰ and -OCF₃;

R^5 is 1-3 substituents attached to the heteroaryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-OR^8$, $-OH$, $-(C_{1-3} \text{ alkyl})OR^8$, $-NR^9R^{10}$, $-(C_{1-3} \text{ alkyl})NR^9R^{10}$ and $-OCF_3$;

R^6 is 1-3 substituents attached to the heterocyclyl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-OR^8$, $-OH$, $-(C_{1-3} \text{ alkyl})OR^8$, $-NR^9R^{10}$, $-(C_{1-3} \text{ alkyl})NR^9R^{10}$ and $-OCF_3$;

R^7 is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-OR^8$, $-OH$, $-(C_{1-3} \text{ alkyl})OR^8$, $-NR^9R^{10}$, $-(C_{1-3} \text{ alkyl})NR^9R^{10}$ and $-OCF_3$;

each R^8 is independently selected from the group consisting of C_{1-9} alkyl, $-$ heteroaryl R^{12} , $-$ heterocyclyl R^{13} , $-$ aryl R^{14} , carbocyclyl R^{11} , $-(C_{1-3} \text{ alkyl})$ heteroaryl R^{12} , $-(C_{1-3} \text{ alkyl})$ heterocyclyl R^{13} , $-(C_{1-3} \text{ alkyl})$ aryl R^{14} and $-(C_{1-3} \text{ alkyl})$ carbocyclyl R^{11} ;

each R^9 is independently selected from the group consisting of H, C_{1-6} alkyl, $-$ heteroaryl R^{12} , $-$ heterocyclyl R^{13} , $-$ aryl R^{14} , carbocyclyl R^{11} , $-(C_{1-3} \text{ alkyl})$ heteroaryl R^{12} , $-(C_{1-3} \text{ alkyl})$ heterocyclyl R^{13} , $-(C_{1-3} \text{ alkyl})$ aryl R^{14} and $-(C_{1-3} \text{ alkyl})$ carbocyclyl R^{11} ;

each R^{10} is independently selected from the group consisting of H and C_{1-6} alkyl;
or

R^9 and R^{10} are optionally linked to form a five or six membered heterocyclyl ring;


R^{11} is 1-3 substituents attached to the carbocyclyl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;


R^{12} is 1-3 substituents attached to the heteroaryl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;

R^{13} is 1-3 substituents attached to the heterocyclyl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;

R^{14} is 1-3 substituents attached to the aryl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;

with the proviso that a compound of Formula I is not a compound selected from the group consisting of:





[0008] Some embodiments include stereoisomers and pharmaceutically acceptable salts of a compound of Formula (I).

[0009] Some embodiments include pro-drugs of a compound of Formula (I).

[0010] Some embodiments of the present disclosure include pharmaceutical compositions comprising a compound of Formula (I) and a pharmaceutically acceptable carrier, diluent, or excipient.

[0011] Other embodiments disclosed herein include methods of inhibiting one or more members of the Wnt pathway, including one or more Wnt proteins by administering to a patient affected by a disorder or disease in which aberrant Wnt

signaling is implicated, such as cancer and other diseases associated with abnormal angiogenesis, cellular proliferation, cell cycling and mutations in Wnt signaling components, a compound according to Formula (I). Accordingly, the compounds and compositions provided herein can be used to treat cancer, to reduce or inhibit angiogenesis, to reduce or inhibit cellular proliferation and correct a genetic disorder due to mutations in Wnt signaling components.

[0012] Non-limiting examples of diseases which can be treated with the compounds and compositions provided herein include a variety of cancers, diabetic retinopathy, pulmonary fibrosis, rheumatoid arthritis, sepsis, anklyosing spondylitis, psoriasis, scleroderma, mycotic and viral infections, osteochondrodysplasia, Alzheimer's disease, lung disease, osteoarthritis, articular cartilage (chondral) defects, degenerative disc disease (or intervertebral disc degeneration), polyposis coli, osteoporosis-pseudoglioma syndrome, familial exudative vitreoretinopathy, retinal angiogenesis, early coronary disease, tetra-ameliasyndrome, Müllerian-duct regression and virilization, SERKAL syndrome, diabetes mellitus type 2, Fuhrmann syndrome, Al-Awadi/Raas-Rothschild/Schinzle phocomelia syndrome, odonto-onycho-dermal dysplasia, obesity, split-hand/foot malformation, caudal duplication syndrome, tooth agenesis, Wilms tumor, skeletal dysplasia, focal dermal hypoplasia, autosomal recessive anonychia, neural tube defects, alpha-thalassemia (ATRX) syndrome, fragile X syndrome, ICF syndrome, Angelman syndrome, Prader-Willi syndrome, Beckwith-Wiedemann Syndrome, Norrie disease, and Rett syndrome.

[0013] Some embodiments of the present disclosure include methods to prepare compounds of Formula (I).

[0014] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

DETAILED DESCRIPTION

[0015] Compositions and methods for inhibiting one or more members of the Wnt pathway, including one or more Wnt proteins are useful.

[0016] Some embodiments relate to a method for treating a disease including, but not limited to, cancers, diabetic retinopathy, pulmonary fibrosis, rheumatoid arthritis, sepsis, anklyosing spondylitism, psoriasis, scleroderma, mycotic and viral infections, bone and cartilage diseases, Alzheimer's disease, lung disease, osteoarthritis, articular cartilage (chondral) defects, degenerative disc disease (or intervertebral disc degeneration), polyposis coli, bone density and vascular defects in the eye (Osteoporosis-pseudoglioma Syndrome, OPPG), familial exudative vitreoretinopathy, retinal angiogenesis, early coronary disease, tetra-amelia, Müllerian-duct regression and virilization, SERKAL syndrome, type II diabetes, Fuhrmann syndrome, Al-Awadi/Raas-Rothschild/Schinzal phocomelia syndrome, odonto-onycho-dermal dysplasia, obesity, split-hand/foot malformation, caudal duplication, tooth agenesis, Wilms tumor, skeletal dysplasia, focal dermal hypoplasia, autosomal recessive anonychia, neural tube defects, alpha-thalassemia (ATRX) syndrome, fragile X syndrome, ICF syndrome, Angelman's syndrome, Prader-Willi syndrome, Beckwith-Wiedemann Syndrome, Norrie disease, and Rett syndrome.

[0017] In some embodiments, non-limiting examples of bone and cartilage diseases which can be treated with the compounds and compositions provided herein include bone spur (osteophytes), craniosynostosis, fibrodysplasia ossificans progressive, fibrous dysplasia, giant cell tumor of bone, hip labral tear, meniscal tears, osteoarthritis, articular cartilage (chondral) defects, degenerative disc disease (or intervertebral disc degeneration), osteochondritis dissecans, osteochondroma (bone tumor), osteopetrosis, relapsing polychondritis, and Salter-Harris fractures.

[0018] In some embodiments, pharmaceutical compositions are provided that are effective for treatment of a disease of an animal, e.g., a mammal, caused by the pathological activation or mutations of the Wnt pathway. The composition includes a pharmaceutically acceptable carrier and a compound as described herein.

Definitions

[0019] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this disclosure belongs. All patents, applications, published applications, and other

publications are incorporated by reference in their entirety. In the event that there is a plurality of definitions for a term herein, those in this section prevail unless stated otherwise.

[0020] As used herein, "alkyl" means a branched, or straight chain chemical group containing only carbon and hydrogen, such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, iso-pentyl, sec-pentyl, and neo-pentyl. Alkyl groups can either be unsubstituted or substituted with one or more substituents, e.g., halide, alkoxy, acyloxy, amino, amido, cyano, nitro, hydroxyl, thio, carboxy, carbonyl, benzyloxy, aryl, heteroaryl, or other functionality that may be suitably blocked, if necessary for purposes of the disclosure, with a protecting group. Alkyl groups can be saturated or unsaturated (e.g., containing $-C=C-$ or $-C\equiv C-$ subunits), at one or several positions. Typically, alkyl groups will comprise 1 to 9 carbon atoms (e.g., 1 to 6, 1 to 4, or 1 to 2 carbon atoms).

[0021] As used herein, "carbocyclyl" means a cyclic ring system containing only carbon atoms in the ring system backbone, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cyclohexenyl. Carbocyclyls may include multiple fused rings. Carbocyclyls may have any degree of saturation provided that at least one ring in the ring system is not aromatic. Carbocyclyl groups can either be unsubstituted or substituted with one or more substituents, e.g., alkyl, halide, alkoxy, acyloxy, amino, amido, cyano, nitro, hydroxyl, mercapto, carboxy, carbonyl, benzyloxy, aryl, heteroaryl, or other functionality that may be suitably blocked, if necessary for purposes of the disclosure, with a protecting group. Typically, carbocyclyl groups will comprise 3 to 10 carbon atoms (e.g., 3 to 6 carbon atoms).

[0022] As used herein, "lower alkyl" means a subset of alkyl having 1 to 3 carbon atoms, and can be linear or branched. Examples of lower alkyl groups include methyl, ethyl, n-propyl, and isopropyl. Likewise, radicals using the terminology "lower" refer to radicals having 1 to about 3 carbons in the alkyl portion of the radical.

[0023] As used herein, "amido" means a H-CON-, alkyl-CON-, carbocyclyl-CON-, aryl-CON-, heteroaryl-CON-, or heterocyclyl-CON group wherein the alkyl, carbocyclyl, aryl or heterocyclyl group is as herein described.

[0024] As used herein, “aryl” means an aromatic radical having a single-ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl or anthryl) with only carbon atoms present in the ring backbone. Aryl groups can either be unsubstituted or substituted with one or more substituents, e.g., amino, cyano, hydroxyl, lower alkyl, haloalkyl, alkoxy, nitro, halo, mercapto, and other substituents. An exemplary carbocyclic aryl is phenyl.

[0025] As used herein, the term “heteroaryl” means a mono-, bi-, tri- or polycyclic group having 5 to 14 ring atoms, alternatively 5, 6, 9, or 10 ring atoms; having 6, 10, or 14 pi electrons shared in a cyclic array; wherein at least one ring in the system is aromatic, and at least one ring in the system contains one or more heteroatoms independently selected from the group consisting of N, O, and S. Heteroaryl groups can either be unsubstituted or substituted with one or more substituents, e.g., amino, cyano, hydroxyl, lower alkyl, haloalkyl, alkoxy, nitro, halo, mercapto, and other substituents. Examples of heteroaryl include thienyl, pyridinyl, furyl, oxazolyl, oxadiazolyl, pyrrolyl, imidazolyl, triazolyl, thiodiazolyl, pyrazolyl, isoxazolyl, thiadiazolyl, pyranyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, thiazolyl, benzothienyl, benzoxadiazolyl, benzofuranyl, benzimidazolyl, benzotriazolyl, cinnolyl, indazolyl, indolyl, isoquinolyl, isothiazolyl, naphthyridinyl, purinyl, thienopyridinyl, pyrido[2,3-d]pyrimidinyl, pyrrolo[2,3-b]pyridinyl, quinazolinyl, quinolyl, thieno[2,3-c]pyridinyl, pyrazolo[3,4-b]pyridinyl, pyrazolo[3,4-c]pyridinyl, pyrazolo[4,3-c]pyridine, pyrazolo[4,3-b]pyridinyl, tetrazolyl, chromane, 2,3-dihydrobenzo[b][1,4]dioxine, benzo[d][1,3]dioxole, 2,3-dihydrobenzofuran, 2,3-dihydrobenzo[b][1,4]oxathiine, and others.

[0026] As used herein, “amide” includes both $\text{RNR}'\text{CO}-$ (in the case of $\text{R} = \text{alkyl}$, alkaminocarbonyl-) and $\text{RCONR}'-$ (in the case of $\text{R} = \text{alkyl}$, alkylcarbonylamino-). In both cases, $\text{R}' = \text{H}$ or alkyl. In some embodiments, R' is H.

[0027] As used herein, the term “ester” includes both $\text{ROCO}-$ (in the case of $\text{R} = \text{alkyl}$, alkoxycarbonyl-) and $\text{RCOO}-$ (in the case of $\text{R} = \text{alkyl}$, alkylcarbonyloxy-).

[0028] As used herein, “acyl” means an $\text{H-CO}-$ or $\text{alkyl-CO}-$, carbocyclyl- $\text{CO}-$, aryl- $\text{CO}-$, heteroaryl- $\text{CO}-$ or heterocyclyl- $\text{CO}-$ group wherein the alkyl, carbocyclyl, aryl or heterocyclyl group is as herein described. In some embodiments, acyls contain a

lower alkyl. Exemplary alkyl acyl groups include formyl, acetyl, propanoyl, 2-methylpropanoyl, t-butylacetyl, butanoyl, and palmitoyl.

[0029] As used herein, "halo", "halide" or "halogen" is a chloro, bromo, fluoro, or iodo atom radical. In some embodiments, the halide is fluorine.

[0030] As used herein, "haloalkyl" means a hydrocarbon substituent, which is a linear or branched or cyclic alkyl, substituted with chloro, bromo, fluoro, or iodo atom(s). In some embodiments, haloalkyls are fluoroalkyls, wherein one or more of the hydrogen atoms have been substituted by fluoro. Haloalkyls can be of 1 to about 3 carbons in length, (e.g., 1 to about 2 carbons). In some embodiments, the haloalkyls are 1 carbon in length.

[0031] As used herein, "heterocyclyl" means a nonaromatic cyclic ring system comprising at least one heteroatom in the ring system backbone. Heterocyclyls may include multiple fused rings. Heterocyclyls may be substituted or unsubstituted with one or more substituents, e.g., alkyl, halide, alkoxy, acyloxy, amino, amido, cyano, nitro, hydroxyl, mercapto, carboxy, carbonyl, benzyloxy, aryl, heteroaryl, and other substituents, and are attached to other groups via any available valence (e.g., any available carbon or nitrogen). In some embodiments, heterocycles have 5-7 members. In six membered monocyclic heterocycles, the heteroatom(s) are selected from one up to three of O, N or S, and wherein when the heterocycle is five membered, it can have one or two heteroatoms selected from O, N, or S. Examples of heterocyclyl include aziriny, aziridinyl, azetidiny, oxetanyl, thietanyl, 1,4,2-dithiazolyl, dihydropyridinyl, 1,3-dioxanyl, 1,4-dioxanyl, 1,3-dioxolanyl, morpholinyl, thiomorpholinyl, piperazinyl, pyranyl, pyrrolidinyl, tetrahydrofuryl, tetrahydropyridinyl, oxazinyl, thiazinyl, thiinyl, thiazolidinyl, isothiazolidinyl, oxazolidinyl, isoxazolidinyl, piperidinyl, pyrazolidinyl, imidazolidinyl, thiomorpholinyl, and others.

[0032] As used herein, "substituted amino" means an amino radical which is substituted by one or two alkyl, cycloalkyl, aryl, heteroaryl, or heterocyclyl groups, wherein the alkyl, cycloalkyl, aryl, heteroaryl or heterocyclyl are defined as above.

[0033] As used herein, "substituted thiol" refers to an RS- group wherein R is an alkyl, a cycloalkyl, an aryl, heteroaryl, or a heterocyclyl group, wherein the alkyl, cycloalkyl, aryl, heteroaryl, or heterocyclyl are defined as above.

[0034] As used herein, when two groups are indicated to be “linked” or “bonded” to form a “ring”, it is to be understood that a bond is formed between the two groups and may involve replacement of a hydrogen atom on one or both groups with the bond, thereby forming a carbocyclyl, heterocyclyl, aryl, or heteroaryl ring. The skilled artisan will recognize that such rings can and are readily formed by routine chemical reactions, and it is within the purview of the skilled artisan to both envision such rings and the methods of their formations. In some embodiments, rings have from 3-7 members (e.g., 5 or 6 members). As used herein the term “ring” or “rings” when formed by the combination of two radicals refers to heterocyclic, carbocyclic, aryl, or heteroaryl rings.

[0035] The skilled artisan will recognize that some structures described herein may be resonance forms or tautomers of compounds that may be fairly represented by other chemical structures, even when kinetically, the artisan recognizes that such structures are only a very small portion of a sample of such compound(s). Such compounds are clearly contemplated within the scope of this disclosure, though such resonance forms or tautomers are not represented herein.

[0036] The compounds provided herein may encompass various stereochemical forms. The compounds also encompass diastereomers as well as optical isomers, e.g. mixtures of enantiomers including racemic mixtures, as well as individual enantiomers and diastereomers, which arise as a consequence of structural asymmetry in certain compounds. Separation of the individual isomers or selective synthesis of the individual isomers is accomplished by application of various methods which are well known to practitioners in the art. Unless otherwise indicated, when a disclosed compound is named or depicted by a structure without specifying the stereochemistry and has one or more chiral centers, it is understood to represent all possible stereoisomers of the compound.

[0037] The term “administration” or “administering” refers to a method of giving a dosage of a compound or pharmaceutical composition provided herein to a vertebrate or invertebrate, including a mammal, a bird, a fish, or an amphibian, where the method is, e.g., orally, subcutaneously, intravenously, intranasally, topically, transdermally, intraperitoneally, intramuscularly, intrapulmonarily, vaginally, rectally,

ontologically, neuro-otologically, intraocularly, subconjunctivally, via anterior eye chamber injection, intravitreally, intraperitoneally, intrathecally, intracystically, intrapleurally, via wound irrigation, intrabuccally, intra-abdominally, intra-articularly, intra-aurally, intrabronchially, intracapsularly, intrameningeally, via inhalation, via endotracheal or endobronchial instillation, via direct instillation into pulmonary cavities, intraspinally, intrasynovially, intrathoracically, via thoracostomy irrigation, epidurally, intratympanically, intracisternally, intravascularly, intraventricularly, intraosseously, via irrigation of infected bone, or via application as part of any admixture with a prosthetic device. The method of administration can vary depending on various factors, e.g., the components of the pharmaceutical composition, the site of the disease, the disease involved, and the severity of the disease.

[0038] A “diagnostic” as used herein is a compound, method, system, or device that assists in the identification and characterization of a health or disease state. The diagnostic can be used in standard assays as is known in the art.

[0039] The term “mammal” is used in its usual biological sense. Thus, it specifically includes humans, cattle, horses, dogs, cats, mice, rats, sheeps, pigs, goats, and non-human primates, but also includes many other species.

[0040] The term “pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” includes any and all solvents, co-solvents, complexing agents, dispersion media, coatings, isotonic and absorption delaying agents and the like which are not biologically or otherwise undesirable. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions. In addition, various adjuvants such as are commonly used in the art may be included. These and other such compounds are described in the literature, e.g., in the Merck Index, Merck & Company, Rahway, NJ. Considerations for the inclusion of various components in pharmaceutical compositions are described, e.g., in Gilman *et al.* (Eds.) (2010); Goodman and Gilman’s: The Pharmacological Basis of Therapeutics, 12th Ed., The McGraw-Hill Companies.

[0041] The term “pharmaceutically acceptable salt” refers to salts that retain the biological effectiveness and properties of the compounds provided herein and, which are not biologically or otherwise undesirable. In many cases, the compounds provided herein are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto. Pharmaceutically acceptable acid addition salts can be formed with inorganic acids and organic acids. Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. Organic acids from which salts can be derived include, for example, acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like. Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases. Inorganic bases from which salts can be derived include, for example, sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum, and the like. In some embodiments, the bases can be derived from the ammonium, potassium, sodium, calcium, and magnesium salts. Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, basic ion exchange resins, and the like, specifically such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine. Many such salts are known in the art, as described in WO 87/05297.

[0042] “Solvate” refers to the compound formed by the interaction of a solvent and a compound provided herein, a metabolite, or salt thereof. Suitable solvates are pharmaceutically acceptable solvates including hydrates.

[0043] “Patient” or “subject” as used herein, means a human or a non-human mammal, e.g., a dog, a cat, a mouse, a rat, a cow, a sheep, a pig, a goat, a non-human primate or a bird, e.g., a chicken, as well as any other vertebrate or invertebrate.

[0044] By “therapeutically effective amount” or “pharmaceutically effective amount” of a compound as provided herein is one which is sufficient to achieve the desired effect and may vary according to the nature and severity of the disease condition,

and the potency of the compound. "Therapeutically effective amount" is also intended to include one or more of the compounds of Formulas (I) and/or (Ia) in combination with one or more other agents that are effective to inhibit Wnt related diseases and/or conditions. In some embodiments, the combination of compounds is a synergistic combination. Synergy, as described, for example, by Chou and Talalay, *Advances in Enzyme Regulation* (1984), 22, 27-55, occurs when the effect of the compounds when administered in combination is greater than the additive effect of the compounds when administered alone as a single agent. In general, a synergistic effect is most clearly demonstrated at sub-optimal concentrations of the compounds. It will be appreciated that different concentrations may be employed for prophylaxis than for treatment of an active disease. This amount can further depend upon the patient's height, weight, sex, age and medical history.

[0045] A therapeutic effect relieves, to some extent, one or more of the symptoms of the disease, and can include curing a disease. "Curing" means that the symptoms of an active disease are eliminated. However, certain long-term or permanent effects of the disease may exist even after a cure is obtained (such as extensive tissue damage).


[0046] "Treat," "treatment," or "treating," as used herein refers to administering a compound or pharmaceutical composition as provided herein for therapeutic purposes. The term "therapeutic treatment" refers to administering treatment to a patient already suffering from a disease thus causing a therapeutically beneficial effect, such as ameliorating existing symptoms, preventing additional symptoms, ameliorating or preventing the underlying metabolic causes of symptoms, postponing or preventing the further development of a disorder and/or reducing the severity of symptoms that will or are expected to develop.

Compounds

[0047] The compounds and compositions described herein can be used as anti-proliferative agents, e.g., anti-cancer and anti-angiogenesis agents, and/or as inhibitors of the Wnt signaling pathway, e.g., for treating diseases or disorders associated with aberrant Wnt signaling. In addition, the compounds can be used as inhibitors of one

or more kinases, kinase receptors, or kinase complexes. Such compounds and compositions are also useful for controlling cellular proliferation, differentiation, and/or apoptosis.

[0048] Some embodiments of the present disclosure include compounds, salts, pharmaceutically acceptable salts or pro-drug thereof of Formula (I):



I

[0049] In some embodiments of Formula I, R^1 is $-\text{heteroaryl}R^3R^4$.

[0050] In some embodiments of Formula I, R^2 is selected from the group consisting of H, halide, C_{1-3} alkyl, $-\text{CN}$, $-\text{OR}^8$, $-\text{OH}$, $-(C_{1-3} \text{ alkyl})\text{OR}^8$, $-\text{NR}^9R^{10}$, $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$, $-\text{heteroaryl}R^5$, $-\text{heterocyclyl}R^6$ and $-\text{aryl}R^7$.

[0051] In some embodiments of Formula I, R^3 is 1 substituent attached to the heteroaryl ring and is selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, $-\text{NR}^9R^{10}$, $-\text{NHC}(=\text{O})R^8$, $-(C_{1-3} \text{ alkyl})\text{heterocyclyl}R^6$ and $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$.

[0052] In some embodiments of Formula I, R^4 is 1 substituent attached to the heteroaryl ring and is selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{OR}^8$, $-\text{OH}$, $-(C_{1-3} \text{ alkyl})\text{OR}^8$, $-\text{NR}^9R^{10}$, $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$ and $-\text{OCF}_3$.

[0053] In some embodiments of Formula I, R^5 is 1-3 substituents attached to the heteroaryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{OR}^8$, $-\text{OH}$, $-(C_{1-3} \text{ alkyl})\text{OR}^8$, $-\text{NR}^9R^{10}$, $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$ and $-\text{OCF}_3$.

[0054] In some embodiments of Formula I, R^6 is 1-3 substituents attached to the heterocyclyl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{OR}^8$, $-\text{OH}$, $-(C_{1-3} \text{ alkyl})\text{OR}^8$, $-\text{NR}^9R^{10}$, $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$ and $-\text{OCF}_3$.

[0055] In some embodiments of Formula I, R^7 is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-OR^8$, $-OH$, $-(C_{1-3} \text{ alkyl})OR^8$, $-NR^9R^{10}$, $-(C_{1-3} \text{ alkyl})NR^9R^{10}$ and $-OCF_3$.

[0056] In some embodiments of Formula I, each R^8 is selected from the group consisting of C_{1-9} alkyl, $-\text{heteroaryl}R^{12}$, $-\text{heterocyclyl}R^{13}$, $-\text{aryl}R^{14}$, $\text{carbocyclyl}R^{11}$, $-(C_{1-3} \text{ alkyl})\text{heteroaryl}R^{12}$, $-(C_{1-3} \text{ alkyl})\text{heterocyclyl}R^{13}$, $-(C_{1-3} \text{ alkyl})\text{aryl}R^{14}$ and $-(C_{1-3} \text{ alkyl})\text{carbocyclyl}R^{11}$.

[0057] In some embodiments of Formula I, each R^9 is selected from the group consisting of H, C_{1-6} alkyl, $-\text{heteroaryl}R^{12}$, $-\text{heterocyclyl}R^{13}$, $-\text{aryl}R^{14}$, $\text{carbocyclyl}R^{11}$, $-(C_{1-3} \text{ alkyl})\text{heteroaryl}R^{12}$, $-(C_{1-3} \text{ alkyl})\text{heterocyclyl}R^{13}$, $-(C_{1-3} \text{ alkyl})\text{aryl}R^{14}$ and $-(C_{1-3} \text{ alkyl})\text{carbocyclyl}R^{11}$.

[0058] In some embodiments of Formula I, each R^{10} is selected from the group consisting of H and C_{1-6} alkyl.

[0059] In some embodiments of Formula I, R^9 and R^{10} are linked to form a five or six membered heterocyclyl ring.

[0060] In some embodiments of Formula I, R^{11} is 1-3 substituents attached to the carbocyclyl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;


[0061] In some embodiments of Formula I, R^{12} is 1-3 substituents attached to the heteroaryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;


[0062] In some embodiments of Formula I, R^{13} is 1-3 substituents attached to the heterocyclyl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;


[0063] In some embodiments of Formula I, R^{14} is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3}

alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{O}(\text{R}^{10})$, $-(\text{C}_{1-3} \text{ alkyl})\text{OR}^{10}$, $-\text{N}(\text{R}^{10})_2$, $-(\text{C}_{1-3} \text{ alkyl})\text{N}(\text{R}^{10})_2$ and $-\text{OCF}_3$;

[0064] In some embodiments of Formula I, there is the proviso that a compound of Formula I is not a compound selected from the group consisting of:





**Ia**

[0066] In some embodiments of Formula **Ia**, R^1 is $-\text{pyridine}R^3R^4$.

[0067] In some embodiments of Formula **Ia**, R^2 is selected from the group consisting of H, halide, C_{1-3} alkyl, $-\text{CN}$, $-\text{OR}^8$, $-\text{OH}$, $-(C_{1-3} \text{ alkyl})\text{OR}^8$, $-\text{NR}^9R^{10}$, $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$, $-\text{heteroaryl}R^5$, $-\text{heterocyclyl}R^6$ and $-\text{aryl}R^7$.

[0068] In some embodiments of Formula **Ia**, R^3 is 1 substituent attached to the heteroaryl ring and is selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, $-\text{NR}^9R^{10}$, $-\text{NHC}(=\text{O})R^8$, $-(C_{1-3} \text{ alkyl})\text{heterocyclyl}R^6$ and $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$.

[0069] In some embodiments of Formula **Ia**, R^4 is H.

[0070] In some embodiments of Formula **Ia**, R^5 is 1-3 substituents attached to the heteroaryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{OR}^8$, $-\text{OH}$, $-(C_{1-3} \text{ alkyl})\text{OR}^8$, $-\text{NR}^9R^{10}$, $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$ and $-\text{OCF}_3$.

[0071] In some embodiments of Formula **Ia**, R^6 is 1-3 substituents attached to the heterocyclyl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{OR}^8$, $-\text{OH}$, $-(C_{1-3} \text{ alkyl})\text{OR}^8$, $-\text{NR}^9R^{10}$, $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$ and $-\text{OCF}_3$.

[0072] In some embodiments of Formula **Ia**, R^7 is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{OR}^8$, $-\text{OH}$, $-(C_{1-3} \text{ alkyl})\text{OR}^8$, $-\text{NR}^9R^{10}$, $-(C_{1-3} \text{ alkyl})\text{NR}^9R^{10}$ and $-\text{OCF}_3$.

[0073] In some embodiments of Formula **Ia**, each R^8 is selected from the group consisting of C_{1-9} alkyl, $-\text{heteroaryl}R^{12}$, $-\text{heterocyclyl}R^{13}$, $-\text{aryl}R^{14}$, $\text{carbocyclyl}R^{11}$, $-(C_{1-3} \text{ alkyl})\text{heteroaryl}R^{12}$, $-(C_{1-3} \text{ alkyl})\text{heterocyclyl}R^{13}$, $-(C_{1-3} \text{ alkyl})\text{aryl}R^{14}$ and $-(C_{1-3} \text{ alkyl})\text{carbocyclyl}R^{11}$.

[0074] In some embodiments of Formula **Ia**, each R^9 is selected from the group consisting of H, C_{1-6} alkyl, $-\text{heteroaryl}R^{12}$, $-\text{heterocyclyl}R^{13}$, $-\text{aryl}R^{14}$, $\text{carbocyclyl}R^{11}$, $-(C_{1-3} \text{ alkyl})\text{heteroaryl}R^{12}$, $-(C_{1-3} \text{ alkyl})\text{heterocyclyl}R^{13}$, $-(C_{1-3} \text{ alkyl})\text{aryl}R^{14}$ and $-(C_{1-3} \text{ alkyl})\text{carbocyclyl}R^{11}$.

[0075] In some embodiments of Formula **Ia**, each R^{10} is selected from the group consisting of H and C_{1-6} alkyl.

[0076] In some embodiments of Formula **Ia**, R^9 and R^{10} are linked to form a five or six membered heterocyclyl ring.


[0077] In some embodiments of Formula **Ia**, R^{11} is 1-3 substituents attached to the carbocyclyl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{O}(R^{10})$, $-(C_{1-3} \text{ alkyl})\text{OR}^{10}$, $-\text{N}(R^{10})_2$, $-(C_{1-3} \text{ alkyl})\text{N}(R^{10})_2$ and $-\text{OCF}_3$;

[0078] In some embodiments of Formula **Ia**, R^{12} is 1-3 substituents attached to the heteroaryl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{O}(R^{10})$, $-(C_{1-3} \text{ alkyl})\text{OR}^{10}$, $-\text{N}(R^{10})_2$, $-(C_{1-3} \text{ alkyl})\text{N}(R^{10})_2$ and $-\text{OCF}_3$;

[0079] In some embodiments of Formula **Ia**, R^{13} is 1-3 substituents attached to the heterocyclyl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{O}(R^{10})$, $-(C_{1-3} \text{ alkyl})\text{OR}^{10}$, $-\text{N}(R^{10})_2$, $-(C_{1-3} \text{ alkyl})\text{N}(R^{10})_2$ and $-\text{OCF}_3$;

[0080] In some embodiments of Formula **Ia**, R^{14} is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-\text{CF}_3$, halide, $-\text{CN}$, $-\text{O}(R^{10})$, $-(C_{1-3} \text{ alkyl})\text{OR}^{10}$, $-\text{N}(R^{10})_2$, $-(C_{1-3} \text{ alkyl})\text{N}(R^{10})_2$ and $-\text{OCF}_3$;

[0081] In other embodiments of Formula **Ia**, there is the proviso that a compound of Formula **Ia** is not a compound selected from the group consisting of:



[0082] In some embodiments of Formula **I**, R^1 is pyridine R^3R^4 .

- [0083] In some embodiments of Formulas **I** and/or **Ia**, R^1 is pyridin-3-yl R^3R^4 .
- [0084] In some embodiments of Formulas **I** and/or **Ia**, R^3 and R^4 are both H.
- [0085] In some embodiments of Formulas **I** and/or **Ia**, R^3 is $-C_{1-2}$ alkyl and R^4 is H.
- [0086] In some embodiments of Formulas **I** and/or **Ia**, R^3 is methyl.
- [0087] In some embodiments of Formulas **I** and/or **Ia**, R^3 is ethyl.
- [0088] In some embodiments of Formulas **I** and/or **Ia**, R^3 is $-\text{CH}_2\text{NR}^9\text{R}^{10}$.
- [0089] In some embodiments of Formula **I**, R^4 is H.
- [0090] In some embodiments of Formulas **I** and/or **Ia**, R^9 is $-C_{1-2}$ alkyl; R^{10} is $-(C_{1-2}$ alkyl); and R^4 is H.
- [0091] In some embodiments of Formulas **I** and/or **Ia**, R^4 and R^9 are both H; and R^{10} is selected from the group consisting of $-C_{1-2}$ alkyl, $-\text{CH}_2\text{phenyl}$ and $-\text{CH}_2\text{carbocyclyl}$.
- [0092] In some embodiments of Formulas **I** and/or **Ia**, R^9 is $-\text{CH}_2\text{cyclopropyl}$.
- [0093] In some embodiments of Formulas **I** and/or **Ia**, R^9 is $-\text{CH}_2\text{cyclobutyl}$.
- [0094] In some embodiments of Formulas **I** and/or **Ia**, R^9 is $-\text{CH}_2\text{cyclopentyl}$.
- [0095] In some embodiments of Formulas **I** and/or **Ia**, R^9 is $-\text{CH}_2\text{cyclohexyl}$.
- [0096] In some embodiments of Formulas **I** and/or (**Ia**), R^9 and R^{10} are linked to form a five or six membered heterocyclyl ring and R^4 is H.
- [0097] In some embodiments of Formulas **I** and/or **Ia**, R^9 and R^{10} are linked to form a morpholine ring.
- [0098] In some embodiments of Formulas **I** and/or **Ia**, R^9 and R^{10} are linked to form a piperidine ring.
- [0099] In some embodiments of Formulas **I** and/or **Ia**, R^9 and R^{10} are linked to form a pyrrolidine ring.
- [0100] In some embodiments of Formulas **I** and/or **Ia**, R^9 and R^{10} are linked to form a piperazine ring.
- [0101] In some embodiments of Formulas **I** and/or **Ia**, R^3 is $-\text{NHC}(=\text{O})\text{R}^8$.
- [0102] In some embodiments of Formulas **I** and/or **Ia**, R^4 is H; and R^8 is selected from the group consisting of $-C_{1-5}$ alkyl, $-\text{CH}_2\text{phenyl}$, phenyl and $-\text{carbocyclyl}$.

[0103] In another aspect of embodiments of Formulas **I** and/or **Ia**, R^8 is a $-C_{1-5}$ alkyl. For example, the $-C_{1-5}$ alkyl can be selected from the group consisting of methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, *tert*-butyl, n-pentyl, iso-pentyl and neo-pentyl. In certain embodiments, R^8 is methyl. In another embodiment, R^8 is ethyl. In another embodiment, R^8 is n-propyl. In another embodiment, R^8 is iso-propyl. In another embodiment, R^8 is n-butyl. In another embodiment, R^8 is iso-butyl. In another embodiment, R^8 is sec-butyl. In another embodiment, R^8 is *tert*-butyl. In another embodiment, R^8 is n-pentyl. In another embodiment, R^8 is iso-pentyl. In another embodiment, R^8 is neo-pentyl.

[0104] In some embodiments of Formulas **I** and/or **Ia**, R^8 is $-\text{CH}_2\text{phenyl}$.

[0105] In some embodiments of Formulas **I** and/or **Ia**, R^8 is phenyl.

[0106] In some embodiments of Formulas **I** and/or **Ia**, R^8 is a carbocyclyl. For example, the carbocyclyl can be selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. In certain embodiments, R^8 is cyclopropyl. In another embodiment, R^8 is cyclobutyl. In another embodiment, R^8 is cyclopentyl. In another embodiment, R^8 is cyclohexyl.

[0107] In some embodiments of Formulas **I** and/or **Ia**, R^3 is $-\text{NR}^9\text{R}^{10}$.

[0108] In some embodiments of Formulas **I** and/or **Ia**, R^3 is $-\text{NH}_2$ and R^4 is H.

[0109] In some embodiments of Formulas **I** and/or **Ia**, R^9 is $-C_{1-2}$ alkyl; R^{10} is $-C_{1-2}$ alkyl; and R^4 is H.

[0110] In some embodiments of Formulas **I** and/or **Ia**, R^9 is methyl and R^{10} is methyl.

[0111] In some embodiments of Formulas **I** and/or **Ia**, R^9 is methyl and R^{10} is ethyl.

[0112] In some embodiments of Formulas **I** and/or **Ia**, R^9 is ethyl and R^{10} is ethyl.

[0113] In some embodiments of Formulas **I** and/or **Ia**, R^4 and R^9 are both H; and R^{10} is $-C_{1-4}$ alkyl.

[0114] In some embodiments of Formulas **I** and/or **Ia**, R^{10} is a $-C_{1-4}$ alkyl. For example, the $-C_{1-4}$ alkyl can be selected from the group consisting of methyl, ethyl, n-

propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl and *tert*-butyl. In certain embodiments, R¹⁰ is methyl. In another embodiment, R¹⁰ is ethyl. In another embodiment, R¹⁰ is n-propyl. In another embodiment, R¹⁰ is iso-propyl. In another embodiment, R¹⁰ is n-butyl. In another embodiment, R¹⁰ is iso-butyl. In another embodiment, R¹⁰ is sec-butyl. In another embodiment, R¹⁰ is *tert*-butyl.

[0115] In some embodiments of Formulas **I** and/or **Ia**, R² is H.

[0116] In some embodiments of Formulas **I** and/or **Ia**, R² is -heteroarylR⁵.

[0117] In some embodiments of Formulas **I** and/or **Ia**, R⁵ is H.

[0118] In some embodiments of Formulas **I** and/or **Ia**, R² is -pyridine.

[0119] In some embodiments of Formulas **I** and/or **Ia**, R² is -pyridin-2-yl.

[0120] In some embodiments of Formulas **I** and/or **Ia**, R² is -pyridin-3-yl.

[0121] In some embodiments of Formulas **I** and/or **Ia**, R² is -pyridin-4-yl.

[0122] In some embodiments of Formulas **I** and/or **Ia**, R² is thiophene.

[0123] In some embodiments of Formulas **I** and/or **Ia**, R² is furan.

[0124] In some embodiments of Formulas **I** and/or **Ia**, R² is -imidazolylR⁵; and R⁵ is 1 substituent selected from the group consisting of H or -C₁₋₂ alkyl.

[0125] In some embodiments of Formulas **I** and/or **Ia**, R⁵ is methyl.

[0126] In some embodiments of Formulas **I** and/or **Ia**, R⁵ is ethyl.

[0127] In some embodiments of Formulas **I** and/or **Ia**, R² is a heterocyclyl.

For example, the heterocyclyl can be selected from the group consisting of morpholinyl, piperazinyl, 1-methylpiperazinyl, piperidinyl and pyrrolidinyl. In certain embodiments, R² is piperidinyl. In another embodiment, R² is piperazinyl. In another embodiment, R² is 1-methylpiperazinyl.

[0128] In some embodiments of Formulas **I** and/or **Ia**, R² is -piperazineR⁶; and R⁶ is 1 substituent selected from the group consisting of H or -C₁₋₂ alkyl.

[0129] In some embodiments of Formulas **I** and/or **Ia**, R⁶ is methyl.

[0130] In some embodiments of Formulas **I** and/or **Ia**, R⁶ is ethyl.

[0131] In some embodiments of Formulas **I** and/or **Ia**, R² is -phenylR⁷.

[0132] In some embodiments of Formulas **I** and/or **Ia**, R⁷ is 1 fluorine atom.

[0133] In some embodiments of Formulas **I** and/or **Ia**, R⁷ is 2 fluorine atoms.

[0134] In some embodiments of Formulas **I** and/or **Ia**, R^1 is pyridin-3-yl R^3R^4 ; R^2 is -phenyl R^7 ; R^3 is selected from the group consisting of H, -C₁₋₂ alkyl, -CH₂NR⁹R¹⁰, -NR⁹R¹⁰ and -NHC(=O) R^8 ; R^4 is H; R^7 is 1-2 fluorine atoms; R^8 is selected from the group consisting of -C₁₋₅ alkyl, -CH₂phenyl, phenyl and -carbocyclyl; R^9 is selected from the group consisting of -C₁₋₃ alkyl, -CH₂phenyl and -CH₂carbocyclyl; R^{10} is H or -C₁₋₂ alkyl; or R^9 and R^{10} are optionally linked to form a piperidine or pyrrolidine ring.


[0135] In some embodiments of Formulas **I** and/or **Ia**, R^1 is pyridin-3-yl R^3R^4 ; R^2 is pyridine; R^3 is selected from the group consisting of -C₁₋₂ alkyl, -CH₂NR⁹R¹⁰, -NR⁹R¹⁰ and -NHC(=O) R^8 ; R^4 is H; R^8 is selected from the group consisting of -C₁₋₅ alkyl, -CH₂phenyl and -carbocyclyl; R^9 is selected from the group consisting of H, -C₁₋₃ alkyl, -CH₂phenyl and -CH₂carbocyclyl; R^{10} is H or -C₁₋₂ alkyl; or R^9 and R^{10} are optionally linked to form a pyrrolidine ring.

[0136] In some embodiments of Formulas **I** and/or **Ia**, R^1 is pyridin-3-yl R^3R^4 ; R^2 is 1-methylpiperazinyl- or piperidinyl-; R^3 is selected from the group consisting of H, -C₁₋₂ alkyl, -CH₂NR⁹R¹⁰, -NR⁹R¹⁰ and -NHC(=O) R^8 ; R^4 is H; R^8 is selected from the group consisting of -C₁₋₅ alkyl, -CH₂phenyl, phenyl and -carbocyclyl; R^9 is H or -C₁₋₂ alkyl; R^{10} is -C₁₋₃ alkyl; or R^9 and R^{10} are optionally linked to form a piperidine or pyrrolidine ring.


[0137] In some embodiments of Formulas **I** and/or **Ia**, R^1 is pyridin-3-yl R^3R^4 ; R^2 is selected from the group consisting of furan, thiophene and -imidazolyl R^5 ; R^3 is selected from the group consisting of -CH₂NR⁹R¹⁰, -NR⁹R¹⁰ and -NHC(=O) R^8 ; R^4 is H; R^5 is H or -C₁₋₂ alkyl; R^8 is selected from the group consisting of -C₁₋₅ alkyl, -CH₂phenyl and -carbocyclyl; R^9 is selected from the group consisting of H, -C₁₋₃ alkyl and -CH₂phenyl; R^{10} is H or -C₁₋₂ alkyl; or R^9 and R^{10} are optionally linked to form a piperidine or pyrrolidine ring.

[0138] In some embodiments of Formulas **I** and/or **Ia**, R^1 is pyridin-3-yl R^3R^4 ; R^2 is H; R^3 is selected from the group consisting of -C₁₋₂ alkyl, -CH₂NR⁹R¹⁰ and -NHC(=O) R^8 ; R^4 is H; R^8 is selected from the group consisting of -C₁₋₅ alkyl, phenyl and -carbocyclyl; R^9 is -CH₂phenyl; and R^{10} is H.

[0139] In some embodiments of Formulas **I** and/or **Ia**, R^2 is selected from the group consisting of:





















[0140] In some embodiments of Formulas I and/or Ia, R³ is selected from the group consisting of:


















[0141] Illustrative compounds of Formulas I and/or Ia are shown in Table 1.
















Table 1.



















1		2		3	
---	---	---	--	---	---
















4		5		6	
7		8		9	
10		11		12	
13		14		15	
16		17		18	
















19		20		21	
22		23		24	
25		26		27	
28		29		30	
31		32		33	
















34		35		36	
37		38		39	
40		41		42	
43		44		45	
46		47		48	
















49		50		51	
52		53		54	
55		56		57	
58		59		60	
61		62		63	
















64		65		66	
67		68		69	
70		71		72	
73		74		75	
76		77		78	
79		80		81	
















82		83		84	
85		86		87	
88		89		90	
91		92		93	
94		95		96	
















97		98		99	
100		101		102	
103		104		105	
106		107		108	
109		110		111	
















112		113		114	
115		116		117	
118		119		120	
121		122		123	
124		125		126	
















127		128		129	
130		131		132	
133		134		135	
136		137		138	
139		140		141	
















142		143		144	
145		146		147	
148		149		150	
151		152		153	
154		155		156	
















157		158		159	
160		161		162	
163		164		165	
166		167		168	
169		170		171	
















172		173		174	
175		176		177	
178		179		180	
181		182		183	
184		185		186	



















187		188		189	
190		191		192	
193		194		195	
196		197		198	
199		200		201	










202		203		204	
205		206		207	
208		209		210	
211		212		213	
214		215		216	

217		218		219	
220		221		222	
223		224		225	
226		227		228	
229		230		231	

232		233		234	
235		236		237	
238		239		240	
241		242		243	
244		245		246	

247		248		249	
250		251		252	
253		254		255	
256		257		258	
259		260		261	

262		263		264	
265		266		267	
268		269		270	
271		272		273	
274		275		276	
277		278		279	

280		281		282	
283		284		285	
286		287		288	

Administration and Pharmaceutical Compositions

[0142] Some embodiments include pharmaceutical compositions comprising:
 (a) a therapeutically effective amount of a compound provided herein, or its corresponding enantiomer, diastereoisomer or tautomer, or pharmaceutically acceptable salt; and (b) a pharmaceutically acceptable carrier.

[0143] The compounds provided herein may also be useful in combination (administered together or sequentially) with other known agents.

[0144] Non-limiting examples of diseases which can be treated with a combination of a compound of Formula (I) or (Ia) and other known agents are colorectal cancer, ovarian cancer, diabetic retinopathy, pulmonary fibrosis, and osteoarthritis.

[0145] In some embodiments, colorectal cancer can be treated with a combination of a compound of either Formula (I) or (Ia) and one or more of the

following drugs: 5-Fluorouracil (5-FU), which can be administered with the vitamin-like drug leucovorin (also called folinic acid); capecitabine (Xeloda[®]), irinotecan (Camptosar[®]), oxaliplatin (Eloxatin[®]). Examples of combinations of these drugs which could be further combined with a compound of either Formula (I) or (Ia) are FOLFOX (5-FU, leucovorin, and oxaliplatin), FOLFIRI (5-FU, leucovorin, and irinotecan), FOLFOXIRI (leucovorin, 5-FU, oxaliplatin, and irinotecan) and CapeOx (Capecitabine and oxaliplatin). For rectal cancer, chemo with 5-FU or capecitabine combined with radiation may be given before surgery (neoadjuvant treatment).

[0146] In some embodiments, ovarian cancer can be treated with a combination of a compound of either Formula (I) or (Ia) and one or more of the following drugs: Topotecan, Liposomal doxorubicin (Doxil[®]), Gemcitabine (Gemzar[®]), Cyclophosphamide (Cytoxan[®]), Vinorelbine (Navelbine[®]), Ifosfamide (Ifex[®]), Etoposide (VP-16), Altretamine (Hexalen[®]), Capecitabine (Xeloda[®]), Irinotecan (CPT-11, Camptosar[®]), Melphalan, Pemetrexed (Alimta[®]) and Albumin bound paclitaxel (nab-paclitaxel, Abraxane[®]). Examples of combinations of these drugs which could be further combined with a compound of either Formula (I) or (Ia) are TIP (paclitaxel [Taxol], ifosfamide, and cisplatin), VeIP (vinblastine, ifosfamide, and cisplatin) and VIP (etoposide [VP-16], ifosfamide, and cisplatin).

[0147] In some embodiments, a compound of either Formula (I) or (Ia) can be used to treat cancer in combination with any of the following methods: (a) Hormone therapy such as aromatase inhibitors, LHRH [luteinizing hormone-releasing hormone] analogs and inhibitors, and others; (b) Ablation or embolization procedures such as radiofrequency ablation (RFA), ethanol (alcohol) ablation, microwave thermotherapy and cryosurgery (cryotherapy); (c) Chemotherapy using alkylating agents such as cisplatin and carboplatin, oxaliplatin, mechlorethamine, cyclophosphamide, chlorambucil and ifosfamide; (d) Chemotherapy using anti-metabolites such as azathioprine and mercaptopurine; (e) Chemotherapy using plant alkaloids and terpenoids such as vinca alkaloids (i.e. Vincristine, Vinblastine, Vinorelbine and Vindesine) and taxanes; (f) Chemotherapy using podophyllotoxin, etoposide, teniposide and docetaxel; (g) Chemotherapy using topoisomerase inhibitors such as irinotecan, topotecan, amsacrine, etoposide, etoposide phosphate, and teniposide; (h) Chemotherapy using cytotoxic

antibiotics such as actinomycin, anthracyclines, doxorubicin, daunorubicin, valrubicin, idarubicin, epirubicin, bleomycin, plicamycin and mitomycin; (i) Chemotherapy using tyrosine-kinase inhibitors such as Imatinib mesylate (Gleevec[®], also known as STI-571), Gefitinib (Iressa, also known as ZD1839), Erlotinib (marketed as Tarceva[®]), Bortezomib (Velcade[®]), tamoxifen, tofacitinib, crizotinib, Bcl-2 inhibitors (e.g. obatoclax in clinical trials, ABT-263, and Gossypol), PARP inhibitors (e.g. Iniparib, Olaparib in clinical trials), PI3K inhibitors (e.g. perifosine in a phase III trial), VEGF Receptor 2 inhibitors (e.g. Apatinib), AN-152, (AEZS-108), Braf inhibitors (e.g. vemurafenib, dabrafenib and LGX818), MEK inhibitors (e.g. trametinib and MEK162), CDK inhibitors, (e.g. PD-0332991), salinomycin and Sorafenib; (j) Chemotherapy using monoclonal antibodies such as Rituximab (marketed as MabThera[®] or Rituxan[®]), Trastuzumab (Herceptin also known as ErbB2), Cetuximab (marketed as Erbitux[®]), and Bevacizumab (marketed as Avastin[®]); and (k) radiation therapy.

[0148] In some embodiments, diabetic retinopathy can be treated with a combination of a compound of either Formula (I) or (Ia) and one or more of the following natural supplements: Bilberry, Butcher's broom, Ginkgo, Grape seed extract, and Pycnogenol (Pine bark).

[0149] In some embodiments, a compound of either Formula (I) or (Ia) can be used to treat pulmonary fibrosis in combination with any of the following methods: oxygen therapy, pulmonary rehabilitation, and surgery.

[0150] In some embodiments, a compound of either Formula (I) or (Ia) can be used to treat osteoarthritis in combination with any of the following methods: (a) Nonsteroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen, naproxen and acetaminophen; (b) physical therapy; (c) injections of corticosteroid medications; and (d) injections of hyaluronic acid derivatives (e.g. Hyalgan, Synvisc).

[0151] Administration of the compounds disclosed herein or the pharmaceutically acceptable salts thereof can be via any of the accepted modes of administration, including, but not limited to, orally, subcutaneously, intravenously, intranasally, topically, transdermally, intraperitoneally, intramuscularly, intrapulmonarily, vaginally, rectally, otopologically, neuro-otologically, intraocularly, subconjunctivally, via anterior eye chamber injection, intravitreally, intraperitoneally,

intrathecally, intracystically, intrapleurally, via wound irrigation, intrabuccally, intra-abdominally, intra-articularly, intra-aurally, intrabronchially, intracapsularly, intrameningeally, via inhalation, via endotracheal or endobronchial instillation, via direct instillation into pulmonary cavities, intraspinally, intrasynovially, intrathoracically, via thoracostomy irrigation, epidurally, intratympanically, intracisternally, intravascularly, intraventricularly, intraosseously, via irrigation of infected bone, or via application as part of any admixture with a prosthetic devices. In some embodiments, the administration method includes oral or parenteral administration.

[0152] Compounds provided herein intended for pharmaceutical use may be administered as crystalline or amorphous products. Pharmaceutically acceptable compositions may be solid, semi-solid, liquid, solutions, colloidal, liposomes, emulsions, suspensions, complexes, coacervates, and aerosols. Dosage forms, such as, e.g., tablets, capsules, powders, liquids, suspensions, suppositories, aerosols, implants, controlled release or the like are provided herein. They may be obtained, for example, as solid plugs, powders, or films by methods such as precipitation, crystallization, milling, grinding, supercritical fluid processing, coacervation, complex coacervation, encapsulation, emulsification, complexation, freeze drying, spray drying, or evaporative drying. Microwave or radio frequency drying may be used for this purpose. The compounds can also be administered in sustained or controlled release dosage forms, including depot injections, osmotic pumps, pills (tablets and or capsules), transdermal (including electrotransport) patches, implants, and the like, for prolonged and/or timed, pulsed administration at a predetermined rate.

[0153] The compounds can be administered either alone or in combination with a conventional pharmaceutical carrier, excipient or the like. Pharmaceutically acceptable excipients include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems (SEDDS) such as d- α -tocopherol polyethylene glycol 1000 succinate, surfactants used in pharmaceutical dosage forms such as Tweens, poloxamers or other similar polymeric delivery matrices, serum proteins, such as human serum albumin, buffer substances such as phosphates, tris, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen

phosphate, potassium hydrogen phosphate, sodium-chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethyl cellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, and wool fat. Cyclodextrins such as α -, β -, and γ -cyclodextrin, or chemically modified derivatives such as hydroxyalkylcyclodextrins, including 2- and 3-hydroxypropyl- β -cyclodextrins, or other solubilized derivatives can also be used to enhance delivery of compounds described herein. Dosage forms or compositions containing a compound as described herein in the range of 0.005% to 100% with the balance made up from non-toxic carrier may be prepared. The contemplated compositions may contain 0.001%-100% of a compound provided herein, in one embodiment 0.1-95%, in another embodiment 75-85%, in a further embodiment 20-80%. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see *Remington: The Science and Practice of Pharmacy*, 22nd Edition (Pharmaceutical Press, London, UK. 2012).

[0154] In one embodiment, the compositions will take the form of a unit dosage form such as a pill or tablet and thus the composition may contain, along with a compound provided herein, a diluent such as lactose, sucrose, dicalcium phosphate, or the like; a lubricant such as magnesium stearate or the like; and a binder such as starch, gum acacia, polyvinylpyrrolidone, gelatin, cellulose, cellulose derivatives or the like. In another solid dosage form, a powder, granule, solution or suspension (*e.g.*, in propylene carbonate, vegetable oils, PEG's, poloxamer 124 or triglycerides) is encapsulated in a capsule (gelatin or cellulose base capsule). Unit dosage forms in which one or more compounds provided herein or additional active agents are physically separated are also contemplated; *e.g.*, capsules with granules (or tablets in a capsule) of each drug; two-layer tablets; two-compartment gel caps, etc. Enteric coated or delayed release oral dosage forms are also contemplated.

[0155] Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, etc. a compound provided herein and optional pharmaceutical adjuvants in a carrier (*e.g.*, water, saline, aqueous dextrose, glycerol, glycols, ethanol or the like) to form a solution, colloid, liposome, emulsion, complexes, coacervate or suspension. If desired, the pharmaceutical composition can also contain

minor amounts of nontoxic auxiliary substances such as wetting agents, emulsifying agents, co-solvents, solubilizing agents, pH buffering agents and the like (e.g., sodium acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine acetate, triethanolamine oleate, and the like).

[0156] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 0.25 mg/Kg to about 50 mg/Kg in humans.

[0157] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 0.25 mg/Kg to about 20 mg/Kg in humans.

[0158] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 0.50 mg/Kg to about 19 mg/Kg in humans.

[0159] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 0.75 mg/Kg to about 18 mg/Kg in humans.

[0160] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 1.0 mg/Kg to about 17 mg/Kg in humans.

[0161] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 1.25 mg/Kg to about 16 mg/Kg in humans.

[0162] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 1.50 mg/Kg to about 15 mg/Kg in humans.

[0163] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 1.75 mg/Kg to about 14 mg/Kg in humans.

[0164] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 2.0 mg/Kg to about 13 mg/Kg in humans.

[0165] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 3.0 mg/Kg to about 12 mg/Kg in humans.

[0166] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 4.0 mg/Kg to about 11 mg/Kg in humans.

[0167] In some embodiments, the unit dosage of compounds of Formulas (I) and/or (Ia) is about 5.0 mg/Kg to about 10 mg/Kg in humans.

[0168] In some embodiments, the compositions are provided in unit dosage forms suitable for single administration.

[0169] In some embodiments, the compositions are provided in unit dosage forms suitable for twice a day administration.

[0170] In some embodiments, the compositions are provided in unit dosage forms suitable for three times a day administration.

[0171] Injectables can be prepared in conventional forms, either as liquid solutions, colloid, liposomes, complexes, coacervate or suspensions, as emulsions, or in solid forms suitable for reconstitution in liquid prior to injection. The percentage of a compound provided herein contained in such parenteral compositions is highly dependent on the specific nature thereof, as well as the activity of the compound and the needs of the patient. However, percentages of active ingredient of 0.01% to 10% in solution are employable, and could be higher if the composition is a solid or suspension, which could be subsequently diluted to the above percentages.

[0172] In some embodiments, the composition will comprise about 0.1-10% of the active agent in solution.

[0173] In some embodiments, the composition will comprise about 0.1-5% of the active agent in solution.

[0174] In some embodiments, the composition will comprise about 0.1-4% of the active agent in solution.

[0175] In some embodiments, the composition will comprise about 0.15-3% of the active agent in solution.

[0176] In some embodiments, the composition will comprise about 0.2-2% of the active agent in solution.

[0177] In some embodiments, the compositions are provided in dosage forms suitable for continuous dosage by intravenous infusion over a period of about 1-96 hours.

[0178] In some embodiments, the compositions are provided in dosage forms suitable for continuous dosage by intravenous infusion over a period of about 1-72 hours.

[0179] In some embodiments, the compositions are provided in dosage forms suitable for continuous dosage by intravenous infusion over a period of about 1-48 hours.

[0180] In some embodiments, the compositions are provided in dosage forms suitable for continuous dosage by intravenous infusion over a period of about 1-24 hours.

[0181] In some embodiments, the compositions are provided in dosage forms suitable for continuous dosage by intravenous infusion over a period of about 1-12 hours.

[0182] In some embodiments, the compositions are provided in dosage forms suitable for continuous dosage by intravenous infusion over a period of about 1-6 hours.

[0183] In some embodiments, these compositions can be administered by intravenous infusion to humans at doses of about 5 mg/m² to about 300 mg/m².

[0184] In some embodiments, these compositions can be administered by intravenous infusion to humans at doses of about 5 mg/m² to about 200 mg/m².

[0185] In some embodiments, these compositions can be administered by intravenous infusion to humans at doses of about 5 mg/m² to about 100 mg/m².

[0186] In some embodiments, these compositions can be administered by intravenous infusion to humans at doses of about 10 mg/m² to about 50 mg/m².

[0187] In some embodiments, these compositions can be administered by intravenous infusion to humans at doses of about 50 mg/m² to about 200 mg/m².

[0188] In some embodiments, these compositions can be administered by intravenous infusion to humans at doses of about 75 mg/m² to about 175 mg/m².

[0189] In some embodiments, these compositions can be administered by intravenous infusion to humans at doses of about 100 mg/m² to about 150 mg/m².

[0190] It is to be noted that concentrations and dosage values may also vary depending on the specific compound and the severity of the condition to be alleviated. It is to be further understood that for any particular patient, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed compositions.

[0191] In one preferred embodiment, the compositions can be administered to the respiratory tract (including nasal and pulmonary) e.g., through a nebulizer, metered-dose inhalers, atomizer, mister, aerosol, dry powder inhaler, insufflator, liquid instillation or other suitable device or technique.

[0192] In some embodiments, aerosols intended for delivery to the nasal mucosa are provided for inhalation through the nose. For optimal delivery to the nasal

cavities, inhaled particle sizes of about 5 to about 100 microns are useful, with particle sizes of about 10 to about 60 microns being preferred. For nasal delivery, a larger inhaled particle size may be desired to maximize impaction on the nasal mucosa and to minimize or prevent pulmonary deposition of the administered formulation. In some embodiments, aerosols intended for delivery to the lung are provided for inhalation through the nose or the mouth. For delivery to the lung, inhaled aerodynamic particle sizes of about less than 10 μm are useful (e.g., about 1 to about 10 microns). Inhaled particles may be defined as liquid droplets containing dissolved drug, liquid droplets containing suspended drug particles (in cases where the drug is insoluble in the suspending medium), dry particles of pure drug substance, drug substance incorporated with excipients, liposomes, emulsions, colloidal systems, coacervates, aggregates of drug nanoparticles, or dry particles of a diluent which contain embedded drug nanoparticles.

[0193] In some embodiments, compounds of Formulas (I) and/or (Ia) disclosed herein intended for respiratory delivery (either systemic or local) can be administered as aqueous formulations, as non-aqueous solutions or suspensions, as suspensions or solutions in halogenated hydrocarbon propellants with or without alcohol, as a colloidal system, as emulsions, coacervates, or as dry powders. Aqueous formulations may be aerosolized by liquid nebulizers employing either hydraulic or ultrasonic atomization or by modified micropump systems (like the soft mist inhalers, the Aerodose[®] or the AERx[®] systems). Propellant-based systems may use suitable pressurized metered-dose inhalers (pMDIs). Dry powders may use dry powder inhaler devices (DPIs), which are capable of dispersing the drug substance effectively. A desired particle size and distribution may be obtained by choosing an appropriate device.

[0194] In some embodiments, the compositions of Formulas (I) and/or (Ia) disclosed herein can be administered to the ear by various methods. For example, a round window catheter (e.g., U.S. Pat. Nos. 6,440,102 and 6,648,873) can be used.

[0195] Alternatively, formulations can be incorporated into a wick for use between the outer and middle ear (e.g., U.S. Pat. No. 6,120,484) or absorbed to collagen sponge or other solid support (e.g., U.S. Pat. No. 4,164,559).

[0196] If desired, formulations of the disclosure can be incorporated into a gel formulation (e.g., U.S. Pat. Nos. 4,474,752 and 6,911,211).

[0197] In some embodiments, compounds of Formulas (I) and/or (Ia) disclosed herein intended for delivery to the ear can be administered via an implanted pump and delivery system through a needle directly into the middle or inner ear (cochlea) or through a cochlear implant stylet electrode channel or alternative prepared drug delivery channel such as but not limited to a needle through temporal bone into the cochlea.

[0198] Other options include delivery via a pump through a thin film coated onto a multichannel electrode or electrode with a specially imbedded drug delivery channel (pathways) carved into the thin film for this purpose. In other embodiments, the acidic or basic solid gacyclidine can be delivered from the reservoir of an external or internal implanted pumping system.

[0199] Formulations of the disclosure also can be administered to the ear by intratympanic injection into the middle ear, inner ear, or cochlea (e.g., U.S. Pat. No. 6,377,849 and Ser. No. 11/337,815).

[0200] Intratympanic injection of therapeutic agents is the technique of injecting a therapeutic agent behind the tympanic membrane into the middle and/or inner ear. In one embodiment, the formulations described herein are administered directly onto the round window membrane via transtympanic injection. In another embodiment, the ion channel modulating agent auris-acceptable formulations described herein are administered onto the round window membrane via a non-transtympanic approach to the inner ear. In additional embodiments, the formulation described herein is administered onto the round window membrane via a surgical approach to the round window membrane comprising modification of the crista fenestrae cochleae.

[0201] In some embodiments, the compounds of Formulas (I) and/or (Ia) are formulated in rectal compositions such as enemas, rectal gels, rectal foams, rectal aerosols, suppositories, jelly suppositories, or retention enemas, containing conventional suppository bases such as cocoa butter or other glycerides, as well as synthetic polymers such as polyvinylpyrrolidone, PEG (like PEG ointments), and the like.

[0202] Suppositories for rectal administration of the drug (either as a solution, colloid, suspension or a complex) can be prepared by mixing a compound provided herein with a suitable non-irritating excipient that is solid at ordinary temperatures but

liquid at the rectal temperature and will therefore melt or erode/dissolve in the rectum and release the compound. Such materials include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, poloxamers, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol. In suppository forms of the compositions, a low-melting wax such as, but not limited to, a mixture of fatty acid glycerides, optionally in combination with cocoa butter, is first melted.

[0203] Solid compositions can be provided in various different types of dosage forms, depending on the physicochemical properties of the compound provided herein, the desired dissolution rate, cost considerations, and other criteria. In one of the embodiments, the solid composition is a single unit. This implies that one unit dose of the compound is comprised in a single, physically shaped solid form or article. In other words, the solid composition is coherent, which is in contrast to a multiple unit dosage form, in which the units are incoherent.

[0204] Examples of single units which may be used as dosage forms for the solid composition include tablets, such as compressed tablets, film-like units, foil-like units, wafers, lyophilized matrix units, and the like. In one embodiment, the solid composition is a highly porous lyophilized form. Such lyophilizates, sometimes also called wafers or lyophilized tablets, are particularly useful for their rapid disintegration, which also enables the rapid dissolution of the compound.

[0205] On the other hand, for some applications the solid composition may also be formed as a multiple unit dosage form as defined above. Examples of multiple units are powders, granules, microparticles, pellets, mini-tablets, beads, lyophilized powders, and the like. In one embodiment, the solid composition is a lyophilized powder. Such a dispersed lyophilized system comprises a multitude of powder particles, and due to the lyophilization process used in the formation of the powder, each particle has an irregular, porous microstructure through which the powder is capable of absorbing water very rapidly, resulting in quick dissolution. Effervescent compositions are also contemplated to aid the quick dispersion and absorption of the compound.

[0206] Another type of multiparticulate system which is also capable of achieving rapid drug dissolution is that of powders, granules, or pellets from water-soluble excipients which are coated with a compound provided herein so that the

compound is located at the outer surface of the individual particles. In this type of system, the water-soluble low molecular weight excipient may be useful for preparing the cores of such coated particles, which can be subsequently coated with a coating composition comprising the compound and, for example, one or more additional excipients, such as a binder, a pore former, a saccharide, a sugar alcohol, a film-forming polymer, a plasticizer, or other excipients used in pharmaceutical coating compositions.

[0207] Also provided herein are kits. Typically, a kit includes one or more compounds or compositions as described herein. In certain embodiments, a kit can include one or more delivery systems, e.g., for delivering or administering a compound as provided herein, and directions for use of the kit (e.g., instructions for treating a patient). In another embodiment, the kit can include a compound or composition as described herein and a label that indicates that the contents are to be administered to a patient with cancer. In another embodiment, the kit can include a compound or composition as described herein and a label that indicates that the contents are to be administered to a patient with one or more of hepatocellular carcinoma, colon cancer, leukemia, lymphoma, sarcoma, ovarian cancer, diabetic retinopathy, pulmonary fibrosis, rheumatoid arthritis, sepsis, ankylosing spondylitis, psoriasis, scleroderma, mycotic and viral infections, bone and cartilage diseases, Alzheimer's disease, lung disease, osteoarthritis, articular cartilage (chondral) defects, degenerative disc disease (or intervertebral disc degeneration), polyposis coli, bone density and vascular defects in the eye (Osteoporosis-pseudoglioma Syndrome, OPPG), familial exudative vitreoretinopathy, retinal angiogenesis, early coronary disease, tetra-amelia, Müllerian-duct regression and virilization, SERKAL syndrome, type II diabetes, Fuhrmann syndrome, Al-Awadi/Raas-Rothschild/Schinzle phocomelia syndrome, odonto-onychodermal dysplasia, obesity, split-hand/foot malformation, caudal duplication, tooth agenesis, Wilms tumor, skeletal dysplasia, focal dermal hypoplasia, autosomal recessive anonychia, neural tube defects, alpha-thalassemia (ATRX) syndrome, fragile X syndrome, ICF syndrome, Angelman's syndrome, Prader-Willi syndrome, Beckwith-Wiedemann Syndrome, Norrie disease, and Rett syndrome.

Methods of Treatment

[0208] The compounds and compositions provided herein can be used as inhibitors and/or modulators of one or more components of the Wnt pathway, which may include one or more Wnt proteins, and thus can be used to treat a variety of disorders and diseases in which aberrant Wnt signaling is implicated, such as cancer and other diseases associated with abnormal angiogenesis, cellular proliferation, and cell cycling. Accordingly, the compounds and compositions provided herein can be used to treat cancer, to reduce or inhibit angiogenesis, to reduce or inhibit cellular proliferation, to correct a genetic disorder, and/or to treat a neurological condition/disorder/disease due to mutations or dysregulation of the Wnt pathway and/or of one or more of Wnt signaling components. Non-limiting examples of diseases which can be treated with the compounds and compositions provided herein include a variety of cancers, diabetic retinopathy, pulmonary fibrosis, rheumatoid arthritis, sepsis, anklyosing spondylitis, psoriasis, scleroderma, mycotic and viral infections, bone and cartilage diseases, neurological conditions/diseases such as Alzheimer's disease, amyotrophic lateral sclerosis (ALS), motor neurone disease, multiple sclerosis or autism, lung disease, osteoarthritis, articular cartilage (chondral) defects, degenerative disc disease (or intervertebral disc degeneration), polyposis coli, bone density and vascular defects in the eye (Osteoporosis-pseudoglioma Syndrome, OPPG), familial exudative vitreoretinopathy, retinal angiogenesis, early coronary disease, tetra-amelia, Müllerian-duct regression and virilization, SERKAL syndrome, type II diabetes, Fuhrmann syndrome, Al-Awadi/Raas-Rothschild/Schinzle phocomelia syndrome, odonto-onycho-dermal dysplasia, obesity, split-hand/foot malformation, caudal duplication, tooth agenesis, Wilms tumor, skeletal dysplasia, focal dermal hypoplasia, autosomal recessive anonychia, neural tube defects, alpha-thalassemia (ATR-X) syndrome, fragile X syndrome, ICF syndrome, Angelman's syndrome, Prader-Willi syndrome, Beckwith-Wiedemann Syndrome, Norrie disease, and Rett syndrome.

[0209] With respect to cancer, the Wnt pathway is known to be constitutively activated in a variety of cancers including, for example, colon cancer, hepatocellular carcinoma, lung cancer, ovarian cancer, prostate cancer, pancreatic cancer and leukemias such as CML, CLL and T-ALL. Accordingly, the compounds and compositions

described herein may be used to treat these cancers in which the Wnt pathway is constitutively activated. In certain embodiments, the cancer is chosen from hepatocellular carcinoma, colon cancer, leukemia, lymphoma, sarcoma and ovarian cancer.

[0210] Other cancers can also be treated with the compounds and compositions described herein.

[0211] More particularly, cancers that may be treated by the compound, compositions and methods described herein include, but are not limited to, the following:

[0212] 1) Breast cancers, including, for example ER⁺ breast cancer, ER⁻ breast cancer, her2⁻ breast cancer, her2⁺ breast cancer, stromal tumors such as fibroadenomas, phyllodes tumors, and sarcomas, and epithelial tumors such as large duct papillomas; carcinomas of the breast including *in situ* (noninvasive) carcinoma that includes ductal carcinoma *in situ* (including Paget's disease) and lobular carcinoma *in situ*, and invasive (infiltrating) carcinoma including, but not limited to, invasive ductal carcinoma, invasive lobular carcinoma, medullary carcinoma, colloid (mucinous) carcinoma, tubular carcinoma, and invasive papillary carcinoma; and miscellaneous malignant neoplasms. Further examples of breast cancers can include luminal A, luminal B, basal A, basal B, and triple negative breast cancer, which is estrogen receptor negative (ER⁻), progesterone receptor negative, and her2 negative (her2⁻). In some embodiments, the breast cancer may have a high risk Oncotype score.

[0213] 2) Cardiac cancers, including, for example sarcoma, e.g., angiosarcoma, fibrosarcoma, rhabdomyosarcoma, and liposarcoma; myxoma; rhabdomyoma; fibroma; lipoma and teratoma.

[0214] 3) Lung cancers, including, for example, bronchogenic carcinoma, e.g., squamous cell, undifferentiated small cell, undifferentiated large cell, and adenocarcinoma; alveolar and bronchiolar carcinoma; bronchial adenoma; sarcoma; lymphoma; chondromatous hamartoma; and mesothelioma.

[0215] 4) Gastrointestinal cancer, including, for example, cancers of the esophagus, e.g., squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, and lymphoma; cancers of the stomach, e.g., carcinoma, lymphoma, and leiomyosarcoma; cancers of the pancreas, e.g., ductal adenocarcinoma, insulinoma, glucagonoma,

gastrinoma, carcinoid tumors, and vipoma; cancers of the small bowel, e.g., adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, and fibroma; cancers of the large bowel, e.g., adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, and leiomyoma.

[0216] 5) Genitourinary tract cancers, including, for example, cancers of the kidney, e.g., adenocarcinoma, Wilm's tumor (nephroblastoma), lymphoma, and leukemia; cancers of the bladder and urethra, e.g., squamous cell carcinoma, transitional cell carcinoma, and adenocarcinoma; cancers of the prostate, e.g., adenocarcinoma, and sarcoma; cancer of the testis, e.g., seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, and lipoma.

[0217] 6) Liver cancers, including, for example, hepatoma, e.g., hepatocellular carcinoma; cholangiocarcinoma; hepatoblastoma; angiosarcoma; hepatocellular adenoma; and hemangioma.

[0218] 7) Bone cancers, including, for example, osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors.

[0219] 8) Nervous system cancers, including, for example, cancers of the skull, e.g., osteoma, hemangioma, granuloma, xanthoma, and osteitis deformans; cancers of the meninges, e.g., meningioma, meningiosarcoma, and gliomatosis; cancers of the brain, e.g., astrocytoma, medulloblastoma, glioma, ependymoma, germinoma (pinealoma), glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, and congenital tumors; and cancers of the spinal cord, e.g., neurofibroma, meningioma, glioma, and sarcoma.

[0220] 9) Gynecological cancers, including, for example, cancers of the uterus, e.g., endometrial carcinoma; cancers of the cervix, e.g., cervical carcinoma, and pre tumor cervical dysplasia; cancers of the ovaries, e.g., ovarian carcinoma, including serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma,

granulosa theca cell tumors, Sertoli Leydig cell tumors, dysgerminoma, and malignant teratoma; cancers of the vulva, e.g., squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, and melanoma; cancers of the vagina, e.g., clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma, and embryonal rhabdomyosarcoma; and cancers of the fallopian tubes, e.g., carcinoma.

[0221] 10) Hematologic cancers, including, for example, cancers of the blood, e.g., acute myeloid leukemia, chronic myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, and myelodysplastic syndrome, Hodgkin's lymphoma, non-Hodgkin's lymphoma (malignant lymphoma) and Waldenström's macroglobulinemia.

[0222] 11) Skin cancers and skin disorders, including, for example, malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, and scleroderma.

[0223] 12) Adrenal gland cancers, including, for example, neuroblastoma.

[0224] Cancers may be solid tumors that may or may not be metastatic. Cancers may also occur, as in leukemia, as a diffuse tissue. Thus, the term "tumor cell," as provided herein, includes a cell afflicted by any one of the above identified disorders.

[0225] A method of treating cancer using a compound or composition as described herein may be combined with existing methods of treating cancers, for example by chemotherapy, irradiation, or surgery (e.g., oophorectomy). In some embodiments, a compound or composition can be administered before, during, or after another anticancer agent or treatment.

[0226] The compounds and compositions described herein can be used as anti-angiogenesis agents and as agents for modulating and/or inhibiting the activity of protein kinases, thus providing treatments for cancer and other diseases associated with cellular proliferation mediated by protein kinases. Accordingly, provided herein is a method of treating cancer or preventing or reducing angiogenesis through kinase inhibition.

[0227] In addition, and including treatment of cancer, the compounds and compositions described herein can function as cell-cycle control agents for treating proliferative disorders in a patient. Disorders associated with excessive proliferation

include, for example, cancers, scleroderma, immunological disorders involving undesired proliferation of leukocytes, and restenosis and other smooth muscle disorders. Furthermore, such compounds may be used to prevent de-differentiation of post-mitotic tissue and/or cells.

[0228] Diseases or disorders associated with uncontrolled or abnormal cellular proliferation include, but are not limited to, the following:

- a variety of cancers, including, but not limited to, carcinoma, hematopoietic tumors of lymphoid lineage, hematopoietic tumors of myeloid lineage, tumors of mesenchymal origin, tumors of the central and peripheral nervous system and other tumors including melanoma, seminoma, and Kaposi's sarcoma.
- a disease process which features abnormal cellular proliferation, e.g., benign prostatic hyperplasia, familial adenomatosis polyposis, neurofibromatosis, atherosclerosis, arthritis, glomerulonephritis, restenosis following angioplasty or vascular surgery, inflammatory bowel disease, transplantation rejection, endotoxic shock, and fungal infections. Fibrotic disorders such as skin fibrosis; scleroderma; progressive systemic fibrosis; lung fibrosis; muscle fibrosis; kidney fibrosis; glomerulosclerosis; glomerulonephritis; hypertrophic scar formation; uterine fibrosis; renal fibrosis; cirrhosis of the liver, liver fibrosis; adhesions, such as those occurring in the abdomen, pelvis, spine or tendons; chronic obstructive pulmonary disease; fibrosis following myocardial infarction; pulmonary fibrosis; fibrosis and scarring associated with diffuse/interstitial lung disease; central nervous system fibrosis, such as fibrosis following stroke; fibrosis associated with neuro-degenerative disorders such as Alzheimer's Disease or multiple sclerosis; fibrosis associated with proliferative vitreoretinopathy (PVR); restenosis; endometriosis; ischemic disease and radiation fibrosis.
- defective apoptosis-associated conditions, such as cancers (including but not limited to those types mentioned herein), viral infections (including but not limited to herpesvirus, poxvirus, Epstein-Barr virus, Sindbis virus and

adenovirus), prevention of AIDS development in HIV-infected individuals, autoimmune diseases (including but not limited to systemic lupus erythematosus, rheumatoid arthritis, sepsis, ankylosing spondylitis, psoriasis, scleroderma, autoimmune mediated glomerulonephritis, inflammatory bowel disease and autoimmune diabetes mellitus), neuro-degenerative disorders (including but not limited to Alzheimer's disease, lung disease, amyotrophic lateral sclerosis, retinitis pigmentosa, Parkinson's disease, AIDS-related dementia, spinal muscular atrophy and cerebellar degeneration), myelodysplastic syndromes, aplastic anemia, ischemic injury associated with myocardial infarctions, stroke and reperfusion injury, arrhythmia, atherosclerosis, toxin-induced or alcohol related liver diseases, hematological diseases (including but not limited to chronic anemia and aplastic anemia), degenerative diseases of the musculoskeletal system (including but not limited to osteoporosis and arthritis), aspirin-sensitive rhinosinusitis, cystic fibrosis, multiple sclerosis, kidney diseases and cancer pain.

- genetic diseases due to mutations in Wnt signaling components, such as polyposis coli, bone density and vascular defects in the eye (Osteoporosis-pseudoglioma Syndrome, OPPG), familial exudative vitreoretinopathy, retinal angiogenesis, early coronary disease, tetra-amelia, Müllerian-duct regression and virilization, SERKAL syndrome, type II diabetes, Fuhrmann syndrome, Al-Awadi/Raas-Rothschild/Schinzel phocomelia syndrome, odonto-onycho-dermal dysplasia, obesity, split-hand/foot malformation, caudal duplication, tooth agenesis, Wilms tumor, skeletal dysplasia, focal dermal hypoplasia, autosomal recessive anonychia, neural tube defects, alpha-thalassemia (ATRX) syndrome, fragile X syndrome, ICF syndrome, Angelman's syndrome, Prader-Willi syndrome, Beckwith-Wiedemann Syndrome, Norrie disease and Rett syndrome.

[0229] Furthermore, the compounds and compositions described herein can be used to treat neurological conditions, disorders and/or diseases caused by dysfunction in the Wnt signaling pathway. Non-limiting examples of neurological

conditions/disorders/diseases which can be treated with the compounds and compositions provided herein include Alzheimer's disease, aphasia, apraxia, arachnoiditis, ataxia telangiectasia, attention deficit hyperactivity disorder, auditory processing disorder, autism, alcoholism, Bell's palsy, bipolar disorder, brachial plexus injury, Canavan disease, carpal tunnel syndrome, causalgia, central pain syndrome, central pontine myelinolysis, centronuclear myopathy, cephalic disorder, cerebral aneurysm, cerebral arteriosclerosis, cerebral atrophy, cerebral gigantism, cerebral palsy, cerebral vasculitis, cervical spinal stenosis, Charcot-Marie-Tooth disease, Chiari malformation, chronic fatigue syndrome, chronic inflammatory demyelinating polyneuropathy (CIDP), chronic pain, Coffin-Lowry syndrome, complex regional pain syndrome, compression neuropathy, congenital facial diplegia, corticobasal degeneration, cranial arteritis, craniosynostosis, Creutzfeldt-Jakob disease, cumulative trauma disorder, Cushing's syndrome, cytomegalic inclusion body disease (CIBD), Dandy-Walker syndrome, Dawson disease, De Morsier's syndrome, Dejerine-Klumpke palsy, Dejerine-Sottas disease, delayed sleep phase syndrome, dementia, dermatomyositis, developmental dyspraxia, diabetic neuropathy, diffuse sclerosis, Dravet syndrome, dysautonomia, dyscalculia, dysgraphia, dyslexia, dystonia, empty sella syndrome, encephalitis, encephalocele, encephalotrigeminal angiomas, encopresis, epilepsy, Erb's palsy, erythromelalgia, essential tremor, Fabry's disease, Fahr's syndrome, familial spastic paralysis, febrile seizure, Fisher syndrome, Friedreich's ataxia, fibromyalgia, Foville's syndrome, Gaucher's disease, Gerstmann's syndrome, giant cell arteritis, giant cell inclusion disease, globoid cell leukodystrophy, gray matter heterotopia, Guillain-Barré syndrome, HTLV-1 associated myelopathy, Hallervorden-Spatz disease, hemifacial spasm, hereditary spastic paraplegia, heredopathia atactica polyneuritiformis, herpes zoster oticus, herpes zoster, Hirayama syndrome, holoprosencephaly, Huntington's disease, hydranencephaly, hydrocephalus, hypercortisolism, hypoxia, immune-mediated encephalomyelitis, inclusion body myositis, incontinentia pigmenti, infantile phytanic acid storage disease, infantile Refsum disease, infantile spasms, inflammatory myopathy, intracranial cyst, intracranial hypertension, Joubert syndrome, Karak syndrome, Kearns-Sayre syndrome, Kennedy disease, Kinsbourne syndrome, Klippel Feil syndrome, Krabbe disease, Kugelberg-Welander disease, kuru, Lafora disease, Lambert-Eaton

myasthenic syndrome, Landau-Kleffner syndrome, lateral medullary (Wallenberg) syndrome, Leigh's disease, Lennox-Gastaut syndrome, Lesch-Nyhan syndrome, leukodystrophy, Lewy body dementia, lissencephaly, locked-in syndrome, Lou Gehrig's disease, lumbar disc disease, lumbar spinal stenosis, Lyme disease, Machado-Joseph disease (Spinocerebellar ataxia type 3), macrencephaly, macropsia, megalencephaly, Melkersson-Rosenthal syndrome, Menieres disease, meningitis, Menkes disease, etachromatic leukodystrophy, microcephaly, micropsia, Miller Fisher syndrome, misophonia, mitochondrial myopathy, Mobius syndrome, monomelic amyotrophy, motor neurone disease, motor skills disorder, Moyamoya disease, mucopolysaccharidoses, multi-infarct dementia, multifocal motor neuropathy, multiple sclerosis, multiple system atrophy, muscular dystrophy, myalgic encephalomyelitis, myasthenia gravis, myelinoclastic diffuse sclerosis, myoclonic Encephalopathy of infants, myoclonus, myopathy, myotubular myopathy, myotonia congenital, narcolepsy, neurofibromatosis, neuroleptic malignant syndrome, lupus erythematosus, neuromyotonia, neuronal ceroid lipofuscinosis, Niemann-Pick disease, O'Sullivan-McLeod syndrome, occipital Neuralgia, occult Spinal Dysraphism Sequence, Ohtahara syndrome, olivopontocerebellar atrophy, opsoclonus myoclonus syndrome, optic neuritis, orthostatic hypotension, palinopsia, paresthesia, Parkinson's disease, paramyotonia Congenita, paraneoplastic diseases, paroxysmal attacks, Parry-Romberg syndrome, Pelizaeus-Merzbacher disease, periodic paralyses, peripheral neuropathy, photic sneeze reflex, phytanic acid storage disease, Pick's disease, polymicrogyria (PMG), polymyositis, porencephaly, post-polio syndrome, postherpetic neuralgia (PHN), postural hypotension, Prader-Willi syndrome, primary lateral sclerosis, prion diseases, progressive hemifacial atrophy, progressive multifocal leukoencephalopathy, progressive supranuclear palsy, pseudotumor cerebri, Ramsay Hunt syndrome type I, Ramsay Hunt syndrome type II, Ramsay Hunt syndrome type III, Rasmussen's encephalitis, reflex neurovascular dystrophy, Refsum disease, restless legs syndrome, retrovirus-associated myelopathy, Rett syndrome, Reye's syndrome, rhythmic movement disorder, Romberg syndrome, Saint Vitus dance, Sandhoff disease, schizophrenia, Schilder's disease, schizencephaly, sensory integration dysfunction, septo-optic dysplasia, Shy-Drager syndrome, Sjögren's syndrome, snatiation, Sotos syndrome, spasticity, spina bifida, spinal cord tumors, spinal muscular

atrophy, spinocerebellar ataxia, Steele-Richardson-Olszewski syndrome, Stiff-person syndrome, stroke, Sturge-Weber syndrome, subacute sclerosing panencephalitis, subcortical arteriosclerotic encephalopathy, superficial siderosis, Sydenham's chorea, syncope, synesthesia, syringomyelia, tarsal tunnel syndrome, tardive dyskinesia, tardive dysphrenia, Tarlov cyst, Tay-Sachs disease, temporal arteritis, tetanus, tethered spinal cord syndrome, Thomsen disease, thoracic outlet syndrome, tic douloureux, Todd's paralysis, Tourette syndrome, toxic encephalopathy, transient ischemic attack, transmissible spongiform encephalopathies, transverse myelitis, tremor, trigeminal neuralgia, tropical spastic paraparesis, trypanosomiasis, tuberous sclerosis, ubisiosis, Von Hippel-Lindau disease (VHL), Viliuisk Encephalomyelitis (VE), Wallenberg's syndrome, Werdnig, Hoffman disease, west syndrome, Williams syndrome, Wilson's disease, and Zellweger syndrome.

[0230] The compounds and compositions may also be useful in the inhibition of the development of invasive cancer, tumor angiogenesis and metastasis.

[0231] In some embodiments, the disclosure provides a method for treating a disease or disorder associated with aberrant cellular proliferation by administering to a patient in need of such treatment an effective amount of one or more of the compounds of Formulas (I) and/or (Ia), in combination (simultaneously or sequentially) with at least one other agent.

[0232] In some embodiments, the pharmaceutical composition comprises a therapeutically effective amount of a compound of Formulas (I) and/or (Ia), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

[0233] In some embodiments, the method of treating a disorder or disease in which aberrant Wnt signaling is implicated in a patient is provided herein, the method comprises administering to the patient a therapeutically effective amount of a compound of Formulas (I) and/or (Ia), or a pharmaceutically acceptable salt thereof.

[0234] In some embodiments, the disorder or disease is cancer.

[0235] In some embodiments, the disorder or disease is diabetic retinopathy.

[0236] In some embodiments, the disorder or disease is pulmonary fibrosis.

[0237] In some embodiments, the disorder or disease is rheumatoid arthritis.

[0238] In some embodiments, the disorder or disease is scleroderma.

[0239] In some embodiments, the disorder or disease is a mycotic or viral infection.

[0240] In some embodiments, the disorder or disease is a bone or cartilage disease.

[0241] In some embodiments, the disorder or disease is Alzheimer's disease.

[0242] In some embodiments, the disorder or disease is osteoarthritis.

[0243] In some embodiments, the disorder or disease is lung disease

[0244] In some embodiments, the disorder or disease is a genetic disease caused by mutations in Wnt signaling components, wherein the genetic disease is selected from: polyposis coli, osteoporosis-pseudoglioma syndrome, familial exudative vitreoretinopathy, retinal angiogenesis, early coronary disease, tetra-amelia syndrome, Müllerian-duct regression and virilization, SERKAL syndrome, diabetes mellitus type 2, Fuhrmann syndrome, Al-Awadi/Raas-Rothschild/Schinzal phocomelia syndrome, odonto-onycho-dermal dysplasia, obesity, split-hand/foot malformation, caudal duplication syndrome, tooth agenesis, Wilms tumor, skeletal dysplasia, focal dermal hypoplasia, autosomal recessive anonychia, neural tube defects, alpha-thalassemia (ATRX) syndrome, fragile X syndrome, ICF syndrome, Angelman syndrome, Prader-Willi syndrome, Beckwith-Wiedemann Syndrome, Norrie disease, and Rett syndrome.

[0245] In some embodiments, the patient is a human.

[0246] In some embodiments, the cancer is chosen from: hepatocellular carcinoma, colon cancer, breast cancer, pancreatic cancer, chronic myeloid leukemia (CML), chronic myelomonocytic leukemia, chronic lymphocytic leukemia (CLL), acute myeloid leukemia, acute lymphocytic leukemia, Hodgkin lymphoma, lymphoma, sarcoma, and ovarian cancer.

[0247] In some embodiments, the cancer is chosen from: lung cancer - non-small cell, lung cancer - small cell, multiple myeloma, nasopharyngeal cancer, neuroblastoma, osteosarcoma, penile cancer, pituitary tumors, prostate cancer, retinoblastoma, rhabdomyosarcoma, salivary gland cancer, skin cancer - basal and squamous cell, skin cancer - melanoma, small intestine cancer, stomach cancers, testicular cancer, thymus cancer, thyroid cancer, uterine sarcoma, vaginal cancer, vulvar cancer, laryngeal or hypopharyngeal cancer, kidney cancer, Kaposi sarcoma, gestational

trophoblastic disease, gastrointestinal stromal tumor, gastrointestinal carcinoid tumor, gallbladder cancer, eye cancer (melanoma and lymphoma), Ewing tumor, esophagus cancer, endometrial cancer, colorectal cancer, cervical cancer, brain or spinal cord tumor, bone metastasis, bone cancer, bladder cancer, bile duct cancer, anal cancer, and adrenal cortical cancer.

- [0248] In some embodiments, the cancer is hepatocellular carcinoma.
- [0249] In some embodiments, the cancer is colon cancer.
- [0250] In some embodiments, the cancer is breast cancer.
- [0251] In some embodiments, the cancer is pancreatic cancer.
- [0252] In some embodiments, the cancer is chronic myeloid leukemia (CML).
- [0253] In some embodiments, the cancer is chronic myelomonocytic leukemia.
- [0254] In some embodiments, the cancer is chronic lymphocytic leukemia (CLL).
- [0255] In some embodiments, the cancer is acute myeloid leukemia.
- [0256] In some embodiments, the cancer is acute lymphocytic leukemia.
- [0257] In some embodiments, the cancer is Hodgkin lymphoma.
- [0258] In some embodiments, the cancer is lymphoma.
- [0259] In some embodiments, the cancer is sarcoma.
- [0260] In some embodiments, the cancer is ovarian cancer.
- [0261] In some embodiments, the cancer is lung cancer - non-small cell.
- [0262] In some embodiments, the cancer is lung cancer - small cell.
- [0263] In some embodiments, the cancer is multiple myeloma.
- [0264] In some embodiments, the cancer is nasopharyngeal cancer.
- [0265] In some embodiments, the cancer is neuroblastoma.
- [0266] In some embodiments, the cancer is osteosarcoma.
- [0267] In some embodiments, the cancer is penile cancer.
- [0268] In some embodiments, the cancer is pituitary tumors.
- [0269] In some embodiments, the cancer is prostate cancer.
- [0270] In some embodiments, the cancer is retinoblastoma.
- [0271] In some embodiments, the cancer is rhabdomyosarcoma.

- [0272] In some embodiments, the cancer is salivary gland cancer.
- [0273] In some embodiments, the cancer is skin cancer - basal and squamous cell.
- [0274] In some embodiments, the cancer is skin cancer – melanoma.
- [0275] In some embodiments, the cancer is small intestine cancer.
- [0276] In some embodiments, the cancer is stomach cancers.
- [0277] In some embodiments, the cancer is testicular cancer.
- [0278] In some embodiments, the cancer is thymus cancer.
- [0279] In some embodiments, the cancer is thyroid cancer.
- [0280] In some embodiments, the cancer is uterine sarcoma.
- [0281] In some embodiments, the cancer is vaginal cancer.
- [0282] In some embodiments, the cancer is vulvar cancer.
- [0283] In some embodiments, the cancer is Wilms tumor.
- [0284] In some embodiments, the cancer is laryngeal or hypopharyngeal cancer.
- [0285] In some embodiments, the cancer is kidney cancer.
- [0286] In some embodiments, the cancer is Kaposi sarcoma.
- [0287] In some embodiments, the cancer is gestational trophoblastic disease.
- [0288] In some embodiments, the cancer is gastrointestinal stromal tumor.
- [0289] In some embodiments, the cancer is gastrointestinal carcinoid tumor.
- [0290] In some embodiments, the cancer is gallbladder cancer.
- [0291] In some embodiments, the cancer is eye cancer (melanoma and lymphoma).
- [0292] In some embodiments, the cancer is Ewing tumor.
- [0293] In some embodiments, the cancer is esophagus cancer.
- [0294] In some embodiments, the cancer is endometrial cancer.
- [0295] In some embodiments, the cancer is colorectal cancer.
- [0296] In some embodiments, the cancer is cervical cancer.
- [0297] In some embodiments, the cancer is brain or spinal cord tumor.
- [0298] In some embodiments, the cancer is bone metastasis.
- [0299] In some embodiments, the cancer is bone cancer.

[0300] In some embodiments, the cancer is bladder cancer.

[0301] In some embodiments, the cancer is bile duct cancer.

[0302] In some embodiments, the cancer is anal cancer.

[0303] In some embodiments, the cancer is adrenal cortical cancer.

[0304] In some embodiments, the disorder or disease is a neurological condition, disorder or disease, wherein the neurological condition/disorder/disease is selected from: Alzheimer's disease, frontotemporal dementias, dementia with lewy bodies, prion diseases, Parkinson's disease, Huntington's disease, progressive supranuclear palsy, corticobasal degeneration, multiple system atrophy, amyotrophic lateral sclerosis (ALS), inclusion body myositis, autism, degenerative myopathies, diabetic neuropathy, other metabolic neuropathies, endocrine neuropathies, orthostatic hypotension, multiple sclerosis, and Charcot-Marie-Tooth disease.

[0305] In some embodiments, the compound of Formulas (I) and/or (Ia) inhibits one or more proteins in the Wnt pathway.

[0306] In some embodiments, the compound of Formulas (I) and/or (Ia) inhibits signaling induced by one or more Wnt proteins.

[0307] In some embodiments, the Wnt proteins are chosen from: WNT1, WNT2, WNT2B, WNT3, WNT3A, WNT4, WNT5A, WNT5B, WNT6, WNT7A, WNT7B, WNT8A, WNT8B, WNT9A, WNT9B, WNT10A, WNT10B, WNT11, and WNT16.

[0308] In some embodiments, the compounds of Formulas (I) and/or (Ia) inhibit a kinase activity.

[0309] In some embodiments, a method for treating a disease or disorder mediated by the Wnt pathway in a patient is provided, the method comprises administering to the patient a therapeutically effective amount of a compound (or compounds) of Formulas (I) and/or (Ia), or a pharmaceutically acceptable salt thereof.

[0310] In some embodiments, the compounds of Formulas (I) and/or (Ia) inhibit one or more Wnt proteins.

[0311] In some embodiments, a method for treating a disease or disorder mediated by kinase activity in a patient is provided, the method comprises administering

to the patient a therapeutically effective amount of a compound (or compounds) of Formulas (I) and/or (Ia), or a pharmaceutically acceptable salt thereof.

[0312] In some embodiments, the disease or disorder comprises tumor growth, cell proliferation, or angiogenesis.

[0313] In some embodiments, the method inhibits the activity of a protein kinase receptor, the method comprises contacting the receptor with an effective amount of a compound (or compounds) of Formulas (I) and/or (Ia), or a pharmaceutically acceptable salt thereof.

[0314] In some embodiments, a method for treating a disease or disorder associated with aberrant cellular proliferation in a patient is provided, the method comprises administering to the patient a therapeutically effective amount of a compound (or compounds) of Formulas (I) and/or (Ia), or a pharmaceutically acceptable salt thereof.

[0315] In some embodiments, the method prevents or reduces angiogenesis in a patient; the method comprises administering to the patient a therapeutically effective amount of a compound (or compounds) of Formulas (I) and/or (Ia), or a pharmaceutically acceptable salt thereof.

[0316] In some embodiments, the method prevents or reduces abnormal cellular proliferation in a patient; the method comprises administering to the patient a therapeutically effective amount of a compound (or compounds) of Formulas (I) and/or (Ia), or a pharmaceutically acceptable salt thereof.

[0317] In some embodiments, a method for treating a disease or disorder associated with aberrant cellular proliferation in a patient is provided, the method comprises administering to the patient a pharmaceutical composition comprising one or more of the compounds of Formulas (I) and/or (Ia) in combination with a pharmaceutically acceptable carrier and one or more other agents.

[0318] Moreover, the compounds and compositions, for example, as inhibitors of the cyclin-dependent kinases (CDKs), can modulate the level of cellular RNA and DNA synthesis and therefore are expected to be useful in the treatment of viral infections such as HIV, human papilloma virus, herpes virus, Epstein-Barr virus, adenovirus, Sindbis virus, pox virus, and the like.

[0319] Compounds and compositions described herein can inhibit the kinase activity of, for example, CDK/cyclin complexes, such as those active in the G₀ or G₁ stage of the cell cycle, e.g., CDK2, CDK4, and/or CDK6 complexes.

Evaluation of Biological Activity

[0320] The biological activity of the compounds described herein can be tested using any suitable assay known to those of skill in the art, see e.g., WO 2001/053268 or WO 2005/009997. For example, the activity of a compound may be tested using one or more of the test methods outlined below.

[0321] In one example, tumor cells may be screened for Wnt independent growth. In such a method, tumor cells of interest are contacted with a compound (i.e. inhibitor) of interest, and the proliferation of the cells, e.g. by uptake of tritiated thymidine, is monitored. In some embodiments, tumor cells may be isolated from a candidate patient who has been screened for the presence of a cancer that is associated with a mutation in the Wnt signaling pathway. Candidate cancers include, without limitation, those listed above.

[0322] In another example, *in vitro* assays for Wnt biological activity may be used, e.g. stabilization of β -catenin and promoting growth of stem cells. Assays for biological activity of Wnt include stabilization of β -catenin, which can be measured, for example, by serial dilutions of a candidate inhibitor composition. An exemplary assay for Wnt biological activity contacts a Wnt composition in the presence of a candidate inhibitor with cells, e.g. mouse L cells. The cells are cultured for a period of time sufficient to stabilize β -catenin, usually at least about 1 hour, and lysed. The cell lysate is resolved by SDS PAGE, then transferred to nitrocellulose and probed with antibodies specific for β -catenin.

[0323] In a further example, the activity of a candidate compound can be measured in a *Xenopus* secondary axis bioassay [Leyns, L. *et al. Cell* (1997), 88(6), 747-756].

[0324] To further illustrate this disclosure, the following examples are included. The examples should not, of course, be construed as specifically limiting the disclosure. Variations of these examples within the scope of the claims are within the

purview of one skilled in the art and are considered to fall within the scope of the invention as described, and claimed herein. The reader will recognize that the skilled artisan, armed with the present disclosure, and skill in the art is able to prepare and use the invention without exhaustive examples.

EXAMPLES

Compound preparation

[0325] The starting materials used in preparing the compounds of the disclosure are known, made by known methods, or are commercially available. It will be apparent to the skilled artisan that methods for preparing precursors and functionality related to the compounds claimed herein are generally described in the literature. The skilled artisan given the literature and this disclosure is well equipped to prepare any of the compounds.

[0326] It is recognized that the skilled artisan in the art of organic chemistry can readily carry out manipulations without further direction, that is, it is well within the scope and practice of the skilled artisan to carry out these manipulations. These include reduction of carbonyl compounds to their corresponding alcohols, oxidations, acylations, aromatic substitutions, both electrophilic and nucleophilic, etherifications, esterification and saponification and the like. These manipulations are discussed in standard texts such as *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure* 7th Ed., John Wiley & Sons (2013), Carey and Sundberg, *Advanced Organic Chemistry* 5th Ed., Springer (2007), *Comprehensive Organic Transformations: A Guide to Functional Group Transformations*, 2nd Ed., John Wiley & Sons (1999) and the like.

[0327] The skilled artisan will readily appreciate that certain reactions are best carried out when other functionality is masked or protected in the molecule, thus avoiding any undesirable side reactions and/or increasing the yield of the reaction. Often the skilled artisan utilizes protecting groups to accomplish such increased yields or to avoid the undesired reactions. These reactions are found in the literature and are also well within the scope of the skilled artisan. Examples of many of these manipulations can be found for example in T. Greene and P. Wuts *Protective Groups in Organic Synthesis*, 4th Ed., John Wiley & Sons (2007).

[0328] Trademarks used herein are examples only and reflect illustrative materials used at the time of filing the present disclosure. The skilled artisan will recognize that variations in lot, manufacturing processes, and the like, are expected. Hence the examples, and the trademarks used in them are non-limiting, and they are not intended to be limiting, but are merely an illustration of how a skilled artisan may choose to perform one or more of the embodiments of the disclosure.

[0329] (^1H) nuclear magnetic resonance spectra (NMR) were measured in the indicated solvents on a Bruker NMR spectrometer (Avance TM DRX300, 300 MHz for ^1H or Avance TM DRX500, 500 MHz for ^1H) or Varian NMR spectrometer (Mercury 400BB, 400 MHz for ^1H). Peak positions are expressed in parts per million (ppm) downfield from tetramethylsilane. The peak multiplicities are denoted as follows, s, singlet; d, doublet; t, triplet; q, quartet; ABq, AB quartet; quin, quintet; sex, sextet; sep, septet; non, nonet; dd, doublet of doublets; d/ABq, doublet of AB quartet; dt, doublet of triplets; td, triplet of doublets; dq, doublet of quartets; m, multiplet.

[0330] The following abbreviations have the indicated meanings:

brine = saturated aqueous sodium chloride

CDCl_3 = deuterated chloroform

CsCO_3 = cesium carbonate

DCE = dichloroethane

DCM = dichloromethane

DHP = 3,4-dihydro-2*H*-pyran

DMF = N,N-dimethylformamide

DMSO-d_6 = deuterated dimethylsulfoxide

ESIMS = electron spray mass spectrometry

EtOAc = ethyl acetate

Et_3SiH = triethylsilane

HCl = hydrochloric acid

HOAc = acetic acid

KOAc = potassium acetate

KOH = potassium hydroxide

K_3PO_4 = potassium phosphate

LAH = lithium aluminum hydride

MeOH = methanol

MgSO₄ = magnesium sulfate

NaBH(OAc)₃ = sodium triacetoxo borohydride

Na₂CO₃ = sodium carbonate

NaHCO₃ = sodium bicarbonate

NaHSO₃ = sodium bisulfite

NaOAc = sodium acetate

NMR = nuclear magnetic resonance

Pd/C = palladium on carbon

Pd₂(dba)₃ = tris(dibenzylideneacetone)dipalladium(0)

PdCl₂(dppf)₂ = 1,1'-bis(diphenylphosphino)ferrocene-palladium(II)dichloride

Pd(PPh₃)₂Cl₂ = dichloro-bis(triphenylphosphine)palladium (II)

Pd(PPh₃)₄ = tetrakis(triphenylphosphine)palladium(0)

PE = petroleum ether

PPTS = pyridinium p-toluenesulfonate

rt = room temperature

SEM = 2-(trimethylsilyl)ethoxymethyl

TFA = trifluoroacetic acid

THF = tetrahydrofuran


TLC = thin layer chromatography

[0331] The following example schemes are provided for the guidance of the reader, and collectively represent an example method for making the compounds provided herein. Furthermore, other methods for preparing compounds of the disclosure will be readily apparent to the person of ordinary skill in the art in light of the following reaction schemes and examples. The skilled artisan is thoroughly equipped to prepare these compounds by those methods given the literature and this disclosure. The compound numberings used in the synthetic schemes depicted below are meant for those specific schemes only, and should not be construed as or confused with same numberings

in other sections of the application. Unless otherwise indicated, all variables are as defined above.

General procedures

[0332] Compounds of Formulas (I) and/or (Ia) of the present disclosure can be prepared as depicted in Scheme 1.




Scheme 1

[0333] Scheme 1 describes a method for preparation of indazole derivatives (IX) by first formylating 5-bromo-1H-indole (I) to produce 5-bromo-1H-indazole-3-carbaldehyde (II) followed by protection with SEM-Cl to give 5-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazole-3-carbaldehyde (III). Bromide (III) is then reacted with bis(pinacolato)diboron to form the borate ester (IV). Suzuki coupling with

various bromides (V) yields indazole derivatives (VI). Aldehyde (VI) is reacted with various 1,2-diamines (VII) to produce (VIII). Final deprotection of the pyrazole nitrogen yields the desired indazole derivatives (IX).

Illustrative Compound Examples

[0334] Preparation of intermediate (IV) is depicted below in Scheme 2.



Scheme 2

Step 1

[0335] A solution of NaNO_2 (110.4 g, 1.6 mol, 8 eq) in water (200 mL) was added dropwise to a solution of 5-bromoindole (I) (39.2 g, 0.2 mol, 1 eq) in acetone (1000 mL) stirred at $-10 \rightarrow 0^\circ\text{C}$, while adding NaNO_2 the solution temperature was maintained below 20°C . An aqueous 2N HCl solution (480 mL) was added slowly to the solution with vigorously stirring while keeping the internal temperature between 0 and 20°C . The solution was further stirred at 20°C for 3 h after the addition. The solution was concentrated under reduced pressure to remove acetone while keeping the temperature

below 35°C. The solid was collected by filtration and transferred to a flask. Cold (-10°C) DCM (200 mL) was added and stirred for 30 min at -5°C, the solids were filtered and dried under vacuum at 40°C to get 5-bromo-1H-indazole-3-carbaldehyde (**II**) (34.0 g, 151 mmol, 76% yield) as a brown solid. ESIMS found for C₈H₅BrN₂O *m/z* 225 (M+H).


Step 2

[0336] To a suspension of NaH (6.6 g, 166 mmol, 1.10 eq) in DMF (500 mL) was added a solution of 5-bromo-1H-indazole-3-carbaldehyde (**II**) (34.0 g, 151 mmol, 1.00 eq) in DMF (50 mL) dropwise at 0°C over a period of 30 min. The mixture was stirred at room temperature for 2 h, then SEM-Cl (26.4 g, 159 mmol, 1.08 eq) was added dropwise and the mixture was stirred at room temperature for another 3 h. Then the mixture was poured into an ice-water mixture (1000 mL) and extracted with EtOAc (300 mL x 3), the organic phases were combined, dried over Na₂SO₄, filtered and concentrated in vacuo, the resultant residue was purified by flash chromatography on silica gel (PE:EtOAc = 20:1→10:1) to afford 5-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazole-3-carbaldehyde (**III**) as a mixture of regioisomers (53.0 g, 151 mmol, 100% yield) as a yellow oil. ESIMS found for C₁₄H₁₉BrN₂O₂Si *m/z* 355 (M+H).

Step 3

[0337] To a solution of the mixed 5-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazole-3-carbaldehyde (**III**) (53.0 g, 151 mmol, 1.0 eq), bis(pinacolato)diboron (38.0 g, 150 mmol, 1.0 eq) and KOAc (44.0 g, 450 mmol, 3.00 eq) in DMF (1000 mL) was added Pd(dppf)Cl₂ (7.7 g, 10.5 mmol, 0.07 eq). The mixture was stirred at 90°C under nitrogen for 10 h. The mixture was filtered; the filtrate was poured onto water (1000 mL) and extracted with EtOAc (500 mL x 3). The combined organic phases were dried, filtered and concentrated in vacuo. The resultant residue was purified by flash chromatography on silica gel (PE:EtOAc = 10:1→1:1) to give the 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazole-3-carbaldehyde (**IV**) as a mixture of regioisomers (42.9 g, 106 mmol, 71% yield) as a yellow oil. ESIMS found for C₂₀H₃₁BN₂O₄Si *m/z* 403 (M+H).

[0338] Preparation of intermediate N-(5-bromopyridin-3-yl)isobutyramide (VII) is depicted below in Scheme 3.




Scheme 3


Step 1

[0339] 3-Amino-5-bromo pyridine (V) (1 eq) was dissolved in DCM and cooled to 0°C before adding pyridine (2.2 eq) and isobutyryl chloride (VI) (1.1 eq). The reaction mixture was stirred at r.t. for 15 h until TLC showed the reaction was complete. The reaction mixture was diluted with DCM and washed with water. The organic extract was dried, concentrated and purified by column chromatography using silica gel (100-200 mesh) to afford N-(5-bromopyridin-3-yl)isobutyramide (VII) as an off-white solid, (71% yield). ¹H NMR (CDCl₃) δ ppm 8.55-8.35 (m, 3H), 7.32 (s, 1H), 2.59-2.48 (m, 1H), 1.28-1.27 (d, 6H); ESIMS found C₉H₁₁BrN₂O *m/z* 243.05(M+H).


[0340] The following compounds were prepared in accordance with the procedure described in the above Scheme 3.




[0341] N-(5-Bromopyridin-3-yl)propionamide (VIII): Off white solid (92% yield). ¹H NMR (DMSO-d₆) δ ppm 1.09 (t, *J*=7.54 Hz, 3H), 2.36 (q, *J*=7.54 Hz, 2H), 8.36 (m, 2H), 8.65 (d, *J*=2.07 Hz, 1H), 10.26 (s, 1H); ESIMS found C₈H₉BrN₂O *m/z* 231 (M+H).




[0342] N-(5-Bromopyridin-3-yl)butyramide (IX): Yellow solid (2.1 g, 8.64 mmol, 88.8% yield). ESIMS found C₉H₁₁BrN₂O *m/z* 243 (M+H).

**X**


[0343] N-(5-Bromopyridin-3-yl)pentanamide (**X**): Yellow solid (2.0 g, 7.78 mmol, 85.3% yield). ESIMS found $C_{10}H_{13}BrN_2O$ m/z 257 ($M+H$).

**XI**


[0344] N-(5-Bromopyridin-3-yl)-3-methylbutanamide (**XI**): Off white solid, (67% yield), ^1H NMR ($CDCl_3$, 400 MHz) δ ppm 8.55-8.42 (m, 3H), 7.62 (s, 1H), 2.31-2.18 (m, 3H), 1.02-1.01 (d, $J = 6$ Hz, 6H); ESIMS found $C_{10}H_{13}BrN_2O$ m/z 258.80 ($M+H$).

**XII**


[0345] N-(5-Bromopyridin-3-yl)-3,3-dimethylbutanamide (**XII**): Yellow solid (1.7 g, 6.27 mmol, 78.6% yield). ESIMS found $C_{11}H_{15}BrN_2O$ m/z 271 ($M+H$).

**XIII**


[0346] N-(5-Bromopyridin-3-yl)pivalamide (**XIII**): Off-white solid (1.082 g, 4.22 mmol, 73.1% yield). ^1H NMR ($DMSO-d_6$, 500 MHz) δ ppm 1.23 (s, 9H), 8.37 (d, $J=2$ Hz, 1H), 8.39 (t, $J=2$ Hz, 1H), 8.80 (d, $J=2$ Hz, 1H), 9.58 (brs, 1H); ESIMS found $C_{10}H_{13}BrN_2O$ m/z 257.0 ($M+H$).

**XIV**


[0347] N-(5-Bromopyridin-3-yl)-2-phenylacetamide (**XIV**): White solid (2.5 g, 8.59 mmol, 77.9% yield). ESIMS found $C_{13}H_{11}BrN_2O$ m/z 291 (M+H).

**XV**


[0348] N-(5-Bromopyridin-3-yl)benzamide (**XV**): White solid (2.7 g, 9.74 mmol, 60% yield). ESIMS found $C_{12}H_9BrN_2O$ m/z 277 (M+H).

**XVI**


[0349] N-(5-Bromopyridin-3-yl)cyclopropanecarboxamide (**XVI**): Off-white solid, (83% yield), ^1H NMR ($CDCl_3$, 400 MHz) δ ppm 8.46-8.39 (m, 3H), 7.54 (bs, 1H), 1.56-1.50 (m, 1H), 1.13-1.07 (m, 2H), 0.96-0.90 (m, 2H); ESIMS found for $C_9H_9BrN_2O$ m/z 240.9 (M+H).

**XVII**

[0350] N-(5-Bromopyridin-3-yl)cyclobutanecarboxamide (**XVII**): Yellow solid (2.1 g, 6.27 mmol, 86.6% yield). ESIMS found $C_{10}H_{11}BrN_2O$ m/z 255 (M+H).

**XVIII**


[0351] N-(5-Bromopyridin-3-yl)cyclopentanecarboxamide (**XVIII**): Yellow solid (1.9 g, 7.06 mmol, 80.2% yield). ESIMS found $C_{11}H_{13}BrN_2O$ m/z 269 (M+H).



XIX

[0352] N-(5-bromopyridin-3-yl)cyclohexanecarboxamide (XIX): Yellow solid (2.0 g, 7.06 mmol, 84.3% yield). ESIMS found $C_{12}H_{15}BrN_2O$ m/z 283 (M+H).

[0353] Preparation of intermediate 5-bromo-N,N-dimethylpyridin-3-amine (XXI) is depicted below in Scheme 4.




Scheme 4

Step 1

[0354] To a solution of 3,5-dibromopyridine (XX) (2.37 g, 10.0 mmol) in dry DMF (20.0 mL) was added K_2CO_3 (4.5 g, 33 mmol) and dimethylamino hydrochloride (1.79 g, 22 mmol). The mixture was heated overnight at 200°C in a sealed tube. The solution was cooled to room temperature and excess DMF was removed under vacuum. The residue was partitioned between EtOAc and water. The organic phase was separated. The aqueous phase was washed with EtOAc and the combined organic phases were dried over $MgSO_4$, and concentrated to afford 5-bromo-N,N-dimethylpyridin-3-amine (XXI) as an off-white solid (1.78 g, 8.85 mmol, 88% yield). ^1H NMR ($DMSO-d_6$, 500 MHz) δ ppm 2.94 (s, 6H), 7.25 (t, $J=2$ Hz, 1H), 7.91 (d, $J=2$ Hz, 1H), 8.07 (d, $J=2$ Hz, 1H); ESIMS found $C_7H_9BrN_2$ m/z 201.1 (M+H).

[0355] Preparation of intermediate 5-bromo-N-isopropylpyridin-3-amine (XXIII) is depicted below in Scheme 5.




Scheme 5

Steps 1

[0356] To a solution of 5-bromopyridin-3-amine (XXII) (535 mg, 3.09 mmol) in MeOH (62 mL) was added acetone (296 μ L, 4.02 mL). The pH was adjusted to 4 using HOAc and stirred for 30 min. NaCNBH₃ (272 mg, 4.33 mmol) was added and stirred at room temperature overnight. The MeOH was removed under vacuum and the residue was partitioned between EtOAc and saturated aqueous NaHCO₃. The organic layer was dried over MgSO₄ and evaporated under vacuum. The crude product was purified on a silica gel column (100% hexane $\rightarrow$ 90:10 hexane:EtOAc) to produce 5-bromo-N-isopropylpyridin-3-amine (XXIII) as an oil which slowly solidified into an off-white solid (309 mg, 1.44 mmol, 47% yield). ¹H NMR (DMSO-d₆, 500 MHz) δ ppm 1.12 (d, J =6.3Hz, 6H), 3.55-3.59 (m, 1H), 6.03 (d, J =7.9Hz, 1H), 7.05-7.06 (m, 1H), 7.75 (d, J =2Hz, 1H), 7.90 (d, J =2Hz, 1H); ESIMS found C₈H₁₁BrN₂ m/z 215 (M+H).

[0357] Preparation of intermediate 1-(5-bromopyridin-3-yl)-N,N-dimethylmethanamine (XXV) is depicted below in Scheme 6.




Scheme 6

Step 1

[0358] To a solution of 5-bromonicotinaldehyde (XXIV) (5.0 g, 26.9 mmol) in DCE (108 mL) was added dimethylamine-HCl (4.39 g, 53.8 mmol) and TEA (7.5 g, 53.8 mmol). The reaction was stirred at room temperature for 1 h. NaBH(OAc)₃ was


added and the reaction was stirred overnight at room temperature. The reaction was diluted with DCM and sat. aq. NaHCO_3 . The organic layer was separated, washed with water, brine, dried and concentrated under vacuum to produce 1-(5-bromopyridin-3-yl)-N,N-dimethylmethanamine (**XXV**) as a brown liquid (5.36 g, 24.9 mmol, 92.6% yield). ^1H NMR (CDCl_3) δ ppm 2.15 (s, 6H), 3.43 (s, 2H), 7.94 (s, 1H), 8.47 (d, $J=1.1\text{Hz}$, 1H), 8.59 (d, $J=2.2\text{Hz}$, 1H); ESIMS found $\text{C}_8\text{H}_{11}\text{BrN}_2$ m/z 215 ($\text{M}^{\text{Br}79}+\text{H}$) and 217 ($\text{M}^{\text{Br}81}+\text{H}$).

[0359] The following intermediates were prepared in accordance with the procedure described in the above Scheme 6.




XXVI

[0360] 3-Bromo-5-(pyrrolidin-1-ylmethyl)pyridine (**XXVI**): Golden liquid (1.35 g, 97% yield). ^1H NMR ($\text{DMSO}-d_6$) 1.68-1.71 (m, 4H), 2.42-2.44 (m, 4H), 3.60 (s, 2H), 7.96 (s, 1H), 8.48 (d, $J=2\text{Hz}$, 1H), 8.58 (d, $J=3\text{Hz}$, 1H); ESIMS found for $\text{C}_{10}\text{H}_{13}\text{BrN}_2$ m/z 242 ($\text{M}+\text{H}$).




XXVII

[0361] 3-Bromo-5-(piperidin-1-ylmethyl)pyridine (**XXVII**): Brown liquid (13.1 g, 94% yield). ^1H NMR ($\text{DMSO}-d_6$) 1.36-1.39 (m, 2H), 1.46-1.51 (m, 4H), 2.31-2.32 (m, 4H), 3.46 (s, 2H), 7.94 (s, 1H), 8.47 (d, $J=2\text{Hz}$, 1H), 8.58 (d, $J=3\text{Hz}$, 1H); ESIMS found for $\text{C}_{11}\text{H}_{15}\text{BrN}_2$ m/z 257 ($\text{M}+\text{H}$).



XXVIII


[0362] N-((5-Bromopyridin-3-yl)methyl)ethanamine (**XXVIII**): Golden liquid (1.29 g, 6.00 mmol, 60% yield). ESIMS found for $\text{C}_8\text{H}_{11}\text{BrN}_2$ m/z 215 ($\text{M}+\text{H}$).



XXIX

[0363] N-Benzyl-1-(5-bromopyridin-3-yl)methanamine (XXIX): Golden liquid (77 mg, 0.28 mmol, 25% yield). ESIMS found for $C_{13}H_{13}BrN_2$ m/z 277 (M+H).

[0364] Preparation of intermediate *tert*-butyl (5-bromopyridin-3-yl)methyl (cyclopentylmethyl)carbamate (XXXIV) is depicted below in Scheme 7.



Scheme 7

Step 1

[0365] To a solution of 5-bromonicotinaldehyde (XXIV) (2.0 g, 10.8 mmol, 1 eq) in MeOH (20 mL) was added $NaBH_4$ (2.4 g, 64.9 mmol, 6 eq) and the reaction mixture was stirred at room temperature for 3 h. The mixture was concentrated in vacuo and the residue was diluted in water (15 mL), the aqueous phase was extracted with DCM (10 mL x 3). The combined organic layers were dried over $MgSO_4$, filtered and concentrated in vacuo to afford (5-bromopyridin-3-yl)methanol (XXX) (1.8 g, 9.57 mmol, 90.0% yield) as a colorless oil. ^1H NMR ($CDCl_3$, 400 MHz) δ ppm 4.73 (s, 2H), 7.90 (s, 1H), 8.47 (s, 1H), 8.57 (s, 1H). ESIMS found for C_6H_6BrNO m/z 188 (M+H).

Step 2

[0366] To a stirred solution of (5-bromopyridin-3-yl)methanol (XXX) (1.60 g, 8.5 mmol, 1 eq), phthalimide (1.24 g, 8.5 mmol, 1 eq) and PPh_3 (3.33 g, 12.75 mmol, 1.5 eq) in anhydrous THF (15 mL) was added DEAD (2.21 g, 12.75 mmol, 1.5 eq) dropwise at $0^\circ C$ under N_2 . Then the reaction mixture was stirred at room temperature for 6 h. The mixture was washed with saturated $NaHCO_3$ solution (15 mL), water (15 mL) and brine

(15 mL) subsequently. The organic layers were dried over MgSO_4 , concentrated under reduced pressure, the resultant residue was purified by flash chromatography on silica gel (PE:EtOAc = 4:1) to give 2-((5-bromopyridin-3-yl)methyl)isoindoline-1,3-dione (**XXXI**) (2.5 g, 7.88 mmol, 82.3% yield) as a white solid. ESIMS found for $\text{C}_{14}\text{H}_9\text{BrN}_2\text{O}_2$ m/z 317 (M+H).

Step 3

[0367] A solution of 2-((5-bromopyridin-3-yl)methyl)isoindoline-1,3-dione (**XXXI**) (1.9 g, 6.0 mmol, 1 eq) and hydrazine hydrate (2.0 g, 40 mmol, 6 eq) in EtOH (20 mL) was heated at 70°C for 3 h. The mixture was filtered through a Celite pad and the filtrate was concentrated in vacuo, the crude product was dissolved in 1N HCl solution (15 mL) and concentrated to dryness, then it was washed with acetone (10 mL x 3), the precipitate was collected by filtration, dried in vacuo to give (5-bromopyridin-3-yl)methanamine (**XXXII**) (1.3 g, 6.95 mmol, 97.7% yield) as a white solid. ^1H NMR (D_2O , 400 MHz) δ ppm 4.34 (s, 2H), 8.56 (s, 1H), 8.75 (d, $J=1.2\text{Hz}$, 1H), 8.91 (d, $J=1.6\text{Hz}$, 1H). ESIMS found for $\text{C}_6\text{H}_7\text{BrN}_2$ m/z 187 (M+H).

Step 4


[0368] A solution of (5-bromopyridin-3-yl)methanamine (**XXXII**) (1.30 g, 5.8 mmol, 1.0 eq), cyclopentanecarbaldehyde (0.57 g, 5.8 mmol, 1.0 eq) and TEA (0.60 g, 5.8 mmol, 1.0 eq) in MeOH (15 mL) was stirred at room temperature for 2 h. Then NaBH_3CN (1.98 g, 34.6 mmol, 6.0 eq) was added and the mixture was stirred at the same temperature for another 3 h. The solvent was removed under reduced pressure and the residue was diluted in water (20 mL) and extracted with DCM (10 mL x 3), combined organic layers were dried over MgSO_4 and concentrated in vacuo to give 1-(5-bromopyridin-3-yl)-N-(cyclopentylmethyl)methanamine (**XXXIII**) (1.23 g, 4.57 mmol, 79.3% yield) as a brown oil. ESIMS found for $\text{C}_{12}\text{H}_{17}\text{BrN}_2$ m/z 269 (M+H).

Step 5

[0369] To a solution of 1-(5-bromopyridin-3-yl)-N-(cyclopentylmethyl)methanamine (**XXXIII**) (1.00 g, 3.7 mmol, 1 eq) and TEA (0.93 g, 9.2 mmol, 2.5 eq) in

DCM (20 mL) was added portionwise (Boc)₂O (0.85 g, 4.0 mmol, 1.1 eq) at 0°C, the reaction mixture was stirred at room temperature for 1 h. The mixture was washed with water (10 mL), brine (10 mL), the organic layer was separated, dried over MgSO₄ and concentrated *in vacuo* to give *tert*-butyl (5-bromopyridin-3-yl)methyl (cyclopentylmethyl)carbamate (**XXXIV**) (1.25 g, 3.38 mmol, 91.9% yield) as a white solid. ESIMS found for C₁₇H₂₅BrN₂O₂ *m/z* 369 (M+H).

[0370] Preparation of intermediate 3-(4-methyl-imidazol-1-yl)-benzene-1,2-diamine (**XXXVII**) is depicted below in Scheme 8.



Scheme 8

Step 1


[0371] A solution of 3-chloro-2-nitro-aniline (**XXXV**) (1.0 g, 5.8 mmol), potassium carbonate (2.4 g, 17.4 mmol), and 4-methylimidazole in dry DMF was heated overnight at 120°C under nitrogen. The reaction was cooled and the solvent was evaporated *in vacuo*. The residue was suspended in a saturated NaHCO₃ solution and extracted with CH₂Cl₂. The combined organic phases were dried over MgSO₄ and concentrated *in vacuo*. The crude product was purified by flash chromatography to provide 3-(4-methyl-imidazol-1-yl)-2-nitro-phenylamine (**XXXVI**). ¹H NMR (CDCl₃, 400 MHz) δ ppm 2.19 (s, 3H), 6.53 (m, 1H), 6.79 (m, 1H), 6.93 (m, 1H), 7.32 (m, 1H), 7.60 (m, 1H).

Step 2

[0372] To a solution of 3-(4-methyl-imidazol-1-yl)-2-nitro-phenylamine (**XXXVI**) in MeOH was added with 5% Pd/C. The combination was stirred under a hydrogen filled balloon at 40°C for 6 h. The solution was then filtered through a pad of Celite. The filtrate was concentrated *in vacuo* to get 3-(4-methyl-imidazol-1-yl)-

benzene-1,2-diamine (**XXXVII**). ^1H NMR (CDCl_3 , 400 MHz) δ ppm 2.17 (s, 3H), 6.54 (m, 1H), 6.80 (m, 1H), 6.97 (m, 1H), 7.28 (m, 1H), 7.56 (m, 1H).

[0373] Preparation of intermediate 2'-fluorobiphenyl-2,3-diamine (**XLI**) is depicted below in Scheme 9.



Scheme 9


Step 1

[0374] A solution of 3-bromo-2-nitroaniline (**XXXVIII**) (2.00 g, 9.30 mmol, 1 eq), 2-fluorophenylboronic acid (**XXXIX**) (1.42 g, 10.14 mmol, 1.1 eq), $\text{Pd(PPh}_3)_4$ (0.35 g, 0.03 mmol, 0.03 eq), Na_2CO_3 (1.95 g, 18.40 mmol, 2 eq) in a mixed solvent of toluene (15 mL), H_2O (9 mL) and EtOH (3 mL) was stirred at 75°C for 15 h under nitrogen atmosphere. Then the reaction mixture was washed with brine (20 mL) and dried over anhydrous Na_2SO_4 , filtered and concentrated in vacuo, the resultant residue was purified by chromatography on silica gel ($\text{PE:EtOAc} = 3:1$) to afford 2'-fluoro-2-nitrobiphenyl-3-amine (**XL**) (1.0 g, 4.30 mmol, 46.6% yield) as a yellow solid. ^1H NMR (DMSO-d_6 , 400 MHz) δ ppm 6.54 (d, $J=6.4\text{Hz}$, 1H), 6.64 (s, 2H), 7.04 (dd, $J=8.8\text{Hz}$, $J=1.2\text{Hz}$, 1H), 7.18-7.31 (m, 2H), 7.33-7.47 (m, 3H); ESIMS found for $\text{C}_{12}\text{H}_9\text{FN}_2\text{O}_2$ m/z 233 ($\text{M}+\text{H}$).

Step 2


[0375] To a solution of 2'-fluoro-2-nitrobiphenyl-3-amine (**XL**) (1.0 g, 3.45 mmol, 1 eq) in MeOH (50 mL) was added Pd/C (0.5 g) under nitrogen atmosphere, the mixture was stirred under 50 psi of H_2 for 6 h at room temperature. Then the mixture was filtered and concentrated in vacuo to afford 2'-fluorobiphenyl-2,3-diamine (**XLI**) (0.8 g, 3.96 mmol, 92% yield) as a black solid. ^1H NMR (DMSO-d_6 , 400 MHz) δ ppm 3.99 (s, 2H), 4.62 (s, 2H), 6.32 (d, $J=7.6\text{Hz}$, 1H), 6.49 (t, $J=7.6\text{Hz}$, 1H), 6.60 (d, $J=7.6\text{Hz}$, 1H), 7.21-7.35 (m, 3H), 7.35-7.45 (m, 1H); ESIMS found for $\text{C}_{12}\text{H}_{11}\text{FN}_2$ m/z 203 ($\text{M}+\text{H}$).

[0376] The following intermediates were prepared in accordance with the procedure described in the above Scheme 9.




XLII

[0377] 3'-Fluorobiphenyl-2,3-diamine (XLII): White solid (2.0 g, 9.89 mmol, 81% yield). ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 4.16 (s, 2H), 4.64 (s, 2H), 6.38 (dd, $J=7.6\text{Hz}$, $J=1.6\text{Hz}$, 1H), 6.51 (t, $J=7.6\text{Hz}$, 1H), 6.60 (d, $J=6\text{Hz}$, 1H), 7.11-7.26 (m, 3H), 7.48 (q, $J=6.4\text{Hz}$, 1H); ESIMS found for $\text{C}_{12}\text{H}_{11}\text{FN}_2$ m/z 203 (M+H).



XLIII


[0378] 4'-Fluorobiphenyl-2,3-diamine (XLIII): White solid (2.4 g, 11.87 mmol, 98% yield). ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 4.07 (s, 2H), 4.60 (s, 2H), 6.34 (dd, $J=7.6\text{Hz}$, $J=1.6\text{Hz}$, 1H), 6.50 (t, $J=7.6\text{Hz}$, 1H), 6.58 (dd, $J=7.6\text{Hz}$, $J=1.6\text{Hz}$, 1H), 7.26 (t, $J=7.6\text{Hz}$, 2H), 7.40 (q, $J=5.6\text{Hz}$, 2H); ESIMS found for $\text{C}_{12}\text{H}_{11}\text{FN}_2$ m/z 203 (M+H).



XLIV


[0379] 3-(Pyridin-3-yl)benzene-1,2-diamine (XLIV): White solid (1.36 g, 7.34 mmol, 92.5% yield). ^1H NMR (CDCl_3 , 400 MHz) δ ppm 1.57 (brs, 2H), 3.42 (brs,

2H), 6.66 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 1H), 6.68-6.72 (m, 2H), 7.31 (dd, $J=8\text{Hz}$, $J=4.8\text{Hz}$, 1H), 7.71 (td, $J=8\text{Hz}$, $J=2\text{Hz}$, 1H), 8.54 (dd, $J=4.8\text{Hz}$, $J=1.6\text{Hz}$, 1H), 8.64 (d, $J=1.6\text{Hz}$, 1H); ESIMS found for $\text{C}_{11}\text{H}_{11}\text{N}_3$ m/z 186 ($\text{M}+\text{H}$).



XLV


[0380] 3-(Thiophen-3-yl)benzene-1,2-diamine (XLV): White solid (1.2 g, 6.31 mmol, mmol, 94% yield). ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 4.19 (s, 2H), 4.59 (s, 2H), 6.47 (dd, $J=4.8\text{Hz}$, $J=1\text{Hz}$, 2H), 6.55 (q, $J=4.8\text{Hz}$, 1H), 7.24 (dd, $J=4.8\text{Hz}$, $J=1\text{Hz}$, 1H), 7.50 (t, $J=1.6\text{Hz}$, 1H), 7.63 (dd, $J=4.8\text{Hz}$, $J=2.8\text{Hz}$, 1H); ESIMS found for $\text{C}_{10}\text{H}_{10}\text{N}_2\text{S}$ m/z 191 ($\text{M}+\text{H}$).



XLVI

[0381] 3-(Furan-3-yl)benzene-1,2-diamine (XLVI): White solid (1.3 g, 7.46 mmol, mmol, 85% yield). ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 4.24 (brs, 2H), 4.57 (brs, 2H), 6.46-6.50 (m, 1H), 6.50-6.56 (m, 2H), 6.72 (s, 1H), 7.74 (t, $J=1.6\text{Hz}$, 1H), 7.87 (s, 1H); ESIMS found for $\text{C}_{10}\text{H}_{10}\text{N}_2\text{O}$ m/z 175 ($\text{M}+\text{H}$).

[0382] Preparation of intermediate (XLIX) is depicted below in Scheme 10.



Scheme 10


Step 1

[0383] A mixture of 1-methylpiperazine (XLVII) (20 mL) and 3-chloro-2-nitroaniline (XXXV) (1.5 g, 8.7mmol, 1 eq) was stirred at 50°C for 1 h under microwave irradiation. The reaction mixture was diluted with water (100 mL) and filtered, the cake washed with water (30 mL x 3), dried in vacuo to give the 3-(4-methylpiperazin-1-yl)-2-nitroaniline (XLVIII) (1.64 g, 6.94 mmol, 80% yield) as a yellow solid. ESIMS found for C₁₁H₁₆N₄O₂ *m/z* 237 (M+H).

Step 2

[0384] A mixture of 3-(4-methylpiperazin-1-yl)-2-nitroaniline (XLVIII) (1.64 g, 6.9 mmol, 1 eq) and Pd/C (0.2 g) in MeOH (20 mL) was stirred under 30 psi of H₂ at room temperature overnight. The reaction was monitored by TLC. The mixture was filtered and the filtrate was concentrated in vacuo to give the 3-(4-methylpiperazin-1-yl)benzene-1,2-diamine (XLIX) (1.31 g, 6.35 mmol, 92% yield) as a black solid. ¹H NMR (CDCl₃, 400 MHz) δ ppm 2.30 (s, 3H), 3.30 (brs, 2H), 3.68 (brs, 2H), 6.46 (dd, *J*=7.2Hz, *J*=2Hz, 1H), 6.54-6.63 (m, 2H); ESIMS found for C₁₁H₁₈N₄ *m/z* 207 (M+H).

[0385] Preparation of intermediate 3-(piperidin-1-yl)benzene-1,2-diamine (LI) is depicted below in Scheme 11.



Scheme 11

Step 1


[0386] To a solution of 3-chloro-2-nitroaniline (XXXV) (2.00 g, 11.6 mmol, 1 eq) and piperidine (2.95 g, 34.7 mmol, 3 eq) in DMF (60 ml) was added K₂CO₃ (4.78 g, 34.4 mmol, 3 eq) in one portion and the mixture was stirred at 120°C under nitrogen overnight. The reaction mixture was diluted with EtOAc (60 ml) and washed with saturated NaHCO₃ solution (50 mL). The organic phases were dried over Na₂SO₄ and concentrated in vacuo, the resultant residue was purified by silica gel column chromatography (PE:EtOAc = 5:1→1:1) to give 2-nitro-3-(piperidin-1-yl)aniline (L) (1.8

g, 8.14 mmol, 70.3% yield) as a black solid. ESIMS found for $C_{11}H_{15}N_3O_2$ m/z 222 (M+H).

Step 2

[0387] A mixture of 2-nitro-3-(piperidin-1-yl)aniline (**L**) (1.64 g, 6.9mmol, 1 eq) and Pd/C (0.50 g) in MeOH (20 mL) was stirred at room temperature under 30 psi H_2 overnight. After the starting material was consumed completely, the mixture was filtered through a Celite pad and the filtrate was concentrated in vacuo to give the 3-(piperidin-1-yl)benzene-1,2-diamine (**LI**) (1.1 g, 5.75 mmol, 76% yield) as a yellow solid. ^1H NMR ($CDCl_3$, 400 MHz) δ ppm 1.59 (brs, 2H), 1.73 (quin, $J=5.6Hz$, 4H), 2.84 (brs, 4H), 3.50 (brs, 4H), 6.52 (dd, $J=6.4Hz$, $J=1.6Hz$, 1H), 6.59-6.75 (m, 2H); ESIMS found for $C_{11}H_{17}N_3$ m/z 192 (M+H).

[0388] Preparation of intermediate 3-(pyridin-4-yl)benzene-1,2-diamine (**LVII**) is depicted below in Scheme 12.



Scheme 12

Step 1

[0389] To a solution of 2-bromoaniline (**LII**) (50 g, 0.29 mol, 1 eq) in acetic anhydride (265 mL) was added dropwise nitric acid (fuming) (36.75 mL, 0.93 mol, 3.2 eq) at 0°C and then stirred at that temperature, when the starting material was consumed,

the mixture was filtered, the filtrate was poured into ice water. The aqueous phase was basified with aqueous solution of sodium bicarbonate to pH=7, then the mixture was extracted with EtOAc (30 mL x 3). The organic layers were combined, dried and concentrated in vacuo to give the N-(2-bromo-6-nitrophenyl)acetamide (**LIII**) (12.6 g, 48.6 mmol, 16.7% yield) as a white solid. ¹H NMR (DMSO-d₆, 400 MHz) δ ppm 2.06 (s, 3H), 7.43 (t, *J*=8Hz, 1H), 7.94 (d, *J*=8Hz, 1H), 8.05 (d, *J*=8Hz, 1H); ESIMS found for C₈H₇BrN₂O₃ *m/z* 259 (M+H).

Step 2

[0390] A degassed mixture of N-(2-bromo-6-nitrophenyl)acetamide (**LIII**) (2.59 g, 10 mmol, 1.0 eq), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (**LIV**) (2.05 g, 10 mmol, 1.3 eq), Na₂CO₃ (2.12 g, 20 mmol, 2 eq) and Pd(PPh₃)₄ (1.16 g, 1 mmol, 0.1 eq) in a mixed solvent of DME (30 mL) and H₂O (10 mL) was heated to reflux under nitrogen overnight, the mixture was poured onto water (40 ml) and extracted with EtOAc (30 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated in vacuo, purification the resultant residue was purified by column chromatography (EtOAc:PE=1:4→100% EtOAc) to afford N-(2-nitro-6-(pyridin-4-yl)phenyl)acetamide (**LV**) (1.42 g, 5.52 mmol, 55% yield) as a yellow solid. ¹H NMR (DMSO-d₆, 400 MHz) δ ppm 1.92 (s, 3H), 7.46 (d, *J*=5.6Hz, 2H), 7.69 (t, *J*=8Hz, 1H), 7.80 (dd, *J*=7.6Hz, *J*=1.2Hz, 1H), 8.06 (dd, *J*=8Hz, *J*=1.6Hz, 1H), 8.73 (d, *J*=6Hz, 2H), 9.96 (s, 1H); ESIMS found for C₁₃H₁₁N₃O₃ *m/z* 258 (M+H).


Step 3

[0391] To a solution of N-(2-nitro-6-(pyridin-4-yl)phenyl)acetamide (**LV**) (3.94 g, 15 mmol, 1 eq) in MeOH (20 mL) was added 2 N aqueous NaOH solution (50 mL) and the mixture was refluxed until the starting material was consumed completely, the precipitate was collected by filtration to afford the 2-nitro-6-(pyridin-4-yl)aniline (**LVI**) (3.0 g, 13.9 mmol, 91% yield) as yellow solid. ESIMS found for C₁₁H₉N₃O₂ *m/z* 216 (M+H).

Step 4

[0392] To a solution of 2-nitro-6-(pyridin-4-yl)aniline (LVI) (3 g, 14 mmol, 1 eq) in EtOAc (350 mL) was added Pd/C (0.3 g) and the mixture was stirred at room temperature under 1 atm of H₂ atmosphere overnight, the mixture was filtered and concentrated in vacuo to give the product 3-(pyridin-4-yl)benzene-1,2-diamine (LVII) (2.4 g, 13.0 mmol, 93% yield) as a white solid. ¹H NMR (DMSO-d₆, 400 MHz) δ ppm 4.35 (s, 2H), 4.75 (s, 2H), 6.45 (dd, *J*=7.6Hz, *J*=1Hz, 1H), 6.58 (t, *J*=7.6Hz, 1H), 6.67 (d, *J*=6.8Hz, 1H), 7.47 (d, *J*=6Hz, 2H), 8.65 (d, *J*=6Hz, 2H); ESIMS found for C₁₁H₁₁N₃ *m/z* 186 (M+H).

[0393] Preparation of intermediate 3-(pyridin-2-yl)benzene-1,2-diamine 3HCl (LXII) is depicted below in Scheme 13.



Scheme 13

Step 1

[0394] To a solution of 2-bromopyridine (LVIII) (10 g, 63 mmol, 1.00 eq) in THF (150 mL) was added *n*-BuLi (25.3 mL, 63 mmol, 1.00 eq) and the mixture was stirred at -70°C for 30 min under nitrogen atmosphere. Then *n*-Bu₃SnCl (21.7 g, 67 mmol, 1.06 eq) was added and the mixture was stirred at the same temperature for another 2 h. Saturated ammonium chloride solution (150 mL) was added to the solution

and extracted with EtOAc (150 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated in vacuo to afford the crude 2-(tributylstannyl)pyridine (**LIX**) (25.9 g, 63 mmol, 100% yield) as a yellow oil. The crude product was used without further purification.

Step 2

[0395] A degassed mixture of N-(2-bromo-6-nitrophenyl)acetamide (**LIII**) (4.8 g, 19 mmol, 1.00 eq), 2-(tributylstannyl)pyridine (**LIX**) (7.5 g, 20 mmol, 1.05 eq) and Pd(PPh₃)₄ (2.1 g, 1.8 mmol, 0.01 eq) in toluene (60 mL) was heated to reflux under nitrogen overnight. Saturated sodium bicarbonate solution (50 mL) was then added to the mixture and it was extracted with EtOAc (50 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated in vacuo, the residue was purified by column chromatography on silica gel (EtOAc:PE=1:2→100% EtOAc) to afford N-(2-nitro-6-(pyridin-2-yl)phenyl)acetamide (**LX**) (4.4 g, 17.1 mmol, 92% yield) as a white-off solid. ¹H NMR (DMSO-d₆, 400 MHz) δ ppm 1.93 (s, 3H), 7.43-7.51 (m, 1H), 7.51-7.65 (m, 1H), 7.67 (d, *J*=7.6Hz, 1H), 7.97 (dd, *J*=7.6Hz, *J*=2.4Hz, 3H), 8.75 (d, *J*=4.4Hz, 1H), 10.52 (s, 1H); ESIMS found for C₁₃H₁₁N₃O₃ *m/z* 258 (M+H).

Step 3

[0396] To a solution of N-(2-nitro-6-(pyridin-2-yl)phenyl)acetamide (**LX**) (4.41 g, 17 mmol, 1 eq) in MeOH (20 mL) was added 2N NaOH aqueous (50 mL) and the mixture was refluxed until the stirring material was consumed completely. The mixture was concentrated in vacuo to remove the MeOH and the precipitate was collected by filtration to afford 2-nitro-6-(pyridin-2-yl)aniline (**LXI**) (2.4 g, 11.2 mmol, 65% yield) as a yellow solid. ESIMS found for C₁₁H₉N₃O₂ *m/z* 216 (M+H).

Step 4

[0397] To a solution of 2-nitro-6-(pyridin-2-yl)aniline (**LXI**) (2.4 g, 0.01 mmol, 1 eq) in EtOAc (350 mL) was added Pd/C (1 g) and the mixture was stirred at room temperature under 1 atm of H₂ atmosphere overnight, filtered and then concentrated


in vacuo, to give 3-(pyridin-2-yl)benzene-1,2-diamine (1.9 g, 10.3 mmol, 89% yield) as a yellow oil. ESIMS found for $C_{11}H_{11}N_3$ m/z 186 (M+H).

Step 5

[0398] To a solution of 3-(pyridin-2-yl)benzene-1,2-diamine (1.86 g, 0.01 mmol) in EtOAc (200 mL) was added HCl in EtOAc (40 mL) and the mixture was stirred at 0°C for 20 min. The precipitate was collected by filtration to give 3-(pyridin-2-yl)benzene-1,2-diamine-3HCl (**LXII**) as a yellow solid. ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 6.89 (t, $J=7.6$ Hz, 1H), 7.33 (brs, 1H), 7.51 (d, $J=7.2$ Hz, 1H), 7.54-7.66 (m, 2H), 7.97 (d, $J=8$ Hz, 1H), 8.16 (brs, 1H), 8.75 (brs, 1H).

Example 1.

[0399] Preparation of N-(5-(3-(4-(4-methylpiperazin-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-2-phenylacetamide (**202**) is depicted below in Scheme 14.



Scheme 14

Step 1

[0400] A solution of N-(5-bromopyridin-3-yl)-2-phenylacetamide (**XIV**) (2.2 g, 5.5 mmol, 1.00 eq), 5-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazole-3-carbaldehyde (**IV**) (1.5 g, 5.5 mmol, 1.00 eq), Pd(dppf)Cl₂ (0.28 g, 0.39 mmol, 0.07 eq) and Na₂CO₃ (0.8 g, 16.5 mmol, 3.00 eq) in a mixed solvent of 1,2-dimethoxyethane (30 mL) and H₂O (5 mL) was refluxed for 3 h under nitrogen atmosphere. The reaction mixture was diluted in water (30 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated in vacuo, the resultant residue was purified by flash chromatography on silica gel (PE/EtOAc = 10:1→3:1) to give N-(5-(3-formyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazol-5-yl)pyridin-3-yl)-2-phenylacetamide (**LXIII**) (2.3 g, 4.73 mmol, 51% yield) as a brown oil. ESIMS found C₂₇H₃₀N₄O₃Si *m/z* 487 (M+H).

Step 2


[0401] To a well stirred solution of N-(5-(3-formyl-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazol-5-yl)pyridin-3-yl)-2-phenylacetamide (**LXIII**) (100 mg, 0.205 mmol, 1.0 eq), 3-(4-methylpiperazin-1-yl)benzene-1,2-diamine (**XLIX**) (46 mg, 0.223 mmol, 1.09 eq) in CH₃CN (2 mL) was added compound 2,3,5,6-tetrachlorocyclohexa-2,5-diene-1,4-dione (60 mg, 1.1 eq). The reaction mixture was refluxed for 3 h, TLC analysis (1:1, PE/EtOAc) showed complete consumption of starting material (**LXIII**). The precipitate was collected by filtration, washed with CH₃CN and dried under reduced pressure to give N-(5-(3-(4-(4-methylpiperazin-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazol-5-yl)pyridin-3-yl)-2-phenylacetamide (**LXIV**) (41 mg, 0.061 mmol, 29.6% yield). ESIMS found for C₃₈H₄₄N₈O₂Si *m/z* 673 (M+H).

Step 3

[0402] To a well stirred solution of N-(5-(3-(4-(4-methylpiperazin-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-indazol-5-yl)pyridin-3-yl)-2-phenylacetamide (**LXIV**) (40 mg, 0.06 mmol, 1.0 eq) in DCM (5 mL) was added

TFA (0.3 mL) at room temperature for 2 h, TLC analysis (10:1, DCM/MeOH) showed complete consumption of compound (**LXIV**). Then the reaction mixture was added NH_4OH (2 mL) and washed with EtOAc and water, dried over Na_2SO_4 , filtration, and concentration, the crude product was obtained as a solid, which was purified by preparative HPLC to give N-(5-(3-(4-(4-methylpiperazin-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-2-phenylacetamide **202** (20 mg, 0.037 mmol, 60.9% yield) as a white solid. ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.98 (s, 3H), 3.22-3.31 (m, 2H), 3.51-3.61 (m, 2H), 3.67-3.76 (m, 2H), 3.88 (s, 2H), 4.11-4.20 (m, 2H), 7.03 (d, $J=7.6\text{Hz}$, 1H), 7.27-7.51 (m, 7H), 7.91 (ABq, 2H), 8.84 (s, 1H), 8.88 (s, 1H), 8.93 (s, 1H), 9.06 (s, 1H); ESIMS found for $\text{C}_{32}\text{H}_{30}\text{N}_8\text{O}$ m/z 543.3 ($\text{M}+\text{H}$).


[0403] The following compounds were prepared in accordance with the procedure described in the above Example 1.



1

[0404] N-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)propionamide **1**

[0405] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.29 (t, $J=7.2\text{Hz}$, 3H), 2.60 (q, $J=7.6\text{Hz}$, 2H), 7.23-7.40 (m, 2H), 7.61-7.77 (m, 5H), 7.94 (d, $J=8\text{Hz}$, 1H), 7.99 (s, 1H), 8.87 (s, 1H), 8.95 (s, 1H), 9.04 (s, 1H), 9.33 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{21}\text{FN}_6\text{O}$ m/z 477.2 ($\text{M}+\text{H}$).




96

4

[0406] 3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-5-(pyridin-3-yl)-1H-indazole 4


[0407] ^1H NMR (CD_3OD , 400 MHz) δ ppm 7.32 (t, $J=8\text{Hz}$, 1H), 7.59-7.73 (m, 4H), 7.78 (t, $J=7.6\text{Hz}$, 1H), 8.02 (d, $J=8.4\text{Hz}$, 2H), 8.09 (d, $J=8.4\text{Hz}$, 1H), 8.29 (brs, 1H), 8.93 (d, $J=5.2\text{Hz}$, 2H), 9.18 (d, $J=7.6\text{Hz}$, 1H), 9.46 (s, 1H); ESIMS found for $\text{C}_{25}\text{H}_{16}\text{FN}_5$ m/z 406.1 (M+H).



5

[0408] 3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-5-(4-methylpyridin-3-yl)-1H-indazole 5

[0409] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.45 (s, 3H), 7.08 (dt, 1H), 7.36 (t, $J=7.6\text{Hz}$, 1H), 7.41-7.53 (m, 4H), 7.62 (d, $J=8\text{Hz}$, 1H), 7.74 (d, $J=8.4\text{Hz}$, 1H), 7.97 (d, 1H), 8.43 (brs, 1H), 8.51 (brs, 1H), 8.60 (s, 1H); ESIMS found for $\text{C}_{26}\text{H}_{18}\text{FN}_5$ m/z 420.1 (M+H).




7

[0410] 5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)-N,N-dimethylpyridin-3-amine 7

[0411] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.30 (s, 6H), 7.32 (t, 1H), 7.58-7.68 (m, 3H), 7.70 (d, $J=7.6\text{Hz}$, 1H), 7.78 (t, $J=7.6\text{Hz}$, 1H), 7.94-8.08 (m, 3H), 8.17 (s,


1H), 8.21 (s, 1H), 8.51 (s, 1H), 8.82 (s, 1H); ESIMS found for C₂₇H₂₁FN₆ *m/z* 449.1 (M+H).



9

[0412] N-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)isobutyramide **9**


[0413] ¹H NMR (CD₃OD, 400 MHz) δ ppm 1.31 (d, *J*=6.8Hz, 6H), 2.83 (sep, *J*=6.8Hz, 1H), 7.24-7.33 (m, 1H), 7.62-7.71 (m, 4H), 7.72 (t, *J*=8Hz, 1H), 7.97 (d, 1H), 8.00 (s, 2H), 8.86 (s, 1H), 9.01 (s, 1H), 9.06 (s, 1H), 9.36 (s, 1H); ESIMS found for C₂₉H₂₃FN₆O *m/z* 491.2 (M+H).



10


[0414] N-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-2-phenylacetamide **10**

[0415] ¹H NMR (CD₃OD, 400 MHz) δ ppm 3.89 (s, 2H), 7.22-7.46 (m, 6H), 7.60-7.77 (m, 5H), 7.95 (d, *J*=8Hz, 1H), 8.00 (s, 2H), 8.81 (s, 1H), 9.01 (s, 1H), 9.03 (s, 1H), 9.29 (s, 1H); ESIMS found for C₃₃H₂₃FN₆O *m/z* 539.1 (M+H).

**13**


[0416] 1-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-N,N-dimethylmethanamine **13**

[0417] ^1H NMR (CD_3OD , 400 MHz) δ ppm 3.12 (s, 6H), 4.80 (s, 2H), 7.36 (dt, $J=8.8\text{Hz}$, $J=2\text{Hz}$, 1H), 7.64-7.77 (m, 4H), 7.83 (t, $J=7.2\text{Hz}$, 1H), 8.06 (d, $J=8.8\text{Hz}$, 1H), 8.12 (d, $J=8\text{Hz}$, 1H), 8.20 (d, $J=8.4\text{Hz}$, 1H), 9.11 (s, 1H), 9.41 (s, 1H), 9.51 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{23}\text{FN}_6$ m/z 463.2 ($\text{M}+\text{H}$).

**16**


[0418] N-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-3,3-dimethylbutanamide **16**

[0419] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.04 (s, 9H), 2.30 (s, 2H), 7.09 (t, $J=8\text{Hz}$, 1H), 7.41-7.52 (m, 3H), 7.57 (d, $J=7.6\text{Hz}$, 1H), 7.60-7.71 (m, 2H), 7.76 (Abq, 2H), 8.69 (s, 1H), 8.71 (s, 1H), 8.83 (brs, 1H), 9.17 (brs, 1H); ESIMS found for $\text{C}_{31}\text{H}_{27}\text{FN}_6\text{O}$ m/z 519.2 ($\text{M}+\text{H}$).

**18**


[0420] N-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **18**

[0421] ^1H NMR (CD_3OD , 400 MHz) δ ppm 0.90 (t, $J=7.2\text{Hz}$, 3H), 1.36 (sex, $J=7.2\text{Hz}$, 2H), 1.66 (quin, $J=7.6\text{Hz}$, 2H), 2.47 (t, $J=7.6\text{Hz}$, 2H), 7.13-7.22 (m, 1H), 7.50-7.59 (m, 4H), 7.63 (t, $J=8\text{Hz}$, 1H), 7.82-7.93 (m, 3H), 8.72 (s, 1H), 8.88 (s, 1H), 8.95 (s, 1H), 9.24 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{25}\text{FN}_6\text{O}$ m/z 505.2 (M+H).

**19**


[0422] N-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopropanecarboxamide **19**

[0423] ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm 0.95 (d, $J=7.6\text{Hz}$, 4H), 1.94-2.05 (m, 1H), 7.22 (dt, $J=6.8\text{Hz}$, $J=1.6\text{Hz}$, 1H), 7.31 (s, 1H), 7.42 (t, $J=7.6\text{Hz}$, 2H), 7.55-7.71 (m, 3H), 7.91 (Abq, 2H), 8.10 (d, $J=6.8\text{Hz}$, 1H), 8.22 (d, $J=10.4\text{Hz}$, 1H), 8.92 (s, 1H), 8.97 (s, 1H), 9.02 (s, 1H), 9.17 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{21}\text{FN}_6\text{O}$ m/z 489.2 (M+H).

**21**


[0424] N-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopentanecarboxamide **21**

[0425] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.66-1.78 (m, 2H), 1.78-1.89 (m, 2H), 1.89-2.01 (m, 2H), 2.01-2.12 (m, 2H), 3.01 (quin, $J=8\text{Hz}$, 1H), 7.26-7.35 (m, 1H), 7.62-7.73 (m, 4H), 7.76 (t, $J=8\text{Hz}$, 1H), 7.98 (d, $J=8\text{Hz}$, 1H), 8.02 (s, 2H), 8.83 (s, 1H), 9.03 (s, 1H), 9.05 (s, 1H), 9.35 (d, $J=1.6\text{Hz}$, 1H); ESIMS found for $\text{C}_{31}\text{H}_{25}\text{FN}_6\text{O}$ m/z 517.2 ($\text{M}+\text{H}$).

**22**


[0426] N-(5-(3-(4-(3-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclohexanecarboxamide **22**

[0427] ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm 1.27 (d, $J=14.8\text{Hz}$, 1H), 1.38 (q, $J=13.6\text{Hz}$, 2H), 1.54 (q, $J=10.4\text{Hz}$, 2H), 1.72 (d, $J=12.4\text{Hz}$, 1H), 1.84 (d, $J=13.6\text{Hz}$, 2H), 1.94 (d, $J=12.4\text{Hz}$, 2H), 2.49 (q, $J=10.8\text{Hz}$, 1H), 7.18-7.28 (m, 1H), 7.54-7.64 (m, 4H), 7.68 (t, $J=8\text{Hz}$, 1H), 7.90 (d, $J=8\text{Hz}$, 1H), 7.95 (s, 2H), 8.76 (s, 1H), 8.94 (s, 1H), 8.97 (s, 1H), 9.27 (s, 1H); ESIMS found for $\text{C}_{32}\text{H}_{27}\text{FN}_6\text{O}$ m/z 531.2 ($\text{M}+\text{H}$).

**26**


[0428] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide **26**

[0429] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.08 (d, $J=6.4\text{Hz}$, 6H), 2.28 (non, $J=6.8\text{Hz}$, 1H), 2.45 (d, $J=7.2\text{Hz}$, 2H), 7.39 (t, $J=8.4\text{Hz}$, 1H), 7.64 (d, 1H), 7.72 (t, 1H), 7.87 (dd, $J=8.4\text{Hz}$, $J=5.2\text{Hz}$, 2H), 7.94 (d, 1H), 8.00 (s, 2H), 8.84 (s, 1H), 9.02 (s, 1H), 9.06 (s, 1H), 9.32 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{25}\text{FN}_6\text{O}$ m/z 505.1 ($\text{M}+\text{H}$).

**29**


[0430] 3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-5-(4-methylpyridin-3-yl)-1H-indazole **29**

[0431] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.69 (s, 3H), 7.30-7.42 (m, 2H), 7.63 (d, $J=6\text{Hz}$, 1H), 7.69-7.87 (m, 4H), 7.94-8.06 (m, 2H), 8.14 (brs, 1H), 8.56 (brs, 1H), 8.80 (brs, 1H), 8.92 (brs, 1H); ESIMS found for $\text{C}_{26}\text{H}_{18}\text{FN}_5$ m/z 420.2 ($\text{M}+\text{H}$).

**32**


[0432] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pivalamide **32**

[0433] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.42 (s, 9H), 7.39 (t, $J=8.4\text{Hz}$, 2H), 7.65 (d, $J=7.2\text{Hz}$, 1H), 7.74 (t, $J=7.6\text{Hz}$, 1H), 7.80-7.91 (m, 2H), 7.98 (d, $J=8\text{Hz}$, 1H), 8.02 (s, 2H), 8.87 (s, 1H), 9.10 (s, 1H), 9.26 (s, 1H), 9.39 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{25}\text{FN}_6\text{O}$ m/z 505.2 (M+H).

**33**


[0434] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)isobutyramide **33**

[0435] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.29 (d, $J=6.8\text{Hz}$, 6H), 2.79 (sep, $J=6.8\text{Hz}$, 1H), 7.37 (t, $J=8.8\text{Hz}$, 2H), 7.62 (d, $J=7.2\text{Hz}$, 1H), 7.70 (t, 1H), 7.82-7.93 (m, 3H), 7.98 (s, 2H), 8.78 (s, 1H), 8.98 (s, 1H), 9.00 (s, 1H), 9.23 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{23}\text{FN}_6\text{O}$ m/z 491.2 (M+H).

**35**


[0436] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)benzamide **35**

[0437] ^1H NMR (CD_3OD , 400 MHz) δ ppm 7.43 (t, 2H), 7.61-7.82 (m, 5H), 7.92 (dd, 2H), 8.00 (d, 1H), 8.08 (Abq, $J=5.6\text{Hz}$, 2H), 8.16 (d, $J=7.6\text{Hz}$, 2H), 8.89 (s, 1H), 9.13 (s, 1H), 9.34 (s, 1H), 9.47 (s, 1H); ESIMS found for $\text{C}_{32}\text{H}_{21}\text{FN}_6\text{O}$ m/z 525.1 (M+H).

**38**


[0438] 3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-5-(5-(pyrrolidin-1-ylmethyl)pyridin-3-yl)-1H-indazole **38**

[0439] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.14 (brs, 2H), 2.26 (brs, 2H), 3.35 (brs, 4H), 3.70 (brs, 2H), 7.36 (t, $J=8\text{Hz}$, 2H), 7.62 (d, $J=7.2\text{Hz}$, 1H), 7.72 (t, 1H), 7.82 (brs, 2H), 7.98 (d, $J=8.4\text{Hz}$, 1H), 8.03 (d, $J=8\text{Hz}$, 1H), 8.12 (d, $J=8.4\text{Hz}$, 1H), 9.02 (s, 1H), 9.03 (s, 1H), 9.32 (s, 1H), 9.38 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{25}\text{FN}_6$ m/z 489.2 (M+H).

**41**


[0440] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzimidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)butyramide **41**

[0441] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.12 (t, $J=7.2\text{Hz}$, 3H), 1.87 (sex, $J=7.6\text{Hz}$, 2H), 2.62 (t, $J=7.6\text{Hz}$, 2H), 7.45 (t, $J=8.4\text{Hz}$, 2H), 7.73 (dd, $J=6.8\text{Hz}$, $J=1\text{Hz}$, 1H), 7.81 (t, $J=8\text{Hz}$, 1H), 7.90 (dd, $J=8.4\text{Hz}$, $J=5.2\text{Hz}$, 2H), 8.02 (d, $J=7.6\text{Hz}$, 1H), 8.07 (d, $J=1\text{Hz}$, 2H), 8.87 (s, 1H), 9.11 (s, 1H), 9.12 (s, 1H), 9.37 (d, $J=1.6\text{Hz}$, 1H); ESIMS found for $\text{C}_{29}\text{H}_{23}\text{FN}_6\text{O}$ m/z 491.2 (M+H).

**42**


[0442] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **42**

[0443] ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 0.92 (t, $J=7.2\text{Hz}$, 3H), 1.38 (sex, $J=7.6\text{Hz}$, 2H), 1.66 (quin, $J=7.6\text{Hz}$, 2H), 2.72 (d, $J=5.2\text{Hz}$, 2H), 7.34-7.46 (m, 3H), 7.50 (d, $J=7.6\text{Hz}$, 1H), 7.63 (d, $J=7.6\text{Hz}$, 1H), 7.90 (s, 2H), 8.29 (brs, 2H), 8.93 (s, 1H), 8.96 (s, 1H), 9.03 (s, 1H), 9.19 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{25}\text{FN}_6\text{O}$ m/z 505.2 (M+H).

**43**


[0444] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopropanecarboxamide **43**

[0445] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.43 (d, $J=7.2\text{Hz}$, 2H), 1.50 (brs, 2H), 1.59 (t, $J=7.2\text{Hz}$, 1H), 7.78 (t, 2H), 8.03 (d, $J=7.2\text{Hz}$, 1H), 8.11 (d, $J=7.6\text{Hz}$, 1H), 8.14 (brs, 2H), 8.35 (d, $J=7.6\text{Hz}$, 1H), 8.40 (s, 2H), 9.24 (s, 1H), 9.42 (brs, 2H), 9.69 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{21}\text{FN}_6\text{O}$ m/z 489.2 (M+H).

**44**

[0446] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclobutanecarboxamide **44**


[0447] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.92-2.02 (m, 1H), 2.07-2.19 (m, 1H), 2.28-2.39 (m, 2H), 2.39-2.51 (m, 2H), 3.42-3.53 (m, 1H), 7.40 (t, $J=8.4\text{Hz}$, 2H), 7.67 (d, $J=7.6\text{Hz}$, 1H), 7.77 (t, $J=8\text{Hz}$, 1H), 7.85 (Abq, $J=8\text{Hz}$, $J=4.8\text{Hz}$, 2H), 8.01 (d, $J=8\text{Hz}$, 1H), 8.03 (s, 2H), 8.87 (s, 1H), 9.10 (s, 2H), 9.38 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{23}\text{FN}_6\text{O}$ m/z 503.2 (M+H).



45

[0448] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopentanecarboxamide **45**

[0449] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.64-1.76 (m, 2H), 1.76-1.87 (m, 2H), 1.87-1.97 (m, 2H), 1.97-2.11 (m, 2H), 3.00 (quin, $J=8\text{Hz}$, 1H), 7.36 (t, $J=8.4\text{Hz}$, 2H), 7.61 (d, $J=7.6\text{Hz}$, 1H), 7.69 (t, 1H), 7.86 (Abq, $J=8\text{Hz}$, $J=5.2\text{Hz}$, 2H), 7.91 (d, $J=8\text{Hz}$, 1H), 7.97 (s, 2H), 8.80 (s, 1H), 9.00 (s, 2H), 9.26 (s, 1H); ESIMS found for $\text{C}_{31}\text{H}_{25}\text{FN}_6\text{O}$ m/z 517.2 (M+H).




46

[0450] N-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclohexanecarboxamide **46**


[0451] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.44 (q, $J=12.8\text{Hz}$, 2H), 1.62 (q, $J=14.8\text{Hz}$, 2H), 1.90 (d, $J=12.8\text{Hz}$, 2H), 2.01 (d, $J=16\text{Hz}$, 2H), 2.50-2.68 (m, 1H), 7.39 (t,

$J=8.8\text{Hz}$, 2H), 7.62 (d, 1H), 7.67 (t, 1H), 7.90 (t, 3H), 7.98 (s, 2H), 8.80 (s, 1H), 8.97 (s, 1H), 8.99 (s, 1H), 9.21 (s, 1H); ESIMS found for $\text{C}_{32}\text{H}_{27}\text{FN}_6\text{O}$ m/z 531.2 ($\text{M}+\text{H}$).

**47**


[0452] N-benzyl-1-(5-(3-(4-(4-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methanamine **47**

[0453] ^1H NMR (CD_3OD , 400 MHz) δ ppm 4.47 (s, 2H), 4.64 (s, 2H), 7.38 (t, $J=8.8\text{Hz}$, 2H), 7.48-7.53 (m, 3H), 7.62-7.69 (m, 3H), 7.76 (t, $J=7.6\text{Hz}$, 1H), 7.85 (Abq, $J=8.4\text{Hz}$, $J=5.2\text{Hz}$, 2H), 8.02 (Abq, $J=10.8\text{Hz}$, $J=8.8\text{Hz}$, 2H), 8.09 (d, $J=8.8\text{Hz}$, 1H), 8.96 (brs, 1H), 8.97 (s, 1H), 9.21 (s, 1H), 9.36 (brs, 1H); ESIMS found for $\text{C}_{33}\text{H}_{25}\text{FN}_6$ m/z 525.2 ($\text{M}+\text{H}$).

**54**


[0454] N-((5-(3-(4-(2-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methyl)ethanamine **54**

[0455] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.49 (t, $J=7.2\text{Hz}$, 3H), 3.29-3.38 (m, 2H), 4.62 (s, 2H), 7.36-7.49 (m, 2H), 7.57-7.73 (m, 3H), 7.79 (t, $J=8\text{Hz}$, 1H), 8.02 (d, $J=8.8\text{Hz}$, 1H), 8.14 (d, $J=7.6\text{Hz}$, 2H), 9.05 (s, 1H), 9.08 (s, 1H), 9.41 (s, 1H), 9.45 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{23}\text{FN}_6$ m/z 463.2 ($\text{M}+\text{H}$).

**57**


[0456] N-(5-(3-(4-(2-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)isobutyramide **57**

[0457] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.30 (d, $J=6.8\text{Hz}$, 6H), 2.84 (sep, $J=7.2\text{Hz}$, 1H), 7.36-7.49 (m, 2H), 7.61 (q, $J=7.2\text{Hz}$, 1H), 6.65 (d, 7.6Hz, 1H), 7.71 (t, $J=7.6\text{Hz}$, 1H), 7.77 (t, $J=7.6\text{Hz}$, 1H), 8.00 (s, 2H), 8.05 (d, $J=8\text{Hz}$, 1H), 8.85 (s, 1H), 9.02 (s, 1H), 9.08 (brs, 1H), 9.36 (brs, 1H); ESIMS found for $\text{C}_{29}\text{H}_{23}\text{FN}_6\text{O}$ m/z 491.2 ($\text{M}+\text{H}$).

**59**

[0458] N-(5-(3-(4-(2-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)benzamide **59**


[0459] ^1H NMR (CD_3OD , 400 MHz) δ ppm 7.36-7.47 (m, 2H), 7.56-7.65 (m, 3H), 7.65-7.76 (m, 3H), 7.78 (t, 1H), 8.00-8.11 (m, 2H), 8.13 (d, $J=7.6\text{Hz}$, 3H), 8.91 (s, 1H), 9.18 (s, 1H), 9.34 (s, 1H), 9.53 (s, 1H); ESIMS found for $\text{C}_{32}\text{H}_{21}\text{FN}_6\text{O}$ m/z 525.2 ($\text{M}+\text{H}$).



60

[0460] 5-(3-(4-(2-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)-N-isopropylpyridin-3-amine **60**


[0461] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.23 (d, $J=5.6\text{Hz}$, 6H), 3.78 (sep, $J=5.6\text{Hz}$, 1H), 7.22-7.35 (m, 2H), 7.49 (q, $J=6.8\text{Hz}$, 1H), 7.53 (d, $J=7.2\text{Hz}$, 1H), 7.59 (t, $J=7.6\text{Hz}$, 1H), 7.67 (t, $J=8\text{Hz}$, 1H), 7.83 (s, 2H), 7.93 (d, $J=8\text{Hz}$, 3H), 8.30 (s, 1H), 8.65 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{23}\text{FN}_6$ m/z 463.1 ($\text{M}+\text{H}$).



63

[0462] 3-(4-(2-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-5-(5-(piperidin-1-ylmethyl)pyridin-3-yl)-1H-indazole **63**

[0463] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.61 (brs, 1H), 1.83-2.07 (m, 5H), 2.17 (s, 1H), 2.69 (s, 1H), 3.13-3.29 (m, 2H), 3.65 (d, $J=9.6\text{Hz}$, 2H), 7.42 (quin, $J=8.8\text{Hz}$, 2H), 7.61 (q, $J=6.8\text{Hz}$, 1H), 7.69 (q, $J=8.4\text{Hz}$, 2H), 7.79 (t, $J=8\text{Hz}$, 1H), 8.01 (d, $J=8.4\text{Hz}$, 1H), 8.16 (t, $J=7.6\text{Hz}$, 2H), 9.10 (s, 1H), 9.48 (s, 1H); ESIMS found for $\text{C}_{31}\text{H}_{27}\text{FN}_6$ m/z 503.2 ($\text{M}+\text{H}$).




66

[0464] N-(5-(3-(4-(2-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **66**


[0465] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.02 (t, $J=7.6\text{Hz}$, 3H), 1.48 (sex, $J=7.2\text{Hz}$, 2H), 1.77 (quin, $J=7.6\text{Hz}$, 2H), 2.51 (t, $J=8\text{Hz}$, 2H), 7.37 (quin, $J=8.4\text{Hz}$, 3H),

7.46 (sex, $J=7.6\text{Hz}$, 3H), 7.77-7.89 (m, 4H), 8.53 (s, 1H), 8.74 (s, 1H), 8.79 (s, 1H), 8.85 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{25}\text{FN}_6\text{O}$ m/z 505.2 (M+H).

**69**


[0466] N-(5-(3-(4-(2-Fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopentanecarboxamide **69**

[0467] ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 1.57-1.69 (m, 2H), 1.66-1.76 (m, 2H), 1.73-1.86 (m, 2H), 1.90-2.02 (m, 2H), 2.95 (quin, $J=8\text{Hz}$, 1H), 7.34-7.55 (m, 5H), 7.76 (brd, $J=7.2\text{Hz}$, 1H), 7.82-8.00 (m, 3H), 8.78 (s, 1H), 8.84 (s, 1H), 8.92 (s, 1H), 9.09 (s, 1H); ESIMS found for $\text{C}_{31}\text{H}_{25}\text{FN}_6\text{O}$ m/z 517.2 (M+H).

**72**


[0468] 1-Cyclopentyl-N-((5-(3-(4-(2-fluorophenyl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methyl)methanamine **72**

[0469] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.16-1.36 (m, 4H), 1.51-1.70 (m, 4H), 1.88 (brs, 2H), 2.80 (s, 1H), 3.10-3.17 (m, 2H), 7.23-7.36 (m, 2H), 7.45-7.55 (m, 1H), 7.54-7.64 (m, 2H), 7.64-7.73 (m, 1H), 7.92 (d, $J=8\text{Hz}$, 1H), 8.04 (d, $J=7.6\text{Hz}$, 2H), 9.00 (brs, 2H), 9.37 (brs, 1H), 9.43 (brs, 1H); ESIMS found for $\text{C}_{32}\text{H}_{29}\text{FN}_6$ m/z 517.2 (M+H).

**75**


[0470] 5-(3-(4-(Pyridin-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-amine **75**

[0471] ^1H NMR (CD_3OD , 400 MHz) δ ppm 7.63 (t, $J=8\text{Hz}$, 1H), 7.77 (d, $J=7.6\text{Hz}$, 1H), 7.88 (s, 2H), 7.93 (d, $J=8\text{Hz}$, 1H), 8.05 (d, $J=2.4\text{Hz}$, 1H), 8.11 (s, 1H), 8.30 (dd, $J=8\text{Hz}$, 1H), 8.45 (s, 1H), 8.88 (s, 1H), 8.94 (d, $J=5.6\text{Hz}$, 1H), 9.36 (d, $J=8\text{Hz}$, 1H), 9.73 (s, 1H); ESIMS found for $\text{C}_{24}\text{H}_{17}\text{N}_7$ m/z 404.1 ($\text{M}+\text{H}$).

**77**


[0472] 5-(4-Methylpyridin-3-yl)-3-(4-(pyridin-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazole **77**

[0473] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.68 (s, 3H), 7.71 (d, $J=8.8\text{Hz}$, 1H), 7.81 (Abq, $J=6.8\text{Hz}$, $J=4\text{Hz}$, 2H), 7.97 (d, $J=8.8\text{Hz}$, 1H), 8.11-8.31 (m, 2H), 8.32 (dd, 1H), 8.69 (s, 1H), 8.78 (d, $J=6.4\text{Hz}$, 1H), 8.92 (s, 1H), 9.01 (d, $J=6\text{Hz}$, 1H), 9.11 (d, 1H), 9.50 (s, 1H); ESIMS found for $\text{C}_{25}\text{H}_{18}\text{N}_6$ m/z 403.2 ($\text{M}+\text{H}$).

**79**

[0474] N,N-Dimethyl-5-(3-(4-(pyridin-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-amine **79**


[0475] ^1H NMR (CD_3OD , 400 MHz) δ ppm 3.10 (s, 6H), 7.42 (t, $J=8\text{Hz}$, 1H), 7.49 (d, $J=7.2\text{Hz}$, 1H), 7.52 (s, 1H), 7.61 (dd, $J=8\text{Hz}$, $J=3.2\text{Hz}$, 1H), 7.67 (d, 1H), 7.73 (d, $J=8.4\text{Hz}$, 1H), 7.79 (d, 1H), 8.07 (d, 1H), 8.23 (s, 1H), 8.30 (s, 1H), 8.57 (d, $J=3.6\text{Hz}$, 1H), 8.61 (brs, 1H), 8.82 (s, 1H), 9.25 (brs, 1H); ESIMS found for $\text{C}_{26}\text{H}_{21}\text{N}_7$ m/z 432.2 (M+H).



84

[0476] N-Isopropyl-5-(3-(4-(pyridin-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-amine **84**

[0477] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.34 (d, $J=6.4\text{Hz}$, 6H), 3.88 (sep, $J=6.4\text{Hz}$, 1H), 7.68 (t, $J=7.6\text{Hz}$, 1H), 7.79 (d, $J=7.6\text{Hz}$, 1H), 7.91 (s, 2H), 7.99 (d, $J=8.4\text{Hz}$, 1H), 8.02-8.07 (m, 2H), 8.32 (dd, $J=8\text{Hz}$, $J=5.6\text{Hz}$, 1H), 8.46 (s, 1H), 8.91 (s, 1H), 8.98 (d, $J=5.6\text{Hz}$, 1H), 9.28 (d, $J=8.4\text{Hz}$, 1H), 9.70 (s, 1H); ESIMS found for $\text{C}_{27}\text{H}_{23}\text{N}_7$ m/z 446.2 (M+H).




88

[0478] 3,3-Dimethyl-N-(5-(3-(4-(pyridin-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)butanamide **88**


[0479] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.19 (s, 9H), 2.36 (s, 2H), 7.41 (t, $J=7.6\text{Hz}$, 1H), 7.50 (d, $J=7.2\text{Hz}$, 1H), 7.65 (quin, $J=5.2\text{Hz}$, 2H), 7.76 (q, $J=6.8\text{Hz}$, 2H),

8.22 (brs, 2H), 8.41 (s, 1H), 8.55 (d, $J=4.4\text{Hz}$, 1H), 8.65 (s, 1H), 8.67 (brs, 1H), 8.82 (s, 1H), 8.85 (s, 1H), 9.31 (brs, 1H); ESIMS found for $\text{C}_{30}\text{H}_{27}\text{N}_7\text{O}$ m/z 502.2 ($\text{M}+\text{H}$).

**89**


[0480] N-(5-(3-(4-(Pyridin-3-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)butyramide **89**

[0481] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.09 (t, $J=7.6\text{Hz}$, 3H), 1.84 (sex, $J=7.2\text{Hz}$, 2H), 2.59 (t, $J=7.6\text{Hz}$, 2H), 7.64 (t, $J=8\text{Hz}$, 1H), 7.81 (d, $J=7.2\text{Hz}$, 1H), 7.88-7.99 (m, 3H), 8.33 (dd, $J=8\text{Hz}$, $J=5.6\text{Hz}$, 1H), 8.94 (d, $J=5.6\text{Hz}$, 1H), 8.98 (s, 1H), 9.08 (t, $J=1.8\text{Hz}$, 2H), 9.29 (d, $J=1.6\text{Hz}$, 1H), 9.36 (d, $J=8.4\text{Hz}$, 1H), 9.90 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{23}\text{N}_7\text{O}$ m/z 474.1 ($\text{M}+\text{H}$).

**90**


[0482] N-(5-(3-(4-(Pyridin-3-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **90**

[0483] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.02 (t, $J=7.6\text{Hz}$, 3H), 1.48 (sex, $J=7.6\text{Hz}$, 2H), 1.77 (quin, $J=7.2\text{Hz}$, 2H), 2.61 (t, $J=7.6\text{Hz}$, 2H), 7.83 (d, $J=6\text{Hz}$, 2H), 8.00 (s, 2H), 8.14 (d, $J=6\text{Hz}$, 1H), 8.38 (t, $J=7.2\text{Hz}$, 1H), 9.05 (t, $J=7.6\text{Hz}$, 3H), 9.14 (d, $J=7.6\text{Hz}$, 1H), 9.18 (s, 1H), 9.43 (s, 1H), 9.57 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{25}\text{N}_7\text{O}$ m/z 488.1 ($\text{M}+\text{H}$).

**92**


[0484] N-(5-(3-(4-(Pyridin-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclobutanecarboxamide **92**

[0485] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.86-1.98 (m, 1H), 2.06 (sex, $J=9.6\text{Hz}$, 1H), 2.19-2.31 (m, 2H), 2.38 (quin, $J=9.2\text{Hz}$, 2H), 3.34 (quin, $J=8.4\text{Hz}$, 1H), 7.31 (t, $J=8\text{Hz}$, 1H), 7.41 (d, $J=7.2\text{Hz}$, 1H), 7.52-7.60 (m, 1H), 7.60 (d, $J=8.4\text{Hz}$, 1H), 7.66 (d, $J=8.4\text{Hz}$, 1H), 8.12 (s, 1H), 8.35 (s, 1H), 8.47 (d, $J=4.8\text{Hz}$, 1H), 8.53 (s, 1H), 8.61 (d, $J=7.6\text{Hz}$, 1H), 8.69 (s, 1H), 8.76 (s, 1H), 9.25 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{23}\text{N}_7\text{O}$ m/z 486.2 ($\text{M}+\text{H}$).

**94**


[0486] N-(5-(3-(4-(Pyridin-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclohexanecarboxamide **94**

[0487] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.45 (q, $J=12.8\text{Hz}$, 2H), 1.61 (q, $J=13.2\text{Hz}$, 2H), 1.79 (d, $J=12.8\text{Hz}$, 1H), 1.91 (d, $J=12.8\text{Hz}$, 2H), 2.03 (d, $J=11.6\text{Hz}$, 2H), 2.60 (t, $J=11.2\text{Hz}$, 1H), 2.77 (d, $J=4.4\text{Hz}$, 1H), 7.71 (t, $J=7.6\text{Hz}$, 1H), 7.83 (d, $J=7.6\text{Hz}$, 1H), 7.97 (s, 2H), 8.01 (d, $J=8\text{Hz}$, 1H), 8.35 (t, $J=6.4\text{Hz}$, 1H), 8.97 (d, $J=5.2\text{Hz}$, 1H), 9.01 (s, 1H), 9.08 (s, 1H), 9.11 (s, 1H), 9.29 (d, $J=8\text{Hz}$, 1H), 9.36 (s, 1H), 9.78 (s, 1H); ESIMS found for $\text{C}_{31}\text{H}_{27}\text{N}_7\text{O}$ m/z 514.3 ($\text{M}+\text{H}$).

**95**


[0488] N-Benzyl-1-(5-(3-(4-(pyridin-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methanamine **95**

[0489] ^1H NMR (CD_3OD , 400 MHz) δ ppm 4.39 (s, 2H), 4.62 (s, 2H), 7.34-7.42 (m, 3H), 7.58 (t, $J=4.4\text{Hz}$, 2H), 7.62 (t, $J=7.6\text{Hz}$, 1H), 7.71 (d, $J=7.6\text{Hz}$, 1H), 7.85 (d, $J=8.8\text{Hz}$, 1H), 7.96 (t, $J=7.2\text{Hz}$, 2H), 8.21 (t, 1H), 8.85 (d, $J=4.4\text{Hz}$, 1H), 8.93 (brs, 1H), 9.04 (s, 1H), 9.11 (s, 1H), 9.36 (s, 1H), 9.37 (brs, 1H), 9.65 (s, 1H); ESIMS found for $\text{C}_{32}\text{H}_{25}\text{N}_7$ m/z 508.2 ($\text{M}+\text{H}$).

**104**


[0490] N-(5-(3-(4-(Pyridin-4-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pivalamide **104**

[0491] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.42 (s, 9H), 7.72 (t, $J=8\text{Hz}$, 1H), 7.92-8.00 (m, 3H), 8.07 (d, $J=8\text{Hz}$, 1H), 8.95 (d, $J=5.6\text{Hz}$, 2H), 9.02 (s, 1H), 9.06 (d, $J=6\text{Hz}$, 2H), 9.12 (s, 1H), 9.28 (s, 1H), 9.41 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{25}\text{N}_7\text{O}$ m/z 488.1 ($\text{M}+\text{H}$).

**106**


[0492] 2-Phenyl-N-(5-(3-(4-(pyridin-4-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)acetamide **106**

[0493] ^1H NMR (CD_3OD , 400 MHz) δ ppm 3.92 (s, 2H), 7.25-7.45 (m, 6H), 7.59 (t, 1H), 7.87-7.98 (m, 4H), 8.92 (d, $J=6.4\text{Hz}$, 2H), 8.98 (s, 1H), 9.03 (s, 1H), 9.12 (s, 2H), 9.15 (d, $J=5.2\text{Hz}$, 1H); ESIMS found for $\text{C}_{32}\text{H}_{23}\text{N}_7\text{O}$ m/z 522.1 ($\text{M}+\text{H}$).

**108**


[0494] N-Isopropyl-5-(3-(4-(pyridin-4-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-amine **108**

[0495] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.23 (d, $J=6\text{Hz}$, 6H), 3.78 (sep, $J=6\text{Hz}$, 1H), 7.68 (t, $J=8\text{Hz}$, 1H), 7.79-7.87 (m, 3H), 7.93 (s, 1H), 7.98 (s, 1H), 8.03 (d, $J=8\text{Hz}$, 1H), 8.36 (s, 1H), 8.67 (d, $J=6.4\text{Hz}$, 2H), 8.83 (s, 1H), 8.96 (d, $J=6\text{Hz}$, 2H); ESIMS found for $\text{C}_{27}\text{H}_{23}\text{N}_7$ m/z 446.3 ($\text{M}+\text{H}$).

**110**


[0496] 3-(4-(Pyridin-4-yl)-1H-benzo[d]imidazol-2-yl)-5-(5-(pyrrolidin-1-ylmethyl)pyridin-3-yl)-1H-indazole **110**

[0497] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.04-2.32 (m, 6H), 3.64 (brs, 2H), 4.67 (s, 2H), 7.56 (t, $J=7.6\text{Hz}$, 1H), 7.82-8.01 (m, 4H), 8.72 (s, 1H), 8.87 (brs, 1H), 8.96 (d, $J=6.8\text{Hz}$, 2H), 9.00 (s, 1H), 9.15 (d, $J=6\text{Hz}$, 2H), 9.27 (brs, 1H); ESIMS found for $\text{C}_{29}\text{H}_{25}\text{N}_7$ m/z 472.1 ($\text{M}+\text{H}$).

**114**


[0498] N-(5-(3-(4-(Pyridin-4-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **114**

[0499] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.00 (t, $J=7.2\text{Hz}$, 3H), 1.47 (sex, $J=7.2\text{Hz}$, 2H), 1.76 (quin, $J=7.6\text{Hz}$, 2H), 2.58 (t, $J=8\text{Hz}$, 2H), 7.58 (t, $J=7.6\text{Hz}$, 1H), 7.86-7.98 (m, 4H), 8.97 (s, 1H), 8.98 (d, $J=6.4\text{Hz}$, 2H), 9.01 (s, 1H), 9.05 (s, 1H), 9.14 (d, $J=6.4\text{Hz}$, 2H), 9.22 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{25}\text{N}_7\text{O}$ m/z 488.1 ($\text{M}+\text{H}$).

**116**


[0500] N-(5-(3-(4-(Pyridin-4-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclobutanecarboxamide **116**

[0501] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.92-2.04 (m, 1H), 2.12 (sex, 1H), 2.27-2.37 (m, 2H), 2.43 (quin, 2H), 3.50 (quin, 1H), 7.81 (t, 1H), 7.94 (d, 1H), 7.99 (s, 2H), 8.16 (d, 1H), 8.75 (d, 2H), 9.01 (s, 1H), 9.06 (s, 1H), 9.10 (d, 2H), 9.12 (s, 1H), 9.41 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{23}\text{N}_7\text{O}$ m/z 486.2 ($\text{M}+\text{H}$).

**117**


[0502] N-(5-(3-(4-(Pyridin-4-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopentanecarboxamide **117**

[0503] ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 1.53-1.64 (m, 2H), 1.64-1.73 (m, 2H), 1.73-1.84 (m, 2H), 1.86-2.01 (m, 2H), 2.91 (quin, $J=7.6\text{Hz}$, 1H), 7.47 (t, $J=7.6\text{Hz}$, 1H), 7.75 (d, $J=8\text{Hz}$, 1H), 7.85 (s, 3H), 8.65 (s, 1H), 8.75 (s, 1H), 8.79 (s, 1H), 8.83 (s, 4H), 8.90 (s, 1H), 10.38 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{25}\text{N}_7\text{O}$ m/z 500.1 ($\text{M}+\text{H}$).

**118**

[0504] N-(5-(3-(4-(Pyridin-4-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclohexanecarboxamide **118**


[0505] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.25-1.50 (m, 3H), 1.58 (q, $J=14.4\text{Hz}$, 2H), 1.76 (d, $J=12.4\text{Hz}$, 1H), 1.88 (d, $J=12.8\text{Hz}$, 2H), 1.98 (d, $J=12\text{Hz}$, 2H), 2.50 (quin, 1H), 7.53 (t, $J=8\text{Hz}$, 1H), 7.78-7.89 (m, 3H), 7.93 (d, $J=7.6\text{Hz}$, 1H), 8.82 (s, 2H), 8.88 (s, 1H), 8.93 (d, $J=6\text{Hz}$, 3H), 9.16 (d, $J=6\text{Hz}$, 2H); ESIMS found for $\text{C}_{31}\text{H}_{27}\text{N}_7\text{O}$ m/z 514.2 ($\text{M}+\text{H}$).

**122**

[0506] 3-Methyl-N-(5-(3-(4-(pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)butanamide **122**


[0507] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.09 (d, $J=6.8\text{Hz}$, 6H), 2.27 (non, $J=6.8\text{Hz}$, 1H), 2.45 (d, $J=7.6\text{Hz}$, 2H), 7.64 (t, $J=8\text{Hz}$, 1H), 7.89 (q, $J=8.8\text{Hz}$, 2H), 7.98

(d, $J=8\text{Hz}$, 1H), 8.05 (brs, 1H), 8.10 (d, $J=7.6\text{Hz}$, 1H), 8.68-8.79 (m, 2H), 8.96 (s, 1H), 9.08 (s, 1H), 9.16 (s, 2H), 9.33 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{25}\text{N}_7\text{O}$ m/z 488.2 (M+H).

**125**


[0508] 5-(4-Methylpyridin-3-yl)-3-(4-(pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazole **125**

[0509] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.72 (s, 3H), 7.69-7.81 (m, 2H), 7.96 (d, $J=8\text{Hz}$, 1H), 8.06 (d, $J=6.8\text{Hz}$, 1H), 8.08-8.22 (m, 3H), 8.66 (brs, 1H), 8.70-8.80 (m, 3H), 9.01 (s, 1H), 9.05 (brs, 1H); ESIMS found for $\text{C}_{25}\text{H}_{18}\text{N}_6$ m/z 403.1 (M+H).

**127**

[0510] N,N-Dimethyl-5-(3-(4-(pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-amine **127**


[0511] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.68 (s, 3H), 2.92 (s, 3H), 7.68 (t, 1H), 7.90 (d, $J=8.8\text{Hz}$, 1H), 7.99 (d, 1H), 8.00-8.10 (m, 2H), 8.10-8.23 (m, 2H), 8.60 (s, 1H), 8.69 (t, $J=8\text{Hz}$, 1H), 8.78 (d, $J=8.4\text{Hz}$, 1H), 8.99 (s, 1H), 9.06 (d, $J=4.8\text{Hz}$, 1H); ESIMS found for $\text{C}_{26}\text{H}_{21}\text{N}_7$ m/z 432.3 (M+H).



129

[0512] N-(5-(3-(4-(Pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)isobutyramide **129**


[0513] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.32 (d, $J=6.8\text{Hz}$, 6H), 2.85 (sep, $J=6.8\text{Hz}$, 1H), 7.63 (t, $J=8\text{Hz}$, 1H), 7.91 (q, $J=8.8\text{Hz}$, 2H), 7.97 (d, $J=8\text{Hz}$, 1H), 8.04 (brs, 1H), 8.12 (d, $J=7.2\text{Hz}$, 1H), 8.71 (t, $J=7.2\text{Hz}$, 1H), 8.78 (d, $J=8\text{Hz}$, 1H), 8.97 (s, 1H), 9.15 (s, 2H), 9.32 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{23}\text{N}_7\text{O}$ m/z 474.1 ($\text{M}+\text{H}$).



133

[0514] N,N-Dimethyl-1-(5-(3-(4-(pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methanamine **133**

[0515] ^1H NMR (CD_3OD , 400 MHz) δ ppm 3.11 (brs, 6H), 4.87 (brs, 2H), 7.50 (brs, 1H), 7.72 (brs, 1H), 7.84 (brs, 2H), 7.99 (brs, 1H), 8.06 (brs, 1H), 8.52 (brs, 1H), 8.66 (brs, 1H), 8.96 (brs, 1H), 9.10 (brs, 1H), 9.15 (brs, 1H), 9.60 (brs, 1H), 9.64 (brs, 1H); ESIMS found for $\text{C}_{27}\text{H}_{23}\text{N}_7$ m/z 446.1 ($\text{M}+\text{H}$).




134

[0516] 3-(4-(Pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-5-(5-(pyrrolidin-1-ylmethyl)pyridin-3-yl)-1H-indazole **134**


[0517] ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm 1.92 (brs, 2H), 2.07 (brs, 2H), 3.23 (brs, 2H), 3.48 (brs, 2H), 4.74 (brs, 2H), 7.53 (brs, 1H), 7.88 (brs, 2H), 8.02 (brs, 1H), 8.14 (brs, 2H), 8.75 (brs, 1H), 9.04 (brs, 1H), 9.12 (s, 1H), 9.17 (brs, 1H), 9.20 (brs,

1H), 9.41 (s, 1H), 9.60 (s, 1H), 12.01 (brs, 1H); ESIMS found for C₂₉H₂₅N₇ *m/z* 472.2 (M+H).

**138**


[0518] N-(5-(3-(4-(Pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **138**

[0519] ¹H NMR (DMSO-d₆, 400 MHz) δ ppm 0.94 (t, *J*=7.2Hz, 3H), 1.20 (sex, *J*=7.2Hz, 2H), 1.39 (quin, *J*=7.6Hz, 2H), 2.52 (t, *J*=8Hz, 2H), 7.21-7.42 (m, 2H), 7.56 (t, *J*=7.6Hz, 1H), 7.65-7.98 (m, 4H), 8.16 (d, *J*=8Hz, 1H), 8.62 (d, *J*=8.8Hz, 1H), 8.95 (s, 1H), 9.01 (s, 1H), 9.03-9.17 (m, 2H); ESIMS found for C₂₉H₂₅N₇O *m/z* 488.2 (M+H).

**140**


[0520] N-(5-(3-(4-(Pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclobutanecarboxamide **140**

[0521] ¹H NMR (CD₃OD, 400 MHz) δ ppm 1.92-2.05 (m, 1H), 2.05-2.20 (m, 1H), 2.26-2.40 (m, 2H), 2.37-2.51 (m, 2H), 3.46 (quin, *J*=8.4Hz, 1H), 7.55 (t, *J*=7.2Hz, 1H), 7.72 (d, *J*=8Hz, 1H), 7.78 (d, *J*=8.4Hz, 1H), 7.87 (d, *J*=8Hz, 1H), 7.96 (d, *J*=8Hz, 1H), 8.06 (brs, 1H), 8.62 (d, *J*=7.6Hz, 1H), 8.71 (brs, 1H), 8.80 (s, 1H), 8.98 (s, 1H), 9.10 (s, 2H), 9.28 (s, 1H); ESIMS found for C₂₉H₂₃N₇O *m/z* 486.2 (M+H).

**142**


[0522] N-(5-(3-(4-(Pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclohexanecarboxamide **142**

[0523] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.37-1.66 (m, 3H), 1.38 (d, $J=12.4\text{Hz}$, 1H), 1.90 (d, $J=12.8\text{Hz}$, 2H), 2.02 (d, $J=11.2\text{Hz}$, 2H), 2.59 (s, 1H), 7.65 (t, $J=8\text{Hz}$, 1H), 7.95 (s, 2H), 8.00 (d, $J=7.6\text{Hz}$, 2H), 8.21 (d, $J=7.5\text{Hz}$, 1H), 8.70 (t, $J=7.6\text{Hz}$, 1H), 8.87 (d, $J=8.4\text{Hz}$, 1H), 9.02 (s, 1H), 9.11 (s, 1H), 9.13-9.20 (m, 2H), 9.25 (s, 1H); ESIMS found for $\text{C}_{31}\text{H}_{27}\text{N}_7\text{O}$ m/z 514.3 ($\text{M}+\text{H}$).

**144**

[0524] 1-Cyclopentyl-N-((5-(3-(4-(pyridin-2-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methyl)methanamine **144**


[0525] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.44 (brs, 2H), 1.68 (brs, 2H), 1.77 (brs, 2H), 2.01 (brs, 2H), 2.45 (brs, 1H), 3.30 (brs, 2H), 7.58 (brs, 1H), 7.86 (d, $J=6\text{Hz}$, 1H), 7.96 (d, $J=6.8\text{Hz}$, 1H), 8.05 (brs, 3H), 8.69 (brs, 2H), 9.13 (brs, 3H), 9.59 (brs, 1H), 9.61 (brs, 1H); ESIMS found for $\text{C}_{31}\text{H}_{29}\text{N}_7$ m/z 500.2 ($\text{M}+\text{H}$).



146


[0526] 3-Methyl-N-(5-(3-(4-(piperidin-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)butanamide **146**

[0527] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.08 (d, $J=6.8\text{Hz}$, 6H), 1.90 (brs, 2H), 2.16-2.33 (m, 5H), 2.44 (d, $J=7.6\text{Hz}$, 2H), 4.21 (brs, 4H), 7.47 (t, $J=8\text{Hz}$, 1H), 7.56 (d, $J=8\text{Hz}$, 1H), 7.80 (d, $J=8\text{Hz}$, 1H), 7.93 (s, 2H), 9.03 (s, 2H), 9.10 (s, 1H), 9.26 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{31}\text{N}_7\text{O}$ m/z 494.3 ($\text{M}+\text{H}$).

**151**

[0528] N,N-Dimethyl-5-(3-(4-(piperidin-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-amine **151**


[0529] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.80 (brs, 2H), 2.10 (brs, 4H), 3.16 (s, 6H), 4.07 (brs, 4H), 7.22-7.29 (m, 1H), 7.31-7.41 (m, 1H), 7.47 (d, $J=8\text{Hz}$, 1H), 7.68 (d, 1H), 7.78 (d, 1H), 8.01 (s, 1H), 8.10 (s, 1H), 8.38 (s, 1H), 8.87 (s, 1H); ESIMS found for $\text{C}_{26}\text{H}_{27}\text{N}_7$ m/z 438.2 ($\text{M}+\text{H}$).

**155**

[0530] N-(5-(3-(4-(Piperidin-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)benzamide **155**


[0531] ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm 1.79 (brs, 2H), 2.17 (brs, 4H), 4.16 (brs, 4H), 7.46 (t, $J=8\text{Hz}$, 1H), 7.52-7.63 (m, 3H), 7.70 (t, 1H), 7.79 (d, $J=8\text{Hz}$, 1H),

7.92 (d, 1H), 7.98 (d, 1H), 8.12 (d, $J=7.2\text{Hz}$, 2H), 9.11 (s, 2H), 9.37 (s, 1H), 9.49 (s, 1H); ESIMS found for $\text{C}_{31}\text{H}_{27}\text{N}_7\text{O}$ m/z 514.0 (M+H).

**156**


[0532] N-Isopropyl-5-(3-(4-(piperidin-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-amine **156**

[0533] ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 1.35 (d, $J=6.4\text{Hz}$, 6H), 1.94 (brs, 2H), 2.25 (brs, 4H), 3.92 (sep, $J=6.4\text{Hz}$, 1H), 4.18 (brs, 4H), 7.54 (t, $J=7.6\text{Hz}$, 1H), 7.68 (d, $J=7.6\text{Hz}$, 1H), 7.85 (d, $J=8.4\text{Hz}$, 1H), 7.91 (s, 2H), 8.06 (s, 2H), 8.45 (s, 1H), 9.04 (s, 1H); ESIMS found for $\text{C}_{27}\text{H}_{29}\text{N}_7$ m/z 452.3 (M+H).

**158**


[0534] 3-(4-(Piperidin-1-yl)-1H-benzo[d]imidazol-2-yl)-5-(5-(pyrrolidin-1-ylmethyl)pyridin-3-yl)-1H-indazole **158**

[0535] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.97 (brs, 3H), 2.16 (brs, 2H), 2.32 (brs, 6H), 2.70 (brs, 1H), 3.43 (d, $J=5.2\text{Hz}$, 2H), 3.76 (brs, 2H), 4.11 (brs, 4H), 7.58 (d, $J=6.8\text{Hz}$, 1H), 7.74 (d, $J=7.2\text{Hz}$, 1H), 7.88 (d, $J=7.6\text{Hz}$, 1H), 7.96 (s, 1H), 8.14 (d, $J=4.4\text{Hz}$, 1H), 9.19 (brs, 1H), 9.37 (brs, 1H), 9.58 (brs, 1H), 9.62 (brs, 1H); ESIMS found for $\text{C}_{29}\text{H}_{31}\text{N}_7$ m/z 478.2 (M+H).

**162**


[0536] N-(5-(3-(4-(Piperidin-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **162**

[0537] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.02 (t, $J=7.2\text{Hz}$, 3H), 1.48 (sex, $J=7.6\text{Hz}$, 2H), 1.66-1.81 (m, 4H), 1.88-2.00 (m, 2H), 2.48 (t, $J=7.6\text{Hz}$, 2H), 3.51 (brs, 4H), 4.63 (brs, 2H), 6.77 (d, 1H), 7.20 (q, $J=7.6\text{Hz}$, 1H), 7.20-7.28 (m, 1H), 7.79 (dq, $J=7.6\text{Hz}$, $J=1.6\text{Hz}$, 2H), 8.51 (s, 1H), 8.71 (d, $J=1.6\text{Hz}$, 1H), 8.77 (d, $J=2.4\text{Hz}$, 1H), 8.93 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{31}\text{N}_7\text{O}$ m/z 494.2 ($\text{M}+\text{H}$).

**166**


[0538] N-(5-(3-(4-(Piperidin-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclohexanecarboxamide **166**

[0539] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.23 (m, 3H), 1.51 (q, $J=10.8\text{Hz}$, 2H), 1.67 (d, $J=11.6\text{Hz}$, 1H), 1.78 (d, $J=10.8\text{Hz}$, 4H), 1.90 (d, $J=11.6\text{Hz}$, 2H), 2.13 (brs, 4H), 2.46 (t, $J=12\text{Hz}$, 1H), 4.08 (brs, 4H), 7.38 (t, $J=8\text{Hz}$, 1H), 7.51 (d, $J=7.6\text{Hz}$, 1H), 7.71 (d, $J=8\text{Hz}$, 1H), 7.82 (s, 2H), 8.97 (d, $J=2.4\text{Hz}$, 2H), 9.08 (s, 1H), 9.17 (s, 1H); ESIMS found for $\text{C}_{31}\text{H}_{33}\text{N}_7\text{O}$ m/z 520.2 ($\text{M}+\text{H}$).

**179**


[0540] N-(5-(3-(4-(4-Methyl-1H-imidazol-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)benzamide **179**

[0541] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.38 (s, 3H), 7.52 (t, 1H), 7.56-7.66 (m, 3H), 7.71 (d, 1H), 7.77 (d, $J=8\text{Hz}$, 1H), 7.83 (d, $J=8.8\text{Hz}$, 1H), 7.91 (d, 1H), 8.15 (s, 1H), 8.20 (d, $J=7.6\text{Hz}$, 2H), 8.93 (s, 1H), 9.07 (s, 1H), 9.14 (s, 1H), 9.51 (s, 1H), 10.12 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{22}\text{N}_8\text{O}$ m/z 511.2 ($\text{M}+\text{H}$).

**184**


[0542] 3,3-Dimethyl-N-(5-(3-(4-(4-methyl-1H-imidazol-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)butanamide **184**

[0543] ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm 1.07 (s, 9H), 2.34 (s, 2H), 2.45 (s, 3H), 7.49 (t, $J=8\text{Hz}$, 1H), 7.69 (d, $J=7.6\text{Hz}$, 1H), 7.75 (d, $J=8\text{Hz}$, 1H), 7.87 (q, $J=7.6\text{Hz}$, 2H), 8.46 (s, 1H), 8.63 (s, 1H), 8.84 (s, 1H), 8.86 (s, 1H), 8.97 (s, 1H), 9.99 (s, 1H), 10.68 (s, 1H); ESIMS found for $\text{C}_{29}\text{H}_{28}\text{N}_8\text{O}$ m/z 505.2 ($\text{M}+\text{H}$).

**186**


[0544] N-(5-(3-(4-(4-Methyl-1H-imidazol-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **186**

[0545] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.03 (t, $J=7.2\text{Hz}$, 3H), 1.49 (sex, $J=7.6\text{Hz}$, 2H), 1.75 (quin, $J=8\text{Hz}$, 2H), 2.36 (s, 3H), 2.50 (t, $J=7.6\text{Hz}$, 2H), 7.26-7.43 (m, 2H), 7.44-7.57 (1H), 7.64 (d, $J=8.4\text{Hz}$, 1H), 7.70 (d, $J=8.4\text{Hz}$, 1H), 7.89 (brs, 1H), 8.22 (s, 1H), 8.41 (d, $J=10.4\text{Hz}$, 1H), 8.56 (s, 1H), 8.61 (s, 1H), 8.65 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{26}\text{N}_8\text{O}$ m/z 491.2 (M+H).

**189**

[0546] N-(5-(3-(4-(4-Methyl-1H-imidazol-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopentanecarboxamide **189**


[0547] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.66-1.80 (m, 2H), 1.78-1.87 (m, 2H), 1.87-2.00 (m, 2H), 2.01-2.12 (m, 2H), 2.55 (s, 3H), 3.04 (quin, $J=8\text{Hz}$, 1H), 7.53 (t, $J=8\text{Hz}$, 1H), 7.66 (d, $J=7.6\text{Hz}$, 1H), 7.81 (d, $J=8\text{Hz}$, 1H), 7.90 (Abq, $J=13.2\text{Hz}$, $J=8.8\text{Hz}$, 2H), 8.21 (s, 1H), 8.93 (s, 1H), 9.04 (s, 2H), 9.32 (s, 1H), 10.04 (d, $J=1.2\text{Hz}$, 1H); ESIMS found for $\text{C}_{29}\text{H}_{26}\text{N}_8\text{O}$ m/z 503.2 (M+H).

**191**

[0548] N-Benzyl-1-(5-(3-(4-(4-methyl-1H-imidazol-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methanamine **191**


[0549] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.33 (s, 3H), 4.36 (s, 2H), 4.44 (s, 2H), 7.36-7.51 (m, 5H), 7.51-7.61 (m, 2H), 7.79 (d, $J=8.8\text{Hz}$, 1H), 7.85 (d, $J=9.2\text{Hz}$, 1H),

8.36 (s, 3H), 8.68 (d, $J=1.6$ Hz, 1H), 8.92 (s, 1H), 9.02 (d, $J=1.6$ Hz, 1H), 9.11 (brs, 1H); ESIMS found for $C_{31}H_{26}N_8$ m/z 511.2 (M+H).

**196**


[0550] 3-(4-(4-Methylpiperazin-1-yl)-1H-benzo[d]imidazol-2-yl)-5-(pyridin-3-yl)-1H-indazole **196**

[0551] ^1H NMR (CD_3OD , 400 MHz) δ ppm 3.02 (s, 3H), 3.58 (brs, 4H), 3.85 (brs, 4H), 6.78 (brs, 1H), 7.24 (t, $J=8$ Hz, 1H), 7.28 (brs, 1H), 7.62 (dd, $J=7.6$ Hz, $J=4.8$ Hz, 1H), 7.79 (d, $J=8.8$ Hz, 1H), 7.85 (d, $J=8.8$ Hz, 1H), 8.28 (d, $J=7.6$ Hz, 1H), 8.59 (d, $J=4.4$ Hz, 1H), 8.85 (s, 1H), 9.02 (s, 1H); ESIMS found for $C_{24}H_{23}N_7$ m/z 410.2 (M+H).

**197**


[0552] 3-(4-(4-Methylpiperazin-1-yl)-1H-benzo[d]imidazol-2-yl)-5-(4-methylpyridin-3-yl)-1H-indazole **197**

[0553] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.39 (s, 3H), 2.46 (s, 6H), 2.74 (brs, 5H), 7.18 (brs, 2H), 7.46 (d, $J=4.8$ Hz, 1H), 7.52 (d, $J=8.4$ Hz, 1H), 7.75 (d, $J=8.8$ Hz, 1H), 8.44 (d, $J=5.6$ Hz, 1H), 8.52 (s, 2H), 8.57 (s, 1H); ESIMS found for $C_{25}H_{25}N_7$ m/z 424.1 (M+H).

**200**


[0554] N-(5-(3-(4-(4-Methylpiperazin-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pivalamide **200**

[0555] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.36 (s, 9H), 2.42 (s, 3H), 2.82 (s, 4H), 3.67 (brs, 4H), 6.68 (brs, 1H), 7.18 (brs, 2H), 7.22-7.33 (m, 2H), 8.50 (brs, 1H), 8.69 (brs, 1H), 8.77 (brs, 1H), 8.84 (brs, 1H); ESIMS found for $\text{C}_{29}\text{H}_{32}\text{N}_8\text{O}$ m/z 509.2 ($\text{M}+\text{H}$).

**207**


[0556] N-(5-(3-(7-(5-fluoropyridin-3-yl)-3H-imidazo[4,5-c]pyridin-2-yl)-1H-indazol-5-yl)pyridin-3-yl)propionamide **207**

[0557] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.52 (brs, 2H), 1.60-1.72 (m, 4H), 2.48 (s, 3H), 2.55 (brs, 5H), 2.87 (brs, 4H), 3.61-3.80 (m, 5H), 6.71 (brs, 1H), 7.21 (brs, 2H), 7.78 (d, 1H), 7.85 (d, 1H), 8.23 (brs, 1H), 8.54 (brs, 1H), 8.88 (brs, 1H), 8.91 (brs, 1H); ESIMS found for $\text{C}_{30}\text{H}_{34}\text{N}_8\text{O}$ m/z 507.2 ($\text{M}+\text{H}$).

**209**


[0558] N-(5-(3-(4-(4-Methylpiperazin-1-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)butyramide **209**

[0559] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.05 (t, $J=7.6\text{Hz}$, 3H), 1.78 (sex, $J=7.6\text{Hz}$, 2H), 2.45 (t, $J=7.6\text{Hz}$, 2H), 2.84 (s, 3H), 3.35 (brs, 4H), 3.85 (brs, 4H), 6.73 (brs, 1H), 7.17-7.28 (m, 2H), 7.77 (Abq, $J=8.8\text{Hz}$, $J=2.8\text{Hz}$, 2H), 8.54 (brs, 1H), 8.69 (brs, 2H), 8.81 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{30}\text{N}_8\text{O}$ m/z 495.2 ($\text{M}+\text{H}$).

**213**


[0560] N-(5-(3-(4-(4-Methylpiperazin-1-yl)-1H-benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopentanecarboxamide **213**

[0561] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.25 (d, $J=8.8\text{Hz}$, 1H), 1.50-1.62 (m, 2H), 1.62-1.83 (m, 6H), 1.82-1.95 (m, 2H), 2.31 (s, 3H), 2.70 (brs, 4H), 2.77 (quin, 1H), 3.53 (brs, 1H), 6.55 (brs, 1H), 7.05 (brs, 2H), 7.60 (Abq, $J=5.6\text{Hz}$, 2H), 8.35 (s, 1H), 8.53 (s, 1H), 8.65 b(s, 1H), 8.68 (s, 1H); ESIMS found for $\text{C}_{30}\text{H}_{32}\text{N}_8\text{O}$ m/z 521.2 ($\text{M}+\text{H}$).

**218**


[0562] N-(5-(3-(1H-Benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-3-methylbutanamide **218**

[0563] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.07 (d, $J=6.4\text{Hz}$, 6H), 2.26 (sep, $J=6.8\text{Hz}$, 1H), 2.43 (d, $J=7.6\text{Hz}$, 2H), 7.68 (dd, $J=6\text{Hz}$, $J=2.8\text{Hz}$, 2H), 7.95 (dd, $J=6.4\text{Hz}$, $J=3.2\text{Hz}$, 2H), 8.01 (s, 2H), 8.98 (s, 2H), 9.09 (s, 1H), 9.35 (s, 1H); ESIMS found for $\text{C}_{24}\text{H}_{22}\text{N}_6\text{O}$ m/z 411.2 ($\text{M}+\text{H}$).

**221**


[0564] 3-(1H-Benzo[*d*]imidazol-2-yl)-5-(4-methylpyridin-3-yl)-1H-indazole
221

[0565] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.41 (s, 3H), 7.29 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.42 (d, $J=4.8\text{Hz}$, 1H), 7.48 (d, 1H), 7.67 (brd, $J=3.2\text{Hz}$, 2H), 7.75 (d, $J=8.4\text{Hz}$, 1H), 8.42 (d, $J=5.2\text{Hz}$, 1H), 8.50 (s, 2H); ESIMS found for $\text{C}_{20}\text{H}_{15}\text{N}_5$ m/z 326.1 ($\text{M}+\text{H}$).

**225**


[0566] N-(5-(3-(1H-Benzo[*d*]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)
isobutyramide **225**

[0567] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.29 (d, $J=6.8\text{Hz}$, 6H), 2.81 (sep, $J=6.8\text{Hz}$, 1H), 7.69 (dd, $J=6.4\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.96 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 8.03 (d, $J=1\text{Hz}$, 2H), 9.01 (s, 1H), 9.04 (t, $J=2\text{Hz}$, 1H), 9.13 (d, $J=1.6\text{Hz}$, 1H), 9.39 (d, $J=2\text{Hz}$, 1H); ESIMS found for $\text{C}_{23}\text{H}_{20}\text{N}_6\text{O}$ m/z 397.1 ($\text{M}+\text{H}$).

**227**

[0568] N-(5-(3-(1H-Benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)benzamide **227**


[0569] ^1H NMR (CD_3OD , 400 MHz) δ ppm 7.31 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.57 (t, $J=7.6\text{Hz}$, 2H), 7.64 (t, 1H), 7.70 (brs, 2H), 7.78 (d, $J=8.4\text{Hz}$, 1H), 7.85 (d, $J=8.4\text{Hz}$, 1H), 8.04 (d, $J=7.6\text{Hz}$, 2H), 8.59 (s, 1H), 8.79 (d, $J=1.6\text{Hz}$, 1H), 8.84 (s, 1H), 9.01 (d, $J=2\text{Hz}$, 1H); ESIMS found for $\text{C}_{26}\text{H}_{18}\text{N}_6\text{O}$ m/z 431.1 ($\text{M}+\text{H}$).



232

[0570] N-(5-(3-(1H-Benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-3,3-dimethylbutanamide **232**


[0571] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.05 (s, 9H), 2.34 (s, 2H), 7.56 (dd, $J=6\text{Hz}$, $J=2.8\text{Hz}$, 2H), 7.85 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.90 (s, 2H), 8.89 (d, $J=2\text{Hz}$, 1H), 8.93 (s, 1H), 9.04 (s, 1H), 9.31 (s, 1H); ESIMS found for $\text{C}_{25}\text{H}_{24}\text{N}_6\text{O}$ m/z 425.2 ($\text{M}+\text{H}$).



234


[0572] N-(5-(3-(1H-Benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pentanamide **234**

[0573] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.02 (t, $J=7.2\text{Hz}$, 3H), 1.49 (sex, $J=7.2\text{Hz}$, 2H), 1.78 (quin, $J=7.6\text{Hz}$, 2H), 2.59 (t, $J=7.2\text{Hz}$, 2H), 7.70 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.98 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 8.04 (s, 2H), 9.02 (s, 1H), 9.05 (s, 1H), 9.15 (s, 1H), 9.42 (d, $J=2\text{Hz}$, 1H); ESIMS found for $\text{C}_{24}\text{H}_{22}\text{N}_6\text{O}$ m/z 411.1 ($\text{M}+\text{H}$).

**235**


[0574] N-(5-(3-(1H-Benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopropanecarboxamide **235**

[0575] ^1H NMR (CD_3OD , 400 MHz) δ ppm 0.90-0.98 (m, 2H), 1.01-1.07 (m, 2H), 1.87 (quin, 1H), 7.32 (dd, $J=6.4\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.71 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.80 (q, $J=8.8\text{Hz}$, 2H), 8.46 (t, $J=0.4\text{Hz}$, 1H), 8.74 (d, $J=2\text{Hz}$, 1H), 8.82 (s, 1H), 8.83 (d, $J=2.4\text{Hz}$, 1H); ESIMS found for $\text{C}_{23}\text{H}_{18}\text{N}_6\text{O}$ m/z 395.1 ($\text{M}+\text{H}$).

**238**

[0576] N-(5-(3-(1H-Benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclohexanecarboxamide **238**


[0577] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.14-1.40 (m, 3H), 1.48 (q, $J=12\text{Hz}$, 2H), 1.66 (d, $J=11.6\text{Hz}$, 1H), 1.78 (d, $J=12.8\text{Hz}$, 2H), 1.89 (d, $J=12.4\text{Hz}$, 2H), 2.60 (t, $J=11.6\text{Hz}$, 1H), 7.57 (dd, $J=6.4\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.85 (dd, $J=6\text{Hz}$, $J=3.6\text{Hz}$, 2H), 7.90 (s, 2H), 8.91 (s, 2H), 9.01 (s, 1H), 9.26 (s, 1H); ESIMS found for $\text{C}_{26}\text{H}_{24}\text{N}_6\text{O}$ m/z 437.1 ($\text{M}+\text{H}$).



239

[0578] 1-(5-(3-(1H-Benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)-N-benzylmethanamine **239**


[0579] ^1H NMR (CD_3OD , 400 MHz) δ ppm 4.25 (s, 2H), 4.32 (s, 2H), 7.33 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.40-7.58 (m, 5H), 7.71 (dd, $J=6\text{Hz}$, $J=3.2\text{Hz}$, 2H), 7.81 (d, $J=8.8\text{Hz}$, 1H), 7.88 (dd, $J=8.8\text{Hz}$, $J=1.6\text{Hz}$, 1H), 8.40 (s, 1H), 8.65 (d, $J=1.6\text{Hz}$, 1H), 8.87 (s, 1H), 9.06 (d, $J=2\text{Hz}$, 1H); ESIMS found for $\text{C}_{27}\text{H}_{22}\text{N}_6$ m/z 431.2 (M+H).



246

[0580] N-((5-(3-(4-(Thiophen-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methyl)ethanamine **246**


[0581] ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm 1.12 (t, $J=7.2\text{Hz}$, 3H), 2.74 (q, $J=7.2\text{Hz}$, 2H), 4.00 (s, 2H), 7.31 (t, $J=7.6\text{Hz}$, 1H), 7.49 (d, $J=7.6\text{Hz}$, 1H), 7.65 (d, $J=7.2\text{Hz}$, 1H), 7.71 (dd, $J=5.2\text{Hz}$, $J=3.2\text{Hz}$, 1H), 7.84 (d, $J=8.4\text{Hz}$, 1H), 7.92 (d, $J=7.2\text{Hz}$, 1H), 8.23 (s, 1H), 8.27 (s, 1H), 8.62 (d, $J=1.6\text{Hz}$, 1H), 8.77 (d, $J=1.6\text{Hz}$, 1H), 8.96 (d, $J=2\text{Hz}$, 1H), 8.99 (s, 1H); ESIMS found for $\text{C}_{26}\text{H}_{22}\text{N}_6\text{S}$ m/z 451.1 (M+H).



249


[0582] N-(5-(3-(4-(Thiophen-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)isobutyramide **249**

[0583] ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm 1.19 (d, $J=6.8\text{Hz}$, 6H), 2.71 (sep, $J=7.2\text{Hz}$, 1H), 7.31 (t, $J=8\text{Hz}$, 1H), 7.49 (d, $J=7.6\text{Hz}$, 1H), 7.64 (d, $J=7.2\text{Hz}$, 2H), 7.84 (s, 2H), 8.16 (d, $J=4.4\text{Hz}$, 1H), 8.60 (brs, 1H), 8.72 (brs, 2H), 8.79 (brs, 1H), 8.93 (brs, 1H); ESIMS found for $\text{C}_{27}\text{H}_{22}\text{N}_6\text{OS}$ m/z 479.1 (M+H).

**250**


[0584] 2-Phenyl-N-(5-(3-(4-(thiophen-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)acetamide **250**

[0585] ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 3.86 (s, 2H), 7.24-7.46 (m, 8H), 7.52 (d, 1H), 7.63 (d, $J=6\text{Hz}$, 2H), 7.89 (s, 2H), 8.11 (brs, 1H), 8.61 (brs, 1H), 8.94 (brs, 2H); ESIMS found for $\text{C}_{31}\text{H}_{22}\text{N}_6\text{OS}$ m/z 527.1 (M+H).

**253**


[0586] N,N-Dimethyl-1-(5-(3-(4-(thiophen-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)methanamine **253**

[0587] ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm 3.02 (s, 6H), 4.62 (d, $J=2.8\text{Hz}$, 2H), 7.42 (t, $J=7.6\text{Hz}$, 1H), 7.66 (t, $J=7.2\text{Hz}$, 2H), 7.78 (dd, $J=4.8\text{Hz}$, $J=3.2\text{Hz}$, 1H), 7.94 (d, $J=8.8\text{Hz}$, 1H), 8.02 (d, $J=4.8\text{Hz}$, 1H), 8.10 (d, $J=8.8\text{Hz}$, 1H), 8.56 (s, 1H), 9.10 (s, 2H), 9.23 (s, 1H), 9.47 (s, 1H), 11.77 (brs, 1H); ESIMS found for $\text{C}_{26}\text{H}_{22}\text{N}_6\text{S}$ m/z 451.1 (M+H).

**254**


[0588] 5-(5-(Pyrrolidin-1-ylmethyl)pyridin-3-yl)-3-(4-(thiophen-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazole **254**

[0589] ^1H NMR (CD_3OD , 400 MHz) δ ppm 2.17 (brs, 2H), 2.30 (brs, 2H), 3.33 (s, 2H), 3.39 (brs, 2H), 3.60 (brs, 2H), 7.61 (d, $J=4\text{Hz}$, 1H), 7.67-7.79 (m, 3H), 7.97-8.10 (m, 3H), 8.18 (d, $J=8.4\text{Hz}$, 1H), 9.13 (s, 1H), 9.15 (s, 1H), 9.52 (s, 1H), 9.61 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{24}\text{N}_6\text{S}$ m/z 477.1 ($\text{M}+\text{H}$).

**260**


[0590] N-(5-(3-(4-(Thiophen-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclobutanecarboxamide **260**

[0591] ESIMS found for $\text{C}_{28}\text{H}_{22}\text{N}_6\text{OS}$ m/z 491.1 ($\text{M}+\text{H}$).

**267**


[0592] 5-(3-(4-(Furan-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-amine **267**

[0593] ^1H NMR (CD_3OD , 400 MHz) δ ppm 7.03 (s, 1H), 7.57-7.69 (m, 2H), 7.75 (s, 1H), 7.81 (d, 1H), 7.93 (Abq, 2H), 8.05 (d, $J=2\text{Hz}$, 1H), 8.10 (s, 1H), 8.33 (s, 1H), 8.43 (s, 1H), 8.77 (s, 1H); ESIMS found for $\text{C}_{23}\text{H}_{16}\text{N}_6\text{O}$ m/z 393.0 ($\text{M}+\text{H}$).

**272**


[0594] N-(5-(3-(4-(Furan-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)pivalamide **272**

[0595] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.40 (s, 9H), 7.20 (brs, 1H), 7.33 (t, $J=7.6\text{Hz}$, 1H), 7.49 (d, $J=6.8\text{Hz}$, 1H), 7.53 (brs, 1H), 7.66 (s, 1H), Abq (7.80 [d, $J=9.2\text{Hz}$, 1H], 7.86 [dd, $J=8.4\text{Hz}$, $J=1.6\text{Hz}$, 1H]), 8.58 (s, 1H), 8.75 (d, $J=2\text{Hz}$, 2H), 8.84 (d, $J=6.4\text{Hz}$, 1H), 8.96 (s, 1H); ESIMS found for $\text{C}_{28}\text{H}_{24}\text{N}_6\text{O}_2$ m/z 477.2 (M+H).

**276**

[0596] 5-(3-(4-(Furan-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)-N-isopropylpyridin-3-amine **276**


[0597] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.32 (d, $J=6.4\text{Hz}$, 6H), 3.83 (sep, $J=6\text{Hz}$, 1H), 7.20 (brs, 1H), 7.32 (t, $J=7.6\text{Hz}$, 1H), 7.41 (t, $J=2.4\text{Hz}$, 1H), 7.49 (d, $J=6.8\text{Hz}$, 1H), 7.50 (brs, 1H), 7.66 (s, 1H), Abq (7.74 [d, $J=8.4\text{Hz}$, 1H], 7.80 [dd, $J=8.4\text{Hz}$, $J=1.6\text{Hz}$, 1H]), 7.95 (d, $J=2.4\text{Hz}$, 1H), 7.17 (d, $J=2\text{Hz}$, 1H), 8.78 (brs, 1H), 8.88 (brs, 1H); ESIMS found for $\text{C}_{26}\text{H}_{22}\text{N}_6\text{O}$ m/z 435.1 (M+H).

**279**

[0598] 3-(4-(Furan-3-yl)-1H-benzo[d]imidazol-2-yl)-5-(5-(piperidin-1-ylmethyl)pyridin-3-yl)-1H-indazole **279**

[0599] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.85-2.03 (m, 4H), 2.90 (s, 4H), 3.62 (d, $J=10.4$, 2H), 4.64 (s, 2H), 7.01 (s, 1H), 7.63-7.73 (m, 2H), 7.76 (s, 1H), 7.92 (d, $J=5.2\text{Hz}$, 1H), Abq (7.99 [d, $J=9.2\text{Hz}$, 1H], 8.11 [d, $J=8.8\text{Hz}$, 1H]), 8.33 (s, 1H), 8.97 (s,


1H), 9.01 (s, 1H), 9.24 (s, 1H), 9.36 (s, 1H); ESIMS found for $C_{29}H_{26}N_6O$ m/z 475.2 (M+H).



281

[0600] N-(5-(3-(4-(Furan-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)butyramide **281**

[0601] ^1H NMR (CD_3OD , 400 MHz) δ ppm 1.08 (t, $J=7.6$ Hz, 3H), 1.82 (sex, $J=7.2$ Hz, 2H), 2.48 (t, $J=7.6$ Hz, 2H), 7.20 (brs, 1H), 7.33 (t, $J=7.6$ Hz, 1H), 7.66 (s, 1H), 7.82 (q, $J=8.8$ Hz, 2H), 8.57 (s, 1H), 7.73 (s, 1H), 8.80 (s, 1H), 8.94 (s, 1H); ESIMS found for $C_{27}H_{22}N_6O_2$ m/z 463.2 (M+H).



283

[0602] N-(5-(3-(4-(Furan-3-yl)-1H-benzo[d]imidazol-2-yl)-1H-indazol-5-yl)pyridin-3-yl)cyclopropanecarboxamide **283**

[0603] ^1H NMR (CD_3OD , 400 MHz) δ ppm 0.73-0.87 (m, 2H), 0.87-0.99 (m, 2H), 1.76 (quin, $J=4$ Hz, 1H), 7.08 (brs, 1H), 7.20 (t, $J=7.6$ Hz, 1H), 7.37 (d, $J=7.6$ Hz, 1H), 7.54 (s, 1H), 7.68 (q, $J=8.8$ Hz, 2H), 8.46 (s, 1H), 8.59 (s, 1H), 8.65 (s, 1H), 8.81 (s, 1H); ESIMS found for $C_{27}H_{20}N_6O_2$ m/z 461.1 (M+H).

Example 2.

[0604] The above synthesized compounds were screened using the assay procedure for Wnt activity described below.

[0605] Reporter cell lines were generated by stably transducing cells of cancer cell lines (e.g., colon cancer) with a lentiviral construct that include a wnt-responsive promoter driving expression of the firefly luciferase gene.

[0606] Lentiviral constructs were made in which the SP5 promoter, a promoter having eight TCF/LEF binding sites derived from the SP5 promoter, was linked upstream of the firefly luciferase gene. The lentiviral constructs included a hygromycin resistance gene as a selectable marker. The SP5 promoter construct was used to transduce SW480 cells, a colon cancer cell line having a mutated APC gene that generates a truncated APC protein, leading to de-regulated accumulation of β -catenin. A control cell line was generated using another lentiviral construct containing the luciferase gene under the control of the SV40 promoter which does not require β -catenin for activation.

[0607] Cultured SW480 cells bearing a reporter construct were distributed at approximately 10,000 cells per well into 384 well multiwell plates. Compounds were then added to the wells in half-log dilutions using a three micromolar top concentration. A series of control wells for each cell type received only buffer and compound solvent. Twenty-four hours after the addition of compound, reporter activity for luciferases was assayed, for example, by addition of the BrightGlo luminescence reagent (Promega) and the Victor3 plate reader (Perkin Elmer). Readings were normalized to DMSO only treated cells, and normalized activities were then used for the IC₅₀ calculations. Table 2 shows the activity of selected compounds as provided herein.

Table 2.


Compound	Wnt inhibition (μM)	Compound	Wnt inhibition (μM)	Compound	Wnt inhibition (μM)
1	0.002	79	0.045	184	0.115
4	0.004	84	0.013	186	0.006
5	0.034	88	0.135	189	0.013
7	0.008	89	0.004	191	0.138
9	0.25	90	0.011	196	0.028
10	0.069	92	0.008	197	0.249
13	0.011	94	0.121	200	0.076
16	0.031	95	0.163	202	0.248
18	0.03	104	0.065	207	0.801
19	0.005	106	1.93	209	0.061

21	0.05	108	0.016	213	0.195
22	0.18	110	0.205	218	0.037
26	0.179	114	0.067	220	0.165
29	0.1	116	0.023	221	0.317
32	0.035	117	0.054	222	0.235
33	0.034	118	1.46	225	0.028
35	0.161	122	0.011	227	0.099
38	0.153	125	0.054	232	0.065
41	0.015	127	0.032	234	0.123
42	0.033	129	0.021	235	0.025
43	0.043	133	0.127	238	0.147
44	0.14	134	0.219	239	0.372
45	0.11	138	0.018	246	0.014
46	0.225	140	0.008	249	0.013
47	0.45	142	0.155	250	0.159
54	0.046	144	1.18	250	0.018
57	0.006	146	0.058	253	0.007
57	0.116	151	0.041	254	0.002
59	0.332	155	0.289	267	4.15
63	0.01	156	0.066	272	0.159
66	4	158	0.116	276	0.016
69	0.023	162	0.14	279	0.418
72	1.14	166	0.445	281	0.025
75	0.006	173	0.046	283	0.057
77	0.063	179	0.01		

[0608] The term “comprising” as used herein is synonymous with “including,” “containing,” or “characterized by,” and is inclusive or open-ended and does not exclude additional, unrecited elements or method steps.

WHAT IS CLAIMED IS:

1. A compound, or pharmaceutically acceptable salt thereof, of Formula I:



wherein:

R^1 is -heteroaryl R^3R^4 ;

R^2 is selected from the group consisting of H, halide, C_{1-3} alkyl, -CN, -OR⁸, -OH, -(C_{1-3} alkyl)OR⁸, -NR⁹R¹⁰, -(C_{1-3} alkyl)NR⁹R¹⁰, -heteroaryl R^5 , -heterocyclyl R^6 and -aryl R^7 ;

R^3 is 1 substituent attached to the heteroaryl ring and is selected from the group consisting of H, C_{1-3} alkyl, -CF₃, -NR⁹R¹⁰, -NHC(=O)R⁸, -(C_{1-3} alkyl)heterocyclyl R^6 and -(C_{1-3} alkyl)NR⁹R¹⁰;

R^4 is 1 substituent attached to the heteroaryl ring and is selected from the group consisting of H, C_{1-3} alkyl, -CF₃, halide, -CN, -OR⁸, -OH, -(C_{1-3} alkyl)OR⁸, -NR⁹R¹⁰, -(C_{1-3} alkyl)NR⁹R¹⁰ and -OCF₃;

R^5 is 1-3 substituents attached to the heteroaryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, -CF₃, halide, -CN, -OR⁸, -OH, -(C_{1-3} alkyl)OR⁸, -NR⁹R¹⁰, -(C_{1-3} alkyl)NR⁹R¹⁰ and -OCF₃;

R^6 is 1-3 substituents attached to the heterocyclyl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, -CF₃, halide, -CN, -OR⁸, -OH, -(C_{1-3} alkyl)OR⁸, -NR⁹R¹⁰, -(C_{1-3} alkyl)NR⁹R¹⁰ and -OCF₃;

R^7 is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, -CF₃, halide, -CN, -OR⁸, -OH, -(C_{1-3} alkyl)OR⁸, -NR⁹R¹⁰, -(C_{1-3} alkyl)NR⁹R¹⁰ and -OCF₃;

each R^8 is independently selected from the group consisting of C_{1-9} alkyl, –heteroaryl R^{12} , –heterocyclyl R^{13} , –aryl R^{14} , carbocyclyl R^{11} , –(C_{1-3} alkyl)heteroaryl R^{12} , –(C_{1-3} alkyl)heterocyclyl R^{13} , –(C_{1-3} alkyl)aryl R^{14} and –(C_{1-3} alkyl)carbocyclyl R^{11} ;

each R^9 is independently selected from the group consisting of H, C_{1-6} alkyl, –heteroaryl R^{12} , –heterocyclyl R^{13} , –aryl R^{14} , carbocyclyl R^{11} , –(C_{1-3} alkyl)heteroaryl R^{12} , –(C_{1-3} alkyl)heterocyclyl R^{13} , –(C_{1-3} alkyl)aryl R^{14} and –(C_{1-3} alkyl)carbocyclyl R^{11} ;

each R^{10} is independently selected from the group consisting of H and C_{1-6} alkyl;

R^9 and R^{10} are optionally linked to form a five or six membered heterocyclyl ring;


R^{11} is 1-3 substituents attached to the carbocyclyl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, – CF_3 , halide, –CN, –O(R^{10}), –(C_{1-3} alkyl)OR ^{10} , –N(R^{10}) $_2$, –(C_{1-3} alkyl)N(R^{10}) $_2$ and –OCF $_3$;


R^{12} is 1-3 substituents attached to the heteroaryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, – CF_3 , halide, –CN, –O(R^{10}), –(C_{1-3} alkyl)OR ^{10} , –N(R^{10}) $_2$, –(C_{1-3} alkyl)N(R^{10}) $_2$ and –OCF $_3$;


R^{13} is 1-3 substituents attached to the heterocyclyl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, – CF_3 , halide, –CN, –O(R^{10}), –(C_{1-3} alkyl)OR ^{10} , –N(R^{10}) $_2$, –(C_{1-3} alkyl)N(R^{10}) $_2$ and –OCF $_3$;

R^{14} is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, – CF_3 , halide, –CN, –O(R^{10}), –(C_{1-3} alkyl)OR ^{10} , –N(R^{10}) $_2$, –(C_{1-3} alkyl)N(R^{10}) $_2$ and –OCF $_3$;


with the proviso that a compound of Formula I is not a compound selected from the group consisting of:







2. A compound, or pharmaceutically acceptable salt thereof, of Formula **Ia**:



Ia

wherein:

R^1 is -pyridine R^3R^4 ;

R^2 is selected from the group consisting of H, halide, C_{1-3} alkyl, -CN, -OR⁸, -OH, -(C_{1-3} alkyl)OR⁸, -NR⁹R¹⁰, -(C_{1-3} alkyl)NR⁹R¹⁰, -heteroarylR⁵, -heterocyclylR⁶ and -arylR⁷;

R^3 is 1 substituent attached to the heteroaryl ring and is selected from the group consisting of H, C_{1-3} alkyl, -CF₃, -NR⁹R¹⁰, -NHC(=O)R⁸, -(C_{1-3} alkyl)heterocyclylR⁶ and -(C_{1-3} alkyl)NR⁹R¹⁰;

R^4 is H;

R^5 is 1-3 substituents attached to the heteroaryl ring and is each independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-OR^8$, $-OH$, $-(C_{1-3} \text{ alkyl})OR^8$, $-NR^9R^{10}$, $-(C_{1-3} \text{ alkyl})NR^9R^{10}$ and $-OCF_3$;

R^6 is 1-3 substituents attached to the heterocyclyl ring and is each independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-OR^8$, $-OH$, $-(C_{1-3} \text{ alkyl})OR^8$, $-NR^9R^{10}$, $-(C_{1-3} \text{ alkyl})NR^9R^{10}$ and $-OCF_3$;

R^7 is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-OR^8$, $-OH$, $-(C_{1-3} \text{ alkyl})OR^8$, $-NR^9R^{10}$, $-(C_{1-3} \text{ alkyl})NR^9R^{10}$ and $-OCF_3$;

each R^8 is independently selected from the group consisting of C_{1-9} alkyl, $-heteroarylR^{12}$, $-heterocyclylR^{13}$, $-arylR^{14}$, $carbocyclylR^{11}$, $-(C_{1-3} \text{ alkyl})heteroarylR^{12}$, $-(C_{1-3} \text{ alkyl})heterocyclylR^{13}$, $-(C_{1-3} \text{ alkyl})arylR^{14}$ and $-(C_{1-3} \text{ alkyl})carbocyclylR^{11}$;

each R^9 is independently selected from the group consisting of H, C_{1-6} alkyl, $-heteroarylR^{12}$, $-heterocyclylR^{13}$, $-arylR^{14}$, $carbocyclylR^{11}$, $-(C_{1-3} \text{ alkyl})heteroarylR^{12}$, $-(C_{1-3} \text{ alkyl})heterocyclylR^{13}$, $-(C_{1-3} \text{ alkyl})arylR^{14}$ and $-(C_{1-3} \text{ alkyl})carbocyclylR^{11}$;

each R^{10} is independently selected from the group consisting of H and C_{1-6} alkyl;

R^9 and R^{10} are optionally linked to form a five or six membered heterocyclyl ring;


R^{11} is 1-3 substituents attached to the carbocyclyl ring and each independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;

R^{12} is 1-3 substituents attached to the heteroaryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;

R^{13} is 1-3 substituents attached to the heterocyclyl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;

R^{14} is 1-3 substituents attached to the aryl ring and each is independently selected from the group consisting of H, C_{1-3} alkyl, $-CF_3$, halide, $-CN$, $-O(R^{10})$, $-(C_{1-3} \text{ alkyl})OR^{10}$, $-N(R^{10})_2$, $-(C_{1-3} \text{ alkyl})N(R^{10})_2$ and $-OCF_3$;

with the proviso that a compound of Formula **Ia** is not a compound selected from the group consisting of:




3. The compound of claim 1, wherein R^1 is pyridine R^3R^4 .
4. The compound of claim 3, wherein R^1 is pyridin-3-yl R^3R^4 .
5. The compound of claim 4, wherein R^3 and R^4 are both H.
6. The compound of claim 4, wherein R^3 is $-C_{1-2}$ alkyl and R^4 is H.
7. The compound of claim 4, wherein R^3 is $-\text{CH}_2\text{NR}^9\text{R}^{10}$.
8. The compound of claim 7, wherein R^9 is $-C_{1-2}$ alkyl; R^{10} is $-(C_{1-2}$ alkyl); and R^4 is H.
9. The compound of claim 7, wherein R^4 and R^{10} are both H; and R^9 is selected from the group consisting of $-C_{1-2}$ alkyl, $-\text{CH}_2\text{phenyl}$ and $-\text{CH}_2\text{carbocyclyl}$.
10. The compound of claim 7, wherein R^9 and R^{10} are linked to form a five or six membered heterocyclyl ring; and R^4 is H.
11. The compound of claim 4, wherein R^3 is $-\text{NHC}(=\text{O})\text{R}^8$.
12. The compound of claim 11, wherein R^4 is H and R^8 is selected from the group consisting of $-C_{1-5}$ alkyl, $-\text{CH}_2\text{phenyl}$, phenyl and $-\text{carbocyclyl}$.
13. The compound of claim 4, wherein R^3 is $-\text{NR}^9\text{R}^{10}$.
14. The compound of claim 4, wherein R^3 is $-\text{NH}_2$ and R^4 is H.
15. The compound of claim 13, wherein R^9 is $-C_{1-2}$ alkyl; R^{10} is $-C_{1-2}$ alkyl; and R^4 is H.
16. The compound of claim 13, wherein R^4 and R^9 are both H; and R^{10} is $-C_{1-4}$ alkyl.
17. The compound as in any one of claims 4-16, wherein R^2 is H.
18. The compound as in any one of claims 4-16, wherein R^2 is $-\text{heteroarylR}^5$.
19. The compound as in any one of claims 4-16, wherein R^2 is $-\text{pyridineR}^5$.
20. The compound as in any one of claims 4-16, wherein R^2 is $-\text{thiopheneR}^5$.
21. The compound as in any one of claims 4-16, wherein R^2 is $-\text{furanR}^5$.
22. The compound as in any one of claims 4-16, wherein R^2 is $-\text{imidazolylR}^5$; and R^5 is 1 substituent selected from the group consisting of H or $-C_{1-2}$ alkyl.


23. The compound as in any one of claims 4-16, wherein R^2 is $-\text{heterocycl}R^6$.
24. The compound as in any one of claims 4-16, wherein R^2 is $-\text{piperidine}R^6$.
25. The compound as in any one of claims 4-16, wherein R^2 is $-\text{piperazine}R^6$; and R^6 is 1 substituent selected from the group consisting of H or $-\text{C}_{1-2}$ alkyl.
26. The compound as in any one of claims 4-16, wherein R^2 is $-\text{phenyl}R^7$.
27. The compound of claim 26, wherein R^7 is 1-2 fluorine atoms.
28. The compound of claim 1 or 2, wherein R^1 is $\text{pyridin-3-yl}R^3R^4$; R^2 is $-\text{phenyl}R^7$; R^3 is selected from the group consisting of H, $-\text{C}_{1-2}$ alkyl, $-\text{CH}_2\text{NR}^9R^{10}$; $-\text{NR}^9R^{10}$ and $-\text{NHC}(=\text{O})R^8$; R^4 is H; R^7 is 1-2 fluorine atoms; R^8 is selected from the group consisting of $-\text{C}_{1-5}$ alkyl, $-\text{CH}_2\text{phenyl}$, phenyl and $-\text{carbocycl}R^9$; R^9 is selected from the group consisting of $-\text{C}_{1-3}$ alkyl, $-\text{CH}_2\text{phenyl}$ and $-\text{CH}_2\text{carbocycl}R^{10}$; R^{10} is H or $-\text{C}_{1-2}$ alkyl; or R^9 and R^{10} are optionally linked to form a piperidine or pyrrolidine ring.
29. The compound of claim 1 or 2, wherein R^1 is $\text{pyridin-3-yl}R^3R^4$; R^2 is pyridine; R^3 is selected from the group consisting of $-\text{C}_{1-2}$ alkyl, $-\text{CH}_2\text{NR}^9R^{10}$, $-\text{NR}^9R^{10}$ and $-\text{NHC}(=\text{O})R^8$; R^4 is H; R^8 is selected from the group consisting of $-\text{C}_{1-5}$ alkyl, $-\text{CH}_2\text{phenyl}$ and $-\text{carbocycl}R^9$; R^9 is selected from the group consisting of H, $-\text{C}_{1-3}$ alkyl, $-\text{CH}_2\text{phenyl}$ and $-\text{CH}_2\text{carbocycl}R^{10}$; R^{10} is H or $-\text{C}_{1-2}$ alkyl; or R^9 and R^{10} are optionally linked to form a pyrrolidine ring.
30. The compound of claim 1 or 2, wherein R^1 is $\text{pyridin-3-yl}R^3R^4$; R^2 is 1-methylpiperazinyl- or piperidinyl-; R^3 is selected from the group consisting of H, $-\text{C}_{1-2}$ alkyl, $-\text{CH}_2\text{NR}^9R^{10}$, $-\text{NR}^9R^{10}$ and $-\text{NHC}(=\text{O})R^8$; R^4 is H; R^8 is selected from the group consisting of $-\text{C}_{1-5}$ alkyl, $-\text{CH}_2\text{phenyl}$, phenyl and $-\text{carbocycl}R^9$; R^9 is H or $-\text{C}_{1-2}$ alkyl; R^{10} is $-\text{C}_{1-3}$ alkyl; or R^9 and R^{10} are optionally linked to form a piperidine or pyrrolidine ring.
31. The compound of claim 1 or 2, wherein R^1 is $\text{pyridin-3-yl}R^3R^4$; R^2 is selected from the group consisting of furan, thiophene and $-\text{imidazol}R^5$; R^3 is selected from the group consisting of $-\text{CH}_2\text{NR}^9R^{10}$, $-\text{NR}^9R^{10}$ and $-\text{NHC}(=\text{O})R^8$; R^4 is H; R^5 is H or $-\text{C}_{1-2}$ alkyl; R^8 is selected from the group consisting of $-\text{C}_{1-5}$ alkyl, $-\text{CH}_2\text{phenyl}$ and $-\text{carbocycl}R^9$; R^9 is selected from the group consisting of H, $-\text{C}_{1-3}$ alkyl and $-\text{CH}_2\text{phenyl}$; R^{10} is H or $-\text{C}_{1-2}$ alkyl; or R^9 and R^{10} are optionally linked to form a piperidine or pyrrolidine ring.

32. The compound of claim 1 or 2, wherein R^1 is pyridin-3-yl R^3R^4 ; R^2 is H; R^3 is selected from the group consisting of $-C_{1-2}$ alkyl, $-\text{CH}_2\text{NR}^9\text{R}^{10}$ and $-\text{NHC}(=\text{O})\text{R}^8$; R^4 is H; R^8 is selected from the group consisting of $-C_{1-5}$ alkyl, phenyl and $-\text{carbocyclyl}$; R^9 is $-\text{CH}_2\text{phenyl}$, R^{10} is H.


33. The compound of claim 1 or 2, wherein R^2 is selected from the group consisting of:





34. The compound of claim 1 or 2, wherein R^3 is selected from the group consisting of:





35. The compound according to claim 1 or 2, having a structure selected from the group consisting of:




































36. The compound according to claim 1 or 2, having a structure selected from the group consisting of:









37. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to claim 1 or 2, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

38. A method of treating a disorder or disease selected from the group consisting of cancer, pulmonary fibrosis, a bone or cartilage disease, a neurological condition/disorder/disease, osteoarthritis, lung disease, a fibrotic disorder, the method comprising administering to the patient a therapeutically effective amount of a compound according to claim 1 or 2, or a pharmaceutically acceptable salt thereof.

39. A method of treating a disorder or disease selected from the group consisting of: polyposis coli, osteoporosis-pseudoglioma syndrome, familial exudative vitreoretinopathy, retinal angiogenesis, early coronary disease, tetra-amelia syndrome, Müllerian-duct regression and virilization, SERKAL syndrome, diabetes mellitus type 2, Fuhrmann syndrome, Al-Awadi/Raas-Rothschild/Schinzel phocomelia syndrome, odonto-onycho-dermal dysplasia, obesity, split-hand/foot malformation, caudal duplication syndrome, tooth agenesis, Wilms tumor, skeletal dysplasia, focal dermal hypoplasia, autosomal recessive anonychia, neural tube defects, alpha-thalassemia (ATRX) syndrome, fragile X syndrome, ICF syndrome, Angelman syndrome, Prader-Willi syndrome, Beckwith-Wiedemann Syndrome, Norrie disease, and Rett syndrome, the method comprising administering to the patient a therapeutically effective amount of a compound according to claim 1 or 2, or a pharmaceutically acceptable salt thereof.

40. The method of claim 38, wherein the disorder or disease is cancer.

41. The method of claim 38, wherein the disorder or disease is pulmonary fibrosis.

42. The method of claim 38, wherein the disorder or disease is a bone or cartilage disease.

43. The method of claim 38, wherein the disorder or disease is a neurological condition, disorder or disease, wherein the neurological condition/disorder/disease is selected from: frontotemporal dementias, dementia with lewy bodies, prion diseases, Parkinson's disease, Huntington's disease, progressive supranuclear palsy, corticobasal degeneration, multiple system atrophy, amyotrophic lateral sclerosis (ALS), inclusion body myositis, autism, degenerative myopathies, diabetic neuropathy, other metabolic

neuropathies, endocrine neuropathies, orthostatic hypotension, multiple sclerosis and Charcot-Marie-Tooth disease.

44. The method of claim 38, wherein the disorder or disease is osteoarthritis.

45. The method of claim 38, wherein the disorder or disease is lung disease.

46. The method of claim 38, wherein the disorder or disease is a fibrotic disorder, wherein the fibrotic disorder is selected from: skin fibrosis; scleroderma; progressive systemic fibrosis; lung fibrosis; muscle fibrosis; kidney fibrosis; glomerulosclerosis; glomerulonephritis; hypertrophic scar formation; uterine fibrosis; renal fibrosis; cirrhosis of the liver, liver fibrosis; adhesions; chronic obstructive pulmonary disease; fibrosis following myocardial infarction; pulmonary fibrosis; fibrosis and scarring associated with diffuse/interstitial lung disease; central nervous system fibrosis; fibrosis associated with proliferative vitreoretinopathy (PVR); restenosis; endometriosis; ischemic disease, and radiation fibrosis.

47. The method of claim 38, wherein the patient is a human.

48. The method of claim 40, wherein the cancer is chosen from: hepatocellular carcinoma, colon cancer, breast cancer, pancreatic cancer, chronic myeloid leukemia (CML), chronic myelomonocytic leukemia, chronic lymphocytic leukemia (CLL), acute myeloid leukemia, acute lymphocytic leukemia, Hodgkin lymphoma, lymphoma, sarcoma, and ovarian cancer.

49. The method of claim 40, wherein the cancer is chosen from: lung cancer - non-small cell, lung cancer - small cell, multiple myeloma, nasopharyngeal cancer, neuroblastoma, osteosarcoma, penile cancer, pituitary tumors, prostate cancer, retinoblastoma, rhabdomyosarcoma, salivary gland cancer, skin cancer - basal and squamous cell, skin cancer - melanoma, small intestine cancer, stomach cancers, testicular cancer, thymus cancer, thyroid cancer, uterine sarcoma, vaginal cancer, vulvar cancer, laryngeal or hypopharyngeal cancer, kidney cancer, Kaposi sarcoma, gestational trophoblastic disease, gastrointestinal stromal tumor, gastrointestinal carcinoid tumor, gallbladder cancer, eye cancer (melanoma and lymphoma), Ewing tumor, esophagus cancer, endometrial cancer, colorectal cancer, cervical cancer, brain or spinal cord tumor, bone metastasis, bone cancer, bladder cancer, bile duct cancer, anal cancer, and adrenal cortical cancer.

50. The method of claim 38, wherein the compound inhibits one or more proteins in the Wnt pathway.

51. The method of claim 50, wherein the compound inhibits signaling induced by one or more Wnt proteins.

52. The method of claim 51, wherein the Wnt proteins are chosen from: WNT1, WNT2, WNT2B, WNT3, WNT3A, WNT4, WNT5A, WNT5B, WNT6, WNT7A, WNT7B, WNT8A, WNT8B, WNT9A, WNT9B, WNT10A, WNT10B, WNT11, and WNT16.

53. The method of claim 38, wherein the compound inhibits a kinase activity.

54. A method of preventing or reducing angiogenesis in a patient, the method comprising administering to the patient a therapeutically effective amount of a compound according to claim 1 or 2, or a pharmaceutically acceptable salt thereof.