

WO 2010/149745 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 December 2010 (29.12.2010)

(10) International Publication Number
WO 2010/149745 A1

PCT

(51) International Patent Classification:
A61K 39/155 (2006.01) C07K 14/135 (2006.01)

(74) Agents: DALTON, Marcus Jonathan William et al.; GlaxoSmithKline, Global Patents CN925.1, 980 Great West Road, Brentford Middlesex TW8 9GS (GB).

(21) International Application Number:

PCT/EP2010/059008

(22) International Filing Date:

24 June 2010 (24.06.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/219,964 24 June 2009 (24.06.2009) US
61/334,568 13 May 2010 (13.05.2010) US

(71) Applicants (for all designated States except US): GLAXOSMITHKLINE BIOLOGICALS S.A. [BE/BE]; rue de l'Institut 89, B-1330 Rixensart (BE). ID BIOMEDICAL CORPORATION OF QUEBEC [CA/CA]; 525 Cartier Boulevard West, Laval, Quebec H7V 3S8 (CA).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BI, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, IIN, IIR, IIU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(72) Inventors; and

(75) Inventors/Applicants (for US only): BAUDOUX, Guy Jean Marie Fernand Pierre [BE/BE]; GlaxoSmithKline Biologicals s.a., rue de l'Institut 89, B-1330 Rixensart (BE). BLAIS, Normand [CA/CA]; GlaxoSmithKline Biologicals Laval, 525 Cartier Boulevard West, Laval, Quebec H7V 3S8 (CA). CYR, Sonya L [CA/CA]; GlaxoSmithKline Biologicals Laval, 525 Cartier Boulevard West, Laval, Quebec H7V 3S8 (CA). RHEAULT, Patrick [CA/CA]; GlaxoSmithKline Biologicals Laval, 525 Cartier Boulevard West, Laval, Quebec H7V 3S8 (CA). RUELLE, Jean Louis [BE/BE]; GlaxoSmithKline Biologicals s.a., rue de l'Institut 89, B-1330 Rixensart (BE).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: RECOMBINANT RSV ANTIGENS

(57) Abstract: This disclosure provides recombinant respiratory syncytial virus (RSV) antigens and methods for making and using them, including immunogenic compositions (e.g., vaccines) for the treatment and/or prevention of RSV infection.

RECOMBINANT RSV ANTIGENS

CROSS-REFERENCE TO RELATED APPLICATIONS

[001] This application claims benefit of the earlier filing dates of United States Provisional Applications 61/334,568, filed 13 May 2010 and 61/219,964, filed 24 June 2009, the disclosures of which are incorporated herein by reference.

COPYRIGHT NOTIFICATION PURSUANT TO 37 C.F.R. § 1.71(E)

[002] A portion of the disclosure of this patent document contains material which is subject to copyright protection. The copyright owner has no objection to the facsimile reproduction by anyone of the patent document or the patent disclosure, as it appears in the Patent and Trademark Office patent file or records, but otherwise reserves all copyright rights whatsoever.

BACKGROUND

[003] This disclosure concerns the field of immunology. More particularly this disclosure relates to compositions and methods for eliciting an immune response specific for Respiratory Syncytial Virus (RSV).

[004] Human Respiratory Syncytial Virus (RSV) is the most common worldwide cause of lower respiratory tract infections (LRI) in infants less than 6 months of age and premature babies less than or equal to 35 weeks of gestation. The RSV disease spectrum includes a wide array of respiratory symptoms from rhinitis and otitis to pneumonia and bronchiolitis, the latter two diseases being associated with considerable morbidity and mortality. Humans are the only known reservoir for RSV. Spread of the virus from contaminated nasal secretions occurs via large respiratory droplets, so close contact with an infected individual or contaminated surface is required for transmission. RSV can persist for several hours on toys or other objects, which explains the high rate of nosocomial RSV infections, particularly in paediatric wards.

[005] The global annual infection and mortality figures for RSV are estimated to be 64 million and 160,000 respectively. In the U.S. alone RSV is estimated to be responsible for

18,000 to 75,000 hospitalizations and 90 to 1900 deaths annually. In temperate climates, RSV is well documented as a cause of yearly winter epidemics of acute LRI, including bronchiolitis and pneumonia. In the USA, nearly all children have been infected with RSV by two years of age. The incidence rate of RSV-associated LRI in otherwise healthy children was calculated as 37 per 1000 child-year in the first two years of life (45 per 1000 child-year in infants less than 6 months old) and the risk of hospitalization as 6 per 1000 child-years (per 1000 child-years in the first six months of life). Incidence is higher in children with cardio-pulmonary disease and in those born prematurely, who constitute almost half of RSV-related hospital admissions in the USA. Children who experience a more severe LRI caused by RSV later have an increased incidence of childhood asthma. These studies demonstrate widespread need for RSV vaccines, as well as use therof, in industrialized countries, where the costs of caring for patients with severe LRI and their sequelae are substantial. RSV also is increasingly recognized as an important cause of morbidity from influenza-like illness in the elderly.

[006] Various approaches have been attempted in efforts to produce a safe and effective RSV vaccine that produces durable and protective immune responses in healthy and at risk populations. However, none of the candidates evaluated to date have been proven safe and effective as a vaccine for the purpose of preventing RSV infection and/or reducing or preventing RSV disease, including lower respiratory infections (LRIs).

SUMMARY

[007] This disclosure concerns recombinant respiratory syncytial virus (RSV) antigens. More specifically, this disclosure concerns antigens including a recombinant F protein that has been modified to stabilize the trimeric prefusion conformation. The disclosed recombinant antigens exhibit superior immunogenicity, and are particularly favorably employed as components of immunogenic compositions (*e.g.*, vaccines) for protection against RSV infection and/or disease. Also disclosed are nucleic acids that encode the recombinant antigens, immunogenic compositions containing the antigens, and methods for producing and using the antigens.

BRIEF DESCRIPTION OF THE DRAWINGS

[008] FIG. 1A is a schematic illustration highlighting structural features of the RSV F protein. FIG. 1B is a schematic illustration of exemplary RSV Prefusion F (PreF) antigens.

[009] FIG. 2 is a line graph illustrating representative results of asymmetrical field flow fractionation (AFF-MALS) analysis of PreF.

[010] FIG. 3 is a bar graph showing neutralization inhibition of human serum by PreF antigen.

[011] FIGS. 4A and B are bar graphs showing serum IgG titers elicited in mice in response to PreF antigen.

[012] FIGS. 5A and B are bar graphs illustrating titers of neutralizing antibodies specific for RSV elicited by PreF antigen.

[013] FIGS. 6A and B are graphs indicating protection against challenge provided by the RSV PreF antigen in mice.

[014] FIG. 7 is a graph evaluating BAL leukocytes following immunization and challenge.

[015] FIG. 8 is a bar graph illustrating serum IgG elicited following immunization with PreF formulated with dilutions of an oil-in-water emulsion (AS03).

[016] FIG. 9 is a bar graph illustrating neutralizing antibody titres following immunization with PreF formulated with dilutions of an oil-in-water emulsion (AS03).

[017] FIG. 10 is a graph illustrating protection against challenge following immunization with PreF formulated with dilutions of an oil-in-water emulsion (AS03).

DETAILED DESCRIPTION

INTRODUCTION

[018] Development of vaccines to prevent RSV infection has been complicated by the fact that host immune responses appear to play a role in the pathogenesis of the disease. Early studies in the 1960s showed that children vaccinated with a formalin-inactivated RSV vaccine suffered from more severe disease on subsequent exposure to the virus as compared to unvaccinated control subjects. These early trials resulted in the hospitalization of 80% of vaccinees and two deaths. The enhanced severity of disease has been reproduced in animal models and is thought to result from inadequate levels of serum-neutralizing antibodies, lack of local immunity, and excessive induction of a type 2

helper T-cell-like (Th2) immune response with pulmonary eosinophilia and increased production of IL-4 and IL-5 cytokines. In contrast, a successful vaccine that protects against RSV infection induces a Th1 biased immune response, characterized by production of IL-2 and γ -interferon (IFN).

[019] The present disclosure concerns recombinant respiratory syncytial virus (RSV) antigens that solve problems encountered with RSV antigens previously used in vaccines, and improve the immunological as well as manufacturing properties of the antigen. The recombinant RSV antigens disclosed herein involve a Fusion (F) protein analog that include a soluble F protein polypeptide, which has been modified to stabilize the prefusion conformation of the F protein, that is, the conformation of the mature assembled F protein prior to fusion with the host cell membrane. These F protein analogs are designated “PreF” or “PreF antigens”, for purpose of clarity and simplicity. The PreF antigens disclosed herein are predicated on the unforeseen discovery that soluble F protein analogs that have been modified by the incorporation of a heterologous trimerization domain exhibit improved immunogenic characteristics, and are safe and highly protective when administered to a subject *in vivo*.

[020] Details of the structure of the RSV F protein are provided herein with reference to terminology and designations widely accepted in the art, and illustrated schematically in FIG. 1A. A schematic illustration of exemplary PreF antigens is provided in FIG. 1B. It will be understood by those of skill in the art that any RSV F protein can be modified to stabilize the prefusion conformation according to the teachings provided herein. Therefore, to facilitate understanding of the principles guiding production of PreF antigens, individual structural components will be indicated with reference to an exemplary F protein, the polynucleotide and amino acid sequence of which are provided in SEQ ID NOs:1 and 2, respectively. Similarly, where applicable, G protein antigens are described in reference to an exemplary G protein, the polynucleotide and amino acid sequences of which are provided in SEQ ID NOs:3 and 4, respectively.

[021] With reference to the primary amino acid sequence of the F protein polypeptide (FIG. 1A), the following terms are utilized to describe structural features of the PreF antigens.

[022] The term F0 refers to a full-length translated F protein precursor. The F0 polypeptide can be subdivided into an F2 domain and an F1 domain separated by an intervening peptide, designated pep27. During maturation, the F0 polypeptide undergoes proteolytic cleavage at two furin sites situated between F2 and F1 and flanking pep27. For purpose of the ensuing discussion, an F2 domain includes at least a portion, and as much as all, of amino acids 1-109, and a soluble portion of an F1 domain includes at least a portion, and up to all, of amino acids 137-526 of the F protein. As indicated above, these amino acid positions (and all subsequent amino acid positions designated herein) are given in reference to the exemplary F protein precursor polypeptide (F0) of SEQ ID NO:2.

[023] The prefusion F (or “PreF”) antigen is a soluble (that is, not membrane bound) F protein analog that includes at least one modification that stabilizes the prefusion conformation of the F protein, such that the RSV antigen retains at least one immunodominant epitope of the prefusion conformation of the F protein. The soluble F protein polypeptide includes an F2 domain and an F1 domain of the RSV F protein (but does not include a transmembrane domain of the RSV F protein). In exemplary embodiments, the F2 domain includes amino acids 26-105 and the F1 domain includes amino acids 137-516 of an F protein. However, smaller portions can also be used, so long as the three-dimensional conformation of the stabilized PreF antigen is maintained. Similarly, polypeptides that include additional structural components (e.g., fusion polypeptides) can also be used in place of the exemplary F2 and F1 domains, so long as the additional components do not disrupt the three-dimensional conformation, or otherwise adversely impact stability, production or processing, or decrease immunogenicity of the antigen. The F2 and F1 domains are positioned in an N-terminal to C-terminal orientation designed to replicate folding and assembly of the F protein analog into the mature prefusion conformation. To enhance production, the F2 domain can be preceded by a secretory signal peptide, such as a native F protein signal peptide or a heterologous signal peptide chosen to enhance production and secretion in the host cells in which the recombinant PreF antigen is to be expressed.

[024] The PreF antigens are stabilized (in the trimeric prefusion conformation) by introducing one or more modifications, such as the addition, deletion or substitution, or one or more amino acids. One such stabilizing modification is the addition of an amino acid sequence comprising a heterologous stabilizing domain. In exemplary embodiments,

the heterologous stabilizing domain is a protein multimerization domain. One particularly favorable example of such a protein multimerization domain is a coiled-coil domain, such as an isoleucine zipper domain that promotes trimerization of multiple polypeptides having such a domain. An exemplary isoleucine zipper domain is depicted in SEQ ID NO:11. Typically, the heterologous stabilizing domain is positioned C-terminal to the F1 domain.

[025] Optionally, the multimerization domain is connected to the F1 domain via a short amino acid linker sequence, such as the sequence GG. The linker can also be a longer linker (for example, including the sequence GG, such as the amino acid sequence: GGSGGSGGS; SEQ ID NO:14). Numerous conformationally neutral linkers are known in the art that can be used in this context without disrupting the conformation of the PreF antigen.

[026] Another stabilizing modification is the elimination of a furin recognition and cleavage site that is located between the F2 and F1 domains in the native F0 protein. One or both furin recognition sites, located at positions 105-109 and at positions 133-136 can be eliminated by deleting or substituting one or more amino acid of the furin recognition sites, such that the protease is incapable of cleaving the PreF polypeptide into its constituent domains. Optionally, the intervening pep27 peptide can also be removed or substituted, *e.g.*, by a linker peptide. Additionally, or optionally, a non-furin cleavage site (*e.g.*, a metalloproteinase site at positions 112-113) in proximity to the fusion peptide can be removed or substituted.

[027] Another example of a stabilizing mutation is the addition or substitution of a hydrophilic amino acid into a hydrophobic domain of the F protein. Typically, a charged amino acid, such as lysine, will be added or substituted for a neutral residue, such as leucine, in the hydrophobic region. For example, a hydrophilic amino acid can be added to, or substituted for, a hydrophobic or neutral amino acid within the HRB coiled-coil domain of the F protein extracellular domain. By way of example, a charged amino acid residue, such as lysine, can be substituted for the leucine present at position 512 of the F protein. Alternatively, or in addition, a hydrophilic amino acid can be added to, or substituted for, a hydrophobic or neutral amino acid within the HRA domain of the F protein. For example, one or more charged amino acids, such as lysine, can be inserted at or near position 105-106 (*e.g.*, following the amino acid corresponding to residue 105 of

reference SEQ ID NO:2, such as between amino acids 105 and 106) of the PreF antigen). Optionally, hydrophilic amino acids can be added or substituted in both the HRA and HRB domains. Alternatively, one or more hydrophobic residues can be deleted, so long as the overall conformation of the PreF antigen is not adversely impacted.

[028] Additionally or alternatively, one or more modification may be made, which alters the glycosylation state of the PreF antigen. For example, one or more amino acids in a glycosylation site present in a native RSV F protein, *e.g.*, at or around amino acid residue 500 (as compared to SEQ ID NO:2) can be deleted or substituted (or an amino acid can be added such that the glycosylation site is disrupted) to increase or decrease the glycosylation status of the PreF antigen. For example, the amino acids corresponding to positions 500-502 of SEQ ID NO:2 can be selected from: NGS; NKS; NGT; and NKT. Thus, in certain embodiments, the PreF antigens include a soluble F protein polypeptide comprising an F2 domain (*e.g.*, corresponding to amino acids 26-105 of SEQ ID NO:2) and an F1 domain (*e.g.*, corresponding to amino acids 137-516 of SEQ ID NO:2) of an RSV F protein polypeptide, in which at least one modification that alters glycosylation has been introduced. The RSV PreF antigen, typically includes an intact fusion peptide between the F2 domain and the F1 domain. Optionally, the PreF antigen includes a signal peptide.

[029] As disclosed above, such F protein polypeptides can include at least one modification selected from: (i) an addition of an amino acid sequence comprising a heterologous trimerization domain (such as a isoleucine zipper domain); (ii) a deletion of at least one furin cleavage site; (iii) a deletion of at least one non-furin cleavage site; (iv) a deletion of one or more amino acids of the pep27 domain; and (v) at least one substitution or addition of a hydrophilic amino acid in a hydrophobic domain of the F protein extracellular domain. As disclosed above, such glycosylation modified RSV PreF antigens assemble into multimers, *e.g.*, trimers.

[030] In exemplary embodiments, the glycosylation modified PreF antigens are selected from the group of: a) a polypeptide comprising or consisting of SEQ ID NO:22; b) a polypeptide encoded by SEQ ID NO:21 or by a polynucleotide sequence that hybridizes under stringent conditions over substantially its entire length to SEQ ID NO:21; c) a polypeptide with at least 95% sequence identity to SEQ ID NO:22.

[031] Any and/or all of the stabilizing modifications can be used individually and/or in combination with any of the other stabilizing modifications disclosed herein to produce a PreF antigen. In exemplary embodiments the PreF protein comprising a polypeptide comprising an F2 domain and an F1 domain with no intervening furin cleavage site between the F2 domain and the F1 domain, and with a heterologous stabilizing domain (*e.g.*, trimerization domain) positioned C-terminal to the F1 domain. In certain embodiments, the PreF antigen also includes one or more addition and/or substitution of a hydrophilic residue into a hydrophobic HRA and/or HRB domain. Optionally, the PreF antigen has a modification of at least one non-furin cleavage site, such as a metalloproteinase site.

[032] A PreF antigen can optionally include an additional polypeptide component that includes at least an immunogenic portion of the RSV G protein. That is, in certain embodiments, the PreF antigen is a chimeric protein that includes both an F protein and a G protein component. The F protein component can be any of the PreF antigens described above, and the G protein component is selected to be an immunologically active portion of the RSV G protein (up to and/or including a full-length G protein). In exemplary embodiments, the G protein polypeptide includes amino acids 149-229 of a G protein (where the amino acid positions are designated with reference to the G protein sequence represented in SEQ ID NO:4). One of skill in the art will appreciate that a smaller portion or fragment of the G protein can be used, so long as the selected portion retains the dominant immunologic features of the larger G protein fragment. In particular, the selected fragment retains the immunologically dominant epitope between about amino acid positions 184-198 (*e.g.*, amino acids 180-200), and be sufficiently long to fold and assemble into a stable conformation that exhibits the immunodominant epitope. Longer fragments can also be used, *e.g.*, from about amino acid 128 to about amino acid 229, up to the full-length G protein. So long as the selected fragment folds into a stable conformation in the context of the chimeric protein, and does not interfere with production, processing or stability when produced recombinantly in host cells. Optionally, the G protein component is connected to the F protein component via a short amino acid linker sequence, such as the sequence GG. The linker can also be a longer linker (such as the amino acid sequence: GGSGGS: SEQ ID NO:14). Numerous conformationally neutral linkers are known in the art that can be used in this context without disrupting the conformation of the PreF antigen.

[033] Optionally, the G protein component can include one or more amino acid substitutions that reduce or prevent enhanced viral disease in an animal model of RSV disease. That is, the G protein can include an amino acid substitution, such that when an immunogenic composition including the PreF-G chimeric antigen is administered to a subject selected from an accepted animal model (e.g., mouse model of RSV), the subject exhibits reduced or no symptoms of vaccine enhanced viral disease (e.g., eosinophilia, neutrophilia), as compared to a control animal receiving a vaccine including that contains an unmodified G protein. The reduction and/or prevention of vaccine enhanced viral disease can be apparent when the immunogenic compositions are administered in the absence of adjuvant (but not, for example, when the antigens are administered in the presence of a strong Th1 inducing adjuvant). Additionally, the amino acid substitution can reduce or prevent vaccine enhanced viral disease when administered to a human subject. An example of a suitable amino acid substitution is the replacement of asparagine at position 191 by an alanine (Asn → Ala at amino acid 191: N191A).

[034] Optionally, any PreF antigen described above can include an additional sequence that serves as an aid to purification. One example, is a polyhistidine tag. Such a tag can be removed from the final product if desired.

[035] When expressed, the PreF antigens undergo intramolecular folding and assemble into mature protein that includes a multimer of polypeptides. Favorably, the preF antigen polypeptides assemble into a trimer that resembles the prefusion conformation of the mature, processed, RSV F protein.

[036] Any of the PreF antigens (including PreF-G antigens) disclosed herein can be favorably used in immunogenic compositions for the purpose of eliciting a protective immune response against RSV. Such immunogenic compositions typically include a pharmaceutically acceptable carrier and/or excipient, such as a buffer. To enhance the immune response produced following administration, the immunogenic composition typically also includes an adjuvant. In the case of immunogenic compositions for eliciting a protective immune response against RSV (e.g., vaccines), the compositions favorably include an adjuvant that predominantly elicits a Th1 immune response (a Th1 biasing adjuvant). Typically, the adjuvant is selected to be suitable for administration to the target population to which the composition is to be administered. Thus, depending on the

application, the adjuvant is selected to be suitable for administration, *e.g.*, to neonates or to the elderly.

[037] The immunogenic compositions described herein are favorably employed as vaccines for the reduction or prevention of infection with RSV, without inducing a pathological response (such as vaccine enhanced viral disease) following administration or exposure to RSV.

[038] In some embodiments, the immunogenic composition includes a PreF antigen (such as the exemplary embodiment illustrated by SEQ ID NO:6) and a second polypeptide that includes a G protein component. The G protein component typically includes at least amino acids 149-229 of a G protein. Although smaller portions of the G protein can be used, such fragments should include, at a minimum, the immunological dominant epitope of amino acids 184-198. Alternatively, the G protein can include a larger portion of the G protein, such as amino acids 128-229 or 130-230, optionally as an element of a larger protein, such as a full-length G protein, or a chimeric polypeptide.

[039] In other embodiments, the immunogenic composition includes a PreF antigen that is a chimeric protein that also includes a G protein component (such as the exemplary embodiments illustrated by SEQ ID NOs:8 and 10). The G protein component of such a chimeric PreF (or PreF-G) antigen typically includes at least amino acids 149-229 of a G protein. As indicated above, smaller or larger fragments (such as amino acids 129-229 or 130-230) of the G protein can also be used, so long as the immunodominant epitopes are retained, and conformation of the PreF-G antigen is not adversely impacted.

[040] Optionally, the immunogenic compositions can also include at least one additional antigen of a pathogenic organism other than RSV. For example, the pathogenic organism is a virus other than RSV, such as Parainfluenza virus (PIV), measles, hepatitis B, poliovirus, or influenza virus. Alternatively, the pathogenic organism can be a bacterium, such as diphtheria, tetanus, pertussis, *Haemophilus influenzae*, and *Pneumococcus*.

[041] Recombinant nucleic acids that encode any of the PreF antigens (including PreF-G antigens) are also a feature of this disclosure. In some embodiments, the polynucleotide sequence of the nucleic acid that encodes the PreF antigen of the nucleic acid is optimized for expression in a selected host (such as CHO cells, other mammalian cells, or insect cells). Accordingly, vectors, including expression vectors (including prokaryotic and

eukaryotic expression vectors) are a feature of this disclosure. Likewise, host cells including such nucleic acids, and vectors, are a feature of this disclosure. Such nucleic acids can also be used in the context of immunogenic compositions for administration to a subject to elicit an immune response specific for RSV.

[042] The PreF antigens are favorably used for the prevention and/or treatment of RSV infection. Thus, another aspect of this disclosure concerns a method for eliciting an immune response against RSV. The method involves administering an immunologically effective amount of a composition containing a PreF antigen to a subject (such as a human or animal subject). Administration of an immunologically effective amount of the composition elicits an immune response specific for epitopes present on the PreF antigen. Such an immune response can include B cell responses (*e.g.*, the production of neutralizing antibodies) and/or T cell responses (*e.g.*, the production of cytokines). Favorably, the immune response elicited by the PreF antigen includes elements that are specific for at least one conformational epitope present on the prefusion conformation of the RSV F protein. The PreF antigens and compositions can be administered to a subject without enhancing viral disease following contact with RSV. Favorably, the PreF antigens disclosed herein and suitably formulated immunogenic compositions elicit a Th1 biased immune response that reduces or prevents infection with a RSV and/or reduces or prevents a pathological response following infection with a RSV.

[043] The immunogenic compositions can be administered via a variety of routes, including routes, such as intranasal, that directly place the PreF antigen in contact with the mucosa of the upper respiratory tract. Alternatively, more traditional administration routes can be employed, such as an intramuscular route of administration.

[044] Thus, the use of any of the disclosed RSV antigens (or nucleic acids) in the preparation of a medicament for treating RSV infection (for example, prophylactically treating or preventing an RSV infection) is also contemplated. Accordingly, this disclosure provides the disclosed recombinant RSV antigens or the immunogenic compositions for use in medicine, as well as the use thereof for the prevention or treatment of RSV-associated diseases.

[045] Additional details regarding PreF antigens, and methods of using them, are presented in the description and examples below.

TERMS

[046] In order to facilitate review of the various embodiments of this disclosure, the following explanations of terms are provided. Additional terms and explanations can be provided in the context of this disclosure.

[047] Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Definitions of common terms in molecular biology can be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew *et al.* (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

[048] The singular terms “a,” “an,” and “the” include plural referents unless context clearly indicates otherwise. Similarly, the word “or” is intended to include “and” unless the context clearly indicates otherwise. The term “plurality” refers to two or more. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Additionally, numerical limitations given with respect to concentrations or levels of a substance, such as an antigen, are intended to be approximate. Thus, where a concentration is indicated to be at least (for example) 200 pg, it is intended that the concentration be understood to be at least approximately (or “about” or “~”) 200 pg.

[049] Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of this disclosure, suitable methods and materials are described below. The term “comprises” means “includes.” Thus, unless the context requires otherwise, the word “comprises,” and variations such as “comprise” and “comprising” will be understood to imply the inclusion of a stated compound or composition (e.g., nucleic acid, polypeptide, antigen) or step, or group of compounds or steps, but not to the exclusion of any other compounds, composition, steps, or groups thereof. The abbreviation, “e.g.” is derived from the Latin *exempli gratia*, and is used herein to indicate a non-limiting example. Thus, the abbreviation “e.g.” is synonymous with the term “for example.”

[050] Respiratory syncytial virus (RSV) is a pathogenic virus of the family Paramyxoviridae, subfamily Pneumovirinae, genus Pneumovirus. The genome of RSV is a negative-sense RNA molecule, which encodes 11 proteins. Tight association of the RNA genome with the viral N protein forms a nucleocapsid wrapped inside the viral envelope. Two groups of human RSV strains have been described, the A and B groups, based on differences in the antigenicity of the G glycoprotein. Numerous strains of RSV have been isolated to date. Exemplary strains indicated by GenBank and/or EMBL Accession number can be found in WO2008114149, which is incorporated herein by reference for the purpose of disclosing the nucleic acid and polypeptide sequences of RSV F and G proteins suitable for use in PreF antigens (including chimeric PreF-G antigens), and in combinations with PreF antigens. Additional strains of RSV are likely to be isolated, and are encompassed within the genus of RSV. Similarly, the genus of RSV encompasses variants arising from naturally occurring (e.g., previously or subsequently identified strains) by genetic drift, or artificial synthesis and/or recombination.

[051] The term “F protein” or “Fusion protein” or “F protein polypeptide” or “Fusion protein polypeptide” refers to a polypeptide or protein having all or part of an amino acid sequence of an RSV Fusion protein polypeptide. Similarly, the term “G protein” or “G protein polypeptide” refers to a polypeptide or protein having all or part of an amino acid sequence of an RSV Attachment protein polypeptide. Numerous RSV Fusion and Attachment proteins have been described and are known to those of skill in the art. WO2008114149 sets out exemplary F and G protein variants (for example, naturally occurring variants) publicly available as of the filing date of this disclosure.

[052] A “variant” when referring to a nucleic acid or a polypeptide (e.g., an RSV F or G protein nucleic acid or polypeptide, or a PreF nucleic acid or polypeptide) is a nucleic acid or a polypeptide that differs from a reference nucleic acid or polypeptide. Usually, the difference(s) between the variant and the reference nucleic acid or polypeptide constitute a proportionally small number of differences as compared to the referent.

[053] A “domain” of a polypeptide or protein is a structurally defined element within the polypeptide or protein. For example, a “trimerization domain” is an amino acid sequence within a polypeptide that promotes assembly of the polypeptide into trimers. For example, a trimerization domain can promote assembly into trimers via associations with other trimerization domains (of additional polypeptides with the same or a different amino acid

sequence). The term is also used to refer to a polynucleotide that encodes such a peptide or polypeptide.

[054] The terms “native” and “naturally occurring” refer to an element, such as a protein, polypeptide or nucleic acid, that is present in the same state as it is in nature. That is, the element has not been modified artificially. It will be understood, that in the context of this disclosure, there are numerous native/naturally occurring variants of RSV proteins or polypeptides, *e.g.*, obtained from different naturally occurring strains or isolates of RSV.

[055] The term “polypeptide” refers to a polymer in which the monomers are amino acid residues which are joined together through amide bonds. The terms “polypeptide” or “protein” as used herein are intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. The term “polypeptide” is specifically intended to cover naturally occurring proteins, as well as those which are recombinantly or synthetically produced. The term “fragment,” in reference to a polypeptide, refers to a portion (that is, a subsequence) of a polypeptide. The term “immunogenic fragment” refers to all fragments of a polypeptide that retain at least one predominant immunogenic epitope of the full-length reference protein or polypeptide. Orientation within a polypeptide is generally recited in an N-terminal to C-terminal direction, defined by the orientation of the amino and carboxy moieties of individual amino acids. Polypeptides are translated from the N or amino-terminus towards the C or carboxy-terminus.

[056] A “signal peptide” is a short amino acid sequence (*e.g.*, approximately 18-25 amino acids in length) that direct newly synthesized secretory or membrane proteins to and through membranes, *e.g.*, of the endoplasmic reticulum. Signal peptides are frequently but not universally located at the N-terminus of a polypeptide, and are frequently cleaved off by signal peptidases after the protein has crossed the membrane. Signal sequences typically contain three common structural features: an N-terminal polar basic region (n-region), a hydrophobic core, and a hydrophilic c-region).

[057] The terms “polynucleotide” and “nucleic acid sequence” refer to a polymeric form of nucleotides at least 10 bases in length. Nucleotides can be ribonucleotides, deoxyribonucleotides, or modified forms of either nucleotide. The term includes single and double forms of DNA. By “isolated polynucleotide” is meant a polynucleotide that is not immediately contiguous with both of the coding sequences with which it is

immediately contiguous (one on the 5' end and one on the 3' end) in the naturally occurring genome of the organism from which it is derived. In one embodiment, a polynucleotide encodes a polypeptide. The 5' and 3' direction of a nucleic acid is defined by reference to the connectivity of individual nucleotide units, and designated in accordance with the carbon positions of the deoxyribose (or ribose) sugar ring. The informational (coding) content of a polynucleotide sequence is read in a 5' to 3' direction.

[058] A "recombinant" nucleic acid is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination can be accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, *e.g.*, by genetic engineering techniques. A "recombinant" protein is one that is encoded by a heterologous (*e.g.*, recombinant) nucleic acid, which has been introduced into a host cell, such as a bacterial or eukaryotic cell. The nucleic acid can be introduced, on an expression vector having signals capable of expressing the protein encoded by the introduced nucleic acid or the nucleic acid can be integrated into the host cell chromosome.

[059] The term "heterologous" with respect to a nucleic acid, a polypeptide or another cellular component, indicates that the component occurs where it is not normally found in nature and/or that it originates from a different source or species.

[060] The term "purification" (*e.g.*, with respect to a pathogen or a composition containing a pathogen) refers to the process of removing components from a composition, the presence of which is not desired. Purification is a relative term, and does not require that all traces of the undesirable component be removed from the composition. In the context of vaccine production, purification includes such processes as centrifugation, dialization, ion-exchange chromatography, and size-exclusion chromatography, affinity-purification or precipitation. Thus, the term "purified" does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified nucleic acid preparation is one in which the specified protein is more enriched than the nucleic acid is in its generative environment, for instance within a cell or in a biochemical reaction chamber. A preparation of substantially pure nucleic acid or protein can be purified such that the desired nucleic acid represents at least 50% of the total nucleic acid content of the preparation. In certain embodiments, a substantially pure nucleic acid will represent at

least 60%, at least 70%, at least 80%, at least 85%, at least 90%, or at least 95% or more of the total nucleic acid or protein content of the preparation.

[061] An “isolated” biological component (such as a nucleic acid molecule, protein or organelle) has been substantially separated or purified away from other biological components in the cell of the organism in which the component naturally occurs, such as, other chromosomal and extra-chromosomal DNA and RNA, proteins and organelles. Nucleic acids and proteins that have been “isolated” include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids and proteins.

[062] An “antigen” is a compound, composition, or substance that can stimulate the production of antibodies and/or a T cell response in an animal, including compositions that are injected, absorbed or otherwise introduced into an animal. The term “antigen” includes all related antigenic epitopes. The term “epitope” or “antigenic determinant” refers to a site on an antigen to which B and/or T cells respond. The “dominant antigenic epitopes” or “dominant epitope” are those epitopes to which a functionally significant host immune response, *e.g.*, an antibody response or a T-cell response, is made. Thus, with respect to a protective immune response against a pathogen, the dominant antigenic epitopes are those antigenic moieties that when recognized by the host immune system result in protection from disease caused by the pathogen. The term “T-cell epitope” refers to an epitope that when bound to an appropriate MHC molecule is specifically bound by a T cell (via a T cell receptor). A “B-cell epitope” is an epitope that is specifically bound by an antibody (or B cell receptor molecule).

[063] An “adjuvant” is an agent that enhances the production of an immune response in a non-specific manner. Common adjuvants include suspensions of minerals (alum, aluminum hydroxide, aluminum phosphate) onto which antigen is adsorbed; emulsions, including water-in-oil, and oil-in-water (and variants therof, including double emulsions and reversible emulsions), liposaccharides, lipopolysaccharides, immunostimulatory nucleic acids (such as CpG oligonucleotides), liposomes, Toll-like Receptor agonists (particularly, TLR2, TLR4, TLR7/8 and TLR9 agonists), and various combinations of such components.

[064] An “immunogenic composition” is a composition of matter suitable for administration to a human or animal subject (e.g., in an experimental setting) that is capable of eliciting a specific immune response, e.g., against a pathogen, such as RSV. As such, an immunogenic composition includes one or more antigens (for example, polypeptide antigens) or antigenic epitopes. An immunogenic composition can also include one or more additional components capable of eliciting or enhancing an immune response, such as an excipient, carrier, and/or adjuvant. In certain instances, immunogenic compositions are administered to elicit an immune response that protects the subject against symptoms or conditions induced by a pathogen. In some cases, symptoms or disease caused by a pathogen is prevented (or reduced or ameliorated) by inhibiting replication of the pathogen (e.g., RSV) following exposure of the subject to the pathogen. In the context of this disclosure, the term immunogenic composition will be understood to encompass compositions that are intended for administration to a subject or population of subjects for the purpose of eliciting a protective or palliative immune response against RSV (that is, vaccine compositions or vaccines).

[065] An “immune response” is a response of a cell of the immune system, such as a B cell, T cell, or monocyte, to a stimulus. An immune response can be a B cell response, which results in the production of specific antibodies, such as antigen specific neutralizing antibodies. An immune response can also be a T cell response, such as a CD4+ response or a CD8+ response. In some cases, the response is specific for a particular antigen (that is, an “antigen-specific response”). If the antigen is derived from a pathogen, the antigen-specific response is a “pathogen-specific response.” A “protective immune response” is an immune response that inhibits a detrimental function or activity of a pathogen, reduces infection by a pathogen, or decreases symptoms (including death) that result from infection by the pathogen. A protective immune response can be measured, for example, by the inhibition of viral replication or plaque formation in a plaque reduction assay or ELISA-neutralization assay, or by measuring resistance to pathogen challenge *in vivo*.

[066] A “Th1” biased immune response is characterized by the presence of CD4+ T helper cells that produce IL-2 and IFN- γ , and thus, by the secretion or presence of IL-2 and IFN- γ . In contrast, a “Th2” biased immune response is characterized by a preponderance of CD4+ helper cells that produce IL-4, IL-5, and IL-13.

[067] An “immunologically effective amount” is a quantity of a composition (typically, an immunogenic composition) used to elicit an immune response in a subject to the

composition or to an antigen in the composition. Commonly, the desired result is the production of an antigen (*e.g.*, pathogen)-specific immune response that is capable of or contributes to protecting the subject against the pathogen. However, to obtain a protective immune response against a pathogen can require multiple administrations of the immunogenic composition. Thus, in the context of this disclosure, the term immunologically effective amount encompasses a fractional dose that contributes in combination with previous or subsequent administrations to attaining a protective immune response.

[068] The adjective “pharmaceutically acceptable” indicates that the referent is suitable for administration to a subject (*e.g.*, a human or animal subject). Remington’s Pharmaceutical Sciences, by E. W. Martin, Mack Publishing Co., Easton, PA, 15th Edition (1975), describes compositions and formulations (including diluents) suitable for pharmaceutical delivery of therapeutic and/or prophylactic compositions, including immunogenic compositions.

[069] The term “modulate” in reference to a response, such as an immune response, means to alter or vary the onset, magnitude, duration or characteristics of the response. An agent that modulates an immune response alters at least one of the onset, magnitude, duration or characteristics of an immune response following its administration, or that alters at least one of the onset, magnitude, duration or characteristic as compared to a reference agent.

[070] The term “reduces” is a relative term, such that an agent reduces a response or condition if the response or condition is quantitatively diminished following administration of the agent, or if it is diminished following administration of the agent, as compared to a reference agent. Similarly, the term “prevents” does not necessarily mean that an agent completely eliminates the response or condition, so long as at least one characteristic of the response or condition is eliminated. Thus, an immunogenic composition that reduces or prevents an infection or a response, such as a pathological response, *e.g.*, vaccine enhanced viral disease, can, but does not necessarily completely eliminate such an infection or response, so long as the infection or response is measurably diminished, for example, by at least about 50%, such as by at least about 70%, or about 80%, or even by about 90% of (that is to 10% or less than) the infection or response in the absence of the agent, or in comparison to a reference agent.

[071] A “subject” is a living multi-cellular vertebrate organism. In the context of this disclosure, the subject can be an experimental subject, such as a non-human animal, *e.g.*, a mouse, a cotton rat, or a non-human primate. Alternatively, the subject can be a human subject.

PreF ANTIGENS

[072] In nature, the RSV F protein is expressed as a single polypeptide precursor 574 amino acids in length, designated F0. *In vivo*, F0 oligomerizes in the endoplasmic reticulum and is proteolytically processed by a furin protease at two conserved furin consensus sequences (furin cleavage sites), RARR¹⁰⁹ (SEQ ID NO:15) and RKRR¹³⁶ (SEQ ID NO:16) to generate an oligomer consisting of two disulfide-linked fragments. The smaller of these fragments is termed F2 and originates from the N-terminal portion of the F0 precursor. It will be recognized by those of skill in the art that the abbreviations F0, F1 and F2 are commonly designated F₀, F₁ and F₂ in the scientific literature. The larger, C-terminal F1 fragment anchors the F protein in the membrane via a sequence of hydrophobic amino acids, which are adjacent to a 24 amino acid cytoplasmic tail. Three F2-F1 dimers associate to form a mature F protein, which adopts a metastable prefusogenic (“prefusion”) conformation that is triggered to undergo a conformational change upon contact with a target cell membrane. This conformational change exposes a hydrophobic sequence, known as the fusion peptide, which associates with the host cell membrane and promotes fusion of the membrane of the virus, or an infected cell, with the target cell membrane.

[073] The F1 fragment contains at least two heptad repeat domains, designated HRA and HRB, and situated in proximity to the fusion peptide and transmembrane anchor domains, respectively. In the prefusion conformation, the F2-F1 dimer forms a globular head and stalk structure, in which the HRA domains are in a segmented (extended) conformation in the globular head. In contrast, the HRB domains form a three-stranded coiled coil stalk extending from the head region. During transition from the prefusion to the postfusion conformations, the HRA domains collapse and are brought into proximity to the HRB domains to form an anti-parallel six helix bundle. In the postfusion state the fusion peptide and transmembrane domains are juxtaposed to facilitate membrane fusion.

[074] Although the conformational description provided above is based on molecular modeling of crystallographic data, the structural distinctions between the prefusion and

postfusion conformations can be monitored without resort to crystallography. For example, electron microscopy can be used to distinguish between the prefusion and postfusion (alternatively designated prefusogenic and fusogenic) conformations, as demonstrated by Calder *et al.*, *Virology*, 271:122-131 (2000) and Morton *et al.*, *Virology*, 311:275-288, which are incorporated herein by reference for the purpose of their technological teachings. The prefusion conformation can also be distinguished from the fusogenic (postfusion) conformation by liposome association assays as described by Connolly *et al.*, *Proc. Natl. Acad. Sci. USA*, 103:17903-17908 (2006), which is also incorporated herein by reference for the purpose of its technological teachings. Additionally, prefusion and fusogenic conformations can be distinguished using antibodies (e.g., monoclonal antibodies) that specifically recognize conformation epitopes present on one or the other of the prefusion or fusogenic form of the RSV F protein, but not on the other form. Such conformation epitopes can be due to preferential exposure of an antigenic determinant on the surface of the molecule. Alternatively, conformational epitopes can arise from the juxtaposition of amino acids that are non-contiguous in the linear polypeptide.

[075] The PreF antigens disclosed herein are designed to stabilize and maintain the prefusion conformation of the RSV F protein, such that in a population of expressed protein, a substantial portion of the population of expressed protein is in the prefusogenic (prefusion) conformation (e.g., as predicted by structural and/or thermodynamic modeling or as assessed by one or more of the methods disclosed above). Stabilizing modifications are introduced into a native (or synthetic) F protein, such as the exemplary F protein of SEQ ID NO:2, such that the major immunogenic epitopes of the prefusion conformation of the F protein are maintained following introduction of the PreF antigen into a cellular or extracellular environment (for example, *in vivo*, e.g., following administration to a subject).

[076] First, a heterologous stabilizing domain can be placed at the C-terminal end of the construct in order to replace the membrane anchoring domain of the F0 polypeptide. This stabilizing domain is predicted to compensate for the HRB instability, helping to stabilize the -prefusion conformer. In exemplary embodiments, the heterologous stabilizing domain is a protein multimerization domain. One particularly favorable example of such a protein multimerization domain is a trimerization domain. Exemplary trimerization

domains fold into a coiled-coil that promotes assembly into trimers of multiple polypeptides having such coiled-coil domains. One favorable example of a trimerization domain is an isoleucine zipper. An exemplary isoleucine zipper domain is the engineered yeast GCN4 isoleucine variant described by *Harbury et al. Science* 262:1401-1407 (1993). The sequence of one suitable isoleucine zipper domain is represented by SEQ ID NO:11, although variants of this sequence that retain the ability to form a coiled-coil stabilizing domain are equally suitable. Alternative stabilizing coiled coil trimerization domains include: TRAF2 (GENBANK® Accession No. Q12933 [gi:23503103]; amino acids 299-348); Thrombospondin 1 (Accession No. P07996 [gi:135717]; amino acids 291-314); Matrilin-4 (Accession No. O95460 [gi:14548117]; amino acids 594-618; CMP (matrilin-1) (Accession No. NP_002370 [gi:4505111]; amino acids 463-496; HSF1 (Accession No. AAX42211 [gi:61362386]; amino acids 165-191; and Cubilin (Accession No. NP_001072 [gi:4557503]; amino acids 104-138. It is expected that a suitable trimerization domain results in the assembly of a substantial portion of the expressed protein into trimers. For example, at least 50% of a recombinant PreF polypeptide having a trimerization domain will assemble into a trimer (e.g., as assessed by AFF-MALS). Typically, at least 60%, more favorably at least 70%, and most desirably at least about 75% or more of the expressed polypeptide exists as a trimer.

[077] In order to stabilize HRB even more, the leucine residue located at position 512 (relative to the native F0 protein) of the PreF can be substituted by a lysine (L482K of the exemplary PreF antigen polypeptide of SEQ ID NO:6). This substitution improves the coiled coil hydrophobic residue periodicity. Similarly, a lysine can be added following the amino acid at position 105.

[078] Secondly, pep27 can be removed. Analysis of a structural model of the RSV F protein in the prefusion state suggests that pep27 creates a large unconstrained loop between F1 and F2. This loop does not contribute to stabilization of the prefusion state, and is removed following cleavage of the native protein by furin.

[079] Third, one or both furin cleavage motifs can be deleted. With this design, the fusion peptide is not cleaved from F2, preventing release from the globular head of the prefusion conformer and accessibility to nearby membranes. Interaction between the fusion peptide and the membrane interface is predicted to be a major issue in the prefusion state instability. During the fusion process, interaction between the fusion peptide and the

target membrane results in the exposure of the fusion peptide from within the globular head structure, enhancing instability of the prefusion state and folding into post-fusion conformer. This conformation change enables the process of membrane fusion. Removal of one or both of the furin cleavage sites is predicted to prevent membrane accessibility to the N-terminal part of the fusion peptide, stabilizing the prefusion state. Thus, in exemplary embodiments disclosed herein, removal of the furin cleavage motifs results in a PreF antigen that comprises an intact fusion peptide, which is not cleaved by furin during or following processing and assembly.

[080] Optionally, at least one non-furin cleavage site can also be removed, for example by substitution of one or more amino acids. For example, experimental evidence suggests that under conditions conducive to cleavage by certain metalloproteinases, the PreF antigen can be cleaved in the vicinity of amino acids 110-118 (for example, with cleavage occurring between amino acids 112 and 113 of the PreF antigen; between a leucine at position 142 and glycine at position 143 of the reference F protein polypeptide of SEQ ID NO:2). Accordingly, modification of one or more amino acids within this region can reduce cleavage of the PreF antigen. For example, the leucine at position 112 can be substituted with a different amino acid, such as isoleucine, glutamine or tryptophan (as shown in the exemplary embodiment of SEQ ID NO:20). Alternatively or additionally, the glycine at position 113 can be substituted by a serine or alanine.

[081] Optionally, a PreF antigen can include one or more modifications that alters the glycosylation pattern or status (*e.g.*, by increasing or decreasing the proportion of molecules glycosylated at one or more of the glycosylation sites present in a native F protein polypeptide. For example, the native F protein polypeptide of SEQ ID NO:2 is predicted to be glycosylated at amino acid positions 27, 70 and 500 (corresponding to positions 27, 70 and 470 of the exemplary PreF antigen of SEQ ID NO:6). In an embodiment, a modification is introduced in the vicinity of the glycosylation site at amino acid position 500 (designated N470). For example, the glycosylation site can be removed by substituting an amino acid, such as glutamine (Q) in place of the asparagine at position 500 (of the reference sequence, which corresponds by alignment to position 470 of the exemplary PreF antigen). Favorably, a modification that increases glycosylation efficiency at this glycosylation site is introduced. Examples of suitable modifications include at positions 500-502, the following amino acid sequences: NGS; NKS; NGT;

NKT. Interestingly, it has been found that modifications of this glycosylation site that result in increased glycosylation also result in substantially increased PreF production. Thus, in certain embodiments, the PreF antigens have a modified glycosylation site at the position corresponding to amino acid 500 of the reference PreF sequence (SEQ ID NO:2), *e.g.*, at position 470 of the PreF antigen exemplified by SEQ ID NO:6). Suitable, modifications include the sequences: NGS; NKS; NGT; NKT at amino acids corresponding to positions 500-502 of the reference F protein polypeptide sequence. The amino acid of an exemplary embodiment that includes an “NGT” modification is provided in SEQ ID NO:18. One of skill in the art can easily determine similar modifications for corresponding NGS, NKS, and NKT modifications. Such modifications are favorably combined with any of the stabilizing mutations disclosed herein (*e.g.*, a heterologous coiled-coil, such as an isoleucine zipper, domain and/or a modification in a hydrophobic region, and/or removal of pep27, and/or removal of a furin cleavage site, and/or removal of a non-furin cleavage site, and/or removal of a non-furin cleavage site). For example, in one specific embodiment, the PreF antigen includes a substitution that eliminates a non-furin cleavage site and a modification that increases glycosylation. An exemplary sequence is provided in SEQ ID NO:22 (which exemplary embodiment includes an “NGT” modification and the substitution of glutamine in the place of leucine at position 112).

[082] More generally, any one of the stabilizing modifications disclosed herein, *e.g.*, addition of a heterologous stabilizing domain, such as a coiled-coil (for example, an isoleucine zipper domain), preferably situated at the C-terminal end of the soluble PreF antigen; modification of a residue, such as leucine to lysine, in the hydrophobic HRB domain; removal of pep27; removal of one or both furin cleavage motifs; removal of a non-furin cleavage site; and/or modification of a glycosylation site can be employed in combination with any one or more (or up to all-in any desired combination) of the other stabilizing modifications. For example, a heterologous coiled-coil (or other heterologous stabilizing domain) can be utilized alone or in combination with any of: a modification in a hydrophobic region, and/or removal of pep27, and/or removal of a furin cleavage site, and/or removal of a non-furin cleavage site, and/or removal of a non-furin cleavage site. In certain specific embodiments, the PreF antigen includes a C-terminal coiled-coil (isoleucine zipper) domain, a stabilizing substitution in the HRB hydrophobic domain, and removal of one or both furin cleavage sites. Such an embodiment includes an intact fusion

peptide that is not removed by furin cleavage. In one specific embodiment, the PreF antigen also includes a modified glycosylation site at amino acid position 500.

[083] The native F protein polypeptide can be selected from any F protein of an RSV A or RSV B strain, or from variants thereof (as defined above). In certain exemplary embodiments, the F protein polypeptide is the F protein represented by SEQ ID NO:2. To facilitate understanding of this disclosure, all amino acid residue positions, regardless of strain, are given with respect to (that is, the amino acid residue position corresponds to) the amino acid position of the exemplary F protein. Comparable amino acid positions of any other RSV A or B strain can be determined easily by those of ordinary skill in the art by aligning the amino acid sequences of the selected RSV strain with that of the exemplary sequence using readily available and well-known alignment algorithms (such as BLAST, *e.g.*, using default parameters). Numerous additional examples of F protein polypeptides from different RSV strains are disclosed in WO2008114149 (which is incorporated herein by reference for the purpose of providing additional examples of RSV F and G protein sequences). Additional variants can arise through genetic drift, or can be produced artificially using site directed or random mutagenesis, or by recombination of two or more preexisting variants. Such additional variants are also suitable in the context of the PreF (and PreF-G) antigens disclosed herein.

[084] In selecting F2 and F1 domains of the F protein, one of skill in the art will recognize that it is not strictly necessary to include the entire F2 and/or F1 domain. Typically, conformational considerations are of importance when selecting a subsequence (or fragment) of the F2 domain. Thus, the F2 domain typically includes a portion of the F2 domain that facilitates assembly and stability of the polypeptide. In certain exemplary variants, the F2 domain includes amino acids 26-105. However, variants having minor modifications in length (by addition, or deletion of one or more amino acids) are also possible.

[085] Typically, at least a subsequence (or fragment) of the F1 domain is selected and designed to maintain a stable conformation that includes immunodominant epitopes of the F protein. For example, it is generally desirable to select a subsequence of the F1 polypeptide domain that includes epitopes recognized by neutralizing antibodies in the regions of amino acids 262-275 (palivizumab neutralization) and 423-436 (Centocor's ch101F MAb). Additionally, desirable to include T cell epitopes, *e.g.*, in the region of

amino acids 328-355. Most commonly, as a single contiguous portion of the F1 subunit (e.g., spanning amino acids 262-436) but epitopes could be retained in a synthetic sequence that includes these immunodominant epitopes as discontinuous elements assembled in a stable conformation. Thus, an F1 domain polypeptide comprises at least about amino acids 262-436 of an RSV F protein polypeptide. In one non-limiting example provided herein, the F1 domain comprises amino acids 137 to 516 of a native F protein polypeptide. One of skill in the art will recognize that additional shorter subsequences can be used at the discretion of the practitioner.

[086] When selecting a subsequence of the F2 or F1 domain (or as will be discussed below with respect to the G protein component of certain PreF-G antigens), in addition to conformational consideration, it can be desirable to choose sequences (e.g., variants, subsequences, and the like) based on the inclusion of additional immunogenic epitopes. For example, additional T cell epitopes can be identified using anchor motifs or other methods, such as neural net or polynomial determinations, known in the art, see, e.g., RANKPEP (available on the world wide web at: mif.dfci.harvard.edu/Tools/rankpep.html); ProPredI (available on the world wide web at: imtech.res.in/raghava/propredI/index.html); Bimas (available on the world wide web at: www-bimas.dcrt.nih.gov/molbi/hla_bind/index.html); and SYFPEITH (available on the world wide web at: syfpeithi.bmi-heidelberg.com/scripts/MHCServer.dll/home.htm). For example, algorithms are used to determine the “binding threshold” of peptides, and to select those with scores that give them a high probability of MHC or antibody binding at a certain affinity. The algorithms are based either on the effects on MHC binding of a particular amino acid at a particular position, the effects on antibody binding of a particular amino acid at a particular position, or the effects on binding of a particular substitution in a motif-containing peptide. Within the context of an immunogenic peptide, a “conserved residue” is one which appears in a significantly higher frequency than would be expected by random distribution at a particular position in a peptide. Anchor residues are conserved residues that provide a contact point with the MHC molecule. T cell epitopes identified by such predictive methods can be confirmed by measuring their binding to a specific MHC protein and by their ability to stimulate T cells when presented in the context of the MHC protein.

[087] Favorably, the PreF antigens (including PreF-G antigens as discussed below) include a signal peptide corresponding to the expression system, for example, a

mammalian or viral signal peptide, such as an RSV F0 native signal sequence (e.g., amino acids 1-25 of SEQ ID NO:2 or amino acids 1-25 of SEQ ID NO:6). Typically, the signal peptide is selected to be compatible with the cells selected for recombinant expression. For example, a signal peptide (such as a baculovirus signal peptide, or the melittin signal peptide, can be substituted for expression, in insect cells. Suitable plant signal peptides are known in the art, if a plant expression system is preferred. Numerous exemplary signal peptides are known in the art, (see, e.g., see *Zhang & Henzel, Protein Sci.*, 13:2819-2824 (2004), which describes numerous human signal peptides) and are catalogued, e.g., in the SPdb signal peptide database, which includes signal sequences of archaea, prokaryotes and eukaryotes (<http://proline.bic.nus.edu.sg/spdb/>). Optionally, any of the preceding antigens can include an additional sequence or tag, such as a His-tag to facilitate purification.

[088] Optionally, the PreF antigen can include additional immunogenic components. In certain particularly favorable embodiments, the PreF antigen includes an RSV G protein antigenic component. Exemplary chimeric proteins having a PreF and G component include the following PreF_V1 (represented by SEQ ID NOs:7 and 8) and PreF_V2 (represented by SEQ ID NOs:9 and 10).

[089] In the PreF-G antigens, an antigenic portion of the G protein (e.g., a truncated G protein, such as amino acid residues 149-229) is added at the C-terminal end of the construct. Typically, the G protein component is joined to the F protein component via a flexible linker sequence. For example, in the exemplary PreF_V1 design, the G protein is joined to the PreF component by a -GGSGGSGGS- linker (SEQ ID NO:14). In the PreF_V2 design, the linker is shorter. Instead of having the -GGSGGSGGS- linker (SEQ ID NO:14), PreF_V2 has 2 glycines (-GG-) for linker.

[090] Where present, the G protein polypeptide domain can include all or part of a G protein selected from any RSV A or RSV B strain. In certain exemplary embodiments, the G protein is (or is 95% identical to) the G protein represented by SEQ ID NO:4. Additional examples of suitable G protein sequences can be found in WO2008114149 (which is incorporated herein by reference).

[091] The G protein polypeptide component is selected to include at least a subsequence (or fragment) of the G protein that retains the immunodominant T cell epitope(s), e.g., in the region of amino acids 183-197, such as fragments of the G protein that include amino

acids 151-229, 149-229, or 128-229 of a native G protein. In one exemplary embodiment, the G protein polypeptide is a subsequence (or fragment) of a native G protein polypeptide that includes all or part of amino acid residues 149 to 229 of a native G protein polypeptide. One of skill in the art will readily appreciate that longer or shorter portions of the G protein can also be used, so long as the portion selected does not conformationally destabilize or disrupt expression, folding or processing of the PreF-G antigen. Optionally, the G protein domain includes an amino acid substitution at position 191, which has previously been shown to be involved in reducing and/or preventing enhanced disease characterized by eosinophilia associated with formalin inactivated RSV vaccines. A thorough description of the attributes of naturally occurring and substituted (N191A) G proteins can be found, *e.g.*, in US Patent Publication No. 2005/0042230, which is incorporated herein by reference.

[092] For example, with respect to selection of sequences corresponding to naturally occurring strains, one or more of the domains can correspond in sequence to an RSV A or B strain, such as the common laboratory isolates designated A2 or Long, or any other naturally occurring strain or isolate (as disclosed in the aforementioned WO2008114149). In addition to such naturally occurring and isolated variants, engineered variants that share sequence similarity with the aforementioned sequences can also be employed in the context of PreF (including PreF-G) antigens. It will be understood by those of skill in the art, that the similarity between PreF antigen polypeptide (and polynucleotide sequences as described below), as for polypeptide (and nucleotide sequences in general), can be expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity); the higher the percentage, the more similar are the primary structures of the two sequences. In general, the more similar the primary structures of two amino acid (or polynucleotide) sequences, the more similar are the higher order structures resulting from folding and assembly. Variants of a PreF polypeptide (and polynucleotide) sequences typically have one or a small number of amino acid deletions, additions or substitutions but will nonetheless share a very high percentage of their amino acid, and generally their polynucleotide sequence. More importantly, the variants retain the structural and, thus, conformational attributes of the reference sequences disclosed herein.

[093] Methods of determining sequence identity are well known in the art, and are applicable to PreF antigen polypeptides, as well as the nucleic acids that encode them (e.g., as described below). Various programs and alignment algorithms are described in: Smith and Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman and Wunsch, *J. Mol. Biol.* 48:443, 1970; Higgins and Sharp, *Gene* 73:237, 1988; Higgins and Sharp, *CABIOS* 5:151, 1989; Corpet *et al.*, *Nucleic Acids Research* 16:10881, 1988; and Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444, 1988. Altschul *et al.*, *Nature Genet.* 6:119, 1994, presents a detailed consideration of sequence alignment methods and homology calculations. The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul *et al.*, *J. Mol. Biol.* 215:403, 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, MD) and on the internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. A description of how to determine sequence identity using this program is available on the NCBI website on the internet.

[094] In some instances, the PreF antigens has one or more amino acid modification relative to the amino acid sequence of the naturally occurring strain from which it is derived (e.g., in addition to the aforementioned stabilizing modifications). Such differences can be an addition, deletion or substitution of one or more amino acids. A variant typically differs by no more than about 1%, or 2%, or 5%, or 10%, or 15%, or 20% of the amino acid residues. For example, a variant PreF antigen (including PreF-G) polypeptide sequence can include 1, or 2, or up to 5, or up to about 10, or up to about 15, or up to about 50, or up to about 100 amino acid differences as compared to the exemplary PreF antigen polypeptide sequences of SEQ ID NOs:6, 8, 10, 18, 20 and/or 22. Thus, a variant in the context of an RSV F or G protein, or PreF antigen (including PreF-G antigen), typically shares at least 80%, or 85%, more commonly, at least about 90% or more, such as 95%, or even 98% or 99% sequence identity with a reference protein, e.g., the reference sequences illustrated in SEQ ID NO:2, 4, 6, 8, 10, 18, 20 and/or 22, or any of the exemplary PreF antigens disclosed herein. Additional variants included as a feature of this disclosure are PreF antigens (including PreF-G antigens) that include all or part of a nucleotide or amino acid sequence selected from the naturally occurring variants disclosed in WO2008114149. Additional variants can arise through genetic drift, or can be produced artificially using site directed or random mutagenesis, or by recombination of two or more preexisting variants. Such additional variants are also suitable in the context

of the PreF (and PreF-G) antigens disclosed herein. For example, the modification can be a substitution of one or more amino acids (such as two amino acids, three amino acids, four amino acids, five amino acids, up to about ten amino acids, or more) that do not alter the conformation or immunogenic epitopes of the resulting PreF antigen.

[095] Alternatively or additionally, the modification can include a deletion of one or more amino acids and/or an addition of one or more amino acids. Indeed, if desired, one or more of the polypeptide domains can be a synthetic polypeptide that does not correspond to any single strain, but includes component subsequences from multiple strains, or even from a consensus sequence deduced by aligning multiple strains of RSV virus polypeptides. In certain embodiments, one or more of the polypeptide domains is modified by the addition of an amino acid sequence that constitutes a tag, which facilitates subsequent processing or purification. Such a tag can be an antigenic or epitope tag, an enzymatic tag or a polyhistidine tag. Typically the tag is situated at one or the other end of the protein, such as at the C-terminus or N-terminus of the antigen or fusion protein.

NUCLEIC ACIDS THAT ENCODE PREF ANTIGENS

[096] Another aspect of this disclosure concerns recombinant nucleic acids that encode PreF antigens as described above. More explicitly, such nucleic acids encode polypeptides that include a soluble F protein polypeptide antigen that includes an F2 domain and an F1 domain of an RSV F protein polypeptide, which includes at least one modification selected from: (i) an addition of an amino acid sequence comprising a heterologous trimerization domain; (ii) a deletion of at least one furin cleavage site; (iii) a deletion of at least one non-furin cleavage site; (iv) a deletion of one or more amino acids of the pep27 domain; and, (v) at least one substitution or addition of a hydrophilic amino acid in a hydrophobic domain of the F protein extracellular domain. Optionally, such a polynucleotide encodes a PreF antigen with a modification in a glycosylation site. The nature and structural details of such polypeptides are disclosed in detail above. One of skill in the art will readily be able to determine nucleotide sequences that encode any and all of the described polypeptide sequences based on the teachings herein, including the exemplary sequences provided in the sequence listing, and otherwise included (e.g., by incorporation by reference) in this disclosure.

[097] In certain embodiments, the recombinant nucleic acids are codon optimized for expression in a selected prokaryotic or eukaryotic host cell. For example, SEQ ID NOs: 5,

12, 17, 19 and 21 are different illustrative, non-limiting, examples of sequences that encode a PreF antigen, which have been codon optimized for expression in mammalian, *e.g.*, CHO, cells. To facilitate replication and expression, the nucleic acids can be incorporated into a vector, such as a prokaryotic or a eukaryotic expression vector. Host cells including recombinant PreF antigen-encoding nucleic acids are also a feature of this disclosure. Favorable host cells include prokaryotic (*i.e.*, bacterial) host cells, such as *E. coli*, as well as numerous eukaryotic host cells, including fungal (*e.g.*, yeast) cells, insect cells, and mammalian cells (such as CHO, VERO and HEK293 cells).

[098] To facilitate replication and expression, the nucleic acids can be incorporated into a vector, such as a prokaryotic or a eukaryotic expression vector. Although the nucleic acids disclosed herein can be included in any one of a variety of vectors (including, for example, bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, pseudorabies, adenovirus, adeno-associated virus, retroviruses and many others), most commonly the vector will be an expression vector suitable for generating polypeptide expression products. In an expression vector, the nucleic acid encoding the PreF antigen is typically arranged in proximity and orientation to an appropriate transcription control sequence (promoter, and optionally, one or more enhancers) to direct mRNA synthesis. That is, the polynucleotide sequence of interest is operably linked to an appropriate transcription control sequence. Examples of such promoters include: the immediate early promoter of CMV, LTR or SV40 promoter, polyhedrin promoter of baculovirus, *E. coli* lac or trp promoter, phage T7 and lambda P_L promoter, and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector typically also contains a ribosome binding site for translation initiation, and a transcription terminator. The vector optionally includes appropriate sequences for amplifying expression. In addition, the expression vectors optionally comprise one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells, such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as kanamycin, tetracycline or ampicillin resistance in *E. coli*.

[099] The expression vector can also include additional expression elements, for example, to improve the efficiency of translation. These signals can include, *e.g.*, an ATG

initiation codon and adjacent sequences. In some cases, for example, a translation initiation codon and associated sequence elements are inserted into the appropriate expression vector simultaneously with the polynucleotide sequence of interest (e.g., a native start codon). In such cases, additional translational control signals are not required. However, in cases where only a polypeptide-coding sequence, or a portion thereof, is inserted, exogenous translational control signals, including an ATG initiation codon is provided for translation of the nucleic acid encoding PreF antigen. The initiation codon is placed in the correct reading frame to ensure translation of the polynucleotide sequence of interest. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. If desired, the efficiency of expression can be further increased by the inclusion of enhancers appropriate to the cell system in use (Scharf *et al.* (1994) Results Probl Cell Differ 20:125-62; Bitter *et al.* (1987) Methods in Enzymol 153:516-544).

[0100] In some instances, the nucleic acid (such as a vector) that encodes the PreF antigen includes one or more additional sequence elements selected to increase and/or optimize expression of the PreF encoding nucleic acid when introduced into a host cell. For example, in certain embodiments, the nucleic acids that encode the PreF antigen include an intron sequence, such as a Human Herpesvirus 5 intron sequence (see, e.g., SEQ ID NO:13). Introns have been repeatedly demonstrated to enhance expression of homologous and heterologous nucleic acids when appropriately positioned in a recombinant construct. Another class of expression-enhancing sequences includes an epigenetic element such as a Matrix Attachment Region (or MAR), or a similar epigenetic element, e.g., STAR elements (for example, such as those STAR elements disclosed in Otte *et al.*, *Biotechnol. Prog.* 23:801-807, 2007). Without being bound by theory, MARs are believed to mediate the anchorage of a target DNA sequence to the nuclear matrix, generating chromatin loop domains that extend outwards from the heterochromatin cores. While MARs do not contain any obvious consensus or recognizable sequence, their most consistent feature appears to be an overall high A/T content, and C bases predominating on one strand. These regions appear to form bent secondary structures that may be prone to strand separation, and may include a core-unwinding element (CUE) that can serve as the nucleation point for strand separation. Several simple AT-rich sequence motifs have been associated with MAR sequences: e.g., the A-box, the T-box, DNA unwinding motifs, SATB1 binding sites (H-box, A/T/C25) and consensus Topoisomerase II sites for vertebrates or Drosophila. Exemplary MAR sequences are described in published US

patent application no. 20070178469, and in international patent application no. WO02/074969 (which are incorporated herein by reference). Additional MAR sequences that can be used to enhance expression of a nucleic acid encoding a PreF antigen include chicken lysozyme MAR, MARp1-42, MARp1-6, MARp1-68, and MARpx-29, described in Girod *et al.*, *Nature Methods*, 4:747-753, 2007 (disclosed in GenBank Accession Nos. EA423306, DI107030, DI106196, DI107561, and DI106512, respectively). One of skill will appreciate that expression can further be modulated by selecting a MAR that produces an intermediate level of enhancement, as is reported for MAR 1-9. If desired, alternative MAR sequences for increasing expression of a PreF antigen can be identified by searching sequence databases, for example, using software such as MAR-Finder (available on the web at futuresoft.org/MarFinder), SMARTTest (available on the web at genomatix.de), or SMARScan I (Levitsky *et al.*, *Bioinformatics* 15:582-592, 1999). In certain embodiments, the MAR is introduced (e.g., transfected) into the host cell on the same nucleic acid (e.g., vector) as the PreF antigen-encoding sequence. In an alternative embodiment, the MAR is introduced on a separate nucleic acid (e.g., in trans) and it can optionally cointegrate with the PreF antigen-encoding polynucleotide sequence.

[0101] Exemplary procedures sufficient to guide one of ordinary skill in the art through the production of recombinant PreF antigen nucleic acids can be found in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press, 1989; Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 3d ed., Cold Spring Harbor Press, 2001; Ausubel *et al.*, *Current Protocols in Molecular Biology*, Greene Publishing Associates, 1992 (and Supplements to 2003); and Ausubel *et al.*, *Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology*, 4th ed., Wiley & Sons, 1999.

[0102] Exemplary nucleic acids that encode PreF antigen polypeptides are represented by SEQ ID NOs: 5, 7, 9, 12, 13, 17, 19 and 21. Variants that include a modification in a glycosylation site, e.g., at the amino acid corresponding to position 500 of SEQ ID NO:2 can be produced by altering (e.g., mutating) the nucleotides in the vicinity of positions 1408-1414 (as compared, for example, to the polynucleotide sequence of SEQ ID NO:12, e.g., SEQ ID NOs:17 and 21). Suitable sequences of nucleotides to encode glycosylation variants (e.g., that increase glycosylation efficiency) include: aacgggt, aacaagt, aacggga, and aacaaga. Alternative sequences, such as cagcagt, which eliminate a glycosylation site

are also possible. Additional variants of can be produced by assembling analogous F and G protein polypeptide sequences selected from any of the known (or subsequently) discovered strains of RSV, *e.g.*, as disclosed in WO2008114149. Additional sequence variants that share sequence identity with the exemplary variants can be produced by those of skill in the art. Typically, the nucleic acid variants will encode polypeptides that differ by no more than 1%, or 2%, or 5%, or 10%, or 15%, or 20% of the amino acid residues. That is, the encoded polypeptides share at least 80%, or 85%, more commonly, at least about 90% or more, such as 95%, or even 98% or 99% sequence identity. It will be immediately understood by those of skill in the art, that the polynucleotide sequences encoding the PreF polypeptides, can themselves share less sequence identity due to the redundancy of the genetic code. In some instances, the PreF antigens has one or more amino acid modification relative to the amino acid sequence of the naturally occurring strain from which it is derived (*e.g.*, in addition to the aforementioned stabilizing modifications). Such differences can be an addition, deletion or substitution of one or more nucleotides or amino acids, respectively. A variant typically differs by no more than about 1%, or 2%, or 5%, or 10%, or 15%, or 20% of the nucleotide residues. For example, a variant PreF antigen (including PreF-G) nucleic acid can include 1, or 2, or up to 5, or up to about 10, or up to about 15, or up to about 50, or up to about 100 nucleotide differences as compared to the exemplary PreF antigen nucleic acids of SEQ ID NOs: 5, 7, 9, 12, 13, 17, 19 and/or 21. Thus, a variant in the context of an RSV F or G protein, or PreF antigen (including PreF-G antigen) nucleic acid, typically shares at least 80%, or 85%, more commonly, at least about 90% or more, such as 95%, or even 98% or 99% sequence identity with a reference sequence, *e.g.*, the reference sequences illustrated in SEQ ID NO:1, 3, 5, 7, 9, 12, 13, 17, 19 and/or 21, or any of the other exemplary PreF antigen nucleic acids disclosed herein. Additional variants included as a feature of this disclosure are PreF antigens (including PreF-G antigens) that include all or part of a nucleotide sequence selected from the naturally occurring variants disclosed in WO2008114149. Additional variants can arise through genetic drift, or can be produced artificially using site directed or random mutagenesis, or by recombination of two or more preexisting variants. Such additional variants are also suitable in the context of the PreF (and PreF-G) antigens disclosed herein.

[0103] In addition to the variant nucleic acids previously described, nucleic acids that hybridize to one or more of the exemplary nucleic acids represented by SEQ ID NOs:1, 3,

5, 7, 9, 12, 13, 17, 19 and/or 21 can also be used to encode PreF antigens. One of skill in the art will appreciate that in addition to the % sequence identity measure discussed above, another indicia of sequence similarity between two nucleic acids is the ability to hybridize. The more similar are the sequences of the two nucleic acids, the more stringent the conditions at which they will hybridize. The stringency of hybridization conditions are sequence-dependent and are different under different environmental parameters. Thus, hybridization conditions resulting in particular degrees of stringency will vary depending upon the nature of the hybridization method of choice and the composition and length of the hybridizing nucleic acid sequences. Generally, the temperature of hybridization and the ionic strength (especially the Na^- and/or Mg^{++} concentration) of the hybridization buffer will determine the stringency of hybridization, though wash times also influence stringency. Generally, stringent conditions are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Conditions for nucleic acid hybridization and calculation of stringencies can be found, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 2001; Tijssen, *Hybridization With Nucleic Acid Probes, Part I: Theory and Nucleic Acid Preparation*, Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Ltd., NY, NY, 1993 and Ausubel *et al.* *Short Protocols in Molecular Biology*, 4th ed., John Wiley & Sons, Inc., 1999.

[0104] For purposes of the present disclosure, “stringent conditions” encompass conditions under which hybridization will only occur if there is less than 25% mismatch between the hybridization molecule and the target sequence. “Stringent conditions” can be broken down into particular levels of stringency for more precise definition. Thus, as used herein, “moderate stringency” conditions are those under which molecules with more than 25% sequence mismatch will not hybridize; conditions of “medium stringency” are those under which molecules with more than 15% mismatch will not hybridize, and conditions of “high stringency” are those under which sequences with more than 10% mismatch will not hybridize. Conditions of “very high stringency” are those under which sequences with more than 6% mismatch will not hybridize. In contrast, nucleic acids that hybridize under “low stringency” conditions include those with much less sequence identity, or with sequence identity over only short subsequences of the nucleic acid. It

will, therefore, be understood that the various variants of nucleic acids that are encompassed by this disclosure are able to hybridize to at least one of SEQ ID NOs: 1, 3, 5, 7, 9, 12, 13, 17, 19 and/or 21 over substantially their entire length.

METHODS OF PRODUCING RSV ANTIGENIC POLYPEPTIDES

[0105] The PreF antigens (including PreF-G antigens, and also where applicable, G antigens) disclosed herein are produced using well established procedures for the expression and purification of recombinant proteins. Procedures sufficient to guide one of skill in the art can be found in the following references: *Sambrook et al., Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 200; and *Ausubel et al. Short Protocols in Molecular Biology*, 4th ed., John Wiley & Sons, Inc., 999. Additional and specific details are provided hereinbelow.

[0106] Recombinant nucleic acids that encode the PreF antigens are introduced into host cells by any of a variety of well-known procedures, such as electroporation, liposome mediated transfection (e.g., using a commercially available liposomal transfection reagent, such as LIPOFECTAMINETM2000 or TRANSFECTINTTM), Calcium phosphate precipitation, infection, transfection and the like, depending on the selection of vectors and host cells. Exemplary nucleic acids that encode PreF antigens (including PreF-G antigens) are provided in SEQ ID NOs:5, 7, 9, 12, 13, 17, 19 and 21. One of skill in the art will appreciate that SEQ ID NOs:5, 7, 9, 12, 13, 17, 19 and 21 are illustrative and not intended to be limiting. For example, polynucleotide sequences that encode the same proteins as SEQ ID NOs:5, 7 and 9, (e.g., represented by SEQ ID NOs: 6, 8 and 10), but that differ only by the redundancy of the genetic code (such as by alternative codon optimization, as shown in SEQ ID NO:12), can easily be used instead of the exemplary sequences of SEQ ID NOs:5, 7, and 9. The same is true of SEQ ID NOs:17, 19 and 21. Similarly, polynucleotide sequences that include expression-enhancing elements, such as internally positioned introns (or by the addition of promoter, enhancer, intron or other similar elements), as illustrated in SEQ ID NO:13, can be employed. One of ordinary skill in the art will recognize that combinations of such modifications are likewise suitable. Similarly, homologous sequences selected from any RSV A or RSV B strain, and/or other sequences that share substantial sequence identity, as discussed above, can also be used to express PreF antigens. Indeed, any of the variant nucleic acids previously disclosed can

suitably be introduced into host cells and used to produce PreF antigens (including PreF-G antigens) and where applicable G polypeptides.

[0107] For example, in certain instances the variant nucleic acids are modified to alter the glycosylation pattern, *e.g.*, as described above by substitution of one or more amino acids in the vicinity of amino acid position 500 (with respect to SEQ ID NO:2, *e.g.*, SEQ ID NO:17). It has been found that modifying the glycosylation pattern, *e.g.*, in combination with modifying a cleavage recognition site, increases production of the PreF antigen in cell culture. In such cases, the methods described hereinbelow for expressing and isolating recombinant PreF antigens provide a process for increasing the production of a PreF antigen by altering the glycosylation pattern of a PreF antigen by substituting one or more amino acids of a glycosylation recognition site, optionally in combination with modifying one or more cleavage sites (such as a non-furin or furin cleavage recognition site) as described above.

[0108] Host cells that include recombinant PreF antigen-encoding nucleic acids are, thus, also a feature of this disclosure. Favorable host cells include prokaryotic (*i.e.*, bacterial) host cells, such as *E. coli*, as well as numerous eukaryotic host cells, including fungal (*e.g.*, yeast, such as *Saccharomyces cerevisiae* and *Pichia pastoris*) cells, insect cells, plant cells, and mammalian cells (such as CHO and HEK293 cells). Recombinant PreF antigen nucleic acids are introduced (*e.g.*, transduced, transformed or transfected) into host cells, for example, via a vector, such as an expression vector. As described above, the vector is most typically a plasmid, but such vectors can also be, for example, a viral particle, a phage, etc. Examples of appropriate expression hosts include: bacterial cells, such as *E. coli*, *Streptomyces*, and *Salmonella typhimurium*; fungal cells, such as *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Neurospora crassa*; insect cells such as *Drosophila* and *Spodoptera frugiperda*; mammalian cells such as 3T3, COS, CHO, BHK, HEK 293 or Bowes melanoma; plant cells, including algae cells, etc. In some instances, transiently transfected cells can be used to produce recombinant PreF antigens. In certain embodiments, cells (*e.g.*, clones) that have stably integrated the PreF nucleic acid are selected and used to produce the recombinant PreF antigen.

[0109] The host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the inserted polynucleotide sequences. The culture conditions, such as temperature, pH and the like,

are typically those previously used with the host cell selected for expression, and will be apparent to those skilled in the art and in the references cited herein, including, *e.g.*, Freshney (1994) Culture of Animal Cells, a Manual of Basic Technique, third edition, Wiley- Liss, New York and the references cited therein. Optionally, the host cells are cultured in serum-free and/or animal product-free medium.

[0110] Expression products corresponding to the nucleic acids of the invention can also be produced in non-animal cells such as plants, yeast, fungi, bacteria and the like. In addition to Sambrook, Berger and Ausubel, details regarding cell culture can be found in Payne *et al.* (1992) Plant Cell and Tissue Culture in Liquid Systems John Wiley & Sons, Inc. New York, NY; Gamborg and Phillips (eds) (1995) Plant Cell, Tissue and Organ Culture: Fundamental Methods Springer Lab Manual, Springer-Verlag (Berlin Heidelberg New York) and Atlas and Parks (eds) The Handbook of Microbiological Media (1993) CRC Press, Boca Raton, FL.

[0111] In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the expressed product. For example, when large quantities of a polypeptide or fragments thereof are needed for the production of antibodies, vectors which direct high level expression of fusion proteins that are readily purified are favorably employed. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the coding sequence of interest, *e.g.*, a polynucleotide of the invention as described above, can be ligated into the vector in-frame with sequences for the amino-terminal translation initiating Methionine and the subsequent 7 residues of beta-galactosidase producing a catalytically active beta galactosidase fusion protein; pIN vectors (Van Hecke & Schuster (1989) J Biol Chem 264:5503-5509). In certain examples, the nucleic acids are introduced into cells via vectors suitable for introduction and expression in prokaryotic cells, *e.g.*, *E. coli* cells. For example, a nucleic acid including a polynucleotide sequence that encodes a PreF antigen can be introduced into any of a variety of commercially available or proprietary vectors, such as the pET series of expression vectors (*e.g.*, pET9b and pET2d). Expression of the coding sequence is inducible by IPTG, resulting in high levels of protein expression. The polynucleotide sequence encoding the PreF antigen is transcribed under the phage T7 promoter. Alternate vectors, such as pURV22 that include a heat-inducible lambda pL promoter are also suitable.

[0112] The expression vector is then introduced (e.g., by electroporation) into a suitable bacterial host. Numerous suitable strains of *E. coli* are available and can be selected by one of skill in the art (for example, the Rosetta and BL21 (DE3) strains have proven favorable for expression of recombinant vectors containing polynucleotide sequences that encode PreF antigens.

[0113] Similarly, in yeast, such as *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH can be used for production of the desired expression products. For reviews, see Berger, Ausubel, and, e.g., Grant *et al.* (1987; *Methods in Enzymology* 153:516-544). In mammalian host cells, a number of expression systems, including both plasmids and viral-based systems, can be utilized.

[0114] In another example, the polynucleotide sequence that encodes the PreF antigen is introduced into insect cells using a Baculovirus Expression Vector System (BEVS). Recombinant baculovirus capable of infecting insect cells can be generated using commercially available vectors, kits and/or systems, such as the BD BaculoGold system from BD BioScience. Briefly, the polynucleotide sequence encoding the antigen is inserted into the pAcSG2 transfer vector. Then, host cells SF9 (*Spodoptera frugiperda*) are co-transfected by pAcSG2-chimeric plasmid and BD BaculoGold, containing the linearized genomic DNA of the baculovirus *Autographa californica nuclear polyhedrosis virus* (AcNPV). Following transfection, homologous recombination occurs between the pACSG2 plasmid and the Baculovirus genome to generate the recombinant virus. In one example, the PreF antigen is expressed under the regulatory control of the polyhedrin promoter (pH). Similar transfer vectors can be produced using other promoters, such as the basic (Ba) and p10 promoters. Similarly, alternative insect cells can be employed, such as SF21 which is closely related to the Sf9, and the High Five cell line derived from a cabbage looper, *Trichoplusia ni*.

[0115] More typically, the polynucleotides that encode the PreF antigens are incorporated into expression vectors that are suitable for introduction and expression in eukaryotic (e.g., insect or mammalian cells). Favorably, such nucleic acids are codon optimized for expression in the selected vector/host cell (for example, the sequences illustrated in SEQ ID NOS:5, 7, 9, 12, 13, 17, 19 and 21 are codon optimized for expression in CHO cells). In one exemplary embodiment, the polynucleotide sequence that encodes the PreF antigen

is introduced into a vector, such as the pEE14 vector developed by Lonza Biologicals. The polypeptide is expressed under a constitutive promoter, such as the immediate early CMV (CytoMegaloVirus) promoter. Selection of the stably transfected cells expressing the polypeptide is made based on the ability of the transfected cells to grow in the absence of a glutamine source. Cells that have successfully integrated the pEE14 are able to grow in the absence of exogenous glutamine, because the pEE14 vector expresses the GS (Glutamine Synthetase) enzyme. Selected cells can be clonally expanded and characterized for expression of the desired PreF polypeptide.

[0116] A host cell is optionally chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the protein include, but are not limited to, glycosylation, (as well as, *e.g.*, acetylation, carboxylation, phosphorylation, lipidation and acylation). Post-translational processing for example, which cleaves a precursor form into a mature form of the protein (for example, by a furin protease) is optionally performed in the context of the host cell. Different host cells such as 3T3, COS, CHO, HeLa, BHK, MDCK, 293, WI38, etc. have specific cellular machinery and characteristic mechanisms for such post-translational activities and can be chosen to ensure the correct modification and processing of the introduced, foreign protein.

[0117] For long-term, high-yield production of recombinant PreF antigens disclosed herein, stable expression systems are typically used. For example, cell lines which stably express a PreF antigen polypeptide are introduced into the host cell using expression vectors which contain viral origins of replication or endogenous expression elements and a selectable marker gene. Following the introduction of the vector, cells are allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. For example, resistant groups or colonies of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. Host cells transformed with a nucleic acid encoding a PreF antigen are optionally cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. As indicated above, if desired, the host cells can be cultured in serum-free (and/or animal product-free) medium.

[0118] Following transduction of a suitable host cell line and growth of the host cells to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period. Optionally, the medium includes components and/or additives that decrease degradation of expressed proteins by proteinases. For example, the medium used for culturing cells to produce PreF antigen can include a protease inhibitor, such as a chelating agent or small molecule inhibitor (e.g., AZ11557272, AS111793, *etc.*), to reduce or eliminate undesired cleavage by cellular, or extracellular (e.g., matrix) proteinases.

[0119] The secreted polypeptide product is then recovered from the culture medium. Alternatively, cells can be harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification. Eukaryotic or microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, or other methods, which are well known to those skilled in the art.

[0120] Expressed PreF antigens can be recovered and purified from recombinant cell cultures by any of a number of methods well known in the art, including ammonium sulfate or ethanol precipitation, acid extraction, filtration, ultrafiltration, centrifugation, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography (e.g., using any of the tagging systems noted herein), hydroxylapatite chromatography, and lectin chromatography. Protein refolding steps can be used, as desired, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed in the final purification steps. In addition to the references noted above, a variety of purification methods are well known in the art, including, *e.g.*, those set forth in Sandana (1997) Bioseparation of Proteins, Academic Press, Inc.; and Bollag *et al.* (1996) Protein Methods, 2nd Edition Wiley-Liss, NY; Walker (1996) The Protein Protocols Handbook Humana Press, NJ, Harris and Angal (1990) Protein Purification Applications: A Practical Approach IRL Press at Oxford, Oxford, U.K.; Scopes (1993) Protein Purification: Principles and Practice 3rd Edition Springer Verlag, NY; Janson and Ryden (1998) Protein Purification: Principles, High Resolution Methods and Applications, Second Edition Wiley-VCH, NY; and Walker (1998) Protein Protocols on CD-ROM Humana Press, NJ.

[0121] In one exemplary embodiment, the PreF proteins are recovered from cells according to the following purification scheme. Following introduction of a recombinant nucleic acid encoding the PreF polypeptide into host CHO cells, transiently transfected host cells or expanded stable populations comprising the introduced polynucleotide sequence are grown in medium and under conditions suitable for growth at an acceptable scale for the desired purpose (e.g., as generally described in Freshney (1994) Culture of Animal Cells, a Manual of Basic Technique, third edition, Wiley- Liss, New York and the references cited therein). Typically, the cells are grown in serum-free medium at 37°C and passaged at 2-3 day intervals in shake flasks or in bioreactors. New cultures established from cells expanded in these conditions, are typically carried out in bioreactors in serum-free medium and incubated at 27 °C with pO2 maintained at 20% for 5 to 7 days in order to produce the preF antigen.

[0122] To recover recombinant PreF antigen, the cell culture is centrifuged and the cell culture supernatant stored at minus 70°C until further use. Following thawing of culture supernatants, the supernatants are diluted 2 x with MilliQ water and adjusted to pH 6.0 with HCl. Diluted supernatant is loaded at 75 cm/h onto a 3 L CM Ceramic HyperD FF resin packed in BPG 140/500 column, equilibrated in 20 mM phosphate pH 6.0. After loading of the sample, equilibration buffer is processed through the column to get back to UV baseline. After washing with 5 column volumes (CV) of 25 mM phosphate pH 7.0 buffer, elution is performed using a 50 mM Phosphate pH 7.0 buffer containing 0.1 M NaCl.

[0123] The CM Hyper D eluate is diluted 3.3 x with 20 mM phosphate, pH 7.0 to be processed onto a 270 ml Hydroxyapatite Type II column (packed in XK 50), equilibrated with 20 mM PO₄ (Na) buffer pH 7.0, at 50 mL/min. After washing the column with the equilibration buffer (~3 CV), elution is performed using a 20 mM PO₄ (Na) pH 7.0 buffer containing 0.5 M NaCl.

[0124] The HA eluate is processed at 15 mL/min (to respect a 10 minutes contact time with the resin), onto a 150 mL Capto Adhere column (packed in XK 26), equilibrated in 20 mM phosphate pH 7.0. After washing with 5 CV of 10 mM phosphate pH 7.0 containing 0.1 M arginine buffer, elution is performed using a 10 mM Phosphate pH 7.0 buffer containing 0.6 M arginine.

[0125] The Capto Adhere eluate is then concentrated approximately 10 x for processing onto a preparative size exclusion chromatography (SEC) column. Concentration is performed using a 50 kD Pellicon polyethersulfone membrane. Before being processed onto the SEC column, the material is filtered through a PLANNOVA 20N 100 cm² filter, used as a viral clearance step. This nanofiltration step can be either placed after or before concentration on Pellicon membrane.

[0126] Preparative SEC is then performed using a 500 mL Superdex S200 column and 10 mM phosphate (Na/K₂), 160 mM NaCl, pH 6.5 buffer (corresponding to final buffer) as mobile phase. A volume of concentrated PreF corresponding to 5% of SEC column volume is loaded onto the resin at ~2.6 mL/min. Typically, fractions of 10 mL are collected. Analytical pools of fractions can be analyzed on SDS gel by silver staining and western blot anti HCP (Host cell proteins) if desired to optimize yields while minimizing HCP levels.

[0127] Purified bulk is obtained after filtration on 0.22 µm Millex filters (alternatively a Sterivex filter can be used). If desired the purified PreF antigen preparation can be stored at minus 70°C prior to use.

[0128] Alternatively, PreF proteins can include a polyhistidine (*e.g.*, six histidine) tag, which can be used to facilitate purification. For such histidine tagged PreF polypeptides, the following purification protocol can be employed. Prior to purification using immobilized metal ion affinity chromatography (IMAC), the cell culture supernatant is diluted twofold in buffer A (20mm Bicine, pH8,5) and pH is adjusted to 8.5. The resulting solution is loaded on a Q sepharose FF column (GE Healthcare), *e.g.*, of 23 ml of column volume, previously equilibrated with Buffer A. PreF proteins are captured on the column, along with some host cell contaminants. The culture media components that would interfere with the IMAC purification step are not retained and are eliminated in the flow through. The proteins are separated and eluted by a stepwise elution of 200mM, 400mM, 600mM, 800mM and 1M NaCl. PreF proteins of interest are eluted during the first step at 200mM NaCl. Optionally, recovery can be monitored using SDS PAGE and western blotting using an anti His-tag antibody to detect the tagged PreF protein. Fractions can be pooled prior to continuing the purification.

[0129] The (pooled) PreF protein containing eluate is diluted threefold in buffer B (20mM Bicine, 500mM NaCl, pH8,3) and pH is adjusted to 8.3. The resulting solution is loaded

on IMAC sepharose FF resin loaded with Nickel chloride (GE Healthcare) (e.g., of 5 ml of column volume), previously equilibrated with buffer B. PreF are bound to the resin and the majority of host cell contaminants are eluted in the flow through. The column is washed with 20mM Imidazole in order to remove weakly bound contaminants. PreF proteins are eluted by a step elution of 250mM Imidazole. SDS PAGE stained with coomassie blue and western blot anti His-tag can be performed to identify positive fractions.

[0130] The pool from IMAC can then be concentrated to a concentration of at least 150 µg/ml using a centricon concentration device (Millipore) and the protein can be dialysed in PBS buffer supplemented with 500mM L-Arginine. Resulting protein is quantified using RCDC protein assay (BioRad) and stored at -70 or -80°C until use.

[0131]

IMMUNOGENIC COMPOSITIONS AND METHODS

[0132] Also provided are immunogenic compositions including any of the PreF antigens disclosed above (such as those exemplified by SEQ ID NOS: 6, 8, 10, 18, 20 and 22) and a pharmaceutically acceptable carrier or excipient.

[0133] In certain embodiments, typically, embodiments in which the PreF antigen does not include a G protein component (such as SEQ ID NO:6), the immunogenic composition can include an isolated, recombinant and/or purified G protein. Numerous suitable G proteins have been described in the art, and include full length recombinant G proteins and chimeric proteins made up of a portion of the G protein (such as amino acids 128-229 or 130-230) and a fusion partner (such as thioredoxin), or a signal and/or leader sequence, that facilitates expression and/or purification. Exemplary G proteins for use in admixture with a PreF antigen can be found in WO2008114149, US Patent No. 5,149,650, US Patent No. 6,113,911, US Published Application No. 20080300382, and US Patent No. 7,368,537, each of which is incorporated herein by reference. As indicated with respect to the chimeric PreF-G proteins, a smaller fragment of the G protein, such as the portion between amino acids 149-229, or the portion between approximately 128 to approximately 229 can favorably be employed in the context of mixtures involving a PreF (without G) and G. As discussed above, the important consideration is the presence of immunodominant epitopes, e.g., included within the region of amino acids 183-197. Alternatively, a full-length G protein can be employed in such compositions.

[0134] Pharmaceutically acceptable carriers and excipients are well known and can be selected by those of skill in the art. For example, the carrier or excipient can favorably include a buffer. Optionally, the carrier or excipient also contains at least one component that stabilizes solubility and/or stability. Examples of solubilizing/stabilizing agents include detergents, for example, laurel sarcosine and/or tween. Alternative solubilizing/stabilizing agents include arginine, and glass forming polyols (such as sucrose, trehalose and the like). Numerous pharmaceutically acceptable carriers and/or pharmaceutically acceptable excipients are known in the art and are described, *e.g.*, in *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, PA, 5th Edition (975).

[0135] Accordingly, suitable excipients and carriers can be selected by those of skill in the art to produce a formulation suitable for delivery to a subject by a selected route of administration.

[0136] Suitable excipients include, without limitation: glycerol, Polyethylene glycol (PEG), Sorbitol, Trehalose, N-lauroylsarcosine sodium salt, L-proline, Non detergent sulfobetaine, Guanidine hydrochloride, Urea, Trimethylamine oxide, KCl, Ca^{2+} , Mg^{2+} , Mn^{2+} , Zn^{2+} and other divalent cation related salts, Dithiothreitol, Dithioerytrol, and β -mercaptoethanol. Other excipients can be detergents (including: Tween80, Tween20, Triton X-00, NP-40, Empigen BB, Octylglucoside, Lauroyl maltoside, Zwittergent 3-08, Zwittergent 3-0, Zwittergent 3-2, Zwittergent 3-4, Zwittergent 3-6, CHAPS, Sodium deoxycholate, Sodium dodecyl sulphate, Cetyltrimethylammonium bromide).

[0137] Optionally, the immunogenic compositions also include an adjuvant. In the context of an immunogenic composition suitable for administration to a subject for the purpose of eliciting a protective immune response against RSV, the adjuvant is selected to elicit a Th1 biased or Th1/Th2 balanced immune response, characterized by the production of interferon-gamma (IFN- γ).

[0138] The adjuvant is typically selected to enhance a Th1 biased immune response (or a Th1/Th2 balanced immune response), characterized by the production and secretion of IFN- γ , in the subject, or population of subjects, to whom the composition is administered. For example, when the immunogenic composition is to be administered to a subject of a particular age group susceptible to (or at increased risk of) RSV infection, the adjuvant is selected to be safe and effective in the subject or population of subjects. Thus, when

formulating an immunogenic composition containing an RSV PreF antigen for administration in an elderly subject (such as a subject greater than 65 years of age), the adjuvant is selected to be safe and effective in elderly subjects. Similarly, when the immunogenic composition containing the RSV PreF antigen is intended for administration in neonatal or infant subjects (such as subjects between birth and the age of two years), the adjuvant is selected to be safe and effective in neonates and infants. In the case of an adjuvant selected for safety and efficacy in neonates and infants, an adjuvant dose can be selected that is a dilution (e.g., a fractional dose) of a dose typically administered to an adult subject.

[0139] Additionally, the adjuvant is typically selected to enhance a Th1 immune response when administered via a route of administration, by which the immunogenic composition is administered. For example, when formulating an immunogenic composition containing a PreF antigen for nasal administration, proteosome and protollin are favorable Th1-biasing adjuvants. In contrast, when the immunogenic composition is formulated for intramuscular administration, adjuvants including one or more of 3D-MPL, squalene (e.g., QS21), liposomes, and/or oil and water emulsions are favorably selected.

[0140] One suitable adjuvant for use in combination with PreF antigens is a non-toxic bacterial lipopolysaccharide derivative. An example of a suitable non-toxic derivative of lipid A, is monophosphoryl lipid A or more particularly 3-Deacylated monophosphoryl lipid A (3D-MPL). 3D-MPL is sold under the name MPL by GlaxoSmithKline Biologicals N.A., and is referred throughout the document as MPL or 3D-MPL. See, for example, US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094. 3D-MPL primarily promotes CD4+ T cell responses with an IFN- γ (Th1) phenotype. 3D-MPL can be produced according to the methods disclosed in GB2220211 A. Chemically it is a mixture of 3-deacylated monophosphoryl lipid A with 3, 4, 5 or 6 acylated chains. In the compositions of the present invention small particle 3D-MPL can be used. Small particle 3D-MPL has a particle size such that it can be sterile-filtered through a 0.22 μ m filter. Such preparations are described in WO94/21292.

[0141] A lipopolysaccharide, such as 3D-MPL, can be used at amounts between 1 and 50 μ g, per human dose of the immunogenic composition. Such 3D-MPL can be used at a level of about 25 μ g, for example between 20-30 μ g, suitably between 21-29 μ g or between 22 and 28 μ g or between 23 and 27 μ g or between 24 and 26 μ g, or 25 μ g. In another

embodiment, the human dose of the immunogenic composition comprises 3D-MPL at a level of about 10 μ g, for example between 5 and 15 μ g, suitably between 6 and 14 μ g, for example between 7 and 13 μ g or between 8 and 12 μ g or between 9 and 11 μ g, or 10 μ g. In a further embodiment, the human dose of the immunogenic composition comprises 3D-MPL at a level of about 5 μ g, for example between 1 and 9 μ g, or between 2 and 8 μ g or suitably between 3 and 7 μ g or 4 and 5 μ g, or 5 μ g.

[0142] In other embodiments, the lipopolysaccharide can be a β (1-6) glucosamine disaccharide, as described in US Patent No. 6,005,099 and EP Patent No. 0 729 473 B1. One of skill in the art would be readily able to produce various lipopolysaccharides, such as 3D-MPL, based on the teachings of these references. Nonetheless, each of these references is incorporated herein by reference. In addition to the aforementioned immunostimulants (that are similar in structure to that of LPS or MPL or 3D-MPL), acylated monosaccharide and disaccharide derivatives that are a sub-portion to the above structure of MPL are also suitable adjuvants. In other embodiments, the adjuvant is a synthetic derivative of lipid A, some of which are described as TLR-4 agonists, and include, but are not limited to: OM174 (2-deoxy-6-o-[2-deoxy-2-[(R)-3-dodecanoyloxytetra-decanoylamino]-4-o-phosphono- β -D-glucopyranosyl]-2-[(R)-3-hydroxytetradecanoylamino]- α -D-glucopyranosyldihydrogenphosphate), (WO 95/14026); OM 294 DP (3S, 9 R) -3-[(R)-dodecanoyloxytetradecanoylamino]-4-oxo-5-aza-9(R)-[(R)-3-hydroxytetradecanoylamino]decan-1,10-diol,1,10-bis(dihydrogenophosphate) (WO 99/64301 and WO 00/0462); and OM 197 MP-Ac DP (3S-, 9R) -3-[(R)-dodecanoyloxytetradecanoylamino]-4-oxo-5-aza-9-[(R)-3-hydroxytetradecanoylamino]decan-1,10-diol,1 -dihydrogenophosphate 10-(6-aminohexanoate) (WO 01/46127).

[0143] Other TLR4 ligands which can be used are alkyl Glucosaminide phosphates (AGPs) such as those disclosed in WO 98/50399 or US Patent No. 6,303,347 (processes for preparation of AGPs are also disclosed), suitably RC527 or RC529 or pharmaceutically acceptable salts of AGPs as disclosed in US Patent No. 6,764,840. Some AGPs are TLR4 agonists, and some are TLR4 antagonists. Both are thought to be useful as adjuvants.

[0144] Other suitable TLR-4 ligands, capable of causing a signaling response through TLR-4 (Sabroe et al, JI 2003 p1630-5) are, for example, lipopolysaccharide from gram-

negative bacteria and its derivatives, or fragments thereof, in particular a non-toxic derivative of LPS (such as 3D-MPL). Other suitable TLR agonists are: heat shock protein (HSP) 10, 60, 65, 70, 75 or 90; surfactant Protein A, hyaluronan oligosaccharides, heparan sulphate fragments, fibronectin fragments, fibrinogen peptides and b-defensin-2, and muramyl dipeptide (MDP). In one embodiment the TLR agonist is HSP 60, 70 or 90. Other suitable TLR-4 ligands are as described in WO 2003/011223 and in WO 2003/099195, such as compound I, compound II and compound III disclosed on pages 4-5 of WO2003/011223 or on pages 3-4 of WO2003/099195 and in particular those compounds disclosed in WO2003/011223 as ER803022, ER803058, ER803732, ER804053, ER804057, ER804058, ER804059, ER804442, ER804680, and ER804764. For example, one suitable TLR-4 ligand is ER804057.

[0145] Additional TLR agonists are also useful as adjuvants. The term “TLR agonist” refers to an agent that is capable of causing a signaling response through a TLR signaling pathway, either as a direct ligand or indirectly through generation of endogenous or exogenous ligand. Such natural or synthetic TLR agonists can be used as alternative or additional adjuvants. A brief review of the role of TLRs as adjuvant receptors is provided in Kaisho & Akira, *Biochimica et Biophysica Acta* 1589:1-13, 2002. These potential adjuvants include, but are not limited to agonists for TLR2, TLR3, TLR7, TLR8 and TLR9. Accordingly, in one embodiment, the adjuvant and immunogenic composition further comprises an adjuvant which is selected from the group consisting of: a TLR-1 agonist, a TLR-2 agonist, TLR-3 agonist, a TLR-4 agonist, TLR-5 agonist, a TLR-6 agonist, TLR-7 agonist, a TLR-8 agonist, TLR-9 agonist, or a combination thereof.

[0146] In one embodiment of the present invention, a TLR agonist is used that is capable of causing a signaling response through TLR-1. Suitably, the TLR agonist capable of causing a signaling response through TLR-1 is selected from: Tri-acylated lipopeptides (LPs); phenol-soluble modulin; *Mycobacterium tuberculosis* LP; S-(2,3-bis(palmitoyloxy)-(2-RS)-propyl)-N-palmitoyl-(R)-Cys-(S)-Ser-(S)-Lys(4)-OH, trihydrochloride (Pam3Cys) LP which mimics the acetylated amino terminus of a bacterial lipoprotein and OspA LP from *Borrelia burgdorferi*.

[0147] In an alternative embodiment, a TLR agonist is used that is capable of causing a signaling response through TLR-2. Suitably, the TLR agonist capable of causing a signaling response through TLR-2 is one or more of a lipoprotein, a peptidoglycan, a bacterial lipopeptide from *M tuberculosis*, *B burgdorferi* or *T pallidum*; peptidoglycans

from species including *Staphylococcus aureus*; lipoteichoic acids, mannuronic acids, *Neisseria* porins, bacterial fimbriae, *Yersina* virulence factors, CMV virions, measles haemagglutinin, and zymosan from yeast.

[0148] In an alternative embodiment, a TLR agonist is used that is capable of causing a signaling response through TLR-3. Suitably, the TLR agonist capable of causing a signaling response through TLR-3 is double stranded RNA (dsRNA), or polyinosinic-polycytidylic acid (Poly IC), a molecular nucleic acid pattern associated with viral infection.

[0149] In an alternative embodiment, a TLR agonist is used that is capable of causing a signaling response through TLR-5. Suitably, the TLR agonist capable of causing a signaling response through TLR-5 is bacterial flagellin.

[0150] In an alternative embodiment, a TLR agonist is used that is capable of causing a signaling response through TLR-6. Suitably, the TLR agonist capable of causing a signaling response through TLR-6 is mycobacterial lipoprotein, di-acylated LP, and phenol-soluble modulin. Additional TLR6 agonists are described in WO 2003/043572.

[0151] In an alternative embodiment, a TLR agonist is used that is capable of causing a signaling response through TLR-7. Suitably, the TLR agonist capable of causing a signaling response through TLR-7 is a single stranded RNA (ssRNA), loxoribine, a guanosine analogue at positions N7 and C8, or an imidazoquinoline compound, or derivative thereof. In one embodiment, the TLR agonist is imiquimod. Further TLR7 agonists are described in WO 2002/085905.

[0152] In an alternative embodiment, a TLR agonist is used that is capable of causing a signaling response through TLR-8. Suitably, the TLR agonist capable of causing a signaling response through TLR-8 is a single stranded RNA (ssRNA), an imidazoquinoline molecule with anti-viral activity, for example resiquimod (R848); resiquimod is also capable of recognition by TLR-7. Other TLR-8 agonists which can be used include those described in WO 2004/071459.

[0153] In an alternative embodiment, a TLR agonist is used that is capable of causing a signaling response through TLR-9. In one embodiment, the TLR agonist capable of causing a signaling response through TLR-9 is HSP90. Alternatively, the TLR agonist capable of causing a signaling response through TLR-9 is bacterial or viral DNA, DNA containing unmethylated CpG nucleotides, in particular sequence contexts known as CpG

motifs. CpG-containing oligonucleotides induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555, WO 99/33488 and U.S. Patent Nos. 6,008,200 and 5,856,462. Suitably, CpG nucleotides are CpG oligonucleotides. Suitable oligonucleotides for use in the immunogenic compositions of the present invention are CpG containing oligonucleotides, optionally containing two or more dinucleotide CpG motifs separated by at least three, suitably at least six or more nucleotides. A CpG motif is a Cytosine nucleotide followed by a Guanine nucleotide. The CpG oligonucleotides of the present invention are typically deoxynucleotides. In a specific embodiment the internucleotide in the oligonucleotide is phosphorodithioate, or suitably a phosphorothioate bond, although phosphodiester and other internucleotide bonds are within the scope of the invention. Also included within the scope of the invention are oligonucleotides with mixed internucleotide linkages. Methods for producing phosphorothioate oligonucleotides or phosphorodithioate are described in US Patent Nos. 5,666,153, 5,278,302 and WO 95/26204.

[0154] Other adjuvants that can be used in immunogenic compositions with a PreF antigens, *e.g.*, on their own or in combination with 3D-MPL, or another adjuvant described herein, are saponins, such as QS21.

[0155] Saponins are taught in: Lacaille-Dubois, M and Wagner H. (1996. A review of the biological and pharmacological activities of saponins. *Phytomedicine* vol 2 pp 363-386). Saponins are steroid or triterpene glycosides widely distributed in the plant and marine animal kingdoms. Saponins are noted for forming colloidal solutions in water which foam on shaking, and for precipitating cholesterol. When saponins are near cell membranes they create pore-like structures in the membrane which cause the membrane to burst. Haemolysis of erythrocytes is an example of this phenomenon, which is a property of certain, but not all, saponins.

[0156] Saponins are known as adjuvants in vaccines for systemic administration. The adjuvant and haemolytic activity of individual saponins has been extensively studied in the art (Lacaille-Dubois and Wagner, *supra*). For example, Quil A (derived from the bark of the South American tree *Quillaja Saponaria Molina*), and fractions thereof, are described in US 5,057,540 and “Saponins as vaccine adjuvants”, Kensil, C. R., *Crit Rev Ther Drug Carrier Syst*, 1996, 12 (1-2):1-55; and EP 0 362 279 B1. Particulate structures, termed Immune Stimulating Complexes (ISCOMS), comprising fractions of Quil A are haemolytic and have been used in the manufacture of vaccines (Morein, B., EP 0 109 942

B1; WO 96/11711; WO 96/33739). The haemolytic saponins QS21 and QS17 (HPLC purified fractions of Quil A) have been described as potent systemic adjuvants, and the method of their production is disclosed in US Patent No.5,057,540 and EP 0 362 279 B1, which are incorporated herein by reference. Other saponins which have been used in systemic vaccination studies include those derived from other plant species such as Gypsophila and Saponaria (Bomford *et al.*, Vaccine, 10(9):572-577, 1992).

[0157] QS21 is an Hplc purified non-toxic fraction derived from the bark of Quillaja Saponaria Molina. A method for producing QS21 is disclosed in US Patent No. 5,057,540. Non-reactogenic adjuvant formulations containing QS21 are described in WO 96/33739. The aforementioned references are incorporated by reference herein. Said immunologically active saponin, such as QS21, can be used in amounts of between 1 and 50 μ g, per human dose of the immunogenic composition. Advantageously QS21 is used at a level of about 25 μ g, for example between 20–30 μ g, suitably between 21–29 μ g or between 22–28 μ g or between 23–27 μ g or between 24–26 μ g, or 25 μ g. In another embodiment, the human dose of the immunogenic composition comprises QS21 at a level of about 10 μ g, for example between 5 and 15 μ g, suitably between 6–14 μ g, for example between 7–13 μ g or between 8–12 μ g or between 9–11 μ g, or 10 μ g. In a further embodiment, the human dose of the immunogenic composition comprises QS21 at a level of about 5 μ g, for example between 1–9 μ g, or between 2–8 μ g or suitably between 3–7 μ g or 4–6 μ g, or 5 μ g. Such formulations comprising QS21 and cholesterol have been shown to be successful Th1 stimulating adjuvants when formulated together with an antigen. Thus, for example, PreF polypeptides can favorably be employed in immunogenic compositions with an adjuvant comprising a combination of QS21 and cholesterol.

[0158] Optionally, the adjuvant can also include mineral salts such as an aluminium or calcium salts, in particular aluminium hydroxide, aluminium phosphate and calcium phosphate. For example, an adjuvant containing 3D-MPL in combination with an aluminium salt (e.g., aluminium hydroxide or “alum”) is suitable for formulation in an immunogenic composition containing a PreF antigen for administration to a human subject.

[0159] Another class of suitable Th1 biasing adjuvants for use in formulations with PreF antigens includes OMP-based immunostimulatory compositions. OMP-based immunostimulatory compositions are particularly suitable as mucosal adjuvants, e.g., for intranasal administration. OMP-based immunostimulatory compositions are a genus of

preparations of outer membrane proteins (OMPs, including some porins) from Gram-negative bacteria, such as, but not limited to, *Neisseria* species (see, e.g., Lowell *et al.*, J. Exp. Med. 167:658, 1988; Lowell *et al.*, Science 240:800, 1988; Lynch *et al.*, Biophys. J. 45:104, 1984; Lowell, in "New Generation Vaccines" 2nd ed., Marcel Dekker, Inc., New York, Basil, Hong Kong, page 193, 1997; U.S. Pat. No. 5,726,292; U.S. Pat. No. 4,707,543), which are useful as a carrier or in compositions for immunogens, such as bacterial or viral antigens. Some OMP-based immunostimulatory compositions can be referred to as "Proteosomes," which are hydrophobic and safe for human use. Proteosomes have the capability to auto-assemble into vesicle or vesicle-like OMP clusters of about 20 nm to about 800 nm, and to noncovalently incorporate, coordinate, associate (e.g., electrostatically or hydrophobically), or otherwise cooperate with protein antigens (Ags), particularly antigens that have a hydrophobic moiety. Any preparation method that results in the outer membrane protein component in vesicular or vesicle-like form, including multi-molecular membranous structures or molten globular-like OMP compositions of one or more OMPs, is included within the definition of Proteosome. Proteosomes can be prepared, for example, as described in the art (see, e.g., U.S. Pat. No. 5,726,292 or U.S. Pat. No. 5,985,284). Proteosomes can also contain an endogenous lipopolysaccharide or lipooligosaccharide (LPS or LOS, respectively) originating from the bacteria used to produce the OMP porins (e.g., *Neisseria* species), which generally will be less than 2% of the total OMP preparation.

[0160] Proteosomes are composed primarily of chemically extracted outer membrane proteins (OMPs) from *Neisseria meningitidis* (mostly porins A and B as well as class 4 OMP), maintained in solution by detergent (Lowell GH. Proteosomes for Improved Nasal, Oral, or Injectable Vaccines. In: Levine MM, Woodrow GC, Kaper JB, Cobon GS, eds, New Generation Vaccines. New York: Marcel Dekker, Inc. 1997; 193-206). Proteosomes can be formulated with a variety of antigens such as purified or recombinant proteins derived from viral sources, including the PreF polypeptides disclosed herein, e.g., by diafiltration or traditional dialysis processes. The gradual removal of detergent allows the formation of particulate hydrophobic complexes of approximately 100-200nm in diameter (Lowell GH. Proteosomes for Improved Nasal, Oral, or Injectable Vaccines. In: Levine MM, Woodrow GC, Kaper JB, Cobon GS, eds, New Generation Vaccines. New York: Marcel Dekker, Inc. 1997; 193-206).

[0161] "Proteosome: LPS or Protollin" as used herein refers to preparations of protosomes admixed, *e.g.*, by the exogenous addition, with at least one kind of lipo-polysaccharide to provide an OMP-LPS composition (which can function as an immunostimulatory composition). Thus, the OMP-LPS composition can be comprised of two of the basic components of Protollin, which include (1) an outer membrane protein preparation of Proteosomes (*e.g.*, Projuvant) prepared from Gram-negative bacteria, such as *Neisseria meningitidis*, and (2) a preparation of one or more liposaccharides. A lipo-oligosaccharide can be endogenous (*e.g.*, naturally contained with the OMP Proteosome preparation), can be admixed or combined with an OMP preparation from an exogenously prepared lipo-oligosaccharide (*e.g.*, prepared from a different culture or microorganism than the OMP preparation), or can be a combination thereof. Such exogenously added LPS can be from the same Gram-negative bacterium from which the OMP preparation was made or from a different Gram-negative bacterium. Protollin should also be understood to optionally include lipids, glycolipids, glycoproteins, small molecules, or the like, and combinations thereof. The Protollin can be prepared, for example, as described in U.S. Patent Application Publication No. 2003/0044425.

[0162] Combinations of different adjuvants, such as those mentioned hereinabove, can also be used in compositions with PreF antigens. For example, as already noted, QS21 can be formulated together with 3D-MPL. The ratio of QS21 : 3D-MPL will typically be in the order of 1 : 10 to 10 : 1; such as 1:5 to 5 : 1, and often substantially 1 : 1. Typically, the ratio is in the range of 2.5 : 1 to 1 : 1 3D-MPL: QS21. Another combination adjuvant formulation includes 3D-MPL and an aluminium salt, such as aluminium hydroxide. When formulated in combination, this combination can enhance an antigen-specific Th1 immune response.

[0163] In some instances, the adjuvant formulation a mineral salt, such as a calcium or aluminium (alum) salt, for example calcium phosphate, aluminium phosphate or aluminium hydroxide. Where alum is present, *e.g.*, in combination with 3D-MPL, the amount is typically between about 100 μ g and 1mg, such as from about 100 μ g, or about 200 μ g to about 750 μ g, such as about 500 μ g per dose.

[0164] In some embodiments, the adjuvant includes an oil and water emulsion, *e.g.*, an oil-in-water emulsion. One example of an oil-in-water emulsion comprises a metabolisable oil, such as squalene, a tocol such as a tocopherol, *e.g.*, alpha-tocopherol, and a surfactant, such as sorbitan trioleate (Span 85TM) or polyoxyethylene sorbitan

monooleate (Tween 80TM), in an aqueous carrier. In certain embodiments, the oil-in-water emulsion does not contain any additional immunostimulants(s), (in particular it does not contain a non-toxic lipid A derivative, such as 3D-MPL, or a saponin, such as QS21). The aqueous carrier can be, for example, phosphate buffered saline. Additionally the oil-in-water emulsion can contain span 85 and/or lecithin and/or tricaprylin.

[0165] In another embodiment of the invention there is provided a vaccine composition comprising an antigen or antigen composition and an adjuvant composition comprising an oil-in-water emulsion and optionally one or more further immunostimulants, wherein said oil-in-water emulsion comprises 0.5-10 mg metabolisable oil (suitably squalene), 0.5-11 mg tocol (suitably a tocopherol, such as alpha-tocopherol) and 0.4-4 mg emulsifying agent.

[0166] In one specific embodiment, the adjuvant formulation includes 3D-MPL prepared in the form of an emulsion, such as an oil-in-water emulsion. In some cases, the emulsion has a small particle size of less than 0.2 μ m in diameter, as disclosed in WO 94/21292. For example, the particles of 3D-MPL can be small enough to be sterile filtered through a 0.22micron membrane (as described in European Patent number 0 689 454).

Alternatively, the 3D-MPL can be prepared in a liposomal formulation. Optionally, the adjuvant containing 3D-MPL (or a derivative thereof) also includes an additional immunostimulatory component.

[0167] The adjuvant is selected to be safe and effective in the population to which the immunogenic composition is administered. For adult and elderly populations, the formulations typically include more of an adjuvant component than is typically found in an infant formulation. In particular formulations using an oil-in-water emulsion, such an emulsion can include additional components, for example, such as cholesterol, squalene, alpha tocopherol, and/or a detergent, such as tween 80 or span85. In exemplary formulations, such components can be present in the following amounts: from about 1-50mg cholesterol, from 2 to 10% squalene, from 2 to 10% alpha tocopherol and from 0.3 to 3% tween 80. Typically, the ratio of squalene: alpha tocopherol is equal to or less than 1 as this provides a more stable emulsion. In some cases, the formulation can also contain a stabilizer.

[0168] When an immunogenic composition with a PreF polypeptide antigen is formulated for administration to an infant, the dosage of adjuvant is determined to be effective and

relatively non-reactogenic in an infant subject. Generally, the dosage of adjuvant in an infant formulation is lower (for example, the dose may be a fraction of the dose provided in a formulation to be administered to adults) than that used in formulations designed for administration to adult (e.g., adults aged 65 or older). For example, the amount of 3D-MPL is typically in the range of 1 μ g-200 μ g, such as 10-100 μ g, or 10 μ g-50 μ g per dose. An infant dose is typically at the lower end of this range, e.g., from about 1 μ g to about 50 μ g, such as from about 2 μ g, or about 5 μ g, or about 10 μ g, to about 25 μ g, or to about 50 μ g. Typically, where QS21 is used in the formulation, the ranges are comparable (and according to the ratios indicated above). In the case of an oil and water emulsion (e.g., an oil-in-water emulsion), the dose of adjuvant provided to a child or infant can be a fraction of the dose administered to an adult subject. A demonstration of the efficacy of immunogenic compositions containing a PreF antigen in combination various doses of an exemplary oil-in-water adjuvant is provided in Example 9.

[0169] An immunogenic composition typically contains an immunoprotective quantity (or a fractional dose thereof) of the antigen and can be prepared by conventional techniques. Preparation of immunogenic compositions, including those for administration to human subjects, is generally described in Pharmaceutical Biotechnology, Vol.61 Vaccine Design-the subunit and adjuvant approach, edited by Powell and Newman, Plenum Press, 1995. New Trends and Developments in Vaccines, edited by Voller *et al.*, University Park Press, Baltimore, Maryland, U.S.A. 1978. Encapsulation within liposomes is described, for example, by Fullerton, U.S. Patent 4,235,877. Conjugation of proteins to macromolecules is disclosed, for example, by Likhite, U.S. Patent 4,372,945 and by Armor *et al.*, U.S. Patent 4,474,757.

[0170] Typically, the amount of protein in each dose of the immunogenic composition is selected as an amount which induces an immunoprotective response without significant, adverse side effects in the typical subject. Immunoprotective in this context does not necessarily mean completely protective against infection; it means protection against symptoms or disease, especially severe disease associated with the virus. The amount of antigen can vary depending upon which specific immunogen is employed. Generally, it is expected that each human dose will comprise 1-1000 μ g of protein, such as from about 1 μ g to about 100 μ g, for example, from about 1 μ g to about 50 μ g, such as about 1 μ g, about 2 μ g, about 5 μ g, about 10 μ g, about 15 μ g, about 20 μ g, about 25 μ g, about 30 μ g, about

40 μ g, or about 50 μ g. The amount utilized in an immunogenic composition is selected based on the subject population (e.g., infant or elderly). An optimal amount for a particular composition can be ascertained by standard studies involving observation of antibody titres and other responses in subjects. Following an initial vaccination, subjects can receive a boost in about 4 weeks.

[0171] It should be noted that regardless of the adjuvant selected, the concentration in the final formulation is calculated to be safe and effective in the target population. For example, immunogenic compositions for eliciting an immune response against RSV in humans are favorably administered to infants (e.g., infants between birth and 1 year, such as between 0 and 6 months, at the age of initial dose). Immunogenic compositions for eliciting an immune response against RSV are also favorably administered to elderly humans (e.g., alone or in a combination with antigens of other pathogens associated with COPD). It will be appreciated that the choice of adjuvant can be different in these different applications, and the optimal adjuvant and concentration for each situation can be determined empirically by those of skill in the art.

[0172] In certain embodiments, the immunogenic compositions are vaccines that reduce or prevent infection with RSV. In some embodiments, the immunogenic compositions are vaccines that reduce or prevent a pathological response following infection with RSV. Optionally, the immunogenic compositions containing a PreF antigen are formulated with at least one additional antigen of a pathogenic organism other than RSV. For example, the pathogenic organism can be a pathogen of the respiratory tract (such as a virus or bacterium that causes a respiratory infection). In certain cases, the immunogenic composition contains an antigen derived from a pathogenic virus other than RSV, such as a virus that causes an infection of the respiratory tract, such as influenza or parainfluenza. In other embodiments, the additional antigens are selected to facilitate administration or reduce the number of inoculations required to protect a subject against a plurality of infectious organisms. For example, the antigen can be derived from any one or more of influenza, hepatitis B, diphtheria, tetanus, pertussis, *Hemophilus influenza*, poliovirus, *Streptococcus* or *Pneumococcus*, among others.

[0173] Accordingly, the use of PreF antigens or nucleic acids that encode them in the preparation of a medicament for treating (either therapeutically following or prophylactically prior to) exposure to or infection by RSV is also a feature of this

disclosure. Likewise, methods for eliciting an immune response against RSV in a subject are a feature of this disclosure. Such methods include administering an immunologically effective amount of a composition comprising a PreF antigen to a subject, such as a human subject. Commonly, the composition includes an adjuvant that elicits a Th1 biased immune response. The composition is formulated to elicit an immune response specific for RSV without enhancing viral disease following contact with RSV. That is, the composition is formulated to and results in a Th1 biased immune response that reduces or prevents infection with a RSV and/or reduces or prevents a pathological response following infection with a RSV. Although the composition can be administered by a variety of different routes, most commonly, the immunogenic compositions are delivered by an intramuscular or intranasal route of administration.

[0174] An immunogenic composition typically contains an immunoprotective quantity (or a fractional dose thereof) of the antigen and can be prepared by conventional techniques. Preparation of immunogenic compositions, including those for administration to human subjects, is generally described in Pharmaceutical Biotechnology, Vol.61 Vaccine Design-the subunit and adjuvant approach, edited by Powell and Newman, Plenum Press, 1995. New Trends and Developments in Vaccines, edited by Voller *et al.*, University Park Press, Baltimore, Maryland, U.S.A. 1978. Encapsulation within liposomes is described, for example, by Fullerton, U.S. Patent 4,235,877. Conjugation of proteins to macromolecules is disclosed, for example, by Likhite, U.S. Patent 4,372,945 and by Armor *et al.*, U.S. Patent 4,474,757.

[0175] Typically, the amount of protein in each dose of the immunogenic composition is selected as an amount which induces an immunoprotective response without significant, adverse side effects in the typical subject. Immunoprotective in this context does not necessarily mean completely protective against infection; it means protection against symptoms or disease, especially severe disease associated with the virus. The amount of antigen can vary depending upon which specific immunogen is employed. Generally, it is expected that each human dose will comprise 1-1000 μ g of protein, such as from about 1 μ g to about 100 μ g, for example, from about 1 μ g to about 50 μ g, such as about 1 μ g, about 2 μ g, about 5 μ g, about 10 μ g, about 15 μ g, about 20 μ g, about 25 μ g, about 30 μ g, about 40 μ g, or about 50 μ g. The amount utilized in an immunogenic composition is selected based on the subject population (*e.g.*, infant or elderly). An optimal amount for a

particular composition can be ascertained by standard studies involving observation of antibody titres and other responses in subjects. Following an initial vaccination, subjects can receive a boost in about 4-12 weeks. For example, when administering an immunogenic composition containing a PreF antigen to an infant subject, the initial and subsequent inoculations can be administered to coincide with other vaccines administered during this period.

[0176] The following examples are provided to illustrate certain particular features and/or embodiments. These examples should not be construed to limit the invention to the particular features or embodiments described.

EXAMPLES

Example 1: Exemplary PreF antigens

[0177] The PreF antigen was modified as compared to a native RSV F protein in order to stabilize the protein in its prefusion conformation, based on the prediction that an immune response generated to the prefusion conformation of F would preferentially include antibodies that would prevent binding, conformation shifting and/or other events involved in membrane fusion, thereby increasing the efficacy of the protective response.

[0178] FIG. 1A and B schematically illustrate features of RSV F0 and exemplary PreF recombinant antigens. FIG. 1A is a representation of the RSV F0 protein. F0 is a pre-protein consisting of 574 amino acids. The F0 pre-protein is proteolytically processed and glycosylated following translation. A signal peptide, which is later removed by a signal peptidase, targets translation of the F0 pre-protein to the reticulum endoplasmic (RE). Nascent peptide in the RE is then N-glycosylated at multiple sites (represented by triangles). Furin cleavage of F0 generates F2 and F1 peptide domains, which are folded and assembled together as a trimer of F2-F1 heterodimers (that is, 3 times F2-F1). In its native state, the F protein is anchored to the membrane by a transmembrane helix in the C-terminal region. Additional features of the F0 polypeptide include, 15 Cysteine residues, 4 characterized neutralizing epitopes, 2 coiled-coil regions, and a lipidation motif. FIG. 1B illustrates features of exemplary PreF antigens. To construct the PreF antigen, the F0 polypeptide was modified to stabilize the prefusion conformation of the F protein, thereby retaining the predominant immunogenic epitopes of the F protein as presented by the RSV virus prior to binding to and fusion with host cells. The following stabilizing mutations were introduced into the PreF antigen relative to the F0 polypeptide. First, a stabilizing coiled-coil domain was placed at the C-terminal end of the extracellular domain of the F0 polypeptide, replacing the membrane anchoring domain of F0. Second, the pep27 peptide (situated between the F2 and F1 domains in the native protein) was removed. Third, both furin motifs were eliminated. In alternative embodiments (designated PreF_V1 and PreF_V2), an immunologically active portion (e.g., amino acids 149-229) of the RSV G protein was added to the C-terminal domain.

[0179] In other embodiments, modifications were introduced to alter (increase or decrease) glycosylation and/or to reduce cleavage by a protease other than furin.

Example 2: Production and Purification of PreF recombinant protein from CHO cells.

[0180] A recombinant polynucleotide sequence encoding an exemplary PreF antigen was introduced into host CHO cells for the production of PreF antigen. Transiently transfected host cells or expanded stable populations comprising the introduced polynucleotide sequence were grown in medium and under conditions suitable for growth at an acceptable scale for the desired purpose (e.g., as generally described in Freshney (1994) Culture of Animal Cells, a Manual of Basic Technique, third edition, Wiley- Liss, New York and the references cited therein). Typically, the cells were grown in serum-free medium in shake flasks at 37°C with 5% CO₂ and passaged at 2-3 day intervals, or in bioreactors at 29°C with pO₂ maintained at 20%.

[0181] To recover recombinant PreF antigen, the cell culture was centrifuged and the cell culture supernatant stored at about -80°C until further use. For further analysis, two liter aliquots of cell culture supernatant were diluted 2x with purified water and adjusted to pH 9.5 with NaOH. The supernatant was loaded at a rate of 14 ml/min. onto a Q Sepharose FF ion exchange column (60 ml, 11.3 cm), equilibrated in 20 mM piperazine pH 9.5. After washing the column with the starting buffer, elution was performed with a NaCl gradient from 0 to 0.5 M NaCl in 20 column volumes (fraction size 10 ml). Fractions were analyzed on SDS PAGE gel by silver staining and western blot. Fractions containing substantial PreF protein were then pooled prior to further processing.

[0182] The pooled elution of the Q step (~130 mls) was subjected to buffer exchange into 10 mM phosphate, pH 7.0 using the bench-scale TFF system from Millipore with the Pelllicon XL PES Biomax 100 (MWCO 10,000 Da) membrane cassette. The resulting material had a pH of 7.0 and a conductivity of 1.8 mS/cm. 100 ml of this sample was loaded at 5 ml/min. on a 10 ml Hydroxyapatite Type II (HA TII) gel (XK 16, height=5cm) equilibrated with 10 mM PO₄ (Na) buffer pH 7.0. After washing the column with the starting buffer, elution was performed with a gradient from 10 mM to 200 mM PO₄ (Na) pH 7.0 in 20 column volumes. Fractions were again analysed on SDS PAGE with silver staining and coomassie blue, and the positive fractions were pooled.

[0183] Following affinity chromatography, the pooled fractions were concentrated and the buffer exchanged into DPBS (pH ~7.4) using a Vivaspin 20 concentrator unit, 10,000 Da MWCO. The final product was about 13 ml. Protein concentration was 195 µg/ml,

assessed by Lowry assay. Purity was greater than 95%. This purified PreF antigen preparation was filter sterilized and stored at -20°C prior to use.

Example 3: Characterization of the PreF recombinant protein produced in CHO cells.

[0184] PreF recombinant protein produced in CHO cells was characterized by asymmetrical field flow fractionation (AFF-MALS) and compared to a chimeric antigen including RSV F and G protein components. AFF-MALS allows separation of protein species according to their molecular size in a liquid flow with minimal matrix interaction and further analysis by multi-angle light scattering for accurate molecular weight determination. FIG. 2A shows that more than 65% of purified FG material is found as high molecular weight oligomers (1000–100 000 KDa) in final PBS buffer while 3% remain in monomeric form.

[0185] FIG. 2B shows that the purified PreF protein is folded in its trimeric form to a proportion of 73% in PBS buffer. 10% of the material is found as 1000 to 20 000 KDa oligomers. These results indicate that the recombinant PreF protein expressed in CHO cells is folded as a trimer as predicted for the native state.

[0186] Purified PreF protein was also crosslinked with glutaraldehyde for the double purpose of confirming the soluble nature of the protein in phosphate buffer solution and of generating aggregates for comparative in vivo evaluation with FG protein (see Example 7 below). Glutaraldehyde crosslinking is known for providing a good assessment of the quaternary structure of a protein, and is described in (*Biochemistry*, 36:10230-10239 (1997); *Eur. J. Biochem.*, 271:4284-4292 (2004)).

[0187] Protein was incubated with 1%, 2% and 5% of glutaraldehyde crosslinking agent for four hours at 4°C and the reaction was blocked by addition of NaBH₄. Excess glutaraldehyde was removed by column desalting in PBS buffer. Resulting protein was quantified by absorbance at 280nm and evaluated by SDS PAGE in denaturing and reducing condition. The majority of purified recombinant PreF was determined to migrate as a trimer in PBS solution. Increasing the incubation temperature to 23°C was required to convert majority of trimeric protein to high molecular weight aggregates, as confirmed by SDS PAGE.

Example 4: *In vitro* Neutralization Inhibition by the PreF antigen

[0188] Human sera obtained from volunteers were screened for reactivity against RSV A by ELISA and used in the neutralization inhibition (NI) assay at relevant dilution based on prior RSV neutralization potential titration established for each serum sample. Briefly, Sera were mixed with inhibitor proteins (PreF or a control protein corresponding essentially to the chimeric FG disclosed in US Patent No. 5,194,595, designated RixFG) at concentrations of 25 μ g/ml in DMEM with 50% 199-H medium, with 0.5% FBS, 2 mM glutamine, 50 μ g/ml genamycin (all Invitrogen), and incubated 1.5 to 2 hours at 37°C on a rotating wheel. 20 μ l the serial dilutions of sera and proteins were mixed in a round bottom 96-well plate with a RSV A titred to optimize the range of inhibition for each serum sample. The resulting mixtures were incubated for 20 minutes at 33°C under 5% CO₂ to maintain pH.

[0189] The sera-inhibitor-virus mixtures were then placed into previously Vero cell-seeded flat bottom 96-well plates and incubated for 2 hours at 33°C prior to addition of 160 μ l of medium. The plates were further incubated for 5-6 days at 33°C with 5% CO₂ until immunofluorescence assay for NI titer detection. Following fixation for 1 hour with 1% paraformaldehyde in phosphate buffered saline (PBS), plates were blocked with 2% milk/PBS and Block biffer. Goat anti-RSV antibody (Biodesign Internation; 1:400) was added to each well without rinsing and incubated for 2 hours at room temperature (RT). Samples were washed 2x with PBS and anti-goat IgG-FITC (Sigma; 1:400) in blocking buffer was added to the wells. The plates were again incubated for 2 hours at RT and washed 2x as above prior to reading. A well was considered positive when \geq 1 fluorescent syncytium was detected. The 50% tissue cultue infective dose (TCID₅₀) calculations were performed using the Spearman-Karber (SK) method and percentages of NI calculated as follow: [(Neut titer of 0 μ g/ml inhibitor – Neut titer of 25 μ g/ml inhibitor)/Neut titer of 0 μ g/ml inhibitor] X 100. Exemplary results shown in FIG. 3 indicate that PreF is superior to FG in NI in 16/21 donors tested.

Example 5: PreF antigen is immunogenic

[0190] To demonstrate immunogenicity of the PreF antigen, mice were immunized twice IM at two weeks interval with preF (6.5, 3.1, 0.63, 0.13, and 0.025 μ g/ml) and a Th1 adjuvant containing 3D MPL and QS21 at 1/20 of human dose (“AS01E”) or preF (1, 0.2

and 0.04 µg/ml) and a Th1 adjuvant containing 3D MPL and alum 1/10 of human dose (“AS04C”) and serum was collected three weeks later.

[0191] Antigen-specific IgG antibody titers were determined on pooled serum samples by ELISA according to standard procedures. Briefly, 96-well plates were coated with purified inactivated RSV A, RSV B and homologous preF protein and incubated overnight at 4°C. Serum samples were serially diluted in blocking buffer starting at an initial concentration of 1:50, along with purified mouse IgG (Sigma, ON) at starting concentrations of 200 ng/ml and incubated for 2h at room temperature. Bound antibody was detected with horseradish peroxidase (HRP)-conjugated anti-mouse IgG (Sigma, ON). 3,3A,5,5A-tetramethylbenzidine (TMB, BD Opt EIA/TM, BD Biosciences, ON) was used as the substrate for HRP. 50µl of 1M H₂SO₄ was added to each well to stop the reaction. Absorbance values for each well were detected at 450 nm with a Molecular Devices microplate reader (Molecular Devices, USA).

[0192] Representative results detailed in FIGS. 4A and 4B show that strong titers are elicited against both RSV A and RSV B following immunization with preF antigen.

Example 6: PreF elicits neutralizing antibodies

[0193] The presence and quantity of neutralizing antibodies was assessed in serum samples of mice immunized as described above in Example 5. Pooled sera from immunized animals were serially diluted from a starting dilution of 1:8 in RSV medium in 96-well plates (20 µl/well). Control wells contained RSV medium only, or goat anti-RSV antibody at 1:50 (Biodesign international). 500-1000 infectious doses of a representative RSV A or B strain were added to the wells, and the plates were incubated for 20 minutes at 33°C, 5% CO₂, before the mixture was transferred to 96-well flat-bottomed plates previously seeded with 1x10⁵ cells/mL Vero cells. Cells were incubated for approximately 2 hours at 33°C, 5% CO₂ and refed, prior to incubation for 5-6 days at the same temperature. Supernatants were removed; plates were washed with PBS and adhering cells fixed with 1% paraformaldehyde in PBS for 1 hour, followed by indirect immunofluorescence (IFA) using a goat anti RSV primary antibody and anti-goat IgG-FITC for detection.

[0194] Representative results shown in FIGS 5A and 5B, respectively, demonstrate that significant neutralizing antibodies against both RSV strains are detected in sera of animals immunized with preF.

Example 7: PreF protects against RSV challenge

[0195] Mice were immunized twice IM at a two week interval as described above, and challenged three weeks after the second injection with RSV A. Protection against RSV was evaluated by measuring the virus present in lungs following challenge. In brief, lungs from immunized animals were aseptically removed following euthanasia and washed in RSV medium using 2 volumes of 10 ml/lung in 15 ml tubes. Lungs were then weighed and homogenized individually in RSV medium with an automated Potter homogenizer (Fisher, Nepean ON), and centrifuged at 2655 x g for 2 minutes at 4°C. The virus present in the supernatants was titered by serial dilution (eight replicates starting at 1:10) on a previously seeded Vero cell (ATCC# CCL-81) monolayer in 96-well plates and incubated for 6 days. RSV was detected by indirect IFA following fixing in 1% paraformaldehyde/PBS pH7.2, with goat anti-RSV primary antibody and FITC labeled anti-goat IgG secondary antibody.

[0196] Representative results shown in FIGS. 6A and B demonstrate that doses equal to or higher than 0.04 μ g when given in presence of adjuvant elicit strong protection against RSV.

Example 8: PreF does not induce pulmonary eosinophil recruitment following challenge

[0197] To assess the potential of the PreF antigen to provoke exacerbated disease following immunization and subsequent challenge, groups of mice (5 mice/group) were immunized twice each with (a) 10 μ g gluteraldehyde-treated preF, (b) 10 μ g preF or (c) 10 μ g FG without adjuvant. Mice were challenged with RSV A 3 weeks post-boost and bronchoalveolar lavage (BAL) was performed 4 days post challenge. Total leukocyte infiltrates in BAL were enumerated per mouse as well as differential enumerations (300 cells) based on cell morphology of macrophages/monocytes, neutrophils, eosinophils and lymphocytes.

[0198] Total cell numbers were multiplied by differential percentages of eosinophils for each animal. Represented are geometric means per group with 95% confidence limits. Representative results shown in FIG. 7 demonstrate that eosinophils are not recruited to the lungs following immunization with preF and challenge. Furthermore, these results suggest that the soluble nature of the preF antigen, as compared to a deliberately aggregated form of preF (gluteraldehyde treatment) or FG antigen (naturally aggregated) does not favour eosinophils.

Example 9: Immunogenicity of PreF antigen formulated with dilutions of an oil-in-water emulsion adjuvant.

[0199] Mice received 250ng preF formulated with the exemplary oil-in-water adjuvant, AS03A, at a 1/10 of the human “full” dose (AS03A) of 10.70 mg squalene, 11.88 mg DL- α -tocopherol, 4.85 mg polysorbate 80, 1/2 dose (AS03B), 1/4 dose (AS03C), or no adjuvant. Control mice received AS03A alone or PBS. The mice were immunized on days 0 and 14. Blood, splenocyte collection and challenge were performed on Day 39 (25 days post dose 2). Lungs were homogenized for titration of RSV 4 days post challenge.

[0200] Antigen-specific IgG antibody titers were determined on individual serum samples by ELISA. Briefly, 96-well plates were coated with purified inactivated RSV A and incubated overnight at 4°C. Serum samples were serially diluted in blocking buffer starting at 1:200, along with purified mouse IgG (Sigma, ON), at starting concentrations of 200 ng/ml and incubated for 2h at 37°C. Bound antibody was detected with horseradish peroxidase (HRP)-conjugated anti-mouse IgG (Sigma, ON). 3,3A,5,5A-tetramethylbenzidine (TMB, BD Opt EIATM, BD Biosciences, ON) was used as the substrate for HRP. 50 μ l of 1M H₂SO₄ was added to each well to stop the reaction. Absorbance values for each well were detected at 450 nm with a Molecular Devices microplate reader (Molecular Devices, USA). Results are illustrated in FIG. 8. Anti- RSV IgG concentration of approximately 250,000 ng/mL was observed in serum from mice immunised with preF in combination with AS03A, AS03B or AS03C, whereas very little (1828 ng/ml) was observed in serum from mice immunised with preF alone.

[0201] Pooled sera from immunized animals were serially diluted from a starting dilution of 1:8 in RSV medium in 96-well plates (20 μ l/well). Control wells contained RSV medium only or goat anti-RSV antibody at 1:50 (Biodesign international). RSV Long strain was added, the plates were incubated for 20 minutes at 33°C and the mixture was transferred to 96-well flat-bottomed plates previously seeded with 1x10⁵ cells/mL Vero cells. After 5-6 days at the same temperature, supernatants were removed; plates were washed with PBS and adhering cells fixed with 1% paraformaldehyde in PBS for 1 hour, followed by indirect immunofluorescence (IFA). FIG. 9 shows that regardless of the amount of AS03 administered in combination with preF the RSV neutralising antibody titres remained the same (~ 11 log2), whereas the RSV neutralising antibody titre in the mice immunised without AS03 were significantly lower (~6 log2).

[0202] Lungs from immunized animals were aseptically removed following euthanasia and washed in RSV medium using 2 volumes of 10 ml / lung in 15 ml tubes. They were then weighed and homogenized individually in RSV medium with an automated Potter homogenizer (Fisher, Nepean ON), and centrifuged at 2655 x g for 2 minutes at 4°C. The virus present in the supernatants was titered. Briefly, lung homogenates were serially diluted in eight replicates starting at a 1:10 on a previously seeded Vero cell (ATCC# CCL-81) monolayer in 96-well plates and incubated for 5-6 days. RSV was detected by indirect IFA. FIG. 10 illustrates results demonstrating that no difference in protection was observed between mice immunised with preF in combination with AS03A, AS03B or AS03C, whereas less protection was observed in mice immunised in absence of AS03.

[0203] These results demonstrate that the PreF antigen can be formulated across a broad range of adjuvant concentrations to produce a composition that elicits an immune response against RSV.

Sequence Listing

SEQ ID NO:1

Nucleotide sequence encoding RSV reference Fusion protein
Strain A2 GenBank Accession No. U50362

SEQ ID NO:2

Amino acid sequence of RSV reference F protein precursor F0
Strain A2 GenBank Accession No. AAB86664

MELLILKANAITTILTAVTFCFASGQNITEEFYQSTCSAVSKGYLSALRTGWYTSVITIELSNIKENKCNGT
DAKVKLICKQELDKYKNAVTELQILMQSTPATNNRARRELPRFMNYTLNNAKTKNTVLSKKRKRRLFGFLGIV
GSAIASGVAVSKVLHLEGEVNKIKSALLSTNKAVVSLSGNVSVLTSKVLDLKNYIDKQLLPVNQKQSCSISN
IATVIEFQQKNNRLLITREFSVNAGVTPVSTYMLTNSELLSLLINDMPITNDQKKLMSNNVQIVRQQSYI
MSIIKEEVLAYVQQLPLYGVIDTPCWKLHTSPLCNTNTKEGSNICLRTDRGWYCDNAGSVSFFPQAETCKV
QSNRVCDFTMNSLTLPSSEVNLCNVDIFNPKYDCKIMTSKTDVSSSVITSLGIAVSCYGKTKTASNKNRGII
KTFSNGCDYVSNKGVDTSVGNTLYYVNQEGKSLYVKEPEIINFYDPLVFPSSDEFDASISQVNEKINQSLA
FIRKSDETLHNVNAGKSTINIMITTIIIVIIVILLSIIAVGLLILYCKARSTPVTLISKDOLSGINNIAFESN

SEQ ID NO:3

Nucleotide sequence encoding RSV reference G protein
Strain Long

SEQ ID NO:4

Amino acid sequence of RSV reference G protein
 MSKNKDQRTAKTLEKTWDTLNHLLFISSGLYKLNLKSTIAQITLSILAMIISTSLIITAIIFIASANHKVTLT
 TAIIQDATSQIKNTTPTYLTQDPQLGISFSNLSEITSQTTILASTTPGVKSNLQPTTVKTKNTTTQTQPS
 KPTTKQRQNKPNNKPNNDHFEVNFVPCSICSNNPTCWAICKRIPNKKPGKTTKPTKKPTFKTTKDLK
 PQTTKPKEVPTTKPTEPTINTTKTNITTLTNNTTGNPKLTSQMFHSTSSEGNLSPSQVSTSEHPSQ
 PSSPPNTRQ

Seq ID NO:5

Nucleotide sequence of PreF analog optimized for CHO
 aagcttgcaccatggagctgctgatcctgaaaaccaacgcacatcaccgcacatcctggccgtgaccctg
 tgcttcgcctcccccagaacatcaccggaggagtctaccatgcaccatgcgttccgcgtgtccaaacatcaaggaaaacaag
 tgcaacggcaccgacgccaaggtgaagctgatcaagcaggagctggacaagaatacaagagcgcgtgaccgg
 ctccagctgtatgcagttccacccctgcaccaacaacaaggttctgggttcctgtgtggcgtggcgtgg
 gccatgcgcctccggcatggccgtgagaagggtctgcacctggagggtgaacaagatcaagagcgc
 ctgctgtccaccaacaaggccgtgtgtccctgtccacccgtgtccgtgtccgtgtaccccaaggtgtgg
 ctgaagaactacatgcacaaggcagctgtccatgtgtaccaagcagttctgtccatctcaacatcgag
 accgtgatcgagttccacccctgtgtccacccatgtgtccacactccgcagatcaccgcgtgtccgtgt
 gtgaccacccctgtgtccacccatgtgtccacactccgcagatcaccgcgtgtccgtgtccgtgt
 accaacgcggaaaaaaactgtatgtccaccaacacgtgcagatcgtgcggcagcgtccatcagcatcatgag
 atcatcaaggaagagggtgtggctacgtggcgtacgtggcgtgtacccgtgtccacccctgtgg
 aagctgcacacccctcccccgtgtccaccaacaacccggctgtggagatcaccgcgtgtccgtgt
 cgggctggactgtgcacaacgcggctccgtgtcccttctccctgtggccgagacccgtcaagggtgcagtc
 aaccgggtttctgcacccatgtacccctgcacccctgcacccgtgtacccctgtgt
 ttcaaccccaagtgactgactgcaagatcatgaccaggcagatcgtgtccctccagcgtgt
 ggcgcacatcgatcgactacgtccacccctgtgtccaccaacaaggagggtccacatctgc
 ttctccaaacggctgtgcactacgtgtccaccaataaggcggtggacccgtgtccgtgg
 gtgaaataaggcaggaggcaagaggcctgtacgtgtacccgtgtacccctgtgt
 ttccctccgcacgcgttgcacccctccatcagccaggtaacccatcactgaccctctgg
 cggaagtccgcacgcgtacgtccacccctgtgtacccgtgtacccctgtgt
 gagaacgcgatcgcggatcaagaagctgatcgccgaggcgtataatctaga

SEQ ID NO:6

Amino acid sequence of PreF analog
 MELLILKTNAITAILAAVTLCFASSQNITEEFYQSTCSAVSKGYLSALRTGWYTSVITIELSNIKENKNGT
 DAKVKLIKQELDKYKSAVTTELQLLMQSTPATNNKFLGFLLGVGSATASGIAVSKVHLGEVNKIKSALLST
 NKAIVSLSNGSVLTSKVLSDLKNYIDKQLLPIVNKQSCSISNIETVIEFQQKNNRLEITREFSVNAGVTP
 VSTYMLTNTSELLSLLNDMPITNDQKKLMSNNVQIVRQQSYSIMSIKEEVLAYVQPLYGVIDTPCWKLHT
 SPLCTTNTKEGSNICLRTDRGWYCDNAGSVSFFPLAETCKVQSNRVCDFTMNSLTPSEVNLCNIDIFNPK
 YDCKIMTSKTDVSSVITSLGAIYSCYGKTKCTASNKNRGIIKTFNSNGCDYVSNKGVDTVSVGNLTYVNQ
 EGKSLYVKGEPIINFYDPLVFPSEFDASISQVNEKINQSLAFIRKSDEKLHNVEDKIEEILSKIYHIENEI
 ARIKLLIGEA

Seq ID NO:7

Nucleotide sequence encoding PreFG_V1 optimized for CHO
 aagcttgcaccatggagctgctgatcctcaagaccaacgcacatcaccgcacatcctggccgtgaccctg
 tgcttcgcctcccccagaacatcaccggagggtctaccatgcaccatgcgttccgcgtgtccaaagggtac
 ctgcttcgcctcccccgtgtggacccctgcgtgtccatgtgtccgtgtccgtgt
 tgcaacggcaccgacgccaaggtcaagctgatcaagcaggaaatggacaatcaaggatcaaggcgcgtgaccgg
 ctccagctgtatgcagttccacccctgcaccaacaacaaggaaatgttctgggttcctgtgtggcgtgg
 tccgcacatcgccatccggatcgccgtgtccacccctgcaccaacaaggccgtgtccgtgt
 gatctgaagaactacatcgacccatgtgtccatgtgtccatgtgt
 gagaccgtgtccacccctgtgtccacccctgcaccaacaaggccgtgt
 ggcgtgaccacccctgtgtccacccatgtgtccacccctgcaccaacaaggccgtgt
 atcaccacgcacccaaaaaaagctgtatgtccaccaacgtgcacatgtgt
 agcatcatcaaggaagaatgtccgtgtccacccctgcaccaacaaggccgtgt
 tggaaatgtgtccacccctgcaccaacaaggccgtgt
 gaccggggctggactgtgtccacccctgcaccaacaaggccgtgt
 gaccggggctggactgtgtccacccctgcaccaacaaggccgtgt

tccaaaccgggtgttctgcgacaccatgaactccctgaccctgccttccgaggtaacctgtgcaacatcgac
atcttcaccccaagtacgactgcaagatcatgaccaggcaagacgcgtgtccctccagcgtgatcacctcc
ctggggccatcgtgtccctgtacggcaagaccaactgcaccgcctccaacaagaaccggggaatcatcaag
accttcaccccaacggctgcgactacgtgtccataaaggcgtggacaccgtgtccgtggcaacacactgtac
tacgtgaataagcaggaaggcaagagacgtgtacgtgaagggcggacgcctatcatcaacttctacgaccctctg
gtgtccctccgacgagttcgcacgcctccatcagccaggtaacgagaagatcaaccaggccctggccct
atccggaaactccgacgagaagctgcataacgtggaggacaagatcgaagagatcctgtccaaaatctaccac
atcgagaacgagatcgcggatcaagaagactgtacgcggcggctctggccggcagcggcggctcc
aagcagcggcagaacaagcctctaacaagcccaacaacgacttccacttcgaggtgttcaacttcgtgct
tgctccatctgtccataacaaccctacctgtggccatctgcagaagagaatccccacaagaagcctggcaag
aaaaccaccaccaagcctaccaagaagcctacccatcaccaccatcaccactgataatcta

Seq ID NO:8

PreFG_V1 peptide for CHO
MELLILKNTNAITAILAAVTLCFASSQNITEEFYQSTCSAVSKGYLSALRTGWYTSVITIELSNIKENCNGT
DAKVKLICKQELDKYKSAVTTELQLLMQSTPATNNKKFLGFLGVGSIAASGIAVSKVLHLEGEVNKIKSALLS
TNKAVVSLNSNGSVLTSKVLSDLKNYIDKQLLPIVNKQSCSISNIETVIEFQQKNNRLEITREFSVNAGVTT
PVSTYMLTNSELLSLINDMPITNDQKKLMSNNVQIVRQQSYSIMSIKEEVILAYVQLPLYGVIDTPCWKLH
TSPLCTNTKEGSNICLRTDRGWYCDNAGSVFFPLAETCKVQSNRVCDFTMNSLTPSEVNLNCNIDIFNP
KYDCKIMTSKTDVSSSVITSLGAIIVSCYGKTKCTASNKNRGIKTFNSNGCDYVSNKGVDTVSVGNTLYYV
QEGKSLYVKGEPIINFYDPLVFPSSDEFDASISQVNEKINQSLAFIRKSDEKLHNVEDKIEEILSKIYHIENE
IARIKKLIGEAGGSGGGSKQRQNKEPNKPNNDHFVFEVFNFVPCSICSNNPTCWAICKRIPNKKPGKTTT
KPTKKPFTKTTKDKHPQTTKPKEVPTTK

Seq ID NO:9

Nucleotide Sequence encoding PreFG_V2 for CHO
aagcttgcacccatggagctgtatccatcaagaccaacgcacccatcaccgcacatcctggccgcgtgaccctg
tgcttcgcctctcccaacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcacc
ctgtccgcctgcggaccggctggtacccatccgtgatcaccatcgcacatcaccgcacatcaccgcacatcacc
tgcaacacgcacccatccgtgatcaccatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
ctccacatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
ctccacatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
tccgcacatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
gcccacatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
gatcgtcaagaactacatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
gatcgtcaagaactacatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
gagaccgtgatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
ggcgtgaccacccctgtgtccacatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
atccacaacgcacaaaaaaactgcacatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
agcatcgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
tggaaagctgcacacccctgtgtcaccaccaacaccaaaaaaggggttccatcgcacatcaccgcacatcacc
gaccggggctgtactgcgacacgcggctccgtgtcccttccctgtgtccggcagacactgtcaaggtgeag
tccaaccgggtgttctgcgacccatgcacactccctgaccctgcaccctggcggcaggtgaacctgtgcaacatcgac
atcttcaccccaactgcacatcgcacactccctgaccctgcaccctggcggcaggtgaacctgtgcaacatcgac
ctggggccatcgtgtccctgcaccgcacatcgcacactccctgaccctgcaccctggcggcaggtgaacctgtgcaacatcgac
accttcaccccaactgcacatcgcacactccctgaccctgcaccctggcggcaggtgaacctgtgcaacatcgac
tacgtgaataagcaggaaggcaagacgcgtgtacgtgaagggcggcagctatcatcaacttctacgaccctctg
gtgtccctccgacgagttcgcacgcctccatcagccaggtaacgagaagatcaaccaggccctggccct
atccggaaactccgacgagaagctgcataacgtggaggacaagatcgaagagatcctgtccaaaatctaccac
atcgagaacgagatcgcggatcaagaagactgtatcggcgaggctggccggcaagcagcggcagaacaacaagc
cctaacaagcccaacaacgcacttccacttcgagggtgttcaacttctgccttgcctcatctgcataaca
cctacctgtggccatctgcacccatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
aagaacgcctacccatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat
accaacgcacccatcaccgcacatcaccgcacatcaccgcacatcaccgcacatcaccgcacat

Seq ID NO:10

PreFG_V2 peptide for CHO
MELLILKNTNAITAILAAVTLCFASSQNITEEFYQSTCSAVSKGYLSALRTGWYTSVITIELSNIKENCNGT
DAKVKLICKQELDKYKSAVTTELQLLMQSTPATNNKKFLGFLGVGSIAASGIAVSKVLHLEGEVNKIKSALLS
TNKAVVSLNSNGSVLTSKVLSDLKNYIDKQLLPIVNKQSCSISNIETVIEFQQKNNRLEITREFSVNAGVTT
PVSTYMLTNSELLSLINDMPITNDQKKLMSNNVQIVRQQSYSIMSIKEEVILAYVQLPLYGVIDTPCWKLH
TSPLCTNTKEGSNICLRTDRGWYCDNAGSVFFPLAETCKVQSNRVCDFTMNSLTPSEVNLNCNIDIFNP

KYDCKIMTSKTDVSSSVITSLGAI VSCYGKTKCTASNKNRGI I KTFNSNGCDYVSNKGVDTSVGNLTYYVNQ
QEGKSLSYVKGEPI INFYDPLVFP SDEF DAS ISQVNEKINQSLAFIRKSDEKLHNVEDKIEEILSKIYHIENE
IARIKKLIGEAGGGKQRQNKP PNKPNNDFHFEVNFVPC SICCSNNPTCWAICKRIPNKKPGKTTTKPTKKPT
FKTTKKDHKPQTTKPKEVPTTK

SEQ ID NO:11

Exemplary coiled-coil (isoleucine zipper)
EDKIEEILSKIYHIENEIARIKKLIGEA

SEQ ID NO:12

PreF antigen polynucleotide CHO2

SEQ ID NO:13

PreF antigen polynucleotide with intron

SEQ ID NO:14
 Synthetic linker sequence
 GGSGGSGGS

SEQ ID NO:15
 Furin cleavage site
 RARR

SEQ ID NO:16
 Furin cleavage site
 RKRR

SEQ ID NO:17
 Nucleotide Sequence encoding PreF_NGTL
 atggagctgtatcctgaaaaccaacgcacatcaccgcacatcctggccgggtgaccctgtgcttcgcctcc
 tcccagaacatcaccgaggagttcaccatcaccgtccacatcgtccgggtgaccctgtgcttcgcctcc
 cggaccggctggtacacccgtatcaccatcggagctgtccacatcaaggaaaacaagtgcacggcacc
 gacccaaggtaagctgtatcagaagcaggagctggacaaggtaacaaaggagcggctgaccgaactccagctgt
 atgcagtccacccctgcccaccaacaacaaatgttctggcttcgtctgggtggctccgcacatcgcctcc
 ggcacatcggctgagcaagggtgctgcacccgtggaggcgagggtgaaacaagatcaagagcgcctgtgtccacc
 aacaaggccgtgtgtccctgtccacccgtgtccgtgtccatgtggatctgaagaactac
 atcgacaaggcagctgtgtccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 ttccagcagaagaacaacccgtgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 gtgtccacccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 aaaaaactgtatgtccaccaacgtgcagatgtggatccatgtggatccatgtggatccatgtggatccatgt
 gagggtgtggctacgtgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 tccccctgtgcacccaccaacacccgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 tgcgacaacgcggctccgtgtcccttcgtggccggatccatgtggatccatgtggatccatgtggatccatgt
 tgcgacaccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 tacgactgtcaagatcatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 tcctgtgtacggcaagatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 tgcgactacgtgtccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 gagggcaagagccgtgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 gagttcgacgcgtccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 gagaagctgtccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 gcccggatcaagaagctgtatcggcgaggcc

SEQ ID NO:18
 Amino Acid Sequence of PreF_NGTL
 MELLILKNTAITAILAAVTLCFASSQNITEEFYQSTCSAVSKGYLSALRTGWYTSVITIELSNIKENKNGT
 DAKVKLIKQELDKYKSAVTTELQLMQSTPATNNKFLGFLGVGSIAASGIAVSKVHLGEVNKIKSALLST
 NKAIVSLSNGSVLTSKVLSDLKNYIDKQLLPIVNQKSCSISNIETVIEFQQKNNRLEITREFSVNAGVTP
 VSTYMLTNSSELLSLINDMPITNDQKKLMSNNVQIVRQQSYSIMSIIKEEVLAYVVQLPLYGVIDTPWKLHT
 SPLCTNTKEGSNICKLRTDRGWYCDAGSVSFFPLAETCKVQSNRVCDFTMNSLTLPEVNLCNIDIFNPK
 YDCKIMTSKTDVSSSVITSLGAIVSCYGKTKCTASNKNRGIIKTFSNGCDYVSNKGVDTVSGNTLYVNQ
 EGKSLYVKGEPIINFYDPLVFPSEFDASISQVNEKINGTLAFIRKSDEKLHNVEDKIEEILSKIYHIENEI
 ARIKKLIGEA

SEQ ID NO:19
 Nucleotide Sequence encoding PreF_L112Q
 atggagctgtatcctgaaaaccaacgcacatcaccgcacatcctggccgggtgaccctgtgcttcgcctcc
 tcccagaacatcaccgaggagttcaccatcaccgtccacatcgtccgggtgaccctgtgcttcgcctcc
 cggaccggctggtacacccgtatcaccatcggagctgtccacatcaaggaaaacaagtgcacggcacc
 gacccaaggtaagctgtatcagaagcaggagctggacaaggtaacaaaggagcggctgaccgaactccagctgt
 atgcagtccacccctgcccaccaacaacaaatgttctggcttcgtccatgtggatccatgtggatccatgt
 ggcacatcggctgagcaagggtgctgcacccgtggaggcgagggtgaaacaagatcaagagcgcctgtgtccacc
 aacaaggccgtgtgtccctgtccacccgtgtccgtgtccatgtggatccatgtggatccatgtggatccatgt
 atcgacaaggcagctgtgtccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 ttccagcagaagaacaacccgtgtggatccatgtggatccatgtggatccatgtggatccatgt
 gtgtccacccatgtggatccatgtggatccatgtggatccatgtggatccatgtggatccatgt
 aaaaaactgtatgtccaccaacgtgcagatgtggatccatgtggatccatgtggatccatgt

gaggtgctggcctacgtggcagctgcctctgtacggcgtgatecgacaccccttgcgttggaaagctgcacacc
tccccccctgtgcaccaccaacaccaggaggctccaacatctgcctgaccggaccggaccggggctggta
tgcgacaacgcggctccgtgtccttcctctggccgagacgtcaagggtcagttcaaccgggtgttc
tgcgacaccatgaactccctgaccctgcctccggaggtaacactgtcaacatcgacatcttcaccccaag
tacgactgtcaagatcatgaccaggcaagacggacgtgtcctccagcgtatcacccctggcgccatcgt
tcctgtacggcaagaccaagtgcaccgcctccacaagaacccgggaaatcatcaagaccccttcaccc
tgcgactacgtgttcaataaggcgtggacaccgtgtccgtggcaacacactgtactacgtgaataagcg
gagggcaagagcctgtacgtgaagggcggaccatcatcaacttctacgaccctctgtgttccctccgac
gatcgcacgcctccatcagccaggtaacgagaagatcaaccaggcccttcatccgaaagtcgcac
gagaagctgcataacgtggaggacaagatcgaggagatcctgtccaaaatctaccacatcgagaacgagatc
gccccggatcaagaagctgatcggcgaggcc

SEQ ID NO:20

Amino Acid Sequence of PreF_L112Q

MELLILKNTNAITAILAAVTLCFASSQNITEEFYQSTCSAVSKGYLSALRTGWYTSVITIELSNIKENKNGT
DAKVLIKQELDKYKSAVTELQLLMQSTPATNNKFLQGVGSIAASGIAVSKVLHLEGEVNKIKSALLST
NKAIVSLSNGVSVLTSKVLSDLKNYIDKQLLPIVNQKSCSISNIETVIEFQQKNNRLLEITREFSVNAGVTP
VSTYMLTNSELLSINDMPITNDQKKLMSNNVQIVRQQSYSIMSIKEEVLAYVVLPLYGVIDTPCWLH
SPLCTTNTKEGSNICKLRTDRGWYCDNAGSVSFFPLAETCKVQSNRVCDFMNSLTLPEVNLCNIDIFNPK
YDCKIMTSKTDVSSSVITSLGAIVSCYGKTKCTASNKNRGIIKTFNSNGCDYVSNKGVDTVSVGNTLYVNQ
EGKSLYVKGEPIINFYDPLVFPSEFDASISQVNEKINQSLAFIRKSDEKLHNVEDKIEEILSKIYHIE
ARIKKLIGEA

SEQ ID NO:21

Nucleotide Sequence encoding PreF_NGTL_L112Q

atggagctgtatccctgaaaaccaacgcacatcaccgcacatcgtggccgggtgaccctgtgttgcctcc
tcccagaacatcaccggagggttctaccaggatccacgttccacgtgtccggccgtgttcaagggttacatgtccggcc
cgaccggctgttacacccgtatcaccatcgagctgttccacatcaaggaaaacaagtgcacccgacc
gaccccaagggttacgttcaaggcaggagctggacaaatgttctggcttctgcaggccgtgggttccatcgcctcc
atgcaggccacccctgcaccaacaacaaatgttctggcttctgcaggccgtgggttccatcgcctcc
ggcatcgcctgttacgttccacatcgacccgtgttccatcgcctgttccatcgcctcc
aacaaggccgtgggttccatcgcacccgtgttccatcgcctgttccatcgcctgttccatcgcctcc
atcgacaaggcactgtgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
ttccagcagaagaacaaacccggctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
gtgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
aaaaaaatgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
gaggtgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
tcccccctgtgcaccaccaacaccaggagggttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
tgcgacaacgcgggttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
tgcgacaccatgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
tacgactgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
tcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
tgcgactacgtgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
gagggcaagagcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
gatcgcacgcctccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctgttccatcgcctcc
gagaagctgcataacgtggaggacaagatcgaggagatcctgtccaaaatctaccacatcgagaacgagatc
gccccggatcaagaagctgatcggcgaggcc

SEQ ID NO:22

Amino Acid Sequence of PreF_NGTL_L112Q

MELLILKNTNAITAILAAVTLCFASSQNITEEFYQSTCSAVSKGYLSALRTGWYTSVITIELSNIKENKNGT
DAKVLIKQELDKYKSAVTELQLLMQSTPATNNKFLQGVGSIAASGIAVSKVLHLEGEVNKIKSALLST
NKAIVSLSNGVSVLTSKVLSDLKNYIDKQLLPIVNQKSCSISNIETVIEFQQKNNRLLEITREFSVNAGVTP
VSTYMLTNSELLSINDMPITNDQKKLMSNNVQIVRQQSYSIMSIKEEVLAYVVLPLYGVIDTPCWLH
SPLCTTNTKEGSNICKLRTDRGWYCDNAGSVSFFPLAETCKVQSNRVCDFMNSLTLPEVNLCNIDIFNPK
YDCKIMTSKTDVSSSVITSLGAIVSCYGKTKCTASNKNRGIIKTFNSNGCDYVSNKGVDTVSVGNTLYVNQ
EGKSLYVKGEPIINFYDPLVFPSEFDASISQVNEKINGTLAFIRKSDEKLHNVEDKIEEILSKIYHIE
ARIKKLIGEA

We claim:

1. A recombinant respiratory syncytial virus (RSV) antigen comprising a soluble F protein polypeptide comprising an F₂ domain and an F₁ domain of an RSV F protein polypeptide, wherein the F protein polypeptide comprises at least one modification that alters glycosylation.
2. The recombinant RSV antigen of claim 1, wherein the F protein polypeptide comprises at least one modification selected from:
 - (i) an addition of an amino acid sequence comprising a heterologous trimerization domain;
 - (ii) a deletion of at least one furin cleavage site;
 - (iii) a deletion of at least one non-furin cleavage site;
 - (iv) a deletion of one or more amino acids of the pep27 domain; and
 - (v) at least one substitution or addition of a hydrophilic amino acid in a hydrophobic domain of the F protein extracellular domain.
3. The recombinant RSV antigen of claim 1, wherein the modification that alters glycosylation comprises a substitution of one or amino acids comprising and/or adjacent to the amino acid corresponding position 500 of SEQ ID NO:2.
4. The recombinant RSV antigen of claim 1 or 3, wherein amino acids corresponding to positions 500-502 of SEQ ID NO:2 are selected from: NGS; NKS; NGT; NKT.
5. The recombinant RSV antigen of any one of claims 1-4, wherein the modification that alters glycosylation comprises a substitution of glutamine at the amino acid corresponding to position 500 of SEQ ID NO:2.
6. The recombinant RSV antigen of any one of claims 1-5, wherein the soluble F protein polypeptide comprises an intact fusion peptide between the F2 domain and the F1 domain.
7. The recombinant RSV antigen of any one of claims 1-6, wherein the at least one modification comprises the addition of an amino acid sequence comprising a heterologous trimerization domain.

8. The recombinant RSV antigen of claim 7, wherein the heterologous trimerization domain is positioned C-terminal to the F₁ domain.

9. The recombinant RSV antigen of any one of claims 1-8, comprising an F2 domain and an F1 domain with no intervening furin cleavage site.

10. The recombinant RSV antigen of any one of claims 1-9, wherein the RSV antigen assembles into a multimer

11. The recombinant RSV antigen of any one of claims 1-10, wherein the RSV antigen assembles into a trimer.

12. The recombinant RSV antigen of any one of claims 1-11, wherein the F₂ domain comprises at least a portion of an RSV F protein polypeptide corresponding to amino acids 26-105 of the reference F protein precursor polypeptide (F₀) of SEQ ID NO:2.

13. The recombinant RSV antigen of any one of claims 1-12 and wherein the F₁ domain comprises at least a portion of an RSV F protein polypeptide corresponding to amino acids 137-516 of the reference F protein precursor polypeptide (F₀) of SEQ ID NO:2.

14. The recombinant RSV antigen of any one of claims 1-13, wherein the F₂ domain comprises an RSV F protein polypeptide corresponding to amino acids 26-105 and/or wherein the F₁ domain comprises an RSV F protein polypeptide corresponding to amino acids 137-516 of the reference F protein precursor polypeptide (F₀) of SEQ ID NO:2.

15. The recombinant RSV antigen of any one of claims 1-14, wherein the RSV antigen is selected from the group of:

- a) a polypeptide comprising SEQ ID NO:22;
- b) a polypeptide encoded by SEQ ID NO:21 or by a polynucleotide sequence that hybridizes under stringent conditions over substantially its entire length to SEQ ID NO:21;
- c) a polypeptide with at least 95% sequence identity to SEQ ID NO:22.

16. The recombinant RSV antigen of any one of claims 1-15, wherein the F2 domain comprises amino acids 1-105 of the RSV F protein polypeptide.

17. The recombinant RSV antigen of any one of claims 1-16, wherein the F₂ domain and the F₁ domain are positioned with an intact fusion peptide and without an intervening pep27 domain.
18. The recombinant RSV antigen of any one of claims 1-17, wherein the heterologous trimerization domain comprises a coiled-coil domain.
19. The recombinant RSV antigen of claim 18, wherein the multimerization domain comprises an isoleucine zipper.
20. The recombinant RSV antigen of claim 19, wherein the isoleucine zipper domain comprises the amino acid sequence of SEQ ID NO:11.
21. The recombinant RSV antigen of any one of claims 1-20, wherein the RSV antigen comprises at least one substitution or addition of a hydrophilic amino acid in a hydrophobic domain of the F protein extracellular domain.
22. The recombinant RSV antigen of claim 21, wherein the hydrophobic domain is the HRB coiled-coil domain of the F protein extracellular domain.
23. The recombinant RSV antigen of claim 22, wherein the HRB coiled-coil domain comprises the substitution of a charged residue in place of a neutral residue at the position corresponding to amino acid 512 of the reference F protein precursor (F₀) of SEQ ID NO:2.
24. The RSV antigen of claim 23, wherein the HRB coiled-coil domain comprises a substitution of lysine or glutamine for leucine at the position corresponding to amino acid 512 of the reference F protein precursor (F₀) of SEQ ID NO:2.
25. The recombinant RSV antigen of claim 21, wherein the hydrophobic domain is the HRA domain of the F protein extracellular domain.
26. The recombinant RSV antigen of claim 25, wherein the HRA domain comprises the addition of a charged residue following the position corresponding to amino acid 105 of the reference F protein precursor (F₀) of SEQ ID NO:2.

27. The recombinant RSV antigen of claim 26, wherein the HRA domain comprises the addition of a lysine following the position corresponding to amino acid 105 of the reference F protein precursor (F₀) of SEQ ID NO:2.

28. The recombinant RSV antigen of any one of claims 21-27, wherein the RSV antigen comprises at least a first substitution or addition of a hydrophilic amino acid in the HRA domain and at least a second substitution or addition of a hydrophilic amino acid in the HRB domain of the F protein extracellular domain.

29. The recombinant RSV antigen of any one of claims 1-28, wherein the RSV antigen comprises at least one amino acid addition, deletion or substitution that eliminates a furin cleavage site present in a naturally occurring F protein precursor (F₀).

30. The recombinant RSV antigen of 29, wherein the RSV antigen comprises an amino acid addition, deletion or substitution that eliminates a furin cleavage site at a position corresponding to amino acids 105-109, a position corresponding to amino acids 133-136, or at both positions corresponding to amino acids 105-109 and 133-136 of the reference F protein precursor (F₀) of SEQ ID NO:2.

31. The recombinant RSV antigen of any one of claims 1-30, wherein the F₁ and F₂ polypeptide domains correspond in sequence to the RSV A Long strain.

32. The recombinant RSV antigen of any one of claims 1-31, wherein the RSV antigen comprises a multimer of polypeptides.

33. The recombinant RSV antigen of any one of claims 1-31, wherein the RSV antigen comprises a trimer of polypeptides.

34. An immunogenic composition comprising the recombinant RSV antigen of any one of claims 1-33, and a pharmaceutically acceptable carrier or excipient.

35. The immunogenic composition of claim 34, further comprising an adjuvant.

36. The immunogenic composition of claim 35, wherein the adjuvant comprises at least one of: 3D-MPL, QS21, an oil-in-water emulsion, and Alum.

37. The immunogenic composition of claim 36, wherein the adjuvant comprises an oil-in-water emulsion.

38. The immunogenic composition of claim 36 or 37, wherein the oil-in-water emulsion comprises a tocol.

39. The immunogenic composition of any one of claims 36-38, wherein the oil-in-water emulsion comprises less than 5 mg squalene per human dose.

40. The immunogenic composition of any one of claims 35-39, wherein the adjuvant is suitable for administration to a neonate.

41. The immunogenic composition of any one of claims 34-40, wherein when the immunogenic composition comprises the recombinant RSV antigen of any one of claims 1-33 the immunogenic composition further comprises a G protein polypeptide comprising an amino acid sequence corresponding to amino acid positions 149 to 229 of an RSV G protein polypeptide.

42. The immunogenic composition of any one of claims 34-41, further comprising at least one additional antigen of a pathogenic organism other than RSV.

43. A recombinant nucleic acid comprising a polynucleotide sequence that encodes the recombinant RSV antigen of any one of claims 1-33.

44. The recombinant nucleic acid of claim 43, wherein the polynucleotide sequence that encodes the RSV antigen is codon optimized for expression in a selected host cell.

45. The recombinant nucleic acid of claim 43 or 44, wherein the nucleic acid comprises a polynucleotide sequence selected from:

- a) a polynucleotide sequence comprising SEQ ID NO:21;
- b) a polynucleotide sequence that encodes SEQ ID NO:22;
- c) a polynucleotide sequence that hybridizes under stringent conditions over substantially its entire length to SEQ ID NO:21;
- d) a polynucleotide sequence with at least 95% sequence identity to SEQ ID NO:21, which polynucleotide sequence does not correspond to a naturally occurring RSV strain.

46. A vector comprising the recombinant nucleic acid of any one of claims 43-45.

47. A host cell comprising the nucleic acid of claim 43 or 44 or the vector of claim 46.

48. The use of the RSV antigen of any one of claims 1-33 in the preparation of a medicament for treating an RSV infection.

49. The use of the RSV antigen or nucleic acid of claim 48, wherein the medicament is administered for the purpose of prophylactically treating an RSV infection.

50. A method for eliciting an immune response against Respiratory Syncytial Virus (RSV), the method comprising:

administering to a subject a composition comprising the recombinant RSV antigen of any one of claims 1-33 or the immunogenic composition of any one of claims 34-42.

51. The method of claim 50, wherein administering the composition comprising the RSV antigen elicits an immune response specific for RSV without enhancing viral disease following contact with RSV.

52. The method of claims 50 or 51, wherein the immune response comprises a protective immune response that reduces or prevents infection with a RSV and/or reduces or prevents a pathological response following infection with a RSV.

53. The method of any one of claims 50-52, wherein the subject is a human subject.

54. The recombinant RSV antigen of any one of claims 1-33 or the immunogenic composition of any one of claims 34-42 for use in medicine.

55. The recombinant RSV antigen of any one of claims 1-33 or the immunogenic composition of any one of claims 34-42 for the prevention or treatment of RSV-associated diseases.

56. A method for producing a recombinant RSV antigen with an altered glycosylation pattern, the method comprising:



- (i) expressing a nucleic acid of any one of claims 43-45 in a host cell; and
- (ii) isolating the recombinant RSV antigen expressed thereby.

57. A method for increasing expression of an RSV fusion protein polypeptide, the method comprising:

expressing in a host cell an a nucleic acid encoding an RSV fusion protein polypeptide comprising at least one mutation that produces an amino acid addition, deletion or substitution, which amino acid addition, deletion or substitution alters a glycosylation site of the encoded RSV fusion protein polypeptide as compared to a naturally occurring RSV fusion protein, whereby altering the glycosylation site increases expression of the RSV fusion protein polypeptide as compared to an RSV fusion protein without the amino acid addition, deletion or substitution.

58. The method of claim 57, wherein the nucleic acid encoding the RSV fusion protein polypeptide further comprises an additional mutation that produces an amino acid addition, deletion or substitution that eliminates a peptidase cleavage site within the encoded RSV fusion protein polypeptide.

59. The method of claim 57 or 58, wherein the RSV fusion protein polypeptide is a recombinant fusion protein polypeptide stabilized in the prefusion conformation.

FIG. 1A**FIG. 1B**




FIG. 2




FIG. 3

FIG. 4A

Serology in immunized mice, CHO-preF




FIG. 4B

Serology in immunized mice, CHO-preF





FIG. 5A

Evaluation of CHO-preF in vivo

Neutralization titers



* Same assay, sera taken from different animal study

FIG. 5B
Evaluation of CHO-preF in vivo
Neutralization titers




FIG. 6A
CHO-preF protection against challenge




FIG. 6B
CHO-preF protection against challenge




FIG. 7
Pulmonary eosinophil recruitment following challenge in mice




FIG. 8
Serum IgG levels in mice immunized with PreF formulated with an oil-in-water emulsion adjuvant





FIG. 9
Neutralizing Antibody Titers in Mice immunized with PreF
formulated with an oil-in-water emulsion adjuvant



PROTECTION AGAINST CHALLENGE IN MICE IMMUNIZED
WITH preF FORMULATED WITH AN OIL-IN-WATER
EMULSION ADJUVANT
RSV TITRATION IN LUNG HOMOGENATES
OF CHALLENGED MICE
SPEARMAN-KARBER FORMULA (WITH GEOMETRIC MEAN)




FIG. 10