



HU000032330T2

(19) **HU**(11) Lajstromszám: **E 032 330**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala**EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA

- (21) Magyar ügyszám: **E 10 702358**
- (22) A bejelentés napja: **2010. 01. 25.**
- (96) Az európai bejelentés bejelentési száma:
EP 20100702358
- (97) Az európai bejelentés közzétételi adatai:
EP 2381797 A1 **2010. 07. 29.**
- (97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2381797 B1 **2017. 03. 08.**
- (51) Int. Cl.: **A23K 20/158** (2006.01)
A23K 50/30 (2006.01)
A23K 50/70 (2006.01)
A23K 50/75 (2006.01)
A23K 50/80 (2006.01)
- (86) A nemzetközi (PCT) bejelentési szám: **PCT/NL 10/050033**
- (87) A nemzetközi közzétételi szám: **WO 10085149**

(30) Elsőbbségi adatok: 09151285 2009. 01. 23. EP	(73) Jogosult(ak): Nutreco Nederland B.V., 5831 JN Boxmeer (NL)
(72) Feltaláló(k): HAN, Yanming, NL-6525 RB Nijmegen (NL) SMITS, Coenraad Henricus Maria, NL-6581 PC Malden (NL) BRENNAN, John, GuelphOntario N1G4T2 (CA) PAGE, Gregory Ian, ParisOntario N3L 4G4 (CA) VAN DAM, Johannes Teunis Pieter, NL-3903 GR Veenendaal (NL)	(74) Képviselő: Danubia Szabadalmi és Jogi Iroda Kft., Budapest

- (54) **Közepes lánchosszúságú zsírsavak alkilésztereit tartalmazó állateledel-adalék és állateledel, és alkalmazásuk állateledelben**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmat az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 2 381 797 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
08.03.2017 Bulletin 2017/10

(51) Int Cl.:
A23K 20/158 ^(2016.01) **A23K 50/70** ^(2016.01)
A23K 50/75 ^(2016.01) **A23K 50/30** ^(2016.01)
A23K 50/80 ^(2016.01)

(21) Application number: **10702358.2**

(86) International application number:
PCT/NL2010/050033

(22) Date of filing: **25.01.2010**

(87) International publication number:
WO 2010/085149 (29.07.2010 Gazette 2010/30)

(54) **ANIMAL FEED ADDITIVE AND ANIMAL FEED COMPRISING ALKYL ESTERS OF MEDIUM CHAIN FATTY ACIDS, AND THEIR USE IN ANIMAL FEED**

TIERFUTTERZUSATZMITTEL UND TIERFUTTER ENTHALTEND ALKYLESTER VON FETTSÄUREN MITTLERER KETTENLÄNGE, UND IHRE VERWENDUNG IN TIERFUTTER

ADDITIF D'ALIMENT DE BÉTAIL ET ALIMENT DE BÉTAIL COMPRENANT DES ESTERS ALCOYLS D'ACIDES GRAS DE LONGUEUR DE CHAÎNE INTERMÉDIAIRE, ET LEUR UTILISATION DANS LES ALIMENTS DE BÉTAIL

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR

• **VAN DAM, Johannes Teunis Pieter**
NL-3903 GR Veenendaal (NL)

(30) Priority: **23.01.2009 EP 09151285**

(74) Representative: **van Someren, Petronella F. H. M. Arnold & Siedsma**
Bezuidenhoutseweg 57
2594 AC The Hague (NL)

(43) Date of publication of application:
02.11.2011 Bulletin 2011/44

(56) References cited:
WO-A-01/52837 WO-A-01/97799
WO-A-2006/002927

(73) Proprietor: **Nutreco Nederland B.V.**
5831 JN Boxmeer (NL)

(72) Inventors:
 • **HAN, Yanming**
NL-6525 RB Nijmegen (NL)
 • **SMITS, Coenraad Henricus Maria**
NL-6581 PC Malden (NL)
 • **BRENNAN, John**
Guelph
Ontario N1G4T2 (CA)
 • **PAGE, Gregory Ian**
Paris
Ontario N3L 4G4 (CA)

• **KABARA J J: "Fatty acids and derivatives as antimicrobial agents: a review. In: Pharmacological effects of lipids, pp. 1-14" 1 January 1978 (1978-01-01), AMERICAN OIL CHEMIST'S SOCIETY , XP000892960 page 1, paragraph 2 page 10, paragraphs 3,6 tables II-III**
 • **KABARA J J: "Combined effects on antibacterial activity of fatty acids and their esters against gram-negative bacteria. In: Pharmacological effects of lipids, pp. 15-24" 1 January 1978 (1978-01-01), AMERICAN OIL CHEMIST'S SOCIETY , XP009115700 abstract page 16, paragraph 1**

EP 2 381 797 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The invention is directed to an animal feed and method for increasing feed efficiency and health in farming animals, including mammals, birds and fish.

[0002] The use of fatty acids as antimicrobial agents to inhibit growth of harmful microbes in living organisms has long been known in literature, see e.g. US-A-2 466 663. A mixture of medium chain fatty acids (MCFAs) and organic acids is currently widely used to inhibit growth of pathogens in the gastrointestinal tract of animals. The presence of such a mixture in animal feed results in a favorable composition of the gastrointestinal microbiota and an improved feed efficiency in animals.

[0003] EP-A-I 059 041 discloses a feed supplement composition comprising at least one triglyceride containing MCFA and at least one exogenous lipolytic enzyme. The addition of this supplement to animal feed results in a physiological environment in the animal's gastrointestinal tract, including its stomach which regulates and stabilizes the gut microbiota and consequently results in improvement of growth in the animal.

[0004] WO-A-01/97799 discloses the use in feed compositions of one or more C6 - C10 medium chain fatty acids, salts, derivatives or mixtures thereof for the inhibition of microbial contamination. However the specific use and effectiveness of alkyl esters of MCFAs as antimicrobial agents is not mentioned.

[0005] US-A-4 526 798 discloses a mixture comprising the ethyl ester of 2- hydroxy-4-methylpentanoic acid, which mixture can be used in flavors that enhance the aroma or taste of tropical flavored foodstuffs. The concentration of the mixture in such flavors ranges from 0.75% to 1.2% by weight of the flavor. The resulting dosage of the ethyl ester in the foodstuff is low, for example 3 - 8 ppm. The effect of the mixture as an antimicrobial agent is not described in this document.

[0006] WO-2006/00297 discloses the use of MCFAs, derivatives thereof or mixtures for inhibiting the growth and/or for reducing the amount of microbial pathogens. This document also does not specifically mention alkyl esters of MCFAs and their effectiveness as an inhibitor of microbial pathogens.

[0007] WO-A-01/52837 discloses the use of a fatty ester for preparing a composition designed to inhibit 5- α -reductase activity in pharmacology, dermatology, cosmetics and as a food additive.

[0008] Karbara J. J., American Oil Chemist's Society, pp. 1-14, 1 January 1978, discloses a review of fatty acids and derivatives as antimicrobial agents. However it does not specifically mention alkyl esters of MCFAs and their effectiveness as an antimicrobial agent.

[0009] Karbara J. J., American Oil Chemist's Society pp. 15-24, 1 January 1978, discloses the combined effects on antibacterial activity of fatty acids and their esters with freezing or heating and the addition of chemicals such as citric acid against gram-negative bacteria. Further this review only discusses monoglycerides of MCFAs and not alkyl esters of MCFAs.

[0010] A disadvantage of the MCFAs and derivatives in animal feed of the prior art is that they are quickly absorbed *in vivo* in the proximal small intestinal tract. Consequently, they are not able to exert microbiota modulating properties in the distal small intestine and hind gut.

[0011] Other disadvantages of MCFAs are their corrosivity, their irritating odor and poor flavor. Ethyl esters and methyl esters are not corrosive, but esters based on the shorter chain MCFAs usually have a low flash point.

[0012] Another disadvantage is that in many cases MCFAs showed less pathogen activity than their ester counterparts because the esters possess nonspecific activity of a surface- active agent (surfactant). Studies have demonstrated that the anti-bacterial, anti-fungal and anti-viral activities of the esters were due to their functions in disrupting cell membranes, see K. Nihei et al, J. Agric. Food Chem., 52 (2004) 5011-5020.

[0013] Object of the present invention is to provide an animal feed that has strong antimicrobial and antipathogenic properties and results in a favorable composition of the gastrointestinal microbiota and improved feed efficiency in animals fed with said animal feed, and which overcomes at least in part one or more of the above-mentioned disadvantages of the prior art.

[0014] In a first aspect, this object is met by providing an animal feed suitable for feeding mammals, birds and fish, comprising an alkyl ester of a fatty acid, wherein said fatty acid has a chain length of 5 - 12 carbon atoms, and wherein the dosage of said ester in said animal feed is 50 ppm by weight or higher, based on the total weight of said animal feed, for use in decreasing the risk of microbial infections in an animal.

[0015] In the present application, with a medium chain fatty acid (MCFA) is meant a fatty acid having a chain length of 5 - 12 carbon atoms, i.e. having a longest continuous chain of 5 - 12 carbon atoms. Preferably, the animal feed of the invention comprises an MCFA having a chain length of 8 - 12 carbon atoms. A chain length longer than 12 carbon atoms is not desirable, because this will have a negative influence on the degree of hydrolysis of the ester. In the present application, an MCFA having a chain length of x is sometimes referred to as Cx. For example, octanoic acid may be referred to as C8.

[0016] The inventors found that the esters of MCFAs showed stronger antimicrobial activity than MCFAs and therefore provide stronger microbiota modulating properties inside the gut compared to MCFAs.

[0017] The inventors further surprisingly found that the esterified forms of MCFA according to the invention have a

prolonged activity in the gastrointestinal tract compared to non-esterified forms of MCFA. In addition, it was found that alkyl esters of MCFAs according to the present invention are even more anti- microbial than their non-esterified counterparts. A further advantage of the animal feed of the present invention is that alkyl esters of MCFAs are still relatively cheap, although they are more expensive than plain MCFAs (C8, C10 and/or C12).

5 **[0018]** The alkyl in the ester in the animal feed of the invention is preferably methyl, ethyl, propyl, butyl or a combination thereof.

[0019] The MCFA in the ester in the animal feed of the invention is preferably octanoic acid, decanoic acid, dodecanoic acid or a combination thereof.

10 **[0020]** The alkyl ester of MCFA in the animal feed of the invention is preferably methyl octanoate, methyl decanoate, methyl dodecanoate, ethyl octanoate, ethyl decanoate, ethyl dodecanoate, propyl octanoate, propyl decanoate, propyl dodecanoate, butyl octanoate, butyl decanoate, butyl dodecanoate or a combination thereof.

[0021] The dosage of the alkyl ester of MCFA in the animal feed of the invention is preferably 50 ppm by weight or higher, more preferably 100 ppm by weight or higher, most preferably 200 ppm by weight or higher, based on the total weight of the animal feed. A dosage lower than 50 ppm by weight may result in an insufficient antimicrobial effect. The dosage of the alkyl ester of MCFA in the animal feed of the invention should preferably be less than 5000 ppm by weight, preferably less than 1000 ppm by weight, more preferably less than 500 ppm by weight, based on the total weight of the animal feed. A disadvantage of using dosages higher than 5000 ppm by weight is that such dosages may have an undesired effect on the microbiotic balance. A further disadvantage of using high dosages is the resulting high cost price.

15 **[0022]** The gastrointestinal tract comprises the stomach, which has a pH of 3-4, the large intestine, which has a pH of 6-7, and the small intestine, which has a pH of about 7. Microorganisms, both pathogens and neutral or beneficial microorganisms, are in particularly present in the large and the small intestine.

[0023] Without wishing to be bound by theory, it is believed that the absorption of MCFAs is delayed by the esterified form according to the present invention. In this way the bioactive form of alkyl esters of MCFA will be able to express its bioactivity as far as the distal small intestinal tract, e.g. the distal end of the small intestines, which is crucial for modulating the local microbiota, resulting in improved efficiency of nutrient utilization (feed efficiency). The feed conversion ratio (FCR) is a measure of an animal's efficiency in converting feed mass into increased body mass and can be defined as the mass of the food eaten divided by the body mass gain, all over a specified period of time. The ability of alkyl esters of MCFA to express its bioactivity in the small intestine may also be beneficial for the control of important potential enteric pathogens that are mainly situated in the distal intestinal tract. Examples of such pathogens are *Clostridium perfringens*, *Streptococcus suis*, *Escherichia coli* and *Salmonella spp.*

25 **[0024]** Microorganisms can be divided into gram-negative and gram-positive microorganisms. An example of gram-negative microorganisms is *Escherichia coli*, which is the causative agent of diarrhea in pigs. An example of a gram-positive microorganism is *Clostridium perfringens*, which microorganism plays an important role in development of *Necrotic Enteritis* in the small intestine of broiler chickens.

30 **[0025]** The balance in the gastrointestinal tract is very important for prevention and treatment of enteric infections in living organism. For example, it is important for the health of a living organism that each part of the gastrointestinal tract has a certain pH range and that there is a favorable composition of different microorganisms present in each part of the gastrointestinal tract. The balance in the gastrointestinal tract may be influenced by adding additives to the animal feed.

[0026] Alkyl esters of MCFA have no significant effect on the pH in the intestines. Alkyl esters of MCFA act on the membrane of microorganisms of both pathogens and neutral and beneficial microorganisms, thus disabling and/or destroying the microorganisms.

35 **[0027]** The animal feed according to the present invention works in two ways. In the first place, pathogens are inhibited by the antimicrobial property of the alkyl esters of MCFA, thereby decreasing the risk of infections. Furthermore, the decrease in microbial activity of microorganisms in general in the gastrointestinal tract results in a decrease in the feed conversion ratio (FCR), corresponding to an improved feed efficiency in the animal.

40 **[0028]** It was found that alkyl esters of MCFA are particularly suitable for inhibition of gram-positive bacteria, while organic acids are particularly suitable for inhibition of gram-negative bacteria. The mode of action on microbes of MCFA and organic acids differs. It was further found that the combination of alkyl esters of MCFA and organic acids may have a synergistic effect on the inhibition of harmful microbes and on lowering the microbial activity in general. Alkyl esters of MCFA, in particular ethyl esters of MCFA, disrupt the barrier properties of the membrane of the microbe, thereby enhancing the passage of organic acids into the microbial cell, which leads to inhibition of essential metabolic pathways. Therefore, it may be preferable to add one or more organic acids to the animal feed according to the present invention. Examples of preferable organic acids are one or more of valeric acid (pentanoic acid), caprylic acid (octanoic acid), capric acid (decanoic acid), formic acid, acetic acid, propionic acid, lactic acid, butyric acid, citric acid, malic acid, fumaric acid, benzoic acid, succinic acid, sorbic acid, tartaric acid, or sodium-, potassium-, ammonium-, or calcium salts thereof.

50 **[0029]** Additionally, gallic acid or an alkyl ester of gallic acid may be added to the animal feed according to the invention. Gallic acid and alkyl ester of gallic acid also have antimicrobial properties. It was found that animal feed comprising an alkyl ester of MCFA in combination with gallic acid and/or an alkyl ester of gallic acid has a synergistic effect on the

antimicrobial properties of these molecules in the gastrointestinal tract, thereby significantly improving the bioactivity of these molecules. Examples of particularly preferred alkyl esters of gallic acid that may be added to the animal feed of the invention are propylgallate, pentyl gallate, octyl gallate, and lauryl gallate.

[0030] The animal feed according to the invention is suitable for all animals, including mammals, fish and birds. It is particularly suitable for animals having a single stomach, for example for pigs or poultry.

[0031] The invention is furthermore directed to an ingredient, premix or supplement for an animal feed suitable for feeding mammals, birds and fish, comprising an alkyl ester of a fatty acid, wherein said fatty acid has a chain length of 5 - 12 carbon atoms, and wherein the dosage of said ester is 1 wt.% or higher. Such an ingredient, premix or supplement may further comprise one or more of the following additives: micro ingredients, such as vitamins and trace elements; MCFA; minerals and organic acids.

[0032] The invention is further directed to a method for increasing the feed efficiency in animals comprising feeding a mammal, bird or fish with the animal feed, ingredient, premix or supplement of the present invention.

[0033] The invention is also directed to the use of an alkyl ester of MCFA or an alkyl ester of MCFA enriched product in a feed supplement for the improvement of feed efficiency in an animal, wherein said MCFA has a chain length of 5-12 carbon atoms and the dosage of said ester in said animal feed is 50 ppm by weight or higher, based on the total weight of said animal feed, and wherein said use is non-therapeutic.

[0034] The invention is also directed to an alkyl ester of a fatty acid or alkyl ester of a fatty acid enriched product for use in an animal feed for decreasing the risk of microbial infections in an animal, wherein said fatty acid has a chain length of 5-12 carbon atoms and the dosage of said ester in said animal feed is 50 ppm by weight or higher, based on the total weight of said animal feed.

[0035] The invention is now elucidated on the basis of some examples, which are not intended to limit the scope of the invention.

Examples

Example 1. Inhibition of *Clostridium perfringens* by various methyl and ethyl esters of Medium chain fatty acids.

[0036] A spectrophotometer was used to measure growth of micro organisms in broth over time (8 hours) by means of optical density (OD) for broths inoculated with specific bacterial strains. The relative decrease of OD is a measure of inhibitory strength. Broths were treated with graded levels of potential inhibitory substances and incubated for 24 hours.

[0037] Treatments used were the following methyl octanoate; ethyl octanoate; methyl decanoate; ethyl decanoate; ethyl dodecanoate.

[0038] The results are given in figure 1. From figure 1 it can be seen that ethyl dodecanoate shows complete inhibition at all tested levels, while methyl octanoate shows nearly the same results. It can be concluded that from all the treatments used ethyl dodecanoate is the most effective inhibitor against *Clostridium perfringens*.

Example 2. Inhibition of *Clostridium perfringens* by Lauric acid (C12) and its ethyl ester (EL) either or not in combination with octanoic/decanoic acid (C8/C10).

[0039] A spectrophotometer was used to measure growth of micro organisms in broth over time (8 hours) by means of optical density of broths inoculated with specific bacterial strains. The relative decrease of OD is a measure of inhibitory strength. Broths are treated with graded levels of potential inhibitory substances and incubated for 24 hours.

[0040] Treatments used were the following: ethyl dodecanoate (ED); dodecanoic acid (C12); blend of octanoic/decanoic acid (C8/C10); combination of C8/C10 and ED (50/50); combination of C8/C10 and C12 (50/50).

[0041] The results are given in figure 2. From figure 2 it can be seen that ethyl ester of dodecanoic acid demonstrated complete inhibition of *Clostridium perfringens* even at the lowest inclusion level of treatment whereas dodecanoic acid was not as effective as effective at 67 ppm. The combination of octanoic/decanoic acid and ethyl dodecanoate also was effective in inhibiting *Clostridium perfringens* at all levels of treatment.

[0042] Therefore it can be concluded that the ethyl ester has a higher inhibitory strength than the free fatty acid, but a comparable inhibitory effect to that of the combination of octanoic/decanoic acid and ethyl dodecanoate.

Example 3. Effect of 1000 ppm of ethyl dodecanoate or dodecanoic acid in the feed of broilers on recovery in the various segments of the gastro intestinal tract.

[0043] Broilers were offered feed supplemented with 1000 ppm of either dodecanoic acid or ethyl dodecanoate throughout the trial. At day 43 of age, 12 birds per treatment were sacrificed and the content of the duodenum, jejunum and ileum was harvested and analysed for levels of ethyl dodecanoate and dodecanoic acid. In figure 3 the mean levels are depicted.

[0044] Figure 3 shows that the level of the ethyl ester is higher than the free fatty acid in the jejunum. Therefore it can be concluded that the ethyl ester is not as quickly adsorbed through the intestinal wall as the free fatty acid.

Example 4. Effect of 1000 ppm of ethyl dodecanoate or dodecanoic acid in the feed of broilers, inoculated with Clostridium, on subsequent Clostridium counts in jejunum.

[0045] Broilers were housed in group pens and offered feed supplemented with 1000 ppm of ethyl dodecanoate or dodecanoic acid. Parallel trials were conducted, one with normal birds and one with birds which were inoculated with 10⁸ CFU *Clostridium perfringens* at day 9 through 11 birds. At day 13 of age, Clostridium counts were measured in fresh chime taken in the jejunum (see table 1). It was found that the feed supplemented by ethyl ester of dodecanoic acid led to an overall significant reduction of Clostridium counts. The effect was largest when the broilers were stressed by Clostridium inoculation.

Table 1. Clostridium perfringens counts in jejunum of infected and normal broilers on day 13 of age.

Animal model	Supplementation in feed		P-level
	C12 free form	C12 ethyl ester	
Non-challenged	1.418	1.048	<i>p</i> = 0.15
Challenged	2.122	0.977	<i>p</i> = 0.24
Overall	1.770	1.013	<i>p</i> = 0.08

[0046] These results demonstrate that supplementation with ethyl dodecanoate provides a stronger inhibition of *Clostridium perfringens* in jejunum than supplementation with equal levels of dodecanoic acid. This effect can most probably be attributed to the higher residual levels of degradation product of ethyl dodecanoate (viz. dodecanoic acid) in the jejunum (see example 3, figure 3).

Example 5. Effect of 1000 ppm of ethyl dodecanoate or dodecanoic acid in the feed of broilers, inoculated with Clostridium, on growth, feed conversion and mortality.

[0047] In broilers infected at day 9-11 with *Clostridium perfringens*, weight gain, feed conversion ratio and mortality was followed for 6 weeks (42 days). The feed was either supplemented with ethyl dodecanoate or with dodecanoic acid at 1000 ppm. An overall improvement was shown for all the parameters measured in the group of ethyl dodecanoate (table 2) as compared to the free fatty acid.

Table 2: Growth, feed conversion and mortality of *Clostridium perfringens* infected broilers

0-42 day performance	1000 ppm Dodecanoic acid	1000 ppm Ethyl dodecanoate
Weight gain, g/day	63.78	65.5
Feed intake, g/day	104.3	106.5
Feed conversion ratio	1.636	1.626
Mortality, %	8.333	4.621

[0048] From table 2 it can be concluded that the ethyl ester of dodecanoic acid results in a better zootechnical performance than dodecanoic acid itself.

Example 6. Effect of 1000 ppm of ethyl dodecanoate or dodecanoic acid in the feed of piglets on zootechnical performance and diarrhoea.

[0049] A trial was conducted wherein the effect of 1000 ppm of dodecanoic acid or 1000 ppm ethyl dodecanoate on the feed of 108 weaned piglets housed in group pens (n=9 per treatment), was tested. The effect on average diarrhoea score (over the total period) per treatment is shown below in table 3:

EP 2 381 797 B1

Table 3: Diarrhoea score of weaned piglets fed with ethyl dodecanoate or dodecanoic acid supplements

Days	Treatment			Overall			
	Control	C12	Ethyl-C12	mean	std	χ^2	<i>P</i> -value ¹
Diarrhoea score	49.84 a	46.64 ab	42-41 b	48.01	20.75	7.55	0.056

[0050] From table 3, it can be seen that ethyl dodecanoate was able to significantly reduce diarrhoea whereas dodecanoic acid did not. Therefore it can be concluded that the effect on pathogenic bacteria in the intestinal tract of piglets of ethyl ester is stronger than of the related fatty acid.

Example 7. Effect of blend of medium chain fatty acids and ethyl dodecanoate in piglet

[0051] In a trial with 48 individually housed piglets the effect of a blend of medium chain fatty acids (octanoic, decanoic, dodecanoic acid) and ethyl dodecanoate in the relative proportion 30% : 37% : 18% : 15% was tested. This blend was dosed at a total level of 1000 ppm in the weaner feed and was offered in the first 4 weeks after weaning. The piglets were infected with β - hemolytic *E.coli* (O149:K88acK91) at day 6 after weaning. At day 21 after weaning faeces samples were collected from 8 piglets per treatment and microbiological counts were measured. Further, each piglet was visually scored for faecal consistency in the same period (20-27 days after weaning) to determine the diarrhoea score.

Table 4: Microbiological counts of faeces and diarrhoea score of piglets infected with β -hemolytic *E.coli*

	Enterobacteria	<i>E.coli</i>	Lactobacillus	Diarrhoea score
1. Control	6.7740 a	6.2431 a	7.7108	38.1%
2. MCFA and esters	5.4452 b	5.2044 b	7.9165	25.0%
Standard deviation	1.4179	1.2158	0.8964	n.a.
<i>p</i> value	<i>p</i> < 0.10	<i>p</i> < 0.10	<i>p</i> > 0.10	<i>p</i> > 0.10

[0052] From table 4 it can be seen that a blend of medium chain fatty acids and esters can reduce microbial contamination by pathogenic species like *E. coli* (part of the family of Enterobacteria) and indeed reduce the number of cases of diarrhoea. Moreover, commensal microbial species like *Lactobacillus* were not inhibited.

Claims

1. Animal feed suitable for feeding mammals, birds and fish, comprising an alkyl ester of a fatty acid, wherein said fatty acid has a chain length of 5-12 carbon atoms, and wherein the dosage of said ester in said animal feed is 50 ppm by weight or higher, based on the total weight of said animal feed, for use in decreasing the risk of microbial infections in an animal.
2. Animal feed according to claim 1, wherein said alkyl is methyl, ethyl, propyl, butyl or a combination thereof.
3. Animal feed according to claim 1 or 2, wherein said fatty acid has a chain length of 8-12 carbon atoms.
4. Animal feed according to any of the previous claims, wherein said fatty acid is octanoic acid, decanoic acid, dodecanoic acid or a combination thereof.
5. Animal feed according to any of the previous claims, wherein said ester is methyl octanoate, methyl decanoate, methyl dodecanoate, ethyl octanoate, ethyl decanoate, ethyl dodecanoate, propyl octanoate, propyl decanoate, propyl dodecanoate, butyl octanoate, butyl decanoate, butyl dodecanoate or a combination thereof.
6. Animal feed according to any of the previous claims, wherein the dosage of said ester in said animal feed is 100 ppm by weight or higher, preferably 200 ppm by weight or higher, based on the total weight of said animal feed.
7. Animal feed according to any of the previous claims, wherein the dosage of said ester in said animal feed is less

than 5000 ppm by weight, preferably less than 1000 ppm by weight, based on the total weight of said animal feed.

8. Animal feed according to any of the previous claims, further comprising organic acids and/or salts thereof.
- 5 9. Animal feed according to claim 8, wherein said organic acid and/or said salt thereof is chosen from the group consisting of pentanoic, octanoic, decanoic, formic, acetic, propionic, lactic, butyric, citric, malic, fumaric, benzoic, succinic, sorbic, tartaric acid and/or salt and combinations thereof, wherein said salt is preferably the sodium, potassium, ammonium or calcium salt of said organic acids.
- 10 10. Animal feed according to any of the previous claims, further comprising gallic acid or an alkyl ester of gallic acid.
11. Animal feed according to claim 10, wherein said alkyl ester of gallic acid is propyl gallate, pentyl gallate, octyl gallate or lauryl gallate.
- 15 12. Animal feed suitable for feeding mammals, birds and fish, comprising one or more organic acids and an alkyl ester of a fatty acid, wherein said fatty acid has a chain length of 5-12 carbon atoms, and wherein the dosage of said ester in said animal feed is 50 ppm by weight or higher, based on the total weight of said animal feed.
- 20 13. Ingredient, premix or supplement for an animal feed suitable for feeding mammals, birds and fish, comprising an alkyl ester of a fatty acid, wherein said fatty acid has a chain length of 5-12 carbon atoms, and wherein the dosage of said ester is 1 wt.% or higher, based on the total weight of said ingredient, premix or supplement, wherein said ingredient, premix or supplement further comprises vitamins, trace elements, minerals and organic acids.
- 25 14. Ingredient, premix or supplement according to claim 13, for use in decreasing the risk of microbial infections in an animal.
15. Method for increasing the feed efficiency in an animal, comprising feeding a mammal, bird or fish with the animal feed as defined in any of claims 1-12 or with the ingredient, premix or supplement according to claim 13.
- 30 16. Use of an alkyl ester of a medium chain fatty acid or an alkyl ester of a medium chain fatty acid enriched product in an animal feed for the improvement of feed efficiency in an animal, wherein said fatty acid has a chain length of 5-12 carbon atoms and the dosage of said ester in said animal feed is 50 ppm by weight or higher, based on the total weight of said animal feed, and wherein said use is non-therapeutic.
- 35 17. Alkyl ester of a fatty acid or alkyl ester of a fatty acid enriched product for use in an animal feed for decreasing the risk of microbial infections in an animal, wherein said fatty acid has a chain length of 5-12 carbon atoms and the dosage of said ester in said animal feed is 50 ppm by weight or higher, based on the total weight of said animal feed.

40 Patentansprüche

- 45 1. Tierfutter, geeignet zum Füttern von Säugetieren, Vögeln und Fischen, umfassend einen Alkylester einer Fettsäure, wobei die Fettsäure eine Kettenlänge von 5-12 Kohlenstoffatomen aufweist und wobei die Dosierung des Esters in dem Tierfutter 50 Gew.-ppm oder mehr beträgt, bezogen auf das Gesamtgewicht des Tierfutters, zur Verwendung der Verringerung des Risikos von mikrobiellen Infektionen bei einem Tier.
2. Tierfutter nach Anspruch 1, wobei das Alkyl Methyl, Ethyl, Propyl, Butyl oder eine Kombination davon ist.
3. Tierfutter nach Anspruch 1 oder 2, wobei die Fettsäure eine Kettenlänge von 8-12 Kohlenstoffatomen aufweist.
- 50 4. Tierfutter nach einem der vorhergehenden Ansprüche, wobei die Fettsäure Octansäure, Decansäure, Dodecansäure oder eine Kombination davon ist.
- 55 5. Tierfutter nach einem der vorhergehenden Ansprüche, wobei der Ester Methyloctanoat, Methyldecanoat, Methyl-dodecanoat, Ethyloctanoat, Ethyldecanoat, Ethyldodecanoat, Propyloctanoat, Propyldecanoat, Propyldodecanoat, Butyloctanoat, Butyldecanoat, Butyldodecanoat oder eine Kombination davon ist.
6. Tierfutter nach einem der vorhergehenden Ansprüche, wobei die Dosierung des Esters in dem Tierfutter 100

EP 2 381 797 B1

Gew.-ppm oder mehr, vorzugsweise 200 Gew.-ppm oder mehr beträgt, bezogen auf das Gesamtgewicht des Tierfutters.

- 5 7. Tierfutter nach einem der vorhergehenden Ansprüche, wobei die Dosierung des Esters in dem Tierfutter weniger als 5000 Gew.-ppm, vorzugsweise weniger als 1000 Gew.-ppm beträgt, bezogen auf das Gesamtgewicht des Tierfutters.
8. Tierfutter nach einem der vorhergehenden Ansprüche, ferner umfassend organische Säuren und/oder Salze davon.
- 10 9. Tierfutter nach Anspruch 8, wobei die organische Säure und/oder das Salz davon ausgewählt ist aus der Gruppe bestehend aus Pentansäure, Octansäure, Decansäure, Ameisensäure, Essigsäure, Propionsäure, Milchsäure, Buttersäure, Zitronensäure, Äpfelsäure, Fumarsäure, Benzoesäure, Bernsteinsäure, Sorbinsäure, Weinsäure und/oder Salze und Kombinationen davon, wobei das Salz vorzugsweise das Natrium-, Kalium-, Ammonium- oder Calciumsalz der organischen Säuren ist.
- 15 10. Tierfutter nach einem der vorhergehenden Ansprüche, ferner umfassend Gallussäure oder einen Alkylester der Gallussäure.
- 20 11. Tierfutter nach Anspruch 10, wobei der Alkylester der Gallussäure Propylgallat, Pentylgallat, Octylgallat oder Laurylgallat ist.
- 25 12. Tierfutter, geeignet zum Füttern von Säugetieren, Vögeln und Fischen, umfassend eine oder mehrere organische Säuren und einen Alkylester einer Fettsäure, wobei die Fettsäure eine Kettenlänge von 5-12 Kohlenstoffatomen aufweist und wobei die Dosierung des Esters in dem Tierfutter 50 Gew.-ppm oder mehr beträgt, bezogen auf das Gesamtgewicht des Tierfutters.
- 30 13. Zutat, Vormischung oder Ergänzung für ein zur Fütterung von Säugetieren, Vögeln und Fischen geeignetes Tierfutter, umfassend einen Alkylester einer Fettsäure, wobei die Fettsäure eine Kettenlänge von 5-12 Kohlenstoffatomen aufweist und wobei die Dosis des Esters 1 Gew.-% oder mehr beträgt, bezogen auf das Gesamtgewicht der Zutat, Vormischung oder Ergänzung, wobei die Zutat, Vormischung oder Ergänzung ferner Vitamine, Spurenelemente, Mineralien und organischen Säuren umfasst.
- 35 14. Zutat, Vormischung oder Ergänzung nach Anspruch 13 zur Verwendung der Verringerung des Risikos von mikrobiellen Infektionen bei einem Tier.
- 40 15. Verfahren zur Erhöhung der Futtereffizienz bei einem Tier, umfassend das Füttern eines Säugetiers, Vogels oder Fisches mit dem Tierfutter nach einem der Ansprüche 1 bis 12 oder mit der Zutat, Vormischung oder Zusatz nach Anspruch 13.
- 45 16. Verwendung eines Alkylesters einer mittelkettigen Fettsäure oder eines mit einem Alkylester mit einer mittelkettigen Fettsäure angereicherten Produkts in einem Tierfutter zur Verbesserung der Futtereffizienz in einem Tier, wobei die Fettsäure eine Kettenlänge von 5-12 Kohlenstoffatomen aufweist und die Dosierung des Esters in dem Tierfutter 50 Gew.-ppm oder mehr beträgt, bezogen auf das Gesamtgewicht des Tierfutters, wobei die Verwendung ist nicht therapeutisch ist.
- 50 17. Alkylester einer Fettsäure oder ein mit einem Alkylester mit einer Fettsäure angereichertes Produkt zur Verwendung in einem Tierfutter zur Verringerung des Risikos von mikrobiellen Infektionen bei einem Tier, wobei die Fettsäure eine Kettenlänge von 5-12 Kohlenstoffatomen aufweist und die Dosierung des Esters in dem Tierfutter 50 Gew.-ppm oder mehr beträgt, bezogen auf das Gesamtgewicht des Tierfutters.

Revendications

- 55 1. Aliment pour animaux adapté à l'alimentation de mammifères, d'oiseaux et de poissons, comprenant un ester alkylique d'un acide gras, dans lequel ledit acide gras présente une longueur de chaîne de 5 à 12 atomes de carbone, et dans lequel le dosage dudit ester dans ledit aliment pour animaux est supérieur ou égal à 50 ppm en poids, sur la base du poids total dudit aliment pour animaux, destiné à être utilisé afin de réduire le risque d'infection microbienne chez un animal.

EP 2 381 797 B1

2. Aliment pour animaux selon la revendication 1, dans lequel ledit alkyle est du méthyle, de l'éthyle, du propyle, du butyle ou un mélange de ceux-ci.
- 5 3. Aliment pour animaux selon la revendication 1 ou 2, dans lequel ledit acide gras présente une longueur de chaîne de 8 à 12 atomes de carbone.
4. Aliment pour animaux selon l'une quelconque des revendications précédentes, dans lequel ledit acide gras est de l'acide octanoïque, de l'acide décanoïque, de l'acide dodécanoïque ou un mélange de ceux-ci.
- 10 5. Aliment pour animaux selon l'une quelconque des revendications précédentes, dans lequel ledit ester est de l'octanoate de méthyle, du décanoate de méthyle, du dodécanoate de méthyle, de l'octanoate d'éthyle, du décanoate d'éthyle, du dodécanoate d'éthyle, de l'octanoate de propyle, du décanoate de propyle, du dodécanoate de propyle, de l'octanoate de butyle, du décanoate de butyle, du dodécanoate de butyle ou un mélange de ceux-ci.
- 15 6. Aliment pour animaux selon l'une quelconque des revendications précédentes, dans lequel le dosage dudit ester dans ledit aliment pour animaux est supérieur ou égal à 100 ppm en poids ou, de préférence, supérieur ou égal à 200 ppm en poids, sur la base du poids total dudit aliment pour animaux.
- 20 7. Aliment pour animaux selon l'une quelconque des revendications précédentes, dans lequel le dosage dudit ester dans ledit aliment pour animaux est inférieur à 5000 ppm en poids, de préférence, inférieur à 1000 ppm en poids, sur la base du poids total dudit aliment pour animaux.
- 25 8. Aliment pour animaux selon l'une quelconque des revendications précédentes, comprenant en outre des acides organiques et/ou des sels de ces derniers.
- 30 9. Aliment pour animaux selon la revendication 8, dans lequel ledit acide organique et/ou ledit sel de celui-ci est choisi à partir du groupe constitué par de l'acide pentanoïque, octanoïque, décanoïque, formique, acétique, propionique, lactique, butyrique, citrique, malique, fumarique, benzoïque, succinique, sorbique, tartrique et/ou un sel ou des mélanges de ces derniers, dans lequel ledit sel est, de préférence, le sel de sodium, de potassium, d'ammonium ou de calcium desdits acides organiques.
- 35 10. Aliment pour animaux selon l'une quelconque des revendications précédentes, comprenant en outre l'acide gallique ou un ester alkylique d'acide gallique.
- 40 11. Aliment pour animaux selon la revendication 10, dans lequel ledit ester alkylique d'acide gallique est du gallate de propyle, du gallate de pentyle, du gallate d'octyle ou du gallate de lauryle.
- 45 12. Aliment pour animaux adapté à l'alimentation de mammifères, d'oiseaux et de poissons, comprenant un ou plusieurs acides organiques et un ester alkylique d'un acide gras, dans lequel ledit acide gras présente une longueur de chaîne de 5 à 12 atomes de carbone, et dans lequel le dosage dudit ester dans ledit aliment pour animaux est supérieur ou égal à 50 ppm en poids, sur la base du poids total dudit aliment pour animaux.
- 50 13. Ingrédient, prémélange ou complément pour un aliment destiné à des animaux adapté à l'alimentation de mammifères, d'oiseaux et de poissons, comprenant un ester alkylique d'un acide gras, dans lequel ledit acide gras présente une longueur de chaîne de 5 à 12 atomes de carbone, et dans lequel le dosage dudit ester est supérieur ou égal à 1 % en poids sur la base du poids total dudit ingrédient, prémélange ou complément, dans lequel ledit ingrédient, prémélange ou complément comprend en outre des vitamines, des oligo-éléments, des minéraux et des acides organiques.
- 55 14. Ingrédient, prémélange ou complément selon la revendication 13, destiné à être utilisé afin d'assurer la réduction du risque d'infection microbienne chez un animal.
15. Procédé d'augmentation de l'efficacité alimentaire chez un animal, comprenant l'alimentation d'un mammifère, oiseau ou poisson avec l'aliment pour animaux selon l'une quelconque des revendications 1 à 12 ou avec l'ingrédient, le prémélange ou le complément selon la revendication 13.
16. Utilisation d'un ester alkylique d'un acide gras à chaîne moyenne ou d'un ester alkylique d'un produit enrichi en acide gras à chaîne moyenne dans un aliment pour animaux afin d'assurer l'amélioration de l'efficacité alimentaire

EP 2 381 797 B1

chez un animal, dans lequel ledit acide gras présente une longueur de chaîne de 5 à 12 atomes de carbone et le dosage dudit ester dans ledit aliment pour animaux est supérieur ou égal à 50 ppm en poids, sur la base du poids total dudit aliment pour animaux, et dans lequel ladite utilisation n'est pas thérapeutique.

- 5 **17.** Ester alkylique d'un acide gras ou ester alkylique d'un produit enrichi en acide gras destiné à une utilisation dans un aliment pour animaux afin de réduire le risque d'infections microbiennes chez un animal, dans lequel ledit acide gras présente une longueur de chaîne de 5 à 12 atomes de carbone et le dosage dudit ester dans ledit aliment pour animaux est supérieur ou égal à 50 ppm en poids sur la base du poids total dudit aliment pour animaux.

10

15

20

25

30

35

40

45

50

55

Figure 1.




Figure 2.





Figure 3.



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 2466663 A [0002]
- EP 1059041 A [0003]
- WO 0197799 A [0004]
- US 4526798 A [0005]
- WO 200600297 A [0006]
- WO 0152837 A [0007]

Non-patent literature cited in the description

- **KARBARA J. J.** *American Oil Chemist's Society*, 01 January 1978, 1-14 [0008]
- **KARBARA J. J.** *American Oil Chemist's Society*, 01 January 1978, 15-24 [0009]
- **K. NIHEI et al.** *J. Agric. Food Chem.*, 2004, vol. 52, 5011-5020 [0012]



1. Állateledel, amely alkalmas emlősök, madarak és halak etetésére, és amely tartalmaz zsírsav-alkilésztert, ahol a zsírsav lánchosszúsága 5-12 szénatom, és ahol az észter dózisa tömegben kifejezve az állateledelben az állateledel teljes tömegére számított 50 ppm vagy magasabb, állatban mikrobiális fertőzés kockázatának csökkentésében történő alkalmazásra.

2. Az 1. igénypont szerinti állateledel, ahol az alkil metil, etil, propil, butil vagy ezek bármilyen kombinációja.

3. Az 1. vagy 2. igénypont szerinti állateledel, ahol a zsírsavlánc hossza 8-12 szénatom.

4. Az előző igénypontok bármelyike szerinti állateledel, ahol a zsírsav oktánsav, dekánsav, dodekánsav vagy ezek bármilyen kombinációja.

5. Az előző igénypontok bármelyike szerinti állateledel, ahol az észter metil-oktanoát, metil-dekanoát, metil-dodekanoát, etil-oktanoát, etil-dekanoát, etil-dodekanoát, propil-oktanoát, propil-dekanoát, propil-dodekanoát, butil-oktanoát, butil-dekanoát, butil-dodekanoát vagy ezek bármilyen kombinációja.

6. Az előző igénypontok bármelyike szerinti állateledel, ahol az észter dózisa tömegben kifejezve az állateledelben az állateledel teljes tömegére számított 100 ppm vagy magasabb, előnyösen 200 ppm vagy magasabb.

7. Az előző igénypontok bármelyike szerinti állateledel, ahol az észter dózisa tömegben kifejezve az állateledelben az állateledel teljes tömegére számított kevesebb, mint 5000 ppm, előnyösen kevesebb, mint 1000 ppm.

8. Az előző igénypontok bármelyike szerinti állateledel, amely tartalmaz szerves savakat és/vagy azok sóit is.

9. A 8. igénypont szerinti állateledel, ahol a szerves sav és/vagy annak sója a következőkből álló csoportból van kiválasztva: pentánsav, oktánsav, dekánsav, metánsav, ecetsav, propánsav, tejsav, butánsav, citromsav, almasav, fűmársav, benzoésav, borostyánkősav, szorbátsav, borkősav és/vagy sója és ezek kombinációi, ahol a só előnyösen a szerves sav nátriumsója, káliumsója, ammóniumsója vagy kalciumsója.

10. Az előző igénypontok bármelyike szerinti állateledel, amely tartalmaz galluszsavat vagy galluszsav-alkilésztert is.

11. A 10. igénypont szerinti állateledel, ahol a galluszsav-alkilészter propil-gallát, pentil-gallát, oktil-gallát vagy lautil-gallát.

12. Állateledel, amely alkalmas emlősök, madarak és halak etetésére, amely tartalmaz egy vagy több szerves savat és zsírsav-alkilésztert, ahol a zsírsav lánchosszúsága 5-12 szénatom, és ahol az észter dózisa tömegben kifejezve az állateledelben az állateledel teljes tömegére számított 50 ppm vagy magasabb.

13. Összetevő, előkeverék vagy kiegészítő emlősök, madarak és halak etetésére alkalmas állateledelhez, amely tartalmaz zsírsav-alkilésztert, ahol a zsírsav lánchosszúsága 5-12 szénatom, és ahol az észter dózisa az összetevő, előkeverék vagy kiegészítő teljes tömegére számítva 1 tömeg% vagy magasabb, ahol az összetevő, előkeverék vagy kiegészítő vitaminokat, nyomelemeket, ásványi anyagokat és szerves savakat is tartalmaz.

14. A 13. igénypont szerinti összetevő, előkeverék vagy kiegészítő, állatban mikrobiális fertőzés kockázatának csökkentésében történő alkalmazásra.

15. Eljárás táplálás hatékonyságának növelésére állatban, amely magában foglalja emlős, madár vagy hal etetését az 1-12. igénypontok bármelyike szerinti állateledellel vagy a 13. igénypont szerinti összetevővel, előkeverékkel vagy kiegészítővel.

16. Közepes lánchosszúságú zsírsav alkilészterének vagy közepes lánchosszúságú zsírsav alkilészterével dúsított termék alkalmazása állateledelben, táplálás hatékonyságának növelésére állatban, ahol a zsírsav lánchosszúsága 5-12 szénatom, és az észter dózisa tömegben kifejezve az állateledelben az állateledel teljes tömegére számított 50 ppm vagy magasabb, és ahol az alkalmazás terápiás alkalmazástól különbözik.

17. Közepes lánchosszúságú zsírsav alkilésztere vagy közepes lánchosszúságú zsírsav alkilészterével dúsított termék állatban mikrobiális fertőzés kockázatának csökkentésére szolgáló állateledelben történő alkalmazásra, ahol a zsírsav lánchosszúsága 5-12 szénatom, és az észter dózisa tömegben kifejezve az állateledelben az állateledel teljes tömegére számított 50 ppm vagy magasabb.