



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/07/03
(87) Date publication PCT/PCT Publication Date: 2020/01/09
(85) Entrée phase nationale/National Entry: 2021/01/05
(86) N° demande PCT/PCT Application No.: US 2019/040496
(87) N° publication PCT/PCT Publication No.: 2020/010197
(30) Priorités/Priorities: 2018/07/05 (US62/694,138);
2018/11/05 (US62/755,845)

(51) Cl.Int./Int.Cl. *C07D 487/04* (2006.01),
A61K 31/4985 (2006.01), *A61P 25/28* (2006.01),
A61P 29/00 (2006.01), *A61P 35/00* (2006.01),
A61P 9/00 (2006.01), *C07D 519/00* (2006.01)


(71) Demandeur/Applicant:
INCYTE CORPORATION, US

(72) Inventeurs/Inventors:
HOANG, GIA, US;
WANG, XIAOZHAO, US;
CARLSEN, PETER NIELS, US;
GAN, PEI, US;
LI, YONG, US;
...

(74) Agent: SMART & BIGGAR LLP

(54) Titre : DERIVES DE PYRAZINE FUSIONNES EN TANT QU'INHIBITEURS D'A2A/A2B

(54) Title: FUSED PYRAZINE DERIVATIVES AS A2A / A2B INHIBITORS



(I),

(57) Abrégé/Abstract:

This application relates to compounds of Formula (I) or pharmaceutically acceptable salts thereof, which modulate the activity of adenosine receptors, such as subtypes A2A and A2B receptors, and are useful in the treatment of diseases related to the activity of adenosine receptors including, for example, cancer, inflammatory diseases, cardiovascular diseases, and neurodegenerative diseases.

(72) **Inventeurs(suite)/Inventors(continued):** QI, CHAO, US; WU, LIANGXING, US; YAO, WENQING, US; YU, ZHIYONG, US; ZHU, WENYU, US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

09 January 2020 (09.01.2020)



(10) International Publication Number

WO 2020/010197 A1

(51) International Patent Classification:

C07D 487/04 (2006.01) A61P 29/00 (2006.01)

C07D 519/00 (2006.01) A61P 35/00 (2006.01)

A61P 9/00 (2006.01) A61K 31/4985 (2006.01)

A61P 25/28 (2006.01)

(21) International Application Number:

PCT/US2019/040496

(22) International Filing Date:

03 July 2019 (03.07.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/694,138 05 July 2018 (05.07.2018) US

62/755,845 05 November 2018 (05.11.2018) US

(71) Applicant: INCYTE CORPORATION [US/US]; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US).

(72) Inventor; and

(71) Applicant: HOANG, Gia [US/US]; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US).

(72) Inventors: WANG, Xiaozhao; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). CARLSEN, Peter Niels; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). GAN, Pei; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). LI, Yong; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). QI, Chao; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). WU, Liangxing; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). YAO, Wenqing; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). YU, Zhiyong; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). ZHU, Wenyu; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US).

(74) Agent: KARNAS, Elizabeth et al.; Fish & Richardson, P.O. Box 1022, Minneapolis, Minnesota 55440 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) Title: FUSED PYRAZINE DERIVATIVES AS A2A / A2B INHIBITORS



(I),

(57) Abstract: This application relates to compounds of Formula (I) or pharmaceutically acceptable salts thereof, which modulate the activity of adenosine receptors, such as subtypes A2A and A2B receptors, and are useful in the treatment of diseases related to the activity of adenosine receptors including, for example, cancer, inflammatory diseases, cardiovascular diseases, and neurodegenerative diseases.

WO 2020/010197 A1

FUSED PYRAZINE DERIVATIVES AS A2A / A2B INHIBITORS

TECHNICAL FIELD

The present invention provides fused pyrazine derivatives that modulate the activity of adenosine receptors, such as subtypes A2A and A2B, and are useful in the treatment of diseases related to the activity of adenosine receptors including, for example, cancer, inflammatory diseases, cardiovascular diseases, and neurodegenerative diseases.

BACKGROUND

Adenosine is an extracellular signaling molecule that can modulate immune responses through many immune cell types. Adenosine was first recognized as a physiologic regulator of coronary vascular tone by Drury and Szent-György (Sachdeva, S. and Gupta, M. *Saudi Pharmaceutical Journal*, 2013, 21, 245–253), however it was not until 1970 that Sattin and Rall showed that adenosine regulates cell function via occupancy of specific receptors on the cell surface (Sattin, A., and Rall, T.W., 1970. *Mol. Pharmacol.* 6, 13–23; Hasko', G., et al., 2007, *Pharmacol. Ther.* 113, 264–275).

Adenosine plays a vital role in various other physiological functions. It is involved in the synthesis of nucleic acids, when linked to three phosphate groups; it forms ATP, the integral component of the cellular energy system. Adenosine can be generated by the enzymatic breakdown of extracellular ATP, or can be also released from injured neurons and glial cells by passing the damaged plasma membrane (Tautenhahn, M. et al. *Neuropharmacology*, 2012, 62, 1756–1766). Adenosine produces various pharmacological effects, both in periphery and in the central nervous system, through an action on specific receptors localized on cell membranes (Matsumoto, T. et al. *Pharmacol. Res.*, 2012, 65, 81–90). Alternative pathways for extracellular adenosine generation have been described. These pathways include the production of adenosine from nicotinamide dinucleotide (NAD) instead of ATP by the concerted action of CD38, CD203a and CD73. CD73-independent production of adenosine can also occur by other phosphates such as alkaline phosphatase or prostate-specific phosphatase.

There are four known subtypes of adenosine receptor in humans including A1, A2A, A2B and A3 receptors. A1 and A2A are high affinity receptors, whereas A2B and A3 are low affinity receptors. Adenosine and its agonists can act via one or more of these receptors and can modulate the activity of adenylate cyclase, the enzyme responsible for increasing cyclic AMP (cAMP). The different receptors have differential stimulatory and inhibitory effects on this enzyme. Increased intracellular concentrations of cAMP can suppress the activity of immune and inflammatory cells (Livingston, M. et al., *Inflamm. Res.*, 2004, 53, 171–178).

The A2A adenosine receptor can signal in the periphery and the CNS, with agonists explored as anti-inflammatory drugs and antagonists explored for neurodegenerative diseases (Carlsson, J. et al., *J. Med. Chem.*, 2010, 53, 3748–3755). In most cell types the A2A subtype inhibits intracellular calcium levels whereas the A2B potentiates them. The A2A receptor generally appears to inhibit inflammatory response from immune cells (Borrmann, T. et al., *J. Med. Chem.*, 2009, 52(13), 3994–4006).

A2B receptors are highly expressed in the gastrointestinal tract, bladder, lung and on mast cells (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842–857). The A2B receptor, although structurally closely related to the A2A receptor and able to activate adenylate cyclase, is functionally different. It has been postulated that this subtype may utilize signal transduction systems other than adenylate cyclase (Livingston, M. et al., *Inflamm. Res.*, 2004, 53, 171–178). Among all the adenosine receptors, the A2B adenosine receptor is a low affinity receptor that is thought to remain silent under physiological conditions and to be activated in consequence of increased extracellular adenosine levels (Ryzhov, S. et al. *Neoplasia*, 2008, 10, 987–995). Activation of A2B adenosine receptor can stimulate adenylate cyclase and phospholipase C through activation of Gs and Gq proteins, respectively. Coupling to mitogen activated protein kinases has also been described (Borrmann, T. et al., *J. Med. Chem.*, 2009, 52(13), 3994–4006).

In the immune system, engagement of adenosine signaling can be a critical regulatory mechanism that protects tissues against excessive immune reactions. Adenosine can negatively modulate immune responses through many immune cell types, including T-cells, natural-killer cells, macrophages, dendritic cells, mast cells

and myeloid-derived suppressor cells (Allard, B. et al. *Current Opinion in Pharmacology*, 2016, 29, 7–16).

In tumors, this pathway is hijacked by tumor micro-environments and sabotages the antitumor capacity of immune system, promoting cancer progression. In the tumor micro-environment, adenosine was mainly generated from extracellular ATP by CD39 and CD73. Multiple cell types can generate adenosine by expressing CD39 and CD73. This is the case for tumor cells, T-effector cells, T-regulatory cells, tumor associated macrophages, myeloid derived suppressive cells (MDSCs), endothelial cells, cancer- associated fibroblast (CAFs) and mesenchymal stromal/stem cells (MSCs). Hypoxia, inflammation and other immune-suppressive signaling in tumor micro-environment can induce expression of CD39, CD73 and subsequent adenosine production. As a result, adenosine level in solid tumors is unusually high compared to normal physiological conditions.

A2A are mostly expressed on lymphoid-derived cells, including T-effector cells, T regulatory cells and nature killing cells. Blocking A2A receptor can prevent downstream immunosuppressive signals that temporarily inactivate T cells. A2B receptors are mainly expressed on monocyte-derived cells including dendritic cells, tumor-associated macrophages, myeloid derived suppressive cells (MDSCs), and mesenchymal stromal/stem cells (MSCs). Blocking A2B receptor in preclinical models can suppress tumor growth, block metastasis, and increase the presentation of tumor antigens.


In terms of safety profile of ADORA2A/ADORA2B (A2A/A2B) blockage, the A2A and A2B receptor knockout mice are all viable, showing no growth abnormalities and are fertile (Allard, B. et al. *Current Opinion in Pharmacology*, 2016, 29, 7–16). A2A KO mice displayed increased levels of pro-inflammatory cytokines only upon challenge with LPS and no evidence of inflammation at baseline (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842-857). A2B KO mice exhibited normal platelet, red blood, and white cell counts but increased inflammation at baseline (TNF-alpha, IL-6) in naive A2B KO mice (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842-857). Exaggerated production of TNF-alpha and IL-6 was detected following LPS treatment. A2B KO mice also exhibited increased vascular adhesion molecules that mediate inflammation as well leukocyte

adhesion/rolling; enhanced mast-cell activation; increased sensitivity to IgE-mediated anaphylaxis and increased vascular leakage and neutrophil influx under hypoxia (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842-857).

In summary, there is a need to develop new adenosine receptor selective ligands, such as for subtypes A2A and A2B, for the treatment of diseases such as cancer, inflammatory diseases, cardiovascular diseases and neurodegenerative diseases. This application is directed to this need and others.

SUMMARY

The present invention relates to, *inter alia*, compounds of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein constituent members are defined herein.

The present invention further provides pharmaceutical compositions comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention further provides methods of inhibiting an activity of an adenosine receptor, comprising contacting the receptor with a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

The present invention further provides methods of treating a disease or a disorder associated with abnormal expression of adenosine receptors, comprising administering to said patient a therapeutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

The present invention further provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, for use in any of the methods described herein.


The present invention further provides use of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, for the preparation of a medicament for use in any of the methods described herein.

5

DETAILED DESCRIPTION

Compounds

The present application provides, *inter alia*, compounds of Formula (I):



(I)

or a pharmaceutically acceptable salt thereof, wherein:

10 R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $C(=NR^{e1})R^{b1}$, $C(=NR^{e1})NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl of R^1 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents;

15 each R^{a1} , R^{b1} , R^{c1} , and R^{d1} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl, wherein the C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl of R^{a1} , R^{b1} , R^{c1} , and R^{d1} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents;

20 each R^{e1} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{1A} is independently selected from OH, CN, halo, C_{1-3} alkyl, C_{2-3} alkenyl, C_{2-3} alkynyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, C_{1-3} haloalkoxy, amino, C_{1-3} alkylamino, and $di(C_{1-3}$ alkyl)amino;

X^3 is N or CR^3 ;

25 R^3 is selected from H, D, halo, OH, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, cyano- C_{1-3} alkyl, HO- C_{1-3} alkyl,

- C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;
- 10 Cy¹ is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of Cy¹ are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R⁷ substituents;
- each R⁷ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a7}, SR^{a7}, NHOR^{a7}, C(O)R^{b7}, C(O)NR^{c7}R^{d7}, C(O)NR^{c7}(OR^{a7}), C(O)OR^{a7}, OC(O)R^{b7}, OC(O)NR^{c7}R^{d7}, NR^{c7}R^{d7}, NR^{c7}NR^{c7}R^{d7}, NR^{c7}C(O)R^{b7}, NR^{c7}C(O)OR^{a7}, NR^{c7}C(O)NR^{c7}R^{d7}, C(=NR^{e7})R^{b7}, C(=NOH)R^{b7}, C(=NCN)R^{b7}, C(=NR^{e7})NR^{c7}R^{d7}, NR^{c7}C(=NR^{e7})NR^{c7}R^{d7}, NR^{c7}C(=NOH)NR^{c7}R^{d7}, NR^{c7}C(=NCN)NR^{c7}R^{d7}, NR^{c7}C(=NR^{e7})R^{b7}, NR^{c7}S(O)NR^{c7}R^{d7}, NR^{c7}S(O)R^{b7}, NR^{c7}S(O)₂R^{b7}, NR^{c7}S(O)(=NR^{e7})R^{b7}, NR^{c7}S(O)₂NR^{c7}R^{d7}, S(O)R^{b7}, S(O)NR^{c7}R^{d7}, S(O)₂R^{b7}, S(O)₂NR^{c7}R^{d7}, OS(O)(=NR^{e7})R^{b7}, OS(O)₂R^{b7}, SF₅, P(O)R^{f7}R^{g7}, OP(O)(OR^{h7})(ORⁱ⁷), P(O)(OR^{h7})(ORⁱ⁷), and BR^{j7}R^{k7}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁷ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{7A} substituents;
- 30 each R^{a7}, R^{b7}, R^{c7}, and R^{d7} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-

C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a7}, R^{b7}, R^{c7} and R^{d7} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{7A} substituents;

or, any R^{c7} and R^{d7}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{7A} substituents;

each R^{e7} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f7} and R^{g7} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h7} and Rⁱ⁷ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j7} and R^{k7} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j7} and R^{k7} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{7A} is independently selected from OH, NO₂, CN, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, cyano-C₁₋₆ alkyl, HO-C₁₋₆ alkyl, C₁₋₆ alkoxy-C₁₋₆ alkyl, C₃₋₅ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, amino, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, thio, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, carbamyl, C₁₋₆ alkylcarbamyl, di(C₁₋₆ alkyl)carbamyl, carboxy, C₁₋₆ alkylcarbonyl, C₁₋₄

alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxycarbonylamino, C₁₋₆ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₆ alkylaminocarbonyloxy, di(C₁₋₆ alkyl)aminocarbonyloxy, C₁₋₆ alkylsulfonylamino, aminosulfonyl, C₁₋₆ alkylaminosulfonyl, di(C₁₋₆ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₆ alkylaminosulfonylamino, di(C₁₋₆ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₆ alkylaminocarbonylamino, and di(C₁₋₆ alkyl)aminocarbonylamino;

R² is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a2}, SR^{a2}, NHOR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)NR^{c2}(OR^{a2}), C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, C(=NR^{e2})R^{b2}, C(=NOH)R^{b2}, C(=NCN)R^{b2}, C(=NR^{e2})NR^{c2}R^{d2}, NR^{c2}C(=NR^{e2})NR^{c2}R^{d2}, NR^{c2}C(=NOH)NR^{c2}R^{d2}, NR^{c2}C(=NCN)NR^{c2}R^{d2}, NR^{c2}C(=NR^{e2})R^{b2}, NR^{c2}S(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)(=NR^{e2})R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, S(O)₂NR^{c2}R^{d2}, OS(O)(=NR^{e2})R^{b2}, OS(O)₂R^{b2}, SF₅, P(O)R^{f2}R^{g2}, OP(O)(OR^{h2})(ORⁱ²), P(O)(OR^{h2})(ORⁱ²), and BR^{j2}R^{k2}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R² are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

R⁴ is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a4}, SR^{a4}, NHOR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)NR^{c4}(OR^{a4}), C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, C(=NR^{e4})R^{b4}, C(=NOH)R^{b4}, C(=NCN)R^{b4}, C(=NR^{e4})NR^{c4}R^{d4}, NR^{c4}C(=NR^{e4})NR^{c4}R^{d4}, NR^{c4}C(=NOH)NR^{c4}R^{d4}, NR^{c4}C(=NCN)NR^{c4}R^{d4}, NR^{c4}C(=NR^{e4})R^{b4}, NR^{c4}S(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)(=NR^{e4})R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4},

$S(O)_2NR^{c4}R^{d4}$, $OS(O)(=NR^{e4})R^{b4}$, $OS(O)_2R^{b4}$, SF_5 , $P(O)R^{f4}R^{g4}$, $OP(O)(OR^{h4})(OR^{i4})$,
 $P(O)(OR^{h4})(OR^{i4})$, and $BR^{j4}R^{k4}$, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀
 aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl,
 C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆
 5 alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally
 substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4A} substituents;

provided that:

(a) when R² is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered
 heteroaryl and 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀
 10 cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of
 R² are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently
 selected R^{2A} substituents;

then R⁴ is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆
 alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl,
 15 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆
 alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered
 heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a4}, SR^{a4}, NHOR^{a4}, C(O)R^{b4},
 C(O)NR^{c4}R^{d4}, C(O)NR^{c4}(OR^{a4}), C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4},
 NR^{c4}R^{d4}, NR^{c4}NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4},
 20 C(=NR^{e4})R^{b4}, C(=NOH)R^{b4}, C(=NCN)R^{b4}, C(=NR^{e4})NR^{c4}R^{d4},
 NR^{c4}C(=NR^{e4})NR^{c4}R^{d4}, NR^{c4}C(=NOH)NR^{c4}R^{d4}, NR^{c4}C(=NCN)NR^{c4}R^{d4},
 NR^{c4}C(=NR^{e4})R^{b4}, NR^{c4}S(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4},
 NR^{c4}S(O)(=NR^{e4})R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4},
 S(O)₂NR^{c4}R^{d4}, OS(O)(=NR^{e4})R^{b4}, OS(O)₂R^{b4}, SF₅, P(O)R^{f4}R^{g4},
 25 OP(O)(OR^{h4})(ORⁱ⁴), P(O)(OR^{h4})(ORⁱ⁴), and BR^{j4}R^{k4}, wherein the C₁₋₆ alkyl,
 C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered
 heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀
 cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10
 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted
 30 with 1, 2, 3, 4, 5, or 6 independently selected R^{4A} substituents;

or, alternatively,

(b) when R^2 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a2} , SR^{a2} , $NHOR^{a2}$, $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$,
 5 $C(O)NR^{c2}(OR^{a2})$, $C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $NR^{c2}R^{d2}$,
 $NR^{c2}NR^{c2}R^{d2}$, $NR^{c2}C(O)R^{b2}$, $NR^{c2}C(O)OR^{a2}$, $NR^{c2}C(O)NR^{c2}R^{d2}$,
 $C(=NR^{e2})R^{b2}$, $C(=NOH)R^{b2}$, $C(=NCN)R^{b2}$, $C(=NR^{e2})NR^{c2}R^{d2}$,
 $NR^{c2}C(=NR^{e2})NR^{c2}R^{d2}$, $NR^{c2}C(=NOH)NR^{c2}R^{d2}$, $NR^{c2}C(=NCN)NR^{c2}R^{d2}$,
 $NR^{c2}C(=NR^{e2})R^{b2}$, $NR^{c2}S(O)NR^{c2}R^{d2}$, $NR^{c2}S(O)R^{b2}$, $NR^{c2}S(O)_2R^{b2}$,
 10 $NR^{c2}S(O)(=NR^{e2})R^{b2}$, $NR^{c2}S(O)_2NR^{c2}R^{d2}$, $S(O)R^{b2}$, $S(O)NR^{c2}R^{d2}$, $S(O)_2R^{b2}$,
 $S(O)_2NR^{c2}R^{d2}$, $OS(O)(=NR^{e2})R^{b2}$, $OS(O)_2R^{b2}$, SF_5 , $P(O)R^{f2}R^{g2}$,
 $OP(O)(OR^{h2})(OR^{i2})$, $P(O)(OR^{h2})(OR^{i2})$, and $BR^{j2}R^{k2}$, wherein the C_{1-6} alkyl,
 C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-
 10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)-
 C_{1-6} alkyl- of R^2 are each optionally substituted with 1, 2, 3, 4, 5, or 6
 15 independently selected R^{2A} substituents;

then R^4 is selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl,
 C_{2-6} alkynyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10
 20 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6}
 alkyl-, CN, NO_2 , OR^{a4} , SR^{a4} , $NHOR^{a4}$, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$,
 $C(O)NR^{c4}(OR^{a4})$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$,
 $NR^{c4}NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$,
 $C(=NR^{e4})R^{b4}$, $C(=NOH)R^{b4}$, $C(=NCN)R^{b4}$, $C(=NR^{e4})NR^{c4}R^{d4}$,
 $NR^{c4}C(=NR^{e4})NR^{c4}R^{d4}$, $NR^{c4}C(=NOH)NR^{c4}R^{d4}$, $NR^{c4}C(=NCN)NR^{c4}R^{d4}$,
 25 $NR^{c4}C(=NR^{e4})R^{b4}$, $NR^{c4}S(O)NR^{c4}R^{d4}$, $NR^{c4}S(O)R^{b4}$, $NR^{c4}S(O)_2R^{b4}$,
 $NR^{c4}S(O)(=NR^{e4})R^{b4}$, $NR^{c4}S(O)_2NR^{c4}R^{d4}$, $S(O)R^{b4}$, $S(O)NR^{c4}R^{d4}$, $S(O)_2R^{b4}$,
 $S(O)_2NR^{c4}R^{d4}$, $OS(O)(=NR^{e4})R^{b4}$, $OS(O)_2R^{b4}$, SF_5 , $P(O)R^{f4}R^{g4}$,
 $OP(O)(OR^{h4})(OR^{i4})$, $P(O)(OR^{h4})(OR^{i4})$, and $BR^{j4}R^{k4}$, wherein the C_{1-6} alkyl,
 C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-
 30 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)-
 C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, 3, 4, 5, or 6
 independently selected R^{4A} substituents;

each R^{a2} , R^{b2} , R^{c2} , and R^{d2} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a2} , R^{b2} , R^{c2} and R^{d2} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

or, any R^{c2} and R^{d2} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

each R^{e2} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{f2} and R^{g2} is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{h2} and R^{i2} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{j2} and R^{k2} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j2} and R^{k2} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

- each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)NR^{c21}(OR^{a21}), C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21}, NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21}, C(=NR^{e21})R^{b21}, C(=NOH)R^{b21}, C(=NCN)R^{b21}, C(=NR^{e21})NR^{c21}R^{d21}, NR^{c21}C(=NR^{e21})NR^{c21}R^{d21}, NR^{c21}C(=NOH)NR^{c21}R^{d21}, NR^{c21}C(=NCN)NR^{c21}R^{d21}, NR^{c21}C(=NR^{e21})R^{b21}, NR^{c21}S(O)NR^{c21}R^{d21}, NR^{c21}S(O)R^{b21}, NR^{c21}S(O)₂R^{b21}, NR^{c21}S(O)(=NR^{e21})R^{b21}, NR^{c21}S(O)₂NR^{c21}R^{d21}, S(O)R^{b21}, S(O)NR^{c21}R^{d21}, S(O)₂R^{b21}, S(O)₂NR^{c21}R^{d21}, OS(O)(=NR^{e21})R^{b21}, OS(O)₂R^{b21}, SF₅, P(O)R^{f21}R^{g21}, OP(O)(OR^{h21})(ORⁱ²¹), P(O)(OR^{h21})(ORⁱ²¹), and BR^{j21}R^{k21}, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{2A} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2B} substituents;
- each R^{a21} , R^{b21} , R^{c21} and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a21} , R^{b21} , R^{c21} and R^{d21} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2B} substituents;
- or, any R^{c21} and R^{d21} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2B} substituents;

each R^{e21} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{f21} and R^{g21} is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{h21} and R^{i21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{j21} and R^{k21} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j21} and R^{k21} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

each R^{2B} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a22} , SR^{a22} , $NHOR^{a22}$, $C(O)R^{b22}$, $C(O)NR^{c22}R^{d22}$, $C(O)NR^{c22}(OR^{a22})$, $C(O)OR^{a22}$, $OC(O)R^{b22}$, $OC(O)NR^{c22}R^{d22}$, $NR^{c22}R^{d22}$, $NR^{c22}NR^{c22}R^{d22}$, $NR^{c22}C(O)R^{b22}$, $NR^{c22}C(O)OR^{a22}$, $NR^{c22}C(O)NR^{c22}R^{d22}$, $C(=NR^{e22})R^{b22}$, $C(=NOH)R^{b22}$, $C(=NCN)R^{b22}$, $C(=NR^{e22})NR^{c22}R^{d22}$, $NR^{c22}C(=NR^{e22})NR^{c22}R^{d22}$, $NR^{c22}C(=NOH)NR^{c22}R^{d22}$, $NR^{c22}C(=NCN)NR^{c22}R^{d22}$, $NR^{c22}C(=NR^{e22})R^{b22}$, $NR^{c22}S(O)NR^{c22}R^{d22}$, $NR^{c22}S(O)R^{b22}$, $NR^{c22}S(O)_2R^{b22}$, $NR^{c22}S(O)(=NR^{e22})R^{b22}$, $NR^{c22}S(O)_2NR^{c22}R^{d22}$, $S(O)R^{b22}$, $S(O)NR^{c22}R^{d22}$, $S(O)_2R^{b22}$, $S(O)_2NR^{c22}R^{d22}$, $OS(O)(=NR^{e22})R^{b22}$, $OS(O)_2R^{b22}$, SF_5 , $P(O)R^{f22}R^{g22}$, $OP(O)(OR^{h22})(OR^{i22})$, $P(O)(OR^{h22})(OR^{i22})$, and $BR^{j22}R^{k22}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered

heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2B} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

each R^{a22}, R^{b22}, R^{c22} and R^{d22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a22}, R^{b22}, R^{c22} and R^{d22} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

or, any R^{c22} and R^{d22}, attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group, wherein the 4-, 5-, 6-, or 7-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

each R^{e22} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f22} and R^{g22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h22} and Rⁱ²² is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j22} and R^{k22} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j22} and R^{k22} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally

substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{2C} is independently selected from OH, NO₂, CN, halo, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, C₁₋₃ haloalkyl, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, C₁₋₃ alkoxy, C₁₋₃ haloalkoxy, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxy carbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxy carbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;

each R^{a4}, R^{b4}, R^{c4}, and R^{d4} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a4}, R^{b4}, R^{c4} and R^{d4} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

or, any R^{c4} and R^{d4}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4A} substituents;

each R^{c4} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f4} and R^{g4} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀

cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h4} and Rⁱ⁴ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j4} and R^{k4} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j4} and R^{k4} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{4A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a41}, SR^{a41}, NHOR^{a41}, C(O)R^{b41}, C(O)NR^{c41}R^{d41}, C(O)NR^{c41}(OR^{a41}), C(O)OR^{a41}, OC(O)R^{b41}, OC(O)NR^{c41}R^{d41}, NR^{c41}R^{d41}, NR^{c41}NR^{c41}R^{d41}, NR^{c41}C(O)R^{b41}, NR^{c41}C(O)OR^{a41}, NR^{c41}C(O)NR^{c41}R^{d41}, C(=NR^{e41})R^{b41}, C(=NOH)R^{b41}, C(=NCN)R^{b41}, C(=NR^{e41})NR^{c41}R^{d41}, NR^{c41}C(=NR^{e41})NR^{c41}R^{d41}, NR^{c41}C(=NOH)NR^{c41}R^{d41}, NR^{c41}C(=NCN)NR^{c41}R^{d41}, NR^{c41}C(=NR^{e41})R^{b41}, NR^{c41}S(O)NR^{c41}R^{d41}, NR^{c41}S(O)R^{b41}, NR^{c41}S(O)₂R^{b41}, NR^{c41}S(O)(=NR^{e41})R^{b41}, NR^{c41}S(O)₂NR^{c41}R^{d41}, S(O)R^{b41}, S(O)NR^{c41}R^{d41}, S(O)₂R^{b41}, S(O)₂NR^{c41}R^{d41}, OS(O)(=NR^{e41})R^{b41}, OS(O)₂R^{b41}, SF₅, P(O)R^{f41}R^{g41}, OP(O)(OR^{h41})(ORⁱ⁴¹),

P(O)(OR^{h41})(ORⁱ⁴¹), and BR^{j41}R^{k41}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{4A} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4B} substituents;

each R^{a41}, R^{b41}, R^{c41} and R^{d41} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-

C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a41}, R^{b41}, R^{c41} and R^{d41} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4B} substituents;

or, any R^{c41} and R^{d41}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4B} substituents;

each R^{e41} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f41} and R^{g41} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h41} and Rⁱ⁴¹ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j41} and R^{k41} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j41} and R^{k41} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{4B} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN,

- NO_2 , $\text{OR}^{\text{a}42}$, $\text{SR}^{\text{a}42}$, $\text{NHOR}^{\text{a}42}$, $\text{C}(\text{O})\text{R}^{\text{b}42}$, $\text{C}(\text{O})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{C}(\text{O})\text{NR}^{\text{c}42}(\text{OR}^{\text{a}42})$,
 $\text{C}(\text{O})\text{OR}^{\text{a}42}$, $\text{OC}(\text{O})\text{R}^{\text{b}42}$, $\text{OC}(\text{O})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{NR}^{\text{c}42}\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$,
 $\text{NR}^{\text{c}42}\text{C}(\text{O})\text{R}^{\text{b}42}$, $\text{NR}^{\text{c}42}\text{C}(\text{O})\text{OR}^{\text{a}42}$, $\text{NR}^{\text{c}42}\text{C}(\text{O})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{C}(=\text{NR}^{\text{e}42})\text{R}^{\text{b}42}$,
 $\text{C}(=\text{NOH})\text{R}^{\text{b}42}$, $\text{C}(=\text{NCN})\text{R}^{\text{b}42}$, $\text{C}(=\text{NR}^{\text{e}42})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{NR}^{\text{c}42}\text{C}(=\text{NR}^{\text{e}42})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$,
5 $\text{NR}^{\text{c}42}\text{C}(=\text{NOH})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{NR}^{\text{c}42}\text{C}(=\text{NCN})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{NR}^{\text{c}42}\text{C}(=\text{NR}^{\text{e}42})\text{R}^{\text{b}42}$,
 $\text{NR}^{\text{c}42}\text{S}(\text{O})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{NR}^{\text{c}42}\text{S}(\text{O})\text{R}^{\text{b}42}$, $\text{NR}^{\text{c}42}\text{S}(\text{O})_2\text{R}^{\text{b}42}$, $\text{NR}^{\text{c}42}\text{S}(\text{O})(=\text{NR}^{\text{e}42})\text{R}^{\text{b}42}$,
 $\text{NR}^{\text{c}42}\text{S}(\text{O})_2\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{S}(\text{O})\text{R}^{\text{b}42}$, $\text{S}(\text{O})\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$, $\text{S}(\text{O})_2\text{R}^{\text{b}42}$, $\text{S}(\text{O})_2\text{NR}^{\text{c}42}\text{R}^{\text{d}42}$,
 $\text{OS}(\text{O})(=\text{NR}^{\text{e}42})\text{R}^{\text{b}42}$, $\text{OS}(\text{O})_2\text{R}^{\text{b}42}$, SF_5 , $\text{P}(\text{O})\text{R}^{\text{f}42}\text{R}^{\text{g}42}$, $\text{OP}(\text{O})(\text{OR}^{\text{h}42})(\text{OR}^{\text{i}42})$,
 $\text{P}(\text{O})(\text{OR}^{\text{h}42})(\text{OR}^{\text{i}42})$, and $\text{BR}^{\text{j}42}\text{R}^{\text{k}42}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,
10 C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered
heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered
heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{4B} are
each optionally substituted with 1, 2, 3, or 4 independently selected R^{4C} substituents;
each $\text{R}^{\text{a}42}$, $\text{R}^{\text{b}42}$, $\text{R}^{\text{c}42}$ and $\text{R}^{\text{d}42}$ is independently selected from H, C_{1-6} alkyl, C_{1-6}
15 haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered
heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl-
 C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered
heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10}
aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl,
20 C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6}
alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of $\text{R}^{\text{a}42}$, $\text{R}^{\text{b}42}$, $\text{R}^{\text{c}42}$ and $\text{R}^{\text{d}42}$
are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4C}
substituents;
or, any $\text{R}^{\text{c}42}$ and $\text{R}^{\text{d}42}$, attached to the same N atom, together with the N atom
25 to which they are attached, form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group,
wherein the 4-, 5-, 6-, or 7-membered heterocycloalkyl group is optionally substituted
with 1, 2, 3, or 4 independently selected R^{4C} substituents;
each $\text{R}^{\text{e}42}$ is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy,
 C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;
30 each $\text{R}^{\text{f}42}$ and $\text{R}^{\text{g}42}$ is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy,
 C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl,
5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-,

C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h42} and Rⁱ⁴² is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j42} and R^{k42} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j42} and R^{k42} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{4C} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a43}, SR^{a43}, NHOR^{a43}, C(O)R^{b43}, C(O)NR^{c43}R^{d43}, C(O)NR^{c43}(OR^{a43}), C(O)OR^{a43}, OC(O)R^{b43}, OC(O)NR^{c43}R^{d43}, NR^{c43}R^{d43}, NR^{c43}NR^{c43}R^{d43}, NR^{c43}C(O)R^{b43}, NR^{c43}C(O)OR^{a43}, NR^{c43}C(O)NR^{c43}R^{d43}, C(=NR^{e43})R^{b43}, C(=NOH)R^{b43}, C(=NCN)R^{b43}, C(=NR^{e43})NR^{c43}R^{d43}, NR^{c43}C(=NR^{e43})NR^{c43}R^{d43}, NR^{c43}C(=NOH)NR^{c43}R^{d43}, NR^{c43}C(=NCN)NR^{c43}R^{d43}, NR^{c43}C(=NR^{e43})R^{b43}, NR^{c43}S(O)NR^{c43}R^{d43}, NR^{c43}S(O)R^{b43}, NR^{c43}S(O)₂R^{b43}, NR^{c43}S(O)(=NR^{e43})R^{b43}, NR^{c43}S(O)₂NR^{c43}R^{d43}, S(O)R^{b43}, S(O)NR^{c43}R^{d43}, S(O)₂R^{b43}, S(O)₂NR^{c43}R^{d43}, OS(O)(=NR^{e43})R^{b43}, OS(O)₂R^{b43}, SF₅, P(O)R^{f43}R^{g43}, OP(O)(OR^{h43})(ORⁱ⁴³), P(O)(OR^{h43})(ORⁱ⁴³), and BR^{j43}R^{k43}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{4C} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents;

each R^{a43}, R^{b43}, R^{c43} and R^{d43} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-

C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a43}, R^{b43}, R^{c43} and R^{d43} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43}, attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group, wherein the 4-, 5-, 6-, or 7-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents;

each R^{e43} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f43} and R^{g43} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h43} and Rⁱ⁴³ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j43} and R^{k43} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j43} and R^{k43} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{4D} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-7 membered heteroaryl)-C₁₋₆ alkyl-, (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN,

- NO_2 , OR^{a44} , SR^{a44} , NHOR^{a44} , $\text{C(O)R}^{\text{b44}}$, $\text{C(O)NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{C(O)NR}^{\text{c44}}(\text{OR}^{\text{a44}})$,
 $\text{C(O)OR}^{\text{a44}}$, $\text{OC(O)R}^{\text{b44}}$, $\text{OC(O)NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{NR}^{\text{c44}}\text{NR}^{\text{c44}}\text{R}^{\text{d44}}$,
 $\text{NR}^{\text{c44}}\text{C(O)R}^{\text{b44}}$, $\text{NR}^{\text{c44}}\text{C(O)OR}^{\text{a44}}$, $\text{NR}^{\text{c44}}\text{C(O)NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{C(=NR}^{\text{e44}})\text{R}^{\text{b44}}$,
 $\text{C(=NOH)R}^{\text{b44}}$, $\text{C(=NCN)R}^{\text{b44}}$, $\text{C(=NR}^{\text{e44}})\text{NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{NR}^{\text{c44}}\text{C(=NR}^{\text{e44}})\text{NR}^{\text{c44}}\text{R}^{\text{d44}}$,
5 $\text{NR}^{\text{c44}}\text{C(=NOH)NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{NR}^{\text{c44}}\text{C(=NCN)NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{NR}^{\text{c44}}\text{C(=NR}^{\text{e44}})\text{R}^{\text{b44}}$,
 $\text{NR}^{\text{c44}}\text{S(O)NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{NR}^{\text{c44}}\text{S(O)R}^{\text{b44}}$, $\text{NR}^{\text{c44}}\text{S(O)}_2\text{R}^{\text{b44}}$, $\text{NR}^{\text{c44}}\text{S(O)(=NR}^{\text{e44}})\text{R}^{\text{b44}}$,
 $\text{NR}^{\text{c44}}\text{S(O)}_2\text{NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{S(O)R}^{\text{b44}}$, $\text{S(O)NR}^{\text{c44}}\text{R}^{\text{d44}}$, $\text{S(O)}_2\text{R}^{\text{b44}}$, $\text{S(O)}_2\text{NR}^{\text{c44}}\text{R}^{\text{d44}}$,
 $\text{OS(O)(=NR}^{\text{e44}})\text{R}^{\text{b44}}$, $\text{OS(O)}_2\text{R}^{\text{b44}}$, SF_5 , $\text{P(O)R}^{\text{f44}}\text{R}^{\text{g44}}$, $\text{OP(O)(OR}^{\text{h44}})(\text{OR}^{\text{i44}})$,
 $\text{P(O)(OR}^{\text{h44}})(\text{OR}^{\text{i44}})$, and $\text{BR}^{\text{j44}}\text{R}^{\text{k44}}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,
10 phenyl, C_{3-7} cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl,
phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6} alkyl-, (5-7 membered heteroaryl)- C_{1-6} alkyl-,
and (4-7 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{4D} are each optionally
substituted with 1, 2, 3, or 4 independently selected R^{4E} substituents;
each R^{a44} , R^{b44} , R^{c44} and R^{d44} is independently selected from H, C_{1-6} alkyl, C_{1-6}
15 haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-7 membered
heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6}
alkyl-, (5-7 membered heteroaryl)- C_{1-6} alkyl-, and (4-7 membered heterocycloalkyl)-
 C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl,
5-7 membered heteroaryl, 4-10 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7}
20 cycloalkyl- C_{1-6} alkyl-, (5-7 membered heteroaryl)- C_{1-6} alkyl-, and (4-7 membered
heterocycloalkyl)- C_{1-6} alkyl- of R^{a44} , R^{b44} , R^{c44} and R^{d44} are each optionally
substituted with 1, 2, 3, or 4 independently selected R^{4E} substituents;
or, any R^{c44} and R^{d44} , attached to the same N atom, together with the N atom
to which they are attached, form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group,
25 wherein the 4-, 5-, 6-, or 7-membered heterocycloalkyl group is optionally substituted
with 1, 2, 3, or 4 independently selected R^{4E} substituents;
each R^{e44} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy,
 C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;
each R^{f44} and R^{g44} is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy,
30 C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-7
membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7}

cycloalkyl-C₁₋₆ alkyl-, (5-7 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h44} and Rⁱ⁴⁴ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-7 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-;


each R^{j44} and R^{k44} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j44} and R^{k44} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl; and

each R^{4E} is independently selected from OH, NO₂, CN, halo, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, C₁₋₃ haloalkyl, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, C₁₋₃ alkoxy, C₁₋₃ haloalkoxy, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxy carbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxy carbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino.

In some embodiments, 1, 2, 3, 4, 5, 6, 7, or 8 hydrogen atoms, attached to carbon atoms of any “alkyl”, “alkenyl”, “alkynyl”, “aryl”, “cycloalkyl”, “heterocycloalkyl”, or “heteroaryl” substituents or “-C₁₋₆ alkyl-”, “alkylene”, “alkenylene”, and “alkynylene” linking groups, are each optionally replaced by a deuterium atom.

In some embodiments, the compound of Formula (I) provided herein, or a pharmaceutically acceptable salt thereof, is a compound of Formula (II):



or a pharmaceutically acceptable salt thereof.

- 5 In some embodiments of Formulas (I) and (II), R¹ is selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₁₋₆ haloalkyl, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₁₋₆ haloalkyl of R¹ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents.

- In some embodiments of Formulas (I) and (II), R¹ is selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₁₋₆ haloalkyl.

In some embodiments of Formulas (I) and (II), R¹ is H or C₁₋₃ alkyl.

In some embodiments of Formulas (I) and (II), R¹ is H.

- In some embodiments of Formulas (I) and (II), R³ is selected from H, D, halo, OH, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino.

- 25 In some embodiments of Formulas (I) and (II), R³ is selected from H, D, halo, OH, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of Formulas (I) and (II), R^3 is CN or H.

In some embodiments of Formulas (I) and (II), R^3 is H.

In some embodiments of Formulas (I) and (II), R^2 is selected from H, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, CN, OR^{a2} , SR^{a2} , C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, and C_{6-10} aryl- C_{1-6} alkyl-, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, and C_{6-10} aryl- C_{1-6} alkyl-, of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (II), R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (II), each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$.

In some embodiments of Formulas (I) and (II), each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$.

In some embodiments of Formulas (I) and (II), each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl-

C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a21}, R^{b21}, R^{c21} and R^{d21} are each optionally substituted with 1 or 2 independently selected R^{2B} substituents.

In some embodiments of Formulas (I) and (II), each R^{a21}, R^{b21}, R^{c21}, and R^{d21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl.

In some embodiments of Formulas (I) and (II), R² is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of R² are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents; and

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21}, NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21}, S(O)R^{b21}, and S(O)₂R^{b21}.

In some embodiments of Formulas (I) and (II), R² is selected from C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl, wherein the C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl of R² are each optionally substituted with 1 or 2 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (II), each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21}, NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21}, and S(O)R^{b21}.

In some embodiments of Formulas (I) and (II), each R^{a21}, R^{b21}, R^{c21}, and R^{d21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl.

In some embodiments of Formulas (I) and (II), R^2 is selected from C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl, wherein the C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl of R^2 are each optionally substituted with 1 or 2 independently selected R^{2A} substituents;

- 5 each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, and $S(O)R^{b21}$; and
- 10 each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl.

- In some embodiments of Formulas (I) and (II), R^2 is selected from cyclopropyl, pyrazolyl, pyridyl, pyrimidinyl, dihydropyridin-(2H)-yl, and pyridinonyl,
- 15 each of which is optionally substituted with 1 or 2 independently selected R^{2A} substituents.

- In some embodiments of Formulas (I) and (II), each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$,
- 20 $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, and $S(O)R^{b21}$.

- In some embodiments of Formulas (I) and (II), each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

- 25 In some embodiments of Formulas (I) and (II), R^2 is selected from cyclopropyl, pyrazolyl, pyridyl, pyrimidinyl, dihydropyridin-(2H)-yl, and pyridinonyl, each of which is optionally substituted with 1 or 2 independently selected R^{2A} substituents;

- each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$,
- 30 $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, and $S(O)R^{b21}$; and

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of Formulas (I) and (II), R^2 is selected from phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl, wherein the phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21}, NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21}, S(O)R^{b21}, and S(O)₂R^{b21}, wherein said phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl of R^{2A} are optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C₁₋₆ alkyl, and C₁₋₆ haloalkyl; and

each R^{2B} is independently selected from halo, OH, C₁₋₄ alkoxy, C₁₋₄ haloalkoxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, amino, C₁₋₃ alkylamino, and di(C₁₋₃ alkyl)amino.

In some embodiments of Formulas (I) and (II), R^2 is selected from C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl, wherein the C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl of R^2 are each optionally substituted with 1 or 2 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C₁₋₄ alkyl, C₁₋₄ haloalkyl, and C(O)NR^{c21}R^{d21}, wherein said C₁₋₄ alkyl of R^{2A} is optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C₁₋₆ alkyl, and C₁₋₆ haloalkyl; and

each R^{2B} is independently selected from halo and OH.

In some embodiments of Formulas (I) and (II), R^2 is selected from cyclopropyl, oxazolyl, triazolyl, pyrazolyl, pyridyl, pyrimidinyl, dihydropyridin-(2H)-yl, and pyridinonyl, each of which is optionally substituted with 1 or 2 independently selected R^{2A} substituents;

- 5 each R^{2A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, and $S(O)R^{b21}$, wherein said C_{1-6} alkyl of R^{2A} is optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;
- 10 each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and
- each R^{2B} is independently selected from halo and OH.

In some embodiments of Formulas (I) and (II), R^2 is selected from 1-(trifluoromethyl)cycloprop-1-yl, 1-ethyl-1H-pyrazol-5-yl, 1-propyl-1H-pyrazol-5-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, pyrimidin-4-yl, 3,6-dihydropyridin-4-yl-1(2H)-carboxamide, pyridin-4-yl, 4-(1-hydroxyethyl)-2-methyloxazol-5-yl, 2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl, 1-ethyl-1H-1,2,3-triazol-5-yl, and 1-methyl-1H-1,2,3-triazol-5-yl.

In some embodiments of Formulas (I) and (II), R^2 is selected from 1-(trifluoromethyl)cycloprop-1-yl, 1-ethyl-1H-pyrazol-5-yl, 1-propyl-1H-pyrazol-5-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, pyrimidin-4-yl, 3,6-dihydropyridin-4-yl-1(2H)-carboxamide, pyridin-4-yl, 4-(1-hydroxyethyl)-2-methyloxazol-5-yl, 2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl, 1-ethyl-1H-1,2,3-triazol-5-yl, and 1-methyl-1H-1,2,3-triazol-5-yl.

25 In some embodiments of Formulas (I) and (II), R^2 is selected from 1-(trifluoromethyl)cycloprop-1-yl, 1-ethyl-1H-pyrazol-5-yl, 1-propyl-1H-pyrazol-5-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, pyrimidin-4-yl, 3,6-dihydropyridin-4-yl-1(2H)-carboxamide, and pyridin-4-yl.

In some embodiments of Formulas (I) and (II), R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered

heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a4}, SR^{a4}, NHOR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, S(O)R^{b4}, and S(O)₂R^{b4}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents.

In some embodiments of Formulas (I) and (II), each R^{4A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO₂, OR^{a41}, SR^{a41}, NHOR^{a41}, C(O)R^{b41}, C(O)NR^{c41}R^{d41}, C(O)OR^{a41}, OC(O)R^{b41}, OC(O)NR^{c41}R^{d41}, NR^{c41}R^{d41}, NR^{c41}C(O)R^{b41}, NR^{c41}C(O)OR^{a41}, NR^{c41}C(O)NR^{c41}R^{d41}, S(O)R^{b41}, and S(O)₂R^{b41}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, of R^{4A} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4B} substituents.

In some embodiments of Formulas (I) and (II), each R^{4B} is independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-.

In some embodiments of Formulas (I) and (II), each R^{4A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO₂, OR^{a41}, SR^{a41}, NHOR^{a41}, C(O)R^{b41}, C(O)NR^{c41}R^{d41}, C(O)OR^{a41}, OC(O)R^{b41}, OC(O)NR^{c41}R^{d41}, NR^{c41}R^{d41}, NR^{c41}C(O)R^{b41}, NR^{c41}C(O)OR^{a41}, NR^{c41}C(O)NR^{c41}R^{d41}, S(O)R^{b41}, and S(O)₂R^{b41}.

In some embodiments of Formulas (I) and (II), R⁴ is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a4}, SR^{a4}, NHOR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4},

$C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$,
 $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6}
 alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered
 heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered
 5 heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are
 each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;
 and

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6}
 alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10
 10 membered heterocycloalkyl, CN, NO_2 , OR^{a41} , SR^{a41} , $NHOR^{a41}$, $C(O)R^{b41}$,
 $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $OC(O)R^{b41}$, $OC(O)NR^{c41}R^{d41}$, $NR^{c41}R^{d41}$, $NR^{c41}C(O)R^{b41}$,
 $NR^{c41}C(O)OR^{a41}$, $NR^{c41}C(O)NR^{c41}R^{d41}$, $S(O)R^{b41}$, and $S(O)_2R^{b41}$.

In some embodiments of Formulas (I) and (II), R^4 is selected from H, D, halo,
 C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, 5-6 membered heteroaryl,
 15 phenyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$,
 $C(O)OR^{a4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, and $NR^{c4}C(O)NR^{c4}R^{d4}$, wherein the C_{1-6}
 alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, 5-6 membered heteroaryl, phenyl- C_{1-6} alkyl-,
 and (5-6 membered heteroaryl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1
 or 2 independently selected R^{4A} substituents.

20 In some embodiments of Formulas (I) and (II), each R^{4A} is independently
 selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl,
 C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN,
 NO_2 , and OR^{a41} .

25 In some embodiments of Formulas (I) and (II), each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is
 independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6}
 alkynyl.

In some embodiments of Formulas (I) and (II), each R^{a41} is independently
 selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

30 In some embodiments, any R^{c4} and R^{d4} , attached to the same N atom, together
 with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl
 group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted
 with 1, 2, 3, or 4 independently selected R^{4A} substituents.

In some embodiments, any R^{c4} and R^{d4} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1 or 2 independently selected R^{4A} substituents.

- 5 In some embodiments of Formulas (I) and (II), R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, 5-6 membered heteroaryl, phenyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, and $NR^{c4}C(O)NR^{c4}R^{d4}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, 5-6 membered heteroaryl, phenyl- C_{1-6} alkyl-,
10 and (5-6 membered heteroaryl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1 or 2 independently selected R^{4A} substituents;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , and OR^{a41} ;

- 15 each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and

each R^{a41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

- In some embodiments of Formulas (I) and (II), R^4 is selected from C_{1-6} alkyl, phenyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, $C(O)NR^{c4}R^{d4}$, and $NR^{c4}C(O)OR^{a4}$, wherein the C_{1-6} alkyl, phenyl- C_{1-6} alkyl-, and (5-6 membered heteroaryl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1 or 2 independently selected R^{4A} substituents;

- 25 In some embodiments of Formulas (I) and (II), each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , and OR^{a41} .

- In some embodiments of Formulas (I) and (II), each R^{a4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

30 or, any R^{c4} and R^{d4} , attached to the same N atom, together with the N atom to which they are attached, form a 4-6 membered heterocycloalkyl group.

In some embodiments of Formulas (I) and (II), each R^{a41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

In some embodiments of Formulas (I) and (II), R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a4} , SR^{a4} , $NHOR^{a4}$, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl, wherein said C_{1-6} alkyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl of R^{a4} , R^{b4} , R^{c4} , and R^{d4} are each optionally substituted by 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, CN, NO_2 , OR^{a41} , SR^{a41} , $NHOR^{a41}$, $C(O)R^{b41}$, $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $OC(O)R^{b41}$, $OC(O)NR^{c41}R^{d41}$, $NR^{c41}R^{d41}$, $NR^{c41}C(O)R^{b41}$, $NR^{c41}C(O)OR^{a41}$, $NR^{c41}C(O)NR^{c41}R^{d41}$, $S(O)R^{b41}$, and $S(O)_2R^{b41}$, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl of R^{4A} are each optionally substituted by 1 or 2 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} , and R^{d41} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein said C_{1-6} alkyl of R^{a41} , R^{b41} , R^{c41} , and R^{d41} is optionally substituted by 1, 2, or 3 independently selected R^{4B} substituents; and

each R^{4B} is independently selected from halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and di(C_{1-3} alkyl)amino.

In some embodiments of Formulas (I) and (II), R^4 is selected from phenyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, and $NR^{c4}R^{d4}$, wherein the phenyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, and (4-7 membered heterocycloalkyl)- C_{1-3} alkyl- of R^4 are each optionally substituted with 1 or 2 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, phenyl, and 4-7 membered heterocycloalkyl, wherein said C_{1-6} alkyl, phenyl, and 4-7 membered heterocycloalkyl of R^{a4} , R^{b4} , R^{c4} , and R^{d4} are optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, 4-7 membered heterocycloalkyl, and OH, wherein said 4-7 membered heterocycloalkyl of R^{4A} is optionally substituted with 1 or 2 independently selected R^{4B} substituents; and

each R^{4B} is independently selected from halo, OH, and C_{1-4} alkyl

In some embodiments of Formulas (I) and (II), R^4 is selected from phenyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, $C(O)NR^{c4}R^{d4}$, and $NR^{c4}C(O)OR^{a4}$, wherein the C_{1-6} alkyl, phenyl- C_{1-6} alkyl-, and (5-6 membered heteroaryl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1 or 2 independently selected R^{4A} substituents;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , and OR^{a41} ;

each R^{a4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

or, any R^{c4} and R^{d4} , attached to the same N atom, together with the N atom to which they are attached, form a 4-6 membered heterocycloalkyl group; and

each R^{a41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

In some embodiments of Formulas (I) and (II), R^4 is selected from $C(O)NHC_{1-6}$ alkyl, $C(O)$ -azetidiny, $C(O)$ -pyrrolidiny, $C(O)$ -piperidiny, $C(O)N(C_{1-6} alkyl)_2$,

NHC(O)OC₁₋₆ alkyl, (azetidiny)-C₁₋₆ alkyl, (pyridyl)-C₁₋₆ alkyl, (phenyl)-C₁₋₆ alkyl, (fluorophenyl)-C₁₋₆ alkyl, 3,6-dihydro-2H-pyran, NH-(phenyl), and pyridyl.

In some embodiments of Formulas (I) and (II), R⁴ is selected from C(O)NHC₁₋₆alkyl, C(O)-azetidiny, C(O)-pyrrolidiny, C(O)-piperidiny, C(O)N(C₁₋₆ alkyl)₂,
 5 NHC(O)OC₁₋₆ alkyl, (azetidiny)-C₁₋₆ alkyl, (pyridyl)-C₁₋₆ alkyl, (phenyl)-C₁₋₆ alkyl, (fluorophenyl)-C₁₋₆ alkyl, 3,6-dihydro-2H-pyran, NH-(phenyl), pyridyl, and (pyrrolo[3,2-b]pyridiny)-C₁₋₆ alkyl,

wherein each C₁₋₆ alkyl and azetidiny group is optionally substituted by 1 or 2 OH groups; and each pyridyl is optionally substituted by a methylpiperaziny group.

10 In some embodiments of Formulas (I) and (II), R⁴ is selected from C(O)NHC₁₋₆alkyl, C(O)-azetidiny, C(O)-pyrrolidiny, C(O)-piperidiny, C(O)N(C₁₋₆ alkyl)₂, NHC(O)OC₁₋₆ alkyl, (azetidiny)-C₁₋₆ alkyl, (pyridyl)-C₁₋₆ alkyl, (phenyl)-C₁₋₆ alkyl, (fluorophenyl)-C₁₋₆ alkyl, 3,6-dihydro-2H-pyran, NH-(phenyl), and pyridyl,

wherein each C₁₋₆ alkyl and azetidiny group is optionally substituted by 1 or 2
 15 OH groups; and each pyridyl is optionally substituted by a methylpiperaziny group.

In some embodiments of Formulas (I) and (II), R⁴ is selected from C(O)NHCH₂CH₃, C(O)N(CH₃)(CH₂CH₃), C(O)N(CH₂CH₃)₂, NHC(O)OCH₂CH₃, C(O)-azetidiny, C(O)-hydroxyazetidiny, C(O)-pyrrolidiny, C(O)-piperidiny, CH₂-azetidiny, CH₂-pyridyl, CH₂-fluorophenyl, CH(OH)-fluorophenyl, NH-phenyl, 3,6-dihydro-2H-pyran, (methylpiperaziny)pyridiny, and (1H-pyrrolo[3,2-b]pyridin-3-yl)methyl
 20

In some embodiments of Formulas (I) and (II), R⁴ is selected from C(O)NHCH₂CH₃, C(O)N(CH₃)(CH₂CH₃), C(O)N(CH₂CH₃)₂, NHC(O)OCH₂CH₃, C(O)-azetidiny, C(O)-hydroxyazetidiny, C(O)-pyrrolidiny, C(O)-piperidiny, CH₂-azetidiny, CH₂-pyridyl, CH₂-fluorophenyl, CH(OH)-fluorophenyl, NH-phenyl, 3,6-dihydro-2H-pyran, and (methylpiperaziny)pyridiny.
 25

In some embodiments of Formulas (I) and (II), R² is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2},
 30 NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, S(O)R^{b2}, and S(O)₂R^{b2}.

In some embodiments of Formulas (I) and (II), each R^{a2} , R^{b2} , R^{c2} , and R^{d2} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of Formulas (I) and (II), R^2 is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, S(O)R^{b2}, and S(O)₂R^{b2}; and

each R^{a2} , R^{b2} , R^{c2} , and R^{d2} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of Formulas (I) and (II), R^2 is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, and CN.

In some embodiments of Formulas (I) and (II), R^2 is selected from H, difluoroethyl, bromo, and CN.

In some embodiments of Formulas (I) and (II), R^4 is selected from C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, S(O)R^{b4} and S(O)₂R^{b4}, wherein the C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents.

In some embodiments of Formulas (I) and (II), each R^{a4} , R^{c4} , and R^{d4} is independently selected from H and C₁₋₆ alkyl.

In some embodiments of Formulas (I) and (II), each R^{4A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO₂, OR^{a41}, SR^{a41}, NHOR^{a41}, C(O)R^{b41}, C(O)NR^{c41}R^{d41}, C(O)OR^{a41}, OC(O)R^{b41}, OC(O)NR^{c41}R^{d41}, NR^{c41}R^{d41}, NR^{c41}C(O)R^{b41}, NR^{c41}C(O)OR^{a41}, NR^{c41}C(O)NR^{c41}R^{d41}, S(O)R^{b41}, and S(O)₂R^{b41}.

In some embodiments of Formulas (I) and (II), R^4 is selected from C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4},

OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4},
 NR^{c4}C(O)NR^{c4}R^{d4}, S(O)R^{b4} and S(O)₂R^{b4}, wherein the C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀
 cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered
 heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1, 2, 3, or 4
 5 independently selected R^{4A} substituents;

each R^{a4}, R^{b4}, R^{c4}, and R^{d4} is independently selected from H and C₁₋₆ alkyl;

and

each R^{4A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆
 alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10
 10 membered heterocycloalkyl, CN, NO₂, OR^{a41}, SR^{a41}, NHOR^{a41}, C(O)R^{b41},
 C(O)NR^{c41}R^{d41}, C(O)OR^{a41}, OC(O)R^{b41}, OC(O)NR^{c41}R^{d41}, NR^{c41}R^{d41}, NR^{c41}C(O)R^{b41},
 NR^{c41}C(O)OR^{a41}, NR^{c41}C(O)NR^{c41}R^{d41}, S(O)R^{b41}, and S(O)₂R^{b41}.

In some embodiments of Formulas (I) and (II), R⁴ is selected from phenyl-C₁₋₆
 alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, C(O)NR^{c4}R^{d4}, and NR^{c4}C(O)OR^{a4},
 15 wherein each C₁₋₆ alkyl is optionally substituted by 1 or 2 OH groups; and the phenyl-
 C₁₋₆ alkyl- is optionally substituted with 1 or 2 independently selected halo groups.

In some embodiments of Formulas (I) and (II), each R^{a4}, R^{b4}, R^{c4}, and R^{d4} is
 independently selected from H and C₁₋₆ alkyl.

In some embodiments of Formulas (I) and (II), each R⁴ is selected from
 20 phenyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and C(O)NR^{c4}R^{d4}, wherein
 the phenyl-C₁₋₆ alkyl- and (5-6 membered heteroaryl)-C₁₋₆ alkyl- of R⁴ are each
 optionally substituted with 1 or 2 substituents independently selected from OH and
 halo; and

each R^{a4}, R^{c4}, and R^{d4} is independently selected from H and C₁₋₆ alkyl.

25 In some embodiments of Formulas (I) and (II), R⁴ is selected from phenyl-C₁₋₆
 alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, C(O)NR^{c4}R^{d4}, NR^{c4}C(O)OR^{a4}, wherein
 each C₁₋₆ alkyl is optionally substituted by 1 or 2 OH groups; and the phenyl-C₁₋₆
 alkyl- is optionally substituted with 1 or 2 independently selected halo groups; and

each R^{a4}, R^{c4}, and R^{d4} is independently selected from H and C₁₋₆ alkyl.

30 In some embodiments of Formulas (I) and (II), R⁴ is selected from
 C(O)NHCH₂CH₃, -CH₂-pyridyl, CH₂-fluorophenyl, and CH(OH)-fluorophenyl.

In some embodiments of Formula (II), R^2 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a2} , SR^{a2} , $NHOR^{a2}$, $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)NR^{c2}(OR^{a2})$, $C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $NR^{c2}R^{d2}$, $NR^{c2}NR^{c2}R^{d2}$, $NR^{c2}C(O)R^{b2}$, $NR^{c2}C(O)OR^{a2}$, $NR^{c2}C(O)NR^{c2}R^{d2}$, $C(=NR^{e2})R^{b2}$, $C(=NOH)R^{b2}$, $C(=NCN)R^{b2}$, $C(=NR^{e2})NR^{c2}R^{d2}$, $NR^{c2}C(=NR^{e2})NR^{c2}R^{d2}$, $NR^{c2}C(=NOH)NR^{c2}R^{d2}$, $NR^{c2}C(=NCN)NR^{c2}R^{d2}$, $NR^{c2}C(=NR^{e2})R^{b2}$, $NR^{c2}S(O)NR^{c2}R^{d2}$, $NR^{c2}S(O)R^{b2}$, $NR^{c2}S(O)_2R^{b2}$, $NR^{c2}S(O)(=NR^{e2})R^{b2}$, $NR^{c2}S(O)_2NR^{c2}R^{d2}$, $S(O)R^{b2}$, $S(O)NR^{c2}R^{d2}$, $S(O)_2R^{b2}$, $S(O)_2NR^{c2}R^{d2}$, $OS(O)(=NR^{e2})R^{b2}$, $OS(O)_2R^{b2}$, SF_5 , $P(O)R^{f2}R^{g2}$, $OP(O)(OR^{h2})(OR^{i2})$, $P(O)(OR^{h2})(OR^{i2})$, and $BR^{j2}R^{k2}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^2 are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents; and

R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a4} , SR^{a4} , $NHOR^{a4}$, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)NR^{c4}(OR^{a4})$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $C(=NR^{e4})R^{b4}$, $C(=NOH)R^{b4}$, $C(=NCN)R^{b4}$, $C(=NR^{e4})NR^{c4}R^{d4}$, $NR^{c4}C(=NR^{e4})NR^{c4}R^{d4}$, $NR^{c4}C(=NOH)NR^{c4}R^{d4}$, $NR^{c4}C(=NCN)NR^{c4}R^{d4}$, $NR^{c4}C(=NR^{e4})R^{b4}$, $NR^{c4}S(O)NR^{c4}R^{d4}$, $NR^{c4}S(O)R^{b4}$, $NR^{c4}S(O)_2R^{b4}$, $NR^{c4}S(O)(=NR^{e4})R^{b4}$, $NR^{c4}S(O)_2NR^{c4}R^{d4}$, $S(O)R^{b4}$, $S(O)NR^{c4}R^{d4}$, $S(O)_2R^{b4}$, $S(O)_2NR^{c4}R^{d4}$, $OS(O)(=NR^{e4})R^{b4}$, $OS(O)_2R^{b4}$, SF_5 , $P(O)R^{f4}R^{g4}$, $OP(O)(OR^{h4})(OR^{i4})$, $P(O)(OR^{h4})(OR^{i4})$, and $BR^{j4}R^{k4}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6}

alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4A} substituents.

In some embodiments of Formulas (I) and (II), Cy¹ is C₆₋₁₀ aryl which is optionally substituted by 1, 2, 3, or 4 independently selected R⁷ substituents.

5 In some embodiments of Formulas (I) and (II), Cy¹ is phenyl which is optionally substituted by 1 or 2 independently selected R⁷ substituents.

In some embodiments of Formulas (I) and (II), each R⁷ is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, and NO₂.

In some embodiments of Formulas (I) and (II), Cy¹ is phenyl which is
10 optionally substituted by 1 or 2 independently selected R⁷ substituents; and
each R⁷ is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, and NO₂.

In some embodiments of Formulas (I) and (II), Cy¹ is phenyl which is substituted by 1 or 2 independently selected R⁷ substituents; and
15 each R⁷ is independently selected from halo and CN.

In some embodiments of Formulas (I) and (II), Cy¹ is unsubstituted phenyl.

In some embodiments of Formulas (I) and (II), Cy¹ is cyanophenyl.

In some embodiments of Formulas (I) and (II), Cy¹ is 3-cyanophenyl.

In some embodiments of Formulas (I) and (II), Cy¹ is 3-cyanophenyl or 3-
20 cyano-2-fluorophenyl.

In some embodiments of Formulas (I) and (II), Cy¹ is phenyl substituted by C(O)NR^{c7}R^{d7}, wherein R^{c7} and R^{d7} are each independently selected from H and C₁₋₆ alkyl.

In some embodiments of Formulas (I) and (II), Cy¹ is phenyl substituted by
25 C(O)NH₂.

In some embodiments of Formulas (I) and (II), Cy¹ is 3-formylphenyl.

In some embodiments of Formulas (I) and (II):

R¹ is H;

Cy¹ is C₆₋₁₀ aryl which is optionally substituted by 1, 2, 3, or 4 independently
30 selected R⁷ substituents; and

R² is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10

membered heteroaryl, and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (II):

R^1 is H;

5 Cy^1 is phenyl which is optionally substituted by 1 or 2 independently selected R^7 substituents; and

R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments of the compounds of Formulas (I) and (II):

R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl;

R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents;

R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a4} , SR^{a4} , $NHOR^{a4}$, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$;

each R^{c21} and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a41} , SR^{a41} , $NHOR^{a41}$, $C(O)R^{b41}$, $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $OC(O)R^{b41}$, $OC(O)NR^{c41}R^{d41}$, $NR^{c41}R^{d41}$, $NR^{c41}C(O)R^{b41}$, $NR^{c41}C(O)OR^{a41}$, $NR^{c41}C(O)NR^{c41}R^{d41}$, $S(O)R^{b41}$, and $S(O)_2R^{b41}$;

Cy^1 is C_{6-10} aryl which is optionally substituted by 1, 2, 3, or 4 independently selected R^7 substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^4 is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

or, any R^{c4} and R^{d4} , attached to the same N atom, together with the N atom to which they are attached, form a 4-6 membered heterocycloalkyl group;

each R^{a41} , R^{b41} , R^{c41} , and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and

each R^7 is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, and NO_2 .

In some embodiments of the compounds of Formulas (I) and (II):

R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl;

R^2 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, NO_2 , OR^{a2} , SR^{a2} , $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $NR^{c2}R^{d2}$, $NR^{c2}C(O)R^{b2}$, $NR^{c2}C(O)OR^{a2}$, $NR^{c2}C(O)NR^{c2}R^{d2}$, $S(O)R^{b2}$, and $S(O)_2R^{b2}$;

R^4 is selected from C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$ and $S(O)_2R^{b4}$, wherein the C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a41} , SR^{a41} , $NHOR^{a41}$, $C(O)R^{b41}$, $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $OC(O)R^{b41}$, $OC(O)NR^{c41}R^{d41}$, $NR^{c41}R^{d41}$, $NR^{c41}C(O)R^{b41}$,
 5 $NR^{c41}C(O)OR^{a41}$, $NR^{c41}C(O)NR^{c41}R^{d41}$, $S(O)R^{b41}$, and $S(O)_2R^{b41}$;

Cy^1 is C_{6-10} aryl which is optionally substituted by 1, 2, 3, or 4 independently selected R^7 substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

10 or, any R^{c4} and R^{d4} , attached to the same N atom, together with the N atom to which they are attached, form a 4-6 membered heterocycloalkyl group;

each R^{a41} , R^{b41} , R^{c41} , and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and

each R^7 is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, and NO_2 .
 15

In some embodiments of the compounds of Formulas (I) and (II):

R^1 is H;

R^2 is selected from C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl, wherein the C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl of R^2 are each optionally substituted with 1 or 2 independently selected R^{2A} substituents;
 20

R^4 is selected from C_{1-6} alkyl, phenyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, $C(O)NR^{c4}R^{d4}$, and $NR^{c4}C(O)OR^{a4}$, wherein the C_{1-6} alkyl, phenyl- C_{1-6} alkyl-, and (5-6 membered heteroaryl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1 or 2 independently selected R^{4A} substituents;
 25

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$,
 30 $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, and $S(O)R^{b21}$;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , and OR^{a41} ;

Cy^1 is phenyl which is optionally substituted by 1 or 2 independently selected R^7 substituents;

each R^{a4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

or, any R^{c4} and R^{d4} , attached to the same N atom, together with the N atom to which they are attached, form a 4-6 membered heterocycloalkyl group;

each R^{a41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and

each R^7 is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, and NO_2 .

In some embodiments of the compounds of Formulas (I) and (II):

R^1 is H;

R^2 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, and CN;

R^4 is selected from phenyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, $C(O)NR^{c4}R^{d4}$, $NR^{c4}C(O)OR^{a4}$, wherein each C_{1-6} alkyl is optionally substituted by 1 or 2 OH groups; and the phenyl- C_{1-6} alkyl- is optionally substituted with 1 or 2

independently selected halo groups;

Cy^1 is phenyl which is optionally substituted by 1 or 2 independently selected R^7 substituents;

each R^{a4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

or, any R^{c4} and R^{d4} , attached to the same N atom, together with the N atom to which they are attached, form a 4-6 membered heterocycloalkyl group; and

each R^7 is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, and NO_2 .

In some embodiments of the compounds of Formulas (I) and (II):

R^1 is H;

R² is selected from cyclopropyl, pyrazolyl, pyridyl, pyrimidinyl, dihydropyridin-(2H)-yl, and pyridinonyl, each of which is optionally substituted with 1 or 2 independently selected R^{2A} substituents;

R⁴ is selected from C(O)NHC₁₋₆alkyl, C(O)-azetidiny, C(O)-pyrrolidinyl, C(O)-piperidinyl, C(O)N(C₁₋₆ alkyl)₂, NHC(O)OC₁₋₆ alkyl, (azetidiny)-C₁₋₆ alkyl, (pyridyl)-C₁₋₆ alkyl, (phenyl)-C₁₋₆ alkyl, (fluorophenyl)-C₁₋₆ alkyl, 3,6-dihydro-2H-pyranyl, NH-(phenyl), and pyridyl, wherein each C₁₋₆ alkyl and azetidiny group is optionally substituted by 1 or 2 OH groups; and each pyridyl is optionally substituted by a methylpiperazinyl group;

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21}, NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21}, and S(O)R^{b21};

Cy¹ is cyanophenyl; and

each R^{a21}, R^{b21}, R^{c21}, and R^{d21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of the compounds of Formulas (I) and (II):

R¹ is H;


R² is selected from H, difluoroethyl, bromo, and CN;

R⁴ is selected from phenyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, C(O)NR^{c4}R^{d4}, NR^{c4}C(O)OR^{a4}, wherein each C₁₋₆ alkyl is optionally substituted by 1 or 2 OH groups; and the phenyl-C₁₋₆ alkyl- is optionally substituted with 1 or 2 independently selected halo groups;

Cy¹ is cyanophenyl; and

each R^{a4}, R^{c4}, and R^{d4} is independently selected from H and C₁₋₆ alkyl.

In some embodiments of Formulas (I) and (II), the compound is a compound of Formula (IIa):



or a pharmaceutically acceptable salt thereof, wherein variables R^2 , R^3 , R^4 , and Cy^1 are defined according to the definitions provided herein for compounds of Formulas (I) and (II).

In some embodiments of Formula (IIa),

R^3 is H or CN;

Cy^1 is phenyl which is substituted by 1 or 2 independently selected R^7

substituents;

each R^7 is independently selected from halo and CN;

R^2 is selected from phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl, wherein the phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$, wherein said phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl of R^{2A} are optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl;

each R^{2B} is independently selected from halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and $di(C_{1-3}$ alkyl)amino;

R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a4} , SR^{a4} , $NHOR^{a4}$, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl, wherein said C_{1-6} alkyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl of R^{a4} , R^{b4} , R^{c4} , and R^{d4} are each optionally substituted by 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, CN, NO_2 , OR^{a41} , SR^{a41} , $NHOR^{a41}$, $C(O)R^{b41}$, $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $OC(O)R^{b41}$, $OC(O)NR^{c41}R^{d41}$, $NR^{c41}R^{d41}$, $NR^{c41}C(O)R^{b41}$, $NR^{c41}C(O)OR^{a41}$, $NR^{c41}C(O)NR^{c41}R^{d41}$, $S(O)R^{b41}$, and $S(O)_2R^{b41}$, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl of R^{4A} are each optionally substituted by 1 or 2 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} , and R^{d41} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein said C_{1-6} alkyl of R^{a41} , R^{b41} , R^{c41} , and R^{d41} is optionally substituted by 1, 2, or 3 independently selected R^{4B} substituents; and

each R^{4B} is independently selected from halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and di(C_{1-3} alkyl)amino.

In some embodiments of Formula (IIa),

5 R^3 is H or CN;

Cy^1 is 3-cyanophenyl or 3-cyano-2-fluorophenyl;

R^2 is selected from C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl, wherein the C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl of R^2 are each optionally substituted with 1 or 2
10 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C_{1-4} alkyl, C_{1-4} haloalkyl, and $C(O)NR^{c21}R^{d21}$, wherein said C_{1-4} alkyl of R^{2A} is optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and
15 C_{1-6} haloalkyl; and

each R^{2B} is independently selected from halo and OH.

R^4 is selected from phenyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, and $NR^{c4}R^{d4}$,
20 wherein the phenyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, and (4-7 membered heterocycloalkyl)- C_{1-3} alkyl- of R^4 are each optionally substituted with 1 or 2 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, phenyl, and 4-7 membered heterocycloalkyl, wherein said C_{1-6} alkyl, phenyl, and 4-7 membered heterocycloalkyl of R^{a4} , R^{b4} , R^{c4} , and R^{d4} are optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, 4-7 membered heterocycloalkyl, and OH, wherein said 4-7 membered heterocycloalkyl of R^{4A} is optionally substituted
30 with 1 or 2 independently selected R^{4B} substituents; and

each R^{4B} is independently selected from halo, OH, and C_{1-4} alkyl.

In some embodiments of Formula (IIa),

R^3 is H or CN;

Cy^1 is phenyl which is substituted by 1 or 2 independently selected R^7 substituents;


each R^7 is independently selected from halo and CN;

5 R^2 is selected from H, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, and CN;

R^4 is selected from phenyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, and $C(O)NR^{c4}R^{d4}$, wherein the phenyl- C_{1-6} alkyl- and (5-6 membered heteroaryl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1 or 2 substituents independently selected from OH and halo; and


10 each R^{a4} , R^{c4} , and R^{d4} is independently selected from H and C_{1-6} alkyl.

In some embodiments of Formulas (I) and (II), the compound is a compound of Formula (IIb):




or a pharmaceutically acceptable salt thereof, wherein n is an integer from 0 to 4, and
15 wherein variables R^2 , R^4 , and R^7 are defined according to the definitions provided herein for compounds of Formulas (I) and (II).

In some embodiments of Formulas (I) and (II), the compound is a compound of Formula (IIc):



or a pharmaceutically acceptable salt thereof, wherein variables R^2 , R^4 , and R^7 are defined according to the definitions provided herein for compounds of Formulas (I) and (II).

In some embodiments, the compound of Formula (I) provided herein, or a pharmaceutically acceptable salt thereof, is a compound of Formula (III):



or a pharmaceutically acceptable salt thereof.

In some embodiments of Formulas (I) and (III), R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl of R^1 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents.

In some embodiments of Formulas (I) and (III), R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl.

In some embodiments of Formulas (I) and (III), R^1 is H or C_{1-6} alkyl.

In some embodiments of Formulas (I) and (III), R^1 is H.

In some embodiments of Formulas (I) and (III), R^2 is selected from H, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (III), R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (III), each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$,
 5 $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$.

In some embodiments of Formulas (I) and (III), R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10
 10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$,
 15 $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$, wherein said C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R^{2A} are each optionally substituted by 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{241} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein said C_{1-6} alkyl of R^{a21} , R^{241} , R^{c21} , and R^{d21} is optionally substituted by 1, 2, or 3 independently selected R^{2A} substituents; and

each R^{2B} is independently selected from D, halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and di(C_{1-3}
 25 alkyl)amino.

In some embodiments of Formulas (I) and (III), R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4
 30 independently selected R^{2A} substituents; and

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10

membered heterocycloalkyl, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21}, NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21}, S(O)R^{b21}, and S(O)₂R^{b21}.

In some embodiments of Formulas (I) and (III), R² is selected from 5-10
 5 membered heteroaryl and 4-7 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-7 membered heterocycloalkyl of R² are each optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, and OH, wherein said C₁₋₆ alkyl of R^{2A} is optionally substituted by 1, 2, or 3
 10 independently selected R^{2B} substituents; and

each R^{2B} is independently selected from D, halo, and OH.

In some embodiments of Formulas (I) and (III), R² is selected from 5-6
 membered heteroaryl and 4-6 membered heterocycloalkyl, wherein the 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl of R² are each optionally substituted
 15 with 1 or 2 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (III), each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21}, NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21},
 20 and S(O)R^{b21}.

In some embodiments of Formulas (I) and (III), each R^{a21}, R^{b21}, R^{c21}, and R^{d21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of Formulas (I) and (III), R² is selected from 5-6
 25 membered heteroaryl and 4-6 membered heterocycloalkyl, wherein the 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl of R² are each optionally substituted with 1 or 2 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21},
 30 NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21}, and S(O)R^{b21}; and

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

In some embodiments of Formulas (I) and (III), R^2 is a 5-6 membered heteroaryl, optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (III), R^2 is a 5-6 membered heteroaryl.

In some embodiments of Formulas (I) and (III), R^2 is a 5-6 membered heterocycloalkyl.

In some embodiments of Formulas (I) and (III), R^2 is selected from 6-oxo-1,6-dihydropyridin-3-yl, pyrimidin-4-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, 1-methyl-6-oxo-1,6-dihydropyridazin-3-yl, 4-methyloxazol-5-yl, 4-ethyloxazol-5-yl, 3-methylpyridin-4-yl, 4-(2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl, 2-methyl-4-(2,2,2-trifluoro-1-hydroxyethyl)oxazol-5-yl, 1-ethyl-1H-pyrazol-5-yl, 6-hydroxypyridin-3-yl, 2,6-dimethylpyridin-4-yl, 3-methyl-1H-pyrazol-4-yl, [1,2,4]triazolo[4,3-*a*]pyridin-6-yl, oxazol-5-yl, imidazo[1,2-*a*]pyridine-6-yl, 3-fluoropyridin-4-yl, and 1-(methyl-*d*3)-6-oxo-1,6-dihydropyridazin-3-yl.

In some embodiments of Formulas (I) and (III), R^2 is selected from 6-oxo-1,6-dihydropyridin-3-yl, pyrimidin-4-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, 1-methyl-6-oxo-1,6-dihydropyridazin-3-yl, 4-methyloxazol-5-yl, 4-ethyloxazol-5-yl, and 3-methylpyridin-4-yl.

In some embodiments of Formulas (I) and (III), R^2 is pyridinonyl.

In some embodiments of Formulas (I) and (III), R^2 is 6-oxo-1,6-dihydropyridin-3-yl.

In some embodiments of Formulas (I) and (III), R^2 is pyrimidin-4-yl.

In some embodiments of Formulas (I) and (III), R^2 is selected from the group consisting of 1-methyl-6-oxo-1,6-dihydropyridazin-3-yl, pyrimidin-4-yl, and 2,6-dimethylpyridin-4-yl.

In some embodiments of Formulas (I) and (III), R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered

heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1 or 2 independently selected R^{4A} substituents.

In some embodiments of Formulas (I) and (III), each R^{4A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO₂, OR^{a41}, SR^{a41}, NHOR^{a41}, C(O)R^{b41}, C(O)NR^{c41}R^{d41}, C(O)OR^{a41}, OC(O)R^{b41}, OC(O)NR^{c41}R^{d41}, NR^{c41}R^{d41}, NR^{c41}C(O)R^{b41}, NR^{c41}C(O)OR^{a41}, NR^{c41}C(O)NR^{c41}R^{d41}, S(O)R^{b41}, and S(O)₂R^{b41}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{4A} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4B} substituents.

In some embodiments of Formulas (I) and (III), R⁴ is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of Formulas (I) and (III), R⁴ is H.

In some embodiments of Formulas (I) and (III), R² is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of Formulas (I) and (III), R² is H.

In some embodiments of Formulas (I) and (III), R² is 6-oxo-1,6-dihydropyridin-3-yl or imidazo[1,2-a]pyridin-6-yl; and R⁴ is H.

In some embodiments of Formulas (I) and (III), R² is H; and R⁴ is –NHC(O)OC₁₋₆ alkyl.

In some embodiments of Formulas (I) and (III), R⁴ is selected from C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, S(O)R^{b4}, and S(O)₂R^{b4}.

In some embodiments of Formulas (I) and (III), each R^{a4}, R^{b4}, R^{c4}, and R^{d4} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl.

In some embodiments of Formulas (I) and (III), R^4 is selected from $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$, and $S(O)_2R^{b4}$; and

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl.

In some embodiments of Formulas (I) and (III), R^4 is selected from $C(O)NR^{c4}R^{d4}$ and $NR^{c4}C(O)OR^{a4}$.

In some embodiments of Formulas (I) and (III), R^4 is $NR^{c4}C(O)OR^{a4}$.

In some embodiments of Formulas (I) and (III), each R^{a4} , R^{c4} , and R^{d4} is independently selected from H and C_{1-6} alkyl.

In some embodiments of Formulas (I) and (III), R^4 is selected from $C(O)NR^{c4}R^{d4}$ and $NR^{c4}C(O)OR^{a4}$, and

each R^{a4} , R^{c4} , and R^{d4} is independently selected from H and C_{1-6} alkyl.

In some embodiments of Formulas (I) and (III), R^4 is $-NHC(O)OC_{1-6}$ alkyl.

In some embodiments of Formulas (I) and (III), R^4 is $-NHC(O)OCH_2CH_3$.

In some embodiments of Formulas (I) and (III), R^4 is selected from C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, OR^{a4} , $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $NR^{c4}R^{d4}$, $S(O)_2NR^{c4}R^{d4}$, and $S(O)_2R^{b4}$, wherein the C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered

heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a4} , R^{b4} , R^{c4} and R^{d4} are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

- each R^{4A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, OR^{a41} , $C(O)R^{b41}$,
5 $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $NR^{c41}R^{d41}$, $S(O)_2NR^{c41}R^{d41}$, and $S(O)_2R^{b41}$, wherein the C_{1-6} alkyl, 5 C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;
- 10 each R^{a41} , R^{b41} , R^{c41} and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl, wherein the C_{1-6} alkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^{a41} , R^{b41} , R^{c41} and R^{d41} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;
- 15 each R^{4B} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, 5-6 membered heteroaryl, phenyl- C_{1-3} alkyl-, (C_{3-6} cycloalkyl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, (5-6 membered heteroaryl)- C_{1-3} alkyl-, CN, OR^{a42} , $C(O)R^{b42}$, $C(O)NR^{c42}R^{d42}$, $C(O)OR^{a42}$, $NR^{c42}R^{d42}$, $S(O)_2NR^{c42}R^{d42}$, and $S(O)_2R^{b42}$, wherein the C_{1-6} alkyl, phenyl,
20 C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, 5-6 membered heteroaryl, phenyl- C_{1-3} alkyl-, (C_{3-6} cycloalkyl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, (5-6 membered heteroaryl)- C_{1-3} alkyl- of R^{4B} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;
- each R^{a42} , R^{b42} , R^{c42} and R^{d42} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6
25 membered heteroaryl, wherein the C_{1-6} alkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^{a42} , R^{b42} , R^{c42} and R^{d42} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;
- each R^{4C} is independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, CN, OR^{a43} ,
30 $C(O)R^{b43}$, $C(O)NR^{c43}R^{d43}$, $C(O)OR^{a43}$, $NR^{c43}R^{d43}$, $S(O)_2NR^{c43}R^{d43}$, and $S(O)_2R^{b43}$, wherein the C_{1-6} alkyl of R^{4C} is optionally substituted with 1 or 2 independently selected R^{4D} substituents;

each R^{a43} , R^{b43} , R^{c43} and R^{d43} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein the C_{1-6} alkyl of R^{a43} , R^{b43} , R^{c43} and R^{d43} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43} , attached to the same N atom, together with the N atom
 5 to which they are attached, form a 4-, 5-, or 6-membered heterocycloalkyl group, wherein the 4-, 5-, or 6-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents; and

each R^{4D} is independently selected from C_{1-3} alkyl and OH.

In some embodiments of Formulas (I) and (III), R^4 is selected from phenyl- C_{1-3}
 10 C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, OR^{a4} , $C(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the phenyl- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl- and (5-10 membered heteroaryl)- C_{1-3} alkyl- of R^4 are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from phenyl, 4-7
 15 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl-, wherein the phenyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl- of R^{a4} , R^{b4} , R^{c4} and R^{d4} are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, 5-6
 20 membered heteroaryl, (4-10 membered heterocycloalkyl)- C_{1-3} alkyl-, OR^{a41} , $S(O)_2R^{b41}$, and $NR^{c41}R^{d41}$, wherein the C_{1-6} alkyl, 5-6 membered heteroaryl, and (4-10 membered heterocycloalkyl)- C_{1-3} alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6}
 25 haloalkyl, and 5-6 membered heteroaryl, wherein the C_{1-6} alkyl and 5-6 membered heteroaryl of R^{a41} , R^{b41} , R^{c41} and R^{d41} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-6}
 cycloalkyl, 4-7 membered heterocycloalkyl, (5-6 membered heteroaryl)- C_{1-3} alkyl-,
 30 CN , OR^{a42} , $C(O)R^{b42}$, $C(O)OR^{a42}$, and $NR^{c42}R^{d42}$, wherein the C_{1-6} alkyl, C_{3-6} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, and (5-6

membered heteroaryl)-C₁₋₃ alkyl- of R^{4B} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{a42}, R^{b42}, R^{c42} and R^{d42} is independently selected from H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, and 5-7 membered heterocycloalkyl, wherein
 5 the C₁₋₆ alkyl, C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, and 5-7 membered heterocycloalkyl of R^{a42}, R^{b42}, R^{c42} and R^{d42} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{4C} is independently selected from C₁₋₆ alkyl, CN, C(O)NR^{c43}R^{d43}, C(O)OR^{a43}, NR^{c43}R^{d43}, and S(O)₂R^{b43}, wherein the C₁₋₆ alkyl of R^{4C} is optionally
 10 substituted with 1 or 2 independently selected R^{4D} substituents;

each R^{a43}, R^{b43}, R^{c43} and R^{d43} is independently selected from H, C₁₋₆ alkyl, and C₁₋₆ haloalkyl, wherein the C₁₋₆ alkyl of R^{a43}, R^{b43}, R^{c43} and R^{d43} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43}, attached to the same N atom, together with the N atom
 15 to which they are attached, form a 4-, 5-, or 6-membered heterocycloalkyl group, wherein the 4-, 5-, or 6-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents;

each R^{4D} is independently selected from C₁₋₃ alkyl and OH.

In some embodiments of Formulas (I) and (III), R⁴ is selected from C₆₋₁₀ aryl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, OR^{a4}, C(O)R^{b4}, and S(O)₂R^{b4},
 20 wherein the C₆₋₁₀ aryl-C₁₋₆ alkyl- and (5-10 membered heteroaryl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents; and

each R^{4A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, OR^{a41}, SR^{a41}, and NR^{c41}R^{d41}, wherein the C₁₋₆ alkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{4A} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4B} substituents.
 30

In some embodiments of Formulas (I) and (III), R^4 is selected from C_{6-10} aryl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, OR^{a4} , $C(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the C_{6-10} aryl- C_{1-6} alkyl- and (5-10 membered heteroaryl)- C_{1-3} alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

5 each R^{4A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-3} alkyl-, C_{3-10} cycloalkyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-3} alkyl-, CN, OR^{a41} , SR^{a41} , and $NR^{c41}R^{d41}$, wherein the C_{1-6} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-3} alkyl-, C_{3-10} cycloalkyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-3} alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, OR^{a42} ,
15 $COOR^{a42}$, and $NR^{c42}R^{d42}$;

each R^{a4} and R^{b4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, and azetidiny1 wherein the phenyl is optionally substituted with 1 or 2 groups selected from halo, C_{1-6} alkyl, and C_{1-6} haloalkyl;

each R^{a41} , R^{c41} and R^{d41} is independently selected from H and C_{1-6} alkyl; and
20 each R^{a42} , R^{c42} and R^{d42} is independently selected from H and C_{1-6} alkyl.

In some embodiments of Formulas (I) and (III), R^4 is selected from C_{6-10} aryl- C_{1-3} alkyl- and (5-10 membered heteroaryl)- C_{1-3} alkyl- wherein the C_{6-10} aryl- C_{1-3} alkyl- and (5-10 membered heteroaryl)- C_{1-3} alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

25 each R^{4A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, and OR^{a41} ; and

each R^{a41} is independently selected from H and C_{1-3} alkyl.

In some embodiments of Formulas (I) and (III), R^4 is C_6 aryl- C_{1-3} alkyl- optionally substituted with 1, 2, or 3 substituents independently selected from OH and
30 halo.

In some embodiments of Formulas (I) and (III), R^4 is selected from pyridin-2-ylmethyl, 1*H*-Pyrrolo[2,3-*b*]pyridin-1-yl, 7*H*-pyrrolo[2,3-*b*]pyridin-7-yl, 2-

fluorophenoxy, hydroxy(pyridin-2-yl)methyl, 2-(1-methyl-1H-pyrazol-4-yl)benzyl,
 (imidazo[1,2-a]pyridin-8-yl)methyl, (pyrazolo[1,5-a]pyridin-7-yl)methyl, (2H-
 indazol-2-yl)methyl, (1H-indazol-1-yl)methyl, (2,6-difluorophenyl)(hydroxy)methyl,
 (2,5-difluorophenyl)(hydroxy)methyl, (2,3-difluorophenyl)(hydroxy)methyl, (2-
 5 fluorophenyl)(hydroxy)methyl, (2-chlorophenyl)(hydroxy)methyl,
 hydroxy(phenyl)methyl, phenylsulfonyl, azetidine-1-carbonyl, benzo[d]oxazol-4-
 ylmethyl, 2-fluoro-6-(1-methyl-1H-pyrazol-5-yl)benzyl, 2-fluoro-6-((6-methyl-5-oxo-
 2,6-diazaspiro[3.4]octan-2-yl)methyl)benzyl, 2-fluoro-6-((6-
 oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)benzyl, 2-fluoro-6-(((2-
 10 oxopyrrolidin-3-yl)amino)methyl)benzyl, 2-fluoro-6-((3-oxopiperazin-1-
 yl)methyl)benzyl, 2-fluoro-6-(((1-methyl-2-oxopyrrolidin-3-yl)amino)methyl)benzyl,
 2-fluoro-6-(((2-methyl-2H-1,2,3-triazol-4-yl)amino)methyl)benzyl, 2-(((2-
 oxopyrrolidin-3-yl)amino)methyl)benzyl, amino(2,6-difluorophenyl)methyl, (2,6-
 difluorophenyl)(methylamino)methyl, (2,6-difluorophenyl)((2-
 15 hydroxyethyl)amino)methyl, amino(2-fluorophenyl)methyl, amino(2,6-
 difluorophenyl)methyl, (3-(oxazol-5-yl)pyridin-2-yl)methyl, 2-fluoro-6-(1-methyl-
 1H-pyrazol-4-yl)benzyl, (1-((1-methyl-1H-imidazol-4-yl)sulfonyl)pyrrolidin-2-
 yl)methyl, 2-((1-acetylpiperidin-4-yl)methyl)-6-fluorobenzyl, (2-(difluoromethoxy)-6-
 fluorophenyl)(hydroxy)methyl, 2-fluoro-6-(1-((1-methyl-1H-pyrazol-4-yl)methyl)-
 20 1H-pyrazol-4-yl)benzyl, (2-((dimethylamino)methyl)-6-
 fluorophenyl)(hydroxy)methyl, 2-fluoro-6-(1-(pyridin-2-ylmethyl)-1H-pyrazol-4-
 yl)benzyl, (2-fluoro-6-(pyrrolidin-1-ylmethyl)phenyl)(hydroxy)methyl, 2-fluoro-6-(1-
 (2-(methylsulfonyl)ethyl)-1H-pyrazol-4-yl)benzyl, 2-fluoro-6-((6-
 oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)phenyl)(hydroxy)methyl, 2-
 25 fluoro-6-(1-((trans)-3-(methylamino)cyclobutyl)-1H-pyrazol-4-yl)benzyl, 2-(1-(2-
 cyanoethyl)-1H-pyrazol-4-yl)-6-fluorobenzyl, 2-fluoro-6-(1-(2-(3-hydroxyazetidin-1-
 yl)-2-oxoethyl)-1H-pyrazol-4-yl)benzyl, (3-methylpyridin-2-yl)methoxy, (3-((1-
 (pyridin-4-ylmethyl)-1H-pyrazol-4-yl)amino)pyridin-2-yl)methyl, (3-((1-(tetrahydro-
 2H-pyran-4-yl)-1H-pyrazol-4-yl)amino)pyridin-2-yl)methyl, (3-(1-methyl-1H-
 30 pyrazol-4-yl)pyridin-2-yl)methyl, 2-((3-hydroxypyrrolidin-1-yl)methyl)benzyl, and
 (6-methoxypyridin-2-yl)methyl.

In some embodiments of Formulas (I) and (III), R⁴ is selected from pyridin-2-ylmethyl, 1*H*-pyrrolo[2,3-*b*]pyridin-1-yl, 7*H*-pyrrolo[2,3-*b*]pyridin-7-yl, 2-fluorophenoxy, hydroxy(pyridin-2-yl)methyl, 2-(1-methyl-1*H*-pyrazol-4-yl)benzyl, imidazo[1,2-*a*]pyridin-8-ylmethyl, pyrazolo[1,5-*a*]pyridin-7-ylmethyl, (2*H*-Indazol-2-yl)methyl, 1*H*-indazol-1-yl)methyl, (2,6-difluorophenyl)(hydroxy)methyl, (2,5-difluorophenyl)(hydroxy)methyl, (2,3-difluorophenyl)(hydroxy)methyl, (2-fluorophenyl)(hydroxy)methyl, (2-chlorophenyl)(hydroxy)methyl, hydroxy(phenyl)methyl, phenylsulfonyl, and azetidine-1-carbonyl.

In some embodiments of Formulas (I) and (III), Cy¹ is C₆₋₁₀ aryl which is optionally substituted by 1, 2, 3, or 4 independently selected R⁷ substituents.

In some embodiments of Formulas (I) and (III), Cy¹ is phenyl which is optionally substituted by 1 or 2 independently selected R⁷ substituents.

In some embodiments of Formulas (I) and (III), each R⁷ is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, and NO₂.

In some embodiments of Formulas (I) and (III), Cy¹ is phenyl which is optionally substituted by 1 or 2 independently selected R⁷ substituents; and each R⁷ is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, CN, and NO₂.

In some embodiments of Formulas (I) and (III), Cy¹ is phenyl which is substituted by 1 or 2 independently selected R⁷ substituents; and each R⁷ is independently selected from halo and CN.

In some embodiments of Formulas (I) and (III), Cy¹ is phenyl which is optionally substituted by 1 or 2 substituents independently selected from C₁₋₆ alkyl, halo, and CN.

In some embodiments of Formulas (I) and (III), Cy¹ is phenyl which is optionally substituted by 1 or 2 substituents independently selected from C₁₋₆ alkyl and CN.

In some embodiments of Formulas (I) and (III), Cy¹ is unsubstituted phenyl.

In some embodiments of Formulas (I) and (III), Cy¹ is cyanophenyl or cyanofluorophenyl.

In some embodiments of Formulas (I) and (III), Cy¹ is cyanophenyl.

In some embodiments of Formulas (I) and (III), Cy¹ is 3-cyanophenyl or 3-cyano-2-fluorophenyl.

In some embodiments of Formulas (I) and (III):

R¹ is H;

5 Cy¹ is C₆₋₁₀ aryl which is optionally substituted by 1, 2, 3, or 4 independently selected R⁷ substituents; and

R² is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R² are each optionally
10 substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (III):

R¹ is H;

Cy¹ is phenyl which is optionally substituted by 1 or 2 independently selected R⁷ substituents; and

15 R² is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R² are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments of Formulas (I) and (III):

20 R¹ is H or C₁₋₆ alkyl;

R² is 5-6 membered heteroaryl, optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

R⁴ is selected from C₆₋₁₀ aryl-C₁₋₃ alkyl-, (5-10 membered heteroaryl)-C₁₋₃ alkyl-, OR^{a4}, C(O)R^{b4}, and S(O)₂R^{b4}, wherein the C₆₋₁₀ aryl-C₁₋₃ alkyl- and (5-10 membered heteroaryl)-C₁₋₃ alkyl- of R⁴ are each optionally substituted with 1, 2, 3, or
25 4 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₃ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₃ alkyl-, (5-10 membered heteroaryl)-C₁₋₃ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₃ alkyl-, CN, OR^{a41}, SR^{a41}, and
30 NR^{c41}R^{d41}, wherein the C₁₋₆ alkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₃ alkyl-, C₃₋₁₀ cycloalkyl-

C₁₋₃ alkyl-, (5-10 membered heteroaryl)-C₁₋₃ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₃ alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, OR^{a42},
5 COOR^{a42}, and NR^{c42}R^{d42}.

Cy¹ is phenyl which is optionally substituted by 1, 2, or 3 independently selected R⁷ substituents;

each R^{a4} and R^{b4} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, phenyl, and azetidiny1 wherein the phenyl is optionally substituted with 1 or 2 groups
10 selected from halo, C₁₋₆ alkyl, and C₁₋₆ haloalkyl;

each R^{a41}, R^{c41}, and R^{d41} is independently selected from H and C₁₋₆ alkyl;

each R^{a42}, R^{c42}, and R^{d42} is independently selected from H and C₁₋₆ alkyl; and

each R⁷ is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, and
CN.

15 In some embodiments of Formulas (I) and (III):

R¹ is H or C₁₋₆ alkyl;

R² is 5-6 membered heteroaryl, optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

R⁴ is selected from C₆₋₁₀ aryl-C₁₋₃ alkyl- and (5-10 membered heteroaryl)-C₁₋₃ alkyl-, wherein the C₆₋₁₀ aryl-C₁₋₃ alkyl- and (5-10 membered heteroaryl)-C₁₋₃ alkyl- of
20 R⁴ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, and
OR^{a41};

25 Cy¹ is phenyl which is optionally substituted by 1 or 2 substituents independently selected from C₁₋₆ alkyl and CN; and

each R^{a41} is independently selected from H and C₁₋₃ alkyl.

In some embodiments of Formulas (I) and (III):

R¹ is selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₁₋₆ haloalkyl;

30 R² is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

R^4 is selected from $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$, and $S(O)_2R^{b4}$;

Cy^1 is C_{6-10} aryl which is optionally substituted by 1, 2, 3, or 4 independently selected R^7 substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and

each R^7 is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, and NO_2 .

In some embodiments of Formulas (I) and (III):

R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl;

R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents;

R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$;

Cy^1 is C_{6-10} aryl which is optionally substituted by 1, 2, 3, or 4 independently selected R^7 substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and

each R^7 is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, and NO_2 .

In some embodiments of Formulas (I) and (III):

R^1 is H;

R^2 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

R^4 is selected from $C(O)NR^{c4}R^{d4}$ and $NR^{c4}C(O)OR^{a4}$;

Cy^1 is phenyl which is optionally substituted by 1 or 2 independently selected R^7 substituents;

each R^{a4} , R^{c4} , and R^{d4} is independently selected from H and C_{1-6} alkyl; and

5 each R^7 is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, and NO_2 .

In some embodiments of Formulas (I) and (III):

R^1 is H;

10 R^2 is selected from 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl, wherein the 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl of R^2 are each optionally substituted with 1 or 2 independently selected R^{2A} substituents;

R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl;

15 each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, and $S(O)R^{b21}$;

20 Cy^1 is phenyl which is optionally substituted by 1 or 2 independently selected R^7 substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H and C_{1-6} alkyl; and

each R^7 is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, and NO_2 .

25 In some embodiments of Formulas (I) and (III):

R^1 is H;

R^2 is H;

R^4 is $NR^{c4}C(O)OR^{a4}$;

Cy^1 is cyanophenyl; and

30 R^{a4} and R^{c4} are each independently selected from H and C_{1-6} alkyl.

In some embodiments of Formulas (I) and (III):

R^1 is H;

R² is selected from 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl;

R⁴ is H; and

Cy¹ is cyanophenyl.

5 In some embodiments of Formulas (I) and (III):

R¹ is H;

R² is selected from 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl, wherein the 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl are each optionally substituted with 1 or 2 C₁₋₃ alkyl groups;

10 R⁴ is phenyl-C₁₋₃ alkyl- or pyridyl-C₁₋₃ alkyl-, wherein the phenyl-C₁₋₃ alkyl- and pyridyl-C₁₋₃ alkyl- are each optionally substituted with 1, 2, or 3 substituents independently selected from OH and halo; and

Cy¹ is cyanophenyl.

In some embodiments of Formulas (I) and (III):

15 R¹ is H;

R² is selected from 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl;

R⁴ is phenyl-C₁₋₃ alkyl- optionally substituted with 1, 2, or 3 substituents independently selected from OH and halo; and

20 Cy¹ is cyanophenyl.

In some embodiments of Formulas (I) and (III):


R¹ is H;

R² is pyrimidin-4-yl;

25 R⁴ is phenyl-C₁₋₃ alkyl- optionally substituted with 1, 2, or 3 substituents independently selected from OH and halo; and

Cy¹ is cyanophenyl.

In some embodiments of Formulas (I) and (III), the compound is a compound of Formula (IIIa):



or a pharmaceutically acceptable salt thereof, wherein variables R^2 , R^4 , and Cy^1 are defined according to the definitions provided herein for compounds of Formulas (I) and (III).

In some embodiments of Formula (IIIa),

Cy^1 is phenyl which is substituted by 1 or 2 independently selected R^7 substituents;

each R^7 is independently selected from halo and CN;

R^2 is selected from H, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$, wherein said C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R^{2A} are each optionally substituted by 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{241} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein said C_{1-6} alkyl of R^{a21} , R^{241} , R^{c21} , and R^{d21} is optionally substituted by 1, 2, or 3 independently selected R^{2A} substituents; and

each R^{2B} is independently selected from D, halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and $di(C_{1-3}$ alkyl)amino.

R^4 is selected from C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, OR^{a4} , $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $NR^{c4}R^{d4}$, $S(O)_2NR^{c4}R^{d4}$, and $S(O)_2R^{b4}$, wherein the C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a4} , R^{b4} , R^{c4} and R^{d4} are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, OR^{a41} , $C(O)R^{b41}$, $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $NR^{c41}R^{d41}$, $S(O)_2NR^{c41}R^{d41}$, and $S(O)_2R^{b41}$, wherein the C_{1-6} alkyl, 5 C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl, wherein the C_{1-6} alkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^{a41} , R^{b41} , R^{c41} and R^{d41} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, 5-6 membered heteroaryl, phenyl- C_{1-3} alkyl-, (C_{3-6} cycloalkyl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, (5-6 membered heteroaryl)- C_{1-3} alkyl-, CN, OR^{a42} , $C(O)R^{b42}$, $C(O)NR^{c42}R^{d42}$, $C(O)OR^{a42}$, $NR^{c42}R^{d42}$, $S(O)_2NR^{c42}R^{d42}$, and $S(O)_2R^{b42}$, wherein the C_{1-6} alkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, 5-6 membered heteroaryl, phenyl- C_{1-3} alkyl-, (C_{3-6} cycloalkyl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, (5-6 membered heteroaryl)- C_{1-3} alkyl- of R^{4B} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{a42} , R^{b42} , R^{c42} and R^{d42} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl, wherein the C_{1-6} alkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^{a42} , R^{b42} , R^{c42} and R^{d42} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{4C} is independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, CN, OR^{a43} , $C(O)R^{b43}$, $C(O)NR^{c43}R^{d43}$, $C(O)OR^{a43}$, $NR^{c43}R^{d43}$, $S(O)_2NR^{c43}R^{d43}$, and $S(O)_2R^{b43}$, wherein the C_{1-6} alkyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl- of R^{4C} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

each R^{a43} , R^{b43} , R^{c43} and R^{d43} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein the C_{1-6} alkyl of R^{a43} , R^{b43} , R^{c43} and R^{d43} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43} , attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, or 6-membered heterocycloalkyl group, wherein the 4-, 5-, or 6-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents; and

each R^{4D} is independently selected from C_{1-3} alkyl and OH.

In some embodiments of Formula (IIIa),

Cy^1 is 3-cyanophenyl or 3-cyano-2-fluorophenyl;

R^2 is selected from H, 5-10 membered heteroaryl and 4-7 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-7 membered

heterocycloalkyl of R^2 are each optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, and OH, wherein said C_{1-6} alkyl of R^{2A} is optionally substituted by 1, 2, or 3

5 independently selected R^{2B} substituents;

each R^{2B} is independently selected from D, halo, and OH;

R^4 is selected from phenyl- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, OR^{a4} , $C(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the phenyl- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl- and (5-10

10 membered heteroaryl)- C_{1-3} alkyl- of R^4 are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from phenyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl-, wherein the phenyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl- of

15 R^{a4} , R^{b4} , R^{c4} and R^{d4} are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, 5-6 membered heteroaryl, (4-10 membered heterocycloalkyl)- C_{1-3} alkyl-, OR^{a41} , $S(O)_2R^{b41}$, and $NR^{c41}R^{d41}$, wherein the C_{1-6} alkyl, 5-6 membered heteroaryl, and (4-10

20 membered heterocycloalkyl)- C_{1-3} alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, and 5-6 membered heteroaryl, wherein the C_{1-6} alkyl and 5-6 membered heteroaryl of R^{a41} , R^{b41} , R^{c41} and R^{d41} are each optionally substituted with 1, 2, or 3

25 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, (5-6 membered heteroaryl)- C_{1-3} alkyl-, CN, OR^{a42} , $C(O)R^{b42}$, $C(O)OR^{a42}$, and $NR^{c42}R^{d42}$, wherein the C_{1-6} alkyl, C_{3-6}

cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, and (5-6

30 membered heteroaryl)- C_{1-3} alkyl- of R^{4B} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{a42} , R^{b42} , R^{c42} and R^{d42} is independently selected from H, C_{1-6} alkyl, C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 5-7 membered heterocycloalkyl, wherein the C_{1-6} alkyl, C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 5-7 membered heterocycloalkyl of R^{a42} , R^{b42} , R^{c42} and R^{d42} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{4C} is independently selected from C_{1-6} alkyl, CN, $C(O)NR^{c43}R^{d43}$, $C(O)OR^{a43}$, $NR^{c43}R^{d43}$, and $S(O)_2R^{b43}$, wherein the C_{1-6} alkyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl- of R^{4C} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

each R^{a43} , R^{b43} , R^{c43} and R^{d43} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein the C_{1-6} alkyl of R^{a43} , R^{b43} , R^{c43} and R^{d43} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43} , attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, or 6-membered heterocycloalkyl group, wherein the 4-, 5-, or 6-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents; and

each R^{4D} is independently selected from C_{1-3} alkyl and OH.


In some embodiments of Formula (IIIa),

R^2 is selected from 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl, wherein the 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl are each optionally substituted with 1 or 2 C_{1-3} alkyl groups;

R^4 is phenyl- C_{1-3} alkyl- or pyridyl- C_{1-3} alkyl-, wherein the phenyl- C_{1-3} alkyl- and pyridyl- C_{1-3} alkyl- are each optionally substituted with 1, 2, or 3 substituents independently selected from OH and halo; and


Cy^1 is 3-cyanophenyl.

In some embodiments of Formulas (I) and (III), the compound is a compound of Formula (IIIb):



or a pharmaceutically acceptable salt thereof, wherein n is an integer from 0 to 4, and
 5 wherein variables R^2 , R^4 , and R^7 are defined according to the definitions provided herein for compounds of Formulas (I) and (III).

In some embodiments of Formulas (I) and (III), the compound is a compound of Formula (IIIc):



10 or a pharmaceutically acceptable salt thereof, wherein variables R^2 , R^4 , and R^7 are defined according to the definitions provided herein for compounds of Formulas (I) and (III).

In some embodiments, the compound is the (*S*)-enantiomer of one of the preceding compounds, or a pharmaceutically acceptable salt thereof. In some
 15 embodiments, the compound is the (*R*)-enantiomer of one of the preceding compounds, or a pharmaceutically acceptable salt thereof.

It is further appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention

which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

At various places in the present specification, divalent linking substituents are described. It is specifically intended that each divalent linking substituent include both the forward and backward forms of the linking substituent. For example, -
 5 NR(CR'R'')_n- includes both -NR(CR'R'')_n- and -(CR'R'')_nNR-. Where the structure clearly requires a linking group, the Markush variables listed for that group are understood to be linking groups.

The term “n-membered” where n is an integer typically describes the number
 10 of ring-forming atoms in a moiety where the number of ring-forming atoms is n. For example, piperidiny1 is an example of a 6-membered heterocycloalkyl ring, pyrazolyl is an example of a 5-membered heteroaryl ring, pyridyl is an example of a 6-membered heteroaryl ring, and 1,2,3,4-tetrahydro-naphthalene is an example of a 10-membered cycloalkyl group.

As used herein, the phrase “optionally substituted” means unsubstituted or
 15 substituted. The substituents are independently selected, and substitution may be at any chemically accessible position. As used herein, the term “substituted” means that a hydrogen atom is removed and replaced by a substituent. A single divalent substituent, *e.g.*, oxo, can replace two hydrogen atoms. It is to be understood that
 20 substitution at a given atom is limited by valency.

As used herein, the phrase “each ‘variable’ is independently selected from” means substantially the same as wherein “at each occurrence ‘variable’ is selected from.”

Throughout the definitions, the term “C_{n-m}” indicates a range which includes
 25 the endpoints, wherein n and m are integers and indicate the number of carbons. Examples include C₁₋₃, C₁₋₄, C₁₋₆, and the like.

As used herein, the term “C_{n-m} alkyl”, employed alone or in combination with other terms, refers to a saturated hydrocarbon group that may be straight-chain or branched, having n to m carbons. Examples of alkyl moieties include, but are not
 30 limited to, chemical groups such as methyl (Me), ethyl (Et), *n*-propyl (*n*-Pr), isopropyl (iPr), *n*-butyl, *tert*-butyl, isobutyl, *sec*-butyl; higher homologs such as 2-methyl-1-butyl, *n*-pentyl, 3-pentyl, *n*-hexyl, 1,2,2-trimethylpropyl, and the like. In some

embodiments, the alkyl group contains from 1 to 6 carbon atoms, from 1 to 4 carbon atoms, from 1 to 3 carbon atoms, or 1 to 2 carbon atoms.

As used herein, “C_{n-m} alkenyl” refers to an alkyl group having one or more double carbon-carbon bonds and having n to m carbons. Example alkenyl groups include, but are not limited to, ethenyl, *n*-propenyl, isopropenyl, *n*-butenyl, *sec*-butenyl, and the like. In some embodiments, the alkenyl moiety contains 2 to 6, 2 to 4, or 2 to 3 carbon atoms.

As used herein, “C_{n-m} alkynyl” refers to an alkyl group having one or more triple carbon-carbon bonds and having n to m carbons. Example alkynyl groups include, but are not limited to, ethynyl, propyn-1-yl, propyn-2-yl, and the like. In some embodiments, the alkynyl moiety contains 2 to 6, 2 to 4, or 2 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkoxy”, employed alone or in combination with other terms, refers to a group of formula -O-alkyl, wherein the alkyl group has n to m carbons. Example alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy (e.g., *n*-propoxy and isopropoxy), butoxy (e.g., *n*-butoxy and *tert*-butoxy), and the like. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “amino” refers to a group of formula -NH₂.

As used herein, the term “aryl,” employed alone or in combination with other terms, refers to an aromatic hydrocarbon group, which may be monocyclic or polycyclic (e.g., having 2, 3 or 4 fused rings). The term “C_{n-m} aryl” refers to an aryl group having from n to m ring carbon atoms. Aryl groups include, e.g., phenyl, naphthyl, anthracenyl, phenanthrenyl, indanyl, indenyl, and the like. In some embodiments, aryl groups have from 5 to 10 carbon atoms. In some embodiments, the aryl group is phenyl or naphthyl. In some embodiments, the aryl is phenyl.

As used herein, “halo” refers to F, Cl, Br, or I. In some embodiments, a halo is F, Cl, or Br. In some embodiments, a halo is F or Cl. In some embodiments, a halo is F. In some embodiments, a halo is Cl.

As used herein, “C_{n-m} haloalkoxy” refers to a group of formula -O-haloalkyl having n to m carbon atoms. Example haloalkoxy groups include OCF₃ and OCHF₂. In some embodiments, the haloalkoxy group is fluorinated only. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} haloalkyl”, employed alone or in combination with other terms, refers to an alkyl group having from one halogen atom to 2s+1 halogen atoms which may be the same or different, where “s” is the number of carbon atoms in the alkyl group, wherein the alkyl group has n to m carbon atoms. In some
 5 embodiments, the haloalkyl group is fluorinated only. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms. Example haloalkyl groups include CF₃, C₂F₅, CHF₂, CH₂F, CCl₃, CHCl₂, C₂Cl₅ and the like.

As used herein, the term “thio” refers to a group of formula-SH.

As used herein, the term “carbamyl” to a group of formula -C(O)NH₂.

10 As used herein, the term “carbonyl”, employed alone or in combination with other terms, refers to a -C(O)- group.

As used herein, the term “C_{n-m} alkylamino” refers to a group of formula -NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

15 As used herein, the term “C_{n-m} alkoxy carbonyl” refers to a group of formula -C(O)O-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkylcarbonyl” refers to a group of formula -C(O)-alkyl, wherein the alkyl group has n to m carbon atoms. In some
 20 embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkylcarbonylamino” refers to a group of formula -NHC(O)-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkylsulfonylamino” refers to a group of
 25 formula -NHS(O)₂-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “aminosulfonyl” refers to a group of formula -S(O)₂NH₂.

As used herein, the term “C_{n-m} alkylaminosulfonyl” refers to a group of
 30 formula -S(O)₂NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “di(C_{n-m} alkyl)aminosulfonyl” refers to a group of formula -S(O)₂N(alkyl)₂, wherein each alkyl group independently has n to m carbon atoms. In some embodiments, each alkyl group has, independently, 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

5 As used herein, the term “aminosulfonylamino” refers to a group of formula -NHS(O)₂NH₂.

As used herein, the term “C_{n-m} alkylaminosulfonylamino” refers to a group of formula -NHS(O)₂NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

10 As used herein, the term “di(C_{n-m} alkyl)aminosulfonylamino” refers to a group of formula -NHS(O)₂N(alkyl)₂, wherein each alkyl group independently has n to m carbon atoms. In some embodiments, each alkyl group has, independently, 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “aminocarbonylamino”, employed alone or in combination with other terms, refers to a group of formula -NHC(O)NH₂.

As used herein, the term “C_{n-m} alkylaminocarbonylamino” refers to a group of formula -NHC(O)NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “di(C_{n-m} alkyl)aminocarbonylamino” refers to a group of formula -NHC(O)N(alkyl)₂, wherein each alkyl group independently has n to m carbon atoms. In some embodiments, each alkyl group has, independently, 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkylcarbonyl” refers to a group of formula -C(O)-NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkylthio” refers to a group of formula -S-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkylsulfinyl” refers to a group of formula -S(O)-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkylsulfonyl” refers to a group of formula -S(O)₂-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “cyano-C_{1-n} alkyl” refers to a group of formula -(C_{1-n} alkylene)-CN, wherein the alkyl group has 1 to n carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms, e.g., -(C₁₋₃ alkylene)-CN.

As used herein, the term “HO-C_{1-n} alkyl” refers to a group of formula -(C_{1-n} alkylene)-OH, wherein the alkyl group has 1 to n carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms, e.g., -(C₁₋₃ alkylene)-OH.

As used herein, the term “C_{1-n} alkoxy-C_{1-n} alkyl” refers to a group of formula -(C_{1-n} alkylene)-O(C_{1-n} alkyl), wherein the alkyl group has 1 to n carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms, e.g., -(C₁₋₆ alkylene)-O(C₁₋₆ alkyl).

As used herein, the term “carboxy” refers to a group of formula -C(O)OH.

As used herein, the term “di(C_{n-m}-alkyl)amino” refers to a group of formula -N(alkyl)₂, wherein the two alkyl groups each has, independently, n to m carbon atoms. In some embodiments, each alkyl group independently has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “di(C_{n-m}-alkyl)carbonyl” refers to a group of formula -C(O)N(alkyl)₂, wherein the two alkyl groups each has, independently, n to m carbon atoms. In some embodiments, each alkyl group independently has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkylcarbonyloxy” is a group of formula -OC(O)-alkyl, wherein the alkyl group has n to m carbon atoms.

As used herein, “aminocarbonyloxy” is a group of formula -OC(O)-NH₂.

As used herein, “C_{n-m} alkylaminocarbonyloxy” is a group of formula -OC(O)-NH-alkyl, wherein the alkyl group has n to m carbon atoms.

As used herein, “di(C_{n-m}alkyl)aminocarbonyloxy” is a group of formula -OC(O)-N(alkyl)₂, wherein each alkyl group has, independently, n to m carbon atoms.

As used herein “C_{n-m} alkoxycarbonylamino” refers to a group of formula -NHC(O)O(C_{n-m} alkyl), wherein the alkyl group has n to m carbon atoms.

As used herein, “cycloalkyl” refers to non-aromatic cyclic hydrocarbons including cyclized alkyl and alkenyl groups. Cycloalkyl groups can include mono- or polycyclic (e.g., having 2 fused rings) groups, spirocycles, and bridged rings (e.g., a bridged bicycloalkyl group). Ring-forming carbon atoms of a cycloalkyl group can be optionally substituted by oxo or sulfido (e.g., C(O) or C(S)). Also included in the definition of cycloalkyl are moieties that have one or more aromatic rings fused (*i.e.*, having a bond in common with) to the cycloalkyl ring, for example, benzo or thienyl derivatives of cyclopentane, cyclohexane, and the like. A cycloalkyl group containing a fused aromatic ring can be attached through any ring-forming atom including a ring-forming atom of the fused aromatic ring. Cycloalkyl groups can have 3, 4, 5, 6, 7, 8, 9, or 10 ring-forming carbons (*i.e.*, C₃₋₁₀). In some embodiments, the cycloalkyl is a C₃₋₁₀ monocyclic or bicyclic cycloalkyl. In some embodiments, the cycloalkyl is a C₃₋₇ monocyclic cycloalkyl. In some embodiments, the cycloalkyl is a C₄₋₇ monocyclic cycloalkyl. In some embodiments, the cycloalkyl is a C₄₋₁₀ spirocycle or bridged cycloalkyl (e.g., a bridged bicycloalkyl group). Example cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcarnyl, cubane, adamantane, bicyclo[1.1.1]pentyl, bicyclo[2.1.1]hexyl, bicyclo[2.2.1]heptanyl, bicyclo[3.1.1]heptanyl, bicyclo[2.2.2]octanyl, spiro[3.3]heptanyl, and the like. In some embodiments, cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

As used herein, “heteroaryl” refers to a monocyclic or polycyclic (e.g., having 2 fused rings) aromatic heterocycle having at least one heteroatom ring member selected from N, O, S and B. In some embodiments, the heteroaryl ring has 1, 2, 3, or 4 heteroatom ring members independently selected from N, O, S and B. In some embodiments, any ring-forming N in a heteroaryl moiety can be an N-oxide. In some embodiments, the heteroaryl is a 5-10 membered monocyclic or bicyclic heteroaryl having 1, 2, 3, or 4 heteroatom ring members independently selected from N, O, S, and B. In some embodiments, the heteroaryl is a 5-10 membered monocyclic or bicyclic heteroaryl having 1, 2, 3, or 4 heteroatom ring members independently

selected from N, O, and S. In some embodiments, the heteroaryl is a 5-6 monocyclic heteroaryl having 1 or 2 heteroatom ring members independently selected from N, O, S, and B. In some embodiments, the heteroaryl is a 5-6 monocyclic heteroaryl having 1 or 2 heteroatom ring members independently selected from N, O, and S. In some
 5 embodiments, the heteroaryl group contains 3 to 10, 4 to 10, 5 to 10, 5 to 7, 3 to 7, or 5 to 6 ring-forming atoms. In some embodiments, the heteroaryl group has 1 to 4 ring-forming heteroatoms, 1 to 3 ring-forming heteroatoms, 1 to 2 ring-forming heteroatoms or 1 ring-forming heteroatom. When the heteroaryl group contains more than one heteroatom ring member, the heteroatoms may be the same or different.

10 Example heteroaryl groups include, but are not limited to, thienyl (or thiophenyl), furyl (or furanyl), pyrrolyl, imidazolyl, thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, isoxazolyl, 1,2,3-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-triazolyl, 1,2,4-thiadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-triazolyl, 1,3,4-thiadiazolyl, 1,3,4-oxadiazolyl and 1,2-dihydro-1,2-azaborine, pyridinyl, pyrimidinyl, pyrazinyl,
 15 pyridazinyl, azolyl, triazolyl, thiadiazolyl, quinolinyl, isoquinolinyl, indolyl, benzothiophenyl, benzofuranyl, benzisoxazolyl, imidazo[1, 2-b]thiazolyl, purinyl, triazinyl, thieno[3,2-b]pyridinyl, imidazo[1,2-a]pyridinyl, 1,5-naphthyridinyl, 1H-pyrazolo[4,3-b]pyridinyl, triazolo[4,3-a]pyridinyl, 1H-pyrrolo[3,2-b]pyridinyl, 1H-pyrrolo[2,3-b]pyridinyl, pyrazolo[1,5-a]pyridinyl, indazolyl, and the like.

20 As used herein, "heterocycloalkyl" refers to monocyclic or polycyclic heterocycles having at least one non-aromatic ring (saturated or partially unsaturated ring), wherein one or more of the ring-forming carbon atoms of the heterocycloalkyl is replaced by a heteroatom selected from N, O, S, and B, and wherein the ring-forming carbon atoms and heteroatoms of a heterocycloalkyl group can be optionally
 25 substituted by one or more oxo or sulfido (e.g., C(O), S(O), C(S), or S(O)₂, etc.). When a ring-forming carbon atom or heteroatom of a heterocycloalkyl group is optionally substituted by one or more oxo or sulfide, the O or S of said group is *in addition* to the number of ring-forming atoms specified herein (e.g., a 1-methyl-6-oxo-1,6-dihydropyridazin-3-yl is a 6-membered heterocycloalkyl group, wherein a
 30 ring-forming carbon atom is substituted with an oxo group, and wherein the 6-membered heterocycloalkyl group is further substituted with a methyl group). Heterocycloalkyl groups include monocyclic and polycyclic (e.g., having 2 fused

rings) systems. Included in heterocycloalkyl are monocyclic and polycyclic 3 to 10, 4 to 10, 5 to 10, 4 to 7, 5 to 7, or 5 to 6 membered heterocycloalkyl groups.


Heterocycloalkyl groups can also include spirocycles and bridged rings (*e.g.*, a 5 to 10 membered bridged biheterocycloalkyl ring having one or more of the ring-forming carbon atoms replaced by a heteroatom independently selected from N, O, S, and B). The heterocycloalkyl group can be attached through a ring-forming carbon atom or a ring-forming heteroatom. In some embodiments, the heterocycloalkyl group contains 0 to 3 double bonds. In some embodiments, the heterocycloalkyl group contains 0 to 2 double bonds.

Also included in the definition of heterocycloalkyl are moieties that have one or more aromatic rings fused (*i.e.*, having a bond in common with) to the non-aromatic heterocyclic ring, for example, benzo or thienyl derivatives of piperidine, morpholine, azepine, etc. A heterocycloalkyl group containing a fused aromatic ring can be attached through any ring-forming atom including a ring-forming atom of the fused aromatic ring.

In some embodiments, the heterocycloalkyl group contains 3 to 10 ring-forming atoms, 4 to 10 ring-forming atoms, 3 to 7 ring-forming atoms, or 5 to 6 ring-forming atoms. In some embodiments, the heterocycloalkyl group has 1 to 4 heteroatoms, 1 to 3 heteroatoms, 1 to 2 heteroatoms or 1 heteroatom. In some embodiments, the heterocycloalkyl is a monocyclic 4-6 membered heterocycloalkyl having 1 or 2 heteroatoms independently selected from N, O, S and B and having one or more oxidized ring members. In some embodiments, the heterocycloalkyl is a monocyclic or bicyclic 5-10 membered heterocycloalkyl having 1, 2, 3, or 4 heteroatoms independently selected from N, O, S, and B and having one or more oxidized ring members. In some embodiments, the heterocycloalkyl is a monocyclic or bicyclic 5 to 10 membered heterocycloalkyl having 1, 2, 3, or 4 heteroatoms independently selected from N, O, and S and having one or more oxidized ring members. In some embodiments, the heterocycloalkyl is a monocyclic 5 to 6 membered heterocycloalkyl having 1, 2, 3, or 4 heteroatoms independently selected from N, O, and S and having one or more oxidized ring members.

Example heterocycloalkyl groups include pyrrolidin-2-one (or 2-oxopyrrolidinyl), 1,3-isoxazolidin-2-one, pyranyl, tetrahydropyran, oxetanyl,

azetidiny, morpholino, thiomorpholino, piperaziny, tetrahydrofurany, tetrahydrothieny, piperidiny, pyrrolidiny, isoxazolidiny, isothiazolidiny, pyrazolidiny, oxazolidiny, thiazolidiny, imidazolidiny, azepany, 1,2,3,4-tetrahydroisoquinoline, benzazapene, azabicyclo[3.1.0]hexany, 5 diazabicyclo[3.1.0]hexany, oxobicyclo[2.1.1]hexany, azabicyclo[2.2.1]heptany, diazabicyclo[2.2.1]heptany, azabicyclo[3.1.1]heptany, diazabicyclo[3.1.1]heptany, azabicyclo[3.2.1]octany, diazabicyclo[3.2.1]octany, oxobicyclo[2.2.2]octany, azabicyclo[2.2.2]octany, azaadamantany, diazaadamantany, oxo-adamantany, azaspiro[3.3]heptany, diazaspiro[3.3]heptany, oxo-azaspiro[3.3]heptany, 10 azaspiro[3.4]octany, diazaspiro[3.4]octany, oxo-azaspiro[3.4]octany, azaspiro[2.5]octany, diazaspiro[2.5]octany, azaspiro[4.4]nonany, diazaspiro[4.4]nonany, oxo-azaspiro[4.4]nonany, azaspiro[4.5]decany, diazaspiro[4.5]decany, diazaspiro[4.4]nonany, oxo-diazaspiro[4.4]nonany, oxo-dihydropyridaziny, oxo-2,6-diazaspiro[3.4]octany, oxohexahydropyrrolo[1,2- 15 a]pyraziny, 3-oxopiperaziny, oxo-pyrrolidiny, oxo-pyridiny and the like. For example, heterocycloalkyl groups include the following groups (with and without N-methyl substitution):



As used herein, “C_{o-p} cycloalkyl-C_{n-m} alkyl-” refers to a group of formula 20 cycloalkyl-alkylene-, wherein the cycloalkyl has o to p carbon atoms and the alkylene linking group has n to m carbon atoms.

As used herein “C_{o-p} aryl-C_{n-m} alkyl-” refers to a group of formula aryl-alkylene-, wherein the aryl has o to p carbon atoms and the alkylene linking group has n to m carbon atoms.

25 As used herein, “heteroaryl-C_{n-m} alkyl-” refers to a group of formula heteroaryl-alkylene-, wherein alkylene linking group has n to m carbon atoms.

As used herein “heterocycloalkyl-C_{n-m} alkyl-” refers to a group of formula heterocycloalkyl-alkylene-, wherein alkylene linking group has n to m carbon atoms.

As used herein, an “alkyl linking group” is a bivalent straight chain or 30 branched alkyl linking group (“alkylene group”). For example, “C_{o-p} cycloalkyl-C_{n-m}

alkyl-”, “C_{o-p} aryl-C_{n-m} alkyl-”, “phenyl-C_{n-m} alkyl-”, “heteroaryl-C_{n-m} alkyl-”, and “heterocycloalkyl-C_{n-m} alkyl-” contain alkyl linking groups. Examples of “alkyl linking groups” or “alkylene groups” include methylene, ethan-1,1-diyl, ethan-1,2-diyl, propan-1,3-diyl, propan-1,2-diyl, propan-1,1-diyl and the like.

5 At certain places, the definitions or embodiments refer to specific rings (e.g., an azetidine ring, a pyridine ring, etc.). Unless otherwise indicated, these rings can be attached to any ring member provided that the valency of the atom is not exceeded. For example, an azetidine ring may be attached at any position of the ring, whereas a pyridin-3-yl ring is attached at the 3-position.

10 As used herein, the term “oxo” refers to an oxygen atom (i.e., =O) as a divalent substituent, forming a carbonyl group when attached to a carbon (e.g., C=O or C(O)), or attached to a nitrogen or sulfur heteroatom forming a nitroso, sulfinyl, or sulfonyl group.

As used herein, the term “independently selected from” means that each
15 occurrence of a variable or substituent, e.g., R⁷ or R^{2A}, are independently selected at each occurrence from the applicable list.

The compounds described herein can be asymmetric (e.g., having one or more stereocenters). All stereoisomers, such as enantiomers and diastereomers, are intended unless otherwise indicated. Compounds of the present disclosure that contain
20 asymmetrically substituted carbon atoms can be isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically inactive starting materials are known in the art, such as by resolution of racemic mixtures or by stereoselective synthesis. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable
25 isomers are contemplated in the present invention. *Cis* and *trans* geometric isomers of the compounds of the present disclosure are described and may be isolated as a mixture of isomers or as separated isomeric forms. In some embodiments, the compound has the (*R*)-configuration. In some embodiments, the compound has the (*S*)-configuration. The Formulas (e.g., Formula (I), (II), etc.) provided herein include
30 stereoisomers of the compounds.

Resolution of racemic mixtures of compounds can be carried out by any of numerous methods known in the art. An example method includes fractional

recrystallizaion using a chiral resolving acid which is an optically active, salt-forming organic acid. Suitable resolving agents for fractional recrystallization methods are, for example, optically active acids, such as the D and L forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid or
5 the various optically active camphorsulfonic acids such as β -camphorsulfonic acid. Other resolving agents suitable for fractional crystallization methods include stereoisomerically pure forms of α -methylbenzylamine (*e.g.*, *S* and *R* forms, or diastereomerically pure forms), 2-phenylglycinol, norephedrine, ephedrine, N-methylephedrine, cyclohexylethylamine, 1,2-diaminocyclohexane, and the like.

10 Resolution of racemic mixtures can also be carried out by elution on a column packed with an optically active resolving agent (*e.g.*, dinitrobenzoylphenylglycine). Suitable elution solvent composition can be determined by one skilled in the art.

Compounds provided herein also include tautomeric forms. Tautomeric forms result from the swapping of a single bond with an adjacent double bond together with
15 the concomitant migration of a proton. Tautomeric forms include prototropic tautomers which are isomeric protonation states having the same empirical formula and total charge. Example prototropic tautomers include ketone – enol pairs, amide – imidic acid pairs, lactam – lactim pairs, enamine – imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, for
20 example, 1H- and 3H-imidazole, 1H-, 2H- and 4H- 1,2,4-triazole, 1H- and 2H- isoindole, 2-hydroxypyridine and 2-pyridone, and 1H- and 2H-pyrazole. Tautomeric forms can be in equilibrium or sterically locked into one form by appropriate substitution.

All compounds, and pharmaceutically acceptable salts thereof, can be found
25 together with other substances such as water and solvents (*e.g.* hydrates and solvates) or can be isolated.

In some embodiments, preparation of compounds can involve the addition of acids or bases to affect, for example, catalysis of a desired reaction or formation of salt forms such as acid addition salts.

30 In some embodiments, the compounds provided herein, or salts thereof, are substantially isolated. By “substantially isolated” is meant that the compound is at least partially or substantially separated from the environment in which it was formed

or detected. Partial separation can include, for example, a composition enriched in the compounds provided herein. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99%
5 by weight of the compounds provided herein, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

The term “compound” as used herein is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted. Compounds herein identified by name or structure as one particular tautomeric form are intended
10 to include other tautomeric forms unless otherwise specified.

The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other
15 problem or complication, commensurate with a reasonable benefit/risk ratio.

The present application also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, “pharmaceutically acceptable salts” refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of
20 pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of the present disclosure include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically
25 acceptable salts of the present disclosure can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous
30 media like ether, ethyl acetate, alcohols (e.g., methanol, ethanol, iso-propanol, or butanol) or acetonitrile (ACN) are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton,


Pa., 1985, p. 1418 and *Journal of Pharmaceutical Science*, 66, 2 (1977), each of which is incorporated herein by reference in its entirety.

Synthesis

5 As will be appreciated by those skilled in the art, the compounds provided herein, including salts and stereoisomers thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes.

Compounds of formula **1-14** and **1-15** can be prepared *via* the synthetic route outlined in Scheme 1. Alkylation of commercially available starting material **1-1** with carbonyl adduct **1-2** (Hal is a halide, such as F, Cl, Br, or I), followed by a condensation reaction at elevated temperature, using an appropriate reagent, such as ammonium acetate, generates bicyclic compound **1-3**. Compound **1-3** can then react with reagents, such as phosphoryl chloride (POCl₃), to give intermediate **1-4**. A
15 nucleophilic aromatic substitution (S_NAr) reaction of intermediate **1-4** with amine adduct **1-5** (PG is a suitable protecting group, such as 2,4-dimethoxybenzyl), followed by reduction of the ester functionality with a suitable reductant (*e.g.*, DIBAL-H), affords alcohol **1-6**. Halogenation of **1-6** with an appropriate reagent, such as phosphorous tribromide (PBr₃), generates intermediate **1-7**. Compound **1-7** can then
20 be cross-coupled with an adduct of formula **1-8**, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is B(OR)₂, Sn(Alkyl)₃, or Zn-Hal], under standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*,
25 in the presence of a palladium catalyst) to afford the cross-coupling product, which undergoes protecting group removal to generate intermediate **1-9**. In some embodiments, Cy⁴ can be a R⁴ or a R⁴-R^{4a}. Halogenation of **1-9** with an appropriate reagent, such as *N*-bromosuccinimide (NBS), affords two isomers **1-10** and **1-11**. The final products **1-14** and **1-15** can then be prepared by reacting the two isomers **1-10**
30 and **1-11** with either adduct **1-12** or **1-13** using reaction conditions similar to that described for the preparation of **1-9** from **1-7**.


Scheme 1.



- Compounds of formula 2-7 can be prepared *via* the synthetic route outlined in Scheme 2. Alkylation of commercially available starting material 2-1 (Hal is a halide, such as F, Cl, Br, or I) with carbonyl adduct 1-2, followed by condensation using an appropriate reagent, such as ammonium acetate, at elevated temperature generates bicyclic compound 2-2. Compound 2-2 can then react with a suitable reagent, such as phosphoryl chloride (POCl_3), to give intermediate 2-3. A nucleophilic aromatic substitution ($\text{S}_{\text{N}}\text{Ar}$) reaction of intermediate 2-3 with amine adduct 1-5 (PG is a suitable protecting group, such as 2,4-dimethoxybenzyl), followed by removal of the protecting group, affords compound 2-4. Halogenation of 2-4 with a suitable reagent, such as *N*-bromosuccinimide (NBS), gives compound 2-5. Intermediate 2-5 can be cross-coupled with an adduct of formula 1-12, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is B(OR)_2 , Sn(Alkyl)_3 , or Zn-Hal], under standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium

catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst) to afford compound **2-6**. Intermediate **2-6** then undergoes a second cross-coupling reaction with compound **1-8**, using a similar procedure as described for the preparation of **2-6** from **2-5**, to generate the product **2-7**.

Scheme 2.




Compounds of formula **3-6** can be prepared *via* the synthetic route outlined in Scheme 3. A nucleophilic aromatic substitution (S_NAr) reaction of compound **1-4** (prepared using procedures described in Scheme 1) with amine adduct **1-5** (PG is a suitable protecting group, such as 2,4-dimethoxybenzyl), followed by removal of the protecting group, affords intermediate **3-1**. Halogenation of **3-1** with an appropriate reagent, such as *N*-bromosuccinimide (NBS), followed by reduction of the ester functionality with a suitable reductant (*e.g.*, DIBAL-H), generates alcohol **3-2**. Intermediate **3-2** can then be oxidized with an appropriate oxidant (*e.g.*, Dess-Martin periodinane) to afford aldehyde **3-3**. An addition reaction between **3-3** and **3-4** (M^1 is a metal group, such as MgBr or Li) then affords secondary alcohol **3-5**. The final product **3-6** can be prepared via a cross-coupling reaction between intermediate **3-5** and an adduct of formula **1-12**, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is B(OR)₂, Sn(Alkyl)₃, or Zn-Hal], under

standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst).


5

Scheme 3.




Compound of formula **4-5** can be prepared *via* the synthetic route outlined in Scheme 4. A halogenation reaction of compound **3-1** (prepared using procedures from Scheme 3) with an appropriate reagent, such as *N*-bromosuccinimide (NBS), affords compound **4-1** (Hal is a halide, such as F, Cl, Br, or I). Compound **4-1** can then be cross-coupled with an adduct of formula **1-12**, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is B(OR)₂, Sn(Alkyl)₃, or Zn-Hal], under standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst) to generate intermediate **4-2**. Hydrolysis of **4-2** with a suitable reagent, such as sodium hydroxide, gives carboxylic acid **4-3**. Compound **4-3** can then react with amine **4-4** under standard amide coupling conditions, such as using HATU as coupling reagent and DIPEA as base, to generate product **4-5**.

Scheme 4.




- Compounds of formula 5-7 can be prepared *via* the synthetic route outlined in
- 5 Scheme 5. A nucleophilic aromatic substitution (S_NAr) reaction of commercially available starting material 5-1 (Hal is a halide, such as F, Cl, Br, or I) with amine 1-5 (PG is a suitable protecting group, such as 2,4-dimethoxybenzyl) affords compound 5-2. Compound 5-2 can then be cross-coupled with an adduct of formula 5-3, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M
- 10 is $B(OR)_2$, $Sn(Alkyl)_3$, or $Zn-Hal$], under standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst) to generate intermediate 5-4. The protecting group in 5-4 can be removed to give
- 15 compound 5-5. Halogenation of 5-5 with an appropriate reagent, such as *N*-bromosuccinimide (NBS), affords intermediate 5-6. Compound 5-6 can then be cross-coupled with an adduct of formula 1-12 to give the product 5-7, using a procedure similar to that described for the preparation of 5-4 from 5-2.

Scheme 5.



Compounds of formula **6-9** can be prepared *via* the synthetic route outlined in Scheme 6. Commercially available starting material **6-1** (Hal is a halide, such as F, Cl, Br, or I) can react with an appropriate reagent, such as *t*-butyl *O*-mesitylene carbamate (*Journal of Heterocyclic Chemistry*, **1975**, 12, 107), to form pyrazinium salt **6-2**. Intermediate **6-2** can then undergo a condensation reaction with an adduct of formula **6-3** to form compound **6-4**. A nucleophilic aromatic substitution ($\text{S}_{\text{N}}\text{Ar}$) reaction of **6-4** with amine **1-5** (PG is a suitable protecting group, such as 2,4-dimethoxybenzyl) affords compound **6-5**. Compound **6-6** can then be prepared via a cross-coupling reaction between intermediate **6-5** and an adduct of formula **5-3**, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is $\text{B}(\text{OR})_2$, $\text{Sn}(\text{Alkyl})_3$, or Zn-Hal], under standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst). The protecting group in **6-6** can be removed to afford compound **6-7**, which undergoes a halogenation reaction using an appropriate reagent, such as *N*-bromosuccinimide (NBS), to form compound **6-8**. The final product **6-9** can be synthesized by coupling **6-8** with an adduct of formula **1-12**, using similar procedures as described for the preparation of compound **6-6** from **6-5**.


Scheme 6.



- 5 Compounds of formula 7-9 can be prepared *via* the synthetic route outlined in Scheme 7. Alkylation of commercially available starting material 7-1 (Hal is a halide, such as F, Cl, Br, or I) with carbonyl adduct 1-2, followed by a condensation reaction at elevated temperature, using an appropriate reagent, such as ammonium acetate, generates bicyclic compound 7-2. Compound 7-2 can then react with reagents, such as
- 10 phosphoryl chloride (POCl₃), to give intermediate 7-3. A nucleophilic aromatic substitution (S_NAr) reaction of intermediate 7-3 with amine adduct 7-4 (PG is a suitable protecting group, such as 4-methoxybenzyl) affords intermediate 7-5. Compound 7-5 can then be cross-coupled with an adduct of formula 7-6, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is
- 15 B(OR)₂, Sn(Alkyl)₃, or Zn-Hal], under standard Suzuki Cross-Coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst) to generate compound 7-7. The protecting groups in 7-7 are removed, and the resulting
- 20 intermediate undergoes a halogenation reaction with an appropriate reagent, such as *N*-Bromosuccinimide (NBS), to afford adduct 7-8. The final product 7-9 can then be

synthesized by coupling **7-8** with an adduct of formula **1-12**, using similar procedures as described for the preparation of compound **7-7** from **7-5**.

Scheme 7




5

Compounds of formula **8-7**, **8-10**, and **8-11** can be prepared *via* the synthetic route outlined in Scheme 8. Compound **7-5** (can be prepared as described in Scheme 7) can first be cross-coupled with reagent of formula **8-1**, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is B(OR)₂, Sn(Alkyl)₃, or Zn-Hal], under standard Suzuki Cross-Coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst) to generate compound **8-2**. A halogenation reaction of **8-2** can then be carried out using an appropriate reagent, such as 1-bromopyrrolidine-2,5-dione, to afford intermediate **8-3**. Another cross-coupling reaction between **8-3** and reagent **1-12** can then be performed using similar conditions as described for the transformation from **7-5** to **8-2** to deliver compound **8-4**. The vinyl group in **8-4** is cleaved under suitable conditions, such as using osmium (VIII) oxide and sodium periodate, and the resulting aldehyde **8-5** is reacted with **3-4** in an 1,2-addition reaction (M¹ is a metal group, such as MgCl or Li) to generate alcohol **8-6**. The protecting group (PG) in **8-6** can then be removed to

generate the desired product **8-7**.

On the other hand, aldehyde **8-5** can undergo a reduction reaction using appropriate reagents, such as NaBH_4 , to afford alcohol **8-8**. A halogenation reaction of **8-8** then affords intermediate **8-9** using reagents such as PBr_3 . A cross-coupling reaction between **8-9** and **1-8** (using conditions described for the synthesis of **8-2** from **7-5**), followed by the removal of protecting groups (PG), will generate product **8-10**. Alternatively, **8-9** can react with amine **4-4** in a nucleophilic substitution ($\text{S}_\text{N}2$) reaction, followed by removal of protecting groups (PG), to afford product **8-11**. The reaction sequence described in this scheme can be rearranged and adjusted accordingly to fit the need of each analogue synthesis.


Scheme 8



Compounds of formula **9-9** can be prepared using the synthetic route outlined in Scheme 9. Commercially available starting material **9-1** (Hal is a halide, such as F, Cl, Br, or I) can be subjected to nucleophilic aromatic substitution ($\text{S}_\text{N}\text{Ar}$) with amine

7-4 (PG is a suitable protecting group, such as 4-methoxybenzyl) to afford compound 9-2. Intermediate 9-2 can react with an appropriate reagent, such as *O*-(mesitylsulfonyl)hydroxylamine (*Journal of Heterocyclic Chemistry*, **1975**, 12, 107), to form pyrazinium salt 9-3. Intermediate 9-3 can then undergo a condensation reaction with an intermediate of formula 9-4 to form compound 9-5. Compound 9-6 can then be prepared using a cross-coupling reaction between intermediate 9-5 and an intermediate of formula 5-3, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is B(OR)₂, Sn(Alkyl)₃, or Zn-Hal], under standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst). Compound 9-6 can undergo halogenation using an appropriate reagent, such as *N*-bromosuccinimide (NBS), to form compound 9-7. Compound 9-8 can be synthesized by coupling 9-7 with an intermediate of formula 1-12, using similar procedures as described for the preparation of compound 9-6 from 9-5. The final product 9-9 can be formed after removal of the protecting group in intermediate 9-8. Certain synthetic steps described herein can be rearranged, and/or omitted, to prepare different analogues.

Scheme 9.



20

Methods of Use

The compounds of the present disclosure can modulate the activity of adenosine receptors, such as subtypes A2A and A2B receptors. Accordingly, the compounds, salts or stereoisomers described herein can be used in methods of inhibiting adenosine receptors (e.g., A2A and/or A2B receptors) by contacting the receptor with any one or more of the compounds, salts, or compositions described herein. In some embodiments, the compounds or salts can be used in methods of inhibiting activity of an adenosine receptor in an individual/patient in need of the inhibition by administering an effective amount of a compound or salt of described herein. In some embodiments, modulating is inhibiting. In some embodiments, the contacting is *in vivo*. In some embodiments, the contacting is *ex vivo* or *in vitro*.

The compounds or salts described herein can be selective. By “selective,” it is meant that the compound binds to or inhibits an adenosine receptor with greater affinity or potency, respectively, compared to at least one other receptor, kinase, etc. The compounds of the present disclosure can also be dual antagonists (i.e., inhibitors) of adenosine receptors, e.g., A2A and A2B adenosine receptors.

Another aspect of the present disclosure pertains to methods of treating an adenosine receptor associated disease or disorder in an individual (e.g., patient) by administering to the individual in need of such treatment a therapeutically effective amount or dose of one or more compounds of the present disclosure or a pharmaceutical composition thereof. An adenosine receptor associated disease or disorder can include any disease, disorder or condition that is directly or indirectly linked to expression or activity of the adenosine receptor, including overexpression and/or abnormal activity levels.

The compounds of the present disclosure are useful in the treatment of diseases related to the activity of adenosine receptors including, for example, cancer, inflammatory diseases, cardiovascular diseases, neurodegenerative diseases, immunomodulatory disorders, central nerve system diseases, and diabetes.

Based on the compelling roles of adenosine, e.g., A2A, A2B, receptors in multiple immunosuppressive mechanisms, developing inhibitors can boost the immune system to suppress tumor progression. Adenosine receptor inhibitors can be used to treat, alone or in combination with other therapies, bladder cancer, lung cancer

(e.g., non-small cell lung cancer (NSCLC), lung metastasis), melanoma (e.g., metastatic melanoma), breast cancer, cervical cancer, ovarian cancer, colorectal cancer, pancreatic cancer, esophageal cancer, prostate cancer, kidney cancer, skin cancer, thyroid cancer, liver cancer, uterine cancer, head and neck cancer, and renal cell carcinoma (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842-857). See also, <https://globenewswire.com/news-release/2017/04/04/954192/0/en/Corvus-Pharmaceuticals-Announces-Interim-Results-from-Ongoing-Phase-1-1b-Study-Demonstrating-Safety-and-Clinical-Activity-of-Lead-Checkpoint-Inhibitor-CPI-444-in-Patients-with-Adva.html>; Cekic C. et al., *J Immunol*, 2012, 188:198-205; Iannone, R. et al., *Am. J. Cancer Res.* 2014, 4:172-181 (study shows that both A2A and CD73 blockade enhance the antitumor activity of anti-CTLA-4 mAb therapy in a B16F10 murine melanoma model); Iannone, R. et al., *Neoplasia*, 2013, 15:1400-1410 and Beavis PA., et al., *Proc Natl Acad Sci. USA*, 2013, 110:14711-14716 (study shows that A2A and CD73 blockade decreased metastasis in 4T1 breast tumor model with high CD73 expression). In some embodiments, the prostate cancer is metastatic castrate-resistant prostate carcinoma (mCRPC). In some embodiments, the colorectal cancer is colorectal carcinoma (CRC).

In some embodiments, the disease or disorder is lung cancer (e.g., non-small cell lung cancer), melanoma, pancreatic cancer, breast cancer, head and neck squamous cell carcinoma, prostate cancer, liver cancer, color cancer, endometrial cancer, bladder cancer, skin cancer, cancer of the uterus, renal cancer, gastric cancer, or sarcoma. In some embodiments, the sarcoma is Askin's tumor, sarcoma botryoides, chondrosarcoma, Ewing's sarcoma, malignant hemangioendothelioma, malignant schwannoma, osteosarcoma, alveolar soft part sarcoma, angiosarcoma, cystosarcoma phyllodes, dermatofibrosarcoma protuberans, desmoid tumor, desmoplastic small round cell tumor, epithelioid sarcoma, extraskeletal chondrosarcoma, extraskeletal osteosarcoma, fibrosarcoma, gastrointestinal stromal tumor (GIST), hemangiopericytoma, hemangiosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, lymphosarcoma, malignant peripheral nerve sheath tumor (MPNST), neurofibrosarcoma, rhabdomyosarcoma, synovial sarcoma, or undifferentiated pleomorphic sarcoma.

In some embodiments, the disease or disorder is mesothelioma or adenocarcinoma. In some embodiments, the disease or disorder is mesothelioma. In some embodiments, the disease or disorder is adenocarcinoma.

MDSC (myeloid-derived suppressor cells) are a heterogeneous group of
5 immune cells from the myeloid lineage (a family of cells that originate from bone marrow stem cells). MDSCs strongly expand in pathological situations such as chronic infections and cancer, as a result of an altered haematopoiesis. MDSCs are discriminated from other myeloid cell types in which they possess strong immunosuppressive activities rather than immunostimulatory properties. Similar to
10 other myeloid cells, MDSCs interact with other immune cell types including T cells, dendritic cells, macrophages and natural killer cells to regulate their functions. In some embodiments, the compounds, etc. described herein can be used in methods related to cancer tissue (e.g., tumors) with high infiltration of MDSCs, including Solid tumors with high basal level of macrophage and/or MDSC infiltration.

15 In some embodiments, the compounds of the disclosure can be used in treating pulmonary inflammation, including bleomycin-induced pulmonary fibrosis and injury related to adenosine deaminase deficiency (Baraldi, et al., *Chem. Rev.*, 2008, 108, 238–263).

In some embodiments, the compounds of the disclosure can be used as a
20 treatment for inflammatory disease such as allergic reactions (e.g., A2B adenosine receptor dependent allergic reactions) and other adenosine receptor dependent immune reactions. Further inflammatory diseases that can be treated by compounds of the disclosure include respiratory disorders, sepsis, reperfusion injury, and thrombosis.

25 In some embodiments, the compounds of the disclosure can be used as a treatment for cardiovascular disease such as coronary artery disease (myocardial infarction, angina pectoris, heart failure), cerebrovascular disease (stroke, transient ischemic attack), peripheral artery disease, and aortic atherosclerosis and aneurysm. Atherosclerosis is an underlying etiologic factor in many types of cardiovascular
30 disease. Atherosclerosis begins in adolescence with fatty streaks, which progress to plaques in adulthood and finally results in thrombotic events that cause occlusion of vessels leading to clinically significant morbidity and mortality. Antagonists to the

A2B adenosine receptor and A2A adenosine receptor may be beneficial in preventing atherosclerotic plaque formation (Eisenstein, A. et al., *J. Cell Physiol.*, 2015, 230(12), 2891-2897).

5 In some embodiments, the compounds of the disclosure can be used as a treatment for disorders in motor activity; deficiency caused by degeneration of the striatonigral dopamine system; and Parkinson's disease; some of the motivational symptoms of depression (Collins, L. E. et al. *Pharmacol. Biochem. Behav.*, 2012, 100, 498–505.).

10 In some embodiments, the compounds of the disclosure can be used as a treatment for diabetes and related disorders, such as insulin resistance. Diabetes affects the production of adenosine and the expression of A2B adenosine receptors (A2BRs) that stimulate IL-6 and CRP production, insulin resistance, and the association between A₂BR gene single-nucleotide polymorphisms (ADORA2B SNPs) and inflammatory markers. The increased A2BR signaling in diabetes may increase
15 insulin resistance in part by elevating pro-inflammatory mediators. Selective A2BR blockers may be useful to treat insulin resistance (Figler, R. A. et al. *Diabetes*, 2011, 60 (2), 669-679).

It is believed that compounds provided herein, e.g., compounds of Formula (I), or any of the embodiments thereof, may possess satisfactory pharmacological profile
20 and promising biopharmaceutical properties, such as toxicological profile, metabolism and pharmacokinetic properties, solubility, and permeability. It will be understood that determination of appropriate biopharmaceutical properties is within the knowledge of a person skilled in the art, e.g., determination of cytotoxicity in cells or inhibition of certain targets or channels to determine potential toxicity.

25 The terms “individual” or “patient”, used interchangeably, refer to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans.

The phrase “therapeutically effective amount” refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in
30 a tissue, system, animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

As used herein, the term “treating” or “treatment” refers to one or more of (1) inhibiting the disease; *e.g.*, inhibiting a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, arresting further development of the pathology and/or symptomatology); and (2) ameliorating the disease; *e.g.*, ameliorating a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, reversing the pathology and/or symptomatology) such as decreasing the severity of disease.

In some embodiments, the compounds of the invention are useful in preventing or reducing the risk of developing any of the diseases referred to herein; *e.g.*, preventing or reducing the risk of developing a disease, condition or disorder in an individual who may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease.

Combination Therapies

I. Immune-checkpoint therapies

In some embodiments, A2A and A2B dual inhibitors provided herein can be used in combination with one or more immune checkpoint inhibitors for the treatment of cancer as described herein. In one embodiment, the combination with one or more immune checkpoint inhibitors as described herein can be used for the treatment of melanoma. Compounds of the present disclosure can be used in combination with one or more immune checkpoint inhibitors. Exemplary immune checkpoint inhibitors include inhibitors against immune checkpoint molecules such as CD20, CD28, CD40, CD122, CD96, CD73, CD47, GITR, CSF1R, JAK, PI3K delta, PI3K gamma, TAM, arginase, HPK1, CD137 (also known as 4-1BB), ICOS, B7-H3, B7-H4, BTLA, CTLA-4, LAG3, TIM3, VISTA, TIGIT, PD-1, PD-L1 and PD-L2. In some embodiments, the immune checkpoint molecule is a stimulatory checkpoint molecule selected from CD27, CD28, CD40, ICOS, OX40, GITR and CD137. In some embodiments, the immune checkpoint molecule is an inhibitory checkpoint molecule selected from A2AR, B7-H3, B7-H4, BTLA, CTLA-4, IDO, KIR, LAG3, PD-1, TIM3, TIGIT, and VISTA. In some embodiments, the compounds of the disclosure provided herein can be used in combination with one or more agents selected from

KIR inhibitors, TIGIT inhibitors, LAIR1 inhibitors, CD160 inhibitors, 2B4 inhibitors and TGFR beta inhibitors.

In some embodiments, the A2A and A2B dual inhibitors provided herein can be used in combination with one or more agonists of immune checkpoint molecules, e.g., OX40, CD27, OX40, GITR, and CD137 (also known as 4-1BB).

In some embodiments, the inhibitor of an immune checkpoint molecule is anti-PD1 antibody, anti-PD-L1 antibody, or anti-CTLA-4 antibody.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-1, e.g., an anti-PD-1 monoclonal antibody. In some embodiments, the anti-PD-1 monoclonal antibody is nivolumab, pembrolizumab (also known as MK-3475), durvalumab (Imfinzi®), pidilizumab, SHR-1210, PDR001, MGA012, PDR001, AB122, or AMP-224. In some embodiments, the anti-PD-1 monoclonal antibody is nivolumab or pembrolizumab. In some embodiments, the anti-PD1 antibody is pembrolizumab. In some embodiments, the anti-PD-1 monoclonal antibody is MGA012. In some embodiments, the anti-PD1 antibody is SHR-1210. Other anti-cancer agent(s) include antibody therapeutics such as 4-1BB (e.g. urelumab or utomilumab).

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-L1, e.g., an anti-PD-L1 monoclonal antibody. In some embodiments, the anti-PD-L1 monoclonal antibody is BMS-935559, MEDI4736, MPDL3280A (also known as RG7446), or MSB0010718C. In some embodiments, the anti-PD-L1 monoclonal antibody is MPDL3280A or MEDI4736.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-1 and PD-L1, e.g., an anti-PD-1/PD-L1 monoclonal antibody. In some embodiments, the anti-PD-1/PD-L1 is MCLA-136.

In some embodiments, the inhibitor is INCB086550.

In some embodiments, the inhibitor is MCLA-145.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of CTLA-4, e.g., an anti-CTLA-4 antibody. In some embodiments, the anti-CTLA-4 antibody is ipilimumab, tremelimumab, AGEN1884, or CP-675,206.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of LAG3, e.g., an anti-LAG3 antibody. In some embodiments, the anti-LAG3 antibody is BMS-986016, LAG525, or INCAGN2385.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of TIM3, e.g., an anti-TIM3 antibody. In some embodiments, the anti-TIM3 antibody is INCAGN2390, MBG453, or TSR-022.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of GITR, e.g., an anti-GITR antibody. In some embodiments, the anti-GITR antibody is TRX518, MK-4166, INCAGN1876, MK-1248, AMG228, BMS-986156, GWN323, or MEDI1873.

In some embodiments, the inhibitor of an immune checkpoint molecule is an agonist of OX40, e.g., OX40 agonist antibody or OX40L fusion protein. In some embodiments, the anti-OX40 antibody is MEDI0562, MOXR-0916, PF-04518600, GSK3174998, or BMS-986178. In some embodiments, the OX40L fusion protein is MEDI6383.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of CD20, e.g., an anti-CD20 antibody. In some embodiments, the anti-CD20 antibody is obinutuzumab or rituximab.

The compounds of the present disclosure can be used in combination with bispecific antibodies. In some embodiments, one of the domains of the bispecific antibody targets PD-1, PD-L1, CTLA-4, GITR, OX40, TIM3, LAG3, CD137, ICOS, CD3, tumor specific antigens (e.g., CD70) or TGF β receptor.

In some embodiments, the compounds of the disclosure can be used in combination with one or more metabolic enzyme inhibitors. In some embodiments, the metabolic enzyme inhibitor is an inhibitor of IDO1, TDO, or arginase. Examples of IDO1 inhibitors include epacadostat, NLG919, BMS-986205, PF-06840003, IOM2983, RG-70099 and LY338196.

As provided throughout, the additional compounds, inhibitors, agents, etc. can be combined with the present compound in a single or continuous dosage form, or they can be administered simultaneously or sequentially as separate dosage forms.

II. Cancer therapies

Cancer cell growth and survival can be impacted by multiple signaling pathways. Thus, it is useful to combine different enzyme/protein/receptor inhibitors, exhibiting different preferences in the targets which they modulate the activities of, to
5 treat such conditions. Targeting more than one signaling pathway (or more than one biological molecule involved in a given signaling pathway) may reduce the likelihood of drug-resistance arising in a cell population, and/or reduce the toxicity of treatment.

The compounds of the present disclosure can be used in combination with one or more other enzyme/protein/receptor inhibitors or one or more therapies for the
10 treatment of diseases, such as cancer. Examples of diseases and indications treatable with combination therapies include those as described herein.

The compounds of the present disclosure can be used in combination with one or more additional pharmaceutical agents such as, for example, chemotherapeutics, immune-oncology agents, metabolic enzyme inhibitors, chemokine receptor
15 inhibitors, and phosphatase inhibitors, as well as targeted therapies such as Bcr-Abl, Flt-3, EGFR, HER2, JAK, c-MET, VEGFR, PDGFR, c-Kit, IGF-1R, RAF and FAK kinase inhibitors. The one or more additional pharmaceutical agents can be administered to a patient simultaneously or sequentially.

For example, the compounds as disclosed herein can be combined with one or
20 more inhibitors of the following kinases for the treatment of cancer and other diseases or disorders described herein: Akt1, Akt2, Akt3, TGF- β R, PKA, PKG, PKC, CaM-kinase, phosphorylase kinase, MEKK, ERK, MAPK, mTOR, EGFR, HER2, HER3, HER4, INS-R, IGF-1R, IR-R, PDGF α R, PDGF β R, CSFIR, KIT, FLK-II, KDR/FLK-1, FLK-4, flt-1, FGFR1, FGFR2, FGFR3, FGFR4, c-Met, Ron, Sea, TRKA, TRKB,
25 TRKC, FLT3, VEGFR/Flt2, Flt4, EphA1, EphA2, EphA3, EphB2, EphB4, Tie2, Src, Fyn, Lck, Fgr, Btk, Fak, SYK, FRK, JAK, ABL, ALK and B-Raf. Non-limiting examples of inhibitors that can be combined with the compounds of the present disclosure for treatment of cancer and other diseases and disorders described herein include an FGFR inhibitor (FGFR1, FGFR2, FGFR3 or FGFR4, e.g., INCB54828,
30 INCB62079 and INCB63904), a JAK inhibitor (JAK1 and/or JAK2, e.g., ruxolitinib, baricitinib or INCB39110), an IDO inhibitor (e.g., epacadostat, NLG919, or BMS-986205), an LSD1 inhibitor (e.g., INCB59872 and INCB60003), a TDO inhibitor, a

PI3K-delta inhibitor (e.g., INCB50797 and INCB50465), a Pim inhibitor, a CSF1R inhibitor, a TAM receptor tyrosine kinases (Tyro-3, Axl, and Mer), a histone deacetylase inhibitor (HDAC) such as an HDAC8 inhibitor, an angiogenesis inhibitor, an interleukin receptor inhibitor, bromo and extra terminal family members inhibitors
 5 (for example, bromodomain inhibitors or BET inhibitors such as INCB54329 and INCB57643) and an adenosine receptor antagonist or combinations thereof.

Example antibodies for use in combination therapy include but are not limited to Trastuzumab (e.g. anti-HER2), Ranibizumab (e.g. anti-VEGF-A), Bevacizumab (trade name Avastin, e.g. anti-VEGF, Panitumumab (e.g. anti-EGFR), Cetuximab
 10 (e.g. anti-EGFR), Rituxan (anti-CD20) and antibodies directed to c-MET.

One or more of the following agents may be used in combination with the compounds of the present disclosure and are presented as a non-limiting list: a cytostatic agent, cisplatin, doxorubicin, taxotere, taxol, etoposide, irinotecan, camptostar, topotecan, paclitaxel, docetaxel, epothilones, tamoxifen, 5-fluorouracil,
 15 methotrexate, temozolomide, cyclophosphamide, SCH 66336, R115777, L778,123, BMS 214662, IRESSATM(gefitinib), TARCEVATM (erlotinib), antibodies to EGFR, intron, ara-C, adriamycin, cytoxan, gemcitabine, uracil mustard, chlormethine, ifosfamide, melphalan, chlorambucil, pipobroman, triethylenemelamine, triethylenethiophosphoramine, busulfan, carmustine, lomustine, streptozocin,
 20 dacarbazine, floxuridine, cytarabine, 6-mercaptopurine, 6-thioguanine, fludarabine phosphate, oxaliplatin, leucovirin, ELOXATINTM (oxaliplatin), pentostatin, vinblastine, vincristine, vindesine, bleomycin, dactinomycin, daunorubicin, doxorubicin, epirubicin, idarubicin, mithramycin, deoxycoformycin, mitomycin-C, L-asparaginase, teniposide 17.alpha.-ethinylestradiol, diethylstilbestrol, testosterone,
 25 Prednisone, Fluoxymesterone, Dromostanolone propionate, testolactone, megestrolacetate, methylprednisolone, methyltestosterone, prednisolone, triamcinolone, chlorotrianisene, hydroxyprogesterone, aminoglutethimide, estramustine, medroxyprogesteroneacetate, leuprolide, flutamide, toremifene, goserelin, carboplatin, hydroxyurea, amsacrine, procarbazine, mitotane, mitoxantrone,
 30 levamisole, navelbene, anastrozole, letrozole, capecitabine, reloxafine, droloxafine, hexamethylmelamine, avastin, HERCEPTINTM (trastuzumab), BEXXARTM (tositumomab), VELCADETM (bortezomib), ZEVALINTM (ibritumomab tiuxetan),

TRISENOX™ (arsenic trioxide), XELODA™ (capecitabine), vinorelbine, porfimer, ERBITUX™ (cetuximab), thiotepa, altretamine, melphalan, trastuzumab, lerozole, fulvestrant, exemestane, ifosfomide, rituximab, C225 (cetuximab), Campath (alemtuzumab), clofarabine, cladribine, aphidicolon, rituxan, sunitinib, dasatinib, 5 tezacitabine, Sml1, fludarabine, pentostatin, triapine, didox, trimidox, amidox, 3-AP, and MDL-101,731.

The compounds of the present disclosure can further be used in combination with other methods of treating cancers, for example by chemotherapy, irradiation therapy, tumortargeted therapy, adjuvant therapy, immunotherapy or surgery.

10 Examples of immunotherapy include cytokine treatment (e.g., interferons, GM-CSF, G-CSF, IL-2), CRS-207 immunotherapy, cancer vaccine, monoclonal antibody, adoptive T cell transfer, Toll receptor agonists, STING agonists, oncolytic virotherapy and immunomodulating small molecules, including thalidomide or JAK1/2 inhibitor and the like. The compounds can be administered in combination with one or more

15 anti-cancer drugs, such as a chemotherapeutics. Example chemotherapeutics include any of: abarelix, aldesleukin, alemtuzumab, alitretinoin, allopurinol, altretamine, anastrozole, arsenic trioxide, asparaginase, azacitidine, bevacizumab, bexarotene, baricitinib, bleomycin, bortezombi, bortezomib, busulfan intravenous, busulfan oral, calusterone, capecitabine, carboplatin, carmustine, cetuximab, chlorambucil, cisplatin,

20 cladribine, clofarabine, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, dalteparin sodium, daunorubicin, decitabine, denileukin, denileukin diftitox, dexrazoxane, docetaxel, doxorubicin, dromostanolone propionate, eculizumab, epirubicin, erlotinib, estramustine, etoposide phosphate, etoposide, exemestane, fentanyl citrate, filgrastim, floxuridine, fludarabine, fluorouracil, fulvestrant, gefitinib,

25 gemcitabine, gemtuzumab ozogamicin, goserelin acetate, histrelin acetate, ibritumomab tiuxetan, idarubicin, ifosfamide, imatinib mesylate, interferon alfa 2a, irinotecan, lapatinib ditosylate, lenalidomide, letrozole, leucovorin, leuprolide acetate, levamisole, lomustine, meclorothamine, megestrol acetate, melphalan, mercaptopurine, methotrexate, methoxsalen, mitomycin C, mitotane, mitoxantrone,

30 nandrolone phenpropionate, nelarabine, nofetumomab, olaparib, oxaliplatin, paclitaxel, pamidronate, panitumumab, pegaspargase, pegfilgrastim, pemetrexed disodium, pentostatin, pipobroman, plicamycin, procarbazine, quinacrine, rasburicase,

rituximab, ruxolitinib, rucaparib, streptozocin, tamoxifen, temozolomide, teniposide, testolactone, thalidomide, thioguanine, thiotepa, topotecan, toremifene, tositumomab, trastuzumab, tretinoin, uracil mustard, valrubicin, vinblastine, vincristine, vinorelbine, vorinostat, niraparib, veliparib, talazoparib, and zoledronate.

5 Additional examples of chemotherapeutics include proteasome inhibitors (e.g., bortezomib), thalidomide, revlimid, and DNA-damaging agents such as melphalan, doxorubicin, cyclophosphamide, vincristine, etoposide, carmustine, and the like.

Example Bcr-Abl inhibitors include imatinib mesylate (GLEEVECTM), nilotinib, dasatinib, bosutinib, and ponatinib, and pharmaceutically acceptable salts.

10 Other example suitable Bcr-Abl inhibitors include the compounds, and pharmaceutically acceptable salts thereof, of the genera and species disclosed in U.S. Pat. No. 5,521,184, WO 04/005281, and U.S. Ser. No. 60/578,491.

Example suitable Flt-3 inhibitors include midostaurin, lestaurtinib, linifanib, sunitinib, sunitinib, maleate, sorafenib, quizartinib, crenolanib, pacritinib, tandutinib, 15 PLX3397 and ASP2215, and their pharmaceutically acceptable salts. Other example suitable Flt-3 inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 03/037347, WO 03/099771, and WO 04/046120.

Example suitable RAF inhibitors include dabrafenib, sorafenib, and vemurafenib, and their pharmaceutically acceptable salts. Other example suitable 20 RAF inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 00/09495 and WO 05/028444.

Example suitable FAK inhibitors include VS-4718, VS-5095, VS-6062, VS-6063, BI853520, and GSK2256098, and their pharmaceutically acceptable salts. Other example suitable FAK inhibitors include compounds, and their 25 pharmaceutically acceptable salts, as disclosed in WO 04/080980, WO 04/056786, WO 03/024967, WO 01/064655, WO 00/053595, and WO 01/014402.

In some embodiments, the compounds of the disclosure can be used in combination with one or more other kinase inhibitors including imatinib, particularly for treating patients resistant to imatinib or other kinase inhibitors.

30 In some embodiments, the compounds of the disclosure can be used in combination with a chemotherapeutic in the treatment of cancer, and may improve the treatment response as compared to the response to the chemotherapeutic agent alone,

without exacerbation of its toxic effects. In some embodiments, the compounds of the disclosure can be used in combination with a chemotherapeutic provided herein. For example, additional pharmaceutical agents used in the treatment of multiple myeloma, can include, without limitation, melphalan, melphalan plus prednisone [MP],
5 doxorubicin, dexamethasone, and Velcade (bortezomib). Further additional agents used in the treatment of multiple myeloma include Bcr-Abl, Flt-3, RAF and FAK kinase inhibitors. In some embodiments, the agent is an alkylating agent, a proteasome inhibitor, a corticosteroid, or an immunomodulatory agent. Examples of an alkylating agent include cyclophosphamide (CY), melphalan (MEL), and
10 bendamustine. In some embodiments, the proteasome inhibitor is carfilzomib. In some embodiments, the corticosteroid is dexamethasone (DEX). In some embodiments, the immunomodulatory agent is lenalidomide (LEN) or pomalidomide (POM). Additive or synergistic effects are desirable outcomes of combining a PI3K inhibitor of the present disclosure with an additional agent.

15 In some embodiments, the compounds of the disclosure can be used in combination with an inhibitor of JAK or PI3K δ .

The agents can be combined with the present compound in a single or continuous dosage form, or the agents can be administered simultaneously or sequentially as separate dosage forms.

20 The compounds of the present disclosure can be used in combination with one or more other inhibitors or one or more therapies for the treatment of infections. Examples of infections include viral infections, bacterial infections, fungus infections or parasite infections.

In some embodiments, a corticosteroid such as dexamethasone is administered
25 to a patient in combination with the compounds of the disclosure where the dexamethasone is administered intermittently as opposed to continuously.

The compounds of Formula (I) or any of the formulas as described herein, a compound as recited in any of the claims and described herein, or salts thereof can be combined with another immunogenic agent, such as cancerous cells, purified tumor
30 antigens (including recombinant proteins, peptides, and carbohydrate molecules), cells, and cells transfected with genes encoding immune stimulating cytokines. Non-limiting examples of tumor vaccines that can be used include peptides of melanoma

antigens, such as peptides of gp100, MAGE antigens, Trp-2, MARTI and/or tyrosinase, or tumor cells transfected to express the cytokine GM-CSF.

The compounds of Formula (I) or any of the formulas as described herein, a compound as recited in any of the claims and described herein, or salts thereof can be used in combination with a vaccination protocol for the treatment of cancer. In some embodiments, the tumor cells are transduced to express GM-CSF. In some embodiments, tumor vaccines include the proteins from viruses implicated in human cancers such as Human Papilloma Viruses (HPV), Hepatitis Viruses (HBV and HCV) and Kaposi's Herpes Sarcoma Virus (KHSV). In some embodiments, the compounds of the present disclosure can be used in combination with tumor specific antigen such as heat shock proteins isolated from tumor tissue itself. In some embodiments, the compounds of Formula (I) or any of the formulas as described herein, a compound as recited in any of the claims and described herein, or salts thereof can be combined with dendritic cells immunization to activate potent anti-tumor responses.

The compounds of the present disclosure can be used in combination with bispecific macrocyclic peptides that target Fe alpha or Fe gamma receptor-expressing effectors cells to tumor cells. The compounds of the present disclosure can also be combined with macrocyclic peptides that activate host immune responsiveness.

In some further embodiments, combinations of the compounds of the disclosure with other therapeutic agents can be administered to a patient prior to, during, and/or after a bone marrow transplant or stem cell transplant. The compounds of the present disclosure can be used in combination with bone marrow transplant for the treatment of a variety of tumors of hematopoietic origin.

The compounds of Formula (I) or any of the formulas as described herein, a compound as recited in any of the claims and described herein, or salts thereof can be used in combination with vaccines, to stimulate the immune response to pathogens, toxins, and self antigens. Examples of pathogens for which this therapeutic approach may be particularly useful, include pathogens for which there is currently no effective vaccine, or pathogens for which conventional vaccines are less than completely effective. These include, but are not limited to, HIV, Hepatitis (A, B, & C), Influenza, Herpes, Giardia, Malaria, Leishmania, Staphylococcus aureus, Pseudomonas Aeruginosa.

Viruses causing infections treatable by methods of the present disclosure include, but are not limit to human papillomavirus, influenza, hepatitis A, B, C or D viruses, adenovirus, poxvirus, herpes simplex viruses, human cytomegalovirus, severe acute respiratory syndrome virus, ebola virus, measles virus, herpes virus (e.g., VZV, HSV-1, HAV-6, HSV-II, and CMV, Epstein Barr virus), flaviviruses, echovirus, rhinovirus, coxsackie virus, cornovirus, respiratory syncytial virus, mumpsvirus, rotavirus, measles virus, rubella virus, parvovirus, vaccinia virus, HTLV virus, dengue virus, papillomavirus, molluscum virus, poliovirus, rabies virus, JC virus and arboviral encephalitis virus.

Pathogenic bacteria causing infections treatable by methods of the disclosure include, but are not limited to, chlamydia, rickettsial bacteria, mycobacteria, staphylococci, streptococci, pneumonococci, meningococci and conococci, klebsiella, proteus, serratia, pseudomonas, legionella, diphtheria, salmonella, bacilli, cholera, tetanus, botulism, anthrax, plague, leptospirosis, and Lyme's disease bacteria.

Pathogenic fungi causing infections treatable by methods of the disclosure include, but are not limited to, Candida (albicans, krusei, glabrata, tropicalis, etc.), Cryptococcus neoformans, Aspergillus (fumigatus, niger, etc.), Genus Mucorales (mucor, absidia, rhizophus), Sporothrix schenkii, Blastomyces dermatitidis, Paracoccidioides brasiliensis, Coccidioides immitis and Histoplasma capsulatum.

Pathogenic parasites causing infections treatable by methods of the disclosure include, but are not limited to, Entamoeba histolytica, Balantidium coli, Naegleria fowleri, Acanthamoeba sp., Giardia lamblia, Cryptosporidium sp., Pneumocystis carinii, Plasmodium vivax, Babesia microti, Trypanosoma brucei, Trypanosoma cruzi, Leishmania donovani, Toxoplasma gondi, and Nippostrongylus brasiliensis.

Methods for the safe and effective administration of most of these chemotherapeutic agents are known to those skilled in the art. In addition, their administration is described in the standard literature. For example, the administration of many of the chemotherapeutic agents is described in the "Physicians' Desk Reference" (PDR, e.g., 1996 edition, Medical Economics Company, Montvale, NJ), the disclosure of which is incorporated herein by reference as if set forth in its entirety.

Pharmaceutical Formulations and Dosage Forms

When employed as pharmaceuticals, the compounds of the disclosure can be administered in the form of pharmaceutical compositions. These compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including transdermal, epidermal, ophthalmic and to mucous membranes including intranasal, vaginal and rectal delivery), pulmonary (*e.g.*, by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal or intranasal), oral, or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal intramuscular or injection or infusion; or intracranial, *e.g.*, intrathecal or intraventricular, administration. Parenteral administration can be in the form of a single bolus dose, or may be, for example, by a continuous perfusion pump. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

This disclosure also includes pharmaceutical compositions which contain, as the active ingredient, the compound of the disclosure or a pharmaceutically acceptable salt thereof, in combination with one or more pharmaceutically acceptable carriers (excipients). In some embodiments, the composition is suitable for topical administration. In making the compositions of the disclosure, the active ingredient is typically mixed with an excipient, diluted by an excipient or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, the active compound can be milled to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it can be milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size can be
5 adjusted by milling to provide a substantially uniform distribution in the formulation, *e.g.*, about 40 mesh.

The compounds of the disclosure may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet formation and for other formulation types. Finely divided (nanoparticulate)
10 preparations of the compounds of the disclosure can be prepared by processes known in the art, *e.g.*, see International App. No. WO 2002/000196.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose,
15 water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents. The compositions of the disclosure can be formulated so as to provide quick, sustained or delayed
20 release of the active ingredient after administration to the patient by employing procedures known in the art.

The compositions can be formulated in a unit dosage form, each dosage containing from about 5 to about 1000 mg (1 g), more usually about 100 to about 500 mg, of the active ingredient. The term "unit dosage forms" refers to physically
25 discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

In some embodiments, the compositions of the disclosure contain from about 5 to about 50 mg of the active ingredient. One having ordinary skill in the art will
30 appreciate that this embodies compositions containing about 5 to about 10, about 10 to about 15, about 15 to about 20, about 20 to about 25, about 25 to about 30, about

30 to about 35, about 35 to about 40, about 40 to about 45, or about 45 to about 50 mg of the active ingredient.

In some embodiments, the compositions of the disclosure contain from about 50 to about 500 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compositions containing about 50 to about 100, about 100 to about 150, about 150 to about 200, about 200 to about 250, about 250 to about 300, about 350 to about 400, or about 450 to about 500 mg of the active ingredient.

In some embodiments, the compositions of the disclosure contain from about 500 to about 1000 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compositions containing about 500 to about 550, about 550 to about 600, about 600 to about 650, about 650 to about 700, about 700 to about 750, about 750 to about 800, about 800 to about 850, about 850 to about 900, about 900 to about 950, or about 950 to about 1000 mg of the active ingredient.

Similar dosages may be used of the compounds described herein in the methods and uses of the disclosure.

The active compound can be effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It will be understood, however, that the amount of the compound actually administered will usually be determined by a physician, according to the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present disclosure. When referring to these preformulation compositions as homogeneous, the active ingredient is typically dispersed evenly throughout the composition so that the composition can be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, about 0.1 to about 1000 mg of the active ingredient of the present disclosure.

The tablets or pills of the present disclosure can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the compounds and compositions of the present disclosure can be incorporated for administration orally or by injection include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. In some embodiments, the compositions are administered by the oral or nasal respiratory route for local or systemic effect.

Compositions can be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device can be attached to a face mask, tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions can be administered orally or nasally from devices which deliver the formulation in an appropriate manner.

Topical formulations can contain one or more conventional carriers. In some embodiments, ointments can contain water and one or more hydrophobic carriers selected from, for example, liquid paraffin, polyoxyethylene alkyl ether, propylene glycol, white Vaseline, and the like. Carrier compositions of creams can be based on water in combination with glycerol and one or more other components, e.g. glycerinmonostearate, PEG-glycerinmonostearate and cetylstearyl alcohol. Gels can be formulated using isopropyl alcohol and water, suitably in combination with other components such as, for example, glycerol, hydroxyethyl cellulose, and the like. In

some embodiments, topical formulations contain at least about 0.1, at least about 0.25, at least about 0.5, at least about 1, at least about 2, or at least about 5 wt % of the compound of the disclosure. The topical formulations can be suitably packaged in tubes of, for example, 100 g which are optionally associated with instructions for the treatment of the select indication, e.g., psoriasis or other skin condition.

The amount of compound or composition administered to a patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions can be administered to a patient already suffering from a disease in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. Effective doses will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors such as the severity of the disease, the age, weight and general condition of the patient, and the like.

The compositions administered to a patient can be in the form of pharmaceutical compositions described above. These compositions can be sterilized by conventional sterilization techniques, or may be sterile filtered. Aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

The therapeutic dosage of a compound of the present disclosure can vary according to, for example, the particular use for which the treatment is made, the manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. The proportion or concentration of a compound of the disclosure in a pharmaceutical composition can vary depending upon a number of factors including dosage, chemical characteristics (e.g., hydrophobicity), and the route of administration. For example, the compounds of the disclosure can be provided in an aqueous physiological buffer solution containing about 0.1 to about 10% w/v of the compound for parenteral administration. Some

typical dose ranges are from about 1 µg/kg to about 1 g/kg of body weight per day. In some embodiments, the dose range is from about 0.01 mg/kg to about 100 mg/kg of body weight per day. The dosage is likely to depend on such variables as the type and extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, formulation of the excipient, and its route of administration. Effective doses can be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

The compositions of the disclosure can further include one or more additional pharmaceutical agents such as a chemotherapeutic, steroid, anti-inflammatory compound, or immunosuppressant, examples of which are listed herein.

Labeled Compounds and Assay Methods

Another aspect of the present disclosure relates to labeled compounds of the disclosure (radio-labeled, fluorescent-labeled, etc.) that would be useful not only in imaging techniques but also in assays, both *in vitro* and *in vivo*, for localizing and quantitating A2A and/or A2B receptors in tissue samples, including human, and for identifying A2A and/or A2B antagonists by inhibition binding of a labeled compound. Substitution of one or more of the atoms of the compounds of the present disclosure can also be useful in generating differentiated ADME (Adsorption, Distribution, Metabolism and Excretion.) Accordingly, the present disclosure includes adenosine receptor (e.g., A2A and/or A2B) assays that contain such labeled or substituted compounds.

The present disclosure further includes isotopically-labeled compounds of the disclosure. An “isotopically” or “radio-labeled” compound is a compound of the disclosure where one or more atoms are replaced or substituted by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature (i.e., naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present disclosure include but are not limited to ²H (also written as D for deuterium), ³H (also written as T for tritium), ¹¹C, ¹³C, ¹⁴C, ¹³N, ¹⁵N, ¹⁵O, ¹⁷O, ¹⁸O, ¹⁸F, ³⁵S, ³⁶Cl, ⁸²Br, ⁷⁵Br, ⁷⁶Br, ⁷⁷Br, ¹²³I, ¹²⁴I, ¹²⁵I and ¹³¹I. For example, one or more hydrogen atoms in a compound of the present disclosure can be

replaced by deuterium atoms (e.g., one or more hydrogen atoms of a C₁₋₆ alkyl group of Formula (I) can be optionally substituted with deuterium atoms, such as –CD₃ being substituted for –CH₃). In some embodiments, alkyl groups in any of the disclosed Formulas, e.g., Formula (I), can be perdeuterated.

5 One or more constituent atoms of the compounds presented herein can be replaced or substituted with isotopes of the atoms in natural or non-natural abundance. In some embodiments, the compound includes at least one deuterium atom. For example, one or more hydrogen atoms in a compound presented herein can be replaced or substituted by deuterium (e.g., one or more hydrogen atoms of a C₁₋₆ alkyl
10 group can be replaced by deuterium atoms, such as –CD₃ being substituted for –CH₃). In some embodiments, the compound includes two or more deuterium atoms. In some embodiments, the compound includes 1-2, 1-3, 1-4, 1-5, or 1-6 deuterium atoms. In some embodiments, all of the hydrogen atoms in a compound can be replaced or substituted by deuterium atoms.

15 In some embodiments, 1, 2, 3, 4, 5, 6, 7, or 8 hydrogen atoms, attached to carbon atoms of any “alkyl”, “alkenyl”, “alkynyl”, “aryl”, “phenyl”, “cycloalkyl”, “heterocycloalkyl”, or “heteroaryl” substituents or “-C₁₋₆ alkyl-”, “alkylene”, “alkenylene” and “alkynylene” linking groups, as described herein, are each optionally replaced by a deuterium atom.

20 Synthetic methods for including isotopes into organic compounds are known in the art (Deuterium Labeling in Organic Chemistry by Alan F. Thomas (New York, N.Y., Appleton-Century-Crofts, 1971; The Renaissance of H/D Exchange by Jens Atzrodt, Volker Derdau, Thorsten Fey and Jochen Zimmermann, Angew. Chem. Int. Ed. 2007, 7744-7765; The Organic Chemistry of Isotopic Labelling by James R.
25 Hanson, Royal Society of Chemistry, 2011). Isotopically labeled compounds can be used in various studies such as NMR spectroscopy, metabolism experiments, and/or assays.

Substitution with heavier isotopes, such as deuterium, may afford certain therapeutic advantages resulting from greater metabolic stability, for example,
30 increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances. (see e.g., A. Kerekes et. al. *J. Med. Chem.* 2011, 54, 201-210; R. Xu et. al. *J. Label Compd. Radiopharm.* 2015, 58, 308-312). In

particular, substitution at one or more metabolism sites may afford one or more of the therapeutic advantages.

The radionuclide that is incorporated in the instant radio-labeled compounds will depend on the specific application of that radio-labeled compound. For example, for *in vitro* adenosine receptor labeling and competition assays, compounds that incorporate ^3H , ^{14}C , ^{82}Br , ^{125}I , ^{131}I or ^{35}S can be useful. For radio-imaging applications ^{11}C , ^{18}F , ^{125}I , ^{123}I , ^{124}I , ^{131}I , ^{75}Br , ^{76}Br or ^{77}Br can be useful.

It is understood that a “radio-labeled” or “labeled compound” is a compound that has incorporated at least one radionuclide. In some embodiments, the radionuclide is selected from the group consisting of ^3H , ^{14}C , ^{125}I , ^{35}S and ^{82}Br .

The present disclosure can further include synthetic methods for incorporating radio-isotopes into compounds of the disclosure. Synthetic methods for incorporating radio-isotopes into organic compounds are well known in the art, and an ordinary skill in the art will readily recognize the methods applicable for the compounds of disclosure.

A labeled compound of the disclosure can be used in a screening assay to identify/evaluate compounds. For example, a newly synthesized or identified compound (*i.e.*, test compound) which is labeled can be evaluated for its ability to bind an adenosine receptor by monitoring its concentration variation when contacting with the adenosine receptor, through tracking of the labeling. For example, a test compound (labeled) can be evaluated for its ability to reduce binding of another compound which is known to bind to an adenosine receptor (*i.e.*, standard compound). Accordingly, the ability of a test compound to compete with the standard compound for binding to the adenosine receptor directly correlates to its binding affinity. Conversely, in some other screening assays, the standard compound is labeled and test compounds are unlabeled. Accordingly, the concentration of the labeled standard compound is monitored in order to evaluate the competition between the standard compound and the test compound, and the relative binding affinity of the test compound is thus ascertained.

Kits

The present disclosure also includes pharmaceutical kits useful, for example, in the treatment or prevention of adenosine receptor-associated diseases or disorders (such as, e.g., cancer, an inflammatory disease, a cardiovascular disease, or a neurodegenerative disease) which include one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a compound of the disclosure. Such kits can further include, if desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, can also be included in the kit.

The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of non-critical parameters which can be changed or modified to yield essentially the same results. The compounds of the Examples have been found to inhibit the activity of an adenosine receptor (e.g., A2A and/or A2B) according to at least one assay described herein.

EXAMPLES

Preparatory LC-MS purifications of some of the compounds prepared were performed on Waters mass directed fractionation systems. The basic equipment setup, protocols, and control software for the operation of these systems have been described in detail in the literature (see *e.g.* “Two-Pump At Column Dilution Configuration for Preparative LC-MS”, K. Blom, *J. Combi. Chem.*, 4, 295 (2002); “Optimizing Preparative LC-MS Configurations and Methods for Parallel Synthesis Purification”, K. Blom, R. Sparks, J. Doughty, G. Everlof, T. Haque, A. Combs, *J. Combi. Chem.*, 5, 670 (2003); and “Preparative LC-MS Purification: Improved Compound Specific Method Optimization”, K. Blom, B. Glass, R. Sparks, A. Combs, *J. Combi. Chem.*, 6, 874-883 (2004)). The compounds separated were typically subjected to analytical

liquid chromatography mass spectrometry (LCMS) for purity analysis under the following conditions: Instrument; Agilent 1100 series, LC/MSD, Column: Waters Sunfire™ C₁₈ 5 μm, 2.1 x 50 mm, Buffers: mobile phase A: 0.025% TFA in water and mobile phase B: acetonitrile; gradient 2% to 80% of B in 3 minutes with flow rate

5 2.0 mL/minute.

Some of the compounds prepared were also separated on a preparative scale by reverse-phase high performance liquid chromatography (RP-HPLC) with MS detector or flash chromatography (silica gel) as indicated in the Examples. Typical preparative reverse-phase high performance liquid chromatography (RP-HPLC)

10 column conditions are as follows:

pH = 2 purifications: Waters Sunfire™ C₁₈ 5 μm, 30 x 100 mm or Waters XBridge™ C₁₈ 5 μm, 30 x 100 mm column, eluting with mobile phase A: 0.1% TFA (trifluoroacetic acid) in water and mobile phase B: acetonitrile; the flow rate was 60 mL/minute, the separating gradient was optimized for each compound using the


15 Compound Specific Method Optimization protocol as described in the literature (see *e.g.* "Preparative LCMS Purification: Improved Compound Specific Method Optimization", K. Blom, B. Glass, R. Sparks, A. Combs, *J. Comb. Chem.*, **6**, 874-883 (2004)).

pH = 10 purifications: Waters XBridge™ C₁₈ 5 μm, 30 x 100 mm column, eluting with mobile phase A: 0.1% NH₄OH in water and mobile phase B: acetonitrile; the flow rate was 60 mL/minute, the separating gradient was optimized for each compound using the Compound Specific Method Optimization protocol as described in the literature (see *e.g.* "Preparative LCMS Purification: Improved Compound Specific Method Optimization", K. Blom, B. Glass, R. Sparks, A. Combs, *J. Comb.*


20 *Chem.*, **6**, 874-883 (2004)).

25

Example 1. 4-Amino-6-(3-cyanophenyl)-N-ethyl-7-(1-ethyl-1H-pyrazol-5-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide




Step 1: Diethyl 1-(2-(3-cyanophenyl)-2-oxoethyl)-1H-pyrazole-3,5-dicarboxylate




To a solution of diethyl 1H-pyrazole-3,5-dicarboxylate (12.9 g, 60.8 mmol), 3-(2-bromoacetyl)benzonitrile (13.62 g, 60.8 mmol) in acetone (253 mL) was added
 5 potassium carbonate (9.24 g, 66.9 mmol). The mixture was stirred at room temperature (rt or RT) for 12 h. The reaction mixture was concentrated and the residue was taken up in water and dichloromethane (DCM). The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as a white solid (21.6
 10 g, 100%). LC-MS calculated for C₁₈H₁₈N₃O₅ (M+H)⁺: m/z = 356.1; found 356.1.

Step 2: Ethyl 6-(3-cyanophenyl)-4-oxo-4,5-dihydropyrazolo[1,5-a]pyrazine-2-carboxylate




15 Diethyl 1-(2-(3-cyanophenyl)-2-oxoethyl)-1H-pyrazole-3,5-dicarboxylate (21.6 g, 60.8 mmol) was dissolved in acetic acid (260 mL), and ammonium acetate (46.9 g, 608 mmol) was added. The mixture was stirred at 110 °C for 36 h. After cooling to rt, the mixture was diluted with water, the precipitate was collected via filtration, washed with water, and dried to give the product. LC-MS calculated for
 20 C₁₆H₁₃N₄O₃ (M+H)⁺: m/z = 309.1; found 309.1.

Step 3: Ethyl 4-chloro-6-(3-cyanophenyl)pyrazolo[1,5-a]pyrazine-2-carboxylate




A mixture of ethyl 6-(3-cyanophenyl)-4-oxo-4,5-dihydropyrazolo[1,5-*a*]pyrazine-2-carboxylate (15.8 g, 51.2 mmol) and POCl₃ (96 mL, 1025 mmol) was heated at 110 °C for 4 h. After cooling to rt, the mixture was slowly added to a flask containing ice and sodium bicarbonate. The resulting precipitate was collected,
 5 washed with water, and dried to give the product (15.8 g, 94%). LC-MS calculated for C₁₆H₁₂ClN₄O₂ (M+H)⁺: m/z = 327.1; found 327.1.

*Step 4. Ethyl 6-(3-cyanophenyl)-4-((2,4-dimethoxybenzyl)amino)pyrazolo[1,5-*a*]pyrazine-2-carboxylate*




A microwave vial was charged with ethyl 4-chloro-6-(3-cyanophenyl)pyrazolo[1,5-*a*]pyrazine-2-carboxylate (1.22 g, 3.73 mmol), (2,4-dimethoxyphenyl)methanamine (0.749 g, 4.48 mmol), N,N-diisopropylethylamine (DIEA, 1.304 mL, 7.47 mmol) and butan-1-ol (13.0 mL). The mixture was heated at
 15 180 °C for 30 min in microwave reactor. The mixture was diluted with water, and the resulting precipitate was collected via filtration, washed with water, and dried to give the product (1.5 g, 88%). LC-MS calculated for C₂₅H₂₄N₅O₄ (M+H)⁺: m/z = 458.2; found 458.2.

*Step 5: 6-(3-Cyanophenyl)-4-((2,4-dimethoxybenzyl)amino)pyrazolo[1,5-*a*]pyrazine-2-carboxylic acid*




A mixture of ethyl 6-(3-cyanophenyl)-4-((2,4-dimethoxybenzyl)amino)pyrazolo[1,5-*a*]pyrazine-2-carboxylate (1.35 g, 2.95 mmol), sodium hydroxide (5.90 mL, 5.90 mmol), and acetonitrile (20 mL) was stirred at room temperature for 2 h. The reaction was diluted with 1 N HCl (6 mL). The precipitate
 5 was collected via filtration, washed with water, and dried to give the product (1.0 g, 79%). LC-MS calculated for $C_{23}H_{20}N_5O_4$ ($M+H$)⁺: m/z = 430.1; found 430.1.

*Step 6. 4-Amino-6-(3-cyanophenyl)-N-ethylpyrazolo[1,5-*a*]pyrazine-2-carboxamide*



To a vial was added 6-(3-cyanophenyl)-4-((2,4-dimethoxybenzyl)amino)pyrazolo[1,5-*a*]pyrazine-2-carboxylic acid (0.5 g, 1.164 mmol), PyBOP (0.727 g, 1.397 mmol), and dimethylformamide (DMF, 1.0 mL), followed by 2.0 M ethanamine in tetrahydrofuran (THF, 1.164 mL, 2.329 mmol) and *N,N*-diisopropylethylamine (1.017 mL, 5.82 mmol). After stirring at room
 15 temperature (rt) for 2 h, the reaction mixture was diluted with water and DCM. The organic layer was dried over Na_2SO_4 , filtered, and concentrated. The crude was treated with trifluoroacetic acid (TFA, 0.5 mL) and heated at 90 °C for 30 min. After removing the solvent, the resulting precipitate was washed with water and ethyl acetate to give the desired product as white solid (0.32 g, 90%). LC-MS calculated for
 20 $C_{16}H_{15}N_6O$ ($M+H$)⁺: m/z = 307.1; found 307.1.

*Step 7. 4-Amino-7-bromo-6-(3-cyanophenyl)-N-ethylpyrazolo[1,5-*a*]pyrazine-2-carboxamide*



To a solution of 4-amino-6-(3-cyanophenyl)-*N*-ethylpyrazolo[1,5-*a*]pyrazine-2-carboxamide (747 mg, 2.439 mmol) in DCM (5 mL) and DMF (1.250 mL) was added *N*-bromosuccinimide (NBS, 421 mg, 2.365 mmol). The resulting mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted with DCM

and water. The organic layer was dried over Na₂SO₄, filtered, and concentrated. The precipitate was collected and washed with ethyl acetate to give the desired product as white solid (0.75 g, 80%). LC-MS calculated for C₁₆H₁₄BrN₆O (M+H)⁺: m/z = 385.0, 387.0; found 385.0, 387.0.


5

Step 8. 4-Amino-6-(3-cyanophenyl)-N-ethyl-7-(1-ethyl-1H-pyrazol-5-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide

A mixture of 4-amino-7-bromo-6-(3-cyanophenyl)-N-ethylpyrazolo[1,5-a]pyrazine-2-carboxamide (14 mg, 0.036 mmol), 1-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (9.69 mg, 0.044 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (2.86 mg, 3.63 μmol) and tripotassium phosphate hydrate (18.41 mg, 0.080 mmol) in 1,4-dioxane (0.6 mL)/water (0.200 mL) was stirred at 80 °C for 1 h. The residue was dissolved in methanol and 1 N HCl and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₁H₂₁N₈O (M+H)⁺: m/z = 401.2; found 401.2. ¹H NMR (600 MHz, DMSO) δ 8.07 (t, *J* = 6.0 Hz, 1H), 7.80 – 7.73 (m, 3H), 7.73 – 7.70 (m, 1H), 7.56 – 7.46 (m, 4H), 6.32 (d, *J* = 1.8 Hz, 1H), 3.87 (m, 1H), 3.75 (m, 1H), 3.27 (m, 2H), 1.16 (t, *J* = 7.2 Hz, 3H), 1.08 (t, *J* = 7.1 Hz, 3H).

20


Example 2. 4-Amino-6-(3-cyanophenyl)-N-ethyl-7-(1-propyl-1H-pyrazol-5-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide



The title compound was prepared using similar procedures as described for Example 1, with 1-propyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole replacing 1-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole in Step 8. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₂H₂₃N₈O (M+H)⁺: m/z = 415.2; found 415.2.


30

Example 3. 4-Amino-6-(3-cyanophenyl)-*N*-ethyl-7-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)pyrazolo[1,5-*a*]pyrazine-2-carboxamide



The title compound was prepared using similar procedures as described for Example 1 with 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2(1*H*)-one replacing 1-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole in Step 8. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₂H₂₀N₇O₂ (M+H)⁺: m/z = 414.2; found 414.2.

Example 4. 4-Amino-6-(3-cyanophenyl)-*N*-ethyl-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazine-2-carboxamide




A mixture of 4-amino-7-bromo-6-(3-cyanophenyl)-*N*-ethylpyrazolo[1,5-*a*]pyrazine-2-carboxamide (Example 1, Step 7; 10 mg, 0.026 mmol), 4-(tributylstannyl)pyrimidine (14.4 mg, 0.039 mmol), and copper(I) chloride (3.1 mg, 0.031 mmol), lithium chloride (1.3 mg, 0.031 mmol) and tetrakis(triphenylphosphine)palladium(0) (3.0 mg, 2.60 μmol) in THF (1.0 mL) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt (4.2 mg, 42%). LC-MS calculated for C₂₀H₁₇N₈O (M+H)⁺: m/z = 385.1; found 385.1.

Example 5. 4-Amino-6-(3-cyanophenyl)-7-(1,1-difluoroethyl)-*N*-ethylpyrazolo[1,5-*a*]pyrazine-2-carboxamide




To a vial was added 4-amino-6-(3-cyanophenyl)-*N*-ethylpyrazolo[1,5-*a*]pyrazine-2-carboxamide (Example 1, Step 6; 20.0 mg, 0.065 mmol), sodium 1,1-difluoroethane-1-sulfinate (59.6 mg, 0.392 mmol), diethyl carbonate (2.0 mL), water (1.3 mL) and *tert*-butyl hydroperoxide (0.090 mL, 0.653 mmol). The resulting mixture was heated at 90 °C for 3 h. The reaction mixture was purified by prep-LCMS (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt (5.6 mg, 23%). LC-MS calculated for C₁₈H₁₇F₂N₆O (M+H)⁺: m/z = 371.1; found 371.1.

Example 6. 4-Amino-6-(3-cyanophenyl)-*N*-ethyl-7-(1-(trifluoromethyl)cyclopropyl)pyrazolo[1,5-*a*]pyrazine-2-carboxamide




The title compound was prepared using similar procedures as described for Example 5 with sodium 1-(trifluoromethyl)cyclopropane-1-sulfinate replacing sodium 1,1-difluoroethane-1-sulfinate. The reaction mixture was purified by prep-LCMS (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₀H₁₈F₃N₆O (M+H)⁺: m/z = 415.1; found 415.1.

Example 7. 3-(4-Amino-2-(azetidine-1-carbonyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




Step 1. Ethyl 4-amino-7-bromo-6-(3-cyanophenyl)pyrazolo[1,5-a]pyrazine-2-carboxylate




To a solution of ethyl 6-(3-cyanophenyl)-4-((2,4-dimethoxybenzyl)amino)pyrazolo[1,5-a]pyrazine-2-carboxylate (Example 1, Step 4; 8.35 g, 18.26 mmol) was treated with TFA (20 mL) and heated at 90 °C for 30 min. After removing the solvent, the resulting precipitate was washed with water and ethyl acetate. The crude product was dissolved in DCM (73.0 mL) and DMF (18.26 mL), to this solution NBS (3.15 g, 17.71 mmol) was added. The resulting mixture was stirred at room temperature for 1 h. The reaction mixture was diluted with DCM and water. The precipitate was collected via filtration and washed with water to give the desired product as a white solid (5.6 g, 79%). LC-MS calculated for $C_{16}H_{13}BrN_5O_2$ (M+H)⁺: m/z = 386.0; found 386.0.

Step 2. Ethyl 4-amino-6-(3-cyanophenyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazine-2-carboxylate



A mixture of ethyl 4-amino-7-bromo-6-(3-cyanophenyl)pyrazolo[1,5-a]pyrazine-2-carboxylate (904 mg, 2.341 mmol), 4-(tributylstannyl)pyrimidine (1.3 g, 3.51 mmol), copper(I) chloride (278 mg, 2.81 mmol), lithium chloride (119 mg, 2.81 mmol), and tetrakis(triphenylphosphine)palladium(0) (270 mg, 0.234 mmol) in THF (15 mL) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction was diluted with ethyl acetate and water, and the aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product (0.34 g, 38%). LC-MS calculated for $C_{20}H_{16}N_7O_2$ (M+H)⁺: m/z = 386.1; found 386.2.

Step 3. 4-Amino-6-(3-cyanophenyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazine-2-carboxylic acid




A mixture of ethyl 4-amino-6-(3-cyanophenyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazine-2-carboxylate (340 mg, 0.882 mmol), 1.0 M sodium hydroxide (4.41 mL, 4.41 mmol), acetonitrile (10 mL), and THF (5.00 mL) was stirred at room temperature for 2 h. The reaction was quenched with 1 N HCl to pH 4. After removing most of the organic solvent, the precipitate was collected via filtration, washed with water, and dried under vacuum to give the desired product as a white solid (295 mg, 94%). LC-MS calculated for C₁₈H₁₂N₇O₂ (M+H)⁺: m/z = 358.1; found 358.1.

Step 4. 3-(4-Amino-2-(azetidine-1-carbonyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile


To a solution of 4-amino-6-(3-cyanophenyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazine-2-carboxylic acid (8.0 mg, 0.022 mmol) and HATU (8.51 mg, 0.022 mmol) in *N,N*-dimethylformamide (1.0 mL), was added azetidine (3.02 μl, 0.045 mmol) and DIEA (7.82 μl, 0.045 mmol). After stirring at rt for 2 h, The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt (2.5 mg, 28%). LC-MS calculated for C₂₁H₁₇N₈O (M+H)⁺: m/z = 397.1; found 397.1. ¹H NMR (500 MHz, DMSO) δ 9.11 (m, 1H), 8.94 (d, *J* = 5.2 Hz, 1H), 7.96 – 7.90 (m, 2H), 7.79 – 7.73 (m, 3H), 7.55 (m, 2H), 7.47 (t, *J* = 7.7 Hz, 1H), 4.29 (t, *J* = 7.7 Hz, 2H), 4.03 (t, *J* = 7.7 Hz, 2H), 2.24 (m, 2H).

Example 8. 3-(4-Amino-7-(pyrimidin-4-yl)-2-(pyrrolidine-1-carbonyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for
 5 Example 7 with pyrrolidine replacing azetidine in Step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₂H₁₉N₈O (M+H)⁺: m/z = 411.2; found 411.2.

Example 9. 3-(4-Amino-2-(piperidine-1-carbonyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



The title compound was prepared using similar procedures as described for
 Example 7 with piperidine replacing azetidine in Step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product
 15 as TFA salt. LC-MS calculated for C₂₃H₂₁N₈O (M+H)⁺: m/z = 425.2; found 425.3.

Example 10. 4-Amino-6-(3-cyanophenyl)-*N,N*-diethyl-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazine-2-carboxamide




The title compound was prepared using similar procedures as described for
 20 Example 7 with diethylamine replacing azetidine in Step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₂H₂₁N₈O (M+H)⁺: m/z = 413.2; found 413.2.

¹H NMR (500 MHz, DMSO) δ 9.16 (d, *J* = 1.3 Hz, 1H), 8.93 (d, *J* = 5.2 Hz, 1H), 8.02-7.95 (m, 2H), 7.87 (m, 1H), 7.83 – 7.75 (m, 2H), 7.58 – 7.38 (m, 3H), 3.50 (q, *J* = 6.8 Hz, 2H), 3.41 (q, *J* = 7.0 Hz, 2H), 1.12 (t, *J* = 7.0 Hz, 3H), 1.00 (t, *J* = 6.9 Hz, 3H).


5

Example 11. 4-Amino-6-(3-cyanophenyl)-*N*-ethyl-*N*-methyl-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazine-2-carboxamide




The title compound was prepared using similar procedures as described for Example 7 with *N*-methylethanamine replacing azetidine in Step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₁H₁₉N₈O (M+H)⁺: m/z = 399.2; found 399.2.

Example 12. 3-(4-Amino-2-(3-hydroxyazetidine-1-carbonyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for Example 7 with azetidin-3-ol replacing azetidine in Step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₁H₁₇N₈O₂ (M+H)⁺: m/z = 413.1; found 413.1.

Example 13. 3-(4-Amino-2-(azetidin-1-ylmethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




5 Step 1. 3-(4-Amino-7-bromo-2-(hydroxymethyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



To a solution of ethyl 4-amino-7-bromo-6-(3-cyanophenyl)pyrazolo[1,5-*a*]pyrazine-2-carboxylate (Example 7, Step 1; 0.547 g, 1.416 mmol) in CH₂Cl₂ (7.08 mL) and THF (7.08 mL) was added 1.0 M DIBAL-H in THF (4.25 mL, 4.25 mmol) at 0 °C. The resulting mixture was warmed to room temperature and stirred overnight. The reaction was diluted with DCM and 1 N NaOH solution. The organic layer was separated and dried over Na₂SO₄, filtered, and concentrated. The crude was used in the next step without purification. LC-MS calculated for C₁₄H₁₁BrN₅O (M+H)⁺: m/z = 344.0; found 344.0.


Step 2. 3-(4-Amino-2-(hydroxymethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



A mixture of 3-(4-amino-7-bromo-2-(hydroxymethyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (487 mg, 1.415 mmol), 4-(tributylstannyl)pyrimidine (575 mg, 1.556 mmol), copper(I) chloride (168 mg, 1.698 mmol), lithium chloride (72.0 mg, 1.698 mmol) and tetrakis(triphenylphosphine)palladium(0) (164 mg, 0.141 mmol) in THF (12 mL) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The

reaction was diluted with ethyl acetate and water, the aqueous layer was extracted with ethyl acetate once. The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified with flash chromatography to give the desired product (0.16 g, 34% for 2 steps). LC-MS
 5 calculated for C₁₈H₁₄N₇O (M+H)⁺: m/z = 344.1; found 344.1.

Step 3. 3-(4-Amino-2-formyl-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(4-amino-2-(hydroxymethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (160 mg, 0.466 mmol), Dess-Martin periodinane (237 mg, 0.559 mmol), and CH₂Cl₂ (4660 μL) was stirred at room temperature for 2 h. The reaction was diluted with DCM and saturated NaHCO₃ solution. After stirring for 30 min, the organic layer was separated and dried over Na₂SO₄, filtered, and
 10 concentrated. The crude was used in the next step without purification. LC-MS calculated for C₁₈H₁₂N₇O (M+H)⁺: m/z = 342.1; found 342.1.


Step 4. 3-(4-Amino-2-(azetidin-1-ylmethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile

To a mixture of 3-(4-amino-2-formyl-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.029 mmol) and azetidine (3.35 mg, 0.059 mmol) in DCM (1 mL) was added sodium triacetoxyborohydride (12.4 mg, 0.059 mmol). After stirring at room temperature for 2.5 h, solvent was removed *in vacuo*. The resulting residue was dissolved in methanol and 1 N HCl (1 N) and purified with prep-LCMS (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt.
 20
 25 LC-MS calculated for C₂₁H₁₉N₈ (M+H)⁺: m/z = 383.2; found 383.2.

Example 14. Ethyl (8-amino-6-(3-cyanophenyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)carbamate




5 *Step 1. tert-Butyl ((mesitylsulfonyl)oxy)carbamate*



To a solution of 2,4,6-trimethylbenzenesulfonyl chloride (9.10 g, 41.6 mmol) and *tert*-butyl *N*-hydroxycarbamate (5.54 g, 41.6 mmol) in methyl *tert*-butyl ether (MTBE, 90 mL) was added triethylamine (TEA, 6.09 mL, 43.7 mmol) dropwise while
 10 stirring at 0 °C. The resulting suspension was stirred at 0 °C for an additional 30 min and then warmed to ambient temperature. The reaction was then diluted with water (90 mL) and adjusted to pH 4 with 1 N HCl. The organic layer was washed with brine, dried over Na₂SO₄, filtered, and concentrated to give the desired product. LC-MS calculated for C₁₄H₂₂NO₅S (M+H)⁺: m/z = 316.1; found 316.1.


15

Step 2. O-(Mesitylsulfonyl)hydroxylamine




To TFA (37.7 mL, 490 mmol) at 0 °C was slowly added *tert*-butyl ((mesitylsulfonyl)oxy)carbamate (12.56 g, 39.8 mmol). The reaction mixture was
 20 stirred at 0 °C for 1.5 h and then quenched with the sequential addition of crushed ice and water. The resulting white suspension was vigorously stirred at ambient temperature for 5 min. Without allowing the filter cake to run dry, the solids were collected by careful vacuum filtration followed by subsequent rinsing with water until the filtrate reached pH 6. The wet filtrate was taken up in DCM and the resulting
 25 biphasic solution was separated. The DCM layer was dried over MgSO₄ for 30 min and then filtered and rinsed with DCM to provide the compound as a solution. LC-MS calculated for C₉H₁₄NO₃S (M+H)⁺: m/z = 216.1; found 216.1.

Step 3. 1,2-Diamino-3,5-dibromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate




To a solution of O-(mesitylsulfonyl)hydroxylamine (2.468 g, 11.47 mmol) in CH_2Cl_2 (50 mL) was added 3,5-dibromopyrazin-2-amine (2.90 g, 11.47 mmol), and the resulting solution was stirred at ambient temperature overnight. The precipitate was collected via filtration and dried under vacuum. LC-MS calculated for $\text{C}_4\text{H}_5\text{Br}_2\text{N}_4$ (M)⁺: m/z = 266.9; found 266.9.

Step 4. Ethyl (6,8-dibromo-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)carbamate



To a suspension of 1,2-diamino-3,5-dibromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate (190 mg, 0.406 mmol) in DCM (1.0 mL) and *N,N*-dimethylformamide (1 mL) was added *O*-ethyl carbonisothiocyanatide (52.7 μL , 0.446 mmol). The resulting mixture was stirred at room temperature for 3 h. The reaction mixture was diluted with DCM and water. The organic layer was separated and dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue was purified with flash chromatography to give the desired product (10 mg, 8%) LC-MS calculated for $\text{C}_8\text{H}_8\text{Br}_2\text{N}_5\text{O}_2$ ($\text{M}+\text{H}$)⁺: m/z = 365.9; found 365.8.


Step 5. Ethyl (6-bromo-8-((2,4-dimethoxybenzyl)amino)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)carbamate



To a mixture of ethyl (6,8-dibromo-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)carbamate (10 mg, 0.027 mmol) and (2,4-dimethoxyphenyl)methanamine (4.58 mg, 0.027 mmol) in DCM (1 mL) was added DIEA (9.57 μL , 0.055 mmol). After

stirring at 40 °C for 2.5 h, the solvent was removed *in vacuo*. The residue was purified with flash chromatography to give the desired product (6.0 mg, 49%). LC-MS calculated for $C_{17}H_{20}BrN_6O_4$ ($M+H$)⁺: m/z = 451.1, 453.1; found 451.1, 453.1.

5 *Step 6. Ethyl (6-(3-cyanophenyl)-8-((2,4-dimethoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)carbamate*




A mixture of ethyl (6-bromo-8-((2,4-dimethoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)carbamate (6.0 mg, 0.013 mmol), (3-cyanophenyl)boronic acid (1.954 mg, 0.013 mmol), dicyclohexyl(2',4',6'-triiisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (1.046 mg, 1.330 μ mol) and tripotassium phosphate hydrate (6.74 mg, 0.029 mmol) in 1,4-dioxane (0.6 mL)/water (0.200 mL) was stirred at 70 °C for 1 h. The resulting residue was dissolved in methanol and 1 N HCl and purified with prep-LCMS (pH 2) to give the desired product as white solid (4.2 mg, 66%). LC-MS calculated for $C_{24}H_{24}N_7O_4$ ($M+H$)⁺: m/z = 474.2; found 474.2.


Step 7. Ethyl (8-amino-6-(3-cyanophenyl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)carbamate

20 A mixture of ethyl (6-(3-cyanophenyl)-8-((2,4-dimethoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)carbamate (6.3 mg, 0.013 mmol) and trifluoroacetic acid (0.3 mL) was stirred at 90 °C for 30 min. The volatiles were removed and the resulting residue was diluted with methanol and purified with prep-LCMS (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt (0.3 mg, 7%). LC-MS calculated for $C_{15}H_{14}N_7O_2$ ($M+H$)⁺: m/z = 324.1; found 324.1.

Example 15. 3-(8-Amino-5-(6-oxo-1,6-dihydropyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




5 Step 1. 6-Bromo-N-(2,4-dimethoxybenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine



A vial was charged with 6,8-dibromo-[1,2,4]triazolo[1,5-a]pyrazine (720 mg, 2.59 mmol), (2,4-dimethoxyphenyl)methanamine (433 mg, 2.59 mmol), DIEA (679 μ l, 3.89 mmol), 2-propanol (6 mL), and *N,N*-dimethylformamide (6 mL). The mixture was heated at 90 °C for 2 h and then diluted with water. The resulting precipitate was collected via filtration (0.94 g, 100%). LC-MS calculated for $C_{14}H_{15}BrN_5O_2$ ($M+H$)⁺: m/z = 364.0, 366.0; found 364.0, 366.0

15 Step 2. 3-(8-((2,4-Dimethoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 6-bromo-N-(2,4-dimethoxybenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine (0.94 g, 2.58 mmol), (3-cyanophenyl)boronic acid (0.417 g, 2.84 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.203 g, 0.258 mmol) and tripotassium phosphate hydrate

(1.486 g, 6.45 mmol) in 1,4-dioxane (9.68 mL)/water (3.23 mL) was stirred at 70 °C for 1 h. After cooling to rt, the mixture was diluted with water. The resulting precipitate was collected via filtration (0.7g, 70%). LC-MS calculated for C₂₁H₁₉N₆O₂ (M+H)⁺: m/z = 387.1; found 387.1.

5


Step 3. 3-(8-Amino-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-((2,4-dimethoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.70 g, 1.812 mmol) and trifluoroacetic acid (4.19 mL, 54.3 mmol) was heated at 90 °C for 30 min. The volatiles were removed *in vacuo*. The resulting solid was washed with water and ethyl acetate and dried under vacuum (0.35 g, 82%). LC-MS calculated for C₁₂H₉N₆(M+H)⁺: m/z = 237.1; found 237.1.

10

Step 4. 3-(8-Amino-5-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




15

To a solution of 3-(8-amino-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (129 mg, 0.546 mmol) in DCM (5 mL) and DMF (1.250 mL) was added NBS (94 mg, 0.530 mmol). The resulting mixture was stirred at room temperature for 1 h. The reaction mixture was diluted with DCM and water. The precipitate was collected via filtration and washed with water and ethyl acetate (135 mg, 78%) LC-MS calculated for C₁₂H₈BrN₆ (M+H)⁺: m/z = 315.0, 317.0; found 315.0, 317.0.

20

Step 5. 3-(8-Amino-5-(6-methoxypyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-amino-5-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (20 mg, 0.063 mmol), (6-methoxypyridin-3-yl)boronic acid (9.71 mg, 0.063 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (5.0 mg, 6.35 μ mol), and tripotassium phosphate hydrate (32.2 mg, 0.140 mmol) in 1,4-dioxane (0.6mL)/water (0.200 mL) was stirred at 70 °C for 1 h. The residue was dissolved in methanol and 1 N HCl and purified with prep-LCMS (pH = 2, acetonitrile/water+TFA) to give the desired product as white solid (12 mg, 55%). LC-MS calculated for $C_{18}H_{14}N_7O$ (M+H)⁺: m/z = 344.1; found 344.1.

Step 6. 3-(8-Amino-5-(6-oxo-1,6-dihydropyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile


A mixture of 3-(8-amino-5-(6-methoxypyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.029 mmol), potassium iodide (14.50 mg, 0.087 mmol), and acetic acid (1.0 mL) was heated at 90 °C for 1 h. The mixture was diluted with methanol and purified with prep-LCMS (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt (3.5 mg, 37%). LC-MS calculated for $C_{17}H_{12}N_7O$ (M+H)⁺: m/z = 330.1; found 330.1.

Example 16. 3-(4-Amino-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




25

Step 1: 3-(4-(2,4-Dimethoxybenzylamino)-2-(hydroxymethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of ethyl 6-(3-cyanophenyl)-4-((2,4-dimethoxybenzyl)amino)pyrazolo[1,5-a]pyrazine-2-carboxylate (Example 1, Step 4; 4.00 g, 8.74 mmol) in THF (200 mL) was added diisobutylaluminum hydride (1.0 M toluene solution, 35.0 mL, 35.0 mmol) at -78 °C. The reaction mixture was warmed to rt and stirred at rt for 30 min. The reaction mixture was quenched by adding 300 mL of saturated Rochelle's salt water solution. The resulting mixture was stirred at rt for 1 h, which was then concentrated and the residue was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified with flash chromatography to give the desired product as white solid (2.1 g, 58%). LC-MS calculated for C₂₃H₂₂N₅O₃ (M+H)⁺: m/z = 416.1; found 416.2.

Step 2: 3-(2-(Bromomethyl)-4-(2,4-dimethoxybenzylamino)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




3-(4-(2,4-Dimethoxybenzylamino)-2-(hydroxymethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (3.0 g, 7.22 mmol) was dissolved in DCM (200 mL), and PBr₃ (1.4 mL, 14.44 mmol) was added. The mixture was stirred at rt for 5 h. After completion, the reaction was quenched by adding sat. NaHCO₃, the mixture was then extracted

with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as white solid (2.4 g, 69%). LC-MS calculated for C₂₃H₂₁BrN₅O₂ (M+H)⁺: m/z = 478.1; found 478.1.

5

Step 3: 3-(4-(2,4-Dimethoxybenzylamino)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(2-(bromomethyl)-4-(2,4-dimethoxybenzylamino)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (700 mg, 1.46 mmol), CuI (55.7 mg, 0.293 mmol), CsF (445 mg, 2.93 mmol), tetrakis(triphenylphosphine)palladium(0) (169 mg, 0.146 mmol), and 2-(tributylstannyl)pyridine (646 mg, 1.756 mmol) in 1,4-dioxane (2 mL) was heated at 140 °C for 1 h in a microwave reactor. The reaction was diluted with water and extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as white solid (428 mg, 62%). LC-MS calculated for C₂₈H₂₅N₆O₂ (M+H)⁺: m/z = 477.2; found 477.2.

Step 4: 3-(4-Amino-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile


A reaction vial was charged with 3-(4-(2,4-dimethoxybenzylamino)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (428 mg, 0.9 mmol) and TFA (1 mL). The mixture was heated at 70 °C for 20 min. The mixture was diluted with water and quenched with sat. NaHCO₃. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired

product as a light yellow solid (260 mg, 89%). LC-MS calculated for $C_{19}H_{15}N_6$ ($M+H$)⁺: m/z = 327.1; found 327.2.

**Example 17. 3-(4-Amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-
a]pyrazin-6-yl)benzonitrile**



Step 1: 3-(4-Amino-7-bromo-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of 3-(4-amino-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 16, Step 4; 260 mg, 0.8 mmol) in DMF (2 mL) was added a DMF (0.5 mL) solution of *N*-bromosuccinimide (122 mg, 0.68 mmol) dropwise at 0 °C. The resulting mixture was stirred at 0 °C for 10 min. The reaction mixture was diluted with water. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na_2SO_4 , filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as a light yellow oil (202 mg, 62%). LC-MS calculated for $C_{19}H_{14}BrN_6$ ($M+H$)⁺: m/z = 405.0; found 405.1.


Step 2. 3-(4-Amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile

A mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (15 mg, 0.037 mmol), CuI (1.4 mg, 0.007 mmol), CsF (11 mg, 0.074 mmol), tetrakis(triphenylphosphine)palladium(0) (4.2 mg, 0.004 mmol), and 4-(tributylstannyl)pyrimidine (16.4 mg, 0.044 mmol) in 1,4-dioxane was heated at 140 °C for 1 h in a microwave reactor. The reaction mixture was concentrated under

vacuum and the resulting residue was dissolved in methanol, added a few drops of TFA, and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{23}H_{17}N_8$ ($M+H$)⁺: $m/z = 405.2$ found 405.2.


5

Example 18. 3-(4-Amino-7-(1-ethyl-1*H*-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




A mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)pyrazolo[1,5-
 10 *a*]pyrazin-6-yl)benzonitrile (Example 17, Step 1; 15 mg, 0.037 mmol), 1-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole (9.69 mg, 0.044 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (2.86 mg, 3.63 μ mol) and Cs_2CO_3 (23.2 mg, 0.071 mmol) in 1,4-dioxane (1 mL)/water (0.200 mL) was stirred at 90 °C for 1 h. The reaction
 15 mixture was concentrated under vacuum and the resulting residue was dissolved in methanol and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{24}H_{21}N_8$ ($M+H$)⁺: $m/z = 421.2$; found 421.2.

20 **Example 19. 4-(4-Amino-6-(3-cyanophenyl)-2-(2-fluorobenzyl)pyrazolo[1,5-*a*]pyrazin-7-yl)-5,6-dihydropyridine-1(2*H*)-carboxamide**




Step 1. 3-(4-(2,4-Dimethoxybenzylamino)-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(2-(bromomethyl)-4-(2,4-dimethoxybenzylamino)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 16, Step 2; 200 mg, 0.42 mmol), (2-fluorophenyl)boronic acid (70.2 mg, 0.502 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (32.9 mg, 0.042 mmol) and Cs₂CO₃ (272 mg, 0.836 mmol) in 1,4-dioxane (2 mL)/water (0.2 mL) was stirred at 90 °C for 3 h. The reaction was diluted by water and extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as white solid (157 mg, 76%). LC-MS calculated for C₂₉H₂₅FN₅O₂ (M+H)⁺: m/z = 494.2; found 494.1.


Step 2. 3-(4-Amino-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a reaction vial was charged with 3-(4-(2,4-dimethoxybenzylamino)-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (157 mg, 0.32 mmol), TFA (1 mL). The mixture was heated at 70 °C for 20 min. The mixture was diluted with water, and quenched with sat. NaHCO₃. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as white solid (96 mg, 87%). LC-MS calculated for C₂₀H₁₅FN₅ (M+H)⁺: m/z = 344.1; found 344.2.

25

Step 3. 3-(4-Amino-7-bromo-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile and 3-(4-amino-3-bromo-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




5 To a solution of 3-(4-amino-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (96 mg, 0.28 mmol) in DMF (1 mL) was added a DMF (0.5 mL) solution of *N*-bromosuccinimide (39.8 mg, 0.22 mmol) dropwise at 0 °C. The resulting mixture was stirred at 0 °C for 10 min. The reaction mixture was diluted with water and extracted with DCM. The organic phase was washed with brine, dried
10 over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give two regioisomers (101 mg, 0.24 mmol 86%), which are used directly in the next step. LC-MS calculated for C₂₀H₁₄FBrN₅ (M+H)⁺: m/z = 422.0; found 422.1.

15 Step 4. 4-(4-Amino-6-(3-cyanophenyl)-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-7-yl)-5,6-dihydropyridine-1(2H)-carboxamide


A mixture of 3-(4-amino-7-bromo-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile, 3-(4-amino-3-bromo-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (15 mg, 0.037 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-
20 3,6-dihydropyridine-1(2H)-carboxamide (13.4 mg, 0.053 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (2.86 mg, 3.63 μmol) and Cs₂CO₃ (23.2 mg, 0.071 mmol) in 1,4-dioxane (0.6 mL)/water (0.200 mL) was stirred at 90 °C for 1 h. The reaction was concentrated under vacuum and the residue was dissolved in methanol and
25 purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give two compounds. Compound with shorter retention time was assigned as title compound as a TFA salt. LC-MS calculated for C₂₆H₂₃FN₇O (M+H)⁺: m/z = 468.2; found 468.2.

Example 20. 4-Amino-6-(3-cyanophenyl)-2-(2-fluorobenzyl)pyrazolo[1,5-*a*]pyrazine-7-carbonitrile




A mixture of 3-(4-amino-7-bromo-2-(2-fluorobenzyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, 3-(4-amino-3-bromo-2-(2-fluorobenzyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (15 mg, 0.037 mmol) (Example 19, Step 3), zinc cyanide (8.3 mg, 0.071 mmol), and tBuXPhos-Pd-G3 (2.8 mg, 3.6 μ mol) in 1,4-dioxane (1 mL)/water (1 mL) was stirred at 100 °C for 4 h. The reaction mixture was concentrated under vacuum and the resulting residue was dissolved in methanol and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give two compounds. Compound with shorter retention time was assigned as title compound as a TFA salt. LC-MS calculated for $C_{21}H_{14}FN_6$ ($M+H$)⁺: m/z = 369.1; found 369.2.


Example 21. 4-Amino-6-(3-cyanophenyl)-2-(2-fluorobenzyl)pyrazolo[1,5-*a*]pyrazine-3-carbonitrile



The title compound was prepared using same procedures as described for Example 20. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give two compounds. Compound with longer retention time was assigned as title compound as a TFA salt. LC-MS calculated for $C_{21}H_{14}FN_6$ ($M+H$)⁺: m/z = 369.1; found 369.2.

Example 22. 3-(4-Amino-7-bromo-2-((2-fluorophenyl)(hydroxy)methyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




Step 1. 3-(4-Amino-7-bromo-2-formylpyrazolo[1,5-a]pyrazin-6-yl)benzonitrile

To a solution of 3-(4-amino-7-bromo-2-(hydroxymethyl)pyrazolo[1,5-
a]pyrazin-6-yl)benzonitrile (Example 13, Step 1; 1.06 g, 3.1 mmol) in DCM (30 mL)
 5 was added Dess-Martin periodinane (1.44 g, 3.39 mmol). The resulting mixture was
 stirred at room temperature for 30 min. The reaction mixture was diluted with sat.
 NaHCO₃. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and
 concentrated. The residue was purified with flash chromatography to give the desired
 product as white solid (0.64 g, 61%). LC-MS calculated for C₁₄H₉BrN₅O (M+H)⁺:
 10 m/z = 342.0; found 342.0.

*Step 2. 3-(4-Amino-7-bromo-2-((2-fluorophenyl)(hydroxy)methyl)pyrazolo[1,5-
a]pyrazin-6-yl)benzonitrile*


To a THF (20 mL) solution of 1-fluoro-2-iodobenzene (0.31 mL, 2.7 mmol)
 15 was added isopropylmagnesium bromide (0.8 mL, 2.3 mmol) in THF dropwise at -20
 °C, and the solution was stirred for 1 h. Then to the solution was added a THF (2 mL)
 solution of 3-(4-amino-7-bromo-2-formylpyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile
 (228 mg, 0.67 mmol). The mixture was stirred for 12 h while warming to room
 temperature. After completion, the reaction was quenched by adding sat. NH₄Cl. The
 20 aqueous phase was extracted with DCM, and the organic phase was washed with
 brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with
 flash chromatography to give the desired product as white solid (0.18 g, 63%). LC-
 MS calculated for C₂₀H₁₄BrFN₅O (M+H)⁺: m/z = 438.0; found 438.1.

Example 23. 3-(4-Amino-2-((2-fluorophenyl)(hydroxy)methyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




A mixture of 3-(4-amino-7-bromo-2-((2-fluorophenyl)(hydroxy)methyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (Example 22; 16.0 mg, 0.037 mmol), CuI (1.4 mg, 0.007 mmol), CsF (11.0 mg, 0.074 mmol), tetrakis(triphenylphosphine)palladium(0) (4.2 mg, 0.004 mmol), and 4-(tributylstannyl)pyrimidine (16.4 mg, 0.044 mmol) in 1,4-dioxane was heated at 140 °C for 1 h in a microwave reactor. The reaction mixture was concentrated under vacuum and the resulting residue was dissolved in methanol, mixed with a few drops of TFA, and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₄H₁₇FN₇O (M+H)⁺: m/z = 438.1 found 438.2.

Example 24. 3-(4-Amino-2-(3,6-dihydro-2*H*-pyran-4-yl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



*Step 1. 3-(2-Chloro-4-oxo-4,5-dihydropyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*




20

To a solution of methyl 3-chloro-1*H*-pyrazole-5-carboxylate (901 mg, 5.61 mmol) and 3-(2-bromoacetyl)benzonitrile (1257 mg, 5.61 mmol) in acetone (253 mL) was added potassium carbonate (853 mg, 6.17 mmol). The mixture was stirred at rt for 12 h. The reaction mixture was then concentrated and the resulting residue was


taken up in water and DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was dissolved in acetic acid (30 mL), and ammonium acetate (4.7 g, 60.8 mmol) was added. The mixture was stirred at 110 °C for 36 h. After cooling to rt, the mixture was diluted with water and the precipitate
 5 was collected via filtration and washed with water to give the desired product as white solid. LC-MS calculated for C₁₃H₈ClN₄O (M+H)⁺: m/z = 271.0; found 271.1.

Step 2. 3-(2,4-Dichloropyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(2-chloro-4-oxo-4,5-dihydropyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (1.6 g, 5.9 mmol) and POCl₃ (2 mL, 20.1 mmol) was heated at 110 °C overnight. After cooling to rt, the mixture was added to a flask containing ice. The resulting precipitate was collected and washed with water to give the desired product as white solid. (1.18 g, 69%). LC-MS calculated for C₁₃H₇Cl₂N₄ (M+H)⁺: m/z =
 15 289.0; found 289.1.

Step 3. 3-(4-Amino-2-chloropyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




A microwave vial was charged with 3-(2,4-dichloropyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (1.4 g, 4.84 mmol), (2,4-dimethoxyphenyl)methanamine (0.90 g, 5.4 mmol), DIEA (1.304 mL, 7.47 mmol) and butan-1-ol (10 mL). The mixture was heated at 150 °C for 30 min in a microwave reactor. The mixture was diluted with water, and the resulting precipitate was collected via filtration. The solid was then treated with TFA (10 mL) and heated at 70 °C for 30 min. The reaction was then
 25 quenched with sat. NaHCO₃, and the resulting solid was collected via filtration and washed with water to give the desired product as a white solid (860 mg, 3.2 mmol, 67%). LC-MS calculated for C₁₃H₉ClN₅ (M+H)⁺: m/z = 270.0; found 270.0.

Step 4. 3-(4-Amino-7-bromo-2-chloropyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



To a solution of 3-(4-amino-2-chloropyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (860 mg, 3.2 mmol) in DMF (5 mL) was added NBS (570 mg, 3.2 mmol). The resulting mixture was stirred at room temperature for 30 min. The reaction mixture was diluted with water. The resulting precipitate was collected and washed with water to give the desired product as white solid (972 mg, 88%). LC-MS calculated for $C_{13}H_8ClBrN_5$ ($M+H$)⁺: m/z = 348.0; found 348.0.

Step 5. 3-(4-Amino-2-chloro-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




A mixture of 3-(4-amino-7-bromo-2-chloropyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (100 mg, 0.288 mmol), CuI (14 mg, 0.07 mmol), CsF (110 mg, 0.74 mmol), tetrakis(triphenylphosphine)palladium(0) (42 mg, 0.04 mmol), and 4-(tributylstannyl)pyrimidine (118 mg, 0.34 mmol) in 1,4-dioxane (3 mL) was heated at 140 °C for 1 h in a microwave reactor. The reaction mixture was concentrated under vacuum and the resulting residue was purified using flash chromatography to give the desired product as white solid (52 mg, 52%). LC-MS calculated for $C_{17}H_{11}ClN_7$ ($M+H$)⁺: m/z = 348.1; found 348.1.

Step 6. 3-(4-Amino-2-(3,6-dihydro-2H-pyran-4-yl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile


A mixture of 3-(4-amino-2-chloro-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (13 mg, 0.037 mmol), 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (8.4 mg, 0.04 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1)

(2.86 mg, 3.63 μ mol), and Cs_2CO_3 (23.2 mg, 0.071 mmol) in 1,4-dioxane (1 mL)/water (0.200 mL) was stirred at 90 °C for 1 h. The reaction mixture was concentrated under vacuum and the resulting residue was dissolved in methanol and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $\text{C}_{22}\text{H}_{18}\text{N}_7\text{O}$ ($\text{M}+\text{H}$)⁺: m/z = 396.2; found 396.1.

Example 25. 3-(4-Amino-2-(phenylamino)-7-(pyridin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile



Step 1: 3-(4-Amino-2-chloro-7-(pyridin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(4-amino-7-bromo-2-chloropyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 24 Step 4; 129 mg, 0.370 mmol), pyridin-4-ylboronic acid (45.5 mg, 0.370 mmol), $\text{PdCl}_2(\text{dppf})\text{-CH}_2\text{Cl}_2$ adduct (30.2 mg, 0.037 mmol), sodium carbonate (78 mg, 0.740 mmol) in 1,4-dioxane (1682 μ L), and water (168 μ L) was purged with N_2 and heated at 95 °C for 5 h. The mixture was then concentrated and purified by silica gel chromatograph eluting with 0 to 13% MeOH in DCM to afford the desired product. LCMS calculated for $\text{C}_{18}\text{H}_{12}\text{ClN}_6$ ($\text{M}+\text{H}$)⁺: 347.1; found 347.1.


Step 2: 3-(4-Amino-2-(phenylamino)-7-(pyridin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile

A mixture of 3-(4-amino-2-chloro-7-(pyridin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.029 mmol), aniline (8.06 mg, 0.087 mmol), cesium carbonate (18.79 mg, 0.058 mmol), chloro[(4,5-bis(diphenylphosphino)-9,9-dimethylxanthene)-2-(2'-amino-1,1'-biphenyl)]palladium(II) (2.56 mg, 2.88 μ mol)

(XantPhos Pd G2) in 1,4-dioxane (144 μ l) was purged with N₂ and heated at 95 °C for 1 h. The mixture was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₄H₁₈N₇ (M+H)⁺: 404.2; found 404.1.


5

Example 26. 3-(4-Amino-2-(2-(4-methylpiperazin-1-yl)pyridin-4-yl)-7-(pyridin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




Chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-
 10 amino-1,1'-biphenyl)]palladium(II) (11.34 mg, 0.014 mmol) was added to a mixture
 of 3-(4-amino-2-chloro-7-(pyridin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile
 (Example 25, Step 1; 50 mg, 0.144 mmol), 1-methyl-4-(4-(4,4,5,5-tetramethyl-1,3,2-
 dioxaborolan-2-yl)pyridin-2-yl)piperazine (52.5 mg, 0.173 mmol), sodium carbonate
 (15.28 mg, 0.144 mmol) in 1,4-dioxane (655 μ l) and water (65.5 μ l). The mixture was
 15 purged with N₂ and heated at 90 °C for 2 h. The resulting mixture was concentrated
 and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the
 product as TFA salt. LCMS calculated for C₂₈H₂₆N₉ (M+H)⁺: 488.2; found 488.1.

**Example 27. 3-(8-Amino-2-(pyridin-2-ylmethyl)-5-(pyrimidin-4-yl)-
 20 [1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile**




Step 1: Methyl 3-bromo-1-(2-(3-cyanophenyl)-2-oxoethyl)-1H-1,2,4-triazole-5-carboxylate




To a solution of methyl 3-bromo-1H-1,2,4-triazole-5-carboxylate (5.0 g, 24.3 mmol), 3-(2-bromoacetyl)benzonitrile (5.44 g, 24.3 mmol) in DMF (100 mL) was added potassium carbonate (3.35 g, 24.3 mmol). The reaction mixture was stirred at ambient temperature for 2 h. The reaction mixture was then diluted with water and DCM. The organic layer was separated, washed with brine, dried over Na₂SO₄, filtered and concentrated. The resulting residue was purified via flash chromatography to give the desired product as a white solid (5.2 g, 61%). LC-MS calculated for C₁₃H₁₀BrN₄O₃ (M+H)⁺: m/z = 349.0; found 349.0.

Step 2: 3-(2-Bromo-8-oxo-7,8-dihydro-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Methyl 3-bromo-1-(2-(3-cyanophenyl)-2-oxoethyl)-1H-1,2,4-triazole-5-carboxylate (10.5 g, 30.1 mmol) was dissolved in acetic acid (100 mL), and ammonium acetate (23.18 g, 301 mmol) was added. The mixture was stirred at 110 °C for 12 h. After cooling to room temperature, the reaction mixture was diluted with water. The resulting precipitate was collected via filtration, washed with water, and dried under vacuum to afford the product (8.4 g, 88%). LC-MS calculated for C₁₂H₇BrN₅O (M+H)⁺: m/z = 316.0; found 316.0.

Step 3: 3-(2-Bromo-8-chloro-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(2-bromo-8-oxo-7,8-dihydro-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (8.4 g, 26.6 mmol) and POCl₃ (49.5 mL, 531 mmol) was stirred at 110 °C overnight. After cooling to room temperature, the reaction mixture was slowly added to a flask containing ice and sodium bicarbonate. The resulting precipitate was collected, washed with water, and dried to afford the product (8.8 g, 99%). LC-MS calculated for C₁₂H₆BrClN₅ (M+H)⁺: m/z = 333.9; found 334.0.

*Step 4. 3-(8-(Bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*



A mixture of 3-(2-bromo-8-chloro-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (8.99 g, 26.9 mmol), bis(4-methoxybenzyl)amine (10.37 g, 40.3 mmol), and DIPEA (9.4 mL, 53.7 mmol) in DMF (134 mL) was stirred at 85 °C overnight. The reaction mixture was cooled to room temperature, and diluted with water. The resulting precipitate was collected via filtration, and dried to afford the product (14.1 g, 94%). LC-MS calculated for C₂₈H₂₄BrN₆O₂ (M+H)⁺: m/z = 555.1; found 555.1.


*Step 5: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*



To a solution of 2-methylpyridine (0.050 g, 0.540 mmol) in THF (0.5 mL) was added 2.5 M *n*-butyllithium (0.216 mL, 0.540 mmol) at -78° C. The resulting solution was stirred at the same temperature for 1 h, before 1.9 M zinc chloride in 2-methyltetrahydrofuran (0.284 mL, 0.540 mmol) was added, and the resulting mixture was stirred at room temperature for 10 min.


A microwave vial charge with 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (0.15 g, 0.270 mmol), palladium acetate (1.1 mg, 4.7 μmol), and 2'-(dicyclohexylphosphino)-*N,N,N',N'*-tetramethylbiphenyl-2,6-diamine (4.1 mg, 9.5 μmol) was evacuated under high vacuum and backfilled with nitrogen. THF (2.0 mL) and toluene (0.5 mL) was then added to the reaction vial. The mixture was cooled to 0 °C and the zinc reagent prepared from previous step was added slowly via a syringe. The reaction mixture was then stirred at 60 °C overnight, cooled to room temperature, and partitioned between ethylacetate and saturated NH₄Cl solution. The layers were separated and the aqueous layer was extracted with ethylacetate. The combined organic layers were washed with water and brine, dried over MgSO₄, and concentrated. The resulting residue was purified via flash chromatography to afford the product (0.11 g, 71%). LC-MS calculated for C₃₄H₃₀N₇O₂ (M+H)⁺: *m/z* = 568.2; found 568.3.

*Step 6. 3-(8-Amino-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (110 mg, 0.194 mmol) and TFA (746 μL, 9.69 mmol) was stirred at 80 °C for 30 min, cooled to room temperature, and concentrated. The resulting residue was purified via prep-LCMS (pH 2) to give the product as a white solid (TFA salt) (57 mg, 90%). LC-MS calculated for C₁₈H₁₄N₇ (M+H)⁺: *m/z* = 328.1; found 328.1.

Step 7. 3-(8-Amino-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



To a solution of 3-(8-amino-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (TFA salt) (35 mg, 0.079 mmol) in DMF (0.5 mL)/DCM (0.5 mL) was added NBS (14.1 mg, 0.079 mmol). The reaction mixture was then stirred at room temperature for 1 h, and concentrated to afford the crude product, which was used in the next step without further purification. LC-MS calculated for C₁₈H₁₃BrN₇ (M+H)⁺: m/z = 406.0; found 406.0.


10

Step 8. 3-(8-Amino-2-(pyridin-2-ylmethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile

A mixture of 3-(8-amino-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (15 mg, 0.037 mmol), 4-(tributylstannyl)pyrimidine (20 mg, 0.055 mmol), and copper(I) chloride (4.4 mg, 0.044 mmol), lithium chloride (1.9 mg, 0.044 mmol) and tetrakis(triphenylphosphine)palladium(0) (4.3 mg, 3.7 μmol) in THF (1 mL) was purged with N₂, and stirred at 90 °C for 2 h. The reaction mixture was then cooled to room temperature, diluted with methanol, and purified via prep-LCMS (pH 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₂H₁₆N₉ (M+H)⁺: m/z = 406.2; found 406.2.

20

Example 28. 3-(8-Amino-5-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-Amino-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (Example 27, Step 7; 10 mg, 0.025 mmol), 1-methyl-5-

25

(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2(1*H*)-one (10 mg, 0.042 mmol), cesium carbonate (37.6 mg, 0.116 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (2.26 mg, 2.88 μ mol) (XPhos Pd G2) in 1,4-dioxane (500 μ l) and water (100 μ l) was purged with N₂ and heated at 95 °C for 1 h. The mixture was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₄H₁₉N₈O (M+H)⁺: 435.2; found 435.2.


Example 29. 3-(8-Amino-5-(1-methyl-6-oxo-1,6-dihydropyridazin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile"




A mixture of 6-chloro-2-methylpyridazin-3(2*H*)-one (30 mg, 0.21 mmol), bis(pinacolato)diboron (53 mg, 0.21 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (15.7 mg, 0.02 mmol) (XPhos Pd G2) and potassium acetate (61.7 mg, 0.63 mmol) in 1,4-dioxane (1 mL) was stirred at 100 °C for 1 h. 3-(8-Amino-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (Example 27, Step 7; 10 mg, 0.025 mmol), cesium carbonate (37.6 mg, 0.116 mmol) and water (0.2 mL) were then added to the reaction mixture. The resulting mixture was heated at 90 °C for 1h. The mixture was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₃H₁₈N₉O (M+H)⁺: 436.2; found 436.2.

¹H NMR (500 MHz, DMSO) δ 8.66 – 8.62 (d, *J* = 5.1 Hz, 1H), 8.09 – 8.02 (d, *J* = 1.8 Hz, 1H), 7.88 – 7.85 (t, *J* = 1.8 Hz, 1H), 7.85 – 7.81 (m, 3H), 7.78 – 7.72 (d, *J* = 9.6 Hz, 1H), 7.66 – 7.51 (m, 4H), 7.10 – 7.06 (d, *J* = 9.6 Hz, 1H), 4.59 – 4.48 (s, 2H), 3.53 – 3.43 (s, 3H).

Example 30. (*S*)-1-(2-((8-Amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)-3-fluorobenzyl)pyrrolidine-3-carboxylic acid




- 5 *Step 1: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*




- A microwave vial was charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (Example 27, Step 4; 0.15 g, 0.270 mmol), palladium acetate (1.1 mg, 4.7 μ mol), and 2'-(dicyclohexylphosphino)-*N,N,N',N'*-tetramethylbiphenyl-2,6-diamine (4.1 mg, 9.5 μ mol), the vial was then evacuated under high vacuum and backfilled with nitrogen. THF (2.0 mL) was then added to the reaction vial. The mixture was cooled to 0 °C and the (2-chloro-6-fluorobenzyl)zinc(II) chloride (0.5 M THF solution, 1.08 mL) was added slowly via a syringe. The reaction mixture was then stirred at 60 °C for 2h, cooled to room temperature, and partitioned between ethylacetate and saturated NH_4Cl solution. The layers were separated and the aqueous layer was extracted with ethylacetate. The combined organic layers were washed with water and brine, dried over MgSO_4 , and concentrated. The resulting residue was purified via flash chromatography to afford the product. LC-MS calculated for $\text{C}_{35}\text{H}_{29}\text{ClFN}_6\text{O}_2$ ($\text{M}+\text{H}$) $^+$: m/z = 619.2; found 619.3.

Step 2: 3-(8-Amino-5-bromo-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (100 mg, 0.161 mmol) and TFA (746 μ L, 9.69 mmol) was stirred at 100 °C for 10 min, cooled to room temperature, and concentrated. The resulting residue was dissolved in DMF, and a DMF (0.5 mL) solution of *N*-bromosuccinimide (28 mg, 0.161 mmol) was added dropwise at 0 °C. The resulting mixture was stirred at 0 °C for 10 min. The reaction mixture was diluted with sat. NaHCO₃. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for C₁₉H₁₂BrClFN₆ (M+H)⁺: m/z = 457.0; found 457.1.

Step 3: 3-(8-Amino-2-(2-chloro-6-fluorobenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-amino-5-bromo-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (51 mg, 0.11 mmol), CuI (4.2 mg, 0.021 mmol), CsF (33 mg, 0.22 mmol), tetrakis(triphenylphosphine)palladium(0) (12 mg, 0.012 mmol), and 4-(tributylstannyl)pyrimidine (49 mg, 0.13 mmol) in 1,4-dioxane (2 mL) was heated at 140 °C for 1 h in a microwave reactor. The reaction mixture was concentrated under vacuum and the resulting residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for C₂₃H₁₅ClFN₈ (M+H)⁺: m/z = 457.1 found 457.1.

Step 4: 3-(8-Amino-2-(2-fluoro-6-vinylbenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-amino-2-(2-chloro-6-fluorobenzyl)-5-(pyrimidin-4-yl)-
 5 [1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (50 mg, 0.11 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (34 mg, 0.22 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (8.5 mg, 10.8 μ mol) and K_3PO_4 (47 mg, 0.22 mmol) in 1,4-dioxane (2 mL)/water (0.4 mL) was stirred at 110 °C for 1 h. The reaction mixture was concentrated under
 10 vacuum and the resulting residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for $C_{25}H_{18}FN_8$ (M+H)⁺: m/z = 449.2; found 449.1.

Step 5: 3-(8-Amino-2-(2-fluoro-6-formylbenzyl)-5-(pyrimidin-4-yl)-
 15 [1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-amino-2-(2-fluoro-6-vinylbenzyl)-5-(pyrimidin-4-yl)-
 [1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (40 mg, 0.089 mmol) in THF (1 mL)
 and water (1 mL) was added 0.157 M osmium tetroxide in water (0.02 mmol). After
 20 2 min, sodium metaperiodate (86 mg, 0.4 mmol) was added. The reaction mixture was heated at 60 °C for 1 h before quenched with sat. $Na_2S_2O_3$. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na_2SO_4 , filtered, and concentrated to afford the product as a light yellow oil. LC-MS calculated for $C_{24}H_{16}FN_8O$ (M+H)⁺: m/z = 451.1; found 451.1

25

Step 6: (S)-1-(2-((8-amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)pyrrolidine-3-carboxylic acid

To a solution of 3-(8-amino-2-(2-fluoro-6-formylbenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.02 mmol) in DCM (0.5 mL) was added (S)-pyrrolidine-3-carboxylic acid (4.6 mg, 0.04 mmol) then acetic acid (4 μ L, 0.08 mmol). After 1h, sodium triacetoxymethylborohydride (8.5 mg, 0.04 mmol) was added to the reaction. The reaction was stirred overnight and the mixture was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for $C_{29}H_{25}FN_9O_2$ (M+H)⁺: 550.2 found 550.2.


Example 31. 1-(2-((8-Amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)azetidine-3-carboxylic acid




The title compound was prepared using similar procedures as described for Example 30, with azetidine-3-carboxylic acid replacing (S)-pyrrolidine-3-carboxylic acid in Step 6. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{28}H_{23}FN_9O_2$ (M+H)⁺: 536.2 found 536.2.

Example 32 and Example 33. 3-(2-((1*H*-Pyrrolo[2,3-*b*]pyridin-1-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile and 3-(2-((7*H*-pyrrolo[2,3-*b*]pyridin-7-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile

5




*Step 1: 3-(8-(Bis(4-methoxybenzyl)amino)-2-vinyl-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*



10 A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (362 mg, 0.66 mmol) (from Example 27, Step 4), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (204 mg, 1.32 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (51 mg, 64.8 μ mol) and K_3PO_4 (282 mg, 1.32 mmol) in 1,4-dioxane (5 mL)/water (1
15 mL) was stirred at 80 °C for 1 h. The reaction mixture was concentrated under vacuum and the resulting residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for $C_{30}H_{27}N_6O_2$ ($M+H$)⁺: m/z = 503.2; found 503.2.


Step 2: 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (241 mg, 0.48 mmol) in DCM (5 mL) was added NBS (84.6 mg, 0.48 mmol). The reaction mixture was then stirred at room temperature for 1 h, and concentrated to afford the crude product, which was used in the next step without further purification. LC-MS calculated for $C_{30}H_{26}BrN_6O_2$ ($M+H$)⁺: m/z = 581.1; found 581.1.

10

Step 3: 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-(hydroxymethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (203 mg, 0.35 mmol) in THF (4 mL) and water (4 mL) was added 0.157 M osmium tetroxide in water (0.1 mmol). After 2 min, sodium metaperiodate (430 mg, 2 mmol) was added. The reaction mixture was heated at 60 °C for 1 h before quenched with sat. $Na_2S_2O_3$. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na_2SO_4 , filtered, and concentrated. The crude material was dissolved in DCM (1 mL) and MeOH (3 mL), which was cooled to -78 °C before $NaBH_4$ (13 mg, 0.35 mmol) was added. The resulting mixture was warmed to 0 °C and stirred at this temperature for 10 min.

20

Water (10 mL) was then added and the mixture was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated to afford the product as a light yellow oil. LC-MS calculated for C₂₉H₂₆BrN₆O₃ (M+H)⁺: m/z = 585.1; found 585.1

5


Step 4: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(hydroxymethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(hydroxymethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (257 mg, 0.44 mmol), CuI (17 mg, 0.084 mmol), CsF (132 mg, 0.88 mmol), tetrakis(triphenylphosphine)palladium(0) (48 mg, 0.048 mmol), and 4-(tributylstannyl)pyrimidine (196 mg, 0.52 mmol) in 1,4-dioxane (4 mL) was heated at 140 °C for 1 h in a microwave reactor. The reaction mixture was concentrated under vacuum and the resulting residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for C₃₃H₂₉N₈O₃ (M+H)⁺: m/z = 585.2 found 585.2.

15

Step 5: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(bromomethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




20

3-(8-(Bis(4-methoxybenzyl)amino)-2-(hydroxymethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (234 mg, 0.4 mmol) was dissolved in THF (5 mL), and PBr_3 (0.077 mL, 0.8 mmol) was added. The mixture was stirred at 60 °C for 4 h. After completion, the reaction was quenched by adding sat. NaHCO_3 , the mixture was then extracted with EtOAc. The organic phase was washed with brine, dried over Na_2SO_4 , filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as white solid. LC-MS calculated for $\text{C}_{33}\text{H}_{28}\text{BrN}_8\text{O}_2$ ($\text{M}+\text{H}$)⁺: m/z = 647.1; found 647.2.


Step 6: 3-(2-((1H-pyrrolo[2,3-b]pyridin-1-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile and 3-(2-((7H-pyrrolo[2,3-b]pyridin-7-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

3-(8-(Bis(4-methoxybenzyl)amino)-2-(bromomethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (20 mg, 0.031 mmol) was dissolved in MeCN (2 mL), K_2CO_3 (8.6 mg, 0.062 mmol) and 1H-pyrrolo[2,3-b]pyridine (7.3 mg, 0.062) were added. The mixture was stirred at 70 °C for 6 h. After completion, the solvent was removed under vacuum and 1 mL of TFA was added to the residue. The mixture was heated at 100 °C for 10 min. The reaction mixture was concentrated under vacuum and the resulting residue was dissolved in methanol, and purified with prep-LCMS (pH 2, acetonitrile/water+TFA). The first peak was isolated as Example 32 as a TFA salt. LC-MS calculated for $\text{C}_{24}\text{H}_{17}\text{N}_{10}$ ($\text{M}+\text{H}$)⁺: m/z = 445.2 found 445.1. The second peak was isolated as Example 33, also as a TFA salt. LC-MS calculated for $\text{C}_{24}\text{H}_{17}\text{N}_{10}$ ($\text{M}+\text{H}$)⁺: m/z = 445.2 found 445.1.

Example 34. 3-(4-Amino-7-(4-(1-hydroxyethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




Step 1: 4-(1-((tert-Butyldimethylsilyl)oxy)ethyl)-2-methyloxazole



In a flame dried round-bottomed flask equipped with a magnetic stir bar, a solution of 1-(2-methyloxazol-4-yl)ethan-1-ol (1 g, 7.87 mmol) in DCM (10 mL) was treated at rt with tert-butylchlorodimethylsilane (1.3 g, 7.88 mmol) followed by imidazole (0.54 g, 7.87 mmol) and the resulting suspension was stirred for 1 h at rt. Water was added. The aq. layer was extracted with DCM and the combined org. layers were dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified with flash chromatography to give the desired product. LC-MS calculated for C₁₂H₂₄NO₂Si (M+H)⁺: m/z = 242.2; found 242.2.

Step 2: 4-(1-((tert-Butyldimethylsilyl)oxy)ethyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole




In a flame dried round-bottomed flask equipped with a magnetic stir bar, was charged with (1,5-cyclooctadiene)(methoxy)iridium(I) dimer (30 mg, 0.045 mmol), 4,4,5,5-tetramethyl-1,3,2-dioxaborolane (190 mg, 1.5 mmol), and pentane (2.0 mL). The mixture was stirred at rt for 10 min. Then 4,4'-di-tert-butyl-2,2'-dipyridyl (24 mg, 0.09 mmol) was added to this mixture and reaction stirred for additional 20 min. 4-(1-((tert-butyldimethylsilyl)oxy)ethyl)-2-methyloxazole (300 mg, 1.24 mmol) dissolved in Et₂O (2 mL) was added to the active catalyst mixture. The reaction was stirred at room temperature until completion. Solvent was removed under reduced pressure, and the crude material was purified with flash chromatography to give the desired product. LC-MS calculated for the corresponding boronic acid C₁₂H₂₅BNO₄Si (M+H)⁺: m/z = 286.2; found 286.2.

Step 3: 3-(4-Amino-7-(4-(1-hydroxyethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile


A mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (Example 17, Step 1; 15 mg, 0.037 mmol), 4-(1-((tert-butyl)dimethylsilyl)oxy)ethyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (29 mg, 0.08 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (2.86 mg, 3.63 μ mol) and Cs₂CO₃ (23.2 mg, 0.071 mmol) in 1,4-dioxane (1 mL)/water (0.200 mL) was stirred at 90 °C for 1 h. The reaction mixture was cooled down to room temperature before 6 N HCl (1 mL) was added and resulting reaction mixture was stirred at 60 °C for 30 min. The reaction was then diluted with methanol and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as a racemic material. Chiral separation was then conducted with chiral HPLC using AM-1 column and 30% EtOH in hexanes (20ml/min) solvent system. Peak 2 was collected as the desired product. LC-MS calculated for C₂₅H₂₂N₇O₂ (M+H)⁺: m/z = 452.2; found 452.2.

15

Example 35. 3-(4-Amino-7-(4-(2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



20 *Step 1: 2,2-Difluoro-1-(2-methyloxazol-4-yl)ethan-1-ol*




To a solution of (bromodifluoromethyl)trimethylsilane (2.4 mL, 15.07 mmol) in dry acetonitrile (10.1 mL) was added 3-bromo-4-methylbenzaldehyde (2 g, 10.05 mmol) and triphenylphosphine (3.16 g, 12.06 mmol). The resulting suspension was cooled to 0 °C and 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (2.4 mL, 20.10 mmol) [DMPU] was added dropwise. The cooling bath was removed and the reaction mixture was warmed to rt and stirred for 1h. With the reaction flask in a rt

25


water bath, aqueous 3 M KOH (1.7 g, 30.1 mmol) was added dropwise via addition funnel. The bath was removed, and the reaction mixture was stirred rapidly for 1.5 h. The reaction was then neutralized by aqueous 2 M HCl (10.1 ml, 20.10 mmol), and the mixture was extracted with MTBE (2 x 75 ml). The combined organic layers were washed with Sat. NaCl solution, dried over Na₂SO₄, filtered, and concentrated to afford the crude product as a brown oil, which was purified with flash chromatography to give the desired product. LC-MS calculated for C₆H₈F₂NO₂ (M+H)⁺: m/z = 164.0; found 164.0.

Step 2: 4-(1-((tert-butyl)dimethylsilyl)oxy)-2,2-difluoroethyl)-2-methyloxazole



In a flame dried round-bottomed flask equipped with a magnetic stir bar, a solution of 2,2-difluoro-1-(2-methyloxazol-4-yl)ethan-1-ol (1.29 g, 7.87 mmol) in DCM (10 mL) was treated at rt with tert-butylchlorodimethylsilane (1.3 g, 7.88 mmol) followed by imidazole (0.54 g, 7.87 mmol) and the resulting suspension was stirred for 1 h at rt. Water was added. The aq. layer was extracted with and the combined org. layers were dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified with flash chromatography to give the desired product. LC-MS calculated for C₁₂H₂₂F₂NO₂Si (M+H)⁺: m/z = 278.1; found 278.1.

Step 3: 4-(1-((tert-Butyl)dimethylsilyl)oxy)-2,2-difluoroethyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole




In a flame dried round-bottomed flask equipped with a magnetic stir bar, was charged with (1,5-cyclooctadiene)(methoxy)iridium(I) dimer (60 mg, 0.09 mmol), 4,4,5,5-tetramethyl-1,3,2-dioxaborolane (834 mg, 3.0 mmol), and pentane (4.0 mL).


The mixture was stirred at room temperature for 10 min. Then 4,4'-di-tert-butyl-2,2'-dipyridyl (48 mg, 0.18 mmol) was added to this mixture and reaction stirred for additional 20 min. 4-(1-((tert-butyldimethylsilyl)oxy)ethyl)-2-methyloxazole (600 mg, 2.4 mmol) dissolved in Et₂O (4 mL) was added to the active catalyst mixture. The reaction was stirred at room temperature until completion. Solvent was removed under reduced pressure, and the crude material was purified with flash chromatography to give the desired product. LC-MS calculated for the corresponding boronic acid C₁₂H₂₂BF₂NO₄Si (M+H)⁺: m/z = 322.2; found 322.1.

10 *Step 4: 3-(4-Amino-7-(4-(2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile*

A mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 17, Step 1; 15 mg, 0.037 mmol), 4-(1-((tert-butyldimethylsilyl)oxy)-2,2-difluoroethyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (32 mg, 0.08 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (2.86 mg, 3.63 μmol) and Cs₂CO₃ (23.2 mg, 0.071 mmol) in 1,4-dioxane (1 mL)/water (0.200 mL) was stirred at 90 °C for 1 h. The reaction mixture was cooled down to room temperature before 6 N HCl (1 mL) was added and resulting reaction mixture was stirred at 60 °C for 30 min. The reaction was then diluted with methanol and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as a racemic material. Chiral separation was then conducted with chiral HPLC using Phenomenex LUX Amylose-1 column and 45% EtOH in hexanes (20ml/min) solvent system. Peak 1 was collected as the desired product. LC-MS
25 calculated for C₂₅H₂₀F₂N₇O₂ (M+H)⁺: m/z = 488.2; found 488.2.

Example 36. 3-(4-Amino-7-(1-ethyl-1H-1,2,3-triazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile



Step 1: 1-Ethyl-5-(trimethylstannyl)-1H-1,2,3-triazole

To a stirred solution of 1-ethyl-1H-1,2,3-triazole (0.58 g, 6.0 mmol) in anhydrous tetrahydrofuran (50 mL) at -78 °C under an atmosphere of nitrogen was added n-butyllithium (2.5M solution in hexanes, 2.6 mL, 6.6 mmol) dropwise over ten minutes. On complete addition the reaction was allowed to warm to -30 °C and stirred for 2 h. A solution of chlorotrimethylstannane (1.3 g, 6.6 mmol) in tetrahydrofuran (2 mL) was added dropwise over 10 minutes then the reaction mixture was allowed to warm to room temperature over 2 h. The reaction was quenched by the addition of saturated ammonium chloride solution (5 mL) then diluted with water (20 mL). The solvent was evaporated in vacuo and the aqueous phase extracted with ethyl acetate (2×30 mL). The combined organic layer was dried over sodium sulfate, filtered and concentrated to afford the desired product as a pale oil. LC-MS calculated for C₇H₁₆N₃Sn (M+H)⁺: m/z = 262.0; found 262.0.


15

Step 2: 3-(4-Amino-7-(1-ethyl-1H-1,2,3-triazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile

A mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 17, Step 1; 15 mg, 0.037 mmol), CuI (17 mg, 0.084 mmol), CsF (132 mg, 0.88 mmol), tetrakis(triphenylphosphine)palladium(0) (48 mg, 0.048 mmol), and 1-ethyl-5-(trimethylstannyl)-1H-1,2,3-triazole (135 mg, 0.52 mmol) in 1,4-dioxane (4 mL) was heated at 140 °C for 1 h in a microwave reactor. The reaction mixture was concentrated under vacuum and the resulting residue was dissolved in methanol, and purified with preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₃H₂₀N₉ (M+H)⁺: m/z = 422.2 found 422.2.


25

Example 37. 3-(4-Amino-7-(1-methyl-1*H*-1,2,3-triazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for
 5 Example 36, with 1-methyl-5-(tributylstannyl)-1*H*-1,2,3-triazole replacing 1-ethyl-5-(trimethylstannyl)-1*H*-1,2,3-triazole in Step 2. The reaction mixture was purified with preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₂H₁₈N₉ (M+H)⁺: m/z = 408.2 found 408.1.

Example 38. 3-(4-Amino-2-((2-fluorophenyl)(hydroxy)methyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




The title compound, an enantiomeric pure compound, was prepared by separating a racemic mixture of Example 23. The chiral separation was conducted
 15 with chiral HPLC using Phenomenex Lux Cellulose-4, 21.2x250mm, 5μm column and 40% EtOH in hexanes (20ml/min) solvent system. Peak 1 was collected as the title compound. LC-MS calculated for C₂₄H₁₇FN₇O (M+H)⁺: m/z = 438.2 found 438.2.

Example 39. 3-(8-Amino-2-(2-fluorophenoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




Step 1: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(3-fluorophenoxy)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A reaction vial was charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-
 5 [1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (25 mg, 0.045 mmol) (from Example 27, Step 4), 3-fluorophenol (7.6 mg, 0.068 mmol), dimethylglycine (4.2 mg, 0.041 mmol), CuI (2.6 mg, 0.014 mmol), cesium carbonate (29 mg, 0.090 mmol) and dioxane (1 mL). The reaction mixture was purged with nitrogen for 5 min before heating to 100 °C and stirring for 15 h. The reaction mixture was then diluted with
 10 water and ethyl acetate. The organic layer was separated, washed with brine, dried over Na₂SO₄, filtered and concentrated. The resulting residue was purified via flash chromatography to give the desired product as a white solid (18 mg, 56%). LC-MS calculated for C₃₄H₂₈FN₆O₃ (M+H)⁺: m/z = 587.2; found 587.2.

Step 2: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(2-fluorophenoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




3-(8-(Bis(4-methoxybenzyl)amino)-2-(3-fluorophenoxy)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (18 mg, 0.025 mmol) was dissolved dichloromethane (1
 20 mL), and NBS (8 mg, 0.045 mmol) was added. The mixture was stirred at rt for 0.5 h before quenching by the addition of aqueous Na₂SO₃ solution. The organic layer was

separated, dried over Na₂SO₄, filtered and concentrated. The crude brominated product was added LiCl (1.3 mg, 0.030 mmol), CuI (5.8 mg, 0.030 mmol), Pd₂(dba)₃ (2.3 mg, 0.003 mmol), PPh₃ (1.3 mg, 0.005 mmol) and 4-(tributylstannyl)pyrimidine (14 mg, 0.038 mmol). The reaction mixture was dissolved in dioxane, and purged
 5 with nitrogen for 5 min, before heating to 100 °C for 15 h. The reaction mixture was then cooled to rt, filtered, concentrated, and purified via flash chromatography to give the desired product as a white solid (10 mg, 60%). LC-MS calculated for C₃₈H₃₀FN₈O₃ (M+H)⁺: m/z = 665.2; found 665.2.


10 *Step 3: 3-(8-Amino-2-(2-fluorophenoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile*

3-(8-(Bis(4-methoxybenzyl)amino)-2-(2-fluorophenoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.015 mmol) was added TFA (0.5 mL), and stirred at 100 °C for 5 min. The reaction mixture was then cooled to
 15 room temperature, solvent removed, diluted with methanol, and purified via prep-LCMS (pH 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₂H₁₄FN₈O (M+H)⁺: m/z = 425.1; found 425.2.

Example 40. 3-(8-Amino-2-(hydroxy(pyridin-2-yl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile
 20




Step 1: 3-(8-(Bis(4-methoxybenzyl)amino)-2-formyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (1.00 g, 1.99 mmol) (From Example 32, Step 1) and sodium periodate (1.92 g, 8.95 mmol) in THF (10 mL) and water (10 mL) was added osmium tetroxide (4% solution in water, 1.21 mL, 0.20 mmol). The reaction mixture was stirred at 60 °C for 1 h. The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The resulting residue was purified using flash chromatography to give the desired product (0.74 g, 74%). LC-MS calculated for C₂₉H₂₅N₆O₃ (M+H)⁺: m/z = 505.2; found 505.2.


Step 2: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(hydroxy(pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of 2-iodopyridine (0.40 g, 1.95 mmol) in THF (2 mL) was added isopropylmagnesium chloride lithium chloride complex solution (1.3 M, 1.3 mL, 1.71 mmol) dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 30 minutes. A solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.25 g, 0.49 mmol) in THF (2 mL) was added dropwise and the

reaction mixture was stirred at 0 °C for 1 h. The reaction mixture was then quenched with saturated aqueous NH₄Cl solution and diluted with DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The resulting residue was purified using flash chromatography to give the desired product (0.19 g, 67%). LC-MS calculated for C₃₄H₃₀N₇O₃ (M+H)⁺: m/z = 584.2; found 584.3.

Step 3: 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-(hydroxy(pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(hydroxy(pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.17 g, 0.29 mmol) in DCM (3 mL) was added a solution of NBS (0.052 g, 0.29 mmol) in DCM (3 mL) dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 1 h. The reaction mixture was concentrated, and the resulting residue was purified using flash chromatography to give the desired product (0.15 g, 78%). LC-MS calculated for C₃₄H₂₉BrN₇O₃ (M+H)⁺: m/z = 662.2; found 662.2.

Step 4: 3-(8-Amino-2-(hydroxy(pyridin-2-yl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile


To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(hydroxy(pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.010 g, 0.015 mmol), 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (0.013 g, 0.060 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.004 g, 0.005 mmol) in dioxane (0.250 mL) and water (0.050 mL) was added potassium phosphate tribasic (0.016 g, 0.075 mmol). The reaction mixture was stirred at 100 °C for 1 h. The reaction mixture

was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The crude material was dissolved in TFA (1 mL) and heated to 80 °C for 20 minutes. The solution was diluted with DMF (4 mL) and purified with prep-LCMS (pH = 2, acetonitrile/water+TFA) to give the desired product as a pair of enantiomers, and as a TFA salt (3.5 mg, 43%). LC-MS calculated for C₂₂H₁₇N₈O₂ (M+H)⁺: m/z = 425.2; found 425.3.

Example 41. 3-(8-Amino-2-(2-(1-methyl-1H-pyrazol-4-yl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Step 1: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A microwave vial was charge with 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (350 mg, 0.630 mmol) (from example 27 step 4), palladium acetate (7.07 mg, 0.032 mmol), and 2'-(dicyclohexylphosphino)-N,N,N',N'-tetramethylbiphenyl-2,6-diamine (27.5 mg, 0.063 mmol) was evacuated under high vacuum and backfilled with nitrogen. (2-chlorobenzyl)zinc(II) chloride (1386 µl, 0.693 mmol) was added via syringe. After addition, the reaction was heated to 60 °C for 1 h. The reaction solution was partitioned between EtOAc and sat. NH₄Cl solution. The layers were separated and the aqueous extracted further with EtOAc (2x). The combined organics were washed


with water and brine, dried over MgSO_4 , and concentrated. The residue was purified with flash chromatography to give the desired product (0.32 g, 82%). LC-MS calculated for $\text{C}_{35}\text{H}_{30}\text{ClN}_6\text{O}_2$ ($\text{M}+\text{H}$)⁺: m/z = 601.2; found 601.2.

- 5 *Step 2. 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-(2-chlorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile*



- To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.379 g, 0.631 mmol) in DCM (6.3 ml) was added 1-bromopyrrolidine-2,5-dione (0.107 g, 0.599 mmol) at 0 °C. After stirring at room temperature for 1 h, the reaction mixture was diluted with water. The organic layer was separated, dried over Na_2SO_4 , filtered and concentrated. The residue was purified with silica gel column to give the desired product (0.38 g, 89%). LC-MS calculated for $\text{C}_{35}\text{H}_{29}\text{BrClN}_6\text{O}_2$ ($\text{M}+\text{H}$)⁺: m/z = 679.1, 681.1; found 679.2, 681.2.
- 10
- 15

- Step 3. 3-(8-(Bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile*




- A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(2-chlorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (381 mg, 0.560 mmol), 4-
- 20

(tributylstannyl)pyrimidine (310 mg, 0.840 mmol), and copper(I) chloride (66.6 mg, 0.672 mmol), lithium chloride (28.5 mg, 0.672 mmol) and tetrakis(triphenylphosphine)palladium(0) (64.7 mg, 0.056 mmol) in THF (6 ml) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction mixture was diluted with ethyl acetate and water. The reaction mixture was filtered through a pad of Celite®. The organic layer was separated and dried over Na₂SO₄, filtered and concentrated. The residue was purified with flash chromatography to give the desired product (0.31 g, 83%). LC-MS calculated for C₃₉H₃₂ClN₈O₂ (M+H)⁺: m/z = 679.2; found 679.2.


Step 4. 3-(8-Amino-2-(2-(1-methyl-1H-pyrazol-4-yl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (10mg, 0.015 mmol), 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (3.68 mg, 0.018 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (1.158 mg, 1.472 μmol) and tripotassium phosphate hydrate (7.46 mg, 0.032 mmol) in 1,4-dioxane (1.0 mL)/water (0.3 mL) was stirred at 80 °C for 1 h. The reaction mixture was diluted with EtOAc and water, the organic layer was separated and concentrated. The residue was treated with TFA (1 mL) at 80 °C for 20 min. The solvent was removed, the residue was dissolved in methanol and DMSO, then purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₇H₂₁N₁₀ (M+H)⁺: m/z = 485.2; found 485.2.

Example 42. 3-(8-Amino-5-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Step 1: 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.697 g, 1.228 mmol) (From Example 27, Step 5) in DCM (12 ml) was added 1-bromopyrrolidine-2,5-dione (0.219 g, 1.228 mmol) at 0 °C. After stirring at room temperature for 1h, the reaction mixture was diluted with water. The organic layer was separated, dried over Na₂SO₄, filtered and concentrated. The residue was purified with silica gel column to give the desired product (0.74 g, 93%). LC-MS calculated for C₃₄H₂₉BrN₇O₂ (M+H)⁺: m/z = 646.2, 648.2; found 646.2, 648.2.


Step 2: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-5-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (390 mg, 0.603 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (111 mg, 0.724 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (47.5 mg, 0.060 mmol) and tripotassium phosphate hydrate (306 mg, 1.327 mmol) in 1,4-dioxane (5.0 mL)/water (1.7 mL) was stirred at 80 °C for 1 h. The reaction mixture was diluted with EtOAc and water. The organic layer

was separated, dried over Na₂SO₄, filtered and concentrated. The residue was purified with silica gel column to give the desired product (0.28 g, 77%). LC-MS calculated for C₃₆H₃₂N₇O₂ (M+H)⁺: m/z = 594.3; found 594.3.

5 *Step 3. 3-(8-(Bis(4-methoxybenzyl)amino)-5-formyl-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile*




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-5-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (277 mg, 0.467 mmol) in THF
 10 (2.3 mL) and water (2.3 mL) was added 4% osmium tetroxide in water (233 μ l, 0.037 mmol) and sodium metaperiodate (449 mg, 2.1 mmol). The reaction mixture was stirred at 60 °C for 1 h. The mixture was filtered through a plug of Celite®, rinsed with THF. The organic layer was concentrated under vacuum. The residue was purified with flash chromatography to give the desired product (0.21 g, 76%). LC-MS
 15 calculated for C₃₅H₃₀N₇O₃ (M+H)⁺: m/z = 596.2; found 596.2.

Step 4. 3-(8-Amino-5-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (38 mg, 0.064 mmol) in methanol (0.3 mL) and 1,2-dichloroethane (0.3 mL) was added 1-((1-isocynoethyl)sulfonyl)-4-methylbenzene (13.35 mg, 0.064 mmol). The resulting mixture was heated at 70 °C overnight. The reaction mixture was diluted with DCM and water, the organic layer was separated and concentrated. The residue was treated
 25 with TFA (1 mL) at 80 °C for 20 min. The solvent was removed, the residue was dissolved in methanol and DMSO, then purified with prep-LCMS (pH 2,


acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{22}H_{17}N_8O$ ($M+H$)⁺: m/z = 409.2; found 409.1.

Example 43. 3-(8-Amino-5-(4-ethyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for Example 42 with 1-((1-isocyanopropyl)sulfonyl)-4-methylbenzene replacing 1-((1-isocyanoethyl)sulfonyl)-4-methylbenzene in Step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{23}H_{19}N_8O$ ($M+H$)⁺: m/z = 423.2; found 423.2.

Example 44. 3-(8-Amino-5-(3-methylpyridin-4-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-amino-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 27, Step 7; 10 mg, 0.025 mmol), (3-methylpyridin-4-yl)boronic acid (4.1 mg, 0.030 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (1.9 mg, 2.5 μ mol) and tripotassium phosphate hydrate (12.5 mg, 0.054 mmol) in 1,4-dioxane (1.0mL)/water (0.3 mL) was stirred at 80 °C for 1 h. The reaction mixture was diluted with DCM and water, the organic layer was separated and concentrated. The resulting residue was dissolved in methanol and DMSO, then purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{24}H_{19}N_8$ ($M+H$)⁺: m/z = 419.2; found 419.2.

Example 45. 3-(8-Amino-2-(imidazo[1,2-*a*]pyridin-8-ylmethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



A microwave vial charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-(bromomethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (20 mg, 0.031 mmol) (from Example 32, Step 5), 8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-*a*]pyridine (15.08 mg, 0.062 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (2.430 mg, 3.09 μ mol) and cesium carbonate (30.2 mg, 0.093 mmol) was sealed and evacuated under high vacuum and refilled with nitrogen (repeated three times). 1,4-Dioxane (2.0 mL) and water (0.67 mL) was stirred at 90 °C overnight. The reaction mixture was diluted with ethyl acetate and water. The organic layer was separated and concentrated. The residue was treated with TFA (1 mL) at 80 °C for 20 min. After removal of the volatile, the residue was dissolved in methanol and DMSO and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{24}H_{17}N_{10}$ ($M+H$)⁺: m/z = 445.2; found 445.2.


Example 46. 3-(8-Amino-2-(pyrazolo[1,5-*a*]pyridin-7-ylmethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



The title compound was prepared using similar procedures as described for Example 45 with 7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazolo[1,5-*a*]pyridine replacing 8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-*a*]pyridine. The reaction mixture was purified by prep-HPLC (pH = 2,


acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{24}H_{17}N_{10}$ ($M+H$)⁺: m/z = 445.2; found 445.2.

Example 47 and Example 48. 3-(2-((2*H*-Indazol-2-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile and 3-(2-((1*H*-Indazol-1-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(bromomethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (15 mg, 0.023 mmol) (from Example 32, Step 5), 1*H*-indazole (4.1 mg, 0.035 mmol), cesium carbonate (22.64 mg, 0.069 mmol) in DMF (300 μ L) was stirred at 50 °C for 15 min. The reaction was diluted with EtOAc and water. The organic layer was separated and concentrated. The residue was treated with TFA (1 mL) at 80 °C for 20 min. The volatile was removed *in vacuo*, the residue was dissolved in methanol and DMSO and purified with prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the two products both as TFA salts. LC-MS calculated for both compounds $C_{24}H_{17}N_{10}$ ($M+H$)⁺: m/z = 445.2; found 445.2.

Example 49. 3-(8-Amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1




Step 1: Methyl 3-bromo-1-(2-(3-cyanophenyl)-2-oxoethyl)-1H-1,2,4-triazole-5-carboxylate




To a solution of methyl 3-bromo-1H-1,2,4-triazole-5-carboxylate (5.0 g, 24.3 mmol), 3-(2-bromoacetyl)benzonitrile (5.44 g, 24.3 mmol) in DMF (100 mL) was added potassium carbonate (3.35 g, 24.3 mmol). The reaction mixture was stirred at ambient temperature for 2 h. The reaction mixture was then diluted with water and DCM. The organic layer was separated, washed with brine, dried over Na₂SO₄, filtered and concentrated. The resulting residue was purified via flash chromatography to give the desired product as a white solid (5.2 g, 61%). LC-MS calculated for C₁₃H₁₀BrN₄O₃ (M+H)⁺: m/z = 349.0; found 349.0.

Step 2: 3-(2-Bromo-8-oxo-7,8-dihydro-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Methyl 3-bromo-1-(2-(3-cyanophenyl)-2-oxoethyl)-1H-1,2,4-triazole-5-carboxylate (10.5 g, 30.1 mmol) was dissolved in acetic acid (100 mL), and ammonium acetate (23.18 g, 301 mmol) was added. The mixture was stirred at 110 °C for 12 h. After cooling to room temperature, the reaction mixture was diluted with water. The resulting precipitate was collected via filtration, washed with water, and dried under vacuum to afford the product (8.4 g, 88%). LC-MS calculated for C₁₂H₇BrN₅O (M+H)⁺: m/z = 316.0; found 316.0.

Step 3: 3-(2-Bromo-8-chloro-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(2-bromo-8-oxo-7,8-dihydro-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (8.4 g, 26.6 mmol) and POCl₃ (49.5 mL, 531 mmol) was stirred at 110 °C overnight. After cooling to room temperature, the reaction mixture was slowly added to a flask containing ice and sodium bicarbonate. The resulting precipitate was collected via filtration, washed with water, and dried to afford the product (8.8 g, 99%). LC-MS calculated for C₁₂H₆BrClN₅ (M+H)⁺: m/z = 336.0; found 336.0.

*Step 4: 3-(8-(Bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*




10

A mixture of 3-(2-bromo-8-chloro-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (8.99 g, 26.9 mmol), bis(4-methoxybenzyl)amine (10.37 g, 40.3 mmol), and DIPEA (9.4 mL, 53.7 mmol) in DMF (134 mL) was stirred at 65 °C overnight. The reaction mixture was cooled to room temperature, and diluted with water. The resulting precipitate was collected via filtration, and dried to afford the product (14.1 g, 94%). LC-MS calculated for C₂₈H₂₄BrN₆O₂ (M+H)⁺: m/z = 555.1; found 555.1.

15


*Step 5: 3-(8-(Bis(4-methoxybenzyl)amino)-2-vinyl-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*

20




A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (10.0 g, 18.0 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (3.88 g, 25.2 mmol), potassium phosphate tribasic (9.55 g, 45.0 mmol) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (567 mg, 0.72 mmol) in 1,4-dioxane (200 mL) and water (50 mL) was stirred at 85 °C for 2 hrs. The reaction mixture was cooled to room temperature, and most of 1, 4-dioxane was removed. The resulting precipitate was collected via filtration, washed with water and dried to afford the crude product (9.1 g), which was used in the next step directly. LC-MS calculated for C₃₀H₂₇N₆O₂ (M+H)⁺: m/z = 503.2; found 503.1.

*Step 6. 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-vinyl-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-vinyl-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (717 mg, 1.43 mmol) in 10 mL of dichloromethane, 1-bromopyrrolidine-2,5-dione (254 mg, 1.43 mmol) was added at 0 °C. The resulting mixture was stirred for 4 hrs, and directly purified by a silica gel column to afford the desired product (780 mg, 94%). LC-MS calculated for C₃₀H₂₆BrN₆O₂ (M+H)⁺: m/z = 581.1; found 581.2.

Step 7: 3-(8-(Bis(4-methoxybenzyl)amino)-5-(pyrimidin-4-yl)-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (260 mg, 0.45 mmol), 4-(tributylstannyl)pyrimidine (215 mg, 0.58 mmol), lithium chloride (28.4 mg, 0.67 mmol), copper(I) chloride (67 mg, 0.67 mmol), and Tetrakis(triphenylphosphine)palladium(0) (52 mg, 0.045 mmol) in THF (5 mL) was stirred at 90 °C for 45 mins. The reaction mixture was quenched with water and extracted with dichloromethane. The combined organic layers were concentrated, and purified by a silica gel column to afford the desired product (176 mg, 67%). LC-MS calculated for $C_{34}H_{29}N_8O_2$ ($M+H$)⁺: m/z = 581.2; found 581.1.

Step 8: 3-(8-(Bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-(pyrimidin-4-yl)-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (176 mg, 0.3 mmol), osmium(VIII) oxide (3 mg in 0.3 mL water, 0.015 mmol), and sodium periodate (292 mg, 1.36 mmol) in THF/water (1:1, 6 mL) was stirred at 65 °C for 1 h. The reaction mixture

was cooled to room temperature, and extracted with dichloromethane. The combined organic layers were concentrated, and purified by silica gel column to afford the desired product (130 mg, 74%). LC-MS calculated for $C_{33}H_{27}N_8O_3$ ($M+H$)⁺: m/z = 583.2; found 583.2.


5

Step 9: 3-(8-Amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

Preparation of the Grignard reagent: To a solution of 1,3-difluoro-2-iodobenzene (142 mg, 0.6 mmol) in tetrahydrofuran (1 mL), isopropylmagnesium chloride solution (296 μ l, 2 M) was added at -10 °C. The resulting mixture was stirred for 1 h, and used directly in the following step.

To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (120 mg, 0.2 mmol) in THF (2 mL), the freshly prepared Grignard reagent from previous step was added at -10 °C. The reaction mixture was stirred for 30 min, quenched with ammonium chloride solution (4 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The resulting material was dissolved in TFA (5 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous $NaHCO_3$ solution. The crude material was directly purified by a silica gel column to afford the desired product (60 mg, 64%) as a racemic mixture. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5 μ m Cellulose-4, 21.2x250mm) and 75% EtOH in hexanes (20 mL/min) solvent system. Peak 1 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{23}H_{15}F_2N_8O$ ($M+H$)⁺: m/z = 457.1; found 457.0.


Example 50. 3-(8-Amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 2



This compound was prepared using the same procedure as described for
 5 Example 49. The product was separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.2x250mm) and 75% EtOH in hexanes (20 mL/min) solvent system. Peak 2 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₅F₂N₈O (M+H)⁺: m/z = 457.1; found 457.0.

10 ¹H NMR (600 MHz, DMSO-*d*₆) δ 9.14 (d, *J* = 1.3 Hz, 1H), 8.95 (d, *J* = 5.2 Hz, 1H), 7.90 (dd, *J* = 5.2, 1.4 Hz, 1H), 7.88 (s, 1H), 7.78 (dt, *J* = 7.6, 1.4 Hz, 1H), 7.74 (t, *J* = 1.4 Hz, 1H), 7.54 (dt, *J* = 7.9, 1.3 Hz, 1H), 7.51 – 7.40 (m, 2H), 7.09 (t, *J* = 8.4 Hz, 2H), 6.27 (s, 1H).


Example 51. 3-(8-Amino-2-((2,5-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (115 mg, 0.2 mmol) in THF (1
 20 mL), the freshly prepared Grignard reagent (prepared using a similar procedure as described in Example 49, Step 9, using 1,4-difluoro-2-iodobenzene instead of 1,3-difluoro-2-iodobenzene) was added at -10 °C. The reaction mixture was stirred for 30 min, quenched with ammonium chloride solution (4 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The


resulting material was dissolved in TFA (5 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (70 mg, 77%) as a racemic mixture. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.2x250mm) and 40% EtOH in hexanes (20 mL/min) as the mobile phase. Peak 1 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₅F₂N₈O (M+H)⁺: m/z = 457.1; found 457.0.

Example 52. 3-(8-Amino-2-((2,5-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 2




This compound was prepared using the same procedure as described for Example 51. The product was separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.2x250mm) and 40% EtOH in hexanes (20 mL/min) as the mobile phase. Peak 2 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₅F₂N₈O (M+H)⁺: m/z = 457.1; found 457.0.

Example 53. 3-(8-Amino-2-((2,3-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1




To a solution of 1,2-difluoro-3-iodobenzene (70 mg, 0.3 mmol) in tetrahydrofuran (1 mL), isopropylmagnesium chloride (150 μ l, 2 M solution) was added at -10 °C, and the resulting mixture was stirred for 1 h before a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-
 5 α]pyrazin-6-yl)benzonitrile (60 mg, 0.1 mmol) in THF (1mL) was added at -10 °C. The reaction mixture was stirred for 30 min, quenched with ammonium chloride solution (4 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The resulting material was dissolved in TFA (5 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room
 10 temperature, concentrated, and basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (30 mg, 70%) as a racemic mixture. The resulting product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5 μ m Cellulose-4, 21.2x250mm) and 40% EtOH in hexanes (20 mL/min) as the mobile phase. Peak 1
 15 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₅F₂N₈O (M+H)⁺: m/z = 457.1; found 457.0.

**Example 54. 3-(8-Amino-2-((2,3-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-
 20 4-yl)-[1,2,4]triazolo[1,5- α]pyrazin-6-yl)benzonitrile, Peak 2**



This compound was prepared using the same procedure as described for Example 53. The product was separated with chiral HPLC using a chiral column (Phenomenex Lux 5 μ m Cellulose-4, 21.2x250mm) and 40% EtOH in hexanes (20
 25 mL/min) as the mobile phase. Peak 2 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₅F₂N₈O (M+H)⁺: m/z = 457.1; found 457.0.


Example 55. 3-(8-Amino-2-((2-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (100 mg, 0.17 mmol) in THF (2 mL), the corresponding Grignard reagent (freshly prepared using a similar procedure as described in Example 49, Step 9, using 1-fluoro-2-iodobenzene instead of 1,3-difluoro-2-iodobenzene) was added at -10 °C. The reaction mixture was stirred for 30 min, quenched with ammonium chloride solution (4 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The resulting material was dissolved in TFA (5 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (60 mg, 50%) as a racemic mixture. The resulting product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.2x250mm) and 60% EtOH in hexanes (20 mL/min) as the mobile phase. Peak 1 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₆FN₈O (M+H)⁺: m/z = 439.1; found 439.0.

20


Example 56. 3-(8-Amino-2-((2-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 2



This compound was prepared using the same procedure as described for Example 55. The product was separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.2x250mm) and 60% EtOH in hexanes (20


mL/min) as the mobile phase. Peak 2 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{23}H_{16}FN_8O$ ($M+H$)⁺: m/z = 439.1; found 439.0.

5 **Example 57. 3-(8-Amino-2-((2-chlorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1**



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (100 mg, 0.17 mmol) in THF (2
10 mL), the corresponding Grignard reagent (freshly prepared using a similar procedure as described in Example 49, Step 9, using 1-chloro-2-iodobenzene instead of 1,3-difluoro-2-iodobenzene) was added at -10 °C. The reaction mixture was stirred for 30 min, quenched with ammonium chloride solution (4 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The
15 resulting material was dissolved in TFA (5 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous $NaHCO_3$ solution. The crude material was directly purified by a silica gel column to afford the desired product (50 mg, 64%) as a racemic mixture. The
20 resulting product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.2x250mm) and 45% EtOH in hexanes (20 mL/min) as the mobile phase. Peak 1 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{23}H_{16}ClN_8O$ ($M+H$)⁺: m/z = 455.1; found 455.1.


Example 58. 3-(8-Amino-2-((2-chlorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 2



This compound was prepared using the same procedure as described for
 5 Example 57. The product was separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.2x250mm) and 45% EtOH in hexanes (20 mL/min) as the mobile phase. Peak 2 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₆ClN₈O (M+H)⁺: m/z = 455.1; found 455.1.


10

Example 59. 3-(8-Amino-2-(hydroxy(phenyl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (20 mg, 0.17 mmol) in THF (2
 15 mL), phenylmagnesium chloride (3 M solution in ethyl ether, 0.17 mL) was added at -10 °C. The reaction mixture was stirred for 30 min, quenched with ammonium chloride solution (4 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The resulting material was dissolved in TFA
 20 (5 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and purified with preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the product as a TFA salt. LC-MS calculated for C₂₃H₁₇N₈O (M+H)⁺: m/z = 421.2; found 421.1.

Example 60. 3-(8-Amino-2-(phenylsulfonyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



Step 1: 3-(8-Amino-2-(phenylsulfonyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




5

A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (33 mg, 0.06 mmol) (from Example 27, Step 4) and sodium benzenesulfinate (100 mg, 0.6 mmol) in 1 mL DMSO was stirred at 110 °C for 48 hrs, the reaction was quenched with ammonium chloride and extracted with dichloromethane. The combined organic layers were concentrated and purified with silica gel column to afford the desired Product (18 mg, 50%), LC-MS calculated for C₃₄H₂₉N₆O₄S (M+H)⁺: m/z = 617.2; found 617.2.

Step 2: 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-(phenylsulfonyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

15



To a solution of 3-(8-Amino-2-(phenylsulfonyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (20 mg, 0.032 mmol) in 1 mL dichloromethane, 1-bromopyrrolidine-2,5-dione (6 mg, 0.032 mmol) was added at 0 °C. The resulting mixture was stirred for 16 hrs before being purified by silica gel column to afford the desired product (18 mg, 80%). LC-MS calculated for C₃₄H₂₈BrN₆O₄S (M+H)⁺: m/z = 695.1; found 695.2.

20


Step 3: 3-(8-Amino-2-(phenylsulfonyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

A mixture of 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-(phenylsulfonyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (10 mg, 0.014 mmol), 4-(tributylstannyl)pyrimidine (11 mg, 0.029 mmol), lithium chloride (1.0 mg, 0.022 mmol), copper(I) chloride (2.1 mg, 0.022 mmol), and

5 tetrakis(triphenylphosphine)palladium(0) (3.3 mg, 2.88 μ mol) in THF (2 mL) was stirred at 90 °C for 45 mins. The reaction mixture was filtered and the organic solvent was removed by vacuo, the crude product was dissolved in 3 mL TFA and stirred at 80 °C for 20 mins. After TFA being removed, the crude product was purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to afford the desired


10 product as a TFA salt. LC-MS calculated for $C_{22}H_{15}N_8O_2S$ ($M+H$)⁺: m/z = 455.1; found 455.1.

Example 61. 3-(8-Amino-2-(azetidine-1-carbonyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile



15


*Step 1: 3-(2-(Azetidine-1-carbonyl)-8-(bis(4-methoxybenzyl)amino)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*



20 A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (99 mg, 0.18 mmol) and azetidine (50 mg, 0.89 mmol), $PdCl_2(dppf)$ (26 mg, 0.036 mmol) and Na_2CO_3 (57 mg, 0.48 mmol) in 5 mL 1,4-dioxane and 1 mL water was stirred at 80 °C for 18 hrs under CO balloon atmosphere, the reaction was quenched with ammonium chloride and extracted with

25 dichloromethane. The combined organic layers were concentrated and purified with silica gel column to afford the desired Product (44 mg, 44%), LC-MS calculated for $C_{32}H_{30}N_7O_3$ ($M+H$)⁺: m/z = 560.2; found 560.2.

Step 2: 3-(2-(Azetidine-1-carbonyl)-8-(bis(4-methoxybenzyl)amino)-5-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(2-(azetidine-1-carbonyl)-8-(bis(4-methoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (40 mg, 0.125 mmol) in 3 mL dichloromethane, 1-bromopyrrolidine-2,5-dione (23 mg, 0.125 mmol) was added at 0 °C. The resulting mixture was stirred for 16 hrs before being purified by silica gel column to afford the desired product (50 mg, 100%). LC-MS calculated for $C_{32}H_{29}BrN_7O_3$ (M+H)⁺: m/z = 638.1; found 638.2.

Step 3: 3-(8-Amino-2-(azetidine-1-carbonyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

A mixture of 3-(2-(azetidine-1-carbonyl)-8-(bis(4-methoxybenzyl)amino)-5-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.016 mmol), 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (7 mg, 0.032 mmol), sodium carbonate (5 mg, 0.048 mmol) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (5 mg, 0.006 mmol) in 4:1 THF/Water (2 mL) was stirred at 100 °C for 75 mins. The reaction mixture was filtered and the organic solvent was removed by vacuo, the crude product was dissolved in 3 mL TFA and stirred at 80 °C for 20 mins. After TFA being removed, the crude product was purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to afford the desired product as a TFA salt. LC-MS calculated for $C_{20}H_{17}N_8O_2$ (M+H)⁺: m/z = 401.2; found 401.1.


Example 62. 3-(8-amino-5-(6-hydroxypyridin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-Amino-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 27, Step 7; 10 mg, 0.025 mmol), (6-methoxypyridin-3-yl)boronic acid (10 mg, 0.042 mmol), cesium carbonate (17.7 mg, 0.116 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (2.26 mg, 2.88 μ mol) (XPhos Pd G2) in 1,4-dioxane (500 μ l) and water (100 μ l) was purged with N₂ and heated at 95 °C for 1 h.

The mixture was concentrated and added MeCN (0.5 mL) and TMSCl (0.5 mL), the mixture was heated at 80 °C for 30 min before concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₃H₁₇N₈O (M+H)⁺: 421.1; found 421.2.


Example 63. 3-(8-amino-2-(benzo[d]oxazol-4-ylmethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(bromomethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 32, Step 5; 10 mg, 0.022 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[d]oxazole (11 mg, 0.044 mmol), cesium carbonate (17.7 mg, 0.116 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (2.26 mg, 2.88 μ mol) (XPhos Pd G2) in 1,4-dioxane (500 μ l)


and water (100 μ l) was purged with N₂ and heated at 95 °C for 1 h. The mixture was concentrated and added TFA (1 mL), the mixture was heated at 100 °C for 30 min before concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₄H₁₆N₉O (M+H)⁺: 446.1; found 446.1.

Example 64. 3-(8-amino-2-(2-fluoro-6-(1-methyl-1H-pyrazol-5-yl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-Amino-2-(2-chloro-6-fluorobenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 30, Step 3; 10 mg, 0.022 mmol), 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (9.2 mg, 0.044 mmol), cesium carbonate (17.7 mg, 0.116 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (2.26 mg, 2.88 μ mol) (XPhos Pd G2) in 1,4-dioxane (500 μ l) and water (100 μ l) was purged with N₂ and heated at 95 °C for 1 h. The mixture was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₇H₂₀N₁₀F (M+H)⁺: 503.2; found 503.1.

Example 65. (R)-1-(2-((8-amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)-3-methylpyrrolidine-3-carboxylic acid




The title compound was prepared using similar procedures as described for Example 30 with (R)-3-methylpyrrolidine-3-carboxylic acid replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{30}H_{27}N_9O_2F$ (M+H)⁺: m/z = 564.2; found 564.2.

Example 66. 3-(8-amino-2-(2-fluoro-6-((6-methyl-5-oxo-2,6-diazaspiro[3.4]octan-2-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



The title compound was prepared using similar procedures as described for Example 30 with 6-methyl-2,6-diazaspiro[3.4]octan-5-one replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{31}H_{28}N_{10}OF$ (M+H)⁺: m/z = 575.2; found 575.2.

Example 67. 3-(8-amino-2-(2-fluoro-6-((6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for Example 30 with hexahydropyrrolo[1,2-a]pyrazin-6(2H)-one replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH

= 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{31}H_{28}N_{10}OF$ ($M+H$)⁺: m/z = 575.2; found 575.2.

Example 68. (S)-3-(8-amino-2-(2-fluoro-6-(((2-oxopyrrolidin-3-


5 **yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile**



The title compound was prepared using similar procedures as described for Example 30 with (S)-3-aminopyrrolidin-2-one replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{28}H_{24}N_{10}OF$ ($M+H$)⁺: m/z = 535.2; found 535.2.


Example 69. 2-((2-((8-amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-

15 **[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)amino)acetamide**




The title compound was prepared using similar procedures as described for Example 30 with 2-aminoacetamide replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{26}H_{22}N_{10}OF$ ($M+H$)⁺: m/z = 509.2; found 509.2.

Example 70. 3-(8-amino-2-(2-fluoro-6-((3-oxopiperazin-1-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for
 5 Example 30 with piperazin-2-one replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{28}H_{24}N_{10}OF$ (M+H)⁺: m/z = 535.2; found 535.2.

Example 71. (1S,3S)-3-((2-((8-amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)amino)cyclobutane-1-carboxylic acid



The title compound was prepared using similar procedures as described for
 15 Example 30 with (1S,3S)-3-aminocyclobutane-1-carboxylic acid replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{29}H_{25}N_9O_2F$ (M+H)⁺: m/z = 550.2; found 550.2.


Example 72. 3-(8-amino-2-(2-fluoro-6-(((1-methyl-2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



5 The title compound was prepared using similar procedures as described for Example 30 with 3-amino-1-methylpyrrolidin-2-one replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{29}H_{26}N_{10}OF$ ($M+H$)⁺: m/z = 549.2; found 549.2.


10

Example 73. 1-(2-((8-amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)azetidine-3-carbonitrile




15 The title compound was prepared using similar procedures as described for Example 30 with azetidine-3-carbonitrile replacing (S)-pyrrolidine-3-carboxylic acid. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{28}H_{22}N_{10}F$ ($M+H$)⁺: m/z = 517.2; found 517.2.

Example 74. 3-(8-amino-2-(2-fluoro-6-(((2-methyl-2H-1,2,3-triazol-4-yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




5 To a solution of 3-(8-amino-2-(2-fluoro-6-formylbenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 30, step 5; 10 mg, 0.02 mmol) in DCM (0.5 mL) was added 2-methyl-2H-1,2,3-triazol-4-amine (4.0 mg, 0.04 mmol) then TFA (4 μ L, 0.08 mmol). After 1h, sodium triacetoxymethylborohydride (8.5 mg, 0.04 mmol) was added to the reaction. The reaction was stirred overnight and the
10 mixture was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for $C_{27}H_{22}FN_{12}$ (M+H)⁺: 533.2 found 533.2.

Example 75. (S)-3-(8-amino-2-(2-(((2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Step 1: 3-(8-(bis(4-methoxybenzyl)amino)-5-(pyrimidin-4-yl)-2-(2-vinylbenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 41, step 3; 149 mg, 0.22 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (68 mg, 0.44 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (17 mg, 21.6 μ mol) and K_3PO_4 (94 mg, 0.44 mmol) in 1,4-dioxane (4 mL)/water (0.8 mL) was stirred at 110 $^{\circ}C$ for 1 h. The reaction mixture was concentrated under vacuum and the resulting residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for $C_{41}H_{35}N_8O_2$ ($M+H$) $^{+}$: m/z = 671.3; found 671.2.

Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-formylbenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-(pyrimidin-4-yl)-2-(2-vinylbenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (60 mg, 0.089 mmol) in THF (1 mL) and water (1 mL) was added 0.157 M osmium tetroxide in water (0.02 mmol). After 2 min, sodium metaperiodate (86 mg, 0.4 mmol) was added. The reaction mixture was heated at 60 °C for 1 h before quenched with sat. Na₂S₂O₃. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated to afford the product as a light yellow oil. LC-MS calculated for C₄₀H₃₃N₈O₃ (M+H)⁺: m/z = 673.3; found 673.3.

Step 3: (S)-3-(8-amino-2-(2-(((2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile


To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-formylbenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (13.4 mg, 0.02 mmol) in DCM (0.5 mL) was added (S)-3-aminopyrrolidin-2-one (4.1 mg, 0.04 mmol) then acetic acid (4 µL, 0.08 mmol). After 1h, sodium triacetoxyborohydride (8.5 mg, 0.04 mmol) was added to the reaction. The reaction was stirred overnight and the mixture was concentrated, then 0.5 mL of TFA was added to the mixture and the mixture was heated at 100 °C for 10 min. The reaction was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₈H₂₅N₁₀O (M+H)⁺: 517.2 found 517.2.

Example 76. (R)-1-(2-((8-amino-6-(3-cyanophenyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)-3-methylpyrrolidine-3-carboxylic acid




25

Step 1: 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole




To a solution of 4-methyloxazole (0.654 g, 7.87 mmol) in heptane (3 mL) and Et₂O (1 mL) was added (1,5-cyclooctadiene)(methoxy)iridium(I) dimer (0.221 g, 0.393 mmol), 4,4'-Di-tert-butyl-2,2'-dipyridyl (0.211 g, 0.787 mmol) and 4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.841 mL, 11.80 mmol). The vial was then evacuated under high vacuum and backfilled with nitrogen. The reaction was stirred overnight, then concentrated and purified via flash chromatography to afford the desired product as a colorless oil. LC-MS calculated for C₁₀H₁₇BNO₃ (M+H)⁺: m/z = 210.1; found 128.0 (as the corresponding boronic acid).

Step 2: 3-(8-amino-2-(2-chloro-6-fluorobenzyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-amino-5-bromo-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 30, Step 2; 500 mg, 1.09 mmol), 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (274 mg, 1.31 mmol), tetrakis(triphenylphosphine)palladium(0) (126 mg, 0.11 mmol) and Cs₂CO₃ (712 mg, 2.185 mmol) in 1,4-dioxane (2 mL) and water (200 µl) was purged with N₂ and heated at 95 °C for 7 h. The mixture was concentrated and purified via flash chromatography to afford the desired product as a white solid. LCMS calculated for C₂₃H₁₆N₇OCIF (M+H)⁺: 460.1; found 460.1.

Step 3: 3-(8-amino-2-(2-fluoro-6-vinylbenzyl)-5-(4-methyloxazol-5-yl)-
[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-amino-2-(2-chloro-6-fluorobenzyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (101 mg, 0.22 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (68 mg, 0.44 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (17 mg, 21.6 μ mol) and K_3PO_4 (94 mg, 0.44 mmol) in 1,4-dioxane (2 mL)/water (0.4 mL) was stirred at 110 $^{\circ}C$ for 1 h. The reaction mixture was concentrated under vacuum and the resulting residue was purified with flash chromatography to give the desired product as a colorless oil. LC-MS calculated for $C_{25}H_{19}FN_7O$ ($M+H$) $^{+}$: m/z = 452.1; found 452.2.

Step 4: 3-(8-amino-2-(2-fluoro-6-formylbenzyl)-5-(4-methyloxazol-5-yl)-
[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-amino-2-(2-fluoro-6-vinylbenzyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (40 mg, 0.089 mmol) in THF (1 mL) and water (1 mL) was added 0.157 M osmium tetroxide in water (0.02 mmol). After 2 min, sodium metaperiodate (86 mg, 0.4 mmol) was added. The reaction mixture was heated at 60 $^{\circ}C$ for 1 h before quenched with sat. $Na_2S_2O_3$. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na_2SO_4 , filtered, and concentrated to afford the product as a light yellow oil. LC-MS calculated for $C_{24}H_{17}FN_7O_2$ ($M+H$) $^{+}$: m/z = 454.1; found 454.1


Step 5: (R)-1-(2-((8-amino-6-(3-cyanophenyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)-3-methylpyrrolidine-3-carboxylic acid

To a solution of 3-(8-amino-2-(2-fluoro-6-formylbenzyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.02 mmol) in DCM (0.5 mL) was added (R)-3-methylpyrrolidine-3-carboxylic acid (4.8 mg, 0.04 mmol) then acetic acid (4 μ L, 0.08 mmol). After 1h, sodium triacetoxyborohydride (8.5 mg, 0.04 mmol) was added to the reaction. The reaction was stirred overnight and the mixture was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for $C_{30}H_{28}FN_8O_3$ (M+H)⁺: 567.2 found 567.2.

Example 77. 3-(8-amino-2-(amino(2,6-difluorophenyl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Step 1: 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-formyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 32, step 2; 174 mg, 0.3 mmol), osmium(VIII) oxide (3 mg in 0.3 mL water, 0.015 mmol), and sodium periodate (292 mg, 1.36 mmol) in THF/water (1:1, 6 mL) was stirred at 65 °C for 1 h. The reaction mixture was cooled to room temperature, and extracted with dichloromethane. The combined organic layers were concentrated, and purified by

silica gel column to afford the desired product. LC-MS calculated for $C_{29}H_{24}N_6O_3Br$ ($M+H$)⁺: m/z = 583.1; found 583.1.


Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



Preparation of the Grignard reagent: To a solution of 1,3-difluoro-2-iodobenzene (142 mg, 0.6 mmol) in tetrahydrofuran (1 mL), isopropylmagnesium chloride solution (296 μ L, 2 M) was added at -10 °C. The resulting mixture was stirred for 1 h, and used directly in the following step.

To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-formyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (120 mg, 0.2 mmol) in THF (2 mL), the freshly prepared Grignard reagent from previous step was added at -10 °C. The reaction mixture was stirred for 30 min, quenched with ammonium chloride solution (4 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum and purified by a silica gel column to afford the desired product as a racemic mixture. LC-MS calculated for $C_{35}H_{28}N_6O_3BrF_2$ ($M+H$)⁺: m/z = 697.1; found 697.1.


Step 3: 3-(8-(bis(4-methoxybenzyl)amino)-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (382 mg, 0.55 mmol), 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole

(137 mg, 0.65 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (17 mg, 21.6 μ mol) and Cs_2CO_3 (356 mg, 1.09 mmol) in 1,4-dioxane (2 mL) and water (200 μ l) was purged with N_2 and heated at 95 $^\circ\text{C}$ for 7 h. The mixture was concentrated and purified via flash chromatography to afford the desired product as a colorless oil. LCMS calculated for $\text{C}_{39}\text{H}_{32}\text{N}_7\text{O}_4\text{F}_2$ ($\text{M}+\text{H}$) $^+$: 700.2; found 700.2.

Step 4: 3-(8-(bis(4-methoxybenzyl)amino)-2-(chloro(2,6-difluorophenyl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (201 mg, 0.29 mmol) in 2 mL of dichloromethane, thionyl chloride (105 μ l, 1.435 mmol) was added at rt. The resulting mixture was stirred for 4h, concentrated and used in next step without any further purification. LC-MS calculated for $\text{C}_{39}\text{H}_{31}\text{N}_7\text{O}_3\text{ClF}_2$ ($\text{M}+\text{H}$) $^+$: m/z = 718.2; found 718.2.

Step 5: 3-(8-amino-2-(amino(2,6-difluorophenyl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(chloro(2,6-difluorophenyl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (40 mg, 0.084 mmol) in 1 mL of DMSO was added ammonia solution (1 mL). The mixture was heated with microwave condition at 100 $^\circ\text{C}$ for 10 h before diluted with water and extracted with EtOAc. The combined organic layers were washed with water and brine, dried over MgSO_4 , and concentrated. The resulting residue was dissolved in TFA (1 mL), and stirred at 80 $^\circ\text{C}$ for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aq. NaHCO_3 solution. The crude material was directly purified by a silica gel column


to afford the desired product as a racemic mixture. The product was then separated with chiral HPLC using a chiral column (AM-1) and 45% EtOH in hexanes (20 mL/min) solvent system. Peak 1 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₇F₂N₈O (M+H)⁺: m/z = 459.1; found 459.0.

Example 78. 3-(8-amino-2-((2,6-difluorophenyl)(methylamino)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



The title compound was prepared using similar procedures as described for Example 77 with methyl amine (2M THF solution) replacing ammonia solution in step 5. In addition, the replacement reaction was conducted at 70 °C for 1h with 2 equivalents of K₂CO₃. After deprotection, the racemic product was separated with chiral HPLC using a chiral column (C2) and 30% EtOH in hexanes (20 mL/min) solvent system. Peak 2 was isolated, and further purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₄H₁₉F₂N₈O (M+H)⁺: m/z = 473.2; found 473.2.


Example 79. 3-(8-amino-2-((2,6-difluorophenyl)((2-hydroxyethyl)amino)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



The title compound was prepared using similar procedures as described for Example 77 with 2-aminoethan-1-ol replacing ammonia solution in step 5. In addition, the replacement reaction was conducted at 70 °C for 1h with 2 equivalents of K₂CO₃. After deprotection, the reaction mixture was purified by prep-HPLC (pH = 2,


acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{25}H_{21}N_8O_2F_2$ ($M+H$)⁺: m/z = 503.2; found 503.2.

Example 80. 3-(8-amino-2-(amino(2-fluorophenyl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for Example 77 with 1-fluoro-2-iodobenzene replacing 1,3-difluoro-2-iodobenzene in step 2. The final reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{23}H_{18}N_8OF$ ($M+H$)⁺: m/z = 441.1; found 441.2.

Example 81. 3-(8-amino-2-(amino(2,6-difluorophenyl)methyl)-5-(1-methyl-6-oxo-1,6-dihydropyridazin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for Example 77 with 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridazin-3(2H)-one replacing 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole in step 3. The final reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{24}H_{18}N_9OF_2$ ($M+H$)⁺: m/z = 486.1; found 486.2.

Example 82. 3-(8-amino-2-((3-(oxazol-5-yl)pyridin-2-yl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




The title compound was prepared using similar procedures as described for Example 122 with 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole replacing 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole in step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{25}H_{17}N_{10}O$ ($M+H$)⁺: m/z = 473.1; found 473.2.

Example 83. 3-(8-amino-2-(2-fluoro-6-(1-methyl-1H-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (13 mg, 0.022 mmol) (from Example 99, step 4), 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (9.5 mg, 0.044 mmol), cesium carbonate (17.7 mg, 0.116 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (2.26 mg, 2.88 μ mol) (XPhos Pd G2) in 1,4-dioxane (500 μ l) and water (100 μ l) was purged with N_2 and heated at 95 $^{\circ}C$ for 1 h. The mixture was concentrated and dissolved in TFA (1 mL) and stirred at 100 $^{\circ}C$ for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and purified via preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{23}H_{17}F_2N_8$ ($M+H$)⁺: m/z = 443.1; found 443.2.

Example 84. (S)-3-(8-amino-2-(2-fluoro-6-(((1-methyl-2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




- 5 *Step 1: 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-fluoro-6-vinylbenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile*



- A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (Example 99, step 4; 70 mg, 0.11 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (34 mg, 0.22 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (8.5 mg, 10.8 μ mol) and K_3PO_4 (47 mg, 0.22 mmol) in 1,4-dioxane (2 mL)/water (0.4 mL) was stirred at 110 °C for 1 h. The reaction mixture was concentrated under vacuum and the resulting residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for $C_{37}H_{31}F_2N_6O_2$ ($M+H$)⁺: m/z = 629.2; found 629.3.

Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-fluoro-6-formylbenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile



20


To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-fluoro-6-vinylbenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (56 mg, 0.089 mmol) in THF (1 mL) and water (1 mL) was added 0.157 M osmium tetroxide in

water (0.02 mmol). After 2 min, sodium metaperiodate (86 mg, 0.4 mmol) was added. The reaction mixture was heated at 60 °C for 1 h before quenched with sat. Na₂S₂O₃. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated to afford the product as a light yellow oil. LC-MS calculated for C₃₆H₂₉F₂N₆O₃ (M+H)⁺: m/z = 631.2; found 631.1.

Step 3: (S)-3-(8-amino-2-(2-fluoro-6-(((1-methyl-2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile


To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-fluoro-6-formylbenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (10 mg, 0.02 mmol) in DCM (0.5 mL) was added (S)-3-amino-1-methylpyrrolidin-2-one (4.6 mg, 0.04 mmol) then acetic acid (4 µL, 0.08 mmol). After 1h, sodium triacetoxyborohydride (8.5 mg, 0.04 mmol) was added to the reaction. The reaction was stirred overnight and the mixture was concentrated, then 0.5 mL of TFA was added to the mixture and the mixture was heated at 100 °C for 10 min. The reaction was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₅H₂₃N₈OF₂ (M+H)⁺: 489.2 found 489.2.

Example 85. 3-(8-amino-2-(2-fluoro-6-((6-methyl-5-oxo-2,6-diazaspiro[3.4]octan-2-yl)methyl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




The title compound was prepared using similar procedures as described for Example 84 with 6-methyl-2,6-diazaspiro[3.4]octan-5-one replacing (S)-3-amino-1-methylpyrrolidin-2-one in step 3. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₇H₂₅N₈OF₂ (M+H)⁺: m/z = 515.2; found 515.2.

Example 86. 3-(8-amino-5-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




- 5 *Step 1: 6-bromo-N,N-bis(4-methoxybenzyl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine*



To a flask charged with 2-(pyridin-2-yl)acetic acid (164 mg, 1.2 mmol), HATU (708 mg, 1.9 mmol) in CH₂Cl₂ (10 ml) was added 1,2-diamino-3-(bis(4-methoxybenzyl)amino)-5-bromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate (400 mg, 0.62 mmol), followed by DIEA (0.65 ml, 3.72 mmol). After stirring at room temperature for 6h, LCMS showed completion of reaction. The reaction mixture was diluted with DCM and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude product was purified with flash chromatography to give the desired product. LC-MS calculated for C₂₇H₂₆BrN₆O₂ (M+H)⁺: m/z = 545.1; found 545.2.

Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




20

A flask charged with 6-bromo-N,N-bis(4-methoxybenzyl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine (108 mg, 0.2 mmol), (3-cyano-2-fluorophenyl)boronic acid (49.7 mg, 0.35 mmol), Cs₂CO₃ (134 mg, 0.41 mmol), Pd-

tetrakis (23 mg, 0.02 mmol), 1,4-dioxane (2 ml) and water (0.2 ml) was evacuated under vacuum and refilled with N₂ (repeated three times). The mixture was heated at 100 °C for 8h. LCMS showed total completion of reaction. The reaction mixture was diluted with DCM and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude product was purified with flash chromatography to give the desired product. LC-MS calculated for C₃₄H₂₉FN₇O₂ (M+H)⁺: m/z = 586.2; found 586.2.

Step 3: 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (146 mg, 0.25 mmol) in 3 mL of dichloromethane, 1-bromopyrrolidine-2,5-dione (46 mg, 0.25 mmol) was added at 0 °C. The resulting mixture was stirred for 16 h before concentrated and purified by silica gel column to afford the desired product. LC-MS calculated for C₃₄H₂₈FBrN₇O₂ (M+H)⁺: m/z = 664.1; found 664.2.

Step 4: 3-(8-amino-5-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile

A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (15 mg, 0.022 mmol), 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (Example 76, step1; 9.1 mg, 0.044 mmol), cesium carbonate (17.7 mg, 0.116 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (2.26 mg, 2.88 μmol) (XPhos Pd G2) in 1,4-dioxane (500 μl) and water (100 μl) was purged with N₂ and heated at 95 °C for 1 h. The mixture was concentrated, then 0.5 mL of TFA was added to the mixture and the mixture was heated at 100 °C for 10 min. The reaction was concentrated and purified by


preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for $C_{22}H_{16}N_8OF$ ($M+H$)⁺: 427.1; found 427.1.

Example 87. 3-(8-amino-5-(4-(2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




The title compound was prepared using similar procedures as described for Example 86 with 4-(1-((tert-butyldimethylsilyl)oxy)-2,2-difluoroethyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (Example 35, step 3) replacing 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole in step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{24}H_{18}N_8O_2F_3$ ($M+H$)⁺: m/z = 507.1; found 507.2.

Example 88. 3-(8-amino-5-(2-methyl-4-(2,2,2-trifluoro-1-hydroxyethyl)oxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




Step 1: 2,2,2-trifluoro-1-(2-methyloxazol-4-yl)ethan-1-ol



To a solution of 2-methyloxazole-4-carbaldehyde (300 mg, 2.70 mmol) in dry THF (10 ml) was added trimethyl(trifluoromethyl)silane (797 μ l, 5.40 mmol)


dropwise followed by adding CsF (820 mg, 5.40 mmol). After stirring at rt for 30 min, TBAF (1M THF solution, 2.70 mmol) was added, and the reaction mixture was stirred further for 10 min before quenched with sat. NH_4Cl . The mixture was extracted twice with EtOAc, The combined organic layers were washed with sat. NaCl solution, dried over Na_2SO_4 , filtered, and concentrated to afford the crude product as a brown oil, which was used directly in next step without further purification. LC-MS calculated for $\text{C}_6\text{H}_7\text{F}_3\text{NO}_2$ ($\text{M}+\text{H}$)⁺: m/z = 182.0; found 182.0.

Step 2: 4-(1-((tert-butyldimethylsilyl)oxy)-2,2,2-trifluoroethyl)-2-methyloxazole



In a flame dried round-bottomed flask equipped with a magnetic stir bar, a solution of 2,2,2-trifluoro-1-(2-methyloxazol-4-yl)ethan-1-ol (1.24 g, 6.87 mmol) in DCM (10 mL) was treated at rt with tert-butylchlorodimethylsilane (1.13 g, 6.88 mmol) followed by imidazole (0.47 g, 6.87 mmol) and the resulting suspension was stirred for 1 h at rt. After completion, water was added to quench the reaction. The mixture was then extracted with EtOAc, the organic layers were dried over Na_2SO_4 , filtered and the solvent was removed under reduced pressure. The residue was purified with flash chromatography to give the desired product. LC-MS calculated for $\text{C}_{12}\text{H}_{21}\text{F}_3\text{NO}_2\text{Si}$ ($\text{M}+\text{H}$)⁺: m/z = 296.1; found 296.1.

Step 3: 4-(1-((tert-butyldimethylsilyl)oxy)-2,2,2-trifluoroethyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole



In a flame dried round-bottomed flask equipped with a magnetic stir bar, was charged with (1,5-cyclooctadiene)(methoxy)iridium(I) dimer (60 mg, 0.09 mmol), 4,4,5,5-tetramethyl-1,3,2-dioxaborolane (834 mg, 3.0 mmol), and pentane (4.0 mL). The mixture was stirred at room temperature for 10 min. Then 4,4'-di-tert-butyl-2,2'-


dipyridyl (48 mg, 0.18 mmol) was added to this mixture and reaction stirred for additional 20 min. 4-(1-((tert-butyldimethylsilyl)oxy)-2,2,2-trifluoroethyl)-2-methyloxazole (708 mg, 2.4 mmol) dissolved in Et₂O (4 mL) was added to the active catalyst mixture. The reaction was stirred at room temperature until completion.

- 5 Solvent was removed under reduced pressure, and the crude material was purified with flash chromatography to give the desired product. LC-MS calculated for the corresponding boronic acid C₁₂H₂₂BF₃NO₄Si (M+H)⁺: m/z = 340.2; found 340.1.

10 *Step 4: 3-(8-amino-5-(2-methyl-4-(2,2,2-trifluoro-1-hydroxyethyl)oxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile*


- A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (Example 86, step 3; 15 mg, 0.022 mmol), 4-(1-((tert-butyldimethylsilyl)oxy)-2,2,2-trifluoroethyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (18 mg, 0.044 mmol),
 15 cesium carbonate (17.7 mg, 0.116 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (2.26 mg, 2.88 μmol) (XPhos Pd G2) in 1,4-dioxane (500 μl) and water (100 μl) was purged with N₂ and heated at 95 °C for 1 h. The mixture was concentrated, then 0.5 mL of TFA was added to the mixture and the mixture was heated at 100 °C for 10 min. The reaction
 20 was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₄H₁₇N₈O₂F₄ (M+H)⁺: 525.1; found 525.1.

25 **Example 89. 3-(8-amino-5-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile**




The title compound was prepared using similar procedures as described for Example 86 with 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2(1H)-one replacing 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole in step 4. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{24}H_{18}N_8OF$ ($M+H$)⁺: m/z = 453.1; found 453.2.

Example 90. 3-(8-amino-2-((1-((1-methyl-1H-imidazol-4-yl)sulfonyl)pyrrolidin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




Step 1: tert-butyl 2-((8-(bis(4-methoxybenzyl)amino)-6-bromo-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)pyrrolidine-1-carboxylate



To a solution of 1,2-diamino-3-(bis(4-methoxybenzyl)amino)-5-bromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate (1.80 g, 2.79 mmol), 2-(1-(tert-butoxycarbonyl)pyrrolidin-2-yl)acetic acid (704 mg, 3.07 mmol), 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethylammonium tetrafluoroborate (TBTU, 1.35 g, 4.19 mmol) in dichloromethane (20 mL) was added *N,N*-diisopropylethylamine (0.980 mL, 5.58 mmol) dropwise. The resultant mixture was stirred at room temp overnight. The resulting mixture was filtered, concentrated under reduced pressure, and purified by Biotage Isolera (with 50 g silica gel column) eluting with 0–50% EtOAc/Hexane to give the product. LCMS calculated for $C_{31}H_{38}BrN_6O_4$ ($M+H$)⁺: m/z = 637.2, 639.2; found: 637.3, 639.3.


Step 2: *tert*-butyl 2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)pyrrolidine-1-carboxylate



A mixture of *tert*-butyl 2-((8-(bis(4-methoxybenzyl)amino)-6-bromo-
 5 [1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)pyrrolidine-1-carboxylate (1.20 g, 1.88 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (XPhos Pd G2) (0.148 g, 0.188 mmol), sodium carbonate (0.299 g, 2.82 mmol) and (3-cyano-2-fluorophenyl)boronic acid (.310 g, 1.88 mmol) in 1,4-dioxane (17 ml)/Water (1.7 ml) in a 40 mL vial was heated at 90 °C
 10 overnight. The mixture was diluted with water and extracted with EtOAc (x3). The organic extracts were dried (anhyd. Na₂SO₄) and concentrated under reduced pressure. The residue was purified by Biotage Isolera (with 50 g silica gel column) eluting with 0–50% EtOAc/Hexane to give the product. LCMS calculated for C₃₈H₄₁FN₇O₄⁺ (M+H)⁺: m/z = 678.3; found: 678.4.

15

Step 3: 3-(8-amino-2-(pyrrolidin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile




A mixture of *tert*-butyl 2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-
 20 fluorophenyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)pyrrolidine-1-carboxylate (1.13 g, 1.67 mmol) in TFA (30 mL) was heated at 70 °C for 1 h. After cooling to room temperature, TFA was evaporated, and the residue was diluted with 1 N NaOH (200 mL). The resultant mixture was extracted with DCM (x3) and the combined organic extracts were dried (anhyd. Na₂SO₄) and concentrated under reduced pressure
 25 to afford the product using without further purification. LCMS calculated for C₁₇H₁₇FN₇⁺ (M+H)⁺: m/z = 338.1; found: 338.1.

Step 4: 3-(8-amino-2-((1-((1-methyl-1H-imidazol-4-yl)sulfonyl)pyrrolidin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile


A mixture of 3-(8-amino-2-(pyrrolidin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (14 mg, 0.043 mmol) and 1-methyl-1H-imidazole-4-sulfonyl chloride (6.0 mg, 0.033 mmol) and triethylamine (14 μ L, 0.099 mmol) in DCM (0.25 ml) was stirred at room temp for 2 h. The resultant mixture was diluted with acetonitrile, filtered, and purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to afford the desired product as a TFA salt. LCMS calculated for $C_{21}H_{21}FN_9O_2S^+$ (M+H)⁺: m/z = 482.1; found: 482.0.

10

Example 91. 3-(2-(2-((1-acetylpiperidin-4-yl)methyl)-6-fluorobenzyl)-8-amino-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile



Step 1: *tert*-butyl 4-(2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzylidene)piperidine-1-carboxylate




A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (193 mg, 0.303 mmol) (from Example 99, step 4), XPhos Pd G2 (23.84 mg, 0.030 mmol), potassium phosphate (193 mg, 0.909 mmol) and *tert*-butyl 4-((4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)methylene)piperidine-1-carboxylate (98.1 mg, 0.303 mmol) in 1,4-dioxane (2.75 mL)/Water (0.550 mL) in a 40 mL vial was flushed with nitrogen for ca. 2 min and

20


heated at 120 °C for 3 h. After cooling to room temp, the mixture was diluted with water and extracted with DCM (x3). The combined organic extracts were dried (anhyd. Na₂SO₄), concentrated under reduced pressure, and purified by Biotage Isolera (with 50 g silica gel column) eluting with 0–100% EtOAc/Hexane to give the product. LCMS calculated for C₄₆H₄₆F₂N₇O₄⁺ (M+H)⁺: m/z = 798.4; found: 798.4.

Step 2: tert-butyl 4-(2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)piperidine-1-carboxylate



A mixture of *tert*-butyl 4-(2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzylidene)piperidine-1-carboxylate (151 mg, 0.188 mmol) and Pd(OH)₂/C (20% wt, 26 mg, 0.038 mmol) in MeOH (2 mL)/DCM (1.000 mL) in a 20 mL vial was stirred at room temp under H₂ balloon overnight. The resultant mixture was filtered, concentrated under reduced pressure, and used without further purification. LCMS calculated for C₄₆H₄₈F₂N₇O₄⁺ (M+H)⁺: m/z = 800.4; found: 800.5.

Step 3: 2-fluoro-3-(2-(2-fluoro-6-(piperidin-4-ylmethyl)benzyl)-8-((4-methoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of *tert*-butyl 4-(2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)piperidine-1-carboxylate (130mg, 0.163 mmol) in DCM (10 mL) was added TFA (5.0 mL) dropwise. The resultant mixture was stirred at room temp for 30 min and transferred to a separated funnel with DCM and add 1N NaOH (ca. 200 mL) was added. The

layers were separated and the aqueous layer was extracted with DCM (x2). The combined organic extracts were dried (anhyd. Na_2SO_4), concentrated under reduced pressure, and used without further purification. LCMS calculated for $\text{C}_{33}\text{H}_{32}\text{F}_2\text{N}_7\text{O}^+$ ($\text{M}+\text{H}$) $^+$: $m/z = 580.3$; found: 580.3.


5

Step 4: 3-(2-(2-((1-acetylpiperidin-4-yl)methyl)-6-fluorobenzyl)-8-((4-methoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile

To a solution of 2-fluoro-3-(2-(2-fluoro-6-(piperidin-4-ylmethyl)benzyl)-8-((4-methoxybenzyl)amino)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (14 mg, 0.025 mmol) and acetyl chloride (25 μL , 0.025 mmol, 1M in DCM) in DCM (0.5 mL) was added triethylamine (10.5 μL , 0.0750 mmol). The resultant mixture was stirred at room temp for 1 h, filtered, and concentrated under reduced pressure. The residue was added TFA (0.5 mL) stirred 70 $^\circ\text{C}$ for 1 h. The resulting mixture was diluted with acetonitrile, filtered, and purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to afford the desired product as a TFA salt. LCMS calculated for $\text{C}_{27}\text{H}_{26}\text{F}_2\text{N}_7\text{O}^+$ ($\text{M}+\text{H}$) $^+$: $m/z = 502.2$; found: 502.1.


15

Example 92. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(1-ethyl-1H-pyrazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 1




20

Step 1: 3-(8-(bis(4-methoxybenzyl)amino)-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of 1,3-difluoro-2-iodobenzene (0.822 g, 3.42 mmol) in THF (2 mL) was added isopropylmagnesium chloride lithium chloride complex solution (1.3 M, 2.3 mL, 3.0 mmol) dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 30 minutes. A solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (0.432 g, 0.856 mmol) (from Example 40, step 1) in THF (2 mL) was added dropwise and the reaction mixture was stirred at 0 °C for 1 h. The reaction mixture was then quenched with saturated aqueous NH₄Cl solution and diluted with DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The crude residue was purified using flash chromatography to give the desired product (0.301 g, 57%). LC-MS calculated for C₃₅H₂₉F₂N₆O₃ (M+H)⁺: m/z = 619.2; found 619.2.

Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (0.301 g, 0.487 mmol) in DCM (2 mL) was added a solution of NBS (0.087 g, 0.487 mmol) in DCM (2 mL) dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 1 h. The reaction mixture was concentrated, and the resulting residue was purified using flash chromatography to give the desired product (0.339 g, 99%). LC-MS calculated for C₃₅H₂₈BrF₂N₆O₃ (M+H)⁺: m/z = 697.1; found 697.1.

Step 3: 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(1-ethyl-1*H*-pyrazol-5-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile

To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (0.210 g, 0.301 mmol), 1-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-

pyrazole (0.267 g, 1.20 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.071 g, 0.090 mmol) in dioxane (2.50 mL) and water (0.50 mL) was added potassium phosphate tribasic (0.320 g, 1.51 mmol). The reaction mixture was stirred at 100 °C for 2 h. The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The crude material was dissolved in TFA (5 mL) and heated to 80 °C for 20 minutes. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (110 mg, 77%) as a racemic mixture. The product was then separated with chiral SFC using a chiral column (ES Industries ChromegaChiral CC4) and 25% MeOH in CO₂ (85 mL/min) solvent system. Peak 1 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₄H₁₉F₂N₈O (M+H)⁺: m/z = 473.2; found 473.2.

Example 93. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(1-ethyl-1H-pyrazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 2




20

This compound was prepared using the same procedure as described for Example 92. The product was separated with chiral SFC using a chiral column (ES Industries ChromegaChiral CC4) and 25% MeOH in CO₂ (85 mL/min) solvent system. Peak 2 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₄H₁₉F₂N₈O (M+H)⁺: m/z = 473.2; found 473.2.


25

Example 94. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(2,6-dimethylpyridin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (0.518 g, 0.638 mmol) (from example 92, step 2), 2,6-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (0.346 g, 1.48 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.058 g, 0.074 mmol) in dioxane (3.0 mL) and water (0.60 mL) was added potassium phosphate tribasic (0.472 g, 2.23 mmol). The reaction mixture was stirred at 90 °C for 1 h. The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The crude material was dissolved in TFA (5 mL) and heated to 80 °C for 20 minutes. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (257 mg, 72%) as a racemic mixture. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5μm Cellulose-2, 21.1x250mm) and 35% EtOH in Hexanes (20 mL/min) solvent system. Peak 1 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₆H₂₀F₂N₇O (M+H)⁺: m/z = 484.2; found 484.2. ¹H NMR (500 MHz, DMSO-*d*₆) δ 7.92 (s, 2H), 7.85 (s, 1H), 7.83 (d, *J* = 7.6 Hz, 1H), 7.56 (d, *J* = 8.0 Hz, 1H), 7.53 – 7.40 (m, 4H), 7.10 (t, *J* = 8.4 Hz, 2H), 6.27 (s, 1H), 2.51 (s, 6H).

Example 95. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(2,6-dimethylpyridin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 2




This compound was prepared using the same procedure as described for
 5 Example 94. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-2, 21.1x250mm) and 35% EtOH in Hexanes (20 mL/min) solvent system. Peak 2 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{26}H_{20}F_2N_7O$ ($M+H$)⁺: m/z = 484.2; found 484.2. ¹H NMR (500 MHz, DMSO-*d*₆) δ 7.92 (s, 2H), 7.85 (s, 1H), 7.83 (d, J = 7.6 Hz, 1H), 7.56 (d, J = 8.0 Hz, 1H), 7.53 – 7.40 (m, 4H), 7.10 (t, J = 8.4 Hz, 2H), 6.27 (s, 1H), 2.51 (s, 6H).

Example 96. 3-(2-((1H-pyrrolo[3,2-b]pyridin-3-yl)methyl)-4-amino-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile



15

Step 1: 3-(4-amino-2-(bromomethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(4-amino-2-(hydroxymethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (0.48 g, 1.40 mmol) (from Example 13, step 2) in dry THF (10 ml) was added PBr₃ (1.14 g, 4.19 mmol) dropwise at room temperature. The reaction mixture was stirred rapidly at 60 °C for 5 hours. The
 5 reaction mixture was cooled and quenched with saturated aqueous NaHCO₃ solution. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The crude material was directly purified by a silica gel column (0 to 100% ethyl acetate/hexanes) to afford the desired product (0.48 g, 84%). LC-MS calculated for
 10 C₁₈H₁₃BrN₇ (M+H)⁺: m/z = 406.0; found 406.1.

*Step 2: 3-(2-((1H-pyrrolo[3,2-*b*]pyridin-3-yl)methyl)-4-amino-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*

To a solution of 3-(4-amino-2-(bromomethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (0.015 g, 0.037 mmol), 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-tosyl-1H-pyrrolo[3,2-*b*]pyridine (0.029 g, 0.074 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.006 g, 0.007 mmol) in dioxane (0.3 mL) and water (0.06 mL) was added potassium phosphate tribasic (0.024 g, 0.111 mmol). The reaction
 20 mixture was stirred at 90 °C for 1 h. The reaction mixture was then diluted with water (0.3 ml) and THF (0.3 ml). NaOH (25 mg) was added to the vial and the vial was stirred at room temperature for 30 minutes. The reaction mixture was diluted with DMF (4 ml) and purified with prep-LCMS (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for C₂₅H₁₈N₉ (M+H)⁺: m/z =
 25 444.2; found 444.2.

Example 97. 3-(8-amino-2-((2-(difluoromethoxy)-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1




5 To a solution of 1-(difluoromethoxy)-3-fluoro-2-iodobenzene (74 mg, 0.26 mmol) in tetrahydrofuran (0.2 mL), isopropylmagnesium chloride lithium chloride (0.2 mL, 1.3 M solution) was added at -10 °C, and the resulting mixture was stirred for 1 hour before a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (20 mg, 0.034 mmol) (from

10 Example 49 step 8) in THF (0.2 mL) was added at -10 °C. The reaction mixture was stirred for 30 min, quenched with ammonium chloride solution (1 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The resulting material was dissolved in TFA (1 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and

15 basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (16 mg, 94%) as a racemic mixture. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5μm Cellulose-4, 21.1x250mm) and 45% EtOH in Hexanes (20 mL/min) solvent system. Peak 1 was isolated, and further purified using


20 preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₄H₁₆F₃N₈O₂ (M+H)⁺: m/z = 505.1; found 505.1.

Example 98. 3-(8-amino-2-((2-(difluoromethoxy)-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 2




5 This compound was prepared using the same procedure as described for Example 97. The racemic product was separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.1x250mm) and 45% EtOH in Hexanes (20 mL/min) solvent system. Peak 2 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product
10 as a TFA salt. LC-MS calculated for $C_{24}H_{16}F_3N_8O_2$ ($M+H$)⁺: m/z = 505.1; found 505.1.

Example 99. 3-(8-amino-2-(2-fluoro-6-(1-((1-methyl-1*H*-pyrazol-4-yl)methyl)-1*H*-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile



15


*Step 1: 6-bromo-*N*²,*N*²-bis(4-methoxybenzyl)pyrazine-2,3-diamine*



To a stirred suspension of 3,5-dibromopyrazin-2-amine (6 g, 23.25 mmol) and
20 bis(4-methoxybenzyl)amine (6.72 g, 25.6 mmol) in *n*-butanol (23.25 ml) at room temperature was added *N,N*-diisopropylethylamine (8.18 ml, 46.5 mmol). The reaction mixture was heated at 120 °C for 72 hours. The reaction mixture was cooled to room temperature. The resulting slurry was stirred at room temperature for 1 hour.


The suspension was filtered to remove the excess 3,5-dibromopyrazin-2-amine. The filtrate was concentrated *in vacuo*. The residue was purified by Biotage Isolera (with 330 g silica gel column) eluting with 0 - 50% EtOAc/Hexane to give the product as a brown very viscous oil (6.282 g, 77% yield). LC-MS calculated for $C_{20}H_{22}BrN_4O_2$ (M+H)⁺: m/z = 429.1; found 429.4.

Step 2: 1,2-diamino-3-(bis(4-methoxybenzyl)amino)-5-bromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate



To a solution of *O*-(mesitylsulfonyl)hydroxylamine (8.88 g, 41.3 mmol) (from Example 14 step 2) in dichloromethane (300 ml) was added 6-bromo-*N*²,*N*²-bis(4-methoxybenzyl)pyrazine-2,3-diamine (16.1 g, 37.5 mmol). The resulting solution was stirred at room temperature overnight. The mixture was concentrated and purified with a silica gel column (eluting with a gradient of 0-100% ethyl acetate in hexanes then 0-20% methanol in DCM) to give the desired product (16 g, 66%). LC-MS calculated for $C_{20}H_{23}BrN_5O_2$ (M-C₉H₁₁O₃S)⁺: m/z = 444.1; found 444.1.

Step 3: 6-bromo-2-(2-chloro-6-fluorobenzyl)-*N,N*-bis(4-methoxybenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-8-amine




To a solution of 1,2-diamino-3-(bis(4-methoxybenzyl)amino)-5-bromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate (2.5 g, 3.88 mmol), *N,N*-diisopropylethylamine (7.52 g, 58.2 mmol), and 2-(2-chloro-6-fluorophenyl)acetic acid (2.93 g, 15.5 mmol) in DMF (20 ml) was added (3-dimethylaminopropyl)-*N*'-ethylcarbodiimide hydrochloride (0.36 g, 1.86 mmol). The reaction mixture was stirred at room temperature for 1 hour. The solvent was removed under reduced pressure and the crude residue was diluted with water and dichloromethane. The layers were separated and the aqueous layer extracted with dichloromethane. The

combined organic fractions were filtered over a plug of magnesium sulfate and concentrated. Purification by automatic flash column chromatography afforded the desired product (1.34 g, 58%). LC-MS calculated for $C_{28}H_{25}BrClFN_5O_2$ ($M+H$)⁺: m/z = 596.1; found 596.1.

5

Step 4: 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




To a solution of 6-bromo-2-(2-chloro-6-fluorobenzyl)-*N,N*-bis(4-methoxybenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-8-amine (1.34 g, 2.25 mmol), 2-fluoro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzonitrile (0.72 g, 2.92 mmol), potassium phosphate tribasic (1.43 g, 6.73 mmol) in dioxane (10 ml) and water (2 ml) was added tetrakis(triphenylphosphine)palladium(0) (0.39 g, 0.38 mmol). The reaction mixture was sparged with nitrogen gas for five minutes, sealed and heated to 90 °C for 1 hour. The reaction mixture was cooled to room temperature, the solvent was removed under reduced pressure, and the crude residue was diluted with water and dichloromethane. The layers were separated and the aqueous layer extracted with dichloromethane. The combined organic fractions were filtered over a plug of magnesium sulfate and concentrated. Purification by automatic flash column chromatography afforded the desired product (0.886 g, 62%). LC-MS calculated for $C_{35}H_{28}ClF_2N_6O_2$ ($M+H$)⁺: m/z = 637.2; found 637.2.

15

20

Step 5: 1-methyl-4-((4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazol-1-yl)methyl)-1H-pyrazole




25

A vial was charged with 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (0.5 g, 2.58 mmol), 4-(bromomethyl)-1-methyl-1H-pyrazole hydrobromide (0.660 g, 2.58 mmol), cesium carbonate (2.52 g, 7.73 mmol), and DMF (6.44 ml). The reaction mixture was stirred at 60 °C for one hour. The solvent was stripped and the crude residue was diluted with water and dichloromethane. The layers were separated and the aqueous layer extracted with dichloromethane. The combined organic fractions were filtered over a plug of magnesium sulfate and concentrated. The crude material was used in the next step without further purification (0.74 g, 99%). LC-MS calculated for C₁₄H₂₂BN₄O₂ (M+H)⁺: m/z = 289.2; found 289.1.

Step 6: 3-(8-amino-2-(2-fluoro-6-(1-((1-methyl-1H-pyrazol-4-yl)methyl)-1H-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile


To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (0.010 g, 0.016 mmol), 1-methyl-4-((4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazol-1-yl)methyl)-1H-pyrazole (0.009 g, 0.031 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.002 g, 0.003 mmol) in dioxane (0.3 mL) and water (0.06 mL) was added potassium phosphate tribasic (0.010 g, 0.047 mmol). The reaction mixture was stirred at 90 °C for 1 h. The reaction mixture was then diluted with water and dichloromethane. The layers were separated and the aqueous layer extracted with dichloromethane. The combined organic fractions were concentrated under vacuum. The resulting material was dissolved in TFA (1 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room temperature, diluted with DMF (4 mL) and purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₇H₂₁F₂N₁₀ (M+H)⁺: m/z = 523.2; found 523.2.

Example 100. 3-(8-amino-2-((2-((dimethylamino)methyl)-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1




5

Step 1: 1-(bromomethyl)-3-fluoro-2-iodobenzene



A round bottom flask was charged with (3-fluoro-2-iodophenyl)methanol (0.445 g, 1.766 mmol), carbon tetrabromide (0.703 g, 2.119 mmol), triphenylphosphine (0.556 g, 2.119 mmol), *N,N*-diisopropylethylamine (0.617 ml, 3.53 mmol), and dichloromethane (17.66 ml). The reaction mixture was stirred overnight at room temperature. The solvent was stripped and the crude residue was purified by automatic flash column chromatography to afford the desired product (0.554 g, 99%).

*Step 2: 1-(3-fluoro-2-iodophenyl)-*N,N*-dimethylmethanamine*




A round bottom flask was charged with 1-(bromomethyl)-3-fluoro-2-iodobenzene (0.554 g, 1.77 mmol), dichloromethane (17 ml), and dimethylamine solution (4.41 ml, 2M in ethanol). The reaction mixture was stirred at room temperature for 1 hour. The solvent was stripped and the crude residue was purified by automatic flash column chromatography to afford the desired product (0.176 g, 36%). LC-MS calculated for C₉H₁₂FIN (M+H)⁺: m/z = 280.0; found 280.1.

Step 3: 3-(8-amino-2-((2-((dimethylamino)methyl)-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 1

To a solution of 1-(3-fluoro-2-iodophenyl)-*N,N*-dimethylmethanamine (0.179 g, 0.642 mmol) in tetrahydrofuran (1.5 mL), isopropylmagnesium chloride lithium chloride (0.726 mL, 1.3 M solution) was added at -10 °C, and the resulting mixture was stirred for 1 h before a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.220 mg, 0.378 mmol) (from Example 49 step 8) in THF (1.5 mL) was added at -10 °C. The reaction mixture was stirred for 60 min, then quenched with ammonium chloride solution (3 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The resulting material was dissolved in TFA (3 mL), and stirred at 80 °C for 20 min. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (87 mg, 47%) as a racemic mixture. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5μm Cellulose-1, 21.1x250mm) and 30% EtOH in Hexanes (20 mL/min) solvent system. Peak 1 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₆H₂₃FN₉O (M+H)⁺: m/z = 496.2; found 496.2.


Example 101. 3-(8-amino-2-((2-((dimethylamino)methyl)-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 2



This compound was prepared using the same procedure as described for Example 100. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5μm Cellulose-1, 21.1x250mm) and 30% EtOH in


Hexanes (20 mL/min) solvent system. Peak 2 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{26}H_{23}FN_9O$ (M+H)⁺: m/z = 496.2; found 496.2.

5 **Example 102. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(3-methyl-1H-pyrazol-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 1**




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.43 g, 0.062 mmol) (from example 92, step 2), *tert*-butyl 3-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole-1-carboxylate (0.076 g, 0.246 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.010 g, 0.012 mmol) in dioxane (0.5 mL) and water (0.1 mL) was added potassium phosphate tribasic (0.065 g, 0.308 mmol). The reaction mixture was stirred at 100 °C for 1 hour. The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The crude material was dissolved in TFA (2 mL) and heated to 80 °C for 20 minutes. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (8 mg, 29%) as a racemic mixture. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.1x250mm) and 40% EtOH in Hexanes (20 mL/min) solvent system. Peak 1 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{23}H_{17}F_2N_8O$ (M+H)⁺: m/z = 459.1; found 459.2.

Example 103. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(3-methyl-1H-pyrazol-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 2



This compound was prepared using the same procedure as described for Example 102. The product was separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.1x250mm) and 40% EtOH in Hexanes (20 mL/min) solvent system. Peak 2 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₇F₂N₈O (M+H)⁺: m/z = 459.1; found 459.2.


Example 104. 3-(5-([1,2,4]triazolo[4,3-a]pyridin-6-yl)-8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.015 g, 0.022 mmol) (from example 92, step 2), 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,2,4]triazolo[4,3-a]pyridine (0.021 g, 0.086 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.005 g, 0.006 mmol) in dioxane (0.3 mL) and water (0.06 mL) was added potassium phosphate tribasic (0.023 g, 0.108 mmol). The reaction mixture was stirred at 100 °C for 1 hour. The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and


concentrated. The crude material was dissolved in TFA (1 mL) and heated to 80 °C for 20 minutes. The reaction mixture was then cooled to room temperature, diluted with DMF (4 ml), and purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired racemic product as a TFA salt. LC-MS calculated for $C_{25}H_{16}F_2N_9O$ (M+H)⁺: m/z = 496.1; found 496.1

Example 105. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(oxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 1



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.43 g, 0.062 mmol) (from example 92, step 2), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (0.048 g, 0.246 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.010 g, 0.012 mmol) in dioxane (0.5 mL) and water (0.1 mL) was added potassium phosphate tribasic (0.065 g, 0.308 mmol). The reaction mixture was stirred at 100 °C for 1 hour. The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The crude material was dissolved in TFA (2 mL) and heated to 80 °C for 20 minutes. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (6 mg, 22%) as a racemic mixture. The product was then separated with chiral SFC using a chiral column (ES Industries CC4 5um 20x250 mm) and 35% MeOH in CO₂ (65 mL/min) solvent system. Peak 1 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{22}H_{14}F_2N_7O_2$ (M+H)⁺: m/z = 446.1; found 446.1.


Example 106. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(oxazol-5-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 2



This compound was prepared using the same procedure as described for
 5 Example 105. The product was then separated with chiral SFC using a chiral column (ES Industries CC4 5um 20x250 mm) and 35% MeOH in CO₂ (65 mL/min) solvent system. Peak 2 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₂H₁₄F₂N₇O₂ (M+H)⁺: m/z = 446.1; found 446.1.

10

Example 107. 3-(8-amino-2-(2-fluoro-6-(1-(pyridin-2-ylmethyl)-1*H*-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-
 15 fluorobenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile (0.01 g, 0.016 mmol) (from Example 99, step 4), 2-((4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazol-1-yl)methyl)pyridine (0.009 g, 0.031 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.002 g, 0.003 mmol) in dioxane (0.3 mL) and water (0.06 mL) was added potassium
 20 phosphate tribasic (0.010 g, 0.047 mmol). The reaction mixture was stirred at 90 °C for 1 hour. The reaction mixture was then diluted with water and dichloromethane. The layers were separated and the aqueous layer extracted with dichloromethane. The combined organic fractions were concentrated under vacuum. The resulting material was dissolved in TFA (1 mL), and stirred at 80 °C for 20 min. The reaction mixture
 25 was then cooled to room temperature, diluted with DMF (4 mL) and purified using

preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{28}H_{20}F_2N_9$ (M+H)⁺: m/z = 520.2; found 520.1.


Example 108. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(3-

5 methylpyridin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 1



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.43 g, 0.062 mmol) (from example 92, step 2), 3-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (0.054 g, 0.246 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.010 g, 0.012 mmol) in dioxane (0.5 mL) and water (0.1 mL) was added potassium phosphate tribasic (0.065 g, 0.308 mmol). The reaction mixture was stirred at 100 °C for 1 hour. The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over $MgSO_4$, filtered and concentrated. The crude material was dissolved in TFA (2 mL) and heated to 80 °C for 20 minutes. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding aqueous $NaHCO_3$ solution. The crude material was directly purified by a silica gel column to afford the desired product (8 mg, 28%) as a racemic mixture. The product was then separated with chiral SFC using a chiral column (ES Industries CC4 5um 20x250 mm) and 35% MeOH in CO_2 (65 mL/min) solvent system. Peak 1 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for $C_{25}H_{18}F_2N_7O$ (M+H)⁺: m/z = 470.2; found 470.2.


Example 109. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(3-methylpyridin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 2



This compound was prepared using the same procedure as described for
 5 Example 108. The product was then separated with chiral SFC using a chiral column (ES Industries CC4 5um 20x250 mm) and 35% MeOH in CO₂ (65 mL/min) solvent system. Peak 2 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₅H₁₈F₂N₇O (M+H)⁺: m/z = 470.2; found 470.2.


10

Example 110. 3-(8-amino-2-((2-fluoro-6-(pyrrolidin-1-ylmethyl)phenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile, Peak 1




15

*Step 1: 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-chloro-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*




To a solution of 1-chloro-3-fluoro-2-iodobenzene (0.335 g, 1.30 mmol) in tetrahydrofuran (1.5 mL), isopropylmagnesium chloride lithium chloride (0.878 mL, 1.3 M solution) was added at -10 °C, and the resulting mixture was stirred for 1 hour before a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (0.190 g, 0.326 mmol) (from Example 5
49 step 8) in THF (1.5 mL) was added at -10 °C. The reaction mixture was stirred for 60 minutes, then quenched with ammonium chloride solution (3 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The crude material was directly purified by a silica gel column to afford the
10 desired product (147 mg, 63%) as a racemic mixture. LC-MS calculated for $C_{39}H_{31}ClFN_8O_3$ (M+H)⁺: m/z = 713.2; found 713.3.

*Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-fluoro-6-vinylphenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile*
15



To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-chloro-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile (.212 g, 0.297 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (0.055 g, 0.357 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.023 g, 0.030 mmol) in dioxane (2.5 mL) and water (0.5 mL) was added potassium phosphate tribasic (0.126 g, 0.595 mmol). The reaction mixture was stirred at 100 °C for 1 h. The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was
20 extracted with DCM, and the combined organic fractions were dried over $MgSO_4$, filtered and concentrated. The crude material was directly purified by a silica gel column to afford the desired product (0.195 mg, 93%). LC-MS calculated for $C_{41}H_{34}FN_8O_3$ (M+H)⁺: m/z = 705.3; found 705.4.

Step 3: 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-fluoro-6-formylphenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



5 A vial was charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-fluoro-6-vinylphenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (.195 g, 0.277 mmol), THF (2.5 ml), water (2.5 ml), sodium periodate (0.266 g, 1.245 mmol), and osmium tetroxide solution (0.176 ml, 4% in water). The mixture was stirred at room temperature for 30 minutes. The reaction mixture was
10 diluted with water and dichloromethane. The layers were separated and the aqueous layer extracted with dichloromethane. The combined organic fractions were filtered over a plug of magnesium sulfate and purified by automatic flash column chromatography to afford the product (0.135 g, 69%). LC-MS calculated for $C_{40}H_{32}FN_8O_4$ ($M+H$)⁺: m/z = 707.2; found 707.3.

15


Step 4: 3-(8-amino-2-((2-fluoro-6-(pyrrolidin-1-ylmethyl)phenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile, Peak 1

A vial was charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-fluoro-6-formylphenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (.013 g, 0.018 mmol), pyrrolidine (0.013 g, 0.178 mmol), acetic acid (2.041 μ l, 0.036 mmol), dichloromethane (0.4 ml), and sodium borohydride (1.3 mg, 0.036 mmol). The reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was quenched with saturated aqueous sodium bicarbonate solution. The solution was diluted with dichloromethane and the layers were separated. The
20 aqueous layer was extracted with dichloromethane and the combined organic fractions were filtered over a plug of magnesium sulfate and concentrated. The crude residue was dissolved into 1 mL TFA and stirred at 80 °C for 20 minutes. The reaction mixture was then cooled to room temperature, concentrated, and basified by adding

aqueous NaHCO₃ solution. The crude material was directly purified by a silica gel column to afford the desired product (6 mg, 65%) as a racemic mixture. The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-1, 21.2x250mm) and 30% EtOH in hexanes (20 mL/min) solvent system.


- 5 Peak 1 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₈H₂₅FN₉O (M+H)⁺: m/z = 522.2; found 522.2.

**Example 111. 3-(8-amino-2-((2-fluoro-6-(pyrrolidin-1-ylmethyl)phenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-
10 a]pyrazin-6-yl)benzonitrile, Peak 2**



- This compound was prepared using the same procedure as described for Example 110. The product was then separated with chiral HPLC using a chiral
15 column (Phenomenex Lux 5um Cellulose-1, 21.2x250mm) and 30% EtOH in hexanes (20 mL/min) solvent system. Peak 2 was isolated, and further purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₈H₂₅FN₉O (M+H)⁺: m/z = 522.2; found 522.2.


Example 112. 3-(8-amino-2-(2-fluoro-6-(1-(2-(methylsulfonyl)ethyl)-1H-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




- To a solution 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (.075 g, 0.118 mmol) (from
25 Example 99, step 4), 1-(2-(methylsulfonyl)ethyl)-4-(4,4,5,5-tetramethyl-1,3-dioxolan-

2-yl)-1*H*-pyrazole (0.071 g, 0.235 mmol), and dicyclohexyl(2',4',6'-triiisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.018 g, 0.024 mmol) in dioxane (1.0 mL) and water (0.2 mL) was added potassium phosphate tribasic (0.075 g, 0.353 mmol). The reaction mixture was stirred at 100 °C for 1 h. The reaction mixture was then diluted with water and dichloromethane. The layers were separated and the aqueous layer extracted with dichloromethane. The combined organic fractions were concentrated under vacuum. The resulting material was dissolved in TFA (2 mL), and stirred at 80 °C for 20 minutes. The reaction mixture was then cooled to room temperature, diluted with DMF (3 ml) and purified using preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₅H₂₁F₂N₈O₂S (M+H)⁺: m/z = 535.1; found 535.1.

Example 113. 3-(8-amino-2-((2-fluoro-6-((6-oxohexahydropyrrolo[1,2-*a*]pyrazin-2(1*H*)-yl)methyl)phenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile



*Step 1: 6-bromo-N,N-bis(4-methoxybenzyl)-2-vinyl-[1,2,4]triazolo[1,5-*a*]pyrazin-8-amine*




20

To a solution of 1,2-diamino-3-(bis(4-methoxybenzyl)amino)-5-bromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate (1.00 g, 1.55 mmol) (from Example 99, step 2) and acryloyl chloride (0.253 ml, 3.10 mmol) in DMF (4 ml) and dichloromethane (4 ml) at 0 °C was added triethylamine (0.540 ml, 3.88 mmol). The reaction mixture was stirred at 0 °C for for 2 hours. The solvent was removed under reduced pressure and the crude residue was purified by automatic flash column

25


chromatography to afford the desired product (0.268 g, 36%). LC-MS calculated for $C_{23}H_{23}BrN_5O_2$ (M+H)⁺: m/z = 480.1; found 480.1.

Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-2-vinyl-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-
2-fluorobenzonitrile



To a solution of 6-bromo-*N,N*-bis(4-methoxybenzyl)-2-vinyl-[1,2,4]triazolo[1,5-*a*]pyrazin-8-amine (0.070 g, 0.146 mmol), (3-cyano-2-fluorophenyl)boronic acid (0.048 g, 0.291 mmol), cesium carbonate (0.142 g, 0.437 mmol) in dioxane (1.3 ml) and water (0.15 ml) was added tetrakis(triphenylphosphine)palladium(0) (0.034 g, 0.029 mmol). The reaction mixture was sparged with nitrogen gas for five minutes, sealed and heated to 90 °C for 4 hours. The reaction mixture was cooled to room temperature, the solvent removed under reduced pressure, and the crude residue purified by automatic flash column chromatography to afford the desired product (0.061 g, 80%). LC-MS calculated for $C_{30}H_{26}FN_6O_2$ (M+H)⁺: m/z = 521.2; found 521.1.


Step 3: 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile



A vial was charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-vinyl-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile (0.287 g, 0.551 mmol), THF (2.5 ml), water (2.5 ml), sodium periodate (0.531 g, 2.481 mmol), and osmium tetroxide solution (0.433 ml, 4% in water). The mixture was stirred at room temperature overnight. The reaction mixture was adsorbed onto silica gel, and the solvent was removed under reduced pressure. The crude material (adsorbed onto silica

gel) was purified by automatic flash column chromatography to afford the product (0.127 g, 44%). LC-MS calculated for $C_{29}H_{24}FN_6O_3$ ($M+H$)⁺: m/z = 523.2; found 523.1.


- 5 *Step 4: 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-chloro-6-fluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile*



To a solution of 1-chloro-3-fluoro-2-iodobenzene (0.491 g, 1.92 mmol) in tetrahydrofuran (2.5 mL), isopropylmagnesium chloride lithium chloride (1.47 mL, 1.3 M solution) was added at -10 °C, and the resulting mixture was stirred for 1 hour before a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-formyl-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile (0.143 g, 0.274 mmol) in THF (2.5 mL) was added at -10 °C. The reaction mixture was stirred for 60 min, then quenched with ammonium chloride solution (3 mL), and extracted with dichloromethane. The combined organic layers were concentrated under vacuum. The crude material was directly purified by a silica gel column to afford the desired product (116 mg, 65%) as a racemic mixture. LC-MS calculated for $C_{35}H_{28}ClF_2N_6O_3$ ($M+H$)⁺: m/z = 653.2; found 653.1.

20

*Step 5: 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-fluoro-6-vinylphenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile*




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-chloro-6-fluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile (0.194 g, 0.297 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-

dioxaborolane (0.093 g, 0.602 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.016 g, 0.021 mmol) in dioxane (1.8 mL) and water (0.2 mL) was added potassium phosphate tribasic (0.170 g, 0.803 mmol). The reaction mixture was stirred at 100 °C for 1 hour.


The reaction mixture was then diluted with water and DCM. The layers were separated, the aqueous layer was extracted with DCM, and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The crude material was directly purified by a silica gel column to afford the desired product (0.097 mg, 75%). LC-MS calculated for C₃₇H₃₁F₂N₆O₃ (M+H)⁺: m/z = 645.2; found 645.3.

Step 6: 3-(8-(bis(4-methoxybenzyl)amino)-2-((tert-butyldimethylsilyloxy)(2-fluoro-6-vinylphenyl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile



A vial was charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-((2-fluoro-6-vinylphenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (0.178 g, 0.276 mmol), DMF (2.76 ml), imidazole (0.150 g, 2.205 mmol), and TBS-Cl (0.166 g, 1.102 mmol). The reaction mixture was stirred at room temperature for 1 hour. The solvent was removed under reduced pressure, and the crude residue was purified by automatic flash column chromatography to afford the desired product (0.166 g, 79%). For the purposes of characterization using LC-MS, the desired product was subjected to deprotection of a single PMB group; an aliquot of pure product was dissolved into 1:1 dichloromethane/trifluoroacetic acid solution (0.1 ml) and allowed to stand at room temperature for 5 minutes, furnishing 3-(2-((tert-butyldimethylsilyloxy)(2-fluoro-6-vinylphenyl)methyl)-8-(4-methoxybenzylamino)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile. LC-MS calculated for C₃₅H₃₇F₂N₆O₂Si (M+H)⁺: m/z = 639.3; found 639.3.

Step 7: 3-(8-(bis(4-methoxybenzyl)amino)-2-((tert-butyldimethylsilyloxy)(2-fluoro-6-formylphenyl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




A vial was charged 3-(8-(bis(4-methoxybenzyl)amino)-2-(((tert-
 5 butyldimethylsilyl)oxy)(2-fluoro-6-vinylphenyl)methyl)-[1,2,4]triazolo[1,5-
 a]pyrazin-6-yl)-2-fluorobenzonitrile (0.109 g, 0.144 mmol), sodium periodate (0.138
 g, 0.646 mmol), THF (0.7 ml), water (0.7 ml), and osmium tetroxide solution (0.113
 ml, 4% in water). The mixture was stirred at 45 °C overnight. The reaction mixture
 was adsorbed onto silica gel, and the solvent was removed under reduced pressure.
 10 The crude material (adsorbed onto silica gel) was purified by automatic flash column
 chromatography to afford the product (0.05 g, 46%). LC-MS calculated for
 $C_{42}H_{43}F_2N_6O_4Si$ (M+H)⁺: m/z = 761.3; found 761.3.

Step 8: 3-(8-amino-2-((2-fluoro-6-((6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-
 15 yl)methyl)phenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-
 fluorobenzonitrile


A vial was charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-(((tert-
 butyldimethylsilyl)oxy)(2-fluoro-6-formylphenyl)methyl)-[1,2,4]triazolo[1,5-
 a]pyrazin-6-yl)-2-fluorobenzonitrile (0.01 g, 0.013 mmol), hexahydropyrrolo[1,2-
 20 a]pyrazin-6(7H)-one (0.018 g, 0.131 mmol), acetic acid (1.505 µl, 0.026 mmol),
 DCM (0.202 ml), and sodium triacetoxyborohydride (5.57 mg, 0.026 mmol). The
 reaction mixture was heated to 40 °C and stirred for 2 hours. The reaction mixture
 was quenched with saturated aqueous sodium bicarbonate solution and diluted with
 DCM. The layers were separated and the aqueous layer was extracted with DCM. The
 25 combined organic fractions were concentrated, and the crude residue was dissolved
 into 1 mL TFA and 0.1 ml MeOH. The solution was stirred at 80 °C for 20 minutes.
 The reaction mixture was diluted with DMF (4 mL) and purified by preparative
 LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired racemic product as a
 TFA salt. LC-MS calculated for $C_{27}H_{25}F_2N_8O_2$ (M+H)⁺: m/z = 531.2; found 531.2.

**Example 114. 2-(4-(2-((8-amino-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-
a]pyrazin-2-yl)methyl)-3-fluorophenyl)-1H-pyrazol-1-yl)-N,N-dimethylacetamide**



Step 1: 2-(4-(2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-fluorophenyl)-

5 *[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorophenyl)-1H-pyrazol-1-yl)acetic acid*




To a solution 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-
[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (0.075 g, 0.118 mmol) (from
10 Example 99, step 4), ethyl 2-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-
pyrazol-1-yl)acetate (0.157 g, 0.559 mmol), and dicyclohexyl(2',4',6'-
triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1)
(0.022 g, 0.028 mmol) in dioxane (2.0 mL) and water (0.4 mL) was added potassium
phosphate tribasic (0.178 g, 0.838 mmol). The reaction mixture was stirred at 100 °C
15 for 1 h. The reaction mixture was cooled to room temperature, diluted with MeOH (2
ml) and water (2 ml), and into this solution was added lithium hydroxide hydrate
(0.176 g, 4.19 mmol). The suspension was stirred at room temperature for 1 hour. The
suspension was made neutral by addition of saturated aqueous ammonium chloride
solution, and diluted with DCM. The layers were separated and the aqueous layer was
20 extracted with DCM. The combined organic fractions were dried over a plug of
magnesium sulfate and concentrated. Purification by automatic flash column
chromatography afforded the desired product (0.050 g, 25%). LC-MS calculated for
C₄₀H₃₃F₂N₈O₄ (M+H)⁺: m/z = 727.3; found 727.2.

25 *Step 2: 2-(4-(2-((8-amino-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-a]pyrazin-
2-yl)methyl)-3-fluorophenyl)-1H-pyrazol-1-yl)-N,N-dimethylacetamide*

A vial was charged with 2-(4-(2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)-3-fluorophenyl)-1*H*-pyrazol-1-yl)acetic acid (0.01 g, 0.014 mmol), DMF (0.46 ml), dimethylamine solution (0.069 ml, 2M in THF), *N,N*-diisopropylethylamine (8.89 mg, 0.069 mmol), and 2-(3*H*-[1,2,3]triazolo[4,5-*b*]pyridin-3-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate(V) (10.46 mg, 0.028 mmol). The reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted with dichloromethane and water. The layers were separated and the aqueous layer was extracted with DCM. The combined organic fractions were dried over a plug of magnesium sulfate and concentrated. The combined organic fractions were concentrated, and the crude residue was dissolved into 1 mL TFA. The solution was stirred at 80 °C for 20 minutes. The reaction mixture was diluted with DMF (4 mL) and purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₆H₂₂F₂N₉O (M+H)⁺: m/z = 514.2; found 514.2.

Example 115. 2-(4-(2-((8-amino-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)-3-fluorophenyl)-1*H*-pyrazol-1-yl)acetamide




To a solution 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile (0.075 g, 0.118 mmol) (from Example 99, step 4), 2-(4-(4,4,5,5-tetramethyl-1,3-dioxolan-2-yl)-1*H*-pyrazol-1-yl)acetamide (0.060 g, 0.235 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (0.018 g, 0.024 mmol) in dioxane (1.0 mL) and water (0.2 mL) was added potassium phosphate tribasic (0.075 g, 0.353 mmol). The reaction mixture was stirred at 100 °C for 1 h. The reaction mixture was cooled to room temperature and diluted with water and DCM. The layers were separated and the aqueous layer was extracted with DCM. The combined organic fractions were dried over a plug of magnesium sulfate and


concentrated. The crude residue was dissolved into TFA (2 ml) and stirred at 80 °C for 20 minutes. The reaction mixture was diluted with DMF (3 mL) and purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₄H₁₈F₂N₉O (M+H)⁺: m/z = 486.2; found

5 486.1.

Example 116. 3-(8-amino-2-(2-fluoro-6-(1-((*trans*)-3-(methylamino)cyclobutyl)-1*H*-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile



10 *Step 1: Tert-butyl methyl((trans)-3-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazol-1-yl)cyclobutyl)carbamate*



A vial was charged with 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole (0.192, 0.988), tert-butyl ((*cis*)-3-hydroxycyclobutyl)(methyl)carbamate (0.1 g, 0.494 mmol), triphenylphosphine (0.285 g, 1.09 mmol), and THF (1 ml). The solution was cooled to 0 °C and diisopropyl (*E*)-diazene-1,2-dicarboxylate (0.234 ml, 1.19 mmol) was added dropwise. The reaction mixture was slowly warmed to room temperature and stirred overnight. The solvent was stripped and the crude residue purified by automatic flash column chromatography to afford the desired product (0.112 g, 60%). LC-MS calculated for C₁₉H₃₃BN₃O₄ (M+H)⁺: m/z = 378.3; found


15 378.3.

20

*Step 2: 3-(8-amino-2-(2-fluoro-6-(1-((trans)-3-(methylamino)cyclobutyl)-1*H*-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile*

To a solution 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile (0.01 g, 0.016 mmol) (from Example 99, step 4), *tert*-butyl methyl((*trans*)-3-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazol-1-yl)cyclobutyl)carbamate (0.012 g, 0.031 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (2.50 mg, 3.14 μ mol) in dioxane (0.24 mL) and water (0.06 mL) was added potassium phosphate tribasic (0.010 g, 0.047 mmol). The reaction mixture was stirred at 100 °C for 1 h. The reaction mixture was cooled to room temperature and diluted with water and DCM. The layers were separated and the aqueous layer was extracted with DCM. The combined organic fractions were dried over a plug of magnesium sulfate and concentrated. The crude residue was dissolved into TFA (1 mL) and stirred at 80 °C for 20 minutes. The reaction mixture was diluted with DMF (4 mL) and purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₇H₂₄F₂N₉ (M+H)⁺: m/z = 512.2; found 512.1.


Example 117. 3-(8-amino-2-(2-(1-(2-cyanoethyl)-1*H*-pyrazol-4-yl)-6-fluorobenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile



To a solution 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chloro-6-fluorobenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile (0.01 g, 0.016 mmol) (from Example 99, step 4), 3-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazol-1-yl)propanenitrile (0.008 g, 0.031 mmol), and dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine-(2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (2.50 mg, 3.14 μ mol) in dioxane (0.24 mL) and water (0.06 mL) was added potassium phosphate tribasic (0.010 g, 0.047 mmol). The reaction mixture was stirred at 100 °C for 1 hour. The reaction mixture was cooled to room temperature and diluted with water and DCM. The layers were separated and the aqueous layer was extracted with DCM. The combined organic fractions were dried over a plug of magnesium sulfate


and concentrated. The crude residue was dissolved into TFA (1 ml) and stirred at 80 °C for 20 minutes. The reaction mixture was diluted with DMF (4 mL) and purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₅H₁₈F₂N₉ (M+H)⁺: m/z = 482.2; found 482.2.

Example 118. 3-(8-amino-2-(2-fluoro-6-(1-(2-(3-hydroxyazetidin-1-yl)-2-oxoethyl)-1H-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




A vial was charged with 2-(4-(2-((8-(bis(4-methoxybenzyl)amino)-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorophenyl)-1H-pyrazol-1-yl)acetic acid (0.01 g, 0.014 mmol) (from Example 114, step 1), DMF (0.46 ml), azetidin-3-ol (10 mg, 0.138 mmol), *N,N*-diisopropylethylamine (8.89 mg, 0.069 mmol), and 2-(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate(V) (10.46 mg, 0.028 mmol). The reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted with dichloromethane and water. The layers were separated and the aqueous layer was extracted with DCM. The combined organic fractions were dried over a plug of magnesium sulfate and concentrated. The combined organic fractions were concentrated, and the crude residue was dissolved into 1 mL TFA. The solution was stirred at 80 °C for 20 minutes. The reaction mixture was diluted with DMF (4 mL) and purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₇H₂₂F₂N₉O₂ (M+H)⁺: m/z = 542.2; found 542.1.

Example 119. 3-(8-amino-2-((3-methylpyridin-2-yl)methoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Step 1: 3-(8-(bis(4-methoxybenzyl)amino)-2-hydroxy-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A reaction vial was charged with 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (50 mg, 0.090 mmol) (from Example 27, Step 4), tBuBrettPhos Pd G3 (3.8 mg, 0.0045 mmol), sodium tert-butoxide (17.3 mg, 0.18 mmol), H₂O (0.1 mL) and dioxane (1 mL). The reaction mixture was purged with nitrogen for 5 min before heating to 110 °C and stirring for 5 h. The reaction mixture was then diluted with water and ethyl acetate. The organic layer was separated, washed with brine, dried over Na₂SO₄, filtered and concentrated. The resulting residue was purified via flash chromatography to give the desired product as a white solid (40 mg, 90%). LC-MS calculated for C₂₈H₂₅N₆O₃ (M+H)⁺: m/z = 493.2; found 493.3.

Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-2-((3-methylpyridin-2-yl)methoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



3-(8-(Bis(4-methoxybenzyl)amino)-2-hydroxy-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (15 mg, 0.030 mmol) was dissolved in acetonitrile (0.5 mL), and 2-(chloromethyl)-3-methylpyridine (13 mg, 0.090 mmol) and potassium carbonate (13 mg, 0.090 mmol) were added. The reaction was stirred at room temperature for 1 h. Upon completion, NH₄Cl saturated aqueous solution was added and the content was extracted with EtOAc (2 mL x 3). The combined organic phase was dried over MgSO₄, filtered, and the solvents removed. The crude product was re-dissolved in dichloromethane (1 mL). NBS (8 mg, 0.045 mmol) was added. The mixture was stirred at rt for 0.5 h before quenching by the addition of aqueous Na₂SO₃ solution. The organic layer was separated, dried over Na₂SO₄, filtered and concentrated. The crude brominated product was added LiCl (1.3 mg, 0.030 mmol), CuI (5.8 mg, 0.030 mmol), Pd₂(dba)₃ (2.3 mg, 0.003 mmol), PPh₃ (1.3 mg, 0.005 mmol) and 4-(tributylstannyl)pyrimidine (14 mg, 0.038 mmol). The reaction mixture was dissolved in dioxane, and purged with nitrogen for 5 min, before heating to 100 °C for 15 h. The reaction mixture was then cooled to rt, filtered, concentrated, and purified via flash chromatography to give the desired product as a white solid (10 mg, 50%). LC-MS calculated for C₃₉H₃₄N₉O₃ (M+H)⁺: m/z = 676.3; found 676.3.


Step 3: 3-(8-amino-2-((3-methylpyridin-2-yl)methoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

3-(8-(bis(4-methoxybenzyl)amino)-2-((3-methylpyridin-2-yl)methoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.015 mmol)

was added TFA (0.5 mL), and stirred at 100 °C for 5 min. The reaction mixture was then cooled to room temperature, solvent removed, diluted with methanol, and purified via prep-LCMS (pH 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₃H₁₈N₉O (M+H)⁺: m/z = 436.2; found


5 436.0.

Example 120. 3-(8-Amino-2-((3-((1-(pyridin-4-ylmethyl)-1H-pyrazol-4-yl)amino)pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile



10


Step 1: Diethyl 2-(3-chloropyridin-2-yl)malonate



The mixture of 3-chloro-2-fluoropyridine (6.25 g, 47.5 mmol), diethyl malonate (18.27 g, 114 mmol), cesium carbonate (37.2 g, 114 mmol) and DMSO (55.9 ml) was heated at 100 °C for 10 h. The mixture was poured onto ice, diluted with ethyl acetate. The organic layer was separated, washed with water, and brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified with silica gel column (eluting with a gradient 0-30% ethyl acetate in hexane) to give the desired product (12.9 g, 100%). LC-MS calculated for C₁₂H₁₅ClNO₄ (M+H)⁺: m/z = 272.1; found 272.1.

20

Step 2: ethyl 2-(3-chloropyridin-2-yl)acetate




A mixture of diethyl 2-(3-chloropyridin-2-yl)malonate (12.9 g, 47.5 mmol), sodium chloride (3.05 ml, 52.2 mmol), water (1.711 ml, 95 mmol) in DMSO (68 ml) was heated at 145 °C for 5 h. LCMS showed completion of reaction. The reaction

25

mixture was diluted with ethyl acetate and washed with water (2x), brine, dried over MgSO_4 , filtered and concentrated. The residue was purified with silica gel column (eluting with a gradient 0-30% ethyl acetate in hexane) to give the desired product (7.8 g, 82%). LC-MS calculated for $\text{C}_9\text{H}_{11}\text{ClNO}_2$ ($\text{M}+\text{H}$)⁺: m/z = 200.0; found 200.0.

5


Step 3: 2-(3-chloropyridin-2-yl)acetic acid



To a solution of ethyl 2-(3-chloropyridin-2-yl)acetate (7.8 g, 39.1 mmol) in THF (130 ml) was added 1.0 M sodium hydroxide solution (78 ml, 78 mmol). The resulting mixture was stirred at rt for 1 h. LCMS showed the completion of reaction. pH of the reaction mixture was adjusted with 1 N HCl to pH 3. The organic solvent was removed *in vacuo*. The resulting precipitate was collected via filtration, washed with water and ethyl acetate and dried under vacuum to give the product as white solid (5.5 g, 82%). LC-MS calculated for $\text{C}_7\text{H}_7\text{ClNO}_2$ ($\text{M}+\text{H}$)⁺: m/z = 172.0; found 172.0.

15

Step 4: 6-Bromo-2-((3-chloropyridin-2-yl)methyl)-N,N-bis(4-methoxybenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine




To a flask charged with 2-(3-chloropyridin-2-yl)acetic acid (0.372 g, 2.168 mmol), HATU (0.907 g, 2.385 mmol) in CH_2Cl_2 (21.68 ml) was added 1,2-diamino-3-(bis(4-methoxybenzyl)amino)-5-bromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate (1.398 g, 2.168 mmol) (from Example 99, Step 2), followed by DIEA (0.757 ml, 4.34 mmol). After stirring at room temperature for 6h, LCMS showed completion of reaction. The reaction mixture was diluted with DCM

25

and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude was purified with flash chromatography (eluting with a gradient 0-40% ethyl acetate in hexanes with 10% DCM) to give the desired product (1.0 g, 80%). LC-MS calculated for C₂₇H₂₅BrClN₆O₂ (M+H)⁺: m/z = 579.1, 581.1;

5 found 579.1, 581.1.

Step 5: 3-(8-(bis(4-methoxybenzyl)amino)-2-((3-chloropyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




10 A flask charged with 6-bromo-2-((3-chloropyridin-2-yl)methyl)-N,N-bis(4-methoxybenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine (4.70 g, 8.11 mmol), (3-cyano-2-fluorophenyl)boronic acid (1.871 g, 11.35 mmol), Cs₂CO₃ (5.28 g, 16.21 mmol), tetrakis (0.937 g, 0.811 mmol), 1,4-dioxane (73.7 ml) and water (7.37 ml) was evacuated under vacuum and refilled with N₂ (repeated three times). The mixture was heated at 90 °C for 4h. Another 0.3 equivalent (3-cyano-2-fluorophenyl)boronic acid (1.871 g, 11.35 mmol) was added and heated at 90 °C for 2h. LCMS showed total completion of reaction. The reaction mixture was diluted with DCM and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude was triturated with hexanes and ethyl acetate, the resulting precipitate was collected via filtration and washed with methanol, dried under vacuum to give the desired product as white solid (4.4 g, 88%). LC-MS calculated for C₃₄H₂₈ClFN₇O₂ (M+H)⁺: m/z = 620.2; found 620.2.

25 *Step 6: 3-(8-Amino-2-((3-((1-(pyridin-4-yl)methyl)-1H-pyrazol-4-yl)amino)pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile*


To a vial 3-(8-(bis(4-methoxybenzyl)amino)-2-((3-chloropyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (15 mg, 0.024 mmol), 1-(pyridin-4-ylmethyl)-1H-pyrazol-4-amine (4.21 mg, 0.024 mmol), Brettphos palladacycle (3.29 mg, 3.63 μ mol), and cesium carbonate (12.44 μ l, 0.073 mmol) were added. The vial was sealed with a teflon screw-cap, evacuated and backfilled with nitrogen (this process was repeated a total of three times). Anhydrous t-butanol (1 ml) was added. The mixture was heated to 90 °C for 2 h. The reaction mixture was filtered through a SiliaPrep-Thiol funnel, the filtrate was concentrated. The residue was treated with TFA (1 mL) at 80 °C for 20 min. The volatile was removed, the crude was dissolved in methanol and purified with prep-LCMS (pH 2, acetonitrile/water+TFA) to give the desired product as TFA salt (4 mg, 33%). LC-MS calculated for $C_{27}H_{21}FN_{11}$ (M+H)⁺: m/z = 518.2; found 518.2.

Example 121. 3-(8-amino-2-((3-((1-(tetrahydro-2H-pyran-4-yl)-1H-pyrazol-4-yl)amino)pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




The title compound was prepared using similar procedures as described for Example 120 with 1-(tetrahydro-2H-pyran-4-yl)-1H-pyrazol-4-amine replacing 1-(pyridin-4-ylmethyl)-1H-pyrazol-4-amine in Step 9. The reaction mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{26}H_{24}FN_{10}O$ (M+H)⁺: m/z = 511.2; found 511.2.

Example 122. 3-(8-Amino-2-((3-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Step 1: 3-(8-(bis(4-methoxybenzyl)amino)-2-((3-chloropyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-chloro-2-methylpyridine (0.367 g, 2.88 mmol) in THF (10 mL) was added 0.65 M (2,2,6,6-tetramethylpiperidin-1-yl)zinc(II) lithium chloride (6.65 mL, 4.32 mmol) at rt. The resulting yellow solution was stirred at same temperature for 1 h, scandium trifluoromethanesulfonate (0.057 g, 0.115 mmol) was added and stirred at room temperature for 15 min. A microwave vial was charge with 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.64 g, 1.152 mmol), palladium acetate (0.021 g, 0.092 mmol) , and 2'-(dicyclohexylphosphino)-N,N,N',N'-tetramethylbiphenyl-2,6-diamine (0.080 g, 0.184 mmol) was evacuated under high vacuum and backfilled with nitrogen . The mixture was cooled to 0 °C and the zinc reagent was added slowly via syringe. After addition, the reaction was heated to 60 °C for 1h. The reaction solution was partitioned between EtOAc and sat. NH₄Cl solution. The layers were separated and the aqueous extracted further with EtOAc (2x). The combined organics were washed with water and brine, dried over MgSO₄, and concentrated. The resulting residue was purified via flash chromatography to afford the product. LC-MS calculated for C₃₄H₂₉ClN₇O₂ (M+H)⁺: m/z = 602.2; found 602.2.

Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((3-chloropyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-((3-chloropyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.218 g, 0.362 mmol), 1-bromopyrrolidine-2,5-dione (0.061 g, 0.344 mmol) and CH₂Cl₂ (4 ml); was stirred at 0 °C for 30 min, The reaction mixture was diluted with sat. NaHCO₃. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for C₃₄H₂₈BrClN₇O₂ (M+H)⁺: m/z = 680.1, 682.1; found 680.1, 682.1.

Step 3: 3-(8-Amino-2-((3-chloropyridin-2-yl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-((3-chloropyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (86 mg, 0.126 mmol), 4-(tributylstannyl)pyrimidine (69.9 mg, 0.189 mmol), and copper(I) chloride (15.00 mg, 0.152 mmol), lithium chloride (6.42 mg, 0.152 mmol) and tetrakis(triphenylphosphine)palladium(0) (14.59 mg, 0.013 mmol) in THF (3 ml) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction was diluted with methanol and purified with prep-LCMS (pH 2, acetonitrile/water with

TFA) to give coupling product, that was treated with TFA (1 mL) at 80 °C for 20 min. The volatile was removed and the resulting residue was dissolved in methanol and purified with prep-LCMS (pH 2, acetonitrile/water with TFA) to give the desired product. LC-MS calculated for $C_{22}H_{15}ClN_9$ ($M+H$)⁺: m/z = 440.1 found 440.1.


5

Step 4: 3-(8-amino-2-((3-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

A mixture of 3-(8-amino-2-((3-chloropyridin-2-yl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (10 mg, 0.023 mmol), 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (5.7 mg, 0.027 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (1.8 mg, 2.3 μ mol) and tripotassium phosphate hydrate (11.5 mg, 0.050 mmol) in 1,4-dioxane (2.0 mL)/Water (0.65 mL) was stirred at 80 °C for 1 h. The mixture was diluted in methanol and DMSO and purified with prep-LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for $C_{26}H_{20}N_{11}$ ($M+H$)⁺: 486.2 found 486.2.


15

Example 123. (S)-3-(8-Amino-2-(2-((3-hydroxypyrrolidin-1-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




20

Step 1: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A microwave vial was charge with 3-(8-(bis(4-methoxybenzyl)amino)-2-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (350 mg, 0.630 mmol),
 5 palladium acetate (7.07 mg, 0.032 mmol), and 2'-(dicyclohexylphosphino)-N,N,N',N'-tetramethylbiphenyl-2,6-diamine (27.5 mg, 0.063 mmol) was evacuated under high vacuum and backfilled with nitrogen . (2-chlorobenzyl)zinc(II) chloride (1.4 mL, 0.693 mmol) was added via syringe. After addition, the reaction was heated to 60 °C
 10 for 1 h. The reaction solution was partitioned between EtOAc and sat. NH₄Cl solution. The layers were separated and the aqueous extracted further with EtOAc (2x). The combined organics were washed with water and brine, dried over MgSO₄, and concentrated. The residue was purified with flash chromatography to give the desired product (0.32 g, 82%). LC-MS calculated for C₃₅H₃₀ClN₆O₂ (M+H)⁺: m/z =
 15 601.2; found 601.2.

Step 2: 3-(8-(Bis(4-methoxybenzyl)amino)-5-bromo-2-(2-chlorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (0.379 g, 0.631 mmol) in DCM (6.3 ml) was added 1-bromopyrrolidine-2,5-dione (0.107 g, 0.599 mmol) at 0 °C. The resulting mixture was stirred at 0 °C for 30 min. The reaction mixture was diluted with sat. NaHCO₃. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue was purified with flash chromatography to give the desired product as a light yellow oil (0.38 g, 89%). LC-MS calculated for C₃₅H₂₉BrClN₆O₂ (M+H)⁺: m/z = 679.1, 681.1; found 679.1, 681.1.

Step 3: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(2-chlorobenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (381 mg, 0.560 mmol), 4-(tributylstannyl)pyrimidine (310 mg, 0.840 mmol), and copper(I) chloride (66.6 mg, 0.672 mmol), lithium chloride (28.5 mg, 0.672 mmol) and tetrakis(triphenylphosphine)palladium(0) (64.7 mg, 0.056 mmol) in THF (6 ml) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction mixture was concentrated under vacuum and the resulting residue was purified with flash chromatography to give the desired product as a light yellow oil (0.31 g, 83%). LC-MS calculated for C₃₉H₃₂ClN₈O₂ (M+H)⁺: m/z = 679.2 found 679.2.

Step 4: 3-(8-(Bis(4-methoxybenzyl)amino)-5-(pyrimidin-4-yl)-2-(2-vinylbenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-2-(2-chlorobenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (300 mg, 0.442 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (82 mg, 0.530 mmol), dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine - (2'-aminobiphenyl-2-yl)(chloro)palladium (1:1) (34.8 mg, 0.044 mmol) and tripotassium phosphate hydrate (224 mg, 0.972 mmol) in 1,4-dioxane (5.0mL)/water (1.7 mL) was stirred at 80 °C for 1 h. The reaction mixture was concentrated under vacuum and the resulting residue was purified with flash chromatography to give the desired product as a light yellow oil. LC-MS calculated for $C_{41}H_{35}N_8O_2$ ($M+H$)⁺: m/z = 671.3; found 671.3.

Step 5: 3-(8-(Bis(4-methoxybenzyl)amino)-2-(2-formylbenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




3-(8-(bis(4-methoxybenzyl)amino)-5-(pyrimidin-4-yl)-2-(2-vinylbenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (296 mg, 0.441 mmol) was mixed with tetrahydrofuran (2.2 mL), 0.16 M osmium tetroxide in water (220 μ L, 0.035 mmol), sodium metaperiodate (425 mg, 1.986 mmol) and water (2.2 mL). The reaction was stirred at 60 °C for 1 h before quenched with sat. $\text{Na}_2\text{S}_2\text{O}_3$. The mixture was extracted with DCM. The organic phase was washed with brine, dried over Na_2SO_4 , filtered, and concentrated to afford the product as a light yellow oil. LC-MS calculated for $\text{C}_{40}\text{H}_{33}\text{N}_8\text{O}_3$ ($\text{M}+\text{H}$) $^+$: m/z = 673.3; found 673.3.

Step 6: (S)-3-(8-Amino-2-(2-((3-hydroxypyrrolidin-1-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

A mixture of (3-(8-(bis(4-methoxybenzyl)amino)-2-(2-formylbenzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (12 mg, 0.018 mmol), (S)-pyrrolidin-3-ol (1.6 mg, 0.018 mmol) in DCM (0.5 mL) and MeOH (0.5 mL) was added sodium triacetoxyborohydride (7.6 mg, 0.036 mmol). After stirring at room temperature overnight, solvent was removed *in vacuo*. The residue was treated with TFA (1 mL) at 80 °C for 20 min. After removal of volatiles. The residue was dissolved in methanol and purified with preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for $\text{C}_{28}\text{H}_{26}\text{N}_9\text{O}$ ($\text{M}+\text{H}$) $^+$: 504.2 found 504.2.


Example 124. 3-(8-Amino-5-(imidazo[1,2-a]pyridin-6-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



The title compound was prepared using similar procedures as described for Example 15 with 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridine replacing (6-methoxypyridin-3-yl)boronic acid in Step 5. The reaction

mixture was purified by prep-HPLC (pH = 2, acetonitrile/water+TFA) to give the desired product as TFA salt. LC-MS calculated for $C_{19}H_{13}N_8$ (M+H)⁺: m/z = 353.1; found 353.1.


5 **Example 125. 3-(8-amino-2-(azetidine-1-carbonyl)-5-(3-fluoropyridin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile**




A mixture of 3-(2-(azetidine-1-carbonyl)-8-(bis(4-methoxybenzyl)amino)-5-bromo-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (20 mg, 0.032 mmol) (from
 10 Example 61, Step 2), 3-fluoropyridine-4-boronic acid (16 mg, 0.13 mmol), sodium carbonate (34 mg, 0.32 mmol) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (5 mg, 0.006 mmol) in 4:1 dioxane/Water (1.3 mL) was stirred at 100 °C for 1.5 hours. The
 reaction mixture was diluted with dichloromethane and water, and the organic solvent
 15 was concentrated *in vacuo*, the crude product was dissolved in 3 mL TFA and stirred at 80 °C for 20 mins. After TFA being removed, the crude product was purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to afford the desired product as a TFA salt. LC-MS calculated for $C_{21}H_{16}FN_8O$ (M+H)⁺: m/z = 415.2; found 415.1.

20

Example 126. 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(2,6-dimethylpyridin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




Step 1: 2-((tert-butyldimethylsilyl)oxy)-2-(2,6-difluorophenyl)acetonitrile




To a stirred solution of 2,6-difluorobenzaldehyde (0.987 ml, 8.97 mmol) in
 5 acetonitrile (9 ml) at rt was added tert-Butyldimethylsilyl cyanide (1.959 g, 13.45
 mmol) (1.5 eq) and cesium fluoride (0.272 g, 1.793 mmol) (0.2 eq). The reaction
 mixture was stirred at rt overnight (16 h). The reaction mixture was filtered to remove
 CsF. The filtrate was concentrated *in vacuo*. The residue was purified by Biotage
 Isolera (with 40 g silica gel column) eluting with 0 - 10% EtOAc/Hexane to give the
 10 product as a colorless oil (2.468 g, 97%). LCMS calculated for C₁₄H₂₀F₂NOSi
 (M+H)⁺: m/z = 284.1; found: 284.1.

Step 2: 2-((tert-butyldimethylsilyl)oxy)-N-(3,5-dibromopyrazin-2-yl)-2-(2,6-
 difluorophenyl)acetimidamide



15 To a stirred solution of 2-((tert-butyldimethylsilyl)oxy)-2-(2,6-
 difluorophenyl)acetonitrile (2.0 g, 7.06 mmol) in anhydrous 1,2-Dichloroethane (10
 ml) (5 volume) at rt was added 3,5-
 dibromopyrazin-2-amine (2.73 g, 10.59 mmol) (1.5 eq) and Tin(IV) chloride (1.264
 20 ml, 10.59 mmol) (1.5 eq). The resulting suspension was heated at 110 °C overnight
 (15 h). The reaction mixture was cooled to rt. It was diluted with dichloromethane (20
 mL), basified with 1N NaOH to pH 10. It was extracted with dichloromethane (50
 mL). The dichloromethane layer was dried over Na₂SO₄, filtered and concentrated *in*
vacuo. The residue was purified by Biotage Isolera (with 120 g silica gel column)
 25 eluting with 0 - 30% EtOAc/Hexane to give the product as a light yellow solid (2.681
 g, 70%). LCMS calculated for C₁₈H₂₃Br₂F₂N₄OSi (M+H)⁺: m/z = 534.9; found:
 534.9.


Step 3: 6,8-dibromo-2-(((tert-butyldimethylsilyl)oxy)(2,6-difluorophenyl)methyl)-[1,2,4]triazolo[1,5-a]pyrazine



To a stirred solution of 2-(((tert-butyldimethylsilyl)oxy)-N-(3,5-dibromopyrazin-2-yl)-2-(2,6-difluorophenyl)acetimidamide (1.0 g, 1.831 mmol) in Hexafluoroisopropanol (18 ml) (HFIPA, 18 volume) at rt was added (Bis(trifluoroacetoxy)iodo)benzene (1.623 g, 3.66 mmol) (2 eq) and Triethylamine (1.023 ml, 7.32 mmol) (4 eq). The reaction mixture was stirred at rt for 2 hours. The reaction was quenched with saturated aqueous NaHCO₃ (20 mL). It was extracted with dichloromethane (50 mL). Dichloromethane layer was dried over Na₂SO₄, filtered and concentrated *in vacuo*. The residue purified by Biotage Isolera (with 120 g silica gel column) eluting with 0 - 20% EtOAc/Hexane to give the product as a viscous light yellow oil (0.98 g, 95%). LCMS calculated for C₁₈H₂₁Br₂F₂N₄OSi (M+H)⁺: m/z = 533.0; found 532.8.

15


Step 4: 6-bromo-2-(((tert-butyldimethylsilyl)oxy)(2,6-difluorophenyl)methyl)-N,N-bis(4-methoxybenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine



To a stirred solution of 6,8-dibromo-2-(((tert-butyldimethylsilyl)oxy)(2,6-difluorophenyl)methyl)-[1,2,4]triazolo[1,5-a]pyrazine (0.98 g, 1.741 mmol) in 2-Propanol (10 ml) at rt was added bis(4-methoxybenzyl)amine (0.594 g, 2.264 mmol) and N,N-Diisopropylethylamine (0.613 ml, 3.48 mmol). The reaction mixture was heated at 80 °C for 2 hours. The reaction mixture was cooled to rt, and concentrated *in vacuo*. The residue was purified by Biotage Isolera (with 120 g silica gel column) eluting with 0 - 30% EtOAc/Hexane to give the product as a white foamy solid (1.190 g, 96%). LCMS calculated for C₃₄H₃₉BrF₂N₅O₃Si: m/z = 710.2, found 710.3.

25


Step 5: 3-(8-(bis(4-methoxybenzyl)amino)-2-(((tert-butyldimethylsilyl)oxy)(2,6-difluorophenyl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile



To a stirred solution of 6-bromo-2-(((tert-butyldimethylsilyl)oxy)(2,6-difluorophenyl)methyl)-N,N-bis(4-methoxybenzyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine (1.37 g, 1.93 mmol) in 1,4-dioxane/H₂O (4 : 1, 13 mL), (3-cyano-2-fluorophenyl)boronic acid (0.413 g, 2.5 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (78 mg, 0.1 mmol) (XPhos Pd G2) and sodium carbonate (0.613 mg, 5.78 mmol) were added at rt. The reaction mixture was heated at 90 °C for 2 hours. The reaction mixture was cooled to rt, extracted with dichloromethane and concentrated *in vacuo*. The residue was purified by Biotage Isolera eluting with 0 - 30% EtOAc/Hexane to give the product as a foamy solid (1.3 g, 90%). LCMS calculated for C₄₁H₄₂F₃N₆O₃Si: m/z = 751.3, found 751.2.

15

Step 6: 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(((tert-butyldimethylsilyl)oxy)(2,6-difluorophenyl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile




To a solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-(((tert-butyldimethylsilyl)oxy)(2,6-difluorophenyl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (690 mg, 0.92 mmol) in 5 mL dichloromethane, 1-bromopyrrolidine-2,5-dione (180 mg, 1.0 mmol) was added at rt. The reaction mixture was stirred for overnight before it was purified by Biotage Isolera eluting with 0 - 30% EtOAc/Hexane to give the product as a foamy solid (700 mg, 92%). LCMS calculated for C₄₁H₄₁BrF₃N₆O₃Si: m/z = 829.2, found 829.3.

25


Step 7: 3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(2,6-dimethylpyridin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile

A mixture of 3-(8-(bis(4-methoxybenzyl)amino)-5-bromo-2-(((tert-butyl)dimethylsilyl)oxy))(2,6-difluorophenyl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile (350 mg, 0.42 mmol), (2,6-dimethylpyridin-4-yl)boronic acid (96 mg, 0.63 mmol), sodium carbonate (134 mg, 1.26 mmol) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (16 mg, 0.02 mmol) in 4:1 dioxane/Water (5 mL) was stirred at 90 °C for 75 mins. The reaction mixture was diluted with dichloromethane and water and the organic solvent was removed *in vacuo*, the crude product was dissolved in 10 mL TFA and stirred at 80 °C for 1 hour. After TFA being removed *in vacuo*, the crude product was basified by sodium bicarbonate solution, and extracted with dichloromethane. The dichloromethane layer was concentrated *in vacuo*, and purified by Biotage Isolera to give the desired product as a racemic mixture (140 mg, 66%)
The product was then separated with chiral HPLC using a chiral column (Phenomenex Lux 5um Cellulose-4, 21.2x250mm) and 75% EtOH in hexanes (20 mL/min) solvent system. Peak 2 was isolated, and further purified by preparative LC/MS (pH = 2, acetonitrile/water with TFA) to give the desired product as a TFA salt. LC-MS calculated for C₂₆H₁₉F₃N₇O (M+H)⁺: m/z = 502.2; found 502.2.

Example 127. 3-(8-amino-5-(1-(methyl-d3)-6-oxo-1,6-dihydropyridazin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile




Step 1: 6-bromo-2-(methyl-d3)pyridazin-3(2H)-one



To a solution of 6-bromopyridazin-3(2H)-one (1.1 g, 6.3 mmol) in 9 mL DMF, iodomethane-d3 (1.0 g, 6.91 mmol) and potassium carbonate (1.3 g, 9.4 mmol) were added at rt and stirred overnight. The resulting mixture was quenched with ammonium chloride solution and extracted with dichloromethane, after concentrated *in vacuo*, the crude product was purified by Biotage Isolera to afford the desired product (0.96g, 80%) as white solid. LC-MS calculated for C₅H₃D₃BrN₂O (M+H)⁺: m/z = 192.0; found 192.1.

Step 2: 2-(methyl-d3)-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridazin-3(2H)-one




To a mixture of 6-bromo-2-(methyl-d3)pyridazin-3(2H)-one (300 mg, 1.56 mmol), potassium acetate (460 mg, 4.69 mmol), and 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (440 mg, 1.7 mmol) in dioxane (5 mL), Tetrakis(triphenylphosphine)palladium(0) (90 mg, 0.08 mmol) was added at rt, the resulting mixture was stirred at 100 °C overnight. The reaction mixture was then quenched with ammonium chloride solution and extracted with dichloromethane, after being concentrated *in vacuo*, the crude product was purified by Biotage Isolera, and the desired product (0.17 g, 46%) was obtained as a white solid. LC-MS calculated for C₁₁H₁₅D₃BN₂O₃ (M+H)⁺: m/z = 240.2; found 240.2.


Step 3: 3-(8-amino-5-(1-(methyl-d3)-6-oxo-1,6-dihydropyridazin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile

To a mixture of 3-(8-amino-5-bromo-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile (Example 27, Step 7; 20 mg, 0.05 mmol), 2-(methyl-d₃)-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridazin-3(2H)-one (24 mg, 0.1 mmol), and sodium carbonate (20 mg, 0.2 mmol) in dioxane/water (4:1, 1.5 mL), chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (4 mg, 0.005 mmol) (XPhos Pd G2) was added. The resulting mixture was heated at 90 °C for 1 hour. The mixture was concentrated and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for C₂₃H₁₅D₃N₉O (M+H)⁺: m/z = 439.2; found 439.2.

Example 128. 3-(8-amino-2-((6-methoxypyridin-2-yl)methyl)-5-(1-methyl-6-oxo-1,6-dihydropyridazin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



Step 1: 6-bromo-N,N-bis(4-methoxybenzyl)-2-((6-methoxypyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine




To a vial charged with 2-(6-methoxypyridin-2-yl)acetic acid (47 mg, 0.28 mmol), HATU (133 mg, 0.35 mmol) in dichloromethane 2 mL was added 1,2-diamino-3-(bis(4-methoxybenzyl)amino)-5-bromopyrazin-1-ium 2,4,6-trimethylbenzenesulfonate (example 99, step 2, 150 mg, 0.23 mmol), followed by N-ethyl-N-isopropylpropan-2-amine (90 mg, 0.7 mmol). After stirring at room temperature for 6 hours, the reaction mixture was diluted with dichloromethane and water. The organic layer was washed with brine, dried over Na₂SO₄, filtered and

concentrated. The crude was purified with flash chromatography (eluting with a gradient 0-10% ethyl acetate in dichloromethane) to give the desired product (100 mg, 75%). LC-MS calculated for $C_{28}H_{28}BrN_6O_2$ ($M+H$)⁺: m/z = 575.1, 577.1; found 575.1, 577.1.

5


Step 2: 3-(8-(bis(4-methoxybenzyl)amino)-2-((6-methoxypyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



To a solution of 6-bromo-N,N-bis(4-methoxybenzyl)-2-((6-methoxypyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-8-amine (75 mg, 0.13 mmol) and (3-cyanophenyl)boronic acid (29 mg, 0.2 mmol) in 2 mL 1,4-dioxane/ H_2O = 4:1, sodium carbonate (42 mg, 0.4 mmol) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (3 mg, 0.004 mmol) (XPhos Pd G2) were added. The reaction mixture was heated to 90 °C and stirred for 1 hour before being diluted with dichloromethane and water. The organic layer was washed with brine, dried over Na_2SO_4 , filtered and concentrated. The crude was purified with flash chromatography (eluting with a gradient 0-10% ethyl acetate in dichloromethane) to give the desired product (60 mg, 77%). LC-MS calculated for $C_{35}H_{32}N_7O_3$ ($M+H$)⁺: m/z = 598.2; found 598.2.

20

Step 3: 3-(8-amino-5-(1-(methyl-d3)-6-oxo-1,6-dihydropyridazin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile



A solution of 3-(8-(bis(4-methoxybenzyl)amino)-2-((6-methoxypyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzotrile (60 mg, 0.1 mmol) in 4 mL TFA was heated to 80 °C and stirred for 20 mins. The reaction mixture was then concentrated *in vacuo*, basified with sodium bicarbonate solution and extracted with dichloromethane. The organic layer was concentrated to get crude product for next step.

To a solution of above product in 3 mL dichloromethane, 1-bromopyrrolidine-2,5-dione (23 mg, 0.13 mmol) was added at rt and the reaction mixture was stirred for overnight before being concentrated *in vacuo*. The crude product was then used for next step directly without any further purification.

To a solution of the above crude product in dioxane/water (4:1, 2.0 mL), 2-methyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridazin-3(2H)-one (35 mg, 1.5 mmol), sodium carbonate (30 mg, 0.3 mmol) chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (4 mg, 0.005 mmol) (XPhos Pd G2) was added. The resulting mixture was heated at 90 °C for 1 hour before being diluted with acetonitrile and methanol and purified by preparative LCMS (pH 2, acetonitrile/water with TFA) to afford the desired product as TFA salt. LCMS calculated for $C_{24}H_{20}N_9O_2$ (M+H)⁺: m/z = 466.2; found 466.2.

Example A. Adenosine A2A Receptor cyclic AMP GS Assay

Stably transfected HEK-293 cells expressing the human adenosine A2A receptor (Perkin Elmer) are maintained in MEM culture medium with 10% FBS and 400 µg/mL Geneticin (Life Technologies). 18 to 24 hours prior to assay, geneticin is removed from culture. The cisbio cAMP-GS Dynamic kit utilizing the FRET (Fluorescence Resonance Energy Transfer) technology is used to measure cAMP accumulation in the cells. Compounds of the present disclosure at an appropriate concentration are mixed with 10000 cells/well in white 96 well half area plates (Perkin Elmer) for 30 min at rt gently shaking. Agonist, CGS21680 (R&D Technologies) at 4 nM is added to each well for 60 min at room temperature gently shaking. Detection reagents, d2-labeled cAMP (acceptor) and anti-cAMP cryptate (donor) are added to each well for 60 min at room temperature gently shaking. Plates are read on Pherastar (BMG Labtech), fluorescence ratio 665/620 is calculated and

EC₅₀ determination is performed by fitting the curve of percent of control versus the log of the compound concentration using GraphPad Prism.

Example B. Adenosine A2B Receptor cyclic AMP GS Assay

5 Stably transfected HEK-293 cells expressing the human adenosine A2B receptor (Perkin Elmer) were maintained in MEM culture medium with 10% FBS and 100 µg/mL Geneticin (Life Technologies). 18 to 24 hours prior to assay, geneticin was removed from culture. The cisbio cAMP-GS Dynamic kit utilizing the FRET (Fluorescence Resonance Energy Transfer) technology was used to measure cAMP
10 accumulation in the cells. Compounds of the present disclosure at an appropriate concentration were mixed with 10000 cells/well in white 96 well half area plates (Perkin Elmer) for 30 min at room temperature gently shaking. Agonist, NECA (R&D Technologies) at 12 nM was added to each well for 60 min at room temperature gently shaking. Detection reagents, d2-labeled cAMP (acceptor) and anti-cAMP
15 cryptate (donor) were added to each well for 60 min at rt gently shaking. Plates were read on Pherastar (BMG Labtech), fluorescence ratio 665/620 was calculated and EC₅₀ determination was performed by fitting the curve of percent of control versus the log of the compound concentration using GraphPad Prism. The EC₅₀ data for the Examples obtained via this method are shown in Table 1.

20

Example C. A2A Tag-lite® HTRF Assay

 Assays were conducted in black low volume 384-well polystyrene plates (Greiner 784076-25) in a final volume of 10 µL. Test compounds were first serially diluted in DMSO and 100 nl added to the plate wells before the addition of other
25 reaction components. The final concentration of DMSO was 1%. Tag-lite® Adenosine A2A labeled cells (CisBio C1TT1A2A) were diluted 1:5 into Tag-lite buffer (CisBio LABMED) and spun 1200 g for 5 mins. The pellet was resuspended at a volume 10.4 X the initial cell suspension volume in Tag-lite buffer, and Adenosine A2A Receptor Red antagonist fluorescent ligand (CisBio L0058RED) added at 12.5
30 nM final concentration. 10 ul of the cell and ligand mix was added to the assay wells and incubated at room temperature for 45 minutes before reading on a PHERAstar FS plate reader (BMG Labtech) with HTRF 337/620/665 optical module. Percent binding

of the fluorescent ligand was calculated; where 100 nM of A2A antagonist control ZM 241385 (Tocris 1036) displaces the ligand 100% and 1% DMSO has 0% displacement. The % binding data versus the log of the inhibitor concentration was fitted to a one-site competitive binding model (GraphPad Prism version 7.02) where the ligand constant = 12.5 nM and the ligand K_d = 1.85 nM. The K_i data for the Examples obtained via this method are shown in Table 1.

Example D. A2B Filter Binding Assay

Assays are conducted in deep well polypropylene plates (Greiner 786201) in a final volume of 550 μ L. Test compounds are first serially diluted in DMSO and 5.5 μ L is then added to the plate wells before the addition of other reaction components. The final concentration of DMSO is 3%. HEK293 cell membranes overexpressing the human adenosine receptor A2B (Perkin Elmer ES-113-M400UA) are diluted to 40 μ g/mL in 50 mM HEPES pH 7.0, 5 mM $MgCl_2$, 1 mM EDTA (Assay buffer). [3H] 8-cyclopentyl-1,3-dipropylxanthine (Perkin Elmer NET974001MC) is diluted in assay buffer + 22% DMSO to 24.2 nM, and then further diluted to 1 nM by addition to the diluted membranes. 545 μ L of the membrane and ligand mix is added to the assay wells and incubated on a shaker at room temperature for 1 hour. The membrane mix is then filtered over a UniFilter GF/C filter plate (Perkin Elmer 6005174) pre-soaked in 50 mM HEPES pH 6.5, 5 mM $MgCl_2$, 1 mM EDTA 0.5% BSA and then washed with 5 mL ice cold 50 mM HEPES pH 6.5, 5 mM $MgCl_2$, 1 mM EDTA 0.2% BSA. 50 μ L MicroScintTM cocktail (Perkin Elmer 6013621) is added and plates are read on a Topcount NXT FS (Perkin Elmer). Percent binding of the [3H] ligand is calculated, where 1000 nM of LUF 5834 (Tocris 4603) control displaces the ligand 100% and 3% DMSO has 0% displacement. The % binding data versus the log of the inhibitor concentration is fitted to a one-site competitive binding model (GraphPad Prism version 7.02) where the ligand constant = 2 nM and the ligand K_d = 13 nM.

Example E. A1 and A3 SPA Binding Assays

Both assays are conducted in white 384-well polystyrene plates (Greiner 781075) in a final volume of 50 μ L. Inhibitors are first serially diluted in DMSO and 100 nL is added to the plate wells before the addition of other reaction components.

The final concentration of DMSO is 2%.

Wheatgerm agglutinin-coated yttrium silicate SPA beads (Perkin Elmer RPNQ0023) and CHO-K1 cell membranes overexpressing each human adenosine receptor are incubated in 50 mM HEPES pH 7.0, 5 mM MgCl₂, 1 mM EDTA (Assay buffer) on a rotary stirrer for 2 hours at 4 °C. The beads are pelleted by centrifugation at 6000 g for one minute, and then the supernatant with unbound membrane is discarded. The beads are re-suspended to the original volume in assay buffer. Each radioligand is diluted in assay buffer + 22% DMSO at 12.2X the final concentration, and then added to the SPA bead suspension. 50 µl of the SPA bead reaction mix is added to the assay wells and the plates shaken at 600 rpm for 1 hour at room temperature. The beads are then allowed to settle for 1 hour before reading on a Topcount NXT FS (Perkin Elmer). Percent binding of the radiolabeled ligand is calculated, where a control at >100X K_i displaces the ligand 100% and 2% DMSO has 0% displacement. The % binding data versus the log of the inhibitor concentration is fitted to a one-site competitive binding model (GraphPad Prism version 7.02). Assay conditions are provided in Table A below.

Table A.

Assay Component	A1	A3
SPA beads in Hepes buffer	3 mg/mL	1.25 mg/mL
Membrane	60 µg/mL Perkin Elmer ES-010	20 µg/mL Perkin Elmer ES-012
Radioligand	1 nM [3H] DP-CPX (Perkin Elmer NET974) K _D = 1nM	0.1 nM [125I] MECA (Perkin Elmer NEX312) K _D = 0.8nM
Control	1 µM DPCPX (Tocris 0439)	0.1 µM IB-MECA (Tocris 1066)

20

The A_{2A}_K_i data and A_{2B}_cAMP_EC₅₀ data are provided below. The symbol “†” indicates A_{2A}_K_i or A_{2B}_cAMP_EC₅₀ ≤ 10 nM, “††” indicates A_{2A}_K_i or A_{2B}_cAMP_EC₅₀ > 10 nM but ≤ 100 nM. “†††” indicates A_{2A}_K_i or A_{2B}_cAMP_EC₅₀ > 100 nM but ≤ 1 µM; and “††††” indicates A_{2A}_K_i or A_{2B}_cAMP_EC₅₀ is greater than 1 µM.

25

Table 1.

Ex. No.	A _{2A} _Ki (nM)	A _{2B} _cAMP_EC ₅₀ (nM)
1	†	††
2	†	††
3	†	†††
4	†	††
5	†	†††
6	††	††††
7	†	†
8	†	††
9	†	††
10	†	†
11	†	†
12	†	††
13	††	†††
14	††	††††
15	†	†
16	†	†
17	†	†
18	†	†
19	†	††
20	†	†
21	†	†††
22	†	†††
23	†	†
24	†	††
25	†	††
26	†	††
27	†	†
28	†	††
29	†	†
30	†	††
31	†	††
32	†	†
33	†	†
34	†	††
35	†	†
36	†	†
37	†	†
38	†	†
39	†	†
40	†	†
41	†	†
42	†	†
43	†	†

44	†	†
45	†	†
46	†	†
47	†	††
48	†	††
49	†	†
50	†	†
51	†	†
52	†	†
53	†	†
54	†	†
55	†	†
56	†	†
57	†	†
58	†	†
59	†	††
60	†	††
61	†	†
62	†	†
63	†	†
64	†	†
65	†	††
66	†	†
67	†	†
68	†	††
69	†	†
70	†	††
71	†	†
72	†	†
73	†	†
74	†	†
75	†	†
76	†	††
77	†	†
78	†	†
79	†	†
80	†	†
81	†	†
82	†	†
83	†	†
84	†	†
85	†	††
86	†	†
87	†	††
88	†	††
89	†	††
90	†	†††


91	†	††
92	†	†
93	†	†
94	†	††
95	†	††
96	†	††
97	†	†
98	†	†
99	†	††
100	†	†††
101	†	†
102	†	†
103	†	†
104	†	††
105	†	†
106	†	†
107	†	††
108	†	†
109	†	†
110	†	†††
111	†	†
112	†	†
113	†	††
114	†	††
115	†	†
116	†	††
117	†	†
118	†	†
119	†	†
120	†	††
121	†	††
122	†	††
123	†	††
124	†	†
125	†	††
126	†	††
127	†	†
128	†	††

Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. Each

reference, including all patent, patent applications, and publications, cited in the present application is incorporated herein by reference in its entirety.

What is claimed is:

1. A compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $C(=NR^{e1})R^{b1}$, $C(=NR^{e1})NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl of R^1 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents;

each R^{a1} , R^{b1} , R^{c1} , and R^{d1} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl, wherein the C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl of R^{a1} , R^{b1} , R^{c1} , and R^{d1} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents;

each R^{e1} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{1A} is independently selected from OH, CN, halo, C_{1-3} alkyl, C_{2-3} alkenyl, C_{2-3} alkynyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, C_{1-3} haloalkoxy, amino, C_{1-3} alkylamino, and $di(C_{1-3} \text{ alkyl})\text{amino}$;

X^3 is N or CR^3 ;

R^3 is selected from H, D, halo, OH, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, cyano- C_{1-3} alkyl, HO- C_{1-3} alkyl, C_{1-3} alkoxy- C_{1-3} alkyl, C_{3-5} cycloalkyl, amino, C_{1-3} alkylamino, $di(C_{1-3} \text{ alkyl})\text{amino}$, thio, C_{1-3} alkylthio, C_{1-3} alkylsulfinyl, C_{1-3} alkylsulfonyl, carbamyl, C_{1-3} alkylcarbamyl, $di(C_{1-3} \text{ alkyl})\text{carbamyl}$, carboxy, C_{1-3} alkylcarbonyl, C_{1-4} alkoxycarbonyl, C_{1-3} alkylcarbonylamino, C_{1-3} alkoxycarbonylamino, C_{1-3} alkylcarbonyloxy, aminocarbonyloxy, C_{1-3} alkylaminocarbonyloxy, $di(C_{1-3}$

alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;

Cy¹ is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of Cy¹ are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R⁷ substituents;

each R⁷ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a7}, SR^{a7}, NHOR^{a7}, C(O)R^{b7}, C(O)NR^{c7}R^{d7}, C(O)NR^{c7}(OR^{a7}), C(O)OR^{a7}, OC(O)R^{b7}, OC(O)NR^{c7}R^{d7}, NR^{c7}R^{d7}, NR^{c7}NR^{c7}R^{d7}, NR^{c7}C(O)R^{b7}, NR^{c7}C(O)OR^{a7}, NR^{c7}C(O)NR^{c7}R^{d7}, C(=NR^{e7})R^{b7}, C(=NOH)R^{b7}, C(=NCN)R^{b7}, C(=NR^{e7})NR^{c7}R^{d7}, NR^{c7}C(=NR^{e7})NR^{c7}R^{d7}, NR^{c7}C(=NOH)NR^{c7}R^{d7}, NR^{c7}C(=NCN)NR^{c7}R^{d7}, NR^{c7}C(=NR^{e7})R^{b7}, NR^{c7}S(O)NR^{c7}R^{d7}, NR^{c7}S(O)R^{b7}, NR^{c7}S(O)₂R^{b7}, NR^{c7}S(O)(=NR^{e7})R^{b7}, NR^{c7}S(O)₂NR^{c7}R^{d7}, S(O)R^{b7}, S(O)NR^{c7}R^{d7}, S(O)₂R^{b7}, S(O)₂NR^{c7}R^{d7}, OS(O)(=NR^{e7})R^{b7}, OS(O)₂R^{b7}, SF₅, P(O)R^{f7}R^{g7}, OP(O)(OR^{h7})(ORⁱ⁷), P(O)(OR^{h7})(ORⁱ⁷), and BR^{j7}R^{k7}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁷ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{7A} substituents;

each R^{a7}, R^{b7}, R^{c7}, and R^{d7} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆

alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a7}, R^{b7}, R^{c7} and R^{d7} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{7A} substituents;

or, any R^{c7} and R^{d7}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{7A} substituents;

each R^{e7} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f7} and R^{g7} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h7} and Rⁱ⁷ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j7} and R^{k7} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j7} and R^{k7} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{7A} is independently selected from OH, NO₂, CN, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, cyano-C₁₋₆ alkyl, HO-C₁₋₆ alkyl, C₁₋₆ alkoxy-C₁₋₆ alkyl, C₃₋₅ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, amino, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, thio, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, carbamyl, C₁₋₆ alkylcarbamyl, di(C₁₋₆ alkyl)carbamyl, carboxy, C₁₋₆ alkylcarbonyl, C₁₋₄ alkoxy carbonyl, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxy carbonylamino, C₁₋₆ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₆ alkylaminocarbonyloxy, di(C₁₋₆ alkyl)aminocarbonyloxy, C₁₋₆ alkylsulfonylamino, aminosulfonyl, C₁₋₆ alkylaminosulfonyl, di(C₁₋₆ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₆

alkylaminosulfonylamino, di(C₁₋₆ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₆ alkylaminocarbonylamino, and di(C₁₋₆ alkyl)aminocarbonylamino;

R² is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a2}, SR^{a2}, NHOR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)NR^{c2}(OR^{a2}), C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, C(=NR^{e2})R^{b2}, C(=NOH)R^{b2}, C(=NCN)R^{b2}, C(=NR^{e2})NR^{c2}R^{d2}, NR^{c2}C(=NR^{e2})NR^{c2}R^{d2}, NR^{c2}C(=NOH)NR^{c2}R^{d2}, NR^{c2}C(=NCN)NR^{c2}R^{d2}, NR^{c2}C(=NR^{e2})R^{b2}, NR^{c2}S(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)(=NR^{e2})R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, S(O)₂NR^{c2}R^{d2}, OS(O)(=NR^{e2})R^{b2}, OS(O)₂R^{b2}, SF₅, P(O)R^{f2}R^{g2}, OP(O)(OR^{h2})(ORⁱ²), P(O)(OR^{h2})(ORⁱ²), and BR^{j2}R^{k2}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R² are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

R⁴ is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a4}, SR^{a4}, NHOR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)NR^{c4}(OR^{a4}), C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, C(=NR^{e4})R^{b4}, C(=NOH)R^{b4}, C(=NCN)R^{b4}, C(=NR^{e4})NR^{c4}R^{d4}, NR^{c4}C(=NR^{e4})NR^{c4}R^{d4}, NR^{c4}C(=NOH)NR^{c4}R^{d4}, NR^{c4}C(=NCN)NR^{c4}R^{d4}, NR^{c4}C(=NR^{e4})R^{b4}, NR^{c4}S(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)(=NR^{e4})R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4}, S(O)₂NR^{c4}R^{d4}, OS(O)(=NR^{e4})R^{b4}, OS(O)₂R^{b4}, SF₅, P(O)R^{f4}R^{g4}, OP(O)(OR^{h4})(ORⁱ⁴), P(O)(OR^{h4})(ORⁱ⁴), and BR^{j4}R^{k4}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆

alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4A} substituents;

provided that:

(a) when R² is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of R² are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

then R⁴ is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a4}, SR^{a4}, NHOR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)NR^{c4}(OR^{a4}), C(O)OR^{a4}, OC(O)R^{b4}, OC(O)NR^{c4}R^{d4}, NR^{c4}R^{d4}, NR^{c4}NR^{c4}R^{d4}, NR^{c4}C(O)R^{b4}, NR^{c4}C(O)OR^{a4}, NR^{c4}C(O)NR^{c4}R^{d4}, C(=NR^{e4})R^{b4}, C(=NOH)R^{b4}, C(=NCN)R^{b4}, C(=NR^{e4})NR^{c4}R^{d4}, NR^{c4}C(=NR^{e4})NR^{c4}R^{d4}, NR^{c4}C(=NOH)NR^{c4}R^{d4}, NR^{c4}C(=NCN)NR^{c4}R^{d4}, NR^{c4}C(=NR^{e4})R^{b4}, NR^{c4}S(O)NR^{c4}R^{d4}, NR^{c4}S(O)R^{b4}, NR^{c4}S(O)₂R^{b4}, NR^{c4}S(O)(=NR^{e4})R^{b4}, NR^{c4}S(O)₂NR^{c4}R^{d4}, S(O)R^{b4}, S(O)NR^{c4}R^{d4}, S(O)₂R^{b4}, S(O)₂NR^{c4}R^{d4}, OS(O)(=NR^{e4})R^{b4}, OS(O)₂R^{b4}, SF₅, P(O)R^{f4}R^{g4}, OP(O)(OR^{h4})(ORⁱ⁴), P(O)(OR^{h4})(ORⁱ⁴), and BR^{j4}R^{k4}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4A} substituents;

or, alternatively,

(b) when R² is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a2}, SR^{a2}, NHOR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)NR^{c2}(OR^{a2}), C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, NR^{c2}R^{d2},

$\text{NR}^{\text{c}2}\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$, $\text{NR}^{\text{c}2}\text{C}(\text{O})\text{R}^{\text{b}2}$, $\text{NR}^{\text{c}2}\text{C}(\text{O})\text{OR}^{\text{a}2}$, $\text{NR}^{\text{c}2}\text{C}(\text{O})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$,
 $\text{C}(=\text{NR}^{\text{e}2})\text{R}^{\text{b}2}$, $\text{C}(=\text{NOH})\text{R}^{\text{b}2}$, $\text{C}(=\text{NCN})\text{R}^{\text{b}2}$, $\text{C}(=\text{NR}^{\text{e}2})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$,
 $\text{NR}^{\text{c}2}\text{C}(=\text{NR}^{\text{e}2})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$, $\text{NR}^{\text{c}2}\text{C}(=\text{NOH})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$, $\text{NR}^{\text{c}2}\text{C}(=\text{NCN})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$,
 $\text{NR}^{\text{c}2}\text{C}(=\text{NR}^{\text{e}2})\text{R}^{\text{b}2}$, $\text{NR}^{\text{c}2}\text{S}(\text{O})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$, $\text{NR}^{\text{c}2}\text{S}(\text{O})\text{R}^{\text{b}2}$, $\text{NR}^{\text{c}2}\text{S}(\text{O})_2\text{R}^{\text{b}2}$,
 $\text{NR}^{\text{c}2}\text{S}(\text{O})(=\text{NR}^{\text{e}2})\text{R}^{\text{b}2}$, $\text{NR}^{\text{c}2}\text{S}(\text{O})_2\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$, $\text{S}(\text{O})\text{R}^{\text{b}2}$, $\text{S}(\text{O})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$, $\text{S}(\text{O})_2\text{R}^{\text{b}2}$,
 $\text{S}(\text{O})_2\text{NR}^{\text{c}2}\text{R}^{\text{d}2}$, $\text{OS}(\text{O})(=\text{NR}^{\text{e}2})\text{R}^{\text{b}2}$, $\text{OS}(\text{O})_2\text{R}^{\text{b}2}$, SF_5 , $\text{P}(\text{O})\text{R}^{\text{f}2}\text{R}^{\text{g}2}$,
 $\text{OP}(\text{O})(\text{OR}^{\text{h}2})(\text{OR}^{\text{i}2})$, $\text{P}(\text{O})(\text{OR}^{\text{h}2})(\text{OR}^{\text{i}2})$, and $\text{BR}^{\text{j}2}\text{R}^{\text{k}2}$, wherein the C_{1-6} alkyl,
 C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-
10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)-
 C_{1-6} alkyl- of R^2 are each optionally substituted with 1, 2, 3, 4, 5, or 6
independently selected $\text{R}^{2\text{A}}$ substituents;

then R^4 is selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl,
 C_{2-6} alkynyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10
membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6}
alkyl-, CN, NO_2 , $\text{OR}^{\text{a}4}$, $\text{SR}^{\text{a}4}$, $\text{NHOR}^{\text{a}4}$, $\text{C}(\text{O})\text{R}^{\text{b}4}$, $\text{C}(\text{O})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$,
 $\text{C}(\text{O})\text{NR}^{\text{c}4}(\text{OR}^{\text{a}4})$, $\text{C}(\text{O})\text{OR}^{\text{a}4}$, $\text{OC}(\text{O})\text{R}^{\text{b}4}$, $\text{OC}(\text{O})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$, $\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$,
 $\text{NR}^{\text{c}4}\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$, $\text{NR}^{\text{c}4}\text{C}(\text{O})\text{R}^{\text{b}4}$, $\text{NR}^{\text{c}4}\text{C}(\text{O})\text{OR}^{\text{a}4}$, $\text{NR}^{\text{c}4}\text{C}(\text{O})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$,
 $\text{C}(=\text{NR}^{\text{e}4})\text{R}^{\text{b}4}$, $\text{C}(=\text{NOH})\text{R}^{\text{b}4}$, $\text{C}(=\text{NCN})\text{R}^{\text{b}4}$, $\text{C}(=\text{NR}^{\text{e}4})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$,
 $\text{NR}^{\text{c}4}\text{C}(=\text{NR}^{\text{e}4})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$, $\text{NR}^{\text{c}4}\text{C}(=\text{NOH})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$, $\text{NR}^{\text{c}4}\text{C}(=\text{NCN})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$,
 $\text{NR}^{\text{c}4}\text{C}(=\text{NR}^{\text{e}4})\text{R}^{\text{b}4}$, $\text{NR}^{\text{c}4}\text{S}(\text{O})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$, $\text{NR}^{\text{c}4}\text{S}(\text{O})\text{R}^{\text{b}4}$, $\text{NR}^{\text{c}4}\text{S}(\text{O})_2\text{R}^{\text{b}4}$,
 $\text{NR}^{\text{c}4}\text{S}(\text{O})(=\text{NR}^{\text{e}4})\text{R}^{\text{b}4}$, $\text{NR}^{\text{c}4}\text{S}(\text{O})_2\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$, $\text{S}(\text{O})\text{R}^{\text{b}4}$, $\text{S}(\text{O})\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$, $\text{S}(\text{O})_2\text{R}^{\text{b}4}$,
 $\text{S}(\text{O})_2\text{NR}^{\text{c}4}\text{R}^{\text{d}4}$, $\text{OS}(\text{O})(=\text{NR}^{\text{e}4})\text{R}^{\text{b}4}$, $\text{OS}(\text{O})_2\text{R}^{\text{b}4}$, SF_5 , $\text{P}(\text{O})\text{R}^{\text{f}4}\text{R}^{\text{g}4}$,
 $\text{OP}(\text{O})(\text{OR}^{\text{h}4})(\text{OR}^{\text{i}4})$, $\text{P}(\text{O})(\text{OR}^{\text{h}4})(\text{OR}^{\text{i}4})$, and $\text{BR}^{\text{j}4}\text{R}^{\text{k}4}$, wherein the C_{1-6} alkyl,
 C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-
10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)-
 C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, 3, 4, 5, or 6
independently selected $\text{R}^{4\text{A}}$ substituents;

each $\text{R}^{\text{a}2}$, $\text{R}^{\text{b}2}$, $\text{R}^{\text{c}2}$, and $\text{R}^{\text{d}2}$ is independently selected from H, C_{1-6} alkyl, C_{1-6}
haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered
heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl-
 C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered
heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10}

aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a2}, R^{b2}, R^{c2} and R^{d2} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

or, any R^{c2} and R^{d2}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

each R^{e2} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f2} and R^{g2} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h2} and Rⁱ² is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j2} and R^{k2} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j2} and R^{k2} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)NR^{c21}(OR^{a21}), C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}NR^{c21}R^{d21},

$\text{NR}^{\text{c}21}\text{C}(\text{O})\text{R}^{\text{b}21}$, $\text{NR}^{\text{c}21}\text{C}(\text{O})\text{OR}^{\text{a}21}$, $\text{NR}^{\text{c}21}\text{C}(\text{O})\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$, $\text{C}(=\text{NR}^{\text{e}21})\text{R}^{\text{b}21}$,
 $\text{C}(=\text{NOH})\text{R}^{\text{b}21}$, $\text{C}(=\text{NCN})\text{R}^{\text{b}21}$, $\text{C}(=\text{NR}^{\text{e}21})\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$, $\text{NR}^{\text{c}21}\text{C}(=\text{NR}^{\text{e}21})\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$,
 $\text{NR}^{\text{c}21}\text{C}(=\text{NOH})\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$, $\text{NR}^{\text{c}21}\text{C}(=\text{NCN})\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$, $\text{NR}^{\text{c}21}\text{C}(=\text{NR}^{\text{e}21})\text{R}^{\text{b}21}$,
 $\text{NR}^{\text{c}21}\text{S}(\text{O})\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$, $\text{NR}^{\text{c}21}\text{S}(\text{O})\text{R}^{\text{b}21}$, $\text{NR}^{\text{c}21}\text{S}(\text{O})_2\text{R}^{\text{b}21}$, $\text{NR}^{\text{c}21}\text{S}(\text{O})(=\text{NR}^{\text{e}21})\text{R}^{\text{b}21}$,
 $\text{NR}^{\text{c}21}\text{S}(\text{O})_2\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$, $\text{S}(\text{O})\text{R}^{\text{b}21}$, $\text{S}(\text{O})\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$, $\text{S}(\text{O})_2\text{R}^{\text{b}21}$, $\text{S}(\text{O})_2\text{NR}^{\text{c}21}\text{R}^{\text{d}21}$,
 $\text{OS}(\text{O})(=\text{NR}^{\text{e}21})\text{R}^{\text{b}21}$, $\text{OS}(\text{O})_2\text{R}^{\text{b}21}$, SF_5 , $\text{P}(\text{O})\text{R}^{\text{f}21}\text{R}^{\text{g}21}$, $\text{OP}(\text{O})(\text{OR}^{\text{h}21})(\text{OR}^{\text{i}21})$,
 $\text{P}(\text{O})(\text{OR}^{\text{h}21})(\text{OR}^{\text{i}21})$, and $\text{BR}^{\text{j}21}\text{R}^{\text{k}21}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,
 C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered
heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered
heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of $\text{R}^{2\text{A}}$ are
each optionally substituted with 1, 2, 3, or 4 independently selected $\text{R}^{2\text{B}}$ substituents;
each $\text{R}^{\text{a}21}$, $\text{R}^{\text{b}21}$, $\text{R}^{\text{c}21}$ and $\text{R}^{\text{d}21}$ is independently selected from H, C_{1-6} alkyl, C_{1-6}
haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered
heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl-
 C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered
heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10}
aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl,
 C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6}
alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of $\text{R}^{\text{a}21}$, $\text{R}^{\text{b}21}$, $\text{R}^{\text{c}21}$ and $\text{R}^{\text{d}21}$
are each optionally substituted with 1, 2, 3, or 4 independently selected $\text{R}^{2\text{B}}$
substituents;

or, any $\text{R}^{\text{c}21}$ and $\text{R}^{\text{d}21}$, attached to the same N atom, together with the N atom
to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein
the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4
independently selected $\text{R}^{2\text{B}}$ substituents;

each $\text{R}^{\text{e}21}$ is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy,
 C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each $\text{R}^{\text{f}21}$ and $\text{R}^{\text{g}21}$ is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy,
 C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl,
5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-,
 C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10
membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{h21} and R^{i21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{j21} and R^{k21} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j21} and R^{k21} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

each R^{2B} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a22} , SR^{a22} , $NHOR^{a22}$, $C(O)R^{b22}$, $C(O)NR^{c22}R^{d22}$, $C(O)NR^{c22}(OR^{a22})$, $C(O)OR^{a22}$, $OC(O)R^{b22}$, $OC(O)NR^{c22}R^{d22}$, $NR^{c22}R^{d22}$, $NR^{c22}NR^{c22}R^{d22}$, $NR^{c22}C(O)R^{b22}$, $NR^{c22}C(O)OR^{a22}$, $NR^{c22}C(O)NR^{c22}R^{d22}$, $C(=NR^{e22})R^{b22}$, $C(=NOH)R^{b22}$, $C(=NCN)R^{b22}$, $C(=NR^{e22})NR^{c22}R^{d22}$, $NR^{c22}C(=NR^{e22})NR^{c22}R^{d22}$, $NR^{c22}C(=NOH)NR^{c22}R^{d22}$, $NR^{c22}C(=NCN)NR^{c22}R^{d22}$, $NR^{c22}C(=NR^{e22})R^{b22}$, $NR^{c22}S(O)NR^{c22}R^{d22}$, $NR^{c22}S(O)R^{b22}$, $NR^{c22}S(O)_2R^{b22}$, $NR^{c22}S(O)(=NR^{e22})R^{b22}$, $NR^{c22}S(O)_2NR^{c22}R^{d22}$, $S(O)R^{b22}$, $S(O)NR^{c22}R^{d22}$, $S(O)_2R^{b22}$, $S(O)_2NR^{c22}R^{d22}$, $OS(O)(=NR^{e22})R^{b22}$, $OS(O)_2R^{b22}$, SF_5 , $P(O)R^{f22}R^{g22}$, $OP(O)(OR^{h22})(OR^{i22})$, $P(O)(OR^{h22})(OR^{i22})$, and $BR^{j22}R^{k22}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{2B} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

each R^{a22} , R^{b22} , R^{c22} and R^{d22} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10}

aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a22}, R^{b22}, R^{c22} and R^{d22} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

or, any R^{c22} and R^{d22}, attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group, wherein the 4-, 5-, 6-, or 7-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

each R^{e22} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f22} and R^{g22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h22} and Rⁱ²² is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j22} and R^{k22} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j22} and R^{k22} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{2C} is independently selected from OH, NO₂, CN, halo, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, C₁₋₃ haloalkyl, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, C₁₋₃ alkoxy, C₁₋₃ haloalkoxy, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃

alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;

each R^{a4}, R^{b4}, R^{c4}, and R^{d4} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a4}, R^{b4}, R^{c4} and R^{d4} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4A} substituents;

or, any R^{c4} and R^{d4}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4A} substituents;

each R^{e4} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f4} and R^{g4} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h4} and Rⁱ⁴ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j4} and R^{k4} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j4} and R^{k4} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a41} , SR^{a41} , $NHOR^{a41}$, $C(O)R^{b41}$, $C(O)NR^{c41}R^{d41}$, $C(O)NR^{c41}(OR^{a41})$, $C(O)OR^{a41}$, $OC(O)R^{b41}$, $OC(O)NR^{c41}R^{d41}$, $NR^{c41}R^{d41}$, $NR^{c41}NR^{c41}R^{d41}$, $NR^{c41}C(O)R^{b41}$, $NR^{c41}C(O)OR^{a41}$, $NR^{c41}C(O)NR^{c41}R^{d41}$, $C(=NR^{e41})R^{b41}$, $C(=NOH)R^{b41}$, $C(=NCN)R^{b41}$, $C(=NR^{e41})NR^{c41}R^{d41}$, $NR^{c41}C(=NR^{e41})NR^{c41}R^{d41}$, $NR^{c41}C(=NOH)NR^{c41}R^{d41}$, $NR^{c41}C(=NCN)NR^{c41}R^{d41}$, $NR^{c41}C(=NR^{e41})R^{b41}$, $NR^{c41}S(O)NR^{c41}R^{d41}$, $NR^{c41}S(O)R^{b41}$, $NR^{c41}S(O)_2R^{b41}$, $NR^{c41}S(O)(=NR^{e41})R^{b41}$, $NR^{c41}S(O)_2NR^{c41}R^{d41}$, $S(O)R^{b41}$, $S(O)NR^{c41}R^{d41}$, $S(O)_2R^{b41}$, $S(O)_2NR^{c41}R^{d41}$, $OS(O)(=NR^{e41})R^{b41}$, $OS(O)_2R^{b41}$, SF_5 , $P(O)R^{f41}R^{g41}$, $OP(O)(OR^{h41})(OR^{i41})$, $P(O)(OR^{h41})(OR^{i41})$, and $BR^{j41}R^{k41}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{4A} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a41} , R^{b41} , R^{c41} and R^{d41} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4B} substituents;

or, any R^{c41} and R^{d41} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{4B} substituents;

each R^{e41} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{f41} and R^{g41} is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{h41} and R^{i41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{j41} and R^{k41} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j41} and R^{k41} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

each R^{4B} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a42} , SR^{a42} , $NHOR^{a42}$, $C(O)R^{b42}$, $C(O)NR^{c42}R^{d42}$, $C(O)NR^{c42}(OR^{a42})$, $C(O)OR^{a42}$, $OC(O)R^{b42}$, $OC(O)NR^{c42}R^{d42}$, $NR^{c42}R^{d42}$, $NR^{c42}NR^{c42}R^{d42}$, $NR^{c42}C(O)R^{b42}$, $NR^{c42}C(O)OR^{a42}$, $NR^{c42}C(O)NR^{c42}R^{d42}$, $C(=NR^{e42})R^{b42}$, $C(=NOH)R^{b42}$, $C(=NCN)R^{b42}$, $C(=NR^{e42})NR^{c42}R^{d42}$, $NR^{c42}C(=NR^{e42})NR^{c42}R^{d42}$, $NR^{c42}C(=NOH)NR^{c42}R^{d42}$, $NR^{c42}C(=NCN)NR^{c42}R^{d42}$, $NR^{c42}C(=NR^{e42})R^{b42}$, $NR^{c42}S(O)NR^{c42}R^{d42}$, $NR^{c42}S(O)R^{b42}$, $NR^{c42}S(O)_2R^{b42}$, $NR^{c42}S(O)(=NR^{e42})R^{b42}$, $NR^{c42}S(O)_2NR^{c42}R^{d42}$, $S(O)R^{b42}$, $S(O)NR^{c42}R^{d42}$, $S(O)_2R^{b42}$, $S(O)_2NR^{c42}R^{d42}$,

OS(O)(=NR^{e42})R^{b42}, OS(O)₂R^{b42}, SF₅, P(O)R^{f42}R^{g42}, OP(O)(OR^{h42})(ORⁱ⁴²), P(O)(OR^{h42})(ORⁱ⁴²), and BR^{j42}R^{k42}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{4B} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4C} substituents;

each R^{a42}, R^{b42}, R^{c42} and R^{d42} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a42}, R^{b42}, R^{c42} and R^{d42} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4C} substituents;

or, any R^{c42} and R^{d42}, attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group, wherein the 4-, 5-, 6-, or 7-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4C} substituents;

each R^{e42} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f42} and R^{g42} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h42} and Rⁱ⁴² is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j42} and R^{k42} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j42} and R^{k42} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

each R^{4C} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a43} , SR^{a43} , $NHOR^{a43}$, $C(O)R^{b43}$, $C(O)NR^{c43}R^{d43}$, $C(O)NR^{c43}(OR^{a43})$, $C(O)OR^{a43}$, $OC(O)R^{b43}$, $OC(O)NR^{c43}R^{d43}$, $NR^{c43}R^{d43}$, $NR^{c43}NR^{c43}R^{d43}$, $NR^{c43}C(O)R^{b43}$, $NR^{c43}C(O)OR^{a43}$, $NR^{c43}C(O)NR^{c43}R^{d43}$, $C(=NR^{e43})R^{b43}$, $C(=NOH)R^{b43}$, $C(=NCN)R^{b43}$, $C(=NR^{e43})NR^{c43}R^{d43}$, $NR^{c43}C(=NR^{e43})NR^{c43}R^{d43}$, $NR^{c43}C(=NOH)NR^{c43}R^{d43}$, $NR^{c43}C(=NCN)NR^{c43}R^{d43}$, $NR^{c43}C(=NR^{e43})R^{b43}$, $NR^{c43}S(O)NR^{c43}R^{d43}$, $NR^{c43}S(O)R^{b43}$, $NR^{c43}S(O)_2R^{b43}$, $NR^{c43}S(O)(=NR^{e43})R^{b43}$, $NR^{c43}S(O)_2NR^{c43}R^{d43}$, $S(O)R^{b43}$, $S(O)NR^{c43}R^{d43}$, $S(O)_2R^{b43}$, $S(O)_2NR^{c43}R^{d43}$, $OS(O)(=NR^{e43})R^{b43}$, $OS(O)_2R^{b43}$, SF_5 , $P(O)R^{f43}R^{g43}$, $OP(O)(OR^{h43})(OR^{i43})$, $P(O)(OR^{h43})(OR^{i43})$, and $BR^{j43}R^{k43}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{4C} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents;

each R^{a43} , R^{b43} , R^{c43} and R^{d43} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a43} , R^{b43} , R^{c43} and R^{d43}

are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43} , attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group, wherein the 4-, 5-, 6-, or 7-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents;

each R^{e43} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{f43} and R^{g43} is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{h43} and R^{i43} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{j43} and R^{k43} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j43} and R^{k43} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

each R^{4D} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6} alkyl-, (5-7 membered heteroaryl)- C_{1-6} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a44} , SR^{a44} , $NHOR^{a44}$, $C(O)R^{b44}$, $C(O)NR^{c44}R^{d44}$, $C(O)NR^{c44}(OR^{a44})$, $C(O)OR^{a44}$, $OC(O)R^{b44}$, $OC(O)NR^{c44}R^{d44}$, $NR^{c44}R^{d44}$, $NR^{c44}NR^{c44}R^{d44}$, $NR^{c44}C(O)R^{b44}$, $NR^{c44}C(O)OR^{a44}$, $NR^{c44}C(O)NR^{c44}R^{d44}$, $C(=NR^{e44})R^{b44}$, $C(=NOH)R^{b44}$, $C(=NCN)R^{b44}$, $C(=NR^{e44})NR^{c44}R^{d44}$, $NR^{c44}C(=NR^{e44})NR^{c44}R^{d44}$, $NR^{c44}C(=NOH)NR^{c44}R^{d44}$, $NR^{c44}C(=NCN)NR^{c44}R^{d44}$, $NR^{c44}C(=NR^{e44})R^{b44}$,

NR^{c44}S(O)NR^{c44}R^{d44}, NR^{c44}S(O)R^{b44}, NR^{c44}S(O)₂R^{b44}, NR^{c44}S(O)(=NR^{e44})R^{b44}, NR^{c44}S(O)₂NR^{c44}R^{d44}, S(O)R^{b44}, S(O)NR^{c44}R^{d44}, S(O)₂R^{b44}, S(O)₂NR^{c44}R^{d44}, OS(O)(=NR^{e44})R^{b44}, OS(O)₂R^{b44}, SF₅, P(O)R^{f44}R^{g44}, OP(O)(OR^{h44})(ORⁱ⁴⁴), P(O)(OR^{h44})(ORⁱ⁴⁴), and BRⁱ⁴⁴R^{k44}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-7 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{4D} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4E} substituents;

each R^{a44}, R^{b44}, R^{c44} and R^{d44} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-7 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-7 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a44}, R^{b44}, R^{c44} and R^{d44} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4E} substituents;

or, any R^{c44} and R^{d44}, attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, 6-, or 7-membered heterocycloalkyl group, wherein the 4-, 5-, 6-, or 7-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4E} substituents;

each R^{e44} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f44} and R^{g44} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-7 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h44} and Rⁱ⁴⁴ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-7 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-7 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-;


each R^{j44} and R^{k44} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j44} and R^{k44} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl; and

each R^{4E} is independently selected from OH, NO₂, CN, halo, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, C₁₋₃ haloalkyl, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, C₁₋₃ alkoxy, C₁₋₃ haloalkoxy, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino.

2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein 1, 2, 3, 4, 5, 6, 7, or 8 hydrogen atoms, attached to carbon atoms of any alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, or heteroaryl substituents or -C₁₋₆ alkyl-, alkylene, alkenylene, and alkynylene linking groups, are each optionally replaced by a deuterium atom.

3. The compound of claim 1 or 2, which is a compound of Formula (IIa):



or a pharmaceutically acceptable salt thereof.

4. The compound of any one of claims 1 to 3, or a pharmaceutically acceptable salt thereof, wherein R^3 is CN or H.

5. The compound of any one of claims 1 to 3, or a pharmaceutically acceptable salt thereof, wherein R^3 is H.

6. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl, wherein the phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$, wherein said phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl of R^{2A} are optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl; and

each R^{2B} is independently selected from halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and di(C_{1-3} alkyl)amino.

7. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl, wherein the C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl of R^2 are each optionally substituted with 1 or 2 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C_{1-4} alkyl, C_{1-4} haloalkyl, and $C(O)NR^{c21}R^{d21}$, wherein said C_{1-4} alkyl of R^{2A} is optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl; and

each R^{2B} is independently selected from halo and OH.

8. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from cyclopropyl, oxazolyl, triazolyl, pyrazolyl, pyridyl, pyrimidinyl, dihydropyridin-(2H)-yl, and pyridinonyl, each of which is optionally substituted with 1 or 2 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, and $S(O)R^{b21}$, wherein said C_{1-6} alkyl of R^{2A} is optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl; and

each R^{2B} is independently selected from halo and OH.

9. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from 1-(trifluoromethyl)cycloprop-1-yl, 1-ethyl-1H-pyrazol-5-yl, 1-propyl-1H-pyrazol-5-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl,

pyrimidin-4-yl, 3,6-dihydropyridin-4-yl-1(2H)-carboxamide, pyridin-4-yl, 4-(1-hydroxyethyl)-2-methyloxazol-5-yl, 2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl, 1-ethyl-1*H*-1,2,3-triazol-5-yl, and 1-methyl-1*H*-1,2,3-triazol-5-yl.

10. The compound of any one of claims 1 to 9, or a pharmaceutically acceptable salt thereof, wherein R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a4} , SR^{a4} , $NHOR^{a4}$, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl, wherein said C_{1-6} alkyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl of R^{a4} , R^{b4} , R^{c4} , and R^{d4} are each optionally substituted by 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, CN, NO_2 , OR^{a41} , SR^{a41} , $NHOR^{a41}$, $C(O)R^{b41}$, $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $OC(O)R^{b41}$, $OC(O)NR^{c41}R^{d41}$, $NR^{c41}R^{d41}$, $NR^{c41}C(O)R^{b41}$, $NR^{c41}C(O)OR^{a41}$, $NR^{c41}C(O)NR^{c41}R^{d41}$, $S(O)R^{b41}$, and $S(O)_2R^{b41}$, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl of R^{4A} are each optionally substituted by 1 or 2 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} , and R^{d41} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein said C_{1-6} alkyl of R^{a41} , R^{b41} , R^{c41} , and R^{d41} is optionally substituted by 1, 2, or 3 independently selected R^{4B} substituents; and

each R^{4B} is independently selected from halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and $di(C_{1-3}$ alkyl)amino.

11. The compound of any one of claims 1 to 9, or a pharmaceutically acceptable salt thereof, wherein R^4 is selected from phenyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, and $NR^{c4}R^{d4}$, wherein the phenyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, and (4-7 membered heterocycloalkyl)- C_{1-3} alkyl- of R^4 are each optionally substituted with 1 or 2 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, phenyl, and 4-7 membered heterocycloalkyl, wherein said C_{1-6} alkyl, phenyl, and 4-7 membered heterocycloalkyl of R^{a4} , R^{b4} , R^{c4} , and R^{d4} are optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, 4-7 membered heterocycloalkyl, and OH, wherein said 4-7 membered heterocycloalkyl of R^{4A} is optionally substituted with 1 or 2 independently selected R^{4B} substituents; and

each R^{4B} is independently selected from halo, OH, and C_{1-4} alkyl.

12. The compound of any one of claims 1 to 9, or a pharmaceutically acceptable salt thereof, wherein R^4 is selected from $C(O)NHC_{1-6}$ alkyl, $C(O)$ -azetidiny, $C(O)$ -pyrrolidinyl, $C(O)$ -piperidinyl, $C(O)N(C_{1-6}$ alkyl)₂, $NHC(O)OC_{1-6}$ alkyl, (azetidiny)- C_{1-6} alkyl, (pyridyl)- C_{1-6} alkyl, (phenyl)- C_{1-6} alkyl, (fluorophenyl)- C_{1-6} alkyl, 3,6-dihydro-2H-pyranyl, NH-(phenyl), pyridyl, and (pyrrolo[3,2-b]pyridiny)- C_{1-6} alkyl,

wherein each C_{1-6} alkyl and azetidiny group is optionally substituted by 1 or 2 OH groups; and each pyridyl is optionally substituted by a methylpiperazinyl group.

13. The compound of any one of claims 1 to 9, or a pharmaceutically acceptable salt thereof, wherein R^4 is selected from $C(O)NHCH_2CH_3$, $C(O)N(CH_3)(CH_2CH_3)$, $C(O)N(CH_2CH_3)_2$, $NHC(O)OCH_2CH_3$, $C(O)$ -azetidiny, $C(O)$ -hydroxyazetidiny,

C(O)-pyrrolidinyl, C(O)-piperidinyl, CH₂-azetidiny, CH₂-pyridyl, CH₂-fluorophenyl, CH(OH)-fluorophenyl, NH-phenyl, 3,6-dihydro-2H-pyranyl, (methylpiperazinyl)pyridinyl, and (1*H*-pyrrolo[3,2-*b*]pyridin-3-yl)methyl.

14. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein R² is selected from H, D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, and CN.

15. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein R² is selected from H, difluoroethyl, bromo, and CN.

16. The compound of any one of claims 14 and 15, or a pharmaceutically acceptable salt thereof, wherein R⁴ is selected from phenyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and C(O)NR^{c4}R^{d4}, wherein the phenyl-C₁₋₆ alkyl- and (5-6 membered heteroaryl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1 or 2 substituents independently selected from OH and halo; and

each R^{a4}, R^{c4}, and R^{d4} is independently selected from H and C₁₋₆ alkyl.

17. The compound of any one of claims 14 and 15, or a pharmaceutically acceptable salt thereof, wherein R⁴ is selected from -C(O)NHCH₂CH₃, -CH₂-pyridyl, CH₂-fluorophenyl, and CH(OH)-fluorophenyl.

18. The compound of claim 3, or a pharmaceutically acceptable salt thereof, wherein:

R³ is H or CN;

Cy¹ is phenyl which is substituted by 1 or 2 independently selected R⁷ substituents;

each R⁷ is independently selected from halo and CN;

R² is selected from phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl, wherein the phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl of R² are each optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$, wherein said phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl and 4-7 membered heterocycloalkyl of R^{2A} are optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl;

each R^{2B} is independently selected from halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and $di(C_{1-3} \text{ alkyl})\text{amino}$;

R^4 is selected from H, D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a4} , SR^{a4} , $NHOR^{a4}$, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $OC(O)R^{b4}$, $OC(O)NR^{c4}R^{d4}$, $NR^{c4}R^{d4}$, $NR^{c4}C(O)R^{b4}$, $NR^{c4}C(O)OR^{a4}$, $NR^{c4}C(O)NR^{c4}R^{d4}$, $S(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl, wherein said C_{1-6} alkyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl of R^{a4} , R^{b4} , R^{c4} , and R^{d4} are each optionally substituted by 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, CN, NO_2 , OR^{a41} , SR^{a41} , $NHOR^{a41}$, $C(O)R^{b41}$,

$C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $OC(O)R^{b41}$, $OC(O)NR^{c41}R^{d41}$, $NR^{c41}R^{d41}$, $NR^{c41}C(O)R^{b41}$, $NR^{c41}C(O)OR^{a41}$, $NR^{c41}C(O)NR^{c41}R^{d41}$, $S(O)R^{b41}$, and $S(O)_2R^{b41}$, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, and 4-7 membered heterocycloalkyl of R^{4A} are each optionally substituted by 1 or 2 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} , and R^{d41} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein said C_{1-6} alkyl of R^{a41} , R^{b41} , R^{c41} , and R^{d41} is optionally substituted by 1, 2, or 3 independently selected R^{4B} substituents; and

each R^{4B} is independently selected from halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and $di(C_{1-3}$ alkyl)amino,

19. The compound of claim 3, or a pharmaceutically acceptable salt thereof, wherein:

R^3 is H or CN;

Cy^1 is 3-cyanophenyl or 3-cyano-2-fluorophenyl;

R^2 is selected from C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl, wherein the C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 4-6 membered heterocycloalkyl of R^2 are each optionally substituted with 1 or 2 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C_{1-4} alkyl, C_{1-4} haloalkyl, and $C(O)NR^{c21}R^{d21}$, wherein said C_{1-4} alkyl of R^{2A} is optionally substituted with 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl;

each R^{2B} is independently selected from halo and OH;

R^4 is selected from phenyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, and $NR^{c4}R^{d4}$, wherein the phenyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, and (4-7 membered

heterocycloalkyl)-C₁₋₃ alkyl- of R⁴ are each optionally substituted with 1 or 2 independently selected R^{4A} substituents;

each R^{a4}, R^{b4}, R^{c4}, and R^{d4} is independently selected from H, C₁₋₆ alkyl, phenyl, and 4-7 membered heterocycloalkyl, wherein said C₁₋₆ alkyl, phenyl, and 4-7 membered heterocycloalkyl of R^{a4}, R^{b4}, R^{c4}, and R^{d4} are optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, 4-7 membered heterocycloalkyl, and OH, wherein said 4-7 membered heterocycloalkyl of R^{4A} is optionally substituted with 1 or 2 independently selected R^{4B} substituents; and

each R^{4B} is independently selected from halo, OH, and C₁₋₄ alkyl.

20. The compound of claim 3, or a pharmaceutically acceptable salt thereof, wherein:

R³ is H or CN;

Cy¹ is phenyl which is substituted by 1 or 2 independently selected R⁷ substituents;


each R⁷ is independently selected from halo and CN;

R² is selected from H, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, and CN;

R⁴ is selected from phenyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and C(O)NR^{c4}R^{d4}, wherein the phenyl-C₁₋₆ alkyl- and (5-6 membered heteroaryl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1 or 2 substituents independently selected from OH and halo; and


each R^{a4}, R^{c4}, and R^{d4} is independently selected from H and C₁₋₆ alkyl.

21. The compound of any one of claims 1 to 3, which is a compound of Formula (IIb):




or a pharmaceutically acceptable salt thereof, wherein n is an integer from 0 to 4.

22. The compound of any one of claims 1 to 3, which is a compound of Formula (IIc):



or a pharmaceutically acceptable salt thereof.

23. The compound of claim 1 or 2, which is a compound of Formula (IIIa):



or a pharmaceutically acceptable salt thereof.

24. The compound of any one of claims 1 to 17 and 23, or a pharmaceutically acceptable salt thereof, wherein Cy^1 is phenyl which is substituted by 1 or 2 independently selected R^7 substituents; and

each R^7 is independently selected from halo and CN.

25. The compound of any one of claims 1 to 17 and 23, or a pharmaceutically acceptable salt thereof, wherein Cy^1 is cyanophenyl.

26. The compound of any one of claims 1 to 17 and 23, or a pharmaceutically acceptable salt thereof, wherein Cy^1 is 3-cyanophenyl or 3-cyano-2-fluorophenyl.

27. The compound of any one of claims 1, 2, and 23-26, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $S(O)R^{b21}$, and $S(O)_2R^{b21}$, wherein said C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R^{2A} are each optionally substituted by 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21} , R^{b21} , R^{c21} , and R^{d21} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein said C_{1-6} alkyl of R^{a21} , R^{b21} , R^{c21} , and R^{d21} is optionally substituted by 1, 2, or 3 independently selected R^{2A} substituents; and

each R^{2B} is independently selected from D, halo, OH, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkyl, C_{1-4} haloalkyl, amino, C_{1-3} alkylamino, and $di(C_{1-3}$ alkyl)amino.

28. The compound of any one of claims 1, 2, and 23-26, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from 5-10 membered heteroaryl and 4-7 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-7 membered heterocycloalkyl of R^2 are each optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, and OH, wherein said C_{1-6} alkyl of R^{2A} is optionally substituted by 1, 2, or 3 independently selected R^{2B} substituents; and

each R^{2B} is independently selected from D, halo, and OH.

29. The compound of any one of claims 1, 2, and 23-26, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from 6-oxo-1,6-dihydropyridin-3-yl, pyrimidin-4-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, 1-methyl-6-oxo-1,6-dihydropyridazin-3-yl, 4-methyloxazol-5-yl, 4-ethyloxazol-5-yl, 3-methylpyridin-4-yl, 4-(2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl, 2-methyl-4-(2,2,2-trifluoro-1-hydroxyethyl)oxazol-5-yl, 1-ethyl-1H-pyrazol-5-yl, 6-hydroxypyridin-3-yl, 2,6-dimethylpyridin-4-yl, 3-methyl-1H-pyrazol-4-yl, [1,2,4]triazolo[4,3-*a*]pyridin-6-yl, oxazol-5-yl, imidazo[1,2-*a*]pyridine-6-yl, 3-fluoropyridin-4-yl, and 1-(methyl-*d*3)-6-oxo-1,6-dihydropyridazin-3-yl.

30. The compound of any one of claims 1, 2, and 23-26, or a pharmaceutically acceptable salt thereof, wherein R^2 is H.

31. The compound of any one of claims 1, 2, and 23 to 26, or a pharmaceutically acceptable salt thereof, wherein R^2 is 6-oxo-1,6-dihydropyridin-3-yl or imidazo[1,2-*a*]pyridin-6-yl; and R^4 is H.

32. The compound of any one of claims 1, 2, and 23 to 26, or a pharmaceutically acceptable salt thereof, wherein R^2 is H; and R^4 is $-NHC(O)OC_{1-6}$ alkyl.

33. The compound of any one of claims 1, 2, and 23-30, or a pharmaceutically acceptable salt thereof, wherein:

R^4 is selected from C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, OR^{a4} , $C(O)R^{b4}$, $C(O)NR^{c4}R^{d4}$, $C(O)OR^{a4}$, $NR^{c4}R^{d4}$, $S(O)_2NR^{c4}R^{d4}$, and $S(O)_2R^{b4}$, wherein the C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^4 are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a4} , R^{b4} , R^{c4} and R^{d4} are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, OR^{a41} , $C(O)R^{b41}$, $C(O)NR^{c41}R^{d41}$, $C(O)OR^{a41}$, $NR^{c41}R^{d41}$, $S(O)_2NR^{c41}R^{d41}$, and $S(O)_2R^{b41}$, wherein the C_{1-6} alkyl, 5 C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl, wherein the C_{1-6} alkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^{a41} , R^{b41} , R^{c41} and R^{d41} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, phenyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, 5-6 membered heteroaryl, phenyl- C_{1-3} alkyl-, (C_{3-6} cycloalkyl)- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3}

alkyl-, (5-6 membered heteroaryl)-C₁₋₃ alkyl-, CN, OR^{a42}, C(O)R^{b42}, C(O)NR^{c42}R^{d42}, C(O)OR^{a42}, NR^{c42}R^{d42}, S(O)₂NR^{c42}R^{d42}, and S(O)₂R^{b42}, wherein the C₁₋₆ alkyl, phenyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, 5-6 membered heteroaryl, phenyl-C₁₋₃ alkyl-, (C₃₋₆ cycloalkyl)-C₁₋₃ alkyl-, (4-7 membered heterocycloalkyl)-C₁₋₃ alkyl-, (5-6 membered heteroaryl)-C₁₋₃ alkyl- of R^{4B} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{a42}, R^{b42}, R^{c42} and R^{d42} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, phenyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl, wherein the C₁₋₆ alkyl, phenyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^{a42}, R^{b42}, R^{c42} and R^{d42} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{4C} is independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, OR^{a43}, C(O)R^{b43}, C(O)NR^{c43}R^{d43}, C(O)OR^{a43}, NR^{c43}R^{d43}, S(O)₂NR^{c43}R^{d43}, and S(O)₂R^{b43}, wherein the C₁₋₆ alkyl of R^{4C} is optionally substituted with 1 or 2 independently selected R^{4D} substituents;

each R^{a43}, R^{b43}, R^{c43} and R^{d43} is independently selected from H, C₁₋₆ alkyl, and C₁₋₆ haloalkyl, wherein the C₁₋₆ alkyl of R^{a43}, R^{b43}, R^{c43} and R^{d43} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43}, attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, or 6-membered heterocycloalkyl group, wherein the 4-, 5-, or 6-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents; and

each R^{4D} is independently selected from C₁₋₃ alkyl and OH.

34. The compound of any one of claims 1, 2, and 23 to 30, or a pharmaceutically acceptable salt thereof, wherein R⁴ is selected from phenyl-C₁₋₃ alkyl-, (4-7 membered heterocycloalkyl)-C₁₋₃ alkyl-, (5-10 membered heteroaryl)-C₁₋₃ alkyl-, OR^{a4}, C(O)R^{b4}, and S(O)₂R^{b4}, wherein the phenyl-C₁₋₃ alkyl-, (4-7 membered heterocycloalkyl)-C₁₋₃ alkyl- and (5-10 membered heteroaryl)-C₁₋₃ alkyl- of R⁴ are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{a4}, R^{b4}, R^{c4}, and R^{d4} is independently selected from phenyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)-C₁₋₃ alkyl-, wherein the

phenyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)-C₁₋₃ alkyl- of R^{a4}, R^{b4}, R^{c4} and R^{d4} are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, 5-6 membered heteroaryl, (4-10 membered heterocycloalkyl)-C₁₋₃ alkyl-, OR^{a41}, S(O)₂R^{b41}, and NR^{c41}R^{d41}, wherein the C₁₋₆ alkyl, 5-6 membered heteroaryl, and (4-10 membered heterocycloalkyl)-C₁₋₃ alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{a41}, R^{b41}, R^{c41} and R^{d41} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, and 5-6 membered heteroaryl, wherein the C₁₋₆ alkyl and 5-6 membered heteroaryl of R^{a41}, R^{b41}, R^{c41} and R^{d41} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, (5-6 membered heteroaryl)-C₁₋₃ alkyl-, CN, OR^{a42}, C(O)R^{b42}, C(O)OR^{a42}, and NR^{c42}R^{d42}, wherein the C₁₋₆ alkyl, C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)-C₁₋₃ alkyl- of R^{4B} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{a42}, R^{b42}, R^{c42} and R^{d42} is independently selected from H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, and 5-7 membered heterocycloalkyl, wherein the C₁₋₆ alkyl, C₃₋₆ cycloalkyl, 5-6 membered heteroaryl, and 5-7 membered heterocycloalkyl of R^{a42}, R^{b42}, R^{c42} and R^{d42} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{4C} is independently selected from C₁₋₆ alkyl, CN, C(O)NR^{c43}R^{d43}, C(O)OR^{a43}, NR^{c43}R^{d43}, and S(O)₂R^{b43}, wherein the C₁₋₆ alkyl of R^{4C} is optionally substituted with 1 or 2 independently selected R^{4D} substituents;

each R^{a43}, R^{b43}, R^{c43} and R^{d43} is independently selected from H, C₁₋₆ alkyl, and C₁₋₆ haloalkyl, wherein the C₁₋₆ alkyl of R^{a43}, R^{b43}, R^{c43} and R^{d43} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43}, attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, or 6-membered heterocycloalkyl group,

wherein the 4-, 5-, or 6-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents;

each R^{4D} is independently selected from C₁₋₃ alkyl and OH.

35. The compound of any one of claims 1, 2, and 23-30, or a pharmaceutically acceptable salt thereof, wherein R⁴ is selected from pyridin-2-ylmethyl, 1*H*-Pyrrolo[2,3-*b*]pyridin-1-yl, 7*H*-pyrrolo[2,3-*b*]pyridin-7-yl, 2-fluorophenoxy, hydroxy(pyridin-2-yl)methyl, 2-(1-methyl-1*H*-pyrazol-4-yl)benzyl, (imidazo[1,2-*a*]pyridin-8-yl)methyl, (pyrazolo[1,5-*a*]pyridin-7-yl)methyl, (2*H*-indazol-2-yl)methyl, (1*H*-indazol-1-yl)methyl, (2,6-difluorophenyl)(hydroxy)methyl, (2,5-difluorophenyl)(hydroxy)methyl, (2,3-difluorophenyl)(hydroxy)methyl, (2-fluorophenyl)(hydroxy)methyl, (2-chlorophenyl)(hydroxy)methyl, hydroxy(phenyl)methyl, phenylsulfonyl, azetidine-1-carbonyl, benzo[d]oxazol-4-ylmethyl, 2-fluoro-6-(1-methyl-1*H*-pyrazol-5-yl)benzyl, 2-fluoro-6-((6-methyl-5-oxo-2,6-diazaspiro[3.4]octan-2-yl)methyl)benzyl, 2-fluoro-6-((6-oxohexahydropyrrolo[1,2-*a*]pyrazin-2(1*H*)-yl)methyl)benzyl, 2-fluoro-6-(((2-oxopyrrolidin-3-yl)amino)methyl)benzyl, 2-fluoro-6-((3-oxopiperazin-1-yl)methyl)benzyl, 2-fluoro-6-(((1-methyl-2-oxopyrrolidin-3-yl)amino)methyl)benzyl, 2-fluoro-6-(((2-methyl-2*H*-1,2,3-triazol-4-yl)amino)methyl)benzyl, 2-(((2-oxopyrrolidin-3-yl)amino)methyl)benzyl, amino(2,6-difluorophenyl)methyl, (2,6-difluorophenyl)(methylamino)methyl, (2,6-difluorophenyl)((2-hydroxyethyl)amino)methyl, amino(2-fluorophenyl)methyl, amino(2,6-difluorophenyl)methyl, (3-(oxazol-5-yl)pyridin-2-yl)methyl, 2-fluoro-6-(1-methyl-1*H*-pyrazol-4-yl)benzyl, (1-((1-methyl-1*H*-imidazol-4-yl)sulfonyl)pyrrolidin-2-yl)methyl, 2-((1-acetylpiperidin-4-yl)methyl)-6-fluorobenzyl, (2-(difluoromethoxy)-6-fluorophenyl)(hydroxy)methyl, 2-fluoro-6-(1-((1-methyl-1*H*-pyrazol-4-yl)methyl)-1*H*-pyrazol-4-yl)benzyl, (2-((dimethylamino)methyl)-6-fluorophenyl)(hydroxy)methyl, 2-fluoro-6-(1-(pyridin-2-ylmethyl)-1*H*-pyrazol-4-yl)benzyl, (2-fluoro-6-(pyrrolidin-1-ylmethyl)phenyl)(hydroxy)methyl, 2-fluoro-6-(1-(2-(methylsulfonyl)ethyl)-1*H*-pyrazol-4-yl)benzyl, 2-fluoro-6-((6-oxohexahydropyrrolo[1,2-*a*]pyrazin-2(1*H*)-yl)methyl)phenyl)(hydroxy)methyl, 2-fluoro-6-(1-((trans)-3-(methylamino)cyclobutyl)-1*H*-pyrazol-4-yl)benzyl, 2-(1-(2-

cyanoethyl)-1H-pyrazol-4-yl)-6-fluorobenzyl, 2-fluoro-6-(1-(2-(3-hydroxyazetidin-1-yl)-2-oxoethyl)-1H-pyrazol-4-yl)benzyl, (3-methylpyridin-2-yl)methoxy, (3-((1-(pyridin-4-ylmethyl)-1H-pyrazol-4-yl)amino)pyridin-2-yl)methyl, (3-((1-(tetrahydro-2H-pyran-4-yl)-1H-pyrazol-4-yl)amino)pyridin-2-yl)methyl, (3-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)methyl, 2-((3-hydroxypyrrolidin-1-yl)methyl)benzyl, and (6-methoxypyridin-2-yl)methyl.

36. The compound of claim 23, or a pharmaceutically acceptable salt thereof, wherein:

Cy¹ is phenyl which is substituted by 1 or 2 independently selected R⁷ substituents;

each R⁷ is independently selected from halo and CN;

R² is selected from H, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R² are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, NR^{c21}C(O)R^{b21}, NR^{c21}C(O)OR^{a21}, NR^{c21}C(O)NR^{c21}R^{d21}, S(O)R^{b21}, and S(O)₂R^{b21}, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of R^{2A} are each optionally substituted by 1, 2, or 3 independently selected R^{2B} substituents;

each R^{a21}, R²⁴¹, R^{c21}, and R^{d21} is independently selected from H, C₁₋₆ alkyl, and C₁₋₆ haloalkyl, wherein said C₁₋₆ alkyl of R^{a21}, R²⁴¹, R^{c21}, and R^{d21} is optionally substituted by 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2B} is independently selected from D, halo, OH, C₁₋₄ alkoxy, C₁₋₄ haloalkoxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, amino, C₁₋₃ alkylamino, and di(C₁₋₃ alkyl)amino;

R⁴ is selected from C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-,

OR^{a4}, C(O)R^{b4}, C(O)NR^{c4}R^{d4}, C(O)OR^{a4}, NR^{c4}R^{d4}, S(O)₂NR^{c4}R^{d4}, and S(O)₂R^{b4}, wherein the C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁴ are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{a4}, R^{b4}, R^{c4}, and R^{d4} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a4}, R^{b4}, R^{c4} and R^{d4} are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, OR^{a41}, C(O)R^{b41}, C(O)NR^{c41}R^{d41}, C(O)OR^{a41}, NR^{c41}R^{d41}, S(O)₂NR^{c41}R^{d41}, and S(O)₂R^{b41}, wherein the C₁₋₆ alkyl, 5 C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{a41}, R^{b41}, R^{c41} and R^{d41} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, phenyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl, wherein the C₁₋₆ alkyl, phenyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^{a41}, R^{b41}, R^{c41} and R^{d41} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, phenyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, 5-6 membered heteroaryl, phenyl-C₁₋₃ alkyl-, (C₃₋₆ cycloalkyl)-C₁₋₃ alkyl-, (4-7 membered heterocycloalkyl)-C₁₋₃ alkyl-, (5-6 membered heteroaryl)-C₁₋₃ alkyl-, CN, OR^{a42}, C(O)R^{b42}, C(O)NR^{c42}R^{d42}, C(O)OR^{a42}, NR^{c42}R^{d42}, S(O)₂NR^{c42}R^{d42}, and S(O)₂R^{b42}, wherein the C₁₋₆ alkyl, phenyl,

C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, 5-6 membered heteroaryl, phenyl-C₁₋₃ alkyl-, (C₃₋₆ cycloalkyl)-C₁₋₃ alkyl-, (4-7 membered heterocycloalkyl)-C₁₋₃ alkyl-, (5-6 membered heteroaryl)-C₁₋₃ alkyl- of R^{4B} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{a42}, R^{b42}, R^{c42} and R^{d42} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, phenyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl, wherein the C₁₋₆ alkyl, phenyl, C₃₋₆ cycloalkyl, 4-7 membered heterocycloalkyl, and 5-6 membered heteroaryl of R^{a42}, R^{b42}, R^{c42} and R^{d42} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{4C} is independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, OR^{a43}, C(O)R^{b43}, C(O)NR^{c43}R^{d43}, C(O)OR^{a43}, NR^{c43}R^{d43}, S(O)₂NR^{c43}R^{d43}, and S(O)₂R^{b43}, wherein the C₁₋₆ alkyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)-C₁₋₃ alkyl- of R^{4C} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

each R^{a43}, R^{b43}, R^{c43} and R^{d43} is independently selected from H, C₁₋₆ alkyl, and C₁₋₆ haloalkyl, wherein the C₁₋₆ alkyl of R^{a43}, R^{b43}, R^{c43} and R^{d43} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43}, attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, or 6-membered heterocycloalkyl group, wherein the 4-, 5-, or 6-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents; and

each R^{4D} is independently selected from C₁₋₃ alkyl and OH.

37. The compound of claim 23, or a pharmaceutically acceptable salt thereof, wherein:

Cy¹ is 3-cyanophenyl or 3-cyano-2-fluorophenyl;

R² is selected from H, 5-10 membered heteroaryl, and 4-7 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-7 membered heterocycloalkyl of R² are each optionally substituted with 1, 2, or 3 independently selected R^{2A} substituents;

each R^{2A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, and OH, wherein said C_{1-6} alkyl of R^{2A} is optionally substituted by 1, 2, or 3 independently selected R^{2B} substituents;

each R^{2B} is independently selected from D, halo, and OH;

R^4 is selected from phenyl- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl-, (5-10 membered heteroaryl)- C_{1-3} alkyl-, OR^{a4} , $C(O)R^{b4}$, and $S(O)_2R^{b4}$, wherein the phenyl- C_{1-3} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-3} alkyl- and (5-10 membered heteroaryl)- C_{1-3} alkyl- of R^4 are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{a4} , R^{b4} , R^{c4} , and R^{d4} is independently selected from phenyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl-, wherein the phenyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl- of R^{a4} , R^{b4} , R^{c4} and R^{d4} are each optionally substituted with 1, 2, or 3 independently selected R^{4A} substituents;

each R^{4A} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, 5-6 membered heteroaryl, (4-10 membered heterocycloalkyl)- C_{1-3} alkyl-, OR^{a41} , $S(O)_2R^{b41}$, and $NR^{c41}R^{d41}$, wherein the C_{1-6} alkyl, 5-6 membered heteroaryl, and (4-10 membered heterocycloalkyl)- C_{1-3} alkyl- of R^{4A} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{a41} , R^{b41} , R^{c41} and R^{d41} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, and 5-6 membered heteroaryl, wherein the C_{1-6} alkyl and 5-6 membered heteroaryl of R^{a41} , R^{b41} , R^{c41} and R^{d41} are each optionally substituted with 1, 2, or 3 independently selected R^{4B} substituents;

each R^{4B} is independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-6} cycloalkyl, 4-7 membered heterocycloalkyl, (5-6 membered heteroaryl)- C_{1-3} alkyl-, CN, OR^{a42} , $C(O)R^{b42}$, $C(O)OR^{a42}$, and $NR^{c42}R^{d42}$, wherein the C_{1-6} alkyl, C_{3-6} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl- of R^{4B} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{a42} , R^{b42} , R^{c42} and R^{d42} is independently selected from H, C_{1-6} alkyl, C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 5-7 membered heterocycloalkyl, wherein the C_{1-6} alkyl, C_{3-6} cycloalkyl, 5-6 membered heteroaryl, and 5-7 membered

heterocycloalkyl of R^{a42} , R^{b42} , R^{c42} and R^{d42} are each optionally substituted with 1, 2, or 3 independently selected R^{4C} substituents;

each R^{4C} is independently selected from C_{1-6} alkyl, CN, $C(O)NR^{c43}R^{d43}$, $C(O)OR^{a43}$, $NR^{c43}R^{d43}$, and $S(O)_2R^{b43}$, wherein the C_{1-6} alkyl, 4-7 membered heterocycloalkyl, and (5-6 membered heteroaryl)- C_{1-3} alkyl- of R^{4C} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

each R^{a43} , R^{b43} , R^{c43} and R^{d43} is independently selected from H, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein the C_{1-6} alkyl of R^{a43} , R^{b43} , R^{c43} and R^{d43} are each optionally substituted with 1 or 2 independently selected R^{4D} substituents;

or, any R^{c43} and R^{d43} , attached to the same N atom, together with the N atom to which they are attached, form a 4-, 5-, or 6-membered heterocycloalkyl group, wherein the 4-, 5-, or 6-membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{4D} substituents; and

each R^{4D} is independently selected from C_{1-3} alkyl and OH.


38. The compound of any one of claims 1, 2, and 23, or a pharmaceutically acceptable salt thereof, wherein:

R^2 is selected from 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl, wherein the 5-6 membered heteroaryl and 4-6 membered heterocycloalkyl are each optionally substituted with 1 or 2 C_{1-3} alkyl groups;

R^4 is phenyl- C_{1-3} alkyl- or pyridyl- C_{1-3} alkyl-, wherein the phenyl- C_{1-3} alkyl- and pyridyl- C_{1-3} alkyl- are each optionally substituted with 1, 2, or 3 substituents independently selected from OH and halo; and


Cy^1 is cyanophenyl.

39. The compound of any one of claims 1, 2, and 23, which is a compound of Formula (IIIb):



or a pharmaceutically acceptable salt thereof, wherein n is an integer from 0 to 4.

40. The compound of any one of claims 1, 2, and 23, which is a compound of Formula (IIIc):



or a pharmaceutically acceptable salt thereof.

41. The compound of claim 1 or 2, selected from:

4-Amino-6-(3-cyanophenyl)-N-ethyl-7-(1-ethyl-1H-pyrazol-5-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide;

4-Amino-6-(3-cyanophenyl)-N-ethyl-7-(1-propyl-1H-pyrazol-5-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide;

4-Amino-6-(3-cyanophenyl)-N-ethyl-7-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide;

4-Amino-6-(3-cyanophenyl)-N-ethyl-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide;

4-Amino-6-(3-cyanophenyl)-7-(1,1-difluoroethyl)-N-ethylpyrazolo[1,5-a]pyrazine-2-carboxamide;

4-Amino-6-(3-cyanophenyl)-N-ethyl-7-(1-(trifluoromethyl)cyclopropyl)pyrazolo[1,5-a]pyrazine-2-carboxamide;

3-(4-Amino-2-(azetidine-1-carbonyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-7-(pyrimidin-4-yl)-2-(pyrrolidine-1-carbonyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-(piperidine-1-carbonyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

4-Amino-6-(3-cyanophenyl)-N,N-diethyl-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide;

4-Amino-6-(3-cyanophenyl)-N-ethyl-N-methyl-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazine-2-carboxamide;

3-(4-Amino-2-(3-hydroxyazetidine-1-carbonyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-(azetidin-1-ylmethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

Ethyl (8-amino-6-(3-cyanophenyl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)carbamate;

3-(8-Amino-5-(6-oxo-1,6-dihydropyridin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

4-(4-Amino-6-(3-cyanophenyl)-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazin-7-yl)-5,6-dihydropyridine-1(2H)-carboxamide;

4-Amino-6-(3-cyanophenyl)-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazine-7-carbonitrile;

4-Amino-6-(3-cyanophenyl)-2-(2-fluorobenzyl)pyrazolo[1,5-a]pyrazine-3-carbonitrile;

3-(4-Amino-7-bromo-2-((2-fluorophenyl)(hydroxy)methyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-((2-fluorophenyl)(hydroxy)methyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-(3,6-dihydro-2H-pyran-4-yl)-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-(phenylamino)-7-(pyridin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-(2-(4-methylpiperazin-1-yl)pyridin-4-yl)-7-(pyridin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(pyridin-2-ylmethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-Amino-5-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-Amino-5-(1-methyl-6-oxo-1,6-dihydropyridazin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

(S)-1-(2-((8-Amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)pyrrolidine-3-carboxylic acid;

1-(2-((8-Amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)azetidine-3-carboxylic acid;

3-(2-((1*H*-Pyrrolo[2,3-*b*]pyridin-1-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(2-((7*H*-pyrrolo[2,3-*b*]pyridin-7-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-7-(4-(1-hydroxyethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-7-(4-(2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-7-(1-ethyl-1*H*-1,2,3-triazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-7-(1-methyl-1*H*-1,2,3-triazol-5-yl)-2-(pyridin-2-ylmethyl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(4-Amino-2-((2-fluorophenyl)(hydroxy)methyl)-7-(pyrimidin-4-yl)pyrazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(2-fluorophenoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(hydroxy(pyridin-2-yl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(2-(1-methyl-1*H*-pyrazol-4-yl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-5-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-5-(4-ethyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-5-(3-methylpyridin-4-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(imidazo[1,2-*a*]pyridin-8-ylmethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(pyrazolo[1,5-*a*]pyridin-7-ylmethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(2-((2*H*-Indazol-2-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(2-((1*H*-Indazol-1-yl)methyl)-8-amino-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-((2,5-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-((2,3-difluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-((2-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-((2-chlorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(hydroxy(phenyl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(phenylsulfonyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-(azetidine-1-carbonyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-amino-5-(6-hydroxypyridin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(benzo[d]oxazol-4-ylmethyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(2-fluoro-6-(1-methyl-1H-pyrazol-5-yl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

(R)-1-(2-((8-amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)-3-fluorobenzyl)-3-methylpyrrolidine-3-carboxylic acid;

3-(8-amino-2-(2-fluoro-6-((6-methyl-5-oxo-2,6-diazaspiro[3.4]octan-2-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(2-fluoro-6-((6-oxohexahydropyrrolo[1,2-*a*]pyrazin-2(1H)-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

(S)-3-(8-amino-2-(2-fluoro-6-(((2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

2-((2-((8-amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)-3-fluorobenzyl)amino)acetamide;

3-(8-amino-2-(2-fluoro-6-((3-oxopiperazin-1-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

(1S,3S)-3-((2-((8-amino-6-(3-cyanophenyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)-3-fluorobenzyl)amino)cyclobutane-1-carboxylic acid

3-(8-amino-2-(2-fluoro-6-(((1-methyl-2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(2-fluoro-6-(((2-methyl-2H-1,2,3-triazol-4-yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

(S)-3-(8-amino-2-(2-(((2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

(R)-1-(2-((8-amino-6-(3-cyanophenyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl)-3-fluorobenzyl)-3-methylpyrrolidine-3-carboxylic acid;

3-(8-amino-2-(amino(2,6-difluorophenyl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((2,6-difluorophenyl)(methylamino)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((2,6-difluorophenyl)((2-hydroxyethyl)amino)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(amino(2-fluorophenyl)methyl)-5-(4-methyloxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(amino(2,6-difluorophenyl)methyl)-5-(1-methyl-6-oxo-1,6-dihydropyridazin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((3-(oxazol-5-yl)pyridin-2-yl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(2-fluoro-6-(1-methyl-1H-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

(S)-3-(8-amino-2-(2-fluoro-6-(((1-methyl-2-oxopyrrolidin-3-yl)amino)methyl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-(2-fluoro-6-((6-methyl-5-oxo-2,6-diazaspiro[3.4]octan-2-yl)methyl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-5-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-5-(4-(2,2-difluoro-1-hydroxyethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-5-(2-methyl-4-(2,2,2-trifluoro-1-hydroxyethyl)oxazol-5-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-5-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)-2-(pyridin-2-ylmethyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-((1-((1-methyl-1H-imidazol-4-yl)sulfonyl)pyrrolidin-2-yl)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(2-(2-((1-acetylpiperidin-4-yl)methyl)-6-fluorobenzyl)-8-amino-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(1-ethyl-1H-pyrazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(2,6-dimethylpyridin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(2-((1H-pyrrolo[3,2-b]pyridin-3-yl)methyl)-4-amino-7-(pyrimidin-4-yl)pyrazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((2-(difluoromethoxy)-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(2-fluoro-6-(1-((1-methyl-1H-pyrazol-4-yl)methyl)-1H-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-((2-((dimethylamino)methyl)-6-fluorophenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(3-methyl-1H-pyrazol-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(5-([1,2,4]triazolo[4,3-a]pyridin-6-yl)-8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(oxazol-5-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(2-fluoro-6-(1-(pyridin-2-ylmethyl)-1H-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(3-methylpyridin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((2-fluoro-6-(pyrrolidin-1-ylmethyl)phenyl)(hydroxy)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(2-fluoro-6-(1-(2-(methylsulfonyl)ethyl)-1H-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-((2-fluoro-6-((6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1*H*)-yl)methyl)phenyl)(hydroxy)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile;

2-(4-(2-((8-amino-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)-3-fluorophenyl)-1*H*-pyrazol-1-yl)-*N,N*-dimethylacetamide;

2-(4-(2-((8-amino-6-(3-cyano-2-fluorophenyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-2-yl)methyl)-3-fluorophenyl)-1*H*-pyrazol-1-yl)acetamide;

3-(8-amino-2-(2-fluoro-6-(1-((*trans*)-3-(methylamino)cyclobutyl)-1*H*-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-(2-(1-(2-cyanoethyl)-1*H*-pyrazol-4-yl)-6-fluorobenzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-(2-fluoro-6-(1-(2-(3-hydroxyazetidin-1-yl)-2-oxoethyl)-1*H*-pyrazol-4-yl)benzyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-((3-methylpyridin-2-yl)methoxy)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-2-((3-((1-(pyridin-4-ylmethyl)-1*H*-pyrazol-4-yl)amino)pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-2-((3-((1-(tetrahydro-2*H*-pyran-4-yl)-1*H*-pyrazol-4-yl)amino)pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-Amino-2-((3-(1-methyl-1*H*-pyrazol-4-yl)pyridin-2-yl)methyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

(*S*)-3-(8-Amino-2-(2-((3-hydroxypyrrolidin-1-yl)methyl)benzyl)-5-(pyrimidin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-Amino-5-(imidazo[1,2-*a*]pyridin-6-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-(azetidine-1-carbonyl)-5-(3-fluoropyridin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile;

3-(8-amino-2-((2,6-difluorophenyl)(hydroxy)methyl)-5-(2,6-dimethylpyridin-4-yl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)-2-fluorobenzonitrile;

3-(8-amino-5-(1-(methyl-*d*3)-6-oxo-1,6-dihydropyridazin-3-yl)-2-(pyridin-2-yl)methyl)-[1,2,4]triazolo[1,5-*a*]pyrazin-6-yl)benzonitrile; and

3-(8-amino-2-((6-methoxypyridin-2-yl)methyl)-5-(1-methyl-6-oxo-1,6-dihydropyridazin-3-yl)-[1,2,4]triazolo[1,5-a]pyrazin-6-yl)benzonitrile,
or a pharmaceutically acceptable salt thereof.

42. A pharmaceutical composition comprising a compound of any one of claims 1 to 41, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient or carrier.

43. A method of inhibiting an activity of an adenosine receptor, comprising contacting the receptor with a compound of any one of claims 1 to 41, or a pharmaceutically acceptable salt thereof.

44. A method of treating a disease or disorder in a patient, wherein the disease or disorder is associated with abnormal expression of A2A or A2B receptors, comprising administering to said patient a therapeutically effective amount of a compound of any one of claims 1 to 41, or a pharmaceutically acceptable salt thereof.


45. The method of claim 44, wherein the disease or disorder is cancer, an inflammatory disease, a cardiovascular disease, or a neurodegenerative disease.

46. The method of claim 45, wherein the cancer is bladder cancer, lung cancer, melanoma, breast cancer, cervical cancer, ovarian cancer, colorectal cancer, pancreatic cancer, esophageal cancer, prostate cancer, kidney cancer, skin cancer, thyroid cancer, liver cancer, uterine cancer, head and neck cancer, or renal cell carcinoma.

47. The method of claim 45, wherein the cancer is non-small cell lung cancer (NSCLC), metastatic castrate-resistant prostate carcinoma (mCRPC), or colorectal carcinoma (CRC).

48. The method of claim 45, wherein the inflammatory disease is pulmonary inflammation.

49. The method of claim 48, wherein the pulmonary inflammation is bleomycin-induced pulmonary fibrosis.
50. The method of claim 45, wherein the inflammatory disease is an adenosine receptor dependent allergic reaction or adenosine receptor immune reaction.
51. The method of claim 50, wherein the adenosine receptor dependent allergic reaction is A2B receptor dependent.
52. The method of claim 45, wherein the inflammatory disease is a respiratory disorder, sepsis, reperfusion injury, or thrombosis.
53. The method of claim 45, wherein the cardiovascular disease is coronary artery disease, cerebrovascular disease, peripheral artery disease, aortic atherosclerosis, or aneurysm.
54. The method of claim 53, wherein the coronary artery disease is myocardial infarction, angina pectoris, or heart failure.
55. The method of claim 53, wherein the cerebrovascular disease is stroke or transient ischemic attack.
56. The method of claim 45, wherein the neurodegenerative disease is Parkinson's disease.
57. The method of claim 44, wherein the disease or disorder is diabetes or insulin resistance.
58. A method of treating or preventing atherosclerotic plaque formation in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a compound of any one of claims 1 to 41, or a pharmaceutically acceptable salt thereof.



(I),